




YALE
MEDICAL LIBRARY



HISTORICAL
LIBRARY

THE GIFT OF
WILLIAM W. PETER

MEDICAL LEXICON.

A
DICTIONARY
OF
MEDICAL SCIENCE;

CONTAINING

A CONCISE EXPLANATION OF THE VARIOUS SUBJECTS AND TERMS OF ANATOMY, PHYSIOLOGY,
PATHOLOGY, HYGIENE, THERAPEUTICS, PHARMACOLOGY, PHARMACY, SURGERY, OB-
STETRICS, MEDICAL JURISPRUDENCE, AND DENTISTRY; NOTICES OF CLI-
MATE, AND OF MINERAL WATERS; FORMULÆ FOR OFFICINAL,
EMPIRICAL, AND DIETETIC PREPARATIONS;

WITH

THE ACCENTUATION AND ETYMOLOGY OF THE TERMS.

AND

THE FRENCH AND OTHER SYNONYMES;

SO AS TO CONSTITUTE

A FRENCH AS WELL AS ENGLISH MEDICAL LEXICON.

BY


ROBLEY DUNGLISON, M.D., LL.D.,

EMERITUS PROFESSOR OF INSTITUTES OF MEDICINE AND MEDICAL JURISPRUDENCE IN THE
JEFFERSON MEDICAL COLLEGE OF PHILADELPHIA.

THOROUGHLY REVISED

AND

VERY GREATLY MODIFIED AND AUGMENTED.



PHILADELPHIA:
HENRY C. LEA.

1868.

Entered, according to Act of Congress, in the year 1865, by

BLANCHARD AND LEA,

in the Clerk's Office of the District Court of the United States for the Eastern District of
Pennsylvania.

COLLINS, PRINTER.



PREFACE.

THE author has again been required to subject his Medical Lexicon to a thorough revision. The progress of Medical Science, and the consequent introduction of new subjects and terms, demanded this; and he has embraced the occasion to render more complete the etymology and accentuation of the terms. On no previous revision has so much time and labor been expended by him. Some idea may be formed of this, from the fact, that although the page has been augmented in all its dimensions, not fewer than between sixty and seventy pages have been added.

As the author has remarked on former occasions, it has ever been his ardent wish to make the work a satisfactory and desirable—if not indispensable—lexicon, in which the inquirer may search, without disappointment, for every term that has been legitimated in the nomenclature of the science; and he confidently presents this edition as having more claims on the attention of the practitioner and student than its predecessors.

Once more the author gladly seizes the opportunity afforded him to express his grateful acknowledgments for the vast amount of favor which has been extended to the Dictionary.

ROBLEY DUNGLISON.

PHILADELPHIA, 1116 Girard St. January, 1865.

P R E F A C E

TO THE SECOND EDITION.

THE present undertaking was suggested by the frequent complaints, made by the author's pupils, that they were unable to meet with information on numerous topics of professional inquiry,—especially of recent introduction,—in the medical dictionaries accessible to them.

It may, indeed, be correctly affirmed, that we have no dictionary of medical subjects and terms which can be looked upon as adapted to the state of the science. In proof of this, the author need but remark, that he has found occasion to add several thousand medical terms, which are not to be met with in the only medical lexicon at this time in circulation in the country.

The present edition will be found to contain many hundred more terms than the first, and to have experienced numerous additions and modifications.

The author's object has not been to make the work a mere lexicon or dictionary of terms, but to afford, under each, a condensed view of its various medical relations, and thus to render the work an epitome of the existing condition of medical science. In its preparation he has freely availed himself of the English, French and German works of the same nature, and has endeavored to add every subject and term of recent introduction, which has fallen under his notice; yet, with all his care, it will doubtless be found that subjects have been omitted. The numerous additions, however, which he has made, and his strong desire to be useful, "by removing rubbish and clearing obstructions from the paths through which learning and genius press forward to conquest and glory," will, he trusts, extenuate these and other objections that might be urged against the work; especially when the toil which every compiler of a dictionary must endure, is taken into consideration; a toil which has been so forcibly depicted by the great English Lexicographer, as well as by the distinguished SCALIGER, [in an epigram written after he had made an index to the "Inscriptions" of GRUTER:

*"Si quem dura manet sententia Judicis olim,
Damnatum ærumnis suppliciisque caput;
Hunc neque fabрили lassent ergastula massâ
Nec rigidas vexent fossa metalla manus,
Lexica contextat; nam cætera quid moror? Omnes
Pœnarum facies hic labor unus habet."*

Thus freely rendered in French:—]

*"Si quelqu'un a commis quelque crime odieux,
S'il a tué son père, ou blasphémé les Dieux,
Qu'il fasse un Lexicon: s'il est supplice au monde
Qui le punisse mieux, je veux que l'on me tonde."*

EXPLANATION.

If the simple synonymy of any term be desired, a mere reference to the term may be sufficient; but if farther information, as to meaning or accentuation, be needed, it may be obtained under the term referred to. For instance, under *Cimicifuga*, reference is made to *Actæa racemosa*, as a synonyme. This may be sufficient for the inquirer; but should he desire to know the accentuation and properties of the plant, he must turn to *Actæa racemosa*. In the same manner, the French word *Tronc* is defined *Trunk*. This may be sufficient: should it not, the requisite information may be found under *Trunk*. In terms of more words than one, the definition is generally given under the more significant word; but, if not, it will be found under the other:—*Respiratory murmur*, for example, is defined under *Murmur*; *Shaking palsy*, under *Palsy*; *Musculus succingens*, under *Musculus*; *Ocular spectrum*, under *Spectrum*.

When the orthography of the French term is the same as that of the English, or with the mere difference of an accented vowel in the former, the term is not always repeated.

ABBREVIATIONS ARBITRARILY EMPLOYED.

Arab.	Arabic.	Old Eng.	Old English.
Belg.	Belgic.	Ord.	Order.
Celt.	Celtic.	P.	Portuguese.
Ch.	Chaussier.	Part.	Participle.
Corr.	Corrupted.	Peruv.	Peruvian.
D.	Dutch.	Ph. B.	British Pharmacopœia.
Da.	Danish.	Ph. D.	Pharmacopœia of Dublin.
Dim.	Diminutive.	Ph. E.	“ Edinburgh.
E.	English.	Ph. L.	“ London.
F.	French.	Ph. P.	“ Paris.
Fah.	Fahrenheit.	Ph. U. S.	“ United States.
Fam.	Family.	Pl.	Plural.
Fem.	Feminine.	Priv.	Privative.
G.	German.	Pron.	Pronounced.
Gael.	Gaelic.	Prov.	Provincial.
Gen.	Genitive.	Q. v.	Quod vide, ‘which see.’
Heb.	Hebrew.	R.	Réaumur.
I.	Italian.	S.	Spanish.
Icel.	Icelandic.	S.g. & Sp.gr.	Specific Gravity.
Imp. Meas.	Imperial Measure.	Sax.	Anglosaxon.
Ind.	Indian.	Sc.	Scottish.
Ir.	Irish.	Sex. Syst.	Sexual System.
L.	Latin.	Sing.	Singular.
Linn.	Linnaeus.	Su. G.	Suiogothic.
Masc.	Masculine.	Sw.	Swedish.
Mex.	Mexican.	Teut.	Teutonic.
Nat. Ord.	Natural Order.	Vulg.	Vulgarly or Commonly.
Neg.	Negative.	W.	Welsh.



A

DICTIONARY

OF

MEDICAL SCIENCE.

A.

A—ABBREVIATION.

A, before a consonant; *An* before a vowel, *a*, *av*, have, in the compound medical terms, a privative negative or debasing signification, like that of the particles *in*, *im*, *un*, *ir*, in English. Thus: *Stheni'a* means strength; *Astheni'a*, want of strength; *Anæmia*, want of blood, etc. *Ab*, *Abs* and *An*, as a prefix, have like signification. Occasionally, in compound words, they have an intensive meaning.

AACHEN, Aix-la-Chapelle.

A, or *AA*. See Abbreviation.

AANDE, Breath.

AARON, *Arum maculatum*.

AARZIL, MIN'ERAL WA'TERS OF. A is in the canton of Berne, in Switzerland. The chief spring contains chlorides of calcium and sodium, sulphates of lime and soda, oxyd of iron, and sulphohydric acid gas.

ASMUS, Anhelatio.

AB, as a prefix, see A.

ABACH, MIN'ERAL WA'TERS OF. A hydro-sulphuretted saline spring not far from Ratisbon or Regensburg, in Bavaria.

ABASSEMENT, (from *bas*, 'low.') Depression; *a. de la Cataracte*, see Cataract—*a. de la Matrice*, Prolapsus uteri.

ABASSEUR DE L'AILE DU NEZ, Depressor alæ nasi—*a. de l'angle des lèvres*, Depressor anguli oris—*a. de la langue*, Glossocatochus—*a. de la lèvre inférieure*, Depressor labii inferioris—*a. de la mâchoire inférieure*, Digastri-cus—*a. de l'œil*, Rectus inferior oculi.

ABALIENATIO MENTIS, Insanity.

ABALIENATUS. *Corruptus*, Corrupted; (*ab*, and *alienus*, 'different.') *Membra abalienata*. Limbs dead or benumbed.

ABAN'GA. Name given by the inhabitants of St. Thomas to the fruit of a palm tree, the seeds of which they consider very useful in diseases of the chest, in the dose of three or four, two or three times a day.

ABAPTIST'A. *Abaptis'ton* or *Abaptis'tum*, (*a*, and *βαπτίζω*, 'to plunge.') The old trepan, the conical shape of which prevented it from plunging suddenly into the cavity of the cranium.

ABAPTISTON, Abaptista.

ABAPTISTUM, Abaptista.

ABAREMO-TEMO. A Brazilian tree, which grows in the mountains, and appears to be a mimosa. Piso relates that the decoction of its bark, which is bitter and astringent, was applied in that country to ulcers of a bad character.

ABARNAHAS, Magnesia.

ABARTICULATIO, (*ab*, and *articulus*, 'a joint.') Diarthrosis, and Synarthrosis.

ABÂTARDISSEMENT, (from *bâtard*, *bastard*.) Degeneration.

ABATTEMENT, (from *battre*, 'to beat.') Prostration.

ABATIS, Giblets.

ABBE COURT, MIN'ERAL WA'TERS OF. A chalybeate spring, six leagues from Paris, and one from Poissy. It was once much frequented, but is now abandoned.

ABBEVILLE, MIN'ERAL WA'TERS OF. An acidulous chalybeate at Abbeville, in the department of Somme, France.

ABBREVIATION, *Abbrévia'tio*, *Brachyn'sis*, *Brachys'mos*, *Abbrévia'tura*. (F.) *Abbréviation*, (*brevis*, 'short.') Abbreviations are chiefly used in medicinal formulæ. They are by no means as frequently employed now as of old, when every article had its appropriate symbol. The following are some of the abbreviations which have been or are employed:

R. *Recipe*, Take.

A. *AA*, ANA, (*ava*) *utriusque*, of each.

ABDOM. *Abdomen*.

ABS. FEBR. *Absente febre*, In the absence of fever.

AD. or ADD. *Adde* or *addatur*.

AD. LIB. *Ad libitum*, At pleasure.

ADMOV. *Admoveatur*, Let it be applied.

ALTERN. HOR. *Alternis horis*, Every other hour.

ALV. ADSTRICT. *Alvo adstrictâ*, The bowels being confined.

AQ. *Aqua*, Water.

AQ. BULL. *Aqua bulliens*, Boiling water.

AQ. COMM. *Aqua communis*, Common water.

AQ. FERV. *Aqua fervens*, Hot water.

AQ. FONT. *Aqua fontis*, Spring water.

AQ. MARIN. *Aqua marina*, Sea water.

B. A. *Balneum arenæ*, A sand-bath.

BALS. *Balsamum*, Balsam.

BB. BBDS. *Barbadensis*, Barbadoes.

BIB. *Bibe*, Drink.

BIS IND. *Bis indies*, Twice daily.

B. M. *Balneum mariæ*. (F.) *Bain marie*, A water bath.

BOL. *Bolus*.

BULL. *Bulliat*, Let it boil.

BUT. *Butyrum*, Butter.

B. V. *Balneum vaporis*, A vapor bath.

C. *Congius*, A gallon.

- CÆRUL. *Cæruleus*, Blue.
 CAP. *Capiat*, Let him take.
 C. C. *Cornu cervi*, Hartshorn.
 C. C. U. *Cornu cervi ustum*, Burnt hartshorn.
 C. M. *Cras mane*, To-morrow morning.
 C. N. *Cras nocte*, To-morrow night.
 C. V. *Cras vespere*, To-morrow evening.
 COCHL. *Cochleare*, A spoonful.
 COCHL. AMPL. *Cochleare amplum*, A large spoonful.
 COCHL. INF. *Cochleare infantum*, A child's spoonful.
 COCHL. MOD. or MED. *Cochleare modicum* or *medium*, A dessertspoonful.
 COCHL. PARV. *Cochleare parvum*, A teaspoonful.
 COL. *Cola*, and *Colaturæ*, Strain, and to the strained.
 COMP. *Compositus*, Compound.
 CONF. *Confectio*, Confection.
 CONS. *Conserva*, Conserve.
 CONT. *Continetur*, Let it be continued.
 COQ. *Coque*, Boil.
 CORT. *Cortex*, Bark.
 CRAFT. *Crastinus*, For to-morrow.
 CUJ. *Cujus*, Of which.
 CUJUSL. *Cujuslibet*, Of any.
 CYATH. *Cyathus*, A glassful.
 CYATH. THEÆ, A teacupful.
 D. *Dosis*, A dose.
 D. et S. *Detur et signetur* (placed at the end of a prescription.)
 D. D. *Detur ad*, Let it be given in or to.
 DEAUR. PIL. *Deauretur pilula*, Let the pill be gilded.
 DEB. SPISS. *Debita spissitudo*, A due consistence.
 DEC. *Decanta*, Pour off.
 DECUB. *Decubitus*, Lying down, going to bed.
 DE D. IN D. *De die in diem*, From day to day.
 DEJ. ALV. *Dejectiones alvi*, Alvine evacuations.
 DEP. *Depuratus*, Purified.
 DEST. *Destilla*, Distil.
 DET. *Detur*, Let it be given.
 DIEB. ALTERN. *Diebus alternis*, Every other day.
 DIEB. TERT. *Diebus tertiis*, Every third day.
 DIG. *Digeratur*, Let it be digested.
 DIL. *Dilutus*, Dilute.
 DIM. *Dimidius*, One-half.
 DIV. *Divide*, Divide.
 DONEC ALV. SOLUT. FUER. *Donec alvus soluta fuerit*, Until the bowels are opened.
 DRACH. *Drachma*, A drachm.
 EJUSD. *Ejusdem*, Of the same.
 ENEM. *Enema*, A clyster.
 EXHIB. *Exhibeatur*, Let it be exhibited.
 EXT. SUPER ALUT. *Extende super alutam*, Spread upon leather.
 F. *Fiat*, Let it be made.
 F. PIL. *Fiat pilula*, Make into a pill.
 F. VENÆS. or F. VS. *Fiat venæsectio*, Let bleeding be performed.
 FEB. DUR. *Febre durante*, The fever continuing.
 FEM. INTERN. *Femoribus internis*, To the inside of the thighs.
 FIST. ARMAT. *Fistula armata*, A bag and pipe—a clyster pipe and bladder fitted for use.
 FL. *Fluidus*, and *Flores*, Fluid, and Flowers.
 FRUST. *Frustillatim*, In small pieces.
 GEL. QUAVIS. *Gelatinâ quavis*, In any kind of jelly.
 G. G. G. *Gummi guttæ Gambiæ*, Gamboge.
 GR. *Gravum*, A grain.
 GT. *Gutta*, A drop. GTT. *Guttæ*, Drops.
 GTT. or GUTT. QUIBUSD. *Guttis quibusdam*, With some drops.
 GUM. *Gummi*, Gum.
 GUTTAT. *Guttatim*, By drops.
 HOR. DECUB. *Horâ decubitis*, At bed-time.
 HOR. INTERM. *Horis intermediis*, At intermediate hours.
 H. S. *Horâ somni*, At bed-time.
 INF. *Infunde*, Infuse.
 IND. *Indies*, Daily.
 INJ. ENEM. *Injiciatur enema*, Let a clyster be given.
 IN PULM. *In pulmento*, In gruel.
 JUL. *Julepus*, A julep.
 LAT. DOL. *Lateri dolenti*, To the pained side.
 LB. and LIB. *Libra*, A pound weight.
 LIB. LLB. *Libræ*, Pounds.
 LIQ. *Liquor*.
 M. *Misce*, Mix.
 MAC. *Macera*, Macerate.
 MAN. *Manipulus*, A handful.
 MAN. PRIM. *Manè primo*, Early in the morning.
 MIC. PAN. *Mica panis*, Crumb of bread.
 MIN. *Minimum*, The 60th part of a drachm by measure.
 MITT. *Mitte*, Send.
 MITT. SANG. *Mittatur sanguis*, Let blood be drawn.
 MOD. PRÆSCRIPT. *Modo præscripto*, In the manner directed.
 MOR. SOL. *More solito*, In the usual manner.
 MUC. *Mucilago*, Mucilage.
 N. M. *Nux moschata*, Nutmeg.
 O. OCTARIUS, A pint.
 OL. *Oleum*, Oil.
 OL. LINI S. I. *Oleum lini sine igne*, Cold-drawn linseed oil.
 OMN. BID. *Omni biduo*, Every two days.
 OMN. BIH. *Omni bihorio*, Every two hours.
 OMN. HOR. *Omni horâ*, Every hour.
 OMN. MAN. *Omni manè*, Every morning.
 OMN. NOCTE, Every night.
 OMN. QUADR. HOR. *Omni quadrante horæ*, Every quarter of an hour.
 O. O. O. *Oleum olivæ optimum*, Best olive oil.
 OV. *Ovum*, An egg.
 OX. *Oxymel*.
 OZ. *Uncia*, An ounce.
 P. *Pondere*, By weight, and *Pars*, A part.
 P. and PUG. *Pugillus*, A pugil.
 P. Æ. *Partes æquales*, Equal parts.
 PART. VIC. *Partitis vicibus*, In divided doses.
 PERACT. OP. EMET. *Peractâ operatione emetici*, The operation of the emetic being over.
 PIL. *Pilula*, A pill. PIL. or PILL. *Pilulæ*, Pills.
 POST. SING. SED. LIQ. *Post singulas sedes liquidas*, After every liquid evacuation.
 POT. *Potio*, A potion.
 P. P. *Pulvis patrum*, Jesuits' bark.
 P. RAT. ÆTAT. *Pro ratione ætatis*, According to the age.
 P. R. N. *Pro re natâ*, As occasion may be.
 PULV. *Pulvis*, A powder.
 Q. P. *Quantum placeat*, As much as may please.
 Q. S. *Quantum sufficit*, As much as is sufficient.
 Q. V. *Quantum volueris*, As much as you wish.
 QUOR. *Quorum*, Of which.
 RAD. *Radix*, Root.
 RAS. *Rasuræ*, Shavings.
 RECT. *Rectificatus*, Rectified.
 RED. or REDIG. IN PULV. *Redactus in pulverem*, or *Redigatur in pulverem*, Powdered, or Let it be powdered.
 REG. UMBIL. *Regio umbilici*, The umbilical region.
 REPET. *Repetatur*, Let it be repeated.
 S. *Signa*, Write, or Give directions.
 S. A. *Secundum artem*, According to art.
 SEM. *Semen*, Seed.
 SEMI-DR. *Semidrachma*, half a drachm.
 SEMI-H. *Semihora*, Half an hour.
 SERV. *Serva*, Keep, preserve.
 SESQUIH. *Sesquihora*, An hour and a half.
 SESUNC. *Sesuncia*, An ounce and a half.
 SI NON VAL. *Si non valet*, If it does not answer.
 SI OP. SIT. *Si opus sit*, If there be need.

Si VIR. PERM. *Si vires permittant*, If the strength will permit.

SING. *Singulorum*, Of each.

SOLV. *Solve*, Dissolve.

SP. and SPIR. *Spiritus*, Spirit.

SS. *Semi*, One half.

ST. STET. Let it stand.

SUB FIN. COCT. *Sub finem coctionis*, Towards the end of the boiling.

SCM. SUMAT. Let him take; also, *Summitates*, The tops.

S. V. SPIRITUS VINI, Spirit of wine.

S. V. R. SPIRITUS VINI RECTIFICATUS, Rectified spirit of wine.

S. V. T. SPIRITUS VINI TENUIOR, Proof spirit of wine.

SYR. SYRUPUS, Syrup.

TEMP. DEXT. *Tempori dextro*, To the right temple.

T. O. TINCTURA OPII, Tincture of opium.

TR. TRA. and TINCT. *Tinctura*, Tincture.

TRIT. TRITURA, Triturate.

V. O. S. or VIR. OV. SOL. *Vitello ovi solutus*, Dissolved in the yolk of an egg.

VS. VENÆSECTIO, Venesection.

Z. Z. Anciently *myrrh*: now *zinziber* or ginger.

lb. Libra, A pound. **lb. Pounds**.

℥. Uncia, An ounce.

℥ss. Fluiduncia, A fluidounce.

℥. Drachma, A drachm.

℥ss. Fluidrachma, A fluidrachm.

℥. Scrupulum, A scruple.

℥. Minimum, A minim.

ss, Semissis, or half; **iss, sesqui**, one and a half.

j, one; ij, two; iij, three; iv, four, &c. See Symbol.

The same system is not always followed in abbreviating. The subjoined will exhibit the usual mode:

R	<i>Infus. Colomb.</i>	f ℥ss.
	<i>Tinct. Gent. Comp.</i>	f ℥iii.
	<i>Syr. Cort. Aurant.</i>	f ℥ii.
	<i>Tinct. Caps.</i>	f ℥l. M.

Capt. coch. ij. p. r. n.

This, written at length, is as follows:

Recipe

Infusi Colombæ fluiduncias quinque cum semisse.

Tincturæ Gentianæ Compositæ fluidrachmas tres.

Syrupi Corticis Aurantiorum fluidrachmas duas.

Tincturæ Capsici fluidrachmam.

Misce.

Capiat cochlearia duo pro re natâ.

In the United States the directions are always written in English.

ABCÈS, Abscess—*a. Aigu*, see Abscess.

ABCÈS EN BOUTON DE CHEMISE, *Shirt-button or Shirtstud Abscess*. A deep-seated mammary abscess, following the septa of the breast, and appearing under the skin, so as eventually to give rise to one or more subcutaneous abscesses, without losing the original character of submammary supuration. A cavity exists between the integuments and the gland—another, larger, between the mamma and the chest—the two communicating by a passage, which is generally narrow—the whole having the exact appearance of a shirtstud.—Velpéau.

ABCÈS CHAUD, see Abscess—*a. Chronique*, see Abscess—*a. par Congestion*, see Abscess—*a. Consécutif*, Abscess, metastatic—*a. Diathésique*, see Abscess—*a. Froid*, see Abscess—*a. Métastatique*, Abscess, metastatic—*a. Rétro-utérine*, see Retro-uterine—*a. Scrofuleux*, see Abscess—*a. Soudain*, see Abscess.

ABDO'MEN, gen. *Abdo'minis*, (from *abdere*, 'to conceal'; because it conceals the viscera; by some, fancifully presumed to be from *abdere*, 'to conceal,'

and *omentum*, 'the caul.' Fr in *abdo* is formed *abdomen*, as from *lego*, *legu-men*.) *Etron*, *Hypogastrium*, *Physcæ*, *Hypocœlium*, *Epis'chion*, *Lap'tura*, *Hypocœ'lium*, *Gaster*, *Ara'ca*, *Mirach*, *Nedy's*, *Neia'ra*, *Nei'ra*, *Abdu'men*, *Venter*, *Venter imus*, *Venter in'fimus*, *Alvus*, *Uterus*, *The belly*, (Sc.) *Kyte*, *Penche*, (Prov.) *Baggie*, *Hag*, *Pook*, (F.) *Ventre*, *V. inférieur*, *Bas ventre*. The largest of the three splanchnic cavities, bounded, above, by the diaphragm; below, by the pelvis; behind, by the lumbar vertebra; and at the sides and fore part, by muscular expansions. It is distinguished into three anterior regions, from above to below; viz. the epigastric, umbilical, and hypogastric, each of which is itself divided into three others, one middle, and two lateral: thus, the *epigastric region* comprises the *epigastrium* and *hypochondria*; the *umbilical*, the *umbilicus* and *flanks* or *lumbar regions*; and the *hypogastric*, the *hypogastrium* and *iliac regions*. None of these regions has its limits well defined. The chief viscera, contained in the cavity of the abdomen—*Cœ'lia*, *Cavum Abdom'inis*—are the stomach, intestines, liver, spleen, pancreas, kidneys, &c. It is lined by the peritoneum.

ABDOMEN, PENDULOUS, *Physconia*.

ABDOM'INAL, *Abdomina'lis*, *Ventra'lis*, *Ventral*. That which belongs to the Abdomen, as *abdominal muscles*, *abdominal viscera*, &c.

ABDOMINAL RING, *Inguinal ring*.

ABDOMINIS EXPLORATIO, *Abdominoscopia*.

ABDOMINOSCOPIA, *Gastrosco'pia*, *Laparosco'pia*, *Abdom'inis Explora'tio*. *Abdom'inoscopy*. A hybrid word, from *abdomen*, and *σκοπεω*, 'I view.' Examination of the lower belly as a means of diagnosis. See Auscultation.

ABDOM'INOUS, *Ventrio'sus*, *Ventro'sus*, (from *abdomen*,) *Big-bellied*, *Big-paunched*, *Ventrip'otent*. Having a large abdomen.

ABDUCENS LABIORUM, *Levator anguli oris*.

ABDUCENTES, *Motor oculi externus*.

ABDUCTEUR, *Abductor*—*a. de l'œil*, *Rectus externus oculi*—*a. de l'oreille*, *Abductor auris*—*a. du gros orteil*, *Abductor pollicis pedis*—*a. du petit orteil*, *Abductor minimi digiti pedis*—*a. court du ponce*, *Abductor pollicis brevis*—*a. long du ponce*, *Abductor longus pollicis*.

ABDUCTION, *Abduc'tio*, (from *abducere*, 'to separate,' *ab* and *ducere*, 'to lead.') The movement which separates a limb or other part from the axis of the body.

The word has also been used synonymously with *Abrup'tio*, *Apag'ma*, *Apoclas'ma*, a fracture near the articular extremity of a bone, with separation of the fragments.

ABDUCTOR, gen. *Abducto'ris*, (F.) *Abducteur*. A muscle which moves certain parts by separating them from the axis of the body.

ABDUCTOR AUCULARIS, *Abductor auris*—*a. Indicis pedis*, *Prior indicis pedis*, *Posterior indicis pedis*—*a. Medii digiti pedis*, *Prior medii digiti pedis*—*a. Minimi digiti*, *Flexor parvus minimi digiti*—*a. Minimi digiti*, *Prior minimi digiti*—*a. Oculi*, *Rectus externus oculi*—*a. Pollicis manus*, and *a. Brevis alter*, *Abductor pollicis brevis*.

ABDUCTOR AU'RIS, *Abductor auricu'laris*. (F.) *Abducteur de l'oreille*. A portion of the *posterior auris*, whose existence is not constant, which passes from the mastoid process to the concha.

ABDUCTOR IN'DICIS, *Semi-interos'seus in'dicis*. A muscle which arises from the os trapezium and metacarpal bone of the thumb, and is inserted into the first bone of the forefinger. Its use is to bring the forefinger towards the thumb.

ABDUCTOR LONGUS POLLICIS, *Abductor pollicis longus*.

ABDUCTOR MIN'IMI DIG'ITI, *Carpophalan'geus min'imi digiti*, *Exten'sor ter'tii interno'dii minimi digiti*—(Douglas.) *Hypoth'enar mi'nor metacar'peus*, (F.) *Carpo-phalangien du petit doigt*. See

Flexor parvus. It originates, fleshy, from the os pisiforme, and from the annular ligament near it; and is inserted, tendinous, into the inner side of the base of the first bone of the little finger. *Use*, to draw the little finger from the rest.

ABDUCTOR MINIMI DIGITI PE'DIS, Calcosubphalan'geus minimi digiti, Calcaneo-phalangien du petit orteil, Parath'énar ma'jor.—(By Winslow, the muscle is divided into two portions.—*Parathénar major* and *metatarsus*.) *Calcaneo-sous-phalangien du petit orteil*—(Ch.) (F.) *Abducteur du petit orteil.* This muscle forms the outer margin of the sole of the foot, and is immediately beneath the plantar aponeurosis. It arises, tendinous and fleshy, from the outer side of the protuberance of the os calcis, and from the root of the metatarsal bone of the little toe, and is inserted into the outer part of the root of the first bone of the little toe. *Use*, to draw the little toe outwards.

ABDUCTOR POL'ICIS BRE'VIS, Abductor Pollicis Ma'nús, Scaphocarpusuperphalan'geus Pollicis, Sus-phalangien du ponce, A. pollicis ma'nús and A. bre'vis al'ter—(Albinus.)—(F.) *Abducteur court du ponce, Carpo-sus-phalangien du ponce*—(Ch.) A short, flat, triangular muscle, which arises from the anterior surface of the os scaphoides and the annular ligament of the carpus, and terminates at the outside of the upper extremity of the first phalanx of the thumb. A particular portion, on the inner side of this muscle, is called, by Albinus, *Abductor brevis alter*.

ABDUCTOR POL'ICIS LONGUS, A. P. L. Manús, Extensor os'sis metacarpi pollicis manús, Extensor pri'mi internodii—(Douglas.) *Extensor pri'mus Pollicis, Cubito-radi-sus métacarpien du ponce, Cubito-sus-métacarpien du ponce*,—(Ch.) (F.) *Abducteur long du ponce.* A long, thin muscle, arising from the posterior surface of the ulna, radius, and interosseous ligament, and inserted at the outer side of the upper extremity of the first metacarpal bone.

ABDUCTOR POL'ICIS PE'DIS, Calcosubphalan'geus Pollicis. (F.) *Abducteur du gros orteil.* This muscle arises, fleshy, from the anterior and inner part of the protuberance of the os calcis, and tendinous from the same bone where it joins with the os naviculare. It is inserted, tendinous, into the internal os sesamoidem and root of the first bone of the great toe. *Use*, to pull the great toe from the rest.

The name *Abductor* has been given also to all those interosseous muscles of the hand and foot, which perform the motion of abduction on the fingers or toes, and to muscles which execute the same function in other parts of the body.

ABDUMEN, Abdomen.

ABEB'E'US, (αἰβ'ε'ιος, from a, neg., and β'ε'βαιος, 'firm,') Infr'mus, Deb'ilis. Weak, infirm, unsteady.

ABE'ILLE, (from apicula, dim. of apis, 'a bee,') Bee.

ABELME'LUCH. One of the names of the Ricinus, according to some authors.—Prosper Alpinus says that a tree, which grows about Mecca, is so called. Its seeds, which are black and oblong, are said to be a most violent cathartic.

ABELMOSCHUS, Hibiscus abelmoschus—a. *Eseulentus, Hibiscus eschulentus*—a. *Moschatus, Hibiscus abelmoschus.*

ABELMUSK, Hibiscus abelmoschus.

ABENSBURG, MIN'ERAL WA'TERS OF. A. is a city of Bavaria, where there is a cold, sulphureous spring.

ABERRATIO, gen. Aberratio'nis, Aberration—a. *Lactis, Galactoplasia*—a. *Loci, Heterotopia*—a. *Mensium, Menstruation, vicarious*—a. *Menstruorum, Menstruation, vicarious*—a. *Temporis, Heterochronia.*

ABERRA'TION, Aberra'tio, (from aberrare,

(ab and errare, 'to stray, 't: wander from.') This word has several meanings.

1. The passage of a fluid of the living body into an order of vessels not destined for it. In this sense it is synonymous with the *Error Loci* of Boerhaave. 2. The flow of a fluid towards an organ different from that to which it is ordinarily directed; as in cases of vicarious hemorrhage. *Aberrations of sense or judgment* are certain errors in the perceptions, or certain derangements of the intellectual faculties.

The word is used in optics to designate the dispersion of the rays of light in passing through a lens.

ABERRATION, CHROMATIC, Aberration of Refrangibility.

ABERRATION OF REFRAINGIBIL'ITY, Chromat'ic aberrat'ion, (F.) Aberration de Réfrangibilité, Chromatisme, exists, when, as in a common lens, the rays that pass near the circumference of the lens are decomposed, so that a coloured image is observed. This aberration in the human eye is corrected by the iris, which does not permit the rays to fall near the circumference of the lens, and also by the crystalline lens itself, which, owing to its structure, serves the purposes of an achromatic glass.

ABERRATION, SPHERICAL, Aberration of sphericity.

ABERRATION OF SPHERIC'ITY or spher'ical aberrat'ion takes place, when the rays, as in a common lens, which pass through the centre of the lens, and those which pass near the circumference, are unequally refracted, so that they do not meet at a common focus.

This aberration of sphericity in the human eye is corrected by the iris and lens.

ABESSI, Realgar.

ABEVACUA'TIO, gen. Abevacuatio'nis, Apoce-no'sis, (ab, and evacuare, 'to empty,') An evacuation. A partial or imperfect evacuation. By some it is applied to an immoderate evacuation.

ABHAL. A fruit well known in India, and obtained from a species of cypress. It passes for an emmenagogue.

ABIES, gen. Abi'etis, Pinus picea—a. *Balsamea, Pinus balsamea.*

ABIES BALSAMIFERA, Pinus Balsamea—a. *Canadensis, Pinus Canadensis*—a. *Excelsa,* see *Pinus abies*—a. *Gallica, Pinus picea*—a. *Larix, Pinus larix*—a. *Pectinata, Pinus picea*—a. *Picea, Pinus picea*—a. *Rubra, Pinus rubra.*

ABIETIQUE, Terebinthinate.

ABIGA, Ajuga, Teucrium Chamæpitys.

ABIOSIS, (a, and β'ιος, 'life,') Death.

ABIOTUS, (αβ'ιωτος) Conium maculatum.

ABIRRITA'TION, Abirrita'tio, (ab, and irritatio, 'irritation,') This word strictly means absence or defect of irritation. The disciples of Broussais used it to indicate a pathological condition, opposite to that of irritation. It may be considered as synonymous with debility, asthenia, &c.

ABLACTATIO, (ab, and lactare, 'to suckle,') Weaning.

ABLASTES, (αβλαστης, from a, and βλαστος, 'a bud,') Sterile.

ABLATIO, (ab, and ferre, latum, 'to take,') Ex-tirpation.

ABLEPH'ARUS, (a, privative, and βλεφαρον, 'eyelid,') One who has no eyelids.

ABLEPSIA (αβλεψια), Cæcitas.

ABLUENTIA, (ablueré, (ab, and luere, 'to wash,') Detergents.

ABLUENTS, Detergents.

ABLUTION, Ablu'tio, Apion'esis, Cataclysmus. A name given to legal ceremonies in which the body is subjected to particular affusions. Ablution (especially of the extremities) with cold or tepid water is employed, therapeutically, to re-

duce febrile heat. Also, the washing by which medicines are separated from the extraneous matters mixed with them.

ABNORMAL, Abnormous.

ABNORMITY, Anomalial.

ABNORMOUS, *Abnor'mis*, *Enor'mis*, *Abnor'mal*, (F.) *Anormal*, (*ab*, and *norma*, 'rule.') Not conformable to rule; irregular.

ABOLITION, *Aboli'tio*, destruction or suppression, (*ab*, and *luere* (?) 'to wash.') A word, often employed, especially by the French, to express the complete suspension of any symptom or function. *Abolition of the sight*, e.g. is the complete loss of sight.

ABOMA'SUS, *Aboma'sum*, *Enys'tron*, *Ventriculus intestinalis*, *Ren'net*, (Sc.) *Roddikin*, (F.) *Cuilette*. The lowermost or fourth stomach of ruminating animals.

ABOMINATIO, Disgust.

ABONDANCE, see *Plethora*.

ABORSIO, Abortion.

ABORSUS, Abortion.

ABORTICIDIUM, Fœticide.

ABORTIF, Abortive.

ABORTIFACIENS, Abortive.

ABORTION, *Abor'tus*, *Abor'eus*, *Abor'sio*, *Dys'to'cia aborti'va*, *Omoto'cia*, *Paracye'sis abor'tus*, *Amblo'sis*, *Amblo'ma*, *Amblos'mus*, *Ec'bole*, *Embryoto'cia*, *Diaph'thora*, *Ectro'sis*, *Examblo'ma*, *Examblo'sis*, *Ectro'e'mus*, *Apopalle'sis*, *Apopal'sis*, *Apoph'thora*, *Phtho'ra*, *Convulsio uteri*, *Deperdi'tio*, (F.) *Avortement*, *Blessure*, *Miscarriage*; (from *ab* and *oriri*, 'to rise,' applied to that which has arisen out of season.) The expulsion of the fœtus before the seventh month of utero-gestation, or before it is *viable*. The causes of this accident are referable either to the mother, and particularly to the uterus; or to the fœtus and its dependencies. The causes, in the mother, may be:—extreme nervous susceptibility, great debility, plethora, faulty conformation, &c.; and it is frequently induced immediately by intense mental emotion, violent exercise, &c. The causes seated in the fœtus are its death, rupture of the membranes, &c. It most frequently occurs between the 8th and 12th weeks of gestation. The symptoms of abortion are:—uterine hemorrhage with or without flakes of decidua, with intermitting pain. When abortion has once taken place, it is extremely apt to recur in subsequent pregnancies about the same period. Some writers have called abortion, when it occurs prior to three months, *Effluxion*. The treatment must vary according to the constitution of the patient and the causes giving rise to it. In all cases, the horizontal posture and perfect quietude are indispensable.

ABORTION is likewise applied to the product of an untimely birth,—*Abor'tus*, *Abor'vus*, *Apoble'ma*, *Apob'ole*, *Ecblo'ma*, *Amblothr'id'ion*, *Ectro'ma*, *Fructus immaturus*, *Abort'ment*, (F.) *Avorton*, *Avortin*.

TO ABORT, *Abori'ri*. To miscarry. (F.) *Avorter*.

ABORTIVE, *Aborti'vus*, *Ecbol'ius*, *Amblo'ticus*, *Amblothr'id'ium*, *Ambol'icus*, *Phthor'ius*, *Apophthor'ius*, *Ectrot'icus*, *Abortifaciens*, *Acyte'rius*, *Expel'lens*, *Phthirot'onus*, *Phthorot'onus*, *Ecbol'icus*, *Contractor uteri*, *Accelera'tor Partûs*, *Parturiens*, *Parturificient*, *Ecbol'ic*. (F.) *Abortif*. A medicine to which is attributed the property of causing abortion. There is probably no direct agent of the kind. See *Ectrotic*.

ABORTMENT, Abortion.

ABORTUS, gen. *Abortûs*, Abortion.

ABOUCHEMENT, (*bouche*, 'mouth,') *Anastomosis*.

ABOULAZA, a tree of Madagascar, used, according to Flacourt, in the practice of the country, in diseases of the heart.

ABOUTISSEMENT, (*aboutir*, 'to come to an end,' from *bout* 'end,') *Suppuration*.

ABOYEUSES, (*aboyer*, 'to bark,') See *Convulsionnaire*.

ABRABAX, *Abracax*, *Abrasax*, *Abraxas*. A mystic term, expressing the number 365, to which the Cabalists attributed miraculous properties. A form of amulet.

ABRACADA'BRA, *Abrasada'bra*, the name of a Syrian idol, according to Selden. This word, when pronounced and repeated in a certain form and a certain number of times, was supposed to have the power of curing fevers and preventing many diseases. It was figured on amulets and worn suspended around the neck.

A B R A C A D A B R A
B R A C A D A B R A
R A C A D A B R A
A C A D A B R A
C A D A B R A
A D A B R A
D A B R A
A B R A
B R A
R A
A

ABRACALAN, A cabalistic term to which the Jews attributed the same virtue as to the word ABRACADABRA.

ABRACAX, Abrabax.

A'BRAHAM MAN. A name given, three centuries ago, to one who, for purposes of imposition, personated a 'Tom of Bedlam' or vagrant lunatic.

To *sham Abraham* or *Abram* is now used for one who feigns sickness or infirmity of any kind.

ABRASADABRA, *Abracadabra*.

ABRASAX, Abrabax.

ABRASIO, gen. *Abrasio'nis*, Abrasion—a. Dentium, *Odontotripsis*.

ABRA'SION, *Abra'sio*, *Aposyr'ma*, *Aposys'mus*, (Prov.) *Fleck*, (from *abradere*, (*ab* and *radere*,) 'to rasp,') A superficial excoriation, with loss of substance, under the form of small *shreds*, in the mucous membranes of the intestines,—(F.) *Racures des Boyaux*. Also, an ulceration of the skin, possessing similar characters. According to Vicq d'Azyr, the word has been used for the absorption of the molecules composing the various organs.

ABRATHAN, *Artemisia abrotanum*.

ABRAXAS, Abrabax.

ABRE, *Abrus precatorius*.

ABRÉVIATION, Abbreviation.

ABRICOT, (from *præcox*, 'precocious,') see *Prunus Armeniaca*—a. *Sauvage*, *Mammea Americana*.

ABRICOTIER, *Prunus Armeniaca*.

ABROSIA, (a, and βρωσις, 'food,') Abstinence.

ABROTANUM, (αβροτανον), *Artemisia abrotanum*—a. *Cathsum*, *Artemisia abrotanum*—a. *Mas*, *Artemisia abrotanum*.

ABROTONUM, *Artemisia abrotanum*.

ABROTONITES, (αβροτονιται), (*uinos*, 'wine,' understood.) Wine impregnated with *Artemisia abrotanum* or Southernwood.

ABROTONUM, (αβροτανον), *Artemisia Abrotanum*.

ABRUPTIO, (*ab* and *rumpere*, *ruptum*, 'to break,') see *Abduction*.

A'BRUS PRECATORIUS, (from *αβρος*, 'elegant,') *Liquorice Bush*, *Red Bean*, *Love pea*, (F.) *Abre*, *Liane à réglisse*. A small ornamental shrub, found from Florida to Brazil, as well as in Egypt and the West Indies; *Nat. Ord.* Leguminosæ. *Sec. Syst.* *Monadelphia Enneandria*; having beautiful scarlet seeds with a black spot. The roots and leaves are sweet mucilaginous demulcents. The seeds of the American kind are considered to

be purgative and poisonous. They are employed to form rosaries, and hence called, in mockery, *Jumble beads*.

ABS, as a prefix, see A.

AB'SCESS, (from *abscedo*, (*abs*, and *cedere*.) 'I depart.' or 'separate from.') *Abscessus*, *Abscessio*, *Aphiste'sis*, *Apostema*, dim. *Apostematium*, *Epye'ma*, *Epye'sis*, *Reces'sus*, *Abscessio*, *Apostemum*, *Impos'thume*, *Impos'tem*, *Gath'ering*, *Ri'sing*; (Old Eng.) *Apostemucion*, *Apost'hume*. (Sc.) *Hat-trel*, (Prov.) *Coul*, *Numpost*, *Postume*, *Addle*. (F.) *Abrès*, *Depôt*. A collection of pus in a cavity, the result of a morbid process. See *Pyogenia*, and *Suppuration*.

The French have various distinctive terms for Abscesses. *Abrès chaud*, *aigu*, *soudain*, is one which follows violent inflammation. *Abrès froid*, *chronique*, *serofuleux*, *Cold*, *chronic*, or *serofulous abscess*, one which is the result of chronic or serofulous inflammation. *Abrès par Congestion*, *A. diathésique*, a symptomatic abscess; one which occurs in a part at a distance from the inflammation by which it is occasioned: e.g. a *lumbar abscess*; in which the inflammation may be in the lumbar vertebrae, whilst the pus exhibits itself at the groin.

ABSCISS, ALVEOLAR, *Parulis*—a. Cold, see Abscess.

ABSCISS, METASTATIC, *Abscessus metastaticus*, (F.) *Abrès métastatique*, *A. consécutif*, an abscess which forms suddenly, and sometimes without any precursory signs of inflammation, in a part of the body remote from one in a state of suppuration, and without presenting a sufficient reason for its development in the place which it occupies. It is a consequence of phlebitis.

ABSCISS, PERFORATING, OF THE LUNG, see Lung, perforating abscess of the —a. Perilaryngeal, see Perilaryngitis—a. *Psoas*, Lumbar abscess—a. Retropharyngeal, see Retropharyngeal—a. *Shirtstud*, *Abrès en bouton de chemise*.

ABSCISSIO, Abscess.

ABSCISSUS, Abscess—a. *Capitis Sanguineus Neonatorum*, *Cephalematoma*—a. *Cerebri*, *Encephalopoyosis*—a. *Gangraenescens*, *Anthrax*—a. *Gangraenosus*, *Anthrax*—a. *Lacteus*, *Mastodynia apostematosa*—a. *Lumborum*, *Lumbar abscess*—a. *Mammae*, *Mastodynia apostematosa*—a. *Metastaticus*, Abscess, metastatic—a. *Nucleatus*, *Furunculus*—a. *Oculi*, *Hypopyon*—a. *Pectoris*, *Empyema*—a. *Pulmonum*, *Pneumapostema*—a. *Renalis*, *Nephropostasis*—a. *Spirituosus*, *Aneurism*—a. *Thoracis*, *Empyema*—a. *Urinosis*, *Urapostema*.

ABSCISSIO, Abscession.

ABSCISSIO, Abscession—a. *Præputii*, *Circumcision*.

ABSCIS'SION, *Abscis'io*, *Abscis'sio*, (from *abscedere* or *abscindere*, 'to cut off,') *Apoc'opē*, *Apothraus'sis*, *Diac'opē*. Excision or extirpation of a part, especially of a soft part. Fracture or injury of soft parts, with loss of substance. Diminution, or loss of voice. Sudden and premature termination of a disease.

ABSCONISIO, (*abs*, and *condere*, 'to hide,') *Sinus*.

ABSENCE DU BRUIT RESPIRATOIRE, see Murrain, respiratory.

ABSENTHE, see *Artemisia absinthium*.

ABSENTHITES, (*αψινθίτης*), *Apsinthi'tes*, Wine impregnated with *Absinthium* or *Wormwood*.

ABSENTHIUM. (Ph. U. S.), (*αψινθιον*), *Artemisia absinthium*—a. *Marinum*, *Artemisia maritima*—a. *Maritimum*, *Artemisia maritima*—a. *Ponticum*, *Artemisia pontica*—a. *Romanum*, *Artemisia pontica*—a. *Santonium*, *Artemisia santonica*—a. *Vulgare*, *Artemisia absinthium*.

ABSORBANT, Absorbent.

ABSORBENT, *Absorbens*, (from *absorbere* (*ab*

and *sorbere*), 'to drink, to suck up,') (F.) *Ab-sorbant*. That which absorbs.

A medicine used for absorbing acidity in the stomach and bowels, as *magnesia*, *chalk*, &c. *Inver'tens*, *Resorbens*, *Saturans*, *Træcipitans*.

Also, any substance, such as *cobweb*, *sponge*, &c., which, when applied to a bleeding surface, retains the blood, and forms with it a solid and adhesive compound, which arrests the hemorrhage.

ABSORBENT SYSTEM is the collection of vessels, *Vasa absorbentia* seu *resorbentia*, and glands, which concur in the exercise of absorption.

ABSORPTIO, Absorption—a. *Sanguinis*, *Hæmorrhophesis*.

ABSORPTION, (same etymon as Absorbent), *Absorptio*, *Resorp'tio*, *Inhalat'io*, *Imbibit'io*, *Anar'rhop'hē*, *Anarrop'hē'sis*, *Catapino'sis*, *Rhabdē'sis*, *Catar'rhop'hē'sis*, *Catar'rhop'hē*. The function of absorbent vessels, by virtue of which they take up substances from without or within the body. Two great divisions have been made of this function.

1. *External absorption*, or the *absorption of composition*, which obtains, from without the organs, the materials intended for their composition; and, 2. *Internal absorption*, or the *absorption of decomposition*, which takes up from the organs the materials that have to be replaced.

By *external absorption* is meant not only that which takes place at the external surface of the body, but also that of the mucous membranes of the digestive and respiratory passages. Hence, again, the division of external absorption into *cutaneous*—*resorptio cutanea* seu *cutis*, *inhalat'io cutis*,—*intestinal* or *digestive*, and *pulmonary* or *respiratory*.

Internal absorption is also subdivided into, 1. *Molecular* or *interstit'ial*, *nutritive*, *organ'ic*, or *decompo'sing*, which takes up from each organ the materials that constitute it, so that the decomposition is always in equilibrio with the deposition. 2. The *absorption of recementit'ial secreted fluids*, such as the fluid of serous membranes, synovia, &c. As these are constantly exhaled on surfaces which have no external outlet, they would augment indefinitely if absorption did not remove them in the same proportion as that in which they are deposited. 3. The *absorption of a part of the recementit'ial secreted fluids*, as they pass over the excretory passages.

The great agents of external absorption are the veins and chyloferous vessels; of internal absorption, probably the lymphatics. In the chyloferous vessels and lymphatics the fluid is always found to possess the same general properties. In them, therefore, an action of elaboration or selection must have taken place. The veins, on the other hand, seem to exert no selection. Any fluid, possessing the necessary tenuity, passes through the coats of the vessel readily by imbibition, and proceeds along with the torrent of the circulation. Watery fluids in this manner enter the blood when they are taken into the stomach. Substances that require digestion, on the other hand, may pass through the chyloferous vessels and thoracic duct.

ABSORPTION OF COMPOSITION, see Absorption—a. *Cutaneous*, see Absorption—a. of *Decomposition*, see Absorption—a. *Digestive*, see Absorption—a. of *Excrementitial Secreted Fluids*, see Absorption—a. *External*, see Absorption—a. *Internal*, see Absorption—a. *Interstitial*, see Absorption—a. *Intestinal*, see Absorption—a. *Molecular*, see Absorption—a. *Nutritive*, see Absorption—a. *Organic*, see Absorption—a. *Pulmonary*, see Absorption—a. of *Recementitial Secreted Fluids*, see Absorption—a. *Respiratory*, see Absorption.

ABSTÈME, Abstemious.

ABSTEMIOUS, *Abstem'ius*, *Aoi'nos*, (from *abs* and *temetum*, 'wine,') (F.) *Abstème*. Used by the ancient writers, as well as by the French, in the

sense only of its roots; one who abstains from wine or fermented liquors in general.

ABSTERGENTIA, (*abs* and *tergere*, *tersum*, 'to clean,') Detergents.

ABSTERSIVA, Detergents.

ABSTERSORIA, Detergents.

ABSTINENCE, *Abstinentia*, (from *abs* and *tenerē*, 'to hold,') *Abrosia*, *Asitia*, *Liman'chia*, Fasting. Privation, usually voluntary, as when we speak of *abstinence from pleasure*, *abstinence from drink*, &c. It is more particularly used to signify voluntary privation of certain articles of food. Fasting is a useful remedial agent in certain diseases, particularly in those of an inflammatory character.

AB'SUS, a kind of cassia, *C. Absus*, which grows in Egypt and in India, the seeds of which, pulverized and mixed with powdered sugar, have been employed in form of a dry collyrium, in the endemic ophthalmia of Egypt.

ABU'LIA, (*αβουλια*, from *α*, and *βουλη*, 'will,') Loss of the will, or of volition.

ABU'LICUS. One who has lost the power of will or of volition.

ABUS DE SOI-MÊME, ('abuse of one's self,') Masturbation.

ABUTA (a Guiana name), *Paraiba brava*.

ABUTILON **AVICEN'NÆ**, *A. Cordatum*, *Si'di Abutilon*, Indian or Yellow Mal'low, *Vell'ret leaf*; *Order*, Malvaceæ, (*α*, 'privative,' *δους*, 'an ox,' and *τιλος*, 'diarrhœa,' that is 'a remedy for the diarrhœa of cattle;') is naturalized, and common in most parts of the United States. It resembles common mallow in its properties, being mucilaginous and demulcent.

ABVACUA'TIO, (from *ab*, and *vacuare*, *vacuatum*, 'to empty,') An excessive or colliquative evacuation of any kind.

AC, as a prefix, same as **Ad**.

ACACIA (Ph. U. S.), *Acaciæ gummi*—a. Catechu, Catechu—a. False, *Robinia pseudo-acacia*—a. Germanica, see *Prunus spinosa*—a. Giraffæ, see *Acaciæ gummi*—a. Horrida, see *Acaciæ gummi*—a. Indica, *Tamarindus*—a. Nilotica, see *Acaciæ gummi*—a. Nostræ, see *Prunus spinosa*—a. Senegal, see *Senegal, gum*—a. Vera, see *Acaciæ gummi*—a. Zeylonica, *Hæmatoxylon Campechianum*.

ACA'CLE GUM'MI, *Aca'cia*, *G. Aca'cia Arab'ice*, *G. Mimo'sæ*, *G. Arab'icum*, *G. Acanth'inum*, *G. Leu'cum*, *G. Theba'icum*, *G. Serapio'nis*, *G. Lamæ*, *G. Sen'ega* or *Sen'eca*, (see *Senegal, gum*), *Gum Arabic*. (F.) *Gomme Arabique*. The gum of the tree *Aca'cia* seu *Mimo'sa Nilot'ica*, *Aca'cia re'na*, *Sp'ina Egypti'aca*, of Upper Egypt, *Nat. Ord.* Mimoseæ. *Sex. Syst.* Polygamia Monœcia, (from *ακν*, 'a point,' so called in consequence of the spines.) It is in irregular pieces, colourless, or of a pale yellow colour, hard, brittle, of a shining fracture, transparent, soluble in water, and insoluble in alcohol, s. g. 1.4317.

It is mucilaginous; but is rarely used, except in pharmacy. Sometimes it is administered alone as a demulcent.

Acaciæ Hor'rida and *A. Giro'f'fæ*, of South Africa, yield a good gum.

ACAJOU, (of Indian origin) *Anacardium occidentale*.

ACAJUBA OFFICINALIS, *Anacardium occidentale*.

ACALYPHA BETULI'NA, (*α*, 'privative,' *αλος*, 'beautiful,' and *ἀσπν*, 'touch,' 'disagreeable to the touch,') *Cau'da fel'is agre's'tia*, is employed by the natives of India as a stomachic and in cholera. It is given in the form of infusion of the leaves.

ACALYPHA HISPIDA, *Caturus spiciferus*.

ACALYPHA IN'DICA, *Cupameti*, *Order* Euphorbiacæ. A common annual in the gardens of India. An infusion of the root, and a decoction of the leaves are cathartic.

ACALYPHA VIRGIN'ICA. *Three seeded mer'cury*,

indigenous, flowering in August, is said to have expectorant and diuretic properties.

ACAM'ATUS, (*α*, and *καμνω*, 'I labour,') This word has been sometimes used for a good constitution of the body. According to Galen, it means that position in which a limb is intermediate between flexion and extension; a position which may be long maintained without fatigue.

ACAMPSIA, (*ακαμψια*, from *α*, and *καμψις*, 'flexion,') Contractura.

ACANOS, (*ακανος*, 'a prickle,') *Onopordium acanthium*—a. Spina, *Onopordium acanthium*.

ACANTHA, (*ακανθα*, 'a spine or thorn,') Vertebral column. Also, Spinous process of a vertebra.

ACANTHAB'OLUS, *Acan'thulus*, *Acanthav'olu*, *Valsell'la*, (*ακανθα*, 'a spine,' and *βαλλω*, 'I cast out,') A kind of forcep. for removing extraneous substances from wounds.

ACANTHALZUCA, *Echinops*.

ACANTHAVOLA, *Acanthabolus*.

ACANTHE FAUSSE, *Heracleum spondylium*.

ACANTHIA LECTULARIA, *Cimex lectularius*.

ACANTHIUM, *Onopordium acanthium*.

ACANTHULUS, *Acanthabolus*.

ACANTHUS MOL'LIS, (*ακανθος*,) (from *ακανθα*, 'a spine or thorn,') *Melampyrl'ium*, *Bran'ca ursi'na* seu *ve'ra*, *Brankur'sine*, *Bear's Breech*. (F.) *Pied d'ours*. This plant is mucilaginous like *Althæa*, and is used as a demulcent.

ACAPATLI, *Piper longum*.

ACAR'DIA, (*α*, and *καρδια*, 'the heart,') The state of a fœtus without a heart.

ACARDIOHÆMIA. (F.) *Acardiôhémie*, (*α*, *καρδια*, 'heart,' and *αίμα* 'blood,') Want of blood in the heart.

ACARDIONER'VIA, (F.) *Acardionervie*: (*α*, *καρδια*, 'heart,' and *νευρον*, 'nerve,') Want of nervous action in the heart as indicated by the sounds ceasing to be audible.

ACARDIOTROPHIA, (*τροφη*, 'nourishment,') Heart, atrophy of the.

ACARE, *Acarus*.

AC'ARICIDE, (*acarus*, and *cædere*, 'to kill,') A destroyer of acari.—as of the *acarus scabiei*.

ACARICO'BA. The Brazilian name for *Hydrocotyl'le umbellatum*, used by the Indians as an aromatic, alexipharmic, and emetic.

ACARON. (*ακαρνος*, 'small, short,') *Myrica gale*.

ACARP'Æ, (*α*, and *καρπος*, 'fruit,') A division of the family of cutaneous diseases by Fuchs, in which there is no "fruit," (Germ. *Frucht*,) or production from the cutaneous surface—tubercles, vesicles or pustules. Lentigo, Chloasma, Argiria, and Pityriasis belong to it.

AC'ARUS, (*α*, and *καρνος*, 'divisible,') (F.) *Acare*. A minute insect, one species of which has been noticed, by several observers, in the itch. The *Acarus Scabiei*, see *Psora*.

ACARUS CIRO, see *Psora*—a. *Comedonum*, *Acarus Folliculorum*.

AC'ARTS CROS'SEI, *Crosse mite*. An insect supposed by Mr. Crosse, of England, to have been developed in a solution of silicate of potassa when submitted to slow galvanic action, for the purpose of obtaining crystals of silex. It did not, however, prove to be a new formation.

ACARUS FOLLICULO'RU, *A. Comedo'num*, *Entozo'on* seu *De'moder* seu *Simo'nea* seu *Steatczo'on folliculo'rum*, *Macroga'ster platypus*. An articulated animalcule, discovered in the sebaceous substance of the cutaneous follicles. According to Professor Owen, it belongs to the Arachnida.

ACARUS SCABIEI, *Acarus*, see *Psora*.

ACATALEP'SIA, (*ακαταληψια*, from *α*, *καταλαμβανω*, 'I comprehend,') Uncertainty in diagnosis. Its opposite is *Catalepsia*.

ACATALIS, see *Juniperus communis*.

ACATAP'OSIS, (*a*, and *καταποσις*, 'deglutition.') Incapacity of swallowing. Vogel has given this name to difficulty of deglutition.

ACATASTATIC, *Acatastat'icus*, (*a*, and *καταστατικός*, 'to determine.') An epithet given to fevers, &c., when irregular in their periods or symptoms.

ACATERA, see *Juniperus communis*.

ACATHAR'SIA, (*a*, and *καθαρσις*, 'I purge;') *Sordes*, Impurities. Omission of a purgative.

ACATSJAVALLI, a Malabar plant, which is astringent and aromatic. A bath of it is used in that country in cases of hemierania. It is supposed to be the *Cassytha filiformis* of Linnaeus.

ACAWERIA, *Ophioxylum serpentinum*.

ACCABLEMENT, Torpor.

ACCÉLÉRATEUR DE L'URINE, (*ac*, and *celer*, 'quick;') Accelerator urine.

ACCELERATOR PARTUS, Abortive.

ACCELERATOR URINÆ, *Bulbocavernosus*, *Bulbo-urétral*—(*Ch*.), *Ejaculator Seminis*, *Bulbosyndesmo-cavernosus*. (*F*.) *Accélérateur de l'urine* *Bulbo-cavernosus*, *Ano-cavernosus*. A muscle of the penis, which arises, fleshy, from the sphincter ani and membranous part of the urethra, and tendinous, from the crus and beginning of the corpus cavernosum penis. In its course it forms a thin, fleshy layer, the inferior fibres of which run more transversely than the superior, which descend in an oblique direction; the muscles of both sides completely enclosing the bulb of the urethra. It is inserted into its fellow by a tendinous line running longitudinally on the middle of the bulb. Its use is to propel the urine or semen forwards.

ACCENT, *So'nus vo'cis*, (*ad* and *canere*, *cantum*, 'to sing;') Inflection or modification of the voice, which consists in raising or dropping it on certain syllables.

The accent exhibits various alterations in disease.

ACCÈS, Accession, Attack, Paroxysm.

ACCESS, Accession, Attack, Paroxysm.

ACCES'SION, *Acces'sio*, (*from accedo*, *accessum*, (*ac*, and *cedere*,) 'to approach;') The invasion, approach, or commencement of a disease.

ACCESSOIRE, Accessory—*a. du long Fléchisseur commun des orteils*: see *Flexor longus digitorum pedis profundus perforans* (*accessorius*)—*a. de l'Obturator interne*, *Ischiochroanterianus*—*a. du Pied d'Hippocampe*: see *Cornu ammonis*—*a. du Sacro-lombaire*: see *Sacrolumbalis*.

ACCESSORIUS, Accessory—*a. Flexor Longus Digitorum Pedis*: see *Flexor longus digitorum pedis profundus perforans* (*accessorius*)—*a. Pedis hippocampi*:—see *Cornu ammonis*.

ACCESSORY, *Accesso'rius*, (*F*.) *Accessoire*, *Annexe*. A consequence or dependence on any thing; as *accessory ligament*, *muscle*, *nerve*, &c.

ACCESSORY OF THE PAROTID is a name given by Haller to a small gland, which accompanies the parotid duct, and is commonly a mere prolongation of the parotid itself. See *Parotid*.

ACCESSORY SCIENCES TO MEDICINE are those which do not relate directly to the science of man in a state of health or disease; as *physics*, *chemistry*, &c.

ACCESSORY OF THE PAR VAGUM, Spinal nerve. The term *accessory* is also given to several muscles.

ACCESSUS, Coition.

ACCIDENS, (*from accidere*, (*ac*, and *cadere*,) 'to happen;') Symptom—*a. Consecutifs*, Consecutive phenomena.

ACCIDENT, *Ac'cidents*. A casualty; an unforeseen event. The French—like older English writers—use the term in nearly the same sense as *symptom*. It means also an unexpected symptom.

ACCIDEN'TAL, *Adventit'ious*. That which happens unexpectedly.

The French give the name *Tissus accidentels* to those adventitious textures, that are the result of a morbid process.

ACCIP'ITER, *Hi'erax*, 'the hawk, (*from accipere*,—*ac*, and *cipio*, 'to take;') *Menec'ratis* *Accip'iter*, (*F*.) *Épervier*. A bandage applied over the nose, so called from its likeness to the claw of a hawk.

ACCLIMATED, *Accli'matised*, *Clima'ti assue'tus*, (*ac*, and *clima*, 'climate;') A word of modern introduction from the French, which means 'accustomed to a climate.'

ACCLIMATATION, Acclimation.

ACCLIMATEMENT, Acclimation.

ACCLIMA'TION, *Acclima'tio*, *Clima'tion*, *Seas'-oning*. (*F*.) *Acclimatement*, *Acclimation*, *Acclimatisation*. The act of becoming acclimated or accustomed to a climate.

The constitution of a person, who goes to live in another and a very different climate, usually experiences changes, which are frequently of an unfavourable character, and the study of which is of considerable importance in medicine.

ACCLIMATISATION, Acclimation.

ACCLIMATISED, Acclimated.

ACCOMPAGNEMENT DE LA CATARACTE, Accompaniment of the cataract.

ACCOMPANIMENT, *Adjunc'tion*. (*F*.) *Accompagnement*, (*compagnon*, 'an associate'). That which is joined to any thing.

Accompagnement to the cataract, (*F*.) *Accompagnement de la cataracte*, is a whitish, viscid substance, which sometimes surrounds the opaque crystalline lens, and remains after the operation for cataract, causing a secondary cataract.

ACCOUCHÉE, (*ac*, and *couche*, 'a bed;') Puerpera.

ACCOUCHEMENT, Parturition—*a. Laborious*, *Dystocia*—*a. contre Nature*, see *Presentation*, *preternatural*—*a. Laborieux*, *Laborious labour*,—*a. Provoqué*, see *Parturition*.

ACCOUCHEUR, (*F*.) *Adj'u'tor Par'tûs*, *Obstet'ricans*, *Obstetric'i'us*, *Maien'ter*, *Maien'tes*. He who practices the art of midwifery. A physician *Accoucheur*, a Surgeon *Accoucheur*, a Midwife, &c.

ACCOUCHEUSE, Midwife.

ACCOUPLEMENT, (*ac*, and *coupler*, 'to couple;') Coition.

ACCOUSTOMANCE, (*ac*, and *coutume*, 'custom;') Habit.

ACCRETION, *Accre'tio*, (*ac*, and *crescere*, 'to increase;') Augmentation; (*F*.) *Accroissement*; also, increase by juxtaposition.

ACCROISSEMENT, (*ac*, and *croissance*, 'growth;') Accretion, Increase.

ACCU'BITUS, (*from accubo*, (*ac*, and *cubo*, 'I lie;') 'I lie by the side of;') Of old, the lying of a person with another, as where a young person was placed in bed with an old one, or a healthy person along with one that was sick.

ACCUSATIO, (*ac*, and *causa*, 'cause;') Indication.

ACE'DIA, (*ακηδεια*, and *ακηδία*,) (*a*, and *κηδος*, 'care;') *Incu'ria*. Want of care, neglect. Also, fatigue.

ACELLA, Axilla.

ACEOGNOSIA, (*ακκομαι*, 'I cure; and *γνωσις*, 'knowledge;') Pharmacognosia.

ACEOLOGIA, (*ακκομαι*, and *λογος*, 'a discourse;') Materia Medica.

ACÉPHALE, Acephalous.

ACEPHALHE'MIA, (*F*.) *Acephalhémie*, (*acephalo*, and *αίμα*, 'blood;') Want of blood in the encephalon.

ACEPHALIA, see Acephalous.

ACÉPHALIE, see Acephalous.

ACEPH'ALO, (from *a*, privative, and *κεφαλη*, 'head,') in composition, devoid of head.

ACEPHALOBRA'CHUS, (*acephalo*, and *βραχιον*, 'arm,') A fetus without head or arms.

ACEPHALOCAR'DIA, (*acephalo*, and *καρδια*, 'heart,') Monstrosity in which there is absence of head and heart.

ACEPHALOCHI'RUS, (*acephalo*, and *χειρ*, 'hand,') A fetus without head or hands.

ACEPH'ALOCYST, *Acephalocystis*, (F.) *Acéphalocyste*, (*acephalo*, and *κυστις*, 'bladder,') A hydatiform vesicle, without head or visible organs, ranked amongst the Entozoa, although possessed of few animated characteristics. In no organ of the body are acephalocysts so frequently found as in the liver. Generally it is the 'multiple acephalocyst,' *A. socialis* seu *prolifer*a, which is met with. At times, however, it is the 'solitary acephalocyst,' *A. eremita* seu *sterilis*.

The *acephalocystis endog'ena* has a firm coat, and is composed of different layers, which have numbers of smaller hydatids within them, and are thrown off from the interior of the parent cyst. This species has hence been termed *endogena*, to distinguish it from the *A. exog'ena* of ruminant animals, in which the young vesicles are developed from the exterior of the parent vesicles.—See Hydatid.

ACEPHALOGAS'TER, *Athoracoceph'alus*, (*acephalo*, and *γαστηρ*, 'the belly,') A name given to monsters devoid of head, chest, and abdomen; or to those which have an abdomen, but no chest or head.

ACEPHALOP'ODUS, (*acephalo*, and *πους*, gen. *ποδος*, 'foot,') A monster devoid of head and feet. The condition is called *Acephalopod'ia*.

ACEPHALORA'CHIA, (*acephalo*, and *ραχις*, 'spine,') Monstrosity in which there is absence of head and spine.

ACEPHALOS'TOMA, (from *acephalo*, and *στομα*, 'mouth,') An acephalus fetus, at the upper part of which there is an opening resembling a mouth.

ACEPHALOTHORA'CIA, (*acephalo*, and *θωραξ*, gen. *θωρακος*, 'thorax,') Monstrosity in which there is absence of head and chest.

ACEPHALOTHOR'US, (*acephalo*, and *θωραξ*, 'chest,') *Apectoceph'alus*. A monster devoid of head or chest.

ACEPH'ALUS, (*ακεφαλος*, from *a*, and *κεφαλη*, 'head,') (F.) *Acéphale*. A monster born devoid of head. The condition is called *Acephalia*. (F.) *Acéphalie*.

ACER, gen. *A'ceris*, *Acrid*.

ACER PALMIFOLIUM, (*acer*, 'the maple,') *A. Saccharinum*.

ACER PENNSYLVANICUM, *Stri'ped Ma'ple*, *Stri'ped Dog'wood*. A decoction of the bark has been used internally and externally in cutaneous affections; and a decoction of the leaves and twigs is said to relieve nausea and vomiting.

ACER RU'BUM, *Red Ma'ple*; Indigenous. The inner bark is a mild astringent. The Indians use a decoction of it for sore eyes.

ACER SACHARI'NUM, *A. palmifol'ium*, *Ma'ple*, *Sugar Ma'ple*. (F.) *Érable*; *Ord.*, *Acercææ*. This tree contains a large amount of sweet sap, whence a considerable quantity of sugar may be extracted. When purified, this sugar can scarcely be distinguished from that obtained from the cane.—See *Saccharum*.

ACERATES LONGIFO'LIA, *Long-leaved green Milk'weed*; *Order*, *Asclepiadacææ*; indigenous, flowering in June and July; has the properties of the order. See *Asclepias*.

ACERATO'SIS, (*a*, privative, and *κερας*, gen. *κερατος*, 'horn,') Defective development of the corneous tissue.

ACERB', *Acerbus*, *Stryph'nos*, (from *acer*, 'sharp,') A savour, or taste, compounded of the

acid, bitter, and astringent; such as is met with in unripe fruits, &c.

ACER'CUS, (*a*, privative, and *κερκος*, 'a tail,') A monster devoid of tail.

ACERIDES, *Acerodes*, (*a*, priv., and *κερος*, 'wax,') Plasters devoid of wax.

ACERODES, *Acerides*.

ACERO'SUS, *Achyrodes*, (*αχυρον*, 'chaff,')

Pityri'nus, *Furfura'ceus*. An epithet used by Hippocrates for the coarsest bread, made of flour not separated from the chaff.

ACERVULUS CEREBRI, (dim. of *acervus*, 'a heap,') See *Pineal Gland*—*a. Glandula Pinealis*, see *Pineal Gland*.

ACES'CENCY, *Acescen'tia*, [*acescere*, 'to grow sour,'] (*akis*, 'a point, *acer*, 'sharp,'] A disposition to acidity. The humourists believed that the animal humours are susceptible of this change.

ACESIA (*ακεις*), *Cure*.

ACESINOSUS (*ακισινοςος*), *Curative*.

ACESIS (*ακεις*), *Curation*, *Cure*, *Medicament*.

ACESMA gen. *Acesma'tis*, (*ακεσμα*), *Medicament*.

ACESMIUS (*ακεσμιος*), *Curable*.

ACESMUS (*ακεσμος*), *Cure*.

ACESODYNES, (*ακεσις*, 'cure,' and *οδυνη*, 'pain,') *Anodyne*.

ACESOPHORUS, (*ακεσις*, 'cure,' and *φερω*, 'I carry,') *Curative*.

ACESTER (*ακεστηρ*), *Physician*.

ACESTIS, *Medicament*.

ACESTOR, *Physician*.

ACESTORIA (*ακεστορια*), *Medicine*

ACESTORIS (*ακεστορις*), *Midwife*.

ACESTOS (*ακεστος*), *Curable*.

ACESTRA (*ακεστρα*), *Needle*.

ACESTRIA (*ακεστρια*), *Midwife*.

ACESTRIS (*ακεστρις*), *Midwife*.

ACESTRUM (*ακεστρον*), *Medicament*.

ACETA MEDICATA, *Acetica*.

ACETABULA UTERINA, (pl. of *acetabulum*,) *Cotyledons*.

ACETABULUM, (F.) *Acétabule* (from *acetum*, because it resembles the old vinegar vessel *oxybaph'ion*.) A measure capable of containing the eighth part of a modern pint.—*Athenæus*. *Galen*. See *Cotylod*. According to *Castelli*, the lobes or cotyledons of the placenta of ruminating animals have been so called *acetabula*.

ACETABULUM, *Cotyle*, see *Cotylod*—*a. Humeri*, see *Glenoid*—*a. Marinum*, *Umbilicus marinus*.

ACETARIA, (from *acetum*.) A salad or pickle.

ACETAS, *Acetate*.

AC'ETATE, *Ac'etas*. (F.) *Acétate*. A salt formed by the union of the acetic acid with an alkaline, earthy, or metallic base. The acetates chiefly used in medicine are the acetates of ammonia, lead, potash, and zinc.

ACE'TICA, *Ac'eta Medicat'a*. (F.) *Vinaigres Médicinaux*. Pharmaceutical preparations of vinegar.

ACETICUM AC'IDUM, *Acidum Aceticum fortius* seu *fort'è* seu *pu'r'um* seu *glacia'tè*, *Ac'etum radica'tè*, *Ox'us*, *Ac'etic Acid*, *Strong Ac'etous Acid*, *Acidum Aceto'sum fort'è*, *Rad'ical Vin'egar*, *Spir'itus Ven'etis*, and *Spirit of Verdigris*, when made from *verdigris*. Concentrated acetic acid, prepared by decomposing an acetate and receiving the acetic acid by distillation, has a very pungent and grateful odour, and an acid and acrid taste. Its s. g. is about 1.046, and it is very volatile.

It is stimulant, rubefacient, and escharotic, and is applied to the nostrils in syncope, asphyxia, headache, &c. It destroys warts.

An *Aromatic Spir'it of Vin'egar*, *Ac'idum Aceticum Camphora'tum*, *A. aceto'sum camphora't*

tum, is formed of this strong acid, f3x; Camphor, 3j; Alcohol, f3i.

A strong Acetic Acid was ordered by the London pharmacopœia, prepared from wood. It was called *Vinegar of wood*, Improved distilled *Vinegar*, *Pyropligneous Acid*, *Acetum Ligno'rum*, and its strength was such, that 87 gr. of crystallized subcarbonate of soda should saturate 100 grains of the acid.

Ac'idum Aceticum Dilu'tum, A. A. ten'ne seu debil'ins, *Acetum destilla'tum*, *Acidum aceticum*, *Acidum Aceto'sum destilla'tum*, *Distill'd vin'egar*, (F.) *Acide Acétique faible*, *Vinaigre distillé*, is prepared by distilling vinegar, until seven-eighths have passed over. An *Acidum aceticum dilutum*, *Diluted acetic acid*, is made by mixing a pint of the strong acetic acid with seven pints of distilled water.—Ph. U. S. Its properties are like those of vinegar.

ACETICUM MARTIALE, Ferri Acetas.

ACÉTOLAT, (F.) (from *acetum*.) A liquid medicine resulting from the distillation of vinegar on one or more aromatic vegetable substances, and which are formed of vinegar and essential oils, or other volatile principles.

ACÉTOLATURE. A tincture formed by digesting a vegetable substance or substances in vinegar.

ACÉTOLÉ, Océolé. A medicine, formed of distilled vinegar and medicinal principles dissolved in it.

ACÉTOMEL, Oxymel.

AC'ETONE, Ac'eton, (from *acetate*.) *Spir'itus pyroaceticus lignosus*, *Pyroacetic spirit*, *Pyroacetic Ether*, *Mesit'ic Al'cohol*, *Bihy'drate of Mesit'ylen*, (F.) *Acétone*; erroneously called *Naph'tha* and *Wood Naphtha*. By others, however, the terms are applied to *Pyroxyl'ic Spirit*. A limpid, colourless liquid, having a peculiarly penetrating and slightly empyreumatic odour. Its density in the liquid state, is almost the same as that of alcohol, 0.7921. Its taste is disagreeable, and analogous to that of peppermint. It is miscible in all proportions with water, alcohol, and ether. It may be prepared by distilling a mixture of two parts of crystallized acetate of lead and one part of quicklime in a salt-glaze jar (gray-beard,) the lower part of the jar being coated with fire-clay; and a bent glass tube, half an inch in diameter, adapted to the mouth by a cork, so as to form a distillatory apparatus. The jar is supported on the mouth of a small furnace, by which the lower part only is heated to redness, and the vapours are conducted into a Liebig's condenser. The product is repeatedly redistilled from quicklime, until its boiling point is constant at 132°.

It was brought forward as a remedy in phthisis pulmonalis; but evidently with unfounded pretensions. It is an excitant, and may be serviceable in chronic bronchitis. The dose is ten to forty drops three times a day, diluted with water.

ACETOSA ALPINA, *Rumex alpinus*—a. *Nosttras*, *Rumex acetosa*—a. *Pratensis*, *Rumex acetosa*—a. *Romana*, *Rumex scutatus*—a. *Rotundifolia*, *Rumex scutatus*—a. *Scutata*, *Rumex scutatus*—a. *Vulgaris*, *Rumex acetosa*.

ACETOSELLA, *Oxalis acetosella*.

ACE'TUM, *Oxus*, *Acetum Vi'ni* seu *Britan'nicum* seu *Gal'licum*, seu *Cru'dum* seu *Communi's*, *Acidum aceto'sum*, *A'legar*, *Acetum Cerevis'ia*, *Com'mon Vin'egar*, *Eisel*, (F.) *Vinaigre*; (from *akis*, 'a point,' *acer*, 'sharp.') A liquor obtained by the acetous fermentation. Vinegar has a pungent odour, and a pleasant acid taste. One fluid ounce of the *Acetum* of the United States Pharmacopœia is saturated by about 35 grains of crystallized bicarbonate of soda. It is refrigerant in

fevers; antiseptic, and antinarcotic; and externally is stimulant and discutient.

Vinegar Whey is made by stirring a small wine-glassful of vinegar, sweetened with a dessert-spoonful of sugar, in a pint of milk; boiling for fifteen minutes, and straining. Like tamarind whey it is an agreeable drink in febrile affections.

ACETUM AROMAT'ICUM, A. *Theriaca'l'e* seu *quat'uor fu'rurum*, *Ac'idum Aceticum Aromaticum*, *Thièces' Vin'egar*, *Vinegar of the four Thièces*, *Marseilles Vinegar*, (F.) *Vinaigre Aromatique*, *V. des quatre voleurs*, (Rorismarin. *cacum*, *sicc.*, *Fol. Salvia* sing. 3j. *Lavand. flor. sicc.* 3iv. *Caryoph. cont.* 3ss. *Acid. Acet. Oij.* Macerate 7 days, and filter.—Ph. E.) Odour, pungent and aromatic. Used as a perfume.

ACETUM BRITANNICUM, *Acetum*.

ACETUM CANTHARIDIS, *Vin'egar of Cantharidis*, *Cantharid.* in pulv. 3ij. *Acid. acet. f3v.*, *Acid. pyrolign. f3xv.* *Euphorb.* in pulv. crass. 3ss. Mix the acids; add the powders; macerate for seven days; strain; express strongly, and filter the liquor.—Ph. E. The London College macerates *cantharid.* 3ij in *acid. acet. Oj.* (Imp. meas.) for eight days; expresses and strains. It is used as a prompt vesicant.

ACETUM COL'CHICI, *Vin'egar of meadow saffron*, *Colchic. rad.* pulv. 3ij. *Acid. acetic. dilut. vel Acet. destillat.* Oij; Ph. U. S. It may also be made by displacement.) It is used as a diuretic, and also in gout. Dose f3ss. to f3iss.

ACETUM COMMUNE, *Acetum*—a. *Crudum*, *Acetum*—a. *Destillatum*; see *Aceticum acidum*—a. *Gallicum*, *Acetum*—a. *Lignorum*: see *Aceticum acidum*.

ACETUM LOBE'LLÆ, *Vin'egar of Lobelia*. (*Lobel.* in moderately coarse powder, 3iv. *Acid. acetic. dilut.* Oij. Made by percolation or maceration.—Ph. U. S., 1863.) Dose 30 to 60 drops and more.

ACETUM MELSUM DULCE, *Oxyglycos*—a. *Opii*, *Gutta Nigra*—a. *Plumbicum* seu *Saturninum*, *Liquor Plumbi subacetatis*—a. *Quatuor furum*, *Acetum Aromaticum*—a. *Radiale*, *Aceticum Acidum*—a. *Rosatum*, *Oxyrrhodinum*.

ACETUM SANGUINARIÆ, *Vin'egar of Bloodroot*, (*Sanguinar.* in moderately coarse powder, 3iv. *Acid. acet. dil.* Oij. Made by percolation or maceration.—Ph. U. S., 1863.) Dose 30 to 60 drops and more.

ACETUM SCIL'LE, *Ac'idum Aceticum Scillit'icum*, *Vin'egar of Squills*, (F.) *Vinaigre Scillitique*, (*Scille* contus. 3iv; *Acet. destillat.* Oij; Ph. U. S. It may also be made by displacement.) Diuretic, expectorant, and emetic. Dose f3ss to f3ij as a diuretic and expectorant. It may also be made by the process of displacement.

ACETUM THERIACALE, *Acetum aromaticum*.

ACEYTE DE SAL. A remedy for bronchocele used in S. America. Roulin found it to contain a portion of iodine.

ACHACANA. A species of cactus, in the province of Potosi in Peru. Its root is thick and fleshy, and of a conical shape. It is a good edible, and is sold in the markets of the country.

ACHALYBHÉMIE, (*a*, *chalys*, 'steel,' and *ápa*, 'blood.') See *Chlorosis*.

ACHANACA. A plant of the kingdom of Mely in Africa. It is used by the natives as an antisyphilitic.

ACHAOVAN, a species of Egyptian chamomile.

—Prosper Alpinus.

ACHAOVAN-ABIAT. The Egyptian name of *Cinera'ria marit'ima*, used in female diseases.

ACHAR, *Atchar*.

ACHE, (*αχος*, Sax. *ace*.) Pain.

ACHEBONE, *Innommatum* os.

ACHE, *Apium graveolens*—a. *des Montagnes*, *Ligusticum levisticum*.

ACHEILIA, *Achi'lia*, (*a*, and *χελος*, 'lip,')

A malformation, consisting in a deficiency of a lip or lips.

ACHEILUS, *Achi'lus*: (αχιλος.) One who is without lips.

ACHEIR, *Achir*, *De'manus*, (αχειρ, from α, and χειρ, 'hand.') One devoid of hands.

ACHEIRIA, *Achi'ria*: (αχειρια.) The state of being devoid of hands.

ACHEROIS, (αχερωις, brought from Acheron,) Populus.

ACHIA, *Achiar*. A name given in India to the pickled shoots of the bamboo.

ACHIA, Atehar.

ACHIAR, Achia.

ACHIC'OLUM, *Achit'olus*, *Hidrote'rion*, *Suda'r-rium*, *For'nix*, *Tho'lus*, *Sudato'rimum*, *Sud'atory*, *Sweat'inghouse*, *Sweat'ingbath*. The sweating-room in the ancient bagnios. Sweating-baths have been called *Hum'mums*,—a word said to be of oriental origin. In the seventeenth century, they were known in England by the name of *hot-houses*, and were much frequented, especially by the ladies.

ACHILIA, Acheilia.

ACHILLE, *TENDON D'*, Achillis tendo.

ACHILLE'A AGE'RATUM, (after Achilles,) *A. visco'sa*, *Balsami'ta femini'ca*, *Eupato'rimum Mes'nes*, *Age'ratum*, *Cos'tus horto'rimum mi'nor*, *Maud'lin*, *Maud'lin Tan'sey*; (F.) *Achillè Visqueuse*; *Ord. Compositæ*; *Sex. Syst. Syngenesia Polygamia Superflua*,—has the same bitter and aromatic properties as tansy, and is used in like affections.

ACHILLEA ATRA'TA, *Her'ba Gen'ipi ve'ri*, (F.) *Achillèe Noire*, has similar virtues.

ACHILLEA MILLEFO'LIUM, *Achillea Myriophyl'lon*, *Chrysoc'oma*, *Millefo'limum*, *Chiliophyl'lon*, *Lum'bus Ven'eria*, *Com'mon Yar'row* or *Mil'foil*. (F.) *Millefeuille*, *Herbe aux charpentiers*. The herb and flowers, *Achillea*, (Ph. U. S.), have an aromatic smell, and a rough, bitterish, somewhat pungent taste. They have been used in dyspepsia, flatulence, hemorrhage, &c. An extract of the plant, made with proof spirit, has been called *Achille'num*; and is used by the Italians in intermittent fever.

ACHILLEA MOSCHA'TA, an Alpine plant, has been esteemed diaphoretic and vulnerary (?) A volatile oil called *Euprit d'Iva*, is obtained from it in Switzerland, which is much prized for its musky odour. It is said by Mérat and De Lens, to be the *Gen'ipi* of the Savoyards, which is usually referred to *Artemisia rupestris*.

ACHILLEA PTAR'MICA, *Ptar'mica*, *P. Vulga'ris*, *Pseudopy'rethrum*, *Pyrethrum sylves'trè*, *Dra'co sylves'tris*, *Sternutament'ria*, *Dra'conculcus Prate'nsis*, *Sneezewort*, *Bas'tard Pell'titory*. (F.) *Herbe à éternuer*. The roots and flowers have a hot, biting taste, approaching that of pyrethrum. Their principal use is as a masticatory and sialagogue.

ACHILLEA VISCOSA, *A. Ageratum*.

ACHILLÉE NOIRE, *Achillea atrata*—*a. Visqueuse*, *Achillea ageratum*.

ACHILLEINUM, see *Achillea Millefolium*.

ACHILLE'IS, (αχιλλυις.) A beautiful species of barley, mentioned by Theophrastus and Galen, called after Achilles, a labourer. The decoction was used in fevers and jaundice.

ACHILLEUM, *Telephium*.

ACHIL'IS TEN'DO, *Fu'nis HIPPOC'RATIS*, *Cor'da seu Chor'da Hippoc'ratis*, *Cor'da magna*, *Ner'vus lat'us*, *Heelstring*. (F.) *Tendon d'Achille*. The strong tendon of the gastrocnemii muscles above the heel: so called, because it was the only vulnerable part of Achilles, or because of its strength. See *Tendon*.

ACHILUS, Acheilus.

ACHIMBASSI. An archiater or chief of phy-

sicians. A name given, at Grand Cairo, to a magistrate who licenses physicians.

ACHIR, Acheir.

ACHIRIA, Acheiria.

ACHITOLUS, Achicolum.

ACHLYS, (αχλυσ, 'a cloud.') Caligo.

ACHMELLA, *Spilanthus acmella*.

ACH'NÈ (αχνη.) Lint. See *Linten* n. Also, small mucous flocculi seen in front of the cornea.

ACHOL'IA, (αχολια, from α, and χολη, 'bile.') Deficiency or want of bile. Also, Asiatic cholera.

A'CHOLUS, (αχολος.) One deficient in bile.

ACHOR, gen. *Acho'ris*, Porrigo larvalis.

ACHO'RES. A term often employed by the ancients to designate both *crusta lac'tea*, and small superficial ulcerations on the skin of the face and head. See *Porrigo Larvalis*.

ACHORES CAPITIS, Porrigo scutellata.

ACHORION LEBERTII, see *Porrigo decalvans*—*a. Schonleini*, see *Porrigo favosa*.

ACHORIS'TUS, (α and χωρίζω, 'I separate.') Any sign which necessarily accompanies a state of health or disease.

ACHOUROU. The Carai'b name for a species of myrtle used in dropsy.

ACHRAS AUSTRALIS, (αχρας, 'the wild pear,') *Sapota*—*a. Sapota*, *Sapota*—*a. Zapota*, *Sapota*.

A'CHROI, (αχροι,) *Achromatis'ti*, *Achro'mati*, *Achro'mi*, (α, and χρωμα, 'colour.') Pale individuals. It is nearly synonymous with *λειψαιμοι*, *leiphæ'mi*, persons without colour; bloodless.

ACHRO'MA, gen. *Achro'matis*, *Chloas'ma al'b-um*, *Leucopath'i'a partial'is acquis'ita*, *Pac'il'ia*. Partial privation of colour of the skin.

Achroma vitili'go of Alibert, *vitil'igo*, as in the pied negro.

ACHROMASIA, Decoloration.

ACHROMATI, (αχρωματοι,) Achroi.

ACHROMATIC, *Achromat'icus*. A lens, so constructed as to correct the aberration of refrangibility of common lenses, is so termed. The *Crystalline* is an achromatic lens.

ACHROMATISTE, (αχρωματιστοι,) Achroi.

ACHROMATOP'SIA, *Achromatop'sis*, *Achromatop'sy*, *Chromatopseudop'sia*, *Dyschromatop'sia*, *Parachro'ma*, *Pseudochro'mia*, *Parora'sis*, *Vit'ius de'color*, *Colour blindness*, *Idiop'tey*, *Dal'tonism*, (α, χρωμα, 'colour,' and οστρομαι, 'I see.') Incapability of distinguishing colours; a defect situate in the cerebral part of the visual organ. Persons so circumstanced have been termed, by Mr. Whewell, *Id'iopts*. See *Acyanoblepsia* and *Anerythropsia*.

ACHROMI, (αχρωμοι,) Achroi.

ACHROMODERMIE, (δερμα, 'skin.') See *Albino*.

ACHROMOTRICHOMIE, (τριξ, gen. τριχος, 'a hair.') See *Albino*.

ACHRYSTALLODIAPHANIE, (α, crystalline, δια, 'through,' and φαivo, 'I shine.') See *Cataract*.

ACHYLO'SIS, (αχυλος, 'juice, chyle,' and *osis*.) Defective chylous or formation of chyle.

ACHYMO'SIS, (α, χυμος, 'juice, chyle,' and *osis*.) Defective chymification.

ACHYRON'THES RE'PENS, (αχυρον, 'chaff,' and αβνη, 'a flower.') *Illice'brum polygonoi'des*, *Forty Knot*; indigenous. *Order*, *Amaranthaceæ*. A decoction of the plant is drunk as a diuretic in dropsy, ischuria, &c.

ACHYRODES, (αχυρωδης,) *Acerosus*.

ACHYRON, (αχυρον,) *Furfur*.

A'CIA, (from αcis, gen. ακιδος, 'a point.') A word used by Celsus, which has puzzled commentators,—some believing it to have meant a needle; others the thread; and others, again, the kind of suture. "*Acia mollis, non nimis torta*." (Chifflet thinks it meant the thread.—Antwerp, 1638.)

AC'ID, *Ac'idus*, *Ox'ys*. (F.) *Acide*, *Aigre*, (from *ακίς*, gen. *ακίδος*, 'a point.') Sharp; sour; especially as applied to odorless or sapid substances. The French also use the term *aigre*, when referring to the voice, in the sense of sharp and shrill:—as *une voix aigre*, *voix as'pera*.

ACID, ACETIC, Aceticum acidum—a. Acetic, of commerce, Pyroligneous acid—a. Acetic, dilute, see Aceticum acidum—a. Acetous, strong, Aceticum acidum—a. Aerial, Carbonic acid—a. Antimonious, Arsenic diaphoreticum—a. Arsenious, Arsenicum album—a. Auric, see Gold—a. Azotic, Nitric acid—a. Benzoic, Benjamin, flowers of—a. Benzoic, Hippuric acid—a. Bezardic, Uric acid—a. Boric, Boracic acid—a. Calcareous, Carbonic acid—a. Calculus, Uric acid.

ACID, CARBAZOTIC, *Ac'idum Carbazot'icum* seu *Picrin'icum* seu *Pic'ricum*, *Carbonit'ric* or *Pic'ric* acid, *Welter's Bit'ter*, (F.) *Acide Carbazotique*, *A. Picrique*, *Amer de Welter*. This acid is frequently formed by the action of concentrated nitric acid on animal and vegetable substances, and especially on oil of tar. It, as well as the carbozates or picrates, is tonic and astringent, and said to be antiperiodic.

ACID, CARBOLIC, Acid, phenic—a. Carbonic, see Carbonic acid—a. Carbonitric, Acid, Carbazotic—a. Carbonous, Oxalic acid—a. Caseic, Lactic acid—a. Chromic, see Chromic acid—a. Citric, Citric acid—a. Crotonic, see Croton tiglium—a. Cyanhydric, Hydrocyanic acid—a. Cyano-hydric, Hydrocyanic acid—a. Delpich, see Acidum Valerianicum—a. Galactic, Lactic acid.

ACID, GAL'LIC, *Ac'idum Gall'icum*. (F.) *Acide Gallique*. This acid, prepared from galls, is found in most of the astringent plants that contain tannic acid of the kind obtained from galls. It is in delicate silky needles, usually somewhat yellowish, inodorous, and of a harsh, somewhat astringent taste. It dissolves in one hundred parts of cold and three parts of boiling water. It is very soluble in alcohol, and but slightly so in ether.

It has been highly extolled in internal hemorrhage, especially from the urinary organs and uterus. Dose from ten to twenty grains.

The Pharmacopœia of the United States directs it to be made by exposing a thin paste of powdered galls and distilled water for a month, adding the water from time to time to preserve the consistence; expressing the paste; boiling the residue in distilled water; filtering through animal charcoal, and crystallizing.

ACID, GASTRIC, see Gastric juice.

ACID, HIPPU'RIC, *Ac'idum Hippur'icum*, *Urobenzo'ic*, *Urinobenzo'ic* or *Benzur'ic* acid. An acid found in the urine of graminivorous animals. It is contained in human urine, especially after benzoic acid has been taken. See Hippuria.

ACID, HYDRIOD'IC, *Ac'idum Hydriod'icum*. This acid may be made by mixing solutions of iodide of potassium and tartaric acid; filtering the liquor to separate the bitartrate of potassa, and adding water to make the resulting hydriodic acid of definite strength.

A formula for the preparation of *acidum hydriodicum dilutum* has been introduced into the Pharmacopœia of the United States of 1863. It is a good solvent for iodine.

It has been used in the same cases as the preparations of iodine in general, but is rarely employed.

ACID, HYDROCHLORONITRIC, Nitro-muriatic acid—a. Hydrocyanic, Hydrocyanic acid—a. Hydrocyanic, dilute, see Hydrocyanic acid—a. Hydro-sulphuric, Hydrogen, sulphurette—a. Hydrothionic, Hydrogen, sulphuretted—a. Igasuric: see Jatropha curcas.

ACID, IOD'IC, *Ac'idum Iod'icum*, (F.) *Acide Iodique*. This is obtained by boiling iodine with nitric acid: or by decomposing iodate of baryta

by dilute sulphuric acid. It is a white, transparent solid, slightly deliquescent, and very soluble in water. It has been given with sulphate of quinia in hoarseness, scrofula, incipient phthisis, chronic inflammation, syphilis, etc. Dose three to six grains, or more.

ACID, LACTIC, see Lactic acid.

ACID OF LEMONS, Citric acid—a. Lithenic, Uric acid—a. Lithiasic, Uric acid—a. Lithic, Uric acid—a. Manganic, see Potassa permanganas—a. Marine, dephlogisticated, Chlorine—a. Marine dulcified, Spiritus ætheris muriatici—a. Mephitic, Carbonic acid—a. of Milk, Lactic acid—a. Muriatic, see Muriaticum acidum—a. Muriatic, dilute, Muriaticum acidum—a. Nanceic, Lactic acid—a. Nitric, see Nitric acid—a. Nitric, dilute, see Nitric acid—a. Nitro-hydrochloric, Nitro-muriatic acid—a. Nitro-muriatic, see Nitro-muriatic acid—a. Nitrous, dephlogisticated, Nitric acid—a. Oxyseptonic, Nitric acid.

ACID, PHEN'IC, *A. Phenyl'ic*, *Acidum phen'icum* seu *phenyl'icum*, *Carbol'ic* Acid, *Phenol*. A hydrated oxide of phenyl, produced in the manufacture of coal gas: (from *φαῖνω*, 'I produce light.') It is a heavy, transparent oil, having the taste and smell of creosote. In the proportion of one part to 40 of acetic acid to 100 of water, it has been applied, by means of a compress wetted with it, in porrigo, scabies, &c.; but its most important use is in the preservation of dead bodies.

ACID, PICRIC, Acid, carbazotic—a. Polygalic, see Polygala senega—a. Prussic, Hydrocyanic acid—a. Pyroligneous, see Aceticum acidum—a. Pyrolignic, Pyroligneous acid—a. Rosacic, see Porphyria—a. of Sorrel, Oxalic acid—a. of Sugar, Oxalic acid—a. Sulphhydric, Hydrogen sulphuretted—a. Sulpho-hydric, Hydrogen sulphuretted—a. Sulphuric, see Sulphuric acid—a. Tannic, Tannin—a. Uric, Uric acid—a. Urinobenzoic, Acid, hippuric—a. Urous, Uric oxide—a. Urylic, Uric acid—a. Valerianic, Acidum valerianicum—a. Valeric, Acidum valerianicum.

ACID, PNEU'MIC, *Ac'idum Pneu'micum*, (*πνεύμων*, 'lung.') An immediate crystallizable principle, found by M. Verdeil in the substance of the lung of the mammalia, and which gives it its acid reaction. By decomposing the carbonate or bicarbonate of soda of the blood, it is said to furnish a part of the carbonic acid expired.

ACIDE ACÉTIQUE FAIBLE, see Aceticum acidum—a. Azotique, Nitric acid—a. Boracique, Boracic acid—a. Chromique, Chromic acid—a. Gallique, Acid, gallic—a. Hydrocyanique, Hydrocyanic acid—a. Hydro-sulphurique, Hydrogen, sulphuretted—a. Iodique, Acid, iodie—a. Lactique, Lactic acid—a. Lithénique ou Lithiasique ou Lithique, Uric acid—a. Nitrique, Nitric acid—a. Phosphorique, Phosphoric acid—a. Picrique, Acid, carbazotic—a. Pneumique, Acid, pneumatic—a. Prussique, Hydrocyanic acid—a. Pyroacétique, Pyroligneous acid—a. Pyrolignique, Pyroligneous acid—a. Sulfureux, Sulphurous acid—a. Sulfurique, Sulphuric acid—a. Sulfurique délayé, Sulphuricum acidum dilutum—a. Tannique, Tannin—a. Tartrique, Tartaric acid—a. de l'urine, Uric acid—a. Urique, Uric acid—a. Valérianique, Acidum valerianicum.

ACIDITATIO, Acidities.

ACIDITIES, *Acid'ies*, *Aciditatio*, *Ac'idum morbo'sum* seu *prima'rum viarum*, *Ox'ytes*, *Sor'des ac'idæ*, (F.) *Aigreurs*. Sourness of the stomach, the result of indigestion, indicated by acid eructations, etc. The affection is very common in children, and must be obviated by absorbents, as magnesia, chalk, etc., and by regulated diet.

ACIDOLOG'IA, (*ακίς*, gen. *ακίδος*, 'a point, a sharp instrument,' and *λόγος*, 'a description,') A description of surgical instruments.

ACIDOMETER, (F.) *Acidomètre*, *Pèse-acide*,

pimple or tubercle on the face.—Gorræus. Fösius thinks the word ought to be *Acne*; and, according to Cassius, it is, at all events, derived from *ακμη*, 'vigour,' the disease affecting those in the vigour of life especially.

Willan and Bateman have adopted the term in their Nosology of cutaneous diseases, and placed it in the order TUBERCULA. *Acne*, with them, is an eruption of distinct, hard, inflamed tubercles, sometimes continuing for a considerable length of time, and sometimes suppurating slowly and partially. They usually appear on the forehead, temples, and chin, and are common to both sexes; but the most severe forms are seen in young men. They require but little management, and consist of four varieties: *Acne indurata*, *A. simplex*, (*Hypochondria*), *A. punctata* (*Don't this varus punctatus*, *Punctata mucosa*, *Varus punctatus*, *Crinoides*, *Comedones* or *Maggot Pimple*, *Worms*, *Grubs*, (F.) *Tanne*), and *A. rosacea*.—See *Gutta Rosea*.

ACNE INDURATA, see *Acne*—*a. Mentagra*, *Sycosis*—*a. Punctata*, see *Acne*—*a. Rosacea*, *Gutta Rosea*, see *Acne*—*a. of the Throat*, *Pharyngitis*, follicular.

ACNÉ, *Acne*—*a. Molluscoïde*, *Molluscum*.

ACNÉMIA, *Acne'mia*, (*a*, and *κνημη*, 'the leg.') Absence of legs.

ACNÉS'TIS, (*ακνηστίς*, from *a*, and *κναιν*, 'to scratch.') The part of the spine which extends, in quadrupeds, from between the shoulders to the loins. According to Pollux, the middle of the loins. The vertebral column.

ACOE, (*ακον*), *Audition*, *Ear*.

ACŒLIS, (*a*, and *κοιλία*, 'belly.') Devoid of belly. One who is so emaciated as to appear to have no belly.

ACOËMETER, (*ακον*, 'hearing,' and *μετρον*, 'measure,') *Acoumeter*.

ACŒMETRUM, *Acoumeter*.

ACOENOSI, (*νοσος*, 'a disease,') *Aconusi*.

ACOESIS, *Audition*.

ACOGNOSIA, (*ακος*, 'a remedy,' and *γνωσις*, 'knowledge,') *Pharmacognosia*.

ACOLASIA, (*a*, and *κολαζω*, 'I restrain,') *Intemperance*.

ACOLOGY, (*ακος*, 'a remedy,' and *λογος*, 'a discourse,') *Materia Medica*. According to some, restricted to surgical instruments and their uses—operative surgery; like *aciurgia*, (from *ακίς*, 'a point, a sharp instrument,')

ACONE, (*ακων*,) *Mortar*.

ACONIT À GRANDS FLEURS, *Aconitum cammarum*—*a. Salutare*, *Aconitum anthora*.

ACONITA, see *Aconitum napellus*.

ACONITE, *Aconitum*.

ACONITI FOLIA, see *Aconitum*—*a. Radix*, see *Aconitum*.

ACONITIA, see *Aconitum napellus*.

ACONITIN, see *Aconitum napellus*.

ACONITINE, see *Aconitum napellus*.

ACONITIUM, see *Aconitum napellus*.

ACONITUM, (from *Ακονίη*, a place in Bithynia, where it is common.) *Cynocotonon*, *Pardalianches*, *Pardalianchum*, *Cnici'da*, *Ac'onite*, *Wolfsbane*, *Monks'hood*, *Ord. Ranunculaceæ*. See *Syst. Polyandria Trigynia*.

Aconitum, *Aconite*, in the Pharmacopœia of the United States, 1842, is the leaves of *Aconitum napellus*, and *a. paniculatum*. In the edition of 1851, *Aconiti folia*, of 1863, *A. folium*, is the official name for the leaves, and *Aconiti radix* for that of the root of *Aconitum napellus*.

ACONITUM ANTHORA, *A. saluti'ferum* seu *Nemoroum* seu *Candollei* seu *Jacquinii* seu *Eulophum* seu *Anthoroïdeum*, *Anthora vulgaris*, *Anthora*, *Antithora*, *Salutary Monks'hood*, *Wholesome Wolfsbane*, *Yellow helmetflower*, (F.) *Aconit salutare*. The root of this variety, as of all the rest, is poisonous. It is used as a cathartic and anthelminthic. Dose ℥ss to ʒj.

ACONITUM ANTHOROIDEUM, *A. anthora*.

ACONITUM CAMMARUM, *A. paniculatum*, *A. macranthum*, *A. Kusnezovii*, (F.) *Aconit à grands fleurs*, resembles *Aconitum Napellus* in properties.

ACONITUM CANDOLLEI, *A. anthora*—*a. Eulophum*, *A. anthora*.

ACONITUM FEROX, a most violent ranunculaceous poison, which grows in Nepaul, and is the *Bish* or *Bikk* poison of that country. Its effects are like those of *Aconitum Napellus*.

ACONITUM JACQUINI, *A. anthora*—*a. Kusnezovii*, *A. cammarum*—*a. Macranthum*, *A. cammarum*.

ACONITUM NAPELLUS, *Aconitum Napellus* *ve'rus*, *Common Monks'hood* or *Wolfsbane*, *A. Nemomontanum*, (F.) *Chaperon de Moine*, *Coqueluchov*, *Tue-loup*. The leaves are narcotic, sudorific, and 'deobstruent' (?). They have been used in chronic rheumatism, scrofula, scirrhus, paralysis, amaurosis, etc. The active principle is called *Aconiti'na*, *Aconiti'na*, *Aconiti'na*, *Aconiti'um* or *Aconiti'ne*. It may be made by treating an alcoholic extract of the root with dilute sulphuric acid; precipitating by solution of ammonia; dissolving the precipitate in dilute sulphuric acid; treating with animal charcoal; again precipitating with solution of ammonia; washing with water, and drying. Ph. U. S., 1851. It requires 150 parts of cold and 50 of boiling water to dissolve it, but is readily dissolved by alcohol and ether. It neutralizes the acids, and forms with them uncrystallizable salts. It has been used internally, and especially applied externally, in neuralgic cases, iatroleptically and endermically.

Dose of *Aconitum*, gr. j. to gr. iij.

ACONITUM NEMOROSUM, *A. anthora*—*a. Nemontanum*, *A. napellus*—*a. Paniculatum*, *A. cammarum*—*a. Racemosum*, *Actæa spicata*—*a. Salutiferum*, *A. anthora*.

ACONITUM UNCINATUM, *Wild Monks'hood*, indigenous, is said to have similar virtues with *Aconitum napellus*.

ACONU'SI, *Acœn'osi*, *Acœn'osi*, (*ακον*, 'audition,' and *νοσος*, 'disease,') *Morbi aurium et auditus*. Diseases of the ear and audition.

ACOËNOSI, *Aconusi*.

AC'OPIS, (*a*, and *κοπος*, 'weariness,') Pliny gives this name to a precious stone, which was boiled in oil and used against weariness.

AC'OPON, *Ac'opum*. A remedy against weariness.

ACOPRIA, (*a*, and *κοπος*, 'excrement,') *Constipation*.

ACOPROSIS, *Constipation*.

ACOR BENZOÏNUS, (*acer*, 'sharp, acid,') Benjamin—*a. Boracicus*, *Boracic acid*—*a. Succineus*, *Succinic acid*—*a. Sulphuris*, *Sulphuric acid*—*a. Tartaricus*, *Tartaric acid*.

ACORE BÂTARD, *Iris pseudacorus*—*a. Faux*, *Iris pseudacorus*—*a. Odorant*, *Acorus calamus*.

ACORES, *Acids*, and *Acidities*.

ACOR'IA, (*ακορία*, from *a*, privative, and *κρεω*, 'I satiate,') *An inordinate or canine appetite*.

ACORITES, (*ακοριτης*,) A wine made of *Acorus*.

ACOR'MUS, (from *a*, privative, and *κορμος*, 'trunk,') A monster devoid of a trunk.

ACORN, *JUPITER'S*, *Fagus castanea*—*a. Oily*, *Guilandina moringa*—*a. Sardinian*, *Fagus castanea*.

ACORNS. See *Quercus alba*.

ACORUS ADULTERINUS, (*ακορος*,) *Iris pseudacorus*.

AC'ORUS CALAMUS, *A. Ve'rus* seu *Braziliensis*, *Calamus Aromaticus* seu *Odoratus* seu *Vulgaris*, *Typha Aromatic*, *Clav'va Rugo'sa*, *Sweetflag* or *Ac'orus*, *Flag-root*, *Sweet cane*, *Myrtle Flag*, *Sweet grass*, *Sweet root*, *Sweet rush*. (F.) *Jonc roseau* ou *Canne aromatique*. *Acore odorant*.

Ord. Aroidæ; Acoraceæ. (Lindley;) *Sec. Syst. Hexandria Monogynia.* The rhizoma—*Calamus* (Ph. U. S.—is stomachic and carminative, but is rarely used. It is regarded as a good adjuvant to bark and quinia in intermittents. The essential oil is used in like cases.

ACORUS PALUSTRIS, *Iris pseudacorus*—a. Vulgaris, *Iris pseudacorus*.

ACOS, (ακος,) Medicament.

ACOSMIA, (ακοσμία, from *a*, and κοσμος, 'order, ornament,') Disorder, irregularity in the critical days, according to Galen, who uses the word κοσμος, for regularity in those days. Others, and particularly Pollux, call bald persons ακοσμοι, because they are deprived of one of their most beautiful ornaments.

ACOU'ETER, *Acoum'eter*, *Acoem'eter*, *Acoë'trum*, *Acu'meter*, *Acusin'eter*, (F.) *Acoumètre*, (ακουω, 'I hear,' and μετρον, 'measure,') An instrument designed by M. Itard for measuring the degree of hearing.

ACOU'ETRE, *Acoumeter*.

ACOU'PHON'IA, *Cupho'nia*; (ακουω, 'I hear,' and φωνη, 'voice,') *Auscultatory Percussion*. A mode of auscultation, in which the observer places his ear on the chest, and analyzes the sound produced by percussion.

ACOUS'MA, gen. *Akous'matia*, (ακουσμα, 'any thing heard,') an imaginary noise. Depraved sense of hearing.

ACOUS'TIC, (ακουστικός,) *Acus'ticus*, (F.) *Acoustique*. That which belongs to the ear; as *Acoustic nerve*, *Acoustic trumpet*.

ACOUS'TIC MEDICINE is one used in diseased audition.

ACOUS'TICO-MALLÉEN, *Laxator tympani*.

ACOUS'TICS, *Acus'tica*. (F.) *Acoustique*. The part of physics which treats of the theory of sounds. It is also called *Pho'nics*.

ACOUSTIQUE, *Acoustic*, *Acoustics*.

ACQUA BINELLI (L), (acqua, 'water,') *Aqua Binelli*—a. Brocchieri, *Aqua Brocchieri*—a. Monterossi, *Aqua Binelli*—a. di Napoli, *Liquor arsenicalis*—a. della Toffana, *Liquor arsenicalis*.

ACQUETTA (L), (dim. of *acqua*,) *Liquor Arsenicalis*.

AQUI, MIN'ERAL WA'TERS OF. These thermal sulphureous springs are in Piémont. Their temperature is 167° Fahr., and they contain sulphohydric acid and chloride of sodium.

ACQUI'RED DISEASES, *Mor'bi acquis'iti*, (M. adventit'ii, M. epictet'ii, *Adventit'ious diseases*, (F.) *Maladies acquises*. Diseases which occur after birth, and which are not dependent upon hereditary predisposition.

ACRAI'PALA, (a, and κραταια, 'drunkenness,') Remedies against the effects of a debauch.

ACRA'LEA, (ακρος, 'at the extremity,') The extreme parts of the body, as the head, hands, feet, nose, ears, &c. See *Acraea*.

ACRA'NIA, (a, and κρανιον, 'the cranium,') Want of cranium, wholly or in part.

ACRA'SIA, (ακρασία, from *a*, and κρασις, 'mixture,') Intemperance. Excess of any kind. Surfeit.

It has been employed to denote debility, synonymously with *Acratia*; but this may have been a typographical inaccuracy.

ACRAT'IA, (ακρατεία, from *a*, and κρατος, 'strength,') Impotence; weakness, fainting.

ACRATIS'MA, gen. *Acratism'atia*, (ακρατισμα, from *a*, and κραννμι, 'to mix,') A breakfast, consisting of bread steeped in wine, not mixed with water.

ACRATOM'ELI, (*acratum*, and μελι, 'honey,') Wine mixed with honey.

ACRATOPE'GÆ, *Akratope'gæ*, (a, κρατος, 'strength,' and πηγη, 'a spring,') Mineral waters having no marked chemical qualities.

ACRATOPOS'IA, (ακρατοποσία, from *acratum*,

and ποσις, 'drink,') The drinking of pure or unmixed wine.

ACRATUM, (ακρατον, from *a*, and κρανννμι, 'to mix,') Unmixed wine, *Acratum vinum*, *Vinum me'rum*.

ACRATURE'SIS, (*acratia*, and ονησις, 'the act of discharging the urine,') Inability to void the urine from paralysis of the bladder.

ACRÊ, (ακρα,) The extremity or tip of any thing, as of the nose.

AC'REA, (ακρεα,) *Aerote'ria*, (ακρον, 'the summit,') The extreme parts of the body, as the feet, hands, ears, &c.

Also the extreme parts of animals that are used as food, *Acroco'lia*.

AC'RID, (ακρος, 'pointed,') or from *axis*, 'a point,') *Acer*. An epithet for substances of a hot, biting taste, which occasion a disagreeable sense of irritation and constriction at the top of the throat; or which, when applied externally, cause heat and irritation of the skin.

ACRID HEAT, (F.) *Chaleur acre*, is one that causes a hot tingling sensation at the extremities of the fingers.

ACRID POISON, See *Poison*.

ACRIDS, in Pathology, are certain, imaginary substances, supposed by the humourists to exist in the humours, and to cause various diseases. See *Acrimony*.

ACRIDOPH'AGI, (ακριόφάγοι, from *ακρίς*, gen. *ακρίδος*, 'a locust,' and φαγω, 'I eat,') *Locust-eaters*. Acridophagous tribes are said to exist in Africa.

AC'RIMONY, *Acu'itas*, *Acrimo'nia*, (*acer*, 'acid,') *akis*, 'a point,') *Acrimony* of the humours. An imaginary acid change of the blood, lymph, &c., which, by the humourists, was conceived to cause many diseases.

ACRIN'IA, (a, and κρινω, 'I separate,') A diminution of the quantity, or a total suspension, of the secretions.

ACRIS, (*ακρίς*,) a sharp bony prominence. Also, the locust.

ACRI'SIA, *Acri'sis*, (from *ακρισία*, a, and *κρισις*, 'judgment,') A condition of disease, in which no judgment can be formed; or in which an unfavourable opinion must be given.

ACRISIS, *Acrisia*.

ACRIT'ICAL, *Ac'ritus*, (F.) *Acritique*. That which takes place without any crisis; or which does not foretell a crisis; as an *acritical symptom*, *abscess*, &c.

ACRITUS, (*ακριτος*,) *Acritical*.

ACRIVIOLA, (*acer*, 'sharp,' and *viola*, 'a violet,') *Tropæolum majus*.

A'CRO, *A'kro*, (from *ακρον*, 'top, extremity,') In composition means the same.

ACROBYS'TIA, *Acropos'thia*, (ακρον, 'top,' and βω, 'I cover,') The prepuce or the extremity of the prepuce. Also, circumcision (?)—Kraus.

ACROBYS'TIOLITH, (*acrobystia*, and λιθος, 'a stone,') Preputial calculus.

ACROBYSTITIS, (*acrobystia* and *itis*,) *Posthitis*.

ACROCHEIR, (*ακροχειρ*,) *Acrochir'*, *Acrocheir'on*, (*acro*, and χειρ, 'the hand,') The forearm and hand. Also, the hand.

ACROCHOR'DON, (*ακροχορδων*, from *acro*, and χορδη, 'a string,') A tumour which hangs by a pedicle. *Verru'ca pens'ilis*. A kind of hard wart.

ACROCHORIS'MUS, (*acro*, and χορευω, 'I dance,') A kind of dance, with the ancients, in which the arms and legs were violently agitated.

ACROCOLIA, (*ακροκωλία*, from *acro* and κωλοι, 'a member,') *Acrea*.

ACROCOLIUM, (*ακροκωλιον*,) *Acermion*.

ACROD'RYA, (*ακροδρυον*, from *acro*, and δρυς, 'a tree,') Fruits, as nuts, apples, &c.

ACRODYN'IA, *Erythe'ma acrod'ymum*, *E*.

acrodyn'ia, *Chiropodal'gia*, (F.) *Acrodynie*, *Mal des pieds et des mains*, (*acro*, and *odynē*, 'pain.') A painful affection of the wrists and ankles especially, which appeared in Paris as an epidemic, in 1828 and 1829. It was supposed by some to be rheumatic, by others to be owing to spinal irritation. It appears to have been the same as Dengue.

ACROLEXION, (*acro*, and *αλεξή*, 'ulna,') Olecranon.

ACROMASTIUM, (*acro*, and *μαστός*, 'breast,') Nipple.

ACROMIA, *Acromion*.

ACRO'MIAL, *Acromia'lis*. Relating to the Acromion.

ACROMIAL ARTERY, *Exter'nal Scap'ular*, *A. Arter'ia Thorac'ica humera'lis*, *Artère troisième des Thoraciques*,—(Ch.) *A. Thoracique humérale*, arises from the anterior part of the axillary artery, opposite the upper edge of the pectoralis minor. It divides into two branches, one, *superior*; the other, *inferior*,—the branches of which are distributed to the subclavius, serratus major antius, first intercostal, deltoid, and pectoralis major muscles, as well as to the shoulder joint, &c. They anastomose with the superior scapular, thoracic, and circumflex arteries.

ACROMIAL NERVES, *Ner'vi acromia'les*. Branches of the fourth cervical nerve, which are distributed to the acromial region.

ACROMIAL VEIN has the same arrangement as the artery.

ACRO'MIOCORACOIDEUS. Belonging to the acromion and coracoid process.

The triangular ligament between the acromion and coracoid process of the scapula is so called.

ACRO'MION, *Acro'mium*, *Acro'mia*, *Acro'mis*, (*ακρωμιον*, from *acro*, and *ωμος*, 'the shoulder,') *Os Acro'mium*, *Hum'erus sum'mus*, *Ar'mus sum'mus*, *Mu'cro hu'meri*, *Ros'trum porci'num*, *Ca'put Scap'u'lae*, *Acroco'lum*. The process which terminates the spine of the scapula, and is articulated with the clavicle.

ACROMIS, (*ακρωμις*,) *Acromion*.

ACROMPHALIUM, (*ακρομφαλιον*,) *Acromphalon*.

ACROM'PHALON, *Acrompha'lium*, (*acro*, and *σφαλος*, 'the navel,') The extremity of the umbilical cord, which remains attached to the fœtus after birth.

ACROMYLE, (*acro*, and *μυλή*, 'patella,') see Patella.

ACRONARCOTIC, see Poison.

ACRONYX, (*acro*, and *ονυξ*, 'the nail,') Growing in of the nail.

ACROPARALYSIS, (*acro*, and *παράλυσις*, 'palsy,') *Paral'ysis extremitat'um*, Palsy of the extremities.

ACROPOSTHIA, (*acro*, and *ποσθη*, 'prepuce,') *Acrobystia*. See Prepuce.

ACROPOSTHITIS, (*acroposthia*, and *itis*,) *Posthitis*.

ACROPSYLON, (from *acro*, and *ψιλος*, 'naked,') The extremity of the glans penis.

ACRO'RIA, (*ακρωρία*, from *ακρον*, 'top,') Vertex.

ACRORRHEU'MA, gen. *Acrorrheum'atis*, *Rheumatis'mus extremitat'um*, (*acro*, and *ῥευμα*, 'defluxion, rheumatism,') Rheumatism of the extremities.

A'CROS, (*ακρος*, 'at the extremity, top,') The strength of the Athleta, and of diseases; the prominences of bones; the extremity of the fingers, &c. See *Acrocheir*, *Acromion*, &c.

ACROSTICHUM POLYPODIOIDES, *Polypodium ineanum*.

ACROTHERIA, *Acreea*. See Extremity.

ACROTHERIASIS, *Acroteriasmus*.

ACROTHERIASMUS, *Acroteri'asis*, (*ακροτηρια*, 'the extremities;' hence *ακροτηριαζειν*, 'to mutilate,') Amputation of the extremities.

ACROTHYM'ION, *Akrothym'ion*, (*acro*, and

θυμον, 'thyme,') A kind of conical, rugous, bloody wart, a "moist wart," compared by Celsus to the flower of thyme.

ACROTICA, (from *Acreea*.) Diseases affecting the excretent functions of the external surface of the body.

Privacy of the fluids or emunctories that open on the external surface; without fever or other internal affection as a necessary accompaniment.

The 3d order of the class *Ecceit'ica* of Good.

ACROTISMUS, (*a*, and *κροτος*, 'pulse,') *Asphyxia*.

ACT, *Ac'tus*, (*actum*, past participle of *agere*, 'to do, 'a thing done,') The effective exercise of a power or faculty. The action of an agent. *Acte* is used by the French, to signify the public discussion, which occurs in supporting a thesis;—thus, *soutenir un Acte aux Écoles de Médecine*, is, 'to defend a Thesis in the Schools of Medicine.'

ACTA, *Gesta*.

ACTEA AMERICANA, (*ακταία*, and *ακτη*, 'the elder,') see *Actæa spicata*—*a. Cimicifuga*—*a. racemosa*.

ACTEA RACEMO'SA, *A. Cimicif'uga*, *Cimicif'uga racemo'sa*, *Macro'tys racemo'sa*, *Bot'rophia Serpenta'ria* (?), *Serpenta'ria ni'gra*, *Black snake-root*, *Rich'weed*, *Cohosh'*, *Squaw root*, *Rattleweed*, *Black Cohosh*. (F.) *Actée à grappes*, *Serpentaire noire*. Ord. Ranunculaceæ. See *Syst. Polyandria* Pentagynia. A common plant in the United States. The root, *Cimicif'uga* (Ph. U. S.), is asstringent; and, according to Barton, has been successfully used, in the form of a decoction, as a gargle in putrid sore throat. A decoction of the root cures the itch. It is acro-narcotic, and has been used in rheumatism, acute and chronic; chorea, &c.

ACTEA SPICA'TA, *Christophoria'na spica'ta*, *Aconi'tum racemo'sum*, *Bane'berry*, *Herb Christoph'er*. (F.) *Herbe St. Christophe*. A perennial herbaceous European plant, the root of which resembles that of the black hellebore. The root is cathartic, and sometimes emetic, and in overdoses may produce dangerous consequences.

Actæa America'na, of which there are two varieties, *A. al'ba* and *A. ru'bra*,—white and red *cohosh*,—is indigenous in the United States. It has the same properties as *A. spicata*.

ACTE, *Sambucus*.

ACTE, *Act*.

ACTÉE à GRAPPES, *Actæa racemosa*.

ACTIF, (*agere*, *actum*, 'to act,') *Active*.

ACTIO, *Action*, *Function*.

AC'TION, *Ac'tio*, *Opera'tio*, *Energi'a*, *Prax'is*. Mode in which one object influences another.

The *animal actions* are those that occur in the animal body: the *vital*, those that are essential to life: the *physiological*, those of a healthy character: the *pathological* or *morbid*, those that occur in disease, &c. The ancients divided the *physiological actions* into *vital*, *animal*, *natural*, *sexual*, *particular*, *general*, &c. See *Function*.

ACTIONES NATURALES, see *Function*.

ACTIVE, *Dras'ticus*, *Acti'vus*. (F.) *Sthénique*, *Actif*. This adjective is used, in *Pathology*, to convey the idea of superabundant energy or strength, *Sthénicus*, *Hypersthen'icus*, (F.) *Sthénique*. *Active symptoms*, *e. g.* are those of excitement. In *Therapeutics*, it signifies *energet'ic*:—as, an *active treatment*. The French use the expression *Médecine agissante*, in contradistinction to *Médecine expectante*. In *Physiology*, *active* has a similar signification, many of the functions being divided into active and passive.

ACTON. A village near London, at which there is a purgative mineral spring, like that at Epsom.

ACTUAL. Same etymon as *actives*. That which acts immediately. A term usually re-

stricted to the red-hot iron, or to heat in any form: in contradistinction to the *potential* or *virtual*, which is applied to caustics or escharotics.

ACTUARIUS. Originally a title of dignity given to the Byzantine physicians.

ACTUS, Act—*a. Parturitionis, Parturition.*

ACUDUCTOR, (acus, and ducere, ductum, 'to lead,') Porte-aiguille.

ACUITAS, (acus, 'a needle,') Acrimony.

ACULEUS LIGNEUS, (acus, 'a needle,') Splinter.

ACUMETER, Acoumeter.

A'CUPRESS'URE, Acupressu'ra, (acus, 'a needle, and premere, pressum, 'to press,') A process for arresting hemorrhage, proposed by Prof. Simpson, of Edinburgh, which consists in passing an appropriate needle or pin twice through the substance of a wound, so as to compress and close, by the middle portion of the needle, the tube of the bleeding vessel a line or two, or more, on the cardiac side of the bleeding point.

A'CUPUNCTURE, Acupunctu'ra, Stylostix'is, Acupuncta'tum, Acupunctu'ation, (acus, 'a needle, and punctura, 'a puncture,') A surgical operation, much in use amongst the Chinese and Japanese, which consists in puncturing parts with a very fine needle. It has been employed, of late years, in obstinate rheumatic affections, &c., and apparently with success. Acupuncture is likewise a mode of infanticide in some countries; the needle being forced into the brain through the fontanelles, or into the spinal marrow, &c.

ACURGIA, (axis, 'a point, a sharp instrument, and εγρον, 'work,') Surgery (operative).

ACUS, Needle—*a. Capitata, Pin*—*a. Invasignata, see Needle*—*a. Ophthalmica, see Needle*—*a. Paracentetica, Trocar*—*a. Paracentica, Trocar*—*a. Triquetra vulgaris, Trocar*—*a. Veneris, Eryngium campestre.*

ACUSIMETER, (acusis, and μετρον, 'a measure,') Acoumeter.

ACUSIS, (ακουσις, Audition.

ACUSTICA, Acoustics.

ACUSTICUS, (ακουστικός, Auditory.

ACUTE, Acut'us, Or'y's, (akis, 'a point,') (F.) Aigu. A disease which, with a certain degree of severity, has a rapid progress, and short duration, is said to be "acute."—*Oxyneose'ma, Oxyneoseos, Oxyneoseos.*

Diseases were formerly subdivided into *Morbi acutis'simi*, very acute, or those which last only three or four days: *M. subacutis'simi*, which continue seven days: *M. subacu'ti*, or those which last from twenty to forty days.

The antithesis to *acute* is *chronic*. *Acute*, when applied to pain, sound, cries, &c., means *sharp*.

ACUTENACULUM, (acus, 'a needle, and tenaculum, 'a holder,') Porte-aiguille.

ACYANOBLEP'SIA, (a, κυανος, 'blue, and βλέπω, 'I see,') Defective vision, which consists in incapability of distinguishing blue. See Achromatopsia.

ACYESIS, (a, and κυησις, 'pregnancy,') Sterilitas.

ACYRUS, Arnica montana.

ACYSTINERVIA, (a, κυστις, 'bladder, and νευρον, 'nerve,') Cystoparalysis.

ACYSTURONERVIA, (a, κυστις, 'bladder, ουρον, 'urine, and νευρον, 'nerve,') Cystoparalysis.

ACYSTUROTROPHIA, (a, κυστις, 'bladder, ουρον, 'urine, and τροφή, 'nourishment,') Cystatrophia.

ACYTERIUS, (ακυτήριος, from α, and κειν, 'to conceive,') Abortive.

AD, as a prefix, in Romanic words, signifies to. ADACA. The *Sphæranthus Indicus*, a Malabar plant, which is acrid and aromatic.

ADAC'RYA, (a, and δακρυα, 'I weep,') Defective secretion of tears.

AD.EMONIA, properly Ademonia, (αδemonia.) Anxiety.

ADAKO'DIEN. A Malabar plant of the family Apocynæ, used in that country in diseases of the eyes.

AD'ALI, Lip'pia. A Malabar plant, which the Orientals regard as an antidote to the bite of the *naja*.

ADAMANT, Diamond, Magnēt.

ADAMANTINE SUBSTANCE, Enamel of the teeth.

ADAMAS, (αδαμας, Diamond.

ADAMI'TA, Adami'tum, (from adamas.) A very hard, white calculus.—Paracelsus.

The first word has been used for stone in the bladder: the second for lithiasis or the calculous condition.

ADAM'S APPLE, Pomum Adami.

ADANSONIA DIGITATA, (after Michel Adanson, the French naturalist.) Baobab.

ADAP'TER, (ad and apto, 'I fit,') A tube employed in pharmaceutical operations for lengthening the neck of a retort; or in cases where the opening of the receiver is not large enough to admit the beak of the retort.

ADAR'CE, (αδαρκη, Adar'cion, Adar'cis. A concretion found about the reeds and grass in the marshy regions of Galatia, and hiding them, as it were: hence the name, from *a* privative, and *δερκοω, 'I see, 'It was formerly in repute for cleansing the skin from freckles, &c.*

ADARIGO, Orpiment.

ADARNECH, Orpiment.

ADARTICULATIO, (ad and articulare, 'to join,') Arthroïdia.

AD CU'TEM ABDOM'INIS (ARTE'RIA). The *superficial artery of the abdomen*,—a branch of the crural or femoral, which arises at the lower part of Poupart's ligament, and ascends towards the umbilicus, being distributed to the integuments.

ADDAD. A Numidian plant; bitter and poisonous.

ADDEPHAG'IA, Adep'hag'ia, αδδφφαγια, αδδφφαγια, (from αδδφν, 'in excess, and φαγειν, 'to eat,') Voraciousness. Galen and Hoffmann have given this name to voracious appetite in children affected with worms. Sauvages refers it to Eulimia. Also, the goddess of gluttony.

ADDER'S TONGUE, Ophioglossum vulgatum.

ADDISON, DISEASE OF, see Capsule, renal.

ADDITAMEN'TUM, (addere, additum, 'to adjoin,') A term once used synonymously with Epiph'ysis. It is now restricted to the prolongation of two cranial sutures, the lambdoidal and squamous.

ADDITAMENTUM COLI, Appendix vermiformis cæci—*a. Necatum, Olecranon*—*a. ad Sacro-lumbalem, see Sacro-lumbalis*—*a. Ulnæ, Radius*—*a. Uncatum ulnæ, Olecranon.*

ADDLE (Prov.), Abscess.

ADDUCENS OCULI, (ad and ducere, ductum, 'to draw,') Rectus internus oculi.

ADDUCTEUR DE L'ŒIL, Rectus internus oculi—*a. du Gros orteil, Adductor pollicis pedis*—*a. Premier ou moyen, Adductor longus femoris*—*a. du Pouce, Adductor pollicis manus*—*a. Second ou petit, Adductor brevis*—*a. Troisième ou grand, Adductor magnus.*

ADDUC'TION, Adduc'tio, Parag'o'ge. The action by which parts are drawn towards the axis of the body.

The muscles which execute this function are called *Adductors*.

ADDUCTOR MEDII DIGITI PEDIS, Posterior medii digiti pedis—*a. Oculi, Rectus internus oculi.*

ADDUCTOR METACAR'PI MIN'IMI DIG'ITI, Metacar'pens, Car'pometacar'pens min'imi dig'iti, is situate between the adductor and flexor, next

to the metacarpal bone. It arises, fleshy, from the unciform process of the os unciforme, and from the contiguous part of the annular ligament of the wrist, and is inserted, tendinous and fleshy, into the fore-part of the metacarpal bone of the little finger, from its base to its head.

ADDUCTOR OCULI, *Rectus internus oculi*.

ADDUCTOR POL'LCIS MANUS, *A. Pol'licis, A. adductor pollicis digiti, Metacarpophalangeus pollicis*—(Ch.) (F.) *Adducteur du ponce*. A muscle which arises, fleshy, from almost the whole length of the metacarpal bone of the middle finger, and is inserted into the inner part of the root of the first bone of the thumb.

ADDUCTOR POL'LCIS PEDIS, *Antithenar, Metatarsophalangeus pollicis*—(Ch.) *Tarso-metatarsi-phalangien du ponce*. (F.) *Adducteur du gros orteil*. Arises by a long thin tendon, from the under part of the os calcis, from the os cuboides, os cuneiforme externum, and from the root of the metatarsal bone of the second toe. It is divided into two fleshy portions, and is inserted into the external sesamoid bone, and root of the metatarsal bone of the great toe.

Bichat has given the general name, *Adductors*, to those of the interosseous muscles of the hand or foot, which perform the action of adduction.

ADDUCTOR TERTII DIGITI PEDIS, *Prior tertii digiti pedis*.

ADDUCTORS OF THE THIGH. These are three in number, which have, by some anatomists, been united into one muscle—the *Triceps Adductor Femoris*.

1. *Adductor longus femoris, A. femoris pri'mus, Triceps minor, Pubiofemor'lis*—(Ch.) (F.) *Premier ou moyen adducteur*. Arises by a strong tendon from the upper and fore part of the os pubis and ligament of the symphysis, at the inner side of the pectinialis. It runs downwards and outwards, and is inserted by a broad, flat tendon into the middle of the linea aspera.

2. *Adductor brevis, A. femoris secundus, Triceps secun'dus, Subpubiofemor'lis*—(Ch.) (F.) *Second ou petit Adducteur*. Arises tendinous from the os pubis, at the side of its symphysis, below and behind the last muscle. It runs obliquely outwards, and is inserted by a short, flat tendon into the inner and upper part of the linea aspera, from a little below the trochanter minor to the beginning of the insertion of the adductor longus.

3. *Adductor magnus, A. femoris tertius et quartus, Triceps magnus, Ischiofemor'lis*—(Ch.) (F.) *Troisième ou grand adducteur*, is much larger than either of the others. It arises from the ramus of the pubis, from that of the ischium, and from the tuber ischii, and is inserted into the whole length of the linea aspera. Near the lower part of the linea aspera it is pierced by a kind of oblique, fibrous canal, through which the crural artery and vein pass.

ADEC. The inner man, according to Paracelsus.

ADECTA, (*a*, and *δακνω*, 'I bite.') Sedatives.

ADELIPARIA, (*αδην*, 'to excess,' and *λιπαρος*, 'fat.') Polysarcia.

ADELODAGAM. A bitter Malabar plant, used in asthma, catarrh, and gout.

ADELPHIA, see Adelpheia.

ADELPHIX'IA, *Adelphix'is*, (*αδελφος*, 'brother.') Consanguinity of parts in health or disease; *Fraternitas, Fratratio*. Hippocrates used the word *Adelphia* for diseases that resemble each other.

ADELPHIXIS, (*αδελφίς*.) Sympathy.

ADEMONIA, (*αδμονία*.) Depression, Nostalgia.

ADEMOSYNE, (*αδμοσυνη*.) Depression, Nostalgia.

ADEX, gen. A'DENIS, (*αδην*, gen. *αδενος*, 'a gland,') in composition, gland.

ADEX ULCER, see Beriberi.

ADENAL'GIA, *Adenodyn'ia*, (*aden*, and *αλγος*, 'pain,') Glandular pain.

ADENECTOP'IA, (*aden*, and *εκτος*, 'removed from its place,') Dislocation of a gland.

ADENEMPHRAX'IS, (*aden*, and *εμφραξις*, 'obstruction,') Glandular obstruction.

ADÉNIES, (from *aden*.) *Adenopathies angibromiques*. Diseases of the glands connected with the digestive tube.

ADENIFORM, *Adeniform'is, Adenoï'des, Adenoid, aden and forma*, 'form or resemblance,') *Glandiform*, or resembling a gland.

ADÉNITE, Adenitis—*a. Lymphatique, Lymphadenitis*.

ADENITIS, (*aden* and *itis*.) *Phlegma'sia adeno'a seu glandulo'sa*. (F.) *Adénite*. Glandular inflammation.

ADENITIS LYMPHATICA, Lymphadenitis.

ADENITIS MESENTER'ICA, *Mesenter'ic Ganglion'itis*. Inflammation of the mesenteric glands.

ADENITIS PALPEBRARUM CONTAGIOSA, see Ophthalmia.

ADENOCE'LE, (*aden*, and *κλη*, 'a rupture,') 'a tumour.' Under this term Mr. Birkett includes tumours of the mammae, which contain structures similar to, if not identical with, the normal secreting tissue of the gland, and more especially the cystosarcoma, and chronic mammary tumours of surgical authors.

ADENOCHIRAPSOLOG'IA, (*aden*, *χειρ*, 'the hand,' *απρω*, 'I lay hold of,' and *λογος*, 'a description,') The doctrine of curing scrofula or the king's evil by the royal touch.

Adenocheradolog'ia and *Adenochoiradolog'ia* have been used with the same acceptance; as well as for a treatise on scrofula.

ADENOCHON'DRIUS, (*aden*, and *χονδρος*, 'a cartilage,') Relating to gland and cartilage,—for example, *Arthrophyma adenocho'n'drium*, a tumefaction of the glands and cartilages of joints.

ADÉNODERMIES SYPHILIT'QUES, *Asyphosiques*, (*aden*, and *δερμα*, 'skin,') Syphilitic affections of the glands of the skin.

ADENODYNIA, (*aden*, and *δυν*, 'pain,') Adenalgia.

ADENOG'RAPHY, *Adenogra'phia*, (*aden*, and *γραφω*, 'I describe,') That part of anatomy which describes the glands.

ADENOID, Adeniform.

ADENOIDES, Adeniform.

ADENOLOGADITIS, (*aden*, and *logaditis*.) Ophthalmia of the new born.

ADENOL'OGY, *Adenolog'ia*, (*aden*, and *λογος*, 'a description,') A treatise on the glands.

ADENOMALA'CIA, (*aden*, and *μαλακία*, 'softening,') Mollescence or softening of a gland.

ADENOMENINGEAL, see Fever, adenomeningeal.

ADENONCOSIS, (*aden*, *ογκος*, 'a tumour,' and *οσις*.) Adenophyma.

ADENONCUS, (*aden*, and *ογκος*, 'a tumour,') Adenophyma.

ADENONERVOUS FEVER, Plague.

ADÉNOPATHIES ANGI'BROMIQUES, (*aden*, and *παθος*, 'affection,') *Adénies*.

ADENOPHARYNGE'US, (*aden*, and *φάρυγξ*, 'a fleshy fibre,') Some fleshy fibres, which pass from the constrictor pharyngis inferior to the thyroid gland, have received this name. Their existence is not constant.

ADENOPHARYNGITIS. Inflammation of the tonsils and pharynx.

ADENOPHTHALMIA, Ophthalmia tarsi.

ADENOPHY'MA, gen. *Adenophym'atis, Adenon'cus, Adenonco'sis*, (*aden*, and *φυμα*, 'a swelling,') Swelling of a gland, or glandiform ganglion. (F.) *Glandage*. *Adenophyma* is used by

some to signify a soft glandular swelling;—*Adenoma*, one of a harder character.

ADENOPHYMA INGINALIS. Bubo.

ADENOSCIRRHUS. *Adenoscirrhos*, (aden, and *scirrhos*, 'induration,') Scirrhus induration of a gland.

ADENOSCLEROSIS, (*σκληρος*, 'hard,') Adenoscirrhus.

ADENOSIS SCROPHULOSA, (from *aden*, Scrofula.

ADENOSUS, (*Ἀδενός*, 'hard,') A hard, glandular abscess, which suppurates slowly.

ADENOSYNCHITONITIS, (aden, *synchiton*, 'the conjunctiva,' and *itis*, 'inflammation,') Ophthalmia Neonatorum.

ADENOTOMY, *Adenotomia*, (aden, and *τομή*, 'I cut,') Dissection of the glands.

ADEPHAGIA, (*ἀδελφία*, from *ἀδελφός*, 'in excess,' and *φαγών*, 'to eat,') Addephagia, Boulimia.

ADEPS, gen. *Adipis*, *Adeps Suillus*, *Oryngium*, *Pinguedo*, *Lardum*, *Lardum*, *Lard*, (Prov.) *Mort*, *Sallis*, *Seacineame*. Pig's flare. The fat of the hog. In the Ph. U. S. the prepared fat of *Sus scrofa*, free from saline matter.

ADEPS ANSERINUS, *Adeps anseris* or *Graisse greuse*, (F.) *Graisse d'Oie*, is emollient. It has been used as an emetic.

ADEPS CANTHARIDIBUS MEDICATUS, Unguentum lytte medicatum—a. *Cortice Daphnes gnidii medicatus*, Unguentum epispasticum de Daphne gnidio—a. *Humanus*, *Liquamunia*—a. *Hydrargyro medicatus*, Unguentum Hydrargyri—a. ex Hydrargyro mitius dictum cinereum, Unguentum oxidi hydrargyri cinereum—a. *Hydrargyri murate oxygenato medicatus*, Unguentum muratis hydrargyri oxygenati medicatum—a. *Hydrargyri nitrate medicatus*, Unguentum hydrargyri nitratis—a. *Hydrargyri oxido rubro et plumbi acetate medicatus*, Unguentum ophthalmicum—a. *Lauro medicatus*, Unguentum laurinum—a. *Myristicea*, see *Myristica*—a. *Ovilli*, *Sevum*—a. *Papavere*, *hyoscyamo*, et *belladonna* medicatus, Unguentum populeum.

ADEPS PRÆPARATUS, *A. Suillus curatus* (see *Cura*,) *A. Sullus Præparatus*, *Hog's lard*, *Barrard's grease*, *Lard*, *Ac'unge*, *Azeungia*, *Azeungia porcina*, (F.) *Graisse de Porc*, *Saindouze*, *Acunge*, is prepared by melting pig's flare, and straining it. This is called *rendering* the lard. Lard is emollient, but is chiefly used for forming ointments and plasters.

ADEPS SULLUS PRÆPARATUS, *Adeps Præparatus*—a. *Sulfure et ammonie murate medicatus*, Unguentum sulphuratum ad seabium—a. *Sulfure et carbonate potassæ medicatus*, Unguentum sulphuratum alcalinum ad seabium—a. *Tartaro stibii medicatus*, Unguentum antimonii tartarizati—a. *Oxido zinci medicatus*, Unguentum oxidi zinci impuri.

ADEPT, (*adip'iscor*, *adeptus*, 'to acquire,') Alchymist.

ADEPTA MEDICINA. Medicine, which treated of diseases contracted by celestial operations, or communicated from heaven.

ADEPTA PHILOSOPHIA, Alchymy.

ADERMONERVIA, (*a, δερμα*, 'skin,' and *νεῦρον*, 'nerve,') Dermatanervia.

ADERMOTROPHIA, (*a, δερμα*, 'skin,' and *τροφή*, 'nourishment,') Extenuation or atrophy of the skin.

ADFLATUS, Afflatus.

ADHERENTIA, (*ad*, and *hære*, hæsum, 'to stick,') Adherence.

ADHESIO, Adherence.

ADHATO'DA, *Justicia adhato'da*. The *Mahabar' Nat Tree*, (F.) *Noyer de Ceylon*. Used in India for expelling the dead foetus in abortion. The word is said to convey this meaning in the Ceylonese.

ADHERENCE, *Adhæsi*, *Adhærentia*, *Con-*

cre'tio, *Pros'physis*, *Procoll'e'sis*, *Adhæ'sis*, (*from adhære*, [*ad* and *hære*,] 'to stick to,') These words are usually employed synonymously. The French often use *adherence* for the state of union, and *adhesion* for the act of adhering.

ADHESION, Adherence.

ADHÉSIF, Adhesive.

ADHÉSIVE, *Adhæ'sivus*, *Adhæ'sivus*, (F.) *Adhésif*; same etymon as Adherence. Sticking, uniting. An epithet,—as '*adhesive plaster*,' see *Emplastrum resinae*.

ADHÉSIVE INFLAMMATION is that inflammation which terminates by an adhesion between inflamed and separated surfaces, and which was, at one time, supposed to be necessary for such adhesion.

ADIANTHUM, Adiantum.

ADIANTUM, (*ἀδίατρον*, from *a*, and *δίατρον*, 'to grow wet,' from the leaves not being easily moistened,) A. pedatum.

ADIANTUM ETHIOPIUM. A South African plant, *Ord. Filices*, an infusion of which is sometimes used as an emollient in coughs, and in diseases of the chest.

ADIANTUM ALBUM, *Asplenium ruta muraria*—a. *Aureum*, *Polytrichum*.

ADIANTUM CAPILLUS VENERIS, *A. Corsan'dri-folium* seu *Nigrum*, *Capillus Veneris*, *Maidenhair*, (F.) *Capillaire de Montpellier*. A European plant, of feeble, aromatic and demulcent properties. It is used for forming the *Sirap de Capillaire* or *Capillaire*.

ADIANTUM CORIANDRIFOLIUM, *A. Capillus Veneris*—a. *Nigrum*, *A. Capillus Veneris*.

ADIANTUM PEDATUM, *A. Canadense* seu *Pa'tens*, *Adiantum*, *Capillus Veneris Canadensis*, *Herba Veneris*, *Filix Veneris*, *Canada Maidenhair*, *American Maidenhair*, *Rock'fern*, *Sweet'fern*, (F.) *Capillaire du Canada*, has the same properties. *Capillaire* was once made from this. See *Adiantum capillus Veneris*.

ADIANTUM RUBRUM, *Asplenium trichomanoides*.

ADIAPHANIE PÉRICHRYSAITIQUE, (*a, dia*, 'through,' and *παύω*, 'I shine,') See *Cataract*.

ADIAPHOROSIS, *Adiaphorosis*, (*a, dia*, 'through,' *πορος*, 'a pore,' and *osis*,) Defect or suppression of perspiration, *Adiapneustia*.

ADIAPHOROUS, *Adiaph'orus*, *Indiff'erens*, *Neut'ral*. A medicine which will neither do harm nor good.

ADIAPNEUSTIA, (*adiapneustia*, from *a, dia*, 'through,' and *πνέω*, 'to breathe,') *Adiaphorosis*.

ADIARRHŒA, (*a*, and *διάρρην*, 'to flow,') Retention of any excretion.

ADIATHESIC, *Adiathes'icus*, (*a*, and *διαθεσις*, 'diathesis,') In the system of *controstimulism*, *adiathesic diseases* are such as occur without any antecedent diathesis.

ADICE, (*ἀδικέω*, 'I injure,') *Urtica*.

ADIPATUS, (*adeps*, gen. *adipis*, 'fat,') *Fatty*.

ADIPEUX, *Adipose*.

ADIPOCERA, *Adipoc'era*—a. *Cetosa*, *Cetaceum*.

ADIPOCIRE, *Adipoc'era*, (*adeps*, gen. *adipis*, 'fat,' and *cera*, 'wax,') The base of biliary calculi, called also *Chal'exterine*. Also, a sort of soap, formed from animal matter under certain circumstances. (F.) *Gras des Caducres*, *Gras des Cimetières*. The human body, when it has been for some weeks in water, assumes this appearance; and it has been a subject of legal inquiry, what length of time is necessary to produce it. This must, of course, depend upon various circumstances, as climate, season, &c.

ADIPOCIRE DE BALEINE, *Cetaceum*.

ADIPOCIRIFORM, (*adipocire*, and *forma*, 'form,') See *Cholesteatoma*.

ADIPOSE, *Adip'ous*, *Adipo'us*, (*adeps*, gen. *adipis*, 'fat,') (F.) *Adipoux*. That which relates to fat—as *Adipose membrane*, *A. vessels*, &c. See *Fatty*.

AD'IPŒSE SARCO'MA of AB'ERNETHY, *Emphy'ma sarco'ma adiposum*, is suetty throughout, and enclosed in a thin capsule of condensed areolar substance, connected by means of minute vessels. It is chiefly found on the fore and back parts of the trunk. See Sarcoma.

ADIPO'SIS, (*adeps*, and *osis*.) See Polysarcia. ADIPO'SIS HEPAT'ICA, *Pimelo'sis* seu *Steatosis* seu *Mala'x'is hepatic'a*, *He'par adiposum*, *Fat'ty liv'er*, *Fat'ty degenera'tion of the liv'er*, (F.) *Dégénérescence graisseuse du Foie*. Fatty disease of the liver.

ADIPOSURIA, (*adeps*, gen. *adipie*, 'fat,' and *ουρον*, 'urine.') Pimeluria.

ADIPOSUS, Fatty.

ADIPOUS, Fatty.

ADIP'SIA, (*a*, and *διψα*, 'thirst,') *Dipso'sis* *xepers*. Absence of thirst.

ADIP'SON, *Adip'sum*. Any substance which relieves thirst. Applied to a decoction of barley to which oxymel was added.

ADIPSOS, Glycyrhiza.

AD'ITUS, 'an entrance,' 'an approach,' (*adire*, *aditum*, 'to go to,') *Pros'odos*. The entrance to a canal or duct, as *Ad'itus ad Aquæduc'tum Fallo'pii*.

ADITUS AD INFUNDIBULUM, Vulva.

ADIULIS'TOS, (*a*, and *διυλιζω*, 'I strain,') Unstrained wine for pharmaceutical purposes.

ADJUNCTUM, (*ad*, and *jungere*, *junctum*, 'to join,') Accompaniment.

ADJUTOR PARTÛS, ('an aider of birth,') Accoucheur.

ADJUVANT, *Ad'juvans*, (*adjuvare*, 'to aid,') A medicine, introduced into a prescription to aid the operation of the principal ingredient or basis. Also, whatever assists in the removal or prevention of disease.

ADMINICULE (F.), *Adminic'ulum*, (*ad*, and *manus*, 'a hand,') A prop, a support. Anything that aids the action of a remedy.

ADNASCENTIA, (*ad*, and *nascor*, 'to grow,') Prosphysis.

ADNATA (TUNICA), (*ad*, and *nascor*, *natus*, 'to grow,') Conjunctiva.

ADNÉE (MEMBRANE), Conjunctiva.

ADOLESCENCE, *Adolescen'tia*, *Juven'ta*, *Juven'tas*, *Juven'tus*, *Æ'tas bo'na*, *Youth*: from *adolescere*, (*ad*, and *olescere*, 'to grow,') (F.) *Jeunesse*. The period between puberty and that at which the body acquires its full development; being, in man, between the 14th and 25th years; and, in woman, between the 12th and 21st.

ADOLESCENS, *Ju'venis*, *Hebe'tes*, *Hebe'ter*, *Hypen'tes*. A youth. A young man in the period of adolescence.

ADO'LIA. A Malsbar plant, whose leaves, put in oil, form a liniment, used in facilitating labour.

ADOR, Zea mays.

ADORION, *Daucus carota*.

ADOUCISSANT, (from *adoucir*, 'to soften,') Demulcent.

AD PON'DUS OM'NIUM, 'To the weight of the whole.' In a prescription it means, that any particular ingredient shall equal in weight the whole of the others.

ADRAGANT, (corr. from *tragacantha*,) *Tragacantha*.

ADRAGANTHE, *Tragacantha*.

ADRA RIZA, (*adros*, *adra*, 'thick,' and *ρίζα*, 'a root,') *Aristolochia clematitis*.

A DROBO'LO'N, *adros*, 'thick,' and *βωλος*, 'mass,') The bellium of India, which is in larger pieces than that of Arabia.

ADROS, (*adros*, 'plump and full,') Applied to the habit of body, and also to the pulse.

ADSARIA PALA, *Dolichos pruriens*.

ADSPIRATIO, (*ad*, and *spiro*, 'I breathe,') Aspiration, Inspiration.

ADSTANS, (*ad*, 'to or near,' and *stare*, 'to stand,') Prostate.

ADSTITES GLANDULOSI, Prostate.

ADSTRICTIO, (*ad*, and *stringere*, *strictum*, 'to bind,') Astriktion, Constipation.

ADSTRICTORIA, Astringtons.

ADSTRINGENS FOTHERGILLI, Kino.

ADSTRINGENTIA, Astringtons.

ADULAS'SO. The *Justic'ia bival'vis*. A small shrub, used in India as a local application in gout.

ADULT, (*ad*, and *olere*, *olitum*, 'to grow,') see Adult age.

ADULT AGE, *Andri'a*, *Viril'itas*, *Æ'tas viril'is*, *Viril'ity*, *Manhood*. The age succeeding adolescence, and preceding old age. In the civil law, an adult is one, who, if a boy, has attained the age of fourteen years; and, if a girl, of twelve. In the common law, one of full age. *Adult*, *Adultus*, is also used for one in the adult age.

ADULTERATIO, Falsification.

ADULTUS, see Adult age.

ADUNCATIO UNGUIUM, (*ad*, and *uncus*, 'a hook,') Onychogryphosis.

ADURENS, (*ad*, and *urere*, 'to burn,') Caustic.

ADURION, *Rhus coriaria*.

ADUST', *Adus'tus*, (from *adurere*, (*ad*, and *urere*,) 'to burn,') The blood and fluids were formerly said to be adust, when there was much heat in the constitution, and but little serum in the blood.

ADUSTIO, Adustion, Burn.

ADUST'ION, *Adus'tio*. State of the body described under Adust. In surgery, it signifies cauterization.

ADVEHENS, (*ad*, and *vehere*, 'to carry,') Affluent.

ADVENTITIOUS DISEASES, Acquired diseases.

ADVENTITIUS, (*ad*, and *venio*, *ventum*, 'to come,') Accidental.

ADYNAMIA, *Impoten'tia*; (*adynamia*, from *a*, and *δυναμις*, 'strength,') *Adyna'sia*, *Adyna'tia*. Considerable debility of the vital powers; as in typhus fever. Some Nosologists have a class of diseases under the name *Adynamia*, *Ec'lyase*, *Mor'bi asthen'ici*.

ADYNAMIA VIRILIS, Impotence.

ADYNAM'IC, *Adynam'icus*, *Hypodynam'ic*, *Hypodynam'icus*; same etymon as *Adynamia*. Appertaining to debility of the vital powers.

ADYNASIA (*adynasia*,) *Adynamia*.

ADYNATIA, (*adynatia*,) *Adynamia*.

ADYNATOCOMIUM, (*adynatus*, and *κομεω*, 'I take care of,') Hospital.

ADYNATODOCHIUM, (*adynatus*, and *δεχομαι*, 'to receive,') Hospital.

ADYNATUS, (*adynarus*,) Sickly.

ÆDŒA, (*aïdia*, from *aïdēs*, 'shame,') Genital Organs.

ÆDŒAGRA, (*ædæa*, and *αγρα*, 'seizure,') Gout in the genitals.

ÆDŒAGRAPHY, *Ædæagrap'hia*, *Ædæog'r'aphy*, (*ædæa*, and *γραφω*, 'I describe,') A description of the organs of generation.

ÆDŒAL'O'GY, *Ædæalog'ia*, *Ædæol'ogy*, (*ædæa*, and *λογος*, 'a description,') A treatise on the organs of generation.

ÆDŒAT'OMY, *Ædæato'mia*, *Ædæoto'mia*, *Ædæot'omy*, *Ædæot'omy*, (*ædæa*, and *τεμνω*, 'I cut,') Dissection of the parts of generation.

ÆDŒECI, (from *ædæa*,) Gennetici.

ÆDŒEITIS, *Ædæoti'tis*, *Medei'tis*, (*ædæa*, and *itis*,) Inflammation of the genital organs.

ÆDŒEULENORRHŒA, Leucorrhœa.

ÆDŒODYN'IA, (*ædæa*, and *δύνη*, 'pain,') Pain in the genitals. Pudendagra.

ÆDEOGARGALUS, (*ædæa*, and *γαργαλιζω*, 'to tickle.') Masturbation. Nymphomania.

ÆDEOGARGARISMUS, Masturbation, Nymphomania.

ÆDEOGRAPHY, Ædæography.

ÆDEOLOGY, Ædæology.

ÆDEOMANIA, (*ædæa*, and *mania*), Nymphomania.

ÆDEOMYCODERMITIS, (*ædæa*, *μυκος*, 'mucous,' *δερμα*, 'skin,' and *itis*.) Inflammation of the mucous membrane of the genital organs.

ÆDEON, (*αιδεον*.) Inguen.

ÆDEOPSOPHESIS, Ædæopsophia.

ÆDEOPSOPHIA, *Ædæopsophia*, (*ædæa*, and *ψοφειν*, 'to make a noise.') Emission of wind by the urethra in man, by the vagina in woman.

ÆDEOPSOPHIA UTERINA, Physometra.

ÆDEOSCOPY, *Ædæosco'pia*, (*ædæa*, and *σκοπειν*, 'I view.') Exploration of the organs of generation.

ÆDEOTITIS, Ædœitis—æ. Gangrænosa, Colpocæ—æ. Gangrænosa puellarum, Colpocæ infantilis—æ. Gangrænosa puerperarum, Colpocæ puerperarum.

ÆDEOTOME, Ædæatomy.

ÆDEOTOMIA, Ædæatomy.

ÆDEOTOMY, Ædæatomy.

ÆDOPTOSIS, (*ædæa*, and *πτωσις*, 'a falling.') Hysteroptosis—æ. Uteri, Prolapsus uteri—æ. Uteri inversa, Uterus, inversion of the—æ. Uteri retroversa, Retroversio uteri—æ. Vaginæ, Prolapsus V.—æ. Vesicæ, Exocyste.

ÆEIG'LUCES, *Æig'luces*, (*αι*, 'always,' and *λυκος*, 'sweet.') A kind of sweet wine or must.

ÆGLE MARMELOS, (from *αγλη*, 'splendor,') Covolam.

ÆGAGROPILA, *Ægagropi'lus*, (*αιγpayos*, 'the rock goat,' and *πιλος*, 'hair,') *Bézoar d'Allemagne*, *Pila Dama'rum* seu *Rupicapra'rum*. A ball composed of hairs, found in the stomach of the goat: once used medicinally.

ÆGEIROS, (*αιγειρος*.) Populus.

ÆGER, Sick.

Æ'GIAS, *Æ'gis*, *Æg'lia*, *Æ'gides*, (from *αιγίς*, 'a goat's skin?') (F.) *Aige* or *Aigle*. There is obscurity regarding the precise meaning of this word. It was used to designate an ulcer, or speck on the transparent cornea.

Maître Jean uses it for a calcareous deposit between the conjunctiva and sclerótica.

ÆGIDES, Ægias.

Æ'GILOPS, *An'chilops*, *An'kylops*, (*ægo*, and *ωψ*, 'the eye.') An ulcer at the greater angle of the eye, which sometimes does not penetrate to the lachrymal sac, but at others does, and constitutes fistula lacrymalis.

ÆGIRINON, (from *ægeiros*.) An ointment of which the fruit or flower of the poplar was an ingredient.

ÆGLIA, Ægias.

ÆGO, (from *αιξ*, gen. *αιγος*, 'a goat.') In composition, goat.

ÆGOCERAS, (*ægo*, and *κερας*, 'horn,') *Trigonella fœnum*.

ÆGOLETHRON, (*ægo*, and *ὀλλυνμι*, 'to destroy,') *Ranunculus flammula*.

ÆGONYCHON, (*ægo*, and *ὄνυξ*, gen. *ονυχος*, 'a hoof,') *Lithospermum officinale*.

ÆGOPHONIA, Egophony.

ÆGOPHONICUS, Egophonic.

ÆGOPODIUM PODAGRARIA, (*ægo*, and *ποδῖον*, 'a little foot,') *Ligusticum podagraria*.

ÆGRIPPA, Agrippa.

ÆGRITUDO, gen. *Ægritu'dinis*, (from *æger*), Disease—æ. Ventriculi, Vomiting.

ÆGROTAT, (L.) 'he is sick,' (*ægroto*, 'I am sick.') A medical certificate in the English colleges, to show that the student is unfit for duty.

ÆGROTATIO, (from *æger*), Disease.

ÆGROTUS, (from *æger*), Sick.

ÆGYPTIA, (*Αιγυπτια*.) An epithet for several medicines, mentioned by Galen, Paulus of Ægina, and Myrepsus.

ÆGYPTIA MOSCHATA, Hibiscus abelmoschus.

ÆGYPTIA STYPTERIA, *Αιγυπτια στυπτηρία*, *Ægyptian alum*. Recommended by Hippocrates.

ÆGYPTIA UL'CERA; *Ægypt'ion ul'cers*. Ulcers of the fauces and tonsils, described by Aretæus, as common in Egypt and Syria.

ÆGYPTIACUM, *Ægypt'ion*, *Mende'sion*, *Mel Ægypti'acum*, *Phar'macum Ægypti'acum*. A preparation of vinegar, honey, and verdigris, scarcely used now, except by veterinary surgeons as a detergent. See Linimentum Æuginis.

ÆGYPTION, (*Αιγυπτιον*), Ægypticum.

ÆGYPTIUM MEDICAMENTUM AD AURES, Pharmacum ad aures.

ÆGYPTIUS PES'SUS: *Ægypt'ian pessary*. A pessary, composed of honey, turpentine, butter, oil of lily or of rose, and saffron, each one part; with sometimes a small quantity of verdigris.

ÆICHRYSON, Sedum.

ÆIPATHEIA, (*αι*, 'ever,' and *παθος*, 'disease,') see Continent (Disease.)

ÆIPATHIA, see Continent (Disease.)

ÆMOPTOICA PASSIO, (*ἀίμα*, 'blood,' and *πτωω*, 'to spit,') Hæmoptysis.

ÆNEA, (*æneus*, 'of brass,') Catheter.

ÆOLECTHYMA, (*αιολλω*, 'to agitate,') 'to variegate' (?), and *ecthyma*.) Variola.

ÆOLLION, Varicella.

ÆOLIUM, Varicella.

ÆON, (*αιων*.) The entire age of a man from birth till death. Also, the spinal marrow. See Medulla Spinalis.

ÆONESIS, (*αιοναω*, 'to moisten,') Fomentation.

ÆONION, Sedum.

ÆO'RA, (*αιωρα*, from *αιωρω*, 'I suspend,') Gestation, swinging.

ÆQUALIS, Equal.

ÆQUA'TOR OC'ULI. The line formed by the union of the upper and under eyelid, when they are closed. It is below the middle of the globe.

ÆQUIVOCUS, Equivocal.

AER, gen. *A'eris*, (*αερ*), Air.

ÆÉRAGE, Ventilation.

ÆÉRATION, Ventilation.

AERATION OF THE BLOOD, Hæmatosis.

AERATUS, Carbonated.

ÆÉRÉ, Carbonated.

ÆERENDOCAR'DIA, (F.) *Æerendocardie*, (*aer*, and *endocardium*.) Presence of air in the endocardium or lining membrane of the heart.

ÆERENTERCTASIA, (*aer*, *εντερον*, 'intestine,' and *εκτασις*, 'dilatation,') Tympanites.

ÆER'OLUM, *Ære'olus*, *Chal'cus*, (from *αε*, *aeris*, 'brass,') The sixth part of an obolus by weight, consequently about 2 grains.

Æ'RESIS, (*αιρεσις*, 'the removal of any thing,') A suffix denoting a removal or separation, as *Aphæ'resia*, *Diæ'resis*, &c.

ÆÉRETHMIE, (*aer*, and *ηθμος*, 'a sieve,') Emphysema.

ÆÉRETHMOPNEUMONIE, see Emphysema of the Lungs.

ÆERGIA, (*αεργια*, from *a*, and *εργον*, 'work,') Torpor.

ÆERHÆMATOX'IA, (F.) *Æérhêmatoxie*; (*aer*, *ἀίμα*, 'blood,' and *τοξικον*, 'poison,') Poisoning by the reception of air into the blood-vessels.

ÆÉRIF'EROUS, *Æerifer*, (F.) *Æerifère*, (*ac*, and *ferre*, 'to carry,') An epithet for tubes which convey air, as the larynx, trachea, and bronchia.

ÆÉRIFLUX'US, (*aer*, and *fluere*, *fluxum*, 'to flow,') The discharge of gas, and the fetid emanations from the sick. Flatulence.

ÆRODERMECTASIA, (*aer*, δερμα, 'skin,' and *ectasis*, 'dilatation.') Emphysema.

ÆRODIAPHTHORA, (*aer*, and διάφθορα, 'corruption.') A corrupt state of the air.

ÆRO-ENTERECTASIA, Tympanites.

ÆROHYDROPATHY, *Ærohydropathia*, (*aer*, ὕδωρ, 'water,' and παθος, 'affection.') A mode of treating disease, in which air and water are the chief agents employed.

ÆROL'OGY; *Ærolog'ia*, *Ærolog'icē*, (*aer*, and λογος, 'a description.') That part of physics which treats of the air, its qualities, uses, and action on the animal economy.

ÆROMAN'CY, *Æromanti'a*, (*aer*, and μαντεία, 'divination.') An art in judicial astrology, which consists in the foretelling, by means of the air, or substances found in the atmosphere.

ÆROMELL, (*aer*, and μέλι, 'honey,') Fraxinus ornus.

ÆROPÉRITONIE, see Tympanites.

ÆROPHOB'IA, (*aer*, and φοβος, 'fear.' Dread of the air.) This symptom often accompanies hydrophobia, and sometimes hysteria and other affections.

ÆROPHOB'ICUS, *Æroph'obus*; same etymon. One affected with aerophobia.

ÆROPHOBUS, Aerophobicus.

ÆROPTHORA, (*aer*, and φθορα, 'corruption,') Aerodiaphthora.

ÆROPLEURIE, (*aer*, and πλευρον, 'the side,') Pneumothorax.

ÆROPNEUMONASIE, (*aer*, and πνευμων, 'the lung,') see Emphysema of the Lungs.

ÆROPNEUMONECTASIE, see Emphysema of the Lungs.

ÆRORA'CHIA, (*aer*, and ραχis, 'spine,') Accumulation of air in the spine.

ÆROSIS, (from *aer*,) Pneumatosis, Tympanites.

ÆROTHORAX, Pneumothorax.

ÆRUCIA, Cupri subacetata.

ÆRUG'INOUS, *Ærugine'us*, *Io'des*, (*ærugo*, 'verdigris,') (F.) *Ærugineux*. Resembling verdigris in colour; as the bile when discharged at times from the stomach.

ÆRUGO, gen. *Æruginis*, *ios*, (from *æs*, *æris*, 'copper,') The rust of any metal, properly of brass. See Cupri Subacetata.

ÆRUGO FERRI, Ferri subcarbonas—æ. Plumbi, Plumbi subcarbonas.

ÆS, see Cuprum.

ÆSCHOS, (*αἰσχος*,) Deformity of the body generally, or of some part.

ÆSCULUS HIPPOCAS'TANUM, (*esca*, 'food,') [?] *Castanea equina*, *Pavia*, *Horse-chestnut*, *Buckeye*. (F.) *Marronnier* or *Marronnier d'Inde*. Ord. Hippocastaneæ. Sez. Syst. Heptandria Monogynia. The bark has been advised as a substitute for cinchona. Both bark and fruit are astringent. Externally, it has been employed, like cinchona, in gangrene.

ÆSECAVUM, (from *æs*,) Brass.

ÆSTATES, (from *æstas*, gen. *æstatis*, 'summer,') Ephelides.

ÆSTHE'MA, gen. *Æsthe'matis*, (*αἰσθημα*, gen. *αἰσθηματος*, 'a sensation, a perception,') See Sensation and Sensibility. In the plural, *Æsthe'mata*, the apparatuses of the senses.

ÆSTHEMATOL'OGY, *Æsthematolog'ia*; (*æsthemata*, and λογος, 'a discourse,') The doctrine of, or a treatise of, the senses, or on the apparatus of the senses.

ÆSTHEMATONU'SI, *Æsthematorganonu'si*, (*æsthemata*, and νοσος, 'diseases,') Diseases affecting sensation.

ÆSTHEMATORGANUSI, *Æsthematonusi*.

ÆSTHESIOGRAPHY, *Æsthesiogra'phia*,

(*æsthesia*, and γραφή, 'a description,') A description of the sensorial apparatuses.

ÆSTHESIOLOG'Y, *Æsthesiolog'ia*, (*æsthesia*, and λογος, 'a discourse,') The doctrine of, or a treatise on, sensation or the senses.

ÆSTHESIOM'ETER, (*æsthesia*, and μετρον, 'measure,') A measurer of sensation. An instrument of the beam-compass kind, contrived by Dr. Sieveking to indicate the distance at which two points can be distinguished, when brought in contact with any portion of the cutaneous surface.

ÆSTHESIS, (*αἰσθησις*,) *Aisthe'sis*, (*αἰσθανομαι*, 'I feel,') The faculty of being affected by a sensation. Perception. Sensibility, as well as the senses themselves. See Sense.

ÆSTHESOD'IC, *Æsthesod'icus*, (*æsthesia*, and ὁδος, 'a way,') Having the faculty of conveying sensation.

ÆSTHETERION, (*αἰσθητηριον*,) (from *æsthesia*,) Sensorium.

ÆSTHETICA, (same etymon.) Diseases affecting the sensations. Dulness, deprivation or abolition of one or more of the external organs of sense. The 2d order, class *Neurotica*, of Good. Also, agents that effect sensation.

ÆSTIVUS, (*æstas*, 'summer,') Estival.

ÆSTUARIUM, (from *æstus*,) Stove.

ÆSTUATIO, Ardor, Ebullition, Fermentation.

ÆSTUS, ('heat,') Ardor.

ÆSTUS VOLAT'ICUS. Sudden heat, scorching or flushing of the face.

ÆTAS, gen. *Ætatis*, Age—æ. Bona, Adolescence—æ. Decrepita, Decrepitude—æ. Mala, Senectus—æ. Provecta, Senectus—æ. Senilis, Senectus—æ. Virilis, Adult age.

ÆTHER, gen. *Ætheris*, *Æther*, (from αἰθήρ, 'the subtle air of the upper regions,' or from αἶθερ, 'I burn,') *Naph'tha*. *Liq'or æthereus*. A volatile liquor obtained by distillation from a mixture of alcohol and a concentrated acid. See *Æther sulphuricus*, and *Ether*.

ÆTHER ACETICUS, see *Æther Sulphuricus—æ. Anæstheticus, see *Æther Muriatricus*—æ. Chloratus alcoholicus, *Spiritus Ætheris muriatrici*—æ. Chloricus, Chloroform, *Ether*, chloric—Fortior, see *Æther sulphuricus*.*

ÆTHER HYDRID'ICUS, *Hydriod'ic* or *Iodhyd'ric* *Æther*, *Iodide of Ethyle*. (F.) *Æther iodhydrique*: obtained by mixing alcohol, parts v.; iodine, parts x.; and phosphorus, one part, and distilling. It is used for iodine inhalation.

ÆTHER HYDROCHLORICUS, *Æther muriatricus*.

ÆTHER HYDROCYAN'ICUS, *Æ. Prus'sicus*, *Hydrocyan'ic*, or *Prus'sic* *Æther*, *Hydrocyan'ate of Eth'erine*, *Cyan'uret of Eth'yle*, (F.) *Æther Hydrocyanique*, has been advised in whooping-cough, and where the hydrocyanic acid is indicated. Dose, 6 drops.

ÆTHER LIGNOSUS, Acetone—æ. Martialis, *Tinctura seu Alcohol sulfurico-æthereus ferri*.

ÆTHER MURIAT'ICUS, *Æ. Hydrochlor'icus*, *Muriat'ic*, *Chlorohyd'ric* or *Marine* *Æther*, *Muriate of Eth'erine*, *Chloride of Eth'yle*. This ether, on account of its volatility, can only be kept in cool places. It has the properties of the other ethers, and when used is generally mixed with an equal bulk of alcohol. It has been employed as an anæsthetic.

A *Chlorinated Chlorohyd'ric* *Æther*, *Æther muriat'icus trichloratus*, *Æther anæsthet'icus*, (F.) *Æther Chlorhydrique chloré*, formed by the action of Chlorine on Chlorohydric *Æther*, has been introduced into practice as a local anæsthetic.

ÆTHER MURIATICUS ALCOHOLICUS, *Spiritus ætheris muriatricus*—æ. *Muriaticus trichloratus*, see *Æther muriatricus*—æ. *Nitrico-alcoholicus*, *Spiritus ætheris nitrici*—æ. *Nitricus*, see *Æther sulphuricus*—æ. *Nitricus Alcoolisatus*, *Spiritus ætheris nitrici*—æ. *Nitrosus*, see *Æther Sulphuricus*—æ. *Pyro-aceticus*, Acetone.

ETHER SULPHURICUS, *Æther* (Ph. U. S.), *Æ. Vitriolicus*, *Naphtha Vitrioli seu Sulphurica*, *Oleum vitriolæ dulcis*, *Sulphuric Ether*. (F.) *Éther, Éther sulfurique, vitriolique ou hydratée*. Ether prepared from sulphuric acid and alcohol.

Rectified Ether, Ether rectificatus, prepared by distilling 12 oz. from a mixture of sulphuric ether, f³xiv, fused potass, f³ss. and distilled water, f³ij, is a limpid, colourless, very inflammable volatile liquor; of a penetrating and fragrant odour, and hot pungent taste. Its s. g. is 0.752.

Æther Sulphuricus, Sulphuric Ether of the Pharmacopœia of the United States (1842), *Æther* of that of 1851 and of 1863, is formed from stronger alcohol, Ovi; sulphuric acid, f³xxxvi; potassa, gr. coelx; distilled water, f³ij; distilling and redistilling according to the process there laid down. The specific gravity of this ether is 0.758.

ETHER FORTIOR, Stronger Ether (Ph. U. S.), is made by agitating ether with chloride of calcium, lime, and an equal quantity of water; decanting the ether after standing, and distilling off half the amount of the ether used. Its s. g. is 0.728. Used for inhalation as an anæsthetic.

Ether is a diffusible stimulant, narcotic and antispasmodic, and is, externally, refrigerant. Dose, grt. xxx to f³ss. When it is inhaled, it is found to be a valuable anæsthetic agent; and is employed with advantage in spasmodic affections, and in surgical operations. See Anæsthetic.

The Parisian Codex had an *Æther aceticus*, *Naphtha Aceti*, (F.) *Éther acétique*; an *Æther muriaticus seu hydrochloricus*, (F.) *Éther chlorhydrique*; an *Æther nitricus seu nitrosus* (F.) *Éther nitrique ou nitreux ou azoteux*; and an *Æther phosphoratus*. They all possess similar virtues. See Anæsthetic.

ETHER SULPHURICUS ACIDUS, *Elixir acidum Halleri*—*æ. Sulphuricus cum alcohole*, *Spiritus ætheris sulphurici*—*æ. Sulphuricus cum alcohole aromaticis*, *Spiritus ætheris aromaticus*.

ETHER TEREBINTHINATUS, *Terebinthinated ether*, made by mixing gradually two pounds of alcohol, and half a pound of spirit of turpentine, with two pounds of concentrated nitric acid, and distilling one-half the mixture with a gentle heat, is employed externally and internally in biliary calculi, rheumatism, &c. Dose, 20 to 40 drops, in honey or yolk of egg.

ETHEREA HERBA, *Eryngium maritimum*.

ÆTHEREAL, *Æthereus*, *Æthereal*, *Æthereus*, (F.) *Éthérée*, (from *æther*.) An ethereal tincture, (F.) *Teinture éthérée*, is one formed by the action of sulphuric ether, at the ordinary temperature, on medicinal substances. An ethereal is a volatile oil. See Olea Volatilia.

ÆTHEREO-OLEOSA (REMEDIA), (from *æthereum*, 'a volatile oil.') Remedies, whose properties are dependent upon the volatile oil they contain.

ÆTHERISATIO, Etherization.

ÆTHERISATUS, Etherized.

ÆTHERISMUS, Etherism.

ÆTHEROLEA, Olea volatilia.

ÆTHIOPICUS LAPIS, *Æthiopian stone*. A stone formerly supposed to be possessed of considerable virtue.

ÆTHIOPIFICATIO, *Æthiopoiosis*, *Æthiopiæmus*, *Æthiopiæsis*, (*Æthiops*, and *facere*, 'to make.') The mummy-like colouring of the skin, induced at times by the use of mercurial ointment; and seen in bodies poisoned by arsenic.

ÆTHIOPIOSIS, *Æthiopicatio*.

ÆTHIOPIS, *Salvia selarea*.

ÆTHIOPISMUS, *Æthiopicatio*.

ÆTHIOPOPOESIS, (*ποιω*, 'I make,') *Æthiopicatio*.

ÆTHIOPS, gen. *Æthi'opis*, (*αιθω*, 'I burn,' and *ωψ*, 'countenance.') A black or burnt countenance. The ancients gave this name to certain oxides and sulphurets of metals, which were of a black colour.

ÆTHIOPS ALBUS, Albino—*æ. Alcalisatus*, *Hydrargyrum cum cretâ*—*a. Animal*, see Choroid.

ÆTHIOPS MARTIALIS, *Martial Etheriops*, *Ferri Deutoxydum nigrum*. The black deutoxide of iron: once in repute as a tonic. See Oxydum Ferri nigrum.

ÆTHIOPS MINERALIS, *Hydrargyri sulphuretum nigrum*—*æ. Narcoticus*, *Hydrargyri sulphuretum nigrum*—*æ. Per se*, *Hydrargyri oxydum cinereum*—*æ. Saccharatus*, *Hydrargyrum saccharatum*—*æ. Vegetabilis*, see *Fucus vesiculosus*.

ÆTHOLICES, (*αιθολικες*, pl. of *αιθολιξ*, from *αιθω*, 'to burn,' and *ηλιος*, 'the sun.') Fiery pustules on the skin. Some have considered them to have been boils.

ÆTHUSA, (from *αιθω*, 'to burn,') *Ammi*, *Sison ammi*.

ÆTHUSA CYNAPIUM, *Fool's Parsley*, (F.) *Faux Persil*, *Petit Ciguë*. Family, Umbelliferae. Sex. Syst. Pentandria Digynia. A poisonous plant, which has been mistaken for true parsley, producing nausea, vomiting, headache, giddiness, sopor, and at times, fatal results. It resembles conium in its action.

ÆTHUSA MEUM, *Meum*, *M. Athaman'ticum*, seu *Anethog'lium*, *Athaman'ta Meum*, *Ligusticum Capilla'ceum* seu *Meum*, *Ses'eli Meum*, *Me'u*, *Radix ursi'na*, *Spignel*, *Baldmoney*, (F.) *Ethuse*, *Méum*. The root has been advised as carminative, stomachic, &c.

ÆTHIOI PHLEBES, (*αιθιοι φλεβες*, 'eagle veins,') Temporal veins.

ÆTIOLOG'IA, *Ætiology*, *Etiology*, *Aitiolog'ia*, (*aitia*, 'cause,' and *λογος*, 'a discourse.') The doctrine of the causes of disease.

ÆTITES, or **ÆTTITES**, (*αιτες*, 'an eagle,') *Eaglestone*, (F.) *Pierre d'Aigle*, *Hydrate de trioxide de fer*. This stone was formerly supposed to facilitate delivery, if bound on the thigh; and to prevent abortion, if bound on the arm. It was also called *La'pis Coll'ymus*.

ÆTOLION, *Cnidia grana*.

Æ, as a prefix, see **Ad**.

AFFADIL, *Narcissus pseudo-narcissus*.

AFFADISSEMENT, (F.) (*fade*, 'insipid,') That condition of the digestive function in which the appetite is diminished, the sense of taste blunted, and the action of the stomach enfeebled; a state usually accompanied by general languor.

AFFAIBLISSEMENT, (from *faible*, 'weak,') Asthenia.

AFFAIRES, *Menses*.

AFFAISSEMENT, (*af*, and *faire*, 'weight,') Collapse.

AFFECT, see **Affection**.

AFFECTIO, gen. *Affectio'nis*, (*officio* or *affectare*, (*ad* and *facere*), 'to move or influence,')

Affection—*a. Arthritica Cordis*, *Cardiagra*—*a. Hypochondriaca*, *Hypochondriasis*—*a. Hysterica*, *Hysteria*—*a. Sarmatica*, *Plica*—*a. Tympanitica*, *Tympanites*.

AFFECTION, *Affectio*. Any mode in which the mind or body is affected or modified.

AFFECTION TYPHOÏDE, see Typhus—*a. Vaporeuse*, *Hypochondriasis*.

AFFECTIONES ANIMI, Affections of the mind.

AFFECTIONS DE L'ÂME. Affections of the mind.

AFFECTIONS OF THE MIND, *Affectus seu Passio'nes seu Affectio'nes seu Conquassatio'nes seu Confusio'nes seu Turbatio'nes seu Perturbatio'nes an'imi*, (F.) *Affections de l'âme* include not only the different passions, as love, hatred, jea-

lousy, &c., but every condition of the mind that is accompanied by an agreeable or disagreeable feeling, as pleasure, fear, sorrow, &c.

In Pathology, *Affection*, *Affect*, *Pa'thos*, *Pathe'ma*, is synonymous with disease: thus we speak of a *pulmonary affection*, a *calculus affection*, &c.

AFFECTIVE. That which affects, touches, &c. Gall gives the term *affective faculties*, (F.) *Facultés affectives*, to functions dependent upon the organization of the brain, comprising the sentiments, affections, &c.

AFFECTUS, gen. *Affectûs*, Passion—a. Fau-cieum pestilens, *Cynanche maligna*—a. Hydr-odes. **Hydrops**—a. Spasmodico-convulsivus la-biorum, *Neuralgia faciei*.

AFFERENT, *Afferens*, *Ad'vehens*, *Centrip'e-tal*, *Erod'ic*, (from *affero*, (ad and *fero*, 'to carry,') 'I bring,') Conveying inwards, as from the peri-phery to the centre. The vessels which convey the lymph or chyle to the lymphatic glands, are called *afferent* or *inferent*, *vasa afferen'tia* seu *in-feren'tia*. Also, nerves that convey impressions towards the nervous centres—*nervi entobanun'tes*.

AFFION, *Aff'ium*, *Of'fium*. Opium. See Pa-paver. The Bantames thus designate an elec-tuary of which opium is the basis, and which they use as an excitant. Mr. Auberger calls the indigenous opium, prepared by him in France, *Affium*.

AFFLATUS, gen. *Afflatûs*, *Adfla'tus*, *Epi-pno'ta*, (af, and *flave*, 'to blow,') Any air that strikes the body and produces disease.

AFFLUENCE, *Aff'lux*, (from *affluere*, (af, and *fluere*, 'to flow,') 'to flow to,') A flow or deter-mination of humours, and particularly of blood, towards any part.

AFFLUXUS, gen. *Affluxûs*, Fluxion.

AFFRODILE, *Narcissus pseudo-narcissus*.

AFFRONTER, (F.) (af, and *frons*, gen. *frontis*, 'front,') To bring together. To approximate the edges of a wound as close as possible. The action is termed *affrontement*.

AFFUSIO, gen. *Affusio'nis*, (ad, and *fundere*, *fusum*, 'to pour,') Affusion—a. Frigida, see Af-fusion—a. Orbicularis, Placenta.

AFFUSION, *Affu'sio*, *Pro'schysis*, *Epich'y-sis*. The action of pouring a liquid on any body. *Af-fusions*, *Rhyptolu'siæ*, cold and warm, are used in different diseases. The cold affusion, *Affu'sio* seu *Perfu'sio frig'ida*, is said to have been ben-eficial in cutting short typhus fever and scarlatina, if used during the first days. It consists in plac-ing the patient in a tub, and pouring cold water over him; then wiping him dry, and putting him to bed. The only precaution necessary is, to use it in the state of greatest heat and exacerbation; not when chilliness, or topical inflammation, is present.

AFIUM, Opium.

AFTER-BIRTH. Secundines.

AFTER-BURTHEN, Secundines.

AFTER-DAMP, see Hydrogen, carburetted.

AFTER-PAINS, see Pains, labour.

AG, in composition, see Ad.

AGACEMENT, (F.) (from *aca'zein*, 'to sharpen,') The setting on edge.

AGACEMENT DES DENTS. A disagree-able sensation experienced when acids are placed in contact with the teeth. *Tooth-edge*. Setting the teeth on edge.

AGACEMENT DES NERFS. A slight irri-tation of the system, and particularly of the organs of sense and locomotion, corresponding nearly to the English *Fid'gets*. The painful rest-lessness of the lower extremities, occasionally witnessed, in nervous affections more especially, has been termed *anri'etas tibia'rum*.

AGAH, Intermittent Fever.

AGALACTATIO. Agalactia.

AGALACTIA, *Agalac'tia*, *Agalac'tio*, *Agalac-*

ta'tio, *Defec'tus lac'tis*, *Oligoga'llia*, *Oligogalac'tia*, (a, and *gala*, gen. *γαλακτος*, 'milk,') Absence of milk in the mamme.

AGALAXIA, Agalactia.

AGALLOCHUM, (αγαλλοχον, from αγαλλω, 'I decorate,') *Calambac*, *Calambouk*, *Lig'nium Agal-lochi ve'ri* seu *Al'oë* seu *Aspal'athi*, *Xyloölöë*, *Al'oëswood*. A resinous and very aromatic wood of the East Indies, from *Excoeca'ria Agal'lochu*, *Cynome'tra Agal'lochum*, *Alöë'yilon Agal'lochum*. Used in making pastiles, &c., according to Dios-corides, Oribasius, and Paulus.

AGAMOUS, (a, and *γαμω*, 'to marry,') See Cryptogamous.

AGARIC, *Agar'icum*, *Agar'icus*. A genus of plants in the Linnaean system, some of which are edible, others poisonous. (It is supposed to have been called *αγαρικον* from Agaria, a town and region of Sarmatia,)—Dioscorides. Among the edible varieties of the Boletus, the following are the chief:—1. The *Agar'icus edu'lis* seu *Arven'sis* seu *Sylva'ticus* seu *Campe'stris*, (F.) *Agaric comestible et champignon de couche*. 2. The *Agar'icus odora'tus*, (F.) *Mousseron*. The most common poisonous varieties are the *Agar'icus nece'tor*, (F.) *Agaric meurtrier*; and, 2. The *Agar'icus a'eris*, (F.) *Agaric äcre*; besides the *Auronite*, a sub-genus, which includes several species. One of the most delicate is the *Agar'icus Auranti'acus*, but care must be taken not to confound it with the *A. Pseudoaurantiacus*, which is very poisonous. The *A. aurantiacus* is called, in French, *Orange*.

AGARIC, see Boletus ignarius—a. *Amadouvier*, Boletus laricis—a. *Blanc*, Boletus laricis—a. *de Chêne*, Boletus ignarius—a. Female, Boletus ignarius—a. of the Oak, Boletus ignarius—a. *Odorant*, *Dædalea suaveolens*—a. White, Boletus laricis.

AGARICUM, (αγαρικον,) Boletus ignarius.

AGARICUS, Boletus ignarius—a. *Albus*, Boletus laricis—a. *Arvensis*, see Agaric—a. *Auran-tiacus*, see *Amanitæ*, *Bolites*—a. *Auriculæforma*, *Peziza auricula*—a. *Campestris*, see Agaric—a. *Chirurgorum*, Boletus ignarius—a. *Ignarius*, Boletus ignarius—a. *Laricis*, Boletus laricis—a. *Pseudo-aurantiacus*, *Amanitæ*—a. *Quercus*, Boletus ignarius—a. *Sylvaticus*, see Agaric.

AGARIFIED, See Feverish.

AGASTRONERVIA, (F.) *Agastronervie*, (a, *γαστηρ*, 'stomach,' and *νευρον*, 'nerve,') Want of nervous action in the stomach.

AGASYLLIS GALBANUM, (αγασυλλis,) Bu-bon galbanum.

AGATHIS DAMARRA, (from *αγadis*, 'a clew of thread,') *Pinus damarra*.

AGATHOSMA CRENATA, (αγαθος, 'good,' and *σμη*, 'odour,') *Diosma crenata*.

AGATHOTES CHIRAYITA, (αγαθους, 'good-ness,') *Gentiana Chirayta*.

AGA'VE AMERICA'NA, (corr. from *ayavos*, 'admirable,') *A. Ramo'sa*, *Americ'an Ag'ave*, *Americ'an al'oe*, *Maquay*. *Nat. Ord.* Bromeliaceæ. *Sex. Syst.* Hexandria Monogynia. This plant has been considered diuretic and antispyllitic. The favourite drink of the Mexicans—*Pulque*—is the fermented juice of the plant.

AGAVE RAMOSA, *A. Americana*.

AGAVE VIRGINICA, *Rattlesnake's mas'ter*, *False al'oe*,—grows in the Southern States. The root is very bitter. It has been used in tincture as a carminative in colic; and as a remedy for bites of serpents.

AGE, ἡλικία, *Heli'kia*, *Æ'tas*; (of uncertain etymon.) Period of life. Time that has elapsed since birth, &c. Five ages are often designated in the life of man: 1. First infancy (*Infan'tia*,) 2. Second infancy (*Pueri'tia*,) 3. Adolescence (*Adolescen'tia*,) 4. The adult or virile age (*Vi-*

virilitas, *Ætas virilis*, (F.) Virilité; 5. Old age, (Old Eng. *Elth*, *Senectus*.)

AGE DE RETOUR, (F.) ('age of return.') The time of life when the vital powers begin to be diminished, as at the approach of old age.

AGENEIOS, (ἀγενεῖος, from *a*, and γενεῖον, 'beard.') Imberbis.

AGENESIS, (α and γενεσις, 'generation.') (F.) *Agénésie*. Imperfect development of any part of the body; as *cerebral agenesi*s, i. e. imperfect development of the brain in the fœtus. See *Monster*.

AGENNÉSIA, (ἀγεννησία,) Impotence, Sterilitas.

AGENNESIS, Impotence, Sterilitas.

AGENOSOMUS, (α, γεννω, 'I generate,' and σωμα, 'body.') A malformation in which the fissure and eversion are chiefly in the lower part of the abdomen; the urinary or sexual apparatus absent, or very rudimentary.

AGENT, *Agens*, (agere, 'to act.') Any power which produces, or tends to produce, an effect on the human body. *Morbific agents*, (F.) *Agens morbosiques*, are the causes of disease;—therapeutical agents, (F.) *Agens thérapeutiques*, the means of treating it.

AGER NATURE, ('field of nature,') Uterus.

AGERASIA, *Isenescentia*, (α, and γηρας, 'old age,') A vigorous and green old age.

AGERATUM, (ἀγερτον,) Achillea ageratum—a. *Altissimum*, *Eupatorium ageratoides*.

AGERATUS LAPIS, (ἀγερτος λίθος.) A stone used by cobblers to polish shoes. It was formerly esteemed diuretic and astringent.

AGES, Palm.

AGEUSIA, (α, and γευσις, 'taste,') Ageusia.

AGEUSTIA, (ἀγευστία,) *Agheus'tia*, *Ageusia*, *Apogeus'tia*, *Apogeus'is*, *Dysæsthesia gustatoria*, *Porogeus'is*. Diminution or loss of taste, *Anæsthesia lingue*.

AGGLÖMERATE, *Agglomeratus*, (from *agglomerare*, (ag, and *glomerare*, 'to wind up yarn in a ball,') 'to collect together,') Applied to tumours or glands in aggregation.

AGGLUTINANT, *Agglutinans*, *Collet'icus*, *Glutinans*, (ag, and gluten, 'glue,') (F.) *Agglutinant*, *Agglutinatif*, *Glutinatif*, *Collétique*. Remedies were formerly so called, which were considered capable of uniting divided parts.

Plasters are called *agglutinants*, (F.) *agglutinatifs*, which adhere strongly to the skin. Certain bandages are likewise so termed; (F.) *Ban-dettes agglutinatives*.

AGGLUTINATE, *Agglutina'rē*. The French use the word *agglutiner* in the sense of 'to reunite; as *agglutiner les lèvres d'une plaie*, 'to reunite the lips of a wound.'

AGGLUTINATIF, *Agglutinant*.

AGGLUTINATIO, *Agglutination*, *Coition*.

AGGLUTINATION, *Agglutina'tio*, *Colle'sis*, *Epicolle'sis*, *Proscolle'sis*, *Glutina'tio*. The first degree of adhesion. Also, the action of agglutinants.

AGGLUTINER, To agglutinate.

AGGREGATE, *Aggregatus*, (from *aggregare*, (ag, and *gregare*, 'to flock together,') 'to assemble together,') Glands are called *aggregate* which are in clusters. See *Peyer's Glandulæ*. *Aggregate pills*, (F.) *Pilules agréatives*, signified, formerly, those which were believed to contain the properties of a considerable number of medicines, and to be able to supply their place.

AGHEUSTIA, *Ageusia*.

AGHOUL, *Agul*.

AGIAHALID or **AGIAHALID** or **AGRAHALID**. An Egyptian and Ethiopian shrub, similar to *Xime'na*. The Ethiopians use it as a vermifuge. The fruit is purgative.

AGIAHALID, *Agiahalid*.

AGISSANT, (*agir*, 'to act,') Active.

AGITATION, *Agita'tio*, *Done'sis*, (*agere* 'to act,') Constant and fatiguing motion of the

body, *Ty'r'bē*, *Ty-ba'sia*, *In'quies*,—or distressing mental inquietude,—*An'imi Agita'tio*.

AGITATORIUS, Convulsive.

AGLITHES, see *Allium*.

AGLOBULIE, See *Anæmia*.

AGLOSSIA, (αγλωσσία, from α, and γλωσσα, 'the tongue,') *Aglost'ia*. A malformation, which consists in the want of a tongue.

AGLOSSOS'TOMA, (*aglossia*, and *στομα*, 'mouth,') A mouth without a tongue.

AGLOSSOSTOMOGRAPHY, (*Aglossostomographia*, (α, γλωσσα, 'the tongue,') *στομα*, 'the mouth,') and *γραφω*, 'I describe,') Description of a mouth without a tongue.

AGLOTTIA, (αγλωττία,) *Aglossia*.

AGLOTT'ION, *Aglutit'io*, (α, and *glutire*, 'to swallow,') A hybrid term, designating impossibility of swallowing.

AGMA, (*agma*, gen. *αγματος*,) Fracture.

AGMATOLOG'IA, (*agma*, and *λογος*, 'a description,') The doctrine of fractures. A treatise on fractures.

AGME, Fracture.

AGMINA DIGITORUM MANUS, (*agmen*, *agminis*, 'a phalanx,') Phalanges of the fingers.—a. *Membrana Amnios*.

AGMINATED GLANDS, *Peyer's glands*.

AGNAIL, Hangnail.

AGNA'THIA, (α, and γναθος, 'jaw,') A malformation, which consists in the want of the jaw, especially of the lower.

AGNINA MEMBRANA, (*agnus*, 'a lamb,') *Amnios*.

AGNOIA, (*agnoia*, from α, and γινωσκω, 'I know,') *Agno'e*. State of a patient who does not recognise individuals.

AGNUS CASTUS, (αγνος,) *Vitex*.

AGO'GĒ, (*agōgē*, from αγω, 'I lead,') or 'drive,') The order or condition of a disease. Likewise the state of the air.

AGOGUE, (*αγωγος*, 'a leader,') A suffix. Hence *Chologogue*, a leader or expeller of bile; *Hydragogue*, &c.

AGOMPHIASIS, *Agompho'sis*, (α, and γομφίον, 'I nail,') Looseness of the teeth. See *Gomphiasis*.

AGOMPHOSIS, *Agomphiasis*.

AGON, (*agōn*, 'a combat,') *Agony*.

AGONE, (from *agonos*; being supposed to cause sterility,) *Hyoseyamus*.

AGONIA, (*agonia*,) *Agony*, (*αγωνία*,) *Sterilitas*.

AGONISMA, gen. *Agonism'atis*, *Agony*.

AGONISMUS, *Agony*.

AGONIS'TICA, (*agōn*, 'a combat,') The part of ancient gymnastics, which had reference to the combats of the *Athletæ*.

Also, very cold water, given internally to calm febrile heat.

AGONIZANS, *Psychorages*.

AGONOS, (*agōnos*, from α, and γωνος, 'seed,') Sterile.

AG'ONY, *Agon'ia*, *A'gon*, *Agonis'ma*, *Agonis-mus*, *Moch'tus*, *Mo'gus*, *Psychorag'ia*, *Psychorag'h'ia*, *An'gor*, *Throe*, *Throv*, (Sc.) *Deadthraw*, *Patient of death*; (*agōn*, 'a combat,') The last struggle of life. The agony, which is of longer or shorter duration, is characterized by great change in the features, gradual abolition of sensation and motion, loss of voice, dryness or lividity of the tongue and lips, rattling in the throat, small and intermittent pulse, and coldness of the extremities. This state is not present in those who die suddenly. See *Facies Hippocratica*.

AGOS'TUS, (*agōstos*, from αγω, 'I lead,') The fore arm from the elbow to the fingers. Also, the palm of the hand. See *Palm*.

AGRA, (*agra*, from αγρω, 'I seize hold of,') A seizure, as *Odontagra*, a tooth seizure, toothache; *Chiragra*, *Podagra*, &c.

AGRAFE DE VALENTIN. A kind of forceps with parallel branches, employed by Valen-

tin in the operation for hare lip, to adect the approximation of the edges of the wound.

AGRAHALID, Agiabalid.

AGREGATIVES PILULES. See Aggregato.

AGRÉMIE, (*agra*, and *αἷμα*, 'blood.') The altered condition of the blood in gout.

AGRIA, (*αγριος*, 'wild.') Herpes exedens, Lichen agrius.

AGRIAMPELOS, (*αγριαμπελος*, from *αγριος*, 'wild,' and *αμπελος*, 'vine.') Bryonia alba.

AGRIELCO'SIS, (*αγριος*, 'wild,' and *ἔλκος*, 'ulceration,') A malignant ulceration.

AGRIE, (from *αγρα*, 'a seizure,') Gout.

AGRIFOLIUM, *Ilex aquifolium*.

AGRIMONIA, (corrupted from Argemone.[?]) Agrimony—a. *Eupatoria*, Agrimony—a. *Odorata*, Agrimony—a. *Officinalis*, Agrimony.

AGRIMONIA PARVIFLORA, *Small-flowered* or *Sweet-scented Agrimony*; indigenous; has similar properties to A. *Eupatoria*. See Agrimony.

AGRIMONY, *Agrimo'nia*, A. *Eupatoria* seu *odorata* seu *officinalis*, *Caf'al*, *Lap'pula hepatica*, *Cuckebur*, *Stick'wort*. (F.) *Aigremoine*. Ord. Rosaceæ. *Sex. Syst.* Icosandria Digynia. A mild astringent and stomachic. *Dose*, in powder, from ʒj to ʒj.

AGRIMONY, HEMP, *Eupatorium cannabinum*—a. *Small-flowered*, *Agrimonia parviflora*—a. *Sweet-scented*, *Agrimonia parviflora*.

AGRIOCASTANUM, (*αγριος*, 'wild,' and *καστανον*, 'chestnut,') *Bunium bulbocastanum*, *Lycoperdon tuber*.

AGRIOCINARA, (*αγριος*, 'wild,' and *κιναρα*, 'artichoke,') *Sempervivum tectorum*.

AGRIOCOCIME'LEA, (*αγριος*, 'wild,' *κοκκος*, 'a berry,' and *μῆλον*, 'an apple,') *Prunus spinosa*.

AGRIORIGANUM, (*αγριorigανον*, from *αγριος*, 'wild,' and *origανον*, 'majoram,') *Origanum majorana*.

AGRIOSELINUM, (*αγριοσελινον*, from *αγριος*, 'wild,' and *σελινον*, 'parsley,') *Smyrnum olusatrum*.

AGRIOTHYM'IA, (*αγριος*, 'wild,' and *θυμος*, 'disposition,') Ferocious insanity.

AGRIPALMA GALLIS, (*αγριος*, 'wild,' and *palma*, 'palm,') *Leonurus cardiaca*.

AGRIP'PA, *Ægrip'pa*, (from *ager partus*, 'difficult birth:' or perhaps from *αγρα*, 'taking or seizure,' and *πους*, 'the foot,') This term has been given to those born by the feet. It is pretended that the family of Agrippa obtained their name from this circumstance. Parturition, where the feet present, is called *Agrip'pæ par'tus*, *Agrippi'nus par'tus*.

AGRIPPINUS PARTUS, see Agrippa.

AGRO DI CEDRO, see Citrus medica.

AGROPYRUM LÆVISSIMUM, (*αγρος*, 'a field,' and *πυρος*, 'wheat,') *Triticum repens*.

AGROSTEMMA GITHAGO, (gen. *Agrostem'matie*) (from *αγρος*, 'a field,' and *στεμμα*, 'a garland,') *Lychnis githago*.

AGROSTIS, (*αγρωστis*, 'grass,') from *αγρος*, 'a field,') *Bryonia alba*—a. *Vorticillata*, *Andropogon muricatus*.

AGRUNA, *Prunus spinosa*.

AGRYPNIA, (*αγρυπνια*, from *αγριος*, 'wild,' and *ἕπνος*, 'sleep,') Insomnia.

AGRYPNOCOMA, (*αγρυνus* and *coma*.) Coma vigil.

AGRYPNODES, (from *αγρυπνος*, 'sleepless,' and *odes*.) Characterized by sleeplessness, as *Fe'bris Agrypnode*, a fever accompanied with sleeplessness.

AGRYPNOTICUS, Anthypnotic.

AGRYPNUS, (*αγρυπνος*, from *αγριος*, 'wild,' and *ἕπνος*, 'sleep,') Sleepless, vigilant.

AQUA DE VERUGA, see Verugas.

AGUARDIENTE, (S.) ('hot water,') Brandy. See also Spirit.—a. *de Italia*, see Spirit.

A'GUE, (from Gothic *agis*, 'trembling,') (?) Intermittent fever; often used in the same sense as chill or rigor.

AGUE AND FEVER, Intermittent fever.

AGUE, BRASS-FOUNDERS'. A series of morbid phenomena, resembling ague, in those who are exposed to the fumes of deffragating zinc in brass founderies.

AGUE, BROW, Neuralgia frontalis.

AGUE CAKE, *Placen'ta febrilis* seu *sple'nica*, *Physco'nia sple'nica*, *Sple'nia Tu'mor*; (F.) *Gâteau fébrile*. A visceral obstruction—generally in the spleen—which follows agues, and is distinctly felt by external examination. To a greater or less degree, it is not uncommon.

AGUE, DEAD, see Fever, masked.

AGUE DROP, TASTELESS, Liquor arsenicalis.

AGUE, DUMB, see Fever, masked—a. Face, Neuralgia, facial.

AGUE FREE, *Laurus sassafras*.

AGUE, LEAPING, see Leaping ague—a. Quartan, Quartan—a. Tertian, Tertian fever—a. Third-day, Tertian.

AGUE TREE, *Laurus sassafras*—a. Weed, *Eupatorium perfoliatum*.

AGU'IOS, (a, and *γινω*, 'limb,') Mutilated or wanting limbs. Weak, feeble.

A'GUL, *Ag'houl*, *Alha'gi*, *Hedysa'rum alhagi*. A thorny shrub of Persia and Mesopotamia, which affords manna. The leaves are purgative.

AGYRIAS, (*αγυρις*, 'a collection,') Opacity of the crystalline.

AGYRTA, (*αγυρτης*, from *αγυρις*, 'a crowd,') Formerly, a stroller who pretended to supernatural powers. Subsequently, a quack or illiterate pretender.—See Charlatan.

AGYRTIA, (*αγυρτια*,) Charlatantry.

AH'ORA, (a, privative, and *ἔρα*, 'youth,') Tardy development of the organs: the opposite to *Hyperho'ra*.

AHOUI, Theretia ahoui.

AHUSAL, Orpiment.

AHPNIA, (a, and *ἕπνος*, 'sleep,') Insomnia.

AI, *Lai*, (F.) A Gascon name for a tumefaction of the fibro-synovial *contusions* of the tendons in general, but most frequently noticed in the course of the external radial tendons of the abductor longus, and extensor brevis pollicis. It is accompanied by a peculiar crepitation; (F.) *Ténalgie* or *Ténosynite crépitante*.—Velpéau.

AIAULT, Narcissus pseudo-narcissus.

AIDE (F.), *Ad'jutor min'ister*. An aid or assistant to a surgeon in his operations.

AIDOICI, Gennetici.

AIDOIOMANIA, (*αἰδοια*, 'genital organs,' and *mania*.) Nymphomania.

AIERSA, Iris Germanica.

AIGE, Ægias.

AIGLE, MIN'ERAL WA'TERS OF. Near the city of this name, in Normandy, is the chalybeate spring of St. Xantin, much used in the 16th and 17th centuries.

AIGLE, Ægias.

AIGRE, (*acer*, 'sharp,') Acidulous—a. *Voix*. See Acid.

AIGRELET, (from *acer*, 'sharp,') Acidulous.

AIGRETTE, (from *acer*, 'sharp,') *Rumex acetosa*, see *Thypha atifolia*.

AIGREMOINE, Agrimony.

AIGREURS, Acidities.

AIGU, Acute.

AIGUES-BONNES, see Bonnes.

AIGUES-CAUDES, MIN'ERAL WA'TERS OF, *Aigues-chaudes*, *Eaux chaudes*, *Aqua cal'idæ*. Sulphurous springs, about six miles from Bonnes, Basses Pyrénées, in France: five in number, the temperature of which is respectively,

about 70°, 81°, 92°, 93°, and 95°. They are visited from the first of July to November.

AIGUILLE, (from dim. of *acus*.) Needle—*a. à Acupuncture*, see Needle—*a. à Appareil*, see Needle—*a. à Bec de Lièvre*, see Needle—*a. à Cataracte*, see Needle—*a. de Deschamps*, see Needle—*a. Engainée*, see Needle—*a. à Fistule*, see Needle—*a. à Gaine*, see Needle—*a. à Ligature*, see Needle—*a. à Manche*, see Needle—*a. à Sétou*, see Needle—*a. à Suture*, see Needle.

AIGUILLON (F.) (from *acus*), *Spi'na Helmon'tii*. A term used since the time of Van Helmont to designate the proximate cause of inflammation. According to him, an inflamed part is in the same condition as if an *aiguillon* or thorn were thrust into it.

AIGUISER, (from *aigu*, 'sharp.') To acidulate.

AIL, Allium.

AIL, (from (Sax.) *arðlian*, 'to be sick.') Disease.

AILAN'THUS GLANDULO'SA, *Chinese' Su'-mach*, *Tree of Heaven*, (from *aylanto*, an Amboyna word, signifying "tree of heaven.") A tree; a native of China, but well known in the United States as a shade tree. *Ord. Xanthoxylacæ*. Its flowers have a strong peculiar odour. The powdered bark, in doses of eight or ten grains, has been used as an anthelmintic.

AILE, Ala, *Aileron*.

AILERON (F.), *Ex'tre'ma A'la seu Pin'nula*, (dim. of *aile*, 'a wing.') The extremity of the wing of a bird, to which the great feathers are attached.

AILERONS DE LA MATRICE. Three folds at the base of the broad ligaments of the uterus, which are occupied by the ovary and its ligament, the Fallopian tube, and the round ligament.

AILING, (see *Ail*.) Sick, Sickly.

AILMENT, (see *Ail*.) Disease.

AIMA, (*άίμα*), see Hæma.

AIMANT, (from *adamas*.) Magnet.

AIMORRHŒA, Hæmorrhagia.

AIMORRHŒIS, Hæmorrhœis.

AINE, Inguen.

AIPATHIA, (*αἰμαθία*, from *αι*, 'ever,' and *παθος*, 'disease.') Continent disease.

AIPI, *Jatropha manihot*.

AIPIA CŒXERA, *Jatropha manihot*.

AIPIPOCA, *Jatropha manihot*.

AIR, (*αἴρ*) *Aér*, (*αἰ*, 'I breathe.') *Pneu'ma*, *Com'mon Air*, *Atmospher'ic air*, (F.) *Air atmosphérique*, is an invisible, transparent, inodorous, insipid, ponderable, compressible, and elastic fluid, which, under the form of the atmosphere, surrounds the earth to the height of 15 or 16 leagues.

Air is essentially composed of two gases, oxygen and nitrogen, in the proportion of about 20 of the former, to 80 of the latter. Oxygen is the vital portion, but the nitrogen is necessary to dilute it. Air also contains a small portion of carbonic acid gas, and has always floating in it aqueous vapour, different terrestrial emanations, etc. Its effects upon the human body vary according to its greater or less density, temperature, moisture, etc. Change of air is found extremely serviceable in the prevention and cure of certain morbid conditions. See Climate, and Respiration.

AIR ACIDE VITRIOLIQUE, Sulphurous acid—*a. Alcalin*, Ammonia—*a. Atmosphérique*, Air

AIRBLAD'DER, *Swimbladder*, *Swim'ming-bladder*, *The Swim*, *Vesi'ca natato'ria*, (F.) *Vesie natatoire*. An abdominal organ in many fishes, sometimes communicating by means of a duct, *Ductus pneumat'icus*, with the alimentary canal, at others, not, which is considered by some to belong to the respiratory system. Its contents are the elements of atmospheric air, but in dif-

ferent proportions; and its chief and general function appears to be to regulate the specific gravity of the fish.

AIR CELLS OF THE LUNGS, Cellules, bronchic—a Chamber, Folliculus aeris.

AIR, DEPHLOGISTICATED, Oxygen—a. Empyreal, Oxygen—a. Factitious, Carbonic acid.

AIR DU FEU, Oxygen.

AIR, FIXED, Carbonic acid—*a. Gaté*, Azote—a. *Hépatique*, Hydrogen sulphuretted—a. Inflam-mable, Hydrogen, Hydrogen carburetted—a. Me-phitic, Carbonic acid.

AIR PASSAGES, (F.) *Voies aériennes*, *V. aéri-fères*. The larynx, trachea, bronchia, etc.

AIR PUANT, Hydrogen, sulphuretted.

AIR, PURE, Oxygen.

AIRSACS, see Pulmo.

AIR, SOLID, of Hales, Carbonic acid—*a. Vicié*, Azote—a. Vital, Oxygen.

AIRAIN, (*æs*, *æris*, 'brass.') Bell-metal, Brass.

AIRE, (*area*, 'a space.') Areola.

AIRELLE ANGULEUSE, Vaccinium myr-tillus—a. *Punctuée*, Vaccinium vitis idæa.

AIRIGNE, Hook.

AIR'THREY, MIN'ERAL WATERS OF. Airthrey is situate about two miles north of Stirling, Scotland. The waters are saline cathartics; containing chloride of sodium, chloride of calcium, sulphate of zinc, and chloride of magnesium.

AISELLE, Axilla.

AISTHESIS, (*αισθησις*) *Æsthesis*.

AITCH, Pain.

AITHOMO'MA, (*αιθος*, 'black.') A black condition of all the humours of the eye.

AITIA, (*αἰτια*) Cause.

AITIOLOGY, *Ætiologia*.

AITION, (*αιτιον*) Cause.

AIX-LA-CHAPELLE, MIN'ERAL WATERS OF. Called by the Germans, *Aachen*. A thermal, sulphureous mineral water, which contains, in 1000 grammes, 28.54 cubic inches of sulpho-hydric acid gas, 18.05 cubic inches of carbonic acid gas, 0.1304 grammes of carbonate of lime, 0.0440 grammes of carbonate of magnesia, 0.5444 grammes of carbonate of soda, 2.3697 grammes of chloride of sodium, 0.2637 of sulphate of soda, and 0.0705 of silica. The temperature is 134° Fahrenheit. The springs are visited at all seasons.

The factitious water of *Aix-la-Chapelle*, *A'qua Aquigranensis*, (F.) *Eau d'Aix la Chapelle*, is made by adding pure water f³ivjss, to hydro-sulphuretted water f³iv., carbonate of soda gr. xx, chloride of sodium gr. ix.—Ph. P.

There are thermal sulphureous springs at Aix in Savoy (98°), and some thermal springs at Aix in Provence (91°).

AIZOON, (*αι*, 'ever,' and *ζωος*, 'alive.') Sem-pervium tectorum.

A'JUGA, (said to be a corruption of *ab'iga*, [?]) *A. pyramidalis*, *Consol'ida me'dia*, *Bu'gula*, *B. pyramidalis*, *Teu'crium pyramidale*, *Up'right Bu'gloss*, *Mid'dle Consound*, (F.) *Bugle pyramidale*. This plant is substringent and bitter.

AJUGA CHAMÆPITYS, *Teucrium chamæpitys*.

AJUGA REP'TANS, *Bu'gula*, *B. reptans*, *Com'mon Bu'gle*, (F.) *Bugle rampant*, has similar properties.

AKATALIS, see *Juniperus communis*.

AKATERA, see *Juniperus communis*.

AKÉRATODIAPHANIE, (*α*, *κερας*, gen. *κερα-τος*, 'horn' (cornea), *δια*, 'through,' and *φαινα*, 'I shine.') See Caligo.

AKINESIA, Acinesia.

AKNEMIA, Acnemia.

AKOLOGY, Acology, *Materia Medica*.

AKRATOPEGÆ, *Acratopægæ*.

AKRO, in composition, see *Acro*.

AKROTHYMION, *Acrothymion*.

AKSIS (Old Eng.), Intermittent Fever

AL, in composition, see *Ad*.

A'LA, 'a wing,' *Pin na, Pte'ryx*, (F.) *Aile*. A term often used by anatomists for parts which project like a wing from the median line; as the *A'la na'si*, *Ala* of the *u'terus*, etc. See *Axilla*, and *Pavilion* of the Ear. Also, *Pterygium*.

ALA EXTREMA, see *Aileron*.

ALABA'MA, MIN'ERAL WA'TERS OF. *Tallahat'ta springs* contain sulphur, and salts of iron, lime, and magnesia. But the most noted are those of *Bladen*. See *Bladen*, Mineral Waters of. *Bailey's spring* is an acidulo-sulphurous chalybeate. It is fourteen miles from *Tuscumbia*.

ALABAST'ER, (αλαβαστρος) *Alabas'trum*, (F.) *Albâtre*, *Alabastri'tes*. A variety of compact gypsum; of which an ointment was once made—the *unguentum alabastri'num*; used as a discutient. Alabaster likewise entered into several dentifrices.

ALABASTRITES, (αλαβαστριτης) (λιθος) *Alabaster*.

ALÆ (pl. of *Ala*), INTERNÆ MINORES CLITORIDIS, *Nymphæ*—a. *Magnæ*, see *Sphenoid bone*—a. *Majores*, *Labia pudendi*—a. *Minores*, *Nymphæ*, see *Sphenoid bone*—a. *Muliebres minores*, *Nymphæ*—a. *Nasi*, see *Nasus*—a. *Pudendi Muliebris* *Labia pudendi*—a. *Pulmonum*, see *Pulmo*—a. of the *Uterus*, see *Ala*—a. *Vespertilionis*, see *Uterus*.

ALAIRE, *Alaris*.

ALAISE, *Alcæ*.

ALALIA, (α, and λαλεω, 'I speak.') *Mutitas*.

ALAMBIC, *Alembic*.

ALANFU'TA. A name given by the Arabians to a vein, situate between the chin and lower lip, which they were in the habit of opening in cases of fœtor of the breath.

ALAQUE'CA. The Hindoostanee name of a stone, found in small, polished fragments, which is considered efficacious in arresting hæmorrhage when applied externally. It is a sulphuret of iron.

ALARES MUSCULI, (pl. of *alaris*.) *Pterygoid muscles*.

ALA'RES VE'NÆ. The superficial veins at the fold of the arm.

ALARIA OS'SA. The wing-like processes of the sphenoid bone.

ALARIS, *Ala'rius*, *Aliform'is*; (from *ala*, 'a wing,' (F.) *Aloire*.) Wing-shaped, winged.

ALATERNUS, COMMON, *Rhamnus alaternus*—a. *Latifolius*, *Rhamnus alaternus*.

ALAT'US, *Pterygo'des*, *Ho'mo ala'tus*. One whose scapula project backward like wings.

ALBA'DARAN, *Alda'baran*. The sesamoid bone of the metatarso-phalangeal joint of the great toe. The Rabbis and Magicians attributed extraordinary virtues to it.

ALBAGIAZI, *Sacrum*.

ALBAMENTUM, (*albus*, 'white.') *Albumen*.

ALBAN, SAINT, MIN'ERAL WA'TERS OF. A French acidulous chalybeate, in the department of the Loire.

ALBARAS ALBA, *Lepa alphoides*—a. *Nigra*, *Lepa nigricans*.

ALBARÆS, *Lepa alphoides*.

ALBAROS, *Lepa alphoides*.

ALBÂTRE, *Alabaster*.

ALBEDO UNGUIUM, (*albus*, 'white.') See *Nail*.

ALBIAN, *Albino*.

ALBIN D'ŒUF, *Albumen ovi*.

ALBINISM, see *Albino*.

ALBINISMUS, see *Albino*.

ALBI'NO, 'white,' *Al'bian*, ([L.] *albus*, 'white,') *Leuca' thiops*, *E'thiops al'bus*, *Don'do*, (F.) *Blafard*, *Nègre-blanc*. A Spanish word applied to individuals of the human race who have the skin and hair white (*Achromodermie* and *Achromotrichomie*, of *Piorry*): the iris very pale, bordering on red;

and the eyes so sensible that they cannot bear the light of day. This condition, which has been called *Leuca' thopia*, *Alpho'sis*, *E'thiop'ica*, *Albinos'mus*, *Albinis'mus*, *Al'binism*, *Kokerlakism*, *Leucopath'i'a*, *Leuco'sis*, (F.) *Leucéthiopie*, is seen more frequently in the Negro. Both sexes are exposed to it. It does not seem to be true that there are tribes of *Albinos* in the interior of Africa.

ALBINOISMUS, see *Albino*.

ALBOR OVI, *Albumen ovi*.

AL'BORA. A kind of itch or complicated leprosy, according to *Paracelsus*.

ALBOT, *Crucible*.

ALBOTIM, *Terebinthina*.

ALBUGINÆ, *Albugineous*.

ALBUGIN'EA, *A. Testis*, *Tu'nica albugin'ea*, *Perites'tis*, *Du'ra ma'ter test'is*, *Membra'na capsula'ristes'tis*, (F.) *Albuginée*, *Tunique albuginée*, (*albus*, 'white,') A strong, fibrous, and resisting membrane, which immediately envelopes the testicle, and has, at its upper part, an enlargement, called *corpus Highmorianum*. From its inner surface it sends off a number of flat, filiform prolongations or septa, between which are contained the seminiferous vessels. Externally, it is covered by the *tunica vaginalis testis*.

ALBUGINEA OCULI, *Sclerotic*.

ALBUGINÉE, *Albuginea*.

ALBUGIN'EOUS, *Albugin'eus*, (F.) *Albuginée*. A term applied to textures, humours, &c., which are perfectly white.

ALBUGIN'EOUS FI'BRE, (F.) *Fibre olbuginée*. A name given by *Chaussier* to what he considers one of the four elementary fibres.

The albugineous fibre is linear, cylindrical, tenacious, elastic, but little extensible, and of a shining, satiny appearance. It forms fasciæ or fasciculi, which constitute the tendons, articular ligaments, and aponeuroses; hence the name *Albugin'eous membranes*, given by *Chaussier* to the fibrous membranes.

Gauthier considered, that the *rete mucosum* consists of four layers, to two of which he gave the names *membra'na albugin'ea profun'da* and *membra'na albugin'ea superficia'lis*, respectively.

ALBUGINI'TIS, (F.) *Albuginite*. A term employed by some authors for inflammation of the albugineous tissue. Thus, *gout* and *rheumatism* are regarded as species of the genus *albuginitis*.

ALBUGO OCULI, gen. *Albu'ginis*, (from *albus*, 'white,') see *Sclerotic*—a. *Oculorum*, *Leucoma*—a. *Ovi*, *Albumen ovi*.

ALBULA, (*albus*, 'white,') *Leucoma*.

ALBUM CANIS, (*albus*, 'white,') *Album græcum*—a. *Ceti*, *Cetaceum*.

AL'BUM GRÆ'CUM, *Al'bum Ca'nis*, *Cynoc'oprus*, *Spo'dium Græco'rum*, *Ster'cus Cani'num* *Al'bum*, The white dung of the dog. It consists almost wholly of *phosphate* and *carbonate of lime*, from the bones used as food. It was formerly applied as a discutient gargle in quinsies, and as an antiepileptic; but is now justly banished from practice.

ALBUM NI'GRUM. The excrement of the mouse.

ALBUM OCULI, see *Sclerotic*.

ALBUM RHA'ZIS. A white ointment made of cerusse and lard, prescribed by the Arabian physician *Rhazes*.

ALBUMEN, gen. *Albu'minis*, *Leuco'ma*, *Ooni'nē*, *Ozemun*, (*albus*, 'white,') (F.) *Albumine*. An immediate principle of animals and vegetables, which constitutes the chief part of the white of egg. It is found in the serum, chyle, synovia, serous fluids, &c. There is not much difference in chemical composition between animal and vegetable albumen, fibrin, and casein. Also, the white of the eye. See *Sclerotic*.

ALBUMEN OVI, *Ovile'con*, *Ovul'bumen*, *Albu'men*, *Albu'mor*, *Albu'go seu* *Al'bor seu* *Can'didum* *O'vi*,

Claretta, O'vi al'lus li'gnor, Albumen'tam, Luc a'rie, Glair, White of egg, (F.) Blanc d'œuf, (Old F.) Albin d'œuf, is used in pharmacy for suspending oils, &c., in water. See Ovary.

ALBUMEN, SALIVARY, see Saliva.

ALBUMINE, Albumen—a. *Cérébrale, Neurine*—a. *du Cerveau, Neurine*.

ALBUMINOID, *Albuminoides*, (albumen, and *eidōs*, 'resemblance.') Resembling albumen, as fibrin, casein, &c. It is often applied to aliments in the same sense as proteinous.

ALBUMINOSE. The soluble principle of fibrin, identical with the dominant principle of 'the white of egg according to Bouchardat. According to others, the ultimate product of the transformation, in the stomach, of albuminous matters, which renders them fit for assimilation and nutrition. Albuminose is not coagulable by heat, and incompletely so by acids; an excess of acid dissolving the precipitate. Robin and Verdel regard it as synonymous with the *casein* of the small intestine of Tiedemann and Gmelin, the *gelatiniform matter of the intestine* of Prévost and Morin, and the *casein of the blood* of Dumas. See Peptone.

ALBUMINO'SIS, (albumen and *osis*.) A condition of the blood, in which the ratio of albumen is increased, as in abdominal typhus, variola, rubella, &c.

ALBUMINURIA. A hybrid term, (albumen, and *ουρον*, 'the urine.') A condition of the urine in which it contains albumen, the presence of which is indicated by its coagulation on the application of adequate heat. See Kidney, Bright's disease of the ure.

ALBUMINURIC, *Albuminuricus*, (F.) *Albuminurique*. Relating or appertaining to albuminuria. One labouring under Albuminuria.

ALBUMINURORRÉE, (albumen, *ουρον*, 'urine,' and *ῥεω*, 'I flow.') Kidney, Bright's disease of the.

ALBUMOR, Albumen ovi.

ALCAEST, *Al'cohest, Al'chaest*, (perhaps from (G.) all, 'all,' and *geist*, 'spirit.') A word invented by Paracelsus to designate a liquor, which, according to him, was capable of removing every kind of swelling.

The same word was used by Van Helmont for a fancied universal solvent, capable of reducing every body to its elements.

ALCAEST OF GLAUBER is a thick liquor obtained by detonating nitrate of potassa on hot coals, which transforms it into subcarbonate of potassa.

ALCAEST OF RESPOUR is a mixture of potassa and oxyd of zinc.

ALCAHEST, Alcaest.

ALCAHOL, Alcohol.

ALCALES'CENCE, *Alkales'cence, Alcalescen'tia*. The condition in which a fluid becomes alkaline.

ALCALESCENCE OF THE HUMOURS was an old notion of the humourists. It can only occur during the putrid fermentation of animal matters, which contain azote, and produce ammonia. ALCALIN'ITY is the quality of being alkaline.

AL'CALI or ALCA'LI, *Al'kali*, (from *al* (Arab.) 'the,' and *kali*, the name of the *Salsola Soda*, a plant which contains a large quantity of one of the principal alkalis—*soda*.) The alkalis are substances soluble in water, possessing generally a urinous, acrid, and caustic taste, turning the syrup of violets green, and restoring to blue infusion of litmus, which has been reddened by acids; reddening the yellow of turmeric, and having the greatest tendency to unite with acids, whose character they modify, and form salts with them. In medicine we understand by this term *Potassa, Soda, or Ammonia*.

Soda and potassa are *fixed alkalis*, ammonia is the *volatile alkali*.

ALCALI AMMONIACUM ACETATUM, Liquor ammoniæ acetatis—a. *Ammoniacum fluidum, Liquor ammoniæ*.

ALCALI CAUS'TIC, *Al'kali Caus'ticum*. A pure alkali. One deprived of its carbonic acid.

ALCALI, FIXED, see Alkali—a. *Fixum tartarizatum, Potassæ tartras*—a. *Minerale sulphuricum, Soda, sulphate of*—a. *Tartari aceto saturatum, Potassæ acetat*—a. *Vegetabile salito-dephlogisticatum, Potassæ murias hyperoxygenatus*—a. *Vegetabile tartarizatum, Potassæ tartras*—a. *Vegetabile vitriolatum, Potassæ sulphas*—a. *Volatile*, see Alkali—a. *Volatile acetatum, Liquor ammoniæ acetatis*—a. *Volatile aeratum, Ammoniæ carbonas*—a. *Volatile ex sale ammoniaco, Ammoniæ carbonas*.

ALCALIGENE, (*γεννω*, 'I generate,') Azote.

AL'CALINE, *Al'kaline, Alkali'nus, Alkali'nus*, (F.) *Alcalin*. Having the properties of an alkali.

ALCALINITY. See Alkalinescence.

ALCALINURIA, Alkalinuria.

AL'CALOID; (*alkali*, and *ειδός*, 'form.') A name given to the organic alkalis to distinguish them from the mineral, from which they differ as regards composition and general properties;—having nothing in common but their basic properties. *Brucia, Emetia, Morphia, Strychnia*, &c., are *alcaloids*.

ALCANA, *Anchusa officinalis*.

ALCANNA MAJOR LATIFOLIA DENTATA, *Prinos*—a. *Orientalis, Lawsonia inermis*—a. *Spuria, Anchusa tinctoria*—a. *Vera, Lawsonia inermis*.

ALCEA, (*αλκω*, 'I help,') *Hibiscus abelmoschus*—a. *Ægyptiaca, Hibiscus abelmoschus*—a. *Indica, Hibiscus abelmoschus*.

AL'CEA RO'SEA, *Common hollyhock, Hoc, Hocks*. Emollient, like Althæa.

ALCHACHENGE, *Physalis*.

ALCHACHIL, *Rosmarinus*.

ALCHAEST, Alcaest.

ALCHEMIL'LA, (said to have been celebrated with the Alchemists [?], supposed by some to be of Arabic derivation.) *A. vulgaris, Com'mon La'dies' Man'tle, Pes Leo'nis, Leontopo'dium*, (F.) *Pied de Lion*. Formerly in great repute as an astringent in hemorrhage.

ALCHEMY, Alchymy.

ALCHITRAM, see *Pinus Sylvestris*.

ALCHITURA, see *Pinus Sylvestris*.

ALCHHOL, Alcohol.

ALCHORNEA LATIFOLIA, see Alcornoque

AL'CHYMY, *Al'chymy, Alchemi'a, Alchimi's, Adept'a Philosoph'ia*, (from *al*, an Arabic particle, signifying, 'superiority, excellence,' and *Chimia*, 'Chymistry.') This word was formerly synonymous with Chymistry; but, from the 7th century it has been applied to the mysterious art of endeavouring to discover a universal remedy, and a mode of transmuting the baser metals into gold: an operation to which they gave the name *O'pus mag'num*, and *Philos'opher's stone*.

Alchymy has also been called *Sci'en'tia vel Philosoph'ia Hermet'ica*, from an idea that Hermes or Mercury was its inventor. Harris has well defined this chimerical art: '*Ars sine arte, cujus principium est mentiri, medium laborare, et finis mendicare.*'

AL'CHYMIST, (Old Eng.) *Alkymistre, Flatua'rius, Adept'*. One pretending to alchymy.

ALCOCALUM, *Cynara scolymus*.

AL'COHOL, gen. *Alcohol'is, Al'cohol, Al'choöl, Al'kol, Al'col, Al'cool, Al'kool*. An Arabic word, formerly used for an impalpable powder, and signifying 'very subtle, much divided.' At the present day it is applied to highly rectified spirit of wine:—see *Spiritus rectificatus or rectified spirit*, distilled from dried subcarbonate of po-

ta.s.a. In the Ph. U. S., Alcohol is rectified spirit of the specific gravity 0.835. The Dublin college has a *spiritus fortior*, used in the preparation of certain essences, whose specific gravity is 0.818, and the Pharmacopœia of the United States an *Alcohol Fortius, strong'er al'cohol*, s. g. 0.817. *Alcohol, absolute alcohol*, of the Edinburgh and Dublin Pharmacopœias, is of specific gravity 0.797.

Alcohol is an inflammable liquor, lighter than water, of a warm, acrid taste, colourless, transparent, and of a pungent, aromatic smell. It is the product of the distillation of vinous liquors; is miscible with water in all proportions, and is the direct solvent of resins, balsams, &c. Various other vegetable principles are soluble in it, and hence it is used, in different states of concentration, in the preparation of *elixirs, tinctures, essences*, &c.

Alcohol acts on the animal body as a powerful stimulus: as such, in a dilute form, it is used in the prevention and cure of disease. Its habitual and inordinate use is the cause of many serious affections, of a chronic character especially, as visceral obstructions, dropsy, &c.

ALCOHOL, ABSOLUTE, see Alcohol.—a. *Æthereus Ferratus*, A. Sulfurico-æthereus ferri.—a. cum Aloe perfoliata, Tinctura aloes.—a. Ammoniae et guaiaci, Tinctura guaiaci ammoniata.—a. Ammoniatum, Spiritus ammoniae.—a. Ammoniatum aromaticum, Spiritus ammoniae aromaticus.—a. Ammoniatum foetidum, Spiritus ammoniae foetidus.—a. Amylicum, Oil, fusel.—a. cum Aromatibus sulphuricatus, Sulphuricum acidum aromaticum.—a. cum Aromatibus compositus, Tinctura cinnamomi composita.—a. Castoriatum, Tinctura castorei.—a. cum Crotone casearilla, Tinctura casearillæ.—a. Dilutum, Spiritus tenuior.—a. Ferratus, Tinctura ferri murialis.—a. cum Sulphate ferri tartarisatus, see Ferrum tartarisatum.—a. cum Guaiaco officinale ammoniatus, Tinctura guaiaci ammoniata.—a. Iodii, Tinctura Iodini.—a. cum Opio, Tinctura opii.—a. Methylic, Spiritus Pyroxylicus.—a. Pyroxylic, Spiritus Pyroxylicus.—a. Sulphuricatum, Elixir acidum Halleri.—a. Sulphuricum, Elixir acidum Halleri.—a. Sulphuris, Carbonis sulphuretum.—a. Vini, Spiritus rectificatus.—a. Wood, Spiritus pyroxylicus.

ALCOHOL'IC, *Alcohol'icus, Spirituosus, Spirituosus*. (F.) *Alcoolique*. Relating to or containing alcohol—as an *alcoholic* drink or remedy.

ALCOHOLISM'US, *Al'coholism*, (F.), *Alcoolisme*. The series of morbid phenomena produced by the use of alcoholic liquors.

ALCOL, Alcohol.

ALCOLÆ, Aphthæ.

ALCOLAT, *Alcoolat*.

ALCOOL, Alcohol.—a. *Camphrè*, Spiritus camphoræ.—a. *de Soufre*, Carbonis sulphuretum.

ALCOLLAT, Spirit. It now generally signifies alcohol, charged, by distillation, with the volatile principles of certain substances, or a distilled spirit.—*Alcolat* (Ph. P.).

ALCOOLATUM, Tincture.—a. Antiscorbuticum, Tinctura de Cochleariis.—a. Carminativum Sylvii, Tinctura de Cochleariis.—a. de Croco compositum, Tinctura de Croco composita.

ALCOOLATURE, *Teinture avec les plantes fraîches*. A name given by M. Béral to certain preparations introduced by him, which consist of alcohol, charged, by maceration, with the soluble principles of fresh plants. See Tincture.

ALCOOLÉ. A name given, in the new French pharmaceutical nomenclature, to medicines resulting from the solvent action of alcohol on different substances, or from its admixture with liquids which unite with it in all proportions. See Tincture.

ALCOOLÉ SUCRÉ, *Ratafia*.

ALCOOLIQUE, Alcoholic.

ALCOOLISER (F.) Formerly, 'to reduce into an impalpable powder.' No longer used.

ALCOOLISME, Alcoholicism.

ALCOLOMETER, (μετρον, 'measure,') Areometer.

ALCOOMETER, Areometer.

ALCORNORQUE, (F.), *Cortex Alcornoco* seu *Chabar'ro*, (after Alchorne, an English botanist.) The bark of *Alchor'nea latifolia*, of Jamaica, which has been considered capable of curing phthisis. It is bitter, tonic, and slightly astringent. Dose of the powder ʒi to ʒss.

AL'CYON, *Hal'cyon*, (αλς, sea, κευιν, 'to conceive,') A swallow of Cochinchina, whose nest is gelatinous and very nutritious. It has been proposed in medicine as analeptic and aphrodisiac.

ALCYO'NIUM, *Bas'tard sponge*. The ashes were formerly employed as dentrifices: they were believed proper for favouring the growth of the hair and beard, and were used in Alopecia.

ALDABARAN, Albadaran.

ALDEHYDE, (abridged from *alcohol dehydro-genatus*, 'alcohol deprived of hydrogen,') see Anæsthetic.

ALDER, AMERICAN, *Alnus serrata*—a. Black, *Prinos*, *Rhamnus frangula*—a. European. *Alnus glutinosa*—a. White, *Clethra alnifolia*.

ALE, Cerevisia.

ALE, WHITE. A drink, used in Devonshire, England, amongst the ingredients of which are milk, spice, and spirit.

ALE'BERRY. A beverage, formerly made by boiling *ale* with *spice* and *sugar*, and *sops* of *bread*. It was given to invalids.

ALECOST, Tanaetum balsamita.

ALECTO'RIOUS LA'PIS, *Alecto'ria*; (αλεκτωρ, 'a cock,') The name of a stone, supposed to exist in the stomach of the cock, or, according to others, in that of the capon, four years old. Many marvellous properties were formerly attributed to it, which are as groundless as its existence. There are no stones in the stomach, except what have been swallowed.

ALEGAR, (*ale*, and (F.) *aigre*, 'sour,') Acetum.

ALEHOOF, *Glechoma hederacea*.

ALEIMMA, gen. *Alein'matis*, (αλειμμα, from αλειψω, 'I anoint,') Liniment.

ALEIPHA, Liniment.

ALEIPTE'RIMUM, (αλειπτηριον, from αλειψω, 'I anoint,') The place in the ancient gymnasium where the combatants anointed themselves.

ALEIP'TRON, (αλειπτρον.) A box for containing ointments.

ALEMA, (αλημα,) Farina.

ALEM'BIC (Arab.), *Moorshead, Capitellum, Capitulum, Am'bicus*, (F.) *Alambic*. A utensil made of glass, metal, or earthen ware, adapted for distillation. A still. It consists of a *body* or *cucurbit*, (F.) *cucurbite, chaudière*, to which is attached a *head* or *capital*, (F.) *chapiteau*, and out of this a *beak* descends laterally to be inserted into the *receiver, worm, condenser*, or *refrigerator*, (F.) *serpentin, réfrigérant*, as the case may be.

ALEM'BROTH (Salt.) *Sal Alembroth*. The alchemists designated by this name, and by those of *Sal sapientia*, *Sal artis*, *Sal vita* and *S. Scien'tia*, the product resulting from the sublimation of a mixture of corrosive sublimate and sal ammoniac. It is stimulant, but not employed.

ALÈSE (F.), *Alèze, Alaise, Lin'teum* (αλεξω, 'I preserve,') A guard. A cloth arranged in several folds, and placed upon a bed, so as to guard it from the lochia or other discharges.

ALETON, (αλεω, 'I grind,') Farina.

ALETRIS, (αλειπα, 'meal,') A. farinosa.

AL'ETRIS, A. *Farino'sa* seu *al'ba*, *Star'grass*, *Star'wort*, *Blazing star*, *Aloeroot*, *Bittergrass*, *Blackroot*, *Unicornroot*, *A'gueroot*, *A'guegrass*, *Col'ieroot*, *Dev'il's bit*, *Meal'y starwort*, (F.) *Alé-tris*. *Nat. Ord.* Asphodelææ. *Sec. Syst.* Hexandria Monogynia. The root—*Aletris* (Ph. U. S.)—is an intense and permanent bitter, and is used as a tonic and stomachic. It is common in the United States.

ALETRIS ALBA, Aletris.

ALETRIS AUREA; indigenous, has similar virtues.

ALEURON, (ἀλευρον,) *Farina*.

ALEURITES TRI'LOBA, (from ἀλευριτης, 'mealy,' from the meal-like substance which covers it,) *Candle nut tree*. *Ord.* Euphorbiaceæ. A plant common in the Sandwich, Society, and other islands in the Southern seas; met with, also, in the East and West Indies. The oil of the fruit, *Kukui oil* in the Sandwich Islands, *Kekune oil* in Ceylon; known in Jamaica as *Spanish Walnut oil*, and, in India, as *Belaum Walnut oil*, is cathartic, and resembles castor oil in its action.

ALEUROTESIS, (from *aleuron*;) see Cribration.

ALEXANDERS, *Smyrniun olusatrum*.

ALEXANDRI ANTIDOTUS AUREA. See *Alexandrine*.

ALEXAN'DRINE, *Emplas'trum Alexan'dri*. A garlic plaster, invented by Alexander, contemporary of Mesuë. Other ancient preparations were called 'Alexandrine,' as the *Alexan'dri antid'otus au'rea*, used in apoplexy; the *Colly'rium sic'eum Alexandri'num*, or 'Colly'rium of King Alexander,' mentioned by Aëtius.

ALEXICACUM, (ἀλεξιν, 'to repel,' κακον, 'evil,') *Amuletum, Alexipharmic*.

ALEXIPHAR'MIC, *Alexiphar'macus*, *Anti-phar'macus*, *Alexi'acus*, *Caco'alexite'ria*, *Leal-phar'macus*, (F.) *Alexipharmaque*, (ἀλεξιν, 'to repel,' and φαρμακον, 'poison,') A term formerly used for medicines which were considered proper for expelling from the body various morbid principles, or for preventing the bad effects of poisons taken inwardly.

ALEXIPYRETICUS, *Febrifuge*.

ALEXIR, *Elixir*.

ALEXITE'RIA, *Cacalexite'ria*, (ἀλεξιν, 'to repel,') Originally, *alexiterium* was used synonymously with remedy. In more modern times it has been applied to a class of medicines, that counteract poisons placed in contact with the exterior of the body, in contradistinction to alexipharmics.

ALEXITERIUM CHLORICUM, see *Disinfection*—a. *Nitricum*, see *Disinfection*.

ALÈZE, *Alèze*.

ALFUSA, *Tutia*.

ALGA MARINA, (*ad* and *ligare*, 'to bind,') *Pila marina*.

ALGALIC, *Catheter*.

ALGALIE, *Catheter*.

ALGAROBA, *Ceratonia siliqua*.

AL'GAROTH, *Al'garot*, *Algaro'thi Pul'vis*, *Pul'vis Angeli'cus*, *Ox'idum* seu *Submu'rias Stib'ii præcipitan'do para'tum*, *Antimo'nii Ox'y'dum*, *Ox'idum antimo'nii Ni'tromuriat'icum*, *Ox'idum Stib'ii Ac'idio Muriat'ico oxygena'to para'tum*, *Mercur'ius Vi'tæ*, *Mercur'ius Mor'tis*, *Flow'ers of Antimony*, (F.) *Oxyde d'Antimoine*, (so called from Victor Algarotti, a Veronese physician,) is separated from the *terchloride of antimony* by pouring the *terchloride* into a large quantity of water. When this, an oxychloride, is washed with abundance of water to separate the muriatic acid, and afterwards by a solution of alkali to remove the *terchloride*, the *teroxide of antimony*, *Antimo'nii Ox'idum* of the Edinburgh Pharmacopœia is left. In the Pharmacopœia of the United States (1863) oxide of antimony is directed to be prepared by

treating sulphuret of antimony with *muriatic acid* and afterwards with *nitric acid*; adding a little water; filtering; pouring the filtered liquid into water; washing the precipitate, and subjecting it to the action of *water of ammonia*. It is used in the preparation of the *tartrate of antimony and potassa*, and was formerly much given as an emetic, purgative, and diaphoretic.

ALGE'DO, gen. *Alge'dinis*, (αλγος, 'pain,') Violent pain about the neck of the bladder, occasionally occurring in gonorrhœa.

ALGEDON, (αλγιδων,) *Pain*.

ALGEMA, (αλγημα,) *Pain*.

ALGESIS, (αλγσις,) *Pain*.

ALGETICUS, See *Algos*.

ALGIA, see *Algos*.

AL'GIDUS, *Alg'id*, (*algor*, 'cold,') (F.) *Algide*. That which is accompanied by coldness.

AL'GIDA FE'BRIS, *Fe'bris horrif'ica*, seu *horrida* seu *quer'quera* seu *crymo'des*, *Bry'ctus*, *Bry'chetus*. (F.) *Fèvre algide*, *Alg'id Fe'ver*. A pernicious intermittent, accompanied by icy coldness, which is often fatal in the second or third paroxysm.

ALGOR, *Rigor*.

AL'GOS, (αλγος, 'pain,') See *Pain*. Hence, *Alge'ticus*, 'painful,' as *Epilep'sia alge'tica*. The suffix *algia* has the same signification—as in *Cephalalgia*, *Pleuralgia*, *Neuralgia*, &c.

ALGOSPAS'MUS, (αλγος, 'pain,' and σπασμος, 'spasm,') Painful spasm or cramp of the muscle.

ALGUE COMMUNE, (L. *alga*.) *Pila maris*—a. *Marine*, *Pila marina*—a. *des Verriers*, *Pila marina*.

ALHAGI, *Agul*.

ALHANDAL, see *Cucumis colocynthis*.

ALHASEF, *Sudamina*.

ALHAUNE, *Lawsonia inermis*.

ALIBILIS, (*alere*, 'to nourish,') *Nutritious*.

AL'ICA, *Hal'ica*, *Farina'rium*, *Chon'drus*, (*alere*, 'to nourish,') A grain from which the ancients made their tisanæ; supposed, by some, to have been the *Triticum spelta*. At times it seems to have meant the tisanæ itself.

AL'ICES, (αλιζω, 'I sprinkle,') Spots which precede the eruption of small pox.

ALIENATIO, (*alienus*, 'another's,' 'strange,') *Anomalia*—a. *Mentis*, *Insanity*.

ALIENATION, MENTAL, *Insanity*.

ALIENATUS, *Insane*.

ALIÉNÉ, *Insane*.

ALIENIST PHYSICIAN, (from *alienus*,) *Psychiater*.

ALIENISTE, *Psychiater*.

ALIENUS, *Delirious*.

ALIFORMES MUSCULI, (*ala*, 'a wing,' and *forma*, 'shape,') *Pterygoid muscles*.

ALIFORMIS, *Alaris*, *Pterygoid*.

ALIGULUS, *Confection*.

ALIMA, (*alere*, 'to nourish,') *Aliment*.

ALIMELLEÆ, *Parotid*.

AL'IMENT, *Alimen'tum*, *Al'ima*, *Harmo'llia*, *Nutri'men*, *Nutriens*, *Sustentac'ulum*, *Ciba'rium*, *Bro'ma*, *Comis'te*, *Ci'bus*, *Es'ca*, *Nutri'tus*, *Nutrimen'tum*, *Si'tos*, *Tro'phæ*. (Prov.) *Bellytimber*. (F.) *Aliment*, *Nouriture*, (*alere*, 'to nourish,') *Food*. Any substance which, if introduced into the system, is capable of nourishing it and repairing its losses.

The study of aliments forms one of the most important branches of hygiene. Properly, all substances, organic or inorganic, that enter into the composition of organized beings are aliments.

As regards the immediate principles which predominate in their composition, they have been classed, but imperfectly, as follows:—

TABLE OF ALIMENTS.

1. <i>Feculaceous.</i>	Wheat, barley, oats, rye, rice, Indian corn, potato, sago, peas, beans, &c.
2. <i>Mucilaginous.</i>	Carrot, salsify, beet, turnip, asparagus, cabbage, lettuce, artichoke, melon, &c.
3. <i>Saccharine.</i>	Sugar, fig, date, raisin, apricot, &c.
4. <i>Acidulous.</i>	Orange, currant, gooseberry, cherry, peach, strawberry, raspberry, mulberry, prune, pear, apple, sorrel, &c.
5. <i>Oleaginous and Fatty.</i>	Cocoa, olive, sweet almond, nut, walnut, animal fat, oil, butter, &c.
6. <i>Caseous.</i>	Different kinds of milk, cheese.
7. <i>Gelatinous.</i>	Tendon, aponeurosis, true skin, cellular texture; very young animals.
8. <i>Albuminous.</i>	Brain, nerve, eggs, &c.
9. <i>Fibrinous.</i>	Flesh and blood.

Dr. Prout has four great classes—the *aqueous*, *saccharine*, *oleaginous*, and *albuminous*:—Dr. Pereira, twelve—*aqueous*, *mucilaginous* or *gummy*, *saccharine*, *amylaceous*, *ligneous*, *pectinaceous*, *acidulous*, *alcoholic*, *oily* or *fatty*, *proteinaceous*, *gelatinous*, and *saline*. Liebig divides them into two classes:—the *NITROGENIZED* or *PLASTIC ELEMENTS* OF NUTRITION, in which he comprises *vegetable fibrin*, *vegetable albumen*, *vegetable casein*, *flesh* and *blood*; and the *NON-NITROGENIZED* OR *ELEMENTS* OF RESPIRATION, in which he comprises *fat*, *starch*, *gum*, *cane sugar*, *grape sugar*, *sugar of milk*, *pectin*, *bassorin*, *wine*, *beer*, and *spirits*. The former alone, in his view, are inservient to the nutrition of organized tissue: the latter are burnt in respiration, and furnish heat.

The following simple arrangement is, perhaps, as little objectionable as any:—

- | | |
|--------------------------------------|---|
| 1. <i>Nitrogenized Aliments,</i> | { Fibrinous (Glutinous?)
Albuminous.
Caseinuous, Gelatinous.
Amylaceous. |
| 2. <i>Non-nitrogenized Aliments,</i> | { Saccharine.
Oleaginous. |
| 3. <i>Inorganic.</i> | |

The second division might be still further simplified, inasmuch as amylaceous aliments are convertible into sugar during the digestive process; and, from both, oleaginous matter may be formed.

Aliments are, at times, divided into the *substantive*, and the *adjective* or *accessory*; the former comprising all the nutritive substances, which are used as food; the latter condiments. See *Condiment*.

ALIMENTAL, Nutritional.

ALIMENTARY, Nutritional—a. Tube, Canal, alimentary.

ALIMENTATION, *Alimenta'tio*. The act of nourishing.

ALIMENTEUX, Nutritional.

ALIMENTUM, Aliment, Pabulum.

ALIMOS, Glycyrrhiza.

ALINDE'SIS, (αλινδής, from αλινδωμαι, 'to be turned about.') A species of exercise, which consisted in rolling in the dust, after having been anointed with oil.

ALIPÆNOS, *Alipæ'num*, *Alipan'tos*, (α, and λιπαρ, 'to be fat.') An epithet formerly given to every external remedy, devoid of fat or moisture; such as powders.

ALIPANTOS, Alipænos.

ALIP'TA, *Alip'tes*, (αλειπτης, from αλειφω, 'I anoint.') He who anointed the Athletæ after bathing. The place where this was done was called *Alip'terium*.

ALIPTE'RIUM, (αλειπτηριον,) see *Alip'ta*.

ALIP'TICA. The part of ancient medicine, which treated of inunction as a means of preserving health.

ALISIER BLANC, *Cratægus aria*.

ALISMA, gen. *Alism'atia*, ([Celt.] from *alis*,

'water.') *A. plantago*, *Arnica montana*—a. *Graminifolia*, *A. plantago*—a. *Lanceolata*, *A. plantago*.

ALIS'MA PLANTA'GO, (αλίσμα,) *Alis'ma*, *A. lanceolata* seu *graminifolia*, *Plantago aquatica*, *Water Plantain*, (F.) *Plantain d'Eau*. Ord. *Alismaceæ*. Ser. *Syst. Hexandria Polygynia*. The fresh root is acrid, and the dried leaves will vesicate. The leaves have been proposed as substitutes for *Uva Ursi*.

ALITURA, (αληρε, 'to nourish,') Nutrition.

AL'KALE, *O'leum Galli'næ*. An ancient pharmaceutical name for pullets' fat.

ALKALESCENCE, Alcalescence.

ALKALI, see *Alcali*—a. *Ammoniacum causticum*, *Ammonia*—a. *Ammoniacum spirituosum*, *Spiritus ammoniæ*—a. *Minerale nitratum*, *Soda*, nitrate of—a. *Minerale phosphoratum*, *Soda*, phosphate of—a. *Minerale salinum*, *Soda*, muriate of—a. *Vegetabile cum aceto*, *Potassæ acetat*—a. *Vegetabile fixum causticum*, *Potassæ fusa*—a. *Vegetable*, *Potash*—a. *Volatile*, *Ammonia*—a. *Volatile causticum*, *Ammonia*—a. *Volatile*, concrete, *Ammonia carbonas*—a. *Volatile*, mild, *Ammonia carbonas*—a. *Volatile nitratum*, *Ammonia nitras*—a. *Volatile tartarizatum*, *Ammonia tartars*—a. *Volatile vitriolatum*, *Ammonia sulphas*.

ALKALINE, Alcaline.

ALKALINURIA, *Alcalinuria*, (from *alkali*, and *ουρον*, 'urine.') A condition of the urine in which it is alkaline.

ALKANET, BASTARD, *Lithospermum officinale*—a. *Dyer's Anchusa tinctoria*—a. *Garden Anchusa officinalis*—a. *Officinalis*, *Anchusa officinalis*.

ALKAR, Medicament.

ALKEKENG, Physalis.

ALKER'MES, *Confectio Alker'mes*, *Alcher'mes*. A celebrated electuary, composed of a multitude of substances. It was so called from the grains of kermes contained in it. It was used as a stimulant. Also, kermes.

ALKERVA, see *Ricinus communis*.

ALKITRAN, Cedria.

ALKOL, Alcohol.

ALKOOL, Alcohol.

ALKYMISTRE, Alchymist.

ALLA, Cerevisia.

ALLAITEMENT, (from *allaiter*,) Lactation.

ALLAITER, (F.) *al*, and *lait*, 'milk,' (L.) *ad lactare*. To suckle.

ALLAMAND'A, *A. Cathartica* seu *grandiflora*, *Ore'lia grandiflora*, *Gal'arips*, *Echi'nus scan'dens*, *Apoc'gynum scan'dens*, (after Dr. Allamand, a Leyden naturalist.) A shrub, native of Guiana, the infusion of whose leaves is said by Linnæus to be useful in Colica Pictonum.

ALLANTODES, Allantois.

ALLANTOIC AC'ID, *Ac'idum allanto'icum*. A peculiar acid found in the liquor of the allantois of the cow.

ALLANTOIDES, Allantois.

ALLANTO'IS, *Allanto'idēs*, *Allanto'des*, *Membra'na urina'ria*, *M. seu Tu'nica Farcimina'lis*, *M. Intestina'lis*, the *Allan'toid Ves'icle*, (αλλας, gen. αλλαντος, 'a sausage,' and ειδος, 'shape.') A sort of elongated bladder, between the chorion and amnion of the fœtus, which is thrown out from the caudal extremity of the embryo, and communicates with the bladder by the urachus. It is very apparent in quadrupeds, but not in the human species. As the allantois is developed, its walls become very vascular, and contain the ramifications of what become the umbilical artery and vein, which, by the elongation of the allantois, are brought through the villi of the chorion, into indirect communication with the vessels of the mother.

ALLANTOTON'ICUM, (αλλας, gen. αλλαντος, 'a sausage,' and τοξικον, 'a poison.') Sausage poison

Wurstgift. The Germans have given this name to a poison developed in sausages formed of blood and liver. See *Trichina spiralis*.

ALLECTUARY. Electuarius.

AL'LEGHANY MOUNTAIN SPRINGS. These are situated at the summit of the Alleghany, in Cambria county, Pennsylvania, near Cresson, where there is an excellent summer retreat, more than two thousand feet above the level of the ocean. The waters are pure. One source, however, *Ignatius's Spring*, is chalybeate.

ALLELUIA. *Oxalis acetosella*.

ALLEN'THESIS, (αλλος, 'another,' and εθεσις, 'introduction,') The entrance into, or presence of extraneous bodies in, the organism.

ALLEVIATOR, (ad, and levare, 'to raise,') An instrument for raising invalids, invented by Mr. Jenks, of Rhode Island. It consists of two upright posts, about six feet high, each supported by a pedestal; of two horizontal bars at the top, rather longer than a common bedstead: of a windlass of the same length, placed six inches below the upper bar; of a cog-wheel and handle; of linen belts from six to twelve inches wide; of straps secured at one end of the windlass; and at the other having hooks attached to corresponding eyes in the linen belts, and of a head-piece made of netting. The patient lying on his mattress, the surgeon passes the linen belts beneath his body, attaching them to the hooks on the ends of the straps, and adjusting the whole at the proper distance and length, so as to balance the body exactly, and then raises it from the mattress by turning the handle of the windlass. To lower the patient again, and replace him on the mattress, the windlass must be reversed.

ALLFLOWER WATER. See *Urine* (of cows).

ALLGOOD, *Chenopodium bonus Henricus*.

ALLHEAL, *Heracleum spondylium*.

ALLIACEOUS, *Alliaceous*, (F.) *Alliacé*, (from *allium*.) Belonging to garlic, as *alliaceous* odour.

ALLIAIRE, *Alliaria*.

ALLIARIA, (from *allium*, its smell resembling garlic.) *A. officinalis*, *Erysimum alliaria* seu *cordifolium*, *Sisymbrium alliaria*, *Jack-in-the-hedge*, *Stinking Hedge-mustard*, *Hedgegarlic*, *Sauce-à-l'herbe*, *Hesperis alliaria*, (F.) *Alliaire*. Ord. Cruciferae. This plant has been sometimes given in humid asthma and dyspnoea. It is reputed to be diaphoretic, diuretic, and antiscorbutic.

The Parisian Codex has a compound syrup of *alliaria*. *Sirap d'erysimum composé*, which is used in hoarseness.

ALLIGATURA, *Fascia*, *Ligature*.

ALLIOTICUS, *Alterative*.

ALLIUM, (Ph. U. S.,) (from *oleo*, 'I smell,') *A. sativum*, *Theriacal rusticorum*, *Ampeloprasum*, *Scorodion*, *Scor'don*, *Garlic*, (F.) *Ail*. Ord. Liliaceae. See *Syst. Hexandria Monogynia*. A native of Sicily, but cultivated for use. The bulbs or cloves, *Ag'lithes*, have a strong, offensive, and penetrating odour, and a sweetish, biting, and caustic taste. Internally, garlic is stimulant, diuretic, expectorant, emmenagogue (?), diaphoretic, and anthelmintic. Externally, it is rubefacient, maturative, and repellent; and a *garlic ointment* is made by digesting, at a moderate heat, for half an hour, two or three cloves of fresh garlic in an ounce of lard, and straining. It is applied as a rubefacient to the chest in croup.

Dose, one to six cloves, swallowed whole, or from fʒss fʒij of the juice.

Taylor's Remedy for Deafness, a nostrum, appears to consist of garlic, infused in oil of almonds, and coloured by *alkonet* root.

ALLIUM ASCALONICUM, *Échalotte*.

ALLIUM CANADENSE, *Mendougarlic*; indigo-nous; has the same properties as *Allium*.

ALLIUM CÆPÆ, *Cæpa Vulgaris*, *Cæpa*, *Cæpa*,

Common Onion, *Cepæla*, *Cressmyon*, (F.) *Oignon*. Acriid and stimulating, and possessing very little nutriment. Onions have been used as stimulants, diuretics, and anthelmintics. The boiled or roasted onion, as a cataplasm, is emollient and maturating. The fresh root is rubefacient. The expressed juice is sometimes used in otalgia and in rheumatism.

ALLIUM GALLICUM, *Portulaca*—a. *Plantaginum*, A. *Victorialis*.

ALLIUM POR'RUM, *Por'rum*, *P. sativum*, *Præsum*, the *Leek* or *Por'ret*; (F.) *Poireau*, *Porreau*. It possesses the same property as the onion.

The virtues of the genus *Allium* depend upon an acrid principle, soluble in water, alcohol, acids, and alkalies.

ALLIUM REBOLENS, *Teucrium scordium*—a. *Sativum*, see *Allium*.

ALLIUM VICTORIA'LE, *A. plantagin'eum*, *Cæpa victorialis*, *Victoria'lis lou'ga*. The root, which, when dried, loses its alliaceous smell and taste, is said to be efficacious in allaying the abdominal spasms of pregnant women (?).

ALLO, (αλλος, 'another,') In composition, 'another.'

ALLOCHET'IA, *Allotrichet'ia*, (allo, and χεζειν, 'to go to stool,') The discharge of extraneous matters from the bowels. The discharge of feces by an abnormal opening.

ALLOCHROMA'SIA, (allo, and χρωμα, 'color,') A change of color.

ALLOPATHIA, (αλλος, 'different,' and παθος, 'affection,') *Allopathy*.

ALLOPATHIC, *Allopathic*.

ALLOEOSIS, (αλλοωσις,) *Alteration*.

ALLOEOTICUS, *Alterative*.

ALLOIOSIS, *Alteration*.

ALLOIOTICUS, *Alterative*.

ALLONGEMENT, *Elongation*.

ALLOPATH, (allo, and παθος, 'affection,') *Allopathist*.

ALLOPATHES, *Allopathic*.

ALLOPATH'IC, *Allopath'ic*, *Allæopath'ic*, *Allæopath'ic*, *Allop'athes*, *Heteropath'ic*. Relating to the ordinary method of medical practice, in contradistinction to the *homæopathic*. See *Allopathy*.

ALLOPATHIE, *Allopathy*.

ALLOPATHIST, *Al'lopath*. One who follows *allopathy*.

ALLOP'ATHY, *Allopath'ia*, *Allæopath'ia*, *Hypenanth'isis*, *Hypenanth'isma*, *Curat'io contrari'orum per contraria* (F.) *Allopathie*. The opposite to *homæopathy*; or, according to Hahnemann, a method of treatment in which remedial agents are employed, the action of which, on healthy man, produces morbid phenomena different from those that are observed in the sick person. The ordinary medical practice is so designated by the *homæopathist*.

ALLOPHASIS, (allo, and φασ, 'I say,') *Delirium*.

ALLOTRIO, (from αλλοτριος, 'pertaining to others; foreign,') In composition, has these significations.

ALLOTRIOCHET'IA, (*allotrio*, and χεζειν, 'to go to stool,') *Allochetia*.

ALLOTRIODON'TIA, (*allotrio*, and οδους, 'a tooth,') *Transplantation of teeth*.

ALLOTRIOËC'CRISIS, (*allotrio*, and εκκρισις, 'separation,') The separation of extraneous matters from the body in disease.

ALLOTRIOEUSTIA, (*allotrio*, and ευσις, 'taste,') *Parægestia*.

ALLOTRIOPHAGIA, (*allotrio*, and φαγω, 'I eat,') *Malacia*.

ALLOTRIOTEX'IS, (*allotrio*, and τεξις, 'parturition,') The bringing forth of an abnormal foetus.

ALLOTRIURIA, (*alotrio*, and *ουρον*, 'urine.') Admixture of foreign matters with the urine.

ALLOTROPISM, or ALLOTROPISM; (F.) *Allotropie*, (*allo*, and *τροπος*, 'a turn or change.') A term introduced into modern chemistry, the object of which is to express the property, possessed by certain simple bodies, of assuming different qualities on being subjected to certain modes of treatment. Carbon, for example, furnishes three forms—plumbago, charcoal, and diamond.

ALL-OVERISH, see Indisposition.

ALLSPICE, see *Myrtus pimenta*—a. Bush, *Laurus Benzoin*—a. Carolina, *Calycanthus*—a. Wild, *Laurus Benzoin*.

ALLUCINATIO, Hallucination.

ALLURE, Influenza.

ALLUS POLLEX, Pollex, see *Digitus*.

ALMA, (*αλμη*,) Water (sea).

ALMARIAB, see *Plumbi oxydum semivitreum*.

ALMEZERION, *Cneorum tricoceum*.

ALMOND, Amygdala.

ALMOND BLOOM. A liquid cosmetic, formed of *Brazil dust* $\mathfrak{z}\text{ij}$, *nutter*, Olij; boil and strain; and add *isinglass* $\mathfrak{z}\text{vj}$, *grana sylvestria* $\mathfrak{z}\text{ij}$, or *cochineal* $\mathfrak{z}\text{ij}$, *alum* $\mathfrak{z}\text{ij}$, *borax* $\mathfrak{z}\text{ij}$; boil again, and strain through a fine cloth.

ALMOND CAKE, see Amygdala—a. of the Ear, Tonsil—a. Earth, *Arachis hypogaea*.

ALMOND, JORDAN, see Amygdala.

ALMOND PASTE, see Amygdala—a. Powder, see Amygdala—a. of the Throat, Tonsil.

ALNUS, *A. glutinosa*—a. *Communis*, *A. glutinosa*.

ALNUS GLUTINOSA, *Alnus*, *A. communis*, *Clethra*, *Betula glutinosa* see *emarginata*, *Europeon Alder*, (Sc.) *Arn*, *Eller*, *Ord*. *Betulaceæ*. A tree which grows in Europe, in moist places. The bark and leaves are astringent and bitter; and hence are employed in intermittents, and as a tonic and astringent.

ALNUS NIGRA, *Rhamnus frangula*.

ALNUS SERRATULA, *American Alder*, has similar properties.

ALOCHIA, (*α*, and *λοχεια*, 'lochia.') Absence of the lochial discharge.

ALOËDARIUM. A compound medicine, containing aloes.

ALOE (Arab.), Aloes—a. *Barbaden'sis*, Aloes hepatica—a. *Capensis*, Aloes, cape—a. False, *Agave Virginica*.

ALOE PURIFICATA, see Aloes Succotorina.

ALOE ROOT, *Aletris farinosa*.

ALOE, SOCOTRINE, Aloes Succotorina.

AL'OËS, *Al'oë*, *Fel Natu'rae*. The inspissated juice of the *Aloe*. *Ord*. *Asphodelæ*. *Sex. Syst*. *Hexandria Monogynia*.

ALOES BARBADENSIS, *A. hepatica*—a. *des Barbades*, *A. hepatica*—a. *Bombay*, *A. hepatica*.

ALOES CABALLINA, *A. Guinien'sis*. *Horscaloes*. Used chiefly for horses. It is collected in Spain and Portugal, and is very coarse.

ALOES EN CALÉBASSES, *A. hepatica*.

ALOES, CAPE, *Al'oë Capen'sis*, *Shining Aloes*; a cheap and excellent form of aloes, collected at the Cape of Good Hope, from *Aloe fer'ox*, *A. Africana*, *A. spicata*, and other species.

ALOES, EAST INDIA, *A. Succotorina*—a. *Guinien'sis*, *A. Caballina*.

ALOES HEPATICA, *A. vulgaria*, *A. Barbaden'sis*, *Al'oë Barbaden'sis*, *Hepat'ic al'oës*, *Bombay al'oës*, *Barbadoes al'oës*, *A. vulgaria extractum*, (F.) *Aloes en calébasses*, *A. des Barbades*. This species has a very disagreeable odour, and an intensely bitter and nauseous taste. Properties the same as the others.

ALOES, HORSE, *A. Caballina*—a. *Lucida*, *A. Succotorina*—a. *Socotrine*, *A. Succotorina*—a. *Spicata extractum*, *A. Succotorina*.

ALOES SUCOTORINA, *Socotrine al'oës*, *Al'oë*

Socotrina, *Turkey al'oës*, *East India al'oës*, *Al'oës lucida*, *A. Zocotrina*, *A. spicata extractum*, *An'ina Al'oës*, is the best species. In the British Pharmacopœia (1864), it is said to be from one or more undetermined species of *Aloe*. Its odour is not unpleasant; taste very bitter, and slightly aromatic; colour reddish-brown, with a shade of purple; mass hard, friable; fracture conchoidal and glossy; soluble in dilute alcohol. Powder of a bright cinnamon-yellow colour. It is cathartic, warm, and stimulating; emmenagogue, anthelmintic, and stomachic. As a cathartic, it affects the rectum chiefly. *Dose*, as a cathartic, gr. v. to $\mathfrak{z}\text{ij}$ in pill.

Purified al'oës, *Al'oë Purificata*, (Ph. U. S., 1863,) is made by melting *Socotrine aloes*; adding alcohol; straining and evaporating.

ALOES, TURKEY, *A. Succotorina*—a. *Vulgaris*, *A. hepatica*—a. *Wood*, *Agalloeum*—a. *Zocotrina*, *A. Succotorina*.

ALOËTIC, *Al'oëticus*. A preparation which contains aloes.

ALOËXYLON, (*ξύλον*, 'wood,') *Agalloeum*.

ALOGOTROPHIA, (*αλογος*, 'disproportionate,' and *τροφή*, 'nutrition,') Irregular nutrition. Used particularly to designate the irregular manner in which the nutrition of bones is effected in rickety individuals.

ALOPECIES, (*αλωπεκς*, from *αλωπηξ*, 'the fox,' being strong in that animal,) *Psoræ*.

ALOPECIA, (*αλωπηξ*, 'a fox,' this animal being said to be subject to the affection,) *Cupillo'rum depil'vium*, *A'thrice depil'is*, *Phalacro'tes*, *Depilatio*, *Tricho'sis A'thrice*, *Gangræ'na Alopec'ia*, *A'trich'ia*, *Depil'vium seu Lap'sus seu Flac'us Pilo'rum*, *Lipso'trich'ia*, *Vul'pis mor'bus*, *Pela'du*, *Baldness*, (F.) *Pelade*. Falling off of the hair; loss of the hair. When this is confined to the crown of the head, it is called *calvit'ies*, although the terms are often used synonymously. When congenital, it is called *A'trich'ia seu Alopec'ia adnata*. The falling off of the hair in old age is termed *Alopec'ia senil'is*.

ALOPECIA ADNATA, see Alopecia—a. *Acreata*, *Porrigo decalvans*—a. *Circumscripta*, *Porrigo decalvans*—a. *Partialis*, *Porrigo decalvans*—a. *Senilis*, see Alopecia.

ALOUCHE, *Cratægus aria*.

ALOUCHI. The name of a gum procured from the *canella alba*.

ALOUCHIER, *Cratægus aria*.

ALOUSENEL, *Cowanina Stansburiana*.

ALPAM. A shrub which grows on the coast of Malabar. Certain parts of this, infused in oil, form an antipsoptic ointment. The juice of the leaves, mixed with that of *calamus*, is employed against the bites of serpents.

ALPHENIC, *Saccharum candidum*.

ALPHITEDON, (from *aliphon*, the bone being broken into small portions like meal,) see *Fracture*.

ALPHITON, (*αλφίτον*,) *Polen'ta*, *Fari'na*. Any kind of meal. Toasted barley-meal. *Polenta* means, also, a food composed of Indian meal, cheese, &c. See *Farina*.

ALPHON'SIN, *Alphon'sinum*. A kind of bullet forceps, similar to a *Porte-crayon*, so-called from the inventor, *Alphonso Ferri*, of Naples.

ALPHOS, (*αλφος*, 'white,') *Lepra alphonides*.

ALPHOSIS ÆTHIOPICA, see *Albino*.

ALPINIA CARDAMOMUM, (after *Prosper Alpinus*,) *Amomum cardamomum*—a. *Exaltata*, *Renecalmia exaltata*—a. *Tubulata*, *Renecalmia exaltata*—a. *Galanga*, *Maranta galanga*.

ALPISIE, *Phalaris Canadensis*.

ALSANDERS, *Smyrnum olusatrum*.

ALSIDUM HELMINTHOCHORTON, *Coralina Corsicana*.

ALSYNE ME'DIA, (*αλσυνή*,) *A. avicula'rum seu vulgaria*, (from *αλσος*, 'a grove,' because growing

abundantly in the woods. [?] *Mor'sus Galli'næ*, *Holos'teum Als'i'næ*, *Stella'ria me'dia*, *Mousecar*, *Chickweed*, *Chickencort*, (F.) *Monron des Oiseaux*, *Morgelina*. Ord. Caryophyllacæ. This plant, if boiled tender, may be eaten like spinach, and forms an excellent emollient poultice. It was formerly regarded as a vulnerary and detergent.

ALSTÓNIA THEÆFOR'MIS, (after Prof. Charles Alston, of Edinburgh.) *Santa Fé Tea*: Nat. Ord. Apocynæ. A plant of New Mexico, the leaves of which are used as tea.

ALSTROEMERIA, (after Alstroemer, of Gothenburg,) see Arrow Root.

ALTAFOR, Camphor.

ALTER, (from *alter*, 'other,') see Unwell.

ALTER SEXUS, 'the other sex,' Sex, female.

ALTERANS, Alterative.

ALTERNANT, Alterative; see, also, Alteration.

ALTERATION, *Alteratio*, (*alter*, 'other,') *Alloio'sis*, *Alloeo'sis*. This word is used in France to express any change which supervenes; for example, in the expression of the countenance (*altération de la face*), or in the structure of an organ (*altération organique*), or in the nature of fluids excreted (*altération de l'urine, des larmes, du lait, &c.*)

Altération is also used in an entirely different sense, to express intense thirst in disease. In this case its etymology is different. It comes from *haléter*, and was formerly written *halétermination*.

Applied to animals more especially, *alteration* is, in common parlance, used for castration.

ALTERATIVE, *Alterans*, *Alloio'ticus*, *Alloeo'ticus*, *Allio'ticus*, *Immutans*. An agent considered to be capable of producing a salutary change in a disease, but without exciting any sensible evacuation. As medicine improves, this uncertain class of remedies becomes, of necessity, diminished in number. See Eutrophic.

(F.) *Altérant*. The French term likewise means, that which causes thirst, — *Siticulo'sus*, *Dipsel'icus*, as *altérer* means both to change, and to cause thirst. *Saltér* is to experience a change for the worse, — (*corrum'pi*.)

ALTERATIVES, RENAL, see Diuretic.

ALTERCANGENON, Hyoscyamus.

ALTERCUM, Hyoscyamus.

ALTERED, see Castratus.

ALTHÆA, (αλθαία, from αλθεῖν, 'to heal,') *A. officinalis*, *Malvavis'cum*, *Aristalthæ'a*, *Hibis'cus*, *Hibis'cum*, *Ibis'chus*, *Ibis'cha mis'mal'va*, *Bismal'va*, *Marshmallow*. (Prov.) *Malice*. (F.) *Guimauve*. Ord. Malvacæ. Sex. Syst. Monadelphia Polyandria. The leaves, *Althæa folia*; and root, *Althæa radix*, contain much mucilage. They are emollient and demulcent, and are employed wherever medicines, possessing such properties, are required. In the Ph. U. S., *Althæa* is the root of *Althæa officinalis*.

ALTHANAIHA, Orpiment.

ALTHEXIS, (αλθξις) Curation.

ALTHEUS, (αλθευς) Physician.

ALTHOS, (αλθος) Medicament.

ALTILIBAT, Terebinthina.

ALU'DEL, *Alu'tel*, *Vitrum sublimato'rium*. A hollow sphere of stone, glass, or earthen ware, with a short neck projecting at each end, by means of which one glass might be set upon the other. The uppermost had no aperture at the top. Aludels were formerly used in the sublimation of various substances.

AL'LULA, (dim. of *ala*, 'a wing,') A little wing.

AL'UM, Alumen, Symphytum—a. Cataplasm, Coagulum aluminosum.

ALUM, EGYPTIAN, *Egyptia stypteria*—a. Ammonia, Aluminæ et ammoniæ sulphas.

AL'UM, AMMONIO-FERRIC. Ferr. et ammoniæ sulphas—a. Potassa, Alumen.

AL'UM, ROCHE, *Alu'men de Rochi*, (F.) *Alun de Roche*. (So called from Roccha in Syria, where there was a manufactory of it.) It is in pieces of the size of an almond, covered with a reddish efflorescence.

Common Roche Alum, *A. Ro'chi Gal'lis*. Fragments of common alum moistened and shaken with prepared bole. It is white when broken.

ALUM ROOT, *Geranium maculatum*, *Heuchera cortusa*.

ALUM, SOLUTION OF, COMPOUND, Liqueur aluminis compositus.

ALUM SPRINGS OF VIRGINIA, see Virginia, mineral waters of.

ALUMEN, gen. *Alu'minis*, (Arabic term, *alum*, or *ḍls*, 'a salt,') *Al'um*, *Hypersul'phas Alu'minæ et Potas'sæ*, *Potas'sæ alu'minosulphas*, *Sul'phas Aluminæ Acid'ulus cum Potas'sâ*, *Sulphas Alu'minæ*, *Sul'phas kal'icoalumin'icum*, *Sulphas alumin'aris*, *Supersul'phas alu'minæ et potas'sæ*, *Argil'la sulphu'rica alcalisa'ta*, *A. vitriola'ta*, *Stypt'eria*, *Supersul'phas Argil'læ alcalisa'tum*, *Argilla kaliæsulphurica*. (F.) *Alun*.

ALUMEN CATINUM, Potash of comnerce.

ALUMEN COMMUNE, *Common alum*, *English alum*, *Rock alum*, *Alumen facti'um*, *A. crystal'linum*, *A. ru'peum*, (F.) *Alun d'Angleterre*, is the variety usually employed. It is in octahedral crystals, but generally in large, white, semitransparent masses; has a sweetish, styptic taste: effloresces in the air, and is soluble in 16 parts of water at 60°. Alum is tonic and astringent, and as such is used internally and externally. Dose, gr. v to xv. In large doses, it is emetic and cathartic.

ALUMEN EXSICCATUM, *Alumen us'tum seu siccatum seu calcinatum*, *Sul'phas aluminæ fus'us*, *Argil'la sulphu'rica us'ta*, *Burnt alum*, *dried alum*. (F.) *Alun calciné*, (*Alum* melted in an earthen vessel until ebullition ceases.) Escharotic.

ALUMEN FIXUM, see Potash—a. *Kinosatum*, *Pulvis sulphatis aluminæ compositus*.

ALUMEN ROMANUM, *Roman alum*, *A. Ru'tilum* seu *Ru'brum*. (F.) *Alun de Rome*. In crystals, which are of a pale red when broken, and covered with a reddish efflorescence.

ALUMEN SICCATUM, *Alumen exsiccatum*.

ALUMINA, ACETATE OF, *Aluminæ Acetas*—a. *Depurata*, *Argilla pura*—a. *Pura*, *Argilla pura*—a. *Sulphate of Aluminæ Sulphas*.

ALU'MINÆ ACE'TAS, *Argil'læ Ace'tas*, *Alu'mina seu Argil'la ace'tica*, *Ac'etate of Alu'mina*. A deliquescent salt, obtained by the addition of acetate of lead to sulphate of alumina and potassa. It possesses the same properties as the sulphate of alumina. *Gannal's Solution* for preserving animal substances is made by dissolving an ounce of acetate of alumina in twenty ounces of water.

ALUMINÆ ET AMMONIÆ SUL'PHAS, *Sulphate of alumina and ammonia*, *Ammonia alum*. A combination of sulphate of alumina and sulphate of ammonia (Ph. U. S., 1863).

ALUMINÆ ET POTASSÆ HYPERSULPHAS, *Alumen*—a. et *Potassæ supersulphas*, *Alumen*—a. et *Potassæ Sulphas*, *Alumen*.

ALUMINÆ SUL'PHAS, *Argil'læ Sul'phas*, *Sul'phate of Alu'mina*. Simple sulphate of alumina may be made by the direct combination of alumina and sulphuric acid, and contains 30 per cent. of the former, to 70 per cent. of the latter. In the Pharmacopœia of the United States (1863), it is directed to be prepared by double decomposition from sulphate of alumina and ammonia, and carbonate of soda; dissolving the precipitate in dilute sulphuric acid. It is a deliquescent salt; and is an excellent antiseptic and detergent to ulcers. It is chiefly used to preserve dead bodies—a strong solution—one pound in two pints of water—being injected into the arteries.

A ben'zinated solution of alumina; formed of sulphate of alumina, Sviii ; dissolved in water Sxxvi , and saturated with hydrated alumina; with which selected benzoin balsam Svi are digested for six hours, and the whole filtered—has been used as a styptic, and, largely diluted with water, as an injection in leucorrhœa and ulcerous affections.

ALUMINÆ SULPHAS ACIDULUS CUM POTASSÂ.
Alumen—a. Sulphas fusus, Alumen exsiccatum.

ALUMINE FACTICE, Argilla pura.

ALUMINII OXIDUM, Argilla pura.

ALUN, Alumen.

ALUNSEL, Gutta.

ALUS, Symphytum.

ALUSIA, (αλυσ, 'I wander,') Hallucination—a. Hypochondriasis, Hypochondriasis.

ALUTEL, Aludel.

ALVAQUILLA, Psoralea glandulosa.

ALVARAS NIGRA, Ichthyosis.

ALVEARIUM, (αλveus, 'a cavity,') Auditory canal, external.

ALVEOLAR, *Alveola'ris*, (diminutive of *alveus*, 'a cavity,') (F.) *Alvéolaire*. That which relates to the alveoli.

ALVEOLAR ABSCESS, Parulis.

ALVEOLAR ARCHES, (F.) *Arcades alvéolaires*, are formed by the margins or borders, *Ju'ga alveola'ria*, of the two jaws, which are hollowed by the alveoli.

ALVEOLAR ARTERY, *Supramaxillary A.*, *Artère sus-maxillaire* of Chaussier, arises from the internal maxillary, descends behind the tuberosity of the upper jaw, and gives branches to the upper molar teeth, gums, periosteum, membrane of the maxillary sinus, and buccinator muscle.

ALVEOLAR BORDER, *Lim'bus alveola'ris*. The part of the jaws that is hollowed by the alveoli.

ALVEOLAR MEMBRANES are very fine membranes, situate between the teeth and alveoli, and formed by a portion of the sac or follicle which enclosed the tooth before it pierced the gum. By some, this membrane has been called the *alve'olodental perios'teum*.

ALVEOLAR VEIN. This has a similar distribution with the artery.

ALVÉOLE, Alveolus.

ALVEOLI DENTIS, see Alveolus.

ALVEOLODENTAL PERIOSTEUM, see Alveolar membranes.

ALVÉOLOLABIAL Buccinator.

ALVEOLUS, (same etymon as Alveolar.) *Bo'thion*, *Odontoboth'rium*, *Odontophat'nē*, *Fre'na* (*Alve'oli*), *Mortari'olum*, *Holmis'cus*, *Præsep'iolum*, *Phat'nē*, *Phat'nion*, *Præsep'ium*, *Pat'nē*, *Path'nē*, (F.) *Alvéole*. The alveoli are the sockets of the teeth, *Alve'oli den'tis*, *Ma'n'ia* seu *Caver'næ den'tium*, into which they are, as it were, driven. Their size and shape are determined by the teeth which they receive, and they are pierced at the apex by small holes, which give passage to the dental vessels and nerves.

ALVEUS, ('a cavity,') Auge—a. Ampullescens, Thoracic duct—a. Ampullus, Receptaculum chyli—a. Communis: see Semicircular canals—a. Utriculosus: see Semicircular canals.

ALVI EXCRETIO, Defecation—a. Fluxus aquosus, Diarrhœa—a. Laxitas, Diarrhœa—a. Profuvium, Diarrhœa.

ALVIDUCUS, (*alvus*, 'the abdomen,' and *ducere*, 'to lead,') Laxative.

ALVINE, *Alci'nus*, (*alvus*, 'the abdomen,') That which relates to the lower belly, as *alvine dejections*, *alvine flux*, *alvine obstructions*, &c.

ALVUM EVACUANS, Cathartic.

ALVUS, Abdomen, Uterus—a. Adstricta seu Adstricta, Constipation—a. Cita, Diarrhœa—a. Dura, Constipation—a. Renum, Pelvis of the kidney—a. Tarda, Constipation—a. Viridis, Dejection.

ALYCE, (ἀλυσ,.) Anxiety.

ALYMPH'IA, (α, and *lymp'ha*, 'lymph,') Want of lymph.

ALYPON, (αλυσον, from α, and λυη, 'pain,') An acrid, purging plant, described by Matthioli. By some it has been supposed to be the *Glab'aria al'gum* of botanists.

ALYSIS, (ἀλυσις,.) Anxiety.

ALYSMUS, (αλυσμος,.) Anxiety.

ALYSSUM PLINII, (α, and λυσσα, 'hydrophobia,') Galium mollugo.

ALYSSUS, (α, and λυσσα,.) Antihydrophobic.

ALYX'IA STELLA'TA, (an Indian name,?) *A. aromati'ca*, (F.) *Alyxie aromati'que*. A plant of the family Apocynaceæ, the bark of which resembles canella alba, and is used in Batavia in pernicious fevers.

AL'ZILAT. In some of the Arabian writers, a weight of three grains.

AM, as a prefix, like *Amphi*.

AMABILE, ('lovely,') *amo*, 'I love,') Lacuna Labii Superioris.

AMADOU, Boletus igniarius.

AMADOUVIER, Boletus igniarius.

AMAGRIR, (from *maigre*, (L.) *mauer*, 'lean,') Emaciate.

AMAGRISSEMENT, Emaciation.

AMAL'GAM, *Amal'gama*, (*ama*, 'with,' and *gama*, 'I unite, marry,') [?] A mixture of mercury and other metals. An amalgam of mercury and silver has been used, in dentistry, for filling teeth.

AMALGAMATION, see Miscogeneration.

AMAND (SAINT). MINERAL WATERS OF, *Saint-Amand-les-Eaux*. These springs, in the department Nord, France, are thermal (77° Fahr.), carbonated chalybeates; and are much used internally, or in the form of *boues*, in chronic affections of the digestive organs and liver, and in rheumatism.

AMANDES, see Amygdala.

AMANI'TÆ, (*amanitai*, from α, and *pavia*, madness: i. e. 'not poisonous,') [?] A name given, by the Greeks and Romans, to the edible *champignons*. *Amanita* forms, at the present day, a genus, some of which are edible, others poisonous. Amongst others, it contains the *Agaricus aurantiacus*, and *A. pseudo-aurantiacus*.

AMARA DULCIS, ('bitter sweet,') *Solanum dulcamara*.

AMARAC'NUM. An ancient and esteemed plaster, containing several aromatics, the marjoram, (*amarakos*,) in particular.

AMARACUS, (*amarakos*,) *Origanum majorana*—a. Tomentosus, *Origanum dictamnus*.

AMARITIES, Bitterness.

AMARITUDO, Bitterness.

AMAROR, Bitterness.

AMARUCACHU, Polyanthus.

AMARUS, *Pi'eros* 'bitter.' (F.) *Amèr*. The bitter principle of vegetables is the great natural tonic, and hence *bitters*, as they are termed collectively, belong to the class of tonics. Several are used in medicine; the chief are, gentian, quassia, cinchona, calumba, dog-wood, &c.

AMAS MURIFORME, see Muriform.

AMASE'SIS, *Amasse'sis*, (α, and *μασσις*, 'mastication,') Mastication when impeded or impracticable.

AMATES, Amethyst.

AMATORII, Oblique muscles of the eye.

AMATORIUM, Lacuna labii superioris—a. Veneficium, Philter.

AMATORIUS MUSCULUS, Obliquus superior oculi.

AMATYSTE, Amethyst.

AMAURO'SIS, *Obfusca'tio*, *Offusca'tio*, (*αμαυρωσις*, from *αμαυρος*, 'obscure,' and *osis*,) *Drop*

serene', Gull'ta serene', Cataract' to nil'gra, Parop'-sis amaurosis, Immobilitas pupill'ae, Suffusio nil'-gra, Black cat'aract, Nervous blind'ness, (F.) Goutte-serene, Cataracte noire, Anopticonervie of Piorry. Diminution, or complete loss of sight, without any perceptible alteration in the organization of the eyes; generally, perhaps, owing to loss of power of the optic nerve or retina—Optic anæsthesia. Counter-irritants are the most successful remedial agents, although the disease is always very difficult of removal, and generally totally incurable.

AMACROSIS DIMIDIATA, Hemipopia—a. Diurnal, Nyctalopia—a. Imperfecta, Hypo-amaurosis—a. Nocturnal, Hemeralopia.

AMAUROT'IC, (αμαυρωτικός,) *Amauro'ticus, (F.) Amaurotique.* Affected with amaurosis.

AMAUROTIC CAT'S EYE, *Galeamauro'sis.* A name given by Beer to an amaurotic affection, accompanied by a remarkable change of colour in the pupil, which presents, apparently in the fundus of the eye, a lighter tint, yellowish or brownish yellow, instead of its natural clear black.

AMA'ZIA, (α, and μᾶζος, 'breast.') A monstrosity, in which there is absence of one or both breasts.

AMB, as a prefix, see Am.

AMBARUM, Ambergis—a. Cineritium, Ambergis.

AM'BÈ, (αβη, from αβαίνω, 'I ascend;') *Ambi.* A superficial eminence on a bone. Also, an old surgical machine for reducing dislocations of the shoulder; the invention of which is ascribed to Hippocrates. It is no longer used. See Crista.

AMBER, Succinum—a. Liquid, see Liquidamber styraciflua.

AM'BERGRIS, 'gray amber,' *Am'bor, Am'bar, Am'bra gri'sea seu cinerea'cea seu ambrosi'aca, Am'barum, Succinum cinereum seu gri'seum, Am'barum cinerit'ium.* A concrete substance, of the consistence of wax, cineritious colour, studded with yellow and blackish spots, and exhaling a very pleasant odour. It seems highly probable that ambergis is formed in the intestines of the whale, and voided with its excrement. Like all aromatic substances, ambergis is slightly antispasmodic and excitant; but it is oftener employed as a perfume than as a medicine.

AM'BIA. A liquid, yellow bitumen, the smell and virtues of which are similar to those of the resin tacamahaca. It is obtained from a spring in India.

AMBI'CUS, Alembic.

AMBIDEX'TER, *Amphidex'tus, (ambo, 'both,' and dexter, 'right.')* One who uses both hands with equal facility. Celsus says the surgeon ought to be 'non minus sinistra quam dextra promptus.' One of the aphorisms of Hippocrates says, that a woman is never ambidexter. This a mistake.

AMBILÆVUS, (ambo, and lævus, 'left,') *Ampharisteros.*

AMBI'POPIA, (ambo and ωψ, 'the eye,') *Diplopia.*

AMBITUS GENITALIS MULIEBRIS, (ambitus, 'circumference,') *Vestibulum.*

AMBL'OMA, gen. *Amblo'matis, (αμβλωμα, from α, and βλωσκω, 'to go, '[?])* Abortion.

AMB O'S, (αμβλωσις,) Abortion.

AMBLOSMUS, Abortion.

AMBLOTHRIDI'ON, (αμβλοθριδιον,) see Abortion.

AMBLOTHRIDIUM, Abortive.

AMBLOTICUS, Abortive.

AM'BLY, (from αμβλυσ, 'obscure;') in composition, obscure.

AMBLYP'HIA, (ambly, and ἄφθ, 'feeling,') *Dulness of the sense of touch.*

AMBLYOGMUS, (αμβλωγμος,) *Amblyopia.*

AMBLYO'PTA, (αμβλωπια, from ambly, and

ωψ, 'the eye,') *Amblyos'mus, Amblyog'mus, Amblyo'pia* (so called by some, according to Castelli, ob ignorantiam Græcæ linguae, Hebetudo visus, Feebleness of sight, (F.) *Amblyopie, Vue faible.* First degree of Amaurosis.

AMBLYOPIA CREPUSCULARIS, Hemeralopia—a. Dissitorum, Myopia—a. Meridiana, Nyctalopia—a. Proximorum, Presbytia.

AMBLYOSMUS, Amblyopia.

AMBOLICUS, (βαλλω, 'I cast out,') Abortive.

AM'BON, (αμβων, 'the raised rim of a shield or dish,') from αβαίνω, 'I ascend,') The fibro-cartilaginous rings or *bourrelets*, which surround the articular cavities, as the glenoid cavity of the scapula, the acetabulum, &c., have been so called. See Crista.

AMBOR, Ambergis.

AMBRA, Succinum—a. Ambrosiaca, Ambergis—a. Cineracea, Ambergis.

AMBAGRISEA, Ambergis.

AMBRE BLANC, Succinum (album)—a. *Jaune, Succinum.*

AMBRETTE, Hibiscus abelmoschus.

AMBRINA AMBROSIOIDES, Chenopodium ambrosioides—a. Anthelmintica, Chenopodium anthelminticum—a. Botrys, Chenopodium botrys.

AMBROISIE DU MEXIQUE, Chenopodium ambrosioides.

AMBRO'SIA, (αμβροσια, from α, and βροτος, 'mortal,') Food which makes immortal, or the food of immortals. Food of the gods.—See, also, Chenopodium botrys.

AMBROSIA ELATIOR, see A. Trifida.

AMBROSIA MARITIMA, Ord. Compositæ. A plant which grows on the shores of the Levant, and has a pleasant, bitter and aromatic taste. It is given in infusion, as a tonic and antispasmodic.

AMBROSIA TRIFIDA, *Horse'weed, Rich'weed, Horse'mint, Horse'cane, Bit'terweed, Great or Tall Rag'weed, Wild Hemp.* This indigenous plant is found in low grounds, and along streams, from Canada to Georgia, and west to Louisiana and Arkansas. It is an annual, and flowers in August and September. An infusion has been recommended locally in mercurial salivation.

Ambrosia Ela'tior, Ragweed, is said by Dr. R. E. Griffith to have much more developed sensible properties.

AMBROSIE DU MEXIQUE, Chenopodium ambrosioides.

AMBULANCE (F.), *Hôpital ambulant, (ambulare, 'to walk,')* A military hospital attached to an army, and moving along with it. Also, a vehicle for the conveyance of the sick and wounded.

AMBULATIO, (ambulare, ambulation, 'to walk,') *Walking.*

AMBULATORY, *Ambulans. Ambulati'vus, Ambulative, (F.) Ambulant.* A morbid affection is said to be 'ambulatory,' (F.) *ambulant*, when it skips from one part to another; as *Erysipêles ambulans*, &c. When blisters are applied successively on different parts of the body, they are called *Vésicatoires ambulants*.

AMBULEIA, Cichorium intybus.

AM'BULI. The Brachmanic name for an Indian aquatic herb, which appears to belong to the family *Lythymachia*. The whole plant has a sweet smell. Its decoction has a very bitter taste, and is an excellent febrifuge. It is also taken in milk in cases of vertigo.

AMBULOFLATULENTUS ET FURIOSUS, *Flat'us furio'sus, Vare'ni.* Painful, mobile, and periodical tumours affecting different parts, which were once considered as the effect of very subtle vapours. Their nature is by no means clear.

AMBUSTIO, (ambi, 'around,' and urere, ustum, 'to burn,') *Burn.*

AMBTUA, Pareira brava.

AMBUYA-EMBO. A very beautiful, creeping

aristoloehia of Brazil, the decoction of which is exhibited successfully in obstructions. It is also used in fumigation and in baths as a tonic.

ÂME, Anima.

AMELI, A Malabar shrub, belonging to a genus unknown. The decoction of its leaves is said to relieve colic. Its roots, boiled in oil, are used to repel tumours.

AMELIA, (αμελία,) Apathy.

AMÉLIE-LES-BAINS, see Arles.

AMENIA, (α, and μην, 'a month,') Amenorrhœa, Emmenagogues.

AMENOMA'NIA, Chairoma'nia, Chæroma'nia, (amēnus, 'agreeable,' and μανία, 'mania,') A gay form of insanity.

AMENORRHŒA, Paramē'nia obstructio'nis, Menœcyp'hia, Menosta'sia, Apophras'tis, Arrhœ'a, Defec'tus seu Reman'sio seu Cessa'tio men'sium, Menstrua'tio impedi'ta, Ischome'nia, Amē'nia, Ametrohœ'mia, (from α, mēnes, and ρεω, 'I flow,') Suppres'sion of the men'ses, (F.) Suppression du flux menstruel. This suppression is most commonly symptomatic, and hence the chief attention must be paid to the cause. Usually, there is an atonic state of the system generally, and hence chalybeates and other tonics are advisable.

Two great varieties of Amenorrhœa are commonly reckoned. 1. A. Emansio'nis, Emans'io men'sium, Menis'chesis, Menos'chesis, Menstrua'tio retenta, Men'sium reten'tio, Reten'tion of the men'ses, when the menses do not appear at the usual age; and, 2. Suppres'sio Men'sium seu Menstrua'tio'nis, Amenorrhœ'a Suppres'sio'nis, Interrup'tio menstruatio'nis, Menstrua'tio suppres'sa, in which the catamenia are obstructed in their regular periods of recurrence. See Emansio Mensium and Menses.

AMENORRHŒA DIFFICILIS, Dysmenorrhœa—a. Emansio'nis, see Amenorrhœa—a. Hymenica, see Hymenica—a. Partialis, Dysmenorrhœa—a. Suppressionis, see Amenorrhœa.

AMENTIA, (α, and mens, mentis, 'mind,') Dementia: see, also, Fatuitas, and Idiotism—a. Senilis, Dementia of the aged.

AMER, Amarus—a. de Bœuf, Bile of the ox—a. de Welther, Acid carbozotic.

AMERICAN, see Homo.

AMERICANUM TUBEROSUM, Solanum tuberosum.

AMERTUME, (from amer, (L.) amarus, 'bitter,') Bitterness.

AMETHYST, Amethys'tus, Am'atyste, Am'ates, (α, and μεθυσ, 'I am drunk,') A precious stone, to which the ancients attributed the property of preventing drunkenness. It was also used as an anti-diarrhœic and absorbent.

AMETHYSUM, (αμεθυσον,) Amethys'tum, (αμεθυσον,) (remedium.) A remedy for drunkenness.

AMET'RIA, (αμετρία, from α, and μετρον, 'measure,') Intemperance. Also, absence of the uterus: (α, and μετρα, 'the uterus,') Amet'ria.

AMETROHŒMIA, (α, μετρα, 'uterus,' and αίμα, 'blood,') Amenorrhœa.

AMIAN'THIUM MUSCETOX'CIUM, (αμιαντος, 'pure,' and ανθος, 'a flower,') Melan'thium musceto'xi'um, Helo'nias erythrosperm'a, Fly-poison, Fall'poison, indigenous. Ord. Melanthaceæ. A narcotic poison, not used, however.

AMICULUM, (amicio, 'I wrap around,') Amnios.

AMIDOLIQUE, (amidum, 'starch,') Appertaining or belonging to starch. M. Bérall calls médicaments amidoliques those, which owe their general properties to starch.

AMIDON, Amylum—a. Iodure d', Starch, Iodide of.

AMIDUM, Amylum.

AMINŒA, Anime.

AMINŒUM (from Aminæa, a region of Italy)

VINUM, Amine'an wine, highly esteemed as a stomachic. Virgil distinguishes it from the Falernian.

AMMA, (appa,) gen. Am'matis, Truss.

AMMI, (appu,) Ammi ma'jus seu cicuta'folium seu vulga'rē seu Bolbe'ri, Am'mias murica'ta, A'pium ammi, Bish'op's weed. The seeds of this plant are aromatic and pungent. They are said to be carminative and diuretic, and are tonic and stomachic.

AMMI BOLBERI, Ammi—a. des Boutiques, see Sison ammi—a. Cicuta'folium, Ammi—a. Majus, Ammi—a. Verum, see Sison ammi—a. Vulgare, Ammi.

AMMION, Hydrargyri sulphuretum rubrum.

AMMIOS MURICATA, Ammi.

AMMISMUS, (ampos, 'sand,') Psammismus.

AMMOCHO'SIA, Ammocho'sis, (ampos, 'sand,' and χρω, 'I pour,') Arenat'io. Putting the human body in hot sand, for the cure of disease.

AMMŌ'NIA, Ammo'nia or Ammoni'acal gas, Volatile al'kali, Al'cali ammoni'acum causticum, A. volatile causticum, Ammo'nia caustica seu pu'ra, Ammoni'acum, A. causticum, Gas ammoniacale, Mephi'tis urino'sa, (F.) Ammoniaque, Air alcalin, Gaz ammoniacal. An alkali, so called, because obtained principally by decomposing sal ammoniac (muriate of ammonia) by lime. This gas is colourless, transparent, elastic, of a pungent, characteristic odour, and an acrid, urinous taste. It turns the syrup of violets green, and its specific gravity is 0.596. When inhaled, largely diluted with common air, it is a powerful irritant. When unmixed, it instantly induces suffocation.

AMMONIA, ACETATE OF, SOLUTION OF, Liquor ammoniac acetatis—a. Arseniate of, Arseniate of ammonia—a. Benzoate of, Ammoniac benzoas—a. Bicarbonate of, see Ammoniac carbonas—a. Caustica liquida, Liquor ammoniac—a. Chlorhydrate of, Ammoniac murias—a. Citrate of, Ammoniac citras—a. Hydriodate of, Ammonium, iodide of—a. Hydrochlorate of, Ammoniac murias—a. Hydrosulphuret of, Ammoniac sulphuretum—a. Hypophosphite of, see Calcis hypophosphis—a. Iodide of, see Iodine—a. and Iron, tartrate of, Ferri et ammoniac tartras—a. Liniment of, strong, Linimentum ammoniac fortius—a. Liquid, Liquor Ammoniac—a. Muriatica, Ammoniac murias—a. Ammoniac nitras—a. Phosphate of, Ammoniac phosphas—a. Præparata, Ammoniac carbonas—a. Pura liquida, Liquor ammoniac—a. Solution of, Liquor ammoniac—a. Solution of, stronger, Liquor ammoniac fortior—a. Tartrate of, Ammoniac tartras.

AMMONIAC GUM, (αμμωνιακον,) Ammoni'acum (Ph. U. S.), Gum'mi Ammoni'acum, Armoni'acum, Moto'rium, (F.) Ammoniac, Gomme ammoniacque, (from Ammonia, in Lybia, whence it is brought.) A gum-resin, the concrete juice of Dore'ma ammoni'acum, of Persia: a species of a genus allied to Ferula; and also the gum-resin of Fer'ula tin-gita'na. The latter is the ammoniacum of commerce. It is in irregular, dry masses and tears, yellow externally, whitish within. Its odour is peculiar, and not ungrateful: taste nauseous, sweet, and bitter. It forms a white emulsion with water: is soluble in vinegar; partially so in alcohol, ether, and solutions of the alkalies.

Gum ammoniacum is expectorant, 'deobstruent' (?), antispasmodic, discutient, and resolvent. It is chiefly used, however, in the first capacity, and in the formation of certain plasters.

Two varieties are met with in the market, Gut'tæ ammoni'aci, the best; and La'pis ammoni'aci, the more impure.

AMMONIAC, SAL, Ammoniac murias.

AMMONIACÆ NITRAS, Ammoniac nitras—a. Sulphas, Ammoniac sulphas.

AMMONIACUM, Ammonia, Ammoniac gum—a. Hydrochloratum ferratum, Ferrum ammoni-

acum—a. Succinatum. Spiritus ammoniæ foetidus —a. Volatile mite, Ammonia carbonas.

AMMONIÆ ACETAS, Liquor ammoniæ acetatis —a. Arsenias, Arseniate of Ammonia.

AMMONIÆ BENZO'AS, Benzoate of Ammonia. A salt formed by the union of benzoic acid and ammonia, which has been prescribed for the removal of gouty depositions of urate of soda in the joints. It is regarded as a good diuretic. It is official in the British Pharmacopœia (1864). Dose, gr. xv to xxx and more.

AMMONIÆ BICARBO'NAS, Bicarbonate of Ammonia was official in the Dublin Pharmacopœia. It has the same properties as the next, and is more palatable. Dose, six to twenty-four grains.

AMMONIÆ CARBO'NAS, A. Subcarbo'nas seu Sesquicarb'onas, Salt of bones, Sal Os'sium, Sal Fulig'inis, Salt of woodsoot, Salt of urine, Volatile Sal Ammoniac, Ba'ker's salt, Al'kali volatilè aëratum, A. volatilè ammoniacalè, A. volatilè ex sa'tis ammoni'aco, Ammoni'acum volatilè mi'te, Ammonium carbo'nicum seu subcarbo'neum, Carbo'nas ammoniæ alkali'nis seu incompletus seu superammoni'acus, Hypocarb'onas ammoniæ, Flores sal'is ammoni'aci, Sal cor'nu cer'vi volatilè, Sal volatil'is sal'is ammoni'aci, Con'crete or mild volatile al'kali, Carbonate or Subcarbonate of ammonia, Ammonia præpara'ta, Sal volatilè, Smelling salt, (F.) Carbonate d'ammoniaque, Sel volatil d'Angleterre, (Ammon. muriat. HJ; Crete Miss. Sublime—Ph. U. S.) (Formerly obtained by the dry distillation of hartshorn, bones, and other animal matters.) A white, striated, crystalline mass; odour and taste pungent and ammoniacal: soluble in two parts of water: insoluble in alcohol: effloresces in the air. It is stimulant, antacid, diaphoretic, and antispasmodic. Dose, gr. v to xv.

Carbonate of ammonia is at times used to form effervescent draughts. One scruple saturates six fluidrachms of lemon juice, twenty-six grains of crystallized tartaric acid, and twenty-six grains of crystallized citric acid.

Carbonate of ammonia is the usual *smelling salts*. When a few drops of Liquor Ammonia fortior are added to it coarsely powdered, and some volatile oil, it forms *Preston Salts*.

AMMONIÆ CITRAS, Citrate of Ammonia. Made by saturating lemon or lime juice, or a solution of citric acid, with carbonate of ammonia. Dose, fʒss. It may be made extemporaneously, and taken in an effervescent state. Seventeen grains of citric acid, or half a fluidounce of lemon juice, will be sufficient for thirteen grains of carbonate of ammonia.

AMMONIÆ CUPROSULPHAS, Cuprum ammoniatum.

AMMONIÆ ET FERRI MURIAS, Ferrum ammoniatum —a. Ferro-citras, Ferri ammonio-citras —a. Hydriodas, Ammonium, iodide of —a. Hydrochloras, Ammonia murias—a. Hydrosulphuretum, Liquor fumans Boylii—a. Hypocarbonas, Ammonia Carbonas.

AMMONIÆ MU'RIAS (Ph. U. S.), Muriate of Ammonia, Hydrochlorate of Ammonia, Chlorohydrate of Ammonia, Sal Ammoni'acum, Sal Ammoniac, Sal Ammoni'acus, Ammonia Muriat'ica, Ammoni'um Muriat'um, Ammoni'æ Hydrochloras, Chloruretum Ammoni'um, Sal Armeni'acum, Sal'mine, Ful'igo Al'ba Philosopho'rum, Misadir, (F.) Muriate d'Ammoniaque, Sel Ammaniac ou Armeniac, (from the temple of Jupiter Ammon, near which it was obtained.) A saline concrete, formed by the combination of muriatic acid with ammonia. In Egypt it is manufactured in large quantities by subliming the soot formed by burning camel's dung—26 pounds of the soot yielding 6 pounds. It is also prepared, in great quantities, by adding sulphuric acid to the volatile alkali obtained from soot, bones, &c., mixing this with common salt, and subliming.

Muriate of ammonia is inodorous, but has an acid, pungent, bitterish, and urinous taste. Three parts of cold water dissolve one. Soluble also in 4.5 parts of alcohol. It is aperient and diuretic, but seldom used internally. Externally, it is employed, producing cold during its solution, in inflammations, &c.

AMMONIÆ NI'TRAS, Ni'trate of Ammonia, Al'kali volatilè nitrat'um, Sal ammoni'acus nitrosus, Ammonia nitrat'a, Ni'tras ammoni'acæ, Ni'trum flammans, (F.) Nitrate d'Ammoniaque. A salt composed of nitric acid and ammonia. It is diuretic and 'deobstruent.' (?) Externally, it is discutient and sialogogue.

AMMONIÆ PHOS'PHAS, Ammoni'um phosphoricum, Phosphate of Ammonia, (F.) Phosphate d'Ammoniaque. This salt has been recommended as an excitant, diaphoretic, and discutient. More recently, it has been proposed as a new remedy for gout and rheumatism, as a solvent of uric acid calculus, and for diseases, acute and chronic, connected directly with the lithic acid diathesis.

AMMONIÆ SESQUICARBONAS, A. carbonas.

AMMONIÆ SUL'PHAS, Sul'phate of Ammonia, Sulphas ammoni'acæ, Ammonium sulphuricum, Al'kali volatilè vitriolat'um, Sal Ammoni'acum secretum Glaube'ri, Sal secretus Glaube'ri, Vitriolum ammoniacalè, (F.) Sulphate d'ammoniaque. Formed by adding sulphuric acid either to sal ammoniac or to ammoniacal liquor. Its properties are like those of the muriate of ammonia.

AMMONIÆ SULPHURETUM, Sul'phuret of Ammonia, Hydrosulphuret of Ammonia, Ammonium Sulphurat'um, Hydrosul'phas Ammoni'æ, Spiritus Beguini, Sp. fulmans Beguini, Sulphuretum ammoni'acæ, Sp. salis ammoni'aci sulphurat'us, Li'quor ammonii hydrothi'odis, Hydrosulphuretum Ammoni'um, Hydrosulph. ammoniacalè aquosum, Hydrog'enosulphuretum ammoni'acæ liquidum, Spiritus sulphuris volatil'is, He'par sulphuris volatilè, Boyle's or Beguine's fuming spirit, (F.) Hydrosulphate sulfurè d'Ammoniaque, Liqueur fumante de Boyle, Sulfure hydrogéné d'Ammoniaque, Hydrosulfure d'Ammoniaque. Odour very fetid; taste nauseous and styptic; colour dark yellowish green. It is reputed to be sedative, nauseating, emetic, disoxygenizing (?), and has been given in diabetes and diseases of increased excitement. Dose, grt. viij to grt. xx.

AMMONIÆ TAR'TRAS, Al'kali volatilè tartarizat'um, Sal Ammoni'acum tartaratum, Tartarus ammoniac, Tartrate of Ammonia, (F.) Tartrate d'Ammoniaque. A salt composed of tartaric acid and ammonia. It is diaphoretic and diuretic; but not much used.

AMMONIÆ VALERIA'NAS, Vale'rianate of Ammonia. This salt is obtained by causing gaseous ammonia, disengaged from muriate of ammonia by lime, to pass into valerianic acid until the acid is neutralized; and crystallizing. (Ph. U. S., 1863). Used like valerian, valerianic acid, and the other valerianates, in neuropathic affections.

AMMONIAQUE, Ammonia—a. Arseniate d', Arseniate of ammonia—a. Hydrosulfate sulfurè d', Ammonia sulphuretum—a. Hydrosulfure d', Ammonia sulphuretum—a. Liquide, Liquor ammoniæ—a. Phosphate d', Ammonia phosphas—a. Sulfure hydrogéné d', Ammonia sulphuretum.

AMMONIUM IODIDUM, Ammonium, iodide of —a. Ioduretum, Ammonium, iodide of.

AMMONIUCUPRICUS SUBSULPHAS, Cuprum ammoniatum.

AMMONION, (appos, 'sand.') An ancient collyrium of great virtue in many diseases of the eye, and which was said to remove sand from that organ.

AMMONIUM ACETICUM LIQUIDUM, Liquor ammoniæ acetatis—a. Arsenicum, Arseniate of ammonia—a. Carbonicum Ammonia carbonas

—a. Hydroiodicum, Ammonium, iodide of — a. Iodatum, Ammonium, iodide of.

AMMO'NIUM, IODIDE OF, *Iod'idum* seu *Iodure'tum ammonii*, *Ammonium Ioda'tum* seu *Hydroiod'i-cium*, *Hydro'iodus ammon'ia*, *Hydro'iodate of am-mo'nia*. This salt is formed by saturating liquid hydroiodic acid with caustic ammonia, and evaporating the solution. It is applied in the form of ointment (℞ *ad adipis* ℥j) in lepra, psoriasis, &c.

AMMONIUM MURIATICUM MARTIATUM SEU MARTIALE, Ferrum ammoniatum — a. Muriatum, Ammonia murias — a. Phosphoricum, Ammonia phosphas — a. Subcarbonicum, Ammonia carbonas — a. Succineum liquidum, Spiritus ammonie succinatus — a. Sulphhydratum, Ammonia sulphuretum — a. Sulphuricum, Ammonia sulphas.

AMNA ALCALIZATA, Water, mineral, saline. AMNEMOSYNE, (α, and μνησιν, 'memory,') Amnesia.

AMNE'STIA, *Amnesia*, *Amnemos'ynē*, (α, and μνησις, 'memory,') *Mo'ria imbec'ilis amne'sia*, *Obliv'io*, *Recollectio'nis jactu'ra*, *Dysasthe'sia inter-talia*, *Debilitas memo'ria*, *Memo'ria dele'ta* (F.) *Perte de Mémoire*, *Loss of Memory*. By some nosologists, amnesia constitutes a genus of diseases. By most, it is considered only as a symptom, which may occur in many diseases.

AMNESTIA, (αμνηστία,) Amnesia.

AMNIC, Amniotic.

AMNIC ACID, Amniotic acid.

AMNIITIS, Amnitis.

AMNIOCLEP'SIS, (amnion, and κλεπω, 'I steal away,') Premature escape of the liquor amnii.

AM'NION, (αμνιον,) *Am'nios*, *Am'nium*, *Hym'nium*, *Char'ta virgin'lea*, *Amnato'ra*, *Aqui'na mem-bra'na*, *Pellu'cida membra'na*, *Gal'lea*, *Scepar'num*, *Indu'sium*, *Amic'ulum*, *Membra'na fa'tum invol'uens*. The innermost of the enveloping membranes of the foetus:—so called because first observed in the sheep (?), (αμνος, 'a sheep,') It is thin, transparent, and is generally considered to be produced by a fold of the external layer of the germinal membrane, rising up, and gradually enveloping the embryo. Its external surface is feebly united to the chorion by areolar and vascular filaments. Its inner surface is polished, and is in contact with the body of the foetus and the liquor amnii.

AMNIORRHŒ'A, (amnion, and ρω, 'I flow,') A premature discharge of the liquor amnii.

AMNIOT'IC, *Amniot'icus*, *Am'nic*, *Am'nicus*, (F.) *Amniotique* ou *Amnique*. Relating, or appertaining, to the amnion.

AMNIOTIC ACID, *Ac'idum am'nicum* seu *amniot'icum*. A peculiar acid, found by Vauquelin and Berthollet in the liquor amnii of the cow.

AMNIQUE, Amniotic.

AMNI'TIS, *Amni'tis*, (amnion, and itis.) Inflammation of the amnion.

AMENOMA'NIA, (amemus, 'agreeable,' and mania.) A form of mania in which the hallucinations are of an agreeable character.

AMOME FAUX, Sison amomum.

AMOMIS, see the next.

AMO'MUM CARDAMO'MUM, *A. verum*, (αμωπος, from α, and ωπος, 'a fault,') *faultless,') Alpini'cia cardamo'mum*, *Caro'pi*, *Mato'nia Cardamo'mum*, *Elletta'ria Cardamo'mum*, *Cardamo'mum Mi'tis*, *Les'ser*, *True*, or *Offic'inal Cardamom*, (F.) *Cardamome de la Côte de Malabar*, *Cardamome*. Ord. Zingiberaceæ. The fruit—*Cardamomum*, (Ph. U. S.)—of this East India plant, *C. Malabaren'sē*, has an agreeable, aromatic odour, and a pungent, grateful taste. It is carminative and stomachic: but is chiefly used to give warmth to other remedies. It is called, also, *Amo'mia*. Dose, gr. v to ℥j.

Amomum Cardamomum of Linnæus, *C. Malabaren'sē* of Round *Cardamom* of Sumatra, Java, and other islands eastward of the Bay of Bengal, yields the round *Cardamom*, *Cardamo'mum rotundum*,

of the shops. The fruits, in their native clusters or spikes, constituting the *Amomum racemosum*, are rarely met with.

AMOMUM CURCUMA, *Curcuma longa*—a. Galanga, *Maranta galanga*.

AMO'MUM GRA'NUM PARADI'SI, *A. max'imum*, *Cardamo'mum majus* seu *piperat'um*, *Molequet'ta*, *Maniquet'ta*. Greater cardamom seeds—*Grana Paradi'si*, *Grains of Par'dise*, (F.) *Graines de Paradi's*—resemble Cardamomum in properties. They are extremely hot, and not much used.

AMOMUM, GREAT-WINGED, *Amomum maximum* — a. Hirsutum, *Costus*.

AMOMUM MAX'IMUM, *Great-wing'ed Amomum*, yields the fruit known in commerce by the names *Ju'va* or *Nepal Cardamoms*, *Benjal Cardamoms* of the Calcutta market, &c.

AMOMUM MONTANUM, see Cassumuniar — a. Pimenta, see Myrtus pimenta — a. Racemosum, see A. cardamomum — a. Sylvestre, see Cassumuniar — a. Zedoaria, *Kæmpferia rotunda*—a. Zerumbet, see Cassumuniar.

AMOMUM ZIN'GIBER, *Zin'giber officina'le* seu *al'bum* seu *ni'grum* seu *commu'ne*, *Zin'ber*, *Gin'ger*, (F.) *Gingembre*. The black and white ginger, *Zin'giber fus'cum* et *al'bum*, *Zin'giber* (Ph. U. S.), are the rhizoma of the same plant, *Zin'giber officina'le*, the difference depending upon the mode of preparing them.

The odour of ginger is aromatic; taste warm, aromatic, and acrid. It yields its virtues to alcohol, and in a great degree to water. It is carminative, stimulant, and sialogogue.

Preserved Gin'ger, *Zingib'ris Ra'dix Condi'ta*, *Ra'dix Zingib'ris condi'ta ex Indiā alla'ta*, is a condiment which possesses all the virtues of ginger.

Gingerbeer Pow'ders may be formed of white sugar ℥j and ℥ij, ginger, gr. x, subcarbonate of soda gr. xxxvj, in each blue paper: acid of tartar ℥iss, in each white paper—for half a pint of water.

Oc'ley's Concentrated Es'sence of Jama'ica Gin'ger is a solution of ginger in rectified spirit.

AMOR, Love.

AMORGE, (αμωργη,) Amurea.

AMORPHUS, (α, and μωρφη, 'form,') Anhistous, Anideus.

AMOSTEUS, Osteocolla.

AMOUR, (amor, 'love')—a. Physique, Appetite, venereal.

AMOUREUX, (muscule.) Obliquus superior oculi.

AMPAC, *Am'pacus*. An East India tree, the leaves of which have a strong odour, and are used in baths as detergents. A very odoriferous resin is obtained from it.

AMPAR, Succinum.

AMPELOCARPUS, (ampelos, and καρπος, 'fruit,') Galium aparine.

AMPELOLEUCE, (ampelos, and λευκος, 'white,') Bryonia alba.

AMPELOPRASUM, (ampelos, and πρασον, 'a leek,') Allium.

AMPELOP'SIS QUINQUEFO'LIA, (ampelos, and πτε, 'appearance,') *Vinifolia Creeper*, *American Ivy*, *Five-leaved Ivy*, *Woody Climber*. An indigenous climbing plant. Ord. Vitaceæ; which flowers in July. It has been advised as an expectorant.

AMPELOS, (αμπελος, 'vine,') *Vitis vinifera*—a. Agria, *Bryonia alba*—a. Idæa, *Vaccinium Vitis Idæa*—a. Melæna, *Tamus communis*—a. Oinophorus, *Vitis vinifera*.

AMPHAMPHOTERODIOPSISIA, (αμφι, αμφοτερος, 'on both sides,') διπλος, 'double,' and πτε, 'sight,') Diplopia.

AMPHARISTEROS, *Ambila'trus*, 'awkward,') (αμφιστερος, from αμφ, and αστερος, 'left,') Opposed to ambidexter.

AMPHEMERINUS, (*amphemerinos*, from *amphi*, and *hēra*, 'a day,') Quotidian.

AMPHEMERUS, (*amphemerus*,) Quotidian.

AMPHI, (*amphi*,) both, around, on all sides.' Hence, a prefix in many terms.

AMPHIAM, Common.

AMPHIARTHRŌSIS, (*amphi*, and *arthrosis*, 'articulation,') A mixed articulation, in which the corresponding surfaces of bones are united in an intimate manner by an intermediate body, which allows, however, of some slight motion. Such is the junction of the bodies of the vertebrae by means of the intervertebral cartilages. This articulation has also been called *Diarthrosis de Continuis*. The motion it permits is but slight.

AMPHIBLESTROÏDES, Reticular—a. Membrana, Retina.

AMPHIBLESTROÏDITIS, (*amphiblaston*, 'a covering, a net or net,' *itēs*, 'resemblance,' and *itis*,) Retinitis.

AMPHIBLESTROMALACIA, (*Amphiblastro-donala'cia*, (*amphiblastoides* (membrana), 'the retina,' and *malacia*, 'softening,') Mollescence or softening of the retina.

AMPHIBRANCHIA, (from *amphi*, and *branchia*, 'the gills,') *Amphibronchia*. The tonsils and neighbouring parts.

AMPHID'EUM, (*amphidion*, from *amphi*, and *ēu*, 'I bind,') The outermost margin of the cervix uteri: the *Lobium uteri*.

AMPHIDEXIUS, (*amphidexios*, from *amphi*, and *dexios*, 'right,') Ambliexter.

AMPHIARTHRŌSIS, (from *amphi*, and *arthrosis*, 'a moveable joint,') A name given by Winslow to the temporo-maxillary articulation, because, according to that anatomist, it partakes both of ginglymus and arthrodia.

AMPHIES'MA (gen. *Amphiesmatia*) CORDIS, (*amphiesma*, 'a covering,') Pericardium.

AMPHEMERINA, Pertussis, (*amphi*, and *hēra*, 'a day,')—a. Hæticæ, Hætic fever.

AMHIMERINOS, Quotidian.

AMPHION, Muslach.

AMPHIPLEX, Perineum.

AMPHIPNEUMA, (*amphi*, and *πνεμα*, 'breath,') Dyspnœa.

AMPHISMELA, Knife, double-edged.

AMPHISMILE, (*amphi*, and *σμιλη*, 'a knife,') Knife, double-edged.

AMPHISPHAL'SIS, *Circumductio*, *Circumductio*, (*amphi*, and *σφαλλω*, 'I wander,') The movement of circumduction used in reducing luxations.

AMPHOPLOPIA, (*amphi*, *επιπλος*, 'double,' and *οπτι*, 'sight,') see Diplopia.

AMPHORA, per synchōn for *amphoræ*, (from *amphi*, and *phōra*, 'I bear,' because it had two handles,) A liquid measure among the ancients, containing above seven gallons. Also called *Quadrantal*, *Cœcianum*, *Cœdus*.

AMPHORIC RESPIRATION, see Cavernous Respiration.

AMPHOTERODIPLOPIA, (*amphoteris*, 'on both sides,') see Diplopia.

AMPLEXATIO, (*am*, and *plectere*, 'to interweave,') Coition.

AMPLIFICATIO, (*amplus*, and *facere*, 'to make large,') Platygnosis.

AMPLIOPIA, see Amblyopia.

AMPOSIS, (*ampositio*,) Anaposis.

AMPOULE, Ampulla—a. de Vater, see Vater, *ampulla de*.

AMPOULES, Essera.

AMPULLA [L.], 'a bottle,') (F.) *Ampoule*. A membranous bag, shaped like a cæthern bottle. See *Cæstus* EIDIPLOS. In pharmacy, a receiver.

AMPULLA CHYLIFERA SEU CHYLI, Receptaculum chyli.

AMPULLE, Phlyctenæ.

AMPUTATION, *Amputatio*, (*amputare*, am-

putatum; *am*, 'about,' and *putare*, 'to cut off,') *Amputatio*, *Amputatio*. The operation of separating, by means of a cutting instrument, a limb or a part of a limb, or a projecting part, as the mamma, penis, &c., from the rest of the body. In the case of a tumour, the term *excision*, *removal*, or *extirpation*, (F.) *Resectio*, is more commonly used.

AMPUTATION, CIRCULAR, is that in which the integuments and muscles are divided circularly.

AMPUTATION, FLAP, (F.) *A. à lambeau* is when one or two flaps are left so as to cover the stump when the limb has been removed.

AMPUTATION, JOINT, *Excision*, (F.) *A. dans l'articulation* or *dans la continuité des membres*, is when the limb is removed at an articulation.

Each amputation requires a different process, which is described in works on operative surgery.

AMPUTATION SÈCHE, see *Écrasé*.

AMPUTATION, SPONTANEOUS, see Spontaneous.

AMULET, Amuletum.

AMULETTE, Amuletum.

AMULETUM, (*amulitum*, 'to remove, to avert,') An *Amulet*, *Perian'ma*, *Apotropa'm*, *Periap'ton*, *Phylact'rium*, *Apoteles'ma*, *Evastes'ma*, *Alexis'mum*, *Preservati'um*, *Probascan'tium*, (F.) *Amulette*. Any image or substance worn about the person for the purpose of averting disease or danger.

AMUR'CA, *Amur'ga*, (*amur'ga*, from *amur'ga*, 'I press out,') The *mare* or grounds remaining after olives have been crushed and deprived of their oil. It has been used as an application to ulcers.

AMURGA, (*amur'ga*,) *Amurea*.

AMUSA, *Musa Paradisiaca*.

ANYCHA, (*anyxa*,) *Amur'is*, (*anyxa*, 'I scratch,') Excoriation, Scarification.

AMYCTICA, (from *anyxa*, 'I scratch,') Medicines which stimulate and vellicate the skin.

AMYDOLÉ, (from *amidon*,) *Fecula*.

AMYDRIASIS, *Mydriasis*.

AMYEL'IA, (*α*, and *μυελος*, 'marrow,') A monstrous formation in which there is an absence of spinal marrow.

AMYELONER'VIA, (*α*, *μυελος*, 'marrow,' and *νερα*, 'nerve,') Paralysis or deficient action of the spinal marrow.

AMYELOTROPH'IA, (*α*, *μυελος*, 'marrow,' and *τροφον*, 'nourishment,') Atrophy of the spinal marrow.

AMYGDALA, (*amygdala*,) *Amygdalum*, (*amygdalos*, 'from *amēco*, 'to tear,' owing to its fissured shell [?].) The *Almond*, of which there are two kinds, *Amygdale amara* and *A. dulcis*, (F.) *Amandes amères*, and *A. douces*, obtained from two varieties of *Amygdalus communis* or *A. sativa*, *Almond tree*, (Old Eng.) *Amygdier*, a native of Barbary. *Ord. Amygdaleæ*. *See. Syst. Iconodoria Monogynia*.

The taste of *Amygdala dulcis*, *Jordan Almond*, is soft and sweet; that of *A. amara*, bitter. Both yield, by expression, a sweet, bland oil. The bitter almond contains Prussic acid. They are chiefly used for forming emulsions.

AMYGDALÉ PASTA, *Almond Paste*, a cosmetic for softening the skin and preventing chaps, is made of bitter almonds, blanched, $\frac{3}{4}$ iv, white of one egg; rose water, and rectified spirits, equal parts, or as much as is sufficient.

AMYGDALÉ PLACEN'TA, *Almond Cake*, is the cake left after the expression of the oil. The ground *Almond Cake*, *Almond Powder*, *Farina Amygdalorum*, is used instead of soap for washing the hands.

AMYGDALA, Tonsil. Also, a lobule or prominence of the cerebellum, so called from its resemblance to an enlarged tonsil. This and its fellow of the opposite side form the lateral boundaries of the anterior extremity of the valley, and are in

great part covered by the medulla oblongata. The Amygdalæ are seated on either side of the uvula, in the fourth ventricle.

AMYGDALATOME, *Amygdalatōmus*, (from *amygdala*, and *tomē*, 'incision;') improperly *Tonsillitome*,—*Tonsil guillotine*, (F.) *Scateur des Amygdales*. An instrument of surgery for excising a portion of the tonsil. See *Kiotome*.

AMYGDALATUM, Emulsio amygdalæ.

AMYGDALE, Tonsil.

AMYGDALIN, *Amygdalīnum*, *Amygdalīna*, *Amygdaline*. A principle contained in bitter almonds, which is prepared by pressing the bruised almonds between heated plates to separate the fat oil; boiling the residue in alcohol; evaporating, and treating with ether, which precipitates the amygdalin in a crystalline powder. A weak solution of it, under the influence of a small quantity of *emulsin* or *synaptase*, which constitutes the larger portion of the pulp of almonds, yields at once oil of bitter almonds and hydrocyanic acid.

AMYGDALITIS, (*amygdala*, and *itis*.) Cynanche tonsillaris.

AMYGDALUM, Amygdala.

AMYGDALUS, (*αμυγδαλος*;) see Amygdala—a. Communis, see Amygdala.

AMYGDALUS PER'SICA, *Per'sica vulga'ris*. The common peach tree, (F.) *Pêcher*. The leaves and flowers have been considered laxative. They are bitter and aromatic, and have been given in hæmaturia, nephritis, &c. The fruit—*peach*, (F.) *pêche*—is one of the pleasant and wholesome summer fruits, when ripe. The kernels, *Amygdalæ Per'sicæ*, as well as the flowers, contain prussic acid.

Peach Brandy is distilled from the fruit, and is much used in the United States.

AMYGMOS, (*αμυγμος*, from *αμύσσω*, 'I scratch,') Scarification.

AMYL, Amylum.

AMYLACÉOUS, *Amyla'ceus*, (F.) *Amylace*, (*amylum*, 'starch,') Having the nature of, or containing starch. Starchy, starch-like.

AMYLACEOUS BODIES, Corpora amylacea.

AMYLE, HYDRATED OXIDE OF, Oil, fusel.

AMYLENE, see Potato oil.

AMYLEON, Amylum.

AMYL IODIDUM, Starch, iodide of—a. Ioduretum, Starch, iodide of.

AMYLIER, see Amygdala.

AM'YLOID, *Amyloi'des*, (*amylum*, and *ειδος*, 'resemblance,') Resembling starch. See Glycogenic matter.

AMYLOID BODIES, Corpora amylacea.

AMYLON, Amylum, Glycogenic matter.

AM'YLU, *A'midum*, *Fœcula*, *Amyl'eon*, *Amyl'ion*, *Am'ylon*, (*αμυλον*, from *α*, and *μύλη*, 'a mill,' because made without a mill,) *Starch*, (Old Eng.) *Am'yl*, (F.) *Amidon*, *Amylon*. *Starch of Wheat*, *Fari'na*, *Trit'ici fari'na*, *Am'ylum tritic'eum* seu *Trit'ici*, *Fœcula Amyla'cea*, is inodorous and insipid, white and friable. It is insoluble in cold water and alcohol, but forms with boiling water a strong, semi-transparent jelly. It is demulcent, and is used as an emollient glyster, and as the vehicle for opium, when given *per anum*. Starch is met with abundantly in all the cereal grains, in the stalks of many of the palms, in some lichens, and in many tuberous roots, particularly in the bulbs of the orchis.

AMYLUM AMERICANUM, see Arrow-root—a. Canaceum, *Tous-les-mois*—a. Iodatium, Starch, iodide of—a. Manihotium, see *Jatropha manihot*—a. Marantaceum, Arrow-root—a. Palmaceum, *Sago*—a. Querneum, *Racahout*.

A'MYON, (from *α*, and *μύων*, 'a muscle,') *Emusculatus*. Without muscle. Applied to the limbs, when so extenuated that the muscles cannot be distinguished.

AMYOSTHENIA, (F.) *Amyosthénie*. (*α*, *μύων*, 'a muscle,' and *σθένος*, 'strength,') Defect of muscular contraction.

AMYOSTHÉNIE CYSTURIQUE, Paralysis of the bladder, of Piorry.

AMYRIS COMPIMP'ORA, (*α*, intensive, and *μύρον*, an odoriferous ointment, 'because amyris enters into the composition of such.) See *Bdellium*.

AMYRIS ELEMIF'ERA, (F.) *Balsamier ou Bannier Elemifère*. Ord. Terebinthaceæ. Sec. Syst. Octandria Monogynia. The plant whence it has been supposed GUM EL'EMI is obtained. This gum or resin is brought from the Spanish East and West Indies. *Brazil'ian El'emi*, according to Dr. Royle, is produced by *Ici'ca Ici'ca'riba*; *Mex'ican El'emi*, by *Ela'phrium elemif'erum*; and *Manilla El'emi*, by *Canarium commune*. It is softish, transparent, of a pale whitish colour, inclining a little to green, and of a strong, though not unpleasant smell. It is only used in ointments and plasters, and is a digestive.

AMYRIS GILEADENSIS, see A. opobalsamum.

AMYRIS OPOBAL'SAMUM, (F.) *Balsamier ou Bannier de la Mecque*, *Bal'sam*, *Bal'samum*. The plant from which is obtained the BALSAM OF MEC'CA, *Bal'samum genuinum antiquorum* seu *Asiat'icum* seu *Juda'icum* seu *Syri'acum* seu *e Meccâ* seu *Alpi'ni*, *Balsamel'on*, *Egypti'acum* *Bal'samum*, *Cocobal'samum*, *O'ileum* *Bal'sami*, *Opobal'samum*, *Xylobal'samum*, *Bal'sam* or *Balm of Gil'ead*, (F.) *Baume Blanc*, *B. de Constantino'ple blanc*, *B. de Galaad*, *B. du Grand Caire*, *B. Vrai*, *Térébinthine de Giléad*, *T. d'Égypte*, *T. du Grand Kaïre*, *T. de Judée*. A resinous juice obtained by making incisions into *Am'yris opobal'samum* and *A. Gileaden'sis* of Linnaeus, *Balsamaden'dron Gileaden'sis* of Kunth. The juice of the fruit is called *Carpobal'samum*; that of the wood and branches *Xylobal'samum*. It has the general properties of the milder Terebinthinates.

AMYRIS TOMENTOSA, *Fagara octandra*.

AMYRON, *Carthamus tinctorius*.

A'MYUS, (*α*, and *μύς*, 'a muscle,') Weak or poor in muscle.

AMYX'IA, (*α*, and *μύξα*, 'mucus,') Deficiency of mucus.

AMYXIS, (*αμυξίς*), *Amycha*, Scarification.

AN, as a prefix. See A, Ad, and Am.

ANA, (*ανα*), a word which signifies 'of each.' It is used in prescriptions as well as *ā* and *āā*, its abbreviations. As a prefix, it means 'in,' 'through,' 'upwards,' 'above,' in opposition to *cata*—also, 'again,' 'repetition,' like the English *re*.

ANAB'ASIS, (*αναβασις*, from *αναβαινω*, 'I ascend,') The first period of a disease, or that of increase. See Augmentation.

ANABEXIS, (*ανα*, and *βησσειν*, 'to cough,') Expectoration.

ANABLEP'SIS, (*αναβλεψις*, from *ανα*, and *βλεπω*, 'I see,') Restoration to sight.

ANABOL'ON, (*αναβολαων*), *Anabole'us*, (*αναβαλλω*, 'I cast up,') An ointment for extracting darts or other extraneous bodies.

ANAB'OLE, (*αναβολη*), *Anagol'gē*, *Anaph'ora*, *Anacine'na*, *Anacine'sis*. An evacuation upwards. An act by which certain matters are ejected by the mouth. In common acceptance it includes, *expectation*, *expectoration*, *regurgitation*, and *vomiting*.

ANABROCHIS'MUS, *Anabron'chismus*, (*ανα*, and *βροχος*, 'a running knot,') An operation for removing the eye-lashes, for example, when they irritate the eye, by means of a hair knotted around them.

ANABRONCHISMUS, *Anabrochismus*.

ANABROSIS, (*αναβρωσις*, from *ανα*, and *βρωσκειω*, 'I eat,') Corrosion, Erosion.

ANACAHUITE WOOD, whose botanical origin is not known, but which is supposed to belong to

Nat. Ord. Papilionaceæ; is Mexican. It is inodorous and insipid; but a decoction of it has been inordinately extolled in the first stages of phthisis.

ANACAMPSEROS, (*ανακαμψερως*), *Sedum telephium*.

ANACARDIUM OCCIDENTALE, (*ανα*, and *καρδία*, 'heart,' from the resemblance of the fruit to a dried heart,) *Acaju'ba occidentalis*, *Cassauvium pomif'erum*, *Cashew* (*W. Indies*.) (F.) *Ac'ajon*. *Ord.* Terebinthaceæ. *Sex. Syst.* Enneandria Monogynia. The Oil of the *Cashew Nut*, *O'leum Anacardi'ii*, *Cardo'leum*, (F.) *Huile d'Ac'ajon*, is an active caustic, and used as such in the countries where it grows, especially for destroying warts, &c. A gum resembling gum Arabic, and called *Cashew gum*, exudes from the bark.

ANACARDIUM ORIENTALE, *Avicennia tomentosa*.

ANACATHARSIS, (*ανακαθαρσις*, from *ανα*, and *καθαρειν*, 'to purge,') Purgation upwards. Expectoration. See, also, *Repurgatio*.

ANACATHARSIS CATARRHALIS SIMPLEX, *Catarrh*.

ANACATHARTICUS, (*ανακαθαρτικός*), Expectorant.

ANACESTOS, (*ανακεστος*, from *αν*, and *ακεομαι*, 'to cure,') Incurable.

ANACHREMPISIS, (*ανα*, and *χερμεισις*, 'spitting,') Expulsion.

ANACHREMPUTUM, Sputum.

ANACHRON, Soda.

ANACINEMA, gen. *Anacine'matis*, (*ανακίνημα*, from *ανα*, and *κινω*, 'I move,') Anabole, Expulsion.

ANACINESIS, (*ανακίνησις*), Anabole, Expulsion.

ANACLASIS, (*ανακλασις*, 'a bending or breaking upwards or backwards,') Repercussion.

ANACLINTERIUM, (*ανακλιντήριον*), *Anaclin'trum*, *Recubito'rium*, (*ανακλινω*, 'I recline,') A long chair or seat, so formed that the person can rest in a reclining posture.

ANACLINTRUM, (*ανακλιντρον*), *Anaclinterium*.

ANACOLLEMA, gen. *Anacolle'matis*, (*ανακολλημα*, from *ανα*, and *κολλω*, 'I glue,') A healing medicine.

ANACOLLEMATA, Frontal bandages.

ANACOLUP'PA. A creeping plant of Malabar, the juice of which, mixed with powdered pepper, passes in India as a cure for epilepsy, and as the only remedy for the bite of the naja. It is supposed to be *Zizipha'nia nodiflora*.

ANACOLUTHIE, (*αν*, and *ακολουθος*, 'consequent,') Incoherence.

ANACOMIDE, (*ανακομιδή*, from *ανα*, and *κομιζω*, 'I bring,') Restaura'tio.

ANACONCHYLASMUS, (*ανακογχυλιασμος*, from *ανα*, and *κογχυλιαζειν*, 'to gargle,') Gargarism.

ANACONCHYLISMUS, Gargarism.

ANACTESIS, (*ανακτησις*, from *ανα*, and *κταμαι*, 'to acquire,') Restaura'tio.

ANACTIRION, *Artemisia*.

ANACYCLEON, (*ανα*, and *κυκλω*, 'I go in a circle,') Charlatan.

ANACYCLUS OFFICINARUM, see *Anthemis Pyrethrum*—a. *Pyrethrum*, *Anthemis pyrethrum*.

ANADESMUS, (*αναδεσμος*, from *ανα*, and *δεσμος*, a bandage,') Fascia.

ANADIPLO'SIS, (*αναδιπλωσις*, from *ανα*, and *διπλω*, 'I double,') *Epanadiplo'sis*, *Epanalep'sis*, *Reduplicatio*. The redoubling which occurs in a paroxysm of an intermittent, when its type is double.

ANADORA, (*αναδωρα*), *Eedora*.

ANAD'OSIS, (*αναδωσις*, from *αναδιδωμι*, 'I give up,') Purgation upwards, as by vomiting. Congestion of blood towards the upper parts of the body. *Anadosis* seems also to have occasionally

meant chylification, whilst *diadosis* meant capillary nutrition.

ANAD'ROMÊ, (*αναδρουν*, from *ανα*, and *δρυνω*, 'I run,') The transport of a humour or pain from a lower to an upper part. Also, the globus hystericus. See *Angone*.

ANÆDE'US, (*αν*, and *αἷδα*, 'organs of generation,') A monster devoid of sexual organs.

ANÆMATOPOE'SIS, (*αν*, 'αἷμα', 'blood,' and *ποιω*, 'I make,') Impeded or obstructed hæmatisation.

ANÆMATO'SIS, *Anhæmato'sis*, (*αν*, 'αἷμα', 'blood,' and *osis*.) Defective hæmatisation or preparation of the blood. *Anæmia*.

ANÆMIA, *Exæ'mia*, *Anæ'masis*, *Anhæ'mia*, *Anhæmato'sis*, *Polyanhæ'mia*, *Anæmo'sis*, *Oligæ'mia*, *Oligohæ'mia*, *Hyperæ'mia*, *Hydroæ'mia*, *Hydræ'mia*, *Anæ'mia*, (F.) *Anémie*, *Anhémie*, *Anhématosie*, *Polyanhémie*, *Hydrohémie*, *Exsanguin'ity*, *Blood'lessness*, (*αναἷμα*, from *αν*, and *αἷμα*, 'blood,') Privation of blood;—the opposite to *plethora*. It is characterized by every sign of debility. Also, diminished quantity of fluids in the capillary vessels:—the opposite to *Hyperæmia*.—The essential character of the blood in *anæmia* is diminution in the ratio of red corpuscles, (F.) *Aglobulie*.

ANÆMIC, *Anæ'mic*, *Anæ'micus*. Appertaining or relating to *anæmia*,—as an "anæmic person." See *Exsanguinus*.

ANÆMOCH'RŌUS, (*αν*, 'αἷμα', 'blood,' and *χρῶα*, 'colour,') Devoid of colour, pale.

ANÆMOSIS, *Anæmia*.

ANÆMOT'ROPHY, *Anæmotroph'ia*, (*αν*, 'αἷμα', 'blood,' and *τροφή*, 'nourishment,') A deficiency of sanguineous nourishment.

ANÆMYDRIA, (*αν*, 'αἷμα', 'blood,' and *ἰδωρ*, 'water,') *Anhydræmia*.

ANÆSTHE'SIA, *Anæsthe'sis*, *Insensibil'itas*, *Analgæ'sia*, *Parop'sis exers*, (F.) *Anesthésie*; (*αναίσθησια*, from *αν*, and *αἰσθάνομαι*, 'I feel,') Privation of sensation, and especially of that of touch according to some. It may be general or partial, and is almost always symptomatic.

ANÆSTHESIA GUSTATORIA, *Agustia*—a. *Ingustæ*, *Agustia*—a. *Olfactoria*, *Anosmia*—a. *Optic*, *Amâusis*.

ANÆSTHESIS, *Anæsthesia*.

ANÆSTHET'IC, *Anæsthet'ic*, *Anæsthet'icus*, *Anæsthet'us*, (F.) *Anesthésique*. Relating to privation of feeling, as an "anæsthetic agent;" one that prevents feeling. The term is, now, almost restricted to agents, which produce such effect by being received into the lungs in the form of vapours or gases, and passing with the blood to the nervous centres on which their action is exerted. Perhaps, as a general rule, the intellectual faculties first feel their influence,—a sort of intoxication supervening, with imperfect power of regulating the movements; the sensory ganglia become afterwards or simultaneously affected, sensation and motion are suspended, and ultimately, if the quantity inhaled be sufficient, the medulla oblongata has its actions suspended or destroyed, respiration ceases, and death is the consequence. Different agents have been used as anæsthetics by way of inhalation—sulphuric ether, chloroform, chloric ether, compound ether, chlorohydric and nitric ethers, bisulphure of carbon, chloride of olefant gas, benzine, aldehyde, light coal-tar naphtha, &c.; but the first four are alone employed. They have been, and are, greatly used in serious surgical operations, and during parturition; and in such cases, as well as in many diseases, especially of a painful nature, produce the most beneficial results.

ANÆSTHETIZATION, *Anæsthetization*, (F.) *Anesthésisation*. The condition of the nervous system induced by anæsthetics. To *an'æstheize*, or *an'esthetize*, (F.) *anesthésier*, is to produce such a condition.

ANÆSTHETUS, (αναίσθητος.) Anæsthetic.

ANAGAL'LIS, (αναγαλλίς, from *ana*, and *gala*, 'milk,' from its power of coagulating milk.) *A. arven'sis*, *A. Phœnic'lea*, *Red Pim'pernel*, *Scarlet Pim'pernel*, *Shep'herd's Sand'al*, *Chickweed*. *Nat. Ord.* Primulaceæ. *Sex. Syst.* Pentandria Monogynia. (F.) *Mouvon rouge*. A common European plant; a reputed antispasmodic and stomachic. A decoction of this in beer is said to form an important part of *Stoy's medicine for hydro-phobia*.

Another species—*Anagal'lis cœru'lea* is a mere variety of the above.

ANAGALLIS AQUATICA, *Veronica Beceabunga*.

ANAGARGALICTON, (αναγαργαλίκτον, from *αναγαργαριζω*, 'to gargle,') Gargarism.

ANAGARGARISMUS, Gargarism.

ANAGARGARISTON, (αναγαργαρίστον,) Gargarism.

ANAGENNE'SIS, (αναγήνυσις, (*ana*, and *γενναω*, 'I generate,') Regeneration; growing again.

ANAGLYPHE, (αναγλύφη, from *ana*, and *γλυφω*, 'I carve,') *Calamus scriptorius*.

ANAGOSTAKIS, (after the Greek physician, who invented it.) See *Ophthalmoscope*.

ANAGOGE, (αναγωγή, from *ana*, and *αγω*, 'I lead,') *Anabole*, *Rejection*.

ANAGRAPHÉ, (αναγραφή, from *ana*, and *γραφω*, 'I write,') *Prescription*.

ANAG'YRIS, (αναγυρίς,) *Anag'yris foetida*, *Anag'yris*, *Ac'orpon*, *Stink'ing Bean Trefoil*. (F.) *Anagyris*. *Ord.* Leguminosæ. Native of Italy. The leaves are powerfully purgative. The juice is said to be diuretic, and the seeds emetic.

ANAGYRUS, (αναγυρίς,) *Anagyris*.

A'NAL, *Ana'lis*. That which refers to the anus;—*Anal region*; *Anal fistula*; *Anal hairs*—*Pi'li ana'les*, &c.

ANAL'DIA, (F.) *Analdie*; (*an*, and *αλδεν*, 'to grow,') *Defective nutrition*.

ANALEMSIA, *Analepsia*.

ANALENTIA, *Analepsia*.

ANALEP'SIA, (αναληψίς,) *Analep'sis*, *Epanalepsis*, *Analen'tia*, *Analem'sis*, (*ana*, and *λαμβάνειν*, (future *λησεται*, 'to take,') *Restoration to strength after disease*. A kind of sympathetic epilepsy, originating from gastric disorder. See *Epilepsy*. Also, the support given to a fractured extremity.

ANALEPSIS, (αναληψίς,) *Convalescence*, *Restauratio*.

ANALEP'TICA, (αναληπτικά,) *Anapys'e'tica*, *Psychot'ica*, *Refect'iva*, *Refect'iva*, *Restauran'tia*, *Analep'tics*. *Restorative medicines or food*; such as are adapted to recruit the strength during convalescence.

ANALEP'TIC PILLS, JAMES'S, consist of *James's Powder*, *Gum Ammoniacum*, and *Pills of Aloës and Myrrh*, equal parts, with *Tincture of Castor*, sufficient to form a mass.

ANALGE'SIA, (αναλγησία,) *Ana'gia*, (*an*, and *αλγος*, 'pain,') *Absence of pain both in health and disease*. See *Anæsthesia*.

ANALGIA, *Analgesia*.

ANALOGOUS TISSUES, see *Tissues*.

AN'ALOGUE, *Analogus*; (*ana*, and *λογος*, 'proportion,') A part in one organized being which has the same function as another part in another organized being.

ANALOSIS, (αναλωσις,) *Atrophy*.

ANALTHES, (αναλθής,) *Incurable*.

ANAMIRTA COCCULUS, *Menispermum cocculus*—*a. Peniculata*, *Menispermum cocculus*.

ANAMNES'TIC, (αναμνηστικόν,) *Anamnes'ticum*, (*ana*, and *μνησκω*, 'I remember,') A medicine for improving the memory. See, also, *Commemorative*.

ANANAS, *Bromelia ananas*—*a. Aculeata*, *Bromelia ananas*—*a. Americana*, *Bromelia pinguin*—*a. Ovata*, *Bromelia ananas*—*a. Wild*, broad-eaved, *Bromelia pinguin*.

ANANAZIP'TA. A word formerly scrawled on amulets to charm away disease.

ANANDRI'A, (αναδρία, from *an*, and *ανηρ*, gen. *ανδρος*, 'a man,') *Want of manliness*. *Impotence in the male*. The state and act of emasculation.

ANANEO'SIS, (αναανεωσις, from *ana*, *νεος*, 'new,' and *οσις*,) *Renova'tio*. *Renovation or renewal*,—as of the blood by the chyliferous vessels and lymphatics.

ANAPETI'A, (αναπετεια,) *Expan'sio med'tuum*, (*ana*, and *πετω*, 'I dilate,') A state opposite to that of the closure of vessels.

ANAPHALANTIASIS, (αναφαλαντίασις,) *Anaphalanto'ma*, (*ana*, and *φαλαντος*, 'bald,') *Loss of the hair of the eyebrows*. Also, *balddness in general*.

ANAPHALANTOMA, gen. *Anaphalanto'matis*, (*αναφαλαντωμα*, from *ana*, and *φαλαντος*, 'bald,') *Anaphalantiasis*.

ANAPHE, *Anaphia*.

ANAPH'IA, *Anaph'ia*, *An'aphē*, (*an*, and 'αφη, 'touch,') *Diminution or privation of the sense of touch*.

ANAPHLASMUS, (αναφλασμος, from *αναφλω*, 'I produce erection,') *Masturbation*.

ANAPHONE'SIS, (αναφωνησις,) (*ana*, and *φωνη*, 'voice,') *Exercise of the voice: vociferation:—the act of crying out; Vocifera'tio*, *Clamor*.

ANAPHORA, (αναφορά, from *ana*, and *φέρω*, 'to carry,') *Anabole*.

ANAPHRODIS'IA, (αναφροδισία, from *an*, and *αφροδιτις*, 'Venus,') *Defec'tus Ven'erie*. *Absence of the venereal appetite*. Sometimes used for *Impotence and Sterility*.

ANAPHRODISIAC, *Antaphrodisiac*.

ANAPHROMELI, (*an*, *αφρος*, 'froth,' and *μελι*, 'honey,') *Mel despumatum*.

ANAPHYSIS, (*ana*, and *φνω*, 'to produce,') *Anagennesis*.

ANAP'LASIS, *Anaplas'mus*, (*αναπλασις*, from *αναπλασσω*, 'I restore,') *Confirma'tio*, *Reposit'io*. *Restoration*. Union or consolidation of a fractured bone.

ANAPLASMATIC, *Anaplastic*.

ANAPLASMUS, (αναπλασμος,) *Anaplasia*.

ANAPLAS'TIC, *Anaplas'ticus*. An epithet applied to the art of restoring lost parts or the normal shape,—as '*Anaplastic Surgery*.' See *Morioplastic*. Also, an agent, that increases the amount of plastic matter—fibrin—in the blood; *Anaplas-mat'ic*.

ANAPLERO'SIS, (αναπληρωσις, from *ana*, *πληρω*, 'I fill,' and *οσις*,) *Repletion*. That part of surgical therapeutics whose object is to supply parts that are wanting. Also, *Apposition* or *Prosthesis*.

ANAPLEROTICUS, *Incarnans*.

ANAPLEU'SIS, (αναπλευσις,) *Fluctua'tio*, *Inna'tio*, (from *αναπλω*, (*ana*, and *πλω*), 'to swim above,') The looseness or shaking of an exfoliated bone; or of a carious or other tooth, &c.

ANAPLOSIS, (αναπλωσις, from *ana*, *πλω*, 'I unfold,' and *οσις*,) *Growth*.

ANAPNEUSIS, (αναπνευσις,) (*ana*, and *πνω*, 'I breathe,') *Respiration*.

ANAPNOË, (αναπνοή,) *Respiration*.

ANAPNOË'USI, (αναπνοή, and *νους*, 'dis-ease,') *Diseases of the respiratory organs*.

ANAPNOMETER, (αναπνομε, and *μετρον*, 'mea-sure,') *Spigrometer*.

ANAPODISIS UTERI, (αναποδισις, from *αναποδίζω*, 'I go back,') *Retroversio Uteri*.

ANAPODISMUS UTERI, (αναποδισμος,) *Retroversio Uteri*.

ANAPODOPHYLLUM CANADENSE, *Podophyllum peltatum*.

ANAP'OSIS, *Am'posis*, (*ana*, and *ποσις*, 'drink,')

A recession of humours from the circumference to the centre of the body.

ANAPSYCTICA, (αναψυχω, 'I refresh.') Analeptica.

ANAPTYSIS, (ανα, and πτω, 'I spit.') Expectoration.

ANAPTYXIS, (αναπτύξις, from αναπτύσσω, 'I unfold.') Growth.

ANARCOTINA, Narcotine.

ANARRHEGNU'MINA, (αναρήγνυμι, 'I break out again.') Fractures are so called when they become disunited; as well as ulcers when they break out afresh.

ANARRHINON, (ανα, and 'ριν, 'the nose.') That which returns by the nose.

According to others, that which issues by the skin: (from ανα, and 'αυς, 'the skin.')

ANARRHINUM, Sternutatory.

ANARRHŒA, (Anarrhoë, Anarrhopia, Anas'tasis, (ανα, and 'ρω, 'I flow.') Afflux of fluid towards the upper part of the body.

ANARRHOPHE, (αναρήρω, 'I suck in.') Absorption.

ANARRHOPHENU'SI, (anarrhophe, and νου-σος, 'disease.') Diseases of the absorbents.

ANARRHOPHESIS, (αναρήρησις,) Absorption.

ANARRHOPIA, (αναρήπια, from ανα, and 'ρω, 'I incline.') Anarrhea.

ANAR'THRUS, (αναρθρος,) (from αν, and αρθρον, 'a joint.') Without a joint. One who is so fat that his joints are scarcely perceptible.

ANASAR'CA, (ανα, 'through,' and σαφ, gen. σαφω, 'the flesh.') Catasar'ca. A qua inter'cus seu in'ter cut'em, Hyposar'ca, Hy'drops cellula'ris to'tus cor'poris, H. Anasar'ca seu inter'cus seu sub-cut'a'neus seu cellulo'us seu cut'a'neus seu te'l'a cellulo'sa, Katasar'ca, Episar'ceid'ium, Hy'deros, Hydaton'cus, Hyderon'cus, Hydron'cus, Hydrosar'ca, Hydroder'ma, Hydrop'icis ve'ca, Sarcot'es, Polygm'phia, Hyposarceid'ius, Leucophlegmat'ia, Gen'eral drop'sy, Drop'sy of the cellular membrane, (F.) Anasarque. Commonly, it begins to manifest itself by swelling around the ankles; and is characterized by tumefaction of the limbs and of the soft parts covering the abdomen, thorax, and even the face, with paleness and dryness of the skin, and pitting when any of these (especially the ankles) are pressed upon. Like dropsy in general, Anasarca may be active or passive; and its treatment must be regulated by the rules that are applicable to dropsy in general. At times, the symptoms are of an acute character, and the effusion sudden, constituting Dermatoc'h'y'sis, Hy'drops Anasar'ca acu'tus, Ede'ma cal'idum, E. acu'tum, E. febr'il'e of some. See Hydrops.

ANASARCA HYSTERICUM, Anathymiasis—a. Pulmonum, Hydropneumonia, E'dema of the Lungs—a. Serosa, Phlegmatia dolens.

ANASARQUE, Anasarca.

ANASISMUS, (ανασισμος, from ανα, and σειω, 'I shake.') Concussion.

ANASPADIA, see Anaspadiaens.

ANASPA'DIAS, Epispad'ias, (ανα, and σπau, 'I draw.') One whose urethra opens on the upper surface of the penis.

ANASPADISIS, see Anaspadiaens.

ANASPADISMUS, see Anaspadiaens.

ANAS'PASIS, Anaspasm'us, (ανασπασις, from ανα, and σπau, 'I draw.') Retract'io. Contraction, especially of the bowels. The condition is called Anaspa'dia, Anaspa'd'isis, and Anaspa'dis'mus.

ANASPASMUS, Anaspasis.

ANASSA, Bromelia ananas.

ANASTALITICA, (ανα, and στελλω, 'I contract.') Styptics.

ANASTASIS, (αναστασις,) Anarrhœa. Also, restoration from sickness. Convalescence; (from ανίστημι, 'I rise up.')

ANASTŒCHEIO'SIS, (ανα, and στοιχειω, 'element.') Re'elementa'tio. Resolution of a body or its parts into their elements.

ANASTOMO'SIS, (αναστομωσις, from ανα, στομα, 'a mouth,' and οεις,) Inosculat'io seu Res'u'tio vaso'rum, Exanastomo'sis, Concu'r'ens, Inoscula'tion, (F.) Anastomose, Abouchement. Communication between two vessels. By considering the nerves to be channels, in which a nervous fluid circulates, their communication, likewise, has been called Anastomosis. By means of anastomoses, if the course of a fluid be arrested in one vessel, it can proceed along others.

ANASTOMOSIS ANEURISMATICA, Telangiectasia—a. Jacobson's. see Petrosal ganglion.

ANASTOMOT'IC, (αναστομωτικός,) Anastomo'ticus, (F.) Anatomotique. Belonging or relating to anastomosis.

ANASTOMOT'ICS, Anastomo'tica. Certain medicines were formerly so called, which were believed to be capable of opening the mouths of vessels, as aperients, diuretics, etc.

ANASTOMOTICUS MAG'NUS (RA'MUS), (F.) Artère collatérale interne, A. collatérale du coude, is a branch of the brachial artery, which comes off a little above the elbow, and bestows branches to the brachialis internus, to the under edge of the triceps, and to the muscles, ligaments, etc., about the elbow joint. See, also, Articular arteries of the knee.

ANASTROPHE UTERI, (αναστροφή, from ανα and στρεφω, 'I turn,') Inversio uteri.

ANATASIS, (αναστασις, from ανα, and τεινω, 'I stretch,') Extension.

ANATHEMA, gen. Anathe'matis, (αναθημα, from ανα, and τιθημι, 'to put,') Tabula votiva.

ANATHERUM MURICATUM, Andropogon muricatus.

ANATHOMIA, Anatomy.

ANATHEREPSIS, (αναθερψις, from ανα, and τρεφω, 'I nourish,')

ANATHYMIAMA, gen. Anathymia'matie, (ανασθυμα,.) Anathymiasis.

ANATHYMIASIS, (ανασθυμιασις,) Anathymi'a'ma, (ανα, and θυμα, 'fumigation,') Ede'ma fu'gaz, Phygged'e'ma, Ede'ma epus'ticum, Ede'ma hys'tericum, Anasar'ca hys'tericum. An uncertain and transient swelling or inflation, said to have been observed at times in nervous and hysterical persons. It also means Exhalation, Fumigation, and Hypochondriasis.

ANATOLE UNGUIUM, (ανατολη, from ανα-τελλω, 'I come forth,') See Nail.

ANATOME, (ανatomy,.) Anatomy—a. Animata, Physiology.

ANATOMIA, Anatomy—a. Animalis, Zootomy—a. Comparata, Zootomy—a. Comparativa, Zootomy—a. Viva, Physiology.

ANATOMICAL, Anatom'icus, (F.) Anatomique; same etymon as Anatomy. Relating or appertaining to anatomy.

ANATOMIE, Anatomy—a. Chirurgicale, see Anatomy—a. Clastique, see Anatomy—a. des Régions. see Anatomy.

ANATOMIQUE, Anatomical.

ANATOMISM, Anatomism'us. The doctrine of those who look into the arrangement of parts, to explain all the phenomena of the organism.

ANATOMIST, Anatom'icus. One who occupies himself with anatomy. One versed in anatomy.

ANAT'OMY, (ανatomy,) Anat'omē, Anatom'i'a. Anathom'ia, Prosect'io, (from ανα, and τεινειν, 'to cut,') (F.) Anatomie. The word Anatomy properly signifies dissection; but it has been appropriated to the study and knowledge of the number, shape, situation, structure, and connection—in a word, of all the apparent properties of organized bodies. Anatomy is the science of organization. Some have given the term a still more extended accep-

tation, applying it to every mechanical decomposition, even of inorganic bodies. Thus *Crystallology* has been termed the *Anatomy* of crystallized minerals. *Anatomy* has also been called *Morphology*, *Samat'ology*, *Samat'omy*, *Orgonol'ogy*, etc. It assumes different names, according as the study is confined to one organized being, or to a species or class of beings. Thus, *Androt'omy*, or *Anthropot'omy*, or *Anthropog'raphy*, or *Anthroposomatol'ogy*, is the *Anatomy of Man*; *Zoöl'omy*, that of the other species of the animal kingdom; and *Vet'inary Anat'omy* is the anatomy of domestic animals; but when the word is used abstractly, it means *Human Anatomy*, and particularly the study of the organs in a physiological or healthy state. *Physiolog'ical Anatomy* is occasionally used to signify the kind of anatomy which investigates structure with a special view to function. The *Anatomy* of the diseased human body is called *Patholog'ical* or *Morbid Anatomy*, and when applied to Medical Jurisprudence, *Foren'sic Anatomy*. Several of the organs possessing a similarity of structure, and being formed of the same tissues, they have been grouped into *Systems* or *Genera of Organs*; and the study of, or acquaintance with, such systems, has been called *General Anat'omy*, *Histol'ogy*, or *Morphol'omy*, whilst the study of each organ in particular has been termed *Descriptive* or *Spec'ial Anatomy*, *Anthropomorphol'ogy*. *Histol'ogy* is however, more frequently applied to the *Anatomy of the Tissues*, which is called, also, *Textural and Microscop'ic Anatomy*, *Micranatom'ia*, see *Histol'ogy*. *Descriptive Anatomy* has been divided into *Skeletal'ogy*, which comprises *Osteol'ogy* and *Syn-desmol'ogy*; and *Sarcol'ogy*, which is subdivided into *Myol'ogy*, *Neuro'ogy*, *Angiol'ogy*, *Adenol'ogy*, *Splanchnol'ogy*, and *Dermol'ogy*. *Surgical Anat'omy*, *Medicochirurgical Anat'omy*, *Topograph'ical Anat'omy*, *Re'gional Anat'omy*, (F.) *Anatomie Chirurgicale*, *A. des Régions*, is the particular and relative study of the bones, muscles, nerves, vessels, etc., with which it is indispensable to be acquainted before performing operations. *Comparative* or *Analog'ical Anatomy* is the comparative study of each organ, with a view to an acquaintance with the modifications of its structure in different animals, or in the different classes of animals; whilst *Homolog'ical Anatomy* examines the relations to each other of the different parts of the same individual. *Transcendent'ial* or *Philosoph'ical Anatomy* inquires into the mode, plan, or model upon which the animal frame or organs are formed: *Artific'ial Anat'omy* is the art of modelling and representing, in wax or other substance, the different organs or different parts, of the human body, in the sound or diseased state; and *Clas'tic Anat'omy* (F.) *Anatomie Classique*, (κλασ, 'to break,') in the language of Auzous, is the formation of models in such sort, that they can be broken or separated into pieces. *Phytol'omy* and *Phytanat'omy* is the anatomy of vegetables, and *Pictorial Anatomy*, anatomy artistically illustrated.

ANATOMY, see **Skeleton**—*a.* **Analogical**, see **Anatomy**—*a.* **Artificial**, see **Anatomy**—*a.* **Clastic**, see **Anatomy**—*a.* **Comparative**, see **Anatomy**, **Zoatomy**—*a.* **Descriptive**, see **Anatomy**—*a.* **Forensic**, see **Anatomy**—*a.* **General**, see **Anatomy**—*a.* **Homological**, see **Anatomy**—*a.* **Human**, see **Anatomy**—*a.* of **Man**, see **Anatomy**—*a.* **Medicochirurgical**, see **Anatomy**—*a.* **Microscopic**, see **Anatomy**—*a.* **Morbid**, see **Anatomy**—*a.* **Pathological**, see **Anatomy**—*a.* **Pathological, microscopic**, see **Histology**—*a.* **Philosophical**, see **Anatomy**—*a.* **Physiological**, see **Anatomy**—*a.* **Pictorial**, see **Anatomy**—*a.* **Practical**, see **Dissection**—*a.* **Regional**, see **Anatomy**—*a.* **Special**, see **Anatomy**—*a.* **Surgical**, see **Anatomy**—*a.* **Textural**, see

Anatomy—*a.* **Topographical**, see **Anatomy**—*a.* **Transcendental**, see **Anatomy**—*a.* **Veterinary**, see **Anatomy**.

ANATRESIS, (αναρρησις, from *ana*, and *τρεω*, 'I bore,') Perforation, Trepanning.

ANATRIIBE, (*ana*, and *τριβω*, 'I rub,') Friction.

ANATRIPSIS, (αντριψις,) Friction.

ANATRIPSOL'OGY, *Anatripsolog'ia*, *Anatriptolog'ia*, (*anatripsis*, and *λογος*, 'a discourse,')

A treatise on friction as a remedy.

ANATRIPTOLOGIA, *Anatripsology*.

ANATRON, *Natron*, Soda.

ANA'TROPÊ, (αναρροπη, from *ana*, and *τρεπω*, 'I turn,') Subversion. A turning or subversion or inverted action of the stomach, characterized by nausea, vomiting, etc. We still speak of the stomach turning against any thing.

ANAUDIA, (αυαυδία, from *an*, and *αυή*, 'speech,') Catalepsy, Mutitas.

ANAXYRIS, *Rumex acetosa*.

ANAZESIS, (αναζεις, from *ana*, and *ζω*, 'I boil,') Ebullition.

ANAZOTIQUE, see **Nitrogenized**.

ANAZOTURIA, (*an*, *azotum*, and *ουρον*, 'urine,') see **Urine**.

ANCHA, Haunch.

ANCHILOPS, (αγκιλωψ, from *αγκι*, 'near,' and *ωψ*, 'the eye,') *Ægilops*.

ANCHONE, (αγκωνη, from *αγκω*, 'I strangle,') *Angone*.

ANCHORALIS PROCESSUS, (*anchora*, 'an anchor,') *Coracoid*.

ANCHUSA ANGUSTIFOLIA, (αγκυσουα,) *A. Officialis*—*a.* *Incarnata*, *A. Officialis*—*a.* *Lycopersoides*, *A. Officialis*.

ANCHUSA OFFICIALIS, *A. Angustifolia* seu *Incarnata* seu *Lycopersoides*, *Euch'usa*, *Alcan'ua*, *Lit'gua Bo'via*, *Buglos'um sylvest're*, *Offic'inal* or *Garden Al'kanet* or *Bu'gloss*, (Old. Eng.) *Langbeef*; *Ord.* *Boraginæ*. *Sex. Syst.* *Pentandria Monogynia*. (F.) *Buglose*. A native of Great Britain. The herb was formerly esteemed as a cordial in melancholia and hypochondriasis; but it is now rarely used. It is also called *Buglos'ea*, *Buglos'um angustifolium majus*, *B. vulgar'ē majus*, *B. sativum*.

ANCHUSA TINCTORIA, *Alcan'ua spur'ia*, *Dy'er's Bu'gloss*, *Ane'bium*, *Buglos'um Tincto'rum*, *Lithospermum villo'sum*, *Dy'er's Al'kanet*, *Orchanet*, (F.) *Orcanette*. A European plant. The medical properties are equivocal. It is used to give a beautiful red colour to ointments.

ANCHYLOSIS, *Ankylosis*.

ANCISTRON, (αγκιστρον,) *Hamulus*.

ANCLE, (from *αγκλος*, 'crooked,') *Astragalus*, *Malleolus*.

ANCLER, *Malleolus*.

ANCLER, *Malleolus*.

ANCLIFF, *Malleolus*.

ANCLOWE, *Malleolus*.

ANCOLIE, *Aquilegia vulgaris*.

ANCON, (αγκων,) *Elbow*, *Olecranon*.

ANCONAD, see **Anconal Aspect**.

ANCONAGRA, (*ancon*, and *αγρα*, 'a seizure,') *Pechyagra*.

ANCONAL or **ANCONAL**, (*ancon*,) *Relating*, or appertaining to the elbow or olecranon.

ANCONAL ASPECT. An aspect towards the side on which the ancon or elbow is situated.—*Barclay*. *An'conad* is used by the same writer adverbially, to signify 'towards the anconal aspect.'

ANCONÊ, *Anconeus*.

ANCONÊUS, (*ancon*,) A term once applied to every muscle attached to the olecranon. Winslow distinguished four—the *great*, *external*, *internal*, and *small*; the first three being portions of the same muscle, the *triceps brachialis*. The last has, alone, retained the name. It is the *Ancone'us minor* of Winslow, the *Ancone'us* seu *Cubitalis*

Riolani of Douglas, the *Epicandylonebitalis* of Chausser, the *Brevis Cubiti*, (F.) *Ancus*, and is situate at the upper and back part of the forearm. It arises from the external condyle of the os humeri, and is inserted into the posterior edge of the upper third of the ulna. Its use is to aid in the extension of the forearm.

ANCONEUS EXTERNUS, see Triceps extensor cubiti—a. Internus, see Triceps extensor cubiti—a. Major, see Triceps extensor cubiti.

ANCTERES, (αγκυρες, pl. of αγκυρα, 'a clasp, from αγκω, 'I strangle.') *Filulae* or *Clasps* by which the lips of wounds were formerly kept together.

ANCTERIASMUS, (αγκυριασμος.) Infibulation. ANCU'BITUS, *Petrifac'tio*. An affection of the eye, in which there is a sensation as if sand were irritating the organ.

ANCUNULEN'TÆ. A name formerly given to menstruating females.

AN'CUS, *An'kus*, (αγκος, from αγκων, 'the elbow.') One who cannot extend his arms completely.

Also, the deformity resulting from a luxation of the humerus or fore-arm.

ANCYLE. Ankylosis.

ANCYLEN'TERUM, (from *aneylo*, and *εντερον*, 'an intestine.') Adhesion or growing together of the intestines.

ANCYLO, (from αγκυλη, 'a curve,' and hence a stiff joint,—*ankylosis*.) In composition, a curved condition: and a growing together.

ANCYLOBLEPHARON, (βλεφαρον, 'eyelid,') Ankyloblepharon.

ANCYLOCHEILIA, Ankylocheilia.

ANCYLOCOLPUS, (κολπος, 'vagina,') Colpatresia.

ANCYLOCORE, Ankylocore.

ANCYLODERE, (δερε, 'neck,') Torticollis.

ANCYLODERIS, Torticollis.

ANCYLODONTIA, Ankylodontia.

ANCYLOGLOSSIA, Ankyloglossia.

ANCYLOMBLE, Ankylomele.

ANCYLOMERISMUS, Ankylomerismus.

ANCYLORRHINIA, Ankylorrhinia.

ANCYLOSIS, Ankylosis.

ANCYLOTIA, Ankylotia.

ANCYLOTOMUS, Ankylotomus.

ANCYRA, (αγκυρα,) Hook.

ANCYROID CAVITY, (from *ancyra*, and *αιδος*, 'shape,') Digital cavity.

ANCYROIDES PROCESSUS, Coracoid.

AND, Breath.

AN'DA. Native name of a tree of Brazil—*Anda Gome'si* seu *Brasilien'sis* seu de *Pison*, *Andas'su*, *Anda-aqu*, *Juane'sia* princeps. Ord. Euphorbiaceae. *Sex. Syst.* Monœcia Monadelphica. An oil—*Oil of An'da*—is obtained from the seeds by pressure, 50 to 60 drops of which act as a cathartic. The fruit is an oval nut, containing two seeds. These have the taste of the chestnut; but are strongly cathartic, and even emetic. The shell is astringent, and is used as such in diarrhoea, etc.

ANDE. Breath. ANDELY, MIN'ERAL WATERS OF. Andely is in France, near Gisors, and eight leagues from Rouen. The water is cold, and a weak chalybeate. It is used in chlorosis and abdominal obstructions.

ANDERSON'S PILLS, see *Pilulae Aloes et Jalapae*.

ANDIRA IBAI, (Brazilian name,) *Geoffræa vermifuga*—a. *Inermis*, *Geoffræa inermis*—a. *Racemosa*, *Geoffræa inermis*—a. *Surinamensis*, *Geoffræa Surinamensis*.

ANDRACHAHARA, *Sempervivum tectorum*.

ANDRACHNE, (ανδραχνη,) *Arbutus unedo*, *Portulaca*.

ANDRÆDÆA, (andro, and *ædæa*,) *Pudibilia*.

ANDRALOGOMÈLE, (andro, a. priv., *λογος*, 'proportion,' and *ανωλον*, 'a domestic animal,') A name given by Malacarne to a monster in which he supposed the presence of the body of a man with the limbs of a brute.

ANDRANATOM'IA, *Andranat'omē*, *Andron'tom'ia*, *Androt'omē*, *Anthropot'omy*, (andro, and *τεμνω*, 'to cut,') The anatomy of man.

ANDRI'A, (ανδρεια,) Adult age. Manhood.

ANDRIA Mc'LIER, *Mulier Hermaphroditi'ca*. A female hermaphrodite.

ANDRO, (from *ανηρ*, gen. *ανδρος*, 'man,') In composition, man.

ANDROARIUM, (andro, and *ovarium*, 'seed-vessel,') Testicle.

ANDROGENTA, (ανδρογενεια, from *andro*, and *γενεσις*, 'generation,') The procreation of males.

ANDROGYNOS, *Hermaphroditic*.

ANDROG'YNUC, (ανδρῳγνος, from *andro*, and *γυνη*, 'a woman,') A hermaphrodite. An effeminate person.

ANDROLEPSIA, (ανδροληψις, from *andro*, and *λαμβάνω*, 'to lay hold of,') Conception.

ANDROMANIA, (from *andro*, and *mania*,) Nymphomania.

ANDROMEDA ARBO'REA, (from *Andromeda*, in antiquity.) *Sorrel Tree*, *Sour Tree*, *Sour Wood*, *Elk Tree*, *Elk Wood*, *Sorrel Wood*, *Sour Leaf*, (F.) *Andromède*, *Andromédier*. A small indigenous tree; Ord. Ericaceae, *Sex. Syst.* Decandria Monogynia; found in the Alleghany mountains and the hills and valleys diverging from them, as far as the southern limits of Georgia and Alabama; but seldom north of Virginia. The sour leaves are refrigerant and astringent, and have been used to make a kind of lemonade, which has been given in fevers.

ANDROMEDA MARIA'NA, *Broad-leaved Mour't-wort*, *Kill Lamb*, *Lamb Killer*, *Stag'gerbush*. A decoction of this American plant is said to have been successfully employed as a wash, in a disagreeable affection—not uncommon amongst the slaves in the southern parts of the United States—called the *Toe Itch*, and *Ground Itch*.

ANDROMEDA NIT'IDA, *Sourwood*, *Sorrel tree*, indigenous; has properties similar to those of *A. arborea*.

ANDROPOGON BICORNIS, (andro, and *πωγων*, 'a beard,') *Junctus odoratus*—a. *Calamus aromaticus*, see *Oleum graminis Indici*—a. *Citratus*, *Junctus odoratus*—a. *Citridiorus*, *Junctus odoratus*, *Nardus Indica*.

ANDROPOGON MURICA'TUS, *A. squarrosus*, *Phalaris zizanioides*, *Agrostis verticillata*, *Ammethum muricatum*, *Vetiveria odorata*, *Vetiver*, *Cuscuta*, *Khus-Khus*, (F.) *Vetiver*, *Vetiver*, *Vetiver*; *Nat. Ord.* Gramineae. The root of this plant, from Bombay, is aromatic and bitterish. It is used as a perfume, and to drive away insects. It is also an excitant and diaphoretic.

ANDROPOGON NARDUS, *Calamus Alexandrinus*, *Nardus Indica*—a. *Schœnanthus*, *Junctus odoratus*—a. *Squarrosus*, *A. Muricatus*.

ANDROSACE, (andro, and *σας*, 'a buckler,' from the appearance of the leaf.) *Umbilicus marinus*—a. *Matthioli*, *Umbilicus marinus*.

ANDROSÆMUM, (andro, and *αιμα*, 'blood,') *Hypericum perforatum*.

ANDROTOMY, *Andranatomia*.

ANDRUM. An East India word, latinized by Kæmpfer, signifying a kind of elephantiasis of the scrotum, endemic in southern Asia.

ANÉANTISSEMENT (F.), (from *anéantir*, (a, and *neant*, 'nothing,') (L.) *annihilare*, 'to annihilate,') *Virium extinctio*. This word is often employed hyperbolically, by patients in France, to signify excessive fatigue, debility or syncope.

ANEBIUM, *Anchusa tinctoria*.

ANEBUS, (ἀνέβω, from *an*, and ἰβη, 'puberty,') Impuler.

ANECYPE'TUS, (ἀνεκπυετός, from *an*, and πυνω, 'I promote suppuration,') That which does not suppurate, or is not likely to suppurate.

ANEGERTICA, (ἀνεγερτω, (*ava*, and *εγερτω*), 'I awaken,') The art of resuscitating the apparently dead.

ANEILE'MA, gen. *Aneile'matis*, *Aneile'sis*, (ἀνείλημα, from ἀνείλω, (*ava*, and εἰλω), 'I roll up,') Applied particularly to the motion of air in the intestines and the tormina accompanying it.

ANEILE'SIS, (ἀνείλησις,) *Aneilema*.

ANEMIA, *Anæmia*.

ANEMO'NE, (ἀνεμώνη,) *Wind Flower*: (from *anemos*, 'the wind,' because it is said not to open its flowers until blown upon by the wind [?].)

ANÉMONE DES BOIS, *Anemone nemorosa*.

ANEMONE COLLINA, *A. Pulsatilla*—*a. Grönlandica*, *Coptis*—*a. Hepatica*, *Hepatica triloba*—*a. Intermedia*, *A. Pulsatilla*.

ANEMONE LUDOVICIANA, *A. patens*.

ANEMONE NEMORO'SA, *Ranunculus albus* seu *nemorosus*, *Wood Anem'ony* or *Anem'ony*, (F.) *Anémone des bois*. *Ord.* *Ranunculacææ*. The herb and flowers are poisonous, acrid, and corrosive. They have been used as rubefacients.

ANEMONE PA'TENS, *A. Ludoviciana*, *Pulsatilla patens*; indigenous: from Illinois and Wisconsin to the Rocky Mountains, is supposed to possess similar properties.

ANEMONE PRATEN'SIS, *A. Sylvestris*, *Pulsatilla nigricans* seu *pratensis*. This plant has similar properties with the last. It is also called *Mead'ow Anem'ony*, (F.) *Pulsatille noire*. *P. des prés*.

ANEMONE PULSATIL'LA, *A. Colli'na* seu *Intermedia* seu *Pratensis* seu *Ru'bra*, *Pulsatilla vulgaris*, *Her'ba ven'tis*, *Pasque flower*, (F.) *Coquelourde*, *Pasce-fleur*, possesses like properties.

ANEMONE RUBRA, *A. Praten'sis*—*a. Rue-leaved*, *Thalictrum anemonoides*—*a. Sylvestris*, *A. Praten'sis*.

ANEMONY, *Anemone hepatica*—*a. Meadow*, *Anemone pratensis*—*a. Wood*, *Anemone nemorosa*.

ANEMOS, (άνεμος,) *Wind*.

ANCEPHEALIA, see *Anencephalus*.

ANCEPHEALOHÆMIA, (αν. εγκεφαλος, 'encephalon,' and ἷμα, 'blood,') Defect of blood in the brain. *Syncope*.

ANCEPHEALONEU'RIA, *Anencephalon'eria*, (αν, εγκεφαλος, 'encephalon,' and νευρον, 'nerve,') Want of nervous action in the encephalon.

ANCEPHEALOTROPH'IA, (αν, εγκεφαλος, 'the encephalon,' and τροφή, 'nourishment,') Atrophy of the encephalon.

ANCEPHE'ALUS, (αν, and εγκεφαλος, 'brain,') A monster devoid of brain. The condition has been called *Anencepha'lia*. Also, one that has a part only of the brain;—*Paraceph'alus*. A weak, silly person.

ANENERGESIA, (ανεργησία, from *an*, *ev*, 'in,' and εργον, 'work,') *Debility*.

ANENERGIA, *Debility*.

ANÉTERONERVIE SATURNINE, (αν, εντέρον, 'intestine,' and νευρον, 'nerve,') see *Palsy*, *lead*.

ANEPISCHEISIS, (αν, and epischesis,) *Incontinencia*.

ANEPITHYM'IA, (αν, and επιθυμία, 'desire,') Many nosologists have used this word for a loss of the appetites, as of those of hunger, thirst, venery, &c.

ANEPITHYMIA CHLOROSIS, *Chlorosis*.

ANER, (ανηρ, gen. ανδρος,) *A man*.

ANERETHIS'IA, *Inirritabil'itas*, (αν, and ερεσις, 'irritability,') Defect of irritability.

ANERVISMIE, (α, and nervus, 'a nerve,') *Paralysis*.

ANERYTHROP'SIA, (αν, ερυθρος, 'red,' and οψις, 'vision,') Defective vision, which consists in an incapacity of distinguishing red.

ANESIS, (ανείσις, from *anistu*, 'to relax, remit,') *Remission*.

ANESON, (ανέσσον,) *Anethum*.

ANESTHÉSIE, *Anæsthesia*.

ANESTHÉSIE EXTATIQUE. The aggregate of phenomena of impaired feeling produced especially by the manipulations of the animal magnetizer.

ANESTHÉSIER, see *Anæsthetization*.

ANESTHÉSIQUE, *Anæsthetic*.

ANESTHETIC, *Anæsthetic*.

ANESTHETIZATION, *Anæsthetization*.

ANESTHETIZE, see *Anæsthetization*.

ANESUM, *Pimpinella anisum*.

ANET, *Anethum*.

ANETH, *Anethum graveolens*.

ANETHUM, (ανηθον, from *ano*, and *θω*, 'to run,' from its quick growth, [?]) *An'e'son*, *An'e'ton*, *An'e'thum* *Fœnic'ulum*, seu *Seg'etum* seu *Piper'itum*, *Fœnic'ulum*, *F. Dul'ce* seu *Officinale* seu *vulga're*, *Ligusticum fœnic'ulum*, *Fœn'iculum*, *Mar'athrum*, *Fœn'el* or *Fœn'kle*, *An'et*, *Sweet Fen'nel*, (Prov.) *Spin'gel*, (F.) *Fenouil* ou *Anis doux*. *Ord.* *Umbelliferae*. *Sex. Syst.* *Pentandria Digynia*. The fruit, *Fœnic'ulum* (Ph. U. S.), has an aromatic odour, and warm, sweetish taste. It is carminative. The oil—*Oleum Fœn'iculi*, oil of Fennel—is official in the Ph. U. S. The root is said to be pectoral and diuretic.

ANETHUM FENICULUM, *Anethum*.

ANETHUM GRAVEOLENS, *An'e'thum*, *A. hortens'e*, *Pastinaca An'e'thum* seu *Grave'olens*, *Fer'ula Grave'olens*, *Dill*, (F.) *Aneth*, *Fenouil puant*. A native of the south of Europe. The seeds are stimulant and carminative. A distilled water—*A'qua ane'thi*, *Dill water*—was official in the London and Edinburgh Pharmacopœias. Dose. gr. xv to ʒi.

O'leum An'e'thi, *Oil of Dill*, (F.) *Huile d'Aneth*, possesses the carminative properties of the plant.

ANETHUM PASTINACA, *Pastinaca Sativa*—*a. Piperitum*, *Anethum*—*a. Segetum*, *Anethum*.

ANETICUS, (see *Anesis*,) *Anodyne*.

ANETON, (ανητον,) *Anethum*.

ANETUS, (ανετος,) *Intermittent fever*—*a. Quartan*, *Quartan*—*a. Quotidianus*, *Quotidian*—*a. Tertianus*, *Tertian fever*.

ANEURAL'GICON, (α, νευρον, 'nerve,' and αλγος, 'pain,') A name given by Dr. C. T. Downing to an instrument used by him to allay pain in nerves. It is a kind of fumigating apparatus, in which dried narcotic and other herbs are burnt, the heated vapour being directed to any part of the body.

ANEURIA, (α, and νευρον, 'a nerve,') *Paralysis*.

ANEURISM, *An'eury'sm*, *Aneurys'ma*, *Aneurys'mus*, *Aneuris'ma*, *Ced'ma*, (ανευρύσμα, from *aneuryon*, (*ava*, and *ευρυειν*), 'to dilate or distend,') *Dilatatio Arteriarum*, *Ectasia*, *Exan'gia aneuris'ma*, *Arterieurys'ma*, *Artereurys'ma*, *Hæmatoce'le arterio'ea*, *Abces'sus spirituosus*, *Arteriec'tasis*, (F.) *Anévrisme*, *Aneurisme*. Properly, *Aneurism* signifies a tumour, produced by the dilatation of an artery; but it has been extended to lesions of arteries, as well as to dilatations of the heart.

There are various kinds of aneurism. The following are the chief:

I. When the blood, which forms the tumour, is enclosed within the dilated coats of the artery. This is the true *Aneurism*, *Aneurys'ma ve'rum*, *Her'nia Arteriarum*, (F.) *Anévrisme vrai*.

II. When the blood has escaped from the opened artery, it is called *spuri'ous* or *false* *Aneurism*, *Aneuris'ma spu'rium*, *Ruptu'ra Arte'rie*, *Arteriorrhex'is*, *Arteriodial'ysis*, *Echymo'ma arterio'cum*, (F.) *Anévrisme faux*. The latter is divided into

three varieties. 1. *Diffused False Aneurism*, (F.) *Ànéisme faux, primitif, diffus, noncirconscrit ou par infiltration*, which occurs immediately after the division or rupture of an artery, and consists of an extravasation of blood into the areolar texture of the part. 2. *Circumscribed False Aneurism*, (F.) *Anérysme faux consécutif, circonscrit ou par épanchement, enkysté ou saciforme, tumeur hémorrhagiale circonscrite*, in which the blood issues from the vessel some time after the receipt of the wound, and forms itself a sac in the neighbouring areolar tissue. 3. *Aneurism by Anastomosis*, *Varicose or Cirroid Aneurism*, *Phlebotriodinal ysis*, *Aneurysma veno-arteriosum*, *A. varicosum*, (F.) *Anérysme par anastomose ou variqueux*, *A. par érosion*, *A. de Pott*, *A. des plus petites artères*, which arises from the simultaneous wounding of an artery and a vein;—the arterial blood passing into the vein, and producing a varicose state of it.

III. *Mixed Aneurism*, (F.) *Anérysme mixte*, is that which arises from the dilatation of one or two of the coats, with division or rupture of the other. Some authors have made two varieties of this:—1. *Mixed external Aneurism*, where the internal and middle coats are ruptured, and the areolar is dilated. 2. *Mixed internal Aneurism*, in which the internal coat is dilated, and protrudes, like a hernial sac, through the ruptured middle and outer coats. This variety has been called *Aneurysma Herniam Arteriarum sistens*.

Aneurisms have been termed *traumatic* or *exogenous*, and *spontaneous*, according as they may have been caused by a wound, or have originated spontaneously. The latter, when originating from lesions of the inner coats of arteries, have been termed *endogenous*. They have also been divided into *internal* and *external*.

The *internal aneurisms* are situate in the great splanchnic cavities, and occur in the heart and great vessels of the chest, abdomen, &c. Their diagnosis is difficult, and they are often inaccessible to surgical treatment.

The *external aneurisms* are situate at the exterior of the head, neck, and limbs, and are distinctly pulsatory.

Aneurisms, especially the internal, may be combated by a debilitating treatment, on the plan of Valsalva, which consists in repeated blood-letting, with food enough merely to support life. In external aneurism, the artery can be obliterated. This is usually done by applying a ligature above the aneurismal tumour.

ANEURISM BY ANASTOMOSIS, see Aneurism—a. Brasdor's operation for, see Brasdor—a. Cirroid, see Aneurism.

ANEURISM, DISSECTING, is one in which, owing to rupture of the inner and middle coats of an artery, the blood makes itself a channel between these coats and the outer coat.

In many cases, the lesion appears to consist in a separation of the laminae of the middle coat, between which the blood forms itself a channel.

ANEURISM, ENDOGENOUS, see Aneurism, and Endogenous—a. Exogenous, see Aneurism, and Exogenous—a. External, see Aneurism—a. False, see Aneurism—a. False, circumscribed, see Aneurism—a. False, diffused, see Aneurism.

ANEURISM OF THE HEART, *Cardionchus*, *Cardiureysma*, *Cardianeureysma*, (F.) *Anérysme du cœur*, has been divided into *active* and *passive*. The former can scarcely be esteemed aneurisms, as they most commonly consist of increased thickness of the parietes of the heart, which diminishes its cavity instead of increasing it. The term *Hypertrophy of the heart* better indicates their character. *Passive aneurism*, *Cardiectasis*, on the contrary, is attended with extenuation of the parietes of the organ, and enlargement of the cavities. The physical signs of *dilatation of the heart* are

the following:—The action of the heart is not visible, and no impulse is conveyed to the hand. On percussion, there is a loss of resonance over a larger surface than usual, but the dullness is much less intense than that which accompanies hypertrophy. On auscultation, the action of the heart is only slightly felt, and communicates at once the impression of its diminished power. The impulse is feeble than usual. Both sounds are widely transmitted over the thorax, and are not much fainter at a distance from their point of origin.

Partial or true aneurism of the Heart—*Cardiectasis partialis*, *Aneurysma consecutum cordis*—is sometimes seen; rarely, however.

The name *Aneurism of the Valves of the Heart* has been given to pouch-like projections of the valves into the auricles.

ANEURISM INTERNAL, see Aneurism—a. Mixed, see Aneurism—a. Mixed, external, see Aneurism—a. Mixed, internal, see Aneurism—a. Spontaneous, see Aneurism—a. Spurious, see Aneurism—a. Traumatic, see Aneurism—a. True, see Aneurism—a. Valsalva's method of treating, see Aneurism—a. Varicose, see Aneurism.

ANEURISMA, gen. *Aneurism'atis*, Aneurism.

ANEURISMAL, *Aneurys'mal*, *Aneurismat'ic*, *Aneurysmat'icus*, *Aneurisma'lis*, (F.) *Anérysmael*, *Anérysmatique*. That which belongs to Aneurism.

ANEURISMAL SAC or CYST, (F.) *Sac ou Kyste anérysmael*, is a sort of pouch, formed by the dilatation of the coats of an artery, in which the blood, forming the aneurismal tumour, is contained.

ANEURISMATIC, Aneurismal.

ANEURYSM, Aneurism.

ANEURYSMA, gen. *Aneurysm'atis*, (*aneurysma*.)

Aneurism—a. *Cordis activum*, Heart, hypertrophy of the—a. *Herniam arteriarum sistens*, see Aneurism—a. *Spurium*, see Aneurism—a. *Varicosum*, see Aneurism—a. *Venoso-arteriosum*, see Aneurism—a. *Verum*, see Aneurism.

ANEURYSME, Aneurism.

ANEURYSMUS, Aneurism, Dilatation.

ANÉVRIA, (*a*, and *νεῦρον*, 'a nerve,') Paralysis.

ANÉVRYSMAL, Aneurismal.

ANÉVRYSMATIQUE, Aneurismal.

ANÉVRYSME, Aneurism—a. *de l'Aorte*, *Aortureysma*—*a. de Pott*, see Aneurism—a. *des plus petites artères*, see Aneurism—a. *Circonscrit*, see Aneurism—a. *Diffus*, see Aneurism—a. *Enkysté*, see Aneurism—a. *Faux*, see Aneurism—a. *Faux consécutif*, see Aneurism—a. *Mixte*, see Aneurism—a. *par Anastomose*, see Aneurism—a. *par Épanchement*, see Aneurism—a. *par Érosion*, see Aneurism—a. *par Infiltration*, see Aneurism—a. *Primitif*, see Aneurism—a. *Sacciforme*, see Aneurism—a. *Variqueux*, see Aneurism—a. *Vrai*, see Aneurism.

ANEYS, see Pimpinella anisum.

ANFION, Maslach.

ANFRACTUOSITÉS, Anfractuosités —*a. Cérébrales*, Anfractuosités, cerebral—a. *Ethmoïdales*, see Anfractuosity.

ANFRACTUOSITY, *Anfrac'tus*, *Sul'cus*, (*am*, 'around,' and *frangere*, *fractum*, 'to break.') (F.) *Anfractuosité*. A groove or furrow. Used in anatomy to signify sinuous depressions or sulci, of greater or less depth, like those which separate the convolutions of the brain from each other. These

ANFRACTUOSITIES, CEREBRAL, *Anfrac'tus Cerebri*, (F.) *Anfractuosités Cérébrales*, are always narrow, and deeper at the upper surface of the brain than at its base; and are lined by a prolongation of the pia mater.

The Ethmoid Cells are, sometimes, called *Anfractuosités ethmoïdales*.

ANFRACTUS, Anfractuosity—a. *Cerebri*, Anfractuosités, (cerebral.)

ANGECTASIA, Angiectasia.

ANGEIA, (pl. of *Angeion*.) see *Angeion*—a. Pneumatica, see Artery.

ANGELIAL, (from *angeion*.) Vascular.

ANGIECTASIA, Angiectasis.

ANGIECTASIS, Angiectasis.

ANGIECTOMA, Angiectasis.

ANGIEOCARDITIS, Angiocarditis.

ANGIEOG'ENY, *Angeiog'enia*, *Angiog'eny* *Angeiog'enia*, (*angeion*, and *γεννω*, 'I generate.') Formation or generation of vessels.

ANGIEOG'RAPHY, *Angiog'raphy*, *Angeiog'raph'ia*, (*angeion*, and *γραφη*, 'a description.') A description of the vessels.

ANGIEOHYDROG'RAPHY, *Angiohydrog'raphy*, *Angiohydrog'raphia*, *Hydrangiograph'ia*, (*angeion*, 'vessel,' and *γραφω*, 'I describe.') A description of the lymphatics.

ANGIEOHYDROTOMY, *Angiohydrot'omy*, *Angiohydrotom'ia*, *Hydrangiotom'ia*, (*angeion*, 'vessel,' and *τεμνω*, 'to cut.') Dissection of the lymphatics.

ANGIEOLEUCITIS, *Angioleucit'is*, (*angeion*, λευκος, 'white,' and *itis*.) *Angiolymph'itis*, *Lymphangi'itis*, *Lymphangi'itis*, *Lymphangi'itis*, *Hydrangi'itis*, *Lymphat'itis*, *Lymphat'itis*, *Inflammatio vasorum lymphaticorum*, (F.) *Inflammation des vaisseaux lymphatiques ou des tissus blancs*. Inflammation of the lymphatics. Lymphatic or serofulous inflammation.

ANGIEOLOG'Y, *Angiol'ogy*, *Angieolog'ia*, (*angeion*, and *λογος*, 'a discourse.') A treatise on the vessels. The anatomy of the vessels. It includes *Arteriolog'ogy* *Phlebolog'ogy*, and *Angieohydrolog'ogy*.

ANGIEOMALA'CIA, *Angiomala'cia*, (from *angeion*, and *μαλακία*, 'softening.') Mollescence or softening of vessels.

ANGIEOMYCES, (*angeion*, and *μυκης*, 'a fungus.') *Hæmatodes fungus*.

ANGIEON, (*αγγειον*, pl. *αγγεία*.) Vessel.

ANGIEONOSUS, (*νοσος*, 'disease,') *Angieopathia*.

ANGIEONUSUS, *Angieopathia*.

ANGIEOPATHI'A, *Angiopathi'a*, *Angieon'osis*, *Angieon'us*, *Angio'is*, (*angeion*, and *παθος*, 'a disease,') Disease of the vessels.

ANGIEOPLA'NIA, *Angiopl'a'nia*, (*angeion*, 'a vessel,' and *πλανη*, 'error,') Anomaly in the structure and distribution of vessels.

ANGIEOPLEROSIS, (*angeion*, and *πληρω*, 'I fill up,') *Plethora*.

ANGIEOPYRA, (*angeion*, and *πυρ*, 'fire, fever,') *Synocha*.

ANGIEORRHAGIA, (*angeion*, and *ρην*, 'rupture,') *Hæmorrhagia activa*.

ANGIEORRHÆ'A, *Angiorrhæ'a*, (F.) *Angieorrhæe*, (*angeion*, and *ρην*, 'I flow,') *Passive hæmorrhage*.

ANGIEOSIS, Angiosis.

ANGIEOSTEGNOSIS, (*angeion*, and *στεγνω*, 'I constrict,') Constriction of vessels.

ANGIEOSTENOSIS, (*angeion*, and *στενος*, 'narrow,') Narrowness of vessels.

ANGIEOSTEO'SIS, *Angiost'o'is*, (*angeion*, and *οστεωσις*, 'ossification,') Ossification of vessels.

ANGIEOSTROPHE, (*angeion*, and *στρεφειν*, 'to turn, to twist,') See *Torsion*.

ANGIEOTELECTASIA, (*angeion*, *τελος*, 'end,' and *εκτασις*, 'dilatation,') *Telangiectasia*.

ANGIEOT'OMY, *Angiot'omy*, *Angieotom'ia*, (*angeion*, and *τεμνω*, 'to cut,') Dissection of vessels.

ANGIE'TIS, *Angi'tis*, *Angioi'tis*, *Inflammatio vasorum*, (F.) *Angéite*, (*angeion* and *itis*.) Inflammation of vessels in general.

ANGEL-BREAD. A kind of purgative cake, formerly made of spurge, ginger, flour and oatmeal.

ANGEL WATER. A fashionable cosmetic in

the 17th century. It is said to have been prepared by simmering together *orangeflower water*, *benzoin*, *storax*, *cinnamon*, *cloves*, and *calamus aromaticus*; adding, also, *musk*. The supernatant fluid was the angel water. Perfumed cakes were made of the dregs, and 'sweet bags' to lay among clothes.

ANGELIC ROOT, *Angelica lucida*.

ANGEL'ICA, *Angel'ica* *Archangel'ica* seu *Hisp'ana* seu *Sati'va*, *Archangel'ica officina'lis*, *Gar'den Angel'ica*, (F.) *Angélique*, *Racine de Saint Esprit*. (So called from its supposed angelic virtues.) *Ord.* Umbelliferae. *Sec. Syst.* Pentandria Digynia. Native of Lapland. The root—*Angelica*, (Ph. U. S.,)—stalk, leaves, and seed, are aromatic and carminative. A sweetmeat is made of the root, which is agreeable.

ANGELICA ARCHANGELICA, *Angelica*.

ANGELICA ATROPURP'UREA, *Angel'ica* (Ph. U. S.), *Masterwort*. An indigenous species, growing over the whole United States, and admitted into the secondary list of the Pharmacopœia of the United States. Virtues, same as those of the *Angelica* of Europe.

ANGELICA LEVISTICUM, *Ligusticum levisticum*.

ANGELICA LU'CIDA, *Angel'ic root*, *Bell'lyache root*, *Neu'do*, *White root*, an indigenous plant, the root of which is bitterish, subacrid, fragrant, aromatic, stomachic, and tonic. Also, *Ligusticum actæifolium*.

ANGELICA NENDO, *Ligusticum actæifolium*.

ANGELICA OFFICINALIS, *Imperatoria*—a. *Paludapifolia*, *Ligusticum levisticum*—a. *Sativa*, *Angelica*, *A. sylvestris*.

ANGELICA SYLVES'TRIS, *A. sati'va*, *Seli'nium Sylves'trè* seu *Angel'ica* seu *Pubes'cens*, *Imperato'ria Sylves'tris* seu *Angel'ica*, *Wild Angel'ica*, (F.) *Angélique sauvage*. Possesses similar properties to the last, but in an inferior degree. The seeds, powdered and put into the hair, are used to destroy lice. Also, *Ligusticum podagraria*.

ANGELICA TREE, *Aralia spinosa*.

ANGELINE COR'TEX. The bark of a Grenada tree, (*Geoffræa iner'mis* [?]), which has been recommended as anthelmintic and cathartic.

ANGÉLIQUE, *Angelica*—a. *Sauvage*, *Angelica sylvestris*.

ANGELOCACOS, *Myrobalanus*.

ANGEMPHRAXIS, *Angiemphraxis*.

ANGIAIRRHAGIE, (*angeion*, *anh*, 'air,' and *ρηγνυμι*, 'to break forth,') *Hæmoptysis*.

ANGIAIRRHÉMIE, (*angeion*, *anh*, 'air,' and *ἄμα*, 'blood,') *Hæmoptysis*.

ANGIAIRRRHYDRIE, (*angeion*, *anh*, 'air,' and *ῥυδω*, 'water,') *Asphyxia* by submersion.

ANGIBROMELMINTHIE, (*angeion*, *βρωμα*, 'food,' and *ἐλμινς*, 'a worm,') *Worms (intestinal)*.

ANGICHOLOLITHE, (*angeion*, *χολος*, 'bile,' and *λιθος*, 'a stone,') *Calculi biliary*.

ANGIDIECTASIA, *Trichangiectasia*.

ANGIDIOSPONGUS, (*αγγειδιον*, 'a small vessel,' and *σπογγος*, 'a fungus,') *Hæmatodes fungus*.

ANGIECTASIA VENOSA, *Varix*.

ANGIEO'TASIS, *Angiecta'sia*, *Angecta'sia*, *Angieury'sma*, *Angiecto'ma*, *Angiodias'tasis*, (*angeion*, and *εκτασις*, 'dilatation,') *Dilatation of vessels*. See *Telangiectasia*.

ANGIECTOPIA, (*angeion*, *ex*, 'out of,' and *τοπος*, 'a place,') A condition in which the vessels are out of their natural situation.

ANGIEMPHRAX'IS, *Angiemphrax'is*, (*angeion*, and *εμφραξις*, 'obstruction,') *Obstruction of vessels*.

ANGIEURY'SMA, gen. *Angieury'sm'atis*, (*angeion*, and *ευρυνω*, 'to dilate,') *Angiectasis*.

ANGIITE, *Angieitis*, *Inflammation*.

ANGIITIS, (*angeion*, and *itis*.) *Angieitis*.

ANGI'NA, *Febris Angino'sa*, *Isthm'i'tis*, *Quin'sy*

or *Sorethroat*; (*angere*, 'to choke.') Inflammation of the supra-diaphragmatic portion of the alimentary canal, and of the air passages. The Latin writers applied the term to every disease in which deglutition or respiration, separately or united, was affected, provided that such affection was above the stomach and lungs. Buerhaave speaks of the angina of the moribund, which is nothing more than the dysphagia or difficult deglutition preceding death. See *Cynanche*.

ANGINA APHTHOSA, *Aphthæ*—*a. Aquea*, *Edema of the glottis*—*a. Bronchialis*, *Bronchitis*—*a. Canina*, *Cynanche trachealis*—*a. Cordis*, *Angina pectoris*—*a. cum Tumore*, *Cynanche tonsillaris*—*a. Epidemica*, *Cynanche maligna*—*a. Epiglottidea*, *Epiglottitis*—*a. Erysipelatosa*, *Erythræche*—*a. Exsudatoria*, *Cynanche trachealis*—*a. Externa*, *Cynanche parotidea*—*a. Faucium*, *Isthmitis*—*a. Faucium exsudativa*, *Pharyngitis*, *diphtheritic*—*a. Faucium Maligna*, *Cynanche maligna*—*a. Folliculosa of the pharynx*, *Pharyngitis*, *follicular*—*a. Gangrenosa*, *Cynanche maligna*—*a. Glandulosa*, *Pharyngitis*, *follicular*—*a. Hogskin*, *Pharyngitis diphtheritic*—*a. Humida*, *Cynanche trachealis*—*a. Inflammatoria*, *Cynanche*, *Cynanche trachealis*—*a. Laryngea*, *Laryngitis*—*a. Laryngea* (*Edematosa*, *Edema of the glottis*—*a. Lingularia*, *Glossitis*—*a. Maligna*, *Angina pellicularis*, *Cynanche maligna*, *Pharyngitis*, *diphtheritic*—*a. Maxillaris*, *Cynanche parotidea*—*a. Membranacea*, *Cynanche trachealis*—*a. Mitis*, *Isthmitis*.

ANGINA NASALIS, *Nasitis postica*. An inflammation of the posterior portion of the Schneiderian membrane lining the nose. Also, *Coryza*.

ANGINA (EDEMATA) SA. (F.) *Angine œdémateuse*, *Edème de la Glotte*. An œdematous swelling of the glottis, the effect of chronic *cynanche laryngea*. See *Edema of the Glottis*.

ANGINA PALATINA, *Hyperoitis*—*a. Paralytica*, *Pharyngoplegia*—*a. Parotidea Externa*, *Cynanche parotidea*.

ANGINA PECTORIS, *A. cordis*, *Sternalgia*, *Asthma spasticœrthriticum inconstans*, *Asthma diaphragmaticum*, *Arthritis diaphragmatica*, *Orthopnea cardiaca*, *Sternodynia syncopica et palpitans*, *S. syncopalis*, *Cardiogenus cordis sinistri*, *Asthenia pectoralis*, *An'gor pectoris*, *Stenocardia*, *Diaphragmatic gout*, *Asthma convulsivum*, *Asthma arthriticum*, *Cardioneuralgia*, *Neuralgia brachiothoracica*, *Hyperæsthesia plexus cordis*, *A. dolorificum*, *Syncope anginosæ seu angens*, *Cardiodyne spasmodica intermittens*, *Phlegmo'gia*, *Prunella*, *Suspensum cardiacum*, *Pneumonalgia*, *Suffocative Breast pang*, (F.) *Angine de Poitrine*, *Nérose du Cœur*. A disease, the precise pathology of which is not known. The principal symptoms are, violent pain about the sternum, extending towards the arms, anxiety, dyspnoea, and sense of suffocation. It is an affection of great danger, and is often connected with ossification, or other morbid condition of the heart. It appears to be neuropathic, and has been termed *Neuralgia of the Heart*. Some, however, employ this last term for an acutely painful intermittent affection of the heart, which seems to differ from *angina pectoris* more in regard to the small number of parts which are drawn into morbid consent with the affected cardiac nerves, than in regard either to its nature or appropriate treatment. The most powerful stimulating and narcotic antispasmodics are required during the paroxysm.

ANGINA PELLICULARIS, *Pharyngitis*, *diphtheritic*—*a. Perniciosa*, *Cynanche trachealis*—*a. Pestilentialis*, *Pharyngitis*, *diphtheritic*—*a. Membranacea*, *Cynanche trachealis*—*a. Polyposa*, *Cynanche trachealis*—*a. Pseudo-membranosa*, *Pharyngitis*, *diphtheritic*—*a. Pulposa*, *Cynanche trachealis*—*a. Sanguinea*, *Cynanche tonsillaris*.

ANGINA SICCA, (F.) *Angine Sèche*, is a chronic

inflammation of the pharynx, with a distressing sense of dryness and heat, in chronic diseases of the stomach and lungs. See *Pædanchone*.

ANGINA SIMPLEX, *Isthmitis*.

ANGINA SQUIRROSA, (F.) *Angine squirreuse*, consists in difficulty of deglutition, caused by scirrhus disorganization of the pharynx or œsophagus, or by enlarged tonsils.

ANGINA STRANGULATORIA, *Cynanche trachealis*—*a. Strepitosa*, *Cynanche trachealis*—*a. Suffocatoria*, *Cynanche trachealis*—*a. Synochalis*, *Cynanche tonsillaris*—*a. Thyreoidica*, *Thyreoiditis*—*a. Tonsillaris*, *Cynanche tonsillaris*—*a. Trachealis*, *Cynanche trachealis*—*a. Ulcerosa*, *Cynanche maligna*—*a. Uvularis*, *Staphyloedema*, *Uvulitis*—*a. Vera et Legitima*, *Cynanche tonsillaris*.

ANGINE, *Angina*—*a. Cœuvenneuse*, *Pharyngitis*, *diphtheritic*—*a. Diphthérique*, *Pharyngitis*, *diphtheritic*—*a. Gutturale*, *Cynanche tonsillaris*—*a. Gutturale cœuvenneuse*, *Pharyngitis*, *diphtheritic*—*a. Laryngée*, *Laryngitis*—*a. Laryngée et trachéale*, *Cynanche trachealis*—*a. Laryngée adémateuse*, *Edema of the glottis*—*a. Œsophagienne*, *Œsophagitis*—*a. Pharyngée*, *Cynanche parotidea*—*a. Plastique*, *Pharyngitis*, *diphtheritic*—*a. de Poitrine*, *Angina pectoris*—*a. Pseudomembraneuse*, *Pharyngitis*, *diphtheritic*—*a. Sèche*, *Angina sicca*—*a. Simple*, *Isthmitis*—*a. Squirreuse*, *Angina Squirrosa*—*a. Tonsillaire*, *Cynanche tonsillaris*.

ANGINETIX, *Angineuse*.

AN'GINOSE, *An'ginous*, *Anginosæ*. (F.) *Angineux*. Relating or appertaining to angina; as *Scarlati'na anginosæ*.

ANGIOCARDITIS, *Angiocarditis*, (*angeion*, and *carditis*.) Inflammation of the heart and great vessels.

ANGIODIASTASIS. (*angeion*, and *διαστασις*, 'separation.') *Angiectasis*.

ANGIOGENY, *Angiogeny*.

ANGIOGRAPHY, *Angiography*.

ANGIOHÉMIE, (*angeion*, and *αἷμα*, 'blood.') *Hyperemia*.

ANGIOHYDROGRAPHY, *Angiohydrography*.

ANGIOHYDROTOMY, *Angiohydrotomy*.

ANGIOITIS, *Angitis*.

ANGIOLEUCITIS, *Angioleucitis*.

ANGIOLOGY, *Angiology*.

ANGIOLYMPHITIS, *Angioleucitis*.

ANGIOMALACIA, *Angiomalacia*.

ANGIOMYCES, (*angeion*, and *μυκῆς*, 'a fungus.') *Hæmatodes fungus*.

ANGIONOSUS, (*angeion*, and *νόσος*, 'a disease.') *Angiopathia*.

ANGIONUSUS, *Angiopathia*.

ANGIOPATHIA, *Angiopathia*.

ANGIOPLANIA, *Angioplania*.

ANGIOPLETHOROSIS, (*angeion*, and *πληρω*, 'I fill.') *Plethora*.

ANGIOPYRA, (*angeion*, and *πῦρ*, 'fire, fever.') *Synocha*.

ANGIORRHAGIA, (*angeion*, and *ῥαγν*, 'rupture.') *Hæmorrhagia activa*.

ANGIORRHEA, *Angiorrhœa*.

ANGIOSIS, *Angiopathia*.

ANGIOSTEGNOSIS, (*στεγνωσις*, 'constriction.') *Angiostegnosis*.

ANGIOSTENOSIS, (*στενωσις*, 'contraction.') *Angiostenosis*.

ANGIOSTOSIS, *Angiostosis*.

ANGIOSTROPHE, (*στροφῶ*, 'I turn.') See *Torsion*.

ANGIOTELECTASIA, (*angeion*, *τελος*, 'end,' and *εκτασις*, 'extension, dilatation.') *Telangiectasis*.

ANGIOTENIC, (*Angiotenic*, *Angioten'ic* seu *Angioten'ic*, (*angion*, and *τενεν*, 'to stretch.') An epithet given to inflammatory fever, owing to its action seeming to be chiefly exerted on the vascular system.

ANGIOTOMY, Angiotomy.

ANGLE, *An'gulus*, (αγκυλος, 'a hook.') The space between two lines which meet in a point.

ANGLE, FA'CIAL, pointed out by Camper, is formed by the union of two lines, one of which is drawn from the most prominent part of the forehead to the alveolar edge of the upper jaw, opposite the incisor teeth—the *facial line*—and the other from the meatus auditorius externus to the same point of the jaw. According to the size of the angle it has been attempted to appreciate the respective proportions of the cranium and face, and, to a certain extent, the degree of intelligence of individuals and of animals. In the white varieties of the species, this angle is generally 80°; in the negro not more than 70°, and sometimes only 65°. As we descend the scale of animals, the angle becomes less and less; until, in fishes, it nearly or entirely disappears. Animals which have the snout long, and facial angle small, such as the snipe, crane, stork, &c., are proverbially foolish, at least they are so esteemed; whilst intelligence is ascribed to those in which the angle is more largely developed, as in the elephant and the owl. In these last animals, however, the large facial angle is caused by the size of the frontal sinuses:—so that this mode of appreciating the size of the brain is very inexact, and cannot be depended upon.

The following is a table of the angle in man and certain animals:

Man.....	from 65° to 85° and more.
Sapajou.....	65
Orange-Utang.....	56 to 58
Guenon.....	57
Mandrill.....	30 to 42
Coati.....	28
Pole-Cat.....	31
Pug-Dog.....	35
Mastiff.....	41
Hare.....	30
Ram.....	30
Horse.....	23

ANGLE, OCCIP'ITAL, OF DAUBENTON, is formed by a line drawn from the posterior margin of the foramen magnum to the inferior margin of the orbit, and another drawn from the top of the head to the space between the occipital condyles. In man, these condyles, as well as the foramen magnum, are so situate, that a line drawn perpendicular to them would be a continuation of the spine; but in animals they are placed more or less obliquely; and the perpendicular is necessarily thrown farther forward, and the angle rendered more acute.

ANGLE, OP'TIC, (F.) *Angle optique ou visuel*, is the angle formed by two lines, which shave the extremities of an object, and meet at the centre of the pupil.

ANGLE, VISUEL, Angle, optic.

ANGNAIL, Hangnail.

ANGOISSE, Angor.

ANGO'LAM. A very tall Malabar tree, which possesses vermifuge properties.

ANG'ONE, properly *An'chonē*, (αγκων, from αγκυ, 'I choke.') *Anad'romē*, *Hysterapopnē'is*, *Hysteropnē'is*, *Præfoca'tio Fau'cium* seu *Uteri'na* seu *Matri'cia*, *Strangula'tio* seu *Præfoca'tio uteri'na*, *Suffoca'tio uteri'na* seu *hyster'ica*, *Glo'bus* seu *Glo'bulus* seu *No'dus hyster'icus*, *Orthopnē'a hyster'ica*, *Dyspha'gia globo'sa* seu *hyster'ica*, *Ner'vous Quin'sy*. A feeling of strangulation, with dread of suffocation. It is common in hysterical females, and is accompanied with a sensation as if a ball arose from the abdomen to the throat.

ANG'OR, gen. *Ango'ris*, *An'guish*, (F.) *Angoisse*. Extreme anxiety, accompanied with painful constriction at the epigastrium, and often with palpitation and oppression. It is frequently an unfavourable symptom.

ANGOR, Agony, Orthopnœa—a. Fau'cium, *Is'th-mitis*—a. Pectoris, *Angina pectoris*.

ANGOS, (*ayyos*), Uterus, Vessel.

ANGOSTURE VRAIE, ('true angustura,') *Cusparia febrifuga*.

ANGOURION, (*αγγουριον*, from *ayyos*, 'a hollow vessel,') *Cucumis sativus*.

ANGRY, see Inflamed.

ANGU'NUM, (*anguis*, 'a snake,') A snake's egg. Supernatural powers were ascribed to it by the ancient Britons.

ANGUIS, Serpent.

ANGUIS, Angor.

ANGUISH, FE'BRILE, *Angor Febrilis*. The combination of weariness, pain, anxiety, and weakness affecting the head and neck, which is so generally observed at the commencement of fever.

ANGULAIRE, Angular—a. *de l'Omo'plate*, *Levator scapulæ*.

ANG'ULAR, *Angula'ris*, (from *angulus*, 'an angle,') (F.) *Angulaire*. That which relates to an angle.

ANGULAR AR'TERY AND VEIN. A name given, 1. to the termination of the facial artery and vein, because they pass by the greater angle of the eye; and, 2. to the facial artery and vein themselves, because they pass under the angle of the jaw. See Facial.

ANGULAR NERVE is a filament furnished by the inferior maxillary, which passes near the greater angle of the eye.

ANGULAR PROC'ESSES of the frontal bone are seated near the angles of the eyes. See Orbitar.

ANGULARIS, *Levator scapulæ*.

ANGULI-SCAPULO-HUMÉRAL, *Teres major*.

ANGULUS COSTÆ, see Costa—a. *Ocularis*, *Canthus*.

ANGURIA, (*αγγουριον*, from *ayyos*, 'a hollow vessel,') *Curebita citrullus*.

ANGUSTATIO, (*angustus*, 'narrow,') *Arctatio*—a. *Cordis*, *Systole*—a. *Intestini recti vel Ani*, *Stricture of the rectum*.

ANGUS'TIA, *Angusta'tio*, *Stenochor'ia*. Anxiety, narrowness, strait, constriction.

ANGUSTIA ABDOMINALIS, *Pelvis*, (Brim)—a. *Perinaealis*, *Pelvis*, (Outlet.)

ANGUSTURA, *Cusparia febrifuga*—a. *False*, *Brucea antidysenterica*, and *Strychnos nux vomica*—a. *Spuria*, *Brucea antidysenterica*, and *Strychnos nux vomica*.

ANGUSTURE, FAUSSE, *Brucea antidysenterica*—a. *Ferrugineuse*, *Brucea antidysenterica*—a. *Vraie*, *Cusparia febrifuga*.

ANGUSTURINE, *Bruceine*.

ANHÆMATOSIA, (an, and *hæmotosis*.) *Asphyxia*, *Anæmia*.

ANHÆMIA, *Anæmia*.

ANHAPHIA, *Anaphia*.

ANHELAT'IO, (*anhelare*, *anhelatum*, 'to pant,') *Anhelitus*, *Aas'mus*, *Pant'ing*, *Gasp'ing*, *Anhelat'ion*, (Sc.) *Hech*, *Hegh*, (Prov.) *Gien*, *Kaw'ing*, *Pech'ing*, (F.) *Essoufflement*. Short and rapid breathing. See *Dyspnœa*.

Anhelatio is sometimes employed synonymously with *asthma*.

ANHELITUS, *Anhelation*, *Breath*.

ANHÆMATOSIE, (an, and *hæmotosis*.) *Anæmia*, *Asphyxia*.

ANHÉMIE, *Anæmia*.

ANHIS'TOUS, (an, and *iseros*, 'organic texture,') *Anorgan'ic*, *Amor'phus*. The tunica decidua uteri is termed by Velpeau the *anhistous membrane*.

ANHUIBA, *Laurus sassafras*.

ANHYDRÆMIA, *Anemyl'ria*, (an, *idw*, 'water,' and *huma*, 'blood,') A condition of the blood in which there is a diminution in the quantity of the serum.

ANHYDROMYEL'IA, (an, *idw*, 'water,' and

αἷμος, 'marrow.') Deficiency or absence, in the spinal cavity, of the cephalo-spinal fluid.

ANICE'TON, *Anice'tum*, *Mesia'nium*, (ἀνικητόν, from *an*, and *νίκη*, 'victory,' 'invincible.') A plaster much extolled by the ancients in cases of achores. It was formed of litharge, cerusse, thus, alum, turpentine, white pepper, and oil.

ANID'EUS, (an, and εἶδος, 'shape.') *Amorphus*. A monster devoid of shape.

ANIDRO'SIS, (ἀνιδρώσις, from *an*, ἰδρώς, 'sweat,' and *osis*.) *Sudo'ris null'itas* vel *priva'tio*. Absence of sweat. Deficiency of perspiration.

ANIL, *Indigofera anil*.

ANILEMA, (ἀνίλημα, from *ana*, and εἰλω, 'I twist.') Borborygmus, Tormina.

ANILESIS, (ἀνίλησις,) Borborygmus, Tormina.

AN'ILINE, *Anil'ina*. An artificial alkaloid, called after *Anil*, *Indigofera anil*, which furnishes indigo. It is an oily liquid, which distils over when finely-pulverized indigo is decomposed by a concentrated solution of potassa in a retort. It is also present in coal tar, from which it is extensively prepared to make certain dyes. The sulphate of aniline has been prescribed in many cases of nervous disorder, as in chorea. Dose, a grain three times a day.

ANILITAS, (*anus*, 'an old woman.') See Dementia.

AN'IMA, *An'imus*, *Mens*, *Psy'chê*. (F.) *Âme*. The mind, breath, &c., (ἀνέμος, 'wind or breath.') The principle of the intellectual and moral manifestations. Also, the principle of life,—the life of plants being termed *An'ima vegetati'va*, (F.) *Âme végétative*; that of man, *An'ima sensiti'va*, (F.) *Âme sensitive*.

The *Anima* of Stahl, *An'ima Stahlia'na*, was a fancied intelligent principle, which he supposed to preside over the phenomena of life,—like the *Archæus* of Van Helmont.

Under the term *Anima Mun'di*, the ancient philosophers meant a universal Spirit, which they supposed spread over every part of the universe.

The precise seat of the mind in the brain has given rise to many speculations. The point is unsettled.

With the ancient chemists, *Anima* meant the active principle of a drug separated by some chemical management.

ANIMA ALOES: see Aloes Succotorina—a. Articulatorum, Hermodactylus—a. Hepatis, Ferri sulphas—a. Pulmonum, Crocus—a. Rhei, Infusum rhei—a. Stahlianæ, see Anima—a. Vegetativa, Plastic force.

AN'IMAL, *Zo'ön*. A name given to every animated being. Most animals have the power of locomotion; some can merely execute partial movements, such as contraction and dilatation. In other respects it is often a matter of difficulty to determine what is an animal characteristic. The study of animals is called *Zoö'logy*.

AN'IMAL, *Anima'lis*, *Zoo'des*. That which concerns, or belongs to, an animal.

ANIMAL HEAT, *Cal'or anima'lis* seu *nati'vus*, *Cal'idum anima'lis* seu *innatum*, *Biolychnion*, *Flam'mula vita'lis*, *Ther'ma em'phytum*, *Ther'mum em'phytum*, *Ig'nis anima'lis* seu *natura'lis* seu *vita'lis*, (F.) *Chaleur animale*, is the caloric constantly formed by the body of a living animal, by virtue of which it preserves nearly the same temperature, whatever may be that of the medium in which it is placed. This formation seems to take place over the whole body, in the production of carbonic acid, and in other oxidizing processes, and to be connected with the changes in nutrition.

A table of the natural temperatures of many animals—that of man being 100°—is contained in the author's Human Physiology, vol. i., Art. Calorification.

ANIMAL KING'DOM, (F.) *Règne Animal*, comprises all animated beings.

ANIMAL LAY'ER, see *Tache embryonnaire*—a. Magnetism, see Magnetism, animal.

ANIMALCULA SEMINALIA, (pl. of *Animalculum*.) Spermatozoa—a. Spermatica, Spermatozoa.

ANIMAL'CULE, *Animal'culum*; (dim. of *animal*.) A small animal. An animal well seen only by means of the microscope.

ANIMALCULES, SEMINAL, Spermatozoa—a. Spermatic, Spermatozoa.

ANIMALCULISM, Spermatism.

ANIMAL'CULIST, *Animal'ist*. One who attempts to explain different physiological or pathological phenomena by means of animalcules. One who believes that the embryo is formed from the so-called spermatic animalcules—*sperm'atist*, *sem'inist*. See Seminist.

ANIMAL'CULOVISM, *Animalculovism'us*, (*animalculum* and *ovum*.) The doctrine, now universal, which maintains that the new being is formed by the concurrence of the spermatic animalcule or spermatozoid with the ovum.

ANIMALCULUM, *Animalcule*.

ANIMALIST, *Animalculist*.

ANIMAL'ITY, *Animal'itas*, *Zo'ism'us*, (F.) *Animalité*. The aggregate of qualities which distinguish that which is animated. That which constitutes the animal.

ANIMALIZA'TION, *Animalisa'tio*. The transformation of the nutritive parts of food into the living substance of the body to be nourished. Assimilation.

To AN'IMATE, *Anima'rē*. To give life to. The French use it in the sense of,—to excite or render active; as, *animer un vésicatoire*: to excite a blister to suppurate.

ANIMATIO, Animation—a. Fœtus, see Quickening.

ANIMA'TION, *Anima'tio*, *Zoö'sis*. The act of animating. The state of being enlivened.

ANIMATION, SUSPENDED, Asphyxia.

AN'IMÉ, *Gum an'imē*, *Aminē'a*, *Gum'mi an'imē*, *Can'camum*, *Can'camy*. A resin obtained from the trunk of *Hymenæ'a cour'baril*. Ord. Fabacææ. It has been given as a cephalic and uterine, but is not used. The plant is also called *Cour'baril*.

ANIMÉ, (F.) (from *animare*, 'to animate.') An epithet applied to the countenance, when florid, in health or disease.

ANIMELLE, Parotid.

ANIMI CASUS SUBITUS, ('sudden loss of mind.') Syncope—a. Deliquium, Syncope—a. Pathemata, Passions.

ANIMISM, see Animist.

AN'IMIST. One who, following the example of Stahl, refers all the phenomena of the animal economy to the soul, *An'imism*.

The *anima* or soul, according to Stahl, is the immediate and intelligent agent of every movement, and of every material change in the body. Stahl therefore concluded, that disease is nothing more than a disturbance or disorder in the economy, or an effort by which the soul, attentive to every morbid cause, endeavours to expel whatever may be deranging the habitual order of health.

ANIMUS, *Anima*, Breath.

ANIRID'IA, (an, and ἰρις, gen. ἰριδος, 'the iris.') Absence of the iris.

ANIS, *Pimpinella anisum*—a. *Aigre*, *Cuminum Cyminum*—a. *de la Chine*, *Illicium anisatum*—a. *Doux*, *Anethum*—a. *Étoilé*, *Illicium anisatum*.

ANISA'TUM, (from *Anisum*.) A sort of medicated wine, formerly prepared with honey, wine of Ascalon, and aniseed.

ANISCALPTOR, (*anus*, and *scalpere*, to scratch.) *Latissimus dorsi*.

ANISCHURIA, (an, and *ischuria*, 'suppression of urine.') Enuresis.

ANISE, *Pimpinella anisum*—a. Star, *Illicium anisatum*, l. *Floridanum*—a. Tree of Florida, *Illicium floridanum*—a. Tree, yellow-flowered, *Illicium anisatum*.

ANISEED, see *Pimpinella anisum*.

ANISI SEMINA, see *Pimpinella anisum*.

ANISO'DUS L'URIDUS, (*ανισος*, 'unequal,' and *οδους*, 'a tooth,' from the unequal divisions of the calyx.) *Nican'dra anom'ala*, *Physalis stramonium*, *White'ya stramonifo'lia*. A plant of Nepal, possessed of narcotic properties, and resembling belladonna and tobacco. It dilates the pupil, and is used in diseases of the eye like belladonna. It is given in alcoholic tincture (*dried leaves* 3j to alcohol f3viii). Dose, 20 drops internally in the 24 hours.

ANISOPHYLLUM IPECACUANHA, (*ανισος*, 'unequal,' and *φυλλον*, 'a leaf.') *Euphorbia Ipecacuanha*.

ANISOS'THENES, *Inequal'i rob'orē pol'lens*. That which is unequal in strength, (*ανισος*, 'equal,' and *σθενος*, 'strength.') An epithet applied particularly to the muscular contractility which, in the sick, is sometimes augmented in certain muscles only,—in the flexors, for example.

ANISOT'ACHYS, (*ανισος*, 'unequal,' and *ταχυς*, 'quick.') An epithet for the pulse, when quick and unequal.

ANISUM, (*ανισον*,) *Pimpinella anisum*—a. Africanum frutescens, *Bubon Galbanum*—a. Fruticosum galbaniferum, *Bubon Galbanum*—a. Officinale, *Pimpinella anisum*—a. Sinense, *Illicium anisatum*—a. Stellatum, *Illicium anisatum*—a. Vulgare, *Pimpinella anisum*.

ANKLE, *Astragalus*, *Malleolus*.

ANKLEY, *Malleolus*.

ANKUS, *Ancus*.

ANKYLO, see *Ancylo*.

ANKYLOBLEPH'ARON, (*Ancylobleph'aron*, *Palpebra'rum coal'itus*, (*ankylo*, and *βλεφαρον*, 'eyelid.') A preternatural union between the free edges of the eyelids. Likewise called *Symbpleh'aron*, *Symbplephar'osis*, and *Pros'physis*.

Also, union between the eyelids and the globe of the eye.

ANKYLOCHEI'LIA, *Ancylochei'lia*, (F.) *Ancylocheilie*, (*ankylo*, and *χειλος*, 'lip.') Accidental union of the lips.

ANKYLOCHILIE, *Ancylocheilia*.

ANKYLOCOLPUS, (*ankylo*, and *κολπος*, 'the vagina.') *Colpatresia*.

ANKYLOCORE, (*ankylo*, and *κορη*, 'the pupil.') *Coreclisis*.

ANKYLODON'TIA, (*ankylo*, and *οδους*, gen. *οδοντος*, 'a tooth.') An irregular position of the teeth in the jaws.

ANKYLOGLOS'SIA, *Ancyloglos'sia*, *Concre'tio lin'gue*, (*ankylo*, and *γλωσσα*, 'the tongue.') (F.) *Soubrelangue*. Impeded motion of the tongue in consequence of adhesion between its margins and the gums; or in consequence of the shortness of the frænum: the latter affection constituting *Tongue-tie*, *Olopho'nia lin'gue fren'a'ta*. It merely requires the frænum to be divided with a pair of scissors.

ANKYLOGLOSSOT'OMUM, (*ancyloglos'sia*, 'tongue-tie,' and *τομη*, 'incision.') An instrument used in the operation for tongue-tie.

ANKYLOME'LĒ, *Ancylome'lē*, (*ankylo*, and *μελη*, 'a probe.') A curved probe.

ANKYLOMERIS'MUS, *Ancylomeris'mus*, (*ankylo*, and *μερος*, 'a part.') Morbid adhesion between parts.

ANKYLOPS, (*αγκυ*, 'near,' and *ωψ*, 'the eye.') *Egriops*.

ANKYLORRHIN'IA, *Ancylorrhin'ia*, (*ankylo*, and *ριν*, 'the nose.') Morbid union of the parietes of the nose.

ANKYLOSED, see *Ankylosis*.

ANKYLO'SIS, *Ancylo'sis*, *Anchylo'sis*, *An'cylē*,

Stiff Joint, (*ankylo* and *osis*.) An affection, in which there is great difficulty or even impossibility of moving a diarthrodial articulation. It is so called, because the limb commonly remains in a constant state of flexion, and a joint thus affected is said to be *ankylo'sed*. Anchylosis is said to be *complete* or *true*, when there is an intimate adhesion between the synovial surfaces, with union of the articular extremities of the bones. In the *incomplete* or *false* anchylosis, there is obscure motion, but the fibrous parts around the joint are more or less stiff and thickened. In the treatment of this last state, the joint must be gently and gradually exercised; and oily, relaxing applications be assiduously employed.

ANKYLOSIS, BARTON'S OPERATION FOR, see *Barton*—a. *Spuria*, *Rigiditas articulorum*.

ANKYLO'TIA, *Ancylo'tia*, (*ankylo*, and *οτις*, gen. *ωτις*, 'the ear.') Morbid union of the parietes of the ear.

ANKYLOT'OMUS, *Ancylo'tomus*, (*ankylo*, and *τομειν*, 'to cut.') Any kind of curved knife. An instrument for dividing the frænum linguæ.

ANNEAU, (*anneau*, 'a ring,') Ring—a. *Crural*, *Crural canal*—a. *Diaphragmaticue*, *Diaphragmatic ring*—a. *Femoral*, *Crural canal*—a. *Inguinal*, *Inguinal ring*—a. *Ombilical*, *Umbilical ring*.

ANNEXE, Accessory, Appendix.

ANNI CRITICI, *Climacteriei* (anni)—a. *Decretorii*, *Climacteriei* (anni)—a. *Fatales*, *Climacteriei* (anni)—a. *Genethliaci*, *Climacteriei* (anni)—a. *Gradarii*, *Climacteriei* (anni)—a. *Hebdomadici*, *Climacteriei* (anni)—a. *Heroici*, *Climacteriei* (anni)—a. *Natalitii*, *Climacteriei* (anni)—a. *Scalares*, *Climacteriei* (anni)—a. *Scansiles*, *Climacteriei* (anni).

ANNOTAT'IO, *Episma'sia*. Under this term some have included the preludes to an attack of intermittent fever—as yawning, stretching, somnolency, chilliness, &c.

ANNOTTO, see *Terra Orleana*.

ANNUAL DISEASES, *Mor'bi an'nuī*, *M. anniver'sarii*, (F.) *Maladies annuelles*. A name given, by some, to diseases which recur every year about the same period. *Fe'bris an'nuā*, (F.) *Fièvre annuelle*, is a term used for a fancied intermittent of this type.

ANNUENS, *Rectus capitis internus minor*.

ANNUIT'IO, *Nod'ding*, (from *an*, and *nutus*, 'a nod.') A gesture, denoting assent in most countries. Also, the state of somnolency, when the individual is in the erect or sitting posture, with the head unsupported, in which the power of volition over the extensor muscles of the head is lost, and the head drops forward.

AN'NULAR, *Annula'ris*, *Anula'ris*, *Cricoi'des*, (F.) *Annulaire*, (*annulus*, 'a ring.') Any thing relating to a ring, or which has the shape or fulfils the functions of a ring.

ANNULAR FINGER, *Ring Finger*, *Ringman*, *Dig'itus annula'ris*, *Param'esos*. The fourth finger, so called from the wedding ring being worn thereon. See *Digitus*. Of old, it was believed, that there is a vein of direct communication between the ring finger and the heart. According to some of the missals, the ring must remain on this finger 'quia in illo digito est quedam vena procedens usque ad cor.'

ANNULAR GANGLION, see *Ciliary ligament*.

ANNULAR LIG'AMENT, *Trans'verse lig'ament*, *Cruc'ial ligament*. A strong ligamentous band, which arches across the area of the ring of the atlas, from a rough tubercle upon the inner surface of one articular process to a similar tubercle on the other. It serves to retain the odontoid process of the axis in connexion with the anterior arch of the atlas.

AN'NULAR LIG'AMENT OF THE RA'DIUS, is a very

strong fibro-cartilaginous band, which forms, with the lesser sigmoid cavity of the cubitus, a kind of ring, in which the head of the radius turns with facility.

AN'NULAR LIG'AMENTS OF THE CAR'PUS, *Articulæ nūl'ares membranosæ*, are two in number.

The one, *ante'rior*, is a broad, fibrous, quadrilateral band, extending transversely before the carpus, and forming the gutter, made by the wrist, into a canal. It is attached, externally, to the trapezium and scaphoides; and internally to the os pisiforme and process of the ulniforme. It keeps the tendons of the flexor muscles, median nerve, &c., applied against the carpus.

The *posterior* ligament is situate transversely behind the joint of the hand, and covers the sheaths of the tendons, which pass to the back of the hand. Its fibres are white and shining, and are attached, externally, to the inferior and outer part of the radius; internally, to the ulna and os pisiforme.

ANNULAR LIG'AMENTS OF THE TAR'SUS are two in number. The *ante'rior* is quadrilateral, and extends transversely above the instep. It is attached to the superior depression of the os calcis, and to the malleolus internus. It embraces the tendons of the extensor muscles of the toes, the *tibialis anticus* and *peroneus anticus*. The *inter'nal* is broader than the last. It descends from the malleolus internus to the posterior and inner part of the os calcis, with which it forms a kind of canal, enclosing the sheaths of the tendons of the *tibialis posticus*, *flexor longus digitorum pedis*, and *F. longus pollicis pedis*, as well as the plantar vessels and nerves.

ANNULAR VEIN, *Vena annula'ris*, is situate between the annular finger and the little finger. Aëtius recommends it to be opened in diseases of the spleen.

ANNULARIS, Cricoid: see Digitus—a. Ani, Sphincter ani.

ANNULI CARTILAGINEI, see Trachea—a. Cartilaginosi Tracheæ; see Trachea.

AN'NULI FIBRO-CARTILAGINEI, *Fibrocartilagin'ous* or *festooned rings*. Tough tendinous tracts in the heart, two of which, anterior, are situate on the right and left of the aortal opening; and one posterior, which runs backwards from the aorta to the border of the auriculoventricular septum, where it splits into two slender crura.

AN'NULI-TENDINO-PHALANGIENS, *Lumbricales manus*.

ANNULUS, see Daetylius, Vulva—a. Abdominis, Inguinal ring—a. Albidus, see Ciliary ligament—a. Cellulosus, Ciliary ligament—a. Ciliaris, Ciliary ligament—a. Fossæ ovalis, see Ovis fossa.

AN'NULUS CONJUNCTIVÆ. A slight circular swelling, formed by the sclerotic conjunctiva at the edge of the cornea, especially in old people, which extends for a little way over the cornea, particularly at its upper margin.

ANNULUS GANGLIFORMIS, see Ciliary ligament—a. Repens, Herpes circinatus—a. Umbilicalis, Umbilical ring—a. Ventriculi, Pylorus—a. Vieussenii, see Ovis fossa.

ANO, *ano*. A prefix denoting 'above, up.'

ANOC'AVERNEUX, (*anus*, and *cavernous* [bodies].) Accelerator urinae.

ANOCHEY'LOX, *Anochi'l'us*, (*ano*, and *χειλος*, 'lip.') The upper lip. Also, one who has a large upper lip.

ANOCHEILOS'CHESIS, (*anochailon*, 'the upper lip,' and *σχισις*, 'a cleaving.') Fissure of the upper lip, as in hare lip.

ANOC'HILUS, *Anochailon*.

ANOC'ELIA, (*ano*, and *κωλια*, 'belly.') Stomach.

ANOC'ELIADELPHUS, *Cœliadelphus*.

ANO'DIA, (*an*, and *ωδη*, 'song.') An unconnected or dissonant mode of speech.

ANOD'IC, *Anod'icus*, (*ano*, and *δός*, 'a way.') Tending upwards. An epithet applied by Dr. Marshall Hall to an ascending course of nervous action.

ANODIN, *Anodyne*.

ANODIN'IA, (*an*, and *ωδιν*, 'a labour pain.') Absence of labour pains.

ANODMIA, (*an*, and *οδμη*, 'smell.') Anosmia.

ANODON'TIA, (F.) *Anodontie*, (*an*, and *οδους*, gen. *οδοντος*, 'a tooth.') Absence of teeth.

ANODUS, *Edentulus*.

AN'ODYNE, (*ανωδυνος*,) *Anod'ynus*, *Antod'ynus*, *Antid'ynous* (improperly), *Paregor'icus*, *Anel'icus*, *Antal'gicus*, *Acesod'ynus*, (F.) *Anodin* ou *Anodyn*, (*an*, and *οδυν*, 'pain.') *Anodynæ* are those medicines which relieve pain, or cause it to cease; as opium, belladonna, &c. They act by blunting the sensibility of the encephalon, so that it does not appreciate the morbid sensation.

ANODYNE, HOFFMANN'S, *Spiritus Ætheris sulphurici compositus*.

ANODYN'IA, (*ανωδυνια*,) *Indolent'ia*. Cessation or absence of pain. Vogel has given this name to a genus of diseases, characterized by a cessation of pain, and the exasperation of other symptoms; as we see in gangrene.

ANODYNUM MINERALE, *Potassæ nitratis sulphatis paucillo mixtus*.

ANØ'A, *Anoia*, (*avoia*, from *a*, and *νοος*, 'mind.') Delirium, imbecility. See Dementia and Idiotism.

ANOESIA, (*ανωσια*,) *Dementia*—a. *Adstricta*, *Melancholy*.

ANOIA, (*ανοια*,) *Anœa*.

ANOMAL, *Anomalous*.

ANOMA'LIA, (*an*, and *μαλος*, 'regular.') *Abnorm'ity*, *Abnorm'itas*, *Alienatio*. (F.) *Anomalie*, *Anomaly*, irregularity. In Pathology, anomaly means something unusual in the symptoms proper to a disease, or in the morbid appearances presented by it.

ANOMALIA NERVORUM, *Nervous diathesis*.

ANOMALIE, *Anomaly*—a. *par Cloissonnement*, see *Cloissonnement*.

ANOMALIS, *Anomalous*.

ANOMALOTROPHIES, (*an*, *μαλος*, 'regular,' and *τροφη*, 'nourishment.') A class of diseases, which consist in modifications of the nutrition of organs.

ANOM'ALOUS, *Anom'alus*, *Anom'alis*; same etymon as *Anomalia*. Irregular; contrary to rule. (F.) *Anomal*. In Medicine, a disease is called *anomalous*, in whose symptoms or progress there is something unusual. Affections are also called *anomalous*, which cannot be referred to any known species. See Irregular.

ANOMMATUS, (*an*, and *ομμα*, 'eye.') *Anophthalmus*.

ANOMOCEPH'ALUS, (*a*, *νομος*, 'rule,' and *κεφαλη*, 'head.') One whose head is deformed.

ANOMOSPLE'NOTOPIE, (*a*, *νομος*, 'rule,' *σπλην*, 'spleen,' and *τοπος*, 'place.') *Splenectopia*.

ANOMOTOPIE, (*a*, *νομος*, 'rule,' and *τοπος*, 'place.') *Ectopia*.

ANOM'PHALUS, (*an*, and *ομφαλος*, 'the navel.') One devoid of navel. Many writers have endeavoured to show that Adam and Eve must have been *ανομφαλοι*, as they could not have had umbilical vessels.

ANONA TRIPET'ALA. A tree of the family *Anonæ* or *Anonacæ*; *Sex. Syst.* Polyandria Polygynia; from fifteen to twenty feet high, native of South America, which bears a delicious fruit called *Chirimoya*. Both the fruit and flowers emit a fine fragrance, which, when the tree is covered with blossoms, is almost overpowering.

ANONA TRILOBA, see *Carica papaya*.

ANONIS, (ἀνώνις,) Ononis.

ANONYCHIA, (ἀν, and ονυχ, gen. ονυχος, 'a nail.') Want of nails,—a rare congenital defect.

ANONYME, Anonymous, see Innominatum.

ANONYMOUS, *Anonymus*, *Innominatus*, (F.) *Anonyme*, (an, and ονομα, 'name.') That which has no name.

The word has been applied to many parts of the body:—to the *Anonymous bone* or *Os innominatum*:—the *Anonymous foramen* or *Foramen innominatum*, &c.

ANOPHTHALMUS, *Anom'matus*, (an, and οφθαλμος, 'an eye.') One who is devoid of eyes. The condition is *Anophthalmia*.

ANOP'IA, (an, and οψις, 'vision.') A case of monstrosity in which the eye and orbit are wanting; also, *Cæcitas*.

ANOPTICNERVIE, Amaurosis.

ANORCHI, (ανωρχοι,) Anorchides.

ANORCHIDES, pl. of *Anorchis*, *Anorchi*, (from an, and ορχις, gen. ορχιδος, 'a testicle.') They who are without testicles.

ANOREX'IA, (an, and ορεξις, 'appetite.') *Inappeten'tia*, *Limo'sis ex'pere*, (F.) *Perte d'appetit*. Absence of appetite, without loathing. Anorexia or want of appetite is symptomatic of most diseases. Also, *Indigestion*, *Dyspepsia*.

ANOREXIA EXHAUSTORUM, Frigidity of the stomach—a. *Mirabilis*, *Fasting*.

ANORGANIC, see Anhistous, and Inorganic.

ANORMAL, Abnormal.

ANORTHOP'IA, (an, ορθος, 'upright,' and οψις, 'vision.') Vision in which persons are unable to judge when objects are not parallel or not symmetrical.

ANORTHOSIS, (an, and ορθος, 'upright.') Erection.

ANOS'IA, (ανωσια, from a, and νοσος, 'disease.') Health. Freedom from disease.

ANOS'MIA, (an, and οσμη, 'odour.') Loss of smell. Called, also, *Anosphre'sia*, *Anosphra'sia*, *Paros'mia*, *Anod'mia*, *Anosmo'sia*, *Olfactus amis'sio seu defic'iens*, *Dysasthes'ia olfacto'ria*, *Anasthes'ia olfacto'ria*, *Odorat'us deper'ditus*, (F.) *Perte de l'Odorat*.

ANOSMOSIA, Anosmia.

ANOSPHERASIA, (an, and σφρασια, 'smell,') Anosmia.

ANOSPHERSIA, (an, and σφρησις, 'smell,') Anosmia.

ANOT'US, (an, and ος, gen. ωτος, 'the ear.') A monster having no ears.

ANOXÉMIE, (an, ox for 'oxygen,' and άίμα, 'blood,') Asphyxia.

ANSA, *Anse*—a. *Intestinalis*, *Anse intestinale*.

ANSE (F.), *Ansa* (L.), signifies, properly, the handle of certain vessels, usually of an arched form. By analogy, it has been applied to that which is curved in the form of such handle. Thus, the French speak of *Anse intestinale*—*Ansa seu Gyrcus intestinal'is*—to signify a portion of intestine, supported by its mesentery, and describing a curved line:—also, of *Anse nerveuse*, *Anse anastomotique*, &c.

Anse de fil is used, in Surgery, to designate a thread, curved in the form of an *Anse*.

ANSERINA, (anser, 'a goose,') *Potentilla anserina*.

ANSÉRINE, *Chenopodium ambrosioides*—a. *Anthelmintique*, *Chenopodium anthelminticum*—a. *Bon Henri*, *Chenopodium Bonus Henrius*—a. *Botrys*, *Chenopodium Botrys*—a. *Fétide*, *Chenopodium vulvaria*—a. *Vermifuge*, *Chenopodium anthelminticum*.

ANTAC'IDS, *Antidids*, *Antiac'ida*, *Inverten'tia*, (anti, and acida.) Remedies which obviate acidity of the stomach. They are chemical agents, and act by neutralizing the acid. Those chiefly

used are ammonia, calcei carbonas, calx, magnes'ia, magnes'ia carbonas, potassa, potassu bicarbonas, p. carbonas, sodæ bicarbonas, and s. carbonas. They are, of course, only palliatives, removing that which exists, not preventing the formation of more.

ANTAG'ONISM, *Antagonis'mus*, *Antis'tasis*, (anti, and αγωνίζομαι, 'to strive.') Action in an opposite direction. It applies to the action of muscles that act in a contrary direction to others. In estimating the force of muscles, this antagonism must be attended to.

ANTAG'ONIST, *Antagonis'ta*. A muscle whose action produces an effect contrary to that of another muscle. Every muscle has its antagonist, because there is no motion in one direction without a capability of it in another.

ANTALGICUS, (anti, and αλγος, 'pain,') Anodyne.

ANTAPHRODIS'IA, *Antaphrodit'ic*, *Antaphrodisi'acus*, *Anaphrodisi'acus*, *Anaphrodisi'iac*, *Antero'ticus*, (anti, and αφροδισιακος, 'aphrodisiac,') A substance capable of blunting the venereal appetite.

ANTAPHRODITIC, Antaphrodisiac.

ANTAPOD'OSIS, (ανταποδοσις, from ανταποδιδωμι, 'I return in exchange,') The succession and return of the febrile periods.

ANTAPOPLECTICUS, Antiapoplectic.

ANTARTHRETIC, Antiarthritic.

ANTASTHENICUS, (anti, and ασθενεια, 'debility,') Tonic.

ANTASTHMATICUS, Antiasthmatic.

ANTATROPH'IC, *Antatroph'icus*, *Antat'rophus*, *Antiatroph'icus*, (anti, and τροφια, 'atrophy,') A remedy opposed to atrophy or consumption.

ANTE (L.), 'before,' (F.) *Avant*, as a prefix, 'before, fore.'

ANTECEDEN'TIA, (ante, and cedo, 'I go,') The precursory or warning symptoms of a disease.

ANTEFLEXIO UTERI, (ante, and flectere, 'to bend,') See Anteversion.

ANTELA'BIA, *Proche'il'a*, (ante, 'before,' and labia, 'the lips,') The extremity of the lips. See Antilabia and Procheilon.

ANTELOPE, Antelope.

ANTEMBALLOMENUM, (αντεμβαλλονεν, from αντεβαλλω, 'I substitute,') Succedaneum.

ANTEM'BASIS, (αντεμβασις, from anti, and εμβασις, 'I enter,') *Mu'tuus ingres'sus*. The mutual reception of bones.

ANTEMETIC, Antiemetic.

ANTENDEIXIS, (anti, and ενδειξις, 'indication,') Counterindication.

ANTENDIXIS, Counterindication.

ANTENEAS'MUS, (anti, and τεινω, 'I extend or stretch,') One furious against himself. Mania, in which the patient attempts his own life.

ANTENNA'RIA DIOICA, *Gnaphal'ium Dioicum*, *Hispid'ula*. *Pes ca'ti*, *El'chrysium montanum*, *Dio'cium* *Everlast'ing*, *Cuts'foot*, (F.) *Pied de chat*. Order, Compositæ. A common European plant, which has been advised in hemorrhage, diarrhœa, &c.

ANTENNARIA MARGARITACEA, *Gnaphalium margaritaceum*.

ANTENNARIA PLANTAGINIFO'LIA, *Gnaphal'ium plantaginifol'ium* seu *plantagin'eum* seu *dioicum*—variety *plantaginifolium*, *Plantain Life everlast'ing*, *Cudweed*, indigenous, has similar properties.

ANTE PARTUM, 'before delivery,' as 'ante-partum hemorrhage.'

ANTEPHIALTIC, Antiephialtic.

ANTEPILEPTIC, Antiepileptic.

ANTEPONENS, (ante, and ponere, 'to place,') Antiepiating.

ANTEREI'SIS, (αντερεισις, from anti, and εριδω, 'I support,') The resistance—the solidity—of bones.

ANTÉRIEUR, Anterior—*a. du Marteau*, Laxator tympani—*a. de l'Oreille*, Anterior auris.

ANTÉRIOR, Anticus, (F.) Antérieur, (ante, 'before,') Situate before. Great confusion has prevailed with anatomists in the use of the terms *before*, *behind*, etc. Generally, the word *anterior* is applied to parts situate before the median line, the body being in the erect posture, with the face and palms of the hands turned forwards, and the feet applied longitudinally together.

ANTÉRIOR AURIS (*Muscle*), *Auricula'ris antérior*, *Al'trahens auriculam*, (F.) *Auriculaire antérieur*, *Antérieur de l'oreille*, *Zygomato-auriculaire*, *Z. auriculaire*. A small muscle, passing from the posterior part of the zygoma to the helix. *Use*, to draw the ear forwards and upwards.

ANTERIOR MALLEI, Laxator tympani.

ANTEROTICUS, (*anti*, and *eros*, gen. *eros*;*s*, 'love,') Antaphrodisiac.

ANTEUPHORBIIUM, *Cacalia antephorbium*.

ANTEVER'SION, *Antever'sio*, *Antrover'sio*, (ante, and *vertere*, *versum*, 'to turn,') Displacement of the uterus, in which the fundus is turned towards the pubes, whilst its orifice is towards the sacrum. It may be caused by extraordinary size of the pelvis, pressure of the viscera on the uterus, etc.; and is recognized by examination *per vaginam*. A simple forward inclination of the body of the uterus, without the os uteri being carried much backwards, is termed *Anteflex'ion of the uterus*, *Anteflex'io uteri*. Not unfrequently, however, it is used synonymously with *Retroversion of the Uterus*. See *Retroversio uteri*.

ANTHEMOPHYTICUS, *Anthemophyticus*, (*anti*, and *hæmoptysis*,) Against spitting of blood. A remedy for spitting of blood—*anthemophyticum* (*remedium*).

ANTHEMORRHAGICUS, *Anthemorrhagic*.

ANTHECTICUS, *Antiectic*.

ANTHELITRAGUS, (F.) *Anthélitragien*. One of the proper muscles of the pavilion of the ear.

ANTHELIX, *Antihelix*, (*anti*, and *ἄλξ*, 'the helix,') An eminence on the cartilage of the ear, in front of the helix, and extending from the concha to the groove of the helix, where it bifurcates.

ANTHELMINTIC, *Anthelmin'ticus*, *Antiscol'icus*, *Anthelmin'thicus*, *Antiscol'ethicus*, *Helmin'thicus*, *Helminthog'yus*, *Antivermino'sus*, *Vermif'ugus*, *Vermifuge*, (*anti*, and *ἄνυς*, *ἀνύσος*, 'a worm,') A remedy which destroys or expels worms, or prevents their formation and development. Anthelmintics may be divided into the *mechanical* and the *true*. To the former belong emetics, cathartics, mucuna, pulvis stanni, &c. To the latter, oleum terebinthine, chenopodium anthelminticum, spigelia, filix mas, radix granati, Brayera anthelmintica, &c. The great object, however, is to prevent the generation of the worms. This must be done by generous diet, fresh air, tonics, and everything that is calculated to invigorate the system. See *Worms*.

ANTHEMA ERUPTIO, gen. *Anthem'atis*, Exanthem.

ANTHEMIS, gen. *Anthem'idis*, *Anthemis nobilis*.

ANTHEMIS COT'ULA, (*ανθεω*, 'I flower,') *A. fetida* seu *Novboracensis*, *Cot'ula*, *C. fetida*, *Cot'a*, *Cynan'themis*, *Chamæ'melum fetidum* seu *Chamomilla spur'ia* seu *fetida*, *Mari'ta Cot'ula*, *Mayflower*, *Mayweed*, *Stinking Cham'omile*, *Wild Cham'omile*, *Dog's fen'el*, *Dilly*, *Dill'weed*, *Fieldweed*, *Piss'weed*. Ord. Compositæ Corymbiferae. See *Syst. Syngenesia Superflua*. (F.) *Maroute*, *Camomille fétide*, *Camomille puante*.

This plant, *Cot'ula* (Ph. U. S.), has a very disagreeable smell, and the leaves have a strong, acrid, bitterish taste. It is reputed to have been useful in hysterical affections.

ANTHEMIS FETIDA, *A. cotula*.

ANTHEMIS NO'BILIS, *Anthemis*, *A. odor'ata*, *Chamæ'melum*, *Ch. Nob'ile* seu *odor'atum*, *Chamomilla Rom'na*, *Euan'themon*, *Leucan'themum*, *Cham'omile*, (F.) *Camomille Romaine*. The leaves and flowers,—*Ant'hemis*, Ph. U. S.—have a strong smell, and bitter, nauseous taste. The flowers are chiefly used. They possess tonic and stomachic properties, and are much given as a pleasant and cheap bitter. A simple infusion is taken to produce, or to assist vomiting. Externally, they are often used in fomentations.

The *O'leum Anthem'idis* possesses the aromatic properties of the plant, but not the bitter and tonic. Consequently, the '*Camomille Drops*,' as sold by the druggists, must be devoid of the latter qualities. They are made by adding *Ol. anthem.* f 3j. to *Sp. vini rectif.* Oj.

ANTHEMIS NOVEBORACENSIS, *A. cotula*—*a. Odorata*, *A. cotula*.

ANTHEMIS PY'RETHRUM, *Py'rethrum*, *P. ve'r'um*, *Anagy'ellus pyrethrum*, *Bupthalamum Coctivum*, *Denta'ria*, *Her'ba saliv'aria*, *Pes Alexandri'aus*, *Span'ish Cham'omile*, *Pell'tory of Spain*, (F.) *Py-rèthre*, *Racine salivaire*, *Salivaire*, *Pied d'Alexandre*. The root, *Pyrethrum* (Ph. U. S.), is hot and acrid, its acrimony residing in a resinous principle. It is never used except as a masticatory in toothache, rheumatism of the face, paralysis of the tongue, etc. It acts as a powerful sialogogue.

The Pellitory of the shops in Germany is said to be derived from *Anagy'ellus officin'um*; a plant cultivated in Thuringia for medicinal purposes.

ANTHEMIS TINCTO'RIA, *Bupthalam' Her'ba*, *Dy'er's Cham'omile*, a European plant, has a bitter and astringent taste, and has been regarded as stomachic and vulnerary. (F.) *Camomille des Teinturiers*, *Ceil de Boeuf*.

ANTHEMIS VULGARIS, *Matricaria Chamomilla*.

ANTHE'RA, (*ανθρ'ος*, 'flourishing,' from its 'florid' colour.) A remedy compounded of several substances—myrrh, sandarac, alum, saffron, etc. It was used under the form of liniment, collyrium, electuary, and powder.

ANTHEREON, (same etymon, because the beard flourishes on the chin?) *Mentum*.

ANTHORA, *Aconitum anthora*—*a. Vulgaris*, *Aconitum anthora*.

ANTHORIS'MA, gen. *Anthorism'atis*, (*anti*, and *ὁραμα*, 'boundary,') *Tu'mor diff'usus*. A tumour without any defined margin.

ANTHOS, (*ανθος*, 'a flower,') see *Rosmarinus*—*a. Sylvestris*, *Ledum sylvestre*.

ANTHRA'CIA, (*ανθραξ*, gen. *α-θρακος*, 'coal,') *Carbuncular Exanthem*. An eruption of tumours, imperfectly suppurating, with indurated edges, and, for the most part, a sordid and sanious core. A genus in the order *Exanthematica*, class *Hæmatica* of Good, and including *Plague* and *Yaws*.

ANTHRACIA, *Anthracosis*—*a. Pestis*, *Plague*—*a. Rubula*, *Frambœsia*.

ANTHRACION, see *Anthrax*.

ANTHRACOID, *Anthracoï'des*, (*ανθραξ*, gen. *α-θρακος*, 'coal,' and *ειδος*, 'resemblance,') (F.) *Charbonneux*. As black as coal. Accompanied by, or resembling anthrax.

ANTHRACOMA, *Anthrax*.

ANTHRACONECROSIS, (*νεκρος*, 'death,') see *Sphacelus*.

ANTHRACOPHLYCTIS, (*phlyctis*, *phlyctæna*,') see *Anthrax*.

ANTHRACOSIA, *Anthrax*.

ANTHRACOSIS, *Anthrac'ia*, *Car'bo Palpebrarum*, (from *anthrax*, and *osis*.) A species of carbuncle, which attacks the eyelids and globe of the eye. Also, a carbuncle of any kind. It has been used for the "black lung of coal miners," which is induced by carbonaceous accumulation in the lungs; *Pseudo-melanotic formation* of Carswell.

Matière noire des Poumons, Charbon pulmonaire. When ulceration results from this cause, *black phthisis*, (F.) *Phthisie avec Mélanose*, exists. See *Melanosis*.

ANTHRACOSIS PULMONUM, see *Melanosis*.

ANTHRACOTYPHUS, (anthrax, and typhus,) Plague.

ANPHRAKOK'ALI, *Lithanthrakok'ali*, (αἰθαλ, gen. αἰθαλός, 'coal,' and kali, 'potassa.') An article introduced as a remedy in cutaneous diseases. It is formed by dissolving carbonate of potassa in 10 or 12 parts of boiling water, and adding as much slacked lime as will separate the potassa. The filtered liquor is placed on the fire in an iron vessel, and suffered to evaporate, until neither froth nor effervescence occurs, and the liquid presents a smooth surface like oil. To this, levigated coal is added in the proportion of 160 grammes to 192 grammes of potassa. The mixture is stirred, and removed from the fire, and the stirring is continued, until a black homogeneous powder results. A sulphuretted anthrakokali is made by mixing accurately 16 grammes of sulphur with the coal, and dissolving the mixture in the potassa, as directed above. The dose of the simple and the sulphuretted preparations is about two grains three times a day.

ANTHRAX, gen. *Anthracis*, (αἰθαλ, gen. αἰθαλός, 'a coal.') *An'trax*, *Car'bo*, *Rubi'us ve'l-rus*, *Cudsel'la*, *Erythe'ma gangræno'sum*, *Grantris'tum*, *Prud'na*, *Py'ra*, *Granatris'tum*, *Phy'ma An'thrax*, *Erythe'ma an'thrax*, *Carbun'culus*, *Anthrac'o'sia*, *Anthrac'o'ma*, *Absee's'sus gangræne's seu gangræno'sus*, *Furun'culus malign'us*, *F. gangræno'sus*, *Car'buncle*, (F.) *Charbon*. An inflammation, essentially gangrenous, of the cellular membrane and skin, which may arise from an internal or external cause. The definition by Dr. Good is sufficiently distinctive. Tumour, common to the surface; flat; firm; burning; penetrant; livid and vesicular; or crusty above, with a sordid and gangrenous core below; imperfectly suppurative. It sometimes prevails epidemically; is often owing to a peculiar cachectic condition, and may be induced by the reception of an animal infection into the organism; when it has received the names *Anthrax malign'us*, *Cacant'hrax*, *Anthrac'ion*, *Vesic'ula gangræne's cens*, *Anthracophyl'itis*, *Pustula malign'na*, (F.) *Pustule maligne*, *Bouton malin*, and is characterized at the outset by a vesication or bleb filled with a sero-sanguinolent fluid, under which a small induration is formed, surrounded by an areolar inflammation, which becomes gangrenous. It has been thought by some to be induced altogether by contact with the matter of the carbuncle of animals, or of the exuviae of the bodies of such as had died of the disease, but is now known to arise primarily in the human subject. This form of carbuncle has received different names, many of them from the places where it has prevailed:—*Carbunc'ulus contagio'sus seu Gall'ic'us seu Hunga'ric'us seu Polon'ic'us seu Septentriona'lis*, *Per'sic'us ign'is*, *Mor'bus pustulo'sus Fin'ic'us*, *Pustula gangræno'sa seu lic'ida Estho'nia*, *Pem'phigus Hunga'ric'us*, *Puce de Bourguigne ou maligne*, *Feu Persique*, *Malcat de Languedoc*.

Anthrax is a malignant boil, and its treatment is similar to that which is required in case of gangrene attacking a part. Laying open the part freely is one of the best remedies.

ANTHRAX MALIGNUS, see Anthrax—a. Pulmonum, *Neuropneumonia*.

ANTHRISCUS CEREFOLIUM, *Scandix cerefolium*—a. *Humilis*, *Chærophyl'ium Sylvestre*—a. *Procureus*, *Chærophyl'ium Sylvestre*.

ANTHROPE. (ανθρωπην.) Cutis.

ANTHROPIATRICA (MEDICINA), (anthropo, and iatrica, 'medicine.') Medicine applied to man, in contradiction to animals.

ANTHROPO, (ανθρωπος, 'man;') in composition, man.

ANTHROPOCHEMIA, Chymistry (human).

ANTHROPOCHYMY, Chymistry (human).

ANTHROPOGEN'IA, *Anthropogen'esis*, *Anthropog'eny*, *Genean'thropy*, (anthropo, and γενεα, 'generation.') The knowledge, or study, or phenomena of human generation.

ANTHROPOG'RAPHY, *Anthropograph'ia*, (anthropo, and γραφη, 'a description.') Anthropology. A description of the human body.

ANTHROPOLD, *Anthropoi'des*, (anthropo, and αἰος, 'resemblance.') Resembling man:—as the ape.

ANTHROPOL'ITHUS, (anthropo, and λιθος, 'a stone.') The petrification of the human body or of any of its parts. Morbid concretions in the human body.

ANTHROPOL'OGY, *Anthropolog'ia*, (anthropo, and λογος, 'a discourse.') A treatise on man. By some, this word is used for the science of the structure and functions of the human body. Frequently, it is employed synonymously with the *Natural History*, and the *Physiology of Man*.

ANTHROPOMAGNETISMUS, see Magnetism, animal.

ANTHROPOMANCY, *Anthropomanti'a*, (anthropo, and μαντεια, 'divination.') Divination by inspecting the entrails of a dead man.

ANTHROPOMET'RY, *Anthropomet'ria*, (anthropo, and μετρον, 'measure.') Measurement of the dimensions of the different parts of the human body.

ANTHROPOMORPHOL'OGY, (anthropo, μορφη, 'form,' and λογος, 'a discourse,') Anatomy (descriptive). A treatise on human anatomy.

ANTHROPOMORPHUS, (anthropo, and μορφη, 'shape,') *Atropa mandragora*.

ANTHROPON'OMY, *Anthroponom'ia*, (anthropo, and νομος, 'law,') A knowledge of the special laws which preside over the functions of the human body.

ANTHROPOPH'AGUS, (F.) *Anthropophage*, (anthropo, and φάγω, 'I eat,') A name given to one who eats those of his own species.

ANTHROPOPH'AGY, *Anthropophag'ia*, (same etymon.) The custom of eating human flesh. A disease in which there is great desire to eat it.

ANTHROPOS, (ανθρωπος,) Homo.

ANTHROPOSCOPIA, Physiognomy.

ANTHROPOTOMY, Andranatomia.

ANTHUS, (ανθος,) Flos.

ANTHYPNOT'IC, *Anthyppnot'icus*, *Antihypnot'ic*, *Agrypnot'ic*, (anti, and ὑπνος, 'sleep,') A remedy for stupor.

ANTHYPOCHON'DRIAC, *Anthyppochondriac'us*, (anti, and ὑποχονδριακος, 'hypochochondriac,') A remedy for hypochondriasis.

ANTHYSTER'IC, *Antihyster'ic*, *Antihyster'ic'us*, (anti, and ὕστερα, 'the uterus,') A remedy for hysteria.

ANTI, (αντι, 'against,') as a prefix, generally means 'opposition.'

ANTIADÉS, (αριαδες,) Tonsils.

ANTIADITIS, (antiades, and itis,) Cynanche tonsillaris.

ANTIADON'CUS, (antiades, and ογκος, 'tumour,') A swelling of the tonsils, *Anti'agri* has a similar meaning.

ANTIADONCUS INFLAMMATORIUS, Cynanche tonsillaris.

ANTIAGRI, (antiades, and agra,) *Antiadoncus*.

ANTIAPOPLEC'TIC, *Antiapoplec'ticus*, *Antapoplec'ticus*, *Apoplec'ticus*, (anti, and αποπληξια, 'apoplexy,') A remedy for apoplexy.

ANTIARIS TOXICARIA, see Upas.

ANTIARTHRIT'IC, *Antiarthrit'ic*, *Antiarthrit'ic'us*, *Antipodag'ric*, (anti, and αρθρις, 'the gout,') (F.) *Antigoutteux*. A remedy for gout.

ANTIASTHEN'IC, *Antiasten'ic'us*, (anti, and ασθεια, 'debility,') A remedy for debility.

ANTI-ASTHMATIC, *Antiasthmat'icus*, *Ant'-asthmatic'icus*, (*anti*, and *asthma*.) A remedy for asthma.

ANTIATROPHICUS, *Antatrophic*.

ANTIBALLOMENUM, (*αντιβαλλειν*, 'to put in place of another,') Succedaneum.

ANTIBELLA, (*anti*, and *βέλλα*, 'a leech,') *Antia sanguisuga*.

ANTIBECHICUS, (*anti*, and *βηξ*, gen. *βηξος*, 'cough,') Expectorant.

ANTIBRACHIAL, *Antibrachia'lis*. That which concerns the fore-arm.—Bichat. J. Cloquet suggests that the word should be written *antebrachial*, from *ante*, 'before,' and *brachium*, 'the arm'—as *antebrachial region*, *antebrachial aponeurosis*, &c.

ANTEBRACHIAL APONEUROSIS, (F.) *Aponévrose antibrachiale*, is a portion of the aponeurotic sheath which envelops the whole of the upper limb. It arises from the brachial aponeurosis, from a fibrous expansion of the tendon of the biceps muscle, from the epicondyle, epitrochlea, and, behind, from the tendon of the triceps brachialis. Within, it is inserted into the cubitus, etc.; and, below, is confounded with the two annular ligaments of the carpus. It is covered by the skin, by veins, lymphatics, and by filaments of superficial nerves; it covers the muscles of the fore-arm, adheres to them, and sends between them several fibrous septa, which serve them for points of insertion.

ANTIBRACHIUM, Fore-arm.

ANTIBROMIC, *Antibromo'micus*, (*anti*, and *βρωμος*, 'fœtor,') A *Deodorizer*, or *Deodorant*. An agent that destroys offensive odours—as chloride of zinc, simple sulphate of alumina, &c.

ANTICACHECTIC, *Anticachect'icus*, *Anticacchym'ic*, (*anti*, and *καχξία*, 'cachexy,') A remedy against cachexy.

ANTICACCHYMIC, (*anti*, *κακος*, 'bad,' and *χυμος*, 'juice,') Anticachectic.

ANTICANCEROUS, (*anti*, and *cancer*,) *Anticancero'us*, *Anticancero'sus*, *Anticarcinom'atous*, *Antiscir'rhous*. Opposed to cancer.

ANTICANCROUS, *Anticancerous*.

ANTICARCINOMATOUS, (*anti*, and *καρκινωμα*, 'cancer,') *Anticancerous*.

ANTICARDIUM, (*anti*, and *καρδία*, 'heart,') *Fossette du cœur*, *Scrobiculus cordis*.

ANTICATARRHAL, *Anticattarrha'lis*, *Anticattarrho'icus*, (*anti*, and *κατάρρως*, 'catarrh,') A remedy for catarrh.

ANTICAUSODIC, *Anticausod'ic*, *Anticausod'icus*, (*anti*, and *καυσος*, 'a burning fever,') A remedy for *causos* or inflammatory fever.

ANTICAUSOTIC, *Anticausodic*.

ANTICHEIR, (*αντιχειρ*, from *anti*, and *χειρ*, 'hand,') Pollex, see *Digitus*.

ANTICHERADICUS, (*anti*, and *χοιρας*, 'scrofula,') *Antiscrofulous*.

ANTICHOLERIC, *Anticholer'icus*, (*anti*, and *cholera*. A remedy against cholera.

ANTICHOLERICA, *Sophora heptaphylla*.

ANTICIPATING, *Anticipans*, *Antepo'nens*, *Prolep'ticus*, (F.) *Anticipant*, (*ante*, 'before,' and *capio*, 'I take,') An epithet for a periodical phenomenon, recurring at progressively shorter intervals. An *anticipating intermittent* is one in which the intervals between the paroxysms become progressively less.

ANTICENMIUM, (*αντικνημιον*, from *anti*, and *κνημη*, 'leg,') Shin.

ANTICŒUR, (*anti*, and *cœur*, 'heart,') *Scrobiculus cordis*.

ANTICOLIC, *Anticol'icus*, (*anti*, and *κολικος*, 'the colic,') That which is opposed to colic.

ANTICOMMA, gen. *Anticom'matis*, (*anti*, and *κομμα*, 'a blow,') *Contre-coup*.

ANTICOPE, (*αντικοπη*,) *Contre-coup*.

ANTICOPTOSCOPIUM, (*αντι, κοπτω*, 'to strike, and *σκοπεω*, 'to inspect,') *Plex'meter*.

ANTICRUSIS, (*αντικρουσις*,) *Contraissura*.

ANTICRUSMA, *Contraissura*.

ANTICUS, (*ante*, 'before,') *Anterior*.

ANTIDARTREUX, (*dartre* (F.), herpes,) *Antierpetic*.

ANTIDEIXIS, (*anti*, and *δειξις*, 'indication,') *Counterindication*.

ANTIDIARRHŒIC, *Antidiarrhœ'icus*, (*anti*, and *diarrhœa*.) A remedy for diarrhœa. Opposed to diarrhœa.

ANTIDIN'IC, *Antidin'icus*, *Din'ic*, *Din'icus*, (*anti*, and *δινος*, 'vertigo,') Opposed to vertigo.

ANTIDOTAL or *Antido'tal*, *Antidota'le*. Relating to an antidote; possessed of the powers of an antidote.

ANTIDOTARIUM. A dispensatory. A pharmacopœia or formulary. *Antidotary* was used formerly for any composition that had the properties of an antidote.

ANTIDOTARY, see *Antidotarium*.

ANTIDOTE, *Antid'otum*, *Antitox'icum*, (*anti*, and *διδωμι*, 'I give,') Originally, this word signified an *internal remedy*. It is now used synonymously with *counterpoison*, *antiphar'macum*, (F.) *Contre-poison*, and signifies any remedy capable of combating the effect of poison.

ANTIDOTE, *Bibron's*, (F.) *Antidote de Bibron*. An antidote to the poison of the rattlesnake, and other serpents, proposed by Professor Bibron. which, when administered early, and continued for a sufficient length of time, it has been conceived, proved effectual. It is composed as follows: *R. Potassii iodidi*, gr. iv; *Hydargyri chloridi corrosivi*, gr. ij; *Brominii*, ʒv. M. Dose, ten drops, in a tablespoonful of wine or brandy; to be repeated if necessary. The mixture must be kept in a well-stopped vial.

ANTIDOTUM, *Antidote*—a. *Heraclidis*, *Enneapharmacos*—a. *Mithridatium*, *Mithridate*.

ANTIDYNAMICA, (*anti*, and *δυναμις*, 'force,') *Debitants*.

ANTIDYNOUS, (badly formed from *anti*, and *οδυνη*, 'pain,') *Anodyne*.

ANTIDYSENTERIC, *Antidysenter'icus*, (*anti*, *δυσ*, 'with difficulty,' and *εντερων*, 'intestine,') Opposed to dysentery.

ANTIEMETIC, *Antemet'ic*, *Antiemet'icus*, (*anti*, and *εμετικος*, 'emetic,') A remedy for vomiting.

ANTIEPHAL'TIC or **ANTEPHAL'TIC**, *Antiephial'ticus*, (*anti*, and *εφιαλτης*, 'nightmare,') A remedy for nightmare.

ANTIEPILEP'TIC or **ANTEPILEP'TIC**, *Antiepilept'icus*, (*anti*, and *επιληψια*, 'epilepsy,') A remedy for epilepsy.

ANTIFEBRILIS, (*anti*, and *febris*, 'fever,') *Febrifuge*.

ANTIGALAC'TIC, *Antigalac'ticus*, *Antilac'teus*, *Antilactes'cent*, (*anti*, and *gala*,) (F.) *Antilacteux*. Opposed to the secretion of milk or to diseases caused by the milk.

ANTIG'ONI COLLYRIUM NI'GRUM, *Black colly'r'ium of Antig'onus*. It was composed of cadmia, antimony, pepper, verdigris, gum Arabic, and water.

ANTIGUA, see *West Indies*.

ANTIHŒMPTYICUS, *Anthæmptyicus*.

ANTIECH'TIC, *Antiech'ticus*, *Antiech'ticus*, (*anti*, and *ἐχς*, 'habit of body,') The *Antiech'ticum* *Pore'rii* is the white oxyd of antimony; also called *Diaphoret'icum Jovial'e*.

ANTHELIX, *Anthelix*.

ANTHELMINTICUS, *Anthelmintic*.

ANTHEMORRHAG'IC, *Antihæmorrhag'icus*, *Anthæmorrhag'icus*, (*anti*, and *αιμορραγια*, 'hemorrhage,') That which is against hemorrhage; an antihemorrhagic remedy.

ANTHEMORRHOIDAL, *Antihemorrhoidalis*, (anti, and *αιμορροϊδης*, 'hemorrhoids.') A remedy for hemorrhoids.

ANTIHERPETIC, *Antiherpeticus*, (anti, and *ερπης*, gen. *ερπητος*, 'herpes.') (F.) *Antidartreux*. A remedy for herpes.

ANTHYDROPHOBIC, *Antihydrophobicus*, *Antihysus*, *Alysus*, (anti, *υδωρ*, 'water,' and *φοβος*, 'dread.') A remedy for hydrophobia.

ANTHYDROPIC, *Antihydropticus*, *Hydropicus*, (anti, and *υδρωψ*, 'dropsy.') A remedy for dropsy.

ANTHYPNOTIC, *Anthypnotic*.

ANTHYSTERIC, *Anthysteric*.

ANTI-ICTERIC, *Anti-ictericus*, *Ictericus*, (anti, and *ικτερος*, 'jaundice.') A remedy for jaundice.

ANTI-IMPETIGENES, SOLOMON'S, see *Liquor Hydrargyri oxymerialis*.

ANTILABIUM, *Prolabium*, see *Antelabia*.

ANTILACTESCENT, *Antilactetic*.

ANTILACTEUS, (anti, and *lac*, gen. *lactis*, 'milk,') *Antilactetic*.

ANTILALITEUX, *Antilactetic*.

ANTILEPSIS, (*αντιληψις*, from *αντιλαμβανω*, 'I take hold of.') *Apprehensio*. The mode of attaching a bandage over a diseased part, by fixing it upon the sound parts. The mode of securing bandages, &c., from slipping. Treatment by revulsion or derivation.

ANTILETHARGIC, *Antilethargicus*, (anti, and *ληθαργικος*, 'affected with lethargy.') A remedy for lethargy.

ANTILITHICS, *Antilithica*, *Lithica*, (anti, and *λιθος*, 'a stone,') A substance that prevents the formation of calculi in the urinary organs.

The chief antilithics—according as the calculi are lithic acid or phosphatic—are alkalies or acids; with revellents, especially change of air; tonics, as *diosma crenata* [?], and *uva ursi* [?].

ANTILY, MINERAL WATERS OF. A celebrated French medicinal spring, near Meaux, in France. The waters have not been analysed; but astonishing and chimerical effects have been ascribed to them.

ANTIOBIUM, (anti, and *λοβιον*, 'the lobe of the ear,') *Antitragus*, *Tragus*.

ANTIOIMIC, *Antioimicus*, *Antilemic*, *Antipestilentialis*, (anti, and *λοιμος*, 'the plague,') A remedy for the plague.

ANTIOPIUS. The *Antelope*. (F.) *Gazelle*, (*αγρος*, 'a flower,' and *ωψ*, 'the eye'—so called from its beautiful eye.) An African animal, whose hoofs and horns were formerly given in hysteric and epileptic cases.

ANTILYSSUS, (anti, and *λυσσα*, 'rabies,') *Antihydrophobic*.

ANTIMELANCHOLIC, *Antimelancholicus*, (anti, and *μελαγχολια*, 'melancholy,') A remedy for melancholy.

ANTIMEPHITIC, *Antimephiticus*, (anti, and *mephitic*.) A remedy against mephitic or deleterious gases.

ANTIMIASMATIC, *Antimiasmaticus* (anti, and *miasma*.) A remedy against miasmatic affections, as quinia.

ANTIMOINE, *Antimonium*—a. *Beurre d'*, *Antimonium muriatum*—a. *Chlorure d'*, *Antimonium muriatum*—a. *Oxide d'*, *Algaroth*—a. *Oxide blanc d'*, *Antimonium diaphoreticum*—a. *Soufre doré d'*, *Antimonii sulphuretum præcipitatum*—a. *Sulfure d'*, *Antimonium*—a. *Sulfuré*, *hydrosulphure rouge d'*, *Antimonii sulphuretum rubrum*—a. *Verre d'*, *Antimonii vitrum*.

ANTIMOIAL, *Antimonia'lis*, *Stibio'lis*, (from *antimonium*.) A composition into which antimony enters. A preparation of antimony.

ANTIMONIAL POWDER, *Pulvis antimonia'lis* seu *antimonii compositus*, *Oxidum antimonii cum phosphate cal'cis*, *Phosphas cal'cis stibiat'us* seu

cal'cium stibiatum, *Pulvis Jame'sii* seu *stibiat'us* seu de *phosphate cal'cis et stibii compositus* seu *Jaco'bi* seu *febrifugus Jaco'bi*, *Facitiosus Jame's Powder*, *Schwan'berg's Fe'ver Powder*, *Chen'e'vix's Antimo'nial Pow'der*, (F.) *Poudre antimoniale composée* ou de *JAMES*. A peroxide of antimony combined with phosphate of lime. (Take of common sulphuret of antimony, ℥ij; kartsborn shavings, ℥ij. Roast in an iron pot, until they form a gray powder. Put this into a long pot, with a small hole in the cover. Keep it in a red heat for two hours, and grind to a fine powder. In the Ph. B. (1864), *Pulvis antimonia'lis* is made by mixing thoroughly one ounce of oxide of antimony with two ounces of precipitated phosphate of lime.)

This preparation has long been esteemed as a febrifuge: but it is extremely uncertain in its action. The ordinary dose is 6 or 8 grains.

The original James's Powder—*Pulvis Jaco'bi*, *P. Jaco'bi ve'rus*—of which the above is a succedaneum, was a patent medicine, so ambiguously described, that the powder of James could not be prepared from the formula. It had an immense reputation, which it did not merit.

ANTIMONIALE CAUSTICUM, *Antimonium muriatum*.

ANTIMONIATUM, SULPHUR, *Antimonii sulphuretum præcipitatum*—a. *Tartar*, *Antimonium tartarizatum*.

ANTIMONII BUTYRUM, *Antimonium muriatum*—a. *Calx*, *Antimonium diaphoreticum*—a. *Cerussa*, *Antimonium diaphoreticum*—a. *Murias*, *Antimonium muriatum*—a. *Oleum*, *Antimonium muriatum*—a. *Oxydum hydrosulphuratum aurantiacum*, *Antimonii sulphuretum præcipitatum*—a. *Oxydum*, see *Algaroth*—a. *Oxydum auratum*, *Antimonii sulphuratum præcipitatum*—a. *Oxydum nitro-muriaticum*, *Algaroth*—a. *Oxydum cum sulphure vitrificatum*, *Antimonii vitrum*—a. *Oxydum sulphuretum vitrificatum*, *Antimonii vitrum*—a. *Oxysulphuretum*, see *A. sulphuretum rubrum*—a. *Potassio-tartaras*, *Antimonium tartarizatum*—a. *Regulus medicinalis*, *Antimonium medicinale*—a. *Sal*, *Antimonium tartarizatum*—a. *Sulphur auratum*, *Antimonii sulphuretum præcipitatum*—a. *Sulphur præcipitatum*, *Antimonii sulphuretum præcipitatum*—a. *Sulphuratum*, *Antimonii sulphuretum præcipitatum*—a. *Sulphuretum*, *Antimonium*.

ANTIMONII SULPHURETUM PRÆCIPITATUM, *Sulphur antimonia'tum*, *Hydrosulphure'tum stibio'sum cum sulph'ure*, *Oxides stibii sulphuratum*, *Oxydum antimonii hydrosulphuratum aurantiacum*, *Oxydum auratum antimonii*, *Sulphure'tum stibii oxydula'ti*, *Hydrosulphure'tum luteum oxydi stibii sulfurati*, *Sulphur antimonii præcipitatum*, *Sulphur auratum antimonii*, *Antimonium sulphuratum aurantiacum*, *Gold'en Sulphur of Antimony*, (F.) *Soufre doré d'Antimoine*.

Antimonii Sulphuretum Præcipitatum, (Ph. U. S., 1851.) *Antimonium Sulphuratum*, (Ph. U. S., 1863,) is made by boiling together *Sulphuret of Antimony*, in fine powder, *Solution of Potassa*, and *distilled water*; straining the liquor while hot, and dropping into it *Diluted Sulphuric Acid* so long as it produces a precipitate. Dose, gr. i to gr. iij.

ANTIMONII SULPHURETUM RUBRUM, *Antimonium sulphuratum rubrum* seu *fus'cum*, *Hydrosulphure'tum stibii rubrum*, *Subhydrosulphure'tum stibii*, *Hydrosulphure'tum rubrum stibii sulphurati*, *Pulvis Carthusiano'rum*, *Red Sulphuret of Antimony*, *Kermes minera'lis* seu *minera'lis*, *Kermes* or *Chermes mineral*, (F.) *Hydrosulphure rouge d'Antimoine sulfuré*, *Vermillon de Provence*.

Antimonii Oxysulphure'tum (Ph. U. S., 1863) is considered to be nearly the same as the old *Kermes* or *Chermes Mineral*. It is prepared by boiling together *sulphuret of antimony* and *carbonate of soda* in water; filtering; and washing and dry-

ing the precipitate. It is a powder of a purplish brown colour, and almost devoid of taste. It is emetic, diaphoretic, and cathartic, according to the dose; and has been chiefly used in chronic rheumatism, and in cutaneous affections. Dose, gr. j. to gr. iij.

ANTIMONII TARTRAS, Antimonium tartarizatum —a. Tersulphuretum, Antimonium.

ANTIMONII VITRUM, *Glass of Antimony*, *Antimonii oxydum sulphuretum vitrificatum*, *Oxydum stibii semivitrificatum*, *Antimonium vitrificatum*, *Oxydum antimonii cum sulphure vitrificatum*, *Vitrum stibii*, *Antimonii vitrum hyacinthinum*, *Oxydum vitrum stibii vitreum*, (F.) *Verre d'Antimoine*. (Formed by roasting powdered common antimony in a shallow vessel, over a gentle fire, till it is of a whitish gray colour, and emits no fumes in a red heat; then melting it, on a quick fire, into a clean, brownish red glass.) It has been used for preparing the tartarized antimony, and antimonial wine.

ANTIMONII VITRUM HYACINTHINUM, *Antimonii vitrum*.

ANTIMONII ET POTASSÆ TARTRAS, Antimonium tartarizatum.

ANTIMONIOUS ACID, Antimonium diaphoreticum.

ANTIMONIUM, (of unknown derivation; by some supposed to be from *anti*, and *monos*, 'alone'; i.e. not found alone: or, according to others, from *anti*, 'against,' and (F.) *moine*, 'a monk; because, it is asserted, certain monks suffered much from it.) *Stibi*, *Stibium*, *Regulus Antimonii*, *Mineralium*, *Gynæceum*, *Mugnesia Saturi*, *Marcesita plumbea*, *Platyophthalmon*, *Stimmi*, *Aurum leprosum*, *Antimonium crudum* seu *sulphuratum nigrum*, *A. sulphuretum* seu *Tersulphuretum*, *Sulphuretum stibii nigrum*, *Common* or *Crude Antimony*, *Sulphuret of Antimony*, (F.) *Antimoine*, *Sulfure d'Antimoine*. Sulphuret of antimony is the ore from which all the preparations of antimony are formed. In Pharmacy, it is the native sesquisulphuret of antimony, purified by fusion. When prepared for medical use, by trituration and levigation, it forms a powder, *Antimonii sulphuretum preparatum*, (Ph. D.) of a black or bluish gray colour, which is insoluble. It is slightly diaphoretic and alterative, and has been used in chronic rheumatism, cutaneous diseases, &c.

ANTIMONIUM ALBUM, Bismuth —a. Calcinatum, Antimonium diaphoreticum —a. Chloratum solutum, Antimonium muriatum —a. Crudum, Antimonium.

ANTIMONIUM DIAPHORETICUM, *Diaphoretic Antimony*, *Antimonious Acid*, *Mineral Bezard*, *Antimonium Calcinatum*, *Mineral Diaphoretic*, *Matière perlée de KERKING*, *Peroxide of Antimony*, *Calx Antimonii*, *Antimonium diaphoreticum lotum*, *Cerusæ Antimonii*, *Calx Antimonii elotæ*, *Oxodes stibii album*, *Oxydum stibiosum*, *Oxydum stibii album medianlê nître confectum*, *Potassæ biantimonias*, *Deutoxide of Antimony*, (F.) *Oxide blanc d'Antimoine préparé par le moyen du nitre*. (Common antimony, libj; purified nitre, liij.—Throw it by spoonfuls into a red-hot crucible; powder and wash. The flowers that stick to the side of the crucible must be carefully separated, otherwise they render it emetic. Dose, gr. x to xxx.

ANTIMONIUM EMETICUM, *A. tartarizatum*.

ANTIMONIUM MEDICINALE, *Regulus Antimonii Medicinalis*, *Medicinal Regulus of Antimony*. (*Antimon. sulphur. 3v Potass. subcarb. 3i Sodii chlorid. 3iv*. Powder, mix, and melt. When cold, separate the scoræ at top, powder the mass, and wash it well.) It is conceived to be more active than common antimony.

ANTIMONIUM MURIATUM, *Antimonii Murias*,

Chloridum seu *Sesquichloridum* seu *Terechloridum* seu *Chloruretum stibii*, *Sputum tritum draconum*, *Deutomurias stibii sublimatus*, *Butter of Antimony*, *Muriate of Antimony*, *Chloride or Chloruret, Sesquichloride or Terechloride of Antimony*, *Butyrum Antimonii* seu *Stibii*, *Oleum Antimonii*, *Cavsticum antimoniale*, *Antimonium salitum* seu *chlora tum solutum*, *Liquor Antimonii murialis*, (F.) *Chlorure d'Antimoine*, *Bear's d'Antimoine*. (Common antimony and corrosive sublimate, of each equal parts: grind together, and distil in a wide-necked retort, and let the butyraceous matter, that comes over, run, in a moist place, to a liquid oil.) A caustic, but not much used as such. Sometimes taken as poison. Dissolved in hydrochloric acid, it forms the *Antimonii Terechloridi Liquor*, *Solution of Terechloride of Antimony*, of the British Pharmacopœia (1864), from which oxide of antimony is prepared.

ANTIMONIUM SALITUM, *Antimonium muriatum* —a. Sulphuratum aurantiacum, *Antimonii sulphuretum præcipitatum* —a. Sulphuratum nigrum, see Antimonium —a. Sulphuratum rubrum, *Antimonii sulphuretum rubrum* —a. Tartaratum, *A. tartarizatum*.

ANTIMONIUM TARTARIZATUM, *A. Tartaratum*, *Tartras antimonii*, *Tartras Antimonii et Potassæ*, *Antimonii et Potassæ Tartras* (Ph. U. S.), *Antimonii potassiotartras*, *Tartris Antimonii*, *Tar'tar Antimoniatum*, *Sol Antimonii*, *Tartras Potassæ stibiosus* seu *stibialis*, *Tartris lixivie stibialis*, *Deutotartras potassæ et stibii*, *Tartarus emeticus* seu *stibialis*, *Tartarum emeticum*, *Antimonium emeticum*, *Tartarized Antimony*, *Tartrate of Antimony and potassæ*, *Potassiotartrate of Antimony*, *Emetic Tar'tar*, *Tar'tar Emetic*, (F.) *Tartre stibié*, *Tartre Émétique*, *Émétique*; in some parts of the United States, vulgarly and improperly, called *Tartar*. (Made by boiling powdered oxide of antimony and powdered bitartrate of potassa in distilled water; filtering and crystallizing [Ph. U. S., 1863].) Tartarized antimony is emetic, sometimes cathartic and diaphoretic. Externally, it is rubefacient, and in the form of ointment more especially, *unguentum antimonii*, is used to cause pustulation. Dose, as an emetic, gr. j to gr. iv in solution; as a diaphoretic, gr. one-sixteenth to gr. one-quarter.

The empirical preparation, called **NORRIS'S DROPS**, consists of a solution of tartarized antimony in rectified spirit, disguised by the addition of some vegetable colouring matter.

ANTIMONIUM VITRIFICATUM, *Antimonii vitrum*.

ANTIMONY, Antimonium —a. Butter of, Antimonium muriatum —a. Chloride of, Antimonium muriatum —a. Chloruret of, Antimonium muriatum —a. Common, Antimonium —a. Crude, Antimonium —a. Deutoxide of, Antimonium diaphoreticum —a. Flowers of, Algaroth —a. Glass of, Antimonii vitrum —a. Golden sulphur of, Antimonii sulphuretum præcipitatum —a. Medicinal regulus of, Antimonium medicinale —a. Muriate of, Antimonium muriatum —a. Oxide of, see Algaroth —a. Peroxide of, Antimonium diaphoreticum —a. Potassio-tartrate of, Antimonium tartarizatum —a. Sesquichloride of, Antimonium muriatum —a. Sulphuret of, red, Antimonii sulphuretum rubrum —a. Tartarized, Antimonium tartarizatum —a. Terechloride of, see Algaroth —a. Teroxide of, Antimonium muriatum —a. Vegetable, Eupatorium perfoliatum.

ANTIMONY AND POTASSA, TARTRATE OF, Antimonium tartarizatum.

ANTINEPHRITIC, *Antinephretic*, *Antinephreticus*, (from *anti*, 'against,' and *nephritis*.) A remedy for inflammation of the kidney.

ANTINEUROPATHIC, *Nervine*.

ANTINEUROTIC, *Nervine*.

ANTINIAD, see Antinial.

ANTINIAL, (*anti*, and *ινω*, 'the ridge of the occiput.') An epithet for an aspect towards the side opposite to the *inion*, or ridge of the occiput. —Barclay. *Antin'iad* is used adverbially by the same writer, to signify 'towards the antinial aspect.'

ANTIOCHI HYERA. A preparation extolled by the ancients in melancholy, hydrophobia, epilepsy, &c. It was formed of germander, agaric, pulp of colocynth, Arabian stoechas, opoponax, sagapenum, parsley, aristolochia, white pepper, cinnamon, lavender, myrrh, honey, &c.

ANTIOCHI THERIACA. A theriac employed by Antiochus against every kind of poison. It was composed of thyme, opoponax, millet, trefoil, fennel, aniseed, nigella sativa, &c.

ANTIODONTALGIC, *Antodontalgic*, *Antiodontalgic*, *Antodontalgic*, *Odontalgic*, *Odontalgic*, (F.) *Odontalgique*, (*anti*, and *οδονταλγια*, 'toothache.') A remedy for toothache.

ANTIORGAS'TIC, *Antiorgas'ticus*, (*anti*, and *οργαω*, 'I desire vehemently.') A remedy for orgasm or erethism, and for irritation in general.

ANTIPARALYTIC, *Antiparalytic*, (*anti*, and *παρλυσις*, 'palsy.') Opposed to palsy.

ANTIPARASITIC, *Antiparasitic*, (*anti*, and *παρσιτος*, 'a parasite.') An agent that destroys parasites, as the different vermin that infest the body. The chief antiparasitics are *Cocculus*, *Staphisagria*, *Veratrum album*, and certain of the mercurial preparations.

ANTIPARASTATITIS, (*ante*, and *παρστανις*, 'the epididymis'; also, 'the prostate, and *itis*.) Inflammation of Cowper's glands.

ANTIPATHIA, (*avtiπαθεια*, from *anti*, and *παθος*, 'passion, affection.') Aversion. A natural repugnance to any person or thing.

ANTIPATHIC, *Antipathic*, (F.) *Antipathique*. Belonging to antipathy. Opposite, contrary,—as *humeurs antipathiques* (F.); humours opposed to each other. Also, palliatives.

ANTIPATRI THERIACA, *The'riac of Antipater*. A farrago of more than 40 articles: used as an antidote against the bites of serpents.

ANTIPEDICULOŒA, *Antiphthierica*.

ANTIPERIODIC, *Antiperiodic*, (*anti*, and *περιδος*, 'a period.') A remedy which possesses the power of arresting morbid periodical movements;—e. g. the sulphate of quinia in intermittents.

ANTIPERISTALTIC, *Antiperistaltic*, (*anti*, and *περισταλτις*, 'I contract.') An inverted action of the intestinal tube.

ANTIPERIS'TASIS, (*avtiπεριστασις*, from *anti*, and *περιστασις*, 'reunion, aggregation.') A union of opposite circumstances: the action of two contrary qualities, one of which augments the force of the other. The peripateticians asserted, that it is by Antiperistasis that fire is hotter in winter than in summer. Theophrastus attributes the cause, which renders man more vigorous, and makes him digest more readily in winter, to the augmentation of heat caused by Antiperistasis.

ANTIPERNIUS, (*anti*, and *περνιο*, 'a chilblain.') A remedy against chilblains;—as *Unguentum antipernium*, an ointment for chilblains.

ANTIPERTUSSIS, (*anti*, and *pertussis*, 'hooping cough,') see *Zinci sulphas*.

ANTIPESTILENTIALIS, *Antiloomic*.

ANTIPHARMACUM, (*avtiφαρμακον*, from *anti*, and *φάρμακον*, 'poison,') Antidote.

ANTIPHARMACUS, *Alexipharmic*.

ANTIPHLOGIS'TIC, *Antiphlogistic*, (*anti*, and *φλεγω*, 'I burn.') Opposed to inflammation; as *Antiphlogistic remedies*, *A. regimen*, &c.

ANTIPH'THERIACA, *Antiphthiriaca*, *Anti-*

pediculo'sa, (*anti*, and *φθιραιω*, 'I am lousy.') A remedy used to destroy lice.

ANTIPHTHIS'ICAL, *Antiphthis'icus*, (*anti*, and *φθισις*, 'consumption.') Opposed to phthisis.

ANTIPHYSETICS, *Carminatives*.

ANTIPHY'SICA, *Carminatives*.

ANTIPHYSICAL, *Antiphys'icus*, (*anti*, and *φυσαιω*, 'I blow.') An expeller of wind: a carminative.

It has also been used for any thing preternatural; here, the derivation is from *anti*, and *φυσις*, 'nature.' The French sometimes say, '*un goût antiphysique*,' 'an unnatural taste.'

ANTIPLAS'TIC, *Antiplas'ticus*, *Plastilytic*, *Plastilytic*, (*anti*, and *πλαστικός*, 'formative.') Antiformative. An agent that diminishes the quantity of plastic matter—fibrin—in the blood.

ANTIPLEURIT'IC, *Antipleuret'icus*, *Antipleuret'ic*, (*anti*, and *πλευριτις*, 'pleurisy.') Opposed to pleurisy.

ANTIPNEUMON'IC, *Antipneumo'nicus*, (*anti*, and *πνευμονια*, 'disease or inflammation of the lungs.') A remedy for disease or inflammation of the lungs.

ANTIPODAGRIC, (*anti*, and *ποδαγρα*, 'gout.') Antiarthritic.

ANTI'PRAX'IS, (*avτιπραξις*, from *anti* and *πρασσω*, 'I act.') A contrary state of different parts in the same patient: e. g. an increase of heat in one organ, and diminution of it in another.

ANTIPSOR'IC, *Antipso'ricus*, *Antiscab'ious*, (*anti*, and *ψωρα*, 'the itch.') (F.) *Antigaleux*. Opposed to the itch.

ANTIPTRID, *Antiseptic*.

ANTI'PY'IC, *Antipy'icus*, (*anti*, and *πυον*, 'pus.') Opposed to suppuration.

ANTI'PYRETIC, (*anti*, and *πυρετος*, 'fever,') Febrifuge.

ANTI'PYROT'IC, *Antipyrot'icus*, (*anti*, and *πυρ*, 'fire.') Opposed to burns or to pyrosis.

ANTIQUARTANA'RUM, *Antiquart'ium*. A remedy formerly used against quartan fever.

ANTIQUUS, ('old.') Chronic.

ANTIRHACHIT'IC, *Antirhachit'icus*, (*anti*, and *rhachitis*, 'rickets or rickets.') Opposed to rachitis or rickets.

ANTIRHEUMAT'IC, *Antirheumat'icus*, (*anti*, and *ρευμα*, gen. *ρευματος*, 'rheumatism.') A remedy for rheumatism.

ANTIRHINUM ACUTANGULUM, (*avti*, 'for or resembling,' and *ριν*, 'the nose.') A Linaria—a. Auriculatum, A. Elatine.

ANTIRHINUM ELAT'NE, A. auriculatum, *Elati'ne*, *E. hasta'ta*, *Lina'ria elati'ne*, *Cymbala'ria elati'ne*, *Fluel'len* or *Fe'male Speed'well*, was formerly used against scurvy and old ulcerations.

ANTIRHINUM HEDERACEUM, A. Linaria—a. Hederæfolium, A. Linaria.

ANTIRHINUM LINA'RIA, A. hederæfolium seu *hederæfolium* seu *acutangulum*, *Lina'ria*, *L. vulgari's* seu *cymbala'ria*, *Elati'ne* *cymbala'ria*, *Cymbala'ria muralis*, *Osy'ria*, *Urina'ria*, *Common Toad-flax*, (F.) *Linaire*. Ord. Scrophulariaceæ. The leaves have a bitterish taste. They are reputed to be diuretic and cathartic. An ointment made from them has been extolled in hemorrhoids.

ANTISCABIOUS, *Antipsoric*.

ANTISCRRHOUS, *Antiscancerous*.

ANTISCOLETICUS, (*anti*, and *σκωληξ*, 'a worm.') Anthelmintic.

ANTISCOLICUS, *Anthelmintic*.

ANTISCORBU'TIC, *Antiscorbu'ticus*, (*anti*, and *scorbutus*.) Opposed to scurvy.

ANTISCROF'ULOUS, *Antiscroph'ulous*, *Antiscroful'ous*, *Antistrumo'ous*, *Antichercad'icus*. Opposed to scrofula.

ANTISEP'TIC, *Antsept'icus*, *Antipn'trid*, (*anti*, and *σπυρος*, 'putrid.') *Antiputredino'ous*. Opposed to putrefaction. The chief antiseptics, internally or externally employed, are *Acidum Muraticum*, *Acidum Nitricum*, *Acidum Sulphuricum*, *Alumina*

sulphas, Carbo Ligni, Calx Chlorinata, Chlorinum, Cinchona and its active principles, *Oreasote, Duacil Radix, Fermentum Cerevisiæ, Plumbi nitras, Potassæ permanganas, Soda Chlorinata, and Zinci Chloridum.*

ANTISIAL/AGOGUE, *Antisialago'gus, Antisi'alus*, (*anti, σιῶλον, 'saliva,' and ἀγω, 'I expel.'*) A remedy against ptyalism.

ANTISPASIS, (*αντισπασίς*, from *anti*, and *σπᾶω*, 'I draw.') Derivation, Revulsion.

ANTISPASMOD'IC, *Antispasmod'icus, Antispas'ticus*, (*anti*, and *σπᾶω*, 'I draw.') Opposed to spasm. The whole operation of antispasmodics is probably revulsive. The following are the chief reputed antispasmodics. *Æther Sulphuricus, Asafœtida, Castoreum, Dracontium, Moschus, Oleum Animale Dippelii, and Valeriana*—with the mental antispasmodics, abstraction, powerful emotions, fear, &c. Of direct antispasmodics, there is no example.

ANTISPASTICUS, (*αντισπαστικός*;) *Antispasmodic*, Derivative.

ANTISQUAM'IC, *Antisqua'micus*, (*anti*, and *squama*, 'a scale.') A name given by Dr. Headland to a medicine, which, by acting on the blood, removes cutaneous affections. The name is inappropriate, as the true scaly diseases—lepra and psoriasis—are not the only ones that require the use of eutrophics.

ANTISTASIS, (*αντιστάσις*), from *anti*, and *σῆω*, 'I place.' Antagonism.

ANTISTER'NUM, (*αντιστερνον*, from *anti* and *σπῆρον*, 'the sternum.') The back.

ANTISTERYG'MA, gen. *Antisteryg'matis*, (*anti* and *στηρυγμα*, 'a support.') A fulcrum, support, crutch.

ANTISTRUMOUS, (*anti* and *struma*, 'scrofula.') Antiscrofulous.

ANTISUDORAL, *Antisudora'lis*, (*anti*, and *sudor*, gen. *sudor'is*, 'sweat.') A remedy, that diminishes sweat.

ANTISYPHILIT'IC, *Antisypphilit'icus*, (*anti*, and *sypphilis*, 'the venereal disease.') Opposed to the venereal disease.

ANTITASIS, (*anti*, and *ταίω*, 'I extend.') Counter-extension.

ANTITHENAR, (*anti*, and *thenar*.) Opponens pollicis, Adductor pollicis pedis.

ANTITHERMA, (*anti*, and *θερμ*, 'heat,') Refrigerants.

ANTITHORA, *Aconitum anthora*.

ANTITOXICUM, (*anti*, and *toxicum*, 'a poison,') Antidote.

ANTITRAG'ICUS, *Antitra'geus*, (F.) *Muscle de l'Antitragus, M. antitragien*.—(Ch.) Belonging to the antitragus. A small muscle is so called, the existence of which is not constant. It occupies the space between the antitragus and anhelix.

ANTITRAGIEN, *Antitragieus*.

ANTITRAGUS, (*anti*, and *τραγος*, 'the tragus,') *Antilo'bium, Oblo'bium*. A conical eminence on the pavilion of the ear, opposite the tragus.

ANTITYP'IA, (*αντιτυπία*, from *anti*, and *τυπῶ*, 'I strike.') Resistance. Hardness. Repercussion.

ANTITYPICUS, *Antiperiodic*.

ANTIVENE'REAL, *Antivene'reus*, (*anti*, and *Venus*, gen. *Veneris*, 'Venus,') The same as *Antisypphilitic*. Formerly it was used synonymously with *Antaphrodisiac*.

ANTIVERMICULAR, *Antiperistaltic*.

ANTIVERMINOSUS, (*anti*, and *vermis*, 'a worm,') Anthelmintic.

ANTIZYMOT'IC, *Antizymo'ticus*. Opposed to zymosis. See *Zymotic*.

ANT'LIA or ANT'L'IA, (*αντλία, αντλία*, from *αντλῶ*, 'to pump out,') A syringe; a pump. Hence, *Antlia lac'tea, Lactis'u'gium*, a breast-pump; and

Antlia sanguis'u'ga, Anti'hæm'la, Hiru'do artificia'lis, the exhausting syringe used in cupping.

ANTLIA GASTRICA, Stomach-pump.

ANTODONTALGIC, *Antiodontalgic*.

ANTODYNUS, (*anti*, and *δύνη*, 'pain,') Anodyne.

ANTOTHESES, (*anti*, and *ωθεω*, 'to push,') Endomose.

ANTOTHISMUS, Endomose.

ANTRAX, Anthrax.

ANTRE, Antrum—a. d'Hyghmore, Antrum of Highmore.

ANTROVERSIO, (*antro*, modern for 'forwards,' and *vertere, versum*, 'to turn,') Anteversio.

AN'TRUM, (*αντρον*, 'a cavern,') *Cavern'a, Bar'athrum*, (F.) *Antre*. A name given to certain cavities in bones, the entrance to which is smaller than the bottom.

ANTRUM AURIS, Tympanum—a. Buccinosum, Cochlea, Labyrinth—a. Dentale, see Tooth—a. Pylori, see Stomach.

ANTRUM OF HIGH'MORE, *Antrum Highmoria'num* seu *Ge'næ seu maxilla'rē seu maxil'læ superio'ris, Genyan'trum, Gnathan'trum, Maxill'ary Sinus, Sinus Ge'næ pituita'rius*, (F.) *Antre d'Hyghmore, Sinus Maxillaire*. A deep cavity in the substance of the superior maxillary bone communicating with the middle meatus of the nose. It is lined by a prolongation of the Schneiderian membrane.

ANTSJAR, Upas.

ANULARIS, Annular.

ANULUS, (dim.,) see *Anus, Fossette*.

ANURESIS, (*an*, and *ουρεω*, 'to pass the urine,') Ischuria.

ANURIA, Ischuria.

ANUROCRINIE, (*an*, *ουρον*, and *κρίνω*, 'to separate,') see *Ischuria*.

ANURORRHÆA, (*an*, *ουρον*, and *ῥέω*, 'to flow,')

ANUS, gen. *Ani*, ('a circle,') *Po'dex, Po'tex, Mol'ynē, Mol'yn'ē, Dactyl'ios, Cath'edra, Cyr'ceon, Cys'saros, Cys'thos, Cy'sus, Cys'sus, A'nulus, Aph'edra, Aph'edron, He'dra, Proc'tos, Ar'chos, Se'des, Cul'us*. The circular opening situate at the inferior extremity of the rectum, by which the excrement is expelled. The *fundament*. The *body*. The *seat*, (F.) *Siège, Fondement*.

ANUS also signifies the anterior orifice of the *A'queduct of Syl'vius*. By some, this *Anus*, called, also, *Fora'men commu'nē poste'rius*, has been supposed to form a communication between the back part of the third ventricle and the lateral ventricles. It is closed up, however, by the tela choroidea, and also by the fornix, which is intimately connected with this. The foramen is situate between the commissura mollis of the optic thalami and the pineal gland.

ANUS, ARTIFIC'IAL. An opening made artificially, to supply the place of the natural anus. The term is often used to include preternatural anus.

ANUS CEREBRI, see *Anus*, and *Aquæductus Sylvii*.

ANUS CONTRACT'ED, (F.) *Anus rétréci*. A state of the anus when, from some cause, it is contracted.

ANUS, IMPER'FORATE. A malformation, in which there is no natural anus. See *Atresia ani adnata*.

ANUS, PRETERNAT'URAL, (F.) *Anus contre nature, A. anormal*. An accidental opening which gives issue to the whole or to a part of the fæces. It may be owing to a wound, or, which is more common, to gangrene attacking the intestine in a hernial sac.

This term, as well as (F.) *Anus devit, de'vions anus*, is also employed to the case where the anus, instead of being in its natural situation, is in some neighbouring cavity, as the bladder, vagina, etc.

ANXIETAS, Anxiety—a. *Præcordiorum*, see *Anxiety*—a. *Tibiarum*, see *Agacement des Nèrfs*.

ANXIETY, *Anxi'etas*, *Anxi'etude*, *Ademo'nia*, *Dyspho'ria anxi'etas*, *Alys'mus*, *Al'yce*, *Al'yis*, *A'se*, (F.) *Anxi'été*, (*angere*, *αγγειν*, 'to strangle, to suffocate.') A state of restlessness and agitation, with general indisposition, and a distressing sense of oppression at the epigastrium, *Anxi'etas p'ecordiorum*. *Inquietude*, *anxiety*, and *anguish*, represent degrees of the same condition.

ANYPNIA, (*an*, and *πνως*, 'sleep,') *Insomnia*.

AOCHLE'SIA, (*a*, and *αχλος*, 'disturbance,') *Tranquillity*, *calmness*.

AOINOS, (*ainos*, from *a*, and *oinos*, 'wine,') *Abstemious*.

AORTA, *Arte'ria mag'na seu cras'sa seu max'ima*, *Hæ'mal Ax'is*, of Owen, (F.) *Aorte*. This name was given by Aristotle to the chief artery of the body, (*ασπασται*, 'I am suspended,') as it seems to be suspended from the heart. [?] It is probable that Hippocrates meant by *ασπασται* the bronchia and their ramifications. The aorta is the common trunk of the arteries of the body. It arises from the left ventricle of the heart, about opposite to the fifth dorsal vertebra, passes upwards (*ascending Aorta*), forms the *great arch or cross of the Aorta*, and descends along the left of the spine (*descending Aorta*), until it reaches the middle of the fourth or fifth lumbar vertebra, where it bifurcates, to give origin to the common iliaes. The aorta is sometimes divided into the *Thorac'ic* or *Pec'toral*, and the *Abdom inal*. For the arteries, which arise from it, etc., see *Artery*.

AORTASIE, *Aorteuryisma*.

AORTECTASIE, (*aorta*, and *εκτασις*, 'dilatation,') *Aorteuryisma*.

AORTEURYSMA, gen. *Aorteuryism'atis*, (*aorta*, and *ευρυς*, 'dilated,') *Aneurism of the Aorta*, (F.) *Aortéuryisme*, *Anéuryisme de l'Aorte*, *Aortectasie*, *Aortasie of Piorry*. By carefully auscultating over the dorsal vertebra, a bellows' sound, with a deep and not always perceptible impulse, may be detected.

AOR'TIC, *Aor'ticus*, (F.) *Aortique*. Relating to the Aorta. The *Aortic ventricle*, (F.) *Ventricule Aortique*, is the left ventricle. The *Aortic valves* are the sigmoid valves at the origin of the Aorta, etc.

AORTITIS, *Inflammat'io Aor'tæ*, (*Aorta* and *itis*,) *Inflammation of the Aorta*.

AORTRON, (*ασπρον*, same etymon as *Aorta*,) *A lobe of the lungs*.

AOTUS, (*a*, and *ους*, gen. *ωτος*, 'an ear,') *A monster devoid of ears*.

AP, as a prefix, in composition, see *Ad*, and *Apo*.

A PAG'MA, gen. *Apag'matis*, *Apoclas'ma*, (*απαγμα*, from *apo*, and *αγω*, 'I remove,') *Separation*, *abduction*. *Separation of a fractured bone*.

APAGOGE, (*απαγωγη*, from *apo*, and *αγω*, 'I lead,') *Defecation*, *Inductio*.

APALACHINE, *Ilex vomitoria*—*a. à Feuilles de Prunier*, *Prinos*—*a. Gallis*, *Ilex vomitoria*.

APAL'LAGE, (*απαλλαγη*, from *απαλρω*, 'I change,') *Apallax'is*, *Mutation*, *change*. It is generally taken in a good sense, and means the change from disease to health.

APALLAXIS, (*απαλλαξις*,) *Apallage*.

APALOTICA, (*απαλότης*, 'softness, tenderness,') *Fortuitous lesions or deformities affecting the soft parts*. The first order in the class *Tychica*, of Good.

APANTHESIS, (*απανθησις*,) *Apanthismus*.

APANTHISMUS, *Apanthisis*, (*απανθησις*, from *apo*, and *ανθη*, 'I flower,') *The obliteration of parts previously inservient to useful purposes, as of the ductus venosus and ductus arteriosus, which are essential to foetal existence, but are subsequently unnecessary*. See also *Stuprum*.

APANTHROP'IA, (*απανθρωπια*, from *apo*, and *ανθρωπος*, 'man,') *Detestation of man; desire for solitude*. One of the symptoms of hypochondriasis.

APAPHRISMUS, (*apo*, and *αφρος*, 'froth,') *De-spumation*.

APARACH'YTUM VT'NUM, (*απαραχυτος* (*αιος*), from *a*, and *παραχω*, 'I pour over,') *The purest wine: that which has not been mixed with seawater*.

APARINE, (*απαρινη*,) *Galium aparine*—*a. Hispida*, *Galium aparine*.

APARTHROSIS, (*ap*, and *αρθρω*, 'to articulate,') *Diarthrosis*.

AP'ATHY, *Apathi'a*, *Ameli'a*, (*απαθεια*, from *a*, and *παθος*, 'affection,') (F.) *Apathie*. *Accidental suspension of the moral feelings*. It takes place in very severe diseases, particularly in malignant fevers.

APECHE'MA, gen. *Apeche'matis*, (*απηχημα*, from *apo*, and *ηχος*, 'sound,') *Properly the action of reflecting sound*. In medicine, it is synonymous with the Latin *Contrafissura*, a counter-fissure, a counter-blow.

APECTOCEPHALUS, *Acephalothorus*.

APEL'LA, *Appel'la*, *Leipoder'mos*, *Recuti'tus*, (*a*, and *pellis*, 'skin,') or more probably, perhaps, from a circumcised Jew, among the Romans, called *Apella*.) *One whose prepuce does not cover the glans*. Retraction or smallness of any other soft appendage. One who is circumcised.

APEPSIA, (*a*, and *πεπτιν*, 'to digest,') *Dyspepsia*.

APERIENT, *Aper'iens*, *Aperiti'vus*, *Aper'tive*, (*aperire* (*ab* and *pario*, 'to bring forth,') 'to open,') *Res'erans*. A laxative. (F.) *Apéritif*. A medicine which gently opens the bowels. The term had formerly a much more extensive signification, and, like *Catalyt'icum*, was given to a substance supposed to have the power of opening any of the passages, and even the blood-vessels.

APERIS'TATON, *Aperis'tatum*, (*απεριστατον*, from *a*, and *περιστημι*, 'I surround,') *An epithet for an ulcer not dangerous nor considerable nor surrounded by inflammation*.

APÉRITIF, (*aperio*, 'I open,') *Aperient*.

APERITIVUS, *Aperient*.

APERTIVE, *Aperient*.

APERTOR OCULI, *Levator palpebræ superioris*.

APERTO'RIMUM. An instrument for dilating the os uteri during labour.

APERTURA, *Mouth*—*a. Anterior ventriculi tertii cerebri*, *Vulva* (*cerebri*)—*a. Pelvis superior*, see *Pelvis*.

APEUTHYSMENOS, (*απευθυσμένος*, from *απευθεναι*, 'to straighten,') *Rectum*.

A'PEX, gen. *A'p'icis*, *M'icro*. The point or extremity of a part,—as the apex of the tongue, nose, etc.

APEX BEAT, see *Heart*.

APEX LINGUE, *Proglossis*.

APH, as a prefix, see *Apo*.

APH'A'CIA, *Aphu'cia*, (*a*, and *φακος*, 'lens,') *Absence of the crystalline lens*.

APHÆRESIS, (*αφαιρεσις*,) *Apheresis*, *Extirpation*.

APHA'GIA, (*a*, and *φαγω*, 'I eat,') *Inability to swallow*.

APHALANG'ASIS, (*ψαλαγγίς*, gen. *φαλαγγος*, 'a phalanx,') *The fourth stage of Oriental leprosy, which is recognised chiefly by a gangrenous condition of the fingers*.

APHASSOM'ENOS, (*αφασσομενος*, from *αφασσω*, 'I touch, I feel,') *The touching of the parts of generation of the female as a means of diagnosis*. See *Espace*.

APHEDRA, (*αφέδρα*, from *aph*, and *εδρα*, 'a seat,') *Anus*, *Lasanum*.

APHEDRIA, (*αφεδρία*,) *Menses*.

APHEDRON, *Anus*, *Lasanum*.

APHEL'IA, (*αφελια*, from *αφελής*, 'simple,') *Simplicity*. The simple manners of the sect of Methodists in teaching and practising medicine.

APHELX'IA, (*αφελκω*, 'I abstract,') *Voluntary inactivity of the whole or the greater part of the external senses to the impressions of sur-*

rounding objects, during wakefulness. *Rev'ery*, (F.) *Rêverie*. Dr. Good has introduced this into his Nosology, as well as *Aphela'ia sol'ens* or *absence of mind*.—*A. inten'ta* or *abstraction of mind*: and *A. otio'sa*, *Stu'dium inu'te*, *Cenosp'u'dia*, *brown study* or *listless musing*.

APHEPSEMA, (ἀφελσμα, from *aph*, and εἶναι, 'to boil,') Decoction.

APHEPSIS, (ἀφεψις,) Decoction.

APHE'RESIS, *Aphæ'resis*, (ἀφαιρεσις, from ἀφαιρω, 'I take away,') An operation by which any part of the body is separated from the other. Hippocrates, according to Pœsius, uses the expression *Aphæ'resis Sanguinis* for excessive hemorrhage; and Sennertus, to express the condition of an animal deprived both of the faculties of the mind and of the mind itself.

APH'ESIS, (ἀφesis, from ἀφηναι, 'I relax,') A remission. This word expresses sometimes the diminution or cessation of a disease; at others, languor and debility of the lower extremities. See Languor, and Remission.

APHILANTHROPY, *Aphilanthro'pia*, (ἀφιλω, 'I love,' and ἀνθρωπος, 'a man,') Dislike to man. Love of solitude. Vogel has given this name to the first degree of melancholy.

APHISTESIS, (ἀφιστημι, 'to remove,' from *aph*, and ιστημι, 'to place,') Abscess.

APHODEUMA, (ἀφoδευμα,) Excrement.

APHODUS, (ἀφoδος, from *aph*, and ὁδος, 'a way,') Excrement.

APHONETUS, (ἀφωντος,) Aphonus.

APHO'NTA, (ἀφωνία,) *Liga'tio lin'guæ*, *Loque'la abol'ita*, *Defec'tus loque'le*, *Dyspho'nia*, (of some,) *Ap'hony*, (F.) *Aphonie*, *Perte de la Voix*, (a, and φωνη, 'voice,') Privation of voice, or of the sounds that ought to be produced in the glottis. When aphonia forms part of catarrh or of 'cold,' it is commonly of but little consequence; but when produced by causes acting on the nervous system, as by some powerful emotion, or without any appreciable lesion of the vocal apparatus, (*Laryngoparalysis*;) it frequently resists all remedies.

APHONIA, Catalepsy—a. Surdorum, Mutitas Surdorum.

APHONICUS, Aphonus.

APHO'NUS, (ἀφωνος,) *Apho'nicus*, *Apho'netus*, *Apho'nic*. Relating to aphonia.

APHONY, Aphonia.

APHORIA, (ἀφορία, from α, and φερειν, 'to bear,') Sterilitas.

APHORICUS, Sterile.

APHOR'ME, (ἀφορμη, 'occasion,') The external and manifest cause of any thing. The occasional cause of a disease.

APHORUS, see Aphoria, Sterile.

APHRO'DES, (ἀφροδης, 'frothy,' from αφρος, 'foam,' and odes,) Applied to the blood and the excrements.

APHRODISIA, (ἀφροδισια,) Coition, Puberty.

APHRODISI'AC, *Aphrodisi'acus*, (ἀφροδιστη, 'Venus,') (F.) *Aphrodisiaque*. Medicine or food believed to be capable of exciting to the pleasures of love; as ginger, cantharides, &c. They are generally stimulants.

APHRODISIACUS, (ἀφροδισιακος,) Venereal.

APHRODISIASMUS, (ἀφροδισιασμος,) Coition.

APHRODISIOG'RAPHY, (ἀφροδιστη, 'Venus,' and γραφω, 'I describe,') Etymologically, this term means a description of the pleasures of love, but it has been placed at the head of a work describing the venereal disease.

APHRODITIC, Venereal.

APHROG'ALA, (ἀφρογαλα, from αφρος, 'foam,' and γαλα, 'milk,') *Loc spurto'num*. A name formerly given to milk rendered frothy by agitation.

APHRONIA, (α, and φρην, 'mind,') Apoplexy.

APHRONITRUM, (αφρος, 'foam,' and νιτρον, 'nitre,') Natrum, Soda.

APHROSYNE, (ἀφροσυνη,) Delirium, Insanity.

APH'THÆ, *Aph'tæ*, *Ap'thæ*, (απτω, 'I inflame,') Thrush or sore mouth, *Aph'ta lactu'cimen* seu *infan'tum*, *Lactu'cimen*, *Lactu'm'ina*, *Lactu'mina*, *Al'colæ*, *Em'phlysis aph'tha*, *Ul'cera serpenti' o'ris*, *Pu'stula o'ris*, *Fe'bris aphtho'sa*, *Angi'na aphtho'sa*, *Vesic'ulæ gingiva'rum*, *Stomatiti's exsudati'va* seu *vesiculo'sa infan'tum*, *Stomati'tis*, *S. aph'tha*, *Prunella*, *White Thrush*, *Milk Thrush*. Aphthæ consist of roundish, pearl-coloured vesicles, confined to the lips, mouth, and intestinal canal, and generally terminating in curd-like sloughs. In France, the Aphthæ of children, *Aphthæ des Enfants*, is called *Muguet*, *Millet*, *Blanchet*, *Catarrhe buccal* and *Stomatite crémense pultacée*, *Pultaceous inflammation of the Mouth*; and generally receives two divisions—the mild or discreet, (F.) *Muguet bévin* ou *discret*, and the malignant, (F.) *Muguet malin* ou *confluent*, the *Black Thrush*. Common Thrush is a disease of no consequence, requiring merely the use of absorbent laxatives. The malignant variety, which is rare, is of a more serious character, and is accompanied with typhoid symptoms.—*Ty'phus aphthoideus*.

A microscopic vegetable growth has been found in aphthæ; and the disease was, thence, called by Gruby, of Vienna, *Aphthophyton*.

APHTHÆ ADULTORUM, Stomatitis, aphthous—a. Præputii, Herpes præputii—a. Serpentes, Cancer aquaticus.

APH'THAPHYTE, *Aphthaphyton*, *Aphthophyton*, *Oid'ium albicans*, *Champignon du Muguet*, (*aphthæ*, and φηρον, 'a vegetable,') A parasitic vegetable growth observed in aphthæ. See Aphthæ.

APHTHES GANGRÊNEUX, Cancer aquaticus—a. *des Enfants*, Aphthæ.

APHTHEUX, Aphthous.

APHTHO'DES, *Aphthoides*, *Aphthoideus*, (from *aphthæ*, and odes,) Aphthous-like, belonging to aphthæ.

APHTHOPHYTON, *Aphthaphyte*, see Aphthæ.

APH'THOUS, *Aphtho'sus*, (F.) *Aphtheux*. Belonging to aphthæ; complicated with aphthæ; as *Aphthous Fever*.

APIASTRUM, (from *apis*, 'a bee,') *Melissa*. APICES (pl. of *aper*) CRURUM MEDULLÆ OBLONGATÆ, Corpora striata—a. Digitorum, Pupulæ. See Apex.

APILEPSIA, (απο, and ληψις, from λαμβανω, 'a seizure,') Apoplexy.

A'PIOL, *Apiolum*, *Pars'ley oil*, (*apium* 'parsley,' and *oleum*, 'oil,') A yellowish oily liquid, obtained from *apium petroselinum* or *parsley*, which, in the dose of 15 grains, has been used as an antiperiodic.

APIONTA, (απιοντα, from απειμι, 'to go away,') see Excretion.

APIOS, (απιος,) *Apios tuberosa*, *Pyrus communis*.

A'PIOS TUBERO'SA, *Glyci'ne Apios*. Ground nut, Wild bean, *Sagaban*. The root is used as food by the Indians of Northwestern America.

APIS, Bee.

API'TES, (απιτης, from απιον, 'a pear,') Perry.

APIUM, (απιον,) A. graveolens—a. Ammi, Ammi—a. Anisum, Pimpinella anisum—a. Carvi, Carum.

A'PIUM GRAVE'OLENS, *A'pium Paluda'pium*, *Eleoseli'num*, *Beli'num*, *Ses'eli grave'olens*, *S'i'num grave'olens* seu *a'pium*, *Small'age*, (F.) *Ache*. Ord. Umbelliferae. *Scar. Syst.* Pentandria Digynia. The plants, roots, and seeds are aperient and carminative. *Cel'ery* is a variety of this.

APIUM HORTENSE, A. graveolens—a. Montanum, Athamanta aureoselinum—a. Paludapium, A. Graveolens—a. Petræum, Bubon Macedonicum.

APIUM PETROSEL'NUM, *A'pium Hor'eu'se* seu

vulgarè, Grielum, Seli'um, Petroselinum, Com'mon Pars'ley, (F.) Persil. The root—*Petroselinum*, (Ph. U. S.)—and seeds are diuretic and aperient. See Apol.

APIUM SIUM, *Sium nodiflorum*—a. Vulgare, A. graveolens.

APLAS'TIC, *Aplast'icus*, (a, and *πλασσω*, 'I form.') That which is not capable of forming; or is not organizable.

APLASTIC EL'EMENT; one which is unsusceptible of any farther amount of organization.—Gerber.

APLESTIA, (*απληστια*, from *α*, and *πλεω*, 'to fill,') Ingluvies, Intemperance.

APLEU'ROS, (*απλευρος*, from *α*, and *πλευρον*, 'a rib.') One without ribs, or pleurae. The condition is *Apleuria*.

APLOT'OMY, *Aplotom'ia*, (*απλος*, 'simple,' and *τεμνω*, 'I cut.') A simple incision.

APNÉE, Apnoea.

APNEUMATOSIS, (a, and *πνευματω*, 'I distend with wind,') see Atelectasis.

APNEU'MIA, (a, and *πνευμων*, 'lung,') A monstrosity in which there is absence of lungs.

APNEUMONER'VIA, *Apneumoneu'ria*, (a, *πνευμων*, 'lung,' and *νευρον*, 'nerve,') Want of nervous action in the lungs.

APNEUSTIA, (*απνευστια*), Apnoea, Asphyxia.

APNEUSTUS, Apnus.

APNE'A, (*απνοια*, from *α*, and *πνέω*, 'I respire,') (F.) *Apnée*, Absence of respiration, *Respiratio abol'ita*; or insensible respiration. Also, Orthopnoea. See Asphyxia.

APNŒA INFANTUM, Asthma thymicum.

APNŒASPHYXIA, Asphyxia.

APNOICUS, Apnus.

AP'NUS, (*απνος*), *Apno'icus*, *Apneust'us*. One devoid of respiration. An epithet applied by authors to cases in which the respiration is so small and slow, that it seems suspended. It is probable, however, that the word was always applied to the patient, not to the disease.

APO, (*απο*), as a prefix denotes 'from, off, out, out of.'

APOBAMMA, gen. *Apobam'matis*, (*απο*, and *βαπτειν*, 'to wash,') Embamma.

APOBIOSIS, (*αποβιωσις*, from *απο*, and *βιωω*, 'to live,') Death.

APOBLEMA, gen. *Apoble'matis*, (*αποβλημα*, from *απο*, and *βαλλω*, 'to cast off,') Abortion.

APOBOLE, (*αποβολη*), (same etymon,) Abortion.

APOBRASMA, gen. *Apobrasm'atis*, (*αποβρασμα*, from *απο*, and *βρασσειν*, 'to winnow,') Furfur.

APOCAPNISMUS, (*αποκαπνισμος*, from *απο*, and *πνέω*, 'smoke,') Fumigation.

APOCATASTASIS, (*αποκαταστασις*, from *αποκα-στημι*, 'to re-establish,') Consistentia, Restauratio.

APOCATHARSIS, (*αποκαθαρσις*, from *απο*, and *καθαρειν*, 'to purge,') Catharsis.

APOCATHARTICUS, Cathartic.

APOCŒSIS, *Apocœno'sis*, (*αποκœνωσις*, from *απο*, and *κœνωσις*, 'evacuation,') A partial evacuation according to some, in opposition to Cœnosis, which signifies a general evacuation.—Cullen and Swediaur apply it to morbid fluxes.

APOCŒNOSIS, *Abœvacuatio*—a. Diabetes mellitus, Diabetes—a. Pytalismus mellitus, see Salivation—a. Vomitus pyrosis, Pyrosis.

APOCHOREON, (*αποχορευ*, 'to go away from,') Excrement.

APOCHREMMMA, gen. *Apochrem'matis*, (*αποχρημα*, from *απο*, and *χρεμα*, 'spit,') Sputum.

APOCHREMPISIS, (*αποχρημψις*), Exspuition.

APOCHYLIS'MUS, *Apochylis'ma*, (*απο*, and *χυλος*, 'juice,') The expression of the juices of vegetables.

APOCHYMA, gen. *Apochym'atia*, (*αποχω*, 'I pour out,') A sort of tar, obtained from old ships, which is impregnated with chloride of sodium. It was used as a disient of tumours.

APOCYN GÖBE-MOUCHE, *Apocynum androsæmifolium*.

APOCLASMA, gen. *Apoclas'matis*, (*αποκλασμα*, from *απο*, and *κλαω*, 'I break,') Abduction, Apagma.

APOCLEISIS, (*αποκλεισις*, from *αποκλειω*, 'I shut out,') Asitia, Disgust.

APOC/OPE, (*αποκοπη*, from *απο*, and *κοπτειν*, 'to cut,') Abseccion. A wound with loss of substance. Fracture with loss of part of a bone.

Amputation.

APOCOPUS, (*αποκοπος*), Castratus.

APOCRISIS, (*αποκρισις*, from *αποκρίνω*, 'I separate,') Contagion, Excrement, Secretion.

APOCROUS'TIC, pl. *Apocroust'ica* seu *Apocroust'ica*, (*remed'ia*,) (*αποκρουστικά*, from *απο*, and *κρουω*, 'I push,') An astringent and repellent.

APOCRUSTICA, Apocroustics.

APOCYESIS, (*αποκυσις*, from *απο*, and *κυω*, 'to be pregnant,') Parturition.

APOCYN, see Apocynum Cannabinum.

APOCYNIN, see Apocynum Cannabinum.

APOC'YNUM ANDROSÆMIFOLIUM, (Ph. U. S.) (*αποκυνον*, from *απο*, and *κυνω*, 'a dog,' because esteemed of old, to be fatal to dogs.) *Dog's Bone*, *Bitter Dog's Bone*, *Milkweed*, *Bitter-root*, *Honeybloom*, *Catch'fly*, *Fly'trap*, *Ip'ecac*, *Ameri-*

ican Ip'ecac, (F.) *Apocin gobe-mouche*, A. *amer.* Ord. Apocynaceae. Sex. Syst. Pentandria Digynia.

The root of this plant is found from Canada to Carolina. Thirty grains evacuate the stomach as effectually as two-thirds of the amount of *Ipecacuanha*, by which name it is known in various parts of the Eastern States. It is in the secondary list of the Pharmacopœia of the United States.

APOCYNUM CANNA'BINUM, (Ph. U. S.) *Indian Hemp*. The root of this American plant, which possesses emetic, cathartic, diaphoretic and diuretic properties, has been strongly recommended in dropsy, and has been given in decoction,—3j boiled in three pints of water to two. A wineglassful for a dose. An active principle, *Apoc'yn* or *Apoc'ynin*, has been extracted from it.

APOCYNUM HYPERICÆFOLIUM, a variety of A. cannabinum, which, as well as A. androsæmifolium, abounds in a milky juice. This, when applied to the skin, produces a troublesome eruption resembling flea-bites: hence, the plants have been termed, by the *voyageurs* in the Hudson's Bay territory, *Herbes à la puce*.

APOCYNUM NOVÆ ANGLIÆ HIRSETUM, *Asclepias tuberosa*—a. Orange, *Asclepias tuberosa*—a. Scandens, Allamanda.

APODACRYTICUS, *Delachrymat'ivus*, (*αποδακρυτικός*, from *απο*, and *δακρυω*, 'I weep,') A substance, supposed to occasion a flow of the tears, and then to arrest them.

APODEMIALGIA, Nostalgia. According to others, the opposite to Nostalgia, or a desire to leave one's country; (from *αποδημια*, 'foreign travel,' and *αλγος*, 'suffering,')

APOD'IA, (*αποδια*, from *α*, and *πους*, gen. *ποδος*, 'a foot,') Want of feet; hence *Apon* or *Apus*, one who has no feet.

APODYTERIUM, (*αποδυτηριον*), *Coniste'rium*, *Spoliato'rium*, *Spoli'd'rium*, (*αποδυω*, 'I strip off,') The ante-room, where the bathers stripped themselves in the ancient gymnasia.

APOGALACTISMUS, (*απογαλακτισμος*, from *απο*, and *γαλαξ*, gen. *γαλακτος*, 'milk,') Weaning.

APOGALACTOS, (*απογαλακτος*), Exuber.

APOGEUSIS, (*απο*, and *γευσις*, 'taste,') Ageustia.

APOGEUSTIA, Ageustia.

APOGLAUCOSIS, (*απογλαυκωσις*), Glaucois.

APOGON, (*απωγων*, from *α*, and *πωγων*, 'beard,') Imberbis.

APOG'ONUM, (*απογονον*, from *απο*, and *γινωμαι*, 'I exist,') A living foetus in utero.

APOLAR, see Neurine.

APOLEPISIS, (*apo*, and *λεπις*, 'a scale.') Desquamation.

APOLEPISMUS, Desquamation.

APOLEP'SIS, *Apolep'sia*, *Apolip'sis*, (*αποληψις*, from *απολαμβάνω*, 'I retain.') Retention, suppression. Also, Asphyxia.

APOLEXIS, (*αποληξις*, from *αποληγω*, 'I cease.') Old age, decrepitude.

APOLINO'SIS, (*απολινωσις*, from *apo*, and *λινον*, 'a flaxen thread.') The mode of operating for fistula in ano, by means of a thread of *Homolinon* or *Linum crudum*.

APOLIPSIS, Apoplepsis.

APOLLINARIS ALTERCUM, *Hyoseyamus*.

APOLYS'IA, *Apol'yis*, (*απολυσις*, from *απολυω*, 'I loosen.') Solution. Relaxation. Debility of the limbs or looseness of bandages. Expulsion of the fetus and its dependencies. Termination of a disease.

APOMATHE'MA, gen. *Apomathe'matis*, *Apomathe'sis*, (*απομαθημα*, *απομαθησις*, from *apo*, and *μαθάνω*, 'I learn.') Forgetfulness of things taught.

APOM'ELI, (*απομελι*, from *apo*, and *μελι*, 'honey.') An oxymel or decoction made of honey.

APOMYLE'NAS, (*απομυληνας*, from *απομύλλαινω*, 'I make a wry mouth.') One who pushes his lips forwards, pressing them against each other. Occasionally a symptom of nervous fever.

APOMYTHO'SIS, *Apomytho'sis*, (*απομυσσω*, 'I snore.') A disease in which there is stertor.

APONEUROGRAPHY, *Aponeurogra'phia*, (F.) *Aponeurographie*, (*aponeurosis*, and *γραφη*, 'a description.') A description of the Aponeuroses.

APONEUROLOGY, *Aponeurolog'ia*, (F.) *Aponeurologie*, (*aponeurosis*, and *λογος*, 'a discourse.') *Aponeurosiology*. The anatomy of the aponeuroses.

APONEUROSIOLOGY, Aponeurology.

APONEURO'SIS, (*απονευρωσις*.) *Aponeuro'sis*, (*apo*, and *νευρον*, 'a nerve.') *Pronerva'tio*, *Denerva'tio*, *Enerva'tio*, *Expan'sio nervi'sa*, (F.) *Aponeurose*, *Aponevrose*. The ancient called every white part *νευρον*, and regarded an Aponeurosis as a nervous expansion. The Aponeuroses are white, shining membranes, very resisting, and composed of fibres interlaced. Some are continuous with the muscular fibres, and differ only from tendons by their flat form. They are called *Aponeuro'ses of inser'tion*, (F.) *Aponevroses d'insertion*, when they are at the extremities of muscles, and attach them to the bone;—*Aponeuro'ses of intersec'tion*, (F.) *Aponevroses d'intersection*, if they interrupt the continuity of the muscle, and are continuous on both sides with muscular fibres. Others surround the muscle, and prevent its displacement: they are called *envel'oping Aponeuro'ses*, (F.) *Aponevroses d'enveloppe*.

APONEUROSIS, Fascia—a. Crural, Fascia lata—a. Femoral, Fascia lata—a. Iliac, Fascia iliaca.

APONEUROSITIS, (from *aponeurosis*, and *itis*.) Inflammation of an aponeurosis.

APONEUROT'IC, *Aponeurotic*, (F.) *Aponevrotique*. What relates to Aponeuroses:—thus, we say *Aponeurotic expansion*, *Aponeurotic muscle*, &c.

APONEUROT'OMY, *Aponeurotom'ia*, (F.) *Aponevrotomie*, (*aponeurosis*, and *τεμνω*, 'I cut.') Anatomy of aponeuroses.

Aponeurotomy has, also, been proposed for the division (*débriderment*) of filaments, &c., in aponeurotic openings, and for the section of fasciæ.

APONÉVROGRAPHIE, Aponeurography.

APONÉVROLOGIE, Aponeurology.

APONÉVROSE, Aponeurosis—a. *Pédieuse*, see *Pedal Aponeurosis*—a. *Superficielle de l'Abdomen et de la Cuisse*, Fascia superficialis.

APONEVROSIS, Aponeurosis.

APONÉVROTIQUE, Aponeurotic.

APONÉVROTOMIE, Aponeurotomy.

APON'IA, (*απονία*, from *a*, and *πονος*, 'pain.') Freedom from pain.

APONIPSIS, (*apo*, and *νιπτιο*, 'I wash.') Ablution.

APOPALLE'SIS, *Apopa'll'sis*, (*αποπαλλσις*, from *αποπαλλω*, 'I throw off.') Expulsion. Protrusion. Also, Abortion.

APOPATE'MA, gen. *Apopate'matis*, *Apopate'thos*, *Apop'atus*, (*αποπατημα*, from *απο*, and *πατος*, 'dirt.') The excrement, and the place where it is deposited.

APOPHLEGMATISANS PER NARES, Erhine—a. per Os, Sialagogue.

APOPHLEGMATISAN'TIA, *Apophlegmatisan'tia*, *Apophlegmatis'mi*, (*apo*, and *φλεγμα*, 'phlegm.') Medicines which facilitate the upward expulsion of mucus from the mucous membrane of the digestive or air passages; as gargles, masticatories, &c.

APOPHLEG'MATISM, (*αποφλεγματισμος*) *Apophlegmatis'mus*. The action of *Apophlegmatisantia*.

APOPHLEGMATISMI, *Apophlegmatisantia*.

APOPH'RADES, (*αποφρας*, 'unlucky.') An epithet applied to unlucky days, (*dies nefundi*.) Days on which a favourable change is not expected to occur in a disease.

APOPHRAXIS, (*αποφραξις*, from *αποφρασσω*, 'I obstruct.') Amenorrhæa.

APOPHTHAR'MA, gen. *Apophtharm'atis*, (*αποφθαγμα*, from *apo*, and *φθειρω*, 'I corrupt.') Abortion, as well as a medicine to procure abortion.

APOPHTHORA, (*αποφθορα*, from *αποφθειρω*, 'to miscarry.') Abortion.

APOPHTHORIUS, Abortive.

APOPHY'ADES, (*αποφυαδες*, from *apo*, and *φυω*, 'I spring.') The ramifications of veins and arteries.

APOPH'YSAR, *Apophysa'ris*, (F.) *Apophysaire*. Relating or appertaining to an apophysis.

APOPHYSE, Apophysis—a. *Basilaire*, Basilar process—a. *Engainante ou vaginale*, Vaginal process—a. *Pétrée*, see Temporal Bone—a. *Pyramidale*, see Temporal Bone.

APOPHYSES ÉPINEUSES, Spinous processes of the vertebæ.

APOPH'YSIS, (*αποφυσις*, from *apo*, and *φυω*, 'I rise') *Ec'physis*, *Proces'sus*, *Appen'dix*, *Prominen'tia os'sis contin'ua*, *A proc'ess of a bone*. (F.) *Apophyse*. When the apophysis is yet separated from the body of the bone by intervening cartilage, it is called *Epiph'ysis*. The apophyses or processes are, at times, distinguished by epithets, expressive of their form: as *A. styloid*, *A. coracoid*, &c. Others are not preceded by the word apophysis; as *Trochanter*, *Tuberosity*, &c.

APOPHYSIS OF INGRAS'SIAS is a term applied to the lesser ala of the sphenoid bone.

APOPHYSIS MAMILLARIS, Mastoid process—a. of *Rau*, *Grêle apophyse du Marteau*: see *Malleus*—a. *Zygomatica*, Zygomatic process.

APIOPIES'MA, gen. *Apiopies'matis*, (*αποπιεσμη*, from *αποπιεζω*, 'I compress.') Hippocrates uses the term to signify a fancied expression or forcing out of humours by the application of bandages in wounds and fractures.

APOPLANESIS, (*αποπλανησις*, from *apo*, and *πλανω*, 'to lead astray.') Error loci.

APOPLEC'TIC, *Apoplec'ticus*. Referring to Apoplexy. This word has various significations. It is applied, 1. To individuals labouring under apoplexy: 2. To remedies proper for combating apoplexy: 3. To the constitution, temperament, or make, *Architectu'ra apoplec'tica*, *Hab'itus apoplec'ticus seu quadra'tus seu toro'sus*, which predisposes to it, and, 4. To the symptoms which characterize apoplexy; as *Apoplectic sleep*, *A. stroke*, *A. stertor*, &c. The jugular veins have

also, by some, been called *Apoplectic veins*, *Venæ apoplecticae*.

APOPLECTIC CELL. *Foetus apoplecticus*. A cavity remaining in the encephalon, after effusion of blood and its subsequent absorption.

APOPLECTICUS, Antiapoplectic, Apoplectic.

APOPLEXIA, Apoplexy—a. *Catalepsia*, *Catalepsia*—a. *Cerebralis*, see *Apoplexy*—a. *Cerebri*, see *Apoplexy*—a. *Cordis*, *Hæmocardiorrhagia*—a. *Hepatica*, *Hepatorrhagia*—a. *Hydrocephalica*, *Hydrocephalus internus*—a. *Interarachnoidealis*, *Apoplexy*, *meningeal*—a. *Intermeningealis*, *Apoplexy*, *meningeal*—a. *Medullaris*, *Apoplexia myelitica*—a. *Meningæa*, *Apoplexy*, *meningeal*.

APOPLEXIA MYELITICA, *A. Medullaris* seu *Spinalis* seu *Rachialis*, *Hæmorhachis*, *Myelorrhagia*, *Myelapoplexia*, (F.) *Apoplexie de la Moëlle épinière*, *Hémorrhagie de la Moëlle épinière*, *Hémato-myélie*, *Hémo-myélorrhagie*, *Hématorrhachie*. Hemorrhage into the spinal marrow.

APOPLEXIA NERVOSA, *Apoplexy*, *nervous*—a. *Nervosa traumatica*, *Concussion of the brain*—a. *Oculi*, *Hæmophthalmia*—a. *Pituitosa*, see *Apoplexy*—a. *Pulmonalis*, see *Hæmoptysis*—a. *Pulmonum*, see *Hæmoptysis*—a. *Rachialis*, *A. myelitica*—a. *Renalis*, *Apoplexy*, *renal*—a. *Sanguinea*, see *Apoplexy*—a. *Serosa*, see *Apoplexy*—a. *Simplex*, *Apoplexy*, *nervous*—a. *Spasmodica*, *Apoplexy*, *nervous*—a. *Spinalis*, *Apoplexia myelitica*—a. *Temulenta*, see *Temulencia*.

APOPLEXIE, *Apoplexy*—a. *Capillaire*, *Molities cerebri*—a. *Cérébrale*, *Apoplexy*, *Hémorrhagie cérébrale*—a. *du Cœur*, *Hæmocardiorrhagia*.

APOPLEXIE FOUDROYANTE, ('Thundering Apoplexy.') A form of apoplexy, which is intense and rapidly fatal.

APOPLEXIE MENINGÉE, *Apoplexy*, *meningeal*—a. *de la Moëlle Épineière*, *Apoplexy*, *spinal*.

AP'OPLEXY, *Apoplexia*, (Sc.) *Poplexy*, (αποπληξία, from ἀποπληττειν, 'to strike with violence.') At the present day, the term apoplexy is employed by many writers to signify *interstitial hæmorrhage*, (F.) *Hémorrhagie interstitielle*, or every effusion of blood, which occurs suddenly into the substance of an organ or tissue. Hence, we speak of cerebral apoplexy, pulmonary apoplexy, &c. &c. Formerly it was always—and still is by many—used in a restricted sense, to signify, in other words, the train of phenomena, which characterize cerebral apoplexy. This disease, *Hæmorrhagia Cerebri*, *Aphro'nia*, *Cerebrus Apoplexia*, *Co'ma Apoplexia*, *Apoplexia cerebri sanguinea* seu *cerebralis*, *Encephalorrhagia*, *Sanguinis ic'tus*, *Hæmatencephalum*, *Pulpezia*, *Sideratio*, *Apilepsia*, *Morbis atton'itus*, *Gut'ta*, *Theoplegia*, *Theoplexia*, (F.) *Apoplexie*, *A. cérébrale*, *Hématoencephalie*, *Coup de sang*, is characterized by diminution, or loss of sensation and mental manifestation; by the cessation, more or less complete, of motion; and by a comatose state,—circulation and respiration continuing. It generally consists in pressure upon the brain; either from turgescence of vessels, or from extravasation of blood: hence the terms *Hæmencephalus*, *Hémorrhagie cérébrale*, and *Hématoencephalorrhagie*, applied to it by some. The general prognosis is unfavourable; especially when it occurs after the age of 35. When Apoplexy is accompanied with a hard, full pulse, and flushed countenance, it is called *Apoplexia sanguinea*, *Cataph'ora Co'ma*; when with a feeble pulse and pale countenance, and evidences of serous effusion, *Apoplexia serosa*, *A. pituitosa*, *Serous Apoplexy*, *Cataph'ora hydrocephalica*, *Encephaloch'ysis senilis*, *Hydroceph'alus acut'us senum*, (F.) *Hydroencephalorrhœe* of Piorry, *Hydropisie cérébrale suraiguë*, *Hydrorrhagie*.

In *Nervous Ap'oplexy*, *Apoplexia nervosa* seu

spasmodica, *A. simplex*, *Simple apoplexy*, no lesion whatever may be perceptible on dissection; although the patient may have died under all the phenomena that are characteristic of apoplexy.

APOPLEXY, HEAT, see *Coup-de-Soleil*—a. of the Heart, *Hæmocardiorrhagia*.

APOPLEXY, MENINGEAL, *Apoplexia meningea* seu *intermeningealis* seu *interarachnoidealis*, (F.) *Apoplexie méningée*, *Hémorrhagie méningée*. Hemorrhage from the meninges of the brain or spinal marrow, generally into the great cavity of the arachnoid.

APOPLEXY, NERVOUS, see *Apoplexy*—a. *Pulmonary*, see *Hæmoptysis*—a. *Simple*, *A. Nervous*.

APOPLEXY, RENAL, *Apoplexia renalis*. A condition of the kidney, characterized by knotty, irregular, tuberculated eminences, some of a deep black colour. Effusion of blood into the substance of the kidney.

APOPLEXY, SEROUS, see *Apoplexy*—a. *Spinal*, *Apoplexia myelitica*.

APOPNEUSIS, (ἀποπνευσις, from ἀπο, and πνεω, 'I breathe.') *Exhalatio*.

APOPNIxis (apo, and πνιγω, 'I strangle.') *Suffocation*.

APOPNOE, (ἀποπνοη, from ἀπο, and πνεω, 'I breathe.') *Expiratio*.

APOPNEA, (ἀποπνοια,) *Expiratio*.

APOPSYCHIA, (apo, and ψυχη, 'the breath of life.') *Syncope*.

APOPTOSIS, (ἀποπτωσις, from ἀποπτωω, 'I fall down.') *A relaxation of bandages*.

APORRHINOSIS, (apo, and ριν, 'the nose.') *A discharge from the nose*.

APORRHŒ, (ἀπορρῆ,) *Aporrhœa*.

APORRHŒA, *Apor'r'hœe*, *Apor'r'hysis*, *Deflu-vium*, (ἀπορρῆα, from ἀπορρῆω, 'I flow from.') An emanation, effluvia, contagion. A falling off of the hair, according to some.

APORRHYSIS, (ἀπορρῆσις,) *Aporrhœa*.

APOSEMMA, gen. *Aposem'matis*, *Aposep-sis*, (ἀποσκημμα, from ἀποσκηπτω, 'I lie down, I direct myself towards.') Afflux of fluids towards a part. *Metastasis*. The first word has been applied to the excrements.

APOSENOSIS, *Apocenososis*.

APOSEPARNIS'MUS, *Deascia'tio*, (ἀποσκηπαρισμος, from ἀπο, and ἀσκηπαρνον, 'a hatchet.') Wound of the cranium, by a cutting instrument, in which a piece of the bone has been cut out, as with a hatchet.

APOSEPSIS, (ἀποσκηψις,) *Aposemma*.

APOS'CHASIS, (ἀποσχασις,) *Aposch'asis*, (ἀποσχαζω, 'I scarify.') *Scarification*. A slight superficial incision in the skin. Also, blood-letting.

APOS'IA, *Sit'is defectus*, (a, and ποσις, 'drink.') Want of thirst, absence of desire for liquids.

APOSTITIA, (ἀποστῖα, from ἀπο, and εἶσος, 'food.') *Aversion for food*. See *Disgust*.

APOSIT'IC, *Aposit'icus*, (ἀποστιτικός.) Any substance which destroys the appetite, or suspends hunger.

AOSPAS'MA, gen. *Apospasm'atis*, (ἀοσπασμα, from ἀοσπασω, 'I tear or lacerate.') (F.) *Arrachement*. A solution of continuity, especially of a ligament; *Rheg'ma ligamenta'rè*, *Lacera'tio ligamenta'ria*.

AOSPASTIC, *Derivative*.

AOSPHACEL'ISIS, (ἀοσφακελις, from ἀπο, and σφακελος, 'mortification.') *Gangrene* in wounds and fractures, owing to the bandages being too tight.

AOSPHINX'IS, (ἀοσφινγίς, 'constriction, compression.') The action of a tight bandage.

AOSPONGIS'MUS, (ἀοσπογγισμός,) the act of sponging for any purpose.

AOSTALAG'MA, gen. *Apostalag'matis*, *Apostag'ma*, (apo, and σταλάω, 'I drop.') The ancient

name for the saccharine liquor which flows from grapes when not yet pressed.

APOS'TASIS, (ἀποστασις, from *apo*, and *ιστημι*, 'I stop.') The ancients had different significations for this word. It was most commonly used for an abscess. The separation of a fragment of bone by fracture. Removal of disease by some excretion, &c.

APOSTAX'IS, (ἀποσταξις, from ἀποσταζω, 'I distil from.') *Stasis*. The defluxion of any humour, as of blood from the nose.

APOSTEM, Abscess, see *Apostema*.

APOSTE'MA, gen. *Apostem'atis*, *Apos'tem*, (ἀποστημα, from *apo*, and *ιστημι*, 'I settle,' or ἀφιστημι, 'I recede from.') This word is used by the ancients somewhat vaguely. It meant an affection in which parts, previously in contact, are separated from each other by a fluid collected between them. The moderns regard it as synonymous with *Ab'scess*. Some, even of the moderns, have applied it to any watery tumour, and even to tumours in general.

APOSTEMA CEREBRI, Encephalopyosis—a. Empyema, Empyema—a. Parulis, Parulis—a. Phalangum, *Fourche*—a. Psoticum, Lumbar abscess.

APOSTEMACION, Abscess.

APOSTEMA'TION, (ἀποστηματιον,) *Apostema'tium*, (dim. of *Apostema*.) A small abscess.

APOSTERIG'MA, gen. *Aposterig'matis*, (ἀποστηριγμα, from ἀποστηρίζω, 'I support.') Anything that supports a diseased part, as a cushion, a pillow, &c. A deep-seated and inveterate disease of the intestines.

APOS'THIA, *Leipoder'mia*, (α, and ποσθια, 'prepuce.') Want of prepuce.

APOSTHUME, Abscess. See *Apostema*.

APOSTOLÉ, Extractum.

APOSTOLO'RU M UNGUEN'TUM, *Dodecaphar'macum*, Oint'ment of the *Apos'tles*. So called, because as many solid ingredients entered into its composition as there were apostles. It contained several resins and gum-resins, yellow wax, oil, vinegar, verdigris, &c., and was formerly employed as a vulnerary.

APOSTROPHE, (ἀποστροφή, from *apo*, and στροφή, 'I turn.') An aversion or disgust for food. Also, the direction of humours towards other parts.

APOSYRMA, gen. *Aposyr'matis*, (ἀποσυρμα, from *apo*, and σνρω, 'I tear,') Abrasion, Desquamation.

APOTELES'MA, gen. *Apoteles'matis*, (ἀποτελεσμα, from *apo*, and τέλεσμα, 'completion.') The result or termination of a disease. See, also, *Amuletum*.

APOTEXIS, (ἀποτηξις,) *Tabes*.

APOTHANASIA, (απο, and θάνατος, 'death,') See *Death*.

APOTHE'CA, (αποθήκη, *Pharmace'um*, *Pharmacopo'lum*, (απο, and τιθημι, 'to place,') Any place where things are kept, and therefore 'a shop,' and particularly a wine cellar. A place or vessel wherein medicines are kept. See *Pharmacopolium*.

APOTH'ECARIES' HALL. The Hall of the Corporation or Society of Apothecaries of London, where medicines are prepared and sold under their direction, &c. This Company obtained a charter of incorporation in the 15th year of James the First. No general practitioner can establish himself in England or Wales, without having obtained a license from the Court of Examiners of the Company.

APOTH'ECARY, *Apotheca'rius*, *Dispensa'tor*, *Pharmacopo'la*, *Pharmacopœ'us*, *Pharma'ceus*, *Pharmaceut'a*, *Confectiona'rius*, *Pharmacter*, *Pharmacur'gicus*, *Pharmacur'gus*, *Pharmaceut'ist*, (Prov.) *Pot'ecary*, (So.) *Pottingar*, (F.) *Apothicaire*, *Pharmacien*, *Pharmacopole*. In every country except Great Britain, Apothecary means one

who sells drugs, makes up prescriptions, &c. In addition to these offices, which, indeed, they rarely exercise, except in the case of their own patients, the Apothecaries in England form a privileged class of practitioners—a kind of sub-physician. See *Surgeon-apothecary*.

APOTHÈME, (from αποτιθημι, 'to deposit,' *apo*, and τιθημι, 'to put,') *Oxidized extractive*, (F.) *Extractif oxygéné*. A deposit which forms, when an extract is redissolved in water, or when vegetable solutions are boiled with access of air.

APOTHERAPEI'A, (αποθεραπεία,) *Apothera'pi'a*, *Apotherapeus'is*, (αποθεραπεύω, from *apo*, and θεραπεύω, 'I cure,') A perfect cure. In the ancient Gymnastics, it meant the last part of the exercises:—the friction, inunction, and bathing, for the purpose of obviating fatigue, or curing disease.

APOTHERAPEUSIS, (αποθεραπεισις,) *Apothera'peia*.

APOTHER'MUM, (αποθερμον, from *apo*, and θερμη, 'heat,') A pickle made of mustard, oil, and vinegar.

APOTH'ESIS, (αποθεσις,) (αποτιθημι, 'I put up,') The position proper to be given to a fractured limb, after reduction.

APOTHICAIRE, Apothecary.

APOTHICAIRERIE (F.). The same as *Apotheca*; also, a gallipot. See *Pharmacopolium*.

APOTHLIM'MA, (αποθλιμμα, from *apo*, and θλιβω, 'I press from,') Anciently, the dregs, and sometimes the expressed juice, *Succus expressus*, of plants.

APOTHRAS'IS, (αποθραυσις, from αποθρανω, 'I break,') Fracture of a bone, with spicula remaining. Extraction of a spiculum of bone. Also, *Ab-scission*.

APOTILMUS, (αποτιλμος, from *apo*, and τιλλω, 'I pluck,') Evulsion.

APOT'OKUS, (αποτοκος, from *apo*, and τικτω, 'I bring forth,') An abortive foetus.

APOTOME, (αποτομη, from *apo*, and τομη, 'incision,') Amputation.

APOTOMIA, (αποτομία,) Amputation.

APOTROPÆUM, (αποτροπαιον, from *apo*, and τρέπειν, 'to turn,') Amuletum.

APOTROPE, (αποτροπή,) *Aversion*. Also, deviation—as of a limb, *Parat'rope*.

APOXYSMUS, (αποξυσμος, from *apo*, and ξωω, 'I rasp,') Abrasion.

APOZEM, (αποζεμα, from *apo*, and ζωω, 'I boil,') Decoction.

APOZESIS, (αποζέσις,) Decoction.

APPARA'TUS, *Parasceu'e*, (απ, and παrare, 'to prepare,') This word signifies a collection of instruments, &c., for any operation whatever. (F.) *Appareil*.

In Surgery, it means the methodical arrangement of all the instruments and objects necessary for an operation or dressing. By extension, the French give the name *Appareil*, *Capso chirurgica*, to the case or drawers in which the apparatus is arranged.

Apparatus has likewise been applied to the different modes of operating for the stone. See *Lithotomy*.

In *Physiology*, *Apparatus*, (F.) *Appareil*, is applied to a collection of organs, all of which work towards the same end. A system of organs comprehends all those formed of a similar texture. An apparatus often comprehends organs of very different nature. In the former, there is analogy of structure; in the latter, analogy of function.

APPARATUS ALTUS, see *Lithotomy*.

APPARATUS IMMOV'ABLE, (F.) *Appareil immobile* ou *immovible*, *Immovable Band'age*, *Per'mnent Bandage*. An apparatus for fractures, which is generally formed by wetting the bandage in

some substance, as starch, (*Collodes'mus*), which becomes solid, and retains the parts in situ.

APPARATUS LATERALIS, see Lithotomy—a. Major, see Lithotomy—a. Minor, see Lithotomy—a. Urinary, Urinary organs—a. Uropoeticus, Urinary organs.

APPAREIL, Apparatus, *Boëtier*—a. *Diapno-gène*, see Perspiration—a. *Grand*, see Lithotomy—a. *Haut*, see Lithotomy—a. *Immobile*, Apparatus, immovable—a. *Inamovible*, Apparatus, immovable—a. *Lacrymal*, see Lacrymal passages—a. *Lateralisé*, see Lithotomy—a. *Locomoteur*, see Locomotive—a. *Névrothèle*, see *Névrothèle*—a. *Petit*, see Lithotomy—a. *Pigmental*, Pigmental apparatus.

APPAREILS DE FORMATION (F.). Gall admits, in the brain, two kinds of fibres; the one, divergent, proceeding from the cerebral peduncles to the convolutions, and constituting what he calls *appareils de formation*: the other, convergent, and proceeding from the convolutions to the centre of the organ, constituting what he calls *appareils de réunion*. The first, as a whole, form the organs of the mental faculties: the latter, are commissures, which unite parts of the organ that are double, and in pairs.

APPAUVRI, (ap, and pauvre, (L.) pauper, 'poor,') Impoverished.

APPENDAGE TO THE OVARY, Parovarium.

APPENDICE, Appendix—a. *Cæcal*, Appendix vermiformis cæci—a. *Digital*, Appendix vermiformis cæci—a. *Sous-sternale*, Xiphoid cartilage—a. *Sus-sphénoïdale du cerveau*, Pituitary gland—a. *Xiphoidé*, Xiphoid cartilage.

APPENDICES, see Appendix—a. *Coli Adiposæ*, Appendiculæ epiploicæ—a. *Épiploïques*, Appendiculæ epiploicæ.

APPENDICULA CEREBRI, (dim. of *Appendix*), Pituitary gland—a. *Epiploica*, Epiploic appendage—a. *Vermiformis cæci*, see Appendix.

APPENDICULÆ PINGUEDINOSÆ, *A. Epiploicæ*, *Epiploicæ appendages*, *Appendices colli adiposæ*, *Fimbrice carnosæ colli*, *Supplementa epiploica*, *Omentula*. (F.) *Appendices Epiploïques*. Prolongations of the peritoneum beyond the surface of the great intestine, which are analogous in texture and arrangement to omenta.

APPEN'DIX, gen. *Appen'dicis*, *Epiph'ysis*, (*appendere*, ap, and *pendere*, 'to hang,' 'to hang to or from,') (F.) *Appendice*, *Annexe*. Any part that adheres to an organ or is continuous with it: seeming as if added to it. An *appendage*; an apophysis.

APPENDIX AURICULÆ, see Auricles of the Heart—a. *Cerebri*, Pituitary gland—a. *ad Cerebrum*, Cerebellum—a. *Cutanea Septi Narium*, Statica Septi Narium—a. *to the Epididymis*, Vasculum aberrans—a. *Ventriculi*, Duodenum.

APPENDIX VERMIFORMIS seu *Appendic'ula Vermiformis Cæci*, *Tubus Vermicula'ris Cæci*, *Ec'phas*, *Additamentum Coli*, *Appen'dix Cæci*, (F.) *Appendice vermiforme*, *A. cæcal* or *digital*. A vermicular process, the size of a goose-quill, which hangs from the intestine cæcum. Its functions are unknown.

APPENDIX VESICÆ, see Bladder, sacculated.

APPETENCE, *Ap'petency*, *Appeten'tia*, (*appetere*, from ap, and *petere*, 'to seek,' 'to desire,') A desire, and especially an ardent, passionate desire for any object.

APPETIT, Appetite—a. *Perte d'*, Anorexia.

APPETITE, *Appeti'tus*, *Appeten'tia*, *Appeti'tia*, (ap, and *petere*, 'to seek,') *Cupi'do*, *Orex'is*, *Ormé*: (same etymology as Appetence.) An internal sensation, which warns us of the necessity of exerting certain functions, especially those of digestion and generation. In the latter case it is called *venereal appetite*, (F.) *Appetit vénérien*: in the former, simply *appetite*, (F.) *Appetit* ou *Appe-*

tition. If the desire for food, occasioned by a real want, be carried to a certain extent, it is called *hunger*, when solid food is concerned; *thirst*, when liquid. *Appetite*, and *hunger* ought not, however, to be employed synonymously: they are different degrees of the same want. Hunger is an imperative desire: it cannot be provoked, like the appetite. It is always allayed by eating: but not so the appetite; for, at times, it may be excited in this manner. They are very generally, however, used synonymously.

APPETITE, MORBID Limosis.

AP'PETITE, VENE'RAL, Venereal desire, (F.) *Le génésique*, *Amour physique*, *Sens génital*. The instinctive feeling that attracts the sexes towards each other to effect the work of reproduction.

APPETITIVE, *Appeti'tive*, (F.) *Appétitif*. Having or causing desire.

APPETITUS, Appetite—a. *Caninus*, *Boulimia*—a. *Deficiens*, *Dysorexia*.

APPLE, see Melum, and *Pyrus malus*—a. *Adam's*, *Pomum Adami*—a. *Bitter*, *Cucumis colocynthis*—a. *Curassoa*, *Aurantium curassavientum*—a. *Dead Sea*, see *Quercus infectoria*—a. *Eye*, see *Melon*—a. of the Eye, *Pupil*—a. *Mad*, see *Quercus infectoria*—a. *May*, *Podophyllum peltatum*—a. of Peru, *Datura stramonium*—a. root, *Euphorbia corollata*—a. of Sodom, see *Quercus infectoria*.

APPLE TEA, *Apple water*. Slice two large, not over-ripe apples, and pour over a pint of boiling water. After an hour, pour off the fluid, and, if necessary, sweeten with sugar.

APPLE TREE, *Pyrus malus*.

APPLICA'TA, (*applicare*, from ap, and *plicare*, 'to fold,' 'to apply,') A word, unnecessarily introduced into medical language, to express the objects which are applied immediately to the surface of the body, as clothes, cosmetics, baths, &c.

APPLICA'TION, *Applica'tio*, in a moral signification, is synonymous with attention. Also, the act of applying one thing to another; as the application of an apparatus, of a bandage, blister, &c., as well as the thing applied—*applicatum*.

APPREHEN'SIO, (ap, and *prehendere*, 'to take,') This word is employed in various senses. It means catalepsy or catoche. A kind of bandage for securing any part. Also, a therapeutical indication.

APPROCHE, (ap, and *proche*, 'near,') Coition.

APPROXIMA'TION, *Approxima'tio*, (ap, and *proximus*, 'nearest,') Ettmüller gave this name to a pretended method of curing disease, by making it pass from man into some animal or vegetable, by the aid of immediate contact.

APRAC'TA, (*απρακτα*, from a, and *πρασσω*, 'I act,') Without action. An epithet for the parts of generation, when unfit for copulation or generation.

APRICATIO, (*apricor*, 'to sun one's self,') Insolation.

APRICOT, *Prunus Armeniaca*.

APROC'TIA, *Atre'sia ani*, (a, and *πρωκτος*, 'anus,') Imperforation of the anus.

APROCTUS, see Atretus.

APRONIA, *Tamus communis*.

APROSO'PIA, *Tricocephalia*, (a, and *προσωπον*, 'the face,') A malformation, which consists in the face being deficient.

APROSOPUS, *Microprosopus*.

APSINTHIA'TUM, (*αψινθιον*, 'wormwood,') A sort of drink made of wormwood.

APSINTHITES, (*αψινθιτης*), Absinthites.

APSYCHIA, (a, and *ψυχη*, 'soul,' 'breath,') Syncope.

APSYCHISME, Idiostism.

APSYCHY, Syncope.

APSYXIA, Syncope.

APTHÆ, Aphthæ.

APTALIA, *Asia'lia*, (a, and *πυαλος*, 'saliva,') Want of saliva.

APTYS'TUS, (απυστος, from α, and πωω, 'I spit.') Devoid of expectoration. An epithet given, of old, to certain pleurisies, in which there is no expectoration.

APULO'SIS, (απουλωσις, from απο, and ουλη, 'cicatatrix'. Cicatrization.

APULOTICA, (απουλωτικά, Cicatrisantia.

APUS, (απος, from α, and πους, 'a foot'.) See Apodia.

APY'ETUS, (α, and πον, 'pus'.) An external affection, which does not end in suppuration.

APY'IQUE, Apyus.

APY'US, (απος, from α, and πον, 'pus'.) (F.) *Apygie*. That which does not afford pus.

APYRECTIC, (απυρεκτικός, from α, and πυρεσσειν, 'to be feverish'.) Apyretic.

APYRENOLE, Apyromele.

APYRETIC, *Apyreticus*, *Apyreetic*, *Apyreeticus*, *Apyretus*, (απυρετικός, from α, and πυρετός, 'fever'.) Without fever. This epithet is given to days in which there is no paroxysm of a disease, as in the case of an intermittent, as well as to some local affections which do not induce fever. Urticaria is sometimes called an *appyretic exanthema*.

APYREX'IA, (απυρεξία, *A'pyrezy*. Absence of fever: *Dialept'ia*, *Dialept'is*, *Dialip'is*, *Tem-pus intercala'rē*, *Interval'lum*, *Intermis'sio*. Apyrexia is the condition of an intermittent fever between the paroxysms. The duration of the apyrexia consequently depends on the type of the intermittent. Occasionally, the term has been applied to the cessation of the febrile condition in acute diseases.

APYROME'LĒ, *Apyrenome'lē*, (απυρομηλη, from α, πυρρον, 'a nut,' and μηλη, 'a sound'.) A sound or probe, without a button or nut. It is the *Melo'tis*, *Speci'illum auricula'rium* or *Auric'ular sound* of Galen.

AQUA, Urine, Water.—a. Acidi carbonici, Acidulous water.—a. Acidula hydrosulphurata, Naples water (factitious).—a. Aeris fixi, Acidulous water (simple).—a. Alkalina oxymuriatica, *Eau de Javelle*.—a. Aluminis composita, Liqueur aluminis compositus.—a. Aluminosa Bateana, Liqueur aluminis compositus.—a. Ammonie, Liqueur ammonie.—a. Ammonie acetatis, Liqueur ammonie acetatis.—a. Ammonie carbonatis, Liqueur ammonie subcarbonatis.—a. Ammonie caustica, Liqueur ammonie.—a. Ammonie fortior, see Liqueur ammonie.—a. Amnii, Liqueur Amnii.

AQUA AMYGDALA'RUM CONCENTRA'TA, *A. amygdal'arum amara'rum*, (F.) *Eau d'Amandes amères*, *Water of bitter almonds*. Made by bruising well two pounds of bitter almonds; adding, whilst triturating, ten pounds of spring water, and four pounds of alcohol; letting the mixture rest in a well-closed vessel, and then distilling two pounds. Used instead of the Aqua Laurocerasi, and the Hydrocyanic acid. It must be given with great caution.

An *Aqua amygdal'arum amara'rum*, *Bitter Almond water*, has been introduced into the Ph. U. S. (*Ol. amygdal. amar. Mxxvj*)—*Magnes. Carbon. ʒj*; *Aque Oij.* Dose. fʒss.

AQUA ANETHI, see *Anethum graveolens*.—a. Anisi, see *Pimpinella anisum*.—a. Anisi fortis, Spiritus anisi.—a. Aquisgranensis, see *Aix-la-Chapelle*.—a. Auditoria, Cotunnus, liqueur of.—a. Aurantii, see *Citrus aurantium*.—a. Azotica oxygenata, Aqua nitrogenii protoxydi.—a. Balsamica arterialis, Aqua Binellii.—a. Bareginensis, *Baréges water*.—a. Barytæ Muriat, see *Baryta*, muriate of.—a. Belliucana, Balaruc waters.—a. Benedicta, Liqueur calcis.—a. Benedicta composita, Liqueur calcis compositus.—a. Benedicta Rulandi, Vinum antimonii tartarizati.

AQUA BINEL'LI, *Aqua Binelli*, *Aqua Balsam'ica arterial'is*, *Binelli's styptic*, (F.) *Eau de Binelli*. A celebrated Italian hæmostatic, invented

by one Binelli. Its composition is unknown, but its virtues have been ascribed to creasote; although there is reason for believing it to possess no more activity than cold water. *Aqua Monterossii*, (F.) *Eau de Monterossi*, appears to be of analogous composition.

AQUA BORVONENSIS, Bourbonne-les-Baines, mineral waters of.—a. Bristolensis, Bristol water.

AQUA BROCCHE'RII, *Aqua Brocchieri*, *Brocchieri* or *Brocchiari water*, *Brocchieri's styptic*, (F.) *Eau de Brocchieri*, *Eau styptique de Brocchieri*. A supposed styptic, which made much noise at Paris at one time. It is devoid of efficacy. Dr. Paris found nothing in it but water perfumed by some vegetable essence.

AQUA CALCARE USTÆ, Liqueur calcis.—a. Calcis, Liqueur Calcis.—a. Calcis composita, Liqueur calcis compositus.—a. Camphoræ, Mistura camphoræ.—a. Camphorata, Bates's, see *Cupri sulphas*.—a. Carbonatis sodæ acidula, Acidulous water, simple.—a. Carmelitana, see *Melissa*.—a. Carui, see *Carum Carui*.—a. Cassiæ, see *Laurus cassia*.—a. Catapultarum, *Arquebuseade*, *eau d'*.—a. Chlorini, see *Chlorine*.

AQUA CINNAMO'MI, *Cinn'amon wa'ter*, (Sc.) *Cannel water*. Distilled water of Cinnamon Bark. Prepared also in the following manner. *Ol. Cinnam. fʒss*; *Magnes. Carbon. ʒj*; *Aq. destillat. Oij.* Rub the oil and carbonate of magnesia; add the water gradually, and filter. (Ph. U. S.)

AQUA CINNAMOMI FORTIS, Spiritus Cinnamomi.

—a. Colcestrensis, Colchester, mineral waters of.

AQUA COLORA'TA, 'coloured water.' A name given to a prescription in which simple coloured water is contained. Used in hospital cases, more especially where a *placebo* is demanded.

AQUA CREASOTI, see *Creasote*.—a. Cupri ammoniati, Liqueur c. a.—a. Cupri vitriolati composita, Liqueur cupri sulphatis composita.—a. Destillata, Water, distilled.

AQUA FABRO'RUM, 'Water of smiths.' Water in which hot iron has been quenched. A feeble chalybeate.

AQUA FLORUM AURANTII, see *Citrus aurantium*.—a. Fluviatilis, Water, river.

AQUA FENIC'ULI, *Fennel Water*. The distilled water of fennel seed. It may be prepared also like the aqua cinnamomi.

AQUA FONTANA, Water, spring.—a. Fortis, Nitric acid.—a. Goulardi, Liqueur Plumbi subacetatis dilutus.—a. Hepatica, Hydrosulphuretted water.—a. Hordeata, Decoctum hordei.—a. Hungarica, Spiritus Rosmarinus.—a. Imbrium, Water, rain.—a. Infernalis, see *Toddy*.—a. Interens, Anasarea.—a. Inter Cutem, Anasarea.—a. Javellensis, *Eau de Javelle*.—a. Juniperi composita, Spiritus juniperi compositus.—a. Kali, Liqueur potassæ subcarbonatis.—a. Kali caustici, Liqueur potassæ.—a. Kali præparati, Liqueur potassæ subcarbonatis.—a. Kali puri, Liqueur potassæ.—a. Kali subcarbonatis, Liqueur potassæ subcarbonatis.—a. Labyrinthi, Cotunnus, liqueur of.—a. Lactis, Serum lactis.—a. ex Lacu, Water, lake.—a. Lauro-cerasi, see *Prunus Lauro-cerasus*.—a. Laxativa Viennensis, Infusum Sennæ compositum.—a. Lithargyri acetati composita, Liqueur plumbi subacetatis dilutus.—a. Lucicæ, Spiritus ammoniæ succinatus.—a. Marina, Water, sea.—a. Medicata, Water, mineral.

AQUA MENTHÆ PIPERITÆ, *Peppermint Water*. The distilled water of peppermint. It may be pared like the Aqua cinnamomi.

AQUA MENTHÆ PIPERITIDIS SPIRITUOSA, Spiritus menthæ piperitæ.—a. Menthæ pulegii, see *Mentha pulegium*.—a. Menthæ viridis, *Spearmint water*, see *Aqua menthæ piperitæ*.—a. Menthæ vulgaris spirituosa, Spiritus menthæ viridis.—a. Mineralis, Water, mineral.—a. Mirabilis, Spiritus pimentæ.—a. Monterossii, see *Aqua Binellii*.—a. Mulsa, Hydromeli.—a. Natri Oxymuriatici, Liqueur sodæ chlorinatæ.—a. Neapolitana, Naples

water (factitious)—a. Nephritica, Spiritus myristica.

AQUA NITROGENII PROTOXYDI, *Aqua azotica oxygenata*, *Protoxide of Nitrogen Water*, *Searle's patent oxygenous aerated water*. A patent solution of protoxide of nitrogen, said to contain five times its own bulk of gas. It has been recommended as a nerve, and excitant in nervous conditions, dyspepsia, &c. It has also been used in cholera, and to counteract the evil consequences of drunkenness. The dose is fʒvj, or ʒviii, two or three times a day; or, in dyspepsia, as a beverage between meals.

AQUA NIVATA, Water, snow—a. Nucis moschatae, Spiritus myristicae—a. Ophthalmica, Liquor zinci sulphatis cum camphorâ—a. Pagliarii, Hæmostatica, Pagliarii's—a. Paludosa, Water, marsh—a. Pedum, Urine—a. Pericardii, see Pericardium—a. Phagedenica, see Hydrargyri Oxymurias—a. Picea, see Pinus sylvestris—a. Picis, see Pinus sylvestris—a. Pimentæ, see Myrtus Pimenta—a. Plumbi, Liquor Plumbi subacetatis dilutus—a. Pluvialis, Water, rain—a. Potassæ, Liquor potassæ—a. Pulegii, see Mentha pulegium—a. Pulegii spirituosus, Spiritus pulegii—a. Pura, see Water—a. Putealis, Water, well—a. ex Puteo, Water, well—a. Rabelli, Elixir acidum Halleri—a. Raphani composita, Spiritus armoracæ compositus—a. Regia, Nitromuriatic acid—a. Reginae Hungariæ, Spiritus Rosmarini.

AQUA RO'SÆ, *Rhodostegma*, *Rose Water*, (*Ros. centifol.* lbviii; *Aquæ cong.* ij. M. Distil a gallon—Ph. U. S.)

AQUA SALUBRIS, Water, mineral—a. Sambuci, see Sambucus—a. Sappharina, Liquor Cupri ammoniati—a. Saturni, Liquor plumbi subacetatis dilutus—a. Sclopetaria, *Arquebuseade eau d'*—a. Seminum anisi composita, Spiritus anisi—a. Seminum carui fortis, Spiritus carui—a. Sodæ effervescens, Acidulous water, simple—a. Soteria, Water, mineral—a. Stygia, Nitro-muriatic acid—a. Styptica, Liquor cupri sulphatis compositus—a. Sulphurata simplex, Hydrosulphuretted water—a. Sulphureti ammoniæ, Liquor fumans Boylii—a. Thediana, *Arquebuseade eau d'*—a. Theriacalis Bezoardica, Chylostagma diaphoreticum Mindereri—a. Tofani, Liquor arsenicalis—a. Tosti panis, Toast water—a. Traumatica Thedenii, *Arquebuseade eau d'*—a. Vegetomineralis, Liquor plumbi subacetatis dilutus—a. Viciensis, Vichy water—a. Vitriolica cærulea, Solutio sulphatis cupri composita—a. Vitriolica camphorata, Liquor zinci sulphatis cum camphorâ—a. Vulneraria, *Arquebuseade eau d'*—a. Zinci vitriolati cum camphorâ, Liquor zinci sulphatis cum camphorâ.

AQUÆ ACIDULÆ, Acidulous waters—a. Badiguae, Bath, Mineral waters of—a. Badizæ, Bath, Mineral waters of—a. Bathoniæ, Bath, Mineral waters of—a. Buxtoniensis, Buxton, Mineral waters of—a. Calidæ, *Aigues chaudes*—a. Cantuariensis, Canterbury, waters of—a. Chalybeatæ, Waters, mineral, chalybeate.

AQUÆ DESTILLATÆ, *Hydrolata*, *Distilled Waters*, (F.) *Hydrolats*. These are made by putting vegetable substances, as roses, mint, pennyroyal, &c., into a still with water, and drawing off as much as is found to possess the aromatic properties of the plant. To every gallon of the distilled water, 5 oz. of spirit should be added to preserve it. The simple distilled waters are sometimes called *Aquæ stillatit'ix simplicis*; the *epirituous*, *Aquæ stillatit'ix spirituosæ*, but more commonly *Spiritus*.

AQUÆ FERROSÆ, Waters, mineral, chalybeate.

AQUÆ MARTIALES, Waters, mineral, chalybeate.

AQUÆ MEDICATÆ, *Mollified waters*, include, in the Pharmacopœia of the United States, preparations consisting of waters impregnated with some medicinal substance, which are not arranged in any other class. Among these are the "Waters,"

and "Distilled waters" of the British pharmacopœias.

AQUÆ METUS, ('dread of water,') *Hydrophobia*—a. Minerales acidulæ, Waters, mineral, gaseous—a. Minerales ferruginosæ, Waters, mineral, chalybeate—a. Minerales sulphuræ, Waters, mineral, sulphureous—a. Solis, Bath, mineral waters of—a. Stillatiæ, *Aquæ destillatæ*.

AQUÆDUCTUS, *Aqueduct* (*aqua*, 'water,' and *ducere*, *ductum*, 'to lead,') (F.) *Aqueduc*. Properly, a canal for conducting water from one place to another. Anatomists have used it to designate certain canals.

AQUÆDUCTUS CEREBRI, Infundibulum of the brain.

AQUÆDUCTUS COCH'LEÆ, (F.) *Aqueduc du Limacon*:—a very narrow canal which proceeds from the tympanic scale of the cochlea to the posterior edge of the *pars petrosa*.

AQUÆDUCTUS COTUNNI, *Aquæductus vestibuli*. **AQUÆDUCTUS FALLOPII**, *Canalis Fallopii*, (F.) *Canal spiroïde de l'os temporal* of Chaussier, *Aqueduc de Fallope*. A canal in the *pars petrosa* of the temporal bone, which extends from the meatus auditorius internus to the foramen stylo-mastoideum, and gives passage to the facial nerve. The opening into this aqueduct is called *Hiat'us Fallopii*, *H. Canalis Fallopii*, *Fissu'ra Canalis Fallopii*, *Foramen Tari'ni*, *F. anony'mum Ferreinii*.

AQUÆDUCTUS SYLVII, *Canalis eminentiæ quadrigeninæ*, *Pter ad quatuor ventriculorum seu à tercio ad quatuor ventriculorum*, *Canalis medius*, (F.) *Aqueduc de Sylvius*, *Canal intermédiaire des ventricules* of Chaussier. A canal forming a communication between the third and fourth ventricles of the brain. The ventricular opening is termed *a'nus cerebri*.

AQUÆDUCTUS VESTIBULI, *Aquæductus Cotunnii*, *Canal' of Cotunnii*, (F.) *Aqueduc du vestibule* ou *Aqueduc de Cotugno*. This begins in the vestibule, near the common orifice of the two semicircular canals, and opens at the posterior surface of the *pars petrosa*.

AQUALICULUS, (diminutive of *aqualis*, 'a water-pot,') That part of the abdomen which extends from the umbilicus to the pubes. See Hypogastrium. It has also been applied to the stomach or intestinal canal.

AQUAS/TER. A word used by Paracelsus to express the visions or hallucinations of patients.

AQUEDUC, *Aqueduct*—a. *de Cotugno*, *Aquæductus vestibuli*—a. *de Fallope*, *Aquæductus Fallopii*—a. *du Limacon*, *Aquæductus cochleæ*—a. *de Sylvio*, *Aquæductus Sylvii*—a. *du Vestibule*, *Aquæductus vestibuli*.

AQUEDUCT, *Aquæductus*.

A'QUEOUS, *A'queus*, *Aquo'sus*, *Hydato'sdes*. *Hydro'sdes*, (*aqua*, 'water,') (F.) *Aqueus*, *Watery*. The absorbents or lymphatics are sometimes called, in France, *Conduits ou Canaux aqueux*.

AQUEOUS HUMOUR OF THE EYE, *Hu'mor aqueus*, *Oëci'sdes*, *Oo'sdes*, *Hydato'sdes*, *Hydato'sdes*, *Oci'tus*, *seu Oci'for'mis hu'mor*, *Albugin'eous hu'mor*, (F.) *Humeur aqueuse*. The limpid fluid which fills the two chambers of the eye, from the cornea to the crystalline, and which is, consequently, in contact with the two surfaces of the iris. Quantity, 5 or 6 grains: s. g. 1.0003. It contains albumen, chloride of sodium, and phosphate of lime in small quantity; and is enveloped in a fine membrane:—the membrane of the aqueous humour, *Tu'nica pro'pria seu Vagi'na seu Membra'na seu Cap'sula hu'moris a'quei seu Membra'na Demuria'na seu Descemet'i*, *Membrane of Demours'* or of *Descemet'*; although these last terms are by some appropriated to a third layer of the cornea.

AQUEUS, *Aqueous*.

AQUIDUCA, (*aqua*, 'water,' and *ducere*, 'to lead,' Hydragogues.

AQUIFOLIUM, (*acus*, 'a needle,' and *folium*, 'a leaf,') *Ilex aquifolium*—a. *Foliis deciduis*, *Prinos*.

AQUILA, *Hydrargyri submuriatis*, Sulphur.

The alchemists used this word for sublimed sal ammoniac, precipitated mercury, arsenic, sulphur, and the philosopher's stone. See *Hydrargyri Submuriatis*, and Sulphur.

AQUILA CELESTIS; a sort of panacea, of which mercury was a constituent.

AQUILA LACRYME; a liquor prepared from several ingredients, especially from calomel.

AQUILA PHILOSOPHICUM. The alchemists, whose terms were always mysterious, called mercury thus, when reduced to its original form.

AQUILA VENERIS; an ancient preparation, made by subliming verdigris and sal ammoniac.

AQUILÆ VENÆ, Temporal veins.

AQUILEGIA, (from *aquila*, 'the eagle,' owing to some fancied resemblance of the spurs to talons.)

A. vulgaris, seu *sylvestris*, seu *Alpina*, *Complanata*, *Colombina* or *Columbine*, (F.) *Ancolie*. Order, Ranunculaceæ. The seeds, herb, and flowers were formerly used in jaundice and cutaneous diseases. They are still retained in many of the Pharmacopœias of continental Europe.

AQUILEGIA ALPINA, *Aquilegia*.

AQUILEGIA CANADENSIS, *Wild Columbine*, is indigenous, and flowers in April and June. The seeds are said to be tonic.

AQUILEGIA SYLVESTRIS, *Aquilegia*—a. *Vulgaris*, *Aquilegia*.

AQUILICIA SAMBUCINA, (*aqua* 'water,' and *elcire*, 'to attract,' because it grows in moist places. [?]) An East India plant, the decoction of whose root is used in indigestion to allay heartburn.

AQUIPAROUS, (F.) *Aquipare*, (*aqua*, 'water,' and *pario*, 'I bring forth,') An epithet for glands which, like the parotid, secrete much water, in contradistinction to the submaxillary glands, which are *muciparous*.

AQUOCAPSULITIS, *Aquomembranitis*.

AQUOMEMBRANITIS, *Keratitidis*, *Aquocapsulitis*, *Hydromeningitis*. Inflammation of the anterior chamber of the eye. A badly compounded term, denoting inflammation of the capsule or membrane of the aqueous humour.

AQUOSUS, *Hydatoid*.

AQUULA, (dim. of *aqua*, 'water,') *Ceratocele*, *Hydatid*, *Hydroa*—a. *Acustica*, *Cotunnus*, liquor of.

AQUULA seu A'QUA MORGAGNI. The minute portion of water which escapes when an opening is made into the capsule of the crystalline.

AQUULA VITREA AUDITIVA, *Vitrine auditivæ*.

AR, in composition, see Ad.

A'RA PARVA, 'a small altar,'—a kind of bandage invented by Sostratus, which represents the corners of an altar.

ARABICA ANTIDOTUS HEPATICA, *Arabice Hepaticæ Antidotæ*. A powder composed of myrrh, costus, white pepper, &c. It was administered in new wine.

ARABICUS LAPIS. A sort of white marble, analogous to alabaster, found in Arabia. It was regarded as absorbent and desiccative, and was employed in hemorrhoids.

ARABIS BARBAREA. *Erysimum barbarea*.

ARABIS MALAGMA. An antiscorbutic medicine, composed of myrrh, olibanum, wax, sal ammoniac, iron pyrites, &c.

ARABISTS, see Arabs, medicine of the.

ARABS, MEDICINE OF THE. The Arabians kept the torch of medical science illuminated during a dark period of the middle ages. Before the year of the Hegira, they had schools

of medicine; but these were most flourishing during the 10th, 11th, and 12th centuries. The chief additions made by them to medical science were in the departments of pharmacy and in the description of diseases. Their principal writers were Avicenna, Serapion, Averrhoes, Hali Abbas, Moses Maimonides, Avenzoar, Rhazes, Albucasis, &c. The disciples of the Arab school were called *Arabists*.

ARACACHA, *Conium moschatum*.

ARACHIDNA HYPOGÆA, *Arachis hypogæa*.

ARACHIS AFRICANA, (*a*, and *ῥαχis*, 'a branch,'—having no branches.) *A. hypogæa*—a. *Americana*, *A. Hypogæa*.

ARACHIS HYPOGÆA, *A. Americana* seu *Africana*, *Arachidna hypogæa*, *Ground nut*, *Peanut*, *Earth almond*, (S.) *Mane*; erroneously called *Pistachio nut* in the South; *Pindars* of the West Indies. Ord. Leguminosæ. Cultivated in the Southern States. The seeds are oily, and are much eaten. A kind of inferior chocolate may be made of them.

ARACHNE, (*αράχνη*, 'a spider,' 'a cobweb,') Hence,

ARACHNITIS, more properly *Arachnoiditis*, *Arachnoiditis*, *Inflammation of the Arachnoid*. A variety of phrenitis.

ARACHNOIDITIS, *Arachnitis*.

ARACHNOID, *Arachnoides*, *Arachnoides*, *Arachnoides*, (*arachne*, and *idos*, 'form, resemblance,') Resembling a spider's web.

ARACHNOID CANAL, see Canal, arachnoid.

ARACHNOID OF THE EYE. The lining membrane of a cavity, supposed by some to exist between the sclerotic and choroid.

ARACHNOID MEMBRANE, *Membrana media seu serosa*, *Tunica arachnoidea seu crystallina*, *Membrana*. A name given to several membranes, which, by their extreme thinness, resemble spider-webs. Celsus and Galen called thus the membrane of the vitreous humour,—the *tunica hyaloidea*. The moderns use it now for one of the membranes of the brain, situate between the dura mater and pia mater. It is a serous membrane, and composed of two layers; the external being confounded, in the greater part of its extent, with the dura mater, and, like it, lining the interior of the cranium and spinal canal; the other being extended over the brain, from which it is separated by the pia mater, without passing into the sinuosities between the convolutions, and penetrating into the interior of the brain by an opening at its posterior part under the corpus callosum. It forms a part of the investing sheath of the nerves, as they pass from the encephalic cavities. Its chief uses seem to be:—to envelop, and, in some measure, protect the brain, and to secrete a fluid for the purpose of keeping it in a state best adapted for the proper performance of its functions.

ARACHNOIDITIS, (*arachnoid*, and *itis*,) *Arachnitis*.

ARACK', *Arrack*, *Rack*; (East Indian.) A spirituous liquor made in India in various ways, often from rice, sometimes from sugar fermented along with the juice of the cocoa nut; frequently from toddy, the juice which flows from the cocoa-nut tree by incision, and from other substances. It is a strong, heating spirit.

ARACK, Mock, is made by adding *Ëij* of Benzoic acid to a quart of rum. The celebrated Vauxhall punch was made with such arack.

ARACOUCHINI, *Icica aracouchini*.

ARACUS AROMATICUS, *Vanilla*.

AR'ADOS, (*αράδος*, from *αράδω*, 'I disturb,') The agitation excited in the stomach by the coction of aliments of different nature. Likewise, the motion produced by cathartics.

ARÆA, (*αῖτα*,) *Abdomen*, *Intestine* (small.)

ARÆOMA, (ἀραιωμα, from ἀραιος, 'porous,') Interstice.

ARÆOMETER, Areometer.

ARÆOTICA, (ἀραιωτικά, from ἀραιω, 'I rarefy,') Medicines supposed to have the quality of rarefying the humours. See Rarefaciens.

ARAKI, see Spirit.

ARALIA CANADENSIS, *Panax quinquefolium*.

ARALIA HISPIDA, *Dwarf Elder*, *Ord. Araliaceæ*, is said to be diuretic, and has been recommended, in decoction, in dropsy.

ARALIA NUDICAULIS, *Nardus Americanus*, *Small Spike-nard*, *Wild Liquorice*, *Sweet root*, *False or Wild Sarsaparilla*, (F.) *Petit nard*. This American plant is said to be a mild stimulant and diaphoretic, and has been recommended as a substitute for sarsaparilla. It is used, also, as a tonic.

ARALIA RACEMOSA, *American Spike-nard*, has the same properties as *A. Nudicaulis*.

ARALIA SPINOSA, *Angelica Tree*, *Prickly Ash*, *Toothache Tree*, *Spike-nard Tree*, *Prickly Elder*, *Shot-bush*, *Pigeon Tree*. Its properties are not clear. The berries, and a tincture of them, have been employed, it is said, successfully in toothache. A spirituous infusion has also been used in colic.

ARANEÆ, *Aranæe Tela*—*a. Tarentula*, see Tarentula.

ARANEÆ TELÆ, *Ara'nea*, *Ara'neum*, *Spider's web*, *Cobweb*, (F.) *Toile d'Araignée*. Formerly, this substance was much employed, and supposed to possess extraordinary virtues, especially when applied to the wrists. It has been used again internally in intermittents. The spider itself, softened into a plaster and applied to the forehead and temples, is said by Dioscorides to prevent ague. Cobweb is a mechanical styptic, and is so applied at times.

ARANEOSA URINA. A term applied to the urine when loaded with filaments like cobwebs.

ARANEOSUS (PULSUS); a term employed to express extreme weakness of pulse; when the movements resemble those of a delicate net raised by the wind.

ARANEUM, *Aranæe Tela*.

ARANEUM ULCUS, *Astaktil'los*. A name given by Paracelsus to a malignant, gangrenous ulcer, extending from the feet to the legs.

ARARA, *Myrobalanus citrina*.

ARATRUM, (L.) ('a plough,') *Vomer*.

ARAUCARIA DOMBEYI, *Dombeya excelsa*.

ARBOISE, *Arbutus unedo*.

ARBOL DE LECHE, (S.) ('milk tree,') *Galactodendron utile*.

ARBOR ('a tree') BENIVI, Benjamin—a. *Indica*, *Laurus cassia*—*a. Maris*, *Coral*—*a. Thurifera*, *Juniperus Lycia*—*a. Uteri Vivificans*, *Palmæ uteri plicatæ*.

ARBOR VITÆ, ('tree of life,') (F.) *Arbre de vie*. A name given to an arborescent appearance, observed on cutting the cerebellum longitudinally; and which results from the particular arrangement of the white substance with the cineritious. Also, the *Thuya occidentalis*.

ARBOR VITÆ, AMERICAN, *Thuya occidentalis*—*a. Vitæ Uterinus*, *Palmæ uteri plicatæ*—*a. Vitæ of the Uterus*, *Palmæ uteri plicatæ*.

ARBORES. A morbid alteration of the skin, which precedes its ulceration, according to Ruland.

ARBORIZATION, *Arborisatio*, (F.) *Arborisation*, (*arbor*, gen. *arboris*, 'a tree,') The figure or appearance of a tree or plant. The capillary vessels, when injected, as in inflammation, frequently appear under the form of Arborizations.

ARBOUSIER, *Arbutus unedo*.

ARBRE DE VIE, ('tree of life,') *Arbor Vitæ*. ARBUSCULA (dim. of *arbor*) GUMMIFERA BRAZILIENSIS, *Hypericum bacciferum*.

ARBUTUS, *A. Unedo*—*a. Trailing*, *A. Ursi*, *Epigæa repens*.

ARBUTUS UNE'DO, *Arbutus*, *Andrach'næ*, *Une'do*, *U. papyracea*, *Com'arus*, (F.) *Arbousier*, *Arboise*. A decoction of the leaves is astringent, and has been used in diarrhœa.

ARBUTUS U'VA UR'SI, *Arctostaphylos U'va ur'si*, *Mair'nia u'va ur'si*. *Ord. Ericaceæ*. *Sex. Syst. Decandria Monogynia*. (F.) *Busserolle* ou *Raisin d'Ours*. The leaves—(U'va Ur'si, Ph. U. S.)—of this plant are tonic and astringent, and have been employed, chiefly, in diseases of the urinary organs, and also as a parturifacient. Dose of the powder from gr. xv to ʒss. The English names are *Trailing Arbutus*, *Bear's Whortleberry* or *Bearberry*, *Mountain-box*, *Redberry*, *Upland Cranberry*, *Foxberry*, *Checkerberry*, (Sc.) *Braw-lins*.

ARC, *Arch*, *Ar'cus*, (F.) *Arc*, *Arcade* (diminutive). Any part of the body resembling an arch in form; as the *Arch of the colon*, (F.) *Arc du colon*,—the transverse portion of that intestine:—*Arch or Cross of the Aorta*, *Ar'cus aor'tæ*, (F.) *Crosse de l'Aorte*, &c., the turn which the aorta takes in the thorax.

ARCA ARCANORUM, *Hydrargyrum*—*a. Cordis*, *Pericardium*.

ARCADE, see *Arc*—*a. Anastomotique*, *Arch*, *anastomotique*—*a. Crurale*, *Crural arch*—*a. Inguinale*, *Crural arch*—*a. Orbitaire*, *Orbital arch*—*a. Pubienne*, *Pubic arch*—*a. Zygomaticque*, *Zygomatic arch*.

ARCADES DENTAIRES, *Dental arches*—*a. Palmaires*, *Palmar arches*.

ARCADI-TEMPORO-MAXILLAIRE, *Temporals*.

ARCE'US or ARCE'US, BALSAM OF. See *Balsam of Arceus*.

ARCANSON, *Colophonia*.

ARCANUM, (*arca*, 'a chest,') A secret, a *nostrum*, a *quack* or *empirical medicine*, (F.) *Arceane*. A remedy whose composition is kept secret; but which is reputed to possess great efficacy.

ARCANUM CORALLINUM, *Hydrargyri nitricooxydum*—*a. Duplicatum*, *Potassæ sulphas*—*a. Tartari*, *Potassæ acetas*.

ARCEAU, *Arculus*, *Cradle*.

ARCEION, (ἀρκιον,) *Arctium*.

ARCEUTHOS, (ἀρκευθος,) *Juniperus communis*.

ARCH, *Arco*—*a. Anostomatic*, (F.) *Arcade Anostomotique*, is the union of two vessels, which anastomose by describing a curved line. The vessels of the mesentery inosculate in this manner.

ARCH OF THE AORTA, see *Arc* and *Aorta*—*a. Crural*, see *Crural arch*—*a. Femoral*, see *Crural arch*—*a. Gluteal*, see *Gluteal aponeurosis*—*a. Hæmal*, see *Hæmal arch*—*a. Inguinal*, see *Crural arch*—*a. Orbital*, see *Orbital arch*—*a. of the Palate*, see *Palate bone*—*a. of the Pubis*, see *Pubic arch*—*a. Subpubic*, see *Subpubic arch*—*a. Superciliary*, see *Superciliary arches*—*a. of a Vertebra*, see *Vertebra*—*a. Zygomatic*, see *Zygomatic arch*.

ARCHES OF THE PAL'ATE. These are two in number on each side of the throat, one of which is termed *anterior*, the other *posterior*.

The *anterior arch* arises from the middle of the *velum palati*, at the side of the *uvula*, and is fixed to the edge of the base of the tongue.

The *posterior arch* has its origin, likewise, from the side of the *uvula*, and passes downwards to be inserted into the side of the pharynx. The *anterior arch* contains the *circumflexus palati*, and forms the *isthmus faucium*. The *posterior arch* has within it the *levator palati*, and between the arches are the *tonsils*.

ARCHÆ'US, *Archæus*, (ἀρχη, 'commencement,') (F.) *Archée*. A word invented by Basil Valentine, and afterwards adopted by Paracelsus

and Van Helmont. The latter used it for the internal principle of our motions and actions. This archæus, according to Van Helmont, immaterial principle, existing prior to fecundation, and presiding over the development of the body, and over all organic phenomena. Besides this chief archæus, whose seat Van Helmont placed in the upper orifice of the stomach, he admitted several of a subordinate character, which had to execute its orders; one, for instance, in each organ, to preside over its functions; each of them being subject to anger, caprice, terror, and every human feeling.

ARCHANGEL NEW, MINERAL SPRINGS. About twenty miles to the north of New Archangel, Sitka Island, on the N. W. coast of North America, are some thermal sulphureous waters, the temperature of one of which is upwards of 153° of Fahr. They are much celebrated, according to Sir Geo. Simpson.

ARCHANGELICA, ('archangelic,' from its virtues,) *Lamium album*—a. *Officinalis*, *Angelica*.

ARCHÊ, (αρχή,) *Initium*, *Principium*, *Primordium*, *Origo*, *Invasio*. The first attack of a disease.

ARCHECPTOMA, (archo, εἶ, 'out of,' and πτωμα, 'a fall,') *Proctocoele*.

ARCHÉE. Archæus.

ARCHELL, CANARY, *Lichen roccella*.

ARCHELOG'IA, (αρχή, 'beginning,' and λογος, 'a discourse,') A treatise on fundamental principles—of medicine, for example.

ARCHEN'DA. A powder of the leaves of the *ligustrum*, used by the Egyptians after bathing, to obviate the unpleasant odour of the feet.

ARCHIA'TER, *Archiatrus*, *Protomedicus*, *Protiatras*, *Superpositus medicorum*, (αρχος, 'a chief,' and ιατρος, 'a physician,') The original signification of this word is a matter of dispute. Some consider, with Mercurialis, that it meant physician to a prince, king, emperor, &c.; others, with C. Hoffmann, apply it to every physician who, by his situation, is raised above his colleagues. The former opinion seems to have prevailed—*Archiatre des Rois de France* being applied to the chief physician to the kings of France.

ARCHIG'ENI MORBI, (αρχή, 'beginning,' and γινωμαι, 'to be,') Acute diseases; because they hold the first rank.

ARCHIMAGIA, (αρχος, 'a chief,' and μαγεια, 'magic,') Chymistry.

ARCHINGEAY, MINERAL WATERS OF. Archingéay is situate in France, three leagues from St. Jean d'Angely. The waters are prized in all diseases. They seem to contain carbonate of lime, a little chloride of sodium, carbonate of iron, and some bitumen.

ARCHITECTURA APOPLECTICA, Apoplectic make.

ARCHITIS, (archo, and itis,) *Proctitis*, *Rectitis*.

ARCHO, (from αρχος, 'the anus, the rectum,') in composition, 'anus or rectum.'

ARCHOCELE, (archo, and κηλη, 'rupture,') *Proctocoele*.

ARCHOCYSTOSYR'INX, (archo, κυστις, 'bladder,' and σφυγξ, 'a pipe,') *Fistula of the anus and urinary bladder*.

ARCHOPTOMA, (archo, and πτωμα, 'a fall,') *Proctocoele*.

ARCHOPTOSIS, *Proctocoele*.

ARCHORRHAG'IA, (archo, and ῥινηναι, 'to break forth,') *Archorrhæa*. Hemorrhage from the anus.

ARCHORRHŒA, (archo, and ῥεω, 'I flow,') *Archorrhægia*.

ARCHOS, (αρχος,) *Anus*, *Rectum*.

ARCHOSTEGNOMA, (archo, στεγνωω, 'I constrict,' and oma,) *Stricture of the Rectum*.

ARCHOSTEGNOSIS, *Stricture of the Rectum*.

ARCHOSTENOSIS, (archo, στενος, 'narrow,' and osis,) *Stricture of the Rectum*.

ARCHOSYRINX, (archo, and σφυγξ, 'a pipe,') *Fistula in ano*.

AR'CIFORM, *Arceiformis*, (arx, gen. arcis, 'a top or ridge,' and forma, 'shape,') An epithet given to certain fibres, *Fibræ arceiformes*, of the anterior pyramids of the medulla oblongata, which take a curved course around the inferior extremity of each corpus olivare and ascend towards the cerebellum.

ARCTA'TIO, *Arctitu'do*, (arcto, arctatuni, 'to make narrow,') *Angusta'tio*, *Coarcta'tio*, (F.) *Rétrécissement*. Contraction of a natural opening or of a canal, and especially of the vulva, of the orifice of the uterus, or of the intestinal canal. Constipation, (see *Stegnosis*.) Reunion by suture or infibulation.

ARCTITUDO, *Arctatio*.

ARCTIUM, (from αρκτος, 'a bear,' owing to the rough, bristly fruit,) *A. lappa*—a. *Bardana*, *A. lappa*.

ARCTIUM LAP'PA. The root and seed of the *Clit'bur*, *Barda'na*, *Arce'ion*, *Arctium*, *A. barda'na*, seu *ma'jus* seu *mi'nus* seu *tomentosum*, *Flaphis*, *Lappa glabra*, *Lappa major* seu *persona'ta*, *Perolla'ta*, *Persolla'ta*, *Persolla'ta*, *Burdock*, (Old Eng.) *Clithe*, *Cuckold*, (F.) *Bardane*, *Glouteron*, *Herbe aux teigneux*. *Ord.* *Compositæ*. *Sex. Syst.* *Syngenesia aequalis*. The Root—*Lappa* (Ph. U. S.)—is diuretic; the seed are cathartic. It has been used in decoction in diseases of the skin and in syphilis.

ARCTIUM MAJUS, *A. lappa*—a. *Minus*, *A. lappa*—a. *Tomentosum*. *A. lappa*.

ARCTOPUS ECHIN'A'TUS, (αρκτος, 'a bear,' and πους, 'a foot,') A South African plant, *Ord.* *Umbellifera*, which is demulcent and diuretic, somewhat approaching sarsaparilla. The decoction of the root is employed in syphilis, lepra, and chronic cutaneous affections of all kinds.

ARCTOSTAPHYLOS UVA URSI, (αρκτος, 'a bear,' and σταφυλη, 'a grape,') *Arbutus uva ursi*.

ARCTU'RA, (from arcto, 'I press close,') The effects of a nail grown into the flesh, *Arctu'ra unguis*. See *Oncyhogryphosis*.

ARCTURA UN'GUIUM. The growing in or inversion of the nails. See *Oncyhogryphosis*.

ARCUA'TIO, (arcus, 'a bow,') *Concava'tio*. An anterior gibbosity or projection of the sternum.

ARCUEIL, MINERAL WATERS OF. Arcueil is about one league south of Paris. The water contains carbonic acid, carbonate of lime, sulphate of lime, chloride of sodium, and some deliquescent salts.

A celebrated society held its meetings at this village, of which Berthollet, Humboldt, Laplace, &c., were members.

ARCULA CORDIS, (dim. of arca, 'a press or chest,') *Pericardium*.

ARCULÆ, (pl. of arcula,) The Orbital Fosse.

ARC'ULUS, (dim. of arcus, 'an arch,') A small arch; a cradle, (F.) *Arceau*, *Archet*. A semicircular box or basket used for preventing the bed-clothes from coming in contact with injured or diseased parts. An ordinance of the Grand Duke of Tuscany forbade mothers to sleep with an infant near them, unless it was put under a solid cradle.

ARCUS, gen. *Arcus*. Arc or Arch—a. *Medullaris*, *Fornix*—a. *Senilis*, *Gerotoxon*—a. *Subpubicus*, *Subpubic arch*—a. *Superciliaris*, *Superciliary arches*—a. *Unguium*, see *Nail*—a. *Zygomatius*, *Zygomatich arch*.

ARDA, (αρδα,) *Excrement*.

ARDALOS, (αρδαλος,) *Excrement*.

ARD'ENT, *Ardens*, (*ardere*, 'to burn.') Hot, fiery.

ARDENT FEVER, (F.) *Fièvre ardente*. *Causus*, *Synocha* or inflammatory fever.

ARDENT or INFLAMED EYES, (F.) *Yeux ardents*. The eyes are so called when injected red.

ARDENT U'RINE, (F.) *Urine ardente*. Urine, hot, or of a deep red.

ARDESIA HIBERNICA, *Hibernicus lapis*.

ARDEUR, Ardor—*a. du Cœur*, Cardialgia—*a. d'Estomac*, Ardor ventriculi, Pyrosis—*a. de la Fièvre*, Ardor Febrilis—*a. d'Urine*, Ardor Urinæ.

ARDISIA HUMILIS, *Badulam*.

ARDOISE, see *Hibernicus lapis*.

ARDOR, gen. *Ardo'ris*, (F.) *Ardeur*. Heat. A feeling of burning, of violent heat; *Æstus*, *Æstusatio*, *Causo'ma*.

ARDOR FEBRI'LIS, (F.) *Ardeur de la Fièvre*. The hot period of fever.

ARDOR STOMACHI, Pyrosis.

ARDOR URI'NÆ, (F.) *Ardeur d'Urine*. A scalding sensation occasioned by the urine in passing over the inflamed mucous membrane of the urethra, or over the neck of the bladder.

ARDOR VENEREUS, Heat, (venereal.)

ARDOR VENTRIC'ULLI, *Ebullit'io Stom'achi*, *Heartburn*, (Sc.) *Heartaxes*, *Heartscald*, *Hersket*, (F.) *Ardeur d'Estomac*. See Cardialgia and Pyrosis.

A'REA, ('a void place,' 'an open surface.') A Latin word used by some authors to designate a variety of Alopecia, in which the hair changes colour, but does not fall off; also, *Porrigio decalvus*.

AREA GERMINATIVA, *Tache embryonnaire*.

AREA PELLU'CIDA. An elliptical depression in the ovum, filled with a pellucid fluid, in the centre of which is the germ.

AREA VASCULO'SA, see *Circulus venosus*.

ARE'CA. The fruit—*Are'ca nut*, *Be'tel nut*—of *Are'ca Cat'echu* seu *Faufel*, *Cann'ga*; *Ord. Palmæ*; *Sex. Syst.* Monoclea Monadelphica; (F.) *Arec*, is astringent and tonic, and enters into the composition of the *Be'tel*, the great masticatory of the Orientals.

ARECA CATECHU, see *Areca*—*a. Faufel*, see *Areca*.

AREFAC'TION, *Arefac'tio*, *Xeran'sis*, (*arefacere*, 'to make dry,' from *arere*, 'to dry,' and *facere*, 'to make.') The process of drying substances, prior to pulverization.

ARENA, (from *areo*, 'to be dry.') See *Gravel*, *Sand*.

ARENAMEN, *Bole Armenian*.

ARENA'TIO, *Inhumatio*, *Cho'sis*, *Sand* or *Earth Bath*, *Saburra'tio*. The application of hot sand to the body. *Pediluv'ia* [?] of sand were formerly used in *Ascites*.

ARENGA SACCHARIFERA, see *Toddy*.

ARENO'SA URI'NA, (*arena*, 'sand.') *Sand'y Urine*. Urine when it deposits a sandy sediment.

ARENO'SUS, *Sabulous*. Also, one who passes sandy urine.

ARENULA, (dim. of *arena*, 'sand.') See *Gravel*.

ARE'OLA, (dim. of *Area*), (F.) *Aire*, *Are'ole*, *Vacuole*. Anatomists understand by *Areole* the interstices between the fibres composing organs, or those existing between laminæ, or between vessels which interlace with each other.

Areola is also applied to the coloured circle *Ha'lo*, *Ha'los*, which surrounds the nipple, *Are'ola papilla'ris*, and which becomes much darker during pregnancy; as well as to the circle surrounding certain vesicles, pustules, &c., as the pustules of the small-pox, the vaccine vesicle, &c.

Chaussier, in such cases, recommends the word *Aure'ola*, (F.) *Auréole*.

AREOLA PAPILLARIS, see *Areola*—*a. Tubercles* of the, see *Mamma*.

ARE'OLAR, *Areola'ris*, (F.) *Aréolaire*. Appertaining to an areola.

AREOLAR EXHALATIONS are those recremental secretions which are effected within the organs of sense, as the aqueous, crystalline, and vitreous humours, &c., or in the areolar tissue of parenchymatous and other structures.

AREOLAR TISSUE, Cellular Tissue.

ARÉOLE, *Areola*.

AREOM'ETER, *Aræom'eter*, *Pycnom'eter*, *Gravim'eter*, *Alcoölom'eter*, *Alcoöm'eter*, *Aërostat'io Bal'ance*, (*ἀραιος*, 'light,' and *μετρον*, 'measure,' i. e. 'measure of lightness.') An instrument so called because first employed to take the specific gravity of fluids lighter than water. The *Areometer* of *Baumé*, which is the most used in Pharmacy, particularly in France, consists of a tube of glass, largely expanded towards its inferior extremity, and terminating below by a small ball, containing mercury or lead, which serves it as a balance, so that it may remain upright in the fluid. This tube is furnished with a graduated scale. If the fluid into which the *Areometer* is plunged be heavier than water, the instrument rises; if lighter, it sinks. There are various *Areometers*, as those of the Dutch, of Fahrenheit, Nicholson, &c. The *Areometer* is also called *Hydrom'eter*, (F.) *Hydromètre*, *Aréomètre*, *Pèse-liqueur*.

There are some hydrometers which have a general application for determining the specific gravities of liquids—as Fahrenheit's, Nicholson's, Guyton de Morveau's, and the common glass hydrometers, including *Baumé's*, *Cartier's*, *Twaddle's*, *Zanetti's*, and the specific gravity beads; others intended for special application—as for estimating the comparative strength of spirits; the comparative densities of syrups, oils, &c.—as *Gay Lussac's*, *Sikes's*, and *Dicat's* hydrometers, and the saccharometer, urinometer, and elæometer.

SCALE OF BAUMÉ'S HYDROMETER, WITH CORRESPONDING SPECIFIC GRAVITIES.

1. Ascending Scale for light liquids

Scale of Baumé.	Specific Gravities.	Substances.
70	700	Pure hydrocyanic acid.— <i>Gay Lussac</i> .
66	715	Very pure sulphuric ether.
60	742	The same concentrated.
50	782	
48	792	Equal parts of alcohol and ether.
42	819	Very pure alcohol for pharmaceutical purposes.
40	827	
36	847	Pure alcohol. Naphtha.
33	863	Alcohol of commerce.
32	868	Essential oil of turpentine.
30	878	
26	900	Hydrocyanic acid of Scheele and pure hydrocyanic acid, mixed with an equal portion of water.— <i>Robiquet</i> .
25	906	
23	915	Acetic ether.
		Nitric ether.
Id.	Id.	Muriatic ether.
22	923	Liquid ammonia.
		Olive oil.
Id.	Id.	
20	935	Brandy.
18	948	
13	980	Burgundy wine.
12	986	
11	993	Bordeaux wine.
10	1000	Distilled water.

1. *Descending Scale for heavy liquids.*

Scale of Baume.	Specific Gravities.	Substances.
0	1000	Common distilled water.
1	1007	Distilled vinegar.
2	1009	
3	1014	
4	1032	Cow's milk.
10	1075	Concentrated acetic acid.
12	1091	
20	1161	
21	1180	Liquid hydrochloric acid.
25	1210	
30	1261	
35	1321	Common nitric acid.
40	1384	
41	1398	
45	1444	Concentrated nitric acid.
Id.	Id.	
50	1532	
60	1714	Phosphoric acid for medical use.
66	1847	
70	1949	

[Scales of the hydrometer and the corresponding specific gravities are given in the United States Dispensatory, and in Parrish's Practical Pharmacy.]

ARES. A term invented by Paracelsus to designate the principle on which depends the form of mercury, sulphur, and salt. These the alchemists regarded as the three bodies that give birth to every other.

AR'ETÉ, (αρετή, 'power.') Mental or corporeal vigour.

ARETHU'SA, *A. bulbosa*; indigenous. Order, Orchidaceæ. The bruised bulbs are used in tooth-ache; and as cataplasms to tumours.

ARGEL, *Cynanchum oleæfolium*.

AR'GEMA, gen. *Argem'otis*, *Ar'gemon*, *Ar'gemus*, (ἀργεμα, ἀργεμον, ἀργεμος; from αἶψος, 'white.') *Fos'sula*, (F.) *Encavure*. A white spot or ulceration of the eye. See Leucoma.

ARGEMO'NE MEXICA'NA, (αργεμώνη), *Thorn Pop'py*, *Prick'ly Pop'py*, *Yell'low Thistle*. A native of Mexico, but naturalized in most parts of the world. Ord. Papaveraceæ. Sex. Syst. Polyandria Monogynia. The juice resembles gamboge, and has been used as a hydragogue. The seeds are employed in the West Indies as a substitute for ipecacuanha. They are also used as a cathartic.

ARGENSON, MIN'ERAL WATERS OF. A chalybeate situate at Argenson in Dauphiny: used in cases of obstruction, jaundice, &c.

ARGENT, (αργος, 'white,') *Argentum*—*a. Chlorure d'*, see *Argentum*—*a. et d'Ammoniaque, chlorure d'*, see *Argentum*—*a. Cyanure d'*, see *Argentum*—*a. Iodure d'*, see *Argentum*—*a. Oxide d'*, see *Argentum*—*a. Petit*, *Platinum*—*a. Vif*, *Hydrargyrum*.

ARGENTERIA, *Potentilla anserina*.

ARGENTI CHLORIDUM, see *Argentum*—*a. et Ammonia chloridum*, see *Argentum*—*a. et Ammonia chloruretum*, see *Argentum*—*a. Cyanidum*, see *Argentum*—*a. Cyanuretum*, see *Argentum*—*a. Iodidum*, see *Argentum*—*a. Ioduretum*, see *Argentum*.

ARGEN'TI NI'TRAS, *A. azo'tas*, *Argentum Nitratum* seu *Nitricum*, *Sal argenti*, (F.) *Nitrate* ou *Azotate d'Argent*, *Nitrate* of *Silver*. This preparation is sometimes kept in crystals, the *Ni'tras Argenti* in crystal'los concre'tus, *Nitrate d'Argent* crystallisé of the Codex of Paris, *Lu'na potab'ilis*, *Crystall'i Lu'ne*, *Argentum nitricum crystallisatum*, *Ni'tras argenti crystallinus*, *Nitrum luna'rè*, *Hydrago'num BOYL'EI*: generally, however, it is in the fused state: and it is this which is admitted

into most Pharmacopœias, and which, besides the name *Argenti Nitras*, is called *Argenti ni'tras fusus* seu *fusa* (Ph. U. S. 1863) seu *sumum*, *Cans'neum luna'rè*, *Lapis inferna'lis*, *Argentum nitricum fusum*, and *lunar caustic*, (F.) *Nitrate d'argent fondus*, *Pierre infernale*. It may be prepared as follows:—Take of *silver*, in small pieces, $\frac{3}{4}$ j; *nitric acid*, f $\frac{3}{4}$ vij, *distilled water*, f $\frac{3}{4}$ ij. Mix the acid with the water, and dissolve the silver in the mixture in a sand bath; then crystallize, or gradually increase the heat, so that the resulting salt may be dried. Melt this in a crucible over a gentle fire, and continue the heat until ebullition ceases; then immediately pour it into suitable moulds.

The virtues of nitrate of silver are tonic, and escharotic. It is given in chorea, epilepsy, &c.: locally, it is used in various cases as an escharotic. Dose, gr. 1–8 to gr. 1–4 in pill, three times a day.

When silver is combined with iodine, it is said to have the same effect as the nitrate, and not to produce the slate colour of the surface, which is apt to follow the protracted use of the latter.

ARGENTI OXIDUM, see *Argentum*.

ARGENTILLA VULGARIS, *Potentilla anserina*.

AR'GENTINE, *Argento'sus*, (from *argentum*.) Pertaining to silver; as an 'argentine solution,' or solution of a salt of silver.

ARGENTINE, *Potentilla anserina*.

ARGENTUM, *Ar'gyrus*, (αργος, 'white,') *Sil'ver*, *Lu'na*, *Dia'na*, (F.) *Argent*. A solid metal of a shining white appearance; insipid; inodorous; highly sonorous; malleable and ductile; somewhat hard; crystallizable in triangular pyramids; fusible a little above a red heat, and volatilizable; s. g. 10.4. Not used in medicine, unless in some places for silvering pills. **SILVER LEAF**, *Argentum foliatum*, is the state in which it is used for this purpose.

The **CHLO'RIDE**, (*Argenti chlo'ridum*, *Argentum muriat'icum* seu *chlora'tum* seu *sali'tum*, *Chlorure'tum Argenti*, *Chlor'uret* ou *Mu'riate* of *Silver*, (F.) *Chlorure d'Argent*); the **CYAN'URET**; the **IODIDE** (*Argenti Io'didum*, *Argentum Ioda'tum*, *Iodure'tum Argenti*, *Iod'uret* of *Silver*, (F.) *Iodure d'Argent*); the **OX'IDE** (*Argenti ox'idum*, *Argentum oxyda'tum*, (F.) *Oxide d'Argent*), and the **CHLO'RIDE** of **AMMO'NIA** and **SIL'VER** (*Argenti et Ammo'niæ chlo'ridum*, *Argentum muriat'icum ammoniacum*, *Chlorure'tum Argenti et Ammo'niæ*, *Chlo'ruret* of *Silver* and *Ammonia*, *Ammonio-chloride* of *Silver*, (F.) *Chlorure d'Argent et d'Ammoniaque*), have been used in syphilis. At first, these different preparations were administered intraleptically on the gums; the chloride, the cyanide and the iodide in the dose of 1–12th of a grain; the chloride of silver and ammonia in the dose of 1–14th of a grain, and the oxide of silver, and divided silver, in the dose of 1–8th and 1–4th of a grain. M. Serre, of Montpellier, who made many trials with them, soon found that these doses were too small; he therefore raised that of the chloride to 1–10th, and of the iodide to 1–8th of a grain, without any inconvenience resulting. The dose of the other preparations was likewise increased in a similar ratio. M. Serre extols the preparations of silver—used internally as well as intraleptically—as antisypilitics, but they are not to be depended upon.

The **Cy'anide** or **Cyan'uret**, *Argenti Cyanure'tum* seu *Cyan'idum*, *Argentum cyanogena'tum*, (F.) *Cyanure d'argent*, is thus directed to be prepared in the Pharmacopœia of the United States.—*Nitrate* of *Silver*, dissolved in *distilled water*, is put into a tubulated glass receiver: *Ferrocyamuret* of *Potassium*, dissolved in *distilled water*, is put into a tubulated retort, previously adapted to the receiver. *Dilute Sulphuric Acid* is added to the solution in the retort; and by means of a sand-bath and a moderate heat, distillation is carried on until the liquid that passes over no longer

produces a precipitate in the receiver. The precipitate is then washed with distilled water, and dried.

Oxide of Silver, Argenti Oxidum, is made by precipitating a solution of the *Nitrate of Silver* by solution of *Potassa*, drying the precipitate. (Ph. U. S.)

ARGENTUM CHLORATUM, see *Argentum*—a. Cyanogenatum, see *Argentum*.

ARGENTUM DIVISUM, *metallic silver*, in very fine powder, has been recommended internally in syphilis.

ARGENTUM FUGITIVUM, *Hydrargyrum*—a. Fumum, *Hydrargyrum*—a. Iodatium, see *Argentum*—a. Liquidum, *Hydrargyrum*—a. Mobile, *Hydrargyrum*—a. Muraticum, see *Argentum*—a. Muraticum Ammoniatum, see *Argentum*—a. Oxydatum, see *Argentum*—a. Salutum, see *Argentum*—a. Vivum, *Hydrargyrum*.

ARGILL. PURE, (*argos*, 'white.') *Argilla pura*. *ARGILE*, see *Argilla pura*—a. *Ochreuse Pâle*, *Bolus Alba*.

ARGILLA, (*αργίλλος* or *αργίλος*, 'white clay,' from *argos*, 'white,') *Argilla pura*—a. *Acetica*, *Alumina acetata*—a. *Bolus Flava*, *Terra Lemnia*—a. *Bolus rubra*, *Bole Armenian*—a. *Ferruginea rubra*, *Bole Armenian*—a. *Kalisulphurica*, *Alumen*—a. *Pallida*, *Bolus alba*.

ARGILLA PURA, *Terra Aluminis* seu *bola'ris*, seu *argilla'cea pura* seu *deputata* seu *hydra'ta*, *Alumina pura* seu *deputata*, *Oxidum aluminii*, pure *Argil* or *Alumina*, (F.) *Argile* ou *Alumine factice*. This substance, which is prepared by drying alum and exposing it, for twenty or twenty-five minutes, to a red heat, until the sulphuric acid is driven off, has been recommended in indigestion as antacid, as well as in vomiting and diarrhoea accompanied with acidity. The dose to a very young child is from 3ss to ʒj; to older children from ʒj to ʒij.

ARGILLA SULPHURICA ALCALISATA, *Alumen*—a. *Sulphurica usta*, *Alumen exsiccatum*—a. *Supersulphur alcahisatum*, *Alumen*—a. *Vitriolata*, *Alumen*.

ARGILLÆ ACETAS, *Alumina acetata*—a. *Sulphas*, *Alumina sulphas*.

ARGOL, RED, *Potassæ supertartas impurus*—a. *White*, *Potassæ supertartas impurus*.

ARGUMENTUM INTEGRITATIS, *Hymen*.

ARGYRIA, (*αργυρος*, 'silver,') The discoloration of the skin occasioned by the internal use of nitrate of silver.

ARGYROCHÆTA, (*αργυρος*, 'silver,' and *χαιτη*, 'hair,') *Matricaria*.

ARGYROPH'ORA, (*αργυρος*, 'silver,' and *φωρ*, 'I bear,') A name given, by Myrepsus, to an antidote which he regarded as extremely precious.

ARGYROTROPHEMA, (*αργυρος*, 'silver,' and *τροπή*, 'food,') *Blancmanger*.

ARGYRUS, (*αργυρος*,) *Argentum*.

ARHEUMATIC, *Arheumat'icus*, *Arheumat'ic*, *Arheumat'icus*, (*α*, and *ῥευμα*, 'fluxion or rheumatism,') One without fluxion or rheumatism.

ARIA, *Cratægis aria*—a. *Cuttica*, see *Miasma*.

ARIC'INA, *Aric'ina*, *Cus'camin*, *Cusco-Cincho'nia*, (from *Arica* in South America, the place where the bark is shipped.) An alkaloid found in *Cusco Bark*, which is very similar in many of its properties to *Cinchonia*. *Cusco* was the ancient residence of the Incas.

ARIC'YMON, (*αρικ'υμων* from *αρι*, an intensive particle, and *κειν*, 'to conceive,') A name given to a female who conceives readily.

ARIDE'NA, (*aridus*, 'dry,') A Latin word employed to designate the leanness of any part.

ARID'ITY, *Arid'itas*, (F.) *Aridité*, (same etymon.) The French use the word *Aridité* to express the dryness of any organ, and particularly of the skin and tongue, when such dryness is so great as to render the organ rough to the touch.

Aridité also means the lanuginous appearance of the hair in some diseases in which it seems covered with dust.

ARIDU'RA, (same etymon.) Wasting or emaciation of the whole or of any part of the body; *Marasmus*, *Atrophy*.

ARIDURA CORDIS, *Heart*, atrophy of the—*a. Hepatis*, *Hepatrophia*.

ARIKA, see *Spirit*.

ARISÆMA ATRORUBENS, *Arum triphyllum*—a. *Dracontium*, *Arum dracontium*.

ARISTALTHÆA, (*αριστος*, 'best,' and *althæa*,) *Althæa*.

ARISTOLOCH'IA or ARISTOLO'CHIA, (*αριστος*, 'best,' and *δοχεια*, 'parturition,' so called, because the different varieties were supposed to aid parturition.) *Birthwort*, (F.) *Aristolochæ*. *Ord.* *Aristolochiaceæ*. Several varieties were once in use.

ARISTOLOCHIA CAVA, *Fumaria bulbosa*.

ARISTOLOCHIA CLEMATITIS, *Aristolochia Vulgaris* seu *Cretica* seu *ten'uis*, *A'dra R'iza*, *Upright Birthwort*, (F.) *Aristolochæ ordinaire*, *Pommerasse*. The root has been considered stimulant and emmenagogue, and as such has been used in amenorrhœa, chlorosis, and cachexia.

ARISTOLOCHIA CRETICA, *A. Clematitis*—a. *Fabacea*, *Fumaria bulbosa*—a. *Hastata*, see *A. serpentaria*—a. *Hirsuta*, see *A. serpentaria*.

ARISTOLOCHIA LONGA, and *A. ROTUNDA*, (F.) *Aristolochæ longue et ronde*, *Long and Round Birthwort*. Virtues the same as the preceding.

ARISTOLOCHIA PISTOLOCH'IA, *Pistoloch'ia Aristolochia*, *Polyrrh'iza*. This variety has an aromatic odour, and an acrid and bitter taste. (F.) *Aristolochæ crénellée*.

ARISTOLOCHIA RETICULATA, see *A. serpentaria* a. *Rotunda*, see *A. Longa*.

ARISTOLOCHIA SERPENTARIA, *Serpentaria*, *S. Virginia'na*, *Viperaria*, *Viperi'na Virginia'na*, *Colubri'na Virginia'na*, *Contrayer'va Virginia'na*, (F.) *Serpentaire* et *Aristolochæ serpentaire* de *Virginie*, *Coluvrine* ou *Coleuvrée* ou *Vipérine* de *Virginie*, *Virgin'ia Snake'root*, *Snake'root Birth'wort*, *Snake'weed*, *Snag'el*. Virtues of the root—*Serpentaria*—tonic, stimulant; and as such, employed in debility, intermittents, &c. *ARISTOLOCH'IA HIRSU'TA*, *A. HASTATA* and *A. RETICULATA*; indigenous, are employed indiscriminately with *A. Serpentaria* under the name *Serpentaria* or *Virginia Snake'root*. The virtues appear to reside in a bitter extractiform matter, called *Serpentinin*.

ARISTOLOCHIA SI'PHO, *Dutchman's pipe*, and *A. TOMENTO'SA*, indigenous, have virtues like those of *A. Serpentaria*.

ARISTOLOCHIA TENUIS, *A. Clematitis*—a. *Tomentosa*, see *A. Siphon*—a. *Trifida*, *A. Trilobata*.

ARISTOLOCHIA TRILOBA'TA, *A. trif'ida*, (F.) *Aristolochæ trilobée*. A plant of Surinam and Jamaica; possessing the general virtues of the *Aristolochiaceæ*. The other varieties of *Aristolochia* have similar properties.

ARISTOLOCHIA VULGARIS ROTUNDA, *Fumaria bulbosa*.

ARISTOLOCH'IC, *Aristoloch'icus*. Same etymon. An old term for remedies supposed to have the property of promoting the flow of the lochia.

ARIS'TON MAG'NUM, and ARIS'TON PAR'VUM, (*αριστος*, 'excellent,') These names were formerly given to pharmaceutical preparations, used in phthisis, tormina, and fever.

ARISTOPHANE'ION, (*αριστοφανειον*,) A sort of emollient plaster, prepared with four pounds of pitch, two of apoc'hyma, one of wax, an ounce of opoponax, and half a pint of vinegar. Not used.

ARKAN'SAS, MINERAL WATERS OF. About five miles from the Washita river, and about a quarter of a degree north of the Louisiana line, there are about 70 of those springs;—*Onachita* or *Washita Springs*. They are thermal, varying

from 138° to 150° Fahrenheit, and are employed in rheumatism, cutaneous affections, &c.

ARKBEIN (Sc.), Pubis os.

ARKEION, (ἀρκιον.) Aretium Lappa.

ARLADA, Realgar.

ARLES, MINERAL WATERS OF. *Bains près Arles* on *Amélie-les-Bains*. Thermal sulphureous springs in the department of Pyrénées Orientales, France. Their temperature is 103° to 145° of Fah., and they contain sulphohydric acid. The season is from the 15th of May to the 15th of October.

ARM, Brachium.

ARMA, Penis—a. Ventrís, Penis.

ARMAMENTARIUM, (*armare*, 'to equip,') Arsenal—a. Chirurgicum, see Arsenal—a. Portabile, see Case.

ARMARIUM, (from *arma*, 'arms,') Arsenal.

ARMATORY UNGUENT, Hoplochrysa.

ARMATURA, Amnios.

ARME, (*apn*, from *apa*, 'I adapt,') Any physiological or mechanical junction or union of parts. A suture, as of the cranium.

ARMENIACA EPIROICA, *Prunus Armeniaca*—a. *Malus*, Apricot, see *Prunus*—a. *Vulgaris*, *Prunus Armeniaca*.

ARMENIAN STONE, Melochites.

ARMENITES, Melochites.

ARMILLE MANUS MEMBRANOSÆ, (*armilla*, 'a bracelet,') Annular ligaments of the carpus.

ARMOISE BLANCHE, *Artemisia rupestris*—a. *Commune*, *Artemisia vulgaris*—a. *Estragon*, *Artemisia dracunculoides*—a. *Ordinaire*, *Artemisia vulgaris*.

ARMONIACUM, Ammoniac, gum.

ARMORACIA, (from *Armorica*, where it was largely cultivated.) In the Pharmacopœia of the United States, the fresh root of *Cochlearia armoracia*.

ARMORACIA RUSTICANA, *Cochlearia armoracia*—a. *Sativa*, *Cochlearia armoracia*.

ARMOUR, Condom.

ARMURE DES JAMBES, see *Cornu ammonis*.

ARMUS, Humerus—a. *Summus*, *Acromion*.

ARN, *Alnus glutinosa*.

ARNALD'IA, *Arnaldie*. A disease, which appears to have prevailed in England at one time, but whose origin and nature are unknown. It was accompanied with loss of hair; and, by some, is supposed to have been Syphilis. "Deinde atterque Rex incidit in ægritudinem, quam *Arnaldiam* vocant, in qua ipsi usque ad mortem laborantes, capillos suos deposuerunt."—Rogerus Hovedenus, in Ricardo I., cited by Du Cange.

ARNICA MONTANA. (Derivation uncertain.) *Arnica*, *Leopard's Bane*, *Doronicum Germanicum* seu *Oppositifolium*, seu *Plantag'inis folio*, *Arnica Plauen'sis*, *Alis'ma*, *Ac'yrus*, *Diuretica*, *Panacea lapso'rum*, *Ptar'mica montana*, *Col'tha* seu *Calendula Alpina*, *Nar'dus Celtica altera*, (F.) *Arnique*, *Bétoine* ou *Tabac des Montagnes*, *Tabac des Vosges*, *Tabac ou Bétoine des Savoyards*, *Doronic d'Allemagne*, *Plantain ou Souci des Alpes*. *Sex. Syst.* Syngenesia Polygamia superflua. *Ord. Compositæ*. The plant and flowers (*Arnica*, Ph. U. S.) and the root—*Arnica*, Ph. B., (1864)—have been considered narcotic, stimulant, emmenagogue, diuretic, &c.; and, as such, have been given in amaurosis, paralysis, all nervous affections, rheumatism, gout, chlorosis, &c. Dose, gr. v to x, in powder. In large doses it is deleterious.

ARNICA MOL'IS, and A. NUDICAUL'IS, are supposed to have medical virtues like the last.

ARNICA PLAUENSIS, *Arnica montana*—a. *Spuria*, *Inula dysenterica*—a. *Suedensis*, *Inula dysenterica*.

ARNIQUE, *Arnica montana*.

ARNOGLOSSUM, (ἀρνοσ, 'a lamb,' and γλῶσσα, 'the tongue,' from the shape of its leaves,) *Plantago*.

ARNOTT'S DILATOR, see *Dilator*, *Arnott's*.

ARNUT, *Bunium bulbocastanum*.

ARO'MA, gen. *Aro'matis*, (*aroma*, 'perfume:' from *ari*, 'intensive,' and *osm* or *odm*, 'odour,' or from *aro*, 'I make fitting or agreeable,') *Ar'tyma*, *Spirit'us Rector*, (F.) *Arôme*. The odorous part of plants. An emanation—frequently imponderable—from bodies, which acts on the organ of smell, and varies with the body exhaling it.

AROMATE, Aromatic.

AROMATIC, *Aromaticus*, (F.) *Aromate*. Any odoriferous substance obtained from the vegetable kingdom, which contains much volatile oil, or a light and expansible resin. Aromatics are used in perfumes, in seasoning, and embalming. In medicine they are employed as stimulants. Ginger, cinnamon, cardamoms, mint, &c., belong to this class.

AROMATIZE, (F.) *Aromatiser*. To add to a mixture or potion some aromatic, to mask its taste or render it more agreeable. Such mixture is then said to be *aromatized*.

AROMATOPO'LA, (*αρωματωποιος*, from *aroma*, and *πωλεω*, 'I sell,') An apothecary or druggist. One who sells spices.

ARON, (*apov*), *Arum*.

AROPH. A barbarous word, which had various significations with the ancients. Paracelsus employed it to designate a lithontriptic remedy. The mandragora, according to some. Also, a mixture of bread, saffron, and wine.

AROPH PARACELSI, *Ferrum ammoniatum*.

ARQUEBUSADE, *EAU D'*, (F.) (*arquebuse*, 'a hand gun,') *A'qua traumatica Thed'ni* seu *Thedia* seu *sclopeta'ria* seu *vulnera'ria* seu *catapulta'rum*, *Mistu'ra vulnera'ria ac'ida*. A sort of vulnerary water, distilled from a farrago of aromatic plants. *Rosemary*, *hiss*; *millefoil*, *thyme*, each *lbss*; *Proof spirit*, 2 gallons—distil a gallon. This is one form.

ARR (Sc.), *Cicatrix*.

ARRACHEMENT (F.) (from *arracher*, 'to tear out,') *Apospas'ma*, *Abruptio*, *Acut'sio*. Act of separating a part of the body by tearing it from the bonds connecting it with others. *Eviscion*. *Laceration*.

Arrachement is applied to certain operations, as to the extraction of a tooth, the extirpation of a *polypus*, &c.

ARRACK, *Arack*. See *Spirit*.

ARRAPHON, (*αρραφον*, from *a*, and *ραφω*, 'a suture,'—'without suture,') A term applied to the cranium when it presents no sutures.

ARRECTIO, (*arri'go*, *arrectum*, 'to set upright,') *Erection*.

ARRECTORES PILORUM, see *Horrida cutis*.

ARRED (Sc.), see *Scar*.

ARREPTIO, *Insanity*.

ARRESTA BOVIS, *Ononis spinosa*.

ARRÊT D'HILDAN, *Remora Hildani*.

ARRÊTE-BŒUF, (*Arresta bovis*), *Ononis spinosa*.

ARRHEUMATIC, *Arheumatic*.

ARRHŒ'A, (*αρρῶα*, from *a*, and *ρῶω*, 'I flow,') The suppression of any flux. *Amenorrhœa*.

ARRHOSTEMA, (*αρρῶστημα*, from *αρρῶστω*, 'to loose strength,') *Disease*.

ARRHOSTIA, (*αρρῶστια*) *Disease*, *Infirmity*.

ARRHYTHMUS, (*αρρhythμος*, from *a*, and *ρhythmos*, 'rhythm,') *Caecorhythmus*.

ARRIBA, *Geoffræa vermicifuga*.

ARRIÈRE-BOUCHE, (*arrière*, (*ar*, and *re-tro*), 'behind,' and *bouche*, 'mouth,') *Pharynx*—a. *Dent*, see *Dentition*—a. *Faux*. *Secundines*.

ARRIÈRE-GORGE (F.), 'back throat.' The

portion of the pharynx situated behind the tonsils, and the velum pendulum, which can be seen by depressing the tongue.

ARRIÈRE-GOUT (F.), 'after taste.' The taste left by certain bodies in the mouth for some time after they have been swallowed, owing perhaps to the papillæ of the mouth having imbibed the savoury substance.

ARRIÈRE-NARINES, Nares, posterior.

ARROCHE, *Atriplex hortensis* — a. *Puant*, *Chenopodium vulvaria*.

ARROSEMENT, (ar, and ros, 'dew,') Aspersión.

ARROW HEAD, *Sagittaria variabilis*.

ARROWLEAF, *Sagittaria variabilis*.

ARROWPOISON. *Poison de fêches*. This differs with different tribes of Indians. By some, the poison capsicum, and infusions of a strong kind of tobacco, and of euphorbiaceæ are mixed together, with the poisonous emmet, and the teeth of the formidable serpent, called, by the Peruvian Indians, *Mimamaru* or *Jer'gon* — *Luche'sis pic'ta* of Tschudi. See *Curare*.

ARROWROOT, *Fœcula Marantæ*, *Amylum marantæ* seu *Americana*. The fecula of the rhizoma of *Maranta Arundinacea*, *Maranta* (Ph. U. S.), *Ord. Marantaceæ*, which, like all feculæ, is emollient and nutritive, when prepared with water, milk, &c.

Dr. Carson has shown, that *Flor'ida arrowroot* is derived from *Zamia integrifolia* or *Z. pu'ila*, *Sugar pine*, *Ord. Cycadaceæ*; *Bermuda arrowroot* being obtained from *Maranta arundinacea*. Florida arrowroot, as well as the farina, is known in the Southern States under the name *Coonti* or *Coontie*. *Talcahua'no arrowroot*, on the same authority, is obtained from *Alstroeme'ria*, *Fam. Amaryllidaceæ*, of tropical America, and extra-tropical Australia.

According to Dr. Ainslie, an excellent kind of arrowroot is prepared in Travancore from the root of *Curcuma angustifolia*, *Ord. Zingiberaceæ*.

Arrowroot mucilage or *pap* is made by rubbing *arrowroot powder* with a little cold water, in a basin, by means of the back of a spoon, until it is completely mixed with the water; then pouring boiling water over it, stirring assiduously until a soft, gelatinous, tenaceous mucilage is formed; and, lastly, boiling for five minutes. A tablespoonful of arrowroot powder is sufficient to make a pint of mucilage. It may be moderately sweetened; and wine or lemon juice may be added.

With milk also it forms a bland and nutritious article of diet.

ARROWROOT, BRAZIL'IAN. The fecula of *Jatropha Manihot*.

ARROWROOT, COMMON, see *Solanum tuberosum*.

ARROWROOT, EAST IND'IAN. The fecula of the tubers of *Curcuma angustifolia*, or narrow-leaved Turmeric.

ARROWROOT, ENGLISH, Arrowroot, common.

ARROWROOT, TAHITI, *Tac'ca starch*, *Otahe'ite Salep*. The fecula of *Tacca Oceanica*.

ARROWROOT, Talcahuana, see *Arrowroot*.

ARROWWEED, *Sagittaria variabilis*.

ARROWWOOD, *Enonymus*, *Viburnum dentatum*.

ARRYTHMIA, Arrhythm.

ARS (gen. *Artis*) **CABALISTICA**, Cabal — a. Chymiatrica, Chymistria — a. Clysmatica nova, Infusion of medicines — a. Coquinaria, Culinary art — a. Cosmetica, Cosmetics — a. Culinaria, Culinary art — a. Empirica, Empiricism — a. Formulas medicas concinnandi, see *Prescription* — a. Hermetica, Chymistry — a. Homœopathica, Homœopathy — a. Hydriatica, Hydrosudotherapeia — a. Infusoria, Infusion of medicines — a. Machaonia, Medicina — a. Macœntica, Obstetrics — a. Majorum, Chymistry — a. Medica, Medicina — a. Obstetricia,

Obstetrics — a. Sanandi, Art, healing — a. Separatoria, Chymistry — a. Signata, Cabal — a. Spagyrica, Chymistry — a. Sphygmica, Sphygmologia — a. Veterinaria, Veterinary Art — a. Zoiatrica, Veterinary Art.

ARSALTOS, Asphaltum.

ARSATUM, Nymphomania.

ARSECOCKLE (Sc.), (synonymous with Teut. aers-bleyn e, *tuberculum in ano*.) Originally, a hot pimple on the hips. Now, in Scotland, a similar pimple on the face or any part of the body.

ARSENAL (F.), *Chirapothe'ca*, *Armamentarium*, *A. Chirur'gicum*, *Armamentum*. A collection of surgical instruments. A work containing a description of surgical instruments. *Armamentarium obstet'ricum* has the same significations in regard to obstetrics.

ARSENIASIS, Arsenicisimus.

ARSEN'iate, *Arsen'ias*, (F.) *Arséniate*, (from *αρσν*, 'vigorous,') A salt formed by a combination of arsenic acid with a salifiable base.

ARSEN'iate of AMMO'NIA, *Arsen'ias Ammo'niæ*, *Ammonium Arsen'icum* seu *Arsenic'icum*, (F.) *Arséniate d'Ammoniaque*. This preparation is highly extolled in cutaneous diseases. A grain of the salt may be dissolved in an ounce of distilled water, and 20 to 25 drops be commenced with as a dose. This constitutes the *Liquor arsenica'tis Biet'tii*.

ARSEN'iate of IRON, *Fer'ri Arsen'ias*, (Ph. B. 1864,) *Fer'rum Arseniatum* seu *Arsenic'icum oxydulat'um*; (F.) *Arséniate de Fer*. This preparation has been applied externally to cancerous ulcers. An ointment may be made of ʒss of the arseniate, ʒij of the phosphate of iron, and ʒvj of spermaceti ointment. The arseniate has also been given internally in cancerous affections, in the dose of one-sixteenth of a grain.

ARSEN'iate of PROTOX'IDE of POTAS'SIUM, *Proto-arsen'iate of Potas'sium*, *Arsen'iate of Potas'sa*, *Arsen'ias Potas'sæ* seu *Ka'li*, (F.) *Arséniate de Potasse*, *Sel arsenical de Macquer*. Properties the same as those of arsenious acid.

ARSEN'iate of QUINIA, *Quiniæ Arsenias*.

ARSEN'iate of SO'DA, *So'dæ arsen'ias*, *Na'trum arsenic'icum*, (F.) *Arséniate de Soude*. Made, according to the codex of Paris, by mixing and heating to redness *arsenious acid* and *nitrate of soda*; dissolving the salt in water; adding *carbonate of soda* until there is an alkaline reaction; filtering, evaporating, and crystallizing. It is the basis of *Pearson's Solution*, *Liquor arsenica'tis Pearso'nii*, which is formed of one grain of the crystallized *arseniate* to one fluidounce of distilled water. Dose, grt. xx. *Liquor so'dæ arseniat'is*, *Solu'tion of Arsen'iate of So'da* (Ph. B. 1864), is four times the strength of this. *Hei'neck's Solu'tion* consists of the arseniate, gr. yj; *peppermint water*, f ʒiiss; *wine of cinnamon*, f ʒss; *laudanum*, f ʒj. Dose, forty to fifty drops.

ARSEN'iate of SOUDE, *Arseniate of Soda*.

AR'SENIC, *Arsenicum*, (Ph. U. S.) (*αρσν* or *αρσν*, 'vigorous,') A solid metal; of a steel-gray colour; granular texture; very brittle; volatilizing before melting; very combustible and acidifiable. It is not dangerous of itself, and only becomes so by virtue of the facility with which it absorbs oxygen.

ARSENIC BLANC, *Arsenicum album*.

ARSENIC, I'ODIDE of, *Arsen'ici Io'didum* (Ph. U. S.) seu *Teriod'idum* seu *Iodure'tum*, *Arsenic'ium Ioda'tum*, *Hydri'odas arsen'ici*, formed by the combination of *arsenic* and *iodine*. This preparation, applied externally, has been highly extolled in various cutaneous affections. An ointment may be made of three grains of iodide to ʒj of lard. It has also been given internally in the dose of a tenth of a grain in similar affections.

ARSENIC, OXIDE of, *Arsenicum album* — a. Ox-

ide of, white, Arsenicum album—a. White, Arsenicum album.

ARSENIC AND MER'CURY, I'ODIDE OF, *Hydragyri et Arsenici Iodidum*, Double I'odide of Mer'cury and Ar'senic, I'odo-ar'senite of Mer'cury. A compound, which has been proposed as more efficacious than either the iodide of arsenic or the iodide of mercury. It is made by triturating 6.08 grains of arsenic; 14.82 grains of mercury; 49 of iodine, with a fluidrachm of alcohol, until the mass has become dry, and from being deep brown has become pale red. Eight ounces of distilled water are poured on, and, after trituration for a few moments, the whole is transferred to a flask; half a drachm of hydriodic acid, prepared by the acidification of two grains of iodine, is added, and the mixture is boiled for a few moments. When the solution is cold, make the mixture up to f $\overline{3}$ viij with distilled water. This is called by Mr. Donovan, the proposer, *Liquor Arsenici et Hydragyri Iodidi*, each drachm of which, by measure, consists of water, $\overline{3}$ i; arsenious acid, gr. 1-8th; peroxide of mercury, gr. 1-4th; iodine, converted into hydriodic acid, gr. 3-4ths. In the Ph. U. S. it is directed to be made of *Arsenici Iodidum* and *Hydragyri Iodidum rubrum*, each gr. xxxv; *Aqua destillata*, Oss; dissolving by rubbing, heating to the boiling point, and filtering.

The dose of *Donovan's Solution*, *Liquor Donovanii*, is from $\overline{M}$ x to f $\overline{3}$ ss, two or three times a day.

It has been used successfully in inveterate cutaneous diseases.

ARSENICAL, *Arsenicalis*, (F.) *Arsénical*. Relating, or appertaining to, or containing, arsenic.

ARSENICAL PASTE, (F.) *Pâte Arsénicale*. This application to cancers is formed of 70 parts of cinnabar, 22 of dragon's blood, and 8 of arsenious acid; made into a paste with saliva, when about to be applied.

ARSENICAL POWDERS OF FRÈRE CÔME, AND OF JUSTAMOND, see Powder, arsenical.

ARSENICAL SOLUTION, see Arseniate of Soda, and Liquor Arsenicalis.

ARSENICI HYDRIODAS, Arsenic, Iodide of—a. Iodidum, Arsenic, Iodide of—a. Ioduretum, Arsenic, Iodide of—a. Teriodidum, Arsenic, Iodide of.

ARSENICIASIS, Arsenicismus.

ARSENICISM'US, *Arsenicis'asis*, *Arsenici'asis*, *Intoxicatio Arsenicalis*. Poisoning by Arsenic.

ARSENICOPHAGE, (arsenic, and φαγω, 'I eat,') see Arsenicum Album.

ARSENICUM, Arsenic.

ARSENICUM ALBUM, *White Ar'senic*, *Oxide of Ar'senic*, *Rats'bane*, *Arsenici oxydum album*, *Calx Arsenici alba*, *Ac'idum Arsenico'sum seu Arsenio'sum* (Ph. U. S.), *Arsen'ious ac'id*, *White oxide of ar'senic*, (F.) *Arsenic blanc*. An acid which is met with in commerce, in compact, white, heavy, fragile masses; of a vitreous aspect, opaque, and covered with a white dust; of an acrid and nauseous taste; without smell when cold; volatilizable by heat, and exhaling the odour of garlic; soluble in water, alcohol, and oil; crystallizable in regular octahedrons. It is this that is meant by the name arsenic, as commonly used.

ARSENICUM ALBUM SUBLIMA'TUM, *Sublimed Oxide of Arsenic*, is the one employed in medicine. It is tonic and escharotic, and is the most virulent of mineral poisons. It is used in intermittents, periodical headaches, neuroses, &c. Dose, gr. one-tenth to one-eighth, in pill. In some countries—as in the mountainous regions of Austria, Styria, and the Tyrol—arsenic is eaten habitually, beginning with small doses and gradually increasing them. It is said to favor nutrition, and to improve the respiration in ascending

heights. Some of the *arsenicephages* can take great quantities with impunity.

ARSENICUM IODATUM, Arsenic, Iodide of—a. Rubrum factitium, Realgar.

ARSENIS, Arsenite—a. Potassæ, Arsenite of protoxide of potassium—a. Potassæ aqueosus, Liquor arsenicalis—a. Potassæ liquidus, Liquor arsenicalis.

AR'SENITE, *Ar'senic*. A salt, formed by a combination of the arsenious acid with a salifiable base.

ARSENITE OF PROTOX'IDE OF POTAS'SIUM, *Proto-ar'senite of Potas'sium*, *Ar'senite of Potas'sæ*, *Ar'senic Potas'sæ*. An uncrystallizable and colourless salt, which forms the basis of the liquor arsenicalis.

ARSENITE OF QUINIA, Quiniæ arsenis.

ARSESMART, BITING, Polygonum hydro-piper.

ART, HEAL'ING. *Ar's Sanan'di*, *Medic'ina*. The appropriate application of the precepts of the best physicians, and of the results of experience to the treatment of disease.

ART, VETERINARY, Veterinary art.

AR'TABE, (*αράβη*.) Name of a Persian measure for dry substances, in use with the ancients, equal at times, to 5 modii; at others, to 3; and at others, again, to 7.

ARTANTHE ELONGATA, (*αρός*, bread, and *ανθή*, 'flower,') see Matico.

AR'TELSHEIM, MIN'ERAL WA'TERS OF. These German waters have been much recommended in hysteria, gout, palsy, &c. Their physical or chemical properties have not been described.

ARTEMIS'IA, *Anacti'rium*. Ord. Compositæ, (after a queen of the name, who first employed it; or from *Αρτεμης*, 'Diana;' because it was formerly used in diseases of women, over whom she presided.) The Gauls called it *Briciumum*.

ARTEMISIA ABROT'ANUM, *Abrot'anum*, *Abrot'onum*, *Abrot'anum Cath'sum seu Mos*, *Abrotham*, *South'ernwood*, *Oldman*, *Slovenwood*, (F.) *Abro-tone*, *Aurone*, *Aurone mâle*, *Aurone des jardins*, *Garderobe*, *Citronelle*. Supposed to be possessed of stimulant properties.

Oil of Southernwood, *O'leum Abrot'ani*, (F.) *Huile d'Aurone*, possesses the aromatic properties of the plant.

ARTEMISIA ABSIN'THIUM, *Absin'thium*, *A. vulgaris*, *Apsin'thium*, *Barypi'eron*, *Com'mun Worm'wood*, *Worm'tit*, (F.) *Absinthe*. Properties:—tonic and anthelminthic. The Oil of Wormwood, *O'leum Absin'thii*, (F.) *Huile d'Absinthe*, contains the aromatic virtues of the plant.

A bitter liqueur, called *Absinthe*, which is much used in France, is made from the leaves infused in brandy.

ARTEMISIA A'FRA, a South African species, is tonic, antispasmodic and anthelminthic; and has been used in debility of the stomach, visceral obstructions, jaundice and hypochondriasis. It is taken in infusion, decoction and tincture. A strong infusion is used by the Cape Colonists as a collyrium in weakness of the eyes; and the pounded leaves and stalks are employed as dis-cutients in oedema and sugillations.

ARTEMISIA ALBA, A. Santonica—a. Balsamita, A. Pontica.

ARTEMISIA BIEN'SIS, *Bi-en'sial Worm'wood*; indigenous.

ARTEMISIA BOTRYS, *Chenopodium ambrosioides*.

ARTEMISIA CAMPESTRIS, *Field Southernwood*, (F.) *Aurone des Champs*. This possesses the same properties as A. Abrot'anum.

ARTEMISIA CANADENSIS, Canada Wormwood—a. Caudata: indigenous.

ARTEMISIA CHENOPODIUM, *Chenopodium botrys*.

ARTEMISIA CHINES' SIS, *A. Indica*, *A. Mox'a*. From this the Chinese form their *poxias*.

ARTEMISIA CON'TRA grows in Persia, Asia Minor, and other parts of the East.

ARTEMISIA DRACUN'CULUS, *Dracuncul'us*, *Tur'agon*, (F.) *Armoise estragon*, *Estragon*. Virtues:—the same as the last. The leaves and the oil are much used to season food, especially in the form of *Estragon vin'equr*.

ARTEMISIA GLACIA'LIS, *Silky Wormwood*;

ARTEMISIA INDICA, *Artemisia Chinensis*, *A. Santonica*;

ARTEMISIA JUDA'ICA, grows in Palestine, Arabia, China, Cochin China, and Northern Africa.

ARTEMISIA LEPTOPHYLLA, *A. Pontica*;

ARTEMISIA MARIT'IMA, *Absinthium Mori'nium* seu *Mari'tinum*, *Sea Worm'wood*, *Mar'itime South-ernwood*;

ARTEMISIA MOXA, *A. Chinensis*;

and

ARTEMISIA PON'TICA, *A. Roma'na* seu *Tenuifolia* seu *Balsamita* seu *Leptophyll'a*, *Absinthium Ponticum* seu *Roma'nium*, *Ro'man Worm'wood*, *Lesser Worm'wood*, possesses like virtues:—as well

as

ARTEMISIA ROMANA, *A. Pontica*;

ARTEMISIA RUBRA, *A. Santonica*; and

ARTEMISIA RUPES'TRIS, *Creeping Worm'wood*, *Gen'ipi al'bum*, (F.) *Armoise blanc*, *Gen'ipi blanc*. This variety has aromatic virtues, and is used in intermittents, and in amenorrhœa.

ARTEMISIA SANTON'ICA, *Santon'icum*, *Can'ni Her'ba*, *Chama'cedrus*, *Chama'cypariss'us*, *Absinthium Santon'icum*, *Xantoli'na*, *Sche'ba* *Ar'abum*, *S'i'na* seu *C'i'na* *Levan'tica*, *Tarta'rian South'ernwood*, (F.) *Barbotine*. From this and other species of absinthium is obtained the so-called *Semen contra vermes* seu *Contra* seu *Zedoi'rix* seu *Santon'ici* seu *Cin'æ*, *Hagiosperm'um*, *Sanctum Semen*, *Wormseed*—*Santonica*, (Ph. U. S.)—which do not consist of seeds, but of broken peduncles. Virtues:—anthelmintic and stimulant. Dose, gr. x. to ʒj in powder. Its active principle is *Santonin*, the dose of which is from 2 to five grains in the day, for a child six or eight years of age.

ARTEMISIA TENUIFOLIA, *A. Pontica*.

ARTEMISIA VULGA' RIS, *Artemisia ru'bra* et *al'ba*, *Cin'gulum Sancti Joan'nis*, *Mother Herba'rum*, *Berenis'cum*, *Bubasteor'dium*, *Canapa'cia*, *Mug'wort*, (F.) *Armoise ordinaire*, *A. Commune*, *Herbe de Saint Jean*, *Ceinture ou Couronne de Saint Jean*. This, as well as some other varieties, possesses the general tonic virtues of the *Artemisia*. *Artemisia vulgaris* has been highly extolled by the Germans in cases of epilepsy. Dose of the powder, in the 24 hours, from ʒss to ʒj.

ARTÈRE, Artery—*a. Brachiale*, *Brachial artery*—*a. Brachio-céphalique*, *Innominate arteria*—*a. Bronchique*, *Bronchial artery*—*a. Cecale*: see *Colic arteries*—*a. Ciliaire*, *Ciliary artery*—*a. Clitorienne*: see *Clitoris*—*a. Collatérale du conde*, *Anastomoticus magnus ramus*—*a. Collatérale externe*, *Arteria profunda humeri*—*a. Collatérale interne*, *Anastomoticus magnus ramus*—*a. Coronaire des lèbres*, *Labial artery*—*a. Coronaire Stomachique*, *Coronary artery*—*a. Crurale*, *Crural artery*—*a. Deuxième des thoraciques*, *Arteria thoracica externa inferior*—*a. Épineuse*, *Meningeal artery*, middle—*a. Femoro-poplitée*, *Ischiatic artery*—*a. Fessière*, *Gluteal artery*—*a. Gastrique droite petite*, *Pyloric artery*—*a. Gutturo-maxillaire*, *Maxillary artery*, internal—*a. Honteuse externe*, *Pubic*, external, artery—*a. Honteuse interne*, *Pubic*, internal, artery—*a. Humérale profonde*, *Arteria profunda humeri*—*a. Hypogastrique*, *Hypogastrio artery*—*a. Riaque primitive*, *Iliac artery*—*a. Innominée*, *Innominate arteria*—*a. Irienne*, *Ciliary artery*—*a. Ischio-pénienne*: see *Pubic*, internal, artery—*a. Médiane antérieure*, *Spinal ar-*

tery, anterior—*a. Médiane postérieure du rachis*, *Spinal artery*, posterior—*a. Méningée moyenne*, *Meningeal artery*, middle—*a. Méningienne postérieure*, *Prevertebral artery*—*a. Mentonnière*, *Mental artery*—*a. Mésocéphalique*, *Basillary artery*—*a. Mésocœlique*: see *Colic artery*—*a. Musculaire du bras*, *Arteria profunda humeri*—*a. Musculaire du bras, grande*: see *Collateral arteries of the arm*—*a. Musculaire grande de la cuisse*, *Arteria profunda femoris*—*a. Opisthogastrique*, *Cœliac artery*—*a. Orbitaire*, *Ophthalmic artery*—*a. de l'Ovaire*, *Spermatic artery*—*a. Pelvi-crurale*, *Iliac artery*—*a. Pelvienne*, *Hypogastric artery*—*a. Première des thoraciques*, *Arteria thoracica externa superior*—*a. Prévertébrale*, *Prevertebral artery*—*a. Radio-carpienne transversale palmaire*, *Radio-carpal artery*—*a. Scrotale*, *Pubic*, external, artery—*a. Sous-clavière*, *Subclavian artery*—*a. Sous-pubienne*, *Pubic*, internal, artery—*a. Sous-pubio fémorale*, *Obturator artery*—*a. Sous-sternal*, *Mammary*, internal—*a. Sphéno-épineuse*, *Meningeal artery*, middle—*a. Stomogastrique*, *Coronary artery*—*a. Sous-carpienne*: see *Sus-carpien*—*a. Sus-maxillaire*, *Alveolar artery*, *Buccal artery*—*a. Sus-métatarsienne*, *Metatarsal artery*—*a. Sus-pubienne*, *Epigastric artery*—*a. Testiculaire*, *Spermatic artery*—*a. Thoracique humérale*, *Acromial artery*—*a. Trachéocœvical*: see *Cervical arteries*—*a. Trochantérienne*, *Circumflex artery of the thigh*—*a. Troisième des thoraciques*, *Acromial artery*—*a. Tympanique*, *Auditory artery*, external—*a. Uvulaire*: see *Ciliary artery*—*a. Vulvaire*, *Pubic*, external, artery.

ARTEREURYSMA, gen. *Artereurysm'atis*, (*arteria*, and *εγρεπα*, 'dilatation,') *Aneurism*.

ARTERIA, (*αρτηρία*,) *Artery*—*Ad Cutem abdominis*, see *Ad Cutem abdominis (arteria)*—*a. Anonyma*, *Innominate artery*—*a. Aspera*, *Trachea*—*a. Cerebralis*, *Carotid*, internal—*a. Cervicalis*, *Basillary artery*—*a. Coronaria dextra*, *Pyloric artery*—*a. Crassa*, *Aorta*—*a. Cruralis*, *Crural artery*—*a. Cubiti externa*, *Radial artery*—*a. Dorsalis metacarpi*, *Metacarpal artery*—*a. Duræ matris media maxima*, *Meningeal artery*, middle—*a. Encephalica*, *Carotid*, internal—*a. Gastrica superior*, *Coronary artery*—*a. Ileo-colica*: see *Colic arteries*—*a. Iliaca interna*, *Hypogastric artery*—*a. Iliaca posterior*, *Hypogastric artery*—*a. Innominate*, *Innominate arteria*—*a. Magna*, *Aorta*—*a. Magna pollicis*, *Princeps pollicis*—*a. Malleolaris externa*: see *Tibial arteries*—*a. Malleolaris interna*: see *Tibial arteries*—*a. Mammaria externa*, *Arthracica externa inferior*—*a. Maxima Aorta*—*a. Media anastomotica*: see *Colic arteries*—*a. Meningæ media*, *Meningeal artery*, middle—*a. Muscularis femoris*, *A. Profunda femoris*—*a. Pharyngea supra*, *Pterygoid artery*—*a. Profunda cerebri*: see *Cerebral arteries*—*a. Pudenda communis*, *Pubic*, internal, artery—*a. Pudica*, *Pubic*, internal, artery—*a. Ramulus ductus pterygoidei*, *Pterygoid artery*—*a. Spheno-spinosa*, *Meningeal artery*, middle—*a. Spinalis*, see *Spinal arteries*, and *Arteria profunda humeri*—*a. Sternalis*, *Mammary*, internal. *Supra-orbitalis*, *Frontal artery*—*a. Sylviana*: see *Cerebral arteries*—*a. Thoracica axillaris vel alaris*, *Scapular artery*, inferior—*a. Thoracica humeralis*, *Acromial artery*—*a. Transversalis colli*: see *Cerebral arteries*—*a. Transversalis humeri*, *Scapular artery*, superior—*a. Ulnaris*, *Cubital artery*—*a. Uterina hypogastrica*, *Uterine artery*—*a. Vasta posterior*, *Profunda femoris (arteria)*.

ARTERIIAC, *Arteri'acus*. A medicine prescribed in diseases of the windpipe. Also, arterial.

ARTERIE ADIPO'SÆ. The arteries which secrete the fat about the kidneys are sometimes so called. They are ramifications of the capsular, diaphragmatic, renal, and spermatic arteries.

ARTERIE APOPLECTICÆ, *Carotids*—*a. Capitaies*,

Carotids—*a. Ciliares*, Ciliary arteries—*a. Corporis callosi cerebri*, Mesolobar arteries—*a. Jugulares*, Carotids—*a. Lethargicæ*, Carotids—*a. Mesolobicæ*, Mesolobar arteries—*a. Preparantes*, Spermatic arteries—*a. Somniferæ*, Carotids—*a. Soporales*, Carotids—*a. Soporariæ*, Carotids—*a. Venosæ*, Pulmonary veins.

ARTE'RIAL, *Arteri'acus, Arterio'sus*. Belonging to arteries.

ARTERIAL BLOOD, (F.) *Sang artériel*. Red blood is so called because contained in the arteries. The pulmonary veins, however, also contain red blood; hence the name *arterial veins*, (F.) *Veines artérielles*, applied to them.

ARTERIAL CONSTITUTION, *Plethora arteriosa*.

ARTERIAL DUCT, Canalis arterio'sus, Ductus arterio'sus seu **Botali** (F.) **Canal artériel, C. Pulmo-aortique**, is the portion of the pulmonary artery which terminates in the aorta in the foetus. When this duct is obliterated after birth, it is called **Arterial Ligament, Ligamentum arterio'sum**. (F.) **Ligament artériel.**

ARTERIAL SYS'TEM includes all the arteries, from their origin in the heart to their termination in the organs. See Vascular System.

ARTERIALIZATION OF THE BLOOD, Hæ-
motosis.

ARTÉRIARCTIE, (αρτηρια, 'artery,' and arcto, 'I straiten.') Contraction of an artery.

ARTERIECTASIS, (*arteria*, and *ectasis*, 'dilatation.') Aneurism.

ARTERIECTOP'IA, (*arteria*, and *εκτοπος*, 'out of place.') Dislocation of an artery.

ARTERIEURYSMA, gen. *Arterieuryam'atis*,
(*arteria*, and *ευρυσμα*, 'dilatation.') Aneurism.

ARTERIODIALYSIS, (*arteria*, and *διαλυσις*, 'dissolution.') See Aneurism.

ARTERIOGRAPHY, *Arteriogra'phia*, (*arteria*, and *γραφη*, 'a description.') A description of the arteries.

ARTERI'OLA, (dim. of Arteria.) A small artery.

ARTERIOLOGY, *Arteriolog'ia*, (*arteria*, and *λογος*, 'a discourse.') A treatise on the arteries.

ARTERIOPHLEBOTOMY, see Bloodletting.

ARTERIOPIITU'ITOUS. An epithet applied to vessels which creep along the pituitous membrane of the nostrils.

ARTERIORRHEXIS, (*arteria*, and ῥήξις, 'rupture.') Rupture of an artery.

ARTERIOS'ITAS, *Arterios'ity*, (from *arteria*.) A condition of the blood in which it preserves in the veins the arterial character. The opposite to Venositas.

ARTERIOSITAS SANGUINIS, Prædominium sanguinis arteriosi.

ARTÉRIOSTEIE. (*arteria*, and *οστέον*, 'a bone.') Ossification of an artery.

ARTERIOSTENOSIS, (*arteria*, στενός, 'contracted,' and *osis*.) Contraction or obliteration of an artery.

ARTERIOTOMY, *Arteriotom'ia*, (*arteria*, and *tomos*, 'I cut.'). This word has been used for the dissection of arteries. Most commonly, however, it means a surgical operation, which consists in opening an artery, to draw blood from it. Arteriotomy is chiefly used in inflammatory affections of the head, when the blood is generally obtained from the temporal artery. See Bloodletting.

ARTERIOTREP'SIS, (*arteria*, and *τρεψις*, 'tor-
sion.') Torsion of an artery with the view of ob-
literating it.

ARTERITIS, *Arterii'tis*, *Inflammat'io Arteriarum*, (F.) *Artérite*, *Inflammation des artères*, (*arteria* and *itis*.) Inflammation of an artery. Inflammation of the inner coat of an artery is termed *Endo-arteri'tis*, or *Endonarteri'tis*; of the outer, *Eso-arteri'tis* or *Exarteri'tis*.

AR'TERY, *Arte'ria*, (F.) *Artère*, (from *apo*, 'air.'

and *ῥησιν*, 'to preserve,' *quasi*, 'receptacle of air,' because the ancients believed that it contained air. They, at first, gave the name *Artery* to the trachea, *ἀρτηρία τραχεια*, because it is filled with air; and afterwards they used the same term *ἀρτηρία* *pneumatika*, ἀρτηρία πνευματικά, for the arteries, properly so called, probably because they commonly found them empty in the dead body. We find, also, *φλεβες* to designate the arteries, called by the Latins *Vense mican'tes pulsat'iles*.) Arteries, with the moderns, are the order of vessels, which arise from the two ventricles of the heart, and have valves only at their origin. They are cylindrical, firm, and elastic canals; of a yellowish-white colour; little dilatable; easily lacerable; and formed, 1. Of an external laminated, or areolar membrane, *Tu'nica externa seu adventitia seu cellularis*, *Vagii cellularis*, of a dense and close character. 2. Of a middle coat, *Tu'nica media seu elastica*, composed of fibres, which do not, however, contract on the application of the galvanic stimulus, formed chiefly of elastic tissue, and also of smooth muscular fibres, and eminently elastic; and, 3. Of an inner coat, *Tu'nica intima seu glabra*, which is thin, diaphanous, reddish, and polished.

The use of the arteries is to carry the blood from the heart to the various parts of the system. It will be obvious, however, that they cannot all convey arterial blood. The pulmonary artery, for example, is destined to convey the *venous* blood to the lungs, there to be converted into *arterial*; whilst the pulmonary veins convey *arterial* blood back to the heart.

TABLE OF THE PRINCIPAL ARTERIES OF THE BODY.

All the arteries take their rise from the Pulmonary Artery, or the Aorta; and the names generally indicate the parts to which they are distributed.

I. ARTERIA PULMONALIS.

The Pulmonary Artery arises from the right ventricle, and soon divides into a right and left branch, one of which is distributed to each lung.

II. ARTERIA AORTA.

The Aorta arises from the left ventricle. It is the common trunk of the arteries of the body, and may be divided into five portions.

a. *Arteries furnished by the Aorta at its origin.*

1. A. Cardiaca or coronaria anterior.
2. A. Cardiaca or coronaria posterior.

b. *Arteries furnished by the Aorta at its arch.*

The arch of the Aorta gives off, to the left, two considerable trunks—the *Arteria carotidea primitiva*, and *A. subclavia*; and, to the right, a single trunk, which is larger—the *A. innominata* or *Brachio-cephalica*, which divides into the *primitivæ carotid* and *subclavian*.

A. ARTERIA CAROTIDEA PRIMITIVA. { Divides into A. Carotidea externa,
and A. Carotidea interna.

Furnishes, 1. *A. Thyroidea superior*.
2. *A. lingualis*, which gives off the *A. dorsalis* linguae and *A. sublingualis*.
3. *A. fucialis* vel *A. maxillaris externa*, which furnishes the *A. palatina inferior*, the *A. submentalis*, and *A. cornaria superior* and inferior.
4. *A. occipitalis*, which gives off the *A. mastoidea posterior*.
5. *A. auricularis posterior*, which gives off *A. stylo-mastoidea*.
6. *A. pharyngea inferior*.

The external carotid ultimately divides into the temporal artery and internal maxillary.

1. A. { Furnishes *A. transversalis faciei*, *A. auricularis anterior*, and *A. temporalis media*.

- Furnishes 12 branches, viz.—*A. meningea media*, *A. dentalis inferior*, *A. temporalis profunda posterior*, *A. masseterina*, *A. pterygoidea*, *A. buccalis*, *A. temporalis profunda anterior*, *A. alveolaris*, *A. suborbitalis*, *A. Vidiani*, *A. pterygopalatina* or *pharyngea superior*, *A. palatina superior*, and *A. sphenopalatina*.
- Furnishes, 1. *A. ophthalmica*, which gives off *A. lacrymalis*, *A. centralis retinae*, *A. supraorbitalis vel superciliaris*, *A. ciliares posteriores*, *A. ciliares longae*, *A. muscularis superior et inferior*, *A. ethmoidalis posterior et anterior*, *A. palpebralis superior et inferior*, *A. nasalis*, and *A. frontalis*. 2. *A. communicans Willisii*. 3. *A. choroidea*. 4. *A. cerebialis anterior*. 5. *A. cerebialis media*.
- Furnishes, 1. *A. vertebralis*, which gives off *A. spinalis anterior et posterior*, *A. cerebello inferior*, and forms—by uniting itself with that of the opposite side—the *A. basilaris*, divided into *A. cerebello superior* and *A. cerebialis posterior*. 2. *A. thyroidea inferior*, which gives off *A. cervicalis ascendens*. 3. *A. mammaria interna*, which gives off the *A. mediastina anterior* and *A. diaphragmatica superior*. 4. *A. intercostalis superior*. 5. *A. cervicalis transversa*. 6. *A. scapularis superior*. 7. *A. cervicalis posterior vel profunda*. Farther on, the subclavian artery continues its progress under the name *A. axillaris*.
- Furnishes, 1. *A. acromialis*. 2. *A. thoracica superior*. 3. *A. thoracica inferior vel longa vel mammaria externa*. 4. *A. scapularis inferior vel communis*. 5. *A. circumflexa posterior*. 6. *A. circumflexa anterior*. Farther on, the axillary artery continues under the name *A. brachialis*.
- Furnishes *A. humeralis profunda vel collateralis externa*. 2. *A. collateralis interna*. It afterwards divides into the *radial* and *cubital* arteries.
- Gives off *A. recurrens radialis*, *A. dorsalis carpi*, *A. dorsalis metacarpi*, *A. dorsalis pollicis*, and terminates in forming the *Arcus palmaris profundus*.
- Gives off *A. recurrens cubitalis anterior* and *posterior*; *A. interossea anterior* and *posterior*, which latter furnishes *A. recurrens radialis posterior*. It terminates in forming the *superficial palmar arcu*, which gives off *A. collaterales digitorum*.
- c. Arteries given off by the Aorta in the Thorax.**
- These arteries are, { 1. *A. bronchica, dextra et sinistra*.
2. *A. oesophagica* (to the number of four, five, or six).
3. *A. mediastina posteriores*.
4. *A. intercostales inferiores vel aorticae* (to the number of eight, nine, or ten).
- d. Arteries furnished by the Aorta in the Abdomen.**
- These branches are, { 1. The *A. diaphragmatica vel phrenica, dextra et sinistra*.
Which divides into three branches, 1. *A. coronaria ventriculi*. 2. *A. hepatica*, which gives off *A. pylorica*, *A. gastro-epiploica dextra*, and *A. cystica*; and, lastly, the *A. splenica*, which gives off *A. gastro-epiploica sinistra* and *Vasa brevia*.
Which gives off at its concavity the *A. colica dextra superior, media et inferior*, and at its convex part from 15 to 20 *Rami intestinales*.
Which gives off *A. colica superior media, et inferior*, and divides into *A. hemorrhoidales superiores*.
5. The *A. Capsulares mediae* (to the number of two on each side).
6. *A. Renales vel Emulgentes*.
7. *A. Spermatice*.
8. *A. Lumbares* (to the number of four and five on each side).
- e. Arteries resulting from the bifurcation of the Aorta.**
- The Aorta, a little above its bifurcation, gives off the *A. sacra media*, and divides into *A. iliaca primitiva*.
A. iliaca { Divides into *iliaca interna* and *A. iliaca externa*.
- Furnishes, 1. *A. ilio-lumbaris*. 2. *A. sacra lateralis*. 3. *A. glutea vel iliaca posterior*. 4. *A. umbilicalis*. 5. *A. vesicalis*. 6. *A. obturatoria*. 7. *A. hamorrhoida media*. 8. *A. uterina*. 9. *A. vaginalis*. 10. *A. ischiatica*. 11. *A. pudenda interna*, which gives off the *A. hamorrhoidales inferiores*, *A. of the septum*, *A. transversa perinei*, *A. corporis cavernosi*, and *A. dorsalis penis*.
- Furnishes, 1. *A. epigastrica*, 2. *A. iliaca anterior vel circumflexa ili*, and is continued afterwards under the name of *Crural Artery*.
- Furnishes, 1. *A. subcutanea abdominalis*. 2. *A. pudenda superficialis et profunda*. 3. *A. muscularis superficialis*. 4. *A. muscularis profunda*, which gives off the *A. circumflexa externa et interna*, and the three *Perforantes*, distinguished into superior, middle, and inferior. Farther on, the crural artery continues under the name *A. Poplitea*.
- Furnishes, 1. *A. Articulares superiores, interna, media, et externa*. 2. *A. Gemelle*. 3. *A. Articulares inferiores, (interna et externa.)* 4. *A. tibialis antica*, which, at the foot, takes the name *A. dorsalis tarsi*, and gives off the *tarsal* and *metatarsal* arteries. In the leg, the popliteal artery divides into the *peroneal* and *posterior tibial*.
- Divides into *A. peronea antica* and *A. peronea postica*.
- Divides into *A. plantaris interna* and *A. plantaris externa*. The latter by anastomosing with the *A. dorsalis tarsi*, forms the *plantar arcu*, whence arise *Rami superiores vel perforantes postici*, *R. inferiores postici et antici*, which give off *Rami perforantes antici*.
- ARTERY, ANGULAR, Facial artery**—*a. Articular, Circumflex artery*—*a. Brachiocephalic, Innominata arteria*—*a. Central of the retina, Central artery of the retina*—*a. Central of Zinn, Central artery of the retina*—*a. Cephalic, Carotid*—*a. Cerebral posterior, Vertebral*—*a. Cervico-scapular, see Cervical arteries*—*a. Coronary of the lips, Labial artery*—*a. Crotaphite, Temporal artery*—*a. Fibular, Peroneal artery*—*a. Gastric inferior, Gastro-epiploic artery*—*a. Gastro-hepatic, see Gastro-epiploic artery*—*a. Genital, Pudic (internal) artery*—*a. Guttural inferior, Thyroideal A. inferior*—*a. Guttural superior, Thyroideal A. superior*—*a. Humeral, Brachial artery*—*a. Iliac posterior, Gluteal artery*—*a. Iliaco-muscular, Ileo-lumbar artery*—*a. Labial, Facial artery*—*a. Laryngeal superior, Thyroideal artery, superior*—*a. Maxillary internal, Facial artery*—*a. Median of the sacrum, Sacral artery, anterior*—*a. Nasal, lateral, large, Spheno-palatine artery*—*a. of Neubauer, see Thyroideal arteries*—*a. Palato-labial, Facial artery*—*a. Periephalic, Carotid (external)*—*a. Pharyngeal, superior, Pterygo-palatine artery*—*a. Phrenic, Diaphragmatic artery*—*a. Posterior, of the brain, see Cerebral arteries*—*a. Scapular external, Acromial artery*—*a. Spinal, Meningeal artery, middle*—*a. Subclavian right, Innominata arteria*—*a. Subscapular, Scapular artery, inferior*—*a. Superficial of the abdomen, Ad cutem abdominis (arteria)*—*a. Supramaxillary, Alveolar artery*—*a. Suprarenal, Capsular artery*—*a. Thoracic, internal, Mammary internal*—*a. Urethro-bulbar, Transverse perineal artery*—*a. Vesico-prostatic, Vesical artery*—*a. Vidian, Pterygoid artery*.
- ARTETIS'CUS, (artus, 'a limb.')** One who has lost a limb.
- ARTHANI'TA, (apros, 'bread;')** the *Cyc'lamen* or *Sonbread*. It was formerly made into ointment, *Unguentum Arthanitæ*, with many other substances, and was employed as a purgative, being rubbed on the abdomen.
- ARTHANITA CYCLAMEN, Cyclamen.**
- ARTHETICA, Teucurium chamaepitys.**
- ARTHRAGRA, (arthro, and agra, 'seizure.')** Gout—*a. Anomala, Gout, anomalous*—*a. Genuina,*

Gout, regular.—a. Legitima, Gout, regular.—a. Normalis. Gout, regular.—a. Vera, Gout, regular.

ARTHRALGIA, (*arthro*, and *algos*, 'pain.') Arthralgia. Gout. See Lead rheumatism.

ARTHRELCO'SIS, (*arthro*, and *ἄλσσις*, 'ulceration.') Ulceration of a joint.

ARTHREMBOLE'SIS, (same etymon as the next.) The reduction of a fracture or luxation.

ARTHREMBOLUS, (*ἀρθρεμβολος*, from *arthro*, *αἰ*, 'in,' and *βάλλω*, 'I cast.') An ancient instrument used in the reduction of dislocations.

ARTHRETICA, Teuericun chamæpitys.

ARTHRIC, *Ar'hricus*, (*ἀρθρικός*,) (F.) *Arthrique*, (from *arthro*). Appertaining or relating to joints.

ARTHRIT'IC, *Arthrit'icus*, (*ἀρθριτικός*, from *arthro*,) *Arthrel'icus*, *Arthrel'icus*, *Gouty*, (F.) *Arthritique*, *Goutteux*. That which relates to arthritis or inflammation of the joints in general, and especially to gout, as *arthritic symptoms*, &c. See Podagrie.

ARTHRITICUS VERUS, Gout.

ARTHRITIFUGUM, (*arthriticus*, and *fugare*, 'to drive away.') A remedy that drives away gout. Heyden terms cold water, internally, the *arthritifugum magnum*.

ARTHRITIS, gen. *Arthrit'idis*, (*ἀρθριτις*, from *arthro*, and *itis*), Gout, Arthrophlogosis, Arthrosia—a. Aberrans, Gout (wandering)—a. Acuta, Gout (regular)—a. Arthrodynia, Rheumatism, chronic—a. Asthenica, Gout (atonic)—a. Atonic, Gout (atonic)—a. Diaphragmatica, Angina Pectoris—a. Erratica, Gout (wandering)—a. Hydrarthros, Hydrarthrus—a. Inflammatoria, Gout (regular)—a. Juvenilis, see Rheumatism, acute—a. Maxillaris, Siagonagra—a. Nodosa, Gout (with nodosities)—a. Planetica, Gout (wandering)—a. Podagra, Gout—a. Retrograda, Gout (retrograde)—a. Rheumatica, see Rheumatism, acute—a. Rheumatismus, Rheumatism, acute—a. Vaga, Gout (wandering).

ARTHRO, from *ἀρθρον*, 'a joint,' in composition, 'joint.'

ARTHROC'ACĒ, (*arthro*, and *κακός*, 'bad.') Disease of the joints; and especially caries of the articular surfaces. *Spina ventosa*.

ARTHROCACE COXARTUM, Coxarum morbus.

ARTHROCACOLOG'IA, (*arthrocacia*—according to Rust, a chronic disease of the joints; and *logos*, 'a description.') A treatise on chronic diseases of the joints.

ARTHROCARCINO'MA, gen. *Arthrocarcino'matis*, (*arthro*, and *καρκίνωμα*, 'cancer.') Cancer of the joints.

ARTHROCHONDRI'TIS, (*arthro*, *χόνδρος*, 'a cartilage,' and *itis*.) Inflammation of the cartilages and joints.

ARTHREDEMA, Hydrarthrus.

ARTHRO'DIA, (*ἀρθρώδια*, from *arthro*.) *Adarticula'tio*, (F.) *Enfonçure*, *Enfonçure*. A movable joint, formed by the head of a bone applied to the surface of a shallow socket, so that it can execute movements in every direction. *Arthro'dium* is also, 'a small joint;' diminutive of *Arthrodia*.

ARTHRODIUM, see Arthrodia.

ARTHRODYN'IA, *Arthronal'gia*, *Arthral'gia*, (*arthro*, and *δύνη*, 'pain.') Articular pain. Pain in the joints. See Rheumatism, chronic.

ARTHRODYNIA PODAGRICA, Gout.

ARTHROG'RAPHY, *Arthrog'raph'ia*, (*arthro*, and *γραφω*, 'I write.') A description of the joints.

ARTHROGYPO'SIS, (*arthro*, and *γυρω*, 'I incurvate.') Malposition of the joints, especially of the hands and feet.

ARTHROHYDRINE, (*arthro*, and *ῥωψ*, 'water.') Synovia.

ARTHROLOGY, *Arthrol'ogy*, (*arthro*, and

λόγος, 'a description.') A description of the joints. The anatomy of the joints.

ARTHROMBOLE, (*arthro*, and *βάλλω*, 'I cast.') Coaptation, reduction. Reduction of a luxated or fractured bone.

ARTHROMENINGITIS, (*arthro*, *μηνιγξ*, gen. *μηνιγγος*, 'membrane,' and *itis*.) Meningarthrocace.

ARTHRON, (*ἀρθρον*, 'a joint.') The ancients used the word *Arthron* for the articulation of bones with motion, in opposition to *Symphysis* or articulation without motion.

ARTHRONALGIA, (*arthro*, and *algos*, 'pain,') Arthrodynia.

ARTHRON'CUS, *Arthrophy'ma*, (*arthro*, and *ογκος*, 'a swelling.') Tumefaction of a joint.

ARTHRONEMPYESIS, (*arthro*, *αἰ*, 'in,' and *πυον*, 'pus.') Arthropyosis.

ARTHROP'ATHY, *Arthrophath'ia*, (*arthro*, and *πάθος*, 'affection.') A disease of the joints.

ARTHROPHLOGO'SIS, (*arthro*, and *φλεγω*, 'I burn;') *Arthrit'is*, *Ostearthro'sis*. Inflammation of the joints.

ARTHROPHLOGOSIS SYNOVIALIS, Synovitis.

ARTHROPHYMA ADENOCHONDRUM, gen. *Arthrophy'matis*, (*arthro*, and *φυμα*, 'a swelling.') See Adenochondrius.

ARTHROPLAS'TIC, *Arthroplas'ticus*, (*arthro*, and *πλασσειν*, 'to form.') An epithet for an operation to form a false joint in cases of ankylosis.

ARTHROPYO'SIS, *Arthronempy'e'sis*, (*arthro*, *πυον*, 'pus,' and *osis*.) Suppuration or abscess of the joints.

ARTHRORHEUMATISMUS, Rheumatism (acute).

ARTHRO'SIA, (*ἀρθρῶσα*, 'I articulate,') *Arthritis*, (of some). Inflammation, mostly confined to the joints; severely painful; occasionally extending to the surrounding muscles. A genus of diseases in the Nosology of Good, including *Rheumatism*, *Gout*, *Articular inflammation*, *Joint-ache*, &c.

ARTHROSIA ACUTA, Rheumatism, acute—a. Chronica, Rheumatism, chronic—a. Lumborum, Lumbago—a. Podagra, Gout—a. Podagra complicata, Gout (retrograde)—a. Podagra larvata, Gout (atonic)—a. Podagra regularis, Gout (regular).

ARTHROSIS, Articulation.

ARTHROSPON'GUS, (*arthro*, and *σπγγος*, 'a sponge.') A white, fungous tumour of the joints.

ARTHROTAPHUS, see Tophus.

ARTHROTRAUM'A, gen. *Arthrotrauma'tis*, (*arthro*, and *τραυμα*, 'a wound.') A wound of a joint.

ART'IA, (*αἰρ*, 'air.') According to some, this word is synonymous with *αἰτηρία*; others use it synonymously with *Trachea*.

ARTICHAUT, *Cynara scolymus*.

ARTICHOKE, *Cynara scolymus*.

ARTICLE, Articulation—a. of Death, see Psychorages.

ARTICOALUS, *Cynara scolymus*.

ARTICULAR, *Articula'ris*, (F.) *Articulaire*, (*artus*, 'a joint;' *articulus*, 'a small joint.') That which relates to the articulations—as the *articular capsules*, &c.

ARTICULAR ARTERIES OF THE ARM, Circumflex arteries of the arm.

ARTICULAR ARTERIES OF THE KNEE arise from the popliteal artery, and surround the tibio-femoral articulation. Although of a small size, they are important, as they furnish blood to the lower extremity after the operation for popliteal aneurism. They are distinguished into *superior* and *inferior*. The *superior articular arteries*, *popliteal articular arteries*, are commonly three in number; one of which is *internal*, another *external*, and another *middle*, the *azygous articular*. The first, *Ramus anastomot'icus magnus*, anastomoses by

one branch with the external circumflex, and by another with the external superior articular. The second anastomoses with the external circumflex, the superior internal articular, and the inferior external articular; and the third is distributed within the joint. The inferior articular arteries are two in number: an internal and an external. The former anastomoses with the internal superior articular and the external inferior articular. The latter anastomoses with the recurrent branch of the anterior tibial, and the external superior articular. To each articular artery there is an articular nerve.

ARTICULAR FACETTES' are the contiguous surfaces, by means of which the bones are articulated.

ARTICULAR PROCESSES, see Vertebrae.

ARTICULAR VEINS of the knee follow the same course as the arteries.

ARTICULATED, *Articulatus*, (F.) *Articulé*, same etymon as Articulation. Provided with, or united by, articulations; as an 'articulated skeleton.' See Skeleton, articulated; and Voice, articulated.

ARTICULATION, Articulation—a. Artificialis, Pseudarthrosis—a. Coxæ, Coxo-femoral articulation—a. Genu, Genu—a. Notha, Pseudarthrosis.

ARTICULATION, *Articulatio*, *Arthro'sis*, *Ar'tis*, dim. *Artic'ulus*, *Junctu'ra*, *Conjunctio*, *Nodus*, *Commissu'ra*, *Compag'es*, *Syntax'is*, *Har'mus*, *Vertic'ula*, *Vertic'ulus*, *Vertic'ulum*, *Joint*, (F.) *Articulation*, *Article*, *Jointure*, (same etymon as Articular.) By some of the French surgeons and anatomists, *article* is restricted more particularly to a movable articulation. The union of bones with each other, as well as the kind of union. Articulations are generally divided into *Diarthro'ses* or movable articulations, and *Synarthro'ses* or immovable.

The articulations are subject to a number of diseases, which are generally somewhat severe. These may be physical, as wounds, sprains, luxations, &c.; or they may be organic, as ankylosis, extraneous bodies, caries, rheumatism, gout, hyarthro'ses, arthroprosis, &c.

ARTICULATION means also the combination of letters which constitute words. See Voice.

ARTICULATION EN CHARNIÈRE, Ginglymus.

ARTICULATION, FALSE, *Pseudarthro'sis*, *Artic'ulus falsus*, (F.) *A. fausse*, *A. accidentelle*, *A. contre nature*, *A. anormale*. A false joint, formed between fragments of bone, that have remained ununited; or between a luxated bone and the surrounding parts.

ARTICULATION DE LA HANCHE, Coxo-femoral articulation.

ARTICULATION SURNUMÉRAIRE, Pseudarthrosis.

ARTICULI DIGITORUM MANÛS, Phalanges of the fingers—a. *Digitum pedis*, Phalanges of the toes.

ARTICULO MORTIS, ('in the article of death,') see Psychorages—a. *Spinalis*, *Semispinalis colli*.

ARTICULUS NOVUS, ('new joint,') Pseudarthrosis—a. *Præternaturalis*, Pseudarthrosis—a. *Spurius*, Pseudarthrosis.

ARTIFICIAL, *Artificialis*, (F.) *Artificiel*, (*ars*, *artis*, 'art,' and *facere*, 'to make,') That which is formed by art.

ARTIFICIAL EYES are usually made of enamel, and represent a sort of hollow hemisphere, which is applied beneath the eyelids, when the eye is lost.

ARTIFICIAL TEETH are made of ivory, the tooth of the hippopotamus, (F.) *Osanore*; porcelain, &c. Porcelain teeth are likewise called *mineral*, *incorruptible*, and *vitrecent* teeth.

Pièces d'Anatomie Artificielles, are prepara-

tions of anatomy, modelled in wax, plaster, paper, &c.

ARTISCOCCUS LÆVIS, *Cynara scolymus*.

ARTIS'CUS, (*aproskos*, dim. of *apros*, 'bread,') See Trochiscus. A troch of the shape of a small loaf. Also, and especially, a troch made of vipers.

ARTOCARP'US. The Bread-fruit Tree, (F.) *Jaquier*, (*apros*, 'bread, and *karpos*, 'fruit,') A Polynesian tree, so called because the fruit, which is milky, and juicy, supplies the place of bread to the inhabitants. It grows to the height of 40 feet.

ARTOCARPUS INTEGRIFOLIA, *Caoutchouc*.

ARTOC'REAS, (*aproskreas*, from *apros*, 'bread,' and *kreas*, 'flesh,') A kind of nourishing food made of various aliments boiled together.

ARTOG'ALA, (*apros*, 'bread,' and *gala*, 'milk,') An alimentary preparation of bread and milk. A poultice.

ARTOM'ELI, (*apros*, 'bread, and *melis*, 'honey,') A cataplasm of bread and honey.

ARTUS, Articulation, Membrum.

ARTYMA, gen. *Artym'atis*, (*artyma*, from *artym*, 'to prepare,') Aroma, Condiment.

ARUM, (*apor*,) A. maculatum, and A. triphyllum—a. Americanum *betæ foliis*, *Dracontium foetidum*.

ARUM DRACON'TIUM, *Arisæ'ma dracon'tium*, *Green Dragon*; indigenous, has probably the same medical properties as A. triphyllum.

ARUM DRACUN'CULUS, A. *polyphyllum*, *Dracunculus polyphyllus*, *Colubrina Dracontia*, *Erva de Sancta Maria*, *Gig'arum serpentaria*, *Serpentaria Gallo'rum*. Family, *Araceæ*. *Sez. Syst.* *Monœcia Polyandria*. The roots and leaves are very acrimonious. The plant resembles the A. maculatum in its properties.

ARUM ESCULENTUM, *Calladium esculentum*, *Ta'ro*, *Ka'lo*. The foliage and roots possess acid qualities, which are dissipated by baking or boiling; in which form it is used as food by the people of Madeira, the Polynesians, &c. The tuberous root of this Arum, called *Co'cos* or *Ed'does*, is said to be the principal dependence for food of the laboring population of the West Indies.

ARUM MACULA'TUM, *A'ron*, *Arum* (of the older writers), *A. vulgare*, *Bar'ba Aaro'nis*, *Serpentaria m'nor*, *Zin'giber German'icum*, *Sacerdotia pe'nis*, *Wake Robin*, *Priest's pintle*, *Cuckow Pint*, (Prov.) *Cockybaby*, *Aaron*, (F.) *Gouet*, *Pied de Veau*, *Pied de lièvre*, *Picotin*, *Giron*, *Amidonnière*. The fresh root is stimulant internally, Dose, ℞j. of the dried root. Externally, it is very acrid. From the root of this Arum a starch is prepared, which is called *Port'land Island Sa'go*, *Gersa serpentaria*, *Cerus'sa serpentaria*, *Fec'ula a'ri maculati*.

ARUM, THREE-LEAVED, *Arum triphyllum*.

ARUM TRIPHYLLUM, *Arisæ'ma atrorubens*, *Three-leaved arum*, (F.) *Pied de Veau triphyllé*, *Ind'ian Tur'nip*, *Drag'on Root*, *Drag'on Tur'nip*, *Pep'per Tur'nip*. This plant grows all over the United States, and is received into the Pharmacopœia under the title *Arum*. The recent root, or Cormus—ARUM, (Ph. U. S.)—is very acrimonious, and has been employed in asthma, croup, and hooping-cough. Boiled in lard, it has been used in tinea capitis, and in milk in consumption.

ARUM VIRGINICUM, *Peltandra Virginica*—a. *Vulgare*, A. maculatum—a. Water, *Calla palustris*.

ARUMARI, Caramata.

ARUNDO BAMBOS, Bamboo—a. *Brachii major*, *Ulna*—a. *Brachii minor*, *Radius*—a. *Indica*, *Sagittarium alexipharmacum*—a. *Major*, *Tibia*—a. *Minor*, *Fibula*—a. *Saccharifera*, see *Saccharum*.

ARUNGZEBE, Delhi boil.

ARVA, *Ava*.

ARVUM, ('a field.') Vulva—a. Naturæ, Uterus.

ARY-ARYTÆNOIDÆUS, Arytænoidæus—a. Epiglotticus, Arytæno-epiglotticus.

ARYTÆ'NA, (ἀρταινα, 'a ladle.') Hence,

ARYTÆ'NOËPIGLOT'TICUS, Arytæ'no-epiglottidæus, Ary-epiglott'icus. That which belongs to the arytenoid cartilages and epiglottis. Winslow gives this name to small, fleshy fasciculi, which are attached, at one extremity, to the arytenoid cartilages, and, by the other, to the free edge of the epiglottis. These fibres do not always exist. They form part of the arytenoid muscle of modern anatomists.

ARYTÆNOIDÆ'US, (F.) Arytænoidien. A small muscle, which passes from one arytenoid cartilage to the other, by its contraction brings them together, and diminishes the aperture of the glottis. Winslow divided the muscle into three portions;—the Arytænoidæus transversus, or Ary-arytænoidæus, and two Arytænoidæi obli'qui.

ARYTÆNOID, Arytænoid'es, Arytænoidæus, (arytæna, and eidos, 'shape.') Ladle-shaped.

ARYTÆNOID CAR'TILAGES, Cartilag'ines arytænoid'es seu guttura'les seu gutturi'næ seu gutturi'for'mes seu triq'uetræ, Guttur'nia, are two cartilages of the larynx, situate posteriorly above the cricoid, which, by approximation, diminish the aperture of the glottis. Their upper extremities or cornua are turned towards each other, and are now and then found loose in the form of appendices, which are considered by some as distinct cartilages, and termed cu'neiform, or tuber'culated Car'tilages, Cornic'ula laryn'gis.

ARYTÆNOID GLANDS, Gland'ulæ Arytænoidæ'æ, are small, glandular, whitish bodies, situate anterior to the A. cartilages. They pour out a mucous fluid to lubricate the larynx.

ARYTHM, Aryth'mia, Arryth'mia, (ἀρρυθμία, from α, and ρυθμος, 'rhythm,' 'measure.') Irregularity. This word is applied chiefly to the pulse.

AS, in composition, see Ad.

AS, gen. Assis, Pound.

ASA, Asafetida: see Assa—a. Dulcis, Benjamin.

ASAFET'IDA, Asasafetida (Ph. U. S.), Asasafetida, Ster'eus diabol'i, Cibus Deo'rum, A'sa. Devil's dung. Food of the Gods. A gum-resin, the concrete juice of Fer'ula Asasafetida, Nar'theæ Asasafetida. Order, Umbellifera. It is in small masses of a whitish, reddish, and violet hue, adnating together. Taste bitter and subacid: smell insupportably alliaceous. The Asiatics use it regularly as a condiment.

Its medical properties are antispasmodic, stimulant, and anthelmintic. Dose, gr. v to xx, in pill.

ASAGRAYA or ASAGRÆA OFFICINALIS (after Dr. Asa Gray), see Veratrina.

AS'APES, (ασαπης,) Asep'ton, properly, not putrid, (α, and σαπιο or σπηω, 'I corrupt.') A term applied by Hippocrates to the sputa, or to other matters evacuated, which do not give signs of coction.

ASAPHA'TUM, Sapha'tum, (α, and σαφης, 'clear.') This term has been applied to collections in the sebaceous follicles of the skin, which may be pressed out like little worms, with a black head. See Acne.

ASAPH'IA, (ασαφεια, same etymon.) Dyspho'nia inmodula'ta palati'na, Parapho'nia guttura'lis seu palati'na. Defective articulation, dependent upon diseased palate.

ASARABACCA, Asarum—a. Broad-leaved, Asarum Canadense.

ASARCIA, (α, and σαξ, gen. σαρκος, 'flesh.') Emaciation.

ASAR'CON, (ασαρκον.) Devoid of flesh. Aristotle uses the term for the head when it is but

little fleshy, compared with the chest and abdomen.

ASARET, Asarum—a. du Canada, Asarum Canadense.

ASARIT'ES, (ασαριτης, from asarum.) A diuretic wine, of which asarum was an ingredient.

AS'ARUM, (ασαρον, from α, and σαρω, 'to adorn,' because not admitted into the ancient coronal wreaths [?],) As'arum Europæum seu officina'le, Nar'dus Monta'na seu Rust'ica, Az'arum, (F.) Asaret ou Cabaret, Azarum Cabaret, Oreille d'homme, Oreillette, Girard-Roussin, Nard Sauvage. Order, Aristolochiaceæ, Sex. Syst. Dodecandria Monogynia. The plant, used in medicine, is the As'arum Europæum, Asarabacca, and of this the leaves. They are emetic, cathartic, and errhine, but are hardly ever employed, except for the last purpose.

ASARUM CANADEN'SË, A. Carolina'num, Can'ada Snake'root, Wild Gin'ger, Colt's Foot, Broad-leaf As'arabacca, In'dian Gin'ger, Heart Snake'root, (F.) Asaret du Canada. The root—As'arum, (Ph. U. S.)—is used as a substitute for ginger, and is said to act as a warm stimulant and diaphoretic.

ASARUM CAROLINIANUM, A. Canadense—a. Europæum, see Asarum—a. Hypocistis, Cytinus hypocistis—a. Officinale, see Asarum.

ASBESTOS SCALL, (ασβεστος, 'inveterate,') see Eczema of the hairy scalp.

ASBOL'ICUS, (ασβολη, 'soot,') Relating or appertaining to soot:—Carcino'ma acro'ti asbol'icum, Chimney-sweepers' Cancer.

ASCAIN, MIN'ERAL WATERS OF. Ascain is a village, situate about a league from St. Jean-de-Luz, in France. The water is a cold chalybeate.

ASCARDAMYC'TUS, (ασκαρδαμυκτος, from α, and σκαρδαμυκτω, 'I twinkle the eyes,') One who stares with fixed eyes, without moving the eyelids.

ASCARICIDA ANTHELMINTICA, (ascarica, and cædere, 'to kill,') Vernonia anthelmintica.

ASCARIDE LUMBRICOIDE, Ascaris lumbricoides—a. Vermiculaire, Ascaris vermicularis.

AS'CARIS, gen. Ascar'idis, (ασκαρις, pl. ασκαριδες, from ασκαριζω, 'I leap,') A genus of intestinal worms, characterized by a long, cylindrical body, attenuated at the extremities; and having a mouth furnished with three tubercules, from which a very short tube is sometimes seen issuing. Formerly there were reckoned two varieties of the Ascaris—the As'caris lombricoï'des seu gi'gas homi'nis, Lumbric'us, L. te're's homi'nis, Sco'lex, (F.) Lombricoïde, Ascaride lombricoïde, Lombric, or long round worm; and the As'caris Vermicula'ris—the Ascaris proper—the thread worm or main worm. The former is alone included under the genus, at present—a new genus having been formed of the A. vermicularis, under the name Oxyu'ris. It is the Oxyu'ris seu Fusa'ria vermicula'ris, (F.) Ascaride, A. vermiculaire, Oxyure vermiculaire.

A new species of entozoa was found by Dr. Bellingham, the As'caris alata.

ASCARIS ALATA, see Ascaris—a. Gigas hominis, see Ascaris—a. Lumbricoides, see Ascaris—a. Renalis, Strongylus gigas—a. Trichuria, Trichocephalus—a. Vermicularis, see Ascaris—a. Visceralis, Strongylus gigas.

AS'CELES, (ασκελης,) As'keles, Ca'rens cru'ribus, (α, and σκελος, 'a leg,') One who has no legs.

ASCELLA, Axilla.

ASCEN'DENS, (ascendere, ascensum, (as, and scandere,) 'to ascend,') (F.) Ascendant. Parts are thus called, which are supposed to arise in a region lower than that where they terminate. Thus, Aorta ascendens is the aorta from its origin to the arch: Vena cava ascendens, the large vein which carries the blood from the inferior

parts to the heart: *Obliquus ascendens* (muscle), the lesser oblique muscle of the abdomen, &c.

ASCENSION DE LA MATRICE. Hysteria.

ASCENSUS MORBI. The period of increase of a disease.

ASCESIS, (ασκη, 'to exercise.') Exercise.

ASCHERSONIAN VESICLES, see Vesicles, Aschersonian.

ASCHIL, Scilla.

ASCHISTODACTYLUS, *Syndactylus*, (α, σχιστος, 'cleft'; and δακτυλος, 'a finger.') A monster whose fingers are not separated from one another.

AS'CIA, 'an axe,' *Axi'nē*, *Secepar'nos*, *Dol'abra*, *Fas'cia epira'lis*. Name of a bandage mentioned by the ancients, and figured by Scultetus, in the shape of an axe or hatchet. See Doloire.

ASCILLA, Axilla.

ASCITE, *Ascites*—*a. Peritonéale*, *Ascites*.

ASCITES, (ασκίτης, from ασκος, 'a leathern sack, a large belly:')—*Asci'tes*, *Hydroce'le Peritonæ'i*, *Hy'drops Abdom'inis* seu *Asci'tes*, *Hydrogas'ter*, *Hydroperitone'um*, *Hydroce'lia*, *Hydré'trum*, *Celi'och'y'sis*, *Drop'sy of the low'er bell'y*, *Drop'sy of the Peritone'um*, (F.) *Ascite*, *A. péritonéale*, *Hydropéri-tonite*, *Hydropisie du Bas-ventre ou du péritoine*. A collection of serous fluid in the abdomen. *Ascites* proper is dropsy of the peritoneum; and is characterized by increased size of the abdomen, by fluctuation and the general signs of dropsy. It is rarely a primary disease; but is always dangerous, and but little susceptible of cure. Most generally, it is owing to obstructed circulation in some of the viscera, or to excitement of the vessels of the abdominal organs. The treatment is essentially the same as that of other dropsies. Paracentesis, when had recourse to, can only be regarded as a palliative.

Dropsy of the peritoneum may also be sacculated or in cysts, and occasionally the fluid accumulates exterior to the peritoneum, *Hydropisgastrium*. When in cysts it is termed *Hydrocyst'is*, *Hy'drops abdom'inis sacca'tus* seu *cys'ticus* and *Asci'tes sacca'tus*.

ASCITES HEPATOCYSTICUS, *Turgescencia vesiculæ felleæ*—*a. Ovarii*, *Hydrops ovarii*—*a. Purulentus*, *Pyocœlia*—*a. Saccatus*, see *Ascites*, *Hydroarion*, and *Hydrops ovarii*.

ASCITIC, *Ascit'icus*, (F.) *Ascitique*. Relating or appertaining to, or affected with, ascites.

ASCLEPIADÆ, (Ἀσκληπιάδαι,) *Asclepi'ades*, (Ἀσκληπιάς, 'Æsculapius.') The priest physicians, who served in the ancient temples of Æsculapius, and who took their name from being his descendants.

ASCLÉPIADE, *Asclepias vincetoxicum*.

ASCLEPIAS, (ασκληπιάς,) *Asclepias tuberosa*—*a. Alba*, *A. vincetoxicum*—*a. Apocynum*, *A. Syriaca*—*a. Asthmatica*, *Tylophora asthmatica*—*a. Cornuti*, *A. Syriaca*—*a. Crispa*, *Gonphocarpus crispus*.

ASCLEPIAS CURASSAVICA, *Bastard Ipecacuan'ha*, *Red'head*, *Blood'weed*. The leaves are emetic in the dose of one or two scruples. It is the *Ipecacuanha blanc* of St. Domingo.

ASCLEPIAS DECUMBENS, the root. Escharotic, cathartic, sudorific, diuretic.

ASCLEPIAS, FLESH-COLOURED, *A. Incarnata*—*a. Gigantea*, *Mudar*.

ASCLEPIAS INCARNATA, *A. pul'chra*, *Flesh-coloured* or *incamp asclepias* or *silk'weed*. The root of this plant, which grows in all parts of the United States, and is official in the Ph. U. S., has the same virtues as *A. Syriaca*, and *A. tuberosa*.

ASCLEPIAS OBOVATA, *A. Syriaca*.

ASCLEPIAS PROC'ERA [?], *Beidelossar*, *Beidel'sar*. An Egyptian plant, the leaves of which are made into a plaster, and applied to indolent tumours. The milky juice is caustic, and is used as such.

ASCLEPIAS PSEUDOSARSA, *Hemidesmus In'dicus*—*a. Pubescens*, *A. Syriaca*—*a. Pulchra*, *A. incarnata*.

ASCLEPIAS SYRIACA, *A. pubes'cens* seu *apoc'y'num* seu *obov'ata* seu *toment'osa* seu *cornu'ti*, *Common Silk'weed*, *Milk Weed*, (F.) *Herbe à la honette*. The cortical part of the root, official in the Ph. U. S., has been given, in powder, in asthmatic and pulmonic affections in general, and, it is said, with success.

ASCLEPIAS SULLIVAN'TII, *Smooth Milkweed*, *Silkweed*: indigenous, possesses the same virtues as *A. tuberosa*.

ASCLEPIAS TOMENTOSA, *A. Syriaca*.

ASCLEPIAS TUBERO'SA, *Butterfly Weed*, *Plen'risy Root*, *Flux Root*, *Wind Root*, *Wind Weed*, *White Root*, *Orange Sial'low Root*, *Silk Weed*, *Can'ada Root*, *Orange Apoc'ynum*, *Tu'berous Rooted Sial'low Root*. Said to have been first recommended by the Asclepiades. In Virginia and the Carolinas, the root of this plant—official in the Ph. U. S., (*Asclepias*, Ph. U. S., 1863)—has been long celebrated as a remedy in pneumonic affections. It is sudorific, and the powder acts as a mild purgative. Its chief powers are said to be expectorant, diaphoretic, and febrifuge. It is occasionally given to relieve pains of the stomach from flatulency and indigestion.

ASCLEPIAS VINETOX'ICUM, *A. Al'ba*, *Cynan'chum Vincetoxicum*, *Vincetoxicum*, *V. Officinale*, *Hirundina'ria*, *Apoc'ynum No'tus*, *Aug'lia hirsu'tum*, &c., *Swa'llow-Wort*, *White Swallow-Wort*, (F.) *Asclépiade*, *Dompte-venin*.

The root is said to be stimulant, diuretic, and emmenagogue, but is hardly ever used.

ASCLEPIASMUS, *Hæmorrhoids*.

ASCLEPION, (ασκληπιον,) *Asclepie'ion*, (ασκληπειον,) from Ἀσκληπιος, *Æsculapius*, 'the god of Physic.' A temple of Æsculapius, used, in antiquity, as a place of worship, and frequented by numbers of the sick, who were there attended to.

ASCO'MA, gen. *Asco'matis*, (ασκωμα, from ασκος, 'a bottle.') The eminence of the pubes at the period of puberty in females.

ASE, (αση,) *Anxiety*, *Satiety*.

ASELLI, (pl. of *Asellus*), *Onisci aselli*.

ASELLUS, (dim. of *asinus*, 'an ass,') *Oniseus*.

ASE'MA CRISIS, gen. *As'e'matis*, (ασμα κρίσεως, from *a*, and *σμμα*, 'a sign.') A crisis occurring unexpectedly and without the ordinary precursory signs.

ASEPTON, (ασηπτον, from *a*, and *σπω*, 'I corrupt,') *Asapes*.

ASH, BITTER, *Bittera febrifuga*, *Quassia*—*a. Blue*, *Fraxinus quadrangulata*—*a. Mountain*, *Sorbus acuparia*—*a. Mountain*, *American*; *Sorbus Americana*—*a. Prickly*, *Aralia spinosa*, *Xanthoxylum clava Herculis*—*a. Prickly*, shrubby, *Xanthoxylum fraxineum*—*a. Stinking*, *Ptelea trifoliata*—*a. Tree*, *Fraxinus excelsior*—*a. White*, *Fraxinus Americana*.

ASIALIA, *Aptyalia*.

ASIALORRHŒA, (F.) *Asialorrhée*, (α, σιαλας, 'saliva,' and *ρρω*, 'to flow.') Diminution in the flow of saliva.

ASIMINA TRILOBA, see *Carica papaya*.

ASIT'IA, (ασιτια, from *a*, and *σιτος*, 'food.') Abstinence from food. Want of appetite,—*Fastid'ium ciborum*, *Apoclei'sis*.

ASIUS LAPIS, *Assius Lapis*.

ASJAGAN, *As'jogam*. An Indian tree, the juice of whose leaves, mixed with powdered cumin seeds, is employed in India in colic.

ASJOGAM, *Asjagan*.

ASKELES, (ασκλης,) *Asceles*.

ASKER, *Eschar*.

ASKITES, (ασκιτης,) *Ascites*.

ASO'DES, *Asso'des*, (ασωδης, from *αση*, 'disgust,')

'satiety,' and *odes*.) A fever accompanied with anxiety and nausea: *Febris aoides* vel *azo'des*.

ASPALASOMUS, (ασπαλαξ, 'a mole,' and σωμα, 'a body,') A genus of monsters in which there is imperfect development of the eyes. Also, a malformation, in which the fissure and eversion extend chiefly upon the lower part of the abdomen; the urinary apparatus, genitals and rectum opening externally by three distinct orifices.

ASPALTUM, Asphaltum.

ASPARAGINE, see Asparagus.

ASPARAGUS, (ασπαργος,) *Asparagus officinalis*. Common *Asparagus*, *Sparagus*, *Speragus*, *Sparagras*, *Spar'roigrass*, *Grass*, *Sperage*. Ord. Asphodelaceæ. Sec. Syst. Hexandria Monogynia. *Asparagi officinalis* Turro'nes, (F.) *Asperge*. The fresh roots are diuretic, perhaps owing to the immediate crystallizable principle, *Asparagine*, which is said to be sedative in the dose of a few grains. The young shoots are a well known and esteemed vegetable diet. They communicate a peculiar odour to the urine. A syrup made of the young shoots and an extract of the roots has been recommended as a sedative in heart affections.

ASPA'SIA, (after Aspasia, the mistress of Pericles. [?]) A ball of wood soaked in an infusion of galls, and used by females for constringing the vagina.

ASPEN, AMERICAN, *Populus tremuloides* — a. European, *Populus tremula*.

ASPERA ARTERIA, Trachea.

ASPERGE, Asparagus.

ASPERGILLI SPECIES, see Fungus.

ASPERITAS, (*asper*, 'rough,') *Asperity* — a. *Arterie asperæ*, *Raucedo*.

ASPÉRITÉ DES PAUPIÈRES, *Trachoma-Asperitudo*, *Trachoma*.

ASPER'ITY, *Asper'itas*. *Asperities* are inequalities on the surfaces of bones, which often serve for the insertion of fibrous structures.

ASPERMATIA, *Aspermatismus*.

ASPERMATISMUS, *Asper'mia*, *Aspermat'ia*, (α, and σπέρμα, 'sperm,') Reflux of sperm from the urethra into the bladder, during the venereal orgasm.

ASPERMIA, *Aspermatismus*.

ASPER'SIO, *Asper'sio*, *Catapasma*, *Fomentation*.

ASPER'SION, *Asper'sio*, (*aspergere*, (from *as*, and *spargere*,) 'to sprinkle,') (F.) *Arroisement*. Act of sprinkling or pouring a liquid *guttatim* over a wound, ulcer, &c.

ASPERULA, (diminutive of *asper*, 'rough,') the seeds being rough,) *Galium asperum*.

ASPER'ULA ODORATA, *Galium odoratum*, *Mat'riay'ca*, *Hepatic'ca stellata*, (F.) *Aspérule odorante* ou *Maquet des bois*, *Hépatique étoilée*. Ord. Rubiaceæ. Sec. Syst. Tetrandria Monogynia. Sweet-scented Woodroof. Said to be diuretic, deobstruent, tonic, and vulnerary.

ASPÉRULE ODORANTE, *Asperula odorata*.

ASPHALITES, (ασφαλτης,) *Nephrit'es*, *Nephrit'is*, *Prima Vertebra lumb'ria*, (same etymon as asphaltum.) A name given by some to the first lumbar vertebra.

ASPHALTUM, *Nep'ta*, *Arsal'tos*, (from ἀσφαλίζω, 'to strengthen,' being used as a comment. [?]) With the Greeks, ἀσφαλ'is, signified any kind of bitumen. *Asphaltum* is now restricted chiefly to the BITUMEN OF JUDEA. B. *Juda'icum*, A. *solidum*, *Jews' Pitch*, *Ca'rabé*, *Ka'rabé* of Sodom, (F.) *Asphalte*, *Bitume de Judée*, *Poix minérale scorifiée*, *Baume de momie* ou *des funérailles*. It is solid, friable, vitreous, black, shining, inflammable, and of a fetid smell. An oil is obtained from it by distillation. It enters into the composition of certain ointments and plasters.

It is collected on the surface of the water of the Dead Sea or Lake Asphaltites, in Judæa.

ASPHARINE, *Galium aparine*.

ASPHODELUS, (ασφodelος,) A. *Ramosus* seu *Al'bus* seu *Ma'ris*, *Has'tula Re'gia*, (F.) *Lis asphodelæ*. The bulbs of this southern European plant have an acrimony which they lose in boiling water. They contain a fecula with which bread has been made, and have been considered diuretic. They have been used as a succedaneum for the squill.

ASPHYC'TIC, *Asphycticus*, *Asphyctus*, (same etymon as *Asphyxia*.) Appertaining or relating to asphyxia.

ASPHYCTUS, (ασφυκτος,) *Asphyctic*. Pulseless.

ASPHYXIA, (ασφύξια, from α, and σφύξις, 'pulse,') *Defectus Pul'sus*, *Aeroti'mus*, *Sidera'tio*, *Sydera'tio*, (F.) *Anhæmatisie*. For a long time, *Asphyxia* was confined to the sense of 'suspension of circulation or syncope.' *Pulselessness*. It now generally means *suspend'ed animation*, produced by the nonconversion of the venous blood of the lungs into arterial, — *Apno'a*, *Apneus'tia*, *Apnoasphyx'ia*, *Anhæmato'sia*, *Ecl'lysis pneumocardi'aca*, (F.) *Apnée*, *Anoxémie*, *Hypoxémie*. Owing to the supply of air being cut off, the unchanged venous blood of the pulmonary artery passes into the minute radicles of the pulmonary veins, but their peculiar excitability requiring arterial blood to excite them, more or less stagnation takes place in the pulmonary capillaries, and death occurs chiefly from this cause, and from the want of arterial blood, and not owing to venous blood being distributed through the system, and 'poisoning' it, as was the idea of Bichat. *Car'us asphyx'ia*, *Mors appa'rens*, *Mors putati'va*, *Pseudothan'atus*, *Apparent death*, *Death trance*, (F.) *Mort appa'rente*, is characterized by suspension or diminution of respiration and circulation, to such a degree as to be scarcely, or not at all, perceptible.

Several varieties of *Asphyxia* have been designated.

1. **ASPHYXIA OF THE NEWBORN**, *A. neonatorum*. (F.) *Asphyxie des nouveau-nés*. This is often dependent upon the feeble condition of the infant, or on torpor of the medulla oblongata, not permitting respiration to be established.

2. **ASPHYXIA BY INHALATION OF GASES**, some of which cause death by producing a spasmodic closure of the glottis; others by the want of oxygen.

3. **ASPHYXIA, FROM MECHANICAL CAUSES**, as by Strangulation, or Hanging.

4. **ASPHYXIA BY SUBMERSION**, *A. by drown'ing*, *A. Immers'um*, *Angiairhydrie* of Piorry, as occurs in the drowned, who perish in consequence of the medium in which they are plunged being unfit for respiration. See Submersion, and Suffocation.

5. **ASPHYXIA FROM TORPOR OF THE MEDULLA OBLONGATA**.

For modes of resuscitation, see Ready method of Marshall Hall, and Sylvester's method.

Mr. Chevalier has used the term *Asphyxia Idiopath'ica*, for fatal syncope owing to relaxation of the heart. See Suffocation.

ASPHYXIA IMMERSORUM, *A. by submersion* — a. Local, — see Gangrene — a. Neonatorum, *A. of the new-born* — a. Pestilential: — see Cholera — a. Pestilential: — see Cholera.

ASPHYX'IAL, *Asphyx'ic*. Relating to asphyxia — as '*asphyxial phenomena*.'

ASPHYXIE, *Asphyxia* — a. *des Nouveau-nés*, *Asphyxia of the newborn* — a. *des Parties*, *Gangrene* — a. *Lente des nouveau-nés*, *Induration of the cellular tissue*.

ASPHYX'IED, *Asphyxia'ted*, *Asphyxia'tus*, In a state of asphyxia.

ASPIC, *Lavandula spica*.

ASPIDISCUS, (ασπίδισκος, 'a small round shield,') *Sphincter ani externus*.

ASPIDIUM ATHAMANTICUM, (ασπίδιον, 'a small shield,' dim. of *aspis*, 'a shield.') A South African fern, *Ord.* Filices, which is possessed of anthelmintic properties. Its caudex, in the form of powder, infusion, or electuary, has been found excellent in helminthiasis, and especially in tape-worm.

ASPIDIUM CORIACEUM, *Calagualæ radix*—a. Depastum, *Polypodium filix mas*—a. Discolor, see *Calagualæ radix*—a. Erosum, *Polypodium filix mas*—a. Ferrugineum, see *Calagualæ radix*—a. *Filix femina*, *Asplenium filix femina*—a. *Filix mas*, *Polypodium filix mas*.

ASPIRATIO, Aspiration, Inspiration.

ASPIRATION, *Aspiratio*, *Aspiratio*, (*aspire*, (*as*, and *aspire*), 'to breathe.') The French sometimes use the term synonymously with inspiration. It also means the act of attracting or sucking like a pump. Imbibition. Also, the pronunciation of a vowel with a full breath. See Derivation.

ASPIS, gen. *Aspidis*, (*aspis*.) A name given by the ancients to a venomous serpent—the *Ægyptian viper* of Lacépède, (F.) *Aspic*. Its bite is very dangerous, and it is supposed to have been the reptile which Cleopatra used for her destruction.

ASPLENIUM, *Asplenium*, (*ασπληνον*, from *a*, and *σπλην*, 'the spleen,' from its reputed action.) *Spleenwort*, *Miltworte*.

ASPLENIUM AUREUM, *A. ceterach*.

ASPLENIUM CETERACH, *A. auræum* seu *latifolium*, *Gymnogramme ceterach*, *Doradilla*, *Blechnum squarrosum*, *Scolopendria*, *Athyrium*, *Ceterach officinarum* seu *Canariensis*, *Grammitis ceterach* seu *auræa*, *Gymnopteris ceterach*, *Vitaria ceterach*, (F.) *Doradille*. *Ord.* Filices. Supposed to be substringent and mucilaginous, and has been recommended as a pectoral. It has also been given in calculous cases.

ASPLENIUM FILIX FEMINA, *Polypodium filix femina* seu *molle* seu *dentatum* seu *incisum* seu *trifidum*, *Aspidium filix femina*, *Athyrium filix femina* seu *molle* seu *ovatum* seu *trifidum*, *Pteris palustris*, *Femalæ fern*, *Spleenwort*, (F.) *Fougère, femelle*. The root of this plant resembles that of the male fern, and is said to possess similar anthelmintic virtues. The name *female fern* is also given to *Pteris aquilina*.

ASPLENIUM LATIFOLIUM, *A. ceterach*—a. Murale, *A. ruta*—a. Obtusum, *A. ruta muraria*.

ASPLENIUM RUTA MURARIA, *A. murale* seu *obtusum*, *Paronychia*, *Phyllitis ruta muraria*, *Scolopendrium ruta muraria*, *Adiantum album*, *Ruta muraria*, *Salvia Vitæ*, *Wallrue*, *White Maidenhair*, *Tentwort*, (F.) *Rue des murailles*, *Sauce-vie*. Used in the same cases as the last.

ASPLENIUM SCOLOPENDRIUM, *Scolopendrium officinarum* seu *lingua seu phyllitis seu vulgare*, *Scolopendra*, *Scolopendria*, *Hart's Tongue*, *Spleenwort*, *Phyllitis*, *Lingua cervina*, *Blechnum hypnifolium*, (F.) *Scolopendre*, *Longue de cerf*. Properties like the last. See Calcifraga.

ASPLENIUM TRICHOMANOIDES, *A. Trichomanes*, *Phyllitis rotundifolia*, *Calyphyllum*, *Trichomanes*, *T. crenata*, *Adiantum rubrum*, *Common Maidenhair*, *Polytrichum commune*, (F.) *Polytrich*. Properties like the last.

ASPLENUM, (*ασπληνον*.) *Asplenium*.

ASPREDO, gen. *Aspreddinis*, (*asper*, 'rough,') *Trachoma*—a. Miliacea, *Miliary fever*.

ASPRELE, *Hippuris vulgaris*.

ASPRITUDO, gen. *Aspritudinis*, (*asper*, 'rough,') *Trachoma*.

ASSABA. A Guinea shrub, whose leaves are considered capable of dispersing buboes.

ASSACOU, *Hura Brasiliensis*.

ASSA DOUX, ('sweet Assa,') Benjamin—a. Duleis, Benjamin—a. O odorata, Benjamin.

ASSAFETIDA, *Asafoetida*.

ASSAFETIDA, *Asafoetida*.

ASSAIERET. A compound of bitter, stomachic, and purgative medicines in the form of pill.

ASSAINISSEMENT (F.), (*as*, and *sauare*, 'to make healthy,') The act of rendering healthy, as by the draining of marshes, the disinfection of the air, &c.

ASSAISONNEMENT, (*as*, and *saison*, 'season,') Condiment.

ASSAKUR, *Saccharum*.

ASSALA, see *Myristica moschata*.

ASSATIO, *Opte'sis*, (*assare*, 'to broil,') The preparation of food or medicines in their own juice, without the addition of any liquid. Various kinds of cooking by heat.

ASSELLA, *Axilla*.

ASSERAC, *Assis*. A preparation of opium, or of some narcotic, used by the Turks as an excitant.

ASSERCULUM, (dim. of *asser*, 'a board,') Splint.

ASSERVATION, (*ad*, and *servare*, 'to keep,') Conservation.

ASSES' MILK, see *Milk*, *asses'*.

ASSES' MILK, ARTIFICIAL, see *Milk*, *asses'*.

ASSIDENS, (*as*, and *sedere*, 'to be seated,') That which accompanies or is concomitant. An epithet applied to the accessory symptoms, *Assidentia signa*, and general phenomena of diseases.

ASSIDENTIA SIGNA, see *Assidens*.

ASSIMILABLE, *Assimilabilis*, (*as*, *similis*, 'like,' and *habilis*, 'able,') That which is susceptible of assimilation.

ASSIMILATING, *Assimilatory*, *Assimilatory*, (F.) *Assimilateur*, (*as*, and *similis*, 'like,') That which effects assimilation, as an *assimilating* or *assimilatory* organ.

ASSIMILATION, *Assimilatio*, *Similatio*, *Appropriatio*, *Exomoi'sis*, *Homoi'sis*, *Threp'sis*. The act by which living bodies appropriate and transform into their own substance matters with which they may be placed in contact. The reverse action—that of separating and eliminating matters already assimilated, in the act of nutrition—is termed *Destructive Assimilation*, (F.) *Désassimilation*.

ASSIMILATION, DESTRUCTIVE, see *Assimilation*.

ASSIS, *Asserac*.

ASSIUS LA'PIS, *A'sius La'pis*. A sort of stone or earth, found near the town of Assa in the Troad, which had the property of destroying proud flesh.

ASSOCIATED MOVEMENTS, see *Instinctive*.

ASODES, (*ασωδης*), *Asodes*.

ASSOUPISSANT, (L. *opor*, 'sleep,') *Somniferous*.

ASSOUPISSEMENT, *Somnolency*.

ASSOURON, see *Myrtus Pimenta*.

ASSUETUDO, gen. *Assuetudinis*, (*as*, and *sueto*, 'to accustom to anything,') *Habit*.

ASSULA, (dim. of *assis*, 'a board,') Splint.

ASSULTUS, (*assilio*, *assultum*, (*as*, and *salio*,) 'to leap upon,') *Attack*.

ASSUMPTIO, (*as*, and *sumere*, *sumptum*, 'to take,') *Prehension*.

ASTACI FLUVIATILIS CONCREMENTA, *Cancerorum chelæ*.

ASTACUS FLUVIATILIS, (*αστακος*;) *Crab*.

ASTAKILLOS, *Araneum ulcus*.

ASTARZOF. An ointment, composed of litharge, frog's spawn, &c. Also, camphor dissolved in rose water,—with *Paracelsus*.

ASTASIA, (*αστασια*, from *a*, and *στασις*, 'to fix,') *Dysphoria*.

ASTER ATTICUS, (from *astron*, 'a star,' after the shape of the flowers,) *Bubonium*.

ASTER CORDIFOLIUS, *Heart-leaved Aster*, *A. Punicæus*, *Rough-stemmed Aster*, and other indigenous species, *Ord.* *Compositæ*, possess aromatic properties.

ASTER DYSENTERICUS, *Inula dysenterica*—a. Heart-leaved, *A. cordifolius*—a. *Helenium*, *Inula Helenium*—a. *Inguinalis*, *Eryngium campestre*—a. *Officinalis*, *Inula Helenium*—a. Rough-stemmed, *A. Puniceus*—a. *Tortifolius*, *Sericocarpus tortifolius*—a. *Undulatus*, *Inula dysenterica*.

ASTERIA GEMMA, (*asteria*,) *Asterius*, *Astroites*, *Astrios*, *Astrobelolus*. The ancients attributed imaginary virtues to this stone,—that of dispersing *Navi Materni*, for example.

ASTERIAS LUTEA, (from *aster*), *Gentiana lutea*.

ASTER'NIA, (*a*, and *στερνον*, 'the sternum.') A monstrosity, in which there is absence of the sternum.

ASTEROCEPHALUS SUCCISA, (*αστηρ*, 'a star,' and *κεφαλη*, 'head,') *Scabiosa succisa*.

ASTHENES, (*ασθενς*,) Infirm.

ASTHEN'IA, *Via imminuta*, (*ασθενια*, from *a*, and *σθενς*, 'force,' 'strength,') Want of strength, debility. (F.) *Affaiblissement*. Infirmary. A word used in this sense by Galen, and employed, especially by Brown, to designate debility of the whole economy, or diminution of the vital forces. He distinguished it into *direct* and *indirect*: the former proceeding from diminution of stimuli; the latter from exhaustion of incitability by the abuse of stimuli.

ASTHENIA DEGLUTITIONIS, Pharyngoplegia—a. Pectoralis, *Angina Pectoris*.

ASTHEN'IC, (*ασθενικος*,) *Asthen'icus*, (F.) *Asthénique*. Relating or belonging to asthenia.

ASTHENICOPYRA, (*ασθενς*, 'weak,' and *πυρ*, or *πυρετος*, 'fever,') Fever, adynamic.

ASTHENICOPYRETUS, Fever, adynamic.

ASTHÉNIQUE, Asthenic.

ASTHENO'PLIA, *Debilitas* seu *Hebetudo visus*, (F.) *Affaiblissement de la Vue*, (*ασθενς*, 'weak,' and *ωψ*, 'the eye,') Weakness of sight; *Weak-sightedness*.

ASTHENOPYRA, (*ασθενς*, 'weak,' and *πυρ*, or *πυρετος*, 'fever,') Fever, adynamic, Typhus.

ASTHENOPYRETUS, Fever, adynamic.

ASTHMA, gen. *Asth'matis*, (*ασθμα*, 'laborious breathing,' from *αω*, 'I respire.') *A spas'ticum adult'rum* seu *bronch'i'tis* seu *senio'rum* seu *convuls'cum* seu *spas'ticum intermit'tens*, *Spas'mus bronch'i'tis*, *Dyspno'e'a* et *orthopno'e'a convuls'i'va*, *Ma'lum Cadu'cum pulmo'num*, *Broken-windedness*, *Nervous asthma*, (F.) *Asthme, A. nerveux*. Great difficulty of breathing, recurring at intervals, accompanied with a wheezing sound and sense of constriction in the chest; cough and expectoration. It is a chronic disease, and not curable with facility. There are no pathognomonic physical signs of it. In some cases, the respiration is universally puerile during the attack. In the spasmodic form, the respiratory murmur is very feeble or absent during the fit; and in all forms percussion elicits a clear pulmonary sound. The disease generally consists in some source of irritation, and occasionally, perhaps, in paralysis of the pneumogastric nerves, *Bronchoparalysis*, *Paralysis ner'vi va'gi in par'te thorac'i'ca*, more frequently of the former—all the phenomena indicating constriction of the smaller bronchial ramifications. The treatment is one that relieves spasmodic action—anæsthetics, narcotics, counter-irritants, change of air, &c.

ASTHMA ACU'TUM, of Millar, *A. Millari acu'tum*, *A. spas'ticum infan'tum*, *Cynan'chè Trachea'lis spasmod'ica*, (F.) *Asthme aigu*. Probably, spasmodic croup [?]. See *Asthma Thymicum*.

ASTHMA AËRIUM, *Pneumothorax*—a. *Aërium ab Emphysemate Pulmonum*, *Emphysema of the Lungs*—a. *Arthriticum*, *Angina Pectoris*—a. *Bronchiale*, *Asthma*.

ASTHMA CAR'DIAC, *Asthma cardi'acum*, *Cardi-asthma*, (F.) *Dyspnée ou Asthme Cardiaque*, *Dyspnœa* dependent upon disease of the heart.

ASTHMA CONVULSIVUM, *Angina pectoris*—a. *Diaphragmaticum*, *Angina Pectoris*—a. *Dolorificum*, *Angina pectoris*—a. *Emphysematicum*, *Pneumothorax*.

ASTHMA, GRIND'ERS', *Grind'ers' Disease* or *Rot*. The aggregate of functional phenomena, induced by the inhalation of particles thrown off during the operation of grinding metallic instruments, &c. The structural changes induced are enlargement of the bronchial tubes, expansion of the pulmonary tissue, and phthisis.

ASTHMA GYPSEUM, *A. pulverulentum*—a. Hay, Fever, hay.

ASTHMA HU'MIDUM, *Hu'mid*, *Com'mon*, or *Spitting asthma*, is when the disease is accompanied with expectoration. It is also called *A. humoral'e*, *A. flatulent'um*, *A. pneumo'nicum*, *Blennothorax chron'icus*, &c.

ASTHMA INSPANDIUM, *Cynanche trachealis*—a. *Infantum Spasmodicum*, *A. Thymicum*—a. *Koppian*, *A. Thymicum*—a. *Millari acutum*, *Asthma acutum*—a. *Montanum*, *A. pulverulentum*—a. *Nervous*, *Asthma*—a. *Nocturnum*, *Incubus*.

ASTHMA PULVERULENTUM, *A. gyp'seum*, *A. montanum*. The variety of asthma to which millers, bakers, grinders, and others are subject.

ASTHMA SIC'CUM, so called when the paroxysm is sudden, violent, and of short duration; cough slight, and expectoration scanty; spasmodic constriction.

ASTHMA SPASTICO-ARTHRITICUM INCONSTANS, *Angina pectoris*—a. *Spasticum Infantum*, *A. Thymicum*.

ASTHMA THY'MICUM, *A. T. Kopp'ii*, *Asthma spas'ticum infan'tum* seu *infan'tum spasmo'dicum*, seu *larynge'um infan'tum* seu *intermit'tens infan'tum* seu *dienten'tium* seu *period'icum acu'tum*, *Thymasthma*, *Cynanchè trachea'lis spasmo'dica*, *Spas'mus glott'idis*, *Kopp'ian Asthma*, *Thy'mic Asthma*, *Laryngis'mus strid'ulus*, *Laryngospas'mus*, *Apno'e'a infan'tum*, *Spasm of the larynx or glottis*, *Croup-like inspiration of infants*, *Child-crowling*, *Spasmodic croup*, *Pseudocroup*, *Spir'ious* or *Cerebral croup*, *Phrenoglott'ism*, *Suffocating ner'vous catarrh*, (F.) *Laryngite stridul'euse*, *Faux Croup*, *Pseudo-croup nerveux*, *Spasme de la Glotte et du Thorax*. A disease of infants, characterized by suspension of respiration at intervals; great difficulty of breathing, with *crowing inspiration*, especially on waking, swallowing, or crying; ending often in a fit of suffocation, with convulsions. These symptoms are often accompanied by rigidity of the fingers and toes; the thumb being frequently drawn forcibly into the palm of the clenched hand, whence the name *Car'po-pe'dal spasm*, applied, at times, to the disease.

The pathology of the disease has been supposed to consist in an enlargement of the thymus gland, or of the glands of the neck pressing on the pneumogastric nerves [?]. The ear, by auscultation, at a distance from the chest, detects an incomplete, acute, hissing inspiration, or rather cry; whilst the expiration and voice are croupal, both at the accession and termination of the paroxysm. The heart's action has been observed to be distinct and feeble.

ASTHMA TY'PICUM. *Asthma* characterized by periodicity.

ASTHMA UTERI, *Hysteria*—a. *Weed*, *Lobelia inflata*.

ASTHMAT'IC, (*ασθματικος*,) *Asthmat'icus*, *Pneocolyt'icus*, *Brokenwinded*, *Bel'loned* (Prov.) (F.) *Asthmatique*. Affected with asthma. Relating to asthma.

ASTHME. *Asthma*—a. *Aigu*, *Asthma acutum*—a. *Cardiaque*, *Asthma*, *cardiac*—a. *Nerveux*, *Asthma*.

ASTIG'MATISM, *Astigmatism'us*, (a. and *στυμα*.

gen. σημῆμα, 'a point.') A state of irregular refraction in the eye, in which the rays are not brought to one focus, but converge at different distances, so as to form two linear images at right angles to each other.

ASTOMUS, (αστομος, from α, and στομα, 'a mouth.') (F.) *Astome*. One without a mouth. Pliny speaks of a people in India without mouths, who live *anhelati et odore!*

ASTRAGALE COL D', Collum astragali.

ASTRAGALOIDES SYPHILITICA, Astragalus exsensus.

ASTRAG'ALUS, (αστραγαλος,) *Talus*, the *Ankle*, *Quadratio*, *Quartio*, *Quaternio*, *Diabebos*, *Peza*, *Cuvicula*, *Cuvilla*, *Tetroros*, *As'trion*, *Oa Balistæ* seu *Tes'sera*, (αστραγαλος, 'a die' which it has been considered to resemble [?].) A short bone, situate at the superior and middle part of the tarsus, where it is articulated with the tibia. It is the *ankle bone*, *sling bone*, or *first bone of the foot*. The anterior surface is convex, and has a well-marked prominence, supported by a kind of neck, and hence has been called the *head of the astragalus*. The astragalus is developed by two points of ossification.

Also, the atlas or first vertebra of the neck.

ASTRAG'ALUS EXSCAPUS, *Astragaloides syphilitica*, *Sten'less Milk'vetch*, (F.) *Astragale à gousses velus*. Ord. Leguminosæ. Ser. Syst. *Dialuphia Decandria*. The root is said to have cured confirmed syphilis.

ASTRAG'ALUS TRAGACANTHUS, see *Tragacantha*.

ASTRAG'ALUS VE'RUS, *Spi'na hirci*, *Astrag'alus aculeatus*, *Goat's thorn*, *Milkvetch*. The plant which affords *Gum Trag'acanth*. See *Tragacantha*.

ASTRANTIA, (from *astro*,) *Imperatoria*—a. *Diapensia*, *Sanicula*.

AS'TRAPE, (αστραπη,) *Coruscat'io*, *Fulgur*, *Fulmen*, *Light'ning*, (F.) *Eclair*. Galen reckons it amongst the remote causes of epilepsy.

ASTRICT'ION, *Astrict'io*, *Styp'sis*, *Adstrict'io*, *Constrict'io*, (astringere, (ad and stringere,) 'to constringe.') Action of an astringent substance on the animal economy.

ASTRICTORIA, *Astringents*.

ASTRINGENT ROOT, *Comptonia asplenifolia*.

ASTRIN'GENTS, *Astringen'tia*, *Adstrictoria*, *Astringen'tia*, *Styp'h'na*, *Catastalt'ica*, *Constringen'tia*, *Contrahe'n'tia*, *Stegno'tica*, *Syncrit'ica*, *Astrictoria*. Same etymon as *Astriction*. Medicines which have the property of constringing the organic textures. External astringents are called *Styptics*.

ASTRION, *Astragalus*.

ASTRIOS, (αστηρ, 'a star.') *Asteria gemma*.

ASTRO, (from αστρον, 'a star.') In composition, a star.

ASTROBLES, (αστροβλης, from *astro*, and βαλλω, 'I strike.') One struck by the stars, (*sideratus*.) One who is in a state of sideration—in an apoplectic state.

ASTROBOLESIA, (αστροβολησια,) *Astrobolismus*.

ASTROBOLIA, (αστροβολια,) *Astrobolismus*.

ASTROBOLIS'MUS, *Astrobol'ia*, *Astrobol'e'sia*, *Helio'sis*, *Helio'sis*. (Same etymon as *Astrobles*.) *Sideration*, or action of the stars on a person. *Apoplexy*.

ASTROBOLUS, *Asteria gemma*.

ASTROITIS, (αστηρ, 'a star.') *Asteria gemma*.

ASTROLOGER, see *Astronomy*.

ASTROL'OGY, *Astrolog'ia*, (αστρολογία, from *astro*, and *logos*, 'a discourse.') The art of divining by inspecting the stars *Astromanti'a*. This was formerly considered to be a part of medicine; and was called *Judic'ial Astrology*, to distinguish it from *astronomy*.

ASTROMANTIA, *Astrology*.

ASTRON'OMY, *Astronom'ia*, (αστρονομια, from *astro*, and *nomos*, 'a law,' 'rule.') A science which makes known the heavenly phenomena, and the laws that govern them. Hippocrates places this and astrology amongst the necessary studies of a physician. In the earlier English writers, *Astronomer* is often used in the sense of *Astrologer*, and conversely.

ASTRUTHIUM, *Imperatoria*.

ASTYPHIA, (α, and στυειν, 'to make erect.') *Impotence*.

ASTYSIA, (same etymon,) *Impotence*.

ASUAR, *Myrobalanus Indica*.

ASULCI, *Lapis lazuli*.

ASYNERGIA, (α, and *synergy*.) Want of correlation between organs.

ASYNESIA, (α, and *συνιμι*, 'to comprehend,') *Dementia*.

ASYNODIA, (α, and *συνος*, 'copulation.') *Impotence*.

ASYSTOLIE, (α, and *συστολη*, 'systole.') A name given by M. Beau to a period, in diseases of the heart, in which the systole is incomplete; or, at least, is insufficient to free the ventricles from the blood in them.

AT, in composition, the same as *Ad*.

ATACTOS, (ατακτος, from α, and *τασσω*, 'to put in order,') *Erratic*.

ATARACTAPOE'SIA, *Ataractopoe'sia*, (αταρακταποιησια, from α, *ταρακτος*, 'troubled,' and *ποιειν*, 'to make,') *Intrepidity*, *firmness*; a quality of which, according to Hippocrates, the physician ought to be possessed in the highest degree.

ATARAX'IA, (αταραξια, from α, and *ταραξις*, 'trouble,' 'emotion.') *Moral tranquillity*, *peace of mind*.

AT'AVISM, *Atavism'us*, (*atavus*, 'an old grand-sire or ancestor indefinitely.') The case in which an anomaly or disease, existing in a family, is lost in one generation and reappears in the following.

ATAX'IA, (αταξια, from α, and *ταξις*, 'order.') Disorder, irregularity. Hippocrates employs the word in its most extensive acceptance. Galen applies it, especially, to irregularity of pulse; and Sydenham speaks of *Ataxia Spirituum* for disorder of the nervous system. *Ataxia*, now, usually means the state of disorder that characterizes nervous fevers, and the nervous condition.

ATAXIA SPIRITUM, *Nervous diathesis*. See *Ataxia*.

ATAX'IC, *Atax'icus*. Having the characters of *ataxia*.

ATCHAR, *A'chia*, *Achar*. A condiment used in India. It is formed of green fruits of various kinds—garlic, ginger, mustard, and pimento—pickled in vinegar.

ATECNIA, (ατεκνια, from α, and *τεκνον*, 'a child.') *Sterilitas*.

ATELEC'TASIS, *Atelek'tasis*, (*atelo*, and *εκτασις*, 'dilatation,') *Imperfect expansion* or *dilatation*; as in

ATELEC'TASIS PULMO'NUM, *Pneumonatelec'tasis*, *Pneumatelec'tasis*. *Imperfect expansion* of the lungs at birth, giving rise to *Cyano'sis pulmonalis*. A similar condition is observed in lungs which have received air, and in a distinctly lobular form. This is regarded by Dr. W. T. Gairdner as, in all probability, a secondary lesion, and dependent, in the majority of instances, on a catarrhal condition of the bronchial tubes. It is called, by him, *pulmonary collapse*, or *collapse of the lung*, and has received the name *apneumato'sis* from Fuchs.

ATEL'IA, (ατελεια, 'imperfection.') (F.) *Atelie*. A monstrosity in which there is a want of some member.

ATELO, (from *ατελης*, 'imperfect,') in composition, defective.

ATELOCAR'DIA, *Cardiatel'ia*, (*atelo*, and *καρδία*, 'the heart.') Imperfect formation of the heart.

ATELOCHEI'LIA, (*atelo*, and *χειλος*, 'lip.') A malformation which consists in an imperfect development of the lip.

ATELOENCEPHAL'IA, (*atelo*, and *εγκεφαλον*, 'the encephalon.') State of imperfect development of the brain.

ATELOGLOS'SIA, (*atelo*, and *γλῶσσα*, 'tongue.') A malformation which consists in an imperfect development of the tongue.

ATELOGNA'THIA, (*atelo*, and *γναθος*, 'the jaw.') A malformation which consists in an imperfect development of the jaw.

ATELOMYEL'IA, (*atelo*, and *μυελος*, 'marrow.') State of imperfect development of the spinal marrow.

ATELOPROSO'PIA, (*atelo*, and *προσωπον*, 'the face.') A malformation which consists in imperfect development of the face.

ATELORACHIDIA, (*atelo*, and *ραχis*, 'spine.') Hydrorachis.

ATELOSTOM'IA, (*atelo*, and *στομα*, 'mouth.') One whose mouth is imperfectly developed.

ATER SUCCUS, ('black juice.') Atrabilis.

ATHAMAN'TA, (Athamas, a place in Thessaly.) A genus of plants.

ATHAMANTA ANNUA, *A. Cretensis*.

ATHAMAN'TA AUREOSILI'NUM, *Oreoseli'num*, *O. legitimum* seu *nitrum*, *Seli'num oreo'seli'num*, *Peuced'anum oreoseli'num*, *Apium montanum*, *Black Mountain Parsley*, (F.) *Persil de Montagne*; *Ord. Umbelliferae*. The plant, seed, and roots, are aromatic. It has been considered attenuant, aperient, deobstruent, and lithontriptic. The distilled oil has been used in toothache.

ATHAMANTA CRETENSIS seu *CRE'TICA*, *A. an'nua*, *Libano'tis an'nua* seu *Creten'sis* seu *hirsu'ta*, *Daucus Cre'ticus* seu *Candidianus*, *Myrr'his an'nua*, *Black Car'rot*. The seeds of this plant are acrid and aromatic. They have been used as carminatives and diuretics.

ATHAMANTA MACEDONICA, *Bubon Macedonicum*—*a. Meum*, *Æthusa meum*.

ATHANASIA, *Tanaetum*.

ATHANA'SIA, (*αθανασία*, from *a*, and *θανατος*, 'death.') An antidote for diseases of the liver, jaundice, gravel, &c. It consisted of saffron, cinnamon, lavender, cassia, myrrh, juncus odoratus, honey, &c., and was esteemed to be sudorific.

ATHARA, (*αθαρα*), *Athera*.

ATHELAS'MUS, (*a*, and *θηλη*, 'a breast or nipple.') Impracticability of giving suck; from want of nipple or otherwise.

ATHE'LES, *Athel'us*, (*αθλης*, *αθλος*, from *a*, and *θηλη*, 'the nipple, the breast.') Without the breast; weaned; devoid of nipple; one that has not sucked.

ATHE'NA, (*αθηνα*.) Name of a plaster, recommended by Aesclepiades, and composed of oxide of copper, sublimed oxide of zinc, sal ammoniac, verdigris, gall nuts, and a variety of resinous and other ingredients.

ATHENIO'NIS CATAPOT'IUM. A pill, composed of myrrh, pepper, castor, and opium; used to allay coughing.

ATHE'RA, *Atha'ra*, (*αθηρα*, from *αθηρ*, 'an ear of corn.') A kind of pap for children; also, a kind of liniment.

ATHERAPEUTUS, (*αθεραπευτος*, from *a*, and *θεραπευω*, 'I treat or cure.') Incurable.

ATHERO'MA, gen. *Athero'matis*, (*αθηρωμα*, from *αθηρα*, 'pap or pulp.') *Emphy'ma encyst'is athero'ma*, *Mollus'cum*, *Pultu'tio*, (F.) *Athérome*. A tumour formed by a cyst containing matter like pap or *bouillie*, or plaster.

ATHEROM'ATOUS, *Atheromato'des*, (F.) *Athéromateux*. Having the nature of Atheroma.

ATHÉROME, Atheroma.

ATHLE'TA, (*αθλητης*, from *αθλος*, 'a combat.') (F.) *Athlète*. Athletæ were men who exercised themselves in combats at the public festivals.

ATHLET'IC, (*αθλητικος*), *Athl'eticus*, (F.) *Athlétique*; concerning *Athletæ*. Strong in muscular powers.

ATHOL PORRIDGE, see Mahogany.

ATHORACOCEPHALUS, (*a*, *θωραξ*, 'thorax,' and *κεφαλη*, 'head.') Acephalogaster.

A'THRIX, *A'trichus*, *Calvus*, (F.) *Chauve*; (*ασιξ*, from *a*, and *θριξ*, gen. *τριχος*, 'hair.') *Bald*, (Sc.) *Beld*, *Bellit*. One who has lost his hair.

ATHRIX DEPILIS, Alopecia.

ATHYM'IA, *An'imi defect'us* et *anxi'etas* seu *demis'sio*, *Tristi'tia*, *Moror*, *Ly'pē*, (*αθυμία*, from *a*, and *θυμος*, 'heart,' 'courage.') Despondency. The prostration of spirits often observable in the sick.—Melancholy. See Panophobia.

ATHYMIA PLEONECTICA, see Pleonectica.

ATHYRION, *Asplenium ceterach*.

ATHYRIUM FILIX FEMINA. *Asplenium filix femina*—*a. Filix mas*, *Polypodium filix mas*—*a. Molle*, *Asplenium filix femina*—*a. Ovatum*, *Asplenium filix femina*—*a. Trifidum*, *Asplenium filix femina*.

ATLANTAD, see Atlantal aspect.

ATLAN'TAL; same etymon as *Atlas*. Relating or appertaining to the atlas.

ATLANTAL AS'PECT. An aspect towards the region where the atlas is situated.—Barclay. *Atlantad* is used by the same writer to signify 'towards the atlantal aspect.'

ATLANTAL EXTREM'ITIES, The upper limbs.

ATLANTION, Atlas.

AT'LAS, gen. *Atlan'tis*, (*ατλας*, gen. *ατλαντος*, from *ατλω*, 'I sustain,') *Atlan'tion*, *Astrag'alus*. The first cervical ver'tebra; so called from its supporting the whole weight of the head, as Atlas is said to have supported the globe on his shoulders. Chaussier calls it *Atloïde*. This vertebra in no respect resembles the others. It is a kind of irregular ring, into which, anteriorly, the *processus dentatus* of the second vertebra is received. Posteriorly, it gives passage to the medulla spinalis.

ATLOD'YMUS, *Atlodid'ymus*, (F.) *Atlodyme*; (*Atlas*, 'the first bone of the neck,' and *ιδυνος*, 'a twin.') A monster which is simple below and double above. M. Isid. Geoffroy Saint-Hilaire applies this term to a monster, which has a single body, with two distinct heads supported on one neck.

ATLOÏDE, Atlas.

ATLOIDO, in composition, Atlas.

ATLOID'OAX'OID, (F.) *Atloïdo-axoïdien*. Relating to both the Atlas and the Axis or Vertebra Dentata.

ATLOIDOAXOID ARTICULATION. The articulation between the first two cervical vertebrae.

ATLOIDOAXOID LIG'AMENTS. These are two in number; one anterior and another posterior, passing between the two vertebrae.

ATLOID'OCCIP'ITAL. Relating to the atlas and occiput. The *Atloïdo-occip'ital Articula'tion* is formed by the condyles of the occipital bone and the superior articular surfaces of the Atlas. The *Atloïdo-occipital* muscle is the *Rectus capitis posticus minor*.

ATLOÏDOSOUSMASTOÏDIEN, *Obliquus superior oculi*—*a. Sous-occipitale*, *Rectus capitis lateralis*.

ATMIATR'IA, *Atmidiat'ricē*, (*αρμος*, 'vapour,' and *ιατρεία*, 'treatment.') Treatment of diseases by fumigation, or inhalation, (F.) *Médecine pneumatique*.

ATMIADIATRICE, (*αρμς*, gen. *αρμδος*, 'vapour,' and *ιατρεία*, 'treatment.') Atmiatria.

ATMISTERION, (ατμιστος, 'turned into vapour.') Vaporarium.

ATMOLUTRON, (ατμος, 'vapour,' and λουτρον, 'a bath.') Vaporarium.

ATMOS, (ατμος, 'Breath.

ATMOSPHERE, *Atmosphæra*, (ατμος, 'vapour,' and σφαῖρα, 'a sphere,'—as it were *Sphere of vapours*.) The atmosphere is a spherical mass of air, surrounding the earth in every part; the height of which is estimated at 15 or 16 leagues. It presses on the surface of the earth, and this pressure has, necessarily, sensible effects on organized bodies. The surface of the human body being reckoned at 15 square feet, it is computed that a pressure of 33,000 pounds or more exists under ordinary circumstances; and this pressure cannot be increased or diminished materially, without modifying the circulation and all the functions.

ATMOSPHERIZATION, Hæmatisis.

ATOCIA, (α, and τοκος, 'a child.') Sterilitas.

ATOLMIA, (ατολμία, from α, and τολμα, 'confidence,') Want of confidence; discouragement. A state of mind unfavourable to health, and injurious in disease. It is the antithesis of *Eutolmia*.

ATOMY, Skeleton.

ATONIA, (ατονία, 'Atony—a. Ventriculi, Gastresthenia.

ATONIC, *Atonicus*, (F.) *Atonique*. Wanting tone. Also, a medicine capable of allaying organic excitement or irritation.

AT'ONY, *Aton'ia*, *Infir'mitas et Remissio vi-rum*, *Lan'guor*, *Lax'itas*, (ατονία, from α, and τῶνος, 'tone,') Want of tone. Weakness of every organ, and particularly of those that are contractile. Violent gastritis has been described by Scribonius Largus under a similar name, *At'onos*, (ατονον.)

ATRABILIARY, *Atrabili'ous*, *Atrabilia'ria*, *Atrabilio'sus*, (F.) *Atrabilaire*, *Atrabileus*, (ater, 'black,' and bilis, 'bile,') An epithet given by the ancients to the melancholic and hypochondriac, because they believed the *Atrabilis* to predominate in such.

ATRILIARY CAP'SULES, ARTERIES and VEINS. The renal capsules, arteries and veins; the formation of *Atrabilis* having been attributed to them.

ATRABI'LIS, (same etymon,) *A'ter suc'cus*, *Black Bile* or *Mel'ancholy*, (F.) *Atrabile*. According to the ancients, a thick, black, acrid humour, secreted, in the opinion of some, by the pancreas; in that of others, by the supra-renal capsules. Hippocrates, Galen, Aëtius, and others, ascribe great influence to the *Atrabilis* in the production of hypochondriasis, melancholy, and mania. There is really no such humour. It was an imaginary creation.

ATRACHE'LOCEPHALUS, (α, τραχηλος, 'neck,' and κεφαλη, 'head,') A monster whose neck is partially or wholly deficient.

ATRACHE'LUS, (ατραχηλος, from α, and τραχηλος, 'neck,') One devoid of neck or who is very short-necked.

ATRACTYLIS GUMMIF'ERA, (ατρακτυλις,) *Car'buncu'piniens*, *Ixi'vne*, *Gummy-rooted Atractylis*, *Pine This'tle*. Ord. Compositæ. The root, when wounded, yields a milky, viscid juice, which concretes into tenacious masses, and is said to be chewed with the same views as mastich.

ATRAGENE, *Clematis vitalba*.

ATRAMENTUM, (ater, 'black,') *A. Suto'rium*, *Ink*, *Chalcant'hum*, (F.) *Encere*. Made of galls and sulphate of iron, it has been advised as an astringent, and as an external application in herpetic affections.

ATRAMENTUM STUTORIUM, Ferri sulphas.

ATRE'SIA, (see Atreto,) (F.) *Atrésie*. Imperforation. See Monster.

ATRESIA ANI, Aprocætia.

ATRE'SIA A'NI A'DNA'TA, *A'nus Imperforat'us*, *Imperforat'io a'ni*, *Atretocys'ia*, (F.) *Imperforation de l'anus*. Congenital imperforation of the intestinal canal.

ATRESIA VAGINÆ, Colpatresia.

ATRETELYTRIA, (atreto, and ελυτρον, 'vagina,') Colpatresia.

ATRETISMUS, (from atreto,) Imperforation.

ATRETO, (α, and τραω, 'to perforate,') in composition, 'imperforate.'

ATRETOCEPH'ALUS, (atreto, and κεφαλη, 'head,') A monster, in which some of the natural apertures of the head are wanting.

ATRETOCOR'MUS, (atreto, and κορμος, 'trunk,') A monster in which the natural apertures of the trunk are wanting.

ATRETOCYSIA, (atreto, and κυσος, 'anus,') *Atresia ani*.

ATRETOMETRIA, (atreto, and μετρα, 'uterus,') *Hysteratresia*.

ATRETOPSIA, (atreto, and ωψ, 'the eye,') *Coreclisis*.

ATRETOSTOM'IA, (atreto, and στομα, 'mouth,') Imperforation of the mouth.

ATRETURE'THRIA, (atreto, and ουρηθρα, 'urethra,') Imperforation of the urethra.

ATRE'TUS, (ατρετος,) *Imperforat'us*, *Imperforate*. One whose anus, (*aproc'tus*,) or parts of generation, for example, are imperforate.

ATRIA, (pl. of atrium, 'the hall of a house,') Auricles of the heart—a. Mortis, see Death.

ATRICAPILLUS, (ater, 'black,' and capillus, 'hair,') *Melanothrix*.

AT'RICES. Small tumours, which appear occasionally around the anus. Some commentators consider the word to be synonymous with condylomata.

ATRICH'IA, (α, and θριξ, gen. τριχος, 'a hair,') *Alpecia*—a. *Adnata*, see *Alpecia*—a. *Senilis*, see *Alpecia*.

ATRICHUS, (ατριχος,) *Atrich*.

ATRICI, (*atricus*, 'a porter at the atrium or hall,') Small sinuses in the vicinity of the anus, not penetrating the rectum.

ATRIOVENTRICULAR, Auriculo-ventricular.

ATRIplex PÆTIDA, *Chenopodium vulvaria*.

AT'RIplex HORTEN'SIS, *A. Sati'va*, (F.) *Arroche*, *Bonne Dame*, *Follette*. Ord. *Chenopodiaceæ*. The herb and seed of this plant have been exhibited as antiscorbutics.

Atriplex al'vina, *A. Portulacoi'des*, and *A. Pat'ula*, are used as pickles, and have similar properties.

ATRIplex LACINIATA, *A. patula*—a. *Mexicana*, *Chenopodium ambrosioides*—a. *Odorata*, *Chenopodium botrys*—a. *Olida*, *Chenopodium vulvaria*.

ATRIplex PAT'ULA, *a. laciniata* seu *Purshia'na*, *Spread'ing Arroche*; indigenous; Family, *Chenopodiaceæ*. The expressed juice is said to be cathartic. It has been used, in place of gamboge, in dropsy and asthma.

ATRIplex PURSHIANA, *A. patula*.

ATRIUM ('a hall') *CORDIS DEXTRUM*, *Sinus dexter cordis*—a. *Cordis sinistrum*, *Sinus pulmonalis*—a. *Vaginæ*, *Vestibulum*.

AT'ROPA, (from ἄρπτος, 'immutable,') 'the goddess of destiny,' so called from its fatal effects.)

ATROPA BELLADON'NA, *Belladonna*, *B. baccif'era* seu *trichot'oma*, *Sola'num letha'le* seu *horten'se* *nig'rum*, *Sola'num man'acum* seu *furiu'sum* seu *melanocer'æus*, *Strych'nos*, *Deadly Nightshade*, *Common Duale*, (Se.) *Mekilwort*, (F.) *Belladone*, *Morrelle furieuse*, *Belle Dame*. Ord. *Solanaceæ*. *Ser. Syst.* *Tetrandria Monogynia*. The leaves—*Belladonna fo'tium*, (Ph. U. S. 1863,)—are powerfully narcotic, and also diaphoretic, and diuretic. They are occasionally used where narcotics are indicated. Sprinkling the powdered leaves over cancerous sores has been found to allay the pain;

and the leaves form a good poultice. Dose, gr. $\frac{1}{2}$ to gr. j of the powdered leaves.

ATROPA MANDRAGORA, *Mandragora*, *M. vernalis* seu *officinalis* seu *acaulis*, *Circæa*, *Anthropomorphus*, *Sentiviva* *Ma'ham terrestris*, *Man-drake*. The boiled root has been used in the form of poultice to indolent swellings.

ATROPHIA, Atrophy, *Tabes*—a. Ablactatorium, Brash, weaning—a. Cerebri, Pnenatrophia—a. Cordis, Heart, atrophy of the—a. Dentis, Odontatrophia—a. Glandularis, *Tabes mesenterica*—a. Hepatis, Hepatrophia—a. Infantum, Pædatrophia, *Tabes mesenterica*—a. Intestini-rum, Enteratrophia.

ATROPHIA LACTANTII, *Ta'bes nutritum* seu *lacten*. The atrophy of nursing women.

ATROPHIA LIENIS, Splenatrophia—a. Mesenterica, *Tabes mesenterica*—a. Testiculi, Orchidatrophia—a. Unguium, Onychatrophia.

ATROPHICI (MORBI) Diseases of nutrition, characterized by atrophy and debility.

ATROPHIE, Atrophy—a. *des Dents*, Odontatrophia—a. *Mésentérique*, *Tabes mesenterica*.

ATROPHIE MUSCULAIRE PROGRESSIVE, *Paralysie musculaire atrophique*, *Atrophie musculaire primitive ou idiopathique ou avec transformation graisseuse*. *Cruveilhier's atrophy*, *Wasting Palsy*. A rare malady, in which the muscles become so greatly atrophied that they cannot perform their functions.

ATROPHIED, see Atrophy.

AT'ROPHY, (*τροφία*, from *a*, and *τροφή*, 'nourishment.') *Marasmus Atrophia*, *Atrophia Marasmus*, *Marantis*, *Ma'cies*, *Contabescencia*, *Ta'bes*, *Marcores*, *Analo'sis*, (F.) *Atrophie*, *Dessechement*. Defective nutrition,—*Hypot'rophy*. Progressive and morbid diminution in the bulk of the whole body or of a part. Atrophy is generally symptomatic. Any tissue or organ thus affected is said to be *atrophied*.

ATROPHY, CRUVEILHIER'S, *Atrophie musculaire progressive*—a. of the Heart, see Heart, atrophy of the.

ATROPIA, Atropine—a. Sulphate of, Atropiæ sulphas.

ATROP'LE SULPHAS, *Sulphate of Atropia*. Formed by dissolving atropia in dilute sulphuric acid; evaporating, and crystallizing. Virtues, the same as those of Atropia.

ATROPINE, *At'ropin*, *Atropina*, *Atropia* (Ph. U. S.), *Atropium*, *Atropinum*, *Belladonna*, (F.) *Atropine*. The active principle of *Atropa Belladonna* may be separated by a process similar to that for procuring morphia. A formula for its preparation has been introduced into the last edition of the Ph. U. S. (1863.) A single drop of a solution, of one grain in f $\frac{3}{4}$ iv of distilled water with a few drops of acetic acid, applied to the inner surface of the lower eyelid, causes dilatation of the pupil, in fifteen or twenty minutes. A grain to a drachm of lard is an application in neuralgia.

ATTACHE, Insertion.

ATTACK, *Invasio*, *Assultus*, *Irrep'tio*, *Invasio*, *Eis'bolé*, *Lep'sis*, *Access*, (Prov.) *Take*, (F.) *Attaque*, *Accès*. A sudden attack, invasion or onset of a disease. A seizure. One attacked or affected with severe disease is often said, in the United States, to be "taken down," or to be "down" with it.

ATTAGAS, (*arrayas*) Attagen.

ATTAGEN, (*arrayn*), *At'tagas*, the *Fran'colin*. Celebrated with the ancients both as food and medicine.

ATTANCOURT, MINERAL WATERS OF. A mineral water in France, at Attancourt, in Champagne; about three leagues north of Joinville. The water is a chalybeate, and contains sulphate of lime. In large doses it is purgative.

ATTAQUE, Attack—a. *des Nerfs*, Nervous attack.

ATTAR OF ROSES, see *Rosa centifolia*.

ATTELLE, Splint.

ATTENÖTING, MINERAL WATERS OF, in Bavaria. The water contains carbonic acid, carbonates of lime and soda, sulphates of lime and magnesia, chloride of sodium, iron and alum. It is much used in skin diseases, fistula, old ulcers, calculi, and hemorrhoids.

ATTENUANTS, *Attemuantia*, *Leptantica*, (F.) *Atténuants*, *Leptontiques*, (*at*, and *tenuis*, 'thin,') Medicines which augment the fluidity of the humours.

ATTENUATION, *Attenuatio*. Thinness, emaciation. A term used by the homœopathists, in the sense of dilution or division of remedies into infinitesimal doses.

ATTRANT, (*at*, and *tirer*, 'to draw,') Attraction.

ATTITUDE, *Si'tus Corporis*. (Low Latin, *aptitudo*; from *aptare*, 'to fit,') Situation, position of the body. The attitudes are the different postures which man is capable of assuming. In *General Pathology*, the attitude will often enable the physician to pronounce at once upon the character of a disease, or it will aid him materially in his judgment. In St. Vitus's dance, in fractures, luxations, &c., it is the great index. It will also indicate the degree of nervous or cerebral power; hence sinking down in bed is an evidence of great cerebral debility in fever. The position of a patient during an operation is also an interesting subject of attention to the surgeon.

ATTOLLENS AUREM, (*at*, and *tollo*, 'to raise,') *A. Auriculæ*, *Levator Auris*, *Superior Auris*, *Attollens Auriculam*, *Auricularis superior*, (F.) *Auricularis supérieure*. *Temporo-auriculare*. A muscle of the ear, which arises, thick, broad, and tendinous, from the tendon of the occipito-frontalis, and is inserted into the upper part of the ear, opposite to the anti-helix. It raises the ear.

ATTOLLENS OCULI, *Rectus superior oculi*—a. *Oculum*, *Rectus superior oculi*.

ATTOUCHEMENT, (*at*, and *toucher*, 'to handle,') Masturbation.

ATTRACTION OF AGGREGATION, Cohesion, force of.

ATTRACTIVUM, (*at*, and *trahere*, 'to draw,')

ATTRACTIVUS, Attrahent.

ATTRACTORIUS, Attrahent.

ATTRAHENS, Attrahent—a. *Auriculam*, *Anterior auris*.

ATTRAHENT, *Attrahens*, *Attractivus*, *Attractivus*, (F.) *Attractif*, *Attrahant*. Remedies are so called, which attract fluids to the parts to which they are applied, as blisters, rubefacients, &c.

ATTRAPE-LOURDAU, (F.) (*attraper*, 'to seize,') A bistoury invented by a French surgeon, called Biennaise, and used in the operation for hernia. See *Bistouri caché*.

ATTRITA, (*at*, and *terere*, *tritum*, 'to rub,') Chafing.

ATTRITIO, Attrition, Chafing.

ATTRITION, *Attritio*, *Ecthléma*. Friction or bruising. Chafing. Also, a kind of ecchymosis. Likewise, a violent contusion.

ATTRITUS, Chafing.

ATYPIC, *Atypicus*, *Atypas*, (F.) *Atypique*, (*a*, and *typos*, 'type,') That which has no type. Irregular. Chiefly applied to an irregular intermittent.—*Febris atypica*.

ATYPOS, (*atypas*) Erratic.

AU, as a prefix, see *A*.

AUANISIS, (from *avavos*, 'desiccation,') Drying.

AUAN'TE. Hippocrates gave this name to a disease, the principal symptom of which was emaciation. Atrophy.

AUBÉPINE, (*aube*, (L.) *albus*, 'white,' and 'pine,' spina, thorn,) *Mespilus oxyacantha*.

AUBÉVIGNE, (*aube*, 'white,' and *vigne*, 'vine,' *Clematis vitalba*.

AUBERGINE, *Solanum Melongena*.

AUBIFOIN, (*aube*, 'white,' and *foin*, *foenum*, hay, grass,) *Centaurea cyanus*, *Cyanus segetum*.

AUCHEN, (*avvyn*,) *Collum*.

AUCHENORRHEUMA, (*auchen*, and *rheuma*, 'rheumatism,') *Torticollis*.

AUCHE TICUS, (*avvyn*, 'the neck,') One affected with stiff neck or torticollis.

AUCTUMNUS, (*augere*, *auctum*, 'to increase,') Autumn.

AUDÉ, (*avén*,) Voice.

AUDINAC, MINERAL WATERS OF. Audinac is situate in the department of Arriège, France. The water contains a small quantity of sulphohydric acid, carbonic acid, sulphates of lime and magnesia, carbonates of lime and iron, and a bituminous substance. Temp. 67° Fah. It is much used in chronic rheumatism, herpes, serofulous diseases, &c. Season, from the 1st of June to the 1st of September.

AUDITIF, Auditory.

AUDITION, (*audire*, *auditus*, 'to hear;') *Auditio*, *Auditus*, *A'coë*, *Acoë'sis*, *Acu'sis*, (F.) *Ouïe*, Hearing. The act of hearing. The sensation arising from an impression made on the auditory nerves by the vibrations of the air, produced by a sonorous body. The physiology of Audition is obscure. It probably takes place:—

1. By the vibrations being communicated from the *membrana tympani* along the chain of small bones to the membrane of the *foramen ovale*. 2. By means of the air in the cavity of the *tympanum*, the membrane of the *foramen rotundum* is agitated. 3. The transmission may be made by means of the bony parietes. In these three ways the vibrations produced by a sonorous body may reach the auditory nerve. Audition may be *active* or *passive*: hence the difference between *listening* and simply *hearing*.

AUDITORY, *Auditorius*, *Auditivus*, *Acus-ticus*, (F.) *Auditif*. That which relates to audition.

AUDITORY ARTERIES AND VEINS, are vessels which enter the auditory canals, and are, like them, distinguished into *internal* and *external*. The *external auditory artery*, *A. Tympanique*—(Ch.), is given off by the *styloid*, a branch of the *external carotid*: the *internal* is a branch of the *basilary artery*, which accompanies the auditory nerve, and is distributed to it. The *Auditory Veins* empty into the *internal* and *external jugulars*.

AUDITORY CANAL, EXTERNAL, *Meatus auditivus externus*, *Porus acusticus externus*, *Alvearium*, *Sca'pha*, *Sca'phus*, (F.) *Conduit auditif externe*, *Conduit auriculaire*, commences at the bottom of the concha, at the *Foramen auditivum externum*, passes inwards, forwards, and a little downwards, and terminates at the *membrana tympani*. It is partly cartilaginous, partly osseous, and partly fibrous.

AUDITORY CANAL, INTERNAL, *Meatus auditivus internus*, *Porus seu S'itus acusticus*, *Cyl'ar*, (F.) *Conduit auditif interne*, *C. labyrinthique*, is situate on the posterior surface of the *pars petrosa* of the temporal bone. From the *Foramen auditivum internum*, where it commences, it passes forwards and outwards, and terminates by a kind of *cul-de-sac*, *macula cribrosa*, perforated by many holes, one of which is the orifice of the *Aqueductus Fallopii*; and the others communicate with the labyrinth.

AUDITORY NERVE, *Nervus auditivus seu acusticus*. *Nerv. labyrinthique*—(Ch.), is the *Portio Mollis* of the seventh pair. It arises from the *truncus restiforme*, from the floor of the fourth

ventricle, and by means of white *striae* from the sides of the *calamus scriptorius*. As it leaves the *encephalon*, it forms a flattened cord, and proceeds with the *facial nerve* through the *foramen auditivum internum*, and as far as the bottom of the meatus, where it separates from the *facial*, and divides into two branches, one going to the *cochlea*, the *coch'lear*; the other to the *vestibule* and *semi-circular canals*, the *vestib'ular*.

AUGÉ, *Al'teus*. Some of the older anatomists gave this name to a reservoir, into which liquids flow in an interrupted manner, so that it is alternately full and empty. Such are the *ventricles* and *auricles* of the heart.

AUGGERE, Intermittent Fever.

AUGMENTATION, (from *augere*, *auctum*, 'to increase;') *Augmentum*, *Incrementum*, *Anab'asis*, *Aug'tio*, *Auz'is*, *Progres'sio*, *Progres'sus*, *Anze'sis*, *Accre'tion*, (F.) *Augment*. The stage of a disease in which the symptoms go on increasing.

AULACOMELE, (*αυλαξ*, gen. *αυλακος*, 'a furrow,' and *μηλη*, 'a probe,') see *Grooved*.

AULISCUS, (*αυλισκος*, dim. of *Aulus*,) Canula. See *Fistula*.

AULNÉE, *Aunée*.

AULUS, (*αυλος*, 'a fistula or pipe,') *Canula*, *Fistula*. See *Foramen*, and *Vagina*.

AUMALE, MINERAL WATERS OF. Aumale is a town of Upper Normandy, in the country of Caux. Several springs of ferruginous mineral waters are found there, whose odour is penetrating, and taste rough and astringent. They are tonic, and employed in debility of the viscera, &c.

AUMURE, (*ai*, and *murus*, 'a wall,') *Parietaria*.

AUME NOIRE, (*almus*,) *Rhamnus frangula*.

AUNÉE, *Inula helenium*—*a. Dysentérique*, *Inula dysenterica*—*a. Officinale*, *Inula helenium*.

AURA, (*αυρα*,) *Pno'ë*, *Pno'a*, *Pnuus*. A vapour or emanation from any body, surrounding it like an atmosphere. Van Helmont regarded the vital principle as a gas and volatile spirit, which he called *Aur'a vitalis*.

In *Pathology*, *Aura* means the sensation of a light vapour, which, in some diseases, appears to set out from the trunk or limbs; and to rise towards the head. This feeling has been found to precede attacks of epilepsy and hysteria, and hence it has been called *Aur'a Epilep'tica*, and *A. hysterica*.

AURA EPILEPTICA, see *Aura*—*a. Hysterica*, see *Aura*.

AURA SANGUINIS. The odour exhaled by blood newly drawn. See *Gaz Sanguinis*.

AURA SEMINIS, *A. semina'lis*, *Spir'itus genita'lis*:—a volatile principle fancied to exist in the sperm, and regarded by some as the fecundating agent. Such is not the case.

AURA VITALIS, Vital principle.

AUR'AL, *Aura'lis*, (from *auris*, 'ear,') Relating or appertaining to the ear.

AURAL MEDICINE AND SURGERY, *Otiatria*.

AURANCUM, See *Ovum*.

AURANITE, see *Agaric*.

AURANTIA CURASSAVENTIA, (from *aurum*, 'gold' [?], *A. Curassav'ica*, *Curasso'a ap'ples* or *or'anges*. Immature oranges, checked, by accident, in their growth. They are a grateful, aromatic bitter, devoid of acidity. Infused in wine or brandy they make a good stomachic. They are also used for *issue peos*. See *Citrus aurantium*.

AURANTII CORTEX, see *Citrus aurantium*—*a. Flores*, see *Citrus aurantium*—*a. Floris Aqua*, see *Citrus aurantium*.

AURANTIUM, (from *aurum*, 'gold,' (?) owing to the color,) *Citrus aurantium*.

AURELIANA CANADENSIS, *Panax quinquefolium*.

AUREOLA, *Areola*.

AURI (gen. of *aurum*) **CHLORETUM CUM CHLORETO NATRII**, see Gold—a. Chloretum, Gold, muriate of—a. Chloridum, Gold, muriate of—a. Cyanidum, see Gold—a. Cyanuretum, see Gold—a. Iodidum, see Gold—a. Ioduretum, see Gold—a. Murias, Gold, muriate of—a. Nitro-murias, see Gold—a. Oxidum, see Gold—a. Terechloridum, see Gold—a. Tereyanidum, see Gold—a. Teroxidum, see Gold—a. et Natri chloruretum, see Gold.

AURICHALCUM, (*aurum*, and *χαλκός*, 'copper,') Brass.

AURICLE, *Auric'ula*, (F.) *Auricule*, *Oricule*, (dim. of *auris*.) The pavilion of the ear. See Pavilion.

AURICLES OF THE HEART, *Auric'ulæ*, *A'tria*, *Cavita'tes innomina'tæ*, (F.) *Oreillettes*, are two cavities, one right, the other left, each communicating with the ventricle of its side. These two cavities receive the blood from every part of the body. Into the *right auricle*, the two *venæ cavæ* and coronary vein open: into the *left*, the four pulmonary veins. Chaussier calls the former the *Sinus of the Ve'næ Cavæ*, *Sinus dex'ter cor'dis*: the latter, the *Sinus of the Pul'monary Veins*, *Sinus sinis'ter cor'dis*. The foliated or dog's ear portion of each auricle is called *Appen'dix Auric'ulæ*. See Sinus.

AURICULA, *Auricle*—a. Judæ, *Peziza auricula*—a. *Muris*, *Hieracium Pilosella*—a. *Muris major*, *Hieracium murorum*.

AURICULAIRE, *Auricular*, see *Digitus*—a. *Postérieur*, *Retrahens auris*—a. *Supérieur*, *Attollens aurem*.

AURICULAR, *Auricula'ris*, *Ori'cular*, (F.) *Auriculaire*. That which belongs to the ear, especially to the external ear.

AURICULAR ARTERIES AND VEINS, *Oreil-laïres*—(Ch.), are divided into *anterior* and *posterior*. The *anterior* are of indeterminate number. They arise from the temporal artery and are distributed to the *meatus auditorius externus*, and to the pavilion of the ear. The *posterior auricular* is given off by the external carotid, from which it separates in the substance of the parotid gland. When it reaches the inferior part of the pavilion of the ear it bifurcates; one of its branches being distributed to the inner surface of the pavilion, the other passing over the mastoid process, and being distributed to the temporal and posterior *auris muscels*, &c. Before its bifurcation it gives off the *stylo-mastoid artery*. The *Anterior and Posterior Auricular Veins* open into the temporal and external jugular.

AURICULAR FIN'GER, (F.) *Doigt auriculaire*, is the little finger, so called because, owing to its size, it can be more readily introduced into the *meatus auditorius*.

AURICULAR NERVES are several. 1. The *auricular branch*, *Zygo'mato-auric'ular*, is one of the ascending branches of the cervical plexus. It ramifies and spreads over the two surfaces of the pavilion. 2. The *auricular or superficial temporal*, *Temporal-cutaneous*—(Ch.), is given off from the inferior maxillary. It ascends between the condyle of the jaw and the *meatus auditorius externus*, sends numerous filaments to the *meatus* and pavilion, and divides into two twigs, which accompany the branches of the temporal artery, and are distributed to the integuments of the head. There is also a *posterior auricular* furnished by the facial.

AURICULARIA SAMBUCI, *Peziza auricula*.

AURICULARIS ANTERIOR, *Anterior auris*—a. *Superior*, *Attollens aurem*.

AURICULE, *Auricle*, *Pavilion of the ear*.

AURICULOVENTRICULAR, *Atrioventric'ular*, *Auriculoventricula'ris*, *Auriculoventricula'ris*, (F.) *Auriculo-ventriculaire*. That which belongs to the auricles and ventricles of the heart. The

communications between the auricles and ventricles are so called. The *Tricuspid and Mitral Valves* are auriculo-ventricular valves.

AURIGA. A species of bandage for the ribs, described by Galen. See, also, *Liver*.

AURIGO, gen. *Aurig'inis*, (*aurum*, 'gold;') from its color,) *Icterus*—a. *Neophytorum*, *Icterus Infantum*.

AURIPIGMENTUM, (*aurum*, and *pigmentum*, 'paint,') *Orpiment*—a. *Rubrum*, *Realgar*.

AURIS, *Ear*.

AURISCALPIUM, (*auris*, and *scalpere*, 'to scrape,') *Earpick*.

AURISCOPE, *Auriscop'ium*, (*auris*, and *σκοπεω*, 'I view,') An instrument for exploring the ear.

AURIST, *Otia'ter*, *Otia'trus*, *Ear-doctor*, *Ear-surgeon*; (from *auris*.) One who occupies himself chiefly with the diseases of the ear and their treatment.

AURIUM FLUCTUATIO, *Bombus*—a. *Marmorata*, *Cerumen*—a. *Sibilus*, *Bombus*—a. *Sonitus*, *Bombus*—a. *Sordes*, *Cerumen*—a. *Susurrus*, *Bombus*.

AURONE (corr. from *Abrotanum*), *Artemisia abrotanum*—a. *des Champs*, *Artemisia campestris*—a. *des Jardins*, *Artemisia abrotanum*—a. *Mâle*, *Artemisia abrotanum*.

AURUGO, gen. *Aurug'inis*, (from *aurum*, 'gold,') from its color,) *Icterus*.

AURUM, (*αυρον*.) *Gold*—a. *Chloratum*, *Gold*, *muriate of*—a. *Chloratum natronatum*, see *Gold*—a. *Foliatum*, *Gold leaf*—a. *Leprosum*, *Antimonium*—a. in *Libellis*, *Gold leaf*—a. *Limatum*, see *Gold*—a. *Muriaticum*, see *Gold*—a. *Muriaticum natronatum*, see *Gold*.

AURUM MUSIVUM, *Aurum Mosai'cum*, *Sulph'uret of Tin*, *Deuto'sulph'uret* or *Persulphuret of tin*. (*Quicksilver*, tin, sulphur, sal ammoniac, &c., equal parts.) The tin being first melted, the quicksilver is poured into it, and then the whole are ground together, and sublimed in a bolthead. The *aurum musivum* lies at the bottom.) It is used in some empirical preparations.

AURUM OXYDATUM, see *Gold*—a. *Oxydulatum muriaticum*, *Gold*, *muriate of*—a. *Nitro-muriaticum*, see *Gold*—a. *Salitum*, *Gold*, *muriate of*.

AUSCULTATE, (*auscultare*, *auscultatum*, 'to listen,') To practise auscultation. 'To auscult' is at times used with the same signification.

AUSCULTATION, *Ausculta'tio*, *Echos'copé*, act of listening. Buisson has used it synonymously with *list'ening*. Laënnec revived *auscultation* to appreciate the different sounds which can be heard in the chest, and in the diagnosis of diseases of the heart, lungs, &c. This may be done by the aid of an instrument called a *stethoscope*, one extremity of which is applied to the ear, the other to the chest of the patient. This mode of examination is called *Me'diate Auscultation*, (F.) *Auscultation médiante*,—the application of the ear to the chest being *Imme'diate Auscultation*.

The act of exploring the chest physically is called *Stethoscop'ia*, and *Thoracoscop'ia*, and, in popular parlance, *Sounding the chest*; of the abdomen, *Abdominoscop'ia*.

AUSCULTATORY, *Auscultato'rius*; *Auscultatory*, *Auscult'ic* (with some.) Belonging or having relation to auscultation.

AUSCULTATORY PERCUSSION, see *Acouphonia*.

AUSTERE, *Auste'rus*. Substances which produce a high degree of acerb impression on the organs of taste.

AUSTRUCHE, (from *Astruthum*,) *Imperatoria*.

AUTALGIA DOLOROSA, (*auto*, and *αλγος*, 'pain,') *Neuralgia*, *facial*, *Pleurodynia*—a. *Pruriginosa*, *Itching*—a. *Vertigo*, *Vertigo*.

AUTARCIA, (*αυτάρκεια*, from *auto*, and *αρκεω*, 'I am satisfied,') *Moral tranquillity*.

AUTEMES'IA, (*auto*, and *emesis*, 'vomiting.') Spontaneous or idiopathic vomiting.

AUTEMPRESMUS, (*auto*, and *εμπρησμος*, 'the art of burning.') Combustion, human.

AUTHIMERON, (*αυθημερον*, from *auto*, and *ημερα*, 'day.') A medicine which cures on the day of its exhibition.

AUTHYGLANSIS, (*auto*, and *υγιαζω*, 'I heal or cure.') *Vis medicatrix naturæ*.

AUTO, (from *αυτος*, 'self,') as a prefix, self.

AUTOCHIR, (*αυτοχειρ*, from *auto*, and *χειρ*, 'hand,') *Autochirus*, *Suicida*. One who has committed suicide. A self-murderer or suicide.

AUTOCHIRIA, (*αυτοχειρια*,) Suicide.

AUTOCINE'SIS, (*Μο'tus voluntarius*, (*αυτοκινησις*, from *auto*, and *κινησις*, 'motion,') Voluntary motion.

AUTOC'RASY, *Autocrati'a*, *Autocrato'ria*, (*auto*, and *κρατος*, 'strength,') Independent force. Action of the vital principle, or of the instinctive powers, towards the preservation of the individual. See *Vis Medicatrix Naturæ*. Also, the vital principle.

AUTOCRATIA, Autocracy, *Vis Medicatrix Naturæ*.

AUTOCRATORIA, (*αυτοκρατορια*,) Autocracy—
a. *Physiatrix*. *Vis medicatrix naturæ*.

AUTOCTONIA, (*auto*, and *κτος*, 'murder,') Suicide.

AUTOGENIA, (*auto*, and *γεννω*, 'to engender,') Generation, spontaneous.

AUTOG'ENOUS, (same etymon.) Relating or appertaining to spontaneous generation. Applied by Mr. Owen to parts or elements that are usually developed from distinct and independent centres; as in the case of the different parts or elements that form a vertebra.

AUTOLITHOTOMUS, (*auto*, *λιθος*, 'a stone,' and *τεμνειν*, 'to cut,') One who operates upon himself for the stone.

AUTOMATIC, *Automat'icus*, *Autom'atus*, (F.) *Automatique*, (from *αυτοματος*, 'spontaneous,') That which acts of itself. Those movements are called automatic which are executed without volition:—involuntary motions, *motus automatici* seu *autom'ati* seu *involunta'rii*.

AUTOMNAL, Autumnal.

AUTONOMIA, (*αυτονομια*, from *auto*, and *νομος*, 'law,') *Vis medicatrix natu'ræ*. The word *Autonomia* is occasionally employed by the French and Germans for the peculiar mechanism of an organized body. Thus, although individuals of the same species may differ in outward conformation, their mechanism or instinctive laws (*Autonomia*) may be the same.

AUTONYCTOBATIA, (*auto*, *νυξ*, gen. *νυκτος*, 'night,' and *βαρυν*, 'to go,') Somnambulism.

AUTOPEP'SIA, (*auto*, and *πεπρω*, 'I concoct,') Self-digestion,—as of the stomach after death.

AUTOPHIA, Autopsia.

AUTOPHONIA, Autophony; also, Suicide, (*auto*, and *φονος*, 'murder,')

AUTOPH'ONY, *Autopho'nia*, (F.) *Autophonie*, *Retentissement autophonique*, (*auto*, and *φωνη*, 'voice,') An auscultatory sign, which consists in noting the character of the observer's own voice, while he speaks with his head placed close to the patient's chest. The voice, it is alleged, will be modified by the condition of the subjacent organs. The resonance, thus heard, is termed *retentissement autophonique*. This diagnostic method Dr. R. G. Latham proposes to term *heautophon'ica*.

AUTOPHOSPHORUS, Phosphorus.

AUTOPLAS'TIC, *Autoplas'ticus*, (*auto*, and *πλαστικός*, 'formative,') Relating to autoplasty or plastic surgery.

AUTOPLASTICE, Morioplastice.

AUTOPLASTY, Morioplastice.

AUTOP'SIA, *Aut'opsy*, (*αυτοψια*, from *auto*, and *ψις*, 'vision,') *Autoph'ia*, *Autoscop'ia*. Inspec-

tion; examination by one's self: self-inspection. Often improperly used for the following:

AUTOP'SIA CADAVER'ICA, (F.) *Autopsie on Ouverture cadavérique*. Attentive examination after death.—*Examination post mortem*, *Sec'tio Cadaver'ica*, *Dissac'tion*, *Necropsopy*, *Necropsy*, *Necropsop'ia*, *Necrop'sia*, *Necrop'sis*,—practised for the purpose of investigating the causes and seat of an affection of which a person may have died, &c.

AUTOP'SIA CADAVER'ICA LEGA'LIS, *Sec'tio seu Abduc'tio lega'lis*, is the examination after death for medico-legal purposes.

AUTOPSY, Autopsia.

AUTOPYROS, (*αυτοπυρος*, from *auto*, and *πυρος*, 'wheat,') Syncomists.

AUTOSCOPIA, (*auto*, and *σκοπεω*, 'I view,') Autopsia.

AUTOSITE, (*auto*, and *σιτος*, 'nourishment,') A single monster, capable of deriving nourishment from its own proper organs, in contradistinction to *Omph'alosite*.

AUTOTHERAPIA, (*auto*, and *θεραπεω*, 'I cure,') *Vis medicatrix naturæ*.

AUTUMN, *Autum'nus*, *Autum'nus*, from (*augere*, *auctum*, 'to increase' [?]), (*Opo'ra*, early part of autumn, *Pthinopo'ra*, end of autumn.) (F.) *Automne*. One of the seasons of the year, between the 23d of September and the 21st of December. In all climates, the Autumn or Fall is liable to disease; a combination of local and atmospheric causes being then present, favourable to its production.

AUTUM'NAL, *Autumna'lis*, (F.) *Autumnal*. Relating to Autumn; as *Autumnal Fruits*, *Autumnal Fevers*, &c.

AUTUMNAL FE'VER generally assumes a bilious aspect. Those of the intermittent kind are much more obstinate than when they appear in the spring.

AUXESIS, (*αυξισ*,) Augmentation, Increase.

AUXIL'IARY, *Auxilia'ris*, (from *auxilium*, 'aid,') (F.) *Auxiliaire*. That which assists, or from which assistance is obtained.

AUXILIARY MEDICINE is one which assists the principal medicine or basis. It is synonymous with Adjuvant.

AUXILIARY MUSCLES are those which concur in the same movement. Some anatomists have applied the term to several ligaments, as well as to the fleshy fibres, which hang from the *sacro-spinalis* muscle.

AUXILIUM, ('aid,') Juvans, Medicament.

AUXIS, (*αυξισ*,) Augmentation, Increase.

A'VA, *Ar'va*, *Ka'va*. An intoxicating narcotic drink, made by chewing the *Pi'per methys'ticum*. It is much used by the Polynesians.

AVAILLES, MIN'ERAL WA'TERS OF. A small village in France, 13 leagues S. S. E. of Poitiers, at which there is a cold saline chalybeate. It contains chlorides of sodium and calcium, sulphate and subcarbonate of soda, iron, &c.

AVANT (F.), in composition, Ante.

AVANT-BOUCHE (F.), ('fore-mouth,') *Os anti'cum*. This name has been applied, by some, to the mouth, properly so called—in contradistinction to the *Arrière bouche* or *Pharynx*. With others, it means the part of the mouth between the lips and teeth.

AVANT-BRAS, Fore-arm.

AVANT-CŒUR, ('fore-heart,') Scrobiculus cordis.

AVANT-GOUT (F.), *Prægesta'tio*. A fore-taste; prægnation.

AVANT-MAIN (F.), ('fore-hand,') *Adrer'ea manus*. The inside of the hand, when extended.

AVANT-PIED (F.), ('fore-foot,') The most advanced part of the foot.

AVANT-POIGNET (F.), ('fore-wrist,') The anterior part of the wrist.

A VELINE, *Corylus avellana*, (nut.)

AVELLANA, *Corylus avellana*.—a. Cathartica, *Jatropha curcas*.

AVENA, *Bro'mos*, Oats, (Prov., North of England) *Hav'er*, *Aits*. The seeds of *Ave'na sati'va*. Ord. Gramineæ. *Sex. Syst.* Triandria Digynia (F.) *Avoine*. Oats are used as food for man in some parts, particularly in the North of England and Scotland. When deprived of the husks, they form *Groats*. Reduced to meal—*Ave'næ Fari'na*, *Oat'meal*, (Ph. U. S.)—they are applied as cataplasms to promote suppuration. The dry meal is sprinkled over erysipelatous parts.

Oat'meal gruel, *Wa'ter gru'el*, is prepared as follows:—Take of *oatmeal*, $\frac{3}{4}$ ij; *soft water*, Oiss. Rub the meal in a basin, with the back of a spoon, in a moderate quantity of the water, pouring off the fluid after the grosser particles have subsided, but whilst the milkiness continues; and repeat the operation until no more milkiness is communicated to the water. Put the washings in a pan, after having stirred them well, in order to suspend any fecula which may have subsided; and boil until a soft, thick mucilage is formed. See *Grit gruel*.

It is a good demulcent, and is used also as a vehicle for clysters.

A thick oatmeal gruel, or hulled oats boiled,—a dish made at sea—is called *Burgoo*, *Burgout*, *Bargout*, and *Loblolly*.

AVENA EXCORTICATA, Groats.

AVENÆ FARINA, see Avena.

AVENHEIM, MINERAL WATERS OF. Avenheim is three leagues from Strasburg: near it is an aperient mineral water.

AVENNES or AVÈNE, MINERAL WATERS OF. Avennes is a village in the department of Hérault, in France: near it is a saline spring, the temperature of which rises to 84° Fahrenheit.

AVENS, COMMON, *Geum urbanum*.—a. Purple, *Geum rivale*.—a. Water, *Geum rivale*.—a. White, *Geum Virginianum*.—a. Yellow, *Geum urbanum*.

AVERICH, Sulphur.

AVERRHO'A BILIM'BI, (after *Averrhoes*), *Bilim'bi*, *Bilimbing teres*. Ord. Oxalidæ. An Indian tree, which has a fruit that is too acid to be eaten alone. It is used as a condiment, and in the form of syrup as a refrigerant.

AVERRHO'A CARAM'BOLA, *Mal'um Coë'n'sê*, *Prunum stella'tum*, *Tam'ara*, *Con'ga*, *Caram'bolo*. An Indian tree, whose fruits are agreeably acid. The bark, bruised, is employed as a cataplasm, and its fruit is used as a refrigerant in bilious fever and dysentery.

AVER'SION, *Aver'sio*, *Apot'ropê*, (*avertere*, *avertum*, (a, and *vertere*), 'to turn from.') Extreme repugnance for anything whatever.

AVERSION (F.) also means, in therapeutics, the action of medicines which turn the afflux of fluids from one organ, and direct them to others; being synonymous with *counter-irritation*, or rather, with *revulsion* or *derivation*.

AVERTIN (F.), (same etymon, or from *vertigo*, 'giddiness.') A disease of the mind, which, according to Lavoisier, renders the patient obstinate and furious.

AVEUGLE, (*ab*, and *oculus*), Cæcus.

AVEUGLEMENT, Cæcitas—a. *de Jour*, Nyctalopia—a. *de Nuit*, Hemeralopia.

AVICEN'NIA TOMENTO'SA, *A. Africa'na* seu *resinif'era* seu *nit'ida*, *Bont'ia ger'minans*: (after *Avicenna*.) Ord. Verbenacæ. The plant which affords the *Malac'ca Bean* or *Anacar'dium Orienta'lè* of the Pharmacopœias, *Semecarpus Anacar'dium*. The oil drawn from the bark of the fruit is corrosive and an active vesicatory, but it is not used.

AVICULA CYPRIA, *Pastil*.—a. *Margaritifera*, see Pearl.

AVOIN, Avena.

AVON, MINERAL WATERS OF. Avon Springs are in the State of New York, on the eastern branch of the Genesee river, 18 miles from Rochester. There are three sulphureo-saline and one iodine spring. They are applicable to the same set of cases as the Sharon Springs.

AVORTMENT, Abortion—a. *Provoqué*, see Parturition.

AVORTER, to Abort.

AVORTIN, Abortion.

AVORTON, Abortion.

AVULSIO, (a, and *vellere*, *vulsum*, 'to pluck,') *Arrachement*.

AVULSION, Evulsion.

AX, MINERAL WATERS OF. Ax is a small town in the department of Arriège, France; where there are several sulphurous springs, the temperature of which varies from 77° to 162° of Fahrenheit. The season is from the first of May to October.

AXE, Axis—a. *de l'Œil*, Axis of the eye.

AX'EA COMMISSU'RA, *Trochoi'des*. A pivot-joint. See Trochoid.

Axen Körper, Corpuscula tactūs.

AXES, Intermittent, Paroxysm.

AXILE BODIES, Corpuscula tactūs.

AXIL'LA, (dim. of *ala*), *A'la*, *Asc'ella*, *Aessel'la*, *Ascil'la*, *Acel'la*, *Cor'dis emuncto'rium*, *Ma'te*, *Hypo'mia*, *Po'vea axilla'ris*, *Mas'chalè*, *Mas'chalis*, (Sc. and Prov.) *Oxter*, *Oxter*, (F.) *Aisselle*. The cavity beneath the junction of the arm with the shoulder; the *armpit*, *Vallis axilla'rum*; (F.) *Creux de l'Aisselle*. It is bounded, anteriorly, by a portion of the pectoralis major; posteriorly, by the latissimus dorsi. It is covered with hair, *pili subaxilla'res*, contains much areolar membrane, lymphatic ganglions, important vessels and nerves, and numerous sebaceous follicles, furnishing an odorous secretion. In consequence of such secretion, the ancients called it *emuncto'rium cordis*.

AX'ILLARY, *Axilla'ris*, *Subala'ris*, *Maschali'us*, (F.) *Axillaire*. Belonging to the armpit.

AXILLARY ARTERY, *Arte'ria axilla'ris*; a continuation of the subclavian, extending from the passage of the latter between the scaleni muscles as far as the insertion of the pectoralis major, when it takes the name of *Brachial*.

AXILLARY GLANDS are lymphatic glands seated in the armpit; into which the lymphatic glands of the upper extremity open.

AXILLARY NERVE, *Circumflex N.*, *Artic'ular nerve*, *Scapulo-hu'meral* (Ch.), *Nerf Circonflexe*; arises from the posterior part of the brachial plexus, particularly from the last two cervical pairs and the first dorsal. It is chiefly distributed to the posterior margin of the deltoid.

AXILLARY VEIN, *Ve'na Axilla'ris*, *Ve'na Subala'ris*. This vein corresponds with the artery; anterior to which it is situate. It is a continuation of the *brachial veins*; and, at its termination assumes the name *Subclavian*.

AXINE, (*αξινς*) *Ascia*.

AXIRNACH. An Arabic word, used by Albucasis to designate a fatty tumour of the upper eyelid observed particularly in children.

AX'IS, *Ax'on* ('an axle'), (F.) *Axe*. A right line which passes through the centre of a body.

AXIS, CEREBRO-SPINAL, see Encephalon—a. of the Cochlea, Modiolus—a. Cranio-Spinal, see Encephalon—a. *Cœliac*, *Cœliac artery*—a. Cylinder of Nerve, see Nerve fibre.

AXIS OF THE EYE, (F.) *Axe de l'œil*, called also, *Vis'ual Axis* and *Op'tic Axis*, is a right line, which falls perpendicularly on the eye, and passes through the centre of the pupil.

AXIS, HEMAL, Aorta—a. Neural, see Encephalon.

Axis, is also the second vertebra of the neck,

Az'on, Epistroph'eus, Epist'rophus, Maschalister: Vertebra Denta'ta, (F.) Essieu. So called, because it forms a kind of axis on which the head moves. Chaussier calls it *Axoïde*, (from *αξων*, 'axis,' and *ειδος*, 'shape,') It is distinguished by its *odontoid process, Proces'sus odontoid'es seu denta'tus, Dens, Dens Epistroph'eï*, which projects from the anterior and superior part of the body.

AXLETEETH, Molar teeth.

AXOÏDE, Axis.

AXOÏDO, in composition, *Axis*, (vertebra.)

AXOÏDO-ATLOÏDEUS. What refers to both the axis and atlas, as *Axoïdo-atloïdean* articulation.

The lesions of the Axoïdo-atloïdean, are, 1. Fracture of the *Proces'sus Denta'tus*. 2. Rupture of the odontoid ligament, and consequently passage and pressure of the process behind the transverse ligament: and, 3. The simultaneous rupture of the odontoid and transverse ligaments. These different accidents are fatal.

AXOÏDO-ATLOÏDIEN, Obliquus inferior capitis—*a. Occipitale, Rectus capitis posticus major.*

AXON, (αξων), Axis.

AXONGE, (axis, 'axle,' and ungere, 'to grease,') Adeps præparatus.

AXUNGE, Adeps præparatus.

AXUNGIA, (axis, 'axle,' and ungere, 'to grease,') Adeps præparatus, Pinguedo—*a. Articularis, Synovia*—*a. Gadi, Oleum Jecoris Aselli*—*a. de Mumiâ, Marrow*—*a. Piscina Marina, Oleum Jecoris Aselli*—*a. Porcina, Adeps præparatus.*

AYPNIA, (α, and ύπνος, 'sleep,') Insomnia.

AZADIRACHTA INDICA, see *Melia Azedarach*.

AZALEA PROCUMBENS, (from *αζαλος*, 'arid,' from the soil it prefers.) *Loiseleuria procumbens.*

AZARNET, Orpiment.

AZARUM, Asarum—*a. Cabaret, Asarum.*

AZEDARACH, Melia Azedarach.

AZEDARACHA AMËNA, Melia Azedarach.

AZOÏC, Azo'icus, (same etymon as Azote.) Devoid of life. The "*Azoic period*" of the geologist is that before any living being appeared.

AZOODYNAMIA, (α, ζων, 'life,' and δυναμις, 'strength,') Privation or diminution of the vital powers.

AZO'RES, CLIMATE OF. The Azores or Western Islands are said to afford one of the best examples of a mild, humid, equable climate, to be met with in the northern hemisphere. It is slightly colder and moister than that of Madeira, but even more equable. Sir James Clark thinks, that a change from the Azores to Madeira, and thence to Teneriffe—one of the Canaries—would prove more beneficial to the phthisical valetudinarian than a residence during the whole winter in any one of those islands.

AZOTAS, Nitrate—*a. Hydrargyricus liquidus*, see *Hydrargyri nitras*—*a. Hydrargyroso-ammoniacus*, see *Hydrargyri oxydum cinereum*—*a. Hydrargyrosus, Hydrargyri nitras*—*a. Potassicus, Potassæ nitras.*

AZOTATE, Nitrate—*a. d'Argent, Argenti nitras*—*a. de Mercure, Hydrargyri nitras*—*a. de Mercure et d'ammoniaque*, see *Hydrargyri oxydum cinereum*—*a. de Plomb, Plumbi nitras*—*a. de Potasse, Potassæ nitras.*

AZOTE, Azo'tum, (α, and ζων, 'life,') Nitro-

gen, Al'kaligene, Gas azo'ticum, Nitrogen'ium, Sep'ton, (F.) Azote, Nitrogène, Mofette, Air gaté, Air vicié, is a gas which is unfit for respiration. It is not positively deleterious, but proves fatal owing to the want of oxygen. It is one of the constituents of atmospheric air, and a distinguishing principle of animals. Vegetables have it not generally diffused, whilst it is met with in most animal substances. It has been variously called, *phlogistic air, vitiated air, &c.*; has been looked upon as sedative, and recommended to be respired, when properly diluted, in diseases of the chest.

AZOTE, PROTOXIDE OF, Nitrogen, gaseous oxide of.

AZOTÉ, Nitrogenized.

AZOTED, Nitrogenized.

AZOTENESES, (azote, and νοσς, 'disease,') Diseases fancied by Baumes to be occasioned by the predominance of azote in the body.

AZOTIC ACID, Nitric acid.

AZOTIZED, Nitrogenized.

AZOTUM, Azote.

AZOTURIA, (azote, and ουρον, 'urine,') See Urine.

AZTEC CHILDREN, see *Nanus*.

AZUR, Coral, Smalt.

AZURIUM. A compound of two parts of mercury, one-third of sulphur, and one-fourth of sal ammoniac.—*Albertus Magnus.*

AZYGES, (αζυγος, from α, and ζυγος, 'a yoke,') *Azygos, Azygous, si'nè pa'ri*, 'not paired.' The *ephenoid bone*, because it has no fellow. Also, a process, *Proces'sus Azyges, Rostrum ephenoida'te*, projecting from under the middle and forepart of that bone.

AZYGOS UVULÆ, Azygous muscle.

AZYGOUS, see *Azyges*—*a. Articular artery*, see *Articular arteries of the knee*—*a. Ganglion*, see *Trisplanchnic Nerve*.

Azygous MUSCLE, Azygus Uvulæ, is the small muscle which occupies the substance of the uvula.—*Morgagni.* The name is, however, inappropriate, as there are two distinct fasciculi, placed along-side each other, forming the *Pal'ato-staphyli'ni, Staphyli'ni* or *Epistaphyli'ni* muscles, *Staphyli'ni me'dii* of Winslow.

AZYGOUS PROCESS OF THE SPHENOID, see *Azyges*.

AZYGOUS VEIN, Ve'na Azygos, Veine Prélobothoracique—(Ch.), *Ve'na si'nè pa'ri, Ve'na pa'ri ca'rens, (F.) Veins sans Paire.* This vein was so called by Galen. It forms a communication between the *V. cava inferior* and *V. cava superior*, permitting the blood to pass freely between the two. It rises from the *vena cava inferior*, or from one of the lumbar or renal veins, passes through the diaphragm, ascends along the spine to the right of the aorta and thoracic duct, and opens into the *V. cava superior*, where it penetrates the pericardium. On the left side, the *SEMI-AZYGOS, Left bron'chial or left superior intercostal vein, Veina demi-azygos, V. hemi-azyga, Veine petite prélobothoracique*—(Ch.), presents, in miniature, nearly the same arrangement.

AZYMIA HUMORUM, (α, and ζευω, 'leaven,') Crudity of the humours.

AZYMUS, (αζυμος.) Azymous bread is unfermented, unleavened bread.

AZZLE-TEETH, Molar teeth.

B.

BABEURRE, (*battre*, 'to beat,' and *beurre*, 'butter.') Buttermilk.

BABILLEMENT, (from *ba, ba*, the earliest infantile attempts at talking, — babbling.) Loquacity.

BABUICARIUS, Incubus.

BABY, (from *βαβύς*, 'to speak inarticulately.') Infans.

BACCÆ (pl. of *bacca*, 'a berry') **BERMUDENSES**, *Sapindus saponaria* — b. seu *Grana Actes*, see *Sambucus ebulus* — b. *Jujubæ*, *Jujube* — b. *Myrtillorum*, see *Vaccinium myrtillus* — b. *Norlandicæ*, *Rubus arcticus* — b. *Piperis glabri*, see *Piper Cubeba* — b. *Piscatoria*, see *Menispermum cocculus* — b. *Zizyphi*, see *Jujube*.

BAC'CAR, *Bac'caris*, *Bac'charis*, *Baccap*, *Bac'kapis*, *Bac'kapis*. An herb used by the ancients in their garlands, to destroy enchantment. By some considered to be *Nardum rusticum*; by others, *Valeriana Celtica*; by others, *Digitalis purpurea*; by others, *Asarum*; by others, a species of *Conyza*; by others, *Inula dysenterica*; and, by others again, *Gnaphalium sanguineum*. An oil was expressed from the root *Βακκαριον μρον*. *Bac'charis* is the name given by Galen not only to the root, but to a *Lydian ointment*, which was sometimes employed in diseases of the womb.

BACCHARIS, (*Bac'kapis*), *Baccar*.

BAC'CHARIS HALIMIFOLIA, *Ground'sel tree*; *Order*, *Compositæ*; indigenous; is used as a demulcent to allay cough, in the form of decoction.

BACCHI'A, (from *Bacchus*, 'the god of wine.') A name applied to the red or pimpled face of the drunkard. See *Gutta rosea*.

BACCHICA, (sacred to *Βαχχος*, 'Bacchus,') *Hedera helix*.

BACCIV'OROUS, *Bacciv'orus*, (F.) *Baccivore*, (*bacca*, 'a berry,' and *voro*, 'I devour.') Living on berries.

BACHELOR'S BUTTONS, see *Strychnos nuxvomica*.

BACHER'S TONIC PILLS, *Pilulæ ex Helleboro et Myrrhâ*.

BACILE, (*Bacillum*), *Crithmum maritimum*.

BACILLAR LAYER OF THE RETINA, *Tunica Jacobi*.

BACIL'LUM, *Bacil'lus*, *Bac'ulus*, *Bac'culus*: 'a stick or staff.' This name has been applied to a kind of troch, composed of expectorants, and having the shape of a stick. Also, a suppository. *Bacillum* was used by the ancient chemists for several instruments of iron. See *Tunica Jacobi*.

BACK, (Sax. *baec*.) *Dorsum*.

BACK-ACHE ROOT, *Liatris*.

BACKBONE, *Vertebral column*.

BACKSIDE, *Nates*.

BACKSPRENT, *Vertebral column*.

BACKSTROKE OF THE HEART, *Impulse, diastolic*.

BACOVE, *Musa sapientum*.

BACTYRILOBIUM, (dim. of *Βακτρον*, 'a cane,' and *καβαν*, 'a pod.') *Cassia fistula*.

BACULUS, *Bacillum*.

BAD, Sick.

BAD'EN, MIN'ERAL WA'TERS OF. *Baden* is a town six miles from Vienna. Here are 12 springs, containing carbonates of lime and magnesia; sulphates of lime, and magnesia, and soda; and chlorides of sodium and aluminum. The water is used in diseases of the skin, rheumatism, &c. The season is from the 1st of July to September.

There are two other towns of the same name; one in *Swabia*, and the other in *Switzerland*, about 12 miles from *Zürich*, where are mineral springs.

Season from the 15th of May to the 15th of September. The waters of the last two are thermal sulphureous.

BAD'EN-BA'DEN, MIN'ERAL WA'TERS OF. Celebrated thermal springs, situate about a league from the high road to *Basle* and *Frankfort*. Their temperature varies from 130° to 154° Fahrenheit. Their situation is beautiful, and they are much frequented. Season from the 1st of June to the 15th of September.

BADENWEILER, MIN'ERAL WA'TERS OF. *B.* is in *Baden*, about a league to the westward of *Müllheim*. The waters are thermal (87° of Fah.), and contain carbonate of lime, sulphuret and chloride of calcium, and chloride of magnesium.

BADER, *Bather*.

BADIA'GA. A kind of sponge, sold in *Russia*, the powder of which is said to take away the livid marks from blows and bruises in a few hours. Its nature is not understood.

BADIANE, *Illicium anisatum*.

BADISIS, (*Βαδισ*), *Walking*.

BADLY, Sick.

BADUKKA, *Capparis badukka*.

BÆOBOTRYS PICTA, (*Βαος*, 'small, and *βορπυς*, 'a grape,') *Saoria*.

BADULAM. The *Ardisia humilis* of *Ceylon*: a small tree whose fruit, made into syrup, is used as a cooling drink.

BAEL, *Covolam*.

BAG, see *Mamma* — b. *Dusting*, see *Dusting-bag*.

BAG OF WATERS, see *Liquor Amnii*.

BAGEDIA, *Pound*.

BAGGIE, *Abdomen*.

BAGNÈRES-ADOUR, OR **BAGNÈRES DE BIGORRE**, MIN'ERAL WA'TERS OF. *Bagnères de Bigorre* is a small town in the department of *Hautes Pyrénées*, having a great number of mineral springs; some, cold chalybeates; others, thermal salines; but the greatest part sulphureous and warm.

BAGNÈRES DE LUCHON, OR **LUCHON**, is a small town in the department of *Haute Garonne*, on the frontiers of *Spain*. It has been for a long time famous for its numerous sulphureous springs, the temperature of which is from 69° to 148° of Fahrenheit. They are visited from May to October.

BAGNIGGE (pron. *Bag'nidge*) **WELLS**. A saline mineral spring of *London*, resembling the *Epsom*.

BAGNIO, *Baignoire*.

BAGNOLES, MIN'ERAL WA'TERS OF. *Bagnoles* is a village in the department of *Orne*. The water resembles that of *Bagnères de Luchon*. Season from the 1st of August to the 1st of September.

BAGNOLS, MIN'ERAL WA'TERS OF. *Bagnols* is a village, two leagues from *Mende*, in the department of *Lozère*. The waters are hydrosulphurous and thermal: 109° Fahrenheit. The season is from the 1st of July to September.

BAGUENAUDIER, *Colutea arborescens*.

BAHAMA ISLANDS, CLIMATE OF. The climate of the *Bahamas* is not considered to be well adapted for consumptive patients, on account of the rapid alternations of temperature, and the prevalence of winds, often of a dry, cold character. Still, the phthisical valetudinarian from most portions of the *United States* might derive advantage from a residence there during the winter months. The accommodations are not, however, good, or numerous.

BAHEL, *Colum'nea longifolia*. A labiated plant of Malabar, whose leaves, bruised, are applied as cataplasms to suppurating tumours.

BAHEL SCHULLI, *Genista spinosa* Indica.

BAIGNEUR, (from *balneum*,) Bather.

BAIGNOIRE (F.), *Baptiste'rium*, a *Bathing tub*, *Baignio*, *So'lum*, *Pisci'na*. The vessel or place in which bathing is performed. *Baignoire oculaire*, an *eye-bath*,—a small vessel for bathing the eyes. See *Scaphium oculare*.

BAILEY'S SPRING, see Alabama, Mineral Waters of.

BAILLEMENT, Yawning.

BAILLON, *Speculum oris*.

BAIN, (from *balneum*,) Bath—*b. Chaud*, Bath, hot—*b. Électrique*, Bath, electric, see Electricity—*b. Entier*, Bath, general—*b. de Fonteuil*, Bath, hip—*b. Frais*, Bath, tepid—*b. Froid*, Bath, cold—*b. très Froid*, Bath, cold—*b. Hygiénique*, see Bath—*b. Marie*, Bath, water—*b. Médicinal*, Bath, medicated—*b. de Mer*, Bath, sea—*b. de Pied*, Bath, foot, *Pediluvium*—*b. de Sable*, Bath, *b. sand*—*de Siège*, Bath, hip—*b. Simple*, see Bath—*b. Tempéré*, Bath, tepid, *B. Temperate*—*b. de Tête*, Bath, head—*b. Tiède*, Bath, tepid—*b. de Vapeur*, Bath, vapour.

BAINE, Bath.

BAINS, MIN'ERAL WA'TERS OF. These are situated at Plombières, department of the Vosges. They are said to be saline and thermal by some; others deny them any medical properties. Season from the 15th of June to the 15th of September.

BAINS PRÈS ARLES, see Arles.

BAIRN, Infans.

BAIRN WORTS, Bellis.

BALAMPULLI, *Tamarindus*.

BALANCE, AEROSTATIC, Areometer.

BALANCEMENT, *Compensa'tion*, ([F.] *balance*, 'a balance,' itself from *bis*, 'twice,' and *lanx*, 'a dish.')

A law of teratogeny, as maintained by Geoffroy St. Hilaire, by which exuberance of nutrition in one organ is supposed to involve, to a greater or less extent, the total or partial atrophy of some other, and conversely.

BALANDA, *Fagus Sylvatica*.

BALANEUM, (*βαλανιον*,) Bath.

BALANISMUS, (*βαλας*, 'a suppository.')

The application of a suppository.

BALANITE CONTAGIEUSE, see Gonorrhœa.

BALANITIS, (*balano* and *itis*,) Gonorrhœa spuria.

BALANO, in composition, Balanus.

BALANOBLENNORRHŒA, (*balano* and *blennorrhœa*,) Gonorrhœa spuria.

BALANOCASTANUM, (*βαλανος*, 'an acorn,' and *καστανον*, 'a chestnut,')

Bunium Bulbocastanum.

BALANOPOSTHITIS, (*balano*, *ποσθη*, 'prepuce,' and *itis*,) Gonorrhœa spuria.

BALANORRHAGIA, (*balano* and *ῥιγηρυμ*, 'to burst forth,')

see Gonorrhœa.

BALANORRHŒA, (*balano*, and *ῥευ*, 'to flow,')

Gonorrhœa spuria.

BALANOS PHENICOS, (*βαλανος φοινικος*,) Date.

BA'LANUS, (*βαλανος*, 'glands,' 'an acorn.')

The glands penis. Hence, *Balanoblenorrhœa*, a Blenorhœa of the glands; and *Balanitis*, Inflammation of the glands. Suppositories and pessaries were called *Balanî*, (*βαλανι*.)

BALANUS, Glands, Suppository—*b. Myrepica*, *Guilandina moringa*.

BALARUC', MIN'ERAL WA'TERS OF. Balaruc is a town in the department of Hérault, France. The waters are saline and thermal. They contain carbonic acid, carbonate of lime, carbonate of magnesia, chlorides of sodium, calcium, and magnesium, sulphate of lime, and a little iron. They are considered tonic, and are

largely used. Their temperature is about 118° Fahrenheit.

BALARUC' WA'TER, FACTIT'IOUS, (F.) *Eau de Balaruc*; *Aqua Belliluea'na* is made of simple acidulous water (containing twice its bulk of carbonic acid) f 3xsss; chloride of sodium, 3iss; chloride of calcium, gr. xviii; chloride of magnesium, gr. lvi; carbonate of magnesia, gr. j.

BALATRO, Bamballo.

BALAUSTINE FLOWERS, see *Punica granatum*.

BAL'BIS, (*βαλβις*, 'a foundation.')

Any oblong cavity. Hippocrates, in his treatise on the joints, gives the name *Balbito'des* to the olecranon cavity of the humerus.

BALBUS, (F.) *Bègue*. One habitually affected with stammering. A stammerer.

BALBUTIEMENT, Balbuties.

BALBU'TIES, *Paellis'mus*, *Psel'lotes*, *Blæ'sitas*, *Baryglos'sia*, *Dysla'tia*, *Mogila'tia*, *Ischopho'nia*, *Battaris'mus*, *Bamba'lia*, *Hæbita'tio*, *Loque'la blæ'sa*, *Tituba'tio Lin'guæ*, (F.) *Balbutiement*, *Bégaïement*, *Bégayement*. Stuttering, Stammering, Hammering, (Prov.) *Tutt'ering*, *St. Vitus's Dance of the Voice*, (Sc.) *Hab'bering*, *Hobbling*. Also, vicious and incomplete pronunciation, in which almost all the consonants are replaced by the letters B and L. *Tran'lusmus*.

BALCHUS, Bællium.

BALD, Athrix.

BALDMONEY, *Æthusa meum*.

BALDNESS, Alopecia, Calvities—*b. Limited*, *Porrigo decalvans*—*b. Partial*, *Porrigo decalvans*.

BALENAS, (*balena*, 'a whale,')

Leviathan penis.

BALIMBAGO, *Hibiscus populeus*.

BALL, ([F.] *balle*, *Da. bol*.) Pila—*b. of the Eye*, Pupil.

BALLISMUS, (*βαλλισμος*,) Chorea.

BALLISTA, *Astragalus*.

BALLOCK-GRASS, *Orchis mascula*.

BALLOCKS, Testes.

BALLON, (augmentative of (F.) *balle*, 'a ball,')

Receiver.

BALLONNEMENT, Tympanites.

BALLO'TA FŒ'TIDA, (*βαλλωτη*,) *B. vulga'ris* seu *ni'gra*, *Marru'bium ni'grum*, *Black Hore'ound*, *Stink'ing H.*, (F.) *Marrube noir*. Ord. Labiata. *Sec. Syst.* *Didynamia Gymnospermia*. This plant is esteemed to be antispasmodic, resolvent, and detersive. (?)

BALLOTA LANA'TA, *Leonu'rus lana'tus*, (F.) *Ballote cotonneuse*. A plant which grows in Siberia. The whole plant, with the exception of the root, has been recommended in dropsy, and in rheumatism and gout, as a diuretic. It is usually given in decoction (3ss to 3j to f 3viij of water.)

BALLOTE COTONNEUSE, *Ballota lanata*.

BALLOTTEMENT (F.), *Mouvement de Ballottement*, *Agita'tion*, *Succus'sion*, *Repercus'sion*, is the motion impressed on the fœtus in utero, by alternately pressing the uterus by means of the index finger of one hand introduced into the vagina; the other hand being applied on the abdomen. It is one of the least equivocal signs of pregnancy.

BALLS, DEER, *Elaphomyces granulatus*.

BALLSTON SPA. This village is situate in Saratoga County, New York. The spring Sans Souci belongs to the class of Acidulous Chalybeates. It contains iodide of sodium. There is also a sulphur spring.

BALM, ((F.) *baume*, contracted from *bdlsam*,) *Melissa*—*b. Apple*, *Momordica balsamina*—*b. Bastard*, *Melitis Melissophyllum*—*b. of Gilead Poplar*, *Populus candicans*—*b. of Gilead*, *Solomon's*, see *Tinctura cardamomi*—*b. of Gilead tree*, *Dracocephalum Canariense*—*b. Indian*, *Trillium latifolium*—*b. Mountain*, *Monarda coccinea*—*b.*

Red. *Monarda coccinea*—b. Scarlet rose, *Monarda coccinea*—b. Stinking. *Hedeoma*.

BALMONY, *Chelone glabra*.

BALNEA CENOSA, (from *balneum*), see *Boue des eaux*.

BALNEARIUM, (from *balneus*), Hypocaustum.

BALNEARIUS, Bathier.

BALNEATOR, Bathier.

BALNEOGRAPHY, *Balneograph'ia*, (*balneum*, and *γραφη*, 'a description.') A description of baths.

BALNEOLOGY, *Balneolog'ia*, (*balneum*, and *λογος*, 'a description.') A treatise on baths.

BALNEOTHERAPIA, (*balneum*, and *θεραπεία*, 'treatment.') Treatment of disease by baths.

BALNEUM, Bath—b. Acidum, Bath, acid—b. Alkalinum, Bath, alkaline—b. Animale, Bath, animal—b. Antipsoricum, Bath, antipsoric—b. Anti-syphiliticum, Bath, antisiphilitic—b. Arenæ, Bath, sand—b. Gelatinosum, Bath, gelatinous—b. Mariæ, Bath, water—b. Marinum, Bath, sea—b. Maris, Bath, water—b. Medicatum, Bath, medicated—b. Sulphuris, Bath, sulphur—b. Tepidum, see Bath, hot—b. Vaporis, see Bath, hot, and Vaporarium.

BAL'SAM, (*βάλσαμον*), *Bal'samum*, *Bal'samus*, *Boleson*, *Bel'son*, (F.) *Baume*. This name is given to natural vegetable substances,—concrete or liquid, but very odorous, bitter, and piquant; composed of resin, benzoic acid, and sometimes of an essential oil, — which allow benzoic acid to be disengaged by the action of heat; readily dissolved in volatile oil, alcohol, and ether; and, when treated with alkalis, afford a soluble benzoate, and throw down resin. We know of five balsams—those of Peru, and Tolu, Benzoin, solid *Styrax* or *Storax*, and liquid *Styrax*. (See those different words.) There are, however, many pharmaceutical preparations and resinous substances, possessed of a balsamic smell, to which the name *balsam* has been given; but they differ essentially in composition and properties: hence the distinction of balsams into *natural* and *artificial*. The *natural balsams* include the five before mentioned: the *artificial* the remainder.

BAL'SAM ACOUSTIC, *Bal'samum Acousticum*, (F.) *Baume acoustique*. A mixture of fixed and essential oils, sulphur, and tinctures of fetid gums. Used in cases of atonic deafness, dropped into the ear. The *acoustic balsam* of Dr. Hugh Smith is made by mixing three drachms of *oxgall*, with one drachm of *balsam of Peru*.

BAL'SAM AMERICAN, see *Myroxylon Peruiferum*—b. Anodyne, Bates's, *Linimentum saponis* et opii.

BAL'SAM, APOPLECTIC, *Bal'samum Apoplecticum*, *Bal'samus Apoplecticus*, (F.) *Baume Apoplectique*. A medicine composed of several *balsams* properly so called, resins, and volatile oils. It is of a stiff consistence, is worn in ivory boxes about the person, and is smelled at in headaches, &c.

BAL'SAM APPLE, *Momordica balsamina*.

BAL'SAM OF ARCEUS, *Unguentum Elemi compositum*.

BAL'SAM, CALABA, see *Fagara octandra*—b. Canada, see *Pinus balsamea*—b. Canary, *Dracocephalum Canariense*—b. Capivi, *Copaiba*.

BAL'SAM OF CARPATHIA, *B. Carpathian*, *Bal'samum Carpathicum*, (F.) *Baume de Carpathie*. The resin of the *Pinus Cembra*, a tree, which grows in Switzerland, Libya, and the Krapac mountains of Hungary.

BAL'SAM, CHALYBEATE, *Bal'samum Chalybeatum*, (F.) *Baume d'acier ou d'aiguilles*. A mixture of nitrate of iron, alcohol, and oil, prepared by dissolving needles in nitric acid. It was formerly employed in friction in pains of the joints.

BAL'SAM, COMMANDER'S, *Tinctura benzoini composita*—b. of Condom, *Balsam of Leicester*.

BAL'SAM, COR'DIAL, OF SENNER'TUS, *Bal'samum Cordia'le Senner'ti*, (F.) *Baume cordiale de Senner't*. A stimulant medicine, composed of the essential oils of citron, cloves, and cinnamon, of musk, and ambergris. Dose, 6 to 15 drops.

BAL'SAM FOR CUTS, *Tinctura benzoini composita*.

BAL'SAM OF FIERABRAS. A celebrated Spanish vulnerary balsam, mentioned by Cervantes; the composition of which was oil, rosemary, salt, and wine. (?)

BAL'SAM, SPIRITUOUS, OF FIORAVENTI, *Bal'samum Fioraven'ti spirituosum*, (F.) *Baume de Fioraven'ti spiritueux*. Different products of the distillation of resinous and balsamic articles, and of a number of aromatic substances, previously macerated in alcohol, have been thus called. The *Spirituous Balsam of Fioraven'ti*, the only one now used in friction, in chronic rheumatism, is the first product of the distillation from a sand-bath. It is entirely alcoholic. The *Oily Balsam of Fioraven'ti* is obtained by removing the residue, and distilling it in an iron vessel, at a white heat. It has the appearance of a citrine-coloured oil. The *Black Balsam of Fioraven'ti* is the black oil, obtained when the temperature is sufficient to carbonize the substances in the cucurbit.

BAL'SAM OF FIR, see *Pinus balsamea*.

BAL'SAM OF FOURCROY or of LABORDE, (F.) *Baume de Fourcroy ou de Laborde*. A kind of liniment composed of aromatic plants, balsams, resins, aloes, turpentine, theriac, and olive oil. Used in chaps of the skin and nipples.

BAL'SAM, FRIAR'S, *Tinctura benzoini composita*.

BAL'SAM OF GENEVIÈVE, (F.) *Baume de Geneviève*. An ointment composed of wax, turpentine, oil, red saunders, and camphor. Used in contused wounds, gangrene, &c.

BAL'SAM OF HONEY (HILL'S). A tincture made of *tolu* and *honey*, āā ʒj , spirit, a gallon. A pectoral, used in coughs. The committee of the New York College of Pharmacy recommend the following formula—*Gum. Benzoin. ʒv, Bals. Tolut. ʒj, Mellis ʒviij, Alcohol. Oij*—digest for 10 days and filter). See *Mel*.

BAL'SAM OF HORE'HOUD (FORD'S). A tincture of *horehound*, *liquorice-root*, *camphor*, *opium*, *benzoin*, *dried squills*, *oil of aniseed*, and *honey*. It has the same properties as the above. See *Marubium*.

BAL'SAM, HUNGARIAN, see *Pinus mughos*.

BAL'SAM, HYPNOTIC, *Bal'samum Hypnoticum*, (F.) *Baume Hypnotique*. A preparation of which opium, hyoscyamus, camphor, and some other sedative substances, form the basis. It is used externally in friction, to provoke sleep.

BAL'SAM, HYSTERIC, *Bal'samum Hystericum*, (F.) *Baume Hystérique*. A preparation made of opium, aloes, asafoetida, castor, distilled oils of rue, amber, &c. It is held to the nose, applied to the navel, or rubbed on the hypogastrium in hysterical cases.

BAL'SAM, INDIAN, see *Myroxylon peruiferum*.

BAL'SAM, IODURETTED, *Bal'samum iodure'tum*, (F.) *Baume hydriodaté, B. ioduré, B. de Lausanne, Géléé contre le goître*. A balsam used in the way of friction, in goitre, at Lausanne, in Switzerland. It may be made as follows—*animal soap*, 60; *iodide of potassium*, 42; *alcohol* at 85°, 500; *essence of lemon*, 4 parts. The iodide is dissolved in the alcohol, which is added to the soap melted in a water-bath. The whole is then filtered, and put into bottles.

BAL'SAM OF LABORDE, B. of Fourcroy.

BAL'SAM OF LEICTOURE, of CON'DOM or VINCE-GRÈRE, *Bal'samum Lectocens'e*. A strongly stimulant and aromatic mixture of camphor, saffron,

musk, and ambergris, dissolved in essential oils. The ancients burnt it for the purpose of purifying the air of a chamber, when infected with a disagreeable odour.

BALSAM OF LIFE, Decoctum aloes compositum.

BALSAM OF LIFE OF HOFFMANN, *Bal'samum Vitis Hoffmanni*, (F.) *Baume de Vie d'Hoffmann*. A tincture, composed of essential oils and ambergris, employed internally and externally as a stimulant. A mixture of essential oils without alcohol constitutes the *Sax'on Bal'sam*, *Bal'samum apoplecticum*, *B. aromaticum*, *B. cephalicum*, *B. Sacramicum*, *B. nervinum*, *B. SCHERZERI*, *B. Stomachicum*. Employed in friction as a stimulant.

BALSAM OF LIFE, TURLINGTON'S, see *Tinctura benzoini composita*.

BALSAM OF LOCATEL'LI or *LUCATEL'LI*, *Bal'samum Lucatel'li*, (F.) *Baume de Lucatel*. A sort of ointment, composed of wax, oil, turpentine, sherry, and balsam of Peru, coloured with red saunders. It was once administered in pulmonary consumption.

BALSAM OF MECCA, see *Amyris opobalsamum*.

BALSAM OF METZ, GREEN, *Bal'samum Viride Meten'sium*, *B. Viride*, *O'leum ox'idi cu'pri viride*, (F.) *Baume vert de Metz*, *Baume de Feuillet*, *Huile verte*. This is composed of several fixed oils, holding, in solution, subcarbonate of copper, sulphate of zinc, turpentine, aloes, and the essential oils of cloves and juniper. It is green and caustic, and is employed to hasten the cicatrization of atonic ulcers.

BALSAM, MEXICAN, see *Myroxylon Peruiferum*.

—b. Natural, see *Myroxylon Peruiferum*.

BALSAM, NEPHRITIC, of FULLER, *Bal'samum Nephriticum Fulleri*. A liquid medicine composed of oils, resins, and balsams, which have experienced an incipient state of carbonization from concentrated sulphuric acid. It was given in the dose of 15 to 30 drops in certain affections of the kidneys.

BALSAM, NERVOUS, *Bal'samum Nervinum*, (F.) *Baume nervin ou nerveux*. A kind of ointment, composed of fatty bodies, volatile oils, balsam of Peru, camphor, &c. It is employed in friction in cases of sprains and rheumatic pains.

BALSAM, PARALYTIC, of MYN'SICHT. A sort of liniment or soft mixture of the essential oils of different aromatic plants, oils of turpentine and amber.

BALSAM OF PAREIRA BRAVA, *Bal'samum Pareiræ bravae*. A soft mixture of balsam, resin, muriate of ammonia, and powder of the root of *Pareira brava*. It is given internally to excite the urinary secretion.

BALSAM, PERUVIAN, see *Myroxylon Peruiferum* —b. of Peru, red, see *Toluifera balsamum* —b. of Peru, white, see *Myroxylon Peruiferum*.

BALSAM OF RACKAST'RA or of RAKAST'RI. This substance is of a yellowish-brown colour; semi-transparent; fragile, when dry, but softening by heat; adhering to the teeth, when chewed. It has a smell similar to that of the Balsam of Tolu, and is slightly bitter. It is brought from India, in gourd shells, and has been employed in diseases of the urinary and genital organs, especially in gonorrhœa.

BALSAM, RI'GA. Prepared from the shoots of the Scotch Fir, macerated in spirit of wine. Internally, stimulant and diuretic; externally, a vulnerary. See *Pinus Cembra*.

BALSAM OF THE SAMARITAN, (F.) *Baume du Samaritain*. A sort of liniment, prepared by boiling together, at a gentle heat, equal parts of wine and oil. It is said to have been the ointment used by the Samaritan of the Gospel to cure a patient covered with ulcers.

BALSAM OF SATURN, *Bal'samum Saturni*. A solution of acetate of lead in spirit of turpentine,

concentrated by evaporation; to which camphor has been added. This balsam was applied to hasten the cicatrization of wounds.

BALSAM, SAXON, Balsam of Life of Hoffmann.

BALSAM OF SULPHUR, *Bal'samum Sulphuris*, (F.) *Baume de Soufre*. A solution of sulphur in oil.—*B. sulph. anisatum*, (F.) *B. de Soufre anisé*; a solution of sulphur in essential oil of aniseed; given as a carminative.—*B. sulph. succinatum*, (F.) *B. de Soufre succiné*; a solution of sulphur in oil of amber.—*B. Sulphuris terebinthatum*, *Common Dutch Drops*, (F.) *B. de Soufre térébinthiné*; a solution of sulphur in essential oil of turpentine, administered as a diuretic.—The *Balsam of Sulphur of RULAND* is a solution of sulphur in linseed oil or nut oil.

BALSAM OF SYMPATHY, *Bal'samum Sympathicum*, (F.) *Baume de Sympathie*. A balsam, used in the days when sympathetic influence was strongly believed in. It was composed of the raspings of a human skull, blood, and human fat, and was applied to the instrument which had inflicted the wound.

BALSAM, THIBAUT'S. A tincture of myrrh, aloes, dragon's blood, flowers of St. John's wort, and Chio turpentine. Internally, diuretic; externally, vulnerary.

BALSAM OF TOLU, see *Toluifera Balsamum*.

BALSAM, TRANQUIL, *Bal'samum tranquillum seu tranquillans*, (F.) *B. tranquille*. A liquid medicine employed externally in the shape of friction: it is prepared by macerating and boiling, in olive oil, narcotic and poisonous plants,—belladonna, mandragora, hyoscyamus, &c.—and afterwards infusing, in the filtered decoction, different aromatic plants. It was employed as an anodyne.

BALSAM, TURKEY, *Dracocephalum Canariense* —b. Turlington's, see *Tinctura Benzoini composita*.

BALSAM OF TURPENTINE, *Bal'samum Terebinthinæ*, *Dutch Drops*. Obtained by distilling oil of turpentine in a glass retort, until a red balsam is left. It possesses the properties of the turpentines.

BALSAM, VEGETABLE, of GODBOLD, see Godbold.

BALSAM, VERVAIN'S, *Tinctura Benzoini composita*—b. of Vinçeguère, Balsam of Leictoure.

BALSAM, VULNERARY, of MINDERER'S, *Bal'samum vulnerarium Mindereri*, (F.) *B. vulnéraire de MINDERER*. A kind of liniment, composed of turpentine, resin elemi, oil of St. John's wort, and wax. Employed in friction, and as a dressing to wounds.

BALSAM WEEB, *Impatiens fulva*.—b. WOUND, *Tinctura Benzoini composita*.

BALSAMADENDRON GILEADENSE, (*Βαλσαμον*, 'balsam,' and *δένδρον*, 'tree,') *Amyris Gileadensis*—b. Myrrha, see Myrrha.

BALSAMARIA INOPHYLLUM, see *Fagara octandra*.

BALSAMELEON, (*Βαλσαμον*, 'balsam,' and *ελεον*, 'oil,') *Myroxylon Peruiferum*.

BALSAMIC, *Balsamicus*, (F.) *Balsamique*, (from Balsam.) Possessing the qualities of balsams. *Balsamic odour*:—a sweet, faint, and slightly nauseous smell. *Balsamic substance*:—one resembling the balsams in property.

BALSAMIC ÉLÉMIÈRE, *Amyris elemifera*—b. *de la Mecque*, *Amyris opobalsamum*.

BALSAMINA, *Momordica balsamina*.

BALSAMINE, *Momordica balsamina*.

BALSAMIQUE, Balsamic.

BALSAMITA FEMINEA, *Achillea ageratum* —b. Major, *Tanacetum balsamita*—b. Mas, *Tanacetum balsamita*.

BALSAMITA SUAVEOLENS, *B. odora* ta see *ma'ris*, *Men'tha Sacracen'ica* see *Roma'na*. Ord. Compositæ. Ser. Syst. Syngenesia Polygamia super-

flua. A plant, common in the south of France, and cultivated in the gardens; where it bears the names of *Menthe coq* ou *romaine*, *Grand baume*, *Baume coq* ou *Costus des Jardins*. Its smell is strong and aromatic, and taste hot. It is used for the same purposes as tansy, i. e. as a stimulant, vermifuge, &c.

BALSAMITA SUAVEOLENS, *Tanacetum balsamita*—b. *Vulgaris*, *Tanacetum balsamita*.

BALSAMOSACCHARUM, *Elæosaccharum*.

BALSAMUM, *Balsam*, *Amyris opobalsamum*—b. *Egyptiacum*, see *Amyris opobalsamum*—b. *Album*, see *Myroxylon Peruiferum*—b. *Alpini*, *Dracocephalum Canariense*—b. *Alpini*, see *Amyris opobalsamum*—b. *Anodynum*, *Linimentum saponis et opii*—b. *Apoclecticum*, *Balsam* of life of Hoffmann—b. *Arcei*, *Unguentum elemi compositum*—b. *Aromaticum*, *Balsam* of life of Hoffmann—b. *Asiaticum*, see *Amyris opobalsamum*—b. *Brazilense*, *Copaiba*—b. *Calaba*, see *Fagara octandra*—b. *Canadense*, see *Pinus balsamea*—b. *Catholicum*, *Tinctura benzoini composita*—b. *Cephalicum*, *Balsam* of life of Hoffmann—b. *Commen-datoris*, *Tinctura benzoini composita*—b. *Copaibæ*, *Copaiba*—b. *Genuinum antiquorum*, see *Amyris opobalsamum*—b. *Hyperici simplex*, see *Hypericum perforatum*—b. *Indicum*, see *Myroxylon Peruiferum*—b. *Ioduretum*, *Balsam*, ioduretted—b. *Judaicum*, see *Amyris opobalsamum*—b. *Libani*, see *Pinus cembra*—b. *Locatelli*, *Balsam* of Locatelli—b. *Marise*, see *Fagara octandra*—b. *e Meccâ*, see *Amyris opobalsamum*—b. *Mercuriale*, *Unguentum hydrargyri nitratis*—b. *Nervinum*, *Balsam* of life of Hoffmann—b. *Nucistæ*, see *Myristica*—b. *Ophthalmicum rubrum*, *Unguentum hydrargyri nitrice-oxydi*—b. *Opodeldoc*, *Linimentum saponis camphoratum*—b. *Persicum*, *Tinctura benzoini composita*—b. *Peruvianum*, see *Myroxylon Peruiferum*—b. *Saturninum*, *Unguentum plumbi superacetatis*—b. *Scherzeri*, *Balsam* of life of Hoffmann—b. *Stomachicum*, *Balsam* of life of Hoffmann—b. *Styracis*, *Styrax*—b. *Styracis benzoini*, *Benjamin*—b. *Succini*, see *Succinum*—b. *Sulphuris Barbardense*, *Petroleum sulphuratum*—b. *Sulphuris simplex*, *Oleum sulphuratum*—b. *Syriacum*, see *Amyris opobalsamum*—b. *Tolutanum*, see *Toluifera balsamum*—b. *Tranquillans seu Tranquillum*, *Balsam*, tranquil—b. *Traumaticum*, *Tinctura benzoini composita*—b. *Universale*, *Unguentum plumbi superacetatis*—b. *Viride*, *Balsam*, green, of Metz; see *Fagara octandra*.

BALSAMUS, *Balsam*—b. *Palustris*, *Mentha aquatica*.

BALSEM, *Amyris opobalsamum*.

BAMBA, *Bamboo*.

BAMBALIA, *Balbuties*.

BAMBALIO, *Bambalo*, *Bala'tro*, (*Βαββαλω*, 'I speak inarticulately.') One who stammers or lisps, or utters inarticulate sounds. According to KRAUS one who speaks as if he had pap in his mouth, or as if his tongue were paralyzed.

BAMBOO, (F.) *Bambou*, *Bambu*. *Fam. Gramineæ*. *Sec. Syst. Hexandria Monogynia*. The young shoots of *Bambos arundinacea*, *Arun'do bambos*, *Bambu'sa arundinacea*, and of *Bambos verticillata*, contain a saccharine pith, of which the people of both the Indies are very fond. They are sometimes made into a pickle.

BAMBOS (name in India) **ARUNDINACEA**, *Bamboo*—b. *Verticillata*, *Bamboo*.

BAMBUSA ARUNDINACEA, *Bamboo*.

BAMIA MOSCHATA, *Hibiscus abelmoschus*.

BAM'MA, gen. *Bamm'atis*, (*Βαμπα*, from *βαρρω*, 'I plunge.') Anciently, liquids were so called, in which certain bodies were plunged, to moisten or soften them. In the case of tea, for instance, into which bread is dipped, the tea would be the *Lamma*.

BANANA, *Musa sapientum*.

BANANIER, *Musa sapientum*.

BANAUSIA, (*Βαναυσία*) Charlatany.

BANC D'HIPPOCRATE, ('bench of Hippocrates.') Bathron.

BANCAL, (F.) One who has deformed legs. It includes *vulgus*, *compervis*, and *varus*.

BANCROCHE, (F.) A vulgar epithet for a rickety individual.

BAND, FLATTENED, see *Nerve Fibre*—b. **PRIMITIVE**, see *Nerve Fibre*.

BAN'DAGE, *Des'ma*, *Syndes'mus*, *Hypodes'mis*, *Hypodes'ma*, *Hypodes'mus*, *Hypod'es'is*, and *Hypod'es'is*, (the last five signify properly "an under bandage"); A *binder*, (Sax. *bindan*, 'to bind.') This word, with the French, is generally used to express the methodical application of rollers, compresses, &c., *Ban'daging*, *Syn'des'is*, to fix an apparatus upon any part,—corresponding to the words *deligatio*, *fasciatio*, *fasciarum applicatio*, *epid'es'is*. With us the noun is usually applied to the result of the application, or to the bandage itself;—a sense in which the French employ the word *Bande*. Bandages are *simple* or *compound*. The simple bandage is *equal*, if the turns are applied circularly above each other; *unequal*, if the turns are not accurately applied upon each other. If each turn of the bandage be only covered one-third, it forms the *doloire* of the French; if the edges touch only slightly, it is the *moussé*; if the turns are very oblique and separated, it is the *spiral*, or *creeping*, (F.) *rampant*; if folded upon each other, it is termed the *reversed*, (F.) *renversé*. By uniting various kinds of bandaging, we have the *compound*; and these compound bandages have received various names expressive of their figure, or of the parts to which they are applied, as *capistrum*, *spica*, &c. Bandages are divided, also, as regards their uses, into *uniting*, *dividing*, *retaining*, *expelling*, *compressing*, &c.

BANDAGE OR ROLL'ER, *Fas'cia*, *Tæ'nia*, *Epides'mus*, *Vin'culum*, the *Bande* of the French. It may be made of linen, flannel, or other stuff capable of offering a certain resistance. The two extremities of a bandage are called *tails*, (F.) *chefs*, the rolled part is termed its *head*, (F.) *globe*, and the rest of the bandage is the *body*, (F.) *plein*. If rolled at both extremities, it is called a *double-headed roller* or *bandage*, (F.) *Bande à deux globes*.

BANDAGE À BANDELETES SEPARÉES, *Bandage* of separate strips.

BANDAGE, *Bod'y*, *Manti'le*, *Mante'le*, (F.) *Bandage de Corps*, is used for fixing dressings, &c., to the trunk. It is formed of a towel, napkin, or some large compress, folded three or four times; the extremities of which are fastened by pins. This is again fixed by means of the *scapulary bandage*, which is nothing more than an ordinary bandage, stitched to the anterior and middle part of the napkin, passing over the clavicles and behind the head, to be attached to the back part of the napkin.

BANDAGE, COMPOUND, see *Bandage*.

BANDAGE, COMPRESS'ING, or ROLL'ER, *Fas'cia compressiva seu convoluta* (F.) *Bandage compressive* ou *roulé*, is the simple *roller* with one head; and is employed in cases of ulcers, varices, &c., of the limbs. Whenever this roller is applied to the lower part of the limbs, it is carried upwards by the *doloire* and *reversed* methods above described.

BANDAGE, COMPRESSIVE, see *Bandage*, compressing—b. *de Corps*, see *Bandage*—b. *Dividing*, see *Bandage*—b. *Divisif*, *Dividing bandage*—b. *à Dix-huit Chefs*, *Bandage*, eighteen-tailed—b. *en Doloire*, *Doloire*—b. *Double-headed*, see *Bandage*.

BANDAGE, EIGHTEEN-TAILED, *Fas'cia octod'ecim capit'ibus*, (F.) *Bandage à dix-huit chefs*. This bandage is made of a longitudinal portion of a common roller; and with a sufficient number

of transverse pieces or tails, to cover as much of the part as is requisite. It is a very useful bandage, inasmuch as it can be undone without disturbing the part.

BANDAGE, EXPELLING, see Bandage.

BANDAGE, GALLEN'S, *B. for the Poor*, *Fascia Galeni seu Pauperum*, (F.) *Bandage de Galien ou des Pauvres*, *Gal'ea*, is a kind of cucullus or hood, (F.) *Couvrechef*, divided into three parts on each side, of which GALEN has given a description. See Cancer Galeni.

BANDAGE OF GENGA, Bandage of Theden—b. Hernial, see TRUSS—b. Immovable, Apparatus, immovable.

BANDAGE, INGUINAL, *Fascia inguinalis*. A bandage for keeping dressings applied to the groin. It consists of a cineture, to which is attached a triangular compress, adapted for covering the groin. To the lower extremity of this, one or two bandages are attached, which pass under the thigh, and are fixed to the posterior part of the cineture. This bandage may be either simple or double.

Other bandages will be found described under their various names.

BANDAGE DES PAUVRES, ('B. of the poor.') Bandage, Galen's—b. Permanent, Apparatus, immovable—b. of the Poor, see Bandage, Galen's, and Cancer Galeni—b. Rampant, see Bandage—b. *Renversé*, see Bandage—b. *Roulé*, see Bandage—b. of Scultetus, Bandage of separate strips.

BANDAGE OF SEPARATE STRIPS, or **B. OF SCULTETUS**, *Fascia fasciolaris separata* *dispositis seu Sculteti*, (F.) *Bandage à bandelettes séparées ou de Scultet*. This is formed of linen strips, each capable of surrounding once and a half the part to which they have to be applied, and placed upon each other, so as to cover successively one-third of their width. It is used chiefly for fractures, requiring frequent dressing.

BANDAGE, SIMPLE, see Bandage—b. Starch, see Apparatus, immovable.

BANDAGE OF THE DEN, *B. of Genga*. A bandage employed, at times, in brachial aneurism, which commences at the fingers, and extends to the axilla.

BANDAGE, UNDER, *Hypodesmis*—b. Unequal, see Bandage—b. *Unitant*, Uniting bandage—b. Uniting, see Bandage, and Uniting bandage.

BANDAGING, see Bandage—b. Doctrine of, *Desmaturgia*.

BANDAGIST, (F.) *Bandagiste*. One whose business it is to make bandages, and especially those for hernia.

BANDE, Bandage. The word *Bande*, in anatomy, is used by the French for various narrow, flat, and elongated expansions.

BANDE À DEUX GLOBES, see Bandage.

BANDE D'HÉLÉODORE, a kind of bandage for supporting the mammae.

BANDEAU, (F.) A kind of simple bandage, which consists of a piece of cloth, folded four times, and applied round the head. There is also the *Bandeau ou Mouchoir en triangle* or *triangular bandage*, a kind of *couvrechef*, made of a square piece of cloth, or of a handkerchief, folded diagonally, and applied round the head.

BANDELETTE, (F.) (dim. of *Bande*) *Fasciola*, *Ten'ola*, *Vit'la*; a narrow bandage, strip, or fillet. Also *Tænia semicircularis*.

BANDELETTE DES CORNES D'AMMON, *Corpus fimbriatum*—b. *des Eminences pyriformes*, *Tænia semicircularis*—b. *de l'Hippocampe*, *Corpora fimbriata*.

BANDELETTE SEMICIRCULAIRE, *Tænia semicircularis*.

BANDELETTES AGGLUTINATIVES. Small strips, covered with a glutinous plaster. *Vitæ agglutinant'es*. See Agglutinant.

BANDELETTES DÉCOUPÉES, are strips of linen, notched on one edge, and covered, on one side, with ointment. They are applied to wounds to prevent the lint from sticking, and the laceration of the cicatrix.

BANDURA, *Nepentha destillatoria*.

BANDY-LEGGED, *Cnemoscotiosis*.

BANEBERRY, *Actæa spicata*.

BANGUE, *Bhang*, *Bang*, *Bangi* or *Beng*, *Sed-hee*, *Subjee*. Adanson believes this to be the *Nepenthes* of the ancients. The largest leaves and capsules, without the stalks, of *Can'nabis Indica*, (F.) *Chanvre Indien*, *Indian Hemp*, probably identical with *C. sativa*. Family, *Urticæ*. Sex. Syst. *Diœcia Pentandria*. The leaves and flowers of *Cannabis* are narcotic and astringent. They are chewed and smoked. The seeds, mixed with opium, arca, and sugar, produce a kind of intoxication, and are used for this purpose by the people of India. An alcoholic extract of the dried tops, *Chur'rus*—*Ectra'ctum Can'nabis* (Ph. U. S.)—has been used in India, and since then in Europe and in this country, as a narcotic and anticonvulsive, in the dose of from half a grain to ten or more. It requires, however, great caution in its administration. The pure resin—*Cannabin*, *Haschischin*—is active in the dose of two-thirds of a grain.

The dried plant, which has flowered, and from which the resin has not been removed, called *Gun'jah*, *Gen'jah*, *Gua'za*, and *Gua'zah*, *Hasch'isch*, *Hasch'ich*, *Hach'isch*, *Has'h'ish*, or of the Arabs, consists of the tops and tender parts only of the plant, collected immediately after inflorescence, and simply dried.

BANICA, *Pastinaca sativa*.

BANILAS, *Vanilla*.

BANILLA, *Vanilla*.

BANILLOES, *Vanilla*.

BANISTERIA ANGULO'SA, (after Rev. John Banister, a botanist.) This plant, in Brazil and the Antilles, passes for a powerful sudorific, and an antidote to the poison of serpents.

BANKSIA ABYSSINICA, (after Sir Joseph Banks, P. R. S.) *Hagenia Abyssinica*—b. *Speciosa*, *Costus*.

BANNIÈRES, MIN'ERAL WA'TERS OF. Bannières is a village in Quercy, diocese of Cahors, France. The waters are probably chalybeate. They are celebrated in amenorrhœa, cachexia, jaundice, &c.

BA'OBAB, *Adan'sonia digitata*, of Africa. *Not. Ord.* *Bombacæ*. One of the largest productions of the vegetable kingdom. Its fruit is called, in the country, *Pain de singe*, ('monkey bread.') The pulp is sourish, and agreeable to eat; and a refreshing drink is made from it, which is used in fevers Prospero Alpini and Dr. L. Frank think that the *Ter'ra Lem'nia* was prepared, in Egypt, from the pulp. All the parts of the Baobab abound in mucilage. The bark has been given as a substitute for cinchona.

BAPTISIA LEUCANTHA, (from *βαπτειν*, 'to dye,') see *Sophora tinctoria*—b. *Tinctoria*, *Sophora tinctoria*.

BAPTISTERIUM, (*βαπτιστήριον*, from *βαπτίζω*, 'to immerse,') *Baignoire*.

BAPTORRHŒA, (*βαττω*, 'to infect,' and *ῥεω*, 'to flow,') See Gonorrhœa.

BARAQUETTE (F.). A name given by Rous, physician at Nîmes, in France, to a catarrhal epidemic, which occurred there in 1761. See Influenza.

BARATHRON, *Juniperus sabina*.

BARATHRUM, (*Βαρᾶθρον*) *Antrum*.

BARBA, Beard—b. *Aronis*, *Arum maculatum*—b. *Capræ*, *Spiræa ulmaria*—b. *Hirci*, *Tragopogon*—b. *Jovis*, *Sempervivum tectorum*.

BARBADOES, see West Indies—b. *Leg*, see Elephantiasis.

BARBAREA ('herb St. Barbary'), *Erysimum* *barbarea*—b. *Stricta*, *Erysimum* *barbarea*.

BARBAROS'SÆ PILULÆ, *Barbaros'sæ's Pills*. An ancient composition of quicksilver, rhubarb, diagridium, musk, &c. It was the first internal mercurial medicine which obtained any real credit.

BARBE, Peard—b. *de Bouc*, *Tragopogon*.

BARBEAU, *Centaurea cyanus*, *Cyanus* *segetum*.

BARBER-CHIRURGEONS. A Corporation of London, instituted by King Edward IV. The barbers were separated from the surgeons, by 18 Geo. II., c. 15; and the latter were erected into a *Royal College of Surgeons* at the commencement of the present century. The *Barber's pole*, of the present day, is a relic of the old staff that was grasped, and the coloured strip passing spirally around it, of the fillet or tape employed in the operation of bleeding.

BARBER'S POLE, see *Barber-chirurgeons*.

BARBERS, ARMY, see *Bathers*.

BARBERIE, MINERAL WATERS OF. These mineral waters are half a league from Nantes. They contain carbonic acid, chlorides of magnesium and sodium, sulphate of magnesia, carbonates of magnesia, lime, and iron. They are used as chalybeates.

BARBERRY, (*berberis*), *Oxycantha* *Galenii*—b. *American*, see *Oxycantha* *Galenii*.

BARBIERS. A variety of paralysis chiefly prevalent in India; and by many considered to be the same as *Beriberi*. *Beriberi* is commonly an acute disease. *Barbiers* is generally chronic.

BARBITIUM, (from *barba*), *Beard*.

BARBONE, *Pubis*, os.

BARBOTINE, *Artemisia* *SANTICINA*.

BARBULA (dim. of *barba*) *CAPRINA*, *Spiræa* *ulmaria*.

BARCLAY'S ANTIBILIOUS PILLS, *Pilulæ* *antibiliosæ*.

BARDADIA, *Pound*.

BARDANA, *Arctium* *lappa*—b. *Minor*, *Xanthium*.

BARDANE PETITE, *Xanthium*.

BARÈGES, MINERAL WATERS OF. *Barèges* is a village in the department of *Hautes Pyrénées*, near which are several springs. They are sulphureous and thermal, the heat varying from 85° to 112° Fahrenheit. They contain chlorides of magnesium and sodium, sulphates of magnesia and lime, carbonate of lime, sulphur, &c. These springs have long enjoyed a high reputation, and are daily advised in cutaneous and scrofulous affections, &c. The season is from the 1st of June to the 15th of September.

A nitrogenized matter was first found in these waters, and afterwards in other sulphureous springs, to which *Longchamp* gave the name *Barégine*.

FACTITIOUS BARÈGES WATER, *Aqua Baréginen'sis* seu *Baréginen'sis*, (F.) *Eau de Barèges*, is made by adding *hydrosulphuretted water*, f 3iv, to *pure water*, f 3xviijss, carbonate of soda, gr. xvj, chloride of sodium, gr. ss. Bottle closely.

BARÉGINE, see *Barèges*.

BARGADA, *Convolvulus* *pes capræ*.

BAR'GOU. An alimentary preparation formed of ground oats, boiled to a proper consistence with water:—see *Avena*.

BARIGLIA, *Soda*.

BARI CHLORIDUM, *Baryta*, *muriate* of—b. *Iodidum*, *Baryta*, *hydriodate* of.

BARILLA, *Soda*—b. *Alicant*, *Soda*—b. *Carthagena*, *Soda*—b. *Turkey*, *Soda*.

BARILLOR, *Soda*.

BAR'RIUM, *Ba'ryum*, *Baryt'ium*, (*Bapros*, 'heavy.') *Pluto'nium*. The metallic base of *baryta*, so called from the great density of its compounds.

BAR'RIUM CHLORATUM, *Baryta*, *muriate* of—b.

Chloride of, *Baryta*, *muriate* of—b. *Chlorure* *de Baryta*, *muriate* of—b. *Iodatium*, and *Iodide* of, *Baryta*, *hydriodate* of—b. *Protoxide* of, *Baryta*.

BARK, *Cinchona*, see *Cortex*—b. *Arica*, see *Cinchonæ cordifoliae* *cortex*—b. *Ash*, see *Cinchona*—b. *Bitter*, *Pinckneya pubens*—b. *Bogota*, see *Cinchona*—b. *Calisaya*, *Cinchonæ cordifoliae* *cortex*—b. *Calisaya*, *spurious*, see *Cinchonæ cordifoliae* *cortex*—b. *Carabaya*, see *Cinchonæ cordifoliae* *cortex*—b. *Caribæan*, *Cinchonæ Caribææ* *cortex*—b. *Carthagera*, see *Cinchona*—b. *Cherry*, *wild*, *Prunus Virginiana*—b. *Coquette*, see *Cinchona*—b. *Crown*, *Cinchonæ lancifoliae* *cortex*—b. *Cusco*, see *Cinchonæ cordifoliae* *cortex*—b. *Doom*, *Sassy-bark*—b. *Elk*, *Magnolia glauca*—b. *Essential salt* of, see *Cinchona*—b. *Florida*, *Pinckneya pubens*—b. *Fusagasuga*, see *Cinchona*—b. *Georgia*, *Pinckneya pubens*—b. *Gray*, see *Cinchona*—b. *Huamiles*, see *Cinchona*—b. *Huanuco*, see *Cinchona*—b. *Indian*, *Magnolia glauca*—b. *Iron*, see *Kino*—b. *Jaen*, see *Cinchona*—b. *Jesuit's*, *Cinchona*—b. *Lima*, see *Cinchona*—b. *Loxa*, *Cinchonæ lancifoliae* *cortex*—b. *Maraçaybo*, see *Cinchona*—b. *Neem*, *Melia azedarach*—b. *Oak*, *white*, *Quercus alba*—b. *Ordeal*, *Sassy bark*—b. *Pale*, *Cinchonæ lancifoliae* *cortex*—b. *Peruvian*, *Cinchona*—b. *Pitaya*, *Cinchonæ Caribææ* *cortex*, see *Cinchona*—b. *Red*, *Cinchonæ oblongifoliae* *cortex*—b. *Royal*, *Cinchonæ cordifoliae* *cortex*—b. of *St. Ann*, see *Cinchonæ cordifoliae* *cortex*—b. *Saint Lucia*, *Cinchonæ Caribææ* *cortex*—b. *Santa Martha*, see *Cinchona*—b. *Sassy*, *Sassy bark*—b. *Seven*, *Hydrangea arborescens*—b. *Silver*, see *Cinchona*—b. *Yellow*, *Cinchonæ cordifoliae* *cortex*.

BARLERIA BUXIFOLIA, (after *Rev. James Barrelier*, M. D.), *Cara schulli*.

BARKING DISEASE, *Mal de Laira*.

BARLEY, *Hordeum*.

BARLEY-BREE, *Cerevisia*—b. *Corn*, *Cerevisia*.

BARLEY, PEARL, see *Hordeum*—b. *Scotch*, *Hordeum*.

BARLEY WATER, *Decoctum hordei*.

BARM, ([Sax.] *beoþm*), *Yest*.

BARN, ([Prov.], [Sax.] *beapn*), *Infans*.

BARNET, MINERAL WATERS OF. *Barnet* is not far from *London*. The water is of a purging quality, like that of *Epsom*, and about half the strength.

BAROMACROMETER, *Pædobaromacrom'eter*, *Pædom'eter*, (*Bapros*, 'weight,' *μακρος*, 'long,' and *μετρον*, 'measure.') An instrument invented by *Stein* to indicate the length and weight of a newborn infant.

BAROMETER, *Baroscop'ium*, *Ba'roscope*, (*Bapros*, 'weight,' and *μετρον*, 'measure.') (F.) *Baromètre*. An instrument which measures the weight of the air. A certain degree of density in this medium is necessary for health. When we ascend high mountains, great inconvenience is experienced, owing to the diminished density. Changes of this character are indicated by the barometer or weather-glass.

BA'ROS, (*Bapros*, 'weight,') Employed by the Greek physicians to designate the feeling of lassitude and heaviness observable in many diseases.

BAROSCOPE, (*Bapros*, 'weight,' and *σκοπεω*, 'I view.') *Barometer*.

BAROSMA BETULINA, *B. Crenata* seu *Crenulata* and *B. Serratifolia*, (*Bapros*, 'heavy,' and *οσμη*, 'odour,') *Diosma crenata*.

BAROTE, *Baryta*.

BAROTES SALITUS, *Baryta*, *muriate* of.

BARRAS, see *Pinus sylvestris*.

BARRE, MINERAL WATERS OF. *Barre* is a small town, six leagues from *Strasbourg*. The waters are thermal, and contain much iron, calcareous salt, &c. They are diuretic and tonic.

BARRE, (F.) 'a bar.' *Barrure*, *Va'ra*. A projection or prolongation of the symphysis pubis;—a deformity rendering delivery difficult.

BARRÉE, (F.) 'barred.' A term applied, in France, to a female whose pelvis has the deformity described under *Barre*.

BARRÉS (**DENTS**), *Barr'ed Teeth*. The molar teeth, when the roots are spread or tortuous, so that they cannot be extracted without being broken; or without a portion of the alveolar arch being removed.

BARREL OF THE EAR, Tympanum.

BARREN, Sterile.

BARRENNES, Sterilitas.

BARROS, Terra Portugallica.

BARRURE, *Barre*.

BARTON'S FRACTURE, see Fracture of the Radius, Barton's.

BARTON'S OPERATION FOR ANKYL'OSIS. This consists in cutting out a V-shaped portion of bone. It was first devised, and carried into execution, by Dr. John Rhea Barton, of Philadelphia, in 1826.

BARY, (from βαρυς, 'heavy.') *Gravis*; in composition, heavy, oppressive.

BARYCOCCALON, (*bary*, and κοκκαλος, 'a nut.') *Datura stramonium*.

BARYCOCTA, *Bradycocti'a*, *Paracusi'a obtusa*, *Dyscocti'a*, *Audi'tus diffic'ilis*, *A. gra'vis*, *A. imminu'tus*, *Obaudi'tus*, *Hypocopho'sis*, (F.) *Durété d'Oreille*, *Dysécécé*, (*bary*, and ακων, 'hearing.') Hardness of hearing, incomplete deafness. See *Cophosis*, and *Deafness*.

BARYGLOSSIA, (*bary*, and γλωσση, 'tongue.') *Balbuties*, *Baryphonia*.

BARY HYDRAS IODATI, *Baryta*, hydriodate of.

BARYLALIA, (*bary*, and λαλειν, 'to speak.') *Baryphonia*.

BARYODYNE, (*bary*, and οδυνη, 'pain.') *A* dull, heavy pain.

BARYPHONIA, *Baryglos'sia*, *Baryla'lia*, *Loque'la impedi'ta*, (*bary*, and φωνη, 'voice.') Difficulty of voice or speech.

BARYPICRON, (*bary*, and πικρος, 'bitter.') *Artemisia abrotanum*.

BARYSOMATIA, (*bary*, and σωμα, 'body.') *Polysarcia adiposa*.

BARYSOMATICA, *Polysarcia adiposa*.

BARYTA, (*βαρυς*, 'weight.') *Ter'ra ponderosa*, *Bary'tes*, *Protoxide of Barium*, *Heavy Earth*, *Ponderous Earth*, (F.) *Baryte*, *Barote*, *Terre pesante*. This earth and its soluble salts are all highly corrosive poisons. It is never employed in medicine in the pure state. When externally applied, it is caustic, like potassa and soda.

BARYTA, CARBONATE OF, *Bary'tæ Carbo'nas*, (F.) *Carbonate de Baryte*, *Craie barotique*, is only used officinally to obtain the muriate.

BARYTA, HYDRIOATE OF, *Bary'tæ Hydri'odas*, *Bary'ta Hydriod'ica*, *Hy'dras Ba'ryi Ioda'ti*:—(in the dry state, *Iodide of Barium*, *Ba'rii Iod'idum*, *Ba'rium Ioda'tum*), has been given in serofulous and similar morbid conditions. It may be administered internally in the dose of one-eighth of a grain three or four times a day, and be applied externally to serofulous swellings, in the form of ointment, (gr. iv to ʒj of lard.)

BARYTA HYDRIODICA, *Baryta*, hydriodate of.

BARYTA, MURIATE OF, *Bary'tæ Muri'as*, *Chlo'ride of Barium*, *Ba'rii Chlo'ridum* (Ph. U. S.), *Ba'rium Chlo'ratum*, *Chlo'ruret of Barium*, *Ter'ra ponderosa sali'ta seu muri'ata*, *Sal muriat'icum barot'icum*, *Baro'tes sali'tus*, (F.) *Chlorure de barium*, is the combination of *baryta* chiefly used. It is made by the action of muriatic acid on carbonate of *baryta*.

It is given in the form of the *Solutio Muriat'is Bary'tæ*, *Li'quor Ba'rii Chlo'ridi* (Ph. U. S.), *A'qua bary'tæ muri'atis*, (F.) *Solutio de Muriate de Baryte*, (*Ba'rii chlorid. ʒj: Aq. destillat. ʒiij*);

and is employed in serofulous cases, warts, and cutaneous diseases. Externally, to fungous ulcers and to specks on the cornea. Dose, grt. v, two or three times a day.

BARYTA, SULPHATE OF, *Bary'tæ sulph'as*, (F.) *Sulfate de Baryte*, is used in pharmacy, like the carbonate, to obtain the chloride of barium.

BARYTÆ CARBONAS, *Baryta*, carbonate of—*b. Hydriodas*, *Baryta*, hydriodate of—*b. Murias*, *Baryta*, muriate of—*b. Sulphas*, *Baryta*, sulphate of.

BARYTE, *Baryta*—*b. Carbonate de*, *Baryta*, carbonate of—*b. Sulfate de*, *Baryta*, sulphate of.

BARYTHYMIA, (*βαρθυμία*, from *bary*, and θυμος, 'mind,') *Melancholy*.

BARYTIUM, Barium.

BARYUM, Barium.

BAS-FOND, (*bas*, 'low,' and *fond*, 'bottom,') see *Urinary Bladder*.

BAS-LASSÉ, *Stocking*, laced.

BAS-VENTRE, ('lower belly,') *Abdomen*.

BASAAL. The name of an Indian tree, the decoction of whose leaves, in water, with ginger, is used as a gargle in diseases of the fauces. The kernels of the fruit are vermifuge.

BASANASTRA-GALA, (*βασανος*, 'torture,' and αστραγαλος, 'the astragalus,') Pain in the ankle joint; gout in the foot.

BASANIS'MUS, (*βασανισμος*, from βασανιζειν, 'to explore,') 'A touch-stone,') Investigation or examination.

BASE, *Ba'sis*, (*βασis*, from βαινω, 'I proceed,') 'I rest,' 'I support myself,') That which serves as a foundation or support. That which enters, as a principal matter, into a mixture or combination. In anatomy, it is employed in the former sense, as *Base of the Cra'nium*, *Base of the Brain*—*Ba'sis seu Pavimentum cereb'ri*; *Base of a process*, &c., *Base of the heart*—*Ba'sis vel coro'na cor'dis*. In dentistry, it means a metallic, ivory, hippopotamus or other plate, which is used as a support for artificial teeth. In the language of prescribing, *Basis* is the chief substance which enters into a compound formula.

BASE-BORN, Illegitimate.

BASEMENT MEMBRANE, see *Membrane*, basement.

BASIATIO, (*basiare*, *basiatum*, 'to kiss,') *Coin*.

BASIATOR, (same etymon,) *Orbicularis oris*.

BASIL BUSH, ([F.] *Basilic*, from βασιλικος, 'royal,') *Ocimum caryophyllatum*—*b. Citron*, *Ocimum basilicum*—*b. Common*, *Ocimum basilicum*—*b. Small*, *Ocimum caryophyllatum*—*b. Wild*, *Chenopodium vulgare*, *Cunila mariana*, and *Pycnanthemum incanum*.

BASILAD, see *Basilar Aspect*.

BASILAR, *Basila'ris*, *Bas'iliary*, (F.) *Basilaire*. That which belongs to the base. This name has been given to several parts, which seem to serve as bases to others. The sacrum and sphenoid have been so called.

BASILAR ARTERY, *A. basila'ris seu cervica'lis*, (F.) *Artère ou Tronc basilare*, *A. mésocephalique* (Ch.) The union of the two vertebral arteries. It ascends along the middle groove on the inferior surface of the tuber, and is supported, beneath, by the *Fossa basilaris*. It terminates in the posterior cerebral arteries.

BASILAR ASPECT. An aspect towards the base of the head.—Barclay. *Basilad* is used adverbially by the same writer to signify 'towards the basilar aspect.'

BASILAR FOS'SA, (F.) *Gouttière ou Fosse basilare*, is the upper surface of the basiliary process,—so called because it is channelled like a *Fossa* or *Gutter*. The *Tuber annulare* rests upon it.

BASILAR PROC'ESS, *Processus basila'ris os'eus occip'itis seu cuneifor'mis os'eus occip'itis*, (F.) *Apophyse Basilaire*, *Prolongement sous-occip'ial*,

Uniform Process, is the bony projection, formed by the inferior angle of the os occipitis, which is articulated with the sphenoid.

BASILAR SINUS, Sinus transversus.

BASILAR SURFACE, (F.) *Surface basilaire*, is the inferior surface of the process. It is covered by the mucous membrane of the pharynx.

BASILAR VERTEBRA. The last vertebra of the loins.

BASILIC, *Basilicus*, (F.) *Basilique*, (*Βασιλικός*, 'royal.') This name was given, by the ancients, to parts which they conceived to play an important part in the animal economy.

BASILIC VEIN, *Veina basilica seu cubiti interior*, (F.) *Veine Basilique*, *Veine cubitale cutanée* of Chaussier. This vein is one of those on which the operation of bloodletting is performed. It is situate at the internal part of the fold of the elbow, in front of the humeral artery, and is formed by the *anterior* and *posterior cubital veins*, and by the *median basilic*. It terminates, in the arm-pit, in the axillary vein. The ancients thought, that the basilic of the right arm had some connexion with the liver, and hence they called it *hepatic*. The vein of the left arm, for a similar reason, they called *splenic*. The *Median Basilic Vein*, (F.) *Veine médiane basilique*, is one of the branches of the preceding vein. It joins the median cephalic at an acute angle, or rather by a transverse branch, and receives some branches of the deep radial and cubital veins, and a considerable subcutaneous vein—the *common median*.

BASILIC COMMUN, *Ocymum basilicum*—*b. Sauvage*, *grand*, *Chenopodium vulgare*.

BASILICON, *Basilicum*, (*Βασιλικόν*, 'royal, or of great virtue,') An ointment, composed of yellow wax, black pitch, and resin, of each one part, olive oil, four parts. Hence it was called *Unguentum Tetrapharmacum*, (*Τετραφαρμακον*, 'four drugs,')

BASILICON, *Basilicum*, of the Parisian Codex, is the *Onguent de Poix et de Cire*. In most Pharmacopœias, it is represented by the *Unguentum* or *Cera-tum Resinæ*. It is used as a stimulating ointment. See *Cera-tum Resinæ*, and *Unguentum Resinæ Nigræ*.

BASILICUM, *Basilicon*, *Ocymum Basilicum*—*b. Citratum*, *Ocymum basilicum*—*b. Majus*, *Ocymum basilicum*.

BASILIQUE, *Basilic*.

BASILISCUS, *Syphilis*.

BASIO, (from *basis*, 'base,') in composition, base.

BASIOCERATOCHONDROGLOSSUS, (*χονδροειδής*, 'cartilage,') *Hyoglossus*.

BASIOCRATOGLOSSUS, (*basio*, and *κρας*, 'cornu,' and *γλωσσα*, 'tongue,') A name given to a part of the hyoglossus, which is inserted into the cornu of the os hyoides and base of the tongue.

BASIOCES'TRUM, (*basio*, and *κεστρα*, 'a dart,') An instrument for opening the head of the fœtus in utero, invented by Mesler, a German.

BASIOGLOSSUS, *Hypoglossus*, (*basio*, and *κρας*, 'cornu,' and *γλωσσα*, 'tongue,') A name formerly given to the portion of the hyoglossus which is inserted into the base of the os hyoides. See *Lingual Muscles*.

BASIOPHARYNGÆUS, (*basio*, and *φάρυγξ*, 'the pharynx,') A name given to some fibres of the constrictor pharyngis medius.

BASIS, Base, see *Prescription*—*b. Cerebri*, Base of the Brain—*b. Cordis*, Radix cordis—*b. Corporis*, Sole—*b. of the Peduncle*, see *Peduncles of the Brain*.

BASSICOLICA. Name of an ancient medicine composed of aromatics and honey, and proposed by Julius Bassus.

BASSIA BUTYRACEA, (after F. Bassi, botanist of Bologna,) see *Spirit*, (Arrack.)

BASSIN, Pelvis—*b. Oculaire*, Scaphium oculare.

BASSINER, *Fore're*, to foment.

BASSINET, Pelvis of the kidney, *Ranunculus bulbosus*.

BASSORA, GUM. A gum, obtained from a plant unknown, which came originally from the neighbourhood of Bassora on the Gulf of Persia, whence its name. It is in irregularly shaped pieces, white or yellow, and intermediate in its transparency between gum Arabic and gum tragacanth. Only a small portion is soluble in water. The insoluble portion is a peculiar principle, called *Bassorin*. It is not used in medicine; but bassorin enters into the composition of several substances.

BASSORIN, see *Bassora gum*.

BASSWOOD, *Tilia Americana*.

BASTARD, (Celt. *bas* or *boas*, 'fornication,' and *tardd*, 'source,') *Illegitimate*.

BATA, *Musa Paradisiaca*.

BATA'TA DE PUR'GA. The Brazilian and commercial name of the purgative, feculent, and gum-resinous roots of two plants of the family *Convolvulaceæ*, the one called *Jeticum*, *Meehoacan*, *Convolvulus mechoacan'na*, the *Piptostegia Pisonis* of Von Martius; the other *Convolvulus operculatus*, the *Ipomœa seu Piptostegia operculata* of Von Martius.

BATA'TAS. The inhabitants of Peru gave this appellation to several tuberous roots, especially to *Convolvulus Batatas* or *Sweet Potato*. Our word *Potato* comes from this. See *Solanum tuberosum*.

BATATAS EDULIS, *Convolvulus batatas*.

BATEMAN'S PECTORAL DROPS, see *Pectoral Drops*, *Bateman's*.

BATES'S ALUM WATER, *Liquor aluminis compositus*—*b. Anodyne Balsam*, *Linimentum saponis et opii*—*b. Camphorated Water*, *Lotion*, camphorated, of Bates—*b. Collyrium*, see *Lotion*, camphorated, of Bates.

BATH, (Sax. *bað*,) *Bal'neum*, *Balane'um*, *Lou'tron*, (Old Eng.) *Baine*, (F.) *Bain*. Immersion, or stay, for a longer or shorter duration, of the whole or a part of the body, in some medium, as water. Act of plunging into a liquid, sand, or other substance, in which it is the custom to bathe, *Plunge Bath*. Also, the vessel in which the water is put for bathing. Also, a public or private establishment for bathing. The common water-bath, used for hygienic as well as for therapeutical purposes, is the *Bain simple* ou *hygiénique* of the French.

In *Pharmacy*, a vessel, placed over a fire, and filled with any substance, into which another vessel is placed, containing matters for digestion, evaporation, or distillation.

BATH, *Ac'id*, *Bal'neum ac'idum* (*Acid. muriat.* ℞ij; *Aquæ* cong. lvi. One half, one third, or one fourth the quantity of acid is more frequently employed.)

BATH, *Acid*, *Scott's*, see *Scott's Acid Bath*—*b. Air*, hot, see *Bath*, hot—*b. Air*, warm, see *Bath*, hot.

BATH, *Al'kaline*, *Bal'neum alkali'num*. This may be made of half a pound or a pound of *pearl-ash* or *carbonate of soda*, to sixty-six gallons of water.

BATH, *An'im'al*, *Bal'neum animal'e*, consists in wrapping an animal recently killed, or its skin, around the body, or some part of it.

BATH, *Antipso'ric*, *Bal'neum antipso'ricum*. Recommended in cases of itch and other cutaneous diseases. (*Potass. Sulphuret.* ℞iv, *Aquæ* cong. lx.)

BATH, *Antisyphilit'ic*, *Bal'neum antisyphilit'icum*, *Mercur'ial bath*. Made by dissolving from

two drachms to an ounce of the corrosive chloride of mercury in sixty gallons of water.

BATH, ARM, *Brachil'vium*. A bath for the arm.

BATH, COLD, see Bath, hot—b. Cool, see Bath, hot.

BATH, DRY, is one made of ashes, salt, sand, &c. The ancients used these frequently for therapeutic purposes.

BATH, EARTH, *Arenatio*—b. Eastern, see Stove.

BATH, ELECTRIC, (F.) *Bain électrique*, consists in placing the person upon an insulated stool, communicating, by a metallic wire, with the principal conductor of the electrical machine in action. The Electric Bath produces general excitement of all the functions, and especially of the circulation and secretions.

BATH, FOOT, *Pedilu'vium*, (F.) *Bain de Pied*, a bath for the feet.

BATH, GELATINOUS, *Bal'neum gelatino'sum*. Made by dissolving two pounds of gelatin or glue in a gallon of water. A gelatino-sulphurous bath may be made by adding a pound of glue, previously dissolved in water, to the sulphur bath.

BATH, GENERAL, (F.) *Bain Entier*, is one in which the whole body is plunged, except the head; in contradistinction to the partial bath, *Merobalane'um*, *Merobal'neum*.

BATH, HALF, *Semich'pium*, *Excathis'ma*, *Inses'sio*, *Inses'sus*, is one adapted for half the body. One, for receiving only the hips or extremities, is also so called.

The *Sitz-bath*, (G.) *Sitzbad*, of the hydropathists, is a tub of cold water, in which the patient sits for a variable period.

BATH, HAND, *Manuil'vium*, (F.) *Bain de Main* ou *Manulvee*, is a bath for the hands.

BATH, HEAD, *Capitul'vium*, (F.) *Bain & Tête* ou *Capitilve*, a bath for the head.

BATH, HIP, *Coxal'vium*, (F.) *Bain de Fau-teuil*, *Bain de Siège*, is one in which the lower part of the trunk and upper part of the thighs are immersed.

BATH, HOT, *Bal'neum Cal'idum*, *Zestolu'sia*, *Thermolu'sia*, (F.) *Bain chaud*, is a bath, the temperature of which is 95° and upwards; the WARM BATH from 92° to 98°; the TEPID BATH, (F.) *Bain Tiède*, *Bal'neum tep'idum*, from 85° to 92°; the TEMPERATE BATH, (F.) *Bain tempéré*, from 75° to 85°; the COOL BATH, (F.) *Bain frais*, from 60° to 75°; the COLD BATH, *Bal'neum frig'idum*, *Frigida'rium*, (F.) *Bain froid*, *Bain très froid*, (of some), from 30° to 60°; and the VAPOUR BATH, *Bal'neum vapo'ris*, (F.) *Bain de Vapeur*, *Étuve Humide*, from 100° to 130°, and upwards. See Vaporarium. A WARM-AIR BATH, or HOT-AIR BATH, consists of air the temperature of which is raised. See Stove.

BATH, MEDICATED, *Bal'neum medica'tum*, (F.) *Bain médicinal*, is a bath, formed of decoctions or infusions of vegetable substances, or of any ingredient introduced into the water for therapeutic purposes.

BATH, MERCURIAL, Bath, antisymphilitic—b. Mud, see *Boue des Eaux*—b. Nitro-muriatic acid, Scott's acid bath—b. Plunge, see Bath—b. Roman, see Stove—b. Russian, see Vaporarium.

BATH, SAND, *Bal'neum Are'næ*, (F.) *Bain de Sable*, consists of a vessel filled with sand, and placed over the fire. Into this vessel, the one is put which contains the substance to be evaporated. See Psammismus.

BATH, SEA, *Bal'neum mari'num* seu *Ma'ris*, *Thalassolu'trum*, (F.) *Bain de Mer*. A sea-water bath.

BATH, SHOWER, *Implu'vium*, *Hydroco'nium*, is one in which the water is made to fall like a shower on the body. See Douche.

BATH, SITZ, see Bath, half.

BATH, SPONGE. The thorough application of water to the surface by means of a sponge.

BATH, STEAM, may be formed by introducing steam into a properly closed vessel in place of water, as in the water bath.

BATH, SUCCESSION, *Transit'ion bath*. A term applied to the rapid succession or transition from a cold to a warm or hot bath, or conversely.

BATH, SULPHUR, *Bal'neum Sulphu'ris*, *Sulphuretted bath*. A bath much used in psora, and other chronic cutaneous affections. It may be composed of two ounces of diluted sulphuric acid, and eight ounces of sulphuret of potassium added to each bath; or simply of the sulphuret of potassium.

BATH, SULPHUR VAPOUR, see Sulphurous Acid—b. SWEATING, *Achicolum*.

BATH, TAN. An astringent bath, prepared, at times, by boiling two or three handfuls of ground oak-bark,—such as is used by tanners—in two or three quarts of water, for half an hour, and then adding the decoction to the water of the bath.

BATH, TEMPERATE, see Bath, hot—b. Tepid, see Bath, hot—b. Transition, Bath, succession—b. Turkish, see Stove—b. Vapour, see Bath, hot, and Vaporarium—b. Warm, see Bath, hot.

BATH, WATER, *Bal'neum Ma'ris* seu *Ma'ris*, (F.) *Bain Marie*, in chemistry, consists of a vessel filled with boiling water, or salt water, in which the vessel is placed, that contains the substance to be evaporated.

Bathing is much employed in the treatment of disease. The cold bath, especially the cold sea bath, is a sedative and indirect tonic: the warm bath a relaxant; and the hot bath a stimulant.

The regular use of the bath is extremely conducive to health; but if too much indulged in, it is apt to produce injurious effects.

BATH ALUM SPRINGS, see Virginia, Mineral Waters of.

BATH, MINERAL WATERS OF, *A'qua Batho'nica* vel *Bad'iaz*, *A'que Sol'is*, *A'que Bad'igæ*. Celebrated thermal springs at Bath, in England. They contain but little impregnation, and are chiefly indebted to their temperature from 112° to 117° Fahrenheit, for their utility. The main ingredients are sulphate of lime, chloride of sodium, sulphate of soda, carbonate of lime, protoxide of iron, free carbonic acid and azote.

These waters are employed in the most heterogeneous cases; and are serviceable where the simple thermal springs are indicated, as in rheumatism, paralysis, &c.

BATH SPRINGS, in Berkley County, Virginia, is a mild carbonated water: temperature 73° Fah. It is a useful bath in cutaneous affections, and is said to contain some of the salts of lime and magnesia.

BATHER, *Balnea'rius*, *Balnea'tor*, (F.) *Baigneur*. One who bathes. Anciently, the name was given to those that administered baths to the diseased,—the *Étuvistes* of the French. At the present day, in remote districts in Germany, the country people call their medical practitioners Bäder or 'bathmen,' and Feldscheeren or 'army barbers.'

BATHING, see Bath.

BATH/MIS, *Bath'mus*, (*Ba'dus*, 'base, support.') The cavity of a bone, which receives the eminence of another; and especially the two *Fossettes* at the inferior extremity of the humerus into which the processes of the ulna are received, during the flexion and extension of the fore-arm.

BA'THRON, (*Ba'dron*, 'a bench,') *Ba'thrum Hippoc'ratis*, *Scam'num Hippoc'ratis*, (F.) *Banc d'Hippocrate*. An instrument, used for the extension of a limb, in cases of fracture or luxation.

The description of it is found in Galen, Oribasius, and Scultetus, with a figure.

BATHRUM HIPPOCRATIS, Bathron.

BATIA, Retort.

BATISSE, MINERAL WATERS OF. Batisse is three leagues from Clermont, in France. The water is tepid, and contains subcarbonate and sulphate of soda, sulphates of lime and iron, muriate of magnesia, and carbonate of lime.

BATONS DE CASSE, see Cassia fistula.

BATOS, (βατος,) Rubus Idæus.

BATRACHIUS, (βατραχος, 'a frog,') Ranula.

BATTALISM'US, *Battarism'us*, (βατταρισμ'ος, 'to stammer,') Balbuties. Stammering with incapacity to pronounce the R.

BAT'TALUS, (βατταλος,) *Bat'tarus*. A stammerer, a stutterer.

BATTARISMUS, (βατταρισμος,) Battalismus.

BATTARUS, (βατταρος,) Battalus.

BATTATA VIRGINIANA, (see *Batatas*), Solanum tuberosum.

BATTEMENS DOUBLES, (*battre*, 'to beat,') See *Bruit du Cœur fétal*.

BATTEMENT, Pulsation.

BAUDRICOURT, MINERAL WATERS OF. Baudricourt is a town of France, two leagues and a half from Mirecourt. The waters are sulphureous.

BAUDRUCHE, ('goldbeaters' skin,' from old (F.) *baudroyer*, 'to dress leather,') Condom.

BAUHIN, VALVE OF, *Valve* of TULPIUS, V. of FALLOPIUS, V. of VAROLIIUS, *Illeo-caecal Valve*, *Ileo-caecal Valve*, *Ileo-caecal Valve*, *Val'cula Col'i* seu *Col'ic* seu *Tulpii*, *Operculum Ilei*, *Sphincter Ilei*. This name is given to the valve situate transversely at the place where the ileum opens into the cœcum, and which Bauhin says he discovered at Paris, in 1759. It had, however, been previously described by several anatomists; as by Vitulus Vidius, Postius, &c.

BAUME, Balsam—b. d'Acier, Balsam, chalybeate—b. d'Aiguilles, Balsam, chalybeate—b. d'Amérique, see *Toluifera balsamum*—b. Apoplectique, Balsam, apoplectic—b. d'Arcæus, Balsam of Arcæus, and Unguentum elemi compositum—b. Aromatique, Balsam, aromatic—b. Benjoin, Benjamin—b. Blanc, see *Amyris Opobalsamum*—b. du Brésil, Copiba—b. de Canada, see *Pinus balsamea*—b. de Cannelle, Laurus cinnamomum—b. de Carpathie, Balsam of Carpathia—b. de Carthagène, see *Toluifera balsamum*—b. de Constantinople blanc, see *Amyris opobalsamum*—b. de Copahu, Copiba—b. Cordiale de Sennerte, Balsam, cordial, of Sennertus—b. d'Enu à feuilles ridées, *Mentha crispa*—b. de Feuillet, Balsam, green, of Metz—b. de Fioraventi spiritueuse, Balsam, spirituous, of Fioraventi—b. de Fourcroy ou de Laborde, Balsam of Fourcroy or Laborde—b. des Feuilles, Asphaltum—b. de Guluad, see *Amyris opobalsamum*—b. de Geneviève, Balsam of Geneviève—b. Grand, Tanacetum balsamita—b. du Grand Caire, see *Amyris opobalsamum*—b. Hydriodaté, Balsam, ioduretted—b. Hypnotique, Balsam, Hypnotic—b. Hystérique, Balsam, hysteric—b. des Indes, see *Myroxylon Peruiferum*—b. Ioduré, Balsam, ioduretted—b. des Jardins, *Mentha viridis*—b. de Lusanne, Balsam, ioduretted—b. de Lucatel, Balsam, Lucatelli's—b. Marie, see *Fagara octandra*—b. de Momie, Asphaltum—b. Nervin, Balsam, nervous—b. de Peron, see *Myroxylon Peruiferum*—b. de Saint-Thomas, see *Toluifera balsamum*—b. du Samaritain, Balsam of the Samaritan—b. Saxon, Balsam, Saxon—b. de Soufre, Balsam of sulphur—b. de Sympathie, Balsam of sympathy—b. de Tolu, see *Toluifera balsamum*—b. de Vanille, Vanilla—b. Tranquille, Balsam, tranquil—b. Vert, see *Fagara octandra*—b. Vert de Metz, Balsam, green, of Metz—b. de Vie d'Hoffman, Balsam of Life, of Hoffman

—b. de Vie de Lelidère, Tinctura aloes composita—b. Vrai, see *Amyris opobalsamum*—b. Vulnéraire de Minderer, Balsam, vulnerary, of Mindererus.

BAUMIER, Balsamier.

BAUNSCHEIDTISM'US. A kind of acupuncture, proposed by an ignorant, unprofessional person, of the name of Baunscheidt, a German, which consists in impelling on the skin, by means of a spring, a brush of wires, but not so as to draw blood, and then rubbing a secret stimulating fluid into the part. It was suggested to the proposer by the relief which the bites of gnats appeared to afford him in an arthritic affection.

BAURAC, (Arab.) Nitre, or salt in general. From this word comes Borax.

BAURIN, MINERAL WATERS OF. Baurin is a village, four leagues from Roye, department of Somme. The waters are strongly chalybeate.

BAVE (F.), *Saliva ex ore pueri*, *Spuma*, *Humor Salivus*. Frothy, thick, viscid saliva, issuing from the mouth. This drieth or slac'ering, we see in children, old people, &c. The term is, also, applied to the frothy liquid, which flows from the mouth of rabid animals. Sauvages uses it synonymously with salivation.

BAVEUX, Baveuse, an epithet, occasionally applied by the French to the spongy flesh of a wound, which suppurates, and exhibits but little tendency to heal.

BAY, CASTOR, *Magnolia glauca*—b. Rose, *Rhododendron chrysanthemum*—b. Rose, American, *Rhododendron maximum*.

BAY RUM, *Spiritus myrciæ*.

BAY, SWEET, Laurus—b. White, *Magnolia glauca* and *M. macrophylla*.

BAYBERRY BARK, see Wax, myrtle—b. Tallow, see Wax, myrtle—b. Tree, see *Spiritus myrciæ*.

BDALSIS, (βδαλσις,) Sucking.

BDELLA, (βδέλλα,) Hirudo.

BDELLIUM, (βδέλλον,) *Myrrha imperfecta*, *Bol'chou*, *Madelon*, *Bal'chia*. A gum-resin, brought from the Levant and India, and supposed to be obtained from a species of *Amyris*, little known. It is solid, brittle, of a deep brown colour, of an acrid and bitter taste, and sweet odour. It was much vaunted by the ancients, but is now little employed. It is considered, by some, to be the *Onycha* of the Scriptures, which was burnt as incense.

Two different gum-resins have been in the shops, distinguished by the names *Indian* and *African bdellium*. Dr. Royle was informed that the former was obtained from *Amyris Commiph'ora*, growing in India and Madagascar. The latter is said to be from *Heudelotia Africana*, which grows in Senegal.

BDELLOMETER, (βδέλλα, 'a leech,' or βέλλω, 'I suck,' and μέτρον, 'measure,') An instrument, proposed as a substitute for the leech; inasmuch as we can tell the quantity of blood obtained by it, whilst we cannot by the leech. It consists of a cupping-glass, to which a scarificator and exhausting syringe are attached.

BDELLUS, (βδέλλος,) Fart.

BDELYGMIA, (βέλγυμα,) Fart.

BDELYGMUS, (βέλγυμος,) Fart.

BDESMIA, (βέσμα,) Flatulence.

BDOLOS, (βδολος,) Fart.

BEAD TREE, *Melia Azedarach*.

BEAL, (like Boil,) Phlegmon, to Suppurate

BEALING, Pregnant, Suppuration.

BEAN, (Sax. bean,) Faba—b. Bush, *Phaseolus*—b. Calabar, Ordeal nut—b. Carthagena, *Habilla de Carthagena*—b. Egyptian, *Nymphaea nelumbo*—b. French, *Phaseolus vulgaris*—b. Garder, common, *Vicia faba*—b. Indian, *Catalpa*—b. Kidney, *Phaseolus vulgaris*—b. Malacca, *Avicennia*

tomentosa—b. Pontic, *Nymphæa nelumbo*—b. Red, *Abrus precatorius*—b. Sacred, *Nelumbium luteum*—b. Snap, *Phaseolus*—b. St. Ignatius's, *Ignatia amara*—b. String, *Phaseolus*—b. Tonka, *Dipterix odorata*—b. Wild, *Apios tuberosa*.

BEAN TREE, *Catalpa*—b. Tree, white, *Cratægus aria*.

BEAN TREFOIL TREE, see *Cytisine*.

BEARBERRY, *Arbutus uva ursi*.

BEARD, (Sax. *beapd.*) *Barba*, *Polygon*, *Geni'on*, *Genys*, *Barbitium*, (F.) *Barbe*. The hair which covers a part of the cheek, the lips, and chin of the male sex, at the age of puberty.

BEARDLESS, *Imberbis*.

BEAR'S BREECH, *Acanthus mollis*—b. Foot, *Helleborus foetidus*—b. Friglit, *Heptallon graveolens*—b. Grass, *Yucca filamentosa*—b. Whortleberry, *Arbutus uva ursi*.

BEARWEED, *Veratrum viride*.

BEASTINGS, (Sax. *býp̃iñz*), *Colostrum*.

BEATLINGS, *Colostrum*.

BEATING OF THE HEART, see *Heart*.

BEAUGENCY, MINERAL WATERS OF. Beaugency is a quarter of a league from Orleans. The waters contain subcarbonate of soda, iron, magnesia, and lime. They are tonic and aperient.

BEAUMONT ROOT, *Gillenia trifoliata*.

BEAUVAIS, MINERAL WATERS OF. These waters are chalybeate. Beauvais is in Picardie, France.

BEAVER, *Castor fiber*—b. Tree, *Magnolia glauca*, *M. macrophylla*—b. Wood, *Celtis occidentalis*, *Magnolia glauca*.

BEBEERIA, see *Bebeeru*.

BEBEERINE, see *Bebeeru*.

BEBERU, *Sipeeri*. A tree of British Guiana, the bark of which yields two alkalies—*Bebeerine*, *Bebeerina*, *Bebeeria*, *Beberia*, or *Bibeerine*, and *Sipeerine*; and in its properties resembles the Cinchonas. It has been referred to *Nectandra Rodiei* seu *Rodiei*. *Ord.* Laurineæ. The timber of the tree is known to shipbuilders by the name *bibiru* or *green-heart*. The Sulphate of *Bebeeria*, *Beberia* sulphas (Ph. B. 1864), is employed in the same cases as sulphate of quinia. Warburg's *Feber Drops*, *Tinctura antifebrilis Warburgi*, an empirical antiperiodic preparation, have, by some, been considered to be a tincture of the seeds of the *Bebeeru*, but this is questionable.

BEBERIA, see *Bebeeru*.

BEC (F.), *Ros'trum*, *Beak*. This name has been applied to various parts.

BEC CORACOÏDIEN (F.), *Cor'acid beak*, is the end of the coracoid process.

BEC DE CULLIER, *Hamulus*. An instrument used for the extraction of balls. It consists of an iron rod, 7 or 8 inches long, having at one extremity a small cavity, into which the ball is received to be drawn outwards. See *Cochleariformis*.

BEC DE GRUE MUSQUÉ, *Geranium Moschatum*—b. de *Grue Robertin*, *Geranium Robertianum*—b. de *Lièvre*, *Harelip*.

BEC DE LA PLUME À ÉCRIRE, (F.) *Beak of the Calamus Scriptorius*, is a small cavity at the superior part of the medulla oblongata, which forms part of the 4th ventricle.

BEC, (L.) MINERAL WATERS OF. Bee is six leagues from Rouen, in Normandy. The water is strongly chalybeate.

BECCABUNGA, *Veronica Beccabunga*.

BECHÆSTHESIS, (βῆξ, gen. βήξ, 'cough,' and αἰσθησις, 'sensation.') The excitement or desire to cough.

BÊCHE DE MER, *Biche de mer*.

BECHIA, (βήχια,) *Tussis*.

BECHIAS, (βήχιας,) *Tussis*.

BECHICS, *Be'chia*, (βήχια,) *Becha*, *Be'chia*, *Be'chita*, (F.) *Béchiqnes*. Medicines adapted for allaying cough.

BECHITA, *Beechie*.

BECHUM, (βήχων,) *Tussilago*.

BECHORTHOPNEA, βῆξ, gen. βήξ, 'cough,' ὀρθας, 'upright,' and πνεω, 'I breathe.') *Pertussis*.

BECUIBA, *Ibicuiba*.

BED, HYDROSTATIC, Water bed—b. Water, *Arnott's*, see *Water bed*.

BED'EGAR, *Bed'eguar*, *Bed'eguard*, *Spon'gia Cynos'bati*, *Fungus Rosa'rum*, *F. Cynos'bati*, (F.) *Pomme mousseuse*, *Éponge d'églantier*. An excrecence, which makes its appearance on different species of wild roses, and which is produced by the puncture of a small insect.—*Cyn'ipsa Ro'sæ*. It was formerly employed as a lithontriptic and vermifuge, but is not now used. It was slightly astringent.

BEDFORD, MINERAL WATERS OF. Bedford is a village, situate on the great Western Turnpike road from Philadelphia to Pittsburg, a few miles east of the chief elevation of the Alleghany mountains. The "Springs" are about a mile and a half from the village. They are saline and sulphureous. The most celebrated—the "Min'eral Spring" or "An'derson's Spring"—contains carbonic acid, sulphate of magnesia, chlorides of sodium and calcium, and carbonate of iron, but none of these articles in great quantity; hence, the main action of the water is diuretic. At some distance from the Springs, there is a chalybeate water; and about 10 miles S. W. of Bedford, at Milliken's Cove, a strong sulphureous spring. The climate of Bedford Springs is agreeable. During the heat of summer, the nights are generally cool. In the morning of July 14, 1853, the author marked the thermometer, at five o'clock, A. M., at 45°. Three days before, however, it was 70° at the same hour.

BEDLAM, (from Bethlehem, a religious house in London, converted into a lunatic hospital,) *Morotrophium*.

BEDLAMITE, *Insane*.

BEDRIDDEN, *Clinic*.

BEDSTRAW, *Galium verum*—b. Ladies, greater, *Galium mollugo*, *Galium verum*—b. Ladies, rough, *Galium asprellum*.

BEE, (Sax. *beo*.) *A'pis*, *A. mellifica* seu *domestica*, *Melis'sa*, *Meli'ta*, (F.) *Abeille*. This insect was formerly exhibited, when dried and powdered, as a diuretic.

BEE IN THE BONNET, see *Insanity*.

BEEBREAD, *Propolis*.

BEECH, *Fagus sylvatica*—b. Albany, *Pteropora Andromedea*—b. Drop, *Orobanche Virginiana*—b. Drops, false, *Hypopitys lanuginosa*.

BEECH MAST, see *Fagus sylvatica*—b. Nut, see *Fagus sylvatica*.

BEDDY'S EYES, *Viola tricolor*.

BEEF ESSENCE, see *Beef tea*.

BEEF TEA, *Jus bov'num*. An infusion of beef, much used in debilitating maladies, and in convalescence. It may be made as follows: Take two pounds and a half of lean beef; cut it, in small pieces, into three parts of water in an earthen pipkin: let this simmer, but never boil, until the liquor is consumed to a pint and a half: then strain carefully. It ought to be entirely free from fat or grease.

Ess'ence of Beef—as it has been called—may be made by putting a pound of good beef, freed from fat, and cut into small pieces, into a porter-bottle, corking lightly. The bottle must be put into boiling water, and kept there until the water has been boiling at least half an hour. As the boiling goes on, the cork may be inserted a little more tightly, to retain the contents of the bottle. The juices

of the beef are thus separated, and constitute the 'essence,' which may be seasoned to the taste. It contains much nutriment.

BEEH, Centaurea behen.

BEER, ([W.] *bir*.) Cerevisia—b. Black, see Fall-tranck—b. Chowder, see Chowder—b. Jews', see Pinus sylvestris—b. Pissisew, see Pyrola umbellata.

BEER SPRINGS, see Water, Mineral (gaseous.)

BEER, SPRUCE, *Cerevisia 'Abietis*, may be prepared as follows. Take of *Essence of Spruce*, half a pint; *Pimento*, bruised; *Ginger*, bruised; *Hops*, of each, four ounces; *water*, three gallons. Boil for five or ten minutes; strain, and add of *warm water* eleven gallons; *yeast*, a pint; *molasses*, six pints. Mix and allow to ferment for twenty hours. Used as an agreeable drink in summer.

BEER, TAR, see Pinus sylvestris.

BEEST, Colostrum.

BEESTINGS, (Sax. *býrting*.) Colostrum.

BEET, Beta.

BEETLEWEED, Galax aphylla.

BÉGALEMENT, Balbuties.

BEG'MA, gen. *Beg'matis*, (*Bnyua*),—according to some, *Breg'ma*, (*βρεγμιν* or *βρεγμιν*, 'to expectorate after coughing.') Coughing; also, the sputum or expectorated matter.

BEGO'NIA, (after Michel Begon, a French botanist, in St. Domingo.) The *Begonia grandiflora* and *B. tomentosa* have astringent roots, which are used in Peru in cases of hemorrhage, scurvy, low fevers, &c.

BÉGUE, Balbus.

BEHEN ABIAD, Centaurea behen—b. Album, Centaurea behen—b. Officinum, Cucubalus behen—b. Rouge, Statice limonium—b. Vulgaris, Cucubalus behen.

BEIAHALALEN, Sempervivum tectorum.

BEIDELSAR, Asclepias procera.

BEIST (Sc.), ([Sax.] *býrt*.) Colostrum.

BEISTINGS (Sc.), ([Sax.] *býrtung*.) Colostrium.

BEISTYN, Colostrum.

BEJUJO, Habilla de Carthagena.

BELA, Covolam.

BELA-AYE or BE-LAHE. A tonic and astringent bark of a Madagascar tree. Du-petit-Thouars and Sonnerat think it may be substituted for the Simarouba.

BELADAMBOC. A species of convolvulus of the Malabar coast, which contains an acrid milky juice. From this a liniment is formed with oil and ginger, which is used against the bites of rabid animals.

BE-LAHE, Bela-aye.

BELA-MODAGAM. A kind of *Scaevola* of the Malabar coast, the leaves of which are considered diuretic and emmenagogue.

BELANDRE, (F.) ([D.] *bijlander*, from *bij*, 'near,' and *land*, 'land; 'a vessel which coasts along the shore.') A litter surrounded with curtains, in which patients are sometimes carried to hospitals.

BELCHING, Eructation.

BELD, Athrix.

BELEMNOID, Belenoid.

BELEMNOIDES PROCESSUS, Styloid processes.

BEL'ENOID, BEL'ONOID, BEL'EMNOID or BEL'OID, *Belenoides* or *Belemnoides Processus*, (*βελος*, 'an arrow,' and *ειδος*, 'shape.') This name has been given to styloid processes in general—*Processus belenoides*.

BÉLESME, see Bellesme.

BELSON, Balsam, Muscæda frondosa.

BELILLA, Muscæda frondosa.

BELINUM, Apium Graveolens.

BELI OCLUS, Belloculus.

BELL, CANTERBURY, Campanula trachelium.

BELLADONE, Atropa belladonna.

BELLADON'NA, (in (I.) 'beautiful lady,' in the Pharmacopœia of the United States, is the official name of the leaves of Atropa Belladonna.

BELLADONNA BACCIFERA, Atropa belladonna—b. Trichotoma, Atropa belladonna.

BELLADONNIN, Atropine.

BELLARIA, (*bellus*, 'sweet,') Dessert.

BELLE-BLOME, ('beautiful bloom,') Narcissus pseudo-narcissus.

BELLE DAME, ('beautiful lady,') Atropa belladonna.

BELLEGU, Myrobalanus.

BELLEREGI, Myrobalanus.

BELLÈSME, MIN'ERAL WA'TERS OF.

Bellème is about three leagues from Montagne, in France. The waters are chalybeate.

BELLEY, MIN'ERAL WA'TERS OF. The waters at Belley, department of Ain, in France, are saline aperients.

BELLFLOWER, Narcissus pseudo-narcissus.

BELLIDOIDES, (*bellis*, and *ειδος*, 'resemblance,') Chrysanthemum leucanthemum.

BELLINI, TUBES OF, see Uriniferous tubes.

BEL'LIIS, *Bel'lia* ('pretty'), *Bel'lia perennis* seu *mi'nor* seu *horten'sis*, *Syn'phytum mi'nimum*, *Bruise'wort*, *Com'mon Da'isy*, *Day's eye*, (Sc.) *Ene-gowan*, (Prov.) *Bairnwort*, (F.) *Paquerette vivace*, *petite Marguerite*. Ord. Compositæ. The leaves and flowers are rather acrid. They were, at one time, considered to cure different species of wounds. See *Osmitopsis asterisoides*.

BELLIS HORTENSIS, Bellis—b. Major, Chrysanthemum leucanthemum—b. Minor, Bellis—b. Perennis, Bellis—b. Pratensis, Chrysanthemum leucanthemum.

BELLIT, Athrix.

BELL MET'AL, *Cal'cocos*, (?), (F.) *Airain*, *Métal des cloches*. An alloy of copper, zinc, tin, and a small quantity of antimony, used for making bells. The mortars of the apothecary are often formed of this material. They require to be kept clean, to avoid the formation of verdigris.

BELLOC'ULUS, *Beli Oculus*. A kind of gem, which the Assyrians considered efficacious in the cure of many diseases. They imagined that the figure of an eye could be seen in it, and hence its name, *Bel's Eye*, from the god Bel.

BELLON, Colic, metallic.

BELLONAT, (Prov.) Asthmatic.

BELLITAS, see Ilex major.

BELLOWS' SOUND, *Bruit de soufflet*—b. s. Encephalic, see *Bruit de soufflet*.

BELLOWS' SOUND, F'u'NIC, *Funic souffle*, (F.) *Souffle funique*, a single murmur of the bellows kind, synchronous with the first sound of the heart; heard by some observers, and referred by them to diminished calibre of the umbilical arteries, either by pressure or stretching of the funis, or both.

BELLOWS' SOUND, PLACENTAL, *Bruit placentaire*.

BELLWORT, SMALLER, *Uvularia perfoliata*.

BEL'LY, *Ven'ter*, (from *lr. bolg*, 'the belly, a bag or pouch' [?]). At the present day, the abdomen, (Sc. and Prov.) *Wayne*, *Wame*, *Weam*, *Wem*. Formerly, all the splanchnic cavities were called *bellies*;—the lower belly, *venter in'finus*, being the abdomen; the middle belly, *venter me'dius*, the thorax; and the upper belly, *venter su-pre'mus*, the head. Also, the womb. See *Abdomen*, *Uterus*, *Venter*.

BELLY-ACHE, Colica.

BELLY-ACHE, DRY, Colica metallica.

BELLY-ACHE ROOT, *Angelica lucida*.

BELLY-BAND, Belt, Russian.

BELLY-BOUND, Constipated.

BELLY-GOD, Glutton.
 BELLY-HARM, Colic.
 BELLY, POT, Physconia.
 BELLY-THRA, Colic.
 BELLY-TIMBER, Aliment.
 BELLY-WARK, Colic.
 BELMUSCHUS, Hibiscus abelmoschus.
 BELNILEG, Myrobalanus.
 BELOID, (βελος, 'an arrow,' and εἶδος, 'shape.')
 Belenoid.
 BELOIDES PROCESSUS, Styloid processes.
 BELONE, (βελονη,) Needle.
 BELONODES, Styloid, see Belenoid.
 BELONOID, (βελονη, 'a needle,' and εἶδος, 'shape.') Belenoid.
 BEL'S EYE, Belloculus.
 BELT, RUSSIAN, (Sax. belt,) *Ventral'le*,—vulgarly, *Bellyband*, *Abdom'inal support'er*, (F.) *Supporteur abdominal*. A broad bandage applied to the abdomen, so as to support and make methodical pressure upon it. Different forms have been termed *obstetric binders*, *utero-abdominal supporters*, &c.
 BELUL'CUM, (βελος, 'a dart,' and ἔκω, 'I draw out.') An instrument used for extracting darts or arrows. Many instruments of this kind have been employed by surgeons.
 BELZOE, Benjamin.
 BELZOIM, Benjamin.
 BELZOINUM, Benjamin.
 BEN, Guilandina moringa —b. of Judæa, Benjamin—b. Nut, Guilandina moringa.
 BENATH, Pustule.
 BENEDICTA SYLVESTRIS, Geum rivale.
 BENEDICTUM LAXATIVUM, ('a blessed laxative,') *Confectio sennæ*.
 BÉNÉFICE DE LA NATURE, ('benefit of nature,') *Beneficium naturæ* —b. *de Ventre*, see *Beneficium naturæ*.
 BENEFIC'IUM NATU'RÆ, ('a benefit of nature,') (F.) *Bénéfice de la nature*. This term is used by the French pathologists for cases in which diseases have got well without medical treatment. With them, *Bénéfice de nature*, or *B. de ventre*, is synonymous also with *Al'vi proflu'vium*; —a spontaneous diarrhœa, often acting favourably either in the prevention or cure of disease.
 BENEL, Croton racemosum.
 BENE'OLENS, (*bene*, 'well,' and *olere*, 'to smell,') *Euo'des*, *Suave'olens*. A sweet-scented medicine, as gums, &c.
 BENG, Bangu.
 BENGAL ROOT, Cassumuniar.
 BENGALÉ INDORUM, Cassumuniar.
 BENGI, Hyoseyamus.
 BENIGN', *Benig'nus*, *Euth'es*, (F.) *Bénin*, *Bénigne*, (from *bonus*, 'good,') Diseases of a mild character are so called, as a *Benign Fever*, *Febris benigna impu'tris*, &c.: as well as medicines whose action is not violent.
 BÉNIN, Benign.
 BEN'JAMIN, corr. from *Ben'zoin*, *Benzo'inum*, (Ph. U. S.), *B. ve'rum*, *A'sa* or *As'sa odora'ta* seu *dul'cis*, *Ben'jui*, *Ben'juin*, *Ben'jooy*, *Benjo'inum*, *Bel'zoë*, *Bel'zoin*, *Ben'zoë*, *Styl'racis Benzo'ini Bal'samum*, *Li'quor Cyreni'acus*, *Cro'ton Ben'zoë*, *Ben of Judæ'a*, *A'cor Benzo'inus*, *Sal Ac'idum* seu *essentia'le* seu *volat'ilè Ben'zoë*, *Resi'na benzoë*, (F.) *Benjoin*, *Baume Benjoin*, *Asa doux*. A resinous, dry, brittle substance, the concrete juice of *Styl'rac Ben'zoin*, *Ar'bor Benivi*, *Lau'rus Ben'zoin*, of Sumatra. The odour is extremely fragrant, and taste slightly aromatic. It is principally used for the preparation of the acid which it contains. It is also employed in some vulnerary tinctures, and as an expectorant. Benzoic Acid, *Ac'idum Benzo'icum*, is obtained from it by sublimation. The purest Benjamin is in amygdaloid masses: hence called (F.) *Benjoin amygdaloïde*.

BENJAMIN BUSH, *Laurus benzoin*.
 BENJAMIN, FLOW'ERS OF, *Ben'zoic Ac'id*, *Ac'idum Benzo'icum* seu *Benzo'il'icum*, *Flo'res Ben'zoës* seu *Benzo'ini*, *Ac'idum Benzo'icum per sublimatio'nem*, (F.) *Acide Benzoïque*, *Fleurs de Benjoin*. This acid exists in all the balsams, but chiefly in Benzoïn, from which it is obtained by sublimation. It is in vanilla, canella, the urine of infants, and of herbivorous animals. Its odour is aromatic and fragrant; taste hot, slightly acidulous, and agreeable. The crystals consist of white, satiny flakes, slightly ductile. It is probably stimulant; and has been used, as such, in chronic catarrh; but it has little efficacy.
 BENJAOY, Benjamin.
 BENJOIN, Benjamin.
 BENJOINUM, Benjamin.
 BENJUI, Benjamin.
 BEN MOENJA. A Malabar tree. An alexipharmic decoction is made of its roots, in the country, which is much praised in cases of malignant fever. Its bark, boiled with *Calamus aromaticus* and salt, forms a decoction used in bites of poisonous serpents.
 BENNE, *Sesamum orientale*.
 BENNET, HERB, (from *benedictus*, 'blessed,') *Geum urbanum*, and *G. Virginianum*.
 BENOÎTE, *Geum urbanum* —b. *Aquatique*, *Geum rivale* —b. *des Ruiss'eaux*, *Geum rivale* —b. *de Virginie*, *Geum Virginianum*.
 BENZENE, Benzole.
 BENZIN, Benzole, see Anæsthetic.
 BENZOATE OF AMMONIA, *Ammonia benzoas*.
 BENZOE, Benjamin.
 BENZOENIL, Vanilla.
 BENZOIN, Benjamin—b. *Odoriferum*, *Laurus Benzoin*.
 BEN'ZOLE, *Ben'zin*, *Benzene*, *Phene*, *Hy'dru-ret of Phen'yl*. A constituent of coal-gas tar, obtained by distilling coal naphtha. It is a colourless, oleaginous liquid of agreeable odour. Sp. gr. 0·85. The vapour, when inhaled, acts as an anæsthetic. Mixed with four parts of lard it has been used, with advantage, in itch; and it is said to be very effective against parasites of the skin. It has, also, been employed in rheumatic and neuralgic cases, —a piece of cloth or paper saturated with it being applied to the part.
 BERBERINE, see *Oxyacantha Galeni*.
 BERBERIS, (from the Arab.,) *Oxyacantha Galeni* —b. *Canadensis*, see *Oxyacantha Galeni*.
 BERBINE, *Verbena officinalis*.
 BERCE, *Heracleum spondylium*.
 BERENDAROS, *Ocimum basilicum*.
 BERNICE, *Succinum*.
 BERENICIUM, *Potassæ nitras*.
 BERENISECUM, *Artemisia vulgaris*.
 BERGAMOT', *Bergamotè*, *Bergamot'ta*, (F.) *Bergamotte*. A small orange, of a very agreeable taste, and peculiar odour. From its rind an oil, *O'leum Bergamit'ii* (Ph. U. S.), is obtained, which is much employed as a perfume, and sometimes in medicine. See *Citrus mella rosa*.
 BERGAMOTTIER, see *Citrus mella rosa*.
 BER'IBERI, *Beribè'ria*, *Syn'clonus Beribè'ria*, *Indosyn'clonus*, *Paral'yis Ber'iberi*, (from *beri* in the Singalese language, which signifies 'weakness,' therefore, *beriberi*, 'great weakness.' This word is also said to be Hindustanee, and to mean a sheep.) *Beriberi* is an Indian disease, little known in Europe. It consists in debility and tremors of the limbs, —sometimes, indeed, of the whole body; with painful numbness of the affected parts, &c. —the patient walking doubled, and imitating the movements of sheep (?) Some authors have esteemed it rheumatic; others, paralytic; others, to be a kind of chorea. It is a form of cachexia, in which there is great muscular de-

bility and feebleness of the reparatory powers. It is, almost always, incurable; is rarely fatal; and is treated by exercise, stimulant friction, sudorifics, &c. It is sometimes called *Bar'biers*, but this would seem to be a different disease.

The *Aden* or *Ye'men Ul'cer* is said to occur in conditions of the system like that in Beriberi. It affects the leg, assuming a disagreeable appearance, and frequently sloughing.

BERICOCCE, (*Βερικοκκ.*) *Prunus armeniaca*.

BERKLEY SPRINGS, see Bath Springs.

BERLE NODIFLORE, *Sium*.

BERLUE, *Metamorphopsia*.

BERMUDAS, CLIMATE OF. Pulmonary invalids are occasionally sent to Bermuda, but the principal objection to a winter residence there, is the prevalence of strong winds; especially of the dry, sharp, and cold north-west winds, during the winter and spring. Still, it affords a good winter retreat for the phthisical, from any part of the United States, provided due care be selected in choosing a suitable locality. The neighbourhood of Hamilton has been strongly recommended with this view.

BERNARD THE HERMIT, *Cancellus*.

BERRIES, INDIAN, see *Menispermum coculus*—b. Turkey, yellow, see *Piper cubeba*.

BERS. A sort of electuary, composed of pepper, seed of the white hyoscyamus, opium, euphorbia, saffron, &c. The Egyptians used it as an excitant.

BERTIN, COLUMNS OF, *Columnæ Bertini*.

BERU, MINERAL WATERS OF. Beru is in Champagne, France. The waters are slightly chalybeate.

BERULA, *Sium nodiflorum*—b. *Angustifolia*, *Sium nodiflorum*.

BESASA, *Ruta*.

BESICLES, (*bis*, 'twice, and *eyclus*, 'a circle,' [?] or from *bis*, and *oculus*, 'an eye' [?]), Spectacles.

BESOIN, (*soin*, 'care,' [?]) Want—b. *de Respirer*, see Want—b. *de la Vie*, Necessary of life.

BESSANEM. A word used by Avicenna, for redness of the skin, limbs, and face, produced by the action of cold.

BESSON, } (from *bis*, 'twice,' and *homo*,
BESSONNE, } 'man,' [?]). See *Gemellus*.

BE'TA, (from Celt. *bett*, 'red.') The *Beet*, *Sic'ula*, (F.) *Bette*, *Betterave*. Family, *Chenopodææ*. *Sec. Syst.* *Pentandria Digynia*. A genus of plants, of which the following are the chief varieties:

BETA HYBRIDA, *Root of Scarc'ity*. Root red, outside; white, within. Very nutritive; yields sugar.

BETA VULGARIS AL'BA, *White Beet*. The root yields sugar, and the leaves are eaten as a substitute for spinach.

BETA VULGARIS RU'BRA, *Red Beet*. Root red and nutritive; yields a small quantity of sugar.

BETAYNE, *Betonica officinalis*.

BETTEL, *Piper Betel*, *Chavi'ca bettle*. A species of pepper, cultivated in several parts of India. The East Indians are in the habit of chewing the leaves with lime and areca; and they give the name *Betel* to this preparation. It is used in all the equatorial countries of Asia. *Betel* is said to be tonic and astringent. It is also called *Bette*, *Bêtre*, *Belle*. See *Areca*.

BETHROOT, *Trillium latifolium*—b. Broadleaf, *Trillium latifolium*.

BÊTISE, (*bête*, originally from (L.) *bestia*, 'silly,') *Dementia*.

BÊTOINE, *Betonica officinalis*—b. *des Montagnes*, *Arnica montana*—b. *des Savoyards*, *Arnica montana*.

BÊTON, *Colostrum*.

BETONICA AQUATICA, *Scrophularia aquatica*.

BETON'ICA OFFICINA'LIS, *B. purpurea*, *Vet-*

ton'ica, *Cestron*, *Psychotrophum*, *Veron'ica purpurea*, *Be'ony*, *Wood Be'ony*, (Old Eng.) *Betayne*, (F.) *Bêtoine*. Order, *Labiata*. *Sec. Syst.* *Didynamia Gymnospermia*. *Betony* was in much esteem amongst the ancients, who employed the flowers and leaves, in decoction, in gout, sciatica, cephalalgia, &c. (It was so called, according to Pliny, from being in great repute among the Vettones or Bettones, an ancient people of Spain.) Antonius Musa is said to have written a volume in praise of it; recommending it in no less than 47 different diseases. It has, however, little or no virtue. The leaves are said to be aperient, and the root emetic.

BETONICA PAULI, *Veronica*.

BETONY, *Betonica officinalis*—b. Paul's, *Lycopus sinuatus*, *Lycopus Virginicus*, *Veronica*—b. Water, *Scrophularia aquatica*—b. Wood, *Betonica officinalis*.

BÊTRE, *Betel*.

BÊTE, *Beta*.

BÊTTERAVE, (*beta* and *rapa*,) *Beta*.

BET'ULA AL'BA, *Semu'da*, *Birch*, (Sc.) *Birk*, (F.) *Bouleau commun*. Order, *Betulacæ*.

The young leaves are slightly odorous, astringent, and bitter. They are applied to wounds and ulcers. They have been regarded as antiscorbutic and anthelmintic. The tree furnishes a saccharine juice, which is considered antiscorbutic and diuretic.

BETULA EMARGINATA, *Alnus glutinosa*—b. *Glutinosa*, *Alnus glutinosa*.

BETULA L'ENTA, *Sweet Birch*, *Black Birch*, *Cher'ry Birch*, *Moun'tain Mahog'any*, is an American species, the bark and leaves of which have the smell and taste of *Gaultheria procumbens*. An infusion is sometimes made of them, and used as an excitant and diaphoretic. The volatile oil is nearly, if not wholly, identical with that of *Gaultheria*.

BEURRE, (contracted from *Beurrepur*, (L.) *butyrum*,) Butter—b. *de Bambouc*, Butter of bambou—b. *de Cacao*, Butter of cacao—b. *de Coco*, Butter of cocoa—b. *Végétale*, *Persea gratissima*.

BEUVRIGNY, MINERAL WATERS OF. Beuvrigny is in the vicinity of Bayeux, in Normandy. The water is chalybeate.

BEVERAGE, (from [F.] *breuvage*,) Drink.

BÊVUE, (*be*, *bis*, 'twice,' and *vue*, 'sight,') *Diplopia*.

BEX, (*βηξ*, gen. *βηχος*,) Tussis—b. *Convulsiva*, *Pertussis*—b. *Humida*, *Expectoration*—b. *Theorides*, *Pertussis*.

BEXIS, (from *bex*,) Tussis.

BEXU'GO. Under this name, a purgative root was formerly introduced into Europe from Peru. It is supposed to have been the root of a *Hippocratea*.

BEZ'OAR, *Bez'aar*, *Bez'ehard*, *Pa'zahar*, (Persian *pa*, 'against,' and *zahar*, 'poison,') *La'pis Bezoar'dicus*, *Cal'culus Bezo'ar*, *Enterolithus Bezoar'dus*, *Bez'oard*. A calculeous concretion, found in the stomach, intestines, and bladder of animals. Wonderful virtues were formerly attributed to these Bezoars. There were two great varieties: the *Bez'oar orient'al'e*, *Animal Bezoar'ticum orient'al'e*, formed in the fourth stomach of the gazelle of India (*Gazell'a Ind'ica*, or rather *Antilop'us cervica'pra*); and the *Bez'oar occident'al'e*, *Animal Bezoar'ticum occident'al'e*, found in the fourth stomach of the wild goat or *chamois* of Peru. These substances were esteemed to be powerful alexipharmics; but the former was the more valued. It was believed that no poison, and no eruptive, pestilential, or putrid disease, could resist its influence. As so many virtues were ascribed to it, other animal concretions were substituted for it; and factitious Bezoards were made of crab's eyes and claws, bruised and mixed with musk, ambergris, &c.

BEZOAR BOVINUM, (F.) *Bézoard de Bœuf*, *Bezoard of the beef*. A concretion formed in the fourth stomach of beeves; also, a biliary calculus found in the gall-bladder.

BEZOAR EQUINUM, *Bezoard of the horse*—b. *Hystrixis*, *Bezoard of the Indian porcupine*.

BÉZOARD D'ALLEMAGNE, *Ægagropila*.

BEZOARD OF CAYMAN. This was once much prized. It is now unknown.

BEZOARD OF THE CHAMOIS, and **B. OF THE HORSE**, *Bez'oar equinum*, *Hippolithus*, &c., exhibit their origin in the name.

BEZOARD OF THE DEER, *B. of the Lach'rymal Fossa of the Deer*, *Deer's Tears*. A moist, highly odorous, fatty matter, found below the anterior canthus of the orbit of the red deer—*Cervus elaphus*. It has been used, like castor, as an antispasmodic, in the dose of from 5 to 15 grains, two or three times a day.

BEZOARD OF THE INDIAN PORCUPINE, *Bez'oar Hystrixis*, *La'pis Porcinus* seu *Mahucensis*, *Petro del Porco*, (F.) *Bézoard de Porc-Épic*, was formerly the dearest of all the Bezoards, and was sold at an enormous price in Spain and Portugal.

BEZOARD, MINERAL, Antimonyum diaphoreticum—b. Vegetable, see Calapitte.

BEZOARDIC, *Bezoardicus*, (F.) *Bézoardique*; concerning the bezoard. Bezoardic medicines are those supposed to possess the same properties with the bezoard; as antidotes, alexiteria, alexipharmics, cordials.

BEZOARDICA RADIX, *Dorstenia contrajerva*.

BEZOARDICUM HUMANUM. Urinary calculi were formerly employed under this name as powerful alexipharmics.

BEZOARDICUM JUVIALE. A sort of greenish powder, used as a diaphoretic, and formed of antimony, tin, mercury, and nitric acid.

BEZOARDICUM LUNARE. A medicine formerly regarded as a specific in epilepsy, convulsions, megrim, &c. It was prepared of nitrate of silver, and butter of antimony.

BEZOARDICUM MARTIALE. A tonic medicine, used by the ancients in diarrhoea. It was prepared from the tritoxide of iron and butter of antimony.

BEZOARDICUM MERCURIALE. A medicine, formerly vaunted as an antisyphilitic, and prepared from the mild chloride of mercury, butter of antimony, and nitric acid.

BEZOARDICUM MINERALE; the deutoxide of antimony; so called because its properties were supposed to resemble those of animal bezoard.

BEZOARDICUM SATURNI. A pharmaceutical preparation, regarded by the ancients as antihysteria. It was formed of protoxide of lead, butter of antimony, and nitric acid.

BEZOARDICUM SOLARE. A diaphoretic medicine, prepared of gold filings, nitric acid, and butter of antimony.

BEZOARDICUM VENERIS. A pharmaceutical preparation, formerly employed in lepra, diseases of the brain, &c.; which was made from filings of copper, butter of antimony, and nitric acid.

BHANG, *Bangue*.

BI, *Bin*, (from *bis*, 'twice,') as a prefix to words, has the same signification as *Di*, twice, double.

BIBERINE, see Bebeeru.

BIBIRU, Bebeeru.

BIBITORIUS, (*bibere*, 'to drink,') because it draws the eye towards the nose so that the drinker can see into the cup, [?] Rectus internus oculi.

BIBLIOGRAPHY, MEDICAL (*βιβλιον*, 'a book,' and *γραφω*, 'I describe,') A description or history, or skill in the knowledge, of medical books. The most distinguished medical bibliographers have been: J. A. VAN DER LINDEN, Amsteld. 1662, octavo, (L.) M. LIPENIUS, Francf. ad Mœn. 1679, fol. (L.) G. A. MERCK-

LEIN, Norimb. 1686, (L.) J. J. MANGET, Genev. 1695 to 1731, (L.) TARIN (anatomical), Paris, 1753, (F.) A. VON HALLER, Zürich, 1774, &c., (L.) VIGILIUS VON CRUTZENFELD (surgical), Vindob. 1781, (L.) C. G. KUHN, Lips. 1794, (L.) C. L. SCHWEIKARD (anat., phys., and legal medicine), Stuttgart, 1796 to 1800, (L.) G. G. PLOUQUET, Tubing. 1808 to 1814, (L.) C. F. BURDACH, Gotha, 1810 to 1821, (G.) J. S. ERSCH, (since 1750), Leipz. 1822, (G.) TH. CH. FR. ENSLIN, (of Germany, -since 1750), Berlin, 1826, (G.) J. B. MONTFALCON, Paris, 1827, (F.) LUDWIG CHOULANT (ancient medicine), Leipzig, 1822, (G.) J. FORBES, M.D., F.R.S., London, 1835. LUDWIG CHOULANT (Bibliotheca medico-historica), Lips. 1842, (L.) JULIUS ROSENBAUM (additamenta ad L. CHOULANTI, Bib. med.-hist.) Halis Saxonom., 1842, and 1847, (L.) A. C. P. CALLISEN, Copenhagen, 1845, (G.) E. MORWITZ, Leipzig, 1849, (G.) **BIBRON'S ANTIDOTE**, see Antidote, Bibron's.

BICAUDALIS, (*bi*, and *cauda*, 'a tail,' 'double-tailed,') *Retrahens auris*.

BICAUDATUS, (same etymon.) A monster having two tails.

BICEPHALUM, *Dicephalum*, (*bi*, and *κεφαλη*, 'head,') Sauvages applies this epithet to a very large sacroma on the head, which seems to form a double head.

BICEPHALUS, *Dicephalus*.

BI'CEPS, gen. *Bicipitis*, (*bi*, and *caput*, 'head,') That which has two heads. This name has been particularly given to two muscles; one belonging to the arm, the other to the thigh.

BICEPS EXTER'NUS MUS'culus. The long portion of the *Triceps Brachialis*.

BICEPS FLEX'OR CRU'RIS, *B. Cru'ris*, *Biceps*, *Biceps Fem'oris*, (F.) *Biceps Crural*, *Is'chio-fem'oro-péronier*—(Ch.) A muscle on the posterior part of the thigh; one head arising from the tuberosity of the ischium, and the other from a great part of the linea aspera. It is inserted into the top of the fibula. It serves to bend the leg on the thigh.

BICEPS FLEX'OR CU'BITI, *Bi'ceps*, *B. Bra'chii* seu *ma'nus* seu *inter'nus* seu *inter'nus hu'meri*, *Cor'acoradialis*, (F.) *Scapulo-radial*, (Ch.)—*Biceps Brachial*. A muscle situate at the anterior and internal part of the arm; extending from the edge of the glenoid cavity and from the top of the coracoid process to the tuberosity of the radius. It bends the fore-arm upon the arm.

BICHE DE MER, *Bêche de mer*, *Trepang*, *Sea Slug*. A molluscous animal, belonging to the genus *Holothuria*, which is caught amongst the islands of the Fiji group, New Guinea, &c., and when prepared finds a ready sale in China, where it is used as an ingredient in rich soups.

BICHET, Terra Orleana.

BICHICH'LE, (probably corrupted from *βηξ*, gen. *βηξος*, 'cough,') Pectoral medicines, composed of liquorice juice, sugar, blanchéd almonds, &c.

BICHIOS, *Draunculus*.

BICHO, *Draunculus*—b. *di Culo*, *Proctocæce*.

BICHOS. A Portuguese name for the worms that penetrate the toes of people in the Indies; and which are destroyed by the oil of the cashew nut.

BICIPITAL, *Bicipitalis*, (*biceps* (*bi* and *caput*), 'two headed,') Relating to the biceps.

BICIPITAL GROOVE, (F.) *Contissæ* ou *Gouttière bicipitale*, *Contissæ humérale*, (Ch.,) is a longitudinal groove, situate between the tuberosities of the os humeri, which lodges the long head of the biceps.

BICIPITAL TU'BERCLE, *Bicipital tuberosity*, (F.) *Tubérosité bicipitale*; a prominence near the upper extremity of the radius, to which the tendon of the biceps is attached.

BICORNE RUDE, (*bi*, and *cornu*, 'a horn.') Ditrachyceros.

BICUSPID, *Bicuspidatus*, (*bi* and *cuspid*, gen. *cuspidis*, 'a spear.') That which has two cusps or points.

BICUSPID TEETH, *Dentes Bicuspidati*, (F.) *Dentes bicuspidates*. The small molares. See Molar.

BIDDY, *Pedicella*.

BIDENS ACMELLA, (*bi*, and *dens*, 'a tooth,' in allusion to the seed.) *Spilanthus acmella*—b. *Fervida*, *Spilanthus oleracea*.

BIDET, (F.) *Bidet*; (pronounced *beeday'*.) A small horse formerly allowed to each trooper for carrying his baggage. Hence, perhaps, applied to a chamber bathing apparatus, which has to be bestridden. It is a useful arrangement, in case of hemorrhoids, prolapsus ani, affections of the sexual organs, &c.

BIECHO, *Bische*.

BIER-RIGHT, TRIAL BY, see Trial by bier-right.

BIÈRE, (Beer,) *Cerevisia*—b. *Medicinale*, see *Cerevisia*.

BIESTINGS. ([Sax.] *býrting*.) Colostrum.

BIFEMORO-CALCANIEN, *Gastrocnemii*.

BIFURCATION, *Bifurcatio*, (*bi*, and *furca*, 'a fork.') Division of a trunk into two branches; as the bifurcation of the trachea, aorta, &c.

BIGARADE, see *Citrus aurantium*.

BIGASTER, (*bi*, and *gaster*, 'belly.') *Digastrius*.

BIGBELLIED, *Abdominous*.

BIG BLOOD, *Magnolia macrophylla*.

BIGEONE, see Kentucky, Mineral Waters of.

BIGEMINE EMINENTIAE, (*bi*, and *gemi*, 'twins.') *Quadrigenina tubercula*.

BIGEMINAL BODIES, *Quadrigenina tubercula*.

BIGGAR. A disease of Bengal, remarkable for the intensity and danger of the cerebral symptoms.

BIGLEAF, *Magnolia macrophylla*.

BIGLES, (*bis*, and *oculus*, 'an eye.' [?]) See *Strabismus*.

BIGNONIA CAPREOLA'TA, (after the Abbé Bignon, librarian to Louis XIV.,) *B. crucig'era*; indigenous. The root and plant are employed, like sarsaparilla, in infusion and decoction, in syphilis, chronic rheumatism, and in cachexia in general.

BIGNONIA CATALPA, *Catalpa*—b. *Crucigera*, *B. capreolata*.

BIGNONIA INDICA. The leaves are employed in India, as emollients, to ulcers.

BIGNONIA LEUCOX'YLOS, *Trum'pet tree*. A South American tree, the sap of which is said to be the surest antidote against the manchineel.

BIGNONIA OPHTHALMICA, *Eye root*, *Eye vine*. Indigenous in Guiana. The pulp or juice of the root, dropped into the eye, has been used in ophthalmia.

BIGNONIA RADICANS, *Tecoma radicans*—b. *Sempervirens*, *Gelsemium nitidum*.

BIGPAUNCHED, *Abdominous*.

BIJOU, see *Pinus sylvestris*.

BIKH, *Aconitum ferox*.

BILATE OF SODA, (from *bilis*), *Picromel*.

BILATERAL SYMMETRY, (*bi*, and *latus*, gen. *lateralis*, 'a side.') See *Symmetry*.

BILAZAY, MIN'ERAL WATERS OF. *Bilazay* is a town in France, two leagues from Thouar, department of Deux Sèvres, near which is a thermal sulphureous spring. Temperature about 77° Fahrenheit.

BILBERRY, ([?] blueberry) *Vaccinium myrtillus*—b. Red, *Vaccinium vitis idaea*.

BILE, *Bilis*, *Fel*, *Cholus*, *Cho'le*, *Cho'la*, *Choler*, (F.) *Bile*, *Fiel*. A yellow, greenish, viscid, bitter, nauseous fluid, secreted by the liver. It is dis-

tinguished into *hepatic* and *cystic*; according as it flows immediately into the duodenum from the liver or from the gall-bladder. It contains, according to Muratori, water; a peculiar fatty matter; colouring matter (*Cholepyrrhin* or *Bilipha'in*, (F.) *Matière Jaune de la Bile*; cholesterolin combined with soda; picromel or *bilin*; extract of flesh, mucus; soda, phosphate of soda; phosphate of lime, and chloride of sodium.

One use of the bile is to remove from the body superfluous hydro-carbon; and it is probably inservient to useful purposes in digestion.

BILE, *Furunculus*.

BILE OF THE BEAR, *Gall of the Bear*, *Fel Ursi*, was thought to be anti-epileptic; and that of the *Eel*, *Fel anguillæ*, to facilitate labour.

BILE, Black, *Atrabilis*—b. *de Boeuf*, see *Bile*.

BILE OF THE OX, *Gall of the Ox*, *Ox Gall*, *Fel Tauri* seu *Bo'vis* seu *Bovinum*, (F.) *Bile ou Fiel ou Amer de Boeuf*, was once reputed cosmetic and detergent, anti-algic and emmenagogue; as well as to possess the power of facilitating labour. It has also been given as a bitter stomachic and anthelmintic; and as a tonic and laxative, in cases of deficiency of the biliary secretion. *Fel Bovinum purificatum*, *Purified Ox Bile* (Ph. B. 1864) is made by mixing one pint of fresh ox bile, with two pints of rectified spirit in a bottle, setting aside for twelve hours until the sediment subsides; then decanting the clear solution and evaporating in a water bath until it has the consistence of a vegetable extract. Dose, gr. ij and more.

BILE REPANDUE, *Icterus*.

BILE OF THE SWINE, *Bilis porci'na*, inspissated, has been used in the same cases as the bile of the ox.

BILIARY, *Bilia'ris*, *Bilia'rius*, *Fel'leus*, (F.) *Biliaire*. That which relates to bile.

BILIARY APPARATUS, *B. organs*, *B. passages*. The collection of parts that concur in the secretion and excretion of bile—viz., the liver, pori biliares or tubuli biliferi; hepatic, cystic, and cholechoic ducts, and gall-bladder.

BILIARY CONCRETIONS are concretions found in some parts of the biliary apparatus.

BILIARY DUCTS, *Pori biliares*.

BILIATION, *Bilia'tio*, (from *bilis*.) The excretion of bile; as *urination* means the excretion of urine.

BILIEUX, *Bilious*.

BILIFULVINE; (*bilis*, and *fulvus*, 'reddish yellow.') The colouring matter of the bile of the ox.

BILIMBI, *Averrhoa bilimbi*.

BILIMBING TERES, *Averrhoa bilimbi*.

BILIN, (from *bilis*), *Picromel*.

BILIOUS, *Bilio'sus*, *Chol'icus*, *Chol'us*, *Cholo'des*, *Fellin'us*, *Epich'olos*, *Picroch'olos*, *Fel'leus*, (F.) *Bilieux*. That which relates to bile, contains bile, or is produced by bile. An epithet given to certain constitutions and diseases, which are believed to be the effect of superabundance of the biliary secretion; as *Bilious temperament*, *B. symptoms*, *B. fever*, but often used, without any definite ideas, as regards the bile, being attached to it.

BILIPHÆIN, (*bilis*, and *φαῖος*, 'of a dusky colour.') See *Bile*.

BILIS, *Bile*—b. *Fluxio*, *Cholera morbus*.

BILITICUS, (from *bilis*), *Cholagogue*.

BILIVERDIN, (*bilis*, and *viridis*, 'green.') On adding an acid to a solution of the yellow colouring matter of bile, a precipitate of green flocculi takes place, which possesses all the properties of chlorophyll or the green colouring matter of leaves. This is the *biliverdin* of Berzelius.

BILNA, MIN'ERAL WATERS OF, see *Püllna*.

BILOCULAR, (*bi*, and *loculus*, 'a small place.') See *Unilocular*.

BILUMBI BITING-BING, *Malus Indica*.
BILVA, *Covolan*.
BIMANUS, *Bim'anus*, (*bi*, and *manus*, 'a hand.') One that has two hands. A term applied only to man, because he is the sole mammiferous animal that possesses two perfect hands.
BIN, in composition, see *Bi*.
BINDER, *Bandage*.
BINDER, OBSTETRIC, see *Belt*, Russian.
BINDING, *Constipating*.
BINDWEED, *Polygonum aviculare*—*b. Fiddle-leaved*, *Convolvulus panduratus*—*b. Great*, *Convolvulus sepium*—*b. Hedge*, *Convolvulus sepium*—*b. Lavender-leaved*, *Convolvulus Cantabrica*—*b. Sea*, *Convolvulus soldanella*—*b. Virginian*, *Convolvulus panduratus*.
BINDWOOD, *Hedera helix*.
BINKOHUMBA, *Phyllanthus urinaria*.
BINOCULAR, *Bino'cularis*; (*bin*, and *oculus*, 'an eye.') Relating to or affecting both eyes—as 'binocular vision'—vision with both eyes; or from impressions made upon both retinae, which are amalgamated into single vision.
BINOCULUS, *Bin'ocle*, *Diophtalmica Fasl'cia*, *Oculus duplex*, (F.) *Œil double*. A bandage applied over both eyes. It was also formerly called *Diophtalmus*.
BIN'SICA. A Rabbinical term for disorder of the mind. According to VAN HELMONT, an atrophy of the organ of imagination.
BIO, (from *bios*, 'life,') in composition, life.
BIOCHYMIA, (*bio*, and *chymia*,) Chemistry, vital.
BIOD, (*bios*, 'life,') *Vis vitalis*.
BIODYNAMICS, *Biodynam'ica*, *Biodynam'ice*, *Biosoph'ia*, (*bio*, and *δυναμις*, 'power,' 'force,') The doctrine of the vital activity or forces.
BIOGAMIA, (*bio*, and *γάμος*, 'marriage,') Magnetism, animal.
BIOL'OGY, (*bio*, and *λογος*, 'a discourse,') The doctrine of life or of living bodies.—Physiology. 'Biological phenomena,' (F.) *Phénomènes biologiques*, are those presented by living bodies.
BIOLYCH'NION, *Biolych'nium*, (*bio*, and *λυχνιον*, 'a lamp,') Innate heat, vital heat, animal heat; *Lych'nium*, *Lychnid'ium*, *Ther'mum em'phytum*, *Flam'ma seu Flam'mula vital'is seu cor'dis*. Also, a secret preparation of which BEGUIN and BURGRAVE make mention.
BIOLYSIS, see *Biolytic*.
BIOLYTIC, *Biolyt'icus*; (*bio*, and *λυσις*, 'solution,') Relating to the destruction of life. A 'biolytic agent' is one that tends to *biolysis*, or destruction of life. A depressing agent.
BIOMAGNETISMUS, Magnetism, animal.
BIONOMY, (*bio*, and *νομος*, 'law,') Physiology.
BIOPHENOMENOLOGIA, (*bio*, *φαινόμενον*, 'an appearance,' and *λογος*, 'a discourse,') Physiology.
BI'OS, (*bios*,) Life. Also, what is necessary for the preservation of life.
BIOSOPHIA, (*bio*, and *σοφία*, 'wisdom,') Biodynamics.
BIOSTATISTICS, (*bio*, and *statistics*,) Statistics, medical.
BIOTAX'Y, *Taxon'omy*, (*bio*, and *τάξις*, 'order,') the arrangement or co-ordination of living organisms, according to their external characters.
BIOTE, (*βιοτη*,) Life.
BIOTHALMIUS, (*βιοθαλμιος*, from *bio*, and *θαλλω*, 'I bloom,') One who is long lived, or enjoys rude health.
BIO THAN'ATI, (*βιοθανatoi*, from *bio*, and *θανος*, 'death,') They who die of a violent death very suddenly, or as if there was no space between life and death.
BIOTIC, (*βιωτικός*,) Vital.
BIOTICS, Physiology.
BIOTOMIA, (*bio*, and *τεμνειν*, 'to cut,') Vivisection.
BIOTOS, (*βιωτος*,) Life.

BIPARI'ETAL, *Biparieta'lis*, (F.) *Bipari'etali*, (*bi* and *parietale* (os), 'the parietal bone,') An epithet for the diameter of the cranium from one parietal fossa to the other.
BIPARIETAL SUTURE, Sagittal suture.
BIPIN'NA, (*bi*, and *pinna*, 'a wing-feather,') A term used by the ancients for a diminutive penis, not exceeding in size two quills.
BIPOLAR, (*bi*, and *polus*, 'the pole,') See *Neurine*.
BIR, Thorax.
BIRA, *Cerevisia*.
BIRCH, ([Sax.] *birce*,) *Betula alba*—*b. Black*, *Betula lenta*—*b. Cherry*, *Betula lenta*—*b. Sweet*, *Betula lenta*.
BIRDS' BREAD, *Sedum acre*.
BIRDS' NEST, *Hypopitys lanuginosa*—*b. Nest*, edible, see *Hirundo esculenta*.
BIRK, (Sc. and Prov.,) *Betula alba*.
BIRTH, Parturition—*b. Cross*, Presentation, preternatural—*b. Live*, see *Born alive*—*b. Plural*, see *Multiparous*.
BIRTHWORT, (*birth* and *wort*,) *Aristolochia*—*b. Snakeroot*, *Aristolochia serpentaria*.
BISCHE, *Biecho*. A malignant kind of dysentery, which has often prevailed in the island of Trinidad.
BIS'CUIT, *Biscoc'tus*; (*bis*, 'twice,' and *coctus*, 'baked,' (F.) *bis* and *cuit*, 'twice baked,') A kind of dry, hard bread, or cake, which is variously made; and, when without eggs or butter, is easy of digestion. It was formerly called *Dipyr'rites*, and *Di'pyros*. MEDIC'INAL OR MED'ICATED
BIS'CUITS, (F.) *Biscuits Médicinaux*, *Massepains*, *Macarons*, are occasionally made by adding to biscuit paste a medicinal solution or powder, and baking the mass.
BIS'CUIT, MEAT. An alimentary preparation, proposed by Mr. G. Borden, Jr., of Texas, which consists in combining the matters, extracted from meat by boiling, with flour, so as to form biscuits; which keep well, and are of course nutritive.
BISCUITS MÉDICINAUX, see *Biscuit*.
BISERMAS, *Salvia sclarea*.
BISFERIENS, (*bis*, and *ferio*, 'to strike,') Di-crotes.
BISH, *Aconitum ferox*.
BISHOP'S WEED, *Ammi*.
BISLINGUA, *Ruscus hypoglossum*.
BISMALVA, *Athæa*.
BIS'MUTH, ([G.] *Wismuth*,) *Antimonium album*, *Chalcitis*, *Lu'na imperfecta*, *Stannum glaciale* seu *cine'reum*, *Bismuthum*, *Wismuthum*, *Reg'ulus of Bis'muth*, *Marcas'ita*, *Tin glass*, (F.) *Étain gris*, *É. de Glace*. A metal, in spicular plates, of a yellowish-white colour; s. gr. 9.822; fusible at 400° Fahrenheit, and volatilizable at a high temperature. It is used only in the preparation of the subnitrate.
BISMUTH, NITRATE OF, Bismuth, subnitrate of—*b. Oxide of*, Bismuth, Subnitrate of—*b. Oxide of*, white, Bismuth, subnitrate of—*b. Regulus of*, Bismuth—*b. Subcarbonate of*, see *Bismuth*, subnitrate of.
BISMUTH, SUBNITRATE OF, *Bismuthi subn'itras* (Ph. U. S.) seu *Nit'ras*, *Marcas'ita al'ba*, *Plum'bum cine'reum*, *Magister'ium Marcas'ite* seu *Bismuthi*, *Bismuthum Nitricum* seu *Subnitricum*, *Nit'ras Subbismuth'icum*, *Calx Vismuthi*, *Bismuthum oxydula'tum album*, *Bismuthum album*, (Ph. B. 1864.) *Subazotas bismuthicus*, *Album Hispan'icum*, *Ox'ide of Bis'muth*, *Nitrate of Bis'muth*, *White Bismuth*, *Trienitrate of Bis'muth*, *White Ox'ide of Bis'muth*, *Mag'istery of Bis'muth*, *Pearl White*, *Span'ish White*. (F.) *Sousnitrate ou sousazotate de bismuth*, *Oxide blanc de B.*, *Blanc de fard*, *de perle* ou *d'Espagne*. (*Bismuth*, in frustulis, ʒj. Acid nitric. f ʒij. Aq. destill. q. s. Mix

a fluidounce of distilled water with the nitric acid, and dissolve the bismuth in the mixture. When the solution is complete, pour the clear liquor into three pints of distilled water, and set the mixture by, that the powder may subside. Lastly, having poured off the supernatant fluid, wash the subnitrate of bismuth with distilled water, wrap it in bibulous paper, and dry with a gentle heat. A modification of this process is contained in the Ph. U. S. for 1863; and, also, a formula for *Bismuthi subcarbonas*, *Subcarbonate of Bismuth*. Both these preparations are considered to be tonic and antispasmodic, and have been chiefly used in gastrodynia.

BISMUTH, TRISNITRATE OF, Bismuth, subnitrate of.

BISMUTH, VALE'RIANATE OF, *Bismuthi valerianae*, *Bismuthum valerianicum*. Prepared by mixing a neutral solution of oxide of bismuth in nitric acid with valerianate of soda; washing, and drying the precipitate. Used in gastrodynia, chronic gastralgia, neuralgia, and chronic palpitation, as a nerve. Dose, $\frac{1}{2}$ a grain to 2 grains, three or four times a day, in pill.

BISMUTH, WHITE, Bismuth, subnitrate of.

BISMUTHI NITRAS, Bismuth, subnitrate of—b. Subcarbonas, see Bismuth, subnitrate of—b. Subnitras, see Bismuth, subnitrate of—b. Valerianas, Bismuth, valerianate of.

BISMUTHUM, Bismuth—b. Album, Bismuthi subnitras—b. Nitricum, Bismuth, subnitrate of—b. Oxydulatum album, Bismuth, subnitrate of—b. Subnitricum, Bismuth, subnitrate of—b. Valerianicum, Bismuth, valerianate of.

BISPIRUS, (*bi*, and *epi*, 'to breathe.') Dip-nous.

BISSUM, *Hydrangea arborescens*.

BISTORT, OFFICIAL, (*bi*, and *torquere*, *tortum*, 'to twist,') Polygonum bistorta—b. Virginian, Polygonum virginianum.

BISTORTA, Polygonum bistorta.

BISTORTE, Polygonum bistorta.

BISTORTIER, (F.) A name given by the Pharmacien to a long wooden pestle used for reducing soft substances to powder, and in the preparation of electuaries.

BISTOURI, (F.) *Pistorien'sis gladius*, *Scalpel'us*, *Scal'pium*, *Bistoury*. A small cutting-knife, used in surgery, (so called from the town of Pistorium, modern Pistoja, in Tuscany, which was formerly celebrated for the manufacture of those instruments. [?]) A bistoury has the form of a small knife, and is composed of a blade and a handle. The blade, which is most commonly movable in the handle, may be fixed by a button, spring, &c. When fixed in the handle, the bistoury is called by the French, *B. à lame fixe* ou *dormante*.

The chief bistouries are:—1. The **STRAIGHT B.**, (F.) *B. droit*, in which the blade and cutting edge are straight, the point being fine, round, or square. 2. The **CONVEX B.**, (F.) *B. convexe*; the blade of which is convex at the cutting edge, concave at the back. 3. The **CONCAVE B.**, (F.) *B. concave*; the blade of which is concave at its edge, and convex at the back. 4. **BLUNT-POINTED B.**, (F.) *B. boutonné*; the blade of which has a button at its extremity. 5. The **BLUNT OR PROBE-POINTED BISTOURY OF POTT**; concave at its cutting edge, and its point blunt; so that it can be carried on the palmar surface of the index finger, to divide the stricture, in strangulated hernia. Sir Astley Cooper has recommended a useful modification of this, to avoid wounding the intestine, should it come in contact with the edge of the knife. His Bistoury has an edge of not more than eight lines in length, situate about five lines from the point. 6. **BISTOURI À LA LIME** (F.) is a straight bistoury; the blade fixed in the handle, the extremity with a button, and the edge made

with a file. It is chiefly used for dilating parts. 7. **BISTOURI ROYAL** (F.) A Bistoury used in operating upon Louis XIV., for fistula in ano. 8. **BISTOURI GASTRIQUE**, (F.) A complicated instrument, invented by Morand, for dilating wounds of the abdomen. 9. **BISTOURI CACHÉ**, *B. herniaire*, ou *Attrape-lourdard de Biennaise*, *For'ceps deceptoria*. A curved bistoury, the blade of which is placed in a canula, whence it issues on pressing a spring.

The word *Bistoury* is used by the French, at times, where we would employ knife.

BISTOURY, Bistoury.

BITCH-DAUGHTER, Incubus.

BITNOBEN, *Salt of Bitu'men, Padnoon, Soucherloon, Khala mimuc*. A white, saline substance, which is a Hindoo preparation of great antiquity, and has been supposed to be the *Sal asphaltites* and *Sal Sodome'nis* of the ancients. It is used by the Hindoo in the prevention or cure of almost all diseases.

BITHNIMAL'CA, *Gas'teranax*. Two unmeaning words, used by Dolæus, to designate an active principle supposed to have its seat in the stomach, and to preside over chymification, &c.

BITIOS DE KIS, *Proctocæce*.

BITTER, (Sax. *biter*), *Amarus*—b. Ash, see *Bittera* febrifuga—b. Bark, *Pinckneya pubens*—b. Bloom, *Chironia angularis*—b. Holy, *Hiera pica*—b. Redberry, *Cornus Florida*—b. Root, *Apocynum androsæmifolium*, *Gentiana Catesbæi*, *Menyanthes verna*—b. Sweet Nightshade, *Solanum Dulcamara*—b. Sweet Vine, *Solanum Dulcamara*—b. Welter's, Acid, carbazotic.

BITTERA FEBRIFUGA, *Quassia*.

BITTERIN, see *Quassia*.

BIT'TERNESSE, *Amaritudo*, *Amarit'ies*, *Amaror*, *P'eria*, (F.) *Amertume*. A particular taste, which belongs to many substances. In some diseases there is a sense of bitterness felt in the mouth.

BITTERS, COLUMBO, *Tinctura Calumbæ*—b. Spirit, *Tinctura gentianæ composita*—b. Wine, *Vinum gentianæ compositum*.

BITTERSWEET, *Solanum dulcamara*.

BITTERWEED, *Ambrosia trifida*.

BITTERWOOD TREE, *Quassia*.

BITTOS. A disease, in which the chief symptom is an acute pain in the anus: see *Bitios*.

BITUME DE JUDEE, *Asphaltum*.

BITUMEN, GLUTINOUS, (*urra*, 'pitch,') *Pis asphaltum*—b. of Judea, *Asphaltum*—b. Judaicum, *Asphaltum*—b. Malta, *Pissasphaltum*—b. Petroleum, *Petroleum*—b. Salt of, *Bitnoben*—b. Solidum, *Asphaltum*.

BIVENTER, (from *bi*, and *venter*, 'belly,') *Digastricus*—b. Cervicis, *Complexus musculus*—b. Maxillæ, *Digastricus*.

BIVENTRAL LOBE OF THE CEREBELLUM, see *Lobe, biventral*.

BIXA AMERICANA, see *Terra Orleana*—b. Orellana, see *Terra Orleana*—b. Orleana, see *Terra Orleana*.

BLABE, (*βλαβη*), *Wound*.

BLACCLE, *Rubeola*.

BLACIA, (*βλακία*), *Debility*.

BLACKBERRY, AMERICAN, see *Rubus fruticosus*—b. High or Standing, see *Rubus fruticosus*—b. Low, *Rubus trivialis*.

BLACK BOTTLE, see *Infusum Sennæ*—b. Damp, see *Hydrogen*, carburetted—b. Dose, see *Infusum Sennæ*—b. Draught, see *Infusum Sennæ*—b. Drop, *Guttæ nigre*.

BLACK LION. A term given to a sloughing syphilitic ulcer, under which the British soldiers suffered greatly in Portugal.

BLACK ROOT, *Aletris farinosa*, *Pterocaulon pycnostachyum*, *Leptandria purpurea*—b. Sugar, *Extractum glycyrrhizæ*—b. Vomit, *Fever, yellow*.

BLACKWATER, Pyrosis.

BLACTIE, Rubella.

BLADDA, Buttermilk.

BLADDER, (Sax. blæþn,) Urinary Bladder, Vesica.

BLADDER, FASCICULATED. A condition of the bladder, in which the interior of the organ is marked by strong reticulated ridges or columns.

BLADDER, GALL, see Gall Bladder—b. Irritable, Cysterethismus.

BLADDER, SAC'ULATED. A condition of the bladder, in which the mucous coat protrudes through the muscular, so as to produce a hernia, which may go on increasing, so as to form a vesical sacculus or appendix vesicæ.

BLADDER, SWIM, Air bladder—b. Urinary, see Urinary Bladder.

BLADEBONE, Scapula.

BLADOCH, Buttermilk.

BLADON SPRINGS. Mineral springs in Alabama, about 85 miles from Mobile, which are said to be similar in composition to those of Seltzer, Spa, and Aix-la-Chapelle, of Europe.

BLADUM (low L.), Blé.

BLAEBERRY, Vaccinium myrtillus.

BLÆ'SITAS, Blæsa lin'gua, (βλαστος, 'lame, stammering,') Some authors have used this word as synonymous with stammering. See Balbuties. Sauvages understands by it a defect in pronunciation, which consists in substituting soft consonants for those that are hard; as the z for s, the d for t, the s for g and j, &c. Also, Lispings, Traulismus, Traulotes, (F.) Blésité, Blé (parler).

BLÆSOPODES, (βλαστος, 'lame,' and πους, gen. ποδος, 'a foot,') see Kyllosis.

BLÆSOPUS, see Kyllosis.

BLÆSUS, (βλαστος, 'lame,') A distortion; especially the outward distortion of the legs. Also, a stammerer.

BLAFARD (F.), ([G.] bleich, 'pale,' and Farbe, 'color,') Pallidus, Pallidulus. This epithet is sometimes given to the skin, when pale and dull; but, more frequently, to the flesh of a wound, when it has lost its colour, and become white. The word is, also, sometimes used synonymously with Albino.

BLAIN, (Sax. blegene,) Bleine. A pustule; a blotch; a blister. In Scotland, a mark left by a wound; the discolouring of the skin after a sore.

BLAK OF THE EIE, Pupil.

BLANC D'ARGENT, ('silver white,') Plumbi subcarbonas—b. de Baleine, Cetaceum—b. de Céruuse, Plumbi subcarbonas—b. de Crems, Plumbi subcarbonas—b. d'Espagne, Bismuth, subnitrate of—b. de Fard, Bismuth, subnitrate of—b. de Kremnitz, Plumbi subcarbonas—b. de Krems, Plumbi subcarbonas—b. de l'Œil, see Sclerotic—b. d'Œuf, Albumen ovi—b. de Perle, Bismuth, subnitrate of—b. de Plomb, Plumbi subcarbonas—b. de Zine, Zinci oxydum.

BLANCMANGER (F.), (blanc, 'white,' and manger, 'eating,') Cibus albus, Leucophagium, Leucophagum, Argyrotophæma. An animal jelly, so called on account of its colour, combined with an emulsion of sweet almonds, to which sugar has been added, and some aromatic. It is at times prescribed as a nutriment in convalescence and chronic diseases.

BLANC-RAISIN, Blanc Rhazis.

BLANC RHAZIS, Blanc-raisin. An ointment composed of cerussa, white wax, and olive oil.

BLANCA, Plumbi subcarbonas.

BLANCH, ([F.] blanchir, 'to whiten, to bleach,') To whiten by depriving of the outer rind; as 'to bleach almonds;' i. e. to peel them.

BLANCHET (F.), 'a blanket.' A term given by the French Pharmaciens to the woollen strainer

through which they filter syrup and other thick fluids. See, also, Aphthæ.

BLANCHING, (from (F.) blanchir, 'to whiten,') Etiolation—b. of the Hair, Canities.

BLANCNON ORIBASII, Polypodium filix mas.

BLAS. An unmeaning term, invented by Van Helmont to designate a kind of movement in the body; at times, local,—at others, under extraneous influence. Thus, he speaks of the Blas meteoron of the heavenly bodies, and the Blas humanum, that which operates in man.

BLAS ALTERNATIVUM, Plastic force.

BLASÉ (F.), 'cloyed.' An epithet given to one whom the abuse of enjoyment has prevented from any longer deriving satisfaction or pleasure from it.

BLAST, (Sax. blapt, G. blasen, 'to blow,') Contagio. The infection of any thing pestilential. The stroke of some sudden plague.

BLASTE'MA, gen. Blaste'matis, Blaste'sis, (F.) Blastème, (βλαστημα, from βλαστανω, 'I bud,') A germ. The sense of this word, which is often used by Hippocrates, is obscure. Castelli thinks it means the eruption of some morbid principle at the surface of the body. Also, the matrix or general formative element of tissues.

BLASTEMA PILI, see Hair.

BLAST'EMAL, or BLASTE'MAL, Blastema'tis. Relating or appertaining to a blastema,—as 'blastemal formations,' those that are formed from a blastema.

BLASTESIS, (βλαστεισις,) Blastema

BLASTOCARDIA, (βλαστος, 'germ,' and καρδιά, 'heart,') see Molecule.

BLASTOGELIS, (βλαστος, 'germ,' and κελίς, 'a spot,') see Molecule.

BLASTOCYSTINX, (βλαστος, 'germ,' and κυστιγξ, dim. of κυστις, 'a bladder,') see Molecule.

BLASTODERMA, (βλαστος, 'germ,' and δερμα, 'skin,') see Molecule.

BLASTODERMIC, Blastoderm'icus, (F.) Blastodermique. Relating or appertaining to the Blastoderma.

BLASTOSTROMA, (βλαστος, 'germ,' and στρώμα, 'a bed,') Tache embryonnaire.

BLAT'TA BYZANT'IA, Un'guis odora'tus, (F.) Blatte de Byzance. This name seems, formerly, to have been given to a marine production from some of the Conchylia. It had an agreeable smell, a reddish tint, and the shape of a nail. It was prescribed in epilepsy, hysteria, and hepatic obstructions. Rondelet affirms, that it was the production of the shell-fish murex or purpura; and that the name Blatta is derived from the Greek βλαστος, 'purple.'

BLA'VELLE, Centaurea cyanus.

BLA'VEÔLE, Centaurea cyanus.

BLA'VEROLLE, Centaurea cyanus.

BLAWING, Swelling.

BLAWORT, Centaurea cyanus.

BLAZING STAR, Chamælorium luteum, Liatriis.

BLÉ, Blad'um. This word answers, in France, to the word Corn in England; i. e. any kind of grain employed for making bread. Wheat being most commonly used for this purpose, Blé is sometimes restricted to it. Blé mêtéil is a mixture of wheat and rye.

BLÉ CORNU, ('horned corn,') Ergot—b. d'Espagne, Zea mays—b. d'Italie, Zea mays—b. Mêtéil, see Blé—b. Noir, Polygonum fagopyrum—b. de Turquie, Zea mays.

BLÉ (PARLER), Blæsitas.

BLEABERRY, ([?] 'blueberry,') Vaccinium myrtillus.

BLEACHING, ([G.] bleichen, 'to whiten,') Dealbation—b. Liquid, Eau de javelle.

BLEAREDNESS, (blear, like Blain, [?]) Lippitudo.

BLEAR-EYE, Lippitudo.

BLEAR-EYED, Lippus.

BLEAR-EYEDNESS, Lippitudo.

BLEB, Bulla.

BLECHNON, (βληκνον,) Polypodium filix mas.

BLECHNUM LIGNIFOLIUM, Asplenium Scolopendrium—b. Squamosum, Asplenium ceterach.

BLECHROPYRA, (blechros and πυρ, 'fever.')

BLECHROPYRUS, Typhus mitior.

BLE'CHROS, (βληκρος, 'weak, feeble, slow.')

An epithet applied to different affections, and particularly to fevers. Hence, *Blechropyra*, and *Blechrophyg'mia*.

BLECHROSPHYGMIA, (βληκρος, and σφυγμος, 'pulse,') a slow pulse. see *Blechros*.

BLE'D, (from bladam.) Corn—b. *Avorté*, Ergot—b. *Farouche*, Ergot.

BLEDOCH, Buttermilk.

BLEEDER, Phlebotomist.

BLEEDING, Bloodletting, Hæmorrhagia.

BLEEDING FROM THE NOSE, Epistaxis.

BLEEDING BOIST, Cupping-glass—b. Heart, Cypripedium luteum.

BLEIB (Sc.), Bulla.

BLEINE, Blain.

BLEME, (F.) (from (G.) bleich, 'pale.' [?]) This word has nearly the same signification as *Blafard*. Generally, however, it includes, also, emaciation of the countenance.

BLENCYSTORRHÉE, (blenna, and cystirrhœa.) Cystirrhœa.

BLENNA, (βλενα.) Mucus—b. *Narium*, Nasal mucus.

BLENNADENITIS, (blenna, αδην, 'a gland,' and *itis*.) Inflammation of mucous follicles.

BLENNELYTRIA, (blenna, and ελυτρον, 'a vagina or sheath,') A discharge of mucus from the vagina. Leucorrhœa.

BLENNEM'ESIS, (Blennoëm'esis, Vom'itus pituito'us, (blenna, and εμεσις, 'vomiting,') Vomiting of mucus.

BLENNENTERIA, (blenna, and εντερον, 'intestine,') Dysentery.

BLENNISTH'MIA, (blenna, and ισθμος, 'the gullet,') Increased flow of mucus from the pharynx and larynx.

BLENNOCHEZIA, (blenna, and χεζειν, 'to go to stool,') Diarrhœa, mucous.

BLENNOCYSTIDES, (blenna, and κυστις, 'a cyst or bladder,') Bursæ mucosæ.

BLENNOCYSTITE, (blenna, κυστις, 'bladder,' and *itis*.) Cystirrhœa.

BLENNODES, (βλενωδής, from blenna and odes.) Mucous.

BLENNOËMESIS, Blennemesis.

BLENNOG'ENOUS, *Blennog'enus*, *Mucif'ic*, *Mucif'icus*, (blenna, and γεννω, 'I form,') Forming or generating mucus. Breschet and Roussel de Vauzème describe an apparatus of this kind for the secretion of the mucous matter that constitutes the cuticle, composed of a glandular parenchyma or organ of secretion situate in the substance of the true skin, and of excretory ducts, which issue from the organ, and deposite the mucous matter between the papillæ.

BLENNOIDES, (blenna, and ειδος, 'resemblance,') Muciform.

BLENNOIDEUS, Muciform.

BLENNOMETRITIS, (blenna, μετρα, 'uterus,' and *itis*.) See *Leucorrhœa*.

BLENNOMETRORRHŒA, see *Leucorrhœa*.

BLENNOPHLOGOSIS, (blenna, and phlogosis,) Mucitis.

BLENNOPHTHALMIA, see *Ophthalmia*, (purulent.)

BLENNOP'TYSIS, (blenna, and πτω, 'I spit,') Expectoration of mucus. Catarrh.

BLENNOP'YRA, *Blennopyr'ia*, (blenna, and

πυρ, 'fire,') Alibert has classed, under this head, various fevers with mucous complications; as *Mesenteric fever*, *Adeno-meningeal fever*, &c.

BLENNORRHAGIA, (blenna, and ρηνωμι, 'to break forth,') Gonorrhœa—b. Genitium, Leucorrhœa—b. Notha, Gonorrhœa spuria—b. Spuria, Gonorrhœa spuria.

BLENNORRHAG'IC, *Blennorrhag'icus*, (F.) *Blennorrhagique*. Relating or appertaining to blennorrhagia or gonorrhœa.

BLENNORRHAGIC EPIDIDYMITIS, Hernia humoralis.

BLENNORRHAGIE FAUSSE, Gonorrhœa spuria—b. *du Gland*, Gonorrhœa spuria.

BLENNORRHÉE, Blenorrhœa.

BLENNORRHINIA, (blenna, and ριν, 'nose,') Coryza.

BLENNORRHŒ'A, *Blennor'rhoë*, *Blennorrhag'ia*, *Phlegmorrhœ'a*, *Phlegmorrhag'ia*, (F.) *Blennorrhée*, (blenna, and ρεω, 'I flow,') Inordinate secretion and discharge of mucus. Also, Gonorrhœa.

BLENNORRHŒA CHRONICA, (gleet,) see *Gonorrhœa*—b. Genitium, Leucorrhœa—b. *Luodes*, Gonorrhœa impura—b. *Nasalis*, Coryza—b. *Oculi*, see *Ophthalmia*—b. *Oculi gonorrhœica*, see *Ophthalmia*—b. *Oculi neonatorum*, see *Ophthalmia*—b. *Oculi purulenta*, see *Ophthalmia*—b. *Urethralis*, Gonorrhœa, Cystirrhœa—b. *Ventriculi*, Gastrorrhœa—b. *Vesicæ*, Cystirrhœa.

BLENNO'SES, (from blenna.) Affections of the mucous membranes.

BLENNOS'TASIS, (blenna, and στας, 'rest,') Suppression of a mucous discharge.

BLENNOTHORAX, gen. *Blennothorac'is*, (blenna, and θωραξ, 'chest,') Catarrh, Peripneumonia notha—b. *Chronicus*, Asthma humidum.

BLENNOTORRHŒA, (blenna, ος, gen. ωτος, 'ear,' and ρεω, 'to flow,') Otirrhœa.

BLENNURETHRIA, (blenna, and ουρηδρα, 'the urethra,') Gonorrhœa.

BLENNURIA, (blenna, and ουρον, 'urine,') Cystirrhœa.

BLENNYMEN, (blenna, and ιμην, 'a membrane,') Membrane (mucous).

BLENNYMENTIS, Mucitis.

BLEPHARADENTITIS, (blepharo, αδην, 'a gland,' and *itis*.) Ophthalmia Tarsi.

BLEPHARANTHRACOSIS, *Blephar'itis gangræno'sa*, *Carbuncula'tio Oculi*, (blepharo, and anthracosis.) Gangrenous inflammation of the eyelids.

BLEPHARELOSIS, (blepharo, and ειλω, 'I roll,') Entropion.

BLEPHARIDES, (βλεφαρις, pl. βλεφαριδες,) Cilia.

BLEPHARIDOPLASTICE, (blepharo, and πλασ, 'I form,') Blepharoplastice.

BLÉPHARIQUE, (from blepharon,) Palpebral.

BLEPHARISMUS, Nictation.

BLEPHARITIS, (blepharo, and *itis*.) Ophthalmia tarsi—b. *Gangrænosa*, *Blepharanthracosis*.

BLEPHARO, (from βλεφαρον, 'an eyelid,') in composition, eyelid.

BLEPHAROBLENNORRHŒA, (blepharo, and blennorrhœa,) Ophthalmia, purulent—b. *Neonatorum*, see *Ophthalmia* (purulenta infantum.)

BLEPHAROCLEI'SIS, (blepharo, and κλεισις, 'a closure,') Occlusion or adhesion of the eyelids.

BLEPH'AROCONJUNCTIVI'TIS, *Blepharo-syndesmi'tis*, (blepharo, conjunctiva, and *itis*.) Ophthalmia affecting the conjunctiva and eyelids.

BLEPHARODYSCHŒ'A, (blepharo, δυσ, 'with difficulty,' and χρος, 'colour,') Discoloration of the eyelid. Nævus of the eyelid.

BLEPHARŒDEMA AQUOSUM, gen. *Blepharœdema'tis*, (blepharo, and οίδημα, 'a swelling,') Hydroblepharon.

BLEPHARON, (βλεφάρων,) Palpebra—b. Atontion, Blepharoptosis.

BLEPHARONCO'SIS, *Blepharon'cus*, *Blepharophy'ma*, *Palpebra'rūm Tu'mor*, (blepharo, and γκος, 'tumour.') A tumour of the eyelid.

BLEPHARONCUS, Blepharoncosis, BLEPHAROPHIMO'SIS, (blepharo, and φῖμος, 'I constrict.') Congenital diminution of the space between the eyelids. See Blepharostenosis.

BLEPHAROPHTHALMIA, Ophthalmia tarsi—b. Neonatorum, see Ophthalmia—b. Purulenta, Blepharopyorrhœa—b. Ulcerosa, see Ophthalmia tarsi.

BLEPHAROPHTHALMITIS GLANDULOSA, Ophthalmia, purulent, of infants.

BLEPHAROPHYMA, (blepharo, and φυμα, 'a swelling,') Blepharoncosis.

BLEPHAROPLAS'TICE, *Blepharidoplas'ticē*, *Insi'tio Cilio'rūm*, *Bleph'aroplasty*, (blepharo, and πλαστικός, 'forming,' 'formative.') The formation of a new eyelid.

BLEPHAROPLEGIA, (blepharo, and πλῆγη, 'a stroke,') Blepharoptosis.

BLEPHAROPTO'SIS, *Blepharople'gia*, *Ca'usa pal'pebræ superio'ris Delap'sus* seu *Prolap'sus* seu *Propto'sis* seu *Pto'sis pal'pebræ*, *Atontion bleph'aron*, (blepharo, and πτωσις, 'fall.') A falling down of the upper eyelid over the eye, caused by a paralysis of the *Levator palpebræ superioris* muscle. This paralysis is an unfavorable symptom, as it is generally connected with a state of the brain favouring apoplexy or palsy.

BLEPHAROPTOSIS ECTROPIUM, Ectropium—b. Entropion, Entropion.

BLEPHAROPYORRHŒA *Blepharophthal'mia purulen'ta*, *Pyorrhœ'a pal'pebræ*, (blepharo, πον, 'pus,' and ρεω, 'I flow,') Secretion of pus from the eyelids.

BLEPHAROPYORRHŒA NEONATORUM, see Ophthalmia (purulenta infantum.)

BLEPHARORRHŒA, (blepharo, and ρεω, 'I flow,') A discharge of mucus from the eyelids.

BLEPHAROSPAS'MUS, (blepharo, and σπασμος, 'spasm,') A spasmodic action or tic of the orbicularis palpebrarum muscle.

BLEPHAROSTENO'SIS, (blepharo, and στενωσις, 'contraction,') Accidental diminution of the space between the eyelids. See Blepharophimosis.

BLEPHAROSYNDESMITIS, (blepharo, συνδεδεστος, 'a bond,' and itis, 'Blepharoconjunctivitis. BLEPHAROTIS, Ophthalmia tarsi—b. Glandularis contagiosa, see Ophthalmia.

BLEPHAROTITIS, Ophthalmia tarsi.

BLEPHAROTIS, Ectropium.

BLEPHAROXYS'TUM, *Blepharoxys'trum*, (blepharo, and ξυω, 'I scrape,') An instrument used, by the ancients, for removing callosities, which made their appearance in the affection, called by the Greeks τραχώμα.

BLEPHIL'IA HIRSUTA, *Oh'i'o Horse'mint*, *Hair'y Horse'mint*, (βλεφαρίς, 'an eyelash,' in reference to its hairy fringed bracts.) An indigenous plant of the Mint family, Labiate, which has the aromatic properties of the Mints.

BLÉSITÉ, Blasitas.

BLESSED HERB, Geum urbanum.

BLESSURE, (from πλησσειν, 'to strike,') Abortion, Wound.

BLESTRIS'MUS, (βληστρισμος,) Restlessness of the sick.

BLETA. A word, used by Paracelsus for white or milky urine, arising from diseased kidneys. *Biota alba* has the same meaning.

BLEU DE BERLIN, ('Berlin blue,') Prussian blue—b. de Prusse, Prussian blue.

BLEUET DES MOISSONS, ('blue bottle of the harvests,') *Cyanus segetum*.

BLÉVILLE, MIN'ERAL WA'TERS OF. Blé-

ville is a village about two miles from Havre. The waters are acidulous chalybeate.

BLIGHT IN THE EYE, Ophthalmia, catarrhal. Palsy of the orbicularis palpebrarum and muscles of the eyebrow is also vulgarly called blight.

BLIGHTS, see Lichen urticatus.

BLINDNESS, (from Sax. blind,) Cæcitas—b. Colour, Achromatopsia—b. Day, Nyctalopia—b. Nervous, Amaurosis—b. Night, Hemeralopia—b. Snow, see Snow blindness.

BLIS'TER, (D. *bluyster*; Sax. *blæstan*, 'to puff,') *Vesicato'rium*, *Emplas'trum Vesicato'rium* seu *Lyt'tæ*, *Epispas'ticum*, *Ves'icant*, *Blis'ter plas'ter*, (F.) *Vésicatoire*, *Vésicant*. Any substance which, when applied to the skin, irritates it, and occasions a serous secretion, raising the epidermis, and inducing a vesicle. Various articles produce this effect, as *cantharides*, *mustard*, *garou*, *euphorbium*, *garlic*, *ammonia*, &c. Blisters are used as counter-irritants. By exciting a disease artificially on the surface, we can often remove another which may be at the time existing internally. A blister applied for a few hours to produce this effect is termed by the French *Vésicatoire volant*, 'a flying blister.' A *perpetual blister* is one that is kept open for a longer or a shorter time by means of appropriate dressings.

BLISTER, (Prov. *Ercle*, also means the *vesicle* or *vesication*, produced by vesicatories or other causes.

BLISTER BEETLE, *Cantharis*.

BLISTER, FEVER, see Fever blister—b. Fly, *Cantharis*.

BLISTER PLASTER, Blister.

BLISTER, MAG'ISTRAL, (F.) *Vésicatoire magistral*. A prompt means of producing vesication recommended by M. Valleix. It is prepared as follows—Take powdered *cantharides* and *wheat-flour*, of each equal parts; *vinegar*, a sufficient quantity to form a soft paste.

BLISTERS, WHITE, *Rupia escharotica*.

BLISTERWEED, *Ranunculus acris*.

BLISTERING FLY, *Cantharis*—b. Paper, see *Sparadrapum vesicatorium*—b. Tissue, *Sparadrapum vesicatorium*.

BLITUM AMERICANUM, *Phytolacca decandra*.

BLOCK TEETH, see Teeth, block.

BLOOD, (Sax. *bloo*, from *blean*, 'to bleed,') *San'guis*, *Cr'uor*, *La'pis anima'lis*, *Hæ'ma*, (Old Eng.) *Blede*, (F.) *Sang*. An animal fluid formed chiefly from the chyle; acquiring important properties during respiration; entering every organ through the circulation; distributing the nutritive principles to every texture, and the source of every secretion. The blood is white in the molluscous and inferior animals, which have been, hence, called *white-blooded*, to distinguish them from the *red-blooded*, which class includes the mammalia, birds, reptiles, and fishes. Human blood is composed of water, albumen, fibrin, an animal colouring substance, a little fatty matter—*hemateleum*—and different salts; as chlorides of potassium and sodium, phosphate of lime, subcarbonate of soda, lime, magnesia, oxide of iron, and lactate of soda, united with an animal matter. *Arterial blood* is of a florid red colour, strong smell; s. g. 1.049. *Venous blood* is of a brownish red; s. g. 1.051. The difference in colour has given occasion to the first being called *red blood*; the latter, *black*. The former, which is distributed from the heart, is nearly the same through its whole extent: the latter is the remains of the arterial blood after the different elements have been taken from it in nutrition, with the addition of the products of absorption. Blood, taken from a vessel and left to itself, becomes solid, and separates into two distinct parts,—the *serum* or watery supernatant

fluid; and the *cruru*, *coagulum*, *crassamentum*, *hepar seu placenta sanguinis*, *placenta cruris*, *insula*, *thrombus*, or *clot*. The serum is chiefly water holding albumen in solution and the salts of the blood. The clot contains the fibrin, colouring matter—*hæmotosin*, a little serum, and a small quantity of salts. M. Le Canu found the blood to be composed—in 1000 parts—of water, 785.590; albumen, 69.415; fibrin, 3.565; colouring matter, 119.626; crystallizable fatty matter, 4.300; oily matter, 2.270; extractive matter soluble in alcohol and water, 1.920; albumen combined with soda, 2.010; chlorides of sodium and potassium; alkaline phosphates, sulphates, and subcarbonates, 7.304; subcarbonate of lime and magnesia, phosphate of lime, magnesia and iron, peroxide of iron, 1.414; loss, 2.586. The four principal components of the blood are fibrin, albumen, corpuscles, and saline matter. In the circulating blood they are thus combined—

Fibrin, }
Albumen, } In solution forming *Liquor Sanguinis*.
Salts, }

Red and white Corpuscles, suspended in the *Liquor Sanguinis*.

In coagulated blood they are thus combined:

Fibrin, } Forming the *crassa-*
Red and white Corpuscles, } *mentum* or clot.
Albumen, } Remaining in solution, forming
Salts, } *serum*.

The following table exhibits the computations of different physiologists regarding the weight of the circulating fluid—arterial and venous.

Harvey,		
Lister,		
Moulin,		
Abildgaard,		
Blumenbach,		
Lobb,		
Lower, and Lewes,		10
Sprengel,	10 to 15	
Günther and Bock	15 to 20	
Blake	16½ to 18½	
Weber and Lehmann	17½ to 19	
Müller, Burdach, and P. Bérard	20	
Wagner	20 to 25	
Quesnai	20 to 27	
F. Hoffmann	28	
Haller	28 to 30	
Young	40	
Hamberger	40	
Keill	80	
	100	

The estimates of Günther, Bock, and Blake are probably approximations to the truth.

The proportion of arterial blood to venous is about 4 to 9.

Much attention has been paid to the varying condition of the blood in disease. The average proportion of each of the organic elements in 1000 parts of healthy blood is as follows, according to Le Canu, and MM. Andral and Gavarret:—fibrin, 3; red corpuscles, 127; solid matter of the serum, 80; water, 790.

Dried human blood was, at one time, considered to be anti-epileptic; that of the goat, dried, *Sanguis hirci siccatus*, sudorific and antipleuritic. The dried blood of the ox—*Extractum Sanguinis Bovini*—and the dried red corpuscles have been given as analeptics, especially where there was a deficiency of red corpuscles.

BLOOD, ARTERIAL, see Blood—b. Black, see Blood—b. Black, vascular system of, see Vascular.

BLOOD CASEIN, Globulin—b. Corpuscles, Globules of the blood—b. Crystals, Hæmatoidin—b. Disease, Hæmatonosis—b. Disks, Globules of the blood.

BLOOD, DRIED, see Blood.

BLOOD FUNKERS, see Hæmatophobia—b. Glands, see Ganglion.

BLOOD, INFLAMMATORY, see Hæmitis—b. Loss of, Hæmorrhagia—b. Menstrual, see Menses—b. Red, see Blood—b. Red, system of, see Vascular—b. Spitting of, Hæmoptysis—b. Venous, see Blood—b. Vomiting of, Hæmatemesis—b. White, Lymph.

BLOODING, Blood-letting.

BLOODLESSNESS, Anæmia.

BLOODLETTING, *Mis'sio seu Detractio Sanguinis*, *Hæmac'is*, *Cataschas'mus*, *Blood'ing*, *Blood'ing*, *Let'ting blood*, (F.) *Saignée*, *Emission sanguine*. A discharge of a certain quantity of blood produced by art: an operation which consists in making an opening into a vessel to draw or let blood from it. When practiced on an artery, it is called *Arteriotomy*; on a vein, *Phlebotomy*, *Venæsectio*, *Venesection*, *Breath'ing a vein*, and on the capillary vessels, *local* or *capillary*, in contradistinction to the former, which is termed *general*. Bloodletting is used both during the existence of a disease, as in inflammation, and in the way of prophylaxis. It is employed to fulfil various indications. 1. To diminish the actual mass of blood;—when it is termed, by the French pathologists, *Saignée évacuative*. In such case, fluids ought not to be allowed too freely afterwards. 2. To diminish the turgescence in any particular organ—(F.) *Saignée révulsive*, *Revulsive bloodletting* or *bleeding*, *Venæsectio revulsoria*, when performed far from the part affected; and *Saignée dérivative*, when near. 3. To diminish the corpuscles of the blood, (F.) *Saignée spoliative*, *Spoliative Phlebotomy*. The immediate effects of blood-letting are: diminution of the mass of blood and of heat; retardation of the pulse, and sometimes syncope. Bloodletting from the veins—*phlebotomy*—is practised on the subcutaneous veins of the neck, the face, the fore-arm, and the leg; sometimes on those of the hand or foot. The necessary apparatus consists of a bandage or riband, a compress of rag, and a lancet or fleam.

The veins selected for the operation, are, 1. *In the fold of the arm*, five;—the cephalic, basilic, the two median, and the anterior cubital. 2. *In the hand*, the cephalic and salivata. 3. *In the foot*, the great and little saphena. 4. *In the neck*, the external jugular. 5. *In the forehead*, the frontal. 6. *In the mouth*, the ramine. The operation of phlebotomy in the limbs is performed by tying a circular bandage round the limb, in order that the subcutaneous veins may become turgid by the course of the blood being obstructed: the bandage not being so tight, however, as to compress the arteries of the limb. A puncture is made into the vein, and the desired quantity allowed to flow. The ligature is now removed, and a compress and retaining bandage applied. *Capillary or local bloodletting*, *arterio-phlebotomy*, is practised on the skin or mucous membranes, by means of leeches, the lancet, or cupping.

BLOODLETTING, CAPILLARY, see Bloodletting—b. Derivative, see Bloodletting—b. Evacuative, see Bloodletting—b. General, see Bloodletting—b. Local, see Bloodletting—b. Revulsive, see Bloodletting—b. Spoliative, see Bloodletting.

BLOODLIKE, Sanguine.

BLOOD LIQUOR, *Liquor Sanguinis*.

BLOODROOT, *Sanguinaria Canadensis*.

BLOODSHOT, Hyperæmia.

BLOODSTONE, Hæmatites.

BLOODSTROKE, *Coup-de-sang*.

BLOODVESICLE, Globule of the blood.

BLOODVESSEL, (F.) *Vaisseau sanguin*.

vessel destined to contain and convey blood.

BLOODVESSEL, BREAKING, BURSTING, RUPTURING, OF A. Hæmorrhagia.

BLOODWEED, *Asclepias curassavica*.

BLOODWORT, *Hieracium venosum*, *Sanguinaria Canadensis*.

BLOOD'Y, *Hæmac'us*, *Hæmat'icus*, *Sanguin'eus*, *Cruen'tus*, *Sanguin'eus*, (F.) *Sanguin*. Having the character of blood. Relating to blood. See Sanguine.

BLOODY FLUX, Dysentery.

BLOOM, **HONEY**, *Apocynum androsæmifolium*.

BLOSS'BURG MIN'ERAL SPRINGS. The town of Blossburg is in Tioga County, 133 miles north of Harrisburg. The waters contain free sulphuric acid, and sulphates of iron, alumina and magnesia, and are astringent and tonic.

BLOSSOM, see Grog-blossom.

BLOTA ALBA, Bleta.

BLOTCH, (from *blot* [?]) (Prov.) *Splav'in*, *Sproté*. A pustule; more commonly, an eruption of a large kind.

BLOW, *Ictus*, *Pla'gè*, (F.) *Coup*. Effect produced by one body striking another. The impression made by any body which strikes us, or against which we strike;—a common cause of wounds, contusions, fractures, &c.

BLOWING SOUND, *Bruit de Souffle*—b. s. Diffused, see Murmur, respiratory.

BLUEBELLS, *Gentiana Catesbæi*.

BLUEBERRY, *Caulophyllum thalictroides*, *Lantana*—b. Low, *Vaccinium Pennsylvanicum*.

BLUEBONNETS, *Centaurea cyanus*.

BLUEBOTTLE, *Centaurea cyanus*, *Cyanus segetum*.

BLUE LICKS, see Kentucky, Mineral Waters of.

BLUE MASS, *Pilulæ Hydrargyri*—b. Stone, *Capri sulphas*.

BLUET, ('blue bottle,') *Centaurea cyanus*.

BLUET DES MOISSONS, ('blue bottle of the harvests,') *Cyanus segetum*.

BLUTER (Sc.), *Borborismus*.

BLUSH, (D. *blos*), see Flush—b. Cutaneous, see Efflorescence—b. Inflammatory, *Erythema*.

BLUTTER (Sc.), *Borborismus*.

BO'A, *Bo'ia*. An eruption of red, ichorous pimples. See, also, *Hidroa* and *Sudamina*.

BOA UPAS, *Upas*.

BOÆ, Syphilis.

BOALA, see Scherlievo.

BOBERRI, *Curcuma longa*.

BOCCHOE, *Diosma crenata*.

BOCHE (Prov.), Swelling.

BOCHET, *Bochetum*.

BOCHE'TUM, (F.) *Bochet*. A term formerly applied to the second decoction of the sudorific woods. The French word has been lately revived.

BOCIUM, *Bronchocele*.

BOCHO, *Diosma crenata*.

BOCIUM, *Bronchocele*.

BOCK (Sc.), *Vomiturition*.

BOCKING (Sc.), *Eruetation*, *Vomiting*.

BOCKLET, MIN'ERAL WATERS OF. The springs of Bocklet, in Bavaria, are acidulous chalybeates.

BOD'Y, *Cor'pus*, *So'ma*, (F.) *Corps*; (Sax. *bodig*, Teutonic *boden*, the 'fundus or bottom.' [?]) The human body, (F.) *Corps humain*, (Old E.) *Liche*, *Corps*, is the collection of organs which compose the frame. At times, however, body is used synonymously with trunk. We say, also, *body of the femur*, of the *sphenoid*, &c., to designate the shaft or middle portion of those bones; *body of the uterus*, &c. Likewise, any distinct portion of matter perceived by the senses. Also, the rectum.

BODY, COMING DOWN OF THE, *Proctocele*—b. of Rosenmüller, *Parovarium*.

BODYSEARCH'ER, *Search'er*. One who formerly examined the bodies of the dead to report as to the cause of death. In the time of the plague this was done by "the chirurgéons," who were allowed "twelve pence," to be paid out of the

goods of the party searched, if able; otherwise, by the parish. Until of late, this office was executed in England by two old women—called 'searchers'; and hence the imperfections of the Bills of Mortality. In plague times, the office was an important one; and, it is stated, that a noted searcher, named Snacks, finding his business increase so much, that he could not transact it alone, offered to any one who would join him in its hazards, half the profits; and they who joined him were said to "go with Snacks." Hence, the saying "to go Snacks," or to divide the spoils.

BODYSNATCHER, Resurrectionist.

BOE, (*βοη*), Cry.

BOELLI, Intestines.

BOETHEMA, (*βοηθημα*, from *βοηθω*, 'to aid,') Medicament.

BOG-BEAN, *Menyanthes trifoliata*.

BOHON UPAS, *Upas*.

BOIA, Boa.

BOIL, (Sax. *býl*.) *Furunculus*—b. Blind, see *Furunculus*—b. Delhi, see *Delhi*—b. Gum, *Parulis*—b. Malignant, see *Furunculus*—b. Wasp's nest, see *Furunculus*.

BOIS AMER, ('bitterwood,') *Quassia*—b. de *Campêche*, *Hæmatoxylinum Campechianum*—b. de *Cervi*, *Cornu cervi*—b. de *Chypre*, *Rhodium lignum*—b. de *Couleuvre*, see *Strychnos*—b. *Doux*, *Glycyrrhiza*—b. de *Marais*, *Cephalanthus occidentalis*—b. de *Plomb*, *Direa palustris*—b. *Puant*, *Prunus padus*—b. de *Rose*, *Rhodium lignum*—b. de *Sappan*, *Cæsalpinia sappan*—b. *Sudorifique*, Wood, sudorific.

BOISSE, MIN'ERAL WATERS OF. These waters are situate about half a league from Fontenay-le-Compte, in France. They are purgative, and seem to contain carbonate and sulphate of lime and chloride of sodium.

BOISSON, (from *boire*, 'to drink,') Drink.

BOIST, (Prov.), Swelling.

BOÎTE, (F.) *A box or case*, *Cap'sa*, *Pyx'is*. An apparatus for the reception of any matters which it may be desirable to preserve. In *Surgery* and *Anatomy*, *Boîtes à dissection*, *B. à amputation*, *B. à trépan*, *B. à cataracte*, &c., mean the cases containing appropriate instruments. *Boîte du Crane* is the bony case which receives the brain. *Boîte* is, also, the portion of the stem of the trephine which receives the pyramid or centre-pin. *Boîte de Petit* is a machine, invented by M. Petit, to retain the fractured portions of bone in apposition, when the leg has been fractured in a complicated manner. *Boîte* is, also, a kind of case put before an artificial anus to receive the fæces, which are continually being discharged. The vulgar, in France, give the name *Boîte*, to various articulations,—*B. de genou*, *B. de la hanche*; "knee-joint, hip-joint."

BOITEMENT, Claudication.

BOITERIE, Claudication.

BOÎTIER, (F.) *Appareil*, *Cap'sula unguenta'ria*, *Cap'sarium*, *Dress'ing-case*. A box containing salves and different apparatus, used more particularly by the dressers in hospitals.

BOK (Sc.), *Vomiturition*.

BOCKING (Sc.), *Eruetation*, *Vomiting*.

BOL, *Bolus*—b. d'Arménie, *Bol*, Armenian—b. Blanc, *Bolus alba*—b. Oriental, *Bol*, Armenian—b. Rouge, *Bol*, Armenian.

BOLA, Myrrha.

BOLCHON, *Bdellium*.

BOLE, *Bolus*, (*βωλος*, 'a clod of earth,') *Bol*, *Terre bolaire*, meant, with the older writers, argillaceous earth, used as an absorbent and alexipharmic. The various boles had different forms given to them, and were stamped, or sealed, hence called *Ter're sigillatæ* (F.) *Terres sigillées*, as in the following:

BOLE ARME'NIAN, *Bole Armeniac*, *B. Ar'me-nic*, *Argil'la ferrugin'ea ru'bra*, *A. Bo'lus ru'bra*, *Arena'men*, *Bo'lus Oriental'is* seu *Armen'aca* seu *Arme'nia* seu *Arme'na* seu *ru'bra*, (F.) *Bol d'Armenie* ou *oriental ou rouge*. A red, clayey earth, found not only in Armenia, but in several countries of Europe,—in Tuscany, Silesia, France, &c. It was once esteemed a tonic and astringent, and was applied as a styptic. It is now, scarcely, or never, used. It consists of argil, mixed with lime and iron.

BOLESIS, Coral.

BOLESON, Balsam.

BOLET ODORANT, *Dædalea suaveolens*.

BOLETUS AGARICUS, (*βωλος*, 'a lump or mass,' referring to its shape,) *B. Laricis*—*b. Albus*, *Boletus laricis*—*b. Cervinus*, *Elaphomyces granulatus*—*b. Discoides*, *Dædalea suaveolens*.

BOLETUS ESCULENTUS, (F.) *Morelle*. *Ord. Fungi*. An eatable mushroom, found in the woods in Europe, and much admired by *Gastro-nomes*. It was formerly esteemed to be aphrodisiac.

BOLETUS FULVUS, *B. ignarius*—*b. Hippocrepis*, *B. ignarius*.

BOLETUS IGNARIUS. The systematic name for the *Ag'aric*, *Agar'icus*, *Agar'icum* of the Pharmacopœias, *Agar'icus Chirurgo'rum* seu *Quer'cus seu ignia'rius*, *Polyp'orus ignia'rius*, *Is'ca*, *Boletus unguilatus* seu *fulvus* seu *hippocrep'is* seu *obtusus*, *Fun'gus Ignia'rius*, *Fun'gus Querc'inus*, *Spunk*, *Am'adou*, *Punk*, (Prov.) *Funk*, *Ag'aric* of the *Oak*, *Touch'wood*, *Touch'wood Boletus*, *Fe'male Ag'aric*, *Tin'der*, (F.) *Agaric de chêne*, *Amadouvier*. It was formerly much used by surgeons as a styptic.

BOLETUS LARICIS, *B. Larici'us* seu *pur'gans* seu *al'bus* seu *agar'icus* seu *officina'lis*, *Fun'gus Lar'icis*, *Polyp'orus officina'lis*, *Agar'icus al'bus* seu *Lar'icis*, *A. Al'bus opt'imus*, *White Ag'aric*, (F.) *Agaric blanc*, *A. Amadouvier*, *Polypore du Mélèze*. On the continent of Europe it has been given as a cathartic and emetic, as well as to moderate the sweats in phthisis. Externally, styptic.

A tincture of the *Boletus Laricis Canadensis* has been highly recommended by Dr. Grant, of Ottawa, in acute rheumatism.

BOLETUS OBTUSUS, *B. ignarius*—*b. Officinalis*, *B. laricis*—*b. Purgans*, *Boletus laricis*—*b. Salicis*, *Dædalea suaveolens*—*b. Suaveolens*, *Dædalea suaveolens*—*b. Touchwood*, *Boletus ignarius*.

BOLI MARTIS, *Ferrum tartarissatum*.

BOLISMOS, (*βωλος*, 'a mass,') *Boulimia*.

BOLITES, (*βωλιτης*, from *βωλος*, 'a lump or mass,') The mushroom; perhaps the *Agar'icus Auranti'acus*. It was so called in consequence of its shape.

BOLLYNGE, Swelling.

BOLNING, Swelling.

BOLT. Used, at times, for to swallow without chewing—as to "bolt one's food."

BO'LUS, (*βωλος*, 'a lump or mass,') A bole, (F.) *Bol*. A pharmaceutical preparation, having a pilular shape, but larger; capable, however, of being swallowed as a pill. Also, bole.

BOLUS AL'BA, *Argil'la pallid'ior*:—commonly made into small cakes or flat masses, and stamped or sealed with certain impressions, and called *Terræ sigillatæ*. (F.) *Bol blanc*, *Argile ochreuse pâte*. It was used like *Bole Armenian*, and was brought from Etruria. See Bole.

BOLUS ALIMENTARY, *Bolus Alimentari'us*. The bole formed by the food, after it has undergone mastication and insalivation in the mouth; and been collected upon the tongue prior to deglutition.

BOLUS ARMENA, Bole, Armenian.

BOLUS ORIENTA' LIS. A kind of bolar earth,

only distinguished from *Bole Armenian* in being brought from Constantinople. See *Bole Armenian*.

BOLUS RUBRA, *Bole Armenian*.

BOMA'REA SALSIL'LA, (after the naturalist, J. C. Valmont de Bomare.) The inhabitants of Chili use this plant as a sudorific. It is given in infusion in cutaneous diseases.

BOMBAX, (*βουβυξ*, 'raw silk,') *Gossypium*.

BOMBEMENT, *Bombus*.

BOMBITATIO, *Tinnitus aurium*.

BOMB'BUS, (*βουβυξ*, 'the humming of bees,') *Aur'ium fluctua'tio*, *A. Sib'ilus*, *A. Son'itus*, *A. Susur'rus*, (F.) *Bombement*. A kind of ringing or buzzing in the ears;—characterized, according to SAUVAGES, by the perception of blows or beating repeated at certain intervals. Also, *Borborygmus*. See *Fart*, *Flatulence*, and *Tinnitus Aurium*.

BOMB'BYX MORI, (*βουβυξ*), see *Sericum*.

BON, *Coffea Arabica*.

BONA, *Phaseolus vulgaris*.

BONA FEVER, see *Fever*, *Bona*.

BONANNIA OFFICINALIS, *Sinapis alba*.

BONDUE, *Gymnoeladus Canadensis*.

BONE, ([Sax.] *ban*), *Os*, *Os'teon*, *Os'teum*, (F.) *Os*. The bones are the solid and hard parts, which form the basis of the bodies of animals of the superior classes; and the union of which constitutes the *skeleton*. The human body has, at the adult age, 240 bones, including the 32 teeth, the ossa Wormiana, and excluding sesamoid bones. Anatomists divide them, from their shape, into 1. *Long bones*, which form part of the limbs, and represent columns for supporting the weight of the body, or levers of different kinds for the muscles to act upon. 2. *Flat bones*, which form the parietes of splanchnic cavities; and, 3. *Short bones*, met with in parts of the body where solidity and some mobility are necessary. Bones are formed of two different textures; *spong'ly*, *Substantia spongio'sa*; and *compact*, *Substantia compacta*. They afford, on analysis, much phosphate and carbonate of lime, a little phosphate of magnesia, phosphate of ammonia, oxide of iron and manganese, some traces of alumina and silica, gelatin, fat, and water. The uses of the bones are mentioned under each bone. They give shape to the body, contain and defend the viscera, and act as levers to the muscles.

The bones of the ox, *Bos Tau'rus*, are employed in pharmacy.

TABLE OF THE BONES.

BONES OF THE HEAD.	Bones of the Cranium or Skull.	Frontal.....	1
		Parietal.....	2
		Occipital.....	1
		Temporal.....	2
		Ethmoid.....	1
		Sphenoid.....	1
		Superior Maxillary.....	2
		Jugal or Cheek.....	2
		Nasal.....	2
		Lachrymal.....	2
	Bones of the Face.	Palatine.....	2
		Inferior Spongy.....	2
		Vomer.....	1
		Inferior Maxillary.....	1
		Incisors.....	8
	Dentes, or Teeth.	Cuspidati.....	4
		Molars.....	20
	Bone of the Tongue.	Hyoïd.....	1
		Malleus.....	2
	Bones of the Ear.	Incus.....	2
		Orbitulare.....	2
		Stapes.....	2
		Cervical.....	7
	Vertebrae.	Dorsal.....	12
		Lumbar.....	5
BONES OF THE TRUNK.	Sacrum.....		1
		Os Coccygis.....	1
	The Thorax.	Sternum.....	1
		Ribs.....	24

TABLE OF THE BONES.—Continued.

TABLE OF THE BONES.—Continued.					
BONES OF THE UP- PER EX- TREMITY.		The Pelvis.....	Innominatum.....	2	
		The Shoulder.....	Clavicle.....	2	
			Scapula.....	2	
		The Arm.....	Humerus.....	2	
		Fore-Arm.....	Ulna.....	2	
			Radius.....	2	
		The Hand. Carpus, or Wrist.	Naviculare.....	2	
			Lunare.....	2	
			Cuneiforme.....	2	
			Oribiculare.....	2	
			Trapezium.....	2	
			Trapezoides.....	2	
			Magnum.....	2	
			Unciforme.....	2	
		Metacarpus.....		10	
	Phalanges.....		28		
	The Thigh.....	Femur.....	2		
BONES OF LOWER EX- TREMITY.		The Leg.	Patella.....	2	
			Tibia.....	2	
			Fibula.....	2	
			Calcis Os.....	2	
		The Foot.	Tarsus, or Instep.	Astragalus.....	2
				Cuboïdes.....	2
				Naviculare.....	2
				Cuneiforme.....	6
			Metatarsus.....		10
		Phalanges.....		28	
		Total.....			240

BONEACHE, Osteocopus — b. Ague, Osteocopus.

BONE ASH. The product of bones, usually of the ox and sheep, burnt white, consisting chiefly of phosphate of lime and a little of the carbonate. Used in pharmacy.

BONE, BACK, Vertebral column — b. Bar, Pubis, os.

BONE BLACK, see Carbo animalis — b. Canals, see Canals, nutritive, and Diploë.

BONE, BLADE, Scapula — b. Boat-like, Os scapuloïdes — b. Breast, Sternum — b. Crupper, Coccyx.

BONE EARTH, see Cornu ustum — b. Fever, see Inflammation.

BONE, HAUNCH, Ilion — b. Interparietal, Interparietal bone.

BONE PHOSPHATE OF LIME, see Cornu ustum.

BONE, RUMP, Coccyx — b. Share, Pubis — b. Splinter, Fibula.

BONEBINDER, Osteocolla.

BONEDOCTOR, Renouëur.

BONENIP'PERS, Osteu'cum, Tenac'ula, (F.)

Tenaïlle incisive. An instrument used for cutting off splinters and cartilages. It is a kind of forceps, the handles of which are strong, and the edges, which touch each other, cutting.

BONESET, Eupatorium perfoliatum — b. Rough, Eupatorium tucrifolium — b. Upland, Eupatorium sessilifolium.

BONESETER, Renouëur.

BONESHAVE, Neuralgia femoro-poplitea.

BONES, BRITTLINESS OF THE, Fragilitas ossium — b. Friability of the, Fragilitas ossium — b. Salt of, Ammonie carbonas — b. Softening of the, Mollities ossium.

BONHOMME ([F.] 'good man,') Verbasum thapsus.

BONICLATTER, Bonnyclabber.

BONIFACIA, Ruscus hypoglossum.

BONITHLOBER, Bonnyclabber.

BONNE DAME ([F.] 'good lady,') Atriplex hortensis.

BONNES or AIGUES BONNES, MIN'ERAL WATERS OF. Bonnes is a village six leagues from Pau, in the department Basses Pyrénées, France. Here are several thermal springs. They were celebrated as early as the time of Francis I., under the name Eaux d'Arquebuse. They contain chlorides of sodium and magnesium, sulphates of magnesia and lime, sulphur, and silica. The temperature is from 78° to 98° Fahrenheit. The season is from the first of May to the middle of September.

The factitious EAU DE BONNES is made of Hydro-sulphuretted water, f 3 iv; pure water, Oj, and f 3 ss; chloride of sodium, gr. xxx; sulphate of magnesia, gr. i.

BONNET, Reticulum.

BONNET À DEUX GLOBES, Bonnet d'Hippocrate.

BONNET D'HIPPOCRATE, Cap of Hippocrates, Mitra Hippocrati'ca, Fascia capituli, Pileus Hippocrati'cus. A kind of bandage, the invention of which is ascribed to Hippocrates. It consists of a double-headed roller, passed over the head, so as to envelop it like a cap. The French also name it Bonnet à deux globes, Capeline de la tête.

BON'NYCLABBER, Cla'ber, (Irish, baine, 'milk,' and clabar, 'mire,') In Ireland, sour buttermilk. In this country, the thick part of sour milk. Boniclatter and Bonithlobber, also, mean cream or "good milk gone thick."

BONPLANDIA ANGUSTURA, (after the French naturalist Bonpland,) Cusparia febrifuga — b. Trifoliata, Cusparia febrifuga.

BONTIA GERMINANS, (after Jacob Bontius, a Dutch physician of Batavia,) Avicennia tomentosa.

BONUS GENIUS, Peucedanum — b. Henricus, Chenopodium bonus Henricus.

BONY, (from bone,) Osseous.

BOOCHO, Diosma crenata.

BOON UPAS, Upas.

BOONA, Phaseolus vulgaris.

BOOTIA VULGARIS, Saponaria.

BOOT'IKIN, (dim. of boot.) A glove with a partition for the thumb, but no separate ones for the fingers—like an infant's glove—made of oiled silk. Horace Walpole speaks in raptures of the benefit he derived from bootikins in gout.

BORAC'IC ACID, Ac'idum Borac'icum, Sal sedati'vus HOMBER'GI, Bo'ric Acid, (F.) Acide boracique. An acid obtained from borax, which was once looked upon as sedative. It was also called Acor Borac'icus, Sal vitri'oli narcoti'cum, Sal volat'ile Borac'is, and Flores Borac'is.

BORAGE, Borago officinalis.

BORA'GO OFFICINA'LIS, Buglos'eum ve'lum seu latifol'ium, Borra'go, Corra'go, Borago hortens'is, Bor'age, Bur'rage, (F.) Bourrache. Ord. Boraginæ. Sex. Syst. Pentandria Monogynia. The leaves and flowers have been considered aperient.

BORAS SUPERSODICUS, Borax.

BO'RATED, Borat'us, (F.) Boraté. Containing borax, as 'a borated solution,' (F.) Solution boratée.

BORATHRON, Juniperus Sabina.

BO'RAX, gen. Borac'is, Bo'ras So'dæ seu supersat'urus so'dæ seu alcal'es'cens seu alcali'num seu superso'dicus, So'dæ Bibo'ras, Subbo'ras So'dæ seu nat'ricum, So'da Borac'ia, Natrum biborac'i'cum seu subborac'i'cum, Chrysocol'la, Capi'strum au'ri, Bo'rax Ven'etus, Borax'irion, Nitrum facti'tium, &c., Subbo'rate or Bibo'rate of Soda, Subbo'rate of protox'ide of So'dium, Subprotobo'rate of So'dium, Bo'rate of So'da, (F.) Borate ou Soursorbaté de Soude, Borate sursaturé de soude. It is found in an impure state in Tibet and Persia. It is inodorous; taste cool, and somewhat alkaline; soluble in 12 parts of water. Borax is seldom used except as a lotion in apthæ.

Borate of Mercury has been recommended as an antisypilitic.

BORAXTRION, Borax.

BORBON'IA RUSCIFO'LIA, (after Gaston de Bourbon, son of Henry IV., of France, who was fond of botany.) A small South African shrub, used in asthma and hydrothorax. In decoction, it is given as a diuretic.

BORBORUS, (Bor'boros.) Fimus.

BORBORYGMUS, (Bor'boru'gus, 'I make a dull noise,') Mur'mur seu Bom'bus seu Mo'tus Intestino'rum, Anile'ma, Anile'sia, Catoproph'ia, Intona'tio intestinal'is, Mur'mur vent'ris seu intestina'le, Bor'borism, (So.) Blutter, Blutter, (Prov.) Crolling, Crolling, (F.) Gargouillement, Grouillement d'Entrailles. The noise made by flatus in

the intestines. This happens often in health, especially in nervous individuals.

BORD (F.), *Mar'go, Edge, Marg'in.* Anatomists have so named the boundaries of an organ. Thus, the bones, muscles, &c. have *bords* as well as bodies. The 'free edge,' *bord libre*, is one not connected with any part; the 'adhering edge,' *bord adhérent*, one that is connected; and the *bord articulaire*, or 'articular margin or edge' of a bone, that which joins it to another bone.

BORD CILIAIRE, Ciliary margin.

BORDEAUX, MIN'ERAL WATERS OF. Near this great city, in the south-west of France, is a saline, chalybeate spring. It contains oxide of iron, carbonate and sulphate of lime, chlorides of sodium and calcium, subcarbonate of soda, and sulphate of magnesia.

BORE, Boron.

BORE-TREE, (from being easily *bored* by removing the pith,) *Sambucus*.

BORGNE (F.), *Coc'les, Unoc'ulus, Lus'cus, Luscio'cus*. One who has only one eye, or sees only with one. The word has been used, figuratively, for *blind*, in surgery and anatomy. See *Cæcus*.

BORIUM, Boron.

BORKHAUSENIA CAVA, (after Moritz Borkhausen, a German botanist,) *Fumaria bulbosa*.

BORN, (past participle of *bear*,) (F.) *né*. Brought forth from the womb.

BORN ALIVE. It has been decided by English judges, that 'to be born alive,' means, that acts of life must have been manifested after the whole body has been extruded; and that respiration *in transitu* is not evidence that a child was born alive. It must be 'wholly born alive,' hence respiration may be a sign of *life*, but not of *live birth*.

BORON, *Bo'rium, Bo'rum*, (F.) *Bore*. A simple substance, the basis of boracic acid; obtained, by heating potassium with boracic acid, as a dark, olive-coloured powder, devoid of taste and smell. Heated in the air or in oxygen, it is converted into boracic acid.

BOR'OSAIL, *Zael*. Æthiopian names for a disease, very common there, which attacks the organs of generation, and appears to have considerable analogy with syphilis.

BORRAGO, *Borago officinalis*.

BORRI, *Curcuma longa*.

BORRIBERRI, *Curcuma longa*.

BORSE, MIN'ERAL WATERS OF. Borse is a village in Béarn. The waters are chalybeate.

BORUM, Boron.

BO'SA. An Egyptian name for a mass, made of the meal of darnel, hemp-seed, and water. It is inebriating. At the present day, it is applied to a kind of acidulated and, sometimes, fermented scherbet. In its commonest form, it is made by fermenting an infusion of millet-seed. A more agreeable Bosa is prepared by the Egyptians from tamarinds and honey; and by the Greeks from the fruit of *Solanum Lycopersicum*. This last is called, by them, *Δροσικόν*, *Drosi'ticon*.

BOSCHESJESMANSTHEE, *Methystophyllum glaucum*.

BOSOM, (Sax. *bofme*), see *Mamma*.

BOSS, *Bosse*.

BOSSA, Plague token.

BOSSE, ([L] *bozza*, 'a bunch,) Hump, Protuberance—b. *Nasale*, Nasal protuberance.

BOSSU, see *Hump*.

BOSWELLIA FLORIBUNDA, (after Dr. John Boswell, of Edinburgh,) see *Juniperus lycia*—b. *Papyrifera*, see *Juniperus lycia*—b. *Serrata*, see *Juniperus lycia*—b. *Thurifera*, see *Juniperus lycia*.

BO'TAL FORA'MEN, *Fora'men Bota'lè* seu *Bota'li*, *Fora'men Ova'lè*, (F.) of *Galen* or of

Botal, (F.) *Trou de Botal ou de Galien ou ovale*. A large opening which exists in the fetus in the partition between the two auricles of the heart; and by means of which the blood passes from one to the other. Its discovery is generally attributed to Leonard Botallus, Botal, or Botalli, who wrote in 1562. It was spoken of, however, by *Vesalius*, and even by *Galen*.

BOTANE, (*Borane*), Herb.

BOTANICAL DOCTOR, Herb-doctor.

BOTANIQUE MÉDICALE, Botany, medical.

BOT'ANY, MED'ICAL, *Botan'ica Med'ica, Medici'na Botan'ica, Phytolog'ia Med'ica*, (from *botane*), (F.) *Botanique Médicale*. The knowledge of the properties, characters, &c. of those vegetables which are used in medicine.

BOTAR'GO, (F.) *Botargue*. A preparation made in Italy and the south of France, with the eggs and blood of the *Mugileceph'alus* or *Mullet*, strongly salted, after it has become putrescent. It is used as a condiment.

BOTARGUE, Botargo.

BOTCH, ([L] *Bozza*, (F.) *Bosse*.) Of old, a swelling of the skin. A phlegmon. A large ulcerous affection.

BOTHOR. An Arabic term for abscess in the nares. It means, also, a tumour in general; especially those which are without solution of continuity.

BOTHRIOCEPH'ALUS, *B, la'tus, Bothrioceph'alum*, (*Βοθρίον*, 'a small pit,' and *κεφαλή*, 'head.') *Botriocceph'alus, Diboth'rion la'tum, Te'nia lata seu vulga'ris seu os'culis lateral'ibus gem'inis seu gri'sea seu membrana'cea seu tenel'la seu denta'ta seu huma'na iner'mis seu pri'ma seu os'culis lateral'ibus solita'ris seu aceph'ala seu os'culis superficial'ibus, Lumbr'icus la'tus, Plate'a, Platyl'mine, Haly'sis membrana'cea seu la'ta*, (F.) *Ténia d'anneaux courts ou non armé, Ver solitaire, Broad Tapeworm*. Common in Switzerland, Russia, and some parts of France. It inhabits the intestines of man, and extends to an enormous length. A broken specimen has been obtained 60 yards long.

BOTH'RION, *Both'rionum*, (*Βοθρίον*, 'a small pit, cavity,' &c.) An *alveolus*, or small fossa. A small deep ulcer on the cornea. See *Fossette*.

BOTHRION, Bothrion, *Fossette*.

BOTHRUS, (*Βοθρος*), Fovea.

BOTIN, Terebinthina.

BOTIUM, Bronchocele.

BOTOTHINUM. An obscure term, used by *Paracelsus* to denote the most striking symptom of a disease:—the *Flos mor'bi*.

BOTOU, Pareira brava.

BOTRIOCEPHALUS, *Bothriocephalus*.

BOTROPHIS SERPENTARIA, *Actæa racemosa*.

BOTRYS, (*Βοτρυς*, 'a grape, a cluster of grapes,' owing to the resemblance of the fructification.) *Chenopodium botrys*, see *Vitis vinifera*—b. *Ambrosioides*, *Chenopodium ambrosioides*—b. *Americana*, *Chenopodium ambrosioides*—b. *Anthelmintica*, *Chenopodium anthelminticum*—b. *Mexicana*, *Chenopodium ambrosioides*.

BOTTINE, ([F.] dim. of *botte*, 'a boot.') A thin boot or bus'kin, *O'crea le'vior*. An instrument which resembles a small boot, furnished with springs, straps, buckles, &c., and used to obviate distortions of the lower extremities in children.

BOTTLE, FEED'ING. A bottle, provided with a projection to which an artificial nipple can be attached. Used for feeding infants with milk.

BOTTLENOSE, *Gutta rosea*.

BOTTLESTOOP. In Pharmacy, an arrangement for giving the proper inclination to a bottle containing a powder, so as to admit of the contents being readily removed by the knife, in dia-

persing medicines. It consists of a block of wood with a groove in the upper surface, to receive the bottle in an oblique position.

BOTTOM, (Sax. *botm*.) Nates.

BOTTRYTREE, Sambucus.

BOU, in composition, see Bu.

BOUBON, (βουβων, 'the groin.') Bubo.

BOUCAGE ANIS, *Pimpinella anisum*—*b. Majeur*, *Pimpinella magna*—*b. Mineur*, *Pimpinella saxifraga*—*b. Petit*, *Pimpinella saxifraga*.

BOUCHE, (bucca, 'the puffed out cheek.') Mouth—*b. Pâteuse*, Clammy mouth.

BOUCLEMENT, ([F.] *boucle*, 'a buckle.') Infibulation.

BOUE, ([F.] 'mire, mud.') See Corruption.

BOUE SPLÉNIQUE. The altered blood extravasated into the splenic cells has been so called.

BOUES DES EAUX, (F.) *Boues Minérales*. The mud or swamp, formed near mineral springs, impregnated with the substances contained in such springs, and consequently possessing similar properties. The *Boues*, *Mud baths*, *Bal'nea Ceno'sa*, are applied generally, and topically, in France, at the springs of St. Amand, Bagnères de Luchon, Bagnols, and Barèges; in the United States, at the White Sulphur, in Virginia, &c.

BOUES MINÉRALES, *Boues des eaux*.

BOUFFE, (F.) The small eminence, formed by the junction of the two lips.

BOUFFE, Eructation.

BOUFFÉE DE CHALEUR, Flush.

BOUFFISSURE, (from *bouffer*, 'to puff up.') Puffiness.

BOUGIE, ([F.] 'a wax candle.') *Candel'ula*, *Cande'la*, *C. cærea seu medica'ta*, *Cæ'reum medica'tum*, *Cere'olus Chirurgo'rum*, *Dæ'dion*, *Specil'lum cæ'reum*, *Vir'ga cæ'rea*, *Cere'olus*. A flexible cylinder, variable in size, to be introduced into the urethra, œsophagus, rectum, &c., for the purpose of dilating these canals, when contracted. A *Simple Bougie* is composed of solid and insoluble substances; as plaster, elastic gum, gutta percha, catgut (*Bougie de Corde à boyau*), &c. It acts of course only mechanically.

BOUGIE, MEDICATED, (F.) *B. Médicamenteuse*, has the addition of some escharotic or other substance to destroy the obstacle; as in the *Caustic Bougie*, (F.) *B. cautérisante*, *B. armée*, *B. emplastique*, which has a small portion of *Lunar Caustic* or *Common Caustic* inserted in its extremity. Ducaup has recommended a *Bougie*, which swells out near its extremity, for the better dilating of the urethra. This he calls *B. à ventre*. The *metallic Bougie*, invented by Smyth, is a composition of metal, allowing of great flexibility; and a *hol-low Bougie* is one with a channel running through it, to be used in the same manner as the catheter, or otherwise.

BOUGIE, UTERINE, Sound, uterine.

BOUGRERIE, Buggery.

BOUILLIE, (F.) *Putic'ula*, *Pap*, (from (L.) *bullire*, (F.) *bouillir*, 'to boil.') Flour, beaten and boiled with milk. It is a common food for infants.

BOUILLON, ([F.] *bouillir*, 'to boil,') *Jus*, *Sorbitt'io*. A liquid food, made by boiling the flesh of animals in water. The osmazome, gelatin, and soluble salts dissolve; the fat melts, and the albumen coagulates. Bouillon is nourishing, owing to the gelatin and osmazome. The *Jus de Viande* is a very concentrated Bouillon, prepared of beef, mutton, veal, &c. See Broth.

BOUILLON, in common language, in France, means a round fleshy excrescence, sometimes seen in the centre of a venereal ulcer.

BOUILLON BLANC, *Verbascum nigrum*.

BOUILLONS MÉDICINAUX ou **PHARMACEUTIQUES**, *Medic'inal* or *Pharmaceutic* *Bouillons*, contain infusions or decoctions of me-

dicinal herbs. The *bouillon aux herbes* is generally composed of *sorrel* or *beet*.

BOUILLON D'OS, (F.) *Bouillon from bones*, is obtained by treating bones with muriatic acid, in order to dissolve the earthy parts. The gelatin, which remains, is then boiled with a little meat and vegetables.—*D'Arcet*. Bouillon, however, can be easily obtained from the bones of roast meat by simple coction.

BOUILLONNEMENT, Ebullition.

BOUIS, Buxus.

BOULE D'ACIER, ('a ball of steel.') *Ferrum tartarizatum*—*b. de Mars*, see *Ferrum tartarizatum*—*b. de Molsheim*, *Ferrum tartarizatum*—*b. de Nancy*, see *Ferrum tartarizatum*.

BOULEAU COMMUN, *Betula alba*.

BOULESIS, (βουλοια, 'to will.') Voluntas.

BOULIMIA, *Bulim'ia*, *Bulim'ius*, *Bul'i'mus*, *Boul'i'mos*, *Bulim'i'asis*, *Bolis'mos*, *Eclim'ia*, *Fa'mes cani'na* seu *bovi'na* seu *lupi'na*, *Appeti'tus cani'nus*, *Appeten'tia cani'na*, *Adepha'gia*, *Cynorez'ia*, *Orex'ia cyno'des*, *Bupi'na*, *Bupe'i'na*, *Phage'na*, *Phagedæ'na*, (βουλα, from *bous*, 'an ox,' and *λωσ*, 'hunger;' or from *bow*, augmentative particle, and *λωσ*, 'hunger,') (F.) *Boulimie*, *Faim canine*, *F. dévorante*, *Polyphagie*. An almost insatiable hunger. A canine appetite. It is sometimes seen in hysteria and pregnancy; rarely under other circumstances.

BOULIMIE, *Boulimia*.

BOULOGNE, MIN'ERAL WA'TERS OF. Boulogne is in the department of Pas-de-Calais, France. The waters are chalybeate.

BOUNCING BET, *Saponaria*.

BOUND, *Costive*.

BOUQUET ANATOMIQUE DE RIOLAN, ([F.] *bouquet*, a collection of flowers or other substances tied together.) A name given, by some anatomists, to the collection of ligaments and muscles, (F.) *Muscles styliens*, inserted into the styloid process of the temporal bone.

BOUQUET FEVER, Dengue.

BOURBILLON, (from *bourbe*, 'mud, dirt.') See *Furunculus* (core).

BOURBON-LANCY, MIN'ERAL WA'TERS OF. Bourbon-Lancy is a small village in the department of Saône-et-Loire, France; where there are thermal saline springs, containing carbonic acid, chloride of sodium, and sulphate of soda, chloride of calcium, carbonate of lime, iron, and silica. Their heat is from 106° to 135° Fahrenheit. Season, from the 15th of May to the 1st of October.

BOURBON L'ARCHAMBAUT, MIN'ERAL WA'TERS OF. This town is in the department of Allier, six leagues west from Moulins, and has been long celebrated for its thermal chalybeate waters. They contain sulphohydric acid, sulphate of soda, magnesia, and lime, carbonate of iron, and silica. Their temperature varies between 136° and 145° Fahrenheit. Season, from the 15th of May to the 1st of October.

BOURBONNE-LES-BAINS, MIN'ERAL WA'TERS OF. These springs are seven leagues from Langres, department of Haute-Marne, France. They are thermal and saline, and have been long celebrated. Temperature from 106° to 133° Fahrenheit. The *Fæctitious water*, (F.) *Eau de Bourbonne-les-Bains*, *Agua Borvonen'sis*, is composed of water, containing twice its bulk of carbonic acid, f $\frac{3}{4}$ xxss; chloride of sodium, $\mathfrak{z}\mathfrak{j}$, chloride of calcium, gr. x.

BOURBOULE, MIN'ERAL WA'TERS OF. A village near Mont d'Or, where there are two thermal saline springs. Season, from the 1st of June to the 15th of October.

BOURDAINE, *Rhamnus frangula*.

BOURDONNEMENT, (*bourdon*, 'a male bee,') *Tinnitus aurium*.

BOURDONNET, *Pulvil'lus*, *P. e linamen'tis*

confertus seu rotundus, Dos'sil. A term in French surgery for charpie rolled into a small mass of an olive shape, which is used for plugging wounds, absorbing the discharge, and preventing the union of their edges. In cases of deep and penetrating wounds, as of the abdomen or chest, a thread is attached to them by which they may be readily withdrawn, and be prevented from passing altogether into those cavities.

BOURGÈNE, *Rhamnus frangula.*

BOURGEON, ('a bud.') Granulation, Papula—*b. Charnu*, Granulation.

BOURGEONNEMENT, see Generation.

BOURGEONS, (pl. of *bourgeon*), Gutta rosea.

BOURGUÉPINE, *Rhamnus.*

BOURRACHE, *Borago officinalis.*

BOURRELET (F.), *A Pad*, a *Bor'der*. A fibro-cartilaginous border, which surrounds certain articular cavities, such as the glenoid cavity of the scapula and the acetabulum; by which the depth of those cavities is augmented. See *Corpus Callosum*.

BOURRELET ROULÉ, *Cornu ammonis.*

BOURSE à **BERGER**, ('shepherd's purse,') *Thalassip bursa*—*b. à Pasteur*, *Thalassip bursa*—*b. Prérotulienne*, see *Knee*, housemaid's.

BOURSES (LES), *Serotum*—*b. Mucilagineuses*, *Bursæ mucosæ*—*b. Muqueuses*, *Bursæ mucosæ*.

BOURSOUFLURE, Puffiness.

BOUTREE, ('boretree,') *Sambucus.*

BOUSSOLE, *Glossanthrax.*

BOUT DE SEIN ([F.], 'end of the breast, nipple,') An instrument of caoutchouc or ivory to place over the nipple when the infant sucks, soon after delivery, or where the nipple is in a morbid condition.

BOUTON ([F.], 'a bud,') *Papula*. Used, however, vaguely for *papula*, *vesicula*, and *pustula*.

BOUTON D'ALEP. A cutaneous affection in the Levant, to which children are liable. It is characterized by one or more tubercles on the face or limbs, which soften, and become covered with a scar. The ulcerations formed are long in healing, and, generally, leave a scar. It is usually left to itself; but is said to be benefited by crucial incisions into the tubercle prior to the period of softening. The disease appears to be anthracoid in its character.

The *Bouton de Biskara* ou *des Zibans*, *Chancre de Sahara*, *Frina* or *Hhabb* of the Arabs, appears to have numerous analogies with the *Bouton d'Alep*.

BOUTON DE BISKARA, see *Bouton d'Alep*.

BOUTON MALIN, see *Anthrax*—*b. d'Or*, *Ranunculus acris*—*b. des Zibans*, see *Bouton d'Alep*.

BOUTONNIÈRE (F.), ('a button hole,') *Fistula*, *Incisio*. A long incision made into the urethra to extract a calculus from the canal, when it is too large to be discharged.

Also, a small incision or puncture, made in the peritoneum, or above the pubis, to penetrate the bladder in certain cases of retention of urine.

BOVACHEVO, *Datura sanguinea.*

BOVILLÉ, *Rubeola.*

BOVISTA, *Lycoperdon.*

BOWEL, *Intestine.*

BOWELS, CONSTIPATED, see *Constipation*—*b. Looseness of*, see *Diarrhoea*.

BOWELS, OPEN, or REGULAR. The state of the intestinal canal when the evacuations are normal in number.

BOWELS, RELAXED, see *Diarrhoea*.

BOWLEGGED, see *Cnemosciosis*.

BOWMAN'S CAPSULE, see *Kidney*.

BOWMAN'S ROOT, *Euphorbia corollata*, *Gil-*

lenia trifoliata, *Leptandria purpurea*

BOXBERRY, *Gaultheria.*

BOX, MOUNTAIN, *Arbutus uva ursi.*

BOX TREE, *Buxus*, *Cornus Florida.*

BOXWOOD, *Cornus Florida.*

BOY, *Puer*, *Païs*, (*païs*, gen. *païses*), *Puiss*; (F.) *Garçon*; of uncertain derivation. A male child under the age of puberty.

BOYAU, *Intestine*—*b. Pollinique*, see *Pollen*.

BRABYION, *Prunum Damascenum.*

BRACHERIOLUM, *Truss.*

BRACHERIUM, *Truss.*

BRACHIA COPULATIVA, see *Peduncles of the Cerebellum*—*b. Pontis*, see *Peduncles of the Cerebellum*.

BRACHLÆUS, *Brachial*—*b. Internus*, *Brachialis anterior*.

BRA'CHIAL, *Brachia'lis*, *Brachii'us*, (from *brachium*.) What belongs to the arm.

BRACHIAL APONEUROSIS. An *aponeurosis*, formed particularly by expansions of the tendons of the *latissimus dorsi*, *pectoralis major*, and *deltoides muclei*, and which completely envelops the muscles of the arm.

BRACHIAL ARTERY, *Art'ria brachia'lis*, *Humeral Ar'tery*, (F.) *Artère ou Tronc brachial*. The artery, which extends from the axilla to the bend of the elbow; where it divides into *A. cubitalis* and *A. radialis*. It passes along the internal edge of the biceps, behind the median nerve and between the accompanying veins. Under the name *Brachial Artery*, *Chaussier* includes the subclavian, axillary, and humeral, the last being the *brachial proper*.

BRACHIAL MUSCLE ANTE'RIOR, *Musculus Brachia'lis Anterior*, *Brachia'lis inter'nus seu anti'cus*, *Brachii'us*, *B. internus*, (F.) *Muscle brachial interne*, *Huméro-cubital*—(Ch.) This muscle is situate at the anterior and inferior part of the arm, and before the elbow-joint. It arises, fleshy, from the middle of the os humeri, and is inserted into the coronoid process of the ulna. *Use*, To bend the fore-arm.

BRACHIAL PLEX'US, *Plexus Brachia'lis*, is a nervous plexus, formed by the interlacing of the anterior branches of the last four cervical pairs and the first dorsal. It is deeply seated in the hollow of the axilla, and extends as far as the inferior and lateral part of the neck. It gives off the *thoracic nerves*, *supra* and *infra scapular*, and the *brachial* (which are six in number), the *axillary*, *cutaneous*, *musculo-cutaneous*, *radial*, *cubital*, and *median*.

BRACHIAL VEINS are two in number, and accompany the artery, frequently anastomosing with each other: they terminate in the axillary. Under the term *Brachial Veins*, *Chaussier* includes the humeral, axillary, and subclavian.

BRACHIALE, ('a bracelet,') from (L.) *brachium*.) *Carpus*.

BRACHIAL'GIA, *Neural'gia Brachia'lis*, (*brachio*, and *algos*, 'pain,') Pain in the arm, neuralgia of the arm.

BRACHIALIS, *Brachial*—*b. Anticus*, *Brachial musculo*—*b. Externus*, see *Triceps extensor cubiti*—*b. Internus*, *Brachial musculo*.

BRACHERIUM, *Truss.*

BRACHILE, *Truss.*

BRACHILUVIUM, (*brachio* and *luere*, 'to wash,') Bath, arm.

BRACHIO, (*βραχιον*, 'arm,') in composition, *brachium* or arm.

BRACHIOCEPHALIC ARTERY, *Innominate arteria*—*b. Veins*, *Innominate vene*.

BRA'CHIOCU'BITAL, *Brachiocubita'lis*. That which belongs both to the arm and cubitus. This name has been given to the internal lateral ligament of the elbow-joint; because it is attached to the os brachii or os humeri, and to the cubitus or ulna.

BRACHIOCYLLO'SIS, (*brachio*, and *κυλλωσις*, 'the act of making crooked,') Curvature of the

arm inwards. Paralysis or loss of power from curvature of the arm.

BRACHION. (βραχιων,) Brachium.

BRACHION'CUS, (*brachio*, and *oikos*, 'a swelling,') A tumour of the arm.

BR'ACHIORA'DIAL, *Brachioradia'lis*. That which belongs to the brachium and radius. This name has been applied to the external lateral ligament of the elbow-joint, because it is attached to the humerus and to the radius. See *Supinator radii longus*.

BRACHIORRHEU'MA, gen. *Brachiorrheu'matis*, *Rheumatis'mus bra'chii*, (*brachio*, and *ρῆμα*, 'diffusion, rheumatism,') Rheumatism of the arm.

BRACHIOT'OMY, *Brachiotom'ia*, (*brachio*, and *τομή*, 'incision,') Amputation of the arm.

BRACHIROLUM, Truss.

BR'ACHIUM, (βραχιον, 'the arm,') *Bra'chion*, *Lacer'tus*, (F.) *Bras*. The arm from the shoulder to the wrist, or the part between the shoulder and elbow. See *Humeri Os*.

BRACHIUM ANTE'RIOUS. A rounded process, which passes from the anterior pair of the corpora quadrigemina (*nates*) obliquely outwards into the thalamus opticus.

BRACHIUM MOVENS QUARTUS, *Latissimus dorsi*.

BRACHIUM POSTE'RIOUS. A rounded process, which passes from the posterior pair of the quadrigemina (*testes*) obliquely outwards into the optic thalamus.

BRACHUNA, Nymphomania, Satyriasis.

BRACHY, (from βραχυς, 'short,') in composition, short.

BRACHYAU'CHEN, (*brachy*, and *αυχνη*, 'neck,') One who has a short neck.

BRACHYCEPH'ALÆ (*Gentes*), 'short heads,' (*brachy*, and *κεφαλη*, 'head,') In the classification of Retzius, those nations of men whose cerebral lobes do not completely cover the cerebellum—as the Selavonians, Fins, Persians, Turks, Tartars, &c.

BRACHYCEPH'ALIC, *Brachycephal'icus*. Having a short head.

BRACHYCHRON'US, (*brachy*, and *χρονος*, 'time,') That which continues but a short time. A term applied to diseases which are of short duration.

BRACHYGNATHUS or *Brachygnathus*, (*brachy*, and *γναθος*, 'the under jaw,') A monster with too short an under jaw.

BRACHYNSIS, (βραχυνω, 'to shorten,') Abbreviation.

BRACHYPNEA, (*brachy*, and *πνεω*, 'to breathe,') Dyspnoea.

BRACHYP'OTI, (*brachy*, and *ποτης*, 'drinker,') They who drink little, or who drink rarely.

BRACHYRHYN'CHUS, (*brachy*, and *ρυγχος*, 'snout,') A monster with too short a nose.

BRACHYSMUS, (from βραχυς, 'short,') Abbreviation.

BRACING, Corroborant.

BRACKEN, (Sax. *brecan*, 'to break,') on account of the broken character of the leaves.) *Pteris aquilina*—b. Rock, Polypodium, P. incanum.

BRACKEN ROOT, Polypodium vulgare.

BRADY, (from βραδύς, 'difficult, 'tardy,') in composition, the same.

BRADYÆSTHES'IA, (*brady*, and *αἴσθησις*, 'sensation,') Impaired sensation.

BRADYBOLISMUS, (*brady*, and *βαλλω*, 'to throw,') Bradyspermatismus.

BRADYECOIA, (*brady*, and *ακον*, 'hearing,') Deafness.

BRADYFIBRIN, see *Corium phlogisticum*.

BRADYCYROTE, *Bradycyrotus*; (from *brady*, and *κροταιν*, 'to strike,') An agent, that diminishes the number of pulsations of the heart,—as *veratrum viride*.

BRADYLOG'IA, *Dysla'lia*; (*brady*, and *λογος*, 'a discourse,') Difficulty of speech.

BRADYMASE'SIS, *Bradymasse'sis*, improperly *Bradymate'sis*, *Manduca'tio diffi'cilis*, (*brady*, and *μασσεις*, 'mastication,') Difficult mastication. See *Dysmasesis*.

BRADYMASTESIS, *Bradymasesis*.

BRADYPEP'SIA, (βραδυπεψια, from *brady*, and *πεπω*, 'I digest,') *Tarda cibo'rum concoct'io*. Slow digestion. See *Dyspepsia*.

BRADYSPERMATIS'MUS, *Bradybolis'mus*, *Ejacula'tio sem'inis impedi'ta*, *Dyspermatis'mus*, (*brady*, and *σπερμα*, 'sperm,') A slow emission of sperm.

BRADYSU'RIA, *Tenes'mus vesic'æ*, (F.) *Ténésie vésical*, (*brady*, and *ουρειν*, 'to pass the urine,') Painful evacuation of the urine, with perpetual desire to void it. *Dysuria*.

BRADYTOCIA, (*brady*, and *τοκος*, 'birth,') *Dys-tocia*.

BRAG'GET, *Brag'gart*, *Brag'wort*. A name formerly applied to a tisan of honey and water, see *Hydromeli*.

BRAI LIQUIDE, see *Pinus sylvestris*—b. *Sec*, *Colophonia*.

BRADISM, see *Magnetism*, animal.

BRAIN, (Sax. *bræcen*,) *Cerebrum*—b. *Fag*, see *Nervous diathesis*.

BRAIN, LITTLE, *Cerebellum*.

BRAIN PAN, *Cranium*—b. *Sand*, see *Pineal gland*.

BRAINE, MINERAL WATERS OF. *Braine* is a small village, three leagues from Soissons, France, which has purgative waters similar to those of Passy.

BRAIRÈTE, *Primula veris*.

BRAKE, COMMON, (same etymon as *Bracken*,) *Pteris Aquilina*—b. *Rock*, Polypodium incanum, Polypodium vulgare.

BRAKE ROOT, Polypodium vulgare.

BRACING, Vomiting, Vomiturition.

BRAMBLE, AMERICAN HAIRY, see *Rubus fruticosus*—b. *Common*, *Rubus fruticosus*.

BRAMBLE BERRIES, see *Rubus fruticosus*.

BRAN, (F.) *Bran*, *Furfur*.

BRANCURSINE BÂTARDE, *Heracleum spondylium*.

BRANCA GERMANICA, ([F.] *branca*, 'a paw,') *Heracleum spondylium*—b. *Ursina*, *Acanthus mollis*—b. *Vera*, *Acanthus mollis*.

BRANCH, ([F.] *Branche*, originally, probably, from βραχυων, 'an arm,') [?] because branches of trees, &c., go off like arms.) A term applied, generally, to the principal division of an artery or nerve. The word is commonly used synonymously with *Ramus*; but often, with the French, *Branche* signifies the great division;—*Rameau*, Lat. *Ramus*, the division of the branches; and *Ramuscules*, (L.) *Ramusculi*, the divisions of these last.

The French, also, speak of the branches of the pubis for the *Rami* of that bone, *branches* of the ischium for the *rami* of the ischium, &c.

BRANCHES DE LA MOËLLE ALLONGÉE (PETITES), *Corpora restiformia*.

BRAN'CHI, *Bran'che*. Swellings of the tonsils, or parotid, according to some;—of the thyroid gland, according to others.

BRAN'CHIA, (Gr.) βραγχια. The gills or respiratory organs of fishes, corresponding to the lungs of terrestrial animals.

BRAN'CHUS, (βραγχος,) *Rauce'do*. A catarrhal affection of the mucous membrane of the fauces, trachea, &c. Hoarseness.

BRANCI, *Cynanche tonsillar*.

BRANCIA, *Vitrum*.

BRANDWINE, Brandy.

BRAN'DY, ([G.] *Branntwein*, Dutch, *Brandwijn*, 'burnt wine,') (Old Eng.) *Brandwine*. *Vinum adustum* *vin crematum*, *Aqua Vitis*, (F.) *Spiritus vini Gall'ici* (Ph. U. S.), *Spiritus Gall'icus*, *Enu de vie*, (S.) *Aguardiente*. The first liquid product obtained by distilling wine. It is composed of

water, alcohol, and an aromatic oily matter, which gives it its flavour, and ought to contain from 48 to 56 per cent. of absolute alcohol. Brandy is a powerful and diffusible stimulant, and as such is used in medicine. It has been also called *Liquor Aquilegii*. See Spirit.

BRANDY, APPLE, see *Pyrus malus*—b. Egg, see Oyum.

BRANDY BOTTLES, see *Nymphæa lutea*—b. Face, Gutta rosea.

BRANDYWINE, MINERAL WATERS OF. These are in Delaware, a few miles from Wilmington, and have been much frequented. They are chalybeate.

BRANKS, (from *branchi* [?]) *Cynanche parotidæa*.

BRANKURSINE, *Acanthus mollis*.

BRANNTWEIN, ([G.] 'burnt wine,') Brandy.

BRAS. See *Oryza*.

BRAS, ([F.] 'arm,') *Brachium*—b. *du Cervelet*, *Corpora testiformia*.

BRASDOR'S OPERATION FOR ANEURISM. An operation by ligature, proposed by Brasdor, which consists in the application of the ligature on the distal side of the tumour.

BRASÉUR, MINERAL WATERS OF. *Braséur* is a place in the diocese of Rhodéz, where there are cathartic waters.

BRASENIA, *B. Hydropeltis*.

BRASENIA HYDROPELTIS, *Brasenîa*, *B. Peltata*, *Hydropeltis purpurea*, *Gelatinâ aquatica*, *Frog's leaf*, *Little Water Lily*, *Water Jelly*, *Water shield*, *Water target*, *Deer's food*. An indigenous plant, *Nat. Ord.* Ranunculaceæ, *Sex Syst.* Polyandria Polygynia, flourishing from Kentucky to Carolina and Florida; and covering the surface of ponds, marshes, &c. The fresh leaves are mucilaginous, and have been used in pulmonary complaints, dysentery, &c., like *Cetraria*.

BRASENIA PELTATA, *B. Hydropeltis*.

BRASH, (Sc.), (like (E.) *brush*.) A transient attack of sickness. In Scotland, a bodily indisposition. A rash or eruption, (Prov. West of England.

BRASH, WATER, Pyrosis.

BRASH, WEAN'ING, (Sc.) *Spa'ning brash*, *Atrophîa Ablactatorum*. A severe form of diarrhœa, which supervenes at times on weaning. The *Maladie de Cruceilhier* appears to be a similar affection.

BRASILETTO, see *Cæsalpinia*.

BRASIUM, Malt.

BRASMOS, (*βραμωσ*, 'to agitate,') Fermentation.

BRASS, (Sax. *brar*, Welsh, *près*.) *Aurichalcum*, *Orichalcum*, *Chrysochalcum*, *Chalcum*, *Escavum*, (F.) *Airain*, *Cuivre jaune*. A yellow metal, formed by mixing copper with calamine. The same general remarks apply to it as to copper.

BRASSFOUNDERS' AGUE, see *Ague*.

BRASSICA, *Crambê*, *B. oleracea* seu *capitata* seu *cuma'na* of the old Romans. The *Cabbage*, (F.) *Chou potager*. *Order*, Cruciferae. *Sex Syst.* Tetradynamia Siliquosa. Cato wrote a book on its virtues. It is a vegetable by no means easy of digestion when boiled; when raw, it appears to be more digestible. When forming a solid globular mass, like a head, it is the *B. Capitata*, (F.) *Chou-Cabus*, *Chou Pommé*.

BRASSICA CANINA, *Mercurialis perennis*—b. *Capitata*, *Brassica*—b. *Cumana*, *Brassica*.

BRASSICA ERU'CA, *B. hispida*, *Eru'ca*, *E. foetida* seu *sati'va*, *Sinapis eru'ca*, *Sisymbrium erucas'trum*, *Gar'den Rock'et*, *Ro'man Rock'et*, *Shin'lock*, &c., (F.) *Chou Roquette*, *Roquette*. This was considered by the Romans an aphrodisiac. The seeds were ordinarily used.

BRASSICA FLORIDA, *Brassica Pompeiana* of the ancients—the *Caul'flower*, *Caul'is Flor'ida*, (F.) *Chou-fleur*, is a more tender and digestible variety.

The *Broccoli*, *B. Sabellica* of the Romans, *B. Italica*, belongs to this variety.

BRASSICA HISPIDA, *B. eruca*—b. *Italica*, *B. Florida*—b. *Marina*, *Convolvulus soldanella*.

BRASSICA NA'PUS, *Na'pus Sylvestris*, *Bu'nias*, *Rape*, (F.) *Navette*. The seed yield, on expression, a quantity of oil, which, at times, forms part of liniments.

BRASSICA NIGRA, *Sinapis nigra*—b. *Oblonga*, *B. rapa*—b. *Oleracea*, *Brassica*—b. *Pompeiana*, *B. Florida*.

BRASSICA RA'PA, *Ra'pa rotunda* seu *oblonga* seu *na'pus*, *Ra'pum na'pus*, *Sinapis tuberosa*, *Tur'nip*, (F.) *Chou navet*, *Navet*, *Rave*. The turnip is liable to the same objection (but to a less extent) as the cabbage.

BRASSICA SABELLICA, *B. Florida*.

BRATHU, (*βραθυ*,) *Juniperus sabina*.

BRAWLINS, *Arbutus uva ursi*, *Vaccinium vitis idæa*.

BRAWN, *Musele*.

BRAYER, *Truss*.

BRAYERA ANTHELMINTICA, (after Brayer, a German physician,) *Hagenia Abyssinica*.

BRAZIL WOOD, *Cæsalpinia echinata*.

BREAD, ([Sax.] *breod*,) see *Triticum*.

BREAD, GLU'TEN. Bread made of wheat dough deprived of the chief portion of its starch by washing. Bread made of gluten only cannot be eaten, on account of its hardness and toughness; hence one-fifth of the normal quantity of starch is allowed to remain, and in this form the bread is said to be tolerably light, eatable, and moderately agreeable.

BREAD, HOUSEHOLD, *Syncomistos*—b. *Indian*, *Lycoperdon tuber*—b. *St. John's*, *Ceratonia siliqua*.

BREADBERRY, *Pap*.

BREADFRUIT TREE, *Artocarpus*.

BREADROOT, *Camassia esculenta*, *Psoralea esculenta*.

BREAKEBONE FEVER, *Dengue*.

BREAST, ([Sax.] *breofc*,) *Thorax*, *Mamma*—b. Abscess of the, *Mastodynia apostematosa*.

BREAST, BROKEN, see *Mastodynia apostematosa*.

BREASTBONE, *Sternum*.

BREASTGLASS, *Milkglass*. A glass applied to the nipple to receive the milk when secreted copiously by the mamma.

BREAST, IRRITABLE, *Neuralgia mammæ*.

BREASTPANG, *SUFFOCATIVE*, *Angina pectoris*.

BREAST, PIGEON, see *Lordosis*.

BREASTPUMP, *Antlia Lactea*.

BREASTWEED, *Saururus cernuus*.

BREATH, (Sax. *b-ase*,) *Halitus*, *Anhelitus*, *An'imus*, *Spiritus*, *Al'mos*, *Pne'ca*, *Pno'e*, *Pno'us*, (Old Eng.) *Aande*, and *Ande*, *Wind*, (F.) *Haleine*. The air expelled from the chest at each expiration. It requires to be studied in the diagnosis of thoracic diseases especially. See *Respiration*.

BREATH, OFFENSIVE, *Fæ'tor O'ris*, *Catostomatophre'sia*, *Hal'itus o'ris foetidus*, *O'zæ*, *Ozostomia*. An offensive condition, which is usually dependent upon carious teeth, or some faulty state of the secretions of the air passages. The internal use of the chlorides may be advantageous.

BREATH, PULSE. An audible pulsation given to the breath, as it issues from the mouth, by each beat of the heart, occasionally heard, according to Dr. Radclyffe Hall, in dry tuberculous cavities with dense walls not separated from the heart by permeable lung tissue.

BREATH, SATURNINE, see *Saturnine*—b. *Short*, *Dyspnœa*.

BREATHING AIR, see *Respiration*.

BREATHING, DIFFICULTY OF, *Dyspnœa*.

BREATHING A VEIN, see *Bloodletting*.

BRECHET, (F.) *Brisket*, ([?] 'breast-cut,')

This name is given in some parts of France to the *cartilago exsiformis*, and sometimes to the sternum itself.

BRECHMA, (βρεχμα,) gen. *Brech'matis*, Bregma.

BRECHMUS, (βρεχμος,) Bregma.

BREDISSURE (F.), *Trismus Capistratus*. Incapacity of opening the mouth, in consequence of preternatural adhesion between the internal part of the cheek and gums; often occasioned by the abuse of mercury.

BREDOUILLEMENT (F.), *Titubantia*. A precipitate and indistinct mode of utterance, in which a part of the words is pronounced, and several of the syllables are viciously changed. This defect is analogous to stuttering, but differs from it in being dependent on too great rapidity of speech; while stuttering is characterized by continual hesitation, and frequent repetition of the same syllables.

BREE (Sc.), (*breno*.) Supercilium.

BREED, (from (Sax.) *bŕeagan*, 'to nourish.') Race.

BREEDING, Generation, Pregnant.

BREED'ING, CROSS. The act of raising or breeding from different stocks or families.

BREEDING-IN-AND-IN. The act of raising or breeding from the same stock or family.

BREELLS, Spectacles.

BREG'MA, gen. *Breg'matis*, *Brech'ma*, *Brech'mus*, (βρεχμ, 'to moisten') *Fontanel'la*, *Sin'ci-pu't*. The top of the head was thus called, because it was believed to be humid in infants; and, according to some, because it was conceived to correspond to the most humid part of the brain.

BREGMATODYMIA, (*bregma*, and *δυα*, 'I enter into,') see Cephalodymia.

BRENNING, (Teut. *brennen*, 'to burn.') Burning.

BREPHOCTONON, (βρεφοκτονον, from *βρεφος*, 'a babe,' and *κτεivo*, 'to kill,') *Conyza squarrosa*.

BREPHOTROPHE'UM, (βρεφοτροφειον, from *βρεφος*, 'a new-born child,' and *τροφειν*, 'to nourish,') *Ethetobrephotrophe'um*, *Ethetotrophe'um*. A founding hospital.

BRESCHE'S BONECANALS, see Diploic—b's Veins, see Diploic.

BRÉSILLET, *Cæsalpina sappan*.

BRE'VIA VA'SA, 'Short Ves'sels.' (F.) *Vaisseaux courts*. This name has been given to several branches of the splenic arteries and veins, which are distributed to the great *cul-de-sac* of the stomach.

BREVIS CUBITI, ('short (muscle) of the elbow,') see *Anconeus*.

BRICK, (F.) *Brique*. Hot bricks are sometimes used to apply heat to a part, as to the abdomen in colic, or after the operation for popliteal aneurism; or, reduced to very fine powder, and mixed with fat, as an application to herpetic and psoric affections.

BRICKS, *Fornacese Tes'tæ* or *Tiles* were formerly bruised in vinegar, and the liquid was used as a specific in cutaneous affections. They entered, also, into a cerate used for herpetic and other eruptions, &c. To the *Terra Fornacum*, or *Brick earth*, the same virtues were assigned.

BRICKLIKE SEDIMENT, see *Lateritious*.

BRICUMUM, *Artemisia*.

BRIDE (F.), 'a bridle.' *Fræ'nulum*, *Retinaculum*. This term is given, in the plural, to membranous filaments, which are found within abscesses or deep-seated wounds, and which prevent the exit of pus. The term is, also, applied to preternatural adhesions, which occur in cicatrices of the skin, in the urethra, or in inflamed serous or synovial membranes.

BRIEF, (a corruption of) Rife.

BRIER, WILD, ([Sax.] *bŕæp*), *Rosa Canina*.

BRIGHT'S DISEASE OF THE KIDNEY. see *Kidney*, *Bright's disease* of the.

BRIGHT'ON, CLIMATE OF. The air of this fashionable watering-place, on the south coast of England, is dry, elastic, and bracing. Its climate appears to the greatest advantage in the autumn and early part of the winter; when it is somewhat milder and more steady than that of Hastings. Accordingly, it is adapted for all cases in which a dry and mild air at this season of the year proves beneficial. In the spring months, owing to the prevalence of, and its exposure to, north-east winds, the climate is cold, harsh, and exciting to the delicate. It is well adapted for convalescents, and for all who require a dry and bracing sea air.

BRIMSTONE, (*brin* or *bren stone*, that is, 'fiery or burning stone' [?],) Sulphur.

BRINE, ([Sax.] *bŕyne*.) Muria.

BRINTON ROOT, *Leptandria purpurea*.

BRINVILLIERS, *Spigelia anthelmia*.

BRION, *Corallina*.

BRIQUE, Brick.

BRIQUEBEC, MINERAL WATERS OF. This town is three leagues from Cherbourg, in France. The water contains chloride of iron.

BRQUETÉ, (*from brique*.) *Lateritious*.

BRISÉCOQUE (F.), (*briser*, 'to break to pieces,' and *coque*, 'a shell.') An instrument designed by Heurteloup for breaking to pieces the shell of a vesical calculus, after it has been hollowed by his *mandrin à virgule*.

BRISÉPIERRE ARTICULÉ (F.), *briser*, 'to break to pieces,' and *pierré*, 'a stone.' An instrument invented by Jacobson for crushing the stone in the bladder.

BRIS'TOL HOT-WELL, *Bristolien'sis A'qua*. Bristol is about thirteen miles from Bath, in England. The water is an almost pure thermal; slightly acidulated. It contains chlorides of magnesium and sodium, sulphate of soda, sulphate of lime, carbonate of lime, carbonic acid, oxygen and azote. Temperature, 74° Fah. The *Hot Well* has been long celebrated. Its action is like that of thermal waters in general. The climate of Bristol is mild, and hence the water has been celebrated for the cure of incipient pulmonary consumption. See *Clifton*.

BRIZOCERAS, Ergot.

BROAD, (Sax. *bræd*.) *Latus*, (F.) *Large*. Any body is so termed whose transverse extent is considerable compared with its length. The *Broad Bones*, as the frontal, parietal, occipital, and iliac, aid in forming the parietes of splanchnic cavities. *Broad Muscles* generally occupy the parietes of cavities, and especially those of the chest and abdomen. The epithet has also been applied to other parts—as to the *broad ligaments* of the womb, &c.

BROCCOLI, *Brassica sabellica*.

BRO'CHOS, (βροχος,) *La'queus*. A bandage.

BROCHT, Vomiting.

BROCH'THUS, (βροχθος,) *Gu'la*. The throat, also a kind of small drinking vessel.

BRO'CHUS, (βροχος.) This name has been given to one who has a very prominent upper lip. According to others, it means one whose teeth project in front of the mouth.

BRO'DIUM. A synonym of *Jus* or *Jusculum*. Broth, or the liquor in which any thing is boiled. *Bro'dium salis*—a decoction of salt.

BROIEMENT, (*from broyer*, 'to grind, to pound,') see *Cataract*, *Laceration*—*b. Linéaire*, see *Écraseur*—*b. de la Pierre*, *Lithroty*.

BROKELEAK, *Rumex hydrolapathum*.

BROKEN BREAST, see *Mastodynia apostematosa*—*b. Doses*, see *Doses*, broken—*b. Winded*, *Asthmatic*—*b. Windedness*, *Asthma*.

BROMA, (βρωμα, 'food,') Aliment; (βρωμος, 'a stench,') Bromine.

BROMATOËCCRISIS, (βρωμα, gen. βρωματος, 'food,' and εκκρισις, 'excretion,') Lientery.

BROMATOG'RAPHY, *Bromatograph'ia*, *Bromog'r'aphy*, *Bromograph'ia*, (βρωμα, gen. βρωματος, 'food,' and γραφή, 'a description,') A description of aliments.

BROMATOL'OGY, *Bromatolog'ia*, *Sitiol'ogy*, (βρωμα, gen. βρωματος, 'food,' and λογος, 'a discourse,') A treatise on food.

BROME, Bromine.

BROMEGRASS, *Bromus ciliatus*—b. Soft, *Bromus ciliatus*.

BROMELIA ANA'NAS, (after Olaus Bromel, a botanist of Sweden,) *Car'duus Brazilia'nus*, *Ana'nas ova'ta* seu *aculea'ta*, *Ana's'sa*, *Ca'pa-Isiak'ka*, *Ana'nas* or *Pine Ap'ple*, *Sour'sop*. Order, Bromeliaceæ. A West India tree, which produces the most delicious of fruits.

BROMELIA PIN'GUIN, *Ana'nas America'na*, *Pin'guin*, *Broad-leaved wild Ana'nas*, &c. The West India plant, which affords the *Pinguin* fruit. The fruit is refrigerant, and the juice, when ripe, very austere. It is used to acidulate punch. A wine is made from the *Pinguin*, which is very intoxicating, and has a good flavour.

BRO'MIC, *Bro'micus*; same etymon as Bromine. Containing bromine.

BROMIDE OF IRON, see Bromine—b. of Mercury, see Bromine—b. of Potassium, see Bromine.

BROMIDRO'SIS, (βρωμος, 'stench,' and ιδρως, 'sweat,') Fetid sweat.

BRO'MINE, *Bro'minum* (Ph. U. S., 1842), now *Bromin'ium*, *Bro'ma*, *Bromin'eum*, *Bro'mium*, *Bro'mina*, *Bro'mum*, *Mu'rina*, *Mu'ride*, *Brome*. A simple body, of a very volatile nature, and highly offensive and suffocating odour, whence its name, (βρωμος, 'a stench,') It is met with chiefly in sea-water, and in many animal and vegetable bodies that live therein. It has likewise been found in many mineral waters of this and other countries. In its chemical relations, it may be placed between chlorine and iodine. With oxygen it forms an acid—the *Bro'mic*, and with hydrogen another—the *Hydrobro'mic*. Bromine may be dissolved in forty parts of distilled water, and six drops be commenced with as a dose.

BROMIDE OF IRON, *Ferri Bro'midum*, *Fer'rum bro'matum*, (F.) *Bromure de Fer*; in solution; *Hydrobro'mate of Iron*, *Fer'ri hydrobromas*, *Fer'rum hydrobro'micum oxydatum*, (dose, gr. i. or ij); and **BROMIDE OF POTASSIUM**, *Potas'sii bro'midum*, (F.) *Bromure de potassium*, have been used medicinally, and chiefly in scrofulosis—internally, as well as applied externally. Bromide of iron has also been given in painful erections, and in spermatorrhœa. **BROMIDES OF MERCURY** (*Hydrargyri Bro'mida*, (F.) *Bromures de Mercure*), have been given in syphilis. The *proto-bromide* and the *bi-bromide* are analogous in composition and medicinal properties to the corresponding iodides of mercury. **CHLORIDE OF BROMINE**, *Bromin'ii chlo'ridum*—made by passing chlorine through bromine, and condensing the resulting vapours by cold, has been prescribed internally, as well as externally, in cancer; but chiefly in the latter mode, in the form of a caustic paste, either alone or with other chlorides—as those of zinc, antimony, and gold.

BROMIUM, Bromine.

BROMOGRAPHY, *Bromatography*.

BROMOSUS, (βρωμεω, 'to stink,') Fetid.

BROMUM, Bromine.

BROMURE DE FER, see Bromine—b. de *Mercur*, see Bromine—b. de *Potassium*, see Bromine.

BROMUS, (βρωμος.) One of the cerealia, supposed, by some, to be oats. See *Avena*.

BROMUS CILIA'TUS, (βρωμος,) *B. pur'gans*, *Brome'grass*; indigenous: Order, Gramineæ; is said to be emetic, and anthelmintic, cathartic and diuretic. It purges cattle.

BROMUS GLABER, *Triticum repens*.

BROMUS MOL'IS, *Soft Brome'grass*. The seeds are said to cause giddiness in man, and to be fatal to poultry.

BROMUS PURGANS, *B. ciliatus*—b. *Temulentus*, *Lolium temulentum*.

BRONCHADENES, (*bronchia*, and αδην, 'a gland,') Bronchial glands.

BRONCHARCTIA, (*bronchia*, and αρctαι, 'to contract,') Bronchiostenosis.

BRONCHECTASIS, (*bronchia*, and εκτασις, 'dilatation,') Bronchia, dilatation of the.

BRONCHES, Bronchia—b. *Ganglions lymphat'iques des*, Bronchial glands.

BRONCHI, Bronchia.

BRON'CHIA, pl. of *bronchion*, (βρογchia,) *Bron'chie*, *Bron'chi*. The Latins used the term *Bronchus* for the whole of the trachea; whilst they called its ramifications *Bronchia*. *Bronchia*, *Bronchiæ*, and *Bronchi*, (F.) *Bronches*, now mean the two tubes, with their ramifications, which arise from the bifurcation of the trachea, and carry air into the lungs—*Can'nulæ pulmo'nium*, *Syrin'ges* seu *Canal'es aërif'eri*.

BRONCHIA, DILAT'ION OF THE, *Dila'ted Bron'chia*, *Bronche'ctasis*, *Bronchie'ctasis*, *Dila'tio bronchio'rum*. The physical signs of this condition are the following: *Percussion* usually clear, but not unfrequently less so than natural, although very seldom quite dull. *Auscultation* detects coarse mucous or gurgling rhonchi, increased by the cough, combined with, or replaced by, bronchial or cavernous respiration, which is often effected as if by a sudden puff or whiff. The resonance of the voice is increased, but it seldom amounts to perfect pectoriloquy. The most common situations for dilated bronchia are the scapular, mammary, and lateral regions. They are almost always confined to one side.

BRONCHIA, OBLITER'ATION OR COMPRES'SION OF THE. The inspiratory murmur on auscultation is weaker or wholly suppressed over a limited portion of the chest; the expiration is generally more distinct and prolonged: all the other conditions are natural.

BRONCHIÆ, see Bronchia.

BRON'CHIAL, *Bron'chie*, *Bronchia'lis*, *Bron'chieus*, (F.) *Bronchial*, *Bronchique*. That which relates to the Bronchia.

BRONCHIAL ARTERIES, (F.) *Artères Bron'chiques*. These are generally two in number, one going to each lung. They arise from the thoracic aorta, and accompany the bronchia in all their ramifications.

BRONCHIAL CELLS, see *Cellules*, bronchie, and *Pulmo*.

BRONCHIAL COUGH, (F.) *Toux bronchique*, *T. tubaire*. This generally accompanies bronchial respiration. Both indicate obstruction to the entrance of air into the air-cells.

BRONCHIAL GLANDS, *Broncha'denes*, *Glan'dule Vesalia'næ*, *Glands of Vesa'tius*, (F.) *Glandes bronchiques*, *Ganglions lymphatiques des bronches*, are numerous glands of an ovoid shape; of a reddish hue in the infant, and subsequently brown and black, seated in the course of the bronchia. Their functions are unknown. The bronchial glands may be presumed to be affected by scrofulosis, when, in addition to the existence of tumours in the neck, percussion gives a dull sound under the upper and central part of the sternum, whilst there is no appreciable lesion of the lungs.

BRONCHIAL NERVES, (F.) *Nerfs bronchiques*, are furnished by the two pulmonary plexuses.

BRONCHIAL PHTHISIS, see Phtthisis, bronchial—b. Respiration, see Murmur, respiratory.

BRONCHIAL VEINS arise from the last divisions of the arteries of the same name, and pass, on the right side, into the vena azygos; on the left, into the superior intercostal.

BRONCHIC, Bronchial.

BRONCHIECTASIS, (*bronchia*, and *εκτασις*, 'dilatation.') Bronchia, dilatation of the.

BRONCHITIS, Bronchitis.

BRONCHIO, *Broncho*, (*βρογχος*, 'the throat, the windpipe,') in composition, bronchia or bronchus.

BRONCHIOLE, *Bronchiolum*, *Bronchiolus*; (dim. of *Bronchium* or *Bronchus*.) A minute bronchial tube.

BRONCHIOPNEUMONIA, *Bronchopneumonia*.

BRONCHIOSTENOSIS, *Bronchiarectia*, (*bronchio*, and *στενωσις*, 'contraction.') Contraction or narrowness of the bronchi.

BRONCHIQUE, Bronchial.

BRONCHISMUS, Spasmodic contraction of the bronchia.

BRONCHITE CONVULSIVE, Pertussis—b. *Pseudomembraneuse*, Polypus bronchialis.

BRONCHITIS, *Bronchitis*, (*bronchia* and *itis*.) *Inflammatio bronchio-rum*, *Catar'rhus Pulmonum seu bronchio-rum*, *Pleuritis humida seu bronchia'lis*, *Bronchostasis*, *Angi'na bronchia'lis*, *Pulmonary catarrh*, (F.) *Inflammation des bronches*. Inflammation of the lining membrane of the bronchial tubes. This is always more or less present in cases of pulmonary catarrh; and is accompanied by cough, mucous expectoration, dyspnoea, and more or less uneasiness in breathing. The acute form is accompanied with all the signs of internal inflammation, and requires the employment of antiphlogistics followed by revulsives. The chronic form, *Tussis senilis*, *Catar'rhus senilis*, *Rheum'ata catarrh'ale*, *Peripneumonia notha*, *Bronchorrhoea acuta*, *Winter Cough*, *Chronic Catarrh*, may be confounded with phtthisis; from which it must be distinguished mainly by the absence of hectic fever and of the physical signs that are characteristic of the latter, as well as by the nature of the expectoration, which is generally mucous, although at times mucopurulent. When the expectoration is little or none, the bronchitis is said to be dry; *dry catarrh*, (F.) *Catarrhe Sec*; when fetid, the disease is termed *fetid bronchitis*.

When bronchitis affects the smaller tubes, it is termed *capillary bronchitis*, *bronchitis capillaris*, *bronchoecac' infantilis* [?], and is often fatal to children. *Vesicular bronchitis* is used by MM. Kiliet and Barthez for the *vesicular pneumonia* of children.

BRONCHITIS, Catarrh—b. Asthenica, Peripneumonia notha—b. Capillary, see Bronchitis—b. Chronic, see Bronchitis—b. Convulsiva, Pertussis—b. Crouposa, Polypus bronchialis—b. Exsudativa, Polypus bronchialis—b. Fetid, see Bronchitis.

BRONCHITIS, MECHANICAL. Inflammation of the lining membrane of the air-tubes, induced by the inhalation of irritating particles.

BRONCHITIS MEMBRANACEA, Polypus bronchialis—b. Plastic, Polypus bronchialis—b. Pseudomembranous, Polypus bronchialis—b. Summer, Fever, hay—b. Vesicular, see Bronchitis.

BRONCHIUS, Sterno-thyroides.

BRONCHLEMMITIS, (*broncho*, *λεμμα*, 'a sheath,' and *itis*), Cynanche trachealis, Polypus bronchialis.

BRONCHO, in composition, see Bronchio.

BRONCHOCACE, (*broncho*, and *κακος*, 'faulty,') *Peripneumonia notha*—b. Infantilis, see Bronchitis.

BRONCHOCATARRHUS, Catarrh.

BRONCHOCELE, (*βρογχοκήλη*, from *broncho*, and *κήλη*, 'tumour,') *Bron'chocele*. An inaccurate name for the affection which is called, also, *Bo'chium*, *Bo'tium*, *Bo'eium*, *Her'nia gutturis seu guttura'lis seu bronchia'lis*, *Gut'tur tu'midum seu glo-bo'sum*, *Trachelophy'ma*, *Thyroce'le*, *Thyroce'le*, *Tracheoce'le*, *Tracheoce'le*, *Thyremphrax'is*, *Thyreophrax'is*, *Thyrophrax'is*, *Thyreon'eus*, *Thyron'eus*, *Deiron'eus*, *Deron'eus*, *Gos'eum*, *Go'tium*, *Ecechebron'chus*, *Gongro'na*, *Stru'ma*, *Glans*, *Tu'ber gutturo'sum*, *Gut'teria*, &c., the *Der'byshire neck*, *Swell'ed neck*, *Wen*, *Goitre*, &c., (F.) *Goître*, *Goître*, *Gouître*, *Hypertrophie du Corps Thyroïde*, *Grosse Gorge*, *Gros Cou*, (S.) *Goto*. This is no rupture, but consists of an enlargement of the thyroid gland. It is common at the base of lofty mountains in every part of the world; and has been supposed to be owing to the drinking of snow-water, but it occurs where there is no snow. The tumour is sometimes very extensive. Iodine has great power over it, and will generally occasion its absorption, when the case has not been of such duration as to have ended in a cartilaginous condition. Shakespeare describes the tumors as '*wallets of flesh*.'—Gonzalo, in *Tempest*, Act III. Scene 3.

BRONCHOCEPHALITIS, (*broncho*, *κεφαλη*, 'head,' and *itis*), Pertussis.

BRONCHOPARALYSIS, Asthma.

BRONCHOPHONY, (*broncho*, and *φωνη*, 'voice,') Resonance—b. Pectoriloquous, Pectoriloquy—b. Strong, Pectoriloquy.

BRONCHOPLASTIC, *Bronchoplas'ticus*, (*broncho*, and *πλασσω*, 'I form,') An epithet given to the operation for closing fistule in the trachea; *Bron'choplasty*.

BRONCHOPNEUMONIA, *Bron'chiopneumonia*, (*broncho*, and *pneumonia*.) Inflammation of the bronchia and lungs.

BRONCHORRHEMIA, (*broncho*, and *ῥημα*, 'blood,') Bronchorrhagia.

BRONCHORRHAG'IA, *Bronchorrhæ'mia*, (*broncho*, and *ῥήγνυμι*, 'I break forth,') Hemorrhage from the bronchial tubes.

BRONCHORRHÆA, (F.) *Bronchorrhée*, *Catarrhe pituiteux*, *Phlegmorrhagie pulmonaire*, *Flux bronchique*, (*broncho*, and *ῥεω*, 'I flow,') An increased secretion of mucus from the air passages, accompanied or not by inflammation,—a gleet, as it were, of the pulmonary mucous membrane, *Pneumorrhæ'a*. When excessive, it may constitute *Phtthisis pituitosa seu mucosa*, *Mycophthi'sis*.

BRONCHORRHÆA ACUTA, Bronchitis (chronic.)

BRONCHOSTASIS, (*broncho*, and *στασις*, 'station,') Bronchitis.

BRONCHOSTENIA, *Bronchostenosis*, (*broncho*, and *στενωσις*, 'narrow,') Contraction or narrowness of the bronchial tubes.

BRONCHOTOME, *Bronchot'omus*, (*broncho*, and *τεμνειν*, 'to cut,') A kind of lancet, with a blunt and rounded point, mounted on a handle, and fitted to a canula, which passes in along with it, and is allowed to remain in the opening made in the trachea.

BRONCHOTOMY, *Bronchotom'ia*, (F.) *Bronchotomie*. (Same etymology.) A surgical operation, which consists in making an opening either into the trachea, (*Tracheot'omy*;) into the larynx, (*Laryngot'omy*;) or into both, (*Tracheolaryngot'omy*;) to extract foreign bodies or to permit the passage of air to the lungs. These different parts are divided transversely or vertically, according to circumstances.

BRONCHOVESICULAR RESPIRATION, see Respiration.

BRONCHUS, (*βρογχος*), see Bronchia, Pharynx, Trachea.

BRONZED SKIN, see Capsule, renal.

BROOKLIME, *Veronica beccabunga*.

BROOM, (Sax. *brom*), *Sophora tinctoria*, *Spartium scoparium*—b. Butcher's, *Ruscus*—b. Clover,

Sophora tinctoria—b. Indigo, *Sophora tinctoria*—b. Rape of Virginia, *Orobancha Virginiana*—b. Spanish, *Spartium junceum*—b. Yellow, *Sophora tinctoria*.

BROSSARDIÈRE, MIN'ERAL WA'TERS OF. Brossardièrre is a chateau in Bas-Poitou, France. The waters contain carbonates of iron and lime, chloride of sodium, and sulphate of lime. They are aperient.

BROSSE. Brush.

BROTH, (Sax. brog, from brīan, 'to boil.') *Bro'dium, Jus'culum*, (F.) *Bouillon*. A weak decoction of meat, to which various vegetables and condiments may be added. A stronger decoction, with various modifications, constitutes *Soup*, *Zonos, Jus*, (F.) *Potage*.

BROTH, BLACK, OF THE LACEDÆMONIANS. *Jusculum nigrum Lacedæmoniorum*.

BROTH, CHICKEN, see Chicken broth.

BROTH, VEG'ETABLE. Take two potatoes, a carrot, and an onion, all cut fine; boil in a quart of water for an hour, adding more water from time to time, so as to keep the original quantity; flavour with salt, and a small quantity of pot-herbs; strain. A little mushroom catchup improves the flavour.

BROTHER, UTERINE, see Uterine.

BROU DE NOIX, ('the green envelope of the walnut,') see Juglans regia.

BROUILLARD, ('a mist,') Caligo.

BROUS'SAÏT. One who is a believer in, and professor of, the physiological and pathological opinions of Broussais. The system itself was called Broussais's or *The Physiological Doctrine*.

BROW, (Sax. bræp,) Front—b. Ague, Neuralgia frontalis.

BROWN RED, Colecothar.

BROWN'IAN, *Brownonian, Bruno'nian*. Relating to the system or opinions of Dr. John Brown.

BROWNISM, *Brownism, Bruno'nianism*. The doctrines of Brown.

BROWNIST, *Brownonian, Bruno'nian*. A follower of the system of Brown.

BRUCIA ANTI-DYSENTERICA, (after Bruce, the Abyssinian traveller.) *B. ferrugin'ea, Angustura spuria*, (F.) *Fausse Angusture, A. Ferrugineuse*. The systematic name of the plant whence was obtained—it was supposed—false *Angustura* or false *Cusparia Bark*. It is really the bark of *Strychnos nux vomica*.

BRUCIA, Brucine.

BRUCINE, *Brucia, Bruci'na, Bruci'num, Brucium, Pseudangusturinum, Canirami'num, Canira'minum, Vom'icine, Angust'urina*. An organic, salifiable base, discovered in the false angustura—*Brucea antidysenterica*—and obtained from *Strychnos nux vomica*. It is of a pearly white; crystallizes in oblique prisms with a parallelogrammatic base; is very bitter, slightly acrid and styptic, and soluble in water, but more so in alcohol. Brucia is a less active poison than strychnia. It resembles it, however, and may be used as a substitute for it and for the extract of *nux vomica*. Dose, half a grain.

BRUCKENAU, MIN'ERAL WA'TERS OF. These springs are in Bavaria, and contain carbonic acid and iron.

BRUCOURT, MIN'ERAL WA'TERS OF. Brucourt is three leagues and a half from Caen, in Normandy. The waters contain carbonic acid, chloride of sodium, and sulphate of soda, much sulphate of lime, &c.

BRUICK (Sc.), Furunculus.

BRUISE, (from (Sax.) brýran,) Contusion.

BRUISEROOT, *Stylophorum diphyllum*.

BRUISEWORT, *Bellis saponaria*.

BRUISSEMENT (F.), *Frem'itus*. This word

has much the same signification as *Bourdonnement*, as well as *Bruit*.

BRUIT (F.) 'noise,' *Strep'itus*. A sound heard on percussion or auscultation.

BRUIT DE CLAQUEMENT, ('clacking sound,') *Claquement*.

BRUIT DU CŒUR FÉTAL, ('sound of the foetal heart,') *Battemens doubles, Double bruit du cœur du Fœtus*. The pulsations of the foetal heart heard in auscultation in the latter half of uterogestation.

BRUIT DE CŒUR NEUF, ('sound of new leather,') *Bruit de craquement*.

BRUIT DE CRAQUEMENT, *B. de Tiraillement, Bruit de cuir neuf, Cri de cuir*, ('sound of crackling, or bursting, or of new leather,') A sound produced by the friction of the pericardium, when dried and roughened by inflammation.

BRUIT DE DIABLE, *Ronflement du Diable, Bruit de souffle à double courant*, ('sound of the diable or humming-top,') *Venous hum*. A high degree of *Bruit de souffle*, heard on auscultating the arteries or veins—probably the latter—of the neck, in chlorosis. It denotes an impoverished state of the blood.

BRUIT DOUBLE DU CŒUR DU FŒTUS, ('double sound of the heart of the foetus,') *Bruit du Cœur fœtal*—b. *d'Expansion pulmonaire*, see *Murmur, respiratory*—b. *de Froissement pulmonaire*, see *Froissement pulmonaire*—b. *de Frôlement*, see *Frôlement*—b. *de Frôlement péricardique*, see *Frôlement péricardique*.

BRUIT DE FROTTEMENT ASCENDANT ET DESCENDANT, ('sound of friction of ascent and descent,') Sounds produced by the rubbing of the lung against the parietes of the chest, as it rises and falls during inspiration and expiration. They are distinctly heard in pleuritis, when the pleura has become roughened by the disease. *Friction sounds, Rub'bing sounds, To-and-fro sounds* are also heard in pericarditis and peritonitis.

BRUIT HUMORIQUE, ('humoric sound,') *B. Hydropneumatique*. The sound afforded on percussion when organs are filled with liquid and air.

BRUIT HYDRO-PNEUMATIQUE, *Bruit humorique*—b. *de Jappement*, see *Sifflement modulé*—b. *de Lime à bois*, see *Bruit de Scie*.

BRUIT DE MOUCHE (F.), ('fly sound,') A sound analogous to the *Bruit de diable*—so called from its likeness to the buzzing of a fly—heard on auscultating the neck in chlorotic cases.

BRUIT MUSCULAIRE, ('muscular sound,') The sound accompanying the first sound of the heart, referred by some to muscular contraction. Called, also, *Bruit rotatoire*, in consequence of its having been thought to resemble the rumbling of distant wheels.

BRUIT MUSICAL, *Sifflement modulé*.

BRUIT DE PARCHEMIN, ('parchment sound,') A sound as if produced by two sheets of parchment applied to each other. It is said to be produced by thickening and rigidity of the valves of the heart.

BRUIT DE PIAULEMENT, ('whining sound,') see *Sifflement modulé*.

BRUIT PLACENTAIRE, *B. de souffle placentaire ou utérin, Souffle utérin ou placentaire, Placent'al bellows' sound, Utero-placent'al murmur, Uterine murmur or rush*. The bellows' sound heard on auscultating over the site of the placenta in a pregnant female. It does not appear to be owing to the placental vessels, but to the uterine tumor pressing upon the large vessels of the mother.

BRUIT DE POT FÉLÉ, ('sound of a cracked vessel,') *Cracked pot sound*. A sound heard on percussion, when a cavern in the lungs is filled

with air, and has a narrow outlet. It is not diastolic, however.

BRUIT DE RACLEMENT, ('sound of scraping.') A sound produced by the scraping of hard, solid membranes, as the pericardium, against each other, very analogous to *Bruit de craquement*.

BRUIT DE RÂPE, ('sound of a rasp.') A sound heard during the contraction of either the auricles or ventricles. It is constant; and the contraction of the cavity is more prolonged than natural, and emits a hard, rough, and—as it were—stifled sound.

It indicates contraction of the valvular orifices by cartilaginous deposits, or ossification, and is better heard near the apex of the heart, if the auriculo-ventricular valves are concerned,—near the base, if the semilunar valves are the seat of the disease.

BRUIT RESPIRATOIRE, Murmur, respiratory.

BRUIT ROTATOIRE, *Bruit musculaire*—*b. de Rouet*, ('sound of the spinning-wheel,') see *Sifflement modulé*.

BRUIT DE SCIE, ('saw sound,') and **BRUIT DE LIME À BOIS**, ('file sound,') resemble the *Bruit de Râpe*.

BRUIT DE SOUFFLE, *Bruit de soufflet*—*b. de Souffle céphalique*, see *Bruit de souffle*—*b. de Souffle à Double courant*, *Bruit de Diable*—*b. de Souffle placentaire*, *Bruit placentaire*—*b. de Souffle utérin*, *Bruit placentaire*.

BRUIT DE SOUFFLET, *Bruit de Souffle*, ('bellows' sound, 'blowing sound,') *Physèche ma*. A sound like that of a bellows, heard occasionally by the ear applied to the chest during the contraction of the ventricles, auricles, or large arteries. It coexists with affections of the heart, but is heard, also, without any disease of that organ,—whenever, indeed, an artery is compressed. An *Encephalic bellows' sound*, (F.) *Bruit de souffle céphalique*, has been described by Drs. Fisher and Whitney. It is heard on applying the ear to the occiput or to the top of the head; and is considered to indicate turgescence of vessels, or inflammation. When such turgescence exists, the vessels are compressed, and the compression gives rise to the sound in question.

BRUIT DE SOUPAPE, ('valvular or flapping sound,') A sound heard in respiration, when a foreign body is in the air passages. It somewhat resembles the flapping of a valve; hence its French name.

BRUIT DE TAFFETAS, ('sound of taffeta, sarcenet sound,') A respiratory sound, so named, by M. Grisolle, from its resembling the sound caused by the tearing of a piece of taffeta; and which he considers to indicate hepatization of the lung, limited to the surface, in pneumonia.

BRUIT DE TIRAILLEMENT, *Bruit de craquement*.

BRUIT TYMPANIQUE, ('tympanic sound,') The clear sound afforded by percussing the stomach and intestines when containing air.

BRUIT UTÉRIN, *B. placentaire*—*b. Vésiculaire*, Murmur, respiratory.

BRUITS DUCŒUR, ('sounds of the heart,') see Heart.

BRUK (Sc.), *Furunculus*.

BRÛLURE, Burn.

BRUNELLE, *Prunella*.

BRUNNER'S GLANDS, *Brunne'ri Glandulæ*, *G. Brunneri'ne seu solitariæ*, *Solitary glands or follicles*, *See'oul pan'creas*. Compound muciparous follicles, seated between the mucous and muscular coats of the stomach, along the two curvatures of that organ, and in the duodenum; so called from their discovery having been generally attributed to Brunner or Brunner. The solitary intestinal follicles are often known, at the present

day, as the glands of Brunner, although Brunner restricted the latter term to the glands of the duodenum.

BRUNONIAN, Brownian.

BRUNONIANISM, Brownism.

BRUNUS, *Erysipelas*.

BRUSCUS, *Ruscus*.

BRUSH, (F.) *Brosse*, *Scop'ula*. A well-known instrument, used in medicine chiefly for the following purposes:—1. To clean the teeth. 2. To remove the saw-dust which adheres to the teeth of the trephine, during the operation of trephining. 3. To rub the surface of the body, for the purpose of exciting the skin, and favouring transpiration. Westring, a Swedish physician, has recommended metallic brushes for the purpose of conveying galvanism to a part. These brushes consist of a plate of ebony fitted to another of gold, in which threads of the same metal are fixed;—the brush being connected with one of the poles of the galvanic pile.

BRUSH, *Stomach*, *Excitua ventriculi*.

BRUTA, *Juniperus sabina*.

BRUTIA. A sort of thick pitch, obtained from Brutia, in Italy. From *Pix Brutia* was obtained the *O'leum Pic'uum*.

BRUTIA, Instinct.

BRUTINO, *Terebinthina*.

BRUTOLÉ, (from *bryton*,) see *Cerevisia*.

BRUXANELLI. A Malabar tree, the bark and leaves of which have a strong smell, and are astringent. On the coast of Malabar, its juice, mixed with butter, is applied to boils. Its bark is esteemed to be diuretic, and its roots anti-arthritic.

BRUYÈRE VULGAIRE, *Erica vulgaris*.

BRUYÈRES, MINERAL WATERS OF. Bruyères is a small village, $7\frac{1}{2}$ leagues from Lunelville. The waters are acidulous and chalybeate.

BRYCETUS, (*βρυχευ*, 'to grind the teeth,') see *Algidus*.

BRYCHETHMUS, (*βρυχηθμος*,) *Rugitus*.

BRYCHETUS, (*βρυχετος*,) see *Algidus*.

BRYGMA, gen. *Bryg'matis*, *Bryg'mus*, (*βρυγμα*, *βρυγμος*,) *Pris'is*, *Pris'mus*, *Odonto pris'is*, *Stri'dor Dentium*, (F.) *Gricement des Dents*. Grinding of the teeth. A common symptom, in children, of gastric or other derangement, but often present when there is no reason to suspect any.

BRYONE, *Bryonia alba*—*b. d'Amérique*, *Convolvulus Mechoacan*.

BRYONTA, (*βρυων*, from *βρω*, 'I bud forth,') *Bryonia alba*.

BRYO'NIA AFRICA'NA, *Af'rican Bry'ony*, A South African plant, common amongst the Hottentots, which, in the form of decoction, acts simultaneously as an emetic, cathartic, and diuretic. It is used by the natives in cutaneous diseases, dropsy, and syphilis. The tincture is a powerful emetic and cathartic.

BRYONIA AL'BA, *Bryo'nia*, *Bryo'nē*, *Vit'is al'ba sylvest'ris*, *Agros'tis*, *Agriam'pelos*, *Ampelulē'cē*, *Am'pelos a'gria*, *Echeto'sis*, *Bryo'nia as'pera seu Dioic'a*, *Cedros'tis*, *Chelido'nium*, *Labrus'ca*, *Melothrum*, *Ophiotaph'ylon*, *Psilo'thrum*, *Bry'ony*, *White bryony*. Ord. *Cucurbitaceæ*. Sex. Syst. *Monœcia Monadelphia*. (F.) *Bryone*, *Contenvrée*, *Vigne vierge*, *V. blanche*, *Nœut du diable ou galant*. The root is large and succulent, and has an acrid, bitter, and disagreeable taste. It is a drastic cathartic. Externally, it has been applied, in form of cataplasm, in gout. When repeatedly washed, a good starch is obtained from it. The active bitter principle has been separated from it, and called *Bry'oniin*.

BRYONTA MECOACANNA NIGRICANS, *Convolvulus jalapa*—*b. Nigra*, *Tamus communis*—*b. Peruviana*, *Convolvulus jalapa*.

BRYONINE, see *bryonia alba*.

BRYONY, *Bryonia alba*—b. Black, *Tamus communis*—b. White, *Bryonia alba*—b. Wild, *Sycios angulatus*.

BRYTIA, (*Burra*.) Mare of grapes.

BRYTOLATURE, (from *Bryton*.) see *Cerevisia*.

BRYTOLÉ, see *Cerevisia*.

BRYTON, (*Βρυτον*.) *Cerevisia*.

BU, (*βου*, abbreviation of *βους*, 'an ox,') in composition, expresses 'excess, greatness.' Hence *Bulimus*, *Buphthalmia*, &c.

BUBASTECORDIUM, *Artemisia vulgaris*.

BUBE, Pustule.

BUBO, (*βουβων*.) *Panotchia*, *Panus inguinalis*, *Adenophyma inguinalis*, *Bubonopanus*, *Bubononcus*, *Cambuca*, *Boubon*, *Codocel'le*, *Codocell'la*, (F.) *Bubon*, *Poulain*. In the works of Hippocrates and Galen, *βουβων*, sometimes signifies the groin—

Inguen; at others, the inguinal glands; and at others, again, swelling or inflammation of these parts, *Bubonadenitis*. The moderns apply the term to an inflammatory tumour seated in the groin or axilla, and they generally distinguish—

1. *Simple or Sympathetic Bubo*, which is independent of any virus in the economy. 2. *Veneral Bubo*,—of old, *Button of Naples*, (F.) *Bubon vénérien*, which is occasioned by the venereal virus.

3. *Pestilential Bubo*, or *B. symptomatic of the Plague*. The last two have by some been called malignant *Bubo*, (F.) *Bubon malin*.

BUBO, PRIMITIVE, *Bubon d'emblée*.

BUBON, *Bubo*, *Inguen*—b. Gummiiferum, see *Ammoniac gum*.

BUBON, (*βουβων*.) In composition, the groin.

BUBON D'EMBLÉE (F.), *Primitive or primary bubo*, (F.) *Bubon primitif*. An enlargement and suppuration of one or more of the inguinal glands, not preceded by any other of the more common forms of venereal disease, nor by any other syphilitic symptom.

BUBON GALBANUM. The systematic name of a plant which has been supposed to afford galbanum; *Melopion*, *Matorium*. The plant is also called *Ferula Africa'na*, *Oreoselinum Africa'num*, *Anisum fruticosum galbaniferum* seu *Africa'num frutescens*, *Selinum Galbanum*, *Agasyll'lis galbanum*, *The long-leaved or lovage-leaved Galbanum*. *Ord. Umbelliferae*. The plant which affords the galbanum can scarcely, however, be considered to be determined. By the Dublin College, it was referred to *Opor'odia Galbanif'era*, *Ord. Umbelliferae*. Galbanum, *Chal'bané*, is a gummy-resinous juice. Its odour is fetid, and taste bitter and acrid: the agglutinated tears are of a white colour, on a ground of reddish-brown. It forms an emulsion when triturated with water, and is soluble in proof spirits of wine, and vinegar: sp. gr. 1.212. It has been given as an anti-spasmodic and expectorant, in pill or emulsion. Dose, from gr. 10 to 60. Externally, it is applied as a cataplasm.

Bubon galbanum is a South African plant; and is reputed to be an excellent diuretic, under the name of *Wild Celery*. A decoction of the leaves is given in dropsy and gravel. According to Pappé, the resinous matter, which exudes from the stem, differs in appearance, smell, and in every respect, from Gummi Galbanum.

BUBON MACEDONICUM, *Athaman'ta Macedonica*, *Petroselinum Macedonicum*, *A'pium petr'um*, *Petra'pium*, (F.) *Persil de Macédoine*, *Macedonian Parsley*. Its properties are similar to those of common parsley, but weaker and less grateful. The seeds are an ingredient in the celebrated compounds, *Mithridate* and *Theriac*.

BUBON PRIMITIF, *Bubon d'emblée*.

BUBONA, Nipple.

BUBONADENITIS, (*bubon*, *aden*, and *itis*.) see *Bubo*.

BUBONAL'GIA, (*bubon*, and *αλγος*, 'pain,') Pain in the groin.

BUBONIUM, *Aster At'ticus*, *Gold'en Starwort*. A plant anciently supposed to be efficacious in diseases of the groin.

BUBONOCÉ'LE, *Bu'bonocèle*, (*bubon*, and *κλήη*, 'tumour,' 'rupture,') *Her'nia inguinalis*, *Inguinal Her'nia*, *Rupture of the Groin*. (F.) *Hernie inguinale*. Some surgeons have confined this term to hernia when limited to the groin, and have called the same affection, when it has descended to the scrotum, *Oscheocel'le*, *Scrotal Her'nia*. The rupture passes through the abdominal ring; and, in consequence of the greater size of the opening in the male, it is more frequent in the male sex.

BUBONONCUS, (*bubon*, and *ογκος*, 'a swelling,') *Bubo*.

BUBONOPANUS, *Bubo*.

BUBONORRHÉ'XIS, *Bubonorex'is*, (*bubon*, and *ρηξις*, 'a rupture,') A name given to bubonocèle when accompanied with a division of the peritoneum, or when, in other words, it is devoid of a sac.

BUBON'ULUS, *Bubun'culus*, (dim. of *Bubo*.) A painful swelling of the lymphatics of the penis, extending along the dorsum of that organ to the groin. It is an occasional accompaniment of gonorrhœa.

BUBUKLE. A word used by Shakspeare for a red pimple on the nose.

BUBUNCULUS, (dim. of *bubo*.) *Bubonulus*.

BUCAROS, Terra Portugallica.

BUC'CA. The mouth. The cheek and hollow of the cheek. Also, the vulva.

BUCCAC'RATON, *Gna'thos*, (*bucca*, 'a mouthful,' 'a morsel,' and *κραα*, 'I mix,') A morsel of bread sopped in wine, which served, of old, for a breakfast.

BUC'CAL, *Buccal'is*, (from *bucca*.) That which concerns the mouth, and especially the cheek.

BUCCAL AR'TERY, *Artère Sous-maxillaire*, (Ch.) arises from the internal maxillary or from some of its branches, as the *Temporalis profunda anti'ca*, or the *Alveolar*. It distributes its branches to the buccinator muscle, and to the buccal membrane.

BUCCAL GLANDS, *Molar Glands*. Mucous follicles, seated in the buccal membrane, opposite the molar teeth. They secrete a viscid humour, which mixes with the saliva, and lubricates the mouth.

BUCCAL MEM'BRANE, (F.) *Membrane Buccale*. The mucous membrane, which lines the interior of the mouth.

BUCCAL NERVE, *Buccina'tor Nerve*, *Buccolabial*—(Ch.), is given off by the inferior maxillary. It sends its branches to the cheek, and especially to the buccinator muscle.

BUCCAL VEIN follows the artery.

BUC'CEA, *Buccell'la*. The fleshy excrecence of nasal polypus, so called because it was believed to proceed from the mouth. Also, a mouthful.

BUCCEL'ATON, *Buccell'atus*. A loaf-shaped cathartic medicine; made chiefly of scammony.

BUCCELLA, (dim. of *bucca*, 'a morsel,') *Bolus*, *Bucca*.

BUCCELLA'TIO. A mode of arresting hemorrhage, by applying a pledget of lint to the bleeding vessel.

BUCCINA, (*βυκων*, 'a trumpet,') Turbinated bones.

BUCCINA'TOR, (*buccinare*, 'to sound the trumpet,') The *Buccina'tor Muscle*, *Retractor Anguli O'ris*, *Manso'rius*, (F.) *Buccinateur*, *Bucco-Alvéolo-maxillaire*, *Alvéolo-labial*—(Ch.), is situated in the substance of the cheeks. It extends between the posterior portions of the alveolar arches of the two jaws and the commissure of the lips, which it draws backward. It assists in mastication, by pushing the food back towards the teeth; and, if the cheeks be distended by air, its contraction forces it out.

BUC'CO. One who is blub-checked, or wide-mouthed. In composition, the mouth or cheek. Also, *Diosma crenata*.

BUCCO-ALVEOLO-MAXILLAIRE, Buccinator.

BUCCOLABIAL NERVE, Buccal nerve.

BUCOPHARYNGE'AL, *Buccopharynge'us*, (F.) *Bucco-Pharyngien*. Belonging to the mouth and pharynx. The *Buccopharynge'al Aponeurosis* or *Intermaxillary Ligament*, extends from the internal ala of the pterygoid process to the posterior part of the lower alveolar arch, and affords attachment, anteriorly, to the buccinator, and, posteriorly, to the constrictor pharyngis superior.

BUC'ULA, (dim. of *bucca*.) A small mouth. The fleshy part beneath the chin.

BUCERAS, (*Βουκερας*.) *Trigonella fœnum* — *b. Fœnum Græcum*, *Trigonella fœnum Græcum*.

BUCHU, *Diosma crenata* — *b. Leaves*, *Diosma crenata*.

BUCKBEAN, *Menyanthes trifoliata* — *b. American*, *Menyanthes verna*.

BUCKBERRY, *Vaccinium stamineum*.

BUCKET FEVER, Dengue.

BUCKEYE, *Æsculus hippocastanum*.

BUCKHO, *Diosma crenata*.

BUCKTHORN, *PURGING*, *Rhamnus*.

BUCKU, *Diosma crenata*.

BUCKWHEAT, (said to be a corruption of *beech wheat*; from a resemblance to the beech nut,) *Polygonum fagopyrum* — *b. Plant*, eastern, *Polygonum divaricatum*.

BUCNEMIA, (*bu*, and *κνήμη*, 'leg,') see Elephantiasis — *b. Tropica*, see Elephantiasis.

BUCTON, Hymen.

BUFF INFLAMMATORY, (*buff*, 'the colour of buffalo leather, a light yellow,') *Corium phlogisticum*.

BUFFY COAT, *Corium phlogisticum*.

BUG, (*BED*), Cimex.

BUGANTIA, Chibblain.

BUG'ERY, *Sod'omy*, *Sodom'ia*, *Col'itus Sodomit'icus*, (I.) *Bugarone*, (F.) *Bougrerie*. Said to have been introduced by the Bulgarians, *Bulgari*, *Bugari*, *Bugeri*, hence the name. A carnal copulation against nature, as of a man or woman with any animal; or of a man with a man, or a man unnaturally with a woman. The unnatural crime.

BUGLE, (*bugula*), *Prunella* — *b. Common*, *Ajuga reptans* — *b. Pyramidale*, *Ajuga* — *b. Rampante*, *Ajuga reptans* — *b. Water*, *Lycopus Virginicus* — *b. Weed*, *Lycopus*.

BUGLOSE, *Anchusa officinalis*.

BUGLOSS, *DYER'S*, *Anchusa tinctoria* — *b. Garden*, *Anchusa officinalis* — *b. Upright*, *Ajuga*.

BUGLOSSA, (*βους*, 'an ox,' and *γλωσσα*, 'tongue,' from the resemblance of its leaves,) *Anchusa officinalis*.

BUGLOSSUM ANGUSTIFOLIUM MAJUS, (*βουγλωσσον*, 'ox tongue,') *Anchusa officinalis* — *b. Latifolium*, *Borago officinalis* — *b. Sativum*, *Anchusa officinalis* — *b. Sylvestris*, *Anchusa officinalis* — *b. Tinctorum*, *Anchusa tinctoria* — *b. Verum*, *Boracae acid* — *b. Vulgare majus*, *Anchusa officinalis*.

BUGRANDE ÉPINEUSE, *Ononis spinosa*.

BUGRANE, *Ononis spinosa* — *b. des Champs*, *Ononis arvensis*.

BUGULA, (dim. of *buglossa* [?]), *Ajuga* — *b. Chamæpitys*, *Teucrium chamæpitys* — *b. Pyramidalis*, *Ajuga* — *b. Reptans*, *Ajuga reptans*.

BUIS, *Buxus*.

BUISARD, MIN'ERAL WA'TERS OF. Buisard is two leagues from Chateau-Thierry, in France. The water contains chloride of calcium and carbonate of lime.

BULB, *Bul'bus*, (*βολβος*.) (F.) *Bulbe*. A name, given by anatomists to different parts which resemble, in shape, certain bulbous roots. The

Bulb of the Aorta is the great sinus of the Aorta. *Bulb of a Tooth* is the vascular and nervous papilla contained in the cavity of a tooth. The *Bulb* or *Root of the Hair* is the part whence the hair originates;—see Hair. The *Bulb of the Ure'thra* is the dilated portion formed by the commencement of the *Corpus spongiosum* towards the root of the penis. We say, also, *Bulb*, for *Globe*, of the eye.

BULB OF THE EYE, see Eye—*b. of the Female*, *Bulbus vestibuli*.

BULB, *RACHIDIAN*, see *Medulla oblongata*.

BULB OF THE THROAT, Tonsil.

BULBE, *Bulb* — *b. Rachidien*, see *Medulla oblongata* — *b. du Vagin*, *Bulbus vestibuli* — *b. de la Voûte à trois Piliers*, Mamillary tubercles.

BULBI FORNICIS, (pl. of *bulbus*.) Mamillary tubercles — *b. Priorum Crurum Fornicis*, Mamillary tubercles.

BULBOCASTANEUM, (*βολβος*, 'a bulb,' and *καστανον*, 'a chestnut,') *Bunium bulbocastanum*.

BULBO-CAVERNEUX, Accelerator urinæ.

BULBOCAVERNOSUS, Accelerator urinæ — *b. Syndesmo-cavernæux*. Accelerator urinæ — *b. Ure'thral*, Accelerator urinæ.

BULBOCODIUM, *Narcissus pseudonarcissus*.

BULBONACH, *Lunaria rediviva*.

BULBUS, (*βολβος*.) *Bulb*.

BUL'BUS ESCULEN'TUS. The *Es'culent Bulb*: a particular kind, so denominated by the ancients. It is supposed to have been the *Ce'pa Ascalon'ica*.

BULBUS GLANDULOSUS, Proventriculus — *b. Medullæ spinalis*, *Medulla oblongata* — *b. Oculi*, see Eye — *b. Olfactorius*, see Olfactory Nerves — *b. Pili*, see Hair — *b. Rachidius*, see *Medulla oblongata* — *b. Vaginae*, *B. vestibuli*.

BULBUS VESTIB'ULI, *B. Vaginae*, *Plex'us retiformis seu reticularis seu cavernosus*, *Crura clitoridis interna*, *Bulb* or *Sem'ibulb of the Female* or of the *vagina*, (F.) *Bulb du Vagin*. A close-packed plexus of intricately anastomosing veins, inclosed in a fibrous investment,—being an immediate continuation and extension of the *pars intermedia*, and occupying the space between the beginning or vestibule of the vagina and the rami of the pubic arch on each side. It is regarded by Lauth, Taylor, Morgagni and Kobelt as the analogue of the male bulb.

BULBUS VOMITO'RIOUS. A plant, said by Dioscorides to be emetic and diuretic. It is the *Musk-grape flower*, according to Ray,—the *Hya-cinth'us Muscari*.

BULESIS, (*βουλησις*, from *βουλομαι*, 'to will,') Voluntas.

BULGA, Vulva.

BULIMIA, Boulimia.

BU' LITHOS, (*βους*, 'an ox,' and *λιθος*, 'a stone,') A bezoar or stone, found in the kidneys, gall-bladder, or urinary bladder of an ox or cow.

BUL'LA, (F.) *Bulle*, *Bleb*, (Sc.) *Bleib*. A portion of the cuticle, detached from the skin by the interposition of a transparent, watery fluid. It forms the 4th order in Willan's and Bateman's arrangement of cutaneous diseases, and includes erysipelas, pemphigus, and pompholyx. By some, Bulla has been used synonymously with *Pemphigus*. See, also, Hydatid.

BULLE ROTUNDÆ CERVICIS UTERI, Nabothi glandulæ.

BULLACE PLUM, *Prunus insitia*.

BULLDOG, *Tabanus*.

BULLFISTS, *Lycoperdon*.

BUL'LOUS, *Bullo'us*, (F.) *Bulleux*. Having relation to a bulla or bleb, as a 'bullous eruption.' Pemphigus has been designated *Maladie bulleuse*, and *Fievre bulleuse*, when accompanied by fever.

BULLSEGG, ('bulsedge,') *Typha latifolia*.

BULL'ULA, diminutive of *bullu*. A small vesicle or bleb.

BUMBLEKITES, see *Rubus fruticosus*.

BUMELIA, (*βουμेलία*), *Fraxinus excelsior*.

BUMGUT, Rectum.

BUNA, *Coffea Arabica*.

BUNCH, Hunch.

BUNCHBACKED, see Hunch.

BUNCOMBE, WARM AND HOT SPRINGS

OF. See North Carolina, Mineral Waters of.

BUNDURH, *Corylus avellana*.

BUNEWAND, *Heracleum spondylium*.

BUNIAS, (*βουνίας*), *Brassica napus*.

BUNIOID, (*βουνιον*, 'a kind of turnip,' and *ειδος*, 'resemblance,') *Napiform*.

BUNION, *Bunyon*.

BUN'ITES VINUM, (*βουνιτης*.) A wine, made by infusing the *Bunium* in must. It is stomachic, but scarcely ever used.

BUN'NIUM CAR'VI, (*βουνιον*, so called, it has been supposed, from growing on hills, from *βουνος*, 'a hill,' or from the tuberosity of its root.) *Carum*.

BENIUM BULBOCAS'TANUM, *B. mi'nus*, *Balanocastanum*, *Si'um bulbocast'anum*, *Scan'dex bulbocast'anum*, *Ca'rum bulbocast'anum*. Order, *Umbelliferae*. The systematic name of a plant, whose root is called *Agriocast'anum*, *Nu'cula terres'tris*, *Bulbocast'anum majus* et *mi'nus*. *Pig-nut*, *Earth-nut*, *Hawk-nut*, *Kipper-nut*, (Sc.) *Arnut*, (F.) *Terrenoiz*. The root is tuberous, and is eaten raw or roasted. It has been supposed to be of use in stranguy. It is not employed in medicine.

BUN'NIAN, *Bunyon*.

BUNWEED, *Senecio Jacobæa*.

BUN'YON, *Bun'ion*, *Bun'nian*, (*βουνος*, 'an eminence.' [?]) An enlargement and inflammation of the bursa mucosa at the inside of the ball of the great toe.

BUOPHTHALMIA, *Buphthalmia*.

BUPEINA, (*bu* and *πεινα*, 'hunger,') *Boulimia*.

BUPHTHALMI HERBA, *Anthemis tinctoria*.

BUPHTHAL'MIA, *Buphthal'mia*, *Buphthal'mus*, *Elephantom'ma*, (*bu*, and *οφθαλμος*, 'an eye.') *Ox-eye*. Under this name, the generality of authors have designated the first stage of hydrophthalmia. Others, with Sabatier, mean by it turgescence of the vitreous humour, which, by pushing the iris forwards, forms around the crystalline a sort of border.

BUPHTHALMUM CRETICUM, *Anthemis Pyrethrum*—*b. Majus*, *Chrysanthemum leucanthemum*.

BUPHTHALMUS, *Buphthalmia*, *Hydrophthalmia*, *Sempervivum tectorum*.

BUPINA, (*bu*, and *πεινα*, 'hunger,') *Boulimia*.

BUPLEUROIDES, *Bupleurum rotundifolium*.

BUPLEU'RUM ROTUNDIFOLIUM, *Bupleu'ron*, *Bupleuroi'des*, (*bu*, and *πλευρον*, 'a rib,' the leaves having large ribs,') (F.) *Buplèvre*, *Percefeuille*, *Round-leaved Hare's Ear*, *Thoracicææ*. Order, *Umbelliferae*. The herb and seeds are slightly aromatic. It was formerly celebrated for curing ruptures, being made into a cataplasm with wine and oatmeal.

BUPLEV'RE, *Bupleurum rotundifolium*.

BURAC (Arab.), *Borax*. Also, any kind of salt.

BURBOT, see *Oleum Jecoris Aselli*.

BURDOCK, *Aretium lappa*—*b. Lesser*, *Xanthium*—*b. Prairie*, *Silphium terebinthaceum*.

BURGOO, see *Avena*.

BURGOUT, see *Avena*.

BURIAL ALIVE, *Zoothapsis*.

BURIS, *Hernia*, accompanied by scirrhus tumefaction; or, perhaps, a scirrhus tumour only.

BURN. (Sax. *bernan* or *býrnan*, 'to burn or bren.') *Us'tio*, *Ambus'tio*, *Adus'tio*, *Tre'sis Cau'sis*, *Erythéma Ambus'tio*, *Cau'sis*, *Encau'sis*, *Pyri-*

caus'tum, *Combustu'ra*, *Catacau'ma*, *Combust'io*, (F.) *Brûlure*. An injury produced by the action of too great heat on the body. Burns are of greater or less extent, from the simple irritation of the integument to the complete destruction of the part. The consequences are more or less severe, according to the extent of injury, and the part affected. Burns of the abdomen, when apparently doing well, are sometimes followed by fatal results. Their treatment varies,—at times, the antiphlogistic being required; at others, one more stimulating.

BURNEA, see *Pinus Sylvestris*.

BURNET, CANADA, *Sanguisorba Canadensis*.

BUR'NETT'S DISINFECT'ING LIQ'UID or **FLU'ID**. A solution of chloride of zinc, first used by Sir William Burnett for preserving timber, canvas, &c., from dry rot, mildew, &c., and afterwards as an antiseptic and antiseptic, especially in the case of dead bodies. The Dublin Pharmacopœia had a *Zinci Chloridi Lignor, Solu'tion of Chloride of Zinc*, which is not near so strong as Sir William's preparation.

BURN'ING, Bren'ning. A disease mentioned by old historians, from which authors have unsuccessfully endeavoured to demonstrate the antiquity of gonorrhœa virulenta and syphilis.

BURNING OF THE FEET, see *Feet*, burning of the.

BURNT HOLES. A variety of *rupia*, popularly known in Ireland under this name; and not unfrequently there amongst the ill-fed children of the poor. See *Rupia escharotica*.

BURR, (by *onomatopœia*.) *Rotacismus*.

BURRAGE, *Borago officinalis*.

BUR-REED, GREAT, *Sparganium ramosum*.

BURRH, (like *Burr*), *Rotacismus*.

BUR'RHI SPIR'ITUS MATRICALIS. The *Spirit of Burrhus* for diseases of the Womb. It is prepared by digesting, in alcohol, equal parts of myrrh, olibanum, and mastic. Boerhaave frequently prescribed it.

BURSA, (*βυρσα*, 'a sac, 'a leathern bottle,')

Scrotum—*b. Cordis*, *Pericardium*—*b. Omentalis*, see *Epiploon*, *gastro-hepatic*—*b. Pastoris*, *Thlaspi bursa*—*b. Testium*, *Scrotum*—*b. Virilis*, *Scrotum*.

BURSÆ MUCILAGINOSÆ, *B. mucosæ*.

BURSÆ MUCO'SÆ, *B. mucosæ vesicula' res*, *Bursæ seu Cap'sule synovial'es*, *Blennocys'tides*, *Sac'ci mucos'i*, *Vesi'cæ unguino'sæ ten'dinum*, *Vagi'næ Synovial'es seu mucilaginosæ*, *Synovial Crypts* or *Fol'dicles*, (F.) *Bursæ Synoviales* ou *mucosæ* ou *mucilaginosæ*. Small membranous sacs, situate about the joints, particularly about the large ones of the upper and lower extremities, and, for the most part, lying under the tendons. They are naturally filled with an oily kind of fluid, the use of which is to lubricate surfaces over which the tendons play. In consequence of bruises or sprains, this fluid sometimes collects to a great extent. The bursæ are, generally, either of a roundish or oval form, and they have been arranged under two classes, the *spherical* and the *vag'inal*.

BURSÆ SYNOVIALES, *Bursæ mucosæ*.

BUR'SAL, *Bursa'lis*. Relating or appertaining to bursæ,—as a 'bursal tumour.'

BURSALIS, *Bursal*, *Obturator internus*.

BURSAL'OGY, *Bursalog'ia*, (from *bursa*.) A treatise on, or description of, the bursæ mucosæ.

BURSER A ACUMINATA, (after Joachim Burser, a Neapolitan botanist.) *B. gummiifera*.

BURSE'RA GUMMI'FERA, *B. acuminata*, *Terebin'thus gummi'fera*, *Jama'ica Bark Tree*. Order, *Terebinthaceæ*. A resin exudes from this tree, which, as met with in the shops, is solid externally; softish internally; of a vitreous fracture; transparent; of a pale yellow colour; tur-

pentine smell, and sweet, perfumed taste. It has been used like balsams and turpentine in general, and is called, by the French, *Cachibou*, *Chibou*, and *Résine de Gomart*.

BURSITIS, (*bursa*, and *itis*), see Synovitis.

BURST, ('ruptured,') Hernia, Hernial.

BURSTEN, see Hernial.

BURSULA, (dim. of *bursa*), Scrotum.

BURTHISTLE, Xanthium.

BURTREE, Sambucus.

BURUNHEM, Monesia.

BURWEED, Xanthium.

BURWORT, Ranunculus acris.

BUSH, JEW, Pedilanthus tithymaloides.

BUSSANG, MINERAL WATERS OF. Bussang is a village in the department of Vosges, France. The waters are acidulous chalybeates.

BUSSEROLLE, Arbutus uva ursi.

BUS'SII SPIRITUS BEZOARTICUS, *Bezoair'dic Spirit of Bussius*. A preparation, regarded as sudorific, diuretic, and antispasmodic; obtained by distilling subcarbonate and muriate of ammonia, amber, oil of cedar or juniper, &c.

BUTE, ISLAND OF, CLIMATE OF. This island is in the Frith of Clyde, about 18 miles below Greenock. The climate is mild and equable, but rather moist; and, as a winter residence, it holds out advantages for those only that appear to demand such a condition of the atmosphere. The climate resembles, in character, that of the S. W. of England and France, and the Channel islands; although its temperature is lower.

BU'TEA FRONDO'SA, (after John, Earl of Bute, a promoter of botany,) *Erythrina monosperma*, *Rudolphia frondosa*. A tree, common in Bengal, and in the mountainous parts of India; *Nat. Ord.* Leguminosæ; from which *gum butea* flows. Dr. Pereira found this gum to be identical with a specimen marked *gummi rubrum adstringens*—the *gomme astringente de Gambie* of M. Guibourt. By some, this gum has been confounded with kino. See Kino.

BUTIGA, Gutta rosea.

BUTOMON, (*Βουτόμων*), Iris pseudacorus.

BUTTER, (*Βουτύρον*; itself from *βους*, 'ox,' and *ρως*, 'anything coagulated,') *Butyrum*, *Pice'tion*, (F.) *Beurre*. A sort of concrete oil, obtained from the cream that forms on the surface of the milk furnished by the females of the mammalia, especially by the cow and the goat. Fresh butter is digestible, whilst the rancid is irritating. The ancient chemists gave the name *Butter* to many of the metallic chlorides. It has also been applied to vegetable substances, which resemble, in some respects, the butter obtained from milk. The essential fatty matter in it is *butyrin* or *butyrate of glycerin*.

BUTTER OF ANTIMONY, Antimonium muriatum.

BUTTER OF BAMBOUC or BAMBOC, (F.) *Beurre de Bambouc* ou *Bambuk*. A vegetable oil obtained from a species of almond, and used in Senegal in neuralgic and rheumatismal pains.

BUTTER OF CA'CAO, Oil of Ca'cao, *O'leum Ca'cao spissatum*, O. Theobroma, O. Theobroma Ca'cao expres'sum, Oil of Theobroma, (F.) *Beurre de Cacao*, *Huile de Cacao*. The concrete oil, of a sweet and agreeable taste, obtained from the kernels of the fruit of *Theobroma cacao* or chocolate nut. Owing to its firmness and fusibility, it is well adapted to serve as the constituent of suppositories.

BUTTER OF CO'COA, (F.) *Beurre de Coco*. A fatty, concrete substance, which separates from the milk of the cocoa nut. It is sweet and agreeable.

BUTTER AND EGGS, Narcissus pseudonarcissus -- b. Kokum, see *Garcinia purpurea* -- b. of Nutmegs, see *Mystica*.

BUTTERBUR, Tussilago petasites.

BUTTERCUPS, Ranunculus acris, and R. bulbosus.

BUTTERFLY-WEED, Asclepias tuberosa.

BUTTERMILK, (Prov.) *Churnmilk*, *Kernmilk*, *Whig*, (Sc.) *Bladoch*, *Bledoch*, *Bladda*, *Kirn-milk*, *Sourmilk*, (F.) *Baburre*, *Lait de Beurre*. The thin, sour milk, separated from the cream by churning. It contains caseum and a little butter. It is a refreshing drink when newly made.

BUTERNUT, Juglans cinerea.

BUTTERTEETH, Incisive teeth.

BUTTERWEED, Erigeron Canadense.

BUTERWORT, Pinguicula vulgaris.

BUTTOCKHUMP, Stenotopyga.

BUTTON OF THE HAIR, see Hair -- b. of Naples, Bubo.

BUTTONBUSH, Cephalanthus occidentalis.

BUTTON SCUR'VY, *Mo'cula*. An epidemic cachectic affection, which has appeared in the Southern counties of Ireland, and is characterized by indolent buttonlike growths of the corpus papillare of the skin. It appears to be allied to Frambœsia.

BUTTONWOOD SHRUB, Cephalanthus occidentalis.

BUTUA, Pareira brava.

BUTYRATE OF GLYCERIN, see Butter.

BUTYRIN, see Butter.

BUTYRUM, (*Βούτυρον*) Butter -- b. Amygdalorum dulcium, Confection (almond) -- b. Saturni, Unguentum plumbi superacetatis -- b. Zinci, Zinci chloridum.

BUVEUR, ('a drinker,') see Bibitorius, Rectus internus oculi.

BUX'TON, MINERAL WATERS OF, *Buxtonien'ses A'quæ*. Buxton is a village in Derbyshire. The springs are thermal, and about 82° Fahrenheit. They contain sulphate of soda, chloride of calcium, chloride of sodium, chloride of magnesium, carbonate of lime, carbonic acid, and azote. They are used in cases in which thermal springs, in general, are recommended. They contain little or no mineral impregnation.

BUX'US, *Buxus sempervirens*. The Box-tree, (F.) *Buis* on *Bois*. The leaves are bitter and aromatic, and, as such, have been used in medicine, in cases of worms, dyspepsia, &c., in the form of decoction. They are sometimes, also, added to beer. The seed was anciently called *Carthe'gon*.

BYNE, (*Βυνη*), Malt.

BYRETHRUM. A sort of cap or *Couvrechef* filled with cephalic substances.

BY'RON AC'ID or SOUR SPRINGS. These Springs are in the town of Byron, Genesee county, New York. The water is a nearly pure dilute sulphuric acid. They are powerfully astringent and tonic.

BYR'SA, (*Βύρσα*, 'a hide,') A leather skin to spread plasters upon.

BYRSODEP'SICON, (*Βύρσα*, 'a hide,' and *δεΐνω*, 'I tan,') A tan stuff, with which CÆLIUS AURELIANUS sprinkled wool, which he applied in certain cases to the umbilical region.

BYRSODEPSICUM PRINCIPIMUM, Tannin.

BYSAU'CHEN, (*βω*, 'I stop up,' and *αυχνη*, 'the neck,') A morbid stiffness of the neck. One with a short neck, -- *Simotrache'lua*.

BYS'SUS, (*Βύσσος*), *Bys'sum*. The ancients gave this name to several vegetable substances, which were used for the fabrication of stuffs prized for their fineness, colour, and rarity of material. It is now chiefly applied to the filaments, by the aid of which the acephalous mollusca attach their shells to the rocks. Byssus meant formerly also the vulva.

BY'THUS, (*βυθός*, 'depth,') An epithet used by Hippocrates for the fundus of the stomach.

C.

C. This letter in the chemical alphabet signifies nitre. It is also sometimes used in prescriptions for calx.

CAA-AP'IA, *Dorstenia Brazilianis* seu *cordifolia* seu *placentoïdes* seu *vittel'la*. Order, Urticæ. The root, according to Piso, is employed as emetic and antidiarrhœic.

CAA-ATAY'A. A plant of Brazil, supposed to be a species of *gratiola*. It is very bitter, and considered to be one of the best indigenous cathartics.

CAACICA, *Euphorbia capitata*.

CAA-GHIYU'YO, *Fru'tex bac'eifer Brazilian'is*. A shrub of Brazil, whose leaves, in powder, are considered detersive.

CAAOPIA, *Hypericum bacciferum*.

CAAPEBA, *Paraiba brava*, *Pothomorphe peltata* and *P. umbellata*.

CAAPONGA, *Crithmum maritimum*.

CAAROBÀ. A Brazilian tree whose leaves, in decoction, promote perspiration. See *Ceratonia*.

CABAL', *Cab'ala*, *Cabal'la*, *Cal'bala*, *Caba'lia*, *Kab'ala*, *Gaballa*. This word is from the Hebrew, and signifies knowledge transmitted by tradition. Paracelsus and several authors of the 16th and 17th centuries have spoken much of this species of magic, which they distinguished into *Juda'ic* or *theolog'ian*, and *Hermetic* or *medic'inal*; the latter, being, according to them, the art of knowing the most occult properties of bodies by an immediate communication with spirits,—the knowledge being thus acquired by inspiration, and incapable of inducing error. It was also called *Are cabalistica* seu *signa'ta*, 'cabalistic art.'

CABAL'HAU. A plant of Mexico, according to Dalechamps, which passes for an antidote to white hellebore, and yet is used for poisoning arrows. It is unknown to botanists.

CAB'ALIST, *Cabalis'ta*. One instructed in the Cabal.

CABALLATION, *Cynoglossum*.

CABARET, *Asarum*.

CABBAGE, (*caput*, (I.) *capuccio*, 'a head.') *Brassica*—c. Cow, *Nymphæa odorata*—c. Irish, *Dracontium foetidum*—c. Skunk, *Dracontium foetidum*—c. Swamp, *Dracontium foetidum*—c. Tree, *Geoffræa inermis*—c. Water, *Nymphæa odorata*.

CABBAGE BARK TREE, *Geoffræa inermis*.

CABBAGIUM, *Geoffræa inermis*.

CABOTZ, *Hagenia Abyssinica*.

CABUREIBA, see *Myroxylon Peruiferum*.

CABUREICIBA, see *Myroxylon Peruiferum*.

CACÆ'MIA, *Cachæ'mia*, (*caco*, and *δύα*, 'blood.') A faulty or morbid condition of the blood.

CACÆSTHE'SIS, *Cacæsthe'sis*, *Cacæsthes'is*, (*caco*, and *αἴσθησις*, 'feeling.') Morbid sensation. Morbid general feeling. Indisposition.

CACAFERRI, *Ferri subcarbonas*.

CACAGOGUE, *Cacago'gus*, *Cac'cagogue*, *Cac'cago'gus*, (from *cacce*, and *αἶψα*, 'to expel.') An ointment, composed of alum and honey; which, when applied to the anus, produced an evacuation.

CACALEXITERIA, (*caco*, and *ἀλεξίω*, 'I ward off.') *Alexiteria*.

CACA'LIA ANTEUPHOR'BIUM, (*κακαλία*.) *Anteuphor'bium*. Ord. *Compositæ*. A plant, which Dodoens and others considered to be capable of tempering the caustic properties of *euphorbium*. It is also called *Klein'ia*.

Many varieties of the *Cacalia* are used, in different countries, chiefly as condiments.

CACANTHRAX, (*caco*, and *anthrax*.) see *Anthrax*.

CA'CAO, *Ca'coa*, *Caca'vi*, *Quahoil*, *Cacava'tu*. The cocoa or chocolate nut; fruit of *Theobroma Ca'cao*, *Co'coa* *Cacarifera*, *Ca'cao mi'nor* seu *sati'va*, *Cacao theobro'ma*, (F.) *Cacaoyer* on *Cacaotier ordinaire*. Family, *Malvaceæ*. Sex. Syst. *Polydelphia Pentandria*.

CACAOTIER ORDINAIRE, see *Cacao*.

CACAOYER ORDINAIRE, see *Cacao*.

CACAPHTHA, (*caco*, and *aphtha*.) see *Aphthæ*.

CACATION, (*cacare*, 'to go to stool.') Defecation.

CACATORIA, *Diarrhœa*.

CAC'ATORY, *Cacato'rius*. *Febris cacato'ria*; a kind of intermittent fever, accompanied by copious alvine evacuations.

CACAVATA, *Cacao*.

CACAVI, *Cacao*, *Jatropha manihot*.

CACCAGOGUE, *Cacagogue*.

CACCE, (*κακῶς*.) Excrement.

CACCION'DE. A sort of pill, chiefly formed of catechu, recommended by Baglivi in dysentery. CACEPHEBOTE'SIA, (*caco*, and *ἐπιβόησις*, 'puberty.') Morbid puberty. Disease occurring at the period of puberty.

CACHALOT, see *Cetaceum*.

CACHANG-PARANG. A sort of bean of Sumatra, mentioned by Marsden, whose seeds are given in pleurisy. Jussieu considered it to be the *Mimo'sa scandens*.

CACHEC'TIC, *Cachec'tes*, *Cachec'ticus*, (F.) *Cachectique*, (same etymon as *Cachexia*.) One attacked with *cachexia*. Belonging to *cachexia*. *Cachectica remedia* are remedies against *cachexia*.

CACHECTIC DISEASES, *Cachectici mor'bi*, are constitutional diseases—gout, scrofula, tuberculous, for example.

CACHELCOMA, (*caco*, and *ἔλκος*, 'ulcer.') *Helcocœce*.

CACHEN-LAGUEN, *Chironia Chilensis*.

CACHET, (F.) ('a seal.') A stamp or medicinal stamp or seal, used by the ancient Romans for marking their drugs, especially those prescribed in diseases of the eyes; and hence called, by some, *oculist stamps*. Some of those more lately discovered amongst Roman antiquities, have been described by MM. Siehel and Duchalais, and by Prof. Simpson, of Edinburgh.

CACHEX'IA, (*κακῆξία*, from *caco*, and *ἔξις*, 'habit.') *Status cachecticus*, *Cachex'y*, *Dys'thesis*, (F.) *Cachexie*. A condition in which the system of nutrition is evidently depraved. A bad habit of body, chiefly the result of scorbutic, cancerous, tuberculous or venereal diseases when in their advanced stage. Hence we hear of a *Scorbutic Cachex'ia*, *Cancerous Cachexia*, &c. Sauvages and Cullen have included under this head a number of diseases—consumptions, dropsies, &c. *Cachexia* has been sometimes confounded with *diathesis*. *Cachexia Icter'ica* is jaundice or icterus itself, or a disposition thereto. Fluor albus is sometimes called *Cachexia Uteri'na*.

CACHEXIA AFRICANA, *Chthonophagia*—c. *Calculosa*, *Lithia*—c. *Cancerous*, see *Cancer*—c. *Chlorotic*, *Chlorosis*—c. *Dysthetica*, *Dyserasia*—c. *Exophthalmica*, see *Exophthalmia*.

CACHEXIA, GAOL. The deteriorated organic actions induced by confinement in prisons, which is so often the precursor of scrofula.

CACHEXIA ICTERICA, *Icterus*.

CACHEXIA LONDINENS'IS. The paleness and

other evidences of impaired health or nutrition presented by the inhabitants of London. A similar cachexia is seen in those of other crowded cities.

CACHEXIA LYMPHATICA FARCINOSA, see Equinia.

CACHEXIA, MARSH, (F.) *Cachexie paludéenne, ou palustre*. The state of cachexy observed in malarious districts.

CACHEXIA SATURNINA, Saturnismus—c. Scorbatic, see Purpura—c. Scrophulosa, see Scrofula.

CACHEXIA SPLE'NICA. The state of scorbutic cachexia, which often accompanies diseases, especially enlargement, of the spleen, *Splenal'gia Bengal'sis*, in India.

CACHEXIA VENEREA, Syphilis—c. Venous, Venosity—c. Virginum, Chlorosis.

CACHEXIE, Cachexia—c. Iodée, see Iodine—c. *Paludéenne*, Cachexia, marsh—c. *Palustre*, Cachexia, marsh.

CACHEXY, Cachexia.

CACHIBOU, see *Bursera gummifera*.

CACHINLAGUA, *Chironia chilensis*.

CACHINNA'TIO, (*cachinno*, 'I laugh aloud.')

A tendency to immoderate laughter, as in some hysterical and maniacal affections.

CACHIRI. A fermented liquor made, in Cayenne, from a decoction of the rasped root of the manioc. It resembles perry.

CACHLEX, (*καχληξ*). A small stone or pebble, found on the sea shore. One of these, when heated in the fire, and cooled in whey, communicates an astringency to the liquid, so that it was anciently esteemed to be useful in dysentery.

CACHOS. An oriental fruit, apparently of a Solanum, which is esteemed lithontriptic.

CACHOU, Catechu.

CACHRYS LIBANO'TIS, (*καειν*, 'to burn'[?].) An umbelliferous plant which grows in Africa and the south of Europe. It is aromatic and astringent. Its seeds are extremely acid.

CACHRYS MARITIMA, *Crithmum maritimum*.

CACHUN'DE. An Indian troch or pastile composed of amber, mastie, musk, cinnamon, aloes, rhubarb, galanga, pearls, rubies, emeralds, garnets, &c. It is regarded by the people of India as an antidote, stomachic and antispasmodic.

CACO, (*κακο*, properly only an abbreviation of *κακος*, 'bad, defective.')

In composition it means something defective; as in the following words:

CACOE'STHESIS, *Cacæsthesi*s.

CACOALEXITERIA, (*caco*, and *αλεξεω*, 'I ward off.')

Alexipharmic.

CACOC'HOL'IA, (*caco*, and *χολη*, 'bile.')

Diseases induced by a depraved condition of the bile.

CAC'OCHROI, *Cac'ochri*, (*κακοχροι*, from *caco*,

and *χρος*, 'colour.')

Diseases in which the complexion is morbidly changed in colour.

CACOC'HYL'IA, (*caco*, and *χυλος*, 'chyle.')

Depraved chylification.

CACOC'HYM'IA (*κακοχυμια*), *Kakochym'ia*, *Corruptio Humo'rum*, (*caco*, and *χυμος*, 'juice, 'humour.')

Cacoch'ymy. Depravation of the humours.

CACOC'HYMIA PLUMBEEA, Lead poisoning—c. Scorbatica, see Purpura—c. Scrophulosa, Scrofula—c. Venerea, Syphilis.

CACOC'HYMUS, (*κακοχυμος*) *Cacochym'icus*. One attacked with cacochymia. Belonging to cacochymia.

CACOC'NE'MUS, *Cacocne'micus*, *Malis Suris præditi'us*; (*caco*, and *κνηση*, 'the leg.')

One who has bad legs.

CACOCORE'MA, gen. *Cacocore'matis*, (*caco*, and *κορεω*, 'I purge, or cleanse.')

A medicine which purges off the vitiated humours.

CACOCY'NANCHE, *Cynanche maligna*.

CACODÆ'MON, (*κακοδαμων*, from *caco*, and

δαμων, 'a spirit.')

An evil spirit, to which many disorders were ascribed. The nightmare.

CACO'DES, (*κακωδης*, from *caco*, and *οζειν*, 'to smell'—*malè olens*.)

Having a bad smell; *Caco'dia*, *Cacos'mia*.

CACODIA, (*κακωδια*), see *Cacodes*.

CACOE'THES, (*κακοηθης*, from *caco*, and *ηθος*,

'disposition, habit.')

Cacoe'thicus, (F.) *Cacoe'the*, Of a bad or vitiated character, as *ulcus cacoe'thes*, an ulcer of a malignant character.

CACOETHICUS, *Cacoe'thes*.

CACOGALAC'TIA, *Cacoga'lia*, (*caco*, and *γαλα*, gen. *γαλακτος*, 'milk.')

A bad condition of the milk.

CACOGALAC'TICA. One who suffers from a bad condition of the milk.

CACOGALIA, *Cacogalactia*.

CACOGEN'ESIS, (F.) *Cacogénèse*, (*caco*, and *γενεσις*, 'generation.')

A morbid or a monstrous formation.

CACOMETRA, (*caco*, and *μητρα*, 'uterus.')

Metrocace.

CACOMORPHIA, (*caco*, and *μορφη*, 'shape.')

Deformation.

CACOMORPHOSIS, Deformation.

CACONYCH'IA, (*caco*, and *ονυχ*, gen. *ονυχος*, 'a nail.')

A morbid condition of a nail.

CACOPARONYCHIA, see *Paronychia*.

CACOPATH'IA, *Pas'sio Mala*, (*κακοπαθεια*, from *caco*, and *παθος*, 'affection.')

A distressed state of mind.

CACOPHO'NIA, (*κακοφωνια*, from *caco*, and *φωνη*,

'voice,' *vitia'ta voc.*)

A dissonant condition of voice.

CACOPLAS'TIC, *Cacoplas'ticus*, *Dysplasmatic'ic*;

(*caco*, and *πλασσω*, 'I form.')

Susceptible of only a low degree of organization, as the indurations

resulting from low or chronic inflammation, fibrocartilage, cirrhosis, &c.

CACOPRA'GIA, *Cacopra'xis*, (*caco*, and *πρασσω*,

'I perform.')

Depraved condition of the organic functions.

CACOPRAXIS, *Cacopragia*.

CACORRHACHI'TIS, (*caco*, and *ραχις*, 'the spine.')

Cacor'rhachis, *Cacor'hachis*, *Cacor'hachi'tis*, *Spondylal'gia*.

Deformity of the spine. Disease of the spine.

Spontaneous luxation of the vertebrae and ribs dependent upon internal causes.

CACORRHYTH'MUS, *Arrhythm'us*, (*caco*, and *ρυθμος*,

'rhythm,' 'order.')

Irregular.

CACO'SIS. *Mala disposit'io*, (F.) *Vice*. A bad condition of body.

A diseased condition in general.

CACOSIT'IA, (*caco*, and *σιτων*, 'aliment.')

Disgust or aversion for food—*Fastid'ium cibo'rum*.

CACOSMIA, (*κακοσμία*), see *Cacodes*.

CACOSOMI'UM, (*caco*, and *σωμα*, 'the body.')

An hospital for leprosy, and incurable affections in general.

CACOSPERMA'SIA, *Cucosperma'tia*, *Cacosper'mia*,

(*caco*, and *σπερμα*, 'sperm.')

A bad condition of the sperm.

CACOSPHYX'IA, (*caco*, and *σφυξις*, 'pulse.')

Vitio'nis pul'sus. Bad state of pulse.

CACOSPLANCH'NIA, (*caco*, and *σπλαγχνον*, 'a viscus.')

Indigestion. The emaciation dependent upon imperfect digestion.

CACOSTOM'ACHUS, (*caco*, and *στομαχος*, 'the stomach.')

What disagrees with the stomach.

CACOSTOMATOSPHRESIA, (*caco*, *στομα*,

'mouth,' and *σφραειν*, 'to smell.')

Breath, offensive.

CACOS'TOMUS, (*caco* and *στομα*, 'a mouth.')

Having a bad mouth, or a bad breath.

CACOTHANASIA, (*caco*, and *θανωσις*, 'death.')

See *Euthanasia*.

CACOTHYML'IA, *Vit'ium An'imi*, (*caco*, and *θυμος*,

'mind,' 'disposition.')

A vicious state of mind.

CACOTRIBULUS, *Centaurea calcitrapa*.

CACOTRICHIA, (*caco*, and *τριχος*, gen. *τριχος*, 'hair.') Disease of the hair.

CACOTROPHIA, (*κακοτροφία*, from *caco*, and *τροφή*, 'nutrition.') *Vitio'sa nutritio*;—disordered nutrition.

CACOU, *Cagot*, Catechu.

CACOU'CIA COCCIN'EA, *Cocuin'ea Coccin'ea*, *Schousboë'a coccin'ea*, *Tikim'ua*. A perennial twining shrub of South America, the plant of which, as well as the fruit, is possessed of emetocathartic properties.

CACTIER, *Cactus opuntia*.

CACTUS, (*κακτος*, (?), *Cynara*).

CACTUS OPUN'TIA, *Opuntia*, *O. vulgaris*. *Ind'ian Fig*, (F.) *Cactier*, *Raquette*, *Figuier d'Inde*. Ord. *Cactaceæ*. This plant grows in South America, Spain, Italy, &c. Its fruit, which has the shape of the fig, is of a sweetish taste, and colours the urine red when eaten. Its leaves are considered refrigerant.

The fruits of different species of cactus are called *Tunas*.

CADA'BA, *Stroë'mia*. A genus of the family *Capparidæ*, natives of India and Arabia. The young shoots of the *Cada'ba furino'sa* are considered to be an antidote against venomous bites.

CADA'VER, *Pto'ma*, *Nec'ros*. A dead body, corpse, carcase, (Sc.) corp; a subject; a carcass, (F.) *Cadavre*. (The word has been supposed to come from *cado*, 'I fall;' and by some to be a contraction from *caro data vermibus*, 'flesh given to the worms.')

CADAVERIC, *Cadaverous*.

CADAVERISA'TION, *Cadaverisatio*. A condition of a part in which, from its paleness, coldness, and loss of vital manifestations, it resembles the dead body.

CADAVEROUS, *Cadav'eric*, *Cadavero'sus*, *Necro'des*, (F.) *Cadavé'reux*, *Cadavérique*. Belonging to the dead body; as *cadaverous smell*. The *Cadav'erous* or *Hippocra'tic face* (see *Face*), is an unfavourable sign in disease, and generally denotes a fatal termination.

CADAV'EROUS or CADAV'ERIC HYPERÆ'MIA. The hypostatic hyperæmia observed in depending parts of the dead body.

CADDY INSECT, see *Ectozaa*.

CADE, *Juniperus oxycedrus*.

CADEJI-INDI, *Malabathrum*.

CADEL-AVANACU, *Croton tiglium*.

CAD'IA. An Egyptian leguminous plant. The Arabs attribute to its fresh leaves the power of relieving colic.

CADIVA INSANIA, *Epilepsy*.

CADMI'A, (*καδμεια* and *καδμια*), *Calamina*, *Tutia*.

CADMI'I SULPHAS, *Cad'mii S.* (Ph. U. S.). *Cadmium sulphuricum*, *Sulphas Cadmicius*, *Meli'ni Sulphas*, *Klapro'thii Sulphas*, *Klapro'thium Sulphuricum*, *Meli'num Sulphuricum*, *Sulphate of Cad'mium*. A formula for its preparation has been introduced into the Ph. U. S. for 1863. It consists in dissolving *cadmium* in *nitric acid*, precipitating by carbonate of soda, dissolving in *sulphuric acid*, and crystallizing.

Used in spots on the cornea, and in chronic torpid inflammation of the conjunctiva, in the quantity of half a grain to a grain to the ounce of water. It does not differ materially in its properties from sulphate of zinc. The nitrate, in the dose of one-eighth of a grain, induces vomiting and purging.

The *Iodide* has been strongly recommended as possessing all the advantages of the preparations of iodine; and in the form of ointment, in the proportion of one part to eight parts of lard, has been used in scrofulous and other tumefactions.

CADMIUM, IODIDE OF, see *Cadmii Sulphas*—c. *Sulphuricum*, *Cadmii Sulphas*.

CADRE DU TYMPAN, ('frame of the tympanum,') *Tympanal*. The portion of the temporal bone which supports the membrana tympani.

CADTCHU, *Catechu*.

CADUCA HUNTERI, (from *cadere*, 'to fall,') *Decidua*—c. *Passio*, *Epilepsy*.

CADU'CITY, *Imbecil'itas*, *Debil'itas*, *Cadu'citas*, (*cadere*, 'to fall,') The French use the word *Caducité* for the portion of human life which is comprised generally between 70 and 80 years. The age which precedes decrepitude. It is so termed in consequence of the limbs not usually possessing sufficient strength to support the body. The precise age must of course vary in individuals.

CADUQUE, *Decidua membrana*—c. *Répléc'ie*, see *Decidua membrana*—c. *Secondaire*, see *Decidua*—c. *Sérotine*, see *Decidua*—c. *Uérine*, *Decidua*—c. *Vraie*, *Decidua membrana*.

CADURCUS, *Vulva*.

CAD'US, (*καδός*,) A Greek measure equal to ten gallons English. *Amphora*.

CÆ'CA FORAM'INA (ANTE'RIUS ET POSTE'RIUS) are situated at the fore and back parts of the tuber annulare of the brain, and at the extremities of the depression made by the vertebral artery. The former is placed between the nerves of the third, and the latter between those of the sixth pair. See *Cæcus*.

CÆ'CÆ HÆMORRHOÏ'DES, *Blind Piles*, (F.) *Hémorrhoides aveugles*, are those unaccompanied by any discharge.

CÆ'CAL, (from *Cæcum*,) *Cæca'lis*. Belonging to the *Cæcum*. The *Cæcal arteries and veins* are the branches of the *Arteriæ et venæ colicæ dextræ inferiores*, distributed to the *Cæcum*. The *Cæcal* or *blind extremity* of a duct is its closed termination. See *Cæcum*.

CÆCATRIX, *Cicatrix*.

CÆ'CITAS, *Cæcitu'do*, (from *cæcus*,) *Ablep'sia*, *Obcæcatio*, *Ocæcatio*, *Anop'sia*, *Typhlotes*, *Typhlo'sis*, *Cæ'city*, *Blind'ness*, (F.) *Aveuglement*, *Cécité*, *Perte de la vue*. *Cæcitas* may be dependent upon many different diseases—as upon amaurosis, specks, hypopyon, cataract, glaucoma, ophthalmia, atrophy of the eye, &c.

CÆCITAS CREPUSCULARIS, *Hemeralopia*—c. *Diurna*, *Nyctalopia*—c. *Nocturna*, *Hemeralopia*.

CÆCITE, *Typhlo-enteritis*.

CÆCITUDO, gen. *Cæcitu'dinis*, *Cæcitas*.

CÆ'CUM, (*cæcus*, 'blind,') *Cæ'cum*, *Intestini cæci*, *Monom'achon*, *Monom'æum*, *Monocæ'lon*, *Monocul'm*, *Typhlot'erum*, *Typhloën'terum*, *Int'ium intest'ni cras'si*, *Sac'cus Intestini cras'si seu Col'i seu Cæ'cus Col'i*, *Cæ'cum Ca'put col'i*, *Ca'put col'i*, *Prima cella coli*, *Int'ium extuberans coli*. The *Blind Gut*, so called from its being open at one end only. That portion of the intestinal canal which is seated between the termination of the ileum and commencement of the colon; and which fills, almost wholly, the right iliac fossa; where the peritoneum retains it immovably. Its length is about three or four fingers' breadth. The *Ileo-cæcal valve* or *Valve of Bauhin* shuts off all communication between it and the ileum; and the *Appendix vermiformis cæci* is attached to it.

CÆCUM FORAMEN of the frontal bone is a small cavity at the inferior extremity of the internal coronal crest or crista.—*Fronto-ethmoidal foramen*, (F.) *Trou aveugle ou borgne*. Morgagni has given the same name to the small cavity in the middle of the upper surface of the tongue, near its base; the sides of which are furnished with mucous follicles—*Lacunes de la langue*—(Ch.)

CÆCUM, PULEGMOUS TUMOUR OF THE, *Typhlo-enteritis*.

CÆ'CUS, ('blind,') One deprived of sight.

Ty'phlops, (F.) *Arengle, Borgue*. In anatomy, it is used to designate certain holes or cavities, which end in a *cul-de-sac*; or have only one opening. See *Cæcal*.

Blind ducts of the Ure'thra, (F.) *Conduits aveugles de l'urèthre*, are the *Mucous Lacu'næ* of the *Ure'thra*.

CÆLA-DOLO, *Torenia Asiatica*.

CÆMENTUM, ('rubble stone,') *Lute*, see *Tooth*.

CÆPA, *Allium cepa*.

CÆRULEUM BEROLINENSE, (*cæruleus*, [from *cælum*, 'heaven,'] 'blue,') *Prussian blue*—*c. Borussicum*, *Prussian blue*.

CÆRULOSIS NEONATORUM, *Cyanopathy*.

CÆSALPINIA, (after *Andreas Cæsalpinus*.) *C. Sap'pan*, *Sap'pan* or *Samp'fen wood*, (F.) *Brésillet*, *Bois de Sappan*. *Ord. Leguminosæ*. A small Siamese tree, the wood of which is used in decoction, in cases of contusion.

Brazil'wood, *Pernambu'co* or *Fernambu'co wood*, formerly used as an astringent; is the wood of *CÆSALPINIA ECHINATA*. This is the proper *Brazil wood*; but another variety in commerce is the *Brasilletto* from *Cæsalp'nia Brasilien'sis* and *C. cris'ta*, which grow in the West Indies.

The *Nicar'gua* or *Peach-wood* is analogous to this, and is said to be derived from a species of *Cæsalpinia*.

The kernel of *CÆSALPINIA BONDUCELLA*, the seed of which is called in India *Kuthuleja* and *Kutoo Kurunja*, is given as a febrifuge tonic. Dose, ten grains.

CÆSAREAN SECTION, *Cæsarean opera'tion*, *Tomoto'cia*, *Cæsareæ sec'tio*, *Par'tus cæsareus*, *Opera'tio cæsareæ*, *Metrotom'ia*, (F.) *Opéra'tion Césarienne*, (*cædere*, *cæsum*, 'to cut'(?).) An incision made through the parietes of the abdomen and uterus to extract the fetus. In this manner, *Julius Cæsar* is said to have been extracted. It is also called *Hysterotom'ia*, *Hysterotocotom'ia*, *Hysterotokotom'ia*, *Gastrometrotom'ia*, *Gasterhysterot'omy*, *Gastrometrot'omy*, *Gastrohysterot'omy*. An incision has been made into the uterus through the vagina, constituting the *Vagi'nal Cæsarean Section*, *Gastroelytrotom'ia*, *Gastroelytrotom'ia*, *Gastrocolpotom'ia*, *Laparacolytrotom'ia*, *Laparocelytrotom'ia*, (F.) *Opération césarienne vaginale*. The Cæsarean section may be required when the mother dies before delivery; when there is some invincible obstacle to delivery from the faulty conformation of the pelvis; or when the child has passed into the abdominal cavity in consequence of rupture of the uterus.

CÆSARIES, ('a head of hair,') *Capillus*, *Scalp*.

CÆSIUS, ('bluish gray,') *Glaucoma*.

CÆSONES, *Cæsares*. Children brought into the world by the Cæsarean operation.

CÆSULIÆ, (*cæsius*, 'bluish gray,') They who have gray eyes.

CÆSURA, (*cædere*, *cæsum*, 'to cut,') *Cut*.

CETCHU, *Catechu*.

CAF, *Camphor*.

CAFAL, *Agrimony*.

CAFAR, *Camphor*.

CAFÉ, *Coffea*.

CAFÉ CITRIN. The aqueous infusion of un-roasted coffee, so called on account of its yellowish tint.

CAFÉ A LA SULTANE. This name has been given to an infusion or decoction of the ground coques or pericarps which surround the coffee.

CAFÉIER and *CAFÉYER*, *Coffea Arabica*.

CAFEIN, see *Coffea Arabica*.

CAFFA, *Camphor*.

CAFEEA, see *Coffea Arabica*.

CAFEIN, see *Coffea Arabica*.

CAFIER, *Coffea Arabica*.

CAFUR, *Camphor*.

CAGAS'TRUM. According to *Paracelsus*, the principal or germ of diseases which are communicable.

CAGNEUX, *Cayot*. See *Kyllosis*.

CAGOSANGA, *Ipecaacuanha*.

CAGOTS (F.), (supposed to be an abbreviation of *Canis Gothus*, 'Dog of a Goth.' [?]) A name given to deformed and miserable beings, met with in the Pyrénées, Bern, and Upper Gascony, in France, where they are also called *Capots*. In other districts, they are called *Gésits*, *Gézitains*, *Crétins*, *Gahets*, *Capons*, *Coliberts*, *Cacous*, *Cagneux*, &c. See *Crétin*.

CAGUESANGUE, *Caguesangué*.

CAHINCÆ RADIX, *Caincæ radix*.

CAI'PUT OIL, *Cai'put oil*, *Kyaput'ty*, *O'leum Cajuputi*. The volatile oil of the leaves of *Melaleuca Cajuputi*, *Ord. Myrtacæ*, a native of the Moluccas. The oil has a strong, fragrant smell, like camphor; taste pungent and aromatic. It is stimulant, and useful where the essential oils in general are employed. It has also been called *Oil of Witneben*, from the person who first distilled it. The British Pharmacopœia (1864) has a *Spir'itus Cajuputi*, *Spir'it of Cajuput*, which consists of *Oil of Cajuput*, f 3j dissolved in *rectified spirit*, f 3ix.

CAIK, *Pleurodynia*.

CAIL-CE'DRA, *Swietenia Senegalensis*.

CAILLE, *Tetrao coturnix*.

CAILLÉ, (from *coagulare*, 'to curdle,') See *Curd*.

CAILLEAU, *Lantana*.

CAILLEBOTTE, see *Curd*.

CAILLELAIT, (*cailler*, 'to curdle,' and *lait*, 'milk,') *Rennet*—*c. Blanc*, *Galium mollugo*—*c. Vraie*, *Galium verum*.

CAILLETTE, (from *cailler*, 'to curdle,') *Abomasus*.

CAILLOT, (from *cailler*, 'to curdle,') *Coagulum*.

CAINANÆ RADIX, *Caincæ radix*.

CAIN'CÆ RA'DIX, *Ra'dix Chiococ'cæ*, *R. Caina'næ* seu *Canina'næ* seu *Cahin'cæ* seu *Kahin'cæ* seu *Serpentariae Brazilien'sis*, *Cain'ca Root*. The bark of the roots of *Chiococ'ca anguif'uga* seu *densifolia*, and, perhaps, *Ch. racemosa*, a plant of the *Order Rubiacæ*. *Sex. Syst.* *Pentandria Monogynia*, of *Linnaeus*. It is bitter, tonic, and diuretic, but has not been long introduced. Dose of the powder, from ʒj to ʒss.

Dr. John H. Griscom, of New York, considers there is a remarkable analogy between the *Cainca* and the *Apocynum camucabinum*.

CAINITO, *Chrysophyllum Cainito*.

CAIPA SCHORA. A cucurbitaceous Malabar plant, the fruit of which has a pyriform shape. The juice is drunk in that country for the purpose of arresting hiccough. The fruit, when unripe, is emetic.

CAISSE, (from *capsa*,) *Case*—*c. du Tambour*, *Tympanum*—*c. du Tympan*, see *Tympanum*.

CAITCHU, *Catechu*.

CAJAN, *Phaseolus creticus*.

CAJUPUTI, *Caieput*.

CAKES, WORM, STORY'S. These were composed of *calomel* and *jalap*, made into cakes, and coloured with *cinnabar*.

CALABASH-TREE, *NARROW-LEAVED*, ([S.] *calabaza*, 'a gourd,') *Crescentia Cujete*.

CALADIUM ESCULENTUM, *Arum esculentum*.

CALADIUM SEGTI'NUM, *Nat. Ord. Aracæ*. A plant of India, whose acrid juice has been given there in gout and rheumatism, and as an anaphrodisiac to women. The tincture has been prescribed in pruritus vulvæ.

CALAF, *Sa'lie Ægypti'aca*. A large-leaved Egyptian willow, called, also, *Ban*. The distilled water of the flowers, called *Mucabalef*, passes, in that country, for an excellent antaphrodisiac. It is also used as an antiloinic, antiseptic, and cordial.

CALAGERI, *Vernonia anthelmintica*.

CALAGIRAH, *Vernonia anthelmintica*.

CALAGUALA, see *Calagualæ radix*.

CALAGUALÆ RADIX, (*καλαρός*, 'a reed.') This word

The root of *Polypodium Calagua'la* seu *adiantum* seu *coriandrum* seu *ammifolium* seu *argenteum* seu *politum*, *Aspidium coriandrum* seu *ferruginum* seu *discolor*, *Tectaria calahualla* seu *ferruginea*, *Calagua'la*, *Calahu'la*. Order, Filices. It has been exhibited in Italy in dropsy, pleurisy, contusions, abscesses, &c. Its properties are not, however, clear.

CALAHUALA, see *Calagualæ radix*.

CALAMANDRINA, *Teucrium chamædryas*.

CALAMBAC, *Agallochum*.

CALAMBOUK, *Agallochum*.

CALAMEDON, (*καλαρός*, 'a reed.') This word has had various significations. Some have used it for an oblique fracture of a bone; the fractured portions having the shape of the nib of a pen. Others have used it for a longitudinal fracture; and others, again, for one that is comminuted.

CALAMENT, *Melissa calamintha*.

CALAMINA, *Cal'amine*, (*calamus*, 'a reed,' so called from its reed-like appearance.) *Cadmi'a*, *C. lapido'sa*, *æro'sa* seu *fos'silis*, *La'pis æro'nus*, *La'pis Calaminaris*, *Calaminaris*, *Carbo'nas Zinci impu'rus*, *Cath'mir*, (F.) *Pierre calaminaire*. Native impure carbonate of zinc. Calamine is chiefly used for pharmaceutical purposes in the form of the *CALAMINA PRÆPARATA* (Ph. U. S.), *La'pis Calaminaris præpara'tus*, *Carbo'nas zinci impu'rus præpara'tus*, *Zinci carbo'nas præpara'tus*, *Præpa'rad Cal'amine*.—Calamine reduced to an impalpable powder by roasting and levigation. In this state it is sprinkled or dusted on excoriated parts, or to prevent excoriation, &c.

CALAMINARIS, *Calamina*.

CALAMINT, *Melissa Calamintha*—c. Field, *Melissa nepeta*—c. Mountain, *Melissa grandiflora*—c. Spotted, *Melissa nepeta*.

CALAMINTA HUMILIOR, *Glechoma hederacea*.

CALAMINTHA, (*καλαμινθη*, from *καλός*, 'beautiful,' and *μινθη*, 'mint,') *Melissa C.*—c. Anglica, *Melissa nepeta*—c. Erecta Virginiana, *Cunila Mariana*—c. Hederacea, *Glechoma hederacea*—c. Magno flore, *Melissa grandiflora*—c. Montana, *Melissa grandiflora*—c. Nepeta, *Melissa nepeta*—c. Parviflora, *Melissa nepeta*—c. Pulegii odore, *Melissa nepeta*—c. Trichotoma, *Melissa nepeta*.

CALAMUS, (*καλαμός*, 'the reed.') In the Pharmacopœia of the U. S., the rhizoma of *acorus calamus*.

CALAMUS ALEXANDRI'NUS. Celsus has thus called a medicine, which was long confounded with *Calamus Aromaticus*. It is not a root, however, but the stalk of a plant of India and Egypt, probably the *Andropogon Nar'dus*. It entered into the theriaca, and has been been regarded as antihysterical and emmenagogue:—*Calamus aromaticus verus*.

CALAMUS AROMATICUS, *Acorus calamus*—c. *Aromaticus verus*, *Calamus Alexandrinus*—c. *Draco*, *C. rotang*—c. *Indicus*, see *Saccharum*—c. *Odoratus*, *Acorus calamus*, *Juncus odoratus*.

CALAMUS ROTANG, *C. Dra'co*. Order, Palmæ. The systematic name of a plant, whence *Dragon's Blood*, *Sa'guis Draco'nis*, *Cinnab'aris Græco'rum*, *Draconthe'ma*, (F.) *Sing-Drang*, is procured. It is the red, resinous juice, obtained in India, from wounding the bark of the *Calamus Rotang*. It has been used as an astringent in hemorrhages, &c.; but is now rarely employed.

CALAMUS SCRIPTOR'IIUS. *Anag'lyphê*, 'a writing-pen,' (F.) *Fosselle angulaire du quatrième ventricule*. A small, angular cavity, situate at the superior extremity of the medulla, in the fourth ventricle of the brain, which has been, by some, supposed to resemble a pen:—see *Centrum ovale*.

CALAMUS VULGARIS, *Acorus calamus*.

CALAPPITE. Rumphius has given this name to calculeous concretions, found in the interior of certain cocoa-nuts. The cocoa-tree itself the Malays called *Calappa*. These stones are, likewise, termed *Veg'etabile Bez'oards*. The Malays attribute potent virtues to them, and wear them as amulets.

CALASAYA, *Cinchonæ cordifoliæ cortex*.

CALBALA, *Cabal*.

CALBIANUM. The name of a plaster in Myrepsus, the composition of which we know not.

CALCADINUM, *Ferri sulphas*.

CALCAIRE, (*calx*, 'lime,') *Calcareons*.

CALCA'NEAL, *Calca'neus*, (*calx*, 'the heel,') Having relation to the calcaneum, as '*calcaneal arteries*.'

CALCANEO-PHALANGIEN DU PETIT ORTEIL, *Abductor minimi digiti pedis*—c. *Phalanginien commun*, *Extensor brevis digitorum pedis*—c. *Sous-phalangettien commun*, *Flexor brevis digitorum pedis*—c. *Sous-phalangien du petit orteil*, see *Abductor minimi digiti pedis*—c. *Sous-Phalanginien commun*, *Flexor brevis digitorum pedis*—c. *Sus-phalangettien commun*, *Extensor brevis digitorum pedis*.

CALCA'NEUM, (*calx*, gen. *calcis*, 'the heel,') *Calca'neus*, *Cal'car*, *Cal'cia*, *Ich'nus*, *Os Cal'cis*, *Pter'na*, *Pter'nium*. The largest of the tarsal bones: that which forms the heel. It is situate at the posterior and inferior part of the foot; is articulated above and a little anteriorly with the astragalus; anteriorly, also, with the os cuboides. Its posterior surface—called *Heel*, *Pa'lus*, *Calx*, (F.) *Talon*—gives attachment to the tendo-achillis: the lower has, posteriorly, two tuberosities, to which the superficial muscles of the sole of the foot are attached. The *small Apoph'ysis* or *lateral Apoph'ysis* of the *Calca'neum*, (F.) *Petit Apophyse* ou *Apophyse latérale du Calca'neum*, is a projection at the upper surface of this bone, on which is formed the posterior portion of the cavity that receives the astragalus. The *great Apoph'ysis*, *anterior Apoph'ysis* of the *Calca'neum*, is the projection which corresponds, on one side, with the cuboides; and on the other forms the anterior part of the facette which receives the astragalus.

CALCAR, gen. *Cal'caris*, *Calcaneum*; (also, 'a spur,') *Ergot*—c. *Avis*, *Hippocampus minor*.

CALCA'REOUS, *Calca'reus*, *Calca'rius*, (*calx*, 'lime,') (F.) *Calcaire*. Containing lime:—as *calcareous concretions*, *C. depositions*, &c.

CALCAREUS CARBONAS, *Creta*.

CALCARIA CARBONICA, see *Creta*—a *Chlorata*, *Calcis Chloridum*—c. *Chlorica*, *Calcis chloridum*—c. *Chlorinea*, *Calcis chloridum*—c. *Hypochlorosa*, *Calcis chloridum*—c. *Hypophosphoria*, *Calcis hypophosphis*—c. *Phosphoria*, see *Cornu cervi*—c. *Pura*, *Calx*—c. *Pura liquida*, *Liquor calcis*.

CALCARIÆ CHLORUM, *Calcis chloridum*—c. *Hypophosphis*, *Calcis hypophosphis*.

CALCATAR, *Ferri sulphas*.

CALCATREPPOLA, *Centaurea calcitrapa*.

CALCENA, **CALCE'NON**, **CALCENO'NIA**, **CALCINO'NIA**. Words employed by Paracelsus to designate the concretions of tartrate of lime which form in the human body.

CALCENOS, *Calcetus*.

CALCEOLA'RIA, (*calceolus*, 'a small slipper,') *Slipperwort*.

CALCEOLARIA PRIMATA is used in Peru as a laxative.

CALCEOLARIA TRIF'IDA is esteemed to be febrifuge.

CALCE'TUS, *Calcenot'nus*, *Cal'cenos*. That which abounds in tartrate of lime. An adjective used by Paracelsus in speaking of the blood; *Sanguis calce'tus*. Hence came the expression, *Cal'cined blood*, *Sang calciné*.

CALCEUM EQUINUM, ('horseshoe,') *Tussilago*.

CALCIA, *Calcanemum*.

CALCIFEROUS CANALS, (*calx*, gen. *calcis*, and *fero*, 'to carry,') *Canaliculi calciferi*.

CALCIFICATION, (*calx*, gen. *calcis*, and *fio*, 'to become,') see *Cretefaction*.

CALCIF'RAGA, (*calx*, gen. *calcis*, 'stone,' and *frangere*, 'to break,') from its supposed power of breaking stone in the bladder. The *Scolopendrium*, (asplenium,) according to Scribonius Largus. The plant of Pliny is supposed to be *Globularia alatum*. Used at times for *Sarcifraga*.

CALCIGEROUS CELL, (*calx*, gen. *calcis*, and *gero*, 'to bear,') see *Tooth*.

CALCIG'RADUS, *Pternob'ates*, (*calx*, gen. *calcis*, 'the heel,' and *gradus*, 'a step,') One who walks on his heels.

CALCII CHLORIDUM, *Calcis murias*—c. *Chloruretum*, *Calcis murias*—c. *Oxychloruretum*, *Calcis chloridum*—c. *Oxydum*, *Calx viva*—c. *Protochloruretum*, *Calcis chloridum*—c. *Sulphuretum*, *Calcis sulphuretum*.

CALCINA'TION, *Calcina'tio*, *Calci'non*, *Concrema'tio*, (*calx*, gen. *calcis*, 'lime,') The act of submitting to a strong heat any infusible mineral substance, which we are desirous of depriving either of its water, or of any other volatilizable substance that enters into its composition; or which we wish to combine with oxygen. *Alum* is calcined to get rid of its water of crystallization;—*chalk*, to reduce it to the state of pure lime, by driving off the carbonic acid; and *certain metals* are subjected to this operation to oxidize them.

CALCINATUM MAJUS POTERII, *Hydrargyrum præcipitatum*.

CALCINONIA, *Calцена*.

CALCIS (gen. of *calx*) BICHLORURETUM, *Calcis chloridum*—c. *Carbonas*, *Creta*—c. *Carbonas durus*, *Creta*, *Marmor*—c. *Carbonas friabilis*, *Creta*.

CALCIS CARBO'NAS PRÆCIPITA'TUS seu PRÆCIPITA'TA, *Præcipitated Car'bonate of Lime*, *Præcipitated Chalk*. Prepared by mixing hot solutions, in distilled water, of carbonate of soda and chloride of calcium; separating the precipitate; washing it with boiling distilled water until the washings cease to be affected by a solution of nitrate of silver, and drying the precipitate. It has the same properties as *creta præparata*, and is preferred to it in certain cases,—for example, as an ingredient in tooth powders, owing to its freedom from gritty particles.

CALCIS CHLO'RIDUM, *Chlo'ride* or *Chlo'ruret* or *Hypochlo'rite* or *Chlo'rite* or *Oxymuri'ate of Lime*, *Calx chlorina'ta* (Ph. U. S.) seu *Chlora'ta* (Ph. B. 1864) seu *Oxymuriat'ica*, *Protorichlor'uret of Calcium*, *Calca'ria chlora'ta* seu *Chlo'rica* seu *Oxymuriat'ica* seu *Hypochloro'sa* seu *Chlorin'ica*, *Subchlo'ris cal'cieus*, *Chlo'rurum seu Chlo're'tum Calca'rie*, *Oxychlorure'tum* seu *Protochlorure'tum Cal'cii*, *Chlorure'tum Ox'idi Cal'cii*, *Bichlorure'tum seu Oxymuri'as Cal'cis*, *Cal'cis Hypochlo'ris*, *Bleach'ing* or *Ten'nant's Pow'der*, (F.) *Protoxichlorure de Calcium*, *Chlorure d'Oxide de Calcium*, *Chlorure ou Oxichlorure* ou *Bichlorure* ou *Oxymuriate* ou *Muriate suroxygéné* ou *oxygéné de Chaux*, *Poudre de Blanchement*, *P. de Tennant*. A compound resulting from the action of chlorine on hydrate of lime. Chloride of lime is a most valuable disinfecting agent, (see *Disinfection*,) when dissolved in the proportion of one pound to six gallons of water. It has likewise been employed both inter-

nally and externally in various diseases, as in scrofula, fætor oris, foul ulcers, &c. &c.

CALCIS HEPAR, *Calcis sulphuretum*—c. *Hydras*, see *Calx*—c. *Hypochloris*, *Calcis chloridum*.

CALCIS HYPOPHOS'PHIS, *Calca'ria hypophosphur'ica*, *Hypophosph'us Calca'rie*, *Hypophosphite of Lime*; obtained by the union of phosphorus with milk of lime, and long boiling, during which phosphuretted hydrogen gas is given off; purifying and evaporating. Highly recommended as a "specific" by Dr. J. F. Churchill; but without foundation. Dose, one to five grains in syrup. The *Hypophosphites of soda*, *potassa*, and *ammonia*, and of *iron*, *quinia*, and *manganese*, have likewise been used.

CALCIS MU'RIAS, *Mu'riate of Lime*, *Calx sali'ta*, *Cal'cii Chlorure'tum* seu *Chlo'r'idum*, *Chlo'ride of cal'cium*, (F.) *Chlorure de calcium*, *Muriate ou Hydrochlorate de Chaux*. This salt has been given, in solution, as a tonic, stimulant, &c., in scrofulous tumours, glandular obstructions, general debility, &c. A *Solutio Muriat'is Cal'cis*, *Liquor Cal'cis Muriat'is*, *Solutio of Muriate of Lime*, *Liquid Shell*, may be formed of *Muriate of Lime*, 3j; dissolved in distilled water, f 3ij. The LIQUOR CAL'CI CHLO'RIDI or *Solutio of Chloride of Calcium*, of the Pharmacopœia of the United States, is prepared as follows;—*Marble*, in fragments, 3vj; *Muriatic acid*, 3xij; *Distilled water*, a sufficient quantity. Mix the acid with a half pint of the water, and gradually add the marble. Towards the close of the effervescence apply a gentle heat, and, when the action has ceased, pour off the clear liquor and evaporate to dryness. Dissolve the residuum in its weight and a half of distilled water, and filter. Dose, from grt. xxx to f 3j, in a cupful of water.

CALCIS OXYMURIAS, *Calcis chloridum*—c. *Phosphas*, see *Cornu cervi*—c. *Phosphas præcipitata*, see *Cornu cervi*.

CALCIS SULPHURE'TUM, *Cal'cii sulphure'tum*, *He'par Cal'cis* seu *Sulphuris calca'reum*, *Sulphuret of Lime*, *S. of Cal'cium*, *Hydrosulphate of Lime*, (F.) *Proto-hydrosulfate de Calcium*, *Hydrosulfate de chaux*, *Foie de soufre calcareux*. Principally used in solution, as a bath, in itch and other cutaneous affections.

CALCITEA, *Ferri Sulphas*.

CALCITEOSA, *Plumbi oxydum semivitreum*.

CALCITHOS, *Cupri subacetas*.

CALCITRAPA, *Centaurea Calcitrapa*, *Delphinium consolida*—c. *Hippophæstum*, *Centaurea calcitrapa*—c. *Stellata*, *Centaurea calcitrapa*.

CALCITRAPPE, *Centaurea calcitrapa*.

CALCIUM, CHLORIDE OF, *Calcis murias*—c. *Chlorure de, Calcis murias*—c. *Chlorure d'oxide de, Calcis chloridum*—c. *Oxydatum*, *Calx viva*—c. *Oxydatum hydratum*, see *Calx*—c. *Protohydrosulfate de, Calcis sulphuretum*—c. *Protoxichlorure de, Calcis chloridum*—c. *Protoxichloruret of, Calcis chloridum*—c. *Protoxide of, Calx*—c. *Sulphuret of, Calcis sulphuretum*.

CALCOSUBPHALANGÆUS MINIMI DIGITI, *Abductor minimi digiti pedis*—c. *Subphalangis pollicis*, *Abductor pollicis pedis*.

CALCOCOS, *Bell-metal*.

CALCOTAR, *Ferri sulphas*.

CALCUL, *Calculus*—c. *Chatonné*, see *Calcul*.

CALCULEUX, *Calculous*.

CALCULI, pl. of *Calculus*, (dim. of *calx*, 'a stone,') *Lithi*, *Lap'ides*, (F.) *Calculus*, *Piérres*. Calculi are concretions which may form in every part of the animal body, but are most frequently found in the organs that act as reservoirs, and in the excretory canals. They are met with in the tonsils, joints, biliary ducts, digestive passages, lachrymal ducts, mammae, pancreas, pineal gland, prostate, lungs, salivary, spermatic and urinary passages, and in the uterus. The causes which give rise to them are obscure.

Those that occur in reservoirs or ducts are supposed to be owing to the deposition of the substances, which compose them, from the fluid as it passes along the duct; and those which occur in the substance of an organ are regarded as the product of some nutritive irritation. Their general effect is to irritate, as extraneous bodies, the parts with which they are in contact; and to produce retention of the fluid, whence they have been formed. The symptoms differ, according to the sensibility of the organ and the importance of the particular secretion whose discharge they impede. Their solution is generally impracticable: spontaneous expulsion or extraction is the only way of getting rid of them.

CALCULI ALTERNATING, see Calculi, urinary—c. Articular, see Calculi, arthritic, and Concretions, articular.

CALCULI ARTHRITIC, *To'phi*, *Tuber'cula arthrit'ica*, *Calculi arthrit'ici seu podagr'ici*, *Chalk-stones*, *Gout-stones*, *Nodes*, (F.) *Pierres crayeuses*, *Calculs arthritiques*, *Nœuds*. Concretions, which form in the ligaments, and within the capsules of the joints, in persons affected with gout. They are composed of uric acid, soda, and a little animal matter; very rarely, urate of lime and chloride of sodium are met with. Similar calculi are found in other parts besides the joints.

CALCULI BILIARY, *Calculi bilio'si seu fel'lei seu bilit'arii*, *Chololithus*, *Cholel'ithus*, *Bil'ary Concretions*, *Gall-stones*, (F.) *Calculs biliaires*, *Pierres au fiel*, *Angichololithes* of Piorry. Some of these contain all the materials of the bile, and seem to be nothing more than that secretion thickened. The greater part are composed of from 88 to 94 parts of *Cholesterin*, and of from 6 to 12 of the yellow matter of the bile. Biliary calculi are most frequently found in the gall-bladder: at other times, in the substance of the liver, in the branches of the *Ductus hepaticus*, or in the *Ductus Communis Choleodochus*. The first are called *Cystic*; the second *Hepatic*: and the last, sometimes, *Hepatocystic*. The causes which give rise to them are very obscure. When quiescent they often occasion no uneasiness. At times, they are rejected by the mouth, or by the bowels, along with a considerable quantity of bile, which had accumulated behind them; at other times they occasion violent abdominal inflammation, abscesses, and biliary fistulæ, rupture of the gall-bladder, and fatal effusion into the peritoneum. The passage of a gall-stone is extremely painful; yet the pulse is not at first affected. Antiphlogistics, when there is inflammatory action, and anæsthetics, and strong doses of opium, to allay the pain and spasm, with the warm bath, are the chief remedies. Solvents are not to be depended upon. They cannot reach the calculi.

CALCULI BONE EARTH, see Calculi, urinary—c. Compound, see Calculi, urinary—c. Cystic, see Calculi, urinary.

CALCULI OF THE EARS, (F.) *Calculs de l'Oreille*. Hard, light, and inflammable concretions, which occur in the *meatus auditorius externus*, and are merely indurated cerumen. They are a frequent cause of deafness. They can be easily seen, and may be extracted by appropriate forceps, after having been softened by injections of soap and water.

CALCULI FELLEI, Calculi, biliary—c. Fusible, see Calculi, urinary.

CALCULI LACHRYMAL, (F.) *Calculs lacrymanæ*. Concretions sometimes, but rarely, form in the lachrymal passages, where they occasion abscess and fistulæ, which do not heal until they are extracted.

CALCULI LITHIC, see Calculi, urinary.

CALCULI OF THE MAMMÆ, (F.) *Calculs des Mamelles*. Haller gives a case of a concretion, of a yellowish white colour, which had the shape

of one of the excretory ducts of the mammary gland, having been extracted from an abscess seated in that organ.

CALCULI MULBERRY, see Calculi, urinary.

CALCULI OF THE PANCREAS, (F.) *Calculs du Pancréas*. These are but little known. Analogy has induced a belief that they resemble the salivary. Some have supposed that certain transparent calculi, rejected by vomiting, or passed in the evacuations, have proceeded from the pancreas, but there seems to be no sufficient reason for this belief.

CALCULI OF THE PINÆAL GLAND, (F.) *Calculi de la Glande Pinéale*. Sabulous concretion are usual in the gland. No phenomena announce their presence during life. They are composed of phosphate of lime.

CALCULI PODAGRICI. Calculi, arthritic.

CALCULI OF THE PROSTATE. *Prostat'ic calculi*. These are not very rare. They have generally the same composition as the preceding. They usually present the symptoms common to every tumefaction of the prostate, and sometimes those of calculi in the bladder.

CALCULI PULMONARY, (F.) *Calculs pulmonaires*. These concretions are very frequently met with in the dead body, without seeming to have produced unpleasant symptoms during life. At other times, they are accompanied with all the symptoms of phthisis, *Lithophth'isis*, *Phthisis calculeuse*, of Bayle. See *Lithiasis pulmonum*. At times they are expectorated without the supervention of any unpleasant symptom. They are usually formed mainly of carbonate of lime and animal matter.

CALCULI SALIVARY, *Calculi Salivales*, *Sialolithi*, (F.) *Calculs salivaires*. Concretions, usually formed of phosphate and carbonate of lime and animal matter, which are developed in the substance of the salivary glands or in their excretory ducts. In the first case, they may be mistaken for a simple swelling of the gland; in the second, they may generally be detected by the touch. They may be extracted by incision in the interior of the mouth. The calculus developed in the sublingual ducts has been called *Calculus sublingualis* and *Ran'ula lapide'a*.

CALCULI SPERMATIC, (F.) *Calculs spermaticques*. These have been sometimes found in the vesiculæ seminales after death. They cannot be detected during life.

CALCULI OF THE STOMACH AND INTESTINES, *Enterolithus*, *E. Calculus*, *Copro'ithus*, *Concretio'nes alvi'nae*, (F.) *Calculs de l'estomac*, *C. intestinaux* ou *stercoraux*, *Pierres stercorales*, *Concretions intestinales*. Calculi of the stomach are rare, and have almost always been carried thither by the antiperistaltic action of the intestines. The symptoms occasioned by them are those of chronic gastritis. It has been imagined that the continued use of absorbent powders, as magnesia, will give occasion to them.

Intestinal concretions, (F.) *Calculs intestinaux*, are not uncommon in animals (see *BEZOARD*): but they are rare in man. The causes which give rise to them are little known: sometimes a biliary calculus affords them a nucleus. Their composition varies. They are light, hard, very fetid, and are formed, ordinarily, between the valvulæ of the small intestines, or in the cells of the large, and sometimes in old herniæ. Whilst they do not obstruct the passage of the alimentary mass, they produce no unpleasant symptoms. At times, the movable tumour which they form may be felt through the parietes of the abdomen. They are generally evacuated *per anum*.

CALCULI OF THE TONSILS. Calculous concretions, which sometimes form in the tonsils. (F.) *Calculs des Amygdales*. They are easily recognised by the sight and touch: sometimes they

are discharged by spitting, either alone or with the pus of an abscess occasioned by their presence.

CALCULI, TRIPLE, see *Calculi, urinary*—c. *Uric*, see *Calculi, urinary*.

CALCULI, URINARY, *Urolithi*, (F.) *Calculs urinaires*, *Pierres urinaires*. Concretions which form from the crystallizable substances in the urine, and are met with not only in the whole course of the urinary passages, but in fistulous openings wherever the urine stagnates naturally or accidentally. Their causes are but little known. They are more common at the two extremities of life than at the middle, and more so in some countries and districts than in others.

At times, a clot of blood, a portion of mucus, &c., form the nucleus. The symptoms and treatment vary according to the seat of the calculus. There is no such thing, probably, as a medical solvent. See *Urinary Calculi*.

Chymists have demonstrated the existence of several components of urinary calculi, viz., *Lithic Acid* and *lithates*, *Phosphate of Lime*, *Ammoniac-Magnesian Phosphate*, *Oxalate of Lime*, *Cystic Oxide*, and *Xanthic Oxide*, with an animal cementing ingredient. The varieties of calculi, produced by the combination or intermixture of these ingredients, were thus represented by Dr. Paris.

A TABULAR VIEW OF DIFFERENT SPECIES OF URINARY CALCULI.

SPECIES OF CALCULI.	EXTERNAL CHARACTERS.	CHYMICAL COMPOSITION.	REMARKS.
1. LITHIC OR URIC.	Form, a flattened oval. S. G. generally exceeds 1.500. Colour, brownish or fawn-like. Surface, smooth. Texture, laminated.	It consists principally of <i>Lithic Acid</i> . When treated with nitric acid, a beautiful pink substance results. This calculus is slightly soluble in water, abundantly so in the pure alkalies.	It is the prevailing species; but the surface sometimes occurs finely tuberculated. It frequently constitutes the nuclei of the other species.
2. MULBERRY, (F.) <i>Calculs muraux</i> .	Colour, dark brown. Texture, harder than that of the other species. S. G. from 1.428 to 1.976. Surface, studded with tubercles.	It is <i>oxalate of lime</i> , and is decomposed in the flame of a spirit lamp swelling out into a white efflorescence, which is <i>quicklime</i> .	This species includes some varieties, which are remarkably smooth and pale-coloured, resembling <i>hempseed</i> .
3. BONE EARTH.	Colour, pale brown or gray; surface, smooth and polished; structure, regularly laminated; the laminae easily separating into concrete crusts.	Principally <i>phosphate of lime</i> . It is soluble in muriatic acid.	
4. TRIPLE.	Colour, generally brilliant white. Surface, uneven, studded with shining crystals, less compact than the preceding species. Between its laminae small cells occur, filled with sparkling particles.	It is an <i>ammoniac-magnesian phosphate</i> , generally mixed with phosphate of lime. Pure alkalies decompose it, extracting its ammonia.	This species attains a larger size than any of the others.
5. FUSIBLE.	Colour, grayish white.	A compound of the two foregoing species.	It is very fusible, melting into a vitreous globule.
6. CYSTIC.	Very like the triple calculus, but it is unstratified and more compact and homogeneous.	It consists of <i>cystic oxide</i> . Under the blowpipe it yields a peculiarly fetid odour. It is soluble in acids, and in alkalies, even if they are fully saturated with carbonic acid.	It is a rare species.
7. ALTERNATING.	Its section exhibits different concentric laminae.	Compound of several species, alternating with each other.	
8. COMPOUND.	No characteristic form.	The ingredients are separable only by chymical analysis.	

1. *Renal Calculi*, (F.) *Calculs rénaux*. These have almost always a very irregular shape: at times, there is no indication of their presence: at others, they occasion attacks of pain in the kidneys, sometimes accompanied by bloody or turbid urine. Often, they cause inflammation of the kidneys, with all its unpleasant results. They are generally formed of uric acid, or oxalate of lime, and at times, phosphates with animal matter. The treatment will have to vary, according to the presumed constituents and to the absence or presence of inflammatory signs,—relieving the irritation by opiates. A surgical operation can rarely be applicable.

2. *Calculi of the Ureters*, (F.) *Calculs des Urètres*. These come from the kidneys, and do not produce unpleasant effects, unless they are so large as to obstruct the course of the urine, and to occasion distension of the whole of the ureters above them; or unless their surface is so rough as to irritate the mucous membrane, and occasion pain, hemorrhage, abscesses, &c. The pain, during the passage, is very violent, extend-

ing, at times, to the testicles of the same side in the male; and occasioning a numbness of the thigh in both sexes. The treatment consists in general or local blood-letting, warm bath, and opiates.

3. *Vesical Calculi*; *Stone in the Bladder*, *Lithia vesicalis*, *Lithiasis cystica* seu *vesicalis*, *Cystolithiasis*, *Dysuria calculosa* seu *irritata*, *Calculus vesicæ*, (F.) *Calculs vésicaux*. These are the most common. Sometimes, they proceed from the kidneys; most commonly, they are formed in the bladder itself. Sense of weight in the perinaeum, and sometimes of a body rolling when the patient changes his position; pain or itching at the extremity of the glans in men; frequent desire to pass the urine; sudden stoppage to its flow; and bloody urine—are the chief phenomena, that induce a suspicion of their existence. We cannot, however, be certain of this without sounding the patient. Sometimes, when of a small size, they are expelled: most commonly, they remain in the bladder, the disorganization of which they occasion, unless removed by a surgical operation.

4. *Urethral Calculi*. These almost always proceed from the bladder. The obstruction, which they cause to the passage of the urine, the hard tumour, and the noise occasioned when struck by a sound, indicate their presence. They are removed by incision. They form, at times, in the prostate gland, *Prostatolithi*.

5. *Calculi of Fistulous Passages*. These arise when there is some fistulous opening into the urethra. They can be readily recognised, and may generally be extracted with facility. (F.) *Calculs placés hors des voies urinaires*. See *Urinary Calculi*.

CALCULI OF THE UTERUS, (F.) *Calculs de l'Utérus*. These are very rare. The signs, which indicate them during life, are those of chronic engorgement of the uterus. Their existence, consequently, cannot be proved till after death.

CALCULIFRAGUS, (*calculus* and *frangere*, 'to break,') *Lithonptic*.

CAL'CULOUS (F.), *Calculeux*, *Graveleux*. That which relates to calculi, especially to those of the bladder.

CALCULOUS OXYD, *Oxyd*, *cystic*.

CALCULUS (pl. of *Calculus*.) *Calculi*. *Calculs biliaires*, *Calculi*, biliary—*c. de l'Estomac*, *Calculi* of the stomach—*c. de la Glande Pinéale*, *Calculi* of the pineal gland—*c. Intestinaux*, *Calculi* of the stomach and intestines—*c. Lacrymaux*, *Calculi*, lacrymal—*c. des Mamelles*, *Calculi* of the mammae—*c. Mûraux*, see *Calculi*, urinary—*c. de l'Oreille*, *Calculi* in the ears—*c. du Pancréas*, *Calculi* of the Pancreas—*c. Placés hors des voies urinaires*, *Calculi* of fistulous passages—*c. Pulmonaires*, *Calculi*, pulmonary—*c. Réniaux*, *Calculi*, renal—*c. Salivaires*, *Calculi*, salivary—*c. Spermatiques*, *Calculi*, spermatic—*c. Stercoraux*, see *Calculi* of the stomach and intestines—*c. des Urèdres*, *Calculi* of the ureters—*c. Urinaires*, *Calculi*, urinary—*c. de l'Utérus*, *Calculi* of the uterus—*c. Vésicaux*, *Calculi*, vesical.

CALCULUS, see *Calculi*—*c. Bezoar*, *Bezoard*—*c. Dentalis*, *Odontolithus*—*c. Encysted*, *Calculus chatonné*—*c. Sublingualis*, see *Calculi*, salivary—*c. Vesicæ*, see *Calculi*, vesical (urinary).

CALDARIUM, (from *calidus*, 'hot,') See *Stove*.

CALDAS, WA'TERS OF. *Caldas* is a small town, ten leagues from Lisbon, where are mineral springs, containing carbonic and hydrosulphuric acid gases, carbonates and muriates of lime and magnesia, sulphates of soda and lime, sulphuret of iron, silica, and alumina. They are much used in atonic gout. They are thermal. Temperature 93° Fahrenheit.

CALDERIÆ ITALICÆ. Warm baths in the neighbourhood of Ferrara, in Italy, much employed in dysuria.

CALEBASSES, *Cucurbita lagenaria*.

CALEDONIA SPRINGS. These are in Pennsylvania, fifteen miles from Chambersburg. The waters are pure; temperature 52° Fah.

CALEDONIA SPRINGS, see *Canada*, *Mineral Waters of*.

CALEFACIENTS, *Calefacientia*, *Therman'tica*, (*calidus*, 'warm,' and *facio*, 'I make,') (F.) *Échauffants*. Substances which excite a degree of warmth in the part to which they are applied, as mustard, pepper, &c. They belong to the class of stimulants.

CALEFACTIO, *Échauffement*.

CALENDULA ALPINA, (called from flowering every *calend.*) *Arnica montana*.

CALENDULA ARVEN'SIS, *Cal'tha Arven'sis* seu *officina'lis*, *Wild Mar'igold*, (F.) *Souci des Champs*. *Order*, *Compositæ*. This is, sometimes, preferred to the next. Its juice has been given, in the dose of $\text{f}\frac{3}{4}$ to $\text{f}\frac{3}{4}\text{iv}$, in jaundice and cachexia.

CALENDULA OFFICINA' LIS, *C. Sati'va*, *Chrysanthemum*, *Spon'sa so'lis*, *Cal'tha vulga'ris*; *Ver-ruca'ria*, *Sin'gle* or *Gar'den Mar'igold*, (Prov.) *Gold'ing*, (F.) *Souci*, *S. ordinaire*, *Fleur de tous les mois*. *Family*, *Synanthereæ*, *Syngenesia necessaria*, *Linn*. The flowers and leaves have been exhibited as aperients, diaphoretics, &c., and have been highly extolled in cancer.

CALENDULÆ MARTIALES, *Ferrum ammoniatum*.

CALENTU'RA, *Cal'enture*, (*calere*, 'to be warm,') The word, in Spanish, signifies fever. A species of furious delirium to which sailors are subject in the torrid zone;—a kind of phrenitis, the attack of which comes on suddenly after a broiling day, and seems to be characterized by a desire in the patient to throw himself into the sea. It is only a variety of phrenitis.

CALENTURA CONTINUA, *Synocha*.

CALENTU'RAS; *Palo de Calenturas*. *Pomet* and *Léméry* say, that these words are sometimes applied to cinchona. *Camelli* says, they mean, also, a tree of the Philippine Isles, the wood of which is bitter and febrifuge.

CALF KILL, *Kalmia latifolia*—*c. Knee*, *Entogonyancon*—*c. of the Leg*, *Sura*.

CALICE, *Calix*.

CALICES RÉNALES, see *Calix*.

CALICO BUSH, *Kalmia latifolia*.

CALIDARIUM, *Stove*.

CALIDUM ANIMALE, (*calidus*, 'warm,') *Animal heat*—*c. Innatum*, *Animal heat*.

CALIFOR'NIA, MIN'ERAL WA'TERS OF. Several mineral springs exist in California, near 40° N. Lat. and 40½° longitude west of Washington, directly upon the California trail, leading from the sink of Humboldt river to Pyramid lake. Their temperature varies from cold to almost boiling. They ebb and flow at irregular intervals; have not been analyzed; but some have the taste of a strong solution of sulphate of magnesia. They have had the singular effect of inducing blennorrhœal symptoms, both in man and animals.

CALIGATIO, (from *caligo*,) *Dazzling*.

CALIGO, gen. *Calig'inis*, ('a mist,') *Ach'lys*, (F.) *Brouillard*. An obscurity of vision, dependent upon a speck on the cornea; also, the speck itself; *Cal'igo cor'næ* seu *d' Nephel'io* seu *d' Leuco'matê*, *Mac'ula cor'næ* seu *semipellucida*, *Phtha'rma cali'go*, *Hebetu'do vi'sûs*, *Neb'ula*, *Opake' cor'nea*, *Web-eye* (?), *Web* (?), *Pin-and-Web* (?), (F.) *Nuage de la cornée*, *Taie* ou *Taye*, *Obscurissement de la vue*, *Akératodiaphanie*.

CALIGO LENTIS, *Cataract*—*c. Pupillæ*, *Syne-zisis*—*c. Synizesis*, *Syne-zisis*—*c. Tenebrarum*, *Hemeralopia*.

CALIHACHA CANELLA, *Laurus cassia*.

CALISAYA BARK, (etymon not clear,) *Cinchonæ cordifoliæ* cortex.

CALISTHEN'ICS, *Callisthen'ics*, *Calisthen'i'a*, *Callisthen'i'a*, (F.) *Callisthénie*, (*καλός*, 'beautiful,' and *σθένος*, 'strength,') The art of promoting, by appropriate exercises, strength of body, and grace of movements.

CAL'IX, gen. *Cal'icis*, *Cal'yx*, *Infundibulum*, (*καλὺς*, 'a cup,' 'a flower cup,') (F.) *Calice*, *Euton-noir*. Anatomists have given this name to small cuplike membranous canals, which surround the papillæ of the kidney, and open into its pelvis, whither they convey the urine:—*Cal'ices ren'al'es*, *Cylind'ri membra'næ rei re'num*, *Fistulæ ure'terum renum*, *Canal'es membra'næ rei re'num*, *Tubuli pelvis renum*. Their number varies from 6 to 12 in each kidney.

CALIX VOMITORIUS, *Goblet*, *emetic*.

CAL'LA PALUS'TRIS, *Wal'ter A'rums*; *indig-nous*. *Ord. Aracæ*: has similar virtues to those of the *Arums*.

CALLA VIRGINICA, *Peltandra Virginiaica*.

CALLEUX, Callous.

CALLIBLEPH'ARUM, (καλλος, 'beauty,' and βλεφαρον, 'eyelid.') A remedy for beautifying the eyelids.

CALLICANTHUS, Calycanthus.

CALLICAR'PA AMERICA'NA, (καλλος, 'beauty,' and καρπος, 'fruit.') *French Mul'berry*; indigenous. Order, Verbenaceæ. Used as a eutrophic in cutaneous diseases. Said to be useful in dropsy.

CALLICOCCA IPECACUANHA, (καλλος, 'beauty,' and κοκκος, 'berry.') *Ipecacuanha*.

CALLIGREAS, (καλλος, 'beauty,' and κρεας, 'flesh.') Pancreas.

CALLIOMARCHUS, Tussilago.

CALLIP'E'DIA, (καλλιπαιδια, from καλλος, 'beauty,' and παις, gen. παιδος, 'a child.') The art of begetting beautiful children. This was the title of a poem by Claude Quillet, in 1655; "*Callipædia sive de pulchre prolis habendæ ratione*." The author absurdly supposes, that the beauty of children is affected by the sensations which the mother experiences during her pregnancy.

CALLIPERS OF BAUDELOCQUE, see Pelvimeter.

CALLIPHYLLUM, (καλλος, 'beauty,' and φυλλον, 'a leaf.') *Asplenium trichomanoides*.

CALLIP'YGOS, (καλλος, 'beauty,' and πυγη, 'buttocks.') A cognomen of Venus, owing to her beautiful nates.

CALLISTHENICS, Calisthenics.

CALLITRICHE AQUATICA, (καλλος, 'beauty,' and τριχος, gen. τριχος, 'hair,' from the capillary and tufted stems.) *C. Verna*—*c. Heterophylla*, *C. Verna*—*c. Intermedia*, *C. Verna*.

CALLIT'RICHE VER'NA, *C. aquatica* seu *intermedia* seu *heterophylla*, *Wa'ter chick'weed*, *Wa'ter starwort*: indigenous; Order, Callitrichaceæ; is diuretic, and used in dropsy, in the form of tincture or of decoction.

CALLITRIS ECKLO'NI, (from καλλος, 'beauty,') A South African tree, Ord, Coniferae, from the branches and cones of which a gum exudes, that resembles Gum Sandarac. This is successfully used in the form of fumigations in gout, rheumatism, oedematous swellings, &c.

CALLITRIS CUPRESSOIDES, a common shrub in the neighbourhood of Cape Town, exudes a similar substance.

CALLOSITAS, Induration—*c. Palpebrarum*, Scleriosis—*c. Vesicæ*, Cystauxe.

CALLOS'ITY, (from *callous*,) *Callos'itas*, *Scy'ros*, *Ty'le*, *Ty'lus*, *Tylo'ma*, *Tylo'sis*, *Dermatoscle'rosis*, *Dermatotylo'ma*, *Dermatotylo'sis*, *Dermatoty'lus*, *Poros*, *Po'rus*, *Ephy'ma Cal'lus*. Hardness, induration, and thickness of the skin, which assumes a horny consistence in places where it is exposed to constant pressure. (F.) *Durillon*. Also the induration, which is observed in old wounds, old ulcers, fistulous passages, &c.

CAL'LOUS, *Callo'sus*, *Ochtho'des*, (*callus*, 'a hard skin,') (F.) *Callex*. That which is hard or indurated. A *Cal'lous Ul'cer* is one whose edges are thick and indurated.

CAL'LUNA ERICA, (καλλυνω, 'to adorn,') *Erica vulgaris*—*c. Vulgaris*, *Erica vulgaris*.

CAL'LUS, *Cal'lus*, *Cal'lum*, (F.) *Cal*. The osseous matter, thrown out between the fractured extremities of a bone, which acts as a cement, and as a new bony formation; *Osteoty'lus*. The words are, likewise, used occasionally in the same sense as *Callosity*.

CALLUS. PROVISION'AL. When the shaft of a long bone has been broken through, and the extremities have been brought in exact juxtaposition, the new matter, first ossified, is that which occupies the central portion of the deposit, and thus connects the medullary cavities of the broken ends, forming a kind of plug, which enters each.

This was termed by M. Dupuytren the provisional Callus.

CALMANTS, Sedatives.

CALME, (F.) The interval that separates the paroxysms of an acute or chronic disease. When the type is intermittent, the word *intermission* is used.

CALOMBA, Calumba.

CAL'OMEL, *Calom'elias*, (καλος, 'beautiful,' and μελας, 'black,') A term, formerly applied to *Æthiops mineral*. By some, the mild chloride of mercury is said to have been so called by Sir Theodore Turquet de Mayerne, in consequence of its having been prepared by a favourite black servant, or, according to others, because it was a good remedy for black bile. It is now applied to the mild chloride of mercury only. See *Hydrargyri submurias*.

CALOMEL STOOLS. A term applied to the green, spinach-like, evacuations occasioned by the internal use of the mild chloride of mercury.

CALOMEL'ANOS TURQUE'TI, (properly *Calomelas*,) *Hydrargyri submurias*. A name, also, given by Riverius to purgative pills, prepared with calomel, sulphur, and resin of jalap.

CALOMELAS, gen. *Calomel'anos*. *Hydrargyri submurias*.

CALO'NIA, (καλωνια.) An epithet formerly given to myrrh. See *Myrrha*.

CALOPHYLLUM INOPHYLLUM, (καλος, 'beautiful,' and φυλλον, 'a leaf,') See *Fagara octandra*.

CALOR, Heat—*c. Animalis*, Animal heat—*c. Mor'dax*, see Heat, internal—*c. Nativus*, Animal heat *CALORIC*, see Heat.

CALORIC'ITY, *Caloric'itas*, (F.) *Caloric'it'*, (from *calor*.) The faculty possessed by living bodies of generating a sufficient quantity of caloric to enable them to resist atmospheric cold, and to preserve, at all times and in every part, a temperature nearly equal. See *Animal Heat*.

CALORIFA'CIENT, *Calorif'iant*, *Calorifa'ciens*, *Calorifi'ans*: (*calor*, and *facere*, 'to make,') Having the power of producing heat. Relating to the power of producing heat.

CALORIFICA'TION, *Calorifica'tio*, (same etymon.) The function of producing heat in organized bodies.

CALORINESES, (from *calor*.) The name under which M. Baumes proposes to arrange all diseases characterized by a sensible change in the quantity of animal heat. The *Calorinèses* form the first class of his Nosology.

CALOTROPIS GIGANTEA, (καλος, 'beautiful,' and τροπις, 'a keel,' alluding to the keel of the flower.) *Mudar*—*c. Mudarii*, *Mudar*.

CALOTTE (F.), *Pile'olum*. Anatomists sometimes give the name, *Calotte aponevrotique*, to the aponeurosis of the occipito-frontalis muscle, which covers it externally; and that of *Calotte du crane* to the skull-cap.

Calotte is also applied to an adhesive plaster, with which the head of a person labouring under tinea capitis is sometimes covered, after the hair has been shaved off. This plaster is pulled suddenly and violently off, in order to remove the bulbs of the hair. It means, also, a sort of coif made of boiled leather, worn by those who have undergone the operation of trepanning, &c.

CALOTTE D'ASSURANCE, Condom.

CALTHA ALPINA, (from καλαθος, 'a basket,' in allusion to the form of the flower [?]). *Arnica montana*—*c. Arvensis*, *Calendula arvensis*—*c. Marsh*, *C. Palustris*—*c. Officinalis*, *Calendula arvensis*.

CAL'THA PALUS'TRIS, *Marsh Mar'yigold*, *Marsh Cal'tha*, *Cow'slips*. Order, Ranunculaceæ: indigenous. A syrup prepared from it is a popular cough remedy.

CALTHA VULGARIS. Calendula officinalis.

CALTRIPS, (from *calceitrapa*,) see Trapanatans.

CALUMB'A, (Ph. U. S., 1863, and Ph. B., 1864.) *Colomb'bo*, *Calom'ba*, *Colom'ba*, *Colum'io*, *Calum'bo*, *Ra'dix Colum'be*, (F.) *Calumbe* ou *Columbe*. The root of *Menispermum palmatum Cocculus palmatus*. Order, Menispermaceæ: indigenous in Mozambique. Its odour is slightly aromatic; taste unpleasantly bitter. It is tonic and antiseptic. Dose, gr. 10 to 3j in powder.

CALUMBA, AMERICAN, *Fraxel'ra Walteri* seu *Carolinen'sis* seu officina'lis seu verticil' lata, *Suer'tia difform'is* seu *Fraxel'ra*, American or *Mariet'ta Colum'ba*, Indian *Let'tuce*, *Yel'low Gent'ian*, *Gold'en Seal*, *Mead'ow pride*, *Pyramis*. The root—*Fraxel'ra* (Ph. U. S.)—is used in the same cases as the true *Calumba*.

CALUS, Callus.

CALVA, *Calva'ria*, (*calvus*, 'bald.') The cranium; the upper part especially; the skull-cap; —the Vault of the Cranium, *Cam'era* seu *Fornix cran'ii*.

CALVARIA, Cranium.

CALVA'TA FERRAMEN'TA. Surgical instruments, which have a smooth head or button.

CALVIT'IES, *Calvit'ium*, (*calvus*, 'bald,') *Phal'acra*, *Phalacro'sis*, *Glabrit'ies*, *Ophi'asis*, *Depila'tio Cap'itis*, *Phalacro'ma*, *Madaro'sis*, *Lip'sotrich'ia*, *Bald'ness*, (Old Eng.) *Ball'edness*, *Cal'vity*, &c., (F.) *Calvitie*, *Chauveté*. Absence of hair, particularly at the top of, and behind, the head.

CALVITIES PALPEBRARUM is loss of the eyelashes.

CALVUS, ('bald,') *Athrix*.

CALX, gen. *Calcis*, *Cal'rium Ter're*, *Protox'ide* of *Cal'cium*, *Calca'ria pu'ra*, *Lime*, (F.) *Chaux*. The lime employed in pharmacy, should be recently prepared by calcination. When water is sprinkled over caustic lime, we have *slaked lime*, *hydrate of lime*, —the *Calcis Hydras* of the London pharmacopœia, *Calx extinc'ta*, *Cal'cium oxyda'tum hydrat'um*, (F.) *Chaux éteinte*.

CALX, see Calcanæum—c. Bismuthi, Bismuth, subnitrate of—c. Chlorinata, *Calcis chloridum*—c. Extincta, see Calx—c. cum Kali puro, Potassa cum calce—c. Oxymuriatica, *Calcis chloridum*—c. Recens, *Calx viva*—c. Salita, *Calcis murias*.

CALX E TESTIS; lime prepared from shells. It has no real medicinal advantages over that prepared from marble.

CALX USTA, *Calx viva*.

CALX VI'VA, *C. re'ens* seu *us'ta* seu *pu'ra* seu *caustica*, *Calx et Calx vi'va*, *Ox'idum Cal'cii*, *Ox'ydum cal'cium*, *Cal'cium oxyda'tum*, *Fu'mans Nix*, *Lime* or *Quick'lime*, (F.) *Chaux vive*. The external operation of calx viva is escharotic, but it is rarely used. Lime is a good disinfecting agent. It is employed internally in the form of *Liquor Calcis*.

CALYCAN'THUS, *C. Flor'idus*, *Callican'thus* [?], *Caroli'na All'spice*, *Sweet-scented shrub*, *Sweet shrub*, (*calvus*, 'a flower-cup,' and *anthos*, 'a flower,') An indigenous plant; Order, Calycanthaceæ; with purplish flowers, of strong, agreeable odour, which appear from March to June. The root is possessed of emetic properties.

CALYPTRANTHES CARYOPHYLLATA, (*καλυπτρα*, 'a veil,' and *ανθος*, 'a flower,' from the appearance of the flowers,) *Myrtus caryophyllata*.

CALYSTEGIA SEPIUM, (*καλυσ*, 'calyx,' and *στεγη*, 'a roof or covering,') *Convolvulus sepium*—c. Soldanella, *Convolvulus soldanella*.

CALYX, (*καλυσ*,) *Calix*.

CAMARA, ('a vaulted room,') *Calva*.

CAMARÈZ, MINERAL WATERS OF. *Camarèz* is a small canton near Sylvanès, in the department of Aveyron, France, where there are acidulous chalybeates. Season from the 15th of June to the 15th of October.

CAMAROSIS, *Camara'tio*, (*καμαρα*, 'a vaulted room,' and *osis*,) *Camara'tio*, *Testudinatio Cra'nii*.

A species of fracture of the skull—*cam'erated fracture*—in which the fragments are placed so as to form a vault, with its base resting on the dura mater.

CAMAS'SIA ESCULEN'TA, *Camass'*, *Kamas'* or *Quamash'* root, *Bread root*; Nat. Order, *Coronariæ*; *Pomme de Prairie* and *Pomme blanche* of the Canadians: *Prairie turnip* of the hunters and trappers of the West, is said to be very extensively used as food by the Digger Indians.

CAMBIE-LEAF, *Nymphaea alba et lutea*.

CAMB'ING. A tree of the Moluccæ Islands, from the bark of which a kind of gum-resin exudes, which has been highly extolled in dysentery. It appears to have some resemblance to the *simarouba*.

CAM'BIUM, ('exchange,') A name formerly given to a fancied nutritive juice, which was supposed to originate in the blood, to repair the losses of every organ, and produce their increase.

CAM'BO, MINERAL WATERS OF. A village in the department of Basses Pyrénées, France, where there are two mineral springs; the one an acidulous chalybeate, the other sulphureous. Temperature, 62° to 69° Fahrenheit. The season is from the 15th of May to the 15th of October.

CAMBODIA, *Cambogia*.

CAMBO'GIA, from Cambodia, in the East Indies, where it is obtained. Hence, likewise, its names *Cambo'dia*, *Cambo'gium*, *Gambo'gia*, *Gambo'gium*, *Gambu'gium*, *Gamboid'ia*. It is called, also, *Gut'ta*, *Gut'ta gamb'ba* seu *Gaman'dræ*, *Gum'mi Gut'ta*, *Catagau'na*, *Cattagau'ma*, *Chry'sopna*, *Laxativus Ind'icus*, *Gum'mi Be'gia* seu *gaman'dræ* seu *de Go'a* seu *de Je'mu* seu *ad Pod'agram*, *Chitta jemoco*, *Camboge* or *Gamboge*, (F.) *Gomme Gutte*. Ord. *Guttiferæ*. A yellow juice obtained from *Hebradendron Cambogioides*, and other plants of the natural family *Guttiferæ*, but it is not known from which of them the official camboge is obtained. The British Pharmacopœia (1864) refers it to an undetermined species of *Garcinia*. It is inodorous, of an orange yellow colour; opaque and brittle; fracture, glassy; is a drastic cathartic, emetic and anthelmintic; and is used in visceral obstructions and dropsy, and wherever powerful hydragogue cathartics are required. Dose from gr. ij to vi, in powder, united with calomel, squill, &c.

CAMBODIA GUTTA, *Garcinia cambogia*.

CAMBU'CA, *Cambuc'ca membra'ta*. Buboës and venereal ulcers, seated in the groin or near the genital organs. See Bubo.

CAMÉLÉE, *Cneorum tricoccum*.

CAMERA, (*καμαρα*,) Chamber, Fornix, Vault—c. Cordis, *Pericardium*—c. Cranii, *Calva*—c. Oculi, Chamber of the eye.

CAMERATIO, (from *camera*,) *Camarosis*.

CAMFOROSMA, *Camphorosma*.

CAMINGA, *Canella alba*.

CAMISIA FËTUS, ('chemise or shirt of the fœtus,') *Chorion*.

CAMISOLE, *Waistcoat*, *strait*.

CAMMARUS, (*καμμαρος*,) *Crab*.

CAMOMILLE FËTIDE, *Anthemis cotula*—c. Puante, *Anthemis cotula*—c. Romaine, *Anthemis nobilis*—c. des Teinturiers, *Anthemis tinctoria*—c. Vulgaire, *Matricaria chamomilla*.

CAMOSIERS, MINERAL WATERS OF. Camosiers is a canton, two leagues from Marseilles, where are two springs containing carbonate of lime, sulphur, chloride of sodium, &c. They are purgative, and used in skin complaints.

CAMOTES, *Convolvulus batatas*.

CAMPAGNE', MINERAL WATERS OF. Campagne is in the department of Aude, France. The waters contain sulphate and chlorohydrate of magnesia. Temperature, 80° Fahrenheit.

CAMPAN'ULA, (dim. of *campana*, 'a bell,' from the shape of the corolla,) The bellflower.

CAMPANULA TRACHE'LIUM, *Can'terbury Bell* or *Throatwort*, was formerly used, in decoction, in relaxation of the fauces. It is, also, called *Cervicaria*.

CAMPE, (καμπη,) Flexion.

CAMPHENE'. The radical of camphor, which, according to Dumas, consists of camphene and one equivalent of oxygen. Camphene or camphine is represented by pure oil of turpentine. That which is used for burning in lamps, and is occasionally employed therapeutically, is oil of turpentine purified by distillation from a solution of caustic potassa.

CAMPHIRE, Camphor.

CAMP'PHOR, (from Arab. *Ca'phur* or *Kam'phur*;) *Camphora*; according to Kraus, more correctly *Campho'ra*, from its East Indian origin, *Caph'ura*, *Caff'a*, *Caf*, *Caj'ur*, *Caph'ora*, *All'afor*, *Camphire*, (F.) *Camphre*. A concrete substance, prepared, by distillation, from *Laurus, Camphora*, *Per'sea Cam'fora*, *Camphora officina'rum*, an indigenous tree of the East Indies. Order, Laurinæ. Two kinds of crude camphor occur in commerce—the one called *tub camphor*, from being imported in tubs. It comes from Batavia, but is, chiefly, the produce of Japan. It is, also, called *Dutch Camphor*, and *Japan Camphor*;—the other, common crude camphor, *China camphor*, *Formo'sa camphor*, from India, in square chests. It is refined by sublimation. Camphor is, also, obtained in Borneo and Sumatra—*Bor'neo* or *Suma'tra camphor*, from *Dryobalanops camphora*. It is very dear, and not used in this country or in Europe. The odour of camphor is strong and fragrant: it is volatile, not easily pulverizable; texture crystalline. Soluble in alcohol, ether, oils, vinegar, and slightly so in water. Its properties are narcotic, diaphoretic, and sedative. Dose, gr. v to ℥j. Dissolved in oil or alcohol, it is applied externally in rheumatic pains, bruises, sprains, &c.

Camphor Oil or *Liquid camphor*, of India, is obtained by piercing the young camphor tree, *Dryobalanops camphora*. The *O'leum camphoræ*, oil of camphor, of the United States Pharmacopœia, is the volatile oil obtained from *Camphora officina'rum*.

CAMP'HORA, Camphor—c. *Officinarium*, see Camphor—c. *Water*, *Mistura Camphoræ*.

CAMP'HORA'CEOUS, *Camphora'ceus*. Relating to or containing camphor; as a 'camphoraceous smell or remedy.'

CAMP'HORATA HIRSUTA, and **C. MONSPELIENSIIUM**, *Camphorosma Monspelica*.

CAMP'HORATED, *Camphoratus*, (F.) *Camphré*. Relating to camphor; containing camphor; as a *camphorated smell*, a *camphorated draught*.

CAMP'HOROS'MA MONSPELI'ACA, *C. Perren'nis*, (camphor, and οσμη, 'odour') *Sela'go*, *Camphorata hirsuta* seu *Monspelien'sium*, *Hair'y Camphoros'ma*, (F.) *Camphrée* de Montpellier. Order, Chenopodiaceæ. *Sex. Syst.* *Tetrandria Monogynia*. This plant, as its name imports, has an odour of camphor. It is regarded as diuretic, diaphoretic, cephalic, antispasmodic, &c. It is also called *Champhen'cæ* and *Stinking Ground Pine*.

CAMP'HOROSMA PERENNIS, *C. Monspelica*.

CAMP'HRE, Camphor.

CAMP'HRE', Camphorated.

CAMP'HRE DE MONTPELLIER, *Camphorosma Monspelica*.

CAMPION, ROSE, *Lychnis githago*.

CAMPOMANE'SIA LINEATIFO'LIA, (after Campomanes, a Spanish naturalist.) A tree, twenty to thirty feet high, which grows in Peru, and whose fruit—*palillo*—of a bright yellow colour, and as large as a moderate-sized apple, has an exceedingly agreeable scent, and is one of

the ingredients in making the perfumed water called *mistura*.

CAMP'SIS, (καψίς,) *Flex'io*, *Curva'tio*, *Inflex'io*. Bone or cartilage, forcibly bent from its proper shape, without breaking.

CAMP'SIS DEPRESSIO, Depression.

CAMPYLOR'RACHIS, (καμπυλος, 'crooked,' and ραχίς, 'spine.') A monster whose spine is crooked.

CAMPYLORRHINUS, (καμπυλος, 'crooked,' and ριν, 'nose.') A monster whose nose is crooked.

CAMPYLOTIS, (καμπυλότης, 'bent,') *Cataclasis*.

CAMPYLIUM, (same etymon,) *Cataclasis*.

CAMUS (F.), *Si'mus*, *Resi'mus*, *Si'mo*, *Si'lo*, *Si'lus*. One who has a short, stumpy nose. The French speak of *Nez camus*, 'short nose.'

CANADA BURNET, *Sanguisorba canadensis*.

CAN'ADA, MIN'ERAL WATERS OF. *St. Cath'érine's*, at the head of Lake Ontario, at the entrance of the Welland Canal, 12 miles from Niagara Falls, and on the Great Western Railroad, has a mineral spring, much frequented, which Dr. Chilton, of New York, analyzed and found to contain chlorides of calcium, magnesium, sodium, and iron, sulphate of lime, carbonates of lime and magnesia, bromide and iodide of magnesium, silica and alumina.

Tuscaro'ra Ac'id Spring is in the township of Tuscarora, about 20 miles north of Pass Dover. It contains a large amount of free sulphuric acid; beside sulphates of the alkalies, lime, magnesia, alumina and iron.

Char'lotteville Sul'phur Spring is a few miles from Dover, on Lake Erie. It contains an unusual quantity of sulphuretted hydrogen.

The *Caledo'nia Springs*—a well known watering place—are a few miles south of the Ottawa river, and 40 miles from Montreal. There are various sources: one—the *Gas Spring*—discharging carburetted hydrogen; a *Saline* and a *Sulphur Spring*; besides an *Intermitting Spring*, which discharges large quantities of carburetted hydrogen; and in which iodine and bromine have been detected.

CANAL, *Cana'lis*, *Ductus*, *Meat'us*, *Sol'ten*, *Pol'ros*, *Och'etus*, (F.) *Canal*, *Conduit*. A channel for affording passage to liquids, or solids, or to certain organs.

CANAL ALIMEN'TARY, *C. Diges'tive*, *Cana'lis* seu *Ductus cibarius* seu *digestivus* seu *alimentarius*, *Tubus alimentarius* seu *intestinalis*, *Digestive Tube*, *Aliment'ary Duct* or *Tube*. The canal extending from the mouth to the anus.

CANAL ARACH'NOID, *Cana'lis Bicha'tii*, *Canal' of Bichat*. A canal, formed by the extension of the arachnoid over the transverse and longitudinal fissures of the brain, which surrounds the vena magna Galeni. The orifice of the canal has been termed the *Foramen of Bichat*.

CANAL ARTERIEL, Arterial duct—c. *de Bartholin*, *Ductus Bartholinus*—c. of Bichat, *Canal*, arachnoid—c. Bullary, of Petit, *Godronné canal*—c. *Carotidian*, *Carotid canal*—c. *Chole'doque*, *Choledoch duct*—c. Ciliary, *Ciliary canal*—c. of Cotunnus, *Aquæductus vestibuli*—c. *Dé'fèrent*, *Deferens vas*—c. of Fontana, *Ciliary canal*—c. *Godronné*, *Godronné canal*—c. *Hépatique*, *Hepatic duct*.

CANAL OF HY'GUIER. A small canal at the inner side of the fissure of Glaser, through which the chorda tympani nerve leaves the cavity of the tympanum.

CANAL HY'ALOID. A cylindrical passage, described by J. Cloquet as formed by the reflection of the hyaloid membrane into the interior of the vitreous body around the nutritious artery of the lens. M. Cruveilhier has never been able to see it.

CANAL, INCIS'SIVE, see *Palatine canals*—c. *In-*

flexe de l'os temporal, Carotid canal.—c. Infra-orbital, Suborbital canal.—c. *Intermédiaire des ventricules*, Aquæductus Sylvii.

CANAL, INTESTINAL, *Canalis* seu *Ductus intestinalis*. The portion of the digestive canal formed by the intestines.

CANAL OF JACOBSON, Canal, tympanic.

CANAL, MEDULLARY. The cylindrical cavity in the body or shaft of a long bone, which contains the marrow.

CANAL, NASAL, Lachrymal canal.

CANAL OF NUCK. A cylindrical sheath formed around the round ligaments of the uterus by a prolongation of the peritoneum into the inguinal canal. It gives rise to hydrocœlic cysts in the labium externum, which might be mistaken for hernia of the labium.

CANAL DE PETIT, *Godronné canal*—c. *de la Pudeur*, Vagina—c. *Pulmo-aortique*, Arterial duct—c. Rachidian. Vertebral canal.

CANAL OF SCHLEMM. A minute circular canal, discovered by Professor Schlemm, of Berlin. It is situate at the point of union of the cornea and sclerótica.

CANAL SPERMATIC, Deferens vas—c. Spinal, Vertebral canal—c. *Spiroïde de l'os temporal*, Aquæductus Fallopii—c. *de Stenon*, Ductus salivaris superior—c. *Thoracique*, Thoracic duct—c. *Veineux*, Canal, venous—c. Vulvo-uterine, Vagina—c. *de Warthon*, Ductus salivaris inferior.

CANAL, TYMPANIC, *Canalis tympanicus*, Canal of Jacobson. A canal which opens on the lower surface of the petrous portion of the temporal bone, between the carotid canal and the groove for the internal jugular vein. It contains Jacobson's nerve.

CANAL, VECTOR, Tube, Fallopian.

CANAL, VEINOUS, *Canalis* seu *Ductus venosus* seu *Aran'tii* seu *D. V. Aran'tii*, (F.) Canal veineux. A canal which exists only in the fœtus. It extends from the bifurcation of the umbilical vein to the vena cava inferior, into which it opens below the diaphragm. At times, it ends in one of the infra-hepatic veins. It pours into the cava a part of the blood, which passes from the placenta by the umbilical vein. After birth, it becomes a fibro-cellular cord.

CANAL OF WIRUNG, see Pancreas.

CANALES AÉRIFERI, pl. of *Canalis*, see Bronchia.

CANALES BRESCHETI. Canals in the diploë for the passage of veins; called after M. Breschet.

CANALES CIRCULARES, Semicircular canals—c. Cochleæ, Scalæ of the cochlea—c. Lacrymales, Lachrymal ducts—c. Membranei renum, see Calix—c. Tubæformes, Semicircular canals.

CANALICULAR, *Canalicula'ris*, (F.) *Canaliculaire*, (from *Canaliculus*.) Having small canals.

CANALICULAR ABSCESS OF THE BREAST, An abscess that communicates with the galactophorous ducts.

CANALICULAR TISSUE, any tissue containing small canals, as that of bone.

CANALICULATED, Grooved, see Canaliculus.

CANALICULATED, *Cannelé*, Grooved.

CANALICULÉ, Grooved.

CANALICULES CALCIFÈRES, Canaliculi calciferi.

CANALICULI, pl. of Canaliculus—c. Biliiferi, Pori biliarii—c. of Bone, see Lacunæ of Bone.

CANALICULI CALCIFERI, *Calcif'eron* canal, (F.) *Canalicules Calcifères*. Minute canaliculi, observed with the microscope, in the cartilages of bones, which are full of calcareous salts.

CANALICULI HAVERSIANI, Canals, nutritive—c. Lacrymales, Lachrymal ducts—c. Limacum, Lachrymal ducts—c. Semicirculares,

Semicircular canals—c. Seminales, see Testicle—c. Seminiferi, see Testicle—c. Vasculosi, Canals, nutritive.

CANALIC'ULUS, (dim. of *canalis*, 'a channel.') A small channel. See Canaliculi.

CANALIS, Canal, Meatus—c. Alimentarius, Canal, alimentary—c. Arteriosus, Arterial duct—c. Bichatii, Canal, arachnoid—c. Canaliculatus, Gorge—c. Caroticus, Carotid canal—c. Cervicis, see Uterus—c. Deferens, Deferens vas—c. Eminentiæ quadrigeminæ, Aquæductus Sylvii—c. Fallopii, Aquæductus Fallopii—c. Intestinorum, Intestinal tube—c. Lacrymalis, Lachrymal or nasal duct—c. Medius, Aquæductus Sylvii—c. Medullæ Spinalis, see Vertebral column—c. Nervus fistulosus renum, Ureter—c. Orbite nasalis, Lachrymal or nasal duct—c. Scalarum communis, Infundibulum of the cochlea—c. Semicircularis horizontalis, see Semicircular Canals—c. Semicircularis verticalis posterior, see Semicircular Canals—c. Semicircularis verticalis superior, see Semicircular canals—c. Tympanicus, Canal, tympanic—c. Urinarius, Urethra—c. Vidianus, Pterygoid canal.

CANALISATION, *Canalisatio*. The conversion of a vessel into a rigid tube.

CANALES, BONE, of Breschet, see Diploë—c. of Havers, Canals, nutritive—c. Haversian, Canals, nutritive.

CANALES, NUTRITIVE, *Canals for the nutrition of bones*, *Ductus nutrit'ii*, *Canaliculi vasculosi* seu *Haversiani*, *Haver'sian Canals*, *Canals of Havers*, (F.) *Canaux nourriciers* ou du *Nutrition des Os*, *Conduits nourriciers* ou *nutriciers*. The canals through which the vessels pass to the bones. They are lined by a very fine lamina of compact texture, or are formed in the texture itself. There is, generally, one large nutritious canal in a long bone, situate towards its middle.

CANAPACIA, Artemisia vulgaris.

CANARIES, CLIMATE OF. The climate of the Canaries greatly resembles that of Madeira. That of the latter, however, is more equable, and the accommodation for invalids much superior.

CANARIUM COMMUNE, (from the Canaries,) see Amyris elemifera.

CANARYSEED, Phalaris Canariensis.

CANAUX (pl. of *Canal*) A QUEUX, see Aqueous—c. *Demicirculaires*, Semicircular Canals—c. *Éjaculateurs*, Ejaculatory ducts—c. *Nourriciers*, Canals, nutritive—c. *de Nutrition des Os*, Canals, nutritive.

CANAUX DE TRANSMISSION. According to Bichat, the bony canals intended to give passage to vessels and nerves going to parts more or less distant—as the *Canalis Caroticus*.

CANAUX VEINEUX, *Venous Canals*. The canals situate in the diploë, which convey venous blood.

CAN'CAMUM, (κακκρον.) A mixture of several gums and resins exported from Africa, where it is used to deterge wounds. Dioscorides so calls the tears from an Arabian tree, which are similar to myrrh, and of a disagreeable taste. He advises it in numerous diseases. This name is given, also, to Anime.

CANCAMY, Anime.

CANCELLED, *Cancel'latus*, (F.) *Cancel'é*; (*cancelli*, 'lattice-work.') Formed of cancelli,—as the 'cancelled structure of bone.'

CANCEL'LI, (pl. of *Cancel'lus*, dim. of *cancer*, 'a bar,') 'Lattice-work.' The Cellular or Spongy Texture of Bones, (F.) *Tissu cellulaire*; consisting of numerous cells, communicating with each other. They contain a fatty matter, analogous to marrow. This texture is met with, principally, at the extremities of long bones; and some of the short bones consist almost wholly of it. It allows of the expansion of the extremities of bones,

without adding to their weight, and deadens con-
cussions.

CANCEL'LUS, (dim. of *cancer*, 'a crab.') A species of crayfish, called the *Wrong Heir*, and *Bernard the Hermit*: which is said to cure rheumatism, if rubbed on the part.

CAN'CER, gen. *Canceri*, 'a crab.' *Car'cinus*, *Lu'pus Cancro'sus*. A disease, so called either on account of the hideous appearance which the ulcerated cancer presents, or on account of the great veins which surround it, and which the ancients compared to the claws of the crab: called also *Carcino'ma*. It generally consists of a scirrhus, livid tumour, intersected by firm,

whitish, divergent bands; and occurs chiefly in the secretory glands. The pains are acute and lincinating, and often extend to other parts. The tumour, ultimately, terminates in a fetid and ichorous ulcer, — *Ulcus cancro'sum*. It is distinguished, according to its stages, into *occult*, *Can'cer occultus*, and *open*, *C. aper'tus*; the former being the scirrhus, the latter the ulcerated condition. At times, there is a simple destruction or erosion of the organ, at others, an *enceph'loid* or *cer'ebriform*, and at others, again, a *colloid* degeneration.

For its production, it requires a peculiar diathesis, or cachexia. The following table, from

ENCEPHALOID.	SCIRRHUS.	COLLOID.
Resembles lobulated cerebral matter. Is commonly opaque from its earliest formation. Is of a dead white colour. Contains a multitude of minute vessels. Is less hard and dense than scirrhus. Is frequently found in the veins issuing from the diseased mass. The predominant microscopical elements are globular, not always distinctly cellular, and caudate corpuscula. Occasionally attains an enormous bulk. Has been observed in almost every tissue of the body. Very commonly co-exists in several parts or organs of the same subject. Is remarkable for its occasional vast rapidity of growth. Is frequently the seat of interstitial hemorrhage and deposition of black or bistre-coloured matter. When softened into a pulp, appears as a dead white or pink opaque matter of creamy consistence. Subcutaneous tumours are slow to contract adhesion with the skin. Ulcerated encephaloid is frequently the seat of hemorrhage, followed by rapid fungous development. The progress of the disease after ulceration is commonly very rapid. It is the most common form under which secondary cancer exhibits itself. Is the species of cancer most frequently observed in young subjects.	Resembles rind of bacon traversed by cellulo-fibrous septa. Has a semi-transparent glossiness. Has a clear whitish or bluish yellow tint. Is comparatively ill-supplied with vessels. Is exceedingly firm and dense. Has not been distinctly detected in this situation. The main microscopical constituents are juxtaposed nuclear cells; caudate corpuscula do not exist in it. Rarely acquires larger dimensions than an orange. Its seat, as ascertained by observation, is somewhat more limited. Is not unusually solitary. Ordinarily grows slowly. Is comparatively rarely the seat of these changes. Resembles, when softened, a yellowish brown semitransparent gelatinous matter. Scirrhus thus situate usually becomes adherent. Scirrhus ulcers much less frequently give rise to hemorrhage; and fungous growths (provided they retain the scirrhus character) are now more slowly and less abundantly developed. There is no such a remarkable change in the rate of progress of the disease after ulceration has set in. Is much less common before puberty.	Has the appearance of particles of jelly inlaid in a regular alveolar bed. The contained matter is strikingly transparent. Greenish yellow is its predominant hue. Its vessels have not been sufficiently examined as yet. The jelly-like matter is exceedingly soft; a colloid mass is, however, firm and resisting. The pulsatious variety has been detected in the veins. Is composed of shells in a state of emboisement. Observes a mean in this respect. Has so far been seen in a limited number of parts only. Has rarely been met with in more than one organ. Grows with a medium degree of rapidity. Undergoes no visible change of the kind. Has so far been observed in adults only.

Dr. Walshe, exhibits the characters of the three great species of cancer.

The use of irritants in cancerous affections is strongly to be deprecated. When the disease is so situate that excision can be practised, and the constitution does not exhibit great contamination, the sooner it is removed the better. See *Carcinoma*, and *Thyrotoblaste*.

CANCER ALVÉOLAIRE, Colloid — c. Alveolaris, Colloid — c. Apertus, see Cancer.

CANCER AQUATICUS, *Gan'grenous stomati'tis*, *Can'crum O'ris*, *Gan'grena'p'is*, *Can'ker of the mouth*, *Gan'grenous sore mouth*, *Gan'grenous ero'sion of the cheek*, *Slough'ing Phaged'e'na of the mouth*, *Wa'ter Can'ker*: called, also, *Aph'thæ serp'en'tes*, *Gan'gr'e'na O'ris*, *No'ma*, *No'më*, *No'mus*, *Pseudocarcino'ma la'bi'i*, *Stoma'cæ gan'grena'sis*, *Chello'ac'e*, *Ulo'ac'e*, *Uli'tis sep'tica*, *Chriloma'la'cia*, *Scorbu'tus O'ris*, *Stomatoma'la'cia pri'tida*, *Stomatosep'is*, *Stomatonecro'sis*, *Carbu'culus*

labio'rum et gena'rum, (F.) *Cancer aquatique*, *Stomatite gangréneuse*, *S. Charbonneuse*, *Gan'grène ou Sphacèle de la Bouche*, *Fégarité*, *Aph'the gan'gréneux*. Certain sloughing or gangrenous ulcers of the mouth — so called, perhaps, because they are often accompanied with an afflux of saliva. The disease is not uncommon in children's asylums, and demands the same treatment as hospital gangrene, — the employment of caustics, and internal and external antiseptics.

CANCER AQUATIQUE, Cancer aquaticus, Stomacæ — c. *Alvéolaire*, Colloid — c. *Astacus*, see *Cancerorum chelæ* — c. *Black*, *Cancer melanotic* — c. of the Breast, *Mastocarcinoma* — c. *Camini-riorum*, *Cancer*, chimney-sweeper's.

CANCER CELL. The cell which, according to some, characterizes cancer. This as well as the nucleus, (F.) *Cellule et noyau cancéreux*, although most diversified in its appearance — polygonal, caudate, fusiform, concentric, compound — is consi-

dered pathognomonic by some, but not to be relied upon by others.

CANCER, CELLULAR, Encephaloid — *c. Cérébri-forme*, see Encephaloid.

CANCER, CHIMNEY-SWEEPER'S, *Soot'wart*, *Can'-cer mundito'rum*, seu *purgato'ris infumicu'li* seu *caminiario'rum*, *Can'cer* seu *carcino'ma scro'ti*, *Oschocarcino'ma*, *Oschocarcino'ma*, (F.) *Cancer des Ramoneurs*. This affection begins with a superficial, painful, irregular ulcer, with hard and elevated edges, occupying the lower part of the scrotum. Extirpation of the diseased part is the only means of effecting a cure.

CANCER COLLOIDE, Colloid.

CANCER, DAVIDSON'S REMEDY FOR, see *Conium maculatum*—*c. Dur*, *Scirrhus*.

CANCER ÉBURNÉ, ('eburneous cancer.') A kind of waxy degeneration of the breast, so called by M. Alibert, but which appears to be in no way allied to cancer.

CANCER ENCÉPHALOÏDE, *Scirrhus* — *c. Epithelial*, see *Epithelial*—*c. Fibreux*, *Scirrhus*—*c. Fibrous*, *Scirrhus* — *c. du Foie*, *Hépatosarcomie*.

CANCER GALE'NI, (F.) *Cancer de Galien*. A bandage for the head, to which Galen gave the name *cancer*, from its eight heads resembling, rudely, the claws of the crab. It is now supplied by the bandage with six *chefs* or heads, which is called the *Bandage of Galen* or *B. of the Poor*.

CANCER DE GALIEN, *Cancer Galeni* — *c. Gelatiniform*, *Colloid* — *c. Gelatinous*, *Colloid* — *c. Gommeux*, *Colloid* — *c. Hard*, *Scirrhus* — *c. Intestinal*, *Enteropathia cancerosa* — *c. des Intestins*, *Enteropathia cancerosa* — *c. of the Lung*, *Phthisis*, *cancerous* — *c. Lupus*, *Lupus* — *c. Mastoid*, see *Mastoid cancer* — *c. Medullaris*, *Encephaloid* — *c. Melæneus*, *Cancer*, *melanotic* — *c. Mélane*, *Cancer*, *melanotic* — *c. Melanodes*, *Cancer*, *melanotic*.

CANCER, MELANOT'IC, *Can'cer melano'des*, *Carcino'ma melano'des* seu *melano'ticum* seu *melæ'-neus*, *Fungus melano'des*, *Mel'anocarcino'ma*, *Black* or *Mel'anoid Cancer*, (F.) *Cancer Mélane*. A combination of cancer and melanosis.

CANCER, MILTLIKE, see *Miltlike*.

CANCER MOLLIS, see *Encephaloid* — *c. Mon*, *Encephaloid* — *c. Munditorum*, *Cancer*, *chimney-sweepers'* — *c. Nephroid*, see *Nephroid* — *c. Occultus*, see *Cancer* — *c. Oculi*, *Scirrhopthalmus* — *c. Oris*, *Stomacæ* — *c. Ossis*, *Spina ventosa* — *c. Pharyngis* et *oesophagi*, *Læmoscirrhus* — *c. Pul-tacé*, *Colloid* — *c. Purgatoris infumiculi*, *Cancer*, *chimney-sweepers'*.

CANCER ROOT, *Orobanche Virginiana*, *Phytolacca decandra*.

CANCER DES RAMONEURS, *Cancer*, *chimney-sweepers'* — *c. Scirrhusos*, *Scirrhus* — *c. Scro'ti*, *Cancer*, *chimney-sweepers'* — *c. Soft*, *Hæmatodes fungus* — *c. Solanoid*, see *Solanoid* — *c. Squirrheux*, *Scirrhus* — *c. of the Stomach*, *Gastrostenosis cardiaca* et *pylorica* — *c. Uteri*, *Metrocarcinoma*.

CANCER WEED, *Goodyera pubescens*, *Salvia lyrata*.

CANCÉREUX, *Cancerous*.

CANCERISM, see *Diathesis*.

CANCEROMA, gen. *Cancero'matis*, (*cancer* and *oma*.) *Carcinoma*.

CANCEROUS, *Cancero'eus*, *Carcino'eus*, (F.) *Cancéreux*, *Thnéoblastique*. Relating to cancer; as *Cancerous ulcer* (*Carcinocoe'lis*), *Cancerous diathesis*.

CANCHALAGUA, *Chironia Chilensis*.

CANCRENA, *Gangrene*.

CANCRODES, (*cancer* and *odes*.) *Canceroid*.

CANCROID, *Kankroid*, *Cancero'i'des*, *Kencero'i'des*, *Cancro'i'des*, *Carcino'i'des*, *Carcino'i'des*, (*cancer* and *oidos*, 'form.') That which assumes a cancerous appearance. *Canceroid* is a name given to certain cutaneous cancers by Alibert: called

also *Che'lloid*, or *Ke'lloid*, *Che'lloï's*, *Cen'croï's*, *Ke-loï'des*; *Ke'lis*, from their presenting a flattish raised patch of integument, resembling the shell of a tortoise. Velpéau terms such a growth, when it forms cicatrices, *Che'lloïde cicatricielle*.

CANCROID OF THE SKIN, see *Epithelial*.

CANCROIS, see *Canceroid*.

CANCROMA, gen. *Cancero'matis*, *Carcinoma*.

CANCRO'UM CHELÆ, (χελαι, 'a claw,') 'erab's claws,' like *Oc'uli* seu *Lap'id'es* seu *Lapilli Can-cro'rum*, *Concrementa As'taci fluvial'is*, *Crab's stones* or *eyes*, (F.) *Yenez d'écrevisse*, *Pierres d'écrevisse* — *Concretions* found, particularly, in the *Cancer As'tacus* or *Cray-fish* — consist of carbonate and phosphate of lime, and possess antacid virtues, but not more than chalk.

CANCROSUS, *Cancerous*, *Chancreuse*.

CANCRO'UM ORIS, *Cancer Aquaticus*, *Stomacæ*.

CANDELA, (*candere*, 'to glow,') *Bougie* — *c. Fumalis*, *Pastil* — *c. Medicata*, *Bougie* — *c. Regia*, *Verbascum nigrum*.

CANDELARIA, (*candela*, 'a candle,' from its stalk,) *Verbascum nigrum*.

CAN'DI, *Can'dum*, *Can'thum*, *Can'tion*, *Can'dy*, (from *candidus* — 'white, bleached, purified,') *Purified* and *crystallized sugar*. See *Saccharum*.

CANDIDUM OVI, ('white of egg,') *Albumen ovi*.

CANDLE (Prov.), *Pupil*.

CAN'DLE, MERC'URIAL. A candle made of *wax* and *vermilion* for mercurial fumigation.

CANDLE'NUT TREE, *Aleurites triloba*.

CANDLE TREE, *Parmentiera cerifera*.

CANDY, *Candi*.

CANDYTUFT, BITTER, *Xeris amara*.

CANE, SUGAR, see *Saccharum* — *c. Sweet*, *Acorus calamus*.

CANÉFICIER, *Cassia fistula*.

CANELA, *Nectandra cinnamomoides*.

CANELÉ, (from *canal*.) *Grooved*.

CANELLA, see *Canella alba*.

CANEL'LA AL'BA, (dim. of *canna*, 'a reed,') A West India tree. *Fam. Magnoliacæ: Ser. Syst. Dodecandria Monogynia*. It is so called because its bark is rolled up like a reed. *Cort'x Winter'a'nus spu'rius*, *Canell'a Cuba'na* seu *Winter'a'na*, *Cinnamo'mum al'bum*, *Cort'x Antiseor-bu'ticus* seu *Aromat'icus*, *Cos'tus cortico'sus*, *Cam'in'ga*, *Canell'a* (Ph. U. S.), *Canell'a Bark*, (F.) *Canelle* ou *Canelle blanche*, *Fausse Écorce de Winter*, *Écorce Caricostine*; *Ord. Guttifera*, is a pungent aromatic. Its virtues are partly extracted by water; entirely by alcohol. It is a stimulant, and is added to bitters and cathartics.

CANELLA CARYOPHYLLATA, *Myrtus caryophyllata* — *c. Cubana*, *C. alba*, *Laurus cassia* — *c. Malabarica* et *Javensis*, *Laurus cassia* — *c. Winterana*, *c. Alba*.

CANELLIFERA MALABARICA, *Laurus cassia*.

CANE'PIN, (F.) A fine lamb's skin or goat's skin, used for trying the quality of lancets.

CANICACEOUS, (from *canis*.) *Furfuraceous*.

CAN'ICÆ. Meal, in which there is much bran. Also, coarse bread; or bread in which there is much bran — *Pa'nis canica'eus*.

CANICIDA, (*canis*, and *cadere*, 'to kill,') *Aconitum*.

CANIC'ULA: the *Dog's-tail*, (dim. of *canis*, 'a dog,') *Σειριος*, *Sirius*, (F.) *Canicule*. This star, which gives its name to the *Dog-days*, *Dies canicula'res*, because they commence when the sun rises with it, was formerly believed to exert a powerful influence on the animal economy. The *Dog-days* occur at a period of the year when there is generally great and oppressive heat, and therefore — it has been conceived — a greater liability to disease.

CAN'IF, (from the Eng. [?] Knife.

CANIN, *Canine*.

CANINANÆ RADIX, Caineæ radix.

CANINE', *Caninus*, *Cynicus*, (*canis*, 'a dog.')(F.) *Canin*. That which has a resemblance to the structure, &c., of a dog.

CANINE FOS'SA, *Fos'sa Canina*, *In'fra-or'bitar* or *Sub'orbitar fos'sa*, (F.) *Fosse Canine*. A small depression on the superior maxillary bone, above the *dens caninus*, which gives attachment to the *caninus* or *levator anguli oris* muscle.

CANINE LAUGH, *Sardon'ic laugh*, *R'vus Caninus* seu *Sardon'icus* seu *Sardo'nus* seu *Sarda'nus* seu de *Sardo'nia* seu *involuntari'us* seu *spas'ticus*, *Tortu'ra* seu *Distor'sio O'ris*, *Gelas'mus*, *Sardi'asis*, *Sardon'asis*, *Tris'mus Sardon'icus* seu *cyn'icus*, *Spas'mus musculo'rum fa'ciei* seu *cyn'icus*, *Pro'sopospas'mus*, (F.) *Ris canin*, *R. Sardonique*, *R. Sardonien*, *R. moqueur*. A sort of laugh, the facial expression of which is produced particularly by the spasmodic contraction of the *Caninus* muscle. Probably, this expression, as well as *Cyn'ic Spasm*, *Spas'mus caninus* seu *cyn'icus*, *Convul'sio canina*, *Tris'mus cyn'icus*, may have originated in the resemblance of the affection to certain movements in the upper lip of the dog. The *Risus Sardon'icus* is said to have been so called from similar symptoms having been induced by a kind of *Ranunculus*—*R. sceleratus*—that grows in Sardinia.

CANINE TEETH, *Dentes canini* seu *Lania'rii* seu *angula'res* seu *cuspidati* seu *columella'res* seu *can'ta'res* seu *morden'tes*, *Cynodon'tes*, *Eye Teeth*, *Lycodon'tes*, *Pug Teeth*, (F.) *Dents canines* ou *lan'aires* ou *angulaires* ou *oculaires* ou *oeillères* ou *conoïdes*. The teeth between the lateral incisors and small molares of each jaw;—so named because they resemble the teeth of the dog.

CANINUS, *Levator anguli oris*—c. *Sentis*, *Rosa canina*—c. *Spasmus*, see *Canine Laugh*.

CANIRAM, *Strychnos nux vomica*.

CANIRAMINUM, *Brucine*.

CANIRUBUS, *Rosa canina*.

CANIS INTERFECTOR, ('dog killer,') *Vera'trum sabadilla*—c. *Ponticus*, *Castor fiber*.

CANIT'IES, (*canus*, 'white.')(F.) *Blanch'ing*, *White'ness* or *Gray'ness of the Hair*, and especially of that of the head—*Trichono'sis polio'sis*. (F.) *Canitie*. When occurring in consequence of old age, it is not a disease. Sometimes it happens suddenly, and apparently in consequence of severe mental emotion. The causes, however, are not clear. See *Poliosis*.

CANKER, (*cancer*), *Stomaceæ*—c. of the Mouth, *Cancer aquaticus*.

CANKER-RASH, see *Cynanche maligna*.

CANKER, WATER, *Cancer aquaticus*.

CANNA, (*kava*, 'a reed,') see *Tous-les-Mois*, *Cassia fistula*, *Trachea*—c. *Brachii*, *Ulna*—c. *Domestica cruris*, *Tibia*—c. *Eduilis*, *Tous-les-Mois*—c. *Fistula*, *Cassia Fistula*—c. *Indica*, *Sagittarium alexipharmacum*—c. *Major*, *Tibia*—c. *Minor*, *Fibula*, *Radius*—c. *Solutiva*, *Cassia fistula*—c. *Starch*, *Tous-les-Mois*.

CANNABIN, *Bangue*.

CANNAB'INA, (*καναβίς*, 'hemp.')(F.) Remedies composed of *Cannabis Indica*.

CANNABINA AQUATICA, *Eupatorium cannabinum*.

CANNABIS INDICA, (*καναβίς*, 'hemp.')(F.) *Bangue*. See, also, *Churrus*.

CAN'NABIS SATI'VA (F.), *Chanvre*, *Chambrée*. The seed of this—*Hemp'seed*, *Sem'ina Can'nabis*, (F.) *Chênevis*—is oily and mucilaginous. The decoction is sometimes used in gonorrhœa.

CANNACORUS RADICE CROCEA, *Curcuma longa*.

CANNAMELLE, (*mel*, 'honey,') see *Saccharum*.

CANNE AROMATIQUE, *Acorus calamus*—c. *Congo*, *Costus*—c. de *Rivière*, *Costus*—c. à *Sucre*, see *Saccharum*.

CANNEBERGE, *Vaccinium oxycoccos*—c. *Ponctué*, *Vaccinium vitis idæa*.

CANNEL, *Laurus cinnamomum*.

CANNEL BONE, *Clavicle*.

CANNEL WATER, *Aqua cinnamomi*.

CANNELÉ ou CANELEE (F.), ([L.] *canalis*, 'a canal.')(F.) *Sulca'tus*, *Stria'tus*, *Canalicula'tus*. Having a canal or groove—as *Muscle cannelé* (Lieutaud), the Gemini; *Corps cannelés* ou *striés*, the *Corpora striata*; *Sonde cannelée*, a grooved sound, &c. See *Grooved*.

CANNELL BAYNE, *Clavicle*.

CANNELLE (dim. of *canna*, 'a reed,') *Laurus cinnamomum*—c. *Blanche*, *Canella alba*—c. de la *Chine*, *Laurus cassia*—c. de *Coromandel*, *Laurus cassia*—c. *Fausse*, *Laurus cassia*—c. *Giroflée*, *Myrtus caryophyllata*—c. des *Indes*, *Laurus cassia*—c. de *Java*, *Laurus cassia*—c. de *Malabar*, *Laurus cassia*—c. *Matte*, *Laurus cassia*—c. *Officinale*, *Laurus cinnamomum*—c. *Poirée*, see *Wintera aromatica*.

CANNULA, *Canula*.

CANNULÉ PULMONUM, (pl. of *Canula*), *Bronchia*.

CANOPUM, see *Sambucus*.

CANOR STETHOSCOPICUS, (*canor*, 'a tune,')(F.) *Tintement métallique*.

CANQUOIN'S CAUSTIC PASTE, *Pasta zinci chloridi*.

CANTABRICA, (from *Cantabri*, a people of Spain.) *Convolutus Cantabrica*.

CANTABRUM, *Furfur*.

CANTARELLUS, *Meloe proscarabæus*.

CANTATIO, (*cantare*, 'to sing,') *Charm*.

CANTEL, *Vertex*.

CANTERBURY, MIN'ERAL WA'TERS OF, *Aquæ Cantuarien'ses*. The waters of Canterbury, in Kent, England, are impregnated with iron, sulphur, and carbonic acid.

CANTERIUM, *Cantherius*.

CANTHARIDE TACHETÉE, *Lytta vittata*.

CANTHARIDINE, see *Cantharis*.

CANTHARIS, gen. *Canthar'idis*; (*κανθαρίς*, gen. *κανθαρίδος*), *C. vesicato'ria*, *Mus'ca Hispan'ica*, *Mel'oe vesicato'ria*, *Ly'tta vesicato'ria*, *Tege'nera medicina'lis*, *Blist'ering Fly*, *Blist'eryfly*, *Blist'erbeetle*, *Span'ish Fly*, *Fly*, (F.) *Cantharides*, *Mouches*, *M. d'Espagne*. This fly—*Order*, *Coleoptera*—originally, perhaps, a native of Italy and Spain, is now found in France, Germany, Hungary, Russia, Siberia and England. It is, however, rare in the last-named country. It is found on species of *Oleaceæ*—as the ash, privet, and lilac; and of *Caprifoliaceæ*—as the elder and lonicera. It is much employed in medicine, and is the most common vesicatory. Given internally, and even when absorbed from the skin, it affects the urinary organs, exciting strangury. This may be prevented, in cases of blisters, by interposing between the blistering plaster and skin a piece of tissue paper. Diluents relieve the strangury. Dose, half a grain to one grain. If kept dry, the flies will retain their activity for many years. Their active principle—*Can'tharidin*, *Cantharidi'na*—has been separated from them.

Can'tharis vesicato'ria is the ordinary blistering insect used in Europe.

Can'tharis seu *Ly'tta vitta'ta*, (which see,) and *C. atra'ta*, *C. margina'ta*, and *C. cine'rea*, of America; *C. atoma'ta*, of Brazil; *C. ru'ficeps*, of Sumatra and Java; *C. gi'gas*, *Ly'tta cæru'lea*, of Guinea and the East Indies; *C. viola'cea*, *Ly'tta gi'gas mas*, of the East Indies; *C. Syri'aca*, *Ly'tta seg'etum*, of Arabia; *My'ta'bris*, *M. puncta'ta*, *M. pustula'ta*, and *M. cicho'rii*, of China and the East Indies; *Mel'oe proscarabæus*, and *M. maja'lis* or *True May'worm*—possess similar properties. Dr. Leidy found, that the vesicatory principle of *Ly'tta vittata* appears to belong to the blood, the peculiar fatty substance of certain accessory glands of the generative apparatus and to the eggs.

CANTHERIUS, (κάντηριος,) *Cante'rium*. The cross-piece of wood in the apparatus used by Hippocrates for reducing luxations of the humerus.

CANTHITIS, (*canthus* and *itis*.) Inflammation of the canthus of the eye.

CANTHIUM CORONATUM, *Randia dume-torum*.

CANTHOPLAS'TICĒ, *Canthoplasty*, (F.) *Canthoplastic*, (κάνθος, 'the angle of the eye,' and πλαστικός, 'formative.') The formation, by plastic operation, of the angle of the eye. An operation proposed by Ammon, when the eyelids are not sufficiently cleft.

CANTHUM, *Candi*.

CANTHOR/RHAPHY, *Canthorrhaphia*, (*canthus*, and ράφη, 'suture.') Suture of the angle of the eye.

CAN'THUS, (κάνθος,) *An'gulus ocula'ris*, *Fons lacryma'rum*. The corner or angle of the eye. The great'er can'thus is the in'ner an'gle, *Hir'cus*, *Hir'gus*, *Rhan'ter*; the les'ser can'thus, the out'er an'gle, *Parop'ia*, *Pe'ga*.

CANTIA'NUS PUL'VIS, 'Kent Pow'der.' A cordial powder, known under the name 'Countess of Kent's Powder,' composed of coral, amber, crabs' eyes, prepared pearls, &c. It was given in cancer.

CANTION, *Candi*.

CAN'ULA, *Can'nula*, *Au'licus*, *An'lus*, (dim. of *canna*, 'a reed;') *Tu'bulus*, (F.) *Canule* ou *Canule*. A small tube of gold, silver, platinum, iron, lead, wood, elastic gum, or gutta percha, used for various purposes in surgery.

CA'OUTCHOUC. The East India name for *Indian Rub'ber*, *Elast'ic Gum*, *Gum Elast'ic*, *Gum'mi elast'icum*, *Ca'uchuc*, *Resi'na elast'ica* seu *Cayenne'sis*, *Cayenne' Re'sin*, *Ca'ut'chuc*, (F.) *Gomme élastique Résine élastique* ou de *Cayenne*. A substance formed from the milky juice of *Hæ'vea* seu *Hé'vea Guianen'sis*, *Jat'tropa elast'ica*, *Sipho'nia Ca'huchu*, *S. elast'ica*, *Seringe Tree*, and of *Ficus Ind'ica*, and *Artocar'pus integrifo'lia*:—South American trees. It is insoluble in water and alcohol; but boiling water softens and swells it. It is soluble in the essential oils and in ether, when it may be blown into bladders. It is used in the fabrication of catheters, bougies, pessaries, &c.

CAP, **PITCH**, see *Depilatory*.

CAPACITY, **LUNG**, see *Vital Capacity*—c. *Pulmonic*, see *Vital Capacity*.

CAPA-ISIAKKA, *Bromelia ananas*.

CAPBERN, **MIN'ERAL WATERS OF**. Capbern is in the department Hautes-Pyrénées, France. The waters contain sulphates and carbonates of lime and magnesia, and chloride of magnesium. Temperature, 75° Fahrenheit. They are purgative.

CAPELET, *Myrtus carophyllata*.

CAPELL'NA, *Capelli'na*, (F.) *Capeline*; A Woman's Hat, in French; *Capistrum*, (*caput*, 'head.') A sort of bandage, which, in shape, resembles a riding-hood. There are several kinds of *Capelines*:—1. That of the head, *C. de la tête*. *Fascia capita'lia*; see *Bonnet d'Hippocrate*. *C. of the clav'icle*, employed in fractures of the acromion, clavicle, and spine of the scapula. *C. of an amputated limb*—the bandage applied round the stump.

CAPELL'NA, *Capelina*.

CAPER BUSH, *Capparis spinosa*.

CAPER PLANT, *Euphorbia lathyris*.

CAPERS, see *Capparis spinosa*.

CAPHORA, *Camphor*.

CAPHURA, *Camphor*.

CAPILLAIRE, (from *capillus*.) *Capillary*, see *Adiantum capillus veneris*—c. *du Canada*, *Adiantum pedatum*—c. *de Montpellier*, *Adiantum capillus veneris*.

CAPILLAMEN'TUM, (from *capillus*.) *Capil-lit'ium*, *Tricho'na*, *Trichoma'tion*. Any villous or hairy covering. Also, a small fibre or fibril.

CAP'ILLARY, *Capilla'ria*, *Capilla'ceus*, (from *capillus*.) (F.) *Capillaire*. Hair-like; small.

CAPILLARY VES'sELS, *Va'sa capilla'ria*, *Micrang'i'a*, *Trichang'i'a*, (F.) *Vaisseaux capillaires*, *Capillicules*, are the extreme radicles of the arteries and veins, which together constitute the *cap'illary*, *interme'diate*, or *periph'eral vascular sys'tem*—the *methæ'mata* or *methæ'matous blood-channels* of Dr. Marshall Hall. They possess an action distinct from that of the heart, but not one of rhythmic contraction and dilatation. Formerly, it was conceived that white vessels—*Vasa sero'sa*—or vessels so minute as not to admit red corpuscles, were distributed to the tissues which do not receive red blood.

CAPILLATIO, *Trichismus*.

CAPILLATUS, *Impuber*.

CAPILLICULES, *Capillary vessels*.

CAPILLITIUM, *Capillamentum*, *Entropion*, *Scalp*.

CAPILLORUM DEFLUVIUM, *Alopecia*.

CAPIL'LUS, (quasi *capitis pilus*.) *Co'ma*, *Chæ'te*, *Cr'in'is*, *Pi'lus*, *Thrix*, *Cæsa'ries*, (F.) *Cheveu*. This term is generally applied to the hair of the head, *Pi'li* seu *Ho'nor cap'itis*, the characters of which vary according to races, individuals, &c., see *Hair*.

CAPIL'US VENERIS, *Adiantum capillus veneris*—c. *V. Canadensis*, *Adiantum pedatum*.

CAPIPLE'NIUM, *Capitple'num*, (*caput*, gen. *capitis*, and *plenum*, 'full.') A word employed with different significations. A variety of catarrh. A heaviness or disorder in the head common at Rome, like the *καρῆβια*, *Carebi'ria* of the Greeks.

CAPISTRATIO, (*capistrum*, 'a halter, a muzzle.') *Phimosi*.

CAPISTRUM, *Capeline*, *Chevestre*, *Trismus*—c. *Auri*, *Borax*.

CAPIS'TRUM, *Phi'mos*, *Ce'mus*. This name has been given to several bandages for the head. See *Capeline*, *Chevestre*.

CAPITALIA REMEDIA, (from *caput*.) *Cephalic remedies*.

CAPITEL'TUM. In some of the older writers, soapy water; in others, a lixivium.

CAPITELLUM, (dim. of *caput*.) *Alembic*, see *Caput*.

CAPITEUX, *Heady*.

CAPITULU'VIUM, (*caput*, and *luere*, 'to wash.') A bath for the head.

CAPITPLENIUM, *Capiplenium*.

CAPITIPURGIA, *Caput purgia*.

CAPITIT'RAHA, (*caput*, and *trahere*, 'to draw.') Instruments which, like the forceps, draw down the head of the fetus when impacted in the pelvis.

CAPITO'NES, (from *caput*, 'the head.') *Macroceph'ali*, *Proceph'ali*. Fetuses whose heads are so large as to render labour difficult. The condition is called *Macrocephalia*.

CAPITULUM, (dim. of *caput*, 'head.') *Alembic*, *Condyle*, see *Caput*—c. *Costæ*, see *Costa*—c. *Laryngis*, *Corniculum laryngis*—c. *Martis*, *Eryngium campestre*—c. *Santorini*, *Corniculum laryngis*.

CAPITULUVIUM, (*caput*, and *luere*, 'to wash.') *Bath (head)*.

CAPNISMA, (καπνισμα, from καπνός, 'smoke.') *Fumigation*.

CAPNITIS, *Tutia*.

CAPNOÏDES CAVA, *Fumaria bulbosa*.

CAPNORCHIS, (*capnus*, and ορχις, 'a testicle,' owing to its bulbous roots.) *Fumaria bulbosa*

CAPNUS, (καπνός, 'smoke.') *Fumaria*.

CAPON, *Cagot*.

CAPON SPRINGS. A pleasant summer retreat, situated in a gorge of the North Mountain, in Hampshire co., Va., 23 miles W. of Winchester.

The waters in the vicinity are sulphurous and chalybeate;—those at the springs alkaline and diuretic.

CAPOT. *Cagot.*

CAPPARIS, (καππαρίς,) *Cap'paris*, *C. spinosa*, *Cap'par*, *Ca'pria*, *Pric'ly Ca'per Bush*, (F.) *Capprier*. *Ord.* Capparidæ. *Sex. Syst.* Polyandria Monogynia. The bark of the root and the buds have been esteemed astringent and diuretic. The buds are a well-known pickle—*Capers*, (F.) *Câpres*.

CAPPARIS BADUC'CA, *Baduk'ka*. A species of caper, cultivated in India on account of the beauty of its flowers. The Orientals make a liniment with its juice, with which they rub pained parts. The flowers are purgative.

CAPPO'NE, MIN'ERAL WA'TERS OF. At Capponne, in the isle of Ischia, are waters containing carbonate of soda, chloride of sodium, and carbonate of lime. Temp. 100° Fah.

CAPREOLA'RES, (*capreolus*, 'a tendril.') *Cis-oi'des*, *Elico'i'des*, (F.) *Capréolaire*. Twisted.

CAPREOLA'RIA VA'SA. Some have called thus the spermatic arteries and veins, on account of their numerous contortions.

CAPREOLUS, ('a tendril,') *Helix*.

CÁPRES, see *Capparis spinosa*.

CAPRIA, *Capparis spinosa*.

CAPRICORNUS, *Plumbum*.

CÁPRIER, *Capparis spinosa*

CAPRIFOLIA, (*caper*, 'a goat,' and *folium*, 'a leaf,') *Lonicera periclymenum*.

CAPRIFOLIUM DISTINCTUM, *Lonicera periclymenum*—*c.* *Periclymenum*, *Lonicera periclymenum*—*c.* *Sempervivens*, *Lonicera sempervivens*,—*c.* *Sylvaticum*, *Lonicera periclymenum*.

CAPRILOQUIUM, (*caper*, 'a goat,' and *loqui*, 'to speak,') *Egophony*.

CAPRIZANS PULSUS, (from *caprizare*, 'to leap,' like *caper*, 'a goat,') see *Pulse*, *caprizant*.

CAPSA, (καψα, 'a chest,') *Boîte*, *Capsule*, *Case*—*c.* *Cordis*, *Pericardium*.

CAPSARIUM, *Boitier*.

CAPSELLA BURSA PASTORIS, (dim. of *capsea*,) *Thlaspi bursa*.

CAPSICUM, see *Capsicum annuum*.

CAPSICUM AN'NUUM, *C. Hispan'icum*, (καπ'ω, 'I bite,') The systematic name of the plant, *Order*, Solanæ. whence, in part, *Cayenne' Pep'per*, *Cap'sicum*, (Ph. U. S.) is obtained,—*Pé'per In'dicum* seu *Hispan'icum* seu *Brazilia'num* seu *Guineen'se* seu *Calec'ticum* seu *Tur'cium* seu *Lusitan'icum*, *Solanum v'rens*, *Siliquas'trum Plin'ii*, *Cayenne'*, *Guin'ea*, *Chilli* or *Chilly Pep'per*, *Gar'den Gin'ger*, (F.) *Piment*, *Poivre d'Inde*, de *Guinée* ou d'*Es-pagne*, *Corail des Jardins*. The British Pharmacopœia (1864) assigns it to *Capsicum fastigiatum*. The pungent aromatic properties of *Baccæ Cap'sici*, *Capsicum Ber'ries*, known in commerce as *Guin'ea pep'per*, *Pod pep'per*, *Chill'ies*, are yielded to ether, alcohol, and water. They are highly stimulant and rubefacient, and are used as a condiment. Their active principle is called *Cap'sicin*.

CAPSICUM HISPANICUM, *Capsicum annuum*.

CAPSIQUE, *Capsicum annuum*.

CAPSTITIS, (*capsea*, and *itis*,) see *Phacitis*.

CAPSULA, (dim. of *capsea*,) *Boitier*, *Capsule*—*c.* *Articularis*, *Capsular ligament*—*c.* *Cordis*, *Pericardium*—*c.* *Dentis*, *Dental follicle*—*c.* *Hæmatica*, *Capsule*, *hæmatic*—*c.* *Humoris aquei*, see *Aqueous humour* of the eye—*c.* *Lentis*, see *Crystalline*—*c.* *Nervorum*, *Neurilemma*—*c.* *Sequestralis*, see *Sequestrum*.

CAPSULÆ SEMINALES, *Vesiculæ S.*—*c.* *Synoviales*, *Bursæ mucosæ*.

CAPSULAIRE, *Capsular*.

CAP'SULAR, *Capsula'ris*, (F.) *Capsulaire*. Relating to a capsula or capsule.

CAPSULAR AR'TERIES, *Supraren'al Ar'teries and Veins*. Vessels belonging to the suprarenal capsules. They are divided into superior, middle, and inferior. The first proceed from the inferior phrenic, the second from the aorta, and the third from the renal artery. The corresponding veins enter the phrenic, vena cava, and renal.

CAPSULAR CATARACT, see *Cataract*.

CAPSULAR LIG'AMENT, *Ligamen'tum capsula'rē*, *Cap'sula articula'ris*, *Articular* or *Fib'rous cap'sule*, (F.) *Ligament capsulaire*, *Capsule articulaire*, *Capsule fibreuse*, &c. Membranous, horous, and elastic bags or capsules, of a whitish consistence, thick, and resisting, which surround joints.

CAP'SULE, *Cap'sula*, *Cap'sa* ([L.] 'a box, or case,') (F.) *Capsule*. This name has been given, by anatomists, to parts bearing no analogy to each other.

CAPSULE ATRABILLIAIRE, *Capsule*, renal—*c.* *Bowman's*, see *Kidney*—*c.* *Cellular*, of the Eye, see *Eye*—*c.* *Fibrous*, *Capsular ligament*.

CAPSULE, GELAT'INOUS, *Cap'sula gelati'na*, *Collocyst'is*, (F.) *Capsule gélatineuse*, *Capsule of gelatin*. A modern invention by which copaiba, disagreeable pills, &c., can be enveloped in gelatin so as to conceal their taste.

CAPSULE OF GLIS'SON, *Cap'sula GLISSO'NI* seu *communis GLISSO'NI*, *Vagi'na Por'tæ*, *V. GLISSO'NI*. A sort of membrane, described by Glisson, which is nothing more than dense areolar membrane surrounding the vena porta and its ramifications in the liver.

CAPSULE, HEMAT'IC, *Cap'sula hemat'ica*, (F.) *Capsule hémétique*. A gelatinous capsule, containing an extract made from the blood of the calf, sheep, or ox; and proposed to be given where there seems to be a deficiency of blood corpuscles—as in chlorosis.

CAPSULE OF THE HEART, *Pericardium*—*c.* *Ocular*, see *Eye*.

CAPSULE, RE'NAL, *Supraren'al* or *Atrabiliary* *C.*, *Re'nal Gland*, *Glan'dula supraren'al'is*, *Cap'sula renal'is* seu *supraren'al'is* seu *atrabil'is*, *Ren succenturiat'us*, *Nephrid'ium*, *Paraneph'rus*, (F.) *Capsule surrénale ou atrabillaire*, (pl.) *Reins succenturiés* ou *succenturiaux*. A flat triangular body, which covers the upper part of the kidney, as with a helmet. A hollow cavity in the interior contains a brown, reddish or yellowish fluid. The renal capsules were long supposed to be the secretory organs of the fancied atrabillis. They are much larger in the fetus than in the adult. They are probably concerned in lymphosis.

A singular condition of cachexia, the leading characteristics of which are anæmia, general languor and debility; remarkable feebleness of the heart's action; irritability of the stomach, and a peculiar bronzed skin, was first described by Dr. Thos. Addison, of London, as connected with a diseased condition of the supra-renal capsules. It has been proposed to call it the *Disease of Addison*, *Melasma supraren'al'e*, (F.) *Maladie d'Addison* ou *bronzé*.

CAP'SULE, SEM'INAL, *Cap'sula Semina'lis*. **BARTHOLINE** thus designates the extremity of the vas deferens, which is sensibly dilated in the vicinity of the vesiculæ seminales. Some anatomists apply this name to the vesiculæ themselves.

CAPSULE SURRÉNALE, *Capsule*, renal.

CAPSULE, SYNO'VIAL, *Cap'sula Synovia'lis*. A membranous bag, surrounding the movable articulations and canals, which gives passage to tendons. Synovial capsules exhale, from their articular surface, a fluid, whose function is to favour the motions of parts upon each other. See *Bursæ mucosæ*, and *Synovia*.

CAPSLITIS, (*capsula*, and *itis*,) see *Phacitis*.

CAPUCHON, ('a hood,') *Trapezius*.

CAPUCINE, *Tropæolum majus*.

CAPULI, *Prunus capulin*.

CAPULUM. A contortion of the eyelids or other parts.

CAPULUS, ('the handle of anything.') Penis.

CAPUT, gen. *Cap'itis*, ('the head.') Also, the top of a bone or other part, (F.) *Tête*. The head of small bones is sometimes termed *capitulum*, *cephalidium*, *cephalis*, *cephalum*. Also, the glans penis.

CAPUT ASPERÆ ARTERIÆ, Larynx—c. Coli, Cæcum—c. Gallinaceum, see Gallinaginis caput—c. Gallinaginis, see Gallinaginis caput—c. Genitale, Glans—c. Lubricum, Penis—c. Major, see Epididymis—c. Minor, see Epididymis—c. Monachi, Leontodon Taraxacum—c. Obstipum, Torticollis—c. Penis, Glans.

CAPUT PUR'GIA, *Capitipurgia*. Remedies, which the ancients regarded as proper for purging the head:—*errhines*, *sternutatories*, *apophlegmatisantia*, &c. Prosper Alpinus makes the *caput purgia* to be the same as *errhines*; and the *apophlegmatisantia* the same as the *masticatories* of the moderns.

CAPUT SCAPULÆ, Aeromion.

CAPUT SUCCEDANEUM, ('a succedaneous head.')

A term sometimes used for the tumefied scalp, which first presents in certain cases of labour.

CAPUT TESTIS, Epididymis.

CAQUE SANGUE, *Cague-Sangue*, (*cacare*, 'to go to stool,' and *sanguis*, 'blood.')

Old French words which signify *Bloody evacuations*, (F.) *Déjections sanguinolentes*. Under this term was comprehended every affection in which blood is discharged from the bowels.

CARA, (*kapa*.) Head.

CARA SCHUL'LI, *Fru'tex Indicus spinosus*, *Barle'ria buxifolia*. A Malabar plant, which, when applied externally, is maturative and resolvent. The decoction of its root is used, in the country, in ischuria.

CARABAC'CIUM. An aromatic wood of India, of a yellowish colour, and a smell like that of the clove. Its decoction and infusion are given as stomachics and antiscorbutics.

CARABE, Asphaltum.

CAR'ABUS, (*kapaβos*.) A genus of coleopterous insects. Two species, the *chrysocephalus* and *ferruginus* have been recommended for the toothache. They must be pressed between the fingers, and then rubbed on the gum and tooth affected.

CARACTÈRE, Character, Symbol.

CARAGNA, Caranna.

CARAMA'TA, *Arum'ri*. A tree in the inland parts of Pomeroun. It furnishes a febrifuge bark, which Dr. Hancock says may be used in typhoid and remittent fevers where cinchona is either useless or pernicious.

CARAMBOLO, Averrhoa carambola.

CAR'AMEL, *Car'omel*, *Sac'charum peroc'tum seu tos'tum*. Sugar subjected to the action of heat, until it is partly decomposed, deliquescent, of a brown colour, and a strong, agreeable and empyreumatic odour.

CARAN'NA, *Caragna*, *Tacamaha'ca Caragna*, *Caran'ne Gum'mi*, *G. Brel'isis*, *Gum Caran'na*, (F.) *Caragne*, *Gomme Caragne* on *Carane*. A gum-resinous substance, which flows from a large tree in New Spain, and is obtained from South America in impure masses. It preserves its softness for a long time, has an aromatic smell, and a slightly acid and bitter taste. It was formerly used as a vulnerary and in plasters.

CARA'PA GUANEN'SIS, *Persoon'ia guareo'ides*, *Xylocarpus cara'pa*, Ord. Meliaceæ. The bark of this South American tree is bitter and astringent; and has been used as an antiperiodic, the virtues depending upon a peculiar bitter principle, *Car'apin*. The oil of the nuts, *O'leum Cara'pe*, is bitter and anthelmintic; and is poisonous to insects.

CARAPA TOULOUOUNA has a bark possessed of like properties, from which a bitter resinoid substance has been obtained, *Toulououna nin*.

CARAPAT, *Ricinus communis* (oil of).

CARAWAY, Carum.

CARAWAY SEEDS, see Carum.

CARBASA, Linteam.

CARBASUS, (*καρσος*.) Linteam.

CARBO, gen. *Carbo'nis*, Carbon, (F.) *Carbone*. An elementary body, extensively distributed in nature, and one of the principal elements of organized bodies. It occurs in different conditions; and the two following forms are official in the Pharmacopœia of the United States. Also, a live coal; charcoal; and anthrax.

CARBO ANIMA'NIS, *C. carnis*, *An'imal char'coal*, (F.) *Charbon animal*, is usually prepared by subjecting bones to a red heat in close vessels. The result is *Bone black*, commonly called *I'vory black*. (F.) *Noir animal ou d'Os*. It is given in the same cases as *Carbo Ligni*, and has been extolled in cancer. Dose, gr. ss to gr. iij.

The Ph. U. S. contains a formula for the preparation of **CARBO ANIMA'NIS PURIFICA'TUS**, *Purified an'imal char'coal* (*Carbon animal*, bj; *Acid muriat.*, *Aque* aa f 3xij). The muriatic acid, previously mixed with the water, is digested with a gentle heat for two days. The undissolved portion is allowed to subside, the supernatant liquor is poured off, and the charcoal, frequently washed with water until it is entirely free from acid, is dried.

CARBO FOS'SILIS, *Lithan'thrax*, Stone coal.

CARBO HUMA'NUM. The human excrement.—*Paracelsus*.

CARBO LIG'NI, *Charcoal*, (F.) *Charbon*. Fresh charcoal is antiseptic. It is used to improve the digestive organs in cases of worms, dyspepsia, &c.; as a cataplasm to gangrenous and fetid ulcers, tinea, &c., and forms a good tooth-powder. Dose, gr. x to 3j.

CARBO LIGNI, Carbo—c. Mineralis. Graphites—c. Palpebrarum, Anthracosis—c. Spongiæ, *Spongia usta*—c. Trichloratus, Carbonis trichloridum.

CARBON, Carbo—c. Bisulphuret of, Carbonis sulphuretum—c. Sesquichloride of, Carbonis trichloridum—c. Sesqui-iodide of, Carbonis sesquiodidum—c. Sulphide of, Carbonis sulphuretum—c. Sulphuret of, Carbonis sulphuretum—c. Terchloride of, Carbonis trichloridum, see Chloroform.

CARBO'NAS, *Car'bonate*, (F.) *Carbonate*. A generic name for a salt, formed by the combination of carbonic acid with a salifiable base.

CARBONAS NATRICUM, *Sodæ carbonas*—c. Plumbeus, Plumbi subcarbonas.

CARBONATE D'AMMONIAQUE, *Ammonia carbonas*.

CARBONATED, *Carbonatus*, *Aërat'us*, (F.) *Carboné*, *Aéré*. That which is impregnated with carbonic acid.

CARBONE, Carbon—c. Trichlorure de, Carbonis trichloridum.

CARBONÉ, Carbonated.

CARBONEUM CHLORATUM, Chloroform—c. Trichloratum, Carbonis trichloridum.

CARBON'IC ACID, *Ac'idum Carbo'nicum*, *Solid Air of Hales*, *Factit'ious* or *Fixed Air*, *Carbonaceous* or *Calcareous* or *Aë'rial* or *Mephit'ic* *Ac'id*, *Mephit'ic Air*, *Spir'itus bath'is*, (F.) *Acide Carbonique*. This gas, which neither supports respiration nor combustion, is not often used in medicine. It is the main agent in effervescent draughts, fermenting poultices, &c. It is often found occupying the lower parts of mines—when it is called the *choke damp*—caverns, tombs, wells, brewers' vats, &c., and not unfrequently has been the cause of death. Lime thrown into such places soon absorbs the acid.

CARBONIC OXIDE, *Carbo'neum Oxyda'tum*, (F.) *Oxyde de Carbone*. The gas resulting from passing carbonic acid over red hot charcoal or metallic iron. It is positively deleterious, and is often the cause of injurious phenomena, which have been ascribed to the inhalation of carbonic acid. When directed on a denuded surface, it produces anæsthetic effects, similar to those of chloroform, but it is not used.

CARBONIS SESQUICHLORIDUM, Carbonis trichloridum.

CARBONIS SESQUI-IODIDUM, *C. Sesqui-Iodure'tum*, *Sesqui-iodide* or *Sesqui-iod'uret* of Carbon. This is made by mixing concentrated alcoholic solutions of iodine and potassa, until the former loses its colour; a solution is obtained from which water throws down a yellow precipitate—the sesqui-iodide of carbon. It has been used in enlarged glands and in some cutaneous affections, applied externally, (3ss to ʒvj of cerate). See Iodoform.

CARBONIS SULPHURETUM, *Sulph'uris Carbu-retum*, *Sulphure'tum* seu *Sul'fidum Carbo'nii*, *Carbo'nium Sulphura'tum*, *Al'cohol Sul'phuris*, *Bisulphure'tum Carbo'nii*, *Sulphuret*, *Sulphide* or *Bisulphuret* of Carbon, *Carburet* of Sulphur, (F.) *Sulfure de Carbone*, *Carbure de Soufre*, *Soufre Carburé*, *Alcool de Soufre*, *Liquueur de Lampadins*. This transparent, colourless fluid, which has a very penetrating, disagreeable odour, and a taste which is cooling at first, but afterwards acrid and somewhat aromatic; is a diffusible excitant. It is diaphoretic, diuretic, and is said to have proved emmenagogue. It is also used in nervous diseases as an antispasmodic. Dose, one drop to four, repeated frequently.

It is used externally, where a cooling influence has to be rapidly exerted, and has been inhaled as an anæsthetic.

CARBONIS TRICHLORIDUM, *C. Sesquichloridum*, *Carbo'neum trichlora'tum*, *Carbo trichlora'tus*, *Terechloride* or *Sesquichloride* of Carbon, (F.) *Trichlorure* ou *Sesquichlorure de Carbone*, is formed by the action of Chlorine on Chlorohydric ether, under the influence of sun-light. It has been given in cholera, and applied to correct the fetor of foul ulcers. Dose, four grains or more.

CARBONIUM SULPHURATUM, Carbonis sulphuretum.

CARBUNCLE, (dim. of *carbo*, 'a live coal.') Anthrax—c. Berry, Terminusus—c. Fungus, Terminusus.

CARBUNCLE OF THE TONGUE, Glossanthrax.

CARBUNCLED FACE, Gutta rosea.

CARBUNCULAR EXANTHEM, Anthracia.

CARBUNCULATIO OCULI, Blepharanthra-
xis.

CARBUNCULUS, (dim. of *carbo*, 'a live coal.') Anthrax—c. Anginosus, Cynanche maligna—c. Contagiosus, see Anthrax—c. Galliens, see Anthrax—c. Hungaricus, see Anthrax—c. Labiorum et genarum, Cancer aquaticus—c. Polonicus, see Anthrax—c. Pulmonum, Necropneumonia.

CARBUNCULUS RUB'NUS. A red, shining, and transparent stone, from the Isle of Ceylon; formerly employed in medicine as a preservative against several poisons, the plague, &c.

CARBUNCULUS SEPTENTRIONALIS, see Anthrax—c. Uleusculosus, Cynanche maligna.

CARBURE DE SOUFRE, Carbonis sulphuretum.

CARBURETTED HYDROGEN, see Hydrogen, carburetted.

CAR'CAROS, (*καρκας*, from *καρκαίρω*, 'I rebound,' 'I tremble.') A fever, in which the patient has a general tremor, accompanied with an unceasing noise in the ears.

CARCINELCO'SIS, (*carcinus*, and *ἄλωσις*, 'ulceration.') Cancerous ulcer.

CARCINODES, (*καρκινώδης*, from *carcinus*, and *odes*.) Canceroid, *Chancreuse*.

CARCINOIDES, (*carcinus*, and *ειδός*, 'resemblance.') Canceroid.

CARCINO'MA, gen. *Carcino'matis*, (*καρκινώμα*.) *Karkino'ma*, *Cancero'ma*, *Cancro'ma*, (F.) *Carcinome*, (*καρκίος*, cancer, and *μα*.) Some authors have called thus indolent tumours different from cancer; others, incipient cancer; and others, again, the species of cancer in which the affected structure assumes the appearance of cerebral substance; but the majority of authors use Carcinoma in the same sense as Cancer.

CARCINOMA ALVEOLARE, Colloid—c. Epithelioides, Epithelial Cancer—c. Fibrosum, Scirrhus—c. Hamatodes, Hamatodes fungus—c. Intestinorum, Enteropathia cancerosa—c. Lingue, Glos-socarcinoma—c. of the Liver, Hepatoscirrhos—c. Medullare, Encephaloid—c. Melanodes, Cancer, melanotic—c. Melanoticum, Cancer, melanotic—c. Scroti, Cancer, chimney-sweepers'—c. Simplex, Scirrhus—c. Spongiosum, Encephaloid, Hamatodes fungus—c. Uteri, Metrocarcinoma, Metro-scirrhos—c. Ventriculi, Gastroscirrhus; see Gastrostenosis cardiaca et pylorica.

CARCINOM'ATOUS, *Carcinomato'eus*, (F.) *Carcinomateux*. Relating to Carcinoma.

CARCINOME, Carcinoma—c. Mou et Spon-gieux, Encephaloid—c. Sanglant, Encephaloid, Hamatodes fungus.

CARCINO'SES, (G.) *Karcinosen*, (from *carcinus* and *osis*.) A family of diseases, according to the classification of Fuchs; which embraces the different forms of Cancer.

CARCINOSUS, Cancerous.

CARCINUS, (*καρκίος*.) Cancer—c. Spongiosus, Encephaloid.

CARDAMANTICA, Cardamine pratensis, Lepidium Iberis.

CARDAMINE FONTANA, (*καρδαμυν*.) *Sisymbrium nasturtium*—c. *Nasturtium*, *Sisymbrium nasturtium*.

CARDAMINÆ PRATEN'SIS, *Cardam'inæ*, *Carda-man'tica*, *Nasturt'ium Aquaticum*, *Card'amom*, *Card'amum*, *Cu'li flos*, *Ibe'ris soph'ia*, *Nasturt'ium pratens'e*, *Ladies-smock*, *Cuckoo-flower*, *Common Bitter Cress*, *Meadow Cress*, (F.) *Cresson élégant* ou *des prés*, *Passerage sauvage*. Ord. Cruciferae. The flowers have been considered useful as anti-spasmodics, in the dose of ʒi to ʒij. They are probably inert.

CARDAMINUM MINUS, *Tropæolum majus*.

CARDAMOM, (*καρδαμωμ*.) *Cardamo'mum* (F.) *Cardamome*. The name of the fruit of various species of Amomum and Elelettaria, respecting the botanical history of which there has been much confusion. In the Pharmacopœia of the United States, it is the fruit of *Elelett'ria Cardamo'mum*.

CARDAMOM, **BENGAL**, *Amomum maximum*—c. Cluster, *Amomum cardamomum*—c. Greater, *Amomum Grana Paradisi*—c. Java, *Amomum maximum*—c. Lesser, *Amomum cardamomum*—c. Nepal, *Amomum maximum*—c. Round, *Amomum cardamomum*.

CARDAMOME, *Amomum cardamomum*—c. *de la Côte de Malabar*, *Amomum cardamomum*.

CARDAMOMUM MAJUS, *Amomum grana paradisi*—c. Minus, *Amomum cardamomum*—c. Piperatum, *Amomum grana paradisi*—c. Rotundum, *Amomum cardamomum*—c. Wild, *Fagaras-trum Capense*.

CARDAMON, (*καρδαμωμ*.) Cardamine pratensis.

CARDAMON, Cardamine pratensis.

CARDÈRE, (*carder*, 'to card,') *Dipsacus syl-vestris*—c. *Cultivé*, *Dipsacus fullonum*.

CAR'DIA, (*καρδιά*, 'the heart.') *Stom'achus*, *Orific'ium sinis'trum* seu *Ingres'sus super'ior* seu *Os'tium œsophage'um ventric'uli*. The superior or

oesophageal orifice of the stomach. Also, the Heart.

CARDIAC, (*καρδιακός*, from *cardia*.) (F.) *Cardiaque*. Relating to the heart. *Cardi'ic*, *Card'ial*, (F.) *Cardinaire*, *Carditique*,—or to the upper orifice of the stomach. Also, a cordial.

CARDIAC ARTERIES, *Cardiacary arteries*, (F.) *Artères cardiaques* on *coronaires*, are two in number. They arise from the aorta, a little above the free edge of the sigmoid valves, and are distributed on both surfaces of the heart.

CARDIAC GANGLION, *Gan'glion cardi'acum*, situated beneath the arch of the aorta, to the right side of the ligament of the ductus arteriosus. It receives the superior cardiac nerves of opposite sides of the neck, and a branch from the pneumogastric, and gives off numerous branches to the cardiac plexuses.

CARDIAC NERVES, (F.) *Nerfs cardiaques*. These are commonly three on each side; a *superior*, *middle*, and *inferior*, which are furnished by corresponding cervical ganglia. Commonly, there are but two on the left side: the upper and middle, which draw their origin from the last two cervical ganglia. Searpa calls the *superior*—*Card'iacus superficial'is*; the *middle*—*C. profun'dus* seu *mag'nus*; and the *inferior*—*C. par'vus* seu *mi'nor*. There are, besides, *Cardiac fil'aments*, (F.) *Fil'eta cardiaque*, furnished by the par vagum or pneumogastric nerve, which become confounded with the above.

CARDIAC PLEX'US, *Plex'us cardi'acus*. There are three cardiac plexuses. 1. The *great cardiac plexus* is situated upon the bifurcation of the trachea. It is formed by the convergence of the middle and inferior cardiac nerves; and by branches from the pneumogastric, descendens noni, and first thoracic ganglion. 2. The *anterior cardiac plexus* is situated in front of the ascending aorta near its origin. It is formed by filaments from the superior cardiac nerves; from the cardiac ganglion; and from the great cardiac plexus. Filaments from this plexus accompany the left coronary artery, and form the *anterior coronary plexus*. 3. The *posterior cardiac plexus* is seated upon the posterior part of the ascending aorta near its origin. It is formed by numerous branches from the great cardiac plexus. It divides into two sets of branches, which together constitute the *posterior coronary plexus*.

CARDIAC VEINS, *Cor'onary veins*, (F.) *Veines Cardiaques*, are commonly four in number; two anterior and two posterior. Among the former—*venæ innomina'tæ* of Vieussens—there is one running along the right border of the heart, which has been called the *vein of Galen*. The cardiac veins open into the right auricle by one orifice, which is furnished with a valve, and is called, by Portal, *Sinus coronairæ du Cœur*.

CARDIACA CRISPA, *Leonurus cardiaca*—c. *Passio*, *Cardialgia*—c. *Trilobata*, *Leonurus cardiaca*—c. *Vulgaris*, *Leonurus cardiaca*.

CARDIACI (*Mor'bi*.) In the classification of Dr. Wm. Farr. means heart diseases.

CARDIACUS, *Cordial*, *Stomachal*.

CARDI'AGRA, *Affectio arthrit'ica cor'dis*; (*cardia*, and *agra*, 'seizure.') Gout or pain of the heart.

CARDIAGRAPHY, *Cardiagra'phia*, (*cardia*, and *γραφη*, 'a description.') An anatomical description of the heart.

CARDIAIRE, see *Cardiac*.

CARDIAL, see *Cardiac*.

CARDIAL'GIA, (*καρδιαλγία*) *Cardi'aca Pas'sio*, *Col'ica Ventric'uli*, *Spas'mus Ventric'uli*, *Pero'dyn'ia*, *Card'itium*, *Card'ia*, *Dyspepsody'n'ia*, *Dyspepiody'n'ia*, *Dyspeptody'n'ia*, *Peratody'n'ia*, *Cardiody'n'ia*, *Gastrody'n'ia*, *Gastral'gia*, *Gasteral'gia*, *Gastrocol'ica*, *Gastrod'yn'ia*, *Pas'sio Card'ia*, *Stomachal'gia*, *Card'iacus Mor'bus*, *Cardiog'mus*,

Cardi'alg'gy; (*cardia*, and *αλγος*, 'pain.') *Pain of the stomach*, F. *Douleur de l'Estomac*, D. *névral'gique de l'Estomac*. Also, *Heartburn*, (F.) *Cardi'algie*, *Ardour d'Estomac*, A. *du Cœur*. Impaired appetite, with gnawing or burning pain in the stomach or epigastrium,—*Mor'bus seu ard'or ventric'uli*, *Morbus stom'achi*, *Sol'da*, *Linn'æa cardi'al'gia mor'dens*, *Rob'io Stom'achi seu Ventric'uli*—a symptom of dyspepsia.

CARDIALGIA INFLAMMATORIA, *Gastritis*—c. *Sputatoria*, *Pyrosis*.

CARDIALOG'IA, (*cardia*, and *λογος*, 'a discourse.') A treatise on the heart.

CARDIANASTROPHE, (*cardia*, and *αναστροφή*, 'a turning round.') *Ectopia cordis*.

CARDIANEURYSMA, *Aneurism of the heart*.

CARDIARTIE, *Heart*, concentric hypertrophy of the. Contraction of the cavities of the heart, (*cardia*, and *αετο*, 'I constrict.')

CARDIA'RIOUS, (from *cardia*.) A name given to a worm, said to have been found in the heart or pericardium.

CARDIASTHMA, *Asthma cardiac*.

CARDIATELIA, *Atelocardia*.

CARDIATOMIA, (*cardia*, and *τομή*, 'to cut.') Dissection of the heart.

CARDIATROPHIA, *Heart atrophy of the*.

CARDIAUXE, (*cardia*, and *αἴξη*, 'increase.') *Heart hypertrophy of the*.

CARDIECHEMATIA, (*cardia*, and *ἡχημα*, 'a sound.') See *Heart (sounds)*.

CARDIECTASIS, (*cardia*, and *εκτασις*, 'extension.') Dilatation of the heart, see *Aneurism of the heart*—c. *Partialis*, *Aneurism of the ulcer*.

CARDIELCOSIS, (*cardia*, and *έλκος*, 'an ulcer.') Ulceration of the heart.

CARDIETHMOLIPOSIS, (*cardia*, *ἡθμος*, 'a sieve,' and *λίπος*, 'fat.') *Steatosis cordis*.

CARDIEURYSMA, (*cardia*, and *εuryσμα*, 'dilatation.') *Aneurism of the heart*.

CARDILÆA, *Cardialgia*.

CARDIM'ELECH, from (*cardia*, and *מלך*, *Melek*, (Hebr.), 'a governor.') A supposititious active principle seated in the heart, and governing the vital functions.

CARDINAL FLOWER, *Lobelia cardinalis*—c. f. *Blue*, *Lobelia syphilitica*.

CARDINAL PLANT, *Lobelia cardinalis*.

CARDINAMENTUM, (*cardo*, gen. *cardinis*, 'a hinge.') *Ginglymus*, *Gomphosis*.

CARDIOBOTANUM, *Centaurea benedicta*.

CARDIOCE'LE or *Car'diocele*, (*cardia*, and *κηλη*, 'rupture.') *Hernia of the heart*, especially into the abdominal cavity.

CARDIOCLASIE, (*cardia*, and *κλαω*, 'to break.') *Cardiorrhexis*.

CARDIODY'NE, *Cardiodyn'ia*; (*cardia*, and *δύνη*, 'pain.') *Pain in the heart*. Also, *Cardialgia*.

CARDIODYNE SPASMODICA INTERMITTENS, *Angina pectoris*.

CARDIOGM'US, (*καρδιωγμος*.) Hippocrates employed this synonymously with *cardialgia*. In the time of Galen it was used, by some writers, for certain pulsations of the heart, analogous to palpitations. Sauvages understood by *Cardiog'mus* an aneurism of the heart or great vessels, when still obscure. Also, *Angina pectoris*.

CARDIOGMUS CORDIS SINISTRI, *Angina pectoris*.

CARDIOMALA'CIA, *Malac'ia seu Mala'cia seu Malax'ia seu Mollit'ies Cor'dis*, (F.) *Ramol'issement du Cœur*, (*cardia*, and *μαλακία*, 'softness.') Softening of the heart caused by inflammation of the organ, or a consequence of some lesion of the function of nutrition.

CARDIOMETER, see *Hæmodynamometer*.

CARDIOM'ETRY, *Cardiomet'ria*, (*ca'rdia*, and

μετρον, 'measure.') Measurement of the heart, as by percussion and auscultation.

CARDIOMYOLIPOSIS, (*cardia*, μυς, 'a muscle,' and λιπος, 'fat.') Steatosis cordis.

CARDIONCHIL, (*cardia*, and ογκος, 'a tumour.') See Aneurism.

CARDIONEURALGIA, Angina pectoris.

CARDION'OSUS, *Cardiopath'ia Mor'bus cor'dis*, (*cardia*, and νοσος, 'disease.') Disease of the heart. Heart disease.

CARDIOPALMUS, (*cardia*, and παλμος, 'palpitation.') Cardiotromus.

CARDIOPATHIA, (*cardia*, and παθος, 'affection.') Cardionosus.

CARDIOPERICARDITIS, see Pericarditis.

CARDIORRHEU'MA, gen. *Cardiorrh'eum'atis*, *Rheumatis mus cor'dis*; (*cardia*, and ρευμα, 'defluxion, rheumatism.') Rheumatism of the heart.

CARDIORRHEX'IS, (*cardia*, and ρηξις, 'laceration,') *Ruptu'ra cor'dis*, (F.) *Rupture du Cœur*, *Cardioclasis*. Laceration of the heart.

CARDIOSCLÉROSIE, (*cardia*, and σκληρος, 'hard,') (F.) *Endurcissement du Cœur*. Induration of the heart.

CARDIOTENO'SIS, *Stenocar'dia*, (*cardia*, and στενωσις, 'contraction.') Contraction of the openings of the heart.

CARDIOTRAU'MA, gen. *Cardiotraum'atis*, (*cardia*, and τραυμα, 'a wound.') A wound of the heart.

CARDIOT'ROMUS, *Palpita'tio Cor'dis trep'idans*, *Cardiopal'mus*, *Trepida'tio Cor'dis*, (*cardia*, and τρομος, 'tremor.') Rapid and feeble palpitation or fluttering of the heart.

CARDIOT'ROTUS, (*cardia*, and περισσῶ, 'I wound,') One affected with a wound of the heart.

CARDIPERICARDITIS, see Pericarditis.

CARDITE, Carditis.

CARDIT'IC, (F.) *Carditique*, Cardiac. Also, relating or belonging to carditis.

CARDIT'IS, (*cardia*, and *itis*.) Inflammation of the fleshy substance of the heart. *Empres'ma Card'itis*, *Inflamma'tio Cor'dis*, I. *Card'itis*, *Can'ma Card'itis*, *Mycocard'itis*, *Card'itis Muscula'ris*, (F.) *Inflammation du Cœur*, *Cardite*. The symptoms of this affection are by no means clear. They are often confounded with those of pericarditis, or inflammation of the membrane investing the heart. Carditis, indeed, with many, includes both the inflammation of the external investing membrane and that of the interior of the heart. See Pericarditis, and Endocarditis.

CARDITIS EXTERNA, Pericarditis — c. Interna, Endocarditis — c. Membranosa, Pericarditis — c. Muscularis, Carditis — c. Polyposa, Polypi of the heart — c. Serosa, Pericarditis.

CARDO, gen. *Card'inis*, Ginglymus.

CARDOLEUM, see Anacardium occidentale.

CARDOPATHEUM, *Carlina acaulis*.

CARDOPERICARDITIS, see Pericarditis.

CARDUUS ATLILIS, (C. 'a thistle,') *Cynara scolymus* — c. *Benedictus*, *Centaurea benedicta* — c. *Brazilianus*, *Bromelia ananas* — c. *Domesticus capite majori*, *Cynara scolymus* — c. *Hæmorrhoidalis*, *Cirsium arvense*.

CARDEUS MARIA'NUS, *C. Ma'rie* seu *lacteus*, *Sily'bum*, *S. Maria'num* seu *maculatum*, *Carthamus maculatus*, *Cirsium maculatum*, *Spina al'ba*, *Common Milk Thistle*, *Ladies' Thistle*, (F.) *Chardon-Marie*. Ord. Composite. The herb is a bitter tonic. The seeds are oleaginous. It is not used.

CARDUUS PINEUS, *Atractylis gummifera* — c. *Sativus*, *Carthamus tinctorius* — c. *Sativus non-spinosus*, *Cynara scolymus* — c. *Solstitialis*, *Centaurea calcitrapa* — c. *Stellatus*, *Centaurea calcitrapa* — c. *Tomentosus*, *Onopordium acanthium* — c. *Veneris*, *Dipsacus fullonum*.

CARE (καρη), Head.

CAREBAREISIS, Carebaria.

CAREBA'RIA (καρηβαρια), *Carebure'is*, (καρη, 'the head,' and βαρος, 'weight,') *Scordine'ma*, *Scordinis'mus*, *Cordine'ma*. Heaviness of the head.

CARE'NA, *Kare'na*. Formerly, the twenty-fourth part of a drop.

CARE-TAKEER, Nurse.

CAREUM, Carum.

CAREX ARENARIA, (*carex*, 'sedge,') *Sarsaparilla Germanica*.

CARIACOU. A beverage, used in Cayenne, and formed of a mixture of cassava, potato, and sugar, fermented.

CAR'ICA, (from *Caria*, where the fig was erroneously supposed to be native.) See *Ficus carica*.

CARICA PAPA'RIA, *Papaw' tree*, *Panepaw*, (F.) *Papayer*. Ord. Cucurbitaceæ. A native of America, India, and Africa. The fruit has somewhat of the flavour of the pumpkin, and is eaten like it. The milky juice of the plant and the seed and root have been regarded as anthelmintic.

The papaw of North America belongs to the order Anonaceæ, (*Ano'na* seu *Asi'mina* seu *Porcell'ia* seu *Uca'ria tri'loba*, *Ficus In'dica*, *Orchidocar'pum ariet'inum*), Custard Apple Family.

CAR'ICUM (καρικον). A detergent application to ulcers: composed of black hellebore, sandarach, copper, lead, sulphur, orpiment, cantharides, and oil of cedar.

CARIE, Caries — c. *des Dents*, Dental gangrene — c. *Écorçante*, see Desoortication.

CARIE, Carious.

CA'RIES, ('decay,') *Carios'ity*, *Nigrit'ies* *Os'sium*, *Ulceration of bone*, *Osteohelen'is*, *Ost'hel'cus*, — *Necrosis* being death of a bone. It resembles the gangrene of soft parts. Hence it has been termed *Car'ies gangren'osa*, *Gangren'na Car'ies* seu *Os'sium*, *Tere'do*, *Arro'sio*, *Eu'rus*, (F.) *Carie*. It is recognized by the swelling of the bone which precedes and accompanies it; by the abscesses it occasions; the fistulæ which form; the sanious character, peculiar odour and quantity of the suppuration, and by the evidence afforded by probing. The most common causes of caries are blows, the action of some virus, and morbid diathesis. When dependent on any virus in the system, this must be combated by appropriate remedies. When entirely local, it must be converted, where practicable, into a state of necrosis or death of the affected part. For this end, stimulants, the actual cautery, &c., are applied.

CARIES DENTIUM, Dental gangrene — c. *Podend'orum*, see Chancre — c. of the Vertebra, Vertebral disease — c. *Vertebrarum*, Vertebral disease.

CARIEUX, Carious.

CARIM CURINI, *Justitia ecbolium*.

CAR'INA, 'a keel.' The vertebral column, especially of the foetus. Also, the breastbone bent inwards. Hence, *Pectus carina'tum* : — the chest affected with such deformity.

CARIOSITY, Caries.

CA'RIOUS, *Cario'sus*, *Enro'des*, (F.) *Carië*, *Cariour*. Affected with caries.

CARIUM TERRE, Calx.

CARIVE, *Myrtus pimenta*.

CARIVILLANDI, *Smilax sarsaparilla*.

CARLINA, (after Charlemagne,) 'Carline Thistle.'

CARL'INA ACAUL'IS, *C. chamæ'leon*, *Chamæ'leon al'bum*, *Cardopa'tium*, (F.) *Carline sans tige*. Ord. Composite, which grows in the Pyrenees, and on the mountains of Switzerland, Italy, &c., has been recommended as a tonic, emmenagogue, and sudorific.

CARLINA CHAMÆLEON, *C. acaulis*.

CARLINE SANS TIGE, *Carlina acaulis*.

CARLISLE' SPRINGS. These are about five

miles from the town of Carlisle, in Pennsylvania. The water is mildly sulphureous.

CARLO SANCTO RADIX. 'St. Charles's Root,' found in Mechoacan, in America. The bark is aromatic, bitter, and acrid. It is considered to be sudorific, and to strengthen the gums and stomach.

CARLSBAD, MIN'ERAL WATERS OF. Carlsbad is a town in Bohemia, 24 miles from Egria, celebrated for its hot baths. The water contains about 47 parts in the 100 of purging salts. It is a thermal saline; temperature 121° to 167° Fahrenheit. The constituents are—carbonic acid, sulphate of soda, carbonate of soda, and chloride of sodium. Season from the first of June to the 15th of October.

Artificial Carlsbad water may be made of *Sodæ sulph.*, 3 gram.; *Sodæ carb.*, 35 centigram.; *Calcii chlorid.*, 4 decigram.; *Sodii chlorid.*, 4 decigram.; *Ferri sulph.*, 1 centigram.; *Aquæ carbonat.* 625 gram.

CARMANTINE, *Justitia pectoralis*—c. *Pectorale*, *Justitia pectoralis*.

CARMEN, gen. *Car'minis*, 'a verse.' An amulet. A charm, which, of old, often consisted of a verse. See Charm.

CARMINANTIA, (from *carmen*.) Carminatives. **CARMINATIVA**, Carminatives.

CARMINATIVES, *Carminantia* seu *Carminativa*, *Antiphytica*, *Physoga'ga*, *Antiphyset'ics*, (F.) *Carminatifs*. Remedies which allay pain, 'like a charm,' by causing the expulsion of flatus from the alimentary canal. They are generally of the class of aromatics.

THE FOUR GREATER CARMINATIVE HOT SEEDS, *Quatuor sem'ina cal'ida majo'ra carminativa*, were, of old, anise, carui, cummin, and fennel.

THE FOUR LESSER CARMINATIVE HOT SEEDS, *Quatuor sem'ina cal'ida mino'ra*, were bishop's weed, stone parsley, smallage, and wild carrot.

CARMOT. A name given, by the alchemists, to the matter which they believed to constitute the Philosopher's stone.

CARNABADIA, *Carum*, (seed.)

CARNABADIUM, *Cuminum cuminum*.

CARNATIO, (*caro*, gen. *carnis*, 'flesh.') *Sy-sarcosis*.

CARNATION, *Dianthus caryophyllus*.

CARNELIAN, *Cornelian*.

CARNEOLUS, *Cornelian*.

CAR'NEOUS, *Carno'us*, *Sarco'des*, *Incarna'tus*, (from *caro*.) (F.) *Charnu*. Consisting of flesh, or resembling flesh.

CARNEOUS COLUMNS, see *Columnæ carneæ*.

CARNEOUS FIBRES, *Flesh'y F'ibres*, *Mus'cular F'ibres*, (F.) *Fibres charnues ou musculaires*, are fibres belonging to a muscle.

CARNEUM MARSUPIUM, ('fleshy pouch,') *Ischiotrochanterianus*.

CARNIC'ULA, (dim. of *caro*, 'flesh.') The gum.

CARNIFICATIO, Carnification—c. *Pulmonum*, Hepatisation of the lungs.

CARNIFICA'TION, *Carnificatio*, (*caro*, and *feri*, 'to become,') *Transformatio into flesh*. A morbid state of certain organs, in which the tissue acquires a consistence like that of fleshy or muscular parts. It is sometimes observed in hard parts, the texture becoming softened, as in *Osteosarcoma*. When it occurs in the lungs, they present a texture like that of liver. Such is the condition of the fetal lung.

CARNIFOR'MIS ABSCESS'US, (*caro*, and *forma*, 'form,') An abscess, which ordinarily occurs in the neighborhood of the articulations, and whose orifice is hard, the sides thick and callos.

CARNIV'OROUS, *Carniv'orus*, *Sarcoph'agus*, *Creatoph'agus*, *Creoph'agus*, *Zoöph'agus*, *Creatoph'agus*, *Creoph'agus*, *Kreatoph'agus*, (F.)

Carnivore; (*caro*, and *voro*, 'I eat,') That which eats flesh. Any substance which destroys excrescences in wounds, ulcers, &c.

CARNOSA CUTIS, ('fleshy skin,') *Panniculus carnosus*.

CARNOS'ITY, *Carnos'itas*, (F.) *Carnosité*, (from *caro*.) A fleshy excrescence.

CARNOS'ITIES OF THE URE'THRA, *Car'unclis in the Ure'thra*, (F.) *Carnosités ou Caroncles de l'urètre*. Small fleshy excrescences or fungous growths, which were, at one time, presumed to exist in the male urethra, whenever retention of urine followed gonorrhœa.

M. Cullérius uses the term *Carnosité vénérienne* for a cutaneous, cellular, and membranous tumour, dependent upon the syphilitic virus. See, also, *Polysarcia*.

CARNOSUS, *Carneous*.

CARO, gen. *Carnis*, *Flesh*—c. *Accessoria*, see *Flexor longus digitorum pedis profundus perforans*, (*accessorius*)—c. *Anserina*, *Horrida cutis*—c. *Contusa*, *Sarcothlasis*—c. *Excrescens*, *Excrescence*—c. *Fungosa*, *Fungosity*—c. *Gallinacea*, *Horrida cutis*—c. *Glandulosa*, *Epiglottic gland*—c. *Luxurians*, *Fungosity*—c. *Orbicularis*, *Placenta*—c. *Parenchymatica*, *Parenchyma*—c. *Quadrata*, *Palmaris brevis*—c. *Quadratus Sylvii*, see *Flexor longus digitorum pedis profundus perforans*, (*accessorius*)—c. *Viscerum*, *Parenchyma*.

CAROB TREE, *Ceratonia siliqua*.

CAROBA ALNABATI, *Ceratonia siliqua*.

CARODES, (*carus*, and *odes*), *Carotic*.

CAROLI, see *Chancre*.

CAROLI'NA, NORTH, MIN'ERAL WATERS

OF. In the counties of Warren, Montgomery, Rockingham, Lincoln, Buncomb, and Rowan, there are mineral springs. They belong generally to the sulphureous or acidulous saline.

CAROLI'NA, SOUTH, MIN'ERAL WATERS **OF.** They are numerous. Pacolet Springs, on the west bank of Pacolet River, contain sulphur and iron. Many, with similar properties, but not held in estimation, are scattered about the State.

CARONCULE, *Caruncle*—c. *Lachrymale*, *Caruncle*.

CARONCULES MYRTIFORMES, *Carunculæ myrtiformes*—c. *de l'Urètre*, *Carnosities of the urethra*.

CARONCULEUX, *Carunculous*.

CAROPI, *Amomum cardamomum*.

CAROSIS, (*xaposis*), *Carus*.

CAROTA, ('a carrot,') See *Daucus carota*.

CAROT'IC, *Carot'icus*, *Carot'id*, *Carot'idus*, *Carotidæ'us*, *Caro'des*, *Com'atose*, (*xapōs*, 'stupor,') (F.) *Carotique*. Relating to stupor or *carus*—as a *carotic state*—or to the carotids, (F.) *Carotidiën*.

CAROTIC ARTERIES, *Carotids*—c. *Ganglion*, see *Carotid Nerve*—c. *Nerve*, *Carotid Nerve*—c. *Plexus*, see *Carotid Nerve*.

CAROTICA, (*xapōtika*), *Narcotics*.

CAROTICUS, *Carotic*.

CAROTID, *Carotic*.

CAROTIDÆUS, *Carotic*.

CAROTIDIËN, *Carotic*.

CAROTIDS, *Carot'idæ*, *Carot'icæ*, *Carotidæ'us*, (*xapōs*, 'stupor,' produced by pressing on them.[?])

Capita'les, *Jugula'les*, *Sopora'les*, *Sopora'ria*, *Soporif'era*, *Somnif'era*, *Apople'ticæ*, *Lethargicæ* (*Arter'ia*), *Carot'id Ar'teries*, *Cephal'ic Ar'teries*, (F.) *Artères Carotides*. The great arteries of the neck, which carry blood to the head. They are divided into, 1. *Prim'itive* or *common*; the left of which arises from the aorta, and the right from a trunk, common to it and the subclavian. 2. *Exter'nal* or *pericephal'ic*, a branch of the primitive, which extends from the last to the neck of the condyle of the lower jaw; and, 3. *Inter'nal*, *Ar'teria cerebr'al'is* seu *encephal'ic*, another branch of the primitive, which, arising at the same place as the external, enters the cranium,

and terminates on a level with the fissures of Sylvius, dividing into several branches.

CAROTID OF CAROT'IC CANAL, *Canalis Carot'icus*, *Canal inflexe de l'os temporal*—(Ch.), *Canal carotidien*, is a canal in the temporal bone, through which the carotid artery and several nervous filaments pass.

CAROTID OF CAROT'IC FORAM'INA, *Foram'ina Carot'ica*, (F.) *Trous carotidiens*, are distinguished into *internal* and *external*. They are the foramina at each extremity of the *Canalis Carot'icus*.

CAROTID GANGLION, see Carotid nerve.

CAROTID NERVE, *Carot'ic nerve*, *Nervus carot'icus*. A branch from the superior cervical ganglion of the great sympathetic, which ascends by the side of the internal carotid. It divides into two portions, which enter the carotid canal, and, by their communication with each other and the petrosal branch of the vidian, form the *carotid plexus*. They also frequently form a small gangliform swelling on the under part of the artery—the *carotic* or *carotid* or *cavernous ganglion*, *ganglion of Læmmonier*.

CAROTID PLEXUS, see Carotid nerve.

CAROTIQUE, Carotic.

CAROTTE, Daucus carota.

CAROUA, Carum (seed).

CAROUBIER, Ceratonium siliqua.

CAROUGE, see Ceratonium siliqua.

CAR'PAL, *Carpa'lis*, (from *καρπος*, 'the wrist.') Belonging or relating to the carpus or wrist.

CARPASA, Carbasa.

CARPASIUM, *Car'pason*, *Car'pasum*, and *Carpe'sium*. Dioscorides, Pliny, Galen, &c., have given these names, and that of *καρπασος* to a plant, which cannot now be determined, and whose juice, called *Opocar'pason*, *σποκαρπασον*, passed for a violent, narcotic poison, and was confounded with myrrh.

CARPATHICUM, see Pinus cembra.

CARPE, Carpus.

CARPENTARIA, Achillea millefolium.

CARPENTER'S LEAF, Galax aphylla.

CARPESIMUM, Carpasium.

CARPHO'DES, *Carphoi'des*, (*καρφος*, 'flocculus,' and *odes*.) Flocculent, stringy;—as *mucus carphodes*, flocculent or stringy mucus.

CARPHOLOG'IA, *Til'mus*, *Carpholog'ia*, *Crocidis'mus*, *Crocydis'mus*, *Flocco'rum vena'tio*, *Floccile'gion*, *Tricholog'ia*, *Crocidis'is*, *Floccila'tion*, *Floccita'tion*, (*καρφος*, 'flocculus,' and *λεγω*, 'I collect,' or 'pluck,') (F.) *Carphologie*. Action of gathering flocculi. A delirious picking of the bed-clothes, as if to seek some substance, or to pull the flocculi from them. It denotes great cerebral irritability and debility, and is an unfavorable sign in fevers, &c.

CARPHOS, (*καρφος*, 'a light dried substance,') *Trigonella fœnum*.

CARPIA, (*carpere*, 'to pluck,') *Linteum*.

CARPIÆUS, Palmaris brevis.

CAR'PIAL, *Car'pian*, *Carpia'nus*, *Carpia'lis*, (F.) *Carpien*. Belonging to the Carpus.

CAR'PIAL LIG'AMENTS, (F.) *Ligaments Carpiens*, are, 1, The fibrous fasciæ, which unite the bones of the carpus; and, 2, The annular ligaments, anterior and posterior.

CARPIAN, Carpial.

CARPIEN, Carpial.

CARPISMUS, Carpus.

CARPOBALSAMUM, see Amyris opobalsamum.

CARPOLOGIA, Carphologia—c. Spasmodica, Subsultus tendinum.

CARPOMETACARPEUS MINIMI DIGITI, Adductor metacarpi minimi digiti—c. *Métacarpien du petit doigt*, Opponens minimi digiti—c. *Métacarpien du ponce*, Opponens pollicis—c. *Phalangæus minimi digiti*, Abductor minimi digiti—c. *Phalangien du petit doigt*, Abductor minimi digiti,

Flexor parvus minimi digiti—c. *Phalangien du ponce*, *Flexor brevis pollicis manus*—c. *Susphalangien du ponce*, *Abductor pollicis brevis*.

CARPOPE'DAL, from *carpus*, and *pes*, gen. *pedis*, 'the foot.' Relating to the wrist and foot.

CARPOPEDAL SPASM, *Cer'ebral spasmod'ic croup*. A spasmodic affection of the chest and larynx in young children, accompanied by general or partial convulsions. The disease commonly occurs between the third and ninth month, and is characterized by excessive dyspnœa, accompanied by a loud croupy noise on inspiration; the thumbs being locked, and the hands and feet rigidly bent for a longer or shorter period. The seat of the disease is evidently in the cerebro-spinal axis, primarily or secondarily: generally, perhaps, it is owing to erethism seated elsewhere, but communicated to the cerebro-spinal centre, and reflected to the respiratory and other muscles concerned. It seems to be connected with dental irritation; and consequently, in the treatment, where such is the case, the gums should be freely divided; after which, cathartics and revulsives, with the use of narcotics and appropriate diet, will generally remove the affection; for although extremely alarming, it is often not attended with great danger. See *Asthma thymicum*.

CARPOPHALANGÆUS MINIMI DIGITI, Abductor minimi digiti.

CARPOS, (*καρπος*,) Fruit.

CARPOT'ICA, (*καρπος*, 'fruit,') Diseases affecting impregnation. Irregularity, difficulty or danger produced by parturition:—the 3d order, class *Genetica*, of Good.

CARPUS, (*καρπος*,) *Carpis'mus*, *Brachii'læ*, *Rasce'ta*, *Raste'ta*, *Ras'cha*, *Rasce'ta*, *Rasce'ta*, *Wrist*, (Prov.) *Hand-wrist*, *Shack'le*, (F.) *Carpe*, *Poignet*. The part between the fore-arm and hand. Eight bones compose it (in two rows). In the superior row there are, from without to within—the *Scaphoïdes* or *navicula'rê*, *Luna'rê* or *semi-luna'rê*, *Cuneifor'mê*, and *Orbicula'rê* or *pisifor'mê*. In the lower row—*Trape'zium*, *Trapezoi'des*, *Magnum* and *Uncifor'mê*.

CARRAGEEN MOSS, (from Carragaheen, in Ireland,) *Fucus crispus*.

CARRAWAY, Carum.

CARRÊ, ('square,') *Quadratus*—c. *de la Cuisse*, *Quadratus femoris*—c. *des Lombes*, *Quadratus lumborum*—c. *du Menton*, *Depressor labii inferioris*—c. *du Pied*, *Extensor brevis digitorum pedis*.

CARREAU, *Tabes mesenterica*.

CARRÉE, see *Flexor longus digitorum pedis profundus perforans* (accessorius).

CARRELET, (F.) *Acus triangula'ris*. A straight needle two or three inches long, the point of which is triangular; and which was formerly used in different operations. Also, a wooden, triangular frame for fixing a cloth through which different pharmaceutical preparations are passed.

CARRON OIL, *Linimentum aquæ calcis*.

CARROT, Daucus carota—c. *Candy*, *Athamanta cretensis*—c. *Deadly*, *Thapsia*.

CARROVAL, see *Curare*.

CARTHAGENA BARKS, see *Cinchona*.

CARTHAMUS MACULATUS, *Carduus marianus*.

CAR'THAMUS TINCTO'RIOUS, *Am'yron*, *Cn'cus*, *Cro'cus German'icus* seu *Siracen'icus*, *Car'thamum officina'rum*, *Car'duus sat'vus*, *Saff'ronum*, *Saff'ron-flower*, *Saff'flower*, *Saff'flow*, *Bas'tard Saff'ron*, *Dy'er's Saff'ron*, (F.) *Carthame*, *Safran bâlard*, *Carthame des Teinturiers*. Ord. Compositæ. Sex. Syst. Syngenesia Polygamia æqualis. The seeds are aromatic, cathartic, and diuretic; yet to the parrot they are an article of food; hence their name, *Graines de Parroquet*. The flowers, *Car'thamus* (Ph. U. S.), are employed as a cosmetic, and are reputed to be diaphoretic. [?]

CARTHEGON, see *Buxus*.

CARTILAGE, *Chondros*, *Cartilago*, *Gristle*, (Sc.) *Gristle*, (F.) *Cartilage*. A solid part of the animal body, of a medium consistence between bone and ligament, which in the fœtus is a substitute for bone, but in the adult exists only in the joints, at the extremities of the ribs, &c. Cartilages are of a whitish colour, flexible, compressible, and very elastic, and some of them apparently inorganic. They are composed, according to Dr. J. Davy, of .44 albumen, .55 water, and .01 phosphate of lime.

CARTILAGE, Cartilage—c. *Anonyme*, Cricoid, (cartilage)—c. *Epiglottic*, Epiglottis—c. *Mucroné*, Xiphoid Cartilage—c. *Supra-arytenoid*, Corniculum laryngis—c. *Tarsal*, see Tarsus.

CARTILAGES, ARTICULAR, *Obducent Cartilages*, invest bony surfaces, which are in contact; hence they are called *investing* or *incrustating cartilages*, (F.) *Cartilages de revêtement ou d'encroûtement*.

CARTILAGES, CUNEIFORM, OF THE LARYNX, see Cuneiform—c. *Eburnés*, see Eburnification.

CARTILAGES, INTERARTICULAR, *Menisci*, are such as are situate within the joints, as in the knee joint.

CARTILAGES, INTERVERTEBRAL, see Intervertebral.

CARTILAGES OF OSSIFICATION are such as, in the progress of ossification, have to form an integrant part of bones; as those of the long bones in the new-born infant. They are termed *temporary*; the others being *permanent*. All the cartilages, with the exception of the articular, are surrounded by a membrane analogous to the periosteum, called *Perichondrium*.

CARTILAGES OF THE RIBS are, in some respects, only prolongations of the ribs. Those of the nose, of the meatus auditorius, and Eustachian tube, present a similar arrangement. Other cartilages resemble a union of fibrous and cartilaginous textures; hence their name *Fibrocartilages*.

CARTILAGES OF SANTORINI, see Corniculum Laryngis—c. *Semilunar*, see Semilunar—c. *Sigmoid*, Semilunar cartilages—c. of Wrisberg, Cuneiform Cartilages of the Larynx.

CARTILAGINES GUTTURALES, (pl. of *cartilago*,) Arytenoid cartilages—c. *Semilunares*. Semilunar cartilages—c. *Sigmoides*, Semilunar cartilages.

CARTILAGINIS ARYTENOIDÆ CAPITULUM, Corniculum laryngis.

CARTILAG'INOUS, *Cartilagin'eus*, *Cartilagin'ous*, *Chondro'des*, *Chondroï'des*, (F.) *Cartilagineux*. Belonging to, or resembling, cartilage.

CARTILAGINOUS TISSUE, see Tissue.

CARTILAGO, gen. *Cartilag'inis*, Cartilage—c. *Clypealis*, Thyroid cartilage—c. *Ensiformis*, Xiphoid cartilage—c. *Gutturalis*, Arytenoid cartilage—c. *Innominata*, Cricoid—c. *Mucronata*, Xiphoid cartilage—c. *Peltalis*, Thyroid cartilage, Xiphoid cartilage—c. *Peltatus*, Thyroid cartilage—c. *Scutiformis*, Thyroid cartilage—c. *Triticea*, see Thyrohyoid membrane—c. *Uvifer*, Uvula—c. *Xiphoides*, Xiphoid cartilage.

CAR'UM, (from Caria, in Asia.) *A'pium* seu *Bu'nium* seu *Ligusticum* seu *Se'seli car'vi* seu *car'rum*, *S'i'um car'vi*, *Car'rum*, *Car'um car'vi*, *Car'vi*, *Cumi'nium pratense*, *Car'na*, *Car'ron*, the *Car'away*, *Car'away*, (F.) *Carvi*, *Cumin des prés*. Order, Umbelliferae. Sex. Syst. Pentandria Digynia. The fruit or seeds, *Car'away seeds*, (Prov.) *Car'vy seeds*, *Carnaba'dia*, *Car'rona*, *Car'rum*, *Car'away* (Ph. U. S.), *Car'ni* (Ph. B.), are carminative. Dose, gr. x to ʒij, swallowed whole or bruised. The oil, *O'leum Car'ui*, (F.) *Huile de carvi*, has the properties of the seeds. Dose, gtt. ij to vj. *A'qua Car'ui*, *Car'away Water*, was official in the Pharmacopœias of London and Dublin.

CARUM BULBOCASTANUM, *Bunium bulbocastanum*.

CAR'UNCLE, *Carun'cula*, (dim. of *caro*, 'flesh,') A small portion of flesh, *Sar'cium*, *Sarcid'ium*. A fleshy excrescence, — *Ephy'ma carun'cula*, (F.) *Caruncule*.

CARUNCLES IN THE URETHRA, see Carnosities.

CARUNCULA, Caruncle.

CARUN'CUA LACRYMA' LIS, (F.) *Caruncule lacrymale*. A small, reddish, follicular body, situate at the inner angle of the eye. It secretes a gummy substance.

CARUNCULA SEMINALIS, Gallinaginis caput.

CARUNCULE CUTICULARES, (pl. of *Caruncula*,) *Nymphæ*.

CARUNCULE MAMMILLA' RES. The extremities of the lactiferous tubes in the nipples. The olfactory nerves have been so called by some.

CARUNCULE MYRTIFORMES, *C. Vagina'les*, *Glandulæ myrtiformes*, (F.) *Caruncules myrtiformes*. Small, reddish tubercles, more or less firm, of variable form, and uncertain number, situate near the orifice of the vagina, and formed by the mucous membrane. They are regarded as the remains of the hymen.

CARUNCULE PAPILLARES, Papillæ of the kidney.

CARUN'CULOUS, *Carun'cular*, (F.) *Carunculeux*. Relating to caruncles or carnosities.

CARUON, Carum.

CAR'US, (*καρὸς*,) *So'por caro'ticus*, *Caro'ris*, *Profound sleep*. The last degree of coma, with complete insensibility, which no stimulus can remove, even for a few instants. *So'por*, *Co'ma*, *Lethargia*, and *Car'us*, are four degrees of the same condition.

CARUS APOPLEXIA, Apoplexy—c. *Asphyxia*, *Asphyxia*—c. *Catalepsia*, *Catalepsy*—c. *Ecstasis*, *Ecstasis*—c. *Hydrocephalus*, *Hydrocephalus internus*—c. *ab Insolatione*, *Coup de soleil*—c. *Lethargus*, *Lethargy*—c. *Lethargus cataphora*, *Somnolency*—c. *Lethargus vigil*, *Coma vigil*—c. *Paralysis*, *Paralysis*—c. *Paralysis paraplegia*, *Paraplegia*—c. *Veternus*, *Lethargy*.

CARVI, Carum.

CARVY SEEDS, see Carum.

CARYA, (*καρπας*,) Hickory, *Juglans regia*—c. *Basilica*, *Juglans regia*.

CARYDION, (*καρυδίων*,) *Corylus avellana*.

CARYEDON CATAGMA, (*καρυδων*, 'resembling a nut' [broken],) see Fracture.

CARYOCOST'INUS, *Caryocostinum*. An electroly prepared of the costus and other aromatic substances, &c. It was cathartic. See Confectio scammonia.

CARYON, (*καρυον*,) see *Juglans regia*—c. *Ponticon*, *Corylus avellana* (nut.)

CARYOPHYLLA, (*καρυον*, 'a nut,' and *φύλλον*, 'a leaf,' from the appearance of the flower-buds.) *Geum urbanum*.

CARYOPHYLLATA AQUATICA, *Geum rivale*—c. *Nutans*, *Geum rivale*—c. *Urbana*, *Geum urbanum*—c. *Vulgaris*, *Geum urbanum*.

CARYOPHYLLUM, *Eugenia caryophyllata*—c. *Rubrum*, *Dianthus caryophyllus*.

CARYOPHYLLUS, see *Eugenia caryophyllata*—c. *Americanus*, see *Myrtus pimenta*—c. *Aromaticus*, *Eugenia caryophyllata*—c. *Hortensis*, *Dianthus caryophyllus*—c. *Pimenta*, *Myrtus Pimenta*—c. *Vulgaris*, *Geum urbanum*.

CARYO'TI, (from *καρυωτις*, 'a date tree,') The best kind of dates.

CAS RARES (F.), 'Rare cases.' This term is used, by the French, for pathological facts, which vary from what is usual. See a celebrated article under this head in the *Dictionnaire des Sciences Médicales*, Vol. IV.

CASAMUM, Cyclamen.

CASAMUNAR, Cassumunar.

CASCADING, Vomiting.

CAS'CARA, CASCARILLA. Spanish words, which signify *bark* and *little bark*, under which appellations the bark (Cinchona) is known in Peru. They are now applied to the bark of *Croton cascarilla*. The bark-gatherers are called *Cascarilleiros*.

CASCARILLA, *Cascara*, *Croton cascarilla*—c. *Carabaya*, see *Cinchona cordifolia* cortex—c. *Colorado*, *Cinchona oblongifolia* cortex—c. *Roxa*, *Cinchona oblongifolia* cortex.

CASCARILLEROS, see *Cascara*.

CASCARILLIN, see *Croton cascarilla*.

CASCHEU, *Catechu*.

CASE, *Cap'sa*, *The'ca*, (F.) *Caisse*. This name is given to boxes for the preservation of instruments, or of medicines necessary in hospital or other service. We say, e. g.—*A case of amputating*, or of *trepanning instruments*. A *pocket case*, *Armamentarium portabile*, (F.) *Trousse*, contains the smaller instruments in constant use with the surgeon.

CASE, *Ca'sus*, (from *cadere*, *casum*, 'to fall.') The condition of a patient;—as a *case of fever*, &c. (F. *Observation*). Also, the history of a disease. *To be in good case or condition* means to be stout or in a proper state of being. At times, *to be in case or condition* is used with a similar acceptation.

CASEARIUS, (*casus*, 'cheese,') *Cheesy*.

CASEIN, *Caseine*, *Ca'seum*, *Galac'tine*, *Ty'rine*, *Lactalbumen*, *Ca'seous matter*, (F.) *Caseïne*, *Matière caséuse*, (from *caseus*, 'cheese,') The great nitrogenized constituent of milk. It is identical in composition with the chief constituents of blood,—fibrin and albumen, all being compounds of protein. A similar principle exists in the vegetable, *Vegetable Casein* or *Legu'min*, *Veg'etable Gluten*. It is chiefly found in leguminous seeds—peas, beans, lentils. Like vegetable albumen, Casein is soluble in water; and the solution is not coagulable by heat.

CASEIN, BLOOD Globulin—c. of the *Saliva*, see *Saliva*—c. of the *Small intestine*, *Albuminose*.

CASEOSUS, *Cheesy*.

CASEOUS MATTER, *Casein*.

CASEUM, *Casein*.

CASEUS, *Cheese*—c. *Equinus*, *Hippace*.

CASEUX, (from *caseus*,) *Cheesy*.

CASHEW, (F.) *Acajou*, *Anacardium occidentale*—c. *Gum*, see *Anacardium occidentale*.

CASHOO. An aromatic drug of Hindostan, said to possess pectoral virtues.

CASHOW, *Catechu*.

CASIA, *Laurus cassia*.

CASMINA, *Cassumuniar*.

CASMONAR, *Cassumuniar*.

CASSA, *Thorax*.

CASSADA ROOT, *Jatropha manihot*.

CASSAVA, BITTER, *Jatropha manihot*—c. *Root*, *Jatropha manihot*.

CASSE AROMATIQUE, *Laurus cassia*—c. *en Bâtons*, *Cassia fistula*—c. *en Bois*, *Laurus cassia*—c. *des Boutiques*, *Cassia fistula*—c. *Séné*, *Cassia senna*.

CASSE-LUNETTE, (*casser*, 'to break,' and *lunettes*, 'spectacles,') *Centaura cyanus*, *Cyanus segetum*, *Euphrasia officinalis*.

CASSEENA, *Ilex vomitoria*.

CASSENOLLES, see *Quercus infectoria*.

CASSIA, (*casia*,) *Laurus cassia*—c. *Absus*, *Absus*—c. *Acutifolia*, *C. senna*—c. *Egyptian*, *C. senna*—c. *Alexandrina*, *C. fistula*—c. *Bonplandiana*, *C. fistula*—c. *Canella*, *Laurus cassia*—c. *Caryophyllata*, *Myrtus caryophyllata*.

CASSIA CHAMÆCHRIS'TA, *Præ'irie sen'na*, *Par'tridge Pea*, *Wild Sen'na*, an indigenous plant. *Order*, *Leguminosæ*, which flowers in August. It resembles *Cassia Marilandica* in properties.

CASSIA CINNAMOMEA, *Laurus cassia*—c. *Egyptian*, *Cassia senna*—c. *Excelsa*, *C. fistula*.

CASSIA FIS'TULA, *C. ni'gra* seu *fistula'ris* seu

Alexandri'na seu *excel'sa* seu *Bonplandia'na*, *Ca'na*, *C. soluti'va* seu *fis'tula*, *Cathartocar'pus*, *Bactyrilo'bium fis'tula*, *Purg'ing Cas'sia*, (F.) *Casse Canéficier*, *Canéficier*, *Casse en Bâtons* ou *des Boutiques*. The pulp of *Cas'sia Fis'tula* or *Cathartocar'pus Fis'tula*, *Pud'ding pipe tree*; *Fam. Leguminosæ*; *Sex. Syst. Decandria Monogynia*, *Pul'pa Cas'sia*, *Cas'sia Artamentum*, *Cassie Fis'tulæ Pul'pa*, (Ph. U. S.), which is obtained in long pods, is black, bright, and shining; sweet, slightly acid, and inodorous. It is laxative in the dose of ʒiv to ʒi.

CASSIA LANCEOLATA, *C. Senna*—c. *Ligneæ*, *Laurus cassia*—c. *Ligneæ Malabarica*, *Laurus cassia*.

CASSIA MARILAN'DICA, *Sen'na America'na*, *Americ'an or Wild Sen'na*, *Lo'cust Plant*, (F.) *Séné d'Amérique*. The leaves of this plant are similar, in virtue, to those of *cassia senna*. They are, however, much inferior in strength.

CASSIA NIGRA, *C. fistula*—c. *Officinalis*, *C. senna*—c. *Orientalis*, *C. senna*—c. *Purg'ing*, *Cassia fistula*.

CASSIA SEN'NA, *C. lanceolata*, seu *acutifolia*, seu *orientalis* seu *officinalis*. The name of the plant which affords *senna*. It is yielded, however, by several species of the genus *cassia*. The leaves of *Sen'na*, *Sen'na Folia*, *Sen'na Alexandri'na* seu *Ital'ica*, *Se'na*, *Sen'na* or *Egyptian Cas'sia*, (F.) *Séné*, *Casse Séné*, have a faint smell, and bitterish taste. The active part, by some called *Cathart'in*, is extracted by alcohol and water. Their activity is injured by boiling water. They are a hydragogue cathartic, and apt to gripe. Dose of the powder, ʒi to ʒi. Infusion is the best form.

The varieties of *senna*, in commerce, are *Tin-nivélly Senna*, *Bombay'* or *Com'mon In'dia Sen'na*, *Alexandrian Sen'na*, *Trip'oli Sen'na*, and *Alep'po Sen'na*.

CASSIÆ ARTAMENTUM, see *Cassia fistula*—c. *Fistulæ pulpa*, see *Cassia fistula*—c. *Flores*, see *Laurus cinnamomum*.

CASSIALA, *Hyssopus*.

CASSIDA GALERICULATA, *Scutellaria galericulata*.

CASSIDE BLEUE, *Scutellaria galericulata*.

CASSINA, *Ilex vomitoria*.

CASSINE CAROLINIANA, (an Indian name,) *Ilex paraguensis*—c. *Evergreen*, *Ilex vomitoria*—c. *Peragua*, *Ilex paraguensis*.

CASSIS, *Ribes nigrum*.

CASSITERUS, (*κασιτερος*,) *Tin*.

CASSUMUNIAR, *Cassamu'nar*, *Casmonar*, *Zerumbet*, *Casmina*, *Ri'sagon*, *Ben'galé In'dorum*, *Bengal' Root*, (F.) *Racine de Bengale*. A root, obtained from the East Indies, in irregular slices of various forms; some cut transversely, others longitudinally. It is an aromatic bitter, and is consequently tonic and stimulant. It was once considered a panacea, and has been referred to *Zin'giber Cassumu'niar* seu *purpu'reum*, *Amo'mum montanum*, and to *Zin'giber Zerum'bet* seu *spu'rium*, *Amo'mum Zerum'bet* seu *syloes'tre*.

CASSUVIUM POMIFERUM, *Anacardium occidentale*.

CAS'SYTA FILIFORM'IS. A South African plant, *Nat. Ord.* *Laurinæ*, which is employed by the Cape colonists as a wash in scald head, and as an antiparasitic.

CAST, *Contortio*, *Caste*—c. in the *Eye*, see *Strabismus*.

CASTALIA SPECIOSA, (from the *Castalian spring*), *Nymphaea alba*.

CASTANEA, (*καστανή*,) *Fagus castanea*: see, also, *Fagus castanea pumila*—c. *Equina*, *Æsculus Hippocastanum*—c. *Pumila*, *Fagus castanea pumila*.

CASTE, *Cast*, (P.) *Casta*, 'race or lineage,') A name given, by the Portuguese in India, 'to classes of society, divided according to occupa-

tions, which have remained distinct from the earliest times. Hence a separate and fixed order or class. See Half-caste.

CASTELLAMARE DI STABIA, MIN'ERAL WA'TERS OF. Castellamare di Stabia is a town in Naples, in the Principato Citra, 15 miles S. S. E. of Naples. There are two springs, the one sulphureous, the other chalybeate.

CASTELLETO ADONO, MIN'ERAL WA'TERS OF. These waters, situate near Aequi, in Italy, are sulphureous.

CASTERA-VIVENT, MIN'ERAL WA'TERS OF. Castera-Vivent is a small village in the department of Gers, near which is a cold acidulous chalybeate, and another which is sulphureous and thermal. Temp. 84° Fahrenheit.

CASTIGANS, (castigare, 'to correct.) Corrigent.

CASTIGLIONE'S A LOBA'TA, (after L. Castiglione, a traveller in S. America.) *Piñoncillo tree.* A tree, which is cultivated in some parts of Peru, and grows wild in abundance. Its beautiful fruit, when roasted, has an agreeable flavour. When an incision is made into the stem, a clear bright liquid flows out, which, after some time, becomes black and horny-like. It is a very powerful caustic.

CASTILLON POWDERS, (called after the proposer, see Powders, Castillon.

CASTING, Vomiting.

CASTJOE, Catechu.

CASTLE-LEOD, WA'TERS OF. A sulphureous spring in Ross-shire, Scotland, celebrated for the cure of cutaneous and other diseases.

CASTOR, see Castoreum—c. Bay, Magnolia glauca.

CASTOR FIBER, (καστωρ, quasi γαστωρ, from γαστρ, 'the belly,' because of the size of its belly. [?] Fib'er, C'ania Pon'ticus, the Be'aver, (F.) Castor. It furnishes the Castor. Rondelet recommends slippers made of its skin in gout. Its blood, urine, bile, and fat, were formerly used in medicine.

CASTOR OIL PLANT, Ricinus communis.

CASTO'REUM, Casto'rium, Cas'tor, Casto'reum Ros'sicum et Canaden'se. A peculiar concrete matter found in the follicles of the prepuce of the beaver, *Cas'tor fi'ber*. Its odour is strong, unpleasant, and peculiar; taste bitter, subacid; and colour orange brown. It is antispasmodic, and often employed. Dose, gr. x to ʒj.

CASTORI'NA, (from Castoreum.) Medicines containing castor.

CASTRANGULA, Scrophularia aquatica.

CASTRAT'. Castratus.

CASTRATIO, Castration.

CASTRAT'ION, Castrat'io, (castrare, castratum, (from castus, 'chaste,' [?]) Ec'tomē, Ectom'ia, Eirrat'io, Eecastrat'io, Ectescula'tio, Ectirpa'tio testiculo'rum, Detesta'tio, Esec'tio viril'ium, Eunuchis'mus, Orchotom'ia, Orchiotom'ia, Orchitom'ia, Orcheot'omy, Orchidot'omy, Geld'ing, Spay'ing, (F.) Châtrée. The operation of removing the testicles. Sometimes the term is employed for the operation when performed on one testicle: hence the division into complete and incomplete castration. Castration renders the individual incapable of reproduction.

The term *Spaying* is usually restricted to the removal of the ovaries.

CASTRATO, Castratus.

CASTRAT'US, (L.) Castra'to, Ec'tomus, Ectom'ius, Emacula'tus, Eirrat'us, Esec'tus, Desc'tus, Ectescula'tus, Ex ma'ribus, Intestab'ilis, Intesta'tus, Spa'do, Apoc'opus, (F.) Castrat, Châtré. One deprived of testicles. In the Southern States an animal thus situated is said to be al'tered. This privation has a great influence on the development of puberty. It is practised to procure a clearer and sharper voice; and in the East, the guardians of the harem, for the sake of security, are con-

verted into *Castra'ti* or *Eunuchs*. *Eunuchs* have generally both testes and penis removed.

CASUS, (cadere, casum, 'to fall, to happen,') Prolapsus, Symptom—c. Palpebræ superioris, Blepharoptosis—c. Uvulae, Staphylœdema.

CATFOOT, Gnaphalium polycephalum.

CATTAIL, Typha latifolia.

CATA, CATH, from kata, 'downwards'—'after,' applied to time: at times, it gives additional force to the radical word. A common prefix.

CATAB'ASIS, (καταβασις, from καταβαίνω, 'I descend,') An expulsion of humours downwards. Also, a descent, *Descen'sus, Descen'sio*,—as of the testicles, *Descen'sus testiculo'rum*.

CATABLE'MA, gen. Catable'matis, (καταβλημα, (from κατα and βαλλειν, 'anything let fall, as a curtain,') Epible'ma, Perible'ma. The outermost bandage which secures the rest.

CATABYTHISMOM'ANIA, (καταβυθισμος, 'submersion,') and mania, 'mania,') Insanity, with a propensity to suicide by drowning.

CATACASMUS, cata, and ακαειν, 'to prick,') Cupping, Searification.

CATACAUMA, gen. Catacaum'atis, (κατακαυμα, from cata, and καειν, 'to burn,') Burn.

CATACAUISIS, (κατακαυσις, 'Combustion, human—c. Ebriosa, Combustion, human.

CATACERASTICUS, (κατακεραστικός, from κατακεραυνω, 'I temper, 'I correct,') The same as *Epicerasticus*. A medicine capable of blunting acrimony of the humours.

CATACHRISIS, (καταχρσις, from cata, and χριω, 'I anoint,') Inunction.

CATACHRISTON, (χαταχριστον, 'Liniment.

CATACH'YSIS, (καταχυσις, from καταχυω, 'I pour upon,') Effu'sio, Perfu'sio. Affusion with cold water. Decantation.

CATAC'LASIS, (κατακλισις, from κατακλῶ, 'I break to pieces,') Distortion, or spasmodic fixation of the eyes; spasmodic occlusion of the eyelids, *Camp'ylum, Campylo'tis*. Also fracture of a bone.

CATACLEIS, gen. Cataclei'dis, (κατακλεις, from cata, and κλεις, 'the clavicle; 'a lock or fastening,') itself from κατακλειω, (κατα and κλειω, 'I lock up,') This term has been applied to many parts, as to the first rib, the acromion, the joining of the sternum with the ribs, &c.

CATACLEP'SIS, (κατακλεψις, 'A locking up. The act of locking up. Morbid union of the eyelids.

CATACLYS'MUS, Cataclys'ma, Cata'clysis, (κατακλυσμος, κατακλυσμα, κατακλυσις, from κατακλυζειν, 'to submerge, inundate,') A Clyster. Others mean, by the term, a shower-bath, or copious affusion of water; *Catane'msis, Ablution, Douche.*

CATEONESIS, (καταιονσις, from cata, and ιονω, 'to moisten,') Cataclysmus, Catantlema.

CATAGAUNA, Cambogia.

CATAGLOSSUM, (cata, and γλωσσα, 'the tongue,') Glossocatochus.

CATAGMA, gen. Catag'matis, (καταγμα, 'Fracture—c. Fissura, Fissure, see Contrafissura—c. Fractura, Fracture.

CATAGMAT'ICS, Catagmat'ica remed'ia, from Catagma. Remedies supposed to be capable of occasioning the formation of callus.

CATAGLOS'SUM, (καταγειν, 'to draw down, 'and γλωσσα, 'the tongue,') An instrument for pressing down the tongue. See Glossocatochus.

CATAGRAPHOLOGIA, (cata, γραφω, 'I write, 'and λογος, 'a discourse,') Pharmacocatagraphologia.

CATAIRE, (from catus, 'a cat, 'because cats are fond of it.) Nepeta, see *Frémissement cataire*.

CATALEN'TIA, Epilepsy, or some disease resembling it.

CATALEPSIA SPURIA, Ecstasis.

CATALEPSY, Catelep'sia, Catelep'is, (καταλαμβάνω, (cata, and λαμβανω, 'I seize hold of,') Cat'ochē, Cat'ochus, Cat'ochē Gale'ni, Mor'bus action

itus Col'si, Hystericus catalepticus, Congelatio, Dementia, Eucataleptosis, Aphonia (Hippocr.), *Anaëdia* (Antigenes), *Apprehensio, Contemplatio, Stupor vigilians, Prehensio, Carus Cataleptosis, Oppressio, Comprehensio* (Cæl. Aurelianus), *Comprehensio, Apprehensio Cataleptica, Trance*. (?) (F.) *Cataleptosis*. A disease in which there is sudden suspension of the action of the senses and of volition; the limbs and trunk preserving the different positions given to them. It is a rare affection, but is seen, at times, as a form of hysteria. Some of the Greek writers have used the word in its true acceptance of a seizure, surprise, &c.

CATALEPTIC, Catalepticus, (F.) Cataleptique. Relating to catalepsy. Affected with catalepsy.

CATALEPTIC METHOD, Methodus Cataleptica. The administration of external agents when internal agents are inapplicable.

CATALOTIC, Cataloticus, (cata, and aloos, 'I grind.') A remedy which removes unseemly cicatrices.

CATALPA, (name in India, C. Arboorea seu Cordifolia seu Arboreus cens seu Bignonioïdes seu Syringæfolia, Bignonina Catalpa, Catawba tree, Bean tree, Indian Bean. A decoction of the pods of the Catalpa, an American tree, of the Order Bignoniaceæ, Didymia Angiospermia, has been recommended in chronic nervous asthma.

CATALEPA ARBOREA, Catalpa—c. Bignonioïdes, Catalpa—c. Cordifolia, Catalpa—c. Syringæfolia, Catalpa.

CATALYSIS, Paralysis, (καταλυσις, from cata, and λυω, 'I dissolve or decompose.') Also, the action of presence in producing decomposition; as when a body which possesses what has been termed *catalytic force*—(F.) *Force catalytique*—resolves other bodies into new compounds by mere contact or presence, without itself experiencing any modification.

CATALYTIC, Catalyticus. A medicine, which is presumed by Dr. Headland to act by the destruction or counteraction of morbid agencies in the blood. See *Catalysis*.

CATALYTIC FORCE, see Catalysis.

CATAMENIA, (καταμηνια, from cata, and μην, 'a month.') Menses—c. Alba, Leucorrhœa.

CATAMENIAL, Catamenialis, Men'struis, Men'strual, Men'struous, (F.) Menstruel. Appertaining or relating to the catamenia.

CATAMENIORUM FLUXUS IMMODICUS, ('immoderate flow of the menses.') Menorrhagia.

CATANANCE, (κατανανχη,) Cichorium intybus. CATANGELUS, Ruscus.

CATANTLEMA, gen. Catantle-matis, Catantle-osis, (καταντλημα, καταντλησις, from cata, and αντλω, 'I pour.') Catcone'sis and Catcone'sis. Ablution with warm water. A fomentation.

CATAPASMA, gen. Catapasmatis, (καταπασμα, from cata, and πασσω, 'I sprinkle.') *Catapasmum, Conspersio, Pasma, Sympasma, Empasma, Diapasma, Xerion, Aspersio, Epispasmum, Pulvis aspersorius.* A compound medicine, in the form of powder, employed by the ancients to sprinkle on ulcers, absorb perspiration, &c.

CATAPASIS, (cata, and πασις, 'sedation.') Sedation.

CATAPHORA, (καταφορα, 'a fall,' from καταφερω, 'I throw down.') A state resembling sleep, with privation of feeling and voice. Somnolency. According to others, Cataphora is simply a profound sleep, which it is difficult to rouse from—in this sense being synonymous with Sopor.

CATAPHORA COMA, see Apoplexy—c. Hydrocephalica, see Apoplexy—c. Magnética, Somnambulism, magnetic.

CATAPHRACTA, Cataphractes, a Cuirass, (καταφρακτα, from καταφρασσω, 'I furnish with armor.') A name given by Galen to a bandage applied round the thorax and shoulders. It was also called *Quadriga*.

CATAPIESIS. (καταπιεσις, from cata, and πιειν, 'to press.') Depression.

CATAPINOSIS, (cata, and πινειν, 'to drink.') Absorption.

CATAPLASIS, (καταπλασις, from καταπλασσω, 'I besmear.') The act of besmearing or overlaying with plaster.

CATAPLASM, (καταπλασμα,) Cataplasma, Epiplasma, Poultice, Pulvis, (Prov.) Pulvey, (same etymon.) (F.) Cataplasme. A medicine applied externally, under the form of a thick pap. Cataplasms are formed of various ingredients, and for different objects. They may be *anodyne, emollient, tonic, antiseptic, irritating, &c.* A simple poultice acts only by virtue of its warmth and moisture. Mealy, fatty substances, leaves of plants, certain fruits, crumb of bread, &c., are the most common bases. The chief poultices which have been official are the following: *Anodyne—c. Cicute, c. Digitalis. Antiseptic—c. Carbonis, c. Dauci, c. Fermenti, c. Acetosæ, c. Cymini. Emollient—c. Lini, c. Panis, c. Mali maturi. Irritating—c. Sinapis, c. Sodii chloridi, c. Quercus Marinæ. Tonic and Astringent—c. Alum, c. Goulard, c. of Roses.*

The Parisian Codex had some other official cataplasms:—1. *Cataplasma ma amblynum*, made of poppy and hyoscyamus. 2. *Cataplasma ma emolliens*, made of meal and pulps. 3. *Cataplasma ad suppuratio-nem promovendam*, of pulps and basilicon. 4. *Cataplasma rubefaciens vel antipleuriticum*, formed of pepper and vinegar.

The only cataplasms, the preparation of which it is important to describe, are some of the following:—

CATAPLASM, ALUM, Coagulum Aluminosum—c. of Beer grounds, see *Cataplasma Fermenti—c.* Bread and milk, see *Cataplasma lini—c.* Carrot, *Cataplasma Dauci—c.* Charcoal, *Cataplasma carbonis ligni—c.* of Chlorinated Soda, *Cataplasma Sodæ chlorinatæ—c.* Flaxseed, *Cataplasma Lini—c.* Hemlock, *Cataplasma Conii—c.* Ice, see *Ice—c.* Ley, see *Lixivium—c.* Linseed, *Cataplasma Lini—c.* Oilcake, see *Cataplasma lini—c.* Mush, see *Cataplasma lini—c.* Slippery Elm, *Cataplasma Ulmi—c.* Yeast, *Cataplasma Fermenti.*

CATAPLASMA, gen. Cataplasmatis, Cataplasma—c. Bynes, see *C. Fermenti.*

CATAPLASMA CARBO'NIS, Char'coal Cat'aplasma or poultice, (F.) Cataplasme au charbon. Made by adding powdered charcoal to a common cataplasma. Used as an antiseptic to foul ulcers, &c.

CATAPLASMA CON'I, Hem'lock Cat'aplasma, made by spreading soft *Extract of Hemlock on a flaxseed poultice.* Used in malignant painful ulcers.

CATAPLASMA CYMINI, Theriaca Londinensis.

CATAPLASMA DAUCI, Car'rot Cat'aplasma or poultice. Made by boiling the root of the Carrot until it is soft enough to form a poultice. Used in fetid ulcers.

CATAPLASMA FÆCULÆ CEREVISIÆ, see C. Fermenti.

CATAPLASMA FERMENT'I, C. efferves'cens, Yeast Cat'aplasma or poultice, (F.) Cataplasme de Levure. (Take of wheaten flour, ℞j; yeast and water, ℞ā, f3v. Expose to a gentle heat.) It is antiseptic, and a good application to bruises. A *Cataplasma of Beer Grounds, Cataplasma Fæculæ Cerevisiæ, C. By'nes*, is used in the same cases.

CATAPLASMA LI'NI, Lin'seed or Flax'seed Cat'aplasma, (F.) Cataplasme de farine de Lin, Cataplasma commun, made by mixing *boiling water with linseed meal*, is an excellent emollient.

The *Bread and Milk cataplasma*, the *ground oilcake cataplasma*, and the *mush cataplasma* or *C. of Indian corn*, are used in similar cases.

CATAPLASMA SINA'PIS, C. Sinapi's, Sin'apism, Mustard cat'aplasma, poultice or plaster, (F.) Cataplasme de Mustard ou Sinapisme. (Mustard and linseed meal or meal, ℞ā, equal parts; warm vinegar or water, q. s.) A rubefacient and stimu-

lant applied to the soles of the feet in coma, low typhus, &c., as well as to the pained part in rheumatism, &c.

CATAPLASMA SODÆ CHLORINATÆ seu **CHLORATÆ**, *Cataplasm of Chlorinated Soda*. Made by adding solution of chlorinated soda to a common farsacæ cataplasm. Applied to foul, offensive ulcers.

CATAPLASMA ULMI, *Slippery Elm poultice*, is made by adding boiling water to the powdered bark of *Ulmus fulva*, stirring constantly. It is a light, agreeable cataplasm.

CATAPLASME, Cataplasm—*c. au Charbon*, Cataplasm carbonis ligni—*c. de Farine de Lin*, Cataplasm lini.

CATAPLEX'IS, *Stupor*, (καταληξίς, from *cata*, and *ληγναι*, 'I strike.') The act of striking with amazement. Appearance of astonishment, as exhibited by the eyes in particular. See *Hæmolia*.

CATAPOSIS, (καταποσις, from *cata*, and *πιναι*, 'to swallow.') Deglutition.

CATAPOTION, (same etymon,) Pihula.

CATAPSYX'IS, (καταψύξις, from καταψύχω, 'I refrigerate.') *Peripayx'is*. Considerable coldness of the body, without rigor and horripilation. *Perfrictio*. Coldness in the extreme parts of the limbs.

CATAPTO'SIS, (*cata*, and *πτειναι*, 'to fall.') *Decidentia*, 'a fall.' This word, at times, expresses the fall of a patient, attacked with epilepsy or apoplexy; at others, the sudden resolution of a paralytic limb.

CATAPULTA VIRILIS, ('male catapult,' καταπέλτης, from *cata*, and *παλλειν*, 'to throw.') Penis.

CATAPUTIA MAJOR, *Ricinus communis*—*c. Minor*, *Euphorbia lathyris*.

CATARACT, *Cataracta*, *Catarrhacta*, *Suffusio Oculi*, *S. Lentis crystallina*, *Phalarma cataracta*, *Caligo lentis*, *Gutta opaca*, *Hypochyma*, *Hypochymia*, *Hypophymia*, *Phacocystoma*, *Parophtalmia cataracta*, *Glaucoma Woulhouxi*, (F.) *Cataractæ*, (καταράκτες, from καταβάσσειν, (κατα and βάσσειν, 'to tumble down.') A deprivation of sight, which comes on as if a veil fell before the eyes. Cataract consists in opacity of the crystalline lens or its capsule, which prevents the passage of the rays of light, and precludes vision. The causes are obscure. *Diagnosis*.—The patient is blind, the pupil seems closed by an opaque body, of variable colour, but commonly whitish:—the pupil contracting and dilating. Cataracts have been divided, by some, into *spurious* and *genuine*. The former, *Pseudocataracta*, where the obstacle to vision is between the capsule of the lens and the *uvea*: the latter, where it is in the lens or capsule.

A *lenticular cataract* is where the affection is seated in the lens:—a *capsular* or *membranous*, in the capsule. The *capsular* is divided again, by Beer, into the *anterior*, *posterior*, and *complete capsular cataract*. When the capsule is rendered opaque, in consequence of an injury, which cuts or ruptures any part of it, it thickens, becomes leathery, and has been called *Cataracta arida siliquosa*. *Cataracta Morgagniana lactea* seu *puriformis* is the *milk* variety, in which the crystalline is transformed into a liquid similar to milk, (F.) *Cataracta lacteæ*; or, as generally defined, in which there is opacity of the fluid situated between the lens and its capsule. The *capsulolenticular* affects both lens and capsule, (*AchrySTALLADIAPHANIE*, *ADIAPHANIE PÉRIHYSTALLIQUE* of Piorry,) and Beer conceives the liquor Morgagni, in an altered state, may contribute to it. Cataracts are also called *hard* (*Sclerocataracta*), *soft* (*Phacomalacia*), *stony* (F. *piérreuse*), *milk* or *cheesy* (*lacteæ* or *caseuse*), *Glaucocataracta*, *Cataracta lacteolor*, according to their density:—white, pearly, yellow, brown, gray,

green, black, (F.) *blanche*, *perlée*, *jaune*, *brune*, *grise*, *verte*, *noire*, according to their colour:—*fixed* or *vacillating*—*cataracta capsulolenticularis fixa* vel *tremula*, (F.) *fixe* ou *branlante*, according as they are fixed or movable behind the pupil. They are likewise called *Cataractæ marmoræ*, *fenestratæ*, *stellatæ*, *punctatæ*, *dimidiatæ*, &c., according to the appearances they present.

They may be *simple* or *complicated* with adhesion, amaurosis, specks, &c.: and *primary* or *primitive*, when *opaque* before the operation:—*secondary*, when the opacity is the result of the operation.

Cataract is commonly a disease of elderly individuals, although, not unfrequently, *congenital*. It forms slowly; objects are at first seen as through a mist; light bodies appear to fly before the eyes, and it may not be until after months or years that the sight is wholly lost. No means will obviate the evil except an operation, which consists in removing the obstacle to the passage of the light to the retina. Four chief methods are employed for this purpose. 1. *Couching* or *Depression*, *Hyalonix'is*, *Hyalonyx'is*, *Crystallocaptation'is*, *Crystallocatactesis*, (F.) *Abaissement ou Déplacement de la Cataracte*. This consists in passing a cataract needle through the sclerotic and subjacent membranes, a little above the transverse diameter of the eye; and at about two lines' distance from the circumference of the transparent cornea, until the point arrives in the posterior chamber of the eye. With this the crystalline is depressed to the outer and lower part of the globe of the eye, where it is left. 2. *By absorption*,—by the French termed *broiement* or *bruising*. This is performed in the same manner as the former; except that, instead of turning the crystalline from the axis of the visual rays, it is divided by the cutting edge of the needle, and its fragments are scattered in the humours of the eye, where they are absorbed. 3. *By extraction*, which consists in opening, with a particular kind of knife, the transparent cornea and the anterior portion of the capsule of the crystalline; and causing the lens to issue through the aperture. Each of the processes has its advantages and disadvantages, and all are used by surgeons. 4. Some, again, pass a cataract needle through the transparent cornea and pupil to the crystalline, and depress or cause its absorption. This is called *Ceratonyxis*. See, also, *Reclination*.

CATARACT, BLACK, Amaurosis—*c. Capsular*, see *Cataract*—*c. Capsulolenticular*, see *Cataract*—*c. Central*, *Centradiaphanes*—*c. Cheesy*, see *Cataract*—*c. Complicated*, see *Cataract*—*c. Congenital*, see *Cataract*—*c. Fixed*, see *Cataract*—*c. Genuine*, see *Cataract*—*c. Hard*, see *Cataract*—*c. Lenticular*, see *Cataract*—*c. Membranous*, see *Cataract*—*c. Milky*, see *Cataract*—*c. Opaque*, see *Cataract*—*c. Primary*, see *Cataract*—*c. Primitive*, see *Cataract*—*c. Secondary*, see *Cataract*—*c. Simple*, see *Cataract*—*c. Soft*, see *Cataract*—*c. Spurious*, see *Cataract*—*c. Stony*, see *Cataract*—*c. Vacillating*, see *Cataract*.

CATARACTA, *Cataract*—*c. Arida siliquosa*, see *Cataract*—*c. Capsulo-lenticularis*, see *Cataract*—*c. Centralis*, *Centradiaphanes*—*c. Dimidiata*, see *Cataract*—*c. Fenestrata*, see *Cataract*—*c. Glauca*, *Glaucoma*—*c. Laticolor*, see *Cataract*—*c. Liquida*, *Hygrocataacta*—*c. Marmoracea*, see *Cataract*—*c. Morgagniana*, see *Cataract*—*c. Nigra*, *Amaurosis*—*c. Punctata*, see *Cataract*—*c. Stellata*, see *Cataract*.

CATARACTE, *Cataract*—*c. Abaissement de la*, see *Cataract*—*c. Blanche*, see *Cataract*—*c. Branlante*, see *Cataract*—*c. Brune*, see *Cataract*—*c. Caséuse*, see *Cataract*—*c. Déplacement de la*, see *Cataract*—*c. Fixe*, see *Cataract*—*c. Grise*, see *Cataract*—*c. Jaune*, see *Cataract*—*c. Laitée*,

res Cataract—c. *Noire*, Amaurosis, see Cataract—*c. Perle*, see Cataract—*c. Pierreuse*, see Cataract—*c. Verte*, see Cataract.

CATARACTE, Cataractus.

CATARACTOUS, *Cataracté*, (F.) *Cataractus*, *Cataractâ vitidus*. One affected with cataract. The French use this term, both for the eye affected with cataract and the patient himself.

CATARIA, (from *catus*, 'a cat,' because cats are fond of it.) See *Nepeta*—*c. Vulgaris*, *Nepeta*.

CATARRH', *Catar'rhus*, *Catar'rhous*, *Catar'rhœma*, *Rheu'ma*, *Deflux'io*, *Catastagnus*, *Phlegmatorrhag'ia*, *Phlegmatorrhœa*, (*καταρρῶς*, from *cata*, and *ρῶ*, 'I flow,') (F.) *Catarrhe*. A discharge of fluid from a mucous membrane. The ancients considered catarrh as a simple flux, and not as an inflammation. Generally it partakes of this character, however. *Catarrh* is, with us, usually restricted to inflammation of the mucous membrane of the air-passages: the French extend it to that of all mucous membranes; (F.) *Flux muqueux*, *Fluxion catarrhale*.

Catarrh, in the English sense, *Bronchocatar'rhus*, *Pul'monary Catarrh'*, *Lung fever* (vulgarly), *Rheu'ma Pectoris*, *Destillatio Pectoris*, *Catar'rhus Pectoris*, *C. Pul'monum*, *C. Pulmonal'is*, *C. Bronchial'is*, *Blennoptysis*, *Tussis catarrhal'is simplex*, *Gravello* (of many), *Feb'ris Catarrhal'is*, *Blennothorax*, *Bronch'itis*, *Catar'rhus à Frigorè*, a Cold, (F.) *Catarrhe pulmonaire*, *Fievre Catarrhale*, *Rhume de Poitrine*, is a superficial inflammation of the mucous follicles of the trachea and bronchi. It is commonly an affection of but little consequence, but apt to relapse and become chronic. It is characterized by cough, thirst, lassitude, fever, watery eyes, with increased secretion of mucus from the air-passages. The antiphlogistic regimen and time usually remove it. Sometimes, the inflammation of the bronchial tubes is so great as to prove fatal.

CATARRH, ACUTE, OF THE UTERUS, see *Metritis*—*c. Chronic*, *Bronchitis*, (chronic)—*c. Dry*, see *Bronchitis*.

CATARRH, EPIDEMIC, *Catar'rhus epidem'icus*, *C. à contagio*, *Rheu'ma epidem'icum*. Catarrh prevailing owing to some particular *Constitutio aëris* [?], and affecting a whole country,—*Influenza*.

CATARRH, PULMONARY, *Bronchitis*, *Catarrh*—*a. Rose*, *Fever*, *hay*—*c. Suffocating nervous*, *Asthma Thymicum*—*c. Summer*, *Fever*, *hay*.

CATARRHACTA, Cataract.

CATARRHAL, *Catarrhal'is*, *Catarrho'icus*, *Catarrhoit'icus*, *Catarrhoët'icus*. Relating to catarrh,—as *Catarrhal Fever*.

CATARRHETIC, *Catarrh*—*c. Aiguë de l'Utérus*, see *Metritis*—*c. Buccal*, *Aphthæ*—*c. Convulsive*, *Bronchitis*—*c. Gastricæ*, *Gastritis*—*c. Gutural*, *Cynanche tonsillaris*—*c. Intestinal*, *Diarrhœa*—*c. Laryngien*, *Laryngitis*—*c. Nasal*, *Coryza*—*c. Oculaire*, *Ophthalmia*—*c. de l'Oreille*, *Otitrhœa*—*c. Pharyngien*, *Cynanche parotidea*—*c. Pituitæa*, *Bronchorrhœa*—*c. Pulmonaire*, *Catarrh*—*c. See*; see *Bronchitis*—*c. Stomacal*, *Gastrorrhœa*—*c. Uterin*, *Leucorrhœa*—*c. Vesical*, *Cystirrhœa*.

CATARRHETICA, (*καταρρηκτικά*, from *καταρρηγνμι*, 'I break down,') Remedies considered proper for evacuating;—as diuretics, cathartics, &c.

CATARRHEUMA, gen. *Catarrheumat'is*, (*cata*, and *ρῆμα*, 'defluxion,') *Catarrh*.

CATARRHEUX, (F.) *Catarrh'ous*. One subject to catarrh; affected with catarrh.

CATARRHEX'IA, *Catarrhet'is*; (same etymon as *Catarrhæctica*.) The action of *Catarrhæctica*. Also, effusion; evacuation of the bowels.

CATARRHESIS, (*καταρρησις*), *Catarrhæsis*, *Excrement*—*c. Vera*, *Hæmatochezia*.

CATARRHŒA, (*cata*, and *ρῶ*, 'I flow,') *Rhematism*.

CATARRHOÏTICUS, (*καταρρητικός*.) An epithet for disease produced by a discharge of phlegm; catarrhal.

CATARRHOPIA PHY'MATA, (*καταρρῶς, καταρρῶς*, 'sloping downwards,') Tubercles tending downwards, or with their apices downwards.

CATARRHOPHE, (*cata*, and *ρῶ*, 'I sip up,') Absorption.

CATARRHOPHESIS, Absorption.

CATARRHOPIA, *Catar'rhysis*, (*καταρρῶς*, from *cata*, and *ρῶ*, 'inclination,') Afflux of fluids towards the inferior parts, and especially towards the viscera of the abdomen. The Greek word *αναρρῶς* expresses an opposite phenomenon, or a tendency towards the upper parts.

CATARRHOS'CHESIS, (*καταρρῶς*, 'catarrh,' and *σχέσις*, 'suppression,') The suppression of a mucous discharge.

CATARRHOUS, Catarrh.

CATARRHUS, Catarrh, Defluxion, Tussis—*c. Æstivus*, *Fever*, *hay*—*c. Bellinsulanus*, *Cynanche parotidea*—*c. Bronchialis*, *Catarrh*—*c. Bronchiorum*, *Bronchitis*—*c. à Contagio*, *Influenza*—*c. Epidemicus*, *Influenza*, *Catarrh*, *epidemic*—*c. à Frigore*, see *Catarrh*—*c. Genitalium*, *Leucorrhœa*—*c. Gonorrhœa*, *Gonorrhœa*—*c. Intestinalis*, *Diarrhœa*—*c. Laryngeus*, *Laryngo-catarrhus*—*c. ad Nares*, *Coryza*—*c. Nasalis*, *Coryza*—*c. Pectoris*, see *Catarrh*—*c. Pulmonalis*, *Catarrh*—*c. Pulmonum*, *Bronchitis*, *Catarrh*—*c. Senilis*, *Bronchitis*, (chronic)—*c. Suffocativus* *Barbadensis*, *C. trachealis*—*c. Trachealis*, *Laryngo-catarrhus*—*c. Urethræ*, *Gonorrhœa pura*—*c. Urethralis*, *Gonorrhœa*—*c. Vesicæ*, *Cystirrhœa*.

CATARRHYSIS, (*cata*, and *ρῶ*, 'I flow,') *Catarrhopia*, *Defluxion*.

CATARTISIS, (*καταρτησις*), *Catartismus*.

CATARTIS'MUS, *Catar'tisis*, (*καταρτισμος*, from *καταρτιζειν*, 'to repair, replace,') The coaptation of a luxated or fractured bone, or hernia.

CATASARCA, (*cata*, and *σαρξ*, gen. *σαρκος*, 'flesh,') *Anasarca*.

CATASCEUE, (*κατασκευη*, from *cata*, and *σκευη*, 'equipment,') *Structure*.

CATASCHASMUS, (*κατασχασμος*, from *cata*, and *σχαζειν*, 'to scarify,') *Bloodletting*, *Scarification*.

CATASTAGMUS, (*κατασταγμος*, from *cata*, and *σταζειν*, 'to drop,') *Catarrh*, *Coryza*.

CATASTALAGMUS, (*cata*, and *σταλαζειν*, 'to drop,') *Coryza*, *Distillation*.

CATASTALTICA, (*κατασταλτικά*, from *cata*, and *στέλλειν*, 'to repress,') *Hæmatostatica*, *Sedatives*, *Styptics*.

CATAS'TASIS, (*καταστασις*, from *καδιστημι*, 'I establish,') The constitution, state, condition, &c., of any thing. Also, the reduction of a bone. See *Constitution*, and *Habit of Body*.

CATAT'ASIS, (*κατατασις*, from *κατατεινω*, 'I extend,') *Extension*. The extension and reduction of a fractured limb.

CATATHLIPSIS, (*cata*, and *θλιβω*, 'I press,') *Oppression*.

CATAWBA TREE, *Catalpa*.

CATAXIS, (*καταξις*), *Fracture*.

CATCH FLY, *Apocynum androsæmifolium*, *Silene Virginica*.

CATCHING, see *Contagious*.

CATCHUP, *Ketchup*.

CATECHU. The extract of the wood of *Aca'cia* seu *Mimo'sa Cat'echu*, *Catt'echu*, and of the leaves of *Nau'clea* seu *Unca'ria gam'bir*, oriental trees. The drug is also called *Extractum Cat'echu*, *Ter'ra Japon'ica*, *Japan Earth*, *Cas'chen*, *Catt'echu*, *Cash'on*, *Cuit'chu*, *Cash'joe*, *Cacoon*, *Cate*, *Kaath*, *Cuti*, *Cutch*, *Caira*, *Suc'cia Japon'ica*, *Gam'bir*, (F.) *Cinchou*. It is a powerful astringent, and is used in diarrhœa, intestinal hemorrhage, &c. Dose, gr. xv to ʒss, in powder.

The British Pharmacopœia (1864) has *Cut'echu nigrum*, *black catechu*: an extract of the heart-wood of *Aca'cia Cut'echu*, and *Cut'echu pallidum*, *pale Cut'echu*: an extract of the leaves and young shoots of *Uncaria Gum'bir*.

CATECHU SQUARE, see *Nauclea ga'nbir*.

CATEADI'ON, (*cata*, and *eia*, 'a blade of grass.') A long instrument, formerly thrust into the nostrils to excite hemorrhage in headache.

CATENÆ MUSCULUS, Tibialis anticus.

CATEONESIS, (*κατανομις*, from *cata*, and *αινωω*, 'I moisten.') Catantlema.

CATERPILLAR, VEGETABLE, *Sphaeria Sinenensis*.

CATGUT, *Galega Virginiana*.

CATH, in composition, see *Cata*.

CATHÆRESIS, (*καθαιρεσις*, 'subtraction, diminution.') Extenuation or exhaustion, owing to forced exercise. The action of catheterics.

CATHERETICUS, (*καθηρειακος*,) Cathartic.

CATHARETICUS, Cathartic.

CATHARISMUS, (*καθαρισμος*, from *καθαρειν*, 'to purge.') Depuration.

CATHARMA, gen. *Catharm'atis*, *Purgamentum*. The matter evacuated by a purgative, or by spontaneous purging; also, a cathartic.

CATHARMUS. A purgation. Also, the cure of a disease by magic, &c.

CATHAR'SIS, (*καθαρσις*, from *καθαρειν*, 'to purge.') *Purgatio*, *Apocathar'sis*, *Coprophor'ia*, *Coprophore'sis*. A natural or artificial purgation of any passage:—mouth, anus, vagina, &c.

CATHARTIC, (*καθαρτικος*,) *Cathart'icus*, *Cathar'eticus*, *Cuthar'ma*, *Coprocril'icum*, *Cuprago'gum*, *Lustra'mentum*, *Pur'gans medicament'um*, *Dejecto'rium Remed'ium*, *Eecathar'ticus*, *Hypac'ticus*, *Hypochore'ticus*, *Al'vum evac'uans*, *Hypel'atus*, *Lapac'ticus*, *Apocathar'ticus*, (F.) *Cathartique*. A medicine which, when taken internally, increases the number of alvine evacuations. Some substances act upon the upper part of the intestinal canal, as *colomet* and *colocyath*; others, on the lower part, as *aloes*; and some on the whole extent, as *saline* purgatives. Hence a choice may be necessary. Cathartics are divided into purgatives and laxatives.

CATHARTIN, see *Cassia Senna*, and *Convolverulus jalapa*.

CATHARTIQUE, Cathartic.

CATHARTOCARPUS, (*καθαρειν*, 'to purge,' and *καρπος*, 'fruit,') *Cassia fistula*.

CATHEDRA, (*καθέδρα*, 'a seat,') *Anus*.

CATHEMERINUS, (*καθημερινος*, from *cath*, and *ημερα*, 'day,') *Quotidian*.

CATHEMERIUS, (*καθημεριος*,) *Quotidian*.

CATHERETIC, (*καθαρειτικος*, from *καθαρειν*, 'I eat, destroy,') *Cuthar'eticus*, *Ectylot'icus*, *Sarcoph'agus*. Substances applied to warts, exuberant granulations, &c., to eat them down. *Mild caustics*.

CATHETER, (*καθετηρ*, from *καθιμι*, (*καθ'*, and *ιμι*, 'to send,') 'to explore,') *Æne'a*, *Alg'alie*, *Alg'olic*, *Cath'et'ria*, *Demi'sor*, *Immis'sor*. A hollow tube, introduced by surgeons into the urinary bladder, for the purpose of drawing off the urine. Catheters are made of silver or elastic gum. See *Bougie*. The French generally use the word *catheter* for the solid *sound* or *staff*; and *alg'olie* or *sonde* for the hollow instrument.

CATHETER, NA'SAL. An instrument, invented by M. Gensoul, of Lyons, for catheterizing the ductus ad nasum. It is hook-shaped; the extremity, bent at a right angle, is about an inch in length, suited to the distance of the lower orifice of the duct from the nostril, and likewise to the length and form of the duct, with a slight spiral turn.

CATHETERIS, Catheter.

CATHETERISIS, (*καθετηρησις*,) *Catheterismus*.

CATHETERIS MUS, (*καθετηρησις*,) *Cathete'ri-*

sis, *Catheteriza'tio*, *Cath'eterism*, *Catheteriza'tion*, *Immis'sio Cathete'ria*. The introduction of a catheter, hollow sound, or tube, into a natural passage, as through the urethra into the bladder into the Eustachian tube, larynx, &c. In the latter cases, it is, at times, called *Tubage*. Also, probing a wound. *Melosis*.

CATHETERIZATION, *Catheterismus*.

CATH'ETERIZE. To perform the operation of catheterism,—in other words, to introduce the catheter: to probe or sound a cavity.

CATHID'YSIS, (*καθιέρωσις*, from *καθιέρωω*, 'I place together,') Reduction of a part to its natural situation.

CATHMIA, *Plumbi oxydum semivitreum*.

CATHMIR, *Calamina*.

CATHOD'IC, *Cathod'icus*; (*καθόδος*, 'a descent,' from *cath*, and *δός*, 'a way,') An epithet applied by Dr. Marshall Hall to a downward course of nervous action.

CATH'OLIC HU'MOURS, (*καθολικος*, 'universal,' from *cath*, and *ολος*, 'the whole,') (F.) *Humeurs Catholiques*, are the fluids spread over the whole body.

CATHOLICON, (*καθολικον*,) *Panacea*.

CATHOL'ICON DU'PLEX. An ancient purging electuary, chiefly composed of cassia, tamarinds, rhubarb, senna, &c.

CATIL'LIA. A weight of nine ounces.

CATILLUS, (dim. of *catinus*, 'a bowl,') *Cup*.

CATINUS FUSORIUS, *Crucible*.

CATLING, *Knife*, double-edged.

CATNEP or CATNIP, *Nepeta Cataria*.

CATO, (*κατω*, 'below, beneath,') This word, in the writings of Hippocrates, is often used for the abdomen, especially the intestines. When he advises a remedy *κατω*, he means a purgative; when *ανω*, 'above or upwards,' an emetic. As a prefix, *Cato* means 'beneath,' as in

CATOCATHAR'TIC, *Catocathar'ticus*, (*cato*, and *καθαρειν*, 'I purge,') A medicine which purges downwards. One that produces alvine evacuations. The antithesis to *Anacathartic*.

CAT'OCHE, (*κατοχη*, from *κατοχω*, 'I take possession of,') *Cat'ochesis*, *Cat'ochus*. This word has, by some, been used synonymously with *Catalepsy*; by others, with *Coma vigil*; by others, with *Tetanus*.

CATOCHE GALENI, *Catalepsy*.

CATOCHEILON, (*cato*, and *χιλος*, 'lip,') *Lip* (lower).

CATOCCHUS, (*καταχος*,) *Catoche*, *Ecstasis* — c. *Cervinus*, *Tetanus* — c. *Holotonicus*, *Tetanus* — c. *Infantum*. Induration of the cellular tissue.

CATOMIS'MUS, (*κατωμισμος*, from *cato*, and *ωμος*, 'shoulder,') *Subhumera'tio*. A mode with the ancients of reducing luxation of the humerus by raising the body by the arm.

CATOPTER, (*cato*, and *πτερω*, 'I see,') *Speculum*.

CATOP'TRIC, (*κατοπτρικος*,) *Catop'tricus*, (F.) *Catop'trique*. Relating or appertaining to *Catop'trics* or the reflection of light.

CATOPTRIC EXAMINA'TION OF THE EYE. When a lighted candle is held before the eye, the pupil of which has been dilated by belladonna, three images of it are seen—two erect, and one inverted:—the former owing to reflection from the cornea and anterior surface of the crystalline; the latter, owing to reflection from the posterior layer of the crystalline, inverted, and between the other two. This mode of examining the eye has been proposed as a means of diagnosis between cataract and amaurosis. In the latter, all the images are seen.

CATOP'TROMANCY, (*κατοπτρον*, 'a mirror,' and *μαντεία*, 'divination,') A kind of divination by means of a mirror.

CATOPTRON, (κατοπτρον, from *cata*, and *πτω*, 'I see,') Speculum.

CATORCHITES, (κατορχιτης.) A kind of sour wine, prepared with the orchis and black grape, or dried figs. It was formerly employed as a diuretic and emmenagogue. Called, also, *Syci'tes*.

CATORETICUS, Purgative.

CATORETICUS, (κατορετικός, from *κατω*, 'below, beneath,') Purgative.

CATO'TICA, (from *κατω*, 'beneath,') Diseases infecting internal surfaces. Pravity of the fluids or emunctories, that open on the internal surfaces of organs. The second order in the class *Eccritica* of Good.

CATOX'YS, (κατοξυς, from *κατα*, 'intensive,' and *οξυς*, 'acute,') *Peracut'us*. Highly acute; as *Morb'us Catoxy's*, *M. Peracut'us*, a very acute disease.

CATS' EYE. A name, under which several morbid conditions of the eye would appear to have been confounded—all agreeing, however, in presenting an opalescent appearance of the pupil or of the bottom of the eye; those parts reflecting the light in various colours, or, at least, with various degrees of intensity, according to the direction in which the eye is turned. This appearance was compared by Beer to the reflection from the tapetum of the eye of the cat. See Amaurotic cat's eye.

CAT'S FOOT, *Antennaria dioica*—c. Purr, *Frémissement Cutané*.

CATSKILL, MIN'ERAL WATERS OF. See New York, mineral waters of.

CATSUP, Ketchup.

CATTAGAUMA, Cambogia.

CATITERUS, (καττιτερος,) Tin.

CATULOTICA, (κατουλωτικα,) *Cicatrissantia*.

CATURUS SPICIFLORUS, (καττα, 'a cat,' and *ουρα*, 'tail,') *Cav'da fel'is*, from the shape of its flowers. *Acal'lypha his'pida*. An East Indian shrub, *Order*, *Euphorbiaceæ*; used, in its native country, in decoction, in diarrhoea and dysentery.

CATU-TRIPALI, Piper longum.

CAUCALIS CAROTA, (καυκαλις,) *Daucus carota*—c. *Sanicula*, *Sanicula*.

CAUCALOIDES, (καυκαλις, 'the *caucalis*,' and *είδος*, 'resemblance,') *Patella*.

CAUCASIAN, see *Homö*.

CAUCHEMAR, Incubus.

CAUCHEVIEILLE, Incubus.

CAUCHUC, Caoutchouc.

CAUDA, ('a tail,') see *Clitorism*, *Coccyx*, *Penis*.

CAUDA EQU'NA. The spinal marrow, at its termination, about the second lumbar vertebra, gives off a considerable number of nerves, which, when unravelled, resemble a horse's tail; hence the name; (F.) *Queue de Cheval*, *Q. de la Moëlle Épinrière*. See *Medulla Spinalis*.

CAUDA FELIS, *Caturus*—c. *Agrestis*, *Acalypha betulina*—c. *Musculi*, see *Muscle*—c. *Pudendi*, *Cercosis*—c. *Salax*, *Penis*.

CAUDAL, *Caud'ate*, *Cauda'lis*, *Cauda'tus*, (from *cauda*, 'a tail,') Relating or appertaining to a tail. Having a tail or tail-like appendage:—as 'caudal or caudate cells or corpuscles, having a tail-like appendage, as in cancerous growths.

CAUDATE, Caudal.

CAUDATIO, (from *cauda*, 'a tail,') *Clitorism*.

CAUDATUS, Caudal.

CADEX CEREBRI, (*caudex*, 'a stem,') *Peduncles* of the brain.

CAUDIEZ, MIN'ERAL WATERS OF. *Caudiez* is a small town, nine leagues from Perpignan, in France, where there is a thermal spring, containing a little sulphate of soda and iron.

CAUDISONA DURISSA, (from *cauda*, 'the tail,' and *sono*, 'to sound,') *Crotalus horridus*.

CAUDLE, *Cow'del*, (F.) *Chaudéan*, (*chaud*, 'warm or hot,') A nourishing gruel given to women during the childbed state. The following

is a form for it: Into a pint of fine gruel, not thick, put, whilst it is boiling hot, the yolk of an egg beaten with sugar, and mixed with a large spoonful of cold water, a glass of wine, and nutmeg. Mix the whole well together. Brandy is sometimes substituted for the wine, and lemon peel or capillaire added. It is also at times made of gruel and beer, with sugar and nutmeg.

CAUL, (*caula*, 'a sheep-fold,') *Pile'us*, *Pile'olus*, *Gal'ea*, *Vitta*, *Lucky hood*, (F.) *Coeffe*, *Coiffe*—(*Être né coiffé*—'to be born with a caul,') When a child is born with the membranes over the face, it is said to be 'born with a caul.' In the catalogue of superstitions, this is one of the favourable omens. The caul itself is supposed to confer privileges upon the possessor; hence the membranes are dried, and sometimes sold for a high price, and even bequeathed. See *Epiploon*.

CAULE'DON, (*καυληδον*, from *καυλος*, 'a stalk,') *Cicye'don*. A transverse fracture.

CAUL'FLOWER, (*caulis*, (G.) *Kohl*, 'cabbage,' and *flower* [?]), *Brassica Florida*.

CAULIFLOWER EXCRES'CENCE, (F.) *Choufleur*. An excrescence, which appears about the origin of the mucous membranes, chiefly about the anus and vulva, and which resembles, in appearance, the head of the cauliflower. It is often syphilitic in its character.

CAULIFLOWER EXCRES'CENCE OF THE U'TERUS, *Metrob'ol'rytes*. A morbid growth from a part, or the whole, of the circumference of the os uteri; and, at times, from the surface of the cavity of the uterus. It is doubtful whether its progress can be arrested except by excision.

CAULIS, (*καυλος*, 'a stalk,') *Penis*—c. *Florida*, *Brassica Florida*—c. *Uteri*, *Collum uteri*.

CAULOPHYL'UM THALICTROI'DES, (*καυλος*, 'a stalk or stem,' and *θαλλων*, 'a leaf,' because the leaf-stalks appear like a continuation of the stem.) *Leon'tice thalictroi'des*, *Blueberry Cohosh*, *Cohosh*, *Cohush*, *Blue Cohosh*, *Blueberry*, *Parpoose Root*, *Squaw Root*, *Blue Gin'seng*, *Yell'low Gin'seng*, a plant of the *Family Berberidæ*; *Scz. Syst.* *Hexandria Monogynia*, which grows all over the United States, flowering in May and June. The infusion of the root is much used by the Indians in various diseases. To it are ascribed emmenagogue and diaphoretic virtues.

CAULOPL'E'GIA, (*caulus*, and *πληγη*, 'a wound,' or 'stroke,') An injury or paralysis of the male organ.

CAULORRHAGIA, (*caulus*, and *ῥήγνυμι*, 'to break forth,') *Stomatosis*—c. *Ejaculatoria*, *Spermatoecystidorrhagia*—c. *Stillatitis*, *Urethrorrhagia*.

CAULORRHŒA BENIGNA, (*caulus*, and *ῥέω*, 'I flow,') *Gonorrhœa pura*.

CAULUS, (*καυλος*,) *Penis*.

CAUMA, gen. *caum'atis*, (*καυμα*, 'a burnt part,' from *καω*, 'I burn,') Great heat of the body or atmosphere. *Synocha*, *Empresma*.

CAUMA BRONCHITIS, *Cynanche trachealis*—c. *Carditis*, *Carditis*—c. *Enteritis*, *Enteritis*—c. *Gastritis*, *Gastritis*—c. *Hæmorrhagicum*, *Hæmorrhagia activa*—c. *Hepatitis*, *Hepatitis*—c. *Ophthalmitis*, *Ophthalmia*—c. *Peritonitis*, *Peritonitis*—c. *Phrenitis*, *Phrenitis*—c. *Pleuritis*, *Pleuritis*—c. *Podagrium*, *Gout*—c. *Rheumatismus*, *Rheumatism*, *acute*.

CAUMATO'DES, *καυματώδης*, *Caumate'rus*, *καυματ'ρος*, (from *cauma*, and *οδον*,) Burning hot. *Feb'ris caumato'des*, *F. causo'des*. Inflammatory fever. *Synocha*.

CAUNGA, *Arcea*.

CAUSA, *Cause*—c. *Conjuncta*, *Cause*, *proximate*—c. *Continens*, *Cause*, *proximate*.

CAUSÆ (pl. of *causa*) *ABDITÆ*, *Causes*, *pre-disponent* or *remote*—c. *Actuales*, *Causes*, *occa-*

sional—c. *Præcipientes*. Causes, procatartetic—c. *Prægumene*, Causes, predisponent.

CAUSE, *Com'mun, A'ct'ua, A'ct'ion*. An act which precedes another, and seems to be a necessary condition for the concurrence of the latter. The causes of disease are generally extremely obscure; although they, sometimes, are evident enough. The *predisponent* and *occasional* causes are the only two on which any stress can be laid; but as authors have divided them differently, a short explanation of each is necessary.

CAUSE, *Ac'cessory*, (F.) *Cause Accessoire*. One which has only a secondary influence in the production of disease.

CAUSES, *ACCIDENT'AL*, *Com'mon Causes*, (F.) *Causes Accidentelles*, are those which act only in certain given conditions; and which do not always produce the same disease. Cold, e.g., may be the accidental cause of pneumonia, rheumatism, &c.

CAUSES *CACHÉES*, C. occult—c. Common, C. Accidental—c. *Déterminantes*, C. Specific—c. *Éloignées*, C. Predisponent—c. Exciting, C. Occasional—c. Essential, C. Specific.

CAUSES, *EXTERNAL*, (F.) *Causes externes*, are such as act externally to the individual; as air, cold, &c.

CAUSES *FORMELLES* (F.) are such as determine the form or kind of disease. They differ from the *Causes Matérielles*, which are common to a set of diseases; as, to the neuroses, phlegmasia, &c.

CAUSES, *HIDDEN*, C. Occult—c. Immediate, C. Essential.

CAUSES, *INTERNAL*, (F.) *Causes Internes*, are those which arise within the body; as mental emotions, &c.

CAUSES, *MECHANICAL*, (F.) *Causes mécaniques*, are those which act mechanically, as pressure upon the windpipe in inducing suffocation.

CAUSES, *NEGATIVE*, (F.) *Causes négatives*, comprise all those things, the privation of which may derange the functions;—as abstinence too long continued. They are opposed to *positive causes*, which, of themselves, directly induce disease;—as the use of indigestible food, spirituous drinks, &c.

CAUSES, *OBSCURE*, C. Occult.

CAUSES, *OC'CASIONAL*, *Exciting Causes*, *Causes actuelles*, (F.) *Causes occasionnelles*, are those which immediately produce disease.

CAUSES, *OCULT*, *Hidden causes*, *Obscure causes*, (F.) *Causes occultes ou cachées ou obscures*. Any causes with which we are unacquainted; also, certain inappreciable characters of the atmosphere, which give rise to epidemics.

CAUSES, *PHYSICAL*, (F.) *Causes physiques*,—those which act by virtue of their physical properties; as form, hardness, &c. All vulnerating bodies belong to this class.

CAUSES, *PHYSIOLOGICAL*, (F.) *Causes physiologiques*, those which act only on living matter;—narcotics, for example.

CAUSES, *PREDISPO'XENT*, C. *predispos'ing*, *Remote causes*, *Causes prægumene*, *Causes ab'dites*, *Causes remotæ*; (F.) *Causes prédisposantes*, *Causes éloignées*,—those which render the body liable to disease. They may be general, affecting a number of people, or particular, affecting only one person.

CAUSES, *PRIN'ICIPAL*, (F.) *Causes principales*,—those which exert the chief influence in the production of disease, as distinguished from the *accessory causes*.

CAUSES, *PROCATARTIC*, *Causes procatarteticæ*, *Causes præcipientes*, (*προκαταρτικα*, from *pro*, 'before,' and *καταρχω*, 'I begin,' 'the origin or beginning of a thing,') These words have been used with different significations. Some have employed them synonymously with *predisponent* or

remote causes; others with *occasional* or *exciting causes*.

CAUSE *PROCHAINE*, C. proximate.

CAUSE, *PROXIMATE*, *Causes proximæ seu continens seu conjunctæ*, (F.) *Cause continente ou prochaine*, may be the disease itself. Superabundance of blood, e.g., is the proximate cause of plethora.

CAUSES, *Remote*, C. predisponent.

CAUSES, *SPECIFIC*, *Essential* or *Immediate causes*, &c., (F.) *Causes spécifiques*, C. *essentielles*, C. *déterminantes*;—those which produce a determinate disease.

CAUSIS, (*καυσις*, from *καω*, 'I burn,') Burn, Ebullition, Fermentation, Incendium, Ustion.

CAUSODES, (*causis*, and *odes*), Inflammatory.

CAUSOMA, (*καυσωμα*, from *causis* and *oma*), Inflammation.

CAUSTIC, *Causticus*, *Cantereticus*, *Diæreticus*, *Erodens*, *Adulens*, *Urens*, *Pyroticus*, (*καυστικός*, 'that which burns,' from *καω*, 'to burn,') (F.) *Caustique*. Bodies which have the property of burning or disorganizing animal substances. The word is also used substantively. The most active are called *Escharotica*. Caustics are also termed 'corrosives.'

CAUSTIC BEARER, *Porte-pierre*.

CAUSTIC, *COMMON*, *Causticum commune*—c. Lunar, Argenti nitras.

CAUSTIC PASTE, Canquoin's, Pasta Zinci chloridi.

CAUSTICA ADUSTIO, ('caustic burning,') Cauterization.

CAUSTICITY, *Caustic'itas*. The impression which caustic bodies make on the organ of taste; or more commonly, the property which distinguishes those bodies.

CAUSTICOPHORUM, (*causticum*, and *φερο*, 'I carry,') *Porte-pierre*.

CAUSTICUM, *Caustic*.

CAUSTICUM ÆTHIOPICUM, *Caustic*, *Unguentum melan'icum causticum*. A sort of paste, made by rubbing powdered saffron with concentrated sulphuric acid, recommended by Velpeau as a caustic in cases of gangrenous and carcinomatous ulcers. The acid is the caustic: the saffron, the constituent merely.

CAUSTICUM ALKALINUM, *Potassa fusa*—c. Americanum, *Veratrum sabadilla*—c. Antimoniale, *Antimonium muriatum*.

CAUSTICUM COMMUNE, *Potent'ial Caut'ery*, *Common Caustic*, *Caust'rium potent'iale*, *Lapis sepi'ticus*, *Causticum commu'nè mit'ius*. This consists of quicklime and black soap, of each equal parts. Also, *Potassa fusa*.

CAUSTICUM COMMUNE ACERRIMUM, *Potassa fusa*—c. *Commune fortius*, *Potassa cum calce*—c. Lunar, Argenti nitras—c. *Potentiale*, *Potassa fusa*—c. Salinum, *Potassa fusa*—c. *Viennense fusum* Filhos, see Powder, Vienna.

CAUSTIQUE, *Caustic*—c. *Ammoniacale*, *Pommade de Gondret*—c. Filhos, see Powder, Vienna—c. *de Vienne*, Powder, Vienna.

CAUSUS, (*καυος*, 'heat,') A highly ardent fever; *Deul'rens*. Pinel regards it as a complication of bilious and inflammatory fever; Broussais, as an intense gastritis, accompanied with bilious symptoms. See Synocha.

CAUSUS, ENDEMIAL, OF THE WEST INDIES, *Fever*, *Yellow*—c. *Tropicus endemicus*, *Fever*, *Yellow*.

CAUTER, (*καυρη*, 'a burner,') *Cauterium*.

CAUTÈRE, *Cauterium*, *Fonticulus*—c. *Inherent*, *Inherent cautery*.

CAUTERETICUS, (from *cauter*,) *Caustic*.

CAUTERETS, MINERAL WATERS OF. Cauterets is a bourg seven leagues from Barèges (*Hautes-Pyrénées*), France. The waters are hydrosulphurous and thermal—temperature 123° Fah. They are used in the same cases as the Barèges water, and are visited from June to October.

CAUTERIASMUS, Cauterization.

CAUTERIUM, *C. actuale*, *Cauter*, *Imustorium*, *Kuptorium*, *Ignis actualis*, *Cautery*, (*καυτήριον*, from *καω*, 'I burn.') (F.) *Cautère Feu actuel*. A substance, used for 'firing,' burning, or disorganizing the parts to which it is applied. Cauteries were divided by the ancients into *actual* and *potential*. The word is now restricted to the red-hot iron; or to positive burning. It was, formerly, much used for preventing hemorrhage from divided arteries; and also with the same views as a blister. The term *Potential Cautery*, *Cauterium potentiale*, *Ignis potentialis*, (F.) *Feu potentiel*, was generally applied to the *causticum commune*, but it is now used synonymously with *caustic* in general. *Cautère* also means an issue.

CAUTERIUM ACTUALE, Cauterium—c. *Potentiale*, Cauterium.

CAUTERIZATION, *Cauterisatio*, *Cauteriasionus*, (from *καυρίζω*, 'I cauterize,') *Excusio*, *Imustio*, *Caustica Adustio*. Firing. The effect of a cautery or caustic. The French, amongst whom cauterization is much used, distinguish five kinds: 1. *Cautérisation Inhérente*, which consists in applying the actual cautery freely, and with a certain degree of force, so as to disorganize deeply. 2. *Cautérisation transcurante*, which consists in passing the edge of the *Cautère cutellaire*, or the point of the *Cautère conique*, lightly, so as not to disorganize deeply. 3. *Cautérisation par points*, which consists in applying on the skin, here and there, the hot point of the conical cautery, with sufficient force to cauterize the whole thickness of the skin. 4. *Cautérisation lente*, *slow cauterization*, by means of the *moxa*. 5. *Cautérisation objective*, which consists in holding the cautery at some distance from the part to be acted upon by it.

CAUTERIZE; *Caustico adu'rere*; (F.) *Cautériser*. To apply a cautery or caustic. To burn with a cautery or caustic.

CAUTERY, Cauterium—c. *Actual*, Cauterium—c. *Galvanic*, see *Moxa*—c. *Potential*, *Causticum commune*, Cauterium.

CAVA, ('hollow,') *Vulva*.

CAVA VE'NA. The hollow or deep-seated vein. (F.) *Veine cave*. A name given to the two great veins of the body, which meet at the right auricle of the heart. The *vena cava superior*, *thoracica seu descendens*, is formed by the union of the subclavians; and receives successively, before its termination at the upper part of the right auricle, the *inferior thyroid*, *right internal mammary*, *superior diaphragmatic*, *azygos*, &c. The *vena cava inferior*, *abdominalis seu ascendens*, arises from the union of the two *primary iliacs*, opposite the fourth or fifth lumbar vertebra, receives the *middle sacral*, *lumbar*, *right spermatic*, *hepatic*, and *inferior diaphragmatic*, and opens at the posterior and inferior part of the right auricle.

CAVAL, *Cavalis*. Appertaining or relating to the *vena cava*.

CAVATIO, (*cavus*, 'hollow,') *Cavity*.

CAVEA, *Cavity*—c. *Narium*, *Nares*.

CAVER'NA, *Antrum*, ('a cavern,' from *caveus*, 'hollow,') This term has been used for the female organs of generation. See *Cavity*, and *Vulva*.

CAVERNA NARIUM, *Nares*.

CAVERNÆ (pl. of *caverna*) *Dentium*, *Alveoli dentium*—c. *Frontis*, *Frontal Sinuses*.

CAVERNEUX, *Cavernous*.

CAVERNOUS, *Cavernosus*, (F.) *Caverneux*. Filled with small cavities or caverns,—as a sponge.

CAVERNOUS BOD'IES, *Corpora Cavernosa* of the penis, *Corpora Nervosa* seu *Nervosopungiosa Penis*, (F.) *Corps Caverneux*. The corpus cavernosum is a kind of cylindrical sac, composed of cells; separated, through its whole extent, by a

vertical, incomplete septum, *Sep'tum pectiniforme*, *Trabecula corporum cavernosorum*, and forming nearly two-thirds of the penis. The corpus cavernosum, on each side, arises from the ascending portion of the ischium, and terminates obtusely behind the glans. The arteries of the corpora cavernosa come from the internal pudic. See *Helicine Arteries*. Nerves are found on the surface of the outer membrane, but they do not appear to penetrate the substance, and the smooth muscular fibre has been traced into the fibrous parietes of the cells, as in the case of all erectile tissues.

J. Müller's researches have led him to infer, that both in man and the horse, the nerves of the corpora cavernosa are made up of branches proceeding from the organic as well as the animal system, whilst the nerves of animal life alone provide the nerves of sensation of the penis.

CAVERNOUS BODIES (*Corpora Cavernosa*) of the Clitoris, are two hollow crura, forming the clitoris.

CAVERNOUS BOD'Y OF THE VAGI'NA, *Corpus Cavernosum Vaginae*, *Plexus retiformis*, *Sinus of Santorini*, is a substance composed of blood-vessels and cells, similar to those of the penis and clitoris, which covers the outer extremity of the vagina, on each side. It serves to contract the entrance to the vagina during coition. See *Bulbus vestibuli*.

CAVERNOUS GANGLION, see *Carotid* or *Carotic Nerve*.

CAVERNOUS RESPIRA'TION. When a cavity exists in the lungs, and one or more ramifications of the bronchia terminate in it, a loud tubal noise is emitted, provided the cavity be not filled with fluid, which is called *cavernous respiration*. In this condition, the cough likewise is *cavernous*, (F.) *Toux Caverneuse*. When the capacity of the cavern is very great, the sound of the respiration is like that produced by blowing into a decanter, with the mouth at a little distance from the neck. This kind of cavernous respiration has been called *amphoric*, from *amphora*, 'a flask;' (F.) *Respiration amphorique*, *Souffle amphorique*, *S. métallique*.

The *Veiled Puff*, (F.) *Souffle voilé*, is a modification of the cavernous respiration, in which, according to Laënnec, "a sort of movable veil interposed between the excavation and the ear" seems to be agitated to and fro. It is a sign which is not much attended to.

CAVERNOUS RHONCHUS, see *Cavernous Respiration*, and *Gurgling*.

CAVERNOUS SI'NUS, *Si'nus Cavernosus* seu *polymorphus* seu *sphenoidalis*, *Receptaculum sellæ equinae lateribus appositum*, (F.) *Sinus caverneux*. The *Cavernous Sinuses* are venous cavities of the dura matter, filled with a multitude of reddish, soft filaments, intersecting each other; and, as it were, reticulated. They commence behind the inner part of the sphenoid fissure, pass backwards on the sides of the *fossa pituitaria*, and terminate by opening into a cavity, common to the superior and inferior petrosal sinuses. They receive some meningeal veins, the ophthalmic veins, &c. The anterior extremity of each cavernous sinus has been named the *ophthalmic sinus*.

CAVERNOUS TEX'TURE or TIS'SUE, *Tella cavernosa*, (F.) *Tissu Caverneux*. The spongy substance which forms the greater part of the penis and clitoris. It seems to consist of a very complicated lace-work of arteries and veins; and, probably, of nervous filaments, with small fibrous plates, which form by their decussation numerous cells communicating with each other. This spongy texture produces erection, by dilating and swelling on the influx of blood; and probably, also, by virtue of some property inherent in it.

CAVERNOUS WHISPER. A blowing sound, heard over cavities when words are whispered.

CAVERNULOUS (dim. of *cavernous*) RHONCHUS, see Gurgling.

CAVIALE, Caviare.

CAVIARE', *Caviar'*, *Caviale'*, *Kaviar'*. A culinary preparation, much prized by certain peoples, and made, on the shores of the Black and Caspian Seas, chiefly from the roe of the sturgeon, mixed with salt and other condiments.

CAVICULA, *Cavilla*, (from *cavus*, 'hollow.') The ankle or space between the malleoli. Some have given this name to the os cuneiforme. See Astragalus.

CAVICULE PEDIS NODUS, Tarsus.

CAVILLA, Astragalus, Cavicula.

CAVITAS, Cavity, (*cavus*, 'hollow,') — c. Antro-a Auris, Tympanum — c. Buccinata, Cochlea — c. Cochleata, Cochlea — c. Digitata ventriculi lateralis, Cornu posterius ventriculi lateralis.

CAVITAS ELLIPTICA, *Ampulla*, *Sinus ampullaceus*. A dilatation at one end of the semicircular canals of the ear.

CAVITAS HUMERI GLENOIDES, see Glenoid — c. Narium, Nares — c. Oculi, Orbit — c. Oris, Mouth — c. Pulpe, see Tooth.

CAVITATES (pl. of *cavitas*) CEREBRI, Ventricles of the brain — c. Dura matris, Sinuses of the dura mater — c. Innominatae, Auricles of the heart — c. Interseapulares, see Interseapularis.

CAVITÉ, Cavity — c. Dentaire, Dental cavity — c. des Épiploons, see Peritonæum — c. du Tympan, Tympanum.

CAVITY, *Car'itas*, *Car'um*, *Cac'lotis*, *Car'lon*, *Car'lea*, *Caver'na*, *Cava'tio*, (F.) *Cavité*. The hollow of anything, as of the cranium, mouth, nasal fossæ, &c.

CAVITIES, SPANCH'NIC, (F.) *Cavités splanchniques*, are those which contain the viscera. They are three in number:—the cranium, chest, and abdomen. The cavities of bones, connected with joints or otherwise, are described under their particular denominations.

CAVUM, Cavity — c. Abdominis, see Abdomen. CAVUM CRA'NI. Ven'ter Supre'mus. The cavity formed by the proper bones of the cranium.

CAVUM DENTIS, see Tooth — c. Mediastini, see Mediastinum — c. Narium, Nares, Nasal fossæ — c. Oris, Mouth — c. Pericardii, see Pericardium — c. Thoracis, see Thorax — c. Tympani, Tympanum.

CAYAN, Phaseolus Creticus.

CAZABI, Jatropha manihot.

CEANOTHUS, (*κεανωθος*,) *Cirsium arvense* — c. Americanus, *Celastrus* — c. Trinervis, *Celastrus*, CEAR, (*κααρ*) Heart.

CEASMA, gen. *Ceas'matis*, (*κεασμα*,) Fissure.

CEBI GALLINÆ. The liver of the fowl, bruised.

CEBIPIRA. A large Brazilian tree, whose bitter and astringent bark is used in making anti-rheumatic baths and fomentations.

CEBOCÉPHALE, *Ceboceph'alus*, (*κεβος*, 'an ape,' and *κεφαλη*, 'a head'.) A genus of monsters, in which the nose does not project, and the interocular region is narrow and plane, so as to give the physiognomy a striking resemblance to that of the American ape.

CECIS, gen. *Ce'cidis*, (*κηκis*, gen. *κηκίος*,) see *Quercus infectoria*.

CECITÉ, *Cæcitas* — c. Nocturne, Hemeralopia.

CECITY, Cæcitas.

CEDAR, RED, *Juniperus Virginiana* — c. White, *Cupressus thyoides*.

CEDEIA, (*κηδεα*, from *κηδεω*, 'to attend to a dead body,') Embalming.

CEDMA, gen. *Ced'matis*, (*κεδμα*,) Aneurism, Varix.

CED'MATA, (*κεδματα*, from *κεδαν*, 'to disperse,') Rheumatic pains of the joints, espe-

cially of the hips, groin, or genital organs. A form of gout or rheumatism.

CÉDRAT, *Citrus medica*.

CÉDRATIER, see *Citrus Medica*.

CEDRELA FEBRIFUGA, (from *κεδρος*, 'the cedar,') see *Swietenia febrifuga*.

CEDRELE'UM, (*κεδρελαιον*, from *κεδρος*, 'the cedar,' and *ελαιον*, 'oil,') The oil of cedar.

CED'RIA, (*κεδρια*, from *κεδρος*, 'cedar,') *Ced'rium*, *Ced'rinum*, *Ced'ri lacryma*, *Albitran*. The oil or resin which flows from the cedar of Lebanon. It was believed to possess great virtues. It has been supposed to be the same as the pyroligneous acid. See *Pinus Sylvestris*.

CEDRINUM VINUM, *Ced'ar Wine*. A wine prepared by steeping half a pound of bruised cedar berries in six French pints of sweet wine. It is diuretic and subastringent.

CEDRITES, (*κεδριτες*, from *κεδρος*, 'cedar,') A wine prepared from the resin of cedar and sweet wine. It was formerly employed as a vermifuge, &c.

CEDRIUM, (*κεδρον*,) *Cedria*.

CEDROMELA, see *Citrus medica*.

CEDRON, see *Simaba cedron*.

CEDRONELLA, *Melissa* — c. *Triphylla*, *Dra-cocephalum canariense*.

CEDROSTIS, *Bryonia alba*.

CEDRUS BACCIFERA, (*κεδρος*,) *Juniperus Lycia* — c. *Juniperus sabina* — c. Mahogani, *Swietenia mahogani*.

CEINTURE, (from *cingere*, *cinctum*, 'to gird,') *Cingulum*. Herpes zoster — c. *Blanche de la charoide*, Ciliary ligament — c. *Dartreuse*, Herpes zoster — c. *de Hildane*, *Cingulum Hildani* — c. *de Saint-Jean*, *Artemisia vulgaris* — c. *de Vif Argent*, *Cingulum mercuriale*.

CELANDINE, *Impatiens* — c. Common, *Chelidonium majus* — c. Lesser, *Ranunculus ficaria* — c. Poppy, *Stylophorum diphyllum*.

CELASTRUS, *Celastrus*, *Ceanothus Americanus* seu *triner'vis*, *New Jersey Tea*, *Red Root*, (F.) *Thé de Jersey*. Used by the American Indians, in the same manner as lobelia, for the cure of syphilis. It is slightly bitter and somewhat astringent. A strong infusion of the dried leaves and seeds has been recommended in aphthæ, and as a gargle in scarlatina.

CELASTRUS SCANDENS, *Climbing Staff-tree*. A climbing American shrub, the bark of which is said to possess emetic, diaphoretic, and narcotic properties.

CÉLATION, (F.) *Concealment*, (*celare*, *celatus*, 'to conceal,') A word used by French medico-legal writers for cases where there has been concealment of pregnancy or delivery.

CE'LE, (*κηλη*, 'a tumour, protrusion, or rupture,') a very common suffix, as in hydrocele, bubonocèle, &c. See Hernia.

CEL'ERY, (F.) *Celeri*. The English name for a variety of *Apium graveolens*.

CELERY, Wild, *Buben galbanum*.

CELETES, (*κηλητες*,) see Hernial.

CELIA, *Cerevisia*.

CÉLIAQUE, *Cœliac*.

CÉLINE, *Melissa*.

CE'LS, (*κηλς*, 'a spot, a stain,') A *macula* or spot on the skin. See *Kelis*.

CELL, *Cella*. A small cavity. The same signification as *cellule*. Also, a vesicle composed of a membranous *cell-wall*, *Seimone'mine*, with, usually, liquid contents. The whole organized body may be regarded as a congeries of cells having different endowments, each set being concerned in special acts, connected with absorption, nutrition, and secretion, wherever an action of selection or elaboration has to be effected. These cells are generally termed *primary*, *elementary*, or *primitive*. When they give rise to other cells, they are

at times, termed *parent* or *mother cells*; the resulting cells being termed *daughter cells*. See *Cytoblast*.

CELL, APOPLECTIC, see *Apoplectic cell*—c. Bone, *Lacuna* of bone—c. Bronchic, *Cellule*, bronchic—c. Caloigerous, see *Tooth*—c. Cancer, see *Cancer cell*—c. Daughtier, see *Cell*.

CELL DOCTRINE, *Cell the'ory*, *Scenothoe'ria*. The doctrine or theory of the formation of cells and of tissues from cells.

CELL, ELEMENTARY, see *Cell*.

CELL, EPIDERMIC or EPITHE'LIAL. The cells or corpuscles that cover the free membranous surfaces of the body, and which form the epidermis and epithelium, are termed '*epidermic* or *epithelial cells*.' They are developed from germs furnished by the subjacent membrane.

CELL, EPITHELIAT, *Cell*, epidermic—c. Fat, see *Fatty vesicles*.

CELL FORCE. The plastic or formative force seated in a cell, by the agency of which the different tissues are developed. See *Cell Life*.

CELL FORMATION, *Cytogeny*—c. Genesis, *Cytogeny*—c. Germ, *Cytoblast*, see *Module*.

CELL, GERMINAL, see *Cytoblast*.

CELL LIFE. The life which is possessed by the separate cells that form the tissues, and by which the nutrition of the tissues is presumed to be effected.

CELL, MOTHER, see *Cell*—c. Nucleated, see *Cytoblast*—c. Osseous, *Lacuna* of bone—c. Parent, see *Cell*.

CELL, PIG'MENT, (F.) *Cellule pigmentaire*. Pigment cells are mingled with the epidermic cells, and are most manifest in the coloured races. They are best seen on the inner surface of the choroid of the eye, where they form the *pigmentum nigrum*.

CELL, PRIMARY, see *Cell*—c. Primordial, see *Cell*—c. Sperm, see *Sperm*.

CELL TERRITORY. According to Virchow, the district of intercellular substance, which is ruled over by the cell, that lies in the middle of it, and exercises influence on the neighbouring parts.

CELL THE'ORY, *Cell doctrine*.

CELL WALL, see *Cell*, and *Cytoblast*.

CELLA, *Cell*—c. Media, see *Ventricles* of the Brain—c. Turcica, *Sella Turcica*.

CEL'LOID, *Celloï'des*, (badly compounded from *cella*, and *adēs*, 'resemblance.') Resembling a cell or cellule.

CELLULA, (dim. of *cella*), *Cellule*.

CELLULÆ AERÆÆ, *Cellules*, bronchic—c. Bronchicæ, *Cellules*, bronchic—c. Coli, see *Colon*—c. Cellulæ Malpighianæ, *Cellules*, bronchic—c. Mastoidæ, *Mastoid cells*—c. Medullares, see *Medullary membrane*—c. Pulmonales, *Cellules*, bronchic, see *Pulmo*—c. Sanguinis, *Globules* of the blood.

CELL'ULAR, *Cellula'ris*, *Cellulo'sus*, *Cell'ulous*, (from *cella* or *cellula*), (F.) *Cellulaire*, *Celluleux*. Relating or appertaining to or composed of cells or cellules.

CELLULAR MEM'BRANE, *Membra'na cellulosa* seu *Cellula'ris*,—*M. adiposa* seu *pinguedino'sa*, of some, *Pannicu'lus adiposus*,—Membrane formed of cellular or areolar tissue, (F.) *Membrane cellulaire*. Generally used for the tissue itself.

CELLULAR SYS'TEM. The whole of the cellular or areolar tissue of the human body.

CELLULAR TIS'SUE, *Te'ta cellul'a'ris* seu *cellulo'sa* seu *mucosa* seu *Hippoc'ratis cribrosa*, *Eth'myphæ*, *Contextus cellul'o'sus*, *Reticu'lated*, *filamentous*, *lam'inated*, *crib'riform*, *porous*, *areolar*, *connective*, and *mucous Tis'sue*, *Reticu'lar* or *cellular substance*, (F.) *Tissu cellulaire*, *réticulé*, *laminéux*, *cribleux*, *poreux*, *aréolaire*, *muqueux*, *conjonctif*, *unitif*, *Ethmose* of *Piorry*, &c., is the most common of all the organic tissues. It contains irregular *areolæ*—not *cells*, and, therefore,

more properly called *areolar tissue*—between the fibres, as well as serum, fat, and the adipous tissue. Of the fibres, some are of the *yellow* elastic kind; but the greater part are of the *white* fibrous tissue, and they frequently present the form of broad flat bands, in which no distinct fibrous arrangement is perceptible. See *Fibrous*.

The cellular tissue or texture unites every part of the body, determines its shape, and by its elasticity and contractility, and by the fluid which it contains in its areolæ, facilitates the motion of parts on each other.

Cellular tissue has been divided by anatomists into the external, general or common cellular tissue—*textus cellul'aris interme'dius seu las'us*, which does not penetrate the organs,—the cellular texture which forms the envelopes of organs—*textus cellul'aris stric'tus*, and that which penetrates into the organs, accompanying and enveloping all their parts—the *textus cellul'aris stipi'tus*, constituting the basis of all the organs. It has likewise been termed *Textus organ'icus seu parenchyma'tis*.

CELLULAR TISSUE OF BONES, see *Cancelli*.

CELL'ULE, *Cell'ula*, (dim. of *cella*.) A small cavity. (F.) *Cellule*. *Cellules* are the small cavities between the laminae of the cellular tissue, corpora cavernosa, &c.

CELLULE À NOYAU, see *Cytoblast*.

CELLULES or CELLS, BRONCHIC, *Vesiculæ* seu *Cell'ulæ Bronchicæ* seu *Pulmona'les* seu *Aë'reæ* seu *Malpighi'anae*, *Po'ri pulmo'num*, (F.) *Cellules bronchiques*. The air-cells of the lungs. See *Pulmo*.

CELLULES BRONCHIQUES, *Cellules*, bronchic—c. *Cancereux*, see *Cancer cell*—c. *des Os*, *Lacunæ* of Bone—c. *Ossenses*, *Lacunæ* of Bone—c. *Pigmentaires*, *Cells*, pigment—c. *Rouges du sang*, see *Globules*, red, of the blood.

CELLULEUX, *Cellular*.

CELLULIN, *Cellulose*.

CELLULITIS, (*cellule*, and *itis*), *Ethmyphitis*—c. *Venenata*, see *Wound*.

CELLULOFIBROUS TUMOUR, *Tumour*, *fibro-cellular*.

CELLULOSE, *Cell'ulin*, (same etymon as *Cellule*.) The substance which is left after the action upon any kind of vegetable tissue of such solvents as are fitted to dissolve out the matter deposited in its cavities and interstices. The tunicated or ascidian mollusca have, in their integuments, a considerable quantity of it, and it exists in the Corpora amylacea of the brain and other parts. It is identical with starch, in the proportion of its constituents, and forms the principal part of the thickness of the walls of the cells, vessels, &c., of which the vegetable organism is composed.

CELLULOSUS, *Cellular*.

CELLULOUS, *Cellular*.

CE'LO, (from *κηλη*, 'a rupture,') in composition, rupture.

CELOGOG'IA, (from *celo*, and *λογος*, 'a discourse.') The doctrine of hernia. A treatise on hernia.

CELOSO'MUS, (F.) *Célosome*, (*celo*, and *σῶμα*, 'body.') A monster in which the trunk is malformed, and evagination or displacement of the viscera exists.

CELOTOM'IA, (*κητοτομία*, from *celo*, and *τεμνειν*, 'to cut,') *Celotom'ia*, *Celot'omy*. An operation, formerly employed for the radical cure of inguinal hernia; which consisted, principally, in passing a ligature round the hernial sac and spermatic vessels. It necessarily occasioned atrophy and loss of the testicle; and did not secure the patient against the return of the disease. The intestines were, of course, not included in the ligature.

Also, the operation for hernia is general.—*Herni-ot'omy*.

CELOTOMUS, *Herni-ot'omus*. A knife used in the operation for hernia. Adjectively, it means relating to celotomy, like *Celotom'icus*.

CELSA. A term, used by Paracelsus for a cutaneous disease, dependent, according to him, on a false or heterogeneous spirit or vapour, concealed under the integuments, and endeavouring to escape. Perhaps the disease was *Urticaria*.

CELSUS, METHOD OF, see *Lithotomy*.

CELTIS OCCIDENTALIS, *Sugarberry*, *Hackberry*, *Nettle-tree*, *Beaverwood*. Order, *Ulmaceæ*; indigenous, flowering in May. The bark is said to be anodyne and cooling; the berries are sweet and astringent. It has been used in dysentery.

CEMERO NUTS, see *Pinus cembra*.

CEMENT, *Cemen'tum*, see *Lute*, and *Tooth*. Also, a glutinous substance introduced into a carious tooth to prevent the access of air or other extraneous matters. The following is an example: *R. Sandarac. ʒij; Mastich. ʒi; Succi. gr. x. Ether. ʒi*; Dissolve with the aid of heat. *Ostentatier's Cement for the teeth* is prepared of finely powdered caustic lime, thirteen parts; anhydrous phosphoric acid, twelve parts. When introduced into a carious tooth, it becomes solid in about two minutes.

CEMENTERIUM, *Crucible*.

CEMENTUM, see *Tooth*.

CEMOS, (*κμος*, 'a muzzle,') *Capistrum*.

CENANGIA, *Ceneangia*.

CENCHRON, (*κενχρον*,) *Panicum miliaceum*.

CENDRE DU LEVANT, *Soda*.

CENDRE, (*cinis*, gen. *cineris*, 'ashes,') *Cin-eritus*.

CENDRES GRAVELÉES, see *Potash—c. de Saumet*, see *Potash*.

CENEANGI'A, *Cenangi'a*, (*κενος*, 'empty,' and *αγγιον*, 'a vessel,') *Inanition*. Empty state of vessels.

CENEMBATE'SIS, (*κενος*, 'empty,' and *εμβαλω*, 'I enter,') *Paracentesis*. Also, the act of probing a wound or cavity, *Melo'sis*.

CENEONES, (*κενωσις*, 'an empty space,') *Flanks*.

CENESTHÉSIE, *Cænæsthesia*.

CENIGDAM, *Ceniplam*.

CENIGOTAM, *Ceniplam*.

CENIPLAM, *Cen'igdam*, *Cen'igotam*, *Cen'ipolam*. The name of an instrument anciently used for opening the head in epilepsy.

CENIPOLAM, *Ceniplam*.

CENO'SIS, (*κενωσις*, from *κενω*, 'to empty,') *Ine'sis*, *Inethm'us*. Evacuation. It is sometimes employed synonymously with inanition, and opposed to repletion.—*Exinanit'io*.

CENOSPUDIA, (*κενωσπυδία*, from *κενος*, 'empty,' and *σπυδιον*, 'application,') *Aphelxia otiosa*.

CENOTIC, (*κενωτικός*, from *κενω*, 'to empty,') *Drastic*.

CENO'TICA. Diseases affecting the fluids. Morbid discharges or excess, deficiency or irregularity of such as are natural. The first order, class *Genetica*, of Good. Also, *Drastics*.

CENTAUREA BE'HEN, (*κενταυρία*, after the *Centaure*, *Chiron*,) *Serrat'ula be'hen*, *Be'hen a'biad*, *Be'hen al'bum*, *Be'hen*, *White Be'hen*. Ord. *Gentianeæ*. Astringent.

CENTAUREA BENEDICTA, *Car'duus benedic'tus*, *Chi'cus sylvest'ris* seu *benedic'tus*, *Cardiobot'anum*, *Bless'd Ho'ly This'tle*, *Virgin-Mary Thistle*, (F.) *Chardon béni't*. Fam. *Cynaroecephalææ*. *Sex. Syst.* Syngenesia Polygamia frustranea. A strong decoction of the herb is emetic:—a strong infusion, diaphoretic; a light infusion, tonic and stomachic. Dose, gr. xv to ʒj of the powder.

CENTAUREA CALCITRA'PA, *Calceitra'pa*, *Calceat'rop'ola*, *Car'duus solstitia'lis* seu *stella'tus*, *Ja'cea*

ramosis'sima, *Cucostil'abus*, *Calceitrap'pa stella'ta* seu *hippophas'sum*, *Stella'ta rupi'na*, *Centaurea stella'ta*, *Common Star-Thistle*, *Star-Knapweed*, (F.) *Centaurea étoilée*, *Chardon étoilé*, *Chausse-trappe*, *Piguerole*. It is possessed of tonic properties, and has been given in intermittents, dyspepsia, &c. It is not much used.

CENTAUREA CENTAURI'UM, *Rhapont'icum vulgare*, *Centauri'um mag'num* seu *ma'jus* seu *officina'le*, *Great'er Centaury*, (F.) *Centauree grande*. It is a bitter; and was formerly used as a tonic, especially the root.

CENTAUREA CY'ANUS, *Cy'anus*, *Blue bottle*, *Cam-pacer*, *Blue bonnets*, (Sc.) *Blawort*, (F.) *Blavelle*, *Blavéole*, *Blavérille*, *Bluet*, *Barbeau*, *Anbifoïn*, *Casse-lunette*. The flowers were once much used as a cordial, tonic, &c., and in ophthalmia. They are now forgotten.

CENTAUREA STELLATA, *Centaurea calcitrapa*.

CENTAURÉE ÉTOILÉE, *Centaurea calcitrapa—c. Grande*, *Centaurea centaurium—c. Petite*, *Chironia centaurium*.

CENTAUREUM, *Chironia centaurium*.

CENTAURIS, gen. *Centaur'idis*, *Chironia centaurium*.

CENTAURIUM MAGNUM, (*κενταυριον*, and *κενταυριον*), *Centaurea centaurium—c. Minus vulgare*, *Chironia centaurium—c. Officinale*, *Centaurea centaurium—c. Parvum*, *Chironia centaurium*.

CENTAURY, AMERICAN, *Chironia angul-aris—c. European*, *Chironia centaurium—c. Great'er*, *Centaurea centaurium—c. Lesser*, *Chironia centaurium*.

CENTESIS, (*κεντησις*, from *κεντεω*, 'I perforate,') *Paracentesis*, *Puncture*.

CENTI, (from *centum*, 'a hundred,') in composition, 'a hundred.'

CENTIGRAMME (F.), (*centi*, and *γραμμα*, 'gramme,') *Centigram'ma*. The hundredth part of a gramme. A centigramme is equal to about the fifth part of a French grain, gr. .1543 Troy.

CENTILITRE, *Centili'tra*, (*centi*, and *λιτρα*, 'litre,') An ancient Greek measure for liquids:—the hundredth part of a litre—equal to nearly 2.7053 fluidrachms.

CENTIMÈTRE, *Centim'eter*; (*centi*, and *μέ-τρε*,) the hundredth part of a metre—equal to about four lines, .3937 English inch.

CENTIMORBIA, (*centi*, and *morbus*, 'a disease,') *Lysimachia nummularia*.

CENTINERVIA, (*centi*, and *nervus*, 'a nerve,') *Plantago*.

CENTINODE, *Polygonum aviculare*.

CENTINODIA, (*centi*, and *nodus*, 'a knot,') *Polygonum aviculare*.

CENTO VIRGINALIS, ('virginal tent covering,') *Hymen*.

CENTRAD, see *Central aspect*.

CENTRADIAPH'ANES, *Cataracta centra'lis*, (*κεντρον*, 'centre,' a, privative, and *διαφανς*, 'transparent,') *Cataract* owing to obscurity of the central portion of the crystalline.

CENTRAL, (from *centre*,) Relating or appertaining to the centre.

CENTRAL ARTERY OF THE RET'INA, *Arte'ria Centra'lis Ret'inæ*, *Central Ar'tery of Zinn*. This artery is given off from the arteria ophthalmica, and penetrates the optic nerve a little behind the ball of the eye; running in the axis of the nerve, and spreading out into many small branches upon the inside of the retina. When the nerve is cut across near the eye, the orifice of the divided artery is observable. This was formerly called *Po'ris Op'ticus*.

CENTRAL AS'PECT. An aspect towards the centre of an organ.—*Barclay*. *Centrad* is used by the same writer adverbially, to signify 'towards the central aspect.'

CENTRE, *Centrum*, (κεντρον,) (F.) Centre. The middle point of a figure or body.

CENTRE OF ACTION. The viscus in which the whole or a great part of any function is executed, and to which several other organs contribute. Thus, the vital activity seems to be wholly centred in the stomach, during chymification; in the duodenum, during chyliification. In like manner, the uterus becomes a centre of action during gestation.

CENTRE, EPIGAS'TRIC. The ganglions and nervous plexuses, formed by the great sympathetic and pneumogastric nerves, in the epigastrium, around the cœliac artery; where the impressions received from various parts of the body seem to be centred.

CENTRE OF FLUX'ION. The part towards which fluids are particularly attracted. An irritated organ is said to be a centre of fluxion.

CENTRES, NER'VEOUS, (F.) *Centres nerveux*. The organs whence the nerves originate; as the brain and spinal marrow.

CENTRE, OPTIC, see Optic centre.

CENTRE, O'VAL, *Centrum Ovale*, C. O. *Vieussennii*, *Tegumentum ventriculorum cerebri*. When the two hemispheres of the brain are sliced away, till on a level with the corpus callosum, the medullary part in each is of an oval shape: hence called *centrum ovale minus*, (F.) *Centre médullaire hémisphérique*. The two centres of the opposite sides, together with the corpus callosum, form the *centrum ovale* of *Vieussens*. *Vieussens* supposed all the medullary fibres to issue from that point, and that it was the great deservatory of the animal spirits.

CENTRE, PHREN'IC, *Tendinous Centre* of the *Diaphragm*, *Centrum Phrenicum* seu *Nerveum seu Tendinosum* seu *Tendinum*, (F.) *Centre phrénique* ou *tendineux* du *Diaphragme*. The central aponeurosis or cordiform tendon of the diaphragm.

CENTRE OF SYMPATHET'IC IRRADIATIONS, (F.) *Centre d'irradiations sympathiques*. Any organ which excites, sympathetically, the action of other organs, more or less distant from it; and with which it seems to have no immediate communication.

CENTRE, TENDINOUS, OF THE DIAPHRAGM, Centre, phrenic.

CEN'TRIUM, (κεντριον, from κεντρω, 'to prick.') A plaster for a stitch in the side, mentioned by Galen.

CENTROMYRINE, *Ruscus*.

CENTROSTAL'TIC, *Centrostalticus*, (κεντρον, and στελλω, 'I send.') An epithet applied by Dr. Marshall Hall to the action of the *vis nervosa* in the spinal centre.

CENTRUM, (κεντρον,) Centre, see *Vertebræ*—c. Commune, Solar plexus—c. Nerveum, Centre, phrenic—c. Opticum, Optic centre—c. *Ovale*, Centre, oval—c. *Ovale minus*, see Centre, oval—c. *Ovale* of *Vieussens*, Centre, oval—c. *Semicirculare geminum*, *Tænia semicircularis*—c. *Tendinosum*, Centre, phrenic.

CEN'TRUM VITA'LE, ('vital centre,') *Nodus* seu *Fons vitalis*, (F.) *Nœud vital*. A term applied, at times, to the medulla oblongata; at others, to the medulla oblongata and the medulla spinalis as far as the second cervical nerve of the spinal marrow, in any part of which a wound appeared to be instantly fatal. It seems to be essentially the nervous centre of respiration and deglutition. M. Flourens, at one time, placed the *vital point* at the apex of the V of gray matter at the nib of the calamus scriptorius; but he afterwards extended it.

CENTRY, *Chironia angularis*.

CENTUM CAPITA, ('a hundred heads,') *Eryngium campestre*.

CENTUMNODIA, (*centum*, 'a hundred,' and *nodus*, 'a knot,') *Polygonum aviculare*.

CEPA, ('an onion,') *Allium cepa*—c. *Ascalonica*, *Bulbus esculentus*, *Échalotte*—c. *Marina*, *Scilla*—c. *Victorialis*, *Allium victorale*—c. *Vulgaris*, *Allium cepa*.

CEPÆA, (κηπος, 'a garden,') *Veronica beccabunga*.

CEPHAËLINUM, *Emetine*.

CEPHAËLIS IPECACUANHA, *Ipecacuanha*. CEPHALÆ'A, (κεφαλαία, from κεφαλή, 'head,')

Head'ache, (F.) *Céphalée*. Some use the term synonymously with cephalalgia; others, for a periodical headache; others, again, for a more violent headache than cephalalgia implies; and others for a chronic headache. The last was its ancient signification.

Cephale'a spasmodica, *Cephalalgia spasmodica*, C. *Nauseosa*, *Sickhead'ache*, is characterized by partial, spasmodic pain; often shifting from one part of the head to another: chiefly commencing in the morning, with sickness and faintness. It is extremely apt to recur, notwithstanding every care.

CEPHALÆA ARTHRITICA, *Cephalagra*—c. *Hemicrania*, *Hemicrania*—c. *Nauseosa*, C. *Spasmodica*—c. *Pulsatilis*, *Crotaphe*.

CEPHALÆMATO'MA, gen. *Cephalæmatomatis*, (F.) *Céphalématome*, (cephale, άμα, 'blood,' and oma;) *Cephalæmatoma neonatorum*, *Ecchymoma capitis*, *E. capitis recens natorum*, *Thrombus neonatorum*, *Abcessus capitis sanguineus neonatorum*, *Tumor capitis sanguineus neonatorum*, *Cephalophyma*, *Craniohematomus*. A sanguineous tumour, sometimes developed between the pericranium and the bones of the head of new-born children. Similar tumours are met with occasionally above other bones, and at all periods of existence.

CEPHALÆMATOMA NEONATORUM, *Cephalæmatoma*.

CEPHALÆ'MIA, (cephale, and άμα, 'blood,') *Cephalohæmia*, *Hyperhæmia cerebri* seu *Capitis*, *Encephalohæmia*, *Hyperæmia* or congestion of the brain, (F.) *Hyperémie* ou *Congestion du cerveau*, *Encephalohémie*, *H. cérébrale*, *Congestion cérébrale*. Accumulation of blood in the vessels of the brain.

CEPHALAGO'GUS, *Cephaloductor*, *Capitiductor*, (cephale, and αγωγος, 'a leader, a driver,') An instrument used for drawing down the foetal head.

CEPH'ALAGRA, (cephale, and αγρα, 'seizure,') *Cephale'a arthritica*, *Meningitis arthritica*. Gout in the head.

CEPHALAGRA'PHIA, (cephale, and γραφή, 'a description,') An anatomical description of the head.

CEPHALAL'GIA, *Cephaloponia*, *Cephalodynia*, *Encephalodynia*, *Homoponia'gia*, (cephale, and αλγος, 'pain,') *Encephalalgia*, *Dolor capitis* seu *cephalicus*, *Solida*, *Pain in the head*; *Hedache*, (Sc.) *Hedeverk*, (Prov.) *Hedewark*, (F.) *Céphalalgie*, *Mal à tête*. Every kind of headache, whether symptomatic or idiopathic, is a cephalalgia. It is ordinarily symptomatic, and has to be treated accordingly.

CEPHALALGIA CONTAGIOSA, *Influenza*—c. *Inflammatoria*, *Phrenitis*.

CEPHALALGIA PERIOD'ICA, *Febris intermittens cephalica* *laren'ta*, *Intermittent head'ache*. Headache which returns periodically; properly, perhaps, a form of neuralgia.

CEPHALALGIA PULSATILIS, *Crotaphe*—c. *Spasmodica*, see *Cephalæa*.

CEPHALALOG'IA, (cephale, and λογος, 'a discourse,') An anatomical dissertation on the head.

CÉPHALANTHE D'AMÉRIQUE, *Cephalanthus occidentalis*.

CEPHALANTHUS OCCIDENTALIS, (*cephale*, and *ανθος*, 'a flower.') *Buttencood shrub*, *Buttombush*, *White Ball*, *Little Snowball*, *Sicampwood*, *Pond Dogwood*, *Globe-flower*, (F.) *Céphanthe d'Amérique*, *Bois de Mûrais*. An ornamental shrub, *Ord. Rubiaceæ*; *Sec. Syst. Tetrandria Monogynia*, which grows all over the United States, near streams and ponds, and flowers in July and August. The bark of the root has been used as an antiperiodic tonic.

CEPHALARTICA, (*cephale*, and *αρτήρ*, 'to make complete.') Cephalic remedies.

CEPHALATOM'IA, *Cephalotom'ia*, (*cephale*, and *τομή*, 'to cut.') Anatomy, or dissection, or opening of the head.

CEPHALE, (*κεφαλή*), Head.

CEPHALÉMATOME, *Cephalématoma*.

CEPHAL'IC, (*κεφαλικός*), *Cephal'icus*, *Capita'lis*, (from *cephale*.) (F.) *Céphalique*. Relating to the head.

CEPHAL'IC REM'EDIES, *Cephal'ica* seu *Capita'lia remedia*, are remedies capable of relieving affections of the head, especially headache:—*Cephalar'tica*.

CEPHAL'IC VEIN, *Ve'na Cephal'ica* seu *Cap'itis*, (F.) *Veine céphalique*, *Veine radiale cutanée* of Chaussier. The great superficial vein at the outer part of the arm and fore-arm. It begins on the back of the hand, by a number of radicles, which unite into a single trunk, called the *Cephal'ic of the Thumb*, *Cephal'ica Pol'licis*, (F.) *Veine céphalique du pouce*. It ascends along the anterior and outer part of the fore-arm, where it forms the *superficial radial*. At the fold of the elbow it receives the *median cephalic*, ascends along the outer edge of the biceps, and opens into the axillary vein. The name *Cephalic* was given to it by the ancients, because they thought it had some connection with the head, and that blood-letting ought to be performed on it in head affections.

Chaussier calls the internal jugular, *Veine céphalique*, and the primary or common carotid artery, *Artère céphalique*.

CEPHALIC VERSION, see Turning.

CEPHAL'ICI, (*MORBI*), in the classification of Dr. Wm. Farr, brain diseases.

CEPHALIDIUM, (dim. of *cephale*.) See *Caput*.

CEPHALIN'E, (*κεφαλίνη*, from *cephale*.) The base or root of the tongue.

CÉPHALIQUE, Cephalic.

CEPHALITIS, (*cephale*, and *itis*), Phrenitis.

CEPHALIUM, (from *cephale*), see *Caput*.

CEPHALOCENTE'SIS, (*cephale*, and *κέντησις*, 'puncturing.') The puncturing of the head,—as for hydrocephalus.

CEPHALODUCTOR, (*cephale*, and *ductor*, 'a leader.') Cephalagogus.

CEPHALODYM'IA, *Encephalodym'ia*; (*cephale*, and *ἔω*, 'I enter into.') A class of double monstrosities, in which the heads are united. It is divided into two genera, *Frontodym'ia* and *Bregmatodym'ia*: in the former the union being between the ossa frontis; in the latter between the bregmata.

CEPHALODYNIA, (*cephale*, and *ὄδυνη*, 'pain.') Cephalalgia.

CEPHALŒDEMA, (*cephale*, and *οἰῆμα*, 'a swelling.') Hydrocephalus.

CEPHALOG'RAPHY, *Cephalograph'ia*; (*cephale*, and *γραφη*, 'a description.') An anatomical description of the head.

CEPHALOHŒMIA, Cephalæmia.

CEPHALOID, (*cephale*, and *εἶδος*, 'resemblance.') Encephaloid.

'CEPHALOL'OGY, *Cephalolog'ia*; (*cephale*, and *λογος*, 'a discourse.') A treatise on the head.

CEPHALOMA, gen. *Cephalo'matis*, (from *cephale* and *ωμα*.) Encephaloid.

CEPHALOM'ELUS, (F.) *Céphalomèle*; (*cephale*, and *μελος*, 'a limb.') A monster having one or two accessory limbs on the head.

CEPHALOMENINGITIS, (*cephale*, *μηνίγξ*, 'a membrane,' and *itis*, 'inflammation.') Meningocephalitis.

CEPHALOM'ETER, (F.) *Céphalomètre*, (*cephale*, and *μετρον*, 'measure.') An instrument for measuring the different dimensions of the fetal head, during the process of accouchement. A kind of forceps.

CEPHALON'OSUS, (*cephale*, and *ὄσος*, 'disease.') This term has been applied to the *Febria Hungar'ica*, in which the head is much affected. See *Fever*, *Hungaric*. Others have so called any cerebral disease or fever.

CEPHALOPAGES, (*cephale*, and *πηνυμ*, 'to join together.') Symphycephalus.

CEPH'ALOPHARYNGŒ'US, (*cephale*, and *φάρυγξ*, 'the pharynx:' belonging to the head and pharynx.) Winslow has given this name to the portion of the constrictor pharyngis superior, which is attached, above, to the inferior surface of the basilar process of the os occipitis. The *Ceph'alo-pharynge'al Aponeuro'sis* is a thin, fibrous membrane, which is attached to the basilar process, and gives insertion to the fibres of the constrictor superior pharyngis.

CEPHALOPHYMA, gen. *Cephalophy'matis*, (*cephale*, and *φύμα*, 'a tumour.') Cephalématoma.

CEPHALOPONIA, (*cephale*, and *πῶς*, 'pain.') Cephalalgia.

CEPHALORACHIDIAN, Cephalospinal.

CEPHALOSOMATODYM'IA, *Encephalosomatodym'ia*; (*cephale*, *σωμα*, 'body,' and *ἔω*, 'I enter into.') A double monstrosity, in which the union is between the heads and the trunks. Of this there are varieties:—for example, *Infra-maxillosternodym'ia*, where the union is with the inferior maxillary bones and sterna; and *Prosoposternodym'ia*, between the faces and sterna.

CEPH'ALOSPINAL, *Cephalospina'lis*, *Ceph'alorachid'ian*, *Enceph'alorachid'ian*, *Cer'ebrospinal*, *Cer'ebroarachid'ian*, *Cra'niospinal*. A hybrid term, (*cephale*, and *spina*.) Belonging to the head and spine.

CEPH'ALOSPINAL FLU'ID, *Ceph'alorachid'ian flu'id*, *Cer'ebrospinal flu'id*, *Flu'idum cer'ebrospinal'e*, *Liquor cer'ebro-spinal'is*, *Subarachnoid'ean flu'id*, (F.) *Liquide céphalorachidien*, is an exhaled fluid, which is found beneath the arachnoid, wherever the pia mater exists in connexion with the brain and spinal cord. It seems to have a protecting office, and to keep up a certain degree of pressure on the organ—at least in the spinal canal.

CEPH'ALOTHO'RACOSTERUM'ENUS, (*cephale*, *θώραξ*, 'the chest,' and *στερεν*, 'to rob.') A monster without head or chest.

CEPHALOTOMIA, Cephalotomia.

CEPHALOTRIBE, (F.) An instrument invented by Baudeloque, the nephew, for crushing the head of the fetus in utero; (*cephale*, and *τριβω*, 'I bruise.') It consists of a strong forceps, the blades of which are solid: 16 lines broad, and 3 thick. The handles are perforated at their extremity to receive a screw with three threads, the direction of which is very oblique, so as to allow great rapidity of rotation, and the screw is moved by a winch 6 inches long, to increase the force of the pressure. The bones of the head are easily crushed by it.

CEPHALOTRIP'SY, *Cephalotrip'sis*; (same etymon.) The operation of crushing the head of the fetus in utero.

CEPHALOTRYPESIS, (*cephale*, and *τρύπησις*, 'perforation.') Trepanning.

CEPHALOLOXIA, (*cephale*, and *λοξος*, 'oblique.') Torticollis.

CEPULLA, (dim. of *cepa*.) Allium cepa.

CER, (*κρρ*.) Heart.

CERA FLA'VA, (*κρρος*.) *Ce'rus*, Yellow Wax, (F.) *Cire Jaune*. A peculiar concrete animal substance prepared by the bee, and by some plants, as the *Cerax'ylon* and *Myri'ca cerif'era*. Its colour is yellow, and smell like that of honey, but both are lost by bleaching. The wax is then called *Cera alba*, White wax, (F.) *Cire Jaune*. It is demulcent and emollient; is sometimes given in the form of emulsion, in diarrhoea and dysentery, but is chiefly used in cerates and ointments.

CERÆÆ, (*κερας*, 'a horn.') The Cornua of the uterus.

CERAM'ICÆ, *Ceramit'is*, (*κεραμος*, 'potter's earth.') A sort of earth used as a cataplasm in peripneumony.

CERAMIUM, (*κεραμιον*.) Amphora—c. Helminthochortus, Corallina Corsicana.

CERAMURIA, (*κεραμος*, 'earth,' and *ουρον*, 'urine.') See Urine.

CERANTHEMUS, (*κρρος*, 'wax,' and *ανθος*, 'a flower.') Propolis.

CERAS, (*κερας*, gen. *κερατος*, 'horn.') Cornu; also, the Cornea.

CERASION, (*κερασιον*.) see *Prunus cerasus*.

CERAS'MA, gen. *Cerasm'atis*, (*κερασμα*, from *κερασιον*, 'to mix: something mixed.') A mixture of hot and cold water. *Metaceras'ma*.

CERASUM, (*κερασιον*.) see *Prunus cerasus*.

CERASUS ACIDA, (*κερασος*.) *Prunus cerasus*—c. Avium, *Prunus avium*, P. nigra—c. Dulcis, *Prunus nigra*—c. Hortensis, *Prunus cerasus*—c. Laurocerasus, *Prunus laurocerasus*—c. Padus, *Prunus padus*—c. Racemosus *sylvestris*, *Prunus padus*—c. Rubra, *Prunus cerasus*—c. Serotina, *Prunus Virginiana*—c. Virginiana, *Prunus Virginiana*—c. Vulgaris, *Prunus cerasus*.

CÉRAT BLANC ou DE GALIEN, Ceratum Galeni—c. de Blanc de Baleine, Ceratum cetacei—c. de Goulard, Ceratum plumbi—c. pour les Lèvres, Cerate for the lips—c. de Plomb composé, Ceratum plumbi compositum—c. de Savon, Ceratum Saponis—c. de Suracétate de plomb, Ceratum plumbi superacétatis.

CERATE, (*κρρωτον*, from *κρρος*, *cera*, 'wax.') *Cero'tê*, *Cerel'sum*, *Cero'ma*, *Cero'nium*, *Cero'tum*, *Cero'leum*, *Oleocera'tum*, *Cera'tomalag'ma*, (F.) *Cérat*, *Oléo-cérolé*, *Éléocérolé*, *Cérolé*, *Liparoidé*. A composition of wax, oil, or lard, without other ingredients.

CERATE, Simple cerate, *Cera'tum*, *Cera'tum simplex*, C. ad'ipis (Ph. U. S. 1863.) (F.) *Cérat Simple*. (White wax, $\mathfrak{Z}$ iv, Lard, $\mathfrak{Z}$ viiij.) It is applied as an emollient to excoriations, &c. See Ceratum Cetacei.

CERATE, BELLEVILLE'S, see Unguentum Hydrargyri nitrico-oxidi. Cerate, blistering, see Emplastrum Lyttæ.

CERATE OF CALAMINE, *Cera'tum Calami'næ* seu *Calami'næ præpara'tæ* seu *Carbona'tis zin'ci impu'ri* seu *Zinci Carbona'tis* seu *Lap'idis Calami'næ* seu *Epulot'icum*, *Ce'rate* of Car'bonate of Zinc, Tur'ner's *Ce'rate*, *Heal'ing Salve*, (F.) *Cérat de Pierre Calaminaire*, C. de Calamine, (*Calamin*, *Ceræ flavæ*, $\mathfrak{a}\mathfrak{a}$ $\mathfrak{Z}$ ijj, *adipis* lbj. Melt the wax and lard together, and, on cooling, add the carbonate of zinc and stir till cool.—Ph. U. S.)

CERATE OF CANTHARIDES, *Cera'tum Canthar'idis* seu *Lyttæ*, *Blis'ter Oint'ment*, *Oint'ment of Span'ish Flies*, *Unguentum ad vesicato'ria* seu *Pul'veris Mel'oës vesicato'rii* seu *epispas'ticum fort'ius*, (F.) *Cérat de Cantharides*. (*Spermaceti cerate* $\mathfrak{Z}$ vj, *Cantharides*, in powder, $\mathfrak{Z}$ j. The cerate being softened by heat, stir in the flies.) This cerate of the European Pharmacopœias is used to keep blisters, issues, &c., open. See Un-

guentum Lyttæ. For the Cerate of Cantharides of the Ph. U. S., see Emplastrum Lyttæ.

CERATE OF EXTRACT OF CANTHARIDES, see Emplastrum Lyttæ.

CERATE, GOULARD'S, Ceratum plumbi compositum.

CERATE, KIRK'LAND'S NEU'TRAL. (*Diachyl*. $\mathfrak{Z}$ viiij, *Olive oil*, $\mathfrak{Z}$ iv, *Prepared chalk*, $\mathfrak{Z}$ iv: when nearly cool, add *Acet. dest.* $\mathfrak{Z}$ iv, *plumb superacet.* $\mathfrak{Z}$ ij.) A cooling emollient.

CERATE, LEAD, COMPOUND, Ceratum plumbi compositum.

CERATE OR POMM'ATUM FOR THE LIPS, *Cera'tum labia'le ru'brum Pomma'tum ad la'bia demulcen'da*. (F.) *Cérat ou Pommade pour les lèvres*, (Wax 9 parts; Oil, 16 parts;—coloured with *alkanet*.—Ph. P.)

CERATE, MAR'SHALL'S. (*Palm oil* $\mathfrak{Z}$ vi, *Calomel* $\mathfrak{Z}$ j, *Sugar of lead* $\mathfrak{Z}$ ss, *Ointment of nitrate of mercury* $\mathfrak{Z}$ ij.)

CERATE, RESIN, COMPOUND, Ceratum Resinæ compositum—c. Savine, Ceratum sabinae—c. Soap, Ceratum saponis—c. Spermaceti, Ceratum cetacei—c. of Superacetate or sugar of lead, Ceratum plumbi superacétatis—c. Turner's, Cerate of calamine—c. of Carbonate of zinc, Cerate of calamine.

CERATECTOM'IA, *Cerectom'ia*, *Kerectomy*, *Ceratectomy*, (*cerato*, and *εκτομος*, 'cut out.') An incision through the cornea. See Ceratotomy.

CERATIA, (*κερατια*.) Ceratonium siliqua.

CERATI'ASIS, *Kerati'asis*, (*κερας*, gen. *κερατος*, 'horn.') A morbid condition, characterized by cornuous growths.

CERATION, (*κερασιον*.) Siliqua.

CERATI'TIS, *Kerati'tis*, (*cerato*, and *itis*.) *Ceratodei'tis*, *Keratodei'tis*, *Ceratomeningi'tis*, *Cornei'tis*, *Inflamma'tio cor'næ*. Inflammation of the cornea.

CERATIUM, Ceratonium siliqua.

CER'ATO, (from *κερας*, gen. *κερατος*.) In composition, a cornu or horn, or the cornea. See Ceras.

CERATOCE'LE, *Cer'atocèle*, *Keratoce'le*, *Aqu'ula*, *Uval'tio*, *Prominent'ia Cor'næ*, *Her'n'ia Cor'næ*, *Cerato'deoc'elê*, (*cerato*, and *κκλη*, 'tumour.') A protrusion of the transparent cornea, or rather of the membrane of the aqueous humour, through an opening in the cornea.

CER'ATOCRI'COID, *Ker'atocri'coïd*, *Cer'ato* seu *Ker'ato-cricoides mus'culus*, *Mer'kel's muscle*. A muscle of the larynx, first described (1857) by Merkel, of Leipsic. It is not always found, and when it is, it exists only on one side. It arises close to the origin of the outer fibres of the crico-arytenoides posticus, passing obliquely upwards and outwards, and is attached to the posterior margin of the inferior cornu of the thyroid cartilage. Its action is to fix the lower cornu backwards and downwards.

CERATODEITIS, Ceratitis.

CERATODEOC'ELE, Ceratocèle.

CERATODEONYXIS, Ceratonyxis.

CERATODES MEMBRANA, (*cerato*, and *odes*.) Cornea.

CER'ATOGEN'ESIS, (*cerato*, and *γενεσις*, 'generation.') The formation of horn or of a ceratoma—*Ceratosis*.

CERATOGL'OS'SUS, *Keratoglos'sus*, (*cerato*, and *γλωσσα*, 'the tongue.') A muscle, extending from the great cornu of the os hyoides to the base of the tongue. It is a part of the hyoglossus.

CERATOIDES, (*cerato*, and *ειδς*, 'resemblance.') Cornea.

CERATOLEUCOMA, gen. *Ceratoleuco'matis*, Leucoma.

CERATO'MA, gen. *Cerato'matis*, (from *cerato*, and *ομα*.) A horny growth or formation.

CERATOMALA'CIA, *Ker'atomala'cia*, (*cerato*, and *μαλακία*, 'softness.') Softening of the cornea, the result of inflammation or of starvation.

CERATOMALAGMA, gen. *Cer'atomalg'matis*, (*cerato*, and *μαλασσειν*, 'to soften.') Cerate.

CERATOMENINGITIS, (*cerato*, *μυμίζ*, 'a membrane,' and *itis*.) Ceratitis.

CERATOMENINX, Cornea.

CERATONIA SIL'IQUA, (*keras*, gen. *κερας*, 'a horn'—from its hornlike pod.) The *Algaroba* or *St. John's bread*, *Carob Tree*, *Cerat'ium*, *Cerati'a*, *Sil'iqua dulcis*, *Caro'ba Alnabati*, *Sweet'pod*, (F.) *Caroubier*, (*Fruit*, *Carouge*.) The fruit of the *Ceratonía siliqua* is mucilaginous, and employed in decoction where mucilages are indicated.

CERATONYX'IS, *Keratonyx'is*, *Cerato'deonyx'is*, (*cerato*, and *νύσσα*, 'I puncture.') An operation by which the crystalline is depressed by means of a needle introduced into the eye through the cornea. Some divide the crystalline into fragments with the needle, and leave them to the action of the absorbents. The operation is as old as the 17th century.

CERATOPHARYNGE'US, *Ker'atopharynge'us*, (*cerato*, and *φαρυγξ*, 'the pharynx.') The great and small *Cer'atopharynge'i* are small fleshy bundles, forming part of the *Hyopharyngeus* of Winslow.

CERATOPLAS'TICĒ, *Cer'atoplasty*, *Ker'atoplasty*, (*cerato*, and *πλαστικός*, 'forming, formative.') The operation for the formation of an artificial cornea. It has not been practised on man.

CERATORRHEXTIS, *Ruptu'ra cor'neæ*, (from *cerato*, and *ῥήξις*, 'rupture.') Rupture of the cornea.

CERATOSIS, see *Ceratogenesis*.

CERATOSTAPHYL'INUS, *Ker'atostaphyl'inus*, (*cerato*, and *σταφυλή*, 'the uvula.') Some fleshy fibres of the *Thyrostaphylinus* of Winslow.

CERATOTOM'IA, *Ceratectomia*, (*cerato*, and *τομήν*, 'to cut.') Section of the transparent cornea. This incision is used in the operation for cataract, to give exit to pus effused in the eye, in case of hypopyon, &c.

CERATOTOMUS, *Keratot'omus*, *Kerat'omus*. A name given by Wenzel to his knife for dividing the transparent cornea, in the operation for cataract. Many modifications of the instrument have been made since Wenzel's time. See *Knife*, *cataract*.

CERATUM, (*κεράρον*.) Cerate—c. *Adipis*, Cerate—c. *Album*, Ceratum cetacei, Ceratum Galeni—c. de *Althæa*, Unguentum de *Althæa*—c. *Calaminæ*, Cerate of Calamine—c. *Cantharidis*, Cerate of *Cantharides*, Emplastrum *Lyttæ*—c. *Extracti Cantharidis*, see *Emplastrum Lyttæ*—c. de *Cerusæ*, Unguentum plumbi subcarbonatis.

CERATUM CETA'CEI, *O. Spermaceti* i seu *al'bum* seu *ce'ti* seu *sim'plex*, *Unguen'tum Ceta'cei* seu *adipoc'e'ræ cet'o'rum*, *Linimen'tum album*, *Emplas'trum Sperm'atis Ce'ti*, *Spermaceti* i *Ce'rate*, (F.) *Cérat de blanc de baleine*, *Onguent blanc*. (*Spermaceti*, 3j; *White wax*, 3ij; *Olive oil*, 3v. Ph. U. S.) A good emollient to ulcers, &c.

CERATUM CETI, Ceratum cetacei—c. *Cicutæ*, Ceratum conii—c. *Citrinum*, Ceratum resinæ.

CERATUM CONI'i, *Ceratum Cicutæ*. (*Ung. conii*, lbj; *Cetacei*, 3ij; *Ceræ albæ*, 3ij.) A formula in Bartholomew's Hospital; occasionally applied to cancerous and scrofulous sores, &c.

CERATUM EPULOTICUM, Cerate of calamine.

CERATUM GALE'NI, *C. al'bum* seu *refrig'erans Gale'ni*, *Unguen'tum cerat'um* seu *amygdali'num* seu *sim'plex*, *Emplas'trum ad fontic'ulos*, *O'leoce'rat'um a'quâ subac'tum*, *Cold Cream*, (F.) *Cérat blanc* ou de *GALIEN*. (*White wax*, 4 parts; *Oil of sweet almonds*, 16 parts; add, when melted,

Water or Rose-water, 12 parts. Ph. P.) A mild application to chaps, &c.

CERATUM HYDRAR'GYRI COMPOS'ITUM, *Com'pound Ce'rate of Mer'cury*. (*Ung. Hydrarg.*, *Cerat. Sapon. Comp.*, āā 3vj; *Camp'hor*, 3iiss. Rub together. Ph. L.) A discutient to indolent tumours.

CERATUM LABIALE RUBRUM, Cerate for the lips—c. *Lapidis calaminaris*, Cerate of calamine—c. *Lithargyri acetati compositum*, Ceratum plumbi compositum—c. *Lyttæ*, Cerate of *cantharides*—c. *Mercuriale*, Unguentum *hydrargyri*—c. *Picatum*, *Pisselæum*.

CERATUM PLUM'BI COMPOS'ITUM, *C. Lithar'gyri Aceta'ti Compositum* seu *Subaceta'ti plum'bi medica'tum* seu *Plum'bi Subaceta'tis* (Ph. U. S.) seu *Satur'ni*, *Com'pound Lead Ce'rate*, *Goulard's Ce'rate*, *Goulard's Ointment*, (F.) *Cérat de Goulard*, *C. de Plomb composé*. (*Liq. plumb. subacet.*, 3iiss; *Ceræ flavæ*, 3iv; *Ol. oliv.*, 3ix; *Camp'horæ*, 3ss. Ph. U. S.) Its virtues are the same as those of the next.

CERATUM PLUMBI SUBACETATIS, *C. Plumbi compositum*.

CERATUM PLUMBI SUPERACETA'TIS, *C. Plum'bi Aceta'tis*, *Unguen'tum Cerusæ Aceta'tæ*, *Ce'rate of Supera'cetate* or *Sugar of Lead*, *Unguen'tum Aceta'tis Plum'bi*, (F.) *Cérat de suracétate de Plomb*. (*Acetate of lead*, 3ij; *White wax*, 3ij; *Olive oil*, lss.) Cooling and astringent.

CERATUM REFRIGERANS GALE'NI, Ceratum Galeni.

CERATUM RESI'NÆ, *C. Resi'næ fla'væ* seu *ci'tri-num*, *Unguen'tum basil'icon fla'vum* seu *Resi'næ fla'væ* seu *Resino'sum*, *Res'in Ce'rate* or *Ointment*, *Yell'low Basil'icon*, *Basil'icon Ointment*. (*Resin*, flav., 3x; *Ceræ flav.*, 3iv; *Adipis*, 3xvj. Ph. U. S.) A stimulating application to old ulcers, &c. Digestive.

DR. SNEELOME'S Ointment for the Eyes consists of finely powdered *verdigris*, 3ss, rubbed with oil, and then mixed with an ounce of ceratum resinæ.

CERATUM RESI'NÆ COMPOS'ITUM, *Com'pound Res'in Ce'rate*, *Deshler's Salve*, (*Resin.*, *Savi*, *Ceræ flavæ*, āā lbj; *Terebinth.*, lss; *Ol. Lini*, 3vij. Melt together, strain through linen, and stir till cool. Ph. U. S.)

CERATUM SABI'NÆ, *Unguen'tum Sabi'næ*, *Sav'ine Ce'rate*, *Savine Ointment*, (F.) *Cérat de Sabine*. (*Savine*, in powder, 3ij; *Resin Cerate*, lbj. Moisten the savine with ether; and, in a percolator, pour on ether until the filtered liquid passes nearly colourless; evaporate to the consistence of syrup, and mix with the melted resin cerate. Ph. U. S.) Irritative, 'drawing.' Used in the same cases as the cerate of *cantharides*.

CERATUM SAPO'NIS, *C. Sapo'nis Compositum*, *Soap Ce'rate*, (F.) *Cérat de Savon*. (*Emp. Sapon.*, 3ij; *Ceræ alb.*, 3iiss; *Ol. Oliv.*, 3iv. Melt together the plaster and wax, add the oil, and mix. Ph. U. S., 1863.) It is applied in cases of sprains or fractures.

CERATUM SATURNI, Ceratum plumbi compositum—c. *Simplex*, Cerate, Ceratum cetacei—c. *Spermaceti*, Ceratum cetacei—c. *Subaceta'ti plumbi medicatum*, Ceratum plumbi compositum—c. *Tetrapharmacum*, *Pisselæum*—c. *Viride*, Unguentum subaceta'tis cupri.

CERATUM ZIN'CI CARBONA'TIS, *Ce'rate of Car'bonate of Zinc*. (*Zinci carbonat. præparat.*, 3ij; *Ung. simpl.*, 3x. Ph. U. S.) Used in the same cases as the Ceratum calaminæ.

CERAUN'ION, (from *κεραυνος*, 'thunder, thunderbolt.') *Lapis fulmin'ens*. A kind of stone, which was believed to be formed during thunder; and to be possessed of the power of inducing sleep, and of numerous prophylactic virtues. It was rubbed on the knee, breast, &c., in swelling of those parts.

CERBERA AHOVAI, (from the fabulous *Kep-βeros, Cerberus*.) See Strychnine.

CERBERA TAN'GHIN, *Tanghin'ia, T. venenifera*. A tree of Madagascar, Order, Apocynaceæ; the kernel of whose fruit is a deadly paralyzing poison; and although not larger than an almond is sufficient to destroy twenty people.

CERBERA THEVETIA, see Strychnine.

CERBERUS TRICEPS, *Pulvis cornachini*.

CERCA'RIA, (*κερκος, 'a tail.'*) A genus of agastric, caudate, infusory animalcules, one of the most curious of which inhabits the tartar of the teeth. The, so-called, spermatozoa have been presumed by some to belong to this genus.

CERCHNASMUS, (*κερχνασμος*.) Cerchnus.

CERCHNOMA, (*κερχνωμα*.) Cerchnus.

CERCH'NUS, *Cerchnus'mus, Cerch'nium, Cerch-no'ma, (κερχνος, from κερχειν, 'I render hoarse.')* A rough voice produced by hoarseness. See Rattle.

CERCIF'IS, *Tragopogon porrifolium*.

CER'CIS, (*κερκις*.) A sort of pestle for reducing substances to powder. Also, the radius or small bone of the arm. See *Pilum*, and *Radius*.

CERCLE, *Circulus—c. de la Choroidé, Ciliary ligament—c. Ciliare, Ciliary ligament*.

CERCO'SIS, (*κερκωσις, from κερκος, 'a tail,' and osis*.) *Men'tula muli'ebria, Clit'oria*. Some authors have employed the word synonymously with nymphomania and elongation of the clitoris; and with *Polypus Uteri*, the *Sarco'ma Cerco'sis* of Sauvages, *Caud'a pudendi* of Paulus Ægineta.

CERCOSIS CLITORIDIS, Clitorism—*c. Externa, Clitorism*.

CEREA, (from *cera*.) Cerumen.

CEREA'LIA, (*CERES, 'goddess of corn.'*) (F.) *Céréales (Plantes)*. The cerealia are gramineous plants, the seed of which serve for the nourishment of man:—as wheat, barley, rye, &c. At times, the same term is applied to some of the leguminous plants.

CEREBEL'LA URINA. Urine of a whitish appearance, of the colour of the brain or cerebellum, from which Paracelsus thought he could distinguish diseases of those organs.

CEREBELI'TIS, (*cerebellum and itis*.) *Par-encephali'tis, Inflamma'tio cerebel'li*. Inflammation of the cerebellum: a variety of phrenitis or encephalitis.

CEREBEL'LOUS, *Cerebello'us, (from cerebel'um)*. (F.) *Cérébelleux*. Chaussier has given this epithet to the vessels of the cerebellum. These are three in number; two of which are inferior: the larger, *inferior cerebel'li*, which arises from the posterior cerebral or vertebral; and the smaller, whose existence is not constant, from the meso-cephalic or basiliary:—the third, called *superior cerebel'li*, (F.) *A. cérébelleuse supérieure*, is also a branch of the basiliary.

CEREBEL'LOUS AP'OPLEXY, *Apoplex'ia cerebel'lo'sa*: apoplexy of the cerebellum.

CEREBEL'LUM, (*dim. of Cerebrum*.) *Cerebrum par'uum seu poste'rius, Appen'dix ad cer'ebrum, Encra'nion, Encra'nis, Epenera'nis, Parenceph'a-lis, Parenceph'alus, Parenceph'alum, Enceph'alium, Enceph'alus opis'thius, Mierenceph'alium, Mierenceph'alum, Cer'ebel, Little brain*. (F.) *Cervelet*. A portion of the medullary mass, contained in the cavity of the cranium. It fills the lower occipital fossæ below the tentorium, and embraces the tuber annulare and medulla. It is composed, like the brain, of vesicular and tubular substance, arranged in laminae, as it were; so that, when a section is made of it, it has an arborescent appearance, called *Ar'bor vi'tæ*. The cerebellum is divided into two lobes or hemispheres or lateral masses, *Lobus superior anterior seu quadrangula'ris*, and *Lobus superior posterior seu semiluna'ris*, and each lobe is again subdivided into *Monticuli* or *Lob'ules*. In the cerebellum are to be observed

the *crura cerebelli*, the *fourth ventricle*, the *valvula magna cerebri*, the *processus vermiculares*, superior and inferior, &c. Its precise functions are not determined. It has been regarded as, but not proved to be, the organ of the instinct of reproduction; and by others as the co-ordinator and regulator of the movements.

CEREBRAL, *Cerebra'tis*, (F.) *Cérébral*, (from *cerebrum*.) Belonging or relating to the brain; similar to brain.

CEREBRAL APOPHYSIS, Pineal gland.

CEREBRAL ARTERIES are three on each side—the *anterior or artery of the cor'pus callo'sum*, and the *middle, arte'ria Sylvia'na*, are furnished by the internal carotid:—the *posterior or poste'rior and infe'rior artery of the brain, A. profun'da cer'ebr'i*, arises from the vertebral. Chaussier calls these arteries *lobaires*, because they correspond with the anterior, middle, and posterior lobes, whilst he calls the trunks, whence they originate, *cérébral*.

CEREBRAL NERVES are those which arise within the cranium, (F.) *Ner'vi craniens*, all of which, perhaps, not excepting the olfactory, originate from the medulla oblongata. See *Nerves*.

In *Pathology*, an affection is called *cerebral*, which specially occupies the brain. *Fièvre cérébrale, Cerebral fever*, is a variety in which the head is much affected.

CEREBRA'TION, *Cerebra'tio*, (from *cerebrum*.) The action of the brain during any mental or moral manifestation.

CÉRÉBRIE, Insanity.

CEREBRIFORM, (*cerebrum*, and *forma*, 'form,') Encephaloid.

CEREBRITIS, (*cerebrum*, and *itis*.) Phrenitis.

CEREBROMALACIA, (*cerebrum*, and *μαλακία*, 'softness,') Mollities cerebri.

CEREBROPATHY, (*cerebrum*, and *παθος*, 'affection,') see *Nervous diathesis*.

CEREBRORACHIDIAN, (*cerebrum*, and *ραχίς*, gen. *ραχιδος*, 'spine,') Cephalospinal.

CEREBROSPINAL, Cephalospinal. A *cer'ebro-sp'inal* or *cer'ebro-sp'inant* is a neurotic, which exercises a special influence over one or more functions of the brain and spinal cord, and their respective nerves.

CEREBRO-SPINAL AXIS, see *Encephalon*.

CEREBRO-SPINANT, *Cerebro-spinal*.

CEREBRUM or CEREBRUM, (*κρα, 'the head.'*) The brain. (F.) *Cerveau, Cerveille*. This term is sometimes applied to the whole of the contents of the cranium: at others, to the upper portion; the posterior and inferior being called cerebellum. The brain, properly so called, extends from the os frontis to the superior occipital fossæ. Anteriorly, it rests on the orbital vault: behind this, on the middle fossæ of the base of the cranium; and, posteriorly, on the tentorium cerebello superextensum. The upper surface is divided by a deep median cleft (*Scissure interlobaire*, —Ch.) into two halves, called *hemispheres*, which are united at the base by the *corpus callosum*. At its surface are numerous convolutions. The inferior surface exhibits, from before to behind, three lobes, distinguished into *anterior, middle, and posterior*. The middle is separated from the anterior by the *fissure of SYLVIVS*; and from the posterior, by a shallow furrow which corresponds to the upper portion of the *pars petrosa*. Internally, the brain has, on the median line, the *corpus callosum*, *septum lucidum*, *forix*, *pineal gland*, and *third ventricle*:—and, laterally, the *lateral ventricles*, in which are the *corpora striata*, *optic thalami*, &c. It is contained in a triple envelope, (see *Meninges*.) Its texture is pulpy, and varies according to age. Two substances may be distinguished in it—the *white, med'ullary, tubular* or *fib'rous—medull'a cer'ebr'i*, (F.) *Pulpe cérébral*,

Substance blanche, and the *cortical*, *cineritious*, *vesic' alar*, or *gray*, (F.) *Substance grise*. The former is white; and occupies all the interior and base of the brain. The latter is grayish and softer. It is situate particularly at the surface of the organ.

The brain receives several arterial vessels, furnished by the internal carotid and vertebral. Its veins end in the sinuses. It is the material organ of the mental and moral manifestations. According to Gall, each part is the special seat of one of those faculties, and the brain and cerebellum, inclusive, are called by him, '*the nervous system of the mental faculties*.' See *Craniology*.

The substance of the nervous system—*Neurine*—was analyzed by Vauquelin, and found to contain water, 80·00; white fatty matter, 4·53; red fatty matter, called *cerebrine*, 0·70; osmazome, 1·12; albumen, 7·00; phosphorus, 1·50; sulphur, acid phosphates of potassa, lime, and magnesia, 5·15.

CEREBRUM ABDOMINALE, Solar plexus—c. Elongatum, Medulla oblongata—c. Parvum, Cerebellum—c. Posterius, Cerebellum.

CERECTOMY, Ceratectomy.

CEREFOLIUM, (corruption of *Charophyllum*), *Scandix cerefolium*—c. Hispanicum, *Charophyllum odoratum*—c. Sylvestre, *Charophyllum sylvestre*.

CERELÆUM, (*κηρος*, 'wax,' and *ελαιον*, 'oil.') Cerate.

CÉRÉOLÉ, Cerate.

CEREOLUS, (*cera*, 'wax,') *Bougie*.

CERERISIA, *Cerevisia*.

CERERIUM, (from the planet *Ceres*), *Cerium*.

CERUM MEDICATUM, ('a medicated taper,') *Bougie*.

CEREUS, (*cera*, 'wax,') *Bougie*.

CEREVIS'IA, quasi *Ceremis'ia*, *Cervis'ia*, *Ce'lia*, *Zy'thus*, *Zy'thum*, *Li'quor Cer'eris*, *Oinos κρηθς*, *Vinum hordeaceum*, *Bi'ra*, *Bry'ton*, *Bar'ley wine*, (from *Ceres*, the goddess of oorn, 'corn;' whence it is made.) *Ale*, *Alla*, (Sc.) *Bar'leybree*, *Bar'ley-corn*, *Beer*, *Port'er*, (F.) *Bière*, *Cerroise*. These fluids are drunk by the inhabitants of many countries habitually, and in Great Britain and Germany more than in others. They are nourishing, but not very easy of digestion. A German beer, under the name *Lager beer*, (G.) *Lager Bier*, ('stock beer,') is much used all over the United States.

The old dispensaries contain numerous medicated ales, which are no longer in use. Some, however, are still admitted. They are the *Brutolés* on *Brytolés* of Chêreau, *Brytolatures* of Bérail;—*Bières Médicinales*.

CEREVISIA ABIETIS, Beer, spruce—c. Nigra, see *Falltranck*.

CEREVISIE FERMENTUM, *Yest*.

CERFEUIL, (corruption of *Charophyllum*), *Scandix cerefolium*—c. *Musqué*, *Charophyllum odoratum*—c. *Sauvage*, *Charophyllum sylvestre*.

CERION, (*κηρυον*, 'honeycomb,') see *Favosus*, *Porrigio favosa*.

CERISIER, *Prunus cerasus*—c. à *Grappes*, *Prunus padus*—c. de *Virginie*, *Prunus Virginiana*.

CERIUM, *Cere'rium*, (after the planet *Ceres*.) A metal which is never found isolated, being usually combined with lanthanum or lantanum and didymium. The salts of cerium are supposed to be sedative and tonic. They have been used in dyspepsia in its various forms; the vomiting of pregnancy, &c. Dose of the oxalate or nitrate, one or two grains.

CRENIN, SAINT, MIN'ERAL WA'TERS OF. St. C, is a parish in the diocese of St. Flour, Upper Auvergne, France. The water is a chalybeate. It is called *Eau du Cambon*.

CEROËNË, *Cerou'nê*, or *Cirouë'nê*, *Ceroë'nun*,

(*cerus*, and *ovos*, 'wine,') A plaster composed of *yellow wax*, *mutton suet*, *pitch*, *Burgundy pitch*, *bole Armeniac*, *thus* and *wine*. It was used as a strengthening plaster. Sometimes it contained neither wax nor wine.

CEROLEUM, (*cera*, and *oleum*, 'oil,') *Cerate*.

CEROMA, gen. *Cero'matis*, (*κηρος*, 'wax,') *Cerate*. See *Lardaceous*.

CEROMANT'IA, (*cerus*, and *μαντ'ια*, 'divination,') The art of foretelling the future, from the figures which melted wax assumes, when suffered to drop on the surface of water.

CERONIUM, *Cerate*.

CEROPIS'SUS, (*cerus*, and *πισσα*, 'pitch,') A depilatory plaster, composed of pitch and wax.

CEROSTROSIS, (*κερας*, 'horn,' and *στρωσις*, 'spreading,') *Hystriasis*.

CEROTE, (*κηρωτη*,) *Cerate*.

CEROTUM, (*κηρωτον*,) *Cerate*.

CEROXYLON, (*κηρος*, and *ξυλον*, 'wood,') see *Cera flava* et *alba*.

CERUA, *Ricinus communis*.

CERU'MEN, gen. *Ceru'minis*, (from *cera*, 'wax,') *C. Au'rium*, *Ce'rea*, *Au'rium Sor'des*, *Sordic'ule au'rium*, *Marmora'ta Au'rium*, *Cyp'selæ*, *Ceru'minous Hu'mour*, *Ear-wax*, *Cyp'selis*, *Fu'gile*, (F.) *Cire des Oreilles*. A name given to the unctuous humour, similar to wax in its physical properties, which is met with in the *meatus auditorius externus*. It is secreted by glands, situate beneath the skin lining the meatus. It lubricates the meatus, preserves the suppleness of the lining membrane, prevents the introduction of bodies floating in the atmosphere, and by its bitterness and unctuousness prevents insects from penetrating.

CERU'MINOUS, *Cerumino'sus*, (F.) *Cérumineux*. Relating to cerumen.

CERUMINOUS GLANDS, *Ceru'minous Follicles*, *Gland'ulæ Cerumino'sæ*, (F.) *Glandes ou Follicles cérumineuses*. Glands or follicles which secrete the cerumen.

CERUS, (*κηρος*,) *Cera*.

CERUSE, *Plumbi subcarbonas*.

CERUSSA, *Plumbi subcarbonas*—c. *Acetata*, *Plumbi superacetas*—c. *Alba Hispanica*, *Plumbi subcarbonas*—c. *Alba Norica*, *Plumbi subcarbonas*—c. *Nigra*, *Graphites*—c. *Psymmithron*, *Plumbi subcarbonas*—c. *Serpentaria*, see *Arum maculatum*.

CERUSSE, *Plumbi subcarbonas*.

CERUS'SEA URINA. A term used by *Paracelsus* for the urine, when of a colour like cerusse.

CERVARIA ALBA, *Laserpitium latifolium*.

CERVEAU, *Cerebrum*—c. *Substance blanche du*, see *Cerebrum*—c. *Substance grise du*, see *Cerebrum*.

CERVELET, *Cerebellum*.

CERVELLE, *Cerebrum*.

CERVI BOLETUS, *Elaphomyces Granulatus*.

CERVI ELAPHI CORNU, *Cornu cervi*, see *Cervus*.

CERVICAL, *Cervica'lis*, *Trache'lian*, (F.) *Trachélien*, (*cervix*, gen. *cervicis*, 'neck,' 'the back of the neck,') Every thing which concerns the neck. *Cervi'al*, in antiquity, meant a pillow or bolster.

CERVICAL AR'TERIES are three in number: 1. The *ascend'ing*, *super'ior*, or *superfic'ial*, a branch of the inferior thyroid, distributed to the scaleni muscles and integuments. 2. The *trans'verse* (*Cervico-scapulaire*—Ch.), a branch of the axillary artery, or of the subclavian: distributed to the *levator scapulae trapezius*, &c. 3. The *posterior* or *profound*, *A. transversa'lis col'li*, *Trachélo-cervicale*—(Ch.) a branch of the subclavian, distributed to the deep-seated muscles on the anterior and posterior parts of the neck. See, also, *Prinæps Cervicis (arteria)*.

CERVICAL GANGLIONS. The three ganglions of the great sympathetic. The *cervical glands* or lymphatic glands of the neck are, also, so called. See Trisplanchnic nerve.

CERVICAL LIG'AMENTS. These are two in number. 1. The *anterior*, extending from the basilary process of the occipital bone to the anterior part of the first cervical vertebra. 2. The *posterior* or *supraspinous*, *Ligamentum Nuchæ*, which extends from the outer occipital protuberance to the spinous process of the seventh cervical vertebra. In animals with large heads it is very strong. See Nucha.

CERVICAL NERVES are eight in number on each side, and form the *eight cervical pairs*, which are the first given off from the spinal marrow.

CERVICAL PLEX'US, *Plexus Trachælo-scutané* (Ch.) The nervous net-work formed by the anterior branches of the first three cervical nerves, above the posterior scalenus muscle, and at the outer side of the pneumogastric nerve, carotid artery, and jugular vein.

CERVICAL RE'GION, ANTE'RIOR DEEP, Prever'tebral re'gion. The region of the neck, occupied by three pairs of muscles placed immediately in front of the cervical and three superior dorsal vertebrae:—viz., the rectus capitis anticus major, the rectus capitis anticus minor, and longus colli;—hence termed *prevertebral muscles*.

CERVICAL TRI'ANGLES. The side of the neck is divided into two large triangles by the sternomastoid muscle—the *anterior triangle* or *triangular space*, and the *posterior triangle* or *triangular space*, which are interesting in their surgical relations. 1. The anterior triangular space is divided into three smaller triangles by the digastric muscle above, and by the anterior belly of the omo-hyoid below. These are named, from below upwards, the *inferior carotid triangle*, the *superior carotid*, and the *submaxillary*. 2. The posterior triangle or triangular space is crossed by the posterior belly of the omo-hyoid, which divides it into two triangles,—an *upper* or *occipital*, and a *lower* or *subclavian*.

CERVICAL VEINS have nearly the same distribution as the arteries.

CERVICAL VERTEBRÆ. The first seven vertebrae of the spine. See Vertebrae.

CERVICALIS DESCENDENS, see Hypoglossus, and Sacro-lumbalis.

CERVICARIA, Campanula trachelium.

CERVICI-DORSO-COSTAL, Serratus posticus superior—c. Dorsomastoidien et dorso-trachælien, Splenius—c. Dorsoscapulaire, Rhomboideus—c. Mastoïdien, Splenius.

CERVICOFA'CIAL, *Cervicofacialis*. Belonging to the neck and face.

CERVICOFACIAL NERVE, *Nervus cervicofacialis*. A branch of the facial nerve, distributed to the neck and face.

CERVISIA, Cerevisia.

CERVISPINA, ('buckthorn,') Rhamnus.

CERVIX, gen. *Cervicis*, *Collum*. The neck, especially the back part. A neck.

CERVIX OBSTIPA, Torticollis—c. Uteri. Collum uteri.

CERVOISE, Cerevisia.

CERV'US, *Cervus*, ('the stag,') The horn of *Cervus Elaphus*, called *Cor'nu*, *Cervi Elaphi Cor'nu*, *Cor'nu Cervinum*, *Harts'horn*, (F.) *Corne de cerf*, contains 27 parts of gelatin in the 100. A jelly made from the shavings is emollient and nutritive.

The *Stag's Piz'le*, *Pria'pus Cervi*, was once considered to be aphrodisiac. Dose, ʒj to ʒj, in powder.

CERVUS AL'CES. The *Elk*, (F.) *Élan*. The hoof of this animal was anciently used as an antiepileptic. The animal, it was asserted, was subject to attacks of epilepsy, and always cured them by

putting its hoof into the ear. The hoof was also worn as an amulet.

Cor'nu U'tum, *Burnt Harts'horn*, has been used as an antacid, but it consists of 57 parts of phosphate, and only one of carbonate of lime. It is, therefore, not of much use.

CÉSARIENNE (OPÉRATION), Cæsarean section.

CESSATIO MENSIIUM, Amenorrhœa.

CESTRI'TES, (*κεστριον*, 'betony,') A wine prepared from betony.

CESTRON, *Betonica officinalis*.

CETA'CEUM, (*κητος*, *cetus*, 'a whale,') *Al'bum* seu *Se'cum Ceti*, *Adipoc'e ra ceto'sa*, *Stearinum ceta'ceum*, *Spermacet'i*, *Parmacit'y*, *Sperm.* (F.) *Blanc de Baleine*, *Cétine*, *Adipocire de Baleine*. An odorous, insipid, white, crystallized, friable, unctuous substance, obtained from *Physe'ter Macroceph'alus* or *Spermacet'i Whale*, (F.) *Cachalot*, and other varieties of whale. S. g. 9433: melts at 112°. It is demulcent and emollient, and has been given in coughs and dysentery, but is mostly used in ointments. Dose, ʒss to ʒiiss, rubbed up with sugar or egg.

CETERACH OFFICINARUM, *Asplenium ceterach*.

CÉTINE, *Cetaceum*.

CETRARIA ISLANDICA, *Lichen islandicus*.

CETRARIUM, see *Lichen islandicus*.

CETRARIUM, see *Lichen islandicus*.

CEVADILLA, see *Veratrum sabadilla*—c. *Hisp-anorum*, *Veratrum sabadilla*.

CÉVADILLE, *Veratrum sabadilla*.

CHAA, *Thea*.

CHACRILLE, *Croton cascarilla*.

CHADLOCK, *Sinapis arvensis*.

CHÆREFOLIUM, *Scandix cerefolium*.

CHÆROMANIA, (*χαίρω*, 'to be pleased,') *Amenomania*.

CHÆROPHYL'LUM, (*χαίρεφυλλον*, from *χαίρω*, 'I rejoice,' and *φυλλον*, 'a leaf,' owing to its luxuriant foliage.) *Scandix cerefolium*.

CHÆROPHYLLUM ANGULATUM, *Ch. sylvestre*—ch. *Cerefolium*, *Scandix cerefolium*—ch. *Monogynum*, *Ch. sylvestre*.

CHÆROPHYLLUM ODORA'TUM, *Scandix* seu *Cicutaria Odorata*, *Myrrhis Odorata* seu *Ma'jor*, *Cerefo'lium Hispanicum*, *Sweet Cic'ely*, (F.) *Cerfeuil musqué* ou d'Espagne; *Order*, *Umbelliferae*; has the smell of aniseed, and is cultivated on account of its aromatic properties.

CHÆROPHYLLUM SATIVUM, *Scandix*.

CHÆROPHYLLUM SYLVE'STRE, *Cicutaria*, *Chærophyl'lum tem'ulum* seu *monog'ynum* seu *angulatum* seu *verticellatum*, *Anthriscus humilis* seu *proc'erus*, *Cerefo'lium Sylvest're*, *Bas'tard Hemlock*, *Wild Cher'vil* or *Cow-weed*, (F.) *Cerfeuil sauvage*, *Persil d'Ane*, is a slightly fetid aromatic, but is not used.

CHÆROPHYLLUM TEMULUM, *Chærophyl'lum sylvestre*—c. *Verticillatum*, *Ch. sylvestre*.

CHÆTE, (*χαῖτη*,) *Capillus*.

CHAFFBONE, *Jawbone*.

CHA'FING, *Erythe'ma Intertri'go*, *Intertri'go*, *Paratri'm'ma*, *Paratri'psis*, *Diatri'm'ma*, *Attri'ta*, *Attri'tio*, *Attri'tus*, (from (F.) *eschauffer*, *échauffer*, 'to heat,') *Fret*, *Gall of the skin*; (F.) *Échauffement*, *Écorechure*. The red excoriations which occur in consequence of the friction of parts, or between the folds of the skin, especially in fat or neglected children. Washing with cold water and dusting with hair-powder is the best preventive. When occurring between the nates and in the region of the perineum, from long walking,—*Intertri'go pod'icis*, *Proctal'gia intertrigin'osa*,—it is vulgarly designated by the French *Entrefesson*.

CHAIR, (from [L.] *caro*,) *Flesh*—c. *de Poule*, *Horrida cutis*.

CHAIR, OBSTETRIC, *Labour chair*

CHAIR, WHIRLING. A kind of chair in which the insane are placed, and which is made to revolve with great rapidity, as a tamer in furious mania.

CHAIROMANIA, (*Cheromania*), Amenomania.

CHAISE ('chair') *PERCÉE*, Lasanum.

CHALASIS, (*χαλασις*, from *χαλασ*, 'I relax,') Relaxation.

CHALASMUS, (*χαλασμος*), Relaxation.

CHALASODERMIA, Chalastodermia.

CHALAS'TICUS, (*χαλαστικός*.) A medicine proper for removing rigidity of the fibres. An emollient or relaxant.

CHALASTODERM'IA, *Chalastoderm'ia*, *Dermatol'ysis*, (*χαλαστος*, 'relaxed,' and *δερμα*, 'skin,') Abnormal extension and relaxation of the skin.

CHALAZA, *Chalaz'ion*, *Chalaz'ium*, *Chalazo'sis*, (from *χαλαξα*, 'hail,') *Poro'sis*, *Grand'o*, *To'phus*, *Hail*, (F.) *Grêle*, *Gravelle*. A hard, round, transparent tumour, developed in different parts of the body, more especially in the substance of the eyelids.—Also, the Cicatricula of the egg. Generally, however, in the language of ovologists, the *chalazæ* or *poles* are, in the egg of the bird, the more dense internal layer of the albumen, which adheres to the yolk, and is continued, in the form of two spirally twisted bands, towards the extremities of the egg. The twisting is considered to be produced by the revolving motion of the egg in its descent through the oviduct.

CHALAZÆ, see Chalaza.

CHALAZION, Chalaza.

CHALAZONEPHRITIS, Kidney, Bright's disease of the.

CHALAZOSIS, Chalaza.

CHALBANE, (*χαλβανη*), see Bubon galbanum.

CHALCANTHUM, (*χαλκανθον*), Atramentum, Ferri sulphas.—c. Album, Zinc sulphas.

CHALCEDONIUS, (from *Chalcedon*, in Asia,) Cornelian.

CHALCITAS, Bismuth.

CHALCITES, (*χαλκίτης*), Coleothar.

CHALCOIDEUM (OS), Cuneiform bone.

CHALCUS, (*χαλκος*), Æreolum, Brass, Cuprum.

CHALEUR, Heat.—c. *Âcre*, see Acrid.—c. *Animale*, Animal heat.—c. *Ervatique*, see Heat.—c. *Extérieure*, see Heat.—c. *Halitense*, see Heat.—c. *Interne*, see Heat.—c. *Mordicante*, see Heat.—c. *Nerveuse*, see Heat.—c. *Sec*, see Heat.—c. *Septique*, see Heat.

CHALEURS DES ANIMAUX, see Heat.

CHALINI, (*χαλινι*), see Lip.

CHALINOPLASTY, *Chalinoplast'icē*; (*χαλινος*, 'frænum,' 'a bridle,' and *πλασσω*, 'I form,') The operation for forming a new frænum.

CHALK, (*calx*, Sax. *ceale*), Creta.—c. Red, *Rubrica febrilis*—c. Stones, Calculi, arthritic.

CHALL, Maxillary bone.

CHALYBE, Chalybeate.

CHALYB'EATE, *Chalybeatus*, *Ferrugin'eus*, *Ferrugin'us*, *Ferratus*, *Martial'is*, *Ferrugin'ous*, *Martial*, *Ferrated*, (F.) *Ferrugineux*, *Chalybé*, *Ferrique*, (from *chalybs*, 'iron or steel,') Of, or belonging to iron; containing iron. Any medicine into which iron enters, as *chalybeate mixtures*, pills, waters, &c. See Waters, Mineral.

CHALYBIS RUBIGO, ('rust of steel,') Ferri subcarbonas.

CHALYBOKRENÆ, (*chalybs*, and *κρηνη*, 'a spring,') Waters, mineral, (chalybeate.)

CHALYBOPEGÆ, (*chalybs*, and *πηγη*, 'a spring,') Waters, mineral, (chalybeate.)

CHALYBS, gen. *chalybis*, (*χαλυσ*, from *Χαλυβς*, *Chalybes*, a people of Pontus, who dug iron out of the earth; *Αΐεις*, *Steel*. The *Proto-carburet of iron*, (F.) *Acier*. As a medicine, steel does not differ from iron.

CHALYBS TARTARIZATUS, Ferrum tartarizatum.

CHAMA, Chema.

CHAMÆ, (*χαμαί*), in composition, 'on the ground.'

CHAMÆACTE, (*χαμαιακτη*, from *χαμαί*, and *ακτη*, 'the elder tree,') *Sambucus ebulus*.

CHAMÆBATOS, (*χαμαιβατος*, from *χαμαί*, and *βατος*, 'a bramble,') *Fragaria*.

CHAMÆCEDRUS, (*χαμαί*, and *κεδρος*, 'cedar,') *Artemisia santonica*.

CHAMÆCISSUS, (*χαμαικισσος*, from *χαμαί*, and *κισσος*, 'ivy,') *Glechoma hederacea*.

CHAMÆCLEMA, (*χαμαί*, and *κλημα*, 'a shoot,') *Glechoma hederacea*—c. *Hederacea*, *Glechoma hederacea*.

CHAMÆCYPARISSUS, (*χαμαί*, and *κυπαρισσος*, 'cypress,') *Artemisia santonica*.

CHAMÆDROPS, Teucrium *chamædrys*.

CHAMÆDRIY'TES. A wine, in which the *Teu'crium Chamæ'drys* has been infused.

CHAMÆDRYS, (*χαμαιδρυς*, from *χαμαί*, and *δρυς*, 'the oak,') *Rubus chamæmorus*, Teucrium Ch., *Veronica*—c. *Incana maritima*, Teucrium marum—c. *Marum*, Teucrium marum—c. *Minor repens*, Teucrium Ch.—c. *Palustris*, Teucrium scordium—c. *Scordium*, Teucrium scordium—c. *Vulgaris*, Teucrium Ch.

CHAMÆGEIRON, Tussilago.

CHAMÆLEAGNUS, *Myrica gale*.

CHAMÆLAT'ITES. A wine impregnated with *Chamælaia*, (*Daph'nē Alpi'na*.)

CHAMÆLÆA, (*χαμελαία*, from *χαμαί*, and *ελαία*, 'the olive tree,') *Daphne alpina*.

CHAMÆLEON ALBUM, (*χαμαί*, and *λεων*, 'lion,') *Carolina acaulis*.

CHAMÆLEUCE, (*χαμαιλευκη*, from *χαμαί*, and *λευκος*, 'white,') Tussilago.

CHAMÆLINUM, (*χαμαί*, and *λινον*, 'flax,') *Linum catharticum*.

CHAMÆLIR'IUM LU'TEUM, (*χαμαί*, and *λειριον*, 'a lily,') (?) *Vera'trum lu'teum*, *Hel'o'nias lu'tea* seu *Dio's'ca*, *Dev'il's Bit*, *Blaz'ing star*, *U'nicorn root*; indigenous; Order, *Melanthaceæ*; flowering in June; is acrid. An infusion of the root has been given as an anthelmintic; a tincture, as a tonic.

CHAMÆMELUM, (*χαμαιμηλον*, from *χαμαί*, and *μηλον*, 'apple,') *Anthemis nobilis*—c. *Fœtidum*, *Anthemis cotula*—c. *Nobile*, *Anthemis nobilis*—c. *Odoratum*, *Anthemis nobilis*—c. *Vulgare*, *Matricaria chamomilla*.

CHAMÆMORUS, (*χαμαί*, and *μορεα*, 'the mulberry tree,') Teucrium *chamæpitys*, *Rubus chamæmorus*—c. *Norwegica*, *Rubus chamæmorus*.

CHAMÆPEUCE, (*χαμαιπευκη*, from *χαμαί*, and *πευκη*, 'pine,') *Camphorosma Monspeliaca*.

CHAMÆPITU'NUM VINUM. A wine, in which the leaves of the *Chamæ'pitys*, *Teu'crium Chamæ'pitys*, have been infused.

CHAMÆPITYS, (*χαμαιπitys*, from *χαμαί*, and *πitys*, 'the pine,') Teucrium *chamæpitys*—c. *Anthyllum*, Teucrium *iva*—c. *Moschata*, Teucrium *iva*.

CHAMÆPLION, (*χαμαιπλιον*), *Erysimum*.

CHAMERAPH'ANUS, (*χαμαιραφανος*, from *χαμαί*, and *ραφανος*, 'the radish,') The upper part of the root of the *Asium*.

CHAMÆROPS SERRAT'ULA, *Saw Pal met'to*, (*χαμαί*, and *ρωψ*, 'a shrub,') A farina is prepared from the roots of this plant, which is used by the Indians, in Florida, as a diet.

CHAMBAR, Magnesia.

CHAMBER, (*χαμαρα*), *Cam'era*, (F.) *Chambre*. A term used in speaking of the eye, in which there are two chambers, *Cam'ere oc'uli*:—an *anterior* and a *posterior*; (F.) *Chambre antérieure* et *Ch. postérieure*. The anterior is the space between the cornea and the anterior part of the iris:—the *posterior*, the space between the iris and anterior surface of the crystalline. They are filled with the aqueous humour, and communicate by the opening in the pupil.

CHAMBERLIE, Urine.

CHAMBERLAIN'S RESTORATIVE PILLS. This nostrum, recommended in scrofula, and all impurities of the blood, has been analyzed by Dr. Paris, and found to consist of *cinnabar, sulphur, sulphate of lime*, and a little vegetable matter. Each pill weighs 3 grains.

CHAMBRE, Chamber.

CHAMBRIE, Cannabis sativa.

CHAMOMILE, (same etymon as Chamæmulum.) *Anthemis nobilis*—c. Dog's, *Anthemis cotula*, *Matricaria chamomilla*—c. Dyers', *Anthemis tinctoria*—c. German, *Matricaria chamomilla*—c. Spanish, *Anthemis pyrethrum*—c. Stinking, *Anthemis cotula*—c. Wild, *Anthemis cotula*, *Matricaria glabrata*.

CHAMOMILLA PÆTIDA, *Anthemis cotula*—c. Nostras, *Matricaria chamomilla*—c. Romana, *Anthemis nobilis*—c. Spuria, *Anthemis cotula*.

CHAMPACA, *Michelia champaca*.

CHAMPIGNON, (from *champ*, [L.] *campus*, 'a field,') *Fungus*—c. de l'Appareil des Fractures, *Clavaria*—c. de Couche, see *Agaric*—c. du Ferment, *Torula cerevisiæ*—c. de Malte, *Cynomorion coccineum*—c. du Muguet, *Aphthaphyte*—c. du Poupon, see *Fungus*.

CHAMPOOING, Shampooing.

CHANCEBONE, Ischion.

CHANCELAGUA, Canchalagua.

CHANCER, (from *cancer*,) *Shank'er*, *Ulcus seu Ulcusculum cancrosum*. A sore, which arises from the direct application of the venereal virus; hence it is almost always seated, in men, on the penis. The French use the word *Chancre*, in popular language, for cancerous ulcers, the malignant aphthæ of children, &c. Formerly, the terms *Caroli* and *Ca'ries pudendum* were used for venereal pustules or sores on the parts of generation.

CHANCER LARVÉ. 'A concealed or masked chancre,'—one, in other words, situate in the urethra, or vagina, or os uteri—such as has been supposed by M. Ricord to give occasion to gonorrhœa virulenta.

CHANCER, PHAGEDENIC, see *Phagedenic*—*Ch. de Sahara*, see *Bouton d'Allep*.

CHANCREUX, *Chancreux* (F.), *Cancro'sus*, *Carcino'des*. Having the nature of chancre, or of cancer.

Bouton Chancreux. A small tumour of a cancerous nature, which makes its appearance on the face—most frequently on the upper lip—*Noli me tangere* [?].

CHAN'CROID, (F.) *Chaneroïde*: (*chancre*, and *idos*, 'resemblance,') Resembling infectious chancre.

CHANDOO. An aqueous extract of opium used by the Chinese for smoking.

CHANGE OF LIFE, see *Menses*.

CHANT ([L.] *cantus*, 'a song') **DES ARTÈRES**, ('song of the arteries,') *Sifflement modulé*.

CHANVRE, Cannabis sativa—c. *Bâtard*, *Galeopsis grandiflora*—c. *Indien*, *Bangue*.

CHAOMANTI'A. The alchymists meant, by this word, the art of predicting the future from observation of the air. The word *Chaos* was used by Paracelsus for the air; (*μαρτυα*, 'divination,')

CHAOSDA, Plague.

CHAPELIÈRE, *Tussilago petasites*.

CHAPERON DE MOINE, ('monks hood,') *Aconitum napellus*.

CHAPETONNADE, *Chappetonnaide*.

CHAPITEAU, (from *capitellum*, dim. of *caput*,) *Alenbic*.

CHAPPEDONADE, *Chappetonnade*.

CHAPPETONNADE, *Chappedonade*, *Chappetonnade* (F.), *Vom'itus rabio'sus*. Vomiting, accompanied by furious delirium, attacking strangers in hot countries.

CHAPS, Rhagades.

CHARACTER, (*χαρακτῆρ*, 'a mark or impression,') (F.) *Caractère*. In Pathology it is used synonymously with stamp or appearance. We say, 'A disease is of an unfavourable character.' 'The prevailing epidemic has a bilious character,' &c. In *Mental philosophy* it means—that which distinguishes one individual from another, as regards his understanding and passions. See *Symbol*.

CHARA'DRIUS, (*χαραδριος*,) Ælian thus calls a bird, which was reputed to cure jaundice. The word now means the plover.

CHARANTIA, *Momordica elaterium*.

CHARBON, (from *carbo*, 'coal,') *Anthrax carbo*, *Carbo ligni*—c. *Animal*, *Carbo animalis*—c. *Minéral*, *Graphites*—c. *Pulmonaire*, *Anthrax cois*.

CHARBONNEUX, *Anthrax cois*.

CHARCOAL, *Carbo*—c. *Animal*, *Carbo animalis*.

CHARDON AUX ÂNES, (*carduus*, 'a thistle') *Onopordium acanthium*—c. *Bénit*, *Centaurea benedicta*—c. à *Bonnetier*, *Dipsacus fullonum*—c. *Étoilé*, *Centaurea calcitrapa*—c. à *Foulon*, *Dipsacus fullonum*—c. *Hémorrhoidal*, *Cirsium arvense*—c. *Marie*, *Carduus Marianus*—c. *Roland*, *Eryngium campestre*.

CHARLATAN, ([L.] *ciarlare*, 'to talk much,') *Circulator*, *Circumforaneus*, *Periodeutes*, *Pseudomedicus*, *Aggr'ta*, *Anacyleton*, a Quack, Quacksalber, *Empirical Pretender*, *Empiric*. Originally, one who went from place to place to sell a medicine, to which he attributed marvellous properties. By extension—any individual, who endeavours to deceive the public by passing himself off as more skillful than he really is. According to *Ménage*, the word comes from *circulatus*, a corruption of *circulator*.

CHARLATANRY, *Aggr'tia*, *Banau'sia*, the conduct or action of a charlatan, *Quack'ery*, *Empiricism*, (F.) *Charlatanerie*, *Charlatanisme*.

CHARLOCK, *Sinapis arvensis*.

CHARLOTTEVILLE SULPHUR SPRING, see *Canada*, *Mineral Waters* of.

CHARM, (*Car'men*, 'a verse,' because charms often consisted of verses, whence comes the Italian, *Ciarma*, (F.) *Charme*, with the same signification,) *Canta'tio*, *Incantamentum*. A trick, a spell, an enchantment. A sort of magic, or superstitious practice, consisting of words, characters, &c., by which, it was believed, individuals might be struck with sickness or death, or be restored to health.

The following are specimens of old charms—verse charms:

For stanching Blood, (Pepys.)

Sanguis mane in te
Sicut Christus fuit in se;
Sanguis mane in tuâ venâ
Sicut Christus in suâ penâ;
Sanguis mane fixus,
Sicut Christus quando fuit crucifixus.

For Cramp, (Pepys.)

Cramp be thou faintless,
As our lady was sinless,
When she bare Jesus.

For the Foot when asleep. (Coleridge.)

Foot! foot! foot! is fast asleep!
Thumb! thumb! thumb! in spittle we steep;
Crosses three we make to ease us.
Two for the thieves, and one for Christ Jesus.

The same charm served for cramp in the leg, with this substitution:

The devil is tying a knot in my leg!
Mark, Luke, and John, unloose it, I beg!—
Crosses three, &c.

For a Burn, (Pepys.)

There came three angels out of the East;
The one brought fire, the other brought frost.
Out fire, in frost.
In the name of the Father and Son and Holy Ghost.
Amen.

CHARNIÈRE, (*cardo*, gen. *cardinis*, 'a hinge,') Uinglymus.

CHARNU, (from *chair*, (L.) *caro*, 'flesh,') Car-neous.

CHARPENTE OSSEUSE, (carpenters' work, 'frame,' extended from low (L.) *carpentum*, 'a carriage,') Skeleton.

CHARPIE, (from (L.) *carpere*, 'to pluck, to tear,') Lintum—*c. Brute*, see Lintum—*c. Rapée*, see Lintum.

CHARTA ('paper') **ANTIARTHRITICA**, Gout paper—*c. Antirheumatica*, Gout paper—*c. Vesicatoria*, see Sparadrapum Vesicatorium—*c. Virginea*, Amnios.

CHARTRE, ('a prison,' from (L.) *carcer*, thence languor, wasting, [?]) *Tabes mesenterica*.

CHAS (F.), *A'cûs for'a'men*, The eye of a needle. Sometimes, this opening is near the point of the instrument, as in the ligature needle.

CHASCHISCH, see Bangue.

CHASME, (*χασμή*), Yawning.

CHASSE (F.), (from *capra*, 'a case,') [?] *Manubrium*. A kind of handle composed of two movable laminae of horn, shell, or ivory, united only at the extremity, which holds the blade of the instrument,—as in the common bleeding lancet. Also, a *liqueur* or alcoholic liquor, taken after a meal; hence *chasse-café*, *chasse-tout*, &c., from (F.) *chasser*, 'to drive,' because presumed to aid digestion.

CHASSIE (F.), *Le'ma*, *Lip'pa*, *Gla'ma*, *Gle'mê*, *Gra'mia*, *Lemos'itaz*, *Se'bum palpebra'te*; *Gum of the eye*, (Prov.) *Gound* or *Gownde*, (from (F.) *chasser*, 'to drive out,' [?] or from (L.) *casus*, 'cheese,' [?]) A sebaceous humour, secreted mainly by the follicles of Meibomius, which sometimes glues the eyelids together.

CHASSIEUX (F.), *Lippus*; covered with *Chassie*—as *Paupières chassieuses*.

CHASTE TREE, Vitex.

CHÂTAIGNE, see *Fagus castanea*—*c. d'Eau*, *Trapa natans*.

CHÂTAIGNIER COMMUN, (*castanea*,) *Fagus castanea*—*c. Nain*, *Fagus castanea pumila*.

CHATEAU-LANDON, MIN'ERAL WA'TERS OF. A town three leagues from Nemours, in France. The waters contain alum and iron.

CHATEAU-SALINS, MIN'ERAL WA'TERS OF. A town in the department of La-Meurthe, France. The waters contain carbonate of lime, sulphates of lime and magnesia, and chlorides of magnesium and sodium.

CHÂTELDON, MIN'ERAL WA'TERS OF. Chateldon is in the department of Puy-de-Dôme, France. The waters contain carbonic acid and iron. Season from the 1st of May to the 1st of October.

CHATEL-GUYON, MIN'ERAL WA'TERS OF. A village in France, in the department Puy-de-Dôme, near which there are five thermal acidulous springs. Temperature, 86° Fahrenheit. Season from the 15th of May to the 15th of September.

CHATON (F.), ('a husk,' (G.) *Kasten*, 'a chest,') In pathology, it means a *funda* or cavity formed by the irregular or *hour-glass contraction*, of the uterus, in which the placenta is often retained or *enchatonné* after the birth of the child. It is detected by passing the fingers along the cord as far as the part which is contracted, when the placenta will not be felt.

The treatment consists in relaxing by a large dose of an opiate, then passing the fingers along

the cord, and gradually dilating the opening through which it passes, until it permits the hand to go through. The placenta must then be grasped and gently withdrawn.

CHATON, see Vaginal process.

CHATONNÉ, *Incarcera'tus*, (from *chaton*,) An epithet in French for anything shut up in a sac, or imprisoned in any manner—as

CALCUL CHATONNÉ (F.) *Calcul enkysé*, *Cal'culus incarceration'tus*, *Encys'ted Cal'culus*. A urinary calculus, adherent to the inner surface of the bladder, so that it is immovable, and cannot pass to the different parts of that organ. This happens when a calculus forms in some natural or accidental cavity of the bladder; or where ulceration gives rise to fungous growths, which surround the calculus; or when it is lodged in the orifice of the ureter or urethra.

CHATONNÉ, **PLACENTA** (F.). The placenta when retained as described under *Chaton*.

CHATONNEMENT (F.), *Enchatonnement*, *Incarcera'tio*, *Chatonnement du placenta*, *Enkystement*.—Hour-glass contraction of the uterus. See *Chaton*.

CHATOUILLEMENT (F.), (from (L.) *catulire*, 'to desire the male,') This word sometimes means the action of tickling or titillation (*Titilla'tio*) and, at others, the sensation, *Pruri'tus*, *Itching*, which gives rise to the action.

CHÂTRÉ, Castratus.

CHÂTRURE, Castration.

CHAUDEAU, (*chaud*, 'warm or hot,') Caudle.

CHAUDEBOURG, MIN'ERAL WA'TERS OF. C. is three quarters of a league from Thionville, in France. The waters contain iron, sulphate of lime, sulphate of magnesia, and carbonate of lime.

CHAUDEPISSE, (*chaud*, 'hot,' and *pisser*, 'to make water,') *Gonorrhœa impura*—*c. Cordée*, *Gonorrhœa cordata*—*c. Tombée dans les Bourses*, *Hernia humoralis*.

CHAUDES-AIGÜES, MIN'ERAL WA'TERS OF. A small town in the department of Cantal, France, where there is a number of saline springs containing carbonic acid, carbonate of soda, and chloride of sodium. Temp. 190° Fahrenheit.

CHAUDIÈRE, (from *chaud*, 'hot,') see *Alem bic*.

CHAUFFOIR (F.), (from *chauffer*, (L.) *calefacere*, 'to make warm,') *Lintum*, *Calefacto'r-ium*. According to the *Académie*, a warmed cloth, used either for the purpose of warming a patient, or to apply to a female recently delivered.

CHAULE, Maxillary bone.

CHAULMOO'GRA. The fruit of the *Chaulmo'gra*, *Gynocar'dia odora'ta*. Order, *Capparidæ*, of India. A bland fixed oil is obtained by expressing the dried seed, which is used by the natives in leprosy. The seed are given internally, in the dose of about ʒss in the day, and the eruption is rubbed with the oil.

CHAUSSE (F.), ('a stocking,' from (L.) *calx*, 'the foot,') *Chausse d'Hippocrate*, *Manche d'Hippocrate*, *Man'ica Hippoc'ratis*, *Man'ica*, *Hippoc'rates' Sleeve*. A conical bag, made of flannel, for straining liquids.

CHAUSSE-TRAPPE, *Centaurea calcitrapa*.

CHAUVETÉ, Calvities.

CHAUX, Calx—*c. Bichlorure de*, *Calcis chloridum*—*c. Chlorure de*, *Calcis chloridum*—*c. Éteinte*, see Calx—*c. Hydrochlorate de*, *Calcis murias*—*c. Hydrosulfate de*, *Calcis sulphuretum*—*c. Muriate de*, *Calcis murias*—*c. Muriate oxigéné de*, *Calcis chloridum*—*c. Muriate surozigéné de*, *Calcis chloridum*—*c. Oxichlorure de*, *Calcis chloridum*—*c. Oximuriate de*, *Calcis chloridum*—

1. *Phosphate de*, see *Cornu cervi*—*c. Vire*, *Calx viva*.

CHAVEL, Maxillary bone.

CHAVICA BETLE, Betel—*c. Roxburghii*, *Piper longum*.

CHAWBONE, Maxillary bone.

CHAWD PYS, (*Chaudepisse*), Gonorrhœa impura.

CHAWING, (from Sax. *ceopan*, (G.) *kauen*, 'to chew or chew.') Mastication.

CHECKER-BERRY, *Arbutus uva ursi*.

CHEEK, (Sax. *ceac*), *Gena*—*c. Teeth*, *Molar teeth*.

CHEESE, (Sax. *cere*), *Ca'seus*, *Ty'ros*, *Pec'tē*, (Sc.) *Furnage*, (F.) *Fromage*. An aliment, prepared from the caseous and oleaginous parts of milk. Fresh cheeses owe their chief medical properties to the immediate principle, essentially cheesy, to which the name *ca'seum* or *ca'sein* has been applied. Those, which have been recently salted, are digested with comparative facility. The flavour of cheese is owing to an ammoniacal caseate. On the whole, cheese itself is not easy of digestion, although it may stimulate the stomach to greater exertion, and thus aid in the digestion of other substances, in accordance with the old saying,

"Cheese is a surly elf
Digesting all things but itself."

SMEARCASE, (D.) *Smerkaas*, (G.) *Schmierküse*; (*schmieren*, 'to spread,' and *Küse*, 'cheese.') *Cottage cheese*, *Potcheese*, is a preparation of curds used as food.

CHEESE, COTTAGE, see *Cheese*—*c. Pot*, see *Cheese*—*c. Rennet*, *Galium verum*.

CHEESELLOPE, (Prov.) (from Sax. *cerelob*, 'curdled milk,') *Rennet*.

CHEE'SY, *Casearius*, *Caseosus*, *Tyro'des*, (F.) *Caseux* ou *Caseux*. Having the nature of cheese.

CHEF, Bandage (tail.)

CHEGOE, *Chique*.

CHEGRE, *Chique*.

CHEILALGIA, *Chilal'gia*, (*cheilo*, and *αλγος*, 'pain,') Pain in the lip.

CHEILITIS, *Chilitis*, (*cheilo*, and *itis*, denoting inflammation.) Inflammation of the lip.

CHEILO, (from *χειλος*, 'a lip,') in composition, lip.

CHEILOAC'CE, (*cheilo*, and *κακος*, 'evil,') *Labrial'cium*. A disease, characterized, it is said, by swelling, induration, and slight redness of the lips without inflammation; reputed, but without any authority, to be common in England and Scotland, amongst children. Also, the thickness of the upper lip of scrofulous children. See *Stomacace*, and *Cancer aquaticus*.

CHELOCARCINO'MA, gen. *Cheilocarcino'matis*, (*cheilo*, and *καρκινωμα*, 'a cancer,') Cancer of the lip.

CHEILOMALACIA, (*cheilo*, and *μαλακος*, 'soft,') Cancer aquaticus, *Stomacace*.

CHEILON, Chilon.

CHEILON'CUS, *Cheilophy'ma*, (*cheilo*, and *γκος*, 'swelling,') A swelling of the lip.

CHEILOPHYMA, gen. *Cheilophy'matis*, (*cheilo*, and *φυμα*, 'a tumour,') *Cheiloncus*.

CHEILOPLAS'TICĒ, *Cheiloplas'ticē*, *Cheiloplasty*, (*cheilo*, and *πλαστικός*, 'forming,') The operation for an artificial lip.

CHEILOS, (*χειλος*,) Lip.

CHEIMA, (*χειμα*,) Cold.

CHEIMETLON, (*χειμετων*, from *χειμα*, 'winter,' and *ταλιν*, 'to suffer,') *Chilblain*.

CHEIMIA, (*χειμη*, 'winter,') *Rigor*.

CHEIR, (*χειρ*, 'hand,') *Manus*, see *Chir*.

CHEIRANTHUS CHEIRI, (*cheir*, and *ανθος*, 'flower,') 'a flower for the hand.' The systematic name of the *Common Yellow Wall Flower*, *Viola lutea*, *Leucan'tium luteum*, *Key'ri*, *Cheiri*,

(F.) *Gérofée*, *Gérofée* ou *Vivier jaune*. *Ord. Cruciferae*. The flowers have been esteemed *nerve*, *narcotic*, and *deobstruent*.

CHEIRAP'SIA. The action of rubbing or scratching, (*cheir*, and *απαι*, 'I touch,') A troublesome symptom in the itch.

CHEIRARTHROAC'CE, (*cheir*, *αρθρον*, 'a joint,' and *κακια*, 'evil,') Inflammation of the articular surfaces of the wrist.

CHEIRIATER, (*χειρ*, and *ιατρος*, 'a physician,') Surgeon.

CHEIRIS'MA, gen. *Cheiriam'atis*, *Cheiris'mus*, (*χειρισμα*, *χειρισμος*,) The act of touching:—*handling*. Any manual operation.

CHEIRIXIS, (*χειριξις*,) Surgery.

CHEIRONOM'IA, *Cheironom'ia*, (*χειρονομια*, from *χειρονομειν*, 'I exercise with the hands,') An exercise, referred to by Hippocrates, which consisted in using the hands, as in exercise with the dumbbells.

CHEIROPLETHES, (*χειροπληθης*, from *χειρ*, and *πληθω*, 'I fill,') Fasciculus.

CHEIROISIS, (*χειροισις*, from *χειροειν*, 'to subdue,') Subactio.

CHE'LA, (*χηλη*,) *Chele*. This word has several significations. *Che'la*, a forked probe used for extracting polypi from the nose. *Che'lae*—chaps, or cracks on the feet, organs of generation, &c. *Che'lae* likewise means claws, especially those of the crab. See *Cancrorum Chelæ*.

CHELE PALPEBRARUM, see *Tarsus*.

CHELAPA, *Convolvulus jalapa*.

CHELE, (*χηλη*,) *Chela*.

CHELIDOINE GRANDE, *Chelidonium majus*—*c. Petite*, *Ranunculus ficaria*.

CHELIDON, (*χελιδων*,) the hollow at the bend of the arm; *Hirundo*.

CHELIDONIA ROTUNDIFOLIA MINOR, (*χελιδων*, 'a swallow,' because its flowering coincides with the appearance of the swallow.) *Ranunculus ficaria*.

CHELIDONIUM, (*χελιδonium*,) *Bryonia alba*—*c. Diphylum*, *Stylophorum diphylum*—*c. Hæmatodes*, *Ch. majus*.

CHELIDONIUM MA'JUS, *Ch. hæmato'des*, *Papaver Corniculatum* seu *lu'teum*, *Common Celandine*, *Tet'terwort*, (F.) *Chélidoine grande*, *L'Eclair*, *Herbe à l'hirondelle*. *Ord. Papaveraceæ*. *Sex. Syst. Polyandria Monogynia*. The root and recent plant have been considered aperient and diuretic. Externally, the juice has been employed in some cutaneous diseases.

CHELIDONIUM MINUS, *Ranunculus ficaria*.

CHELIDONIUS LAPIS, (from *χελιδων*, 'a swallow,') A name given to stones, which, it was pretended, existed in the stomach of young swallows. They were formerly believed capable of curing epilepsy.

CHELIS, (from *chele*,) *Kelis*.

CHELOID, (*χελυς*, 'a tortoise,' and *ειδος*, 'resemblance,') *Canceroid*.

CHELOIDE CICATRICE'ELLE, see *Canceroid*.

CHELOIS, see *Canceroid*.

CHELO'NĒ, (*χελωνη*, 'a tortoise,') An instrument for extending a limb; so called, because in its slow motions it resembled a tortoise. See *Testudo*.

CHELO'NĒ GLA'BRA, *Com'mon Snake-head*, *Turtle-head*, *Turtlebloom*, *Shell-flower*, *Balmoney*. An indigenous plant, *Sex. Syst. Didynamia Angiospermia*; blossoming from July to November. The leaves are bitter and tonic; without any aromatic smell, and with very little astringency.

CHELO'NIA MY'DAS, *The Green Turtle*. This species of turtle abounds on the coast of Florida. It is the one so prized by the epicure.

CHELO'NION, *Che'lodium*, (from *χελωνη*, 'a tortoise,') from its resembling in shape the shell

of the tortoise. The upper, gibbous part of the back. The scapula.

CHELONOPHAGI, (χελων, 'a tortoise,' and φαγω, 'I eat.') An ancient name for certain tribes, who dwelt on the coast of the Red Sea, and who lived only on tortoises.

CHELSEA PEN'SIONER. An empirical remedy for rheumatism and gout, sold under this name. (Gum guaiac, 3i; powdered rhubarb, 3ij; cream of tartar, 3j; flowers of sulphur, 3j; one nutmeg, finely powdered; made into an electuary with a pound of clarified honey). Dose, two spoonfuls.

CHEL'TENHAM, MIN'ERAL WA'TERS OF. C. is a town in Gloucestershire, England, nine miles from Gloucester, and 94 W. of London. Its water is one of the most celebrated natural purgatives in England. It is a saline, acidulous chalybeate, and is much frequented. Its main constituents are chloride of sodium, sulphate of soda, sulphate of magnesium, carbonic acid and carbonate of iron.

CHEL'TENHAM SALTS. These are sometimes made from the waters; at others, factitiously. The following is a formula: Sodii chlorid., magnes. sulphat., sodæ sulph., aa 1b; dissolve, filter, evaporate to dryness: then add Ferri sulph., 3ss.

CHEL'TENHAM WA'TER, ARTIFICI'AL, may be made of Epsom salts, gr. xij; iron filings, gr. j; Glauber's salt, 3iv; water, 4 gallons; impregnated with the gas from marble powder and sulphuric acid, aa 3ij.

CHELYS, (χελυς,) Thorax.

CHELYS'CION, (χελυσιον, from χελυς, 'the thorax,') A short dry cough.

CHEMÊ, (χημη,) Chama, Che'ramis. An ancient measure equivalent to about two teaspoonfuls.

CHEMEUTICE, (χημειτικη,) Chymistry.

CHEMIA, (χημια,) Chymistry.

CHEMIATER, Chymiatier.

CHEMIATRIA, Chymiatria.

CHEMICOHISTOLOGY, see Chymicohistology.

CHEMICUS, Chymical, Chymist.

CHEMISSETTE, (dim. of chemise,) Condom.

CHEMISMUS, Chymism.

CHEMIST, Chymist.

CHEMISTRY, Chymistry.

CHEMO'SIS, Chymo'sis, (χημοσις, from χημη, 'an aperture,' or rather, perhaps from χυμος, 'a humour,') A name given to ophthalmia, when the conjunctiva, surrounding the cornea, forms a high ring, making the cornea seem, as it were, at the bottom of a well. By some, it is used synonymously with ophthalmia membranarum. See Ophthalmia.

CHEMOTICE, Chymistry.

CHEMOTICUS, Chymical.

CHENAY, MIN'ERAL WA'TERS OF. Chenay is a town in France, two leagues from Rheims. The waters are chalybeate.

CHÊNE, Quercus alba—c. Marin, Fucus vesiculosus—c. Petit, Teucrium chamaedrys—c. Vert, Ilex aquifolium.

CHÊNETTE, (dim. of chène,) Teucrium chamaedrys.

CHÊNEVIS, see Cannabis sativa.

CHENOBOSCON, (χην, 'a goose,' and βοσκη, 'food,') Potentilla anserina.

CHENOC'OPRUS, (χην, 'a goose,' and κοπος, 'dug,') The dung of the goose is so designated in some old Pharmacopœias. It was formerly employed as a febrifuge and diuretic.

CHÊNOPODE, Chenopodium.

CHENOPODIUM AMBROSIIACUM, (χην, 'a goose,' and ποδ, gen. ποδος, 'a foot,') Ch. Ambrosioides.

CHENOPO'DIUM AMBROSIOI'DES, Ch. suffruticosa.

sum seu ambrosi'acum seu Mexica'num. At'riplex Mexica'na, Ambri'na ambrosioi'des, Bo'trys, B. Mexica'na seu America'na, Ambrosioides Mexica'na, Artemis'ia Bo'trys, Mex'ico or Span'ish Tea, (F.) Anserine, Thé du Mexique, Ambrosie ou Ambrosie du Mexique. Ord. Chenopodiaceæ. The infusion was once drunk as tea. It has been given in paralytic cases; and in the United States is said to be used as an anthelmintic indiscriminately with Ch. anthelminticum.

CHENOPODIUM ANTHELMINTICUM, Chenopodium, Bo'trys seu Ambri'na anthelmintica, Worm'seed, Worm goose's foot, Worm'seed goose's foot, Jeru'salem Oak of America, Goose's foot, Stink'weed, (F.) Anserine anthelmintique, A. vermifuge. This plant grows plentifully in the United States. The fruit—Chenopodium, (Ph. U. S.)—is much used in cases of worms. Dose of the powder, from a teaspoonful to a tablespoonful or more. The oil, O'leum Chenopo'dii, (Ph. U. S.), from 8 to 10 drops, is more frequently exhibited. It is as much used in America as the Semen Santonici is in England.

CHENOPODIUM BO'NUS HENRI'CUS, Chrysolach'anum, Mercuria'lis, Bo'nus Henri'cus, To'ta bo'na, Lap'athum unctuosum, Chenopodium, Ch. sagittatum, Pes anserinus, English Mer'cury, All'good, Angular-leaved goose's foot, Mer'cury goose's foot, Good King Hen'ry, (F.) Anserine Bon Henri, Épinard sauvage. The leaves are emollient, and have been applied to ulcers, &c. It has also been considered refrigerant and ecoprotic.

CHENOPODIUM BO'TRYS, Ambri'na Bo'trys, Bo'trys, B. vulgaria, Ambrosia, Artemis'ia Chenopodium, At'riplex odora'ta seu suave'olens; Jeru'salem Oak, (Eng.), (F.) Anserine Botrys, possesses anthelmintic properties, and was once given in diseases of the chest, palsy, &c. It is useless.

CHENOPODIUM FETIDUM, Chenopodium vulvaria—c. Oidium, Ch. vulvaria.

CHENOPODIUM QUI'NOA, Qui'nua. A nutritious, wholesome, and agreeable article of food with the Peruvians. The leaves, before the plant attains maturity, are eaten as spinach: but the seeds are most generally used as food, boiled in milk or broth, and sometimes cooked with cheese and Spanish pepper.

CHENOPODIUM SAGITTATUM, Ch. Bonus Henricus—c. Suffruticosum, Ch. ambrosioides.

CHENOPODIUM VULVARIA seu foetidum seu ol'idum, At'riplex foetida seu ol'ida, Vulva'ria, Garos'mum, Ra'phex, Stink'ing Or'ache or Goose's foot, (F.) Vulvaire, Arroche puant, Anserine fetide. The fetid smell has occasioned it to be used as an antispasmodic and nervine.

CHEOPINA, Chopine.

CHEOPLASTIC, Cheoplasticus; (χεω, 'to pour,' and πλαστικός, 'formative,') Applied to a process for mounting artificial teeth, which consists in pouring an alloy of metals into a properly prepared matrix.

CHEQUERBERRY, Gaultheria.

CHERAGRA, Chiragra.

CHERAMIS, (χηραμς,) Cheme.

CHERBACHEM, Veratrum album.

CHERBAS, Lactuca.

CHERMES, Kermes—c. Mineral, Antimonit sulphuretum præcipitatum, and A. sulphuretum rubrum.

CHERNIBIUM, (χερνιβιον,) Urinal.

CHERRY, see Prunus cerasus—c. Bird, Prunus padus—c. Choke, Prunus Virginiana—c. Ground, Physalis viscosa.

CHERRY TREE, BLACK, Prunus avium—c. Tree, red, Prunus cerasus—c. Tree, wild, Prunus Virginiana—c. Water, Kirschwasser.

CHERRY, WILD CLUSTER, Prunus padus—c. Winter, Physalis.

CHERSÆ, Fœces.

CHERVI, *Sium sisarum*.

CHERVIL, (from *cherophyllum*), Scandix cerefolium—c. Wild, *Charophyllum sylvestre*.

CHE/SIS, (χεῖσιν, 'to go to stool.') A more frequent desire to evacuate the bowels.

CHEST, (κιστή, Sax. cete, or cýrē,) Thorax—ch. Explorer, see Explorer, chest.

CHESTMEASURER, SIBSON'S. An instrument, described by Dr. Sibson, which is used for determining the mobility of the chest. It is a species of spring, which, when applied to the parietes of the chest, measures the modifications of its diameters, and indicates, by the motion of the index hand on a dial, any movement of respiration, to the hundredth of an inch.

CHEST, NARROW, see Lordosis.

CHESTNUT TREE, *Fagus castanea*.

CHEVAUCHEMENT (F.), *Os sium superpositio* seu *equitatio*, *Parallaxis*, *Parallagma*. (*chevaucher*, (*cheval*, 'a horse'), 'to ride on horseback.') The riding of one bone over another after fracture, giving rise to shortening of the limb. See Riding of Bones.

CHEVELURE, (from *cheveu*), Scalp.

CHEVESTRE, *Chevêtre*, *Capistrum*. A bandage, applied round the head in cases of fracture or luxation of the lower jaw. According to the mode in which it is made, it is called *simple*, *double*, *oblique*, &c.

CHEVEU, Capillus.

CHEVILLE DU PIED, Malleolus.

CHÈVREFEUILLE, ('*caprifolium*'), *Lonicera periclymenum*.

CHEVROTAIN PORTE-MUSC, (from *chèvre*, 'a goat.') See Musk.

CHEVROTANTE (VOIX), (from *chèvre*, 'a goat.') Egophony.

CHEWING, Mastication.

CHEYLETES SCABIEI, see Psora.

CHEZANAN'CÈ, (χεζαναγκη, from χεζω, 'I go to stool,' and αναγκη, 'necessity.') An ointment composed of honey and alum, and rubbed on the anus to occasion evacuations.

CHIA, *Chî'a ter'ra*, (from *Chios*, an island where it was found.) A kind of white earth, formerly used for burns.

CHIACUM COLLYRIUM. A collyrium consisting of several drugs and Chian wine.

CHIADUS, Furunculus.

CHIASM, *Chias'mus*, *Chias'ma*, (χιασμος, χιασμα, from χιαζω, I mark like the letter χ.) The crucial union of parts,—as the optic commissure or *chiasm* of the optic nerves, *Chias'mus* seu *Chias'ma nervorum optico'rum*.

CHIASTER, (χιαζω, 'I mark like the letter χ') Kiaster.

CHIAS'TUS, (χιαστρος.) A bandage, so called because it resembles the letter χ.

CHIBOU, see *Bursera gummiifera*.

CHU'CHA. A drink made in Peru with Indian meal dried in the sun, and fermented with water. Its taste is that of bad cider. It is also made from rice, peas, barley, &c. In many places the grain is chewed—generally by old women—and the joint product is allowed to ferment to form the drink.

CHICK'S SPRINGS, see South Carolina, mineral waters of.

CHICKEN-BREADED, see Lordosis.

CHICK'EN-BROTH. When chicken-tea is boiled down one-half, with the addition of a little *parsley* or *celery*, and the yolk of an egg previously beaten up in two ounces of soft water, it forms a soup much relished by the convalescent.

CHICKEN-PEPPER, *Ranunculus abortivus*.

CHICKENPOX, *Varicella*.

CHICKEN-TEA, *Chicken-water*. This may be prepared as follows: Take a small chicken, freed

from the skin and fat between the muscles; and having divided it longitudinally, remove the lungs, liver, and every thing adhering to the back and side-bones: cut the whole—bones and muscles—into very thin slices; put into a pan with a sufficient quantity of boiling water; cover the pan; and simmer with a slow fire for two hours. Put the pan upon the stove for half an hour, and strain through a sieve.

Used where the lightest animal diet is indicated.

CHICKEN-WATER, *Chicken-tea*.

CHICKENWORT, *Alsine media*.

CHICKWEED, *Alsine media*, *Anagallis arvensis*—c. *Water*, *Callitriche verna*.

CHICORÉE DES JARDINS, *Cichorium endivia*—c. *Sauvage*, *Cichorium intybus*.

CHICORY, *Cichorium intybus*.

CHICOT, Stump (of a tooth).

CHIOOTIN (F.) (corr. from *Socotrine aloes* [?]). The bitter powder or juice of the colocynth, with which the nurse, at times, rubs her nipple when desirous of weaning an infant.

CHIENDENT, ('dogtooth,') *Triticum repens*—c. *Rouge*, *Sarsaparilla Germanica*.

CHIGGO, *Chique*.

CHIGGRE, *Chique*.

CHIGOE, *Chique*.

CHILALGIA, *Chilalgia*.

CHILBLADDER, *Chilblain*.

CHILBLAIN, (Prov.) *Chilbladder*, *Per'nio*, *Bugan'tia*, *Eryth'ma Per'nio* seu à *Frigore*, *Chemet'on*, *Chimet'um*, *Chî'mon*, *Mal'cè*, (*chill*, 'cold,' and *blain*, 'a pustule.'). (F.) *Engelure*. An erythematous inflammation of the feet, hands, &c., occasioned by cold. It is very common in youth—not so in the adult or in advanced age. It is apt to degenerate into painful, indolent ulcerations, called *Kibes* (Sc.) see *Mulea*. Chilblains are prevented by accustoming the parts to exposure; and are treated by stimulant, terebinthinate, and balsamic washes, ointments, and liniments.

CHILD, (Sax. eild,) *Infans*. *Puer*.

CHILDBEARING, *Parturition*.

CHILDBED, *Parturient*, *Parturition*.

CHILDBED FEVER, *Puerperal fever*.

CHILDBIRTH, *Parturition*.

CHILDCROWING, *Asthma thymicum*.

CHILDHOOD, *Infancy*.

CHILDILL, *Parturition*.

CHILDING, see *Parturient*.

CHILDMURDER, *Infanticide*.

CHILDREN'S BANE, *Cicuta maculata*.

CHILI, CLIMATE OF, see Lima.

CHIL', MIN'ERAL WA'TERS OF. The most celebrated mineral springs of Chili, in South America, are those of Peldehués and Cauquenes. The former are not far from St. Jago. They consist of two springs, one thermal, the other cold. The hot spring is clear, inodorous, and contains soda and carbonic acid. The cold spring contains iron and sulphate of soda. Cauquenes is much resorted to by invalids during the summer. Mineral waters are very common in Chili. Those of Apoquindo, Colina, and Cauquenes, have been analyzed by Prof. J. Lawrence Smith, M. D.

CHILIOGRAMMA, (χίλιοι, 'a thousand,' and *gramma*, 'a gramme.'). Kilogramme.

CHILIOPHYLLON, (χίλιοι, 'a thousand,' and *φύλλον*, 'a leaf.'). *Achillea millefolium*.

CHILITIS, *Cheilitis*.

CHILL, (Sax. cèle,) Rigor—c. Dumb, Fever, masked.

CHILLI, *Capsicum*.

CHILO, in composition, see *Cheilo*.

CHI'LO, *Cheî'lon*, *Cheili'tis*, (χελος, 'a lip.'). An inflammatory swelling of the lips. One who has a thick lip: *La'beo*, *La'bes*.

CHILOPLASTICE, *Cheiloplastice*.

CHI'LOSTOMATOPLASTY, *Chil'stomato-*

plast'icæ, *Chilo-stomato-plas'ticæ*, (*χειλος*, 'a lip,' *στόμα*, 'mouth,' and *πλασσειν*, 'to form.') A surgical operation for restoring the buccal opening, after the removal of epithelioma of the lower lip.

CHIMAPHILA, (*χειμα*, 'winter,' and *φιλω*, 'I love.') *Pyrola umbellata*—c. *Umbellata*, *Pyrola umbellata*.

CHIMETLUM, (*χιμετλον*, from *χειμα*, 'winter,' and *πασσειν*, 'to suffer.') *Chilblain*.

CHIMIA, Chymistry.

CHIMIATER, (*chimia*, and *ιατρος*, 'a physician.') Chymiatér.

CHIMIATRIA, Chymiatría.

CHIMIE, Chymistry.

CHIMISME, Chymism.

CHIMISTE, Chymist.

CHIMON, (*χιμων*, from *χειμα*, 'winter.') *Chilblain*, Cold.

CHIN, (Sax. cýn.) *Mentum*.

CHINA, *Cinchona*, (from *quinquina*.) *Smilax china*, (from *china*),—c. *American* or *West India*, *Smilax pseudo-china*—c. *Calisaya*, *Cinchona cordifolia* cortex—c. *Occidentalis*, *Smilax pseudo-china*—c. *Orientalis*, *Smilax china*—c. *Ponderosa*, *Smilax china*—c. *Regia*, *Cinchona cordifolia* cortex—c. *Root*, *Smilax china*—c. *Spuria nodosa*, *Smilax pseudo-china*—c. *Vera*, *Smilax china*.

CHINCAPIN, *Fagus castanea pumila*—c. *Water*, *Nelumbium luteum*.

CHINCHE, ([S.] a bed bug,) *Cimex*.

CHINCHINA, *Cinchona*.

CHINCHONA, *Cinchona*.

CHINCHUNCHULLI, *Iodidium marceei*.

CHINCOUGH, (by *onomatopœia*: from *chin*, Teut. *kink*.) *Pertussis*.

CHINESE MEDICINE OF THE, *Medicína Sínica*. Medicine has been long, but most imperfectly, practised by the Chinese. From their therapeutics we have obtained the old operations of acupuncturè and moxibustion.

CHING'S WORM LOZENGES, see *Worm Lozenges*, *Ching's*.

CHINIDIN, *Quinidia*.

CHININUM, *Quinine*. See *Chinium*.

CHINIOID'INE, *Chinoid'ine*, *Thinoid'ina*, *Chinoid'in*, *Quinoid'ine*, *Quinoid'in*, *Quinod'in*; (from *china*, 'cinchona.') A substance presumed to be an alkaloid by Sertürner, who separated it from cinchona. It has been supposed to be a mixture of quinia, cinchona, and a peculiar resinous matter, but according to Liebig it is simply the alkaloid quinia in an amorphous state—*amorphous quinia*.

CHINIUM, *Quinine*—c. *Aceticum*, *Quinix acetat*—c. *Arsenicum*, *Quinix arseniat*—c. *Citricum*, *Quinix citrat*—c. *Ferrocyanogenatum*, *Quinix Ferrocyanas*—c. *Hydrochloricum*, *Quinix murias*—c. *Hydroiodicum*, *Quinix hydriodas*—c. *Lacticum*, *Quinix lactas*—c. *Muriaticum*, *Quinix murias*—c. *Nitricum*, *Quinix nitras*—c. *Phosphoricum*, *Quinix phosphas*—c. *Salutum*, *Quinix murias*—c. *Sulphuricum*, *Quinix sulphas*—c. *Tannicum*, *Quinix et Cinchonix tannas*—c. *Valerianicum*, *Quinix valerianas*.

CHINNEYWEED, *Lichen roccella*.

CHINOLEINUM, (*chinium*, and *ελαιον*, 'oil') *Leukoleinum*.

CHINQUAPIN, *Fagus castanea pumila*.

CHINWELK, (from *chin*, and *whelk*.) *Syncochis*.

CHIOCOCCÆ RADIX, (*χιων*, 'snow,' and *κοκκος*, 'berry.') *Cainxæ radix*.

CHIOLI, *Furunculi*.

CHION, (*χιων*.) *Snow*.

CHIONANTHUS VIRGINICA, *Fringetree*, *Snowdrop Tree*; indigenous; *Nat. Ord. Oleaceæ*; (*χιων*, 'snow,' and *ανθος*, 'a flower.') An infusion of the root has been used as an antiperiodic and tonic.

CHIONYPHE CARTERI, (*χιων*, 'snow,' and *ψη*, 'texture,') See *Mycetoma*.

CHIQUE (F.), *Puce pénétrante*, *Pullex Penetrans*, *Dermatoph'illus*, *Sarcosyll'a penetrans*, *Tick*, *Chig'gre*, *Chig'oe*, *Chig'go*, *Cheg're*, *Cheg'ok*, *Jig'ger*, (from (S.) *chigo*, *chiquito*, 'small.') A small insect in America and the Antilles, which gets under the epidermis, and excites great irritation.

CHIR, (*χειρ*.) *Manus*, ('hand.') See *Choir*.

CHIRAETA, *Gentiana chirayta*.

CHIRAGRA, *Che'ragra*, (*chir*, and *αγρα*, 'a seizure.') *Gout* in the hand.

CHIRAPOTHECA, (*chir*, and *αποθηκη*, 'a store, a closet.') *Arsenal*.

CHIRAPSIA, (*chir*, and *απειν*, 'to handle') *Friction*.

CHIRARTHRI'TIS, (*chir*, *αρθρον*, 'joint,' and *itis*, 'inflammation,') *Inflammation of the joints of the hand*.

CHIRAYITA, *Gentiana chirayta*.

CHIRAYTA, *Gentiana chirayta*.

CHIRETTA, *Gentiana chirayta*.

CHIRHEUMA, *Chirrhœma*.

CHIRIATER, (*chir*, and *ιατρος*, 'a physician') *Surgeon*.

CHIRIATRIA, *Surgery*.

CHIRIMOYA, *Anona tripetala*.

CHIRISIS, (*χειρ*is, from *chir*.) *Surgery*.

CHIRISMUS, (*χειρ*is, from *chir*.) *Surgery*.

CHIRIXIS, (*χειρ*is, from *chir*.) *Surgery*.

CHIRO, (*χειρ*, 'the hand.') In composition, hand.

CHIROCYRTO'SIS, (*chiro*, and *κυρτωσις*, 'crook-edness,') *Crookedness of the hand*.

CHIROMANCY, *Chiromanti'a*, *Cheiromanti'a*, (*χειρομαντεια*, from *chiro*, and *μαντεια*, 'divination,') *Vaticin'ium chiroman'ticum*, *Palm'etry*, (F.) *Chiromancie*. Art of divining by inspection of the hand.

CHIROMANIA, (*chiro*, and *mania*.) *Masturbation*.

CHIRONAX, (*χειρωναξ*, from *chir*.) *Surgeon*.

CHIRONIA, (after *Chiron*, *Χειρον*, the Centaur, who is said to have discovered its use.) A genus of plants. *Ord. Gentianacæ*. Also, *Tamus communis*.

CHIRONIA ANGELARIS, *Sabbatia*, *S. Angul'aris*, *Amer'ican Centaury*, *Rose'pink*, *Wild Sw'eory*, *Bitterbloom*, *Cent'ry*. Every part of this plant is a pure and strong bitter, which property is communicated alike to alcohol and water. It is used as a tonic and stomachic.

CHIRONIA CENTAURI'UM, *Centauri'um mi'num* seu *vilga'rè* seu *par'ven*, *Gentia'na centauri'na* seu *Gent'i'di*, *Erythræ'a Centauri'um*, *Hippocentaure'a centauri'um*, *Centaure'um*, *Centa'ria*, *Smaller* or *Les'ser Cent'ury*, *Europe'an Cent'ury*, (F.) *Centaure'e petite*, *Herbe au centaure* ou *à la fièvre* ou *à Chiron*. The tops of the lesser Centaury, *Centauri'ri Cac'u'mina*, are aromatic and tonic, and are sometimes employed as such.

CHIRONIA CHILEN'SIS, *Gentia'na Cachenta'huen*, *Erythræ'a Chilensis*, *Gentia'na Peruv'iana*, *Cachenta'huen*, *Chachinla'gua*, *Chancela'gua*. A very bitter plant, indigenous in Chili. It possesses the virtues of the *Chironææ*. Given in infusion—(3j, to water Oj.)

CHIRONI'UM, (*χειρωνειον*, from *χειρων*, 'bad, malignant,' [?]) An ulcer difficult of cure:—of a swollen, hard, and callous nature. Some have supposed the word to come from *Chiron*, the Centaur, who was unable to cure such ulcers.

CHIRONOMIA, *Cheironomia*.

CHIROPODALGIA, (*chiro*, *πους*, gen. *ποδ*is, 'foot,' and *αλγος*, 'pain,') *Acrodynia*.

CHIROPODIST, (F.) *Pédi'cour*, (*chiro*, and *πους*, gen. *ποδ*is, 'the foot,') One who treats diseases of the hands and feet, or rather whose profession it is to remove corns and bunyons.

CHIORRHŒMA, Chirrhœma.

CHIOSIS, (χειρῶσις, from χειρῶσις, 'to subdue.') Subactio.

CHIROTROPHOSIS, (chiro, and τροφῶσις, 'to turn.') Distortion of the hand.

CHIROTHER'CA, (chiro, and θῆκη, 'a sheath.') A bandage for the hand. A kind of bandage in which the fingers and hand are enveloped in spiral turns. When the whole hand and fingers are covered, it is called the *double or complete Chirotherca*, *Ch. complete*, *Vinctu'ra om'nibus dig'itis*; and when only a finger is covered, the *half or incomplete*, *Ch. incomplete*, *Vinctu'ra pro uno dig'ito*. See *Gantelot*.

CHIROTIB'IA, (chiro, and τριβῶ, 'I rub.') Friction with the hand. According to others, dexterity in an art.

CHIRRHEU'MA, gen. *Chirrhœu'matis*, *Chirrhœu'ma*, *Rheumatismus ma'nus*, (chir, and ῥευμα, 'flux.') Rheumatism of the hand.

CHIRURGEON, Surgeon.

CHIRURGI PHYSICI, see Surgeon.

CHIRURGIA, (χειρουργία, from chir, and ἔργον, 'work.') Surgery.—c. *Anaplastica*, *Morioplastice*—c. *Curtorum*, *Morioplastice*—c. *Infusoria*, *Infusion of Medicines*—c. *Militaris*, *Surgery, military*—c. *Transfusoria*, *Transfusion*.

CHIRURGICAL, Surgical.

CHIRURGICUS, Surgical.

CHIRURGIE, Surgery.—c. *Conservatrice*, *Surgery, salvation*—c. *Militaire*, *Surgery, military*.

CHIRURGIEN, Surgeon—c. *Consultant*, *Consulting Surgeon*—c. *Hernieur*, see *Hernial*.

CHIRURGIENNE. A female surgeon.

CHIRURGIQUE, Surgical.

CHIRURGUS, Surgeon.

CHIST. An Arabic word which signifies the sixth part. The sixth part of the *Congius* or gallon.

CHITON, (χιτών,) Tunic.

CHITONISCUS, (χιτωνισκος, dim. of Chiton.) Indusium.

CHITTICK'S NOSTRUM, see *Nostrum*.

CHI'UM VINUM. From χιός, *Chios*, the island where it was produced; *Chi'an wine*; used by the physicians of antiquity in cases of *defluxions* and *ophthalmiæ*.

CHLI'AROS, (χλῆρος, 'tepid.') A name anciently given to slight fevers in which the heat is not great.

CHLIAS'MA, gen. *Chlias'matis*, (χλῆσµα.) A tepid and moist fomentation.

CHLOAS'MA, gen. *Chloas'matis*, *Pityriasis versicolor*, *Mac'ula hepatic'a*, *Pan'nus hepaticus*, *Hepatic'izon*, *Pha'zē*, *Pha'cea*, *Pha'cus*, (F.) *Taches hépatiques*, *Chaleurs du foie*, *Éphélide scorbutique*, *Liver spots*, (χλῶς, 'a greenish-yellow colour, verdure.') A cutaneous affection, characterized by one or more broad, irregular-shaped patches, of a yellow or yellowish-brown colour, occurring most frequently on the front of the neck, breast, abdomen, and groins. The patches do not generally rise above the surface. There is usually some degree of itching. A cryptogamous growth has been found in it, to which M. Robin gives the name *Microsporion fur'fur*.

The causes are not very evident. Sulphur externally—in any and every form—generally removes it speedily. Should there be difficles, the external use of the remedy in baths or fumigations may succeed.

CHLOASMA ALBUM, *Achroma*.

CHLORA, Chlorine.

CHLORAS KALICUS DEPURATUS, *Potassæ murias hyperoxygenatus*.

CHLORASMA, (χλωρος, 'green.') Chlorosis.

CHLORE, Chlorine—c. *Liquide*, see *Chlorine*.

CHLORÉTHÉRIDE, *Chloroform*.

CHLORETUM CALCULÆ, *Calci chloridum*.

CHLORIASIS, (χλωρος, 'green.') Chlorosis.

CHLORIC ETHER, CONCENTRATED, see *Ether, chloric*—c. *Ether, strong*, see *Ether, chloric*.

CHLORIDE OF BROMINE, see *Bromine, chloride of*.

CHLO'RINE, (χλωρος, 'green.') *Chlorinum*, *Chlorin'ium*, *Chlorum*, *Chlo'ra*, *Halogen'ium*, *Oxy-muriatic Ac'id Gas*, *Oxygenated Muriatic Ac'id Gas*, *Dephlogisticated Marine Ac'id*, *Hal'ogene*, *Mu'rigene*, (F.) *Chlore*. So far as we know, this is an elementary substance. It is a greenish, yellow gas, of a strong suffocating smell, and disagreeable taste; incapable of maintaining combustion and respiration, and very soluble in water. One of its characteristics is, that of destroying, almost immediately, all vegetable and animal colours. It is employed in fumigations as a powerful disinfecting agent. A dilute solution, *A'qua seu Li'quor Chlo'ri*, *Chlorin'ii Li'quor*, *Chlorin'ei aqua*, *Aqua Chlorin'ii* (Ph. U. S., 1863), *Liquor Chlo'ri*, *Chlo'rine water*, *Solution of Chlo'rine*, (F.) *Chlore liquide*, has been administered internally, in certain cases of diarrhœa and chronic dysentery. A formula for its preparation, by disengaging chlorine from black oxide of manganese, by means of *muriatic acid*; and causing it to pass into water, has been admitted into the *Pharmacopœia of the United States for 1863*.

Immersion of the hands and arms in chlorine has often removed itch and other cutaneous affections. It has also been inhaled in a dilute state in the early stage of phthisis, but it is of doubtful efficacy, and is better adapted for chronic bronchitis.

CHLORINE, BISULPHURET OF, Sulphur, chloride of.

CHLOROÂNÆMIA, (χλωρος, 'green,' and *anæmia*.) Chlorosis.

CHLO'RODYNE. A secret preparation, proposed, in 1848, by a Dr. Browne of India; and said, by him, to consist of "perchloric acid with a new alkaloid." It has been given as an anodyne chiefly; but being a nostrum, is only worthy of mention for its notoriety.

CHLO'ROFORM, *Chloroform'um*, *Chloroform'ium*, (from *chlorine*, and *formyl*.) *Carbo'neum seu For'mylum chlorat'um*, *Superchlor'idum formyl'icum*, *Perchloride and Terchloride of For'myl*, *Chloroform'yl*, called also, but not correctly, *Terchloride of Carbon*, and *Chlo'rie ether*, *Æ'ther chlo'ricus*;—(F.) *Chloroforme*, *Chloréthérde*, is a colourless, oleaginous liquid, of a sweetish ethereal odour, hot, aromatic, and peculiar taste. The specific gravity of that of the Ph. U. S., *Chloroform'um vena'le*, is 1.45 to 1.49. It may be obtained by distilling from a mixture of chlorinated lime and alcohol,—rectifying the product by redistillation, first from a great excess of chlorinated lime, and afterwards from strong sulphuric acid. It has been used with advantage in asthma, and in diseases in which a grateful soothing agent is required. Dose, gtt. x to fʒss or more, diluted with water. It is largely prescribed, and with great success, as an anæsthetic agent in painful diseases; and to obtund sensibility in surgical operations and in parturition,—especially in the way of inhalation; but its use requires caution. The usual quantity with this view is from fʒss to fʒij. See *Anæsthetic*.

The *Pharmacopœia of the United States (1863)* has a *Chloroformum purifica'tum*, which is made by adding sulphuric acid to commercial chloroform; decanting the lighter liquid, and mixing it with stronger alcohol; adding carbonate of potassa, previously heated to redness, and rubbed, while warm, into powder. The mixture is, then, thoroughly agitated, and, by means of a water bath,

distilled to dryness. Its s. g. varies from 1.490 to 1.494.—s. g. (Ph. B. 1864) 1.496.

CHLOROFORM, TINCTURE OF, Ether, chloric.

CHLOROFORMIZATION, *Chloroformisatio*. The aggregate of anæsthetic phenomena occasioned by the inhalation of chloroform. To *chloroformize* a patient is to induce such phenomena.

CHLOROFORMUM, Chloroform—ch. Venale, see Chloroform.

CHLOROFORMYL, Chloroform.

CHLORO'MA, gen. *Chloro'matis*, (χλωρος, 'green,' and *oma*.) A peculiar form of tumour containing a yellow green substance, which takes the place, at times, of muscles and peritoneum.

CHLOROPHYLL, (χλωρος, 'green,' and φύλλον, 'a leaf.') *Fecula*, green.

CHLOROSIS, (χλωρος, 'green,' and *osis*.) *Pallidus Morbus*, *Fœdus Virginum color*, *Pallidus color virginæus*, *Pallor virginum*, *Icteric'ia alba*, *Icterus albus*, *Leucopathia*, *Morbus virginæus seu Parthenis'us*, *Parthenon'osis*, *Chloro-anæmia*, *Fœdi color'es*, *Dyspeps'ia chlorosis*, *Fœbris amot'ria*, *Cachex'ia seu Fœbris virginum*, *Fœbris alba*, *Anepithym'ia chlorosis*, *Chloras'ma*, *Chloros'ma*, *Chlori'asis*, *Citto'is*, *Green-sick'ness*, (F.) *Chlorose*, *Pâles-couleurs*, *Chloro-anémie*. A disease which affects young females, more particularly those who have not menstruated. It is characterized by a pale, lurid complexion, languor, listlessness, depraved appetite and digestion, palpitation, &c. The disease generally goes off on the occurrence of the menstrual flux; but sometimes it is long before this is established, and, at times, the catamenia are in much larger quantity than usual. To this last form M. Trousseau has given the name *chlorose hémorrhagique*.

The blood of chlorosis is generally thin, light-coloured and deficient in red corpuscles, and in iron,—*achalybhémie* of Piorry, and the clot is in less proportion to the serum than in health. On auscultation, a bellows' sound has been almost invariably detected over the heart, and a continuous blowing sound in the larger arteries, (especially the carotids and subclavians,) reinforced by each systole of the ventricle, and resembling the buzzing of a humming-top, the cooing of doves, the whistling of air through a key-hole, &c., (see *Bruit*.) Very similar sounds are heard in the arteries after copious hemorrhage: they seem, therefore, to coincide with enfeebled circulation.

Tonics—as iron—are usually required in the treatment,—the disease most commonly occurring in those in whom there is considerable torpor of the system.

CHLOROSIS ÆTHIOPUM, Chthonophagia—c. *Amatoria*, Hectic fever.—c. *Gigantea*, see *Polysarcia*.

CHLOROSMA, gen. *Chloros'matis*, (from χλωρος, 'green.') Chlorosis.

CHLOROTIC, *Chlorot'icus*, (F.) *Chlorotique*. Affected with chlorosis, or pertaining to chlorosis;—as a *chlorotic female*, *chlorotic symptoms*, &c.

CHLORUM, Chlorine.

CHLORURE DE CARBON, Chloroform—c. *d'Or*, Gold, Muriate of.

CHLORURETUM AMMONICUM, Ammoniac murias—c. *Oxidi Calcii*, Calcis chloridum.

CHOAK, (Sax. *aceocean*.) Cynanche trachealis—c. Wolf, Lyncanche.

CHOANA, (χωαν, 'a funnel.') Pelvis—c. *Cerebri*, Infundibulum of the brain.

CHOANÆ, pl. of Choana. The posterior nares.

CHOANE, Infundibulum.

CHOANORRHAGIA, (*choanæ*, and *ῥήγνμι*, 'to break forth.') See *Epistaxis*.

CHOAVA, *Coffea Arabica*.

CHOCOLAT, see Chocolate.

CHOCOLATA, Chocolate—c. *cum Osmazomâ*, see *Osmazome*—c. *Salutis*, see *Chocolate*—c. *Simplex*, see *Chocolate*.

CHOCOLATE, *Jac'alatt*, *Chocola'tum*, *Chocola'ta*, *Succola'ta*, *Succocolla'ta*. (Dr. Alston says, that this word is compounded from two East Indian words:—*choco*, 'sound,' and *atte*, 'water,' because of the noise made in its preparation [?] Another derivation is from *Choco*, a province in Mexico, where the tree abounds, and whence it was imported into Europe about 1520.) An alimentary paste prepared from the kernels of *Theobroma cacao* or *Cacao*, with sugar, and often aromatics. (See *Cacao*.) The chocolate thus simply prepared—as it is met with, indeed, in commerce—is called in France, *Chocolat de santé*, *Chocola'ta simplex seu Salu'tis*. It is not very easy of digestion.

The *chocolat à la vanille* contains three ounces of vanilla and two of cinnamon to twenty pounds of common chocolate. The addition of the aromatic renders it somewhat more digestible. Chocolates may likewise be medicated.

CHOCOLATE, OSMAZOME, see *Osmazome*.

CHOCOLATE ROOT, *Geum rivale*, G. Virginianum.

CHENICIS, (χοινικis, 'the nave in which the axle turns,') Trepan.

CHERADO'DES, (χοιρας, gen. χοιραι'ος, 'a swine,' 'serofula,' and *odes*.) Of the nature of serofula. Affected with serofula.

CHÆRAS, gen. *Chæ'radis*, (from χοιρας, 'a swine,') Serofula.

CHÆROCNE'S MUS, (*chærnus*, 'vulva,' and *κνησμος*, 'itching,') Itching of the vulva.

CHERUS, (χοιρας,) Vulva.

CHOKE CHERRY, *Prunus Virginiana*—c. Damp, Carbonic acid. See Hydrogen, carburetted.

CHOKES, Throat.

CHOLA, (χολη,) Bile.

CHOLEMIA, *Cholhæ'mia*, *Choli'hæ'mia*, (*chole*, and *αιμα*, 'blood,') A morbid state, in which bile exists in the blood. Jaundice.

CHOLAGO, Cholas.

CHOLAGOGUE, *Cholago'gus*, (χολαγωγus, from *chole*, and *αγω*, 'I expel,') *Felliducus*, *Bilit'icus*. The ancients give this name to cathartics, which were reputed to cause the flow of bile.

CHOLANSIS, (χολαντις, from χωλος, 'lame,') Cholosis.

CHOLAS, (χολας, plur. χολαδες,) *Chola'go*. The epigastric region. The intestines.

CHOLESMA, gen. *Cholasm'atis*, Cholosis.

CHOLE, (χολη,) Bile.

CHOLEC'CHYSIS, *Cholen'chysis*; (*chole*, and *εγχευσις*, 'effusion,') Effusion of bile.

CHOLECYST, (*chole*, and *κυστις*, 'bladder,') Gall-bladder.

CHOLECYSTECTASIA, (*cholecystis*, and *εκτασις*, 'dilatation,') Cholecysteurysma.

CHOLECYSTIS, Gall-bladder.

CHOLECYSTEURYSMA, gen. *Cholecysteurysm'atis*, *cholecystectasia*, (*chole*, *κυστις*, 'bladder,' and *ευρυσμα*, 'dilatation,') Dilatation of the gall-bladder.

CHOLECYSTITIS, *Inflamma'tio Vesicæ fell'æ*, *I. cyst'idis fell'æ*, *Cysti'tis fell'æ*, *Hepati'tis cyst'ica*, (*chole*, *κυστις*, 'bladder,' and *itis*), (F.) *Inflammation de la Vésicule du Fiel*, *Cholécystite*. Inflammation of the gall-bladder.

CHOL'EDOCH, *Choléd'ochus*, (χοληδόχος, from *chole*, and *δοχος*, 'containing or receiving,') (F.) *Cholédogue*. The *Duct'us choled'ochus* seu *hep'atocyst'icus*, *Duct'us commun'is choled'ochus*, (F.) *Conduit on Canal Cholédogue*, is the duct formed by the union of the hepatic and cystic ducts, which pours the hepatic and cystic bile into the duodenum.

CHOLEDOCHITIS, (*choledochus*, and *itis*;) Inflammation of the choledoch duct.

CHOLEDOGRAPHY, (*Choledogra'phia*, *Cholegraph'ia*, *Colograph'ia*, (*chole*, and *γραφειν*, 'to describe.')

A description of what relates to the bile and biliary organs.

CHOLEDOLOGY, (*Choledolog'ia*, *Cholelog'ia*, *Chololog'ia*, (*chole*, and *λογος*, 'a discourse.')

A treatise on the bile and biliary organs.

CHOLEDOQUE, *Choledoch*.

CHOLEGRAPHIA, *Choleodography*.

CHOLEHEMIA, (*chole*, and *αἷμα*, 'blood,') *Icterus*.

CHOLEIA, (*χολος*, 'lame,') *Claudication*.

CHOLELITHIA, (*chole*, and *λίθος*, 'a stone,')

Cysthepatolithiasis—*c. Icterus*, *Icterus*.

CHOLELITHIASIS, *Cysthepatolithiasis*.

CHOLELITHUS, see *Calculi*, *biliary*.

CHOLELOGIA, *Choleodology*.

CHOLEMESIA, (*Cholem'esis*, (*χολεμεια*, from *chole*, and *εμεσις*, 'vomiting,')

Vomiting of bile.

CHOLENCHYSIS, *Cholecchysis*.

CHOLEPYRA, (*chole*, and *πυρ*, 'fire, fever,')

Fever, bilious.

CHOLEPYRETUS, *Fever, bilious*.

CHOLEPYRRHIN, (*chole*, and *πυρρος*, 'of a golden colour,')

see *Bile*.

CHOL'ER, *Bile*. Anger was supposed to be produced by a superabundance of bile; hence the term *Choler* for anger.

CHOL'ERA, *Cholera morbus*, *Cholera nos'tras*

seu *vulgaris*, *Sporadic Cholera*, *Cholerrhag'ia*, *Pas'sio choleric'a*, *Fellif'lua pas'sio*, *Morbus fellif'luis*, *Hol'era*, *Bil'is flux'io*, (F.) *Choléra*, *Choladrée lymphatique*, *Hydrocholadrée*, *Choléra-morbus sporadicus*, *Ch. Européen*, *Trousse-galant*, (*χολερα* (*νοσος*), or from *chole*, and *ῥέω*, 'I flow,' or from *χολαίς*, 'intestines,' or *χολερα*, 'the gutter of a house to carry off the rain,')

The higher degrees have been called *Centrogangli'itis*, *Myelogangli'itis*, and *Hologangli'itis*. A disease characterized by anxiety, gripings, spasms in the legs and arms, and by vomiting and purging (generally bilious:) vomiting and purging are, indeed, the essential symptoms. The disease is most common in hot climates,—and in temperate climates, during summer. *Spasmodic*, *Asiat'ic*, *Malignant*, *Ind'ian*, *Epidemic*, *Pestilential*, *Convulsive Nervous*, *Eastern*, or *Oriental Cholera*. *Asphyx'ia pestilentialis*, *Pestilential asphyx'ia*, *Choleric Pestilence*, *Cholera orientalis* seu *Indica* seu *Epidemica* seu *Asphyx'ia*, *Typhus Bengalen'sis*, *Chol'ero-typhus*, *Ganglion'itis peripher'ica* et *medullaris*, *Hymenogangli'itis*, *Pantogangli'itis*, *Cholerrhæa lymphat'ica*, *Psorenter'ia*, *Achol'ia*, *Typhoid Fever of India*, *Hyperanthrax'is*, *Enterop'yra Asiatic'a*, *Trisplanchn'ia*, *Trisplanchn'itis*, *Hæmataporrh'ois*, *Hæmatorrh'ois*, *Morbus ory'zeus*, *Rice disease*, (because supposed by Tytler to be caused by damaged rice,) (F.) *Mort de Chien*, is frightful in the rapidity of its progress, the patient sometimes dying in a few hours from the first onset. In temperate climates, common cholera is not usually a disease of much consequence. It requires that the stomach and bowels should be cleared, and afterwards the irritation be allayed by full doses of opium. In malignant cholera, large doses of calomel, with opiates, form the great reliance of many practitioners. In the worst forms, however, it runs its course, unmodified by treatment.

CHOL'ERA EUROPÉEN, see *Cholera*—*c. Indian*, see *Cholera*—*c. Indica*, see *Cholera*.

CHOLERA INFAS'TUM, *Chol'eric Fever of Infants*; *Cholera of Infants*, (F.) *Cholera infantile*. A disease so termed by American physicians. It occurs, generally, in the middle states of the Union, in June or July, and continues during the hot weather, hence called the 'summer complaint.'

The chief symptoms are vomiting, purging of green

or yellow matter, slime or blood, attended with pain or uneasiness, and swelling of the abdomen, with some pyrexia, generally. The heat of the weather seems to be the predisposing, if not the exciting, cause. It is a fatal disease in towns; differing little, if at all, from what is vulgarly called the *Watery Gripes* in England. Clearing, gently, the alimentary canal, so as to remove the offending matter, and then exhibiting chalk mixture and laudanum, with counter-irritants, as spice cataplasms to the abdomen, is the most satisfactory plan of treatment.

CHOLERA, *MALIGNANT*, see *Cholera*—*c. Morbus*, *Cholera*—*c. Morbus Sporadicus*, see *Cholera*—*c. Nostras*, *Cholera morbus*—*c. Oriental*, see *Cholera*—*c. Pestilential*, see *Cholera*—*c. Poison*, see *Poison*—*c. Sec*, *Cholera sicca*.

CHOLERA SICCA, (F.) *Choléra sec*. *Pneumatosis* of the digestive passages in which there is a copious discharge of gas upwards and downwards.

CHOLERA, *SPASMODIC*, see *Cholera*—*c. Sporadic*, see *Cholera*.

CHOLERA VULGARIS, *Cholera morbus*.

CHOLERAIC, *Choleric*.

CHOLERAMANIA, *Cholermania*.

CHOLERAPHOBIA, (*cholera*, and *φοβος*, 'dread,')

Cholermania.

CHOLERAPHONIA, (*cholera*, and *φωνη*, 'voice,')

Voice, choleric.

CHOL'ERIC, (*χολερικός*,) *Chol'ericus*, (F.) *Cholérique*. Belonging to bile; also to cholera; *Cholera'ic*,—as the 'choleric or choleraic poison.'

The French use the term *Fèvre cholérique* for the fever accompanying cholera.

CHOLERIC FEVER OF INFANTS, *Cholera infantum*.

CHOLERIC TEMPERAMENT. The bilious temperament.

CHOL'ERINE, (F.) *Cholérine*, (dim. of *cholera*.) The first stage of epidemic cholera; also, the precursory symptoms of cholera.

CHOL'ERIQUE, *Choleric*.

CHOL'EROÏDE, *Choleraï'des*, (*cholera*, and *ειδός*, 'resemblance,')

Resembling cholera; as a 'choleroïd affection.'

CHOLEROMANIA, *Cholermania*, *Cholera-pho'bia*, *Cholero-pho'bia*, (*cholera*, and *μανία*,)

A dread of cholera to such extent that the individual fancies himself affected or threatened by it.

CHOLEROPHOBIA, (*cholera*, and *φοβος*, 'dread,')

Cholermania.

CHOLEROPHONE, (*cholera*, and *φωνη*, 'voice,')

Voice, choleric.

CHOLEROPROSPON, *Cholerop'ois*, (*cholera*, and *προσσωπον*, 'countenance,')

The facial expression of cholera.

CHOLEROPSIS, (*cholera*, and *οψις*, 'countenance,')

Choleroprosopon.

CHOLERTYPHUS, *Cholera*, (*spasmodic*.)

CHOLERRHAGIA, (*chole*, and *ῥήγνυμι*, 'to break forth,')

Cholera.

CHOLERRHÆA LYMPHATICA, (*chole*, and *ῥέω*, 'I flow,')

Cholera.

CHOLESTEATINOMA, (*chole*, *στέαρ*, 'suet,')

and *ομα*,) *Cholesteatoma*.

CHOLESTEATINE, *Cholesterin*.

CHOLESTEATOMA, gen. *Cholesteato'matis*, (*chole*, and *stéatoma*.) An encysted tumour, which presents upon the interior of the cyst several superimposed layers of cholesterol, which glisten like pearl. Hence it was named by Cruveilhier the *laminated na'reous fatty tumour*, and it is, sometimes, termed the *adipociriform*.

CHOL'ESTERIN, *Chol'esterine*, *Cholesteri'na*, *Cholosteri'na*, *Cholest'arine*, *Cholest'atine*, (*chole*, and *στέαρ*, 'suet,')

Ad'ipocire, *Chol'ine*. An inodorous, insipid substance, in white, shining scales: fusible and crystallizing, on cooling, in radiated fibres. Soluble in alcohol. It forms the crystalline part of certain biliary calculi, is con-

tained in neurine, and in various tumours, as well as in the seeds of many plants, olive oil, &c. See Adipocire.

CHOLESTERIN DISEASE, see Lardaceous.

CHOLEUMA, gen. *Choleum*'atis, (χολος, 'lame,')

Cholosis.

CHOLHÆMIA, (chole, and 'αἷμα, 'blood,')

Cholæmia.

CHOLIA, Cholosis.

CHOLICE'LE, (from χολος, 'lame,' and κλη, 'tumour,') A swelling formed by the bile morbidly accumulated in the gall-bladder.

CHOLICUS, (χολικός,) Biliary.

CHOLICYSTIECTASIE, (chole, κυστις, 'bladder,' and εκτασις, 'dilatation,') Turgescencia vesicæ felleæ.

CHOLICYSTITE, Cholecystitis.

CHOLIHÆMIA, (chole, and 'αἷμα, 'blood,')

Cholæmia.

CHOLINE, Cholesterin.

CHOLIUS, (χολη, 'bile,') Biliary.

CHOLOCATARRHOPHÊ, *Cholocatarrhophe'sis*, *Cholocatarrhophe'ia*, (chole, and καταρροφη, 'to sup up,') The absorption of bile.

CHOLODES, (χολώδης,) Biliary.

CHOLODIARRHŒA, see Diarrhœa.

CHOLOGRAPHIA, (chole, and γραφειν, 'to describe,') Choleodography.

CHOLOID, *Cholois'des*, (χολοειδης; from chole, and ειδος, 'resemblance,') Having the appearance of bile.

CHOLOLITHUS, (chole, and λιθος, 'a stone,') see Calculi, biliary.

CHOLO'MA, gen. *Cholo'matis*, (χωλος, 'lame, maimed,' and oma,) Distortion of a limb; incapacity of moving it. Also the act of limping, claudication.

CHOLOPLANIA, (chole, and πλανω, 'I lead astray,') Icterus.

CHOLOPOËT'IC, *Cholopoet'icus*, (χολοποιος, from chole, and ποιω, 'to make,') Relating or appertaining to the formation of bile.

CHOLORRHŒA, (chole, and ρεω, 'I flow,') An abnormous discharge of bile.

CHOL'OSSES, (from chole, or cholus,) Alibert includes, under this head, every morbid affection of the liver and spleen.

CHOLO'SIS, (χολος, 'lame,' and osis.) *Chololotes*, *Cholas'ma*, *Cholan'sis*, *Choleu'ma*, *Chol'ia*, *Claudica'tion*. Hippocrates employs this word particularly for a lameness of the hand, which renders a person one-handed, (F.) *manchet*. Also, icterus; bilious dyscrasy.

CHOLOSIS AMERICANA, Fever, yellow.

CHOLOSTERINA, Cholesterin.

CHOLOTES, (χολος, 'lame,') Cholosis.

CHOLOZEMIA FEBRILIS, (chole, and ζημια, 'detriment,') Fever, bilious.

CHOLUS, (χολος,) Bile.

CHONDRA'LGIA, (*chondrus*, and αλγος, 'a pain,') Pain in a cartilage.

CHONDRIFICA'TION, *Chondrifica'tio*, (*chondro*, and *facere*, 'to make,') Formation of, or conversion into cartilage.

CHON'DRIN, (χονδρος, 'cartilage,') (F.) *Gélatine des cartilages*. A variety of gelatin, obtained by boiling the cornea, the permanent cartilages, and the bones before they are ossified.

CHONDRI'TIS, *Inflammat'io Cartilag'inis*, (F.) *Chondrite*, (*chondrus*, and *itis*.) Inflammation of cartilage.

CHONDROCLASIS, (*chondrus*, and κλασις, 'fracture,') Fracture of a cartilage.

CHONDRODES, (*chondrus*, and *odes*.) Cartilaginous, *Chondroid*.

CHONDROGENES'IA, *Chondrogen'esia*, (*chondrus*, and γενεσις, 'formation,') Formation of cartilage. A morbid conversion of parts into cartilage.

CHONDROGLOS'SUS, (*chondrus*, and γλωσσα,

'the tongue,') A fasciculus of fleshy fibres, passing from the lesser cornu of the os hyoides to the tongue. It forms part of the hyoglossus.

CHONDROGRAPHY, *Chondrogra'phia*, (*chondrus*, and γραφη, 'a description,') A description of cartilages.

CHON'DROID, *Chondroi'des*, *Chondro'des*, (*chondrus*, and ειδος, 'resemblance,') Cartilagini-form. Resembling cartilage.

CHONDROL'OGY, *Chondrolog'ia*, (*chondrus*, and λογος, 'a discourse,') A treatise on cartilages.

CHONDRO'MA, gen. *Chondro'matis*, *Enchondro'ma*, *Osteochondrophyte*, (*chondrus*, and oma.) A cartilaginous growth proceeding from bones, including spina ventosa, osteosarcoma, &c., with many. The term, as well as *Enchondro'ma*, is also applied to growths, which, histologically, resemble cartilage. To a pouched cartilaginous growth the term *Cylindro'ma* has been given.

CHONDROPHARYNGE'US, (*chondrus*, and φάρυγξ, 'the pharynx,') Fibres of the muscular coat of the pharynx, which arise from the lesser cornua of the os hyoides, and form part of the *Constrictor medius*. See Constrictor pharyngis.

CHONDRO'SIS, (*chondrus*, and osis.) A morbid formation or condition of cartilage.

CHONDROSTERN'AL, *Chondrosterna'lis*, (*chondrus*, and στεινον, 'the breast-bone,') Having relation to the cartilages of the ribs and breast-bone,—as the *chondrosternal* ligament, articulations, &c.

CHONDROSYNDES'MUS, (*chondrus*, and συνδεσμος, 'a ligament,') Union of bones, by means of fibro-cartilage. Synchondrosis.

CHONDROT'OMY, *Chondrotom'ia*, (*chondrus*, and τεμνειν, 'to cut,') Dissection of cartilages.

CHONDROXI'PHOID, *Chondroxiphoi'des*, from χονδρος, 'a cartilage,' and xiphoid, the cartilage so called. Having relation to the xiphoid cartilage, as '*Chondroxiphoid lig'ament*.'

CHOND'RUS, (χονδρος,) Cartilage. *The Xiphoid cartilage*, in particular.

CHOND'RUS, (χονδρος,) *Alica*, *Fucus crispus*—c. *Crispus*, *Fucus crispus*—c. *Polymorphus*, *Fucus crispus*.

CHON'US, (χωνος,) Infundibulum.

CHOPINE, *Chopi'na*, *Cheopi'na*. A measure, which contains 16 ounces. A pint measure.

CHORA, (χωρα, 'region,') Any void space. Also, the orbit of the eye.

CHORD, TESTICULAR, Spermatie chord.

CHOR'DA, *Cor'da*, (χορδη, 'a gut,' and hence, a string made of gut.) A cord, (F.) *corde*. It has several meanings:—as a *tendon*, and the intestines (*Chordæ*). Paracelsus calls the genital organs *Chordæ*. *Chorda seu Funic'ulus Tym'pani*, a branch of the 7th pair of nerves, according to some; of the 5th pair, according to others, which passes through the tympanum, (F.) *Nerv Tympanique*—(Ch.) *Corde du Tambour ou du Tympan*. *Chorda Willis'ii*, *Trabec'ula Willis'ii*, the small fibres which cross the sinuses of the dura mater; so called from their first describer, Willis. *Chorda Tendin'ea*; the tendinous and cord-like substances which connect the *columnæ carnea*e of the ventricles of the heart to the valves, &c.

CHORDA, Intestine.

CHORDA DORSA'LIS, *Dor'sal cord*. The axis, around which, according to Von Baer, the first parts of the fœtus are formed,—the rudiment of the future vertebral column.

CHORDA HIPPOCRATIS, *Achillis Tendo*—c. *Longitudinalis Corporis Callosi*, see *Raphe*—c. *Penis*, *Chordee*—c. *Tympani*, see *Chorda*—c. *Veneris*, *Chordee*.

CHORDÆ, *Chordee*—c. *Ferreri*, *Cords*, vocal—c. *Longitudinales Lancisii*, see *Raphe*—c. *Tendinæ*, see *Chorda*—c. *Transversales Willisii*, see *Corpus callosum*—c. *Vocales*, *Cords*, vocal—c. *Willisii*, see *Chorda*.

CHORDAP'SUS, (*χορδῶν*, 'intestine,' and *ἀρτεν*, 'to tie,') *Intususcep'tio*. Constriction or twisting of the intestines. Also, *Colica Ileus*. See *Enteritis* and *Ileus*.

CHORDEE', *Cordee'*, ((*F.*) *corder*, 'to bind with a cord,') *Chor'da Ven'eri's* seu *Pe'nis*, *Phal-lanculo'sis*, *Phallocamp'sis*, *Chor'dæ*, *Chordureth-ris'tis*. A painful affection of the penis, attending gonorrhœa, and happening chiefly at night. It occurs when the violence of the inflammation is such, that the urethra cannot expand as much as the corpora cavernosa during erection, and consequently the glans seems painfully drawn downwards, *Phal-lanculo'sis*. The application of cold water topically, and the administration of opium internally, palliate it.

CHORDURETHRITIS, (*chordee* and *urethri-tis*.) Gonorrhœa cordata.

CHORE'A, (*χορεία*, 'a dance,') *Ch. Sancti Vi'ti* seu *Sancti Modes'ti* seu *gesticulato'ria*, *Syn'clonus saltans*, *Tripudia'tio spas'tica*, sometimes called *Scelotyrbê*, *S. Tarantismus*, *Vi'ti Saltus*, *Saltat'io SANC'TI Vi'ti*, *Ballis'mus*, *Convul'sio habitu'al-is*, *Siph'ita pra'va*, *Syn'clonus Chore'a*, *Myotyrbê*, *Chorema'nia*, *Chorema'nia*, *Chorema'nia*, *Orchestra'ma'nia*, *Epilep'sia Saltato'ria*, *Paraly'sis vac'u-illaus*, (S. H. Dickson), *Mor'bus Saltato'rius* seu *gesticulato'rius* seu *Lasci'vus*, *Hieran'osus*, *St. Vi'tus's Dance*, *Viper's Dance*, because its movements resemble dancing; (*F.*) *Chorée*, *Danse de SAINT WITT*, *Danse de St. GUY*. The characteristics are:—irregular and involuntary motions of one or more limbs, and of the face and trunk. It is a disease which usually occurs before puberty; and is generally connected with torpor of the system, and of the digestive organs in particular. Its duration is long, but it is usually devoid of danger; although frequently but little under the control of medicine. The spasms do not continue during sleep. The indications of treatment are: to strengthen the general system, and stimulate the intestinal canal. Purgatives, once or twice a week, and chalybeates, with appropriate regimen, will fulfil these.

CHOREA, **CHRONIC**, see *Tic*.

CHOREA, **ELEC'TRIC**, *Convul'sio cer'ebral'ty'phus*, *Myel'tis convuls'ra*. An affection, seen in certain districts of Lombardy, which is so called in consequence of the similarity between the convulsive shock of the limbs and the shocks caused by the action of the induction coil. It is very fatal.

CHOREA FACIEI, *Tic*—*c. Gesticulatoria*, *Chorea*—*c. Partial*, see *Tic*—*c. Sancti Modesti*, *Chorea*—*c. Sancti Valentini*, *Tarantismus*—*c. Sancti Viti*, *Chorea*.

CHORÉE, *Chorea*.

CHORE'GIA, (*χορηγία*, from *χορος*, 'a troop of dancers and singers,' and *αγω*, 'I lead,') Hippocrates uses this term for the whole of the apparatus necessary for a physician or surgeon.

CHORE'IC, *Chore'icus*, (*F.*) *Choréique*, (from *chorea*.) Relating or appertaining to, or having the characters of, *chorea*. One affected with *chorea*.

CHOREMANIA, (*χορεία*, 'a dance,' and *μανία*, 'mania,') *Chorea*.

CHORTODEI'TIS, *Chorioidei'tis*, *Chorioidei'tis*, *Chorioidei'tis*, (*F.*) *Chorioïdite*, (from *choroïdes*, and *itis*.) Inflammation of the choroid coat of the eye.

CHORTODES, *Choroid*.

CHORIOIDEA, *Choroid*.

CHORIOIDES, *Choroid*.

CHOR'ION, (*χοριον*, 'skin,' from *χορειν*, 'to contain,' (include); [*?*]) *Comis'sia Fe'tis*. A skin, transparent membrane, formerly confounded with the decidua, which surrounds the fœtus in utero on every side, and appears to be developed

from nucleated cells formed in the Fallopian tube. Some histologists, however, consider, that it exists in the ovary. The general opinion is, that it is formed as above described; and perhaps, also, from the zona pellucida, which disappears in the tube. In the uterus, villous prolongations are formed on its surface, which have given it, with more recent writers, the name 'shaggy chorion.' These villi are probably the agents of the absorption of nutritive matter furnished from the lining membrane of the uterus.

By many anatomists, the chorion is considered to be formed of two layers; the outer, called by Burdach *Exocho'rion*; the inner, *Endocho'rion*. By others, the distinction of laminae is denied. It is exterior to the amnion.

CHORION also means the true skin. See *CUTIS*. By some, the Zona pellucida is so called.

CHORION FUNGOSUM, *Decidua*—*c. Reticulatum*, *Decidua*—*c. Shaggy*, see *Chorion* and *Decidua*—*c. Spongiosum*, see *Decidua*—*c. Spongy*, see *Decidua*—*c. Tomentosum*, see *Decidua*.

CHORIONITIS, (*chorion*, and *itis*), *Cytitis*.

CHOR'OID, *Choroï'deus*, *Choroï'des*, *Choroï'des*, *Choroï'des*, (*χοροΐδης*, from *χοριον*, 'the chorion,' 'the skin,' *corium*, and *αἶδος*, 'resemblance,') Several parts are so called, which resemble the corium, in the multitude of their vessels.

CHOROID MUSCLE, *Ciliary muscle*.

CHOROID PLEX'US, *Plex'us choroï'des* seu *choroï'deus* seu *reticulat'us*, *Ver'mes cer'ebr'i*. Two membranous and vascular duplicatures of the pia mater, situate in the lateral ventricles. They are fixed to the *Tela choroïdea* by one edge, and are loose and floating at the other.

CHOROIDEA TE'LA, *Plex'us choroï'deus me'dius*, *Velum interpos'itum*, (*F.*) *Toile choroïdienne*. A kind of vasculo-membranous prolongation of the pia mater, which lines the lower surface of the fornix united with the corpus callosum. It is stretched above the third ventricle, and covers the posterior commissure and corpora quadrigemina. Anteriorly, the tela choroïdea is continuous with the plexus choroïdes.

CHOROIDEA seu **CHOROIDEA TU'NICA**, *Ch. Membrana*, or simply the *Chor'oid*, *Tu'nica vasculo'sa* seu *aciniform'is* seu *rhagoïdes Oculi*, (*F.*) *Membrana choroïde*, *Choroïde*. A thin membrane, of a very dark colour, which lines the sclerótica internally. The part behind the iris is called *U'lea*. It is situate between the sclerótica and retina, has an opening, posteriorly, for the passage of the optic nerve; and terminates, anteriorly, at the great circumference of the iris, where it is continuous with the ciliary processes. Its outer surface is pretty intimately attached to the sclerótica by bloodvessels and nerves, as well as by a delicate brown tissue, which has been called *Lam'ina fus'ca*.

According to Ruysch, the choroid consists of two layers, to the innermost of which his son gave the name *Tu'nica* seu *Membrana Ruyschiana*, *M. choriocapilla'ris*, (*F.*) *Membrana Ruyschienne*. The internal surface of the membrane is covered with a dark pigment, consisting of several layers of pigment cells, called *Pigmentum nigrum*, *Stratum pigmenti*, *Ophthalmochroï'tes*, *Æthiops animal*, *Mélaine*, *Mélanine*, *Matière ou Principe de la Mélanose*, (*F.*) *Enduit choroïdien*. Its use seems to be, to absorb the rays of light after they have traversed the retina.

CHOROIDEA VE'NE, *Ve'næ Gale'ni*, (*F.*) *Veines choroïdiennes*. Two veins, that creep along the tela choroïdea; into which almost all those of the lateral ventricles, of the upper part of the cerebellum, of the pineal gland, and the corpora quadrigemina open. The *Ve'næ Galeni* open into the *sinus quartus* or *fourth sinus*.

CHOROÏDE, CEINTURE BLANCHE DE LA, Ciliary ligament—*c. Commissure de la,* Ciliary ligament.

CHOROÏDES, Choroid.

CHOROÏDEUS, Choroid.

CHOROÏDEITIS, Chorioiditis.

CHOROÏDORETINI'TIS. Inflammation of the choroid and retina.

CHOROMANIA, (χορος, 'a dance,' and μανια, 'mania.) Choreia.

CHORTASTHMA, (χορτος, 'hay.) Fever, hay.

CHOSSES CONTRE NATURE, Res contra naturam—*c. Naturelles,* Res naturales—*c. non Naturelles,* Res non naturales.

CHOSIS, (χωσις, from χωω, 'I heap up.') Are-natio.

CHOU, (from caulis, 'a potherb,' especially Kale or Colewort,) Brassica—*c. Cabus,* Brassica capitata—*c. Croûte,* Sauer Kraut—*c. Fleur,* Brassica Florida, Cauliflower exorescence—*c. Marin,* Convolvulus soldanella—*c. Navet,* Brassica rapa—*c. Pommé,* Brassica capitata—*c. Potager,* Brassica—*c. Roquette,* Brassica eruca.

CHOW'DER, a favorite New England dish, made of fish, pork, onions, and biscuit, stewed together. In Devonshire, England, *chowder* is a fishseller. In the west of England *chowder beer* is a liquor made by boiling black spruce in water, and mixing it with molasses.

CHREMA, gen. *Chrem'matis,* (χρεμμα, from χρεμπτομαι, 'to spit up.') Sputum.

CHREMPSIS, (χρεμψις,) Exspuition.

CHRISIS, (χρσις, from χρωω, 'I anoint.') The action of anointing. Inunction.

CHRISMA, gen. *Chrim'atis,* (χρσμα,) The act of anointing. The salve or liniment used, *Chriem.*

CHRIS'TIMA'NUS, ('hand of Christ.') Troches prepared from refined sugar boiled in rose-water with or without prepared pearls.

CHRISTOPHER, HERB, Actaea spicata.

CHRISTOPHORIANA SPICATA, Actaea spicata.

CHRIS'TUS, (χριστος, from χρωω, 'I anoint.') A liniment or ointment.

CHRO'A, (χροα,) *Chroc'a,* *Chro'ma.* Colour in general. The surface of the body. The skin.

CHROMA, gen. *Chro'matis,* (χρωμα,) Chroa, (colour.)

CHROMATISME, (from chroma,) Aberration of refrangibility.

CHROMATOG'ENOUS, (chroma, and γεννωω, 'I make.')

CHROMATOG'ENOUS APPARA'TUS. A supposititious glandular apparatus for producing the colouring matter of the skin, composed of a glandular or secreting parenchyma, situate a little below the papillæ, and presenting special excretory ducts, which pour out the colouring matter on the surface of the true skin.

CHROMATOPHO'BIA, (chroma, and φοβος, 'dread.') Morbid sensibility to certain colours.

CHROMATOPSEUDOPSIA, (chroma, ψευδης, 'false,' and οψις, 'vision.') Achromatopsia.

CHROMATOPSIA, Chromopsia.

CHROMATU'RIA, (chroma, and ουρον, 'urine.') Discharge of urine of an unnatural colour.

CHRO'MIC ACID, *Ac'idum Chro'micum,* (F.) *Acide Chromique.* Obtained by crystallization from a mixture of *bichromate of potassa,* and oil of *vitriol.* It has been used as an escharotic in external hemorrhoids.

CHROMIDRO'SIS, (chroma, and ιδρως, 'sweat.') Abnormal coloration of the perspiratory secretion.

CHROMOP'SIA, *Chromop'ia,* *Chromatop'sia,* *Chrotop'sia,* *Chrup'sia,* *V'isus coloratus,* *Suffusio colorans,* *Chroôp'sy,* *Chromop'ey,* *Irides'cent vision,* (chroma, and οψις, 'vision.') A state of vision in

which a coloured impression, as of the rainbow, is made on the retina. Said to be occasionally observed in jaundice.

CHRON'IC, (χρονικος, from χρονος, 'time.) *Chron'icus,* *Chro'nus,* *Polychro'nus,* *Inveteratus,* *Antiquus,* (F.) *Chronique.* Of long duration.

CHRONIC DISEASES, *Mor'bi chron'ici,* *Macronos'ia,* *Macro'sia,* *Chronon'osi,* *Chronopath'ia,* (F.) *Maladies Chroniques,* are those whose duration is long, or whose symptoms proceed slowly. The antithesis to *chronic* is *acute.*

CHRONIC'ITY, *Chronic'itas,* (F.) *Chronicité.* The state of a chronic malady.

CHRONIQUE, Chronic.

CHRONIUS, (χρονιος,) Chronic.

CHRONOGYNIA, (χρονος, 'a time,' and γυνη, 'a woman.') Menstruation.

CHRONOHÉPATITE, Hepatitis, chronic.

CHRONONÉPHRITE, Nephritis, chronic.

CHRONONOSI, (χρονος, and νοσος, 'disease.') Chronic diseases.

CHRONOPATHIÆ, (χρονος, 'time,' and παθος, 'affection.') Chronic diseases.

CHRONOTHERMAL, (χρονος, 'time,' and θερμη, 'heat.) Relating to time and temperature. An epithet given by Dr. Samuel Dickson, to a fanciful 'system' which maintains, that there can be no increase or diminution of temperature without motion; no motion without time; that motion consists in attraction and repulsion; that attraction and repulsion are peculiar to electric action; and hence, that medicines must change the motions of the system, and be electrical in their operation.

CHROOPSY, (χροα, 'colour,' and οψις, 'vision.') Chromopsia.

CHROTIC, (from χρωω, 'the skin,) Diseases of the skin.

CHROTOSIA, (χρωω, gen. χρωωτος, 'colour,' and οψις, 'vision.') Chromopsia.

CHRUSIA, (χροα, 'colour,' and οψις, 'vision.') Chromopsia.

CHRYSA FIBRAUREA, (χρυσους, 'of a golden colour,') Coptis.

CHRYSALEA, Nitro-muriatic acid.

CHRYSANthemUM, (chryso, and ανθεμον, 'a flower.') *Calendula officinalis.*

CHRYSANthemUM LEUCANthemUM. The *Three eye da'isy,* *Da'isy,* *White'weed,* *Gold'ens,* *Maud'linwort,* *Bell'is ma'jor seu pratensis,* *Bupthal'mum ma'jus,* *Leucan'themum vulga'rè seu Bellidii'des,* *Matricaria,* *Consol'ida me'dia,* *Oculus Bo'vis,* (F.) *Chrysanthème,* *Chrysène,* *Grand Marguerite des prés.* Ord. Compositæ. The flowers and herb are slightly acid; and were once used in pulmonary diseases.

CHRYSANthemUM PARTHENIUM, *Matricaria parthenium.*

CHRY'SE, (χρυσος, 'golden.') The name of an ancient yellow plaster, composed of thus, alum, lead, colophony, resin, oil, and orpiment, boiled in vinegar.

CHRY'SENE, *Chrysanthemum leucanthemum.*

CHRY'SITIS, (χρυσιτης, 'golden,') see *Plumbi oxidum semivitreum.*

CHRY'SO, (χρυσος, 'gold,') in composition, gold.

CHRYSOBALANUS GALENI, (χρυσοβαλαος, from *chryso,* and *balanos,* 'a nut.') See *Myristica moschata.*

CHRYSOCHALCUS, (chryso, and χαλκος, 'copper.') Brass.

CHRYSOCOLLA, (χρυσocolλα, from *chryso,* and *κολλα,* 'a glue or cement,') Borax.

CHRYSOCOMA, (χρυσοκομη, from *chryso,* and *κομη,* 'hair,') *Achillea Millefolium.*

CHRYSO LACHANUM, (χρυσολαχανον, from *chryso,* and *λαχανον,* 'a potherb,') *Chenopodium bonus Henricus.*

CHRYSOLITHUS, *Chrysolite*, (*chryso*, and λιθος, 'stone.') A precious stone, of a golden colour, regarded by the ancients as cardiac, cephalic, &c.

CHRYSOMELIA, (*chryso*, and μελον, 'an apple.') see *Citrus aurantium*.

CHRYSOPHYLLUM BURANHEM, *Monesia*.

CHRYSOPHYLLUM CAINITO, (*chryso*, and φυλλον, 'a leaf.') *Cainito*, *Sideroxylon*, *Broad-leaved Star-apple*. A tree of the Antilles, *Ord.* Sapotaceæ, which produces one of the best fruits of the country. There are several varieties of it.

CHRYSOPHYLLUM GLYCIPHLEUM, *Monesia*.

CHRYSOPUS, (from χρυσος, 'of a golden colour.') *Cambogia*.

CHRYSOS, (χρυσος,) *Gold*.

CHRYSOSPERMUM, (*chryso*, and σπερμα, 'seed,') *Sempervivum tectorum*.

CHTHONOPHAGIA, *Ch. Anémica*, *Cuchec'ia Africa'na*, *Malac'ia seu Pi'ca Africano'rum*, *Leucophlegma'tia seu Chloro'sis Ethio'pum*, *Dirt-eat*, (*F.*) *Mal d'Estomac*, (χθων, 'earth,' and φαγω, 'I eat.') A disorder of the nutritive functions observed amongst the negroes of the South, and of the West Indies, in which there is an irresistible desire to eat dirt. It is accompanied by most of the signs of chlorosis.

The Ottomans on the Orinoco, and the natives of portions of the Hudson's Bay territory, are *Chthonophagi* or 'earth eaters.' The earth is a form of steatite. See *Geophagism*.

CHUS, (χους, χους.) A liquid measure amongst the Greeks, answering to the *Congius* of the Romans, and containing six sextarii, or twelve Attic cotylæ, or nine pints.

CHUMOSIS, *Chemosis*.

CHURNMILK, *Buttermilk*.

CHURRUS, see *Bangue*.

CHUTE, *Prolapsus*—*c. du Fondement*, *Proctocoele*—*c. de la Matrice*, *Procidencia uteri*—*c. des Oeufs*, see *Parturition*—*c. du Rectum*, *Proctocoele*.

CHYLAIRE, *Chylous*.

CHYLAR, *Chylous*.

CHYLARION, (χυλαριον,) see *Chyle*.

CHYLE, *Chylus*, *Succus nutritivus*, (χυλος, 'juice,' from χυω, 'I pour.') The word, in Hippocrates, means *Tisane* or *Decoction of barley*, *Chylarion*. Galen first used it in its present sense;—i.e. for a nutritive fluid, extracted by intestinal absorption, from food which has been subjected to the action of the digestive organs. It is of a whitish appearance; and is formed from the chyme in the duodenum, and the rest of the small intestines, by the chyliferous vessels, which arise at the mucous surface of the intestine. Along these it passes through the mesenteric glands to the thoracic duct, and is finally poured into the left subclavian. It is composed, like the blood, of a fluid—*liquor chylus*—and of *chyle corpuscles* or *globules*, the average size of which is about 1-4600th of an inch. They, who regard the chyle to be nothing more than fatty matter of the food, apply the term to the emulsified oil in the intestinal canal even before it is taken up by the chyliferous vessels. See *Chyme*.

CHYLE CORPUSCLES, see *Chyle*.

CHYLEUX, *Chylous*.

CHYLIFEROUS, *Chylifer*, *Chyliferus*, *Chyliferus*, (*F.*) *Chylifère*, (*chylus*, and *ferre*, 'to carry,') *Chyle-bearing*.

CHYLIFEROUS VESSELS, *Vasa Chylifera* seu *Chylof'era*, *Viæ chylif'ere*, *Ve'næ lact'eres*, *Va'sa lact'ea*, *Lact'ea*, (*F.*) *Vaisseaux chylifères*, *V. Lactés*. Vessels which convey the chyle from the intestines to the thoracic duct. They have also been called *galactoph'orous vessels*.

CHYLIFICATION, *Chylifica'tio*, *Chylo'sis*, *Chylopoë'sis*, *Præparatio chyli*, (*chylus*, and *facere*, 'to make,') Formation of chyle during the digestive processes. It has, at times, been ap-

plied to the changes produced on the aliment in the small intestine; but is now restricted to the act of forming chyle at the radicles of the chyliferous vessels.

CHYLINE, *Cyclamen*.

CHYLISMA, gen. *Chylism'atie*, (χυλισμα, from χυλιζω, 'to press out juice,') *Succus expressus*.

CHYLISMUS, (χυλισμος.) The act of expressing the juice of vegetables, &c.

CHYLOCYSTIS, (*chylus*, and κυστις, 'bladder,') *Receptaculum chyli*.

CHYLODES, (χυλωδης, from *chylus*, and *odes*,) *Chylous*.

CHYLODIABETES, *Chyluria*.

CHYLODIARRHŒA, *Cœliac flux*.

CHYLODOCHIUM, (*chylus*, and δεχομαι, 'to receive,') *Receptaculum chyli*.

CHYLOGRAPHY, (*chylus*, and γραφη, 'a description,') A description of the anatomy, &c., of the chyliferous vessels.

CHYLOPOËTIC, *Chylopoë'ticus*, *Chylopoëus*. Relating to or connected with the formation of chyle. Chiefly applied to the organs immediately concerned in it; as the stomach, intestines, omenta, and mesentery. *Assist'ant Chylopoetic*:—applied to viscera which aid in the formation of chyle, as the liver and pancreas.

CHYLOPOIESIS, (*chylus*, and ποιω, 'I make,') *Chylification*.

CHYLORRHŒA, (*chylus*, and ρεω, 'I flow,') *Cœliac flux*—*c. Pectoris*, *Chylorhœax*—*c. Renalis*, *Chyluria*—*c. Urinalis*, *Chyluria*.

CHYLOSIS, (χυλωσις,) *Chylification*.

CHYLOSTAGMA (gen. *Chylostag'matis*) **DIA-PHORETICUM MINDERERI**, (from χυλος, 'juice,' and σταζω, 'I distil'.) A compound prepared by distilling the *theriac* of Andromachus, the *mithridate* of Damocrates and other alexipharmics, &c. It is nearly the same preparation as the *Aqua Theriacalis Bezoardica*.

CHYLOTHORAX, gen. *Chylothora'cia*, *Pleurorrhœa chylo'sa*, *Chylorrhœa Pectoris*, *Hydrothorax chylo'sus*; (*chylus*, and θωραξ, 'the chest,') Effusion of chyle into the chest, owing to the rupture of a chyliferous vessel.

CHYLOUS, *Chylar*, *Chylo'sus* seu *Chyla'ris*, *Chylo'des*, (*F.*) *Chyleux*, *Chytaire*. Relating to the chyle; or having some analogy to that fluid.

CHYLURIA, *Diabètes lacteus* seu *chylo'sus*, *Chylodiabètes*, *Galacturia*, *Fluxus cœliacus per Renes*, *Pyluria lactea* seu *Chylo'sa*, *Calicula urinæ* seu *renalis*, *Chylorrhœa urinæ* seu *renalis*, (*chylus*, and ουρον, 'urine,') (*F.*) *Diabète chyleux*. A discharge of milky urine, without any apparent lesion of the kidneys or bladder.

CHYLUS, (χυλος,) *Chyle*, *Decoction*, *Succus*.

CHYME, *Chymus*, (χυμος, 'juice,' from χυω, 'I pour,') The pulp, formed by the food, mixed with the supra-diaphragmatic and gastric secretions, after it has been for some time in the stomach. In this state it continues until it reaches the biliary and pancreatic ducts, which open into the duodenum, where the conversion into chyle occurs, which is absorbed by the chyliferous vessels,—the excrementitious portion of the food traversing the large intestine to be evacuated *per anum*. Castelli asserts, that *Chyme* and *Chyle* were used in an inverse sense by the ancients to that accepted at present.

CHYMI, (χυμοι,) *Humours*.

CHYMIA, *Chymistry*—*c. Organica*, *Chymistry*, *organico*—*c. Pharmaceutica*, see *Chymistry*.

CHYMIAT'ER, *Chimia'ter*, *Chemia'ter*, (*chymia*, and ιατρος, 'a physician,') *Iatrochymicus*. A chemical physician.

CHYMIATRIA, *Chimiatri'a*, *Chemiatri'a*, *Iatrochemi'a*, *Medici'na spagyrica*, *Ars Chymiatri'ca*, (*F.*) *Chimiatrie*, *Chimisme* (of some,) (from *chymia*, and *ιατρεια*, 'cure,') The art of curing by chemical means.

CHYMICAL, *Chem'ical*, *Chem'icus*, *Chemo'ticus*. A medicine formed by the aid of chymistry, in contradistinction to *Galenic*.

CHYMICOHISTOLOGY, *Chym'icohistolog'ia*, *Chem'icohistolog'gy*. The doctrine of the organic chemistry and morphology of tissues.

CHYMICOPHANTA, (*chymia*, and *φαντα*, 'to manifest,') *Chymist*.

CHYMICUS, *Chymical*, *Chymist*.

CHYMIE, *Chymistry*.

CHYMIFICATION, *Chymified'tio*, *Chymo'sis*, (*chymus*, and *facere*, 'to make,') Formation of chyme.

CHYMISM, *Chymism'us*, *Chemism'us*, (F.) *Chimisme*, (from *chymia*.) The abuse of chymistry in its application to the science of organized bodies. By some used synonymously with *Chymiat'ia*. Also, the aggregate of chemical actions appertaining to a function—as the *chemismus* of respiration.

CHYMIST, *Chem'ist*, *Chem'icus*, *Chymicophan'ta*, *Chym'icus*, (F.) *Chimiste* ou *Chymiste*. One acquainted with chymistry. In Great Britain it has, also, the signification of "one who sells chemicals."

CHYMISTE, *Chymist*.

CHYMISTRY, *Chem'istry*, *Chem'ia*, *Chym'ia*, *Chim'ia*, *Chemue'ticē*, *Chemo'ticē*, *Philosoph'ia per ignem*, *Spagyr'ia*, *Pyrotech'nia*, *Pyrosoph'ia*, *Ars hermet'ica seu mago'rum seu separato'ria seu spagyr'ica*, *Archima'gia*, (*χυμος*, 'juice,' or from Arab. *chemā*, 'a secret,') (F.) *Chimie* ou *Chymie*. A branch of the natural sciences, whose object is to investigate the nature and properties of bodies, simple and compound, inorganic and organized; and to study the force or power, by virtue of which every combination is effected. It investigates the action between the integrant molecules or atoms of bodies.

Organ'ic Chemistry, *Chym'ia organ'ica*, *Organochem'ia*, is the chymistry of organized substances, —animal and vegetable. *Animal Chym'istry*, *Zoöchem'y* or *Zoöch'ymy*, *Zoöchem'ia*, is the chymistry of substances afforded by the dead or living animal body. This branch of chymistry has been farther subdivided into *physiolog'ical*, when it considers the changes produced in organized bodies in health,—*patholog'ical*, when it regards those produced by organic or other diseases. *Anthropoch'ymy*, *Anthropochem'ia*, is the chymistry of the human body. Chymistry is called *Therapeut'ical* or *Pharmaceut'ical*, *Pharmacochym'ia*, *Chym'ia pharmaceut'ica*, when it is engaged in the analysis of simple medicines; in improving the prescribing and preparing of chemical and Galenic medicines; in the means of preparing them, and detecting adulterations, &c. *Hygien'ic Chym'istry* is that which is applied to the means of rendering habitations healthy, of analyzing the air we breathe, preventing the occurrence of disease, pointing out healthy aliments, and appreciating the influence of professions, &c., on the health of man. All these different subdivisions, with vegetable chymistry, *Phytochym'istry*, are, at times, included under the head of *Med'ical Chym'istry*; at others, the term comprehends only the *An'im'al*, *Veget'able*, and *Pharmaceut'ical* subdivisions. *Vital Chemistry*, *Biochym'ia*, is that which is exerted under the influence of vitality. *Histochem'y*, *Histochem'ia*, *Histochem'istry*, is the chemistry of the tissues, and *Phlegmatochem'y*, that of the animal humours.

A knowledge of chemistry is of great importance to the physician. Many of the functions are of a chemical nature: many diseases require a chemical mode of treatment; and, without an acquaintance with it, two or more substances might be given in combination, which, by forming a chemical union, might give rise to other

compounds, possessing very different virtues from the components taken singly, and thus the prescriber be disappointed in the results.

CHYMISTRY, **ANIMAL**, see *Chymistry*—c. Hygienic, see *Chymistry*—c. Medical, see *Chymistry*—c. Organic, see *Chymistry*—c. Pharmaceutic, see *Chymistry*—c. Therapeutical, see *Chymistry*—c. Vegetable, see *Chymistry*—c. Vital, see *Chymistry*.

CHYMOCEZIA, (*chymus*, and *χεῖω*, 'to go to stool,') *Cœliac flux*.

CHYMOPLA'NIA, (G.) *Chymoplanien*, *Dyschymosen*, (*chymus*, and *πλανη*, 'a wandering,') A transposition of secretions:—a family of diseases in the classification of Fuchs, which includes icterus, uroplania, menoplania and galactoplania.

CHYMORRHEEA, (*chymus*, and *ῥέω*, 'to flow,') *Cœliac flux*, *Lientery*.

CHYMOSIN, *Pepsin*.

CHYMOSIS, *Chemosis*, *Chymification*.

CHYMOUS, *Chymo'sus*, (F.) *Chymeux*. Relating to, or resembling chyme.

CHYMOZEMIA, (*chymus*, and *ζημία*, 'loss,') *Hypererinia*.

CHYMUS, *Chyme*.

CHYT'LEN, **RADIX**. A cylindrical root, bitter and inodorous, brought from China. It is held by the Chinese to be stomachic.

CHYT'LON, (*χυτλον*, from *χυω*, 'I pour out,') A liquid formerly used for rubbing the body after bathing.

CIBARIOUS, *Esculent*.

CIBARIUM, (from *cibus*.) *Aliment*.

CIBA'RIOUS PA'NIS, 'Coarse bread.' Bread made of second flour.

CIBA'TIO, *Tro'phē*. The taking of food. In Pharmacy, it is the same as *Incorporation*.

CIBUS, *Aliment*—c. *Albus*, *Blancmanger*—c. *Deorum*, *Asafoetida*.

CICATRICE, *Cicatrix*.

CICATRICES OVARIORUM, (pl. of *Cicatrix*.) *Stigmata ovariorum*.

CICATRIC'ULA, (dim. of *Cicatrix*.) A small cicatrix, *Stigma*, (F.) *Cicatriceule*. The term is, also, applied to a small white spot, called the *tread*, *chala'za*, *chala'zium*, observable at the surface of a fecundated egg. See *Molecule*.

CICATRISAN'TIA, *Epulo'tica*, *Synulo'tica*, *Apu'lo'tica*, *Catulo'tica*, *Ulo'tica*, (F.) *Cicatriscants*. Remedies formerly considered to be capable of producing cicatrization.

CICA'TRIX, gen. *Cicatri'eis*, *Cœcat'rix*, *U'lē*, *Ou'lē*, *Cic'atrice*, (from *cœcare*, 'to conceal,' because it conceals the wound.) (F.) *Cicatrice*. A scar. A seam, (Sc.) *Arr*. The union of parts, which have been divided. A *scar* or *formation*, of a reddish colour, afterwards whitish, and of variable thickness, which takes place at the surface of wounds or ulcers after their cure. A cicatrix may vary much in shape, consistence, and thickness. The cicatrix of a bone is called *Calus*. A *vic'ious cicatrix*, (F.) *Cicatrice vicieuse*, is one which interferes with the action of the parts on which it occurs. The *scars*, (F.) *Coutures*, after small pox, are called *Pits* or *Pockmarks*, (Sc.) *Pockmarks*, (F.) *Coutures par la petite vérole*. See *Pockmark*.

CICATRIZ VARIOLE, *Pockmark*.

CICATRIZA'TION, *Cicatriza'tio*, *Ulo'sis*, *Apu'lo'sis*, *Epulo'sis*, *Synulo'sis*. The process by which a cicatrix is formed. Every tissue, except the nails, epidermis, hair, and enamel is, probably, capable of cicatrization.

CICELY, **SWEET**, *Chærophyl'ium odoratum*, *Osmorrhiza longistylis*, *Scandix odorata*.

CYCER ARIETINUM. The *Cicer* plant, *Erebin'thus*, (F.) *Cicérole*, *Pois Chiche*, *Girvanee*, *Peasette*. The seeds are ground into flour, and used as bread in some countries.

CICER LENS, *Ervum lens*.

CICERA TAR'TARI. Small pills of turpentine and cream of tartar—of the size of a vetch or cicer.

CICÉROLE, *Cicer arietinum*.

CICHORE, (κίχωρον), *Cichorium intybus*.

CICHO'R'IUM ENDIV'IA. The systematic name of the *Endive*, *Endiv'ia*, *Endiv'ca*, *Intubum*, *Intybum* (Antiq.), *Scari'ola*, *Intybus hortensis*, (F.) *Chicorée des Jardins*, *Scariola*. Ord. *Cichoraceæ*. *Sex. Syst.* Syngenesia Polygamia æqualis. It is a common pot herb, and is eaten as salad.

CICHO'R'IUM INTYBUS, *Cichorium*, (κίχωρον), *Chicor'è*, *Se'ris*, *Seri'ola*, *Intubum erraticum*. The systematic name of the *Wild Succory*, *Wild Chic'ory*, *Chick'ory*, *Chic'ory*, *Wild Endive*, *Amblet'ca*, *Heliotro'pion*, *Catanan'cè*, *Cichoreum*, (F.) *Chicorée sauvage*. It is bitter, hence its name *Pi'cris*, (from πικρός, 'bitter'), and was once given as a tonic. The root, roasted and ground, is often used instead of, or mixed with, coffee.

CICHORY, *Cichorium intybus*—c. *Wild*, *Cichorium intybus*.

CICI, (κικι), *Ricinis communis*.

CICINDE'LA, (dim. of *candela*, 'a candle,' [?]) *Lam'pyris Noctilu'ca*, *Nited'ula*. The *Glow-worm*. (F.) *Ver luisant*. This insect was once thought to be anodyne and lithontriptic.

CICINNUS, (κικίννος), *Cinnamus*.

CICINUS, (κικίνος), *Cincinnati*.

CICIS, (κικίς), see *Quercus infectoria*.

CICON'GIUS; an ancient measure, containing 12 pints.

CICUTA, *Conium maculatum*.

CICU'TA AQUAT'ICA, *C. viro'sa*, *Cicuta'ria aquat'ica*, *Coriandrum cicuta*, *Water Hem'lock*, *Cowbane*, (F.) *Ciguè aquatique* ou *vireuse*, *Cicuta'ria aquatique*. Family, Umbelliferae. *Sex. Syst.* Pentandria Digynia. A violent poison, often eaten by mistake for *Wild Smal'lage*, *A'pium Grave'olens*. It produces tremors, vertigo, burning at the stomach, and all the symptoms occasioned by the *Narcotico-acrid* class of poisons.

CICUTA MACULA'TA, (F.) *Ciguè d'Amérique*, *American water hem'lock*, *American Hem'lock*, *Snake'weed*, *Death of man*, *Water pars'ley*, *Poi'son root*, *Wild Hem'lock*, *Child'ren's bane*, *Spotted Cowbane*, is analogous in botanical character and medical properties to the European species. See *Conium maculatum*.

CICUTA MAJOR, *Conium maculatum*—c. *Major foetida*, *Conium maculatum*—c. *Stoerkii*, *Conium maculatum*—c. *Terrestris*, *Conium maculatum*—c. *Virosa*, *Cicuta aquatica*—c. *Vulgaris*, *Conium maculatum*.

CICUTAIRE AQUATIQUE, *Cicuta aquatica*.

CICUTARIA, *Cherophyllum sylvestris*—c. *Aquatica*, *Cicuta aquatica*, *Phellandrium aquaticum*—c. *Odorata*, *Cherophyllum odoratum*.

CICUTINE, see *Conium*.

CYDER, *Cy'der*, *Poma'ceum*, (F.) *Cidre*. This word is said to have been formerly written *sidre*, and to have come from *Sic'era*, *αίκερα*, which signifies any kind of fermented liquor other than wine. It is made from the juice of apples, and, when good, is a wholesome drink.

A liquor, similarly prepared from pears, is called *Per'ry*, (F.) *Poiré*.

CIDRE, *Cider*.

CIERGE DE NOTRE DAME, ('wax taper of our lady,') *Verbascum thapsus*.

CIGAR', *Cigar'ra*, (S.) *Cigarro*, (F.) *Cigare*. A small tubular roll of tobacco, used for smoking.

Medic'inal or *Medic'ated Cigars*, (F.) *Cigares Médicinaux*, are made of dried plants, with, or without, the addition of medicinal articles. When rolled in paper, they form *cigarettes*, which may be used with or without combustion, as in the case of the *Cigarettes d' Camphre*. The *Cigarettes*

arsénicales de Boudin are made of paper impregnated with an aqueous solution of arsenious acid, or of arseniate of soda, and dried.

CIGARETTE, see *Cigar*—c. *Arsénicate de Boudin*, see *Cigar*—c. *de Camphre*, see *Cigar*.

CIGARRA, *Cigar*.

CIGNUS; an ancient measure, which contained about two drachms.

CIGUÈ D'AMÉRIQUE, *Cicuta maculata*—c. *Aquatique*, *Cicuta aquatica*—c. *d'Eau Phellandrium aquaticum*—c. *Grande*, *Conium maculatum*—c. *Officinale*, *Conium*—c. *Ordinaire*, *Conium maculatum*—c. *Petite*, *Æthusa cynapium*—c. *Vireuse*, *Cicuta aquatica*.

CIL'IA (singular *Cil'ium*), *Blephar'ides*, *Epil'co'tides*, *Pili palpebra'rum*, *Eyelashes*, (Sc.) *Wink'ers*. The hairs on the eyelids. (F.) *Cils*. Their use seems to be, to prevent the entrance into the eye of light bodies flying in the atmosphere; and to diminish, in certain cases, the intensity of light. Also, the tarsi, eyelids. Likewise, a peculiar sort of moving organs, resembling small hairs, *vibratory* or *vibratile cil'ia*, *Cil'ia vibrato'ria*, (F.) *Cils vibratils*, *Tromyles*, which are visible with the microscope in many animals. These organs are found on parts of the body which are habitually in contact with water, or other more or less fluid matters, and produce motion in these fluids, impelling them along the surface of the parts. Cilia have been found to exist in all vertebrated animals except fishes, having been discovered on the respiratory, uterine, and other membranes of mammalia, birds, and reptiles.

The terms '*vibratory motion*' and '*ciliary motion*' have been used to express the phenomena exhibited by the moving cilia; and it is probable, that this motion is concerned in the progression of fluids along the membranes. As yet, the motion has only been observed in the direction of the outlets of canals.

CILIAIRE, *Ciliary*.

CIL'ARY, *Ciliar'is*, (F.) *Ciliaire*. Relating to the eyelashes, or to *cilia*. This epithet has, also, been applied to different parts, which enter into the structure of the eye; from the resemblance between some of them (the *ciliary processes*) and the eyelashes.

CILIARY ARTERIES, *Arte'rie cilia'res*, (F.) *Artères ciliaires*. These are furnished by the ophthalmic artery. They are distinguished into, 1. *Short or poste'rior* (*Art. uéales*—Chauss.). 30 or 40 in number, which are distributed to the ciliary processes. 2. *Long*, (*Art. Iriennes* of Chauss.) two in number, which, by the anastomosis of their branches, form two arterial circles at the anterior surface of the iris: and, 3. The *ante'rior*, *Arte'rie cilia'res antero'ries* of Haller, the number of which is variable. These pierce the sclerotic a few lines from its union with the cornea; and are principally distributed to the iris.

CILIARY BODY, *Cor'pus Cilia're*, *Nexus Stam'in'eus Ocul'i*, *Cor'ona Ciliar'is*, *Ciliary Disc*, (F.) *Corps ciliaire*. A ring of the choroid surrounding the crystalline in the manner of a crown, placed behind the iris and the ciliary circle. It resembles the disk of a radiated flower, and is formed by the union of the ciliary processes. See *Ciliary Muscle*.

CILIARY CANAL', *Canal of Fonta'na*. A small, extremely narrow circular space, formed between the ciliary circle, the cornea, and the sclerotic. It can be filled with injection, and it is not certain that it is not the cavity of a blood-vessel.

CILIARY CIRCLE, *Ciliary ligament*—c. *Disc*, *Ciliary body*—c. *Ganglion*, *Ophthalmic ganglion*.

CILIARY LIG'AMENT, *C. Circle* or *Ring*, *Ligamentum seu Interstit'ium cilia're* seu *Pridis*, *Plex'us cilia'ris*, *An'nulus* seu *Circ'ulus* seu *Orb'iculus cilia'ris*, *A. cellulo'sus*, *Com'missu'.* of the *Uvea*,

(F.) *Commissure de la Choroidé*—(Ch.), *Ligament o. Circle ciliare, Circle de la Choroidé, Ceinture blanche de la Choroidé*. A species of greyish ring, of a pulpy consistence, situate between the choroid, iris, and sclerótica. The internal surface of the choroid is uniform, until it approaches within ten lines and a half of the edge of the cornea; here a dentated line is observed, termed *o. r. serrata*. The outer surface presents the *anulosa albidus seu gangliiformis*, the anterior edge of which unites to the inner surface of the sclerótica and constitutes the *ciliary ligament*.

CILIARY or TAR'SAL MAR'GIN of the eyelids; (F.) *Bord ciliaire*. The edge in which the cilia or eyelashes are situate.

CILIARY MOTION, see *Ciliary*.

CILIARY MUS'CLE, *Musculus ciliaris*. The part of the orbicularis palpebrarum in the vicinity of the ciliary margin. Also, the greyish, semi-transparent structure behind the ciliary ligament, and covering the outside of the ciliary body. By its contraction the ciliary processes, and with them the lens, must be drawn towards the cornea. It appears to be the same muscle as the *Ten'sor choroidæ*, *Pro'trahens lentis*, *Choroid muscle* of some anatomists.

CILIARY NERVES, (F.) *Nervi ciliares, Nervi Iriens*. These are 12 to 16 in number. They arise from the nasal nerve, and particularly from the anterior part of the ophthalmic ganglion; and unite in two fasciculi, which pass around the optic nerve, and pierce the sclerótica near the entrance of that nerve into the eye. They are lost in the ciliary ligament.

CILIARY PLEXUS, C. Ligament.

CILIARY PROC'ESSES, *Processus seu Rad'ii seu Striæ ciliares*, (F.) *Process ciliares, Rayons sous-iriens*—(Ch.). Triangular folds, sixty or eighty in number, placed at the side of each other, and radiating so as to resemble the disk of a radiated flower. They are lodged in depressions at the anterior part of the vitreous humour. The uses of these processes are not known.

CILIARY RING, *Ciliary ligament*.

CILIARY STRIÆ are numerous pale, radiated striæ in the posterior portion of the *Corpus ciliare*, but so covered by the *Pigmentum nigrum* as not to be distinctly seen till the paint is removed. The ciliary processes are formed by these striæ.

CILIARY VEINS, (F.) *Veines ciliares*, follow nearly the same course as the arteries. In the choroid they are so tortuous, that they have received the name *Vasa vorticosa*. They open into the ophthalmic vein.

CILIARY ZONE, *Zo'na seu Zo'nula seu Lam'ina Ciliaris, Membran'ula Coro'næ Ciliaris*. Under the corpus ciliare, the capsule of the vitreous humour sends off an external lamina, which accompanies the retina, and is inserted, with it, into the forepart of the capsule of the lens, a little before its anterior edge. This is the *Zo'nula ciliaris, Zo'nula Zinnii* or *Zo'nula of Zinn, Coro'na Ciliaris, Orbiculus Ciliaris*. It is of a striated appearance and circular form, and assists in fixing the lens to the vitreous humour.

CIL'IATED, *Cilia'tus*, (F.) *Cilié*. Provided with cilia—as '*ciliated epithelium*,' the epithelium to which vibratory cilia are attached.

CILIUM, see *Cilia*.

CILLEMENT, (from *cilia*), *Nictation*.

CIL'LO, (*κίλλω*, 'to move') A name given by some authors to those whose upper eyelid is perpetually tremulous;—a trembling, which in some cases is called *Life's blood*. 'To have life's blood in the eye,' in other words, is to have this affection. Vogel calls it *Cillo'sis*.

CILLOSIS, *Cillo*.

CILS, *Cilia*—c. *Vibratilis*, see *Cilia*.

CIMEX, gen. *Cim'icis*, *C. lectularius*, *Acan'thia lectularia*, *Co'ris*, *Ko'ris*. The Bug; Wall, House

or Bed Bug or Chinche. (F.) *Punaise*. Six or seven of these, given internally, are said to have prevented ague! There is scarcely anything which is sufficiently disgusting, that has not been exhibited for this purpose, and with more or less success. The bug has also been esteemed emmenagogue.

CIMICIF'UGA, (*cimex*, and *fugare*, 'to drive away,') *Actæa racemosa*.

CIMO'LIA PURPUS'S/CENS, (*κιμωλία*), *Ter'ra Saponaria seu Fullonica, Fuller's Earth*. A compact bolar earth, employed in the arts. Used at times as a cooling application to inflamed nipples, &c.

CIMO'LIA TER'RA, *Cimo'lia al'ba, Cimo'lis, Smect'is, Smect'ris, Cimo'litæ*; from *κιμωλος*, an island in the Cretan Sea, where it was procured. It was formerly used as an astringent. Probably, the same as the last.

CINA CINA, *Cinchona*—c. *Levantica, Artemisia Santonica*.

CINABARIS, *Hydrargyri sulphuretum rubrum. CINABARIUM*, *Hydrargyri sulphuretum rubrum*.

CIN'ABRA, (*κίναβρα*), *Gra'sus*. The smell of a he-goat. A rank smell, like that of the armpit, *Hir'cus ala'rums*.

CINABRE, *Hydrargyri sulphuretum rubrum*.

CINÆDIA, (*κιναιδία*), *Masturbation*.

CINARA HORTENSIS, (*κινάρα*), *Cynara scolymus*—c. *Scolymus*, *Cynara scolymus*.

CINCHO'NA or **CHINCHO'NA**. (So called from the Spanish Viceroy's lady, the Countess de Chinchon, who was cured of fever by it at Lima, about 1638.) Called also *Cor'tex seu Pul'vis Jesuit'icus, Jes'uit's Bark* or *Pow'der, Cor'tex Pa'trum*, because it was introduced into Europe by the Jesuits; also *Pul'vis Comit'isæ* or the *Count'ess's Pow'der*, and *Cardinal del Lu'go's Pow'der, Cor'tex Cardinal'is de Lu'go*, because he introduced it at Rome; and in France, *Tal'bor's Pow'der* and *Eng'lish rem'edy*, because successfully used there by Sir Robert Talbor, who kept it a secret. It is the pharmacopoeial name of several kinds of barks from various species of *Cinchona*, from the western coast of South America. *Order, Cinchonaceæ. Sex. Syst. Pentandria Monogynia*. The richest in the antiperiodic alkaloids is said to be the bark of *Cinchona succirubra* of Ecuador, which, with other species, has been transplanted to India and other countries. Called, also, *Cor'tex, Bark, Peruvian Bark, Cor'tex Chinæ seu Chi'na, Chinch'ina, Pa'lo de Calcuti'ras, Ki'na Ki'na* or *Qu'ina Qu'ina*, (*Bark of Barks*), *Kink'ina, Ci'na Ci'na, Quinqu'ina, Mug'num De'i do'nun*, (F.) *Quinquina*; and in Peru *Cascara* and *Cascarrilla*. See *Cascara*.

CINCHONA AMYGDALIFOLIA, see *Cinchonæ cordifoliæ cortex*—c. *Boliviana*, see *Cinchonæ cordifoliæ cortex*—c. *Calisaya*, see *Cinchonæ cordifoliæ cortex*—c. *Condaminea*, see *Cinchonæ lancifoliæ cortex*—c. *Micrantha*, see *Cinchonæ lancifoliæ cortex*—c. *Officinalis* (*Cortex Flavus*), *Cinchonæ cordifoliæ cortex*—c. *OVATA*, see *Cinchonæ cordifoliæ cortex*—c. *PALLIDA*, *Cinchonæ lancifoliæ cortex*—c. *PITAYA*, *Pitaya bark*—c. *PUBESCENS*, see *Cinchonæ cordifoliæ cortex*—c. *SCROBICULATA*, see *Cinchonæ cordifoliæ cortex*—c. *SUCCIRUBRA*, see *Cinchona*, and *Cinchonæ oblongifoliæ cortex*—c. of *Virginia, Magnolia glauca*.

CINCHONÆ CARIBÆÆ COR'TEX, from *Exosent'ma Caribæum, Caribæan* or *Saint Lucie's Bark*, (F.) *Écorce de Saint Lucie, Quinquina Piton*, from *Exosent'ma* or *Exosent'ma, floribundum*; and the *Pitaya Bark, Quinquina bi'color*, from an *exosentema* [?] or from *strychnos pseudoquina* [?], are useful substitutes for the cinchona of Peru. These are the most important spurious barks. They contain neither quinia nor cinchonia.

CINCHONÆ CORDIFOLIE COR'TEX, *Cor'tex flu'vus, Cincho'næ officina'lis cor'tex flu'vus, Chi'nu*

re gia seu Calisaya, *Cortex Ch'ina re'gius seu fl'vus seu lu'teus*, *Yellow, Royal or Calisaya's cinchona Bark*, (F.) *Quinquina jaune ou jaune royal, Calisaya*. Odour aromatic; taste strong, bitter, astringent. Not rolled; often without the epidermis, which is very thick and inert; light, friable; fracture fibrous. Active principle *Quinine*. The false or *spurious Calisaya barks* are those of *Cinchona Calisaya*, variety *Josephina*; *C. Bolivia'na*, *C. ova'ta*, variety *rufiner'vis*, called, in Peru, *Cascarilla Carabaya*—the *Carabay'a bark of commerce*;—of *C. scrobiculata*, two varieties, *Cus'co bark* and *Bark of St. Ann*; of *C. pubes'cens*, which furnishes the *Cus'co or Arit'ca bark*, and of *C. amygdalifolia*.

CINCHONÆ LANCEFO'LIÆ CORTEX, *Cortex Peruvia'nus seu pal'lidus*, *Cincho'næ officina'lis cortex commun'is*, *Cincho'na pal'lida*, *Pale Cincho'na Bark*, *Pale, Lox'a or Crown Bark*, (F.) *Quinquina gris de Loxa, Quinquina Orange*. The bark of *Cincho'na Condamine'a* and *C. micran'tha*. Its odour is aromatic; taste pleasant, bitter, and astringent. The pieces are rolled in double or single quills. Epidermis brown, cracked; fracture resinous. Internally of a cinnamon colour. Its active principle is *Cincho'nia*.

CINCHONÆ OBLONGIFO'LIÆ CORTEX, *Cortex rub'er*, *Cincho'næ officina'lis cortex rub'er*, *Red Cinchona Bark*, *Red Bark*, (F.) *Quinquina rouge*, (S.) *Cascarilla roxa* and *C. colorada*, of S. America. Odour and taste the same as the pale, but more intense: in large flat pieces, solid, heavy, dry; fracture short and smooth; of a deep brownish-red colour. Although this variety of bark is assigned to the *Cinchona oblongifolia* and *C. Succir'bra* by some, it would seem, that nothing is sufficiently known as to its source. Active principles, *Cincho'nia* and *Quinine*.

The last three are the only official varieties in the Pharmacopœia of the United States. There are many other varieties, however, which are genuine cinchona barks, and yet have not been considered worthy of an official position. The Edinburgh Pharmacopœia admitted, indeed, *Cincho'na cine'ra*, *Grey bark*, *Sil'ver bark* or *Huanu'co or Li'ma bark*, which is obtained around Huanuco in Peru, and, with the *Jae'n or Ash bark* and the *Huamiles bark*, belongs to the class of pale or Loxa barks. Amongst the genuine but inferior barks are those brought from the northern Atlantic ports of South America, which, in commerce, are variously called *Pitay'a*, *Bogota'*, *Carthage'na*, *Maracay'bo*, and *San'ta Mar'tha barks*. The *Bogota* is also called *Fusagasuga* and *Coquetta bark*.

All these barks are bitter, astringent, tonic, and eminently febrifuge. The yellow bark has been thought equal to any of the others, but the red contains more active principle. The discovery of their alkaloids is one of the most important gifts of modern chemistry. Still, in pernicious intermittents, the bark, in substance, is often demanded. It is employed in every disease in which there is deficient tone; but in cases where the stomach is much debilitated, the powder had better be avoided, in consequence of the woody fibre, which might disagree. Externally, it is used in enemata, gargles, &c., and in gangrenous ulcerations. When it excites nausea, an aromatic may be added to it; if purging, opium; if costiveness, rhubarb, &c. Dose, ʒss to ʒj or more.

ESSENTIAL SALT OF BARK, as it is called, is an extract, prepared by macerating the bruised substance of bark in cold water, and submitting the infusion to a very slow evaporation.

CINCHONIA, *Cinchonine*—c. Sulphate of, see *Cinchonine*—c. Tannate of, *Quinix et cinchonix annas*.

CINCHONLE SULPHAS, see *Cinchonine*.

CINCHON'IC, *Cincho'nicus*, *Qu'nicus*, (F.) *Quinine*. Appertaining or relating to cinchona, as '*Cinchonic* or *Quinic* preparations.'

CINCHON'ICINE, *Cincho'nic'ina*, *Cinchonic'ia*. When a salt of cinchona is exposed to heat, with certain precautions, the alkaloid is changed into another, isomeric with itself, to which this name has been given. *Cinchonine*, and its sulphate, possess the same medical properties as cinchona and its salts. Dose, gr. v to gr. xv.

CINCHONINE, *Cincho'ni'na*, *Cinch'onin*, *Cincho'nia*. The active principle of *Cincho'na lancefo'lia*. An organic, crystalline alkali; of a white colour, and bitter, slightly astringent taste; very soluble in alcohol and ether, but almost insoluble in water.

Cincho'nie Sulphas, *Sul'phate of Cincho'nia*,—which is formed directly from cinchona, by dissolving cinchona, obtained from the mother-water remaining after the crystallization of sulphate of quinia by precipitation by means of solution of soda; treating it frequently with alcohol to remove other alkaloids; adding sulphuric acid to saturation, and crystallizing,—is soluble in water and alcohol. The action of the sulphate of cinchona is similar to that of the sulphate of quinia; but it is less energetic, and consequently requires to be given in a larger dose.

CINCHONINE, **TARTRATE OF**, see *Quinine*, tartrate of.

CINCHONISM, *Quininism*.

CINCHONOLGY, *Quinology*.

CINCHOTIN, *Quinidia*.

CINCIN'ULUS, (dim. of *Cincinnus*.) A little lock or curl of hair.

CINCIN'NUS, (κικινος, κικινος.) A curled or frizzled lock. The hair on the temples.

CINC'LISIS, *Cincli'smus*, (κικλκλκλς, 'agitation; rapid and frequent motion.') The movement of the thorax in dyspnoea. It has been used, also, synonymously with *nicitation*.

CINCLISMUS, (κικκλκλκλς,) *Cinclisis*.

CINEFACTIO, (*cinis*, pl. *cineres*, 'ashes,' and *facere*, 'to make,') *Incineration*.

CINE'MA, gen. *Cine'matis*, (κινημα,) *Cine'sis*, (κινω, 'I move,') *Motion*.

CINERARIA MARITIMA, (*cineres*, 'ashes,' from the white down that covers the leaves.) *Achoa'ron*, *Abiat*.

CINERATION, (*cineres*, 'ashes,') *Incineration*.

CINERES CLAVELLATI, see *Potash*—c. *Gravellati*, see *Potash*—c. *Russici*, *Potash of commerce*.

CINEREUS, *Cineritious*.

CINERIT'IOUS, *Ciner'eus*, (*cineres*, 'ashes,') (F.) *Cendrê*. Of the colour of ashes. The cortical substance of the brain, and the vesicular neurine in general, have been so called. See *Cortex Cerebri* and *Neurine*.

CINESIPATHY, (κινησις, 'motion,' and παθος, 'affection,') *Kinesipathy*.

CINESIS, (κινησις,) *Cinema*, *Motion*.

CINTH'MICS, (κινω, 'I move,') The science of movements in general.

CINETIC, (κινητικός,) *Motory*.

CINETICA. Diseases affecting the muscles, and characterized by irregular action of the muscles or muscular fibres, commonly denominated *Spasm*. The 3d order in the class *Neurotica* of Good. Also, agents that affect the voluntary or involuntary motions.

CINETUS, (κινω, 'I move,') *Diaphragm*.

CINGULARIA, (from *cingulum*, because of its shape.) *Lycopodium*.

CIN'GULUM, *Zone*, (*cingere*, *cinctum*, 'to gird,') (F.) *Ceinture*. A cincture. A girdle. The part of the body, situate below the ribs, to which the girdle is applied. The *waist*. *Herpes zoster*.

CINGULUM HILDA'NI, *Zo'nula Hilda'ni*, (F.) *Ceinture de Hildane*. A leathern girdle formerly

used for the reduction of luxations and fractures of the extremities.

CINGULUM MERCURIALE, *C. Sapientia* seu *Stultitia*. A woollen girdle, containing mercurial ointment. It was used as an antisymphilitic, and in diseases of the skin. (F.) *Ceinture de vif argent*.

CINGULUM SANCTI JOANNIS, *Artemisia vulgaris*.

CININ, *Santonine*.

CINIS FÆCUM, gen. *Cin'eria*, ('ashes of lees,') see Potash—*c. Infectorius*, see Potash.

CINNABAR, (*κινναβάρη*.) Hydrargyri sulphuretum rubrum—*c. Græcorum*, see *Calamus rotang*.

CINNABRE, Hydrargyri sulphuretum rubrum.

CINNAMOMUM, (*κινναμόμων*.) *Laurus cinnamomum*—*c. Album*, *Canella Alba*—*c. Aromaticum*, see *Laurus cinnamomum*—*c. Culilawan*, *Laurus Culilawan*—*c. Indicum*, *Laurus cassia*—*c. Magellanicum*, *Winters aromatica*—*c. Malabaricum*, *Laurus cassia*—*c. Sylvestre Americanum*, *Nectandra cinnamomoides*—*c. Zeylanicum*, *Laurus cinnamomum*.

CINNAMON, (*κινναμόμων*) see *Laurus cinnamomum*—*c. Malabar*, *Laurus cassia*—*c. Wild*, *Laurus cassia*.

CINON'OSI, (*κινω*, 'I move,' and *νοσος*, 'a disease.') Diseases of motion.

CINOPER, Hydrargyri sulphuretum rubrum.

CINOPLANE'SIS, (*κινω*, 'I move,' and *πλανησις*, 'a wandering about,') Irregularity of motion.

CINQUEFOIL, (*quinque*, 'five,' and *folium*, 'a leaf,') *Potentilla reptans*—*c. Marsh*, *Comarum palustre*—*c. Norway*, *Potentilla Norvegica*.

CINZILLA, (*Cingulum*.) *Herpes zoster*.

CION, (*κίον*, 'a column,') *Uvula*.

CI'ONIS, (*κίονις*.) The *Uvula*. Also, tumefaction, or elongation of the uvula; *Staphylo'dial'ysis*.

CIONITIS, (*cion*, and *itis*.) Inflammation of the uvula, *Uvul'itis*.

CIONORRHAPHIA, (*cion*, and *ράφη*, 'a suture,') *Staphylo'raphy*.

CIONOTOME, (*cion*, and *τομή*, 'incision,') Curved scissors for cutting off the uvula.

CIONOTOMY, *Cionotom'ia*. Excision of the uvula when too long.

CIPIPA, see *Jatropha manihot*.

CIRCEA, (after *Circe*, the Enchantress,) *Atropa mandragora*, *Circeæ Lutetiana*.

CIRCEA LUTETIANA, *Circe'a*, *Paris'ian Circe'a*, *Enchant'ers' Nightshade*, (F.) *Herbe de Saint Etienne ou aux Sorcières*. This plant, common in the vicinity of Paris, was formerly considered to be resolute and vulnerary. It was also supposed to possess wonderful magical and enchanting properties.

CIRCINUS, (dim. of *κίρκος*, 'a circle,') *Herpes zoster*.

CIRCLE, *Circulus*—*c. Ciliary*, *Ciliary ligament*—*c. of Willis*, see *Circulus*.

CIRCOCELE, *Cirsocele*.

CIRCON (F.), as a prefix, *Circum*.

CIRCUNCISION, *Circumcision*.

CIRCUNFLEXE, *Circumflexus*.

CIRCUNSCRIT, *Circumscribed*.

CIRCUNVOLUTION, *Convolution*.

CIRCUIT, *Circu'itus*; (*circumeo*, 'to go around,') in pathological language, generally means 'period, course.'

CIRCUITUS, *Circuit*, *Period*.

CIRCULAR, *Circula'ris*, (*circulus*, 'a circle,') (F.) *Circulaire*. Having the form of a circle; as *Circular Amputation*, &c.

The French use the expression '*Une circulaire*, for a turn of a bandage around any part.

CIRCULAR SINUS of *Ridley*, *Sinus coronarius*.

CIRCULA'TION, *Circula'tio*, *Cyclophor'ia*, *Cyclo'sis*, *Peri'odus san'guinis*, (*circum*, 'around,' and *ferre*, *latum*, 'to carry,') (F.) *Circulation*. Phy-

siologists give this name to the motion of the blood through the different vessels of the body—*sanguimotion*;—to that function, by which the blood, setting out from the left ventricle of the heart, is distributed to every part of the body by the arteries through the capillaries;—proceeds into the veins, returns to the heart, enters the right auricle, and passes into the corresponding ventricle, which sends it into the pulmonary artery to be distributed to the lungs, whence it issues by the pulmonary veins, and passes into the left auricle; from this it is sent into the left ventricle, and is again distributed by means of the arteries.

CIRCULATION, *CAP'ILLARY*, *C. des Parenchymes*, is that which takes place in the capillary vessels; and is, in some measure, independent of the action of the heart. See *Capillary Vessels*.

CIRCULATION, *PULMON'IC* or *LESS'ER*, is the circle from the right to the left side of the heart by the lungs.—The *GREAT'ER* or *SYSTEMAT'IC* or *SYSTEM'IC*, is that through the rest of the system.

CIRCULATOIRE, *Circulatory*.

CIRCULATOR, *Charlatan*.

CIRCULATORY, *Circula'torius*, (F.) *Circulatoire*. Relating to the circulation of the blood;—*sanguimot'ory*.

CIRCULUS, (dim. of *circus*, *κίρκος*, 'a circle,') A circle or ring; *Cy'clus*, (F.) *Cercle*. Any part of the body, which is round or annular, as *Circulus Oculi*—the *globe*, *bulb*, or *orb* of the eye. It is, also, applied to objects, which by no means form a circle,—as to the *Circle of Willis*, *Circulus arterio'sus Willis'ii*, (F.) *Hexagone artérielle*, which is an *anastomotic circle* at the base of the brain, formed by the anterior and the posterior cerebral arteries and the communicating arteries of Willis.

CIRCULUS ARTERIO'SUS IRIDIS. The artery which runs round the iris, and forms a circle.

CIRCULUS ARTERIOSUS WILLISII, *Circle of Willis*, see *Circulus*.

CIRCULUS CALLO'SUS HALLE'RI. *Ten'do cor'dis veno'sus*. The fibro-cartilaginous ring, around each auriculo-ventricular opening of the heart, to which the tricuspid and mitral valves are attached.

CIRCULUS CILIARIS, *Ciliary ligament*—*c. Membranous*, *Hymen*.

CIRCULUS QUAD'RUPLEX; a kind of bandage used by the ancients.

CIRCULUS TONSILLA'RIS. A plexus formed by the tonsillitic branches of the glossopharyngeal nerve around the base of the tonsil.

CIRCULUS VENO'SUS, *Fig'u'ra veno'sa*, *Ve'na seu Si'nus termina'tis*. The venous circle in the embryo, which bounds the *A'rea Vasculo'sa* or *Vas'cular Area*.

CIRCULUS VENO'SUS ARE'OLE. The venous circle, formed by the union of the veins around the nipple. It embraces, however, only two-thirds of the circuit.

CIRCUM (L.), (F.) *circon*, as a prehx, 'around.'

CIRCUMAGENTES, (*circum*, and *agere*, 'to act,') Oblique muscles of the eye.

CIRCUMCALUALIS MEMBRANA, *Conjunctiva*.

CIRCUMCISED, *Circumcisis*.

CIRCUMCISIO, (*circum*, and *cædere*, 'to cut,') *Circumcision*—*c. Fœminarum*, see *Circumcision*.

CIRCUMCIS'ION, *Circumcis'io*, *Posthet'omy*, *Præcis'io* seu *Abcis'io Præpu'tii*, *Circumcis'u'ra*, *Circumsec'tio*, *Peri'tomê*, (F.) *Circuncision*. An ancient operation, performed by some nations as a religious ceremony. It consists in removing circularly a portion of the prepuce of infants;—a custom, which was probably suggested with a view to cleanliness. In cases of extraordinary length of prepuce, or when affected with disease, the operation is sometimes undertaken by sur-

geons. A similar operation is performed, amongst the Egyptians, Arabians, and Persians, on the female, *Circumcisio fœminarum*, by removing a portion of the nymphæ, and at times the clitoris.

CIRCUMCISURA, Circumcision.

CIRCUMCISUS, *Verpus*. One who has been circumcised.

CIRCUMDUCTIO, (*circum*, and *ducere*, *ductum*, 'to lead,') Perisphalxis.

CIRCUMDUCTIONIS OPIFEX, ('a worker of circumduction,') Obliquus superior oculi.

CIRCUMFLEX, *Circumflexus*, (*circum*, and *flectere*, *flexum*, 'to bend,') (F.) *Circonflexe*. Curved circularly. A name given to several organs.

CIRCUMFLEX OR ARTICULAR ARTERIES of the arm are distinguished into *anterior* and *posterior*. They arise from the axillary, and are distributed around the shoulder.

CIRCUMFLEX ARTERIES OF THE THIGH are distinguished into *external* and *internal*.—*A. Sous-trochantériennes*.—Ch. They are given off from the *Profunda*, and surround the head of the thigh bone.

CIRCUMFLEX MUSCLE, *Circumflexus Musculus*, *C. Palati Mol'lis*, *Ten'sor Palati*, *Peristaphyl'ius externus* seu *inferior*, *Sph'no-salpin'go-staphyl'ius* seu *Staphyl'ius externus*, *Musculus tub'no-væ*, *Palato-salpin'geus*, *Pter'gyo-staphyl'ius*, *Pe'tro-salpin'go-staphyl'ius*, *Sph'no-pter'gyo-palati'us*, *Salpingo-staphyl'ius*, (F.) *Palato-salpingien*. A muscle, which arises from the spinous process of the sphenoid bone, and is inserted into the *velum pendulum palati*. Its use is to stretch the velum.

CIRCUMFLEX NERVE, Axillary Nerve.

CIRCUMFLEX VEINS follow the arteries.

CIRCUMFORANEUS, (*circum*, and *foris*, 'a door,') Charlatan.

CIRCUMFUSA, (*circum*, and *fundere*, *fusum*, 'to pour,') Hallé has thus designated the first class of subjects that belong to Hygiene—as atmosphere, climate, residence, &c.; in short, every thing which acts constantly on man externally and internally.

CIRCMGYRATIO, (*circum*, and *gyrus*, 'a circle,') Vertigo.

CIRCUMLIGATURA, (*circum*, and *ligare*, 'to bend,') Paraphimosis.

CIRCUMLIT'IO, (*circum*, and *linere*, *litum*, 'to anoint,') *Perich'risis*, *Perichris'tion*. A term formerly used for liniments, but especially for those applied to the eyelids.

CIRCUMOSSALE, (*circum*, and *os*, 'a bone,') Periosteum.

CIRCUMSCRIBED, *Circumscrip'tus*, (F.) *Circonscrit*, (*circum*, and *scribere*, 'to write, to make lines,') A term applied, in pathology, to tumours, which are distinct at their base from the surrounding part.

CIRCUMSECTIO, gen. *Circumsectio'nis*, (*circum*, and *seco*, *sectum*, 'to cut,') Circumcision.

CIRCUMVALLATÆ PAPILLÆ, (*circum*, and *vallo*, *vallatum*, 'to intrench,') see *Papillæ* of the Tongue.

CIRCUS, (*κίρκος*, 'a circle,') A circular bandage.

CIRE, Cera—*c. Jaune et blanche*, Cera flava et alba—*c. des Oreilles*, Cerumen.

CIRON, Acarus, Psora.

CIRRHAGRA, (*cirrus*, 'a curl,' and *αγρα*, 'a seizure,') Plica—*c. Polonorum*, Plica.

CIRRHONOSIS, (from *κίρρος*, 'yellow,' and *οσις*.) Cirrhosis.

CIRRHONOSUS, (*κίρρος*, 'yellow,' and *νοσος*, 'disease,') A disease of the fetus, in which there is a yellow coloration of the serous membranes.

CIRRHOSE. Cirrhosis—*c. du Foie*, Cirrhosis. **CIRRHOSIS**, *Kirrho'sis*, *Cirrhono'sis*, *Kirrhono'sis*, (F.) *Cirrhose*, *Kirrhose*, (*κίρρος*, 'yel-

low,') A yellow colouring matter, sometimes secreted in the tissues, owing to a morbid process.

CIRRHOSIS HE'PATIS, *Granulated*, *granular*, *man'millated*, *tuber'culated*, and *lobu'at'ed liver*, (F.) *Cirrhose du Foie*. It appears to be dependent upon repletion of the terminal extremities of the biliary ducts with bile, along with altered nutrition of the intervening parenchyma, by which the liver becomes smaller in size or atrophied. See Hepatostrophia.

CIRRHOSIS OF THE LUNG, *Cirrho'sis pulmo'num*.

Dr. Corrigan has described a condition of the lung under this name, the general character of which he considers to be a tendency to consolidation or contraction of the pulmonary tissue, with dilatation of the bronchial tubes.

CIRRHOSIS PULMONUM, C. of the Lung.

CIRRHOT'IC, *Cirrho'ticus*. Affected with, or having the character of, cirrhosis.

CIRSIUM ARVEN'SÊ, (from *cirsus*, because formerly used in varix,) *Carduus hemorrhoida'lis*, *Oeano'thus*, (F.) *Chardon hémmorrhoidal*. A common plant, used in France in the form of cataplasm in hemorrhoids; and worn as at amulet.

CIRSIUM MACULATUM, *Carduus marianus*.

CIRSOCELE, *Cir'socèle*, improperly *Circocele*, *Cirso'scheum*, (*cirsus*, and *κηλη*, 'hernia,') *Var'icose Her'nia*. The greater part of authors have employed the term synonymously with *Var'icocele*. Pott gives it a different signification. *Var'icocele*, he calls the tumour formed by the veins of the scrotum; *Cir'cocele*, *Funic'ulus vari'cosus*, the varicose dilatation of the spermatic veins. The scrotum feels as if it contained earthworms. It is commonly an affection of no consequence, demanding merely the use of a suspensory bandage, but sometimes, requires an operation. See Varicocele.

CIRSOIDES, *Cirso'des*, (*cirsus*, and *ειδός*, 'resemblance,') Varicose, or resembling a varix. A term once applied to the upper part of the brain, as well as to the spermatic vessels.

CIRSOM'PHALUS, (*cirsus*, and *ομφαλος*, 'navel,') Varicose dilatation of the veins surrounding the navel. The term has, likewise, been applied to the aneurismal dilatation of the arteries of that region; called also, *Varicomph'alus*, (F.) *Hargne anévrysma'le*, *Aneurism'al Her'nia*.

CIRSOPHTHALMIA, *Cirso'pthalm'us*, (*cirsus*, and *ὀφθαλμος*, 'the eye,') *Telangiectasia oculi*, *Var'icose ophthalm'ia*, *Ophthalm'ia varico'sa*, *Varico'sitas conjunctivæ*. A high degree of ophthalmia, in which the vessels of the conjunctiva are considerably injected.

CIRSOSCHEUM, (*cirso*, and *σχεον*, 'scrotum,') *Cirsocele*.

CIRSOT'OMY, *Cirso'tom'ia*, (*cirsus*, and *τομή*, 'an incision,') Any operation for the removal of varices by incision.

CIRSUS, (*κίρκος*,) Varix.

CIRSYDROSCHEOCELE, (*cirsus*, 'ύδωρ', 'water,' *σχεον*, 'scrotum,' and *κηλη*, 'rupture,') Varicocele with water in scrotum.

CISEAU (F.), *For'fec*, (*scindere*, *scissum*, 'to cut,') An instrument, composed of a flattened rod of metal, sharpened at one extremity, and used in anatomical preparations, and in certain surgical operations, for dividing bones.

CISEAUX, Scissors.

CISSA, (*κίσα*,) Malacia.

CISSAMPELOS CAPEN'SIS, (*κίσσος*, 'ivy, and *αμπελος*, 'vine,') *Nat. Ord. Menispermaceæ*, grows in almost every mountainous part of the Cape of Good Hope. The root is used as an emetic and cathartic by the Boers.

CISSAMPELOS GLABERRIMA, see *Pareira brava*—*c. Pareira*, *Pareira brava*.

CISSINUM, (*κίσσιον*, from *cissus*,) Name of

a plaster of ivy, used in wounds of the nerves or tendons.

CISSOÏDES, (*cissus*, and *αῖσος*, 'resemblance') Capreaularis.

CISSUS, (*κισσός*), *Hedera helix*.

CISTERNA, LUMBAR, Receptaculum chyli.

CISTER'NA, (*κίστην*, (L.) *Cista*, 'a chest.') (F.)

Cisterna. This term has been applied to various parts of the body, which serve as reservoirs for different fluids. The fourth ventricle of the brain has been so called.

CISTERNA CHYLI, Receptaculum chyli.

CISTOCELE, Cystoceles.

CISTUS CANADENSIS, *Helianthemum Canadense*.

CIS'TUS CRE'TICUS, (*κιστός*), *C. salvifolius* seu *tauricus*. *Dorycinum*, *Gum Cis'tus*. *Ord.* Cystineæ; *Sex. Syst.* Polyandria Monogynia. The systematic name of the plant whence the *Labda-num*, *Labda'men* or *Lada-num*, *Gum'mi Labda-num*, is obtained. *Lada-num* is a gum-resinous substance, of a very agreeable smell, found in the shops in great masses. Its colour is blackish-green; taste, warm and bitter. It is but little used now. Formerly, it was a component of warm plasters, and was prescribed internally as a stomachic, and is, by some, regarded as the *On'ycha* of the Scriptures, which was burnt as incense. *Ladanum* is also obtained from *Cis'tus ladanif'erus*, and *C. laurifolius*.

CISTUS, GUM, *Cistus Creticus*—c. *Salvifolius*, *C. Creticus*—c. *Tauricus*, *C. Creticus*.

CITERNE LOMBAIRE, Receptaculum chyli.

CITHARUS, (*κίσθα*, and *κίσθαρος*), *Thorax*.

CITRAGO, (*citrus*, 'a citron'), *Melissa*.

CITRARIA, *Melissa*.

CITRAS CHINICUS, *Quiniaz citras*.

CITREA MALUS, see *Citrus medica*.

CITKEOLUS, *Cucumis sativus*.

CITRIC AC'ID, *Ac'idum citricum*, *Ac'idum Limo'nis*, *Ac'id of Lem'ons*, *Ac'idum Lim'onum*, (F.) *Acide citrique*. This acid is found in the juice of the lemon, lime, &c., from which it is obtained. It is in rhomboidal prisms, which slightly effloresce on exposure to the air. It dissolves in a twelfth part of its weight of boiling water, and has an extremely acid but agreeable taste. It is employed in medicine as antiseptic, refrigerant and diuretic. Rubbed up with sugar and with a little of the essence of lemon, it forms the *dry Lemonade*, (F.) *Limonade sèche*.

CITRION, (*κίτρον*), see *Citrus medica*.

CITRON, see *Citrus medica*—c. *Tree*, see *Citrus medica*.

CITRONADE, *Lemonade*.

CITRONELLE, *Artemisia abrotanum*, *Melissa*.

CITRONNIER, *Citrus medica*.

CITROUILLE, (from its citron colour), *Cucurbita*.

CITRUL, SICILIAN, *Cucurbita citrullus*.

CITRULLUS, (dim. of *citrus*, from its colour), *Cucurbita citrullus*.

CITRULLUS AMA'RUS. An African plant, *Ord.* Cucurbitaceæ, called by the Boers *Bitterappel* or *Wild Watermelon*, the pulp of which, like that of colocynth, is a drastic cathartic.

CITRULLUS COLOCYNTHIS, *Cucumis colocynthis*.

CITRUS, (after *Citron*, in Judea, [?]) see *Citrus medica*—c. *Acida*, see *Lime*.

CITRUS AURANTIUM. The systematic name of the *Orange Tree*, *Aurantium*, *A. Hispanicæ*, *Ma'lus Aurant'ia*, *Ma'lus Aurant'ia ma'jor* seu *vulga'ris*, *Aurantium vulga're*, *Citrus vulga'ris*. *Ord.* Aurantiaceæ. *Sex. Syst.* Polyadelphia Icosandria. The fruit are called *Ma'ta Au'rea*, *Chrysos'melia*, *Po'ma Aurant'ia* seu *Chinen'sia*, *Aurant'ia Curassav'ica*, *Oranges*. The flowers of the *Orange*, *Aurant'ii flores*, *Flo'res Naphæ*, are highly odoriferous, and used as a perfume. On distillation, they yield a small quantity of essential oil—

O'leum Aurant'ii, *O'leum seu Essen'tia Nero'li*, *Nero'li*,—with spirit and water, the *A'qua Flo'rum Aurant'ii*, *Aurant'ii flo'ris a'qua*, *A'qua aurant'ii*, *Or'ange-flow'er wa'ter*. They were once used in convulsive and epileptic cases. The leaves, *Fo'lia Aurant'ii*, have a bitterish taste, and furnish an essential oil. They have been used for the same purposes as the flowers. The yellow rind of the fruit, *Aurant'ii Cor'tex* (Ph. U. S. 1842), *Citrus Bigara'dia* (Ph. L.), *Bigarade'* or *Bitter Or'ange*, *Or'ange Peel*, is an aromatic bitter, and is used in dyspepsia, and where that class of remedies is required. The *Juice*, *Suc'cus Aurant'ii*, *Or'ange juice*, is a grateful acid, and used as a beverage in febrile and scorbutic affections.

The Ph. U. S. 1863 has *Aurant'ii ama'ri cor'tex*, *Bitter or'ange peel*, the rind of the fruit of *Citrus vulga'ris*; and *Aurant'ii dul'cis cor'tex*, *Sweet or'ange peel*, the rind of the fruit of *Citrus Aurantium*.

CITRUS BERGAMIA, *Citrus mella rosa*—c. *Bigaradia*, see *Citrus aurantium*—c. *Decumana*, *Shaddock*—c. *Limetta*, see *Citrus mella rosa*, and *Lime*—c. *Limonum*, *C. medica*.

CITRUS MED'ICA, *C. Lim'onum*, (F.) *Citronnier*, *Cédratier*. The systematic name of the *Lemon Tree*. The *Lem'on*, *Li'mon*, (Ph. U. S.), *Limo'num Ma'lum* seu *Bac'ca*, *Ma'lus Med'ica*, *M. Lim'onia* *Ac'idia*, *Cit'ra Ma'lus*, *Cit'rus*, (F.) *Citron*, *Cédrat*, has a fragrant odour, depending upon the essential oil, *O'leum Lim'onis*, of the rind. The outer rind, *Cor'tex Lim'onum*, *Limo'nis Cor'tex*, *Lem'on Peel*, *Zest*, *Flave'do Cor'ticum Cit'ri*, is used in the same cases as the *Cortex Aurant'ii*.

The juice, *Cit'rion*, *Suc'cus Lim'onis*, *Limo'num Suc'cus* (Ph. L.), (F.) *Suc du Limon*, *Suc de Citron*, is sharp, but gratefully acid, the acidity depending upon the citric acid it contains, and is given as a refrigerant beverage in febrile affections. In doses of half an ounce to an ounce, three times a day, it has appeared to some to exert a sedative influence on the circulation, and has been given, apparently with benefit, in acute rheumatism and rheumatic gout. Alone, or combined with wine, it is prescribed in scurvy, putrid sore throat, &c. Its general properties are refrigerant and antiseptic. Sweetened and diluted, it forms *Lemonade*. *Artific'ial lem'onjuice* is made by dissolving an ounce of citric acid in fourteen fluidounces of water; adding a few drops of essence of lemon.

Lemon'peel tea or *wa'ter* is made by paring the rind of one lemon, previously rubbed with half an ounce of sugar: the peelings and sugar are then put into a jar, and a quart of boiling water is poured over them. When cold, the fluid must be poured off, and a tablespoonful of lemon juice be added.

It is an agreeable drink in fevers.

Cit'ron Tree is likewise considered to belong to the same species—*Citrus Med'ica*. Its fruit is called *cedrome'la*. It is larger and less succulent than the lemon. *Cit'ron juice*, when sweetened with sugar, is called by the Italians *Agro di Cedro*.

CITRUS MEL'ITA Ro'sa of De Lamarck, another variety of *Citrus Med'ica*, affords the Bergamot, as also do *Citrus Limel'ta* and *C. Berga'mia*, (F.) *Limellier* and *Bergamottier*.

CITRUS VULGARIS, *Citrus aurantium*.

CITTA, (*κίττα*), *Malacia*.

CITTARA, MIN'ERAL WA'TERS OF. These springs are in the Isle of Ischia, near the sea. They contain carbonate and sulphate of lime, and chloride of sodium. Their temperature is 100° Fahrenheit.

CITTOS, (*κίττος*), *Hedera helix*.

CITTOSIS, (*citta* and *osis*), *Chlorosis*, *Malacia*.

CIVETTA, *Zib'ethum*, *Civ'et*, (F.) *Civette*. An unctuous perfume, of a very penetrating odour,

obtained from different mammalia of the *Viver'ra* kind, particularly from *Viver'ra civetta*. It is contained in a fold of the skin, situate between the anus and the organs of generation.

CIVITAS HIPPOCRATICÆ, Salernum.

CLABBER, Bonnyclabber.

CLABBERGRASS, Galium verum.

CLACKING (onomatopœia), *Claquement*.

CLADES GLANDULARIA, ('glandular scourge.') Plague.

CLADISCUS, (from *κλαδος*, 'a young shoot,') Ramusculus.

CLADONIA ISLANDICA, (from *cladus*, 'a branch,' because branched,) Lichen Islandicus.

CLADONIA RANGIFERINA. The ancients regarded this European plant as pectoral and stomachic. It enters into the composition of the *Poudre de Chypre*.

CLADRAS'TIS TINCTO'RIA, *Virgil'ia*, *Yell'low Ash*, *Fus'ic Tree*, *Yell'low Lo'cust*. An indigenous tree, which flourishes from Kentucky to Alabama. The bark of the tree and the roots are cathartic.

CLADUS, (κλαῦς,) Ramus.

CLAIRET, (from *clarus*, 'clear,') Claret.

CLAIRVOYANCE (F.), 'clear-seeing.' A clearness of sight, said to be communicated by animal magnetism; which not only enables the magnetized person to see in the dark, through stone walls, &c., but even to observe prospects, which he has never seen previously, whilst he may fancy he is flying in the air. It need hardly be said, that the possession of such powers is fabulous.

CLAM'MY, (Sax. clam, (D.) *Klam*, 'moist,') *Viscidus*, *Glutinosus*, (F.) *Pâteux*. Viscous; glutinous. A clammy mouth, (F.) *Bouche pâteuse*, is one that is covered with a mucous coat.

CLAMMY WEED, *Polanisea graveolens*.

CLAMOR, Cry.

CLANGOR, gen. *Clango'ris*, (κλαγγη, 'a loud clang or noise,') *Oxyphonia*.

CLAP, *Gonorrhœa impura*. ['a leper with a *clap*-*ish*, to give notice he is infectious,' Massinger.]

CLAP'IER (F.), 'a clapper or rabbit burrow,' (κλαπ'ειν, 'to conceal,') *Latibulum*. A purulent *foyer* of disease; concealed in the flesh or under the skin. See *Sinus*.

CLAPWORT, *Orobanche Americana*.

CLAQUEMENT (F.), *Clacking*. A sound—*Bruit de claquement*—produced by the bringing together of valves, as of those at the mouth of the pulmonary artery and the aorta during the dilatation of the ventricles; or of the auriculo-ventricular valves, during the contraction of the ventricles. Also, *Odontosynerismus*.

CLARE'NDON, MIN'ERAL WA'TERS OF. Mild acidulous springs in Vermont; temperature from 48° to 54° Fah. The gas, given off from the water, consists of carbonic acid, oxygen, and nitrogen. The springs have a reputation in cutaneous diseases, chronic bronchitis, &c.

CLAR'ET, (*Vin clair* [?]), *Claretum*, *Vin de Bordennx* (F.) *Clairret*. A pleasant French wine, which may be used whenever wine is required. Also, a wine impregnated with spice and sugar, called likewise *Vinum Hippocraticum* seu *Medicatum*, *Potus Hippocraticus*, *Hippocras*, *Hyp'poeras*. Schröder speaks of a *Claretum al'terans*, and a *C. pur'gans*.

CLARETA, (*clarus*, 'clear,') Albumen ovi.

CLARETUM, (*clarus*, 'clear,') Claret.

CLARIFICATION, *Clarificatio*, *Depuratio*, (*clarus*, 'clear,' and *facio*, 'I make,') A pharmaceutical operation, which consists in separating from a liquid every insoluble substance, held in suspension by it that affects its transparency. Decanting and filtering are the operations necessary for this purpose.

CLARY, COMMON, *Salvia sclarea*.

CLASIS, (κλασις,) Fracture.

CLASMA, gen. *Clas'matis*, (κλασμα,) Fracture.

CLASS, *Clas'sis*, (F.) *Classe*. An assemblage of a certain number of objects. In *Natural History*, and in *Medicine*, a group of objects or individuals having one or more common characters. The *classes* are divided into *orders*, the *orders* into *genera*, the *genera* into *species*, and these last into *varieties*.

CLASSIFICATION, *Classificatio*, (*classis*, 'a class,' and *facio*, 'I make,') The formation of classes. A methodical distribution of any objects whatever—as in *Natural History*—into classes, orders, genera, species, and varieties. See *Nosography*, and *Nosology*.

CLASSY, MIN'ERAL WA'TERS OF. Classy is near Laon, in Picardy, France. The waters are chalybeate.

CLASTIC ANATOMY, see *Anatomy*.

CLAUDICATIO, (*claudicare*, *claudicatum*, 'to halt,') Claudication—c. *Anaticæ*, *Veallatio*.

CLAUDICA'TION, *Claudica'tio*. The act of halting or limping, *Lameness*, (Prov.) *Hirpling*, *Clau'ditas*, *Cholo'sis*, *Chole'ia*, *Cholo'ma*, (F.) *Claudication*, *Boitement*, *Boiterie*. This condition does not constitute any special disease, but is produced by different causes or affections. It may be the result of the shortening or elongation of one of the lower limbs, of ankylosis of the joints, palsy of the muscles, pain, &c.

CLAUDITAS, Claudication.

CLAUSTRUM GUTTURIS, (*claudere*, 'to shut,') Isthmus of the fauces—c. *Palati*, *Velum pendulum palati*—c. *Virginittatis* seu *Virginalis*, *Hymen*.

CLAUSU'RA, (*claudere*, 'to shut,') An imperforation of any canal or cavity.

CLAUSU'RA U'TERI. Preternatural imperforation of the uterus.

CLAVA RUGOSA, (*clava*, 'a knotty branch,') *Acorus calamus*.

CLAVALIER À FEUILLES DE FRÊNE, *Xanthoxylum clava Herculis*.

CLAVARIA CORALLOÏDES, *Coralloïdes Fungus*, *Cor'alboort*. Ord. *Fungi*; (*clava*, 'a knotty branch, a club,') Said to be corroborant and astringent. A kind of clavaria, called (F.) *Digital blanc*, *Digital humain*, *Champignon de l'appareil des fractures*, formed of digitations, grouped together, and two or three inches in length, is said to have been often found, formerly, at the *Hôtel Dieu* of Paris, on the splints of white wood used in the treatment of fractures, in autumn.

CLAVATIO, (from *clavus*, 'a nail,') Gomphosis.

CLAVEAU, (from *clavus*, 'a nail,') Murr.

CLAVELÉE, Murr.

CLAVES CALVARIÆ, (pl. of *clavis*, 'a key,') *Wormiana Ossa*.

CLAVICLE, *Clav'is*, *Clavic'ula*, *Clavic'ulus*, *Lig'ula*, *Fur'cula*, *Os Jug'uli*, *Jug'ulum*, *Cleis*, *Cle'idion*, (dim. of *clavis*, 'a key,') (F.) *Clavicule*, the *Collar-bone*, (Old Eng.) *Can'nelbone*, *Chaw'nelbone* (Sc.), *Can'nelbayne*, *Crag'bane*, *Hals'bane*. The clavicle is shaped like the letter S, and is placed transversely at the upper part of the thorax. It is articulated, at one extremity, with the sternum; at the other with the acromion process of the scapula. It gives attachment, above, to the *Sternocleidomastoides*; below, to the *Subclavius*; before, to the *Pectoralis major* and *Deltoides*; and behind, to the *Trapezius*. It serves as a point of support for the muscles of the arm, and protects the vessels and nerves passing to that extremity.

The fibres, connecting the lamellæ or plates of bones, have also been called *Clavic'uli* or *Nails*.
CLAVICULA, Clavicle.

CLAVICULAR, *Clavicula'ris*, (F.) *Claviculaire*; same etymon. Relating to the clavicle or collar-bone.

CLAVICULAR NERVES, *Nervi clavicula'res*. Branches of the fourth cervical nerve, which are distributed to the clavicular region.

CLAVICULE, Clavicle.

CLAVICULI, see Clavicle.

CLAVIS, ('a key,') Clavicle, Key—*c. Anglica*, Key.

CLAVUS, ('a nail,') *Helus*, *Gomphus*. (F.) *Clou*. This word is employed in medicine in various senses. It means, 1. A *Corn*, from its resemblance to the head of a nail. 2. Certain condylomatous excrescences of the uterus. 3. A callous tumour which forms on the white of the eye, and resembles a nail, the *Clavus Oculi* (F.) *Clou de l'œil*. This last, by some, is considered to be synonymous with staphyloma; by others, with staphyloma of the cornea. Also, the penis.

CLAVUS HYSTERICUS, *Monopa'gia*, *Monope'gia*, (F.) *Clou Hystérique*. An acute pain, confined to a small point of the head, described by the sick as resembling that which would be produced by a nail driven into the head. It has been particularly noticed in hysterical females;—hence its name. It is called *O'vum hystericum*, when the pain occupies a greater extent.

CLAVUS SECALINUS, Ergot—*c. Secalis*, Ergot—*c. Siliginis*, Ergot.

CLAYEATER, Geophagist.

CLAYEATING, Geophagism.

CLEANSINGS, Lochia.

CLEARSEEING, *Clairvoyance*.

CLEARWEED, *Pilea pumila*.

CLEAV'AGE, (Sax. *cleofan*, 'to split,') The natural line of separation exhibited by certain substances, as minerals, when subjected to mechanical force. The term has been applied to the separation of muscles into longitudinal and circular striæ, when mechanical violence is used.

CLEAVERS, (*cleave*, 'to adhere,') *Galium aparine*.

CLEAVERS' BEES, *Galium aparine*.

CLEAVEWORT, *Galium verum*.

CLEF, (from *clavis*,) Key—*c. du Crane*, see Wormiana ossa—*c. de Garengot*, Key—*c. à Noix*, see Key—*c. à Pivot*, see Key—*c. à Pompe*, see Key.

CLEFT (cleaved,) Rima, see Monster—*c. Palate*, see Harelip.

CLEIDAGRA, (*cleis*, and *agra*, 'a seizure,') *Cleisagra*.

CLEIDARTHRTIS, (*cleis*, and *arthritis*,) *Cleisagra*.

CLEIDION, (*κλειδιον*,) Clavicle.

CLEIDOCOSTAL, Costoclavicular.

CLEIDOMASTOÏDEUS. Albinus thus designates the posterior portion of the cleidomastoidæus, which he considers a separate muscle. It has been corrupted into *clinomastoïdeus*.

CLEIS, (*κλεις*, gen. *κλειός*,) Clavicle, Key.

CLEIS'AGRA, *Cleid'agra*, *Cleidarthrit'is*, (*cleis*, and *agra*, 'a seizure,') Gout in the clavicle.

CLEITHORION, Clitoris.

CLEITHORIS, Clitoris.

CLEITORION, Clitoris.

CLEITORIS, Clitoris.

CLEMATIS, gen. *Clemat'idis*, (*κλημα*, gen. *κλήματος*, 'a branch of a vine, a tendril,') A genus of plants in the Linnæan system.

CLEMATIS CORYMBOSA, *C. erecta*—*c. Crispa*, see *Clematis vitalba*—*c. Daphnoides major*, *Vinea major*.

CLEMATIS ERECTA, *C. rec'ta seu corymb'osa*, *Clematis erecta*, *Flam'mula Jo'veis*, *Up'right Vir'gin's Bow'er*, (F.) *Clématite droite*. Order, Ranunculaceæ. Sex. Syst. Polyandria Poly-

gynia. The leaves contain an acrid principle. They have been esteemed anti-venereal; and, in the form of powder, have been used as an escharotic.

CLEMATIS FLAMMULA, see *C. vitalba*—*c. Recta*, *C. erecta*,—*c. Sepium*, *c. Vitalba*—*c. Viorna*, see *C. Vitalba*—*c. Virginica*, see *C. Vitalba*.

CLEMATIS VITAL'BA, *C. se'pium seu sylves'tris*, *Vital'ba*, *Vior'na*, *Atra'genè*, *Trav'eller's Joy*, *Com'mon Vir'gin's Bow'er*, (F.) *Clématite*, *Herbe aux gueux*, *Aubevigne*. It has been used in the same cases as *C. erecta*. In infusion it has been applied in cases of itch.

The leaves of **CLEMATIS CRIS'PA**—*C. FLAM'MULA*, *sweet-scented Vir'gin's bow'er*—*C. VIRGINICA*, *com'mon Vir'gin's bow'er*—and *C. VIOR'NA*, *Leath'er-flower*, have similar properties.

CLEMATITE, *Clematis vitalba*—*c. Droite*, *Clematis recta*.

CLEMATITIS ERECTA, *Clematis erecta*.

CLEONE PENTAPHYLLA, *Gynandropsis pentaphylla*.

CLEONIS COLLYR'TUM. A collyrium described by Celsus, composed of equal parts of Samian earth, myrrh, and thus, mixed with white of egg; (called, doubtless, after its inventor.)

CLEONIS GLUTEN. An astrigent formula of myrrh, frankincense, and white of egg.

CLEPSY'DRA, (*κλεψύδρα*, from *κλεπτω*, 'I conceal,' and *ὕδωρ*, 'water,') An instrument contrived by Paracelsus to convey fumigations to the uterus.

CLEPTOMANIA, (*κλεπτω*, 'to steal,') *Kleptomania*.

CLETHRA, (*κληθρα*,) *Alnus*.

CLE'THRA ALNIFO'LIA, *Sweet pep'perbush*, *White al'der*, indigenous. Order, Ericaceæ. The leaves and flowers are diaphoretic and excitant.

CLETORIS, Clitoris.

CLEVES, MINERAL WATERS OF. This spring is a quarter of a league from Cleves, in Westphalia. It contains carbonate and sulphate of iron.

CLICKING, (onomatopœia,) see *Râle sibilant*.

CLIDER, *Galium aparine*.

CLIFFORT'IA ILICIFO'LIA, (after George Clifort, a burgomaster of Amsterdam.) Order, Rosaceæ. A common South African plant, used by the Boers as an emollient expectorant in catarrh.

CLIF'TON, CLIMATE OF. The vicinity of Clifton and of Bristol, England, appears to be the mildest and driest climate in the west of England: and, consequently, the best winter residence in that part of the country, for invalids. It is, also, a favorable summer climate, and is surrounded by numerous places of agreeable resort, suited for those who may pass the season there.

For the mineral waters of Clifton, see Bristol Hot Well.

CLIGNEMENT, (from *κλωω*, 'to depress,') Seardamynus.

CLIGNOTEMENT, Nictation.

CLIMA, gen. *Cl'imatis*, (*κlima*,) Climate.

CLIMAC'TERIC or **CLIMACTERIC**, *Climacter'icus*, *Climat'er'icus*, *Climater'ic*, *Climater'icus*, (*κλιμακτηρ*, 'a step,') (F.) *Climactérique* ou *Climatérique*. A word, which properly signifies 'by degrees.' It has been applied to certain times of life, regarded to be critical; but is now chiefly applied to certain periods of life, at which great changes occur, independently of any numerical estimate of years. Such are the period of puberty in both sexes; that of the cessation of the menses in women, &c.

CLIMACTERIC YEARS, *Anni Climacter'ici*, are, according to some, all those in the life of man, which are multiples of the number 7, *Septen'niads*.

Others have applied the term to years, resulting from the multiplication of 7 by an odd number. Some have admitted only three *climacterics*; others, again, have extended them to multiples of 9. Most, however, have considered the 63d year to be the *Grand Climacteric*;—63 being the product of the multiplication of 7 by 9, and all have thought that the period of three, seven, or nine, which they respectively adopted, was necessary to the entire renewal of the body; so that there was, at these times, in the economy, none of the parts of which it had previously consisted. The climacteric years have also been called, *Anni hebdomad'ici, scala'res, grada'rii, scan'siles, genethli'aci, natalit'ii, fata'les, erit'ici, decreto'rii, her'ici, &c.* All the notions on the subject are essentially allied to the doctrine of numbers of Pythagoras.

CLIMATAL, Climatic.

CLIMATE, *Cli'ma, Inclina'tio cœ'li*, (F.) *Climat*, (κλίμα, 'a region.') In geography, the word *climate* is applied to a space on the terrestrial globe, comprised between two circles parallel to the equator, and arbitrarily measured according to the length of the days. In a hygienic point of view, it means, since Hippocrates, a country or region, which may differ from another in respect to season, qualities of the soil, heat of atmosphere, &c. Climate, indeed, embraces, in a general manner, all the physical circumstances belonging to each region,—circumstances which exert considerable influence on living beings. The dark complexion of the inhabitants of the torrid zone is easily distinguishable from the paleness of those of the frigid,—so are the diseases. They are all modified, more or less, by climate or locality. Hot climates predispose to abdominal complications in febrile affections; cold climates to thoracic, &c.

One of the most important considerations with regard to climates is their comparative fitness for the residence of invalids, and especially of those who are liable to, or suffering under, bronchitic or consumptive affections. The great object, in such cases, is to select a climate which will admit of regular and daily exercise in the open air, so that the invalid may derive every advantage which this form of revulsion is capable of effecting. To an inhabitant of the northern and middle portions of the United States—and the same applies to Great Britain, France, and the northern parts of the old world—a more southern climate alone has generally been considered to afford these advantages in an eminent degree. During the summer months there are few, if any, diseases, which require a milder climate than that of the United States, or of the milder districts of Europe. The temperature of the winter months is, consequently, the most important object of attention. Equability of temperature

is essential, inasmuch as all sudden changes interfere with the great desideratum—exercise in the open air. In the whole continent of North America the changes are very sudden and extensive. It is not uncommon for the range to be 40°, between two successive days. So far, therefore, as this applies, the American climate is not well adapted to the invalid. In the southern portions, however, of the Union, this objection is counterbalanced by many advantages; and in the North there are many situations which afford a favorable residence to the valetudinarian during the winter. Observation has, indeed, shown, that in Minnesota, at Mackinac, and in Canada, where the air, although very cold, is dry, a sojourn there has proved at least as efficacious as in the warmer climate of the southern portion of the United States, or of Cuba; and hence, many invalids, instead of subjecting themselves to the inconveniences of temporary expatriation, pass the winter, often advantageously, in Minnesota, for example. From the Army Reports it would seem, that the dry air of New Mexico is especially favorable to those who are threatened with consumption.

The following tables exhibit the mean temperature of the year, and of the different seasons—with the mean temperature of the warmest and coldest months at different places in America, Europe, Africa, &c., as deduced from the excellent paper of Von Humboldt on Isothermal Lines, the Meteorological Registers kept by the surgeons of the United States army, under the direction of Surgeon-generals Lovell and Lawson, 1822 to 1860, the work of Sir James Clark on Climate, &c., &c. Recent, full, and, doubtless, accurate details, as regards the climate of different portions of the United States, are contained in the "Army Meteorological Register for Five Years, from January, 1855, to January, 1860"—an appendix to the "Statistical Report on the Sickness and Mortality in the Army of the United States," from January, 1855, to January, 1860: prepared under the direction of Surgeon-general Lawson, by Dr. R. H. Coolidge.

Certain of the following tables show the mean monthly temperature, maximum, minimum, and range, as well as the greatest daily, and mean daily range during the corresponding months—but of different years—at some of the prominent retreats for the valetudinarian in Great Britain, on the continent of Europe, and in the African islands. It is proper, however, to remark, that in no situations, except in those to which an asterisk is affixed, was the register thermometer used. In the others, the observations were made during the *day* only, and consequently the numbers given are far below the real range throughout the twenty-four hours. The places are ranged in the order of their mean temperature.

TABLE OF MAXIMUM, MINIMUM, AND RANGE OF TEMPERATURE.

PLACES.	DECEMBER			JANUARY.			FEBRUARY.			MARCH.			APRIL.		
	Max.	Min.	Range	Max.	Min.	Range	Max.	Min.	Range	Max.	Min.	Range	Max.	Min.	Range
Sidmouth	54	25	29	47	21	26	52	27	25	56	26	30	60	31	29
Penzance	56	34	22	54	28	26	55	33	22	59	34	25	62	36	26
Pau	56	25	31	56	21	35	60	35	25	65	35	30	71	43	28
Montpellier	57	32	25	53	27	26	55	30	25	58	35	23	64	41	23
Nice	59	40	19	58	27	31	58	37	21	65	41	24	69	46	23
Rome	60	31	29	58	29	29	60	33	27	65	37	28	74	44	30
Naples	61	34	27	58	29	29	60	31	29	69	38	31	78	43	35
Madeira	68	52	16	69	50	19	68	51	17	69	51	18	72	55	17

TABLE OF MEAN TEMPERATURE.

PLACES.	DECEMBER.	JANUARY.	FEBRUARY.	MARCH.	APRIL.
Sidmouth.....	43.00	36.30	42.00	45.00	51.00
Penzance.....	46.50	43.00	44.50	46.50	48.50
Pau.....	41.53	38.89	44.96	46.80	55.79
Montpellier.....	46.00	42.00	45.00	47.00	53.00
Nice.....	48.60	45.85	49.00	51.45	57.00
Rome.....	49.62	47.65	49.45	52.05	56.40
Naples.....	50.50	46.50	48.50	52.00	57.00
Madeira.....	60.50	59.50	58.50	61.06	62.50

TABLE OF DAILY RANGE OF TEMPERATURE.

PLACES.	DECEMBER.		JANUARY.		FEBRUARY.		MARCH.		APRIL.	
	Mean daily range.	Greatest daily range.	Mean daily range.	Greatest daily range.	Mean daily range.	Greatest daily range.	Mean daily range.	Greatest daily range.	Mean daily range.	Greatest daily range.
Sidmouth.....		13		13		12		12		13
Penzance.....	3		4		6		8		9	
Pau.....	7	13	7	16	9	16	9	17	8	18
Montpellier.....	9		8		9		14		14	
Nice.....	6	14	8	16	9	18	9	17	11	18
Rome.....	9	15	11	16	10	18	12	19	13	20
Naples.....	9	13	9	14	11	19	11	18	14	20
Madeira.....	11	14	11	17	9	13	10	14	9	13

1. AMERICA, &c.

PLACES.	Latitude.	Mean temperature of several years.	Mean temperature of different seasons.				Mean temperature of	
			Winter.	Spring.	Summer.	Autumn.	Warmest month.	Coldest month.
Nain.....	57° 08'	26° 42'	0° 60	23° 60	48° 38	33° 44	51° 80	11° 20
Fort Brady, Mich...	46.39	41.37	14.09	37.89	61.83	43.94	62.87	12.65
Quebec, L. C.....	46.47	41.74	14.18	38.04	68.00	46.04	73.40	13.81
Eastport, Me.....	44.54	42.44	23.44	38.58	60.54	45.43	63.52	20.91
Fort Howard, Mich..	44.40	44.50	20.82	41.40	68.70	45.18	73.67	17.95
Fort Crawford, Miss.	43.03	45.52	23.76	43.09	69.78	46.74	71.34	20.14
Cambridge, Mass....	42.21	50.36	33.98	47.66	70.70	49.82	72.86	29.84
Council Bluffs, Mo..	41.25	50.82	27.38	46.38	72.84	48.60	75.92	27.19
Newport, R. I.....	41.30	51.02	33.82	46.87	68.70	53.83	71.46	32.14
Philadelphia.....	39.56	53.42	32.18	51.44	73.94	56.48	77.00	32.72
New York.....	40.40	53.78	29.84	51.26	79.16	54.50	80.78	25.34
Cincinnati.....	39.06	53.78	32.90	54.14	72.86	54.86	74.30	30.20
Monticello, Va.....	37.58	55.40	37.67	54.67	73.33	56.50	75.00	36.00
Washington, D. C...	38.53	55.56*	36.80	53.83	75.90	56.59	79.13	34.66
Smithville, N. C....	34.00	58.88	53.44	64.76	80.46	68.15	82.93	50.69
Charleston, S. C....	32.47	60.18	51.09	66.73	80.89	67.55	82.81	49.43
Natchez, Miss.....	31.34	64.76	48.56	65.48	79.16	66.02	79.70	46.94
Pensacola, Flor.....	30.28	68.77	55.13	69.67	82.57	69.05	83.55	53.80
St. Augustine, do....	29.48	72.23	59.29	71.47	82.73	75.15	83.94	56.60
Tampa Bay, do....	27.57	72.37	61.24	72.93	80.14	75.28	80.72	58.70
Vera Cruz.....	19.11	77.72	71.96	77.90	81.50	78.62	81.86	71.06
Havana.....	23.10	78.08	71.24	78.98	83.30	78.98	83.84	69.98
Bahamas.....	26.40							
to		78.3*	71.	77.	83.	80.	90.	64.
Barbadoes.....	27.5							
Cumana.....	13.10	79.3	76.7	19.	81.	80.		
	10.27	81.86	80.24	83.66	82.04	80.24	84.38	79.16

St. Louis, Missouri, Lat. 38° 46'. Mean temp. 55° 86. New Harmony, Lat. 38° 11'. Mean temp. 56° 74.
 New Orleans, Lat. 30°. Mean temp. 69° 01. Baton Rouge, Lat. 30° 26'. Mean temp. 68° 07.
 Jamaica coast, Mean temp. 80° 6.

RANGE OF MEAN TEMPERATURE IN THE U. STATES FOR THE EXTREME MONTHS, SEASONS, AND YEAR.
(Army Meteorological Register, 1855.)

STATIONS.	YEARS.	Latitude.	January.	July.	Spring.	Summer.	Autumn.	Winter	Yea.
Hancock Barracks, Maine.....	17	46° 07'	16.0	8.7	6.4	6.8	7.2	7.1	5.0
Fort Sullivan, Maine.....	25	44 54	14.0	9.9	7.7	6.8	4.7	7.8	3.7
Fort Constitution, New Hampshire.....	25	43 04	14.0	9.0	7.0	7.4	5.7	11.9	6.7
Fort Independence, Massachusetts.....	17	42 20	9.9	8.6	7.8	4.9	6.9	11.0	4.7
Fort Columbus, New York.....	33	40 42	12.6	11.8	7.8	7.1	9.9	10.3	7.3
Watervliet, New York.....	31	42 43	13.3	10.5	8.8	6.7	9.2	12.8	7.4
Alleghany Arsenal, Pennsylvania.....	22	40 32	18.9	14.5	12.9	7.8	12.0	11.2	7.8
Fort McHenry, Baltimore, Md.....	24	39 17	14.2	6.8	8.9	6.0	7.2	11.2	4.9
Fort Monroe, Virginia, (Norfolk,).	30	37 00	16.9	7.4	10.1	5.9	7.6	16.0	9.0
Fort Moultrie, S. C., (Charleston.).....	28	32 45	18.3	6.1	7.1	7.3	9.5	16.7	6.9
Fort Marion, St. Augustine, Florida.....	20	29 48	16.1	8.6	11.3	6.7	7.1	15.4	6.7
Key West, (Florida,).	14	24 32	12.6	5.9	3.5	3.2	4.2	8.2	3.5
Fort Brooke, Florida.....	25	28 00	17.7	6.8	6.8	5.4	4.8	11.9	4.0
Fort Barrancas, Pensacola, Florida.....	17	30 18	18.2	5.3	7.1	3.2	4.5	9.7	3.0
Mount Vernon Arsenal, Mobile, Ala.....	14	31 12	14.2	6.1	4.7	4.3	5.6	6.8	3.5
Fort Pike, Louisiana.....	14	30 10	10.7	6.2	6.2	3.0	7.0	13.3	4.8
New Orleans, Louisiana.....	20	29 57	18.0	6.8	7.4	5.2	5.5	8.9	4.7
Baton Rouge, Louisiana.....	24	30 26	15.4	7.1	9.3	4.0	6.9	11.1	4.2
Fort Jessup, Louisiana.....	23	31 33	17.5	7.8	9.0	4.6	7.6	10.6	6.6
Fort Gibson, Indian Territory.....	27	34 47	19.5	7.8	9.2	8.6	11.3	14.4	6.9
Jefferson Barracks, Missouri.....	26	38 28	17.6	12.0	15.0	9.1	10.2	19.2	7.0
St. Louis, Missouri.....	12	38 40	14.0	7.2	7.2	4.7	7.4	7.3	3.7
Fort Gratiot, Michigan.....	17	42 55	13.6	11.8	11.8	8.2	8.9	10.7	6.1
Fort Mackinac, Michigan.....	24	45 51	12.3	10.3	8.1	5.8	7.9	9.9	6.1
Fort Brady, Michigan.....	31	46 30	13.0	13.3	13.0	8.5	9.8	9.0	7.1
Fort Howard, Wisconsin.....	21	44 30	14.5	11.8	9.2	7.6	9.9	16.6	5.1
Fort Crawford, Wisconsin.....	19	43 05	19.6	10.4	17.7	8.1	9.2	16.4	8.9
Fort Armstrong, Illinois.....	11	41 30	18.5	10.3	6.8	8.3	6.2	17.4	7.5
Fort Snelling, Minnesota.....	35	44 53	27.9	14.6	17.3	11.0	10.4	16.0	8.6
Fort Leavenworth, Kansas.....	24	39 21	24.7	10.0	17.8	8.0	9.4	13.3	8.0
Fort Kearney, Nebraska.....	6	40 38	19.9	5.2	5.8	4.1	7.0	13.0	5.3
Fort Laramie, Nebraska.....	6	42 12	13.2	4.5	6.4	2.6	10.8	4.8	5.8
Fort Brown, Texas.....	7	25 54	10.0	2.9	2.5	2.2	3.1	4.2	1.6
Fort McIntosh, Texas.....	6	27 31	10.2	3.7	4.3	5.5	3.1	3.6	1.9
Santa Fé, New Mexico.....	5	35 41	4.3	6.9	2.3	1.8	5.0	3.7	0.8
San Diego, California.....	5	32 42	3.7	6.9	40.7	2.7	3.6	2.6	2.3
Benicia, California.....	6	38 03	6.9	4.9	4.6	2.9	3.4	3.2	3.2
Fort Vancouver, Oregon.....	6	45 40	10.1	3.4	4.0	2.5	2.2	2.3	1.4
Fort Steilacoom, Washington Ter.....	6	47 10	12.6	3.8	2.3	1.8	2.8	4.0	2.0

2. EUROPE, AFRICA, &c.

PLACES.	Latitude.	Mean tem- perature of several years.	Mean temperature of different seasons.				Mean temperature of	
			Winter.	Spring.	Summer.	Autumn.	Warmest month.	Coldest month.
Geneva.....	48° 12'	49° 28'	34° 70	47° 66	64° 94	50° 00	66° 56	34° 16
Gosport.....	48 1	50 24*	40 44	47 63	62 00	50 88		
Newport, Isle of Wight.....	50 40	51 00	40 31	49 00	63 09	51 63		
Sidmouth.....		52 10	40 43	50 66	63 83	53 50		
Penzance.....	52 11	51 80	44 03	49 63	60 70	53 36		
Undercliff.....		51 11	42 14	29 26	60 28	52 76		
Hastings.....	50 52	57 00	40 11	45 77	60 45	51 00		
Bute.....	55 42	48 25	39 62	46 66	58 02	48 59		
Cove of Cork.....	51 54	51 58	43 90	49 43	61 26	51 73		
Jersey.....	49 13	53 06	43 82	50 97	62 84	54 63		
Paris.....	45 50	51 08	38 66	49 28	64 58	51 44		
Pau.....	43 7	54 95	41 79	54 96	67 41	55 64	65 30	36 14
Sienna.....	43 24	55 60	40 50	54 10	70 80	57 10		
Nantes.....	47 13	55 62	42 23	53 10	70 73	56 41		
Bordeaux.....	44 50	56 48	42 08	56 46	70 88	56 30	70 52	39 02
Montpellier.....	43 36	57 60	44 20	53 33	71 30	61 30	73 04	41 00
Avignon.....		58 20	42 60	57 13	74 66	60 00		
Florence.....	43 46	59 00	44 39	56 00	74 00	60 70		
Nice.....	43 42	59 48	47 82	56 23	72 26	61 63		
Marseilles.....	43 17	59 50	45 50	57 56	72 50	60 08		
Toulon.....	43 07	59 90	43 30	53 70	74 30	59 00		
Leghorn.....	43 33	60 00	46 30	57 60	74 10	62 00		
Genoa.....	44 25	60 37	44 57	58 60	74 03	62 94		
Pisa.....	43 43	60 60	46 03	57 20	75 15	62 80		
Rome.....	41 53	60 40	45 86	57 74	75 20	62 78	77 00	42 26
Naples.....	40 54	61 40	48 50	58 50	70 83	64 50		
St. Michaels, Azores.....	37 47	62 40	57 83	61 17	68 33	62 33		
Cadiz.....	36 32	62 88	52 90	59 53	70 43	65 35		
Madeira, Funchal.....	32 37	64 56	59 50	62 20	69 33	67 23		
Algiers.....	36 48	69 98	61 52	65 66	80 24	72 50	82 76	60 08
Canaries, Santa Cruz.....	28 23	70 94	64 65	68 87	76 68	74 17		
Cairo.....	30 02	72 32	58 46	73 53	85 10	71 42	85 82	56 12

London, Lat. 51° 30'. Mean temp. 50° 36.
Perpignan, Mean temp. 59° 54.
Nismes, Mean temp. 60° 26.

Environs of London. Mean temp. 48° 81.
Lyons, Mean temp. 55° 76.

In the United States, one of the most favourable regions for the phthisical invalid was considered to be Florida. St. Augustine was frequently chosen, but it is liable to north-east storms, which interfere with the out-door movements of the valetudinarian, and are the source of much discomfort. Still, great benefit has often been derived from it as a winter residence. Of the Atlantic Isles, Madeira appears to be best adapted for the consumptive, and those affected with chronic bronchitis. In Italy, Rome, and Pisa, and in England, Torquay and Undercliff, are to be preferred. Chronic rheumatism and gout are benefited by a warm climate, which, again, is unfavourable to those who are predisposed to cerebral diseases, especially to such as are characterized by debility and mobility of the nervous system—as paralysis, epilepsy, mania, &c. Hypochondriasis and dyspepsia require rather change of climate and travelling exercise than a sojourn in any one. (See the Author's Human Health, Philada., 1844).

CLIMATERIC, Climacteric.

CLIMATIC or CLIMATIC, *Climaticus*, *Climatal*. Belonging to, or dependent upon climate.

CLIMATIC DISEASES, *Morbi climatici*, are such as are caused by climate.

CLIMATION, Acclimation.

CLIMATOLOGY, *Climatologia*, (F.) *Climatologie*. (κλίμα, 'a region,' and λόγος, 'a discourse.') A treatise on climates, their effects on organized bodies, &c., &c.

CLIMBER, WOODY, *Ampelopsis quinquefolia*.

CLINE, (κλινη, 'a bed.') Hence:

CLINIATER, (*cline*, and ιατρος, 'a physician.') Clinical physician.

CLINIATRIA, (*cline*, and ιατρεία, 'the healing art.') Clinical medicine.

CLINIATRUS, Clinical physician.

CLIN'IC, *Bedridden*. One confined to bed by sickness. See, also, *Clinique*.

CLINICA, Clinical medicine.

CLIN'ICAL, (κλινικος, from *cline*), *Clin'icus*, (F.) *Clinique*. Relating to a bed.

CLINICAL LECTURE, (F.) *Leçon Clinique*. One given at the bed-side, or on a particular case or cases.

CLINICAL MEDICINE, *Cliniatri'a*, *Clin'ica*, *Clin'icæ*, (F.) *Médecine clinique*. That which is occupied with the investigation of diseases at the bed-side, or individually.

CLINICAL PHYSIC'IAN, *Cliniater*, *Cliniaterus*, (F.) *Clinicien*. One who practices or teaches clinical medicine.

CLINICE, Clinical medicine.

CLINICIEN, Clinical physician.

CLINICUM, *Clinique*.

CLINIQUE (F.), Clinical. *Clinique*, *Clin'icum*, is also used substantively for *École Clinique* or *Clinical School*: a school in which medicine is taught by examining diseases on the patients themselves before the class. Thus, the French say, — 'La Clinique de la Charité,' 'The Clinical School of the Hospital La Charité.' The term has been introduced into this country, and anglicised *Clin'ic*.

CLINOCEPHALIA, *Klino-cephalia*, (*cline*, and κεφαλή, 'head,') Monstrosity in which the head is unusually flattened, owing to synostosis of the parietal and sphenoid bones.

CLINODES, (*cline* and *odes*), *Clinoid*.

CLINOID, *Clinoides*, *Clinodes*, *Clinoides*, (*cline*, and εἶδος, 'form,') Resembling a bed.

CLINOID PROC'ESSES, (F.) *Apophyses clinoides*, are four processes at the upper surface of the sphenoid bone, which have been compared to the posts of a bed. On them the pituitary gland rests. They are divided into *anterior* and *posterior*.

Each of the anterior terminates in a point called *Transverse Sphenoid Process*.

CLINOIDEUS, *Clinoid*.

CLINOPODIA, *Thymus mastichina*.

CLINOPODIUM ARVENSE, (*cline*, and ποῦς, gen. ποός, 'a foot,' so called from the shape of its flowers,) C. vulgare — c. Incanum, *Pycnanthemum incanum* — c. Majus, c. vulgare.

CLINOPODIUM VULGARĒ, *Clinopodium arvense* seu *ma'jus*, *Oc'lymum sylvæstris*, *Melissa clinopodium*, *Thymus sylvaticus*, *Wild Basil*, (F.) *Grand Basilic sauvage*. A European plant, *Ord. Labiata*, which was formerly considered to be an antidote to the bites of venomous animals, to facilitate labour, relieve strangury, stop diarrhoea, &c.

CLIQUE'TIS, (*cliquer*, 'to clack, to click,') Crepitation — c. *Métallique*, *Tintement métallique*.

CLISEOMETER, *Clisio'm'eter*, (F.) *Klisiéomètre*, (κλίσις, 'inclination,' and μέτρον, 'a measure,') An instrument, intended to measure the degree of inclination of the pelvis and to determine the relation between the axis of the pelvis and that of the body.

CLITBUR, *Arctium lappa*.

CLITHE, *Arctium lappa*.

CLITHEREN, *Galium aparine*.

CLITORICARCINO'MA, (*clitoris*, and καρκίνωμα, 'cancer,') Cancer of the clitoris.

CLITORIDAUX'Ē, (*clitoris*, and αυξη, 'augmentation,') Enlargement of the clitoris.

CLITORIDES INFERIORES LATI ET PLANI MUSCULI, *Constrictores cunni*.

CLITORIS, gen. *Clit'idis*, *Clit'oris*, (κλητωρ, 'a servant who invites guests' [?]) *Cleit'oris*, *Cleit'oris*, *Cleit'horis*, *Cleit'orion*, *Cleit'horion*, *Dulcedo Amoris*, *Venus*, *Myr'ton*, *Mur'ton*, *Mur'tum*, *Es'trum*, *Es'trus*, *Col'es* seu *Colis femine'rum*, *Penis mul'eb'ris* seu *femine'us*, *Men'tida mul'eb'ris*, *Mem'brum mul'eb're*, *Superla'b'ia*, *Cerco'sia*, *Hypoder'mis*, *Nym'pha*, *Nym'phē*, *Epider'rhis*, *Lib'i'dinis* *Sc'des*, *Tunic'ula*, *Cris'ta*. A small round organ, situate at the upper part of the vulva, and separated by a small space from the anterior commissure of the labia. Its base is surrounded by a slight fold, similar to the prepuce; and it is, internally, of the same structure as the *corpora cavernosa penis*. The artery of the clitoris, (F.) *A. Clitorienne* ou *Clitoridienne*, is a branch of the internal pudic. The vein communicates with the internal pudic, and the nerve with the pudic.

CLITORISM, *Clitoris'mus*. The abuse made of the clitoris. Also, an unusually large clitoris, *Cerco'sis extern'a* seu *Clitor'idis*, *Cauda'tio*, *Cauda*.

CLITORITIS, *Clitoriti'tis*, (*clitoris*, and *itis*,') Inflammation of the clitoris.

CLITORITITIS, *Clitoritis*.

CLITORIUM, *Vulva*.

CLIVEE, (*cleave*, 'to adhere,') *Galium aparine*.

CLIVUS BLUMENBACH'II, (*clivus*, 'a slope,') An inclining surface behind the dorsum ephippii of the sphenoid bone, which is continuous with the basilar process of the occipital bone.

CLOA'CA, (F.) *Cloaque*, (akin to κλύω, 'I wash' [?], 'a common sewer,') The pouch at the extremity of the intestinal canal, in which the solid and liquid excretions are commingled in birds, fish and reptiles. In the male, it gives exit to the excrements, sperm and urine: in the female, to the eggs, fecal matters, and urine.

CLOANX, *Orpiment*.

CLOAQUE, *Cloaca*.

CLOCHE, (F.), 'a bell.' A popular expression in France for a blister or other vesicle.

CLOISON, (*claudere*, *clausum*, 'to shut,') Septum — c. *des Fosses nasales*, *Septum narium* — c. *Transparente*, *Septum lucidum*.

CLOISONNEMENT, see Generation. Also, the division of a hollow organ — as the bladder,

vagina, and uterus, into two portions by a septum, — a malformation — *anomalie par cloisonnement*.

CLON'IC, *Clon'icus*, (κλονος, 'tumultuous movement.') (F.) *Clonique*. Irregular convulsive motions. Convulsion with alternate relaxation: in contradistinction to *tonic*, which signifies a constant rigidity; — *Clonus*, *Clonospas'mus*, *Clon'ici partiid'les*.

CLONIC SPASM, see **Spasm**.

CLONICI UNIVERSALES, *Synclonus*.

CLONIQUE, *Clonic*.

CLONIS, (κλονις,) *Sacrum*.

CLON'ISM, *Clonis'mus*. *Clonic spasm*.

CLONO'DES, (κλονος, 'tumultuous movement,' and *odes*.) *Convulsive*. Galen applies this term to the pulse, when convulsive, as it were, and unequal. He compares the sensation it communicates to the finger to that produced by a bundle of sticks or of rods in vibration.

CLONOS EPILEPSIA, *Epilepsy* — *c. Hydrophobia*, *Hydrophobia*.

CLONOSPAMUS, (κλονος, 'agitation,' and *spas-mus*, 'spasm.') See **Clonic**.

CLON'US, (κλονος, 'tumultuous movement.') *Synclonus* — *c. Nictitatio*, *Nictation* — *c. Palpitation*, *Palpitation* — *c. Pandiculatio*, *Pandiculation* — *c. Pandiculatio maxillarum*, *Yawning* — *c. Singultus*, *Singultus* — *c. Sternutatio*, *Sneezing* — *c. Sub-sultus*, *Subsultus tendinum*.

CLOPEMANIA, (κλοπη, 'theft,' and *mania*.) *Kleptomania*.

CLOPORTES ORDINAIRES, *Onisci aselli*.

CLOSESTOOL, *Lasanum*.

CLOT, ([D.] *kloot*, 'a ball;' *kluut*, 'a clod.')

Coagulum — *c. of Blood*, see **Blood**.

CLOT, HEART, see **Polypus**.

CLOTBUR, *Xanthium*.

CLOTHING, *Vestitus*.

CLOT'TY, *Grumo'eus*, (F.) *Grumeleux*. Composed of clots.

CLOU, *Clavus*, *Furunculus* — *c. de Girofle*, see *Eugenia caryophyllata* — *c. Hystérique*, *Clavus hystericus* — *c. de l'Œil*, *Clavus oculi* — *c. de Seigle*, *Ergot*.

CLOUDBERRIES, *Rubus chamæmorus*.

CLODBERRY TREE, *Rubus chamæmorus*.

CLOVE, (*clavus*, 'a nail,') see *Eugenia caryophyllata* — *c. Bark*, see *Myrtus caryophyllata* — *c. July flower*, *Dianthus caryophyllus* — *c. Pink*, *Dianthus caryophyllus*.

CLOVE-TONGUE, *Helleborus niger*.

CLUBFEET, *Kyllosis*.

CLUBMOSS, *Lycopodium* — *c. Common*, *Lycopodium complanatum* — *c. Fir*, *Lycopodium selago* — *c. Upright*, *Lycopodium selago*.

CLUNES, (pl. of *clunis*, 'the buttock.') *Nates*.

CLUNE'SIA, (*clunes*, 'the nates.') *Proctal'gia*, *Proctitis*. An inflammation of the buttocks.

CLUNIS, *Sacrum*, see **Clunes**.

CLUPEA THRY'SA. The *Yellow-billed Sprat*; a very poisonous fish of the West Indies.

CLUTIA ELUTERIA, (after T. A. Cluyt, (L.)

Clutius, an apothecary of Leyden.) *Croton cas-carilla*.

CLYDON, (κλυδων, 'a wave,') *Flatulencia*.

CLYPEALIS (*Cartilago*.) *Thyroid cartilage*.

CLYSANTLIUM, (κλυζειν, 'to wash away,') to give a clyster, and *πυλινον*, 'a pump or syringe,') *Syringe*, *Clyster*.

CLY'SIS, (κλυσις,) *Clysmus*. The application of a clyster. Washing out by means of a clyster.

CLY'SMA, gen. *Cly'smatis*, (κλυσμα,) *Clysis*, *Clyster* — *c. Tonicum*, *Enema fetidum*.

CLYSMA'TION, (dim. of *clyasma*.) A small clyster.

CLYSMUS, (κλυσμος,) *Clysis*.

CLYSOIR ou **CLYSOIRE** (F.), (from κλυζειν, to wash away,') An apparatus for administer-

ing enemata, consisting of a long funnel-shaped tube of elastic gum, furnished with a pipe at the apex. An instrument, similar to this, with a syringe attached, is called a *Clysopompe*.

CLYSOPOMPÉ, *Clysoir*, *Syringe*.

CLY'STER, (κλυστηρ,) *Clysterium*, *Cly'sma*, *Encly'sma*, *Cly'smus*, *En'ema*, *Lavamentum*, *Lo'tio*, *Glyster*, *Glistér*, *Injection*, (from κλυζειν, 'to wash away,') (F.) *Clystère*, *Lavement*, *Remède*. A liquid, thrown into the large intestines by means of a syringe, or bladder and pipe properly prepared, &c.; the nozzle of the syringe or pipe being introduced into the anus. See **Enema**.

CLYSTER OF ALOES, *Enema aloes* — *c. Anodyne*, *Enema anodynum* — *c. of Asafoetida*, *Enema asafoetida* — *c. Cathartic*, *Enema catharticum* — *c. of Colocynthis*, *Enema colocynthis* — *c. Common*, *Enema commune* — *c. Domestic*, *Enema commune* — *c. Fetid*, *Enema fetidum* — *c. Pipe*, *Elasma* — *c. Purging*, *Enema catharticum* — *c. Starch* and *opium*, *Enema anodynum* — *c. Tobacco*, *Enema tabaci* — *c. Turpentine*, *Enema terebinthinæ* — *c. Uterinus*, *Sparallium*.

CLYSTÈRE, *Enema*.

CLYSTERION, (κλυστηριον,) *Clyster*.

CNEME, (κνημη,) *Leg*, *Tibia*.

CNEMODACTYLEUS, (κνημη, 'leg,' and *δακτυλος*, 'a finger or toe,') *Extensor communis digitorum pedis*.

CNEMOLOORD'SIS, (*cneme*, and *λορδωσις*, 'the state of being bent forward,') *Bending of the leg forward*.

CNEMOSCOLIO'SIS, (*cneme*, and *σκολιωσις*, 'bending,' especially sideways.) *Bending of the leg sideways*. The state of being *bow-legged*, or *ban'dy legged*, (Prov.) *Scrog-legged*, *shackle-ham-med*, (L.) *Valgus*.

CNEORON, (κνεωρος,) *Daphne gnidium*. [?]

CNEORUM TRICOC'CUM, *Almeze'rian*, *Wid'ow-wail*, *Spurge Olive*, (F.) *Camélee*. This plant, *Ord. Terebinthaceæ*, a native of Southern Europe, contains a very irritating acrid principle. The ancients employed its leaves as a powerful purgative. It is now sometimes used for deterring ulcers.

CNESIS, (κνησις,) *Cnes'mos*, *Cnis'mos*. A painful itching.

CNESMA, gen. *Cnesm'atis*, (κνησμα,) *Itching*.

CNESMOS, (κνησμος,) *Cnesis*, *Itching*, *Prurigo*.

CNICELÆ'UM, (*cnicus*, and *ελαιον*, 'oil,') *Oil of carthamus*.

CNICUS, (κνικος,) *Carthamus tinctorius* — *c. Sylvestris*, *Centaurea benedicta*.

CNIDE, (κνιδη,) *Urtica*.

CNIDELÆ'ON, *Cnidelæ'um*, (κνιδελαιον, *cnid'ion*, and *ελαιον*, 'oil,') *Oil made from the grana cnidia or mezereon berries*.

CNID'IA GRA'NA, *Cnid'ii Ooc'ci*, *Cocco-gnid'ia*, *Æto'lion*, *Coc'cum*. The berries of the *Daphne gnidium*.

CNIDIUM SILAUS, *Peucedanum silaus* — *c. Tenuifolium*, *Sison ammi*.

CNIDO'SIS, (κνιδωσις, from *Cnide*, and *osis*.) A pungent itching, compared to that produced by the Nettle. *Urticaria*, *Urticatio*.

CNIP'OTES, (κνιποτης,) *Pruritus*. *Itching*. The dry ophthalmia, *Xerophthal'mia*.

CNISMOREGMIA, (κνισμος, 'a vellication,' and *ορεγω*, 'I put forth,') *Pyrosis*.

CNISMOS, (κνισμος,) *Cnesmos*.

CNISSA, (κνισσα,) see **Nidoros**.

CNISSOREG'MIA, (κνισσα, 'the smell of burnt fat or flesh,' and *ορεγω*, 'I put forth,') *Ruc'tus nidorosus*. A nidorous eructation, as of rotten eggs.

CNY'MA, gen. *Cny'matis*, (κνυμα.) A slight itching. Also, a puncture or vellication.

CO, as a prefix, like *cum*, with.

COA'CON (EMPLASTRUM NIGRUM.) A black plaster, mentioned by Celsus, and composed of litharge and resin,—the litharge being first boiled in oil.

COACTUS, (from *cogo*, *coactum*, (co, and *ago*, *actum*), 'I bring together,') see *Curds*.

COAGMENTATIO, (*coagmentare*, 'to join together,') *Gomphosis*.

COAGULABLE LYMPH, Fibrin, Liquor sanguinis.

COAG'ULANTS, *Coagulan'tia*, (*coagulare*,—(itself from *co*- and *agere*), 'to bring together,') or from *cogo*, *coactum*, 'to drive or bring together,') Remedies, or poisons, supposed to possess the power of coagulating the blood, or to give consistency to animal fluids. The word and the absurd notions connected with it are equally abandoned.

COAGULA'TION, *Coagula'tio*, *Thrombo'sis*. The conversion of a liquid into a more or less soft and tremulous mass. Many animal and vegetable fluids are capable of coagulation.

COAGULATUS, see *Curds*.

COAG'ULUM, *Gru'mus*, (F.) *Caillot*, *Grumeau*, (same etymon as *Coagulants*.) A soft mass formed in a coagulable liquid. The Clot of the Blood is particularly so called—the *Cru'or*, *In'sula*, *Placen'ta*, *He'par Sanguinis*, *Crassamen'tum Sanguis concretus*;—the red mass, composed of fibrin, serum, and colouring matter, which separates when the blood is left to itself. See *Blood*. The word is also applied, in pathology, to the sanguineous concretions, which form in different natural and accidental cavities; and which, when they occur at the mouth of a divided artery, sometimes suspend the flow of blood. This is, indeed, one of the means by which hemorrhage is arrested.

COAGULUM ALU'MINIS, *Coagulum Aluminosum*, *Cataplas'ma aluminis*, *Alum eurd* or *cataplasma*. This is made by beating the white of egg with a little alum, until a coagulum is formed. It is applied in cases of ophthalmia, where an astringent is necessary.

COALES'CENCE, *Coalescen'tia*, *Coalit'io partium*, (*coalescere*, 'to grow together,') *co*, and *alere*, 'to nourish,') *Syn'physis*, *Pros'physis*. The adhesion or union of parts previously separated, as in case of wounds and preternatural adhesions or malformations. See *Monster*.

COALFISH, see *Oleum jecoris aselli*.

COALITIO, *Coalition*, *Symphysis*—c. *Partium*, *Coalescence*.

COALIT'ION, *Coalit'io*, (same etymon as *Coalescence*.) It has been used in the same cases; as well as to express the action of several parts of the frame, which have the same nutrition.

COALITUS, *Symphysis*.

COALTER'NÆ FEBRES, (*co*, and *alternare*, 'to alternate,') A name given to two intermittents, which attack a person at the same time, but whose paroxysms are distinct: so that the attack of one supervenes when the other has ceased. The term *Double Intermittent* expresses the same meaning.

COÄPTATION, *Coapta'tio*, (*co*, and *aptare*, 'to adjust, adapt,') *Parago'ge*. The act of adapting the two extremities of a fractured bone to each other; or of restoring a luxated bone to its place. Coaptation must be effected gently. Usually, extension and counter-extension are, in the first place, necessary.

COARCTATIO, *Aretatio*, *Coarctation*, *Stricture*—c. *Ventriculi*, *Stricture of the Stomach*.

COARCTA'TION, *Coarcta'tio*, (*co*, and *arctare*, *arctatum*, 'to straiten,') *Stricture*. *Avicenna* speaks of *Coarctation of the Pulse*.

COARCTOTOMIE, *Coarctotom'ia*, a hybrid term, (from *coarctare*, 'to straiten,') and *τομή*, 'incision,') *Urethrotomy*, when limited to the simple division of a stricture.

COARTICULATIO, (*co*, and *articulus*, 'a joint,') *Diathrosis*, *Synarthrosis*.

COAT, ([F.] *Cotte*,) see *Enduit*—c. *Buffy*, *Corium phlogisticum*.

COATED, see *Enduit*.

COATING OF PILLS, see *Deaurentur pilulæ*.

COBALT, PROTOXIDE OF, *Smalt*.

COBHAM, MIN'ERAL WATERS OF. *Cobham* is seven miles from Kingston, in Surrey, England. The waters are weak saline purgatives.

COBRA DI CAPELLO, *Cobra*, *C. Capello* (P.), 'serpent of the hood,' (F.) *Serpent à lunettes*. *Hood'ed snake*, *Spee'tacled snake*. A very venomous reptile of India, of the family *Viperidæ*—the *Naja tripudians*.

COBWEB, *Araneæ tela*.

COCAINE, see *Erythroxylon coca*.

COCASH, *Erigeron Philadelphicum*.

COCCARIUM, (κοκκος, 'a pill,') A very small pill.

COCCHIA, *Cochia*.

COCCI GRANUM, *Kermes*—c. *Orientales*, see *Menispermum cocculus*.

COCIGIO-CUTANÉ SPHINCTER, *Sphincter ani externus*.

COCCINEL'LA, (dim. of κοκκος, *coccus*, 'a berry,') which it resembles,') *Coccus cacti*.

COCCINELLA SEPTEMPUNCTATA, *La'dy-bird*, *La'dy-con*, *La'dy-bug*, *Cow-lady*, *Cush'y-cow-la'dy*, (Prov.) *Fly-gold'ing*, *Gold'en-bug*, *Gold'en-knop*. This insect, bruised upon an aching tooth, has long been regarded as anti-dontalgic!

COC'CION, (κοκκιον.) A weight, mentioned by *Myrepsus*; the same as the *siliqua*.

COCCIONELLA, *Coccus cacti*.

COCCOBALSAMUM, see *Amyris opobalsamum*.

COCCOGNIDIA, *Cnidia grana*.

COCCOLOBA UVIFERA, see *Kino*.

COCCONES, (κοκκωνες,) see *Punica granatum*.

COCCULI OFFICINARUM, see *Menispermum cocculus*.

COCCULUS CRISPUS, (dim. of *Coccus*), *Menispermum tuberculatum*—c. *Indi aromaticus*, see *Myrtus pimenta*—c. *Indicus*, see *Menispermum cocculus*—c. *Palmatus*, *Columba*—c. *Suberosus*, *Menispermum cocculus*.

COCCUM, (κοκκος, 'a berry,') see *Cnidia grana*, *Kermes*—c. *Baphicum*, *Kermes*—c. *Infectorium*, *Kermes*—c. *Scarlatinum*, *Kermes*—c. *Tinctorum*, *Kermes*.

COCCUS, (κοκκος, 'a berry,') 'a pill,') *Coccus cacti*, *Pilula*—c. *Americanus*, *Coccus cacti*.

Coc'cus Cac'ti. The systematic name of the *Coch'ineal Insect*, *Coccus*, (Ph. U. S.), *Coccinell'la*, *Cochinil'la*, *Coccinell'la*, *Coccinil'la*, *F'icus Indie grana*, *Coc'cus Polon'icus* seu *America'nus* seu *Ind'icus Tincto'rius*, *Scarabae'olus hemisphaer'icus*, *Cochinilif'era cochinil'la*, *Coc'cus America'nus*, *Cochinelle*; the *Coch'ineal Animal*, (F.) *Cochenille*, *Graine d'Écarlate*. *Class*, *Insecta*. *Order*, *Hemiptera*. The cochineal insects have a faint, heavy odour; their taste is acrid, bitterish, and astringent; colour blackish-red externally, — purple-red within. They are used chiefly for giving a red colour to tinctures, &c. They were, at one time, esteemed astringent, stimulant, diuretic, and anodyne.

Coccus ILICIS, *Kermes*—c. *Indicus*, see *Menispermum cocculus*—c. *Indicus tinctorius*, *Coccus cacti*—c. *Polonius*, *Coccus cacti*.

COCCYALGIA, (*coccyx*, and *αλγος*, 'pain,') *Coccydynia*.

COCCYCEPH'ALUS, *Coccy'goceph'alus*, (F) *Coccycephale*, (*coccyx*, and *κεφαλή*, 'the head,') A monster whose head has the shape of the os coccygis.

COCCYGALGIA, Coccyodynia.

COCCYGEAL, Coccygeus.

COCCYGEAL GLAND, *Glandula coccygea*. A ductless gland, about the size of a pea, situate in front of the top of the coccyx. Its functions are not known.

COCCYGEAL NERVE, see Sacral Nerves.

COCCYGEOTOMY, *Coccygeotomia*, (coccyx, and *exom*, 'excision,') Excision of the coccyx, or of a portion of it.

COCCYGEUS, *Coccygeus*, (F.) *Coccygien*, (inserted into the coccyx;) *Ischio-Coccygeus*, (F.) *Ischio-coccygien*. Belonging both to the ischium and coccyx. The muscle *Ischio-coccygeus*, *Levator Coccygis*, *Triangularis Coccygis*. It arises from the spinous process of the ischium, and is inserted into the extremity of the sacrum; and into nearly the whole length of the os coccygis laterally. It retains the coccyx in place, and prevents it from being forced backwards during the evacuation of the feces.

COCCYGEARTHROSIS, (coccyx, and *exarthrosis*, 'dislocation,') Dislocation of the coccyx.

COCCYGO-ANAL, Sphincter ani externus.

COCCYGIO-CUTANÆ SPHINCTER, Sphincter ani externus.

COCCYGOCEPHALUS, Coccycephalus.

COCCYMELEA, Prunus domestica.

COCCYME'LON, (κοκκυμῶν,) Prune.

COCCYODYN'IA;—better *Coccygodyn'ia*; *Coccygal'gia*, *Coccyal'gia*, (coccyx, and *odyn*, 'pain,') Pain in the os coccygis.

COC'CYX, gen. *Coccy'gis*, (κοκκυξ, gen. κοκκυγος, 'a cuckoo,' whose bill it is said to resemble;) *Os Coccy'gis* seu *Al'agas*, *Cav'da*, *Os'sis* *ac'ri* *acumen*, *Cul'culus*, *Uropy'gium*, *Rump* or *Cru'pper Bone*. An assemblage of small bones, attached to the lower part of the sacrum; the curvature of which it prolongs, and of which it seems to be an appendage. Generally, it consists of four bones. Behind the base of the coccyx are two small tubercular eminences. These are called *Cor'nea of the Coccyx*.

COCHEMAR, Incubus.

COCHENILLE, Coccus cacti.

COCHERIA, Cochia.

COCH'IA, *Coch'chia*, *Coch'e'ria*, (from κοκκος, 'berry,' 'a pill,' or from κοχυω, 'I flow profusely,') An ancient name for several officinal purgative pills.

COCHIN LEG, see Elephantiasis.

COCHINEAL, (from *coccinella*,) Coccus cacti.

COCHINELIFERA COCHINILLA, Coccus cacti.

COCHINILLA, Coccus cacti.

COCH'LEA, 'a snail's shell,' (κοχλίας, 'a snail,') (F.) *Limaçon*, *Coquille*, *Cochléc*. Anatomists have given this name to the most anterior of the three cavities, which constitute the labyrinth of the ear, the *Pel'vis An'trium*, *Con'cha au'ris inter'na* seu *Labyrinthi*, *Cav'itas cochlea'ta* seu *buccina'ta*, *An'trium buccino'sum*, *Troch'lea labyrinthi*;—and that of *Sea'ta of the Coch'lea*, (F.) *Rampes du limaçon*, to two spiral cavities in the interior of the cochlea. One of these scale terminates at the *Fene'stra rotun'da*, and is called *Sea'ta tym'pani*: the other opens at the anterior and inferior part of the vestibule: it is called *Sea'ta vesti'buli*.

COCHLEA, SCALE OF THE, see Cochlea.

COCH'LEAR, *Cochlea'ria*, (F.) *Cochl'éaire*, *Cochl'éarien*. Relating or appertaining to the cochlea.

COCHLEAR, Cochleare—c. Auriculare, Ear-pick—c. Nerve, see Auditory Nerve.

COCHLEAR'È, *Coch'lear*, *Cochlea'rium* (from *cochlea*; its bowl resembling a shell.) A spoonful, (F.) *Cuillèrte*; abbreviated in prescriptions usually to *coch*. See Abbreviation. Also, a scoop.

COCHLEARÈ MAGNUM, a tablespoonful; *C. Me'*

dium, a desert or pap-spoonful: and *C. Min'im'um*, a teaspoonful.

COCHLEA'RIA, (cochleare, 'a spoon,' so called from the resemblance of its leaves.) *C. officinalis*.

COCHLEARIA ARMORA'CIA, *Raph'anus rusticus* seu *mar'inus* seu *ajac'es'tris*, *Armora'cia*, *A. sati'va* seu *rustica'na*, *Horse'radish*; (Sc.) *Rot-coll. Fam.*, Cruciferae. *Sex. Syst.* Tetradynamia Siliculosa. (F.) *Raifort sauvage*, *Cran*, *Cran de Bretagne*, *Crançon*, *Moutarde des Moines* ou *des Allemands*, *Radis de cheval*. The root of horseradish is frequently used at table; and has long been in the *Materia Medica*. It is stimulant and diuretic. Externally, it is rubefacient.

COCHLEARIA CORON'OPUS, *Coron'opus*, *Coron'opus* *Ruel'lii* seu *depressus* seu *vulgar'is*, *Lepid'ium squama'tum*, *Senebie'ra coron'opus*, *Wild Scur'vy Grass*, *Swine's Cress*, (F.) *Corne de Cerf*. This European plant is considered to be diuretic and antiseptic. The term *Coron'opus* was given, by the ancients, to various plants.

COCHLEARIA HORTENSIS, Cochlearia officinalis.

COCHLEARIA OFFICINA' LIS, *Cochlea'ria*, *C. horten'sis* seu *pyrena'ica* seu *vulgar'is*, *Lem'on Scur'vy Grass*, *Com'mon Scur'vy Grass*, (Sc.) *Scrub-grass*, (Prov.) *Scrooby-grass*, (F.) *Crançon*, *Herbe aux cuillers*. It has been considered a powerful antiscorbutic. It is sometimes eaten in salad.

COCHLEARIA PYRENAICA, *C. officinalis*—c. *Vulgaris*, *C. officinalis*.

COCHLEARIFORM'IS, (cochleare, 'a spoon,' and *forma*, 'shape,') (F.) *Bec de Cuiller*. Spoon-shaped.

COCHLEARIFORMIS PROCES'SUS, *Cochlear'iform process*. A small, very thin plate, which separates the bony portion of the Eustachian tube from the canal for the passage of the tensor tympani.

COCHLEA'RIS. A gelatinous-looking tissue, seen on opening the cochlea, by which the membranous zone is connected, at its outer or convex margin, with the outer wall. It is supposed by Todd and Bowman to be muscular; and to have a preservative office, being placed to defend the cochlear nerves from undue vibrations of sound, in a way analogous to that in which the iris protects the retina from excessive light.

COCHLEARIUM, Cochleare.

COCHON'È, (κοχων.) The junction of the ischium, near the seat or breech. The breech proper, from the hip-bones to the anus. The perineum. The coccyx.

COCIL'IO. A weight of eleven ounces.

COCK, (onomatopœia,) Phasianus gallus.

COCYEYE, see Luscitas.

COCKGRASS, *Lolium temulentum*.

COCKLE, CORN, *Lychnis githago*.

COCKLE-BUR, *Agrimony*, *Xanthium*.

COCKLES, INDIAN, see *Menispermum cocculus*.

COCKMINT, *Tanacetum balsamita*.

COCKSPUR, Ergot—c. Rye, Ergot.

COCKUP HAT, *Stillingia*.

COCKYBABY, *Arum maculatum*.

COCLES, ('blind of one eye,') *Borgne*.

COCO, *Cocos nucifera*.

Co'co of the MALDIVES!, *Co'cos de Mald'ivá*. The fruit of a palm, called *Lodoice'a* by Comerson. It was formerly termed, in the shops, *Nux Med'ica*, and enjoyed great reputation.

COCOA, Cacao, *Cocos nucifera*—c. *Cacavifera*, Cacao.

COCOBAY, Mal de San Lazaro.

COCO-OLEIN, see *Cocos nucifera*.

COCOS, see *Arum esculentum*.

Co'cos BUTYRA'CEA. The name of the plant which affords the palm oil, *O'leum pal'mæ*, obtained chiefly from the fruit, by bruising and mixing the kernels with water, without the aid

of heat, by which the oil is separated and rises to the surface. It is of the consistence of an ointment, with little taste, and a strong, though not disagreeable, smell. It is applied to sprains, &c.: but has no advantage over other oils. It has been used as the excipient for suppositories, and has been called, also, *O'leum Pal'mæ sebæ'ceum*, *O. fir'm nū'eum co'cos butyra'ceæ* and *Mackaw' fat*. It is procured likewise from the *Elæ'is Guineen'sis*, and *Elæ'is Occidentā'lis*, two species of palms. (Cocos is said to be contracted from (P.) *macoco* or *macoco*, 'a monkey,' the three holes at the end of the cocoa-nut shell causing it to resemble somewhat a monkey's head.)

COCOS NUCIF'ERA, *Pal'ma cocos*, (F.) *Cocotier*. Order, Palmæ. The systematic name of the plant whose fruit is in the *co'coa-nut*. (F.) *Coco*. It has an agreeable kernel, but not easy of digestion. *Emulsions, orgeat'*, &c., are made from it. The juice of the cocoa, *Tod'dy*, when fermented, forms wine, and arrack is distilled from it. The expressed oil—*Co'co-o'lein*—has been found almost as efficacious as cod-liver oil in tuberculosis.

COCOTE, Influenza.

COCOTIER, *Cocos nucifera*.

COCOWORT, *Thlaspi bursa*.

COCIOTI, Coction, Digestion—c. Ciborum, Digestion—c. Morbi, Coction.

COC'TION, *Coc'tio*, *Pep'sis*, *Pepan'sis*, *Pepas'mus*, *Symp'ep'sis*, *Concoct'ion*, (*coquere, coctum*, 'to boil.') This word has been employed in various senses. 1. With the ancients, *coction* meant the particular kind of alteration, which the food experiences in the digestive organs, particularly in the stomach. It meant a preparation from its crude state. 2. It expressed the maturation or change, which the humoral pathologists believed morbid matter experiences before elimination. It was considered that coction, *Coc'tio mor'bi*, was produced during the violence of the disease; and hence this was called the *Period of Coction*. See Humorism.

CO'CYTA, (F.) *Cocyste*, (from *κωκυτος*, 'lamentation.') The pain caused by the introduction of poison under the skin.

COD or **CODS**, (Sax. *codde*, 'a bag,') Scrotum.

COD-OIL, *Oleum jecoris aselli*.

COD-LIVER OIL, *Oleum jecoris aselli*—c. l. Olein, see *Oleum jecoris aselli*.

CODAGAM, *Hydrocotyle Asiatica*.

CODAGAPALA BARK, *Nerium antidysentericum*.

CODE, *Codex*.

CODEÏA, *Codeine*.

CO'DEINE, *Codei'na*, *Codei'num*, *Papaveri'num*, (from *κωδία*, 'a poppy head.') An alkaloid discovered by Robiquet in opium, in 1832. It is soluble in water, alcohol and ether, and unites readily with acids. As a hypnotic, Magendie thinks one grain of codeia equal to half a grain of morphia. The muriate of codeia appears to be stronger than the pure codeia.

CODESELLA, *Anthrax*.

CO'DEX, gen. *Co'dicis*, 'a book.' A collection of laws. (F.) *Code*. By extension, a collection of approved medical formulæ, with the processes necessary for forming the compounds referred to in it. The Parisian Pharmacopœia is styled *Co-dex medicamentarius*.

CODEx MEDICAMENTARIUS, Pharmacopœia, see *Codex*.

CODIA, (*κωδία*, or *κωδεία*.) *Papaver* (capsule.)

CODOCELE, (*κωδία*, 'a head,' and *κηλη*, 'a tumour.') Bubo.

CODLING, (dim. of *Cod*), Testicle.

CODOSCELLA, Bubo.

CÆCAL, *Cæcal*.

CÆCITAS, *Cæcitas*.

CÆCUM, *Cæcum*.

COEFFE, *Caul*.

CÆ'LIA, (*καίλια*, *κοίλη*, 'a hollow place.') This word generally meant a cavity in the body:—the abdomen, in particular. It has also been used for the alimentary canal, &c.:—*ανω κοίλια*, 'the stomach,' *κατω κοίλια*, 'the abdomen.' Also, an alvine evacuation; excrement.

CÆ'LIAC, *Cæli'acus*, *Gastrocæli'acus*, *Gastrocæ'licus*, (F.) *Cæ'liaque* ou *Céliaque*, (from *cælia*.) Relating to the cavity of the abdomen.

CÆLIAC AR'TERY, *Ar'teria Cæli'aca*, *Co'liac Ar'tis*, *Artère opistogastrique*, (Ch.) *A. Cæliaque*, *Tronc cæliaque*, is situate above the pancreas, and behind the upper part of the stomach. It arises from the anterior part of the abdominal aorta, where it passes between the pillars of the diaphragm, and soon divides into the *coronaria ventriculi*, *hepatic*, and *splenic* arteries:—the division being called *Tri'pus cæli'acus* seu *Halle'ri*, (F.) *Trépied cæliaque* ou *de la Cæliaque*.

CÆLIAC AXIS, C. Artery.

CÆLIAC FLUX, *Cæliac Pas'sion*, *Cæliac'a chylo'sa*, *Diarrhœ'a chylo'sa* seu *chymo'sa*, *Flux'us chylo'sus* seu *cæli'acus*, *Pas'sio cæli'aca* seu *Ventriculo'sa*, *Chymochœ'zia*, *Flu'or al'bus intestino'rum*, *Chylorrhœ'a*, *Chymorrhœ'a*, *Cæli'aca lac'tea*, *Mor'bus cæli'acus*, *Chylodiarrhœ'a*, *Galactodiarrhœ'a*, *Sêdes lactescen'tes*, *Gastrorrhœ'a*, (F.) *Flux cæli-aque*. A species of diarrhœa, in which the food is discharged by the bowels in an undigested condition. By some, defined to be diarrhœa attended with discharges of chyle or chyme. It is, in general, symptomatic of tubercular disease of the mesenteric glands. See Lientery.

CÆLIAC PASSION, *Cæliac flux*.

CÆLIAC PLEX'US, *So'lar Plex'us*, *Plex'us mesenterii pro'prius* et *ma'ximus*, *P. ganglior'mis semi-lunaris*, (F.) *Plexus médian ou opistogastrique*, (Ch.), *Pl. Cæliac* ou *solaire*, *Ganglion de VIEUSSENS*, is formed of numerous nervous filaments, which proceed from the semilunar ganglia of the great sympathetic. It is strengthened by several branches of the right and left pneumogastric nerves; is seated around the trunk of the cæliac artery, behind the stomach, and furnishes the secondary plexuses—the *diaphragmatic*, *coronary of the stomach*, *splenic* and *hepatic*, which accompany the arteries of the same name.

CÆLI'ACA, (from *cælia*.) Diseases of the digestive organs; the 1st class in Good's *Nosology*. It comprises two orders, *Enterica* and *Splanchnica*. Also, medicines that act on the digestive organs.

CÆLIACA CHYLOSA, *Cæliac flux*—c. *Lactea*, *Cæliac flux*—c. *Renalis*, *Chyluria*—c. *Urinalis*, *Chyluria*.

CÆLIADÉLPH'US, (*cælio*, and *ἀδελφος*, 'brother.') *Anocæliadelph'us*. A double monster, in which the two bodies are united at the abdomen.

CÆLLE'MIA, *Hyperæ'mia abdomin'is*, *Conges'tio abdomina'lis*, (*cælio*, and *αἷμα*, 'blood.') Hyperæmia or congestion of the blood-vessels of the abdomen.

CÆLI'AGRA, (*cælio*, and *αγρα*, 'a seizure.') *Gastritis* seu *Enteritis* seu *Co'liaca* seu *Diarrhœ'a arthrit'ica*. Gout of the abdomen.

CÆLIALGIA, (*cælio*, and *αλγος*, 'pain.') *Tormina*.

CÆLIAQUE, **TRÉPIED DE LA**, *Cæliac artery*.

CÆLIO, (from *κοίλια*, 'the abdomen.') In composition, abdomen.

CÆLIOCELE, (*cælio*, and *κηλη*, 'rupture.') See *Hernia*, hypogastric.

CÆLIOCHYSIS, (*cælio*, and *χυσ*, 'to pour.') *Ascites*.

CÆLIOCYESIS, (*cælio*, and *κυειν*, 'to conceive.') Pregnancy, abdominal.

CÆLIODYNIA, (*cælio*, and *δωνη*, 'pain.') *Colic*.

CÆLIOLYSIS, (*cælio* and *λυσις*, 'solution.') Diarrhoea.

CÆLIOMYAL'GIA, *Cæliomyodyn'ia*, (*cælio*, *μυς*, 'a muscle,' and *αλγος*, 'pain.') Pain in the muscles of the abdomen.

CÆLION'CUS, *Cæliophy'ma*, (*cælio*, and *ογκος*, 'a tumour.') A tumour of the abdomen.

CÆLIOPHYMA, (*cælio*, and *φυμα*, 'a tumour.') Cælioncus.

CÆLIOPHY'MATA, pl. of *Cæliophyma*, *Tubercula peritonæ'i*. Tubercles of the peritoneum.

CÆLIOPSOPHIA, (*cælio*, and *ψοφος*, 'a noise.') Borborygmus.

CÆLIOPYO'SIS, (*cælio*, and *πυσις*, 'suppuration.') Suppuration of the abdomen or its parietes.

CÆLIORRHÆA, (*cælio*, and *ῥευ*, 'to flow.') Diarrhoea.

CÆLIORRHEUM'A, gen. *Cæliorrhæum'atis*, *Rheumatismus abdom'inis*, (*cælio*, and *ῥευμα*, 'defluxion, rheumatism.') Rheumatism of the muscles of the abdomen.

CÆLIOSPAS'MUS, (*cælio*, and *σπασμος*, 'spasm.') Spasm or cramp of the abdomen.

CÆLO, (from *κοιλος*, 'hollow.') In composition, hollow, *cavus*.

CÆLO'MA, gen. *Cælo'matis*, (*κοιλος*, 'hollow.') A round ulcer of the cornea, broader and deeper than that described under the name *Bothrion*.

CÆLOMELE, (*κοιλος*, 'hollow,' and *μηλη*, 'a probe.') Grooved probe.

CÆLON, (from *κοιλος*, 'hollow.') Cavity.

CÆLOPHLEBITIS, (*cælio*, *φλεβ*, gen. *φλεβος*, 'a vein,' and *ιτις*.) Inflammation of the vena cava.

CÆLOPHTHALMUS, (*cælio*, and *οφθαλμος*, 'eye.') One who is hollow-eyed.

CÆLOSTOM'IA, (*cælio*, and *στομα*, 'mouth.') A defect in speaking, when the voice seems as if it came from a cavern;—that is, obscure, and as if at a distance.

CÆLOS'TOMUS, same etymon. One who has a hollow voice.

CÆLOTES, (from *κοιλος*, 'hollow.') Cavity.

CÆNADELPH'US, (*κοινος*, 'common,' and *ἀδελφος*, 'brother.') A double monster, the two portions having, in common, one or more organs indispensable to life.

CÆNÆSTHESIS, (*κοινος*, 'common,' and *αισθησις*, 'feeling.') *Cœnæsthesi*, (F.) *Cœnæsthési*. 'Common feeling.' Some German writers mean, by this, a sixth sense. It is the feeling of self-existence or individuality, and is manifested by the sense of buoyancy or depression, which we experience without any known cause;—by involuntary shuddering, feeling of chill or glow, &c.

CÆNO, from (*κοινος*, 'common.') In composition, common.

CÆNOLOGIA, (*κοινος*, 'common,' and *λογος*, 'a discourse.') Consultation.

CÆNOTES, (*κοινωνη*, from *κοινος*, 'common.') The physicians of the methodic sect asserted, that all diseases arise from *relaxation*, *stricture*, or a mixture of both. These were called *Cœnotes*: or what diseases have in common.

CÆTUS, Coition.

CÆUR, (*κρη*, *cor*.) Heart.

COFFEA ARAB'ICA, (*Caffè'a*, Ph. U. S. 1863.)

C. vulg'aris, *Jas'minum Arab'icum*, (F.) *Caffier*, *Caffèier*, and *Caffeyer*. Order. Rubiaceæ. *Ser. Syst.* Pentandria Monogynia. The plant which affords coffee, *Chaava*, *Bon*, *Buna*, (F.) *Café*. Originally from Yemen.

The infusion of coffee is an agreeable and wholesome article of diet. It is, to a certain extent, a tonic, and is employed as such in convalescence, especially from fevers, &c. In cases of poisoning by opium, and in old asthma, its use has been extolled. For this purpose the *Moka* is the best. It ought to be newly torrefied, but not too much

burnt; should be strong, and drunk soon after grinding.

A highly nitrogenized alkaloid was first found in coffee, and called *Caff'ein* or *Caff'èin*. It was afterwards discovered in the leaves of the tea plant, and called *Thein*, and in the Guarana of Brazil, and called *Guaranin*. In doses of from 2 to 10 grains, it induces violent nervous and vascular excitement. Caffein—as well as its various salts—*the citrate, lactate, malate, and muriate*, has been given as an antiperiodic in grain doses.

Facit'ious Coffees have been, from time to time, recommended, but they are infinitely inferior to the genuine. Various substitutes have been proposed; *wheat, barley, hollyberries, acorns, sunflower seeds, beech-mast, peas, beans, succory-root, seeds of gooseberries and currants* left in making wine, and washed,—*sliced turnips, &c.* These have been roasted, with the addition of a little butter or oil; but they have not the aroma of coffee. The best substitute is said to be the seeds of the *Yellow water flag, Gladi'olus luteus* or *Iris pseudac'orus*.

Hunt's *Econom'ical Break'fast Powder* consisted of rye, roasted with a little butter.

COFFEE, see *Coffea Arabica*.

COFFEE-BEAN TREE, KENTUCKY, *Gymnocladus Canadensis*.

COFFEE-TREE, *Gymnocladus Canadensis*.

COFFEE, WILD, *Triostemum*.

COGNASSIER, *Pyrus cydonia*.

COGNITIO PHYSIOLOGICA, (*cognoscere*, *cognitum*, 'to know,') 'physiological knowledge.' Physiology.

COHABITATIO, Cohabitation, Coition.

COHABITATION, *Cohabitatio*, (*co*, and *habitare*, 'to dwell,') The act of dwelling together. In legal medicine, it means the consummation of marriage.—*Coipulation*.

COHESION, *Cohæsi*, (*co*, and *hæere*, *hæsum*, 'to stick,') *Vis cohæsi'nis* seu *adhæsi'nis* seu *attracti'nis*, *Force of cohæsi*, *Attraction of cohæsi* or of *aggrega'tion*, is that force in the particles of matter, whereby they are connected in such a way as to resist any attempt towards their removal or separation. This force has to be attended to, in the management of disease. Emollients, rubbed into a part, act by diminishing the cohesion.

COHIBENS, 'holding together,' (from *co*, and *habere*, 'to have,') Epichetivus.

COHOBAT'ION, *Cohobatio*, *Cohob'ium*, *Cô'hob*, *Co'hoph*, (*cohob*, a Paracelsian term for repetition.) Distillation of a liquid, already distilled, from the same substances. When this is repeated three or four times, it is called *Recohobation*.

COHOL, Syn. of Alcohol. Also, a dry collyrium.

COHOSH, *Actæa racemosa*, *Caulophyllum thalictroïdes*—c. Black, *Actæa racemosa*—c. Blue, *Caulophyllum thalictroïdes*—c. Blueberry, *Caulophyllum thalictroïdes*—c. Red, *Actæa rubra*—c. White, *Actæa alba*.

COHUSH, *Caulophyllum thalictroïdes*.

COIFFE, *Caul*.

COIGN, see *Pyrus cydonia*.

COIGNASSIER, *Pyrus cydonia*.

COINCIDEN'TIA, (*co*, and *incido*, (*in*, and *cado*,) 'to fall or happen,') Some authors have translated, by this term, the word *parenptosis*, used by Galen to designate the occlusion of the foramen opticum by a humour, proceeding from the base of the brain, and occasioning blindness! COIN'DICANT, *Coin'dicans*, (*co*, and *indico*, 'I indicate,')

COIN'DICANT SIGNS are those which furnish the same indications; or which confirm the indication afforded by another sign—*συνειδικνυμενα*.

COIRA, Catechu.

COIRAS. (χοίρας, from χοίρος, 'a hog,') Scrotula.

COIT, Coition.

COIT'ION, (Coit'io, Coitus, Cætus, coïzo, (co, and eo, 'to go,') 'to go together,') Copula'tio, Cop'ula Carna'tio, Gemina'tio, Aphrodis'ia, Aphrodisiaem'us, Epip'locë, Acces'sus, Basia'tio, Amplex-a'tio, Amplex'us, Conven'tus, Complex'io, Conju-gium, Agglutina'tio, Ocheu'ma, Futu'tio, Patra'tio, Lagne'a, Lagneu'ma, Lagnei'a, Mix'is, Permis'sio, Permis'tio, Syndyus'mus, Syndu'cia, Concu'bitus, Congres'sus, Cohabita'tio, Ve'nus, Res Ven'e'ren, Concu'bitus ven'e'rens, Præ'tium, Duellum ven'e'reum, Noctur'na bella, Concil'ia corpora'tia, Homi'lia, Co'iture, Copula'tion, Ven'ery, Sex'ual in-tercourse, (F.) Coit, Approche, Accouplement. The carnal union of the sexes.

COITURE, Coitus.

COITUS, gen. Coitis, Coition—c. Difficilis, Dys-synodus—c. Sodomiticus, Buggery.

COL, as a prefix, the same as co, cum, 'with.'

COL, Collum—c. de la Matrice, Collum uteri—c. Uterin, Collum uteri.

COLATIO, see Colatura.

COLATO'R IUM, Hylister, (colare, 'to strain,') A strainer of any kind. (F.) Couloir. A term by which the ancient physicians described every canal or conduit through which the excrementitious humours of the body are evacuated. Ulcers, fistulæ, setons, caustics, &c., have been called artificial or accidental Colatoria, because they were considered to be passages by which the animal economy is freed from some morbid matter.

COLATUM, see Colatura.

COLATU'RA (colare, 'to strain,') Colat'um, (F.) Colature. A filtered or strained liquor. It likewise means straining a liquid—Colat'io. Diy-liaia, Diyli'smus, Hy'lisia, Hylis'mus.

COLCHESTER, MINERAL WATERS OF. Aqua Colcestren'sis. The waters of Colchester are of the bitter purging kind, similar to those of Epsom.

COLCHICIA, see Colechicum autumnale.

COLCHICUM, (κολχικον, from Κολχis, Colchis, in Asia, where it is said to have been common.) Colechicum autumnale.

COLCHICUM AUTUMNA'LE, Col'chicium, Co'm, Theri'aca articulo'rum, Mead'ow Saff'ron, (F.) Colechique, Tue-chien, Mort aux chiens, Safran des prés ou bâtar, Narcisse d'autonne, Viellotte. Nat. Ord. Colchicaceæ. Class, Hexandria. Order, Monogynia. The bulb or root or cormus, Col'chici rad'ix, (Ph. U. S.) Colchici cor'mus, Colchic'um corm, (Ph. B. 1864.) and the seeds, Col'chici semen, are the parts used in medicine. The taste is acrid, excoxiating the mouth; but the acrimony is lost by drying. It is narcotic, diuretic, and cathartic; and has been given in dropsy, gout, and rheumatism. Dose, gr. j to vj of the fresh bulb. It is supposed to be the active ingredient of the Eau médicinale d'Husson. The active principle is called Colchie'ia. Colechicum, in an over-dose, is an aro-narcotic poison.

Dr. Wilson's Tincture for the Gout is said to be merely a tincture of the seeds, or of Colchicia.

COLCHICUM ZEYLANICUM, Kæmpferia rotunda.

COLCHIQUE, Colechicum autumnale.

COLCOT'HAR, C. Vitri'oli, Henri'cus ru'bens, Chalc'i'tia, Cro'cus, Ox'idum Fer'ri ru'brum, Tri-toxy'dum Fer'ri, Sul'phas Fer'ri calcina'tum, Fer'rum vitriola'tum us'tum, Ter'ra vitri'oli dul'cis, Cro'cus mart'is vitriola'tus seu adstrin'gens, Brown red, Rouge, (F.) Rouge d'Angleterre ou de Prusse, Safran de Mars astringent. The red oxide of iron, obtained by calcining sulphate of iron to redness, with or without the contact of air. It possesses the general properties of the preparations of iron, and has been applied to staunch blood, &c.

COLD, (Sax. colb,) Fri'gus, Pay'chus, (F.) Froid. The sensation produced by the abstraction of

caloric from our organs,—Chelm'a, Chi'mon. See Heat.

Three degrees of cold are generally distinguished in disease. 1. The simple feeling of cold (Al'gor), 2. Chil'liness (Horror), and 3. Shiver'ing (Ri'gor). Cold is employed in medicine, chiefly, as a refrigerant and sedative.

COLD is, also, used for a morbid affection induced by cold. 'To take cold,' (F.) s'enrhumer, is to be affected by a disease presumed to be caused by cold. See Catarrh and Coryza.

COLDCHILL, Intermittent fever.

COLD IN THE EYE, Ophthalmia, catarrhal — c. in the Head, Coryza.

COLD SLAW OR COLE SLAW, probably a corruption from (G.) Kohlsalat, (D.) Kool salade, (from (G.) Kohl, 'Cole, cabbage,') Cabbage salad. A salad formed of the hard head of white cabbage cut into small strips, and dressed like ordinary salad.

COLE SLAW, Cold slaw.

COLEITIS, (κολεις, 'a vagina or sheath,' and itis,) Colpositis.

COLEOCELE, (κολεις, 'a vagina or sheath,' and κηλη, 'rupture,') Hernia, vaginal.

COLECEDEMA, (κολεις, and οιδημα, 'swelling,') Elytroedema.

COLEOPTOSIS, (κολεις, and πωσις, 'a fall,') Prolapsus vaginae.

COLEORRHÆXIS, (κολεις, and ρηξις, 'rupture,') Laceration or rupture of the vagina.

COLEOSTITIS, (κολεις, and itis,) Leucorrhœa.

COLEOSTEGNO'SIS, Colpostegno'sis, Colpos-tenochor'ia, Colposynize'sis, (κολεις, and στεγνωσις, 'constriction,') Narrowness of the vagina, natural or acquired.

COLERE, (from κολη, 'bile' [?]), Rage.

COLES, Penis—c. Feminarum, Clitoris.

COLEWORT, Col'lard, (cole and wort, A species of cole or cabbage,

COLEUS, Scrotum, Testicle, Vagina.

COLIBERT, Cagot.

COL'IT, Col'licus, (κολικος, from κολον, the colon,') (F.) Colique. Relating to the colon or to colic.

COLIC ARTERIES, Arte'rix col'licæ, (F.) Artères Coliques, are six in number, three of which, given off by the superior mesenteric, are called Colicæ dextræ; and three, proceeding from the inferior mesenteric, Colicæ sinistræ. All proceed towards the colon, passing between the two laminae of the peritoneum, which form the mesocolon, where they ramify and anastomose with each other. The first, Col'lica dex'tra, Ra'mus col'licus dex'ter, is called C. dex'tra super'ior, (Mésocolique,—Ch.) The second, C. dex'tra me'dia, Col'lica me'dia, Ra-mus col'licus me'dius, Arte'ria me'dia anastomot'ica, (C. droite, Ch.), and the third, C. dex'tra inf'erior or Ileocol'lica, (A. cécale,—Ch.) Of the three Col'licæ sinistræ, the first or super'ior is called, by Chaussier, Col'lica magna sinistræ; the second or me'dia is not distinguished from the preceding, as they often arise from a common trunk; and the third is called by Chaussier Col'lica par'va sinistræ. To these arteries as many veins correspond, which open into the great and little mesenteric.

COLIC, (κολικη νόσος,) Col'ica, C. Par'va, Col'licæ, Colicodyn'ia, Caliodyn'ia, Dolo'res intestino'rum, Dol'or col'licus, Dysenteronerr'ia, Enter'al'gia, Gripes, Mulligrubs, Bellyache, (Sc.) Bellythro', Weam-ill, (Prov.) Bellyharm, Bellycark, Gull-ion. In its etymological acceptation, Colic signifies an affection or pain in the colon. But it is employed in a more extensive signification. It includes every acute pain of the abdomen, aggravated at intervals. The word has often, however, epithets affixed to it, which render it more or less precise. See Termina.

COLIC, BILIOUS, Colica biliosa—c. Crapulent, Colica crapulosa—c. Devonshire, Colica metallica—c. Horn, Priapismus—c. Lead, Colica metallica.

COLIC LOBE OF THE LIV'ER is the great lobe of that organ.

COLIC MADRID, *Colica Madridensis*—c. Menstrual, *Colica menstrualis*—c. Metallic, *Colica metallica*—c. Nephretic, *Colica nephretica*—c. Normandy, *Colica metallica*—c. Painters', *Colica metallica*—c. Plumbers', *Colica metallica*—c. of Poitou, *Colica metallica*—c. of Prostitutes, *Colica scortorum*—c. Renal, *Colica nephretica*—c. Root, *Aletris*—c. Saturnine, *Colica metallica*—c. of Surinam, *Colica metallica*—c. Uterine, *Colica uterina*—c. Vegetable, *Colica metallica*—c. Weed, *Dicentra Canadensis*—c. Worm, *Colica verminosa*.

COLICA, *Colic*—c. Accidental, *Colica crapulosa*—c. Acute, *Enteritis*—c. Arthritic, *Coeliagra*.

COLICA BILIOUSA, *Bilious Colic*, (F.) *Colique Bileuse*. Colic, occasioned by an accumulation of bile in the intestines or in its own passages. The treatment required resembles that proper for *Colica convulsiva*.

COLICA CALLO'SA. Colic attended with sense of stricture in some parts of the intestinal canal; often with flatulency and pain; the flatus gradually passing off by the stricture; the bowels tardy; and at length discharging small liquid stools.

COLICA CONVULSIVA, *C. Spasmodica* seu *pituitosa* seu *nervosa* seu *idiopathica*, *Enterospasmus*. Colic, not the symptom of any other affection. It is characterized by griping pain in the bowels, chiefly about the navel, with vomiting and costiveness—the pain increasing at intervals. The indications are to clear the intestines, and allay spasm. Calomel and opium—castor oil and opium—emollient and cathartic enemata, with fomentations, wet or dry, to the abdomen, usually succeed.

COLICA CRAPULOSA, *C. accidentalis* seu *heluonum*, *Cropulent colic*, *Surfeit Colic* from overeating, (F.) *Colique d'Indigestion*. A colic, arising from eating indigestible aliments, or digestible aliments in too great abundance. The remedy is obvious.

COLICA DAMNIOURUM, *Colic, metallic*—c. Febrile, *Colica inflammatoria*—c. Figulorum, *Colica metallica*.

COLICA FLATULENTA, *Inflatio, Gastrodynia flatulenta, Physospermus, Pneumosis enterica*, (F.) *Colique flatulente ou flatueuse ou ventreuse*. Colic, arising from an accumulation of air in the intestines. It is very common in infants, and may be relieved by aromatics, especially when combined with antacids,—for example, oil of aniseed with magnesia.

COLICA HÆMORRHOIDALIS, *Hemorrhoid'al Colic*, (F.) *Colique hémorrhoidale*. A kind of colic, supposed to precede hemorrhoids or to supervene on their suppression.

COLICA HELLUONUM, *Colica crapulosa*.

COLICA HEPATICA, *Hepatalgia, Hepatalgia Calculosa, Hepat'ic colic*, (F.) *Colique hépatique*. Pain in the region of the liver, chiefly about the gall-bladder, occasioned by the passing of a biliary calculus through the cystic and choledoch ducts.

COLICA IDIOPATHICA, *Colica convulsiva*—c. Ileus, Ileus.

COLICA INFLAMMATO'RIA, *C. Phlogistica* seu *plethorica* seu *febricosa* seu *pulsatilis*, *Inflammatory colic*. The same as enteritis.

COLICA LAPPONICA, see *Seta equina*—c. Lochialis, Dyslochia.

COLICA MADRIDEN'SIS, *Col'ic of Madrid', Madrid Colic*. A species of colic, endemic in several provinces of Spain, whose nature is not clear. Its symptoms resemble those occasioned by lead. See *Colica metallica*.

COLICA MENSTRU'ALIS, *Menstrual Colic*, (F.) *Colique menstruelle*. Colic, which precedes or accompanies the menstrual evacuation, or which is owing to the suppression of that evacuation.

COLICA METAL'LICA, *Metal'lic Col'ic, Painters' Col'ic, Col'ic of Poitou, Col'ic of Surinam', Bellow, Devonshire Col'ic, Dry Bell'gache, Dry Gripes, Saturnine Col'ic, Lead Col'ic, Plumbers' Col'ic, Rachial'gia, R. Pictonum* seu *Pictavien'sium, Morbus Metal'licus, Colicople'gia, Colique Rachial'gia* seu *Damnion'rum* seu *Plumbario'rum* seu *Pictonum* seu *Satur'na* seu *Figulo'rum* seu *nervosa* seu *Pictor'um* seu *sic'ca, Paralysis rachial'gia, Pal'mus Plumbarius, Molybdocolica*, (F.) *Dysenternervie Saturnine, Colique de Poitou ou végétale ou des peintres ou de plomb ou métallique ou Saturnine ou des barbouilleurs ou sèche*. Under this head is generally described the colic produced by lead, as well as the other colics mentioned in the synonymy; and they certainly resemble each other greatly, although some of them are more like bilious colic. There is not much to distinguish this variety of colic from others. The same violent pain about the navel is present, with retraction of the abdomen towards the spine. It is apt also to occasion palsy. The only difference of treatment is in the necessity for employing more opium along with the purgative. The paralytic sequelæ must be met by change of air, rubbing the spine, exercise, &c. Treating the disease upon general principles is infinitely more philosophical, and more successful than the former empirical management at La Charité, Paris, which it is unnecessary to detail.

The *Colic of Madrid, of Poitou, Devonshire, Normandy, &c.*, has been called *vegetable colic*, (F.) *Colique végétale*, because ascribed to the use of new wine or cider, unripe fruit, &c., although really, perhaps, owing to the adulteration of wines and cider by litharge. All the symptoms, indeed, are those of *Colica metallica*.

COLICA NEPHRETICA, *Nephrocolica, Col'ica renal'is, Nephret'ic or Ren'al Col'ic*, (F.) *Colique Néphrétique*. Acute pains, which accompany nephritis, and especially calculous nephritis, or the passage of a calculus into the ureter.

COLICA NERVOSA, *Colica convulsiva, Colic, metallic*—c. Passio, *Colica*—c. Phlogistica, *Colica inflammatoria*—c. Pictonum, *Colica metallica*—c. Pictorum, *Colica metallica*—c. Pituitosa, *Colica convulsiva*—c. Plethorica, *Colica inflammatoria*—c. Plumbariorum, *Colica metallica*—c. Pulsatilis, *Colic, inflammatory*—c. Rachialgia, *Colica metallica*—c. Renalis, *Colica nephretica*—c. Saturnina, *Colica metallica*.

COLICA SCORTORUM, *Col'ic of Prostitutes*. A form of colic said, by Dr. Martin Hassing, of Copenhagen, to have been frequently observed by him amongst that unfortunate class of beings. It may well be doubted whether any special affection of the kind appertains to them.

COLICA SICCA, *C. metallica*—c. Spasmodica, *Colica convulsiva, Ileus*.

COLICA STERCO'REA, *Colica Stip'ita, Stercoraceous Col'ic*, (F.) *Colique stercorale*. A species of colic, ascribed to the retention of fecal matters in the intestines. The retention is itself, however, an effect, that may be caused in the same manner as the colic pains themselves.

COLICA STIPATA, *Colica stercorea*.

COLICA UTERINA, *Hystero'colica, U'terine Colic*, (F.) *Colique utérine*. Pain seated in the uterus, sometimes called *Hysteral'gia*.

COLICA VENTRICULI, *Cardialgia*.

COLICA VERMINOSA, *Helminthocolica, Worm Col'ic*, (F.) *Colique vermineuse*. Abdominal pain, occasioned by the presence of worms in the intestines.

COLICODYNIA, (*colic*, and *δύνη*, 'pain.') Colica.

COLICOPLEGIA, (*colic*, and *πληγη*, 'a stroke.') Colic, metaplic.

COLIMAÇON, *Helix pomatia, Limax*.

COLIQUE, Colic—c. *des Barbonnières*, Colica metallica—c. *Hépatique*, Colica hepatica—c. *d'Indigestion*, Colica erapulosa—c. *Métallique*, Colica metallica—c. *de Misère*, Pleus—c. *de Miséricorde*, Pleus—c. *des Peintres*, Colica metallica—c. *de Plomb*, Colica metallica—c. *de Poiton*, Colica metallica—c. *Saturinée*, Colica metallica—c. *Sèche*, Colica metallica—c. *Végétale*, Colica metallica—c. *Ventuse*, Colica flatulenta.

COLIQUES, Pains, [after.]

COLIS, Penis—c. *Feminarum*, see Clitoris.

COLITIS, (colon, and *itis*.) *Coloni'tis*, *Enteritis collica*, (F.) *Inflammation du colon*. Inflammation of the peritoneal or mucous membrane of the colon. The former is termed *Serocolitis*, and *Eecocolitis*; the latter *Endocolitis* and *Dysentery*.

COLIX, (κολιξ.) Trochiscus.

COLLA PISCUM, ('glue of fish.') Ichthyocollo.

COLLAPSE, Collapsus—c. of the Lung, see Atelectasis—c. *Pulmonary*, see Atelectasis.

COLLAP'SUS, (collabor, *collapsus*, *col*, and *labor*, *lapis*, 'to fall down,') *Collapsé*, *Coincidence'tia*; (F.) *Affaissement*. A complete prostration of strength, either at the commencement or in the progress of a disease.

COLLAR-BONE, Clavicle.

COLLARD, Colewort, *Dracontium fetidum*—c. Cow, *Dracontium fetidum*—c. Polecat, *Dracontium fetidum*.

COLLARTUM SALINUM, (from *collum*, 'the neck,') Halidraemum.

COLLATERAL, *Collatera'lis*, (F.) *Collatéral*, (*col*, and *latus*, gen. *lateris*, 'side,') *Collat'el*, which accompanies or proceeds by the side of another.

COLLATERAL ARTERIES OF THE ARM, *Arte'rie Collatera'les Bra'chii*, (F.) *Artères collatérales du bras*. They are given off by the brachial, and are distinguished into, 1. The *collateral—superior* or *external* *Grandes musculaires du bras*—Ch.) which arise from the inner part of the brachial, and extend as far as the inferior and external part of the arm. 2. The *collateral—inferior* or *internal* (*Collatérales du Coudé*—Ch.) which arise from the brachial, near the elbow-joint, and descend towards the upper part of the fore-arm.

The vessels which pass to the fingers and toes are also called *collateral*.

Speaking generally, *collateral branches* are those which follow nearly the same course as the vessel whence they emanate.

COLLATERALIS, *Ischio-cavernosus*.

COLLE-CHAIR, ('glue-flesh,') *Sarcocolla*.

COLLE DE POISSON, (glue of fish, from *κόλλα*, 'glue,') *Ichthyocolla*.

COLLECTION, *Collectio*, (*colligere*, *collectum*, *col*, and *lego*, 'to gather,') This word is often united to the epithet *purulent*, *serous*, &c., to express a *gathering* of pus, serum, &c.

COLLESIS, (καλλοσις, from *κόλλα*, 'glue,') *Agglutination*.

COLLET DES DENTS, (from *collum*, 'neck,') *Collum dentium*, see Tooth.

COLLETICUS, (καλλητικός.) *Agglutinant*.

COLLEY'S DEPILATORY, see Depilatory.

COLLICULE, (Same etymon as *Collection*; or from *col* and *liquere*, 'to be liquid,') 'Drains to collect and convey away water.' See *Gouttière*. Union of the ducts passing from the puncta lacrymalia, *Collic'ix puncto-rum lacrymalium*.

COLLICULI (pl. of *Colliculus*) **BULBI**, see *Corpus spongiosum urethrae*—c. *Nervi Ethmoidalis*, *Corpora striata*—c. *Nervorum optico-rum*, *Thalami nervorum optico-rum*—c. *Vaginae*, *Nymphæ*.

COLLICULUS, (dim. of *collis*, 'a hill,') See *Optic nerve*.

COLLICULUS CAVEÆ POSTERIORIS VENTRICULO-

RUM LATERALIVM, *Hippocampus minor*—c. *Seminalis*, *Gallinaginis caput*.

COLLIER, (F.) *A collar or necklace*, from *collum*, 'the neck,') A name given to certain eruptions which surround the neck like a collar.

COLLIER ANODIN, *Necklace*, *anodyne*.

COLLIGAMEN, (*col*, and *ligare*, 'to bind,') *Ligament*.

COLLIGATIO, *Syndesmosis*.

COLLINSONIA, *C. Canadensis* seu *decussata*, *Horse-weed*, *Horse-balm*, *Rich-weed*, *Rich-leaf*, *Hen'all*, *Stone-root*, *Knot-root*, *Knot-weed*, *Knob-weed*. *Order*, *Labiata*, [after Peter Collinson.] This indigenous plant is possessed of diuretic properties, which seem to reside in a volatile oil. Infusion is the best form of administration. The leaves in domestic practice are applied to wounds and bruises.

COLLINSONIA DECUSSATA, *C. Canadensis*.

COLLIQUAMENTUM, (*colliguo*, (*col*, and *liquo*,) 'I melt,') The first rudiments of an embryo.

COLLIQUATION, *Colliqua'tio*, *Eliqua'tio*, *Syntes'is*, *Ectes'is*, *Solutio*, *Dissolutio*. 'The act of melting.' (F.) *Fonte*. The ancients meant, by this term, the progressive diminution of the solid parts, with copious excretion of liquids by one or more passages. They thought, that all the solids melted; and that the liquids, and particularly the blood, lost a portion of their density.

COLLIQUATIVE, *Colliquati'vus*, *Colligues-cens*, (F.) *Colligatif*. An epithet given to various discharges, which produce rapid exhaustion. Hence we say, *Colligative sweats*, *Colligative diarrhoea*, &c.

COLLIQUESCENS, (*col*, and *liquescere*, 'to grow liquid,') *Colligative*.

COLLISIO, (*collidere*, *collisum*, (*col*, and *la-dere*,) 'to dash together,') *Contusion*.

COLLIX, (κολιξ.) Trochiscus.

COLLOCYSTIS, (κόλλα, 'glue,' and *κυστις*, 'a bladder,') Capsule, gelatinous.

COLLODES, (κόλλα, 'glue,' and *odes*,) *Glutinous*.

COLLODESMUS, (κόλλα, 'I cement,' and *δεσμος*, 'a bandage,') *Apparatus*, *immoveable*, and *Bandage*, *starch*.

COLLODION, *Collo'dium*, *Liquor sulphurico-æthe'reus constrin'gens*, *Liquid cuticle*, *Ethe'real Solution of Gun-cotton*, *Maynard's Adhesive Liquid*, (from *κόλλα*, 'glue,') A solution obtained by dissolving *Gun-cotton*, *Gossypium fulminans*, *Pyrox'lin*, (F.) *Fulmicoton*, *Coton-poudre*, *Poudre-coton*, *Coton fulminant*, *Coton azotique*, in a mixture of rectified ether and alcohol, in the proportion of about 16 parts of the former to 1 of the latter. In the British Pharmacopoeia (1864) it is prepared of *Pyrox'lin*, $\frac{3}{3}$; *Ether*, $f\frac{3}{3}$ xxxvj; *Rectified spirit*, $f\frac{3}{3}$ xij. When applied to a part, the ether evaporates, and the solid adhesive material is left, which contracts. Hence it is used in cases of wounds, to keep their edges together. It forms, also, a coating, and has been applied in abrasions, and in cases of burns. In various chronic cutaneous diseases, it has been applied with advantage; and has been employed to give a coating to pills, which it deprives of their taste, without interfering with their action.

A process for obtaining collodion is in the Ph. U. S.

COLLODION CANTHARIDAL, *Collo'dium vesica'ns* seu *cantharida'le* seu *cantharida'tum*, *Liquor vesica'ns*. Prepared by exhausting, by displacement, a pound of coarsely-powdered *cantharides* with a pound of sulphuric ether, and three ounces of acetic ether. In two ounces of this saturated solution of cantharides, twenty-five grains of gun-cotton are dissolved. A process for this preparation is in the Ph. U. S 1863.

By painting the surface with a little of this solution, vesication is produced, as with the plaster of eaptharides.

COLLODION, RICINATED, see Collodion, terebinthinated.

COLLODION, TEREBINTHINATED or RICINATED, *Collodium terebinthinatum* seu *ricinatum*, (F.) *Collodion térébinthiné ou riciné*. A preparation of *Collodion*, $\frac{3}{4}$; *Ol. ricini* vel *Terebinth. venet.*, gr. x. Used by the Germans and French as a covering in erysipelas, burns, bedsores, and chronic cutaneous diseases.

COLLODIDUM, Collodion—c. *Cantharidale*, *Collodion cantharidal*—c. *Ricinatum*, see *Collodion*, terebinthinated—c. *Terebinthinatum*, *Collodion*, terebinthinated—c. *Vesicans*, *Collodion*, cantharidal.

COLLOID, *Colloïdes*, (κωλλα, 'glue,' and εἶδος, 'resemblance,') *Can'cer alveola'ris*, *Carcino'ma alveola'ris*, (F.) *Cancer alvéolaire ou colloïde ou gélatiniforme*, *Cancer alvéolaire ou aréolaire ou pulsaté ou gommeux*, *Gelat'inous cancer*. An epithet applied to a product of morbid secretion, resembling glue, or particles of jelly inlaid in a regular alveolar bed. The three species of cancer or carcinoma are,—*Encephaloid*, *Scirrhus*, and *Colloid*.

COLLO'MA, (κωλλα, 'glue,' and ομα.) A name proposed by Dr. Walshe for the gelatiniform matter, which is of common occurrence in cysts. It is transparent, amorphous, and devoid of vessels and nerves.

COLLONE'MA, gen. *Collone'matis*, (κωλλα, 'glue,') A peculiar gelatinous tumour, consisting of a remarkably soft gelatiniform tissue, which trembles on being touched.

COLLOSTRUM, Colostrum.

COLLOURION, (κωλλουριον,) Collyrium.

COL'LUM, *Cer'vix*, *Trache'us*, *An'chen*, *Neck*, *Halae*, (Prov.) *Hause*, *Hose*, (F.) *Col*, *Con*. The part of the body situate between the head and chest. Also, a part of an organ resembling the neck, as in the following cases.

COLLUM ASTRAG'ALI, *Cer'vix Astrag'ali*, *Neck of the Astragalus*, (F.) *Col de l'astragale*. A depression, which separates the anterior extremity of the astragalus from its body.

COLLUM COSTA'RUM, *Cer'vix Costa'rum*, *Neck of the Ribs*, (F.) *Col des Côtes*. The narrow part of the ribs, between the head and tubercle.

COLLUM DEX'TIUM, *Cer'vix Den'tium*, *Neck of the Teeth*, (F.) *Col ou Collet des Dents*. The part of the teeth between the corona and fang, which is embraced by the gum.

COLLUM FEM'ORIS, *Cer'vix Fem'oris*, *Neck of the Thigh-bone*, (F.) *Col du Fémur*. The long, narrow, and oblique portion of the os femoris, which separates the head from the two trochanters.

COLLUM FIB'ULÆ, *Cer'vix Fib'ulæ*, *Neck of the Fib'ula*, (F.) *Col du Péroné*. A slight narrowness seated below the head or upper extremity of the fibula.

COLLUM GLANDIS, see Glans.

COLLUM HU'MERI, *Cer'vix Hu'meri*, *Neck of the Hu'merus*. A circular, depressed portion, which separates the head of the os humeri from its two tuberosities. Some surgeons place the neck below the tuberosities, no precise line of demarcation indicating its extent.

COLLUM MANDIB'ULÆ seu MAXIL'Æ INFERIO'RS, *Cer'vix Mandib'ulæ seu Maxil'æ Inferio'ria*, *Neck of the lower jaw*, (F.) *Col de l'os maxillaire inférieure*. A depression observable on each side of the lower jaw, immediately below the condyles.

COLLUM OBSTIPUM, Torticoillis.

COLLUM OS'NIS MAG'NI seu CAPITA'TI, *Cer'vix ossis mag'ni seu capitati*, *Neck of the Os Mag'num*,

(F.) *Col du grand Os*. A circular depression beneath the head of this bone of the carpus.

COLLUM PEDIS, Intep.

COLLUM RA'DII, *Cer'vix Ra'dii*, *Neck of the Ra'dius*, (F.) *Col du Ra'dius*. A narrow portion of the radius, which supports the head of the bone.

COLLUM SCAP'ULÆ, *Cer'vix Scap'ulæ*, *Neck of the Scap'ula*, (F.) *Col de l'Omoplate*. A narrow portion, observable below the glenoid cavity of the scapula, which seems to separate it, as it were, from the rest of the bone.

COLLUM U'TERI, *Cer'vix seu Jug'ndum seu Can'lis U'teri*, *Neck of the U'terus*, (F.) *Col de la Matrice*, *Col utérin*. A narrow, cylindrical, and flattened portion of the uterus, which terminates inferiorly, and opens into the vagina by the *Os Uteri* or *Os Tincæ*. This neck is perceptible on examination *per vaginam*, until near the end of utero-gestation. As the uterus enlarges, however, it becomes shorter and wider, so that a manifest difference exists between its condition at seven and nine months.

COLLUM VESI'CÆ, *Cer'vix vesicæ*, *Neck of the Blad'der*, (F.) *Col de la Vessie*. The anterior part of the base of the bladder, which resembles the neck of a bottle, and is continuous with the urethra.

COLLURIUM, (κωλλυριον,) Collyrium.

COLLUTIO, (*col*, and *luere*, *lutum*, 'to wash,') Gargarism.

COLLUTION, Lotion.

COLLUTORIUM, (same etymon as *Collutio*), (F.) *Collutoire*. A mouth wash.

COLLUTORIUM ADSTRINGENS, Mel boracis.

COLLU'VIES, Filth, excrement. Discharge from an old ulcer.

COLLUVIES GASTRICA, *Embarras gastrique*.

COLLYRIUM, *Collu'rium*, *Collu'rium*, (F.) *Collyre*, (from κωλλω, 'I check,' and ρέω, 'I flow,' or from κωλλα, 'glue,' and ορπα, 'tail,' [?] or from κωλλυρα, 'a cylindrical cake,') (F.) *Collyre*. The ancients designated, by this term, a solid medicine, of a long, cylindrical shape; proper to be introduced into the vagina, anus, or nostril. They are said to have given it this name because it was shaped like a *Rat's Tail*, and because there entered into its composition powders and glutinous matters.—At the present day, *Collyrium* means an application to the eye. Some are *puberulent* and dry, *Ophthalmopas'mata*, *Colly'ria sic'ca*, but the greatest part are *liquid*, *Hygrocolly'ria*; and receive different epithets, as *astringent*, *emollient*, &c. The term is now little more than synonymous with *Eye-water*. *Collyria* are generally extemporaneous formulæ. When the wash is applied to the closed eyelids, it is sometimes called *Lava'erum ophthalmicum*.

COLLYRIUM AMMONIACA'LE, *Pulvis Leayso'ni*, (F.) *Collyre ammoniacale*, *Poudre de Leayson*. A preparation, much used in France in amaurosis and nervous debility of the eye, and which consists of a union of *slacked lime*, *muriate of ammonia*, *charcoal*, *cinnamon*, *cloves*, and *bole Armeniac*, moistened, and kept in a well-stopped bottle. In employing it, the stopper being removed, the bottle should be moved beneath the affected eye.

COLLYRIUM BATEANUM, see Lotion, camphorated, of Bates—c. *Siccum*, see *Collyrium*—c. *Siccum Alexandrinum*, see *Alexandrine*.

COLOBOMA, gen. *Colobo'matis*, (κολοβωμα, from κολβω, 'I maim,' 'any thing truncated or shortened,') A mutilated or maimed organ.

COLOBOMA I'RDIS, *Frido-colobo'ma*. A congenital peculiarity of the iris, consisting of a fissure of its lower portion, and a consequent prolongation of the iris to the margin of the cornea. See *Iridorrhagas*.

COLOBOMA ΠΑΛ'ΕΒΡÆ. A vertical wound of the eyelid, passing through it so as to divide it into two flaps, somewhat resembling the two portions of a harelip.

COLOBO'SIS, (κολωβωσις.) The act of curtailing or mutilating; mutilation.

COLOCHOL'SIS, *Cholodyenter'ia*, (κωλον, 'the colon,' χολη, 'bile,' and οσις.) Bilious dysentery, *Dysenter'ia bilio'sa*.

COLOCLET'SIS, (*colon*, and κλεισις, 'occlusion.') Occlusion of the colon.

COLOCYNTH, (κολοκυνθις.) *Cucumis colocynthis*.

COLOCYNTHIN, see *Cucumis colocynthis*.

COLOCYNTHITE, see *Cucumis colocynthis*.

COLOMBA, Calumba.

COLOMBINE, COMMON, *Aquilegia vulgaris*.

COL'ON, (κωλον.) *C. Cæcum*, *Monen'terum*, *Col'um*, *Intesti'nium majus*, *I. cellula'tum*, *I. cras'sum* et *ple'nium*, *I. gran'de*, *I. lax'um*, *Phys'scæ*. That portion of the large intestines which extends from the cæcum to the rectum. (It is said to have been so called from κωλος, 'hollow,' or from κωλον, 'I arrest,' because the excrements are arrested, for a considerable time, in its sacs,—*cæ'lulæ seu haust'ra seu locula'men'ta col'i*.) The colon is usually divided into four portions. 1. The right lum'bar or ascend'ing, *Col'on dex'trum*, situate in the right lumbar region, and commencing at the cæcum. 2. Transverse colon, *Col'on transver'sum*, transverse arch of the colon, the portion of the colon which crosses from the right to the left side, at the upper part of the abdomen. 3. The left lum'bar or descend'ing colon, *Col'on sinis'trum*, extending from the left part of the transverse arch, opposite the outer portion of the left kidney, to the corresponding iliac fossa. 4. The Iliac colon or Sig'moid flex'ure of the colon, (F.) *Colon ili'aque ou S. du colon*, the portion of the intestine which makes a double curvature in the left iliac fossa, and ends in the rectum.

The muscular fibres, as in the cæcum, are in three flat stripes, *Tæ'nix seu Fas'ciæ Ligamen'to'sæ Col'i*, *Tæ'nix longitudina'les seu Vasa'l'væ*, *Ligamen'ta col'i*.

COLON, (κωλον.) Membrum—c. Inflammation of the, *Colitis*—c. Occlusion of the, *Colocleisis*.

COLON, TOR'POR OF THE. A disease in which the muscular coat of the colon acts with deficient energy; giving occasion to distension of the intestine, which, by pressing upon the other organs, may interfere so much with their functions, as to lead to distressing gastric, cardiac and other disorders.

COLONITIS, (*colon*, and *itis*.) *Colitis*, *Dysentery*.

COLONNE, Columna—c. *Vertébrale*, Vertebral column.

COLONNES CHARNUES DU CŒUR, Columnæ carneæ.

COLOPHANE, Colophonia.

COLOPHANY, Colophonia.

COLOPHO'NIA, (from Colophon, a city of Ionia,) *Fric'ta*, *Pix Græ'ca*, *Resi'na n'gra*, *Col'ophany*, *Col'ophony*, *Black Ros'in*, *Pitch*, *Brown Ros'in*, (F.) *Colophone*, *Colophane*, *Arcanson*, *Brai sec*. The black resin which remains in the retort, after the distillation of common turpentine, by means of a strong fire. It is used like the turpentine in general, and in some pharmaceutical preparations.

COLOPHONY, Colophonia.

COLOQUINTE, *Cucumis colocynthis*.

COLOQUINTIDA, *Cucumis colocynthis*.

COLOR VIRGINEUS PALIDUS, ('pale color of virgins,') *Chlorosis*—c. *Virginum fœdus*, *Chlorosis*.

COLORRECTITIS, (*colon*, *rectum*, and *itis*.) *Dysentery*.

COLOSTRA, *Colostrum*.

COLOSTRATIO, *Colostration*.

COLOSTRA'TION, *Colostratio*. Disease in new-born children, attributable to the colostrum.

COLOS'TRUM, *Colos'tra*, *Collus'trum*, *Colus'trum*, *Troph'alis*, *Protog'ala*, *Neog'ala*, *Prim'mum Puer'peræ Lac*, *Pyos*, *Pyar*, *Py'tia*, *Pye'tia*, *Green Milk*, *Beast'lings*, *Beist'ings*, *Beest'ings*, *Beist'ings* or *Beast'ings*, *Beest*, (Sc.) *Beist*, *Beist'yn*, (Prov.) *Fore'milk*, *Poad'milk*, *Po'red milk*, *Pow'd'milk*, in the cow, &c., (from κωλον, 'food' [?], or κωλλα, 'glue,' on account of the viscosity of the first milk. [?]) (F.) *Béton*. The first milk after accouchement. It contains more serum and butter, and less casein than common milk, and seems to possess a laxative property, which renders it fit to aid in the expulsion of the meconium.

Colostrum formerly meant an emulsion prepared of turpentine and yolk of egg.

COLOT'OMY, *Colotom'ia*, (*colon*, and τομή, 'incision.') The operation of cutting into the colon, as for the formation of an artificial anus.

COLOUR-BLINDNESS, *Achromatopsia*.

COL'OURS, ACCIDENTAL, *Op'posite col'ours*, *Complement'ary* or *Harmon'ic col'ours*, (F.) *Couleurs Complémentaires*. If the eye has been for some time regarding a particular colour, the retina becomes insensible to this colour; and if, afterwards, it is turned to a sheet of white paper, the paper will not seem to be white, but will be of the colour that arises from the union of all the rays of the solar spectrum, except the one to which the retina has become insensible. Thus, if the eye be directed for some time to a red wafer, the sheet of paper will seem to be of a bluish-green, in a circular spot of the same dimensions as the wafer. This bluish-green image is called an oc'ular spec'trum, because it is impressed upon the eye and may be retained for a short time; and the colour bluish-green, is said to be the accidental colour of the red. If this experiment be made with wafers of different colours, other accidental colours will be observed, varying with the colour of the wafer employed, as in the following table:—

Colour of the Wafer.	Accidental colour, or colour of the ocular spectrum.
Red.....	Bluish green.
Orange.....	Blue.
Yellow.....	Indigo.
Green.....	Violet with a little red.
Blue.....	Orange red.
Indigo.....	Orange yellow.
Violet.....	Yellow green.
Black.....	White.
White.....	Black.

If all the colours of the spectrum be ranged in a circle, in the proportions they hold in the spectrum itself, the accidental colour of any particular colour will be found directly opposite. Hence, the two colours have been termed *op'posite col'ours*. It will follow from what has been said, that if the primary colour, or that to which the eye has been first directed, be added to the accidental colour, the result must be the same impression as that produced by the union of all the rays of the spectrum—white light. The accidental colour, in other words, is what the primitive colour requires to make it white light. The primitive and accidental colours are, therefore, *com'plements* of each other; and hence accidental colours have also been called *complement'ary col'ours*. They have likewise been termed *harmon'ic*, because the primitive and its accidental colour *harmonize* with each other in painting.

COLPAL'GIA, (*colpo*, and αλγος, 'pain,') *Vaginodyn'ia*. Pain in the vagina.

COLPATRE'SIA, *Elyt'ratre'sia*, *Ancylocol'pus*, *Atre'tytr'ia*, *Atre'sia vagi'næ*, (*colpo*, and αρησος, 'without opening,') Imperforation of the vagina.

COLPEMPHRAX'IS, (*colpo*, and ἐμπράξις, 'ob-

struction.') Obstruction of the vagina by foreign bodies.

COLPEURYN'TER, (*colpo*, and *εὐρυνω*, 'I dilate, I enlarge,') Speculum vaginae. Also, an instrument for dilating the os uteri, and upper part of the vagina. The operation is called *Colpeurynsis*.

COLPITIS, (*colpo*, and *itis*), Colpositis.

COLPO, (*κολπος*, 'vagina,') In composition, vagina.

COLPOC'ACE, *Ædæotitis gangrenosa*, *Gangrena genitalium et vaginæ*, (*colpo*, and *κακος*, 'evil,') Putrescency or gangrene of the vagina and labia.

COMPOACE INFANT'ILIS, *Ædæotitis gangrenosa puella rum*, *Noma pudendo rum*. Gangrene or putrescency of the vagina and genitals in young children.

COMPOACE PUERPERA'RUM, *Ædæotitis Gangrenosa puerpera rum*, *Tocodomyeodoritis maligna vagina lis*. Sloughing of the vagina and genitals in puerperal women.

COLPOCELE, (*colpo*, and *κλη*, 'rupture,') Elytrocele.

COLPOCYSTOTOM'IA, (*colpo*, *κυστις*, 'bladder, and *τομή*, 'incision,') *Sectio vaginovescicalis*. Lithotomy through the vagina.

COLPODESMORRAPH'IA, (*colpo*, *δεσμος*, 'ligament, and *ρῆψις*, 'suture,') The removal of a part of the mucous membrane of the vagina for the radical cure of prolapsus vaginae et uteri.

COLPEDEMA, gen. *Colpædematis*, (*colpo*, and *οἰδημα*, 'a swelling,') Elytroedema.

COLPOPHLEGHYMENTIS, (*colpo*, *φλεγμα*, mucus, 'ὑμη', 'a membrane, and *itis*,) Encolpitis.

COLPOPTOSIS, (*colpo*, and *πτωσις*, 'a fall,') Prolapsus vaginae.

COLPORRHA'GIA, *Elytrorrhagia*, (*colpo*, and *ῥήγνυμι*, 'I break forth,') Discharge of blood from the vagina.

COLPORRHAPHY, (*colpo*, and *ρῆψις*, 'a suture,') Elytrorrhaphy.

COLPORRHEX'IS, *Ruptura vaginæ*, (*colpo*, and *ῥήξις*, 'rupture,') Rupture of the vagina. Also, colporrhagia.

COLPORRHEA, (*colpo*, and *ῥεω*, 'I flow,') Leucorrhœa.

COLPO'SIS, (*κολπος*, 'vagina, and *osis*,) Inflammation of the vagina. Synonymous with *Elytritis*, *Elytritis*, *Oleitis*, *Colpitis*, *Vaginitis*. See Leucorrhœa.

COLPOSTEGN'OSIS, *Coleostegno'sis*, (*colpo*, and *στεγνωω*, 'I close,') Atresia, or obliteration of the vagina.

COLPOSTENOCHORIA, (*colpo*, *σπρος*, 'narrow, and *χωρος*, 'space,') Coleostegnosis.

COLPOSYNZESIS, (*colpo*, and *συνζωω*, 'falling together,') Coleostegnosis.

COLPOT'OMY, *Colpotom'ia*, (*colpo*, and *τομή*, 'incision,') An incision of the vagina in parturition.

COLPUS, (*κολπος*,) Mamma, Sinus, Vagina.

COLT'S FOOT, *Asarum Canadense*, Tussilago.

COLUBER ('a serpent,') BERUS. The systematic name of the viper. *Vipera*, (F.) *Couleuvre*, *Vipère*. *Class*, Reptilia, *Ord*. Ophidia. A poisonous reptile—the poison lying in small sacs near its teeth. The flesh is innocent, and has been often taken in scrofula, and in cutaneous disorders in general, but it is inefficacious.

COLUBRINA, *Polygonum bistorta*—c. Dracontia, *Arum dracunculus*—c. Lusitania, *Euphorbia capitata*—c. Virginiana, *Aristolochia serpentaria*.

COLUM, Colon.

COLUMBIA (N. Y.), MIN'ERAL WA'TERS OF. See New York, mineral waters of.

COLUMBINE, *Aquilegia vulgaris*—c. Wild, *Aquilegia Canadensis*.

COLUMBO, Calumba—c. American, see Calumba—c. Marietta, see Calumba.

COLUMELLA, (dim. of *columna*,) Pillar, Uvula—c. Cochleæ, Modiolus.

COLUM'NA, *Columna*, (F.) *Colonne*. Anatomists use this word for parts which resemble a column or pillar; hence, for the penis.

COLUMNA ADSTANS INGUINIBUS, Penis—c. Dorsi, Vertebral column—c. Foraminis ovalis, see *Ovalis fossa*—c. Fornicis, see Fornix.

COLUM'NA NASI. The cartilaginous part of the septum of the nostrils. See Nares.

COLUMNA ORIS, Uvula—c. Spinalis, Vertebral column—c. Valvulae Viessensii, see *Valvula Viessensii*—c. Vertebralis, Vertebral column—c. Virginitalis, Hymen.

COLUM'NÆ BERTINI, *Columnæ of Bertin*. Processes of the cortical substance of the kidney, which extend between the pyramids of Malpighi, as far as the hilus.

COLUM'NÆ CAR'NEÆ, ('fleshy columns,') *C. Cordis*, *Lacer'ti* seu *Lacer'tuli* seu *Fu'nies* seu *Fasciculi ter'etes Cordis*, *Trabes* seu *Trabeculæ Cordis* seu *Car'neæ*, (F.) *Colonnes charnues du cœur*. Small, fleshy columns, which project, more or less, into the auricles and ventricles of the heart, whose use appears to be to prevent too great dilatation of those cavities. A few of these *columnæ*—see *Musculi papillares*—are attached by one extremity to the walls of the heart, and, by the other, give insertion to chordæ tendineæ. See Carneous columns.

COLUMNÆ CARNEÆ of the Rectum, see Rectum—c. Papillares, see *Columnæ Carneæ*.

COLUMNÆA LONGIFOLIA, Babel.

COLUMNS OF BERTIN, *Columnæ Bertini*.

COLUMNS, CARNEOUS, see *Columnæ carneæ*—c. Posterior Median, of the medulla oblongata, *Funiculi graciles*—c. of Morgagni or of the Rectum, see Rectum—c. of the Spinal Marrow, see Vertebral Nerves.

COLUS JOVIS, ('staff of Jove,') *Salvia sclarea*.

COLUSTRUM, Colostrum.

COLU'TEA, (*κολυτεια*,) *C. Arborescens* seu *hir-suta*, *Senna Germanica*, *Bladder Senna*, (F.) *Baguenaudier*, *Faux Séné*, *Séné indigène*. *Ord*. Leguminosæ. *Sec. Syst*. *Diadelphia Decandria*. The leaves are slightly purgative, and are often mixed with those of the cassia senna.

COLUVRINE DE VIRGINIE, *Aristolochia serpentaria*.

COLYMB'ADES, (*κολυμβάδες*,) *Pickled Olives*. These, when bruised and applied to a burnt part, were supposed to be able to prevent vesication.

COLYMBIFERA MINOR, *Mentha crispa*.

COM, as a prefix, like *cum*, 'with.'

CO'MA, gen. *Co'matis*, (*κωμα*,) *Semisom'nia*, *Semisop'itus*, *Semisopo'rus*, *Subeth*. A profound state of sleep, from which it is extremely difficult to rouse the individual. It is a symptom which occurs in many diseases. Two varieties are distinguished, 1. *Coma vi'gil* seu *agrypnod'es*, *Pervigilium*, *Vigil'ie nim'ie*, *Typho'nia*, *Veter'nia*, *Veterno'sitas*, *Agrypnoco'ma*, *Carus lethargus vigil*, *Typhoma'nia*, which is accompanied with delirium. The patient has his eyes closed, but opens them when called; and closes them again immediately. This state is accompanied with considerable restlessness. 2. *Coma Somnolentum*, *C. Comato'des*;—in which the patient speaks when roused, but remains silent and immovable in the intervals. Coma is a deeper sleep than *sopor*, but less so than *lethargy* and *carus*.

COMA, gen. *Comæ*, (*κομή*, from *κομω*, 'I adorn,') *Capillus*—c. *Agrypnodes*, see *Coma*—c. *Apoplexia*, *Apoplexy*—c. *Cæsarea*, *Plica*—c. *Comatodes*, see *Coma*—c. *Somnolentum*, *Somnolency*, see *Coma*—c. *Vigil*, see *Coma*.

COMACON, *Myristica moschata*.

COMAN'DRA UMBELLA'TA, (*κομή*, 'hair,')

and *anthes*, 'stamina,' in reference to the hairy tufts attached to the anthers,) *Bas'tard Toadflax*: indigenous; *Order*, Santalaceæ: flowering in May and June; has been used in fevers by some of the Indian tribes.

COM'ARUM PALUS'TRĒ, *Potentilla palustris*, *Marsh Cinquefoil*. An indigenous plant, *Family*, Rosaceæ, which flowers in June. It is possessed of astringent virtues.

COMARUS, (κομᾶρος,) *Arbutus unedo*.

COM'ATA, (from κομᾶ.) Diseases characterized by diminution of the powers of voluntary motion, with sleep or impaired condition of the senses.

COMATEUX, Comatose.

COMATODES, (καταρῶδες,) Comatose.

COMATOSE, *Comato'des*, *Caro'ticus*, *Caro'des*, (F.) *Comateux*. Relating to or resembling coma; —as *comatose sleep*, *comatose fever*, &c.

COMBUSTIBILITY, PRETERNATURAL, Combustion, human.

COMBUSTIO, (comburare, combustum, (com, and urere, ustum,) 'to burn,') Burn—c. Spontanea, Combustion, human.

COMBUSTION, HU'MAN, *Spontaneous Combustion* or *Preternatural Combustibility*, *Catacau'sis*, *Incedium spontaneum*, *Combustio spontanea*, *Autempra'mus*, *Idiopocau'sis*, *Idiempre'sis*, *Catacau'sis ebrio'ea*, (F.) *Combustion humaine*, *C. humaine spontanée*. These terms have been applied to the extraordinary phenomenon of a rapid destruction of the human body, by being reduced to ashes either spontaneously or by the contact of an ignited substance. It is said to have occurred in the aged, and in those that were fat and hard drinkers. In such, Dr. Traill has found a considerable quantity of oil in the serum of the blood. Vieq d'Azyr, Lair, and Dupuytren think it necessary, that the body should be placed in contact with an ignited substance. Le Cat, Kopp, and Marc are of opinion that this is not necessary. The former appears to be the more probable view.

COMBUSTURA, Burn.

COMEDONES, (comedere, 'to waste,' or comedo, 'a glutton,') see Aene.

COMES ARCHIATRORUM, *Comes*, gen. *Com'itis*, 'an associate,') see Exarchiater.

COMESTIBLE (F.), *Edu'lis*, (com, and edere, 'to eat,') *Eatable*, (F.) *Édule*. Esculent. When this word is used substantively, in French, as in English, it means solid food.

COMÉZ. Half a drop.

COMFIT, (conficere (com, and facere) 'to make,') Confection.

COMFREY, *Symphytum*—c. Spotted, *Pulmonaria*—c. Wild, *Cynoglossum Virginicum*.

COMISTE, (κομίστη,) Aliment.

COMMANDUCATIO, (com, and manducare, 'to chew,') Mastication.

COMMEMORATIVE, *Commemorati'vus*, (F.) *Commemoratif*, (commemorare (com, and memor) 'to cause to remember,') That which recalls, or rather which is recalled.

COMMEMORATIVE SIGNS, (F.) *Signes commémoratifs*, are those deduced from circumstances which have gone before; or rather, according to others, those which point out the preceding condition of the patient. They are also called *Anamnetic signs*.

COMMENSUM, (com, and metior, mensus, 'to measure,') Symmetry.

COMMERCIUM, 'communication,' (from com, and merc, mercis, 'merchandise,') Sympathy.

COMM, (κομᾶ,) Gummi.

COMMUNUTED, *Comminu'tus*, (comminuere, (com, and minuo, 'to break to pieces,') (F.) *Comminutif*. A comminuted fracture is one in which the bone is broken into a number of pieces.

COMMUNU'TION, *Comminu'tio*, *Thryp'sis*, *Lei-*

o'sis. Same etymon. Fracture of a bone into a number of pieces.

COMMISSURA, (committo, commissum, (com, and mitto,) 'I join together,') Articulation, Commissure.

COMMISSURA ANTERIOR CEREBRI, Commissure, anterior, of the Brain.

COMMISSURA BRE'VIS. A lobule or prominence of the inferior vermiform process of the cerebellum, situate in the incisura posterior, below the horizontal fissure.

COMMISSURA LABIORUM, Prostomia—c. Magna cerebri, Corpus callosum—c. Maxima cerebri, Corpus callosum—c. Nerveæ, Syndesmosis—c. Ossium carnea, Syssarcosis.

COMMISSURA POSTERIOR CEREBRI, Commissure, posterior, of the Brain.

COMMISSURA SIM'PLEX. A small lobule or prominence of the superior vermiform process, near the incisura posterior of the cerebellum.

COMMISSU'RAL. Of or belonging to a commissure.

COM'MISSURE, *Commissu'ra*, *Compa'ges*, *Compagina'tio*, *Sym'physis*, *Sym'bole*. A point of union between two parts: thus, the commissures of the eyelids, lips, &c., are the angles, which they form at the place of union. See *Fibres*, converging.

COMMISSURE, ANTERIOR, OF THE BRAIN, *Commissu'ra anterior cer'ebri*, (F.) *Commissure antérieure du cerveau*. A small medullary fasciculus, situate transversely at the anterior part of the third ventricle, and uniting the two hemispheres.

COMMISSURE, GREAT, OF THE BRAIN, *Commissu'ra Mag'na Cer'ebri*, (F.) *Grande commissure du cerveau*, which unites the two hemispheres for some extent, is the *Corpus callo'sum*.

COMMISSURE, MIDDLE, OF THE BRAIN. A layer of gray substance uniting the thalami optici.

COMMISSURE, OBLIQUE OR INTERCEREBRAL, see *Valvula Vieussenii*—c. Optic, see *Chiasmus*.

COMMISSURE, POSTERIOR, OF THE BRAIN, *Commissu'ra poste'rior cer'ebri*. A medullary band, situate at the posterior part of the third or middle ventricle.

COMMISSURE OF THE UVEA, Ciliary ligament.

COMMISSURE DE LA CHOROÏDE, Ciliary ligament.

COMMOS'IS, (κομμοσις, from κομω, 'I adorn,') The art of concealing natural deformities, as by painting the face. See, also, *Propolis*.

COMMOTICE, (κομμοτική (τεχνή,) in the older writers, meant the art of communicating factitious beauty to a person. Painting the face; *Compor'ia*.

COMMOTIO, (com, and motio,) Motion, Concussion—c. Cerebri, Concussion of the brain.

COMMOTION, Concussion—c. du Cerveau, Concussion of the brain.

COMMUNICABLE, (communicans, and habilis, 'able,') See *Contagious*.

COMMUN'ICANS, (communicare, 'to make common,') *Con'jgens*. That which communicates or establishes a communication. Communicant. There are two *Arteriæ Communican'tes*, both within the cranium;—the one anterior, very short, and extending transversely from one anterior cerebral artery to the other,—the other posterior, called also *Communicans Willis'ii*, which passes from the internal carotid to the posterior cerebral artery. It is a branch of the basiliary.

COMMUNICANS NO'NI. A long slender nervous branch, formed by filaments from the first, second and third cervical nerves, which descends upon the outer side of the internal jugular vein, and forms a loop with the descendens noni over the sheath of the carotids.

COMMUNICANS PERONEI, see *Communicans poplitei*.

COMMUNICANS POPLITEI, *C. tibialis* (*ner'vus*.) A large nerve, which arises from the popliteal, and, at a variable distance below the articulation of the knee, receives the *communicans peronei* from the peroneal nerve,—the two forming the *external saphenous nerve*.

COMMUNICANS TIBIALIS, *Communicans poplitei*.

COMPACT, *Compac'tus*, (*com*, and *pangere*, *pac-tum*, 'to strike, to fix.') Solid, close. (F.) *Com-pacte*. The term *Compact Tissue* is given to the hardest and closest parts of bone.

COMPAGES, ('a setting together.') Articulation, Commisura—*c. Ossium per Lineam Simpliciem*, Harmony—*c. Verteborum*, Vertebral column.

COMPAGINATIO, Commisura.

COMPAS D'ÉPAISSEUR, (*com*, and *pandere*, *passum*, 'to extend,') see *Pelvimeter*.

COMPASS PLANT, *Silphium laciniatum*.

COMPASSIO, (*com*, and *pator*, *passus*, 'to suffer,') Sympathy.

COMPEBA, *Piper cubeba*.

COMPENSATION, (*com*, and *pensare*, *pensa-tum*, 'to weigh,') *Balancement*.

COMPEPER, *Piper cubeba*.

COMPERNIS. One who has his knees turned inwards. A case of distortion of the legs.

COMPETENTIA MEMBRORUM OMNIUM, ('fitness of all the members,') Symmetry.

COMPLAINT, (*com*, and *plangere*, *plantum*, 'to beat the breast for grief,') Disease—*c. Family*, see *Hereditary*.

COMPLEMENTAL AIR, (*com*, and *pleo*, 'to fill,') See *Respiration*.

COMPLEMENTARY AIR, see *Respiration*.

COMPLETIO, *Plethora*.

COM'PLEX, *Complex'us*, (*com*, and *plectere*, *plexum*, 'to twist,') Embracing several distinct things. *Chaussier* uses this term, in his anatomical descriptions, for *complicated*.

COMPLEXIO, *Coition*, *Complexion*, *Confusio*, *Temperament*.

COMPLEX'ION, *Complexio*. Often employed, in English, for the colour of the face, as 'He has a good complexion,'—a 'sallow complexion,' &c. It formerly had a more extensive signification, and still has in France. It signifies the aggregate of physical characters presented by any individual, considered with respect to his external arrangement or condition. It means more than *constitution*, for which it is used synonymously in many cases; and differs from *temperament*, which is less the external condition of the body than the state or disposition of the organs in health.

COMPLEXUS, *Complex*.

COMPLEXUS M'JOR, *Mastoidæus lateralis*, *Trachælo-mastoidæus*, (F.) *Trachælo-mastoidien*, *Muscle petit Complexus*. It arises from the transverse processes of the last four cervical vertebrae, and is inserted into the mastoid process.

COMPLEXUS MUS'culus, *Bil'venter Cervicis*, *Complexus Ma'jor*, *Dorso-trachælo-occipital*, *Trachælo-occipital* (Ch.), (F.) *Muscle grand complexus*. A muscle, situate at the hind part of the neck, where it extends from the interval that separates the two prominent ridges on the posterior surface of the os occipitis to the transverse and articular processes of the last six cervical vertebrae, as well as to those of the first five dorsal. It serves to straighten, incline, and turn the head.

COMPLICAT'ION, *Complicatio*, (*com*, and *plicare*, *plicatum*, 'to fold,') In medicine, it means the presence of several diseases, *morb'i complicati seu perplex'i*, or of several adventitious circumstances foreign to the primary disease.

COMPOSIT'ION, *Composit'io*, (*componere*, (*com* and *ponere*, *positum*, 'to place,') 'to place together,') *Syn'thesis*. The act of composing or compounding,—of medicines for example; *lama-*

toyntax'is. Also, the *Com'pound*, *Compos'itum*, or thing compounded. Likewise, a combination.

COMPOSITUM, *Compound*.

COMPOUND. To mix medicines. To mingle different ingredients into one whole. Used adjectively, *com'pound* signifies the result of the union of several medicinal agents, as 'a compound medicine.'

COMPREHENSIO, (*com*, and *prehendere*, *pre-hensum*, 'to take hold of,') *Catalepsy*.

COMPRESSIO, *Catalepsy*.

COMPRESS, *Compres'sa*, *Sple'nium*, *Spleni'ola*, *Splenis'cus*, *Pla'gula*, *Penicil'lum*, *Penic'ulum*, (*comprimere*, (*com* and *primere*, *pressum*, 'to press,') 'to press together,') (F.) *Compresse*. Folded pieces of lint or rag, so contrived as, by the aid of a bandage, to make due pressure upon any part. According to their shape, direction, and use, compresses have been called *long* ((F.) *longuettes*,) *square* (*carrées*,) *triangular*, *prismat'ic*, *graduated* (*graduées*,) *split* (*fendues*,) *fend'ées*, *criblées*, *croix de Malte*, *oblique*, *cir'cular*, *divi'ding* (*divisives*,) *uni'ting* (*unisantes*,) *crib'ri-form*, &c.

The *Umschlag* or compress of the hydropathists is a cloth, well wetted with cold water, applied to the surface near the supposed seat of disease, securely covered with a dry cloth, and changed as often as it becomes dry. It is sometimes covered with a layer of oiled silk, to prevent evaporation.

COMPRESSEUR DE DUPUYTREN, *Compressor of Dupuytren*—*c. du Nez*, *Compressor nasi*—*c. de la Prostata*, *Compressor prostatae*.

COMPRESSIF, *Compressive*.

COMPRESSIO, (same etymon as *Compress*,) *Compression*, *Thlipsis*.

COMPRES'ION, *Compressio*, (same etymon.) Pressure; methodical compression. An agent frequently had recourse to in surgery. We *compress* a limb, affected with *œdema*, *varices*, *hydrops articuli*, *callous ulcer*, &c. The abdomen is *compressed* after delivery, after *paracentesis abdominis*, &c. The compression is produced by means of the roller, laced stocking, &c., according to the part, and to the particular case. *Moderate pressure* aids the contractility of parts, and modifies their nutritive action; so that large tumours at times disappear after it has been used for some time. A greater degree of pressure occasions, still more, the emaciation of the part, but it is apt to impede the circulation. *Pressure* is often used to stop or moderate the flow of blood in cases of aneurism or wounds of arteries and veins. In such cases, the compression may be *immediate*, when applied upon the artery itself, or it may be *mediate*, when applied through the integuments and soft parts. The French use the term *Compression immédiate latérale* for that which is exerted perpendicularly to the axis of a vessel, so as to flatten its sides. It is practised with the finger, forceps, tourniquet, &c.

Methodical Compression is pressure applied *secundum artem*. *Dig'ital Compression* is pressure made by the fingers.

COMPRESSION OF THE BRAIN. This may arise either from coagula of blood, a soft tumour, a bony excrescence, a depressed portion of the skull, or the presence of some foreign body. The effects vary, according as the compression takes place *suddenly* or *gradually*. When *suddenly*, the symptoms are of the comatose or apoplectic character. When *gradually*, mania, convulsions, &c., are more likely to occur. Compression, arising from a depressed portion of skull, requires the use of the trephine.

COMPRES'SIVE, *Compress'ious*, (F.) *Compressif*. That which compresses. A *compressive bandage* is one that compresses the parts to which it is applied.

COMPRESSOR or COMPRESSORIUM OF DUPUYTREN, (F.) *Compresseur de Dupuytren*. An instrument for compressing the femoral artery, invented by Dupuytren. It is constructed on the same principles as the tourniquet of J. L. Petit, from which it only differs in this respect;—that, instead of being maintained in its place by a strap, which always compresses more or less the circumference of the limb, the two pads are placed at the extremities of a semicircle of steel, which, by passing from one to the other without touching the parts, limits the pressure to two opposite points of the thigh, and permits the collateral circulation to go on.

COMPRESSOR NA'RIS, Rina'us, Nasa'lis, Transversalis Na'si, Myrtiformis, Dilato'res alu'rum na'vi, Constrictor Na'si seu Na'rrium seu Na'ris, Triangula'ris Na'si, (F.) *Mazillo-narinal, Sus-mazillo-nasal*—(Ch.,) *Compresseur du nez, Transversal du nez*. A muscle, situate at the sides of the nose; flat and triangular. It arises from the inner part of the fossa canina, and passes to the dorsum of the nose; where it is confounded with that of the opposite side.

COMPRESSOR or CONSTRICTOR OF URETHRA. An instrument for compressing the urethra, to obviate incontinence of urine. It consists of a girdle of iron, which surrounds the pelvis, to which is fixed a plate of the same metal, that compresses the urethra in *perineo*.

COMPRESSOR PROST'ATE, Prostat'icus super'ior, Pubioprostat'icus, Subpubioprostat'icus, (F.) *Compresseur de la prostate*. A muscle, admitted by ALBINUS, which is formed of the anterior fibres of the *Levator ani*, that embrace the prostate. It is the *Prostat'icus super'ior* of WINSLOW.

COMPRESSOR URE'THRE. A muscle consisting of two portions—one of which is *transverse* in its direction, and, in consequence of its having been particularly described by Mr. Guthrie, has been called *Guthrie's muscle*. It arises from the ramus of the ischium, and passes inwards to embrace the membranous urethra. The other portion is *perpendicular*, descending from the pubis and passing down to be inserted into the transverse portion of the muscle;—*Pub'io-urethra'lis*. This portion has been considered by many to be only the anterior fibres of the levator ani; and having been described by Mr. Wilson, it has been called *Wilson's muscle*.

COMPRESSOR VE'NÆ DORSALIS PE'NIS. A small muscle, distinctly seen in animals, less distinctly in man, which arises from the ramus of the pubis, and ascending in a direction forwards is inserted above the vena dorsalis, joining with its fellow of the opposite side on the mesial line. Its use is supposed to compress the vein in erection. It is sometimes called the *muscle of Houston*, after Dr. Houston of Dublin.

COMPRESSORIUM, Compressor.

COMPTONIA ASPLENIFO'LLIA, (after Compton, Bishop of London.) *Liquidambar peregrina seu asplenifolia, Myrica asplenifolia, Sweet Fern, Shrub by Sweet Fern, Sweet bush, Fern bush, Fern gale, Spleenwort bush, Meadow Fern, Astringent root*. An indigenous shrubby plant, which grows in sandy or stony woods from New England to Virginia. It possesses tonic and astringent properties, and is used as a domestic remedy in diarrhoea, &c., in the form of decoction.

COMPUNCTIO, (com, and pungere, punctum, 'to puncture,') Paracentesis, Puncturing.

CON, as a prefix, like cum, 'with.'

CONÆSTHESIS, Conæsthesia.

CONARIUM, (κων, 'a cone,') Pineal gland.

CONATUS, gen. Conat'us, (conor, conatus, 'to endeavour,') Effort.

CONCARNATIO, (con, and caro, gen. carnis, 'flesh,') Syssarcosis.

CONCASSER (F.), (conquassare, (con, and

quassare, 'to shake much,') 'to break to pieces' 'to comminute.'). To reduce roots, woods, &c. into small fragments, in order that their active principles may be more readily separated from them.

CONCAU'SA, Concaus'sa, (con, and causa,) *Synæ'tia*. A cause which co-operates with another in the production of disease.

CONCAVITAS CONCHULARIS CEREBRI, Infundibulum of the brain.

CONCAVUM, PEDIS (con, and cavus, 'hollow,') Sole.

CONCENTRANT'IA, (con, and centrum, 'a centre,') A name once given to absorbents of acids.

CONCENTRATION, Concentra'tio. A word sometimes used, in medical theories, to express an afflux of fluids, or a convergence of sensibility or of vital force, towards an organ. It is applied, also, to the pulsation of arteries, when not easily felt under the finger. *Pouls Concentré* is a term applied by the French to a pulse of the above character.

CONCEPTACULA SEMINARIA, Vesiculæ seminariæ.

CONCEPTACULUM, ('a receptacle,') Uterus, Vessel.

CONCEPTIO, Conception—c. *Vitiosa, Pregnancy, extra-uterine.*

CONCEPTION, Concep'tio, Concep'tus, (from concipio, (con, and capio, 'to receive,') 'to bring together,') *Oye'sis, Syllep'sis, Androlep'sia*. The impregnation of the ovum by the positive contact of the male sperm, whence results a new being. The whole subject of conception is most mysterious. It seems to occur as follows. During the sexual union, the male sperm passes along the uterus and Fallopian tubes: the fimbriated extremities of the latter size hold of the ovarium; and the sperm in this manner comes in contact with a matured ovum, and impregates it. The fecundated ovum remains some time in the ovarium, but at length bursts its covering, is laid hold of by the fimbriated extremity of the Fallopian tubes, and passes along the tube into the cavity of the uterus, where it remains for the full period of utero-gestation. Some are of opinion, that the ovum is not impregnated until it has entered the Fallopian tube, or uterus.

CONCEPTION, FALSE, Falsus Concep'tus, Spu'rium germen, (F.) *Fausse Conception, Faux germe*. An irregular, preternatural conception, the result of which is a mole or some similar production, instead of a properly organized foetus. See Mole.

CONCEPTUS. The first rudiments of the foetus, after conception. Also, conception.

CONCEPTUS FALSUS, Conception, false.

CONCHA, Conchus, (κογχη, κογχος, 'a concave shell,') A liquid measure, amongst the Athenians, equal to half an ounce. Anatomists apply this term to several hollow parts of the body;—as the *Concha of the Ear*,—*Concha Auris, Concha Auriculæ*; (F.) *Conque*—the hollow part of the cartilage of the external ear. It has, also, been applied to the genital organs of the female; to the patella, &c.

CONCHA, Patella, Turbinate bone (middle,) *Vulva*—c. *Auris interna, Cochlea*—c. *Cerebri, Infundibulum of the brain*—c. *Genu, Patella*—c. *Inferior, Turbinate bone, (inferior)*—c. *Labyrinthi, Cochlea*—c. *Morgagniana, Turbinate bone, (superior)*—c. *Narium superior, Turbinate bone, (superior.)*

CONCHE, see Ostrea—c. *Narium, Turbinate bones*—c. *Veneris, Turbinate bones.*

CONCHA'RUM ANTIFEBRI'LE. A febrifuge and sudorific preparation in Bates's *Pharmacopœia*. It was composed of vinegar, mussel-shells (*conchas*), and water of *Carduus benedictus*.

CONCHOHELIX. A small, fleshy fascio-

lus, attached to the concha of the ear and helix. It is also called the *small muscle of the helix*.

CON'CHUS, *Concha*, (κογχος, κογχη.) The cranium. In the plural, it means the orbital cavities.

CONCHUS OCULI, Orbit.

CONCHYLIA, (from *concha*, 'a shell,') Turbinate bones.

CONCIDENTIA, (*con*, and *cadere*, 'to fall,') Collapse.

CONCILIA CORPORALIA, (pl. of *concilium*, 'a union,') Coition.

CONCIL'IUM. A milky plant, referred to by Pliny as aphrodisiac and antipthisical. Adanson considers it to be a *Campanula*.

CONCOC'TED, *Concoc'tus*, *Matura'tus*, *Pepei'rus*; (*con*, and *coquere*, *coctum*, 'to boil,') Brought to maturity; ripe; concocted; digested.

CONCOCTIO, Coction—c. *Tarda*, *Dyspepsia*.

CONCOCTION, Coction.

CONCOMBRE ORDINAIRE, *Cucumis sativus*—c. *Sauvage*, *Momordica elaterium*.

CONCOMITANS, Concomitant.

CONCOM'ITANT, *Concom'itans*, (*con*, and *comitare*, (itself from *comire*—*cum* and *ire*,) 'to go with,') That which accompanies. A symptom which accompanies others.

CONCREMATIO, (*con*, and *cremare*, *crematum*, 'to burn,') Calcination.

CONCREMENTA ZOOHYLICA, see *Zoohy-lia*.

CONCREMENTUM, Concretion.

CONCRETIO, Adherence, see *Concretion*—c. *Palpebrarum cum bulbo oculi*, *Symblepharosis*.

CONCRE'TION, *Concre'tio*, *Concremen'tum*, [*concre'scere*, *concretum*, (*con*, and *cre'scere*,) 'to condense, thicken, become solid,'] *Pec'ia*, *Symp'ea'is*, (F.) *Conerétion*. The act of becoming thick or solid. It was once used synonymously with adhesion or growing together—as '*concretion of the toes*.' Most commonly, it is applied to extraneous and inorganic bodies, of a solid character, met with in different textures, after certain chronic inflammations; or which make their appearance in the joints or in the reservoirs for excremental fluids. *Concretion* is, therefore, frequently synonymous with *Calculus*, and is then rendered, in Latin, by the word *Concrementum*. But *Concretion* has a more extensive signification than *Calculus*; thus, accidental ossifications or deposits of phosphates of lime in certain organs, and especially in the liver and lungs, are properly called *osseous concretions*. They could not well be called *osseous calculi*.

CONCRETION, FIBRINOUS, SANGUINEOUS, POLY-PYFORM, or POLYPOUS, OF THE HEART, see *Polypus*—c. *Intestinalis*, *Calculus of the Stomach and Intestines*—c. *Ossiforme*, see *Osteoid*.

CONCRETIONES ALVINÆ, see *Calculi of the stomach and intestines*.

CONCUBITUS, (*con*, and *cubare*, 'to lie,') Coition—c. *Venerens*, Coition.

CONCUR'SUS, gen. *Concur'sus*, (*concurrere*, (*con*, and *currere*, *cursum*, 'to run,') 'to meet together,') *Syn'dromē*. The congeries or collection of symptoms, which constitute and distinguish a particular disease.

CONCUS'SION, *Commo'tio*, (*concute*, *concussum*, (*con*, and *quater*, 'to shake,') 'I shake together,') *Concus'sio*, *Thlasma Concus'sio*, *Commo'tio*, *Anas'ismus*, *Tina'ismus*, (F.) *Commo'tio*. In Surgery, it is used for the agitation often communicated to one organ by a fall upon another; as to the brain from a fall on the breech, &c.

In all severe injuries, in sudden encephalic hemorrhage, and in overwhelming emotions, a *commotion or shock* is felt to a greater or less extent in the nervous system, which requires the careful attention of the physician.

CONCUSSION OF THE BRAIN, *Commo'tio Cerebri*,

Apoplex'ia nervo'sa traumati'ca, *Encephalosis'mus*, (F.) *Commo'tio du Cerveau*, sometimes gives rise to alarming symptoms, even to abolition of the functions of the brain, yet without any sensible organic disease. Slight concussion of the brain, called *stunning*, consists in vertigo, tinnitus aurium, loss of memory, and stupefaction; all these being temporary. When more severe, there is instant loss of sensation and volition, vomiting, the patient being as if in a sound sleep, but there is no stertorous breathing. Pulse variable, generally more rapid and feeble than in compression; extremities cold. Little can be done here, till reaction has occurred: after this, the case must be treated according to general principles,—by bleeding, blisters, cold applied to the head, &c. After severe concussion, a patient, although apparently well, is not safe till some time after the accident.

CONDENSAN'TIA, (*con*, and *densare*, *densatum*, 'to thicken,') *Incassan'tia*, *Syncret'ica*. Medicines esteemed proper for inspissating the humours.

CONDENSATIO, Condensation. Stegnosis.

CONDENSA'TION, *Condensa'tio*, *Inspissa'tio*. Increase in density of the blood and other liquids, or of the solids, which enter into the composition of the human body.

CONDENSER, see *Alembic*.

CONDEN'SER, LIEB'IG'S. A distillatory arrangement, invented by Liebig, in which the tube conveying the vapour is made to pass through another tube, the calibre of which is such as to leave a space between the two, through which a stream of water may be made to run.

CONDIMENT, *Condimen'tum*, *Ar'tyma*, *Hedys'ma*, *Conditu'ra*, (*condire*, *conditum*, 'to season,') (F.) *Assaisonnement*. Some substances are called, at times, *aliments*, and at others, *condiments*, according as they constitute the basis or the accessory to any dish: such are cream, butter, mushrooms, olives, &c. Others are always *condiments*, as they are only used to improve the savour of food, and contain little or no nutritive matter. Such are pepper, salt, cinnamon, &c. Almost all condiments are possessed of stimulant properties.

CONDIT, Confectio.

CONDITUM. A pharmaceutical compound of wine, honey, and some aromatics, especially pepper. Also, a confection.

CONDITURA CADAVERUM, Embalming.

CON'DOM, (from its proposer, Dr. Condom,) *Armour*, *Posthocalyp'tron*, French letter, *Cythe'rean Shield*, (F.) *Baudruche*, *Redingote Anglaise*, *Gant des Dames*, *Calotte d'assurance*, *Peau divine*, *Chemisette*. The *intestinum cæcum* of a sheep, soaked for some hours in water, turned inside out, macerated again in weak, alkaline ley, changed every twelve hours, and scraped carefully to abstract the mucous membrane, leaving the peritoneal and muscular coats exposed to the vapour of burning brimstone, and afterwards washed with soap and water. It is then blown up, dried, cut to the length of seven or eight inches, and bordered at the open end with a riband. It is drawn over the penis prior to coition, to prevent venereal infection and pregnancy.

CONDUCTIO, (*con*, and *ducere*, *ductum*, 'to lead,') Tonic spasm, Convulsion.

CONDUCTION, VIBRATIONS OF, see *Sound*.

CONDUCTOR, *Direct'or*, *Itinerarium*, (F.) *Conducteur*. That which conducts. The *Conductor* was an instrument, formerly used in the high operation for the stone, for directing the forceps into the bladder.

CONDUIT, (*con*, and *ducere*, *ductum*, 'to lead,') Canal—c. *Auditif externe*, Auditory canal, external—c. *Auditif interne*, Auditory canal, internal c. *Auriculaire*, Auditory canal, external—c. *Cholédouque*, *Choledoch duct*—c. *Déférent*, De-

ferens.—*c. d'Eustache*, Eustachian tube.—*c. Gutturale de l'oreille*, Eustachian tube.—*c. Labyrinthique*, Auditory canal, internal.—*c. de Pecquet*, Thoracic duct.—*c. Pterygoïdien*, Pterygoid canal.—*c. de la Pudeur*, Vagina.—*c. Sousorbitaire*, Suborbital canal.—*c. Spermatique*, Deferens (vas).—*c. Thoracique*, Thoracic duct.—*c. Vidua*, Pterygoid canal.

CONDUITS ADIPEUX, Fatty canals.—*c. Aqueux*, see Aqueous.—*c. Acengles de l'urèthre*, see Cæcus.—*c. Dentaires*, Dental canals.—*c. Ejaculateurs*, Ejaculatory ducts.—*c. Lacrymaux*, Lacrymal ducts.—*c. Lactifères*, Lactiferous vessels.—*c. Nourriciers*, Canals, nutritive.—*c. Nutriciers*, Canals, nutritive.

CONDYS DISINFECT'ING LIQUID or **FLUID**. This is supposed to be a concentrated solution of the permanganate of potassa. It is a good antiseptic.

CONDYLARTHROSIS, (κονύλος, 'a condyle,' and αρθρον, 'a joint.') Articulation by condyles. An elongated head or condyle, received into an elliptical cavity.

CONDYLE, *Con'dylē*, *Con'dylus*, *Capitulum*, (κονύλος, 'a knot, eminence.') An articular eminence, round in one direction, flat in the other. A kind of process, met with more particularly in the ginglymoid joints,—such as the condyles of the occipital, inferior maxillary bone, &c. Some anatomists have applied the term, however, to eminences that are not articular,—as to the lateral tuberosities at the inferior extremity of the os humeri, and even to certain depressions,—as to the concave articular surfaces at the upper extremity of the tibia. Chaussier calls the transverse root of the zygomatic process *Condyle of the temporal bone*.

CONDYLI DIGITORUM MANÛS, Phalanges of the fingers.

CONDYLIEN, see Condyloid Foramina.

CONDYLIUS, Condyloid.

CONDYLOID, *Condyloi'deus*, (κονυλοειδής,) *Condyloides*, *Condyllus*, (condyle, and εἶδος, 'shape.') Having the shape of a condyle.

CONDYLOID FORAMINA, *Foram'ina Condyloi'dea*, (F.) *Trous condyloïdiens*, *Fosses condyloïdiennes*. They are distinguished into *anterior* and *posterior*. They are four in number, seated in the occipital bone;—two anterior, and two posterior to the *condyles or condyloid processes* of the same bone, and to depressions, which are called *Fossæ Condyloides*.

As the word *Condyloïdien* means, 'that which has the shape of a condyle,' it has been judiciously proposed by some French anatomists that *condylien* should be used in preference, in the cases of the foramina and fossæ.

CONDYLOIDIEN, see Condyloid Foramina.

CONDYLOMA, gen. *Condylo'matis*, (κονδυλωμα,) *Condylo'sis*, *Con'dylus*, *Verru'ca carno'sa*, (κονύλος, 'a knot,' 'an eminence,' and the termination *oma*.) A soft, fleshy excrescence, of an indolent character, which appears near the orifice of the genital organs and rectum, and occasionally on the fingers and toes. It is a consequence of the syphilitic virus. Such tumours are also called *Dermophyma venerea*.

CONDYLOSIS, Condylooma.

CONDYLUS, (κονύλος,) Condylooma.

CONFLOWER, *Rudbeckia laciniata*.—*c. Purple*, *Echinacea purpurea*.

CONFINE, see Conium.

CONESSI CORTEX, *Nerium antidysentericum*.

CONFEC'TIO, *Confection*, *Confit*, (*conficere*, *conficere*, 'to make,') 'I make up.' *Alig'ulus*, (F.) *Confit*, *Confiture*, *Condit*. In general, it means anything made up or preserved with sugar. In the Pharmacopœias of the United States, London, and Dublin, it includes the arti-

cles before called electuaries and conserves. *Confec'tio* or *Confectum* also means *Confectionery*, *Cupé'die*.

CONFECTIO ALKERMES, Alkermes.

CONFECTIO AMYGDALA'RUM, *C. Amygd'alæ*, *Conser'va amygdala'rum*, *Almond Confection*, *Almond Paste*, *Pa's'ta reg'ia seu Amygdal'na seu Emul'si'va*, *Buty'rum Amygdala'rum Dul'cium*, (F.) *Confection d'Amandes*. (Sweet almonds, 3viij; gum acacia, 3j; White sugar, 3iv. Blanch the almonds, and beat into a paste.) A good mode of keeping almonds in a state fit for making emulsions. In the British Pharmacopœia (1864) the ingredients are rubbed to a coarse powder, which is called *Pulvis Amygd'alæ compos'it'us*.

CONFECTIO ARCHIG'ENI, *C. Pauli'na*. (*Castor*, long pepper, black pepper, storax, galbanum, costus, opium, of each 3ss; saffron, 3ij; syrup of wormwood, 3ij.) It was much recommended as a stimulant in nervous affections.

CONFECTIO AROMAT'ICA, *C. Card'i'aca seu Raleigh'na*, *Electua'rium Aromat'icum*, *Aromatic Confection*, *Sir Wal'ter Raleigh's Cordial*. The following is the formula, (Ph. U. S., 1863): *Pulv. aromat. 3vss; croci*, in pulv. 3ss; *syrup. aurant. 3vj; Mel despumat. 3ij*. Rub the aromatic powder with the saffron; then add the syrup and honey, and beat together until thoroughly mixed. Dose, ʒj to 3j.

Under the name *Pulvis cre'tæ aromat'icus*, *Aromatic powder of chalk*, the British Pharmacopœia (1864) has a substitute for the *Confectio aromat'ica* (Ph. L.), which is made by mixing thoroughly prepared chalk, ʒij, with aromatic powder, ʒiij, and passing the powder through a sieve.

CONFECTIO AURANTIO'RUM, *C. Aurant'ii Cor'ticis*, (Ph. U. S.), *Conser'va Cor'ticum Aurantio'rum seu Cor'ticis Exterio'ris Aurant'ii Hispan'icis seu Flaved'inis Cor'ticum Aurantio'rum Hispan'sium seu Aurant'ii seu Cit'ri Aurant'ii*, *Confection of the Orange*, *Con'serve of Orange Peel*. (Yellow part of the peel of the orange, ʒij; rectified sugar, ʒiij; beat into a conserve.) It is stomachic, and an agreeable vehicle, corrigent, and adjuvant for other remedies. Dose, 3ss.

CONFECTIO CARDIACA, *Confectio aromatica*.

CONFECTIO CAS'SLE, *Conser'va cas'siæ*, *Electua'rium Cas'siæ Fis'tulæ seu Cas'siæ seu Cas'siæ seu laxativum seu Cas'siæ tamarindatum seu lenitivum seu Cas'siæ*, *Diaca'siæ cum Man'nâ*, *Cas'siæ Confection*. (*Cassia pulp*, ʒss; manna, 3ij; tamarind pulp, 3j; syrup of roses, f3viii.) Bruise the manna, dissolve by heat, mix the pulp, and evaporate.—Ph. L.) It is gently laxative. Dose, 3ss.

CONFECTIO CAT'ECHU COMPOS'ITA, *Electua'rium Cat'echu*, *Compound Confection of Cat'echu*, (*Pulv. catechu comp. 3v* (avoidupois); *Syrup. f3v*, mix. (Ph. D.) The Edinburgh confection was formed of catechu and kino, each 3iv; cinnamon and nutmeg, each 3j; opium, diffused in a little sherry, 3iss; *Syrup of Red Roses*, of the consistence of honey, Oiss (*Imp. meas.*): mixed and thoroughly beaten together. A grain of opium is contained in about 200 grains of the mass. Dose, as an astringent, 3ss to 3j.

CONFECTIO DAMOCRATIS, *Mithridate*.

CONFECTIO HAMEC—so called from an Arabian physician—was composed of the bark of the yellow myrobalans, black myrobalans, violets, pulp of colocynth, polypodium of the oak, leaves of wormwood, thyme, aniseed, fennel, red roses, pulps of prunes, raisins, sugar, honey, senna, rhubarb, &c. It was used as a purgative in glysters.

CONFECTIO HYACIN'THI, *Hyacin'thi Confection*, *Electua'rium seu Confectio de Cro'co emendat'a olim dict'a de hyacin'this*, (Ph. P.) A tonic and slightly astringent confection, composed, according to Baumé, of terra sigillata, crabs' stones, cinnamon, leaves of the dittany of Crete, myrrh, saf-

iron, syrup of lemon, camphor, Narbonne honey, oil of lemon, &c.: and formerly the hyacinth was added, whence its name.

CONFECTION O'PII, *C. opia'ta*, *Electua'rium Opia'tum* seu *O'pii*, *Philo'nium Londinen'se* seu *Roma'nium*, *Theri'aca Edinen'sis*, *Electua'rium Theba'icum*, *O'pium Confection*, (*Opium*, in powder, $\mathfrak{z}$ ivss; *Pulv. aromat.* $\mathfrak{z}$ vj; *Mel despumat.* $\mathfrak{z}$ xiv; mix thoroughly. One grain of opium in 36:—Ph. U. S.) It is narcotic and stimulant. Dose, gr. x to $\mathfrak{z}$ j.

CONFECTION EX CAPITIBUS PAPAVERIS, *Diacodium*—*c. Paulina*, *Confectio Archigeni*.

CONFECTION PIP'ERIS, *C. Piperis ni'gri*, *Electua'rium Pip'eris*, *Ward's Paste for Fis'tula*. (*Black pepper*, *elecampane root*, each $\mathfrak{h}$ j; *fennel seed*, $\mathfrak{h}$ ij; *honey and sugar*, each $\mathfrak{h}$ ij. (Ph. U. S.) Dose, size of a nutmeg, three or four times a day.

CONFECTION RALEIGHANA, *Confectio aromatica*.

CONFECTION ROS'Æ, *C. Ros'æ Gal'licæ* seu *Ros'æ rub'ræ*, *Conser'va Ros'æ* seu *Ros'æ Gal'licæ* seu *Flo'rum Rosa'rum Rubra'rum* seu *Flo'rum Ros'æ Rub'ræ*, *Rhodosa'charum*, *Sac'charum Rosa'ceum*, *Confection of the Red Rose*. (*Red roses*, in powder, $\mathfrak{z}$ iv; *Sugar*, in powder, $\mathfrak{z}$ xxx; *Clarified honey*, $\mathfrak{z}$ vj; *Rose water*, $\mathfrak{f}$ zviij. Rub the roses with the rose water at a boiling heat; then add gradually the sugar and honey, and beat until thoroughly mixed.—Ph. U. S.) It is astringent, and chiefly used as a vehicle for other remedies.

CONFECTION ROS'Æ CANI'NÆ, *Conser'va Ros'æ Cani'næ* seu *Ros'æ Fruct'us* seu *Cynos'bati* seu *Cynos'r'hodi*, *Rub Cynos'batos*, *Confection or Con'serve of Dog Rose*. (*Pulp of dog rose*, $\mathfrak{h}$ j; *sugar*, $\mathfrak{z}$ xx. Incorporate. Ph. L.) It is chiefly used as a vehicle for other remedies.

CONFECTION RU'TÆ, *Confection of Rue*. (*Rue leaves*, dried, *carraway seeds*, *bay berries*, of each $\mathfrak{z}$ iss; *sagapenum*, $\mathfrak{z}$ iv; *black pepper*, $\mathfrak{z}$ ij; *honey*, $\mathfrak{z}$ xvj. Mix.—Ph. L.) It is given in clysters, as an antispasmodic and carminative.

CONFECTION DE SAN'TALIS, *Confection of Sand'ers*, (*F.*) *Confection de Sandaux*. (*Sandal wood*, *red coral*, *bole armeniac*, *terra sigillata*, *kermes berries*, *tormentil root*, *dittany*, *saffron*, *myrtle*, *red roses*, *calcined hartshorn*, and *cloves*.) It was formerly used as an astringent.

CONFECTION SCAMMO'NI, *Electua'rium Scammo'ni* seu *e Scammo'nio*, *Caryocost'inum*, *Confection of Scammony*. (*Pounded scammony*, $\mathfrak{z}$ iss; *bruised cloves*, *powdered ginger*, $\mathfrak{a}$ ã $\mathfrak{z}$ vj; *oil of caraway*, $\mathfrak{z}$ ss; *syrup of roses*, q. s.—Ph. L.) A stimulating cathartic. Dose, $\mathfrak{z}$ ss to $\mathfrak{z}$ j.

CONFECTION SEN'NÆ, *Electua'rium Cas'siæ Sen'næ*, *E. Sen'næ cum Pul'pis* seu *Sen'næ compos'itum* seu *ape'riens* seu *cathol'icum commu'nè* seu *diap'urum* seu *eccepro'ticum* seu *Sen'næ seu e Sen'nâ seu leni'tivum*, *Benedic'tum Laxati'vum*, *Confection of Sen'na*, *Len'itive Elec'tuary*, &c. (*Senna leaves*, $\mathfrak{z}$ vij; *figs*, $\mathfrak{h}$ j; *tamarind pulp*, *cassia pulp*, *pulp of French prunes*, each $\mathfrak{h}$ ss; *coriander seed*, $\mathfrak{z}$ iv; *liquorice root*, $\mathfrak{z}$ ij; *sugar*, $\mathfrak{h}$ iss; *water*, *Qiv*. Rub the senna and coriander together; separate 10 ounces of the powder with a sieve; boil the residue with the figs and liquorice root in the water to one-half; press out the liquor and strain. Evaporate the liquor by means of a water bath to a pint and a half; add the sugar, and form a syrup. Rub the pulps with the syrup, gradually added; throw in the sifted powder, and beat till thoroughly mixed.—Ph. U. S. 1861.) The formula in the Ph. U. S. 1863 differs slightly from this. It is a laxative, and is used in habitual constipation, and in constipation during pregnancy. Dose, $\mathfrak{z}$ ss to $\mathfrak{z}$ ss.

CONFECTION SULPH'URIS, *Confection of Sulphur*. (*Sulphur. sublim.* $\mathfrak{z}$ ij; *Potass. Bitart.* $\mathfrak{z}$ j; *Mel. despumat.* $\mathfrak{z}$ j; *Syrup. Zingib.*, *Syrup. Croci*, $\mathfrak{a}$ ã

$\mathfrak{f}$ zss. M.—Ph. D.) As a laxative and alterative. Dose, $\mathfrak{z}$ j to $\mathfrak{z}$ ij.

CONFECTION TEREBINTHINÆ, *Confection of Tur'pentine*. (*Ol. Tereb.* $\mathfrak{f}$ zj; *Pulv. Glycyrrhiz.* $\mathfrak{z}$ j; *Mel. despumat.* $\mathfrak{z}$ ij.—Ph. D.) An agreeable mode of administering turpentine. Dose, a scruple to a drachm.

CONFECTION DE THU'RÈ, *Frank'incense Confection*, (*F.*) *Confection d'Encens*. A compound of *coriander seeds*, *nutmeg*, *thus*, *liquorice*, *mastic*, *cubebs*, *prepared hartshorn*, *conserve of red roses*, *sugar*, &c.

CONFECTION, *Confectio*—*c. d'Amandes*, *Confection*, *almond*—*c. of Catechu*, compound, *Confectio catechu compositum*—*c. d'Encens*, *Confectio de Thure*—*c. Frankincense*, *Confectio de Thure*—*c. of the Orange*, *Confectio aurantiorum*—*c. of the Red rose*, *Confectio ros'æ Gallicæ*—*c. of Rue*, *Confectio rutæ*—*c. de Sandaux*, *Confectio de santalis*—*c. of the Sanders*, *Confectio de santalis*—*c. of Sulphur*, *Confectio sulphuris*—*c. of Turpentine*, *Confectio terebinthina*.

CONFECTIONARIUS, *Apothecary*.

CONFECTUM, *Confection*.

CONFERTA HELMINTHOCORTOS, (*conferere*, 'to knit together.') *Corallina Corsicana*.

CONFER'VA RIVA'LIS. This species of *River Weed* has been recommended in cases of spasmodic asthma, phthisis, &c.

CONFINEMENT, *Parturient state*.

CONFIRMANTIA, (*con*, and *firmare*, 'to strengthen.') *Tonics*.

CONFIRMED, *Consummatus*.

CONFIT, *Confectio*.

CONFUTURE, *Confectio*.

CONFLUENT, *Con'fluens*, (*con*, and *fluere*, *fluxum*, 'to flow.') 'Running together.' An epithet for certain exanthematous affections, in which the pimples, pustules, &c., run together. It is particularly applied to small-pox, so circumstanced. Some authors have called scarlatina or scarlet fever *Con'fluent Measles*, *Morbil'li Con'fluen'tes*.

CONFLUENT DES SINUS, *Torcular Heropbili*.

CONFLUENTIA, *Confederatio*. A term employed by Paracelsus to express the concordance between a disease and its remedies.

CONFLUXIO, *Sympathy*.

CONFEDERATIO, (*con*, and *foedus*, gen. *foederis*, 'a treaty.') *Confluentia*.

CONFORMATIO, *Conformare*, (*con*, and *formare*, *formatum*, 'to form,' 'to arrange,' 'to dispose.') *Conformation*, *Structure*.

CONFORMATION, *Conformatio*, *Diap'lasis*, *Diaplas'mus*, *Struc'ture*. The natural disposition or arrangement of the body.

Fault'y conformation, (*F.*) *Vice de conformation*, is vice of original formation; existing, of course, from birth. In French surgery, *Conformation* is used synonymously with *Cooptation*, and both it, *Diap'lasia*, and *Anap'lasia* mean, also, restoration to the original form—as in fractures, &c.

CONFORTANTIA, (*con*, and *fortis*, 'strong.') *Tonics*.

CONFORTATIVA, *Tonics*.

CONFORTER, (*F.*) *Confirma'rè*, *Conforta'rè*, *Corroborat'rè*. To make stronger—to give energy. *Conforter l'estomac*, 'to strengthen the stomach.'

CONFRICTIO, *Confricatio*, (*confricare*, (*con*, and *fricare*, *fricatum*), 'to rub.') The action of reducing a friable substance to powder, by rubbing it between the fingers; and of expressing the juice of a plant with the hand.

CONFRICTRIX, A female who practises masturbation.

CONFUSÆ FEBRES. Intermittents, whose paroxysms are irregular and confused.

CONFUSIO, (*confundo*, (*con* and *fundere*, *fusum*, 'to pour,') 'I mix together,') *Syn'chysis*. A

disease of the eye, which consists in the mixture of the humours. In modern times, Synchysis has been applied to a morbid state of the vitreous body, in which it is reduced to a diffuent condition. Occasionally, sparkling bodies are perceptible, which seem to be seated behind the crystalline, in the vitreous humour: hence, the disease has been called *sparkling synchysis* or *spintheropia*, (from *σπῆρ*, a spark, and *ωπ*, 'the eye;') (F.) *Spinthéropie*, *Scintillement de l'œil*. *Confusio* has, also, been used synonymously with *Complexio*.

CONFUSIONES ANIMI, 'Confusions of the mind,' see Affections of the mind.

CONGEE DISCHARGES, Rice-water discharges.

CONGELANTIA, (*congelato*, (*con*, and *gelare*, *gelatum*), 'I congeal,' 'I freeze.') Congelativa.

CONGELATIO, Catalepsy, Congelation.

CONGELATION, *Congelatio*, *Conglacatio*, *Gelatio*. The action of congealing, of passing to the solid state by the abstraction of heat; as *congelation of water*, *mercury*, &c. The term had once other acceptations. 1. It was synonymous with *concretion*. 2. With *coagulation*, in which sense it is still occasionally employed. 3. The ancients called all diseases, produced by cold, *congelations*, as well as those in which there was a kind of stupor or torpor—particularly catalepsy. Also, *Frostbite*, (Prov.) *Keenbite*.

Congelation of a part, by the application of powdered ice or of a freezing mixture, in a bladder or gauze bag, has been employed to induce anæsthesia in the lesser surgical operations. It has also been used in external inflammation.

CONGELATIVA MEDICAMENTA, *Conglutinantia*, *Congelan'tia*. Medicines, considered capable of uniting or consolidating wounds, &c.

CON'GENER, gen. *Congen'eri*, *Congen'erous*, (*con*, and *genus*, gen. *generis*, 'kind.') (F.) *Con-génère*. Of the same kind or species. Resembling each other in some manner. When applied to muscles, it means, that they concur in the same action; in opposition to the word *antagonist*, or that which acts in an opposite direction.

In France *Congénères* is applied to those who join in the dissection of the same subject.

CONGENIAL, Congenital.

CONGENIALIS, Congenital.

CONGENITÆ NOTE, Nævus.

CONGENITAL, *Con'genite*, *Congenial'is*, *Con-gen'itus*, *Syngen'icus*, (*con*, and *genitus*, 'begotten.') (F.) *Congénital* ou *Congénital*. Diseases which infants have at birth: hence, *Congenital affections* are those that depend on faulty conformation; as *congenital hernia*, *congenital cataract*, &c. See *Connate*.

CONGESTED, Hyperæmic.

CONGESTIO, (*congerere*, (*con*, and *gerere*, *gestum*, 'to carry,') 'to amass,' 'accumulate,' &c.) Congestion—c. Abdominalis, Cœliæmia—c. Pectoris, Stethæmia—c. Pulmonum, Stethæmia—c. Sanguinis, Congestion.

CONGESTION, *Congest'io*, *Rho'pē*, *Symph'ora*, *Hæmatepago'gē*, *Hæmatosymphore'sis*, *Hæmatosy-nago'gē*, *Hæmorme'sis*, *Symphore'ma*, *Symphore'sis*, *Synathrois'mus*, *Synathro'is'is*, *San'guinis Conges-tio*, *Engorge'ment*. Accumulation of blood in an organ. It is an important symptom in febrile and other disorders. It may arise either from an extraordinary flow of blood by the arteries, or from a difficulty in the return of blood to the heart by the veins. More often, perhaps, it is owing to the latter cause, and is termed *venous congestion*, *stasis* or *stagnation*—being not unusually attended with symptoms of oppression and collapse. See *Hyperæmia*.

CONGESTION OF THE ABDOMEN, Cœliæmia—c. of the Brain, Cephalæmia—c. *Cérébrale*, Cephalæmia—c. *du Cerveau*, Cephalæmia—c. *Hyposta-*

tique, see *Hypostatic*—c. of the Lungs, Stethæmia—c. *des Poumons*, Stethæmia—c. *Sanguine rachidienne*, Hypermyelohæmia.

CONGESTIVE, *Congest'ivus*, (F.) *Congestif*. Belonging or relating to, or affected with, congestion—as 'congestive fever.'

CONGESTIVE FEVER, see *Fever*, congestive.

CONGLACIATIO, (*con*, and *glacies*, 'ice.') Congelation.

CONGLO'BATE, or CON'GLOBATE, *Conglo-ba'tus*, (*conglobare*, (*con*, and *globus*, 'a ball,') 'to collect,' 'to gather into a ball.') (F.) *Conglobé*.

CONGLOBATE GLAND, *Glan'dula congloba'ta*, *Hy'draden*, *Glo'bate gland*, *Lymphat'ic gan'glion*, (F.) *Glande Conglobée*, *Ganglion lymphatique*. A round body, formed of lymphatic vessels, connected together by cellular structure, but having neither a cavity nor excretory duct. The mesenteric, inguinal and axillary glands are of this class.

CONGLOBE, Conglobate.

CONGLOMERATE, *Conglomera'tus*, (*con*, and *glomerare*, *glomeratum*, 'to gather in a heap.') *Glom'erate*, *Glomera'tus*, (F.) *Congloméré*. Heaped together.

CONGLOMERATE GLANDS, *Glan'dulæ conglome-ra'tæ* are those whose lobules are united under the same membrane; as the liver, kidney, testicle, mamma, &c.

CONGLOMERATIO INTESTINORUM, Epiploce intestinalis.

CONGLUTINANTIA, (*con*, and *gluten*, gen. *glutinis*, 'glue,') Congelativa.

CONGRES (F.), *Con'gress*, *Congres'sus*, (*con-gredi*, *congressus*, (*con*, and *gradi*, 'to go,') 'to go together,') This term, which has often been used synonymously with *Coition*, means, also, the ocular proof, formerly ordered by judicial authority, in the presence of surgeons and matrons, to test the impotence or capabilities of parties;—a most unsatisfactory and indecent exhibition. It was forbidden by the Parliament of Paris in the year 1667.

CONGRESSUS, Coition.

CONI VASCULO'SI, *Con'pora pyramida'llia test'is*. Conical bundles, formed by the vasa efferentia of the testis; having their base towards the epididymis, into the tube of which they enter.

CONIA, (*κονία*.) A wine, prepared by fermenting the must of the grape on tar previously washed in sea-water. See, also, *Conium*, and *Lixivium*.

CONIASIS, (*κονιασις*, 'art of plastering,' from *κονία*, 'plaster or stucco,') Incrustation.

CONICINE, see *Conium*.

CONII FOLIA, see *Conium*—c. *Fructus*, see *Conium*—c. *Semen*, see *Conium*.

CONIINE, see *Conium*.

CONINE, see *Conium*.

CONIOSTOSIS, (from *conis*.) Pulverization.

CONIS, (*κονίς*, 'dust,') Pulvis.

CONISTERIUM, (*κονιστήριον*, from *conis*.) Apodyterium.

CONI'UM, (*κωνιον*, *κωνιον*, *κωνιον*.) *C. macula'tum*, *Abi'otus*, *Corian'drum macula'tum*, *Cien'ta* major seu *macula'ta* seu *Stoerkii* seu *terres'tris* seu major *fœtida* seu *vulgar'is*, *Common* or *Poison Hem'lock*, *Hemlock*, *Poison pars'ley*, *Spotted par-ley*, (Sc.) *Humloik*, (F.) *Ciguë ordinaire*, *C. grande* ou *officinale*. Ord. Umbelliferae. Sex. Syst. Pentandria Digynia. The leaves, *Coni'i Fo'lia*, *Coni'um*, (Ph. U. S., 1863,) and seeds, *Coni'i Semen*, are narcotic and poisonous in an overdose. Light destroys the virtues of the leaves; and, therefore, the powder should be kept in opaque bottles, well corked. It has been used as a palliative in cancer and other painful affections; but is not equal to opium. Externally, it has been applied in fomentation to cancerous and serofulous ulcers. Dose, gr. ij to x.

Its active alkaloid principle is *Conia*, *Conine*, *Coniine*, *Conicine*, *Cicutine*. It, or its

salts, have been given as sedatives to the nervous centres in various neuralgic and spasmodic diseases. It is a most energetic poison.

The ripe fruit dried—*Coni'i fructus*, *Hemlock fruit*—is official in the British Pharmacopœia, (1864.)

Davidson's Remedy for Can'cer, is said to consist of powdered hemlock and arsenious acid.

CONIUM MOSCHA'TUM, *Aracacha*. A very agreeable and nutritive kind of tuberous vegetable, in flavour not unlike celery, which grows on the coast of Peru, but is more abundant on the projecting ridges of the Cordilleras, and on the eastern declivity of the Andes. It is cooked by being either simply boiled in water, or made into a kind of soup.

CONJUNCTIVAL, Conjunctival.

CONJUNCTIVE, Conjunctiva.

CONJUNCTIVITE, see Ophthalmia—c. *Blennorrhagique*, see Ophthalmia.

CONJUGAISON, Conjugation.

CONJUGA'TION, *Conjugatio*, (*conjugare*, *conjugatum*, (con, and jugum, 'a yoke,') 'to yoke together,') (F.) *Conjugaison*. Assemblage, union, — *Conjugium*.

CONJUGATIO'NIS FORAM'INA, (F.) *Trous de conjugaison*. The apertures at the sides of the spine, formed by the union of the notches of the vertebrae. They give passage to the nerves of the spinal marrow, and to the vessels which enter or issue from the vertebral canal.

CONJUGIUM, Coition, Conjugation.

CONJUNC'TI (MOR'BI), (*conjungere*, (con, and *ungere*, *junctum*,) 'to join together,') Diseases joined together. Authors have distinguished two kinds of these: one, in which the diseases go on simultaneously — *morbi connex'i*; the other, in which they succeed each other — *morbi consequentes*.

CONJUNCTIO, Articulation.

CONJUNCTIVA, CUTICULAR, Xerophthalmia — c. *Granular*, *Trachoma*.

CONJUNCTI'VA MEMBRA'NA, *Ophthalmodesmum*, *Syn'chion*, *Circumcinalia*, *Tu'nica agna'ta seu adna'ta seu conjuncti'va*, *Tun'da oculi*, (F.) *Conjunctive*, *Membrane adnæ*. A mucous membrane, so called because it unites the globe of the eye with the eyelids. It covers the anterior surface of the eye, the inner surface of the eyelids, and the *caruncula lacrymalis*. It possesses great general sensibility, communicated to it by the fifth pair.

CONJUNCTIVAL, *Conjunctivalis*, (F.) *Conjunctival*. Appertaining or relating to the conjunctiva.

CONJUNCTIVITIS, (*conjunctiva*, and *itis*,) Ophthalmia — c. *Egyptiaca*, Ophthalmia, purulent — c. *Blennorrhagica*, see Ophthalmia — c. *Catarrhalis*, Ophthalmia, catarrhal — c. *Gonorrhœica*, see Ophthalmia — c. *Puro-mucosa catarrhalis*, Ophthalmia, catarrhal — c. *Puro-mucosa contagiosa seu Egyptiaca*, Ophthalmia, purulent.

CONJUGENS, Communicians.

CON'NATE, *Conna'tus*, (F.) *Conné*, (con, and *natus*, 'born') as:—

CONNATE DISEASES, (F.) *Maladies connées*, *Morbi connati*, such as an individual is born with: — *connate* having the same signification as *congenital*. A difference has been made by some, however; those diseases or conditions, which are dependent upon original conformation, being called *congenital*; — whilst the diseases or affections that may have supervened during gestation or delivery, are termed *connate*.

CONNÉ, Connate.

CONNECT'ICUT, MINERAL WATERS OF. There is a mineral spring at Stafford, in this State, twenty-four miles from Hartford, which has obtained more celebrity than any one in New England. Its principal ingredients are iron and car-

bonic acid. It, consequently, belongs to the class of acidulous chalybeates. There are other springs in the State, of which, however, little that is accurate is known.

CONNECTIVE TISSUE, (con, and *nectere*, 'to bind,') Cellular tissue.

CONNERVATIO, (con, and *nervus*, 'a sinew,') Syndesmosis.

CONNEXIO OSSIUM CARNOSA, (con, and *nectere* *nezum*, 'to bind,') *Syssarcosis* — c. *Cartilaginea*, *Synchondrosis* — c. *Ligamentosa*, *Syndesmosis*.

CONNUTRI'TUS, *Syn'trophus*, (con, and *nutrior*, *nutritus*, 'to be nourished,') A disease is so called which has grown up, as it were, with an individual, or has been connate with him.

CO'NOID, *Conoi'deus*, (*κωνοειδής*,) *Conoi'des*, (*κωνος*, 'a cone,' and *eidos*, 'shape,') (F.) *Conoïde*. Resembling a cone.

CONOID LIG'AMENT, *Ligamentum conoi'des*. A ligament, passing from the coracoid process to the scapula, and forming part of the coraco-clavicular ligament of some anatomists.

CONOIDAL SUBSTANCE OF THE KIDNEY, see Kidney.

CONOIDES CORPUS, ('conoid body,') Pineal gland.

CONOPHTHALMIA, (*κωνος*, 'a cone,' and 'ophthalmia,') *Staphyloma corneæ*.

CONQUASSANT (F.), *Conquassans*, (con, and *quassare*, 'to shake,') *Douleurs conquassantes* are the pains of parturition, at the time of their greatest intensity, when the head is engaged in the pelvis.

CONQUASSA'TION, *Conquassa'tio*, *Quassa'tio*, *Quassatu'ra*. A pharmaceutical operation, which consists in dividing, with a pestle, fresh vegetables, fruits, &c. See Confrication.

CONQUASSATIONES ANIMI, Affections of the mind.

CONQUE, Conecha.

CON'SCIOUSNESS, DOUBLE, (con, and *scio*, 'to know,') A somnambulistic condition, in which the individual leads, as it were, two lives, recollecting in each condition what occurred in previous conditions of the same character, but knowing nothing of the occurrences of the other. See Duality of the Mind.

CONSEC'UTIVE, *Consecuti'vus*, (con, and *sequor*, *secutus*, 'to follow,') Following in order.

CONSECUTIVE PHENOM'ENA OR SYMP'TOMS, (F.) *Phénomènes ou accidens consécutifs*, are such as appear after the cessation of a disease, or according to others, during its decline; but without having any direct connexion with it.

CONSENSUAL, (con, and *sensire*, *sensum*, 'to feel,') see Instinctive.

CONSENSUS, Consent of parts, Sympathy.

CONSEN'SUS OCULO'RUM. The intimate association between the two eyes, as exemplified in their consensual action in health, and often in disease.

CONSENT' OF PARTS, *Consen'sus*, *Consen'sus partium*, *Sympathi'a*, (F.) *Consentement des Parties*. That relation of the different parts of the body with each other which is more commonly denominated sympathy.

CONSE'NTEMENT DES PARTIES, Consent of parts.

CONSER'VA, *Con'serve*, (F.) *Conserve*, *Electuaire simple*, *Saccharol. mou*. A pharmaceutical preparation, composed of a vegetable substance and a sufficient quantity of sugar beaten into a uniform mass. See Confection.

CONSERVA ABSIN'THII, *C. absinthii maritimi*, *Conserve of Wormwood*. (Leaves *thj*, Sugar *℥iij*.) It has been employed as a tonic, stomachic, and vermifuge.

CONSERVA AMYGDALARUM, *Confectio amygdalarum.*

CONSERVA ANGELICÆ (Ph. P.), *Conserve d'Angelique, C. of Angelica.* (Pulp of root 250 parts; white sugar, boiled in a decoction of the root, and reduced to the consistence of a solid electuary, 1000 parts.) It is tonic, aromatic, and stomachic.

CONSERVA DE APIO GRAVEOLENTĒ (Ph. P.), *Conserve d'Ache, Conserve of Smallage.* Prepared like the preceding. Reputed to have the same properties.

CONSERVA A'RĪ, *Conserve of Arum.* (Fresh root, lbss, Sugar lbss.) Esteemed to be diuretic and stimulant.

CONSERVA AURANTII, *Confectio aurantiorum*—c. Cassie, *Confectio cassiæ*—c. Citri aurantii, *Confectio aurantiorum.*

CONSERVA COCHLEA'RĪE HORTEN'SIS, *Conserve of Lemon Scurvy Grass.* (Leaves lbj, Sugar lbij.) Reputed to be stimulant and antiscorbutic.

CONSERVA CORTICIS EXTERIORIS AURANTII HISPALENSIS, *Confectio aurantiorum*—c. Flavedinis corticis Aurantiorum Hispalensium, *Confectio aurantiorum*—c. Corticium Aurantiorum, *Confectio aurantiorum*—c. Cynorrhodi, *Confectio rosæ caninæ*—c. Cynosbati, *Confectio rosæ caninæ.*

CONSERVA LU'JULÆ, *C. Folio'rum lu'julæ, Conserve of Woodsorrel.* (Leaves lbj, Sugar lbij.) Gratefully acid and refrigerant.

CONSERVA MENTHÆ, *C. Men'thæ folio'rum, C. Men'thæ sativæ, Conserve of Mint.* (Mint leaves lbj, Sugar lbij.) Stomachic in nausea and vomiting.

CONSERVA PRU'NI SYLVES'TRIS, *C. Pru'næ sylvestris, Pul'pa pruno'rum sylvestrium condita, Conserve of Sloes.* (Pulp 1 part, Sugar 3 parts.) Possessed of astringent properties.

CONSERVA RO'SÆ, *Confectio rosæ Gallicæ*—c. Rosæ caninæ, *Confectio rosæ caninæ*—c. Florum Rosarum rubrarum, *Confectio rosæ Gallicæ*—c. Rosæ fructūs, *Confectio rosæ caninæ*—c. Rosæ Gallicæ, *Confectio rosæ Gallicæ.*

CONSERVA SCI'LLÆ, *Conserve of Squill.* (Fresh squills 3i, Sugar 3x.) Possesses the diuretic and other virtues of the squill.

CONSERVATION, *Conservatio, Phylaxis*, (*conservare*, (con, and *servare*, *servatum*), 'to preserve;') (F.) *Conservation, Asservation.* The art of preserving any object of pharmacy, any remedial agent, &c., from decay.

CONSERVE, *Conserve*, see *Confectio*—c. d'Ache, *Conserve de Apio graveolente*—c. of Aloes, *Conserve pruni sylvestris*—c. of Lemon scurvy grass, *Conserve cochleariæ hortensis*—c. of Mint, *Conserve menthæ*—c. of Orange, *Confectio aurantiorum*—c. of Roses (red), *Confectio rosæ gallicæ*—c. of Smallage, *Conserve de apio graveolente*—c. of Woodsorrel, *Conserve lujulæ*—c. of Wormwood, *Conserve absinthii.*

CONSERVES, Spectacles.

CONSIDENTĪA, (*considerare*, (con and *sedo*), 'to settle;') This word has two acceptations. 1. It is synonymous with *Apocatastasis*, and means restoration or cessation, and the subsiding of a humour; and 2. It signifies contraction of any cavity or canal.—See *Synezeisis*.

CONSISTENTĪA, (*consistere*, (con, and *sistere*), 'to stand still;') A term employed in two senses. 1. When joined to the word *Morbi* or *Ætiæ*, it expresses the acme of a disease, or the age at which the constitution has acquired its full strength. 2. By *Consistentia humoris* is meant the density of a humour.

CONSOLIDA MAJOR, *Symphytum*—c. Media, *Ajuga*, *Chrysanthemum leucanthemum*—c. Minor, *Prunella*—c. Regalis, *Delphinium consolida*—c. Rubra, *Tormentil*—c. Saracenica, *Solidago virgaurea.*

CONSOLIDANTĪA, *Consolidativa Medicamenta*, (con, and *solidare*, 'to make solid;') Substances formerly given for the purpose of consolidating wounds, or strengthening cicatrices.

CONSOLIDATIVA, *Consolidantia.*

CONSUMMĒ (F.), *Consumma'tum*, (*consummo*, (con, and *summa*, 'a sum;') 'I add together;') *Zomos.* Soup strongly charged with gelatin, and consequently very nutritious, although not proportionably easy of digestion.

CONSUMPTION, *Consumption.*

CONSONANCE, (*consono*, 'I sound together;') (con, and *sono*, 'I sound;') see *Sound*.

CONSONANT, *Consonans*, (same etymon.) An alphabetic sound, which cannot be properly expressed, except when conjoined with a vowel. Physiologically, a breath, or sound produced in the larynx, which suffers more or less interruption in its passage through the vocal tube.

CONSORTIUM, (con, and *sors*, gen. *sortis*, 'lot;') *Sympathy.*

CONSOUE GRANDE, ([L.] *consolida*), *Symphytum.*

CONSOUND, MIDDLE, *Ajuga.*

CONSPERSIO, (con, and *spargere*, *sparsum*, 'to sprinkle;') *Catapasma.*

CONSPICILLA, (con, and *specio*, 'I look at;') *Spectacles.*

CONSPIRATIO, (con, and *spirare*, 'to breathe;') *Sympathy.*

CONSTELLATUM UNGUEN'TUM. An ointment composed of cleansed earthworms dried and pulverized; and of the fat of the bear or wild boar. It was formerly employed in toothache, and to hasten the cicatrization of wounds.

CONSTERNATIO, (*consternare*, 'to confuse;') *Stupor.*

CON'STIPATED, *Constipatus, Dyscœlius, Steganus*, (*constipare* (con, and *stipare*, *stipatum*) 'to cram close;') (Prov.) *Maw'bound*, (F.) *Constipé*. Affected with constipation; *Costive; Bound, Bellybound.*

CON'STIPATING, *Constipans, Binding*, (F.) *Reserrant.* Rendering the bowels less open; having the power to induce constipation.

CONSTIPATIO, *Constipation, Stegnosis.*

CONSTIPATION, *Constipatio, Obstipatio, Adstrictio, Arctatio, Obstipatio seu Retentio alvina, Ischocœlia, Alvus astricta seu adstricta seu tar'da seu du'ra seu segnis, Obstructio seu Suppressio alvi, O. Ductus alimentarii, O. intestinalis, Tor'por intestino'rum, Styp'sis, Constipatio alvi, Copros'tasis, Acop'ria, Acopro'sis, Coprop'ris'chesis, Ischocop'ria, Dyscœlia, Costiveness, Fœcal retention, Alvine obstruction, Stopping or Stoppage of the Bowels or Guts or Stomach*, (F.) *Échauffement, Ventre reserré.* A state of the bowels, in which the evacuations do not take place as frequently as usual; or are inordinately hard, and expelled with difficulty. It may be owing either to diminished action of the muscular coat of the intestines, or to diminished secretion from the mucous membrane, or to both. Cathartics will usually remove it; after which its exciting and predisponent causes must be inquired into and obviated, to render the cure permanent.

CONSTIPATUS, *Constipated.*

CONSTIPÉ, *Constipated.*

CONSTITUENS, *Vehicle*, see *Prescription*.

CONSTITUTIO, (*constituere*, *constitutum*, 'to set up;') from *con*, and *statuere*, itself from *stare*, *statum*, 'to stand;') *Constitution, Habit of body*—c. *Aeris*, *Constitution of the atmosphere*—c. *Epidemica*, *Constitution, epidemic*—c. *Nervosa*, *Nervous diathesis*.

CONSTITUTION, *Constitutio, Catâstasis, Stâtus.* A collection of several parts, forming a whole. In medicine, *Constitution* means the state of all the organs of the human body considered

in regard to their special and relative arrangement, order, or activity. A *good constitution* is one in which every organ is well developed, and endowed with due energy, so that all perform their functions with equal facility. Any want of equilibrium in their development and energy forms a difference in the constitution. We say that a man is of a *good or robust, a delicate or weak constitution*, when he is commonly healthy, or commonly labouring under, or unusually susceptible of, disease.

CONSTITUTION, ARTERIAL, *Plethora arteriosa*.

CONSTITUTION OF THE ATMOSPHERE, *Constitutio A'ëris*, (F.) *Constitution Atmosphérique*. The condition of the atmosphere, as regards dryness and humidity, temperature, heaviness, direction of the winds, &c., considered as respects its influence on the animal economy.

CONSTITUTION, EPIDEMIC, *Constitutio epidemica*, *Med'ical Constitution*, (F.) *Constitution médicale*, *C. épidémique*. The aggregate of meteorological conditions, during which diseases prevail epidemically.

CONSTITUTIONAL, (F.) *Constitutionnel*. Belonging to the constitution of an individual,—to his manner of being; as *constitutional phthisis*, *c. gout*, &c. By some, this epithet has been given to diseases, produced by the constitution of the atmosphere; but this acceptance is not common.

CONSTRICTEUR D'HERBINEAUX, see *Serre-nœud*—*c. du Vagin*, *Constrictores cunni*—*c. de la Vulve*, *Constrictores cunni*.

CONSTRIC'TIO, (*constringere*, *constrictum* (*con*, and *stringere*, *strictum*) 'to bind.')

Astriction, Systole.

CONSTRUCTIVA, *Styptics*.

CONSTRIC'TOR, gen. *Constrictor'is*. (F.) *Constricteur*. That which binds in a circular direction. A sphincter. Different muscles are so called.

CONSTRUCTOR ANI, Sphincter ani externus—*c. of Nuck*, Compressor of Nuck.

CONSTRICTORES ALARUM NASI, Depressor alæ nasi.

CONSTRICTORES CUN'NI, *C. Vag'inæ seu Vul'væ*, *Clitor'idia inferiora lat'i et pl'a'ni mus'culi*, *Sphinc'ter Vag'inæ*, (F.) *Constricteurs du vagin*, *C. de la Vulve*. Small muscles, which originate beneath the clitoris, descend along the sides of the vagina, and terminate by becoming confounded with the *transversus perinæi* and *external sphincter ani* muscles. Their use is to contract the entrance of the vagina.

CONSTRICTORES ISTHMI FAUCIUM, see *Glossostaphylinus*.

CONSTRICTORES LARYN'GIS, Lieutaud describes, under the name *Grand constricteur du Larynx*, the muscle *Cricocarytenoïdeus latera'lis* with the *Thyroarytenoïdeus*.

CONSTRICTORES NASI, Compressor naris.

CONSTRICTOR ŒSOPH'AGI, *Constrictor of the Œsoph'agus*, (F.) *Constricteur de l'Œsophage*, *Muscle œsophagien*. A fasciculus of fleshy, circular fibres, at the upper part of the œsophagus.

CONSTRICTORES ORIS, Orbicularis oris—*c. Palpebrarum*, Orbicularis palpebrarum.

CONSTRICTORES PHARYN'GIS, *Constrictors of the Pharynx*, *Sphincter Gu'lae*. Muscular expansions which assist in forming the parietes of the pharynx. Three of these are generally admitted. 1. The *Constrictor Pharyn'gis inf'erior*, *Cricopharynge'us* and *Thyro-pharynge'us*, (F.) *Cricothyro-pharyngien*. It is broad, very thin, quadrilateral, seated superficially, extending from the thyroid and cricoid cartilages as far as the middle of the pharynx, and uniting, on the median line, with that of the opposite side. 2. The *Constrictor Pharyn'gis Medi'us*, *Hyopharynge'us* and *Chondropharynge'us*, *Syndes'mopharyngeus*, (F.) *Hyo-glossobasi-pharyngien*, occupies the middle

part of the pharynx. It is triangular, and attached anteriorly to the great and little cornu of the os hyoides; to the stylohyoid ligament; and terminates, posteriorly, by joining its fellow of the opposite side. 3. The *Constrictor Pharyn'gis sup'erior*, *Ceph'alopharynge'us*, *Glossopharynge'us*, *Mylopharynge'us*, *Pter'ygopharynge'us*, (F.) *Pter'gyo-syndesmo-staphyli-pharyngien* is quadrilateral, and extends from the internal ala of the pterygoid process, from the inter-maxillary ligament, from the internal oblique line of the jaw, and from the base of the tongue, to the posterior part of the pharynx.

The constrictors of the pharynx contract it. They can likewise approximate the ends to each other. Chaussier considers those and the *stylo-pharyngeus* as but one muscle, and includes all under this last name.

CONSTRICTORES VAGINÆ, *C. cunni*—*c. Vulvæ*, *C. cunni*.

CONSTRINGENTIA, (same etymon as *Constrictio*.) *Astringents*, *Styptics*.

CONSUETUDO, gen. *Consuetu'dinis*, (*con*, and *suescere*, *suetum*, 'to become accustomed,') *Habit*—*c. Menstrua*, *Menses*.

CONSULTANT, Consulting physician or surgeon.

CONSULTA'TION, *Consulta'tio*, *Delibera'tio*, *Cœnolog'ia*, *Content'tio*, *Symbolen'sis*, (*consulere*, *consultum*, 'to hold council.')

This word has several acceptations. In English, it means, almost always, the meeting of two or more practitioners, to deliberate on any particular case of disease. In France, it signifies the written result of such deliberations, as well as the opinion of a physician, given to a patient, who consults him, either personally or by writing.

CONSULT'ING PHYSIC'IAN or SUR'GEON, (F.) *Médecin ou Chirurgien consultant*, *Consultant*. One who consults with the attending practitioner regarding any case of disease. Some physicians, surgeons, and accoucheurs confine themselves to consulting practice.

CONSUMMATUM, *Consummé*.

CONSUMMAT'US, (*consummare* (*con*, and *sumus* 'the whole,') 'to accomplish.')

Confirmed; established; developed,—as *Phthis'is consummata*, 'confirmed consumption.'

CONSUMPTIO, *Consumption*.

CONSUMPTION, (*consumere* (*con*, and *sumere*, *sumptum*) 'to waste away,') *Consump'tio*, *Consump'tio*, *Syntex'is*, *Wasting*, *Wearing*, (F.) *Consumption*. Progressive emaciation or wasting away. This condition precedes death in the greater part of chronic diseases, and particularly in *phthisis pulmonalis*: on this account it is, that phthisis has received the name *consumption*—See *Phthisis*. *Fidère de Consumption*, *Consump'tive fé'ver*, is the same as *Hectic fever*.

CONSUMPTION OF THE BOWELS, *Enterophthisis*.

CONSUMPTION, GAL'LOPING, *Phthi'sis acuta*, (F.) *Phthisis galopante ou aiguë*. *Phthisis pulmonalis*, which rapidly runs through its course to a fatal termination.

CONSUMPTION, PULMONARY, *Phthisis pulmonalis*—*c. Tubercular*, *Phthisis pulmonalis*.

CONSUMPTIVA. Caustics, used for the destruction of fungous growths. *Burnt caustic*, *lunar caustic*, &c., were formerly so called.

CONSUMPTIVE, *Phthisicus*.

CONSUMTIO, *Consumption*.

CONTABESCEN'TIA, (*contabescere* (*con*, and *tabescere*) 'to grow lean,')

Consumption, *marasmus*, *atrophy*, &c.

CON'TACT, *Contact'us*, (*con*, and *tangere*, *tactum*, 'to touch,')

The state of two bodies that touch each other. In the theory of contagious diseases, we distinguish *immediate or direct contact*, as when we touch a patient labouring under one of those diseases; and *mediate or indirect*

contact, when we touch, not the patient himself, but objects that have touched or emanated from him. The air is, most commonly, the medium by which this last kind of contact is effected.

CONTA'GION, *Contag'io*, *Contag'ies*, *Contagium*, *Apoerhu'a*, *Apoer'ysis*. The transmission of a disease from one person to another by direct or indirect contact. The term has, also, been applied, by some, to the action of miasmata arising from dead animal or vegetable matter, bogs, fens, &c., but in this sense it is now abandoned. Contagious diseases are produced either by a virus, capable of causing them by inoculation, as in small-pox, cow-pox, hydrophobia, syphilis, &c., or by miasmata, proceeding from a sick individual, as in plague, typhus gravior, and in measles and scarlatina. [?] Scrofula, phthisis pulmonalis, and cancer, have, by some, been esteemed contagious, but apparently without foundation. Physicians are, indeed, by no means unanimous in deciding what diseases are contagious, and what not. The contagion of plague and typhus, especially of the latter, is denied by many. It seems probable, that a disease may be contagious under certain circumstances and not under others. A case of common fever, arising from *common causes*, as from cold, if the patient be kept in a close, foul situation, may be converted into a disease, capable of producing emanations, which may excite a similar disease in those exposed to them. *Contagion* and *infection* are generally esteemed synonymous. Frequently, however, the former is applied to diseases not produced by contact; as measles, scarlet fever, [?] &c., whilst *infection*, (Prov.) *Smit*, is used for those that require positive contact; as itch, syphilis, &c., and conversely. Diseases which cannot be produced in any other way than by contagion, are said to have their origin in *specific contagion*; as small-pox, cow-pox, measles, hydrophobia, syphilis, &c. Those which are produced by contagion, and yet are supposed to be sometimes owing to other causes, are said to arise from *common contagion*; as typhus, cynanche parotidea, &c.

CONTA'GIONIST, (F.) *Contagioniste*. One who believes in the contagious character of a particular disease,—as of yellow fever.

CONTA'GIOUS, *Contag'iosus*, (F.) *Contagieux*. Capable of being transmitted by mediate or immediate contact, *Communicable*,—as a *contagious disease*, *contagious fever*, *contagious effluvia*, &c. Commonly, the epithet *infectious*, (F.) *Infectieux*, (Prov.) *Ta'king*, *Smit'tle*, *Catch'ing*, is applied to those that are communicated by immediate contact, as itch, syphilis, &c.

CONTAGIUM, Contagion, Miasm.

CONTEMPLABLES DIES, Critical days.

CONTEMPLATIF (F.), *Contem'plative*, *Contemplativus*, (con, and *templum*, 'a place for observation.') Appertaining to contemplation. The predominant idea of the melancholic—of the monomaniac—is sometimes called *contemplative*.

CONTEMPLATIO, Catalepsy.

CONTEMPLATIVE, *Contemplatif*.

CONTENTIO, (con, and *tendere*, *tentum*, 'to stretch,' 'to strive to maintain,') Consultation, Tension.

CONTEXTURE, *Contextu'ra*, *Context'us*, (con, and *texere*, 'to touch, to be near,') see *Continuity*. *textum*, 'to weave,' 'to make a web,') Arrangement of parts;—texture. A name given, metaphorically, to the structure of organized bodies; as the *texture of muscles, fibres*, &c. See *Tissue*, and *Texture*.

CONTIGUITY, **DIARTHROSIS OF**, (con, and *tangere*, 'to touch, to be near,') see *Continuity*.

CON'TINENCE, *Continen'tia*, (*continere* (con, and *tenere*, 'to hold or keep') 'to contain one'sself,' 'to restrain,') Restraint. Abstinence from, or moderation in, the pleasures of physical love.

CONTINENS, gen. *Continen'tis*, Continent.

CON'TINENT, *Cont'ineus*. Restrained. This word is synonymous, also, with *Continued*; (F) *Continu*.

CONTINENT CAUSE, *Cau'sa conjuncta*, is a cause, real or presumed, which having given rise to a disease, continues to act during the whole of its duration. It may be considered synonymous with proximate cause. A *continent fever*, *Febris continens*, is one which preserves during its whole course, the same degree of intensity, without any remission or sensible exacerbation. A disease which continues uninterruptedly, has been also called *Ætiopathet'a*, *Æiopathet'a* or *Æiopathi'a*.

CONTINENTIA, Contineence.

CONTINUED FEVER, *Febris contin'ua* seu *continens* seu *anabat'ica* seu *assid'ua*. A fever which presents no interruption in its course. Continued fevers form a division in the class *Pyrexia* of Cullen, and include three genera,—*Synocha*, *Synochus*, and *Typhus*. It is proper to remark, that some of the older writers make a distinction between the continual fever, *συνεχης*, *febris contin'ua*, and the *synochus* or *febris continens*. Thus, Rhases states that the *synochus* or *continens* is a fever, which consists of one paroxysm from beginning to end; whilst the *continua* is allied to intermittents.

CONTINU'ITY, *Continu'itas*. An adhesion of two things between each other, so that they cannot be separated without fracture or laceration.

CONTINUITY, **DIARTHROSES OF**, (F.) *Diarthroses de Continuité*, are movable joints, in which the bones are continuous, but do not touch immediately, there being between them a ligamentous substance, whose flexibility permits motion. The vertebral articulations are examples of this. **DIARTHROSES OF CONTIGU'ITY**, *Diarthroses de Contiguité*, on the other hand, are movable articulations, in which the bones are not continuous, but touch by surfaces covered with a cartilaginous layer, which is always moistened by synovia.

CONTINUITY, **LESION OF**, see *Continuity*, solution of.

CONTINUITY, **SOLU'TION OF**, *Solu'tio contin'ui*, is any division of parts, previously continuous. Wounds and fractures are *solutions of continuity*. The word *Continuity* is opposed to *Contiguity*: the latter meaning the condition of two things which are near each other, or touch without uniting. There is *contiguity* between the head of the humerus and the glenoid cavity of the scapula, but not *continuity*.

A *lesion of continuity*, *Læsio contin'ui*, is a destruction of continuity caused by disease.

CONTONDANT, Contunding.

CONTOR'SION, *Contor'tio*, (*contorqueo*, (con and *torquere*, *tortum*, 'to wring,') 'I twist about,') Violent movement of a part, accompanied with a kind of torsion, twist or cast; as *contortion of the face*.

CONTORTIO, Contorsion—c. *Columnæ vertebralis*, *Rhachistrophosis*.

CONTRA (L), (F.) *Contre*, in composition, counter, against.

CONTRAAPERTURA, (*contra*, and *aperio*, 'I open,') A *contraopening*, *Contra-incis'io*, *Incis'io prio'ri opposita*, (F.) *Contre-ouverture*. An incision, made in the most depending part of a wound or abscess, when the first opening is not situate in a manner favorable for the discharge of the pus.

CONTRACTILE, *Contractilis*, (*contrahere*, (con, and *trahere*, *tractum*,) 'to draw together,') Capable of contracting. The fibre of muscles is *contractile*.

CONTRACTILITÉ, Contractility—c. *par Défant d'Extension*, *Elasticity*—c. *de Tissu*, *Elasticity*.

CONTRACTILITY, *Contractilitas*, (F.) *Contractilité*. That vital property, which gives, to certain parts, the power of contracting. The muscles of locomotion are endowed with a power of *voluntary contractility*, or one dependent immediately on the action of the brain;—the muscles of the viscera of digestion, and other internal organs, enjoy an *involuntary contractility*. *Contractility* and *irritability* are frequently used synonymously to signify the property possessed by any tissue of *contracting* on the application of an appropriate stimulus.

CONTRACTILITY, *Irritability*.

CONTRACTIO, *Contractio*—c. Cordis, Systole.

CONTRACTION, *Contractio*, *Systole*. Action of contracting. When we speak of the *contraction of a muscle*, we mean the phenomenon it exhibits during the time it is in action.

CONTRACTOR UTERI, Abortive.

CONTRACTURA, *Acampsis*, *Entesis articularis*, *Rigor artium*, *Muscular Stiffness*, (F.) *Contracture*. A state of permanent rigidity and progressive atrophy of the flexor muscles, which prevents the motions of extension beyond a certain limit. The affected muscles form hard cords beneath the skin. On dissection, they are found converted into tendinous fibres, the fleshy fibres having almost disappeared, when the disease has been of any duration. It frequently succeeds other diseases, particularly rheumatism, neuralgia, convulsions, syphilis, colica pictonum, &c. The warm bath, vapour bath, or thermal waters, oleaginous embrocations, mechanical extension of the limbs, &c., are the chief means of treatment.

CONTRÆXTENSIO, Counterextension.

CONTRAFISSURA, (*contra*, and *findere*, *fissum*, 'to cleave;') *Repercusio*, *Rosorius*, *Cutagma Fissura contrajectans*, *Apeche'ma*, *Anticom'ma*, *Antiope*, *Antiorisis*, *Antierus'ma*, *Infertunium*, *Contenterstroke*, (F.) *Contre-coup*, *Contre-fente*, *Contre-fracture*. A fracture, contusion, or injury, produced, by a blow, in a part distant from that which is struck. Five species of *contrafissura* or *contre-coups* may occur in the skull. 1. When the internal table yields and fractures. 2. When the bone breaks in any other part than the one struck. 3. When a bone, which has been struck, remains uninjured, and its neighbour is fractured. 4. When the bone is fractured in a place diametrically opposite to that struck, as in fractures at the base of the cranium, from a fall on the vertex; and lastly, when the violence of the blow produces a separation of the neighbouring or distant sutures. These fractures of the skull are also called *Fractures par résonance*.

CONTRAHENTIA, (*con*, and *trahere*, 'to draw;') Astringents. Styptics.

CONTRAÏNCISIO, *Contra-apertura*.

CONTRAÏNDICATIO, Counter-indication.

CONTRAÏRRITATIO, Counter-irritation.

CONTRALUNA'IS, (*contra*, and *luna*, 'the moon;') An epithet for a woman who conceives during the menstrual discharge.

CONTRASTIMULANS, *Contro-stimulant*.

CONTRASTIMULUS, *Controstimulus*.

CONTRAYERVA, (*contra*, and (S.) *yerba*, 'an herb;') *Dorstenia contrayerva*—c. *Balls*, *Pulvis contrayervæ compositus*.

CONTRAYERVA, *LISON*, *Dorstenia contrayerva*—c. *Mexican*, *Psoralea pentaphylla*—c. *Nova*, *Psoralea pentaphylla*—c. *Virginiana*, *Aristolochia serperitaria*.

CONTRE, in composition, see *Contra*.

CONTRECOUP, ((F.) *coup*, 'a blow;') *Contrafissura*.

CONTRÆXTENSIO, Counterextension.

CONTREFENTE, ((F.) *fente*, 'a fissure;') *Contra-fissura*.

CONTREFRACTURE, *Contra-fissura*.

CONTREINDICATION, *Counter-indication*.

CONTREOUVERTURE, ((F.) *ouverture*, 'an aperture;') *Contra-apertura*.

CONTREPOISON, *Antidote*.

CONTESTIMULISME, *Controstimulus*.

CONTREXEVILLE, *MINERAL WATERS* OF. *Contrexville* is a town in France in the department of Vosges, four leagues from Neufchâteau. The waters contain carbonate of iron, chloride of calcium, and carbonate of lime, chloride of sodium, a bituminous substance, and free carbonic acid. They are frequented by those labouring under cutaneous, scrofulous, and calculeous affections.

CONTRITIO, *Syntrim'ma*, *Syntrip'sis*, *Tritura*, *Tri'tus*, *Tri'tio*, *Tri'p'sis*, (*contra*, and *tritum*, (*con*, and *tero*,) 'to bruise or make small;') *Comminution*, *Trituration*.

CONTRISTIMULANT, *Contrastimulans*, *Hypothen'ic*, (F.) *Hypothen'ique*, (*contra*, and *stimulus*, 'that which excites;') A substance that possesses a particular, debilitating property, acting upon the excitability in a manner opposite to stimulus. A name given to therapeutical agents, which, according to the Italian theory of *contro-stimulus*, are endowed with the property of diminishing excitement by a specific action. These agents are by no means clearly defined.

CONTRISTIMULUS or **CONTRASTIMULUS**, *DOCTRINE OF*, (F.) *Contestimulisme*. Same etymon. The name given by Rasori, half a century ago, to a new medical doctrine, of which he was the originator—*La nuova Dottrina Medica Italiana*. It is founded on the *contro-stimulant*—*Rasorian*—property attributed to a certain number of medicines. In this doctrine, as in that of Brown, under the name *excitability* a fundamental principle of physiology is admitted, by virtue of which living beings are endowed with an aptitude for feeling the action of external agents or exciting influences, and of reacting on these influences. When this excitability is too great, there is excess of stimulus or *Hyperthen'ia*: when too little, there is deficiency or *Hypothen'ia*. Diseases, *general* and *local*, are divided into three great classes, or into, 1. *Hyperthen'ic*; 2. *Hypothen'ic*; 3. *Irritative*. The *contro-stimulant* physicians admit only two classes of medicines—*stimulants* and *contro-stimulants*.

CONTUNDING, *Contundens*, *Contundens*, (*contundere*, (*con* and *tundere*, *tisum*,) 'to bruise;') (F.) *Contondant*. That which causes contusions. An epithet given to round, blunt, vulnerating projectiles, which bruise or lacerate parts without cutting them.

CONTUS, *Contused*.

CONTUS, ((*κωτος*,) 'a pole;') *Penis*.

CONTUSED, *Contusus*, (F.) *Contus*. Affected with contusion. Thus we say—a *contused wound*.

CONTUSION, *Bruise*, (Prov.) *Duke*, *Contusio*, *Collisio*, *Phlas'ma*, *Thlas'is*, *Thlas'ma*, *Th. contusio*, *Rhe'ge*, *Rheg'ma*, *Rheg'mus*, (F.) *Mentruisance*, (Same etymon as *Contunding*,) An injury or lesion—arising from the impulse of a body with a blunt surface—which presents no loss of substance, and no apparent wound. If the skin be divided, the injury takes the name of *contused wound*. The differences of contusions, as to extent, are of course infinite. When slight, the blood stagnates in the capillaries of the skin, or is effused into the subcutaneous areolar tissue. Time and cold applications remove it. When the texture has been lacerated, there is effusion of blood, with more or less torpor in the part affected. Cooling applications, general or topical bleeding, emollients, poultices, &c., are here necessary, according to circum-

stances. In the severest kinds of contusion, all the soft and solid parts, except the skin, are bruised, and, at times, reduced to a kind of pulp. Where the disorganization is to this extent, there is no hope except in amputation. A deep contusion of the soft parts has been called *Sarcotylasis*, and *Sarcotylasma*.

CONTUSION, WIND, see Wind of a Ball.

CONTUSUS, Contused.

CONUS ARTERIOSUS, ('arterial cone,') *Infundibulum* of the heart. The portion of the right ventricle from which the pulmonary artery proceeds, forms a prominence on the right side of the anterior furrow of the heart, and is prolonged towards the left, becoming narrower at the same time, so as to form a funnel-shaped projection, which extends a little beyond the base of the ventricles. This is the *Conus arteriosus*.

CONUS VASCULOSUS, ('vascular cone,') see *Coni Vasculosi*.

CONVALESCENCE, *Convalescentia*, *Analepsis*, *Anastasis*, *Reconvalescentia*, (*convalescere*, (con and *valescere*,) 'to grow well,') *Exanastrophē*. Recovery of health after disease. The time which elapses between the termination of a disease and complete restoration of strength.

CONVALLARIA ANGULOSA, (*convallis*, (con and *vallis*,) 'a deep valley,' in which it abounds.) *C. Polygonatum*.

CONVALLARIA CANALICULATA, *Polygounatum canaliculatum* seu *latifolium*, *Great Sol'omon's Seal*; and CONVALLARIA PUBESCENS, *Polygounatum pubescens*, *Smaller Sol'omon's Seal*: indigenous. A decoction of the roots is said to be cathartic, diuretic, and diaphoretic.

CONVALLARIA MAIALIS. *Lilium Convallium*, *Convallaria*, *C. Mop'pi*, *Maianthemum*, *Lily of the Valley*, *May Lily*, (F.) *Muguet*, *Muguet de Mai*. The recent flowers are reputed to be aromatic, cephalic, errhine, and cathartic. They are not used.

CONVALLARIA MAPPI, *C. Maialis*.

CONVALLARIA MULTIFLORA, *Polygounatum multiflorum*, which grows in this country and in Europe, has analogous properties.

CONVALLARIA POLYGONATUM. The systematic name of *Sol'omon's Seal*, *Convallaria angulosa*, *Polygounatum uniflorum* seu *anceps* seu *vulgarē*, *Sigil'um Salomonis*, *Polygounatum*, (F.) *Seau de Salomon*, *Genouillet*. The root was once used as an astringent and tonic. It is, also, a reputed cosmetic.

CONVALLARIA PUBESCENS, see *Convallaria canaliculata*.

CONVENTUS, gen. *Conventus*, (con, and *venire*, *ventum*, 'to come,') Coition.

CONVERSION, *Conversio*, (con and *vertere*, *versum*, 'to turn,') Change from one state into another.

CONVERSION OF DISEASES, (F.) *Conversion des maladies*, is the change or transformation of one disease into another.

CONVOLUTION, *Convolutio*, (*convolvere*, (con and *volvere*, *volutum*) 'to entwine,') *Epispheerion*, *Gy'rus*, *Helig'mus*, (F.) *Circconvolution*. The rolling of any thing upon itself.

CONVOLUTIONS, CEREBRAL, *Gy'ri* seu *Plicaturæ* seu *Spi'ræ* seu *Processus enteroides* seu *Intest'ula Cerebri*, are the round, undulating, tortuous projections observed at the surface of the brain. In them, Gall's organs, of course, terminate.

CONVOLUTION INTER'NAL, *C. of the Corpus Callosum*, *Convolution d'Ourllet* (Foville). A cerebral convolution of great extent, the principal portion of which is found on the inner surface of each hemisphere above the corpus callosum. In front it bends downwards and backwards to the fissure of Sylvius, and behind, it extends to the middle lobe and forms the hippocampus major.

CONVOLUTIONS, INTES'TINAL, are the turns made by the intestines in the abdomen.

CONVOLUTION D'OURLLET, ([F.] *ourlet*, 'a hem, a border,') Convolution, internal.

CONVOLUTION SUPRA-ORBITAR. A convolution of the brain, which exists on the inferior surface of the anterior lobe, and rests upon the roof of the orbit.

CONVOLVULUS, (same etymon as *Convolution*.) *Ileus*, *Intussusceptio*.

CONVOLVULUS BATATAS. *C. Indicus*, *Batatas edulis*, *Camotes*; the *Sweet Potato*, *Spanish Potato*, *Caroli'na potato*, (F.) *Patate*. Ord. *Convolvaceæ*. This is the only esulent root of the genus *Convolvulus*. It is much eaten in the United States.

CONVOLVULUS CANTAB'RICA, *Cantab'rica*, *Lav'enderleaved Bind'weed*, has been considered antihelmintic and actively cathartic.

CONVOLVULUS INDICUS, *C. Batatas*.

CONVOLVULUS JALAPA. The name of the plant once thought to be the *Jalap* plant; now referred to *Exogonium pur'ga*, *Ipomœa Jalapa* seu *pur'ga* seu *Scidea'na* seu *macrorrhiza*, *Jalapium*, *Jalo'pa*, *Mechoaca'na* *ni'gra*, *Jalap'pa*, *Jalo'pa*, *Jal'ap Root*, *Gialap'pa*, *Xalap'pa*, *Bryonia Mechoacan'na* *ni'gricans*, *Bryonia Peruvia'na*, *Chela'pa*, *Rhabar'barum* *Ni'grum*, *Gelap'pium*, (F.) *Jalap*, is procured from Mexico. Its odour is nauseous; taste sweetish and slightly pungent. It is solid, hard, heavy, brittle; fracture, resinous; internally, light gray; externally, covered with a deep brown, wrinkled bark. Its operation is cathartic, the resinous part griping violently. Dose, 10 gr. to 3ss. A drop or two of any essential oil may prevent it from griping. An active principle has been separated from *Jalap*, to which the names *Jal'apin* and *Cathart'in* have been given.

An inferior kind, called *light*, *fusiform* or *male jalap*, is yielded by *Ipomœa* seu *Convolvulus Orizabensis*, *I. Batatoï'des* seu *Mestitan'ica*, which grows near Orizaba and Mestitlan, in Mexico.

CONVOLVULUS MAJOR ALBUS, *Convolvulus sepium*—c. *Maritimus*, *Convolvulus soldanella*.

CONVOLVULUS MECOACAN, *Mechoaca'næ Radix*, *Jalap'pa alba*, *Rhabar'barum album*, *Piptostegia Piso'nis*, *Mechoacan*, *Jetiencu*, (F.) *Rhabarbe* ou *Rhubarbe blanche* ou *des Indes*, *Jalap blanc*, *Bryone d'Amérique*. A Mexican convolvulus, the root of which possesses aperient properties, and was once extensively used instead of *jalap*. See *Batata de purga*.

CONVOLVULUS MEGALORRHIZUS, *C. Panduratus*.

CONVOLVULUS NIL, *Ipomœa nil*, *Blue Morning-glory*. An East Indian plant, naturalized in some of the Western States. The seeds are sold in Calcutta as a cathartic. The roots have the same properties.

CONVOLVULUS OPERCULATUS, see *Batata de purga*—c. *Orizabensis*, see *Convolvulus Jalapa*.

CONVOLVULUS PANDURATUS, *C. Megalorrhizus*, *Pseudomechoaca'na*, *Fiddle-leaved Bindweed*, *Hog Potato*, *Virginian Bindweed*, *Wild Pota'to*, *Mech'ameck*, *M. Bindweed*, *Wild Jalap*, *Man in the ground*, *Man of the Earth*, *Wild Rhubarb*, *Wild Potato-Vine*, *Kassau'der*, *Kassau'der*, *Kussau'der*, (F.) *Liseron Mechemec*. In Virginia, and some other parts of the United States, the root of this plant has been much recommended in cases of gravel. It is used either in powder or decoction.

CONVOLVULUS PERENNIS, *Humulus lupulus*.

CONVOLVULUS PES CA'PRE, *Burgula*. A plant used in India as a cataplasm in arthritic cases.

CONVOLVULUS REPENS, *C. sepium*.

CONVOLVULUS SCAMMO'NEA seu *SCAMMO'NIA*, *C. Syri'acus*. The systematic name of the *Sem'mony Plant*. A Syrian and Mexican plant; the concrete gummi-resinous juice of which, *Scam'monia*, *Scammoniz Gummi-Res'na*, *Scammonium*,

S. Syriacum, *Diagrydium*, *Dacrydion*, *Scam'mony*, *Mahmoudy*, (F.) *Scammonée d'Alep*, comes in blackish-gray fragments, becoming whitish-yellow when touched with wet fingers. It is a drastic, hydragogue cathartic. Dose, gr. iij to gr. xv, triturated with sugar. See *Extractum Scammonii*. The dried root of the plant *Scammonia Radix*, is official in the Ph. B. (1864.)

CONVOLVULUS, SEA, *Convolvulus Soldanella*.

CONVOLVULUS SEPIMUM, *Convolvulus major albus* seu *repens* seu *Tuguriorum*, *Calyte'gia sepium*, *Great or Hedge Bindweed*, (F.) *Liseron des Haies*, *Grand Liseron*. The concrete juice of this plant is the *German Scammony*. It is violently purgative, and is given in dropsical affections.

CONVOLVULUS SOLDANEL'LA. The systematic name of the *Sea Convolvulus*, *Convolvulus Maritimus*, *Bras'sica Mari'na*, *Calyte'gia Soldanella*, *Sea Bindweed*, *Soldanel'la*, (F.) *Chou Marin*. The leaves of this plant are said to be drastic cathartic; but they are not much, if at all, used.

CONVOLVULUS SYRIACUS, *Convolvulus scammonia*—c. *Tuguriorum*, *C. sepium*.

CONVOLVULUS TURPE'THUM, *Turpe'thum*. The systematic name of the *Turbith* plant. (F.) *Racine de Turbith*. The cortical part of the root of this species of convolvulus is brought from the East Indies. It is a cathartic, but not used.

CONVULSÉ, *Convulsed*.

CONVULSED, *Convulsus*, (*convellere*, (*con*, and *vellere*, *vulsum*, 'to tear,') 'to pull together,') (F.) *Convulsé*.

CONVULSIBILITY, *Spasmophilia*, see *Subsultus tendinum*.

CONVULSIF, *Convulsive*.

CONVULSIO, *Convulsion*—c. *Canina*, see *Canine laugh*.

CONVULSIO CEREALIS, *Convulsio ab Ustilag'inè* seu *Soloniens'is*, *Ergotis'mus epasmodicus*, *Myrmec'iasis*, *Myrmecias'mus*, *Myrmecis'mus*, *Myrmeco'sis*, *Cé'real Convuls'ion*, (F.) *Convulsion céréale*, *Ergotisme convulsif*, *Convulsion de Sologne*. A singular disorder of the convulsive kind, attended with a peculiar tingling and formation in the arms and legs: hence called by the Germans *Kriebelkrankheit*. It is said to be endemic in some parts of Germany, and to arise often from the use of spoiled corn. See *Raphania*.

CONVULSIO HABITUALIS, *Chorea*—c. *Indica*, *Tetanus*—c. *Raphania*, *Raphania*—c. *Soloniensis*, *C. cerealis*, *Ergotism*—c. *Tonica*, *Tonic Spasm*—c. *Uteri*, *Abortion*.

CONVULSION, *Spas'mus*, *Convuls'io*, *Conductio*, *Hieran'osus*, *Disten'tio nervo'rum*, *Spas'mus clonicus*, *Convuls'io clon'ica*, *Eclamps'ia*, *Synsps'ia* *Convuls'io*, *Hyperspasm'ia*, *Clon'ic Spasm*. This word has several acceptations. It means any violent perversion of the animal movements. The word *Convuls'ions* generally, however, signifies alternate contractions, violent and involuntary, of muscles, which habitually contract only under the influence of the will. This alternate contraction, when slight, is called *tremor*; when strong and permanent, *tetanus*, *trismus*, &c. *Spasms*, *Cramp*, *Risus Sardonicus*, and *St. Vitus's Dance* are convulsions.

CONVULSION, SALAAM', *Eclamps'ia* seu *Spas'mus sal'tans*. A name given by Sir Charles Clarke to a singular kind of convulsion in children, in which there is a peculiar bobbing of the head forward.

CONVULSION DE SOLOGNE, *Convulsio cerealis*.

CONVULSIONNAIRE (F.). A name given, during the last century, to individuals who had, or affected to have, convulsions, produced by religious impulses. The epithet was first given to fanatics, who exhibited the most wonderful and varied seizures, at the tomb of Paris, a Jansenist, who died in the year 1727, and was buried in the

cemetery of St. Médard. Some of these were called *Sautseuses* or *Jumpers*, of whom there have been examples in more recent times: others, from barking like a dog, were called *aboyseuses*; and others, from mewing like a cat, *miaulantes*.

A form of convulsion, induced by religious frenzy, has been vulgarly called the 'jerks,' and the gymnastic movements of the *Shakers* may be placed in the same category.

CONVULSIONS OF CHILDREN, *Eclampsia*—c. *des Enfants*, *Eclampsia*—c. *des Femmes enceintes et en couches*, *Eclampsia gravidarum et parturientium*—c. *Puerperal*, *Eclampsia gravidarum et parturientium*. See *Mania*, *dancing*.

CONVULSIVE, *Convulsivus*, *Spasmodes*, *Agitatorius*, (F.) *Convulsif*. That which is accompanied by, or is analogous to, convulsions, as *convulsive cough*, *convulsive disease*.

CONYZA, (κονίζα,) *Inula dysenterica*—c. *Bifoliata*, *Sericocarpus tortifolius*—c. *Coma aurea*, *Solidago Virgaurea*—c. *Major*, *C. squarrosa*—c. *Medica*, *Inula dysenterica*—c. *Pycnostachya*, *Pterocaulon pycnostachyum*.

CONYZA SQUARRO'SA, *C. Major*, *Brephoc'tonon*, *In'ula squarro'sa*, *Great Flea'bane* or *Spikel'nard*, (F.) *Herbe aux mouches*. *Ord. Compositæ*. A European plant, whose strong and disagreeable odour was formerly considered to be destructive to flies, fleas, &c. Its infusion in wine was once used as an emmenagogue and anti-icteric; and in vinegar as an anti-epileptic.

COOKERY, (from *coquus*, 'a cook,') *Culinary art*.

COOLWEED, *Pilea pumila*.

COOLWORT, *Tiarella cordifolia*.

COONTIE or COONTI, see *Arrowroot*.

COOPER'S WELL, see *Mississippi*, *Mineral Waters* of.

COÛPERCULUM OCULI, (*co*, and *operculum*, 'a cover,' 'a lid,') *Palpebra*.

COÛPERTORIUM, (*co*, and *opertorium*, 'a cover,') *Thyroid cartilage*.

COÛSTRUM. The middle part of the diaphragm.

COOT, (Sc. and Prov.) *Malleolus*.

COPAHIER, *Copaiba*.

COPAHU, *Copaiba*.

COPALBA or Copai'ba. The resinous juice or oleoresin of *Copaif'era officinalis* seu *Jacqui'ni*, *C. Multijuga*, *Copaiva officinalis*, and other species of *Copaifera*; (F.) *Copahier*. *Ord. Leguminosæ*. *Sex. Syst.* *Decandria Monogynia*. It is the *Copaif'era officinalis* *Resi'na*, *Balsamum Copaibæ* seu *Copaivæ* seu *Brazilien'sè* seu *de Copai'bâ* seu *Capi'vi*, *Balsam of Copaiba* or *Copaiva*, (vulgarly pronounced *cape'ry*,) (F.) *Copahu*, *Baume de Copahu*, *B. du Brésil*, *Térébinthe de Copahu*. Its odour is peculiar, but not unpleasant; taste pungent, bitter; consistence syrupy; colour yellowish, and transparent. It is soluble in alcohol, ether, and the expressed oils. S. g. 0.950. Its properties are stimulant and diuretic; in large doses it is purgative. It acts on the lining membrane of the urethra, and on mucous membranes in general. It is given in gonorrhœa, gleet, leucorrhœa, &c., in the dose of grt. x to ʒj, twice or thrice a day. Should symptoms of urticaria or diarrhœa arise, the dose is too large, and must be diminished. It can be in-spissated by means of magnesia so as to be made into pills, and a plan has been devised for enveloping it in gelatin, so that its taste is entirely concealed. See *Capsules*, *gelatinous*.

COPAIFERA JACQUINI, (*copaiba*, and *fero*, 'to bear,') see *Copaiba*—c. *Multijuga*, see *Copaiba*—c. *Officinalis*, see *Copaiba*.

COPAIVA OFFICINALIS, see *Copaiba*.

COPAL', *Copale*, *Resi'na Copal'*, *Gumi'ni copal'i'tum*. A resinous substance brought from the

East Indies, South America, and the western coast of Africa, which flows spontaneously from *Elaeocar'pus Copallif'erus*, *Vat'eria Ind'ica*, and probably from different species of *Hymen'a*. *Ord.* Leguminosæ. It is a stimulant like all the resins, and, dissolved in rectified spirits of wine, has been used in cases of spongy gums and looseness of the teeth; but it is now only employed in varnishes.

COPALCHE BARK, see *Croton pseudo-china*.

COP-BONE, *Patella*.

COPE, (κοπη,) *Cut*.

COPHOMA, gen. *Cophom'atis*, (κωφωμα,) *Cophosis*.

COPHONIA, *Acouphonia*.

COPHOSIS, (κωφωσις,) *Copho'ma*, *Coph'otes*, *Sur'ditas*, *Paracou'sis*, *Dysæsthe'sia audit'o'ria*; (from κωφος, 'deaf,' and *osis*.) (F.) *Surdité*. Diminution or loss of hearing. Cullen uses the word synonymously with *Dysæcæa*, and Pinel with *Paracou'sia* or *Parac'oë*. According to Sauvages, *Cophosis* differs from both,—from *Dysæcæa*, because in it the sonorous rays cannot reach the labyrinth; and from *Paracou'sia*, which consists in a confused state of hearing.

Cophosis seems, usually, to be synonymous with deafness,—*Paracou'sis*, *Sur'ditas*.

COPHOTES, (κωφωσις,) *Cophosis*.

COPIOPIA, *Coryopia*.

CO'POS, (κοπος,) *Lassitu'do*, *Fatiga'tio*, *Las'situde*, *Fatigue'*. A state of body in which most of the animal functions are exerted with less promptitude and vigour than common. The ancients admitted three species: 1. That arising from plethora, *Lassitu'do tensi'va* seu *tono'des*; 2. From plethora and increased heat combined, *Lassitu'do phlegmono'sa*, *L. æstiu'sa* seu *phlegmono'des*; and, 3. Owing to a morbid condition of the humours, *Lassitu'do ulcero'sa* seu *helo'des*.

COPPER, Cuprum — c. Ammoniated, *Cuprum ammoniatum* — c. Ammonio-sulphate of, *Cuprum ammoniatum* — c. Rose, *Papaver rheas* — c. Subacetate of, *Cupri subacetat* — c. Sulphate of, *Cupri sulphas* — c. Ammoniacal Sulphate of, *Cuprum ammoniatum*.

COPPERAS, [F.] *Couperose*, *Cupri'rosa* [?], [D.] *Koper rood*, 'red copper,' *Ferri sulphas* — c. Blue, *Cupri sulphas* — c. White, *Zinci sulphas*.

COPPERNOSE, *Gutta rosea*.

COPRACRASIA, (*copro*, a. priv., and κρατος, 'strength,') *Scoracrasia*.

COPRACRATIA, *Scoracrasia*.

COPRAGO'GUM, (*copro*, and αγω, 'I bring away,') *Ster'cus è pri'mis vi'sis edu'cens*. A cathartic. The name of a laxative electuary, mentioned by Ruland.

COPRANON, (κοπρανον,) *Excrement*.

COPRECCRITICUS, *Coproecriticus*.

COPREM'ESIS, *Copriëm'esis*, (*copro*, and εμεω, 'I vomit,') *Vom'itus fæculen'tus* seu *ster'coris*: Vomiting of fæces.

COPREM'ETUS, *Coproëm'etus*, *Copriëm'etus*, *Merdiu'omys*. One who vomits fæces.

COPRIEMESIS, *Copremesis*.

COPRIËMETUS, *Copremetus*.

COPRO, (κοπρος, 'excrement,') in composition, *excrement*.

COPROCRIT'ICUS, *Coproecrit'icus*, (*copro*, and κρινω, 'I separate,') A mild cathartic; an ecoprotic.

COPROËMETUS, *Copremetus*.

COPROËPISCHESSIS, (*copro*, and επισκεσις, 'retention,') *Constipation*.

COPROLITHUS, (*copro*, and λιθος, 'a stone,') see *Calculi* of the stomach and intestines.

COPROPHORESIS, (*copro*, and φερω, 'I carry forward,') *Catharsis*.

COPROPHORIA, *Catharsis*.

COPROFRHËA, (*copro*, and ρεω, 'I flow,') *Diarrhœa*.

COPROSCLEROMA, (*copro*, σκληρος, 'hard,' and oma,) *Coprosclerosis*.

COPROSCLERO'SIS. Induration of fæcal matters; *Coprosclero'ma*.

COPROSTASIS, (*copro*, and στασις, 'standing,') *Constipation*.

COPRUS, (κοπρος,) *Excrement*.

COP'TE, (κοπτη,) *Copton*; (κοπτω, 'I beat or pound,') A sort of cake, composed of vegetable substances, which the ancients administered internally, and applied to the epigastric region in the form of cataplasim.

COP'TIS, (κοπτιω, 'to cut,' in reference to the divided leaves.) *C. trifol'ia*, *Nigell'la*, *Helleb'orus trifol'ius*, *Fil'bra au'rea*, *Chry'sa fibrau'rea*, *Anemo'në Grönlän'dica*, *Goldthread*, *Mouthroot*. The root of this evergreen—*Coptis*, (Ph. U. S.)—is much used in Massachusetts, in aphthous and other ulcerations of the mouth, as a local application. It is a pure bitter, and can be used wherever such is indicated.

CORTIS TEE'TA, *Mish'me Tee'ta*, (Upper Assam), *Honglane* (Chinese). The root of this plant is considered to be a powerful tonic and stomachic.

COPTON, (κοπτον,) *Copte*.

COPULA, ('a band,') *Ligament* — c. *Carnalis*, *Coition* — c. *Cartilaginea*, see *Synchondrosis* — c. *Magna cerebri*, *Corpus callosum*.

COPULATION, (*copulare*, 'to bind together,') *Coition*.

COPYO'PIA, *Copio'pia*, *Kopyo'pia*, *Kopio'pia*, *Ophthalmoco'pia*, (F.) *Lassitude oculaire*, (κοπος, 'fatigue, and ωψ, 'the eye,') *Fatigue* of vision. Weakness of sight. Inability of the eye to sustain continued exertion.

COQ, (onomatopœia,) *Phasianus Gallus* — c. *des Jardins*, *Tanacetum*.

COQUE [L.] *concha*, 'a shell' *DU LEVANT*, see *Menispermum cocculeus* — c. *Levant*, see *Menispermum cocculeus* — c. *d'Æuf*, see *Ovu n*.

COQUELICOT, *Papaver rheas*.

COQUELOURDE, *Anemone pulsatilla*.

COQUELUCHE, *Influenza*, *Pertussis*.

COQUELUCHON [F.], 'a hood,' *Accutium napellus*.

COQUEN'TIA MEDICAMENT'A, (*coquere*, 'to digest,') Medicines which were formerly believed to be proper for favouring the coction or digestion of food.

COQUERET, *Physalis*.

COQUETTE, *Influenza*.

COQUILLE, *Cochlea* — c. *d'Æuf*, see *Ovum*.

COR, as a prefix, *cum*, 'with.'

COR, a Corn.

COR, gen. *Cordis*, (κρη,) *Heart* — c. *Aorticum*, see *Heart* — c. *Arteriosum*, see *Heart* — c. *Bovinum*, *Heart*, hypertrophy of the — c. *Hirsutum*, *Cor villosum* — c. *Hispidum*, *Cor villosum* — c. *Pulmonale*, see *Heart* — c. *Taurinum*, *Heart*, hypertrophy of the — c. *Tomentosum*, *Cor villosum* — c. *Venosum*, see *Heart*.

COR VILLO'SUM, *C. tomentosum* seu *hirsu'tum* seu *hisp'idum*. The condition of the heart, at times, after pericarditis, when it is studded with projections of fibrous matter from its surface.

COR'ACO. In composition, appertaining or relating to the coracoid process.

CORACOBACHILËUS, *Coracobrachi'alis*.

CORACOBRA'CHIAL, *Cor'acobrachi'alis*. Belonging both to the coracoid process and arm.

CORACOBACHIA'LIS (Muscle), *Coracobrachi'us*, *Coraco-humeral* — (Ch.), *Perforat'us CASSE'RII*, *Perforat'us*, is situated at the inner and upper part of the arm. It arises from the coracoid process of the scapula, and is inserted at the middle part of the inner side of the humerus. It carries the arm forwards and inwards, raising the humerus a little. It can, also, by acting inversely, depress the shoulder.

CORACOCALVICULAR, *Coracoclaviculāris*. Belonging to the coracoid process and clavicle.

CORACOCALVICULAR LIGAMENT—called, also, *Omo-claviculār*—serves to unite the clavicle to the coracoid process. It is very irregular, and formed of two fasciculi, the greater part of anatomists have described as particular ligaments, under the names *Coinoid* and *Trapezoid*.

CORACODES, (κορακώδης), Coracoid.

CORACOHUMERALIS, *Coracohumeralis*—c. *Hyoides*, *Omo-hyoides*—c. *Radialis*, *Biceps flexor cubiti*.

CORACOID, *Coracoïdes*, *Coracoïdes*, *Coracoïdes*, (κοραξ, gen. κορακός, 'a crow,' and εἶδος, 'resemblance,') (F.) *Coracoïde*. Resembling the beak of a crow. A name given by Galen—and still retained—to the short, thick process, situate at the anterior part of the upper margin of the scapula, which has some resemblance to the beak of a crow. *Processus Coracoïdeus* seu *uncinatus* seu *corniculatus* seu *anchoralis* seu *rostriformis* seu *ancyroïdes*, *Crow's beak-like process*. This process gives attachment to the *Coracoclaviculār* and *Coraco-acromial* ligaments, and to the *Coracobrachialis*, *Pectoralis minor*, and *Biceps* muscles.

CORACOID LIGAMENT, *Ligamentum Coracoïdeum*, (F.) *L. Coracoïdien*. This name is given to the small fibrous fasciculus, which converts the notch, at the superior margin of the scapula, into a foramen.

CORAIL, Coral—c. *des Jardins*, *Capsicum annuum*.

CORAL, *Corallum*, *Corallus*, *Courallium*, *Arbor Maris*, *Azur*, *Bole'sis*, (κορα, 'I adorn,' and 'als,' 'the sea,' or, according to some, from κορη, 'a daughter,' and 'als,' 'the sea,') (F.) *Corail*. One of the most beautiful productions of the deep. It is fixed to submarine rocks, in the form of a shrub, and is of a bright red colour. It is the habitation of a multitude of animals, of the Zoophyta order, and is formed of a calcareous substance, secreted by the animals themselves. It is in very hard, concentric layers; covered, externally, by a species of porous bark full of cellulæ, each of which contains one of these animals. Linnæus calls the red coral *Pis nobiles*, and M. de Lamarck, *Corallium rubrum*. It is much fished for on the coasts of Barbary and Sicily. Coral was formerly esteemed tonic, absorbent, astringent, &c.; but analysis has shown, that it contains only carbonate of lime and a little gelatin. *Corallium album* is a hard, white, calcareous, brittle substance, the nidus of the *Madrepora oculata*. It has been given as an absorbent.

CORALLIUM, (κοράλιον), Coral.

CORALLINA, (dim. of *corallium*, 'coral,') *Musculus maritimus*, *Corallina officinalis*, *Brion*, *Corallina alba*, *Sea Coralline*, *White Worm-weed*, (F.) *Coralline blanche*. The production of an animal, which belongs to the class *Polypi*, and which is found in all the seas of Europe;—particularly in the Mediterranean. It has the appearance of a plant, is homogeneous, an inch or two in height, of a white, reddish, or greenish colour, salt taste, and marine smell. It contains gelatin, albumen, chloride of sodium, phosphate, carbonate, and sulphate of lime, carbonate of magnesia, silica, oxide of iron, and a colouring principle. It was once much used as a vermifuge; but is not now employed. Dose, ʒss to ʒj, in powder.

CORALLINA CORSICANA, *C. rubra* *Helminthochorton*, *Helminthochorton*, *Elminthochorton*, *Musculus helminthochortus*, *Musculus corallinus* seu *Marinus* seu *Corvicius*, *Conferva* *Helminthochortus*, *Spherocecus helminthochortus*, *Gigartina* seu *Plocaria* seu *Fucus* seu *Alsidium* *helminthochorton*, *Corallina melitochorton*, *Lemnischorton*, *Ceramium helminthochorton*, *Corvici-*

can *Worm-weed* or *Moss*, (F.) *Coralline de Corse* ou *Noire*, *Mousse de Corse*. Corsican moss, of commerce, is a mixture of several marine plants and zoophytes, as the *fucus*, *ceramium*, *ulva*, *coralline*, *conferve*, &c., and has gained great reputation for destroying all kinds of intestinal worms, when given in strong decoction. The Geneva Pharmacopœia directs an official syrup—the *Sirup de Coralline*.

CORALLINE BLANCHE, *Corallina*—c. *de Corse*, *Corallina Corsicana*—c. *Noire*, *Corallina Corsicana*.

CORALLIUM, (κοράλλιον), Coral.

CORALLOIDES FUNGUS, *Clavaria coralloides*.

CORALLUS, Coral.

CORALWORT, *Clavaria coralloides*.

CORCULUM GERMINIS, (dim. of *cor*, 'a heart,') See *Molecule*.

CORD, *Fu'nis*, *Funiculus*, (Latin *chor'da*, itself from χορδή, 'intestine,' afterwards applied to musical cords or strings, made of the intestines of animals.) See *Chorda*.

CORD, SPERMATIC, see *Spermatic cord*—c. *Umbilical*, *Funiculus umbilicalis*.

CORDS, *Vo'cal*, *Cords of Ferrein*, *Chor'dæ vocales*, *Ch. Ferre'nii*. A name given to the ligaments of the glottis, which Ferrein compared to stretched cords, and to which he attributed the production of voice. See *Thyreo-arytenoid Ligament*.

CORDA, *Chorda*, *Cord*—c. *Hippocratis*, *Achillis tendo*—c. *Magna*, *Achillis tendo*—c. *Spinalis*, *Medulla spinalis*.

CORDE, *Chorda*—c. *du Tambour* *Chorda tympani*—c. *du Tympan*, *Chorda tympani*.

CORDEE, *Chordee*.

CORDIA AFRICANA, (after Cordius, a German botanist,) *Sebestina*—c. *Domestica*, *Sebestina*—c. *Myxa*, *Sebestina*—c. *Obliqua*, *Sebestina*—c. *Sebestina*, *Sebestina*—c. *Smoothleaved*, *Sebestina*.

CORDIAL, *Cordialis*, *Cardiacus*, (*cor*, gen. *cordis*, 'heart,') A tonic or excitant medicine, judged to be proper for exciting the heart. A warm stomachic.

CORDIAL, GODFREY'S, see *Godfrey's Cordial*—c. *Nervous*, *Brodum's*, see *Tinctura gentianæ composita*—c. *Sir Walter Raleigh's*, *Confectio aromatica*—c. *Warner's*, see *Tinctura rhei et sennæ*.

CORDIFORM TENDON OF THE DIAPHRAGM, (*cor*, gen. *cordis*, 'heart,' and *forma*, 'form,') *Centre phrenic*.

CORDINE/MA, gen. *Cordine/matis*, (κορήνημα, from κορα, 'the head,' and δύνω, 'I turn round,') *Headache*, accompanied with vertigo. See *Cerebaria*.

CORDIS EMUNCTORIUM, ('emunctory of the heart,') *Axilla*.

CORDOLIUM, (*cor*, 'heart,' and *dolor*, 'pain,') [?] *Cardialgia*.

CORDON (F.), (*corda*, 'a cord,') *Funiculus*. A term applied to many parts, which resemble a small cord.

CORDON NERVEUX, *Ramus Nervosus*. A principal division of a nerve, or the nervous trunk itself.

CORDON OMBILICAL, *Funiculus umbilicalis*.

CORDON SANITAIRE. A line of soldiers or others to prevent communication with an infected district.

CORDON SPERMATIQUE, *Spermatic chord*—c. *Testiculaire*, *Spermatic chord*.

CORDONS SUS-PUBIENS, *Round ligaments of the uterus*—c. *Vasculaires*, *Round ligaments of the uterus*.

CORDS, see *Cord*.

CORE, (κορη), *Pupul* see *Furunculus*:—in composition, the pupil.

CORECLISIS, *Coreoclisis*, *Ankyioclisis*, *Atrypoclisis*, (from *core*, and *κλειω*, 'I close.') Closure or obliteration of the pupil.

CORECTASIS, *Coradialis*, *tolis*, (*core*, and *εκτασις*, 'dilatation.') Dilatation of the pupil.

CORECTOMIA, (*core*, *εκ*, 'out of,' and *τομή*, 'incision.') See *Coretomia*.

CORECTOP'IA, (*core*, *εκ*, 'out of,' and *τοπος*, 'place.') Displacement of the pupil. A condition of the iris in which one segment is larger than the other; so that the pupil is not in the centre.

COREDIALYSIS, (*core*, and *διαλυσις*, 'separation.') See *Coretomia*.

CORE'MATA, pl. of *Corema*, (*κορεω*, 'I cleanse.') Remedies proper for cleansing the skin.

COREMETAMORPHOSIS, (*core*, and *μεταμορφωσις*, 'transformation.') *Dyscoria*.

COREMORPHOMA, gen. *Coremorphomatis*, (*core*, *μορφή*, 'shape,' and *ομα*.) A morbid change in the shape of the pupil.

COREMORPHOSIS, *Coreomorphosis*, *Conformatio pupillæ artificialis*, *Cor'eplasty*, (*core*, and *μορφωσις*, 'formation.') The operation for artificial pupil. See *Coretomia*.

CORENCLEISIS, (*core*, and *εγκλειω*, 'I include.') Operation for artificial pupil, which consists in drawing out, through an incision in the cornea, a portion of the iris, and cutting it off.

COREON'CION, *Coron'cion*, *Coreon'cium*, (*core*, and *ογκος*, 'a hook.') An instrument, used by Langenbeck for the formation of artificial pupil. It is hooked at its extremity. A double-hooked forceps, used by Von Gräfe, is similarly named.

COREOP'SIS TRICHOSPER'MA, *Tick'weed sunflower*, *Tick'seed sunflower*, (*κορψ*, 'the bug,' and *οψις*, 'appearance,' applied to the seeds.) An indigenous plant, of the Composite Family, with large golden yellow rays, which flowers in September. It is said to have been used as an alterative.

COREPLASTY, (*core*, and *πλασσω*, 'to form.') *Coremorphosis*.

CORETODIALYSIS, (*core*, and *διαλυσις*, 'separation.') See *Coretomia*.

CORETOMEDIALYSIS, (*core*, *τομή*, 'incision,' and *διαλυσις*, 'separation.') See *Coretodialysis*.

CORETOM'IA, (*core*, and *τεμνω*, 'to cut.') *Iridotom'ia*, *Coretotom'ia*, *Corotom'ia*, *Iridectom'ia*, *Coretometom'ia*, *Corectom'ia*, *Iridodialysis*, *Corodialysis*, *Coretodialysis*, *Coretomedialysis*, *Iridotomedialysis*, *Iridectomedialysis*.

Various operations for the formation of artificial pupil are so termed. The first four signify the simple incision of the iris for that purpose; the next three, excision of the iris, or incision with loss of substance; the next four, the simple separation or tearing asunder of the iris from the ciliary ligament; the next two, such separation with incision of the iris; and the last, such separation with excision of the part of the iris detached. When a portion of the iris is left strangulated in the wound, it is termed *Iridencleisis*, *Iridencleismus*, and *Iridotomencleisis*.

CORETONECTOMIA, See *Coretomia*.

CORETOTOMIA, *Coretomia*.

CORIAN'DER, (*κορις*, 'the bug,' in allusion to the smell of the fresh herb.) *Coriandrum sativum*.

CORIANDRUM, *C. Sativum*—*c. Cicuta*, *Coriandrum sativum*—*c. Cicuta virosa*—*c. Maculatum*, *Conium maculatum*.

CORIAN'DRUM SATIVUM, *Corian'der*, *Corian'non*, *Coria'non*, (F.) *Coriandre*. Family, Umbelliferae. *Sex. Syst* Pentandria Digynia. The seeds of the coriander—*Coriandrum* (Ph. U. S.), *Coriannon*—have an aromatic odour, and grateful, pungent taste. They are carminative, but are chiefly used to cover the taste of other medicines.

CORIANNON, (*κοριαννον*.) *Coriandrum sativum*.

CORIGEEN, *Fucus crispus*.

CORIITIS, (*corium*, and *itis*.) *Cytitis*.

CORINTHIACÆ, (relating or appertaining to Corinth.) See *Vitis Corinthiaca*.

CORION, (*κοριον*.) *Corium*, *Hypericum perforatum*.

CORIS, (*κορις*.) *Cimex*—*c. Monspelienis*, *Symphylum Petrum*.

CO'RIMUM, *Cho'rion*, *Co'rion*, (F.) *Cuir*. The skin of animals is so called, especially when tanned. The *cutis vera*, or the thickest part of the human skin.

CORIUM PHLOGIS'TICUM, *Crus'ta pleuret'ica seu inflammato'ria seu phlogis'tica*, *Peg'min*, *Inflam'matory Crust or Buff*, *Buffy Coat*, (F.) *Couenne*, *C. Pleurétique*, *C. Inflammatoire*. The grayish crust or buff, varying in thickness, observed on blood drawn from a vein during the existence of violent inflammation, pregnancy, &c. It is particularly manifest in pleurisy, and hence one of its names. For its production, it appears to be requisite that there should be an increase in the proportion of the fibrinous element of the blood over that of the red corpuscles, with augmented aggregation of those corpuscles. The blood is then longer in coagulating, and the red corpuscles have time to subside. In highly inflammatory conditions, the buffy coat assumes a concave appearance on its upper surface, and the blood is, therefore, said to be *cupped*. Polli gave to the coagulative substance the name *brady'brin*. It is the ordinary fibrin of the blood,—the only self-coagulable ingredient in it.

When the blood presents the above appearance, it is said to be *buffy* or *sizey*.

CORK, ([G.] *Kork*; originally, perhaps, from *cortex*, (S.) *corcho*, 'bark.') *Suber*

CORM, *Cormus*.

CORMIER, *Sorbus domestica*.

COR'MUS, (*κορμος*.) *Corm*. In botany, when the stem of a plant, without creeping or rooting, is distended under ground, retaining a round or oval form, it is so called. The *Cormus* is vulgarly termed a root,—*radix*.

CORN, (Saxon *corn*.) (G.) *Kern*. In England, this word means the *Cerealia*, or those seeds which grow in ears, not in pods. In the United States *Corn* always means *Indian Corn*. Its English sense corresponds to the French *Blé* or *Bled*.

CORN, (from *cornu*, 'a horn,') *Ola'vus*, *Clavus*, *Ped'is*, *Ecphy'ma*, *Ola'vus*, *Gemur'sa*, (F.) *Cor*, *Ognon*. A small, hard, corneous tumour, which forms upon the foot, generally on the toes; and is commonly produced on the most projecting parts by the pressure of too tight shoes. A part of the corn is raised above the skin, and forms a round tumour, like the head of a nail: the other portion, which serves as its base, is buried more or less deeply in the integuments, and occasionally extends as far as the tendons and periosteum. Corns may, sometimes, be removed, by immersing the feet in warm water, but commonly they return. They can, likewise, be destroyed by the knife or caustic, or by paring them down and pulling them out by the roots; but these operations are not always as simple as they seem. In the way of palliation, they must be constantly pared; and for the purpose of preventing pressure, any soft plaster, spread upon linen or leather, may be applied, with a hole in the centre to receive the corn; and layer after layer of plaster be added, until they attain the level of the corn. When very irritable, lunar caustic, rubbed over the surface, will generally diminish irritability surprisingly, and in a mode not easy of explanation.

CORN. *Zea mays*.

CORN COCKLE, *Lychnis githago*—c. Flag, *Glaucolus vulgaris*—c. Flower, *Centaurea cyanus*.

CORN, GUINEA, *Panicum Italicum*—c. Indian, *Zea mays*.

CORN SALAD, *Valeriana dentata*.

CORN, SQUIRREL, *Dicentra Canadensis*—c. Wild, *Matricaria chamomilla*.

CORNALINE, Cornelian.

CORNE, Cornu—c. d'Ammon, Cornu ammonis—c. de Bétier, Cornu ammonis—c. de Cerf, Cervus, Cornu cervi, Cochlearia coronopus—c. de Chamois, Cornu rupicaprae—c. Descendante ou inférieure du ventricule latéral, Cornu descendens ventriculi lateralis—c. de la Peau, Cornu—c. Postérieure du ventricule latéral, Cornu posterius seu posticum ventriculi lateralis.

CORNÉ, Corneous.

COR'NEA, (from cornu,) *C. transpa'rens* seu pellucida seu lucida, *Ce'ras*, *Sclerot'ica cerato'id'es*, *Cerato'id'es* seu *Cerato'des membra'na*, *Ceratome'ninx*, *Membra'na cor'nea*. The transparent cor'nea. (F.) Cornée. One of the coats of the eye, so called because it has some resemblance to horn. It is termed transparent to distinguish it from the opaque—*Cornica opa'ca* or *Sclerot'ic*. It is convex, anteriorly; concave, posteriorly; forming nearly one-fifth of the anterior part of the eye, and representing a segment of a sphere, about seven lines and a half, or in. 0.625 in diameter. It seems to be constituted of laminae in superposition, but of the precise number anatomists are not agreed. Henle assigns it four; the third, a very solid cartilaginous lamella, being called *Membrane de Demours* or *M. de Descemet*: see *Aqueous humour*. Messrs. Todd and Bowman assign it five layers.

CORNEA, CONICAL, Staphyloma of the cornea—c. Opaca, Sclerotic—c. Opake, Caligo—c. Sugar-loaf, Staphyloma of the cornea.

CORNEAL, *Cornea'tis*, (F.) *Cornéal*, *Cornéen*. Relating or appertaining to the cornea.

CORNÉE, Cornea,

CORNÉEN, Corneal.

CORNEITIS, (cornea, and *itis*), Ceratitis.

CORNEL, ALTERNATE-LEAVED, (from cornus, 'horny,' the wood), *Cornus alternifolia*—c. American Red-rod, *Cornus sericea*—c. Large-flowered, *Cornus Florida*—c. Panoieled, *Cornus paniculata*—c. Round-leaved, *Cornus circinata*.

CORNEL TREE, *Cornus mas*.

CORNEL, WHITE, *Cornus paniculata*.

CORNE'LIAN, *Carne'lian*, *Chalcedo'nins*, *Carne'olus*, *La'pis Carne'olus*, *Corne'tus*, *Corne'olus*, *La'pis Sard'ius*, (F.) *Cornatine*, (from caro, gen. carnis, 'flesh,' from its flesh colour; or more probably from cornu, 'horn,' because of the colour of the finger-nail; hence *Onyx*, ονύξ, 'the nail.') A precious, semi-transparent stone, found in Sardinia. The ancients ascribed to it a number of absurd properties.

CORNELUS, Cornelian.

CORNEOLUS, Cornelian.

COR'NEOUS, *Cor'neus*, *Hor'ny*, (F.) *Corné*, (cornu, 'a horn.') Having the nature or appearance of horn.

CORNEOUS TISSUE, *Tela cor'nea*, is that which forms the nails. The corneous membrane is the cornea.

CORNES (pl. of corne,) DU CARTILAGE THYROÏDE, *Cornua cartilaginis thyroideæ*—c. du Coccyx, *Cornua coccygis*—c. de l'Hyoïde, *Cornua hyoidei ossis*.

CORNES DE LIMAÇON (F.), *Snail's Horns*. A name given by Anel to the lachrymal puncta and ducts.

CORNES DE LA MATRICE, *Cornua uteri*—c. de la Peau, Horny excrescences.

CORNESTA, (from cornu, 'a horn,') Retort.

CORNET ACOUSTIQUE, (from cornu, 'a horn,') Ear-trumpet—c. de Morgagni, Turbinated bone, superior—c. Møgen, Turbinated bone, middle.

CORNETS, Turbinated bones—c. de Bertin, Sphenoidal cornua—c. *Sphénoïdaux*, Sphenoidal cornua.

CORNEUS, Corneous.

CORNICHON, (dim of corne, 'a horn,') see *Cucumis sativus*—c. de Cerf, Cornu cervi.

CORNICULARIS PROCESSUS, Coracoid process.

CORNIC'ULUM, (dim. of cornu,) A species of cupping instrument, shaped like a trumpet, having a hole at the top for sucking the air out, to diminish the pressure in its interior.

CORNICULUM LARYNGIS, *Capitulum Santorini* seu *Laryng'is* seu *Cartilag'inis arytenoïd'es*, *Subpra-aryt'enoid Cartilage*, *Cartilage* of Santorini. A small, very movable, cartilaginous tubercle observed on the arytenoid cartilages.

CORNIER, Cornus Florida.

CORNIFICATION, (cornu, 'horn,' and fieri, 'to become,') *Racornissement*.

CORNINE, see *Cornus Florida*.

CORNOUILLE, (from cornus, 'horny,' the wood,) Cornus Florida, *C. mas*—c. à Feuilles arrondies, *Cornus circinata*—c. à Grandes fleurs, *Cornus Florida*—c. *Soyeux*, *Cornus sericea*.

COR'NU, *Ce'ras*, *Corn*, *Horn*, (F.), *Corne*. A conical, hard, epidermic projection, which grows on the heads of certain animals, serving them as a weapon of offence and defence. Anatomists have given this name to parts of the human body, which have nearly the same shape as the horns of animals.

CORNU. A horny excrescence; a corneous wart, which occasionally forms on the skin, and requires the use of the knife; (F.) *Corne de la Peau*.—See *Corn*. Also, *Cornu Cervi*.—See *Cervus*. Also, a Retort.

CORNU ACUSTICUM, Ear-trumpet.

CORNU AMMO'NIS, *C. Ari'etis*, *Hippocam'pus maj'or*, *Pes hippocam'pi maj'or*, *Pes Hippopot'ami maj'or*, *Protuberan'tia cylind'rica*, *Ver'mis Bombyc'inus*, *Processus cer'e'ri latera'tis*, (F.) *Corne d'Ammon* ou de Bétier, *Grande Hippocampe*, *Pied de cheval marin* ou d'*Hippocampe*, *Protubérance cylindroïde* (Ch.), *Bourrelet roulé*. A broad, considerable eminence, curved on itself, and situate at the posterior part of the lateral ventricle. Its surface presents two or three tubercles separated from each other by shallow grooves.

The *Accessorius Pe'dis Hippocamp'i*, (F.) *Accessoire du pied d'hippocampe*, is a prominence, usually formed by the base of the inferior cornu of the lateral ventricle of the brain. It is merely a fold of the hemisphere, and was by Malacarne called *Cuisseart* on *Armure de Jambes*.

CORNU ANTE'RIOUS seu ANTI'CUM VENTRIC'ULI LATERA'LIS; *Anterior Cor'nu* of the *Lateral Ventricle*. The portion of the lateral ventricle of the brain, which is lodged in the middle of the lobe, and forms the commencement of that cavity.

CORNU CERVI, *Cor'nu*, *C. Cervi'num*, *Cer'ri Elaphi* *Cor'nu*, *Hartsthorn*, (F.) *Corne de cerf*, *Bois de cerf*, *Cornichon de cerf*. The horns of various species of the stag. They contain about 27 per cent. of gelatin. The *Shavings*, *Raspat'ra* seu *Rasu'ra Cor'nu Cer'vi*, *C. C. raspat'um*, boiled in water, have, consequently, been esteemed emollient and nutritive.

Hart's horn Jelly may be made as follows:—*Hartsthorn shavings*, ℥vj; boil in water Oiv to Oij; strain, and add, whilst hot, of lemon-juice, two tablespoonfuls; white sugar, ℥vj; and *Sherry wine*, two glasses. A good nutriment for the sick, where wine is not improper.

When burnt, the shavings constitute the *Cor'nu cervi calcinat'um*, *Cor'nu ust'um*, *Phos'phas Cal'cia*, *Calca'ria phosphor'ica*, (F.) *Corne de cerf*

calcinée, *Phosphate de chaux*, which consists of 57.5 per cent. of phosphate of lime. It has been used as an antacid, but is wholly inert, as its composition would indicate. It contains only 1 per cent. of carbonate of lime.

Phosphate of lime has also been given in osteomalacia, from a presumption that the disease is owing to a deficiency of earthy matter in the blood, which is doubtless an error—the fault being in the agents of assimilation or the cells of nutrition. With as little propriety, it has been proposed as an excitant to the healthy action of the cells in tuberculosis.

Bone earth is usually sold in the shops for burnt hartshorn.

The British Pharmacopœia (1864), and the Pharmacopœia of the United States, have a *Cornu Phosphas Precipitata*, *Precipitated Phosphate of Lime*, prepared by dissolving burnt ox bones in muriatic acid, and precipitating by solution of ammonia. The dried precipitate, called *bone phosphate of lime*, has no virtues not possessed by the cornu ustum.

Hartshorn was once supposed to possess a bezoardic power.

CORNU CERVINUM, see *Cervus*, *Plantago coronopus*.

CORNU DESCENDENS VENTRICULI LATERALIS, (F.) *Corne descendante ou inférieure du ventricule latéral*, *Dig'ital Cav'ity*, *Descending or inferior cornu of the lateral ventricle*. The termination of the lateral ventricle in the middle lobe of the brain, behind the fissure of Sylvius.

CORNU, ETHMOIDAL, Turbinate bone, middle—c. Middle, Turbinate bone, middle.

CORNU POSTERIUS seu POSTICUM VENTRICULI LATERALIS, *Cav'itas dig'itala*, *Posterior Cornu of the Lateral Ventricle*, (F.) *Corne postérieure du ventricule latéral*. The triangular prolongation of the lateral ventricle of the brain into the substance of the occipital lobe.

CORNU RUPICA'PRE, (F.) *Corne de Chamois*. The horn of the chamois. It has the same properties as the *Cornu Cervi*.

CORNUA, (pl. of *Cornu*), see Turbinate bones.

CORNUA CARTILAG'INIS THYROIDÆ, (F.) *Cornes du Cartilage thyroïde*. Eminences on the thyroid cartilage, distinguished into *great or superior*, which are articulated with the great cornu of the os hyoides;—and into *small or inferior*, united with the cricoid cartilage.

CORNUA COCCY'GIS, *Cornua of the Coccyx*, (F.) *Cornes du Coccyx*. Two small, tubercular eminences at the base of the coccyx, which are articulated with those of the sacrum.

CORNUA CUTANEA, Horny excrescences.

CORNUA HYOIDÆI OS'IS, *Radi'ces ossis hyoid'is*, *Cornua of the Hyoid Bone*, (F.) *Cornes de l'hyoïde*. Four fragments of the os hyoides situate above the body of the bone, and distinguished into the *small or superior*, and the *great or lateral*.

CORNUA LACRYMALIA, Lachrymal ducts—c. *Lac'macum*, Lachrymal ducts, see Lachrymal puncta.

CORNUA SACRA'LIA, *Cornua of the Sacrum*. Two tubercles, situate at the posterior and inferior surface of the sacrum, which are sometimes united.

CORNUA, SPHENOIDAL, see Sphenoidal cornua—c. *Styloid*, see *Hyoides*, os.

CORNUA UTERI, *Cornua of the Uterus*, *Cervæe*, *Plec'tane*, (F.) *Cornes de la Matrice*. The angles of the uterus, where the Fallopian tubes arise. Sometimes applied to the Fallopian tubes themselves.

CORNUE, (from *cornu*, 'a horn,') Retort—c. *Tubulée*, see Retort.

CORNUMUSA, Retort.

COR'NUS ALTERNIFO'LIA, (from *cornu*, 'horn,' owing to the hardness of the wood,') *Alternate-leaved Cor'nel*; indigenous. *Nat. Ord.* Cornaceæ. The bark is one of the "Shaker medi-

cines," being considered diaphoretic and astringent.

CORNUS AMOMUS, *C. sericea*—c. *Blueberried, C. sericea*.

CORNUS CIRCINA'TA, *C. tomento'sa*, *Round-leaved Cor'nel* or *Dog'wood*, (F.) *Cornouiller à feuilles arrondies*. *Order*, Cornaceæ. The bark of this variety has been used for similar purposes with that of *Cornus Florida*.

CORNUS FEMINA, *C. sericea*.

CORNUS FLORI'DA, *Dogwood*, *Dog'tree*, *Box'tree*, *Bit'ter Red'berry*, *Large-flow'ered Cor'nel*, *Mule Great-flow'ered Dog'wood*, *Flor'ida Dog'wood*, *Virgin'ian Dog'wood*, *Box'wood (New Eng'land)*, (F.) *Cornouiller*, *Cornier*, *C. à grandes fleurs*. The bark of this beautiful tree, which grows everywhere in the United States, has been long employed as a substitute for cinchona. Dose, from ʒss to ʒj. Its active principle has been separated from it, and received the name of *Cornine*.

CORNUS MAS, *C. mas'cula*, (F.) *Cornouiller*, *Corneltree*, *Corne'lian cherry*, indigenous in Greece. The fruit, called *Krani'a*, is agreeably acid, and prized by the Turks. The flowers are astringent.

CORNUS MAS ODORATA, *Laurus sassafras*.

CORNUS PANICULA'TA, *Pan'icled or White Cor'nel* or *Dog'wood*, indigenous, has been used as a substitute for *Cornus Florida*.

CORNUS RUBIGINOSA, *C. sericea*—c. *Sanguinea*, *Sebestena, C. sericea*.

CORNUS SERI'CEA, *C. amo'mus* seu *fe'mina* seu *rubigino'sa* seu *sanguine'a*, *Sicamp Dog'wood*, *Red Will'ow*, *Rose Will'ow*, *New Eng'land Dog'wood*, *Fe'male Dog'wood*, *Silk'y-leaved Dog'wood*, *American Red-red Cor'nel*, *Red-red*, *Red O'sier*, *Blueberried Dog'wood*, *Blueberried Cor'nus*, (F.) *Cornouiller soyeux*. The bark, it is said, has been found little inferior to *Cinchona Pallida* in intermittents.

CORNUS TOMENTOSA, *C. circinata*.

CORNUTA, (from *cornu*, 'a horn,') Retort.

CORO, in composition, see *Core*.

COROCLEISIS, *Coreclisis*.

CORODIALYSIS, (*coro*, and *διαλυσις*, 'separation,') see *Coretoma*.

CORODIASTOLE, (*coro*, and *διαστολη*, 'dilatation,') *Coreclisis*.

COROLLA INFUNDIBULIFORMIS, (dim. of *corona*,) see *Tube*, *Fallopian*.

COROMIO'SIS, *Coromeio'sis*, (*coro*, and *μειωσις*, 'diminution,') *Diminution of the pupil*.

COROMORPHOSIS, *Coremorphosis*.

COROMYDRIASIS, (*coro*, and *μυδριασις*,) see *Mydriasis*.

CORONA, Crown—c. *Ciliaris*, *Ciliary body*—c. *Dentis*, *Crown of a tooth*—c. *Glandis*, *Crown of the glans*—c. *Posterior ulnæ*, *Olecranon*.

CORONA RA'DIANS, *Ra'diating Crown of Reil*. Fasciculi of white fibres radiate in all directions from every part of the surface of the optic thalamus, excepting its inner side, which is free and corresponds to the third ventricle; the anterior of these fibres pass directly forwards, the middle fibres outwards, and the posterior backwards, forming the *coro'na ra'dians*.

CORONA REGIA, *Trifolium melilotus*—c. *Ulnæ*, *Olecranon*—c. *Veneris*, *Crown of Venus*.

CORONAD, see *Coronal Aspect*.

CORONÆ CILIARIS MEMBRANULA, (pl. of *corona*,) *Ciliary zone*—c. *Palpebrarum*, see *Tarsus*.

CORON'AL or COR'ONAL, *Corona'lis*, *Coro'na'rius*, (from *corona*,) *Relating to the crown*. A name formerly given to the frontal bone, because on it partly reposes the crown of kings.

CORONAL ASPECT. An aspect towards the plane of the *corona* or crown of the head. *Coronad* is used adverbially to signify 'towards the coronal aspect.'

CORONAL SUTURE, *Sut'ra coron'al'is*, *S. Frontopariet'al'is*, *Pup'pis Sut'ra*, *S. Aven't'is*. The suture of the head, which extends from one temporal bone to the other, over the *crown* of the head, and unites the parietal bones with the frontal. The *Suture Coron'al* of the French anatomists is the suture which unites the two halves of the os frontis at the early period of life. It is a prolongation of the sagittal.

CORONALE, MINERAL WATERS OF. These waters are found near Lucca in Italy. They received their name from an erroneous notion, that they are particularly adapted for curing diseases of the os frontis. Their temperature is 95° Fahrenheit. They contain free carbonic acid, sulphates of lime and magnesia, chlorides of sodium and magnesium, &c., and some iron.

CORONARIUS, Coron'al, Coronary.

CORONARY, *Corona'rius*, (F.) *Coronaire*, (from *corona*, 'a crown,') Resembling a crown.

CORONARY ARTERIES OF THE HEART, Cardiac Arteries.

CORONARY ARTERY OF THE STOM'ACH, *Ar'te'ria Corona'ria Ventric'uli* seu *Gast'rica supe'rior*, (F.) *Artère coronaire stomac'hique*, *A. Stomogastrique* (Ch.), is one of the branches of the cœliac, which passes towards the superior orifice of the stomach, extends along its lesser curvature, and ends by anastomosing with the pyloric branch of the hepatic. This artery furnishes branches to the inferior part of the œsophagus, the cardiac orifice, the great *cul-de-sac*, the middle part of the stomach, the lesser omentum, and anastomoses with the other arteries of those organs.

CORONARY LIG'AMENT OF THE LIVER is a reflection of the peritoneum, which surrounds the posterior margin of the liver. The same term is likewise applied to ligaments which unite the radius and ulna.

CORONARY PLEX'US OF THE HEART. The *anterior* and *posterior coronary plexuses* of the heart are derived from the anterior and posterior cardiac plexuses. See Cardiac Plexus.

CORONARY PLEX'US OF THE STOM'ACH. This plexus of nerves is given off from the upper part of the solar plexus.

CORONARY SINUS OF THE HEART, Sinus, coronary, venous—c. Sinus of Ridley, Sinus coronarius.

CORONARY TEN'DONS, *Ten'dines corona'rie*. The fibrocartilaginous rings at the orifices of the coronary arteries and aorta.

CORONARY VEIN OF THE STOM'ACH, *Ve'na Corona'ria Ventric'uli*, accompanies the artery, and terminates in the vena porta. Sümmering, and some other anatomists, call all the four arteries of the stomach, *Corona'rie Stomach'icæ*.

CORONARY VEINS (of the heart), Cardiac veins.

CORONCION, Coreoncion.

CORONE, (*κορωνη*, 'a crown,') The coronoid process of the lower jaw.

CORONOID, *Coronoï'des*, (*κορωνη*, 'a crown,' and *ειδος*, 'resemblance,') Resembling the beak of a crown. The name has been given to two processes. One, situate at the anterior and superior part of the ramus of the os maxillare inferius, and affording attachment to the temporal muscles: the other, called, also, *sharp process*, situate at the superior part of the ulna, anterior to the great sigmoid fossa, and forming a part of the hinge of the elbow-joint.

CORONOPODIUM, (*κορωνη*, 'a crown,' and *πους*, gen. *ποδος*, 'a foot,' in allusion to the shape, &c., of the leaves,) *Plantago coronopus*.

CORONOPUS, Cochlearia coronopus—c. *Depressus*, Cochlearia coronopus—c. *Ruellii*, Cochlearia coronopus—c. *Vulgaris*, Cochlearia coronopus.

COROTOMIA, (*coro*, and *τεμνειν*, 'to cut,') *Corotomia*.

CORP, (from *ecrpus*,) Cadaver.

CORPORA, pl. of Corpus—c. Albicantia, Mamillary eminences.

CORPORA AMYLA'CEA, *Corpus'cula amyla'cea*, *Amyla'ceous* or *Am'yloid bod'ies*, *Starch corpuscles*, *Animal starch*. A name given by Virchow to certain bodies of microscopic size, found chiefly in the human brain and spinal marrow—in the *epen'dyma ventriculo'rum* and its prolongations—mingled with the proper nerve elements; and having most of the chemical characters of cellulose.

CORPORA ARANTII, Noduli Arantii, Tubercula A.—c. Bigemina, Quadrigemina corpora—c. Candicantia, Albicantia corpora, Mamillary eminences—c. Caverosa, Cavernous bodies—c. Fibrosa, *Corps Fibreux*.

CORPORA FIMBRIA'TA, *Tæ'nia Hippocam'pi*, *Fim'briated or fring'ed bodies*, (F.) *Corps Frangés ou Bordés*, *Bandelettes des Cornes d'Ammon*, *Bandelettes de l'Hippocampe*. The thin, flattened, and very delicate band, situate along the concave edge of the cornu ammonis, which is a continuation of the posterior crura of the fornix.

CORPORA GENICULA'TA, (F.) *Corps Géniculés*, *C. génouillés*. Eminences situate at the lower and outer part of the optic thalami. Each optic tract commences at the *corpus geniculatum externum*. The *corpus geniculatum internum* is merely a tubercle inserted into the bend or knee formed by the corpus geniculatum externum.

CORPORA GLOBOSA CERVICIS UTERI, Nabothi glandulæ—c. Lutea, see Corpus luteum.

CORPORA MALPIGHIA'NA, *Malpigh'ian Bod'ies*, *Ac'ini*, *Corpuscles* or *Glomerules* of *Malpighi*, *Glomer'uli ren'al'es*, *Corpus'cula Malpigh'ii*. Scattered through the plexus formed by the blood-vessels and uriniferous tubes in the kidney, a number of small dark points may be seen with the naked eye, which received their name from Malpighi, their describer. Each of these, under the microscope, is found to consist of a convoluted mass of blood-vessels, which constitutes the true *glandule, corpuscle*, or *glomerule* of Malpighi. As they have been traced into the commencement of the urinary tubes, in which they lie uncovered, it has been supposed that their office may be to separate the watery portions of the blood to be mixed with the proper urinous matter. See Kidney.

CORPORA MAMILLARIA, Mamillary eminences—c. Nervio-spongiosa Penis, Corpora cavernosa—c. Nervosa, Corpora cavernosa.

CORPORA OLIVA'RIA, *C. ova'ta*, *Eminent'ie Oliva'res* seu *ova'les later'al'es*, *Oli'væ*, *Prominent'ie Semi'ova'les Medul'læ Oblonga'tæ*, (F.) *Corps olivaires*, *Olives*, *Eminences latérales*, Ch. Oblong, whitish eminences, situate at the occipital surface of the medulla oblongata, exterior to the corpora pyramidalia.

CORPORA OVATA, Corpora olivaria—c. Pisiformia, Mamillary tubercles.

CORPORA PYRAMIDA'LIA, *Eminent'ie pyramida'les* seu *media'næ inter'næ*, *Pyra'mides*, (F.) *Corps pyramidaux*, *Pyramides antérieures* (Gall), *Eminences pyramidales* (Ch.) Two medullary bundles, placed on either side of the anterior fissure. These bodies have also been called *Corpora Pyramidalia anti'ca*, to distinguish them from the *C. olivaria*, which have been called *Corpora Pyramidalia later'al'ia*.

Over a space, commencing 8 or 10 lines below the pons and extending to the lower end of the medulla oblongata, a portion of each pyramid crosses to the opposite side—*decussation of the pyramids*, *Decussatio pyram'idum*.

CORPORA PYRAMIDALIA POSTERIORA, see Corpora restiformia—c. Pyramidalia testis, Coni vacuosi—c. Quadrigemina, Q. tubercula.

CORPORA RESTIFORMIA, *Cor'ra* seu *Peduncul' medull' oblongat'a*, *Cor'pora pyramidalia posteriora*, *Processus à cerebell'o ad medull'am oblongat'am*, *Posterior pyramida*, *Peduncles of the medulla oblongata*, *Inferior peduncles of the cerebellum*, (F.) *Corps restiformes*, *Cuisses postérieures*, *Pyramides postérieures*, *Racines*, *Bras ou Jambes du cercelet*, *Pétites branches de la moëlle allongée*, *Pédoncles du cercelet*. Two medullary projections, oblong, and of a whitish appearance, which proceed from each side of the upper extremity of the medulla oblongata, and contribute to the formation of the cerebellum. The *posterior pyramids*—*fasciculi graciles*—are described as distinct from the restiform bodies. They are situate in contact with each other, one on each side of the posterior median fissure. They become closely applied to the restiform bodies, and have been considered to be blended with them, and to proceed to form the cerebellum. According to some, they ascend to the cerebrum.

CORPORA STRIATA, *Eminentie Lenticulares*, *Colliculi Nervi Ethmoidalis*, *Apices Cru'rum medull' oblongat'e*, *Ganglion cer'ebri antérieur*, *Anterior cerebral ganglion*, (F.) *Corps striés*, *Grand ganglion supérieur du cerveau* (Gall), *Couches des nerfs ethmoïdaux*, *Corps cannelés*. Pyriform eminences of a slightly brownish-gray colour, which form part of the floor of the lateral ventricles of the brain. When cut, a mixture of gray and white substances is seen, arranged alternately, to which they owe their name. The tract of fibres that ascends from the anterior pyramids passes chiefly into them. Willis considered that the soul resided there.

CORPORA STRIATA SUPERNA POSTERIORA, *Thalami nervorum optico'rum*—c. *Triticea*, see *Hyaides os*—c. *Wolfiana*, see *Corpus Wolffianum*.

CORPS, (from *corpus*.) *Body*—c. *Bordés*, *Corpora fimbriata*—c. *Callex*, *Corpus callosum*—c. *Cannelés*, *Corpora striata*—c. *Caverneux*, *Corpora cavernosa*—c. *Cendré*, *Corpus dentatum*—c. *Ciliaire*, *Corpus dentatum*—c. *Dentelé*, *Corpus dentatum*—c. *Étranger*, *Extraneous body*—c. *Festonné*, *Corpus dentatum*.

CORPS FIBREUX, *Cor'pora Fibrosa*. Bayle has given this name to adventitious fibrous productions of a round shape, more or less adherent, and sometimes having a pedicle, which form in certain parts of the body, particularly in the substance of the uterus.

CORPS FOLLICULAIRE VAGINALE, Cowper's glands in the female—c. *Frangés*, *Corpora fimbriata*—c. *Géniculés*, *Corpora geniculata*—c. *Genouillés*, *Corpora geniculata*—c. *Godronné*, *Fascia dentata*—c. *Humain*, see *Body*—c. *Hyalôide*, *Corpus vitreum*—c. *d'Hygmore* ou *d'Highmore*, *Corpus Highmori*—c. *Innominé de Giraldès*, *Organ of Giraldès*—c. *Jaune*, *Corpus luteum*—c. *Mugueux*, *Corpus mucosum*—c. *Mûriforme*, see *Muri-form*—c. *d'Oken*, *Corpus Wolffianum*—c. *Olivaires*, *Corpora olivaria*—c. *Organisés*, *Organized bodies*—c. *Pampiniforme*, *Corpus pampiniforme*—c. *Phacôide*, *Crystalline*—c. *Pituitaire*, *Pituitary gland*—c. *Pyramidaux*, *Corpora pyramidalia*—c. *Restiformes*, *Corpora restiformia*—c. *Rhombôide*, *Corpus dentatum*—c. *Striés*, *Corpora striata*—c. *Thyrôide*, *Thyroid gland*—c. *Variciforme*, *Epididymis*—c. *Varigieux*, *Corpus pampiniforme*, *Epididymis*—c. *Vitré*, *Corpus vitreum*—c. *de Wolf*, *Corpus Wolffianum*.

CORPSE, (from *corpus*.) *Body*, *Cadaver*.

CORPULENT, *Corpulentus*. *Obesus*, *Cras'sus*, *Fat*, *Fleshy*, (Sc.) *Cor'sy*, (Prov.) *Sob'by*, (F.) *Corpulent*, *Pimelotique*, (*corpus*, 'the body,' and *lentus*, 'thick.') Having an unusual development of fat or flesh in proportion to the frame of the body.

CORPULENTIA, *Cor'pulence*, is synonymous with *Obesity* and *Polysarcia*.

CORPULENTIA CARNOSA, *Torositas*.

CORPUS, gen. *Cor'poris*, *So'ma*, *Bod'y*. Any object which strikes one or more of our senses. Gases, liquids, metals, vegetables, animals, are so many bodies. Natural bodies have been divided into *animal*, *vegetable*, and *mineral*; or into *inorganic*, including the mineral kingdom; and *organised*, including the animal and vegetable. The chief differences between organized and inorganic bodies consists in the former having an *origin by generation*, *growth by nutrition*, and *termination by death*; the latter a *fortuitous origin*, *external growth*, and a *termination by chemical or mechanical force*.

Many parts of the frame have, also, been distinguished by this name, as *Corpus Callosum*, *C. Mucosum*, &c. See *Body*, and *Corpora*.

CORPUS ADENIFORME, *Prostate*—c. *Adenoides*, *Prostate*—c. *Adiposum*, *Pinguedo*—c. *Alienum*, *Extraneous*—c. *Annulare*, *Pons varolii*.

CORPUS CALLOSUM, *Commisura Magna* seu *maxima cer'ebri*, *Trabs Medullaris* seu *Cer'ebri*, *Trabecula* seu *Cop'ula magna cer'ebri*, *Mesolobus*, (F.) *Corps Calleux*, *Voûte médullaire*, *Plafond des ventricules du cerveau*, *Mésolobe* (Ch.) A white, medullary band, perceived on separating the two hemispheres of the brain, which it connects with each other. La Peyronie regarded it as the seat of the soul. 'On it are seen longitudinal and transverse fibres—*Striæ longitudinales Lancis'ii*, (see *Raphe*.) and *Striæ* seu *Chordæ transversales Willis'ii*. The anterior portion, which bends downwards, is termed *genu*; the posterior flexure, *splenium*, *tuber*, (F.) *bourrelet*. The fibres which curve backwards into the posterior lobes from the posterior border of the corpus callosum, have been termed *Forceps*; those which pass directly outwards into the middle lobes from the same point, *Tape'tum*; and those which curve forwards and inwards from the anterior border to the anterior lobe, *forceps anterior*.

CORPUS CAVERNOSUM, see *Cavernous bodies*—c. *Ciliare*, *Ciliary Body*, *Corpus dentatum*, see *Ciliary*—c. *Cinereum*, *Corpus dentatum*, c. *Conicum Rosenmulleri*, *Parovarium*—c. *Conoides*, *Pineal gland*—c. *Crystalloides*, *Crystalline*.

CORPUS DENTATUM, *C. Denticulatum* seu *Cilia'rè*, (F.) *Corps dentelé ou festonné*. A central, oval nucleus, of cineritious substance, met with in the cerebellum; the circumference of which exhibits a number of indentations, surrounded by medullary substance. It is seen by dividing the cerebellum vertically into two equal parts.—The same body has been called *Corps cendré* ou *ciliare* ou *rhombôide*, *Cor'pus Cinereum* seu *Rhombôideum* seu *Rhombôidale*, *Ganglion du cercelet*, *Noyau central des Pédoncles du cercelet*, *Nu'cleus dentatus* seu *fimbriatus* seu *centralis seu rhomboidalis*, *Substantia rhombôidea*, *Gan'glion cilia'rè*, *G. cerebelli*. The term *Cor'pus dentatum*. *Nu'cleus olivæ*, is also given to the ganglion of the corpus olivare, which, like that of the cerebellum, is a yellowish-gray dentated capsule, open behind, and containing medullary matter, from which a fasciculus of fibres proceeds upwards to the corpora quadrigemina and thalami optici.

CORPUS DENTICULATUM. c. *Dentatum*—c. *Discoïdes*, *Crystalline*—c. *Externum*, *Extraneous body*—c. *Extraneum*, *Extraneous body*—c. *Geniculatum*, see *Corpora Geniculata*—c. *Glandiforme*, *Prostate*—c. *Glandosum*, *Prostate*—c. *Glandulosum*, *Prostate*.

CORPUS GLANDULOSUM MULIERUM, *Gland'ulæ Prostatae mulierum*. A vascular spongy eminence, which surrounds the orifice of the urethra, and projects at its under part.

CORPUS HIGHMORI, *C. Highmoria'num*, *Mediastinum testis*, *Meatus semina'rius*, (F.) *Corps d'Hygmore* ou *d'Highmore*, *Sinus des Vaisseaux*

seminifères, (Ch.) An oblong eminence, along the superior edge of the testicle, which seems formed of a reflection of the tunica albuginea, through which the principal trunks of the seminiferous vessels pass before they reach the epididymis.

CORPUS HYALOÏDEUM, *C. vitreum*—c. Incomprehensible, Thymus.

CORPUS LU'TEUM, (F.) *Corps jaune*, 'yellow body.' *Ovariule*, *Oöriule*. A small yellowish body perceived in the ovary, and left after the rupture of one of the vesicles. It was, for a long time, considered an evidence of previous impregnation; but *Corpora lutea* are met with in unquestionable virgins; although the corpora lutea of virgins have been generally regarded to differ materially, in size and character, from those of impregnation, which have been called *true corpora lutea* in contradistinction to the other, called *false corpora lutea*.

CORPUS MU'COSUM, *C. reticulatè*, *Re'tè muco'sum*, *Mu'cus* seu *Re'tè* seu *Stra'tum Malpighii*, *Retic'ulum cutaneum* seu *mucosum*, *Mesoderm'm*, *Mu'cus* *ver*, (F.) *Corps muqueux*, *Couche de Malpighi*. The second layer of the skin has been so called. It is situate between the *cutis vera* and *cuticle*, and gives colour to the body. In the white varieties of our species it is colourless; in the negro, black. By most anatomists the existence of such a layer, distinct from the epidermis, is denied.

CORPUS OKENSE, *Corpus Wolffianum*.

CORPUS PAMPINIFORM'E, (*pampinus*, 'a tendril.') *Hedera'ceus plex'us*, *Plexus pampiniform'is* seu *vasculo'sus funiculi spermatici* *pampiniformis*, (F.) *Corps pampiniforme*, *C. Variqueux*. The plexus or retiform arrangement of the spermatic arteries and veins in the cavity of the abdomen, anterior to the *psaos* muscle.

CORPUS PAPILLA'RE, *Tectus Papilla'ris*, *Cor'pus reticulatè*. The nervous and vascular papillæ situate beneath the epidermis, called by Breschet *Neurothel'ic apparatus*.

CORPUS PHACOIDES, Crystalline—c. Pituitare, Pituitary gland—c. Psalloides, *Lyra*—c. Pyramidale testis, see *Coni vasculosi*—c. Reticulare, *Corpus mucosum*, *Corpus Papillare*—c. Rhomboidale, *Corpus dentatum*—c. Rhomboideum, *Corpus dentatum*.

CORPUS SPONGIO'SUM URE'THRÆ, *Substantia spongio'sa ure'thræ*. This substance arises before the prostate gland, surrounds the urethra, and forms the bulb. It then proceeds to the end of the corpora cavernosa, and terminates in the glans penis, which it forms. Where the urethra enters the bulb, it is surrounded by a layer of erectile tissue, called, by Kobelt, *Colliculi bulbi*. Kobelt describes, in the female, as the analogue to the corpus spongiosum of the male, a venous plexus, which, as it lies between the glans clitoridis, and the part that corresponds, in the male, to the bulb of the urethra, he terms *pars intermedia*, (F.) *Réseau intermédiaire*.

CORPUS STRIATUM, see *Corpora striata*—c. Thymianum—c. Thymus—c. Thymicum, Thymus—c. Thyroideum, Thyroid gland—c. Trigonum, see Urinary Bladder—c. Turbinatum, Pineal gland—c. Varicosum, *Corpus pampiniforme*, Spermatic cord—c. Varicosum testis, Epididymis.

CORPUS VARICOSUM, see Spermatic Veins.

CORPUS VIT'REUM, *C. Hyaloï'deum*, *Hu'mor Vit'reus* seu *Hyaloï'des* seu *hyal'inus*, *Glacia'lis hu'mor*, *Vit'reous hu'mor*, *Vit'rina ocula'ris*, (F.) *Corps vitré* ou *hyaloïde*, *Humeur hyaloïde*. The transparent mass, of a gelatinous consistence, which fills the eye, behind the crystalline. It is contained in cells, formed by the *tunica hyaloïdea*.

CORPUS WOLFFII'NUM, *C. Oken'sè*, *Wolffian body*, *Protoneph'rium*, (F.) *Corps de Wolff* ou *d'Oken*. At a very early period of foetal forma-

tion, bodies are perceptible, which were first described by Wolff, as existing in the fowl, and in the mammalia by Oken. According to Müller, they disappear in man very early, so that but slight remains of them are perceptible after the 9th or 10th week of pregnancy. They cover the region of the kidneys and renal capsules, which are formed afterwards; and they are presumed to be the organs of urinary secretion during the first periods of foetal existence.

CORPUSCLE, *Corpusculum*, (F.) *Corpuscule*, (dim. of *corpus*, 'a body.') A small body or particle.

CORPUSCLES, BLOOD, Globules of the blood—c. Bone, see *Lacunæ* of Bone—c. Caudate, see Caudate—c. Chyle, see *Chyle*—c. Cytoid, see *Cytoid*.

CORPUSCLES, EXUDATION. The organizable nuclei contained in fibrinous fluids, which are the origin of the new tissues formed from such fluids.

CORPUSCLES, GANGLION, see *Neurine*—c. Glandiform, *Acinus*—c. Lymph, see *Lymph*—c. of Malpighi, *Corpora Malpighiana*—c. of Meissner, *Corpuscula tactus*—c. Mucous, see *Mucus*—c. Nerve, see *Neurine*—c. Osseous, see *Lacunæ* of bone—c. of Pacchioni, *Pacchioni*, glands of.

CORPUSCLES, PACINIAN, (F.) *Corpuscles* of *Pacini*, so called from Filippo Pacini, an Italian physician, who, it is generally conceived, first noticed them in 1830. They appear, however, to have been depicted in 1741 by Lehmann, from a preparation by A. Vater, who called them *Papillæ* and *P. nervæ*. Hence, it has been proposed by J. C. Strahl (1848) to call them *Vaterian corpuscles* or *Corpuscles* of *Vater* (*Vater'sche Körperchen*). Small bodies connected with the cutaneous nerves of the palm and sole. They have also been found sparingly and incessantly in nerves at the wrist and elbow, the upper arm, fore-arm, and thigh, and in the inter-costal nerve, the sacral plexus, solar plexus and the plexuses adjacent to it. In each corpuscle there is the termination of a nervous filament. Their uses are not known.

CORPUSCLES OF PURKINJE, see *Canaliculus*—c. Pus, see *Pus*—c. Pyoid, see *Pus*—c. Splenic, see *Spleen*—c. Starch, *Corpora amylacea*—c. Tactile, *Corpuscula tactus*—c. Touch, *Corpuscula tactus*—c. of Vater, *C. Pacinian*—c. of Wagner, *Corpuscula tactus*—c. Weberian, see *Ejaculatory Ducts*—c. White, see *Globules* of the blood—c. White granulated, see *Globulin*.

CORPUSCULA ARANTII, (pl. of *corpusculum*,) *Tubercula A.*, see *Sigmoid valves*—c. Glandularum similia intestinorum, *Peyer glandulæ*—c. Globosa cervicis uteri, *Nabothi glandulæ*.

CORPUSCULA LAËNNEC'II. Fibroid granulations, occasionally observed on thickened portions of the pericardium.

CORPUSCULA MALPIGHII, *Corpora Malpighiana*—c. Ossium, see *Lacunæ* of bone—c. Sanguinis, *Globules* of the blood—c. Seminis, *Spermatozoa*—c. Sesamoidea, see *Sigmoid valves*.

CORPUSCULA TACT'IS, *Tactile* or *Touch Corpuscles*, *Corpuscles* of *Wagner* or of *Meissner*, *Ax'ile bodies*, (F.) *Corpuscules du tact*, (G.) *Axen Körper*. Bodies, composed of a horizontally laminated mass of areolar tissue, which are found in the papillæ of parts endowed with great tactile sensibility. The nerves of touch communicate with them, and their function is probably to render the sense more acute.

CORPUSCULA TRITICEA, see *Hyoides*, os—c. Weberiana, see *Ejaculatory Ducts*.

CORPUSCULAR, *Corpuscularis*. Relating to, or containing, corpuscles, as *corpuscular fibrin*. Fibrin which tends to disintegration, and consists of granules in considerable proportion.

CORPUSCULE, *Corpuscule*.

CORPUSCULES CALCAIRES, *Lacunæ* of

bone—c. *Noirs*, Lacunæ of bone—c. *Des Os*, Lacunæ of bone—c. *Ramifiés*, Lacunæ of bone.

CORPUSCULUM, Corpuscle—c. *Arantii*, see Sigmoid valves—c. *Conium Rosenmülleri*, Parovarium—c. *Sesamoideum*, see Sigmoid valves.

CORRAGO, (cor, 'the heart,' and ago, 'to move.' [?]) *Borago officinalis*.

COR'RE, *Cor'sē*, (κορρη, κορρη) (κερω, 'I shave.' [?]) The temple, the hair.

CORRECTIF, Corrigent.

CORRECTION, *Correctio*, (*corriger*, (cor, and *regere*, *rectum*, 'to rule or order,' 'to correct.') The act of correcting medicines; that is, of diminishing their energy or obviating unpleasant effects, by mixing them with substances which modify their operation.

CORRECTORIUS, Corrigens.

CORRELATION, (cor, and *referre*, *relatum*, 'to accomplish.') See Synergy.

CORRIGEEEN MOSS, *Fucus crispus*.

COR'RIGENT, *Cor'rigens*, *Correcto'rius*, *Cas-ti-gans*, *Infrin-gens*, *Emen-dans*: (same etymon as *Correction*.) (F.) *Correctif*. That which corrects. A corrigent, in a pharmaceutical formula, is a substance added to a medicine to mollify or modify its action. In the following formula, the aloes, if not corrected, might induce tormina. The *Oleum Menthe* is added as a corrigent.

R Aloes	3j
Olei Menthe	gtt. v
Syrup q. s. ut fiant Pilulæ xvj	

CORRIG'IA, (akin to *corium*, 'leather.') Leather straps. By extension, the term has been applied to the tendons and ligaments.

CORROBORANT, *Corrob'orative*, *Corrob'orans*, *Corrob'orativus*, *Rob'orans*, *Muscula'ris*, *Restau-rans*, *Restor'ative*, *Bra'cing*, (*corrob'orare*, (cor, and *rob'ur*, 'strength,') 'to strengthen.') (F.) *Corrobo-rant*, *Corroboratif*, *Fortifiant*. Any substance which strengthens and gives tone. Wine, for example, is a corroborant. See Tonic.

CORROBORANTIA, Tonics.

CORROBORATIF, Corroborant.

CORROBORATIVE, Corroborant.

CORRODANT, Corrosive.

CORRODENS, (cor, and *rodere*, *rosum*, 'to gnaw.') Corrosive.

CORROSIF, Corrosive.

CORROSION, *Corro'sio*, *Diabro'sis*, *Anabro'sis*, *Ero'sio*. The action or effect of corrosive substances.

CORRO'SIVE, *Corro'dens*, *Diabro'ticus*, *Cor-ro-sivus*, (F.) *Corrosif*, *Corrodant*. Corrosives are substances, which, when placed in contact with living parts, gradually disorganize them. *Caus-tic alkalies*, *Mineral acids*, *Corrosive sublimate*, are *corrosives*. They act either *directly*, by chemically destroying the part,—or *indirectly*, by causing inflammation and gangrene.

CORROSIVE POISON, see Poison.

CORRUGATIO, Corrugation—c. *Cutis*, Wrinkle.

CORRUGATION, *Corrugatio*, *Syn'e'rema*, (cor, and *ruga*, 'a wrinkle.') Wrinkling, Frowning, (F.) *Froncement*; the contraction of the *Corrugato-res Supercil'ii* muscles. Corrugation of the skin is often owing to the application of styptic medicines: it is rendered by them unequal and rugous.

CORRUGATOR COITERIL, (gen. *Corrugato'ris*.) Corrugator supercili.

CORRUGATOR SUPERCIL'II, *Musculus supercil'ii seu frontalis vel'rus seu Corrugator Coite'rii*, *Supercilio'ris*, (F.) *Cutaneo-sourcilier*, *Muscle Sourcilier ou Sourcilier*, *M. Fronto-Sourcilier*. A muscle situate in the eyebrows. It is attached, by its inner extremity, to the superciliary ridge, and is confounded, externally, with the occipito-frontalis and orbicularis palpebrarum. It carries the eyebrow inwards, and wrinkles the skin of the forehead.

CORRUPTION, *Corrup'tio*, *Phtho'ra*, *Diaph-thora*, (*corrumpere*, *corruptum*, (cor, and *rumpere*, 'to break,') 'to destroy.') Act of corrupting. State of being corrupted. Reaction of the particles of a body upon each other. It is probable that something like corruption may take place even in the living body. The word is used, at times, like *Boue* with the French, in a vague manner for a state of puriform, putrid, or other breaking down of parts or humours. 'To spit corrup-tion,' is a vulgar expression.

CORRUPTUS, Abalienatus.

CORSE, (from (L.) *corpus*.) Cadaver.

CORSE (κορση), Corpe.

COR/SET, ([F.] *corps*, 'the body.') *Stetho-desm'ium*, *Stethodesm'is*, *Stethodesm'us*, *Tu'nica Thora'cis*, *Tho'rax*, *Pectora'lē*. An article of dress, which closely embraces the trunk, and is much used by females in civilized countries. When corsets or stays are worn very tight, many serious evils result from the unnatural compression.

Different bandages, more or less complicated, which embrace the greater part of the trunk, are likewise so called.

An *orthope'dic cor'set* is one intended to obviate deviation of the stature.

CORSET DE BRASDOR. The name of a bandage invented by one Brasdor, to keep *in situ* the fragments of a fractured clavicle.

CORSSY, (from *corpus*.) Corpulent.

CORTALON, Senecio.

COR'TEX, gen. *Cor'ticis*, *Phloios*, *Phlo'os*, *Phlo'us*, *Lem'ma*, *Bark*, (F.) *Écorce*. This word has often been applied exclusively to *Cinchona*: thus, we say *Bark*—the *cortex* or *bark* *kar' éforyn*. It means, also, any bark.

CORTEX ABDSTRINGENS BRASILIENSIS. An astringent bark introduced from Brazil into Germany in the year 1828. It is said to be obtained from *Mimosa cochleacarp'a seu virginalia*. It has been used with advantage in all cases in which astringent barks in general are indicated. Dose of the powder ʒj to 3ss.

CORTEX ALCOGNOCO, *Alcornoque*—c. *Anisi stel-lati*, see *Illicium anisatum*—c. *Antiscorbuticus*, *Canella alba*—c. *Aromaticus*, *Canella alba*—c. *Aurantii*, see *Citrus aurantium*—c. *Brayeræ an-thelmintica*, see *Rottlera Schimperii*—c. *Canellæ Malabarica*, *Laurus cassia*—c. *Cardinalis del Lugo*, *Cinchona*—c. *Caryophyllatus*, see *Myrtus caryophyllata*—c. *Caryophylloides*, *Laurus culila-wan*.

CORTEX CEREBRI. The *Cortical*, *Cinerit'ious*, *Vesic'ular* or *Gray substance of the Brain*. The gray portion observed at the exterior of the cere-brum and cerebellum; so called because it forms a kind of bark to the medullary substance. Gall considers, that this substance forms the nerves; and therefore calls it *Substance matrice des Nerfs*.

The name *cortical* is likewise given to the external substance of the kidneys, because it is of a deeper colour than the inner part of the organ, and forms a kind of envelope to it.

CORTEX CHABARRO, *Alcornoque*—c. *Chacarrilla*, *Croton Cascarrilla*—c. *China*, *Cinchona*—c. *Chinæ flavus*, *Cinchonæ cordifolia* cortex—c. *Chinæ luteus*, *Cinchonæ cordifolia* cortex—c. *Chinæ regius*, *Cinchonæ cordifolia* cortex—c. *Crassier*, *Laurus cassia*—c. *Culilaban*, see *Laurus Culila-wan*—c. *Culilawan*, *Laurus Culilawan*—c. *Den-tium*, *Enamel*—c. *Eleutheriæ*, *Croton cascarrilla*—c. *Flavus*, *Cinchonæ cordifolia* cortex—c. *Lavola*, see *Illicium anisatum*—c. *Magellanicus*, see *Wint-tera aromatica*.

CORTEX MUSENÆ, see *Rottlera Schimperii*.

CORTEX O'VI, *Cortical mem'brane*. This mem-brane, so called by Boer and Granville, is usually regarded as a uterine production, and designated *Decid'ua reflex'a*. They consider it to surround the ovule, when it descends into the uterus, and

to enclose the shaggy chorion. It is absorbed during the first months of utero-gestation, so as to expose the next membrane to the contact of the decidua, with which a connexion takes place at the part where the placenta is to be formed. In that part, Boer and Granville consider, that the Cortex Ovi is never altogether obliterated, but only made thinner, and in process of time is converted into a mere pellicle or envelope, which not only serves to divide the filiform vessels of the chorion into groups or cotyledons, in order to form the placenta, but also covers those cotyledons. This, Dr. Granville calls *membra'na pro'pria*. See *Decidua membrana*.

CORTEX PALLIDUS, *Cinchona lanceifolia* cortex—c. *Patrum*, *Cinchona*—c. *Peruvianus*, *Cinchona*—c. *Profluvii*, *Nerium antidysentericum*—c. *Ruber*, *Cinchona oblongifolia* cortex—c. *Striata dentium*, Enamel of the teeth—c. *Thuris*, *Croton cascarrilla*—c. *Winteranus*, see *Winters aromatica*—c. *Winteranus spurius*, *Canella alba*.

CORTICAL, *Cortica'lis*; (from *cortex*, 'bark,') Belonging to the bark.

CORTICAL MATTER OF THE BRAIN, *Cortex Cerebri*—c. *Membrane*, *Cortex Ovi*—c. *Substance of the Teeth*, see *Tooth*.

CORTUSA AMERICANA, (after Prof. J. A. Cortusi, botanist of Padua), *Heuchera cortusa*.

CORU. An Indian tree, the bark of whose root furnishes a milky juice, which is employed in diarrhoea and dysentery. It is also called *Coru Cana'rica*.

CORUSCATIO, (*coruscare*, 'to flash,') *As-trape*.

CORVISARTIA HELENIIUM, (after Corvisart, physician to Napoleon the 1st.) *Inula helenium*.

CORYBAN'TIASM. *Corybantias'mus*, *Corybantias'mus*, (from *Κορυβας*, one of the Corybantes.) A name formerly given to a kind of frenzy, in which the sick were tormented by fantastic visions, and perpetual want of sleep.

CORYDALIS BULBOSA, (*κορυδαλιν*), *Fumaria bulbosa*—c. *Canadensis*, *Dicentra Canadensis*—c. *Cava*, *Fumaria bulbosa*—c. *Tuberosa*, *Fumaria bulbosa*.

CORYLUS, (*κορυλος*, from *κορυς*, a 'helmet,' from the involucre,) *Corylus avellana*.

CORYLUS AVELLA'NA, *Corylus*, *Bundurh*, *Avella'na*, *Coryd'ion*, *The Ha'zelnut Tree*, (F.) *Coudrier*, *Noisetier*; *Ord. Amentaceæ*; *Sex. Syst. Monocia Polyandria*. The nut,—*Fil'bert*, (F.) *Av'eline*; *Ha'zelnut*, (F.) *Noisette*,—of this tree, is much eaten in many countries. Like all nuts, it is by no means easy of digestion. It is the *Nux avella'na*, *καρυον Παντικον*, of the ancients.

CORYLUS ROSTRA'TA, *Beak'ed Ha'zel*. An indigenous shrub, *Nat. Ord. Amentaceæ*; *Sub-order*, *Cupuliferæ*; *Sex. Syst. Monocia Polyandria*; the nut of which is surrounded by a coriaceous and scaly involucre, terminating in a tube covered with short and thick bristles. These bristles have been given as an anthelmintic in the same cases and doses as *mucuna*.

CORYMBETHRA, (*κορυμβηθρα*), *Hedera helix*.

CORYNE, (*κορυνη*, 'a club,') *Penis*.

CORYPHE, (*κορυφη*), *Acme*, *Vertex*—c. *Cordis*, see *Mucosæ*.

CORYS, (*κορυς*), *Vertex*.

CORY'ZA, (*κορυζα*, from *κορυς*, or *καρυς*, 'the head,' and *ζω*, 'to boil,') *Grave'do*, *Rhini'tis*, *Catag'imus*, *Catastalag'mus*, *Stillicid'ium Na'rium*, *Phlegmatorrhag'ia*, *Blennorrhæ'anasa'lis*, *Blennorrhin'ia*, *Des'tillatio*, *Catar'rhus ad Na'res seu nasa'lis*, *Rhinocatar'rhus*, *Angi'na nasa'lis*: vulgarly *Running at the nose*, *Cold in the head*; (Prov.) *Snurle*; in French, *Rhume de cerveau*, *Catar'rhe nasal*, *Enchifrènement*; in old English, *Pose* or *Mur*; whence *Mur'en* and *Mur'rain*. Inflammation, attended with increased discharge, of the

Schneiderian membrane lining the nose, and the sinuses communicating with it. The affection generally subsides without any medical treatment.

CHRONIC coryza is termed, also, *Oza'na be'ni'na*.

CORYZA ENTONICA, *Ozæna*—c. *Maligna*, *Ozæna*—c. *Ozænosa*, *Ozæna*—c. *Purulenta*, *Ozæna*—c. *Scarlatiosa*, *Rhinocæce*—c. *Ulcerosa*, *Ozæna*—c. *Virulenta*, *Ozæna*.

COSCINISMUS, (*κοσκινον*, 'a sieve,') *Cribration*.

COSCINA, (*κοσκινά*), see *Cribration*.

COSMESIS, (*κοσμησις*), *Cosmetics*.

COSMETICS, *Arts cosmet'ica*, *Cosme'sis*, (F.) *Cosmétique*, (*κοσμεω*, 'I adorn,' 'I embellish,') The art of improving the beauty. *Cosmetic*, (*κοσμητικος*), *Stilbo'ma*, is, also, used for the different means employed for that purpose; as the compounds, into which enter the oxides of lead, bismuth, mercury, arsenic, &c. All these, however, injure the skin, and often give rise to unpleasant cutaneous affections. Frequent ablution with cold water, and bathing are the best cosmetics. Essences, soaps, and all the preparations intended for the toilet, fall, also, under this head.

COSMÉTIQUE, *Cosmetics*.

COSMETOL'OGY, *Cosmetolog'ia*, (*κοσμεω*, 'to adorn,' and *λογος*, 'a discourse,') A treatise on the dress, and cleanliness of the body.

COS'MOS, (*κοσμος*, 'the world,' 'order,' &c.) The order which was supposed to preside over critical days. Hippocrates and others have termed *κοσμοι*, 'bracelets,' employed, not only as ornaments (*κοσμεν*, 'to adorn,') but as therapeutical agents.

COSSA, *Haunch*.

COS'SUM. A malignant ulcer of the nose, often of a syphilitic character.

COS'SUS, *Cos'sia*. A sort of white, short, thick worm or larve, found in trees, logs of wood, &c., and used by the Romans as a great article of *gourmandise*. They were, also, applied to ulcers. The term has, likewise, been given to small vermiform pimples on the face—acne—which arise from inflammation of the sebaceous follicles.

COST, *Costa*.

COS'TA, *Pleu'ra*, *Pleu'rurum*, *Pleuro'ma*, *Cost*, *Rib*, (*custodire*, 'to guard,' 'to defend,') (F.) *Côte*, old French, *Coste*. The ribs are 24 in number;—12 on each side. They are irregular, long, bony curves: slightly flattened, and situate obliquely at the sides of the chest. The intervals between them are called *Intercos'tal spaces*, (F.) *Espaces intercostaux*, and they are numbered *first*, *second*, *third*, &c., reckoning from above to below. They have been distinguished into 1. *Cos'tæ ve'ræ*, *Pleu'ræpoph'yses* of Owen, *True ribs*, (F.) *Vraies Côtes*, *Côtes sternales*, *Côtes vertébro-sternales* (Ch.), and 2. *Cos'tæ spi'ri'æ*, *Mendo'sæ* seu *No'tiæ Cos'tæ*, *False ribs*, *Côtes asternales* (Ch.), *Faussees Côtes*. The *true* or *sternal ribs*, as they have also been called, are the first 7; which are articulated at one extremity to the spine, and at the other, by means of their cartilages, *hæmapoph'yses* of Owen, to the sternum. The *false* or *short ribs* are the remaining 5: the uppermost three being united, by means of their cartilages, to the cartilage of the last true rib. The others are free at their sternal extremity, and, hence, have been called *Float'ing ribs*, *Cos'tæ fluctuantes*, (F.) *Côtes flottantes*. The vertebral extremity of each rib is slightly expanded. It is called the head of the rib—*Capit'ulum Cos'tæ*: the space between this and the tubercle is the *collum* or *neck*. Anterior to the tubercle is the *an'gle*—*An'gulus* seu *Os'bitus Cos'tæ*. The *angle* is the part where the bone bends to form the lateral part of the thorax.

COSTÆ (gen. of *Costa*) CAPITULUM, see *Costa*—c. *Fluctuantes*, see *Costa*—c. *Mendosæ*,

see Costa—c. Spuriar, see Costa—c. Veræ, see Costa.

COSTAL, *Costalis*. Appertaining or relating to a rib—as *costal cartilage*.'

COSTE, Costa.

COSTIVE, Constipated.

COSTIVENESS, Constipation.

COSTMARY, Tanacetum balsamita.

COSTO, (*costa*, 'a rib.'). In composition, relating or appertaining to a rib.

COSTOARDOMINAL, Obliquus externus abdominis—c. *Basi-scapularis*, Serratus magnus—c. *Claviculaire*, Costoclavicular Subclavian muscle.

COSTOCLAVICULAR, Costoclavicularis. Belonging to the ribs and clavicle.

COSTOCLAVICULAR LIGAMENT, *Cleid'ocostal Ligament*, is a fibrous, flattened fascia, which extends, obliquely, from the cartilage of the first rib to the inferior surface of the clavicle.

COSTOCHORACOÏDIEN, Pectoralis minor—c. *Scapularis*, Serratus magnus.

COSTOSTERNAL, *Costosternalis*. Relating to the ribs and sternum. The articulation of the sternum with the anterior extremity of the first seven ribs.

COSTOTRACHELIA'NUS. Relating to the ribs and to the tracheal or transverse process of the neck. Under the name *Costo-trache'lian*, Chaussier designates the anterior and posterior *sealen*.

COSTOTRANSVERSA'RIOUS. Relating to the ribs, and to the transverse processes.—Bichat gave this name to the articulation of the tuberosities of the ribs with the transverse processes of the spine.

COSTOVERTEBRAL, *Costovertebra'lis*. Belonging to the ribs and vertebrae. Bichat gave this name to the articulation of the head of the ribs with the vertebrae, and to the ligaments connected with it.

COSTOXI'PHOID, *Costoxiphoi'deus*. The name of a ligament, which unites the cartilage of the seventh rib to the xiphoid or ensiform cartilage.

COSTUS. (From the Arabic.) The ancients seem to have given this name to several plants. One has borne the appellation, since the time of Dioscorides,—the *Costus Arabicus* seu *Indicus* seu *speciosus* seu *amarus*, *dulcis*, *orientalis*. *Amomum hirsutum*, *Helle'nia grandiflora*, *Bank'sia speciosa*, *Tsida'na*, (F.) *Canne Congo*, *Canne de Rivière*: Family, Amomæ: Ser. Syst. Monandria Monogynia; the root of which is aromatic, and has been considered tonic, carminative, diuretic, emmenagogue, &c. The virtues of the ancient costus are highly extolled by Theophrastus, Dioscorides, Pliny, and Galen.

COSTUS AMARUS, &c., Costus—c. *Arabicus*. Costus—c. *Corticæus*, *Canella alba*—c. *Hortorum*, *Tanacetum balsamita*—c. *Hortorum minor*, *Achillea ageratum*—c. *Indicus*, Costus—c. *des Jardins*, *Balsamita suaveolens*—c. *Nigra*, *Cynara scolymus*—c. *Speciosus*, Costus.

COSTYLE, Cotyle.

COTA, Anthemis cotula.

COTARONIUM; an obscure term, used by Paracelsus for a universal solvent: such a thing as does not exist.

CÔTE, Costa.

CÔTES ASTERNALES, see Costa—c. *Col des*, *Collum costarum*—c. *Fausses*, see Costa—c. *Flottantes*, see Costa—c. *Sternales*, see Costa—c. *Vertébrosternales*, see Costa—c. *Fraies*, see Costa.

COTIGNAC, pron. *cotinniac*. (Called after the city of Cotignac.) A kind of conserve or preserve, prepared from quinces not entirely ripe, and sugar. It is esteemed stomachic and astringent.

COTON, Gossypium—c. *Azotique*, see Collo-

dion—c. *Fulminant*, see Collodion—c. *Poudre*, see Collodion.

COTONIA, Pyrus codynia.

COTTON, (Arab. *kutn* or *kutun*.) Gossypium.

COTULA, (from *κωυλη*, 'a cup,' from the shape of the flowers,) *Anthemis cotula*—c. *Fœtida*, *Anthemis cotula*.

COTULA MULTIFIDA. A South African plant, which is used by the Hottentots in rheumatism, scalds, and cutaneous diseases.

COTURNIUS, LIQUOR OF, *Liquor Coturni'nis*, L. of *Cotugno*, *Aqu'ula acutica*, *Aqua audit'oria*, *Perilymph'a*, *Per'ilymph*, (F.) *Lymph'e ou Humeur de Cotugno*. A transparent, slightly viscid fluid, which surrounds the membranous labyrinth, and of which Cotugno made mention. It is also called *Aqua Labyrinthi*.'

COTURNIX, Tetrao coturnix.

COTYLE, (*κωυλη*), *Cot'yla*, *Cotyle'don*, *Cos'tylé*; the same as *Acetabulum*. A hollow cavity of a bone, which receives the head of another bone, and particularly 'the cotyloid cavity.' *Κωυλη* signified a drinking cup, and, indeed, anything hollow, as the hollow of the hand.

COTYLE'DON, gen. *Cotyledonia*, (*κωυλη'ων*.) Cotyle—c. *Marina*, *Umbilicus marinus*.

COTYLEDON ORBICULA'TA. A plant of South Africa, *Nat. Ord.* Crassulaceæ. The fresh juice is used in epilepsy; and Dr. Pappe speaks well of it. The leaves form a good application to hard corns. *Crass'ula arborescens* has the same properties.

COTYLEDON UMBILICUS, C. U. *Veneria*, *Navel-cort*, *Vénus's Navel-cort*, (F.) *Nombil de Venus*. A plant which grows in Europe on old walls and rocks. The leaves are emollient, and are applied externally to piles, inflamed parts, &c. Internally, the juice has been given in epilepsy. The flowers have been used in calculus cases, and in dropsy.

COTYLEDONS, *Acetab'ula uteri'na*. The lobes which, by their union, form the placenta.

COTYLOID, *Cotyloï'des*, (*κωυλοειδής*, from *κωυλη*, 'a drinking cup,' and *ειδός*, 'form.') *Cotyloï'deus*, *Cotylo'des*, (*κωυλωδής*.) Resembling the ancient *κωυλη*. The name of a hemispherical cavity, situate in the os innominatum, which receives the head of the os femoris;—*Fossa cotyloidea*, *Sinus Coxæ*, *Acetabulum*, *Pyræ*.

COTYLOID LIGAMENT, *Ligamentum Cotyloï'deum*, (F.) *Ligament cotyloïdien*, is a very thick, fibro-cartilaginous ring, surrounding the cotyloid cavity, the depth of which it increases.

COTZ, Hagenia Abyssinica.

COU, Collum—c. *Gros*, *Bronchocele*—c. *du Pied*, *Instep*.

COUCH GRASS, *Triticum repens*.

COUCHE, *Couches*, (F.) (from *coucher*, 'to lie down.'). This word is used, 1. For *parturition*, *accouchement* or *delivery* (*puerperium*, *parturi'tio*;) hence, *une couche heureuse*, 'a happy delivery'; *une fausse couche*, 'a premature delivery'; and, 2. For the time during which a female remains in bed on account of delivery, — *Temps puerperii*, the *child-bed state*. The *Lochia* have been termed *Suites de couches*.

Couche also means a layer, bed or thalamus, as, *COUCHE CELLULEUSE*, *Membrana granulosa*—c. *de Malpighi*, *Corpus mucosum*.

COUCHES DES NERFS ETHMOÏDAUX, *Corpora striata*—c. *des Nerfs oculaires*, *Thalami nervorum optico'rum*—c. *des Nerfs optiques*, *Thalami nervorum optico'rum*—c. *Optiques*, *Thalami nervorum optico'rum*.

COUCHER, Decubitus.

COUCHING, see Cataract.

COUCINEA COCCINEA, *Cacoc'ia coccinea*.

COUDE, (L.) *cubitus*, (.) Elbow.

COU-DE-PIED, *Instep*.

COUDE-PIED, *Instep*.

COUDÉE, Cubitus.

COUDRIER, *Corylus avellana*.

COUENNE (F.), (from [L.] *Cutaneus*, itself from *Cutis*, 'skin.') *Cutis enill'a*. A term given to various parts of the human skin, which are prominent, hard, brownish, and often covered with rough hairs, so as to form patches not very unlike the skin of the hog. It is a malformation, occurring during intra-uterine existence, and remaining for life. See, also, *Nævus*, and *Corium phlogisticum*.

COUENNE INFLAMMATOIRE, *Corium phlogisticum*—c. *Pleurétique*, *Corium phlogisticum*.

COUF, *Tussis*.

COUGH, ([D.] *Kuch*), *Tussis*—c. *Bronchial*, *C. tubal*—c. *Root*, *Trillium latifolium*.

COUGH, *Tu'bal*, *Bron'chial cough*, *Tus'sis bron'chica*, (F.) *Toux tubaire ou bronchique*. Cough is so termed, when the succussion communicated by it to the parietes of the chest is very energetic, and a sensation is experienced by the ear as if a column of air was traversing tubes with solid walls, with much noise, strength, and rapidity.

COUGH, *Winter*, *Bronchitis*, (chronic.)

COUGHING, *Tussis*.

COUL, Abscess, Swelling.

COULER, ([L.] *colare*), To strain.

COULEURS COMPLEMENTAIRES, *Colours*, accidental.

COULEUVRE, *Coluber berus*.

COULEUVREE, *Bryonia alba*—c. *de Virginie*, *Aristolochia serpentaria*.

COULEUVRINE, *Polygonum bistorta*.

COULISSE (F.), (*couler*, 'to flow,') 'a groove, a gutter.' Anatomists designate, by this name, every deep groove or channel in a bone, in which a tendon plays;—such as the *Coulisse Bicipitale* or *Bicipital Groove* of the Humerus.

COULISSE BICIPITALE, *Bicipital groove*—c. *Humérole*, *Bicipital groove*.

COULOIR, ([L.] *colare*, 'to strain,') *Colatorium*.

COUM, *Colchicum autumnale*.

COUMAROUNA ODORATA, *Dipterix odorata*.

COUNTENANCE, *Vultus*, *Vol'tus*, *Opsis*, *Vis'age*. (F.) *Figure*, *Visage*: (*cum*, and *teneo*, 'I hold.') The form of the face. The system of the features.

COUNTEREXTENSION, *Contraexten'sio*, *Antit'asis*, (F.) *Contre-extension*, (*contra-extendere*, 'to extend in a contrary direction,') It consists in retaining firmly and immovably the upper part of a limb, whilst extension is practised on the lower, in cases of fracture or luxation.

COUNTERINDICATION, *Contraindica'tio*, *Antendeix'is*, *Antendix'is*, *Antideix'is*. An indication contrary to another. (F.) *Contre-indication*. Any circumstance, which acts as an obstacle to the employment of such therapeutical means as seem to be indicated by other circumstances.

COUNTERIRRITANT, see *Counterirritation*—c. *Granville's*, (*Lotion*), *Granville's Counterirritant*.

COUNTERIRRITATION, *Contraindica'tio*. An irritation, excited in a part of the body, with the view of relieving one existing in another part. The remedies used for this purpose are called *Counterirritants*, and form a most valuable class of remedial agents. See *Derivation*.

COUNTEROPENING, *Contra-apertura*.

COUNTERPOISON, *Antidote*.

COUNTERSTROKE, *Contraffissura*.

COUNTING, *METHOD OF*, *Method*, numerical.

COUP, (from low L., *Colpus*; itself abridged from *κολαπες*, 'a cuff.') Blow—c. *de Feu*, *Wound*, gunshot.

COUP DE MAÎTRE, *Tour de Maître* (F.). 'A

master stroke or performance.' Applied to a mode of introducing the sound or catheter into the bladder, which consists in first passing it with its convexity towards the abdomen of the patient, and giving it a half turn towards the right groin, when its extremity has reached the root of the penis under the symphysis pubis. There is no advantage in this mode of introduction.

COUP DE SANG, (F.) 'Blood-stroke.' A common term, used by some physicians in France, to designate the loss of sensation and motion, which results from hemorrhage in the brain, or from simple congestion in the vessels of that organ. See *Apoplexy*. Some authors have comprehended, under the same denomination, different hemorrhages, which occur in the areolar texture of the face, lungs, skin, &c.

COUP DE SOLEIL, (F.) *Siri'asis*, *Seiri'asis*, *Siri'asis Egypti'aca*, *le'tus sol'is seu sol'a'ris*, *Insola'tio*, *Encephali'tis insola'tio'nis*, *Phreni'tis calen'tu'ra*, *Sunstroke*, *Stroke of the sun*, *Egypt'tian Star'stroke* or *Sun'stroke*. Any affection produced by the action of the sun on some region of the body;—head, hands, arms, &c. A presumed effect of exposing the naked head to the sun is inflammation [?] of the brain or its meninges, *Helien-cephali'tis*, which Sauvages calls *Cereus ab Insola'tio'ne*, *Morbus solstitia'lis*. *Coup de soleil* has, also, been called *heat apoplexy*; although the morbid phenomena observable after death are generally not confirmatory of either inflammation or apoplexy. Like effects have, also, been ascribed to the moon, *Selenoplex'ia*, *Selenople'gē*, *Moonstroke* and to the stars, *Starstroke*.

COUP DE VENT. 'A wind stroke.' The aggregate of morbid phenomena ascribed to exposure to a blast of air.

COUPE-BRIDE, *Kiotome*.

COUPER ROSE, *Copperas*, *Gutta rosea*—c. *Blanc*, *Zinci sulphas*—c. *Bleu*, *Cupri sulphas*.

COUPURE, (*couper*, 'to cut,') *Cut*, *Wound* (incised.)

COURALIMUM, (*κουραλιον*), *Coral*.

COURANTE, (*courir*, 'to run,') *Diarrhœa*.

COURAP. A distemper, very common in India, in which there is a perpetual itching of the surface, and eruption. It is of an herpetic character, and appears chiefly on the axilla, groins, breast, and face.

COURBAIL, see *Anime*.

COURBATURE (F.), (from *courbe*, 'a curve,' itself from *curvare*, *curvatum*, 'to bend,') *Acer'ba lassitu'do*, *Vi'olent las'situde*, *Fatigue*. An indisposition, characterized by a sensation, as if the limbs were bruised; general feeling of debility; extreme lassitude; and, sometimes, slight fever. It appears immediately after severe exercise, but sometimes not till the next day. Rest removes it.

COURBURE, (from *curvus*, 'crooked,') *Curvature*.

COURGE, ([L.] *Curvus*, 'curved,') *Cucurbita pepo*.

COURMI or **CURMI**, (*کورمی*, *کورمی*.) A fermented liquor, made from barley. A kind of ale or beer.

COURO-MOELLI. An Indian tree, the bark of which is said to be antivenomous.

COURON'DI, *Couron'do*. An evergreen tree of India, the juice of which, mixed with warm whey, is said to cure dysentery.

COURONNE, (from *corona*,) *Crown*—c. *Ciliare*, *Guadranné conal*—c. *du Dent*, *Crown of a tooth*—c. *du Gland*, *Crown of the glans*—c. *de Moine*, *Leontodon taraxacum*.

COURONNE RAYONNANTE (F.) A term given by Reil to the fan-shaped terminations of the cruri cerebri in the brain.

COURONNE DE SAINT-JEAN, *Artemisia vulgaris*—c. *du Trépan*, *Crown of the trepan*—c. *de Vénus*, *Crown of Venus*.

COURONNEMENT, (F.), *Courona*. A vulgar expression, sometimes used to designate the circular ring, which the os uteri forms around the head of the child at a certain period of accouchement—the head seeming to be surrounded, as it were, by a crown. The head is then said to be *en couronnement*.

COURS DE VENTRE, (*currere, cursum*, 'to run.') Diarrhœa.

COURSE, Running.

COURSES, Menses.

COURT, ((*L. curtus*.) Short.

COURT HALEINE, ('short breath.') Dyspnœa.

COUSIN, *Culex*.

COUSSINET, (from *culcita*, 'a cushion.') Pad.

COUSSO, *Hagenia Abyssinica*.

COUTEAU, (from *cutellus*, dim. of *culter*, 'a knife.') Knife—*c. à Amputation*, Knife, amputation—*c. à Cataracte*, Knife, cataract—*c. Courbe*, Knife, crooked—*c. à Crochet*, Knife, crotchet—*c. Désarticulateur*, Knife, double-edged—*c. Droit*, Knife, amputation—*c. de Feu*, *Cutellaire*—*c. Interoesneux*, Knife, double-edged—*c. Lenticulaire*, Knife, lenticular—*c. Lithotome*, Knife, lithotomy—*c. en Serpette*, Knife *en serpette*—*c. Symphysien*, see Symphyseotomy—*c. à deux Tranchans*, Knife, double-edged.

COUTOUB'EA AL'BA, *Er'acum spica'tum*, *Pit'crium spica'tum*. A plant of Guiana, which is very bitter, and is considered, in the country, to be emmenagogue, anthelmintic, and antidyseptic.

COUTURES PAR LA PETITE VEROLE, see Cicatrix.

COUTURIER, (from (F.) *coudre*, 'to sew.') Sartorius.

COUVRECHEF (F.), *Cucul'lus, Fascia'tio cuculla'ta, Scepaster'ium, Scepas'trum*. A bandage, applied to the head for retaining certain dressings, &c., *in situ*. The French surgeons distinguish two kinds: 1. The *Grand Couvrecchef* ou *Servette en carré* (*Cucul'lus ma'ior*), which is formed of a napkin or large square compress; and, 2. The *Petit Couvrecchef* ou *Mouchoir en triangle* (*Cucul'lus mi'ior*), formed of a napkin or other square compress, folded from one angle to the other diagonally opposite to it. See, also, Bandage, Galen's.

COUVERCLE, ((*L. Cooperculum*, (*L. Coperchio*, 'a cover.') See Crucible.

COVE. CLIMATE OF. On the northern side of Cork Harbour, in Ireland, is the Island of Cove. The town of Cove is on the southern acclivity of a hill running from east to west. It is, consequently, sheltered from the north winds, and receives the full force of the sun. It is one of the mildest climates in Great Britain, and corresponds in its influence on disease with the south-west of England. It is well adapted as a winter retreat for the phthisical.

CO'VOLAM, *Crata'eva* seu *Æ'glë Mar'melos*, *Fero'nia pelli'cida*, *Bilœa, Mahura, Bael, Bela*. The bark of this East India tree, *Order, Aurantiaceæ*, is tonic; the unripe fruit is astringent, and has been used in Europe in diarrhœa and dysentery (see *Extractum Belæ liquidum*); but, when ripe, is delicious.

COWANIA STANSBURIA'NA, (Mex.) *Alon'senel*, (called after Mr. James Cowan, an English merchant and botanist, and Captain Stansbury, of the United States army.) A plant of the *Nat. Ord. Rosaceæ*, which grows extensively in the vicinity of Salt Lake, and is a general astringent and local styptic.

COWBANE, *Cicuta aquatica*—*c. Spotted, Cicuta maculata*.

COWBERRY, *Vaccinium vitis idæa*.

COWCUMBER, *Cucumis sativus*.

COWDEL, *Caudle*.

COWHAGE, *Delichos pruriens*.

COWITCH, *Delichos pruriens*.

COWLADY, *Coccinella septempunctata*.

COW'LUCK. A tuft of hair on the head, pressed out of its place, and seeming as if it had been licked by a cow; giving rise to pain, when pressed into its natural position.

COWMUMBLE, *Heracleum spondylium*.

COWPARSNAP, *Heracleum lanatum*, *H. spondylium*.

COW'PER'S GLANDS, *Accessory glands*, (F. *Glandes accessoires* ou de *Couper*, *Prostates inférieures* ou *Petites prostates*. (Called after their discoverer.) Two small ovoid groups of mucous, reddish follicles, situate behind the bulb of the urethra, before the prostate, the excretory ducts of which open into the bulbous portion of the urethra. They are also called *Gland'ulæ antepros'tatæ*, and *G. pros'tatæ infério'res*. The terms *anterior pros'tatæ*, and *antepros'tat'ic gland*, are, generally, however, given to a third glandular body, occasionally found in front of, and between, Cowper's glands.

COW'PER'S GLANDS IN THE FEMALE, *Gland'ulæ Cowperi, Glands of Duvern'ey* or of *Bartholin*, *Pro'stata muli'ebris* seu *Bartholi'ni*, (F.) *Glandes de Cowper* ou de *Méry*, *G. vulvaires conglomérées* ou *vulvo-vaginales, Corps folliculaires vaginaux*, are situate at each side of the entrance of the vagina, beneath the skin covering the posterior or inferior part of the labia. They are rounded, but elongated, flat, and bean-shaped; their long diameter, varying from five to ten lines; their transverse, from two and a half to four and a quarter; and their thickness from two and a quarter to three lines. Like Cowper's glands in the male, they are not invariably present. The secretion from them is a thick, tenacious, grayish-white fluid, which is emitted in great quantity during sexual intercourse; and is probably the fluid supposed, of old, to be female sperm.

COWPOCK, *Vaccina*.

COWPOX, *Vaccina*—*c. Inoculation. Vaccination*—*c. Itch*, see *Itch*, cowpox—*c. Vesicle*, see *Vaccination*.

COWRAP, *Impetigo*.

COWSLIP, *Caltha palustris*, *Primula veris*—*c. Jerusalem, Pulmonaria*.

COWSTRIPINGS, *Primula veris*.

COWSTROPPLE, *Primula veris*.

COWTHWORT, *Matricaria*.

COWTREE, *CEYLON*, *Gymneura lactiferum*.

COWTREE OF SOUTH AMERICA, *Galactodendron utile*.

COWWEED, *Chærophylllum sylvestre*.

COXA, *Haunch*.

COXÆLUVIUM, (*coxa*, and *luere*, 'to wash.') Bath, hip.

COXAGRA, (*coxa*, and *aypa*, 'a seizure.') Neuralgia femoro-poplitea.

COXAL'GIA, *Merocoxal'gia, Oosphyal'gia, Oosphyalge'ma*. A word of hybrid origin. (*coxa*, and *algos*, 'pain.') Pain in the hip. A sign of rheumatic or other inflammation—*Coxi'tis*—or of some disease about the hip-joint. See *Neuralgia femoro-poplitea*, and *Coxarum morbus*.

COXARTHRITIS, (*coxo*, and *arthrit'is*, 'inflammation of a joint.') Coxitis.

COXARTHROACE, (*coxa*, *arthron*, 'a joint,' and *kakos*, 'evil.') Coxarum morbus.

COXA'RUM MOR'BUS, *Coxen'dicus* seu *Coxa'rius mor'bus*, *Arthro'acë coxa'r'um, Coxarthro'acë, Oospharthro'acë, Coxal'gia* (of some), *Hip Disease*. A serofulous caries, and often spontaneous luxation of the head of the os femoris, occasioning permanent shortening of the limb, and not unfrequently hectic and death—*Is'chiophthi'isis*.

COXEN'DIX, gen. *Coxen'dicis*, *Haunch*. This word has been used synonymously with *ischium*; and anatomists have also applied it to the *ilia*, *Ossa Coxen'dicis*.

COXITIS, *Coxarthriti* is a hybrid term, (*coxa*, and *itis*.) Inflammation of the hip joint.

COXOFEMORAL, *Coxofemorialis*. Belonging to the coxal bone or ileum, and to the femoris. *Hiofemor*al has the same signification.

COXOFEMORAL ARTICULATION, *Hiofemor*al A., *Articulatio coxae*, (F.) *Articulation de la Hanche*, *Hipjoint*, (Prov.) *Hunch*, *Lock'ing*, *Lonk*. The head of the femur and the articular cavity are covered by a *diarthrodial cartilage* and *synovial capsule*; and by a very strong *capsular or articular ligament* attached to the circumference of the cotyloid cavity, and to the neck of the femur. There is, also, a *round or interarticular ligament*, passing from the inferior notch of the cotyloid cavity to the rough depression at the top of the caput femoris; and a *cotyloid ligament*—a sort of thick, cartilaginous ring, situate on the edge of the cotyloid cavity, and converting the inferior notch into a foramen. These are the great means of union in this extensive articulation.

CRAB, (*καράβος*, *Car'abus*, Sax. *crabba*). *As'tacus fluviatilis*, *Can'cer*, *Gam'marus*, *Gam'barus*, *Cim'marus*, (F.) *Crabbe*, *Écrevisse*, *Ecrevice*. *Class*, Crustacea. A shell-fish much used as an article of diet. Like other shell-fish, it is apt to disagree, and excite urticaria, &c.

CRABE, Crabyaws.

CRABLOUSE, *Pediculus pubis*.

CRABS' EYES, *Cancerorum chelæ* — c. Stones, *Cancerorum chelæ*.

CRABYAWS, (F.) *Crabe*. A name in the West Indies for a kind of ulcer on the soles of the feet, with edges so hard, that they are difficult to cut.

CRACHAT, (*cracher*, 'to spit.') Sputum.

CRACHEMENT, Excretion—c. *de Pus*, Vomica, Phthisis pulmonalis — c. *de Sang*, Hæmoptysis.

CRACHOTEMENT (F.). The frequent spitting of a small quantity of saliva.

CRACKED POT SOUND, *Bruit de pot fêlé*.

CRACKLING, (onomatopœia.) *Craquement*, Crepitant, Crepitation—c. of Leather, see *Craquement de cuir*.

CRACOW GROATS, (from Cracow.) Semolina. **CRADLE**, (Sax. *crædel*,) *Ar'culus*, *Solen*, (F.) *Arceau* ou *Archet*. A semicircle of thin wood, or strips of wood, used for preventing the contact of the bed-clothes in wounds, fractures, &c. An ordinance of the Grand Duke of Tuscany forbade mothers and nurses to sleep with a child near them, unless it was placed under a solid cradle of this kind, in order that no accident might arise from *overlay'ing*.

CRÆPALÈ, (*κραπαλη*, from *κρα*, 'the head,' and *παλλω*, 'I agitate;') *Cræ'pula*, 'drunkenness, surfeit.' A derangement of the functions of the brain, &c., produced by wine or any other fermented liquor.

CRÆPALIA TEMULENTA, *Lolium temulentum*.

CRAGBANE (Sc.), (from (G.) *Kragen*, 'collar,' and *Bein*, 'bone.') *Clavicle*.

CRATÈ, Creta—c. *Barotique*, *Baryta*, carbonate of—c. *Magnésienne*, *Magnésie carbonas*—c. *de Plomb*, *Plumbi subcarbonas*.

CRATSEY, *Ranunculus acris*.

CRAMA, (*κραμα*, from *κραω*, 'I mix.') A mixture of any kind. Dilute wine.

CRAMBE, (*κραμβη*,) Cabbage.

CRAMBION, (*κραμβιον*,) A decoction of cabbage.

CRAMERIA, *Krameria*.

CRAMP, ([G.] *krampfen*, 'to contract,') *Cramp'us*, *Entu'sia agstrom'ma*, *Tet'anus dolorif'icinus*, *Myo'spas'mus*, *Spas'mus muscula'ris*, *Myal'gia*, *Rhe'gè*, *Rheg'ma*, *Rheg'mus*, (F.) *Crampe*. A sudden, involuntary, and highly painful contraction of a muscle or muscles. It is most frequently experienced in the lower extremities, and is a

common symptom of certain affections—as of *Colica Pictonum* and *Cholera Morbus*. Friction and compression of the limb, by means of a ligature applied around it above the muscles affected, will usually remove the spasm.

CRAMP RINGS, Rings of different form, solemnly consecrated, which were given out by the English kings, of old, for the prevention or cure of cramp.

CRAMP, SCRIVENERS', Cramp, writers'.

CRAMP OF THE STOMACH, (F.) *Crampe de l'Estomac*. A sudden, violent, and most painful affection of the stomach, with sense of constriction in the epigastrium. It seems to be the effect of the spasmodic contraction of the muscular coat of the stomach, and requires the use of the most powerful stimulants and antispasmodics, of which opium is the best.

CRAMP, WRITERS', Graphospas'mus, *Mogigraph'ia*, *Scriveners' Cramp*, *Scribe pal'sy*, *Stammering of the Fingers*, (F.) *Crampe des Écrivains*. A condition of the fingers, in which they are unable to hold the pen, or in which one or more of the muscles of the fingers are irregularly and irresistibly contracted.

CRAMPE DES ÉCRIVAINS, Cramp, writers'.

CRAMPE DE L'ESTOMAC, Cramp of the stomach.

CRAN DE BRETAGNE, *Cochlearia armoracia*.

CRANBERRY, *Vaccinium oxycoccos*—c. Alpine, *Vaccinium vitis idæa*—c. American, common, see *Vaccinium oxycoccos*—c. Upland, *Arbutus uva ursi*.

CRANE, *Cranium*—c. *Humain*, *Cranium humanum*.

CRANE'S BILL, *Geranium*—c. Spotted, *Geranium maculatum*—c. Stinking, *Geranium Robertianum*.

CRANIAL, *Cranialis*, *Crania'nus*, (F.) *Crânien*; (from *cranium*.) Relating or appertaining to the cranium—as '*cranial nerves*.'

CRÂNIEN, Cranial.

CRANIO, in composition, the cranium.

CRANTOCELE, (*cranio*, and *κληη*, 'a tumor,') *Encephalocele*.

CRANIOTOMY, *Cranioclas'mus*, (*cranio*, and *κλαω*, 'to break,') The breaking up of the child's head, as by the craniotomy forceps.

CRANIOHÆMATONCUS, (*cranio*, *αίμα*, 'blood,' and *ογκος*, 'a swelling,') *Cephalæmatoma*.

CRANIOL'OGY, *Craniology*, *Cranilog'ia*, *Craniomancy*, *Phrenol'ogy*, *Cranioscop'ia*, *C. Gallia'na*, *Encephaloscop'ia*, *Doctri'na Gallia'na*. (F.) *Cranilogie*, *Cranioscopie*. Words introduced, since Gall, into medical language. (The first two terms are respectively derived from *cranio*, *λογος*, 'a discourse,' and *σκοπειν*, 'to examine,') They signify a description, or simply an examination, of the different parts of the external surface of the cranium, in order to deduce from it a knowledge of the different intellectual and moral dispositions. Strictly speaking, it is by *Craniology* or *Organology* that we acquire a knowledge of *Craniology*, *Organology*, or *Cranology*, as it has been variously termed. These words are generally, however, used in the same sense. The cranium being moulded to the brain, there are as many prominences on the bone as there are projections at the surface of the brain. According to Gall, each projection, which he calls an *organ*, is the seat of a particular intellectual or moral faculty, and all persons endowed with the same faculty, have, at the same part of the brain, a prominence, which is indicated, externally, by a bump or projection in the bony case. The *System* of Gall is made to comprise 27 prominences, which answer to 27 primary faculties.

A fundamental principle with the *Cranilogists* or *Phrenologists* is,—that the brain does not act as a single organ, but that it consists of a plurality of organs; but, were we able to admit this, the assignment of the seat of different faculties could not but be considered premature.

CRANIOMANCY, (*cranio*, and *μαντεία*, 'divination.') Craniology.

CRANIOMETRY, *Cranimet'ria*; (from *κρανιον*, 'the cranium,' and *μετρον*, 'measure.') Measurement of the skull. A form of compasses used for this purpose is called a *craniometer*, (F.) *Cranimètre*.

CRANION, (*κρανιον*.) Cranium.

CRANIOPATHIA, (*cranio*, and *παθος*, 'disease.') Disease of the cranium.

CRANIOPLASTY, *Cranioplas'ticē*, (*cranio*, and *πλασσω*, 'I form.') An operation for the restoration of the soft parts and bones of the cranium.

CRANIOSCOPY, (*cranio*, and *σκοπεω*, 'I view.') Craniology.

CRANIOSPINAL, Cephalo-spinal.

CRANIOSPINAL AXIS, see Encephalon.

CRANIOSTENOSIS, (*cranio*, and *στενω*, 'contracted.') See Microcephalus.

CRANIO TABES, (*cranio*, and *tabes*, 'wasting.') A softening of the bones of the cranium, and a consequent thinness of those bones,—as the occiput, (*soft occiput*),—which are much exposed to pressure. It is considered to be a variety of rickets.

CRANIOTOMY, *Craniotom'ia*, (F.) *Crâniotomie*; (*cranio*, and *τομή*, 'incision.') The operation of opening the head in parturition.

CRANIOTOMY FORCEPS. An instrument, resembling the lithotomy forceps, for laying hold of and breaking down the bones of the head in parturition.

CRANIOTRYPESIS, (*cranio*, and *τρύπω*, 'I perforate.') Trepanning.

CRANIA, (*κρανιον*.) *Cra'nion*, *Cra'num*, *Cal'varia*, *Scap'hion*, *Con'chus*, *Cal'va*, *O'la cap'itis*, *The'ca cer'e'br*i, *Brain'pan*, (Old Eng.) *Pan'nikell*, (Prov.) *Harn'pan*, the skull; (from *κρανος*, 'a helmet,' or from *κρανον*, 'head.') (F.) *Crâne*. The collection of bones which form the case for lodging the brain and its membranes, as well as their vessels, and some of the nerves. These bones are eight in number,—the *frontal*, *occipital*, *two parietal*, *two temporal*, the *sphenoid*, and *ethmoid*. Besides these, there might be considered, as belonging to the cranium, the *cornua sphenoidalia*, the bones of the ear, and the *ossa Wormiana*. Dr. Prichard has described the primitive forms of the skull according to the width of the *Breg'ma*, or space between the parietal bones: 1. The *Ste'nobreg'mate*, (*στενος*, 'narrow,') or Ethiopian variety. 2. The *Me'sobreg'mate*, (*μεσος*, 'middle,') or Caucasian variety; and 3. The *Pla'tybreg'mate*, (*πλατος*, 'broad,') or Mongolian variety.

CRANIUM HUMA'NUM, *Hu'man Cra'nium* or *Skull*. (F.) *Crâne humain*. This was anciently much used in prescriptions, and was considered anti-epileptic, alexipharmic, antiloinic, &c.

CRA'NOMANCY, *Cranomanti'a*, (*κρανον*, 'the head,' and *μαντεία*, 'divination.') (F.) *Cranomancie*. The art of divining—from the inspection of the head or cranium—the moral dispositions and inclinations of individuals.

CRANSON, *Cochlearia armoracia*, C. officinalis.

CRANSSAC, MINERAL WATERS OF. *Cranssac* is a village in the department of Aveyron, six leagues from Rhodéz, which possesses acidulous chalybeate springs, that have been known for a long time.

CRANTER, gen. *Crante'ria*, (*κραντηρ*, from *κρανιον*, 'to finish,' 'render perfect.') The *Dens Sapien'tie* has been so called.

CRANUM, (*κρανον*.) Cranium.

CRAPULA, Cræpale.

CRAQUEMENT (like *crack*, formed by onomatopœia) *DE CUIR NEUF* (F.). *Nouveau crâne*, 'crackling of new leather.' A sound like the crackling of new leather, sometimes heard on examining the heart with the stethoscope. It is symptomatic of pericarditis.

CRAQUEMENT PULMONAIRE (F.), 'Pulmonary crackling.' This *bruit* or sound consists of a succession of small cracklings, heard during inspiration, and almost always at the top of the lung. It is heard at the commencement of phthisis.

CRA'SIS, (*κρασις*, from *κραω*, 'to mix.') A mixture of the constituents of a fluid: as the *crasis* of the blood, humours, &c. The word has also been employed in a more extensive signification, as synonymous with *Constitution*, *Temperament*, &c.

CRASPEDON, (*κρασπεδον*.) Staphyloedema.

CRASSAMENTUM, (*crassum*, 'thick.') The thick part or deposit of any fluid. It is particularly applied to the clot of the blood.

CRASSAMENTUM SANGUINIS, see Blood.

CRASSE (F.), (from (L.) *crassus*, 'thick' [?]). 'Dirt or impurity.' A sort of layer or *enduit*, which covers the skin, where cleanliness is not attended to; and which is sometimes the result of the cutaneous exhalation; at others, of extraneous matter adhering to the surface of the body.

CRASSE'NA. A term by which Paracelsus designated the saline, putrescent, and corrosive principles, which, he thought, gave rise to ulcers and tumours of different kinds.

CRASS'ULA, (dim. of *crassus*, 'thick,' owing to the thickness of its leaves.) Sedum telephium.

CRASSULA ARBORESCENS, see Cotyledon orbiculata—c. Portulacaceæ, c. Tetragona.

CRASSULA TETRAGO'NA. A South African succulent plant, *Ord.* Crassulaceæ, which is somewhat astringent. Boiled in milk it is used in diarrhœa. *Crassula portulacaceæ* is said to be used in similar cases.

CRASSUS, ('thick,') Corpulent.

CRATE'GUS A'RIA, (*κραταιγος*, from *κρατος*, 'strength,' owing to the hardness of the wood.) *Mes'pilus*, *M. A'ria*, *Aria* seu *Sorbus*, *S. Aria* seu *Alpi'na*, *White Beantree*, (F.) *Alisier blanc*, *Alouche*, *Alouchier*. The fruit, which is of the size of a small pear, is slightly astringent, and somewhat agreeable to the taste. It has been employed in diarrhœa.

CRATEVA MARMELOS, (after *Crateva*, an ancient rhizotomist,) *Covolan*.

CRATERAUCH'EN, (*κρατερος*, 'strong,' and *αυχνη*, 'the neck.') One with a strong neck: as well as a strong neck itself.

CRATEVÆSIUM, *Sisymbrium nasturtium*.

CRATUS, (*κρατος*.) Force.

CRAUTE, MINERAL WATERS OF. *Craute* is a village five leagues from Autun, in France, where are mineral springs, containing sulphhydrate of magnesia, chloride of sodium, chloride of lime, sulphate of lime, and carbonate of magnesia.

CRAVAT', (F.) *Cravate*, (said to be derived from a regiment of Croats in the service of France in the 17th century.) A form of necktie or neckcloth. Also, a name given to different bandages not limited to the neck.

CRAVATE SUISSE (F.), 'Swiss cravat.' A band of the longitudinal or superficial layer of the muscular fibres of the stomach, along the lesser curvature of the organ, the shape of which it assists in preserving;—so called, on account of a fancied resemblance to a Swiss cravat.

CRAW-CRAW. A species of itch, common on the African coast.

CRAYEUX, (from *craie*, 'chalk,') Cretaceous.

CRAYON NOIR, (from *craie*, 'chalk,') Graphites.

CRAZINESS, Insanity.

CRAZY, ([F.] *érasé*, 'crushed,') Insane, Pain.

CREA, Shin.

CREAM, *Cre'mor seu Flos lac'tis*, (Prov.)

Ream, (F.) *Crème*. A thick, unctuous matter, of a yellowish-white colour, and sweet, agreeable taste, which rises to the surface of milk, if kept at rest. It composed of butter, serum, and casein.

Crème is a name given in France to *bouillies* prepared with farina of different kinds; as the *Cream of Rice*, (F.) *Crème de Riz*; *C. of Barley*, (F.) *C. d'Orge*, &c. The same term, and '*medic'i-ni-cream*,' (F.) *Crème médicinale*, are used by M. Béral for a preparation, resulting from the union of *yolk of egg* and *sugar* with *milk*, with or without the addition of medicinal agents.

CREAM, Rigor—c. Cold, Ceratum Galeni, Unguentum Aquæ rosæ—c. of Tartar, Potassæ supertartaras.

CREAS, (*κρεας*,) Flesh; also, Rubeola, (Prov.)

CRE'ASOTE, *Cre'osote, Creaso'tum, Kre'asote, Creaso'ton, Creazo'ton*; *Oxyhydrocarbure'tum ex oleo pyroac'yl'ico para'tum* (Ph. L.), (*κρεας*, 'flesh,' and *σωτηρ*, 'a preserver,') A colourless, oily, neutral liquid of a strong, characteristic odour, and acid, burning taste, discovered by Reichenbach. It is obtained from tar by distillation, and appears to be the active, antiseptic and medicinal agent in tar-water and crude pyrolygneous acid. It is a colourless transparent fluid, of a penetrating and disagreeable odour, and is freely soluble in alcohol and acetic acid. Its taste is bitter. S. g. 1.046. It coagulates albumen, whence its hemostatic power. It is a most powerful antiseptic; and has been largely administered in hemorrhages both internally and externally. As an external application, it has been used in burns, ulcers, especially those of a sloughing character, chronic cutaneous affections, and has been applied to tapeworm when protruded. Internally, it has been administered in phthisis, vomiting, diabetes mellitus, nervous diseases, chronic glands, &c. The dose internally is one or two drops. Externally, it is sometimes applied pure; at others, diluted, and commonly with water, (f℥ss to f℥vj;) or in the form of ointment, (f℥ss to ℥i) of cerate.)

Aqua Creaso'ti, Kre'asote water (Ph. U. S., 1863), is made of *creasote*, f℥j; *distilled water*, Oj.

CREASOTE PLANT, *Larrea Mexicana*.

CRE'ATIN, *Cre'atine, Creat'i'na, Kre'atin*, (*κρεας*, gen. *κρεατος*, 'flesh,') A nitrogenized crystallizable neutral substance, obtained by the agency of water and heat in making broths and soups. It does not combine either with acids or alkalis.

By the action of strong acids, creatine is converted into *Cre'atinin* or *Kre'atinin*, a substance which has a strong alkaline reaction, and forms crystallizable salts with acids. It preëxists to a small extent in the juice of flesh; and is found in conjunction with creatin in urine.

CREATININ, see Creatin.

CREATOPHAGUS, (*κρεας*, gen. *κρεατος*, 'flesh,' and *φαγω*, 'I eat,') Carnivorous.

CREAZOTON, *Creasote*.

CREEPER, TRUMPET, *Tecoma radicans*—c. Virginia, *Ampelopsis quinquefolia*, *Tecoma radicans*.

CREEPERS, Fidgets.

CRE'MASON, Pyrosis.

CREMASTER, *Crema'stes, Suspendi'culum, Eleva'tor Testic'uli, Mus'culus Testis seu Testicu'lus*, (*κρεαστηρ*, 'that which suspends,' from *κρεαω*, 'I suspend,') The Greeks designated, by this term, the spermatic cord, or all the parts by which the testicle is, as it were, suspended; but, since Vesalius, it has received its present limited meaning. The cremaster is a very thin muscular fascia—sometimes hardly perceptible—which de-

taches itself from the internal oblique muscle; passes through the abdominal ring, and vanishes around the tunica vaginalis; serving to draw up the testicle, and to move it slightly. It has been, also, called *Tu'nica Erythro'i'des* and *Suspensoi-rium Testis*.

CREMASTERIC or CREMASTER'IC, *Cre-master'icus*. Appertaining or relating to the cremaster,—as '*cremasteric artery*,'—a branch of the epigastric.

CREMASTES, (*κρεαστης*,) Cremaster.

CRÈME, Cream—c. *Médicinal*, see Cream—c. *de Soufre*, Sulphur præcipitatum—c. *de Tartre*, Potassæ supertartaras.

CREMER. The common name for a disease frequent in Hungary, which is produced by excess in eating and drinking.

CREMNOCE'LE, *Cre'mnoccele, (cremnoi, and κηλη, 'a rupture,') Hernia of the labia pudendi.*

CREM'NOI, (pl. of *κρηνος*, 'a steep bank,') The lips of an ulcer. The *Labia puden'di*.

CREMNON'CUS, (*cremnoi, and ογκος, 'a tumour,')* A swelling of the labia pudendi.

CREMOR LACTIS, gen. *Cre'moris*, Cream—c. Tartari, Potassæ supertartaras—c. Urinæ, see Urine.

CRE'NA, ('a notch,') *Crenatu'ra, Dentic'ulus*, (F.) *Crênelure*. In the plural, the small teeth or projections in the bones of the cranium, by means of which an accurate junction is formed at the sutures. *Crênelure* has also been used in surgery, for the gutter or groove in certain instruments, which is generally intended to secure the passage of cutting instruments; the groove, for example, of a director.

CRENATURA, Crena.

CRÊNELURE, Crena.

CRE'OLE, (S.) *Criollo*; ([S.] *criar*, 'to create or foster,') A native of America, or of the West Indies, whose ancestors emigrated from the Old World, or from Africa. Hence there may be black as well as white creoles. In the South, it is used in the same sense as *native*.

CREOPHAGUS, (*κρεας*, 'flesh,' and *φαγω*, 'I eat,') Carnivorous.

CREOSOTE, *Creasote*.

CREPALIA TEMULENTA, *Lolium temulentum*.

CREPANELLA, *Plumbago Europæa*.

CREPA'TIO, *Crepatu'ra, (crepare, crepatum, 'to make a noise,')* The act of bursting any seed by ebullition. *Coque ad crepatu'ram*, 'Boil till it bursts.'

CREPATURA, Hernia.

CREPIDINES PALPEBRARUM, (*crepido*, 'a border,') see Tarsus.

CREP'ITANT, *Crep'itating, Crep'itans*, (from *crepitare*, 'to crackle,') Crackling.

Râle Crépitant Sec à Grosses Bulles, 'dry crackling noise with formation of large bubbles,' is heard in pulmonary emphysema, when the distension becomes greater and greater, and is followed by rupture of the vesicles. The air forcing itself a passage in the interlobular areolar tissue, gives rise to this *Râle* during inspiration.

Râle Crépitant Sec de Laënnec, R. vésiculaire. The *crep'itant râ'tle, Crepita'tio vésicula'ris*, heard, during respiration, in severe pneumonia and in oedema of the lung; so termed on account of the analogy between the sound and that occasioned by pressing a healthy lung between the fingers. It resembles the sound produced by rubbing slowly and firmly between the finger and thumb a lock of hair near the ear. The *Rhon'chus crep'itans re'dux*, (F.) *Râle crépitant re'dux*, is the sound heard in respiration coexistent with the resolution of pneumonia. It indicates the return of the cells to the puerious condition.

CREPITATIO, Crepitation—c. Vesicularis, Crepitation.

CREPITA'TION, *Crepita'tio*, *Crep'itus*, (*crepitare*, 'to make a noise.') *Crack'ling*. Crepitation or *crep'itus*, (F.) *Cliquetis*, has been used. in *Surgery*, to designate the noise occasioned by the friction of fractured bones, when the surgeon moves them in certain directions. When it cannot be heard at a distance, it may be detected by the immediate application of the ear, or by the use of the stethoscope. *Crepitus* or *crackling* is, likewise, met with in cases of gangrene, when air is effused into the areolar membrane—provided the part be carefully examined with the fingers. The same term is used for the cracking of the joints in health or disease.

CREPITUS, gen. *Crepitūs*, Crepitation, *Fart*—c. *Lupi*, *Lycoperdon*.

CRESCENTIA, (*creescere*, 'to grow,') Growth.

CRESCENTIA CUJETE; (called after Crescentio, an Italian writer on agriculture;) *Cujete*, *Narrow-leaved Calabash Tree*. Ord. Solanææ. The pulp of the fruit of this West India plant is acidulous. It is used in diarrhoea and headache; and, in syrup, for diseases of the chest.

CRESCENTILÆ, Waxing kernels.

CRESERA, (*κρησρα*, 'a sieve,') see Cribration.

CRISPINUS, *Oxyacantha Galeni*.

CRSS, BITTER, COMMON, (Sax. *cærrre*, 'a celeritate crescendi,' [?]) *Cardamine pratensis*—c. *Garden*, *Lepidium sativum*—c. *Indian*, *Tropæolum majus*—c. *Meadow*, *Cardamine pratensis*—c. *Penny*, *Thlaspi*—c. *Swines'*, *Cochlearia coronopus*—c. *Water*, *Sisymbrium nasturtium*—c. *Water*, *marsh*, *Nasturtium palustre*.

CRÉSSES, DOCK, Lapsana—c. *Sciatica*, *Lepidium Iberis*.

CRÉSSI, *Sisymbrium nasturtium*.

CRESSON, see Alleghany Mountain Springs.

CRESSON ('*cress*') **ALÉNOIS**, *Lepidium sativum*—c. *Élégant*, *Cardamine pratensis*—c. *de Fontaine*, *Sisymbrium nasturtium*—c. *des Indes*, *Tropæolum majus*—c. *des Jardins*, *Lepidium sativum*—c. *du Mexique*, *Tropæolum majus*—c. *de Para*, *Spilanthus oleraceus*—c. *des Près*, *Cardamine pratensis*.

CRÉST, ([F.] *Crête*,) *Crista*—c. of the *Ilium*, see *Crista* of the *Ilium*—c. of the *Pubis*, see *Crista* of the *Pubis*—c. of the *Tibia*, see *Crista* of the *Tibia*—c. of the *Urethra*, see *Crista urethralis*.

CRE'TA, (from *Creta*, the island,) *Carbo'nas cal'cis friabilis*, *Carbo'nas cal'cis creus*, *Glimomari-go*, *Melia Ter'ra*, *Calca'ria carbo'nica*, *Chalk*, *Carbonate of Lime*, (F.) *Craie*. Native friable carbonate of lime.

CRETA PREPARA'TA, *Carbo'nas Cal'cis præpara'tus*, *Prepared Chalk*. (Prepared by levigation.) Used externally, as an absorbent; internally, as an antacid. Dose, gr. x. to ʒj or more.

CRETA'CEOUS, *Creta'ceus*, (F.) *Crayeux*, *Cré-tacé*, (from *creta*, 'chalk,') Chalky. Containing, or relating to, or having the characters of, chalk; as 'cretaceous mixture, cretaceous tubercles.' (F.) *Tubercules cretaccés*.

CRÊTE, (old (F.) *Creste*,) *Crista*—c. *de Coq*, *Crista galli*—c. *de l'Ethmoïde*, *Crista galli*—c. *Uréthrale*, *Gallinaginis caput*.

CRETEFACT'ION, *Cretefact'io*, (*creta*, 'chalk,' and *facere*, *factum*, 'to make,') Formation of, or conversion into, chalk, as the conversion of tubercles into calcareous or cretaceous concretions; *Calcification*.

CRÊTES DE COQ, see *Crista*.

CRÉTIN. One affected with cretinism. The word is said to come from *Chrétien*, 'Christian,' because the Crétin, being in a state of idiocy, is incapable of committing any sin [?]; others derive it from the Romanesque *cretina*, 'a miserable creature.' See *Cagot*.

CRETINISM, *Cretinis'mus*, *Cret'inage*, *Cyrto'-sis Cretinis'mus*, *Kretinis'mus*, *Idiotia endem'ica*,

Idiotism'us endem'icus, *Fatu'itas convall'inus* seu *Alpicola'rum*, *Micrenceph'alon*, (F.) *Crétinisme*. An epidemic affection in the low, deep, narrow situations of the Valais; in the valley of Aost, Maurienne, a part of Switzerland, the Pyrenees, Tyrol, &c. It is a state of idiocy, commonly accompanied by an enormous goitre, and is often hereditary. The unfortunate crétin is little better than the animals around him. He rarely attains an advanced age. Like idiocy, cretinism has been divided into complete and incomplete.

CREUSET, (from *creux*, 'hollow,' see *Crucible*.) *Crucible*.

CREUX DE L'AISSSELLE, ('hollow of the axilla,') See *Axilla*—c. *de l'Estomac*, *Fossette du cœur*—c. *de la Main*, *Palm*.

CREUZNACH, MINERAL WATERS OF. The springs of Creuznach are in Germany. They contain iodine, bromine, and the chlorides of sodium and calcium.

CREUZOT, MINERAL WATERS OF. C. is a mountain to the north-east of Mount Cenis. Near it is a saline chalybeate spring.

CREVASSE, (F.) *Ri'nia*, (*crever*, 'to break or crack,') A crack, a cleft. The words *crevasse*, *gerçure*, *fissure*, and *rhagade* are often used synonymously for small longitudinal cracks or chaps of a more or less painful character. Sometimes, *crevasse* is employed to designate the solutions of continuity or ruptures, which supervene in distended parts, in the urinary passages, uterus, &c.: it is then synonymous with rupture.

CREWDS, *Rubeola*.

CREWEL, *Primula veris*.

CREWELS, ([F.] *Écrouelles*,) *Scrofula*.

CRI, *Cry*—c. *de Cuir*, *Bruit de Craquement*, *Craquement de cuir neuf*.

CRIBLÉ, (from *cribrum*, 'a sieve,') *Cribratus*.

CRIBLEUX, *Cribratus*.

CRIBRA, pl. of *cribrum*, see *Cribration*.

CRIBRA'TION, *Cribrat'io*, *Coscini's'mus*, (*cribrare*, *cribratum*, 'to sift,') *Sift'ing*. A pharmaceutical operation, which consists in separating the finer parts of drugs from the coarser. *Sieves*, *Drumsieves*, are used for this purpose. These were formerly called *Cri'bra*; *Coscina*. A sieve for separating the bran from meal was termed *Cre'sera*, *Aleurote'sis*, *Cribrum Pollina'rium*.

CRIBRA'TUS, *Cribro'sus*, (F.) *Criblé* ou *Cri-bleux*. Having holes like a sieve. *Lame criblée*, *Cribriform plate*, is the horizontal lamina of the ethmoid bone—so called because it is perforated like a sieve. Through the perforations, *Foramina cribro'sa*, the olfactory nerves pass.

La'mina Cribro'sa of Albinus; *Cribriform la-mel'la*, a circular spot, perforated with small holes, seen when the optic nerve is regarded from the inside, after removing the retina and choroid. From these holes the medullary matter may be expressed.

CRIB'RIFORM, *Cribriform'is*, (*cribrum*, 'a sieve,' and *forma*, 'form,') The ethmoid bone was formerly so called—*Os Cribriform'e*.

CRIBRIFORM COM'PRESS. A square piece of linen pierced with a number of holes. This is spread with cerate and applied to a suppurating surface, the holes being intended for the escape of the pus.

CRIBRIFORM FASCIA, see *Fascia cribriform*—c. *Lamella*, see *Cribratus*—c. *Plate of the Ethmoid*, see *Cribratus*.

CRIBROSUS, *Cribratus*.

CRIBRUM pl. *cribra*, 'a sieve,' see *Cribration*—c. *Pollinariun*, see *Cribration*.

CRICARYTENODES, *Crico-arytenoid*.

CRICELLA'SIA, *Cricella'sia*, (*κρικηλαστια*, (from *κρικος*, 'a circle,' and *ελαυνω*, 'I drive,') An exer

cise with the ancients, which appears to have been the same as the childish play of rolling the hoop.

CRICK IN THE NECK, (Prov.) *Shoot, Hod.* A painful rheumatic affection of the muscles of the neck, which causes a person to hold his head to one side in a characteristic manner.

CRICO, in composition, cricoid cartilage.

CRICOARYTENOID, *Cricoarytenoïdeus*, *Cricarytenoïdes*. Relating to the cricoid and arytenoid cartilages.

CRICOARYTENOID MUSCLE, *Cricoarytenoïdeus lateralis*, *Crico-latéro-arytenoïdien*. A muscle which proceeds from the lateral part of the superior edge of the cricoid cartilage to the outer and anterior part of the base of the arytenoid cartilage. It carries the arytenoid cartilage outward and forward.

CRICOARYTENOID, POSTERIOR; *Cricoarytenoïdeus posticus*, *Crico-crèti-arytenoïdien*, *Dilatateur postérieur du Larynx*. A small, triangular muscle, seated in the back part of the larynx. It extends from the prominent line, at the middle of the posterior surface of the cricoid cartilage, to the outer and posterior part of the base of the arytenoid cartilage.

CRICOARYTENOIDEUS SUPERIOR, of Winslow, forms part of the *Arytenoïdeus* of modern anatomists.

CRICOCRÉTIARYTÉNOÏDIEN, *Cricoarytenoïdeus, posterior*—*c. Latéro-arytenoïdien*, *Cricoarytenoid*.

CRICOPHARYNGEAL, *Cricopharyngeus*. Belonging to the cricoid cartilage and pharynx.

CRICOPHARYNGEUS muscles of Winslow, is a fleshy bundle, which forms part of the *Constrictor Pharyngis inferior*. He calls the other part *Thyropharyngeus*, and proposes to call the whole *Thyrocricopharyngeus*.

CRICOTHYREOIDES, *Cricothyroidean*.

CRICOTHYROIDEAN, *Cricothyroid*, *Cricothyroïdes*, *Cricothyroïdes*, *Cricothyreoides*. Belonging to the cricoid and thyroid cartilages.

CRICOTHYROID MEMBRANE is of a fibrous nature, and extends from the upper edge of the cricoid cartilage to the inferior edge of the thyroid.

CRICOTHYROID MUSCLE, *Cricothyroïdeus*, (F.) *Dilatateur antérieur du larynx*, is a small fleshy bundle, of a triangular shape, at the anterior and inferior part of the larynx. It extends from the outer surface of the cricoid cartilage to the lateral parts of the inferior margin of the thyroid cartilage; and its use is to approximate, anteriorly, the corresponding margins of the cricoid and thyroid cartilages; and thus to stretch the ligaments of the glottis, which it contracts by separating the thyroid cartilage from the arytenoid.

CRICOTHYROPHARYNGIEN, *Constrictor pharyngis*.

CRICOÏD, *Cricoi'deus*, *Cricoi'des*, *Cricoides*, *Annularis*, *Cymbalaris*, (κρικος, 'a ring,' and εϊδός, 'form,') Having the form of a ring. A name given to one of the cartilages of the larynx, *Cartilago innominata*, (F.) *Cartilage anonyme*, which is situate at its lower part, between the thyroid and first ring of the trachea. It is much higher behind than before.

CRICOIDES, *Annular*, *Cricoid*.

CRICOS, (κρικος,) *Ring*.

CRIMNO'DES, (κρινωιδης,) *Crimnoï'des*. Resembling meal or bran (*Crimmon*). Applied to urine which deposits a bran-like sediment, *Crimnoïdes Urina*.

CRIMNON, (κρινων,) *Farina*.

CRINALĒ, ('*crinis*,' 'hair,') An instrument formerly used to compress in cases of fistula lacrymalis. It has its name from the circumstance

of its having at one end a small cushion stuffed with hair.

CRINA'TUM. A species of fumigation, used by Paulus of Ægina. The roots of lilies entered into the process; hence its name,—(from κρινος, 'a lily,')

CRINES, pl. of *crinis*, *Hair*.

CRINIS, *Capillus*.

CRINLECRANKLE, *Wrinkle*.

CRINOM'YRON, (κρινωμυρον, from κρινος, 'a lily,' and μυρον, 'ointment,') *Unguentum liliaceum*, *U. Ægyptium album*, *U. Susianum*. An ointment, composed of lilies and some aromatic plants.

CRINON, (κρινων,) *Lilium candidum*.

CRINONES, (from *crinis*, 'hair,') See *Acne punctata*.

CRINOSSES, (from κρινω, 'I separate,' and οσις,) *Diaereses*.

CRIO'LO, *Creole*.

CRIO'MYX'US, (κριομυξος, from κριος, 'a ram,' and μυξα, 'mucus,') In antiquity, one who had much mucus flowing from his nasal fossæ, like the ram.

CRISIS, *Dia'crisis*, *Dijudicatio*, 'decision;' (κρίσις, from κρινω, 'I decide,') This word has been used in various acceptations. Some mean by *crisis* of a disease, when it augments or diminishes considerably, becomes transformed into another, or ceases entirely. Some have used the word to signify only the favourable changes which supervene in disease; others, for the change going on in the acme or violence of the disease. Others, again, have given this name only to a rapid and favourable change, joined to some copious evacuation or eruption; whilst others have applied the term to the symptoms that accompany such change, and not to the change itself;—thus including, under the same denomination, the critical phenomena and the *crisis*.

CRISPA'TION, *Crispatura*, (*crispare*, *crispatum*, 'to wrinkle,') A contraction or spasmodic constriction, which supervenes in certain parts, either spontaneously or by the influence of some morbid cause or therapeutical agent. The capillary vessels of a wound are, by the French, termed *crispés*, when, immediately after an operation, the blood does not flow from them. The skin is said to be *crispée*, when contracted, and the bulbs of the hair become more prominent. *Crispation* of the nerves is a slight convulsive motion of external or internal parts, much less than that which occurs in convulsion.

CRIS'TA, *Ant'be*, *Ant'bon*. A crest. (F.) *Crête*. A name given to several bony projections; also, to the clitoris.

The word *Crêtes*, (F.) *Cristae*, is also used, in France, for fimbriated excrescences, which form at the anus, and near the genital organs; and are commonly owing to the syphilitic virus. *Crêtes de Coq* are syphilitic excrescences, resembling, in form, the crest of the cock.

CRISTA GAL'LI; two Latin words, signifying the comb of a cock, (F.) *Crête de l'ethmoïde*, *C. de Coq*. A flat, triangular process, rising above the cribriform plate of the ethmoid bone, and resembling a cock's comb. It gives attachment to the anterior part of the falx cerebri.

CRISTA OR CREST OF THE IL'IUM is the superior margin of the ilium:—the *Crista* or *Crest* of the *Tibia*, the anterior edge, the shin; and the *Crista* or *Crest* of the *Pubis*, the posterior sharp edge on the upper surface of the bone. Winslow calls the nymphæ *Crista* of the *clitoris*; and Chaussier, the verumontanum, *Crista Urethralis*, *Crest* of the *Urethra*.

CRISTA INTERNA. Frontal Spine—*c. Sphenoidalis*, *Sphenoidal spine*—*c. Tibiæ*, *Shin*—*c. Urethralis*, *Gallinaginis caput*.

CRISTA VESTIB'ULI, *Eminentia pyramidalis*. A crest which divides the vestibule of the ear into two fossæ,—one inferior and hemispherical,

called *Fo'rea hemisphærica*, *Si'nus rotundus*; the other superior and semi-elliptical. *Fo'rea elliptica* seu *semi-elliptica* seu *hemi-elliptica*, *Si'nus ovalis*. Morgagni has described a third groove-like fossa, *Recessus* seu *Fo'rea sulciformis*, situated at the mouth of the common orifice of the two superior semicircular canals.

CRISTÆ CLITORIDIS, Nymphæ.

CRISTALLIN, Crystalline.

CRISTALLINE, Mesembryanthemum crystallinum.

CRISTALLOIDITIS, Phacitis, see Crystalloid.

CRITHE, (κρίθη, 'barley.') Hordeolum, Hordeum, Fenis.

CRITHIDION, (κρίθιδιον, dim. of *erithe*.) Hordeolum.

CRITH'NUM MARIT'IMUM, *Crith'mum*, *Calchrys maritima*, *Fœniculum maritimum*, *Samphire*, (F.) *Passé-pierre*, *Perce-pierre*, *Fenouil marin*, *Bacile*. Family, Umbelliferae. Sex. Syst. Pentandria Digynia. A plant which grows on the seacoast, has a spicy, aromatic flavour, and is used, pilled, as a condiment.

The *Caaponga* of Brazil is a kind of crithmum.

CRITICAL, (κρίτικος,) *Criticus*, (F.) *Critique*, (from *crisis*.) Belonging to a crisis.

CRITICAL DAYS, *Dies Critici* seu *Judicatorii* seu *Decretorii* seu *Principes* seu *Radicales* seu *Contemplabiles* seu *Internuntii*, (F.) *Jours critiques*, are those on which a crisis, it is imagined, is most likely to happen. According to Hippocrates and Galen, the greatest number of fevers terminate favourably on the 7th day; and many on the 14th;—these two days being the most propitious. Next to these come, in order of efficiency, the 9th, 11th, 20th or 21st, 17th, 5th, 4th, 3d, 18th, 27th, and 28th. The sixth day was called, by Galen, the *Tyrant*, τυραννία, because the crises that happened then were generally unfavourable. After this, the most unfavourable were the 8th, 10th, 12th, 16th, and 19th. The 13th was a sort of neutral day; the crises which happened on it being neither favourable nor unfavourable. Days were, also, divided into *Intercalary*, on which the crises happened less frequently, and were less complete than on the *critical* or *indicatory*;—and into *vacant* and *non-decretory*, on which a crisis hardly ever occurred. According to this division, they were enumerated as follows:—

Critical days.....7th, 14th, 20th, 27th, 34th, 40th, 60th, &c.

Indicatory days.....4th, 11th, 17th, 24th, &c.

Intercalary days.....3d, 5th, 6th, 9th, &c.

Non-decretory days....2d, 8th, 10th, 12th, 13th, &c.

Fortunate crises were considered to be indicated by favourable signs appearing three days before.

CRITICAL PERIOD, see Menses.

CRITIQUE, Critical.

CROCHET, Crotechet.

CROCIDISMUS, (κροκίδειν, 'to pick wool;' κροκίς or κροκος, 'a lock of wool.') Carphologia.

CROCIDIXIS, Carphologia.

CROCINUM, (κροκινόν, from κροκος, 'saffron.') Made with saffron; coloured with saffron.

CROCO'DES, (κροκώδης, from κροκος, 'saffron,' and *odes*.) Certain troches into which saffron entered as an ingredient.

CROCODILEA, (κροκοδείλεια,) *Stercus Laceratæ*. The excrements of the crocodile, which the Arabists extolled as a remedy for cutaneous diseases, and which was long used as a cosmetic.

CROCODILIUM, Echinops.

CROCOMAG'MA, gen. *Crocomag'matis*, (κροκος, 'saffron,' and μαγμα, 'a kneaded or squeezed mass.') A kind of troch, composed of saffron and spices.

CROCUS, (κροκος,) *C. sativus* seu *officinalis* seu *Austriacus* seu *Orientalis*, *Medicina Tristitia*,

Panacea vegetabilis, *Zaff'ran*, *Anima Pulmonum*, *Jo'vis flos*, *Saff'ron*, (F.) *Safran*. Order, Iridaceæ. The stigmata, *Crocus* (Ph. U. S.) are the parts used in medicine. They are brought from the East. The odour is aromatic, and the taste aromatic, pungent, and bitter:—the colour deep orange red. Its virtues are yielded to alcohol, wine, vinegar, and water. Its operation has been considered stimulant, exhilarating and diaphoretic. It is not much used.

Cake Saff'ron is sometimes met with. It consists of one part of saffron and nine of marigold, made into a cake with oil, and pressed.

CROCUS, Coleothar—c. Antimonii, Oxydum stibii sulphuratum—c. Austriacus, Crocus—c. Germanicus, Carthamus tinctorius—c. Indicus, Curcuma longa—c. Martis adstringens, Coleothar—c. Martis aperiens, Ferri subcarbonas—c. Martis vitriolatus, Coleothar—c. Metallorum, Oxydum stibii sulphuratum—c. Officinalis, Crocus—c. Orientalis, Crocus—c. Saracenicus, Carthamus tinctorius—c. Sativus, Crocus.

CROCXYDISMUS, (κροκνίδειν, 'to pick wool') Carphologia.

CROIL (Sc.), Nanus.

CROISÉ (F.), (from *croix*, 'a cross,') *Crucia'tus*, *Cro'sed*. An epithet given to paralysis, when it attacks the arm of one side and the leg of another.

CROISEMENT, Crossing.

CROISSANCE, (from *croître*, [L.] *creescere*, 'to grow,') Growth.

CROISSANCES, Waxing kernels.

CROISSANTS, Waxing kernels.

CROIX DE MALTE, ([L.] *crux*, gen. *crucis*, 'a cross,') *Splenium Cruciatum*, *Maltese' Cross*, *Cross of Mal'ta*. A compress, having the form of the Maltese Cross. It is made of a piece of square linen, folded in four, and divided with scissors from each angle to a small distance from the centre. It is used especially after amputation of the fingers, penis, and limbs.

When the compress is cut in two of its angles only, it is called the *Half Maltese Cross*, *Demi-croix de Malte*.

CROK (Sc.), Nanus.

CROMMYON, (κρομμύον,) *Allium cepa*.

CROMMYOXYREG'MIA, (κρομμύον, 'an onion,' *oxys*, 'acid,' and *ῥήγνυμι*, 'to break forth.') Sour, fetid eructations, exhaling a smell similar to that of onions.

CROOPEACKED, see Hump.

CROOPY, Croupy.

CROPALÉ, Nerium antidysentericum.

CROSS OF THE AORTA, see Aorta:— Maltese, *Croix de Malte*.

CROSS-EYE, Strabismus.

CROSSE DE L'AORTE, see Arc.

CROSSE MITE, Acarus Crossii.

CROSSES PALMAIRES, Palmar arches.

CROSSFOOT, see Kylosis.

CROSS'ING, (F.) *Croisement*, *Métissage*. The union of an animal of one race with one of another, in order to improve the breed. In man, as well as in animals, the union of near relatives is apt to deteriorate the offspring, and hence crossing is important in his case.

CROSSWORT, (*cross*, and *wort*.) *Eupatorium perfoliatum*, *Lysimachia quadrifolia*.

CROTALOPHORUS, (κροταλον, 'a rattle,' and *φέρω*, 'I carry,') *Crotalus horridus*.

CROTALUS HORRIDUS, *C. durissus*, *Crotalophorus*, *Caudis'ona durissus*. The *Rattlesnake*, (F.) *Crotale*, *Serpent à Sonnettes*. Class, Reptilia, Ord. Ophidia. A venomous reptile of North America. Its poison is virulent. It is so called from the rattle in its tail, (κροταλον, 'a rattle,' a small bell.)

CROTAPHÆ, *Crotaphium*, (κροταφος, 'the temple,') *Cephalalgia Pulsatilis*, *Cephalæ'a Pulsat*.

vis, *Sphgmoccephalus*. A pulsatory pain, chiefly in the temples, with drumming in the ears.

CROTAPHITE ARTERIES, Temporal arteries.

CROTAPHITES, Temporal muscle.

CROTAPHIUM, (from *Crotaphus*), Crotaphe.

CROTAPHUS, (*κροταφός*), Temple, Temporal bone.

CRGT'CHET, *Ha'mus*. A small hook or crook. (F.) *Crochet*, (dim. of *croc*, 'a hook.') An obstetrical instrument, whose name indicates its shape, and which is used in the extraction of the fetus, when it becomes necessary to destroy it to expedite delivery. Crotchets are differently formed; some are sharp, others blunt; some contained in a sheath, others naked.

CROTON BENZOË, gen. *Crotonis*, (from *κροτων*, 'a tick,' which the seeds of certain species resemble), Benjamin.

CRO'TON CASCARILLA, *Cor'tex Eleutheria*, *Cro'ton Eleutheria* seu *Elute'ria*, *Clu'tia Elute'ria*, *Thus Judeo'rum*, *Chacarill'æ cor'tex*, *Cascarill'a*, *Eleute'ria*, *Gascarill'a*. Order, Euphorbiaceæ. Ser. Syst. Monœcia Monadelphia. (F.) *Quinquina aromatique*, *Cascarille*, *Chacarrille*. The bark of *Cro'ton Cascarill'a* of the Bahamas, *Cor'tex Thu'ris*, *Cascarill'a* (Ph. U. S.), (F.) *Écorce éluthérienne*, has a slightly aromatic odour, and bitterish aromatic taste. The smoke has the odour of musk. The active parts are an essential oil and bitter extractive. They are completely extracted by proof spirit. It is tonic and stomachic. Dose, gr. xv to ʒss or more, in powder. The bitter principle is called *Cascarill'in*.

CROTON ELEUTHERIA, *Croton cascarilla*—c. *Jamalgota*, *C. tiglium*—c. *Lacciferum*, see *Lacca*—c. *Oil*, *Croton tiglium*—c. *Malambo*, see *Matias*.

CROTON PSEUDOCHE'NA, *Copal'che* bush. A South American plant, which yields a bark—*Cortex Copal'che* seu *Copal'chi*, very like that of *Cascarilla*.

CROTON, PURGING, *Croton tiglium*.

CROTON RACEMOSUM, *Bee'nel*. A small Malabar tree, whose aromatic root, boiled in oil of sesame, is employed, by the orientals, as a liniment in headach and rheumatism.

CROTON SEEDS, see *Croton tiglium*.

CROTON TIGLIUM, *C. Jamalgot'a*, *Purg'ing croton*, *Cadel-Avanacu*. (F.) *Petit Pignon d'Inde*. A Ceylonese plant, every part of which is endowed with medicinal power. The root is a drastic cathartic: the wood, *Lignum Pava'næ*, *Pava'na wood*, and the seeds, have like virtues. The seeds have been long known under the names *Gra'na Moluc'æ*, *Til'ii Gra'na*, and *Gra'na Tig'lii* seu *Tig'lia*, *Cro'ton seeds*, *Purg'ing nuts*. (F.) *Graines de Moluques*, ou de *Tilly*. From these seeds the *Cro'ton Oil*, *O'leum Tig'lii*, is expressed. It is of a pale brownish-yellow colour, and hot, biting taste; owing to *Croton'ic Acid*, and is a most powerful drastic cathartic. Dose, from half a drop to three drops, made into pills with crumb of bread. It is also applied externally as a rubefacient, 3 to 5 drops being rubbed on the part; or one part of the oil and three parts of olive oil may be added together, and a little of this be rubbed on.

CROTONË, (*κροτων*, from *κροτων*, 'a tick.') A fungus, which grows on trees, and is produced by an insect. By extension, applied to a fungous tumour developed on the periosteum.

CROUP, (formed from the breathing and cough, by onomatopœia,) *Cynanche trachealis*—c. *Bronchial*, *Polypus bronchialis*—c. *Cerebral*, *Asthma thymicum*—c. *Cerebral spasmodic*, *Carpo-pedal spasm*—c. *Chronic*, *Polypus bronchialis*—c. *False*, *Cynanche trachealis*—c. *Faux*, *Asthma thymicum*, see *Cynanche trachealis*.

CROUP, HYSTERIC. A spasmodic affection of

the laryngeal muscles by no means unfrequent in hysterical females,—the paroxysm consisting in a long protracted, loud and convulsive cough, followed at times by crowing respiration, and by dyspnœa so great as to threaten suffocation. The treatment is that advised for hysteria.

CROUP, PSEUDO, *Asthma thymicum*—c. *Pseudo-nervæ*, *Asthma thymicum*—c. *Spasmodic*, see *Asthma thymicum*, and *Cynanche trachealis*—c. *Spurious*, *Asthma thymicum*.

CROUPAL, *Croupy*.

CROUPE, see *Croupion*.

CROUPION (F.), *Uropygium*, *O'rus*, *Or'rhos*, the *Rump*, the *Crupper*, (F.) *Croupe*. The region of the coccyx.

CROUPLIKE INSPIRATION OF INFANTS, *Asthma thymicum*.

CROUPOUS, *Croupy*.

CROUPY, *Croup'ous*, *Crup'ous*, (F.) *Croupal*. Relating, or appertaining to, or resembling croup; as a 'croupy cough.' Hoarse, (Prov.) *Croupy*.

CROUPY INFLAMMATION, *Inflammati'o crup'osa*. Inflammation of a mucous membrane, attended with a pseudo-membranous exudation.

CROÛTE, old (F.) *Crouste*, *Crusta*, *Scab*—c. *de Lait*, *Porrigo larvalis*—c. *Laitense*, *Porrigo larvalis*.

CROÛTEUX, *Crustacé*.

CROWD POISONING, *Ochlesis*.

CROWFOOT, *Geranium maculatum*, *Ranunculus*—c. *Bristly*, *Ranunculus Pennsylvanicus*—c. *Bulbous*, *Ranunculus bulbosus*—c. *Bulbous-rooted*, *Ranunculus bulbosus*—c. *Celery-leaved*, *Ranunculus sceleratus*—c. *Crushed*, *Ranunculus sceleratus*—c. *Marsh*, *Ranunculus sceleratus*—c. *Meadow*, *Ranunculus acris*—c. *Small flowered*, *Ranunculus abortivus*—c. *Tall*, *Ranunculus acris*—c. *Water*, smaller, *Ranunculus flammula*.

CROWLING, (onomatopœia,) *Borborgmus*.

CROWN, *Coro'na*, *Steph'anæ*, (G.) *Krone*, (F.) *Couronne*. In anatomy, this name is given to parts of a circular form, which surmount other portions of the same body. Thus, the *Crown of a Tooth*, *Coro'na Dent'is*, *Men'sa*, (F.) *Couronne du Dent*, is the portion of the tooth which projects above the gum.

CROWN OF THE GLANS, *Coro'na seu Tor'ulus glan'dis*, (F.) *Couronne du Gland*, is the round, almost circular, ring, which circumscribes the base of the glans.

CROWN OF THE HEAD, *Vertex*.

CROWN OF THE TREPAN, *Mod'i'olus*, (F.) *Couronne du Trépan*, is a species of saw, in form of a crown, or rather of a portion of a cylinder, having grooves on its external surface, and teeth at its lower extremity; the other being fitted to the handle of the trepan.

CROWN OF VE'NUS, *Coro'na Ven'ëris*, *Gul'ta Ros'acea Syphilit'ica*, (F.) *Couronne de Vénus*. Red, rosy pustules, dry or moist, on the face; but particularly on the forehead and temples, owing to constitutional syphilis.

CROWS' BEAK-LIKE PROCESS, *Coracoid process*.

CRU, *Crude*.

CRUCIAL, (*crux*, gen. *crucis*, 'a cross.') *Crucial'is*, *Crucia'tus*. Having the shape of a cross. Appertaining to a cross.

CRUCIAL BANDAGE, *T. Bandage*.

CRUCIAL INCISION, *Incisi'o'ra crucial'is*, (F.) *Incision cruciale*. An incision made in the form of a cross. Often employed for exposing the cranium, for the purpose of applying the trepan.

CRUCIAL LIGAMENT OF THE ATLAS, *Annular ligament of the Atlas*.

CRUCIAL LIGAMENTS, (F.) *Ligaments croisés ou cruciformes*. Two strong ligaments within the knee-joint. The anterior passes obliquely from a depression anterior to the spine of the tibia to the posterior and inner part of the exter-

nal condyle of the femur:—the other the posterior, extends from the posterior part of the spine of the tibia to the anterior and outer part of the internal condyle.

CRUCIA'TI DOLO'RES, (pl. of *cruciatius*, from *crux*, gen. *crucis*, 'a cross,' 'torment,') Excruciating pains.

CRUCIATUS, *Croiséé*.

CRUCIBLE, (from *crux*, gen. *crucis*, so called —it has been supposed— from being made in the shape of a cross, or from having a cross impressed upon it, [?] or from (F.) *creux*, 'a cavity,' 'a pit,') *Crucibulum*, *Catinus fusorius*, *Albot*, *Cementarium*, (F.) *Creuset*. A vessel of earth, silver, platinum, gold, blacklead, &c., for receiving substances, which have to be exposed to a strong heat. It is sometimes covered with a top or lid. (F.) *Convercle*.

CRUCIFORM, (*crux*, gen. *crucis*, and *forma*, 'shape,') *Cruciformis*, *Cross-shaped*. A name given to the ligaments which close the articulations of the phalanges; and likewise to the crucial ligaments.

CRUDE, *Cru'dus*, *O'mus*, (F.) *Cru*. Raw, unripe, not concocted.

CRUDITAS, Crudity—c. Morbi, see Crudity—c. Ventriculi, see Crudity.

CRUDITY, *Cru'ditas*, *Om'otes*, *Sta'tus Crudi'tatis*, (from *crudus*,) *Incoc'tus*. Rawness, Crudeness. This has received several acceptations. 1. It expresses the quality of certain aliments, which have not experienced the action of fire. 2. The condition of matters in the digestive tube, which have not undergone the digestive changes—*cruditas ventriculi*; and, 3. In the language of the Humorists, it means the condition of the morbid matter in a sick individual, *cruditas morbi*, when it has not yet been prepared or concocted by the action of the organs—*Azym'ia humorum*.

The word is used in the plural, *Cru'dities*, synonymously with *crude matters*. (F.) *Mat'eres crues*. It is applied to those, when contained in the stomach and intestines.

CRUDUS, Crude.

CRUE, (from part. of *croître*, 'to grow,') Growth.

CRUELS, Herpes Zoster, Scrofula, ([F.] *Écoulees*).

CRUENTA (from *cruur*) EXSPUITIO, Hæmoptysis.

CRUES, pl. of *cru*, (*Mat'ères*,) see Crudity.

CRUMENA VESICÆ, (*crumena*, 'a small pouch,') The cavity of the urinary bladder.

CRU'OR, gen. *Cru'oris*, (from *κρυος*, 'congelation,') Coagulated blood, gore. The signification of this word is very vague. It has been used to designate blood in general, venous blood, extravasated or coagulated blood, and the colouring matter of the blood.

CRUOR SANGUINIS, see Blood.

CRUORIN, Hæmatin.

CRUPPOSUS, Croupy.

CRUPPER, ([F.] *Croupière*, from *croupe*, 'the lump,') *Croupion*—c. Bone, Coccyx.

CRUPSIA, (*κρυψίς*, 'concealment, deception,') Metamorphopsia.

CRU'RA, (pl. of *crus*, 'a leg,') (F.) *Cuisse*. Applied to some parts of the body, from their resemblance to legs or roots; as the *Cru'ra cer'ebri*, *Cru'ra cerebel'li*, &c.

CRURA ANTERIORA MEDULLÆ OBLONGATÆ, Peduncles of the brain—c. Cerebelli, Peduncles of the cerebellum—c. Cerebelli ad Corpora Quadrigemina, see Peduncles of the cerebellum—c. Cerebelli ad Pontem, see Peduncles of the cerebellum—c. Cerebri, Peduncles of the brain—c. Clitoridis interna, Bulbus vestibuli—c. Meçullæ oblongatæ, Corpora restiformia, Thalami nervorum opticorum, see Peduncles of the cerebellum

—c. Posteriora medullæ oblongatæ, Peduncles of the cerebellum.

CRURÆUS, Cruralis.

CRURAL, *Crura'lis*, (*crus*, gen. *cruris*, 'the thigh, and lower limb,') What belongs to the thigh or lower limb.

CRURALIS, Triceps cruris.

CRURAL ARCH, *In'guinal or Fem'oral Arch*, (F.) *Arcade crurale ou inguinale*, (Ch.), *Poupart's Ligament*, *L. of Fallo'pius*. This arch is formed by the internal portion of the inferior edge of the aponeurosis of the obliquus externus muscle, which is attached, at one end, to the pubis; at the other, to the anterior and superior spinous process of the ilium. At its posterior and inner part, the aponeurosis, forming the arch, sends off a falciform reflection, which is attached along the crest of the pubis, and is known under the name of GIMBERNAT'S Ligament. Beneath this arch, the vessels, nerves, and muscles make their exit from the pelvis to the thigh.

CRURAL ARTERY, *Arte'ria Crura'lis*, *Fem'oral Ar'tery*, (F.) *Artère crurale*, is the continuation of the external iliac. It extends from the crural arch to the aperture in the triceps, or to the ham. Chaussier applies the name *Artère crurale* to the trunk, which extends from the primitive or common iliac to the tibial arteries—embracing, of course, the external iliac, femoral, and popliteal.

CRURAL CANAL, *Crural Ring*, *Fem'oral canal' or ring*. M. Jules Cloquet has described this canal with minuteness, and given it the name *Anneau crural*, *Anneau femoral*. It is nearly an inch long, triangular, more spacious above than below, and shorter and broader in the female than in the male. Its upper orifice is bounded, anteriorly, by the crural arch; posteriorly, by the crista of the pubis; on the outer side by the psoas and iliacus muscles, covered by the iliac aponeurosis, and at the inner, by Gimbernats's ligament. This orifice is covered by the peritoneum, and, according to M. Cloquet, is closed by a more or less resisting septum, which he has named *Septum crura'le*, *Crura'ral Sep'tum*. In its course, the crural canal has its anterior parietes formed by the superficial expansion of the fascia lata: the posterior by the pectineus, covered by the deep-seated expansion of the fascia; and more externally by the psoas and iliacus muscles, covered by an expansion of the fascia iliaca. Its inferior orifice is formed by the foramen of the fascia lata, which gives passage to the vena saphæna. It is at the upper orifice of this canal, that *Fem'oral or Crura'ral Hernia*, *Hernie inguinale* of Chaussier, occurs; which would be more common, were it not for the fibrous cellular septum there situate.

CRURAL NERVE proceeds from the lumbar plexus, and is situate at the outer side of the psoas muscle and crural artery. After it has passed under the crural arch, it divides into cutaneous and muscular branches. One of the branches, which is larger than the rest, is called the *Saphæ'na nerve*. It gives off filaments to the integuments of the knee, to the inner part of the leg, and to the dorsal surface of the foot. The remainder of the branches of the crural are distributed to the anterior and inner part of the thigh. The *Ac'cessory of the Crural Nerve* is a term given to the 4th and 5th pairs of lumbar nerves.

CRURAL PLEX'US of Chaussier is the union of the anterior branches of the last four pairs of lumbar nerves, and the first four sacral; forming the lumbar and sacral plexuses of most anatomists.

CRURAL RING, Crural canal—c. Septum, see Crural canal.

CRURAL VEIN, *Fem'oral vein*, has the same arrangement as the artery. It receives only one great branch, the saphæna.

CRURA' LIS, *Crura'lis*. A part of the *Triceps*

crural of the French, or of the *Trièmororotulien* of Chaussier. The *crural* is situate at the anterior, outer, and inner part of the thigh. It arises, fleshy, from between the two trochanters, adheres firmly to most of the fore part of the os femoris, and is inserted, tendinous, into the upper part of the patella, behind the rectus. Its use is to assist the vasti and rectus in the extension of the leg. Under *Muscle Triceps Crural*, the French describe the *cruralis* and the two vasti. Some small muscular slips, sometimes found under the *crureus* muscle, and inserted into the capsular ligament of the knee-joint, have been called *Sub-crurei*.

CRURIS RADIUS, Fibula.

CRUS, gen. *Cruris*, Leg, Thigh—c. Fornicis, see Fornix.

CRUST, Eschar—c. Milk, Porrigo larvalis—c. of the Peduncle, see Peduncles of the brain.

CRUST'IA. A crust or scab. (F.) *Croûte*. An assemblage of small flakes, formed by the drying up of a fluid secreted by the skin.

The lining membrane of the stomach and intestines has been called *Crusta villosa*.

CRUSTA ADAMANTINA DENTUM, Enamel of the teeth.

CRUSTA GENU EQUI'Æ, *Hippogonyolæpus*, *Siccat* or *Knee Scab*, *Mack* or *Envelled Hoof*, *Knees*, *Hang'ers*, *Dev* *Claws*, *Night Eyes*, *Horse Crust*. This morbid secretion from the horse has been advised in cases of epilepsy. It is used in the form of powder (gr. ij to gr. xx), and of tincture.

CRUSTA INFLAMMATORIA, Corium phlogisticum—c. Lactea, Porrigo larvalis, P. lupinosa—c. Membranacea, Peristroma—c. Petrosa, see Tooth—c. Phlogistica, Corium Phlogisticum—c. Pleuretica, Corium phlogisticum—c. Pruriginosa, Gutta rosea—c. Serpiginosa, Gutta rosea—c. Vermicularis, Peristroma—c. Villosa lingue, see Tongue—c. Villosa ventriculi, see Stomach.

CRUSTACÉ (F.), (from *crusta*.) Alibert has substituted this word, in his Nosology, for *croûteux*, 'crusty.' Having crusts or scabs; as *Dartre crustacée*.

CRUSTULA, (dim. of *Crusta*.) A small shell or scab. An ecchymosis of the conjunctiva.

CRUSTUMINATUM. A rob, prepared from the *Pyra Crustumi'na* or *Crustumeri'na*, (from *Crustuminum*, a town in Italy, where they grew,) boiled with honey or in rain-water.

CRUT, Nanus.

CRUX CERVI, (gen. *Crucis*.) An ancient appellation for the bone in the heart of the stag. It was once considered useful in diseases of the heart.

CRY, *Gla'mor*, *Bo'ë*, (F.) *Cri*, (an example of onomatopœia.) The sound of the unarticulated voice. The native voice, which the idiot and deaf possess equally with the man of genius and hearing. The cry of the new-born child has been called *Vagi'tus*, (F.) *Vagissement*. We say, 'A cry of joy, of pleasure, of pain,' &c., according to the expression which it may convey to the hearer.

CRYING, Parturition.

CRYING-OUT, Parturition.

CRYMO'DES, (κρυμδός, from κρυμς, 'cold,' and odes.) A continued fever, *Fe'bris crymo'des*, in which the internal parts feel hot, and the external cold; and which, of old, was attributed to an erysipelatos inflammation of the lungs.

CRYMODYN'IA, (κρυμος, 'cold,' and οδυνη, 'pain,') Chronic rheumatism, and all its modifications.

CRYMO'SES, (κρυμος, 'cold,' and οσις.) Diseases caused by the action of cold.

CRYP'SOR'CHIS, gen. *Crypsor'chidis*, (κρυψορχις, from κρυπτος, 'I conceal,' and ορχις, 'a testicle,') *Cryptor'chis*, *Testicon'dus*. One in whom the testes

have not descended into the scrotum. This state is called, *Cryptorchidism'us*, *Parorchid'ium*.

CRYPT, Crypta.

CRYPT'IA, (κρυπτη,) *Crypt*, *Follic'ulose gland*, *Lacu'na*, *Follic'ulus*, *Follicle*, (F.) *Crypte*, *Follicule*. A *crypt* or *follicle* is a small secreting cavity, formed by a depression of the skin or mucous membranes—*Crypta mucosa*, *Follic'ulus mucosus*.

The use of the cryptal or follicular secretion is to keep the parts on which it is poured supple and moist, and to preserve them from the action of irritating bodies with which they have to come in contact.

The little rounded appearances at the ends of the small arteries, in the cortical substance of the kidney, are also called *Crypte*.

CRYPTA MUCOSA, see Crypta.

CRYPTÆ (pl. of *crypta*) LIEBERKÜHNIANÆ, Lieberkühn's glands—c. Præputiales, see Sebaceous glands—c. Sebaceæ, Sebaceous glands.

CRYPTE, Crypta.

CRYPTO, (κρυπτος, 'concealed,') in composition, concealed.

CRYPTOCEPH'ALUS, (*crypto*, and κεφαλη, 'head,') A monster whose head is excessively small, and does not appear externally.

CRYPTOCOCCUS CEREVISIÆ, (*crypto*, and κοκκος, 'a berry,') *Torula cerevisiæ*—c. Fermentum, *Torula cerevisiæ*.

CRYPTODID'YMUS, (*crypto*, and διδωμις, 'a twin,') *Fœtus in fœtu*. A monstrosity, in which one fetus is found contained in another.

CRYPTOG'AMOUS, *Cryptogam'ic*, *Cryptogam'icus*; (*crypto*, and γαμος, 'marriage,') An epithet applied by botanists to plants whose organs of fructification are concealed or not manifest. *Ag'amous* plants are those whose sexual organs are not known.

CRYPTOPY'IC, *Cryptopy'icus*, (*crypto*, and πυον, 'pus,') A state of disease, kept up by an occult abscess.

CRYPTOPYICUS, *Cryptopyic*.

CRYPTORCHIDISMUS, see *Crypsorchis*.

CRYPTORCHIS, *Crypsorchis*.

CRYPTS (pl. of *crypt*) OF LIEBERKÜHN, see Intestine—c. Synovial, *Bursæ mucosæ*.

CRYSTAL, MINERAL, Potassæ nitras fusus sulphatis paneillo mixtus.

CRYSTALLI, pl. of *Crystallus*, (κρυσταλλος, 'a crystal,') Vesicles filled with a watery fluid. They are also called *crystal'line*. Probably the pemphigus of modern writers.

CRYSTALLI LUNÆ, Argenti nitras.

CRYSTALLIN, Crystalline.

CRYSTALLINA, (κρυσταλλος, 'a crystal,') A vesicle or phlyctæna, filled with serum, and appearing on the prepuce or in the vicinity of the anus, surrounded by a reddish extravasated aureola. It may be syphilitic or not. See Crystallii.

CRYSTALLINA TUNICA, Arachnoid membrane.

CRYSTALLINE, (κρυσταλλινος,) *Crystal'linus*. Having the appearance or nature of crystal.

CRYSTALLINE, *Cry'stalline hu'mour* or *Lens* or *Body*, *Cry'stall'linus*, *Lens crystal'lina* seu *crystal'loï'des*, *Cor'pus Crystal'linum* seu *Diacoï'des* seu *Crystallloï'des* seu *Phacoï'des*, *Hu'mor crystal'linus* seu *glaci'lis*, *Pha'cē*, *Pha'cea*, *Pha'cus*, *Gem'mi Oculi*, (F.) *Cristallin* ou *Crystallin*, *Humeur crystal'line*, *Corps crystal'lin* seu *phacoï'de*, *Lentille crystal'line*. A lenticular, transparent body, situate between the vitreous and aqueous humours of the eye, at the union of the anterior third with the two posterior thirds of the organ. It is composed of a soft exterior substance; and an interior, composed of an albuminoid substance, called *Cry'stalline* or *Kry'stalline* by Hünefeld, (F.) *Crystalline*, forming a solid nucleus, in which a number of elliptical layers is perceptible. It is contained in

a capsule, called *Tu'nica ara'nea seu crystallina*, *Cap'sula len'tis*, *Periph'aena*, *Phacocys'tis*, and receives, at its posterior surface, a small branch of the central artery of the retina, which is always readily distinguishable in the foetus, prior to the seventh month of utero-gestation.

The use of the crystalline is to refract the rays of light, and to serve as an achromatic glass; for which its laminae or layers, increasing in refractive power from the circumference to the centre, admirably adapt it.

CRYSTALLINOCAPSULITIS, see Phacitis.

CRYSTALLION, *Plantago psyllium*.

CRYSTALLITIS, (*crystalline and itis*), Phacitis.

CRYSTALLOCATAPIESIS, (*καταπιεσις*, 'depression.') See Cataract.

CRYSTALLOCATATHESIS, (*καταθεσις*, 'a laying down.') See Cataract.

CRYSTALLOID, *Crystallōides*, (*κρυσταλλος*, 'crystal,' and *ειδος*, 'form,' 'resemblance.') Resembling crystal or the crystalline. The capsule or membrane of the crystalline. Also, the crystalline itself.

CRYSTALLUS MINERALIS, *Potassæ nitras fusus sulphatis paucello mixtus*.

CRYSTALS, BLOOD, *Hæmatoidin*.

CRYSTAUX D'HÉMATINE, *Hæmatoidin*.

CTEDON, (*κτεδων*), Fibre.

CTEIS, (*κτεις*, (L.) *pecten*, 'a comb.') Pubes.

CTESIPHON'TIS MALAG'MA. A plaster employed and described by Celsus.

CUBA, see Havana.

CUBAL SINI, *Piper cubeba*.

CUBATIO, (*cubare*, 'to lie down.') *Decubitus*.

CUBEBA, *Piper Cubeba*.

CUBIFORMIS, (*cubus*, 'a cube,' and *forma*, 'shape.') Cuboid.

CUBIT, *Cubitus*, *Ulna*—c. Top of the, *Olecranon*.

CUBITAL, *Cubita'lis*, *Ul'nar*, *Ulna'ris*, (F.) *Ulnaire*. Connected with or relating to the *cubitus*, or to the inner and posterior part of the forearm.

CUBITAL ARTERY, *Arteria cubita'lis*, *A. ulna'ris*, arises from the humeral a little below the bend of the elbow; proceeds along the anterior and inner part of the forearm; passes anterior to the ligamentum annulare of the carpus, and goes to form, in the palm of the hand, the superficial palmar arch. Besides the numerous muscular branches, which it gives off in its course, it sends posteriorly the common trunk of the *interosceus arteries*, and internally, the two *cubital recurrents*, anterior and posterior,—articular branches, which anastomose on the inside of the elbow with the divisions of the humeral artery.

CUBITAL MUSCLES are two in number. 1. The *Cubita'lis anterior seu inter'nus*, *Flex'or car'pi ulna'ris*, *Ulna'ris inter'nus*, *Epitrochlo-cubito-carp'ien*, *Cubito-carp'ien*—(Ch.), is a long muscle, situate at the anterior and inner part of the forearm. It arises from the inner condyle of the os humeri, at the inner side of the olecranon, and from the posterior edge of the ulna, and is inserted by a tendon into the os pisiforme. Its use is to bend the hand on the forearm, by directing it slightly inwards. 2. The *Cubita'lis poste'rior seu exter'nus*, *Extensor car'pi ulna'ris*, *Ulna'ris exter'nus*, *Cubite'us exter'nus*, *Cubito-us-métacarp'ien*—(Ch.), is situate at the posterior part of the forearm. It arises from the external condyle of the os humeri, and is inserted into the superior extremity of the fifth bone of the metacarpus. Its use is to extend the hand, inclining it a little inwards.

CUBITAL NERVE, *Ul'nar nerve*, *Cubito-digital*—(Ch.), is furnished by the last two or three nerves of the brachial plexus, and is distributed to the inner and anterior side of the forearm; to the inner part of the palm and of the back of the hand, and to the last two or three fingers.

CUBITAL VEINS, DEEPSEAT'ED, and the *Recur'rent cu'bital veins*, accompany the corresponding arteries. The superficial cubital veins belong to the basilic. Chaussier calls them *cutaneous cubital*.

CUBITALE (OS), Cuneiform bone.

CUBITALIS RIOLANI, see Anconeus.

CUBITEUS EXTERNUS, see *Cubital muscles*.

CUBITO, in composition, the cubitus or ulna.

CUBITO-CARPIEN, see *Cubital muscles*—c. *Cutaneé (nerf)*, Cutaneous nerve—c. *Phalangettien commun*, *Flexor profundus perforans*—c. *Radialis-métacarp'ien du pouce*, *Abductor longus pollicis*—c. *Radial*, *Pronator radii quadratus*.

CUBITOSUPRAPALMA'RIS. Belonging to the cubitus and to the supra-palmar or dorsal surface of the hand. Chaussier gives this name: 1. To a small artery, which is given off by the cubital or ulnar, a little above the wrist. 2. To a vein, which accompanies this artery.

CUBITO-SUS-MÉTACARPIEN, see *Cubital muscles*—c. *Sus-métacarp'ien du pouce*, *Abductor longus pollicis*—c. *Sus-phalangettien de l'index*, *Extensor proprius indicis*—c. *Sus-phalangettien du pouce*, *E. longus pollicis*—c. *Sus-phalangien du pouce*, *E. pollicis brevis*.

CUBITON, (*κυβιτων*), *Cubitus*.

CUBITUM, *Cubitus*.

CUBITUS, *Cu'bition*, *Cu'bitum*, *Cy'biton*. The *El'bow*. Also, one of the bones of the forearm. See *Ulna* and *Forearm*.

CUBITUS, (F.) *Coudée*, the ancient name of a measure 18 inches long.

CUBITUS COSTÆ, see *Costa*—c. *Supinus*, see *Decubitus*.

CU'BOID, *Cuboï'des*, *Cuboï'deus*, *Cubo'des*, *Cubifor'mis*, *Cy'boïd*, *Grandino'sum os*, *Tes'sara*, *Tes'sera*, *Os va'r'ium*, (*κυβος*, 'a cube,' and *ειδος*, 'form.') Having the form of a cube. This name was given by Galen to one of the bones of the tarsus, and is still retained. It is situate at the anterior and outer part of the tarsus; and is articulated, behind, with the calcaneum; before, with the last two metatarsal bones, and within, with the third os cuneiforme, and sometimes with the scaphoides. Its inferior surface has an oblique groove for the tendon of the *peroneus longus*.

CUCKOLD, *Aretium lappa*.

CUCKOO FLOWER, *Cardamine pratensis*.

CUCKOW BREAD, *Oxalis acetosella*.

CUCKOW PINT, *Arum maculatum*.

CUC'BALUS BE'HEN, *Be'hen officina'rium seu vulga'ris*, *Sile'nē infl'ata seu crassifolia seu Tho'rei*, *Visca'go be'hen*. This plant was once considered alexipharmic and cordial. It is the *Spitting Poppy*.

CUCULA'TUM MA'JUS. A barbarous term, used by Ruland for brandy and spirit of wine.

CUCULLARIS, *Trapezius*.

CUCULLATA, *Sanicula*.

CUCULLUS, ('a hood,' *Ouvrechef*, *Cucupha*, Infundibulum of the cochlea.

CUCULUS, ('a cuckoo,') see *Coccyx*. Pertussis.

CUCUMBER, ([F.] *concombre*), *Cucumis sativus*—c. *Asses*, *Momordica elaterium*—c. *Indian*, *Medeola Virginia*—c. *Squirting* or wild, *Momordica elaterium*—c. *Star*, one-seeded, *Sycios angulatus*—c. *Tree*, *Magnolia acuminata*.

CUCUMER, *Cucumis sativus*.

CUCUMIS, gen. *Cu'cumis* and *Ouen'merie*, 'a gourd,' *Cucumis sativus*, *Penis*, *Sicyos*—c. *Agrestis*, *Momordica elaterium*—c. *Caffer*, see *Cucurbita citrullus*.

CUCUMIS COLOCYN'THIS, *Citrull'us Colocyn'this*. Official names of the *Colocyn'th* or *Bit'ter Ap'ple*, *Colocyn'this*, *Coloquint'ida*, *Athan'dal* (Arab.), *Bit'ter Gourd*, *Bit'ter Cuc'umber*, (F.) *Coloquinte*, *Ord.* *Cucurbitaceæ*. *Sex. Syst.* *Monœcia Mona-*

delphin. A Turkey and Cape of Good Hope plant. The spongy part or medulla of the fruit, *Colocynthis* seu *Cucumeris Colocynthis* Pullpa, *Colocynthis* (Ph. U. S.), has a bitter, nauseous, and acrimonious taste, and is a strong cathartic, acting chiefly on the upper part of the intestines. It is scarcely ever used, except in combination. Its virtues are dependent upon a bitter principle — *Colocynthin*, *Colocynthin*.

CUCUMIS ME'LO. The systematic name of the Melon Plant, *Me'lo*, *Com'mon Melon*, (F.) *Melon*. The fruit is an agreeable article of diet, but not very digestible, unless when ripe. The seeds possess mucilaginous properties.

CUCUMIS SATIVUS. The systematic name of the *Cucumber plant*, *Cucumis*, *Angour'ion*, *Citrelolus*, *Cucumer* or *Cur'vimer*, from its curved shape, (*curvus*, 'crooked.') (F.) *Concombre ordinaire*. The *cucumber*, *cov'cumber*, is used, when young, as a pickle, when it is called a *Gher'kin*, (F.) *Cornichon*. It is not a fruit easy of digestion. The seeds are mucilaginous.

An ointment of green cucumbers — *cucumber ointment* — is used as a cooling salve. It is made by combining, by means of heat, the juice of green cucumbers with lard and veal suet.

CUCUMIS SYLVESTRIS, *Momordica elaterium*.

CUCUPHA, *Cucull'us*, *Pileus*, *Sacculus cephal'vus*. A sort of coif or cap, with a double bottom, between which is enclosed a mixture of aromatic powders, having cotton for an excipient. It was formerly used as a powerful cephalic.

CUCURBITA, (*à cucuritate*, owing to its shape.) A gourd. (F.) *Citrouille*, *Gourde*. See *Cupping-glass*.

CUCURBITA ANGURIA, *C. citrullus*.

CUCURBITA CITRUL'US, *C. Anguria* seu *pinnatifida*. The systematic name of the *Water-melon plant*; *Citrul'us*, *Anguria*, *Tetrangu'ria*, *Sicilian Cit'ul*, *Watermelon*. *Ord. Cucurbitaceæ*; *Ser. Syst. Monœcia Monadelphia*. (F.) *Melon d'eau*, *Pastèque*. The juice of the fruit is very abundant, whence its name. The *Watermelon* is extremely refreshing and agreeable, when made cool, and is eaten like the common melon. It is very much used in the United States; and in South Africa the *Cucumis Caffr*, the *Kengive* or *Keme*, — *watermelon* of the country — animals of every sort, including man, revel on it.

CUCURBITA LEUCANTHA, *C. Lagenaria* — *c. Melo-pepo*, see *Cucurbita pepo*.

CUCURBITA PE'PO, *Pe'po vulga'ris*, *Pe'po*. The systematic name of the *Com'mon Pom'pion* or *Pump'kin*, *Cucur'bita*, (F.) *Potiron* *courge*. The seeds of this plant, *Pepo* (Ph. U. S.), as well as those of *Cucur'bita lagenaria*, *Bottle-gourd*, contain a large proportion of oil, capable of forming an emulsion. They have been recommended in cases of tapeworm, in the dose of $\mathfrak{z}$ ss to $\mathfrak{z}\text{ij}$. The fixed oil of the seeds has been given with success, in the same cases, in the dose of half a fluidounce.

The fruit of **CUCURBITA LAGENARIA**, *C. leucantha*, *Pe'po lagenarius*, (F.) *Calebasses*; of *C. MEL'OPEPO*, *Round* or *Pattypun Squash*, *Cym'ling*; and of *C. VERRUCOSA*, *Warted* or *Long-necked Squash*, are eaten.

CUCURBITA PINNATIFIDA, *C. Citrullus*.

CUCURBITAIN, *Cucurbitinus*.

CUCURBITATIO, (see *Cucurbita*), *Cupping*.

CUCURBITE, see *Alembic*.

CUCURBITINUS, (F.) *Cucurbitin*. This name was formerly given to *Tenia solium*, because composed of rings which resemble the seeds of the gourd, — *cucurbita*. The ancients believed that the rings, which are sometimes discharged, were so many separate worms. See *Tænia*.

CUCURBITULA, (dim. of *cucurbita*), *Cupping-glass*.

CUCURBITULÆ CRUENTE, (pl. of *cucurbitula*), *Cupping with the scarificator* — *c. cum Ferro*, *Cup-*

ping with the scarificator — *c. Sicce*, *Cupping*, *dry*.

CUDWEED, *Antennaria plantaginifolia*, *Gnaphalium margaritaceum*.

CUILLERÉE, *Cochleare*.

CUIR, *Corium* — *c. Chevelu*, *Scalp*.

CUISSART, (from *cuisse*.) A wooden leg. See *Cornu ammonis*.

CUISSE, (from *coxa* [?]) *Thigh*. *Crus* — *c. Postérieure du cercelet*, *Corpus testiforme*.

CUISSON (F.), (from *cuire*, 'to smart, to burn.') A smarting, burning pain.

CUIVRE, ([G.] *Kupfer*), *Cuprum* — *c. Ammoniacal*, *Cuprum ammoniatum* — *c. et Anemoniaque*, *sulfate de*, *Cuprum ammoniatum* — *c. Jaune*, *Brass* — *Limailles de*, see *Cuprum* — *c. Sous-acétate de*, *Cupri subacetates* — *c. Sulfate de*, *Cupri sulphas*.

CUIVREUX (F.), *Coppercoloured*. A tint of complexion, (F.) *Teint cuivreux*, observed in cancerous affections. Syphilitic ulcers of the throat, &c., are often coppercoloured.

CUJETE, *Crescentia cujete*.

CULBIC'IO, (A Latin word, employed by old writers as synonymous with *ardor urinæ* and *gonorrhœa*.)

CULBUTE (F.), (*cul*, 'the ramp,' and *bute*, 'to stumble;') 'a tumble head-over-heels.' A movement which the foetus has been supposed to execute at the 7th month of utero-gestation; and by means of which, it was presumed, the head presented towards the orifice of the uterus; a change of situation which is impracticable in ordinary circumstances.

CULCITA SALINA, ('a salt pillow,') *Halo-type*.

CUL-DE-SAC, ([F.] 'a blind alley') *DOUGLAS'S*, see *Uterus*.

CULBUS. A measure containing 20 barrels, or 40 urns, equal to 180 gallons. Also, the *Scrotum*.

CULEX, *gen. Culicis*, 'a gnat,' (F.) *Cousin*.

A genus of insects, unhappily too well known in almost every part of the world, on account of their bites, which give rise to painful, local inflammation. The *gnats* and *mosquitoes* belong to this genus.

CULEX MOSQUITO, *Mosquito*.

CULI FLOS, *Cardamine pratensis*.

CULLAWAN, see *Laurus culilawan*.

CULINARY ART, (*culi'na*, 'a kitchen,') *Cook'ery*, *Res culina'ria* seu *coqua'ria*, *Ars culina'ria* seu *coquina'ria*, *Mage'ricæ*. The art of preparing meats for the table. In judging of the dietetic properties of various kinds of aliment, the culinary process to which they have been subjected will always have to be considered. Many of the writers on the culinary art have been physicians.

CULITLAWAN, see *Laurus culilawan*.

CULMINATIO, (*culmen*, *gen. culminis*, 'the top,') *Acme*.

CUTELLAIRE (F.), (*cutellus*, 'a little knife,') *Couteur cutellaire*, known also under the name *Couteau de feu*, *Fire-knife*, is used for what the French term the *Cautére transcurrente*. (See *Cauterization*.) It is shaped like a small hatchet.

CUTELLUS, (dim. of *culter*), *Culter*, *Knife* — *c. Aneeps*, *Knife*, double-edged — *c. Uncus*, *Knife*, *crotchet*.

CULTER, *Cultell'us*, (*colo*, *cultum*, 'I cultivate,') *A coult'er*, *a knife*, *scal'pel*, *mach'e'ra*, *mach'e'ron*, *mach'e'ris*. Also, the third lobe of the liver, so called, from some fancied resemblance, by Theophilus Protospatarius.

CULTER CURVUS, *Knife*, crooked — *c. Falcatus*, *Knife*, crooked — *c. Lenticularis*, *Knife*, *lenticu'lar* — *c. Rectus*, *Knife*, *amputation* — *c. Tonsorius*, *Razor*.

CULTRIVOROUS, *Cultrivorus*, (*culter*, 'a knife,' and *vorare*, 'to devour.') Individuals, who have seemed to swallow knives with impunity, have been so called;—*Knife-eaters*.

CULUS, Anus.

CULVER'S PHYSIC, *Leptandra Virginica*.

CULVER'S ROOT, *Leptandra purpurea*.

CUMAMUS, *Piper cubeta*.

CUMIN, *Cuminum cyminum*—*c. Faux*, *Nigella*—*c. des Prés*, *Carum*.

CUMINUM CYMINUM, (*κuminum*). The systematic name of the *Cumin* or *Cumin plant*, *Cuminum minimum* seu *Romanum*, *Cyminum*, *Cumnum*, *Carnabadium*, *Ord.* Umbelliferae. *Scr. Syst.* Pentandria Digynia. (F.) *Cumin*, *Anis aigre*. The seeds of cummin, which is a native of Egypt, have a warm, bitterish, disagreeable taste. Water extracts their odour, and spirit takes up both odour and taste. They are not used, except in a plaster, which bears their name. When drunk in wine, the ancients believed they produced paleness; hence, Horace called cummin *exsangue*; and Juvenal, *pal lens*.

CUMINUM MINUTUM, *C. cyminum*—*c. Nigrum*, *Nigella*—*c. Pratense*, *Carum*—*c. Romanum*, *C. cyminum*.

CUMMIN, *Cuminum cyminum*.

CUMULUS, 'a heap or pile.' A rounded prominence, in the centre of the proligerous disk, in which there is a small opaque cavity that contains the ovum. See *Tache embryonnaire*.

CUMULUS, *GERMINAL*, *Tache embryonnaire*—*c. Germinativus*, *Tache embryonnaire*—*c. Proligerus*, *Tache embryonnaire*.

CUNEA LIS SUTURA, (*cuneus*, 'a wedge.') The suture formed between the great and little alæ of the sphenoid bone and the os frontis.

CUNÉEN (F.), *Cunea'nus*. Relating to the cuneiform bones.

Articulations Cunéennes;—the joints between the cuneiform bones, as well as between them and other parts.

Ligaments Cunéennes;—the ligaments which hold the cuneiform bones together.

CUNEIFORM, *Cuneiformis*, *Sphenoïdes*, (*cuneus*, 'a wedge,' and *forma*, 'shape.') *Wedge-shaped*. This name has been given to several bones. 1. To the sphenoid. 2. To a bone of the carpus, situate between the os lunare and os orbiculare. It is, also, called *Os Pyramida'le*, *Os Triquetrum*, and *Os Cubital'e*. 3. To the basilar process of the occipital bone; and, 4. To three of the bones of the tarsus, which are distinguished, according to situation, reckoning from within outwards, into *first*, *second* and *third*,—or *internal*, *middle*, and *external*: and according to size, reckoning in the same order, into *great*, *small*, and *middle-sized*. The posterior surface of these bones is united to the anterior face of the scaphoides: the anterior surface with the corresponding metatarsal bones; and, in addition, the external surface of the third is articulated, behind, with the cuboides. They are also called *Chalcoïdea os'ea* seu *ossic'ula*.

CUNEIFORM CARTILAGES OF THE LARYNX, *Cartilages of Wrisberg*, are two small cylinders of fibro-cartilage, about seven lines in length, and enlarged at each extremity. By the base, the cartilage is attached to the middle of the external surface of the arytenoid; and its upper extremity forms a prominence on the border of the aryteno-epiglottidean fold of membrane. They are sometimes wanting.

CUNEO. In composition, the cuneiform bones.

CUNEOCUBOID, *Cuneocuboïdeus*. Belonging to the cuneiform bones and cuboides.

CUNEOCUBOID ARTICULATION is formed by the third cuneiform bone and cuboides. It is fur-

nished with a synovial capsule, and two *cuneocuboid ligaments*:—a *dorsal* and a *plantar*.

CUNEOSCAPHOID, *Cuneoscaphoïdes*. Belonging to the cuneiform bones and scaphoid.

CUNEOCAPHOID ARTICULATION is formed by the posterior surfaces of the three ossa cuneiformia, and the anterior surface of the scaphoid. It is furnished with a synovial capsule and ligaments, some of which are *dorsal*, others *plantar*.

CUNILA, *Satureia capitata*—*c. Bubula*, *Origanum*.

CUNILA MARIA'NA, *Satureia a origanoides*, *Calamintha erecta* Virginia'na, *Dittany*, *American* or *Mountain Dittany*, *Mint-leaved* or *Maryland Cuni'la*, *Stone-mint*, *Wild Bas'il*, *Sweet Horse-mint*, (F.) *Cunille d'Amérique*. A small indigenous herb, *Order*, Labiate; growing on dry, shady hills, from New England to Georgia, and flowering in June and July. Its medical properties are dependent upon essential oils, like the mints.

CUNILA, *MARYLAND*, *Cunila mariana*—*c. Mint-leaved*, *C. mariana*—*c. Muscula*, *Inula dysenterica*—*c. Pulegioides*, *Hedecna pulegioides*—*c. Sativa*, *Satureia hortensis*.

CUNILAGO, *Inula dysenterica*.

CUNILE D'AMÉRIQUE, *Cunila mariana*.

CUNNUS, (*κυνε*, 'to conceive,' (?) (Sax.) *epen*, 'a woman,') *Vulva*.

CUP, ([Sax.] *copp*), *Scutell'a*, *Catil'lus*, *Palet'la*, *Excipulum*, (F.) *Palette*, *Poëlette*, *Poilette*, *Vase à saigner*. A small vessel of determinate size, for receiving the blood during venesection. It has usually contained about four ounces. A bleeding of two cups is, consequently, one of eight ounces. See *Cupping*.

CUPAMENI, *Acalypha Indica*.

CUPEDLÆ, see *Confectio*.

CUPIDITAS, *Voluntas*—*c. Desedendi*, *Voluntas descendendi*.

CUPIDO, gen. *Cupi'dinis*, 'desire,' Appetite. Also, Cupid, the god of love, in ancient mythology, *Deus copulatio'nis*.

CUPOLA (L), 'a dome,' see *Infundibulum* of the cochlea.

CUPPED, see *Corium phlogisticum*.

CUPPING, *Catacasm'us*, ([F.] *couper*, 'to cut,' or to draw blood in vessels resembling cups.) *Applicatio cucurbitarum* seu *cucurbitularum*, *Cucurbitatio*. A species of bloodletting, performed by a *scarificator*, and glass, called a *cupping-glass*, *Cucurbit'ula*, *Cucurb'ita*, *Sic'ua*, (Old. Engl.) *Bleeding-boist*, (F.) *Ventouse*. The lancets are placed in such a manner in the scarificator, that, when it is applied upon the affected part, the whole are, by means of a spring, forced suddenly into it. After scarification, the cupping-glass, which has been previously exhausted by heat, or by an exhausting syringe, is applied. The pressure of the air within the glass being thus diminished, the necessary quantity of blood may be drawn. See *Bellometer*. *Dry cupping*, *Cucurbit'ula sic'ue*, is the application of the glasses, *dry cups*, (F.) *Ventouses sèches*, without previous scarification. It is used to prevent the activity of absorption from any wounded part; occasionally, to excite suppuration in indolent abscesses; and to remove the pus when an abscess is opened. *Cupping*, without any epithet, means the abstraction of blood by means of the scarificator and cups:—(F.) *Ventouses scarifiées*, (L.) *Cucurbit'ula cruent'is*, *C. cum Ferr'o*, *Wet cupping*, or *Wet Cups*, *Cut Cups*. The verb 'to cup,' signifies to draw blood by cupping.

CUPPINGGLASS, see *Cupping*.

CUP PLANT, *Silphium perfoliatum*.

CUPRESSUS, (*κρηταίρεος*,) *C. sempervirens*—*c. Arbor vitæ*, *Thuya occidentalis*—*c. Disticha*, *Taxodium distichum*.

CUPRESSUS SEMPERVIRENS, *C. pyramida'lis*. The systematic name of the *Cupressus*, *Cyparis-*

cus, Cy'press, (F.) Cyprès. Nat. Ord. Coniferae. The berries, leaves, and wood have been considered astringent and useful in intermittents. The whole plant abounds with a bitter, aromatic, and terebinthinate fluid.

CUPRESSUS THYOIDES. *White cedar*, indigenous. An infusion of the tops is said to be stomachic, and, when taken warm, diaphoretic.

CUPRI AMMONIOSULPHAS. *Cuprum ammoniatum*—c. *Diacetas*, *C. subacetas*—c. et *Ammonie sulphas*, *Cuprum ammoniatum*.

CUPRI LIMAT'RA. *Fillings of Cop'per, (F.) Limailles de Cuivre*, have been used in hydrophobia. It is remarked under *Cuprum*, that they are inert.

CUPRI RUBIGO. ('rust of copper,') *Cupri subacetas*.

CUPRI SUBACE'TAS. *C. Diacet'as seu Rubi'go, Æru'go, Hispani'cum viride, Pra'sinum viride, Viride Æris. Æru'ca, Cal'ceithos, Verdigris, Crystals of Venus, Subac'etate of Copper, (F.) Sous-acétate de Cuivre, Vert-de-gris, Crystaux de Venus, Verdet.* Impure subacetate of copper. This, as usually met with, is in masses, difficult to break; not deliquescent; foliaceous; of a fine bluish-green colour, and salt taste. It is tonic, emetic, escharotic, and detergent; but scarcely ever employed internally. Chiefly used in detergent ointments. Dose, as a tonic, under $\frac{1}{2}$ gr.: emetic, from gr. j to gr. ij. Powdered verdigris appears to be the active ingredient in *Smellome's Eye-salve*, which may be imitated by rubbing half a drachm of finely-powdered verdigris with a little oil, and then mixing it with an ounce of yellow basilicon.

An ointment composed of one drachm of finely-powdered verdigris, with an ounce of lard or spermaceti ointment, is used in psoriasis, tetter, &c.

CUPRI SULPHAS. *Vitriolum Cup'ri seu Ven'is seu Cyp'rium seu Cyp'rinum seu Ceru'leum seu Roma'nium, Cup'rum Vitriola'tum, La'pis Cara'leus, Sul'phate of Cop'per, Blue Stone, Blue Vit'riol, Blue Cop'peras, Ro'man Vit'riol, Mor'tooth, (F.) Sulfate de Cuivre, Couperose bleu*, is in rhomboidal, rich, blue, semi-transparent, efflorescing crystals. The taste is harsh, styptic, and corrosive. It is soluble in four parts of water at 60°, and is tonic, emetic, astringent, and escharotic. As a tonic, it has been used in epilepsy, intermittents, &c. Dose, as a tonic, gr. $\frac{1}{4}$ to gr. ij, in pill; as an emetic, gr. ij to x, in water. A very weak solution is sometimes used in ophthalmia and in gleet; and it forms the basis of *Bares's Aqua camphora'ta*, which has been recommended, diluted with sixteen parts of water, in the purulent ophthalmia of children.

CUPRI SULPHAS AMMONIACALIS. *Cuprum ammoniatum*—c. *Vitriolum, Cupri sulphas*.

CUPRUM. (properly *Cyp'rium*, being originally from *Cypros*, Cyprus,) *Chalcus, Æs, Ven'us* of the Alchemists, *Cop'per, Viride montanum, (F.) Cuivre.* Its odour is peculiar and sensible when rubbed; taste disagreeable and metallic; colour red yellow. S. g. 7.87; ductile; very malleable; less hard than iron; easily oxidized. In its metallic state, it exerts no action on the system. When swallowed, it undergoes no deleterious change. Copper cannot be dissolved whilst tin is in the mixture, and hence the utility of tinning copper vessels. Copper culinary vessels are harmless under ordinary cleanliness, provided the substances be not suffered to remain in them till cold. The salts of copper are very deleterious.

CUPRUM ALUMINATUM, Lapis divinus.

CUPRUM AMMONIATUM. *C. Ammoniacal'e seu Ammoni'acum seu ammoni'acosphu'ricum, Ammoni'ated Cop'per, Ammoni'acal Sul'phate of Cop'per, Ammonure'tum Cup'ri, Sol antiepilep'ticus of WEISS'MAN, Sul'fas Cup'ri ammoniacal'is seu*

ammoni'acæ cupra'tus, Sul'fus Cup'ri et Ammo'niæ, Subsul'fus ammoni'acop'ricus, Dentosul'fus Cup'ri et Ammo'niæ, Cup'ri Ammo'ni'osul'fas, Ammo'niæ Cup'rosul'phas, Ammo'ni'osul'phate of Cop'per, (F.) Sulfate de cuivre et d'ammoniaque, Cuivre ammoniacal. (Cupri sulph. 3ss.; Ammon. Carb. 3vj. Rub in a glass mortar till effervescence ceases. Wrap the ammoniated copper in bibulous paper, and dry with a gentle heat. Keep in a well-stopped glass bottle.—Ph. U. S.) A crystalline powder of a rich violet colour, and hot, styptic taste. By exposure to air, it becomes partly converted into carbonate of copper. It is tonic and astringent, and has been chiefly employed in epilepsy, and other obstinate spasmodic diseases. Dose, gr. $\frac{1}{4}$ gradually increased to gr. iv, in pill.

CUPRUM VITRIOLATUM, Cupri sulphas.

CUPULAR, Cupular'is, (F.) Cupulaire; (cupula, dim. of cupa, 'a vat or tun.') Of or belonging to a cupule, cup, or husk.

CUPULAR CAUT'ERY, (F.) Cautère cupulaire, is an iron in the shape of a cupule, formerly used to cauterize the skin of the cranium in certain diseases, as epilepsy, chronic headache, &c.

CURA, ('care,') Cura'tio, Merim'na, Merim'nē. Attention to, or treatment, or cure of, a disease. Keuchen defines *Cura*, 'medicine,' and *Curator*, 'the physician.' *Curatio*, also, sometimes means purification; as, *Adeps ouillus cura'tus*.

CURA AVENA'CEA, ('oat cure.') A decoction of oats and succory root, in which a little nitre and sugar are dissolved. Used as a refrigerant.

CURA DERIVATIVA, Derivation—c. *Fumigatoria, Thymiatechny.*

CURA MAG'NA, 'great cure.' A term employed, at times, for a method of treatment preferable to all others. Thus pytalism has, by many, been considered the '*cura magna*' for syphilis.

CURA MEDEANA, (after Dr. Mead.) Transfusion—c. *Palliativa*, see *Palliative*—c. *Radicalis*, see *Palliative*—c. *Revulsoria*, see *Derivation*.

CURABLE, Sanab'ilia, Aces'tos, Aces'mus, Ias'mus, Medicab'ilis, Medic'able, San'able, (F.) Guérissable. That which is susceptible of cure. An epithet applied to both patients and diseases.

CURAGE, Polygonum hydropiper.

CURA'RE, Woura'li, Woorali, Woor'a, Woura'ri, Woor'a'ri, Wura'li, Ura'ri, Urali, Oura'ri, Voor'a'ra. A very energetic vegetable poison, employed by the South American Indians to poison their arrows. See *Arrow poison*. It is said to be obtained from the bark of a species of convolvulus, called *Vejuco de Maracur*, but is referred by Martius to *Strychnos Guianensis*, and by Dr. Schomburg to *S. toxicaria seu toxifera*. The juice of *Echi'tes suberect*a, another apocynaceous plant, is said to enter into its composition. Dr. Brainard thinks it contains the venom of serpents as its main ingredient. Two new varieties of *Curare* have been investigated by Dr. Hammond of the United States Army, and Dr. S. W. Mitchell, *carroval* and *vao*,—the latter of which is less energetic, but both appear to contain the same active alkaloid—*carroval'ia*. They seem to be of vegetable origin, and to contain no trace of the venom of serpents.

CURATIE, Curative.

CURA'TIO, Mede'la, Sana'tio, A'cesis, Althex'is, Therape'a, Ii'sis, Curation, Treatment (F.) Traitement. The aggregate of means employed for the cure of disease. See *Therapeutics*.

CURATIO, Cura, Cure—c. *Contrarium per Contraria, Allopathy*—c. *Morbi per Inediam, Limotherapia.*

CURATION, Curatio.

CURATIVE, San'ative, San'atory, Heal'ing, Acesoph'orus, Acesin'ous, (F.) Curatif. Relating to the cure of a disease. *Curative Indications* are those which point out the treatment to be

employed. *Curative Treatment*, that employed in the cure of disease, in opposition to *preservative* or *prophylactic treatment*. We say, also, *Curative Process*, *Curative Means*, &c.

CURATOR INFIRMORUM, ('a care taker of the sick,') *Infirmer*.

CURCAS PURGANS, *Jatropha curcas*.

CURCULIO, ('a mite,') *Penis*.

CURCUMA ANGUSTIFOLIA, (Arab. *curcum*), see Arrow-root, *Curcuma longa*—c. *Aromatica*, *Kæmpferia rotunda*.

CURCUMA or CURCUMA LONGA, *Curcuma*, *Amomum Curcuma*, *Borri*, *Borriberri*, *Coberri*, (Hindustan,) *Crocus Indicus*, *Terra Mari'ta*, *Cannaclorus radi'cē cro'ceo*, *Mayella*, *Kua Kaha*, *Cyperus Indicus*, *κνυπος*, *Ινδικος*, of Dioscorides [?], *Turmeric*, (F.) *Racine de Safran*, *Safran des Indes*, *Souchet des Indes*. *Ord.* *Zingiberaceæ*. *Turmeric root*—the rhizoma of *curcuma longa*—*Curcuma* (Ph. U. S.), is brought from the East Indies; but is possessed of very little, if any, medicinal efficacy. It is one of the ingredients in *Curry Powder*.

CURCUMA ZEDOARIA, see *Kæmpferia rotunda*—c. *Zerumbet*, see *Kæmpferia rotunda*.

CURD, (Old E. and Prov.) *Crud*, *Curd of Milk*, *Lac pres'm*, (F.) *Caillé*, *Lait caillé*. The coagulated or curdled (*Coactus*, *Coagultus*, (F.) *Caillé* ou *Caillébotté*), part of milk.

CURD, ALUM, *Coagulum aluminosum*.

CURE, (*cura*, 'care,') *Ace'sia*, *A'cesis*, *Aces'mus*, *Cura'tio*, *C. felix*, *Sana'tio*, *San'titas*, (F.) *Guerison*. A restoration to health; also, a remedy; a restorative.

CURE-ALL, *Geum Virginianum*, *Oenothera biennis*.

CUREDENT, (*curer*, 'to cleanse,' and *dent*, 'a tooth,') *Denticulapium*.

CURE, GRAPE, see *Grape cure*.

CURE-LANGUAGE (F.), (*curer*, 'to cleanse,' and *langue*, 'the tongue,') *Lingue scalpium*, *Tonguescraper*. An instrument of ivory, tortoise-shell, &c., shaped like a knife or rasp, for removing the mucous coating which covers the tongue after sleep, &c.

CURE, MOVEMENT, *Kinesipathy*.

CURE-OREILLE, (*curer*, 'to cleanse,' and *oreille*, 'the ear,') *Ear-pick*—c. *de Petit lait*, *Whey cure*—c. *de Raisin*, *Grape cure*.

CURE, WATER, *Hydrosudotherapeia*—c. *Whey*, see *Whey cure*.

CURETTE, (from *curer*, 'to cleanse,') *Scoop*.

CURMI, (*कूर्म*), *Courmi*.

CURMBERRIES, see *Vitis Corinthiaca*.

CURRENT, BLACK, *Ribes nigrum*—c. *Red*, *Ribes rubrum*.

CURRENTS, (from *Corinth*,) see *Vitis Corinthiaca*.

CURRY or CUR'RIE POW'DER. A condiment, formed of various spices, and eaten with rice, particularly in India. The following is one of the forms of its preparations: *Sem. coriand.* 3xviij, *pip. nigr.* 3ij, *cayen.* 3j, *rad. curcumæ*, *sem. cumini*. aa 3ij, *sem. fenugr.* 3iv: mix.

CURSUMA HÆMORRHOIDALIS HERBA, *Ranunculus ficaria*.

CURSUS, (*currere*, *cursum*, 'to run,') *Running*—c. *Matricis*, *Leucorrhœa*—c. *Menstruus*, *Menses*.

CURVAMEN, (*curvare*, *curvatum*, 'to bend,') *Curvature*.

CURVATEUR DU COCCYX, *Curvator Coccygis*.

CURVATIO, *Campsis*.

CURVATOR COCCY'GIS, (F.) *Curvateur du Coccyx*. Sömmering gives this name to a small bundle of fleshy fibres, which descends on the middle of the coccyx, uniting on each side with the ischio-coccygei muscles. It is a part of those muscles.

CURVATURE, *Curvatu'ra*, *Curva'men*, *Flexu'*

ra, *Gnamp'ais*, *Cyrtu'ma*, (F.) *Courbure*. The condition of a line or surface, which approximates more or less to the form of an arc; as the curvatures of the spine, duodenum, stomach, &c. Accidental curvatures of bones are the effect of rickets, or *Mollities ossium*. The Greeks called the curvature of the spine, *ἰσος*, *σκολιός*, and *καρῖος*, according as the deviation was backwards, laterally, or forwards.

CURVATURE, POTT'S, *Vertebral disease*.

CURVE OF CARUS, see *Pelvis*.

CURVED LINES, *Lin'ee semicircula'res exter'næ* seu *arcuatæ*, (F.) *Courbes Lignes*. Two crooked lines or projections on the posterior surface of the occipital bone. They are distinguished into *superior* and *inferior*. Some lines on the os innominatum are also so called.

CURVIMER, (from *curvus*, 'bent,') *Cucumis sativus*.

CUSCOCINCHONIA, *Aricina*.

CUSCUS, *Andropogon muricatus*.

CUSCUTA, (from the Arabic,) C. *Europæa*—c. *Chlorocarpa*, see C. *Glomerata*.

CUSCUTA EPITHYMIUM, C. *minor*. The systematic name of the *Dodder of Thyme*, *Epithy'mum*, *Epithy'mum Cuscuta* seu *Cre'ticum*. *Ord.* *Convolvulaceæ*. A parasitical plant, possessed of a strong, disagreeable smell, and a pungent taste very durable in the mouth. It was once used as cathartic in melancholia.

CUSCUTA EUROPÆA seu *major* seu *vulgaris* seu *tetrandra* seu *filiformis*, *Cuscuta*, *Epithy'mum officina'rum*, was conceived to possess similar properties.

CUSCUTA FILIFORMIS, C. *Europæa*.

CUSCUTA GLOMERATA, and CUSCUTA CHLOROCARPA, *Dodder*, *American Dodder*, indigenous plants, are bitterish, subastringent, tonic, and antiperiodic.

CUSCUTA MAJOR, C. *Europæa*—c. *Minor*, C. *Epithy'mum*—c. *Tetrandra*, C. *Europæa*—c. *Vulgaris*, C. *Europæa*.

CUSHIA, *Heraclium spondylium*.

CUSHY-COW-LADY, *Coccinella septempunctata*.

CUSPARIA BARK, see C. *Febrifuga*—c. *Bark*, *False*, *Brucea antidisenterica*, and *Strychnos nux vomica*.

CUSPARIA FEBRIFUGA, *Bonplan'dia trifolia'ta* seu *angustura*, *Angustura* (Ph. U. S.), *Galipe'a febrif'uga* seu *Cusparia*. *Order*, *Rutaceæ*. The South American tree which furnishes the *Cusparia* or *Angustura Bark*, (F.) *Angusture* ou *Angosture vraie*, *Quinquina faux de Virginie*. According to Dr. Hancock, however, the *Angustura bark* is derived from *Galipe'a officinalis*, *Sex. Syst.* *Dicandria Monogynia*, *Nat. Ord.* *Diosmeæ*; and this view has been adopted in the *Pharmacopœia* of the United States. Its odour is peculiar; taste intensely bitter and slightly aromatic; pieces thin; externally, gray and wrinkled; internally, yellowish-fawn; fracture, short and resinous. It yields its virtues to water and to proof spirit. It is tonic, stimulant, and aromatic. Dose, gr. v to xx or more, in powder.

CUSP'IS, gen. *Cus'pidis*, 'a spear, a point or cusp.' The glans penis. It meant, also, a kind of bandage.

CUSSO, *Hagenia Abyssinica*.

CUSTODIA VIRGINITATIS, ('guard of virginity,') *Hymen*.

CUSTOS, gen. *Custo'dis*, 'a guardian,' *Vulva*.

CUT, ([F.] *conteau*, 'a knife,' or West Gothic, *kota*, 'to cut,' or from *κατα*, 'I cut,') [?] *Cesu'ra*, *To'mē*, *Inci'sio*, *Vulnus simplex*, *Tresis vulnus simplex*, *Cu'p'is*, (F.) *Coupure*. A common expression for the division or solution of continuity made by a sharp instrument.

CUTAMBULUS, (*cutis*, 'skin,' and *ambulo*, 'I walk,') 'Walking in the skin.' An epithet given

to certain parasitical animals, which creep under the skin—such as the Guinea-worm; and to certain pains felt between the skin and flesh, as it were.

CUTANÉ, Cutaneous.

CUTANEAL, (*cutis*, 'skin,') Cutaneous.

CUTANEO-SOURCILLIER, Corrugator supercilii.

CUTA'NEOUS, *Cuta'neal*, *Cuta'neus*, (F.) *Cutané*, (from *cutis*.) Belonging to the skin.

CUTANEOUS DISEASES, *Eruptions*, *Epiphy'mata*, are the numerous affections of a morbid character to which the skin is liable—*Dermatopathi'a*, *Dermatosis*.

Chronic cutaneous diseases have been imperfectly arranged thus. It is difficult to classify them satisfactorily.

- | | | |
|---------------------|---|-------------|
| 1. Exanthematous | { | Urticaria. |
| | { | Roseola. |
| | { | Erythema. |
| | { | Pemphigus. |
| 2. Vesicular..... | { | Rupia. |
| | { | Herpes. |
| | { | Scabies. |
| | { | Eczema. |
| 3. Pustular..... | { | Impetigo. |
| | { | Ecthyma. |
| | { | Lichen. |
| 4. Papular..... | { | Strophulus. |
| | { | Prurigo. |
| 5. Squamous..... | { | Lepra. |
| | { | Psoriasis. |
| | { | Pityriasis. |
| | { | Acne. |
| | { | Sycosis. |
| 6. Folliculous..... | { | Ichthyosis. |
| | { | Trichosis. |
| | { | Favus. |

CUTANEOUS EXHALATION AND ABSORPTION are those which take place from the skin.

CUTANEOUS, MIDDLE POSTERIOR, see Sciatic nerve, lesser.

CUTANEOUS NERVES, *Cuta'neal Nerves* so called, of the upper extremity, are two in number. They are given off from the brachial plexus. The *internal cutaneous*, *Cubito-cutané*—(Ch.) descends along the inner part of the arm, and divides above the elbow into two branches; the outermost of which follows the outer edge of the biceps, and spreads over the anterior and inner part of the forearm; and the innermost accompanies the basilic vein, passing to the corresponding side of the hand and little finger. The *external cutaneous*, *Radio-cutané* (Ch.) *Musculo-Cuta'neus*, *Perforans CASSE'RII*, passes outwards; perforates the coraco-brachialis; descends along the anterior and outer part of the arm; passes as far as the middle fold of the elbow under the median cephalic; and descends along the anterior and outer edge of the forearm. At some distance above the wrist, it divides into two branches; an *outer*, which is distributed on the back of the hand, the thumb and index finger; and an *inner*, which descends on the muscles of the thumb into the palm of the hand, and divides into fine filaments, which may be traced to the fingers.

Cutaneous nerves, *Mid'dle cutaneous*, two in number, are branches of the crural or femoral nerve. They are distributed to the integument of the middle and lower part of the thigh and of the knee.

Nervus Cuta'neus Mi'nor, *Les'ser inter'nal cutaneous nerve*, *Nerve of Wris'berg*, takes its origin from the axillary plexus, but is more particularly connected with the ulnar nerve. It soon separates from the ulnar, running afterwards between it and the inner side of the arm. A little below the axilla it divides into two branches, which are distributed to the arm.

Chaussier calls the *cephalic* and *basilic veins* the *Ra'dial* and *Cu'bial Cuta'neous*.

CUTCH, Catechu.

CUTE, Malleolus.

CUTI, Catechu.

CUTICLE, (dim. of *cutis*,) Cuticula—c. of the Enamel, see Enamel.

CUTICLE, Liquid, C. Collodion.

CUTICULA, Cuticle, Epidermis.

CUTIO, Oniscus.

CUTIS, *Pell'is*, *Pell'a*, *Co'rium*, *Der'ma*, *De'ria*, *Do'ra*, *Anthro'pe*, *Anthro'pen*, *Skin*, *Derm*, (F.) *Peau*, *Derme*. A dense, resisting membrane, of a flexible and extensible nature, which forms the general envelope of the body; and is continuous with the mucous membranes, through the different natural apertures. It is generally considered to be formed of three distinct layers—the *epidermis*, *rete*, or more properly *corpus mucosum*, and *corium* (*cutis vera*, *derma*). Some anatomists, however, separate it into several others. Its outer surface is covered by a number of small eminences, called papillæ, which are generally regarded as essentially nervous and vascular. The skin is a medium of communication with external bodies. It protects the subjacent parts; is the seat of touch; and through it are exhaled the watery parts of the blood, which are not needed in the nutrition of the body. The state of the skin, as regards heat and dryness, affords useful information in pathological investigations. Its colour, too, requires attention: the paleness of disease is as characteristic as the rosy complexion of health. The colour of the skin varies according to the age, sex, &c. As a general rule, it is finer in the female and child than in the male and adult. In old age it becomes light-coloured, thin, and dry. It likewise varies according to races, &c.

CUTIS ANSERINA, *Horrida cutis*—c. Carnosa, *Panniculus carnosus*—c. Estima, *Epidermis*—c. Gallinacea, *Horrida Cutis*—c. Linguae, see Tongue—c. Suilla, *Couenne*—c. Summa, *Epidermis*—c. Tensa Chronica, *Induration of the cellular tissue*—c. Ultima, *Epidermis*—c. Vera, *Corium*, see *Cutis*.

CUTITIS, (*cutis*, and *itis*,) *Cytitis*, *Erysipelatous inflammation*.

CUTLINS, Groats.

CUTTING ON THE GRIPE, see Lithotomy.

CUTTLE FISH, Sepia.

CUTTUBUTH, *Cut'ubuth*, *Kut'ubuth*, *Leucomoria*, *Melancho'lia errabunda*. The Arabian physicians gave this name to a species of melancholy, accompanied with so much agitation, that the patients could not remain tranquil for the space of an hour.

CUURDO CANELLA, *Laurus cinnamomum*.

CYAN'IC, *Cyan'icus*, (F.) *Cyanique*; (from *κωανος*, 'blue,') An epithet for the blue stage of a malady—as cholera.

CYANEPHIDROSIS, (*cyano*, *eri*, an augmentative preposition, and *ἵδρω*, 'I sweat,') Copious perspiration, which colours the linen blue.

CYANO, (*κωανος*, 'blue,') in composition, blue.

CYANODERMIA, (*cyano*, and *δερμα*, 'skin,') *Cyanopathy*.

CYANOGEN, (F.) *Cyanogène*; (*cyano*, and *γεννω*, 'I generate,') So called from its being an ingredient in Prussian blue. It forms, with oxygen, the cyanic and other acids; with hydrogen, the hydrocyanic acid.

CYANOPATHY, *Cyanopathi'a*, *Cyano'sis*, *C. cordi'aca*, *Cyanoder'mia*, *Hæmatocyano'sis*, *Hæmatocyanoopathi'a*, *Kyano'sis*, *Mor'bus seu lcterna cœru'leus*, *Cœru'le'sis neonato'rum*, (*cyano*, and *παθος*, 'affection,') *Blue Jaw'dice*, (F.) *Cyanose*, *Maladie bleu*, *lctère bleu*. A disease in which the surface of the body is coloured blue. It is often symptomatic, and commonly depends on a direct com-

munication remaining between the cavities of the right and left side of the heart; *Pa'tency of the for'a-men au'te*, *Acleitocardie* of Piorry. Such communication does not, however, always occasion the *blue disease*, but it is generally thought that the disease never exists without this state of parts; or without some obstacle to the circulation in the right side of the heart. The blueness* does not seem to be owing to the admixture of black and red blood. A child affected with blueness is said to be *cyano'sed*, (F.) *Cyanosé*.

CYANOSÉ, see Cyanopathy.

CYANOSIS, (κυανωσις, 'blueness,') Cyanopathy — c. Pulmonalis, Atelectasis pulmonum.

CYANOTIC, *Cyanot'icus*, (F.) *Cyanotique*. Relating or appertaining to cyanosis.

CYANURETUM FERROZINCICUM, Zinci ferro-hydrocyanas.

CYANUS, (κυανος, 'blue,') *Centaurea cyanus* — c. *Ægyptiacus*, *Nymphæa nelumbo*.

CY'ANUS SEG'ETUM, *Blue'bottle*, (F.) *Bluet* ou *Bluet des Moissons*, *Barbeau*, *Aubifoin*, *Casse-Linettes*. Ord. Compositæ. The blue flowers (κυανος, 'blue,') of this European plant, when distilled with water, have been used in ophthalmia.

CY'AR, (κυαρ, 'the eye of a needle,' 'a small hole,') The *Meatus auditorius internus*. See Auditory canal (internal).

CYATHIS'CUS, (dim. of κυθος, 'a bowl,') The concave part of a sound, made like a small spoon, as in the case of the ear-pick.

CY'ATHUS, (κυαθος, 'a bowl,') A measure, both of the liquid and dry kind, equal to about an ounce and a half, or to the tenth part of a pint. According to Pliny and Galen, about 10 drachms.

CYATHUS CEREBRI, Infundibulum of the brain.

CYBE, (κυβη,) Head.

CYBITON, (κυβιτον,) Cubitus.

CYBOID, (κυβος, 'a cube,' and εἶδος, 'resemblance,') Cuboid.

CYCAS (κυκας) CIRCINALIS, see Sago — c. *Revoluta*, see Sago.

CYC'EON, (κυκων, from κυκω, 'I mix together,') The ancient name of a medicine of the consistence of pap, composed of wine, water, honey, flour, barley meal, and cheese.

CYCLAMEN EUROPÆUM, *Arthani'ta*, *A. cyc'lamen*, *Cyclam'inus*, (κυκλαμης, κυκλαμινος, κυκλαμινον, from κυκλος, 'a circle,' from the shape of the leaves.) *Cysson'themum*, *Cyssonophyl'lon*, *Pa'nis porci'nus*, *Cas'amum*, *Chylli'ne*, *Sowbread*. Ord. Primulacææ. Sex. Syst. Pentandria Monogynia. (F.) *Pain du Porceau*. The fresh root is said to be acrid, bitter, drastic, and anthelmintic. Dose, ʒi. For external use, see Arthanita.

CY'CLE, *Cy'clus*, (κυκλος, 'a circle,') A period or revolution of a certain number of years or days. The Methodists gave this name to an aggregate of curative means, continued during a certain number of days. Nine was the usual number.

CYCLE, HEBDOM'ADAL or HEP'TAL. A period of seven days or years, which according to some, either in its multiple or submultiple, governs an immense number of phenomena of animal life.

CYCLISCUS, (κυκλιασος, dim. of *Cyclus*,) *Cyclis-mus*.

CYCLIS'MUS, (from *cyclus*.) A *Troch*, *Trochis'cus*. This name, as well as *Cyclis'cus*, was also given to a circular kind of rasp. They have the same etymon as cycle.

CYCLOCEPH'ALUS, (F.) *Cyclocephale*, (*cyclus*, and κεφαλη, 'head,') A monster whose eyes are in contact, or united in one.

CYCOGANGLIONIC NERVES, see Nerves.

CYCLOPHOR'IA, (*cyclus*, and φερω, 'I bear,') The circulation of the blood or other fluids.

CYCLO'PIA. Same etymon as *Cyclops*. State of a monster that has both eyes united into one. Called, also, *Monops'ia* and *Rhinencephal'ia*.

CYCLOPIA GENISTOI'DES; *Nat. Ord.* Leguminosæ. A South African plant, the decoction and infusion of which are used as expectorants in chronic bronchitis and phthisis.

CY'CLOPS, gen. *Cyclo'pis*, (κυκλωψ, from *cyclus*, and ωψ, 'an eye,') *Monoc'e'ulus*, *Mon'ops*, *Mon'ophthal'mus*, *Monom'matus*, *Unio'e'ulus*, *Uno'e'ulus*. A monster having but one eye, and that placed in the middle of the forehead, as in the fabulous Cyclops.

CYCLOSIS, (κυκλος, 'a circle,') Circulation.

CYCLOTOME, *Cyclo'tomus*, (from *cyclus*, and τεμνω, 'to cut,') An instrument, composed of a ring of gold and a cutting blade, by means of which the ball of the eye can be fixed, whilst the cornea is cut. It was invented by a surgeon of Bordeaux, named Guérin, for extracting the cataract. It is not used.

CYCLUS, (κυκλος,) Circulus.

CYDER, Cider.

CYDONIA, (from *Cydon*, in Crete,) *Pyrus cydonia* — c. *Maliformis*, *Pyrus cydonia* — c. *Vulgaris*, *Pyrus cydonia*.

CYDONIA'TUM, *Cydonia'tum*, *Diacydoni'tes*. A composition of the *Cydonia mala* or quinces, with the addition of spices.

CYDONIUM, see *Pyrus Cydonia* — c. *Malum*, see *Pyrus cydonia*.

CYE'MA, gen. *Cye'matis*, (κυημα, from κυω, 'I conceive,') *Concep'tion*. Likewise the product of conception. See Embryo, and Foetus.

CYESIOGNO'MON, (*cyesis*, and γνωμων, 'a sign, a token,') A sign of pregnancy.

CYESIOGNO'SIS, (*cyesis*, and γνωσις, 'knowledge,') Diagnosis of pregnancy.

CYESIOL'OGY, *Cyesiology'ia*, (*cyesis*, and λογος, 'a description,') The doctrine of gestation.

CY'ESIS, (κυησις,) Conception, Fecundation, Pregnancy.

CYESTEINE, Kiesteine.

CYLICH'NE, *Cyllich'nis*, *Cyllich'nion*, (κυλικηνη, κυλικηνης, κυλικηνιον, dim. of κυλιξ, gen. κυλικος, 'a cup,') A pillbox or earthenware pot, or small cup.

CYLINDRI, (κυλινδρος, pl. κυλινδροι, 'cylinder,') see Villous membranes — c. *Membranacei renum*, see Calix.

CYLINDROID, *Cylindroi'des*, (κυλινδρος, 'cylinder,' and εἶδος, 'form,') Having the form of a cylinder. Chaussier calls the Cornu Ammonis, *Protuberance cylindroïde*.

CYLINDROIDES, Teres.

CYLINDROMA, (κυλινδρος, 'a cylinder,' and ομα,) see Chondroma.

CYLLOEPUS, (κυλλος, 'crooked,' and πους, gen. ποδος, 'a foot,') see Kylosis.

CYLOPODA, see Kylosis.

CYLOPODION, see Kylosis.

CYLOSIS, (κυλλωσις,) Lameness, mutilation, or vicious conformation. See Kylosis.

CYLOSOS'MUS, (κυλλος, 'crooked,' and σωμα, 'body,') A malformation by defect, in which the fissure and eventration are lateral, chiefly in the lower part of the abdomen, the inferior extremity of the side affected with the fissure absent, or very little developed.

CYMATO'DES, *Undo'sus* (κυματωδης, from κυμα, 'a wave,' and οδος.) The vacillating, undulatory character of the pulse in adynamic individuals.

CYMEA, (κυμβος, 'a cavity,' or κυμβη, 'a canoe,') Vulva.

CYMBALARIA ELATINE, Antirrhinum elatine — c. *Muralis*, Antirrhinum linaria.

CYMBALARIS, (from κυμβαλον, 'a cymbal,') Cricoid.

CYMBECEPHALIC, (κυμβη, 'a canoe,' and κεφαλη, 'a head,') Kumbeccephalia.

CYMBIFORME OS. (κύβη, 'a canoe,' and *forma*, 'shape.') Scaphoid bone.

CYMOPOGON SCHÖENANTHUS, *Juncus odoratus*.

CYMINUM, *Cuminum cuminum*.

CYMLING, see *Cucurbita pepo*.

CYMOGRAPHION, (κύμα, 'a wave,' and γράφειν, 'to describe,') Kymographion.

CYNANCHĒ, *Angi'na*, (cyno, and αγγ, 'I suffocate' [?] 'dog choak,') *Sorethroat*; *Paracynanchē*, *Synanchē*, *Prunell'a*, *Empres'ma Paristhmī'tis*, *Cau'ma Paristhmī'tis*, *Isthmī'tis*, *I aristhmī'tis*, *Inflammatio Fau'cium*. Inflammation of the supradiaphragmatic portion of the alimentary canal, and of the lining membrane of the upper part of the air-passages:—(F.) *Angine*, *Esquinancie*, *Mul de Gorge*, *Synancie*, *Squinancie*, *Kinancie*. It comprises the following chief varieties.

CYNANCHE EPIDEMICA, *Cynanche maligna*—c. *Externa*, *C. Parotidæa*—c. *Faucium*, *C. tonsillaris*—c. *Gangrenosa*, *Cynanche maligna*—c. *Laryngea*, *Laryngitis*.

CYNANCHE MALIG'NA, *C. gangrenosa* seu *epidem'ica* seu *pur'puroparotidæ'a* seu *ulcero'sa*, *Cacocynanchē*, *Pharyngoc'acē*, *Cacopharyng'ia*, *Fel'bris epidem'ica* cum *Angi'nā*, *Empres'ma Paristhmī'tis Malig'na*, *Tonsillarum gangrē'sa*, *Tonsill'æ pestilent'es*, *Epidem'ica gut'turis lu'es*, *Pestilens fau'cium affect'us*, *Pædan'chonē* [?], *Gangrē'sa Tonsillarum*, *Ulcus Syri'acum*, *Morbus Syri'acus seu suffocati'vus seu suff'ocans*, *Garotil'lo*, *Carbun'culus angino'sus*, *Angi'na ulcero'sa* seu *ulcusculosa* seu *epidem'ica* seu *gangrē'sa* seu *maligna* seu *fau'cium maligna*, *Isthmot'yphus*, *Pu'trid* or *Ulcero'sus Sorethroat*, *Gan'grenous Inflammation of the Pharynx*, (F.) *Angine gangrēneuse* ou *Maligne*. It is characterized by crimson redness of the mucous membrane of the fauces and tonsils; ulcerations, covered with mucus; and spreading sloughs, of an ash or whitish hue; the accompanying fever typhus. It is often epidemic, and fancied to be contagious [?]; and is frequently found accompanying scarlet fever,—giving rise to the variety, *Scarlati'na maligna*, *Canker rash*. *Cynanche maligna* has been made to include both diphtheritic and gangrenous pharyngitis. See *Pharyngitis*, *diphtheritic*.

The general treatment is the same as in typhus; and stimulant antiseptic gargles must be used, consisting, for example, of the decoction of bark and muriatic acid, a solution of chlorate of potassa, &c.

CYNANCHE MAXILLARIS, c. *Parotidæa*—c. *Æsophaga*, *Æsophagitis*.

CYNANCHE PAROTIDÆ'A, *C. Parotidis* seu *exter'na* seu *maxilla'ris*, *Empres'ma Parotit'is*, *Parotit'is*, *P. epidem'ica* seu *erysipelato'sa* seu *contagio'sa* seu *spu'ria* seu *sero'so glutinē tu'mens*, *Parot'ia*, *Parotidit'is*, *Angi'na maxilla'ris* seu *exter'na* seu *parotidæ'a exter'na*, *Gis'sa*, *Erythrochæ'ras*, *Gencycynanchē*, *Gnathocynanchē*, *Inflammatio parotidum*, *Inflatio parotidum*, *Catar'rhus Bellinsul'danus*, *Mumps*, *Branks* (Scotch), (F.) *Inflammation de la Parotidē*, *Oreillons*, *Ourles*. The characteristic symptoms are:—a painful tumor of the parotid gland, or of the areolar tissue surrounding it, or of both, not of the suppurative kind; frequently extending to the maxillary gland. It is very conspicuous externally, and is often accompanied with swelling of the testis in the male, and of the mamma in the female; the testes being sometimes absorbed afterwards. It is generally epidemic, and apparently contagious. [?] The treatment is very simple; the adoption merely of the antiphlogistic plan, under which it usually soon disappears. When inflammation of the mamma or testes supervenes, it must be treated as if idiopathic.

Epidem'ic Parotit'is or *Mumps* is also termed *Parotit'is polymor'pha* seu *epidem'ica* seu *specif'ica*.

CYNANCHE PHARYNGE'A, *Empres'ma Paristhmī'tis Pharyng'e'a*, *Isthmī'tis*, *Pharyngit'is*, *Inflammatio Pharyng'is*, *Paracynanchē*, *Angi'ne inflammat'oria*, *Sp. iv. of Boerhaave*. (F.) *Angine Pharyngée*, *Catarrhe pharyngien*. Inflammation of the pharynx.

This disease can hardly be said to differ, in pathology or treatment, from *Cynanche tonsillaris*. The same may be remarked of *Cynanchē* (*Æsophaga*), *Æsophagitis*.

CYNANCHE PRUNELLA, *C. Tonsillaris*—c. *Purpuro-parotidæa*, *Cynanche maligna*—c. *Simplex*, *Isthmitis*—c. *Stridula*, *C. trachealis*.

CYNANCHE TONSILLARIS, *C. fau'cium* seu *Prunell'a*, *Empres'ma Paristhmī'tis Tonsilla'ris*, *Synanchē*, *Amygdalit'is*, *Bran'ci*, *Bran'chi*, *Hyanchē*, *Tonsillit'is*, *Tonsill'a*, *Squinanth'ia*, *Paristhmia*, *Paristhmī'tis*, *Dysphag'ia inflammat'oria*, *Antiadit'is*, *Angi'na inflammat'oria* seu cum *tumore seu tonsilla'ris* seu *synocha'lis* seu *sanguinea* seu *ve'ra et legit'ima*, *Inflammatio Tonsillarum*, *Antiadon'cus inflammat'orius*, *Inflammatory Sorethroat*, *Com'mon Squin'ancy*, *Squin'zey*, *Squin'sy* or *Quin'sy*, (Old E.) *Svensie*, *Sicinacie*, (F.) *Amygdalite*, *Inflammation des Amygdales*, *Angine tonsillaire*, *Pharyngite tonsillaire*, *Esquinancie*, *Squinancie*, *Catarrhe guttural*, *Angine gutturale inflammatoire*. The characteristic symptoms of this affection are, swelling and florid redness of the mucous membrane of the fauces, and especially of the tonsils; painful and impeded deglutition, accompanied with inflammatory fever. It is generally ascribed to cold, and is one of the most common affections of cold and temperate climates. It usually goes off by resolution, but frequently ends in suppuration.

Common sore throat is an affection of no consequence. It requires merely rest, and the observance of the antiphlogistic regimen. When more violent,—in addition to this,—bleeding, local or general, or both,—purgatives, inhalation of the steam of warm water; acid, or emollient gargles; rubefacients externally, or sinapisms or blisters. When suppuration must inevitably occur, the continued use of the inhaler must be advised, and an opening be made into the abscess as soon as pus shall have formed. If the patient be likely to be suffocated by the tumefaction, bronchotomy may be necessary.

CYNANCHE TRACHEALIS, *Cynanchē larynge'a* of some, *C. strid'ula*, *Suffocatio strid'ula*, *Asth'ma infantum*, *Catar'rhus suffocati'vus Barbaden'sis*, *Angi'na polypo'sa* seu *membrana'cea* seu *pulpo'sa* seu *pernici'osa* seu *inflammat'oria* seu *suffocato'ria* seu *strepito'sa* seu *cani'na* seu *exsudato'ria* seu *trachealit'is*, *Empres'ma bronchit'is seu bronchlemmit'is*, *Laryngit'is et Tracheit'is infantilis*, *Laryngit'is exsudati'va* seu *membrana'cea* seu *polypo'sa*, *Laryngos'tasis* seu *Laryngotracheit'is*, *Laryngotracheitis* with *diphtherit'ic exudation*, *Laryngoc'acē*, *Orthopnœ'a cynanch'ica*, *Cynanch-orthopnœ'a*, *Pædan'chonē* [?], *Morbus Strangulato'rius*, *Trachit'is*, *Tracheit'is*, *Trachelit'is*, *Tracheop'gra*, *Tracheal'ia*, *Diphtherit'is tracheal'ia*, *Expectoratio Sol'idæ*, *Canni'a Bronchit'is*, *Croup*, *Roup*, *Hives*, *Bold Hives*, *Chokk*, *Stiffing*, *Rising of the lights*, (F.) *Angine laryngée et trachéale*, *Laryngite avec production de fausses membranes*, *Laryngite pseudo-membraneuse*. A disease characterized by sonorous and suffocative breathing; harsh voice; cough, ringing, or like the barking of a dog; fever, highly inflammatory. It is apt to be speedily attended with the formation of a false membrane, which lines the trachea beneath the glottis, and occasions violent dyspnoea and suffocation, but is sometimes expectorated. The dyspnoea, as in all other affections of the air-passages, has evident exacerbations. It differs in its character in different situations; being infinitely more inflammatory in some than in others,

and hence the success obtained from different modes of treatment. It chiefly affects children, and is apt to recur, but the subsequent attacks are usually less and less severe.

As a general rule, it requires the most active treatment; bleeding from the arm or neck, so as to induce paleness; leeches applied to the neck, &c., according to the age,—the warm bath, blisters to the neck or chest, and purgatives. Formidable as the disease may be, if this plan be followed early, it will generally be successful. Many specifics have been recommended, but the search has been idle and fruitless. The majority of cases of what are called croup are not of this inflammatory cast; but are more of a spasmodic character, and have been termed by the French *faux croup*, false croups, and with us are occasionally termed *spasmodic croup*. They generally yield to an emetic and the warm bath.

CYNANCHE TRACHEALIS SPASMODICA, Asthma acutum, A. Thymicum—c. Ulcerosa, Cynanche maligna.

CYNANCHICA. Medicines used in cases of quinsy were formerly so called.

CYNANCHORTHOPNŒA, (*cynanche*, *orthos*, 'up-right,' and *pnœa*, 'I breathe.') Cynanche trachealis.

CYNANCHUM. Same etymon as Cynanche. A genus of plants, *Ord.* Asclepiadaceæ, of which the following are used in medicine.

CYNANCHUM ARGEL, C. oleæfolium—c. Ipecacuanha, Tylophora asthmatica.

CYNANCHUM MONSPELIACUM, *Scam'mony of Montpellier*. The plant furnishes a blackish kind of gum-resin, which is purgative, and but little used.

CYNANCHUM OLEÆFOLIUM, C. argel, *Solenostem'ma argel*, *Argel*. An Egyptian, Nubian and Arabian shrub, the leaves of which form a portion of most samples of Alexandrian senna. They resemble senna in their action.

CYNANCHUM TOMENTOSUM, Tylophora asthmatica—c. Vincetoxicum, *Asclepias vincetoxicum*—c. Viridiflorum, Tylophora asthmatica—c. Vomitorium, Tylophora asthmatica.

CYNANTHEMIS, (*cyno*, and *anthesis*), *Anthemis cotula*.

CYNANTHROPIA, (*cyno*, and *anthropos*, 'man.') A variety of melancholia, in which the patient believes himself changed into a dog; and imitates the voice and habits of that animal.

CYN'ARA, (*κύναρ*, 'dog-briar.') C. *Scolymus*, *Cin'ara scolymus*. The systematic name of the Artichoke, *Alcoec'olum*, *Articoec'ulus*, *Artisoc'cus le'vis*, *Costus nigra*, *Car'duus sativus non spinosus*, *Cinara hortens'is*, *Scolymus sativus*, *Car'duus sativus seu domest'icus cap'itè majo'rè seu al'tilis*, anciently *Cactus*, *κακτος*, (F.) *Artichaut*. *Ord.* Compositæ. *Sex. Syst.* Syngenesia Polygamia æqualis. Indigenous in the southern parts of Europe. Much used as an agreeable article of diet. The juice of the leaves, mixed with white wine, has been given in dropsies.

CYN'ICUS, (*κύνικος*), *Cyno'des*, *Cyn'ic*, (F.) *Cynique*. Relating to, or resembling a dog. *Cyn'ic spasm* is a convulsive contraction of the muscles of one side of the face,—dragging the eye, cheek, mouth, nose, &c., to one side. See Canine.

CYNIPS (*κύνιψ*) QUERCUS FOLII, see Quercus infectoria.

CYNO, (*κύν*, gen. *κύνος*, 'a dog.') In composition, dog.

CYNOCOPRUS, (*cyno*, and *κοπρος*, 'excrement.') *Album Græcum*.

CYNOCRAMBE, (*κυνοκράμβη*, from *cyno*, and *κράμβη*, 'cabbage.') *Mercurialis perennis*.

CYNOCTONON, (*cyno*, and *κτενω*, 'I kill.') *Aconitum*.

CYNOCYTISUS, (*cyno* and *cytissus*), *Rosa canina*.

CYNODEC'TOS, (*κυνόδεκτος*, from *cyno*, and *δεκτω*, 'I bite.') One who has been bitten by a dog.

CYNODES, (*κυνώδης*, from *cyno* and *odes*.) *Cynicus*.

CYNODESMION, (*cyno*, and *δεσμος*, 'a band.') *Frænum penis*.

CYNODESMUS, *Frænum penis*.

CYNODON DAC'TYLOŒ, (*cyno*, and *οδους*, gen. *οδοντος*, 'a tooth,') *Pan'icium seu Digita'ria dactylon*, *Bermu'da grass*; indigenous. *Ord.* Gramineæ. The roots have been used as a substitute for sarsaparilla.

CYNODONTES, Canine teeth.

CYNOGLOS'SUM, (*κυνόγλωσσον*, from *cyno*, and *γλωσσα*, 'a tongue.') *Cynogloss'um officina'le seu bi'color*, *Lingua Cani'na*, *Hound's tongue*, *Cabal'la'tion*, (F.) *Langue de Chien*. *Ord.* Boraginææ. *Sex. Syst.* Pentandria Monogynia. It is aromatic and mucilaginous, and has been supposed to be possessed of narcotic properties.

CYNOGLOSSUM AMPLEXICAULE, C. *Virginicum*.

CYNOGLOSSUM VIRGIN'ICUM, C. *amplexicaul'e*, *Wild Com'frey*; indigenous; has mucilaginous properties. The leaves are smoked, and are said to produce effects like those of tobacco. [?]

CYNOLOPHOI, (*cyno*, and *λοφος*, 'an eminence.') Spinous processes of the vertebrae.

CYNOLYSSA, (*κυνόλυσσα*, from *cyno*, and *λυσσα*, 'madness.') *Hydrophobia*.

CYNOMETRA AGALLOCHUM, *Agallochum*.

CYNOMOIA, *Plantago psyllium*.

CYNOMORIUM COCCIN'EUM, (*cyno*, and *μαριον*, 'penis.') (F.) *Champignon de Malte*. *Ord.* *Balanophoræ*. Improperly called *Fungus Meliten'sis*, or *Fungus of Malta*, as it is not a fungus. The powder has been given as an astringent in hemorrhage, dysentery, &c.

CYNOREXIA, (*cyno*, and *ορεξις*, 'appetite.') *Boulimia*.

CYNORRHODON, (*κυνορρόδον*, from *cyno*, and *ροδον*, 'a rose.') *Rosa canina*.

CYNOBATOΣ, (*κυνοβατος*, from *cyno*, and *βατος*, 'a bramble.') *Rosa canina*.

CYNOSORCHIS, (*cyno*, and *ορχις*, 'a testicle.') *Orchis mascula*.

CYNOSPASTUM, (*cyno*, and *σπασω*, 'I draw.') *Rosa canina*.

CYON, (*κύν*.) The word sometimes signifies the frænum of the prepuce; at others, the penis.

CYOPHORIA, (*κυφορια*, from *κυος*, 'a fœtus,' and *φερω*, 'I carry.') *Pregnancy*.

CYOTROPHY, *Cyotroph'ia*, *Embryot'rophy*, *Embryotroph'ia*, *κυος*, 'fœtus,' and *τροφειν*, 'to nourish.') *Nutrition of the embryo*. *Fœtal nutrition*.

CYPARISSUS, (*κυπαρισσος*), *Cupressus sempervirens*.

CYPERUS ANTIQUORUM, (*κυπερος*), *Lawsonia inermis*—c. *Indicus*, *Curcuma longa*.

CYPERUS LONGUS, C. *Roma'nus*, *Galangale*, (F.) *Souchet odorant*. *Ord.* *Cyperaceæ*. It possesses aromatic and bitter properties, but is not used. See also, *Dorstenia contrayerva*.

CYPERUS ODORUS, *Dorstenia contrayerva*—c. *Peruanus*, *Dorstenia contrayerva*—c. *Romanus*, C. *longus*.

CYPERUS ROTUNDUS, C. *Tetras'tachys*, the *Round Cyper'us*, (F.) *Souchet rond*. It is a more grateful aromatic bitter than the C. *longus*.

CYPERUS TETRASTACHYS, C. *rotundus*.

CYPHEL, *Sempervivum tectorum*.

CYPHO'MA, gen. *Cypho'matis*, (*κυφωμα*, from *κυφος*, 'gibbous,' and *ομα*), *Cyphos*, *Cypho'sis*, *Cyrto'ma*, *Cyrto'sis*, *Opisthocypho'sis*. *Gibbosity of the spine*. *Cyphoscolio'sis*, *Kyphoskolio'sis*. See *Cyphoma*, *Gibbositatis* and *Vertebral disease*. *Cyrtosis* forms a genus in the order *Dysthet'ica*, and class *Hæmat'ica* of Good, and is defined: 'head

bulky, especially anteriorly; stature short and incurvated; flesh flabby, tabid, and wrinkled.' It includes *Cretinism*, and *Rickets*.

CYPHOSCOLIOSIS, (*κυφωσις*, 'gibbous,' *σκολιως*, 'crooked,' and *osis*.) Cyphoma.

CYPIHOSIS, (*κυφωσις*), Gibbosity, see **Hump**.

CYPRËS, *Cupressus sempervirens*.

CYPRESS, *Cupressus sempervirens*, *Taxodium distichum*—c. American, *Schubertia disticha*—c. Bald. *Taxodium distichum*—c. Tree, *Liriodendron*.

CYPRINUM (O'LEUM), (*κυπρινον* (*ελαιον*)) *Oil of Cypress*; prepared of olive oil, cypress flowers, calamus, myrrh, cardamom, inula, bitumen of Judea, &c. It was formerly employed in certain diseases of the uterus, pleurisy, &c., and was regarded to be both stimulant and emollient.

CYPRIPEDIUM ACAUL'Ë, (*Κυπρις*, 'Venus,' and *πόδιον*, 'a sock or buskin.') *C. lu'mile*, *Stem-less Ladies' Slipper*; *Mocassin Flower*; *No'ah's Ark*. Ord. Orchidaceæ. Indigenous; flowers in May and June. The roots are used by steam-doctors in nervous diseases, like valerian.

CYPRIPEDIUM CALCEOLUS, *C. Luteum*, *C. spectabile*—c. *Canadense*, *C. spectabile*—c. *Flavescens*, *C. Luteum*—c. *Humile*, *C. acaule*.

CYPRIPEDIUM LUTEUM, *C. Calceolus* seu *Flavescens* seu *Pubescens* seu *Parviflorum*, *Yellow Ladies' slipper*, *Mocassin flower*, *Moccasin plant*, *Yellow's*, *Bleeding heart*, *American valerian*, *Yellow umbil*, *Male Mer'vine*, *No'ah's ark*, (F.) *Sabot de Vénus jaune*. An indigenous plant, which is found all over the United States; blossoming in May and June. The root *Cypripedium* (Ph. U. S.), is considered to be antispasmodic, and is used in the same cases as valerian.

CYPRIPEDIUM PARVIFLORUM, *C. Luteum*—c. *Pubescens*, *C. Luteum*.

CYPRIPEDIUM SPECTAB'ILE, *C. Calceolus* seu *Canadense*, *Show'y ladies' slipper*, is the most beautiful of the genus. It is said to possess properties identical with those of *C. Luteum*.

CYPRIMUM, (from *Cyprus*), *Cuprum*.

CYPSLE, (*κυψελη*), *Cerumen*.

CYRCOM, *Anus*.

CYRTOMA, gen. *Cyrto'matis*, (*κυρτωμα*, from *κυρτος*, 'curved,' and *oma*.) *Curvature*, *Cyphoma*.

CYRTONOSUS, (*κυρτος*, 'curved,' and *nosus*, 'disease,') *Rachitis*.

CYRTOSIS, *Cyphoma*.

CYSSANTHEMON, (*κυσσανθος*, 'ivy,' and *ανθεμον*, 'a flower,') *Cyclamen*.

CYSSARUS, (*κυσσαρος*), *Anus*, *Rectum*.

CYSSOPHYLLON, (*κυσσοις*, 'ivy,' and *φυλλον*, 'a leaf,') *Cyclamen*.

CYS'SOTIS, gen. *Cyssot'idis*, (*κυσος*, 'the anus,') The lower part of the rectum. *Tenesmus*.

CYS'SUS, (*κυσσοις*), *Anus*.

CYST, *Kyst*.

CYSTAL'GIA, *Cystidal'gia*, (*cystis*, and *αλγος*, 'pain,') *Pain in the bladder*.

CYSTANENCEPHALIA, (*cystis*, and *anencephalia*, 'absence of brain,') A monstrosity, in which, in place of a brain, a bladder is found filled with fluid.

CYSTATROPH'IA, (*cystis*, and *ατροφια*, 'atrophy,') *Acysturorophic*. *Atrophy* or wasting of the bladder.

CYSTAU'CHEN, (*cystis*, and *αυχην*, 'the neck,') The neck of the bladder.

CYSTAUCHENITIS, (from *cystauchen*, and *itis*.) *Inflammation of the neck of the bladder*.

CYSTAUCHENOTOM'IA, *Cystotrachelotom'ia*, *Cystidotrachelotom'ia*, (*cystis*, *αυχην*, 'the neck,' and *τομη*, 'incision,') An incision into the neck of the bladder. See *Lithotomy*.

CYSTAUX'Ë, *Hypertroph'ia vesicæ urinariæ*, *Enlargement vesicæ: cystis*, and *αυξη*, 'increase,') *Hypertrophy of the coats of the urinary bladder*. See *Cysthypersarcosis*.

CYSTE, *Kyst*.

CYSTECTASY, (*cystis*, and *εκτασις*, dilatation,') *Lithectasy*.

CYSTENCEPH'ALUS, (*cystis*, and *κεφαλη*, 'head,') A monster having a head with a vesicular brain.

CYSTEOL'ITHUS, (*cystis*, and *λιθος*, 'a stone,') *Stone in the bladder*. Also, a medicine, employed to dissolve or break stone.

CYSTERETHIS'MUS, *Irritabilitas* seu *Impatient'ia Vesicæ*, (*cystis*, and *ερεθιζω*, 'I irritate,') *Irritability of the bladder*.

CYSTHEPAT'ICUS, (*cystis*, and *ήπαρ*, 'the liver,') Belonging to the gall-bladder and liver. This name was given, formerly, to imaginary excretory ducts for the bile, which were supposed to pass directly from the liver to the gall-bladder.

CYSTHEPATOLITHIASIS, *Cystidepatolithiasis*, *Cholelith'ia*, *Cholelith'iasis*, (*cystis*, *ήπαρ*, 'the liver,' and *λιθιασις*, 'the formation of calculus,') The aggregate of phenomena caused by the presence of biliary calculi. See *Calculi*, *biliary*.

CYSTHITIS, *Kysthitis*.

CYSTHUS, (*κυσθος*), *Anus*, *Vulva*.

CYSTHYPERCARC'OSIS, (*cystis*, *ιμπερ*, 'over,' and *σάρκωσις*, 'a fleshy growth,') *Excrescent'ia Vesicæ urinariæ*. A fleshy thickening of the coats of the bladder.

CYST'IC, *Cysticus*, (F.) *Cystique*, *Kystique*, (from *cyst*.) Belonging to a cyst or cysts or to the gall-bladder. Also, containing cysts, *Cystous*.

CYSTIC ARTERY, *Arteria Cystica*, is given off from the right branch of the hepatic, and divides into two branches, which proceed to the gall-bladder. It is accompanied by two *cystic veins*, which open into the *vena porta abdominalis*.

CYSTIC BILE. Bile contained in the gall-bladder.

CYSTIC CALCULI. Calculi formed in the gall-bladder. See *Calculi*.

CYSTIC DUCT, *Ductus cysticus*, *Men'tus cysticus*. The duct proceeding from the gall-bladder, which, by its union with the hepatic, forms the *ductus communis choledochus*.

CYSTIC OXYD, *Oxyd*, *cystic*—c. *Oxyd Calculi*, see *Calculi*, *urinary*.

CYSTIC SARCO'MA, of AR'ERNETHY, *Emphy'ma Sarco'ma cellul'um*. Tumour, cellular or cystous; cells oval, currant-sized or grape-sized, containing serous fluid; sometimes caseous. Found in the thyroid gland (forming bronchocele), testis, ovary, &c.

CYSTICA, *Cystic Remedies*. Such medicines as were formerly believed proper for combating diseases of the bladder.

CYSTICER'CUS, (*cystis*, and *κερας*, 'a tail,') A genus of entozoa of the family of the hydatids, distinguished by the caudal vesicle in which the cylindrical or slightly depressed body of the animal terminates. The *Cysticer'cus cellul'us*, (F.) *Cysticerque du Tissu cellulaire*, *C. cellulaire*, *Hydatid'is finna*, has been often found in the areolar or cellular membrane. It is now considered to be the offspring of the tapeworm in one stage of its growth, having the power, however, of developing a large number of individuals resembling itself. See *Generation* (alternate), and *Metagenesis*.

CYSTICERCUS CELLULOSÆ, see *Cysticercus*.

CYSTICERQUE DU TISSU CELLULAIRE, *Cysticercus cellulose*.

CYSTIDALGIA, (*cystis*, and *αλγος*, 'pain,') *Cystalgia*.

CYSTIDELCO'SIS, (*cystis*, and *ελκος*, 'ulceration,') *Suppuration or ulceration of the urinary bladder*.

CYSTIDEPATICUS, *Hepatocystic*.

CYSTIDEPATOLITHIASIS, *Cysthepatolithiasis*.

CYSTIDOBLENNORRHOEA, (*cystis*, and *blennorrhœa*.) Cystirrhœa.

CYSTIDOCATARRHUS, (*cystis*, and *catarrhus*, 'catarrh.') Cystirrhœa.

CYSTIDOCELE, Cystocele.

CYSTIDOPLEGIA, (*cystis*, and *πληγη*, 'a stroke.') Cystoparalysis.

CYSTIDORRHAGIA, Cystorrhagia.

CYSTIDORRHEXIS, Cystorrhœxis.

CYSTIDORRHOEA, Cystirrhœa.

CYSTIDOSOMATOTOMIA, Cystosomatotomia.

CYSTIDOSPASMUS, see Cystospastic.

CYSTIDOSTENOCHORIA, (*cystis*, *στενος*, 'narrow,' and *χωρος*, 'place.') Stricture of the urinary bladder.

CYSTIDOTOMIA, Cystotomia.

CYSTIDOTRACHELOTOMIA, (*cystis*, *τραχηλος*, 'neck,' and *τομή*, 'incision.') Cystauchenotomia.

CYSTINE, Oxyd, cystic.

CYSTINURIA, Urine, cystinic.

CYSTINX, (*κυστιγξ*, dim. of *κυστις*, 'a bladder.') Vesicula.

CYSTIPHLOGIA, (*cystis*, and *φλογω*, 'to inflame.') Cystitis.

CYSTIQUE, Cystic.

CYSTIRRHAG'IA, (*cystis*, and *ῥήγνυμι*, 'I break forth.') Hemorrhage from the bladder, (F.) Hémorrhagie de la Vessie. By some used synonymously with cystirrhœa.

CYSTIRRHEUMA, Cystorrhœuma.

CYSTIRRHŒA, (*cystis*, and *ῥέω*, 'I flow.') *Paru'ria Stillati'tia Mucosa*, *Blennu'ria*, *Blennorrhœa urina'tis seu vesicæ*, *Cystidoblenorrhœa*, *Cystorrhœa*, *Cystidoblenorrhœa*, *Ischu'ria Cystophlegmatica*, *Tenesmus Vesicæ Mucosus*, *Uri'na mucosa*, *Cystocatar'rhus*, *Cystidocatar'rhus*, *Cystidorrhœa*, *Urocystocatar'rhus*, *Dysu'ria Mucosa*, *Pyu'ria Mucosa seu viscidula seu serosa*, *Morbus cystophlegmaticus*, *Catar'rhus vesicæ*, (F.) *Cystite muqueuse*, *Blennocystite*, *Flux muqueux de la vessie*, *Catarrhe vésical*, *Blencystorrhée* of Piorry. A copious discharge of mucus from the bladder, passing out with the urine, and generally attended with dysuria. It is commonly dependent upon an inflammatory or subinflammatory condition of the lining membrane. The treatment must be regulated by the cause. If it be not produced by an extraneous body, the antiphlogistic plan, the exhibition of warm diluents, and keeping the surface in a perspirable state, by wearing flannel, are indicated. Some of the turpentine may, at times, be given with advantage; and astringent or other appropriate injections be thrown into the bladder.

CYSTIS, gen. *Cyst'idis*, Follicle, Urinary bladder—c. *Choledochus*, Gall-bladder—c. *Fellea*, Gall-bladder—c. *Serosa*, Hygroma.

CYSTITE, Cystitis—c. *Muqueuse*, Cystitis.

CYSTITIS, (*cystis*, and *itis*.) *Inflammati'o Vesicæ*, *Empres'ma Cysti'tis*, *Cysti'tis u'rica*, *U'rocysti'tis*, *Cystiphlo'gia*, *Cystiphlo'gia*, *Inflammati'o of the blad'der*, (F.) *Cystite*, *Inflammation de la vessie*, *Cysturite*, of Piorry; characterized by pain and swelling in the hypogastric region; discharge of urine painful or obstructed, and tenesmus. It may affect one or all of the membranes; but commonly it is confined to the mucous coat. (F.) *Cystite muqueuse*. In the chronic condition, it appears in the form of cystirrhœa. It must be treated upon the same energetic principles as are required in other cases of internal inflammation; venesection, general and local, the warm bath, warm fomentations, warm, soothing enemata, diluents, &c. Cantharides must be avoided, even in the way of blisters, unless with precautions, as the disease is often occasioned by them.

CYSTITIS FELLEA, Cholecystitis—c. *Urica*, Cystitis.

CYSTITOME, *Cystit'omus*, *Cyst'otome*, *Kibis'titome*, *Kys'titome*, *Kystotome*, (*cystis*, and *τεμνειν*, 'to

cut.') An instrument invented by Lafaye for dividing the anterior part of the capsule of the crystalline, in the operation for extracting cataract. It was formed like the *Pharyngotome*.

CYSTOBLAST, (*κυστις*, 'a bladder,' and *βλαστος*, 'a germ.') Cytoblast.

CYSTOBLENNORRHOEA, Cystirrhœa.

CYSTOBUBONOCE'LE, (*cystis*, *βουβων*, 'the groin,' and *κηλη*, 'a tumour.') Hernia of the bladder through the abdominal ring.

CYSTOCATARRHUS, Cystirrhœa.

CYSTOCE'LE, *Cystocele'lia*, *Cistoce'lë*, *Cystidocel'ë*, (*cystis*, and *κηλη*, 'a tumour.') *Hernia vesicæ urinariæ*, *Zo'nia of the blad'der*, (F.) *Hernie de la Vessie*. It is not common. It occurs, most frequently, at the abdominal ring; less so at the crural arch, perinæum, vagina, and foramen thyroideum. It may exist alone, or be accompanied by a sac containing some abdominal viscous. The tumour is soft and fluctuating; disappears on pressure, and increases in size when the urine is retained. It must be reduced and kept in position by a truss. Vaginal cystocele is kept in place by a pessary.

CYSTOCELE BILIOSA, Turgescencia vesiculæ fellæ.

CYSTODYN'IA, (*cystis*, and *δύνη*, 'pain.') Pain of the bladder; particularly rheumatic pain.

CYSTOHÆMIA, (F.) *Cystohémie*, (*cystis*, and *αἷμα*, 'blood.') Congestion of blood in the bladder.

CYSTOID, *Cystoi'deus*, *Cyst-like*, (*cystis*, and *εἶδος*, 'resemblance.') That which resembles a cyst, as 'cystoid tumour.'

CYSTOLITHIASIS, (*cystis*, and *λίθιασις*, 'the formation of calculi.') See Calculi, vesical.

CYSTOLITHIC, *Cystolith'icus*, (F.) *Cystolithique*, (*κυστις*, 'the bladder,' and *λίθος*, 'a stone.') Relating to stone in the bladder.

CYSTOMA, see Kyst.

CYSTOMEROCE'LE, (*cystis*, *μερος*, 'the thigh,' and *κηλη*, 'hernia.') Femoral hernia, formed by the bladder protruding beneath the crural arch.

CYSTON'CUS, (*cystis*, and *ογκος*, 'tumour.') Swelling of the bladder.

CYSTOPARALYSIS, *Cystoparal'ysis*, *Cystidople'gia*, *Cystople'gia*, *Cystoplex'ia*, *Acystiner'via*, *Acysturonervia*, *Amyosthénie Cysturique* of Piorry, (*cystis*, and *παράλυσις*, 'palsy.') Paralysis of the urinary bladder. See Enuresis.

CYSTOPHLEGMATIC, *Cystophlegmat'icus*, (*cystis*, and *φlegμα*, 'mucus, phlegm.') Belonging to the vesical mucus. *Morbus cystophlegmaticus*. Cystirrhœa.

CYSTOPHLOGIA, (*cystis*, and *φλογω*, 'I inflame.') Cystitis.

CYSTOPHTHISIS, *Phthisis vesica'lis*, (*cystis*, and *φθιω*, 'I consume.') Consumption from ulceration of the bladder,—*Ul'cera seu Helco'sis vesicæ*.

CYSTOPLAS'TIC, *Cystoplas'ticus*, (*cystis*, and *πλασσω*, 'I form.') An epithet for an operation for the cure of fistulous openings into the bladder, *Cystoplasty*, (F.) *Cystoplastic*; sometimes restricted to the cure by translation of skin from a neighbouring part.

CYSTOPLEGIA, (*cystis*, and *πληγη*, 'a stroke.') Cystoparalysis.

CYSTOPLEG'IC, *Cystople'gicus*. Belonging to paralysis of the bladder.

CYSTOPLEXIA, Cystoparalysis.

CYSTOPTO'SIS, (*cystis*, and *πτειναι*, 'to fall.') Relaxation of the inner membrane of the bladder, which projects into the canal of the urethra.

CYSTOPY'IC, *Cystopy'icus*, (F.) *Cystopyique*, (*cystis*, and *πυον*, 'pus.') Relating to suppuration of the bladder.

CYSTORRHAG'IA, *Hæmatu'ria cyst'ica*, *Hæmorrhag'ia vesicæ*, *Strangu'ria cruen'ta*, *Sanguinis flu'or vesicæ*, (*cystis*, and *ῥαγναι*, 'rupture.')

A discharge of blood from the vessels of the urinary bladder.

CYSTORRHEUM'A, gen. *Cystorrhœum'atis*, *Cystirrhœu'ma*, *Rheumatism'us vesicæ urinariæ*; (*cystis*, and *ῥευμα*, 'defluxion,' 'rheumatism.') Rheumatism of the bladder.

CYSTORRHEX'IS, *Cystidorrhœx'is*; (*cystis*, and *ῥήξις*, 'rupture.') Rupture of the urinary bladder.

CYSTORRHŒA, *Cystirrhœa*.

CYSTOSARCO'MA, gen. *Cystosarco'matis*, (*cystis*, and *sarcoma*.) A tumour consisting of a combination of cysts and cystoids.

CYSTOSCOPE, *Cystoscop'ium*, (*cystis*, and *σκοπεω*, 'I examine.') A catheter, with a flattened extremity like the stethoscope, to enable the sound to be heard when the instrument strikes a stone in the bladder.

CYSTOSOMATOM'IA, *Cystosomatotom'ia*, *Cystidosomatotom'ia*, (*cystis*, *σωμα*, 'body,' and *τομή*, 'incision.') An incision into the body of the bladder.

CYSTOSPASMUS, see *Cystospastic*.

CYSTOSPAS'TIC, *Cystospas'ticus*, (*cystis*, and *σπασω*, 'I contract.') Relating to spasm of the bladder, and particularly of its sphincter;—*Cystidospas'mus*, *Spas'mus Vesicæ*, *Ischur'ia spasmod'ica*. (F.) *Spasme de la Vessie*.

CYSTOSPERM'TIS, (F.) *Cystospermite*, (*cystis*, *σπέρμα*, 'sperm,' and *itis*.) Inflammation of the vesiculæ seminales.

CYSTOSTENOCHO'RIA, *Cystidostenochor'ia*, *Strictur'a vesicæ*, *Vesic'a sacca'ta*, (*cystis*, *στενός*, 'narrow,' and *χωρὸς*, 'place.') A stricture, narrowness, inequality, or saccated condition of the urinary bladder.

CYSTOTHROM'BOID, *Cystothromboi'des*, (*cystis*, and *θρομβός*, 'a clot.') Relating to the presence of clots in the bladder.

CYSTOTOME, *Cystot'omus*, (*cystos*, and *τεμνειν*, 'to cut.') An instrument intended for cutting the bladder. Instruments of this kind have been more frequently, although very improperly, called *Lithotomes*.

CYSTOTOMY, *Kystot'omy*, *Cystotom'ia*, *Cystidotom'ia*. Incision of the bladder; *Sec'tio vesicæ'tis*. *Cystotomy* means cutting into the bladder for any purpose; (F.) *Incision de la vessie*. Commonly, it is applied to the puncturing of the bladder for the purpose of removing the urine; whilst *Lithotomy* has been employed for the incisions made with the view of extracting calculi from the bladder. See *Lithotomy*.

CYSTOTRACHELOTOMIA, (*cystis*, *τραχηλος*, 'neck,' and *τομή*, 'incision,') *Cystauchenotomia*. See *Lithotomy*.

CYSTOUS, *Cystic*.

CYSTOVARIUM, (*cyst*, and *οαριον*, 'ovarium.') A morbid condition, which consists in the formation of cysts in the ovary.

CYSTURITE, (*cystis*, *ουρον*, 'urine,' and *itis*), *Cystitis*.

CYSUS, (*κυσος*), *Anus*.

CYTHEREAN SHIELD, *Condom*.

CYTHEROMANIA, (*Κυθέρεια*, 'Venus,' and *μανία*), *Nymphomania*.

CYTINUS, *C. Hypocist'is*, *Hypocist'is*, *As'a-rum Hypocist'is*. *Ord.* *Cistineæ*. A small parasitical plant, which grows in the south of France

and in Greece, on the roots of the woody *cistus*. The juice of its fruit is acid, and very astringent. It is obtained by expression, and converted into an extract, which was called *Sucus Hypocist'idis*, and was formerly much used in hemorrhages, diarrhœa, &c.

CYTIS'INA, *Cyt'isine*. An immediate vegetable principle, discovered by Chevalier and Lassaigne in the seeds of *Cyt'isus Laburnum* or *Bean-Tref'oil Tree*. *Sec. Syst.* *Diadelphia Decandria*. *Cytisine* has analogous properties to emetine. In the dose of one or two grains it produces vomiting and purging; and, in a stronger dose, acts as an acrid poison. The seeds of the *Cyt'isus Laburnum*, (F.) *Ambours*, have been long known to produce vomiting and purging.

CYTISMA ECZEMA, *Eczema*—*c.* *Herpes*, *Herpes*.

CYTISOGENISTA, *Spartium scoparium*.

CYTISUS LABURNUM, (*κυθισος*), see *Cytisina*—*c.* *Scoparius*, *Spartium scoparium*.

CYTIT'IS, *Scyti'tis*, *Dermat'itis*, *Dermi'tis*, *Cutit'is*, *Chorionit'is*, *Cori'itis*, (*κυττις*, *cutis*, 'the skin,' and *itis*.) Inflammation of the skin.

CYTOBLAST, *Cyst'oblast*, (*κυτος*, 'cell,' and *βλαστος*, 'germ,') *Cellgerm*, *Nu'cleus*, (F.) *Noyau*. A granule from which all animal and vegetable bodies are presumed to be formed. When the nucleus or cytoblast forms a cell, and is attached to its walls, the *germinal cell*, thus formed, is called a *nucleated cell*, (F.) *Cellule à noyau*. When the nucleus contains a simple granule, the latter is termed a *nucleolus*, (F.) *Nucléole*, *Nucléolule*. In the cell nomenclature of Professor Agassiz, the cell-wall is termed *ectoblast*: the nucleus, *mesoblast*; the nucleolus, *entoblast*; and when a smaller body exists in this, *entostoblast*.

CYTOBLASTE'MA, gen. *Cytoblaste'matis*, (F.) *Cytoblastème*, same etymon. *Intercell'ular substance*, *Hyaline substance*, *Substantia vit'rea seu hyal'ina*, *Mat'rix*. The gum or mucus in the vegetable, and probably the liquor sanguinis after transudation from the vessels in the animal, in a state fully prepared for the formation of the tissues.—Schwann and Schleiden. By many *Blutstema* is preferred, inasmuch as it does not convey the idea of cell development.

CYTOGENET'IC, *Cytogenet'icus*, (F.) *Cytogénétique*. Relating or belonging to cytogeny.

CYTOG'ENY, *Cytogen'esis*, (*κυτος*, 'cell,' and *γενεσις*, 'generation,') Cell formation, cell genesis. The continuous development of cells, in tissue formation, has been called by Virchow *Proliferation of cells*, (G.) *Zellenwucherung*.

CYTOID, *Cytoi'des*, (*κυτος*, 'a cell,' and *εἶδος*, 'resemblance,') Cell-like. A term applied by Henle to corpuscles—as those of lymph, chyle, pus, &c.—which seem to resemble each other essentially in their chemical and microscopical characters, and which have been termed, in the aggregate, by some, *Leucocytes*.

CYTOS'TASIS, (*κυτος*, 'a cell,' and *στασις*, 'station,') Arrest of red or white blood cells or corpuscles in the small vessels, owing to greater force of aggregation than in health. [?]

CYTAROS, (*κυτταρος*), *Glands*.

CYZEMER. A swelling of the wrists.

CYZICE'NUS, (*κυζικηνος*.) An ancient plaster for chronic ulcers and wounds of tendons.

D.

The figure of the Greek Δ, according to Galen, was the sign for quartan fever.

DABACH, Viscum album.

DACNE'RON, (δακνρον, from δακνειν, 'to bite.')

An ancient name for a collyrium, composed of oxide of copper, pepper, cadmia, myrrh, saffron, gum Arabic, and opium.

DACRY, *Dacryma*, *Dacryon*, (δακρυ, 'a tear.')

DACRYADENAL'GIA, *Dacryoadenalgia*, (dacrý, δαήν, 'a gland,' and αλγος, 'pain.')

Disease or pain in the lachrymal gland.

DACRYADENI'TIS, *Dacryoadenitis*, (dacrý, δαήν, 'a gland,' and itis.) Inflammation of the lachrymal gland.

DACRYALLOO'SIS, (dacrý, and αλλοιωσις, 'change.')

A morbid condition of the tears.

DACRYDION, (from δακρυ, 'a tear,' from the gummy exudation.)

Convulsus scammonia.

DACRYGLO'SIS, (dacrý, and γλω, 'I laugh.')

A kind of insanity, in which the patient weeps and laughs at the same time.

DACRYHEMORRHYSIS; *Sanguineous lachrymation*, (dacrý, and αιμορρυσις, 'hemorrhage.')

A flow of bloody tears.

DACRYMA, (δακρυμα,) see Daery.

DACRYOADENALGIA, *Dacryadenalgia*.

DACRYOADENITIS, *Dacryadenitis*.

DACRYOBLENNORRHEA, (dacrýon, βλεηνα, 'mucus,' and βρω, 'I flow.')

Discharge of tears mixed with mucus.

DACRYOCYSTAL'GIA, (dacrýon, κυστις, 'a sac,' and αλγος, 'pain.')

Disease or pain in the lachrymal sac.

DACROCYSTIS, (dacrýon, and κυστις, 'a sac,')

Lachrymal sac.

DACROCYSTI'TIS, (dacrocyctis, and itis.) Inflammation of the lachrymal sac.

DACRYOCYSTOSYRINGOKATAKLEY'SIS; (dacrýocystis, σπριγξ, 'a pipe, a fistula,' and κατακλεισις, 'a locking up.')

A term proposed by Dieffenbach for the healing of lachrymal fistulae by transplantation. The operation consists in paring the edges of the fistula, loosening the borders, and assisting the requisite tegumental displacement by lateral incisions.

DACRYODES, (δακρυωδης, from dacrýon, and οδης,) *Lacrymosus*.

Weeping; resembling tears:—hence, *Ulcus dacrýodes*. A sanious ulcer, a weeping sore.

DACRYOLITE, *Dacryolithus*, *Dacryolith*, (F.) *Dacryolithes*, (dacrýon, and λιθος, 'a stone.')

A concretion found in the lachrymal passages. When in the nasal duct, it is termed *Rhinolite*, *Rhinolithus*, *Rhinodacryolite*, *Rhinodacryolithus*.

DACRYOLITHI'ASIS, (dacrýon, and λιθιασις, 'formation of calculi.')

The formation of concretions in the tears

DACRYOLITHUS, *Dacryolite*.

DACRYOMA, gen. *Dacryomatia*. Epiphora. (from dacrýon, and oma.)

The effusion of tears, occasioned by an occlusion of the puncta lachrymalia.

DACRYON, (δακρυον,) Tear.

DACRYOPHE'US, (dacrýon and ποιω, 'I make.')

A substance which excites the secretion of tears,—as the onion, horse-radish, garlic, &c.

DACRYOPS, gen. *Dacryopsis*, from dacrýon, and ψ, 'the eye.')

A weeping eye. A tumefaction of the lachrymal ducts, or lachrymal cyst.

DACRYOPYORRHEA, *Pyorrhoea lacrymalis*, (dacrýon, πυορ, 'pus,' and βρω, 'to flow.')

A discharge of tears mixed with purulent matter.

DACRYORRHEA, *Dacryrrhoea*, *Dacryor-*

rhoea, *Dacryrrhoea*, (dacrýon, and βρω, 'to flow.')

A morbid flux of tears.

DACRYORRHYSIS, (dacrýon, and ρησις, 'a flow.')

Dacrýorrhoea.

DACRYOOLEN, (dacrýon, and ωλην, 'a canal,')

Lachrymal canal.

DACRYOLENITIS, (dacrýosolen, and itis.) Inflammation of the lachrymal ducts.

DACRYOSYRINX, (dacrýon, and σπριγξ, 'a fistula,')

Fistula lacrymalis.

DACRYRRHEA, *Dacryorrhoea*.

DACRYRRHYSIS, *Dacryorrhoea*.

DACTYLE'THRA, *Dactylithra*, (δακτυληθρα, 'a covering for a finger.')

A name given by the ancients to different topical applications having the form of a finger, and proper for being introduced into the throat to excite vomiting.

DACTYLETUS, *Hermodactylus*.

DACTYLION, *Dactylium*, (F.) *Doigts palmés*, *Webbed fingers*, (δακτυλος, 'a finger.')

The union of the fingers with each other. This affection is generally congenital; but it may be owing to burns, ulcerations, inflammation of the fingers, &c.

DACTYLITIS, (dactylo, and itis,) Paronychia.

DACTYLUS, (δακτυλιος,) *Annilus*. A ring. A troch or lozenge, when shaped like a finger. The anus.

DACTYLUS ACULEA'TUS. A worm of a light colour, annulated, cylindrical, but tapering slightly towards both extremities, from two-fifths to four-fifths of an inch long, which has been found in the urine.

DACTYLO, (δακτυλος, 'a finger,')

in composition, finger.

DACTYLODOCHME, (δακτυλοδοχη,) *Dochme*.

DACTYLOLOGY, *Dactylogogia*, (dactylo, and λογος, 'a discourse.')

Conversing with the fingers, —as by the deaf and dumb.

DACTYLOSYPHYSIS, (dactylo, and συμψυσις, 'union.')

Adhesion of the fingers to each other.

DACTYLOTHE'KĒ, (dactylo, and θεκη, 'a case or sheath.')

An instrument for keeping the fingers extended when wounded.

DACTYLUS, (δακτυλος,) *Dig'itus*. A finger.

The smallest measure of the Greeks, the sixth part of a foot. Also, the Date.

DÆDALEA SUAVEOLENS, *Boletus seu Fungus Salicis*, *Boletus discoides seu suaveolens*, *Fungus albus saliginosus*, (F.) *Agaric odorant*, *Bolet odorant*. Ord. Fungi.

A champignon, which grows on the trunks of old willows. It has a smell of anise, which is penetrating and agreeable; and has been recommended in phthisis pulmonalis in the dose of a scruple four times a day.

DÆDALUS, Hydragryum.

DÆDION, (dim. of δαίς, 'a torch,')

Bougie.

DÆMONIA, Demonomania.

DÆMONIACUS, (δαιμων, 'a demon,')

Possessed.

DÆMONOMANIA, Demonomania.

DÆMONOPATHIA, Demonomania.

DÆS, (δαις,) Tæda.

DAFFING, Insanity.

DAFFODIL, *Narcissus pseudonarcissus*.

DAFFY'S ELIXIR, (after the proposer,) *Tinctura sennæ composita*.

DAFT, (Sc. and Prov.,) Insane.

DAG, Stitch.

DAISY, (Sax. *dæcerege*,) *Chrysanthemum leucanthemum*, *Erigeron Philadelphicum*—d. Common, Bellis—d. Ox-eye, *Chrysanthemum leucanthemum*.

DALBY'S CARMINATIVE, (after the proposer.)

A celebrated empirical remedy, much used as a carminative for children. The follow

ing is a form for its preparation: (*Magnesia alb.* ʒij; *Ol. mentha piper.* gtt. j; *Ol. nuc. moschat.* gtt. iij; *Ol. anisi.* gtt. iij; *Tinct. castor.* gtt. xxx; *Tinct. asafetid.* gtt. xv; *Tinct. opii.* gtt. v; *Sp. pulegii.* gtt. xv; *Tinct. cardom.* c. gtt. xxx; *Aque mentha pip.* ʒij. M.) A Committee of the Philadelphia College of Pharmacy recommended the following form:—*Aque Ox.*; *Sacchar. alb.* ʒxxxij; *Carbon. Potass.* ʒss; *Carb. Mag.* ʒxij; *Tinct. Opii.* fʒvj; *Ol. menth. pip.*, *Ol. Anethi cœniol.* aa ʒij. M.

DALCOP, Idiot.

DALTONIAN. An absurd name given to one who cannot distinguish colours; because the celebrated physicist Dalton had the defect. See *Achromatopsia*.

DALTONISM, *Achromatopsia*.

DAMP, AFTER, ([G.] *Dampf*, 'vapor,') see Hydrogen, carburetted—d. Black, see Hydrogen, carburetted—d. Choke, Carbonic Acid, see Hydrogen, carburetted—d. Fire, Hydrogen, carburetted.

DAMSON, (from Damascus,) *Prunum Damasacenum*, see *Prunum*—d. Mountain, *Quassia simarouba*—d. Tree, *Prunus domestica*.

DANCE, see Mania, dancing—d. St. John's, see Mania, dancing—d. St. Vitus's, Chorea, see Mania, dancing.

DANC'ING, *Saltatio*, (F.) *Dance*, (G.) *Tanz*. A kind of exercise and amusement, composed of a succession of motions, gestures, and attitudes, executed by measured steps to the sound of the voice or musical instruments. It is a healthy exercise.

DANCING MANIA, see Mania, dancing—d. Plague, see Mania, dancing.

DANDELION, ([F.] *dent de lion*, 'lion's tooth,') *Leontodon taraxacum*.

DANDER, (from *Dandriff*,) *Pityriasis*.

DANDRUFF, (Sax. *tan*, 'a spreading eruption,' and *drof*, 'filthy,') *Pityriasis*.

DANDRUFF, *Pityriasis*.

DANDY, *Dengue*.

DANEVERT, MIN'ERAL WA'TERS OF. A spring, a league and a half from Upsal, in Sweden. The waters contain carbonic acid, holding in solution carbonate of iron, sulphate of iron, sulphates of soda and lime, chloride of sodium, and silica. It is frequently employed in medicine.

DANEWORT, *Sambucus ebulus*.

DANICH; an Arabic word, signifying the weight of 8 grains.

DANIEL-LIA THURFF'ERA. The *Frankincense Tree* of Western Africa, which furnishes the product termed *African Frankincense*.

DANSE, Dancing—d. *de St. Guy*, Chorea—d. *de St. Witt*, Chorea.

DAOUN SETAN, see *Urtica*.

DAPHNE, (*δαφν*, after the nymph *Daphne*, *Laurus*).

DAPHNE ALPI'NA, *Chamæle'a. Chamæle'a, Wildonwail. Ord. Thymelææ. Ser. Syst. Octandria Monogynia*. A sort of dwarf-olive. An acrid, volatile, alkaline principle was separated from the bark of this plant by M. Vauquelin, to which he gave the name *Daph'nine*. The plants of the genus owe their vesicating property to this principle.

DAPHNÉ BOISGENTIL, *Daphne mezereum*—d. Flax-leaved, *Daphne gnidium*—d. *Garou*, *Daphne gnidium*.

DAPHNE GNID'IUM, *D. paniculata, Thymele'a, Th. Monspeli'aca, Thymele'a, Cneo'nor, Spurge Flax, Flax-leaved Daph'né*. The plant which furnishes the *Garou Bark*, (F.) *Daphné Garou, Sain-bois*. It is chiefly used, when used at all, for exciting irritation of the skin. The *Gra'na Gnid'ia*, (see *Cnid'ia gra'na*), are acrid poisons, like all the plants of this genus, when taken in quantity. According to others, the *garou bark*

and *grana gnidia* are obtained from the *daphne laureola*.

DAPHNE LAURE'OLA, *D. major, Thymele'a laureola*. The systematic name of the *Spurge Laurel, Laurel-ola*. The bark of this plant has similar properties to the last.

DAPHNE LIOTTARDI, *D. Mezereum*—d. Major, *D. laureola*.

DAPHNE MEZE'REUM or MEZERE'UM, *D. Liottardi, Thymele'a meze'reum*. The systematic name of the *Meze'reon, Meze'reum, Spurge olive*, (F.) *Daphné Boisgentil, Solibois*. The bark of the *meze'reon, Meze'reum*, possesses analogous properties to the other varieties of *Daphne*. In the Pharmacopœia of the United States, *Mezereum* is the bark of *Daphne mezereum*, and of *Daphne gnidium*. It is considered stimulant and diaphoretic; and, in large doses, is emetic. It has been employed in syphilitic cases, but its efficacy is doubtful. Soaked in vinegar,—*meze'reum acetatum*,—like the other varieties of *daphne*, it has been employed to irritate the skin, especially to keep issues open.

DAPHNE PANICULATA, *D. Gnidium*.

DAPHNELÆ'ON, *O'leum Lauri'nium*, (*δαφνελαιον*, from *δαφν*, 'the laurel or bay tree,' and *ελαιον*, 'oil,') *Oil of Bay*.

DAPHNINE, see *Daphne Alpina*.

DARNEL, *Lolium temulentum*.

DARSENÍ, *Laurus cinnamomum*.

DARSINI, *Laurus cinnamomum*.

DAR'SIS, (*δαρσις*, from *δέρω*, 'I excoriate,' 'I skin,') The Greek physicians seem to have used this word to designate the anatomical preparation, which consists in removing the skin for exposing the organs covered by it.

DARTA, *Impetigo*—d. *Excoriativa*, *Herpes exedens*—d. *Maligna, Herpes exedens*.

DAR'TOS, (*δαρτος*, 'skinned,') *Membrana carnosa, Tunica muscularis seu rubicunda scro'ti, T. dar'tos, Marsupium musculo'sum*. A name given to the second covering of the testicle, which the ancient anatomists conceived to be muscular, but which is merely areolar. Its external surface is towards the scrotum; the internal towards the tunica vaginalis. Frederick Lobstein and Breschet consider, that it proceeds from an expansion of the fibrous cord, known by the name *Gubernaculum testis*.

DARTRE, (*Darta*, [?]) *Herpes, Impetigo, Pityriasis*—d. *Croûteuse, Impetigo*—d. *Crustacée, Ecthyma impetigo*—d. *Crustacée flavescens, Porrigo lupinosa*—d. *Écailleuse, Psoriasis*—d. *Fongueuse, Ecthyma*—d. *Furfuracée arrondie, Lepra, Lepra vulgaris*—d. *Furfuracée volante, Lichen, Pityriasis*—d. *Phlycténôide, Herpes, phlycténoides*—d. *Pustuleuse conopseuse, Gutta rosea*—d. *Pustuleuse, disséminée, Acne*—d. *Pustuleuse mentagra, Syccosis*—d. *Rongeante, Herpes exedens*; see *Esthiomenus*—d. *Squammeuse humide, Eczema*—d. *Squammeuse lichénôide, Psoriasis*—d. *Tonsurante, Porrigo decalvans*—d. *Vive, Eczema*.

DARTREUX (F.), *Herpet'ic*. Participating in the characters of *Dartre* or *Herpes*. Also, one affected with *dartre*. *Dartre* has been used, at one time or other, for almost every disease of the skin. See *Herpes*.

DASY'MA, (*δαρς*, 'rough,' 'hairy,') A disease of the eye—the same as *trachoma*, but less in degree.

DASY'TES, (*δαρτης*,) Roughness, particularly of the tongue and voice. Hairiness, *Hirsuties*.

DATE, *Pal'mula, Dactylus, Bal'anos, Phœ'nicos*. The fruit of the *Phœ'nix dactylif'era* seu *ex-cel'sa, Pal'ma dactylif'era*, (F.) *Datte*. The unripe date is astringent. When ripe, it resembles the fig. The juice of the tree is refrigerant.

DATE PLUM, INDIAN, *Diospyrus lotus*.

DATTE, *Date*.

DATURA, (Arab. *tatorah*,) *D. Stramonium*.

DATU'RA SANGUIN'E. *Red Thornapple*: called by the Indians of Peru *Huacachu*, *Yerba de Huaca* (*huaca*, 'a grave') or *Grave-plant*, and *Bocueche*. *Ord.* Solanaceæ. A plant from which the Peruvian Indians prepare a narcotic drink called *Tungu*.

DATURA STRAMO'NIUM, *Stramo'nia*, *Baryoccalon*, *Sola'nium foetidum*, *Stramo'num*, *S. majus album* seu *spinosum* seu *vulgarium* seu *foetidum*, *Po'mum* seu *Ma'lum spinosum*, *Nux methel.* *Datu'ra*, *Dutro'a*, *Daty'ra*, *Stram'ony*, *Thornapple*, *Apple of Peru*, *James' town Weed*, *Jim's town* or *Simpson Weed*, *Stink'weed*, (F.) *Stramonie*, *Pomme épineuse*, *Endormie*, *Herbe aux sorciers* ou *de diable*. The leaves, *Stramo'nii Fo'lia*, *S. Folium* (Ph. U. S., 1863), the seeds, *Stramo'nii Se'men*, and the root, *Stramo'nii Ra'dix*, are the parts used in medicine. They are narcotic and poisonous; are given internally as narcotics and antispasmodics; and applied externally as sedatives, in the form of fomentation. The seeds are smoked like tobacco, in asthma. The dose of the powder is, gr. j to gr. viii.

Datu'ra tat'ula, has similar properties.

DAT'URINE, *Datur'ina*, *Datu'ria*, *Datur'ium*. The active principle of *Datura Stramonium*, first separated by Brandes, a German chemist. It has not been rendered available in medicine.

DATURIUM, *Daturine*.

DATYRA, *Datura stramonium*.

DAUCI RADIX, see *Daucus Carota*.

DAUCITES VINUM. Wine, of which the *Daucus* or *Carrot* was an ingredient. The seeds were steeped in must. It was formerly used in coughs, convulsions, hypochondriasis, diseases of the uterus, &c.

DAUCUS CANDIANUS, *Athamanta cretensis*.

DAUCUS CARO'TA, (*daucus*). The systematic name of the *Carrot Plant*; *Daucus*, *D. sylvestris* seu *vulgaris* seu *sativus*, *Cavea'lis caro'ta*, *Pastina'ca sylvestris tenuifolia officinarum*, *Ado'rium*; *Order*, Umbelliferae. (Sc.) *Meeran*, *Mirrot*, (F.) *Carotte*; *Ord.* Umbelliferae. The fruit, *Caro'ta*, *Carrot seed*,—(Ph. U. S.)—and the root, *Dau'ci ra'dix*, have been used in medicine. The root is sweet and mucilaginous; and the seeds have an aromatic odour, and moderately warm, pungent taste. The root has been used, as an emollient, to fetid and ill-conditioned sores. The seeds have been regarded as stomachic, carminative, and diuretic; but they have little efficacy. The seeds of the wild plant are, by some, preferred to those of the garden.

DAUCUS CRETICUS, *Athamanta cretensis*—*d.* *Cyanopus*, *Pimpinella magna*—*d.* *Sativus*, *D. carota*—*d.* *Seprinius*, *Scandix cerefolium*—*d.* *Sylvestris*, *D. carota*—*d.* *Vulgaris*, *D. carota*.

DAUPHINELLE, *Delphinium staphisagria*.

DAVIDSON'S REMEDY FOR CANCER, see *Conium maculatum*.

DAVIER, (after the inventor, [?]) *Dentagra*.

DAWLISH, CLIMATE OF. A town in Devonshire, frequented by phthisical invalids during the winter. It is well protected from northerly winds, and also from the violence of the south-westerly gales. It offers, however, but a confined space.

DAWPIT (Sc.), see *Fatuitas*.

DAX, MINERAL WATERS OF. Dax is a city two leagues from Bordeaux, where there are four springs, that are almost purely thermal; containing only a very small quantity of chloride of magnesium, and sulphate of soda and lime. Temperature 70° to 133° Fahrenheit.

DAYBLINDNESS, *Nyctalopia*.

DAYMARE, *Incubus vigilantium*.

DAYNETTLE, *Lamium album*.

DAYS' EYE, *Daisy*.

DAYLIGHT, *Hemeralopia*—*d.* *Vision*, *Hemeralopia*.

DAZZ'LING, (dim of *dazing*), *Obtignis*, (F.) *Éblouissement*. A momentary disturbance of sight, occasioned either by the sudden impression of two powerful a light or by some internal cause,—as plethora.

DE, (F.) *dés*, 'from,' as a prefix, signifies, at times, a downward movement; and at others a privative power, whilst at others again, it has an intensive signification.

DE VENTRE INSPICIEN'DO, 'of inspecting the belly.' Where there is reason to suppose that a woman feigns herself pregnant, a writ *de ventre inspiciendo* may be issued to determine whether she is so or not. Until recently, in England, the decision was left to twelve matrons and twelve respectable men, according to the strict terms of the ancient writ.

DEAD, ((Sax. *dead*), *Mor'tuus*, (F.) *Mort*. Deprived of life, exanimate.

DEADHOUSE, *Morgue*.

DEADLY, *Lethiferous*, *Mortal*.

DEAD MEN'S BELLS, *Digitalis*.

DEADNETTLE, *Lamium album*.

DEADTHRAW, *Agony*.

DEAF, (Sax. *deaf*), *Sur'dus*, (F.) *Sourd*, (Sc.) *Dull*. Devoid of hearing. Hard of hearing.

DEAF-DUMBNES, *Mutitas surdorum*.

DEAF-MUTISM, *Mutitas surdorum*.

DEAFNESS, *Sur'ditas*, *Cophosis*, *Dyscœcia*, *Dyscœcia organica*, *Barycœcia*, *Hardness of hearing*, *Hypocophosis*, *Subsur'ditas*, *Bradycœcia*, *Bradycœcia*, (F.) *Surdité*, *Dureté de l'ouïe*, *Dyscœcie*. Considerable diminution or total loss of hearing. It may be the effect of acute or chronic inflammation of the internal ear, closure of the Eustachian tube—throat deafness; paralysis of the auditory nerve or its pulpy extremity, or of some mechanical obstruction to the sonorous rays. In most cases, however, the cause of the deafness is not appreciable, and the treatment has to be purely empirical. Syringing the ears, dropping in slightly stimulating oils, fumigations, &c., are the most likely means to afford relief.

DEAFNESS, TAYLOR'S REMEDY FOR, see *Allium*—*d.* *Throat*, see *Deafness*.

DEALBATIO, (*de*, and *albus*, 'white,') *Dealbation*, *Paleness*.

DEALBATION, *Dealbatio*, *Bleach'ing*. Applied, particularly, to the whitening of bones for anatomical purposes.

DEAMBULATIO, (*de*, and *ambulare*, 'to walk,') *Walking*.

DEARCH (Sc.), (Sax. *ðreorðh*.) *Napus*.

DEARTICULATIO, (*de*, and *articulus*, 'a joint,') *Diarthrosis*.

DEASCIATIO, (*de*, and *ascia*, 'an axe,') *Apo-seeparnismus*.

DEATH, (Sax. *deað*), (Sc.) *Dede*, *Deid*; *Apobiosis*, *Abiosis*, *Eztinctio*, *Obitus*, *Psychorag'ia*, *Psychorrag'ia*, *Le'thum*, *Le'tum*, *Mors*, *Inter'itus*, *Than'atus*, *Decease*, (F.) *Mort*. Definitive cessation of all the functions, the aggregate of which constitute life. *Real Death*, *Apothanasia*, is distinguished from *asphyxia* or *apparent death*—the latter being merely a suspension of those same functions. But it is often difficult to judge of such suspension, and the only certain sign of *real death* is the commencement of putrefaction. At times, therefore, great caution is requisite to avoid mistakes. Death is commonly preceded by some distressing symptoms, which depend on lesion of respiration, circulation, or of the cerebral functions, and which constitute the *agony*. That which occurs suddenly, and without any, or with few, precursory signs, is called *sudden death*. It is ordinarily caused by disease of the heart, apoplexy, the rupture of an aneurism, or by some other organic affection. Death is *natural*, (Sc.) *Stræ-death*, when it occurs as the result of dis-

case violent, when produced by some forcible agency. It may likewise affect the whole body, or a part only; hence the difference between somatic and molecular death.

The chief varieties of the modes of death usually admitted may be thus given

Death beginning at the heart,	{ sudden — Syncope: gradual — Asthenia.
“ “ in the lungs — Apnoea.	
“ “ in the brain — Apoplexy.	
“ “ in the gray matter of the medulla,	{ Paralysis of pneumogastrics, &c.
“ “ in the blood [?] —	necræmia.

As it was at one time conceived, with Bichat, that death must commence in the heart, lungs, or brain, these organs were called *a'tria mor'tis*, 'the halls of death.'

DEATH, APPARENT, Asphyxia.

DEATH, BLACK, *Melanothan'atus*, *Pes'tis ni'gra*, *Black Plague*, (F.) *Peste noire*. The plague of the 14th century was so called, which is supposed to have proved fatal in Europe to 25,000,000 of people.

DEATH OF MAN, *Cicuta maculata*—d. Rate, see Mortality—d. Stiffening, Rigor mortis—d. Trance, see Asphyxia.

DEAURENTUR PILULÆ, 'let the pills be gilded.' Formerly it was the practice to coat pills with silver or gold leaf, and this is still done occasionally.

DÉBANDER (F.), (*de* and *bande*, 'a bandage.') To remove a bandage or dressing,—as *débunder une plaie*.

DEBILIS, ('weak,') Infirm. See the next word.

DEBILITANTS, *Debilitan'tia*, *Antidynam'ica*, *Philadynam'ica*, (*debilitare*, itself from *debilis*, quasi *debilis*, 'weak,' 'to weaken.') Remedies exhibited for the purpose of reducing excitement. Antiphlogistics are, hence, debilitants.

DEBILITAS, gen. *Debilitat'is*, Debility — d. Erethisis, see Irritable — d. Nervosa, Neurasthenia, see Irritable — d. Visûs, Asthenopia.

DEBILITY, *Debilit'as*, *Asthen'ia*, *Hyposthen'ia*, *Blaci'a*, *Anenerge'sia*, *Anenergi'a*, *Weakness*, (F.) *Faiblesse*. A condition, which may be induced by a number of causes. It must not be confounded with *fatigue*, which is temporary, whilst debility is generally more permanent.

Debility may be *re'al*, or it may be *appa'rent*; and, in the management of disease, it is important to attend to this. At the commencement of fever, for example, there is often a degree of apparent debility, which prevents the use of appropriate means, and thus is the cause of much evil. Excitement is more dangerous than debility.

DÉBOITEMENT, (*de*, and *boiter*, 'to be lame,') Luxation.

DEBORDEMENT (F.), *Déborde*, (*déborder*, 'de, and border,') 'to overflow.' A popular term for one or more sudden and copious evacuations from the bowels. It is chiefly applied to bilious evacuations of this kind.—*Débordement de Bile*.

DEBRIDEMENT (F.) *Fræno'rum solu'tio*, (*débrider*, (*de*, and *brider*,) 'to unbridle,') The removal of filaments, &c., in a wound or abscess, which prevent the discharge of pus. In a more general acceptation, it means the cutting of a soft membranous, or aponeurotic part, which interferes with the exercise of any organ whatever; thus, in paraphimosis, *débridement* of the prepuce is practised to put an end to the inflammation of the glans; in strangulated hernia, *débridement* of the abdominal ring is had recourse to, to remove the stricture of the intestine, &c.

DECAGRAMME, (*δεκα*, 'ten,' and *γραμμα*, 'a gramme,') The weight of ten grammes, 154.34 grains Troy.

DECAM'YRON, (*δεκα*, 'ten,' and *μυρον*, 'ointment,') An ancient cataplasm, composed of *malabathrum*, *mastic*, *euphorbium*, *spikenard*, *styrac calamita*, carbonate of lime, common pepper, *unguentum nardi*, *opobalsamum*, and *resac*.

DECANTA'TION, *Decanta'tio*, (from *decantare*, (*de*, and *cantare*,) 'to sing or repeat over and over,' or from *de*, and (D.) *kant*, 'a corner;' hence 'to cant,' 'to set on edge in order to decant or pour off,' [?]) *Defu'sio*, *Metangiam'us*, *Catuch'ysis*. A pharmaceutical operation, which consists in pouring off, gently, by inclining the vessel, any fluid which has a deposit.

DECAPITATIO, Decollation—d. *Articulatorum*, see Resection.

DECAPITATION, (*de*, and *caput*, gen. *capitis*, 'head,') Decollation.

DECARBONIZATION, *Hæmotosis*.

DECEASE, (*decedere*, *decessum*, (*de*, and *cedo*,) 'to depart,') Death.

DECESSIO, *Ecpiesma*.

DÉCHAPELLEMENT, (*de*, and *chapeau*, from *caput*, 'head,') An old French term, employed by Paré and others for the operation of cutting and removing, with a strong forceps, the corona of a carious tooth, where it is desired to preserve the fang.

DÉCHARNÉ, (*de*, and *chair*, 'flesh,') *Demusculatus*.

DÉCHAUSSEMENT (F.), (*de*, and *chausser*, 'to put on shoes and stockings,') The state in which the gums have fallen away from the teeth, as in those affected by mercury, in old persons, &c. Also, the operation of lancing the gums. See Gum lancet.

DÉCHAUSSOIR, Gum lancet.

DÉCHIRÉ, *Lacerum*.

DÉCHIREMENT, (from *dilacerare*, (*di*, and *lacerare*, 'to tear,') 'to tear to pieces,') *Laceration*.

DÉCHIRURE, Wound, lacerated—d. *de l'Utérus*, Uterus, rupture of the.

DECIDENTIA, (*decidere*, 'to fall off,' *de*, and *cadere*, 'to fall,') Catapsois, Epilepsy.

DECIDUA, *Decidua membrana*.

DECID'UA MEMBRA'NA, *Decid'ua*, *Decid'uons Mem'brane*. So called on account of its being considered to fall off from the uterus at parturition. A membrane, formerly defined to be the outermost membrane of the fœtus in utero; and still so defined by some—as by Dr. Lee. Chaussier calls it *Epicho'rium*; by others, it has been called *Perio'na*, *Membra'na cadu'ca Hunteri*, *M. flocculen'ta seu cellul'osa seu sinu'osa seu commun'is seu præexis'tens*, *Decid'ua exter'na seu spongi'osa*, *Tu'nica exte'rior o'vi seu cadu'ca seu erus'sa*, *Membra'na cribro'sa seu o'vi mater'na seu muco'sa seu u'teri inter'na evolu'ta*, *Epio'nê*, *Placen'ta u'teri succenturia'ta*, *Subplacen'ta*, *Nidamen'tum*, *Anhis'tois mem'brane*, (F.) *Caduque*, *C. craie*, *Membrane caduque*, *Épione*, *Périone*;—prior to the time of the Hunters, called *Cho'rium spongi'osum*, seu *tomento'sum seu fung'osum seu reticul'atum*, *Tu'nica filamento'sa*, *Shag'gy Cho'rium*, *Spon'gy Cho'rium*, &c. Great diversity has prevailed regarding this membrane and its reflected portion. It exists before the germ arrives in the uterus—and it has been met with in tubal and ovarian pregnancies; and is occasioned by a new action, assumed by the uterine vessels at the moment of conception. Chaussier, Lobstein, Gardien, Velpeau, and others, consider it to be a sac, without apertures, completely lining the uterus, and that when the ovum descends through the tube it, pushes the decidua before it, and becomes envel-

oped in it, except at the part destined to form the placenta. That portion of the membrane which covers the ovum, forms the *Membra'na decid'ua reflex'a*, (F.) *Caducue réfléchie*, *Membrane caducue réfléchie*, according to them; the part lining the uterus being the *Decid'ua uteri*, (F.) *Caducue utérine*. Towards the end of the fourth month, the decidua reflex disappears. The very existence of a *Tu'nica decid'ua reflex'a* has, however, been denied. This last membrane has received various names. Dr. Granville, regarding it as the external membrane of the ovum, has termed it *Cor'tex o'vi*. It has also been termed *Invol'ucrium membra'naceum*, *Membra'na retiform'is cho'rii seu filamento'sa seu adventit'ia seu cras'sa*, *O'vuline* and *Decid'ua protrusa*. To the membrane which, according to Bojanus and others, is situate between the placenta and the uterus, and which he considers to be produced at a later period than the decidua vera, he gave the name *membra'na decid'ua sero'tina*, (F.) *Membrane utéroëpichoriale Caducue séro'tine ou secondaire*.

Histological researches seem to show, that the decidua is an altered condition of the lining membrane of the uterus, with a whitish secretion filling the uterine tubular glands. The decidua reflexa is probably formed by the agency of nucleated cells from the plastic materials thrown out from the decidua uteri; in the same manner as the chorion is formed in the Fallopian tube from plastic materials thrown out from its lining membrane. That the decidua reflexa is not a mere inverted portion of the decidua uteri is shown by the fact, that the texture of the two is by no means identical.

DECIDUA PROTRUSA, Decidua reflexa.

DECIDUOUS MEMBRANE, Decidua (membrana).

DEC'IGRAMME, *Decigram'ma*, (*decimus*, 'the tenth part,' and *γρᾱμμα*, 'gramme.') The tenth part of the gramme in weight; equal to a little less than two grains, French; 1.543 Troy.

DECIMA'NA FE'BRIS, (*decem*, 'ten.') An intermittent, whose paroxysms return every 10th day or every 9 days. It is supposititious.

DECLAMA'TION, *Declama'tio*, (*de*, and *clamare*, *clamatus*, 'to cry out.') The art of depicting the sentiments by inflections of the voice, accompanied by gestures, which render the meaning of the speaker more evident, and infuse into the minds of the auditors the emotions with which he is impressed. Declamation may become the cause of disease; the modification, produced in the pulmonary circulation,—accompanied by the great excitement, sometimes experienced,—is the cause of many morbid affections; particularly of pneumonia, hæmoptysis, and apoplexy. In moderation, it gives a healthy excitement to the frame.

DÉCLIN, Decline.

DECLINATIO, (*de*, and *clinare*, *clinatus*, 'to bend.') Decline.

DECLINE', *Declina'tio*, *Inclina'tio*, *Decrementum*, *Remissio*, *Parac'mē*, *Parac'nasis*, (F.) *Déclin*. That period of a disorder or paroxysm, at which the symptoms begin to abate in violence, (F.) *Décours*. We speak, also, of the decline of life, or of the powers, (F.) *Déclin de l'age*, *L'age de déclin*, when the physical and moral faculties lose a little of their activity and energy. See Phthisis, and Tabes.

DÉCLIVE (F.), *Decl'vis*, (*de*, and *clivus*, 'acclivity.') Inclining downwards. This epithet is applied to the most depending part of a tumour or abscess.

DÉCOCTÉ, Decoction.

DECOCTION, *Decoct'io*, (*decoquere*, (*de*, and *coquere*, *coctum*), 'to boil.') *Epse'sis*, *Apoz'em*, *Ze'sis*, *Hepse'sis*, *Aphep'sis*, (F.) *Décoction*. The

operation of boiling certain ingredients in a fluid, for the purpose of extracting the parts soluble at that temperature. Decoction, likewise, means the product of this operation, to which the terms *Decoctum*, *Ze'ma*, *Aphepse'ma*, *Ap'ozem*, *Apoz'em*, *Hepse'ma*, *Chyl'us*, and *Epse'ma*, (F.) *Décocté*, *Hydrolé*, *Hydrolite*, have been applied according to ancient custom, in order to avoid any confusion between the operation and its product;—as *præparatio* is used for the act of preparing; *præparatum*, for the thing prepared.

DECOCTION OF ALOES, COMPOUND, Decoction aloes compositum—d. of Bark, Decoction Cinchonæ—d. of Bark, pale, see Decoction Cinchonæ—d. of Bark, red, see Decoction Cinchonæ—d. of Bark, yellow, see Decoction Cinchonæ—d. of Barley, Decoction hordei—d. of Barley, compound, Decoction hordei compositum—d. Bitter, Decoction amarum—d. *Blanche*, Mistura cornu cervi—d. of Broom, Decoction scoparii—d. of Broom, compound, Decoction scoparii compositum—d. of Cabbage tree bark, Decoction geoffrææ inermis—d. of Cassia, Decoction cassiæ—d. of Chamomile, Decoction anthemidis nobilis—d. of Cinchona, Decoction Cinchonæ—d. of Cinchona, compound laxative, Decoction kinæ kinæ compositum et laxans—d. of Colomba, compound, Decoction Colombæ compositum—d. of Dandelion, Decoction Taraxaci—d. of Dogwood, Decoction Cornûs Floridæ—d. of Elm bark, Decoction ulmi—d. of Flaxseed, compound, Decoction Lini compositum—d. of Foxglove, Decoction digitalis—d. *de Guayac composée*, &c., Decoction de Guayac compositum—d. of Guaiacum, compound, Decoction Guaiaci compositum—d. of Guaiacum, compound purgative, Decoction de Guayac compositum—d. of Hartshorn, burnt, Mistura cornu uesti—d. of Hellebore, white, Decoction veratri—d. of Iceland moss, Decoction cetrariæ—d. of Logwood, Decoction hæmatoxyli—d. of Marshmallows, Decoction althææ—d. *de Mauve composée*, Decoction malvæ compositum—d. of Oak bark, Decoction quercûs—d. *d'Orge*, Decoction hordei—d. *d'Orge composée*, Decoction hordei compositum—d. of Pareira brava, Decoction Pareiræ—d. of Pipsissewa, Decoction chimaphilæ—d. of Poppy, Decoction papaveris—d. of Quince seeds, Decoction cydoniæ—d. *de Quinquina composée et laxative*, Decoction kinæ kinæ compositum et laxans—d. *de Salsepareille composée*, Decoction sarsaparillæ compositum—d. of Sarsaparilla, Decoction sarsaparillæ—d. of Sarsaparilla, compound, Decoction sarsaparillæ compositum—d. of Sarsaparilla, false, Decoction araliæ nudicaulis—d. of Squill, Decoction scilla—d. of Tormentil, Decoction Tormentillæ—d. of Uva ursi, Decoction uvæ ursi—d. White, Mistura cornu cervi—d. of the Woods, Decoction Guaiaci compositum—d. of Woody nightshade, Decoction dulcamaræ—d. of Zittmann, Decoction Zittmanni.

DECOCTUM, Decoction—d. Album, Mistura cornu uesti.

DECOCTUM AL'OE'S COMPOS'ITUM, *Decoctum Al'oës*, *Bal'sam of Life*, Compound Decoction of Aloes. (*Est. glyc. 3vij. Potass. carb. ʒi, Aloes, Myrrhæ contrit., Croci aa ʒiiss, Aquæ Oij (Imp. meas)*. Boil to Oj: strain, and add *Tinct. card. c. 3vij. Ph. L.*) The gum and extractive are dissolved in this preparation. The alkali is added to take up a little of the resin. The tincture prevents it from spoiling. It is gently cathartic. Dose, fʒss to ʒij.

DECOCTUM ALTHÆÆ, *Mistura Althææ*, *D. althææ officinalis*: Decoction or Mixture of Marshmallows, (F.) *Décoction de Guimauve*. (*Rad. althææ sic. ʒiv. Uvæ. passar. ʒij, Aquæ Ovij*. Boil to Ov. Pour off the clear liquor. *Ph. E.*) It is used as a demulcent.

DECOCTUM AMARUM; Bitter Decoction. (*Rad. gentian. ʒj, Aquæ Oijss*. Boil for a quarter of an

hour; add *Species amare* ℥ij. Infuse for two hours and filter, without expressing. *Ph. P.*) It is tonic. Dose, f℥ss to ℥ij.

DECOCTUM ANYLI, *Mucilago amyli.*

DECOCTUM ANTHEMIDIS NOBILIS, *D. Chamæ-melli*; *Decoction of Chamomile.* (*Flor. anthemid. nobil.* ℥j, *Sem. carui* ℥iv, *Aquæ Ov.* Boil for fifteen minutes, and strain. *Ph. E.*) It contains bitter extractive and essential oil, dissolved in water. It is used, occasionally, as a vehicle for tonic powders, pills, &c., and in fomentations and gylsters; but for the last purpose, warm water is equally efficacious. The Dublin college had a compound decoction.

DECOCTUM ARA'LIÆ NUDICAUL'IS; *Decoction of False Sarsaparilla.* (*Araliæ nudicaul.* ℥vj, *Aquæ Oviij.* Digest for four hours, and then boil to four pints: press out and strain the decoction. *Former Ph. U. S.*) It is used as a stomachic, but it is an unnecessary and laborious preparation.

DECOCTUM CAS'SLE; *Decoction of Cassia.* (*Cassia pulp.* ℥ij, *Aquæ Oij.* Boil for a few minutes, filter without expression, and add *syrup. violar.* ℥j, or *mannæ pur.* ℥ij. *Ph. P.*) It is laxative, in the dose of ℥vj.

DECOCTUM CETRA'RIÆ, *Decoction of Lichen Islandicus.* (*Decoction of Iceland Moss.* (*Cetrar.* ℥ss, *Aquæ Oiss.* Boil to a pint, and strain with compression. *Ph. U. S.*) It consists of bitter extractive and fecula, dissolved in water, and its operation is tonic and demulcent. Dose, f℥j to ℥iv. It is also nutrient, but hardly worthy of the rank of an official preparation.

DECOCTUM CHAMÆMELI, *Decoction of anthemidis nobilis.*

DECOCTUM CHIMAPH'ILÆ, *Decoction of Pipsissewa.* (*Chimaph. contus.* ℥j, *Aquæ Oiss.* Boil to a pint, and strain. *Ph. U. S.*) A pint may be given in the course of the twenty-four hours.

DECOCTUM CINCHO'NÆ, *D. Corticis Cincho'næ seu Ki'næ Ki'næ, Decoction of Cincho'na or of Bark.* (*Cinchon. cort. contus.* ℥j, *Aquæ Oj.* Boil for ten minutes in a slightly-covered vessel, and strain while hot, adding water to make a pint. It contains quinia and resinous extractive, dissolved in water. Long coction oxygenates and precipitates the extractive. It can be given where the powder does not sit easy, &c. Dose, f℥j to ℥iv.

The Pharmacopœia of the United States has a *Decoction Cincho'næ flavæ, Decoction of yellow bark*, and a *Decoction Cincho'næ rubræ, Decoction of red bark*, both of which are prepared as above. The London, Edinburgh, and Dublin Colleges have a *Decoction Cinchonæ pallidæ, Decoction of Pale Bark.*

DECOCTUM COLOM'BÆ COMPOSITUM; *Compound Decoction of Colomba.* (*Colomb. contus.* ℥ss, *Quassia* āā ℥ij, *Cort. aurant.* ℥j, *Rhei pulv.* ℥j, *Potassæ carbonat.* ℥ss, *Aquæ* ℥xxx. Boil to a pint, and add *tinct. lavand.* f℥ss. *Former Ph. U. S.*) Given as a tonic, but not worthy an official station.

DECOCTUM COMMUNE PRO CLYSTERE, *D. Malvæ compositum*—*d. Cornu cervini, Mistura cornuusti.*

DECOCTUM COR'NUS FLORIDÆ, *Decoction of Dogwood.* (*Cor'nus Florid.* cont. ℥j, *Aquæ Oj.* Boil for ten minutes in a covered vessel, strain while hot, and add water to make a pint. *Ph. U. S.*) Dose, f℥ss.

DECOCTUM CYDO'NIÆ, *Mucilago Seminis Cydoniæ Mal'i seu Seminum Cydoniorum*; *Decoction or Mucilage of Quince seeds.* (*Cydoniæ sem.* ℥ij, *Aquæ Oj.* Boil for ten minutes over a gentle fire, and strain. *Ph. L.*) It is merely a solution of mucilage in water, and is used as a demulcent.

DECOCTUM DAPH'NES MEZE'REI, *Decoction Me-*

ze'rei, Decoction of Meze'reon. (*Meze'rei* ℥ij, *Glycyrrh.* cont. ℥j, *Aquæ Oij.* Boil over a gentle fire to Oiss, and strain. *Ph. E.*) The acrimony of the meze'reon and the saccharine mucilage of the liquorice root are imparted to the water. It is somewhat stimulant, and has been used in secondary syphilis; but is devoid of power. Dose, f℥ij to ℥vj.

DECOCTUM DIAPHORETICUM, *D. Guaiaci compositum.*

DECOCTUM DIGITA'LIS, *Decoction of Fox-glove.* (*Fol. digit. sicc.* ℥j, *Aquæ q. s. ut colentur* f℥viij. Let the liquor begin to boil over a slow fire, and then remove it. Digest for fifteen minutes, and strain. *Ph. D.*) It possesses the properties of the plant. Dose, f℥j to ℥ij.

DECOCTUM DULCAMA'RÆ, *Decoction of Wood'y Nightshade.* (*Dulcamaræ cont.* ℥j, *Aquæ Oiss.* Boil to Oj, and strain. *Ph. U. S.*) This decoction has been considered diuretic and diaphoretic, and has been administered extensively in skin diseases. It is probably devoid of efficacy.

DECOCTUM PRO FOMENTO, *D. papaveris.*

DECOCTUM GEOFFRÆ'Æ INERMIS; *Decoction of Cab'bagetree Bark.* (*Cort. geoffr. inermis,* in pulv. ℥j, *Aquæ Oij.* Boil over a slow fire to a pint, and strain. *Ph. E.*) It is possessed of anthelmintic, purgative, and narcotic properties, and has been chiefly used for the first of these purposes. Dose, to children, f℥ij—to adults, f℥ss to ℥ij.

DECOCTUM GUA'ACI COMPOSITUM, *D. Guaiaci officinalis compositum seu Ligno'rum seu de Guy'aco compositum seu eudorificum seu diaphoreticum*; *Compound decoction of Guaiacum, Decoction of the Woods.* (*Lign. guaiac. rasur.* ℥ij, *fruct. sicc. vitis vinifer.* ℥ij, *rad. lauri sassafr. concis.* rad. glycyrrh. āā ℥j, *aquæ Ox.* Boil the Guaiacum and raisins over a slow fire to Ov; adding the roots towards the end, then strain. *Ph. E.*) It is possessed of stimulant properties, and has been given in syphilitic, cutaneous, and rheumatic affections. The resin of the guaiacum is, however, insoluble in water, so that the guaiac wood in it cannot be supposed to possess much, if any, effect. The Parisian Codex has a

DECOCTUM DE GUY'ACO COMPOSITUM ET PUR'GANS, (*F.*) *Décoction de Gayac Composée et Purgative*; *Compound purgative Decoction of Guaiacum.* (*Lign. guaiac. rasp., rad. sarsap. āā ℥j, potassæ carbonat. gr. xxv.* Macerate for twelve hours, agitating occasionally in water Oiv, until there remain Oij. Then infuse in it *fol. sennæ* ℥ij, *rhei* ℥j, *lign. sassafr., glycyrrh. rad. āā ℥ij, sem. coriand.* ℥j. Strain gently, suffer it to settle, and pour off the clear supernatant liquor. The title sufficiently indicates the properties of the composition.

DECOCTUM HEMATOXYLI, *Decoction of Logwood.* (*Hæmatozyl. rasur.* ℥j, *aquæ Oiss.* Boil to a pint, and strain. *Ph. U. S.*)

DECOCTUM HELLEBORI ALBI, *D. veratri.*

DECOCTUM HOR'DEI, *Decoction Hordei Distichi, P'tis'ana Hippocrat'ion, Tipta'ria, Tapsa'ria, A'qua Horde'ia, Hydroceri'the, Bar'ley Water, Decoction of Bar'ley.* (*F.*) *Décoction d'Orge, Tisane Commune.* (*Hord.* ℥ij, *Aquæ Oivss.* Wash the barley well, boil for a few minutes in water, Oss: strain this, and throw it away, and add the remainder, boiling. Boil to Oij, and strain. *Ph. L.*) It is nutritive and demulcent, and is chiefly used as a common drink, and in gylsters.

DECOCTUM HOR'DEI COMPOSITUM, *Decoction pectoral'is, Mistura Hor'dei, P'tis'ana communis*; *Compound decoction of Bar'ley.* (*F.*) *Décoction d'Orge composée.* (*Decoct. hord. Oij, caricæ fruct. concis.* ℥iiss, *glycyrrh. rad. concis. et contus.* ℥vj, *uværum pass. demptis acinis,* ℥ijss, *aquæ Oj.* Boil to Oij, and strain. *Ph. L.*) It has similar properties to the last.

DECOCTUM KINÆ KINÆ, D. Cinchonæ.

DECOCTUM KINÆ KINÆ COMPOSITUM ET LAXANS, *Décoction de quinquina composée et laxative*; *Com'pound lax'ative decoction of Cincho'na.* (Cort. cinchon. $\mathfrak{z}\mathfrak{j}$, aquæ Oij. Boil for a quarter of an hour, remove from the fire; then infuse in it, for half an hour, fol. senæ, sodæ sulph. aa $\mathfrak{z}\mathfrak{j}$: add *syrup. de sennâ* f $\mathfrak{z}\mathfrak{j}$.) Its title indicates its properties.

DECOCTUM LICHENIS ISLANDICI, Decoction Cetrariæ—d. Lignorum, D. Guaiaci compositum.

DECOCTUM LINI COMPOSITUM, *Com'pound Decoction of Flaxseed.* (Lini $\mathfrak{z}\mathfrak{j}$, Glycyrrhiz. contus. $\mathfrak{z}\mathfrak{ss}$, aquæ Oiss. Boil for ten minutes, and strain. Ph. D.)

DECOCTUM MALVÆ COMPOSITUM, *Decoction pro enem'atē seu commu'nē pro clyst'ere*; *Com'pound Decoction of Mal'low.* (F.) *Décoction de Mauve composée.* (Malvæ exsic. $\mathfrak{z}\mathfrak{j}$, anthemid. flor. exsic. $\mathfrak{z}\mathfrak{ss}$, aquæ Oj. Boil for fifteen minutes and strain.) It consists of bitter extractive, and mucilage in water, and is chiefly used for clysters and fomentations. It is unworthy a place in the pharmacopœias.

DECOCTUM MEZEREI, D. daphnes mezerei.

DECOCTUM PAPAV'ERIS, *Decoction pro Fomen'to, Fo'tus commu'nis*; *Decoction of Pop'py.* (Papav. somnif. capsul. concis. $\mathfrak{z}\mathfrak{i}\mathfrak{v}$, aquæ Oiv. Boil for fifteen minutes, and strain. Ph. L.) It contains the narcotic principle of the poppy, and mucilage in water: is anodyne and emollient, and employed as such in fomentation, in painful swellings, ulcers, &c.

DECOCTUM PAREIRÆ, *Decoction of Parei'ra Bra'va.* (Pareir. brav. $\mathfrak{z}\mathfrak{x}$, Aq. Destillat Oiss. (Imp. meas.) Boil to a pint and strain. Dose, f $\mathfrak{z}\mathfrak{j}$ to $\mathfrak{z}\mathfrak{i}\mathfrak{j}$.)

DECOCTUM PECTORALE, D. hordei compositum —d. Polygalæ Senegæ, D. Senegæ.

DECOCTUM PRO ENEMATE, D. Malvæ compositum.

DECOCTUM QUERCUS ALBÆ, D. Quercus Roboris; *Décoction of White Oak Bark.* (Quercus cort. $\mathfrak{z}\mathfrak{j}$, aquæ Oiss. Boil to a pint, and strain. Ph. U. S.) It is astringent, and used as such, in injections, in leucorrhœa, uterine hemorrhage, &c., as well as in the form of fomentations to unhealthy ulcers.

DECOCTUM SARSÆ, Decoction Sarsaparillæ.

DECOCTUM SARSAPARILLÆ, D. Smilacis Sarsaparillæ, D. Sar'æ, D. Sar'æ; *Décoction of Sarsaparill'a.* (Sarsaparill. rad. concis. $\mathfrak{z}\mathfrak{v}$, aq. fervent. Oiv. (Imp. meas.) Macerate for four hours near the fire, in a lightly-covered vessel; then bruise the root; macerate again for two hours; then boil to Oij, and strain. Ph. L.) It contains bitter extractive, and mucilage in water; is demulcent, and has been used, although it is doubtful with what efficacy, in the sequelæ of syphilis. Dose, f $\mathfrak{z}\mathfrak{i}\mathfrak{v}$ to Oss.

DECOCTUM SARSAPARILLÆ COMPOSITUM; *Com'pound Decoction of Sarsaparill'a.* (F.) *Décoction de Salsepareille Composée.* (Sarsaparill. concis. et contus $\mathfrak{z}\mathfrak{v}\mathfrak{j}$, Sassafr. Cort. concis., Lign. Guaiac. rasur., Rad. Glycyrriz. contus., aa $\mathfrak{z}\mathfrak{j}$, Mezerei, concis. $\mathfrak{z}\mathfrak{i}\mathfrak{v}$, Aquæ Oiv. Boil fifteen minutes, add water to make the whole Oiv, and strain. Ph. U. S.) This is considered to be possessed of analogous properties to the celebrated *Lis'bon Diet-Drink*, *Decoction Lusitan'icum*, which it resembles in composition.

DECOCTUM SARZÆ, Decoction Sarsaparillæ.

DECOCTUM SCILLÆ; *Décoction of Squill.* (Scillæ. $\mathfrak{z}\mathfrak{i}\mathfrak{ij}$, juniper. $\mathfrak{z}\mathfrak{i}\mathfrak{v}$, Senegæ. $\mathfrak{z}\mathfrak{i}\mathfrak{v}$, Aquæ Oiv. Boil till one half the liquor is consumed; strain, and add *spirit of nitrous ether*, f $\mathfrak{z}\mathfrak{i}\mathfrak{v}$. Ph. U. S.) Dose, f $\mathfrak{z}\mathfrak{ss}$.

DECOCTUM SCOPARII, *Décoction of Broom* (Scopar. exsicat. $\mathfrak{z}\mathfrak{ss}$, (avoidupois); Aquæ Oss. (Imp. meas.) Boil for ten minutes and strain.—Ph. D.)

DECOCTUM SCOPARII COMPOSITUM, D. Scoparii (Ph. E.), *Com'pound Decoction of Broom* (Scopar., Junip. cont., Taraxac. cont. aa $\mathfrak{z}\mathfrak{ss}$; Aq. destillat. Oiss. (Imp. meas.) Boil to a pint, and strain. Ph. L.) Dose, half a pint to a pint, in the day; as a diuretic in dropsy.

DECOCTUM SENEGÆ, D. Polygalæ Senegæ; *Décoction of Senega.* (Seneg. cont. $\mathfrak{z}\mathfrak{j}$, aquæ Oiss. Boil to Oj, and strain. Ph. U. S.) It is reputed to be diuretic, purgative, and stimulant, and has been given in dropsy, rheumatism, &c.

DECOCTUM SMILACIS SARSAPARILLÆ, D. Sarsaparillæ—d. Sudorificum, D. Guaiaci compositum.

DECOCTUM TARAXACI, *Décoction of Dan'delion.* (Taraxac. contus. $\mathfrak{z}\mathfrak{i}\mathfrak{v}$, aquæ Oiss. (Imp. meas.) Boil to a pint, and strain. Ph. L.) Dose f $\mathfrak{z}\mathfrak{i}\mathfrak{ss}$.

DECOCTUM TORMENTILLÆ, *Décoction of Tormentil.* (Tormentil. cont. $\mathfrak{z}\mathfrak{i}\mathfrak{j}$, Aquæ destillat. Oiss. (Imp. meas.) Boil to a pint, and strain.) Dose, as an astringent, one or two fluid ounces.

DECOCTUM ULMI, D. Ulmi Campest'ris, *Décoction of Elm Bark.* (Ulm. cort. recent. cont. $\mathfrak{z}\mathfrak{i}\mathfrak{ss}$, aquæ Oij. (Imp. meas.) Boil to Oj, and strain. Ph. L.) It is a reputed diuretic, and has been used in lepra and herpes; but, probably, has no efficacy. Dose, f $\mathfrak{z}\mathfrak{i}\mathfrak{v}$ to Oss.

DECOCTUM URSI URSI, *Décoction of u'ra ur'si.* (Ursi ur'si $\mathfrak{z}\mathfrak{j}$, aquæ f $\mathfrak{z}\mathfrak{xx}$. Boil to a pint and strain. Ph. U. S.) Dose, f $\mathfrak{z}\mathfrak{i}\mathfrak{ss}$.

DECOCTUM VERA'TRI, *Décoction Helleb'ori al'bi*; *Décoction of White Hel'lebore.* (Veratri rad. cont. $\mathfrak{z}\mathfrak{j}$, aquæ Oij, spir. rect. f $\mathfrak{z}\mathfrak{i}\mathfrak{j}$. Boil the watery decoction to Oj, and, when it is cold, add the spirit. Ph. L.) It is stimulant, acrid, and cathartic; but is used only externally. It is a useful wash in *tinea capitis*, *psora*, &c. Should it excite intense pain on being applied, it must be diluted.

DECOCTUM ZITTMAN'NI, *Zitt'man's Decoction.* A most absurd farrago, extolled by Theden in venereal diseases, the formula for which, according to Jourdan, "some blockheads have lately reproduced among us as a novelty." It is composed of *sarsaparilla*, *pulvis stypticus*, *calomer* and *cinnamon*, boiled in water with *aniseed*, *fennel seed*, and *liquorice root*. A stronger and a weaker decoction were directed by Zittman. Formulæ for its preparation are contained in Jourdan's *Pharmacopœia Universalis*; and in Lincke's *Vollständiges Recept-Taschenbuch*. Leipz., 1841.

DECOLLATION, *Decollatio*, *Decapitatio*, *Decapitatio*, *Obtruncatio*, (*de* and *collum*, 'the neck.') The act of separating the head from the trunk, by the section of the neck, as in cases of dystocia.

DÉCOLLEMENT, (F.) *Deglutina'tio*, (*de* and *coller*, 'to glue.') The state of an organ that is separated from the surrounding parts, owing to destruction of the areolar membrane which united them. The skin is *décollée*, i. e. separated from the subjacent parts, by a burn, subcutaneous abscess, &c.

DÉCOLLEMENT DU PLACENTA, is the separation or detachment of the whole or a part of the placenta from the inner surface of the uterus. *Décollement*, from *de*, 'from,' and *collum*, 'the neck,' *obtruncatio*, also means the separation of the head of the fœtus from the trunk, the latter remaining in the uterus.

DECOLORATION, *Decolora'tio*, *Discolora'tio*, (*de*, and *colorare*, *coloratum*, 'to colour.') Loss of the natural colour; *Parachro'sis*. Devoid of colour, —*Achroma'sia*.

In Pharmac' any process by which liquids, or solids in solution, are deprived wholly or in part of their colour,—as by passing them through animal charcoal.

DÉCOMPOSÉE (F.), (*de*, and *composere*, *compositum*, 'to compose.') *Decomposed*, *Dissolutio*. An epithet, applied to the face when extensively

changed in its expression and colour; as in the choleric or moribund.

DECORTICATION, *Decortica'tio*, (*de*, and *cortex*, gen. *corticis*, 'bark.') An operation, which consists in separating the bark from roots, stalks, &c. The detachment of portions of enamel from a tooth is sometimes termed the *decor'ticating proc'ess*. (F.) *Cavie écorchante*.

DECOSTIS, *Apleu'ros* (*de*, and *costa*, 'a rib.') One who has no ribs.

DÉCOURS, Decline.

DECREMENTUM, (*de*, and *crescere*, 'to grow.') Decline.

DECREPIT, (*de*, and *crepere*, *crepitum*, 'to creak.') See Decrepitude.

DECREPITUDE, *Decrepitu'do*, *Æ'tas decrep'ita*, *Ultima senec'tus*, *Senec'ta decrep'ita* seu *extrema* seu *sum'ma* seu *ult'ima*. The last period of old age, and of human life; which ordinarily occurs about the eightieth year. It may, however, be accelerated or protracted. Its character consists in the progressive series of phenomena which announce the approaching extinction of life. One in this stage of existence is said to be *decrep'it*, *decrep'itus*, (F.) *décépité*.

DECRETORII DIES, (*decretorius*, 'decisive.') Critical days.

DECUBITUS, gen. *Decubitûs*, (*decumbere*, (*de*, and *cubere*, *cubitum*, 'to lie,') 'to lie down.') *Cubatio*. Lying down. Assuming the horizontal posture, *cubitus sup'inus*; (F.) *Concher*. *Ho'râ decu'bitûs*, 'at bed time.' The French say—*Decubitus horizontal*, *sur le dos*, *sur le côté*, for, lying in the horizontal posture, on the back or side.

DECURTATUS, *Meiu'rus*, (*de*, and *curtus*, 'short,') 'shortened,') 'curtailed,') 'running to a point.' When applied to the pulse, it signifies a progressive diminution in the strength of the arterial pulsations, which, at last, cease. If the pulsations return and gradually acquire all their strength, it is called *Pul'sus decurtatus recip'ro-curs*.

DECUSSATIO, Decussation—d. Pyramidum, see Corpora pyramidalia.

DECUSSATION, *Decussatio*, *Chias'mus*, *Incrucia'tio*, *Intersec'tio*, *Intracatu'ra*, (from *decussis*, that is, *decem asses*; also, the figure of the letter X.) Union in the shape of an X or cross. Anatomists use this term chiefly in the case of the nerves—as the *decussation of the optic nerves*, which cross each other within the cranium.

DECUSSATION OF THE PYRAMIDS, see Corpora pyramidalia.

DECUSSORIUM, (*decutio*, *decussum*, (*de*, and *quatio*, 'I shake down,') (F.) *Décu'ssoire*. An instrument used by the ancients, for depressing the dura mater, and facilitating the exit of substances effused on or under that membrane. It is described by Scultetus, Paré, &c. See Meningophylax.

DÉDAIGNEUX, ('disdainful,') from *de*, and *dignor*, 'to esteem worthy.' Rectus superior oculi.

DEDE (Sc.). (Sax. *dead*), Death.

DEDENTITION, see Dentition.

DEDOLATION, *Dedola'tio*, (*dedolare*, (*de*, and *dolare*, *dolatum*,) 'to cut and hew with an axe.') This word has been used by surgeons to express the action by which a cutting instrument divides obliquely any part of the body, and produces a wound with loss of substance. It is commonly on the head, that wounds by dedolation are observed. When there was a complete separation of a portion of the bone of the cranium, the ancients called it *Aposeparnismus*.

DEERBALLS, *Elaphomyces granulatus*.

DEERBERRY, *Gaultheria*, *Vaccinium stamineum*.

DEERFOOD, *Brasenia hydropeltis*.

DEERS' TEARS, Bezoar of the Deer.

DEFÆCATION, (*de*, and *fæx*, gen. *fæcis*, 'excrement.') Defecation.

DÉFAILLANCE, (from *defaillir*, (*de*, and *failir*,) 'to fail.') Syncope.

DEFÆCATION, *Defæca'tio*. The act by which the excrement is extruded from the body. *Caca'tio*, *Ezere'tio al'vi* seu *alvi'na* seu *fæcum alvina'rûm*, *Defec'tio al'vi*, *Seces'sio*, *Ecpul'sio* seu *Ejec'tio fæcum*, *Apago'gê*, *Hypochore'sis*, *Eechore'sis*, *Eccop'o'nis*, *Eges'tio*. The fæces generally accumulate in the colon, being prevented by the annulus at the top of the rectum from descending freely into that intestine. In producing evacuations, therefore, in obstinate constipation, it is well, by means of a long tube, to throw the injection, into the colon.

In *Pharmacy*, defecation means the separation of any substance from a liquid in which it may be suspended. See Clarification.

DEFECTIO ANIMI, (*deficere*, *defectum*, (*de*, and *facere*,) 'to forsake,') Syncope.

DEFECTUS LOQUELÆ, Aponia—d. Veneris, Anaphrodisia.

DEFENSIVUM, (*defendere*, *defensum*, 'to defend,') (F.) *Défensif*. A preservative or defence. The old surgeons gave this name to different local applications, made to diseased parts, for the purpose of guarding them from the impression of extraneous bodies, and particularly from the contact of air.

DEFERENS, (*defero*, (*de*, and *fero*,) 'I bear away,') 'I transport,') Carrying away.

DEFERENS VAS, *Duct'us* seu *Canalis def'erens*, *Vibrato'r*, *Spermat'ic duct*, *Sper'miduct*, (F.) *Canal ou Conduit spermatique*, *Canal ou Conduit déférent*, is the excretory canal of the sperm, which arises from the epididymis, describes numerous convolutions, and with the vessels and nerves of the testicles concurs in the formation of the spermatic chord, enters the abdominal ring, and terminates in the ejaculatory duct.

DEFIBRINATED, (*de*, and *fibrin*,) Defibrinated.

DEFIBRATION, see Defibrinized.

DEFIBRINIZED, (F.) *Définé*, *Defib'rinated*. Deprived of fibrin. A term applied to blood from which the fibrin has been removed, as by whipping. The act of removing fibrin from the blood has been termed *defibrination*.

DEFIGURATIO, (*de*, and *figurare*, *figuratum*, 'to shape,') Deformation.

DEFLAGRATIO, *Deflagra'tio*, (*deflagrare*, (*de*, and *flagrare*, *flagratum*, 'to burn,') 'to set on fire,') In pharmacy, the rapid combustion of a substance with flame; great elevation of temperature; violent motion, and more or less noise. Thus, we speak of the deflagration of the nitrate and the chlorate of potassa, of gunpowder, &c.

DEFLECTENS, (*de*, and *flecto*, *flexum*, 'to bend,') Derivative.

DEFLECTIO, Derivation.

DEFLORATION, (*deflorescere*, (*de*, and *florescere*,) 'to wither,') *Depucella'tion*, *Depudica'tion*, *Devirgination*, *Defloratio*, *Devirgination*, *Vir'ginis vitia'tio*, *Virgin'itas deflorata*, (F.) *Défloration*, *Dépucellement*. The act of depriving a female of her virginity. Inspection of the parts is the chief criterion of defloration having been forcibly accomplished; yet inquiry must be made, whether the injury may not have been caused by another body than the male organ. Recent defloration is infinitely more easy of detection than where some time has elapsed. See Stuprum.

DEFLUVIUM, (*de*, and *fluere*, *fluxum*, 'to flow,') Aporrhœa—d. Capillorum, Alopecia—d. Pilorum, Alopecia.

DEFLUXIO, Catarrh, Defluxion, Diarrhœa—d. Catarrhalis, Influenza.

DEFLUXION, *Deflux'io*, *Deflux'us*, *Catar'rh-*

sis, *Hyporrhœia*, *Emorrhœia*. A falling down of humours from a superior to an inferior part. It is sometimes used synonymously with inflammation.

DEFLUXUS DYSENTERICUS, Dysentery.

DEFORMA'TION, *Deformatio*, (de, and *forma*, 'form,') *Cacomor'phia*, *Cacomorpho'sis*, *Dysmor'phê*, *Dysmor'phia*, *Dysmorpho'sis*, *Deform'itus*, *In-form'itas*, *Defigura'tio*, *Disfigura'tion*, *Deform'ity*, (F.) *Difformité*. Morbid alteration in the form of some part of the body, as of the head, pelvis, spine, &c. A deformity may be natural or accidental.

DEFUR'TUM, (from *Defervitum*, (?) *de*, and *ferreo*, 'to boil,') (F.) *Vin cuit*. Grape must, boiled down to one half and used as a sweetmeat. See Rob.

DEFURFURATIO, (de, and *furfur*, 'bran,') See Desquamation.

DEFUSIO, (de, and *fundere*, *fusum*, 'to pour,') Decantation.

DEGENERATIO, Degeneration—d. *Adiposa cordis*, Steatosis cordis.

DEGENERATION, *Degenera'tio*, *Nothen'sis*, *Nothi'a*, (from *degener*, 'unlike one's ancestors,' [de, and *genus*, *genêrîs*, 'family,'] *Degen'eracy*, (F.) *Dégénération*, *Abâtardissement*. A change for the worse—degradation—in the intimate composition of the solids or fluids of the body. In pathological anatomy, *degeneration* means the change which occurs in the structure of an organ, when transformed into a matter essentially morbid; as a *cancerous*, or *tubercular*, degeneration. Virchow employs the term *neerobio'sis* in a similar signification; and for morbid metamorphosis in general.

Dégénérescence is, by the French pathologists, employed synonymously with Degeneration.

DÉGÉNÈRESCENCE, Degeneration—d. *Graisseuse du Foie*, *Adiposis hepatica*—d. *Granule du Rein*, *Kidney*, Bright's disease of the—d. *Noire*, *Melanosis*.

DEGLUTINATIO, (de, and *gluten*, gen. *glutinis*, 'glue,') *Décollement*.

DEGLUTITIO, Deglutition—d. *Difficilis*, *Dysphagia*—d. *Impedita*, *Dysphagia*—d. *Læsa*, *Dysphagia*.

DEGLUTIT'ION, *Degluti'tio*, *Catap'osis*, (de, and *glutire*, *glutitum*, 'to swallow,') The act by which substances are passed from the mouth into the stomach, through the pharynx and œsophagus. It is one of a complicated character, and requires the aid of a considerable number of muscles; the first step being voluntary, the remainder executed under spinal and involuntary nervous influence.

DEGMUS, (*δῆμος*, from *δῆκω* or *δάκω*, 'I bite,') *Dex'is*, *Mor'sus*, (F.) *Morsure*,—a bite in general. A gnawing sensation about the upper orifice of the stomach, which was once attributed to acrimony of the liquids contained in that viscous.

DÉGORGEMENT, (de, and *gorge*, 'the throat,') Disgorgement.

DÉGOUT, (de, and *gout*, (L.) *gustus*, 'taste,') Disgust.

DEGRADATION, (*degradior*, (de, and *gradior*, 'to step,') 'to descend,') Degeneration.

DEGREE', ([F.] *dégré*, originally from *gradus*, 'a step,') A title conferred by a college,—as the 'degree of Doctor of Medicine.' Galen used this expression to indicate the qualities of certain drugs. Both he and his school admitted *cold*, *warm*, *moist*, and *dry* medicines, and four different 'degrees' of each of those qualities. Thus, *Apium* was warm in the first degree, *Agrimony* in the second, *Roche Alum* in the third, and *Garlic* in the fourth. *Bedegar* was cold in the first, the flower of the Pomegranate in the second, the *Sempervivum* in the third, *Opium* in the fourth, &c. The French use the term *degré* to indicate, 1. The in-

tensity of an affection: as a burn of the first, second, third, degree, &c. 2. The particular stage of an incurable disease, as the third degree of phthisis, cancer of the stomach, &c.

DEGUSTA'TION, *Degusta'tio*, (de, and *gustare*, *gustatum*, 'to taste,') *Gusta'tion*. The appreciation of sapid qualities by the gustatory organs.

DEHIS'CENCE, *Dehiscen'tia*, (*dehiscere*, (de, and *hiscere*, 'to gape,') 'to break open in chinks,') Separation of a shut organ along the line of union. A regular and determinate separation, which takes place in a shut organ.

DELECTIOUS, (*δελτικός*, from *δεικνυμι*, 'to point out,') Index.

DEID, (Sc.) ([Sax.] *deað*.) Death.

DEIL'S SNUFFBOX, *Lycoperdon*.

DEIRONCUS, (*δειρῆ*, 'neck,' and *ογκος*, 'a swelling,') *Bronchocele*.

DEJECTEDNESS, Depression.

DEJECTIO, Dejection—d. *Alvi*, *Defecation*—d. *Alvina*, *Excrement*.

DEJECTION, (*deicere*, *dejectum*, (de, and *jacere*,) 'to cast down,') *Dejec'tio*, *Subduc'tio*, *Hypago'gê*, *Hypochore'sis*, *Hyechore'sis*, *Apop'atus*, *Hypop'atus*, *Hypoph'ora*. The expulsion of the fæces;—*Ejec'tio*, *Eges'tio*. Also, a fecal discharge or stool,—generally, however, with *alvini* prefixed,—as an *alvine Dejection*, *Alvus vir'idis*, *Dejec'tio alvi'na*, a passage; an evacuation; a movement. Also depression of spirits.

DEJECTIONES NIGRÆ, ('black dejections,') *Melena*.

DEJECTORIUM, Cathartic.

DELACRYMATIO, (de, and *lacryma*, 'a tear,') *Epiphora*.

DELACRYMATIVUS, *Apodaeryticus*.

DELAPPIO, (de, and *labi*, *lappus*, 'to fall,') *Prolapsus*.

DELAPSUS, *Prolapsus*—d. *Palpebræ*, *Blepharoptosis*.

DELATIO, (*deferre*, *delatum*, 'to announce,') Indication.

DÉLA YANTS, *Diluentia*.

DELAEROIX'S DEPILATORY, (after the proposer,) see *Depilatory*, *Colley's*.

DELETE'RIOUS, *Delete'rius*, *Pernicio'sus*, *Phthar'cticus*, (F.) *Délétère*, *Pernicieux*, (*δηλεω*, 'I injure,') That which produces destructive disorder in the exercise and harmony of the functions.

DELETERIUM, Poison.

DELHI BOIL. An inveterate cutaneous affection, prevalent at Delhi, in India, which assumes the form of a boil, and may, afterwards, ulcerate, forming the *Delhi ulcer*. It is called, there, *Arungzebe*, owing to the emperor of that name having died with it. The treatment is chiefly local, and consists of the ordinary eutrophic ointments. Eutrophics may, also, be given internally.

DELHI ULCER, see *Delhi Boil*.

DELIBRET, Delirious.

DELIGATION, Deligation, Ligature, see *Bandage*.

DELIGA'TION, *Deliga'tio*, *Epid'esis*, *Deligatu'ra*, *Vul'nerum deliga'tio* seu *vinctu'ra*, *Fascia'tum* *Applica'tio*, *Plaga'rum* *Vinctu'ra*, *Fascia'tio*, (from *deligare*, *deligatum*, (de, and *ligo*,) 'to bind,') The deligation of wounds formerly embraced the application of apparatus, dressings, &c.,—the denomination *Deliga'tor* *Plaga'rum* being synonymous with *Med'icus Vulnera'rius*, and in derivation, with the *Wundarzt*, 'wound physician' or surgeon, of the Germans. Deligation is hardly ever used now as an English word. In France, it is applied to the regular and methodical application of bandages, and to the ligature of arteries, and, occasionally, in the latter sense, with us.

DELIGATURA, Deligation.

DELIQUESCENT, *Deliquescent*, (*deliquescere*, (*de*, and *liquescere*.) 'to melt,' 'to dissolve.') Any salt which becomes liquid by attracting moisture from the air. The deliquescent salts require to be kept in bottles, well stopped. Chloride of lime, acetate of potassa, and carbonate of potassa, are examples of such salts. The ancient chemists expressed the condition of a body, which had become liquid in this manner, by the word *Deliquium*.

DELIQUIUM ANIMI, (*delinquere*, *delictum*, 'to leave.') Syncope.

DELIRAMENTUM, Delirium.

DELIRANS, Delirious.

DELIRANT, Delirious.

DELIRATIO, Delirium—*d. Senum*, see Dementia.

DÉLIRE, Delirium—*d. Crapuleux*, Delirium tremens—*d. des Ivrognes*, Delirium tremens—*d. Nervex*, Delirium nervosum.

DÉLIRE DES PERSÉCUTIONS. Mania, in which the morbid hallucination is, that the individual is persecuted.

DÉLIRE TRAUMATIQUE, Delirium nervosum—*d. Tremblant*, Delirium tremens.

DELIRIA, Insanity.

DELIRIE, Delirious.

DELIRIOUS, *Delirans*, *Delirus*, *Excerebratus*, *Alienus*, *Paraleurus*, *Raving*, (Sc.) *Delie'ret*, *Delirie*, (F.) *Déli-rant*. One who is in a state of delirium. That which is attended by delirium. The French use the term *Fièvre pernicieuse délirante* for a febrile intermittent, in which delirium is the predominant symptom.

DELIRIUM, (perhaps, from *de*, and *lira*, 'a ridge between two furrows': more probably from *de*, and *ληρημα*, 'a silly saying or action.') *Paracopé*, *Phrenitis*, *Phledonia*, *Desipientia*, *Aphrosyne*, *Paralere'ma*, *Paralere'sis*, *Paralog'ia*, *Phanta'sia*, *Paraphrosyne*, *Emotio*, *Le'tos*, *Parana'a*, *Alloph'asis*, *Deliratio*, *Deliramentum*, (Sc.) *Rave'ry*, *Ro'ving*, (F.) *Délire*, *Égarement d'esprit*, *Transport*, *Idéosynchysie*, &c. Straying from the rules of reason; wandering of the mind. Hippocrates used the word *pavia*, *mania*, for *delirium si'ne fe'bre*, and the Greek words given above for *delirium cum fe'bre*. In different authors, also, we find the words, *Paraph'ora*, *Paraphrene'sis*, *Paraphren'ia*, *Phrene'sis*, *Phreneti'asis*, &c., for different kinds of delirium. Delirium is usually symptomatic.

DELIRIUM EBRIOSITATIS, 'delirium of drunkenness' also *D. tremens*—*d. Epileptic*, see Epilepsy—*d. Furiosum*, *Mania*—*d. Maniacum*, *Mania*.

DELIRIUM NERVO'SUM, (F.) *Délire nerveux* or *traumatique*. A form of delirium which supervenes after surgical operations, or severe injuries.

DELIRIUM POTATORUM, *D. Tremens*.

DELIRIUM TRE'MENS, *D. potatorum* seu *ebriositatis* seu *tremifaciens* seu *vig'ilans*, *Mania a potu* seu *e temulentia*, *Enomonia*, *Erethismus ebriosorum*, *Dipsomonia*, *Meningitis* seu *Phrenitis potatorum*, *Meningitis phantasmatorum*, *Tromomonia*, *Tromoparanc'a*, *Potoparanc'a*, *Potomonia*, *Pototroparanc'a*, (F.) *Encéphalopathie crapuleuse*, *Délire tremblant* ou *crapuleux* ou *des Ivrognes*, *Folie des Ivrognes* ou *Alcoolique*. A state of delirium and agitation,—often termed 'the horrors'—peculiar to those addicted to spirituous liquors, with great sleeplessness. It is preceded by indisposition, lassitude, watchfulness, headache, and anorexia; the delirium and tremors, *Tremor potatorum*, most commonly recurring in paroxysms. It is caused by the habitual and intemperate use of ardent spirits or of opium or tobacco; or by abandoning them after prolonged use. The treatment is various. Many have regarded stimulants, with large doses of

opium to induce rest, to be indispensable. It is certain, however, that the expectant system will often, if not generally, be successful; and, it is probable, a cure effected in this manner will be more permanent than when produced by excitants.

DELIRIUM SENILE, see Dementia—*d. Tremifaciens*, *D. tremens*.

DELIRUS, Delirious.

DÉLITESCENCE, (*delitescere*, 'to abscond, to lie hid,' from *de*, and *lateo*.) See Repercussion.

DELITESCENTIA, see Repercussion.

DELIV'ER, (Prov.) *Lay*, (*de*, and *liberare*, (F.) *livrer*, 'to set free.') 'To put to bed.' To disburden of a child.

DELIVERY, Parturition—*d. False*, *False water*.

DÉLIVRANCE (F.). An expression, which, in common language, signifies the action of delivering, *libera'tio*, but in the practice of obstetrics means the extrusion of the secundines, either spontaneously or by the efforts of art. This completion of delivery—*par'tus secundarius*, *secundinarum expul'sio* seu *extra'ctio*—is produced by the same laws as the expulsion of the fœtus. Sometimes, the after-birth follows the child immediately; at others, it is retained; and requires manual interference to remove it. The following are the chief cases in which it has been practised. 1. Intimate adhesion between the placenta and paries of the uterus. 2. Spasmodic contraction of the orifice of the uterus. 3. Hour-glass contraction. 4. Torpor or hemorrhage after the expulsion of the child, and,—5. Insertion of the placenta at the orifice of the uterus.

DÉLIVRE, Secundines.

DELOCATIO, (*de*, and *locus*, 'a place.') *Luxatio*.

DELPHINIA, see Delphinium staphisagria.

DELPHININE, see Delphinium staphisagria.

DELPHINIUM, (from *δέλφις* or *δέλφιν*, 'a dolphin,' which the flowers resemble.) *D. consolida* seu *Seg'letum* seu *Versicolor*, *Calcitra'pa*, *Consolida Rega'lis*, *Branch'ing Lark'spur*, *Stag'gerweed*. *Ord. Ranunculaceæ*. *Sex. Syst.* Polandria Trigynia. (F.) *Pied d'alouette des champs*. It has been employed as a vermifuge. The flowers have been used in ophthalmia, and the seed *Delphinium*, (Ph. U. S.) have the same property as those of *Stavesacre*.

DELPHINIUM CONSOLIDA, Delphinium—*d. Segetum*, Delphinium.

DELPHINIUM STAPHISA'GRIA. The systematic name of the *Stavesacre*, *Staphisagria*, *Phtheirot-tonum*, *Phthe'rium*, *Sta'phis*, *Pedicularia*, (F.) *Staphisaigre*, *Herbe aux Poux*, *Danphinelle*. The seeds, *Staphisa'gria*, (Ph. L. and E.) have a disagreeable smell; a nauseous, bitterish, hot taste; and are cathartic, emetic, and vermifuge: owing, however, to the violence of their operation, they are seldom given internally, and are chiefly used in powder, mixed with hair powder, to destroy lice. An alkaloid of this plant has been separated, and received the name of *Delphin'in*, *Delphia*, (F.) *Delphine*, *Delphinine*. It is extremely acrid, and has been recently used, like verataria, in tic douloureux, paralysis and rheumatism. It is used in the form of ointment, or in solution in alcohol, (gr. x to xxx, or more, to ʒi.) applied externally.

DELPHINIUM VERSICOLOR, Delphinium.

DELPHYS, (*δέλφης*.) Uterus. Vulva.

DELTA, (like the Greek letter Δ.) Vulva.

DELTIIFORMIS, (*delta*, Δ, and *forma*, 'shape,') Deltoid.

DELTOID, *Deltoides*, *Deltoides*, *Deltoidens*, *Deltiformis*, (F.) *Deltóide*, (*clavi*, Δ, and *oides*, 'resemblance.') *Sous-acromio-clavi-huméral* of Du-mas; *Sous-acromio-huméral* of Chaussier. A triangular muscle forming the fleshy part of the

shoulder, and covering the shoulder-joint. It extends from the outer third of the clavicle, from the acromion and spine of the scapula to near the middle and outer part of the os humeri, where it is inserted by means of a strong tendon. This muscle raises the arm directly upwards, when the shoulder is fixed, and carries it anteriorly or posteriorly, according to the direction of the fibres, which are thrown into action. If the arm be rendered immovable, the deltoid acts inversely and depresses the shoulder.

DELTOIDE, Deltoid.

DELUSION, (*de*, and *ludere*, *lusum*, 'to play,') Hallucination.

DEM EL MUCA. A name given by Prosper Alpinus to a disease, which, he says, is proper to Egypt. He considers it to be inflammation of the brain; but others describe it as a pernicious intermittent.

DEMAGNETIZATION, *Demagnetisatio*. The act of removing the condition of magnetization.

DÉMANGEAISON, Itching.

DEMANUS, (*de*, and *manus*, 'a hand,') Acheir.

DÉMENCE, Dementia—*d. Imée*, Idiotism.

DEMENTED, Insane, see Dementia.

DEMEN'TIA, *Amen'tia*, *Fatu'itas*, *Anœ'a*, *Para-nœ'a*, *Mo'ria de'mens*, *Noïsthen'i'a*, *Incohe'rency*, *Imbecil'ity*, *Incohe'rent Insan'ity*; (*de*, and *mens*, gen. *mentis*, 'mind,' or 'reason,') (F.) *Démence*, *Bêtise*. In common parlance, and even in legal language, this word is synonymous with insanity. Physicians, however, have applied it to those cases of unsound mind which are characterized by a total loss of the faculty of thought, or by such an imbecility of intellect that the ideas are extremely incoherent, there being at the same time a total loss of the power of reasoning. Mania and melancholy are apt to end in this, if possible, more deplorable state. *Do'tage* is the Dementia of the aged—*Mo'ria de'mens lere'ma*, *Lere'ma*, *Lere'sis*, *Le'rus*, *Delir'io Se'nium*, *Amen'tia Sen'il'is*, *Pueril'itas*, *Delir'ium Sen'il'e*, *Anil'itas*, *Se'nile Insan'ity*, *Se'nile Dement'ia*, *Insan'ity of the a'ged*, (Sc.) *Doittrie*,—a form of insanity, in which there is a hopeless decadeney or loss of the mental faculties.

DEMENTIA ACCIDENTALIS, see Lunatic—*d. Adventitia*, see Lunatic—*d. Naturalis*, Idiotism—*d. Paralytica*, Paralysis of the Insane.

DEMI, ('half,' *hū*, and *Semi*)—*d. Bain*, Semicupium—*d. Circulaire*, Semi-circular—*d. Épineux*, Semi-spinalis colli—*d. Épineux du dos*, Semi-spinalis dorsi—*d. Gantelet*, *Gantelet*—*d. Interrosseux du ponce*, *Flexor brevis pollicis manus*—*d. Lunaire*, Semi-lunar—*d. Membraneux*, Semi-membranosus—*d. Nerveux*, Semitendinosus—*d. Orbicularis*, Orbicularis oris—*d. Tendineux*, Semitendinosus—*d. Tierce*, Hemitritæa (febris).

DEMISSIO ANIMI, (*demittere*, *demissum*, 'to lower,') Depression.

DEMISSOR, Catheter.

DEMODEX FOLLICULORUM, (*ἐμπος*, 'lard,' and *ὄψ*, 'a boring worm,') *Acarus folliculorum*.

DEMONOMANIA, *Demonomania*, *Demonopath'ia*, *Dæmonopath'ia*, *Dæmo'nia*, *Theoma'nia*, *En-theoma'nia*, (*δαμων*, 'demon,' a spirit of good or evil, and *mania*, 'madness,') A variety of madness, in which the person conceives himself possessed of devils, and is in continual dread of malignant spirits, the pains of hell, &c. Also, religious insanity, *Sebastoma'nia*.

DEMONOPATHY, (*δαμων*, 'demon,' and *πάθος*, 'affection,') Demonomania.

DEMONSTRATOR, (*demonstrare*, *demonstratum*, (*de*, and *monstrare*,) 'to show,' 'exhibit,') One who exhibits. The index finger. See Digitus.

DEMONSTRATOR OF ANAT'OMY. One who ex-

hibits the parts of the human body. A teacher of practical anatomy.

DEMOTIVUS LAP'SUS, *Repenti'na mors*, (*de-movere*, (*de*, and *movere*,) 'to move off,' 'remove,' and *lapsus*, 'a fall,') Sudden death.

DEMUL'CENTS, *Demulcent'ia*, *Involvent'ia*, *Obvolvent'ia*, *Lubrican'tia*, (*demulcere*, (*de*, and *mulcere*,) 'to soothe,' 'to assuage,') (F.) *Adoucissants*. Medicines supposed to be capable of correcting certain acrid conditions imagined to exist in the humours. Substances of a mucilaginous or saccharine nature belong to this class. Demulcents may act directly on the parts with which they come in contact; but in other cases, as in catarrh, their effect is produced by contiguous sympathy; the top of the larynx being soothed by them first, and indirectly the inflamed portion of the air-passages. In diseases of the urinary organs, they have no advantage over simple diluents.—See Diluents and Emollients. The following are demulcents;—Acacia Gummi; Althæa Folia et Radix; Amygdalæ; Amylum; Avena Farina; Cera; Cetaceum; Cydonia Semina; Fucus Crispus; Glycyrrhiza; Hordeum; Lichen; Linum; Olivæ Oleum; Sassafras Medulla; Sesamum; Sevum; Tragacantha, and Ulmus.

DEMUSCULAT'US, (*de*, and *musculus*, 'a muscle,') (F.) *Décharné*. Emaciated, devoid of flesh.

DENARIUS, (*deni*, 'ten,') a Roman coin, equal in value to about 10 cents, or 8 pence English. It was marked with the letter X, to signify 10 asses.

Also, the 7th part of the Roman ounce.

DENDROLIBANUS, (*δενδρόν*, 'a tree,' and *λίβανος*, 'frankincense,') Rosmarinus.

DENERVATIO, (*de*, and *nervus*, 'a tendon,') Aponeurosis.

DENGUE, ([S.] 'an affected person,') (*Di'ngée*, *Dun'ga*, *Dan'dy*, *Bou'quet*, *Buck'et Fe'ver*, *Eruptive articu'lar fe'ver*, *E. rheumatic fe'ver*, *Break'bone*, *Neural'gic*, *Solar* or *Sun Fe'ver*, *Rheumatismus febr'ilis*, *Scarlati'na rheumatica*, *Exanth'es'is arthro'sia*, *Plantari'ria*, *Fe'bris exanthemat'ica articu'laris*, (F.) *Giraffe*. A disease, which first appeared in the years 1827 and 1828, in the West Indies, and in the southern states of North America. It was extremely violent in its symptoms, but not often fatal. It usually commenced with great languor, chilliness, and pain in the tendons about the smaller joints. To these symptoms succeeded burning heat and redness of the skin, pains in the muscles of the limbs or in the forehead, with vomiting or nausea. The fever continued for one, two or three days, and usually terminated by copious perspiration. In different places, it put on different appearances; but seems in all to have been a singular variety of rheumatic fever. The usual antiphlogistic treatment was adopted, and successfully.

DEN'IACH, MIN'ERAL WATERS OF. Deniach is a village in Swabia, at the entrance of the Black Forest. The waters contain carbonic acid, carbonate of iron, carbonate of lime, sulphate of magnesia, and carbonate of soda.

DENIGRATIO, (*denigrare*, (*de*, and *niger*, 'black,') 'to become black,') The act of becoming black, as in cases of sphacelus, sugillation, &c.

DENS, gen. *Dentia*, Tooth, see Axis—*d. Epistrophei*, see Odontoid—*d. Leonis*, Leontodon taraxacum.

DENS PRO'LIFER. A term used by Bartholin for a supernumerary tooth, which appears to grow upon a primitive or parent tooth.

DENT, Tooth—*d. de Lion*, Leontodon taraxacum—*d. de Sagasse*, see Dentition.

DENTACIOGUM, (*dens*, and *aye*, 'I expel,') Dentagra.

DEN'TA PRA, *Den'ticeps*, *Densæ'paga*, *Denti-*

du'cum, *Dentago'gum*, *Dentalis for'fec*, *Fu'fec*, *dental'ria*, *Odonthar'paga*, *Odont'agra*, *Odontago'gon*, (*dens*, and *appa*, 'a seizure.') (F.) *Davies*.
A tooth-forceps.

DENTAIKE, Dental.

DEN'TAL, *Dental'lis*, *Dental'rius*, *Denticula'tus*, (F.) *Dentaire*. That which concerns the teeth. See *Dentiformis*.

DENTAL ARCH'ES, (F.) *Arcades dentaires*, are the arches formed by the range of alveoli in each jaw.

DENTAL ARTERIES are those arteries which nourish the teeth. They proceed from several sources. The teeth of the upper jaw, e. g., receive their arteries from the *infraorbital* and *superior alveolar*,—themselves branches of the *internal maxillary*. The teeth of the lower jaw receive their branches from the *inferior dental* or *inferior maxillary*, which is given off by the *internal maxillary*, and runs through the dental canal, issuing at the mental foramen, after having given numerous ramifications to the teeth and jaw.

DENTAL CANALS', (F.) *Conduits dentaires*, *Maxillary canals*. The bony canals, through which the vessels and nerves pass to the interior of the teeth.

DENTAL CAR'TILAGE, *Cartila'go dental'is*. The cartilaginous elevation, divided by slight fissures, on the biting margins of the gums in infants, prior to dentition. It is a substitute for the teeth.

DENTAL CAV'ITY, (F.) *Cavité dentaire*. A cavity in the interior of the teeth in which is situate the dental pulp. See *Tooth*.

DENTAL EXCAVATOR. An instrument for the removal of the decayed part of a tooth, preparatory to the operation of filling.

DENTAL FOL'LICLE, *Follic'ulus seu Cap'sula dent'is*, *Odonthel'ca*, (F.) *Follicule dentaire*, *Odonthel'que*. A membranous follicle, formed of a double lamina, in which the teeth are contained before they issue from the alveoli, and which, consequently, aids in the formation of the alveolodental periosteum, and of the membrane that envelops the pulp of the teeth. The dental follicles are lodged in the substance of the jaws.

DENTAL GANGRENE, *Car'ies Dent'ium*, *Odontal'gia cario'sa*, *Odontonecro'sis*, *Odontophacell'is*, *Odontophacell'ismus*, *Necro'sis Dent'ium*, (F.) *Carie des Dents*. Gangrene or caries of the teeth. See *Caries*.

DENTAL NERVES, (F.) *Nerfs dentaires*. Nerves which pass to the teeth. Those of the upper incisors and canine are furnished by the *infra-orbital* nerve, a branch of the superior maxillary, and those of the molars by the trunk of the same nerve. The teeth of the lower jaw receive the nerves from the inferior maxillary, which, as well as the superior maxillary, arises from the 5th pair.

DENTAL PULP, *Pul'pa seu Nu'cleus seu Substan'tia pulpo'sa dent'is*, *Ger'men dental'e*, (F.) *Pulpe dentaire*. A pulsatious substance, of a reddish-gray colour, very soft and sensible, which fills the cavity of the teeth. It is well supplied with capillary vessels.

DENTAL SURGEON, Dentist.

DENTAL VEINS have a similar distribution with the arteries.

DENTALIS, *Odontoid*—d. *Forfex*, *Dentagra*.

DENTALIUM. A genus of shells in the shape of a tooth. They formerly entered into several pharmaceutical preparations, but were useless ingredients; consisting—like shells in general—of carbonate of lime and gelatin.

DENTARIA, *Anthemis Pyrethrum*, *Plumbago Europæa*.

DENTARIUS, Dentist.

DENTARPAGA, (*dens*, and *appa'ze*, 'I force away.') *Dentagra*.

DENTATA VERTEBRA, Axis.

DENTATE, *Dentat'el*, *Denta'tus*, *Ser'rat'ed*, *Ser'rat'us*, (F.) *Dentel'è*. Cut in the shape of teeth.

DENTATURE, Digitation.

DENTATUS, Dentate.

DENTELAIRE, *Plumbago Europæa*.

DENTELARIA, *Plumbago Europæa*.

DENTEL'è, Dentate—d. *Antérieur petit*, *Pectoralis minor*—d. *Grand*, *Serratus magnus*—d. *Postérieur et inférieur*, *petit*, *Serratus posticus inferior*—d. *Supérieur*, *petit*, *Serratus posticus superior*.

DENTELURE, see *Dentate*, Digitation.

DENTES, (pl. of *dens*,) Teeth, see *Tooth*—d. *Angulares*, *Canine teeth*—d. *Bicuspidati*, *Molar teeth*—d. *Buccales*, *Molar teeth*—d. *Canini*, *Canine teeth*—d. *Clavales*, *Molar teeth*—d. *Columnellares*, *Canine teeth*—d. *Ctenes*, *Incisive teeth*—d. *Cuspidati*, *Canine teeth*—d. *Dichasteres*, *Incisive teeth*—d. *Gelasini*, *Incisive teeth*—d. *Gomphii*, *Molar teeth*—d. *Incisores*, *Incisive teeth*—d. *Lactei*, see *Dentition*—d. *Laniarii*, *Canine teeth*—d. *Maxillares*, *Molar teeth*—d. *Molares*, *Molar teeth*—d. *Mordentes*, *Canine teeth*—d. *Oculares*, *Canine teeth*—d. *Opsigoni*, see *Dentition*—d. *Primores*, *Incisive teeth*—d. *Rasorii*, *Incisive teeth*—d. *Sapientiae*, see *Dentition*—d. *Serotini*, see *Dentition*—d. *Sophonisti*, see *Dentition*—d. *Sophroretici*, see *Dentition*—d. *Sophonisteres*, see *Dentition*—d. *Tomici*, *Incisive teeth*.

DENTICEPS, (*dens* and *capi'o*, 'I seize.') *Dentagra*.

DENTICOLA HOMINIS, (*dens*, and *colare*, 'to inhabit,') see *Odontolithus*.

DENTICULATUM, (*Ligamentum*,) *Ligamentum dentat'um*, *Membrana dentat'a*, (F.) *Ligament dentel'è* or *denticul'è*. A slender cord, situate between the anterior and posterior fasciculi of the spinal nerves, and between the tunica arachnoidea and pia mater. It is attached to the dura mater, where that membrane issues from the cranium, and accompanies the spinal marrow to its inferior extremity. It sends off, from its outer edge, about twenty slender processes, in the form of *denticuli*, each of which passes outwards and connects itself with the dura mater in the intervals between the anterior and posterior roots of the nerves.

DENTICULATUS, Dental.

DENTICULUS, (dim. of *dens*, 'a tooth,') *Crena*.

DENTIDUCUM, (*dens*, and *ducere*, *ductum*, 'to draw,') *Dentagra*.

DENTIER, (F.) (from *dens*,) *Denture*. A set or row of teeth, mounted on metal or ivory, to be adjusted to the alveolar margin. *Dentiers* are simple or double. To the latter, that is, to the full set, the name *Râtelier* is given.

DENTIFORM, *Dentiform'is*, (*dens*, and *forma*, 'form,') Resembling a tooth in shape.

DENTIFORMIS, *Odontoid*.

DENTIFRICE, *Dentifric'ium*, *Remed'ium dentifric'ium*, *Odontotrin'mus*, *Odontomes'gma*, *Tooth powder*, *Tooth paste*, (*dens*, and *fricare*, 'to rub,') A name given to different powders and pastes proper for cleansing the enamel of the teeth, and removing the tartar which covers them. Powdered bark and charcoal, united to any acidulous salt—as cream of tartar—form one of the most common *dentifrices*.

Electuaire, or *Opiate Dentifrice*, of the Parisian Codex, consists of coral, ʒiv, cuttlefish bones and cinnamon, each ʒij, cochineal, ʒss, honey, ʒx, alum, gr. iv. or v.

Poudre Dentifrice of the Codex is composed of *Bole Armeniac*, red coral [prepared,] and cuttlefish bones, each ʒvj, dragon's blood, ʒij, cochineal, ʒj, cream of tartar, ʒix, cinnamon, ʒij, cloves, gr. xij, well mixed. The one used by the author, is composed of charcoal and orris root, each one part; prepared chalk, two parts.

Charcoal, finely powdered and mixed with chalk, forms as good a dentifrice as any.

DENTIG'EROUS, *Dentiger*, (*dens*, and *gero*, 'I carry.') Tooth carrying, as *dentigerous cyst*: one containing teeth.

DENTILIOUN, *Leontodon taraxacum*.

DENT'INAL, *Dentinalis*, (from *dentine*.) Relating to the dentine of the teeth.

DENTINE, see *Tooth*—*d*. Secondary, see *Tooth*.

DENTISCAL'PIUM, (*dens*, and *scalpere*, 'to scrape.') *Odontoglyphum*. This word has been applied to the instrument used for scaling the teeth; to the *toothpick*, or *toothrake*, (F.) *Cure-dent*; and to the *gum-lancet*,—the *déchaussoir* of the French.

DENT'IST, *Dentista*, *Odontia'ter*, *Odontia'trus*, *Dentarius*, *Dental Surgeon*, *Surgeon dentist*, (F.) *Dentiste*. One who devotes himself to the study of the diseases of the teeth, and their treatment.

DENT'ISTRY, *Odontotech'ny*, *Odontia'tria*, *Odontotherapi'a*, *Dental Surgery*. The art of the dentist.

DENTITIO, *Dentition*—*d*. *Difficilis*, *Dysodontiasis*.

DENTIT'ION, *Teeth'ing*, *Dentit'io*, *Dentit'is*, *Odontophy'ia*, *Odont'asis*, *Odonto'sis*, (*dentire*, *dentium*, 'to breed teeth.') The exit of the teeth from the alveoli and gums; or rather the phenomena which characterize the different periods of their existence. The germs of the first teeth, *dentes lac'tei* or *milk teeth*, (F.) *dents de lait*, are visible in the fœtus about the end of the second month; and they begin to be ossified from the end of the third to that of the sixth month. At birth, the corona of the incisors is formed, but that of the canine is not completed; and the tubercles of the molares are not yet all united. Gradually, the fang becomes developed; and at about six or eight months begins, what is commonly called, the *first dentit'ion*, *Odont'ia dentitio'nis lactant'ium*. The two middle incisors of the lower jaw commonly appear first; and, some time afterwards, those of the upper jaw; afterwards, the two lateral incisors of the lower jaw; and then those of the upper, followed by the four anterior molares; the *canine*, or *eye-teeth*, at first, those of the lower, and, afterwards, those of the upper jaw, next appear; and, subsequently and successively, the first 4 molares—2 above and 2 below, 1 on each side. The whole number of the *primary*, *temporary*, *deciduous*, *shedding* or *milk-teeth*, (*dentes tempora'rii*.) (F.) *Dents de lait*, is now 20.

The eruption of the milk teeth takes place, approximately, in the following order:

Central incisors.....	6th to 8th month.
Lateral incisors.....	7th to 10th month.
First molar.....	12th to 14th month.
Canines.....	15th to 20th month.
Second molar.....	20th to 30th month.

The *second dentit'ion* or *shed'ding of the teeth*, *odont'ia dentitio'nis puer'ilis*, *Dedentit'ion*, begins about the age of 6 or 7. The germs or membranous follicles of these second teeth—to the number of 32—as well as the rudiments of the teeth themselves, are visible, even in the fœtus, with the exception of those of the small molares, which do not appear till after birth. They are contained in alveoli of the same shape as those of the first dentition. Their ossification commences at from 3 to 6 months after birth, in the incisors and first molares; at eight or nine months, in the canine; about three years, in the molares; 3½ in the second great molares, and about 10 years in the last. As the alveolus of a new tooth becomes gradually augmented, the septum between it and that of the corresponding milk tooth is absorbed, and disap-

pears. The root of the milk tooth is likewise absorbed; its corona becomes loose and falls out, and all the first teeth are gradually replaced by the *permanent teeth*, *Dent'es serot'ini*. This second dentition becomes necessary in consequence of the increased size of the jaws. The new teeth have neither the same direction nor the same shape as the old; and they are more numerous, amounting until the age of 25, (sooner or later,) to 28. About this period, a small molaris appears at the extremity of each jaw, which is called *wis'dom tooth*, *wit tooth*, *Dens sapien'tiæ* seu *serot'ianus* seu *sophroret'icus* seu *sophronis'tus* seu *sophronis'ter* seu *opsi'gonus*, *Ops'iodous*, (F.) *Arrière dent*, *Dent de sagesse*, making the whole number of permanent teeth 32.

The eruption of the permanent teeth is remarkable for its general regularity; so that it constitutes an important means for ascertaining the age of the individual during the early period of life.

First molars.....	7th year.
Central incisors.....	8th year.
Lateral incisors.....	9th year.
First bicuspid.....	10th year.
Second bicuspid.....	11th year.
Canines.....	12th year.
Second molars.....	13th year.

The teeth of the lower jaw generally precede by a few weeks those of the upper.

During the period of dentition, that is, of the first dentition, the infant is especially liable to disease;—the irritation, produced by the pressure of the tooth on the superincumbent gum, sometimes occasioning pyrexia, convulsions, diarrhœa, &c., which symptoms are often strikingly relieved by a free division of the distended gum. This disordered condition is called *Teeth'ing*, *Odont'ia dentitio'nis*, *Odont'iasis*, *Odontal'gia*, *dentitio'nis*, *Odax'ismus*.

DENTITIS, (*dens*, and *itis*.) *Dentition*.

DENTIUM CORTEX, ('bark of the teeth,') *Enamel of the teeth*—*d*. *Dolor*, *Odontalgia*—*d*. *Nitor*, *Enamel of the teeth*—*d*. *Scalptura*, (*Lancing the gums*), see *Gum lancet*—*d*. *Vacillantia*, *Odontoseisis*.

DENTO, (from *dens*.) One whose teeth are prominent.

DENTOIDEUS, (*dens*, and *eidos*, 'resemblance,') *Odontoid*.

DENTS (pl. of *dent*) *BICUSPIDÉES*, *Bicuspid teeth*—*d*. *Col des*, *Collum dentium*—*d*. *Conoïdes*, *Canine teeth*—*d*. *de Lait*, see *Dentition*—*d*. *Laniaries*, *Canine teeth*, *Laniarii Dentes*—*d*. *Mâchelières*, *Molar teeth*—*d*. *Molaires*, *Molar teeth*—*d*. *Multicuspidées*, *Molar teeth*, great—*d*. *Éillières*, *Canine teeth*.

DENTURE, *Dentier*.

DENUDA'TION, *Denuda'tio*, *Gymno'sis*, (*denudare* (*de*, and *nudare*, *nudatum*) 'to lay bare,') *Condition of a part*, deprived of its natural coverings, whether by wounds, gangrene, or abscess. It is particularly applied to the bones, when deprived of their periosteum, and to the teeth when they lose their enamel or dental substance, or when the gums recede from them, and their sockets are destroyed.

DEOB'STRUENT, *Deob'struens*, (*de*, and *obstruere* (*ob*, and *struere*) 'to obstruct,') *Dephract'icum*, *Deoppi'lans*, *Deoppi'lavum*, *Ecphract'ic*, (F.) *Désobstruant*, *Désobstructif*, *Désopilatif*, *Désopilant*. A medicine given with the view of removing any obstruction. The word corresponds to *aperient*, in its general, not in its particular sense. It is now almost abandoned, and, when used, conveys by no means definite ideas.

DEODORANT, (*de*, and *ovor*, 'smell,') *Antibromic*.

DEODORIZER, *Antibromic*.

DEONTOL'OGY, *Deontology* 'ia, (α δοντα, 'what are due,' and λογος, 'a description.') A word introduced by Bentham to signify morals, or the science of duties.

MEDICAL DEONTOL'OGY, *Deontology* 'ia med'ica, *Medical eth'ics*, *Medical etiquette*, (F.) *Déontologie médicale*. The duties and rights of medical practitioners.

DEOPILANS, (de, and *oppilo*, 'I stop up.')

DEOPILATIVUM, Deobstruent.

DEPASCENS, (de, and *pascere*, 'to feed,') Phagedenic.

DEPAUPERATUS, (de, and *pauperare*, *pauperatum*, 'to make poor,') Impoverished.

DEPERDITIO, (de, and *perdere*, *perditum*, 'to lose,') Abortion.

DEPÉRISSEMENT, (de, and *perire*, 'to perish,') Emaciation.

DEPHRACTICUM, (de, and *φρασσω*, 'to obstruct,') Deobstruent.

DEPILATIF, Depilatory.

DEPILATIO, (de, and *pilus*, 'hair,') Alopecia, Depilation—d. Capitis, Calvities.

DEPILATION, *Depilatio*, *Epilatio*, *Dropacismus*, *Made'isis*, *Mad'isis*, *Psilo'isis*. Loss of hair, either spontaneously or by art.

DEPILATORY or DEPILATORY, *Depilato'rium*, *Psilo'thron*, *Ectillot'jeus*, *Epilato'rium*, (F.) *Dépilatoire*, *Dépilatif*, *Epilatoire*. Any thing which causes the loss of the hair. Depilatories are usually caustic applications, in which quicklime or some other alkaline substance, sulphuret of iron, &c., enter.

DEPILATORY, COL'LEY'S, seems to consist of quicklime and a portion of sulphuret of potassa.

Delerois's depilatory, and *Plencé's depilatory*, have a similar composition. They all, probably, contain arsenic, in the form of orpiment. The following formula has been given for that of Colley: *quicklime*, 30 parts; *soap lees*, 125 parts; *sulphur*, 4 parts; *nitre*, 4 parts; *orpiment*, 12 parts. Another depilatory by F. Boudet is formed of *sulphuret of sodium*, 3 parts; *quicklime*, in powder, 10; *starch*, 10; *mix*. Rub a little of this powder with water; apply it to the part, and remove the hair in a minute or two with a wooden knife. Dorvault recommends the *Depilatory of Martins*, which is a sulphuretted sulphuret of calcium; made by passing *sulphohydric acid* into *quicklime* to saturation. A little of this is applied to the part, and in eight or ten minutes it becomes solid. It must then be washed off with cold or hot water, when the skin will be found denuded. Another, given by Redwood, is a strong solution of the *sulphuret of barium*, made into a paste with *starch powder*. It is used like the one advised by Boudet. Those that do not contain arsenic are, of course, safer.

A *pitch plaster*, *Pitchcap*, *Dro'pax*, is sometimes used as a *depilatory*. It of course pulls the hair out by the roots. Depilation by it has been termed *Dropacismus*, *Pisso'isis*, *Pitto'isis*, *Picacismus*, and *Picatio*.

DE'PILIS. Devoid of hair. Hairless.

DEPLACEMENT DE LA CATARACTE, ('displacement of the cataract,') see Cataract.

DEPLÉTIF, Depletory.

DEPLETION, *Deple'tio*, (*depleo*, *depletum*, 'to unload,') The act of emptying or unloading,—for example, the vessels, by blood-letting and the different evacuations. Also, inordinate evacuation.

DEPLETORY, *Deple'ting*, (F.) *Dépletif*. Having relation to depletion—as 'a *depletory* or *depleting agent*.'

DEPLUMATIO, *Pti'los*, *Ptilo'isis*, (*deplumis* (de, and *pluma*) 'without feathers.')

A disease of the eyelids, in which they are swollen, and the eyelashes fall out. See Madarosis.

DEPOS'IT, *Depos'itum*, (*depono* (de, and *pono*, *positum*, 'to lay or put') 'to lay or put down.')

(F.) *Dépôt*. Any thing laid or thrown down. In physiology and pathology, a structureless substance, separated from the blood or other fluid, as the typhous, tuberculous, purulent, melanlic, diphtheritic, and urinary *deposits*.

DEPOSIT, Feeculence.

DEPOSITIVE, *Depositi'vus*. An epithet used by Mr. Erasmus Wilson to express that condition of the membrane in which plastic lymph is exuded into the tissue of the derma, so as to give rise to the production of small hard elevations of the skin, or pimples. Under "*depositive inflammation of the derma*," he comprises strophulus, lichen, and prurigo.

DEPOSITUM, Deposit.

DÉPÔT, Abscess, Deposit, Sediment—d. *Lait-euse sur la Cuissée*, Phlegmatia dolens—d. *de l'Urine*, Sediment of the Urine.

DEPRAVATION, *Deprava'tio*, (de, and *pravus*, 'bad,') Perversion, corruption—as depravation of the taste, &c.

DEPREHENSIO, (*deprehendere*, *deprehensum*, 'to seize upon,') Diagnosis, Epilepsy.

DEPRESSANTS, Sedatives.

DEPRESSION, *Depres'sio*, *Impres'sio*, (*deprimere*, *depressum* (de, and *premere*, 'to depress.')

In *Anatomy*, it means an excavation, hollow, or fossa. In *Surgery*, it is applied to a fracture of the cranium, in which the portions of fractured bone are forced inwards; (F.) *Subgrondation*; called, also, *Es'phlasia*, *En'thlasis*, *Cotapi'ceis*, *Camp'sis* *Depres'sio*, *Thla'sis* *Depres'sio*.

Depression, (F.) *Abaissement*, means Couching.—See Cataract.

Depression also means *dejection*, or *deject'edness*, *Ademon'ia*, *Ademo'syné*, *Demis'sio* *an'mi*.

DEPRESSION, NERVOUS, Shock, nervous—I. Vital, Shock, nervous.

DÉPRESOIRE, Meningophylax.

DEPRESSOR, (F.) *Abais'seur*. Several muscles have been so termed, because they depress the parts on which they act.

DEPRESSOR ALÆ NASI, D. *la'bii superio'ris alæque nasi* seu *la'bii superio'ris pro'prius*, *Inci'sivus me'dius*, *Myrtifor'mis*, *Constricto'res ala'rum nasi* ac *depresso'res la'bii superio'ris*, *Maxillo-alveoli-nasal*—part of the *labialis* (Ch.), (F.) *Abais'seur de l'aile du nez*. It arises from the superior maxillary bone immediately above the junction of the gums with the two incisor and canine teeth; and passes upwards to be inserted into the upper lip and root of the ala nasi, which it pulls downwards.

DEPRESSOR ANGULI O'RIS, *Triangula'ris*, *Depressor labio'rum communis*, D. *labio'rum*, (F.) *Sousmaxillo-labial*, *Maxillo-labial* (Ch.), *Abais'seur de l'angle des lèvres ou Muscle triangulaire*. A muscle, situate at the lower part of the face. Its form is triangular. It arises from the outer oblique line on the lower jawbone, and terminates in a point at the commissure of the lips, which it pulls downwards.

DEPRESSOR LABII INFERIORIS, *Quadratu's*, Q. *Men'ti*, *Depressor la'bii inferio'ris pro'prius*—(F.) *Mentonniër-labial*, *Mento-labial* (Ch.), *Corré du Menton*, *Houppé du Menton*, *Abais'seur de la lèvre inférieure*. A small, thin, and quadrilateral muscle, which arises from the external oblique line of the lower jaw, and ascends to the lower lip, where it becomes confounded with the orbicularis oris. It pulls the lip downwards and outwards.

DEPRESSOR LABII SUPERIORIS ALÆQUE NASI, D. *Alæ nasi*—d. *Labii superioris proprius*, D. *Alæ nasi*—d. *Labiorum communis*, D. *Anguli oris*—d. *Oculi*, *Rectus inferior oculi*.

DEPRESSOR PALPEBRÆ INFERIORIS. A fleshy

bundle, which forms part of the palpebralis muscle. Heister describes it separately, but it is not admitted now.

DEPRESSORIUM, Meningophylax.

DEPRIMENS AURICULÆ. (*deprimere*, 'to depress,') *Retrahens auris*—d. *Maxillæ biverter*, *Digasticus*—d. *Oculi*, *Rectus inferior oculi*.

DEPRIMENTIA. Sedatives.

DEPUCELLEMENT (F.) (*de*, and *pucelle*, 'a maid,') Defloration.

DEPUCELATION, Defloration.

DEPUDICATION, (*depudicare*, (*dē*, and *pudicus*, 'chaste,') 'to dishonour,') Defloration.

DEPURANTS, *Depurantia*, *Depuratives*, (*depurare*, *depuratum*, (*dē*, and *purus*,) 'to purify,') (F.) *Depurans*, *Depuratifs*. Medicines were formerly so called, which were supposed to possess the power of removing, from the mass of blood or humours, those principles which disturbed their purity; and of directing them towards some one of the natural emunctories. The juices of what are called antiscorbutic herbs, sulphur, and many other medicines, were ranked under this class.

DÉPURATIFS, *Depurantia*.

DEPURATION, *Depuratio*, *Catharsismus*, *Mundatio*. In Pathology, depuration has been used for the process by which nature purifies the animal economy, either by the agency of some eruptive disease, or some spontaneous evacuation, or by the assistance of medicine. See Clarification, and Depuratory.

DEPURATIVES, *Depurantia*.

DEPURATORY, *Depuratio*, (F.) *Depuratoire*. That which causes depuration, as by the urinary and cutaneous depurations. Applied, also, to diseases which have been considered capable of modifying the constitution advantageously, by acting on the composition of the fluids—such as certain eruptions, intermittents, &c. The word is, also, appropriated to medicines and diet, by which the same effect is sought to be induced.

DERADELPHUS, (F.) *Déradelphie*, (*dere*, and *ἀδελφος*, 'a brother,') A double monocephalous monster, whose two trunks are united above the umbilicus, and separated below.

DER'ADEN, gen. *Deraad'enis*, (*dere*, and *αδην*, 'a gland,') A gland in the neck.

DERADENITIS, (*dere*, *αδην*, 'a gland,') and *itis*,) Inflammation of the glands of the neck.

DERADENON'CUS, (*dere*, *αδην*, 'a gland,') and *ογκος*, 'a swelling,') Tumefaction of the glands of the neck.

DERANENCEPHALIA, *Thlipsencephalia*, (*dere*, and *anencephalia*, 'absence of brain,') A monstrosity in which only a small portion of the brain exists, resting on the cervical vertebrae—more properly *encephalia*. According to Is. G. St. Hilaire, a monstrosity characterized by the total absence of encephalon, and of the medulla spinalis into the cervical region.

DERANGED, (F.), (*de*, and *ranger*, 'to put in order,') Insane.

DERANGEMENT, Insanity.

DÉRATÉ (F.), *Lie'ne privatus*, (*de*, and *rate*, 'spleen,') Deprived of spleen; as by an operation.

DERBIA, Impetigo.

DERCH, (So.) (Sax. *dreoph*,) Nanus.

DERE, (*δερμ*,) Collum.

DERENCEPHALIA, *Deranencephalia*.

DERENCEPH'ALUS, (*dere*, and *κεφαλη*, 'head,') A monster whose brain is in the neck.

DERIS, (*δερμ*,) Cutis.

DERIVANS, (*derivare*, *derivatum*, (*dē*, and *rivus*, 'a river,') 'to turn water from its regular course,') Derivative.

DÉRIVATIF. Derivative.

DERIVATION, *Derivatio*, *Deflectio*, *Para-*

chetesis, *Antilep'sis*, *Antispas'sis*, *Revol'sion*, *Cu'ra derivat'iva seu revuls'o'ria*. When a 'centre of fluxion' is established in a part, for the purpose of abstracting the excited vital manifestations for some other, a derivation is operated. To such agency the term *médication substitutive* has been given.

The term DERIVATION has—like *Aspiration*—been applied to the suction power of the heart—a presumed agency in the circulation of the blood.

DERIVATIVE, *Revel'lent*, *Revol'sive*, *Deflectens*, *Derivans*, *Derivatio'rius*, *Antispas'ticus*, *Apospas'tic*, *Revuls'ivus*, *Revuls'o'rius*, (F.) *Dérivatif*, *Révulsif*. A remedy, which by producing a modified action in some organ or texture derives from the morbid condition of some other organ or texture. Revellents are amongst the most important remedies: they include, indeed, every physical and moral agent which is capable of modifying the function of innervation, and therefore almost every article of the materia medica.

DERIVATORIUS, Derivative.

DERMA, gen. *Der'matis*, (*δερμα*, gen. *δερματος*,) Cutis.

DERMAD, see Dermal Aspect.

DER'MAL, *Der'mic*, *Derma'lis*, *Der'micus*, (from *derma*,) Relating or belonging to the skin.

DERMAL ASPECT. An aspect towards the skin or external surface.—Barclay. *Dermad* is used adverbially by the same writer to signify 'towards the dermal aspect.'

DERMAL'GIA, properly *Dermatal'gia*, *Dermatodyn'ia*, (*derma*, and *αλγος*, 'pain,') Pain in the skin. Neuralgia of the skin. *Rheumatic Dermalgia* or *Rheumatism of the skin* is a form of neuralgia, which is referred, at times, to the nervous trunks, muscles, &c., but appears to be seated in the cutaneous nerves.

DERMATAGRA, (*derma*, and *αγρα*, 'a seizure,') Pellagra.

DERMATALGIA, (*derma*, and *αλγος*, 'pain,') Dermalgia.

DERMATANEUR'IA, *Adermoner'via*, (*derma*, *a*, priv., and *νευρον*, 'nerve,') Paralysis or loss of sensibility of the skin.

DERMATAUXE, (*derma*, and *αυξη*, 'increase,') Dermatophyma.

DERMATIATRIA, (*derma*, and *ιατρεία*, 'healing,') Healing of cutaneous diseases. The treatment of diseases of the skin—*Diadermiat'ria*, *Dermatocrati'va*.

DERMATITIS, (*derma*, and *itis*,) Cytitis, Erysipelatous inflammation.

DERMATOCHOLOSIS, (*derma*, and *χολη*, 'bile,') Icterus.

DERMATOCHYSIS, (*derma*, and *χυσις*, 'effusion,') Anasarca.

DERMATOCRATIA, (*derma*, and *κρατω*, 'to have power over,') see *Dermatiatria*.

DERMATODES, (*derma*, and *odes*,) Dermatoid.

DERMATODYNIA, (*derma*, and *οδυν*, 'pain,') Dermalgia.

DER'MATOID, *Dermatoï'des*, *Dermato'des*, *Dermoï'des*, *Dermo'des*, *Der'moid*, (*derma*, and *ειδος*, 'form,') That which is similar to the skin. This name is given to different tissues, which resemble the skin. The dura mater has been so called by some.

Mor'bi dermatoid'es, chronic cutaneous diseases.

DERMATOL'OGY, *Dermatolog'ia*, *Dermol'ogy*, (*derma*, and *λογος*, 'a discourse,') A discourse or treatise of the skin.

DERMATOLYSIS, (*derma*, and *λυσις*, 'loosing,') Chalastodermia.

DERMATOPATHIA, (*derma*, and *παθος*, 'affection,') Cutaneous disease.

DERMATOPERISCLERISMUS, (*derma*, *περι*, 'around,') and *σκληρος*, 'hard,') Induration of the cellular tissue.

DERMATOPERICLSEROSIS, Induration of the cellular tissue.

DERMATOPHILUS, (*derma*, and *φιλω*, 'I love,') *Chique*.

DERMATOPHY'MA, gen. *Dermatophy'matis*, *Dermatanz'ē*, (*derma*, and *φύμα*, 'tumour,') A tumefaction of the skin.

DERMATOPHYMATA VENEREA, Condylomata.

DERMATOPHY'TÆ, properly *Dermatoph'ytī* (*Mor'bi*), (*derma*, and *φυτον*, 'a vegetable,') An order of cutaneous diseases, including those that depend on, or are characterized by, the presence of parasitic plants. It is made to include *Porrigō* and *Sycosis*.

DERMATORRHAG'IA, (*derma*, and *ῥαγν*, 'rupture,') A discharge of blood from the skin.

DERMATORRHŒ'A, (*derma*, and *ῥεω*, 'I flow,') A morbidly increased secretion from the skin.

DERMATOSCLEROSIS, (*derma*, and *σκληρος*, 'hard,') Callosity.

DERMATOSĒS, (*derma*, and *osis*,) Cutaneous diseases.

DERMATOSIES VÉROLEUSES, *Syphilides*.

DERMATOSPASMUS, (*derma*, and *σπασμος*, 'spasm,') Horrida cutis.

DERMATOTYLOMA, gen. *Dermatotylo'matis*, (*derma*, *τύλος*, 'callosity and oma,') Callosity.

DERMATOTYLOSIS, Callosity.

DERMATOTYLUS, Callosity.

DERME, (*derma*,) Cutis.

DERMIC, *Der'micus*, *Dermat'icus*, *Derma'tinus*, *Derma'tis*, *Derma'tic*, *Der'mal*, (F.) *Der'mien*. Relating to the derma or skin.

DERMITIS, (*derma*, and *itis*,) Cytitis.

DERMODES, (*derma*, and *odes*,) Dermatoid.

DERMOG'RAPHY, *Dermograph'ia*, (*derma*, and *γραφω*, 'I describe,') An anatomical description of the skin.

DERMOHŒ'MIA, (*derma*, and *αἷμα*, 'blood,') Hyperæmia or congestion of the skin.

DERMOID, Dermatoid.

DERMOLOGY, Dermatology.

DERMOPHYTE, *Der'matophyte*, (*derma*, and *φυτον*, 'a vegetable,') A parasitic vegetable growth on the skin.

DERMO-SKELETON, see Skeleton.

DERMO-SYPHILIDES, *Syphilides*.

DERMOT'OMY, *Dermotom'ia*, (*derma*, and *τεμνειν*, 'to cut,') The part of anatomy which treats of the structure of the skin.

DERODYMUS, (*dere*, and *ἄδνμος*, 'a twin' or *δουιν*, 'to mingle,') see Diciphalus.

DERONCUS, (*dere*, and *ογκος*, 'a swelling,') Bronchocele.

DER'TRON, (*δερτρον*,) This word, which is used by Hippocrates, signified, according to some, the omentum or peritoneum, but according to others, the small intestine. See Epiploon.

DĒS (F.), as a prefix, *De*.

DĒSALTERANTS (F.), (*dés*, and *altération*, 'thirst,') Refrigerants.

DĒSARTICULATION (F.), (*dés*, and *articulus*, 'a joint,') *Disjoint'ing*. A word used to express the part of the operation, in amputation at an articulation, which consists in dividing the ligaments, and separating the articular surfaces. The word has, also, been used for that kind of anatomical preparation, the object of which is to separate the different bones of the skeleton, and especially those of the head.

DĒSASSIMILATION, see Assimilation.

LĒSCALORINĒSES, (*de*, and *calor*, 'heat,') A name given by Baumes to diseases which are characterized by diminished heat.

DESCEMET, MEMBRANE OF, see Aqueous Humour, and Cornea.

DESCEMETITIS, (*Descemet*, and *itis*,) A term improperly formed, and really signifying 'inflammation of Descemet,' the anatomist. Inflammation of the membrane of Descemet.

DESCENDENS NONI, see Hypoglossus.

DESCENSIO, (*decendere*, *descensum*, (from *de*, and *scandere*, 'to climb,') 'to descend,') Catapasis.

DESCENSUS, gen. *Descensus*, Catapasis — d. Testiculorum, Orchidocatabasis.

DESCENTE, Hernia — d. *de la Matrice*, Procidencia uteri.

DESECTUS, (*de*, and *secare*, *sectum*, 'to cut,') Castratus.

DĒSENFLURE, (*dés*, and *enfler*, (L.) *inflare*, 'to swell,') *Détumescence*.

DESHLE'R'S SALVE, (after the proposer,) *Ceratum resinæ compositum*.

DESSICANTIA, (*de*, and *siccare*, 'to dry,') Desiccativa.

DESSICATIO, Drying, Drying — d. *Herbarum*, Drying of plants.

DESSICATION, Drying.

DESSICAT'IVA, *Desican'tia*, *Siccan'tia*, *Exsiccati'va*, (F.) *Déssicatif*. Remedies, which, when applied externally, dry up the humours or moisture from a wound.

DĒSINFECTANT, Disinfectant.

DĒSINFECTION, Disinfection.

DESIPIENTIA, (*de*, and *sapio*, 'to be wise,') Delirium.

DESIRE, (*desiderium*, (F.) *désir*,) Libido — d. Venereal, Appetite, venereal, see Libido.

DES'MA, (*δεσμα*,) *Des'mē*, *Des'mus*. A bandage, a ligament.

DESMATUR'GIA, (*desma*, and *εργον*, 'work,') The doctrine of the application of bandages. Bandaging.

DESME, (*δεσμη*,) *Desma*.

DESMEDION, (dim. of *desme*,) Fasciculus.

DESMEUX, Ligamentous.

DESMIOGNA'THUS, (F.) *Desmiognathe*, from *δεσμιος*, 'bound,' and *γναθος*, 'jaw.' A double parasitic monster, *Order*, Polygnathiens, characterized by a supernumerary imperfect head, which is united, under the neck, by muscular and cutaneous, but no bony attachments, to the principal subject.

DESMITIS, *Desmophlogo'sis*, *Desmophlog'ia*, (from *desma*, and *itis*,) Inflammation of ligaments.

DESMOCHAUNO'SIS, (*desma*, and *χαυνωσις*, 'relaxation,') Relaxation of an articular ligament.

DESMODYN'IA, (*desma*, and *δυνν*, 'pain,') Pain in the ligaments.

DESMOG'RAPHY, *Desmograph'ia*, (*desma*, and *γραφη*, 'a description,') An anatomical description of the ligaments.

DESM'OID, *Desmo'idēs*, *Desmo'dēs*, (*desma*, and *ειδος*, 'resemblance,') Having the character of, or resembling, ligament.

DESMOID TIS'SUE, *Ligamen'tous Tissue*, *Tect'us desmo'ius*. This tissue is very generally diffused over the human body; has a very close connexion with the areolar tissue, and is continuous with it in divers places. It constitutes the ligaments, aponeuroses, &c.

DESMOL'OGY, *Desmolog'ia*, (*desma*, and *λογος*, 'a discourse, a treatise,') That part of anatomy which describes the ligaments. Also, a treatise on bandages.

DESMOP'ATHY, *Desmopath'ia*, (*desma*, and *παθος*, 'disease,') Disease of the ligaments.

DESMOPHLOGIA, (*desma*, and *φλογω*, 'I inflame,') Desmitis.

DESMORRHŒ'IS, (*desma*, and *ῥηξις*, 'rupture,') Rupture of an articular ligament.

DESMOSUS, (from *desmus*,) Ligamentous.

DESMOT'OMY, *Desmotom'ia*, (*desma*, and *τεμνειν*, 'to cut,')

nar, 'to cut.') The part of anatomy which teaches the mode of dissecting the ligaments.

DESMUS, (*desmus*, Ligament.

DÉSOSTRUCTANT, Deobstruent.

DÉSOSTRUCTIF, Deobstruent.

DÉSOPILATIF, (*dés*, and *opilo*, 'to stop up.') Deobstruent.

DÉSORGANISATION, Disorganization.

DÉSOXYGÉNÈSES, (*dés*, and *oxygène*,) M. Baumes includes under this title an order of diseases, which he considers dependent upon a diminution in the quantity of the oxygen necessary for the animal economy.

DESPOSTATS (F.), (*dés*, and *portare*, 'to carry away,') Infirm soldiers, formerly charged with the office of removing the wounded from the field of battle. The class of *Despotats* was introduced by Leo VI., at the commencement of the 9th century.

DESPUMATION, *Despumatio*, *Apaphris'mus*, *Epyphris'mus*, (*despumare*, *dés*, and *epumare*, *epumatum*, 'to skim, to remove the froth,') The separation of the froth and other impurities, which rise, by the action of the fire, to the surface of any fluid.

DESQUMATION, *Desquamation* *Cutis*, *Eclepsis*, *Amorpha*, *Apolepis*, *Apolepis'mus*, *Moulting*, (*desquamare*, *desquamatum*, (*dés*, and *equam*, 'a scale,') 'to scale off,') Exfoliation, or separation of the epidermis, in the form of scales, of a greater or less size, *Forfura'tio*, *Defurfura'tio*. This affection is a common consequence of exanthematous diseases.

DESQUMATORIUS, Exfoliative.

DÉSSECHEMENT, ([L.] *dés* and *seccare*, 'to dry,') Atrophy, Drying.

DESSERT, *Bella'ria*, *Trage'mata*, (F.) *Dessert*, (from [I.] *desseoir*, (*dés*, and *servir*, 'to serve,') 'to clear the table,') The last course at table, consisting of fruits, sweetmeats, &c. In the United States, it means the puddings, pies, &c., which, in England, precede the true dessert.

DESSICATIFS, Desiccative.

DÉSSICATION, Desiccation.

DESTILLATIO, (*dés*, and *stillare*, *stillatum*, 'to drop,') Coryza, Distillation—d. *Pectoris*, Catarrh—d. *Uteri*, Leucorrhœa.

DESTRUCTIO, (*dés*, and *struere*, *structum*, 'to build,') Diaphthora.

DESUDATIO, *Ephidro'is*, *Sudan'ina*, *Hidros'a*, (*desudare*, (*dés*, and *sudare*,) 'to sweat,') *Desudation* means a profuse and inordinate sweating, 'a mackswat,') but, most commonly, the term is applied to an eruption of small pimples, similar to millet seed, which appears chiefly on children, and is owing to want of cleanliness.

DÉSYMPHYSER, (*dés*, and *symphyseis*,) see *Symphysotomy*.

DÉTENTIO, (*détineo*, *detentum*, 'to hold down,') Catalepsy.

DÉTÉRGENS, (*detergere*, *detersum*, (*dés*, and *tergere*, 'to clean,') 'to cleanse,') Abstergent.

DÉTÉRGENTS, *Detergentia*, *Detersoria*, *Extergentia*, *Abstergentia*, *Abstersiva*, *Abstersoria*, *Abluentia*, *Transluentia*, *Sme'tica*, *Emundantia*, *Mundificantia*, *Mundificantia*, *Rhy'tica*, *Abstersiva*, (F.) *Detersifs*, *Mundificantifs*, *Abluents*. Medicines which possess the power to deterge or cleanse parts, as wounds, ulcers, &c. They belong to the class of stimulants, or to that of emollients.

DETERMINATION, *Determinatio*, (*dés*, and *terminus*, 'a boundary,') Strong direction to a given point;—as 'a determination of blood to the head,')

DÉTÉRSIFS, Detergents.

DÉTÉRSORIA, Detergents.

DÉTÉRSORIUM, The place, in ancient bath-

ing establishments, where the bathers were cleansed and dried.

DÉTÊSTATIO, (*dés*, and *testis*, 'a testicle,') Castration.

DÉTÔRSE, (*dés*, and *torquere*, *tortum*, 'to twist,') Sprain.

DETRACTIO SANGUINIS, Bloodletting.

DETRITUS, (*deterere*, (*dés*, and *terere*, *tritum*, 'to rub,') 'to bruise or wear out,') The residuum, occupying the place of the organic texture of parts which have undergone disorganization.

DÉTROIT ABDOMINAL, (*dés*, and *strictus*, 'narrow,') Pelvis (brim)—d. *Inferieur*, Pelvis (outlet)—d. *Périnéal*, Pelvis (outlet)—d. *Supérieur*, Pelvis (brim).

DÉTRONCATION, Detruncatio.

DETRUNCATION, *Detruncatio*, (F.) *Detruncation*, (*dés*, and *truncus*, 'a trunk,') Separation of the trunk from the head of the foetus, the latter remaining in the uterus.

DETRUSION, *Detrusio*, (*detrudere*, (*dés*, and *trudere*, *trusum*, 'to thrust,') 'to thrust down or from,') The act of thrusting or forcing down or away. Applied by Dr. Walshe to lateral displacement of the heart by extraneous pressure.

DETRUSOR URINÆ, *Detrusor*. The muscular coat of the urinary bladder was formerly so called. It was, also, named *Constrictor Vesicæ Uring'ria*.

DÉTUMESCENCE (F.) *Detumescencia*, (*detumescere*, (*dés*, and *tumere*, 'to swell,') 'to cease to swell,') A diminution of swelling. This word has nearly the same signification as the French word *Désenfleur*. The latter is, however, more particularly applied to the diminution of œdema or anasarca; the former, to the resolution of a tumour properly so called.

DEUNX. The ancient name of a weight of 11 ounces, supposing the pound to consist of 12.

DEURENS (FEBRIS), (*deuro*, 'to burn up,') Causus.

DEUS COPULATIONIS, ('god of copulation,') Cupido.

DEUTERANGEA, (*deuteros*, 'the second,') and *anæa*,) Dementia.

DEUTERIA, (*deuteros*, 'the second,') Vogel has used this term for the symptoms produced by retention of the secundines. The word was also applied, by the Greeks, to a second or inferior wine.

DEUTERION, Secundines.

DEUTEROPATHIA, *Hysteropathia*, *Men'hys secundaria*, (*deuteros*, 'the second,') and *pathos*, 'disease,') A secondary disease. One produced by another, and of which it is only, in some measure, symptomatic, or the sympathetic effect.

DÉTÉROSCOPY, *Deuteroscopia*, (*deuteros*, 'the second,') and *scopos*, 'I view,') Second sight. A fancied power of seeing future things or events. Also, a form of hallucination, in which the patient sees a spectral image of himself.

DEUTO, (*deuteros*, 'second,') A prefix denoting two, or double,—as *deutoxide*, having two degrees of oxidation.

DEUTOAZOTATE ACIDE LIQUIDE DE MÉRURE, see *Hydrargyri nitras*.

DEVENS, (*dés*, and *vehere*, 'to carry,') Effluent.

DEVELOPMENT, *Evolutio*, ([F.] *développer*, 'to unfold,') (F.) *Développement*. In Physiology, it means growth or increase; and in Pathology, its signification is similar. It—as well as *Différentiation*—is at times used, in Physiology, for the kind of increase which takes place by the formation of new tissues from those already existing, as in the embryo, whilst growth is restricted to the increase of the same tissues. By development of the pulse is understood an increase in its

strength and fulness. Diseases of development, *Morbi evolutio'nis*, are such as are peculiar to the period of growth.

'TAKING DEVELOPMENTS.' A term used by practical craniologists to signify the act of measuring prominences of the skull, which are regarded by them as indicating the size of corresponding cerebral organs.

DEVELOPMENT, VESICLE OF, see Vesicle of Development.

DEVERTICULUM, Diverticulum.

DEVIA'TION, *Devia'tio*, (*de*, and *via*, 'the way.') Out of the way. By this word is meant—a vicious curvature of the spine or other bones;—faulty direction of the teeth or other part;—the passage of blood, bile, urine, milk, &c., into vessels not natural to them.

DÉVIATION DES RÈGLES, Menstruation (vicarious).

DEVIL BIT, *Veratrum viride*—d. in a Bush, Nigella.

DEVIL'S BIT, *Aletris farinosa*, *Scabiosa succisa*, *Chamelirium luteum*—d. Bite, *Liatris*—d. Dung, *Asafetida*—d. Shoestrings, *Galega Virginiana*.

DEVIRGINATION, (*de*, and *virgo*, gen. *virginis*, 'a virgin.') Defloration, Stuprum.

DEVISCERATION, (*de*, and *viscus*, *visceris*, 'an entrail.') Exenterismus.

DÉVOIEMENT, (*de*, and *voie*, (L.) *via*, 'way.') Diarrhoea.

DEWBERRY, AMERICAN, see *Rubus cæsius*—d. Plant, *Rubus cæsius*.

DEWCLOCKS, *Crusta genu equinæ*.

DEXIOCAR'DIA, (*δεξις*, 'right,' and *καρδία*, 'the heart.') A case in which the heart is found to beat on the right side. It is met with occasionally in pleurisy and pneumothorax.

DEXIS, (*δεξις*.) Degmus.

DEX'TANS. A weight of 10 ounces, supposing the pound to consist of 12.

DEXTERINA, Dextrine.

DEX'TRAD, (*dexter*, 'right-handed.') A term used adverbially by Dr. Barclay to signify 'towards the dextral aspect.' See Mesial.

DEXTRAL ASPECT, see Mesial.

DEXTRALITY, (*dexter*, 'right.') The state of being on the right side. Right-handedness. The state of being right-handed.

DEX'TRIN, *Dex'trine*, *Dextri'num*, *Dexteri'na*, *Brit'ish gum*, *Artific'ial gum*, (*dexter*, 'right-handed.') (So called, from its refracting the rays, in the polarization of light, more to the right hand than any substance known.) A substance obtained by the continued action of diluted sulphuric acid upon starch at the boiling point. It is used in the treatment of fractures, by the 'immovable apparatus.' The bandages are soaked in a solution, in water, of the dextrine—previously moistened thoroughly with tincture of camphor, to prevent it from leaking when the water is added. The solution should be of the consistence of mottos.

DEXTRINUM, Dextrine.

DI, DIS, (*di*, *dis*.) in composition, *bis*, twice, double. Also, as well as *Dis* and *Dis*, separation or division, and, at times, negation and privation.

DIA, (*dia*.) in composition, 'through, asunder, out of, separated.' When prefixed to any therapeutical agent, it meant, in ancient pharmacy, a preparation into which that agent entered.

DIABEBOS, *Astragalus*, *Malleolus*.

DIABÈTE, Diabetes—d. *Chyleux*, Chyluria—d. *Faux*, see Diabetes—d. *Inspide*, see Diabetes—d. *Sucré*, Diabetes (mellitus).

DIABETES, (*διαβητης*, from *dia*, and *βαινω*, 'I pass.') *Urinæ profluvium*, *Hyperdiure'sis*, *Siphoniz'is*, *Urorrhag'ia*, *Polyur'ia*, *Hydrops ad Mat'ur'is* seu *Mat'ell'e*, *Polyur'e'sis*, *Uroze'mia*, *Ureor-*

rhœa, *Dip'sacus*, *Diarrhœa* in *Urinâ* seu *urino'sa*, (F.) *Diabète*, *Flux d'Urine*. A disease characterized by great augmentation, and often manifest alteration in the secretion of urine; with excessive thirst and progressive emaciation. Cullen has described two species—*Diabètes insip'idus* and *D. Mell'itus*; the former, (F.) *Diabète faux* on *insipide*, *Diabète* being, simply, a superabundant discharge of limpid urine, of its usual, urinary taste; the latter, *D. Mell'itus*, called, also, *Paru'ria Mell'ita*, *Diabètes An'glicus* seu *vel'us*, *Melitu'ria*, *Mellithyperu'ria*, *Glucon'ria*, *Glycyrrhœa urino'sa*, *Uroze'mia mell'ita*, *Saccharorrhœa urino'sa*, *Phthisu'ria*, *U'rophthi'sis*, *Ta'bes diuret'ica* seu *diab'etica*, *Dyspep'cia saccharin'ena*, *Apoeno'sis Diab'etes Mell'itus*, *Saccharig'ie diab'etes*, (F.) *Diabète sucré*, *Hyperuororrhœe saccharine*, *Phthisurie sucrée*,—falls under the definition given above. The quantity of urine, discharged in the 24 hours, is sometimes excessive, amounting to 30 pints and upwards; each pint containing sometimes 2½ oz. saccharine matter. This replaces the urea, which is not found in quantity in the urine of those laboring under diabetes. Where the disease is situate is not clear. The whole system of nutrition, however, seems to be morbidly implicated. A part of the urine must be formed at the expense of the system, as the egesta frequently far exceed the solid and liquid ingesta. On dissection, no morbid appearance is met with sufficient to enable us to fix on the seat of this distressing affection.

All the remedies that have been tried have usually been found insufficient in *D. Mellitus*.

D. insip'idus, *Hyperu'e'sis aquo'sa*, *Hydru'ria*, *Hydrur'e'sis*, *Paru'ria incontinens aquo'sa*, *Diab'etes spu'rius*, *Uororrhœa*, *U'real Diab'etes*, (F.) *Polyurie*, *Hyperuororrhœe*, *Diabète insipide*, *Faux diabète*, which occurs in hysterical habits, and has, hence, been called *D. hyster'icus*, is of comparatively trifling moment. Exclusive diet, and attention to the state of the cutaneous transpiration, which have sometimes produced good effects in *D. Mellitus*, have most commonly failed.

DIABETES ANGLICUS, see Diabetes—d. Chylousus, Chyluria—d. Insidious, see Diabetes—d. Lacteus, Chyluria—d. Mellitus, see Diabetes—d. Spurius, see Diabetes—d. Ureal, see Diabetes—d. Verus, Diabetes (mellitus).

DIABET'IC, *Diab'eticus*, (F.) *Diabétique*. Relating to diabetes.

DIABETIC SUGAR, see Glucose.

DIABLOTIN, [F.] (from *diable*, 'the devil' who tempts us.) A name given to a pastil reputed to be aphrodisiac. The principal ingredient of the *Diablotins d'Italie* was powdered cantharides.

DIABOT'ANUM, (*dia*, and *βοταν*, 'an herb.') A medicine, prepared with herbs.

DIABROSIS, (*διαβρωσις*, from *dia*, and *βρωσκειν*, 'to eat.') Erosion, Corrosion.

DIABRO'TICUS, (*διαβρωτικός*.) A substance, capable of causing erosion of the part to which it is applied. It ordinarily means a medicine, whose activity places it between escharotics and caustics. See Corrosive.

DIACAR'YON, (*dia*, and *καρυον*, 'a nut.') *Rob nu'eum*. The rob of nuts or of walnuts.

DIACASSIA CUM MANNÂ, Confectio cassiæ.

DIACATHOLICON, *Diacathol'icum*, (*dia*, and *καθολικος*, 'universal'.) The name of a purge, so called from its general usefulness. It was an electuary, and composed of the pulp of cassia, tamarinds, leaves of senna, root of polypody, flowers of the violet, rhubarb root, aniseed, sugar, liquorice, and fennel.

DIACAUS'IS, (*διακαυσις*, from *dia*, and *καω*, 'I burn'.) Excessive heat. Over-heating.

DIACAUST'IC, *Dicaust'icus*. That which is caustic by refraction; as a double convex lens,

which has been sometimes used for cauterizing an ulcer by directing the sun's rays upon it.

DIACELTATESSON PARACELSI, Pulvis Cornuacini.

DIACHALASIS, (ἀχαλασις, from *dia*, and *χάλασις*, 'to be open or relaxed.') Hippocrates uses this word for fracture of the bones of the skull; or for relaxation and separation of the sutures, in consequence of a wound of the head.

DIACHALCITEOS, (*dia*, and *χάλκισ*, 'chalcitis or colcothar.') A plaster, whose composition is the same as that of the diapalma, except that, in place of the sulphate of zinc, a mixture of oil and colcothar is substituted.

DIACHEIRIS'MUS, (ἀαχειρισμός,) *Diacheir'isis*, (ἀαχειρισίς,) *Tractatio manual'ria*, (*dia*, and *χειρ*, 'the hand.') The preparation, administration, and dispensing of medicines.

DIACHORE'MA, gen. *Diachore'matis*, *Diachore'is*, (ἀαχωρημα, from *διαχωρεω*, (*dia*, and *χωρεω*, 'I retire,') 'I separate from.') Every kind of excreted matter and excretion; but more particularly the fæces and alvine excrement. See Excrement.

DIACHOREMA XYSMATODES, see Ramenta intestinum.

DIACHORESIS, (διαχωρησις,) Excretion.

DIACHRISIS, (*dia*, and *χρῖω*, 'I anoint.') Inunction.

DIACHRISMA, gen. *Diachris'matis*, (ἀαχρημα,) Unguentum.

DIACHRIST'A. Medicines, applied as abstergents to the velum palati, the palate itself, the tongue, &c. Probably gargles.

DIACH'YLOX, *Diach'yllum*, *Emplastrum diach'yloxi*, (*dia*, and *χῶλος*, 'juice'; i. e. composed of juices.) The plaster of this name was formerly made of certain juices. The term is now confined to the EMPLASTRUM PLUMBI or *Lead Plaster*.

DIACH'YLOX GOMMÉ, *Emplastrum eum gummi-resinis*—d. Gum, *Emplastrum gummosum*—d. Gummi, *Emplastrum gummosum*—d. Magnum eum gummi, *Emplastrum galbani comp.*—d. Simplex, *Emplastrum plumbi*—d. White, *Emplastrum plumbi*—d. Yellow, *Emplastrum gummosum*.

DIACHYTICA, (διαχυτικός, from *dia*, and *χωω*, 'I pour out,' 'I melt.') Medicines which discuss tumours.

DIACINE'MA, gen. *Diacine'matis*, (διακίνημα, from *dia*, and *κινεω*, 'I move.') A slight dislocation. A subluxation.

DIACLASIS, (ἀακλασις, from *dia*, and *κλαειν*, 'to break.') Refraction.

DIAC'LYSIS, (ἀακλυσίς,) *Diaclys'mus*, (ἀακλυσμός, from *dia*, and *κλύειν*, 'to wash out.') Rinsing or cleansing—especially of the mouth.

DIACLYSMA, gen. *Diaclys'matis*, (ἀακλυσμα,) Gargarism.

DIACOCYMELON, (*dia*, and *coccy-melon*,) Diaprunum.

DIACODION, (*dia*, and *κωδία*, 'a poppyhead.') Syrupus papaveris.

DIACOD'UUM, *Fæculetio ex Capit'ibus Papav'ris*, (F.) *Diacode*. The ancients had various forms for preparing it. The Syrup of Poppies—*Syrupus Papav'ris* seu *Diacod'dion*—is now substituted for it.

DIACOPE, (διακοπή, from *dia*, and *κοπτειν*, 'to cut.') Abcession, Dissection, Intersection.

DIAC'OPÉ, *En'copé*. A cut, incision, fissure, or longitudinal fracture. When used, since Galen, it generally signifies an oblique incision, made in the cranium by a sharp instrument, without the piece being removed. It is not now employed.

DIACOPÉ CRA'NIU, *Priecis'io seu Dissect'io Cranii*. Opening the head; and separation of the bones of the cranium.

DIACOPRÆ'GIA, (*dia*, *κοπος*, 'excrement,' and *αἶς*, gen. *αἶγος*, 'a goat.') A name given, in Blau-

card's Lexicon, to a medicine, composed of goat's dung, which the ancients praised in diseases of the spleen, parotides, &c.

DIACRISIS, *Nosocrin'ies*, *Crino'ses*, (*dia*, and *κρινω*, 'I separate.') A class of diseases characterized by alterations of secretion.

DIACRISIOG'RAPHY, *Diacrisiograph'ia*, (*diacrisis*, 'separation,' and *γραφη*, 'a description.') A description of the organs of secretion.

DIACRIT'ICA SIG'NA. (Same etymon as *Diacrisis*.) Signs by which one disease can be accurately discriminated from another:—*differential diagn'o'sis*.

DIACYDONITES, Cydoniatum.

DIADÉ'MA, gen. *Diadé'matis*, (διαδήμα,) *Fascia cap'itis*, *Redimic'ulum*, (διαδῶ, *dia*, and *δέν*, 'to bind.') 'I bind round.') A sort of bandage, advised in headach, in which relaxation of the sutures was apprehended.

DIADERMATRIA, (*dia*, *δερμα*, 'skin,' and *ιατρεία*, 'medical treatment.') Endermic method.

DIADÉ'XIS, (διαδέξις,) *Diad'oché*, *Metap'osis*, (διαδεχ'ομαι, (*dia*, and *δεχ'ομαι*, 'to take or receive,' 'to transfer,' 'to succeed to.') A transformation of a disease into another, differing from the former both in its nature and seat.

DIADOCHE, (διαδοχή,) *Diadexis*.

DIAD'OSIS, (διαδosis, from *διαδίδωμι*, 'to distribute.') In some authors, it means the distribution of nutritive matter over the whole body,—in other words, *nutrition*; whilst, in others, it is synonymous with the remission or cessation of a disease.

DIÆDÉ'US, (*di*, and *αἰδία*, 'the parts of generation.') A monster whose organs of generation and urinary bladder are double. It has only been observed in animals.

DIÆRESIS, (διαίρεσις, from *διαίρω*, (*dia*, and *αίρω*, 'I take away,' 'I divide,' 'I separate.') A division or solution of continuity. A surgical operation, which consists in dividing any part of the body. *Hæmorrhag'ia per diæ'resin* is hemorrhage owing to separation or division of vessels.

DIÆRESIS UNGULÆ, Onychoptosis.

DIÆRETICUS, Caustic.

DIÆTA, (*diæta*.) Diet—d. Juris, see Diet—d. Juseuli, see Diet—d. Lactea seu Lactis, Galatodiæta, see Diet.

DIÆTE'MA, gen. *Diæte'matis*, (διαίτημα,) has the same signification as *liet*, with most authors. Galen gives it a more extensive meaning, comprising under it what constitutes Hygiene.

DIÆTETICA, (διαίτητικη,) Dietetics.

DIAGNOSE, Diagnosticate.

DIAGNO'SIS, (διαγνωσις, from *dia*, and *γινωσκω*, 'I know.') *Digno'tio*, *Diagnos'tice*, *Deprehen'sio*, *Discrimina'tion*, (F.) *Diagnose*, *Diagnostique*. That part of medicine whose object is the discrimination of diseases, and the knowledge of the pathognomonic signs of each. It is one of the most important branches of general pathology. See Symptom.

DIAGNOSIS, DIFFERENTIAL, see Diacritica signa.

DIAGNOSIS, PHYS'ICAL. The discrimination of diseases by the physical signs afforded by auscultation, percussion, &c. See Symptom.

DIAGNOS'TIC, (διαγνωστικός,) *Diagnos'ticu*, *Discreti'vus*; (F.) *Diagnostique*. A symptom which is characteristic of a disease.

DIAGNOS'TICATE, sometimes *Diagnose*, (F.) *diagnostiques*. To discriminate one disease or phenomenon from another.

DIAGRYDIUM, (*diacrydion*,) *Convolvulus scammonia*.

DIAGRYD'UUM CYDONIA'TUM. A pharmaceutical preparation, obtained by inspissating and drying, by means of heat, two parts of *scammony*, and one of *quince juice*. It was formerly used as an energetic purgative.

DIAGRYDIUM GLYCERHIZA'TUM. An analogous

preparation, containing extract of liquorice in place of quince juice.

DIAGRYDIUM SULPHURATUM, Scammony, which has been exposed to the vapour of burning sulphur. These *diagrydia* are not now used.

DIALEIPSIS, (διαλειψις, from διαλειπειν, 'to intermit,') *Apyrexia*, *Intermission*.

DIALEIPYRA, (διαλειπειν, 'to intermit,' and πυρ, 'fever,') *Intermittent fever*.

DIALEMMA, gen. *Dialemmatis*, (διαλεμμα, 'an interval,') *Apyrexia*.

DIALEPSIS, διαλεψις, from διαλαμβάνω, 'I intercept,') *Interceptio*. Hippocrates employs this word for the interstices, or intervals, left between the turns of a bandage.

DIALIPSIS, (διαλειπειν, 'to intermit,') *Apyrexia*, *Intermissio*.

DIALYSIS, *Dissolutio*, (διαλυσις, from δια, and λυσις, 'solution,') A dissolution or loss of strength; *Resolutio virium*. Weakness of the limbs. Also, a solution of continuity.

DIAMANT, Diamond.

DIAMASMEA, (διαμασσημα, from δια, and μασσαι, 'to chew,') *Masticatory*.

DIAMBRE SPECIES. A name given by the ancients to powders, one of which bore the name—*Species diambrysi nē odora tis*,—the other, that of *Species diambrym cum odora tis*. The former was composed of cinnamon, angelica root, cloves, mace, nutmeg, galanga, cardamom, and numerous other substances; the latter, besides, had ambergris and musk. These powders were used as tonics, in cases of debility of the stomach, and in certain nervous affections.

DIAMETERS OF THE FŒTAL HEAD, see Fœtal head—d. of the Pelvis, see Pelvis.

DIAMNES, *Euresis*.

DIAMOND, *Adamas*, *Ad' amant*, (ἀδάμας, ἀδαμω, from α, privative, and δαμω, 'I conquer,' 'invincible,') So called from its hardness. (F.) *Diamond*. It is the most precious of all stones, and was formerly conceived to possess extraordinary cordial virtues.

DIAMORUM, *Rob ex moris*, (dia, and μωρον, 'a mulberry,') An ancient syrup prepared with honey and mulberry juice. It was employed as a gargle in sore throat.

DIAMOTOSIS, (dia, and μωτος, 'charpie,' 'lint,') The introduction of lint into an ulcer or wound.

DIANA, (after the Goddess Diana,) *Argentum*.

DIANANCAS MUS, (διαναγκασμος,) (dia, and αναγκω, 'I force,') Coaptation, reduction of a fractured or luxated limb.

DIANOËMA, gen. *Dianoematis*, (διανομη, from δια, and νοω, 'I perceive,') *Imagination*.

DIANTHUS CARYOPHYLLUS, (Δις, Δις, 'Jove,' or from διος, 'divine,' ανθος, 'flower,' and caryophyllum, 'the clove,') *Clove Pink*. Also called *Caryophyllum rubrum*, *Tu'nic*, *T. hortensis* seu *rubra*, *Caryophyllum hortensis*, *Clove July flower*, *Gill'flower*, *Carnation*, (Sc.) *Jeroffleria*, *Geraflouris*. Order, *Caryophyllæ*. (F.) *Œillet giroflée*. The flowers were once much used; but are now only employed in syrup, as a useful and pleasant vehicle for other medicines.

DIAPALMA, *Phœnicium Emplastum*. A plaster composed of equal parts of litharge, olive oil, axunge, water, a certain quantity of sulphate of zinc dissolved in water, and white wax. It is classed amongst the topical, desiccative, emollient, resolvent, detersive, and cicatrizing medicines. Mixed with a quarter of its weight of olive oil, it acquires the consistence of an ointment, and forms the *Cerate of Diapalma*.

DIAPASMA, gen. *Diapasmatis*, (διαπασμα, from δια, and πασσω, 'I sprinkle,') *Catapasma*.

DIAPEDESIS, *Transudatio*, *Pseudatio*, *Pervulatio*, (διαπρησις, from διαπρηω, (δια, and πρηω,

'I leap through,') *Exhalation*, as of blood, in the form of dew, at the surface of the skin, or of any membrane; *Sweating of blood*, (F.) *Sueur de Sang*, *Hæmorrhagia per diapedesin*, *Hæmatopedesis*, *Hæmorrhœia*.

DIAPENSIA CORTUSA, *Sanicula*.

DIAPENTES or **DIAPENTĒ**, (δια, and πεντε, 'five,') A medicine composed of five ingredients. See *Diatessaron*.

DIAPHEMETRIC, *Diaphemetricus*, (from δια, αψη, 'touch,' and μετρον, 'measure,') Relating to the measurements of the tactile sensibility of parts;—as *diaphemetric compasses*.

DIAPHENICON, *Diaphenix*, *Medicamentum ex Pal'mulis*, (dia, and φεινιξ, gen. φαινικος, 'a date,') A drastic electuary, of which the date was the chief excipient.

DIAPHORETIC, (διαφορησις, from διαφορω, δια, and φωρω, 'I convey,' 'I dissipate,') (F.) *Diaphoretic*. A greater degree of perspiration than natural, but less than in sweating. Every kind of cutaneous evacuation.

DIAPHORETIC, (διαφορητικός,) *Diaphoreticus*, *Diaphoreticus*, (F.) *Diaphorétique*. A medicine which excites diaphoresis. Diaphoretics are very uncertain as a class. They are all indirect agents; some sedative; others excitant, and cannot, therefore, be used indiscriminately in disease.

The epithet *Diaphoretic* has also been given, by some, to continued fever, accompanied with constant perspiration.

DIAPHORETIC, MINERAL, *Antimonium diaphoreticum*.

DIAPHORETICUM JOVIALE, see *Antihæctic*.

DIAPHHRAGM, gen. *Diaphragmatis*, (διαφραγμα, from δια, and φρασσω, 'I close,') *Diaphragma*, *Diaphragma*, *Respiratorium Ventris*, *Diaphragmum*, *Phrenicus*, *Septum transversum*, *Diaphragma Thoracis et Ventris*, *Cine'tus*, *Diazoma*, *Præcinctus*, *Diazoma*, *Diazot'ra*, *Hypozoma*, *Hypozoma*, *Perizoma*, *Dissip'tum*, *Dissip'tum*, *Præcordia*, *Succin'gens membr'na seu mus'culis*, *Succinctus*, *Succinctura*, the *Mid'riff*, (Prov.) *Heart-scirts*. A large, azygous muscle; stretched transversely between the thoracic and abdominal cavities, which it separates from each other; tendinous in the centre; thin, almost circular, and unequally convex, upwards. It is fleshy at its circumference, which is attached to the cartilago ensiformis, to the last six ribs, to the aponeurosis stretched from the last rib to the transverse process of the first lumbar vertebra; and, lastly, to the bodies of the first three or four lumbar vertebrae. When it contracts, its fibres become straight, the chest is enlarged, and the abdomen diminished. It is then an inspiratory muscle. It may, also, diminish the capacity of the chest, and be an expiratory muscle. The muscle plays an important part in sighing, yawning, coughing, sneezing, laughing, sobbing, crying, hiccupping, singing, vomiting, and in the excretion of the feces and urine, the expulsion of the fœtus, &c.

DIAPHHRAGMA, *Diaphragm*—d. Anris, see *Tympanum*—d. Cerebri, *Tentorium*—d. Narium, *Septum narium*—d. Oris, *Mylohyoideus*—d. Pelvis, *Levator ani*—d. Ventriclelorum lateraliur cerebri, *Septum lucidum*.

DIAPHHRAGMAL'GIA, *Diaphragmat'gia*, (from diaphragm, and αλγος, 'pain,') Pain in the diaphragm.

DIAPHHRAGMAT'ALGIA, *Diaphragmalgia*.

DIAPHHRAGMAT'IC, *Diaphragmaticus*, *Phren'ic*, *Phren'icus*, (F.) *Diaphragmatique*, *Phrénique*. Belonging to the diaphragm. A name given to several vessels and nerves.

DIAPHHRAGMATIC OF PHREN'IC ARTERIES. These are distinguished into *superior* and *inferior*. The former, called, also, *supradiaphragmatic*, are two in number, one on each side. They arise from

the internal mammary, and descend along the phrenic nerve, to be distributed on the upper surface of the diaphragm. The latter or *infradiaphragmatic* are also two in number. They arise from the upper part of the abdominal aorta, or from the coeliac artery, and divide into two principal branches, which are distributed on the lower surface of the diaphragm and in its substance.

The *superior diaphragmatic veins* follow the same course as the arteries, and empty themselves—the *right*, into the vena cava superior; the *left*, into the corresponding subclavian vein. The two *inferior diaphragmatic veins* open into the vena cava inferior.

DIAPHRAGMATIC HERNIA, Phren'ic Hernia, Diaphragmatocele. The abdominal viscera are occasionally protruded through the diaphragm, either through some of the natural apertures in the muscle, or through deficiencies, or wounds, or lacerations in it.

DIAPHRAGMATIC OR PHREN'IC NERVES, Inter-nal res'piratory of Sir Charles Bell, are two in number; one on the left side, the other on the right. They arise from the second and third nerves of the cervical plexus, about the middle of the neck, and receive two or three filaments from the brachial plexus, after which they descend into the chest at the sides of the pericardium, and are distributed to the diaphragm.

DIAPHRAGMATIC PLEX'USES are two in number; one right, and the other left. They arise from the upper part of the solar plexus, by a small number of branches, which are distributed to the diaphragm, following exactly the branches of the inferior diaphragmatic arteries.

DIAPHRAGMATIC RING, (F.) Anneau diaphragmatique of Chaussier, is a name given to the irregularly quadrilateral aperture by which the vena cava inferior passes through the diaphragm.

DIAPHRAGMATITIS, Diaphragmitis.

DIAPHRAGMATOCELE, (diaphragm, and κηλη, 'a rupture,') see Hernia.

DIAPHRAGMITIS, (diaphragm, and itis,) *Diaphragmatitis, Inflammatio septi transversi, Paraphrenitis, Empresma Pleuritidis Diaphragmaticæ, Paraphrenesis Diaphragmaticæ, Inflammation of the Diaphragm.* The terms, *Paraphrenitis* and *Paraphrenesis* have been obtained from the Peripatetic philosophy, which supposed the seat of the φρην or soul to be the præcordia. The essential symptoms of diaphragmitis are:—painful constriction around the præcordia, with small, quick, laborious breathing. It is a rare disease.

DIAPHRAGMODYN'IA, (diaphragm, and δυν, 'pain,') Muscular or other pain of the diaphragm.

DIAPHRATTON HYMEN, (διαφραττων 'υμην, 'separating membrane,') Mediastinum—d. Membrana, Mediastinum.

DIAPHRAXIS, Diaphragm.

DIAPH'THORA, Destructio, (διαφθορα, from δια, and φθειρειν, 'to corrupt,') Corruption in general; more especially corruption of the foetus in utero. Also, corruption of the blood in the stomach. See Abortion.

DIAPHYLACTIC, (διαφυλακτικός, from δια, and φυλασσειν, 'to preserve,') Prophylactic.

DIAPHYSIS, (διαφysis, from διαφνω, 'I rise between,') *Interstitium, Discriminatio*: 'an interstice, interval, division.' Any thing that separates two bodies. Also, the middle part or body of a long bone, *Corpus ossis*. One of the ligaments of the knee.

DIA'PIA. Some lexicographers use this word synonymously with Diapysis or Suppuration; others have employed it in opposition to Myopia.

DIAPLASIS, (διαπλασις, from δια, and πλασσειν, 'to form,') Conformation, Reduction.

DIAPLASMUS, (διαπλάσμα, from δια, and πλασσειν, 'to form,') Conformation, Reduction.

DIAPNEUSIS, (διαπνευσις, from δια, and πναι, 'I breathe, or exhale,') Perspiration.

DIAPNOE, (διαπνοη,) Perspiration.

DIAPNOEA, (διαπνοια,) Perspiration.

DIAPNOGENOUS APPARATUS, (diapnoe, and γεννω, 'I generate,') see Perspiration.

DIAPNOICUS, Diaphoretic.

DIAPOPHYSES, Transverse processes of the vertebrae.

DIAPORE'MA, gen. Diapore'matis, (διαπορημα, from διαπορειν, 'I doubt,') Anxiety, jactitation.

DIAPRU'NUM, Diaococymelon. A purgative electuary, of which the *pulps of prunes* and *rhubarb* formed the basis. By adding to the *diaprunum simplex* a 24th part of powdered scammony, the *Diaprunum resolutivum* seu *compositum* was formed. It was more active than the former.

DIAPYE'MA, gen. Diapye'matis, (διαπυημα, from δια, and πυον, 'pus,') Empyema, Suppuration.

DIAPYEMATA, Diapyetica.

DIAPYESIS, (διαπυησις,) Suppuration—d. Oculi, Hypopyon.

DIAPYE'TICA, pl. of Dyapye'ticus, Dyapye'ticata. Medicines which promote suppuration.

DIAPYETICUS, (διαπυητικος,) Suppurative.

DIARÆ'MIA, (F.) Diarémie, (δια, ρεω, 'I flow,') and *αιμα, 'blood,')* A pathological condition, said to be common in sheep, in which the globules of the blood are diminished in quantity; the blood itself thinner, and transuding through the coats of the vessels into the cavities.

DIARÆ'MIE, Diaræmia.

DIARIA, (from dies, 'a day,') Ephemera.

DIARRHAGE, (διάρρην, from δια, and ρηγνυμι, 'to break,') Fracture,

DIARRHÉE, Diarrhœa.

DIARRHŒ'A, (διάρρεια, from δια, and ρεω, 'I flow,') *Enterorrhœ'a, Incontinentia alvi, Alvi profusivum seu fluxus aquosus, Ventris profusivum, Cœliorrhœ'a, Cœliolysis, Alvus cit'a, Caca-to'ria, Coporrhœ'a, Catar'hus intestinalis, Alvi fluxus, Rheuma, Epiphora alvi, Fluxus alv'nis, Lax'itas alvi, Desflux'io, Lax, Looseness, Purging, Scouring, (Old Eng.) Lusk, (Sc.) Scour, (Prov.) Ray, Scutter, Skitter, (F.) Diarrhée, Dévoisement, Catarrhe intestinal, Flux de Ventre, Cours de Ventre, Ventre relâché, Courante.* A disease characterized by frequent liquid alvine evacuations, and generally owing to inflammation or irritation of the mucous membrane of the intestines. It is commonly caused by errors in regimen, the use of food noxious by its quality or quantity, &c., constituting the *Diarrhœ'a stercora'ria, D. Crapulosa* of writers. It may be acute or chronic. Many varieties have been made by some nosologists—e. g. mucous, — *Diarrhœ'a mucosa, Blennoche'sia, Blennoche'zia, Mycodiarrhœ'a*; bilious, — *Sleo-cholo'sia, Cholodiarrhœ'a, Diarrhœ'a biliosa*; serous, — *Hydroche'zia, Hydrodiarrhœ'a, Orrhoche'zia*; dependent upon the matters evacuated. Diarrhœa requires different treatment, according to its nature. If caused, as it often is, by improper matters in the intestinal canal, these must be evacuated; and the astringent plan of treatment must not be adopted, unless the discharges seem kept up by irritability of the intestines, or unless they are colliquative. The indiscriminate use of astringents is to be deprecated.

A very fatal diarrhœa prevails amongst the native inhabitants of India, to which Mr. Tytler has given the name *Diarrhœ'a hec'tica*, because, like hectic fever, it seems to obtain habitual possession of the constitution, to operate upon it with scarcely any perceptible intermission, and in general, to defy the most powerful remedies.

DIARRHŒA ADIPOSA, Gras-fondure—d. cum Apepsiâ, Lientery—d. Arthritica, Cœliagra—d. Biliosa, see Diarrhœa—d. Carnosa, Dysentery—d. Chylosa, Cœlic flux—d. Chymosa, Cœlic flux

—d. Crapulosa, see Diarrhœa — d. Cruenta, Hæmatochezia — d. Dyspeptica, Lientery — d. Hepatica, Hepatirrhœa — d. Ingestorum, Lientery — d. Lienteria, Lientery — d. Mucosa, see Diarrhœa — d. Purulenta, Pyochezia — d. Sanguinolenta, Hæmatochezia — d. Stercoraria, see Diarrhœa — d. in Urinâ, Diabetes — d. Urinosa, Diabetes.

DIARRHOÏS'CHESIS, (*diarrhœa*, and *σχέσις*, 'arrest.') Arrest of a diarrhœa.

DIARTHRO'DIAL, *Diarthrodial'lis*. Relating to diarthrosis or movable articulations; as *diarthro'dial articulation*. *Diarthro'dial cartilages* or *incrusted cartilages* are the cartilages which invest the articular extremities of bones.

DIARTHRO'SIS, (*διαρθρωσις*, from *διαρῶ*, (*dia*, and *αρθρω*, 'I articulate,' and *osis*,) *Dearticulation*, *Proarthro'sis*, *Aparthro'sis*, *Abarticulation*, *Coarticulation*, *Perarticulation*, *Rotatio*, (F.) *Emboiture*. A movable articulation. One which permits the bones to move freely on each other in every direction, as in the case of the shoulder joint.

DIASATYR'ION, (*δια*, and *σατυριον*, 'the orchis mascula.') An electuary, of which this plant formed the basis. The ancients attributed to it the faculty of exciting the organs of generation.

DIASCINCI ANTIDOTUS, (*δια*, and *σκινγκος*, 'a kind of lizard,' Mithridate.

DIASCORDIUM, (*δια*, and *σκορδιον*, 'the water germander.') An electuary, so called because this plant entered into its composition. The Parisian codex had a formula for its preparation, under the title, *Electuarium opium astrin'gens seu diascordium*. (R. fol. scord. $\mathfrak{z}$ iss, rosar. rubr., bistort. rad., gentiane, tormentillæ, sem. berber. $\mathfrak{a}\mathfrak{a}$ $\mathfrak{z}$ ss, zingib., piper. long. $\mathfrak{a}\mathfrak{a}$ $\mathfrak{z}\mathfrak{i}\mathfrak{j}$, cassiæ lignæ, cinnamom., dictamn. Cretens., styrac. calamit., galban., gum. acaciæ $\mathfrak{a}\mathfrak{a}$ $\mathfrak{z}$ ss, bol. oriental. præpar. $\mathfrak{z}\mathfrak{i}\mathfrak{j}$, extract. vinos. opii. $\mathfrak{z}\mathfrak{i}\mathfrak{j}$, mel. rosat. præp. lbj, vin. hispan. lbss: fiat electuarium.) In place of the *etyrax calamita*, the balsam of tolu or benjamin may be used. The opium is, in this preparation, in the proportion of 1 to 184. The diascordium is employed in diarrhœa and dysentery, as a tonic, stomachic, and astringent. The common dose is from a scruple to a drachm and a half. See *Pulvis cretæ compositus*.

DIASOSTIC, (*διασωστικός*, from *δια*, and *σῶζειν*, 'to preserve,') Prophylactic.

DIASPASIS, (*διασπασίς*, from *δια*, and *σπᾶω*, 'I draw,') Divulsio.

DIASPERMATON, (*δια*, and *σπέρμα*, gen. *σπερματος*, 'seed,') The ancient name of two cataplasms, composed of seeds.

DIASPHYX'IS, (*διασφύξις*, (*δια*, and *σφύξω*, 'I strike, I beat,') Pulse. Also, a violent beat of the pulse.

DIASTAL'TIC, *Diastalticus*; (*δια*, and *στέλλω*, 'I send,') An epithet applied by Dr. Marshall Hall to the reflex or excitomotory system of nerves; because the actions they induce are performed 'through' the spinal marrow as their essential centre.

DIASTALTICUS, Diastolic.

DIASASÉ'MIA, (F.) *Diastasémie*, (*διασπασίς*, 'separation,' and *αἷμα*, 'blood,') A pathological condition, characterized by a separation of the elements of the blood globules:—the fibrin and albumen separating also from the colouring matter, whilst the fibrin attaches itself to the valves of the heart.

DIASTASE. (Same etymon as Diastasis.) A vegetable principle, allied in its general properties to gluten, which appears in the germination of barley and other seeds, and, by its presence, converts the starch into sugar and gum.

DIASASÉ'MIE, Diastasæmia.

DIAS'TASIS, (*διασπασίς*, (*δια*, and *ιστημι*, 'to place,') 'separation,' 'division,') *Diastéma*, *Dissidentia*. A separation of bones, and parti-

cularly of the bones of the cranium, from each other; of the radius from the ulna, and the fibula from the tibia. The ancients used this word to designate the three dimensions of the body,—length, breadth, and thickness; for the interval separating the patient from the physician; the swelling of varicose veins; the time at which some change occurred in disease, &c.

DIASTEMA, gen. *Diastematis*, (*διαστημα*,) Diastasis, Interstice.

DIASTEMATELYT'RIA, (*diastema*, and *ἐλυτρον*, 'vagina,') An organic deviation, characterized by a longitudinal division or fissure of the vagina.

DIASTEMATENCEPHA'LIA, (*diastema*, and *ἐγκεφαλος*, 'the brain,') An organic deviation, consisting in a longitudinal division of the brain.

DIASTEMA'TIA, (from *diastema*,) A term employed by Breschet for an organic deviation, characterized by the presence of a fissure in the mesial line of the body.

DIASTEMATOCAUL'IA, (*diastema*, and *καυλος*, 'trunk,') An organic deviation, characterized by a longitudinal division of the trunk.

DIASTEMATOCHEI'LIA, (*diastema*, and *χείλος*, 'the lip,') An organic deviation, consisting in a longitudinal division or fissure of the lip.

DIASTEMATOCRA'NIA, (*diastema*, and *κρανιον*, 'the cranium,') An organic deviation, consisting in a longitudinal division of the cranium.

DIASTEMATOCYS'TIA, (*diastema*, and *κύστις*, 'bladder,') An organic deviation, characterized by a longitudinal division of the urinary bladder.

DIASTEMATOGAS'TRIA, (*diastema*, and *γαστήρ*, 'the stomach,') An organic deviation, characterized by a longitudinal division of the stomach.

DIASTEMATOGLOS'SIA, (*diastema*, and *γλῶσσα*, 'tongue,') An organic deviation, characterized by a longitudinal division or fissure of the tongue.

DIASTEMATOGNA'THIA, (*diastema*, and *γνάθος*, 'jaw,') An organic deviation characterized by a longitudinal division of the jaw.

DIASTEMATOME'TRIA, (*diastema*, and *μήτρα*, 'womb,') An organic deviation, characterized by a longitudinal division or fissure of the womb.

DIASTEMATOPYEL'IA, (*diastema*, and *πυελος*, 'pelvis,') An organic deviation, characterized by a longitudinal division or fissure of the pelvis.

DIASTEMATORACH'IA, (*diastema*, and *ράχις*, 'spine,') An organic deviation, characterized by a longitudinal division or fissure of the spine.

DIASTEMATORHI'NIA, (*diastema*, and *ῥιν*, 'the nose,') An organic deviation, characterized by a longitudinal division of the nose.

DIASTEMATOSTAPHYL'IA, (*diastema*, and *σταφυλή*, 'uvula,') An organic deviation, characterized by a longitudinal division of the uvula.

DIASTEMATOSTERN'IA, (*diastema*, and *στερνον*, 'the sternum,') An organic deviation, characterized by a longitudinal division of the sternum.

DIASTEMENTER'IA, (*diastema*, and *εντέρον*, 'intestine,') An organic deviation, characterized by a longitudinal division of the intestine.

DIASTOLĒ, (*διαστολή*, from *διαστέλλω*, (*δια*, and *στέλλω*, 'I send,') 'I dilate,' 'I open,') *Relaxatio seu Remissio cordis et arteriarum*. Dilatation of the heart and arteries, when the blood enters their cavities. It is the opposite movement to *sy'stolē*, in which the heart and arteries contract to send forth the blood. *Dias'tolē* and *sy'stolē* are, consequently, successive movements. *Diastole*, *Mo'tus cor'dis diastalticus*, like systole, occurs simultaneously in the two ventricles. The almost

Inappreciable time, which elapses between the diastole and systole has been called *periyole*, and that which succeeds to the diastole, *peridiastole*. When we speak of the contraction or systole of the heart, as well as of its diastole or dilatation, we mean only that of the ventricles. This dilatation is active.

DIASTOLEUS, (διαστολεὺς,) Dilator.

DIASTOLIC or DIAS'TOLIC, *Diastol'icous*, *Diastol'icous*. Belonging to the diastole of the heart—as 'diastolic impulse of the heart.'

DIASTOLIC IMPULSE OF THE HEART, see Impulse, diastolic.

DIATOMOTRIS, (διαστομοτρίς, from διαστομω, (δια, and στωμα, 'mouth,') 'I dilate an aperture,') Dilating instruments, such as the different kinds of specula for the mouth, anus, vagina, &c.

DIASTREMA, gen. *Diastrem'atis*, (διαστρεμμα, from δια, and στρεφω, 'I turn,') Distortion. Perversion. SpRAIN.

DIASTROPHE, (διαστροφή,) Distortion, Perversion. SpRAIN.

DIAT'ASIS, *Diaten'sio*, (διατασις, from διατείνω, (δια, and τέννω, 'I stretch,') 'I distend,') *Ten'sion*. The reduction of a fracture by extension and counterextension.

DIATESARON, (δια, and τεσσαρες, 'four,') An electuary, into the composition of which entered four medicines; viz. the roots of *gentian*, *aristolochia rotunda* and *bay-berries*, each ʒij. honey Rbj. The whole was incorporated with extract of *juniper*. The *diapentes* is nothing more than this electuary, mixed with two ounces of *ivory shavings*. The *diatesaron* is tonic, and it was formerly employed in cases of stings and bites of venomous animals. It was regarded as emmenagogue, alexiterial, and alexipharmic.

DIATHESIQUE, Diathetical.

DIATHESIS, (διαθεσις, from διατίθημι, (δια, and τίθημι, 'to place,') 'I dispose,') (F.) *Diathēse*. Disposition, constitution, affection of the body: predisposition to certain diseases rather than to others;—(F.) *Inimicence morbide*. The principal diatheses, mentioned by authors, are the *scorbutic*, (*scorbutism*,) *scorbutulous*, (*scorbutulus*,) *scorbutic*, *rheumatic*, *gouty*, (*podagrium*,) and *cal'culous*.

DIATHESIS HEMORRHOIDALIS, see Hemorrhoids—d. Hemorrhagic. Hematophilia—d. Nervosa. Nervous diathesis—d. Purulent, see Pyoemia—d. Rheumatic, see Rheumatic—d. Sthenica, Sthenia.

DIATHETIC, *Diathet'ical*, *Diathet'icous*, (F.) *Diathétique*, *Diathésique*. Relating or appertaining to a diathesis:—as "diathetical diseases," *Diathet'ici* (*morbii*)—gout and cancer, for example.

DIATRAGACANTHUS. A powder, composed of *gum tragacanth*, ʒij. *gum Arabic*, ʒj and ʒij. starch, ʒss. *liquorice*, ʒij. as much of the seeds of the melon, and white poppy; seeds of the water-melon, cucumber and gourd, and sugar candy ʒij. It was used as a demulcent.

DIATRESIS, (διάρρησις, from δια, and τρώω, 'I perforate,') Perforation.

DIATRIMMA, gen. *Diatrium'atis*, (διατριμμα, from δια, and τριβα, 'I rub,') Chafing.

DIATRINSANTALON, see Diatrion.

DIATRION, *Diatrium*, (δια, and τρεις, 'three,') A medicine composed of three ingredients. There were formerly two kinds. The first was called *Diatrium Piper'con spe'cies*, and was formed of black and long pepper, aniseed, thyme, and ginger. It was highly stimulating. The second species was known under the name, *Diatrium santal'um*, *Diatrium Santal'orum pul'vis*, or *Pow'der of the three Sant'als*. They were considered diaphoretic.

DIATRITRIL, (δια, and τρεις, 'third,') The Methodists were so called, who pretended to cure all diseases by subjecting the patient to treatment every third day.

DIATRITOS (διάρητος). Relating to every

third day, and the means then used by the Methodists to cure diseases.

DIAZOMA, gen. *Diazom'atie*, (διαζωμα, from δια, and ζωμα, 'to girdle,') Diaphragm.

DIAZOSMA, gen. *Diazos'matie*, (διαζωσμα,) Diaphragm.

DIAZOS'TER (διαζωστηρ), The twelfth vertebra of the back, because it corresponds to the girdle, ζωστηρ.

DIAZOSTRA, (διαζωστρα,) Diaphragm.

DIBOTHRUM LATUM, (δι, and βόθρον, 'a small pit,') *Bathrioccephalus latus*.

DICEN'TRA CANADEN'SIS, *Cory'dalis Canadensis*, (di, and κεντρος, 'a spur,') *Squid'cel corn*, *Col'ic weed*, *Family*, *Fumariaceae*. An indigenous plant, growing from Maine to Wisconsin, which flowers in May, the flowers having the odour of hyacinths. It has been given internally in syphilis, and applied externally in syphilis and gonorrhoea.

DICEPHALIUM, (di, and κεφαλη, 'head,') Bicephalum.

DICEPH'ALUS, *Biceph'alus*, *Dicod'yimus* (where the duplication extends to the neck), *Janus*. A monster with two heads.

DICERAS RUDIS, (di, and κέρας, 'horn,') *Ditrac'hyaeras rudis*, see Worms.

DICALCON (διχάλκον). A weight, equal to a third part of the obolus.

DICHOPHY'IA, (διχα, 'double, and φυω, 'I grow,') A disease of the hairs, in which they split and grow forked.

DICHROMUS, (διχρ'μος, from di, and χρωμα, 'color,') *Verbena officinalis*.

DICIATRIA, (δίκη, 'justice, and ιατρεια, 'medicine,') Medicine, legal.

DICLIDOSTO'SIS, (διχλιδ, and οστωσις, 'ossification,') Ossification of the valves—as of the heart.

DICLIS (διχλιδ), Valve.

DICOR'YPHUS, *Dicor'anus*, (di, and κορυφη, 'the crown of the head,') A monster with a double vertex or cranium.

DICORYPHUS DIPHYGASTRIUS, Hemipages.

DICRANUS, (di, and κρανιον, 'skull,') *Dicoryphus*.

DICROTUS, *Dicr'otus*, *Dicrotatus*, *Dicrot'ic*, (di, and κροτω, 'I strike,') An epithet given to the pulse, when it seems to beat twice as fast as usual. It is, also, synonymous with the term rebounding, the artery rebounding after striking, so as to convey the sensation of a double pulsation. It has been considered, and with truth, to frequently foretell hemorrhage. In bad cases of typhus, it certainly announces such a tendency. See Pulse, dirotic.

DICTAMNE, *Dictamnus albus*—d. *de Crete*, *Origanum dictamnus*.

DICTAMNUS ALBUS, (δικταμνος, from Δικτε, a mountain of Crete; *D. Fraxinell'ia*, *Fraxinella*, *F. Dictam'nus*, *White Fraxinell'a*, *Bas'tard Dil'tany*, (F.) *Dictamn*, *Fraxinella*). The fresh root has been considered nervine, anthelmintic, and emmenagogue. It is not used.

DICTAMNUS CRETICUS, *Origanum dictamnus*—d. *Fraxinella*, *Dictamnus albus*.

DICTOIDES, (δικτωρ, 'a net, and ειδος, 'resemblance,') Reticular.

DICTYITIS, (*dictyon*, and *itis*,) Retinitis.

DICTYON, (δικτυον,) Rete.

DIDELPHYS, (δι, and ειλως, 'uterus,') *Di-hysteria*.

DIDYMAL'GIA, (διδυμοι, 'the testicles, and αλγος, 'pain,') Pain in the testicles.

DIDYMIS, (from διδυμος,) Epididymis.

DIDYMITIS, (*didymus*, and *itis*,) *Hernia humoralis*.

DIDYMUS, (διδυμος,) Gemellus, Testicle—d. *Symphygastrius*, *Gastrodidymus*—d. *Symphygogastrius*, *Hypogastrodidymus*—d. *Symphyo-*

perinaeus, Pygodidymus—d. Symphyothoracogastrius, Thoracogastroididymus.

DIECBOL'ION, (διαβολιον, from *di*, and *εβαλλω*, 'ε and βαλλω, 'I cast out.') A name given, by the ancients, to a remedy which they believed capable of producing abortion.

DIERENBACH, MIN'ERAL WATERS OF. Dierenbach is a city in Bavaria, two leagues from which is a sulphurous spring.

DIERVIL'LA TRIF'IDA, *D. Canadensis* seu *lu'tea* seu *hu'milis* seu *Tournefort'i*, *Lonicera Diervill'a*, (after Dierville, a French Surgeon.) *Bush hon'ysuckle*. An indigenous plant of the Honey-suckle tribe—Lonicereæ. *Ord.* Caprifoliaceæ—whose flowers appear from June to August. It has been used as a diuretic; and in gonorrhœa and syphilis.

DI'ES, He'mera, 'A Day; (F.) *Jour*. The day is, properly, the period during which the solar light illumines our horizon; but commonly, also, we designate by the word *day* the period of 24 hours, *Nycthe'meron*, which is frequently divided into four parts—morning, midday, evening, and midnight. In antiquity, great importance was attached to the observation of days in disease. The medical day is usually reckoned at 24 hours,—universally, in estimating the duration of a disease. In parts of the United States, it comprises only the time when the sun is above the horizon, as regards the administration of medicine, so that if a medicine be ordered to be taken four times a day, it is understood to mean during the 12 hours of day.

DIES CANICULARES, see *Canicula*—d. *Contemplabiles*, *Critical days*—d. *Contemplantes*, *Indicating days*—d. *Critici*, *Critical days*—d. *Decretorii*, *Critical days*—d. *Indicantes*, *Indicating days*—d. *Indicatorii*, *Indicating days*—d. *Indices*, *Indicating days*—d. *Internuntii*, *Critical days*—d. *Judicatorii*, *Critical days*—d. *Radicales*, *Critical days*.

DI'ET, (ἀιτα,) *Diet'a*, *Diet'e'ma*, (F.) *Diète*. Originally, this word signified nearly the same thing as *Hygiene* and *Regimen*,—that is, *Diet* was the employment of every thing necessary for the preservation of health and life. At the present day, it usually signifies a particular kind of food and drink. Also, abstinence:—to put any one upon diet, (F.) *mettre quelqu'un à la diète*, means to deprive him of his usual nourishment. The terms *milk diet*, *Diet'a lactis*; *flesh or meat diet*, *Diet'a carnis*; *soup or broth diet*, *Diet'a juj'ris vel jusculi*, &c., explain themselves. See *Aliment*, *Hygiene*, and *Regimen*.

DIET, BROTH, see *Diet*.

DIET DRINK. A decoction or potion, variously composed, and used in considerable quantity, for the purpose of purifying the blood. The *Decoctum Lusitanicum* or *Lisbon Diet-drink* is one of the most celebrated. See *Decoctum Sarsaparillæ Compositum*.

DIET, DRY. A diet in which there is but a small or no allowance of fluid.

DIET, FLESH, see *Diet*—d. *Meat*, see *Diet*.

DIET SCALE, see *Dietary*.

DIET, SLOP, SLOPS—d. *Soup*, see *Diet*.

DIETARY, *Victus Ratio*. A regulated diet or allowance of food. Every large hospital, military or civil; every prison and eleemosynary institution of every kind, has certain dietetic regulations, and many of them have *Tables of Dietary* or *Diet Scales*, which are more or less implicitly followed. See *Diet*.

The Diet Scale of the British Navy allowed from 31 to 35½ ounces of dry nutritious matter daily; of which 26 ounces were vegetable, and the rest animal—9 ounces of salt meat, or 4½ ounces of fresh. That of the Navy of the United States has been as follows:—*Three days in the week*—Pork, 16 oz.; beans or peas, 7 oz.; biscuit,

14 oz.; pickles or cranberries, 1 oz.; sugar, 2 oz.; tea, ¼ oz.;—40½ oz. *Two days in the week*—Beef, 16 oz.; flour, 8 oz.; fruit, dried, 4 oz.; biscuit, 14 oz.; tea and sugar, 2½ oz.; pickles or cranberries, 1 oz.;—45½ oz. *Two days in the week*—Beef, 16 oz.; rice, 8 oz.; butter, 2 oz.; cheese, 2 oz.; biscuit, 14 oz.; tea and sugar, 2½ oz.; pickles or cranberries, 1 oz.;—45½ oz. The daily Diet allowance to the British soldier has been one pound of bread, and three-quarters of a pound of meat. In the Army of the United States, the quantity of beef, salt or fresh, issued per day, is a pound and a quarter; of pork or of bacon, three-quarters of a pound. Ordinarily, the soldiers have fresh meat four days in the week, and salt meat the other three days; but every diet scale is subject to mutations in the individual articles. It must combine, however, a due admixture of nitrogenized, nonnitrogenized, and inorganic materials. See *Aliment*. The work of Dr. Pereira on 'Food and Diet,' and the Treatise on Hygiene by Dr. W. A. Hammond, of the United States Army, 1863, contain numerous Dietaries, civil and military. See, also, *Ration*.

DIÈTE, *Diet*.

DIETET'ICS, (διαιτητικα τέχνη,) *Dietet'icæ*, *Dietet'ica*, *Medic'ina Dietet'ica*, (F.) *Diététique*. A branch of medicine, comprising the rules to be followed for preventing, relieving, or curing diseases by diet. Dietetics is diet administered according to principle. It is an important part of Hygiene. A well regulated system of diet has great power in checking disease, and likewise in preventing it. A proper knowledge of dietetics is, indeed, as important as that of the *Materia Medica*, strictly so called.

Dietetics has been used, also, synonymously with Hygiene.

DIETÉTIQUE, *Dietetics*.

DIETET'ISTS, *Dietet'istæ*, (F.) *Diététistes*. Physicians who apply only the rules of dietetics to the treatment of disease.

DIET'IC OR DI'ETIC, *Diet'icæ*, (from *Diet*.) Relating or appertaining to diet.

DIETIC DISEASES, *Diet'ici morbi*, according to Dr. Wm. Farr are such as are produced by diet in its enlarged science—as scurvy, rickets, alcoholism, &c.

DIEU-LE-FILT, MIN'ERAL WATERS OF. The waters of Dieu-le-filt, in France, are chalybeate, and much sought after.

DIEURYSMUS, (*δια*, and *ευρυνω*, 'to dilate,') *Dilatation*.

DIEX'ODUS, (διαξόδος, from *dia*, and *εξόδος*, 'an exit or way out,') *Di'odus*. Any opening by which an excretion takes place.

DIF, as a prefix, see *Di*.

DIFFERENTIAL DIAGNOSIS, see *Diacritica*.

DIFFERENTIATION, (*différo* (*di*, and *fero*), 'to be different,') *Development*.

DIFFICULTAS INTESTINORUM, ('difficulty of the intestines,') *Dysentery*.

DIFFLATIO, (*dif*, and *flare*, *flatum*, 'to blow away,') *Perspiration*.

DIFFORMITÉ, (*dif*, and *forma*, 'form,') *Deformation*.

DIFFU'SIBLE (stimulants), (*diffundere* (*dif*, and *fundere*, *fusum*, 'to pour'), 'to pour apart or abroad,') Those stimulating medicines are so called, which augment the action of the vascular and nervous systems in an acute but transitory manner.

DIGAS'TRICUS, (*di*, and *γαστήρ*, 'a belly,') *Biven'ter*, *B. Moxil'is*, *Dop'rimentæ Moxil'is Biven'ter*, *Bigas'ter*. (F.) *Μαστοδομωγένης*, *Μαστοδογένης*—(Ch.), *Digastrique*, *Abaisseur de la mâchoire inférieure*. The name *Digastricus* was formerly given to several muscles. It is now restricted to one of the muscles of the superior hyoid

region. The digastricus is thick and fleshy at its extremities, thin and tendinous at its middle. It is attached to the mastoid groove of the temporal bone, and to a fossette at the side of the symphysis menti. Its tendon passes through an aponeurotic ring, which is attached to the os hyoides.

The use of the digastricus is to depress the lower jaw, or to raise the os hyoides, and to carry it forwards or backwards, as in deglutition.

The strong double-bellied muscle, which forms the gizzard of birds, is also called *Digastricus*.

DIGASTRICUS CRANII, Occipitofrontalis.

DIGASTRIQUE, Digastricus.

DIGENY, (*di*, and *γενος*, 'generation,') see Generation.

DIGERENTIA, (*digere* (*di*, and *gerere*, 'to carry'), 'to digest,') Digestives.

DIGESTIBLE, (*digere*, *digestum*, 'to digest,' and *habilis*, 'able,') *Concoctus habilitas*. Capable of being digested. All food is not equally digestible, and some of the most important is the least so:—the fat of meat, for example. Certain substances, again, are entirely rebellious. As a general rule, albuminous aliments—as tender mutton and beef, are the most easily appropriated.

DIGESTIF, Digestive.

DIGESTION, Digestion—*d. Depravata*, Dyspepsia—*d. Difficilis*, Dyspepsia—*d. Laesa*, Dyspepsia.

DIGESTION, *Digestio*, (from *digere*, *digestum*, 'to digest,') *Cocctio*, *C. Ciborum*, *Pepsis*, *Digestive Process*. Digestion is a function, by means of which alimentary substances, when introduced into the digestive canal, undergo different alterations. The object of it is to convert them into two parts; the one, a reparatory juice, destined to renew the perpetual waste occurring in the economy; the other, deprived of its nutritious properties, to be rejected from the body. This function is composed of a series of organic actions, differing according to the particular organization of the animal. In man they are eight in number, viz.: 1. Prehension of food. 2. Mastication. 3. Insalivation. 4. Deglutition. 5. Action of the stomach. 6. Action of the small intestine. 7. Action of the large intestine. 8. Expulsion of the feces.

DIGESTION is also a *pharmaceutical* operation, which consists in treating certain solid substances with water, alcohol, or other menstruum, at a slightly elevated temperature,—in a sand-bath, for example, or by leaving them exposed for some time to the sun.

DIGESTIVE, see Digestives.

DIGESTIVE PRINCIPLE, Pepsin—*d. Process*, Digestion.

DIGESTIVE TEXTURE. The particular organic condition of substances which affects their digestibility.

DIGESTIVE TUBE, Canal, alimentary.

DIGESTIVES, *Digestiva*, *Digerentia*; (F.) *Digestiva*. A term given, by surgeons, to substances, which, when applied to a wound or ulcer, promote suppuration; as the *ceratum resinae*, warm cataplasms, fomentations, &c. Such substances are vulgarly termed *draw'ing*.

DIGESTIVUS, (F.) *Digestif*. Relating or appertaining to digestion; see, also, Digestives.

DIGITAL, *Digitalis*, (from *digitus*.) Having the shape of a finger; *digitated*. Belonging to the fingers.

The *Appendix vermiformis caeci* is sometimes called DIGITAL APPENDIX.

DIGITAL ARTERIES, VEINS, and NERVES, are those distributed to the fingers.

DIGITAL BLANC, Clavaria.

DIGITAL CAVITY, *An'cyroid cavity*, *Cor'nu descendens ventriculi lateralis*. The occipital portion of the lateral ventricle of the brain.

DIGITAL COMPRESSION, see Compression.

DIGITAL HUMAIN, Clavaria.

DIGITAL IMPRESSIONS, (see Impressions, digital,) are the slight depressions observable on the inner surface of the bones of the cranium, which correspond to the cerebral convolutions.

DIGITALE PETITE, *Gratiola officinalis*.

DIGITALE, (F.) *Doigtier*. A finger stall. The term *Doigtier d'Asdrubali* has been given to a small iron instrument used for measuring the dimensions of the pelvis. Placed at the end of the index finger, it adds to its length and enables it to reach the promontory of the sacrum.

DIGITALINE, see Digitalis.

DIGITALIS, (from *digitale*, because its flower resembles a finger stall,) *Digitalis purpurea*, *Baccharis*, *Bacchar*, *Bacchar* [?], *Fozglove*, (Sc.) *Deadmen's bells*, (Prov.) *Flap'dock*. Ord. Scrophularinæ. *Sez. Syst. Didynamia Angiosperma*. (F.) *Digitale*, *Gants de notre dame*, *Doigtier*. The leaves of this plant, which are indigenous in Great Britain, are powerfully sedative,—diminishing the velocity of the pulse, diuretic, and sorbefacient. In over-doses, Digitalis causes vomiting, purging, dimness of sight, vertigo, delirium, hicough, convulsions, and death:—all the symptoms, in short, which characterize the acro-narcotic class of poisons. Its active principle has been called *Digitaline* or *Digitalin*, *Digitalinum* (Ph. B. 1864). It is said to be a hundred-fold stronger than the most active preparation of digitalis.

Digitalis has been administered in inflammatory diseases, phthisis, active hemorrhage, dropsy, &c.; but although it is a powerful remedy, it has not been as much employed as it probably would have been in particular cases, owing to the overstrained eulogiums, which many have passed upon it in almost all diseases. The average dose is one grain, in the form of pill, which may be repeated every six or eight hours.

DIGITALIS MINIMA, *Gratiola officinalis*.

DIGITARIA DACTYLON, *Cynodon dactylon*.

DIGITATION, *Digitalio*, *Productio dentata*, *Insertio denticulata*, *I. digitata*, (F.) *Dentature*, *Dentelure*. A division into processes having the form of fingers. Several muscles, as the *serrati*, exhibit digitations, similar to those which the fingers form, when held separate.

DIGITATIONES TUBARUM FALLOPII, see Tube, Fallopian.

DIGIT' IUM. Desiccation or atrophy of the fingers. Sauvages calls the same affection *Paronychia Digit'ium*, see Paronychia.

DIGITORUM TENSOR, *Extensor brevis digitorum pedis*.

DIGIT'ITUS, *Dactylus*, *Finger*, (F.) *Doigt*. A name given to the prolongations which form the extremity of the hand. There are five on each hand: the first, the thumb, *Anticheir*, *Pollex*, *Al'lus Pollex*, *Ma'nus par'va majori adjutrix*, *Dig'itus pri'mus seu mag'nus*, *Pro'manus*, (F.) *Pouce*; the second, the *In'dex*, *Dig'itus in'dex seu secundus seu saluta'ris seu saluta'rius seu demonstrato'rius seu demonstrati'vus*, *Indica'tor*, *Demonstrato'r*, *Li'chanos*; the third, *D. me'dius seu tert'ius seu famo'sus seu impud'i'cus seu obsc'u'us seu infam'is seu ver'pus*, *mid'dle fing'er or long fing'er*, (F.) *Doigt du milieu*; the fourth, the *ring fing'er*, *Dig'itus annula'ris*, *Para'm'eos*, *D. quar'tus seu medica'm'lis seu medic'us seu cor'dis*, *Jat'ricus*, (F.) *Annulaire*: and the little *fing'er*, *Oti'tes*, *Dig'itus auricula'ris seu min'imus seu par'vus seu quin'tus seu my'ops seu oti'tes*, *Ear finger*, (F.) *Auriculaire*, *Petit doigt*. All of these have three phalanges, except the first, which has only two.

DIGITUS ANNULARIS, Annular finger—*d. Auricularis*, see *Digitus*—*d. Cordis*, see *Digitus*—*d. Demonstrativus*, see *Digitus*—*d. Demonstratorius*, see *Digitus*—*d. Famosus*, see *Digitus*—*d. Hippocraticus*, *Finger*, *Hippocratic*—*d. Impudens*, see *Digitus*—*d. Index*, *Index*, see *Digitus*—*d. Indicativus*, *Index*, see *Digitus*—*d. Infamis*, see *Digi'*

tus — d. Magnus, Pollex, see Digitus—d. Medicinalis, see Digitus—d. Medicus, see Digitus—d. Medius, see Digitus—d. Minimus, see Digitus—d. Myops, see Digitus—d. Obscenus, see Digitus—d. Otites, see Digitus—d. Parvus, see Digitus.

DIGITUS PE'DIS, *Ortil'us, Toe, (Sc.) Toe, (F.) Orteil.* The toes are five in number, and distinguished numerically, reckoning from within to without. The first is, also, called *great toe, tom toe, (F.) gros orteil*; the fifth, the *little toe, petit orteil*. They have nearly the same organization as the fingers.

DIGITUS PRIMUS, Pollex, see Digitus—d. Quartus, see Digitus—d. Quintus, see Digitus—d. Salutaris, see Digitus—d. Salutaris, see Digitus—d. Salutariorius, Index—d. Secundus, Index, see Digitus.

DIGITUS SEMIMOR'TUUS, 'half dead finger.' A torpid condition of the fingers, met with in connection with chlorosis, and cerebro-spinal affections.

DIGITUS SUPERNUMERA'RIVS, *Supernu'merary fin'ger, (F.) Doigt surnuméraire.* An occasional monstrosity occurs, in which there are supernumerary fingers; and at times owing to hereditary transmission.

DIGITUS TERTIVS, see Digitus—d. Verpus, see Digitus.

DIGNA'THUS, (*di*, and *γναθος*, 'lower jaw.') A monster having two lower jaws.

DIGNOTIO, (*dignoscere, dignotum*, 'to discriminate,') Diagnosis.

DIHYPOGAS'TRIUS, (*di*, and *ὑπογαστριον*, 'the hypogastrium.') A monster whose pelvis, together with the lower portion of the abdomen, is double.

DIHYSTERIA, *Dime'tra, Didel'phys, U'terus du'plex, (di*, and *ὑέρων*, 'uterus.') The state in which there is a double uterus.

DIJUDICATIO, (*di*, and *judicare, judicatum*, 'to judge,') Crisis.

DILACERATIO, (*di*, and *lacerare, laceratum*, 'to tear,') Laceration.

DILATANTS, Dilating agents.

DILATATEUR, Dilator—d. *Antérieur du larynx*, Crico-thyroid muscle—d. *Postérieur du larynx*, Crico-arytenoid, posterior.

DILATATIO, Dilatation—d. Bronchiorum, Bronchia, dilatation of the—d. Intestinorum, Enterectasis—d. Ventriculi, Gastrectasis.

DILATATION, *dilatatio*, (*dilatare, dilatatum, (latum facere)*, 'to enlarge;') *Eurys'mus, Aneurys'mus, Dieurys'mus.* Augmentation of the bulk of a body, occasioned by a separation of some of its molecules. Caloric has the property of dilating all bodies. In Surgery, it means the accidental or preternatural augmentation of a canal or opening; as in aneurisms, varices, &c., or the process of enlarging any aperture or canal. When used so as to obtain a view of parts, as by the speculum, it is termed *Dioptris'mus*.

DILATATOIRE, Dilator.

DILATATORIUM, Dilator.

DILATATORIVS, Dilator.

DILATING A'GENTS, *Dilatan'tia, (F.) Dilatants.* Certain substances used in surgery, either to keep parts separate which have a tendency to unite—as after opening an abscess, to prevent the edges of the incision from uniting; or to increase and dilate openings of canals, either when natural, or formed accidentally or artificially. These agents differ from each other: the chief are—*prepared sponge tents, gentian root, bougies, sounds, dried peas for issues, &c.*

DILATOR, *Dilatato'rius, Dias'toleus, (F.) Dilatateur ou Dilatoire, (di*, and *ferre, latum*, 'to carry.') A muscle, whose office it is to dilate certain parts; as the inspiratory muscles, which dilate the chest.

DILATOR, *Dilatato'rium, (F.) Dilatateur.* An instrument, used for dilating a wound, excretory

canal, or other natural or artificial opening. When employed to obtain an inspector of internal parts, it is termed *Speculum, Diop'tra, or Diop'tron*. There are several instruments of this kind, each taking its name from the part to which it is applied; as *Speculum Oris, S. Nasi, S. Uteri, &c.*

DILATOR, ARNOTT'S. A modification of the old dilators for strictures of the urethra. It consists of a tube of oiled silk, lined with the thin gut of some small animal to make it air-tight, and fixed on the extremity of a small canula, by which it is distended with air or water, from a bag or syringe at the outer end, whilst a stop-cock or valve serves to keep the air or water in, when admitted. As soon as the bag is passed within the stricture or strictures, as much air is to be injected into it as the patient can easily bear. The instrument is not much used.

DILATORES ALARUM NASI, Compressor Naris. DILATRIS TINCTORIA, Lachnanthes tinctoria.

DILL, (Sax. *dile*,) *Anethum graveolens*. DIL'DOO, *Di'doe, Phallus, Fas'cinum, Pe'nis succeda'neus, Mentula factit'ia.* An artificial penis.

DILLY, *Anthemis cotula*.

DILUEN'TIA, pl. of *Diluens, (dihuo, (dis*, and *luere)*, 'I wash away.') (F.) *Délayants.* Medicines which have been conceived proper for augmenting the fluidity of the blood and other animal liquids. All aqueous drinks are diluents. They are administered, with great advantage, in various diseases. In fever, water, which is the most familiar diluent, may be freely allowed; the only precaution being to give it *hot* in the cold stage, *cold* in the hot, and *tepid* in the sweating. In diseases, where it is considered necessary to abstract blood largely, diluents should not be given too freely. The abstraction of blood occasions activity of absorption, and the mass is speedily restored. It is also obvious, that in cases of inflammation of the mammae, in nurses, diluents should not be freely allowed, as they increase the secretion of milk, and add to the irritation. When demulcents are exhibited in cases of urinary disease, they act simply as diluents; their mucilaginous portion is digested in the stomach and small intestine,—the watery portion alone being separated by the kidney.

DILWEED, *Anthemis Cotula*.

DIMETRA, (*di*, and *μητρα*, 'uterus,') Dihysteria.

DINANT, MIN'ERAL WATERS OF. Dinant is a small town, six leagues from St. Malo, in France, where are mineral waters, containing carbonate of iron, chloride of sodium, &c. They are much esteemed.

DINGEE, Dengue.

DINIC, (from *dinus*,) Antidinic.

DINKHOLD, MIN'ERAL WATERS OF. A rich carbonated water, situate near the junction of the Lahn with the Rhine, in the duchy of Nassau. It contains sulphate of soda, chloride of sodium, carbonate of soda, sulphate of lime, carbonate of lime, and sulphate of magnesia.

DINOMANIA, (*dinos*, 'a whirl,' and *μανια*, 'mania,') Tarantismus.

DINUS, (*dinos*,) Vertigo—d. *Scotoma, Scotodynia*—d. Vertigo, Vertigo.

DIOBOLON, (*διωβολον*, from *di*, and *ωβολον*,) Scruple.

DIODONCEPHALUS, (F.) *Diodoncephale, (di*, *odous*, gen. *οδους*, 'a tooth,' and *κεφαλη*, 'head,') A monster with a double range of teeth, or a double jaw.

DIODUS, (*διόδος*, from *dia*, and *οδος*, 'a way,') Diexodus.

DIECESIS, (*διοικησις*, from *dia*, and *οικειν*, 'to manage a house,') Dispensation.

DIONCOSIS. (*διονκωσις*, from *dia*, and *oikos*, 'a tumour.') The Methodists applied this name to a sort of tumefaction or plethora, occurring either directly from too great a quantity of fluid circulating in the system, or owing to the retention of substances which ought to be excreted. It is the antithesis to *symptosis*. See Intumescence.

DIONYSIA'NUS. (F.) *Dionysien*, (*Διονυσος*, 'Bacchus,' who is represented by the poets as wearing horns.') One who has long or horn-like excrescences.

DIONYSIS/CUS, (same etymon.) One who has a long horn-like excrescence on the frontal region.

DIOPHTHALMUS, (*δι*, and *οφθαλμος*, 'an eye,') Binocular.

DIOPTRA, (*διοπτρα*, from *dia*, and *οπτομαι*, 'to see,') Speculum, see Dilator.

DIOPTRISMUS, (from *dioptron*,) see Dilatation.

DIOPTRON, (*διοπτρον*,) Speculum, see Dilator.

DIORTHO'SIS, (*διορθωσις*, from *διορθω*, 'I make straight,') The reduction of a fractured or luxated limb.

DIOSCOREA, (after Dioscorides,) see Yam.

DIOSCORE'A VILLO'SA, *Wild Yam'root*; indigenous: Order, Dioscoriaceæ; flowering in July. A decoction of the root has been prescribed in bilious colic. It is said to be expectorant, diaphoretic, and, in large doses, emetic.

DIOSCURI, (*διοσκουροι*, from *Δις*, gen. *Διός*, 'Jupiter,' and *κουροι*, 'boys,') Castor and Pollux, —so called from their twin-like resemblance. See Parotitis.

DIOSMA, *D. crenata*.

DIOS'MA CRENA'TA, (*διος*, 'divine,' and *οσμη*, 'odour,') *Baros'ma crenata*, *Agathos'ma crenata*. A South African plant, the powder of whose leaves—*Buch'u Leaves*, *Dios'ma* (Ph. U. S., 1842, *Buch'u*, 1851), *Buck'u*, *Boc'choe*, *Boch'u*, *Boo'cho*, *Buch'ho*, (F.) *Diosmæ crénulæ*. Ord. Rutaceæ—is used by the Hottentots to perfume their bodies. It has been employed in chronic affections of the bladder and urinary organs in general; and has also been given in cholera. It is often adulterated in commerce, by the substitution of less potent plants of the same family, as *Diosma serratifolia*, and *Eupleurum serrulatum*.

Buc'co, *Buchu*, is referred by the British Pharmacopœia (1864), to *Baros'ma betul'ina*, *B. crenulata*, and *B. serratifolia*.

DIOSMA SERRATIFOLIA, *D. crenata*.

DIOSMÉE CRÉNELÉE, *Diosma crenata*. **DIOSPYROS LOTUS**, (*διοςτροπος*, from *διος*, 'divine,' and *τροπος*, 'wheat,') *Fa'ba Græ'ca*, *Indian Date Plum*, (F.) *Plaqueminier d'Europe*. Ord. Ebenaceæ. This tree grows in some of the southern parts of Europe. Its fruit is very astringent, and has been recommended in dysentery and hemorrhage.

DIOSPYROS VIRGINIA'NA, *Lotus Virginia'na*, *Persim'mon*. A common tree in the middle parts of the United States. The fruit, *Persim'mons*, *Yellow Plums*, *Winter Plums*, *Seed'ed Plums*, which is only eatable after frost, (when it is tolerable,) is sometimes made into cakes with bran. These, being dried in an oven, are kept to make beer. When bruised in water, fermentation takes place. The unripe fruit, *Dios'pyrus* (Ph. U. S.), is distressingly acerb and astringent. It has been used as an astringent. The bark of the tree is extremely bitter, and may be used where bitters are indicated.

DIO'TA, *Dyo'ta*, (*διωτος*, from *δι*, and *οις*, gen. *ωτος*, 'ear,') Two-eared, two-handled. Applied to a wooden cup, lined with a composition of resin, cinnamon, cloves, and ginger, to give more flavour to beer. It was formerly much used in the north of Europe.

DIPHORUS, *Diphrus*.

DIPH'RUS, *Di'phorus*, (*διφρος*, from *δι*, and

φωρ, 'I carry,') Properly a seat for two. A close stool.

DIPH'RUS MAIEU'TICUS, (*διφρος μαιευτικος*,) *Scl'la obstetric'ia*. An obstetric chair.

DIPH'RYGES, (*διφρυγες*, from *δι*, 'twice,' and *φρυγα*, 'I torrefy,') The oxide of copper, more or less pure. The ancients reckoned three kinds of diphryges, which they used as astringents.

DIPHTHERIA, (*διφθερα*, 'a skin or membrane,') Diphtheritis.

DIPHTHER'IC, *Diphther'icus*, *Diphtherit'ic*, *Diphtherit'icus*, (F.) *Diphthérique*, *Diphthéritique*. Same etymon as Diphtheritis. Relating or appertaining to diphtheria.

DIPHTHERITIC, *Diphtheric*.

DIPHTHÉRITE, *Diphtheritis* — *d. Buccale*, Stomatitis, pseudomembraneous.

DIPHTHERIT'IS, (from *διφθερα*, 'a skin, or membrane,' and *itis*,) *Diphther'ia*, *Diphtherit'ic* Inflammation or *Phlegma'sia*, *Pellie'ular Inflammation*, (F.) *Diphthérite*, *Diphthéropathie*. A name given by M. Bretonneau to a class of diseases, which are characterized by a tendency to the formation of false membranes; and which affect the dermoid tissue,—as the mucous membranes, and even the skin. See Cynanche maligna, and Pharyngitis, diphtheritic.

DIPHTHERITIS OF THE THROAT, Pharyngitis, diphtheritic—*d. Trachealis*, Cynanche trachealis.

DIPHTHÉROPATHIE, (*διφθερα*, 'a skin or membrane,' and *παθος*, 'affection,') Diphtheritis.

DIPLASIASMUS, (*διπλασιασμος*, from *διπλασιος*, 'double,') Duplication.

DIPLECOIA, (*diplo*, and *ακη*, 'hearing,') Double hearing.

DIPLOCEPHA'LIA, (*diplo*, and *κεφαλη*, 'head,') A monstrosity characterized by the presence of two heads on the same body.

DIPLO, (*διπλος*, 'double,') in composition, double.

DIP'LOË, (*διπλην*, from *διπλος*, 'double,') *Diplo'sis*, *Mediull'ium*, *Medium Culv'riae Discri'men*. The areolar structure, which separates the two tables of the skull from each other. The ancients applied the term, also, to the proper coat of the uterus [?].

The Diploë has the same use as the cellular structure of the bones in general. See Cancelli.

DIPLOËTIC, *Diploic*.

DIPLOGEN'ESIS, (*diplo*, and *γενεσις*, 'generation,') (F.) *Diplogénese*. An organic deviation, which consists in the duplication of parts,—monstrosity by duplication, *Duplic'itas monstro'sa*, constituting double monsters.

DIPLOÏC, *Diplo'icus*, *Diploët'ic*, *Diploët'icus*, (F.) *Diploïque*, *Diploëtique*. Relating or appertaining to the diploë.

DIPLOIC OR DIPLOËT'IC VEINS, *Vena diplo'icæ* seu *diploët'icæ*. Veins in the flat cranial bones, the trunks and larger branches of which, *veins of Breschet*, run mostly separately in special arborescent larger canals—*Breschet's bone canals*, which terminate in the *Emissoria Santorini*.

DIPLOLEPIS GALLÆ TINCTORIÆ, see Quercus infectoria.

DIPLO'MA, gen. *Diplo'matis*, (*διπλωμα*, from *διπλω*, 'I fold double,') A letter or writing conferring some privilege: usually applied to the document, certifying that a person has obtained the title of Doctor. It was so called because formerly written on waxed tables, folded together. Diploma is also used in pharmacy for a vessel with double walls,—as a water-bath.

DIPLO'PIA, (*diplo*, and *οπτομαι*, 'I see,') *Vi'sus Duplicatus*, *Ditto'ria*, *Dittop'ria*, *Ambio'ria*, *Double Vi'sion*, (F.) *Bévue*. An affection of the sight, in which two distinct sensations are produced by the same object, which consequently seems double. Sometimes more than two are

seen; but still the disease is termed *Diplopia* and *Suffusio Multiplicans*. The affection arises from some derangement in the visual axes, in consequence of which, the images are not impressed on corresponding parts of the retina of each eye. At times, the diplopia affects one eye, or each eye when used alone, owing to some irregularity in the curvature, density or position of the dioptric media. It is then called *Unicocular Diplopia*, *Monodiplopia*.

The diplopia of both eyes has been termed *Amphodiplopia*, *Amphoterodiplopia*, and *Amphamphoterodiplopia*.

DIPLOSIS, (διπλωσις,) *Diploë*, Duplication.

DIPLOSO'MA, gen. *Diplosoma'tis*, (*diplo*, and *soma*, 'body,') The *Diplosoma crenatum* is an imperfectly described entozoon, which has been passed from the urinary bladder. It varies in length from four to six or eight inches, and is thinnest in the middle, where it is bent at an acute angle upon itself, so that the two halves hang nearly parallel, and give to it an appearance as if two worms had been tied together by their heads. It has been confounded with the *Spiroptera hominis*.

DIPLOSOMA'TIA, *Diplosoma'tia*. An organic deviation characterized by the presence of two complete bodies, united by one or more of their parts.

DIPNOUS, (διπνους, from *di*, and *πνω*, 'breath,') Having two vent holes; *Bispirus*. An epithet applied to wounds which pass through a part, and admit the air at both ends.

DIPROSO'PUS, *Iriod'ymus* et *Opod'ymus*, *Opodid'ymus*, (*di*, and *προσωπον*, 'countenance,') A monster having a double face.

DIPROSOPI'S *DIÆDE'US*, (*di*, and *προσωπον*, 'countenance,; and *di*, and *αἶδω*, 'parts of duplication,') A double monster, in which the duplication affects superiorly the face, and inferiorly the anterior pelvic region.

DIPROSOPI'S *DIHYPOGAS'TRIUS*, (*di*, and *προσωπον*, 'countenance,; and *di*, and *υπογαστριον*, 'the hypogastrium,') A double monster, in which the duplication affects superiorly the face and inferiorly the lower part of the body, four lower extremities being always present—*Tetrascelus*.

DIPSA, (διψα,) Thirst.

DIPSACUS, (διψακος,) *Diabetes*.

DIP'SACUS FULLONUM, (from *dipsa*, said to be so called, owing to the leaves being so placed as to hold water,) *Herba Cardui Ven'eris*, *Carduus Ven'eris*, (F.) *Cardère cultivé*, *Chardon à foulon*, *Chardon à bonnetier*, has had similar properties with the next ascribed to it. Both have been also regarded as stomachic.

DIPSACUS SYLVES'TRIS, *Cultivated Ten'sel*, (F.) *Cardère*. *Ord.* *Dipsacæ*. The roots of this European plant are diuretic and sudorific. The water, which collects at the base of the leaves, has been recommended as an eye-water.

DIPSE'TICUS, (διψητικός, from *dipsa*,) (F.) *Dipsétique*. A remedy believed to be capable of exciting thirst. See *Alterative*.

DIPSO'DES, (διψωδης, *dipsa*, and *odes*,) *Sit'iens*, *Siticulo'sus*, (Old Eng.) *Af'forst'*, (Prov.) *Thursty*. Thirsty. Causing thirst.

DIPSOMA'NIA, (*dipsa*, and *mania*. Really, 'thirst-mania.' Often, however, applied to habitual drunkenness and to delirium tremens. An insatiable desire for intoxicating liquors.

DIPSOPI'ATHY, *Dipsopathi'a*, (*dipsa*, and *παθος*, 'affection,') A mode of treatment, which consists in abstaining from drinks.

DIPSO'SIS, (from *dipsa*,) *Morbid thirst*. The desire for drinking, excessive or impaired. A genus in the class *Celiaca*, order *Enterica*, of Good.

DIPSOSIS *AVENS*. *Polydypsia*—d. *Expers*, *Adipsia*.

DIP'TERIX ODOR'ATA, *Commamon'na odorata*. The fruit of this large tree of Guiana is a pod, inclosing a single seed. The bean, *Tonka bean*, has a strong, agreeable, aromatic odour, and is used to flavour snuff.

DIPY'GUS, (*di*, and *πυγν*, 'nates,') A double monster, the duplication being confined to the posterior portion of the lower end of the trunk—the coccygeal region.

DIPY'RE'NON, (διπυρνον, from *di*, and *πυρν*, 'a kernel,') A *specil'lum* or probe with two buttons or kernels—one at each end.

DIPYRITES, (διπυρης, 'twice baked,') Biscuit.

DIPYROS, (διπυρος, from *di*, and *πυρ*, 'fire,') Biscuit.

DIR'CA PALUS'TRIS, (after *Δικην*, a fountain at Thebes, because it grows in moist places.) *Leath'erwood*, *Swamp Leath'erwood*, *Moose'wood*, *Rope'bark*, *Bois de plomb*, (Canada.) *Ord.* *Thymeleaceæ*. An indigenous shrub, which grows in boggy woods and low wet places throughout the United States. It is analogous to mezereon in its action—six or eight grains of the fresh bark producing violent vomiting, preceded by a sense of heat in the stomach, often followed by purging. Applied to the skin, the bark vesicates.

DIREC'TOR, *Itinera'trium*, (*dirigere*, *directum*, (*di*, and *regere*, 'to rule,') 'to direct,') A *Conductor*. A grooved sound for guiding a knife, in dividing any part, (F.) *Sonde cannelée*.

DIRECTOR PENIS, *Ischioavernosus*.

DIRIBITORIUM, (*diribere*, 'to separate,') *Mediastinum*.

DIRLING, Tingling.

DIRTEATING, *Clathonophagia*, *Geophagism*.

DIRUPTIO, (*di*, and *rumpere*, *ruptum*, 'to break,') *Rhexis*.

DIS, as a prefix, see *Di*.

DISC, see *Disk*.

DISCHARGE', (*dis*, and *charge*,) *Es'roë*, *Flux'us*, *Proflu'vium*, (F.) *Ecoulement*. In pathology, an increased discharge from any part, that naturally secretes a fluid.

DISCHROA, *Dyschroæ*.

DISCRET, *Discrete*.

DISCRETE', (*discernere*, *discretum*, (*dis*, and *cernere*,) 'to separate,') *Discret'us*, *Intertinct'us*, *Sep'arated*, (F.) *Discret*. This epithet is given to certain exanthemata, in which the spots or pustules are separated from each other. It is opposed to confluent.

DISCRE'TA PURGA'TIO. The purgation or expulsion of some particular matter.

DISCRETIVUS, *Diagnostic*.

DISCRETORIUM, *Diaphragm*.

DISCRI'MEN, 'separation, division.' A bandage, used in bleeding from the frontal vein; so called, because, in passing along the sagittal suture, it divides the head into two equal parts.

DISCRIMEN CALVARIÆ MEDIUM, *Diploë*—d. *Narium*, *Septum N*.

DISCRIMEN NASI; a bandage, in the form of the letter X, intended to support the nose, in transverse wounds of the organ.

DISCRIMEN THORACIS ET VENTRIS, *Diaphragm*.

DISCUS, (δίσκος,) *Disk*, (F.) *Disque*. A quoit; a plate or platter, or anything resembling it in form.

DISCUS PROLIGERUS, *Proligerous disc*—d. *Vitellinus*, *Proligerous disc*.

DISCUSSIFS, *Discutients*.

DISCUSSIO, (*discutere*, *discussum*, (*dis*, and *quater*, 'to shake,') 'to shake apart,') *Resolution*.

DISCUSSIVA, *Discutients*.

DISCUSSORIA, *Discutients*.

DISCU'TIENTS, *Discutient'ia*, *Resolvent'ia*, *Discussiv'ra*, *Discussoria*, (F.) *Discussifs*, *Résolutifs*. Substances which possess the power of repelling or resolving tumours.

DISEASE, old French *désaise*, (*dis*, and *ease*.) *Morbus*, *Nosos*, *Noseu'ma*, *Noseu'ma*, *Nusus*, *Pathos*, *Pathe'ma*, *Lues*, *Malum*, *Passio*, *Ægritu'do*, *Æpro-ta'tio*, *Vit'ium*, *Arrhos'tia*, *Arrhos'te'ma*, *Valetu'do ad'versa*, *Mal'ady*, *Complaint*, *Sickness*, *Distem-per*, *Distem'perature*, *Eg'ritude*, *Ail*, *Ailment*, *Ill-ness*, (F.) *Maladie*. An opposite state to that of health, consisting in a change either in the posi-

tion and structure of parts, or in the exercise of one or more of their functions, or in both.

By some *Disease* is applied to structural change, whilst *Disor'der* is restricted to functional derange-ment.

The following table, essentially that of Dr. C. J. B. Williams, comprises the chief elements of structural disease, all of which are dependent on modified cell action.

DISEASED NUTRITION.	{	Increased—Hypertrophy, Diminished—Atrophy.			
		Inflammation, Induration, Softening, Transformation and Degeneration.			
		Deposits.....	{	Euplastic.....	Cicatrices, False membranes, Cirrhosis.
				Cacoplastic.....	Fibro-cartilage, Gray tubercle, Atheroma.
			Aplastic.....	Yellow tubercle, Calcareous matter.	
Perverted.....	{	Growths.....	{	Nonmalignant...	Cysts, Tumours, Hydatids.
			Malignant.....	Carcinoma, Encephaloma, Melanosis.	

ALTERED MECHANISM.	{	Contraction, Dilatation, Obstruction, Compression, Displacement, Rupture, &c.
-----------------------	---	--

DISEASE OF ADDISON, see Capsule, renal—d. Barking, *Mal de Laira*—d. Blue, Cyanopathy—d. of Bright. Kidney, Bright's Disease of the—d. English, Rachitis—d. Extrinsic, External disease—d. Family, see Hereditary—d. Foul, Syphilis—d. Fungoid, Encephaloid—d. Gastro-enteric, Gastro-enteritis—d. Grinders', Asthma, grinders'—d. Hæmatocerebriform, Encephaloid—d. Hereditary, see Hereditary—d. Lucifer match makers, see Phosphorus—d. Phosphorus jaw, see Phosphorus—d. Pretended, Feigned disease—d. Simulated, Feigned disease—d. Surgical, External disease.

DISEASED, Sick.

DISFIGURATION, (*dis*, and *figurare*, *figura-tum*, 'to form,') Deformation.

DISGORGE'MENT, (F.) *Dégorgement*, (*dis*, and *gorge*, 'the throat.') An opposite condition to that of *Engorgement*. The discharge or abstraction of a certain quantity of fluid, which had previously collected in a part; as *Disgorgement of bile*. It also means vomiting.

DISGUST, (*dis*, and *gustare*, 'to taste,') *Cibi fastid'ium*, *Apoeti'ia*, *Asit'ia*, *Apoclei'sis*, *Abomina'tio*, *Siccha'sia*, *Hor'ror Cibo'rum*, *Loath'ing*. An aversion for food; (F.) *Dégout*. *Disgust* is not the same as *Anorexia*. The latter is only a want of appetite; the former consists in real repug-nance for food.

DISINFEC'TANT, *Disinfectans*, (*dis*, and *in-ficio*, *infectum*, (*in*, and *facio*), 'I infect,') (F.) *Désinfectant*. This term has been restricted by some to agents that are capable of neutralizing morbid effluvia; but the author includes under it, also, antiseptics or agents that are capable of removing any incipient or fully formed septic condition of the living body, or of any part of it.

DISINFECTANT OF ELLERMAN, see Ferri chloridum—d. of Mr. James Young, see Man-ganese, sulphate of.

DISINFECTING LIQUID, SIR WILLIAM BURNETT'S, see Burnett's Disinfecting Liquid—d. Liquid of Condy, see Condy's Disinfecting Fluid—d. Liquid, Labarraque's, Liqueur sodæ chlorinatæ—d. Liquid of Larnaudès, see Zincæ Sulphas—d. Liquid of Ledoyen, see Ledoyen's disinfecting liquid.

DISINFEC'TION, *Disinfectio*, (F.) *Désinfect-*

tion. The act of attempting to destroy mias-mata, with which the air, the clothing, &c., may be infected. Various means have been proposed for this purpose. Those most commonly em-ployed are: chlorine, sulphurous and chlorohydric acid gases, vapours of vinegar, nitrous acid, and what is, perhaps, the most manageable of any, chlorinated lime, or any of the chlorides of the alkalis. It is more than questionable whether any chemical action occurs between these agents and the miasmata, whence results a compound which is harmless, or in other words, whether there is any agent that is capable of destroying morbid miasms. Disinfection also includes the action of antiseptics.

The following fumigations have been largely employed as disinfectants.

Chlo'rine or *Oxy-muriat'ic Fumiga'tion*, *Solu'tio alexite'ria Gaub'iana*, *Fumiga'tio antiloin'ica Gau'bii*, *Alexite'rium chlo'ricum*, *Fumiga'tio Gny-tonien'sis*, is made by adding common salt ʒij, to black oxide of manganese ʒj, sulphuric acid ʒj, and water f ʒij. This may be carried through an apartment, or be shut up in it.

Nit'rous Fumiga'tion, *Alexite'rium Ni'tricum*, *Fumiga'tio Smyth'iana*, may be formed by placing nitrate of potass ʒiv, and sulphuric acid ʒij, in a saucer on hot sand.

DISJUNC'TI MOR'BI (*dis*, and *jungere*, *junc-tum*, 'to join,') *Disjoin'ed diseases*. Fernelius has thus denominated diseases which occupy different organs, and are not produced by each other.

DISK, *Discus*—d. Blood, Globule of the blood—d. Intervertebral, Intervertebral Cartilage—d. Proligerous, see Proligerous Disk.

DISLOCATIO, (*dis*, and *locare*, *locatum*, 'to place,') Dislocation, Luxation—d. Lienis seu Splenis, Splenectopia.

DISLOCATION, Luxation.

DISMAL, Melancholy.

DISOD'ICUS, (*dis*, and *odos*, 'threshold,') Hav-ing a double opening.

DISOMATOUS, see Disomus.

DISO'MUS, (F.) *Disome*, (*di*, and *σωμα*, 'body,') A monster with two bodies. Such a monster is said to be *disom'atous*.

DISORDER, (from *dis*, and *order*,) see Dis-ease.

DISORDERED, Sick.

DISORGANIZA'TION, *Organo'rum destructio*, (F.) *Désorganisation*. A complete morbid change in the structure of an organ, or even total destruction of its texture. In sphacelus and sloughy ulcers, we have examples of this change.

DISPENSARE, Dispensary, Dispensary.

DISPEN'SARY, *Dispensarium*, (*dispendere*, (*dis*, and *pendere*, *pensum*, 'to weigh,') 'to take from a weight or mass,') 'to distribute,') The shop or place in which medicines are prepared. Also, an institution in which the poor are furnished with advice and necessary medicines. (F.) *Dispensaire*. This word is used by the French synonymously with Dispensary.

DISPENSAT'ION, *Dispensa'tio*, *Diace'sis*, *Epine'mesia*. The composition and distribution of medicines according to prescription. In France, it is more commonly applied to the weighing, measuring, and arranging of the articles which have to enter into a formula, prior to combining them.

DISPENSATOR, Apothecary.

DISPEN'SATORY, *Dispensato'rium*; *Antidota'rium*, *Lib'rum Pharmaceut'icum*, (F.) *Dispensaire*. A book which treats of the composition of medicines. A Dispensary differs from a Pharmacopœia, in containing the physical and medical history of the various substances; whilst the Pharmacopœia is mainly restricted to the mode of preparing them. The Pharmacopœia too, is published under the authority of, and by the members of a college or association; whilst a Dispensary contains the whole of the Pharmacopœia or Pharmacopœias, with additions by the author, under whose authority alone it usually appears. The Pharmacopœia, in other words, is *officinal*; the Dispensary, generally, *private*. Formerly, the words were used indiscriminately. See Formulary, and Pharmacopœia.

DISPLACED, see Percolation.

DISPLACEMENT, Percolation.

DISPOSIT'ION, *Disposit'io*, (*dis*, and *ponere*, *positum*, 'to put or set,') A particular condition of the body, which renders it susceptible of altering, suddenly, to a state of health or disease;—of improving, or becoming changed for the worse. The *disposition* to phthisis pulmonalis is sometimes so strong, owing to original conformation, that the disease will make its appearance, notwithstanding every care. See Diathesis.

DISPUTATIO, (*dis*, and *putare*, 'to reason,') Thesis.

DISQUE, Discus—d. *Proligère*, *Proligerous* Disk.

DISQUES INTERVERTÉBRAUX, Intervertebral cartilages—d. *Sanguis*, Globules of the blood.

DISRUP'TIO, (*disrumpere*, (*dis*, and *rumpere*,) 'to break off,') A puncture, which interests deep-seated parts.

DISSECTING ANEURISM, See Aneurism.

DISSECTIO, Dissection—d. *Tendinum*, Tenotomy.

DISSECTION, *Dissect'io*, (*dissecare*, (*dis*, and *secare*, *sectum*, 'to cut,') 'to cut open,') *Diace'opé*, *Sec'tio Anatom'ica*, *Practical Anatom'omy*, *Neurot'omy*. An operation, by which the different parts of the dead body are exposed, for the purpose of studying their arrangement and structure. Dissection has received various names, according to the organ concerned;—as *Osteotomy*, *Syndesmotomy*, *Myotomy*, *Angiotomy*, *Neurotomy*, *Desmotomy*, &c.

DISSECTOR, *Prosect'or*, *Pro'tomus*, (F.) *Prosecteur*. A practical anatomist. One who prepares the parts for the anatomical lecture.

DISSEPIMENTUM NARIUM, (*dissepio*, (*dis*, and *sepere*, 'a fence,') 'I separate,') *Septum narium*—d. *Thoracis*, *Mediastinum*.

DISSEPTUM, (*dissepio*, 'I separate,') *Dia-phragm*.

DISSERTATION, INAUGURAL, Thesis.

DISSIDENTIA, (*dis*, and *sedere*, 'to sit,') *Dia-stasis*.

DISSIPIUM, (*dissepio*, 'I separate,') *Dia-phragm*.

DISSOLUTIO, Dissolution, Solution—d. *Sanguis*, *Hæmatelysis*—d. *Ventriculi*, *Gastromalacia*.

DISSOLU'TION, *Dissolu'tio*, *Dial'ysis*, (*dissolvere*, (*dis*, and *solvere*, *solutum*,) 'to loosen,') 'to melt,') This word is in frequent use, in the writings of the humorists. It is employed, particularly, in speaking of the blood,—to designate, not its entire decomposition or putrefaction, but a diminution in its consistence.

DISSOLUTION, Solution.

DISSOLVANTS, Dissolventia.

DISSOLVEN'TIA, *Sol'vents*, (F.) *Dissolvants*. Medicines believed to be capable of dissolving swellings, concretions, &c. *Discutients*, *Resolvents*.

DISTAD, (*disto*, (*di*, and *sto*,) 'to stand apart,') Away from a centre. Towards the far extremity. In anatomy, used by Dr. Barclay adverbially, to signify 'towards the distal aspect.'

DISTAL ASPECT. An aspect of a bone from the trunk, or towards the extremity.

DISTEMPER, (*dis*, and *temperare*, 'to proportion duly,') Disease.

DISTEMPERANTIA, Dyserasia.

DISTEMPERATURE, Disease.

DISTEMPERED, Sick.

DISTENSIO, (*distendere*, *distensum*, 'to stretch,') *Diatasis*, *Tension*, *Tetanus*—d. *Nervorum*, *Convulsion*.

DISTENSION DE LA VÉSICULE DU FIEL, Turgescentia vesicæ fellææ.

DISTICHI'ASIS, *Distich'ia*, *Districh'ia*, *Districh'iasis*, *Disto'chia*, *Distach'iasis*, (*ιστιχιασος*, from *dis*, and *στιχος*, 'a row,') An increased number of eyelashes; some turned towards the eye, and irritating the organ; whilst others preserve their natural direction. See Entropion.

DISTILLATE, see Distillation.

DISTILLA'TION, *Destilla'tio*, *Stalag'mus*, (*destillare*, (*de* and *stillare*,) 'to drop, little by little,') *Catastalag'mus*. An operation, by which—by the aid of heat and in closed vessels—the volatile are separated from the fixed parts of bodies, with the view of collecting the former, when condensed in appropriate receivers. The product of distillation is the *distill'ate*, (G.) *Destillat*.

The ancients distinguished distillation *per latus*, *per ascen'sum*, and *per descen'sum*, according to the direction which the volatilized matters were made to take. In *distillation per latus*, the apparatus is so arranged, that the vapour passes horizontally through a succession of spaces before reaching the receiver. *Distillation per ascensum* is the ordinary method by the still. In *distillation per descensum*, the fire is applied above and around the top of the apparatus; and it is so arranged that the vapour must pass downwards. *Fractional distillation* is when ingredients volatilized at different degrees of heat are separated from each other during the process. When dry organic matter is placed in an apparatus for distillation, and heat is applied until all volatile matter is driven out, the process is called *dry* or *destructive distillation*.

DISTILLATION, DESTRUCTIVE, see Distillation—d. *Dry*, see Distillation—d. *Per Ascensum*, see Distillation—d. *Per Descensum*, see Distillation—d. *Per Latus*, see Distillation.

DISTOC'IA, *Distoc'ia*, (*dis*, and *τοκος*, 'birth,') Delivery of twins.

DISTÆCHIA, Distichiasis.

DISTÆCHIASIS, (διστοχίας,) Distichiasis.

DISTOMA HEPATICUM, gen. *Distomatæ*, *Distomum hepaticum*, (di, and στομα, 'mouth.') *Fasciola hepatica* seu *humana* seu *lanceolata*, *Planaria latiuscula*, *Gourd-worm*, *Fluke*, *Liver Fluke*, (F.) *Douve*, *D. du Foie*. An obovate, flat worm, an inch in length, and nearly half an inch broad; sometimes found in the gall-bladder of man, whence it occasionally passes into the intestinal canal. It is one of the most common varieties of worms, infesting the livers of the sheep, goat, ox, stag, fallow-deer, horse, ass, hog, hare, &c. The treatment is like that required for other worms. **DISTOMA** seu **DISTOMUM LANCEOLATUM**, a different species, has been found in the biliary ducts of man, and of different phytivorous mammalia.

DISTOMA OCULI HUMANI, *Distomum oculi humani*. A parasite observed by Geseheid in the eye of a child who had suffered from lenticular cataract.

DISTOMUM HÆMATOBIUM, see Worms—d. Hepaticum, *Distoma hepaticum*—d. *Oculi humani*, *Distoma oculi humani*.

DISTORTIO, Distortion, Sprain, Strabismus—d. Oris, Canine laugh.

DISTORTION, *Distorsio*, *Intorsio*, (*distorque*, (*dis*, and *torquere*, *tortum*), 'to wrest aside.') *Diastremma*, *Dias'trophæ*, (F.) *Distortion*. Usually applied to the preternatural curvature of a bone; as *distortion of the spine, limbs*, &c. It signifies, also, a morbid state of the muscles of the eye, constituting squinting or strabismus.

DISTORTOR ORIS, *Zygomaticus major*.

DISTRIBUTORIA LACTEA THORACICA, Thoracic duct.

DISTRICHIA, (*dis*, and τριχ, gen. τριχες, 'hair.') Distichiasis.

DISTRICHIASIS, Distichiasis.

DISTRIX, gen. *Districis*, (*dis*, and τριχ, gen. τριχες, 'hair.') *Trichosis distrix*, *Fissura capitolium*. Hairs of the scalp, weak, slender, and splitting at their extremities.

DITOCIA, Distocia.

DITRACHYERAS, (*di*, τραχες, 'rough,' and κερας, 'horn.') A genus of intestinal-worms. The *Ditrachyeras rudis*, *Dic'eras rudis*, (F.) *Bicorne rude*. This fancied entozoon was first observed in the evacuations of a female, by M. Sultz, of Strasburg. It is of a fawn colour, from 3 to 5 lines in length, with an oval head, flattened and terminating in a point posteriorly; contained in a membranous sac, and furnished anteriorly, with a bifurcated, rugous horn. Laënnec considers it a vesicular worm or hydatid. According to Von Siebold, there is no such entozoon. The undigested seed of the mulberry, which had been eaten, was mistaken for it.

DITTANDER, *Lepidium sativum*.

DITTANY, (see Dictamnus.) *Cunila Mariana*—d. American, *Cunila Mariana*—d. Bastard, *Dictamnus albus*—d. of Crete, *Origanum dictamnus*—Mountain, *Cunila Mariana*.

DITTECOIA, (διττος, 'double,' and ακοη, 'hearing.') Double hearing.

DITTOPIA, (διττος, 'double,' and ωψ, 'the eye.') Diplopia.

DITTOPSIA, Diplopia.

DIURESIS, *Diuresis*, *Diureticæsthesis*, *Ureæsthesis*: (διουρω, 'to pass the urine,' and αίσθησις, 'feeling.') The desire or want to pass the urine.

DIURESIS, *Diurism*, (*dia*, and ουρω, 'I pass the urine.') (F.) *Diuræse*. An abundant excretion of urine.

DIURETIC, *Diureticus*, *Ischuret'ic*, *Uret'icus*, *Urina'lis*, (F.) *Diurétique*. A medicine which has the property of increasing the secretion of urine. Diuretics act by producing a discharge of the

watery and other parts of the blood; and, by such discharge, indirectly promote absorption over the whole system. Hence, they are employed in dropsy. To this class belong squill, juniper, digitalis, &c. Some of them appear to affect the system of nutrition, by modifying the metamorphosis of tissue, and giving occasion to the separation of tissue materials from the blood by the kidneys. Hence they were called, by Dr. Golding Bird, *Renal alteratives*. To this class belong the alkalies, alkaline salts, &c.

DIURETICA, *Arnica Montana*.

DIURETICÆSTHESIS, *Diuresisæsthesis*.

DIURISM, *Diuresis*.

DIVARICATIO, (*di*, and *varicare*, 'to straddle.') Ectropion.

DIVERSORIUM CHYLI, *Receptaculum chyli*.

DIVERTICULA SPIRITUM ANIMALIUM, Ganglions, nervous.

DIVERTICULUM, *Deverticulum*. (F.) *Diverticule*. 'A turning;' (*divertere*, (*di*, and *vertere*), 'to turn aside.') A blind tube branching out of the course of a longer one. An organ which is capable of receiving an unusual quantity of blood, when the circulation is obstructed or modified elsewhere, is said to act as a diverticulum. Also, a malformation or diseased appearance of a part, in which it passes out of its regular course. It is sometimes applied to such a condition of the alimentary canal. Also, a hole to get out at. A by-passage. See Ectrope.

DIVERTICULUM NUCULI. The opening through which the round ligaments of the uterus pass.

DIVERTICULUM PHARYNGIS, *Pharyngocele*—d. *Vateri*, *Plica longitudinalis Vateri*.

DIVIDING, *Div'idens*, (*dividere*, *divisum*, (*di* or *dis*, and the *Hetruscan* verb *iduo*, 'to part or portion.') (F.) *Divisif*. That which divides or separates.

DIVIDING BANDAGE, *Fas'cia div'idens*, (F.) *Bandage divisif*, is a bandage employed for the purpose of keeping parts separated from each other. It is used particularly to prevent improper union; as in cases of burns of the neck or other parts.

DIVINATIO, (*divino*, *divinatum*, (from *divus*, 'divine,' 'to foresee,' 'divine.') *Mantia*.

DIVISIF, *Dividing*.

DIVISION, *Divisio*, *Div'esis*. Same etymon as *Dividing*. The accidental separation of parts naturally united; in other words, a wound or solution of continuity. Most frequently, however, it means an operation, which consists in cutting certain parts, with the view of fulfilling some therapeutical indication.

DIVULSIO, *Divulsio*, (*divellere*, (*di*, and *vellere*, *vulsum*), 'to pull asunder.') A term used in Surgery, to express the rupture or laceration of organs by external violence.

DIVULSIO URINÆ, Cloudiness of urine.—See ENOËREMA.

DIXON'S ANTIBILIOUS PILLS, *Pilulæ antibiliosæ*.

DIYLISIS, (διυλισις,) *Colatio*.

DIYLSMUS, (διυλισμος,) *Colatio*.

DIZZINESS, *Vertigo*.

DIZZY, (Sax. *dirri*, *dirrig*.) *Vertiginous*.

DOCCIONE, MINERAL WATERS OF. This spring is at Lucca, in Italy. It is a thermal saline.

DOCH' MÈ, (δοχμ.) A measure of the Greeks, equal to about four fingers' breadth: *Dactylo-doch'mè*.

DOCIMASIA, (δοκιμασία,) *Docimas'ticæ*, *Docimasiolog'ia*, (δοκιμαζω, 'I try or prove the quality of any thing.') The act of assaying.

DOCIMASIA MEDICAMENTO'RUM ET VENENO'RUM. The testing of medicines and poisons.

DOCIMASIA PULMONUM, *D. Pulmonalis*, *Pneobiomanti'a*. *Pneobiomanti'a*, *Pneobiomant'ica*. Lung proof, *Respiration proof*. Different proofs to which the organs of respiration of a new-born child are subjected, for the purpose of detecting whether it has or has not respired after birth; in other words, whether it was born alive or dead; —*Pneubiobion'tis*, *Pneustobios'copē*. These consist, chiefly, 1. In testing the lungs with water, for the purpose of seeing whether they are specifically heavier or lighter than that fluid. This is called *Docimasia Pulmonum hydrostatica*, or the *Hydrostatic Test*. If lighter, it would be some evidence that the fetus had respired: 2. In comparing the weight of the lungs with that of the whole body; the weight of lungs in which respiration has taken place being nearly twice as great. This is *Docimasia Pulmonum Stat'ica*, or *Plouquet's Stat'ic Test*; and, 3. By measuring the circumference of the thorax and lungs; and comparing their dimensions with those of an infant which has not respired. That is *Daniel's Test*. These tests, singly, afford only probable evidence; but when united, the deductions may be more conclusive.

DOCIMASIOLOGIA, (*docimasia*, and *logos*, 'a treatise.') *Docimasia*.

DOCKMACKIE, *Viburnum acerifolium*.

DOCK, (Sax. *doce*.) *Rumex*—*d.* Bitter, *Rumex obtusifolius*—*d.* Bloody, *Rumex sanguineus*—*d.* Bloody-veined, *Rumex sanguineus*—*d.* Blunt-leaved, *Rumex obtusifolius*.

DOCK CRESSSES, *Lapsana*.

DOCK, CURLED, *Rumex crispus*—*d.* Sour, *Rumex acetosus*, *R. crispus*—*d.* Sour, boreal, *Oxyria reniformis*—*d.* Spatter, *Nuphar advena*—*d.* Water, *Rumex hydrolapathum*—*d.* Wild, sharp-pointed, *Rumex acutus*—*d.* Yellow, *Rumex crispus*.

DOCKEN (Sc. and Prov.) ('dock.') *Rumex*.

DOCK'NA SURA, MIN'ERAL WATERS OF. This spring is in the Krapach mountains. The water contains sulphate of soda, carbonate of soda, chloride of sodium, carbonate of lime, silica, and iron.

DOCTEUR, Doctor, Physician.

DOCTOR, *Med'icus*, (*doctus*, 'learned; 'un homme qui devrait être docteur.') (F.) *Docteur*. Frequently applied to any one who practises medicine; although properly confined to him who has received his degree of Doctor of Medicine. See Physician.

'To Doctor' is sometimes used, vulgarly, for 'to treat with drugs.'

DOCTORS' STUFF, see Medicament.

DOCTRINA GALLIANA ('doctrine of Gall.') *Craniology*—*d.* Soteria, *Medicina*.

DOCTRINE HOMŒO-ORGANIQUE, *Homœopathy*.

DOCTRINE, MED'ICAL, *Doctrina Med'ica*. The principles or positions of any medical sect or master. Medicine has been too full of doctrines. One of the first was that of Herodicus of Selivra, who recommended gymnastic exercises in disease. The chief founders of doctrines have been Hippocrates, Serapion of Alexandria, Philinus of Cos, Herophilus, Asclepiades, Themison of Laodicea, Thessalus of Tralles, Soranus of Ephesus, Leonides of Alexandria, Athenæus of Attalia, Archigenes of Apamea, Agathinus of Sparta, Galenus, Paracelsus, John Baptist van Helmont, Sylvius de le Boe, Keill, Hamburger, Piteairne, H. Boerhaave, J. D. Stahl, Frederick Hoffmann, George Baglivi, Cullen, Darwin, John Brown, Beddoes, Girtanner, Reil, Rush, Rasori, and Broussais.

DOCTRINE, PHYSIOLOGICAL, *Broussaïs*.

DODDER, *Cuscuta glomerata*—*d.* American, *Cuscuta glomerata*—*d.* Thyme, *Cuscuta epithymum*.

DODECADACTYLITIS, (*dodecadactylon*, and *itis*.) *Duodenitis*.

DODECADACTYLON, (*δωδεκα*, 'twelve,' and *δακτυλος*, 'a finger.') *Duodenum*.

DODECAPHAR'MACUM, (*δωδεκα*, 'twelve,' and *φάρμακον*, 'a medicine.') An ancient name given to all medicines which consisted of twelve ingredients. See *Apostolorum Unguentum*.

DODECATHŒON, (*δωδεκα*, 'twelve,' and *θεός*, 'I put;' or from *θεός*, a god, after the twelve Grecian gods.) An antidote, consisting of twelve simple substances.

DODECATHEON, *Sanicula*—*d.* Plinii, *Pinguicula vulgaris*.

DODONÆA, (after R. Dodoens, of Leyden,) *Myrica gale*.

DODONÆA THUNBERGIANA. A shrub of the *Nat. Ord.* Sapindaceæ, which grows at the Cape of Good Hope. A decoction of the root is used as a gentle cathartic in fever.

DODRANS, *Spithama*.

DOGCHOAK, *Cynanche*.

DOG'DAYS, (F.) *Jours Caniculaires*. During these days, comprised between the 24th of July and the 23d of August, the temperature of the air in Europe is generally high and oppressive. They have received this name from the dogstar, *Canicula*, *Σείριος*, *Sirius*—a very brilliant star in the constellation of the great dog, which rises and sets, at this time, with the sun. It was formerly believed to be a period particularly unpropitious to health; that it was dangerous to purge during it; and other phantasies were indulged in regard to it.

DOGGRASS, *Triticum repens*.

DOGHIP, see *Rosa canina*.

DOGMAT'IC, (*δογματικός*; from *δογμα*, gen. *dogmares*, from *δοkein*, 'I think.') *Dogmat'icus*, (F.) *Dogmatique*. The name of an ancient medical sect; so called, because its members endeavoured, by reasoning, to discover the essence of diseases and their occult causes; whilst the *Empirics*, their rivals, confined themselves strictly to experience; i. e., to the observation of facts. The union of the two modes of investigation makes the rational physician. These sectarians are likewise called *Dog'matists*, and their doctrine *Dog'matism*. The founders of the sect were Hippocrates, Thessalus, Draco, and Polybius; and the most celebrated of its supporters were Diocles of Carysta, Praxagoras of Cos, Chrysippus of Soli, Herophilus, and Erasistratus.

DOGS' BANE, *Apocynum androsæmifolium*—*d.* Bane, bitter, *Apocynum androsæmifolium*.

DOGS' GRASS, *Triticum repens*.

DOGS' STONES, *Orechis mascula*.

DOGSTAR, *Canicula*.

DOGTREE, *Cornus Florida*.

DOGWOOD, *Cornus Florida*—*d.* Blueberried, *Cornus sericea*—*d.* Female, *Cornus sericea*—*d.* Florid, *Cornus Florida*—*d.* Great flowered, *Cornus Florida*—*d.* Jamaica, *Piscidia erythrina*—*d.* New England, *Cornus sericea*—*d.* Panicled, *Cornus paniculata*—*d.* Pond, *Cephalanthus occidentalis*—*d.* Round-leaved, *Cornus circinata*—*d.* Silky-leaved, *Cornus sericea*—*d.* Striped, *Acer Pennsylvanicum*—*d.* Swamp, *Cornus sericea*, *Ptelea trifoliata*—*d.* Virginian, male, *Cornus Florida*—*d.* White, *Cornus paniculata*.

DOIGT, *Digitus*—*d.* Auriculaire, Auricular finger—*d.* Hippocraticæ, Finger, Hippocratic—*d.* Milien, see *Digitus*—*d.* Petit, see *Digitus*—*d.* Supernuméraire, *Digitus supernumerarius*.

DOIGTIER, (from *doigt*.) *Digitale*, *Digitalis*—*d.* *d'Asdrubati*, see *Digitale*.

DOIGTS PALMÉS, *Dactylon*.

DOITRE, (Sc.), see *Dementia*.

DOKE, *Contusio*.

DOLABRA, (*dolare*, 'to hew.') *Ascia*, *Doloire*.

DOLIC, *Dolichos*.

DOLICHOCEPHALÆ (GENTES); (*δολιχός*, 'long,' and *κεφαλή*, 'head.') 'Longheads.' *Na-*

tions of men whose cerebral lobes completely cover the cerebellum—as the Kelts, Germans, negroes, &c.—Retzius.

DOLICHOCEPHALIA. Monstrosity, in which the head is abnormally long.

DOLICHOS (δολίχος, 'long,' as applied to the pods.) A genus of plants of the leguminous family, (F.) *Dolic*, *Dolique*. It includes a number of species, generally indigenous in India or America. The *Dolichos Lablab*, or *Lablab*, is found in Egypt. (Prospero Alpini.) Its fruit is eaten there, like the haricot with us. The *Dolichos Sinen'sis*, is eaten in China, and is stored up as a provision for long voyages. The *Dolichos Tuberosus* (F.) *Pois Patate* of Martinique, has tuberous roots of the size of both fists, and has the consistence and taste of the turnip. The *Dolichos Bulbosus* of the West Indies resembles the turnip;—and from the seeds of the *Dolichos Soja*, the Japanese prepare the well-known sauce *Soy*, which they term *Sooja*. The most important in medicine is the

DOLICHOS PRURIENS, *Dolichos*, *Stizolobium*, *Mucuna pruriens*, *Negretia pruriens*, *Cac'hage*, *Cac'hitch*, *Adsa'ria Pa'ta*. Order, Leguminosæ. (F.) *Pois à gratter*. The stiff hairs of the *Dolichos Pods*, called *Dolichos Pubes*, *D. Pruriens* *pubes*, *Dol'ichi Se'te Legu'minum*, *Lanu'go Sil'i-quæ hirsu'te*, *Muc'na* (Ph. U. S.), are the parts used in medicine. They excite an intolerable, prurient sensation, when applied to the skin; but do not irritate the mucous membrane over which they pass, when administered internally. The *Dolichos* is a mechanical anthelmintic, and is useful in cases of ascarides lumbricoides, and oxyures vermiculares. Dose, gr. v to x, of the pubes, in molasses.

Muc'na pruri'ta, a distinct species, but possessing similar properties, grows in the East Indies.

DOLIQUE, *Dolichos*.

DOLOIRE (F.), *As'cia*, *Dola'bra*, 'a carpenter's axe,' (from *dolare*, 'to hew.') A *Bandage en doloire* is one in which the turns are so placed, that each one covers two-thirds of that which is immediately beneath it. It has received the name *Doloire* from its representing the obliquity of the edge of the instrument whose name it bears. See *Bandage*.

DOLOR, gen. *Dolo'ris*, Pain.—d. Ani, Proctalgia.—d. Capitis, Cephalalgia.—d. Cephalicus, Cephalalgia.—d. Colicus, Colic.—d. Crucialis faciei, Neuralgia, facial.—d. Dentium, Odontalgia.—d. Dentium à stridore, Hæmodia.—d. Faciei, Neuralgia, facial.—d. Ischiadicus nervosus, Neuralgia femoro-poplitea.—d. Lenis, Hypodynina.—d. Mitis, Hypodynina.—d. Nephreticus, Nephralgia.—d. Pectoris externus, Pleurodynia.—d. Pudendorum, Pudendagra.—d. Rheumaticus et arthriticus, Rheumatism.

DOLORES AD PARTUM, (pl. of *Dolor*), Pains, labour.—d. Intestinorum, Colic.—d. Parturientis, Pains, labour.—d. Post partum, Pains, after.—d. Puerperarum, Pains, after.—d. Rodentes, Pains, gnawing.

DOLORIFIC, (*dolor*, 'pain,' and *facere*, 'to make.') Odynopoetic.

DOLP (Sc.), (Icel. *doef*.) Nates.

DOMBEY'A EXCEL'SA, (after Dombey, a French botanist.) *Arauc'ria Dombey'i*. A tree of Chili, which furnishes *Dombeya turpentine*; a glutinous, milky-looking fluid of a strong odour and taste.

DOMESTIC, *Domes'ticus*, (*domus*, 'a house.') The term *Domestic* or *Popular Medicine* has been given to treatises written for the purpose of enabling those who are not of the profession to treat diseases, which may occur in their families, without the necessity of calling in a physician. The term, likewise, signifies—medicine, when thus

practised. It is probable, that such works have been attended with mischievous as well as advantageous results.

DOMINARUM AQUA, ('ladies' water.') A medicine described by Myrepsus, which he considered emmenagogue.

DOMPTE-VENIN, ('poison-tamer, alexipharmic.') *Asclepias vincetoxicum*.

DOMUS LEPROSARIA, ('a house for the leprous.') *Laderrie*.

DONDO, Albino.

DONESIS, (*δονεσις*.) Agitation.

DOORWEED, Polygonum.

DORA, (*δορα*.) Cutis.

DORADILLA, *Asplenium ceterach*.

DORCADIZON, (*δορκαδίζω*, 'to leap like a roe'—*δορκας*.) Caprizans.

DOREA, Hæmeralops.

DOREMA AMMONIACUM, gen. *Dore'matis*, see Ammoniac, Gum.

DORMITATIO, (*dormitare*, *dormitatum*, 'to be sleepy.') Somnolency.

DORMITO, (*dormire*, 'to sleep.') Sleep.—d. Lucumoriana, see Lucumorianus.

DORONIC, *Doronicum pardalianches*—d. d'Allemagne, *Arnica montana*.

DORONICUM ARNICA, *Arnica montana*—d. Cordatum, D. Pardalianches—d. Germanicum, *Arnica montana*—d. Officinale, D. Pardalianches—d. Oppositifolium, *Arnica montana*.

DORONICUM PARDALIANCHES, *D. Romanum* seu *Corda'tum* seu *Officina'le*, *Ro'man Leopard's Bane*, (F.) *Doronic*, *Doronique*, *Mort aux Panthères*. The root of this plant resembles *Arnica Montana* in its properties.

DORONICUM PLANTAGINIS FOLIO, *Arnica montana*—d. Romanum, D. Pardalianches.

DORSAD, see Dorsal Aspect.

DOR'SAL, *Dorsa'lis*, (from *dorsum*.) *Notia'us*, *No'tal*, *Ter'gal*. Relating to the back of the body, or of one of its parts; as the *Dorsal vertebrae*, nerves, &c.; *Dorsal artery of the tongue*, penis, &c.; *Dorsal region of the foot*, hand, &c.; *Dorsal Consumption*, &c.

DORSAL ASPECT. According to Dr. Barclay, an aspect towards the *dorsum* or *backbone*. *Dorsad* is used by the same writer adverbially, to signify 'towards the dorsal aspect.'

DORSAL LONG, *Longissimus dorsi*.

DORSALIS, Dorsal.

DORSALIS PE'NIS (Nervus.) The branch of the internal pudic nerve, which is distributed to the upper part of the male organ; and to the clitoris of the female.

DORSCH, see *Oleum jecoris aselli*.

DORSE, see *Oleum jecoris aselli*.

DORSO-COSTAL, *Serratus posticus superior*—d. *Lombo-costal*, *Serratus posticus inferior*—d. *Lombo-sacro-humeral*, *Latissimus dorsi*—d. *Sacro-cromien*, *Trapezius*—d. *Trachélon-occipital*, *Complexus*.

DORSTENIA BRASILIENSIS, (after Dr. Dorsten, a botanist of Marburg.) *Ca-a'pia*.

DORSTENIA CONTRAYER'VA, *D. Houstonii*, *Contrayer'va*, *Drake'na*, *Cype'rus longus odoratus* seu *Peruv'anus*, *Bezoardica Ra'diz*, *Lis'bon Contrayer'va* (F.) *Racine de Charcis ou de Drake* (Drake) ou *des Philippines*. Order, *Urticæ* *Sax. Syst.* Tetandria Monogynia. A plant of South America, whose root, *Contrayer'va* (Ph. U. S.), is aromatic, bitter, and astringent. It has been given as a tonic, stimulant and sudorific. Dose, gr. xij to ʒss.

DORSTENIA CORDIFOLIA, D. *Brasiliensis*—d. *Placentoides*, D. *Brasiliensis*—d. *Vitella*, D. *Brasiliensis*.

DOR'SUM, *No'tus*. The back. *Metaph'renon*, *Ter'gum*, (Prov.) *Rig*, (F.) *Dos*. The posterior part of the trunk, extending from the inferior and posterior region of the neck as far as the

loins. The back of the foot, *Dorsum pedis*, is the upper part of that organ, opposite the sole: the back of the hand, *Dorsum manus*, the part opposed to the palm. In the same sense, we say *Dorsum penis*, *Dorsum nasi*, for the upper part of those organs.

DORSUM, Vertebral column—d. Ephippii, see *Sella turcica*—d. Manús, *Opisthenar*.

DOS, Dorsum.

DOSAGE, (from dose,) Posology.

DOSE, *Dosis*, (*dosís*, from *didomi*, 'to give,') *Præbium*, (F.) *Prise*. The quantity of any substance, which ought to form part of a compound medicine, or ought to be exhibited singly, to produce a desired effect. Many circumstances influence the doses of medicine. Women require smaller doses, as a general principle, than men. *Habit* has a great effect, as well as *climate*, *age*, and *idiosyncrasy*: all these, and other circumstances, must be taken into account; and every general rule on the subject will be found to have numerous exceptions. Some of the mechanical physicians laid it down as a rule, that the doses of medicines must always be as the square of the constitution!—A matter not easy of calculation.

The following Tables will exhibit an approximation, but only an approximation, to the proper doses (according to age) of most substances.

TABLE OF DOSES ACCORDING TO AGE.

Age 24	Let the full dose be	1	1
18	will require	2-3ds.	2 drachm.
14	- - - -	half.	$\frac{1}{2}$ drachm.
7	- - - -	1-3d.	1 scruple.
4	- - - -	1-4th.	15 grains.
3	- - - -	1-6th.	10 grains.
2	- - - -	1-8th.	8 grains.
1	- - - -	1-12th.	5 grains.

The table of doses, according to age, recommended by Dr. Thomas Young, differs in some respects from the above. It, also, affords a general approximation. His rule is, that

For children, under twelve years of age, the doses of most medicines must be diminished in the proportion of the age to the age increased by twelve:

Thus, at two years, to 1-7th; i.e. $1-7th = \frac{2}{2+12}$

At twenty-one the full dose is given.

DOSES, BROKEN, (F.) *Doses refractées*. When an agent is given in small portions it is said to be in broken doses,—*refractis dosibus*, (F.) à *Doses refractées*.

DOSIA POWDER, see Powder, dosia.

DOSIOLOGIA, (*dosís*, and *logos*, 'a discourse,') Posology.

DOSIS, (*dosís*,) Dose.

DOSSIL, (Old. F. *dosil*, 'a stopple,') Bourdonnet.

DOTAGE, ([F.] *radoter*, [D.] *dutten*, 'to rave,') See Dementia.

DOTHIEU, (*δοθιου*,) Furunculus.

DOTHIENTERIA, (*dothien*, *εντερον*, 'intestine,' and *itis*,) Dothienteritis.

DOTHIENTÉRIE, Dothienteritis.

DOTHINENTERIA, Dothienteritis.

DOTHINENTERITIS, properly *Dothiententeritis*, *Dothienteritis*, *Dothienteritis*, *Enteritis pustulosa*, *Enterodothienteritis*, *Helcenteritis*, *Follicular Gastroenteritis*, (F.) *Dothienterite*, *Dothientérie*, *Dothientérie*, *Dothientérie*. An inflammation and ulceration of the glands or follicles of Peyer and Brunner, which Bretonneau considered to be the essence of a large class of fevers, particularly of those of the typhoid character. See Typhus.

DOTHION, (*δοθιον*,) Furunculus.

DOTTRINA MEDICA ITALIANA, ('Italian medical doctrine,') *Contrastimulus*, (doctrine of.)

DOUBLE, Ingluvies.

DOUBLE CONSCIOUSNESS, see Consciousness, double.

DOUBLE HEARING, *Diplecoia*, *Dittecioia*, *Paracousis duplicata seu imperfecta*. The action of the one ear unaccordant with that of the other: sounds heard doubly, and in different tones or keys.

DOUBLE-MONSTERS, see Duplication.

DOUBLE-QUARTE, see Quartan.

DOUBLE-QUOTIDIENNE, see Quotidian.

DOUBLE-TIERCE, see Tertian.

DOUBLING GAP, MIN'ERAL WA'TERS OF. They are situated in Cumberland County, Pennsylvania, eight miles from Newville. They contain hydrosulphuric acid, carbonates of soda and magnesia, sulphates of soda and magnesia, and chloride of sodium. A chalybeate spring contains carbonate of iron, sulphate and carbonate of magnesia, and chloride of sodium.

DOUCE-AMÈRE, ('sweet bitter,') *Solanum dulcamara*.

DOUCHE (F.), (modern Latin, *ducia*, (I.) *docciare*, 'to pour,') *Cataclysmus*, *Douse*. This term is applied to a column of fluid, of a determinate nature and temperature, let fall upon the body. Pump'ing is a variety of the *Douche*. In using this kind of bath, the fluid is directed upon the part on which we are desirous of acting. The *douches descendantes* are those in which the fluid falls from a height,—the *douches ascendantes*, those administered in diseases of the uterus,—the *douches horizontales*, where the fluid is impelled horizontally, &c. They may be cold or warm, according to circumstances. The apparatus consists of a reservoir of water having a pipe or plug, by means of which the water can be directed as the practitioner may desire. The *Douche* communicates a considerable and peculiar shock to the nervous system; and is one of the most successful means for taming the furious maniac. It is, also, useful in chronic rheumatism, stiff joints, &c.

Douches of air are occasionally used, as in cases of obstruction of the Eustachian tube by mucus. They are sent from an air-press—of which Deleau and Kramer have invented one each—through a catheter introduced through the nose into the tube.

DOUCHE ÉCOSSAISE, *Douche*, transition—d. Scotch, *Douche*, transition.

DOUCHE, TRANSITION, Scotch *douche*, (F.) *Douche Écossaise*. A *douche*, which consists in the successive use of hot and cold water.

DOUGLAS'S CUL-DE-SAC, (after Dr. James Douglas, the obstetrician and anatomist,) see Uterus.

DOULEUR, (from *dolor*,) Pain, d. de *Coté*, *Pleurodynia*—d. des *Dents*, *Odontalgia*—d. de l'*Estomac*, *Cardialgia*—d. *Névralgique de l'Estomac*, *Cardialgia*—d. *Pulsative*, see Throbbing.

DOULEURS, (pl. of *Douleur*,) Pains, labour—d. *Conquassantes*, see *Conquassant*.

DOULOUREUX, Painful.

DOUNT (Sc.), (onomatopœia,) Palpitation.

DOUP (Sc.), ([Icel.] *doef*,) Nates.

DOUSE, *Douche*.

DOUVE, *Distoma hepaticum*—d. du *Foie*, *Distoma hepaticum*—d. *Petite*, *Ranunculus flammula*.

DOWN, see Attack.

DOWP (Sc.), ([Icel.] *doef*,) Nates.

DOYLE, Strabismus.

DRACHION, Pugillus.

DRACHM, *Drachma*, (*δραχμη*,) *Dram*, (F.) *Gros*. The ancient name of a piece of money, weighing the eighth part of an ounce. At the present day it is used for this weight.

DRACO, (δράκων, gen. δράκοντος, 'a dragon,') in composition, a dragon or large serpent.

DRACON MITIGATUS, Hydrargyri submuriatis-d. Sylvestris, Achillea ptarmica.

DRACOCEPHALUM CANARIENSE, *D. Moldavicum*, (*draco*, and κεφαλή, 'head,' from the shape of the corolla.) *Melissa Turcica* seu *Canariensis*, *Cedronella triphylla*, *Alpinia Balsamum*, *Turkey* or *Canary Balsam*, *Balm of Gil'ead Tree*, (F.) *Mélinse* de *Moldavie*. Ord. Labiatae. A Turkish and Siberian plant, which has an aromatic taste, joined with an agreeable flavour. It has been used as a tonic.

DRACONTHÆMA, (*draco*, and αἷμα, 'blood,') see *Calamus rotang*.

DRACONTIASIS, see *Dracunculus*.

DRACONTIUM, *Dracunculus*. See, also, *Dracunculum foetidum*.

DRACONTIUM FOETIDUM, (from *draco*), *Ictodes foetidus*, *Symplocarpus foetidus*, *Pothos foetida* seu *Putorii*, *Arum Americanum betulaefolium*, *Spathyema foetida*, *Skunk-cabbage*, *Skunkweed*, *Polecatcollard*, *Stinkcollard*, *Collard*, *Itchweed*, *Stinkpoke*, *Swampcabbage*, *Polecatweed*, *Hellebore*, *Ellebore*, *Irish cabbage*. Ord. Aroideæ or Araceæ. *Sex. Syst.* Tetrandria Monogynia. This indigenous plant, as some of its names import, is extremely fetid. The property on which its medical virtues are dependent, resides in a volatile principle, which is impaired by long keeping, especially in powder. Dose, of the dried root, *Dracontium*, (Ph. U. S.,) ten to twenty grains. It resembles asafœtida and other fetid gums in its properties; in other words, belongs to the class of reputed antispasmodics.

DRACONTIUM ANGUSTIPATHA, *Symplocarpus*, *Angustipatha*, *Narrouspathæ* *Skunkcabbage*, is possessed of similar properties.

DRACUNCULUS, (dim of *draco*), *D. Gordius*, *Dracontium*, *Veina Mediana Arabum*, *Veina* seu *Gordius Medinensis*, *Medinagordius*, *Vermiculus Capillaris*, *Filaria Medinensis* seu *Guineensis*, *Malaria Dracunculus* seu *Gordii*, *Helminthioncus Medinensis*, *Muscular Hairworm*, *Bichios*, *Bicho*, *Guinea Worm* or *Threadworm*, (F.) *Dragonneau*, *Ver de Guinée*, *Ver Filaire*, *V. de Médine*, *V. cutané*, *Veine ou Filare de Médine*. A genus of worms, frequently met with in Indian and African climes. They are characterized by a filiform body, and are smooth and nearly of equal thickness throughout. The Guinea worm, when small, insinuates itself through the cutaneous pores, and penetrates into the areolar membrane and between the muscles; especially between those of the lower limbs, where it occasions a tumour like a boil, which subsequently suppurates, and the head of the worm appears and emerges gradually. The head must then be seized, and the worm be cautiously rolled round a small cylinder of linen or other substance. Care must be taken not to break it, as great pain and obstinate suppuration might be the consequence.

Considerable obscurity rests on this subject. Some even deny that the cases of *Dracunculus*, on record, are really those of worms.

The morbid phenomena induced by the *dracunculus* have been termed *Dracontiasis*, (F.) *Dracontiae*.

DRACUNCULUS, *Artemisia dracunculus*—d. Polyphyllus, *Arum dracunculus*—d. *Pratensis*, *Achillea ptarmica*.

DRAGÉES (F.), (corr. from τραγναρα, 'sweetmeats?') Almonds or dried preserves, covered with white sugar; *Sugarplums*. Used in pharmacy for sugar-coated medicines.

DRAGÉES DE KEYSER. A pharmaceutical preparation, formerly much celebrated in syphilis. It was composed of acetate of mercury, mianna, starch, mucilage, and gum tragacanth.

DRAGMA, gen. *Dragmatia*, (δράγμα,) *Pugillus*. DRAGMIS, (δράγμις,) *Pugillus*.

DRAGON (F.), *Dragon*. Some authors have given this name to opaque spots on the cornea, and to cataract.

DRAGON CLAW, SCALY, *Pterospora andromedea*—d. Green, *Arum dracontium*—d. Root, *Arum triphyllum*, *Pterospora andromedea*—d. Turnip, *Arum triphyllum*.

DRAGON'S BLOOD, see *Calamus rotang*.

DRAGONNEAU, (dim. of *dragon*,) *Dracunculus*.

DRAIN'AGE. The act of depriving parts gradually of their redundant humidity. In Surgery, it means the gradual drawing off of purulent fluids from deep-seated abscesses, by means of hollow sounds,—the *Drainage chirurgicale* of Chassaignac.

DRAIN'ING, (Sax. *dræhningean*, or from (F.) *trainer*, 'to draw,') *Dessicat'io*, (F.) *Dessèchement*. The act of drawing off the water from marshes, for the purpose of destroying the unhealthy emanations which proceed from them. It is a well known fact, that from marshes arise certain emanations or miasmata, with the nature of which we are, however, totally unacquainted, but which are the fertile source of intermittents and other diseases. Draining the soil and converting it into arable land changes its character, and the malaria ceases to be given off. It has happened, however, that although in some such situations intermittents have been got rid of, consumptions have taken their place.

DRATTSCH WATER, Godesberg, mineral waters of.

DRAKENA, (after Sir Francis Drake,) *Dorstenia contrayerva*.

DRAM, Drachm; also (Sc.) *Melancholic*.

DRAP FANON, see *Fanon*.

DRAPEAU (F.), from *drap*, 'cloth,') *Vexillum*. A bandage, used in wounds, &c., of the nose, to keep the dressings *in situ*. It is composed of a small triangular compress, having two openings at its inferior part, corresponding to those of the nostrils. It is fixed by means of a bandage, passing from its superior angle over the head, and another passing round the head, under the orbits, so as to join the first at the nape of the neck. Also, *Pterygion*.

DRASTIC, *Dras'ticus*, (δραστικός, from δραω, 'I am active,') *Cenot'ic*, (F.) *Drastique*. Active. An epithet given to purgatives, which operate powerfully; as *elaterium*, *oil of croton*, &c.

DRAUGHT, (past participle of Sax. *dragan*, 'to draw,') *Haus'tus*. A term also applied by nurses to the sudden rush of blood to the mammae, occasioned in the mother by the sight, or even thought, of her infant, and which occasions a greatly increased secretion of milk.

DRAUGHT, BLACK, see *Infusum sennæ compositum*.

DRAWING, see *Digestives*.

DRAX, (δραξ,) *Pugillus*.

DREAM, ([G.] Traum,) *Somnium*—d. *Waking*, see *Hallucination*.

DRÊCHE, Malt.

DREGISTER, Druggist.

DREGS, ([G.] Dreck 'dirt,') *Feculence*.

DRENCH, (Sax. *dræncean*, 'to soak,') To give liquid physic by force. A liquid medicine, given by violence, is called "a drench."

DREPANOIDES, (δρεπανοειδής, from δρεπανον, 'a falx or scythe, and αἶδος, 'resemblance,') *Falciform*.

DRESS, Vestitus.

DRESS'ER, ([F.] dresser, 'to put right,') An hospital assistant, whose office it is to dress wounds, ulcers, &c. He corresponds in function to the *Élève externe* of the French hospitals.

DRESS'ING, *Cu'ra*, *Cura'tio*, (F.) *Pansement*.

The methodical application of any remedy or apparatus to a diseased part. Also, the remedy or apparatus itself.

DRIBURG, MIN'ERAL WA'TERS OF. At the small town of Driburg, near Pyrmont, in Germany, there are nine springs, which are rich in saline ingredients, iron, and carbonic acid gas. Their action on the economy is like that of Pyrmont water. They contain chloride of sodium, sulphate of magnesia, sulphate of lime, carbonate of magnesia, carbonate of iron, chloride of calcium, and chloride of magnesium.

DRIMYPHAGIA, (δριμυφαγια, from δριμυς, 'acid,' 'aromatic,' and φαγω, 'I eat.') An aromatic and exciting diet.

DRIMYS WINTERI, (δριμυς, 'sharp, acid,') *Winters aromaticæ*.

DRINK, (Sax. *drincan*,) *Po'ma*, *Po'tio*, *Po'tus*, *Be'verage*, (F.) *Boisson*. Every liquid introduced into the alimentary canal for the purpose of repairing the loss of the fluid parts of the body. The necessity for its use is indicated by the sensation of thirst. Fluid, taken during a meal, aids in the digestion of the food. Some drinks are exciting and tonic, as the different varieties of beer, wine, and spirits, which we use at table. In a therapeutical point of view, drinks are used to appease the thirst which prevails in febrile affections, or to act as diluents in those and other cases.

The ordinary drinks, according to their material composition, are—1. *Water*, spring water, river water, well water, &c.—2. *Juices and infusions of Animal and Vegetable substances*, lemon juice, currant juice, whey, tea, coffee, mattee, &c.—3. *Fermented Liquors*, wines, ale, beer, cider, perry, &c.—4. *Spirituous Liquors*, brandy, alcohol, ether, kirschwasser, rum, arack, gin, whisky, ratafias, cordials, &c.

DRIVELLING, (from 'drip.' [?]) Slaverling.

DROAT (Prov.), Throat.

DRODDUM (Prov. and Sc.), Nates.

DROGUE, Drug.

DROGUIER (F.). A collection of different simple medicinal substances. A methodical collection of specimens of drugs.

DROGUISTE, Druggist.

DROICH (Sc.), Nanus.

DROIT, (from *dirigere*, *directum*, 'to make straight,') *Rectus*—*d. de l'Abdomen*, *Rectus abdominis*—*d. Antérieur de la cuisse*, *Rectus femoris*—*d. Antérieur de la tête*, *Rectus capitis internus minor*—*d. Antérieur de la tête, grand*, *Rectus capitis internus major*—*d. Externe de l'œil*, *Rectus externus oculi*—*d. Inférieur de l'œil*, *Rectus inferior oculi*—*d. Interne de la cuisse*, *Gracilis*—*d. Interne de l'œil*, *Rectus internus oculi*—*d. Latéral de la tête*, *Rectus capitis lateralis*—*d. Postérieur de la tête, grand*, *Rectus capitis posticus major*—*d. Postérieur de la tête, petit*, *Rectus capitis posticus minor*—*d. Supérieur*, *Rectus superior oculi*.

DROP, (Sax. *droppa*,) *Gutta*—*d. Ague*, tasteless, *Liquor arsenicalis*—*d. Black*, *Guttæ nigræ*—*d. Red*, *Ward's*, see *Vinum antimonii tartarizati*—*d. Serene*, *Amaurosis*.

DROPACISMUS, (δρῶπακισμος,) see *Depilation*.

DROPAX, (δρῶπαξ,) see *Depilatory*.

DROPPED HANDS, see *Hands*, *dropped*.

DROPS, (pl. of *Drop*,) see *Guttæ*—*d. Abbé Rousseau's*, *Laudanum Abbatis Rousseau*—*d. Anodyne*, *Liquor morphinæ acetatis*—*d. Antiscorbutic*, *Marsden's*, see *Infusum gentianæ compositum*—*d. Antivenereal*, see *Tinctura ferri muriatis*—*d. Bateman's*, see *Bateman's pectoral drops*—*d. Chamomile*, see *Anthemis nobilis*—*d. Dutch*, *Balsam of sulphur*, *Balsam of Turpentine*—*d. Fit*, *Spiritus ammoniæ fetidus*, see *Fuligo*—*d. Golden*, *de la Motte's*, *Tinctura seu alcohol sulfurico-æthereus ferri*—*d. Green's*, see *Liquor Hydrargyri oxy-muriatis*—*d. Jesuit's*, *Tinctura ben-*

zoini composita—*d. Lavender*, *Spiritus Lavandulæ compositus*—*d. Nitre*, *Spiritus atheris nitrici*—*d. Norris's*, see *Antimonium tartarizatum*—*d. Norton's*, *Liquor Hydrargyri oxy-muriatis*—*d. Pectoral*, *Pectoral drops*, *Bateman's*—*d. Peppermint*, *Pastilli de menthâ piperitâ*—*d. Soot*, see *Fuligo*—*d. Wade's Tinctura benzoini composita*.

DROPS, WARBURG'S. A secret preparation used in Demarara under the name of *Warburg's Fever Drops*. See *Bebeera*.

DROPS, WHITE, WARD'S, see *Hydrargyri nitras*.

DROPSICAL, *Hydropic*.

DROPSIED, *Hydropic*.

DROPSY, *Hydrops*—*d. of the lower Belly*, *Ascites*—*d. of the Brain*, *Hydrocephalus*.

DROP'SY, CARDIAC. *Hy'drops cardi'acus*, (F.) *Hydropsiæ cardiæque*. Dropsy, dependent on disease of the heart.

DROPSY OF THE CELLULAR MEMBRANE, *Anasarca*—*d. of the Chest*, *Hydrothorax*—*d. of the Eye*, *Hydrophthalmia*.

DROPSY, FIBRINOUS. Dropsy in which the effused fluid contains fibrin.

DROPSY, GENERAL, *Anasarca*—*d. of the Head*, *Hydrocephalus*.

DROPSY, HEPAT'IC, *Hy'drops hepaticus*, (F.) *Hydropsiæ hépatique*. Dropsy, dependent on disease of the liver.

DROPSY OF THE LACHRYMAL SAC, *Fistula lacrymalis*, *Lachrymal hernia*.

DROPSY, OVARIAN, *Hydrooarian*—*d. of the Pericardium*, *Hydropericardium*—*d. of the Peritoneum*, *Ascites*—*d. of the Pleura*, *Hydrothorax*.

DROPSY, RENAL, *Hy'drops renalis*, (F.) *Hydropsiæ rénale*. Dropsy, dependent on disease of the kidney.

DROPSY, WATER OF, *Serum of Serous membranes*—*d. Wind*, *Emphysema*.

DROPSY OF THE WOMB, *Hydrometra*.

DROWORT, (from *drop* and *wort*,) *Spiræa filipendula*—*d. Hemlock*, *Ænanthe*—*d. Water*, *Hemlock*, *Ænanthe*—*d. Western*, *Gillenia trifoliata*.

DROPYK (Prov.), *Hydrops*.

DROSE'RA ROTUNDIFO'LIA, (δρῶσερος, 'dewy,' δρῶος, 'dew,' the leaf glands exuding a clear fluid like dew-drops.) The systematic name of the *Sun'dew*, *Ros soalis*, *Drosobotanon*, *Dro'sium*, *Rorell'ia*, (F.) *Rosalis*, *Rosée du soleil*, *Herbe à la rosée*. *Ord.* *Droseraceæ*. This plant has a bitter, acrid, and caustic taste. It has been used as a rubefacient, and to destroy warts and corns. It has, also, been regarded as a pectoral.

DROSISTICON, see *Bosa*.

DROSIMUM, *Drosera rotundifolia*.

DROSOBOTANON, (δρῶος, 'dew,' and βῶταν, 'plant,') *Drosera rotundifolia*.

DROSOMELI, (δρῶος, 'dew,' and μελι, 'honey,') *Fraxinus ornus*.

DROW (Sc.), *Syncope*.

DROWNING, ASPHYXIA BY, see *Asphyxia* by submersion.

DRUG, *Medicamentum*, (F.) *Droque*. (*Ménage* derives it from *droga*, and this from the Persian *droa*, 'odour,' because many drugs have a strong odour. It is, doubtless, from Teutonic *trocken*, Sax. *drigan*, 'to dry,') A name ordinarily applied to simple medicines, but, by extension, to every substance employed in the cure of disease.

DRUG. To prescribe or administer drugs. Most commonly, perhaps, to dose to excess with drugs. One who so doses is sometimes called "a druggier." "To drug," also means to tincture with some medicinal article.

DRUG DISEASE. A morbid condition, which is—or is presumed to be—caused or kept up by the administration of drugs.

DRUG GRINDER, see *Pulverization*—*d. Mill*, see *Pulverization*.

DRUGGER, Druggist, see Drug.
DRUGGING, Pharmacomania.
DRUGGIST, *Drug'ger*, *Drug'ster*, (Prov.) *Dre-gister*, *Pharmacopola*, *Materialista*, *Seplasia'rius*, *Pharmacopaeus*, (F.) *Droguiste*. One who sells drugs.

DRUGSTER, Druggist.
DRUM (Sc.), Melancholic—d. of the Ear, Tympanum.

DRUMBELLY, Tympanites.

DRUMSIEVE, see Cribration.

DRUNKENNESS, Temulentia.

DRUNKEWORT, Nicotiana tabacum.

DRY'ING, (Sax. *drigan*, 'to dry.') *Desicca'tio*, *Sicca'tio*, *Desicca'tio*, *Esicca'tio*, *Xeran'sis*, *Anan'sis*, same etymon as Drug, (F.) *Dessiccation*. Evaporation or removal of the superfluous humidity in a body.

DRYING OF PLANTS, *Desicca'tio herbarum*, *Desiccation of Plants*. Those which are very succulent should be dried quickly. They must be spread upon lattice work, covered from the light, and exposed to the heat of the sun or to that of a stove, not exceeding 110° Fahrenheit. The drying of less succulent plants can be effected at a lower temperature. *Flowers* must be dried very carefully, excluding light. *Seeds* are dried in a place where there is a free current of air. *Pulpy fruits* may be exposed to a gentle heat in a stove, which may be gradually elevated. *Roots* may also be dried in the stove: the tuberous require to be cut in slices.

DRY NURSE, see Nurse.

DRYOBALANOPS CAMPHORA, see Camphor.

DTHOEK, Framboesia.

DUALITY, *Dual'itas*, (duo, 'two.') The state or quality of being two.

DUALITY OF THE MIND OR BRAIN. As the organ consists of two hemispheres, they have been regarded by some as separately and distinctly concerned in the mental and moral manifestations.

DUALM (Sc.), Syncope.

DUCKFOOT, see Kylosis.

DUCKLEGGED. Having short legs like a duck.

DUCKSFOOT, *Podophyllum montanum*.

DUCT, *Ductus* — d. Alimentary, Canal, alimentary, Thoracic duct.

DUCT OF BELLINI, Uriniferous tube.

DUCT, NASAL, Lachrymal duct—d. Pancreatic, see Pancreas.

DUCT OF PECQUET, Thoracic duct.

DUCT, SPERMATIC, Deferens vas.

DUCT OF STENO, see Parotid.

DUCT, VITELLINE, see Vesicula umbilicalis.

DUCT OF WIRSUNG, see Pancreas.

DUCTIO PRÆPUTII, Masturbation.

DUCTLESS GLANDS, see Ganglion.

DUCTOR CANALICULATUS, Gorget — d. Urinæ, Ureter.

DUCTS OF MÜLLER. Two canals, which descend at the outer border of the Wolffian bodies, of which the vesicula prostatica is the relic in the male, as the Fallopian tubes are in the female.

DUCTULI RECTI, (dim. of *Ductus*), see Rete testis.

DUCTUS, gen. *Ductûs*, (*ducere*, *ductum*, 'to lead,') Canal, Meatus — d. Arantii, Canal, venous—d. Arteriosus, Arterial duct—d. Auris palatinus, Eustachian tube.

DUCTUS BARTHOLINI'NUS, BARTHOLIN'S *Duct*, (F.) *Canal de Bartholin*. The excretory duct of the sublingual gland.

DUCTUS BELLINIANI, Uriniferous tubes — d. Biliarii, Pori biliarii—d. Biliiferi, Pori biliarii—d. Botalli, Arterial duct—d. Choleodochus, Choleodoch duct—d. Chyliferus, Thoracic duct—d. Cibarius, Canal, alimentary—d. Ejaculatorii, Ejaculatory ducts—d. Excretorius, Excretory duct—d. Fer-

reini, Ferrein, canal of—d. Hepaticus, Hepatic duct—d. Incisivus, Palatine duct (anterior)—d. Intestinalis, Canal, alimentary—d. Lacrymalis, Lachrymal duct—d. Lacrumalis, Lachrymal duct—d. Lacteus, Thoracic duct—d. Lactiferi, Lactiferous vessels—d. Nasalis orbitæ, Lachrymal or nasal duct—d. Nasalo-palatinus, Palatine duct (anterior)—d. ad Nasum, Lachrymal or nasal duct—d. Nutritii, Canals, nutritive—d. Omphalo-mesentericus, see Vesicula umbilicalis—d. Pancreaticus, see Pancreas—d. Pecqueti, Thoracic duct—d. Pneumaticus, see Air-bladder—d. Punctorum lacrymalium, Lachrymal ducts—d. Riviniani, see Sublingual gland—d. Rorifer, Thoracic duct.

DUCTUS SALIVARIS INFÉRIOR, *D. Whartoniana*, (F.) *Canal de WARTON*. The excretory duct of the submaxillary gland.

DUCTUS SALIVARIS SUPÉRIOR, *D. Stenoniana*, *STE'NO'S Duct*, (F.) *Canal de STÉNON*. The excretory duct of the parotid gland which opens into the mouth opposite the second upper molar tooth.

DUCTUS SEMICIRCULARES LABYRINTHI, Semicircular canals—d. Serosi, Lymphatic vessels—d. Spirales cochleæ, Scala of the cochlea—d. Stenonianus, *Ductus salivaris superior*—d. Thoracicus, Thoracic duct—d. Urinæ, Ureter—d. Urinari, Urethra—d. Uriniferi Bellini, Uriniferous tubes—d. Varicosi uteri, Tubæ Fallopiæ—d. Venosus, Canal, venous—d. Vitellarius, see Vesicula umbilicalis—d. Vitello-intestinalis, see Vesicula umbilicalis—d. Waltheriani, see Sublingual gland—d. Whartonianus, *Ductus salivaris inferior*—d. Wirsungianus, see Pancreas.

DUELECH, Dulech.

DUELLA. The ancient name of a weight, which was equivalent to eight scruples.

DUELLUM ('war,') *VENEREUM*, Coition.

DUG, (Sw. *dugga*, 'to give suck,') Nipple.

DULCAMARA, (*dulcis*, 'sweet,' and *amarus*, 'bitter,') *Solanum dulcamara*—d. Flexuosa, *Solanum dulcamara*.

DULCEDO AMORIS, (gen. *Dulce'dinis*) ('sweetness of love,') Clitoris—d. Sputatorium, see Salivation.

DUL'LECH, *Du'lelech*. A term employed by Paracelsus and Van Helmont to designate a pretended tartarized substance, which forms in the human body, and produces acute pain, accompanied with great danger.

DULL, (Sax. *dele*,) Deaf.

DULSE, *Rhodomela palmata*—d. Pepper, *Laurientia pinnatifida*.

DUMB, (Sax. *dumb*,) see Mutitas.

DUMBNESS, Mutitas.

DUNBLANE, MINÉRAL WATERS OF. The springs of Dunblane, in Scotland, contain sulphate of soda, chlorides of sodium and calcium, and carbonate of iron.

DUNG, (Sax. *duns*,) Fimus—d. Cow's, Fimus Vaccæ—d. Goose, Fimus anseris—d. Stone-horse, Fimus equinus.

DUNGA, Dengue.

DUNT (Sc.), (onomatopœia,) Palpitation.

DUODENAL, *Duodena'tis*, (F.) *Duodenal*. Relating or appertaining to the duodenum.

DUODENITIS, *Dodecadaetyli'tis*, (F.) *Duodénite*, (*duodenum* and *itis*.) Inflammation of the duodenum, characterized by white tongue, bitter taste, anorexia, fulness and tenderness in the region of the duodenum, and often yellowness of skin, along with the ordinary signs of febrile irritation.

DUODENUM, (*duodeni*, 'twelve,') *Ventriculus Succenturiatus*, *Ec'physis* seu *Appen'dix* seu *Processus Ventriculi*, *Portonarium*, *Dodecadactylon*. The duodenum is the first part of the intestinal canal, commencing at the pyloric orifice of the stomach, and terminating in the jejunum.

Its length was estimated, by the ancients, to be about twelve fingers' breadth. As it is only partially covered by the peritoneum, it is susceptible of considerable dilatation; hence its name *Ventriculus succenturiatus*. In the duodenum, *chylification* takes place after the admixture of the biliary and pancreatic fluids with the chyme.

DUOSTERNAL. Béclard thus names the second osseous portion of the sternum, which corresponds to the second intercostal space.

DUPLICATION, (*duo*, 'two,' and *plicare*, *placatum*, 'to fold.') *Duplicatio*, *Duplicitas*, *Diplasiasmus*, *Diploisis*, (F.) *Duplicité*. A species of malformation or monstrosity, characterized by the parts concerned being doubled. See *Diplogenesis*.

DUPLICATION OF THE FÆTUS, Evolution, spontaneous.

DUPLICATION, *Duplicatura*, *Reflexion*. The folding or reflection of a membrane upon itself; as *Duplicature* or *reflection of the pleura*, *peritoneum*, &c.

DUPLICITAS MONSTROSA, (*duplex*, gen. *duplicitis*, 'double.') See *Diplogenesis*.

DUPLICITE, Duplication.

DUPONDIIUM. A weight of four drachms.

DUR, (from *durus*.) Hard.

DURA MATER, ('hard mother,' called *dura*, because of its great resistance; and *mater*, because it was believed to give rise to every membrane of the body.) *Crassea* seu *Dura meninx*, *Scleromeninx*, *Pachymeninx*, *Perimeninx*, *Méninx extérior* seu *scléris* seu *dermatodes* seu *pachéris* of Galen, *Dura membra'na cerébrum am'biens*, *Cuticula'ris membra'na*, (F.) *Dure mère*, *Dure taye* (Paré), *Méninge*, *Périméninge* (Ch). It is a fibrous, semi-transparent membrane, of a pearly-white colour, thick, and very resisting; lines the cavity of the cranium, and contains the spinal marrow; protects the brain and marrow; and, by its various expansions,—the *falx cerebri*, *tensorium*, &c.,—supports the different parts of the cerebral mass. The largest artery of the dura mater is the *A. meningéa média*.

DURA MATER, LATERAL PROCESSES OF THE, Tentorium—d. m. Testis, Albuginea.

DURATION OF LIFE, see *Life*—d. of Pregnancy, see *Pregnancy*.

DURE-MÈRE, Dura mater.

DURE-TAYE, Dura mater.

DURÉE DE LA VIE, ('duration of life,') see *Longevity*.

DURETÉ D'OREILLE, ('hardness of ear,') *Baryecolia*—d. de l'Ouie, Deafness.

DURILLON, (from *durus*.) Callosity.

DURRA, (in India,) *Panicum Italicum*.

DURUS, Hard.

DUSTING BAG. In pharmacy, a kind of sieve, which consists of a bag made of lawn or other like material, hung inside of a wide-mouthed bottle or tin canister, to the mouth of which it is secured. The powdered substance is put into the bag, and the mouth being closed with a cover, the apparatus is shaken, and the finer particles pass into the bottle or canister.

DUTCHMAN'S PIPE, *Aristolochia hirsuta*.

DUTROA, *Datura stramonium*.

DWALE, ([D.] *dwalen*, 'to wander,') was formerly used to express a lethargic condition. In Devonshire, England, 'to *dwale*,' means to mutter deliriously.

DWALE, Common, *Atropa belladonna*.

DWARF, (Sax. *ƿƿeops*.) *Nanus*—d. Flag, *Iris lacustris*.

DWARFISH, see *Nanus*.

DWARFISHNESS, *Nanosomus*.

DYERS' BROOM, *Genista tinctoria*—d. Weed, *Genista tinctoria*, *Reseda luteola*.

DYNAMETER, *Dynamometer*.

DYNAMIC, *Dynamical*, *Dyn.'micus*, (F.) *Dynamique*, (from *dynamis*.) In Mechanics, *Dynamics* investigates the powers whereby bodies are put in motion, and the laws connected therewith. In Biology, that which relates to the vital forces, to the organism in action, *Vital Dynamics*. The influences of agents on the organism, which are ascribable to neither mechanical nor chemical causes, are sometimes termed *dyn'mic*. In this sense, it is a favorite term with the homeopaths, who maintain that, by certain processes, called, in the aggregate, *dynamization* and *potenti'zing*, the dynamic powers of a medicine may be set free and developed—as by shaking the bottle in which the article is contained, or by mixing an unlimited number of unmedicated globules with one that is medicated, and shaking them together. In this way, the former become *potentized*.

DYNAMIC DISEASES, see *Organic*.

DYNAMICS, VITAL, see *Dynamic*.

DYNAMICUS, *Dynamic*.

DYNAMIOMETER, *Dynamometer*.

DYNAMIS, (*δυναμις*), Faculty, Force.

DYNAMIZATION, see *Dynamic*.

DYNAMOMETER, *Dynamiom'eter*, *Myodynamiom'eter*, *Myodynamom'eter*, (F.) *Dynamomètre*, (*dynamis*, and *μετρον*, 'measure,') An instrument, contrived by M. Regnier for measuring the comparative muscular strength of man and animals. It consists of a spring, which, when pressed upon, causes a needle to move upon a portion of a circle, furnished with a scale of kilogrammes and one of myriagrammes. To measure the strength of the hands, the two branches of the spring are firmly grasped, and brought as near together as the force can carry them. This effort makes the needle traverse, and indicates, on the scale of kilogrammes, the strength of the experimenter's hands. A man, 25 or 30 years of age, exerts a force commonly equal to 50 kilogrammes or 100 pounds.

The strength of the loins of a man, about 30 years of age, as indicated by this instrument, is usually about 30 myriagrammes, or 265 pounds, which shows the weight he is capable of raising.

From experiments made by Péron, in his voyage, on 12 individuals of Van Diemen's Land, 17 of New Holland, 56 of the Island of Timor, 17 Frenchmen belonging to the expedition, and 14 Englishmen in the colony of New South Wales, he found their comparative strength, indicated by the dynamometer, to be as follows:

NATIVES OF	STRENGTH.			
	of the arm.		of the loins.	
	Kilog.	lbs.	Myri.	lbs.
1. Van Diemen's Land.	50.6	101.2		
2. New Holland.	50.8	101.6	10.2	208.08
3. Timor.	58.7	117.4	11.6	238.64
4. France.	69.2	138.4	15.2	310.08
5. England.	71.4	142.4	15.2	332.52

DYNAMOMETER OR DYNAM'ETER, MEDIC'INAL. An instrument devised by Dr. Paris, for the purpose of showing the quantity of active matter contained in a given weight or measure of any official compound, with the dose of any preparation, which will be equivalent in strength to a given quantity of any other of the same class. The instrument is more ingenious than useful.

DYNAMOPATHY, (*dynamis*, and *πάθος*, 'affection,') *Homœopathy*.

DYOTA. Diota.

DYS, *dys*, as a prefix, 'difficult, faulty;' sometimes privative; mostly answering to the particles *dis*, *in*, *mis*, or *un*, in English.

DYSÆMIA, (*dys*, and *αἷμα*, 'blood.') A morbid condition of the blood.

DYSESTHESIA, (*δυσαισθησία*,) *Dysæsthesia*, (*dys*, and *αἰσθάνομαι*, 'I feel.') Obscure, diminished, or even abolished sensation.

DYSESTHESIA AUDITORIA, Cophosis—d. Gustatoria, Ageusia—d. Interna, Amnesia—d. Olfactoria, Anosmia.

DYSESTHESIA VISUALIS, *Parop'sis*. Sense of sight vitiated or lost.

DYSESTHESIS, *Dysæsthesia*.

DYSANAGOGUS, (*δυσαναγωγός*, from *dys*, and *αναγω*, 'I bring up.') That which is expectorated with difficulty. An epithet given, by the Greek writers, to the sputa, when expectorated with difficulty on account of their viscidities.

DYSANARRHOPHESIS, *Dysanarrhophia*; (*dys*, and *αναρροφῆσις*, 'absorption,') Diminished absorption from morbid causes.

DYSAPHÈ, *Dysaphia*, (*dys*, and *ἀφή*, 'feeling,') Morbid touch.

DYSAPULOTUS, *Dysapulus*, *Dysepulo'tus*, *Dysepulo'ticus*, *Dysulotus*, (*dys*, and *ἰσουλω*, 'to heal,') Healing with difficulty.

DYSARTHRITIS, (*dys*, and *arthritis*, 'gout,') Gout, irregular.

DYSCATABROSIS, (*dys*, and *καταβρωσις*, 'the act of eating up,') Dysphagia.

DYSCATAPOPOSIS, (*dys*, and *καταποσις*, 'deglutition,') Dysphagia.

DYSCATAPO'TIA. This term is recommended by Dr. Mead as a substitute for hydrophobia, which means dread of liquids. The dread seems to be partly caused by the difficulty of deglutition.

DYSCHÆZIA, (*dys*, and *χρῆμα*, 'to go to stool,') Difficult and painful defecation.

DYSHRŒA, (*δυσχρῶα*, from *dys*, and *χρῶα* or *χρῶα*, 'colour,') *Dyschro'a*. Sickly and unhealthy colour of the skin. Used, also, synonymously with the *maculæ* of Willan.

DYSCHROMATOPsia, (*dys*, *χρῶμα*, 'colour,' and *οψις*, 'vision,') Achromatopsia.

DYSCHYMOSEN, (*dys*, and *χυμος*, 'juice,') Chymoplasia.

DYSCINE'SIA, (F.) *Dyscinésie*, (*δυσκίνησις*, from *dys*, and *κινῶ*, 'I move,') Difficulty or utter incapability of moving.

DYSCÆLIA, (*dys*, and *καῖα*, 'the abdomen,') Constipation.

DYSCÆLIUS, (*δυσκαίλιος*,) Constipated.

DYSCOPHO'SIS, (*dys*, and *κοφῶ*, 'I am deaf,') A defect in the sense of hearing.

DYSCORIA, *Coremetamorpha'sis*, (*dys*, and *κορη*, 'the pupil,') Irregularity of shape of the pupil.

DYSCRA'SIA, (*δυσκρασία*, from *dys*, and *κράσις*, 'temperament,') *Intemper'ies*, *Distemperan'tia*, *Dys'crasy*. A bad habit of body. See *Dyssthetica*.

DYSCRASIA POTATO'RUM. 'Dyscrasia of drinkers.' The aggregate of cachectic phenomena exhibited by those who indulge in alcoholic liquors.

DYSCRASIA SATURNINA, Saturnismus—d. Serosulosa, Serofula—d. Tuberculosa, see *Tubercle*.

DYSCRASIACUM, Spanæmic.

DYSCRASY, Dyscrasia—d. Bilious, Cholosis, Icterus.

DYSCRITOS, (*δυσκρίτος*, from *dys*, and *κρίσις*, 'judgment,') That which it is difficult to judge of.

DYSDA'RIA, *Dysdacryo'sis*, (*dys*, and *δακρῦον*, 'a tear,') A morbid condition of the tears.

DYSDACRYOSIS, Dysdacria.

DYSECCRIS'IA, (*dys*, and *εκκρίσις*, 'excretion,') Difficult or defective excretion.

DYSECOIA, (*δυσήκοια*, from *dys*, and *ἀκοή*, 'audition,') Barycoia, Deafness.

DYSEL'CIA, (*δυσέλκεια*, and *δυσέλκεια*, from *dys*, and *έλκος*, 'an ulcer,') An ulcer difficult to heal.

DYSEMERISIA, (*dys*, and *εμεσις*, 'vomiting,') Vomiturition.

DYSENTERIA, (*δυσεντερία*,) Dysentery—d. Biliosa, Colocolosis—d. Hæmatera, Dysentery—d. Hepatica, Hepatirrhœa—d. Maligna, Enterocæce—d. Putrida, Enterocæce—d. Scorbatica, Enterocæce—d. Splenica, Melæna—d. Typhodes, Enterocæce.

DYSENTER'IC, (*δυσεντερικός*,) *Dysenter'icus*, *Dysen'terus*, (F.) *Dysentérique*, *Dysentérique*. Relating to dysentery.

DYSENTERIE, Dysentery.

DYSENTERIUM, Dysentery.

DYSENTERONERIA, (*dys*, *εντέρον*, 'intestine, and *νεύρον*, 'a nerve,') Colic—d. Saturnina, Colica metallica.

DYS'ENTERY, *Dysenter'ia*, *D. hæmate'ra*, *Dysenter'ium*, (*dys*, and *εντέρον*, 'intestine,') *Difficul'tas intestino'rum*, *Dissolutus mor'bus*, *Diarrhœa carno'sa*, *Coli'tis*, *Coloni'tis*, *Colorecti'tis*, *Endocoli'tis*, *Esocoli'tis*, *Deflux'us Dysenter'icus*, *Feb'ris Dysenter'ica*, *Flu'men dysenter'icum*, *Flux'us dysenter'icus* seu *cruent'us cum tenes'mo*, *Rheumatis'mus intestino'rum cum ul'cere*, *Tor'mina*, *T. Cell'i*, *Blennenter'ia*, *Mor'bus dissolut'us*, *Se'des cruent'æ*, *Lux'es dysenter'ica*, *Blood'y Flux*; *Flux*, (F.) *Dysentérie*, *Dysentérie*, *Flux dysentérique*, *Flux de Sang*. Inflammation of the mucous membrane of the large intestine; the chief symptoms of which are:—fever, more or less inflammatory, with frequent mucous or bloody evacuations; violent tormina and tenesmus. When the evacuations do not contain blood, it has been called *Dysenter'ia al'ba* or *sim'ple Dys'entery*, *My'codysenter'ia*. The seat of the disease is, generally, in the colon and rectum. It occurs, particularly, during the summer and autumnal months, and in hot climates more than in cold: frequently, also, in camps and prisons, in consequence of impure air, and imperfect nourishment: and is often epidemic. Sporadic cases of dysentery are, generally, easily managed; but when the disease occurs epidemically, it often exhibits great malignancy. Generally, it yields to mild laxatives, as castor oil, combined with diaphoretic narcotics, such as the *pulvis ipecacuanhæ compositus*, and counter-irritants to the abdomen; but, at times, the inflammation runs on so speedily to ulceration, that, unless a new action be rapidly excited, death will be the consequence. In such cases, mercury must be rapidly introduced into the system, and narcotics may be combined with it.

The whole management in acute dysentery, must, of course, be strictly antiphlogistic.

An epidemic dysentery prevailed in the 10th century, under the name *Scitta*.

DYSENTERY, BILIOUS, Colocolosis.

DYSEPULOTICUS, (*dys*, *επι*, 'upon,' and *ουλω*, 'to heal,') Dysapulotus.

DYSEPULOTUS, Dysapulotos.

DYSGALACTIA, Dysgalia.

DYSGA'LIA, *Dysgalac'tia*, (*dys*, and *γάλα*, gen. *γαλακτός*, 'milk,') An unhealthy condition or depravation of the milk.

DYSGENNE'SIA, (*dys*, and *γεννησις*, 'generation,') Lesion of the generative organs or functions.

DYSGEU'SIA, *Dysgeu'stia*, (*dys*, and *γεύσις*, 'taste,') A morbid condition of the sense of taste.

DYSHÆMIA, (*dys*, and *αἷμα*, 'blood,') A morbid condition of the blood.

DYSHÆMORRHŒA, (*dys*, *αἷμα*, 'blood,' and *ῥεω*, 'to flow,') Difficulty in the flow of blood,—according to some, of the hemorrhoidal flux. Also, symptoms occasioned by its diminution or suppression.

DYSHAPH'IA, *Dysaph'ia*, (*dys*, and *ἀφή*,

'touch.') A morbid condition of the sense of touch.

DYSIDRIA, *Dysidria*, (*dys*, and *ιδρως*, 'sweat.') A morbid state of the perspiration.

DYSIATOS, (*δυσιατος*, *dys*, and *ιασαι*, 'to heal.') *Cura tu difficilis*. Difficult to cure.

DYSLALIA, (*dys*, and *λαλειν*, 'to speak,') Balbities, Bradylalia.

DYSLOCHIA, *Colica lochialis*, *Hysteralgia lochialis*, (*dys*, and *λοχος*, 'relating to parturition.') Diminution or suppression of the lochial discharge.

DYSMASESIS, *Dysmasse'sis*, *Bradymasse'sis*, (*dys*, and *μασσειν*, 'mastication.') Difficult or impeded mastication.

DYSMENIA, (*dys*, and *μηνες*, 'the menses,') Dysmenorrhœa.

DYSMENORRHÉE, Dysmenorrhœa.

DYSMENORRHŒA, (from *dys*, and *menorrhœa*.) *Dysmenia*, *Paramenia difficilis*, *Menorrhagia stillatitia*, *Menstrua difficilis* seu *Dolorosa*, *Amenorrhœa difficilis* seu *partialis*, *Menstrues dolorifici*, *Menorrhagia difficilis*, *Menstrua difficilis* seu *dolorifica*, *Laborious* or *Difficult Menstruation*, (F.) *Dysménorrhée*, *Menstruation difficile*, *Règles difficiles*, *Strangurie menstruelle*. Catamenia passed with great local pain, especially in the loins:—with sometimes a membranous discharge. Dysmenorrhœa is very difficult of removal, and prevents conception. In the married female, if she should be able to pass one period without pain, and subsequently become pregnant, the morbid action may be broken in upon by gestation, and a perfect cure be obtained. Change of air, soon after marriage, will sometimes give occasion to this desirable result. The affection generally depends upon erethism of the interior of the uterus, called into action at each catamenial period. The violence of the pain requires the liberal use of narcotics.

DYSMNE'SIA, (*dys*, and *μνησις*, 'memory.') Defective memory.

DYSMORPHE, (*dys*, and *μορφή*, 'shape,') Deformation.

DYSMORPHIA, (*δυσμορφία*), Deformation.

DYSMORPHOSIS, Deformation.

DYSNÉPHRONERVIE, (*dys*, *νεφρος*, 'kidney,' and *νευρον*, 'a nerve,') Nephralgia.

DYSODES, (*δυσωδης*), Fetid.

DYSODIA, (*δυσωδία*, from *dys*, and *οδω*, 'I smell,') *Dysodmia*, *Dysosmia*, *Fætor*, (F.) *Puanteur*. Sauvages has given this generic name to all diseases, characterized by fetid emanations, from whatever part proceeding;—from the mouth, nasal fosse, bronchia, stomach, axillæ, groins, &c. Also, a stench or stink, *Pso'a*.

DYSODMIA, (*δυσωδμία*), Dysodia.

DYSODONTIASIS, *Dentitio difficilis*, (*dys*, and *οδοντασις*, 'dentition.') Difficult dentition.

DYSONEIR'US, (*dys*, and *ονειρος*, 'a dream,') Insomnia, with restlessness.

DYSOPIA, properly *Dysopia*, *Dysora'sis*, (*dys*, and *οπταται*, 'I see,') Difficulty of seeing; obscurity of vision.

DYSOPIA DISSITORUM, Myopia.

DYSOPIA LATERALIS, *Paropsia lateralis*, *Skuesight*, *Skewsight*, *Sight askew*. Vision only accurate when the object is placed obliquely. This state is generally caused by some opacity of the cornea.

DYSOPIA LUMINIS, Nyctalopia—d. Proximorum, Presbytia—d. Tenebrarum, Hemeralopia.

DYSPSIA, Dysopia.

DYSORASIS, (*dys*, and *ορασις*, 'vision,') Dysopia.

DYSOREX'IA, *Inappetentia*, *Appetitus deficiens*, (*dys*, and *ορεξις*, 'appetite,') Diminution of appetite.

DYSOS'MIA, (*δυσωσμία*, from *dys*, and *οσμη*, 'smell,') Diminution of smell.

DYSOSPHERE'SIA, *Dysosphre'sis*, *Dysosphre'sia*, and *οσφρησις*, 'the sense of smell,') A morbid state of the sense of smell.

DYSOSTO'SIS, (*dys*, *οστων*, 'a bone,' and *οσις*.) A faulty conformation or morbid condition of bone.

DYSPATHIA, (*δυσπαθεια*, from *dys*, and *παθος*, 'an affection,') See Serious.

DYSPESPIA, (*δυσπεψία*, from *dys*, and *πεπρω*, 'I concoct,') *Limo'sis* *Dyspepsia*, *Anorexia*, *Apepsia*, *Bradypepsia*, *Gastroataxia*, *Digestio depravata* seu *difficilis* seu *læsa*, *Gastro-atonia*, *Concoctio tarda*, *Stomachi resolutio*, *Cru'ditas*, *Pas'sio stomachica*, *Dyspepsy*, *Indigestion*, *Difficility of Digestion*. A state of the stomach, in which its functions are disturbed, without the presence of other diseases, or when, if other diseases be present, they are of but minor importance. The symptoms of dyspepsia are very various. Those affecting the stomach itself are:—loss of appetite; nausea; pain in the epigastrium or hypochondrium; heart-burn; sense of fulness, or weight in the stomach; acid or fetid eructations; pyrosis, and sense of fluttering or sinking at the pit of the stomach. The sympathetic affections are of the most diversified character. Dyspepsia, being generally of a functional nature, is devoid of danger. When arising from disease of the stomach itself, it is, of course, more serious.

It is usually dependent on irregularity of living, either in the quantity or quality of the food taken: and the most successful treatment is, to put the patient on a diet easy of digestion; to combat the causes, where such are apparent; and, by proper remedies and regimen, to strengthen the system in every practicable manner. A great error exists in regarding it as always a disease of debility. It is often connected with an inflammatory or subinflammatory condition of the mucous lining of the stomach, and of course a very different plan of treatment is required in the two cases. Dyspepsia is often attended with too great a secretion of the gastric acids; but, on other occasions, they would appear to be too small in quantity, so as to constitute *alkaline indigestion* or *neutral indigestion*.

DYSPESPIA CHLOROSIS, Chlorosis—d. Hypochondriasis, Hypochondriasis—d. Pyrosis, Pyrosis.

DYSPESPIODYNIA, (*dyspepsia*, and *οδυνη*, 'pain,') Cardialgia.

DYSPESPODYNIA, Cardialgia.

DYSPESPY, Dyspepsia.

DYSPET'IC, *Dyspepticus*, *Dyspeptus*, *Apeptic*, *Apepticus*. Having relation to dyspepsia, as '*dyspeptic bread*.' One who suffers from dyspepsia.

DYSPETICUS, Dyspeptic.

DYSEPTODYNIA, (*dys*, *πεπρω*, 'I concoct,' and *οδυνη*, 'pain,') Cardialgia.

DYSEPTUS, (*δυσπεπτος*), Dyspeptic.

DYSPERMA'SIA, *Dyspermatismus*, (*dys*, and *σπερμα*, 'sperm,') Difficulty—sometimes incapacity—of voiding the sperm.

DYSPERMATISMUS, (*dys*, and *σπερματισμος*, 'discharge of sperm,') Bradyspermatus.

DYSPHAGIA, *Dyscatbro'sis*, *Dyscatapno'sis*, *Deglutitio difficilis*, *D. læsa*, *D. impedita*, (*dys*, and *φαγειν*, 'I eat,') Difficulty of deglutition. Dysphagia is almost always symptomatic, either of inflammation or of other disease of the organs of deglutition, or of incomplete obstruction of the œsophagus, by some obstacle within it, or by a neighbouring tumour. At times, it is produced by spasms or paralysis of the œsophagus. The prognosis and treatment vary according to the cause.

DYSPHAGIA ATONICA, Pharyngoplegia—d. Callosa, D. constricta.

DYSPHAGIA CONSTRICTA. *D. Pharyngea* seu *Œsophagea* seu *Calloea*, *Strictura Pharyngis* seu *Œsophagi* *e'ra* seu *Callo'sa*, *Stenochœria* *Œsophagi*, *Œsophagiarectia*, *Œsophagostenoma*, *Læmostenosis*. Stricture of the pharynx and œsophagus is an affection which may be the result of pharyngitis or œsophagitis; but more frequently of malignant disease in the parietes of the tube. The only remedy is the bougie.

DYSPHAGIA GLOBOSA. Angone—*d. Hysterica*, Angone—*d. Inflammatoria*, Cynanche tonsillaris—*d. Linguosa*, Paraglossa—*d. Nervosa*, Œsophagismus—*d. Œsophagea*, *D. Constricta*—*d. Paralytica*, Œsophagoplegia, Pharyngoplegia—*d. Pharyngea*, *D. Constricta*—*d. Ranula*, *Ranula*—*d. Scirrhusa*, *Læmoscirrhus*—*d. Spasmodica*, Œsophagismus—*d. Spastica*, Œsophagismus—*d. Torpida*, Pharyngoplegia—*d. Uvulosa*, Staphylœdema.

DYSPHONIA, *Dys'phony*, (*δυσφωνία*, from *dys*, and *φωνη*, 'the voice.') Difficulty of producing and articulating sounds: voice imperfect or depraved. *Aphonia* (of some).

DYSPHONIA IMMOBULATA NASALIS, Rhinophonia—*d. Immodulata palatina*, Asaphia.

DYSPHORIA, (*δυσφωρία*, from *dys*, and *φωρῶ*, 'I bear.') *Inquietudo*, *Astasia*. Dissatisfaction; restlessness; suffering; indisposition.

DYSPHORIA ANXIETAS, Anxiety—*d. Nervosa*, Fidgets—*d. Simplex*, Fidgets.

DYSPHOTIA, (*dys*, and *φως*, gen. *φωτος*, 'light.') Myopia.

DYSPIONIA, (*dys*, and *πιων*, 'fat.') A morbid condition of the adipous substance.

DYSPASMATIC, (*dys*, and *πλάσσειν*, 'to form.') Cacoplastic.

DYSPLASTICUM, Spanæmic.

DYSPNÉE, Dyspnœa—*d. Cardiaque*, Asthma, cardiac.

DYSPNŒA, (*δυσπνοία*, from *dys*, and *πνέω*, 'I breathe.') *Pseudo-asthma*, *Amphipneuma*, *Respiratio difficilis* seu *brevis et raris*, *Brachypnœa*, *Retentio aërea*, *Anhelation*, *Short breath*, *Difficulty of breathing*, (F.) *Dyspnée*, *Courte Hæleine*. Dyspnœa may be idiopathic or symptomatic. The latter accompanies almost all thoracic diseases. Urgent dyspnœa has been called *Amphipneuma*, *αμφιπνεύμα*.

DYSPNŒA CONVULSIVA, Asthma—*d. Hydrothoracica*, *Hydrothorax*—*d. Montana*, *Puna*—*d. Phythoracica*, *Pneumothorax*—*d. Pinguedinosa*, *Pursiness*—*d. Pneumatica*, *Pneumothorax*—*d. Pyothoracica*, *Empyema*.

DYSSENTÉRIE, Dysentery.

DYSSENTÉRIQUE, Dysenteric.

DYSSIALIA, (*dys*, and *σάλω*, 'saliva.') A morbid condition of the saliva.

DYSSYNODUS, *Dys'synusia*, (*δύς, ἐν συνόδῳ*, 'coition.') *Coitus difficilis*. Difficulty in coition.

DYSSYNUSIA, (*dys*, and *συνουσία*, 'coition.') Dysynodus.

DYSTHAN'ATUS, (*δυσθανάτος*, from *dys*, and *θανάτος*, 'death.') That which causes a slow and painful death. One who experiences this kind of death.

DYSTHELA'SIA, (*dys*, and *θηλαζω*, 'I give suck.') Inaptitude for suckling.

DYSTHERAPEUT'US, (*δυσθεραπεύς*, from *dys*, and *θεραπεία*, 'medical treatment,') *difficiliter curabilis*. That which is difficult of cure.

DYSTHE'SIA, (*δυσθεία*, from *dys*, and *θῆμη*, 'I am situate.') *Dys'thesia*, *Cachexia*. Morbid habit. Bad humour. Impatience in disease.

DYSTHET'ICA, *Cachexia*, *Dyscrasia*, *Cachexies*. A morbid condition of the system of nutrition, connected with a faulty state of the blood, producing a diseased habit. The fourth order in the class *Hæmatia* of Good, including *Plethora*, *Hæmorrhagia*, &c.

DYSTHYM'IA, (*δυσθυμία*, from *dys*, and *θυμός*, 'mind.') *Depression*, *Despondency*. A bad sign in acute diseases. Also, Melancholy.

DYSTOC'IA, *Mogostocia*, *Bradynocia*, *Retentio foetus*, (*δυστοκία*, from *dys*, and *τοκος*, 'accouchement.') (F.) *Dystocie*. *A laborious accouchement*, *Laborious or morbid*: or *difficult labour*, *Parodinia*, *Partus difficilis*. See Laborious.

DYSTOCIA ABORTIVA, Abortion—*d. Dyscyesis*, *Pregnancy*, morbid—*d. Dyscyesis extrauterina*, see Pregnancy, preternatural.

DYTECHIASIS, *Hispidity*, (*dys*, and *τεταχας*, 'order.') Irregular position of the eye-lashes.

DYSTON'IA, (*dys*, and *τῶς*, 'tone.') Morbid condition of the tone of a tissue or organ.

DYSTR'AUM'IA, (*dys*, and *τραύμα*, 'a wound.') The condition of wounds when they heal with difficulty, as in some persons and climates.

DYSTROPH'IA, (*dys*, and *τροφή*, 'nourishment.') Imperfect or defective nutrition.

DYSULOTUS, (*dys*, and *ουλη*, 'a cicatrix.') Dysapulotus.

DYSURE'SIA, *Dysure'sis*; (*dys*, and *ουρησις*, 'passing the urine.') Defective secretion and evacuation of the urine.

DYSUR'IA, (*δυσουρία*,) *Dys'ury*, *Urinee difficilis excretio*, *Strangury* (of some). (F.) *Dysurie*. Difficulty of passing the urine. In this affection the urine is voided with pain, and a sensation of heat in some part of the urethra. Dysuria is the first degree of retention of urine. It differs from strangury, in which the urine can only be passed in drops and with great straining.

DYSURIA CALCULOSA, Calculi, vesical—*d. Irritata*, Calculi vesical—*d. Mucosa*, Cystirrhœa.

DYSURY, Dysuria.

E.

E, (L.) as a prefix, 'out of,' 'from.'

EAGLESTONE, *Ætites*.

EAR, (Sax., *eape*.) *Aur'is*, *Ous*, *Ac'oē*, (Prov.)

Lug. (F.) *Oreille*. The organ of audition. It is composed of a series of more or less irregular cavities, in which the sonorous rays are successively received and reflected, until they agitate the nerves which are destined to convey the impression to the brain. The ear is contained partly in the substance of the temporal bone; and a part projects externally, behind the joint of the lower jaw. It may be divided into three portions;—

the *outer or external ear*, formed by the auricle and meatus auditorius; the *middle ear*, comprising the cavity of the tympanum and its dependencies; and the *internal ear*, comprehending the three semicircular canals, the cochlea and the vestibule; which, together, constitute the *osseous labyrinth*. Within the cavity of this labyrinth are contained membranes having nearly the shape of the vestibule and semicircular canals, but not extending into the cochlea. These membranes form the *membranous labyrinth*. Between the osseous and the membranous labyrinth is situate

the liquor of Cotunnus, and within the membranous labyrinth is a fluid, termed, by De Blainville, *vitrine auditiva*, from its supposed analogy to the vitreous humour of the eye. The form of the membranous vestibule is not an exact imitation of the osseous cavity, being composed of two distinct sacs, which open into each other,—the one termed the *Sacculus vestibuli*; the other *Sacculus*. Each sac contains in its interior a small mass of white calcareous matter resembling powdered chalk, which seems to be suspended in the fluid of the sacs by means of a number of nervous filaments proceeding from the auditory nerve. These are the *otococci* and *otolithes* of Breschet.

The auditory nerve is distributed to the cavities of the internal ear.

EARDOCTOR, Aurist—*e.* Drum, Tympanum—*e.* Finger, see Digitus—*e.* Flap, Proptoma auricularum—*e.* Kecker, Tonsil.

EARNING, (from [G.] *gerinnen*, 'to coagulate' [?]). Rennet.

EARPICK, *Otoglyphis*, *Otoglyphum*, *Ooch'lear auricula'rē*, *Auriscalpium*, (F.) *Cure-oreille*. A species of small scoop, used for extracting hardened cerumen from the meatus auditorius externus; or to remove foreign bodies from the ear. If carelessly used, it is apt to excite inflammation of the tube.

EARPOWDER, see Otolith.

EARSURGEON, Aurist—*e.* Surgery, Otiatria.

EARTRUMPET, *Tubus acusticus*, *Acusticum Cor'nu*, *Otopho'nē*, (F.) *Cornet acoustique*. An instrument for collecting sound and increasing its intensity, used by those who are hard of hearing. It is, commonly, a kind of cone, formed of silver, tin, or elastic gum, the base of which is turned towards the person who is speaking, and the apex placed in the entrance of the meatus auditorius externus.

EARWAX, Cerumen.

EARWIG, (*veruca*, 'a caterpillar,' Sax. *eappigsa*, 'earworm.') *Forficula auricularia*.

EARTH CLUB, *Orobancha Americana*—*e.* Eater, Geophagist—*e.* Eating, Geophagism.

EARTH, FULLER'S, *Cimolia purpurescens*.

EARTH GALL, *Veratrum viride*.

EARTH, HEAVY, *Baryta*—*e.* Japan, *Catechu*—*e.* Lemnian, *Terra Lemnia*.

EARTH NUT, *Bunium bulbocastanum*, *Pignut*.

EARTH, PONDEROUS, *Baryta*—*e.* Samian, *Sami terra*—*e.* Sealed, *Terra sigillata*—*e.* Talc, *Magnesia*.

EASTINGWORT, *Scabiosa*.

EATABLE, Esculent.

EATIN-BERRIES (Sc.), see *Juniperus communis*.

EAU, ([L.] *aqua*.) Water—*e.* d'Aix-la-Chapelle, see Aix-la-Chapelle.

EAU D'ALIBOUR. This compound is made of sulphate of zinc, and sulphate of copper, each 3j; camphor, ten grains; saffron, four grains; water, four fluidounces. Employed in chronic inflammation of the eyelids, and as a vulnerary.

EAU D'AMANDES AMÈRES, *Aqua amygdalarum concentrata*—*e.* de l'Amnios, *Liquor amnii*.

EAU ANTIPUTRIDE DE BEAUFORT. Mineral lemonade prepared with sulphuric acid.

EAU D'ARMAGNAC, *Tinctura cinnamomi composita*—*e.* du *Balaruc*, *Balaruc waters*—*e.* de *Barèges*, *Barèges water*—*e.* de *Binelli*, *Aqua Binellii*—*e.* *Blanche*, *Liquor plumbi subacetatis dilutus*—*e.* de *Bonferme*, *Tinctura cinnamomi composita*—*e.* de *Bonnes*, *Bonnes*, mineral waters of—*e.* de *Boule*, see *Ferrum tartarizatum*—*e.* de *Bourbonne-les-Bains*, *Bourbonne-les-Bains*, mineral waters of—*e.* de *Brocchieri*, *Aqua Brocchi-*

erii—*e.* des *Carmes*, see *Melissa*—*e.* de *Chaux*, *Liquor calcei*—*e.* de *Chaux composée*, *Liquor calcei compositus*.

EAU DE COLOGNE, *Cologne' water*. A celebrated perfume, so called from the place where it is made. The following is one formula: *Oil of bergamot*, 3ij; *Oil of lemon*, 3ij; *Oil of Lavender*, 3iiss; *Oil of neroli*, 3iiss; *Oil of origanum*, 3ij; *Oil of rosemary*, 3j; *Essence of vanilla*, 3ij; *Musk*, ten grains; *Rectified spirit*, Oxxij; *Rose-water*, Oij; *Orange-flower water*, Oj. Macerate for fourteen days, and filter.

EAU DE CUIVRE AMMONIACALE, *Liquor cupri ammoniati*—*e.* *Destillée*, Water, distilled—*e.* de *Fontaine*, Water, spring—*e.* des *Fontaines de la Marquerie*, Rouen, mineral waters of—*e.* contre la *Gangrène*, *Liquor hydrargyri nitrici*—*e.* de *Goudron*, see *Pinus sylvestris*—*e.* de *Goulard*, *Liquor plumbi subacetatis dilutus*.

EAU HÉMASTATIQUE DE TISSERAND, *Eau de Tisserand*. A hæmstatic water, reputed to possess the same properties as the *Aqua Brocchieri*. It may be prepared by digesting dragon's blood, and turpentine of the *Vosges* in water.

EAU DHUSSON, *Vinum colchici*—*e.* des *Hydropiques*, Serum of serous membranes—*e.* *Hydrosulfurée simple*, *Hydrosulphuretted water*.

EAU DE JAVELLE, (after the mill of Javelle, where it was first made.) *Bleaching liquid*, *Aqua alkali'na oxymuriat'ica seu Javelle'nsis*, *Labarraque's Solution*, (Common salt, ʒij; *Black oxide of manganese*, ʒij; water, ʒij. Put into a retort, and add, gradually, oil of vitriol, ʒij. Pass the vapour through a solution of subchlorate of potassa 3ij in water 3xxix, applying heat towards the last. S. g. 1.087.) It is stimulant, detergent, and antiseptic,—applied externally.

EAU DE LAC, Water, lake.

EAU DE LÉCHELLE. A hæmstatic water, prepared by macerating, in water, various aromatic and astringent herbs, flowers and roots, with tar, and distilling the same. It has been given internally, in hemorrhage, and used, also, as an external styptic.

EAU DE LUCE, *Spiritus ammoniæ succinatus*—*e.* *Magnéiienne*, *Magnesia*, fluid—*e.* de *Marais*, Water, marsh—*e.* *Médecinale d'Husson*, see *Colchicum autumnale*, *Vinum colchici*, and *Veratrine*, sulphate of—*e.* *Médecinale naturelle*, Water, mineral—*e.* de *Mer*, Water, sea—*e.* *Mercurielle*, *Liquor hydrargyri nitrici*—*e.* *Minérale*, Water, mineral—*e.* de *Monterossi*, *Aqua Binellii*—*e.* de *Naples*, *Naples water*, factitious—*e.* de *Neige*, Water, snow—*e.* de *Pagliari*, *Hæmstatic*, *Pagliari's*—*e.* *Panée*, *Toast water*—*e.* de *Pluie*, Water, rain—*e.* de *Potasse*, *Liquor potassæ*—*e.* de *Puit*, Water, well—*e.* de *Robel*, *Elixir acidum Halleri*—*e.* *Régale*, *Nitromuratic acid*—*e.* *Salée*, Water, sea—*e.* de *Source*, Water, well—*e.* *Styptique de Brocchieri*, *Aqua Brocchieri*—*e.* *Sucrée*, *Hydro-saccharum*—*e.* de *Tisserand*, *Eau hémastatique de Tisserand*—*e.* *Végéto-minérale*, *Liquor Plumbi subacetatis dilutus*—*e.* de *Vichy*, *Vichy water*—*e.* de *Vie*, *Brandy*—*e.* de *Vie Allemande*, *Tinctura jalapii composita*—*e.* de *Vie Camphrée*, *Spiritus camphoræ*.

EAUX LES (pl. of eau), *Liquor amnii*—*e.* *Chaudes*, *Aigues-caudes*—*e.* *Hépatiques*, Waters, mineral, sulfureous—*e.* *Minérales artificielles*, Waters, mineral, artificial—*e.* *Minérales factices*, Waters, mineral, artificial—*e.* *Minérales ferrugineuses*, Waters, mineral, gaseous, &c.—*e.* *Minérales gaseuses ou acides*, Waters, mineral, gaseous, &c.—*e.* *Minérales salines*, Waters, mineral, saline—*e.* *Minérales sulfureuses*, Waters, mineral, sulfureous—*e.* *Sulfurées*, Waters, mineral, sulfureous.

ÉBARBEMENT (F.), (*e*, and *barbe*, 'beard.') The removal by the knife or scissors of a superficial portion of a tumour leaving the rest.

ÉBEAUPIN, MINERAL WATERS OF. An acidulous chalybeate, in the department of *Loire Inférieure*, near Nantes.

ÉBLOUISSEMENT, Dazzling.

EBRIECA'SUM, (*ebrius*, 'drunk.') A term employed by Paracelsus to denote a disturbance of the reason, similar to what occurs in drunkenness.

EBRIETAS, Temulencia.

EBRIOSITAS, Temulencia.

EBULLITIO, Ebullition, Strophulus—c. Stomachi, Ardor ventriculi, Pyrosis.

EBULLIT'ION, *Ebullit'io*, *Æstuat'io*, *Anaz'esis*, *Caus'is*, (*ebullire*, (*e*, and *bullire*,) 'to bubble up.')

Boiling; (F.) Bouillonnement. The motion of a liquid, by which it gives off bubbles of vapour, by heat or fermentation. The boiling point of liquids varies according to the pressure to which they are subjected. For the point of ebullition of different substances, see Heat.

Ebullition is used in France, in a vague manner, for every kind of transient eruption of the skin, occurring without fever or with a slight febrile attack.

EBULUS, Sambucus ebulus.

EBUR, gen. *Eb'oris*, Ivory, see Tooth.

EBURNEATIO, Eburnification.

EBURNIFICATION, *Eburnificatio*, *Eburneat'io*, *Eburnat'ion*, (*ebur*, 'ivory,' and *fio*, 'to be made.')

An incrustation of the cartilages of articular surfaces of bones with phosphate of lime, which gives them the whiteness and hardness of ivory.—*Cartilages éburnés*.

EC—before a vowel, Ex—(*ex*, *ἐξ*), as a prefix, 'out of, from, of.'

ÉCAILLE, (Sax. *reylan*, 'to separate,') Scale.

ÉCAILLES D'HUITRES, *Ostræa testæ*.

ÉCAILLEUX, Squamous.

ECBALIA ELATERIUM, (*ec*, and *βαλλειν*, 'to throw,')

Momordica elaterium.

ECBALIN, see Momordica elaterium.

ECBALIUM AGRESTE, Momordica elaterium—e. *Officinarium*, Momordica elaterium.

ECBESOMENON, (from *ἐκβαίνω*, 'to come out.')

Eventus.

ECBLOMA, (*ec*, and *βαλλειν*, 'to throw,')

See Abortion.

ECBOLE, (*ἐκβολή*), Abortion.

ECBOLIC, Abortive.

ECBOLICUS, Abortive.

ECBOLIUM ELATERIUM, (*ec*, and *βαλλειν*, 'to throw,')

Momordica elaterium.

ECBOLIUS, (*ἐκβολιός*), Abortive.

ECBRAS'MATA, pl. of *Echrasma*, (*ἐκβρασματα*, from *ἐκβράω*, 'I boil up.')

Echym'ata. Hippocrates uses the word for certain burning eruptions.

EC CATHARTICUS, Cathartic.

ECCEPHALO'SIS, *Eccerebra'tio*, *Cephalotom'ia*, (*ec*, *κεφαλή*, 'head,' and *osis*,) The removal of the brain of the child to facilitate delivery.

ECCHELYSIS, (*ec*, and *χελευσσειν*, 'to cough,')

Expectoration.

ECCHORESIS, (*ec*, and *χωρεω*, 'to make room,')

Defecation.

ECCHYLOMA, (*ec*, and *χυλος*, 'juice,')

Succus expressus.

ECCHYMATA, (*ec*, and *χυω*, 'to pour,')

Echymata.

ECCHYMO'MA, gen. *Echym'o'matis*, (*ec*, *χυμός*, 'humour,' and *oma*,) *Echym'o'sis*, *Pelidno'ma*, *Pelio'ma*, *Hyp'e'ma*, *Hypoe'ma*, *Hypoe'mia*, *Effu'sio*, *Exsuec'tio*, *Suffu'sio sanguinis*. A livid, black, or yellow spot, *Li'vor sanguineus*, produced by blood effused into the areolar tissue from a contusion. Spontaneous effusions, occurring as the result of disease or after death, are called *sugillations*.

ECCHYMOMA Aneuriosum, see Aneurism—e.

Capitis recens natorum, Cephalæmatoma—e. Hyponychon, Hyponychon—e. Lymphaticum, Phlegmatia dolens—e. Melasma, Melasma.

ECCHYMOSIS, (*ἐκχυμωσις*,) Ecchymoma.

ECCHYSIS, (*ἐκχυσις*, from *ec*, and *χυω*, 'I pour,')

Effusion.

ECCLISIS, (*ἐκκλισις*, from *ec*, and *κλινω*, 'I bend or give way,')

Luxation.

EC'COPE, *Ec'tomē*, *Ectom'ia*, (*ἐκκοπή*, from *ec*, and *κοπεῖν*, 'to cut.')

The act of cutting out; also, a perpendicular division of the cranium by a cutting instrument. See *Entaille*.

ECOP'EUR, (*ἐκκοπεύς*,) A knife or instrument for cutting. An ancient instrument—the *raspatory*—used in trepanning.

ECOPROSIÆTHES'IS, (*ec*, *κοπρος*, 'excrement,' and *αἴσθησις*, 'sensation,')

The sensation or desire to evacuate the bowels.

ECCOPROSIS, (*ἐκκοπρωσις*, from *ec*, and *κοπρος*, 'excrement,')

Defecation.

ECCOPROT'IC, *Eccoprot'icus*, *Ectoprot'ic*. Mild purgatives or laxatives, whose operation is confined to simply clearing out the intestinal canal.

ECCORTHAT'ICUS, (*ec*, and *κορθαω*, 'I collect,')

An ancient epithet for remedies to which was attributed the property of evacuating collections of humours.

ECRINOL'OGY, *Eccrinolog'ia*, *Eccrisiolog'ia*, (*ἐκκρινω*, *ec*, and *κρινω*, 'I separate,' and *λογος*, 'a discourse,')

A treatise on the secretions.

ECRISIOLOGIA, (*eccrisiis*, and *λογος*, 'a discourse,')

Eccrinology.

ECRISION'OSI, *Eccrisionou'sis*, (*eccrisiis*, and *νοσος*, 'disease,')

Disease of excretion.

ECRISIOS'CHESIS, (*eccrisiis*, and *σχέσις*, 'retention,')

Arrest or retention of excretions: or of a critical evacuation.

ECRISIS, (*ἐκκρίσις*, from *ec*, and *κρινειν*, 'to separate,')

Excretion.

ECRITICA. Disease of the excrent functions. The 6th class in Good's Nosology. Also, medicines that act on the excrent system. *Eliminantia*, *Eliminatives*.

ECYYESIS, (*ec*, and *κυνειν*, 'to conceive,')

Pregnancy, extra-uterine—e. Abdominalis, Pregnancy, abdominal—e. Ovaria, Pregnancy, ovarian—e. Tubalis, Pregnancy, tubal.

ECYLIOS'IS, (*ec*, and *κυλειν*, 'to turn round,')

Morbus evolutio'nis. A disease of evolution or development.

ECDEMIOMANIA, *Ecdemion'osus*, (*ἐκδημω*, 'I travel about,' and *μανία*, 'mania,')

A morbid desire to be travelling about, frequently observed, according to Kraus, "in rich Englishmen and poor Germans," (*"häufig beobachtet an reichen Engländern und armen Teutschen."*) It is the antitheton to Nostalgia.

ECDEMIIONOSUS, (*ἐκδημω*, 'I travel about,' and *νοσος*, 'a disease,')

Ecdemiomania.

EC'DORA, (*ἐκδωρα*, from *ec*, and *δωρω*, 'I flay,')

Anad'ora, *Eccoria'tio*. Excoriation in general, but more especially of the urethra.

EC'DYSIS, (*ἐκδύσις*, from *ἐκδύω*, 'I put off,')

Moulting of the skin of animals. Desquamation.

ÉCHALOTTE, *Eschalotte* (F.), [G.] *Schallotten*, from *Ascalon*, in Syria, *Allium Ascalon'icum*, *Ce'pa Ascalon'ica*. The *Eschalot* or *shallot*. A species of allium, employed in culinary preparations.

ÉCHANCRURE (F.), (*chancre*, [L.] *cancer*, 'a crab' [?]), *Emargina'tio*, *Emarginatu'ra*, *Inci'su'ra*. A word employed by anatomists to designate depressions and notches of various shapes, observed on the surface or edges of bones.

ECHANCRURE ETHMOÏDALE is on the

nasal bone, which unites with the ethmoid. See Ethmoid.

ÉCHANCRURE NASALE, *Nasal Notch*, belongs to the os frontis, and is articulated with the bones of the nose.

ÉCHANCRURE PAROTIDIENNE, is a triangular space, comprised between the parotid edge of the inferior maxillary bone and the mastoid process, so called because it lodges the parotid gland.

ÉCHANCRURE SCAPULAIRE, *Notch, scapular*—*é. Sciatique grande*, see *Sciatic Notch*—*é. Sciatique petite*, see *Sciatic Notch*.

ÉCHARDE, Splinter.

ÉCHARPE, Sling.

ÉCHARPE GRANDE et É. MOYEN, see *Sling*—*é. de J. L. Petit*, see *Sling*—*é. Petite*, see *Sling*.

ÉCHAUBOULURES (F.), *Échanbouillures*, (*chaud*, 'warm,' and *boule*, 'bulla' [?]). *Sudamina*, *Hidro'a*. A word whose meaning is not fixed. It is applied to any eruption on the surface of the body, accompanied with pricking and other uneasy sensations.

ÉCHAUFFANTS, (*échauffer*, [L.] *calefacere*, 'to make warm,') *Calefacients*.

ÉCHAUFFEMENT (F.), *Calefactio*, *Excalfactio*. Augmentation of heat in the animal economy; the symptoms of which are a more than ordinary sensation of heat, disposition to perspiration, great thirst, general indisposition, flushed countenance, &c. It goes off by the use of antiphlogistics and abstinence. In the vulgar language it is often used synonymously with *constipation*, and sometimes for simple gonorrhœa, and for chafing.

ECHECOLLON, (*εχεκollon*, from *εχω*, 'I have,' and *κόλλα*, 'glue,') *Echecollum*. Any topical glutinous remedy.

ECHELLES DU LIMAÇON, *Scalæ of the cochlea*.

ECHENEIS, (*εχενης*,) *Remora* *Hildani*.

ECHETROSIS, (*εχετρωσις*,) *Bryonia alba*.

ECHID'NA (*εχιδνα*) *OCELLA'TA*, *Brown ten-inch-long viper*. A most formidable viper in the forests of Peru, the bite of which is so rapidly fatal, that it kills a strong man in two or three minutes.

ECHINA'CEA PURPU'REA, (*εχινος*, 'the hedge hog,' owing to the spiny chaff of the disk.) *Purple Cone/flower*, *Black Samp/son*; of the *Composite Family*; indigenous in Ohio and westward; its dull purple flowers appearing in July. The root is aromatic, and used popularly as a carminative.

ÉCHINE, *Vertebral column*.

ECHINOCOCCUS HOMINIS, (*εχινος*, 'a hedgehog,' and *κοκκος*, 'a grain or berry,') See *Worms*—*e. Humanus*, *Hydatid*.

ECHINODERMI, (*εχινος*, 'hedgehog,' and *δερμα*, 'skin,') *Porcupine men*.

ECHINOGLOSSUM, (*εχινος*, 'hedgehog,' and *γλωσσα*, 'tongue,') *Ophioglossum vulgatum*.

ECHINOPHTHALMIA, (*εχινος*, 'a hedgehog,' and *οφθαλμια*, 'inflammation of the eye,') *Ophthalmia of the eyelids*, in which the cilia project like the quills of the hedgehog.

ECHINOPS, (*εχινος*, 'hedgehog,' and *ωψ*, 'appearance,') *Crocodilia*, *Acanthalu'ca*, *Scabiosa carduifolia*, *Sphæroceph'ala elatior*, *Echi'nopus*, *Echi'nops Sphæroceph'alus*, *Globe thistle*. The root and seeds are reputed to be moderately diuretic.

ECHINOPUS, (*εχινος*, 'hedgehog,' and *πους*, 'a foot,') *Echinops*.

ECHINUS SCANDENS, *Allamanda*.

ECHITES DIFFORMIS, (from *εχis*, 'the viper,'

owing to its smooth, twining shoots,) *Forsteria difformis*—*e. Suberecta*, see *Curare*.

ECHIOS, (*ηχος* and *αχη*,) *Sound*, *Tinnitus aurium*.

ECHOSCOPE, (*ηχος*, 'sound,' and *σκοπεω*, 'I examine,') *Auscultation*.

ECHOSCOPIUM, *Stethoscope*.

ECHTHYEROCYESIS, (from *ecto*, *βτερα*, 'the uterus,' and *κυησις*, *pregnancy*,) *Pregnancy*, *extra-uterine*.

ECLACTIS'MA, gen. *Eclactism'atis*, *Eclamp'sis*, *Eclamp'sia*, *Epilamp'sia*, *Effulgescent'ia*, (*εκλακτιζω*, 'I kick,') *Epilepsy* is often accompanied with flashings of light; and hence Hippocrates has used the last two words for epilepsy. They have all been applied to convulsions.

ÉCLAIR, *Astrape*.

ÉCLAIRE, *Chelidonium majus*—*é. Petite*, *Ranunculus ficaria*.

ECLAMP'SIA, (*εκλαμψις*, (*ec*, and *λαμβω*, 'I shine,') 'brilliance,') *Flashes of light* before the eyes. See *Eclactisma*. *Convulsion*, as the convulsions of children, *Eclamp'sia infantum*, *Epilep'sia acuta infantum* seu *fibrilis infantum* seu *puerilis*, (F.) *Convulsions des Enfants*, *Eclampsie*; (probably from *εκ*, and *λαμβανω*, *λαμβάνω*, 'to seize hold of,')

ECLAMP'SIA GRAVIDA'RUM ET PARTURIEN'TIUM, *Puer'peral Convulsions*, (F.) *Convulsions des femmes enceintes et en couche*. *Convulsions of pregnant and parturient women*.

ECLAMP'SIA INFANTUM, see *Eclampsia*—*e. Nutans*, *Convulsion*, *salaam*—*e. Typhodes*, *Raphania*.

ECLAMP'SIE DES ENFANS, *Eclampsia infantum*.

ECLECTIC PHYSIC'IAN'S, *Eclect'ici Med'ici*, (*εκλεκτικος*, 'selecting,' from *εκλεγω*, (*εκ*, and *λεγω*,) 'I choose,') A sect of physicians, who professed to choose, from other sects, all the opinions which appeared to them best founded. Agathinus of Sparta, master of Archigenes of Apamea, in Syria, was its reputed founder; and Archigenes and Areteus were its greatest ornaments. The doctrine was called *Eclect'ism*, *Eclectis'mus*, *Medic'ina eclectica*, *Eclectic medicine*. Every judicious physician must be an eclectic.

The term *Eclectic* is, however, not unfrequently applied, in the United States, to one who 'chooses' to adopt exclusive views:—the so called *Eclectic physicians* generally belonging to the class of *botanical physicians*.

ECLECTISM, see *Eclectic*.

ECLECTUS, (*εκλεκτος*,) *Eclig'ma*, *Elig'ma*, *Elix'is*, *Eclig'ma*, *Lambit'ium*, *Linctus*, *Linctu'rium*, (*εκλειχω*, (*ec*, and *λειχω*,) 'I lick,') (F.) *Looch*. A medicine, of a thick, syrupy consistence, chiefly used to allay cough, and consisting of pectoral remedies. It was formerly sucked from the end of a liquorice stick, made into a kind of pencil; hence its name *Linctus*, from *lingere*, 'to lick.' Although the linctus is usually exhibited in thoracic affections, it may have tonic virtues combined with it.

ECLEGMA, gen. *Eclig'matis*, (*ελεγμα*,) *Eclectos*—*e. Album*, *Looch album*—*e. Gummoso-oleosum*, *Looch album*.

ECLIPSIS, (*εκλεισις*, from *ec*, and *λειπις*, 'a scale,') *Desquamation*, *Exfoliation*.

ECLIPSITREPANON, (*ecleipsis*, and *τρεπανον*, 'a trepan,') *Exfoliative trepan*.

ECLIGMA, gen. *Eclig'matis*, *Eclectos*.

ECLIMIA, (*εκλιμια*, from *ec*, and *λιμος*, 'hunger,') *Boulimia*.

ECLIPSIS, (*εκλεισις*, from *ec*, and *λειπω*, 'I leave,') *Syncope*.

ÉCLISSE, *Splint*.

ECLYSES, (pl. of *Eclysis*,) *Adynamia*.

ECLYSIS, *Exsolutio*; (ἐκλυσis, from ἐκλω, 'I loosen.') Resolution, prostration of strength; faintness.

ECLYSIS PNEUMOCARDIACA, Asphyxia.

ECMYZESIS, (ἐκμύζησις, from ἐκ, and μύζειν, 'to suck.') Exsuctio.

ECNŒA, (ἐκνῶια, from ἐκ, and νοός, 'mind.') Dementia.

ÉCOLE, (old (F.) *Eschola*, from *Schola*.) School—e. *Organicienne*, see *Organicism*.

ECON'OMY, (οικονομία, *Economía*, (οικία, 'a house,' a family,' and νέμω, 'I rule.') By the term *animal economy* is understood,—the aggregate of the laws which govern the organism. The word *economy* is, also, used for the aggregate of parts which constitute man or animals.

ÉCORCE, Cortex—*É. Caricocastine*, Canella alba—*É. Éluthérienne*, see *Croton cascarilla*—*É. de Saint Lucie*, Cinchonæ Caribææ cortex—*É. de Winter*, see *Winters aromatica*—*É. Fausse de Winter*, Canella alba.

ÉCORCHURE, (ex, and cortex,) Chafing, Excoriation.

ÉCOULEMENT, (ex, and colare, 'to strain or flow.') Discharge, Gonorrhœa—*É. Blanc*, Leucorrhœa—*É. de Sang par l'intestin*, Hæmatochezia.

ÉCOUVILLON, see *Écouvillonnement*.

ÉCOUVILLONNEMENT (F.), (*écouvillon*, 'a kind of mop, the sponge of a gun.') A term used by the French therapeutists for the act of cleansing or applying remedies to a part by means of a mop or brush fixed to the end of a piece of whalebone. Such mop or brush is termed *Écouvillon*.

EPHLOGOSIS, (ἐφλογωσις, from ἐκ, and φλογω, 'I set on fire.') Inflammation.

EPHLYSIS, (ec, and φλυω, 'I boil.') Vesicula—e. Herpes, Herpes—e. Herpes circinatus, Herpes circinatus—e. Herpes exedens, Herpes exedens—e. Herpes miliaris, Herpes phlyctenodes—e. Herpes zoster, Herpes zoster—e. Pompholyx, Pompholyx—e. Rhyphia, Rupia.

EPHRACTIC, (ἐφρακτικός,) Deobstruent.

EPHRAX'IS, (ἐφραξαι, 'I remove obstruction.') The action of ephractic or deobstruent remedies.

EPHRONIA, (ec, and φρον, 'the mind.') Insanity—e. Mania, Mania—e. Melancholia, Melancholy.

EPHYAS, (ἐκφύας, from ἐκ, and φύω, 'I grow.') Appendix vermiformis cæci.

EPHYMA, gen. *Ephy'matis*, (ἐφυμα, from ἐκ, and φύω, 'I grow.') Excrecence, Tumour—e. Callus, Callosity—e. Caruncula, Caruncle—e. Clavus, Corn—e. Edematium, Edema, Phlegmatia dolens—e. Physconia, Physconia—e. Trichoma, Plica—e. Verruca, Verruca.

EPHYMATA, (pl. of *Ephy'ma*.) Rubeola.

EPHYSE'SIS, *Effl'a'tio*, *Effl'a'tus*, (ἐφύσησις, from ἐκφύσσω, 'I breathe through.') *Exsuffl'a'tio*. A quick and forced expulsion of air from the lungs.

EPHYSIS, (ἐφύσις, from ἐκ, and φύω, 'I grow.') Apophysis—e. Ventriculi, Duodenum.

EPHIES'MA, gen. *Ephies'matis*, (ἐπιέσιμα, from ἐπιέω, (ec, and πιέω,) 'I compress.') *Effr'a'ctu'ra*, *Impac'tion*, *Deces'sio*. A fracture of the cranium, with depression of the fragments and compression of the brain.

ECPIES'MUS, (ἐκπιέσιμος,) *Expres'sio*. Celsus uses these words to signify the forcing of the eye from the orbital cavity, with apparent but not real augmentation of the organ. See *Exophthalmia*.

ECPLERO'MA, gen. *Ecplero'matis*, (ἐκπληρωμα, from ἐκ, and πληρω, 'I fill.') A cushion, a pad. Hippocrates means, by this term, a small pad or ball of leather, or other substance intended to fill the hollow of the arm-pit; used probably in reducing luxations of the shoulder.

ECPLEXIA, (ἐκπληξια, from ἐκπλησσω, 'to stun.') Stupor.

ECPLEXIS, (ἐκπληξίς,) Stupor.

ECPNEUMATOSIS, (ec, and πνευμα, 'air.') Expiration.

ECPNEUSIS, (ἐκπνευσις,) Expiration.

ECPNOE, (ἐκπνοή,) Expiration.

ECPTO'MA, gen. *Ecpto'matis*, *Ecpto'sis*, *Ecce'dentia*, (ἐκπτομα, from ἐκπτεω, (ec, and πτεω,) 'I fall out.') This word has been used in various senses, 1. Synonymously with luxation. 2. For the separation of gangrenous parts. 3. For the expulsion of the secundines. 4. For the prolapsus of the womb: and 5. For intestinal or omental hernia.

ECTOSIS, (ἐκπώσις,) Luxation.

ECPYCTICA, (ec, and πυκνάζειν, 'to thicken.') Incrassantia.

ECPYEMA, gen. *Ecpye'matis*, (ἐκπυμα, from ἐκ, and πυν, 'pus,') Abscess, Suppuration, Empyema.

ECPYESIS, (ἐκπυσις,) Abscess, Empyema, Pus-tule—e. Impetigo, Impetigo—e. Porrigo, Porrigo—e. Porrigo crustacea, Porrigo larvalis—e. Porrigo favosa, Porrigo favosa—e. Porrigo furfuracea, Porrigo furfurans—e. Porrigo galeata, Porrigo scutulata—e. Porrigo lupinosa, Porrigo lupinosa—e. Scabies, Psora.

ECPEYETICUS, (ἐκπυητικός,) Suppurative.

ÉCRASEUR, (F.) ('a crusher,' from *écraser*, 'to crush.') *E. Linéaire*, *Histotriteur*, *Sarcotripteur*, *Ligature articulée*, *Sécateur par écrasement*. A surgical instrument invented by M. Chassaingne, of Paris, which consists of a steel chain, like that of a chain-saw devoid of teeth, attached at both ends to a steel mandril, which passes through a hollow steel cylinder. The tightening of the chain is made slowly and with great force, and in the last modification of the instrument is effected by an endless screw with a lever handle, working on a nut coggled on its outer side, which plays on a thread cut on the mandril. The slow bruising it makes is rarely followed by hemorrhage, even in the case of hemorrhoids, and other vascular tumours. The operation with the *Écraseur* has been termed *Écrasement* ou *Broiement linéaire*, *Sarcotripsie*, *Incision ou Amputation sèche*, and *Histotritie*.

ÉCREVISSE, ((G.) Krebs,) Crab.

ECREX'IS, *Ruptu'ra*, (ἐκρηξις, from ἐκρήγνυμι, (ec, and ρήγνυμι,) 'to break.') Rupture, laceration.

ECRHYTH'MUS, (ec, and ρυθμός, 'rhythm.') A term applied to the pulse, particularly when irregular.

ECROË, (εκρη, from ἐκ, and ρέω, 'to flow.') Discharge.

ÉCROUELLES, Scrofula—*É. Mésentériques*, *Tabes mesenterica*.

EC'RYSIS, (εκρυσις, from εκρῶ, 'I run from.') A discharge.

EC'SARCOMA, gen. *Ecsarco'matis*, (εκσάρκωμα, from ἐκ, and σαρξ, 'flesh.') A fleshy excrecence of various kinds. See *Fungosity*, and *Sarcoma*.

ECSES'MA, gen. *Eceses'matis*, *Eczema*.

EC'STASIS, (ἐκστασις, (ec, and ἵστημι, 'I am beside myself.') An *ec'stasy* or *trance*, *Carus ec'sta-sis*, *Cat'ochus*, *Ec'stasis*, *Catalep'sia spu'ria*, *Hyperplexia*, (F.) *Extase*. A state in which certain ideas so completely absorb the mind, that the external sensations are suspended, the voluntary movements arrested, and even the vital actions retarded. In catalepsy, there is, in addition, complete suspension of the intellectual faculties. This last condition is in general described as *trance*. See, also, *Luxation*.

ECSTROPHE, (ἐκστροφή,) *Extrophia*.

ECTASIA, Aneurism—e. Venarum, Varix.

ECTASIS, (εκτασις, from ἐκ, and τέννω, 'stretch.') Extension, Expansion.

ECTASIS IRIDIS is the extension or expansion of the iris, which occasions diminution of the pupil.

ECTEXIS, (εκτῆσις, from *ec*, and *τεκειν*, 'to melt.') Colligation.

ECTHETOBREPHOTROPHEUM, (εκθετος, 'placed out,' and *brephephotropheum*.) Brephephotropheum.

ECTHETOTROPHEUM, (εκθετος, 'placed out,' and *τροφειν*, 'to nourish.') Brephephotropheum.

ECTHILIMMA, gen. *Ecthilim'atis*, *Evulceratio*, (εκθλιμμα, from εκθλιβω, 'I express,' 'I bruise.') Attrition. Chafing, or excoriation, produced by external violence.

ECTHILIPSIS, (εκθλιψις,) Expression.

ECTHYMA or **ECTHY'MA**, gen. *Ecthy'matis*, (εκθυμα, from εκθω, 'I break out.') *Eccy'e'sis*, *Phly'sis* *ecthy'ma*, *Phlyza'cia* *a'gr'ia*, *Thy'ma*, *Scabies* *Ve'ra*, *Furuncul'ic* *aton'ic*, *Pap'ulous* *Scall*, (F.) *Dartre crustacea*, *D. fungusea*. A cutaneous eruption, characterized by large round pustules, always distinct and seated upon an indurated and highly inflamed base. In the course of a day or two the pustules generally break, and olive-brown incrustations are formed which adhere firmly to the skin. These separate in about a fortnight. The disease requires the antiphlogistic treatment. Under *Ecthy'mata*, Vogel has designated certain hard, unequal tumours which appear transitively on the skin. See Efflorescence, Exanthem, and Pustule.

ECTHYSTEROCYESIS, (ecto, *τετρα*, 'uterus,' and *κυσις*, 'pregnancy.') Pregnancy, extra-uterine.

ECTILOLOTICUS, (ec, and *τλλειν*, 'to pluck.') Depilatory.

ECTILMUS, Evulsion.

ECTO, (εκτος, 'outside,') as a prefix, 'outside.'

ECTOBLAST, (ecto, and *βλαστος*, 'a germ.') See Cytoblast and Molecule.

ECTOME, (εκτομη, from *ec*, and *τομη*, 'incision.') Castration, *Eccope*, *Entaille*, *Excision*.

ECTOMIA, Castration, *Eccope*.

ECTOMIAS, (εκτομιας,) Castratus.

ECTOMON, *Helleborus niger*.

ECTOMOS, (εκτομος,) Castratus.

ECTOPA'GIA, (F.) *Ectopagie*, (ecto, and *πηγνυμι*, 'to put together.') A genus of double monsters, having a common umbilicus, comprising those which are united laterally throughout the whole extent of the thorax.

ECTOPARASITES, (ecto, and *parasite*,) Epizoa.

ECTOPHYTE, *Ectoph'yton*, (ecto, and *φυτον*, 'a vegetable.') A vegetable parasite, which grows on the surface of the body.

ECTOP'IA, *Ectop'isia*, *Ectopism'us*, *Entoce'le*, *Si'tus alie'nus* seu *perver'sus* seu *muta'tus*, *Anomotopie* of Piorry, (ec, and *τοπος*, 'out of place.') Morbid displacement of parts. See Luxation.

ECTOPIA ANI, Proctocele.

ECTOPIA CORDIS, *Cardianas'trophē*. Displacement, dislocation, or unnatural position of the heart.

ECTOPIA HERNIOSA, *Hernia*—e. *Splenis*, *Splenectopia*.

ECTOPOCYST'ICUS, (εκποσος, 'out of place,' and *κυστις*, 'bladder.') A disease dependent upon displacement of the bladder.

ECTOPROTIC, (improperly for) *Eccoprotic*.

ECTOZO'A, (sing. *Ectozo'on*, *Ectozo'a*, *Ectozoa'ria*, (F.) *Ectozoa'ires*; (ecto, and *ζωον*, 'an animal.') Parasitic animals that infest the exterior of the body,—as lice: a term which, like *Helminthia errat'ica*, *Pseudohelmin'thes*, and *Pseudopar'asites*, is applied, also, to worms or larvae of insects that have been introduced into the intestinal canal by accident. Animalcules, most frequently swallowed, are the hairworm, leech, grub of the fly, caddy insect—*Phalæ'na pinguin'alis*; the

larve of the bee, the spider, the *triton palu'stris*, *lacer'ta aquat'ica*, &c. In animals, bots are produced by swallowing the ova of the *centrus* or gad-fly. See *Helminthia erratica*.

ECTRIM'MA, gen. *Ectrim'matis*, (εκτριμμα, from *εκτριβω*, (ec, and *τριβω*,) 'I rub off.') Ulceration of the skin; and particularly that which arises from the pressure of the bed on different parts of the body, after a protracted confinement.

ECTRODACTYL'IA, (εκτροσις, 'abortion,' and *δακτυλος*, 'a finger.') A malformation, in which one or more fingers or toes are wanting.

ECTROG'ENY, *Ectrogen'ia*, *Ectrogen'esia*, (εκ-τροσις, 'abortion,' and *γενεσις*, 'generation.') Monstrosity by defect.

ECTRO'MA, gen. *Ectro'matis*, (εκτωμα,) Abortion.

ECTRO'MELES, (εκτροσις, 'abortion,' and *μελος*, 'a limb.') A genus of monsters, in which the limbs are nearly or altogether deficient, as in the ordinary catæcia.

ECTROPE, *Divertie'ulum*, (εκτροπω, (from *ec*, and *τροπω*,) 'I turn off,' 'divert.') Any duct by which peccant or morbid matter was supposed to be drawn off.

ECTROP'ION, *Ectrop'ium*, *Evert'sio pal'pebræ*, *Blepharopto'sis* *Ectrop'ium*, *Blepharoto'sis*, *Pal'pebræ infe'rior extro'r'sum flex'a*, *Divaricat'io* seu *Reflex'io* seu *Reclina'tio palpebra'r'um*, (F.) *Éva'issement des Paupières*, *Renversement des Paupières*. Eversion of the eyelids, so that they do not completely cover the globe of the eye. It happens more commonly to the lower than to the upper eyelid. It may be owing to the retraction of the skin, after the cure of an ulcer, wound, or burn of the eyelid; or it may depend on tumefaction or relaxation of the conjunctiva. In the majority of cases, removal of a portion of the conjunctiva will effect a cure; but there are many which defy the efforts of art. The ancients called Ectropion of the upper eyelid *lagophthal'mia*.

ECTROSIS, (εκτροσις,) Abortion.

ECTROSMUS, (εκτροσμος,) Abortion.

ECTROT'IC, (from *ectroma*, (ec, and *τιρωσκω*, 'I wound,' 'abortion.') *Ectro'ticus*, *Abort'ive*. An epithet applied to methods for preventing the development or causing the abortion of any disease—as of chancres by the use of caustic; small-pox pustules by the use of mercurial ointment, &c.

ECTYLOTICUS, (ec, and *τυλος*, 'a protuberance.') Cathartic.

ECTYMPANOSIS, (εκτυμπωσις,) Tympanites.

ECUMEUX, (same root as *scum*,) Frothy.

ÉCUSONS, (from *ecu*, old (F.) *écu*, (L.) *scutum*, 'a shield.') 'Escutcheons or shields.' Plasters spread upon the skin; or small bags—*sachets*—of the shape of escutcheons, filled with odorous powders, which are applied on the skin. See *Sachets*.

EC'ZEMA, gen. *Eczem'atis*, (εκζεμα, from εκζεω, (ec, and *ζεω*,) 'I boil out,' 'I effervesce.') *Eczes'ma*, *Eczes'ma*, *Pus'tula ar'dens*, *Cytis'ma* *Ec'zema*, *Hu'mid Scall* or *Tet'ter*, *Run'ning Scall*, (F.) *Dartre squameuse humide*, *D. vive*, *Gale épidémique*. Heat eruption. An eruption of small vesicles on various parts of the skin, usually set close or crowded together; with little or no inflammation around their bases, and unattended by fever.

ECZEMA CAP'ITIS, E. of the hairy scalp.

ECZEMA OF THE FACE, at an advanced stage, and occurring in young children, has been described under the names *Crusta lactea* and *Por-rigo larvalis*.

ECZEMA OF THE HAIR'Y SCALP, *Ec'zema cap'i-tis*, *Vesic'ular scall*, is often confounded with other affections, under the names *Por-rigo* and *Tinea*, which are pustular, not vesicular, in their form. It occurs during dentition, and even after-

wards, and the discharge is so profuse, that the head appears as if dipped in some glutinous liquid. By and by, the secretion dries into crusts and mats the hair into little separate tufts. The scalp gives evidence of inflammatory excitement, and the lymphatic ganglions of the neck are apt to become inflamed and suppurate.

A variety of humid scalled head, in which the humour from the exoriated surface runs down upon the hairs, and encloses them in little silvery pellicles or sheaths, has received the name *Asbestos Scall*.

ECZEMA IMPETIGINODES, *Gall, Gro'cers'* or *Bricklayers' Itch*. Produced by the irritation of sugar or lime.

ECZEMA MERCURIALĒ, *E. ru'brum, Erythema mercuria'le* seu *ichoro'sum, Hydrargyria, Hydrargyrosia, Hydrargyri'asis, Morbus mercuria'lis, Mercu'rial lepra* or *rash*. A variety of eczema, arising from the irritation of mercury. The treatment is chiefly palliative, consisting in ablation with mucilaginous infusions or decoctions; mild dressings, where the cuticle has exfoliated; avoiding all irritation; keeping the bowels open; with the use of sulphuric acid and cinchona.

ECZEMA RUBRUM, *Eczema mercuriale*.

ECZEMATOSSES, (*G.*) *Eczematosen*. A family of diseases, in the classification of Fuchs, including morbid conditions of the cutaneous secretions—as of the perspiration, sebaceous and colouring matters, &c., and hence many chronic cutaneous affections. His subdivisions are, *ephidro'ses, smegmorrhœ'a, acar'pæ, polycar'pæ*, and *monocarp'pæ*.

ECZESIS, (*εκζέσις*.) Effervescence.

ECZESMA, gen. *Ecze'smatis*, (*εκζεσμα*.) *Eczema*, *Lichen tropicus*.

EDDOES, see *Arum esculentum*.

EDEMATOUS, *Edematous*.

EDENTATUS, *Edentulus*.

EDENTULI, *Nefrendes*.

EDENTULOUS, *Eden'tated, Eden'tulus, Edenta'tus*, (*e*, and *dens*, gen. *dentis*, 'a tooth.') *An'odus, Ca'rens dentibus, No'des, No'dus*, (*F.*) *Édenté*. One without teeth. This defect can only be remedied by artificial means. See *Nefrendes*.

EDERA, *Hedera helix*.

EDIBLE, (from *edere*, 'to eat,' and *habilis*, 'able,') *Esculent*.

ÉDOCEPHALE, (*αἰδοῖα*, 'the sexual organs,' and *κεφαλή*, 'head.') A monster, which has the two ears near each other, or united under the head, the jaws atrophied; no mouth, and above the eye a tube (*trompe*) resembling a penis.

EDROPPIT (*Sc.*), *Hydropic*.

EDUCATIO INFANTUM, ('bringing up of children,') *Pædia*.

EDULCORA'TION, *Glycan'sis, Edulcora'tio*, (*e*, and *dulcis*, 'sweet.') An operation, the object of which is to deprive a substance of its acrid and disagreeable taste, or at least to disguise it. Also, the addition of a saccharine substance to a medicine, whose taste it is desirable to modify agreeably.

ÉDULE, (*edere*, 'to eat,') *Comestible*.

EDULIS, *Comestible*.

EE (*Sc.* and *Prov.*), *Eye*.

EEL GRASS, *Pila marina*.

EESKIN (*Sc.*), *Singulus*.

BEWINKER (*Sc.*), *Palpebra*.

EE, as a prefix, like *e*, and *ex*.

EFFERENT, *Eff'ferens, De'vehens, Centrif'ugal, Exod'ic, (effero* (*e*, and *fero*), 'I carry,' 'transport,') Conveying outwards, as from the centre to the periphery.

Vasa efferen'tia are those lymphatic or chyliferous vessels which issue from the glands to convey their lymph to the thoracic duct; so called to distinguish them from those which pass to those

glands, and which have been termed *vasa afferen'tia* seu *inferen'tia*. Also, nerves that convey the nervous influence from the nervous centres to the circumference. See *Aferent*.

At the upper extremity of the mediastinum testis, the ducts of the rete testis terminate in from 9 to 30 small ducts, called *vasa efferen'tia* seu *Graafia'na*, (*F.*) *Vaisseaux éfférents*, which form the *Co'ni vasculo'si*.

EFFERVES'CENCE, (*Effervescentia, Ze'sis, Ec'zesis, (effervescere* (*ef*, and *fervescere*), 'to grow hot,') That agitation, which is produced by the escape of gas through a liquid, independently of the heat of the mixture; such, for instance, as results from the mixture of acetic acid and carbonate of potassa.

In *Pathology* it has a similar signification. It expresses, in the language of the humorists, a sort of ebullition in the fluids of the living body, produced either by elevation of temperature or by the reaction on each other of the principles contained in the fluids in circulation.

EFFETUS, (*ef*, and *fetus* or *fatus*, 'producing,') *Impoverished*.

EFFICA'CIOUS, *Eff'ficax, (efficere* (*ef*, and *facere*), 'to accomplish,') That which produces a great effect,—as 'an efficacious remedy.'

Medic'ina ef'ficax, La Médecine efficace, is a term sometimes applied to surgery.

EFFILA, *Ephelides*.

EFFLATIO, *Eophyses*.

EFFLATUS, gen. *Effla'tūs*, (*ef*, and *flare, flatum*, 'to blow,') *Eophyses*.

EFFLORATIO, (*ef*, and *floreo*, 'to bloom,') *Exanthem*.

EFFLORESCENCE, *Efflora'tio, Efflorescentia*, (from *efflorescere* (*ef*, and *florescere*), 'to blow as a flower,') *Stribili'go, Ec'thyma*.

In *Pathology*, efflorescence has the same meaning as *exanthema*; and, in the nosology of Sauvages, the name is given to that order of diseases. Sometimes, it is confined to the cutaneous blush, the *exanth'e'sis* of Good.

Efflorescence is, also, the conversion of a solid substance into a pulverulent state by exposure to the air. In salts this is generally owing to the loss of a part of their water of crystallization.

EFFLORESCENCE ÉRYSIPÉLATEUSE, *Roseolæ*.

EFFLORESCENTIA, *Efflorescence, Exanthem*.

EFFLORESCENTIÆ, *Élevures*.

EFFLUVIUM, (*ef*, and *fluere, fluxum*, 'to flow,') *Emanation*—*e. Latrinarium, Mitte*—*e. Palustre, Miasm, marsh*.

EFFLUXION, see *Abortion*.

EFFORT, (*ef*, and *fortis*, 'strong,') *Ni'sus, In'nat'us, Pei'ra*. A muscular contraction of greater or less strength, the object of which is, either to resist an external force, or to accomplish a function, which has become naturally laborious:—such are, the act of pushing away, or of drawing a body towards us, and the more or less painful efforts used by the mother to cause the expulsion of the fetus. In France, the word *effort* is often used synonymously with *hernia*; and signifies, likewise, the painful twitches of muscles, occasioned by over-exertion, or by the rupture of some of their fleshy fibres. Sauvages calls *Efforts des reins, Lumbago à ni'su*, the pain in the loins occasioned by bearing too heavy a burden.

EFFORT, *Hernia*—*e. des Reins*, see *Effort*.

EFFOSSIO, (*ef*, and *fodere, fossum*, 'to dig,') *Exhumation*.

EFFRACTURA, (*ef*, and *frangere, fractum*, 'to break,') *Ecpiesma*.

EFFRACTURA CRA'NI, *En'thlasis Cra'nii, Fracture of the Cranium*, with depression.

EFFRENTATIO, (*effrenis*, 'unbridled,') *Hypercatarsis*.

EFFUSIO, *Effusion*—*e. Seminis, Ejaculation*.

EFFU'SION, *Effu'sio*, *Ecchysis*, (*effundere* (*ef*, and *fundere*, *fuum*), 'to pour out,') (F.) *Épanchement*, (*Infiltration* is the term generally employed for effusion into the areolar membrane. The pouring out of blood or of any other fluid into the areolar membrane, or into the cavities of the body. The *effusion of serum* or of *coagulable lymph*, for instance, is a common result of inflammation of serous membranes.

ÉGARÉ, Wild.

ÉGAREMENT D'ESPRIT, Delirium, Insanity.

EGER, MIN'ERAL WA'TERS OF THE. In the valley of the Eger, at the western extremity of Bohemia, there are several acidulous springs. One of the most frequented of these is Franzensbad.

EGE'RIA. In ancient mythology, a nymph to whom pregnant females offered sacrifices *ut conceptus albus facilius egeretur*. By some, supposed to have been identical with Lucina.

EGESTA, (*e*, and *gero*, *gestum*, 'to carry,') see Excretion.

EGESTAS, Want.

EGESTIO, Defecation, Dejection, Excretion.

EGG, (Sax. æs.) Ovum—e. Bed, Ovarium—e. Brandy, see Ovum—e. Plant, *Solanum melongena*.

Egg, WHITE OF, Albumen ovi.

EGLANTIER DE CHIEN, *Rosa canina*—e. *Sauvage*, *Rosa canina*.

EGOBRONCHOPHONY, see Egophony.

ÉGOPHONÉ, Egophony.

EGOPHON'IC, or **EGOPH'ONIC**, *Ægophonicus*, (F.) *Égophonique*. Having the character of, or relating to, egophony.

ÉGOPHONIQUE, Egophonic.

EGOPH'ONY, *Ægopho'nia*, (αἴ, gen. αἴος, 'a goat,' and φωνή, 'voice,') *Caprilo'quium*, *Tragopho'nia*, *Goat's Voice*, *Bleat'ing Voice*, (F.) *Égophonie*, *Voix chévro'tante* ou *égophonique* ou de *Polichinelle* ou *sénile*, *Pectorilo'quie chévro'tante*. Laënnec has designated by this name, the kind of resonance of the voice heard through the stethoscope, when we examine the chest of one labouring under moderate effusion into one of the pleuræ. The voice, which strikes the ear through the cylinder, is more sharp and harsh than usual, and tremulous and broken, like that of the goat. The patient himself is called *Égophone*. Bouillaud affirms, that the 'bronchial and bleating voice' (*égobronchophonie*), is the principal symptom of pleuropneumonia. Egophony exists, however, in cases of hepatization where there is no pleural disease.

EGREGOR'SIS, *Vigil'ia*, *Vigil'ie*, *Vigilant'ia*, *Vigila'tio*, *Vigil'ium*, (*εγρηγορῆς*, from *εγρηγορῶ*, 'I watch,') Watchfulness. A morbid want of sleep.

EGRESSUS VENTRICULI, Pylorus.

EGRITUDE, (*ægritudo*, from *æger*, 'sick,') Disease.

EIDOS, (*εἶδος*, 'form, resemblance,') The *εἰ* is sometimes changed into *ω*, at the termination of a word. Thus, *Hæmato'des* for *Hæmato'i'des*. See Odes.

EILAMIDES, (*εἰλαμίδες*, from *εἰλεω*, 'I involve,') Meninges.

EILE'MA, gen. *Eile'matis*, (*εἰλημα*, from *εἰλεω*, 'I roll,') 'a convulsion.' Vogel has given this name to a fixed pain, occupying some portion of the intestinal canal, which the patient compares to the sensation that would be produced by a nail driven into the part.

EILEON, (*εἰλεον*,) Ileon.

EILEUS, Ileus.

EILOID (*Tu'mour*,) *Eilo'i'des*, (*εἰλεω*, 'I roll,')

and *eidos*,) A morbid growth of the cutis, coiled or folded.

EILSEN, MIN'ERAL WA'TERS OF. Eilsen is about six German miles from Hanover, at the foot of the Harrelberg. It has eleven springs, of which seven are sulphureous and four chalybeate.

EISANTHE'MA, gen. *Eisanthe'matis*, *Exanthe'ma inter'num*, *Eutanthe'ma*, (*εἰς*, 'within,' and *αὔθημα*, 'efflorescence,') An eruption on a mucous membrane;—aphthæ, for example.

EISBOLE, (*εἰσβολή*, from *εἰς*, 'within,' and *βάλλειν*, 'to throw,') Attack, Injection.

EISEL, Acetum.

EISPNOE, (*εἰσπνοή*, from *εἰς*, 'within,' and *πνέω*, 'I breathe,') Inspiration.

EJACULATIO, Ejaculation—e. *Seminis Impedita*, *Bradyspermatismus*.

EJACULA'TION, *Ejacula'tio*, (*e*, and *jaculo* (itself from *jacere*,) 'I throw out,') *Gonobol'ia*, *Gonobol'ismus*, *Ejaculatio* seu *Profusio* seu *Effusio Sem'inis*, *Expatria'tio*, *Patra'tio*, *Spermob'olē*. The emission of sperm. That which occurs during coition has been termed *Insemination*. The act, by which the fluid is projected out of the urethra.

EJACULA'TOR. That which effects the emission of sperm. See *Transversus perinæ*.

EJACULATOR SEMINIS, Accelerator urinæ.

EJAC'ULATORY, *Ejaculato'rius*, *Ejac'ulans*; (F.) *Éjaculateur*, *Éjaculatoire*. Concerned in the ejaculation of sperm.

EJACULATORY DUCTS OR CANALS, *Ductus ejaculato'rii*, (F.) *Conduits ou Canaux éjaculateurs*, are formed by the union of the vasa deferentia with the ducts of the vesiculæ seminales. They open at the lateral and anterior parts of the verumontanum, and convey into the urethra the sperm which is discharged from the vesiculæ, as well as that which comes directly from the testicle by the vas deferens. Between them there is often a depression, sometimes of a large size, which is termed *Utriculus*, *U. prostat'icus*, *Vesic'ea* seu *Vesic'ula prostat'ica*, *Sinus pocula'ris* seu *prostat'æ*, *Corpusculum Weberianum*, *Weberian organ* or *corpuscle*, which has been regarded as the analogue to the uterus in the female, and thence called *Uterus mascul'inus*.

EJECTIO, Ejection, Excretion—e. *Fæcum*, Defecation.

EJECTION, *Ejec'tio*, (*ejicere*, *ejectum* (*e*, and *jacere*), 'to throw out or eject,') The excretion of the fæces, urine, sputa, &c.

ELNISPERO, Sapota.

ELABORATION, *Elabora'tio*, (*e*, and *laborare*, *laboratum*, 'to work,') This word is used, by physiologists, to signify the various changes which substances susceptible of assimilation undergo, through the action of living organs, before they are capable of serving for nutrition. The food is said to be *elaborated* in the stomach during the formation of chyme; the chyme is *elaborated* in the small intestine before it is chyle, &c.

ELÆA, (*ελαία*,) Olea Europæa.

ELÆAGNUS, (*ελαία*, 'olive,' and *αγνος*, 'chaste,') Myrica gale—e. Cordo, Myrica gale.

ELÆOCARPUS COPALLIFERUS, (*ελαία*, 'olive,' and *καρπος*, 'fruit,') Copal.

ELÆOCÉROLÉ, (*ελεων*, and *κηρος*, 'wax,')

ELÆOM'ELI, (*ελεων*, and *μελί*, 'honey,') Dioscorides means, by this, an oil thicker than honey, and of a sweet taste, which flows from the trunk of a tree in Syria. It is acrid and purgative; and sometimes occasions serious symptoms, according to that author.

ELÆOM'ETER, (*ελεων*, and *μετρον*, 'measure,') A very delicate glass hydrometer, for testing the purity of olive or almond oil by determining their densities. The 0 or zero of the scale is the

point at which the instrument floats in the oil of puppy seeds. In pure olive oil, it floats at 50°, and the space between these points is divided into 30 equal parts, and numbered accordingly. It floats at 38° or 38½° in pure oil of almonds.

ELÆON, (ἐλαῖον,) Oil.

ELÆOPH'ANES, (ἐλαῖον, and φαῖνομαι, 'I appear.') That which has the appearance of oil; as ἑλαιοφάνης.

ELÆOSAC'CHARUM, (ἐλαῖον, and σάκχαρ, 'sugar.') Ὀλέοςaccharum, Βαλ'σαμῶοςaccharum, (F.) Oléosaccharolé. A medicine, composed of essential oil and sugar. It was made by pouring upon an ounce of sugar, reduced to an impalpable powder, 3ss or ʒj of an essential oil; the mixture being triturated until the two substances were perfectly united.

ELÆOTHES'TON, Uctua'rium, (ἐλαῖον, and θέσις, 'the act of putting or placing.') The oil chamber in the ancient gymnasia, where the body was rubbed with oil before commencing the exercises.

ELAIN, (from ἐλαῖον, 'oil.') See Fat.

ELAIS GUINEENSIS, (from ἐλαῖα,) see Cocos butyrea—e. Occidentalis, see Cocos butyrea.

ÉLAN, ('a leap,') Cervus alces.

ÉLANC'EMENT, (e, and lancer, 'to dart.') See Lancing.

ELAPHOBOSCUM, (ἐλαφος, 'a deer,' and βοσκειν, 'to eat.') Pastinaca sativa.

ELAPHOM'YCES GRANULA'TUS, E. officina'lis, (ἐλαφος, 'a stag,' and μυκή, 'a fungus.') Lycoperdon cervinum, Cer'vi Bole'tus, Bole'tus cervinus, Tu'ber cervinum, Hart's Truf'fles, Deer Balls, Lycoper'don nuts. Nat. Order, Fungi. Formerly used in the preparation of a Bal'samus apople'cticus, and to promote labour and the secretion of milk. Given, also, as an aphrodisiac. Dose, ʒiss, in powder.

ELAPHURIM ELEMIFERUM, (from ἐλαφος, 'light,' applied to the wood.) See Amyris elemifera—e. Tomentosum, Fagara octandra.

ELAPSUS, (e, and labi, lapsus, 'to fall.') Luxation.

ELAS'MA, gen. Elas'matis, (ἐλαυνω, 'I impel.') A clyster pipe.

ELASTES, (ελαστος, 'a driver.') Elasticity.

ELASTIC MEMBRANE, see Membrane.

ELASTICINE, see Fibrous.

ELASTIC'ITY, Elastic'itas, El'ater, Elas'tes, El'ates, Vis elas'tica, Tone, Tonic'ity, (F.) Contractilité de Tissu, Contractilité par défaut d'extension, (ἐλαυνω, 'I impel,' ελαστος, 'a driver.') The property by which certain bodies return to their proper size and shape, where these have been modified by pressure or otherwise. It is possessed by the dead as well as by the living solid.

ELATE, (ελατη,) Pinus picea—e. Theleia, Pinus rubra.

ELATER, (ελατηρ,) Elasticity.

ELATERIN, see Momordica elaterium.

ELATERIUM, (ελατηρ, 'a driver.') Extractum elaterii, Momordica elaterium—e. Cordifolium, Momordica elaterium.

ELATES, (ελαστος,) Elasticity.

ELATIN, see Momordica elaterium.

ELATINE, Antirrhinum elatine—e. Cymbalaria, Antirrhinum linaria—e. Hastata, Antirrhinum elatine.

ELBOCK (Sc.), Elbow.

EL'BOW, (Sax. elboȝa) the bend of the ulna. Cu'bitus, An'con, Pe'chys, Ul'na, Um'bo, (Sc.) El'bock, El'buck, (F.) Coude. This word, abstractedly, means the angle formed by the union of two straight bodies. It is particularly applied to the articulation of the arm with the forearm, and especially to the projection formed by the olecranon process at the posterior part of the

joints. The anterior part of the joint is called the bend of the elbow, (F.) Pli du coude.

ELBOW, MINERS', see Miners' Elbow.

ELBUCK (Sc.), Elbow.

ELCAJA, Trichilia emetica.

ELCOMA, gen. Elco'matis, Ulcer.

ELCOPLASTY, Heleoplasty.

ELCO'SIS, Heleco'sis, Elco'ma, Helco'ma, (ἐλεος, 'an ulcer,' and osis.) Ulceration in general. An ulcer. A deep ulceration of the cornea, in consequence of a blow, or of violent inflammation.

Sauvages applies the term to a state of the body, in which there are numerous ulcerations of an obstinate character, complicated with caries, putrescency, low fever, &c.

ELCUSTER, (ἐλευστηρ, from ἐλεω, 'I draw.') Embryuleus.

ELDER, (Sax. ellarn, (G.) Holder,) Sambucus—e. Common, Sambucus Canadensis—e. Dwarf, Aralia hispida, Sambucus elulus—e. Prickly, Aralia spinosa—e. Redberried, Sambucus pubens.

ELECAMPANE, (Eoula campana,) Inula helianum.

ELECTARIUM, Electuarium.

ELECTARY, Electuarium.

ELECTRICITAS, Electricity—e. Animalis, Galvanism—e. Galvanica, Galvanism—e. Metallica, Galvanism.

ELECTRIC'ITY, Electric'itas, (from ηλεκτρον 'amber,'—the substance in which it was first noticed.) Electricity is used medicinally as an excitant. It has been occasionally employed with success in paralysis, rheumatism, accidental deafness, amaurosis, amenorrhœa, &c, but is not extensively used; and the cases are not always clear in which it could be of service. It may be communicated by means of the electric bath—Bain électrique, as it has been called; which consists in placing the patient upon an insulated stool, and connecting him with the prime conductor, of which he thus becomes a part. The fluid may be communicated by points, sparks, or by shocks, according to the required intensity. See Electrization.

ELECTRICITY, CHEMICAL, Galvanism—e. Contact, Galvanism—e. Magnetic, Electromagnetism—e. Voltaic, Galvanism.

ELEC'TRIFY, Elec'trize, (electrical, and fio, 'to become,') (F.) Électriser. To produce the electrical condition in a body, or to render it susceptible of producing electrical phenomena. It is often used to signify the act of communicating the electric fluid to man.

ÉLECTRISER, Electrify.

ELECTRIZA'TION, Electriza'tio, (F.) Électrization. The medical application of electricity.

Localized electrization, (F.) Electrization localisée, of Duchenne, of Boulogne, consists in concentrating the electric action in particular parts without electro-puncture or any surgical operation.

It is founded on the greater or less resistance presented by a tissue to electric currents, according to its state of dryness or of moisture; and may be practised by using wet sponges or a metallic knob covered with a piece of wet kid for deep-seated structures, as muscles or any internal viscus; and dry metallic bodies of varying shape, or a metallic brush, for the purpose of cutaneous electrization. Electrization may also be employed for diagnostic purposes; for example, the atrophied muscle of rheumatism contracts well under the galvanic stimulus, whilst the opposite is the case in fatty transformation of the muscular tissue. See Galvanization, localized.

ELECTRO, as a prefix, Electricity.

ELEC'TRODE, (electro, and δῶς, 'way,')

(F.) *Rhétrophore* ou *Réophore*. The poles of a galvanic battery are called electrodes.

ELECTRO'DES, (ηλεκτροῦδες, from ηλεκτρον, 'amber,' and οδες. An epithet for evacuations, which shine like amber.

ELECTROLITHOT'RY. The disintegration of calculi in the bladder by the mechanical force of the electrical discharge, applied by an appropriate apparatus.

ELECTROMAGNETISM. An electro-magnetic apparatus is occasionally used in cases of paralysis, or when a powerful excitant has to be applied to a part of the organism. A convenient form consists of a battery of six curved permanent magnets, and an intensity armature, around whose cylinders 1500 yards of fine insulated copper wire are coiled. The ends of the wire communicate respectively, with a pair of directors, each of which holds a piece of sponge, dipped in vinegar or a solution of common salt. When the armature is rotated, and a portion of the body is interposed between the directors, a succession of shocks is received.

ELECTROPUNCTURE, *Electropunctura'tion*, *Electropunctu'ra*, *Electrostix'is*, and *Galvanopuncture*, *Galvanopunctu'ra*. The operation of inserting two or more wires, and then connecting them to the electrodes of the electric or galvanic apparatus. It has been employed therapeutically in cases in which electricity, galvanism, and acupuncture have been indicated.

ELECTROSTIXIS, (electro, and σιξίς, 'punctuation,') *Electropuncture*.

ELECTRUM, (ηλεκτρον,) *Succinum*.

ELECTUAIRE, *Electuarium*—*e. Dentifrice*, *Dentifrice*—*e. Simple*, *Conserva*.

ELECTUARIUM, *Electa'rium*, *Opiat'um*, (Old Eng.) *Allectuary*, *Electuary*, *Electuary*, (F.) *Electuaire*, *Saccharolé mou*, (*eligere*, *electum*, 'to make choice,') A pharmaceutical composition of a soft consistency, somewhat thicker than honey, and formed of powders, pulps, extracts, syrup, honey, &c. In the London and American Pharmacopœias, electuaries are classed under Confections.

ELECTUARIUM DE ALOË, *Opiatum mesentericum*—*e. Aperient*, *Confectio sennæ*—*e. Aromaticum*, *Confectio aromatica*—*e. Cassiæ*, *Confectio cassiæ*—*e. Cassiæ fistulæ*, *Confectio cassiæ*—*e. Cassiæ sennæ*, *Confectio sennæ*—*e. Cassiæ tamarindatum* seu *lenitivum*, *Confectio cassiæ*—*e. Catechu*, *Confectio catechu compositum*—*e. Catholicum commune*, *Confectio sennæ*—*e. Cinchonæ*, *Opiatum febrifugum*—*e. de Croco emendatum*, *Confectio hyacinthi*—*e. Diaprunum*, *Confectio sennæ*—*e. Eecoproticum*, *Confectio sennæ*—*e. de Kinā kinā*, *Opiatum febrifugum*—*e. Laxativum*, *Confectio cassiæ*—*e. Lenitivum*, *Confectio sennæ*—*e. Opiatum*, *Confectio opii*—*e. Piperis*, *Confectio piperis*—*e. Scammonii*, *Confectio Scammonie*—*e. à Scammonio*, *Confectio scammonie*—*e. Sennæ*, *Confectio sennæ*—*e. Sennæ compositum*, *Confectio sennæ*—*e. Sennæ cum pulpis*, *Confectio sennæ*—*e. Solidum*, *Tabella*—*e. Thebaicum*, *Confectio opii*.

ELECTUARY, *Electuarium*—*e. Lenitive*, *Confectio sennæ*.

ELÉENCÉPHALE, (ελαῖον, 'oil,' and 'encephalon,') A fatty matter found by Couerbe in the encephalic neurine.

ELELISPHACUS, (ελελίσφακος,) *Salvia*.

ELEMENT, *Elementum*, (akin to ancient *eleo*, for *oleo*, 'to grow' [?],) *Principium*, *P. Primitivum*, *Stoicheion*. A simple, ultimate constituent or principle in the human body, which forms the basis of a fibre or tissue. Also, a constituent of a compound organ. The *inorganic elements* are simple principles. An *organic element*, *proximate principle* or *compound of organization*, results from the union of certain inorganic elements.

Oxygen, hydrogen, carbon, and nitrogen, are inorganic elements; fibrin, albumen, casein, &c, organic elements.

ELEMENTS, *ORGANIC*, Principles, immediate—a Sarcous, see Sarcous.

ELEMENTARY CELL, see Cell.

ELEMENTUM, *Element*—*e. Acidificum*, *Oxygen*.

ELEMI, *Amyris elemifera*.

ELENIMUM, *Inula helenium*.

ELEO'CHARIS PALUS'TRIS, (ἐλεος, 'a marsh,' and χαίρω, 'to delight in,') *Scirpus palustris*; indigenous. *Ord.* Cyperacæ. The flowering tops, seeds, and roots are astringent, and have been used in diarrhœa and hemorrhage.

ÉLÉOLÉS, (ελαῖον, 'oil,') *Olea medicinalia*.

ELEOSELINUM, (ἐλεος, 'a marsh,' and σελινον, 'parsley,') *Apium petroselinum*.

ELEPHANTIASIS, (ελεφαντίας,) *Elephantia*, *Elephantismus*, *El'ephas*, *Laz'ari mor'bus* seu *ma'lum*, *Pachyder'mia*, *Phœnic'eus mor'bus*, *Phymato'sis* *Elephantiasis*, (ελεφας, gen. ελεφαντος, 'an elephant,') Various affections have been described under this name, by adding an epithet. It is ordinarily and generally applied to a condition, in which the skin is thick, livid, rugous, tuberculate, and insensible to feeling.

ELEPHANTIASIS OF THE ANTILLES, *Barba'does Leg*, *Glandular disease' of Barba'does*, (F.) *James de Barbade*, is the Elephantiasis of many writers, *Buene'mia*. It is characterized by the leg being enormously tumid and misshapen; skin thickened, livid, and rugous, often scaly; scrotum, arms, or other parts sometimes participating in the affection. The *Buene'mia Tropica*, *Cochin leg*, is an affection of this kind.

ELEPHANTIASIS ARABICA, *Tyri'asis*, *El'ephas*, *Elephantiasis*, *E. Indica*, *Elephantia Arabum*, *Le'pra Arabum* (of some), *Maladie glandulaire*, *Laderrie*, *Yava Skin* of the Polynesians [?]. In this the tubercles are chiefly on the face and joints. There is loss of hair, except on the scalp; voice, hoarse and nasal; and the disease is said to be contagious and hereditary. It most frequently attacks the feet; and gives the lower extremity a fancied resemblance to the leg of an elephant, whence its name. The seat of the disease seems to be in the lymphatic vessels and glands, and in the subcutaneous areolar tissue: the inflammatory condition of which is accompanied with general morbid symptoms. Medicine has little or no effect on this loathsome complaint. See *Lepra*.

ELEPHANTIASIS OF CAYENNE, *Mal rouge de Cayenne*, seems to be a variety of *Lepra*, characterized by red and yellow spots, occupying the forehead, ears, hands, loins, &c., afterwards extending and becoming scaly, with deformity of the parts where they are seated, particularly of the face; and ultimately producing cracks, ulcers, caries, and sometimes death.

ELEPHANTIASIS OF THE GREEKS, *E. Græcorum*, is probably the same disease as *Lepra*.

ELEPHANTIASIS OF INDIA is characterized by red, livid, or yellowish spots, slightly prominent, to which succeed indolent tumours, formed in the areolar texture. At a more advanced period, the phalanges swell, and become ulcerated; the bones of the nose carious, the lips thickened; and emaciation gradually carries off the patient. For this form [?] the term *Spilopla'zie* was proposed by Duchassaing. It, likewise, belongs to *lepra*.

ELEPHANTIASIS ITALICA, *Pellagra*.

ELEPHANTIASIS OF JA'VA is likewise a variety of *lepra*, characterized by large white tumours on the toes and fingers, resembling scrofulous tumefactions. These tumours ulcerate, and the ulcerations spread from the extremities towards the trunk, destroying even the bones. Amputation

can alone arrest its progress. The disease is often accompanied by alopecia, and by an eruption of red spots.

ELEPHANTINUM EMPLASTRUM, (Ἐλεφαντινόν.) An ancient plaster, composed of 30 parts of *ceruus*, 45 of wax, oil lbs, water lbj. Oribasius and Celsus both mention a plaster of this name, but they are by no means similar in composition.

ELEPHANTOMMA, gen. *Elephantom'matis*, (ἑλεφας, 'elephant,' and ὄμμα, 'eye.') Buphthalmia.

ELEPHANTOPUS, (F.) *Éléphantope*; (ἑλεφας, gen. ἑλεφαντος, 'elephant,' and πους, 'foot.') One affected with elephantiasis.

ELEPHANTUSIA, (ἑλεφας, 'ivory.') *Phytelephas macrocarpa*.

ELEPHAS, gen. *Elephan'tis*, (ἑλεφας,) Elephantiasis. Ivory.

ELETTARIA CARDAMOMUM, *Amomum cardamomum*, and *Cardamomum*.

ELEUTHERIA, *Croton cascarilla*.

ÉLEVATEUR, Elevator.—*É. Commun de l'aile du nez et de la lèvre supérieure*, Levator labii superioris alæque nasi.—*É. de l'Œil*, Rectus superior oculi.—*É. de la Paupière supérieure*, Levator palpebræ superioris.

ELEVATIO UTERI. Ascent of the uterus. This displacement occasionally occurs, in the unimpregnated state, in connection with fibrous uterine, and enlarged ovarian tumours.

ÉLEVATOIRE, Elevator.

ELEVATOR, *Leva'tor*, (elevare, (e, and levare,) 'to lift up.') (F.) *Élévateur*. A muscle, whose function it is to raise the part into which it is inserted. See Levator.

ELEVATOR, *Elevato'rium*, *Vec'tis elevato'rius*, *Élévatoire*. A name given to different surgical instruments employed for raising portions of bone which have been depressed, for raising and detaching the portion of bone separated by the crown of the trepan, and for removing stumps of teeth. See Lever.

ELEVATOR ANI, *Levator ani*.—*e. Labii inferioris*, Levator labii inferioris.—*e. Labii superioris proprius*, Levator labii superioris proprius.—*e. Labiorum communis*, Levator anguli oris.—*e. Oculi*, Rectus superior oculi.—*e. Patientiæ*, Levator scapulæ.—*e. Scapulæ*, Levator scapulæ.—*e. Testiculæ*, Cremaster.—*e. Urethræ*, see Transversus perinei.

ELEVATOR COM'MON. This is a mere lever, the end of which is somewhat bent and made rough, in order that it may less readily slip away from the portion of bone which is to be raised. The instrument is used, by forming a fulcrum for it, either on the hand which holds it, or upon the fingers of the other hand; or by making a fixed point for it on the edge of the opening made with the trephine.

ELEVATOR OF LOUIS differed from the last only in the circumstance of the screw-peg being united to the bridge by a kind of pivot instead of hinge, so that greater extent of motion was permitted.

ELEVATOR OF PETIT consists of a lever mounted on a handle, and straight throughout its whole length, except at its very end, which is slightly curved, in order that it may be more conveniently put under the portion of bone intended to be elevated. The lever is pierced at various distances from its butt-end with several holes, intended for the reception of a movable screw-peg, fixed on the top of a kind of bridge. This part of the instrument consists of an arch, the ends of which are long, and covered with small pads, and on its centre is the screw-peg already mentioned. By means of these holes the arm of the lever can be lengthened at pleasure.

ELEVATOR TRIP'LOID, *Vec'tis triploï'des*. This was so called from its consisting of three branches,

uniting above in one common trunk. The latter part was traversed by a long screw, having below a kind of hook, and above a handle for turning it. By turning the screw, the hook was drawn up, and the bone thus elevated.

The simple lever is the only one now used, owing to the want of facility and advantages in the use of the others.

ELEVATOR, UTERINE, see Pessary intrauterine.

ÉLÈVE EXTERNE, (e, and levare, 'to raise,') see House-surgeon.—*É. Interne*, House-surgeon.

ÉLEVURES (F.), *Efflorescentiæ*. A generic name, including all the exanthemata, in which there is tumefaction of the tissue of the skin. See Exanthem.

ELFLOCK. A condition of the hair in which it is matted and twisted into locks, as in plica. Supposed, in olden times, to have been the work of elves or fairies.

ELIASTER, *Ilech*.

ELICHRYSUM, (ἐλκυσσος, from ἐλγ, 'light of the sun,' and χρυσος, 'gold,') *Solidago virgaurea*.—*e. Montanum*, *Antennaria dioica*.

ELICOIDES, (ἐλικοειδης, from ἐλκξ, 'a tendril,' and αἶδος, 'resemblance,') *Capreolaris*.

ELIGMA, gen. *Elig'matis*, *Eclectos*.

ELIMINANTIA, (eliminare, (from e, and limen, 'a threshold,') 'to turn out of doors,') *Ecercitica*.

ELIMINATIVES, *Ecercitica*.

ELIQUATIO, (e, and liquare, *liquatum*, 'to melt,') *Colligation*.

ELIXA'TIO, *Elix'a'tion*, *Eps'e'is*, (from *elixus*, (e, and lix, 'water, liquor, lye,') 'boiled,' 'sodden,') This word has been used synonymously with Decoction. The act of boiling.

ELIX'IR, (Arab.) *I'sir*, *I'ir*, *Quel'les*, *Alex'ir*. (The etymology of this word is not clear. Lemery derives it from *ἐλκω*, 'I extract,' and also from *αλεξω*, 'I aid.' Others believe it to be from Arabic, *al-ecsir*, or *al-eksir*, 'chymistry,') An elixir is a medicine composed of various substances held in solution in alcohol. The name has been used, however, for preparations which contain no spirit of wine.

ELIXIR AC'IDUM HALLE'RI seu **DIPPEL'II**, *E. Antipodag'ricum* seu *Antinephret'icum* seu *Sulphurico-ac'idum*, *Gul'tæ ac'idæ tonicæ*, *A'qua Rabelii*, *Liq'or ac'idus Halle'ri*, *Mistu'ra sulphurico-ac'ida*, *Æ'ther sulphuricus ac'idus*, *Ac'idum sulphuricum alcoholisa'tum*, *A. vitrioli'cum vino'sum*, *Al'cohol Sulphurica'tum*, *A. sulphu'ricum*, is a mixture of concentrated sulphuric acid and alcohol—in the *Eau de Rabel*, of one part of the former to three of the latter. It is employed as an astringent in hemorrhages, &c.

ELIXIR ALOES, *Tinctura aloes composita*.—*e. Anthelminticum Succorum*, *Tinctura jalapii composita*.

ELIXIR, ANTIASTHMAT'IC, OF BOERHAAVE. This elixir is composed of alcohol, aniseed, camphor, orris, asarabacca root, *calamus aromaticus*, liquorice, and elecampane. It is given in asthma, in the dose of 20 or 30 drops.

ELIXIR ANTIHYPOCHONDRICUM, *Tinctura cinchonæ amara*.—*e. Antinephreticum*, *E. acidum Halleri*.

ELIXIR, ANTISCROF'ULOUS, OF PEYRIEHE, is composed of weak alcohol, subcarbonate of potassa, and gentian root. It is administered in scrofula.

ELIXIR, APERITIVUM, *Tinctura aloes composita*.

ELIXIR, BOERHAAVE'S, see *Tinctura aloes composita*.

ELIXIR OF CALISA'YA. This is made of *Calisaya bark* ʒj; fresh orange peel ʒss; cinnamon, coriander, and angelica seeds, ʒā ʒijj; caraway, aniseed, and cochineal, ʒā ʒj; brandy and water, of each a sufficient quantity; *symp* fʒx. Treat the Calisaya and aromatics with the brandy by percolation, until ʒx have been obtained: continue the

percolation with equal parts of brandy and water until $\mathfrak{z}$ xxij have been obtained; then add the syrup to make two pints. A tonic and cordial.

ELIXIR, CARMINATIVE, OF SYLVIVS, Tinctura carminativa Sylvii.

ELIXIR CLAUDE'RI, CLAU'DER'S *Elix'ir*. Made by macerating for a few days carbonate of potassa $\mathfrak{z}$ j; aloe, guaiacum, myrrh, saffron, rhubarb bruised, of each $\mathfrak{z}$ j; water $\mathfrak{f}$ xxvij; and decanting. Dose, a tablespoonful as a laxative and emmenagogue.

ELIXIR, DAFFF'S, Tinctura sennæ comp.—e. Danorum, E. pectorale regis Daniæ—e. Fœtidum, Tinctura castorei composita—e. of Garus, Tinctura de croco composita—e. Guaiaci volatilis, Tinctura Guaiaci ammoniata—e. Jalapæ compositum, Tinctura jalapii composita—e. of Long Life, Tinctura aloes composita—e. de *Longue vie*, Tinctura aloes composita—e. d'Or de M. le Général de la Motte, Tinctura seu Alcohol sulfurico-æthereus—e. Paregoric, Edinburgh, Tinctura opii ammoniata—e. Paregoricum, Tinctura camphoræ composita—e. Pectorale dulce, E. pectorale regis Daniæ.

ELIXIR PECTORA'LE RE'GIS DA'NLE, E. *Dano-rum seu Ringelmanni* seu *ex succo glycyrrhizæ* seu *pectora'le dulcè* seu *è succo liquoritizæ*: (Succ. glycyrrhiz. p. 1; Aq. Fœnicul. p. 2; Alcohol ammoniat. p. 6. A formula in many of the Pharmacopœias of continental Europe.) With the addition of opium it constitutes the *Elixir ammoniato-opiatum*, *Extractum thebaticum ammoniacale*, of some Pharmacopœias.

ELIXIR PROPRIETATIS, Tinctura aloes composita—e. Proprietas Paracelsi, Tinctura aloes composita—e. Purgans, Tinctura jalapii composita—e. Radcliffe's, see Tinctura aloes composita—e. Rhei dulce, Vinum rhei palmati—e. Ringelmanni, E. pectorale regis Daniæ—e. Roborans Whyttii, Tinctura cinchonæ amara—e. Sacrum, Tinctura rhei et aloes—e. Salutis, Tinctura sennæ comp.—e. Squire's, see Tinctura camphoræ composita—e. Stomachicum, Tinctura gentianæ composita—e. Stomachicum spirituosum, Tinctura cinchonæ amara—e. Stoughton's, see Tinctura gentianæ composita—e. ex Succo glycyrrhizæ, E. pectorale regis Daniæ—e. ex Succo liquoritizæ, E. pectorale regis Danorum—e. Sulphurico-acidum, E. acidum Halleri—e. Traumaticum, Tinctura Benzoini composita—e. Viscerale Hoffmanni, Tinctura Gentianæ composita.

ELIXIR VI'TÆ OF MATH'OLUS; composed of alcohol and 22 aromatic and stimulating substances. It was formerly employed in epilepsy.

ELIXIR OF VITRIOL, Sulphuric acid, dilute—e. of Vitriol, sweet, Spiritus ætheris aromaticus—e. of Vitriol, Vigan's, Spiritus ætheris aromaticus—e. Vitrioli, Sulphuricum acidum aromaticum—e. Vitrioli acid-aromaticum, Sulphuricum acidum aromaticum—e. Vitrioli dulce, Spiritus ætheris aromaticus, Sulphuricum acidum aromaticum—e. Vitrioli Edinburgensium, Sulphuricum acidum aromaticum—e. Vitrioli Mynsichti, Sulphuricum acidum aromaticum—e. Vitrioli cum Tincturâ aromaticâ, Sulphuricum acidum aromaticum—e. Whyttii, Tinctura cinchonæ amara.

ELIXIRIUM ANTISEPTICUM DOCTORIS CHAUSSIER, Tinctura cinchonæ ætherea composita.

ELIXIS, Eelectos.

ELIXIVIATION, (e. and *lixivium*, 'a lye,') Lixivation.

ELK, (Sax. ælc.) Cervus alces—e. Bark, Magnolia macrophylla—e. Tree, Andromeda arborea—e. Wood, Andromeda arborea, Magnolia macrophylla.

ELKOPLASTY, Helcoplasty.

ELLARNE, (Sax. ellapn.) Sambucus.

ELLEBORASTER, Helleborus fœtidus.

ELLEBORE, Dracontium fœtidum. See Hellebores.

ELLEBORISMUS, Helleborismus.

ELLEBORUM ALBUM, Veratrum album, see Helleborus.

ELLEBORUS ALBUS, Veratrum, see Helleborus—e. Niger, Helleborus niger.

ELLEN'TREE, (Prov.), Sambucus.

ELLER, (Prov.), Alnus glutinosa.

ELLERMAN'S DEODORIZING FLUID. An antiseptic or remover of odours—said to be a solution of a persalt (chloride) of iron.

ELLET, (Prov.), Sambucus.

ELLIPTIC, Oval.

ELLYCHNIA, Stellulæ Verheyntii.

ELLYCHNIO'TUS, (ἐλλυχνίωσις, from ἐλλυχνίον 'the wick of a lamp,') A sort of tent, used by the ancient surgeons, so called because it was shaped like a wick, or because it was made of a similar material.

ELM, COMMON, Ulmus—e. Red, Ulmus Americana—e. Rough-leaved, Ulmus Americana—e. Slippery, Ulmus Americana.

ELMINTHES, see Helmins.

ELMINTHOCORTON, (Helminthocorton,) Corallina Corsicana.

ELNORNE, Sambucus.

ELO'DES, *Helo'des*, (ελαδης, from ελος, 'a marsh,' and odes,) *Paludosa*, *Palustris*, *Paludal*, *Marshy*, (F.) *Marécageux*, *Paludéen*, *Palustre*, *Febbris elo'des* seu *helo'des* seu *paludosa*, *Holop'gra*, *Holop'gr-etos*, *Potamop'gra*, (F.) *Fièvre intermittente paludéenne*, *F. paludéenne* ou *Maremattique*, so called because it prevails in the Maremma district of Italy.) Marsh fever. Also, a kind of fever, characterized by great moisture or sweating.

ELOME, Orpiment.

ELONGA'TION, *Elonga'tio*, (*elongare*, *elongatum*, (e. and *longus*,) 'to lengthen,' 'extend,') An incomplete luxation, in which the ligaments of an articulation are stretched and the limb lengthened, without total luxation. The word has also been used for the extension required in the reduction of fractures and luxations, and for the increased length of limb, (F.) *Allongement*, in diseases and dislocations of the hip-joint.

ELREN, (Prov.) Sambucus.

ELTH, (old E.,) see Age.

ELUTRIATIO, Decantation, Elutrition.

ELUTRIA'TION, *Elutria'tio*, (originally *eluo*, (e. and *luo*, 'I wash,') 'I wash away, I rinse,') In pharmacy a process by which the finer particles of a powder are separated from the coarser. It consists in diffusing the powder in water, allowing the larger and heavier particles to subside, and then decanting the liquor, that the finer particles may subside.

ELU'VIES, ('a washing away,') An inordinate discharge of any fluid, and also the fluid itself. In the works of some authors it is particularly applied to the mucus which flows from the vagina in cases of leucorrhœa.

ELUXATION, Luxation.

ELYTRA, (pl.,) see Elytron.

ELYTRATRESIA, (*elytro*, and *ατρητος*, 'imperfected,') Colpatresia.

ELYTREURYNTER, (*elytro*, and *εuryντω*, 'I dilate,') Speculum vaginæ.

ELYTRITIS, (*elytro*, and *itis*,) Colpositis, Leucorrhœa.

ELYTRO, (*ελυτρον*, 'a vagina or sheath,') In composition, the vagina.

ELYTROBLENNORRHŒA, (*elytro*, and *blennorrhœa*,) Leucorrhœa (vaginal.)

ELYTROCE'LE, (*elytro*, and *κηλη*, 'a tumour,') Vogel has given this name to vaginal hernia, *Colpuce'le*.

ELYTRODES, (tunica), (*elytro*, and *odes*,) Vaginal coat of the testicle.

ELYTRŒDE'MA, *Coleced'e'ma*, *Colpced'e'ma*, (*elytro*, and *οἰδημα*, 'oedema,') Edema of the vagina.

ELYTROITIS, (*elytro*, and *itis*,) Colpositis.

ELYTRON, (*ελυτρον*, *ελυω*, 'I involve,') A sheath. The vagina. The membranes which envelope the spinal marrow are called *elytra*.

ELYTRON'CUS, *Elytroph'y'ma*, (*elytro*, and *ογκος*, 'a tumour,') A swelling of the vagina.

ELYTROPAP'PUS RHINOCERO'TIS. A South African bush, *Nat. Ord.* Compositæ, the whole of which is bitter and resinous. The tops of the branches, infused in wine or brandy, make excellent bitters. The tops are also given in powder to children affected with diarrhœa.

ELYTROPHY'MA, gen. *Elytroph'y'matis*, (*elytro*, and *φυμα*, 'a swelling,') Elytroncus.

ELYTROPLASTY, *Elytropolas'ticē*, (*elytro*, and *πασσω*, 'I form,') The operation for restoration of the vagina.

ELYTROPTOSIS, (*elytro*, and *πτωσις*, 'a falling,') Prolapsus vaginae.

ELYTRORRHAGIA, (*elytro*, and *ῥαγη*, 'a breaking forth,') Colporrhagia.

ELYTRORRHAPHY, *Elytorrhaph'ia*, *Colpor'r'haphy*, *Kolpor'r'haphy*, (*elytro*, and *ῥαγη*, 'a suture,') The operation of closing the vagina by suture in cases of procidentia uteri.

EM and **EN**, *ε*, 'in, into, within,') also 'excess,') frequently used in this last sense by Dr. Good. A common prefix, generally answering to the prefixes *im* and *in*, in English. In composition, before *β*, *π*, *φ*, *ψ*, and *μ*, the *ν* is changed into *μ*; before *γ*, *κ*, *ξ*, and *χ*, into *γ*; before *λ*, into *λ*; and before *ρ* generally into *ρ*.

EMAC'IMATE, *Maces'cere*, *Tabes'cere*, (F.) *Amaigrir*; (*emaciare*, *emaciatum*, (*e*, and *maceo*, 'to be lean,') 'to grow lean,') To lose flesh, to become lean, to waste away.

EMACIATION, *Emacia'tio*, *Extenua'tio*, *Ma'cies*, *Macritu'do*, *Ma'ror*, *Marco'res*, *Skeleti'a*, *Leptym'us*, *Leptyn'sis*, *Pinguet'inis diminutio*, (F.) *Amaigrissement*, *Dépérissement*. That condition of the body, or of a part of the body, in which it grows lean. The state of one who is lean — *Leanness* *Isch'notes*, *Asar'cia*, (F.) *Mai-greur*.

EMAIL DES DENTS, Enamel of the teeth.

ÉMAILLOTAGE (F.) (*maillot*, 'swathing or swaddling clothes,') The 'wrapping up' or 'packing up' in dry or wet sheets, which is practised in hydropathic establishments to induce sweating.

EMANA'TION, *Emana'tio*, *Efflu'vium*, (*emanare*, *emanatum*, (*e*, and *manare*, 'to issue or flow from,') The term is applied to a body which proceeds or draws its origin from other bodies—such as the light which *emanates* from the sun; the miasm which arises from the putrid decomposition of animal or vegetable substances, &c. See *Miasm*.

EMAN'SIO MEN'SIUM, (*e*, and *manere*, *man-sum*, 'to stay,') This term has been applied to amenorrhœa or obstruction of the menses before they have been established. Some have used it for the retention which occurs even after they have been established. The former is the general acceptance.

EMARGINATIO, (*e*, and *margo*, gen. *marginis*, 'a margin,') *Échancre*, *Notch*.

EMARGINATURA, *Échancre*, *Notch*.

EMASCULA'TION, *Emascula'tio*, (*emasculare*, (*e*, and *masculus*, 'a male,') 'to render impotent,') The act of removing or destroying the generative organs of a male animal.

EMASCULATUS, *Castratus*, *Eunuch*.

EMBALMING, (*em*, and *balsamum*, 'balsam,') *Balsama'tio*, *Smyrni'smus*, *Cede'a*, *Pollinctu'ra*, *Neeroced'a*, *Conditu'ra Cadav'erum*, (F.) *Embau-*

ment, *Imbalsamation*. An operation which consists in impregnating the dead body with substances capable of preventing it from becoming putrid, and thus putting it in a condition to be preserved.

EMBA'MMA, gen. *Emban'matia*, (*εμβαμμα*,) *Apoban'ma*, *Bamma*, (*εμβαπτω*, (*em*, and *βαπτω*, 'I dip,') 'I immerse,') A medicated pickle or sauce.

EMBARRAS, (*em*, and *barre*, 'to bar,') *Emphraxis*.

EMBARRAS GASTRIQUE (F.), *Gas'tric disorder* or *impediment*, *Collu'vies gas'trica*, *Sor'des prima'rum via'rum*, *Sta'tus gas'tricus*, *Gaster-atax'ia*, (F.) *Gastricitē*. Disorder of the stomach, in which there is loss of appetite, with bitterness and clamminess of mouth, white and yellow tongue, feeling of oppression in the epigastrium, and sometimes pain in that region, nausea and bilious or bitter vomiting; this state being accompanied with headach, lassitude, and pain in the back and limbs.

EMBARRAS GASTRO-INTESTINAL, *Gas'tro-intes'tinal disorder*. Slight gastro-enteritis, according to the Broussaists, in which the symptoms of the *Embaras gastrique* and *E. intestinal* are united.

EMBARRAS INTESTINAL, *Intes'tinal disorder*. The principal characters assigned to this are:—tension of the abdomen, colic, borborygmi, discharge of flatus *per anum*, and constipation or diarrhœa.

EMBARRURE, (*em*, and *barré*, 'barred,') *Engisma*.

EMBAUMEMENT, (*em*, and *baume*, 'balsam,') *Embalming*.

EMBOÎTEMENT (F.), 'Encasing,') (*em*, and *boîte*, 'a box,') *Enadel'phia*, *Endadel'phia*, *Enteradel'phia*. This term has been applied to the theory of generation which considers that the germs are encased in each other in the ovary of the female, in such sort that they are developed in succession after impregnation. It is the theory of *encasing of germs*. One, who believes that all organized bodies are the result of the development of germs encased in each other *ad infinitum*, is said to be an *Infin'itivist*.

Monstros'ities by inclu'sion, (F.) *Monstruosités par inclusion*, are supposed to arise from an accidental *emboîtement* of germs.

EMBOÎTURE, *Diarthrosis*.

EM'BOLE, (*εμβολω*, (*εμ*, and *βαλλω*,) 'I put in place,') Reduction of a luxated bone. *Repo-sit'io*. Also, a wedge or plug—*εμβολη*, *εμβολον*, *εμβολος*—hence.

EMBOL'IA, (F.) *Embolies*. Diseases produced by plugging or obturation of the bloodvessels. *Em'bolism*, *Embolism'us*.

EMBOL'IC, *Embol'icus*, (from *εμβολη*, 'a wedge or plug,') Relating to plugging or obturation of the vessels, as '*embolic apoplexy*.'

EMBOLIMOS, (*εμβολιμος*, from *em*, and *βαλλω*, 'I cast,') *Intercalary*.

EMBOLISM, see *Embolia*.

EMBOLUM CEREBRI, *Infundibulum* of the brain.

EMBONPOINT (F.). 'In good point or plight,') *Bo'na cor'poris habitu'do*. The state of the animal body when in full health. Excessive *embonpoint* constitutes corpulence and obesity, and may become a morbid condition.

EMBONPOINT EXCESSIF, *Polysarcia adiposa*.

EMBOWELLING, *Exenterismus*.

EMBREGMA, *Embrocation*.

EMBROCATIO, *Embrocation*, *Irrigation*.

EMBROCA'TION, *Embroca'tio*, *Embreg'ma*, *Em'broché*, *Implu'vium*, (*εμβρεχω*, (*εμ*, and *βρεχω*, 'I wet,') 'I dip in,') A fluid application to be rubbed on any part of the body. It is often used

synonymously with liniment. Originally it was employed in the sense of *Fomentation*.

EMBROCATIO, GUESTONIA, see *Oleum Terebinthinæ rectificatum*.

EMBROCHE, (εμβροχή,) Embrocation.

EMBRYO, gen. *Embryonis*, *Em'bryon*, (εμβρυον, from *em*, and *βρω*, 'I grow.') *Cye'ma*. The fecundated germ, in the early stages of its development in utero. At a certain period of its increase, the name *fœtus* is given to it, but at what period is not determined. Generally, the *embryo state* is considered to extend to the period of quickening.

EMBRYOTONY, *Fœtûs trucidatio*, (*embryo*, and *τονος*, 'destruction.') The act of destroying a fœtus in utero, when insurmountable obstacles—as certain deformities of the pelvis—oppose delivery.

EMBRYOG'ENY, *Embryogen'ia*, (F.) *Embryogénie*; (*embryo*, and *γεννω*, 'to generate.') The generation or formation of the embryo.

EMBRYOGRAPHY, *Em'bryogra'phia*, (*embryo*, and *γραφω*, 'a description.') The part of anatomy which describes the embryo.

EMBRYOL'OGY, *Embryolog'ia*, (*embryo*, and *λογος*, 'a description.') The doctrine of, or a treatise on, embryonic development.

EMBRYON, (εμβρυον,) Embryo.

EMBRYONAL, Embryonic.

EMBRYONIC, *Embryon'ic*, (F.) *Embryonique*, *Em'bryonnaire*, *Em'bryonal*. Relating or appertaining to an embryo:—as 'embryonic life.'

EMBRYONIQUE, Embryonic.

EMBRYONNAIRE, Embryonic.

EMBRYOTHLASTA, *Embryothlas'tes*, *Em'bryothlas'tum*, (F.) *Embryotome*, (*embryo*, and *θλαω*, 'I break.') An instrument for dividing the fœtus piecemeal, in order to effect delivery. A crotchet or other instrument, used, in certain cases of laborious parturition, to break the bones of the fœtus, for the purpose of extracting them with greater facility.

EMBRYOTOCIA, (*embryo*, and *τοκος*, 'delivery,') Abortion.

EMBRYOTOME, Embryothlasta.

EMBRYOT'OMY, *Embryotom'ia*, (εμβρυοτομία, from *embryo*, and *τεμνω*, 'to cut.') A division of the fœtus into fragments, to extract it by piecemeal, when the narrowness of the pelvis or other faulty conformation opposes delivery.

EMBRYOTROPHY, *Embryotroph'ia*, (*embryo*, and *τροφω*, 'nourishment.') Fœtal nutrition: *Cyot'rophy*.

EMBRYUL'CIA, *Embryuterul'cia*, (*embryo*, and *ἐκλω*, 'I extract,' 'I draw.') A surgical operation, by which the fœtus is extracted by means of appropriate instruments, when faulty conformation or other circumstances prevent delivery by the natural efforts.

EMBRYUL'CUS, (εμβρυουλκος,) *Eleus'ter*. An iron hook or crotchet, described by Fabricius ab Aquapendente, which was used to extract the fœtus in some cases of laborious labor.

EMBRYUSTERULCIA, (*embryo*, *βρετρα*, 'uterus,' and *ἐκλω*, 'I extract,') Embryulcia.

EMENDANS, (*emendo*, (e, and *mendum*, 'a fault,') 'to amend,') Corrigent.

ÉMERAUDE, Smaragdus.

EMERAUDES, Hæmorrhoids.

EMERODS, Hæmorrhoids.

EMEROIDS, Hæmorrhoids.

EMESIA, (εμεσια,) Vomiturition.

EMESIS, (εμεσις,) Vomiting.

EMETATROPH'IA, (εμεω, 'I vomit,' and *τροφω*, 'want of nourishment,') Atrophy induced by vomiting.

EMETIA, Emetine.

EMET'IC, *Emet'ic*, *Emetopœ'um*, *Vomito'rium*, *Vom'itory*, *Vom'it*, *Puke*, (Prov.) *Pick*, (εμετικός, from *εμεω*, 'I vomit,') A substance capa-

ble of producing vomiting. (F.) *Vomitif*, *Émetique*. [This last term is also given by the French to *tartarized antimony*—the emetic, as it were, *par excellence*.]

Tartarized antimony, *ipecacuanha*, and *sulphate of zinc*, are the chief emetics. They are valuable agents in disease, and may either act *primarily* on the stomach, or *secondarily* on other parts of the system,—the sympathy between the stomach and other parts of the body being very extensive, and an important object of study.

EMETIC ROOT, *Euphorbia corollata*—e. *Tartar*, *Antimonium tartarizatum*—e. *Weed*, *Lobelia inflata*.

EMETICOLOGIA, Emetology.

EMETINE, *Emet'ina*, *Emet'ia*, *Vom'itine*, *Cephæli'num*. A vegetable alkali, discovered by M. Pelletier in *ipecacuanha*, to which it owes its emetic power. It is obtained from different *ipecacuanhas*, but chiefly from *psychot'ria emet'ica*, *callicoc'ca ipeacacuan'ha*, and *viol'la emet'ica*. It is in transparent scales, of a reddish-brown colour, almost inodorous, of a bitter, slightly acid taste; is unchangeable in the air, soluble in water and alcohol, and insoluble in ether. Three grains of impure emetia or one grain of pure are equal to eighteen of *ipecacuanha*. See *Ipeacacuanha*.

ÉMÉTIQUE, *Antimonium tartarizatum*, Emetic.

ÉMÉTISER (F.). To cause vomiting by emetics. In French pharmacy, it is, likewise, used to signify the addition of *émétique*—tartar of antimony and potassa—to a medicinal preparation.

EMETOCATHARSIS, see *Emetocathartic*.

EMETOCATHARTIC, *Em'etocathar'ticus*, (*emetus*, and *καθαρτικος*, 'a purgative,') A remedy, which at the same time excites vomiting and purging—*Em'etocathar'sis*.

EMETOL'OGY, *Emetolog'ia*, *Emeticolog'ia*, (*emetus*, and *λογος*, 'a discourse,') A treatise on vomiting, and on emetics.

EMETOPŒUM, (*emetus*, and *ποιω*, 'to make,') Emetic.

EMETUS, (εμετος,) Vomiting.

EM'INENCE, (e, and *mineo*, 'to project,') *Eminent'ia*, *Protuberant'ia*, *Ex'ochæ*. A projection at the surface of a healthy or diseased organ.

ÉMINENCE PORTE ANTÉRIEURE, *Lobulus anonymus*—*é. Porte postérieure*, *Lobulus Spigelii*—*é. Sus-pubienne*, *Mons veneris*.

ÉMINENCES BIGÉMINÉES, *Quadrigenima corpora*—*e. Latérales*, *Corpora olivaria*—*é. Pyramidales*, *Corpora pyramidalia*—*é. Vermiformes du cercelet*, *Vermiformes processus*.

EMINENTIA, Eminence, *Protuberantia*—*e. Annularis*, *Pons varolii*—*e. Pyramidalis*, *Crista vestibuli*—*e. Thyroidea*, *Pomum Adami*—*e. Tympani*, *Pyramid*.

EMINENTIÆ BIGEMINÆ, *Quadrigenima tubercula*—*e. Candicantes*, *Mammillary Tubercles*—*e. Lenticulares*, *Corpora striata*—*e. Magnæ cerebri*, *Thalami nervorum optico'rum*—*e. Medianæ internæ*, *Corpora Pyramidalia*—*e. Olivares*, *Corpora olivaria*—*e. Ouales laterales*, *Corpora olivaria*—*e. Pyramidales*, *Corpora pyramidalia*—*e. Quadrigenimæ*, *Quadrigenima tubercula*—*e. Teretes*, *Processus teretes*.

ÉMISSAIRE, (same etymon as *Emission*,) Emenatory.

EMISSARIA SANTORINI, (F.) *Émissaires de Santorini*. A name given to some small veins, which communicate with the sinuses of the dura mater, by apertures in the cranium. Such are the parietal, posterior condyloid, mastoid veins, &c.

EMISSION, Emission—e. Seminis, Spermatismus.

EMISSION, *Emis'sio*, (*emittere*, (e, and *mittere*, *missum*), 'to send out,' 'drive out.') The act by which any matter whatever is thrown from the body. Thus, we say *Emission of urine*, *Emission of semen*, &c.

EMISSION SANGUINE, Bloodletting.

EMMANUEL. The name of an ointment, much used in the latter part of the 16th century, which was formed by boiling various herbs, as dittany, centaury, plantain, spurge, &c., in wine and afterwards adding pitch, resin, olibanum, turpentine, suet, mastich, myrrh, &c. to form an ointment.

EMMENAGOGUES, *Emmenago'ga*, *Emmeni-ago'ga*, *Ame'nia*, (*emmenia*, and *αἷω*, 'I drive,' or 'expel.') *Men'agogues*. A name given to medicines believed to have the power of favouring the discharge of the menses. There is probably no substance which possesses this power directly. According to different conditions of the system, the most opposite remedies may act as emmenagogues. *Black hellebore*, *savin*, *madder*, *polygala senega*, and *ergot*, are by some, reputed to be specific emmenagogues.

EMMENAGOGU'IA, (*emmenagogue*, and *λογος*, 'discourse.') A treatise of emmenagogues.

EMMENIA, (*εμμηνα*, from *em*, and *μην*, 'a month,') Menses.

EMMENIAGOGA, *Emmenagogues*.

EMMENIOP'ATHY, *Emmeniopathi'a*, (*emmenia*, and *πάθος*, 'affection,') An affection or disorder of menstruation.

EMMENOLOG'IA, (*emmenia*, and *λογος*, 'a discourse,') A treatise on menstruation.

EMMENOLOG'ICAL, *Emmenolog'icus*. Relating or appertaining to menstruation.

EMMYXIIUM ARTICULARE, (*em*, and *μυξα*, 'mucus,') Hydrarthrus.

EMOLLIENTIA, *Emollients*.

EMOL'LIENTS, *Emollient'ia*, *Malthac'tica*, *Relaxan'tia*, *Epiceras'tica*, *Malac'tica*, *Lubrican'tia*, *Malacopa'a*, *Mollien'tia*, (from *emollire*, (e, and *mollire*, 'to soften,') Substances which relax and soften parts that are inflamed, or too tense. They are used both internally and externally: as the former, however, consist of mucilaginous substances, they are generally reckoned as demulcents: the latter, or proper emollients, consist of oils, cataplasms, fomentations, &c. Oleaginous bodies, rubbed on a part, act by diminishing its cohesion. Fomentations, in cases of internal inflammation, act probably through contiguous sympathy. The following are the chief emollients:—*Adeps*, *Amygdalæ Oleum*; *Avenæ Farina*; *Cera*; *Cetaceum*; *Linum*; *Olivæ Oleum*; *Sesamum*; *Tritici Farina*, and *Ulmus*.

EMOLLITIES, *Mollities*—e. *Morbosa*, *Mollities*—e. *Ossium*, *Mollities ossium*—e. *Uteri morbosa*, *Hysteromalacia*.

EMOLLITIO, *Mollities*—e. *Ventriculi*, *Gastro-malacia*.

EMONCTOIRE, *Emunctory*.

EMOPTOE, *Hæmoptysis*.

EMOTIO, (e, and *moveo*, *motum*, 'to move,') *Delirium*, *Luxation*, *Passion*.

EMO'TIONAL. Relating to emotion or passion independently of the will:—hence an 'emotional' or instinctive impulse.

EMPASMA, gen. *Empasmat'is*, (*em*, and *πασσιν*, 'to sprinkle,') *Cataplasma*.

EMPATEMENT, (*empâter*, (*em*, and *pâte*, 'paste,') 'to render pasty or doughy,') A non-inflammatory engorgement, which retains, more or less, the impression of the finger.

EMPATHE'MA, gen. *Empathe'matis*, *Ma'nia a pathe'matis*, (F.) *Munie sans délire*; ungovernable passion; (*em*, and *πάθος*, 'suffering,') Fixed delusion.

EMPEIRIA, (*εμπειρία*, from *em*, and *πειρα*, 'a trial,') Experience.

EMPETRUM, (*εμπέτρον*, from *em*, and *πετρα*, 'a rock,') *Herniaria glabra*.

EMPHLYSIS, (*em*, and *φλυσις*, 'a vesicular tumour or eruption,') *Ich'orous Exan'them*. An eruption of vesicular pimples, filled progressively with an acrid and colourless, or nearly colourless, fluid, terminating in scurf or laminated scabs. A genus in the order *Ecanthematica*, class *Hæmatica* of Good.

EMPHLYSIS APHTHA, *Aphthæ*, *Stomatitis*, *aphthous*—e. *Erysipelas*, *Erysipelas*—e. *Miliaria*, *Miliary fever*—e. *Pemphigus*, *Pemphigus*—e. *Vaccina inserta*, *Vaccina*—e. *Variella*, *Variella*.

EMPHRACTIC, *Emphracticus*.

EMPHRACTICA, *Physconia*.

EMPHRACTICUS, *Emphractic*, (*εμφρακτικός*, from *εμφράω*, (*em*, and *φράω*, 'I shut up,') 'I close,' 'I obstruct,') *Emphas'ticus*, *Emplatton'enus*. Any substance which, when applied to the skin, was presumed to close the pores.

EMPHRAG'MA, gen. *Emphrag'matis*, (*εμφραγμα*.) *Obturamentum*, *Impedimen'tum*. Anything that obstructs. Hippocrates uses this term to designate the obstacle to delivery on the part of the fœtus, when the presentation is preternatural.

EMPHRAGMA LACRYMALE, *Fistula lacrymalis*—e. *Salivare*, *Ranula*.

EMPHRAX'IS, (*εμφραξις*.) *Obstruc'tio*, *Obtura'tio*, *Oppila'tio*, *Infarc'tus*, *Infarc'tio*, *Farc'tus*, *Fartus*, *Infarc'tion*. 'Obstruction.' An Embarras or detention in canals or cavities of any substance, which is either morbid from quantity or quality.

EMPHRAXIS HEPATIS, *Hepatemphrax*is.

EMPHYMA, gen. *Emphy'matis*, (*em*, and *φύμα*, 'a swelling,') *Tumour*—e. *Encystis*, *Encystis*—e. *Encystis atheroma*, *Atheroma*—e. *Encystis ganglion*, *Ganglion*, *Testudo*—e. *Encystis meliceris*, *Meliceris*—e. *Encystis steatoma*, *Steatoma*—e. *Exostosis ossea*, *Exostosis*—e. *Exostosis periosteæ*, *Node*—e. *Sarcoma*, *Sarcoma*—e. *Sarcoma adiposum*, *Adipose sarcoma*—e. *Sarcoma cellulosum*, *Cystic sarcoma*—e. *Sarcoma mammæ*, *Mammary sarcoma*—e. *Sarcoma pancreæ*, *Pancreatic sarcoma*—e. *Sarcoma scirrhus*, *Scirrhus sarcoma*—e. *Sarcoma tuberculosum*, *Tuberculate sarcoma*—e. *Sarcoma vasculosum*, *Sarcoma vascular*.

EMPHYSE'MA, gen. *Emphyse'matis*, (*εμφύσημα*, from *εμφύω*, (*em*, and *φύω*, 'I blow,') 'I inflate,') *Infla'tio*, *Empneumatosis*, *Sarci'tes flatuosisus*, *Emphyse'ma cellula'rè seu pneumatosis*, *Physon'cus*, *Tum'or flatulent'is*, *Pneumatosis*, *Infla'tion*, *Winddropsy*, (F.) *Emphyse'me*, *Aërethmie* of *Piorry*. This term is commonly applied to any white, crepitant, shining, elastic, indolent tumor of the integuments, caused by the introduction of air into the areolar texture, *Aërodermectasia*. Injuries of the larynx, trachea, or lungs; fractures of the ribs, or wounds penetrating the chest, are the most frequent causes of this affection, which is owing to the air escaping from the air-passages, and insinuating itself into the areolar texture surrounding the wound. There are some cases of emphysema, which are owing to internal causes; and hence a division has been made into the *accidental* and *symptomatic*, and the *spontaneous* and *idiopathic*.

EMPHYSEMA ABDOMINIS, *Tympanites*.

EMPHYSEMA OF THE LUNGS, *Emph'ema*, *Asth'ma æthereum* ab *Emphyse'matè Pulmo'num*, *Pneumatosis Pulmo'num*, *Pneumonec'tasis*, *Pneumec'tasis*, (F.) *Pneumœctasie*, *Emphyse'me du Poupon*. A considerable pressure or contusion of the chest, or any violent concussion of the lung, may produce a laceration in that viscus, without injury being done to the parietes of the thorax, and may give occasion to the infiltration of air into the areolar texture, *interlob'ular emphysema*. La-

önnec has described another species of emphysema of the lungs, *Vesicular emphysema*, *Aëropneumonie*, *Aëropneumonectasie* of Piorry, consisting in excessive dilatation of the air-cells, some of which become as large as hemp-seed, ultimately break, and give rise to irregular vesicles at the surface of the lung, some of which are as large as a hazelnut. *Physical signs*.—The thorax externally, generally or locally, appears unnaturally convex and prominent. The intercostal spaces are widened, but depressed. The inspiratory efforts are increased. The sound on percussion is morbidly clear, but not tympanitic. On auscultation, the inspiratory murmur is feeble or suppressed. The expiration, which is more frequently audible, is prolonged, laborious and wheezing. There are no certain physical signs which can distinguish *interlobular emphysema* from the *vesicular*.

Emphysema under the pleuro-pulmonalis is termed, by Piorry, *Aërethorax*.

EMPHYSEMA PECTORIS, Pneumothorax—e. Seroti, Physocèle—e. Tympanites, Tympanites—e. Uteri, Hysterophys, Physometra.

EMPHYSEM'ATOUS, *Emphysematosus*. Relating to or appertaining to emphysema.

EMPHYSEME, Emphysema—e. du Poumon, Emphysema of the lungs.

EMPHYTUM THERMUM, (ἐμφύτον θερμὸν, (*em*, and *pus*, 'to have naturally,') 'innate heat.' Biolychnium.

EMPIMELIUM POLYSARCIA, (*em*, and *πιελη*, 'fat.') Polysarcia adiposa.

EMPIRIC or *Empiric*, *Empiricus*, (ἐμπειρικός, from *em*, and *πειρά*, 'a trial,') 'experimental.' One who follows only experience. A sect of physicians, who rejected all theory, and took for their guide experience alone. It was opposed to the dogmatic sect. The Empiric sect prevailed till near the time of Galen. Among its most eminent members, after Philinus and Serapion, were Apollonius, Glaucias, Baccchius of Tanagra, and Zeuxis, both disciples of Herophilus.—Heraclides of Tarentum, Cleophantus (master of Asclepiades,) Menodotus of Nicomedia, and Theudas of Laodicea. They occupied themselves, chiefly, with discovering the properties of drugs, and did important service, in this manner, to medicine.

At the present day, the word *Empiric* is only taken in a bad sense, being employed in nearly the same signification as *charlatan* or *quack*.

EMPIRICE, (ἐμπειρικὴ,) Empiricism.

EMPIRICISM, *Empirica Ars*, *Empiricē*, (F.) *Empirisme*. Medicine founded on experience. It generally, at the present day, signifies *quackery*.

EMPIRISME, Empiricism.

EMPLASTICUS, (ἐμπλαστικός, from *επιπλασσω*, (*em*, and *πλασσειν*, 'to form,') 'I spread upon.') A remedy which adheres, as a plaster, to the surface, and in this manner may obstruct the pores (?): an emphractic.

EMPLASTROËNDERMIC, Endermic.

EMPLASTRUM, (ἐμπλαστρὸν, same etymon as Emplasticus.) (F.) *Emplâtre*, *Stéarate*. A solid and tenacious compound, adhesive at the ordinary heat of the human body. Some of the plasters owe their consistence to wax and resin; and others to the chemical union which takes place between the semivitreous oxide of lead and oil. Most of them become too consistent by age. When such is the case, they may be remelted by a gentle heat, and oil be added to them.

EMPLASTRUM ADHÆSIVUM, Emplastrum resinæ—e. Adhæsivum Woodstockii, see Sparadrapum Adhæsivum—e. Alexandri, Alexandrine—e. de Althææ, Unguentum de Althææ.

EMPLASTRUM AMMONIACI, *Ammoniacum Plaster*, (*Ammoniac. pur.* 3v, *Acidi aceticum dil.* Oss. Dissolve the ammoniac in the vinegar and strain: then evaporate the solution by means of a water-bath, stirring constantly until it acquires a proper

consistence.—*Ph. U. S.*) It is used as a discutient plaster.

EMPLASTRUM AMMONIACI CUM HYDRARGYRO, *Ammoniacum plaster with mercury*, (*Ammoniaci lbj, Hydrarg. 3ijj, Olei oliv. gr. x, Sulphur. gr. viij.*) Rub the mercury with the oil until the globules disappear, then add the ammoniac, previously melted, and mix. *Ph. U. S.*)

EMPLASTRUM ANDRÉE DE CRUCE, E. de Pice et resinis glutinans—e. Anglicum, see Sparadrapum adhæsivum.

EMPLASTRUM ANTIMONII, *Plaster of Antimony*, (*Antimon. et Potass. tart. in pulv. subtil. 3j, Picis Burgundic. 3iv.* Melt the pitch in a water-bath and strain: add the powder, and stir well on cooling. *Ph. U. S.* 1863.) An excitant or warm plaster.

EMPLASTRUM ARNICÆ, *Plaster of Ar'nica*, *Ar'nica plaster*, (*Ext. Arnicæ alcoholic. 3iss, Emp. Resinæ, 3ijj.* Add the extract to the plaster, previously melted, and mix. *Ph. U. S.* 1863.) A warm strengthening plaster.

EMPLASTRUM AROMATICUM, *Aromatic Plaster*, (*Thuris, 3ijj, Cereæ flavæ, 3ss, Pulv. cort. cinnam. 3vj, Ol. ess. Piment., Ol. ess. Limon. aa 3ij.* Melt the frankincense and wax together, and strain; then add, as it cools, the cinnamon, previously rubbed with the oils, and form a plaster. *Ph. D.*) Used as a stimulating plaster.

A *spice plaster*, made by incorporating powdered cinnamon and any other spices with melted *suët*, has been used in cholera infantum. It is spread on leather or linen, and is renewed twice in the twenty-four hours. Another, for extemporaneous use, is to mix powdered spices with meal or flour and spirits. One, advised by Dr. Parrish, senior, of Philadelphia, consisted of powdered capsicum, cinnamon, and cloves, each 3ij; rye meal, spirits, and honey, q. s. To be rubbed together on a plate, spread upon a close fabric, and applied as a cataplasm.

EMPLASTRUM ASSAFETIDÆ, *E. antihystericum*, *Assafetida plaster*, (*Emp. plumb., Assafet. sing. lbj, Galban., Cereæ flavæ, aa lbss, Alcohol. dilut. Oij.* Dissolve the assafetida and galbanum in the alcohol, in a water-bath, strain while hot, and evaporate to the consistence of honey; add the lead plaster and wax previously melted together, stir the mixture well, and evaporate to the proper consistence. *Ph. U. S.*) Used as an antispasmodic (?) and anodyne (?) plaster.

EMPLASTRUM ATTRAHENS, E. cereæ—e. Auriculare, Hypotium.

EMPLASTRUM BELLADONNÆ, *Plaster of Belladonna*, (*Empl. resin. 3ij, Ext. Belladon. alcoholic. 3i.* Add the extract to the plaster, previously melted in a waterbath, and mix. *Ph. U. S.*) An anodyne application in neuralgia and rheumatism.

EMPLASTRUM CALEFA'CIENS, *E. Picis cum cantharidæ*, (*Ph. U. S.*), *Calefacient Plaster*, *Warm Plaster*, *Warming plaster*, (F.) *Emplâtre échauffant*. (*Cerat. cantharid. (Ph. U. S.), 3iv, Picis abiet. 3lvijj.* Melt together, and form into a plaster. *Ph. U. S.*) It is rubefacient and stimulant.

EMPLASTRUM CANTHARIDIS, E. lyttæ—e. Cantharidis vesicatorii compositum, E. lyttæ comp.—e. Cephalicum, E. picis compositum.

EMPLASTRUM CERÆ, E. simplex, E. attrahens, *Wax plaster*, (*Cereæ flavæ, Sevi Præp. aa lbijj, resinæ flavæ, lbj.* Melt them together and strain. *Ph. L.*) It has been considered drawing. It is stimulating.

EMPLASTRUM CICUTÆ, E. de Cicuta seu conii maculati, *Unguentum sol'idum de cicuta*, *Hemlock plaster*, (F.) *Emplâtre de Ciguë*. (*Resinæ abiet. 960 p. Cereæ flav. 640 p. Picis albæ, 448 p. Ol. cicute per decoct. præparat. 128 p. Fol. cicut. recent. 2000 p.* Melt the resins, wax and oil; add the cicuta leaves, and boil; strain and add,

after having dissolved it in vinegar of squills and scienta juice, *gum ammoniac*. 500 p. *Ph. P.*) It is used as a discutient, especially to scirrhus tumours.

EMPLASTRUM CINEREUM, E Hydrargyri—e. Citrinum, Unguentum de althæa—e. Commune, E. Plumbi—e. Commune cum resinâ, E. resinæ—e. Conii, E. ciutæ.

EMPLASTRUM CUMINI, Cum'min plaster, (Cumin. semin., Carui sem., Lauri bac., sing. ʒiij, Picis aridæ, ʒiij, Cere flavæ, ʒiij. Melt the pitch and wax together, then add the other ingredients. *Ph. L.*) It is used as a warm discutient plaster.

EMPLASTRUM DIACHYLON, Diachylon, E. plumbi—e. Divinum Nicolai, see Magnet—e. Emolliens, Unguentum de althæa—e. Epispasticum, E. lyttæ.

EMPLASTRUM FERRI, E. roborans, Iron Plaster, Strengthening Plaster, (Ferri subcarb. ʒiij, Emp. plumbi, ʒiij, Picis Burgund. ʒss. Add the subcarbonate of iron to the lead plaster and Burgundy pitch previously melted together, and stir until they thicken. *Ph. U. S.*)

EMPLASTRUM FERRI RUBRI, E. oxidi ferri rubri—e. Flavum, Unguentum de althæa—e. ad Fonticulos, Ceratum Galeni.

EMPLASTRUM GALBANI, Gal'banum plaster, (Empl. litharg. ʒiij, Gum. galban. ʒss, Cere flavæ, ʒiv. Melt the galbanum before adding the plaster and wax, then melt all together. *Ph. D.*) A stimulant and discutient.

EMPLASTRUM GALBANI COMPOSITUM, E. lithargyri compositum, Diach'ylon magnum cum gummi, Compound gal'banum plaster, (Galb. ʒviiij, Emp. plumbi, ʒiij, Tereb. ʒj, Picis Burgund. contus. ʒiij. Melt the galbanum and turpentine together, then mix in the resin, and afterwards the plaster, previously melted. *Ph. U. S.*) It is stimulant and discutient.

EMPLASTRUM GLUTINOSUM, see Sparadrapum adhaesivum—e. cum Gummatibus, E. gummosum.

EMPLASTRUM E GUMMATIBUS RESINOSIS, E. gummosum.

EMPLASTRUM CUM GUMMIRESTINIS, (F.) Emplâtre de gomme résine, Diachylon gommé, Plaster of gum resins, (Emplastr. simpl. p. 1600, Cere flavæ, p. 96, Picis albæ, p. 96, Terebinth. p. 96. Melt by a gentle heat, and add Gum ammoniac p. 32, Bdellium, p. 32, Galban, p. 32, and Sagapenum, p. 32. Dissolve in a sufficient quantity of alcohol, evaporate to the consistence of honey, and mix carefully all together. *Ph. P.*) A discutient.

EMPLASTRUM GUMMOSUM, Emplastrum lithargyri cum gummi seu cum gummatibus seu gummatibus resinosis seu Oxidi Plumbi semivirei gummosum, Gum plaster, Diach'ylon cum gummi, Yellow or Gum diach'ylon, (Emp. plumbi, p. viii, Ammoniac., Galbani, Cere flavæ, ʒā p. i. *Ph. E.*) Properties and uses like the last.

EMPLASTRUM HYDRARGYRI, Emplastr. lithargyri cum hydrargyro, E. cinereum, Mercurial plaster, (F.) Emplâtre mercuriel, (Hydrarg. ʒvj, Ol. oliv., Resin. ʒā ʒij, Emplast. plumbi, ʒj. Melt the oil and resin together, and when cool rub the mercury with them till the globules disappear; then gradually add the lead plaster, previously melted, and mix all. *Ph. U. S.*) It is stimulant, resolvent, and discutient. Applied as a discutient to venereal and other tumours.

EMPLASTRUM HYDRARGYRI COMPOSITUM, E. de hydrargyro compositum seu de Vigo cum mercurio emendatum, Compound plaster of mercur. (*Empl. simpl. p. 1250, Cere flavæ, p. 64, Resinæ, p. 64.* Melt, and before it congeals add Pulv. g. ammoniac., p. 20, Bdellium, p. 20, Oliban, p. 20, Myrrh, p. 20, Saffron, p. 12. Mix carefully. Take of Mercury, p. 380, Pure turpentine, p. 64, Liquid and Pure Storax, p. 192. Triturate in an iron mortar till the globules disappear: melt all to-

gether, and before congealing add essential oil of Lavender, p. 8. *Ph. P.*) The same properties as the last.

An *Emplastrum de Vigo sine mercurio seu Viginis sine mercurio*, (F.) *Emplâtre de Vigo* was, at one time, much celebrated as an adhesive plaster or spardrap. In the Pharmacopœia of Wirtemberg, it was directed to be formed of living frogs and living earthworms boiled with various ordinary plaster ingredients, and other substances, and was termed *Emplastrum rana'rum simplex seu de Ranis sine mercurio*.

EMPLASTRUM ICHTHYOCOLLÆ TELÆ INDUCTUM, see Sparadrapum adhaesivum—e. Irritans, E. lyttæ—e. Lithargyri, E. Plumbi—e. Lithargyri compositum, E. galbani compositum—e. Lithargyri cum gummi, E. gummosum—e. Lithargyri cum hydrargyro, E. hydrargyri—e. Lithargyri cum resinâ, E. resinæ—e. Lithargyricum cum resinâ pini, E. resinæ.

EMPLASTRUM LYTTÆ, E. cantharidis seu cantharidis vesicatorii seu mel'ois vesicatorii, E. vesicatorium, E. epispasticum seu e cantharidibus epispasticum solidius et tenacius hærens seu irritans seu rubefaciens, Plaster of the Span'ish or blistering fly, Blistering Plaster, Fly Plaster, (Cantharides, in powder, ʒj, Emp. cere, ʒss, Adipis. præp. ʒj. Melt the plaster and lard together, and as the mixture becomes thick, on cooling, sprinkle in the flies, and mix. *Ph. L.*) This is the common blistering plaster. Too much heat must not be used in its preparation. It requires to remain on six or eight hours before its full effect is induced, and it acts sufficiently well, provided even a piece of thin gauze or tissue paper be placed between it and the skin, whilst absorption of the flies is thus prevented. See Blister. The *Blistering Cerate* of the *Ph. U. S.*, *Ceratum Cantharidis*, is made as follows:—*Cantharid* in pulv. subtiliss., ʒxij, *Cere flavæ, Resinæ, ʒā ʒvij, Adipis, ʒx.* To the wax, resin, and lard, previously melted, add the Spanish flies, and stir the mixture constantly until cool.

The *Ph. U. S.* (1863) has a *Ceratum Extracti Cantharidis*, *Cerate of extract of cantharides*, formed by adding to an alcoholic extract of cantharides, resin, yellow wax and lard.

EMPLASTRUM LYTTÆ COMPOSITUM, E. cantharidis vesicatorii compositum seu mel'ois vesicatorii compositum, Compound plaster of cantharides or Span'ish flies, (Resin. liq. pini laricis. p. 18, Resinæ concret. pini abietis, Meloës vesicat. ʒā p. 12, Cere flavæ, p. 4, Subacet. cupri, p. ij, Semin. sinapis alb., Fruct. pip. nigr. ʒā p. j. Melt the pitch and wax, then add the turpentine, and, as these cool, sprinkle in the other substances, in the form of powder, so as to make a plaster. *Ph. E.*) The same properties as the last, but more energetic and speedy in its action.

EMPLASTRUM MELOES VESICATORII, E. lyttæ—e. Meloës vesicatorii comp., E. lyttæ comp.—e. Minio, see Emplastrum Plumbi—e. Mucilaginosum, Unguentum de althæa—e. Nigrum, of Augsburg, see Magnet.

EMPLASTRUM NORIMBERGENSÆ, E. ex. oxido plum'bi rubro camphora'tum, (F.) Emplâtre de Nu'remberg, Nu'remberg plaster. (Oxid. plumb. rubr. p. 300, Ol. oliv. p. 600, Aquæ, q. s. Boil until the oxide is dissolved, and almost to dryness. Remove the vessel from the fire, and add yellow wax, p. 500. Put the vessel again on the fire, and after the wax is melted, add, before it congeals, camphor, p. 24: mix. *Ph. P.* It is considered a desiccative, and has been employed in gangrene.

EMPLASTRUM NYGMATICUM, E. resinæ.

EMPLASTRUM OPII, O'pium Plaster. (Ext. opii, ʒi. Picis Burgund. ʒiij; Emplast. plumbi. ʒj, Aq. bullient. f ʒiv. Melt together the lead plaster and Burgundy pitch; then add the extract of

opium previously mixed with the water, and boil over a gentle fire to the proper consistence.—*Ph. U. S.*) It is employed as an anodyne, and to give support.

EMPLASTRUM OXIDI FERRI RUBRI, *E. Ferri rubri seu Rubraus, Plaster of red oxide of iron, Strengthening Plaster.* (*Empl. oxid. plumb. semivit. p. xxiv, Resinæ pin. p. vi, Cerae flavæ, Olei oliv. Europ. sing. p. iij, Oxidi ferri rubri. p. viij.* Rub the red oxide of iron with the oil, and add the other ingredients melted. *Ph. E.*) It is employed as a strengthening plaster.

EMPLASTRUM OXIDI PLUMBI SEMIVITREI, *E. plumbi—e. Oxidi plumbi semivitrei gummoso, E. gummoso—e. ex Oxido plumbi rubro compositum, E. Norimbergense—e. ex Oxido plumbi semivitreo, E. plumbi—e. Phœniceum, Diapalma—e. Picis, E. Picis Burgundicæ, E. Picis compositum.*

EMPLASTRUM PICIS BURGUNDICÆ, *E. Picis, Bur'gundy Pitch or Pitch Plaster, (Picis Burgund. lbvj, Cerae flavæ. lbss.—Ph. U. S.)*

EMPLASTRUM PICIS CANADENSIS, *Plaster of Canada Pitch, Hemlock Pitch Plaster, (Picis Canadensis. lbvj; cerae flavæ, 3vj. Melt and strain, stirring whilst it thickens. Ph. U. S. 1863.)*

EMPLASTRUM PICIS COMPOSITUM, *E. picis cephalicæ, Cephalic plaster, Compound pitch plaster. (Picis arid. lbij, Abietis resinæ lbj, Resinæ flavæ, Cerae flavæ, aa 3iv, Myrist. ol. 3j, Ol. oliv., Aquæ, aa f3ij. To the pitch, resin, and wax, melted together, add the other matters and mix. Ph. L.)* It is stimulant and rubefacient. Sometimes used in headache,—applied to the temples. See Depilatory.

EMPLASTRUM PICIS CUM CANTHARIDE, *E. Calefaciens.*

EMPLASTRUM DE PICÆ ET RESINIS GLUTINANS, *E. Andréæ de Cruce, (F.) Emplâtre d'André de la Croix, ou collant de poix et de résines, Adhesive plaster of pitch and resins. (Picis albæ. p. 128, Resin. elemi. p. 32, Terebinth. pur. p. 16, Ol. laur. p. 16. Melt with a gentle heat, and pass through linen. Ph. P.)* Used in contusions and fractures as a support.

EMPLASTRUM PLUMBI, *E. lithargyri seu commu'næ seu diach'ylon seu oxidi plumbi semivitrei seu ex oxido plumbi semivitreo, Diach'ylon simplex, White Diach'ylon, Lead or Diach'ylon plaster. (Plumbi oxid. in pulv. 3xxx, Olei oliv. 3lvi, Aquæ q. s. Sift the oxide into the oil; add half a pint of boiling water, and boil until a plaster is formed; adding, from time to time, a little boiling water as that first added is consumed. Ph. U. S.)*

A plaster of red lead, *Emplastum e min'i*, was formerly officinal. It was prepared much in the same way as *Emplastum Plumbi*, but was kept on the fire and stirred, until it became black, when it was called *black plaster*. See *Emplastum oxidi ferri rubri*.

EMPLASTRUM POLYCHRESTUM, *E. Resinæ.*

EMPLASTRUM POTASSII IOIDII, *Plaster of Potass. iodid. (Potass. iodid. 3j, Thuris, 3vj; Cerae, 3vj, Ol. oliv. f3ij. Ph. L.)* This plaster, spread on leather, is applied to indolent tumours.

EMPLASTRUM RANARUM SIMPLEX, see *Emplastum hydrargyri compositum—e. de Ranis sine mercurio, Emplastum hydrargyri compositum.*

EMPLASTRUM RESINÆ, *E. adhæsivum seu lithargyri cum resinâ seu resino'sum seu nymmat'icum seu commu'næ cum resinâ seu Polychrestum seu lithargyricum cum resinâ pi'ni, Res'in, Adhæsive or Stick'ing plaster. (Resin, lbss. Emp. plumb. lbij. Melt the plaster, with a gentle heat, add the resin, and mix. Ph. U. S.)* Employed in wounds and ulcers.

Baynton's adhesive plaster is made by melting one pound of lead plaster and six drachms of resin together.

EMPLASTRUM RESINOSUM, *E. Resinæ—e. Robo-*

rans, E. Ferri (Ph. U. S.), E. oxidi ferri rubri—e. Rubefians, E. Lyttæ.

EMPLASTRUM SAPONIS, *E. saponis acumi, Soap plaster. (Saponis concis. 3iv, Emplast. plumb. lbij. Mix the soap with the melted plaster, and boil to a proper consistence. Ph. U. S.)* It is a mild discutient. Applied to tumours, corns, &c.

EMPLASTRUM SIMPLEX, *E. ceræ—e. Spermatisceti, Ceratum cetacei.*

EMPLASTRUM THURIS, *Frankincense plaster. (Emp. lithargyri, lbij, Thuris, lbss, Oxid. ferri rubri. 3ij, Ph. D.) Use:—the same as the plaster of red oxide of iron.*

EMPLASTRUM VESICATORIUM, *Blister, E. lyttæ—e. de Vigo cum mercurio emendatum, E. hydrargyri compositum.*

EMPLÂTRE, *Emplastum—e. d'André de la Croix, Emplastum de pice et resinis glutinans—e. de Ciqué, Emplastum cicute—e. Collant de poix et de résines, Emplastum de pice et resinis glutinans—e. de Gomme-résine, Emplastum cum gummi-resinis—e. Mercuriel, Emplastum hydrargyri—e. de Nuremberg, Emplastum Norimbergense—e. de Vigo, see Emplastum hydrargyri compositum—e. de Vigo sine mercurio, see Emplastum hydrargyri compositum.*

EMPLATTOMENUS, (*εμπαττομενος*, same etymon as *Emplasticus*.) *Emphracticus.*

EMPNEUMATOSIS, (*εμπνευμασις*, from *em*, *πνευμα*, gen. *πνευματος*, 'air,' and *osis*.) *Emphysema, Inspiration.*

EMPOISONNEMENT, *Poisoning.*

EMPORIUM, (*εμποριον*, from *em*, and *πορεω*, 'I transport or send.') The brain was so called, of old, because there all the mental affairs are transacted.

EMPORIUM SPIRITUM, *Sensorium.*

EMPOSIS, (*em*, and *ποσις*, 'drink.') *Imbibition.*

EMPREINTE, (*em*, and *premere*, 'to press.') *Impression.*

EMPRESIS, (*εμπρησις*, from *em*, and *πρησις*, 'burning.') *Empresina.*

EMPRESMA, gen. *Empresmatis, Empresmus, Empresia*, *Inflammation.* A genus in the class hæmatica, order phlogotica, of Good. *Phlegm'aticæ membranosæ et parynchymatosæ, Phlogis'tici, Fe'bres contin'æ inflammato'rie, Inflammatio inter'na, Cav'ma, Inter'nal inflamm'ation.*

EMPRESMA BRONCHITIS, *Cynanche trachealis—e. Bronchlemmitis, Cynanche trachealis—e. Carditis, Carditis—e. Cephalitis, Phrenitis—e. Cystitis, Cystitis—e. Enteritis, Enteritis—e. Gastritis, Gastritis—e. Hepatitis, Hepatitis—e. Hysteritis, Metritis—e. Nephritis, Nephritis—e. Orchitis, Hernia humoralis—e. Otitis, Otitis—e. Paristhmitis, Cynanche—e. Paristhmitis tonsillaris maligna, Cynanche maligna—e. Paristhmitis pharyngea, Cynanche pharyngea—e. Paristhmitis tonsillaris, Cynanche tonsillaris—e. Parotitis, Cynanche parotidæa—e. Peritonitis, Peritonitis—e. Peritonitis mesenterica, Mesenteritis—e. Peritonitis omentalis, Epiploitis—e. Pleuritis, Pleuritis—e. Pleuritis diaphragmatica, Diaphragmitis—e. Pneumonitis, Pneumonia—e. Splenitis, Splenitis.*

EMPRESMOMANIA, (*εμπρησμος*, 'burning,' and *mania*.) *Pyromania.*

EMPRESMUS, (*εμπρησμος*.) *Empresma.*

EMPRION, (*εμπριων*, from *em*, and *πριων*, 'a saw.') *Serrated.* Galen has given this name to the pulse, when the sensation produced by the artery under the fingers is analogous to that which would be caused by the unequal teeth of a saw.

EMPROSTHOCYRTOMA, (*εμπροσθεν*, 'forwards,' *κυρτος*, 'bent,' and *ωμα*.) *Lordosis.*

EMPROSTHOTONUS, *Emprosthotonia, Entasia tetanus anti'cus, Tetanus anti'cus, (εμπροσθονος, from εμπροσθεν, 'forwards,' and τενω, 'I stretch,' 'I extend.') A variety of tetanus, in*

which the body is drawn forwards by the permanent contraction of the muscles.

EMPSYCHOSIS, (ἐμψύχωσις, (em, and ψυχή, 'life.') 'I animate.' 'I vivify.') A word formerly used for the act of animating. The union of the soul with the body.

EMPTOE, Hæmoptysis.

EMPTOICA PASSIO, Hæmoptysis.

EMPTYSIS, Hæmoptysis.

EMPTYSMA, Sputum.

EMPYE, (ἐμπεύω, Empyema.

EMPYE'MA, gen. *Empyematis*, *Empyem'is*, *Em'pyé*, (ἐμπεύμα, from em, and πύω, 'pus.') *Apostema empyema*. A collection of blood or pus, and, conventionally, of other fluid, in some cavity of the body, and particularly in that of the pleura. Empyema is one of the terminations of inflammation of the pleura, and is called also, *Ptyothorax cævus*, *Phlegmonia purulenta*, *Diaptyema*, *Empyema*, *Empyem'is*, *Empyem'is* seu *Ptyem'is* seu *Hydrothorax purulens*, *Dyspnoea pyothorac'ica*, *Pneumonia suppurativa*, *Abscessus pectoris* seu *Thorac'is*, *Phlegmonia*, *Pleuraptyem'is*, (F.) *Empyema*, *Pyopleuritis*.

The *Operation* for *empyema* properly means the making of an opening into the thorax for the purpose of giving issue to the matter collected in the cavity of the pleura, although it has been used for the operation required for the evacuation of any fluid from the chest, or synonymously with *Paracentesis thoracis*.

EMPYESIS, (ἐμπίσις,) Suppuration. An eruption of phlegmonous pimples, gradually filling with a purulent fluid, and terminating in thick scabs, frequently leaving pits or scars:—*Pustulosa Eczem'ica*. A genus in the order *Eczematologica*, class *Hæmatica* of Groud. See *Emyema*.

EMPHYESIS OCULI, Hypopyon—e. Pectoris, Empyema—e. Variola, Variola.

EMPHYOCE'LE, *Em'phyocele*, (em, πύω, 'pus,' and πύλη, 'tumour,' 'hernia.') A tumour, formed by an accumulation of pus in the serotum. Different diseases have been described under this name, such as suppuration of the testicle, empyema of the tunica vaginalis, accumulation of pus in the cavity of a hernial sac, abscesses of different kinds formed in the areolar texture of the serotum, &c.

EMPYOM'PHALUS, (em, πύω, 'pus,' and ὀμφαλός, 'the navel.') This word has been used to designate a suppurating tumour at the umbilicus; or, at times, umbilical hernia, the sac of which is filled with blood.

EMPYREU'MA, gen. *Empyreumatis*, (ἐμπεύρωμα, from em, and πύω, 'fire.') The burnt smell and acid taste, which volatile products—gaseous and liquid—contract, when animal or vegetable substances are decomposed by a strong heat. The cause of this smell is seated in an oil, called *empyreumatic*, which does not exist in the substance subjected to the operation, but is the result of its decomposition. If the empyreuma occurs when the organic substance is placed in a still with a liquid, it is owing to the solid matter touching the bottom of the vessel to which the fire is applied.

EMPYREUMAT'IC, *Empyreumaticus*. Belonging to empyreuma,—as an *empyreumatic* odour.

EM'PYRUS, (ἐμπίρος, from em, and πύρ, 'fire.') *Febri'ficus*. One who has fever.

EMPUYUS, (ἐμπίρος, from em, and πύω, 'pus.') *Purulent*.

EMS, MIN'ERAL WATERS OF. Celebrated springs on the river Lahn, duchy of Nassau. They are thermal (from 83° to 115° Fahrenheit.) and carbonated salines, containing carbonic acid, bicarbonate of soda, and chloride of sodium; and are much used in gastric and intestinal affection; &

&c. Season from the 1st of June to the 15th of September.

EMUL'GENT, *Emul'gens*, (*emulgere*, (e, and mulgere, vulturn.) 'to milk out,' 'to draw out.') A name given to the renal artery and vein, because the ancients imagined they milked, as it were, the urine from the kidneys. See *Renal*.

EMUL'SIFY, (F.) *Émulsionner*, (*émulsion*, and faire, 'to become.') To make or form an emulsion.

EMULSIN, see *Amygdalin*.

EMUL'SIO, *Emul'sion*, *Emul'sion*; same etymon as *Emulgent*. A pharmaceutical preparation, of a milky-white opaque appearance, which is composed of oil, divided and held in suspension in water by means of mucilage. Emulsions have been divided into the *true* and *oily*, and into the *false* or *not oily*; the latter being composed of resinous substances, balsams, or camphor, rubbed up with dilute alcohol, mucilage, or yolk of egg.

EMULSIO ACA'CLE, *E. Arab'ica*, *Gum Ar'abic Emul'sion*, (*Nucleor. amygd. comm.* ʒi. aquæ Brijss., mucilag. acac. ʒij. sacch. ʒiv. While beating the decorated almonds with the sugar and water, add the mucilage.—*Ph. E.*) Used in the same cases as the next. Under the name *Mistura Acac'ia*, *Gum Ar'abic Mixture*, the Edinburgh Pharmacopoeia had the following. *Mucilag. Acac'ie* f ʒijj; *Amygdal. dulc.* ʒi. and ʒij; *Sacchar.* ʒv; *Aquæ Oij.* (Imper. meas.) Blanch the almonds; peel them; beat to a smooth pulp first with the sugar, and then with the mucilage; add the water gradually, stirring constantly, and strain through linen or calico.

EMULSIO AMYGD'ALE, *E. Sin'plex*, *Lac amygd'ale*, *Amygdalatum*, *Mistura amygd'ale*, *Emul'sio* sive *Lac Amygdalatum*, *Almond emul'sion*, *mixture* or *milk*, (F.) *Lait d'amandes*, (*Amygdal. dulc.* ʒss; *Acac'ie* in pulv. ʒss; *Sacchar.* ʒij; *Aquæ destillat.* f ʒviij. Macerate the almonds in water, and having removed their external coat, beat them with the gum Arabic and sugar, in a marble mortar, till they are thoroughly mixed; then rub the mixture with the distilled water gradually added, and strain.—*Ph. U. S.*) It is used as a diluent and demulcent.

EMULSIO ANTHYSTERICA, *Mistura asafetidae*—e. *Arabica*, *Emulsio acacie Arabice*.

EMULSIO CAM'PHORE, *E. Camphorac'ia*, *Mistura Cam'phora*; *Cam'phor Emul'sion*, (*Camphora* ʒij, *amygd. com. decoctice*, *sacch. dur.* ʒā ʒiv, *aquæ* ʒvj.—*Ph. E.*) A convenient form for giving camphor.

EMULSIO GUAIACINA, *Mistura guaiaci*—e. *Leniens*, *Looch* ex ovo.

EMULSIO O'LEI AMYGDALA'RUM; *Emul'sion of Oil of Almonds*, (*Ol. amygd. ʒi. gum acac. pulv.* ʒij, *syrrup.* ʒi, *aquæ destill.* ʒiv. *Mix.*) A good pectoral or cough mixture.

EMULSIO O'LEI RIC'INI; *Castor Oil Emul'sion*, (*Ol ricini* ʒss, *vitelli ovi* q. s., *aquæ destillat.* ʒj.) An aperient draught.

EMULSIO O'LEI TEREBIN'THINE; *Emul'sion of Oil of Turpentine*, (*Ol. tereb. rect.* ʒij, *sacch. alb.* ʒss, *vitell. ovi* j, *emula. amygd.* vel *aquæ destillat.* ʒvj. *Mix.*) In rheumatic and nephritic affections. Dose f ʒiss.

EMULSIO OLEOSA, *Looch* ex ovo.

EMULSIO PUR'GANS CUM JALAP'E RESI'NA, *Purg'ing Emul'sion with Res'in of Jalap.* (*Jalapae resin.* gr. xij. *sacch. alb.* ʒij. Triturate for some time, and add gradually half the yolk of an egg; continue to triturate, adding by degrees *emula. simpul.* ʒv, *aq. flor. aurant.* ʒij.—*Ph. P.*)

EMULSIO PUR'GANS CUM SCAMM'ONIA; *Purg'ing Emul'sion with Scamm'ony*. It is prepared like the preceding, substituting only *Scammony* for the jalap.

EMULSIO SIMPLEX, *E. Amygdale*.

EMULSION. *Emulsiō*—*c.* Almond. *Emulsiō amygdalæ*—*c.* Campher. *Emulsiō Camphoræ*—*c.* Castor oil. *Emulsiō olei ricini*—*c.* Gum Arabic. *Emulsiō acetiæ Arabicæ*—*c.* *Huileuse*. Looch ex ovo—*c.* of Oil of Almonds. *Emulsiō olei amygdalarum*—*c.* of Oil of Turpentine. *Emulsiō olei terebinthinæ*—*c.* Purging, with resin of jalap. *Emulsiō purgans cum jalapæ resinâ*—*c.* Purging, with scammony. *Emulsiō purgans cum scammonio*.

ÉMULSIONNER, *Emulsify*.

EMULSIVE, *Emulsi'vus*. An epithet given to seeds whence oil may be extracted by expression; such as almonds, apricots, peaches, hemp, rape, melons, gourds; those of the nut kind, and cucurbitaceous and cruciferous plants in general.

EMULSUM, *Emulsion*.

EMUNCTORIUM, *Emunctory*—*c.* Cerebri, *Nasus*.

EMUNCTORY, *Emuncto'rium*, (*emungere* (*e*, and *ungere*, *munctum*), 'to drain off,' 'to cleanse.') *Emissio'rium*, (*F.*) *Emonctoire*, *Émissaire*. Any organ whose office it is to give issue to matters which ought to be excreted. The ancients thought that some organs were more particularly destined to serve as emunctories of others: the nasal fosse, for example, they believed to be the emunctories of the brain.

EMUNDANTIA, (*e*, and *mundare*, 'to cleanse,') *Detergens*.

EMYS PALUS'TRIS, (*εμς*,) *Salt Water Ter-rapin*. This species of turtle is found exclusively in salt or brackish waters, near the sea shore, along the whole Atlantic coast. It is much prized by the epicure, and is nutritious and sufficiently easy of digestion when dressed plain.

EN, in composition, see *En*.

ENADELPHIA, (*en*, and *ἀδελφος*, 'a brother,') *Emboîtement*.

ENÆMA, (*εναμα*, pl. of *εναμων*, from *en*, and *αμα*, 'blood,') *Hæmatostatica*.

ENÆMON, (*εναμων*), *Styptic*.

ENÆORE'MA, gen. *Enæore'matis*, (*εναωρημα*, from *en*, and *αωρεω*, 'I lift up,') 'that which hangs or floats in,') (*F.*) *Enéorème*. *The Neph-elé*, *Nu'ber*, *Sublimatio*, *Subli'mé*, *Suspensum*, *S. Urinæ*, *Suspensæ*, *Nubecula* or cloud which is suspended in the urine as it cools.

ENAM'EL, *E. of the Teeth*, (*en*, and (*F.*) *émail*,) seu *Nitor seu Cortex seu C. striatâ seu Substantia filamentosa* (of Malpighi,) *S. Adamantina seu Vitrea seu Crusta adamantina Dentium*, *Odontophalus*, *Adamantine* or vitreous substance. (*F.*) *Substance vitrée ou émaillée*, *Émail des Dents*. The substance which covers the coronæ of the teeth. The enamel is of a milky white colour, and sufficiently hard to strike fire with steel. Its surface is very smooth and polished, and it forms a thicker layer towards the part where the teeth come in contact, and becomes thinner towards the cervix. The fibres of the enamel are perpendicular to the surface of the teeth, on the ivory of which they seem planted, as it were. This gives them a velvety appearance, when examined by the microscope. The enamel has no blood vessels, and is not renewed when removed. It is formed of phosphate of lime, and a very small portion of animal matter. A delicate membrane covers the enamel—*cuticle of the enamel*, *Nu'amyth'a membrane*; but so closely united with it, that it can be demonstrated only by the use of chlorohydric acid.

ENAMEL, *Cuticle of the*, see *Enamel*.

ENANTHE'MA, gen. *Enanthe'matis*, (*en*, and *ανθω*, 'I bloom,') A name recently given to certain eruptions of the mucous membrane, as *exanthema* is to certain eruptions of the skin.

ENANTHESIS, *Rash exanthem*. Eruption of red, level, or nearly level, patches, variously figured, irregularly diffused, often confluent, and

terminating in cuticular exfoliations. A genus in the order *Exanthematosa*, class *Hemiptera*, of Cullen, including scarlet fever, measles, &c.

ENANTHESIS ROSALIA, *Scarlatina*—*c.* *Urticaria*, *Urticaria*.

ENANTHOPATHIC, (*εναντιος*, 'opposite, and παθος, 'affection,') *Palliative*.

ENARTHRO'SIS, *Articulatio*, *Ball and Sock*, *et-joint*, (*en*, and *αρθρωσις*, 'an articulation,') A kind of diarthrodial articulation, in which the head of a bone is received into the cavity of another, and can be moved in all directions. The joint of the os femoris with the os innominatum is one of this character.

ENARTHURUM, (*en*, and *αρθρον*, 'a joint,') A foreign body in a joint.

ENASMA, gen. *Enas'matis*, (*ενασμα*, 'that which kindles, from *en*, and *αωω*, 'to kindle,') *Fomites*.

ENCAN'THIS, (*εγκανθις*, from *en*, and *κανθος*, 'the angle of the eye,') A tumour, formed by an increase in size, or a degeneration, of the caruncula lacrymalis. Any morbid growth in the inner angle of the eye.

ENCANTHIS BENTIG'NA. Simple excrecence of the caruncula. It commonly yields to astringent collyria.

ENCANTHIS FUNGO'SA. A condition of the semilunar fold and lachrymal caruncle in which they are the seat of morbid growths.

ENCANTHIS INFLAMMATO'RIA. Inflammation with enlargement—swelling—of the semilunar fold and lachrymal caruncle.

ENCANTHIS MALIGNA has often a cancerous character, and requires extirpation before it has attained any considerable size.

ENCARDITIS, *Endocarditis*.

ENCARPUS, (*εγκαρπος*, from *en*, and *καρπος*, 'fruit,') *Pregnant*.

ENCASING, *Emboîtement*.

ENCATALEPSIS, *Catalepsy*.

ENCATHIS'MA, gen. *Encathis'matis*, *Semion-pium*. With the ancients, *Encathis'ma* (*εγκανθισμα*, from *en*, and *καθηω*, 'I seat upon,') meant a vapour-bath taken sitting.

ENCAUMA, gen. *Encau'matis*, (*εγκαυμα*, from *en*, and *καυω*, 'I burn,') *Epicauc'mis*, *Epicauc'sis*, *Encau'sis*. A tumour produced by a burn. A burn. Also, an ulcer of the transparent cornea, occasioning loss of the humours.

ENCAUSIS, (*εγκανσις*,) *Burn*, *Encauma*, *Moxibustion*.

ENCAUSSE, *MINERAL WATERS OF*. Encausse is a village in the department of the Haute-Garonne, which possesses several saline acidulous springs.

ENCAVURE, (*en*, and *caver*, 'to hollow,') *Arcema*.

ENCEINTE, (*sine cinetu*, 'without a zone,') (*Isidore*,) or from *en*, and *cingere*, *cinetum*, 'to surround,') *Pregnant*.

ENCENS, (*incendere*, *incensum*, 'to burn,') *Juniperus lycia*.

ENCENSIER, *Rosmarinus*.

ENCEPHALALGIA, (*encephalon*, and *αλγος*, 'pain,') *Cephalalgia*—*c.* *Hydropica*, *Hydrocephalus internus*.

ENCEPHALIC, *Encephal'icus*, (from *encephalon*,) That which is situate in the head. A name given to several parts which relate to the encephalon, as the *encephalic membrane*, *muscles*, &c.

ENCEPHALITIS, (*encephalon*, and *itis*,) This term has been used, by some nosologists, synonymously with *Cephal'itis* and *Phren'itis*. By others, it has been appropriated to inflammation of the brain, in contradistinction to that of the membranes.

ENCEPHALITIS EXCUDATORIA, *Hydrocephalus internus*—*c.* *Infantum*, *Hydrocephalus internus*

—e. Insolationis, *Coup-de-soleil*—e. Membranosa, Meningitis—e. Peripherica, Meningitis—e. Pustulorum, Delirium tremens.

ENCEPHALIUM, (dim. of *encephalon*.) Cerebellum.

ENCEPHALOCELE, (*enceph'alocele*, (*encephalon*, and *κῆλη*, 'a tumour') *Her'nia seu Fun'gus Cer'e'bri*, *H. Cerebra'lis*, *Craniocele*, *Her'nia of the Brain*. This may be a congenital or accidental affection. In the former case, it is dependent upon tardy ossification of the fontanelles or some faulty conformation. In the latter, it is owing to some considerable loss of substance in the parietes of the cranium, produced by fractures, wounds with a cutting instrument, caries, the application of the trephine, &c. In slight congenital encephalocele, gentle pressure may be exerted upon the protruded portion. When the disease is of considerable extent, it is fatal. In accidental encephalocele, similar means must be used for confining the tumour, and preserving it from the action of external bodies.

ENCEPHALOCYISIS, (*encephalon*, and *χυειν*, 'to pour out,') Hydrocephalus internus—e. Senilis, Apoplexy, serous.

ENCEPHALODYNIA, (*encephalon*, and *δύνη*, 'pain,') Cephalalgia.

ENCEPHALOHÆMIA, (*encephalon*, and *αἷμα*, 'blood,') Cephalæmia.

ENCEPHALOHÉMIE, Cephalæmia.

ENCEPHALOID, *Encephaloï'des*, *Ceph'aloid*, *Ceph'aloï'des*, *Ceph'alo'ma*, (*encephalon*, and *εἶδος*, 'resemblance,') Lænnec has given the term *Enceph'aloid* or *Cer'ebriform* matter to one of the morbid substances commonly formed by scirrhous or cancerous tumors. It is either encysted, in irregular masses without cysts, or infiltrated into the texture of the diseased organ. This name was given to it in consequence of its resemblance, when fully developed, to the medullary substance of the brain. It is also called *Fun'gus medulla'ris seu cancro'sus medulla'ris seu cerebra'lis*, *Sarco'ma medulla'rè*, *Spon'goid inflammation*, *Can'cer mol'lis seu medulla'ris*, *Carcino'ma medulla'rè seu spongio'sum*, *Carci'nus spongio'sus*, *Stru'ta fungo'sa*, *Myelo'ma*, *Myelom'yces*, *Myelospo'ngus*, *Galactom'yces*, *Galactospon'gus*, *Tela accidenta'lis medulla'ris*, (F.) *Matière cérébriforme*, *Carcinome mou et spongieux*, *Tumeur encéphaloïde ou spongieuse ou osseuse ou pulpeuse ou fonguïde ou fongueuse ou spléniforme*, *Fongus médullaire*, *Carcinôme sangiant*, *Cancer mou ou encéphaloïde ou médullaire ou cérébriforme*; *Milt'like tu'mour*, *Med'ullary sarco'ma*, *Cell'ular can'cer*, *Fun'goid* or *Hematoc'er'ebriform* disease.

ENCEPHALOID TUMOUR OF THE LUNG, Phthisis, cancerous.

ENCEPHALOLOG'IA, (*encephalon*, and *λογος*, 'a description,') A description of the encephalon.

ENCEPHALOMALACIA, (*encephalon*, and *μαλακία*, 'softness,') Mollities cerebri.

ENCEPHALOMALACOSIS, Mollities cerebri.

ENCEPHALOMALAXIS, Mollities cerebri.

ENCEPHALOMENINGITIS, Meningocephalitis.

ENCEPH'ALON, (*εγκεφαλον*, from *en*, and *κεφαλή*, 'head,') *Enceph'alum*, *Enceph'alus*, *Enkeph'alon*. That which is situate in the head. This name has generally been given to the *brain*, *cerebellum*, and *mesocephalon*. At times, it includes likewise the *medulla spinalis*, when it is also called the *Cer'ebro-spinal*, and *Neur'al axis*. Under the term *Cra'nio-spinal axis* or *centre of automa'tic actions*, Dr. Carpenter includes the sensory ganglia—the respiratory and stomato-gastric ganglia forming the centre of the *medulla oblongata* and the *spinal ganglia* or true *spinal cord*.

ENCEPHALOPATHIA, (F.) *Encéphalopathie*,

(*encephalon*, and *παθος*, 'disease,') A disease of the encephalon.

ENCEPHALOPATHIA LITERATO'RUM. The condition of brain, supposed to be produced by over-taking it.

ENCEPHALOPATHIA PUERPERALIS, Mania, puerperal—e. Saturnina, see *Encéphalopathie saturnine*.

ENCÉPHALOPATHIE, Encephalopathia—e. *Crapuleuse*, Delirium tremens.

ENCÉPHALOPATHIE SATURNINE, *Encephalopathi'a Saturni'na*. Encephalic disorder occasioned by the poison of lead.

ENCEPHALOPHTHISIS, (*encephalon*, and 'phthisis,') see Encephalopyosis.

ENCEPHALOPHY'MATA, pl. of *Encephalophy'ma*, *Phymato'es seu Strumo'es seu Tuber'cula cer'e'bri*, (*encephalon*, and *φύμα*, gen. *φύματος*, 'a tumour,') Tubercles of the brain.

ENCEPHALOPYO'SIS, (*encephalon*, and *πύω*, 'pus,') *Apost'e'ma seu Absces'sus seu Helco'sis cer'e'bri*. Suppuration of the brain. When accompanied with emaciation and hectic, it is called *Encephalophthi'sis*.

ENCEPHALORACHIDIAN, Cephalospinal.

ENCEPHALORRHAGIA, (*encephalon*, and *ῥαγή*, 'a breaking forth,') see Apoplexy.

ENCEPHALOSCOPIA, (*encephalon*, and *σκοπεω*, 'I examine,') Craniology.

ENCEPHALOSEPSIS, (*encephalon*, and *σπενν*, 'to corrupt,') Mollities cerebri.

ENCEPHALOSIS OF THE LIVER, (*encephalon*, and *osis*,) Hepatoscirrhosis.

ENCEPHALOSISMUS, (*encephalon*, and *σεισμος*, 'a shaking,') Concussion of the brain.

ENCEPHALOSTRUMOSIS, (*encephalon*, and *struma*, 'scrofula,') see Hydrocephalus internus.

ENCEPHALUM, Encephalon.

ENCEPHALUS, (*εγκεφαλος*,) Encephalon.

ENCEPHALUS OPISTHIUS, (*εγκεφαλος* *οπισθιος*, [*οπισθιος*, 'behind,']) Cerebellum.

ENCEROSIS, (*en*, and *κηρος*, 'wax,') Inceration.

ENCHARAXIS, (*εγχαράξις*, from *en*, and *χαράσσειν*, 'to scarify,') Scarification.

ENCHATONNEMENT, Chatonnement.

ENCHEIRE'SIS, *Enchire'sis*, *Enchei'ria*, (*εγχειρησις*, from *en*, and *χειρ*, 'the hand,') Galen uses this term as a part of the title to one of his works, which treats of dissection. It means the manual treatment of any subject.

ENCHEIRIA, Encheiresis.

ENCHIFRÈNEMENT, Coryza.

ENCHIRESIS, Encheiresis.

ENCHONDROMA, gen. *Enchondro'matis*, (*en*, *χόνδρος*, 'a cartilage,' and *ομα*,) Chondroma.

ENCHORIONOSUS, (*en*, *χωρος*, 'a place,' and *νοσος*, 'a disease,') Endemic.

ENCHORIUS, (*εγχωριος*,) Endemic.

ENCHRISTON, (*εγχριστον*, from *en*, and *χρίειν*, 'to anoint,') Liniment.

ENCHUSA, Anchusa.

ENCHYLO'SIS, (*εγκυλωσις*, from *εγκυλιζω*, 'I press out juice,') Extraction.

ENCHYMA, gen. *Enchym'atis*, (*εγχυμα*, from *en*, and *χυειν*, 'to pour,') Infusum, Plethora.

ENCHYMO'MA, gen. *Enchymo'matis*, *Enchymo'sis*, (*εγχυμαμα*, from *en*, *χυω*, 'I pour,' and *ομα*,) By the ancients this word was used to designate the sudden effusion of blood into the cutaneous vessels, which arises from joy, anger, or shame; in the last instance constituting blushing. It differs from ecchymosis in there being, in the latter, extravasation of blood into the areolar texture, and its being produced by an external cause; a contusion, for example.

ENCHYSIS, (*εγχυσις*,) Infusion.

ENCLAVE, (*en*, and (L.) *claudere*, 'to shut,') Wedged.

ENCLAVEMENT, see Wedged.

ENCLUME, Incus.

ENCLYDASTICUS, *intus fluctuans*, (εγκλυδάστικός, from εγκλνδάσσειν (εν, and κλνζειν, 'to wash'), 'to float in.') Floating in or between;—applied to liquids, e. g., to serum, pus, &c., contained in any cavity.

ENCLYSMA, gen. *Enclys'matis*, (εγκλυσμα, from εν, and κλνζειν, 'to wash,') Clyster.

ENCELALIA, (εγκελια, from εν, and κοιλια, 'the belly,') The abdominal viscera. The entrails.

ENCELIALGIA, (*encelalia*, and αλγος, 'pain,') Tortura—e. Inflammatoria, Encelitis.

ENCELITIS, *Encelitis*, *Encelialg'ia* inflammatoria, *Inflammatio abdominalis*; (*encelalia*, and *itis*.) Inflammation of any of the abdominal viscera.

ENCOLEO'SIS, (εν, κολωσ, 'a sheath,') and *osis*.) Invagination.

ENCOLPISMUS, (εν, and κολπος, 'the vagina,') Injection or introduction of any thing into the vagina.

ENCOLPITIS, *Col'pophlegmihymenitis*; (εν, κολπος, 'the vagina,') and *itis*.) Inflammation of the lining membrane of the vagina.

ENCOPE, (εγκοπη, from εν, and κοπτειν, 'to cut,') Diacope, Incision.

ENCRANION, (εν, and κρανιον, 'the skull,') Cerebellum.

ENCRANIS, Cerebellum.

ENCRE, Atramentum.

ENCYESIS, (εγκνσις, from εν, and κνσις, 'pregnancy,') Fecundation, Pregnancy.

ENCYMON, (εγκμων, from εν, and κνειν, 'to be pregnant,') Pregnancy.

ENCYMOSIA, Fecundation, Pregnancy.

ENCYSIS, (εγκνσις,') Pregnancy.

ENCYSTED, *Cyst'idē obduc'tus*, *Sacca'tus*, *Saccula'tus*, *Sac'cated*, *Sac'culated*, *Pouch'ed*. Enclosed in a kyst or cyst, or pouch; (εν, and κνστις, 'a bladder,') (F.) *Encystēd*. An epithet given to certain tumours, or solid or fluid collections enclosed in a particular envelope or cyst. They are movable and often elastic to the touch.

ENCYSTIS, gen. *Encys'tidis*, *Encys'tes*, *Tul'mor tunica'tus* seu *cys'ticus*, *Emph'y'ma encys'tis*. *Lut'p'ia*, *Gland'ula Avicen'næ*, *No'dus*. An encysted tumour.

ENADELPHIA, (*endo*, and αδελφος, 'a brother,') see *Emboitement*.

ENDANGIUM; properly *Endangi'on* or *Endangi'um*, *Membra'na vasoru'm communi's*, (F.) *Tunique commune de Bichat*; (*endo*, and αγγειον, 'a vessel,') The serous or lining membrane of vessels.

ENDARTERITIS, (*endo*, and arteritis 'inflammation of an artery,') Inflammation of the inner coat of an artery.

ENDEICTICUS, (ενδεικτικός,') Indicant.

ENDEIXIOLOGY, *Endeixiolog'ia*, (*endeixis*, and λογος, 'a discourse,') The doctrine of indications.

ENDEIXIS, (ενδειξις, from εν, and δεικνμι, 'to show,') Indication.

ENDEMIA, Endemic.

ENDEMIC, *Endem'ical*, *Ende'mia*, *Regiona'lis mor'bus*, *Ende'micus*; (εν, and δημος, 'the people,') *Encho'rius*, *Enchorion'osus*, *Vernac'ulus* seu *Ende'mius Mor'bus*, *En'demy*, (F.) *Endémie*, *Maladie régionale*. A disease is said to be endemic, (F.) *Endémique*, or to arise from endemic'ity, (F.) *endémicité*, when it is owing to some peculiarity in a situation or locality. Thus, ague is endemic in marshy countries; goitre at the base of lofty mountains, &c. Some authors use the term in the same sense as epidemic. We have no accurate knowledge of the emanations or other circumstances which give occasion to endemic affections. We seem to know that some emanation from marshy lands does produce intermittents: but we are ignorant of the nature of such emanation.

ENDÉMICITÉ, see Endemic.

ENDEMICITY, see Endemic.

ENDEMIOL'OGY, *Endemiolog'ia*, (*endemic*, and λογος, 'a discourse,') The doctrine of endemic diseases.

ENDÉMIQUE, Endemic.

ENEMY, Endemic.

ENDEPIDERMIS, Epithelium.

ENDERMIC, *Ende'matic*, *Ende'micus*, *Endermat'icus*, *Emplas'troēnder'mic*, (F.) *Endermique*, (εν, and δερμα, 'the skin,') An epithet given to the method of treating diseases by placing the therapeutical agent in contact with the skin, especially after the cuticle has been removed; *Enderm'ism*, *Endermis'mus*, *Endermo'sis*, *Meth'odus ende'mica seu endermat'ica*, *Diadermat'ia*. Morphia, strychnia, &c., are often administered in this way.

ENDERMIS, see Endermic.

ENDERMISMUS, see Endermic.

ENDERMOSIS, see Endermic.

ENDESIS, (εν, and δεω, 'I bind,') A ligature, bandage, connexion. Hippocrates has so termed the ankle-joint.

ENDIVE, *Cichorium endivia*—e. Wild, *Cichorium intybus*,

ENDIVIA, *Cichorium endivia*.

ENDIXIS, (εν, and δεικνμι, 'to show,') Indication.

EN'DO, (ενδον, 'within,') as a prefix, within.

ENDOARTITIS, (*endo*, and *aortitis*, 'inflammation of the aorta,') Inflammation of the lining membrane of the aorta.

ENDOARTERITIS, (*endo*, and *arteritis*,) see Arteritis.

ENDOBLAST, (*endo*, and βλαστος, 'a germ,') see Periblast.

ENDOCARDE, Endocardium.

ENDOCARDIAC, *Endocard'ial*, *Endocardi'acus*. Relating to the endocardium, or to the interior of the heart; as '*endocardiac* sound or murmur,'—a sound produced within the cavities of the heart, in contradistinction to *exocardiac*, *exocard'ial*, or *extraocardial* sounds or murmurs, which are induced by conditions of the external surface of the organ.

ENDOCARDITIS, *Encardi'tis*, *Cardi'tis interna*, *Inflammatio superfic'iei inter'næ cord'is*, *Internal Cardi'tis*, *Inflammation of the internal membrane of the heart*, (F.) *Endocardite*: (*endocardium*, and *itis*.) In this disease, the heart's action is visibly increased, and very manifest to the touch; the hand is strongly repelled, and, at moments, is sensible of a trembling vibratory motion. Percussion gives a dull sound over a surface of several inches, owing, according to Bouilland, to the inflammatory turgescence of the heart, and the engorged state of its cavities. On auscultation, a *bruit de soufflet* is generally heard, masking one or both sounds of the heart; and the ear is sensible of a metallic ringing with each systole of the ventricle. The pulsations are rapid as well as strong, and, with few exceptions, irregular, unequal and intermittent. The pulse, by the way, does not always indicate the force or number of the heart's contractions.

ENDOCARDIUM, (*endo*, and καρδια, 'the heart,') (F.) *Endocar'de*. The membrane that lines the interior of the heart.

ENCHORION, see Chorion.

ENDOCOLITIS, (*endo*, *colon*, and *itis*,) Dysentery.

ENDOCYMIEN, (*endo*, and κυμα, 'the product of conception,') A double monster by inclusion. See *Emboitement*.

ENDODONTITIS, (F.) *Inflammation de la Pulpe dentaire*, *Inflammation of the Dent'al membrane*, (*endo*, *odous*, gen. *odontos*, 'a tooth,') and *itis*.) Inflammation of the lining membrane of a tooth.

ENDOENTERITIS. see Enteritis.

ENDOGASTRITIS, *Esogastri'tis*, (*endo*, and *gastritis*, 'inflammation of the stomach.') Inflammation of the lining membrane of the stomach.

ENDOGEN. see Endogenous.

ENDOG'ENOUS, (*endo*, and *γενω*, 'I engender.') A term first applied to plants—hence called *Endo'gens*—in which the new woody matter is deposited within the old, and towards the centre. In the animal, cells are often formed *endog'*-enously, or within the cells, as in the case of the sperm vesicles.

An *endog'*enous an'eurism is one that originates spontaneously from lesions of the inner coat of the arteries.

ENDOLYMPH, *Vitrine auditive*.

ENDOMETRITIS, (*endo*, and *metritis*, 'inflammation of the uterus.') Inflammation of the lining membrane of the uterus. When accompanied with a diphtheritic exudation, it is sometimes called *Endometri'tis croupo'sa*.

ENDONARTERITIS, see Arteritis.

ENDONEPHRITIS, (*endo*, and *nephritis*.) Inflammation of the lining membrane of the pelvis of the kidney.

ENDOPERICARDITIS, (F.) *Endopéricardite*, (*endo*, and *pericarditis*.) Inflammation of the endocardium and pericardium.

ENDOPHLEBITIS, (*endo*, and *phlebitis*.) Phlebitis. Inflammation of the inner coat of a vein.

ENDOPLAST, (*endo*, and *πλασσειν*, 'to form,') see Periplast.

ENDORMIE, (*en*, and *dormir*, 'to sleep,') *Datura stramonium*.

ENDOSIS, (*ενδοσις*, from *en*, and *ιδωμι*, 'to give up,') Remission.

ENDOSKELETON, see Skeleton.

ENDOSMIC, Endosmotic.

ENDOSMOSE', *Endosmo'sis*, *Antoth'esis*, *Antothism'us*, *Petrus'io mutua*, *Double Imbibit'ion*, (*endo*, and *ωσμος*, 'impulse,') A term first used by Dutrochet, to express the action by which fluids pass from without to within organic membranes. The action of two fluids on each other, when separated by a membrane. The general conditions of the phenomena are—first, that they should have an affinity for the interposed membrane; and secondly, that they should have an affinity for each other, and be miscible.

At times, *endosmose* is used to signify the passage of the more transmissible fluid, whilst *exosmose* signifies that of the least transmissible.

The rapidity with which endosmose is accomplished varies according to the nature of the septum or tissue and of the penetrating body, and to the penetrability of the tissue.

ENDOSMOTIC, *Endosmot'icus*, *Endos'mic*, *Endos'micus*; same etymon. Belonging to endosmose:—as an 'endosmotic current.'

ENDOSQUELETTE, see Skeleton.

ENDOSTEITIS, (from *endosteum* and *itis*.) Inflammation of the internal periosteum.

ENDOSTEUM, (*endo*, and *οστεον*, 'a bone,') Medullary membrane.

ENDOÛTERITIS, (*endo*, and *uteritis*;) see Metritis.

ENDUIT (F.), ([L.] *induere*, 'to put upon, 'to put on,') *A coat; a fur*. This term is often applied to a layer of greater or less thickness which covers the surface of certain organs, and particularly of the tongue and the interior of the mouth. The *enduit* is designated variously, according to its appearance,—*enduit bilieux*, *jaune*, *blanc*, &c.—a *bilious*, *yellow*, *white coat or fur*, &c. It is at times owing to the evaporation of the watery portions of the secretions; at others, to a morbid condition of the secretions;—generally, to both causes combined. The tongue, in

such a condition, is said to be *coated* or *furred*; sometimes, *loaded*.

ENDUIT CHOROÏDIEN, see Choroid.

ENDURCISSEMENT, (*en*, and *durcir*, ([L.] *durus*, 'hard,') 'to harden,') Induration—*e. du Cerveau*, *Sclerencephalia*—*e. du Cœur*, *Cardiosclerosis*—*e. Rouge*, see Hepatization—*e. du Tissu cellulaire*, Induration of the cellular tissue.

ENDYMA VENTRICULORUM, (*ενδυμα*, 'a garment,' from *en*, and *δυνειν*, 'to put on,') *Ependyma ventriculorum*.

ENECEMA, (*ενηχημα*, from *en*, and *ηχημα*, 'a sound,') *Tinnitus aurium*.

ENECIA, (*ηνεκσις*, 'continued,') *Synocha*—*e. Cauma*, *Synocha*—*e. Synochus*, *Synochus*—*e. Synochus Puerperarum*, see *Peritonitis*—*e. Typhus*, *Typhus*.

EN'EMA, gen. *Enem'atis*, (*ενημα*, from *ει ημι*, (*ει*, and *ημι*, 'to send in,') *Clyster*.

ENEMA ALOES, *Clyster of Alo'es*. (*Aloës*, $\mathfrak{J}$ ij; *Potass. carbon. gr. xv*; *Decoct. Hordei*, *Oss*, (Imp. meas.,) *M. Ph. L.*) Used in cases of ascariides and amenorrhœa.

ENEMA ANOD'YNUM, *Enema Opii*; *An'odyne* or *Starch* and *O'pium Clyster*. (*Decoct. amyli f3iv*; *Tinct. opii M. xxx. M.*) Exhibited in cases of severe diarrhœa or dysentery.

ENEMA ASSAFETIDÆ, *E. Foetidum*.

ENEMA CATHARTICUM, *Cathart'ic* or *Purg'ing Clyster*. (*Olei. oliv. f3j*; *Sulph. Magnes. 3j*; *Mucilag. acaciae f3xvi*; *M.*)

ENEMA COLOCYNTHIDIS, *Clyster of Col'ocynth*. (*Ext. Colocynth. 3ss*; *Sapon. mollis 3j*; *Aque. Oj*. (Imp. meas.,) *M. Ph. L.*) An efficient cathartic clyster.

ENEMA COMMUNE; *Com'mon* or *Domestic Clyster*. (*Water gruel* or *mollasses* and *water Oss* or *Oj*; add a little oil or lard, and a spoonful of common salt.) Given as a cathartic enema; and, without the common salt, as an emollient.

ENEMA FETIDUM, *E. assafet'idæ*, *Fet'id Clyster*, *Clyster of assafet'idæ*, *Mistu'ra assafet'idæ pro clyst'mate*, *Clyst'ma tonicum* et *antispasmodicum* seu *incitans* et *sedans*, (F.) *Lavement antispasmodique*. (*The last, with the addition of 3j of the tincture of assafetida.*) Given as an antispasmodic and anodyne.

ENEMA MAGNESIÆ SULPHATIS, *E. Catharticum*.

ENEMA NICOTIANÆ; *E. Tabaci*, *Tobac'en Clyster*. This generally consists of from half a pint to a pint of the *Infusum Tabaci*. It is employed in cases of strangulated hernia; but occasionally acts as a fatal poison when given in this way. The smoke of tobacco is sometimes thrown up the rectum to produce the same medicinal effects as the infusion.

ENEMA OPII, *E. Anodynum*.

ENEMA TABACI, *E. Nicotianæ*.

ENEMA TEREBINTHINÆ, *Tur'pentine Clyster*. (*Ol. tereb. 3j*. *Decoct. hordei f3xix*, one yolk of egg. Incorporate the turpentine with the yolk, then add the gruel.) To be administered in cases of ascariides, (*oxyures*.)

ENEPIDERMIC, *Enepiderm'icus*, (*en*, *επι*, 'upon,' and *δερμα*, 'the skin,') An epithet given to the method of treating diseases, which consists in the application of medicines, as plasters, blisters, &c., to the skin.

ENERGIA, *Action*, *Energy*, *Force*.

ENERGY, *Energ'ia*, (*ενεργια*, from *en*, and *εργον*, 'action,') *Action*. *Acting power*. Also, *vigour*; as the '*muscular energy*;' the '*brain acts with energy*.'

ENERVATIO, *Aponeurosis*, *Enervation*.

ENERVATION, *Enervat'io*, (*e*, and *nervus*, 'strength,') The act of weakening—the state of being weakened. See *Debility*.

ENERVATIONES TENDINEÆ, *Inscriptiones tendineæ musculorum*.

ENERVITAS, Inertia.

ENFANCE, Infancy.

ENFANT, Infans—*e. Légitime*, see Legitimate—*e. à Terme*, see Fœtus.

ENFANTEMENT, Parturition.

ENFLURE, (*en*, and *flare*, 'to blow,') Swelling, Edema—*e. des Jambes et des cuisses de la femme accouchée*, Phlegmatia dolens.

ENFONCEMENT SCAPHOÏDE, (*en*, and *fonce*, 'to find the bottom, fond,') Scaphoides fossa.

ENFONCEURE, Arthrodia.

ENFONCURE, Arthrodia.

ENGASTRIMISME, see Engastrimyth.

ENGASTRIMYSME, see Engastrimyth.

ENGAS'TRIMYTH, (*εγγαστρίμυθος*,) *Engastrimythus*, *Englottogastor*, *Gastrilogo'us*, *Ventriloq'us*, *Gastril'oquist*, *Eurycles*, *Eurycl'itus*, *Enteromantis*, *Gastromant'is*, (*en*, *γαστήρ*, 'the belly,' and *μῆσος*, 'I discourse,') *Ventril'oquist*. (*F.*) *Ventriloque*, *Ventriloque*. One who possesses the art of modifying his natural voice, so that it seems to come from a greater or less distance, and from different directions. It was formerly believed that such persons spoke from the belly; hence their name. It is an imitative art, and is called VENTRILQUIZM; (*F.*) *Engastrimisme*, *Engastrimysme*.

ENGELSOMA, Engisoma.

ENGELURE, (*en*, and *geler*, 'to be frozen,') Chilblain.

ENG'HIEU MONTMOR'ENCY, MIN'ERAL WATERS OF. *Engbien-les-Bains*. A hydro-sulphurous water, four leagues from Paris, near Montmorency, which is possessed of some celebrity. It contains chloride of sodium, chloride of magnesium, sulphate of magnesia, sulphate of lime, carbonate of magnesia, carbonate of lime, silica, sulphohydrate of lime and magnesia, sulphohydric acid, carbonic acid and nitrogen. Season, from the first of May to October.

ENGISO'MA, *gen. Engiso'mutis*, *Engiso'ma*, *Engizo'ma*, (*εγγίζω*, 'I approximate,') (*F.*) *Embarre*. A species of fracture of the skull, in which a splinter passes beneath the sound portion of the bone, and compresses the brain. Also, a kind of instrument used in fractures of the clavicle.

ENGLISH DISEASE, Rachitis.

ENGLOTTOGASTOR, (*εγγλोटτογαστωρ*, from *en*, *γλῶττα*, 'the tongue,' and *γαστήρ*, 'the stomach,') *Engastrimyth*.

ENGOMPHOSIS, (*εγγομφωσις*,) Gomphosis.

ENGORGEMENT, (*en*, and [*F.*] *gorge*, 'the throat,') An obstruction occurring in the vessels of a part, giving rise to augmentation of volume. Congestion.

ENGORGEMENT DES MEMBRES ABDOMINAUX À LA SUITE DES COUCHES, Phlegmatia dolens—*e. Hepatic*, *Hepatohæmia*—*e. Latente des membres abdominaux*, Phlegmatia dolens.

ENGOUEMENT (*F.*), *Obstructio*, *Iner'tia*, (*from* [*L.*] *angere*, 'to choke,') Accumulation in a hollow organ, of the matters secreted by it or carried into it. There is said to be *Engouement des bronches*, when the mucus accumulates in the bronchia; and *Engouement des intestins*, when the matters which ought to pass through the intestines are detained; as in a case of strangulated hernia.

ENGOUEMENT DES POUMONS, *E. of the lungs*, in Pathological Anatomy, signifies that state of the lungs, in which a mixture of air and thin fluid escapes from them when cut into.

ENGOURDISSEMENT, Torpor.

ENGRAFT, Graft.

ENGRENURE, Suture.

ENHEMATOSIS, Hæmatosis.

ENIXA, (*enitor*, *enixus*, 'to bring forth,') Puerpera.

ENIXIO FŒTÛS, Parturition.

ENIXUS FŒTÛS, Parturition.

ENKEPHALON, (*εγκεφαλον*,) *Encephalon*.

ENKYSTÉ, Encysted.

ENKYSTEMENT, Chutonnement.

ENMETRITIS, see Metritis.

ENNEAPHAR'MACUS, (*εννεα*, 'nine,' and *φαρμακον*, 'a medicine,') A medicine, composed of nine simple ingredients. A pessary, so formed. The *Antidotus Heracleidis*, described by Galen, and some plasters by Aëtius and Celsus, are, likewise, termed *enneapharmaca*.

ENNEUROSIS, (*en*, and *νευρον*, 'nerve,') Innervation.

ENORMIS, (*e*, and *norma*, 'rule,') Abnormalous.

ENORMITAS CORDIS, Heart, hypertrophy.

ENOR'MON, (*ενορμων*,) *Hor'mon*, *Im'petum fa'ciens*, (*en*, and *ορμω*, 'I rouse,' 'excite,') A word used by Hippocrates in the same sense as *vital principle* is by modern physiologists.

ENOSIS, (*ενωσις*, from *en*, and *ως*, 'one,') In-sertion.

ENOSTO'SIS, *Entostosis*, (*en*, *οσσειον*, 'a bone,' and *osis*,) A morbid growth of bone inward—the opposite to exostosis.

ENRAGE, (*en*, and *rage*, 'rabies,') Rabid.

ENRHUMER, (*en*, and *rhume*, 'rheuma,') See Cold.

ENROUEMENT, (*en*, and (*L.*) *raucus*, 'hoarse,') *Rauco*.

ENRYTH'MOS, (*ενρυθμος*, from *en*, and *ρυθμος*, 'number,') Having rhythm. An epithet applied to the pulse when its pulsations occur with some degree of order. It is opposed to *Aryth'mos*, and differs from *Euryth'mos*, which signifies 'regular.'

ENS. A being. Paracelsus meant, by this term, the power exerted by certain beings on the body. He speaks of the *Ens Del'i*, *E. Astro'rum*, *E. natura'le*, *E. virtut'is*, *E. morbo'rum*, *E. de poten'tibus*, *spirit'ibus*, &c. These absurd denominations suggested to some of the ancient chymists a name for certain chymical preparations. The muriate of ammonia and iron was called *Ens martis*; the muriate of ammonia and copper, *Ens ven'eris*; and *Ens pri'mum* was, with the alchemists, the name of a tincture which they considered possessed of the power of transmuting metals.

ENS MARTIS, *Ferrum ammoniatum*—*e. Veneris* Boyle, *Ferrum ammoniatum*.

EN'SIFORM, *Ensiform'is*, (*ensis*, 'a sword,' and *forma*, 'form,') *Sword-like*.

ENSIFORM APOPHY'SES or PROC'ESSES are the lesser alæ of the sphenoid bone.

ENSIFORM APPEN'DIX or CAR'TILAGE is the xiphoid appendix of the sternum. See Xiphoid.

ENSIFORMIS, Xiphoid.

ENSISTERN'AL. Relating to the ensiform process of the sternum. Bèclard gave this name to the last osseous portion of the sternum. He also called it *l'os ultimi-sternal*. See Ultimosternal.

ENSOMATO'SIS, (*ενσωματωσις*, from *en*, and *σωμα*, *gen. σωματος*, 'a body,') Incorporation.

ENSTALAX'IS, (*en*, and *σταλαζειν*, 'to drop,') Instillation.

EN'STOPHÉ, (*en*, and *στροφω*, 'I turn,') Inversion of a part, as of the eyelids.

ENTAILLE (*F.*), (*en*, and *tailler*, 'to cut,') *Excis'io*, *Ec'copé*, *Ec'tomé*. A deep wound made by a sharp instrument obliquely. *Entailles* and *Taillades* are, also, used to designate deep scarifications, made for the purpose of producing a speedy disorgement of any 'unmedicated part, such, for

ENTERHELCOSIS, Enterelcosis—e. Nervosa, Typhus abdominal.)

ENTEROCĒLĒ, *Enterocele*, *Hernia intestinalis*, (εἰσροσκήλη, from *enteron*, and *σκήλη*, 'a hernia, tumour.') Abdominal hernia, which contains only a portion of intestine, is so called.

ENTEROCELETES, (εντεροκηλη-της,) *Enterocel'*

Ulus (εἰτεροκηλικός.) Affected with intestinal hernia.

ENTEROCEPHALOPYRA INFANTUM, (*enteron*, κεφαλή, 'head,' and πυρ, 'fire or fever.') Hydrocephalus Internus.

ENTEROCE'LICUS, (*enteron*, and κοιλία, 'the abdomen.') Relating to the cavity of the abdomen.

ENTEROCOLITIS, (F.) *Entéro-colite*, (*enteron*, and *colitis*, 'inflammation of the colon.') Inflammation of the small intestine and colon.

ENTEROCYSTOCE'LE, *Enteroceystocele*, (*enteron*, κύστις, 'a bladder,' and κήλη, 'a tumour.') Hernia formed by the bladder and a portion of intestine.

ENTERODAR'SIS, (*enteron*, and δαρσις, 'skinning.') Excoriation of the mucous membrane of the intestines.

ENTERODOTHIENIA, (*enteron*, and δόζηνη, 'a pustule.') Dothineritis.

ENTERODYNIA, (*enteron*, and ὁδυνή, 'pain.') Enteralgia.

ENTEROËPIPLOCE'LE, *Enteroëpiplocele*, *Enteroploce'le*, *Her'nia intestin'lis oment'lis*, (*enteron*, επιπλόν, 'the omentum,' and κήλη, 'tumour.') Hernia, formed by intestine and omentum.

ENTEROËPIPLOM'PHALUS, (*enteron*, επιπλόν, 'the omentum,' and ὀμφαλός, 'the umbilicus.') Umbilical hernia, containing intestine and omentum. Almost all umbilical herniæ are of this kind.

ENTEROGRAPHY, *Enterographia*, (*enteron*, and γραφή, 'description.') The part of anatomy which describes the intestines.

ENTEROHEMORRHAGIA, *Hæmatochezia*.

ENTEROHELCO'SIS, *Enterelco'sis*.

ENTEROHYDROCE'LE, *Enterohy'drocele*, (*enteron*, ὕδωρ, 'water,' and κήλη, 'tumour.') Intestinal hernia complicated with hydrocele.

ENTEROHYDROM'PHALUS, (*enteron*, ὕδωρ, 'water,' and ὀμφαλός, 'the navel.') Umbilical hernia, in which the sac contains, along with a portion of intestine, a quantity of serum.

ENTEROÏSCHIOCE'LE, *Enteroïsch'ioccele*, (*enteron*, ἰσχίον, 'ischium,' and κήλη, 'tumour.') *Ischia'ic her'nia*, formed of intestine.

ENTEROLITHI'ASIS, (*enteron*, and λιθίασις, 'the formation of stone.') The formation of concretions in the intestines.

ENTEROLITHUS, see *Calculi of the stomach and intestines*—e. *Bezoardus*, *Bezoar*—e. *Seybalum*, *Seybala*.

ENTEROLOG'Y, *Enterolog'ia*, (*enteron*, and λόγος, 'a discourse.') The part of anatomy which treats of the intestines.

ENTEROMALA'CIA, *Enteromalax'is*, (F.) *Ramollissement de l'Intestin*, (*enteron*, and μαλασσω, 'I soften.') Softening of the mucous or other coats of the intestine.

ENTEROMALAXIS, *Enteromalacia*.

ENTEROMANTIS, (*enteron*, and μαντις, 'a diviner.') Engastrimyth.

ENTEROMEROCE'LE, *Ent'rome'rocele*, (*enteron*, μηρός, 'the thigh,' and κήλη, 'tumour.') Cru'al hernia, formed of intestine.

ENTEROMESENTER'IC, *Enteromesenter'icus*, (F.) *Entéro-mésentérique*. Relating to the intestine and mesentery.

ENTEROMESENTER'ICA FEBRIS. MM. Petit and Serres have given this name to the typhoid form of adynamic fever, in which the intestines are ulcerated, with enlargement of the corresponding mesenteric glands. See Typhus.

ENTÉROMÉENTERITE, *Tabes mesenterica*—e. *Mésentérite typhoïde*, see Typhus.

ENTEROMOR'PHA COMPRES'SA, (*enteron*, and μορφή, 'shape.') One of the algae, used by the Sandwich Islanders as an esculent.

ENTEROMPHALOCELE, *Ent'rom'phalocèle*, *Enteromphalus*.

ENTEROM'PHALUS, *Enteromphalocèle*, (*enteron*, and ὀμφαλός, 'umbilicus.') Umbilical hernia, formed of intestine.

ENTEROMYCODORITIS, (*enteron*, μύκος, 'slime,' ὄρα, 'skin,' and ἰτις, 'inflammation of the mucous membrane of the intestines.

ENTERON, (*εντερον*, from *ενω*, 'within.') Intestine.

ENTEROPARAL'YSIS, *Enterople'gia*, *Paral'ysis intestin'rum*, (*enteron*, and παραλυσις, 'paralysis.') Paralysis of the intestines.

ENTEROPARISAGOGE, (*enteron*, παρα, *eis*, 'in,' and αγω, 'to lead.') Intussusception.

ENTEROPATH'IA, *Enterop'athy*, (*enteron*, and παθος, 'disease.') Disease of the intestines in general.

ENTEROPATH'IA CANCERO'SA, *Enteroscirrhus*, *Enterostenosis scirrhus* seu *organ'ica*, *Enterosarcoma*, *Indura'tio seu Scirrhus seu Can'cer seu Carcino'ma Intestino'rum*, (F.) *Cancer des Intestins*. Cancer of the intestines.

ENTEROPERISTOLE, *Enterosp'ig'ma*, *Hernia incarceration'ia*, *Incarcera'tio intestin'rum*, (*enteron*, and περιστρέλλω, 'I contract.') Constriction or obstruction of the intestines, from a cause which acts either within the abdomen, or without it, as in strangulated hernia.

ENTEROPHLOGIA, (*enteron*, and φλεγω, 'I inflame.') Enteritis.

ENTEROPHLOGOSIS, *Enteritis*.

ENTEROPHTHISIS, (*enteron*, and φθίσις, 'consumption.') Consumption of the bowels. Consumption owing to suppuration in the intestines.

ENTEROPHYMATA, (*enteron*, and φυμα, 'a swelling.') *Tuber'cula intestin'rum*. Tubercles of the intestines.

ENTEROPLASTY, *Ent'roplas'ticé*, (*enteron*, and πλασσω, 'I form.') A plastic operation for the restoration of an intestine.

ENTEROPLÉGIA, (*enteron*, and πλῆγη, 'a stroke.') Enteroparalysis.

ENTEROPYRA, (*enteron*, and πυρ, 'fire or fever.') Enteritis. Also, enteromesenteric fever, *Entéropyrrie* of Alibert.

ENTEROPYRA ASIATICA, *Cholera*—e. *Biliosa*, *Fever*, *bilious*.

ENTERORRHAGY, *Enterorrhagia*.

ENTERORRHAPHIA, *Enterorrhaphia*.

ENTERORRHAG'IA, *Enteror'rhagy*, *Enteror'hagy*, (*enteron*, and ραγή, 'violent rupture.') Hemorrhage, or excessive discharge from the bowels.

ENTERORRHAGIA SIMPLEX, *Hæmatochezia*.

ENTERORRHAGY, *Enterorhagia*.

ENTERORRHA'PHIA, *Enteror'rhap'hē*, *Enteroraph'ia*, *Enteror'rhapsy*, (*enteron*, and ράφη, 'a suture.') Suture of the intestines, for the relief of injuries done to them.

ENTERORRHEU'MA, gen. *Enterorrheum'atis*, *Rheumatismus Intestino'rum*, (*enteron*, and ρευμα, 'defluxion, rheumatism.') Rheumatism of the intestines.

ENTERORRHEXIS, (*enteron*, and ῥήσσω, 'to break.') Enterobrosis.

ENTERORRHEA, (*enteron*, and ῥέω, 'so flow.') Diarrhoea, Melæna.

ENTEROSARCOCE'LE, *Ent'erosar'cocele*, (*enteron*, σαρξ, 'flesh,' and κήλη, 'a tumour.') Intestinal hernia, complicated with fleshy excrescence, or rather sarcocele.

ENTEROSARCOMIA, (*enteron*, and σαρξ, gen. *σαρκος*, 'flesh.') Enteropathia cancerosa.

ENTEROSCHEOCE'LE, *Enteros'cheocele*, *Enteroschoce'le*, (*enteron*, σκην, 'the scrotum,' and κήλη, 'tumour.') Scrotal hernia consisting of intestine.

ENTEROSCIRRUS, *Enteropathia cancerosa*.

ENTERO'SES, (*εντερον*, 'intestine.') A class of

diseases, comprehending all those that are seated in the intestines.

ENTEROSPHERMA, (*enteros*, and *sphera*, 'to bind tightly.') Enterospiracle.

ENTEROSTENOSIS, (*enteron*, and *stenosis*, 'stricture.') Stricture or constriction of the intestines. See *Ileus*.

ENTEROSTENOSIS ORGANICA, Enteropathia cancerosa—c. *Sclerosa*, Enteropathia cancerosa—c. *Volvulus*, *Ileus*.

ENTEROTOME, *Enterotomus*, (*enteron*, and *temo*, 'I cut.') A name given by J. Cloquet to an instrument for promptly opening the intestinal canal through its whole extent. It consists of a pair of scissors, one blade of which is much longer than the other, and rounded at its extremity. This is passed into the intestine.

ENTEROTOMIA, *Enterotomia*. In anatomy, it signifies dissection of the intestines. In surgery, it means an operation, little used, which consists in opening the intestines, in order to evacuate the fecal matters accumulated in it; for example, in certain cases of hernia, accompanied with contraction of the intestinal canal; in operations for an artificial anus, performed on the newborn, where the rectum is imperforate or not existing, &c.

ENTEROTOMY, Enterotomia.

ENTEROTYPHUS, see Typhus.

ENTEROZOA, (*entero*, and *zoo*, 'an animal.') Worms.

ENTERYDROCELE, *Enterohydrocele*, (*enteron*, 'water,' and *cele*, 'rupture.') Intestinal hernia, with hydrocele.

ENTHELMINTHES, (*ento*, and *helmos*, gen. 'helmos,' 'a worm.') Worms.

ENTHEOMANIA, (*enthos*, *en*, and *theos*, 'God,') 'inspired,' and *mania*, 'Demonomania.

ENTHETIC, *Entheticon*, (*enthos*, *en*, and *thetis*, 'placed, introduced.') Put in, introduced,—as *Enthetia* diseases, *Enthetici* (*Morbi*) of Dr. William Farr, those produced by inoculation or infection.

ENTHLASIS, *Enthlasis*, *Illio*, (*en*, and *thlao*, 'I break.') A fracture of the cranium, with comminution, in which the bone is depressed, or has lost its level.

ENTELASIS CRANII, *Efractura Cranii*.

ENTITÉ, Entity.

ENTITY, *Entitas*, (*F.*) *Entité*, (*ens*, gen. *entis*, 'a being.') The being or essence of anything. It has been used somewhat vaguely, in modern French medicine more especially, to signify apparently a general or essential disease, the nature and seat of which cannot be determined.

ENTIC, (*en*, 'within.') In composition, within.

ENTOBLAST, (*ento*, and *blastos*, 'a germ.') See Cytoblast, and Molecule.

ENTOCELE, (*ento*, and *cele*, 'rupture.') Ectopia.

ENTOCÉLÉ LENTIS, Dislocation of the crystalline into the anterior chamber of the eye.

ENTOGONYAN'CON, *En'au on'con*, *Calif'knee*, *X leg*, (*ento*, *gon*, 'the knee,' and *con*, 'a bend,') Bending of the knees inwards. The state of being *knuckled* or *in-kneed*. [*Prov.*] *Knophweed*, (*Sc.*) *Nitcheed*.

ENTOHYALOID MUSCLE, see Metamorphopsia.

ENTONIA, (*en*, and *tonia*, 'Tension, Tonic spasm.

ENTONIC, *Entonicus*, *En'tonne*, (*en*, and *tonos*, 'tone,') Having great tension or exaggerated action.

ENTONNOIR, (*en*, and *tonne*, 'a tun,') Calix. Infundibulum—*du Ventricle moyen du Cerveau*, Infundibulum of the brain.

ENTONUS, (*en*, and *tonos*, 'Tonic.

ENTOPARASITES, Worms.

ENTOPHTHALMIA, (*ento*, and *ophthalmia*,) Ophthalmia affecting the interior of the eye.

ENTOPHYTE, *Entophyton*, pl. *Entophyton*, (*ento*, and *phyton*, 'a vegetable,') A vegetable parasite, and especially one growing in the interior of the body. For the chief entophytes, see Fungus.

ENTOPHYTIC, *Entophyticus*. Of or belonging to an entophyte,—as 'an entophytic growth.'

ENTOPTERYGOID, see Pterygoid.

ENTOPTIC, *Entopticus*, (*ento*, and *opsis*, 'vision,') Relating or appertaining to the vision of objects within the organ of sight,—as of the bloodvessels of the retina.

ENTORSE, (*en*, and *torquere*, *torquem* 'to twist,') Sprain.

ENTOSTHIDIA, (*en*, and *sthesia*,) Entrails.

ENTOSTHIA, (*en*, and *sthesia*,) Entrails.

ENTOSTHIOBLAST, (*en*, and *sthesia*, 'within,') and *blastos*, 'a germ,') See Cytoblast, and Molecule.

ENTOSTOSIS, (*ento*, and *stosis*, 'a bone,') Enostosis.

ENTOTORRHEA, (*ento*, and *torrhoea*,) See Otorrhoea.

ENTOZENA, (*ento*, and *ozena*,) See Ozena.

ENTOXICISMUS, (*en*, and *toxicum*, 'a poison,') Poisoning.

ENTOXISMUS, Poisoning.

ENTOZOA, (*ento*, and *zoo*, 'an animal,') pl. of *Entozoon*, Worms.

ENTOZOAIRES, Worms.

ENTOZOARIA, Worms.

ENTOZOGENESIS, (*entozoa*, and *genesis*, 'generation,') Helminthiasis.

ENTOZON, see Entozoa—c. *Folliculorum*, *Acarus Folliculorum*.

ENTRAILLES, Entrails.

ENTRAILS, *Splanchna*, *Ence'lia*, *Viscera*, *Entostidia*, *Entosthia*, *Entosthesidia*, *Intera'na*, (from *entera*, a word of low Latin, coming from *entron*, 'intestine,') It is used for the viscera enclosed in the splanchnic cavities, and especially for those in the abdomen; *Bauchs*, *Guts*, (*F.*) *Entrailles*, *Viscères*. *Enca* has had this latter signification, but it was most frequently used by the ancients, as by Pliny, for the thoracic viscera.

ENTRAINEMENT, Training.

ENTRECROISSEMENT, 'Intercrossing.' The reciprocal passage of fibres, and especially of nerve fibres, from one side of a median line to the other.

ENTREFESSON, (*entre*, 'between,' and *fesses*, 'the buttocks,') Chafing.

ENTRICHOMA, (*en*, and *trichoma*, (*Sci*, gen. *trich*, 'hair,') The tarsal cartilage, and the edge of the eyelids, in which the cilia are implanted.

ENTROPION, *Entropion*, (*en*, and *trion*, 'I turn,') *Inversio palpebrarum*, *Oculitis*, *Trichiasis*, *Intramissionis* seu *Biophthalmos* *entropion*, *Biophthalmos*, *Trichiasis*, *Trichiasis*. A name given to the inversion or turning inwards of the eyelids, so that the eye-lashes are directed towards the globe of the eye; irritate and inflame it, and give rise to the affection called *Trichiasis*.

The contact of the hair with the surface of the eye occasions considerable irritation of the conjunctiva, which is soon followed by chemosis, ulceration of the eye, and other symptoms, such as fever, watchfulness, &c.

If the disease be *entropion*, as above defined, that is, dependent upon the inversion of the eyelids, it must be remedied,—either by dispersing the oedema or swelling of the eyelids, or by cutting out a portion of the skin. When the disease is dependent on a vicious direction of the cilia, they must be plucked out, and the bulbs be cauterized.

ENTROPIUM, Entropion.

ENTYPOSIS, (εντυπωσις,) see Glene.

ENUCLEATION, *Enucleatio*, (enucleare, *enucleatum*, (e, and *nucleus*,) 'to take out a kernel.') This term has been proposed for the operation of removing tumours, &c., without cutting into them.

The word is used in *Pharmacy* for the operation of shelling or removing the kernels of any nut.

ENULA CAMPANA, *Inula Helenium*.

ENU'LON, (ενουλόν, from *en*, and ουλόν, 'the gum.') The inner part of the gums. The external part was called ουλόν, and that between the teeth *apnos*.

ENURE'SIS, (ενουρεω, (en, and ουρεω,) 'I void the urine in bed.') *Paru'ria incont'ens*, *Incont'inentia urinae*, *Excre'tio Urinae involunta'ria seu inopportuna*, *Mictio involunta'ria*, *Uracra'tia*, *Uorrhoe'a*, *Perirrhe'a*, *Anischu'ria*, *Hyperure'sis*, *Dian'nes*, *Involuntary discharge' of u'rine*, *Incont'inentine of u'rine*. This affection is most common in advanced life. It may depend on too great irritability of the bladder, or on distension, or injury of the fibres about its neck, paralysis of the organ, *Cystoparal'ysis*, *Cystople'gia*; the presence of an irregularly-shaped calculus impacted in the urethra near its commencement; rupture of the bladder and urethra; renal disease; or on pressure exerted on the bladder by the distended womb or by a tumour. It often occurs after difficult labour, but generally yields in the course of a week or ten days; the catheter being introduced twice a day in the mean time.

The treatment must of course vary according to the cause; and when the affection on which it is dependent cannot be removed, the discharge of urine may be prevented, by exerting a degree of compression on the urethra by means of appropriate instruments; or a urinary may be employed to receive the urine as it passes off.

ENVIE, (from *Li* *invidia*.) Hangnail, Malacia, *Nævus*—*c. de Vomir*, *Nausea*.

ENYPNION, (ενυπνιον, from *en*, and 'νπιος, 'sleep.') *Somnium*.

ENYSTRON, (ηνυστρον, from *ανειν*, 'to complete.') *Abomasus*.

ENZOÏTIA, *Enzooty*.

ENZOÏTY, *Enzoïtia*, (F.) *Enzoïtie*, (en, and ζων, 'animal.') An endemic disease attacking animals.

EP, EPH, EPI, (επ, εφ, επι, 'upon, above:') in composition, generally means 'augmentation, addition, increase, reciprocal action, repetition.' Hence:

EPACMAS'TICUS, (επακμαστικός, from *ep*, and ακμαζω, 'I increase.') An epithet applied to fevers, *Fe'bres epacmas'ticæ*, whose symptoms go on augmenting in violence, from the commencement to the termination. Such fevers are, also, called *Epanadidon'tes*.

EPAGOGITIS, (επαγωγιον, 'prepuce,' and *itis*, inflammation.) *Posthitis*.

EPAGOGIUM, (επαγωγιον, from *ep*, and αγειν, 'to draw.') *Phimosis*, *Prepuce*.

EPANADIPOLOSIS, (επαναδιπλωσις, from *ep*, and αναδιπλωσις.) *Anadiplosis*.

EPANALEPSIS, (επαναληψις, from *ep*, and αληψις.) *Anadiplosis*, *Analepsia*.

EPANASTASIS, (επαναστασις, from επανιστημι, 'to cause to rise up.') The formation of a tumour. The breaking out of an eruption—*Eruptio exanthem'atilis*. A pustule.

EPANASTEMA, *gen. Epanaste'matis*, (επαναστημα.) *Exanthem*, *Swelling*.

EPANCHEMENT, (ε, and pandere, *pansum*, 'to spread.') *Effusion*.

EPANESIS, (επανεσις, from επανιστημι, 'to remit.') *Remission*.

EPANETUS, *Remittent*—*c. Hecticus*, *Hectic*

fever—*c. Malignus flavus*, *Fever*, *yellow*—*c. Mitis*, *Remittent fever*.

EPANORTHOSIS, (*ep*, and αναρθω, 'to make straight.') *Restauratio*.

EPANTHEMA, *gen. Epanthe'matis*, (*ep*, and ανθος, 'a flower.') *Exanthem*.

EPANTHISMA, *gen. Epanthism'atis*, *Exanthem*.

EPAPH'RESIS, (επαραιρω, (ep, απο, and αρω, 'I take,') 'I take away.') *Repetit'io sublat'io seu evacuat'io*. A repeated abstraction or evacuation. It was formerly used synonymously with *repeated blood'letting*; *Phlebotomia iterata*.

EPAPHRISMUS, (*ep*, and αφρος, 'froth.') *Despumation*.

EPARMA, (επαρμα, from *ep*, and αρειν, 'to use up.') *Tumour*.

EPARSIS, (επαρσις,) *Tumour*.

ÉPAULE, (old (F.) *espaule*, from σπαθη, *spatha*, 'the scapula.') *Humerus*.

EPENCRA'NIS, (*ep*, en, and κρανιον, 'the skull.') *Cerebellum*.

EPENDYMA, *gen. Ependym'atis*, (*ep*, and ενδυω, (επ, and δυω,) 'I enter:' hence, επενδυμα, 'an upper garment or cloak.') *Membrane*.

EPENDYMA VENTRICULO'RUM, *Endy'ma seu Indumen'tum ventriculo'rum*. The lining membrane of the ventricles of the brain, formed by a prolongation of the pia mater, and probably also of the arachnoid.

ÉPERON, (old (F.) *esperon*, 'spur,') *Hip-pocampus minor*—*é. des Artères*, *Spur*.

ÉPERVIER, *Accipiter*.

ÉPERVIERE DES MURAILLES, *Hieracium murorum*.

EPH, in composition, see *Ep*.

EPHEBÆ'ON, (εφηβαϊον,) *Epheba'tum*, (*eph*, and 'ηβη, 'pubes,') The parts of generation; also, the region of the pubes, as well as the hair upon the pubes. It meant, likewise, a part of the ancient gymnasium in which the rhetoricians, philosophers, &c., disputed.

EPHEBÆUM, *Mons veneris*.

EPHEBIA, or EPHE'BIA, (εφηβεια,) *Pubertas*. The age of puberty.

EPHEBOSYNE, (εφηβοσυνη,) *Puberty*.

EPHEBOTES, (εφηβοτης,) *Puberty*.

EPHEBUS, (εφηβος,) *Pu'bens*, *Pu'ber*, (F.) *Pu-bère*, *Éphèbe*. One who has attained the age of puberty.

EPHEDRANA, (εφεδρανα, from *eph*, and έτρα, 'a seat,') *Nates*.

EPHEL'CIS, (*eph*, and 'ελκος, 'an ulcer,') The crust or scab of an ulcer.

ÉPHELIDE SCORBUTIQUE, *Chloasma*.

EPHELIDES, (εφηλίδες, from *eph*, and 'ηλιος, 'the sun,') *Epichro'sis*, *Mac'ula sola'ris*, *Mac'ule fus'cæ seu lenticula'res*, *Vitilig'ines*, *Pha'ci*, *Pan'nus lenticula'ris*, *Lentic'ula*, *Lentigo*, *Eph'e'lis Lenti'go seu Lentic'ula seu d' sol'is*, *Nigre'do d' sol'is*, *Spilo'sis eph'e'lis*, *Æsto'tes*, *Eff'lu*, *Freck'les*, *Sun-burn*, (Prov.) *Fernitickles*, *Funtickles*, *Fornitickles*, *Fernfreckles*, *Fanfreckles*, *Fornpeckles*, *Freckens*, *Murples*, (F.) *Taches de rousseur*. This term includes not only the yellow *lentiginæ*, which appear on persons of a fair skin, and the larger brown patches, which also arise from exposure to the direct rays of the sun, but also those large dusky patches which are very similar in appearance, but occur on other parts of the surface, that are constantly covered. See *Chloasma*. They do not extend farther than the skin. Many cosmetics have been recommended for their removal. Simple spirituous lotions or weak mineral acids, applied a few times in the day, are as effectual as any other means.

One, affected with freckles, is said to be *freckled*, (Sc.) *Faintickled*, *Fernitickled*, (Prov.) *Fam'tickled*, *Funtickled*, *Fernfreckled*, *Fornfreckled*, *Freckened*, *Murpled*, (F.) *Tacheté de Rousseurs*.

EPHELIS, (ἐφήλις,) see Ephelides—e. Lentigo, Ephelides—e. à Sole, Ephelides.

EPHELOTES, (ἐφηλότης, from *eph*, and ἦλος, 'a nail.') Leucoma.

EPHEMERA, *Diurnia*, (ἐφήμερος, 'lasting a day,' from *eph*, and ἡμέρα, 'a day.') That which continues a day. An epithet given to diseases, and particularly to fevers, which last only a day.—*Fébris diurnia* seu *simplex* seu *Ephemera*, *Ephemeroptera*, *Fébricula*, *Ephemeral*, *D'ary* or *Simple Féver*, (F.) *Fievre éphémère*. The term *prolonged ephemerata* is sometimes used for fevers, which cease after two or three days' duration.

EPHEMERA ANGLICA PESTILENS, *Sudor anglicus*—e. Britannica, *Sudor Anglicus*—e. Maligna, *Sudor Anglicus*—e. Mortifera, *Plague*—e. Pestilentialis, *Plague*—e. Sudatoria, *Sudor Anglicus*.

EPHEMEROPYRA, (ephemera, and πυρ, 'fire,' 'fever.') Ephemera.

EPHEMERUS, (ἐφήμερος,) Quotidian.

EPHIALTES HYPOCHONDRIACA, (ἐφιάλης, from *eph*, and ἄλλομαι, 'to leap.') Incubus vigilantium—e. Nocturnus, Incubus—e. Vigilantium, Incubus vigilantium.

EPHIDRO'SES, (G.) *Ephidrosen*. A division of the family *Ekzematosen*, of Fuchs, comprising morbid varieties of sweating.

EPHIDROSIS, (ἐφίδρωσις, from *eph*, and ἵδρω, 'I sweat.') *Hidro'sis*, *Idro'sis*, *Suda'tio*, Sweating. Of old, it meant, at times, a slight sweating, *sudation'cula*, of no value, and of bad augury; at others, a good and critical sweat. It has likewise been used for a copious sweat.

EPIDROSIS, Desudatio—e. Cruenta, *Sudor cruentus*—e. Saccharata, *Sudor dulcis*.

EPIDROSIS PROFUSA, *E. spontanea* seu *idiopathica*, *Hydropedesis*, *Exsudatio*, *Exudatio*, *Hyperphidrosia*, *Hyperhydro'sis*, *Hydroorches'a*, *Hydropedesis*, (F.) *Flux de Sueur*. An excessive sweating. A colliquative sweat. Such copious perspirations are generally owing to debility, and require the use of tonics, especially of the mineral acids, opium, &c.

EPIDROSIS SACCHARATA, *Sudor dulcis*.

EPHIPPIUM, (ἐπιππιον, from *eph*, and ἵππος, 'a horse.') *Sella Turcica*.

EPH'ODOS, (εφ'οδος, from *eph*, and 'οδος, 'a way,' 'a road or avenue to a place.') Hippocrates uses this word in three senses: 1. For the vessels or canals, which give passage to the excrements of the body. 2. For a periodical attack or accession of fever; and 3. For the approach of similar or dissimilar things which may be useful or hurtful to the body.

EPH'RATA SPRINGS. These are in Lancaster County, Pennsylvania. The water is pure; and facilities are afforded for bathing—warm and cold.

EPI, in composition, see Ep.

EPI, (old (F.) *epi*.) Spica.

EPI'ALOS, (ἐπι'αλος,) *Hepi'alos*, *Quer'cera*, *Quer'quera*, a name given by the ancients to fever, when the hot stage was mingled with irregular chills. Also, the cold stage of fever.

EPIALTES, (ἐπι'αλης, from *epi*, and ἄλλομαι, 'to leap.') Incubus.

EPIAN, *Frambesia*.

EPIBLEMA, gen. *Epible'matis*, (ἐπιβλημα,) *Catoblema*.

EPIBOLE, (ἐπιβολή,) Incubus.

EPICANTHIS, *Epican'thus*, (from *epi*, and κανθός, 'the angle of the eye.') A defective formation, in which a fold of skin passes from the root of the nose over the inner canthus of the eye. In the plural *epican'thides*—the angles of the eye.

EPICARDIUM, (ἐπί, and καρδιά, 'heart.') See Pericardium.

EPICARPIUM, *Pericar'pium*, (ἐπί, and καρπός, 'the wrist.') An application made to the wrists, or to the region of the pulse, was so called.

EPICAUMA, (ἐπικαυμα, from *epi*, and καίω, 'to burn.') Encauma.

EPICAUSTIS, (ἐπικαυσίς,) Encauma.

EPICERASTICA, (ἐπικεραστικός, 'soothing,' from *epi*, and κεράσνυμι, 'to temper.') Emollients.

EPICHOLOS, (ἐπιχολός, from *epi*, and χολή, 'bile.') Bilius.

EPICHOORDIS, (ἐπιχορδός, from *epi*, and χορδή, 'a gut.') Mesentery.

EPICHORION, (ἐπί, and χορίον, 'skin.') See Decidia, Epidermis.

EPICHORIUM, Epidermis.

EPICHRISIS, (ἐπί, and χρίω, 'to anoint.') Inunctio.

EPICHO'R'SIS, *Spilo'sis*, (ἐπί, and χρωμα, 'colour.') Simple discoloration of the surface.—Ephelides.

EPICHRISIS SPILLI, see Nævus.

EPICHYSIS, (ἐπί, and χέω, 'to pour.') Affusion.

EPICELIDES, (ἐπί, and κελίς, 'the eyelash' [?]) Cilia, see Palpebra (superior.)

EPICØ'MASIS, (ἐπικωμασις, (ἐπί, and κοιμάμαι, 'to lie down to rest,' 'to sleep.') *Decubitus ad dormiendum*, et *Dormititio*. The position of sleeping, as well as sleeping itself.

EPICOL'IC, (ἐπί, and κολών, 'the colon.') The epiploic regions, *regiões epiploicas*, are those parts of the abdomen which correspond to the colon.

EPICOLLESIS, (ἐπί, and κολλά, 'glue.') Agglutination.

EPICON'DYLE, *Epicon'dylus*, (ἐπί, and κοίτυλος, 'a condyle.') A name given by Chaussier, to an eminence at the outer part of the inferior extremity of the os humeri; so called, because it is seated above the condyle. The epicondyle gives attachment to the outer lateral ligament of the elbow joint, and to a very strong tendon to which several of the muscles of the posterior part of the forearm are attached.

EPICONDYLO CUBITALIS, see Anconeus—e. *Radial*—Supinator radii brevis—e. *Sus-métacarpien*, see Radialis—e. *Sus-phalangietien commun*, Extensor digitorum communis—e. *Sus-phalangietien du petit doigt*, Extensor proprius minimi digiti.

EPICOPHO'SIS, (ἐπί, and κωφωσις, 'deafness.') Deafness supervening on another disease.

EPICRA'NIUM, (ἐπί, and κρανιον, 'the cranium.') A name given to different parts seated on the cranium. The skin, aponeurosis between the occipital and frontal muscles, and the whole of the scalp, have been so called. Portal includes under this name the occipito-frontalis muscle, pyramidalis nasi, and superior and anterior auricular muscles. The pericranium.

EPICRANIUS, Occipitofrontalis.

EPICRASIS, *Contempera'tio*, (ἐπικρασις, from *epi*, and κραννυμι, 'I mix,' 'I temper.') A term used by the humourists for an amelioration of the humours. They called *Cure by Epicrasis*—per *epicrasin*, a mode of treatment by imagined alteratives which they supposed to be possessed of the power of gradually correcting the vitiated humours.

EPICROUSIS, (ἐπί, and κρουω, 'I strike.') Percussion.

EPICTENIUM, (ἐπικτενιον, from *epi*, and κτείς, 'pubes.') Pubes.

EPICTETI MORBI, (ἐπικτητος, from *epi*, and κταομαι, 'to acquire.') Acquired diseases.

EPICYEMA, gen. *Epicy'e'matis*, (ἐπικνημα, from *epi*, and κνειν, 'to conceive.') Superfœtation.

EPICYESIS, Superfœtation.

EPICYSTOTOMIA, (ἐπί, and cystotomia.) See Lithotomy.

EPIDEM'IC, *Epidem'icus*, *Epidem'icus*, (F.) *Epidémique*, (ἐπί, and δημος, 'the people.') Appertaining to an epidemic. An epidemic

EPIDEMIC CONSTITUTION, Epidemy—c. Disease, Epidemy—e. Influence, Epidemy.

EPIDÉMICITÉ, see Epidemy.

EPIDEMICUS, Epidemy.

EPIDEMIOGRAPHY, *Epidemiograph'ia*, (epidemy, and γράφω, 'a description.') A history or description of epidemics.

EPIDEMIOLOGY, *Epidemiolog'ia*, (epidemy, and λόγος, 'a description.') The doctrine of epidemics.

EPIDÉMIQUE, Epidemy.

EPIDEMIUS, (ἐπίδημος,) Epidemy.

EPIDEMY, *Epidem'ie*, *Epidém'ia*, *Mor'bus publicus* seu *popula'ris* seu *epidem'icus* seu *epidém'ius*, *Epidém'ie disease*, (F.) *Epidémie*. A disease which attacks at the same time a number of individuals, and which is referred to some particular *constitutio aeris*, or condition of the atmosphere, *E. influence*, *E. constitution*, with which we are utterly ignorant. It differs from *endemic*, the latter being owing to locality; but it is obvious, that should a particular epidemic constitution of the air—*epidemic'ity*, (F.) *épidémicité*—exist along with a favouring endemic condition, these combined influences may act in the causation of several of those serious and fatal complaints, which at times visit a district, and are never afterwards met with, or at least not until after the lapse of a considerable period.

EPIDENDRUM VANILLA, (epi, and δένδρον, 'a tree.') See Vanilla.

EPID'ERIS, (ἐπίδερμος,) *Epider'rhis*, *Epider'rhion*, *Hypod'eris*, *Hypoder'mis*, *Hypoder'rhis*, (epi, and δερμα, 'a skin.') This word, with some, means the nymphæ; with others, the clitoris; with others, again, the prepuce of the clitoris.

EPIDERMA, gen. *Epiderm'atis*, Epidermis.

EPIDERMATIS, Epidermis.

EPIDERMEOUS, Epidemic.

EPIDERMIC, *Epiderm'icus*, *Epiderm'al*, *Epiderm'al'is*, *Epider'meous*, *Epider'midal*, (F.) *Epidermique*. Belonging or relating to the epidermis, —as

EPIDERMIC CELLS, see Cell, epidermic.

EPIDERMIDAL, Epidermic.

EPIDERMIDON'OSI, *Epidermidon'osi*, (epidermis, and νόσος, 'disease.') Diseases of the epidermis.

EPIDERMION, Epidermis.

EPIDER'MIS, gen. *Epider'midis*, (ἐπίδερμος, gen. ἐπίδερμίδος,) *Epider'mion*, *Epider'matis*, *Epider'ma*, *Epicho'ron*, *Epicho'rium*, *Sum'matis cut'is*, *Cutic'ula*, *C. ex'tima* seu *extre'ma*, *Cut'is ex'tima* seu *sum'ma* seu *ul'tima*, *Pel'lis sum'ma*, *Pellic'ula sum'ma* seu *superior*, *Lam'ina pri'ma cut'is*, *Oper'culum cut'is*, (epi, and δερμα, 'the skin.') *Scarf'skin*, *Cut'icle*. (F.) *Surpeau*, *Cuticule*. A transparent, dry, thin membrane, devoid of nerves and vessels, which covers all the surface of the body, except the parts that correspond to the nails. It appears to consist of minute scales, placed one above the other. Chaussier considers it to be formed and reproduced by an excretory action of the true skin; to act like a dry varnish—*Stratum Cor'neum*—which prevents the immediate contact of bodies with the nervous papillæ, and consequently to deaden tactile impressions, which, without its intervention, might be painful. The *Epider'mic*, *Epider'meous*, or *Epider'moid System*, (as Bichat called it) in general anatomy, comprises three parts. 1. External Epidermis. 2. Epidermis spread over the mucous membranes. 3. The Nails and Hair.

EPIDERMIS LINGUÆ, see Tongue.

EPIDERMOID, *Epidermo'id'es*, (epidermis, and εἶδος, 'resemblance.') Resembling the epidermis.

EPIDERRHION, (epi, and δερμός, 'skin.') Epiteris.

EPIDERRHIS, (ἐπίδερμης,) Clitoris, Epideris.

EPIDERRHITIS, (epiderrhis, and ἰtis.) Clitoritis, Nymphitis.

EPIDESIS, (ἐπίδεσις, from epi, and δέω, 'to bind.') Deligation, see Bandage—c. *Hæmostasia*, Ligature.

EPIDESMIS, Epidesmus.

EPIDESMUM, Epidesmus.

EPIDESMUS, (ἐπίδεσμος,) *Epides'mis*, *Epides'mum*, *Superliga'men*. A fascia, bandage or ligature, by which dressings are secured.

EPIDIDY'MIS, gen. *Epididym'idis*, (ἐπίδιδυμος, gen. ἐπίδιδυμδος, from epi, and δίδυμος, 'a testicle.') *Ep'ididymus*, *Did'yms*, *Paras'tata*, *Testic'ulus accesso'rius*, *Cap'ut test'is*, *Cor'pus varico'sum* seu *varicifor'mē test'is*, *Supergenina'lis*, (F.) *Corpus variqueux* ou *variciforme*. That which is placed upon the testicle. A small, oblong, vermiform, grayish body, lying along the superior margin of the testicle. The Epididymis is a canal formed by the union of all the seminiferous vessels folded several times upon themselves, after having traversed the *Corpus Highmorianum*. Its lower portion or tail, *Cauda* seu *globus* seu *caput minor*, is curved upwards, and is continuous with the vas deferens; the opposite extremity is called the head, *globus* seu *caput major*. The length of this canal, folded as it is upon itself and describing numerous curvatures, is, according to Munro, 32 feet.

EPIDIDYMITIS, (epididymis, and ἰtis.) Parastatitis—c. Blennorrhagic, Hernia humoralis.

EPID'OSIS, (ἐπίδωσις, from epi, and δίδωμι, 'to give.') Augmentation, increase. A word applied to the natural increase of the body, or to the increase of a disease.

EPID'ROME, (ἐπίδρομη,) *Epidrom'ia*, *Epiph'ora*, (epi, and δρεπω, 'I run.') An afflux or congestion of humours.

EPIDROMIA, (ἐπίδρομια,) Epidrome.

EPIFAGUS AMERICANUS, (epi, and fagus, 'the beech.') *Orobanche Virginiana*—c. *Virginianus*, *Orobanche Virginiana*.

EPIGÆ'A, EPIGE'A or EPIG'I'A RE'PENS, (επιγεια, from epi, and γη, 'the earth,' from its trailing growth.) *Trailing Arbutus*, *Ground Lau'el*, *May'flower*, called, by the Shakers, *Grav'elpant*. A small, trailing plant, of the Family Ericacæ, which grows in sandy woods, or in rocky soil, especially in the shade of pines. Its flowers appear in early spring, and exhale a rich, spicy fragrance. The leaves and stems are prepared and used like uva ursi.

EPIGASTRAL'GIA, (epigastrium, and αλγος, 'pain.') Pain at the epigastrium.

EPIGAS'TRIC, *Epigas'tricus*, *Epigas'trius*, (epi, and γαστρν, 'the stomach.') This name has been given to several parts.

EPIGASTRIC AR'TERY, *Ar'teria epigas'trica*, *Ar'tère sus-pubienne*, (Ch.) It arises from the *external iliac*, on a level with the crural arch; sometimes by a trunk proper to it, at others by one common to it and the *Obturator artery*. It ascends upwards and inwards, behind the spermatic cord, follows the outer edge of the rectus abdominis muscle, and anastomoses, towards the umbilicus, with the internal mammary. The epigastric artery anastomoses, also, with the obturator, spermatic, lumbar, and intercostal vessels.

EPIGASTRIC REGION, *Re'gio epigas'trica* seu *cardi'aca* seu *stomach'ica* seu *stomacha'lis*, is the superior region of the abdomen, comprised between the false ribs on each side, and extending from the lower surface of the diaphragm to two fingers' breadth above the umbilicus. It is divided into three parts, one *middle*, the *epigas'trium*,—and two *lateral*, the *hypochon'dria*.

EPIGASTRIC VEIN follows nearly the same course as the artery.

EPIGASTRICUS, Epigastric.

EPIGASTRIUM, (ἐπιγαστήριον, same etymon as

Epigastric.) The belly; the epigastric region.

EPIGASTRIUS, (ἐπιγαστήριος,) Epigastric.

EPIGASTROCELE, *Epigastrocele*, (ἐπι, γαστήρ, 'the stomach,' and κήλη, 'a tumour.') Hernia, formed by the stomach; *Gastrocele*. This name has been given more especially to hernia, which occurs towards the upper part of the linea alba or in the epigastric region, whether formed or not by the stomach.

EPIGEA, Epigæa.

EPIGENEMA, gen. *Epigene'matis*, (ἐπιγενημα,) Epigenesis.

EPIGENESIS, (ἐπι, and γένεσις, 'generation.') A theory of conception, according to which the new being is created entirely anew; and receives at once from each parent the materials necessary for its formation. Also, a new formation.

EPIGENESIS, as well as *Epigene'ma* or *Epigene'ma*, is, also, applied to any symptom occurring during a disease, without changing its nature. An accessory symptom. *Epiginom'enos*, *Superve'nientis*, has a similar acceptance.

EPIGINOMENOS, (ἐπιγινόμενος, from ἐπι, and γίνομαι, 'to happen,') see Epigenesis.

EPIGLOTTIC, *Epiglotticus*, (ἐπι, and γλωττις, 'the glottis,') Relating to the epiglottis.

EPIGLOTTIC GLAND, *Periglottis*, *Ca'ro glandulo'sa*. A collection of small, glandular granula, situate in the adipose, areolar texture at the base of the anterior surface of the epiglottis, in a triangular space, bounded anteriorly by the thyroïd membrane and thyroid cartilage. It pours out an unctuous and mucous fluid, which lubricates the epiglottis, keeps it supple and movable, and prevents the larynx from being irritated by the constant passage of the air, in the act of respiration.

EPIGLOTTIS, gen. *Epiglott'idis*, (ἐπιγλωττις, gen. ἐπιγλωττιδος.) *Epiglottic cartilage*, *Lig'ula Oper'culum laryn'gis*, *Lin'gua Ex'igua*, *Lin'gua fist'ula*, *Subling'uum*, *Superlig'ula*. A fibro-cartilage, situate at the upper part of the larynx, behind the base of the tongue. Its form is oval; texture elastic; thickness greater below than above, and greater in the middle than at the sides. By its smaller extremity, which is the lower, it is attached to the thyroid cartilage; its two surfaces are covered by the mucous membrane of the pharynx and larynx. The special use of the epiglottis would seem to be to cover the glottis accurately at the moment of deglutition, and, thus, to assist in opposing the passage of alimentary substances into the air tubes.

EPIGLOTTITIS, (epiglottis, and itis,) *Inflam'matio Epiglott'idis*, *Angi'na epiglottide'a*, (F.) *Inflammation de l'Epiglote*. Inflammation of the epiglottis.

EPIGLOTTUM. An instrument, mentioned by Paracelsus, for raising the eyelids. It resembled the epiglottis in shape.

EPIGLOUTIS, (ἐπι, and γλoutος, 'the buttocks,') The superior region of the nates.

EPIGNATHUS, (ἐπι, and γnaθος, 'the jaw,') A double monster, in which an incomplete fœtus is rooted, with its blood vessels in the palate of one more complete.

EPIGONATIS, (ἐπιγονατις, from ἐπι, and γονυ, 'the knee,') Patella.

EPIGONĒ, (ἐπιγονή,) *Go'nē*, *Hypoph'yesis*, *Pro-gen'ies*, *Pro'les*, *Sob'oles*, *Sub'oles*, (ἐπι, and γονη, 'seed,') Offspring. Progeny. Also, Superfœtation.

EPIGONION, Fœtus.

EPIGONIS, (ἐπι, and γονυ, 'the knee,') Patella.

EPIGONON, (ἐπι, and γονος, 'progeny,') Superfœtation.

EPIGOUNIS, (ἐπιγουνις, from ἐπι, and γουνος, 'the knee,') Patella.

EPIGUNIS, Patella.

EPIILATION, (ἐ, and pilus, 'a hair,') Depilation.

ÉPILATOIRE, Depilatory.

EPILATORIUM, Depilatory.

EPILEPSIS, Epilepsy.

EPILENTIA, Epilepsy.

EPILEPSIA, Epilepsy—e. *Acuta Infantum*, *Eclampsia*—e. *Algetica*, see *Algos*—e. *Febrilis infantum*, *Eclampsia*—e. *Nocturna*, *Incubus*—e. *Puerilis*, *Eclampsia infantum*—e. *Saltatoria*, *Chorea*—e. *Uterina*, *Lorind matricis*.

ÉPILEPSIE, Epilepsy—e. *Utérine*, *Hysteria*—e. *Vertige*, see *Epilepsia*.

EPILEPSY, (ἐπιληψία, ἀπὸ πόνου, σελήναια νοσους,) *Epilep'sia*, *Epilep'sia*, *Epilemp'sis*, *Epilen'tia*, *Syspa'sia Epilep'sia*, *Catalen'tia*, *Deciden'tia*, *Pas'sio cadu'ca*, *Heracle'us mor'bus*, *Hieran'onus*, *Deprehen'sio*, *Olo'dius epilep'sia*, *Prehen'sio*, *Perdit'io*, *Epilep'tica pas'sio*, *Mor'bus comit'ialis* seu *cadu'cus* seu *Hercule'us* seu *ad'cer seu astra'lis* seu *sidera'tus* seu *dæmoni'acus* seu *Sanct'i Johan'nis* seu *fœ'dus* seu *puer'ilis* seu *mag'nus* seu *ma'jor* seu *interlu'nus* seu *divi'nus* seu *dæmo'nus* seu *zon'ticus* seu *seleni'acus* seu *lunat'icus* seu *mensa'lis* seu *deif'icus* seu *ecles'us* seu *vitriola'tus* seu *viridel'us* seu *Sanct'i Valenti'ni*, *Ma'tum cadu'cum*, *Analep'sia*, *Catapto'sis*, *Insan'ia cadu'ca*, *Apoplex'ia par'ca*, *Cadu'ca pas'sio*, *Lu'es divi'na* seu *deif'ica*, *Vit'ium cadu'cum* seu *divi'nium* seu *Hercule'um*, *Fall'ing sick'ness*, *Fall'ing-down*, (Old E.) *Fallandevyl*, (F.) *Mal caduc* ou *divin* ou *Saint-Jean* ou *de Terre*, *Haut mal*, *Goutte caduque*. (The word is derived from ἐπιλαμβάνω, 'future, ἐπιληψω, (ep, and λαβανα, 'I seize,') 'I seize upon,') It is a cerebrospinal disease which may be idiopathic or symptomatic, spontaneous or accidental, and which occurs in paroxysms, with uncertain intervals between. These paroxysms are characterized by loss of consciousness and by convulsive motions of the muscles. Frequently, the fit attacks suddenly; at other times, it is preceded by indisposition, vertigo, and stupor. At times, before the loss of consciousness occurs, a sensation of a cold vapour is felt, hence called *aur'a epilep'tica*. This appears to rise in some part of the body, proceeds towards the head; and as soon as it has reached the brain the patient falls down. The ordinary duration of a fit is from 5 to 20 minutes. Sometimes it goes off in a few seconds; at others, it is protracted for hours. In all cases, there is a loss of sensation, sudden falling down, distortion of the eyes and face; countenance of a red, purple, or violet colour; grinding of the teeth; foaming at the mouth; convulsions of the limbs; difficult respiration, at times stertorous; with, sometimes, involuntary discharge of fæces and urine. After the fit, the patient retains not the least recollection of what has passed, but remains, for some time, affected with headache, stupor, and lassitude.

The disease is cerebrospinal, and is generally organic; but it may be functional and symptomatic of irritation in other parts, as in the stomach, bowels, &c. The prognosis, as to ultimate recovery, is unfavourable. It does not, however, frequently destroy life, but is apt to lead to mental imbecility. Dissection has not thrown light on its pathology.

To the attacks of epilepsy which are unaccompanied by convulsions, as is sometimes the case, the French give the name *Petit-mal*, and *Epilepsie Vertige*, *Cerebral Epilepsy*. Fully formed epilepsy is the *Grand-mal* of the French, *Spi'nal Epilepsy*. When furious mania succeeds to a

paroxysm, it is termed *Ma'nia epilep'tica* and *Epilep'tic Delir'ium*.

In the treatment, the cause must be sought after, and if possible removed. In the paroxysm, but little can be done; but as the tongue is liable to be injured by the teeth, the jaws may be kept open by putting a cork or piece of wood between them. If the fit has been brought on by indigestible food, the stomach must be cleared. It is between the paroxysms that the great effort must be made. Generally, there is considerable irritability and debility of the nervous system, and hence tonics have been found the best remedies. Of these, perhaps, the most powerful, in epilepsy, is the *argenti nitras*, given regularly and continued for months, if necessary. Preparations of iron, copper, and zinc, have also been used, and vegetable tonics and antispasmodics in general. Counter irritants, as blisters, moxa, &c., may be employed, if necessary, along with this course.

Unfortunately, in many cases, these means are found insufficient, and all that can be done is to palliate, removing carefully the exciting causes; such as the use of spirituous liquors, strong emotions, violent exercise, &c.; and regulating the diet.

EPILEPSY, CEREBRAL, see Epilepsy—e. Spinal, see Epilepsy.

EPILEP'TIC, (*επειληπτικός*.) *Epilep'ticus*, *Epilep'tus*, *Lunat'icus*, *Caduc'eans*. One affected with epilepsy. Any thing relating to epilepsy; as an *epilep'tic individ'ual*; an *epilep'tic symp'tom*. Some authors also speak of *epilep'tic remedies*.

EPILEPTIFORM, Epileptoid.

EPILEP'TOID, *Epileptoi'des*, *Epilep'tiform*, *Epileptiform'is*, (*epilepsy*, and *eidos*, 'resemblance.') Resembling epilepsy—as '*epileptoid symptoms*.'

ÉPILEUR (F.). One employed in the French hospitals to practise epilation in affections involving the hair follicles.

EPILO'BIIUM ANGUSTIFOLIUM, (*epi*, and *λοβος*, 'a pod,' the flowers seeming as if on a pod.) *Great Willow herb*. An indigenous plant, *Order*, *Onagraceae*, which flowers in July. The root is emollient and slightly astringent.

EPILOBIUM COLORATUM, *Purple-veined Willow herb*, has similar properties.

EPILO'BIIUM VILLOSUM, a South African plant, is used at the Cape of Good Hope as a domestic remedy for cleansing foul ulcers.

EPIM'ANES, (*επιμανής*, from *epi*, and *μαινωμαι*, 'I am in a rage at.') A name given to a maniac when in a paroxysm.

EPIMELE'TAI, (*επιμελητής*, 'a care taker,' from *epi*, and *μελω*, 'I take care.') A name by which the ancient Persians called the *curers of wounds*, who followed their armies.

EPIMOR'IOS, (*επιμοριος*, 'unequal,' from *epi*, and *μειρω*, 'I divide.') An epithet applied to the pulse when unequal.

ÉPINARD, ([Old F.] *epinard*, from *épine*, (L.) *spina*, 'a thorn or prickle,') *Spinacia*—*é. Sauvage*, *Chenopodium bonus Henricus*.

ÉPINE, ([Old F.] *épine*,) *Spine*—*é. de Cerf*, *Rhamnus*—*é. du Dos*, *Vertebral column*—*é. Cutanéale*, *Nasal spine*, *inferior and posterior*—*e. Sphénoïdale*, *Sphenoidal spine*—*é. Vinette*, *Oxyacantha Galeni*.

EPINEMESIS, (*επινημσις*, from *epi*, and *νημι*, 'to divide,') *Dispensation*.

EPINEPH'ELOS, (*επινεφελος*, from *epi*, and *νεβηλη*, 'a cloud,') *Nubilo'us*. Presenting a cloud. Applied to the urine.

ÉPINEUX, *Spinous*—*é. du Dos, grand*, *Spinialis dorsi major*: see *Interspinales dorsi et lumborum*—*é. du Dos, petit*, see *Interspinales dorsi et lumborum*.

ÉPINGLE, ([Old F. *épingle*; from (L.) *spinula*, dim. of *spina*, 'a thorn,') *Pin*.

ÉPINIÈRE, ([Old F.] *epinière*.) *Spiral*.

EPINOTION, (*επινώτιον*, from *epi*, and *νωτος*, 'the back,') *Scapula*.

EPINYCTIDES, pl. of *epinyctis*, (*επινυκτις*, gen. *επινυκτιδος*, from *epi*, and *νυξ*, gen. *νυκτος*, 'the night,') *Eruptions which appear on the skin in the night, and disappear in the day. Pus'tula livens et noctibus inq'uiescens*.

EPINYCTIS PRURIGINOSA, *Urticaria*.

ÉPIONE, (*epi*, and *ων*, 'ovum,') *Decidua*.

EPIPAROXYSMUS. A paroxysm superadded to an ordinary paroxysm.

EPIPASTUM, (*επιπαστον*, from *epi*, and *πασσι*, 'to sprinkle,') *Cataplasma*.

EPIPE'CHU, (*επιπεχον*, from *epi*, and *πηχυς*, 'the elbow,') The upper part of the elbow. The part above the elbow.

EPIPHÆNOMENON, *Post appa'rens*, *Epiphenomenon*; (same etymon as the next.) Any kind of adventitious symptom which occurs during the progress of an affection; and which is not as intimately connected with the existence of the disease as the symptoms properly so called.

EPIPHANI'A, (*επιφανεια*, from *epi*, and *φανω*, 'I appear,') The external appearance of the body.

EPIPHYGUS VIRGINIANA, (*epi*, and *φηγος*, 'the beech,') *Orobanche Virginiana*.

EPIPHENOMENON, *Epiphenomenon*.

EPIPH'LEBUS, (*επιφλεβος*, from *epi*, and *φλεβ*, gen. *φλεβος*, 'a vein,') An epithet given to those whose veins are very apparent.

EPIPHLEGIA, (*epi*, and *φλεγω*, 'I inflame,') *Inflammation*.

EPIPHLOGISMA, *Erysipelatous inflammation*.

EPIPHLOGOSIS, *Hyperphlogosis*.

EPIPH'ORA, (*επιφωρα*, from *epi*, and *φωρω*, 'I carry,') *Dacryoma*, *Lacrymat'io*, *Delacrymat'io*, *Ilacrymat'io*, *Stillicid'ium lacryma'rum*, *Oculus lacrymans*, *Weeping*. (F.) *Larmoiement*. The *watery eye*;—an involuntary and constant flow of tears upon the cheek. It is almost always symptomatic of some disease of the lachrymal passages, and occurs when the tears cannot pass into the *ductus ad na'sum*. Occasionally, it is owing to the tears being secreted in too great quantity; as in certain cases of ophthalmia. The treatment must of course be directed to the removal of the cause.

EPIPHORA, *Epidrome*, *Ophthalmia*—e. *Alvi*, *Diarrhoea*—e. *Ptyalismus*, *Salivation*.

EPIPHYMATA, (*epi*, and *φωω*, 'to grow,') *Cutaneous diseases*.

EPIPH'YSIS, (*επιφωσις*, same etymon.) *Additamentum*. Any portion of a bone, separated from the body of the bone by a cartilage, which becomes converted into bone by age. The *epiph'ysis* of the fœtus becomes the *apoph'ysis* of the adult.

EPIPHYSIS CRURIS LONGIORIS INCUDIS, *Os orbiculare*.

EP'IPHYTE, *Epiph'yton*, pl. *Epiph'ytæ*, (*epi*, and *φυτον*, 'a plant,') *Phy'toparasite*. A parasite derived from the vegetable kingdom. Epiphytes are divided by Vogel into, 1. Vegetations in the human fluids, as the *tor'ula cerevisiæ* in vomited fluids, and fecal evacuations, and the *sar'cina ventriculi*. 2. Vegetations on the external skin and its appendages, as in *tinca favosa*, *mentagra*, *herpes tonsurans* and *plica polonica*; and, 3. Vegetations on the mucous membrane,—as in the aphthæ of children; in the cicatrices of the mucous membrane after typhus, &c. By some, the term *epiphytes* is restricted to those vegetable parasites which grow on the surface of the body.

EPIPLAS'MA, gen. *Epiplasm'at'is*, (*επιπλάσμα*, from *επιπλασσω*, (*epi*, and *πλάσσω*, 'I smear,') 'I plaster,') *Cataplasma*. Galen uses it especially for an application of wheaten flour, boiled in *hydrelæum*, and applied to wounds.

EPIPLEGIA, (*ἐπι*, and *πληγή*, 'a stroke.') Paralysis of the upper extremities.

EPIPLEROSIS, (*ἐπι*, and *πληρωσις*, 'repletion.') Excessive repletion; distension.

EPIPLO, from *επιπλουν*, (*ἐπι*, and *πλεω*, 'I swim or float,') in composition, 'epiploon.'

EPIPLOCE, (*επιπλοκή*, from *ἐπι*, and *πλεω*, 'I twine,') Coition. Entanglement.

EPIPLOCE INTESTINALIS, *Conglomeratio intestinum*. Conglomeration of the intestines.

EPIPLOCELĒ, *Epiplocele*, (*epiplo*, and *κηλη*, 'tumour,') *Epiploic hernia*, *Zirba'lis hernia*, *Hernia omentalis*, *Omental hernia*, (F.) *Hernie de l'épiploon*. Omental hernia is recognised—but at times with difficulty—by a soft, unequal, pasty tumour, in which no gurgling sound is heard on pressing or reducing it. It is less dangerous than hernia of the intestines.

EPIPLOENTEROCELĒ, *Epiploenterocele*, (*epiplo*, *εντερον*, 'intestine,' and *κηλη*, 'a tumour,') Hernia, formed by the epiploon and a portion of the intestine. It is more commonly called *Enteroëpiplocele*.

EPIPLOENTEROSCHEOCELĒ, *Epiploenteroschecocele*, (*epiplo*, *εντερον*, 'intestine,' *σχχιον*, 'scrotum,' and *κηλη*, 'a tumour,') Hernia with omentum and intestine in the scrotum.

EPIPLOIC, *Epiploicus*. Relating to the epiploon.

EPIPIOIC APPENDAGES, *Appendiculæ pinguedinosæ*.

EPIPIOIC ARTERIES; *Arteriæ Epiploicæ*. The arterial branches, which are distributed to the epiploon, and which are given off by the gastroepiploicæ.

EPIPIOISCHIOCELĒ, *Epiploischiocele*, (*epiplo*, *ισχιον*, 'the ischium,' and *κηλη*, 'a tumour,') Hernia formed by the epiploon through the ischiatic notch.

EPIPIOÏTIS, (*epiplo*, and *itis*,) Inflammation of the omentum; *Omentitis*, *Empres'na peritonitis omentalis*, *Omentitis inflammatoria*, *Enteritis epiploitis*, *Omentosis*. A form of partial peritonitis.

EPIPLOMEROCELĒ, *Epiplomerocele*, (*epiplo*, *μηρος*, 'the thigh,' and *κηλη*, 'a tumour,') Femoral hernia, formed by the epiploon.

EPIPLOMPHALOCELE, (*κηλη*, 'a tumour,') Epiplophalon.

EPIPLOMPHALON, *Epiploömp'halon*, *Epiplophalocelē*, *Hernia umbilicalis*, (*epiplo*, and *ομφαλος*, 'the navel,') Umbilical hernia, formed by the epiploon.

EPIPLOÖCOMISTES, (*epiplo*, and *κομίζω*, 'I carry,') One who has the omentum morbidly large. Also, one labouring under epiplocele.

EPIPLOÖN, (*επιπλουν*,) *Epiploom*, (*epi*, and *πλεω*, 'I swim or float,') *Omentum*, *Re'te, Reti-culum*, *Der'tron*, *Zir'bus*, *Gan'gamē*, *Gan'gamum*, *Omentum intestinum*. *Sac'cus epiploicus*, *Sage'na*, *The Gaul*, (Prov.) *Flick*, *Web*. A prolongation of the peritoneum, which floats above a portion of the intestines, and is formed of two membranous layers, with vessels and fatty bands distributed through it. Anatomists have described several epiploons. The uses of the omentum are but little known. The chief one seems to be, to retain the viscera *in situ*, and to give passage to vessels.

EPIPLOON, *Col'ic*, *Col'ic Omentum*, *Omentum colicum*, *Third Epiploon* or *Omentum*, is a duplication of the peritoneum, situate along the ascending portion of the colon, as far as its junction with the transverse portion. It is behind the great epiploon, and fills up the angle formed by the junction of the ascending with the trans-

verse colon. Its two laminae are separated by colic arteries and veins.

EPIPLOON, *GASTROCOL'IC*, *Great Omentum*, *Omentum gastrocolicum* seu *ma'jus*, *Peritonæum duplicatum*, *Re'te ma'jus*, *Zir'bus adipi'mus*, consists of an extensive duplicature, which is free and floating on the folds of the intestines. It is quadrilateral, and longer on the left side than on the right:—its base being fixed, anteriorly, to the great curvature of the stomach; and, posteriorly, to the arch of the colon. It is formed of two laminae, each consisting of two others. In this epiploon a number of vessels is distributed, and there is much fat.

EPIPLOON, *GASTROEPAT'IC*, *Lesser Omentum*, *Omentum hep'atogast'ricum* seu *mi'nus*, *Membra'na macilentior*, is a duplicature of the peritoneum which extends transversely from the right side of the cardia to the corresponding extremity of the fissure of the liver, and downwards from this fissure to the lesser curvature of the stomach, the pylorus and duodenum. Below it is the *foramen* or *hiatus* of *WINSLOW*; and between its laminae are lodged the biliary and hepatic vessels. It contains but little fat. If air be blown in at the foramen of Winslow, the cavity or sac of the omentum, *Sac'cus Peritonæi retroventricula'ris*, *Bur'sa omentalis*, will be rendered perceptible.

EPIPLOON, *GASTROSPLE'NIC*, *Omentum Gastrople'nicum*. A reflection of the peritoneum, which passes between the concave surface of the spleen and the stomach, from the cardiac orifice to near its great curvature, and which lodges the vasa brevia and splenic vessels between its laminae.

EPIPILOSARCOMPHALUS, (*epiplo*, *σαρξ*, gen. *σαρκος*, 'flesh,' and *ομφαλος*, 'navel,') Umbilical hernia, formed of indurated omentum.

EPIPILOSCHEOCELĒ, *Epiploschecocele*, *Epiploschocelē*, (*epiplo*, *σχχιον*, 'the scrotum,' and *κηλη*, 'tumour,') Scrotal hernia, formed by a displacement of the epiploon.

EPIPILOUM, Epiploon.

EPIPNOIA, (*επιπνοια*, from *epi*, and *πνεω*, 'I breathe,') *Aflatus*.

EPIPOROMA, gen. *Epiporomatis*, (*επιπρωω*, (*epi*, and *πρωω*,) 'I harden,') A tophaceous concretion which forms in the joints. The callus of fracture.

EPIRRHOĒ, (*επιρρῶν*,) *Epirrhe'a*, (*epi*, and *ῥεω*, 'I flow,') Afflux or congestion of humours.

EPIRRHOEA, (*επιρροια*,) *Epirrhe*.

EPISARCIDIUM, (*epi*, and *σαρξ*, 'flesh,') *Anasarca*.

EPISCHESIS, (*επισχισις*, from *επισχω*, 'I restrain,') Retention. A suppression of excretions.

EPISCHET'ICUS, (*επισχετικός*,) *Sistens*, *Re'primens*, *Co'hidens*; 'restraining.' An agent that diminishes—secretion, for example.

EPISCHION, (*epi*, and *ισχιον*, 'ischium,') Abdomen, Pubes, Pubis os.

EPISCOPALES VALVULÆ, (from *episcopus*, 'a bishop,') Mitral valves.

EPISEION, (*επισειον*,) Pubis (os) Vulva, Labium Pudendi, Tressoria.

EPISEMA'SIA, (*επισημασια*, from *επισημαινω*, (*epi*, and *σημα*, 'a sign,') 'I afford some sign,') *Sema'sia*. It has been used synonymously with *sign*, and also with the invasion of an attack of fever. See *Annotatio*.

EPISI'O, (*επισειον*, (modern) the labium pudendi,) in composition, labium pudendi.

EPISEOCELE, (*episto*, and *κηλη*, 'rupture,') see *Hernia*, vulvar.

EPISCEDE'MA, (*episio*, and *οιδημα*, 'oedema,') Oedema of the labia pudendi.

EPISEIOÏTIS, (*episio*, and *itis*,) Inflammation of the labia pudendi.

EPISION, (*επισειον*,) *Episeion*.

EPISION'CUS, *Episiophy'ma*, (*episio*, and *γκος*, 'a tumour.') *Tumor labio'rini*. A swelling or tumour of the labia pudendi.

EPISIOPHYMA, gen. *Episiophy'matis*, (*episio*, and *φυμα*, 'a swelling,') Episioncus.

EPISTORRHAG'IA, (*episio*, and *ρρηγ*, 'rupture,') Hemorrhage from the labia pudendi.

EPISTIOR'RHAPHY, *Epistiorrhaph'ia*, (from *episeion*, and *ρρηγ*, 'suture,') An operation practised in cases of prolapsus uteri, which consists in paring the opposing surfaces of the labia pudendi, bringing them together and uniting them by suture, so as to diminish the outlet of the vulva.

EPISPA'DIA, (*epi*, and *σπaw*, 'I draw,') *Hyperasp'adia*. The condition of an Epispadias.

EPISPA'DIAS, *Epispadias'us*, *Anuspa'dias*, *Anuspa'di'us*, *Hyperasp'adias*, *Hyperaspadi'us*, *Episp'adian*. The opposite to Hypospadias. One who has a preternatural opening of the urethra at the upper part of the penis.

EPISPAST'ICS, (*επισπαστικά*,) *Epispas'tica*, *Helc'tica*, (F.) *Épispastiques*; (same etymon as *Episp'adia*.) An epithet for every medicinal substance which, when applied to the skin, excites pain, heat, and more or less redness, followed by separation of the epidermis,—which is raised up by effused serum,—or by suppuration. Now usually restricted to blisters.

EPISPASTICUM, pl. *Epispastica*. Blister.

ÉPISPASTIQUES, Epispastics.

EPISPH'E'RIA, (*epi*, and *σφαيرا*, 'a sphere,') Some physiologists have applied this term to the different convolutions and sinuosities presented by the external surface of the brain.

EPISPLENTITIS, see Splenitis.

EPISTACTIS'CHESIS, (*epistaxis*, and *ισχειν*, 'to restrain,') Arrest of bleeding from the nose.

EPISTAPHYLINI, (*epi*, and *σταφυλη*, 'the uvula,') see *Azygos Muscæ*.

EPISTASIS, *Insident'ia*, (*επιστασις*, from *epi*, and *σπaw*, 'I rest,') A substance which swims at the surface of urine. It is opposed to the *hypostasis* or sediment. Epistasis has also been employed synonymously with suppression.

EPISTATION (F.), *Epistatio*, *Pistat'ion*, *Pistat'io*, (*e*, and *pistare*, 'to pound,') consists in bruising in a mortar soft and parenchymatous substances.

EPISTAX'IS, (*epi*, and *σταζω*, 'I flow drop by drop,') *Hæmorrhag'ia acti'ca na'rini*, *H. nari'ca* seu *na'rini*, *Epistax'is jmi'o'rini* seu *arte'ri'ca*, *Hæmorrhag'ia*, *Hæmorrhino'rha'g'ia*, *Hæmat'imus*, *Rhinorrhag'ia*, *Stillicid'ium* seu *Stillat'io Sang'ui'is na'r'ibus*, *Hæmorrhage* from the pituitary men'brane, *Bleed'ing* at the nose, *Nose'bleed*, (F.) *Hémorrhagie nasale*, *Saignement du nez*, *Hémorrhinie*. This is one of the most common varieties of hemorrhage; the organization of the Schneiderian membrane being favourable to it, as the blood-vessels are but slightly supported. It does not generally flow from both nostrils, and is very apt to recur. Puberty is the period at which it is most common. *Its exciting causes* are:—any thing that will induce local congestions, as running, coughing, blowing the nose, &c., provided there is a disposition to it. A common case of epistaxis requires but little treatment, especially if in a young person. Light diet and a dose or two of the sulphate of magnesia will be sufficient. In more severe attacks, cold and astringent washes of alum, sulphate of zinc, weak sulphuric acid, or creasote, may be used, and the nostrils be plugged anteriorly; but if the bleeding goes on posteriorly, —*Chæmorrhag'ia*—the posterior nares must be plugged likewise,—the patient being kept with the head raised, and on dry diet. The flow of blood has been arrested by directing the patient to stand up with his head elevated, compressing

the nostril whence the blood flows with the finger, raising the corresponding arm perpendicularly, and holding it in this position for about two minutes. A less vigorous circulation through the carotids appears to result from the increased force required to carry on the circulation through the upper extremity when raised. To prevent the recurrence, strict diet must be inculcated.

EPISTER'NAL, *Episterna'lis*, (*επι*, and *στερνον*, 'the sternum,') An epithet applied to two bones which form part of the sternum, and are situate upon its superior and lateral part. In the young subject, they are attached to the sternum by a synovial membrane, and a fibrous capsule. They have somewhat the appearance of the pisiform bones, but are of a greater size.

According to Professor Owen, the piece of a segment of an articulate animal, which is immediately above the middle inferior piece or sternum.

EPISTHOT'ONUS, (*επισθεν*, 'forwards, and *τενω*, 'I extend,') A spasm of the muscles, drawing the body forwards. The word *emprosthotonus* is more commonly employed.

EPISTROPHE, (*επιστροφή*, from *epi*, and *στρεφω*, 'I turn,') Relapse.

EPISTROPHEUS, (*επιστροφεις*,) Axis.

EPISTROPHUS, (*επιστροφος*,) Axis.

EPISYNAN'CHE, (*epi*, and *synanche* or *cynanche*,) *Episyngangi'na*. A Greek word, used by a modern writer for spasm of the pharynx, by which deglutition is prevented, and the solid and liquid food driven back towards the mouth or nasal fossæ.

EPISYNTHET'IC, *Episynthet'icus*, (*επισυνθεσις*, (*επι*, *συν*, and *τιθημι*, 'to collect, accumulate,') The name of a medical sect, whose object it was to reconcile the principles of the Methodists with those of the Empirics and Dogmatics. Leonides of Alexandria seems to have been one of the first partisans of this sect; of which we know little.

EPIT'ASIS, (*επιτασις*, from *epi*, and *τεινω*, 'to extend,') The period of violence of a fever, (*Feb'ris inten'sio*,) attack or paroxysm; sometimes it is used in the same sense as epistasis, for 'suppression.'

EPITHELIUM, Epithelium.

EPITHE'LIAL, *Epithelia'lis*, (*epi*, and *θηλη*, 'a nipple,') Appertaining or relating to the epithelium,—as epithelial cells, and scales.

EPITHELIAL CAN'CE'R, *Canc'roid* of the skin, *Epithelio'ma*, *Carcino'ma epithelio'des*, is a morbid growth of epithelial cells; forming an *epithelial* tumour, which, by some, is regarded as *carcinoma*. See *Canceroid*.

EPITHELIOMA, gen. *Epithelio'matis*, (*epithelium*, and *oma*,) Epithelial cancer.

EPITHE'LIUM, *Epithelium*, *Epithe'lis*, *Epithe'lia*, *Endepiderm'is*. The thin layer of epidermis, which covers parts deprived of *derma* properly so called,—as the nipple, mucous membranes, lips, &c. Modern histological researches have shown that it exists, in different forms, *pavement*, *cyl'inder*, and *vi'brate* or *cil'iated* *epithelium*. *Tes'selated*, *pavement*, *scaly*, *lam'ellar*, *tubular*, or *flat'tened* *epithelium*, *Epithelium polyed'ricum*, and *E. lamello'sum*, (F.) *Épithélium pavimenteur*, covers the serous and synovial membranes, the lining of blood-vessels, and the mucous membranes, except where cylinder epithelium exists. It is spread over the mouth, pharynx and œsophagus, conjunctiva, vagina, and entrance of the female urethra. *Cyl'inder*, *colum'nar* or *prismat'ic* *epithelium*, *Epitheliu' cylin'dricum*, (F.)

Épithélium cylindrique is found in the intestinal canal, beyond the cardiac orifice of the stomach; in the larger ducts of the salivary glands; in the ductus communis choledochus, prostate, Cowper's glands; vesiculæ seminales, vas deferens, tubuli uriniferi, and urethra of the male; and lines the

urinary passages of the female, from the orifice of the urethra to the beginning of the tubuli uriniferi of the kidneys. In all these situations, it is continuous with tessellated epithelium, which lines the more delicate ducts of the various glands. For the uses of the ciliated epithelium, *Epithelium ciliatum*, see Cilia.

Mr. Bowman has applied the term *spheroidal* to a variety of epithelium, the cells of which, for the most part, retain their primitive roundness; or, being flattened where they touch, acquire a polyhedral figure. At times, these show a disposition to pass into the columnar, or the tessellated variety; and, hence, this form has been called the *transitional*. The spheroidal epithelium is found in the urinary passages, where it succeeds the columnar epithelium of the urethra at the internal orifice of that canal, and lines the whole of the bladder, ureters and pelvis of the kidneys. It is found, also, in the excretory ducts of the mammary, perspiratory, and of many mucous, glands; and a modification of it lines the inmost secreting cavities or commencing ducts of gland generally.

EPITHĒM, *Epithēma* or *Epithēma*, gen. *Epithēmatis*, (*epithēma* or *epithēma*, from *epi*, and *thēma*, 'to place'). This term is applied to topical applications, which cannot be classed either under ointments or plasters. Three sorts of epithēms have been usually distinguished,—the *liquid*, *dry*, and *soft*, which comprise *fundamentals*, *bags filled with dry substances*, and *emplastica*.

EPITHESIS, (*ἐπιθεσις*.) (Same etymen.) The rectification of crooked limbs by means of instruments.

EPITHYM'IAE. (= *ἐπιθυμία*, and *θυμός*, 'desire.') Mor-
bid desires or longings.

EPITHYMIAMA, gen. *Epithymiama*, (ἐπιθυμία, from *epi*, and *θυμάω*, 'tumigation,') Fumigation.

EPITHYMUM, (*epi*, and *thymos*, 'thyme,')
Cuscuta epithymum — e. Officinatum, Cuscuta
Europæa.

ΕΠΙΤΡΟΧΗΛΕΑ, (*epi*, and *trochlea*, 'a pulley,') (F.) *Epitrochlea*. That which is situate above a pulley. Chaussier gives this name to the unequal, round protuberance situate on the inside of the inferior extremity of the humerus above its articular trochlea. It is the part usually called the *inner* or *lesser condyle* of the humerus. From it originate several muscles of the forearm, to which it has given part of their names, according to the system of Chaussier.

EPITROCHLO-CARPI-PALMAIRE, Palmaris longus—*℥. Cubito-carpien*, see Cubital muscles—*℥. Mito-carpien*, Palmaris magnus—*℥. Phalangien commun*, Flex. r. sublimis perforatus—*℥. Radial*, Pronator radii teres.

EPIZOA, pl. of *Epizoon*, (F.) *Epizootes*. *Ectoparasites*, (*epi*, and *zoon*, 'an animal.') Parasitic animals, which infest the surface of the body, or the common integument.

EPIZOOTIA, *Epizootie*, *Epizootic*, (F.) *Epizootie*. (Same etymon.) A disease which reigns amongst animals. It corresponds, in the veterinary art, to epidemic in medicine.

ÉPIZOOT'IC, *Epizoot'icus*. Relating or appertaining to an epizooty,—as an 'epizootic aptha.' An Epizooty.

EP'ŌCHE, (*ἐπιόχη*, from *ἐπι*, and *ἔχω*, 'to have or to hold.') Doubt, suspension of judgment — *Suspensio sententiæ*. Sometimes employed in the same sense as *ἐπιόχῃς* or suppression.

EPOCHETEUSIS, from *ἐποχέτω*, 'to turn water from its course.') Derivation.

EPODE. ἐπὶ, from *epi*, and ὠδή, 'a song.')

EPO'MIS, (ἐπ'ωμῖς,) *Superhumera'lē*, (ep, and ὠμῖς, 'the shoulder.') The acromion. The upper part of the shoulder.

EPOMPHALICUM, Epomphalum.

EPOMPHALION. (*επομφαλίων*.) Uterus.

EPOMPHALUM, *Epomphalum*, *lep.* and *ομφαλός*, 'the navel.' A medicine which, when placed upon the umbilicus, moves the belly.

ÉPONGE, old (F.) *Espouge*, *Spongia* — *l.*
d'Éplautier, Bedegar.

ÉPOQUES. ('periods,') Menses.

EPOSTOMA, *ep.*, *ostea*, and *oma*.

EPOSTOSIS. (ep, *extern*, and *osis*.) Exostosis.

ÉPREINTES, (e, and *premere*, 'to press,')

Tenesmus.

EPSEMA, gen. *Epsē'matis*. (εψμα, from εψω, 'to boil,') Decoction.

EPSESIS, Decoction, Elixatio.

EPSOM, MINERAL WATERS OF. Epsom is a town in Surrey, about 15 miles from London.

The water is a simple saline;—consisting chiefly of sulphate of magnesia, which has consequently received the name of *Epsom salts*, though no longer prepared from the Epsom water.

ÉPUISEMENT, (e, and *puiser*, 'to draw,')
Exhaustion.

ΕΡΥΘΡΙΣ. (ερυθρος, from ερ, and ουλος, 'the gum.') *Odonitium erythraeum*. *Sarcodon erub.*

Ulon'cus. *Excrecentia gingivæ*. (F.) *Épulis*, *Épulis*. An excrecence on the gum, sometimes ending in cancer.

EPULOSIS, (επουλσις, from *ep*, and *euλω*, 'to cicatrize,') Cicatrization.

EPULOTICA, (επυλωτική,) Cicatrisantia.

ÉPURGE, (e. and *purgare*, 'to purge,') *Euphorbia lathyris*.

ÆQUAL. *Æqualis.* An epithet applied particularly to the pulse and to respiration. The pulse

equally to the pulse and respiration. The pulse and respiration are equal, when the pulsations and inspirations which succeed each other are alike in every respect.

EQUILIB'RUM. (*æquus*, 'equal,' and *libere*, 'to weigh,') In medicine, this word is sometimes used to designate that state of organs, fluids and forces, which constitutes health.

EQUINA CAUDA, see Cauda Equina.

EQUINIA. (*equinus*, 'belonging to a horse.')

Glanders, (L.) *Morve*. A dangerous contagious disorder, accompanied by a pustular eruption, which arises from inoculation with certain diseased fluids generated in the horse, the ass and the mule. Two forms are met with,—*E. mellea*, contracted from horses affected with grease, and *E. glandulosa*, a dangerous and commonly fatal disease communicated to man, either in the acute or chronic form, from the glandered horse. The veterinary surgeons make two varieties of the disease in the horse—*Glanders*, *Melina*, *Melismus*, *Mal'ie*, *Mal's*, *Mallens*; and *Farcy glanders*, *Farcina*, *Mal'ia*, *Farcina*, *Mal'bus farcinarius*, *Mal'bus farcinarius*, *Carbunc'ia* *typica*, *Carbunc'ia farcinaria*, (F.) *Morve farcinosa*, *Farcia*;—the former affecting the pituitary membrane and occasioning a profuse discharge from the nostrils, with pustular eruptions or small tumours, which soon suppurate and ulcerate, being attended by symptoms of malignant fever and by gangrene of various parts;—the latter being the same disease, but appearing in the shape of small tumours about the legs, lips, face, neck, &c. of the horse; sometimes very painful, suppurating, and degenerating into foul ulcers. They are often seen together.

The mild cases require little treatment but rest; the more severe generally resist all remedies.

EQUISETUM ARVENSE, *arvense*, 'a horse,' and *seta*, 'a bristle,' see Hippuris vulgaris—*e. Minus*, Hippuris vulgaris.

EQUITATION, *Equitatio*, *Hippi'a*, *Hippa'-*
ria, *Hippen'eis*, (*equitare*, *equitatum*, 'to ride on
 horseback.') *Horseback exercise*. A kind of
 exercise, advisable in many diseases, although

inproper in others—in uterine affections, for instance. It has been much recommended in phthisis pulmonalis. It is less fatiguing than walking, and consequently more available in disease. But to prevent disease, where much exercise is required, the latter is preferable.

EQUIVOCROUS, *Equiv'ocus*, *Hippoph'agous*, (*equus*, 'a horse,' and *vora*, 'I devour.') Feeding or subsisting on horseflesh.

EQUIVOCAL, *Equiv'ocus*, (*equus*, 'equal,' and *vox*, gen. *vocis*, 'voice.') Those symptoms are occasionally so denominated which belong to several diseases. Generation is said by some to be 'equivocal,' when it is, or seems to be, spontaneous. See Generation.

EQUUS, ('a horse,') Hippus.

ER, in composition, see *En*.

ÉRABLE, *Acer saccharinum*.

ERADICATION, *Eradicatio*, (*e*, and *radix*, gen. *radicis*, 'a root.') The act of rooting out, or completely removing a disease.

ERADICATIVE, *Eradi'cans*. Any thing possessed of the power of completely rooting out a disease.

ÉRAILLEMENT DES PAUPIÈRES, ([*L*]) *e*, and *radius*, 'a ray,') Ectropion.

EROLE, Blister.

EREBINTHUS, *Acer arietinum*.

ERECTEUR DU PÉNIS, Ischio-cavernosus.

ERECHTHITES HIERACIFOLIA, (after *Erechthens* [?]), *Sene'cio hieracifolius*, Fireweed. An indigenous plant, Order, Compositæ, which flowers from July to September. It is said to be an acrid tonic, and astringent, and in large doses emetic.

ERECTILE TISSUE, *Tel'a seu Text'us erectilis*, (*erigere*, *erectum*, (*e*, and *rigere*,) 'to erect,' 'to become erect,') A tissue, whose principal character is to be susceptible of dilatation, i. e. of turgescence and increase of size. It is formed of a collection of arteries and veins, intermixed with nervous filaments; the veins varicose, and contained in spaces formed by trabeculæ of fibrous tissue, being prolongations from the fibrous envelope. Smooth muscular fibres have been detected in it. This tissue exists in the corpora cavernosa of the penis and clitoris, at the lower and inner surface of the vagina, in the spongy part of the urethra, the lips, nipples, iris, &c. Sometimes it is developed accidentally, and constitutes a kind of organic transformation.

ERECT'ION, *Erectio*, *Arrectio*, *Anortho'sis*. The state of a part, in which, from having been soft, it becomes stiff, hard and swollen by the accumulation of blood in the areolæ of this tissue, as in the case of the penis—*Sty'ma*, *Sty'sis*.

ERECTOR CLITORIDIS, Ischio-cavernosus—*e*. Penis, Ischio-cavernosus—*e*. Spinæ, Sacrospinalis.

EREMACAU'SIS, (F.) *Érémacausie*, (*ηρεμα*, 'slowly,' and *καυσis*, 'combustion,') A term applied by Liebig to the slow combustion, oxidation, or decay of organic matters in the air.

EREMIA, (*ηρεμα*, 'rest,') Acinesia.

ERESIPÈLE, Erysipelas.

ERETHILYTICUM, (*ερεθος*, 'redness,' and *λυσις*, 'solution,') Spanæmic.

ERETHISM, *Erethis'mus*, *Erethis'ia*, *Irrita'mentum*, *Irita'tio*, (F.) *Érêthisme*, *Érêtisme*, (*ερεσιζω*, 'I irritate,') Irritation. Augmentation of the vital phenomena in any organ or tissue. *Or'gasm*. Under this name, Mr. Pearson has described a state of the constitution produced by mercury acting on it as a poison. He calls it *Mercur'ial Er'ethism* or *Erethis'mus*. It is characterized by great depression of strength; anxiety about the præcordia; irregular action of the heart; frequent sighing; tremors; small, quick, sometimes intermitting pulse; occasional vomit-

ing; pale, contracted countenance, and sense of coldness: but the tongue is seldom furred, nor are the vital and natural functions much disturbed. In this state any sudden exertion may prove fatal.

ERETHISMA, gen. *Erethis'matia*, (*ερεθισμα*,) Rubefacient.

ERETHISMUS, (*ερεθισμος*,) Erethism, Irritation—*e*. Elbriosorum, Delirium tremens—*e*. Hydrophobia, Hydrophobia—*e*. Oneirodynia, Incubus, Paroniria—*e*. Simplex, Fidgets.

ERETHISTICUS, (*ερεθιστικος*,) Erethiticus, Rubefacient.

ERETHIT'IC, *Erethit'icus*, *Erethis'ticus*. Belonging or relating to erethism—as "erethitic phenomena," or phenomena of irritation.

ÉRÊTISME, Erethism.

EREUGMOS, (*ερευγμος*,) Eructation.

EREUXIS, (*ερευξις*,) Eructation, Flatulence.

ERGASTERION, (*εργαστηριον*, from *εργον*, 'work,') Laboratory—*e*. Spiritûs, Pulmo.

ER'GOT, ([*F*]) *ergot*, ('a spur,') *Er'gota* or *Ergo'ta*, *Secal'è cornu'tum* seu *mater'num* seu *tur'gidum* seu *tenu'entum* seu *lucul'rians* seu *clav'atum*, *Clav'us secal'inus* seu *secal'is* seu *Silig'ini's*, *Spermo'dia* *clav'us*, *Ma'ter secal'is*, *Cal'ear*, *Se-cal'ia* *ma'ter*, *Ustil'go*, *Fru'men'tum cornu'tum* seu *cornicula'tum* seu *lucul'rians* seu *tur'gidum* seu *tenu'entum*, *Brizoc'eras* [?], *Melanoph'ia*, *Spur'ed* or *Horn'ed* Rye, *Spur*, *Horn'seed*, *Cock'spur* rye, *Cock'spur*, (F.) *Seigle ergoté ou cornu ou corrompu* ou *à Éperon* ou *Ergotisé*, *Blé cornu*, *Ergot* ou *Clou* ou *Mère de seigle*, *Bled avorté ou farouche*, *Faux seigle*. Ergot is generally considered to be the result of a disease in rye—*Secal'è cereal'è*—occurring most frequently when a hot summer succeeds a rainy spring. Decandolle, however, regards it as a parasitic fungus, and calls it *Sclero'tium clav'us*; whilst Leveillé esteems it to be a fungus giving a coating to the diseased grain; the medical virtues residing in the coating. This parasitic fungus he calls *Sphae'lia seg'etum*. More recently, it has been maintained, that it is a diseased state of the grain occasioned by the growth of a fungus not previously detected, to which the names *Ergot'ia abort'ans* seu *abortif'iciens*, *O'idium abortif'iciens*, *Ergotmould*, have been given. It is found projecting from among the leaves of the spike or ear, and is a long, crooked excrescence, resembling the spur of a cock, pointed at its extremities, of a dark brown colour externally, and white within.

Ergot has been long used in Germany to act on the uterus, as its names Mutterkorn and Gebärpulver ('womb-grain,' *pul'vis partu'riens*, *pul'vis ad partum*) testify. Upwards of fifty years ago, it was recommended in this country for accelerating parturition; and since that period, numerous testimonials have been offered in its favour. Half a drachm of the powder is gently boiled in half a pint of water, and one-third part given every 20 minutes, until proper pains begin. Some recommend the clear watery infusion; others advise the powder; others the oil; and others the wine. It is obvious that, in many cases, the uterine efforts would return in the period which would necessarily elapse in the administration of ergot; so that several of the cases, at least, of reputed efficacy, may not have been dependent upon the assigned agent. Ergot also possesses, it is affirmed, narcotic virtues, which have rendered it useful in hemorrhagic and other affections of excitement.

Bread, made of spurred rye, has been attended with the effects described under *ERGOTISM*.

ERGOT, ('a spur,') Hippocampus minor—*e*, *de Seigle*, Ergot.

ERGOTMOULD, see Ergot.

ERGOTA, Ergot.

ERGOTETIA ABORTANS, (*ergot*, and *area*, 'cause,') see *Ergot*—e. Abortifaciens, see *Ergot*.

ERGOTIN, *Ergotinum*, *Extractum Ergoti*, *Extrait of Ergot*, (F.) *Ergotine*, *Extrait hémotique de Baume*. The extract of ergot has been found an excellent hemostatic in the dose of two grains several times a day. It has also been used externally.

ERGOTINE, *Ergotin*.

ERGOTISM, *Ergotismus*, from *ergota*, 'ergot.' *Mortuus cerealis*. Poisoning by ergot. An affection produced by the use of spurred rye. At times, the symptoms are limited to vertigo, spasms, and convulsions. See *Convulsio cerealis*. Most commonly, there is torpor with numbness of the hands and feet, which waste away, lose sensation and the power of motion, and separate from the body by dry gangrene.—*Ergotismus Sphaecelatus*, *Sphaecelatus cerealis*, *Neurosis cerealis*, *Gangraena ustilaginacea*, *Neurosis ustilaginacea*, *Ergot or Mildew mortification*, (F.) *Ergotisme gangréneux*, *Gangrène des Semences*, *Mol de Saligne*.

ERGOTISME CONVULSIVE, *Convulsio cerealis*.

ERGOTISMUS, *Ergotismus*—e. Spasmodicus, *Convulsio cerealis*—e. Sphaecelosis, *Ergotismus*.

ERICA VULGARIS, (*erica*,) *Calluna vulgaris* see *Erica*, *Common Heath*, (Sc.) *Heath'er*, (F.) *Beydre vulgaire*. This has been regarded as diuretic and diaphoretic.

ERIGERON AMBIGUUM, (*erigeron*, from *er*, 'spring,' and *geron*, 'an old man,' owing to the hoary appearance of some of the vernal species,) *E. Philadelphicum*—e. *Annuum*, see *E. Canadense*.

ERIGERON BELLIDIFOLIUM, *Robin's plantain*, *Ruby Betty*. An indigenous plant of the Composite family; flowering in May. It is said to possess properties like those of *E. Philadelphicum*.

ERIGERON CANADENSE, *Canada Fleabane*, *Pride-weed*, *Horse-weed*, *Butter-weed*, is considered to be allied in properties to *E. Philadelphicum*; and the same may be said of *Erigeron Heterophyllum*, *E. annuum*, *Succet Soabione* or various-leaved *Fleabane*; which is in the secondary list of the Pharmacopœia of the United States.

ERIGERON HETEROPHYLLUM, see *E. Canadense*—e. *Integrifolium*, *E. Philadelphicum*.

ERIGERON PHILADELPHICUM, *E. strigosum* seu *ambiguum* seu *integrifolium*, *Soabione*, *She'eisch*, *Philadelphica* or *She'eisch* or *Nor'one-leaved Fleabane*, *Da'ieg*, *Cucash*, *Frost-weed*, *Field-weed*, *Square-weed*, (F.) *Vergerette de Philadelphie*. The plant has been used in decoction or infusion for gouty and gravelly complaints, and is said to operate powerfully as a diuretic and sudorific. The essential oil has been given in uterine hemorrhage. Dose, five drops.

In the Ph. U. S. 1863, *Erigeron* means the herb of *E. Heterophyllum* and *E. Philadelphicum*.

ERIGERON STRIGOSUM, *E. Philadelphicum*.

ERIGERUM, *Senecio*.

ÉRIGNE, (from *erige*, 'to raise,') *Hook*.

ÉRINE, *Hook*.

ERIOCEPHALUS UMBELLULATUS, (*erion*, 'wool,' and *kephal*, 'head,' from the character of its flowers.) *Wild Rose-mary*. A South African shrub, *Ord. Composite*, which is diuretic, and used by the farmers and Hottentots in various forms of dropsy.

ERIOSPERMUM LATIFOLIUM, (*erion*, 'wool,' and *erosum*, 'seed,' from the woolly envelope of the seeds.) A South African plant, the tuber of which is muculent, and used externally in abrasions of the skin, and in superficial ulcers. It is also employed by the Mohammedans, in decoction, in amenorrhœa.

ERODENS, (*er*, and *rodere*, 'to gnaw,') *Cautic*.

ERODIUM MOSCHATUM, (from *erodios*, 'a heron,') *Geranium moschatum*.

EROMANIA, (*eromania*, from *eros*, and *mania*,) *Erotomania*.

ERONOSUS, *Eronosus*, (*er*, 'spring,' and *nosos* or *noctos*, 'disease,') A disease which appears chiefly in spring.

EROS, (*eros*, gen. *erosus*,) *Love*.

EROSIO, *Corrosion*, *Erosion*—e. et *Perforatio spontanea Ventriculi*, *Gastromalacia*.

EROSION, *Erosio*, *Diabodis*, *Acidosis*, *Corrosio*, (*erodere*, *erodere*, (*er*, and *rodere*,) 'to eat away,') The action of a corrosive substance, or the gradual destruction of a part by a substance of that kind. It is often employed in the same sense as ulceration,—*spontaneous erosion*.

EROSION, GANGRENOUS, OF THE CHEEK, *Cancer aquaticus*.

EROTIC, *Eroticus*, (*eroticus*, from *eros*,) That which is produced by love—as *Erotic melancholy*, *E. delirium*, &c.

EROTICOMANIA, *Erotomania*.

EROTOMANIA, *Erotomania*, *Mania erotica*, (from *eros*, and *mania*, 'mania,') A species of mental alienation caused by love. *Erotic melancholy* is *Love melancholy*. Some authors write it *Erotico-mania*. Also, *Nymphomania*.

ERPES, (*erps*,) *Herpes*.

ERRABUNDUS, (*errare*, 'to wander,') *Planctes*.

ERRATIC, *Erraticus*, *Atactos*, *Atypos*; *Wandering*, *Irregular*. That which disappears and recurs at irregular intervals. This name is given to intermittents, *Febres eraticæ*, which observe no type. Most commonly, it is applied to pains or to any diseased manifestations which are not fixed, but move from one part to another, as in gout, rheumatism, erysipelas, &c.

ERREUR DE LIEU, *Error loci*.

ERRHINE, (*erhine*,) *Erbinum* or *Erhinum*, *Sternutatorium*, *Sternutatory*, *Purmicum*, *Aphlognaticum per nares*, *Nasale*, (*er*, and *rhine*, 'the nose,') A remedy whose action is exerted on the Schneiderian membrane, exciting sneezing and increased discharge. *Erhines* have been mainly used as local stimulants in head affections.

ERRHINUM, *Erhine*, *Sternutatory*.

ERRHYSIS, (*er*, and *rhesis*, 'I flow,') With some, a draining of blood. A trifling hemorrhage.

ERROR LOCI, ('error of place,') *Apophanesis*, (F.) *Erreur de Lieu*. A term frequently used by Boerhaave to express deviation of fluids, when they enter vessels not destined for them. Boerhaave admitted several orders of capillary vessels, the diameters of which went on decreasing. The largest received red blood, those which came next received only white, others only lymph, whilst the last of all were destined for yet more subtle fluids—for a kind of vapour. When the red globules entered vessels destined for the white, or the white those intended for the thinner fluids, there was, according to the theory of Boerhaave, an *error loci*, the proximate cause of inflammation and other mischiefs. See *Aberration*.

ERUCA, *Brassica eruca*, *Sinapis*—e. *Barbarea*, *Erysimum barbarea*—e. *Fœtida*, *Brassica eruca*—e. *Sativa*, *Brassica eruca*.

ERUCTATION, (*eructare*, *eructum*, (*er*, and *ructare*,) 'to belch,') *Ructatio*, *Eructum*, *Eructio*, *Ructus*, *Ructus men*, *Ructatio*, *Eructio*, *Resurgatio*, (Old Eng.) *Belke*, *Bouffe*, *Belke*, (Sc.) *Bock'ing*, *Bok'ing*, *Ruck'ing*, *Efti*, (Prov.) *Kelk'ing*, *Rup'ing*, *Rup'ing*, *R'ing*, *Ruf'ing* (N. of England); *Belching*, (F.) *Rot*, *Rapport*. A sonorous emission, by the mouth, of flatus proceeding from the stomach. When so frequent as to occa-

sion a diseased condition, this is termed *Ructuo-sitas*, *Morbus ructuosus*.

ERUGATORIA, (e, and ruga, 'a wrinkle.') Tetanortha.

ÉRUGINEUX, Érugineus.

ERUPTIO, (erumpere, eruptum, (e, and rumpere, 'to issue with violence,' 'to break out.')

Eruption—e. Exanthematis, Epanastasis—e. Sanguinis, Hæmorrhagia.

ERUPTION, *Eruptio*. This word has several meanings. 1. The sudden and copious evacuation of any fluid from a canal or cavity—of serum, blood, pus, &c., for example; 2. The breaking out of an exanthem; and 3. The exanthem itself, whether simply in the form of a rash, or of pustules, vesicles, &c.

ÉRUPTION ANOMALE, Roseolæ—e. Roseace, Roseolæ—e. Violet, Ionthus.

ERUPTIVE, *Eruptivus*. That which is accompanied by an eruption. Thus small-pox, measles, scarlet fever, miliaria, &c., are *eruptive fevers*. The term *eruptive disease* is nearly synonymous with cutaneous disease.

ERVA DE SANTA MARIA, Arum Dracunculus—e. do Rato, Palicourea Maregraavii.

ERVALENTA, see Eryum.

ERVUM, *E. Ervil'ia*, *E. plicatum*, *Vie'ia ervil'ia*, *Orob'us*, (F.) *Lentille Ers* ou *Orobe*. In times of scarcity, the seeds have been made into bread, but it is said not to be wholesome. The meal was once much used in poultices; it was formerly called *Orob'ion*. Under the name *Ervalent'a*, a secret dietetic preparation has been introduced. It is a vegetable powder, which, when, mixed with milk and soup and taken night and morning, is said to have succeeded in removing old and obstinate obstructions! Buchner thinks it is bean-meal, Ditterich, the meal of the seed of Eryum Ervilia.

A specimen of Warton's *Ervalent'a*, obtained at the dépôt, and analyzed by Dr. Hassall, was found to contain a mixture of the French and German *lentil*, reduced to powder, including portions of the shells or husks, and of a substance very closely resembling, in its microscopic characters, *corn meal*. A sample of Du Barry's *Revalent'a Arab'ica* was found to consist of a mixture of the *Egyptian* or *Arab'ian lentil* and *barley meal*.

ERVUM ERVILIA, Eryum.

ERVUM LENS, *Ci'cer Lens*, *Lens esculen'ta*, *Pha'cos*, the *Lent'il*, (F.) *Lentille*. The seed, *Pha'cè*, *Pha'cea*, *Pha'cus*, is eaten in many places like peas: they are flatulent and difficult of digestion.

ERVUM Plicatum, Eryum,

ERYGE, (ερυγη), Eructation.

ERYNGIUM AQUATICUM, (ερυγγιον,) *E. yuccefolium*, *Water Eryn'go*, *But'ton Snake'root*. Order, Umbelliferae. This root, *Eryn'gium*, (Ph. U. S.), is nearly allied to the contrayerva of the shops, and acts more especially as a sudorific. It is a secondary article in the Pharm. U. S.

ERYNGIUM CAMPESTRÈ, *E. vulga'rè*, *Cent'ium Cap'ita*, *Aster inquin'alis*, *Capitulum Mar'tis*, *A'cus Ven'is*, (F.) *Panicaut*, *Chardon-Roland*, is sometimes used for *E. maritimum*.

ERYNGIUM FETIDUM; indigenous, is said to possess properties like those of valerian and contrayerva.

ERYNGIUM MARITIMUM seu MARINUM, (ερυγγιον, from ερυγομαι, 'to belch.') The *Sea Holly* or *Eryn'go*, *Æthe'rea Her'ba*, (F.) *Panicaut Maritime*. *Eryngo* root has a slightly aromatic odour, and sweetish and warm taste. It is considered to be diuretic and expectorant, but its powers are so weak, that it is scarcely ever used.

ERYNGIUM VULGARE, *E. Campestre*—e. *Yuccæfolium*, *E. Aquaticum*.

ERYNGO, SEA, *Eryngium maritimum*—e. *Water*, *Eryngium aquaticum*.

ERYSIMUM, *E. officina'le*, (ερω, 'to draw,' from its power of raising blisters.) *Sisymbrium officina'le*, *Chama'plion*, *Hedgemustard*, (F.) *Vélar*, *Tortelle*, *Herbe aux Chantres*. This was, formerly, much used for its expectorant and diuretic qualities. The seeds greatly resemble mustard.

ERYSIMUM ALLIARIA, *Alliaria*.

ERYSIMUM BARBARE'A, *E. lyra'tum*, *Arabis barbare'a*, *Sisymbrium barbare'a*, *Eru'ca barbare'a*, *Barbare'a*, *B. stric'ta*, (F.) *Herbe de Sainte Barbe*, has been ranked amongst the antiscorbutic plants. See *Alliaria*.

ERYSIMUM CORDIFOLIUM, *Alliaria*—e. *Lyratum*. *E. barbarea*.

ERYSIPELA'CEA. (Same etymon as the next.) A family of eruptive diseases, comprising erysipelas, variola, rubella, and scarlatina.

ERYSIPE'LAS, gen. *Erysipel'atis*, (ερυσίπelas, from ερω, 'I draw,' and πelas, 'near,' from its tendency to spread, or from ερως, 'red.' [?]) *Febris erysipelato'sa* seu *erysipela'cea*, *Erythropyra*, *Em'phlysis Erysip'elas*, *St. Anthony's fire*, *Wild'fire*, *Ignis Sancti Ant'o'nii* seu *Columell'æ* seu *Sa'cer*, *Hieropyr*, *Icterit'ia ru'bra*, *Ru'bea icterit'ia*, *Vicini'taha*, *Vicini'trac'tus*, *Bru'nus*, *Lug'dus*, *Rose*, (F.) *Érysipèle*, *Érésipèle*, *Feu St. Antoine*, *Feu sacré*, *Mal Saint Antoine*. A disease, so called because it generally extends gradually to the neighbouring parts. Superficial inflammation of the skin, with general fever, tension and swelling of the part; pain and heat more or less acrid; redness diffused, but more or less circumscribed, and disappearing when pressed upon by the finger, but returning as soon as the pressure is removed. Frequently, small vesicles appear upon the inflamed part, which dry up and fall off, under the form of branny scales. Erysipelas is, generally, an acute affection: its medium duration being from 10 to 14 days. It yields, commonly, to general refrigerant remedies. Topical applications are rarely serviceable. At times, when the disease approaches the phlegmonous character, copious bleeding and other evacuations may be required, as in many cases of erysipelas of the face; but this is not commonly necessary. In most cases, indeed, the general action seems deficient, and it becomes necessary to give tonics.

When erysipelas is of a highly inflammatory character, and invades the parts beneath, it is termed *Erysip'elas phlegmono'des*, (F.) *Érysipèle phlegmoneux*: when accompanied with phlyctenæ, and the inflammation terminates in gangrene, *Erysip'elas gangren'o'sum*, (F.) *Érysipèle gangrèneux*; and when associated with infiltration of serum, *Erysip'elas œdema'to'sum*, (F.) *Érysipèle œdémateux*.

At times, the surface in erysipelas has a bronzed rather than a red hue—*bronzed erysipelas*.

ERYSIPELAS, BRONZED, see Erysipelas—e. Gangrenosum, see Erysipelas—e. *œdema'to'sum*, see Erysipelas—e. *Periodica nervosa chronica*, *Pellagra*—e. *Phlegmonodes*, see Erysipelas—e. *Phlyctenoides*, *Herpes zoster*—e. *Pustulosa*, *Herpes zoster*—e. *Zoster*, *Herpes zoster*.

ERYSIPELATODES, (ερυσίπελατώδης), Erysipelatous.

ERYSIPEL'ATOUS, *Erysipelato'des*, *Erysipelato'sus*. Belonging to erysipelas; as an *erysipel'atous affection*.

ERYSIPEL'ATOUS INFLAMMA'TION, *Inflamma'tio erysipelato'sa*, *Epiphlogis'ma*, *Cuti'tis*, *Dermati'tis*, is the character of inflammation which distinguishes erysipelas, *Erythe'ma erysipelato'sum*.

ÉRYSIPE'LE, Erysipelas—*é. Ambulant*, see *Ambulant*—*é. Gangrèneux*, see Erysipelas—*é. Œdémateux*, see Erysipelas—*é. Phlegmoneux*, see Erysipelas—*é. Pustuleux*, *Herpes zoster*—*é. Serpiginéux*, see *Serpiginous*.

ERYSOS, ερως, ('red.') Erythema.

ERYTHE'MA, gen. *Erythematia*, (ερυθμας, *epuroas*, 'red, rose-coloured.') *Inflammatory blush*, *Erythre'ma*, (F.) *Érythème*. This name is, sometimes, given to erysipelas, especially when of a local character. It is, also, applied to the morbid redness on the cheeks of the hectic and on the skin covering bubo, phlegmon, &c. It forms the 6th genus of the 3d order, *Exanthemata*, of Bate-man's classification of cutaneous diseases; and is defined, 'a nearly continuous redness of some portion of the skin, attended with disorder of the constitution, but not contagious.' Many varieties are described by dermatologists,—for example, *E. intertrigo*, *E. ful'gax*, *E. papula'tum*, *E. tubercu-la'tum*, *E. nodosum*, *E. læ'vè*, *E. centrif'ugum*, *E. margina'tum*, and *E. acrody'nia*.

ERYTHEMA ACRODYNIA, Acrodynum—e. Ambustio, Burn—e. Anthrax, Anthrax.

ERYTHEMA CENTRIFUGUM. A variety of erythema attacking the face, commencing with a small red spot, and spreading around, so as, at times, to affect the whole face.

ERYTHEMA CIRCINATUM, see E. Marginatum.

ERYTHEMA ENDEMICUM, Pellagra—e. à Frigore, Chilblain.

ERYTHEMA FU'GAX. Patches of erythema, that sometimes appear on the body in febrile diseases, teething children, &c., and which are of brief duration.

ERYTHEMA GANGRENOsum, Anthrax—e. Icho-rosum, Eczema mercuriale—e. Intertrigo, Chafing.

ERYTHEMA LÆ'VÈ, (F.) *Érythème léger*. Slight erythema affecting the skin, as in oedema.

ERYTHEMA MARGINA'TUM, (F.) *Érythème marginal*. A form in which the prominent patches of erythema are distinctly separated from the skin at their margins. When they can be distinctly measured or composed we have *E. circina'tum*.

ERYTHEMA MERCURIALE, Eczema mercuriale.

ERYTHEMA NODO'SUM, (F.) *Érythème noueux*. A more severe form of Erythema tuberculatum.

ERYTHEMA PAPULA'TUM, (F.) *Érythème papuleux*. Patches of erythema, which are at first papulated, appearing on the face, neck, breast, &c., of females, and young persons more especially.

ERYTHEMA PELLAGRUM, Pellagra—e. Pernio, Chilblain.

ERYTHEMA SOLA'RE. A name given by the Italians to Pellagra, owing to its being attended with an eruption of small red spots or patches on the parts that are exposed to the sun.

ERYTHEMA TUBERCULA'TUM, *E. tuberculo'sum*, (F.) *Érythème tuberculeux*. A form of erythema in which the patches are in small lumps.

ERYTHEMATOUS, *Erythemat'ic*, *Erythema-tous*, *Erythemat'icus*, (F.) *Erythémateux*. Appertaining or relating to Erythema.

ÉRYTHÈME, Erythema—*é. Léger*, Erythema læve—*é. Marginal*, Erythema marginatum—*é. Noureux*, Erythema nodosum—*é. Papuleux*, Erythema papulatum—*é. Tuberculeux*, Erythema tuberculatum.

ERYTHRÆA CENTAURIUM, (from *epudpos*, 'red,' 'the flower.') *Chironia centaarium*—e. Chilensis, Canchalagua, Chironia Chilensis.

ERYTHRÆMIA, (erythro, and *huma*, 'blood.') Plethora.

ERYTHRAN'CHÈ, *Angi'na erysipelato'sa*, (erythro, and *ayxw*, 'I suffocate.') Erysipelatous or erythematous sore throat.

ERYTHREMA, gen. *Erythre'matis*, Erythrema.

ERYTHRINA MONOSPERMA, (epudpos, 'red,'—the flowers.) *Butea frondosa*.

ERYTHRO, (from *epudpos*, 'red,') in composi-tion, red.

ERYTHROCHERAS, (erythro, and *chæras*.) *Cynanche parotidea*.

ERYTHRODANUM, (epudpos, 'red,') *Rubia*.

ER'YTHROID, *Erythro'des*, *Erythroi'des*, (erythro, and *aios*, 'resemblance.') Reddish.

ERYTHROID COAT, *Tu'nica erythro'des*. Vagi-nal coat of the testis.

ERYTHROID VES'ICLE, *Vesic'ula Erythroi'dea*. A vesicle of the fœtus described by Pockels and others. It is pyriform, and much longer than, although of the same breadth as, the umbilical vesicle. Velpeau, Weber, and others, doubt its being a physiological condition.

ERYTHRONIUM, (from *epudpos*, 'red,' which does not, however, apply to the American spe-cies,) *E. Americanum*.

ERYTHRONIUM AMERICA'NUM, *E. fla'tum* seu *dens ca'nis seu lanceola'tum seu longifo'lium*, *Com'mon Erythro'nium*, *Yel'low Snake'leaf*, *Yel'low Ad'der's tongue*, *Ad'der'leaf*, *Dog vi'olet*, *Dog's tooth vi'olet*, *Rat'tlemake vi'olet*, *Lamb's tongue*, *Scrof'ula root*, *Yel'low Snow'drop*. Ord. Liliaceæ. This plant is possessed of emetic properties, but is rarely, if ever, used. The root and herb, *Erythro'nium*, (Ph. U. S.), are in the secondary list of the Pharmacopœia the United States.

ERYTHRONIUM DENS CANIS, *E. Americanum*—e. Flavum, *E. Americanum*—e. Lanceolatum, *E. Americanum*—e. Longifolium, *E. Americanum*.

ERYTHROPHLEUM JUDICIALE, (erythro, and *phleu*, 'to flow'—from the red juice.) *Sassy bark tree*.

ERYTHROPYRA, (erythro, and *pur* 'fire, fever.') Erysipelas.

ERYTHRORRHIZA ROTUNDIFOLIA, (erythro, and *ῥιζα*, 'root,') *Galax aphylla*.

ERYTHROSIS, (erythro, and *osis*.) Ple-thora.

ERYTHROXYLON COCA, *Co'ca*, (erythro, and *ξύλον*, 'wood,') A plant. Ord. Violaceæ, which grows on the eastern slopes of the Andes, is prized in Peru, Chili, and Bolivia, both as a medicine and an article of diet; and serves as a substitute for the tea, coffee, betel, tobacco, hashish, and opium of other peoples. An active alkaloid—*Co'caïne*, *Cocai'na*—has been separated from it.

ES'APHE, (εσάφω, from (εἰς, and ἀφω, 'I feel,') 'I feel inside,') (F.) *Le Toucher*, *Le Toucher vaginal*. The introduction of a finger or fingers into the vagina, for the purpose of examining the condition of the uterus. It is employed to verify the existence or non-existence of pregnancy and its different stages; to detect certain affections or faults of conformation in the genital organs and pelvis; and, during the process of accouchement, to examine into the progress of labour, &c.

ESCA, Aliment.

ESCALDAS, MIN'ERAL WATERS OF, in the department Pyrénées-orientales, France, are thermal sulphureous, containing sulphuret of sodium. Temperature about 108° Fah. The season is from June to the 15th of September.

ESCARA, Eschar.

ESCARGOT, Helix pomatia, Limax.

ESCHALOT, Échalotte.

ES'CHAR, *Es'chara*, (εσχα, *Es'cara*, (Old Eng.) *Asker*; a slough, a crust or scab. The crust or disorganized portion, arising from the mortification of a part, and which is distinguishable from the living parts by its colour, consistence, and other physical properties. The inflammation which it excites in the contiguous parts gives occasion to a secretion of pus between the living and dead parts, which ultimately completely separates the latter.

ESCHARA, (εσχα, *Vulva*.

ESCHAROTIC, *Escharot'icum*, (εσχαρωτικον, from *eschar*.) Any substance which, when applied upon a living part, gives rise to an eschar, such as caustic potassa: the concentrated mineral acids, &c.

ESCHAROTIC POISON, see Poison.

ESCHELLOHE, MINERAL WATERS OF. This water rises at the foot of a mountain near the convent of Etal in Bavaria. It is a sulphuro-saline.

ESCHID'NA OCELLA'TA. A very poisonous viper, which inhabits the sugar-cane fields of Peru. Its bite is almost instantaneously fatal.

ESCORZONERA, Scorzonera.

ESCREVISSE, (G. Krebs,) Crab.

ESCLAPIAN, (from *Æsculapius*.) Medical.

ES'CULENT, *Esculentus*, *Ed'ible*, *Eat'able*, *Ciba'rious*, (*esca*, 'food,') (F.) *Comestible*. Such plants or such parts of plants or animals as may be eaten for food.

ESÈRE, Ordeal nut.

ESKIN, (Sax. *gīfcian*, 'to hiccuph.') Singsults.

ESO, (*eso*, 'within,') in composition, within.

ESOC'HAS, (*εσος*, from *eso*, and *εχειν*, 'to have,') Esoche.

ES'OCHE, *Es'ochas*, A tumour—as a hemorrhoid—within the anus.

ESOCOLITIS, (*eso*, and *colitis*.) Dysentery.

ESODIC, (*eso*, and *ὅς*, 'a way,') Afferent.

ESOFENTERITIS, see Enteritis.

ESOGASTRITIS, Endogastritis.

ESOTERIC, (*εσωτερος*, 'interior,' from *eso*.) Private; interior; in contradistinction to *Esoter'ic*, public, exterior. Hence, we speak of '*esoter'ic* and *exoter'ic* causes of disease.'

ES'OTERISM, MEDICAL. A term used by M. Simon, in his *Déontologie médicale*, for that esotery, or mystery and secrecy, with which the practitioner performs his daily duties, and which, he conceives, he is compelled to adopt by the prejudices and ignorance of his patients.

ESPACES INTERCOSTAUX, (from *spatium*, 'a space,') See Costa.

ESPÈCES, Species, see Pulvis—e. *Pulvéru-lentes*, see Pulvis—e. *Vulnéraires*, Falltranck.

ES'PHLASIS, (*es*, *eis*, 'inwards,' and *φλαω*, 'I break, bruise,' &c.) A fracture of the skull by comminution, the fragments being depressed. See Depression, and Enthlasia.

ESPRIT, Spirit—e. *d'Ammoniaque*, Spiritus ammoniac—e. *d'Anis*, Spiritus anisi—e. *de Bois*, Spiritus Pyroxylicus—e. *de Camphre*, Spiritus camphoræ—e. *de Cannelle*, Spiritus cinnamomi—e. *de Carvi*, Spiritus carui—e. *d'Éther sulfurique*, Spiritus ætheris sulphurici compositus—e. *de Genièvre composé*, Spiritus juniperi compositus—e. *d'Ica*, Achillea moschata—e. *de Lavande*, Spiritus lavandulæ—e. *de Lavande composé*, Spiritus lavandulæ compositus—e. *de Menthe poivrée*, Spiritus menthæ piperitæ—e. *de Muscade*, Spiritus myristicæ—e. *de Poultot*, Spiritus pulegii—e. *Pyroxylique*, Spiritus pyroxylicus—e. *de Ruifort composé*, Spiritus armoraciæ compositus—e. *de Romaria*, Spiritus rosmarini—e. *de Vin délayé*, Spiritus tenuior—e. *de Vin rectifié*, Spiritus rectificatus.

ESPRITS ANIMAUX, ('animal spirits,') Nervous fluid.

ESQUILLE, (*σχιδή*, [L.] *squida*, dim. *squidilla*, 'a splinter,') Splinter.

ESQUINANCIE, Cynanche, Cynanche tonsillar.

ESQUINE, Smilax China.

ES'SENCE, *Essen'tia*, (corrupted from *existentia*, 'standing out,' [?] or, from *esse*, 'to be,') By some, used synonymously with *volatile oil*; by others, with *simple tincture*. The oil of pepper-mint is the oil obtained by distillation; the *essence*, the oil diluted with spirit. See Tinctura olei menthæ.

ESSENCE OF ANISE, *Essentia anisi*—e. of Bar-laza, Hill's, Tinctura guaiaci ammoniata—e. of Caraway, *Essentia Carui*—e. of Cinnamon, *Essen-*

tia Cinnamomi—e. of Coltsfoot, see Tinctura ben-zoini composita—e. of Fennel, *Essentia Fœniculi*—e. of Lemon, Spiritus limonis—e. of Musk, Tinctura moschi—e. of Mustard pills, White-head's, see Sinapis—e. of Mustard, Whitehead's, see Sinapis—e. of Pennyroyal, *Essentia menthæ pulegii*—e. of Peppermint, Spiritus menthæ piperitæ—e. of Pimento, *Essentia Pimentæ*—e. of Rosemary, *Essentia Rosmarini*—e. of Roses, see Rosa centifolia—e. of Spearmint, Spiritus menthæ viridis—e. of Spruce, see Pinus Abies.

ESSENCES, Olea volatil.

ESSENTIA, Essence, Tincture—e. Abietis, see Pinus abies—e. Aloes, Tinctura aloes—e. Absinthii amara, Tinctura A. composita—e. Absinthii composita, Tinctura A. composita.

ESSENTIA ANI'SI, *Es'sence of An'ise*. (Ol. anisi f 3j; Sp. rectif. f 3ix. M. Ph. D.) Dose, ℥xx to f 3j, as a carminative.

ESSENTIA ANTIMONII SEU STIBII, Vinum antimonii tartarizati—e. Antiseptica Huxhami, Tinctura cinchonæ composita—e. Aromatica, Tinctura cinnamomi composita.

ESSENTIA CARUI, *Es'sence of Car'away*. (Ol. carui, f 3j; Sp. rectif. f 3ix. Ph. D.) Dose, gtt. xx to f 3j, as a carminative.

ESSENTIA CASTOREI, Tinctura castorei—e. Catholica purgans Rothii, Tinctura jalapii composita—e. de Cedro, Oleum cedrinum—e. Chinæ Tinctura cinchonæ composita—e. Cinnamomi, Tinctura cinnamomi.

ESSENTIA CINNAMO'MI, *Es'sence of Cin'namon*. (Ol. cinnam. f 3j; Sp. rect. f 3ix. M.) Dose, gtt. x to xx.

ESSENTIA FENIC'ULI, *Es'sence of Fen'iculi*. (Ol. fœnic. f 3j; Alcohol. f 3ix. M. Ph. D.) Dose, gtt. x to f 3j.

ESSENTIA MENTHÆ PIPERITÆ, Tinctura olei menthæ piperitæ.

ESSENTIA MENTHÆ PULE'GII, *Es'sence of Pen'nyroyal*. (Ol. puleg. f 3j; Sp. rectif. f 3ix. Ph. D.) Dose, gtt. xv to xxx.

ESSENTIA MENTHÆ VIRIDIS, Tinctura olei menthæ viridis.

ESSENTIA MYRISTICÆ MOSCHA'TÆ, *Es'sence of Nut'meg*. (Ol. myrist. f 3j; Sp. rectif. f 3ix. M. Ph. D.) Dose, gtt. xx—xxx.

ESSENTIA NEROLI, see Citrus aurantium—e. Corticis Peruviani antiseptica Huxhami, Tinctura cinchonæ composita—e. Corticis Peruviani composita, Tinctura cinchonæ amara.

ESSENTIA PIMEN'TÆ, *Es'sence of Pimen'to*. (Ol. Piment. f 3j; Sp. rectif. f 3ix. M. Ph. D.) Dose, gtt. xxx ad lx.

ESSENTIA ROSMARI'NI, *Es'sence of Rose'mary*. (Ol. Rosmarini f 3j; Sp. rectif. f 3ix. M. Ph. D.) Dose, gtt. xxx ad gtt. lx.

ESSENTIAL, *Essential'is*. The word *essential* has been applied to the immediate or active principles of vegetables, which were believed to be endowed with the properties of the plants from which they were extracted. *Essential oil* was so called because it was regarded as the only immediate principle which was *essential*. This expression is retained. *Essential salts* are true salts or extracts, which exist, ready formed, in vegetables; and which are obtained by distillation, incineration, or some other process.

An *essential disease* is synonymous with a general disease; that is, one not depending on any local affection,—not symptomatic. See Entity.

ES'SERA, *Es'sere*, *So'ra*, *Sai'r'e*, *Sa'r'e*, *Mor'bus porci'nus*, *Ro'sa salt'ana*, *Urtica'ria porcell'ana*, (F.) *Ampoules*, *Porcelaine*. (Of Arabic derivation.) A species of cutaneous eruption, consisting of small, reddish tubercles over the whole body, accompanied by a troublesome itching. It seems to be a variety of lichen or urticaria. See Lichen tropicus.

ESSIEU, (from *axiculus*, dim.) Axis.

ESSOUFFLEMENT, (e, and *souffle*, 'breath.') Anhelatio.

ESTHEMA. gen. *Esthe'matis*, (εσθημα, 'a garment.') Vestitus.

ESTHESIS, (εσθησις, 'clothing.') Vestitus.

ESTHIOMÈNE, Esthiomenus.

ESTHIOMENUS, *Esthiom'enus*, (εσθιω, 'I eat.') *Es'edens*, (F.) *Esthiomène*. That which devours or eats away. Certain ulcers and eruptions—*Dartres rongeantes*—are so called.

ESTIVAL, *Esti'veus*; (*æstas*, gen. *æstatis*, 'summer.') Happening in summer, belonging to summer. This epithet is given to *summer diseases*, so called because they reign at that season, and appear to depend on the influence exerted by it on the animal economy. In the United States, the term *summer disease* or *complaint* means disorder of the bowels; as *diarrhœa*, *cholera morbus*, &c.

ESTOMAC, Stomach.

ESTRAGON, *Artemisia dracunculæ*.

ESULA CYPARISSIAS, *Euphorbia cyparissias*—e. Major, *Euphorbia palustris*—e. Minor, *Euphorbia cyparissias*.

ÉSULE GRANDE, *Euphorbia lathyris*.

ESURIES, Hunger.

ESURIGO, gen. *Esurig'iniis*, Hunger.

ESURITIO, Hunger.

ETARISTRIA, (εταριστρια, from εταρσω, 'to indulge unnatural desires.') Tribas.

ÉTAGE, (old [F.] *estage*,) Stage.

ÉTAÏN, [old F.] *estaïn*, from (L.) *stagnum*, *stagnum*. Tin—é. *de Glace*, Bismuth—é. *Gris*, Bismuth.

ÉTAT, [old F.] *estat*, from *stare*, *statum*, 'to stand.' Acme—é. *Granuleux du Rein*, Kidney, Bright's Disease of the.

ÉTERNUEMENT, [old F.] *esternuer*, from [L.] *sternutare*, 'to sneeze.' Sneezing.

ETESTICULATIO, (e, and *testiculus*, 'a testicle.') Castration.

ETHEMOSYNE, (εθμοσυνη,) Habit.

ETHER, *Æther*, *Æther sulphuricus*—e. Acetic, see *Æther*—e. *Acétique*, see *Æther sulphuricus*—e. Azotew, see *Æther sulphuricus*—e. *Azotew alcoholisé*, *Spiritus ætheris nitrici*.

ETHER CHLORIC, see *Chloroform*. Under the names *concentrated chloric ether* and *strong chloric ether*—a compound of pure *chloroform* and nearly absolute *alcohol*, in the proportion of one-third of the former to two-thirds of the latter—has been used as an anæsthetic by inhalation in the same cases as *sulphuric ether* and *chloroform*. It is properly an *alcoholic solution* or *tincture of chloroform*. See *Spiritus Chloroformi*.

ETHER, CHLOROHYDRIC, CHLORINATED, see *Æther muriaticus*—é. *Chlorohydrique*, see *Æther sulphuricus*—é. *Chlorohydrique chloré*, see *Æther muriaticus*.

ETHER, COM'FOUND. A preparation consisting of a solution of *chloroform* in *sulphuric ether*, which has been used as an anæsthetic by inhalation, in the same cases as *chloroform*.

ETHER, HYDRATIQUE, *Æther sulphuricus*—e. Hydriodic, *Æther hydriodius*—é. *Hydrique alcoolisé*, *Spiritus ætheris sulphurici compositus*—e. Hydrochloric, see *Æther*—e. Hydrochloric, spirit of, *Spiritus ætheris muriatici*—e. Hydrocyanic, *Æther hydrocyanicus*—e. Iodhydric, *Æther hydriodicus*—e. Marine, *Æther muriaticus*—e. Muriatic, *Æther muriaticus*—e. Muriatic, spirit of, *Spiritus ætheris muriatici*—é. *Nitreux*, see *Æther sulphuricus*—e. Nitric, see *Æther*—e. *Nitrique*, see *Æther sulphuricus*—e. Nitrous, see *Æther*—e. Phosphoric, see *Æther*—e. Prussic, *Æther hydrocyanicus*—e. Rectified, *Æther sulphuricus*—e. Stronger, see *Æther sulphuricus*—é. *Sulfurique*, *Æther sulphuricus*—é. *Sulfurique al-*

coolisé, *Spiritus ætheris sulphurici compositus*—e. *Sulphuric*, *Æther sulphuricus*—e. *Terebinthinated*, *Æther terebinthinatus*—é. *Vitriolique*, *Æther sulphuricus*.

ÉTHÉRA *Éthérolat*.

ÉTHÉRÉ, *Æthereal*.

ETHEREAL, *Æthereal*.

ETHEREOUS, *Æthereal*.

ETHERINE, HYDROCYANATE OF, *Æther hydrocyanicus*—e. *Muriate of*, *Æther muriaticus*.

ÉTHÉRISÉ, *Etherized*.

ETHERISM. The aggregate of phenomena induced by the inhalation of ether—*Ætherism'us*.

ETHERIZATION, *Ætherisat'io*. The administration of ether by inhalation, as an anæsthetic.

ETHERIZED, *Ætherisat'us*, (F.) *Éthérisé*. Presenting the phenomena induced by the inhalation of ether.

ÉTHÉROLAT, *Éthérat*. The product of the distillation of ether on aromatic substances.

ÉTHÉROLATURE, see *Tincture*.

ÉTHÉROLÉ. A liquid medicine formed of ether and medicinal principles united with it by direct solution or simple mixture.

ÉTHÉROLÉ D'ESSENCE DE TÉRÉBINTHINE. See *Liquor anodynus terebinthinatus*.

ÉTHÉROLIQUE, *Etherol'ic*. A medicine which has ether for excipient.

ÉTHÉROLOTIF. An etherolic medicine which is exclusively employed externally.

ETHICS, MEDICAL, (ἠθός, 'moral conduct.') Deontology, medical.

ETHIK, Hectic.

ETHIOPIAN, see *Homo*.

ETHIOPS, see *Æthiops*.

ETHISIS, (ἠθισις, from ἠσσω, 'to sift.') Filtration.

ETHISMUS, (from ἠθμος, 'a sieve.') Filtration.

ETH'MO, (ἠθμος, 'a sieve,') in composition, a sieve; also, the ethmoid bone.

ETHMOCÉPHALE, *Ethmoceph'alus*; (*ethmo*, 'the ethmoidal portion of the nose, and κεφαλή, 'head,') A genus of monsters, with eyes close together, but distinct; the nasal apparatus atrophied and rudimental.

ETH'MOID, *Ethmo'des*, *Ethmoï'des*, *Ethmoï'dens*, (*ethmo*, and εἶδος, 'form,') Shaped like a sieve.

ETHMOID BONE, *Os ethmoïdeum seu multifor'me seu spongio'eum seu spongoides seu cribro'sum seu cribrifor'me seu cul'bicum seu cristat'um seu 'coram'inulent'um seu colifor'me seu colato'rium*, (F.) *Os criblenx*. One of the eight bones which compose the cranium; so called, because its upper plate is pierced by a considerable number of holes. The ethmoid bone is situate at the anterior, inferior, and middle part of the base of the cranium. It seems to be composed of a multitude of thin, fragile, semi-transparent laminae, which form more or less spacious cells, called the *Eth'moid'al lab'yrinth* or *cells*, distinguished into *anterior* and *posterior*. These communicate with the nose, and are lined by a prolongation of the pituitary membrane. The ethmoid bone is constituted of compact tissue, and is surrounded by the *Eth'moid al'ature*. It is articulated with the *frontal* and *sphenoidal bones*, the *cornua sphenoidal'ia*, the *superior maxillary bones*, the *palate bones*, the *ossa turbinata inferiora*, the *coner*, the *proper bones of the nose*, and the *lacrimal bones*.

ETHMOID'AL, *Ethmoïda'lis*. Belonging to the ethmoid bone; as, *Ethmoid'al cells*, *E. al'ature*, &c. The *Ethmoid'al apoph'ysis* or *proc'ess* is the advanced part of the sphenoid bone, which articulates with the ethmoid. The *ethmoid'al ar'teries* are two in number, the anterior of which arises from the ophthalmic artery. The origin of the

other varies. The *Échancrure ethmoïdale* of the os frontis receives the ethmoid bones. The *Ethmoïdal veins* correspond to the arteries.

ETHMOSE, (ἠθμός, 'a sieve.') Cellular tissue.

ETHIMYPHE, (ἠθμος, 'a sieve,' and ἦψη, 'tissue.') Cellular tissue.

ETHIMYPH'TIS, *Inflammat'io te'læ cellul'osæ, Cellulit'is*, (*ethmyphæ*, and *itis*.) Inflammation of the cellular membrane.

ETHIMYPHOTYLOSIS, (*ethmyphæ*, and τυλώσις, 'hardness.') Induration of the cellular tissue.

ETHNOG'RAPHY, *Ethnograph'ia*; (ἔθνος, 'nation,' and γράφω, 'a description.') A history of the races and families of man.

ETHNOLOGY, *Ethnology'ia*, (ἔθνος, 'nation,' and λόγος, 'a discourse.') The science of the races and families of man.

ETHIOS, (ἔθος.) Habit.

ETHUSE, *Æthusa cynapium*—*É.Méum*, *Æthusa menum*.

ETHYLE, CHLORIDE OF, *Æther muriaticus*—*e*. Cyanuret of, *Æther hydrocyanicus*—*e*. Iodide of, *Æther hydriodicus*.

ETICK, Hectic.

ETIOLA'TION, *Blanch'ing*, (F.) *Étiolément*. That state of plants which occurs in consequence of privation of light, when they become pale and watery. In *pathology*, it is sometimes used to denote the paleness produced in those persons who have been kept long without light; or a similar paleness, the result of chronic disease.

ÉTIOLEMENT, Etiolation.

ETIOLOGY, *Ætiology*.

ÉTIQUE, Hectic.

ETIQUETTE, MEDICAL. Deontology, medical.

ÉTISE, Hecticis—*É. Mésentérique*, *Tabes menterica*.

ETITA. A singular cutaneous affection described by Du Chaillu as existing amongst the Ashira, in Africa. It consists in the rapid appearance of blotches over the body, the parts being swollen, and as if they had been badly scalded. The skin is raised and a thin matter collects beneath it. It is attended with intolerable itching.

ETNAGHBERRIES, see *Juniperus communis*.

ÉTOILE, ([old F.] *estole*), Stella.

ÉTONNÉE, ([old F.] *estonnée*), Stunned.

ÉTOUFFEMENT, ([old F.] *estouffement*, from (G.) *stopfen*, 'to stop.') Suffocation.

ÉTOUPE, ([old F.] *estoupe*), Stupa.

ÉTOURDISSEMENT, Vertigo.

ÉTRANGLE-LOUP, ('strangle wolf') Paris.

ÉTRANGLEMENT, ([old F.] *étranglement*, from (L.) *strangulare*, 'to strangle.') Strangulation, Hysteria—*É. des Intestins*, *Entéremphraxis*.

ÉTRANGUILLON, (from *étrangler*, 'to strangle.') An ancient French appellation for a violent angina accompanied by symptoms of imminent asphyxia.

ÉTRIÉR, ([old F.] *estrier*, [L.] *strepæ*.) Stapes. Also, a bandage, *Huit du Cou-de-pied*, applied, after bleeding from the foot, to compress the sapheua.

ETRON, (ἡτρον,) Abdomen, Hypogastrium.

ETRON'CUS, (*etron*, and ογκος, 'a tumor.') A tumour of the lower belly.

ETTICK, Hectic fever.

ETILE, Urtica.

ÉTUI DE L'HIPPOCAMPE. A name given by Vicq d'Azyr to the upper part of the sphenoidal portion of the lateral ventricle of the brain, which is bounded by the lateral cornu of the corpus callosum.

ÉTUVE, ([old F.] *estuve*.) Stove—*e*. *Humide*, Bath, vapour.

ÉTUVER, to Foment. See Stupa.

EU, (ευ,) 'good, proper,' when prefixed to words.

EUÉMIA, (eu, and ἄμα, 'blood.') A good condition of the blood.

EUÆSTHES'IA, (eu, and αἰσθησις, 'perception.') Vigorous perception. A good condition of the perceptive faculties.

EUÆSTHET'US, (ευαισθητος.) One whose senses are in full vigor.

EUANALEP'SIS, (eu, and αναληψις, 'recovery.') Rapid restoration to strength.

EUNTHEMON, *Anthemis nobilis*.

EUCALYP'TUS MANNIF'ERA, (eu, and καλυπτος, 'covered,' the flower by the calyx.) From this Australian species, *Order*, Myrtaceæ, a manna-like substance exudes, which has similar properties to manna.

EUCALYPTUS RESINIFERA, see Kino.

EUCHRŒA, (ευχροῖα, from eu, and χροῖα, 'colour.') A good or healthy colour of the skin. A good appearance of the surface.

EUCHYMIA, (ευχymia, from eu, and χυμος, 'juice.') A good state of the humours.

EUCRA'SIA, (ευκρασία, from eu, and κρασις, 'temperament.') A good temperament.

EUDIAPNEUS'TIA, (eu, and διαπνοη, (δια, and πνοη, 'breath;') 'one who breathes well.') Easy transpiration.

EUÉCTICA (medicina,) (ευεκτικός, (eu, and εχω, 'to hold,') 'conducive to health.') Gymnastics.

EUÉL'CES, (ευελκος, from eu, and ἔλκος, 'an ulcer.') One in whom wounds and ulcers are readily healed.

EUÉMES, *Euëm'etos*, (ευemes, from eu, and εμεω, 'I vomit.') That which readily excites vomiting. One who vomits with facility.

EUETHES, (eu, and εθος, 'habit,') Benign.

EUEX'IA, (ευexia, from eu, and ἔξις, 'constitution.') A good constitution.

EUFRAISE, *Euphrasia officinalis*.

EUGENIA ACRIS, see *Spiritus myrciæ*.

EUGENIA CARYOPHYLLA'TA, (after Prince Eugene of Savoy,) *Garryophyll'us*, *Caryophyll'us aromaticus*, *Myrt'us caryophyll'us*. The Indian tree which affords the clove. *Order*, Myrtaceæ. The *Clove*, *Caryophyll'us*, *C. aromaticus*, *Caryophyll'um*, is the unexpanded bud, (F.) *Girofle*, *Gérofle*, *Gyrophle*, *Clou de Girofle*. Its odour is strong, aromatic, and peculiar; taste, pungent and acrid. Like all substances whose virtue depends upon essential oil, it is stimulant and carminative. It is generally used as a corrigent to other remedies, and in cases where substances containing the essential oils are demanded. The oil—(F.) *Huile de Girofle*; *O'leum caryophyll'iti*, *Oil of Cloves*—has the properties of the cloves.

EUGENIA PIMENTA, *Myrtus pimenta*.

EUGION, and **EUGIUM**, (eu, and γη, 'earth,' ευγίος, 'fertile.') Hymen, Uterus, Vulva.

EU'LÊ, (εულê, 'a worm.') *Eulæ*, (ευλα,) with Hippocrates, meant especially worms bred in a wound or ulcer.

EULOPHIA, see Salep.

EUNUCH, *Eunuchus*, (ευνοχος, from ευνη, 'the bed,' and εχω, 'I keep.') *Hemian'drus*, *Hemia'nor*, *Hemianthro'pus*, *Sem'imus*, *Seminas'culus*, *Semimascula'tus*, *Sem'ivir*, *Thla'dias*, *Thla'nias*, *Thlib'ias*, *Castra'tus*, *Gal'tus*, *Eccastra'tus*, *Evira'tus*, *Emascula'tus*, *Excac'tus*, *Extesticula'tus*, *Extom'ius*, *Deli'acus*, (Old Eng.) *Gelding*. (F.) *Eunogue*. One whose organs of generation have been removed, or so altered, that he is rendered incapable of reproducing his species, or of exercising the act of venery. Eunuchs were common with the ancient Romans. With them, *Castra'tus* meant one from whom the external organs had been removed—the

Spada was deprived of the testicles only; whilst the *thio'sias* and *thla'sias* preserved the organs, but the testicles were crushed so as to render them unfit for procreation.

In Italy, this horrible mutilation still takes place to improve the voice; and in the East, eunuchs have the surveillance of the seraglio.

EUNUCHISMUS, Castration.

EUNUCHIUM MECONIS, Lactuca.

EUNUCHUS, Castratus, Eunuch.

EUNUQUE, Eunuch.

EUÔDES, (*εὐὼδης*, from *eu*, and *ὀζειν*, 'to smell.') Beneolentia.

EUONYMUS, (*eu*, and *ὄνυμα*, 'name.') Quassia simarouba.

EUONYMUS AMERICA'NUS, (*eu*, and *ὄνυμα*, 'name.') *Strawberry bush*, *Strawberry tree*, *Burn'ing bush*, *In'dian ar'row-wood*. A shrub of the *Ord. Celastraceæ*, *Sex. Syst.* Pentandria Monogynia, which, like the next, is found throughout the United States and Canada; flowering from May to June.

EUONYMUS ATROPURPUREUS, *Burn'ing bush*, *Spin'dle tree*, *In'dian Ar'row-wood*, flowers from June to July.

From these varieties of *Euonymus* the *Wahoo'* or *Whahoo'* bark is said to be obtained. In the *Pharmacopœia* of the United States, it is referred to *Euonymus atropurpureus*. It is diuretic, antiperiodic, tonic, and a hydragogue cathartic, and has been used in dropsy in infusion, in the proportion of an ounce to a pint of water.

EUPATHIA, (*ευπαθεια*, from *eu*, and *παθος*, 'affection.') A disposition for being affected by pain easily. Also, a good state of health—*Euphor'ia*.

EUPATOIRE D'AVICENNE, *Eupatorium cannabinum*—*e. Percefeuille*, *Eupatorium perfoliatum*.

EUPATORIUM, (after Mithridates *Eupator*.) *Eupatorium perfoliatum*.

EUPATORIUM AGERATOÏDES, *E. Urticifolium*, *Age'ratum altissimum*, *White Snake'root*; indigenous; flowering in August and September; has the same properties as *Eupatorium perfoliatum*.

EUPATORIUM AROMAT'ICUM, *Aromat'ic Eupatorium* or *Hemp'weed*, has similar properties.

EUPATORIUM CANNABINUM, *Eupatorium*, *E. Arab'icum* seu *Japon'icum* seu *trifoliatum*, *E. of Avicenn'na*, *Cannab'ina aquatica*, *Trifo'lium cervinum*, *Orig'anum aquaticum*, *Hemp ag'rimum*, *Herb of Saint Cu'negonde*, (F.) *Eupatoire d'Avicenne*. *Ord. Compositæ*. *Sex. Syst.* Syngenesia regularis. The juice of this plant proves violently emetic and purgative, if taken in sufficient quantity. It has been considered diuretic, cathartic, and emetic.

EUPATORIUM CONNATUM, *E. perfoliatum*—*e. Guaco*, *Guaco*—*e. Japonicum*, *E. Cannabinum*—*e. Mesues*, *Achillea ageratum*.

EUPATORIUM PERFOLIATUM, *E. Conna'tum*, *Bone'set*, *Thor'oughwort*, *Thor'oughstem*, *Thor'ough-wax*, *Veg'table ant'imony*, *Cross'wort*, *A'gueweed*, *F'erwerwort*, *In'dian sa'ge*, *Joe-pye*, *Te'azel*, *Sweat'ing plant*, (F.) *Eupatoire percefeuille*, is a plant which grows in low, wet meadows throughout the United States. The tops and leaves, gathered after flowering has commenced—*Eupatorium*, (Ph. U.S.)—are considered to be stimulating, tonic, emetic, purgative, diuretic, and sudorific. Dose, gr. xv of the powder, as a gentle purgative.

EUPATORIUM PURPLE-STALKED, *E. purpureum*. **EUPATORIUM PURPUREUM**, *Mo'hawk tas'sel*, *Pur'ple-stalk'ed Eupatorium*, *Trump'etweed*, *Joe Pye'weed*, *Grav'elroot*, is used in similar cases, as well as

EUPATORIUM ROTUNDIFOLIUM, *Round-leaved Hemp'weed*, *Wild hore'hound*;

EUPATORIUM SESSILIFOLIUM, *Up'land Bone'set*, and

EUPATORIUM TEUCRIFOLIUM, *Wild hore'hound*, *German'der-leav'ed hore'hound*, *Rough bone'set*.

EUPATORIUM TRIFOLIATUM, *E. Cannabinum*—*e. Urticifolium*, *E. ageratoïdes*.

EUPEP'SIA, (*ευπεσια*, from *eu*, and *πεπω*, 'I digest.') A good digestion. The antithesis to *dyspepsia*.

EUPEP'TIC, *Eupep'tus*, (*ευπεπτος*,) *Eupep'ticus*. Relating to a good digestion. One endowed with a good digestion.

EUPHLO'GIA, (*eu*, and *φλεγω*, 'I burn.') Mild inflammation.

EUPHONY, *Eupho'nia*, (*ευφωνια*, from *eu*, and *φωνη*, 'voice.') An agreeable or regular voice.

EUPHORBIA CYPRESS, *E. cyparissias*—*e. des Marais*, *E. palustris*—*e. Vomitive*, *E. ipecacuanha*.

EUPHORBIA CAPITATA, *Cacicia*. *Ord. Euphorbiaceæ*, (after Euphorbus, physician to Juba.) A Brazilian plant, which is strongly astringent, and not poisonous. It is considered to be one of the best remedies for the bites of serpents. It is, also, called *Colubri'na Lusitanica*.

EUPHORBIA CA'PUT MEDU'SÆ. A South African plant, with which, according to Bruce, the wild inhabitants poison their arrows. The Ethiopians use, for the same purpose, *Euphor'bia heptag'ona*, *E. viro'sa*, and *E. cereiform'is*; the savages, in the most southern part of America, the sap of *E. cotinifolia*.

EUPHORBIA CEREIFORMIS, see *Euphorbia caput Medusæ*.

EUPHORBIA COROLLA'TA, *Large flow'ering spurge*, *Milk'weed*, *Snakes' milk*, *Ipecacuan'ha*, *Hip'po*, *Picac'*, *Ip'cac'*, *Milk pur'slain*, *Purge'root*, *Emet'ic root*, *Bone'man's root*, *Ap'ple root*, *In'dian Phys'ic*; indigenous; has similar properties to the last. The dose of the root is the same.

EUPHORBIA COTINIFOLIA, see *Euphorbia caput Medusæ*.

EUPHORBIA CYPARIS SIAS, *E. Cypress'ina*, *Es'ula mi'nor* seu *Cypariss'ias*, *Tithym'alus Cypariss'ias*, *Cy'press spurge*, *Wel'come to our house*, (F.) *Euphorbe cypres*. This, like most of the spurges, is very acrimonious. Amongst the rustics, it was formerly called *poor man's rhubarb*, and was, consequently, a laxative. It is not used.

EUPHORBIA CYPRESSINA, *E. Cyparissias*.

EUPHORBIA HELIOSCOP'IA, *E. obtusa'ta*, *Sum'spurge*, *Wart'wort spurge*; indigenous. The juice has been given in syphilis. It is said to be cathartic.

EUPHORBIA HEPTAGONA, see *Euphorbia caput Medusæ*.

EUPHORBIA HYPERICIFOLIA, *Large spot'ted spurge*, *Eye'bright*. A native of the United States. It is astringent and tonic. Used in infusion—f℥ss to Oj of boiling water. Dose, a tablespoonful. **EUPHORBIA MACULA'TA**, *Milk Pur'slane*, *Spot'ted spurge*, has similar properties.

EUPHORBIA IPECACUAN'HA, *Anisophyl'um Ipecacuanha*, *Ipecacuan'ha spurge*, *Wild ip'cac'*. (F.) *Euphorbe vomitive*. This species of spurge is common in the southern and middle parts of the United States. The root is a powerful emetic, in the dose of from five to fifteen grains: twenty grains act as a cathartic likewise. In large doses, it acts as a narcotico-acrid.

EUPHORBIA LATH'YRIS. The systematic name of the plant which affords the *les'ser catapu'tia seeds*, *Catapu'tia mi'nor*, *Ca'per Spurge*, *Lath'yris*, *Gar'den spurge*, *Ca'per plant*, *Mole'plant*, *Tithym'alus latifol'ius* seu *lath'yris*, *Galarrhæ'us Lath'yris*, (F.) *Épurge* ou *grande Ésule*. The seeds—(F.) *Graines de Catepue* ou *d'Épurge* ou *Royales mineures*—possess cathartic properties; and an expressed oil of the seeds—*O'leum Euphor'bie Lathyr'idia*, (F.) *Huile d'épurge*—has been given as a cathartic in the dose of six to twelve drops.

EUPHORBIA MACULATA, see *E. hypericifolia*—

c. Myrtifolia, Pedilanthus tithymaloides—e. Ob-turata, E. helioscopia.

EUPHORBIA OFFICINARUM. The systematic name of the plant which affords the *Euphorbium*, *Euphorbia gummi-resi'na*, *Gummi euphor'bium*, in greatest abundance. The euphorbium is an in-odorous gum-resin, in yellow tears, which have the appearance of being worm-eaten. It enters into the composition of some plasters, and has been used as an emmenagogue.

EUPHORBIA PALUS'TRIS, *E. panicula'ta*, *Great'er spurge*, *Es'ula ma'jor*, *Tithym'alus palus'tris*, *Galarrhæ'us palus'tris*, *Marsh spurge*, *Wret'weed*, (F.) *Euphorbe des marais*. The juice is given, in Russia, as a common purge. It is used, too, as an irritant in tinea, warts, &c.

EUPHORBIA PALUSTRIS and **E. VILLO'SA**, or **E. PILO'SA**, have been brought forward as preven-tives of hydrophobia—the bitten part being washed with a decoction, and, at the same time, the decoction being taken internally.

EUPHORBIA PANICULATA, *E. palustris*.

EUPHORBIA PARAL'IAS, *Tithym'alus paral'ias*, *Sea spurge*. This is violently cathartic and irri-tating, and is seldom used.

EUPHORBIA PEPLUS, *Pell'ty spurge*: indigenous. The powdered root is actively cathartic. It has been given in drops.

EUPHORBIA PILOSA, see *Euphorbia palustris*—e. *Tithymaloides*, *Pedilanthus tithymaloides*—e. *Villosa*, see *Euphorbia palustris*—e. *Virosa*, see *Euphorbia caput Medusæ*.

All the spurge are vesicant and rubefa-cient, when applied externally.

EUPHORBUM, *Euphorbia officinarum*.

EUPHORIA, (*ευφορία*, from *eu*, and *φέρω*, 'I carry,') *Eupathia*.

EUPHRAGIA, *Euphrasia officinalis*.

EUPHRA'SIA, (*ευφρασία*, 'cheerfulness; ' so called from its effects,) *E. Officina'tis* seu *min'ima* seu *imbrica'ta*, *Ocula'ria*, *Euphra'gia*, *Ophthal'mica*, *Eyebright*, (F.) *Euphrase*, *Euphrase*, *Casse-Lunette*. It has been recommended in diseases of the eye, but is unworthy of notice.

EUPLASTIC, *Euplast'icus*, (F.) *Euplastique*: (*eu*, and *πλάσσω*, 'I form,') Having the capacity of becoming organizable in a high degree,—as in false membranes resulting from acute inflamma-tion in a healthy person.

EUPLEURUM SERRULATUM, *Diosma crenata*.

EUPNŒ'A, (*ευπνοία*, from *eu*, and *πνέω*, 'I re-spire,') Freedom or facility of respiration.

EURHYTH'MIA, *Eurhyth'mia*, (*ευρύθμια*, from *eu*, and *ῥυθμός*, 'rhythm,') Regularity of the pulse.

EURIBAL, *Juribali*.

EURODES, (*ευρώδης*, from *eurus*, and *odes*,) *Carious*.

EURODON'TICUS, (*ευρώς*, 'putrefaction,' and *ὀδούς*, gen. *ὀδοντός*, 'a tooth,') Suffering from carious teeth.

EURRYTH'MIA, *Eurhythmia*.

EURUS, (*ευρος*, 'mould, putrefaction,') *Mu'cor*, *Ca'ries*. Corruption of the humours.

EURYCHO'RIA, (*ευρύς*, 'extensive,' and *χωρός*, 'space,') *Sinus*.

EURYCLES, (*Ευρυκλής*, a celebrated soothsayer and ventriloquist,) *Engastrimyth*.

EURYCLITUS, *Engastrimyth*.

EURYSMUS, (*ευρυσμεν*, 'to dilate,') *Dilatation*.

EUSAR'CUS, (*ευσάρκος*, from *eu*, and *σάρξ*, gen. *σαρκος*, 'flesh,') One who is fleshy, robust, mus-cular.

EUSEM'IA, (F.) *Eusémië*, (*ευσεμεία*, from *eu*, and *σημείον*, 'a sign,') A collection of good signs.

EUSOMPHALIEN, (*eu*, and *ομφαλός*, 'navel,') A double monster in which each being has a separate navel, and is able to execute, independ-ently, almost all the vital functions.

EUSPLANCH'NIA, (*eu*, and *σπλάχνον*, 'a vis-cus,') A healthy state of the viscera.

EUSTA'CHIAN TUBE, *Tu'ba Eustachia'na* seu *Eusta'chii* seu *Aristotel'ica*, *Syrin'ga*, *Syr'inx*, *Meat'us ear'eus*, *Duct'us aur'is palati'nus*, *Pter a Palat'o ad Aur'em*, (F.) *Trompe* ou *Conduit d'Eus-tache*, *Conduit guttural de l'oreille*, (Ch.,) (called after Eustachius.) It is partly bony and partly cartilaginous: extending from the cavity of the tympanum to the upper part of the pharynx. Its length is about two inches; the bony portion, which belongs to the temporal bone, is about three-fourths of an inch long. It is lined, inter-nally, by a prolongation of the lining membrane of the pharynx. Its nerves are furnished by the palatine branches of the ganglion of Meckel, and its vessels proceed from those of the pharynx and velum pendulum. The use of the tube seems to be to permit the renewal of air in the cavity of the tympanum.

EUSTA'CHIAN VALVE, *Valve of Eusta'chius*, *Valvule d'Eustachi* ou *d'Eustache*. A membranous, semilunar fold, which corresponds to the opening of the vena cava inferior into the right auricle of the heart.

EUSTHEN'IA, *Vig'or*, *Exuberan'tia*, (*ευσθενεία*, from *eu*, and *σθένος*, 'strength,') Flourishing, exuberant health.

EUSTOM'ACHUS, (*ευστομαχος*, from *eu*, and *στομαχος*, 'stomach,') Digesting rapidly. Having a good stomach.

EUSTRONGYLUS GIGAS, *Strongylus gigas*.

EUTAX'IA, (*ευταξία*, from *eu*, and *τάσσω*, 'I arrange,') *Euthe'sia*. A well-ordered consti-tution, in which every part has its proper relation. The ready return of a rupture, or of a luxated bone.

EUTHANA'SIA, (*ευθανασία*, from *eu*, and *θανά-tos*, 'death,') An easy death. The opposite to this is *Cacothana'sia*.

EUTHESIA, (*ευσθεσία*, from *eu*, and *σθεσις*, 'po-sition,') *Eutaxia*.

EUTHYENTERON, (*ευθύς*, 'straight,' and *εν-τερον*, 'intestine,') *Rectum*.

EUTHYENTEROSTENOMA, (*euthyenteron*, and *στενός*, 'narrow,') Stricture of the rectum.

EUTHYM'IA, *An'imi tranquil'itas*, (*ευθυμία*, from *eu*, and *θυμός*, 'mind,') Tranquillity of mind. A good state of the mental faculties.

EUTHYPNOE, (*ευθύς*, 'straight,' and *πνοή*, 'breath,') *Orthopnoea*.

EUTHYPNŒA, *Orthopnoea*.

EUTOC'IA, (*ευτοκία*, from *eu*, and *τοκος*, 'de-livery,') An easy labour. Fecundity.

EUTROPH'IA, (*ευτροφία*, from *eu*, and *τροφή*, 'nourishment,') A good state of nutrition.

EUTROPH'IC, *Eutroph'icus*. A term intro-duced into medical terminology, by the author, for an agent whose action is exerted on the sys-tem of nutrition, without necessarily occasioning manifest increase of any of the secretions. The chief eutrophics are,—mercurials, the preparations of iodine, bromine, cod liver oil, the preparations of gold and silver, sulphur, sugar, and sarsapa-rilla.

EUZOODYNAMIA, (*eu*, *ζωον*, 'animal,' and *δυναμς*, 'power,') *Sanitas*.

EVAC'UANTS, *Evacuanti'a*, (*e*, and *vacuare*, 'to empty,') (F.) *Evacuati'fs*. Medicines are so called which occasion a discharge by some emunc-tory: such as purgatives, emetics, &c.

EVACUAT'IFES, *Evacuants*.

EVACUATIO, *Evacuation*, *Excretion*.

EVACUATION, *Evacuati'o*, *Lapax'is*. The discharge of any matter whatever by the natural passages or by an artificial opening. See *Dejec-tion*.

EVANOUISSEMENT, (from *evanescere*, (*e*, and *vanescere*,) 'to fade away,') *Syncope*.

EVAN ROOT, Geum Virginianum.

EVAPORATION, *Evaporatio*, *Vapora'tio*, *Echala'tio*, (e, and *evaporare*, *vaporatum*, 'to emit a vapour.') Transformation of a liquid into vapour, in order to obtain the fixed matters contained in it dry and separate from the liquid. When the vapour is received into a proper vessel and condensed, the process is called distillation. Evaporation produces cold, and this is one of the processes by which the body is cooled, through the evaporation of the perspiratory fluid.

ÉVAUX, MIN'ERAL WATERS OF. *Evaux* is situate in Auvergne, in France. The waters are hydrosulphurous and thermal. Season, from the 15th of May to the 1st of October.

EVENTRATION, *Eeventra'tio*, *Hypogastror-rhex'is*, (c, and *venter*, 'the belly.') A tumour, formed by a general relaxation of the parietes of the abdomen, and containing a great part of the abdominal viscera. Also, *ventral hernia*, or that which occurs in any other way than through the natural openings of the abdominal parietes. Lastly, any very extensive wound of the abdominal parietes, with issue of the greater part of the intestines.

EVENTUS, (e, and *venire*, *ventum*, 'to come.') *Termina'tio mor'bi*. The event or issue of a disease, either favourable or unfavourable.

EVERLASTING, DICICIOUS, *Antennaria dioica*.

EVERRICULUM, *Specil'lum*, (e, and *ververe*, 'to clean out.') A sort of sound or scoop, used for extracting sand or fragments of stone or clots of blood from the bladder, after or during the operation of lithotomy.

EVERSIO PALPEBRÆ, (e, and *vertere*, *ver-sum*, 'to turn,') *Ectropion*.

ÉVIGILATION, ([L.] *evigilare*, *evigilatum*, 'to awake,') see Sleep.

EVIL, THE, Scrofula—e. King's, Scrofula.

EVIRATIO, (e, and *vir*, 'a man,') Castration.

EVIRATUS, Castratus, Eunuch.

EVISCERATION, (e, and *viscus*, gen. *visceris*,) *Exenterismus*.

EVOLUTIO, (e, and *volvere*, *volutum*, 'to roll,') Development—e. Spontanea, Evolution, spontaneous.

EVOLUTION OF GERMS, see Generation.

EVOLU'TION, SPONTA'NEOUS, *Ver'sio* seu *Evolu'tio sponta'nea*, *Sponta'neous ver'sion*, *Sponta'neous expul'sion*, *Duplica'tion of the fœ'tus*. A term, applied, by Dr. Denman, to what he considered to be a spontaneous turning of the fœtus in utero, in an arm presentation, in consequence of powerful uterine contractions forcing out the breech and feet, whilst the arm receded. It is now usually considered to be a doubling of the fœtus, so that the arm changes its position but little, whilst the breech is forcibly expelled before the upper extremity;—the case becoming similar to a breech presentation.

EVOLUTION, VESICLE OF, Vesicle of development.

EVOMITIO, (e, and *vomitio*,) Vomiting.

EVONYMUS, Euonymus.

EVULSION, *Avul'sion*, *Evul'sio*, (*evellere*, *evulsum*, (e, and *vellere*,) 'to pluck out,') *Apotil'mus*, *Ectil'mus*, *Til'mus*. The action of plucking out; forcible extraction. (F.) *Arrachement*. This word is oftener used by the French than by the English surgeons, for the act of extracting certain parts, the presence of which is injurious,—as the teeth, &c.

EW-GOWAN, or *Ewe-Gowan*, ([Sc.] *gowan*, 'a daisy,') *Bellis*.

EX, as a prefix, out of. See *Ec*.

EX MARIBUS, (ex, and *mas*, gen. *maris*, 'a male,') Castratus.

EXACERBATIO, (ex, and *acerbare*, 'to embitter,') Exacerbation, Paroxysm.

EXACERBATION. *Exacerba'tio*, *Exa'pera'tio*, *Exerescen'tia*. (F.) *Redoublement*. An increase in the symptoms of a disorder. Often used synonymously with paroxysm.

EXACUM SPICATUM, (ex, and *ago*, 'to drive,') from its supposed virtue as an expeller of poison,) *Coutoubæa alba*.

EXÆMATOSIS, (ex, *haima*, 'blood,' and the suffix *osis*,) *Hæmatisis*.

EXÆMIA, (ex, and *haima*, 'blood,') *Anæmia*.

EXÆMUS, *Exanguisus*.

EXÆ'RESIS, (*ἐκάρσις*, from *ex*, and *αἶρω*, 'I take away,') (F.) *Exérèse*. A surgical operation, which consists in drawing, extracting, or removing, from the human body, any thing that is useless, noxious, or extraneous. It is a generic term, which includes extraction, evulsion, evacuation, excision, ablation, amputation, &c.

EXÆRESIS, Extraction.

EXAG'GERATED, *Exaggera'tus*, (F.) *Exagéré*, (*exaggero*, (ex, and *aggeo*, 'I heap up,') 'I magnify,') Heightened. Increased by expression. An epithet for sounds heard on auscultation and percussion when much increased—*Hypereche'sis*, *Hypereche'ma*.

EXAG'GION, (*ἐξαγιον*,) *Exa'gium*. The sixth part of an ounce:—four scruples.

EXAL'MA, gen. *Exal'matis*, (*ἐξάλμα*, from *ex*, and *ἀλλομαι*, 'I leap,') *Exal'is*. Hippocrates calls thus the displacement of the vertebræ.

EXALSIS, (*ἐξάλσις*,) *Exalma*, Leap.

EXALTA'TION OF THE VITAL FORCES, *Exalta'tio vi'tium*, (*exalto*, (ex, and *alto*,) 'I raise,') This expression has been used, by modern pathologists, to designate a morbid increase in the action of organs, and partly that which occurs in an inflamed organ. Some use *exaltation of the vital forces*, and *inflammation*, synonymously.

EXAMBLOMA, gen. *Examblo'matis*, (*ἐξαμβλωμα*, from *ex*, and *αμβλωω*, 'I abort,') Abortion.

EXAMBLOSIS, (*ἐξαμβλωσις*,) Abortion.

EXA'MEN RIGOROSUM, ('a rigorous examination,') An examination of a candidate for the degree of Doctor of Medicine in the German Universities, which, like the *Tentamen medicum*, is conducted in Latin, and takes place before the medical faculty on all branches of medicine.

EXANASTOMOSIS, Anastomosis.

EXANASTROPHE, (ex, *ana*, and *στροφή*, 'I turn,') Convalescence.

EXANGI'A, (ex, and *αγγειον*, 'a vessel,') *Exange'i'a*. An enlargement or rupture of a blood-vessel, without external opening. A genus in the order *Dysthetica*, class *Hæmatica* of Good. It comprises *aneurism* and *varix*.

EXANGIA ANEURISMA, Aneurism.

EXAN'GUIOUS, *Exan'guine*, *Exsanguin'eus*, *Exæ'mus*, *Exsan'guis*, (F.) *Exsangue*, (ex, and *sanguis*, 'blood,') Seeming to be bloodless; as a female, who has suffered largely from uterine hemorrhage. See *Anæmie*.

EXANIA, (ex, and *anus*,) Proctocelæ.

EXANIMA'TION, *Exanima'tio*, (ex, and *anima*,) This word has two acceptations. Sometimes, it means real death, corresponding with the Greek *Σάρατος*, *mors*. At others, it signifies apparent death, corresponding with the Greek *ἀψυχία*, *ἐκψυχία*, *εκλυσίς*, *An'imi deli'quium*.

EXANTHEM, *Exanth'e'ma*, *Anthe'ma eruptio*, *Epanthe'ma*, *Ejflor'escentia*, *Exanthis'ma*, *Ec'thy-ma*, *Epanaste'ma*, *Epanthe'ma*, *Epanthis'ma*, *Ejflora'tio*, (*ἐξανθήμα*, from *ἐξ*, and *ανθῶ*, 'I bloom,') A rash. (F.) *Exanthème*, *Éléveure*. Under this term is comprehended, by some, every kind of eruption, of which the skin is the seat. Others understand by it those eruptions that are accompanied by fever, *Fe'bres exanthemat'icæ*; includ-

ing, under the head of the *major exanthemata*, those which attack a person but once in his life, and which are communicated by specific contagion: and, under the *minor exanthemata*, those which are not marked by these characteristics. Small-pox, measles, cow-pox, &c., belong to the major:—chicken-pox, herpes, lichen, &c., to the minor. The general acceptance of Exanthem is, however, a more or less vivid, circumscribed, or diffuse redness of the skin, which diminishes, or disappears transiently, under the pressure of the finger.

EXANTHEM. CARBUNCULAR, Anthracia—e. Ichorous, Emphylys—e. Pustulous, Empyesis—e. Rash, Eanthesis.

EXANTHEMA, gen. *Exanthem'atis*, Eminence, Exanthem—e. Antivariolosum, Vaccina—e. Inter-
num, Eisanthema.

EXANTHEMA IOD'ICUM. An eruption of dark red definite spots, of various sizes, spreading over the whole body, without the formation of scales, and disappearing only after a long time, and which seems to be produced occasionally by the use of iodine.

EXANTHEMA MILIARIA, Miliary fever—e. Pestis, Plague—e. Serosum, Pemphigus—e. Strophulus, Strophulus—e. Urticatum, Urticaria—e. Vaccina, Vaccina—e. Variellata, Variella.

EXANTHEMATICA, *Eruptive fevers*. Cutaneous eruptions, essentially accompanied with fever;—the third order in the class *Hemastica* of Good.

EXANTHEMATIS/CHESIS, *Exanthematos'-chesis*, (*exanthem*, and *αχειν*, 'to withhold,') Suppression of a cutaneous eruption.

EXANTHEMATOLOG'Y, *Exanthematolog'ia*, (*exanthem*, and *λογος*, 'a discourse,') The doctrine of cutaneous eruptions.

EXANTHEMATOPHTHAL'MIA, *Ophthalm'ia exanthemat'ica*, (*exanthem*, and *οφθαλμια*, 'inflammation of the eye,') Ophthalmia in the course of, or succeeding to, a cutaneous eruption.

EXANTHEMATOUS, *Exanthematos'us*, *Exanthemat'icus*, (F.) *Exanthématique*. Relating or appertaining to exanthem, as "an *exanthematous eruption*," (F.) *Éruption exanthématique*.

EXANTHEME INTESTINALE, see Typhus.

EXANTHESIS, (*εξανthesis*,) see Efflorescence—e. Arthrosia, Dengue—e. Roseola, Roseola—e. Rubeola, Rubeola.

EXANTHISMA, (*εξανθισμα*,) Exanthem.

EXANTHROPIA, (*ex*, and *ανθρωπος*, 'man,') Misanthropia.

EXAPSIS, (*εξαψις*, from *ex*, and *απειν*, 'to inflame,') Inflammation.

EXARAG'MA, (*εξαργμα*, from *εξαρασσω*, 'I tear away,') 'I break,') Collision, violent fracture, a fragment.

EXARCHIA'TER, *Exarchiat'ros*. Chief of the *archiatri*, or chief of physicians, a title, like that of *Comes Archiatro'rum*, given to the chief physician of an emperor or king. Archiater appears to have meant, at times, the same thing.

EXARMA, gen. *Exar'matis*, (*εαρμα*, from *ex*, and *αρειν*, 'to raise up,') Swelling.

EXAR'SIO, (*ex*, and *ardeo*, *arsum*, 'to burn,') A burning heat.

EXARTEMA, gen. *Exarte'matis*, (*ex*, and *αραιω*, 'I suspend,') Amuletum.

EXARTERITIS, *Exarteri'tis*, (*ex*, and *arteria*, 'an artery,') and *itis*,) Inflammation of the outer coat of an artery.

EARTHREMA, gen. *Exarthre'matis*, (*εαρρημα*, from *ex*, and *αρρηρον*, 'a joint,') Luxation.

EXARTHROMA, gen. *Exarthro'matis*, (*εαρρρωμα*,) Luxation.

EARTHROSIS, (*εαρρρωσις*,) Luxation—e. Paretica, see Pareticus.

EXARTICULATIO, (*ex*, and *articulus*, 'a joint,') Amputation, joint. Luxation.

EXARYSIS, (*εξαρυσις*,) Exhaustion.

EXASPERATIO, (*ex*, and *asperare*, *asperatum*, 'to sharpen,') Exacerbation.

EXIBITION, see Imbibition.

EXCÆCARIA AGALLOCHA, (*εκακαρια*, 'to make blind,') its acrid juice, Agallochum.

EXCALEFACTIO, (*ex*, *calidus*, 'warm,') and *facere*, 'to make,') *Échauffement*.

EXCARNA'TION, *Excarna'tio*, (*ex*, and *caro*, gen. *carnis*, 'flesh,') A mode of making anatomical preparations, which consists in separating injected vessels from the parts in which they are situate. This is done by means of corrosion by an acid or by putrefaction.

EXCASTRATIO, Castration, Eunuch.

EXCATHISMA, gen. *Excathis'matis*, (*ex*, and *καθισμα*, 'a seat,') Bath, half; Semicupium.

EXCAVATOR DENTAL, (*ex*, and *cavare*, *ca-vatum*, 'to make hollow,') see Dental.

EXCEREBRATIO, (*ex*, and *cerebrum*, 'brain,') Eccephalosis.

EXCERBRATUS, Delirious.

EXCERNENT, (*ex*, and *cernere*, 'to sift,') Secreting.

EXCIDENTIA, (*ex*, and *cadere*, 'to fall,') Ecptoma.

EXCIP'IENT, *Excip'iens*, (*excipere*, (*ex*, and *capere*,) 'to receive,') (F.) *Intermède*. A substance, which, in a medicinal prescription, gives form and consistence to it, and serves as a *vehicle* or *medium* for the exhibition of the other ingredients.

EXCIPULUM, Cup.

EXCISIO, *Entaille*, Excision,

EXCIS'ION, *Excis'io*, (*excidere*, *excisum*, (*ex*, and *cadere*,) 'to cut off,') *Éctomè*. A surgical operation, by which parts of a small size are removed with a cutting instrument.

EXCITABIL'ITY, *Excitabil'itas*. Irritability. (From *excitare*, (*ex*, and *citare*, *citatum*,) 'to excite,') The faculty possessed by living beings, of being sensible to the action of excitants. The doctrine of excitability forms an important part of the Brunonian system: see Excitation.

EXCITANT, Stimulant.

EXCITA'TION, *Excita'tio*, *Excitement*. The act of exciting; the state of an organ or organs excited. Excitement is, sometimes, used synonymously with augmented arterial action. The effect of the exciting powers acting on the excitability, according to Brown, constitutes *excitement*. Cullen used the term to express the restoration of the energy and action of the brain, which had been interrupted by sleep or some debilitating cause,—a state opposite to that of *collapse*. Not unfrequently, it is employed in the sense of excessive action,—*Superexcita'tio*, (F.) *Sur-excitation*.

EXCITED DISEASES, Feigned diseases.

EXCITO-MOTION, see Excito-motory.

EXCI'TO or EX'CITO-MO'TORY, (F.) *Excito-moteur*. An epithet applied by Dr. Marshall Hall to a division of the nervous system—comprising the gray matter of the spinal marrow, with the afferent and efferent nerves connected with it;—all of which are concerned in *reflex* actions; or those by which impressions are transmitted to a centre, and reflected so as to produce muscular contraction without sensation or volition. They constitute the *reflex system of nerves*. See Nerves.

The term *excito-motion* has also been employed to signify motion, no matter how excited, by the reflex nerves or by volition.

Excitonutrient and *Excitosecretory* are reflex actions in which nutrition and secretion are modified in parts by impressions made elsewhere and reflected to them, as where tumours disappear or secretions of tears, milk, saliva, for example, are increased, by impressions made on the cerebro-spinal nervous centres.

EXCORIATIO, (ex, and corium, 'skin,') Ecdora, Excoriation.

EXCORIATION, *Excoria'tio*, *Excoriatu'ra*, *Am'yôchê* (F.) *Écorchure*. A slight wound, which removes only the skin.

EXCREATURA, Excretion.

EXCREATION, *Excrea'tio*, *Exscrea'tio*, *Screa'tus*, *Rasca'tio*, (ex, and screare, 'to spit,') Act of spitting. (F.) *Crachement*. See Exspuitio.

EXCREMENT, *Excrementum*, *Retrimen'tum*, *Excre'tum*, *Excre'tio*, *Perit'toma*, *Diachore'ma*, *Aph'odos*, *Aphodeu'ma*, *Apoch'o'reon*, *Apoc'risis*, *Arda*, *Ar'dalus*, (excernere, excretum, (ex, and cernere, 'to separate,' 'cleanse,') Everything, which is evacuated from the body of an animal by the natural emunctories as superfluous; such as the fecal matters, the urine, perspiration, nasal mucus, &c. Generally, however, the term is restricted to the fecal evacuations;—*Purgamen'ta*, *He'dra*, *Se'des*, *Fæ'ces*, *Ster'cus*, *Cac'cê*, *Spat'ilê*, (especially when liquid,) *Deje'ctio alvi'na*, *O'nera al'vi seu ven'tris*, *Sor'des ven'tris*, *Hypochore'sis*, *Cœ'lia*, *Hypochore'sis*, *Mer'da*, *Mer'dus*, *Catar-rhex'is*, *Co'prus*, *Co'pranon*, *Scor*.

EXCREMENT, HUMAN, *Stercus humanum*.

EXCREMENTITIAL, *Excrementit'ious*, *Excrementit'ius*, (F.) *Excrémentieux*, *Excrémentiel*. That which is similar to excrement, and forms part of it. Excrementitious humours or parts are those destined to be evacuated as incapable of administering to the nutrition of the body.

EXCREMENTTORECREMENTITIAL, *Excrementorecrementit'ious*. Animal fluids intended to be partly absorbed and partly rejected.

EXCRESCENCE, *Excresec'entia*, *Ephy'ma*, *Phymato'sis*, *Hypersarc'o'sis*, *Sarcophy'ia*, *Ca'ro excre'scens*, (excrecere, (ex, and crescere, 'to grow outwards,') (Sc.) *Squalme*. (F.) *Excroissance*. A tumour, which forms at the surface of organs, and especially on the skin, mucous membranes, or ulcerated surfaces. Excrecences differ in size, shape, cause, &c., and each requires its own treatment. Warts, *condylomata*, *polypi*, *hemorrhoids*, belong to this head.

EXCRESCENTIA, Exacerbation, Protuberance, Tumour—e. *Carnosa*, *Sarcoma*—e. *Fungosa*, *Fungosity*—e. *Gingivæ*, *Epulis*—e. *Vesicæ urinariæ carnosæ*, *Cysthypersarcosis*.

EXCRETA, see Excretion.

EXCRÉTEUR, Excretory.

EXCRETIO, Excretion, Excrement—e. *Alvina*, Defecation—e. *Fæcum alvinarum*, Defecation—e. *Urinæ involuntaria*, *Enuresis*.

EXCRETION, *Excre'tio*, *Ec'crisis*, *Evacua'tio*, *Eje'ctio*, *Ecpul'sio*, *Eges'tio*, *Diachore'sis*, (excernere, excretum, (ex, and cernere, 'to separate,') The separation or throwing off of those matters, *Excre'ta*, *Eges'ta*, *Ion'ta*, *Apion'ta*, from the body of an animal, which are supposed to be useless, as the urine, perspiration, and fæces.

EXCRÉTOIRE, Excretory.

EXCRETORY or EXCRETORY, *Excreto'rius*, (F.) *Excréteur* ou *Excrétoire*. An Excretory vessel or duct, *Ductus excreto'rius*, is one which transmits the fluid secreted by a gland either externally or into the reservoirs into which it has to be deposited. The existence of an excretory duct was regarded as a distinctive character of the glands properly so called.

EXCRETORY ORGAN means any one charged with the office of excreting; thus, the skin is said to be an excretory organ, because through it the perspiration takes place.

EXCRETUM, Excrement.

EXCROISSANCE, Excrescence.

EXCU'TIA VENTRICULI, (ex, and quatio, 'to shake,') *Stom'ach Brush*. An instrument, composed of iron or brass wire, at one of the extremities of which is a pencil of bristles. Some

ancient authors proposed this to extract foreign bodies from the œsophagus, as well as to cleanse the stomach of viscid and tenacious matters adhering to it.

EXECHEBRONCHUS, (ἐξεχειν, 'to project,' and βρογχος, 'the throat,') *Bronchocele*.

EXECHEGLUTI, (ἐξεχειν, 'to project,' and γλουτοι, 'the nates,') *Exischioi*.

EXEDENS, (ex, and edere, 'to eat,') see *Herpex exedens*.

EXELCOSIS, (ex, and ἔλκος, 'an ulcer,') see *Ulceration*.

EXELCYS'MUS, (ἐξελκυσμος, from ex, and ἔλκω, 'I draw,') *Extraction*. Also the act of breaking out into ulcers.

EXENCÉPHALE, (ex, and ἐγκεφαλον, 'the encephalon,') A genus of monsters, in which the encephalon is situate in a great measure out of the cranium, the upper portion of which is almost wholly wanting.

EXENTERATION, Exenterismus.

EXENTERIS'MUS, *Exenter'isis*, *Exentera'tio*, *Exenteration*, *Eviscera'tion*, *Deviscera'tion*, *Unbow'elling*, *Embow'elling* or *Imbow'elling*, *Viscera'tion*, (ex, and εντερον, 'an intestine,') The operation of taking out the bowels.

EXERA'MA, gen. *Exera'matis*, (ἐξεραια, from ex, and ερω, 'I pour out,') Any thing cast out. Vomiting; or the matter vomited.

EXERCISE, *Exercita'tio*, *Exercit'ium*, *Asce'sis*, *Gymna'dion*, (exercere, exercitum, 'to work,') Every motion of the body arising from the contraction of muscles subjected to the will. Also, the action of any organ whatever. Exercises may be active or passive. The passive are referred to under the head of Gestation. The chief active exercises are:—walking, running, dancing, hunting, fencing, playing at ball, cricket, racket, quoits, swimming, declamation, and singing. Exercise is an important prophylactic, particularly for those disposed to be plethoric. It improves the digestion; augments the secretions; and, when used in moderation, gives strength to the body; but when carried to excess, produces debility and disease.

EXÉRÈSE, Exæresis.

EXERHO'SIS, (ex, and ῥέω, 'I flow,') The discharge which takes place by insensible perspiration.

EXFETATION, see *Pregnancy*.

EXFOLIATIF, Exfoliative.

EXFOLIATION, *Exfolia'tio*, *Desquama'tio*, *Eclep'isis*, (ex, and folium, 'a leaf,') By this is meant the separation of the dead portions of a bone, tendon, aponeurosis, or cartilage, under the form of lamellæ or small scales. Exfoliation is accomplished by the instinctive action of the parts, and its object is to detach the dead portion from those subjacent, which are still alive. For this purpose the latter throw out fleshy granulations, and a more or less abundant suppuration occurs, which tends to separate the exfoliated part—now become an extraneous body. The ancients distinguished exfoliation into *sensible* and *insensible*, according as the dead portions of bone were detached in fragments of greater or less size, or in very thin pieces, and in an almost insensible manner. When the dead part embraces all or almost all the substance of a bone, it takes the name *Sequestrum*.

EXFO'LIATIVE, *Exfoliati'vus*, *Desquamato'rius*, (F.) *Exfoliatif*. That which takes away by leaves or scales. The term has been applied to certain medicines, which were regarded as proper to hasten exfoliation, such as alcohol, oil of turpentine, tincture of myrrh, &c.

EXFOLIATIVE TREPAN, *Eclepisitrep'anum*. An ancient raspatory, or instrument for scraping exfoliating portions of bone.

EXHALAISON, Exhalation.

EXHA'LANT, *Exha'lent, Exha'lans*, (*exhalare, exhalatum*, (*ex*, and *halare*, 'to breathe,') 'to exhale, throw out.')

EXHALANT VES'SELS, *Ves'sa exhalant'ia*, are very minute, and rise from the arterial capillary system. They are situate in every tissue of the body, and on the surface of the mucous and serous membranes and skin; on which each pours its particular fluid. Bichat distinguished three sets. 1. The *external*, terminating on the mucous and external dermoid system, where they pour the matter of perspiration. 2. The *internal*, comprising those of the areolar and medullary tissues, and of synovial surfaces; and, 3. The *nutritive exhalants*, which vary in each organ where they are found, and preside over the phenomena of composition and increase of every part of the body. The exhalants are the antagonists of the absorbents. They are imaginary, as a distinct set from the capillary vessels.

EXHALATIO, Evaporation, Exhalation.

EXHALATION, *Exhalatio*; (same etymon as Exhalant,) *Anathym'asia, Apopne'usis*, (F.) *Exhalaison*. A function, by virtue of which certain fluids, obtained from the blood, are spread, in the form of dew, in the areolæ of the different textures, or at the surface of membranes; either for the sake of being thrown out of the body, or to serve ulterior purposes. The sweat is a *liquid excrementitious exhalation*; the serous fluid of the pleura, a *liquid recrementitious exhalation*.

Exhalation is also applied to that which exhales from any body whatever, organic or inorganic, dead or living.

EXHALATION, PULMONARY, see Perspiration.

EXHAUST'ION, *Excar'ysis, Vires exhaust'æ*, (*exhaustire, exhaustum*, (*ex*, and *haurire*,) 'to draw out.')

(F.) *Épuisement*. Loss of strength, occasioned by excessive evacuations, great fatigue or privation of food, or by disease.

EXHIL'ARANT, *Exhil'arans*, (*ex*, and *hilaro*, 'I make merry.')

An agent that exhilarates or enlivens.

EXHUMA'TION, *Exhuma'tio, Effos'sio*, (*ex*, and *humus*, 'the ground.')

The disinterment of a corpse. The circumstances which render this necessary are:—1. Judicial investigations relative to the body of the person inhumed. 2. The removal of a body from one cemetery to another; and, 3. The evacuation of cemeteries or sepulchral vaults. The operation is attended with much unpleasant smell and annoyance, and requires the use of disinfecting agents, of which one of the most available is chlorinated lime. See Disinfection. The putrid effluvia from animal substances are not, however, found to excite endemic disease.

EXIDIA AURICULA JUDÆ, *Peziza auricula*.

EXINANIT'ION, *Exinanit'io*, (*ex*, and *inanis*, 'empty.')

Extreme exhaustion; complete evacuation.

EXIS'CHIUM, (*ex*, and *ischion*, 'the ischium.')

Prominence of the hips.

EXIS'CHIUS, (*ἐξίσχυος*) *Exis'chus*. A luxation of the os femoris. Those with large nates, and prominent hips, were formerly called *Exis'chioi* and *Exechegh'iti*.

EXITU'RA, (*exire, exitum*, (*ex*, and *eo*,) 'to go out.')

According to some, any abscess which discharges. Paracelsus calls thus every kind of putrid excrement.

EXITUS, gen. *Ex'itūs*. The outer termination or exit of a canal. The termination of a disease.

EXO, (*ἐξω*,) in composition, without, outwards.

EXOARTERITIS, see Arteritis.

EXOCARDIAC, see Endocardiac.

EXOCARDIAL, see Endocardiac.

EXOCARDITIS, Pericarditis.

EX'OCHAS, *Ex'ochē*, (*ἐξοχος*, from *ἐξω*, (*ἐξ*, and *χω*,) 'I project.')

A soft tumour—as a hemorrhoid—outside of the anus. An outward pile.

EXOCHE, (*ἐξοχη*,) Eminence, Exochas.

EXOCHORION, see Chorion.

EXOCOLITIS, see Colitis.

EXOCULATIO, (*ex*, and *oculus*, 'an eye.')

Want of eyes. Want of vision. Blindness.

EXOCYST'Ē, *Exocyst'ia*, (*exo*, and *κυστις*, 'the bladder.')

A prolapsus of the bladder into the urethra. Also called *Prolap'sus vesic'æ*, *Ædopto'sis vesic'æ*, (F.) *Renversement de la vessie*.

EXOCYST'Ē NOELIA'NA. Protrusion of the inner membrane of the bladder. So called from M. Noel, who first accurately described it.

EXOCYST'Ē SOLINGENIA'NA. Protrusion of the neck of the bladder. Called after M. Solingen, who first accurately described it.

EXODIC, Efferent.

EXGEDESIS, (*ex*, and *αἰδω*, 'to swell.')

Swelling.

EXOENTERITIS, see Enteritis.

EXOG'ENOUS, (*exo*, and *γεννω*, 'I engender.')

A term first applied to plants—hence called *Ex'ogens*—in which the wood increases by annual additions to the outside.

In animal anatomy, processes which shoot out from any part are termed *exog'enous*. An *exogenous aneurism* is one caused by division, from without, of the arterial walls.

EXOGENS, see Exogenous.

EXOGONIUM PURGA, *Convolvulus jalapa*.

EXOGONYAN'CON, (*exo*, *γονν*, 'the knee,' and *αγκων*, 'an elbow.')

Bowing of the knees outwards.

EXOINE, *Exoïne*, (low [L.] *exonium*, 'an excuse' from *exonerare*, 'to exonerate' [?].) In France, a certificate of excuse, exemption, or dispensation, given to those summoned to appear before a court of justice, and who are unable to do so.

EXOLCE, (*ἐξολκη*, from *ex*, and *ελκω*, 'I draw.')

Extraction.

EXOLUTION, Syncope.

EXOMETRA, (*exo*, and *μητρα*, 'uterus.')

Prolapsus uteri.

EXOMOIOSIS, (*ἐξομοιωσις*, from *ex*, and *ὁμοιος*, 'like.')

Assimilation.

EXOMPHALOCÉLE, (*exomphalus*, and *κηλη*, 'rupture.')

Exomphalus.

EXOM'PHALUS, *Exom'phalum*, (*ἐξομφαλον*,) *Exumbilica'tio, Exomphaloc'elē, Omphaloc'elē, Hern'ia umbilica'tis, Omphaloc'ochē, Omphaloprop'to'sis, Prolap'sus umbilic'i*, (F.) *Hernie ombilicale, H. du nombril, Umbilical hern'ia*, (*ex*, and *ομφαλος*, 'the navel.')

Hernia occurring at the navel. This affection happens more frequently in infants, and takes place by the umbilical ring. In adults, it occurs more commonly in females than in males; and, when it does so, the sac passes in the vicinity of the umbilicus. The organs found in this kind of hernia are particularly—the epiploon, the jejunum, the arch of the colon, and sometimes the stomach. The tumour is, in general, round, and presents all the characters of hernia. It is, commonly, readily reducible, and not subject to strangulation. It must be reduced, and retained by an elastic bandage, made in the form of a girdle, and furnished with a pad at its middle part. When strangulated, the stricture may be divided upwards and towards the left side.

EXONCO'MA, gen. *Exonco'matis*, (*ἐξογκωμα*,) *Exonco'sia*, (*ex*, *ογκος*, 'a tumour,' and *ομα*,) A large, prominent tumour. Used, by Galen, for protuberance of the vertebræ after luxation.

EXONCOSIS, (εξ, ογκος, 'a tumour,' and *osis*.)
Exoncoma—e. Linguae, Glossonecus.

EXONEIROGSMUS, (ἐξονειρωγμος, from *ex*, and *νειρωσσω*, (ονειρος, 'a dream,') 'to dream lasciviously.') Pollution, nocturnal.

EXONEIROISIS, Pollution, nocturnal.

EXONEURISM, (εξο, and νευρον, 'a nerve.')
Magnetism, animal.

EXOPHTHALMIA, *Pto'sis* seu *Prolap'sus* seu *Prociden'tia* *Oc'uli* seu *Bul'bi* *Oc'uli*, *Exor'bitis-mus*, *Ophthalmoc'e'lē*, *Ophthalmopto'sis*, *Oc'uli to'ti'us promin'en'tia*, (εξ, and οφθαλμος, 'eye.') (F.) *Procidence de l'œil*. A protrusion of the eye from its orbit, occasioned by an abscess or tumour in the areolar texture of the orbit; by exostosis of the parietes of the orbit, &c. In exophthalmia, the eye is pressed forwards; the eyelids are raised and separated, so that they can no longer cover the eye and defend it from the action of extraneous bodies; it becomes inflamed, and the sight is disturbed or destroyed. The treatment of course depends upon the cause. An anæmic condition, accompanied by protrusion of the eyeballs, palpitation of the heart and arteries, and tumefaction of the thyroid gland, has been described under the names *Cachex'ia exophthal'mica*, *Exophthal'mic goitre*, *Maladie de Basedow*, and *Anæmic protrusion of the eye-balls*, (F.) *Cachexie exophthalmique*.

EXOPHTHALMIA FUNGOSA, Sarcosis bulbi—e. Sarcomatica, Sarcosis bulbi.

EXOPHTHALMIC, *Exophthal'micus*, (F.) *Exophthalmique*. Appertaining or relating to exophthalmia.

EXOPHTHALMUS, (εξοφθαλμος, Goggle-eyed, (Prov.) *Grozet-eyed*. One whose eyes are very prominent. The opposite to Cœlophthalmus.

EXORBITISMUS, (εξ, and orbita, 'orbit.')
Exophthalmia.

EXORMIA, (εξ, and ορμαιν, 'to break forth,')
Papula—e. Lichen, Lichen—e. Prurigo, Prurigo—e. Strophulus, Strophulus.

EXORTUS UNGUIUM, ('the beginning of the nails,') See Nail.

EXOSIS, (εξωσις, from *ex*, and ωδω, 'to impel,')
Luxation.

EXOSKELETON, see Skeleton.

EXOSMIC, Exosmotic.

EXOSMOSE', *Exosmo'sis*, *Transuda'tion*, (εξ, and ωσμις, 'impulse,') The opposite to Endosmose. The act by which substances transude from within to without an animal or other membrane.

EXOSMOTIC, *Exosmot'icus*, *Ecos'mic*, *Exos'micus*. Belonging to Exosmose—as an exosmotic current.

EXOSSATIO, (εξ, and ος, gen. *ossis*, 'a bone,')
Exostosis.

EXOSTEMMA or EXOSTEMA CARIBÆUM, gen. *Exostemmat'is*, (εξο, 'without,' and στεμμα, 'a crown,') *Cinchona caribæa* cortex.

EXOSTOMA, Exostosis.

EXOSTOSE, Exostosis—e. *des Dents*, *Exostosis dentium*—e. *Sous-unguëale*, see Subungual.

EXOSTO'SIS, *Hyperosto'sis*, *Emphy'ma exostosis*, *Exosto'ma*, *Epосто'ma*, *Epосто'sis*, *Osteo'ma*, *Osteoph'ya*, *Osteonco'sis*, *Exossa'tio*, *Os'sis Eminen'tia*, *Os'teophyte*; (F.) *Exostose*, (εξ, and οστέον, 'a bone,') An osseous tumour, which forms at the surface of bones, or in their cavities. Various kinds have been enumerated. *Ivory Exosto'sis*, (F.) *E. eburnée*;—that which has the appearance and consistence of ivory. *Laminar Exosto'sis*, (F.) *E. Laminée*;—that which is formed of laminae in superposition, or of distinct filaments. The *Spongy* *Exosto'sis* is that whose structure is analogous to the spongy tissue of bones. Exostoses are sometimes distinguished into the *true*, which seem to be a projection of the osseous substance, and have the same or-

ganization and hardness as that substance; and the *false* or osteo-sarcoma. Exostosis may depend on syphilis, scrofula, rickets, gut, &c. In such cases, it is important to get rid of the primary disease.

Those *exostoses*, which occur within the bones, have by some been called *Enosto'ses*.

Exostosis, Spina ventosa.

EXOSTOSIS DENTITUM, (F.) *Exostose des Dents*. Exostosis of the teeth.

EXOSTOSIS STEATOMATODES, Osteosteatomia—e. Subungual, see Subungual.

EXOTERIC, (εξωτερικ, 'outer,') See Esoteric.

EXOTHESIS, (εξωθησις, from εξωδω, 'I put out,') Luxation.

EXOTIC, *Eco'ticus*, (εξωτικος, from εξω, 'with-out,') That which comes from abroad. Plants or drugs which are procured from abroad are so called. It is opposed to indigenous.

EXOTICADEN'IA, (*exot'ie*, and αἰσω, 'I dislike,') Aversion for exotic drugs.

EXOTICHEMATOSIS, (*exotic*, and *hæmatisis*.) Transfusion.

EXOTICOMANIA, (*exotic*, and *mania*.) The opposite to exoticadenia. Fondness for exotic remedies.

EXOTICOSYMPHYSIS, (*exotic*, and συμψις, 'a growing together,') A union or growing together of foreign bodies, as of a foreign body with the human.

EXPANSIO, (*expandere*, *expansum*, (εξ, and pandere, 'to open,') 'to spread out,') Expansion—e. Musculosa, Platysma myoides.

EXPANSION, *Expan'sio*. A prolongation or spreading out presented by certain organs. Thus, we say an *aponeurotic expansion*, &c.

EXPATRATIO, (εξ, and patratio, 'copulation,') Ejaculation (of Sperm).

EXPECTANT, *Expectans*, (*expectare*, *expectatum*, (εξ, and spectare, 'to look,') 'to wait,') That which waits:—as *Expectant Medicine*,—*La Médecine expectante*. See Expectation.

EXPECTATION, *Expecta'tio*. The word *expectation* has been applied, in medicine, to that method, which consists in observing the progress of diseases, and removing deranging influences, without prescribing active medicines, unless such shall be imperiously required. It consists, in fact, in leaving the disease almost wholly to the efforts of nature, and has been termed the *art of curing diseases by expectation or waiting*, *Ars sanan'di cum expecta'tio'nē*.

EXPECTORANT, *Expectorative*, *Expectorans*, *Anacathar'ticus*, *Sputato'rius*, *Antib'e'chicus*, *Pty'smagogue*, (εξ, and pectus, gen. *pectoris*, 'the breast,') A medicine capable of facilitating or provoking expectoration. There is probably no such thing as a *direct* expectorant. They all act through the system, or by impressions made on parts at a distance, which, through the medium of general, continuous, or contiguous sympathy, excite the secretory vessels of the air passages into action. The following are the chief reputed expectorants:—Ammoniacum; Asafetida; Galbanum; Ipecacuanha; Myroxylon; Myrrha; Inhalations of Iodine, Stramonium, Tar, Burning Wool, Tobacco, &c.; Scilla; Senega, and Tolutanum.

EXPECTORATIO, Expectoration—e. Sanguinis, Hæmoptysis—e. Solida, Cynanche trachealis.

EXPECTORATION, *Expectora'tio*, *Ecch'lysis*, *Bex hu'mida*, *Anap'tysis*, *Prop'tysis*, *Stethocathar'sis*, *Anacathar'sis*, *Anabex'is*, *Rais'ing*. The act of expelling from the chest matters or secretions collected or existing there. It is, likewise, used for the expectorated matter.

EXPECTORATION, PRUNE-JUICE, see Prune-juice.

EXPECTORATION DE SANG, Hæmoptysis.

EXPECTORATIVE, Expectorant.

EXPELLENS, (*expellere*, (*ex*, and *pellere*), 'to drive away.') Expulsive.

EXPERIENCE, *Experientia*, *Poir'a*, *Empir'a*, (*ex*, and *πειρα*, 'a trial.') A knowledge of things acquired by observation. In medicine, this knowledge can be obtained both by the practitioner's own experience, and by that obtained from tradition and from books. To profit by experience requires a mind capable of appreciating the proper relations between cause and effect; and hence it happens, that *false experience*, *Experientia fallax*, is extremely common; and that a man had better, in many instances, trust to that which he has learned from others, than to his own fallacious observation.

The union of accurate observation by the physician with that handed down by medical writers constitutes perfect experience, so far as it is attainable in any individual case.

EXPÉRIENCE, Experiment—*e. de Mariotte*, Mariotte, experiment of.

EXPERIENTIA, Experience.

EXPERIMENT, *Experimentum*, (F.) *Expérience*. A trial, made on the bodies of men or animals, for the purpose of detecting the effect of a remedy, or of becoming better acquainted with their structure, functions, or peculiarities. In a more general sense, it means any trial instituted with the intent of becoming better acquainted with any thing. By experiments on living animals, we have obtained much valuable information in the various departments of medicine; but particularly in physiology and toxicology.

EXPERIMENT OF MARIOTTE, see Mariotte.

EXPERIMENT OF WEPFER, *Experimentum Wepferianum*. An experiment attributed to Wepfer, which consisted in injecting air into the blood-vessels of animals, with the view of observing its effects on the economy.

EXPERTS NUPTIARUM, ('ignorant of marriage rites,') Virgin.

EXPERT, (F.) *Expertus*, (*ex*, and *peritus*, 'skilled,') Expert. Skillful or of good experience. A physician, charged with the duty of making a report, (F.) *Expertise*, upon any case of legal medicine.

EXPERTISE, see Expert.

EXPIRATION, *Expiratio*, *Expiratio*, *Ec'pnoë*, *Ecpeumatosis*, *Ecpeunis*, *Aponeunis*, *Aponeunis*, *Aponeunis*, (*ex*, and *spirare*, 'to breathe out,') The act of expelling from the chest, the air received during inspiration. 'To expire,' is, also, to emit the last breath, or 'to die.' See Vital capacity.

EXPIRATORY, *Expiratio'ni inser'viens*. Relating or appertaining to expiration. The expiratory muscles are all those which contribute to diminish the cavity of the chest, for the purpose of expelling the air contained in the lungs or of producing expiration. These muscles are, chiefly, the intercostals, triangularis sterni, quadratus lumborum, serratus posticus inferior, the oblique and recti muscles of the abdomen, and the sacrolumbalis.

EXPIRE, see Expiration.

EXPLORATIO, (*explorare*, *exploratum*, 'to search into,') Exploration—*e. Abdominis*, Abdominoseopia.

EXPLORATION, *Exploratio*, *Recongnit'io*. The act of observing and attentively examining or investigating every thing connected with a case of disease. The word is chiefly used in this sense by the French practitioners.

EXPLORATOR, CHEST. An instrument, proposed by Dr. B. Babington for exploring the chest in cases of empyema. It consists of a needle, contained in the smallest sized canula. This is passed between the ribs into the chest.

The needle is then withdrawn, and the escape of fluid indicates the nature of the case.

EXPLORATORIUM, Sound, Specillum.

EXPRES'SION, *Expres'sio*, *Ecpi'es'mus*, *Ecthlip'sis*, (*ex*, and *premere*, *pressum*, 'to press,') The act of compressing a substance, for the purpose of separating from it the fluids which it contains. Also, the manner in which impressions made upon us are depicted; especially in the traits of the countenance.

EXPRESSION OF SWEAT, (F.) *Sueur d'expression*, is a term given to the passive perspiration observable in very debilitated individuals.

EXPUITION, Exspuition.

EXPULSIF, Expulsive.

EXPULSIO, (*expellere*, *expulsum*, (*ex*, and *pellere*), 'to drive away,') Excretion—*e. Fæcum*, Defecation—*e. Fœtus*, Parturition.

EXPULSION, SPONTANEOUS, Evolution, spontaneous.

EXPULSIVE, *Expellens*, *Expulso'rius*, (F.) *Expulsif*. Driving out.

An *expulsive bandage*, (F.) *Bandage expulsif*, is one constructed with the view of compressing a part, from which we are desirous of expelling pus, serum, &c. Certain medicines were formerly called *expulsives*, which were believed to have the power of driving the humours towards the skin;—as diaphoretics, and sudorifics.

EXPULTRIX, see Vis expultrix.

EXSANGUE, (*ex*, and *sanguis*, 'blood,') Exanguous.

EXSANGUINE, Exanguious.

EXSANGUINITY, Anamia.

EXSANGUIS, Exanguious. *Exsang'uis* is used by Ausonius for one exhausted by venery;—as *sanguis* meant sperm as well as blood. See Sperm.

EXSARCOMA, gen. *Exsarco'matis*, Sarcoma.

EXSCREATIO, Excretion.

EXSECTIO, Exsection—*e. Virilium*, Castration.

EXSECTION, *Exsectio*, (*ex*, and *secare*, *sectum*, 'to cut,') The cutting out of any thing, as of a nerve.

EXSECTUS, Castratus, Eunuch.

EXSICCATIO, (*ex*, and *siccare*, *siccatum*, 'to dry,') Drying.

EXSICCATIVA, Desiccativa.

EXSOLUTIO, (*ex*, and *solvere*, *solutum*, 'to loosen,') Eclysis.

EXSPIRATION, Expiration.

EXSPUIT'ION, *Exspuit'io*, *Sput'io*, *Sputatio*, *Anachremp'sis*, *Apochremp'sis*, *Chremp'sis*, *Pty'sis*, *Anacine'ma*, *Anacine'sis*, (F.) *Expuition*, *Sputation*, (*ex*, and *spuo*, 'I spit,') Rejection of the matters accumulated in the pharynx and larynx; spitting.

EXSTASIS, Ecstasis.

EXTINCTIO, see Extinctio.

EXSTIRPATIO, Extirpation—*e. Lingue*, Glossostere'sis—*e. Testiculorum*, Castratio.

EXTROPH'IA, *Ex'trophy*, *Ec'trophë*, (*ex*, and *τροφή*, 'turning,') *Extrover'sio*, *Extrover'sion*. Eversion or turning out of a part—as of the eyelids. A term used by M. Chaussier for certain displacements of organs, and especially of the urinary bladder.

EXTROPH'IA OR EX'TROPHY OF THE BLAD'DER, (F.) *Renversement de la Vessie*, is a faulty conformation, in which the organ opens above the pubes; so that in the hypogastric region there is a red, mucous surface, formed by the inner coat of the bladder; on which two prominences are distinguishable, corresponding to the openings of the ureters.

EXSUCCATIO, (*ex*, and *succus*, 'juice,') Echy-moma.

EXSUC'TIO, *Suc'tio*, *Ecmyze'sis*, (*ex*, and *sugere*, *suctum*, 'to suck,') The action of sucking.

EXSUDATIO, (*ex*, and *sudare*, *sudatum*, 'to sweat,') Ephidrosis, Exudation.

EXSUFFLATIO, (*ex*, and *sufflare*, *sufflatum*, 'to blow.') Emphysema.

EXTA, see *Entails*.

EXTASE, Ecstasie.

EXTEMPORANEUS, *Extempora'neus*, *Extempora'lis*, (F.) *Extemporane*, (*ex*, and *tempus*, *gen. temporis*, 'time.') Those prescriptions are called 'extemporaneous,' or 'magistral,' which are made on the spot, and composed according to the prescription of the physician.

EXTENSEUR COMMUN DES DOIGTS, (*extendere*, (*ex*, and *tendere*, *tensum*.) 'to stretch out.') Extensor digitorum communis—*e. Commun des orteils*, Extensor communis digitorum pedis—*e. Court du Pouce*, Extensor pollicis brevis—*e. Long du Pouce*, Extensor longus pollicis—*e. Petit des orteils*, Extensor brevis digitorum pedis—*e. Propre de l'Index*, Extensor proprius indicis—*e. Propre du petit doigt*, Extensor proprius minimi digiti.

EXTENSIBILITY, *Extensibil'itas*, (*extensio*, and *habilis*, 'able.') A property, possessed by certain bodies, of being capable of extension or elongation.

EXTENSIO, Extension, Tetanus.

EXTENSION, *Extensio*, *Tensio*, *Ec'tasis*, *Cat'asis*, *Ana'asis*. An operation in surgery, in which either by the hands alone, or by straps, a fractured or luxated limb is pulled strongly, to restore it to its natural position. It is the opposite of *Counterextension*.

EXTENSOR. (F.) *Extenseur*. A muscle, whose office is to extend certain parts.

EXTENSOR BREVIS DIGITORUM PE'DIS, *Ped'i-cus*, (F.) *Calcaneo-phalangien commun*, of *DE-MAS*; *Muscle pédonier ou petit extenseur des orteils*, *Calcaneo-sus-phalangien commun*—(Ch.) *Carré du pied*; *Short Extensor of the Toes*. A muscle, situate on the dorsal region of the foot. It arises from the external surface of the calcaneum, and at the anterior edge of a ligament, which unites that bone to the astragalus. Anteriorly, each of its divisions terminates by a small tendon, which is fixed successively, at the superior part of the posterior extremity of the first phalanx of the great toe, and to the second and last phalanges of the next three toes. Its use is to extend the first four toes, and to direct them a little outwards.

EXTENSOR CARPI RADIALIS BREVIOR, see *Radialis*—*e. Carpi ulnaris*, see *Cubital muscles*.

EXTENSOR COMMUNIS DIGITORUM PE'DIS, *E. Longus Digitorum Pe'dis*, *E. Digitorum longus*, *Cnemodactyl'eus*, (F.) *Péronéo-tibi-sus-phalangien commun*, *Péronéo-sus-phalangien commun*, *Extenseur commun des Orteils*. This muscle is situate at the anterior part of the leg. It is long, thin, flattened, simple, and fleshy above, and divided into four tendons below. It arises from the outer tuberosity of the tibia and the anterior surface of the fibula, and is inserted into the superior part of the posterior extremity of the second and third phalanges of the last four toes. It extends the three phalanges of these toes.

EXTENSOR DIGITORUM COMMUNIS, *E. digitorum communis ma'nus cum extensor'e pro'prio auricu-laris*, *Digito'rum Ten'or*; (F.) *Épicondyl'o-sus-phalangien commun*—(Ch.) *Extenseur commun des Doigts*, *Common Extensor of the Fingers*. A long, flattened muscle; simple above, and divided into four portions inferiorly. It is situate at the posterior part of the forearm; arises from the external tuberosity of the humerus, from the aponeurotic septa situate between it and the neighbouring muscles, and is inserted at the posterior surface of the second and third phalanges of the last four fingers. This muscle extends the phalanges of the last four fingers upon each other, and upon the metacarpal bone. It can, also, extend the hand on the forearm.

EXTENSOR DIGITORUM LONGUS, *E. communis*

digitorum pedis—*e. Dorsi communis*, *Sacro-spi-nalis*—*e. Indicis*, *E. proprius indicis*—*e. Inter-nodii ossis pollicis*, *E. longus pollicis*—*e. Minor Pollicis manus*, *E. pollicis brevis*—*e. Ossis meta-carpi pollicis manus*, *Abductor longus pollicis*—*e. Longus digitorum pedis*, *E. communis digito-rum pedis*.

EXTENSOR POL'LICIS BRE'VIS, *E. mi'nor pol'licis ma'nus seu pri'mi in'terno'dii seu pol'licis pri'mus seu secu'ndi interno'dii* (DOUGLAS.) *seu secu'ndi interno'dii os'is pol'licis*; (F.) *Cubito-sus-phalan-gien du pouce*,—(Ch.) *Court extenseur du pouce*. Seated at the posterior and inferior part of the forearm. It is thin, long, and broader at its middle than at the extremities. It arises from the posterior surface of the ulna and the inter-osseous ligament, and is inserted behind the su-perior extremity of the first phalanx of the thumb. It extends the thumb, and aids in supi-nation.

EXTENSOR POL'LICIS LONG'US; *E. ma'ior pol'li-cis ma'nus seu secu'ndi interno'dii seu pol'licis secu'ndus seu ter'tii interno'dii* (DOUGLAS.) *seu interno'dii os'is pol'licis*, (F.) *Cubito-sus-phalan-gien du pouce*,—(Ch.) *Muscle long extenseur du pouce*. This muscle is long, flat, and fusiform; and is seated at the posterior part of the forearm. It arises from the posterior surface of the ulna and the interosseous ligament, and is inserted at the posterior part of the superior extremity of the first phalanx of the thumb. It extends the last phalanx of the thumb upon the first. See *Ex-tensor proprius pollicis pedis*.

EXTENSOR POLLICIS SECUNDUS, *E. Pollicis longus*—*e. Primi Internodii*, *Abductor longus pollicis*, *E. pollicis brevis*—*e. Proprius hallucis*, *E. proprius pollicis pedis*.

EXTENSOR PRO'PRIUS IN'DICIS, *E. pro'prius pri'mi dig'iti ma'nus*, *E. In'dicis*, *Indica'tor seu secu'ndi interno'dii in'dicis pro'prius*, (F.) *Cu-bito-sus-phalangien de l'Index*,—(Ch.) *Exten-seur propre de l'Index*. This muscle is long and thin; broader in the middle than at the extremi-ties; and is situate at the posterior part of the forearm. It arises from the posterior surface of the ulna, and is inserted at the posterior part of the upper extremity of the second and third pha-langes of the index-finger. It extends the three phalanges of the index-finger; and has, besides, the same uses as the other extensors of the fingers.

EXTENSOR PRO'PRIUS MIN'IMI DIG'ITI, (F.) *Épicondyl'o-sus-phalangien du petit doigt*,—(Ch.) *Extenseur propre du petit doigt*. Situate on the inside of the *Extensor communis digito'rum*. It arises from the external condyle of the os humeri and the aponeurotic septa seated between it, the extensor communis digitorum, and the extensor carpi ulnaris; and is inserted into the last two pha-langes of the little finger. Its use is to extend the little finger, and even the hand upon the fore-arm.

EXTENSOR PRO'PRIUS POL'LICIS PE'DIS, *E. pro'prius Hall'ucis seu Lon'gus* (DOUGLAS.) *seu pol-li-cis longus*, *Péronéo-sus-phalangien du pouce*, *Péronéo-sus-phalangien du pouce*,—(Ch.) This muscle is situate at the anterior part of the leg. It is fleshy, broad, flat above; small and tendi-nous below. It arises from the anterior part of the middle third of the fibula, and is inserted into the posterior part of the superior extremity of the last phalanx of the great toe. It extends the last phalanx of the great toe upon the first, and the first upon the first metacarpal bone.

EXTENSOR SECUNDI INTERNODII, *E. pollicis brevis*, *Extensor pollicis longus*—*e. Secundi in-ternodii indicis proprius*, *E. proprius indicis*—*e. Tertii internodii*, *E. pollicis longus*—*e. Tertii in-ternodii indicis*, *Prior annularis*—*e. Tertii in-ternodii minimi digiti*, *Abductor minimi digiti*.

EXTENSOR TARSII MAGNUS, *E. Tarsi suralis*. A name given, by some anatomists, to the gastrocnemius and soleus combined.

EXTENSOR TARSII MINOR, Plantar muscle.

EXTENUATIO, (*extenuare, extenuatum, (ex, and tenuis,)* 'to make thin,') Emaciation.

EXTERGENTIA, (*extergere, (ex, and tergere,)* 'to clean,') 'to wipe clean,') Detergents.

EXTER'NAL DISEASES, (*exter, 'without,')* *Extrin'sic Diseases, Mor'bi exter'ni seu extrin'seci, Surg'ical diseases, (F.) Maladies externes.* Those diseases which occupy the surface of the body, and form the object of surgical pathology, requiring, generally, external means, or surgical operations.

EXTERNAT. The post or office of an *externe*.

EXTERNE, see Housesurgeon.

EXTERNUS AURIS, Laxator tympani — e. Mallei, Laxator tympani.

EXTESTICULATUS, (*ex, and testiculus, 'a testicle,')* Castratus, Eunuch.

EXTINCTIO, Death — e. Hydrargyri, Extinction of Mercury — e. Mercurii, Extinction of Mercury.

EXTINCTIO VO'cis, (F.) *Extinction de voix*. The French use this term for cases in which the voice is not wholly suppressed, but produces only feeble sounds:—*Incomplete aphonia*.

EXTINCTIO MERCURI, *Extinction of Mercury seu Hydrargyri*. Trituration of mercury with lard or other substance, until the metallic globules disappear. The mercury is then so divided, that it forms a black powder, generally considered to be a protoxide of mercury, but perhaps, erroneously.

EXTINCTION OF MERCURY, Extinctio Mercurii — e. de Voix, Extinctio vocis.

EXTIRPATION, *Extirpa'tio, Abla'tio, Aphæ'resis, (extirpare, (ex, and stirps, 'a root,')* 'to root out,') The complete removal or destruction of any part, either by cutting instruments or the action of caustics. Thus, we speak of the extirpation of cancer, polypus, encysted tumour, &c.

EXTOMIUS, (*ex, and temere, 'to cut,')* Eunuch.

EXTOZOA, (pl. of *Extozo'on,*) Ectozoa.

EXTOZOAIRES, Ectozoa.

EXTOZOARIA, Ectozoa.

EXTRA (L.), in composition, 'on the outside, without.'

EXTRA-PELVIO-PUBI-TROCHANTÉRIEN, Obturator externus.

EXTRACARDIAL, see Endocardiac.

EXTRACT, (*ex, and trahere, tractum, 'to draw,')* Extractum — e. of Aconite, Extractum aconiti — e. of Aconite, alcoholic, Extractum aconiti alcoholicum — e. Alcoholie, see Extractum — e. of Aloes, Extractum aloes — e. Extract, alcoholic of Arnica, Extractum Arnice alcoholicum — e. of Bael, liquid, Extractum Belæ liquidum — e. of Bark, Extractum cinchonæ — e. of Bark, resinous, Extractum cinchonæ resinosum — e. of Belladonna, Extractum belladonnæ — e. of Belladonna, alcoholic, Extractum belladonnæ alcoholicum — e. of Bittersweet, Extractum dulcamaræ — e. of Bittersweet, fluid, see Extractum dulcamaræ — e. of Broom-tops, Extractum cæcinum genistæ — e. Buchu, fluid, Extractum buchu fluidum — e. of Butternut, Extractum juglandis — e. of Cascarilla, resinous, Extractum cascarillæ resinosum — e. of Chamomile, E. anthemidis — e. of Cinchona, Extractum cinchonæ — e. of Colchicum, Extract of Colchicum — e. of Colchicum, acetic, Extractum colchici aceticum — e. of Colchicum, fluid, Extractum Colchici fluidum — e. of Coleoynth, Extractum coleoynthidis — e. of Coleoynth, compound, Extractum coleoynthidis compositum — e. of Cubebs, fluid, Extractum cubebæ fluidum — e. of Dandelion, Extractum taraxaci — e. of Dandelion, fluid, Extractum Taraxaci fluidum — e. of Digitalis, alcoholic, Extractum digitalis alcoholicum — e.

of Elaterium, Extractum elaterii — e. of Ergot, Ergotæ — e. of Ergot, fluid, see Extractum Ergotæ — e. Ethereal, see Extractum — e. of male Fern, see Polypodium filix mas — e. of Foxglove, Extractum digitalis — e. of Gentian, Extractum gentianæ — e. of Hellebore, alcoholic, Extractum Hellebori alcoholicum — e. of Hellebore, American, fluid, Extractum Veratri viridis fluidum — e. of Hellebore, black, Extractum hellebori — e. of Hemlock, Extractum conii — e. of Hemlock, alcoholic, Extractum conii alcoholicum — e. of Hemlock Fluid, Extractum Conii fluidum — e. of Hemp, purified, Extractum cannabis purificatum — e. of Henbane, Extractum hyoscyami — e. of Henbane, alcoholic, Extractum hyoscyami alcoholicum — e. of Henbane, fluid, Extractum hyoscyami fluidum — e. of Hops, Extractum humuli, E. lupuli — e. of Ignatia, alcoholic, Extractum Ignatiæ alcoholicum — e. of Ipecacuanha, fluid, Extractum Ipecacuanhæ fluidum — e. of Jalap, Extractum jalapæ — e. of Lead, Goulard's, Liquor plumbi subacetatis — e. of Lettuce, Extractum lactuæ — e. of Liqueurice, Extractum glycyrrhizæ — e. of Logwood, Extractum hæmatoxyli — e. of Lupulin, fluid, Extractum Lupulini fluidum — e. of Mayapple, Extractum podophylli — e. of Meat, saponaceous, Osmazome — e. of Nux Vomica, Extractum nucis vomicæ — e. of Oak-bark, Extractum corticis quercilæ — e. of Opium, fluid, Extractum opii liquidum — e. of Pareira Brava, Extractum pareiræ — e. of White Poppy, Extractum papaveris — e. of Quassia, Extractum quassiæ — e. of Quinia, Extractum quiniæ — e. of Rhatany, Extractum krameris — e. of Rhubarb, Extractum rhei — e. of Rhubarb, fluid, Extractum rhei fluidum — e. of Rhus, Extractum Rhusii — e. of Rue, Extractum rutæ — e. of Sarsaparilla, Extractum sarsaparillæ — e. of Sarsaparilla, fluid, Extractum sarsaparillæ fluidum — e. of Savine, Extractum foliorum sabinæ e. of Senega, alcoholic, Extractum Senegæ alcoholicum — e. of Senna, fluid, Extractum sennæ fluidum — e. of Serpentaria, fluid, Extractum Serpentariæ fluidum — e. of Spigelia, fluid, Extractum Spigeliæ fluidum — e. of Spigelia and Senna, Extractum spigeliæ et sennæ — e. of Stramonium, Extractum stramonii — e. of Stramonium, alcoholic, Extractum Stramonii alcoholicum — e. of Uva Ursi, Extractum uvæ ursi — e. of Uva Ursi, fluid, see Extractum Uva Ursi — e. of Valerian, Extractum valerianæ — e. of Valerian, alcoholic, Extractum valerianæ alcoholicum — e. of Wild Cherry Bark, Extractum Pruni Virginianæ fluidum — e. of Wormwood, Extractum cæcinum absinthii.

EXTRACTA, see Extractum — e. Fluida, see Extractum.

EXTRACTIF, Extractive — e. *Oxygéné, Apothème*.

EXTRACTIO, Extraction, Extractum.

EXTRACT'ION, *Extract'io, Extr'esis, Extr'cē, Exelcys'mus, Enchylo'sis*. The act of removing an extraneous substance from any part of the body. Thus, a splinter is said to be extracted. It is also applied to the removal of certain parts. The cataract is said to be *extracted*; a tooth is *extracted*, when carious, &c.

EXTRACT'IVE, (F.) *Extractif*. A peculiar, presumed principle, which has been admitted in extracts. Thus, *bitter extractive* is the immediate principle of bitter vegetables, &c. The term is, also, applied to ill defined animal matters met with on analysis.

EXTRACTIVE, OXIDIZED, *Apothème*.

EXTRACTS, see Extractum — e. Fluid, see Extractum.

EXTRACT'UM, *Enchylo'ma, Extract'io, Extr'act, (F.) Extrait, Apostolē*. An extract is prepared by evaporating vegetable solutions, till a tenacious mass is obtained. When prepared from an infusion or decoction, it is called a *watery*;

from ether, an *etheral*; and from alcohol an *alcoholic* or a *spirituous extract*. Both kinds contain all the principles of the vegetable that are soluble in the menstrua with which they are prepared; but the volatile parts are dissipated, and some of the fixed parts are decomposed; the proper extractive is oxygenized, and the virtues of the vegetable substance are altered or destroyed. Extracts are *hard, soft, or fluid*: the consistence of the soft being such as to retain the pilular form without the addition of a powder. A patent was taken out, many years ago, by a Mr. Barry, of London, for preparing them in *vacuo*; and, as the temperature is much lower than in the ordinary method, the virtues of the plant are less altered, and the extracts are generally green. Extracts are also prepared by displacement or percolation. They have, likewise, received different names, according to their predominant principle. The *gummy* or *mu'cous*, or *mucilag'inous*, are those which are mainly composed of gum or mucilage. Gum tragacanth may be considered a pure gummy extract. *Gelat'inous extracts* are those composed especially of gelatin; *res'inous extracts*, those of a resinous character; *extract'ores'inous*, those composed of extractive or colouring matter and resin; *gum-res'inous*, those containing gum and resin; and *saponaceous* or *saponaceous saline*, those containing a notable quantity of saline substances and a resinous matter, so combined with mucus and other soluble substances, that they cannot be separated. Under the name *Fluid Extracts*—*Extracta Fluida* seu *Liquida*—a class of preparations has been introduced into the Pharmacopœias, the distinctive character of which is, the concentration of the active ingredients of medicinal substances into a small bulk, in the liquid form.

EXTRACTUM ABSINTHII CACUMINUM, *Extract of Wormwood*. (A decoction defecated and evaporated.) The flavour is dissipated along with the essential oil. It is a bitter tonic. Dose, gr. x to ʒj, in pill.

EXTRACTUM ACONITI, *Extract of Aconite*, *Succus episcopus aconiti napellii*, (from the inspissated juice without defecation.) It is esteemed to be narcotic and diuretic; and has been given in the cases referred to under Aconitum. Dose, gr. j, gradually increasing it.

EXTRACTUM ACONITI ALCOHOLICUM, *Alcoholic Extract of Aconite*. (Aconit. in pulv. crass. ʒj; Alcohol. dilut. Oiv. Moisten the aconite with half a pint of diluted alcohol: let it stand for 24 hours: transfer it to a displacement apparatus, and gradually add the remainder of the diluted alcohol. When the last portion of this has penetrated the aconite, pour in from time to time water sufficient to keep the powder covered. Stop the filtration when the liquid which passes begins to produce a precipitate, as it falls, in that which has already passed. Distil off the alcohol, and evaporate to a proper consistence.—Ph. U. S. 1851. A modified formula has been received into the Ph. U. S. 1863.) Dose, half a grain or a grain.

EXTRACTUM AL'OEË, *Extract of Aloes*, *E. aloës aquosum* seu *purificatum*, *Purified extract of aloes*. (The gummy part extracted by boiling water, defecated and inspissated.) Dose, gr. v. to gr. xv.

EXTRACTUM ALOES AQUOSUM, *E. aloës*—*E. Aloes purificatum*, *E. aloës*.

EXTRACTUM ANTHEMIDIS, *E. anthemidis nobilis* seu *chamemeli* seu *florum chamemeli*, *Extract of Chamomile*. The volatile oil is dissipated in this preparation. It is a pure, grateful bitter, and is tonic and stomachic. Dose, gr. x. to gr. xx, in pill.

EXTRACTUM AR'NICE ALCOHOLICUM, *Alcoholic Extract of Ar'nica*. (Ph. U. S. 1863.) Prepared

in a similar manner to the Extractum Aconiti alcoholicum.

EXTRACTUM ASPRAGI, see Asparagus.

EXTRACTUM BE'LE LIQ'UIDUM, *Liquid Extract of Ba'el*. A preparation introduced into the British Pharmacopœia (1864). Astringent, in diarrhoea and dysentery. Dose, f ʒss, or more.

EXTRACTUM BELLADON'NE, *Succus episcopus atropæ belladonnae*, *Extract of Belladonna* (an expressed juice inspissated). Properties same as those of the plant. Dose, gr. 1, gradually increased. It dilates the pupil when applied to the eye.

EXTRACTUM BELLADON'NE ALCOHOLICUM, *Alcoholic extract of Belladonna*. (Prepared like the extractum aconiti alcoholicum.—Ph. U. S.) Dose, half a grain or a grain.

EXTRACTUM BUCHU FLU'IDUM, *Fluid Extract of Buchu*. This is official in the Ph. U. S. 1863.

EXTRACTUM CANNABIS, see Bangué.

EXTRACTUM CAN'NABIS PURIFICA'TUM, *Purified Extract of Hemp*, of the Ph. U. S. 1863, is made by rubbing *Extract of Hemp* in *Alcohol*; adding alcohol and macerating; filtering and evaporating to dryness. Dose, half a grain.

EXTRACTUM CAR'NIS, *Extract of Flesh*. An extract formed by evaporating beef tea in a water-bath. It can be kept for sale. An ounce of it may represent thirty-two ounces of meat.

EXTRACTUM CASCARIL'LE RESINO'SUM, *Res'inous extract of cascari'lla*. (Cort. cascari'llæ, in pulv. crass. ʒj; Sp. vini rect. ʒiv. Digest for four days; then decant and strain; boil the residuum in ten pints of water to two; filter and evaporate the decoction, and distil the tincture in a retort, till both are thickened; then mix and evaporate to a pilular consistence.) Dose, gr. x. to gr. xx, in pills.

EXTRACTUM CATECHU, Catechu—*e. Catharticum*, *E. Colocyntidis compositum*.

EXTRACTUM CATHOL'ICUM, (F.) *Extrait Catholique*. This epithet is given to pills composed of aloes, black hellebore, and colocynth, resin of jalap, and scammony. See, also, Extractum Colocyntidis compositum.

EXTRACTUM CHAMEMELI, *E. Anthemidis*—*e. Cicutæ*, *E. Conii*.

EXTRACTUM CIMICIF'UGÆ FLU'IDUM, *Fluid Extract of Cimicif'uga*, is official in the Ph. U. S. (1863). Dose, mxxv. to xx.

EXTRACTUM CINCHO'NE, *E. Corticis Peruvia'ni* seu *Cinchonæ Mol'le*, *Extract of Cinchona* or of *Bark*. (A decoction evaporated.) The active principles are similar to those of the bark in substance; but it is not so effectual, owing to the chymical change induced in the drug during the boiling. When reduced, by drying, to a state fit for being powdered, it is called *Hard Extract of Bark*, *Extractum Corticis Peruvia'ni durum*, *E. Cinchonæ durum*. Dose, gr. x to ʒss.

EXTRACTUM CINCHONÆ (Ph. U. S. 1863.) *E. Cinchonæ Flavæ*, of the Pharmacopœia of the United States (1842). An alcoholic extract of yellow cinchona made by exhausting the bark by means of alcohol in a percolator, and evaporating to the proper consistence. *Fluid Extract of Cinchona*, *Extractum Cinchonæ Flu'idum*, is official in the Pharmacopœia of the United States, 1863. Dose, f ʒss.

EXTRACTUM CINCHONÆ PAL'IDÆ, may be prepared in the same manner.

EXTRACTUM CINCHONÆ RESINO'SUM, *E. Cinchonæ lanceifolia* seu *Cinchonæ Rubræ resinosum*, *Res'inous Extract of Bark*, *E. Cinchonæ Resinæ*. The aqueo-spirituous extract contains both the extractive and resin of the bark. Dose, gr. x to xxx.

EXTRACTUM CINCHONÆ RU'BRÆ, may be prepared in the same manner as Extractum cinchonæ flavæ.

EXTRACTUM COLCHICI, *Extract of Colchicum*, is prepared in the same manner as Extractum Aconiti. Dose, one or two grains.

EXTRACTUM COLCHICI ACETICUM, *Acetous or Acetic Extract of Colchicum*. (Colchic. rad. in pulv. ℥j, Acid. acet. f℥iv, Aquæ q. s. To the acid add a pint of water, and mix this with the root. Put the mixture in a percolator, and pour on water until the liquid that passes has little or no taste. Evaporate to a proper consistence. Ph. U. S.) Dose, gr. j to gr. iij.

EXTRACTUM COLCHICI RADICIS FLUIDUM, *Fluid Extract of Colchicum Root*; and *Extractum Colchici Seminis Fluidum*, *Fluid Extract of Colchicum Seed*, are official in the Ph. U. S. 1863. Dose, ℞.

EXTRACTUM COLOCYNTHIDIS, *Extract of Colocynth*. (Ph. L. & E.) (A watery infusion or decoction of colocynth evaporated.) Cathartic, in the dose of from gr. v to ℥ss.

EXTRACTUM COLOCYNTHIDIS ALCOHOLICUM, *Alcoholic Extract of Colocynth*. (Obtained by the action of diluted alcohol on colocynth, by maceration, percolation, and evaporation.) (Ph. U. S. 1863.)

EXTRACTUM COLOCYNTHIDIS COMPOSITUM, *E. Catharticum seu Catholicum seu Quercetani Compound Extract of Colocynth*. (Colocynth. pulp. concis. ℥vj, Aloes, pulv. ℥xij, Scammon, pulv. ℥iv, Cardamon. pulv. ℥j, Saponis ℥ij, Alcohol. dilut. cong. Macerate the pulp in the spirit at a gentle heat for four days; strain; add the aloes and scammony; then distil off the spirit and mix in the cardamon seeds. Ph. U. S.)

In the Ph. U. S. of 1863, it is directed to be made of *Alcoholic Extract of Colocynth*, in fine powder, ℥iiss; *Socotrine Aloes*, in fine powder, ℥xij; *Resin of Scammony*, in fine powder, ℥ij; *Cardamon*, in fine powder, ℥i; *Soap*, in fine powder, ℥iii; mixed well and kept in a well-stopped bottle. It is an active cathartic, and is used in obstinate visceral obstructions, &c. Dose, gr. vj. to ℥ss.

EXTRACTUM CONII, *E. Cicuta*, *Succus cicutæ spissatus*, *S. spissatus conii maculati*, *Extract of Hemlock*. (In the Ph. U. S. of 1863, fresh hemlock is sprinkled with water, and the juice expressed, filtered, and evaporated.) Employed in the same cases as the conium. Dose, gr. iij to ℥ss.

EXTRACTUM CONII ALCOHOLICUM, *Alcoholic Extract of Hemlock*. (Prepared like the Extractum aconiti alcoholicum.—Ph. U. S.) Dose, one or two grains.

A *Fluid Extract of Hemlock*, *Extractum Conii Fluidum*, is in the Ph. U. S. (1863.) Dose, ℞.

EXTRACTUM CONVULVULI JALAPÆ, *E. jalapæ* —e. *Corticis Peruviani*, *E. Cinchonæ*.

EXTRACTUM CUBEÆ FLUIDUM, *Oleoresina Cubeæ*, (Ph. U. S. 1863.) *Fluid Extract of Cubebs*. (Cubebs exhausted by ether through percolation, and the solution evaporated. Dose, from 5 to 30 minims.)

EXTRACTUM DIGITALIS, *Extract of Foxglove*. (Prepared like Extractum conii.) Dose, from half a grain to a grain.

EXTRACTUM DIGITALIS ALCOHOLICUM, *Alcoholic Extract of Digitalis* (Ph. U. S. 1863), is prepared like Extractum aconiti alcoholicum.

EXTRACTUM DULCAMARÆ, *Extract of Bittersweet*. (Prepared by displacement, by means of diluted alcohol, from bittersweet in moderately fine powder.—Ph. U. S.) Dose f℥j.

A *Fluid Extract of Bittersweet*, *Extractum Dulcamaræ Fluidum*, is in the Ph. U. S. for 1863.

EXTRACTUM ELATERII, *Elaterium*, *Extract of Elaterium*. (The fecula of the expressed juice of the fruit of *Momordica elaterium*.) It is violently cathartic, hydragogue, and sometimes emetic. Dose, gr. ss every hour till it operates.

EXTRACTUM ERGOTÆ, *Ergotin*. A *Fluid Ex-*

tract of *Ergot*, *Extractum Ergotæ Fluidum*, is in the Ph. U. S. 1863. Dose, ℞xxx to f℥j.

EXTRACTUM FILICIS ÆTHEREUM, see *Polypodium filix mas*.

EXTRACTUM GENISETÆ CACUMINUM, *Extract of Broom Tops*. Diuretic and stomachic. Dose, ℥ss to ℥j.

EXTRACTUM GENTIANÆ, *E. Gentianæ luteæ seu Radicis Gentianæ*, *Extract of Gentian*. (The evaporated decoction.) Prepared also by displacement.—Ph. U. S.) Properties like those of Gentian. Dose, gr. x to ℥ss.

A *Fluid Extract of Gentian*, *Extractum Gentianæ Fluidum*, is in the Ph. U. S. for 1863. Dose, f℥j.

EXTRACTUM GLYCYRRHIZÆ, *Succus Glycyrrhizæ inspissatus seu Liquiritiæ*, *Spanish Juice*, *Italian Juice*, *Black Sugar*, *Extract of Liquorice*. (The evaporated decoction.) It is demulcent, taken ad libitum.

Refined Liquorice, which is sold in the form of cylinders, is made by gently evaporating a solution of the pure extract of liquorice with half its weight of gum Arabic, rolling the mass and cutting it into lengths, and then polishing, by rolling them together in a box.

EXTRACTUM GRAMINIS, *Extract of Grass*. An extract prepared from *Triticum repens*. It is considered by the Germans to be a mild tonic; and is greatly used, especially in convalescence from fever. It is probably devoid of all injurious properties, and, as probably, totally inefficacious.

EXTRACTUM HEMATOXYLI, *E. Hematoxyli Campechiani* seu *Scohis Hematoxyli*, *Extract of Logwood*. (The evaporated decoction.) It is astringent. Dose, gr. x to ℥ss.

EXTRACTUM HELLEBORI, *E. Hellebori nigri*, *Extract of Black Hellebore*. (The evaporated decoction.) In large doses, this is cathartic; in smaller, diuretic, resolvent, [?] and emmenagogue. [?] Dose, as a cathartic, gr. x to ℥j; as an emmenagogue, gr. iij to gr. x.

Extractum Hellebori, (Ph. U. S. 1851.) *E. H. Alcoholicum*, *Alcoholic Extract of Black Hellebore*, (Ph. U. S. 1863), is prepared from Black Hellebore, in coarse powder, like the Extractum aconiti alcoholicum. Dose, like the last.

EXTRACTUM HUMULI, *E. Lupuli*, *Extract of hops*. (The evaporated decoction.) It is tonic, anodyne, [?] diuretic. [?] Dose, gr. v to ℥j.

EXTRACTUM HYOSCYAMI, *Succus spissatus Hyoscyami*, *Extract of Henbane*. (The expressed juice, inspissated without defecation, or the expressed juice of the bruised leaf sprinkled with water, heated to the boiling point; strained, and evaporated. Ph. U. S. 1863.) Its virtues are narcotic. Dose, gr. iij to ℥ss.

EXTRACTUM HYOSCYAMI ALCOHOLICUM, *Alcoholic Extract of Henbane*. (Prepared from leaves of Hyoscyamus, in coarse powder, like the Extractum aconiti alcoholicum, Ph. U. S.) Dose, one or two grains. A *fluid Extract of Henbane*. *Extractum hyoscyami fluidum* is official in the Ph. U. S. (1863.) Dose, ℞.

EXTRACTUM IGNATIE ALCOHOLICUM, *Alcoholic Extract of Ignatia*. (Ignatia exhausted by alcohol in a percolator: and the solution evaporated to a proper consistence. Ph. U. S. 1863.) Dose, half a grain to a grain.

EXTRACTUM IPECACUANHÆ FLUIDUM, *Fluid Extract of Ipecacuanha*. Admitted into the Ph. U. S. for 1863. Dose, ℞.v. to xx.

EXTRACTUM JALAPÆ, *E. Convolvuli Jalapæ* seu *Jalapii* seu *Jalapæ*, *Extract of Jalap*. (A spirituous tincture distilled; and an aqueous decoction evaporated; the residua being mixed together: kept both soft and hard.) It is cathartic and hydragogue. Dose, gr. x to ℥j.

EXTRACTUM JALAPÆ RESINOSUM, *Resina Jalapæ*, (Ph. U. S.), *Resinous Extract of Jalap*.

(Made by exhausting *Jalap* powder, by displacement by alcohol; throwing down the resin from the fluid by means of water, and drying the precipitate.) Properties like the last.

EXTRACTUM JUGLANDIS, *Extract of Butternut*. (Prepared by displacement from butternut, in moderately coarse powder.—Ph. U. S.) Dose, gr. x to xx.

EXTRACTUM KRAMERIE, *Extract of Rhatany*. (Prepared by displacement from rhatany, in moderately fine powder.—Ph. U. S.) Dose, gr. x to xx.

EXTRACTUM LACTUCE, *Succus spissatus Lactuce sativæ, Extract of Lettuce*. (Leaves of fresh lettuce 1*lb*; beat them in a stone mortar, sprinkling them with water; then express the juice and evaporate, without allowing it to subside until it acquires a proper degree of consistence.) It is said to be narcotic and diaphoretic. Dose, gr. iij to gr. x.

An extract is, sometimes, made from the juice of the wild lettuce, *Lactuca virosa*, which is regarded as diuretic.

EXTRACTUM LUPULINÆ FLUIDUM, *Fluid Extract of Lupulin*. Admitted into the Ph. U. S. 1863. Dose, $\mathfrak{m}$ v to x.

EXTRACTUM MARTIS ACETICUM, Ferri Acetas.

EXTRACTUM NUCIS VOMICÆ ALCOHOLICUM, *Alcoholic Extract of Nuc Vomica*, see *Strychnos nux vomica*. Dose, $\frac{1}{2}$ a grain to a grain.

EXTRACTUM OPII, *E. Opii aquosum seu Thebaicum seu Opii gummosum, Extract of Opium, Laudanum opiatum seu simplex, Opium colatum seu depuratum*. (A watery solution defecated and evaporated.) Dose gr. ss to gr. iss,—about half that of opium.

A *Fluid Extract of Opium, Extractum opii liquidum*, is official in the Ph. B. 1864.)

EXTRACTUM PANCHYMOGUM. A drastic medicine, composed of colocynth, bruised with its seeds; sena bruised; black hellebore root, Agaric, Scammony, in powder, *Extract of Aloes*, and *Powder of Diarrhodon*, into which roses entered.

EXTRACTUM PAPAVERIS, *E. Papaveris somniferi seu Papaveris albi, Extract of white poppy*. (The decoction evaporated.) It possesses nearly the same virtues as opium, but is much weaker. Dose, gr. ij to $\mathfrak{ij}$.

EXTRACTUM PAREIRÆ, *Extract of Pareira Bra'va*. Prepared like *Extractum Hamatoxyli*. Dose, ten grains to half a drachm.

A *Fluid Extract of Pareira, Extractum Pareiræ liquidum*, is official in the Ph. B. 1864. Dose, f $\mathfrak{z}$ j.

EXTRACTUM PIPERIS FLUIDUM, *Oleoresina Piperis*. (Ph. U. S. 1863.) *Fluid Extract of Black Pepper*. (Black pepper exhausted by ether through percolation, the solution evaporated, and the piperin in crystals separated by expression.) Dose, one to five minims.

EXTRACTUM PODOPHYLLI, *Extract of Mayapple*. (Prepared from podophyllum, in coarse powder, in the same manner as the *Extract of cinchona*.—Ph. U. S.) Dose, gr. v to gr. xv, as a cathartic.

EXTRACTUM PRUNI VIRGINIANÆ, *Fluid Extract of Wild Cherry Bark*. Admitted into the Ph. U. S. for 1863. Dose, f $\mathfrak{z}$ j.

EXTRACTUM PURGANS, see *Hedera helix*—e. *Quercetani*, E. *Colocyntidis* compositum.

EXTRACTUM QUASSIÆ, *Extract of Quassia*. (Prepared by displacement from Quassia in moderately fine powder.—Ph. U. S.) Dose, as a tonic, five grains and more.

EXTRACTUM CORTICIS QUERCUS, *Extract of oak bark*. (The decoction evaporated.) It is astringent and tonic.

EXTRACTUM QUININÆ, *Quininæ sulphas impuræ*. (This is made by evaporating the liquor poured off the crystals of sulphate of quinia to the consistence of a pilular mass.) Twenty-four grains will generally arrest an intermittent.

EXTRACTUM RHEI, *E. Rh. alcoholicum, Extract of Rhubarb*. (A solution in diluted alcohol by percolation evaporated.) Uses like those of the powdered root. Dose, gr. x to $\mathfrak{z}$ ss.

EXTRACTUM RHEI FLUIDUM, *Fluid Extract of Rhubarb*, is in the Pharmacopœia of the United States. Dose, f $\mathfrak{z}$ j.

EXTRACTUM RU'DII, *Extract of Ru'dius*, (F) *Extrait de Rudius*. (Pills made of colocynth, agaric, scammony, roots of black hellebore and jalap, socotrine aloes, cinnamon, mace, cloves, and alcohol.) See *Pilulæ Aloes et Colocyntidis*.

EXTRACTUM RUTÆ GRAVEOLENTIS, *E. Rutæ seu foliorum Rutæ, Extract of Rue*. (A decoction evaporated.) The volatile oil being dissipated in the boiling, this is not a good preparation. Tonic, stomachic. Dose, gr. x to $\mathfrak{ij}$.

EXTRACTUM FOLIORUM SABINÆ, *Extract of Savine*. (A decoction evaporated.) The same remarks may be made on this preparation as on the last. Tonic. Dose, gr. x to $\mathfrak{z}$ ss.

EXTRACTUM SANGUINIS BOVINI, see *Blood*.

EXTRACTUM SARSAPARILLÆ, *Extract of Sarsaparilla*. (A strained decoction evaporated.) Virtues the same as those of the powdered root. Dose, gr. x to $\mathfrak{z}$ j. *Extractum Sarsaparillæ* of the United States Pharmacopœia is prepared from Sarsaparilla, in coarse powder, like the *Extractum aconiti alcoholicum*.

EXTRACTUM SARSAPARILLÆ FLUIDUM, *Fluid Extract of Sarsaparilla*, Ph. U. S. for 1863, is a simple fluid extract.

EXTRACTUM SARSAPARILLÆ FLUIDUM COMPOSITUM, *E. Sarsæ Liquidum, E. Sarsæ Fluidum, Compound Fluid Extract of Sarsaparilla, Extractum Sarsaparillæ Fluidum, Fluid Extract of Sarsaparilla*, Ph. U. S. 1851.) (*Sarsaparilla*, concis. et contus. $\mathfrak{z}$ xvj, *Glycyrrhiz*, contus., *Sassafras*, rad. contus., aa $\mathfrak{z}$ ij, *Mezerei* concis. $\mathfrak{z}$ vj, *Sacchar*. $\mathfrak{z}$ xij, *Alcohol*, dilut. Oviij. Macerate, with the exception of the sugar, for 14 days; express and filter; evaporate to f $\mathfrak{z}$ xij; and add the sugar. Ph. U. S. 1863.) Dose, a fluidrachm.

EXTRACTUM SATURNI, GOULARD'S, *Liquor plumbi subacetatis*—e. *Scobis Hamatoxyli*, E. *hamatoxyli*.

EXTRACTUM SCAMMONII, *Resina Scammonii, Scammonia Resina*. The resinous cathartic principle of scammony obtained from scammony root by means of alcohol.

EXTRACTUM SENEGÆ ALCOHOLICUM, *Alcoholic Extract of Senega*. (Prepared by displacement from Senega and diluted alcohol;—the tincture being evaporated to the proper consistence. Ph. U. S. 1863.) Dose, gr. v to x.

EXTRACTUM SENNÆ FLUIDUM, *Fluid Extract of Sen'na*, is prepared from senna, by displacement by diluted alcohol, adding sugar, and evaporating. (Ph. U. S. 1863.) Dose, f $\mathfrak{z}$ ss.

EXTRACTUM SERPENTARIÆ FLUIDUM, *Fluid Extract of Serpentaria*. (Prepared by displacement by diluted alcohol.) Ph. U. S. 1863. Dose, f $\mathfrak{z}$ ss.

EXTRACTUM SPIGELLÆ FLUIDUM, *Fluid Extract of Spigelia*. (Prepared from *Spigelia* by displacement by diluted alcohol, and evaporation.) Ph. U. S. 1863.) Dose, f $\mathfrak{z}$ j.

EXTRACTUM SPIGELLÆ ET SENNÆ FLUIDUM, *Fluid Extract of Spigelia and Sen'na*. (A mixture of the fluid extracts of spigelia and senna; carbonate of potassa and oils of anise and caraway. Ph. U. S. 1863.) Dose, f $\mathfrak{z}$ j.

EXTRACTUM STRAMONII, (Ph. U. S. 1863.) *E. Stramonii foliorum, Extract of Stramonium or of Stramonium leaves*. (The expressed juice inspissated.) The *Extractum Stramonii Seminis, Extract of Stramonium Seed*, is made from the powdered seed by means of diluted alcohol, with the aid of the percolator;—the solution being evaporated.

EXTRACTUM STRAMONII ALCOHOLICUM, *Alcoholic*

Extract of Stramonium, (Ph. U. S. 1863.) is made in like manner from the recently dried leaves. These Extracts are used as narcotics in asthma and other spasmodic affections. Dose, of Extractum Stramonii, gr. ij to gr. v, of E. S. Alcoholicum, gr. $\frac{1}{2}$ to gr. i.

EXTRACTUM STYRACIS, see STYRAX.

EXTRACTUM TARAX'ACI, *E. Her'bae et Rad'is Tarax'aci*, *Ex'tract of Dan'delion*. (The strained juice evaporated.) It has been considered deobstruent, laxative, and diuretic. Dose, gr. x to $\mathfrak{z}$ j.

A fluid extract of *Dan'delion*, *Extractum Tarax'aci flu'idum*, is official in the Ph. U. S. for 1863.

EXTRACTUM THEBAICUM, E. Opii.

EXTRACTUM U'VÆ UR'SI, *Ex'tract of U'va Ur'si*. (The Decoction evaporated.) Dose, gr. v to gr. xxx.

A fluid Extract of *Uva Ursi*, *Extractum Uvæ Ursi flu'idum*, is official in the Ph. U. S. for 1863. Dose, $\mathfrak{z}$ j.

EXTRACTUM VALERIA'NÆ, *Ex'tract of Vale'rian*. (The expressed decoction evaporated.) The virtues of the valerian being dependent upon its essential oil, this is an objectionable preparation. Dose, gr. x to $\mathfrak{z}$ j.

EXTRACTUM VALERIANÆ ALCOHOL'ICUM, *Alcohol'ic Extract of Valerian*. (Made from Valerian root, in fine powder, by displacement with diluted alcohol, and evaporation. Ph. U. S. 1863.) Dose, gr. ij to v.

A fluid extract of *Vale'rian*, *Extractum Valerianæ flu'idum*, is in the Pharmacopœia U. S. It is prepared by exhausting the valerian by ether and alcohol, through the percolator, and evaporating. Dose, $\mathfrak{z}$ j to $\mathfrak{z}$ ij.

EXTRACTUM VERA'TRI VIR'IDIS FLU'IDUM, *Fluid Extract of American Hellebore*, is official in the Ph. U. S. for 1863. Dose, $\mathfrak{m}$ v to x.

EXTRACTUM ZINGI'ERIS FLU'IDUM, *Fluid Extract of Ginger*, is prepared by exhausting ginger with alcohol by displacement and evaporation. Dose, $\mathfrak{m}$ x to xv.

EXTRAIT, Extract — *e. des Fruits*, Rob — *e. Hémostatique de Bonjean*, Ergotin — *e. de Kino*, Kino — *e. de Viande*, Osmazome.

EXTRA'NEOUS BOD'Y, *For'eign bod'y*, *Cor'pus extra'neum*, *C. exter'num*, *C. alie'num*, (F.) *Corps étranger*. Any solid, liquid, or gaseous substance, inanimate or animate, proceeding from without, or formed in the body; which constitutes no part of the body, but occupies, in the substance of the textures, or some of the cavities, a place foreign to it.

EXTRAVASA'TION, *Extravasa'tio*, *Extrava'sion*, (*extra*, and *vasa*, 'vessels.') Escape of a fluid — *extravasa'tum* — from the vessel containing it, and infiltration or effusion of the fluid into the surrounding textures. Such fluid is said to be *extrav'asated*, (F.) *Extravasé*.

EXTRAVASATUM, see Extravasatio.

EXTRAVASÉ, see Extravasation.

EXTRAVASION, Extravasation.

EXTREM'ITY, *Extrem'itas*, (F.) *Extrémité*; (*extremus*, 'the outermost.') The end or termination of a thing. The limbs, *acrote'ria*, have been so called, — as the *upper* and *lower extremities*. See Membrum. It has been, also, used to express the last moments of life; as when we say, a patient is in extremity, (F.) *le malade est à l'extrémité*, à toute extrémité.

EXTRIN'SIC, *Extrin'secus*, ('from without.') That which comes from without. This term has been used for muscles, which surround certain organs and attach them to the neighbouring parts; in order to distinguish them from other muscles, which enter into the intimate composition of those organs, and which have been named *intrin'sic*. Thus, there are extrinsic and intrinsic muscles of the tongue, ear, &c.

EXTROVERSIO, (*extra*, and *vertere*, *versum*, to turn.') Extrophia.

EXTUBERANTIA, (*ex*, and *tuber*, 'a prominence.') Protuberance.

EXTUBERATIO, Protuberance.

EXTUMEFAC'TIO, (*ex*, *tumere*, 'to swell,' and *facere*, *factum*, 'to make.') Swelling.

EXTUS'SIO, (*ex*, and *tussis*, 'a cough.') 'I cough with expectoration.'

EXUBER, (*ex*, and *ubera*, 'breasts.') *Apogalac'tus*. A child which has been weaned.

EXUDATION, *Exsuda'tion*, *Exsudat'io*, (*ex*, and *sudare*, *sudatum*, 'to sweat.') The oozing of a material through the pores of a membrane. Also, the material which issues in this manner. See Diapedesis, Exhalation, and Endosmose.

EXUDATION CORPUSCLES, see Corpuscles, exudation.

EXULCERATIO, (*ex*, and *ulcus*, gen. *ulceris*, 'an ulcer.') Ecthlima, Ulceration — *e. Uteri*, Hysterolecosis — *e. Ventriculi*, Gastrelcosis.

EXUMBILICATIO, (*ex*, and *umbilicus*, 'the navel.') Exomphalus.

EXUSTIO, (*ex*, and *urere*, *ustum*, 'to burn.') Cauterization.

EXUTOIRE, Exutory, Fonticulus.

EXUTORIUM, (*exuere*, *exutum*, 'to draw or put out.') Fonticulus.

EXUTORY, Fonticulus.

EXUVIÆ, see Exuviation.

EXUVIA'TION, *Exuvia'tio*, (*exuere*, 'to put off.') The shedding or casting off of parts, as the deciduous teeth, the skins of serpents, &c., which are hence called *Exuvia*.

EYE, (Sax. *ea3*, Teuton. *Auge*,) *Oculus*, *Ops*, *Omma*, *Ophthal'mos*, *Illos*, *Op'tilos* (Doric.) *Vis'orium* *Org'anum*, (Sc.) *Ee*, plural *Een*, (F.) *Œil*. The eye is the immediate organ of vision. It is seated in the orbit, while its dependencies, called by Haller *Tutam'ina Oculi*, occupy the circumference of the cavity, and are composed of the eyebrows, the eyelids, cilia, glands of Meibomius, &c. The *Ball*, *Globe*, or *Bulb of the Eye*, *Bulbus Oculi*, is covered anteriorly by the tunica conjunctiva; is moved by six muscles, four straight, two oblique; and is constituted of membranes, as the *sclerotic*, *cornea*, *choroid*, *tunica Jacobi*, *retina*, *iris*, *hyaloid*, and, in the fetus, the *membrana pupillaris*; and of fluids, called *Humours*, or *Media*, — the *aqueous*, *crystalline*, and *vitreous*. The eyeball is invested with a membranous tunic, which separates it from the other structures of the orbit, and forms a smooth, hollow surface, by which its motions are facilitated. This investment has been called *cell'ular cap'sule of the eye*, *oc'ular capsule*, *tu'nica vagin'al'is oculi*, *vag'in'al coat*, and *submus'cular fascia of the eye*. The vessels of the eye proceed from the ophthalmic artery. The nerves, except the optic, are chiefly furnished from the ophthalmic ganglion. The following are the dimensions, &c., of the organ, on the authority of Petit, Young, Gordon, and Brewster:

Length of the antero-posterior diameter of the eye.....	0.91
Vertical chord of the cornea.....	0.45
Versed sine of the cornea.....	0.11
Horizontal chord of the cornea.....	0.47
Size of pupil seen through the cornea, 0.27 to	0.13
Size of pupil diminished by magnifying power of cornea, from.....	0.12
Radius of the anterior surface of the crystalline.....	0.30
Radius of posterior surface.....	0.22
Principal focal distance of lens.....	1.73
Distance of the centre of the optic nerve from the <i>foramen centrale</i> of Sömmerring.....	0.11
Distance of the iris from the cornea.....	0.10
Distance of the iris from the anterior surface of the crystalline.....	0.02

Field of vision above a horizontal line, 50°	120°
Field of vision below a horizontal line, 70°	
Field of vision in a horizontal plane.....	150°
Diameter of the crystalline in a woman above fifty years of age.....	0.378
Diameter of the cornea.....	0.400
Thickness of the crystalline.....	0.172
Thickness of the cornea.....	0.042

EYE, APPLE, see Melum—e. Apple of the, Pupil.

EYE BALL, see Eye—e. b. Anæmic protrusion of the, see Exophthalmia.

EYE BALM, Hydrastis Canadensis—e. Breck, Palpebra—e. Breen, see Supercilium—e. Bright, Euphrasia officinalis, Lobelia Veronica—e. Brow, Supercilium.

EYE, CAT'S, Amaurotic, see Amaurotic—e. Cellular capsule of the, see Eye.

EYE, CYLINDRICAL. An eye is so called in which the rays of light, on entering the organ, are refracted to a nearer focus in a vertical than in a horizontal plane.

EYE DROP, Tear.

EYE GLASS, Scaphium oculare, see Spectacles. Also, a glass adapted for the application of collyria to the eye.

EYE, GUM OF THE, *Chuskie*.

EYE LASHES, Cilia—e. Lid, Palpebra—e. Lid, granular, Trachoma.

EYE, MELON, see Melum—e. Purulent, Ophthalmia, purulent, of infants.

EYE ROOT, Bignonia ophthalmica—e. Salve, Singleton's, Unguentum Hydrargyri nitrico-oxydi—e. Salve, Smellome's, see Cupri subacetat.

EYE, SIGHT OF THE, Pupil.

EYE STONE. The shelly operculum of small turbinides. Used at Guernsey to get things out of the eyes. Being put into the inner corner of the eye, under the eyelid, it works its way out at the outer corner, and brings out any foreign substance with it.

EYE TEETH, Canine Teeth—e. of Typhon, Scilla—e. Vine, Bignonia ophthalmica—e. Water, Collyrium—e. Water, blue, Liquor cupri ammoniati—e. Water, common, Liquor zinci sulphatis cum camphorâ.

EYE, WATERY, Epiphora—e. White of the, see Sclerotic.

F.

FABA, ('a bean,') Phaseolus, Vicia faba—f. Egyptiaca, Nymphaea nelumbo—f. Cathartica, Jatropha curcas—f. Crassa, Sedum telephium—f. Febrifuga, Ignatia amara—f. Græca, Diospyros lotus—f. Indica, Ignatia amara—f. Major, Vicia faba—f. Pechurei, Pichurim Beans—f. Pichurim, Pichurim Beans—f. Purgatrix, Ricinus communis—f. Sancti Ignatii, Ignatia amara—f. Suilla, Hyoscyamus—f. Vulgaris, Vicia faba.

FABÆ, Onisci aselli.

FABAGELLE, Zygophyllum fabago.

FABARIA CRASSULA, (from *faba*), Sedum telephium.

FABRICA ANDROGYNA, ('androgynous make,') Hermaphrodity.

FACE, *Facies*, *Vultus*, *Vultus*, *Prosoπον*, *Stoma*, (F.) *Face*, (*facere*, 'to make or form.') The face is the anterior part of the head. It is formed of 13 bones,—the two superior maxillary, the two malar, the two ossa nasi, the two ossa unguis, the vomer, the two ossa spongiosa inferiora, the two palate bones, and the inferior maxillary, without including the frontal portion of the os frontis, and the 32 teeth, which may be considered to form part of it. Its numerous muscles are chiefly destined for the organs of sight, hearing, taste, and smell. Its arteries proceed from the external carotid; its veins end in the jugular, and its nerves draw their origin immediately from the brain.

The face experiences alterations in disease, which it is important to attend to. It is yellow in jaundice; pale and puffy in dropsy; and its expression is very different according to the seat of irritation, so that, in infants, by an attention to medical physiognomy, we can often detect the seat of disease. Hippocrates has well depicted the change which it experiences in one exhausted by long sickness, by great evacuations, excessive hunger, watchfulness, &c., threatening dissolution. Hence this state has been called *Facies Hippocratica*, *Facies Cadaverica*, *F. Tortuata*. In this, the nose is pinched; the eyes are sunk; the temples hollow; the ears cold, and retracted; the skin of the forehead tense, and dry; the com-

plexion livid; the lips pendent, relaxed, and cold, &c.

The term *Face* (F.) is likewise given to one of the aspects of an organ; thus, we say, the superior face of the stomach.

FACE AGUE, Neuralgia, facial.

FACE INJECTÉE, see *Vultueux*—f. *Vultueuse*, see *Vultueux*.

FACIET, (F.) *Facette*, (dim. of *Face*.) *Facicula*. A small face. A small, circumscribed portion of the surface of a bone, as the articulation of a bone.

FACHINGEN, MINERAL WATERS OF. These springs are at no great distance from those of Geilenau, and two miles north of Wisbaden. They contain free carbonic acid; carbonate, sulphate, and phosphate of soda; chloride of sodium, carbonate of lime, magnesia, and iron.

FA'CIAL, *Facialis*, (from *facies*.) Belonging to, or connected with, the face. See Facial vein.

FACIAL ANGLE, see Angle, facial.

FACIAL ARTERY, *Labial artery*, *Antiquar* or *external maxillary artery*, *A. palato-labial*—(Ch.), is a branch of the external carotid, which rises beneath the digastricus, and is distributed to almost every part of the face. It furnishes the inferior palatal, submental, superior labial, inferior labial, and dorsalis nasi. Chaussier calls the external carotid artery, *Facial Artery*.

FACIAL LINE, see Angle, facial.

FACIAL NERVE, *Ramus du'rior septimæ conjugationis*, *Sympatheticus minor*, *Par septimium seu faciale*, *Communicans faciei nervus*, *Portio du'ra of the 7th pair*, *Respiratory nerve of the face*. This nerve arises from the inferior and lateral part of the tuber annulare, in the groove which separates it from the medulla oblongata, external to the corpora olivaria, and by the side of the auditory nerve. It issues from the cranium by the *meatus auditorius internus*; enters the aqueduct of Fallopius; receives a branch of the Vidian nerve; forms a gangliform swelling—*Intumescencia gangliformis*;—sends off filaments to the internal muscles of the malleus and stapes; furnishes, according to many anatomists, that

called *Clorda Tympani*; makes its exit at the foramen stylo-mastoideum, and divides into two branches—the *temporo-facial*, and *cervico-facial*. On the face it is termed *Pes anserinus*, *Plexus nervorum anserinus*. See Portio Wrisbergii.

FACIAL VEIN, *Palato-labial*—(Ch.), arises between the skin and frontalis muscle, and bears the name *Vena Frontalis*, *V. Prepara'ta*, (F.) *Veine frontale* ou *V. Préparée*. It then descends, vertically, towards the greater angle of the eye, where it is called *Angulà'ris*; and afterwards descends, obliquely, on the face, to open into the internal jugular, after having received branches, which correspond with those of the facial artery. It is only in the latter part of its course that it is called *Facial Vein*.

FACIECULA, (dim. of *Facies*), Facet.

FACIENT, (*faciens*, 'making,' *facio*, 'I make,') A suffix, as in *Calefacient*, *Rubefacient*, &c., 'warm making,' 'red making.'

FACIES, Face—f. Cadaverica, see Face—f. *Concava pedis*, Sole—f. *Hippocratica*, see Face—f. *Inferior pedis*, Sole—f. *Tortualis*, see Face.

FACTICE, Factitious.

FACTITIOUS, *Factiti'us*, (F.) *Factice*, (*facere*, *factum*, 'to make,') Artificial. That which is made by art, in opposition to what is natural, or found already existing in nature. Thus, we say, *factitious mineral waters* for artificial mineral waters.

FACULTAS, Faculty—f. *Auctrix*, Plastic force—f. *Formatrix*, Plastic force—f. *Nutrix*, Plastic force—f. *Vegetativa*, Plastic force—f. *Zotica*, Vis vitalis.

FACULTATES NATURALES, ('natural faculties,') see Function.

FACULTY, *Facultas*, *Dy'namis*, *Vir'tus*, *Po'tency*, *Pow'er*, *Vir'tue*, (F.) *Faculté* (*facilis*, (itself from *facere*, 'to make,') 'what can be made or done,') The power of executing any function or act. The collection of the intellectual faculties constitutes the understanding. We say, also, *vital faculties*, for *vital properties*, &c. Faculty likewise means the whole body of the medical profession, and, also, a body of medical or other professors.

FÆCAL, (from *fæces*,) *Stercoraceous*—f. *Retention*, Constipation.

FÆCES, (pl. of *Fæx*), *Feces*, *Ossææ*, 'the dregs of any thing,') *Fæculence*, (F.) *Fèces*. The alvine evacuations are so called; (F.) *Garderober*; the excrements, *Impuritates alvinae*, *Fæcal matter*, (F.) *Matière fécale*. See Excrement.

FÆCES INDURATÆ, ('indurated feces,') *Scybala*.

FÆCOSITAS, (*fæx*, gen. *fæcis*), *Fæculence*.

FÆCULA, *Fecula*.

FÆCULENTIA, *Fæculence*.

FÆX, *Fæculence*. See *Fæces*.

FAGARA OCTAN'DRA, (*fagus*, 'the beech,' which it resembles,) *Xanthoxylum octan'dra*, *Elaph'rium tomento'sum*, *Am'yris tomento'sa*. Ord. Rutaceæ. The systematic name of the plant, which affords *Tacamahac*, a resinous substance, that exudes from the tree *Tacamahaca*, which has a fragrant, delightful smell, was formerly in high estimation, as an ingredient in warm, stimulating plasters, and was given internally, like the balsams generally. The *East India Tacamahac*, *Balsamum Vir'idè* seu *Cal'aba* seu *Ma'rie*, *O'leum Ma'rie*, (F.) *Bume de Calaba*, *Bume vert*, *B. Marie*, is yielded by *Calophyllum inophyllum* *Balsamaria inophyllum*.

The name *Tacamahac* is also given to a resin furnished by *Populus balsamifera* seu *tacamahaca*, (F.) *Peuplier baumier* ou *de la Caroline*, which grows in the northern parts of America and Siberia.

FAGARA PIPERITA, *Xanthoxylum piperitum*, *Jap'aneæ pepper*, (F.) *Fagariæ poirè*, a native of Japan, possesses the qualities of pepper, and is

used as such by the Japanese. It is also, employed as a rubefacient cataplasm.

FAGARAS'TRUM CAPEN'SE, (after *Fagara*.) *Nat. Ord. Xanthoxylæ*. A South African plant, the fruit of which is known to the Colonists as *wild Card'amom*; and, on account of its aromatic qualities, is prescribed in flatulency and paralysis.

FAGARIER POIVRÈ, *Fagara octandra*.

FAGOPYRUM, (*fagus*, 'the beech,' and *pyros*, 'wheat,') *Polygonum fagopyrum*.

FAGUS, (*φῡς*, from *φαγεω*, 'to eat,' [?] F. *sylvatica*).

FAGUS CASTA'NEA. The systematic name of the *Chestnut Tree*; *Castanea*, *C. vulgaris* seu *ves'ca*, *Lo'pinda*, *Glans Jo'vis Theophrast'i*, *Ju'piter's* or *Sardin'ian A'corn*; *Com'mon Chestnut*, (F.) *Châtaigner commun*, of the Oak Family. Ord. Cupuliferae. *Sex. Syst.* Monocæcia Polyandria. The *Chestnut*, *Castanea nux*, (F.) *Châtaigne*, is farinaceous and nutritious, but not easy of digestion.

FAGUS CASTA'NEA PU'MILA. The *Chin'capin* or *Chin'quapin*, *Castanea Pu'mila*, (F.) *Châtaigner nain*. The nut of this American tree is eaten like the chestnut. The bark, *Castanea* (Ph. U. S.), has been used in intermittents.

FAGUS PURPUREA, F. *sylvatica*.

FAGUS SYLVATICA. The systematic name of the *Beech*, *Fagus*, F. *Sylvestris* seu *purpurea*, *Ox'ya*, *Balan'da*, *Valan'ida*; the *Beech Tree*, (F.) *Hêtre*. The *Beech-nut* or *Beech-mast*, (F.) *Faine*, affords an oil, by expression, which is of a palatable character, and is eaten in some places instead of butter. It has been supposed to be a good vermifuge, but it is no better than any mild oil.

FAGUS SYLVESTRIS, F. *sylvatica*.

FAIBLESSE, (from *febilis*, 'deplorable,' or from *debilis*, 'weak,' [?]) Debility.

FAIM, (from *fames*,) Hunger—f. *Canine*, *Boulimia*—f. *de Loup*, *Fames lupina*.

FAINE, see *Fagus sylvatica*.

FAINT, (from *Sax. fȳnigean*, 'to decay,') see Syncope.

FAINTING, Syncope—f. *Fit*, Syncope.

FAINTNESS, see Syncope.

FAINTNESS, *Languor*, Syncope.

FAIRBURN, MIN'ERAL WATERS OF. The mineral waters at this place, which is in the county of Ross, in Scotland, are sulphureous, and frequented.

FAIRNTICKLED, see *Ephelides*.

FAISANDÉES (VIANDES), (from *faisan*, 'a pheasant,' hence, game in general kept until it is putrescent.) See *Hyposaprus*.

FAISCEAU, *Fasciculus*—f. *Intermédiaire de Wrisberg*, *Portio Wrisbergii*—f. *Petit*, *Fasciculus*.

FAIX, ('a burthen,' from *fascia*, 'a bundle,') *Fœtus*.

FALCADINA, see *Scherlievo*.

FALCES, (pl.), see *Falx*.

FAL'CIFORM, *Falciform'is*, *Drepanoï'des*, (*falx*, gen. *falcis*, 'a scythe,' and *forma*, 'shape,') Having the shape of a scythe. This term has been applied to different parts. See *Falx*, and *Sinus*.

FALCIFORM EXPAN'SION OF THE FAS'CIA LA'TA is the scythe-shaped reflection of the *fascia lata*, which forms, outwards and upwards, the opening for the vena saphæna, and is attached to the crural arch by its superior extremity, forming the anterior paries of the canal of the same name.

FALLACIA, (*fallax*, 'deceitful,') *Hallucination*.

FALLACIA OPTICA, An optic illusion.

FALLAND-EVYL, *Epilepsy*.

FALLING DOWN, *Epilepsy*.

FALLING SICKNESS, *Epilepsy*.

FALLOPIAN TUBE, see *Tube*, *Fallopian*.

FALLOPOISON, *Amianthum musætoxicum*.

FALLTRANCK, *Faltranck* (G.), literally,

a drink against falls. A vulnerary. It is a mixture of several aromatic and slightly astringent plants, which grow chiefly in the Swiss Alps, and hence the name—*Vulnéraire Suisse*—given to such dried plants cut into fragments. They are called, also, *Espèces Vulnéraires*, and *Thé Suisse*. Within the present century, in England, a kind of vulnerary beer was often prescribed, in country practice, in all cases of inward bruises. It bore the name *Cerevisia nigra*, or *black beer*, and was formed by infusing certain reputed vulnerary herbs in beer or ale.

The infusion of the *Falltranck* is aromatic, and slightly agreeable, but of no use in the cases for which it has been particularly recommended.

FALMOUTH, CLIMATE OF. The climate of Falmouth, in Cornwall, England, resembles that of Penzance; and, like it, is in many respects, a favourable retreat for the phthisical during the winter months.

FALSA VIA, False passage.

FALSE, Fal'sus, No'thus, Pseudo' (in composition,) *Spurious, Bas'tard*, (F.) *Faux, Fausse*, (from *fallere, falsum*, 'to deceive.') This epithet has been frequently added to peripneumony, pleurisy, &c., to designate a disease similar to these, but less severe. Most commonly, a severe catarrh or pleurodynia has received the name. See *Peripneumonia notha*.

FALSE PASS'AGE, Fal'sa Via, (F.) Fausse Route. An accidental passage, made in surgical operations, and particularly in introducing the catheter. The catheter is sometimes passed through into the rectum.

FALSE WATERS, False Deliv'ery, (F.) Hydralante. Water, which sometimes collects between the amnion and chorion, and is commonly discharged before the birth of the child.

We say, also, *False Ribs, False Rhubarb, &c.*

FALSETTO VOICE, (I.] dim of falso, 'false,') see Voice.

FALSIFICA'TION, Adultera'tio, (falsus, 'false,' and facere, 'to make.') A fraudulent imitation or alteration of an aliment or medicine by different admixtures. It is synonymous with *adultera'tion* and *sophistica'tion*. By Dr. Hassall, the practice of adulteration is defined as the intentional addition to an article, for purposes of gain or deception, of any substance or substances, the presence of which is not acknowledged in the name under which the article is sold.

The following table was prepared to indicate common falsifications practised with some of the most useful drugs. Of late years, increased attention has been drawn to the alarming extent to which the adulterations of both drugs and aliments have been carried. The object has been seriously examined and reported on by a committee of the British House of Commons, and two elaborate works have been published by Dr. Hassall,—the one "On Food and its Adulterations," (1855); and the other entitled "Adulterations Detected," (1857), both in food and medicines, comprising the Reports of the Analytical Sanitary Commission of "the Lancet," &c. &c., (1857.) The principal means for the detection of such adulterations are afforded by chemistry and the microscope. The former has always been employed for the purpose; but the latter only recently. It is especially adapted for the detection of animal and vegetable admixtures, on the presence of which chemistry can shed little or no light. The distinctive characters of many of the pure articles employed in medicine are given in the *Pharmacopœias*, as in that of the United States (1833.)

MEDICINES.	ADULTERATIONS.	MODE OF DETECTION.
ACACIE GUMMI. <i>Gum Arabic.</i>	{ <i>Gum Senegal</i>	{ Gum Senegal is clammy and tenacious. Gum Arabic is perfectly soluble in water, and its solution limpid.
ACETUM DESTILLATUM. <i>Distilled Vinegar.</i>	{ <i>Sulphuric Acid</i>	{ Acetate of barytes causes a white precipitate.
	{ <i>Nitric Acid</i>	{ By evaporating it, the residuum deflagrates, when thrown on burning charcoal.
	{ <i>Copper</i>	{ Supersaturate with ammonia—a blue colour is produced.
	{ <i>Lead</i>	{ Sulphuretted hydrogen causes a dark precipitate.
ACIDUM ARSENIOSUM. <i>Arsenious Acid.</i>	{ <i>Chalk, Sulphate of Lime, Sulphate of Baryta.</i>	{ Not volatilizable by heat.
ACIDUM CITRICUM. <i>Citric Acid.</i>	{ <i>Tartaric and Oxalic Acids</i>	{ Their presence is indicated by forming a granular sediment in a concentrated solution of a neutral salt of potassa.
ACIDUM MURIATICUM. <i>Muriatic Acid.</i>	{ <i>Sulphuric Acid</i>	{ Deposits by evaporation the salts it may contain; precipitates with solution of hydrochlorate of baryta if it contains sulphuric acid.
ACIDUM NITRICUM. <i>Nitric Acid.</i>	{ <i>Muriatic and Sulphuric Acids.</i>	{ The presence of chlorine is indicated by a precipitate with nitrate of silver: that of sulphuric acid by the same result with hydrochlorate of baryta.
ACIDUM SULPHURICUM. <i>Sulphuric Acid.</i>	{ <i>Muriatic and Nitric Acids</i>	{ The presence of muriatic and nitric acid is indicated by the smell, when the acid tested is strongly heated.
ÆTHER RECTIFICATUS. <i>Rectified Ether.</i>	{ <i>Too dilute.</i>	{ The s. g. detects this.
	{ <i>Sulphuric Acid</i>	{ By acetate of baryta. Precipitate white.
	{ <i>Alcohol</i>	{ With phosphorus a milky instead of limpid solution is formed.
AMMONIÆ AQUA. <i>Water of Ammonia.</i> AMMONIÆ CARBONAS. <i>Carbonate of Ammonia.</i>	{ <i>Carbonic Acid</i>	{ A precipitation occurs on adding a solution of muriate of lime. It should be capable of complete volatilization by heat.
AMMONIACUM. <i>Ammoniac.</i>	{ <i>The Guttæ Ammoniaci</i> are white, clear, and dry. <i>The lump Ammoniacum, lapis Ammoniaci</i> , is often adulterated with common resin.	
ANTIMONII PHURETUM. <i>Sulphuret of Antimony.</i>	{ <i>Lead.</i> <i>Arsenic</i>	{ It ought to be entirely volatilized by a red heat. Imparts a foliated texture, and is not vaporizable. A smell of garlic is emitted when thrown on live coals.
	{ <i>Manganese and Iron.</i>	{ Are not vaporizable.

MEDICINES.	ADULTERATIONS.	MODE OF DETECTION.
ARGENTI NITRAS <i>Nitrate of Silver.</i>	Copper	The solution assumes a blue colour, when supersaturated with ammonia. It may be suspected when the salt deliquesces.
	Nitrate of Potassa.....	The adulteration with nitrate of potassa is easily recognized by the fracture of a stick of it, which is radiated when pure, and granular if adulterated; or by precipitating a solution of the salts with a sufficient quantity of muriatic acid, and evaporating the clear liquor: the nitrate of potassa or other salts will remain.
BALSAMUM PERUVIANUM. <i>Balsam of Peru.</i>	A mixture of Resin and some Volatile Oil, with Benzoin.	Not easily detected.
CALUMBA. <i>Columbo.</i>	A menisperm from Ceylon, tinged Bryony root, and the root of <i>Frasera Walleri</i> .	The true is distinguishable from the false Columbo by adding to an infusion of the root, a few drops of solution of sulph. iron, which gives to the infusion of the false Columbo a greenish black colour; but produces no change in the other.
CAPSIUM. <i>Cayenne Pepper.</i>	Chloride of Sodium	This disposes it to deliquesce.
	Red Lead.....	Digest in acetic acid, and add a solution of sulphuret of ammonia—a dark-coloured precipitate will be produced.
CARTOREUM. <i>Castor.</i>	A mixture of dried blood, gum ammoniac, and a little real castor, stuffed into the scrotum of a goat.	Smell and taste will generally detect the fraud.
CERA ALBA. <i>White Wax.</i>	White Lead.....	Melt the wax, the oxide will subside.
	Tallow.....	The cake has not its ordinary translucency.
	Earth or Peasmeal.....	May be suspected when the cake is brittle and colour grayish.
CERA FLAVA. <i>Yellow Wax.</i>	Resin.....	Put it in cold alcohol, which will dissolve the resin, without acting on the wax.
	Tallow.....	Is known by the greater softness and unctuousity, and its smell when melted. Turmeric is generally added in this case to obviate the paleness.
CINCHONA. <i>Bark.</i>	This is variously adulterated, but generally with the <i>Carthagena</i> and other inferior barks.	Can only be detected by practice, and examining into the quantity of quinia, or cinchona, or both, which it contains.
COCCUS. <i>Cochineal.</i>	Pieces of dough formed in moulds, and coloured with cochineal.	Throw it into water, the adulteration will appear.
COPAIBA. <i>Balsam of Copaiva.</i>	Fixed Oils.	If it does not retain its spherical form when dropped into water, its adulteration may be inferred.
	Fibres of smoked Beef	Mix one part of strong liquid ammonia of 22°, with three parts of copaiba. If pure, the mixture will, in a few minutes, become transparent; if not, it will remain opaque.
CROCUS. <i>Saffron.</i>	Petals of <i>Calendula officinalis</i> , and <i>Carthamus Tinctorius</i> .	Affords an unpleasant odour when thrown on live coals.
CUBEBA. <i>Cubebs.</i>	Turkey Yellow Berries, or the dried fruit of <i>Rhamnus Catharticus</i> .	Infuse the specimen in hot water, and the difference will be perceptible.
CUSPARIE CORTEX. <i>Angustura or Cusparia Bark.</i>	False <i>Angustura</i> sometimes sold for it.	Detected by attentive examination.
	Common Resin.....	The epidermis of the true <i>Cusparia</i> is characterized by being covered with a matter resembling the rust of iron. False <i>cusparia</i> bark yields brucia and strychnia; the true yields neither.
GUAIACA RESINA. <i>Resin of Guaiacum.</i>	Manchineel Gum.....	Detected by the turpentine smell emitted when thrown upon hot coals.
	Corrosive Sublimate and Sub-nitrate of Bismuth.	Add to the tincture a few drops of the spirit of nitre, and dilute with water; the guaiacum is precipitated—the adulteration floats in white striae.
HYDRARGYRI CHLORIDUM MITE. <i>Calomel.</i>	Carbonate of Lead, Sulphate of Baryta, and Carbonate of Baryta.	A precipitation will be produced by the carbonate of potassa, from a solution made by boiling the suspected sample with a small portion of muriate of ammonia in distilled water; or, the presence of deutochloride of mercury is indicated by warming gently a small quantity of calomel in alcohol, filtering and adding to the clear liquor some limewater, by which a reddish yellow precipitate is afforded. When calomel is rubbed with a fixed alkali, it ought to become intensely black, and not exhibit any orange hue.
	Red Lead, Red Oxide of Iron, Brick Dust.	Not volatilized by heat,—the carbonates effervesce with acids.
HYDRARGYRI OXIDUM RUBRUM. <i>Red Oxide of Mercury.</i>	Ivory Black.....	Digest in acetic acid: add sulphuret of ammonia, which will produce a dark coloured precipitate. It should be totally volatilized by heat.
HYDRARGYRI SULPHURETUM NIGRUM.	Red Lead, Red Oxide of Iron, Brick Dust.	Throw a suspected portion on hot coals—the residuum will detect the fraud.
HYDRARGYRI SULPHURETUM RUBRUM. <i>Red Sulphuret of Mercury.</i>	Plumbago, Charcoal, and Oxide of Manganese.	Digest in acetic acid, and add sulphuret of ammonia—a black precipitate will be produced. Does not wholly sublime like the genuine article.
IODINIUM. <i>Iodine.</i>	Bryony Root, spurious or false Jalap Root, and Liquorice Root.	The tests of its purity are—that it is freely soluble in ether and alcohol. Heated on a piece of glass or porcelain, it sublimes without residuum, rising in purple vapour. With starch in cold solution it produces a blue colour.
JALAPÆ RADIX. <i>Jalap Root.</i>		Bryony root is of a paler colour, and less compact texture, and does not easily burn at the flame of a candle. Liquorice is detected by the taste.

MEDICINES.	ADULTERATIONS.	MODES OF DETECTION.
MAGNESIA.	<i>Lime</i> <i>Sulphuret of Lime</i>	Detected by the solution in dilute sulphuric acid affording a precipitate with oxalate of ammonia. Gives off, when moistened, the smell of sulphuretted hydrogen.
MAGNESIE CARBONAS. Carbonate of Magnesia.	<i>Chalk</i> <i>Gypsum</i>	Detected by adding dilute sulphuric acid to the suspected substance, when, if chalk be present, there will be a white insoluble precipitate. Boil in distilled water, and test the solution by a barytic and oxalic reagent.
MANNA.	<i>A factitious article, consisting of honey or sugar, mixed with scammony, is said to be sometimes sold for it.</i>	The colour, weight, transparency, and taste detect it.
MORPHIA ET EJUS SALES. Morphia, and its Salts.		Morphia and its salts, when placed in contact with nitric acid, are coloured red; with persalts of iron, blue. They are perfectly soluble in warm alcohol, and acidulated warm water. When morphia is mixed with narcotina, the adulteration is ascertained by mixing them with sulphuric ether, which dissolves the narcotina, without sensibly affecting the morphia. The bag must not appear to have been opened. This may be suspected, if it emits a fetid smoke when inflamed.
MOSCHUS. Musk.	<i>Dried Blood</i> <i>Asphaltum</i> <i>Fine particles of Lead</i>	Discovered by its melting and running before it inflames. Rub with water. The metallic particles will subside.
OLEA DESTILLATA. Essential Oils.	<i>Fixed Oils</i> <i>Alcohol</i>	Touch writing paper with it, and hold it before the fire: fixed oil leaves a stain of grease. Add water; a milkiness and increase of temperature occurs.
OLEUM RICINI. Castor Oil.	<i>Olive or Almond or Poppy Oil</i> <i>Extract of Liquorice, Bull's and Snouts sometimes; Extract of Poppy, of Chelidonium majus, and other vegetables. Melissa, G. Arabic, G. Tragacanth, Linseed Oil, Owl's Dung.</i>	Alcohol S. g. 820 will mix with any proportion of castor oil, whilst it dissolves very little of the others.
OPIUM.		The best opium is covered with leaves and the reddish capsules of a species of Rumex. The inferior kinds have capsules adherent. It is bad when soft and friable, when intensely black or mixed with many impurities, and when sweet. The quantity of morphia affords the best test.
POTASSII IODIDUM. Iodide of Potassium.	<i>Chlorides of Potassium, and Sodium, Nitrate and Carbonate of Potassa.</i>	The adulteration is ascertained by precipitating a solution of the salt with nitrate of silver, and treating the precipitate with ammonia, which dissolves the chloride of silver, without acting upon the iodide of this metal.
QUINIE SULPHAS. Sulphate of Quinia.	<i>Mannite, Sugar of Milk, Starch, Chalk.</i>	The genuine salt leaves no residue when submitted to calcination: it is perfectly soluble in warm alcohol, and in water slightly acidulated with sulphuric acid.
STRYCHNIE ET EJUS SALES. Strychnia and its Salts.	<i>Brucea</i>	By the red color produced by contact with nitric acid.
ZINCI OXYDUM. Oxide of Zinc.	<i>Chalk and Carbonate of Magnesia.</i> <i>White Lead</i>	Sulphuric acid excites an effervescence. Sulphuric acid forms an insoluble sulphate of lead.

The following table by Dr. Hassall is a classified list of various substances ascertained by him to be employed for the adulteration of different dietetic articles,—to imitate bulk and weight, and give colour, smell, or taste, or other proper-

ties. It sufficiently exhibits the considerable number of substances possessing qualities more or less injurious, which are employed in the adulteration of food.

	For Bulk and Weight.	For Colour.	For taste, Smell, and other Properties.
ARROW-ROOT.....	Sago, Potato, and Tapioca Starches, and various mixtures and combinations of these with the inferior arrow-roots.		
ANCHOVIES	Dutch, French and Sicilian Fish.	Ble Armenian, Venetian Red.	
BREAD	Mashed Potatoes.....		Alum, Hards and Stuff.*
BUTTER	Water		
BOTTLED FRUITS AND VEGETABLES.		Certain Salts of Copper, usually the Acetate.	Salt.
CINNAMON.	Cassia, and most of the articles mentioned under Spices.		
COLOURED CONFECTIONERY.	East India Arrow-root, Wheat and Potato Flour, Hydrated Sulphate of Lime.	Cochineal, Lake, Indigo, Prussian Blue, Antwerp Blue, Artificial Ultramarine, Carbonate of Copper or Verditer, Carbonate of Lead or White Lead, Red Lead, Vermilion; Chrome Yellow or Chromates of Lead; Lemon, Orange, and deep; Gamboge; the three Brunswick Greens, Emerald Green or Arsenite of Copper, Indian red; brown ferruginous earths, chiefly Umber, Sienna, and Vandyke Brown, and various combinations of the above pigments.	

* 'Hards' and 'Stuff' are mixtures of alum and salt used by bakers to whiten bread.

	For Bulk and Weight.	For Colour.	For Taste, Smell, and other Properties.
COFFEE.....	Chicory, Roasted Wheat, Rye and Potato Flours, Roasted Beans, Mangel-wurzel, Acorns.	Burnt Sugar or Black Jack.	
CHICORY.....	Roasted Wheat and Rye Flours, Burnt Beans and Acorns, Burnt Sugar, Sawdust, Mahogany Sawdust, Carrot, Mangel-wurzel.	Ferruginous earths, as Venetian Red and Umber, Burnt Sugar or Black Jack.	
COCOA AND CHOCOLATE.....	Maranta, East India, and Tacea or Tahiti Arrow-roots; Tous les Mois; the Flours of Wheat, Indian Corn, Sago, Potato, and Tapioca, and various Mixtures of these; Sugar, Chicory.	Venetian Red, Red Ochre, and other ferruginous earths.	
CATENNE PEPPER.....	Ground Rice, Mustard Husk, Deal Sawdust, Salt.	Red Lead, Vermilion or Bisulphuret of Mercury, Venetian Red, Turmeric	
CUSTARD AND EGG POWDERS.....	Wheat, Potato, and Rice Flours....	Chrome Yellow, or Chromate of Lead, Turmeric.	
CURRY POWDER.....	Ground Rice, Potato-farina, Salt....	Red Lead.....	Salt.
FLOUR.....	Wheat, Sago, and Potato Flours.	Turmeric Powder.	Alum.
GINGER.....	Ground Rice, Mustard Husks.		
GIN.....	Water, Sugar.		Cayenne, Cassia or Cinnamon, Sugar, and flavouring of different kinds. For fining, Alum, Salt of Tartar.
ISINGLASS.....	Gelatin.		
LARD.....	Potato-flour, Water.....		Salt. Carbonate of Soda, Caustic Lime.
MUSTARD.....	Wheat-flour, Turmeric.....	Turmeric.	
MILK.....	Water.....	Annatto.	
MARMALADE.....	Pulp of Apple or Turnip.		
OATMEAL.....	Barley-flour, and the integuments of Barley called Rubble.		
PORTER AND STOUT.....	Water.....	Sugar, Treacle.....	Sugar, Treacle, Salt.
PICKLES.....	Flour, probably Wheat-flour boiled.	Salts of Copper, usually the Acetate of Copper.	
POTTED MEATS AND FISH.....	Salts of Copper, including the Acetate.	Bole Armenian, and sometimes Venetian Red.	
PRESERVES.....	Wheat and Pea Flour, Ground Rice, Ground Mustard Seeds, Linseed Meal, P. D. or Pepper Dust.		
SNUFF.....		The Chromates of Potash, Chromate of Lead, Ferruginous earths, chiefly Umbers, Red and Yellow Ochre, Red Lead, or Oxide of Lead.	The Chromates of Potash, Carbonate of Ammonia, Lime, Powdered Glass or Silica, Powdered Orris-root.
SUGAR.....	Wheat-flour in two cases only, Potato-flour, and Tapioca-starch.		
SPICES:			
CLOVES.....	Powdered Clove-stalks in one case.		
CINNAMON.....	Cassia, Wheat-flour, Sago-meal, and mixtures of these; East India Arrow-root, Potato-flour.		
PIMENTO.....	Mustard Husk in one instance.		
MIXED SPICE.....	Wheat, Sago, and Potato Flours, Ground Rice, Two Vegetable Substances, one of which resembled Linseed.		
SAUCES, as the Essences of Anchovies, Lobsters, and Shrimps, and Tomato Sauce.		Red ferruginous earths, as Bole Armenian and Venetian Red.	
TEA.....	Exhausted Tea Leaves; Leaves, other than those of Tea, British and Foreign.—as, amongst the former, those of Sycamore, Horse-chestnut, and Plum; Lie Tea,* Paddy Husk, Sand, Starch.	Plumbago, or Black Lead, Gum, Indigo, Prussian Blue, Turmeric, Chinese Yellow, China Clay, Soapstone or French Chalk.	Sulphate of Iron, Catechu, Gum, La Veno Beno, Chinese, Botanical Powder.†
TOBACCO.....	Water, Sugar, Treacle, and Salts....	Sugar, Treacle.....	Oil.
VINEGAR.....	Water.....	Burnt Sugar.....	Sulphuric Acid.

* 'Lie Tea' is so termed from being spurious. It is formed of the dust of tea leaves, and at times of other leaves and sand made into small masses by means of starch or gum; which are painted and coloured so as to resemble either black or green gunpowder tea.

† Called, also, the 'Chinese Tea Improver,' is said to be a mixture of a very small portion of tea powder, with upwards of 90 per cent. of catechu. The 'Chinese Botanical Powder' is an imitation of this.

FALTRANCK, Falltranck.

FALX, gen. *Falcis*, ('a scythe.') Anatomists have given this name to several membranous reflections having the shape of a falx or scythe.

FALX CEREBEL'LI, *Falcis minor*, *Septum Cerebelli seu parvum occipitale*, *Processus falciformis Cerebelli*, (F.) *Faux du cervelet*, *Septum medianum du cervelet*, (Ch.), is a triangular process of the dura mater opposite the internal occipital protuberance. Its base is attached to the middle of the tentorium, and its top or apex bifurcates, to proceed to the sides of the foramen magnum. Its convex surface is towards the cranium, and its concave in the fissure or groove, which separates the two lobes of the cerebellum.

FALX CEREBRI, *F. major*, *Vertical superior longitudinal process*, *Mediastinum cerebri*, *Septum Cerebri*, *Processus falciformis duræ matris*, (F.) *Faux du cerveau*, *Repli longitudinal de la méninge*, (Ch.). The greatest process of the dura mater. It extends from the fore to the hind part of the skull, on the median line; is broad behind and narrow before, and is lodged in the groove which separates the hemispheres from each other—the *interlobular fissure*. At its superior part is situated the longitudinal sinus (*superior*), and at its lower, corresponding to the edge of the scythe, the *inferior longitudinal sinus*. Its anterior extremity is attached to the crista galli; its posterior is continuous with the tentorium cerebelli, and contains the straight sinus.

FALX MAJOR, Falx cerebri—f. Minor, Falx cerebelli.

FALX OF THE PERITONE'UM, GREAT, *F. peritonei maxima*, (F.) *Grande faux du péritoine*, *Faux de la Veine Omphalique*, *Falz of the umbilical vein*, is a reflection of the peritoneum, which ascends from the umbilicus to the anterior and inferior surface of the liver.

FALCES OF THE PERITONE'UM, LESS'ER, *Falces Peritonei minores*, (F.) *Petites faux du péritoine*, are the lateral ligaments of the liver, and the reflections which the peritoneum forms, raised up by the umbilical arteries.

FALX OF THE UMBILICAL VEIN, Falx, great, of the Peritoneum.

FAME, Lancet.

FAMELIUS, (from *fames*), Hungry.

FAMELICA FEBRIS, Fever accompanied with insatiable hunger.

FAMES, Hunger—f. Bovina, Boulimia—f. Canina, Boulimia.

FAMES LUPINA, ('wolf hunger,') *Lycorex'ia*, (F.) *Faim de Loup*. Authors have described, under this name, a kind of boulimia, or depravation of the digestive functions, in which the patient eats voraciously, and passes his food, almost immediately afterwards, *per anum*.

FAMIGERATISSIMUM EMPLASTRUM, (*fama*, 'fame,' and *gero*, 'I wear.') A plaster extolled in ague, and made of aromatic, irritating substances. It was applied to the wrist.

FAMILIARICA SELLA, (*familia*, 'a household,' and *sella*, 'a seat or stool.') Close stool.

FAMILY DISEASES, see Hereditary.

FANCULUM, Anethum.

FANCY MARK, Nævus.

FANFECKLES, Ephelides.

FANG, (from Sax. *fangan*, 'to seize.') Radix.

FANON, (F.), (G.) *Fahne*, 'a banner, ensign, standard,' (L.) *pannus*, 'a cloth,' *Ferula*, *Leetulus stramineus*, *Thorulus stramineus*. A splint of a particular shape, employed in fractures of the thigh and leg to keep the bones in contact.

The Fanons were divided into *true* and *false*. The *true fanon* consists of a cylinder of straw, strongly surrounded with a cord or riband, in the centre of which a stick is usually placed to insure its solidity. The *false* consists of a thick piece of linen, made flat like a compress, and folded at the

extremities. It was placed between the fractured limb and the true fanon. The *Drap-fanon* is a large piece of common cloth placed between the fractured limb, in which the fanon or lateral splints are rolled.

FANTICKLES, Ephelides.

FAN'TOM, *Phantom*, *Phantom's*, (*φάντασμα*, 'a spectre,') (F.) *Phantom*, *Fantôme*. This word has two acceptations. It means the spectres and images which the imagination presents to the sick, when asleep or awake; and, also, the figure called *Man'ikin*, *Man'ikin*, and *Man'akin*, (F.) *Mannequin*, on which surgeons practice the application of bandages, or the accoucheur the manual part of midwifery—*Phantom's obstetricium*.

Fan'tom or Phan'tom Tu'mour is one that forms, in the abdomen for example, simulating organic disease, and occasionally exceedingly deceptive.

FAR'ADIC, (F.) *Faradique*. Relating or appertaining to faradisation or electricity of induction.

FARADISA'TION, *Far'adism*, (after *Faraday*, the celebrated physicist.) A term proposed by Duchenne, of Boulogne, for electricity of induction, or the localised application, therapeutically, of induction currents:—the currents, in other words, which are instantaneously developed in metallic conductors under the influence of voltaic or magnetic currents, or even of the earth.

FARCIMEN, (*farceire*, 'to stuff.') See Equinia.

FARCIMINALIS MEMBRANA SEU TUNICA, (*farcimen*, 'a sausage,') Allantois.

FARCIN, ([Celt.] *farcilh*, from *fare*, 'a horse,') See Equinia.

FARCINOMA, Equinia.

FARCTURA, *Factura*, (*farceire*, *farctum*, or *factum*, 'to stuff.') The operation of introducing medicinal substances into the cavities of animals or of fruits, which have been previously emptied.

FARCTUS, ('stuffed,') Emphraxia.

FARCY GLANDERS, see Equinia.

FARD, ([G.] *farbe*, 'color,') Paint.

FARDEAU, ([Celt.] *fardeall*, 'a burden,') Mole.

FARFARA, Tussilago—f. Bechium, Tussilago.

FARIGOLE, Thymus.

FARINA, (*far*, 'corn,' or grain of any kind, of which it is made.) *Al'hiton*, *Crim'non*, *Al'eton*, *Aleur'on*, *Al'ema*, (F.) *Farine*, *Meal* or *Flour*. The powder, obtained by grinding the seeds of the gramineous, leguminous, and cucurbitaceous plants in particular. It is highly nutritious and much used, dietetically as well as medicinally. In the pharmacopœias of London, Edinburgh, and Dublin, *Farina* means wheat flour—*Farina Tritici*.

Leath's Alimentary Farina or *Homœopathic Farina'ceous Food*, is said to consist principally of wheat flour, slightly baked, and sweetened with sugar, together with potato flour and a very small quantity of Indian corn meal and tapioca.

FARINA AMYGDALARUM, see Amygdala.

FARINA, COMPOUND'ED, EAST'ER'S, is said to consist of wheat flour, sweetened with sugar.

FARINA, NUTRIT'IOUS, MAID'MAN'S, is said to consist of potato flour, artificially coloured of a pink or rosy hue, the colouring matter being probably rose pink.

FARINA TRIT'ICI, Wheaten flour; *F. Secalis*, Rye flour or meal; *F. Hor'dei*, Barley meal; *F. Avena'cea*, Oat meal, &c. See Amylum.

FARINÆ RESOLVENTES, (F.) *Farines Résolutives*. This name was formerly given to a mixture of the farina of four different plants; Lupine, *Lupinus albus*; Ervum *Ervilia*; Vicia *Jaba*, and Barley, *Hordeum distichum*. They were recommended to form cataplasms.

FARINA'CEOUS, *Farina'ceus*, *Farino'us*, (F.) *Farinacé*, *Farineux*, *Meal'y*. Having the appear-

ance or nature of farina. A term given to all articles of food which contain farina. The term *Farinacea* includes all those substances called *cerealium*, *legumina*, &c., which contain farina, and are employed as nutriment.

Hard's farina'ceous food is fine wheat flour, which has been subjected to some heating process. *Brad'en's farina'ceous food* is said to be wheat flour, baked.

In *Pathology*, the epithet *farina'ceous*, (F.) *farineux*, is applied to certain eruptions, in which the epidermis exfoliates in small particles similar to farina.

FARINACEOUS FOOD, BRADEN'S, see *Farinacea*—f. Food, *Hard's*, see *Farinaceous*—f. Food, homoeopathic, see *Farina*.

FARINACEOUS FOOD, PLUMBE'S, is said to consist principally of bean or pea flour, most probably the former, with a little *Tacea* arrowroot, some potato flour, and a very little *Maranta* arrowroot.

FARINACEOUS FOOD, PRINCE ARTHUR'S, for infants and invalids of all ages, is said to consist entirely of wheat flour, slightly baked.

FARINARIUM, *Alicia*.

FARINE, *Farina*.

FARINES RÉSOLUTIVES, *Farinæ* resolutives.

FARINEUX, *Farinaceous*.

FARINOSUS, *Farinaceous*.

FARMERY, *Hospital*.

FARNTICKLES, *Ephelides*.

FARRIER, (from *ferrarius*, relating to *ferrum*, 'iron,' 'a blacksmith.') *Hippiater*.

FARSIGHTEDNESS, *Presbytia*.

FART, (Sax. *fart*, Teut. *fahren*, 'to go;—*fahrt*, 'a voyage,' or by onomatopœia; [G.] *Furz*.)

Bdelus, *Bd'olus*, *Bdelyg'mia*, *Bdelyg'mus*, *Bom'lus*, *Por'de*, *Phy'sa*, *Phy'sē*, *Fla'tus*, *Crep'itus*, *C. vent'ris*, (F.) *Pet*. A sonorous or other discharge of wind from behind. A low word, but of respectable parentage.

FARTURA, *Fartura*.

FARTUS, (*farcire*, *fartum* or *fartum*, 'to stuff.') *Emphraxia*.

FAS'CIA, (a band, from *fas*, 'that which is binding.') *Liga'tio*, *Ligatu'ra*, *Alligatu'ra*, *Anades'mus*, *Vin'culum*, *Spar'ganon*, *Epides'mus*, *Vinctu'ra*. A bandage, *fil'let*, *rol'er*, *tig'ature*. The aponeurotic expansions of muscles, which bind parts together, are likewise termed *Fasciæ*. See *Tænia*.

FASCIA APONEUROTICA FEMORIS, *Fascia lata* aponeurosis—f. *Capitalis*, *Bonnet d'Hippocrate*, *Capelina*—f. *Capitis*, *Diadema*—f. *Cooperi*, *F. Transversalis*.

FASCIA, CRIE'RIFORM, *F. Cribriform'is*. The sieve-like portion of the *fascia lata*; so called from its being pierced by numerous openings for the passage of lymphatic vessels.

FASCIA DENTA'TA, *F. D. Tanti'ni*, *Corps godronné*, of *Vicq d'Azyr*. A band of gray matter seen beneath the *tænia hippocampi* on raising it up, which runs along the inner border of the *cornu ammonis*. It is, as it were, crenated by transverse furrows.

FASCIA DIGITALIS, *Gantelet*—f. *Diopthalmica*, *Binculus*—f. *Dividens*, *Dividing bandage*—f. *Heliadori*, *T. bandage*.

FASCIA ILI'ACA, *Il'iac aponeuro'sis*. An aponeurosis which proceeds from the tendon of the *psaos minor*, or which arises from the anterior surface of the *psaos magnus*, when the former muscle does not exist. It is attached, externally, to the inner edge of the crest of the *ilium*, below and anteriorly,—on one side to the crural arch, sending an expansion to the *fascia transversalis*; and on the other, continuous with the deep-seated lamina of the *fascia lata*, which forms the posterior paries of the crural canal. Within and behind,

the *fascia iliaca* is attached to the brim of the pelvis, and is continuous with the aponeurosis, which *M. Jules Cloquet* has called *Pel'vian*. The *iliac aponeurosis* covers the *iliac* and *psaos* muscles, which it separates from the peritoneum.

FASCIA INFUNDIB'ULIFORMIS, is a funnel-shaped offset from the *fascia transversalis*, which is prolonged as a sheath upon the *vas deferens* and spermatic vessels as they pass out of the abdomen.

FASCIA INGUINALIS, *Spica*.

FASCIA LA'TA. A name given by anatomists to an aponeurosis, and to a muscle.

FASCIA LATA APONEURO'SIS, *F. aponeurotica fem'oris*, *Vagi'na fem'oris*, *Crur'al* or *Fem'oral Aponeuro'sis*, is the most extensive in the body, and envelopes all the muscles of the thigh. Above, it is attached to the outer edge of the *ilium*; before, it arises from the crural arch by two distinct laminae, separated by the femoral vessels, and becoming confounded a little below the part where the great *vena saphæna* opens into the crural vein. Of these two laminae, the one is more anterior and thicker than the other, and may be considered as a prolongation of the aponeurosis of the external oblique: it is intimately united to *Poupart's ligament*. The other, which is thinner, is behind, and deeper seated, and, after its union with the former, proceeds to be inserted into the pubis. Inferiorly, the *fascia lata* becomes confounded with the tendon of the *triceps*, and is attached to the external tuberosity of the tibia. The use of the *fascia lata*, like that of other aponeuroses, is to strengthen the action of the muscles, &c.

FASCIA LA'TA MUSCLE, *Ten'sor vagi'næ fem'oris*, *Fascia'lis*, *Membrano'sus*, *Mus'culus aponeuro'sis seu fas'ciæ la'tæ seu fem'oris membrano'sus*, (F.) *Il'io-aponeuro'si-fem'oral*, *Il'io-aponeuro'ti-fem'oral*—(Ch.) *Tenseur de l'aponeurose fémorale*. A muscle, situate at the upper and outer part of the thigh. It arises, above, from the outer part of the anterior and superior spine of the *ilium*; and is inserted, below, between the two laminae of the *fascia lata*, which it stretches and raises when it contracts.

FASCIA, OBTURATOR, see *Pelvic aponeuroses*—f. *Pelvic*, *Internal*, see *Pelvic aponeuroses*—f. *Pelvic*, *Lateral*, see *Pelvic aponeuroses*—f. *Pelvic*, *Superior*, see *Pelvic aponeuroses*.

FASCIA PRO'PRIA. A layer of areolar tissue derived from the sheath of the femoral vessels—or, according to some, from the *cribriform fascia*. It is one of the coverings of femoral hernia, and is generally pretty dense about the neck of the hernia; but thin or even wanting on its fundus.

FASCIA REPENS, *Spica*—f. *Scapularis*, *Scapulary*—f. *Sculteti*, *Bandage of separate strips*—f. *Semicircularis*, *Tænia semicircularis*—f. *Spiralis*, *Ascia*—f. *Stellata*, *Stella*—f. *Submuscular*, see *Vaginal* (of the eye).

FASCIA, SUBPERITONE'AL, *Subperitone'al aponeuro'sis*. A thin tendinous layer on the outer surface of the peritoneum.

FASCIA SUPERFICIA'LIS, *Superf'cial aponeuro'sis of the abdo'men and thigh*, (F.) *Aponeurose superficielle de l'abdomen et de la cuisse*. A very thin aponeurosis, which covers the muscles and aponeuroses of the abdomen; passes before the crural arch, to which it adheres with some degree of force; sends a membranous sheath, which surrounds the spermatic cord, and is continuous with the *dartos*, to which it assists in forming. The *fascia superficialis* presents, beneath the crural arch, very distinct fibres, whose direction is parallel to the fold of the thigh. It is applied over the *fascia lata* aponeurosis, and is attached, internally, to the ascending ramus of the *ischium*, near the root of the *corpus cavernosum*. Before the descent of the testicle from the abdomen, the *fascia super-*

facialis is very manifestly continuous with the *Gubernaculum testis*.

FASCIA FORMIS, T. bandage—f. Tortilis, Tour-niquet.

FASCIA TRANSVERSA' LIS, *F. Cooperi*. An aponeurosis, which separates the transversalis muscle from the peritoneum in the inguinal region. It arises above the posterior edge of the crural arch, where it seems to be continuous with the aponeurosis of the greater oblique muscle. Above, it is lost in the areolar tissue at the internal surface of the transversalis abdominis. Within, it is continuous with the outer edge of the tendon of the rectus muscle and Gimbernat's ligament; below, it is continuous with the aponeurosis of the greater oblique, and receives an expansion from the *Fascia Iliaca*. Towards its middle and a little above the crural arch, the fascia transversalis has the wide orifice of a canal, which is occupied, in the female, by the round ligament of the uterus; and in man, furnishes an expansion, that serves as a sheath to the spermatic vessels.

FASCIÆ (pl. of *Fascia*), Also, Swathing clothes—f. Coli, see Colon—f. Ligamentosæ Coli, see Colon.

FASCIALIS, *Fascia lata* muscle, Sartorius—f. Longus, Sartorius—f. Sutorius, Sartorius.

FASCIARUM APPLICATIO, Deligation.

FASCIATIO, Deligation, see Bandage—f. Cucullata, *Couvrechef*.

FASCICLE, Fasciculus.

FASCICULATED PORTION OF THE PEDUNCLE, see Peduncles of the Brain.

FASCICULI (pl. of *Fasciculus*) CORTICALES, Ferrein, pyramids of—f. Graciles, see Corpora testiformia.

FASCICULI INNOMINATI. Two large bundles of fibres in the interior of the medulla oblongata, behind the corpora olivaria, and more or less apparent between those bodies and the corpora testiformia. They ascend, and become apparent in the fourth ventricle, under the name *Fasciculi seu Processus teretes*.

FASCICULI, MEDIAN POSTERIOR, OF THE MEDULLA OBLONGATA, Funiculi graciles—f. Muscularum, see Muscular fibre—f. Pyramidales, Ferrein, Pyramids of—f. Teretes, Processus teretes, see Fasciculi innominati—f. Teretes cordis, Columnæ carneæ.

FASCICULUS, *Phacellus*, *Phacelus*, *Fascicle*, (dim. of *fascia*, 'a band,' 'a small bundle.' *Dessmedion*. In *Anatomy*, it is employed in this sense, as 'a fasciculus of fibres.' (F.) *Fascicule*, *Faisceau* ou *Petit Faisceau*, *Trousseau*. In *Pharmacy*, it means *Manipulus*, *Cheuplethes*, 'a handful.' *Musa Brassavolus* says,—as much as can be held in two fingers.

FASCICULUS CUNEATUS, Reinforcement, fasciculus of—f. of Reinforcement, Reinforcement, F. of.

FASCICULUS UNCINATUS. A white bundle, seen on the lateral aspect of the cerebral hemispheres, passing across the bottom of the fissure of Sylvius, and connecting the anterior with the middle and posterior lobes. The fibres of this fasciculus expand at each extremity, and the superficial portions of them curve or hook sharply between the contiguous parts of the anterior and middle lobes, whence the epithet 'uncinatus.'

FASCINOSUS, (from *fascinum*), Membrosus.

FASCINUM, Dildo, Penis.

FASCIOLA, (dim. of *fascia*, 'a band,') *Bandlette*—f. Cinerea, Tuberculum cinereum—f. Hepatica, Distoma hepaticum—f. Humana, Distoma hepaticum—f. Lanceolata, Distoma hepaticum.

FASELUS, Phaseolus vulgaris.

FASÉOLE, Phaseolus.

FASTIDIUM, (abridged from *fatis tedium*.) *Fastidiousness*, *Squeamishness*, or the condition of a stomach that is really affected with nausea.

FASTIDIUM CIBI, Asitia, Disgust—f. Potius, Loathing of drink.

FASTIGIUM, 'the top,' (from *fastus*, 'elevation.') Acme.

FASTING, (from Sax. *fæstan*), *Limo'sis ex-pers protrac'ta*, *Inedia*, *Nesti'a*, *Jejunium*, (F.) *Jeûne*. Loss or want of appetite without any other apparent affection of the stomach; so that the system can sustain almost total abstinence for a long time without faintness. Some wonderful cases of this kind, *Anorexia mirabilis*, are on record. See Abstinence.

FAT, (G. fett, Sax. *fæt*), *Pingue'do*, *Pim'elê*, *Pi'ar*, *Li'pê*, *Li'pos*, *Ste'ar*, *A'deps*, *Se'cum*, *Se'b-um*, *Cor'pus adipo'sum*, *Azun'gia*, (F.) *Graisse*. A soft, white, animal substance; inodorous; insipid; oily; inflammable; easy to melt; spoiling in the air, and becoming rancid by union with oxygen; almost insoluble in alcohol; insoluble in water; soluble in fixed oils. Fat is formed of the immediate principles, *stearin*, *margarin*, and *olein* or *elain*, all of which are regarded as salts composed of stearic, margarin and oleic acids, and a common base, to which, from its sweetish taste, the name *Glyc'erine*, *Glyceri'na*, *Glyceri'num*, (F.) *Glycérine* has been given. To these are, almost always, joined an odorous and a colouring principle. GLYCERIN, *Glyceri'na*, was introduced into the edition of the Pharmacopœia of the U. S. (1851), in which it was directed to be prepared as follows:—*Lead plaster*, recently prepared and yet fluid, and *boiling water*, of each a gallon; mix; stir briskly for 15 minutes; allow it to cool, and pour off the liquid. Evaporate until it has the s. g. 1.15, and pass slowly through it a current of sulphohydric acid until a black precipitate is no longer thrown down. Filter and boil until the sulphohydric acid is driven off, and evaporate the liquid until it ceases to lose weight.

In the last edition of the Pharmacopœia (1863), it has been transferred to the primary list of the *Materia Medica*.

Glycerin is a colourless or straw-coloured syrupy fluid; s. g. 1.25. It is soluble in water and in alcohol, but not in ether. It is used in the form of lotion, composed of half an ounce to ten fluid-ounces of water, in cutaneous diseases, as psoriasis, pityriasis, lepra, and ichthyosis. It keeps the part moist, as it does not evaporate readily. A *Glycerin ointment* may be made of *Spermaceti* 3ss; *White wax* 3i; *Oil of almonds* f3i; *Glycerin* f3i. Melt the spermaceti and wax with the oil, and incorporate the glycerin in a mortar. It may be used wherever a mild ointment is demanded.

Fat is found in a number of animal tissues, and is very abundant in the neighbourhood of the kidneys and in the epiploon. It is generally fluid in the cetacea; soft, and of a strong smell in the carnivora; solid, and inodorous in the ruminating animal; white, and abundant in young animals; and yellowish in old. It generally forms about a twentieth part of the weight of the human body. The fat, considered physiologically, has, for its function, to protect the organs; maintain their temperature; and to serve for nutrition in case of need; as is observed in torpid animals.

FAT, Corpulent—f. Cells, Fatty vesicles.

FAT, MACKAW, see Cocos butyrea.

FATIGATIO, (*fatigare*, *fatigatum*, 'to fatigue.') Copos.

FATIGUE, Copos.

FATITY, (Sax. *fæt*; past participle of *fætan*, 'to nourish.') *Adipo'us*, *Adipa'tus*, *Pimel'icus*, *Pimelo'des*, *Liparo'des*, *Li'parus*, *Pici'rous*, *Ad'i-pous*, *Pin'guid*, *Pingued'inous*. Relating to fat. Resembling or containing fat. The cellular membrane has been called *fatty* or *ad'ipous*; from an opinion that, in its areolæ, the fat is deposited. The areolar membrane, however, merely lodges, be-

tween its lamellæ and filaments, the vesicles in which the fat is contained.

FATTY DEGENERATION OF THE LIVER, Adiposis hepatica.

FATTY LIG'AMENT, Ad'ipous lig'ament. This name has been given to a reflection of the synovial membrane of the knee joint, which passes from the ligamentum patellæ towards the cavity that separates the condyles of the femur.

FATTY LIVER, Adiposis hepatica.

FATTY MEM'BRANE, Ad'ipous mem'brane, Adipous tis'sue. The subcutaneous areolar tissue, or that containing the fatty or adipose vesicles.

FATTY VES'ICLES, Sac'culi adipo'si, Ad'ipous ves'icles, Fat cells. This name is given to small brown or membranous vesicles which enclose the fat, and are found situate in the areolæ of the areolar tissue. These vesicles vary much in size. Generally, they are round and globular; and, in certain subjects, receive vessels which are very apparent. They form so many small sacs without apertures, in the interior of which are filaments arranged like septa. In fatty subjects, the adipose vesicles are very perceptible, being attached to the areolar tissue and neighbouring parts by a vascular pedicle. Raspail affirms that there is the most striking analogy between the nature of the adipose granules and that of the amylaceous grains.

FATTY VES'SELS, Ad'ipous ves'sels. The vessels connected with the fat. Some anatomists have called *Adipous canals*, (F.) *Conduits adipeux*, the vessels to which they attribute the secretion of fat.

FATUISME, Fatuitas.

FATUITAS, Moro'sis, Stultit'ia, Stu'por men'tis, Ament'ia; (F.) *Fatuisme, Fatuité*, (from *fatuus*, 'foolish.') Mental imbecility. Idiotism. Dementia. One affected with fatuity is said to be *fatuuus*, (Sc.) *Dæcipit*.

FATUITAS ALPICOLARUM, Cretinism—f. Convallinus, Cretinism.

FATUOUS, see Fatuity.

FAUCES, Isthmus, Pharynx, Throat.

FAUCETTE VOICE, (dim. of (F.) faux, (L.) falsus, 'false.') See Voice.

FAUCIAL, (from fauces.) Pharyngeal.

FAUCITIS, (fauces, and itis.) Isthmitis.

FAUNORUM LUDIB'RIA, 'The sports of the Faunæ.' Some authors have called thus incubus; others, epilepsy.

FAUQUIER SPRINGS, see Virginia, Mineral Waters of.

FAUSSE, (from falsus,) False—f. Conception, Conception, false.

FAUSSE COUCHE (F.), ('false birth.') *Vanum parturium*. Some authors have used this term for the expulsion of different bodies constituting false conceptions; such as moles, hydatids, clots of blood, &c. Most accoucheurs use the term synonymously with abortion.

FAUSSE GROSSESSE, ('false pregnancy.') Pregnancy, false—*f. Membrane, Membrane, false—f. Pleurésie, Pleurisy, false—f. Pneumonic, Peripneumonia notha.*

FAUSSE POSITION (F.), False position. The French use this term, in vulgar language, to indicate any attitude in which torpor, tingling, and loss of power over the motion of a part, are produced by too strong contraction or painful compression.

FAUSSE RHABARBE, ('false rhubarb.') *Thaliatron—f. Route, False passage—f. Variole, Variella.*

FAUSTINI PASTIL'LI, 'Faustinus's Loz'enges.' These were once celebrated. They were composed of burnt paper, quicklime, oxide of arsenic, sandarach, lentils, &c.

FAUX, (from (L.) falsus,) False—f. du Cerveau, Falx cerebri—f. du Cervelet, Falx cerebelli—f.

Germe, Conception, false—f. Grande du péritoine, Falx, great, of the peritoneum—f. Petite du péritoine, Falx, lesser, of the peritoneum—f. de la Veine ombilicale, Falx, great, of the peritoneum.

FAVEUX, Favosus.

FAVIFORMIS, (favus, 'honeycomb,' and forma, 'shape.') Favosus.

FAVOSUS, favus, similar to a honeycomb. Faviform'is, Fa'vus, (F.) Favenc. An epithet given to a species of porrigo. *Ce'riou, K'e'riou, Favus*; it means also a state of ulceration resembling a honeycomb.

FA'VULUS; (dim. of favus, 'a honeycomb.') *Favuli* is used by Dr. S. G. Morton, of Philadelphia, for the honeycomb-like depressions in the lining membrane of the stomach—the *stomach-cells* of Messrs. Todd and Bowman.

FAVUS, ('a honeycomb.') Porrigo, Porrigo favosa, see Favosus—*f. Confertus, Porrigo scutulata—f. Dispersus, Porrigo lupinosa—f. Disséminé, Porrigo lupinosa—f. en Groupes, Porrigo scutulata.*

FAXWAX, see Nucha.

FEABERRY, Ribes uva crispa.

FEATHERFEW, ([?] 'fewerfew,') *Matricaria.*

FEBRIC'ITANS, Feb'riens, Empyros, (F.) Fébricitant, (febricitare, 'to have a fever—febris.') One attacked with fever.

FEBRICITATIO, Feverishness.

FEBRICOSUS, (from febris.) Feverish.

FEBRIC'ULA, Fe'veret, (F.) Fébricule; (dim. of febris.) A term employed to express a slight degree of fever. *Ephemera.*

FEBRICULOSITY, Feverishness.

FEBRIENS, (febris, 'to have febris, a fever.') *Febriticans, Feverish.*

FEBRIFICIENT, (febris, 'a fever,' and facere, 'to make.') See Feverish.

FEBRIF'EROUS, Feb'rifer, (febris, and fero, 'I carry.') Fever-bearing, as a *febriferous locality.*

FEBRIFIC, Febrificient, see Feverish.

FEB'RIFUGÉ, Lexipyr'et'icus, Lexipyr'etus, Pyret'icus, Alexipyr'et'icus, Antifebri'lis, Antipyret'ic, Febri'fugus, (febris, and fugare, 'to drive away.') A medicine which possesses the property of abating or driving away fever.

FEBRIFUGUM, Febrifuge—f. Lignum, Quassia.

FE'BRILE, Febri'lis, (from febris,) Pyret'icus. Relating to fever, as *febrile movement, febrile pulse, &c.*

FEBRIS, Fever—f. Ac mastica, Synocha—f. Acuta, Synocha—f. Acuta continua, Synocha—f. Adeno-meningea, Fever, adeno-meningeal—f. Adeno-nervosa, Plague—f. Adynamica, Typhus—f. Africana, Fever, African—f. Agrypnodes, see Agrypnodes and Agrypnos—f. Alba, Chlorosis—f. Algida, see Algidus—f. Amatoria, Chlorosis, Hectic fever—f. Americana, Fever, yellow—f. Amphemera, Quotidian—f. Amphimerina hectica, Hectic fever—f. Amphimerina latica, Latica (febris)—f. Ampullosa, Pemphigus—f. Anabatica, Continued fever—f. Anginosa, Angina—f. Angiotenica, Synocha—f. Annua, see Annual diseases—f. Anomala, Fever anomalous—f. Aphonica, Fever, aphonic—f. Aphthosa, Aphtha—f. Apoplectica, Fever, apoplectic—f. Ardens, Synocha—f. Arte promota, Fever, artificial—f. Arthritica, Gout—f. Asodes, Fever, bilious, see Asodes—f. Assidua, Continued fever—f. Asthenica, Fever, asthenic, Typhus—f. Asthmatica, Fever, asthmatic—f. Ataxo-adynamica, Fever, ataxo-adynamic—f. Azodes, see Asodes—f. Biliosa, Fever, bilious—f. Bullosa, Pemphigus—f. Cardialgia, Fever, cardialgic—f. Catarrhalis, Catarrh—f. Catarrhalis epidemica, Influenza—f. Caumatodes, Synocha—f. Causodes, Synocha—f. Cephalalgica, Fever, cephalalgic—f. Cephalica, Fever, cephalic—f. Cholepyretica, Fever, bilious—f. Choleric, Fever, bilious, Fever, choleric—f. Chronica, Fever, chronic—f.

Coalterna, see Coalternae febres—f. Colliquativa, Fever, colliquative—f. Comatodes, Fever, apoplectic—f. Communicans, see Subintrans F.—f. Confusa, see Confusæ febres—f. Continens, Typhus—f. Continens non putrida, Synocha—f. Continens putrida, Typhus—f. Continua inflammatoria, Empresma—f. Continua putrida, Synochus—f. Continua putrida icterodes Caroliniensis, Fever, yellow—f. Continua non putris, Synocha—f. Continua sanguinea, Synocha—f. Convulsiva, Fever, convulsive—f. Crymodes, see Crymodes, and Algid fever—f. Culicularis, Miliary fever—f. cum Delirio, Fever, delirious—f. Depuratoria, Fever, depuratory—f. Deurens, Synocha—f. Diaphoretica, Fever, diaphoretic—f. Diaria, Ephemera—f. Duodecimana, Fever, duodecimane—f. Dysenterica, Dysentery—f. Elodes, see Elodes—f. Elodes icterodes, Fever, yellow—f. Enterica, see Typhus—f. Enteromesenterica, Enteromesenteric—f. Epacnastica, see Epacnasticus—f. Ephemera, Ephemera, Quotidian Fever—f. Epidemica cum angina, Cynanche maligna—f. Epileptica, Fever, epileptic—f. Erotica, Fever, erotic—f. Erratica, Fever, anomalous, Fever, erratic, see Erratic and Planetes—f. Erronea, see Planetes—f. Erysipelacea, Erysipelas—f. Erysipelatosa, Erysipelas—f. Esserosa, Miliary fever—f. Exanthematica articularis, Dengue—f. a Fame, Fever, famine—f. Exquisita, Fever, regular—f. Famelica, see Famelica Febris—f. Flava, Fever, yellow—f. Flava Americanorum, Fever, yellow—f. Gangrænodes, Fever, gangrenous—f. Gastrica, Fever, bilious, Fever, gastric—f. Gastro-adyamica, Fever, gastro-adyamic—f. Hæmoptoica, Fever, hæmoptoic—f. Hebdomadana, Octana—f. Hæctica, Hæctic fever—f. Hæctica infantum, Tabes mesenterica—f. Hæctica maligna nervosa, Typhus mitior—f. Hemeresia, Quotidian—f. Hemitritea, Hemitritea—f. Hepatica, Fever, bilious—f. Hepatica inflammatoria, Hepatitis—f. Horrida, see Algidus—f. Horrida, see Algidus—f. Humoralis, Fever, humoral—f. Hungarica, Fever, Hungaric—f. Hydrocephalica, Hydrocephalus internus—f. Hydrophobica, Fever, hydrophobic—f. Hysteretica, see Postpositio—f. Hysterica, Fever, hysteric—f. Ictericæ, Fever, icteric—f. Iliaca inflammatoria, Enteritis—f. Infantum remittens, Fever, infantile remittent—f. Inflammatoria, Synocha—f. Intensio, Epitasis—f. Intermittens, Intermittent fever—f. Intermittens cephalica larvata, Cephalalgia periodica—f. Intestinalis ulcerosa, see Typhus—f. Intestinarum, Enteritis—f. Irregularis, Fever, anomalous—f. Lactea, Fever, milk—f. Larvata, Fever, masked—f. Lenta, Fever, infantile remittent, Hæctic fever, Synochus—f. Lenta nervosa, Typhus mitior, Fever, nervous—f. Lenticularis, Miliary fever—f. Lethargica, Fever, apoplectic—f. Lochialis, Fever, lochial—f. Lygmodes, Fever, singultuous—f. Lyngodes, Fever, singultuous—f. Maculosa, see Typhus—f. Maligna, Fever, malignant—f. Maligna biliosa America, Fever, yellow—f. Maligna cum Soporæ, Typhus—f. Maligna flava Indiæ occidentalis, Fever, yellow—f. Marasmodus, Hæctic fever, Marasmodia—f. Meningo-gastricus, Fever, gastric—f. Mesenterica, Fever, adeno-meningeal, Fever, mesenteric—f. Methemerina, Quotidian—f. Miliaris, Miliary fever—f. Minuta, Fever, synocephal—f. Morbillosa, Rubella—f. Mucosa, Fever, adeno-meningeal—f. Mucosa verminosa, Fever, infantile remittent—f. Nautica pestilentialis, Typhus gravior—f. Nephritica, Fever, nephritic—f. Nervosa, Fever, nervous—f. Nervosa enterica, see Typhus—f. Nervosa epidemica, Typhus—f. Nervosa exanthematica, Typhus—f. Nervosa gastrica, see Typhus—f. Nervosa mesenterica, see Typhus—f. Nervosa petechialis, Typhus—f. Neurodes, Fever, nervous—f. Nocturnus, see Nocturnal—f. Nonana, Fever, nonane—f. Nosocomiorum, Typhus gravior—f. Nycternus, see Noctur-

nal—f. Octana, Fever, octane—f. Oscitans, Oscitant fever—f. Paludosa, see Elodes—f. Pannonica, Fever, Hungaric—f. Pemphigodes, Pemphigus—f. Pemphingodes, Pemphigus—f. Pempta, Quintan—f. Periodica, Fever, periodic—f. Perniciosa, Fever, pernicious—f. Pestilens, Plague—f. Pestilens maligna, Typhus gravior—f. Pestilentialis, Fever, pestilential—f. Pestilentialis Europæ, Typhus gravior—f. Phthisica, Hæctic fever—f. Planetes, see Planetes—f. Pleuritica, Pleuritis—f. Pneumonica, Fever, pneumonic, Pneumonia—f. Podagrica, Gout—f. Polycholica, Fever, bilious—f. Puerperalis, Fever, puerperal—f. Puerperalis biliosa, Metrocholosis—f. Puerperarum, Puerperal fever—f. Puncticularis, Miliary fever, Typhus gravior—f. Purpurate rubra et alba miliaris, Miliary fever—f. Purulenta, Fever, purulent—f. Putrida, Typhus gravior—f. Putrida nervosa, Typhus mitior—f. Quartana, Quartan—f. Quæquæ, see Algidus—f. Quinta, Quintan—f. Quintana, Fever, quintan, Quintan—f. Quotidiana, Fever, quotidian, Quotidian—f. Recurrens, Relapsing fever—f. Regularis, Fever, regular—f. Remittens, Remittent fever—f. Remittens infantum, Fever, infantile remittent—f. Rheumatica inflammatoria, Rheumatism, acute—f. Rubra, Scarlatina—f. Rubra pruriginosa, Urticaria—f. Sanguinea, Synocha—f. Sapropyræ, Typhus gravior—f. Scarlatina, Scarlatina—f. Scorbatica, Fever, scorbutic—f. Semitertiana, Hemitritea—f. Septana, Fever, septan—f. Sesquialtera, Hemitritea—f. Sextana, Fever, sextan—f. Simplex, Ephemera, Fever, simple—f. Singultosa, Fever, singultuous—f. Soporosa, Fever, apoplectic—f. Stercoralis, Fever, stercoral—f. Sthenica, Synocha—f. Stomachica inflammatoria, Gastritis—f. Subintrans, Fever, subintrans—f. Sudatoria, Fever, diaphoretic, Hydropyretus, Sudor anglicus—f. Syncopalis, Fever, synocephal—f. Syphilitica, Fever, syphilitic—f. Tabida, Fever, colliquative, Hæctic fever—f. Tertiana, Fever, tertian, Tertian fever—f. Tonica, Synocha—f. Topica, Neuralgia, facial—f. Toxicæ, Fever, yellow—f. Tragica, Fever, tragia—f. Traumatica, Fever, traumatic—f. Tropica, Fever, yellow—f. Typhodes, Typhus—f. Urticata, Urticaria—f. Uterina, Metritis—f. Vaga, Fever, anomalous, see Planetes—f. Variolosa, Variola—f. Verminosa, Fever, infantile remittent, F. verminous, Helminthopyra—f. Vernalis, Fever, vernal—f. Vesicularis, Miliary Fever, Pemphigus—f. Virginum, Chlorosis.

FEB'RUA, (*februus*, 'I purge.') In ancient mythology, a goddess who presided over menstruation.

FECAL, Stercoraceous—f. Matter, see Fæces.

FÆCES, Fæces.

FÉCONDATION, Fecundation.

FÉCONDITÉ, Fecundity.

FEC'ULA, *Fæcula*, (dim. of *fæces*.) An immediate principle of vegetables, composed of hydrogen, oxygen, and carbon. It exists in several plants, and has different names, according to that which furnishes it. (F.) *Fécule*, *Amygdolé* When extracted from wheat or barley, it is called *starch*, *Amylum*. When from *Cyclus circinalis*, *Sa'go*;—from *Orchis mo'rio*, *Salep*. We say, also, *Fecula of the Potato*, *Bryonia*, *Arum*, *Manioc*, &c.

FECULA AMYLACEA, *Amylum*.

FECULA GREEN. This name is given to a green, solid matter, of variable character, which is believed to be resinous, and which renders turbid several kinds of juices extracted from vegetables. It is also called *Chlorophyll*.

FECULA MARANTÆ, Arrow-root—f. *Tapicka*, see *Jatropha manihot*.

FÉCULE, Fecula.

FEC'ULENCE, *Fæculentia*, *Fæculositas*, *Fæx*, plural *Fæces*, *Lem'ma*, *Lee*, *Depos't*, *Dregs*. In

Pharmacy, feculent, albuminous, or other substances, which are deposited from turbid fluids.

FEC'ULENT, *Fæculent'us*, *Hypot'rygus*, *Trygo'des*. 'Foul, dreggy, excrementitious;' as a *feculent fluid*, *feculent evacuations*, &c.

FEC'UND, *Fæcund'us*, *Fæcun'dus*, (F.) *Fécond*. Fruitful, prolific.

FECUNDA'TION, *Fæcunda'tio*, *Impregna'tio*, *Impregna'tio*, *Ingravida'tio*, *Pregna'tio*, *Gravida'tio*, *Pregna'tus*, *Fructifica'tio*, *Cy'e'sis*, *Ency'e'sis*, *Eucymo'sia*, *Procrea'tion*, (F.) *Fécondation*. The act by which, in organized beings, the material furnished by the generative organs of the female, unites with that prepared by those of the male, so that a new being results.

FECUN'DITY, *Eutoc'ia*, *Polygon'ia*, *Polytoc'ia*, *Fæcun'ditas*, *Productiv'itas*, (F.) *Fécondité*. The faculty of reproduction, possessed by organized bodies.

It has been estimated that, throughout a country, taking one marriage with another, not more

than 4 children are the result; and in towns only 35 children in 10 marriages.

FEE, *Pysechorages*, *Sostrum*.

FEET, BURN'ING OF THE. A singular cachectic disease, described by Mr. Malcolmson as occurring in India, the prominent symptom of which was a sense of burning in the feet.

FÉGARITE, *Cancer aquaticus*, *Stomatitis*, pseudo-membranous.

FEIGNED DISEASES, *Mor'bi dissimula'ti seu simula'ti seu cela'ti seu infita'ti seu pseudalei*, *Sim'ulated or Pretend'ed diseases*, (F.) *Maladies dissimulées ou simulées ou feintes ou supposées*. The tricks employed, by impostors, to induce a belief that they are attacked with diseases when they are not. These are generally assumed by beggars to obtain alms; by criminals to escape punishment; and by soldiers to be exempt from duty.

The following table exhibits the chief feigned, pretended, simulated or excited diseases, or disqualifications, with the means of detection:—

DISEASES, &c.	HOW FEIGNED.	HOW DETECTED.
ABORTION.	{ By staining the clothes and body with borrowed blood.	{ By constant and minute attention.
ABSTINENCE.		{ Amaurosis is characterized by dilated pupil. Where these substances have been applied, the effects will go off in ten days or a fortnight.
AMAUROTIC BLINDNESS.	{ By applying the extract of belladonna or datura stramonium to the eye.	{ By anæsthetics, and, under their use, or otherwise, taking the person off his guard.
APHONIA.		{ By powerful stimulants; an electric shock; application of hot water, sternutatories, actual cautery, &c.
APOPLEXY.	{ By falling down as if deprived of sensation and consciousness.	{ By examining if the pulse be strong, and the skin hot, and whether there be loss of appetite or of strength, or swelling of the limbs, or unmistakable feebleness of the vital actions.
CACHEXIA, ANEMIA, AND DEBILITY.	{ Using substances to make the face appear pale and livid. Indulging freely in wine, and privation of sleep prior to examination.	{ By the aid of chymistry. We are acquainted with the chymical composition of urinary calculi.
CALCULI, EXCRETION OF.	{ Putting sand, pebbles, &c. into the urine.	{ By noticing whether there be signs of cachexia, and by attentive examination of the part.
CANCEROUS ULCER.	{ By gluing on a portion of a spleen with the smooth side to the skin, leaving on the outside the appearance of an ulcerated surface.	{ By powerful stimulants, as recommended under apoplexy. Letting fall a drop of boiling water on the back. Proposing to use the actual cautery, and seeing whether the pulse rises.
CATALEPSY	{ By seeming to be suddenly motionless, the joints remaining flexible, and external objects making no impression.	{ By examining the patient whilst he may imagine himself unobserved, and seeing whether the convulsive motions go on. By anæsthetics, as under Aphonia.
CHOREA.	{ By assuming the convulsive motions of a part which characterize chorea.	{ Mode of discrimination sometimes so obscure as to deceive the most practised and attentive. By anæsthetics, as under Aphonia.
CONTRACTION OF JOINTS IN GENERAL.		{ Introduce a cord between the fingers and the palm of the hand, and gradually apply weights so as to expand the fingers. Confine him so that he cannot obtain his food without using his clenched hand. By anæsthetics, as under Aphonia.
CONTRACTION OF THE FINGERS.		{ When feigned, they do not present the rigidity of muscles or the rapidity of action which characterize the real. The mode of detection must be the same as in epilepsy.
CONVULSIONS.		{ The existence of the opacity can be detected by attentive observation.
CORNEA, OPAKE.	{ Produced by the application of a strong acid, by acrid powders, as quicklime, &c.	{ By careful watching and examination on the part of practitioner and nurse.
CUTANEOUS DISEASES.	{ Some articles of diet will bring on <i>urticaria</i> or <i>nettle-rash</i> , in particular individuals, as shellfish, bitter almonds, &c. By acrids, acids, or any irritants applied to the surface. An ointment of tartarized antimony causes a painful pustular eruption. See Porriego, in this list.	{ The really deaf and dumb acquire an expression of countenance and gestures which it is difficult to assume. By anæsthetics, as under Aphonia.
DEAFDUMBNESS		

DISEASES, &c.	HOW FEIGNED.	HOW DETECTED.
DEAFNESS.	It may be assumed or excited by putting a pea in the ear, or by inserting irritants, so as to induce inflammation and temporary loss of function.	Make a noise when not expected, and see if the countenance varies or pulse rises. Put to sleep by opium, and then fire a pistol close to the ear, when he may be thrown off his guard. Examine the ear to see if any trick has been played there. By anæsthetics, as under Aphonia.
DEATH.	Some persons possess the power of suspending or moderating the action of the heart.	If suspected, the plan recommended under apoplexy will be found most efficacious.
DEBILITY, see CACHEXIA.		
DEFORMITY.		
DELIRIUM.		Examine the part and its articulation naked, and compare it with the opposite. Real delirium is accompanied by bodily disease; with general wandering and incoherency. The feigned is more confined to special subjects.
DELIVERY.	After enlargement produced artificially, a subsidence of the tumefaction; the parts being moistened by borrowed blood, and the child of another substituted as the female's own.	Can only be positively detected by examination <i>per vaginam</i> . Soon after delivery, the vagina will be relaxed and the lochial discharge be flowing in greater abundance, the shorter the time that may have elapsed since delivery.
DIARRHŒA.	Said to have been caused by a mixture of vinegar and burnt cork. May be occasioned by the use of any of the purgative roots, &c.	When diarrhœa is feigned by the lower classes, inspect the linen; if clean, the bowels are probably not much out of order. Let every individual have a close stool of his own; and inspect the evacuations, taking care that one suffering under the disease does not lend his evacuations to another.
DROPSY.	May be feigned, like pregnancy, by wearing pads. The anasarcons condition of the lower limbs has been caused by applying a ligature round them. By inflating the cellular membrane of the abdomen.	Can be detected by attentive examination. There will be a want of that leucophlegmatic habit which accompanies and characterizes dropsy.
DYSENTERY.	May be feigned, like diarrhœa, by adding a little blood to the evacuations, or by introducing a soap, or some more irritating, suppository.	Same rules as under diarrhœa.
EPILEPSY.	The foaming of the mouth has been produced by keeping a piece of soap in it.	Sensation in epilepsy is totally abolished. If any remain, disease probably feigned. Incontractility of pupil, which occurs in epilepsy, cannot be feigned. Same means to be used as in feigned apoplexy.
FEVER.	By various stimulants, as wine, brandy, pepper; swallowing a small quantity of tobacco, or introducing it into the anus. Flour or chalk used to whiten the tongue. Redness of skin, caused by friction with a hard brush.	This deceit is generally developed by an examination of a day or two. Where flour or chalk has been used to whiten the tongue, the line of demarcation between the whitened part and the clean, healthy margin of the tongue, is too well marked to escape observation.
FISTULA IN ANO.	By making an incision near the verge of the anus, and introducing into it an acrid tent, such as the <i>root of white heliobore</i> .	By careful examination.
FRACTURES.	There is generally nothing but the man's own testimony. He complains of pain in the part; if fracture of the skull be feigned, he states, perhaps, that he becomes deranged on tasting liquor.	By attentive examination.
HÆMATEMESIS.	By drinking the blood of some animal, or using some coloured liquid, and then throwing it up.	By cutting off the supply of the fluid and careful examination.
HÆMOPTYSIS.	By secreting bullock's blood for the purpose of colouring the saliva; making small incisions in the mouth; using bole armeniac or paint of vermillion.	Blood from the lungs is frothy and light-coloured. Mouth and fauces must be carefully inspected, and the individual be observed.
HÆMORRHOIDS.	By introducing bladders of rats or of small fish partly into the rectum. The linen has also been stained with borrowed blood.	The means are obvious.
HEART, DISEASES OF, see PALPITATION.		
HEPATITIS.		Unless the person be a well-educated man, acute inflammation of the liver will be detected by the absence of marks of strong inflammatory action. <i>Chronic liver disease</i> is, frequently, not characterized by wellmarked symptoms, and hence, when assumed, is difficult of detection.
HERNIA.	In the same manner as hydrocele;—by inflation.	The detection is easy.
HYDROCELE.	By puncturing the skin of the scrotum, and inflating the cellular membrane.	Do.
HYDROCEPHALUS.	By opening the integuments of the head near the vertex, e.g. and blowing in air.	Do.
HYSTERIA.		Does not easily resist the application of strong sternutatories to the nostrils. Attentive examination necessary.

DISEASES, &c.	HOW FEIGNED.	HOW DETECTED.
INSANITY.		The expression of countenance cannot easily be feigned. Nor can the affection be kept up so long as in real mental alienation. The individual cannot do so long without food, sleep, &c.
JAUNDICE.	By colouring the skin with an infusion of turmeric or tincture of rhubarb. Clay-coloured stools produced by taking daily a small quantity of muriatic acid. High-coloured urine by rhubarb taken internally.	The eyes cannot be coloured, although smoke has been used for this purpose. The skin must be washed to remove the colouring matter if any exist, and the supply of acid and rhubarb be prevented.
LAMENESS.	By keeping the limb in a contracted state, and resisting any efforts to move it.	By two persons taking hold of the individual and moving rapidly along with him; and when they are tired having relays. The impostor will generally give in. By anæsthetics, as under Aphonia.
MENSTRUATION.	By staining the clothes and body with borrowed blood.	By cutting off the supply.
MYOPIA.		Present an open book, and apply the leaves close to the nose. If it cannot be read distinctly, when thus placed, or when glasses proper for short-sightedness are used, the disease is feigned.
OPHTHALMIA.	Excited by a variety of acrid and corrosive substances applied to the eye; as lime, &c. A portion of black muslin, spread over the cornea. The eyelashes are sometimes extracted, and caustic applied to excite disease in the palpebræ.	When ophthalmia is thus excited, its progress is ordinarily very rapid, arriving at its height within a few hours.
OZENA.	By impregnating a piece of sponge with some offensive juices or oils, mixed with decayed cheese, and putting the imbued sponge into the nostrils.	
PAINS.		The detection is here often difficult. The non-existence of pain cannot be proved, and great pain may be present without any appearance externally.
PALPITATION.	White hellebore, given in the dose of 10 or 12 grains, and repeated, will occasion general indisposition, and undue action of the heart.	The imposition is more frequently detected by inconsistencies and contradictions in the patient's history of the case, than in any other manner.
SHAKING PALSY.		Cut off the supply.
PARALYSIS.		May be suspected, if the person be in an ordinary state of vigour. Use violent remedies and means, recommended under Chorea.
PHTHISIS PULMONALIS.	Individuals with long necks and contracted shoulders have simulated phthisis, by covering the chest with blisters, cicatrices of issues, &c., and by taking drugs which cause paleness.	Violent remedies are here required. Cold affusion, actual cautery, electric shocks, &c. By anæsthetics, as under Aphonia.
POLYPUS NASI.	By introducing the testicle of a young cock, or the kidney of a rabbit, into the nostril, and retaining it there by means of a sponge fastened to it.	By attentive examination of the symptoms.
PORRIGO OR SCALDED HEAD.	By applying nitric acid to the head, after protecting the face with fatty substances; but the chronic state is imitated by the use of depilatories of different kinds applied sometimes in patches, so as to resemble <i>Porrigo decalvans</i> .	
PREGNANCY.	By wearing pads, and assuming the long-ing after particular articles of diet, &c.	By the absence of the areola; the presence of a pad; and, if necessary, by examination <i>per vaginam</i> .
PROLAPSUS ANI.	By a portion of the intestine of the ox, in which a sponge filled with a mixture of blood and milk is placed.	
PROLAPSUS UTERI.	By a similar fraud.	
PULSE, WEAKNESS OR DEFECT OF.	By ligatures applied to the corresponding arm. Slowness of pulse may be caused by the use of <i>Veratrum viride</i> .	By examining whether the arteries of the two arms beat alike; and if a ligature be placed on the arm.
RHEUMATISM, CHRONIC.	By complaining of great suffering on moving the limbs, and of pain and stiffness of the joints.	By careful observation, and the use of anæsthetics, as under Aphonia. It is frequently feigned, and not easy of detection. See Pains.
SCROFULA.	By exciting ulcers below the angles of the jaw.	By examining the general habit, and observing whether the ulcerations be glandular, and the discharge of a scrofulous character.

DISEASES, &c.	HOW FEIGNED.	HOW DETECTED.
SCURVY.	By covering the teeth with wax, and then applying acid corrosive substances to the gums.	By examining the general habit; whether debilitated, cachectic, and possessing the usual general signs of scorbutus. Simulators of this defect generally state, that it is connate, or ascribe it to a fit of apoplexy or severe fever. Where the organs of speech were perfect, and the moral evidence of the previous existence of the infirmity was not satisfactory, the French authorities used to confine the soldiers, and not supply them with food, until they called for it without stammering.
STAMMERING.		If the person be roused suddenly from sleep, the eyes will be right, as they also are during sleep. Under anæsthetics they would be so likewise.
STRABISMUS.		By passing a bougie.
STRICTURE OF THE URETHRA.		Examine the limbs uncovered.
SWELLING OF THE LEGS.	By ligatures round the thighs.	By using sternutatories. By the absence of some of the symptoms of syncope. Examine the naked arms. Wash the face.
SYNCOPE.	Ligatures are sometimes used to prevent the pulse being felt. By applying lotions to the face to make it pale.	
TUBERCULOSIS, see Phthisis.		
TYMPANY.	Persons have possessed the power of swallowing air, so as to distend the stomach, and simulate tympany.	
ULCER OF THE EAR.	By introducing a tent, imbued with blistering plaster into the ear, and repeating the application, until the tube becomes ulcerated, and a discharge of puriform matter is established. The fetid smell is imitated, by dropping into the ear a mixture of empyreumatic oil, asafoetida, and old cheese. Also, by introducing a little honey into the meatus.	By careful examination.
ULCERS OF THE LEGS, &c.	By corrosives, or irritants. Sometimes by abrasion, by rubbing sand on the shin-bone. At others, they are pretended, by gluing on a piece of spleen or the skin of a frog.	Artificial ulcers have, usually, a more distinct margin, and are more readily healed than others; the latter being generally indicative of an impaired constitution.
URINE, BLOODY.	The fruit of the Indian Fig (<i>Cactus opuntia</i>) colours the urine as red as blood. Cantharides will cause it. Blood may also be procured and mixed with the urine.	By making the patient pass his urine in the presence of the physician, and examining the vessel before and after. By cutting off the supply of any substance which could cause the appearance.
URINE, INCONTINENCE OF.		Difficult, at times, of detection. Give the person a full dose of opium, and introduce the catheter when he is asleep. If there be urine, the incontinence is feigned.
VARICOSE VEINS.	By a ligature, placed tightly round the limb. They may be excited in this manner, or aggravated if already existing.	By examining the limb.

FEL, gen. *Fellis*, Bile—f. Anguilla, see Bile—f. Bovinum, Bile of the Ox—f. Bovis, see Bile—f. Naturæ, Aloes—f. Tauri, see Bile—f. Terræ, Ludus Helmontii—f. Ursi, see Bile—f. Vitri, see Vitrum.

FELDSCHEREN, ('army barbers,') see Bathers.

FELINEUS, (from *fel*,) Bilious.

FELLEUS, Biliary, Bilious.

FELLIDUCUS, (*fel*, and *ducere*, 'to lead,') Cholagogue.

FELLIS OBSTRUCTIO, ('obstruction of bile,') Icterus—f. Suffusio, Icterus—f. Superfusio, Icterus.

FELON, 'a great criminal,' (from Sax. *Fællan*, 'to offend,') Paronychia.

FELTING, (Sax. *felt*, 'cloth or stuff made without weaving,') *Tangling*. A term applied to the hair when inextricably interlaced, as occurs occasionally in women from inattention.

FÊLURE DE GLASER, 'fissure of Glaser,' (Old [F.] *feslure*; from [L.] *findere*, *fissum*, 'to split' [?]), Fissure, glenoid.

FELWORT, *Gentiana lutea*.

FEMALE, *Fœmina*, *Fœmina*, *Femell'a*, *Gynē*, (F.) *Femelle*, (dim. of *femme*, 'woman,') In animals, the one that engenders and bears the young. It is, also, used adjectively:—as, the *Female Sex*, &c.

FEMALE PARTS, Vulva.

FEMELLA, Female.

FEMELLE, (from *femme*,) Female.

FEMEN, Thigh (inner part), Vulva.

FEMINA, Female.

FEMINAL, Pubes.

FEMINES'CENCE, *Fæminescen'tia*, (*fæmina*, 'a female,') The possession or assumption of certain male characteristics by the female.

FEMME, Woman—f. *en couche*, Puerpera.

FEMORAL, *Femoral'is*, (*femur*, gen. *femoris*, 'the thigh,') Belonging or relating to the thigh; as *Femoral artery*, *Femoral hernia*, &c. See Crural.

FEMORALIS, Femoral, *Triceps cruris*.

FÉMORO-CALCANIEN PETIT, Plantar muscle—f. Popliteal, great, Sciatic nerve, great—f. *Popliti-tibial*, *Popliteus* muscle.

FÉMOROTIB'IAL, *Femorotibial'is*. Belonging to the femur and tibia. The *Femorotibial articulation* is the knee-joint.

FEMUR, *Mc'rus*, *Mc'rium*. The thigh. Also, the *thighbone*, *Os fem'oris*, (F.) *L'os de la Cuisse*. The strongest and longest of all the bones of the body, extending from the pelvis to the tibia, and forming the solid part of the thigh. The femur is cylindrical, slightly curved anteriorly, unsymmetrical and oblique downwards and inwards.

The body of the bone is prismatic, and has, behind, a prominent ridge, the *linea aspera*.

PROCESSES.	{	<i>Upper or</i>	{	Head supported on a
		<i>Pelvic</i>		column or neck.
		<i>Extremity.</i>		Great trochanter.
				Lesser trochanter.
		<i>Inferior or</i>		External condyle.
		<i>Tibial</i>		Internal condyle.
		<i>Extremity.</i>		External tuberosity.
				Internal tuberosity.

The femur ossifies from five points:—one on each process of the pelvic extremity; one in the body of the bone; and two on the condyles. It is articulated with the pelvis, tibia, and patella. See Thigh.

FÉMUR, COL DU, Collum femoris—f. Moventium septimus, Iliacus internus—f. Moventium sextus, Psoas magnus—f. Summum. Vulva.

FENES'TRA, 'a window' (akin to *φανω*, 'I show,') (F.) *Fenêtre*. Anatomists have given this name to two apertures, situate in the inner paries of the cavity of the tympanum. The one of these is the *Fenes'tra ova'lis seu vestibula'ris, Fora'men ova'le*. It is oval-shaped; situate at the base of the stapes, and corresponds with the cavity of the vestibule. The other is the *Fenes'tra rotun'da seu cochlea'ris seu triquetra, Fora'men rotun'dum*. It is closed by a fine, transparent membrane, called *Membra'na Tym'pani secundaria, Tym'panum mi'nus seu secundarium, Membra'na Fenes'træ rotun'dæ*, and corresponds to the inner scala of the cochlea.

FENESTRA OCULI, Pupil—f. Ovalis, see Fenestra
—f. Rotunda, see Fenestra.

FENES'TRAL, *Fenestra'tus*, (from *fenestra*.) (F.) *Fenêtré* et *Fenêtré*. Bandages, compresses, or plasters with small perforations or openings, are so called. The openings prevent the detention of the discharge.

FENESTRATUS, Fenestral.

FENESTRÉ, Fenestral.

FENÊTRE. Fenestra.

FENÊTRÉ, Fenestral.

FENNEL, (Sax. fenöl.) *Anethum* — f. Dogs',
Anthemis cotula — f. Flower, *Nigella* — f. Hog's,
Peucedanum — f. Sweet, *Anethum* — f. Water,
Phellandrium aquaticum.

FENOUIL, Anethum—*f. d'Eau*, Phellandrium aquaticum—*f. Marin*, Crithmum maritimum—*f. de Pore*, Peucedanum—*f. Puant*, Anethum graveolens.

FENTE, (from *fendre*, [L.] *findere*, 'to split,')
Fissure—*f. Capillaire*, see *Pilatio*—*f. Glénoïdale*,
Fissure, glenoid—*f. Orbitaire*, Orbital fissure—*f.*
Orbitaire inférieure, Spheno-maxillary fissure—*f.*
Sphénoïdale, Sphenoidal fissure—*f. Sphéno-maxil-*
laire, Spheno-maxillary fissure.

FÉNUGREC, (*fœnum Græcum*, 'Greek hay,')
Trigonella fœnum.

FENUGREEK, *Trigonella foenum*.

FERRE, Ferrum — *f. Acétate* de, Ferri acetas — *f. Bromure* de, see Bromine — *f. Carbonate* de, Ferri protocarbonas — *f. Carbure* de, Graphites — *f. Chaud*, Pyrosis — *f. Hydrate* de, tritoxide de, Aëtes — *f. Hydratide* de, see Ferri iodidum — *f. Iodure* de, Ferri iodidum, see Iodine — *f. Lactate* de, Ferri Lactas — *f. Limaille* de, Ferri limatura — *f. Perchlorure* de, Ferri chloridum — *f. Peroxide* de, Ferri subcarbonas — *f. Permanganate* de, see Ferri Sulphas — *f. et de Potasse, tartrate* de, Ferrum tartarizatum — *f. et de Potassium, cyanure* de, Potassii Ferrocyanuretum — *f. et de Potassium, protocyanure* de, Potassii Ferrocyanuretum — *f. Réduit par l'hydrogène*, Ferri pulvis — *f. Rouge*, Pyrosis — *f. Sulfate* de, Ferri Sulphas — *f. Sulfure* de, Ferri Sulfuretum — *f. Tannate* de, Ferri tannas — *f. Trichlorure* de, Ferri chloridum — *f. Valérienate* de, Ferri Valerianas.

FERALIS PEDICULUS, see Pediculus.

FE'RI'NE, *Fer'inus*; ('savage, brutal;' from *fera*, 'a wild beast.') *Therio'dea*. A term, applied to any malignant or noxious disease. In France it is used only when joined to *Toux*; *Toux férine*, a dry, obstinate, and painful cough.

FERMENT, *Fermentum*, (quasi *fermentum*, from *ferveo*, 'I boil, I am hot.') *Zy'ma*, *Zymo'ma*, *Lev'en*, (F.) *Lévain*. The Iatrochymists applied this name to imaginary substances, *Fermenta mor'bi*, to which they attributed the power of producing disease, by exciting a fermentation in the humours.

FERMENTA'TION, *Fermenta'tio, Zymo'sis, Æstua'tio, Caus'is, Bras'mos.* An intestinal movement, occurring spontaneously in a liquid; whence result certain substances, that did not previously exist in it.

The chemical physicians attributed all diseases to an imaginary fermentation of the humours.

FERMENTATION, PUTREFACTIVE, Putrefaction.

FERMENTOLEUM SOLANI, Oil, fusel.

FERMENTUM CEREVISIÆ, Yest—f. Morbi,
Ferment—f. Ventriculi, Gastric juice.

FERN BUSH, (Feapn.) *Comptonia asplenifolia*—f. *Cinnamon*, *Osmunda cinnamomea*—f. *Eagle*, *Pteris aquilina*—f. *Female*, *Asplenium filix foemina*, *Pteris aquilina*—f. *Flowering*, *Osmunda regalis*—f. *Gale*, *Comptonia asplenifolia*—f. *Male*, *Polypodium filix mas*—f. *Meadow*, *Comptonia asplenifolia*—f. *Rock*, *Adiantum pedatum*.

FERN ROOT, *Polypodium vulgare*.

FERN, SWEET, *Adiantum pedatum*, *Comptonia asplenifolia*—f. Sweet, shrubby, *Comptonia asplenifolia*.

FERNAMBUCO WOOD, *Ca salpinia echinata*.
FERNFEBCKLED, *Ephelides*.

FERNITICKLES, Ephelides.

FERONIA ELEPHANTUM, (from the goddess *Feronia*.) A tree of India, Order Aurantiaceæ, which yields a gum closely resembling Gum Arabic. Dr. Pereira thinks it not improbable, that a part of the *East India gum* taken to England may be the produce of this tree. The leaves are aromatic, of an odour resembling anise, and are used as a carminative.

FERONIA PELLUCIDA, Covolam.

FERRAMENTUM, *Side'ion, Instrumen'tum fer'reum*, (from *ferrum*.) Any surgical instrument made of iron. By the vulgar, in France, the word *ferrements* means the instruments used in difficult labours.

FERRARIA, *Scrophularia aquatica*.

FERRATED, Chalybeate.

FERRATUS, (from *ferrum*,) Chalybeate.

FERREIN', CANAL' OF, *Ductus Ferrei'ni*. A triangular channel, which Ferrein supposed to result from the approximation of the free edges of the eyelids applied to the globe of the eye; and which he considered adapted for directing the tears towards the puncta lacrymalia, during sleep. The canal is, probably, imaginary.

The same name is likewise given to the *cortical canals*,—the first portions of the uriniferous ducts, whilst still in the cortical substance of the kidney.

FERREIN, PYRAMIDS OF, *Pyramides Ferrei*'ni. Each of the papillæ of the kidney, according to Ferrein, consists of, at least, 700 subordinate cones or pyramids. To these last the names *Pyramides of Ferrein*, *Pyramides renales Ferrei*'ni, *Fusculi pyramidal'es*, *Fus'ciculi cortical'es*, *Lobuli renum*, have been given, in contradistinction to the Pyramids of Malpighi.

FERRI ACETAS, *Extractum martis acetici-*
cum, Aceticum martiis, Acetate of Iron, Iron
Liquor, (F.) Acétate de fer. A preparation of the
Dublin Pharmacopœia, made by digesting 1 part
of carbonate of iron in 6 parts of acetic acid for
three days, and filtering. Dose, as a tonic and
astringent, ℥v to ℥xx, in water.

FERRI ÆRUGO, F. Subcarbonas.

FERRI ET ALUMINÆ SULPHAS, *Sulphate of Iron and Alumina*, is made by treating bicarbonated solution of soft iron and carbonated solution of pure washed alumina, with sulphuric acid. It has been recommended as a valuable astringent. Dose, from five to ten grains.

FERRI ET AMMONIÆ MURIAS, *Ferrum ammoniatum*.

FERRI ET AMMONIÆ SULPHAS, *Sulphate of Iron and Ammonia*, *Ammonio-fer'ric alum.* (Liq. ferri tersulphat. Oij; Ammon. sulph. ℥ivss. Add the sulphate of ammonia to the solution boiling hot; stir until it is dissolved, and crystallize. Ph. U. S. 1863.)

FERRI ET AMMONIÆ TARTRAS, *F. Ammoniotartras*, *Tartrate of Ammonia and Iron*, (Ph. U. S. 1863,) is formed by dissolving hydrated oxide of iron in a solution of bitartrate of ammonia; evaporating, and drying to form scales. Dose, iv to x grains.

FERRI AMMONIOCHLORIDUM, *Ferrum ammoniatum*.

FERRI AMMONIOCI'TRAS, *Ammonie ferrocitras*, *Ferrum ammonioaceticum*, *Ammonioacit'rate of iron*, *Cit'rate of ammonia and iron*. (Prepared by adding ammonia to citrate of iron, so as to neutralize the excess of acid.) In the Ph. U. S. 1863, *Ferri et Ammonie Cit'ras*, *Cit'rate of Iron and Ammonia*, is prepared by evaporating to the consistency of syrup, and drying on plates of glass a mixture of the official *Liquor Ferri Citratis*, Oj; and *Liquor Ammonie*, f℥vj. Dose, gr. v to gr. viij. A POTASSIOCI'TRATE and a SO'DIOCI'TRATE OF IRON have also been introduced; and a CITRATE of the MAGNETIC OXIDE OF IRON prepared by combining the magnetic oxide with citric acid.

FERRI AMMONIOTARTRAS, *Ammoniotartrate of Iron*. (This salt is best made by dissolving to saturation freshly precipitated hydrated oxide of iron in a solution of bitartrate of ammonia, and evaporating to dryness.) It is very soluble. The dose is five grains or more in pill or solution.

FERRI ARSENIAS, *Arsenate of Iron*—f. Borussias, Prussian blue—f. Bromidum, see Bromine—f. Carbonas, F. subcarbonas—f. Carbonas saccharatum, see F. Protocarbonas—f. Carbonas cum Saccharo, see Ferri Protocarbonas—f. Carbonatum, Graphites—f. Carburetum, Graphites.

FERRI CHLORIDUM, *F. Sesquichloridum seu Perchloridum*, (F.) *Trichlorure ou Perchlorure de fer*, *Chloride*, *Sesquichloride*, *Perchloride* or *Muriate of iron*, is obtained by dissolving sesquichloride of iron in muriatic acid, or by the action of muriatic acid and nitric acid on iron. (Ph. U. S. 1863.) It is generally prescribed in the form of *Tinctura Ferri Chloridi*. A concentrated solution of the perchloride has been used as an injection for the radical cure of aneurism, and it has been employed wherever a powerful styptic has been needed. It forms the *Liquor Ferri Perchloridi*, (Ph. B. 1864,) *Liquor Ferri sesquichloridi*, *Liquor Ferri muriatici oxydati*, *Oleum Martis*, *Liquor stypticus Looffi*; and is the basis of *Ellerman's Disinfectant*.

FERRI CI'TRAS, *Ferrum citricum*, *Cit'rate of Iron*. Two citrates of iron are prepared—the *Sesquicitrate* or *Citrate of the Sesquioxide*; and the *Citrate of the Protoxide*. The former, *Ferri citras*, is official in the Ph. U. S., in which it is directed to be prepared (Ph. U. S. 1863) from the official *Liquor Ferri Citratis*. The citrates resemble, in their medical properties, the tartrate and the lactate of the metal. Dose, five grains or more.

FERRI CYANURETUM, Prussian blue—f. Deuto-carbonas fuscus, F. subcarbonas—f. Deutoxydum nigrum, Æthiops martial—f. Ferrocyanas, Prus-

sian blue—f. Ferrocyanidum, Prussian blue—f. Ferrocyanuretum, Prussian blue.

FERRI FULM. The pharmacopœial name (Ph. U. S. 1851) of iron wire.

FERRI HYDRIODAS, F. Iodidum—f. Hydrocyanas, Prussian blue—f. Hypercarburetum, Graphites.

FERRI IO'DIDUM, *F. Ioduretum*, *Ferrum iodatum*, *Iodide*, or *Ioduret of Iron*, (F.) *Iodure de fer*, *Iodure ferreux*; in solution, *Hydriodate of Iron*, *Fer'ri hydriodas*, *Ferrum hydriodatum*, (F.) *Hydriodate de fer*. (Iodin. ℥ij, Ferri rament. ℥j, Aq. destillat. Oiss.) Mix the iodine with a pint of the distilled water, in a porcelain or glass vessel, and gradually add the iron filings, stirring constantly. Heat the mixture gently until the liquid acquires a light greenish colour; then filter; and, after the liquid has passed, pour upon the filter half a pint of the distilled water, boiling hot. When this has passed, evaporate the filtered liquor, at a temperature not exceeding 212°, in an iron vessel, to dryness; keep the dry iodide in a closely stopped bottle. Ph. U. S.) Dose, gr. iij to gr. x or more.

FERRI IODURETUM, F. Iodidum.

FERRI LACTAS, *Ferrum lacticum*, *Lactate of Iron*, L. of *Protoxide of Iron*, (F.) *Lactate de Fer*. (Prepared by digesting in a sand-bath, at a low temperature, diluted lactic acid with iron filings.) It is employed in the same cases as the precipitated subcarbonate of iron, and especially in chlorosis. Twelve grains of the lactate may be given in the 24 hours, in the form of lozenges.

FERRI LIMATU'RA PURIFICATA, *Purified Iron Filings*:—purified by means of the magnet, for internal use. The filings are, also, called *Ferri Scobs*, *F. Ramenta et Fila*, *Fer'ri seu Martis Limatu'ra*, *Speculum Indicum*, *Fer'ri in pulverem resolutio*, (F.) *Limaille de Fer*. They are considered to possess the general properties of iron;—the iron becoming oxidized.

FERRI ET MAGNE'SIÆ CI'TRAS, *Cit'rate of iron and magnesia*, is prepared by dissolving hydrated oxide of iron in a solution of citric acid, saturated with carbonate of magnesia, and evaporating to dryness. It is soluble in water, and does not constipate. Dose, from four grains to fifteen, in solution.

FERRI NITRAS, see *Liquor Ferri nitratis*—f. Nitratis *Liquor*, see *Liquor Ferri nitratis*—f. Oxidum fuscum, F. subcarbonas—f. Oxidum hydratum, *Ferrum oxydatum hydratum*—f. Oxidum Magnetium, *Oxydum Ferri nigrum*—f. Oxidum Nigrum, *Oxydum Ferri nigrum*—f. Oxidum rubrum, *Colcothar*, *Ferri subcarbonas*—f. Perchloridum, see *Ferri chloridum*—f. Percyanidum, Prussian blue—f. Peroxydum hydratum, *Ferrum oxydatum hydratum*—f. Persequintras, see *Liquor Ferri nitratis*—f. Persulphas, see *Ferri sulphas*.

FERRI PHOS'PHAS, *Phosphate of Iron*, (Ferri Sulphat. ℥v, Sodæ Phosphat. ℥vi, Aque cong. Dissolve the sulphate of iron and phosphate of soda, severally, in four pints of the water; then mix the solutions, and set the mixture by, that the powder may subside; lastly, having poured off the supernatant liquor, wash the phosphate of iron with water, and dry it with a gentle heat.—Ph. U. S.) Dose, 5 to 10 grains as a chalybeate. Rarely used.

The British Pharmacopœia (1864) has a *Syrupus Ferri Phosphatis*, *Syr'up of Phosphate of Iron*—gr. iijss of the phosphate to f℥j of syrup—the dose of which is a fluidrachm or more.

FERRI PYROPHOS'PHAS, *Pyrophosphate of Iron*, has been introduced into the Ph. U. S. for 1863.

FERRI ET POTASSÆ TARTRAS, *Ferrum tartarizatum*—f. Potassio-citras, see *Ferri ammonio-citras*—f. Potassio-tartras, *Ferrum tartarizatum*.

FERRI PROTOCARBO'NAS, *Protocar'bonate of Iron*, (F.) *Carbonate de Fer*. The protocarbonate of iron, thrown down from a solution of the sulphate of iron by the carbonate of soda, readily attracts oxygen and becomes converted into the sesquioxide. To prevent this, it may be associated with honey and sugar, and the mixture be reduced by evaporation to a pillular consistence. The mass constitutes the *Ferruginous Pills*, *Pilules ferrugineuses*, of Vallet, *Vallet's Pills*. See *Pilule Ferri Carbonatis*. The Edinburgh Pharmacopœia contained the *Ferri Carbo'nas Sacchara'tum Ferri Carbo'nas cum Sac'charo*, *Sac'charine Car'bonate of Iron*, prepared in this manner. It is the *Ferrum Carbo'nicum Sacchara'tum* of Klauer, the *Ferri Carbo'nas Sacchara'ta*, *Sac'charated Car'bonate of Iron* of the Ph. B. (1864.)

The protocarbonate is given in the same diseases as the lactate of iron; ten or fifteen grains in the course of the twenty-four hours.

FERRI PROTOSULPHAS VIRIDIS, *F. Sulphas—f. Prussias*, Prussian blue.

FERRI PUL'VIS, (Ph. U. S. 1851,) *Ferrum pulverat'um seu reduct'um seu reduct'um* (Ph. U. S. 1863,) *seu metal'licum*, *Powder of iron*, *Redu'ced Iron*, *Queenie's Iron*, (F.) *Fer réduit*, *F. réduit par l'hydrogène*, prepared by passing a stream of hydrogen gas over the sesquioxide of iron, contained in an iron or porcelain tube heated to low redness. It is very liable to become oxidated, and must be kept in a dry, well-stopped bottle. It has been prescribed in anæmic, and especially in chlorotic, cases. The ordinary dose is three to six grains, or more, three times a day, in pill made with sugar and gum. A formula for its preparation from subcarbonate of iron, is given in the Pharmacopœia of the United States, 1863.

FERRI PYROPHOSPHAS, see *Ferri phosphas*.

FERRI ET QUIN'LE CITRAS, *Fer'rum et Chini'num cit'ricum*, *Cit'rate of Iron and Quin'ia*, formed by the union of four parts of citrate of iron and one part of citrate of quinia, has been prescribed in cases where a combination of these tonics was indicated. In the Ph. U. S. 1863, it is formed by decomposing an acid solution of sulphate of quinia by water of ammonia, and adding the precipitated quinia to a solution of citrate of iron until it is dissolved; evaporating to the consistence of syrup, and drying on plates of glass in scales.

FERRI RAMENTA, *F. limatura—f. Rubigo*, *F. Subcarbonas—f. Scobs*, *F. limatura—f. Sesquichloridum*, *Ferri chloridum—f. Sesquinitratum*, *Liquor Ferri nitratis—f. Sesquioxidum*, *F. subcarbonas precipitatus—f. Sodio citras*, see *Ferri ammonio-citras*.

FERRI SUBCARBO'NAS, *F. Carbo'nan*, *F. Rubi'go*, *Fer'rum precipita'tum*, *Chalybis Rubi'go præpara'ta*. (The last two terms, as well as *Ferru'go*, are applied to the subcarbonate or rust, *Eru'go Ferri*, *Cacaferri*, formed by the action of moist air on metallic iron.) A protoxide of iron, oxidized by the decomposition of water; the carbonic acid being attracted from the air. Dose, gr. v to 3ss and more.

Ferri Subcarbo'nas, *Subcar'bonate of iron*, *Precipitated car'bonate of iron*, may be precipitated from sulphate of iron by carbonate of soda. The following is the formula of the Pharmacopœia of the United States:—*Ferri sulph. 3viij; Sodæ carb. 3ix; Aquæ bullient. cong.* Dissolve the sulphate of iron and carbonate of soda severally in four pints of the water, then mix the solutions, and having stirred the mixture, set it by that the powder may subside: having poured off the liquor, wash the subcarbonate with hot water, wrap it in bibulous paper, and dry with a gentle heat. It has, also, been called *Ferri sesquiox'idum*, *Ox'idum ferri fuscum*, *O. ferri ru'brum*, *Deutocarbo'nas Ferri fuscus*, *Cro'cus martis ap'eriens*, *Ses-*

quioxide or peroxide of iron, (F.) *Peroxiide de fer*, *Safran de Mars apéritif*.

FERRI SUL'PHAS, *Sal Martis*, *Vitri'olum Martis* seu *Fer'ri*, *Fer'rum Vitriola'tum*, *Sulphas seu Protosulphas Ferri viridis*, *Calceid'num*, *Cal'catar*, *Cal'catar*, *Chalcant'hum*, *Calceit'a*, *Atrament'um auto'rium*, *Vit'riol*, *Vitri'olum viride*, *An'ima Hel'patis*, *Sul'phate of Iron*, *Green Vit'riol*, *Cop'peras*, (F.) *Sulfate de fer*. The Pharmacopœia of the United States directs it to be made by the action of sulphuric acid 3xviij, on iron wire, cut in pieces, 3xiij—water, a gallon,—evaporating, crystallizing, and drying the crystals on bibulous paper. This salt is inodorous, and of a strong styptic taste. The crystals are light green, transparent, rhomboidal; and soluble in two parts of water. It is tonic and anthelminthic. Dose, gr. j to vj and more.

Ferri Persul'phas, *Persul'phate of Iron*, *Mon'sel's salt*, (F.) *Persulfate de fer*, *Sel de Monsel*, is a powerful coagulant and styptic, introduced by M. Monsel. A formula for the *Liquor Ferri Subsulpha'tis*, *Solution of Subsul'phate of Iron*, *Solution of Persul'phate of Iron*, *Monsel's solution*, is in the Ph. U. S. 1863. (*Ferri sulphat. in pulv. crass. 3ij; Acid sulphur. 3j and gr. xxx; Acid nitric. 3ij and gr. ccc; Ag. destillat. q. s.* To the acids, mixed with half a pint of the water, and heated to the boiling point, add the sulphate gradually, stirring until effervescence ceases. Boil briskly until nitrous vapours are no longer perceived, and the colour is deep ruby red. When nearly cold, add enough water to make it measure f3xiij.)

FERRI SULPHAS CALCINATUM, *Colcothar*.

FERRI SULPHAS EXSICCAT'UM, *F. S. Exsiccata*, (Ph. U. S. 1863,) *Fer'ri Sul'phas sicca'tum*, is *Sulphate of Iron* exposed to a moderate heat, until it is capable of being reduced to powder. In this form it can be made into pills. Three grains may be esteemed equivalent to five of the crystallized salt.

FERRI SULPHURE'TUM *Sul'phuret of Iron*, *Iron pyri'tes*, (F.) *Sulfure de fer*, may be made by heating one part of sublimed sulphur over three parts of iron filings in a crucible until the mass begins to glow. It is employed as a ready means for obtaining hydrosulphuric acid gas by the addition of sulphuric or chlorohydric acid.

FERRI TAN'NAS, *Fer'rum tan'nicum*, *Tan'nate of Iron*, (F.) *Tannate de fer*. This salt is usually obtained by adding a solution of a salt of sesquioxide of iron, as the persulphate, to a decoction of nut-galls. It is possessed of tonic and astringent properties, and has been extolled, especially in chlorosis.

FERRI TARTARUM, *Ferrum tartarizatum—f. Tritoxydum*, *Colcothar*.

FERRI VALERIA'NAS, *Fer'rum Valeria'nicum*, *Vale'rianate of Iron*, (F.) *Valérianate de fer*. (Formed by the action of valerianic acid on oxide of iron.) It is a dark brick-red powder, insoluble in water, and has been given in hysteria complicating chlorosis. Dose, from two to four grains. The Dublin Pharmacopœia of 1850 has a formula for this salt.

FERRICUS HYDRAS, (from *ferrum*), *Ferrum oxydum hydratum*.

FERRIQUE, (from *fer*), *Chalybeate*.

FERROMANGAN'IC, *Ferromangan'icus*. Composed of iron and manganese—as a “ferromanganic preparation.”

FERROSALINE MIXTURE, see *Mixture, ferrosaline*.

FERRUGINEOUS, *Chalybeate*.

FERRUGINOUS, *Chalybeate*.

FERRUGO, gen. *Ferru'ginis*, (*ferrum* and *ru'bigo*, “rust,”) *Ferri subcarbonas*, *Ferrum oxydatum hydratum*.

FERRUM, *Mars*, *Sider'ros*, *Metal'lum hæmatopœt'icum*, *Prun*, (F.) *Fer*. A metal of a bluish-gray colour; fibrous texture; brilliant and fine-

grained fracture. Specific gravity 7.606 to 7.800; hard, ductile, malleable, and magnetic. The medicinal virtues of iron are tonic; producing fetid eructations, when it takes effect, owing to its meeting with acid in the stomach, which oxidizes it, and causes the evolution of hydrogen gas. When given in the metallic state, the filings are chiefly used; but the oxides and salts are most commonly employed.

Ferrum (Ph. U. S. 1863) means *Ferri sulphurum*, and *F. ramentum*.

FERRUM AMMONIACALE, *F. ammoniacum*—*f. Ammoniaco-citricum*, *Ferri ammonio-citras*.

FERRUM AMMONIATUM, *F. ammoniacale*, *Murias Ammoniacae et Ferri*, *Ferri ammonio-chloridum*, *Flores martiales seu salis ammoniaci martiales*, *Ens martis seu Venenae Boylei*, *Sal martis martiale*, *Ammoniacum hydrochloratum ferratum*, *Aloph. Paracelsi*, *Calendulae mineralis*, *Ammoniated Iron*, *Ammonio-chloride of Iron*, (*F.*) *Muriate of ammonia* and *de fer.* A mixture of the hydro-chlorates of ammonia and iron. [*F.*] (*Ferri Subcarb.* 3ij; *Acid muriat.* f3x; *Ammon. muriat.* Biss; *Aq. destillat.* Oiv. Mix the subcarbonate with the acid in a glass vessel and digest for two hours; then add the muriate previously dissolved in distilled water, and having filtered the liquor, evaporate to dryness. Rub to powder.—Ph. U. S.) Dose, gr. iij to gr. xv.

FERRUM ARSENIATUM, *Arsenate of iron*—*f. Arsenicum oxydulatum*, *Arsenate of iron*—*f. Borussicum*, Prussian blue—*f. Bromatum*, see Bromine—*f. Carbonicum Saccharatum*, *Ferri protocarbons*—*f. et Chininum citricum*, *Ferri citras*—*f. Citricum*, *Ferri citras*—*f. Cyanogenatum*, Prussian blue—*f. Haematites*, *Haematites*—*f. Hydriodatum*, see *Ferri iodidum*—*f. Hydrobromicum oxydatum*, see Bromine—*f. Lacticum*, see *Ferri lactas*—*f. Magnes attractorium*, Magnet—*f. Metallicum*, *Ferri pulvis*.

FERRUM OXYDATUM HYDRATUM, *Ferri Oxidum Hydratum* (Ph. U. S.) seu *Peroxidum Hydratum*, *Ferrugo*, *Hydras Ferriens*, *Hydroxide of Iron*, *Hydrated peroxide of Iron*, *Hydrated Tritoxide of Iron*. It may be prepared by taking a solution of sulphate of iron, increasing its dose of oxygen by heating it with nitric acid, and precipitating the oxide by adding pure ammonia in excess, washing the precipitate, and keeping it moist.

In the Ph. U. S. 1863, it is directed to be prepared by adding water of ammonia to a solution of tersulphate of iron; washing the precipitate until the washings pass nearly tasteless. It is then mixed with water and transferred to a wide-mouthed bottle, which must be kept well-stopped.

It has been brought forward as an antidote to arsenic; and many cases of its efficacy have been published.

From 10 to 20 parts of the hydrated oxide would seem to be more than sufficient to convert 1 part of arsenious acid into the basic salt of iron.

FERRUM OXYDULATUM HYDROCYANICUM, Prussian blue—*f. Potabile*, *F. tartarizatum*—*f. Precipitatum*, *Ferri subcarbonas*—*f. Pulveratum*, *Ferri pulvis*—*f. Reductum*, *Ferri pulvis*.

FERRUM TANNICUM, *Ferri tannas*.

FERRUM TARTARIZATUM, (*F.*) *Tartaratum*, *Tartarus Potassae et Ferri*, *Tartarum Ferri*, *Ferri et Potassae Tartaras* (Ph. U. S.), *Tartarus chalybeatus* seu *martialis*, *Mars solubilis*, *Chalybs tartarizatus*, *Ferri potassio-tartaras*, *Tartaras calicoferriens* seu *Potassae ferruginosus*, *Ferri potabile*;—when formed into balls or pellets, called *Globi martiales* and *Globuli Tartari martiales* seu *martiales soluti* seu *martialis* seu *tartarici ferri et lixivii*, *Boli Martis*, and *Pyri martialis*.

les; *Tartarized Iron*, *Tartrate of Potassa and Iron*, (*F.*) *Tartrate de potasse et de fer*; in balls or pellets, *Boules de Mars ou de Nancy ou d'Acier*, (formed from the union of hydrated oxide of iron and bitartrate of potassa, Ph. U. S.) It is one of the mildest of the salts of iron, and not unpalatable. Dose, gr. x to 3ss.

The *Tinctura Martis Aperitiva*, *Tincture of Ludwig*; *Alcohol cum Sulphate Ferri tartarizatus*; *Tinctura Martis Glauberi* is, essentially, a solution of this salt. *Eau de Boule*, a solution of the salt in water, is used in contusions.

The *Boules de Molsheim* are prepared by adding to five parts of the *Boules de Mars* one part of benzoïn and as much turpentine.

Helvetius's Styptic was composed of the filings of iron and tartar, mixed to a proper consistence with French brandy. It was called in England *Eaton's Styptic*; but this is now formed of *Sulphate of Iron*.

FERRUM VALERIANICUM, *Ferri valerianas*—*f. Vitriolatum*, *Ferri sulphas*—*f. Vitriolatum ustum*, *Colcothar*—*f. Zooticum*, Prussian blue.

FERSÆ, Rubeola.

FERTILITY, (*feto*, 'to bear,') Fecundity.

FERULA, 'a staff,' (from *ferio*, 'to strike,') *Fanon*, *Palette*, *Spint*—*f. Africana*, *Bubon galbanum*—*f. Asafoetida*, see *Asafoetida*—*f. Graveolens*, *Anethum graveolens*—*f. Opoponax*, *Pastinaca opoponax*—*f. Persica*, see *Sagapenum*—*f. Tingitana*, see *Ammoniac*, gum.

FERUS, ('savage,') Homicidal.

FESISIAN, Physician.

FESSES, (from *findere*, *fiassum*, 'to divide,') Nates.

FESSIER, Gluteal—*f. Grand*, *Glutæus maximus*—*f. Mojen*, *Glutæus medius*—*f. Petit*, *Glutæus minimus*.

FESTER. A word of unknown etymology, which signifies to corrupt or rankle; and, when applied to a sore, means to suppurate, or rather to discharge a thin fluid as an evidence of its being irritated.

FETID, *Fœtidus*, (*fæto*, 'to stink,') *Dysodæ*, *Cacoæder*, *Bromoësus*, *Graveolens*. Having a bad smell.

FETUS, Fœtus, Pregnant.

FEU ACTUEL, 'fire,' (from [L.] *focus*, [G.] *Feuer*), *Cauterium*—*f. Persique*, *Herpes zoster*, see *Anthrax*—*f. Potentiel*, see *Cauterium*—*f. Sacré*, *Erysipelas*, *Mal des Ardens*—*f. St. Antoine*, *Erysipelas*—*f. Sauvage*, *Ignis sylvaticus*—*f. Volage*, *Ignis sylvaticus*.

FEUILLET, (from *folium*, 'a leaf,') *Omasum*; see *Tache embryonnaire*.

FEUX (pl. of feu) DE DENTS, *Strophulus*—*f. Volages*, *Porrigo larvalis*.

FÈVE, (from *faba*,) *Vicia faba*—*f. de Carthage*, *Habilla de Carthage*—*f. à Cochon*, *Hyoscyamus*—*f. du Calabar*, *Ordeal nut*—*f. Épaise*, *Sedum telephium*—*f. Épreuve du Calabar*, *Ordeal nut*—*f. des Marnais*, *Vicia faba*—*f. Purgatif*, *Ricinus communis*—*f. de Sainte Ignace*, *Ignatia amara*.

FEVER, *Fœbris*, (*feritas*, 'wildness,' or from *feror*, 'heat,' or from *februo*, 'I purify,') *Pyr*, *Pyræus*, (*F.*) *Fièvre*. One of the most frequent and dangerous affections to which the body is liable. A person has an attack of fever, when he is affected with rigors, followed by increased heat of skin, quick pulse, languor, and lassitude. Rigors, increased heat, and frequency of pulse have each been assumed as the essential character of fever. It is not characterized, however, by any one, but depends upon the coexistence of many symptoms. Fevers have been usually divided into the *idiopathic* or *essential*, and the *symptomatic*. The idiopathic arise without any obvious local cause. The symptomatic are dependent

upon local irritation. Idiopathic fevers may be divided into three classes: 1. Those attended with distinct paroxysms:—intermittents. 2. Remittent and continued fevers: and 3. Fevers complicated with eruptions,—the exanthematous. These divisions admit of great variety, owing to climate, season, soil, age, &c. All ages and climates are liable to fever; and its exciting causes are very numerous. These causes may be common; as irritations in the intestines; external injuries, stimulants, &c.; or they may be *specific*; as miasmata, contagion, &c. The greatest diversity has prevailed regarding the theory of fever. Its primary seat has been placed in the brain, mucous membrane of the stomach and intestines, skin, nerves, blood-vessels, liver, vena cava, pancreas, &c. It would seem, however, that although, in fever, the whole of the functions are morbidly impressed, the arguments in favour of the impression being first made on the nervous system and the system of nutrition are the strongest. The exciting cause of fever, whatever it may be, produces an irregular action in the system of nutrition, which is soon conveyed to the rest of the system, owing to the extensive sympathy which exists between every part of the body; and it is probable, that all those local inflammations and congestions are the consequence, rather than the cause, of this disordered condition of the system. The general character of fever is clearly shown by examination of the blood. When fever is devoid of inflammatory complication, the quantity of fibrin is in no case augmented. It frequently remains in the healthy proportion, and at times diminishes to an extent not met with in any other acute disease. The alteration of the blood in fevers, which consists generally in a diminution of the fibrinous element, is the reverse of what occurs in inflammation.

Many phenomena of fever are influenced by that *periodicity*, which we notice in the exercise of several of the functions of the body. The types of intermittents are strong evidences of such an influence.

In the treatment of fever, it is important to bear in mind, 1. Its tendency, particularly in the case of the exanthemata, to run a definite course, and terminate in restoration to health. 2. The disposition to local determination or hyperæmiæ:—the most frequent cause of the fatal termination of fever; a circumstance requiring the vigilant attention of the physician. 3. That the symptoms must be attentively studied, in order to deduce, as far as possible from them, the indications of cure. Lastly, attention must be paid to the prevalent epidemic. There are particular seasons in which fevers are very malignant; and others in which they are as mild; circumstances which necessarily have an effect upon the treatment.

FEVER AND AGUE, Intermittent Fever—f. Acellimating, F. strangers'.

FEVER, A'DENOMENINGE'AL, *Fe'bris a'denomeninge'a seu mesenterica* (BAGLIVI) *seu mucosa*, *Morbus mucosus*, *Gastroduodenop'gra*, *Mucous* or *Pituitous* or *Catar'hal Fe'ver*, *Phlegmap'gra*, *Phlegmop'gra*, *Phlegmatop'gra*, *Gastrobronchitis*. Fever, accompanied with considerable mucous secretion; especially from the digestive tube.

FEVER, ADENOMERVOUS, Plague.

FEVER, ADYNAM'IC, *Fe'bris adynamica*, *Asthenicop'gra*, *Asthenicoppyr'etus*, *Asthenop'gra*, *Asthenopyr'etus*. Fever attended with great prostration of the vital powers—as Typhoid and Typhus fever.

FEVER, AF'RICAN, *Fe'bris Africa'na*. The malignant bilious remittent fever, which prevails on the western coast of Africa.

FEVER, ALGID, see Algidus.

FEVER ANOM'ALOUS, *Fe'bris Anom'ala seu er-*

rat'ica seu irregula'ris seu va'ga. A fever, whose progress and type are irregular.

FEVER, APHON'IC, *Fe'bris apho'nica*. A variety of intermittent, in which the voice is lost during the paroxysm.

FEVER, APOPLEC'TIC, *Fe'bris apoplec'tica seu comato'des seu lethar'gica seu soporo'sa*, *Comatose Fe'ver*. An intermittent or continued fever, attended with apoplectic symptoms.

FEVER, ARTHRIT'IC. Fever, characterized by inflammation of the joints,—as in gout and rheumatism.

FEVER, ARTICULAR ERUPTIVE, Dengue—f. Articular rheumatic, Dengue.

FEVER, ARTIFIC'IAL, *Fe'bris artificia'lis seu ar'te promo'ta*. Fever produced designedly by the internal or external use of stimulants.

FEVER, ASTHEN'IC, *Fe'bris asthen'ica*. Fever accompanied with debility. It may include every variety of fever under certain circumstances, but is generally appropriated to typhus.

FEVER, ASTHMAT'IC, *Fe'bris asthma'tica*. A pernicious intermittent, accompanied with symptoms of asthma.

FEVER, ATAXOADYNAM'IC, *Fe'bris atax'oüdyn'amica*. Fever characterized by symptoms of ataxia and adynamia.

FEVER, BARCELONA, Fever, yellow—f. Bastard, Illegitimate fever.

FEVER, BIL'IOUS, *Fe'bris bilio'sa seu polychol'lica seu aso'des seu choler'ica seu gas'trica seu hepat'ica seu cholepyret'ica*, *Cholep'gra*, *Cholepyr'etus*, *Hepatogastrocholo'sis*, *Enterop'gra bilio'sa*, *Cholozoe'mia febrilis*, *Syn'ochus bilio'sa*. The common remittent fever of summer and autumn; generally supposed to be owing to, or connected with, derangement of the biliary system.

FEVER, BILIOUS REMITTING, YELLOW, Fever, yellow—f. Bladdery, Pemphigus.

FEVER BLIS'TER. A term, vaguely applied to vesicular or pustular eruptions, which appear particularly about the mouth, during or after febrile disturbance.

FEVER, BOÂ VISTA. A malignant bilious remittent fever, greatly resembling yellow fever, which was very fatal at Fernando Po, and in ships in its waters, in the year 1845.

FEVER, BONA. A malignant paludal fever, which prevailed amongst the troops of the garrison at Bona in Algeria, from 1832 to 1835.

FEVER, BONE, see Inflammation—f. Bouquet, Dengue—f. Brain, F. cerebral, *PLR'itis*—f. Brain, water, *Hydrocephalus internus*—f. Breakbone, Dengue—f. Bucket, Dengue—f. Bulam, Fever, yellow.

FEVER, CAMP. Any continued fever occurring in camp or in the army. Formerly it meant the typhus of armies. See Typhus gravior.

FEVER, CARDIAL'GIC, *Fe'bris Cardial'gica*. A variety of pernicious intermittent, accompanied with violent cardialgia during the paroxysm.

FEVER, CARDIT'IC. Intermittent fever, accompanied with pain at the heart.

FEVER, CATARRHAL, F. adenomeningeal.

FEVER, CEPHALAL'GIC, *Fe'bris cephalal'gica*. A pernicious intermittent, accompanied with intense pain of the head. Also, intermittent cephalalgia.

FEVER, CEPHAL'IC, *Fe'bris cephal'ica*. A febrile affection of children—intermediate between the acute form of active cerebral congestion and the chronic form—which is attended by fever, pain in the head, disorder, or more generally constipation of the bowels, and a train of phenomena often supposed to be premonitory of an attack of hydrocephalus.

FEVER, CEREBRAL, *Brain fe'ver*. Fever, generally of an ataxic character, in which the brain is considerably affected.

FEVER, CHA'GRES, *Panama' Fe'ver*. A severe

form of endemic fever, common on the isthmus of Panama.

FEVER, CHILDBED, *Febris puerperal*, *Puerperal Fever*—f. Childbed, low, see Peritonitis.

FEVER, CHOLERIC, *Febris choleric*. A variety of pernicious intermittent, accompanied with symptoms of cholera morbus.

FEVER, CHOLERIC, OF INFANTS, *Cholera infantum*.

FEVER, CHRONIC, *Febris Chronica*. Some authors apply this name to protracted fevers; others to hectic fever.

FEVER, COLLIQUATIVE, *Febris tabida seu colliquativa*. Fever, characterized by rapid emaciation, copious evacuations, and rapid prostration of strength.

FEVER, COMATOSE, *F. apoplectic*.

FEVER, CONGESTIVE, *Febris congestiva*. Fever accompanied by obscure symptoms; or by great oppression and depression; in which it is difficult—and often impossible—to induce reaction. Congestive fevers occur in various parts of this country, especially in the fall; and they are very common in India. The term congestive fever is often used in some parts of the south of the United States very indefinitely,—to include winter typhus, and typhoid fevers, typhoid pneumonia, as well as intermittents and autumnal remittents.

FEVER, CONTINENT, see Continent and Continued fever—f. Continual, see Continued fever—f. Continued, common, Synochus.

FEVER, CONVULSIVE, *Febris convulsiva*. A pernicious intermittent or remittent, accompanied by convulsions.

FEVER, COUNTRY. Fever occasioned by exposure to the miasms of a highly malarious locality. A term much used in the Southern States.

FEVER, DELIRIOUS, *Febris cum delirio*, (*F.*) *Fièvre délirante*. A pernicious intermittent, characterized by delirium in the paroxysms.

FEVER, DEPURATORY, *Febris depuratoria*. A fever, to which was attributed the property of purifying the blood; or which indicated, that such a supposed depuration had occurred.

FEVER, DIAPHORETIC, *Febris diaphoretica*, *Febris sudatoria*. A pernicious intermittent, with excessive sweating during the fit.

FEVER, DIARY, Ephemera.

FEVER, DIGESTIVE. The chilliness, followed by increased heat and quickness of pulse, which frequently accompanies digestion.

FEVER, DOUBLE, (*F.*) *Fièvre double ou doublée*. An intermittent, which has two paroxysms in a given time, instead of one.

FEVER, DOUBLE QUARTAN. A fever, whose paroxysms occur two days in succession, and fail the third day; the first paroxysm resembling the 4th, and the second the 5th.

FEVER, DOUBLE QUOTIDIAN. An intermittent, whose paroxysms return twice every day at corresponding hours.

FEVER, DOUBLE TERTIAN. An intermittent, whose paroxysms return every day; the first corresponding with the 3d, the second with the 4th, and so on.

FEVER, DUODECIMANE, *Febris duodecima'na*. A supposititious intermittent, whose paroxysms recur on the 12th day, or every 11 days.

FEVER, DYNAMIC, Synocha—f. Endemial, Remittent *F.*—f. Endemic, Remittent *F.*—f. Endemical, Remittent *F.*—f. Enteric, see Typhus—f. Ephemeral, Ephemera.

FEVER, EPILEPTIC, *Febris epileptica*. A variety of pernicious intermittent, accompanied with attacks of epilepsy.

FEVER, EROTIC, *Febris erotica*, (*F.*) *Fièvre d'amour*. A chronic fever, occasioned by unpropitious love.

FEVER, ERRATIC, see Erratic—f. Eruptive, Ex-

anthematica—f. Eruptive, articular, Dengue—f. Exacerbating, Remittent Fever.

FEVER, FAINTING, OF PERSIA. A singular and fatal epidemic, presenting some points of analogy with cholera, which prevailed at Teheran in the autumn of 1842.

FEVER, FAMINE, *Febris à faim*. The febrile phenomena induced by want of food.

FEVER, FIVE DAY, see Relapse.

FEVER, GANGRENOUS, *Febris gangrenodes*. Fever, accompanied by gangrene of various parts, and especially of the limbs and genitals. Sénac describes an intermittent of this kind.

FEVER, GASTRALGIC. An intermittent accompanied with acute burning lacerating pain at the stomach.

FEVER, GASTRIC, *Febris gastrica seu gastrica biliosa*, *Gastropyræ*, *Gastropyrætic*, *Gastrocholosis*, *Hepato-gastrocholosis*, *Stomachic fever*, *Synochus Biliosa*. A name given by some to bilious fever, which has appeared to them to be dependent on an affection of the stomach. Also, called, *Meningogastric fever*, *Febris meningogastrica*, *Meningogastri'tis*, *Gastromeningi'tis*, *Har'vest fever*, (*F.*) *Fièvre de la Moisson*, *F. Meningo-gastrique*.

FEVER, GASTRO-ADYNAMIC, *Febris gastro-ady-namica*. A fever, in which the symptoms of bilious fever are joined with those of adynamic fever.

FEVER, GASTRO-ANGIOTENIC. A fever, in which the symptoms of bilious are united with those of inflammatory fever.

FEVER, GASTRO-ATAXIC. A fever, in which the symptoms of bilious fever are united with those of ataxic fever.

FEVER, GIBALTAR, Fever, yellow—f. *Hæmagastria*, *F. yellow*.

FEVER, HÆMORRHOIC, *Febris hæmoptica*. A variety of masked intermittent, in which periodical hæmoptysis is the chief symptom.

FEVER, HARVEST, *F. Gastric*.

FEVER, HAY, *Catarrhus æstivus*, *Chortanth'ma*, *Summer Catarrh*, *Hay Asthma*, *Rose Catarrh*, *Summer Bronchitis*. A catarrh to which certain persons are subject in summer, and which has been ascribed in England to the effluvium of hay, but this is not the probable cause. It is a catarrh with sneezing, headache, weeping, snuffling and cough, with, at times, fever and general discomfort. It is not uncommon in this country. It disappears spontaneously,—to recur on subsequent years about the same period.

FEVER, HEBDOMADAL. A supposititious fever, whose paroxysms return weekly, and on the same day.

FEVER, HECTIC, see Consumption, and Hectic Fever.

FEVER, HEPATIC or HEPATALGIC. A pernicious intermittent, with violent pain in the right hypochondrium.

FEVER, HILL. A modification of remittent, occurring in the hilly districts of India.

FEVER, HOSPITAL, Typhus gravior.

FEVER, HUMORAL, *Febris humoralis*. Fever in which an alteration or deterioration of the humors is suspected.

FEVER, HUNGARY, *Febris Hungarica seu Pan-nonica*, *Lu'es Pannonia*, *Morbus Ungaricus*, *Cephalonismus*, *Ver'nis cer'ebri*, (*F.*) *Fièvre Hongroise ou de Hongrie*. An epidemic typhus, common amongst the soldiers in barracks, in Hungary.

FEVER, HYDROPHOBIC, *Febris hydrophobica*. Pernicious intermittent, with dread of liquids.

FEVER, HYSTERIC, *Febris hysterica*. Hysteria, accompanied by fever. Hysteria, occurring with each paroxysm of an intermittent.

FEVER, ICTERIC, *Febris icterica*. Fever, accompanied with jaundice. Some intermittents exhibit this complication at each paroxysm.

FEVER, ICTERIC, REMITTING, see Relapse.

FEVER, INFANTILE REMITTING, *Febris infantium remittens*, *F. verminosa*, (of many), *F. mucosa verminosa*, *F. lenta*, *Spurious wormfever*, *Remitting fever of children*, *Hectica infantilis*. A fever occurring in childhood, which often assumes many of the characters of hydrocephalus. It appears generally to be dependent upon a morbid condition of the stomach and bowels.

FEVER, INFLAMMATORY, Synocha.

FEVER, INSIDIOUS. Fever, which, at first, seems devoid of danger, but may subsequently become of a more or less malignant character.

FEVER, INTERMITTENT, see Intermittent fever—*f.* Irritative, Irritation, morbid—*f.* Jail, Typhus gravior—*f.* Jungle, see Jungle fever.

FEVER, LAKE. Fever produced by exposure to the malaria in the neighbourhood of the northern lakes of this country.

FEVER, LENTICULAR, Miliary fever.

FEVER, LOCHIAL, *Febris lochia*. That acceleration of the circulation which sometimes occurs during the discharge of the lochia.

FEVER, LUNG, Catarrh, Pneumonia—*f.* Maculated, Typhus.

FEVER, MALARIAL or MALARIOUS. Fever induced by malarious or miasmatic emanations; hence also called *Miasmatic fever*. Intermittent and remittent fever belong to this class.

FEVER, MALIGNANT, *Febris maligna*. Fever which may make its approaches insidiously, and subsequently becomes formidable. Any fever which exhibits a very dangerous aspect. *Typhus gravior*.

FEVER, MALIGNANT PESTILENTIAL, Fever, yellow—*f.* Marsh, Elodes (febris).

FEVER, MASK'ED, *Febris larvata*, (*F.*) *Fièvre larvée*; *Dead or Dumb Ague*, *Dumb Chill*. An anomalous intermittent, the paroxysms of which have not the regular stages, or are *masked*.

FEVER, MENINGOGASTRIC, Fever gastric.

FEVER, MESENTERIC, *Febris mesenterica*. A name given, by Baglivi, to a species of fever which appears to have belonged either to the *mucous* or *bilious*.

FEVER, MIASMATIC, *F.* Malarial.

FEVER, MILK, *Febris lactea*, *Galactopyra*, *Galactopyretus*, (*F.*) *Fièvre de lait ou lacteuse ou lactée*. The fever which precedes or accompanies the secretion of milk in women recently delivered. It comes on generally about the third day after delivery, and is characterized by quick pulse; increased heat; redness of face; diminution or temporary suspension of the lochial discharge; tumefaction and tension of the breasts. It commonly terminates in twenty-four hours, and often with profuse perspiration. It requires the use of antiphlogistics, with dry diet.

FEVER, MIXED, Synochus.

FEVER, MOUNTAIN. Fever belonging to mountainous regions. A form of fever, said to be peculiar to the elevated regions of the Rocky Mountains; and seen only at an altitude of 7000 feet and upwards.

FEVER, MUCOUS, *F.* Adenomeningeal.

FEVER, NEPHRITIC, *Febris nephritica*. Intermittent fever, accompanied with nephritic pain during the paroxysm.

FEVER, NERVOUS, *Febris nervosa* seu *neurodes* seu *lenta nervosa*, *Neuropyra*, *Neuropyræticus*. A variety of Typhus; the *Typhus mitior* of Cullen; now, generally, however, it is esteemed a distinct disease. See Typhoid fever.

FEVER, NERVOUS, WITH EXANTHEMATOUS ERUPTION, Typhus.

FEVER, NEURALGIC, Dengue.

FEVER, NIGER. A malignant fever, of the bilious remittent kind, which proved fatal to many in the expeditions sent out by the British

government to explore the Niger, in the years 1841-42, and previously.

FEVER, NON'ANE, *Febris non'ana*. A supposititious fever, whose paroxysms recur every ninth day, or every eight days.

FEVER, OCTANE, *Febris oct'ana*. An intermittent, whose paroxysms recur every eighth day.

FEVER, PALUDAL, Intermittent—*f.* Panama, Fever, Chagres—*f.* Paroxysmal, Remittent fever—*f.* Periodic, *F.* Periodical, Intermittent.

FEVER, PERIODICAL or PERIODIC, *Febris period'ica*. An intermittent or remittent fever.

FEVER, PERNICIOUS, *Febris pernicios'a*. Intermittent fever, when attended with great danger, and which destroys the majority of those affected by it in the first four or five paroxysms; sometimes in the very first.

FEVER, PESTILENTIAL, *Febris pestilential'is*. The *Plague*. Also a severe case of typhus. The yellow fever and sweating sickness have, likewise, been thus designated.

FEVER, PESTILENTIAL, OF CATTLE, Murr—*f.* Pettechial, Typhus—*f.* Ptituitous, *F.* adenomeningeal.

FEVER, PLEURITIC. An intermittent or remittent, accompanied with inflammation of the pleura.

FEVER, PNEUMONIC, *Febris pneumon'ica*. An intermittent, accompanied with inflammation of the lungs. Also, pneumonia.

FEVER POISON, see Poison.

FEVER, PSEUDO, Irritation, morbid.

FEVER, PUERPERAL, *Febris puerper'al'is*, *Child-bed fever*, (*F.*) *Fièvre puerpérale*. This name has been given to several acute diseases, supervening on delivery. It means, generally, a malignant variety of peritonitis, which runs its course very rapidly, and passes into a typhoid condition, unless met, at the very onset, by the most active sedative measures. By the generality of practitioners, it is esteemed to be eminently contagious; some, however, deny that it is so. See Peritonitis, and Puerperal fever.

FEVER, PUERPERAL, ADYNAMIC or MALIGNANT, see Peritonitis—*f.* Puking, Milk sickness.

FEVER, PU'RULENT, *Febris purulen'ta*. Fever, which accompanies suppuration.

FEVER, PUTRID, Typhus gravior—*f.* Pythogenic, Typhoid fever.

FEVER, QUINT'AN, *Febris quinta'na*. A fever, whose paroxysms return every fifth day. It is rarely or never seen.

FEVER, QUOTIDIAN, see Quotidian.

FEVER, QUINIC, (*F.*) *Fièvre quinqué*, *F. de Quinquina*. A fever, accompanied by a cutaneous eruption, said to affect those who are engaged in the preparation of the sulphate of quinia. The French use the term *Fièvres à quinquina* for intermittents, which cinchona alone cures.

FEVER, RECURRENT, see Relapse—*f.* Red Tongue, see Typhus.

FEVER, REG'ULAR, *Febris regula'ris* seu *exquisi'ta*. An intermittent whose paroxysms follow a determinate type. It is opposed to *atypic*. Sometimes opposed to *anomalous*.

FEVER, RELAPSING, see Relapse.

FEVER, REMITTENT, see Remittent Fever—*f.* Remittent, Bilious, see Relapse—*f.* Remittent of children, *F.* infantile remittent—*f.* Remittent, infantile, see Fever, infantile remittent—*f.* Remitting icteric, see Relapse—*f.* Rheumatic, Rheumatism, acute.

FEVER ROOT, *Triosteum perfoliatum*.

FEVER, SCORBU'TIC, *Febris scorbut'ica*. The febrile movement, which sometimes accompanies scorbutus or scurvy.

FEVER, SEASONING, *F.*, strangers'.

FEVER, SEC'ONDARY. A febrile condition which recurs in certain affections after having ceased; such as the secondary fever, which comes on at the time of the maturation of the variolous pus-

tules, or as the eruption of scarlatina, &c., disappears.

FEVER, SEPT'AN, *Febris septuana*. An intermittent, whose paroxysms recur every six days, and consequently on the seventh.

FEVER, SEVEN DAY, see Relapse.

FEVER, SEX'TAN, *Febris sexta-na*. A fever, which recurs every five days, and consequently on the sixth.

FEVER, SHIP, see Typhus.

FEVER, SIM'PLE, *Febris Sim'plex*. Simple fever is that which has no predominant character—bilious, inflammatory, or nervous; and which is unaccompanied by any local determination, hyperæmia, or complication. It may be continued, remittent, or intermittent.

FEVER, SIMPLE CONTINUED. This is the most favourable form of continued fever, and has a tendency to wear itself out, provided only the *lætentia* be avoided. The prognosis is consequently favourable, and the treatment simple; consisting in perfect repose of body and mind, abstinence, and relieving the thirst by cold drinks.

FEVER, SINGUL'TOUS, *Febris singultuosa* seu *lygmulosa* seu *lyngo-dea*. Fever, accompanied with singultus or hiccough.

FEVER, SOLAR, Dengue—f. Spotted, Typhus gravior.

FEVER, STER'CORAL, *Febris stercoralis*. Fever, produced by an accumulation of feces in the intestines.

FEVER, STOMACHIC, Gastric Fever.

FEVER, STRA'NGERS', *Acclimating* or *Seasoning fever*. Yellow, or remittent fever, which is endemic in certain places, and to which strangers are especially liable.

FEVER, SUBCONTINUAL, Remittent Fever.

FEVER, SUBIN'TRANT, *Febris subintranea*. An intermittent, in which one paroxysm is scarcely finished before the other begins.

FEVER, SUN, Dengue—f. Sweating, Sudor Anglicus.

FEVER, SYN'OPAL, *Febris synopalis* seu *minuta*, (F.) *Fèvre synopale*. A variety of pernicious intermittent, in which there is, in every paroxysm, one or more faintings.

FEVER, SYNCHOID, Synchosis.

FEVER, SYPHILITIC, *Febris syphilitica*. Fever, accompanying syphilis, or supposed to be owing to a syphilitic taint.

FEVER, TER'TIAN, *Febris tertiana*. A fever, whose paroxysm returns on the third day, and consequently every two days.

FEVER, THIRD DAY, Tertian.

FEVER, TRAG'IC, *Febris Trag'ica*. A low fever, in which the patient declaims like an actor during the delirium.

FEVER, TRAUMAT'IC, *Febris traumatica*. The fever, which supervenes on wounds or great surgical operations.

FEVER, TREMBLING, Intermittent fever.

FEVER, TY'PHOMALARIAL, *Miasmatic typhoid fever*. A fever in which there is a mixture of the phenomena of typhous and malarial fevers;—a combination not unfrequently seen in active military service.

FEVER, TYPHOID, see Typhus—f. Typhoid, of India, Cholera—f. Typhous, Typhus—f. Varicellar, see Varioloid—f. Varioloid, see Varioloid.

FEVER, VER'MINOUS, *Febris verminosa*, *Helminthopyra*, *Worm fever*. Fever, produced by the presence of worms in the digestive tube, or accompanied by their expulsion.

FEVER, VER'NAL, *Febris vernalis*. An intermittent or other fever occurring in the spring. Vernal intermittents were formerly considered salubrious.

"An ague in the spring
Is physic for a king."

FEVER, VESICULAR, *Femphigus*.

FEVER, WAL'CHEREN, *Gallisch-reuss*. The remittents and intermittents to which the British troops were exposed, who were attached to the expedition to Walcheren, in 1809.

FEVER, WATER BRAIN, Hydrocephalus internus—f. Winter, see Tongue, black—f. Worm, Verminous F.—f. Worm, spurious, Fever, infantile remittent.

FEVER, YEL'LOW, *Febris fla'va*, *F. seu Pestis Americana*, *F. fla'va Americana*, cum, *F. tropica*, *F. continua putrida ictero'des Carolinensis*, *F. elod'es ictero'des*, *F. maligna biliosa America*, *F. maligna fla'va India Occidentalis*, *F. tor'ica*, *Cholæria Americana*, *Ochrotyphus*, *Laimocholæria*, *Pestilentia hamagutrica*, *Pestis occidentalis* seu *intertropica*, *Vomitus niger*, *Epar'ius malignus fla'vus*, *Remittens ictero'des*, *Tritropica Americana*, *Typhus ictero'des*, *Ochrotyphus*, *Synochus ictero'des*, *Endemial Causus of the West Indies*, *Causus tropicus endemius*, *Bilious remitting yellow fever*, *Malignant pestilential fever*, (F.) *Fèvre jaune d'Amérique*, *F. matelote*, *F. gastrodynamique*, *F. de la Barbade*, *F. de Siam*, *F. Ictérique*, *F. Gastro-hépatique*, *Typhus miasmatic atarique putride jaune*, *T. jaune*, *T. d'Amérique*; *Vomito prieto*, *Vomito negro*, *Mal de Siam*, *Hamagutric Fever* or *Pestilence*, *Black vomit*, *Bulam*, *Gibral'tar*, and *Barcelona Fever*,—only, in the West Indies, *Malat' to Jack* and *Yellow Jack*. A very acute and dangerous febrile affection; so called, because complicated, in its second stage, with a jaundiced appearance, accompanied by vomiting of black matter. Sauvages, Cullen, and others regard it as a variety of typhus; and Pinel, as a species of gastro-dynamic or bilious putrid fever. It occurs, endemically, annually within the tropics; but is met with epidemically in the temperate regions. Broussais regards the disease as gastro-enteritis, exasperated by atmospheric heat; so that it runs through its stages with much greater rapidity than the gastro-enteritis of our climates. The yellow colour of the skin, according to him, is owing to the inflammation of the small intestine,—and especially of the duodenum,—augmenting the secretion of the liver, and at the same time preventing its discharge into the duodenum. The pathology of this affection, as well as its origin, is still unsettled. The treatment must generally be of the most active nature at the onset; consisting in bleeding largely, and exhibiting mercury, so as to excite a new action, if possible:—the other symptoms being combated according to general principles. It must vary, however, according to the epidemic.

FEVER, YELLOW, MILD, see Relapse.

FEVERBUSH, *Laurus benzoin*, *Prinos*.

FEVER DROPS, WARBURG'S, see Beheert.

FEVERET, *Febricula*.

FEVERFEW, (*Febris* and *fugare*, 'to drive away,') *Matricaria*.

FEVERFOULLIE, *Matricaria*.

FEVERISH, *Feverens*, *Febriens*, *Febriacus*, (F.) *Fibreux*. That which causes fever, or is febrificient or febrific; as *feverish food*, *feverish diathesis*, &c. Also, the state of one labouring under fever, *Feverick*. If affected with intermittent fever, he is said, in Suffolk, England, to be *a'garified*.

FEVERISHNESS, *Febriçitatio*, *Febriçulatio*. The state of having fever. A slight febrile disorder.

FEVEROUS, *Feverish*.

FEVERROOT, *Pterospora andromeda*.

FEVERISH, see *Feverish*.

FEVERTREE, *Pinckneya pubens*.

FEVERWOOD, *Laurus benzoin*.

FEVERWORT, *Eupatorium perfoliatum*, Triosteum.

FEWS, *Sempervivum tectorum*.

FEY, Psychorages.

FEYK, Fidgets.

FIBER, ([G.] Biber,) Castor fiber.

FIBRA, Fibre—*f. Aurea*, Coptis—*f. Nervea*, Nerve-fibre—*f. Sanguinis*, Fibrin.

FIBRÆ ARCIFORMES, see Arciform.

FIBRE, *Fibra*, *Is*, *Cle'don*, *Filum*. An organic filament, of a solid consistence, and more or less extensible, which enters into the composition of every animal and vegetable texture. The *simple* or *elementary fibre* of the ancients, from a particular assemblage and arrangement of which every texture of the body was conceived to be constituted, is entirely ideal. The moderns usually admit, with Haller and Blumenbach, *three elementary fibres or tissues*. 1. The *cellular* or *areolar*, formed chiefly of thin plates, of a whitish colour, and extensible, which seems to consist of concrete gelatin. 2. The *nervous*, *pulpy*, or *medullary*, formed of a soft substance, contained in a cellular sheath, and consisting of albumen united to a fatty matter. 3. The *muscular*, composed of round filaments, of a grayish or reddish colour, and formed of fibrin. Chaussier has added to these the *albugeous fibre*, but it seems to differ from the cellular fibre only in greater condensation of the molecules. See Fibrous.

A very small or ultimate fibre is called a *Fibril* or *Fibrilla*.

FIBRE, ALBUGINEOUS, see Albuginea—*f. Lisse*, see Muscular fibre—*f. Striée*, see Muscular fibre.

FIBRES, CONVERG'ING. Nervous fibres, whose office it is to associate different portions of the nervous centres with each other. They form the *Commissures*.

FIBRES, DIVERG'ING. The fibres composing the columns of the medulla oblongata, which separate in their progress to the periphery of the cerebrum and cerebellum.

FIBRES OF MÜLLER, see Tunica Jacobi.

FIBRES OF REMAK'. Fibres described by Remak as peculiar to the sympathetic nerve, but which Valentin considers to be neurilemma, and to consist of fibrocellular bundles.

FIBREUX, Fibrous.

FIBRIL, see Fibre.

FIBRILLA, (dim. of *fibra*,) Fibril—*f. Muscularis*, Muscular fibre.

FIBRILLAR, *Fibrilla'ris*. Relating or appertaining to fibrils.

FIBRILLATION, *Fibrilla'tio*. The formation of fibrils.

FIBRIN, (from *fibre*,) *Fibrine*, *Fibrina*, *Fibrinè*, *Fibra'san'guinis*, *Mat'eria fibro'sa*, *Lymph'a plast'ica*, *Mat'ière fibreuse du sang*. An immediate animal principle—solid, white, and inodorous; insipid; heavier than water; without action on the vegetable blues; elastic, when moist; hard and brittle when dry. It enters into the composition of the chyle and the blood; is the self-coagulable material in them, and is produced in the retrograde metamorphosis of the tissues. It is in small proportion in the blood; but its cipher increases in inflammation, whilst in continued fever it is diminished. *Muscular fibrin*, *Syn'tonin*, *Musculin*, has been shown, however, to be different from that of the blood. In certain diseased actions, *Fibrin* or *Coagulable lymph*, *Gluten*, is separated from the blood, and is found in considerable quantity on the surfaces of membranes, and in the cavities of the body. See Liquor Sanguinis.

Fibrin is likewise a proximate principle of vegetables, and differs but little in chemical composition from animal fibrin; nor does it differ much from albumen and casein.

Fibrin is very nutritious.

FIBRINATION, *Fibrina'tio*. The act of add-

ing fibrin to the blood. The opposite to *defibrination*.

FIBRINOGENOUS. (A hybrid term, from *fibrin*, and *γενω* or *γεννω*, 'I engender.') Producing fibrin. The fibrinogenous or fibrin-producing material appears to be a product of the metamorphosis of tissue.

FIBRINOUS, *Fibrino'sus*, (F.) *Fibrineux*. That which is composed of fibrin, or has the appearance of fibrin.

FIBRO, in composition, appertaining or relating to fibrous tissue.

FIBROAREOLAR, Fibrocellular.

FIBROBRONCHITIS. Fibrous or rheumatic inflammation of the bronchial tubes.

FIBROCAR'TILAGE, *Fibrocartila'go*. An

organic tissue, partaking of the nature of the fibrous tissue, and of that of cartilage. It is dense, resisting, elastic, firm, supple, and flexible. Fibrocartilages are distinguished into,—1. *Membrani-form*, or those which serve as moulds to certain parts, as the alæ nasi and eyelids. 2. *Vag'uniform*, or those which form sheaths for the sliding of tendons. 3. *Interartic'ular*, those which are met with in the movable articulations. 4. *Unif'ing*, (F.) *Fibro-cartilages d'union*, which form a junction between two bones, as the symphysis pubis. Fibrocartilages are sometimes formed adventitiously, as the result of a morbid process in different organs.

FIBROCAR'TILAGES, TARSAL, see Tarsus.

FIBROCAR'TILAGINES INTERVERTEBRALES, Intervertebral cartilages.

FIBROCELLULAR, *Fibrocellula'ris*, *Fibroareolar*, *Fibroareola'ris*. That which partakes of the fibrous and of the cellular or areolar tissue, as the fascia transversalis.

FIBROCELLULAR TUMOUR, Tumour, fibrocellular.

FIBROMA, gen. *Fibro'matis*, (*fibro*, and *oma*,) Tumour, fibrous.

FIBROMU'COUS, *Fibromuco'sus*, (F.) *Fibromuqueux*. Possessing the nature of fibrous and of mucous membranes. A term applied to fibrous membranes, which are intimately united with other membranes of a mucous nature, as the *pituitary membrane*, the *membrane of the urethra*, &c.

FIBROPLAS'TIC, *Fibro-plas'ticus*, (F.) *Fibroplastique*, (from *fibro*, and *πλασσω*, 'I form.') An epithet for a morbid formation, constituted of the elements of areolar tissue transformed, in part, into fibre.

FIBROPLASTIC TUMOUR, Sarcoma.

FIBROSE'ROUS, *Fibrosero'sus*, (F.) *Fibroseréux*. Possessing the nature of fibrous and serous membranes. Membranes, composed of a fibrous, and a serous sheet, intimately united:—as the *Dura Mater*, *Pericardium*, *Tunica albuginea testis*, &c.

FIBROID, Fibrous, Tumour, fibrous.

FIBROUS, *Fibro'sus*, *Fibroid*, (F.) *Fibreux*. Composed of fibres. Certain membranes, as the *dura mater*, *periosteum*, *ligamentous capsules of the joints*, &c., are *fibrous*. The *fibrous system* of Bichat includes the system of organs formed by the *albugeous fibre* of Chaussier. It comprises, particularly, the *periosteum* and *perichondrium*; the *articular capsules* and *ligaments*; the *tendons*; the *dura mater*, *pericardium*, *tunica sclerotica*, *tunica albuginea testis*, *outer membrane of the spleen*, &c. Under *simple fibrous tissues*, *Tel'ia fibro'sa*, certain writers have classed the *white* and *yellow fibrous tissues*, and areolar tissue. Both the *yellow* and the *white* may be detected in the areolar tissue. The *white* is said to exist alone in *ligaments*, *tendons*, *fibrous membranes*, *aponeuroses*, &c. The *yellow*, (F.) *Tissu jaune*, exists separately in the middle coat of the arteries, the *chorde vocales*, *ligamentum nuchæ* of quadrupeds, &c. It differs from the *white* in possessing a high degree of elasticity, owing to the presence of a

distinct principle, called by MM. Robin and Ver-deil *Elasticine*.

FIBROUS GROWTH, Tumour, fibrous—*f.* Matter of the Brain, see *Cerebrum*—*f.* Membranes, see *Membranes*, fibrous.

FIB'ULA, (for *Fibigula* [?], from *figo*, 'to fix,' 'a clasp, or brace,') *Cru'ris ra'dius*, *Can'tua mi'nor*, *Os per'onæ*, *Per'onæ*, *Perone'um*, *Foc'ilè mi'nus*, *Su'ra*, *Ar'u'do mi'nor*, *Fist'ula Cru'ris*, *Tib'ia mi'nima*, *Os tib'ie mi'nus*, *Spellbone*, The splinter bone of the leg. (F.) *Péroné*. The long, small bone, situate at the outer part of the leg. The superior or tibial extremity of the fibula is rounded and forms the *caput* or head. It is articulated with the tibia. Its inferior or tarsal extremity is broader than the superior. It is articulated with the tibia and astragalus, and forms the *malleolus externus* or outer ankle by means of its *coronoid process*. The body of the bone has three faces, having more or less prominent edges. It is separated from the tibia by the interosseous space, and is developed by three points of ossification; one at the body, and one at each extremity. It prevents the foot from turning outwards.

FIBULAD, see *Fibular aspect*.

FIBULAR, *Peroneal*.

FIB'ULAR AS'PECT. An aspect towards the side on which the fibula is situated. *Fib'ulad* is used by the same writer adverbially, to signify 'towards the fibular aspect.'

FIBULATIO, (*fibula*, 'a clasp,') *Infibulatio*.

FIC, *Ficus*.

FICARE, *Ranunculus ficaria*.

FICARIA, (from *ficus*, 'a fig,') *Scrophularia aquatica*—*f.* *Communis*, *Ranunculus ficaria*—*f.* *Ranunculoides*, *Ranunculus ficaria*—*f.* *Verna*, *Ranunculus ficaria*.

FICATIO, *Ficus*.

FICICION, *Physician*.

FIC'OID, *Ficoi'des*, *Ficoïdens*, (*ficus*, 'a fig,' and *eidos*, 'resemblance,') Resembling a fig in form or character.

FICOSA EMINENTIA, *Ficus*.

FICOSIS, (*ficus*, and the suffix *osis*), *Ficus*.

FIC'OUS, *Fico'us*. Relating or appertaining to *ficus*.

FICUS, *Sy'cè*, *Sy'cea*, *Sy'cum*, *Syco'sis*, *Syco'ma*, *Fico'sis*, *Fica'tio*, *Fico'sus* *Tu'mor*, *Fico'sa eminent'ia*, *Maris'ca*, (F.) *Fic*. A fleshy excrescence, often soft and reddish, sometimes hard and scirrhus, hanging by a peduncle, or formed like a fig, occurring on the eyelids, chin, tongue, anus (*Ischias*), or organs of generation. The *fici* seated on the last-mentioned parts are generally of a syphilitic character.

FICUS, *F. Carica*.

FICUS CAR'ICA, *F. vulga'ris* seu *commu'nis*. The systematic name of the fig-tree, (F.) *Fiquier*. *Ord.* *Urticaceæ*. *Car'ica*, *F'icus*, *Sy'cè*, (F.) *Figue*, the fig, is a pleasant fruit when ripe, as well as when dried in the state in which it is found in the shops—*Ficus*, (Ph. U. S.) It is used, at times, in place of a cataplasm; especially in gum-boils.

FICUS COMMUNIS, *F. Carica*—*f.* *Indiæ grana*, *Coccos cacti*—*f.* *Indica*, *Musa paradisiaca*, see *Caoutchouc*, *Carica papaya*, and *Lacca*—*f.* *Religiosa*, see *Lacca*.

FIDERIS, *MIN'ERAL WATERS OF*. These waters, in the Canton of the Grisons, are strong, acidulous, and alkaline, and possess all the virtues of the class.

FID'GETS. (Of doubtful etymology.) *Dysphor'ia sim'plex*, *D. nervo'sa*, *Erethis'mus sim'plex*, *Tituba'tio*, *Creep'ers*, (Sc.) *Feyk*, (F.) *Frétillement*. General restlessness and troublesome uneasiness of the nerves and muscles, with increased sensibility, and inability of fixing the attention, accompanied

with a perpetual desire for changing the position. See *Agacement des Nerfs*.

FIDICINALES, (from *fidicen*, 'a harper,')

Lumbricales manus.

FIE, *Psychorages*.

FIEL, ([L.] *fel*,) *Bile*—*f. de Bauf*, *Bile of the ox*—*f. de Terre*, *Fumaria*.

FIELDWEED, *Anthemis cotula*, *Erigeron Philadelphicum*.

FIENTE, *Fimus*.

FIÈVRE, (from *febris*,) *Fever*—*f. d'Accès*,

Intermittent fever—*f. Adynamique*, *Typhus*—*f.*

Algide, see *Algidus*—*f. d'Amour*, *Fever, erotic*—*f.*

Angeioténique, *Synocha*—*f. Annuelle*, see *Annual Diseases*—*f. Ardent*, *Ardent fever*, *Synocha*—*f.*

Ataxique, *Typhus*—*f. de la Barbade*, *Fever, yellow*—*f. Bulleuse*, *Pemphigus*—*f. des Camps*,

Typhus gravior—*f. Catarrhale*, *Catarrh*—*f. Ca-*

tarrhale épidémique, *Influenza*—*f. Cérébrale*, *Cerebral fever*—*f. Cérébrale des Enfants*, *Hydroce-*

phalus internus—*f. Cérébro-spinale*, *Meningitis, cerebro-spinal*—*f. Cholérique*, *Choleric fever*—

f. de Consomption, see *Consumption*—*f. Contine-*

nente inflammatoire, *Synocha*—*f. Délirante*,

Fever, delirious—*f. Double*, *Fever, double*—*f.*

Doublee, *Fever, doublé*—*f. Entéro-mésentérique*,

Typhoid fever—*f. Éphémère*, *Ephemera*—*f.*

Étique, *Hectic fever*—*f. Gastro-adynamique*,

Fever, yellow—*f. Gastro-hépatique*, *Fever, yellow*

—*f. Hectique*, *Hectic fever*—*f. de Hongrie*, *Fever, Hungarian*—*f. d'Hôpital*, *Typhus gravior*—*f. Ie-*

térique, *Fever, yellow*—*f. Inflammatoire*, *Synocha*

—*f. Intermittente*, *Intermittent fever*—*f. Inter-*

mittente paludéenne, see *Elodes*—*f. Irritative*,

Synocha—*f. Jaune d'Amérique*, *Fever, yellow*—

f. Lactée, *Fever, milk*—*f. de Lait*, *Fever, milk*—

f. Laitéuse, *Fever, milk*—*f. Larvée*, *Fever, masked*—*f. Lenticulaire*, *Typhus gravior*—*f. du*

Levant, *Plague*—*f. des Marais*, *Intermittent*

fever—*f. Maremmatique*, see *Elodes*—*f. Matelote*,

Fever, yellow—*f. Méningo-gastrique*, *Fever gastro-*

trique—*f. Mésentérique*, see *Typhus*—*f. de la Mois-*

son, *Fever, gastric*—*f. Morbilleuse*, *Rubeola*—*f.*

Nerveuse, *Typhus mitior*—*f. Nosocomiale*, *Typhus*

gravior—*f. Ortée*, *Urticaria*—*f. Oscitante*, *Oscit-*

ant fever—*f. Paludéenne*, see *Elodes*—*f. Pério-*

dique, *Intermittent fever*—*f. Péripleurmonique*,

Pneumonia—*f. Perniciense délirante*, see *Delir-*

ious—*f. Pleurétique*, *Pleurisy*—*f. Pneumonique*,

Pneumonia—*f. Pourprée*, *Scarlatina*—*f. des Pri-*

sons, *Typhus gravior*—*f. Puerpérale*, *Fever, puer-*

peral, see *Peritonitis*—*f. Quarte*, *Quartan*—*f.*

Quinque, *Fever, quinic*—*f. à Quinquina*, see

Fever, quinic—*f. de Quinquina*, *Fever, quinic*—*f.*

Quotidienne, *Quotidian*—*f. à Rechute*, see *Re-*

lapse—*f. Rémittente*, *Remittent fever*—*f. Rhuma-*

tismale, *Rheumatism, acute*—*f. Rouge*, *Roseolæ*,

Scarlatina—*f. Sanguine*, *Synocha*—*f. Semitierce*,

Hemitritæa—*f. de Siam*, *Fever, yellow*—*f.*

Suante, *Sudor Picardicus*—*f. Syncopeale*, see

Fever syncopeal, and *Syncopeal*—*f. Tierce*, *Tertian*

fever—*f. Typhoïde*, see *Typhus*—*f. Vésiculaire*,

Pemphigus.

FIÈVREUX, *Feverish*.

FIG, ([L.] *ficus*, *Sax. pic.*) See *Ficus carica*—

f. Indian, *Cactus opuntia*.

FIGUE, see *Ficus carica*.

FIGUIER, *Ficus carica*—*f. d'Inde*, *Cactus*

opuntia.

FIGURA VENOSA, ('venous figure,') *Circu-*

lus venosus.

FIGURATIO, (*figurare*, *figuratum*, 'to fash-

ion,') Imagination.

FIGURE, *Countenance*.

FIGURÉ, (F.) An epithet for a compressive

bandage, applied over the head after bleeding

from the frontal vein. It has also been called

bandage royal.

FIGWORT, *Scrophularia nodosa*—f. Water, greater, *Scrophularia aquatica*.

FILA NERVEA, Nerve-fibres—f. Spermatica, Spermatozoa.

FILACEOUS, Filamentous.

FILAIRE DE MEDINE (*Filaria*), Dracunculus.

FILAMENT, *Filum*, *Filamentum*, (dim. of *filum*, 'a thread.') This word is used synonymously with *filibril*; thus, we say a *nervous* or *cellular filament* or *filibril*. Also, the glairy, thread-like substance which forms in the urine in some diseases, and which depends on a particular secretion from the mucous membrane of the urinary passages.

FILAMENT OF MÜLLER, see Tunica Jacobi.

FILAMEN'TOUS, *Filamento'sus*. Filaceous; thread-like; filiform. Containing thread-like substances, as the *tunica filamentosa* or decidua:—*Filamentous urine*; urine containing thread-like substances.

FILAMENTUM, Filament, Frænum.

FILARIA GUINEENSIS. (from *filum*, 'a thread.') Dracunculus—f. *Hominis* bronchialis, see Worms—f. *Medinensis*, Dracunculus—f. *Oculi*, see Worms.

FILBERT, *Corylus avellana*.

FILE, DENTAL, (Sax. reol.) Lima dentaria.

FILLELUM, (dim. of *filum*, 'a thread.') Frænum.

FILET, (DE LA LANGUE). (from *filum*, 'a thread,') Frænum—f. *Opération du*, see Frænum—f. *de la Verge*, Frænum penis.

FILETUM, (from *filum*, 'a thread.') Frænum lingue.

FILICULA, (dim. of *filix*, 'fern.') Polypodium filix mas—f. Dulcis. Polypodium vulgare.

FILIFORM, *Filiformis*, (*filum*, 'a thread,' and *forma*, 'form,') Having the shape of a thread; as the "*filiform papillæ* of the tongue;" a "*filiform pulse*." See *Papillæ*.

FILING, Limatio.

FILIPENDULA, (*filum*, 'a thread,' and *pendere*, 'to hang,') *Spiræa filipendula*.

FILIUS ANTE PATREM, 'the son before the father.' A plant whose flower appears before the leaf: as *Tussilago*.

FILIX FEMINEA, (gen. *Filicis*.) *Pteris aquilina*—f. *Florida*, *Osmunda regalis*—f. *Mas*, Polypodium filix mas—f. *Non ramosa dentata*, Polypodium filix mas—f. *Nymphæa*, *Pteris aquilina*—f. *Pinnata*, Polypodium filix mas—f. *Veneris*, *Adiantum pedatum*.

FILLE, (from *filia*, 'a daughter,') Girl—f. *de Joie*, see Prostitution—f. *Publique*, see Prostitution.

FILLET, (from *filum*, 'a thread,') Fascia, Laqueus—f. of the *Corpus Callosum*, *Gyrus fornicatus*.

FILLING, (from Sax. *pyllan*, 'to fill,') A term employed in dentistry for the operation of stuffing the cavity of a carious tooth with some indestructible substance, to prevent the access of extraneous matters. Also called *stopping*, *stuffing*—*Ornithopteroides*. The materials employed for filling are gold foil, sponge or crystalline gold, tin foil, lead, mastic, gutta percha, &c. See *Plugging*.

FILTRATION, *Filtratio*, *Percollatio*, *Ethesis*, *Ethismus*, (from *filtrum*.) A pharmaceutical operation, which consists in passing a fluid through a filter or strainer, for the purpose of clarifying it. In ancient physiology, it meant the action by which the different humours of the body are separated from the mass of the blood.

FILTRUM. A *filter*, (F.) *Filtre*; (from low Latin *feltrum*, (F.) *feutre*, 'felt or fullied cloth,' which was used to form it [?]). Any po-

rous material; such as sand, some kinds of *freestone*, powdered charcoal, pounded glass, flannel, unsized paper, &c., through which a fluid is passed for the purpose of separating it from the matters suspended in it.

FILUM, ('a thread,' akin to *pilus*, 'a hair,') Filament—f. Muscular, Muscular fibre—f. *Tæniaforme*, *Tæniola*.

FILUM TERMINALE. A slender ligament, prolonged from the nervous sheath, formed by the spinal pia mater, which descends through the centre of the cauda equina, and is attached to the dura mater, lining the canal of the coccyx.

FIMBRIA, *Parar'ma*. A band; a fringe; as the *fimbria* or *fimbriated* or fringed extremity of the Fallopian tube.

FIMBRIÆ CARNOSÆ COLI, Appendiculæ pinguedinosæ—f. *Tubarum Fallopii*, see Tube, Fallopian.

FIMBRIATED, see Fimbria.

FIMUS, *Bor'borus*, *On'thus*, *Ster'cus*, (F.) *Fiente*. Dung, excrement.

FIMUS seu STER'CUS AN'SERIS, *Goosedung*, was applied as a poultice to the feet in malignant fever. See *Chenocopus*.

FIMUS EQU'NUS, *Ster'cus e'qui non castrat'i*. *Stonchorse dung* was once thought antileptic.

FIMUS VACCÆ, *Cow dung*, was employed as a cataplasm, especially in gout.

FINCKLE, (Fœniculum.) Anethum.

FINGER, Sax. *pingep*, from *peng*, 'to take,') *Digitus*—f. Ear, see *Digitus*.

FINGER, HIPPOCRATIC, *Dig'it'us Hippocrat'icus*, (F.) *Doigt Hippocratique*. A fusiform tumefaction of the last phalanx of the index finger especially, with curving of the nails, mentioned by Hippocrates as characteristic of phthisis; but present, perhaps, in all cases of profound and protracted disturbance of hæmatisis.

FINGER, MIDDLE, *Medius digitus*—f. Ring, *Annular finger*.

FINGER STALL, *Digitale*.

FINGER, SUPERNUMERARY, see *Monster*.

FINIS ASPERÆ ARTERIÆ, ('end of the trachea,') *Larynx*.

FIÖLE, *Phiala*.

FIPPLE, Lip, under.

FIR BALSAM, *Pinus balsamea*.

FIR MOSS, UPRIGHT, *Lycopodium selago*.

FIR, SCOTCH, *Pinus sylvestris*—f. *Spruce*, *Norway*, *Pinus abies*.

FIR TREE, SILVER, EUROPEAN, *Pinus picea*.

FIR, YEW-LEAVED, *Pinus abies*.

FIRE DAMP, ([G.] *Dampf*, 'vapor,') Hydrogen, carburetted—f. *Persian*, see *Anthrax*—f. *St. Anthony's*, *Erysipelas*.

FIRE WEED, *Erechtithes hieracifolia*, *Senecio*.

FIRING, *Cauterization*.

FIRST INTENTION, see *Intention*.

FISHSKIN, *Ichthyosis*.

FISHTONGUE. (F.) *Langue de Carpe*. An instrument—so called from its shape—used by some dentists for the removal of the dentes sapientie.

FISSICULATIO, (*fissiculare*, *fissiculatum*, 'to cut off, open, make incisions,') An old word for an opening made with a scalpel.

FISSIPARITÉ, see *Generation*.

FISSIPAROUS, (*fissere*, *fissum*, 'to split,' and *parire*, 'to bring forth,') See *Generati'o*.

FISSURA, Fissure—f. *Canalis Fallopii*, see *Aqueductus Fallopii*—f. *Capillare*, *Distrix*—f. *Cerebri longitudinalis*, *Fissure*, longitudinal, of the Brain—f. *Contrajacens*, *Contra-fissura*—f. *Interlobularis*, *Interlobular fissure*.

FISSURA LONGITUDINALIS, ANTERIOR ET POSTERIOR. Two vertical fissures in the median line, in front of and behind the medulla oblongata, which divide it superficially into two symmetrical lateral columns.

FISSURA MAGNA VULVÆ. see Rima—f. Palpebrarum, see Palpebra—f. Pilaris, Trichismus.

FISSURA TRANSVERSA MAGNA CEREBRI, Fissure, transverse, of the Brain.

FISSURE, *Fissu'ra*, *Schis'ma*, *Scissu'ra*, *Rhe'gē*, *Rheg'ma*, *Rheg'mus*, (*findere*, *fissum*, 'to cleave.') A long and narrow cleft or opening in a bone—*Rha'gē*, *Ceas'ma*, (F.) *Fissure*, *Fente*, *Seissure*.

FISSURE has various acceptations. A fracture, *Catag'ma fissu'ra*, in which the bone is cracked, not separated, as in fracture. A narrow, long, and superficial solution of continuity, around the external openings of the mucous membranes. A sort of chap—(F.) *Gerçure*—observed on the hands, particularly on the callous hands of workmen in certain mechanical employments. Small chapped ulcerations, sometimes noticed in young children, owing to the contact of the fæces and urine with the fine delicate skin of the thighs, nates, and genital organs. Clefts of a more or less deep nature, occurring on the genital organs in the vicinity of the anus, in those labouring under syphilis. These are usually called *rhagades*. See Monster.

FISSURE OF BICHAT, Fissure, transverse, of the Brain—f. Capillary, see Pilatio.

FISSURE, CENTRAL. The aggregate of the cavities or ventricles of the brain. Meckel considers them but one cavity, in the form of a cross.

FISSURE OF GLASER, Fissure, glenoid—f. of Glaserius, F. Glenoid.

FISSURE, GLE'NOID, *Fis'sure* of *Gla'ser* or *Gla'ser'ius*, (F.) *Fissure* ou *Scissure* de *Glaser*, *Fente glénoïdale* ou *Fêlure* de *Glaser*, divides the glenoid cavity of the temporal bone into two parts, and gives passage to the chorda tympani, &c.

FISSURE OF THE HE'LI'X. A small vertical fissure of the helix of the ear, a little above the tubercle for the attachment of the *atrahens aurem* muscle.

FISSURE, INFRAORBITAR, Suborbitar fissure.

FISSURE, LONGITU'DINAL, OF THE BRAIN, *Fissu'ra cerebri longitudina'lis*. The space which separates the two hemispheres of the brain.

FISSURE, ORBITAR, see Orbitar fissure—f. Orbitar, inferior, Spheno-maxillary fissure—f. Orbitar, superior, Sphenoidal fissure—f. Portal, see Liver.

FISSURE OF ROLAN'DO. A transverse fissure placed between two superior cerebral convolutions, which are met with above the fissure of Sylvius.

FISSURE, SEMILU'NAR. A notch at the anterior edge of the cerebellum, where it receives fibres which connect it to the cerebellum and mesocephalon.

FISSURE OF SYLV'IVUS, *Fissu'ra* seu *Fos'sa Mag'n'a Syl'vii*. A deep, narrow sulcus, which ascends obliquely backwards from the temporal ala of the sphenoid bone, near to the middle of the parietal bone, and which parts the anterior and middle lobes of the cerebrum on each side.

FISSURE OF THE TRA'GUS. A fissure on the anterior surface of the tragus of the ear.

FISSURE, TRANS'VERSE, OF THE BRAIN, *Great transverse fis'sure*, *Fis'sure* of *Bi'chat*, *Fissu'ra* seu *Ri'ma transver'sa mag'n'a cerebri*. A fissure which passes beneath and behind the edge of the middle lobe of the brain, and extends beneath the hemisphere of one side to the same point of the opposite side.

FISSURES OF THE LIVER, see Liver.

FIST, (Sax. *fýrre*.) *Pygmē*, *Pug'nus*, (F.) *Poing*. The clenched hand.

FIS'TULA, ('a pipe or reed,') *Syr'inx*, *Syrin'ga*, *Aul'us*, *Fis'tule*: when of a small size, *Aulis'eus*. A solution of continuity, of greater or less depth and sinuosity; the opening of which is

narrow, and the disease kept up by an altered texture of parts, so that it is not disposed to heal. A fistula is *incomplete* or *blind*, when it has but one opening; and *complete* when there are two, the one communicating with an internal cavity the other externally. It is lined in its whole course by a membrane which seems analogous to mucous membranes. *Incomplete fistulæ* may be *internal* or *external*. The former are those which open *internally*; the latter those which open *externally*. *External incomplete fistulæ* are kept up by caries or necrosis of bones, by extraneous bodies in any of the living textures, or by purulent cavities, the walls of which have not become united. *Internal incomplete fistulæ* generally soon become complete, since the discharge that escapes from them into the cavities into which they open has a constant tendency to make its way outwardly, and soon occasions ulceration of the integuments. Fistulæ have received different names, according to the discharge which they afford, and the organs in which they are seated,—as *lachrymal*, *biliary*, *salivary*, *synovial*, and *urinary*—*Fis'tula ur'inae*, *U'rias*. The great object of treatment in fistulous sores is to bring on an altered condition of the parietes of the canal, by astringent or stimulating injections, caustics, the knife, pressure, &c. Those which are dependent on diseased bone, cartilage, tendon, &c., do not heal until after the exfoliation of the morbid part. *Fistulæ* of excretory ducts are produced either by an injury of the duct itself or by the retention and accumulation of the fluids to which they have to give passage. Thus, *Fis'tula lachrymal'is*, *Dacryosyr'inx*, *Emphrag'ma lachrymal'e*, *Hy'drops sac'ci lachrymal'is*, *Drop'sy* of the *lachrymal sac*, commonly proceeds from the obliteration of the nasal ducts, or from atony of the lachrymal sac; which circumstances prevent the tears from passing into the nostrils.

FISTULA IN A'NO, *A'nal fistula*, *Archosyr'inx*, generally occurs from some mechanical pressure or impediment. The principal indication in the treatment of these fistulæ of the excretory canals being to put a stop to the constant discharge of the secretions, &c., through the preternatural channel, the fistulous passage is at times laid open, and a communication established with the natural excretory canal; at others, strong pressure is employed to procure its obliteration.

FISTULA BELLINIANA, Uriniferous tube—f. Cibal'is, *Œsophagus*—f. Cruris, *Fibula*—f. Durae matris, Sinus of the dura mater—f. Lacrymalis, see *Fistula*—f. Mammæ, *Mastosyrinx*—f. Nervorum, *Neurilemma*—f. Sacra, *Medulla spinalis*, *Vertebral column*—f. Spiritualis, *Trachea*—f. Ureterum renum, see *Calix*—f. Urinæ, see *Fistula*—f. Urinaria, *Urethra*—f. Ventriculi, *Œsophagus*.

FISTULE, *Fistula*.

FISTULES STERCORAIRES, see *Stercoraceous*.

FISTULOUS, *Fistulo'sus*, *Syring'i'eus*, *Syring'o'des*, (F.) *Fistuleux*. Relating to, or resembling, a fistula; as 'a fistulous opening.'

FIT, Paroxysm, Pes.

FITROOT, *Monotropa uniflora*.

FITS, NINE DAY, *Trismus nascentium*.

FIVE FINGERS, *Panax quinquefolium*.

FIXATIO MONONŒA, Melancholy, see *Mononœa*.

FIX'ED, *Fix'us*, (*figere*, *fixum*, 'to fasten.') A body not capable of being volatilized by fire is said to be fixed. Thus, we say *fixed oils*, in contradistinction to *volatile oils*.

FIXEN, MINERAL WATERS OF. This spring is four leagues from Waldsassen, in Bavaria. It contains carbonic acid, holding in solution carbonates of lime and magnesia; chlorides of lime and magnesia, carbonate of soda and silica. In Bavaria, it replaces the Seltzer water.

FIXI DENTES. The teeth of the second dentition.

FLABELLATION, Flabellatio, (flabellare, flabellatum, (from flabrum, 'a blast of wind,' 'to agitate the air,') An operation recommended by Ambrose Paré, which consists in keeping fractured limbs cool, as well as the dressings surrounding them, by the renewal of the air around them, either by the use of a fan, or the repeated change of position of the parts affected.

FLABELLUM ET VENTILABRUM CORDIS, ('the fan and winnowing machine of the heart,') Pulmo.

FLACCIDITY, Flacciditas, (flaccidus, 'flabby,' 'soft,') Softness of a part, so as to offer little resistance on pressure.

FLAG, BLUE, Iris versicolor—f. Blue, slender, Iris Virginica—f. Dwarf, Iris lacustris—f. Myrtle, Acorus calamus—f. Root, Acorus calamus.

FLAGELLANTS, (from flagellare, 'to scourge,') see Mastigosis.

FLAGELLATIO, (flagellum, dim. of flagrum, 'a scourge,') Mastigosis.

FLAGELLATION MANIA, see Mastigosis.

FLAMBE, Iris Germanica—f. Bâtard, Iris pseudacorus.

FLAMBOISE, (from flammula, 'a small flame' [?]) Ignis sylvaticus.

FLAME, VITAL, Vital principle.

FLAMED, Inflamed.

FLAMMA, ('a flame,' from its shape.) Fleam—f. Cordis, Biolychnium—f. Vitalis, Biolychnium.

FLAMME, Fleam—f. Vitale, Vital principle.

FLAMMETTE, (dim. of flamme,) Fleam.

FLAMMON, Lachesis rhombeata.

FLAMMULA, (dim. of flamma,) Fleam, Ramunculus ficaria—f. Cordis, Biolychnium—f. Jovis, Clematis recta—f. Vitalis, Animal heat, Biolychnium, Vital principle.

FLANC'S, Flanks.

FLANKS, Il'ia, Il'ea, La'gones, Lap'ara, Ceneo'nes, (Prov.) Lists, (F.) Flances, Les Îles. The regions of the body which extend, on the sides, from the inferior margin of the chest to the crista ili.

FLAP, (F.) Lambeau. A portion of the soft parts of the body separated from those beneath, but still attached by the base. Hence there may be 'flap wounds,' (F.) Plaies à lambeaux, and 'flap operations,' (F.) Opérations à lambeaux. When the flap is taken from one part, and applied to another, in plastic operations, the operation is said to be by migration of the flap, (F.) Migration du Lambeau. At times, the flap is made to slide over the part to which it has to adhere:—the operation is then said to be by the sliding of the flap, (F.) Glissement du Lambeau. The flap is also, at times, rolled, inverted, &c., (F.) Roulement ou Inversion du Lambeau.

FLAP DOCK, Digitalis.

FLAP OPERATION OF AMPUTATION, Amputation à lambeaux.

FLAT TOP, Vernonia noveboracensis.

FLATURIUS, (from flare, flatum, 'to blow,') Alchymist.

FLATUEUX, Windy.

FLATULENCE, Flatulentia, Fla'tus, Flatu-ventriculi, Aëripluvium, Pneumatosis, Pneumatosis ventriculi et enterici, Bdes'ma, Hyperpneustia, Cly'don, Phy'sa, Polyphy'sia, Limo'sis Fla'tus, Ereux'ia, Bom'bus, Flatuos'ity, Wind, Windiness, Ventos'ity, Va'pour, (F.) Ventosité, Flatuosité, Vents. Wind emitted from, or accumulated in, the digestive tube especially.

FLATULENT, Windy.

FLATULENTIA, Flatulence.

FLATUOSITÉ, Flatulence.

FLATUOSITY, Flatulence

FLATUS, (from flare, flatum, 'to blow,') Crepitation, Flatulence—f. Furiosus, Ambulo-flatulentus—f. Spinae, Spina ventosa.

FLAVED CORTICUM CITRI, (from flavesc, 'to be yellow,') see Citrus medica.

FLA'VOUR, (F.) flairer, 'to smell,') The quality of a sapid body, which is appreciated by the taste and smell combined, and more especially by the latter. Some physiologists consider that flavour is effected through the smell alone.

FLAVUS, Yellow.

FLAX, COMMON, (Sax. pleax, plex, (G.) Flach.s.) Linum usitatissimum—f. Purging, Linum catharticum.

FLAXSEED, see Linum usitatissimum—f. Seed Meal, see Linum usitatissimum—f. Seed tea, Infusum lini compositum.

FLAX, TOAD, Antirrhinum linaria.

FLEA, COMMON, (Sax. Flea.) Pulex irritans.

FLEABANE, CANADA, Erigeron Canadense—f. Great, Conyza squarrosa—f. Narrow-leaved, Erigeron Philadelphicum—f. Philadelphia, Erigeron Philadelphicum—f. Skevish, Erigeron Philadelphicum—f. Various-leaved, Erigeron heterophyllum.

FLEAM, Flam'ma, Flam'mula, ('a flame,' from its shape [?]) Schaste'rium, Fosa'rium, Phlebotomum. (F.) Flamme, Flammette. A surgical instrument used for the operation of phlebotomy. It consists of a small metallic box, containing a spear-pointed cutting instrument, which, by means of a spring, can be forced into the vein. It is much used in Germany and some other European countries, and is not unfrequently employed in America; but is scarcely ever seen in France or Great Britain.

FLEAWORT, Plantago psyllium.

FLÉCHISSEUR, (from flectere, 'to bind,')—f. Court commun des orteils, Flexor brevis digitorum pedis—f. Court du gros orteil, Flexor brevis pollicis pedis—f. Court du petit doigt, Flexor parvus minimi digiti—f. Court du petit orteil, Flexor brevis minimi digiti pedis—f. Court du ponce, Flexor brevis pollicis manus—f. Grand commun des orteils, Flexor longus digitorum pedis profundus perforans—f. Long commun des orteils, Flexor longus digitorum pedis profundus perforans—f. Long du gros orteil, Flexor longus pollicis pedis—f. Profond des doigts, Flexor profundus perforans—f. Sublime des doigts, Flexor sublimis perforatus—f. Superficiel des doigts, Flexor sublimis perforatus.

FLECK, ((G.) Fleck, 'a spot,') Abrasion.

FLECTENS PAR LUMBORUM, (flectere, 'to bend,') Quadratus lumborum.

FLEGMEN, (from phlegmone [?]) Flemen.

FLEMEN, gen. FLEM'ina, Fleg'men. A tumour about the ankles. Also, a chap on the feet and hands.

FLERECIN, Gout.

FLESH, (Flærc, flerc,) Cre'aa, Caro, Sarr, (F.) Chair. Every soft part of an animal is so named; but more particularly the muscles, which are called muscular flesh.

FLESH, PROUD, Fungosis.

FLESHY, Corpulent.

FLETUS (flere, fletum, 'to weep,') Lacrymatio.

FLEUR DE COUCOU, (from flos, gen. floris, 'a flower,' 'cuckoo flower,' Primula veris—f. Sainte Catherine, Nigella—f. de Sainte Jacques, Thlaspi bursa—f. de Tous les mois, Calendula officinalis.

FLEURS, (pl. of fleur.) Flowers. Menses—f. de Benjoin, Benjamin—f. Blanches, Leucorrhœa—f. de Muscade, Mace—f. de Soufre, Sulphur sublimatum.

FLEXIBILITY, Flexibil'itas, (flectere, flexum, 'to bend,') Capability of being bent. A physical property of the tissues, which varies greatly ac-

cording to the structure. The tendons exhibit this property in a marked manner.

FLEXIO, Campsis, Flexion.

FLEX'ION, *Flex'io, Cam'pē*. The state of being bent. The action of a flexor muscle.

FLEX'OR, 'a bender,' (F.) *Fléchisseur*. A muscle, whose office it is to bend certain parts.

FLEXOR ACCESSORIUS, see Flexor longus digitorum pedis.

FLEXOR BRE'VIS DIGITO'RUM PE'DIS PERFORA'TUS, *F. Sublimis seu bre'vis seu digito'rum bre'vis seu perfora'tus pe'dis, Perfora'tus seu flex'or secundi interno'dii digito'rum pe'dis, (F.) Calcaneo-sous-phalangettien commun, Calcaneo-sous-phalangien commun*,—(Ch.), *Muscle court fléchisseur commun des orteils*. A muscle, placed at the middle of the sole of the foot. It is narrower and thicker behind than before, where it is divided into four portions. It arises from the posterior part of the inferior surface of the os calcis, and is inserted at the inferior surface of the second phalanx of the last four toes. It bends the second phalanges of the toes on the first, and the first on the metatarsal bones; in this manner augmenting the concavity of the vault of the foot.

FLEXOR BREVIS MINIMI DIGITI, *F. Parvus minimi digiti*.

FLEXOR BRE'VIS MIN'IMI DIG'ITI PE'DIS, *Parv'us min'imus, (F.) Court fléchisseur du petit orteil, Tarsosous-phalangien du petit orteil*—(Ch.) A muscle, situate at the anterior and outer part of the sole of the foot. It arises from the posterior extremity of the fifth metatarsal bone, and is inserted into the posterior part of the first phalanx of the little toe, which it bends.

FLEXOR BRE'VIS POL'LICIS MA'NUS, *F. secundus interno'dii, F. pri'mi et secund'i ossis pol'licis, The'nar, (F.) Court fléchisseur du ponce, Carpo-phalangien du ponce*—(Ch.), *Demi-interosseus du ponce*. A muscle, situate at the outer part of the palm of the hand. It is divided into two portions by the tendon of the *Flexor longus pol'licis*. It arises from the os magnum, the anterior annular ligament of the carpus and the third metacarpal bone; and is inserted into the superior part of the first phalanx of the thumb, and into the two ossa sesamoida at the articulation of the first phalanx with the first metacarpal bone. Its use is to bend the first phalanx of the thumb on the first metacarpal bone, and the latter upon the trapezium.

FLEXOR BRE'VIS POL'LICIS PE'DIS, *F. brevis seu hallucis seu bre'vis pol'licis, (F.) Tarsos-phalangien du ponce, Court fléchisseur du gros orteil, Tarsosous-phalangettien du premier orteil*—(Ch.) It is situate at the anterior and inner part of the sole of the foot; is thin and narrow behind; thick and divided into two portions before. It arises from the inferior part of the os calcis and the last two cuneiform bones, and is inserted at the inferior part of the base of the first phalanx of the great toe, and into the two sesamoid bones of the corresponding metatarso-phalangian articulation. It bends the first phalanx of the great toe on the first metacarpal bone.

FLEXOR CARPI RADIALIS, *Palmaris magnus*—f. Carpi ulnaris, see Cubital (muscles)—f. Hallucis, *F. brevis pollicis pedis*—f. Hallucis longus, *F. longus pollicis pedis*.

FLEXOR LONGUS DIGITO'RUM PE'DIS PROFUNDUS PERFORANS, *Perforans seu Flex'or profundus, Perodactyl'us, Peronodactyl'us, Peronodactyl'us, Peronodactyl'us, Flex'or digito'rum longus seu Perforans pe'dis, Perforans seu Flex'or tertii interno'dii digito'rum pe'dis; (F.) Tibio-phalangettien*—(Ch.), *Grand ou long fléchisseur commun des orteils*. A muscle, situate at the posterior and deep-seated part of the leg. It is broader at its middle than at its extremities, the inferior of which is divided into four portions. It arises

from the posterior surface of the tibia, and its tendons are attached to the posterior part of the lower surface of the three phalanges of the last four toes. It bends the three phalanges on each other, and the toes on the metatarsus, and extends the foot on the leg.

The *Accessorius Flexoris Longi Digito'rum Pedis, Flex'or accessorius, Ca'ro quadrata Syl'vii, C. accessorius, Massa car'nea Jaco'bi Syl'vii, Planta'ris ve'rus, (F.) Accessoire du long fléchisseur commun des orteils, Carré (Chair)*, is a small muscle of the sole of the foot, which passes obliquely from the os calcis to the outer edge of the flexor longus, whose force it augments, and obliquity it corrects.

FLEXOR LONGUS POL'LICIS MANUS, *F. longus pol'licis, F. tertii interno'dii seu longissimus pol'licis; (F.) Radio-phalangettien du ponce*,—(Ch.) Situate at the anterior and profound part of the forearm. It arises from the upper three quarters of the anterior surface of the radius and interosseous ligament, and is inserted, by a tendon, into the anterior surface of the last phalanx of the thumb. It bends the second phalanx of the thumb on the first; the first on the corresponding metacarpal bone, and this upon the radius. It can, also, bend the hand on the forearm.

FLEXOR LONGUS POL'LICIS PE'DIS, *F. Hallucis seu Pol'licis longus, (F.) Péronéo-phalangien du gros orteil, Long fléchisseur du gros orteil, Péronéo-sous-phalangettien du ponce*,—(Ch.) It is situate at the posterior and profound part of the leg. It arises from the posterior surface of the fibula and the interosseous ligament, and is inserted, by means of a long tendon, into the inferior part of the first phalanx of the great toe. It bends the third phalanx on the first, and this upon the corresponding metatarsal bone. It augments the concavity of the sole of the foot, and extends the foot on the leg.

FLEXOR OSSIS METACARPI POLLICIS, *Opponens pollicis*.

FLEXOR PAR'VUS MIN'IMI DIG'ITI, *F. brevis minimi dig'iti ma'nus, Abductor minimi dig'iti, Hypoth'enar Riola'ni seu minimi dig'iti, (F.) Carpo-phalangien du petit doigt*—(Ch.), *Court fléchisseur du petit doigt*. It arises from the anterior annular ligament of the carpus and the process of the os unciniforme, and is inserted at the inner side of the superior extremity of the first phalanx of the little finger. It bends the first phalanx of the little finger.

FLEXOR PERFORANS, *F. profundus perforans*—f. Perforatus, *F. sublimis perforatus*—f. Perforatus pedis, *F. brevis digitorum pedis*—f. Primi internodii, *Opponens pollicis*—f. Primi internodii digitorum manus, *Lumbricalis manus*—f. Primi et secundi ossis pollicis, *F. brevis pollicis manus*.

FLEXOR PROFUNDUS PERFORANS, *F. Profundus seu Perforans seu tertii interno'dii digitorum manus, Perforans ma'nus; (F.) Cubito-phalangettien commun*—(Ch.) *Fléchisseur profond des doigts*. A thick, flat, long muscle, seated beneath the *Flexor sublimis perforatus*. Its upper extremity is simple, and arises from the anterior surface of the ulna and from the interosseous ligament. Its inferior extremity terminates by four tendons, which, after having passed through slits in the *sublimis*, are inserted into the anterior surface of the last phalanges of the four fingers. It bends the third phalanges on the second, and, in other respects, has the same use as the flexor sublimis perforatus.

FLEXOR SECUNDI INTERNODII DIGITORUM PEDIS, *F. brevis digitorum pedis*—f. Tertii internodii, *F. longus pollicis manus*—f. Tertii internodii digitorum manus, *F. profundus perforans*—f. Tertii internodii digitorum pedis, *F. longus digitorum pedis profundus perforans*—f. Sublimis, *F. brevis digitorum pedis*.

FLEXOR SUBILI-MIS PERFORATUS, *F. Perforatus*, (F.), *Fléchisseur sublime ou superficiel des doigts*, *Épitrôchlo-phalangien commun*, — (Ch.) It is a thick, flat, muscle, seated at the anterior part of the forearm. Its upper extremity, which is simple, arises from the internal condyle of the os humeri;—from the coronoid process of the ulna, and from the anterior edge of the radius. Its lower extremity divides into four tendons, which slide under the anterior annular ligament of the carpus, and are inserted into the second phalanges of the last four fingers, after having been slit to allow the tendons of the flexor profundus to pass through them. This muscle bends the second phalanges on the first; these on the carpal bones, and the hand on the forearm.

FLEXURA, Curvature.—*f. Iliaca*, Sigmoid flexure.—*f. Sigmoidea*, Sigmoid flexure.

FLICK, Epiploon.

FLIXWEED, *Sisymbrium sophia*.

FLOATING, (Sax. *pleotan*, 'to float;—itself from plopan, 'to flow,') *Hæmorrhagia*.

FLOCCI, (*flocus*, 'a flock of wool,') see Villous membranes.—*f. Volitantes*, see *Metamorphopsia*.

FLOCCILATION, (*flocus*, 'a flock of wool,' and *legere*, 'to collect,') *Carphologia*.

FLOCCILEGIUM, *Carphologia*.

FLOCCITATION, *Carphologia*.

FLOCCORUM VENATIO, (*venor*, *venatus*, 'to hunt,') *Carphologia*.

FLOCCULI, pl. of *Flocculus*, see Villous membranes.

FLOCCULUS; (dim. of *flocus*, 'a flock of wool,') *Pneumogastria lobule*, *Lobulus pneumogastrius*. A long and slender prominence, extending from the side of the vallicula around the corpus testiforme to the crus cerebelli, lying behind the filaments of the pneumogastric nerves.

FLOODING, (from Sax. *plœb*, from *plopan*, 'to flow,') *Metrorrhagia*.

FLOR DE MISTELA, see *Mistura*.

FLOR'ENCE, CLIMATE OF. This agreeable Italian city is by no means a favourable residence for the phthisical invalid. Sir James Clark affirms, indeed, that he does not know any class of invalids for whom Florence offers a favourable residence. It is subject to sudden vicissitudes of temperature, and to cold, piercing winds during the winter and spring.

FLORES BENZOËS, Benjamin, flowers of—*f. Boracis*, Boracic acid.—*f. Cosso*, see *Hagenia Abyssinica*—*f. Macidis*, see *Myristica moschata*—*f. Macis*, Mace—*f. Martiales*, *Ferrum ammoniatum*—*f. Salis ammoniaci martiales*, *Ferrum ammoniatum*.

FLORIDA, CLIMATE OF, see Saint Augustine.

FLORIDA, WATERS OF. Near Long Lake, in Florida, United States, which communicates with St. John's River by a small creek, there is a vast fountain of hot mineral water, issuing from a bank of the river. From its odour it would seem to be sulphureous.

FLORION, *Influenza*.

FLOS, gen. *Floris*, *An'thos*. A flower. Also, the finest and noblest part; and virginity.

FLOS JOVIS, *Croceus*—*f. Lactis*, *Cremor lactis*—*f. Salis*, Soda, subcarbonate of—*f. Sanguineus monardi*, *Tropeolum majus*—*f. Trinitatis*, *Viola tricolor*—*f. Virginitatis*, *Hymen*.

FLOUR, COLD, (from *flor*,) *Pinoli*.

FLOUR, PATENT, JONES'S. A farinaceous preparation, which is said to consist of wheat-flour, with tartaric acid and carbonate of soda.

FLOUR, POTATO, see *Solanum tuberosum*.

FLOW, (Sax. *plopan*, 'to flow,') *Flux*.

FLOWER DE LUCE, *Iris Germanica*.

FLOWERS, *Floris*, (F.) *Flours*. The ancient chymists gave this name to different solid and volatile substances obtained by sublimation. The

term is not yet entirely banished from chymical and medical language, as *Flowers of Benjamin*, *Flowers of Sulphur*, &c.

FLOWERS, Menses.

FLOWERS, FOUR CARMINATIVE, *Quatuor flores carminativi*, were chamomile, dill, feverfew, and melilot.

FLOWERS, FOUR COR'DIAL, *Quatuor flores cordiales*, were formerly, borage, bugloss, roses, and violets.

FLUCTUATIO, (*fluctuare*, *fluctuatum*,) (*fluctus*, 'a wave,') 'to undulate,') *Fluctuation*—*f. Aurium*, *Tinnitus aurium*.

FLUCTUATION, *Fluctuatio*, *Undulatio*, (F.) *Undulation*. The undulation of a fluid collected in any natural or artificial cavity, which is felt by pressure or by percussion, properly practised—*peripheric fluctuation*. In ascites, the fluctuation is felt by one of the hands being applied to one side of the abdomen whilst the other side is struck with the other hand. In abscesses, fluctuation is perceived by pressing on the tumour, with one or two fingers alternately, on opposite points.

FLUCTUATION, PERIPHERIC, see *Fluctuation*—*f. Rhonchal*, see *Rhonchal*—*f. by Succussion*, see *Succussion*.

FLUELLEN, *Antirrhinum elatine*.

FLUELLIN, *Veronica*.

FLUEURS, (from *fluere*, 'to flow,') *Menses*—*f. Blanchæ*, *Leucorrhæa*.

FLUID, *Fluidus*, (*fluere*, 'to flow,') A body whose particles move easily on each other; but have not sufficient cohesion to remain united when subjected to pressure. The human body is largely composed of fluids.

The mean results of experiments by M. Chevreul, which consisted in drying the human body in *vacuo*, with a very moderate degree of heat, gave the proportion of water about 6.667 in the 10.000.

FLUID, BURNETT'S DISINFECTING, see *Burnett*—*f. Cephalorachidian*, *Cephalospinal fluid*—*f. Cephalospinal*, *Cephalospinal fluid*—*f. Cerebrospinal*, *Cephalospinal fluid*—*f. Condy's disinfecting*, see *Condy's disinfecting fluid*—*f. Deodorizing*, *Ellerman's*, see *Ellerman*—*f. Ledoyen's disinfecting*, see *Ledoyen*—*f. of Scarpa*, *Vitrine auditee*—*f. Subarachnoidean*, *Cephalospinal fluid*.

FLUIDE SÉMINAL, *Sperm*.

FLUIDUM CEREBRO-SPINALE, *Cephalospinal fluid*—*f. Nerveum*, *Nervous fluid*.

FLUKE, (from Sax. *plœc*, 'the fluke fish,') *Distoma hepaticum*—*f. Liver*, *Distoma hepaticum*.

FLUMEN DYSENTERICUM, (*flumen*, 'a stream,' from *fluere*, 'to flow,') *Dysentery*.

FLUM'MERY, (corrupted from *frumenty* [?]), (Scotch) *Sov'ens*. A preparation of oatmeal, which forms a light article of food during convalescence. It may be made as follows:—Take of oatmeal or groats, a quart. Rub with two quarts of hot water, and let the mixture stand until it becomes sour; then add another quart of hot water, and strain through a hair sieve. Let it stand till a white sediment is deposited: decant, and wash the sediment with cold water. Boil this with fresh water till it forms a mucilage, stirring the whole time.

FLUOR, (from *fluere*, 'to flow,') *Flux*—*f. Albus intestinorum*, *Celiac flux*—*f. Albus malignus*, *Gonorrhœa impura*—*f. Muliebris*, *Leucorrhœa*—*f. Muliebris non Gallicus*, *Leucorrhœa*—*f. Sanguinis pulmonum*, *Hæmoptysis*—*f. Sanguinis vesicæ*, *Cystorrhagia*.

FLUSH, *Flush'ing*, (F.) *Rougeurs*, *Bouffés de Chaleur*; ([G.] *fliessen*, 'to flow,') The temporary redness and heat produced by accumulation of blood in the capillaries of the face; as the

and len 'flush' or 'flush of emotion': the 'flush of hectic.'

FLUX, *Fluxus*, *Profluxum*, *Fluxio*, *Flux*, (*fluere*, *fluxum*, 'to flow'.) A discharge. *Rhyxis*. In nosology, it comprises a series of affections, the principal symptom of which is the discharge of fluid. Generally it is employed for dysentery.

FLUX, *Bilious*, *Fluxus biliosus*. A discharge of bile, either by vomiting or by stool, or by both, as in cholera.

FLUX, *Bleedy*, Dysentery — *f. de Bouche*, Salivation — *f. Bronchique*, Bronchorrhœa — *f. Dysentérique*, Dysentery — *f. Hemorrhoidal*, see Hemorrhoids — *f. Hépatique*, Hepatirrhœa — *f. Menstrual*, Menses — *f. Muqueux*, Catarrh — *f. Muqueux de l'estomac*, Gastrorrhœa — *f. Muqueux de la vessie*, Cystirrhœa — *f. Root*, Aselepias tuberosa — *f. Salivaire*, Salivation — *f. de Sang*, Hemorrhagia, Dysentery — *f. Sebaceus*, Stearrrhœa — *f. de Sperme*, Spermatorrhœa — *f. de Sueur*, Ephaidrosis — *f. d'Urine*, Diabetes — *f. de Ventre*, Diarrhœa — *f. Weed*, Sysimbrium sophia.

FLUXIO, (*fluere*, *fluxum*, 'to flow'.) Fluxion — *f. Alba*, Leucorrhœa — *f. Arthrica*, Gout — *f. Vulvæ*, Leucorrhœa.

FLUXION, *Fluxio*, *Affluxus*. A flow of blood or other humour towards any organ with greater force than natural. A *determination*. Thus we say, in those disposed to apoplexy, there is a *fluxion* or *determination* of blood to the head.

FLUXION CATARRHALE, Catarrh — *f. sur les Dents*, Odontalgia.

FLUXION DE POITRINE, (F.) By this name, the French often understand *acute pulmonary catarrh*, and *pleurisy*, but most commonly *peripneumony*.

FLUXUS, gen. *Fluxus*, Discharge — *f. Alvinus*, Diarrhœa — *f. Capillorum*, Alopecia — *f. Chylosus*, Coeliac flux — *f. Coeliacus*, Coeliac flux — *f. Coeliacus per renes*, Chyluria — *f. Cruentus cum tenesmo*, Dysentery — *f. Dysentericus*, Dysentery — *f. Hepaticus*, Hepatirrhœa, see Hepateros — *f. Lientericus*, Lientery — *f. Lunaris*, Menses — *f. Matricis*, Leucorrhœa — *f. Menstrualis*, Menstruation — *f. Menstruus*, Menses — *f. Muliebris*, Leucorrhœa — *f. Salivæ*, Salivation — *f. Sebaceus*, Stearrrhœa — *f. Splenicus*, Melæna — *f. Venereus*, Gonorrhœa imputra — *f. Ventriculi*, Gastrorrhœa.

FLYGOLDING, *Coccinella septempunctata*.

FLYPOISON, *Amianthium muscetoxicum*.

FLYTRAP, *Apocynum androsaemifolium*, *Sarcocœnia*.

FOALFOOT, *Tussilago*.

FOCILLÆ. The name was formerly given to the bones of the leg, as well as to those of the forearm.

FOCILE INFERIUS SEU MAJUS, Ulna — *f. Majus*, Tibia — *f. Minus*, Fibula — *f. Minus seu superius*, Radius.

FOCUS APOPLECTICUS, Apoplectic cell.

FÆCUNDATIO, Fecundation.

FÆCUNDITAS, Fecundity.

FÆCUNDUS, Fecund.

FÆDI COLORES, ('filthy colors,') Chlorosis.

FÆMEN, Femen, Perineum.

FÆMINA, Female, Woman.

FÆMINESCENTIA, Feminescence.

FÆNICULUM, (dim. of *fœnum*, 'hay,') Anethum — *f. Aquaticum*, Phellandrium aquaticum — *f. Erraticum*, Peucedanum silaus — *f. Marinum*, Crithmum maritimum — *f. Officinale*, Anethum — *f. Porcinum*, Peucedanum — *f. Vulgare*, Anethum.

FÆNUGREEK, *Trigonella fœnum*.

FÆNUM CAMELORUM, ('hay of camels,')

Juncus odoratus — *f. Græcum*, *Trigonella fœnum*.

FÆTABULUM, (*fæteo*, 'to smell badly,') An encysted abscess.

FÆTAL, *Fœtalis*. Relating to the fœtus. A

name given to the parts connected with the fœtus. Thus we say — the *fœtal surface* of the placenta, in contradistinction to the *uterine* or *maternal surface*.

FÆTAL CIRCULATION differs from that of the adult in several respects. Commencing with the placenta, where it probably undergoes a change analogous to what occurs in the lungs in extra-uterine existence, the blood proceeds by the umbilical vein as far as the liver, where a part of it is poured into the vena porta; the other proceeds into the vena cava inferior; the latter, having received the suprahepatic veins, pours its blood into the right auricle. From the right auricle, a part of the blood is sent into the right ventricle; the rest passes directly through into the left auricle, by the foramen ovale. When the right ventricle contracts, the blood is sent into the pulmonary artery; but as the function of respiration is not going on, little blood passes to the lungs; the rest goes through the ductus arteriosus into the aorta. The blood, received by the left auricle from the lungs, as well as that which passed through the foramen ovale, is transmitted into the left ventricle, by the contraction of which it is sent into the aorta, and by means of the umbilical arteries, which arise from the hypogastric, it is returned to the placenta.

FÆTAL HEAD. The diameters of this at the full period are as follows:—1. The *Biparietal* or *transverse*, extending from one parietal protuberance to the other, and measuring $3\frac{1}{2}$ inches. 2. The *Temporal*, from one temple to another, 3 inches. 3. The *Occipito-mental*, from the occiput to the chin; the greatest of all, 5 inches. 4. The *Occipito-frontal* or *antero-posterior*, $4\frac{1}{2}$ or $4\frac{1}{4}$ inches. 5. The *Cervico-bregmatic*, from the nape of the neck to the centre of the anterior fontanelle. 6. The *Fronto-mental*, from the forehead to the chin, about $3\frac{1}{2}$ inches. 7. The *Trachelo-bregmatic*, from the front of the neck to the anterior fontanelle, $3\frac{1}{2}$ inches. 8. The *Vertical diameter*, from the vertex to the base of the cranium.

FÆTAL NUTRITION, Cytotrophy.

FÆTATION, Pregnancy.

FÆTICIDE, *Fœticidium*, (*fætus*, and *cadere*, 'to kill;') *Abortivum*. Criminal abortion.

FÆTIDUS, Fetid.

FÆTOR, ('a bad smell,') Dysodia.

FÆTOR ALARUM. The disagreeable rank odour exhaled from the axillæ.

FÆTOR ORIS, Breath, offensive.

FÆTUS, *Fœtus*, *Cyēma*, *Ovulus ventris*, *Sarcocœnia*. The young of any creature. The unborn child. (F.) *Fœtus*, *Faix*, *Fruit*. By *œmpta*, *Cyēma*, Hippocrates meant the fecundated, but still imperfect, germ. It corresponded with the term *embryo*, as now used; whilst *εμβρυον*, 'embryo,' signified the fœtus at a more advanced stage of utero-gestation. The majority of anatomists apply to the germ the name *embryo*, which it retains until the third month of gestation, and with some until the period of quickening; whilst *fœtus* is applied to it in its latter stages. The terms are, however, often used indiscriminately. When the ovule has been fecundated in the ovarium, it proceeds slowly towards, and enters the uterus, with which it becomes ultimately connected by means of the placenta. When first seen, the fœtus has the form of a gelatinous flake, which some have compared to an ant, a grain of barley, a worm curved upon itself, &c. The fetal increment is very rapid in the first, third, fourth, and sixth months of its formation, and at the end of nine months it has attained its full dimensions—*Enfant à terme*. Generally, there is but one fœtus in utero: sometimes, there are two; rarely three. The fœtus presents considerable difference in its shape, weight, length, situation in the womb, pro-

portion of its various parts to each other, arrangement and texture of its organs, state of its functions at different periods of gestation, &c. All these differences are important in an obstetrical and medico-legal point of view.

FÆTUS, Pregnant—*f.* in Fœtu, Cryptodidymus.

FÆTUS SEPTIMES'TRIS, A seven months' fœtus.

FÆTUS ZEPHYRIUS, Mole.

FOIE, Liver—*f.* d'Antimoine, Oxidum stibii sulphuretum—*f.* de Soufre, Potassæ sulphuretum—*f.* de Soufre calcuire, Calcis sulphuretum.

FOIN DE CHAMEAU, ([L.] *fœnum*, 'hay.')

Juncus odoratus.

FOIROLLE, Mercurialis.

FOLIA APALACHINES, see Ilex Paraguensis—*f.* Peraguæ, Ilex Paraguensis.

FOLIACEUM ORNAMENTUM, (*folium*, 'a leaf.')

See Tuba Fallopiæ.

FOLIE, Heat, Insanity—*f.* Alcoolique, Delirium tremens—*f.* des Ivrognes, Delirium tremens.

FOLLETTE, Atriplex hortensis, Influenza.

FOLLICLE, *Folliculus*, *Cystis*, (dim. of *follius*, 'a bag.')

See Crypta.

FOLLICLES, CILIARY, Meibomius, glands of—*f.* of De Graaf, Folliculi Graafiani—*f.* Hair, see Hair—*f.* of Lieberkühn, see Intestine—*f.* Nail, see Nail—*f.* Palpebral, Meibomius, glands of—*f.* Solitary, Brunner's glands—*f.* Synovial, Bursæ mucosæ.

FOLLICULAR, *Follicularis*, *Folliculose*, *Folliculosus*, (F.) *Folliculeux*. Relating or appertaining to a follicle,—as 'follicular inflammation,' inflammation affecting crypts or follicles, *Folliculitis*.

FOLLICULAR ELEVATIONS, Tubercles, pearly.

FOLLICULE, Crypta—*f.* Céruminæuse, Ceruminous gland—*f.* Ciliaire, Meibomius, gland of—*f.* Dentaire, Dental follicle—*f.* Palpebral, Meibomius, gland of.

FOLLICULES, DE DE GRAAF, Folliculi Graafiani.

FOLLICULI GRAAFIANI, (pl. of *Folliculus*,) *F. Ovarii*, *O'va seu O'vula Graafiana*, *Vesiculæ Graafianæ*, *Follicles* or *Vesicles* of *De Graaf*, *Graafian* or *Ovarian vesicles*, (F.) *Follicules* ou *Vesicules* ou *Œufs de De Graaf*. Small spherical vesicles in the stroma of the ovary, which have at least two coats; the outer termed *ovicap'sule* and *tunica of the capsule*, *tunica fibrosa* and *tunica folliculi*; the latter, *oviscia* and *membrana propria*; *vésicule ovulifère* of M. Pouchet. See *Membrana granulosa*. They exist in the fœtus. The ovum—*ovule* of some—is contained in, and formed by, them. Kölliker terms the follicles themselves, *Ovisacs*, *Ovisac'ci*. The interior of the follicles contains a clear, light-yellowish fluid—*liquor folliculi*—of the density of the serum of the blood, in which are, almost always, isolated granules, nuclei and cells, which appear to be detached portions of the *membrana granulosa*.

FOLLICULI OVARIJ, Folliculi Graafiani—*f.* Rotundi et Oblongi cervicis uteri, Nabothi glandulæ—*f.* Sanguinis, Globules of the blood—*f.* Sebacei, Sebaceous glands—*f.* Sebacei coronæ glandis, see Sebaceous glands.

FOLLICULITIS, (*follicle*, and *itis*.) Follicular inflammation.

FOLLICULOSE GLAND, Crypta.

FOLLICULUS, Follicle, Vulva.

FOLLICULUS A'ERIS, Air-chamber. A space at the larger end of the bird's egg, formed by a separation of the two layers of the shell membrane, which is inservient to the respiration of the young being.

FOLLICULUS DENTIS, Dental Follicle—*f.* Fellis, Gall-bladder—*f.* Genitalis, Scrotum—*f.* Mucosus, see Crypta—*f.* Pili, see Hair.

FOMENT', (*Fove're*, 'to keep warm,') (F.)

Étuver, *Bassiner*. To apply a fomentation to a part.

FOMENTA'TION, *Fomentatio*, *Fœtus*, *Py'ria*, *Thermas'ma*, *Chlias'ma*, *Æone'sis*, *Perf'usio*, *Asper'sio*, *Fomentum* (quasi *fomentum*.) (F.) *Hydrolé*, *Hydrolitif*. A sort of partial bathing, by the application of cloths which have been previously dipped in hot water, or in some medicated decoction. They act, chiefly, by virtue of their warmth and moisture, except in the case of narcotic fomentations, where some additional effect is obtained.

A dry fomentation, *Fomentum sic'cum*, *Lectulus medicatus*, is a warm, dry application to a part;—as a hot brick, wrapped in flannel; a bag, half filled with chamomile flowers made hot, &c.

FOMENTATION HERBS, *Her'bæ pro fœtu*. The herbs, ordinarily sold under this title by the English apothecary, are—*southernwood*, *tops of sea wormwood*, *chamomile flowers*, each two parts; *bay leaves*, one part. ʒiijss of these to Ovj of water.

FOMENTUM, (quasi *Forimentum*.) Fomentation—*f.* Siccum, see Fomentation.

FOMES MOR'BI, gen. *Fo'mitis*, *F. Ma'li*, (*fomes*, 'fuel.')

The focus or seat of any disease.

(F.) *Foyer*.

FOMES VENTRICULI, Hypochondriasis, Spleen.

FOMITES, (pl. of *fo'mes*, gen. *fo'mitis*, 'tinder, fuel, any thing which retains heat.')

Evans'ma, *Zop'ylon*. A term applied to substances which are supposed to retain contagious effluvia; as woollen goods, feathers, cotton, &c.

FUNCTION, Function.

FUNCTIONNEL, Functional.

FOND, Fundus.

FONDAMENTAL, Fundamental.

FONDANT, (*fundere*, 'to fuse.')

Solvent.

FONDAMENT, (from *fond*,) Anus.

FONDLING, Idiot.

FONGIFORME, Fungoid.

FONGOÏDE, Fungoid.

FONGOSITÉ, Fungosity.

FONGUEUX, Fungous.

FONGUS, Fungus—*f.* Médullaire, Encephaloid.

FONS, gen. *fontis*, 'a fount,' Fontanella—*f.* Lacrymarum, see Canthus—*f.* Medicatus, Water, mineral—*f.* Pulsans, Fontanella—*f.* Pulsatilis, Fontanella—*f.* Salutaris, Water, mineral—*f.* Soterius, Water, mineral—*f.* Vitalis, Centrum vitale.

FONSANGE, MIN'ERAL WA'TERS OF. Fonsange is situated near Nîmes in France. The water is sulphuretted.

FONTAINE DE LA TÊTE, ('fountain of the head.')

See Fontanella.

FONTA'NA, CANAL' OF. A canal of a triangular shape, at the inner side of the ciliary circle; partly formed by the groove at the inner edges of the cornea and scleroticæ.

FONTANEL'LA. A *fontanel*; (dim. of *fons*, gen. *fontis*, 'a fountain.')

Fons, *F. pulsatilis* seu *pulsans*, *Ver'tex pal'pitans*, *Bregma*. *Fonticulus*, *Lacu'na*, *Mould*. (F.) *Fontanelle*. The opening of the head. A name given to a space occupied by a cartilaginous membrane, in the fœtus and new-born child, and situate at the union of the angles of the bones of the cranium. There are six fontanels. 1. The great or *sincipital* or *anterior*, *fonticulus quadrangulus*, *Fontaine de la Tête*, situate at the junction of the sagittal and coronal sutures. 2. The small or *bregmatic* or *posterior*, situate at the part where the posterior and superior angles of the parietal bones unite with the upper part of the os occipitis. 3. The two *sphenoidal*, in the temporal fosse; and, 4. The two *mastoid* or of *Casseri*, *Fonticuli mastoïdes* seu *Casseri*, at the union of the parietal, occipital, and temporal bones.

FORTE, (from (L.) *fundere*, 'to fuse.') Colliquation.

FORTE, pl. of *Fons*—f. *Medicati Plumbarii*, Plumbicæ, mineral waters of—f. *Sulphurei calidi*, Waters, mineral, sulphureous.

FORTECULE À POIS, see *Fonticulus*.

FORTECULUS, *Fontanelle*, (dim. of *fons*, gen. *fontis*, 'a fountain.') *Erutarium*, *Edutory*, *Idæus*, (F.) *Fonticulus*, *Cautére*, *Exutoire*. A small ulcer produced by art, either by the aid of caustics or of cutting instruments; the discharge from which is kept up with a view to fulfil certain therapeutic indications. The *Pea issue*, (F.) *Fonticulus à pois*, is kept up by means of a pea placed in it. This *pea*, (F.) *Pois à cautère*, is sometimes formed of wax; at others, the young, blasted fruit of the orange is employed. The common dried garden pea answers the purpose. The seton is also an issue.

FORTECULUS CASSERI, see *Fontanella*—f. *Mastoidæus*, see *Fontanella*—f. *Quadrangularis*, see *Fontanella*.

FOOD, (Sax. *foode*,) *Aliment*—f. *Farinaceous*, see *Farinaceous*—f. of the Gods, *Asafetida*.

FOOD, PRINCE OF WALES'S. A farinaceous preparation, which is used in the same cases as arrow-root, is said to consist entirely of potato-flour.

FOOL'S STONES, *MALE*, *Orchis mascula*.

FOOT, (Sax. *foet*,) *Pes*—f. *Flat*, see *Kyllosis*—f. *Griffon's*, *Gryphus pes*.

FOOTLING CASE. A presentation of the foot or feet in parturition.

FORAMEN, gen. *Foramina*, *Tre'ma*, *Opê*, (*foro*, 'I pierce.') *Audus*, (F.) *Trou*. Any cavity, pierced through and through. Also, the orifice of a canal.

FORAMEN ALVEOLARE ANTERIUS, see *Palatine canals*—f. *Alveolare posterius*, see *Palatine canals*—f. *Amplum pelvis*, *Obturatorium foramen*—f. *Anonymum Ferreii*, see *Aquæductus Fallopii*—f. *Aquæductus Fallopii*, f. *Stylomastoideum*—f. *Auditorium externum*, see *Auditory canal*, external—f. *Auditorium internum*, see *Auditory canal*, internal—f. of *Bichat*, see *Canal*, *arachnoid*—f. of *Botal*, see *Botal foramen*.

FORAMEN CÆCUM OF THE FRONTAL BONE, see *Cæcum foramen*.

FORAMEN CÆCUM OF THE MEDULLA OBLONGATA or of *VICQ D'AZYR*; (Fr.) *Trou borgne*. A tolerably deep fossa at the point where the median furrow at the anterior surface of the medulla oblongata meets the pons.

FORAMEN CÆCUM OSSIS MAXILLARIS SUPERIORIS, see *Palatine canals*—f. *Cæcum of the Tongue*, see *Cæcum foramen*, and *Tongue*—f. *Carotid*, see *Carotica foramina*.

FORAMEN CENTRALE seu Foræa centralis et Limbus Luteus Retinæ. The central foramen and yellow spot of the retina, discovered by Sömmerring. *Mucula lutea*, *flava seu aurea*, *Punctum aureum*, (F.) *Tache jaune*. It is situate about two lines to the outside of the optic nerve, and in the direction of the axis of the eye.

FORAMEN COMMUNE ANTERIUS, *Valva*—f. *Commune posterius*, see *Anus*—f. *Condyloid*, see *Condyloid*—f. *Conjugationis*, see *Conjugation*—f. *Ethmoideum*, *Orbital foramen*, internal—f. of *Galen*, *Botal foramen*—f. *Incisivum*, see *Palatine canal*—f. *Infracarbitarium*, see *Suborbital canal*—f. *Infrapubianum*, *Obturatorium foramen*—f. *Jugulare*, *Lacerum posterius foramen*—f. *Lacerum in basi cranii*, *Lacerum posterius foramen*—f. *Lacerum inferius*, *Sphenomaxillary fissure*—f. *Lacerum superius*, *Sphenoidal fissure*—f. *Magnum*, see *Occipital bone*—f. *Mastoid*, see *Mastoid foramen*—f. *Mental*, see *Mental foramen*.

FORAMEN OF MONRO, *Foramen Monroianum*. An opening behind the anterior pillar of the for-

nix, somewhat above the anterior commissure, by which the third ventricle communicates with the lateral ventricle; so called after the second Monro.

FORAMEN OF MORGAGNI, see *Cæcum foramen* and *Tongue*—f. *Obturatorium*, *Obturator foramen*—f. *Oculi*, *Pupili*—f. *Œsophageum* (of the diaphragm,) see *Œsophagus*, and *Œsophageal aperture of the diaphragm*—f. *Oodes*, *Ovale foramen*—f. *Opticum*, *Optic foramen*—f. *Opticum Chorioideæ*, see *Optic nerves*—f. *Orbitarium internum*, *Orbital foramen*, internal—f. *Orbitarium superius*, *Orbital foramen*, superior, f. *supra-orbitarium*—f. *Ovale*, *Botal foramen*, *Fenestra ovalis*, *Obturatorium foramen*, *Ovale foramen*—f. *Ovale*, *patency of the*, see *Cyanopathy*—f. *Palatinum anterius*, see *Palatine canals*—f. *Palatinum posterius*, see *Palatine canals*—f. *Palato-maxillare*, see *Palatine canals*—f. *Rotundum*, *Fenestra rotunda*—f. *Sphenospinosum*, *Spinal foramen*—f. *Spinale*, *Spinal foramen*—f. *Spinotum*, *Spinal foramen*—f. *Stylomastoideum*, *Stylomastoid foramen*—f. *Tarini*, see *Aquæductus Fallopii*—f. *Thyroideum*, *Obturatorium foramen*—f. of *Winslow*, *Hiatus of Winslow*.

FORAMINA CRIEROSA, see *Cribratus*.

FORAMINA THERE'SH. Openings resembling vascular orifices, found below the orifice of the vena cava superior in the right auricle, which are supposed to be the openings of veins.

FORATIO, (*forare*, *foratum*, 'to bore,') *Trepanning*.

FORCE, (*fortis*, 'strong,') *Vis*, *Potentia*, *Energia*, *Dynamis*, *Cratus*. Any power which produces an action. Those powers which are inherent in organization are called *vital forces*. We say, also, *organic force*, and *muscular force*, to designate that of the organs in general, or of the muscles in particular. To the latter the word *Dynamis*, *énergia*, corresponds; and the absence of this force is termed *adynamia*. The *vital forces* have to be carefully studied by the pathologist. The doctrine of diseases is greatly dependent on their augmentation or diminution; freedom or oppression, &c.

FORCE OF ASSIMILATION, *Plastic force*—f. *Catalytic*, see *Catalysis*—f. *Cell*, see *Cell force*—f. of *Formation*, *Plastic force*—f. *Germ*, *Plastic force*—f. *Mediatricæ*, *Vis mediatrix*—f. *Metabolicæ*, see *Metabolic force*—f. of *Nutrition*, *Plastic force*—f. *Osmotic*, see *Osmotic force*—f. of *Vegetation*, *Plastic force*—f. *Vital*, *Vis vitalis*—f. *Vitale*, *Vis vitalis*.

FORCE-RÉAL, MINERAL WATERS OF. The name of a mountain, situate four leagues from Perpignan in France. The water is chalybeate.

FORCEPS, gen. *Forcipis*, (quasi *ferrineps*, from *ferrum*, 'iron,' and *capio*, 'I take,') *Pincers*, *Lat'his*, *Volsell'is*. An instrument, for removing bodies which it would be inconvenient or impracticable to seize with the fingers. (F.) *Pince*, *Pincette*. There are various kinds of forceps, 1. The ordinary kind, contained in every dressing-case, for removing lint, &c., from wounds or ulcers. (F.) *Pince à pincement ou à anneau*. 2. *Dissecting or Ligature or Arterial Forceps*, (F.) *Pince à dissection*, *P. à ligature*, to lay hold of delicate parts. 3. *Polypus Forceps*, *Tooth Forceps*, *Forceps of Muenz*, (F.) *Pince de Muenz*, for laying hold of the tonsils or other parts to be removed. 4. *The Bullet Forceps*, (F.) *Pince-balle*. 5. *The Lithotomy Forceps*, *Lithot'abon*, *Lithot'gum*, *Tenaculum*, *Volsell'is*, which resembles the *Craniostrum Forceps*, (F.) *Tenette*. 6. *The Cataract Forceps*, (F.) *Pince à cataracte*, resembling the dissecting forceps, but much finer. 7. *The Pince à graine*, the branches of which can be brought together, and retained by a movable sheath.

Forceps is also an instrument used by obstetrical practitioners to embrace the head, and bring it through the pelvis. It consists of two branches, blades or levers; one of which, in the case of the *short forceps*, is passed over the ear of the child, and the other opposite the former, so that the blades may lock. When the head is securely inclined between the blades, the operation of extraction can be commenced. See Parturition.

FORCEPS, see *Corpus callosum*—*f.* Anterior, see *Corpus callosum*—*f.* Arterial, see *Forceps*—*f.* Bullet, see *Forceps*—*f.* Craniotomy, see *Forceps*—*f.* Lithotomy, see *Forceps*—*f.* Museux, see *Forceps*—*f.* Polypus, see *Forceps*—*f.* Tooth, see *Forceps*.

FORD'S BALSAM OF HOREHOUND, see *Balsam of horehound*.

FORE, *Ante*.

FORE'ARM, *Antibrah'chium*, *Pyg'mē*, *Pars inferior bra'chii*, *Cu'bitus*, (*F.*) *Avant-bras*. The part of the upper extremity, comprised between the arm and the hand. It is composed of two bones—*radius* and *ulna*—and 20 muscles.

FOREFRONT, *Front*.

FOREHEAD, *Front*.

FOREMILK, *Colostrum*.

FOREIGN BODY, *Extraneous body*.

FORENSIC ANATOMY, (*forum*, 'a court of justice,') see *Anatomy*—*f.* Medicine, Medicine, legal.

FORES, (pl. of *foris*, 'a gate,') *Vulva*.

FORESKIN, *Prepuce*.

FORETOP, *Front*.

FOREPEX, *Ciseau*, *Scissors*—*f.* Dentaria, *Dentagra*.

FORFIC'ULA AURICULA'RIA, (dim. of *forfex*, 'scissors,') *Ear'wig*. An insect of the Order Orthoptera, which occasionally enters the meatus auditorius externus, and excites intense pain. It may be destroyed by tobacco-smoke, or by oil poured into the meatus.

FORGES, MIN'ERAL WATERS OF. Forges is situate four leagues from Gournay, in the department of Seine Inférieure, France. There are three springs, which are acidulous chalybeates. These are called *Royal*, *Reinette*, and *Cardinal*, in honor of Louis XIII., Queen Anne of Austria, and Cardinal Richelieu, who used them.

FORMATIVE, (*formare*, *formatum*, 'to form,') *Plastic*.

FORMICA, *Myrmex*. 'The ant or pismire.' (*F.*) *Formica*. It contains an acid juice and gross oil, which were formerly extolled as aphrodisiacs. The chrysalides of the animal are said to be diuretic and carminative; and have been used in dropsy. Also the name of a black wart, *verruca formica'ria*, with a broad base and cleft surface; so called because the pain attending it resembles the biting of an ant, *μυρμηκία*, *myrmecia*. A varicose tumour on the anus and glans penis. Miliary herpes.

FORMICA AMBULATORIA, *Herpes circinatus*—*f.* *Corrosiva*, *Herpes exedens*.

FORMICANT, *Formic'ans*, *Myrmec'izon*, (from *formica*,) (*F.*) *Fourmillant*. An epithet given to the pulse, *Pulsus formic'ans*, when extremely small, scarcely perceptible, unequal, and communicating a sensation like that of the motion of an ant felt through a thin texture.

FORMICA'TION, *Formica'tio*, *Myrmecias'mus*, *Myrmec'asis*, *Myrmecias'mus*, *Myrmeco'sis*, *Stupor formic'ans*. (*F.*) *Fourmillement*. A pain, compared with that which would be caused by a number of ants creeping on a part.

FORMIX, (from *formica*, [?]) *Herpes esthiomenus*.

FORMULA, (dim. of *forma*, 'form or manner,') *Prescription*—*f.* *Extemporanea*, see *Magistral*—*f.* *Magistralis*, see *Magistral*—*f.* *Medica*, *Prescription*—*f.* *Officinalis*, see *Magistral*.

FORM'ULARY, *Formula'rium*, '*Narthecium*, *Narthecium*, *Narthex*. One skilled in preparing writs or forms of process. A collection of medical formulæ or receipts.

FORM'ULE, *Prescription*.

FORMYL, PERCHLORIDE OF. Chloroform.

FORMYLE, IODURE DE, Iodoform.

FORMYLUM CHLORATUM, Chloroform.

FOR'NIX, gen. *For'nicis*, 'an arch or vault.' *F. cer'ebri seu trila'terus seu tricuspid'a'tis*, *Ps'a'lis*, *Ps'alid'ium*, *Cam'erus*, *Testu'do cer'ebri*, *Ar'cus medulla'ris*, (*F.*) *Trigone cé'r'ébral*—(*Ch.*) *Voûte à trois piliers*, *Triangle Médullaire*. A medullary body in the brain, below the corpus callosum and above the middle ventricle, on the median line. This body, which is curved upon itself, terminates anteriorly by a prolongation, which constitutes its *anterior pillar*, *Crus seu Colum'na for'nicis*, (*F.*) *Pilier antérieur*; and posteriorly by two similar prolongations, called *posterior pillars* or *crura*. See *Achiculum*, and *Vault*.

FORNIX CEREBRI, *Fornix*—*f.* *Cranii*, *Calva*.

FORNPECKLES, *Ephelides*.

FORPEX, gen. *For'picis*, *Scissors*.

FORRET, ('forehead,') *Front*.

FORSTERA'NIA DIFFORM'IS, *Ech'ites difform'is seu puber'ula*; indigenous; *Ord.* *Apocynaceæ*; is said so be used as a wash with milk to remove freckles. The juice destroys warts.

FORTIFIANT, (*fortis*, 'strong,' and *facere*, 'to make,') *Corroborant*, *Tonic*.

FORTAITUDE, *Hysteralgia*.

FORTYKNOT, *Achyranthes repens*.

FOS'SA, *Fos'sea*, (*fodio*, *fossus*, 'I dig,') *Scam'ia*, (*F.*) *Fossa*. A cavity of greater or less depth, the entrance to which is always larger than the base. The fossæ of bones have been called *simple*, when they belong to one bone only, as the *parietal fossæ*; and *compound*, (*F.*) *Fossæ composées*, when several concur in their formation, as the *orbital fossæ*, *temporal fossæ*, &c.

FOSSA, AMYGDALOID, *Amygd'aloid excava'tion*. The space between the anterior and posterior pillars of the fauces, which is occupied by the tonsils.

FOSSA AMYN'TÆ. A kind of bandage, used in fractures of the nose; so called, by Galen, from Amyntas of Rhodes, its inventor. It consisted of a long band, applied round the head, the turns of which crossed at the root of the nose.

FOSSA CANINA, *Canine fossa*.

FOSSA CEREBEL'LI, (*F.*) *Fosse cé'r'ébelleuse*. The inferior occipital fossa, which lodges the corresponding portion of the cerebellum.

FOSSA CORONA'LIS, *Cor'on'al or fron'tal fos'sa*.

A depression on the orbital plate of the frontal or coronal bone, which supports the anterior lobe of the brain.

FOSSA CORONOI'DEA, *Cor'onoid fos'sa*. A cavity before the inferior extremity of the humerus, in which the coronoid process of the ulna is engaged during the flexion of the forearm.

FOSSA COTYLOIDEA, see *Cotyloid*.

FOSSA, DIGAS'TRIC, *Fos'sa digas'trica*. A deep groove on the mastoid portion of the temporal bone, which gives origin to the digastric muscle.

FOSSA ETHMOIDA'LIS, *Eth'moid fos'sa*. A shallow gutter on the upper surface of the cribriform plate of the ethmoid bone, in which is lodged the expanded portion of the olfactory nerves.

FOSSA OF THE GALLBLADDER, see *Gallbladder*—*f.* *Genu*, *Poples*—*f.* *Glandis*, see *Glans*.

FOSSA GUTTURA'LIS, *Gut'tural fos'sa*. The depression which forms the guttural region of the base of the cranium, between the foramen magnum and posterior nares.

FOSSA HYALOIDEA, see *Hyaloid (Fossa)*—*f.* *Iliac*, see *Iliac fossæ*—*f.* *Infrarorbital*, *Canine fossa*—*f.* *Infraspinous*, see *Infraspinata fossa*.

FOSSA INNOMINATA. The space between the helix and antihelix of the ear.

FOSSA, ISCHIO-RECTAL, Perineal fossa—*f. Jugularis*, Jugular fossa—*f. Lacrymalis*, Lacrymal fossa—*f. Lenticularis*, see Hyaloid membrane—*f. Magna Muliebris*, Vulva—*f. Magna Sylvii*, Fissura Sylvii.

FOSSA, MENTAL, *Fos'sa mentalis*. A small depression on each side of the symphysis on the anterior surface of the body of the maxilla inferior, for the attachment of muscles.

FOSSA NAVICULARIS, Navicular fossa.

FOSSA OCCIPITALIS, *Occipital fos'sa*. The occipital fossæ are four in number: the superior or cerebral, and the inferior or cerebellous. They are separated by a groove, which lodges the lateral sinus.

FOSSA OVALIS, see Ovalis fossa—*f. Palatina*, Palate—*f. Patellaris*, see Hyaloid membrane—*f. Perinaei*, Perineal fossa—*f. Pituitaria*, Sella Turcica.

FOSSA POPLITEA, *Poplite'al fos'sa*. The hollow of the ham;—the popliteal region.

FOSSA, PORTAL, see Liver—*f. Scaphoides*, Navicularis fossa.

FOSSA SIGMOÏDEA, *Sul'cus sinûs latera'lis*. A broad deep groove on the inner surface of the mastoid portion of the temporal bone, which supports part of the lateral sinus.

FOSSA, SUBORBITAR, Canine fossa.

FOSSA, SUBPYRAMIDAL, *F. subpyramida'lis*. A deep fossa under the pyramid and behind the fenestra rotunda in the middle ear, remarkable for its constancy, and pierced by several foramina at the bottom.

FOSSA SUPRAPHENOIDALIS, Pituitaria fossa—*f. Supraspinata*, see Supraspinatus—*f. of Sylvius*, Fissure of S.—*f. Temporalis*, Temporal fossa—*f. Transversa*, see Liver—*f. Umbilicalis*, see Liver.

FOSSÆ CEREBRALES, *Cer'ebral fos'sæ*. Fossæ or excavations at the base of the cranium. They are nine in number: three occupy the median line, and three are placed at each side. They are distinguished into anterior, middle, and posterior.

FOSSÆ DIGITALES, see Impression.

FOSSÆ, *Fos'sæ*—*f. Basilaire*, Basillary fossa—*f. Sous-épineuse*, Infra-spinata fossa—*f. Sous-épineuse*, Fossa supra-spinata—*f. de la Vésicule biliaire*, see Gallbladder.

FOSSES CONDYLOÏDIENNES, (pl. of *fosse*.) Condyloidea foramina—*f. Nasales*, Ouvertures postérieures des, Nares, posterior.

FOSSETTE (F.), (dim. of *fossa*.) *Scrobic'ulus*, *Both'rion*. Several depressions are so called. A dimpled chin, *Fossette du menton*, consists in a slight depression, which certain persons have on the chin. A dimple of the cheek, (F.) *Fossette des joues*, a depression which occurs on the cheeks of certain persons when they laugh. *Scrobic'ulus cordis*, *Antica'd-ion*, *Præcor'dium*, (F.) *Fossette du cœur*, is the depression observed on a level with the xiphoid cartilage at the anterior and inferior part of the chest. It is, also, called *pit of the stomach*, (F.) *Cœur de l'estomac*.

FOSSETTE, *Fos'sula*, *A'nulus*, *Bothrium*, is also a small ulcer of the transparent cornea, the centre of which is deep.

FOSSETTE ANGULAIRE DU QUATRIÈME VENTRICULE, *Calamus scriptorius*—*f. du Cœur*, *Scrobiculus cordis*.

FOSSORIUM, (*fossor*, 'a digger,') Fleam.

FOSSULA, (dim. of *fossa*, 'a ditch,') Argema, *Fossette*, *Fovea*.

FOTHERGILL'S PILLS, see *Pilule aloes et colocynthidis*.

FOTUS, (*fovere*, *fotum*, 'to keep warm,') Fo-mentation—*f. Communis*, Decoctum papaveris.

FOU, *Fol*, Insane.

FOUETS, *Sempervivum tectorum*.

FOUGÈRE DOUCE, *Polypodium*—*f. Femelle*, *Asplenium filix femina*, *Pteris aquilina*—*f. Grande*, *Pteris aquilina*—*f. Mâle*, *Polypodium filix mas*.

FOUL DISEASE, Syphilis.

FOULURE, Sprain.

FOUNDLING HOSPITAL, *Brephotropheum*.

FOURCHE, (F.), (*fourché*, 'cleft' from *furca*, 'a fork,') *Apostema Phalan'gum*. A French provincial term for small abscesses which form on the fingers and hands of working people. Also, an instrument, invented by M. J. L. Petit, for compressing the ranine artery in cases of hemorrhage from that vessel.

FOURCHETTE, (dim. of *fourche*, 'a fork,') *Furcilla*, a little fork, *Fur'cula*. A surgical instrument used for raising and supporting the tongue, during the operation of dividing the frænum.

Fourchette, in anatomy, is the posterior commissure of the labia majora, called, also, *Frænum*, *Frænum pudendi*, *Fur'cula Labio'rûm*; The cartilago ensiformis: so called from its being sometimes cleft like a fork. Also, the semilunar notch at the superior or clavicular extremity of the sternum.

FOURMI, *Formica*.

FOURMILLANT, (from *fourmi*,) Formicant.

FOURMILLEMENT, Formication.

FOUSEL OIL, ((G.) *Fusel*, 'bad brandy,') see Oil, *fusel*.

FOVEA, dim. *Fove'ola*, (*fodio*, 'I dig,') *Bo'thrus*, a slight depression, *Fos'sula*. The *pudendum muliebre*; see Vulva. The *fossa navicularis*, A vapour-bath for the lower extremities.

FOVEA AXILLARIS, Axilla—*f. Centralis* retinae, *Foramen centrale*—*f. Elliptica*, see *Crista Vestibuli*—*f. Hemisphærica*, see *Crista Vestibuli*—*f. Lacrymalis*, *Lachrymal fossa*, see *Fossa*—*f. Navicularis*, see *Urethra*—*f. Oculi*, *Orbit*—*f. Ovalis*, *Ovalis fossa*—*f. Semi-elliptica*, see *Crista Vestibuli*—*f. Sulciformis*, see *Crista Vestibuli*.

FOVEOLA, see *Fovea*.

POWER, Syncope.

POXBERRY, *Arbutus uva ursi*.

POXGLOVE, *Digitalis*.

FOYER, ((L.) *focus*, 'a fire place,') *Fomes morbi*.

FRACTIONNEMENT, (*frangere*, *fractum*, 'to break,' 'to bruise,') Segmentation.

FRACTURA, *Fracture*—*f. Dentis*, *Odontoclasia*.

FRACTURE, *Fractu'ra*, *Catag'ma*, *C. Fractu'ra*, *Catax'is*, *Clas'is*, *Clas'ma*, *Ag'mè*, *Ag'ma*, *Di-ar'ragè*. A solution of continuity in a bone, *Osteoc'lasis*. A simple fracture is when the bone only is divided. A compound fracture is a division of the bone with a wound of the integuments communicating with the bone,—the bone, indeed, generally protruding. In a comminuted fracture, *Alphite'don*, ἀλφειτῶν, *Carye'don* *Catag'ma*, καρυδὸν κατὰ γμα, the bone is broken into several pieces; and in a complicated fracture there is, in addition to the injury done to the bone, a lesion of some considerable vessel, nervous trunk, &c. Fractures are also termed *transverse*, *oblique*, &c., according to their direction. The treatment of fractures consists, in general, in reducing the fragments when displaced; maintaining them when reduced; preventing the symptoms which may be likely to arise; and combating them when they occur. The reduction of fractures must be effected by extension, counter-extension, and coaptation. The parts are kept in apposition by position, rest, and an appropriate apparatus. The position must vary according to the kind of fracture. Commonly, the fractured limb is placed on a horizontal or slightly inclined plane, in a

state of extension: or rather in a middle state between extension and flexion, according to the case.

FRACTURE, CAMERATED, Camerosis.

FRACTURE OF THE RA'DIUS, BAR'TON'S, *Bar'ton's fracture*. A term applied to a fracture of the lower extremity of the radius, which commences at the articular surface, and extends upwards for an inch or more, to terminate on the dorsal aspect. Owing to the extensor muscles drawing up the separated portion of the bone, and with it the carpus, a deformity results, which has been confounded with simple dislocation. In consequence of the fracture having been well described by Dr. John Rhea Barton, of Philadelphia, it is often called after him.

FRACTURE OF THE RA'DIUS, COL'LES'S, *Col'les's fracture*. A name given to a fracture of the radius at the distance of about an inch and a half above the radio-carpal articulation, well described by Professor Colles, of Dublin.

FRACTURE EN RAVE, Raphanodon.

FRACTURES PAR RÉSONNANCE, see *Contraffissura*.

FRÆNA MORGAGNII, (pl. of *frænum*), F. of the Valve of Bauhin.

FRÆNA OF THE VALVE OF BAU'HIN, (F.) *Freins de la valvule de Bauhin*. A name given by Morgagni to the projecting lines formed by the junction of the extremities of the two lips of the ileo-cæcal valve. They are also called *Fræna* and *Retinacula Morgagni*.

FRÆNULUM, (dim. of *frænum*), see *Frænum*, *Bride* — f. *Clitoridis*, *Frænum clitoridis* — f. *Epiglottidis*, *Glossoepiglottic ligament* — f. *Labiolum*, *Fourchette* — f. *Novum*, *Tænia semicircularis* — f. *Pudendi*, *Fourchette*.

FRÆNULUM VE'LI MEDULLA'RIS ANTERIO'RIS. A narrow slip, given off by the commissure of the encephalic testes, which strengthens the junction of the testes with the valve of Vieussens.

FRÆNUM, *Fre'num*, (F.) *Frein*, 'a bridle.' — *Frænulum*, *File'tum*, *Filamentum*. 'A small bridle.' — A bridle. Names given to several membranous folds, which bridle and retain certain organs.

FRÆNUM CLITO'RIDIS, *Frænulum Clito'ridis*, (F.) *Frein du Clitoris*. A slight duplicature formed by the union of the internal portions of the upper extremity of the nymphæ.

FRÆNUM EPIGLOTTIDIS, *Glosso-epiglottic ligament* — f. *Glandis*, F. *Penis*.

FRÆNUM LABIO'RUM, (F.) *Frein des lèvres*. There are two of these; one for the upper, the other for the lower lip. They unite these parts to the maxillary bone, and are formed by the mucous membrane of the mouth. Also, the *Fourchette*.

FRÆNUM LIN'GUÆ, *Frænulum* seu *Vinculum Linguae*, *Glossodes'mus*, *File'tum*, (F.) *Filet* ou *Frein de la langue*, is a triangular reflection formed by the mucous membrane of the mouth, and situate between the inferior paries of that cavity and the inferior surface of the tongue. When the frænum extends as far as the extremity of the tongue, it cramps its movements, interferes with sucking, &c. This inconvenience is remedied by carefully snipping it with a pair of scissors. The French call this *l'opération du filet*.

FRÆNUM PE'NIS, F. seu *Frænulum* seu *Vinculum Præputii*, F. *Glandis*, *Cynodes'mion*, *Cynodes'mus*, *Vinculum Can'num*, (F.) *Filet* ou *Frein de la verge*, is a membranous reflection which fixes the prepuce to the lower part of the glans. When too short, it prevents the prepuce from sliding over the glans.

FRÆNUM PRÆPUTII, F. *Penis*.

FRAGA'RIA, (*frago*, 'I smell sweetly.') The *Strawberry*, *Fraga'ria vesca* seu *vulga'ris* seu *seu-*

perfol'rens, *Chama'batus*, (F.) *Fraisier*. Ord. Rosaceæ. The fruit is agreeable and wholesome, and the roots have been esteemed tonic and slightly diuretic. The fruit is the *Fragrum*, *κορυμνον*, of the ancients; (F.) *Fraise*.

FRAGARIA ANSERINA, *Potentilla anserina* — f. *Pentaphyllum*, *Potentilla reptans* — f. *Tormentilla officinalis*, *Tormentilla*.

FRAGARIA VIRGINIA'NA, Wild *Strawberry*. An indigenous plant, which has astringent leaves.

FRAGILE VITREUM, *Fragilitas ossium*.

FRAGILITAS, *Ruptibilitas*, *Ruptilitas*, (*fragilis*, 'brittle,' from *frango*, 'I break.') *Fragility*, *Brittleness*. The state of being easily broken or torn.

FRAGILITAS OS'SIUM, *Osteopathyrosis*, *Paros'tia frag'ilis*, *Brittleness of the bones*, *Friability of the bones*, *Frag'ile vit'reum*. Pathologists have given this name to the extreme facility with which bones break in certain diseases of the osseous texture. It is owing to a deficiency of the animal matter.

FRAG'MENT, *Fragmentum*, *Frag'men*, *Ramen'tum*, (*frangere*, *fractum*, 'to break.') The French use this term for the two portions of a fractured bone; thus, they speak of the *superior* and the *inferior fragment*.

FRAGMENTS, PREC'IOUS. A name formerly given, in *Pharmacy*, to the garnet, hyacinth, emerald, sapphire and topaz. The Arabs falsely attributed to them cordial and alixiterial properties.

FRAGON, *Ruscus*.

FRAGUM, see *Fragaria*.

FRAISE, see *Fragaria*.

FRAISIER, *Fragaria*.

FRAMBÆ'SIA, *Frambæ'sia*, *Le'pra fungif'era*, ([F.], *framboise*, 'a raspberry,') *Symph'itis Ind'ica*, *Anthra'cia ru'bula*, *Thym'osis*, *Th.* seu *Lu'es Ind'ica*, *Vari'ola Amboinen'sis*, *Le'pra fungif'era*, *Scroph'ula Molluca'na*. The *Yaws*, *Epian*, *Pi'an*. A disease of the Antilles and of Africa, characterized by tumours, of a contagious character, which resemble strawberries, raspberries, or champignons; ulcerate, and are accompanied by emaciation. The *Pi'an*, for so the Indians call it, differs somewhat in America and Africa.

Pi'an of America, *Frambæ'sia America'na*, *Anthra'cia Ru'bula Americana*, occurs under similar circumstances with the next, and seems to be transmitted by copulation. The tumours have a similar form, and are greater in proportion to their paucity. In some cases they are mixed with ulcers.

Pi'an of Guinea, *Frambæ'sia Guineen'sis*, *Anthra'cia Ru'bula Guineen'sis*, is common amongst the negroes, especially in childhood and youth. It begins by small spots, which appear on different parts, and especially on the organs of generation and around the anus; these spots disappear, and are transformed into an eschar, to which an excrescence succeeds, that grows slowly, and has the shape above described.

The treatment is nearly the same in the two varieties. The tumours will yield to mercurial friction, when small. When large, they must be destroyed by caustic. In both cases, mercury must be given to prevent a recurrence.

An endemic disease resembling yaws was observed in the Feejee Islands by the medical officers of the United States' Exploring Expedition. It is called by the natives *Dhoko*.

FRAMBÆSIA ILLYRICA, *Scherlievo* — f. *Scotica*, *Sibbens*.

FRAMBOISE, *Rubus idæus*.

FRANCOLIN, *Attagen*.

FRANGES SYNOVIALES, ('synovial fringes,') *Synovial glands*.

FRANGIPANE. An article of food, prepared

by evaporating to dryness, in a water-bath, milk mixed with almonds and sugar.

FRANGULA ALNUS, (*frangula*, 'to break,' from its brittleness.) *Rhamnus frangula*.

FRANKFORT, MINERAL WATERS OF. These springs are in Beaver County, Pennsylvania, near the village of Frankfort. Cane spring and Loper spring contain carbonic acid, carbonates of iron and magnesia, hydrosulphuric acid, chloride of sodium, and a small portion of bitumen. They are recommended in dyspepsia, rheumatic and cutaneous affections, and in a variety of diseases.

FRANKINCENSE, COMMON, (*frank*, and 'incense,') see *Pinus abies*—*f. African*, *Daniellia thurifera*—*f. Tree*, *Daniellia thurifera*—*f. True*, *Juniperus lycia*.

FRANZENBAD, FRANZBAD, or FRANZENBRUNN, MINERAL WATERS OF. A celebrated water at Eger, in Bohemia, which contains sulphate of soda, carbonate of iron, and carbonic acid gas. The springs are called *Franzenbrunn*.

FRANZO, *Spiræolus*.

FRAPPER, ('to strike,') Percuss.

FRASERA, (after John Fraser, a botanical collector,) *Calumba*, American—*f. Carolinensis*, see *Calumba*—*f. Officinalis*, see *Calumba*—*f. Verticillata*, *Calumba*, American—*f. Walteri*, see *Calumba*.

FRATER UTERINUS, ('uterine brother,') see *Uterinus frater*.

FRATERNITAS, (*frater*, 'a brother,') *Adelphixia*.

FRATRATIO, *Adelphixia*.

FRAXINELLA DICTAMNUS, (dim of *fraxinus*), *Dictamnus albus*—*f. White*, *Dictamnus albus*.

FRAXINUS AMERICANA, *White Ash*; and **FRAXINUS QUADRANGULATA**, *Blue Ash*, indigenous; *Ord*, *Oleaceæ*; have bitter and astringent barks, and have been used as antiperiodics.

FRAXINUS APETALA, *F. excelsior*—*f. Aurea*, *F. excelsior*—*f. Crispa*, *F. excelsior*—*f. Excelsa*, *F. excelsior*.

FRAXINUS EXCELSIOR. The systematic name of the *Ash tree*, *Fraxinus excelsior*, *F. apetal* seu *an'ra seu oris'pa seu areol'ra seu pau'dala seu verruc'ra*, *Bume'ra* seu *Macedon'ra* *Fraxinus*, *Bume'ra*, (*F. Ficus*). The fresh bark has a moderately strong, bitterish taste. It has been said to possess resolvent and diuretic qualities, and has been given in intermittents, and in gout and rheumatism, and as an antidote to serpent bites. The seeds, called *Bird's tongue*, *Lingua asis*, *Ornithoglossa*, have been exhibited as diuretics, in the dose of a drachm. Its sap has been extolled against deafness.

FRAXINUS FLORIFERA, *F. ornus*—*f. Macedonica*, *F. excelsior*.

FRAXINUS ORNUS, *F. mel'ra* seu *pantoda'ra* seu *florifera*, *Ornus mannifera* seu *rotundifolia*. The systematic name of the tree whence *mannu* flows. This substance is also called *Manna Calabrina*, *Ros Calabritana*, *Almond's*, *Broom'els*, *Dryan'els*, *Mel ad'rium*, *Succus orni concortus*. In Sicily, not only the *Fraxinus Ornus*, but also the *F. rotundifolia* and *F. excelsior* are regularly cultivated for the purpose of procuring manna, which is their condensed juice. Manna is inodorous, sweetish, with a very slight degree of bitterness; in friable flakes, of a whitish or pale yellow colour; opaque, and soluble in water and alcohol. It is laxative, and is used as a purgative for children, who take it readily on account of its sweetness. More generally it is employed as an adjunct to other purgatives. Dose, $\frac{5}{32}$ to $\frac{3}{8}$. Its immediate principle is called *Mannite* or *Mannin*. This has been recommended by Ma-

gendie as a substitute for manna. Dose, $\frac{5}{32}$, for children.

FRAXINUS PANICULATA, *F. Ornus*—*f. Quadrangulata*, see *F. Americana*—*f. Rotundifolia*, *F. ornus*.

FRAYEUR NOCTURNE, ('nocturnal fright,') *Panophobia*.

FRECKENS, ([*G.*] *Fleck*, a 'spot,') *Ephelides*.

FRECKLED, see *Ephelides*.

FRECKLES, *Ephelides*.

FREEMAN'S BATHING SPIRITS, see *Linimentum saponis compositum*.

FREIN, *Frænum*—*f. du Clitoris*, *Frænum clitoridis*—*f. de la Langue*, *Frænum lingue*—*f. des Lèvres*, *Frænum labiorum*—*f. de la Vierge*, *Frænum penis*.

FREINS DE LA VALVULE DE BAUHN, *Freins* of the valve of Bauhin.

FREINWALDE, MINERAL WATERS OF. These springs are in Brandenburg, twelve leagues from Berlin. They contain chloride of sodium, sulphate of magnesia, chloride of magnesium, sulphate of lime, carbonates of lime and magnesia, iron, &c., and are used in asthenic diseases.

FREMISSEMENT, *Fremitus*, Shuddering.

FREMISSEMENT CATAIKE, (*F.* 'Cat's purr, or fremitus,' *Pur'ring Tremor*, *Laënnec* has given this name to the agitation which is sensible to the hand when applied on the præcordial region, and which he considers a sign of ossification or other contraction of the auriculo-ventricular openings. The name was chosen by him from the analogy of the sound to the purring of a cat.

FREMITUS, (*fremere*, *fremitum*, 'to murmur,') *Brimissement*, *Fremissement*, Shuddering.

The **Pæctoral or Vo'cal FREMITUS**, *Pæ'ctoral Vibration*, *Tactile vibration* is an obscure diffused resonance of the voice which is felt when the hand is applied to the chest.

A **FRICTION FREMITUS** is perceptible to the hands in many cases of pleurisy, when the surfaces of the pleura have become roughened from deposition, as well as in rare cases of pericarditis. In the former cases, it is called the *pleural*; in the latter, the *pericardial friction fremitus*.

FREMITUS HYDATID, *Son hydatique*.

FRENA, (pl. of *frenum*,) see *Alveolus*.

FRENCH CRUST, Syphilis—*f. Distemper*, Syphilis—*f. Letter*, Condom—*f. Nut*, Juglans regia—*f. Pox*, Syphilis.

FRÈNE, (Old [*F.*] *Freene*,) *Fraxinus excelsior*—*f. Épinera*, *Xanthoxylum clava Herculis*—*f. Sanguis*, *Sorbus aucuparia*.

FRENETIC, *Phrenetic*.

FRENGA, see *Scherlieve*.

FRENULUM, see *Frænum*.

FRENUM, *Frænum*.

FRENZY, *Phrenitis*.

FRET, (Sax. *freosan*, 'to rub,' and *fretan*, 'to eat or gnaw,') Chafing, Herpes.

FRÉTILLEMENT, Fidgets.

FRETTING, *Tormina*.

FRICTATIO, (*fricare*, *friectum*, 'to rub,') *Friction*.

FRICTATOIRES, (from *fricare*, 'to rub,') See *Shampooing*.

FRICTATORIUM, *Liniment*.

FRICTATRIX, *Tribas*.

FRI'CE, *Fricum*, *Friconium*. A medicine which the ancients employed under the form of friction. They distinguished the *Fric' siccum* and *F. molle*.

FRICONIUM, *Frice*.

FRICTA, *Colophonia*.

FRICTIO, *Frictio*—*f. Humida*, see *Friction*—*f. Sicca*, see *Friction*.

FRICTION, *Fric'tio*, *Fric'tio*, *Anat'ribē*, *Anat'rip'sia*, *Tryp'sia*, *Chirap'sia*. The action of rubbing—as of a part of the surface of the body—more or less forcibly, with the hands, a brush, flannel, &c., constituting *Xerotrib'ia*, *Xerotrip'sis*, *Fric'tio sic'ca* or *dry fric'tion* or *rub'bing*, or with ointments, liniments, tinctures, &c., constituting *moist fric'tion* or *rub'bing*, *Fric'tio hu'mida*. It is a useful means for exciting the action of the skin.

FRICTION FREMITUS, see *Fremitus*—f. Sound, *Bruit de frottement*.

FRICTRIX, Tribas.

FRICTUM, Liniment.

FRICTUM, Frice.

FRIENDBACK, Hangnail.

FRIGEFACIENTIA, (*frigus*, 'cold,' and *facere*, 'to make,') Refrigerants.

FRIGID, *Frig'idus*, (F.) *Froid*, (*frigeo*, 'to be cold,') Cold. Not easily moved to sexual desire; *Imbell'is ad ven'erem*. Impotent.

FRIGIDARIUM, Bath, cold.

FRIGIDITY, *Frigid'itas*. A sensation of cold. Also, impotence and sterility. *Frigidity of the stomach* is a state of debility of that organ, imputed to excessive venery,—the *Anorex'ia exhausto'rum* of Sauvages.

FRIGIDUS, Frigid.

FRIGORIFIC, (*frigus*, 'cold,' and *frigo*, 'I become,') That which has the power of producing cold. The best **FRIGORIFIC MIXTURES** are the following. Their effects are owing to the rapid absorption of heat when solids pass into the liquid state.

FRIGORIFIC MIXTURES WITH SNOW.

Mixtures.	Therm. falls.
Snow, or pounded ice, two parts by weight;	From any temperature. to—50°
Chloride of Sodium..... 1.	
Snow or pounded ice..... 5.	
Chloride of Sodium..... 2.	to—12°
Muriate of Ammonia..... 1.	
Snow or pounded ice..... 24.	
Chloride of Sodium..... 10.	to—18°
Muriate of Ammonia..... 5.	
Nitrate of Potash..... 5.	
Snow or pounded ice..... 12.	to—25°
Chloride of Sodium..... 5.	
Nitrate of Ammonia..... 5.	
Dilute Sulphuric Acid..... 2.	from +32°
Snow..... 3.	to—23°
Concentrated Muriatic Acid..... 5.	from +32°
Snow..... 8.	to—27°
Concentrated Nitrous Acid..... 4.	from +32°
Snow..... 7.	to—30°
Chloride of Calcium..... 5.	from +32°
Snow..... 4.	to—40°
Crystall. Chloride of Calcium..... 3.	from +32°
Snow..... 2.	to—50°
Fused Potash..... 4.	from +32°
Snow..... 3.	to—51°

Frigorific Mixtures may also be made by the rapid solution of salts, without the use of snow or ice. The salts must be finely powdered and dry.

FRIGORIFIC MIXTURES WITHOUT SNOW.

Mixtures.	Therm. falls.
Muriate of Ammonia..... 5.	from +50° to—10°
Nitrate of Potash..... 5.	
Water..... 16.	
Muriate of Ammonia..... 5.	from +50° to—10°
Nitrate of Potash..... 5.	
Sulphate of Soda..... 8.	
Water..... 16.	from +50° to—4°
Nitrate of Ammonia..... 1.	
Water..... 1.	
Nitrate of Ammonia..... 1.	from +50° to—7°
Carbonate of Soda..... 1.	
Water..... 1.	
Sulphate of Soda..... 3.	from +50° to—3°
Dilute Nitrous Acid..... 2.	
Sulphate of Soda..... 6.	
Muriate of Ammonia..... 4.	from +50° to—10°
Nitrate of Potash..... 2.	
Dilute Nitrous Acid..... 4.	

Mixtures.

Therm. falls

Sulphate of Soda..... 6.	from +50° to—14°
Nitrate of Ammonia..... 5.	
Dilute Nitrous Acid..... 4.	
Phosphate of Soda..... 9.	from +50° to—12°
Dilute Nitrous Acid..... 4.	
Phosphate of Soda..... 9.	
Nitrate of Ammonia..... 6.	from +50° to—21°
Dilute Nitrous Acid..... 4.	
Sulphate of Soda..... 8.	
Muriatic Acid..... 5.	from +50° to—0°
Sulphate of Soda..... 5.	
Dilute Sulphuric Acid..... 4.	

FRIGUS, gen. *Fri'goris*, Cold—f. Tenue, see *Rigor*.

FRINGE TREE, *Chionanthus Virginica*.

FRINGED, ([F.] *frange*, 'a fringe,') Laciniated.

FRISSEON, (from *frigere*, 'to be cold,') *Rigor*.

FRISSEONNEMENT, Horripilation.

FROGLEAF, *Brasenia Hydrophilis*.

FROG TONGUE, *Ranula*.

FROGS' SPAWN, *Sperma ranarum*.

FROID, (from *frigidus*,) Cold, Frigid—f. *Gla'cial*, Ice cold.

FROISSEMENT, [F.] (from low L. *fressare*, 'to bruise,') Rubbing, bruising.

FROISSEMENT PULMONAIRE, *Bruit de froissement pulmonaire*, *Pul'monary crumpling sound*. A name given by M. Fournet to a respiratory sound, which communicates to the ear the sensation of the rubbing (*froissement*) of a texture compressed against a hard body. It is by no means well defined.

FRÔLEMENT, (F.) 'Grazing or touching lightly.'

FRÔLEMENT PÉRICARDIQUE, *Bruit de frôlement péricardique*. Rustling noise of the pericardium. A sound resembling that produced by the crumpling of a piece of parchment or of thick silken stuff, accompanying the systole and diastole of the heart. It indicates roughness of the pericardium induced by disease.

FROMAGE, Cheese.

FROMENT, (from [L.] *frumentum*,) Triticum.

FRONCEMENT, Corrugation.

FRONCLE, Furunculus.

FRONDE, Funda.

FRONS, gen. *Frontis*, Front.

FRONT, *Frons*, *Metop'ion*, *Sto'ma*, *Forehead*, *Forefront*, *Fore'top*, *Brow*, (Sc.) *Forret*. That part of the visage which extends from one temple to the other, and is comprised in a vertical direction, between the roots of the hair and the superciliary ridges.

FRON'TAL, *Fronta'lis*. Relating or belonging to the front. This name has been given to several parts. Winslow, Sömmerring, and others, call the anterior part of the occipito-frontalis—the *frontal muscle* or *fronta'lis*, *Musculo'sa Front'is Cutem mo'vens substan'tia* Par of Vesalius.

FRONTAL ARTERY, *Arte'ria supra-orbita'lis*, is given off by the ophthalmic, which is itself a branch of the internal carotid. It makes its exit from the skull at the upper part of the base of the orbit, and ascends the forehead between the bone and the orbicularis palpebrarum; dividing into three or four branches, which are distributed to the neighboring muscles.

FRONTAL BONE, *Os fron'tis seu corona'le seu inverecundum seu pup'pis seu synep'itis seu pro'næ seu ratio'nis*, *Metop'ion*. A double bone in the foetus, single in the adult, situate at the base of the cranium, and at the superior part of the face. It forms the vault of the orbit; lodges the ethmoid bone in a notch at its middle part; and is articulated, besides, with the sphenoid, parietal, and nasal bones, the ossa unguis, superior maxillary, and malar bones.

FRONTAL FUR'ROW extends upwards from the frontal spine, and becomes gradually larger in its

course, to lodge the upper part of the superior longitudinal sinus, and to give attachment to the falx cerebri.

FRONTAL NERVE. *Palpébro-frontal*—(Ch.), is the greatest of the three branches of the ophthalmic nerve,—the first division of the fifth pair. It proceeds along the superior paries of the orbit, and divides into two branches:—the one *internal*, which makes its exit from the orbital fossa, passing beneath the pulley of the oblique muscle:—the other, *external*, issuing from the same cavity by the foramen orbitarium superius.

FRONTAL PROTUBERANCE, *F. tuberosity*, *Tu'ber fronta'le*, *Supgrund'um supercilio'rum*. The protuberance of the frontal bone above the superciliary arch.

FRONTAL SINUSES, *Si'nus Fronta'les*, *Metopop'tra*, *Pneupop'tra*, *Si'nus Supercilia'res* seu *pituita'rii fron'tis*, *Cavernæ fron'tis*, are two deep cavities in the substance of the frontal bone, separated from each other by a median septum, and opening, below, into the anterior cells of the ethmoid bone.

FRONTAL SPINE, *Cris'ta inter'na*, is situate in the middle of the under part of the bone, and is formed by the coalescence of the inner tables for the attachment of the falx cerebri.

Surgeons have given the name **FRONTALIS** to a bandage or topical application to the forehead. Such have, also, been called *παραμετωπία* and *ανακόλλημα*.

FRONTAL TUBEROSITY, Frontal protuberance.

FRONTALĒ, (from *frons*.) A medicine applied to the forehead.

FRONTALIS ET OCCIPITALIS, Occipitofrontalis.

FRONTALIS VERUS, Corrugator supercilii.

FRONTODYMIA, see Cephalodymia.

FRONTOETHMOID FORAMEN, Cæcum foramen—*f. Nasal*, *Pyramidalis nasi*—*f. Sourcilier*, Corrugator supercilii.

FROSTBITE, Congelation.

FROSTPLANT, *Helianthemum Canadense*.

FROSTWEED, *Erigeron Philadelphicum*, *Helianthemum Canadense*.

FROSTWORT, *Helianthemum Canadense*.

FROTH'Y, (*αἶρος*, 'froth.' [?]) *Spu'mous*, (F.) *Spumeux*, *Écumeur*, *Mousseux*. An epithet given to the fæces or sputa when mixed with air.

FROTTEMENT, (from low L. *fricare*, itself from *fricare*, 'to rub.'). See *Bruit de frottement*.

FROTTEMENT GLOBULAIRE. A name given by M. Simonnet to the pulse in aortic regurgitation, when it is jerking; and, in well-marked cases, appears as if the blood consisted of several little masses, which passed in succession under the finger applied to the artery.

FRUCTIFICATIO, (*fructus* and *facere*, 'to make.'). Fecundation.

FRUCTUS, Fruit—*f. Horsei*, Fruit (summer)—*f. Immaturus*, Abortion.

FRUGIVOROUS, *Frugiv'orus*, (*fruges*, 'fruits,' and *oro*, 'I eat.'). One that eats fruits.

FRUIT, *Fructus*, (*fruo*, *fructus*, 'to enjoy.'). *Car'pos*. In botany, the seed with its enclosing pericarp. In the ordinary acception, it means exclusively seed cases, which are eatable; and, generally, such as require no preparation to render them fit for food. The effects of fruits on the body, in a medical as well as a dietetical point of view, are various. They may be distinguished into classes; for, whilst the *Cerealia*, for example, afford fruits which are highly nutritious, the *Summer Fruits* (*Fructus Horæ'i*), which include strawberries, cherries, currants, mulberries, raspberries, figs, grapes, &c., are refrigerant and grateful, but afford little nourishment.

Fruit, (F.), is the foetus, whilst contained in the womb of the mother.

FRUIT-SUGAR, Glucose.

FRUMENT, *Fru'menty*, *Fur'menty*, (*frumentum*, 'wheat or grain,' quasi *frugimentum*, from *fruges*, 'fruits.'). Pottage made of wheat. Food made of wheat boiled in milk.

FRUMENTUM. *Sitos*. Any kind of grain from which bread was made; especially wheat.

FRUMENTUM, *Triticum*—*f. Corniculatum*, *Ergot*—*f. Cornutum*, *Ergot*—*f. Luxurians*, *Ergot*—*f. Temulentum*, *Ergot*—*f. Turgidum*, *Ergot*.

FRUSTRATOIRE (F.). Any liquor, taken a short time after eating, for the purpose of assisting digestion when difficult. Sugared water, *eau sucrée*, or water with the addition of a little brandy, or some aromatic substance, is commonly used for this purpose.

FRUTEX BACCIFER BRAZILIENSIS, *Caa-gihivuyo*—*f. Indicus spinosus*, *Cara schulli*.

FUCUS, (*φύκος*, 'sea weed,') *F. vesiculosus*, *Paint*.

Fucus *AMYLACEUS*, *Jaff'na*, *Ed'ilde* or *Ceylon Moss*, *Ploca'ria can'dida*, *Gracila'ria seu Sphaerococcus seu Gigarti'na seu Fucus lichenoides*, *Marine' Moss*. This moss belongs to the natural order Algæ. It was introduced some years ago into England, from India. It is white, filiform and fibrous, and has the usual odour of sea-weeds. Its medical properties are similar to those of Irish moss.

Fucus, *BLADDER*, *F. vesiculosus*.

Fucus *CRISPUS*, *F. Irlandicus*, *Lic'h'en Car'ra-geen*, *Chou'drus* (Ph. U. S.), *Ch. crispus* seu *polymor'phus*, *Sphaerococcus crispus*, *U'va crisp'a*, *Irish Pearl*, *Curraheen*, *Curraheen* or *Corriggen' moss*, (F.) *Mousse d'Irlande* ou *perlée*. This *Fucus* is found on the coasts of England, Ireland, Western France, Spain, and Portugal, and as far as the tropics. It is also a native of the United States. In Ireland, it is used by the poor as an article of diet. As met with in America, it is of light yellow colour, and resembles plates of horn, crisped and translucent. An agreeable jelly is obtained from it by boiling it in water or milk, which forms a good article of diet in consumptive cases. Its properties are indeed exactly like those of the *Iceland Moss*.

Fucus *HELMINTHOCORTON*, *Corallina Corsicana*—*f. Inflatus*, *F. vesiculosus*—*f. Irlandicus*, *Fucus crispus*—*f. Lichenoides*, *F. amylaceus*—*f. Saccharine*, *Rhodomela palmata*.

Fucus *VESICULOSUS*, *Fucus*, *F. inflatus*, *Hal'idrya vesiculosa*, *Quer'cus seu Lactu'ca Mari'na*, *Blad'der Fu'cus*, *Sea Oak*, *Sea Wrack*, *Yell'ow Blad'derwrack*, (F.) *Varec écisiculeux*, *Chêne marin*. It has been said to be a useful assistant to seawater in the cure of disorders of the glands. When the wrack, in fruit, is dried, cleaned, exposed to a red-heat in a crucible with a perforated lid, and is reduced to powder, it forms the *Æ'thiops vegetabilis*—the *Pulvis Quer'cus Mari'næ* of the Dublin Pharmacopœia—which is used, like the burnt sponge, in bronchocœle and other serofulous swellings. Its efficacy depends on the iodine it contains. Dose, gr. x to ℥ij, mixed in molasses or honey. See *Soda*.

FUGA DÆMONUM, ('aversion of the demons,') *Hypericum perforatum*.

FUGACIOUS, *Fugax*, gen. *fugacis*, (F.) *Fugace*, (*fugere*, 'to fly.'). An epithet given to certain symptoms, which appear and disappear almost immediately afterwards; as a *fugacious redness*,—a *fugacious swelling*, *Tumor fugax*, &c.

FUGAX, *Fugacious*.

FUGE, (*fugo*, 'I expel,') 'an expeller.' A common suffix. Hence, *Febri'fuge*, *Vermi'fuge*, &c.

FUGILĒ, *Fugil'la*. This term has several acceptations. It means, 1. The cerumen of the ear. 2. The nebulous suspension in, or deposition from, the urine. 3. An abscess near the ear. 4. Abscess in general.

FUGILLA, Fugile.

FULGUR, gen. *Fulguris*, 'lightning,' (from *fulgere*, 'to flash,') *Astrape*.

FULIG'INOUS, *Fuliginosus*, *Lignyo'des*, (F.) *Fuliginæus*, (*fuligo*, gen. *fuliginis*, 'soot,') Having the colour of soot. An epithet given to certain parts, as the lips, teeth, or tongue, when they assume a brownish colour, or rather are covered with a coat of that colour,—*Fuliginos'ity*.

FULIGO, gen. *Fulig'inis*, *Lign'ys*, *Soot*, (F.) *Suic*, &c., *Wood'soot*, *Ful'go Lign'i*, consists of volatile alkaline salt, empyreumatic oil, fixed alkali, &c. A tincture, *Tinctura Fulig'inis*, prepared from it, has been recommended as a powerful antispasmodic in hysterical cases. (*Fulig. lign. 3j; Potass. subcarb. lss; Ammon. murat. 3j; Aquæ fluviat. Oij.* Digest for three days.) This tincture bears the name *Soot drops* and *Fit drops*. An ointment of soot has been used in various cutaneous diseases.

FULIGO ALBA PHILOSOPHORUM, *Ammoniaë murias*.

FULIGO'KALI, (*fuligo*, 'soot,' and *kali*, 'potassa,') This is an analogous preparation to anthrakokali; soot being used in the place of coal. It is employed in the same diseases. What might be regarded as a weak solution of fuligokali has been used for many years in Philadelphia, under the names *medic' lye*, *soot tea*, *alkaline solution*, *dyspeptic lye*, and *Physick's lye tea*. It is made by mixing a quart of hickory ashes; half a pint of soot; and a gallon of boiling water; allowing the mixture to stand for twenty-four hours, and decanting.

FULLERS' EARTH, *Cimolia purpurescens*.

FULMEN, ('lightning,' from *fulgere*, 'to flash,') *Astrape*.

FULMICOTON, (*fulmen* and *cotton*,) see *Colloidon*.

FUL'NESS, *Reple'tio*, *Plen'itude*, *Pletho'ra*, *Reple'tion*. The state of being filled. Also, a feeling of weight or distension in the stomach or other part of the system.

FUMARIA, (from *fumus*, 'smoke,' from its smell,) *Fumaria officinalis* seu *med'ia*, *Fum'us ter'ræ*, *Cap'nos*, *Herba melancholif'uga*, *Sola'men Scabios'rum*, *Fumitory*, *Com'mon Fumitory*, *Fumiter'ra*, (F.) *Fumeterre*, *Fiel de terre*, *Pisse-sang*. *Ord.* *Fumariaceæ*. The leaves are extremely succulent, and have a bitter, somewhat saline, taste. The infusion of the dried leaves and the expressed juice of the fresh plant have been extolled for their property of clearing the skin of many disorders of the leprous kind.

FUMARIA BULBO'SA, *F. ca'va* seu *ma'jor*, *Borckhausen'ia* seu *Capno'i'des ca'va*, *Aristolochi'a fab'acea* seu *ca'va* seu *vulga'ris rotun'da*, *Coryd'al'is bulbo'sa* seu *ca'va* seu *tubero'sa*, *Capno'r'chis*. (F.) *Fumeterre bulbeuse*. The root of this plant was formerly given as an emmenagogue and anthelminthic.

FUMARIA CAVA, *F. bulbosa*—f. *Major*, *F. bulbosa*—f. *Media*, *Fumaria*—f. *Officinalis*, *Fumaria*.

FUMETERRE, *Fumaria*.

FUMIGATIO, (*fumigare*, *fumigatum*, 'to make smoke,'—*fumus*,) *Fumigation*—f. *Antiloeimica* *Gaubii*, see *Disinfection*—f. *Guytoniensis*, see *Disinfection*—f. *Smythiana*, see *Disinfection*.

FUMIGATION, *Fumiga'tio*, *Suffi'tus*, *Suffi'tio*, *Suffumina'tio*, *Suffumig'i'tum*, *Suffimen'tum*, *Apo-capn'is'mus*, *Thymia'ma*, *Epithymia'ma*, *Hypothymia'ma*, *Hypothymia'sis*, *Thymia'sis*, *Capn'is'ma*, *Hypocapn'is'mus*, *Hypocapn'is'ma*, *Hypat'mus*, *Hypat'mis'mus*, *Anathymia'sis*. An operation, the object of which is to fill a circumscribed space with gas or vapour, with the intention either of purifying the air, or of perfuming it, or of charging it with a substance proper for acting upon a part of the surface of the human body. See *Sulphurous acid*. Hence, *fumigations* have been dis-

tinguished into *aqueous*, *aromatic*, *sulphureous*, *mercurial*, *disinfecting*, *Guytonian*, &c. *Benzoin* generally constitutes the chief ingredient in the *Fumigating Pastilles*, to which any variety of odoriferous substances may be added. The following are formulæ.

R. *Benzoin. 3j; Cascarilla 3ss; Myrrh. ʒj; Ol. myrist. Ol. caryoph. aā gtt. x; Potassæ nitrat. 3ss; Carbon. lign. 3vj; Mucil. trag. q. s.*—or

R. *Benzoin. p. xvj; Balsam. Tolut. p. iv; Santal. alb. p. iv; Labdan. p. j; Carbon. Lign. p. xlvij; Potass. nitrat. p. ij; Tragacanth. p. j; Acaciæ p. ij; Ag. cinnam. p. xij.* Reduce the solid ingredients to powder, and mix the whole into a plastic mass, which must be formed into cones, flattened at the base, and dried, first in the air, and afterwards in a stove.

FUMIGATION, CHLORINE, see *Disinfection*—f. *Nitrous*, see *Disinfection*—f. *Oxymuratic*, see *Disinfection*.

FUMITERRA, *Fumaria*.

FUMITORY, *Fumaria*.

FUMUS ALBUS, *Hydrargyrum*—f. *Citrinus*, *Sulphur*—f. *Terræ*, *Fumaria*.

FUNAMBULA'TIO, (*funis*, 'a cord,' and *ambulare*, 'to walk,') An exercise with the ancients, which consisted in scaling ropes.

FUNCTION, *Func'tio*, *Act'io*, (F.) *Fonction*; (*fungor*, *functus*, 'to act,' 'perform,') The action of an organ or set of organs. Any act, necessary for accomplishing a vital phenomenon. A *function* is a special office in the animal economy, which has, as its instrument, an organ or apparatus of organs. Thus *respiration* is a function. Its object is the conversion of venous into arterial blood, and its instrument is the lungs. The ancient physiologists divided the functions into *vital*, *animal*, and *natural*. They called *vital functions* those which are essential to life, as innervation, circulation, respiration; *animal functions*, those which belong to the encephalon; viz.: the functions of the intellect, the affections of the mind, and the voluntary motions; and *natural functions*, *Facultat'es* seu *Actio'es natura'les*, those relating to assimilation, such as the actions of the abdominal viscera, of the absorbent and exhalant vessels, &c. Bichat divided the functions into those which relate to the preservation of the individual, and those that relate to the preservation of the species. The former he subdivided into *animal* and *organic*. The *animal functions* or *functions of relation* are those of the intellect, sensation, locomotion, and voice. The *organic functions* include digestion, absorption, respiration, circulation, secretion, nutrition, and calorification. The *functions*, whose object is the preservation of the species—the *organic, nutritive*, or *vegetative functions*—are all those that relate to generation;—such as conception, gestation, accouchement, &c. Each of these admits of numerous subdivisions in a complete course of *Physiology*;—for so the doctrine of the functions is called.

FUNCTIONAL, *Functiona'lis*, (F.) *Fonctionnel*. Relating or belonging to the function of an organ, or to the functions in general.

FUNCTIONAL DISEASES, see *Organic Diseases*.

FUNCTIONAL OR VITAL PHENOMENA are those produced by some modification in the action of an organ or organs, in contradistinction to those of a material or physical character;—thus we distinguish the *functional phenomena* of phthisis from those indicated by physical diagnosis—the *physical phenomena*.

FUN'DA, ('a sling,' *Sphen'donē*, (F.) *Fronde*. A bandage, composed of a fillet or long compress, cleft at its extremities to within about two inches of its middle. It is used in diseases of the nose and chin, and especially in cases of fracture of the lower jaw. In such case it has, also, been

called *Mentonnière*, because placed beneath the chin; from (F.) *Menton*, 'the chin.'

FUNDAMENT, (*fundare*, 'to found; to lay the *fundus* or bottom of anything.') Anus—f. Falling down of the, Proctocoele.

FUNDAMEN'TAL, (F.) *Fondamental*. Some anatomists have called the sacrum *Os Fundamentale*, because it seems to serve as a base to the vertebral column. The *sphenoid bone* has likewise been so denominated, from its being situate at the base of the cranium.

FUN'DUS, 'the bottom.' (F.) *Fond*. The base of any organ which ends in a neck, or has an external aperture, as the *Fundus vesicæ*, F. uteri, &c. Also, the *Vulva*.

FUNDUS VAGINÆ, *Laquear vaginæ*.

FUNES CORDIS, *Columnæ carnae*—f. Semicirculares, Semicircular canals—f. Ventriculi, see stomach.

FUNGIFORM PAPILLÆ, (*fungus*, 'a mushroom,' and *forma*, 'shape,') see *Papillæ* of the Tongue.

FUN'GOID, *Fungoides*, *Mycodes*, *Fungiformis*, *Fungiform*, (F.) *Fongoïde*, *Fongiforme*, (*fungus*, and *eidos*, 'resemblance,') That which has the shape of, or grows in some measure like a mushroom, as the *fungoid* or *fungiform papillæ* of the tongue.

FUNGOID DISEASE, Encephaloid.

FUNGOSITY, *Fungositas*, *Carbo luxurians* seu *fungo'sa*, *Ecsarco'ma*, *Hypersarco'ma*, *Hypersarco'sis*, *Proud Flesh*, (F.) *Fongosité*. The quality of that which is fungous:—fungous excrescence, *Excrescentia fungo'sa*. The fungosities which arise in wounds or ulcers are easily repressed by gentle compression, dry lint, the sulphas cupri, or other gentle caustics. At times, the more powerful are necessary, and sometimes excision is required.

FUN'GOUS, *Fungo'sus*, (F.) *Fongueux*. Having the characters of fungosity or fungus—as a *fungous wound*.

FUN'GUS, *My'ces*, (F.) *Fongus*, *Champignon*. The mushroom order of plants; class Cryptogamia, in the Linnæan system. The fungi or *champignons*, *Ord.* *Mucoraceæ*, (F.) *Mucédinées* ou *Mucorinées*, which are found as parasites in man, and the mammalia, are the following, according to M. Robin,—*Trichophyton tonsurans*, on the hairy scalp; *T. sporuloides*, on ulcers; *Microporum Audouini*, in the hair follicles; *M. mentagrophytes*, at the roots of the hair; *M. furfur*, in the skin; *Mucor mucedo*, in a cavity of gangrene of the lung; *Acho'ria Schönleini*, in the hairy scalp and hair follicles; *Aspergillus* species, [?] in the meatus auditorius externus; *Puccinia favi*, in favus; *Oidium albicans*, in aphthæ; and *Champignon du pommou*, in the lung.

In Pathology, Fungus is commonly used synonymously with fungosity, *myco'sis*. M. Breschet has proposed to restrict the term *fungosity* to vegetations which arise on denuded surfaces, and to apply the term *fungus* to the tumours which form in the substance of the textures, without any external ulceration. Fici and warts, for example, would be fungi of the skin.

FUNGUS ALBUS SALIGNEUS, *Dædalea suaveolens*—f. *Articuli*, *Spina ventosa*—f. Bleeding, *Hæmatodes fungus*—f. *Cancerosus hæmatodes*, *Hæmatodes F.*—f. *Cancerosus medullaris*, see Encephaloid—f. *Cerebralis*, see Encephaloid—f. *Cerebri*, Encephalocoele—f. *Chirurgorum*, *Lycoperdon*—f. *Cynosbati*, *Bedeguar*—f. *Faginosus*, *Morchella esculenta*—f. *Hæmatodes*, *Hæmatodes fungus*—f. *Ignarius*, *Boletus ignarius*—f. *Laricis*, *Boletus laricis*—f. of Malta, *Cynomorion coccineum*—f. *Medullaris*, see Encephaloid—f. *Melanoides*, *Cancer*, *melanotic*—f. *Melitensis*, *Cynomorion coccineum*—f. *Petræus marinus*, *Umbilicus marinus*—f. *Porriginis*, see *Porrigo favosa*—f.

Quercinus, *Boletus ignarius*—f. *Rosarum*, *Bedeguar*—f. *Salicis*, *Dædalea suaveolens*—f. *Sambucinus*, *Peiza auricula*.

FUNIC, *Funicus*, (F.) *Funique*, (from *funis*, 'a cord,') Relating or appertaining to the *Funis umbilicalis*.

FUNIC BELLOWS' SOUND or SOUFFLE, see *Bellows' Sound*, *funic*.

FUNICULI GRAC'ILES, (pl. of *Funiculus*,) (dim. of *funis*, 'a cord,') *Posterior Median Collumns* or *Fasciculi of the medulla oblongata*. Along the posterior border of each corpus testiforme, and separated from it by a groove, is a narrow white cord, separated from its fellow by the *fissura longitudinalis posterior*. The pair of cords are the *funiculi graciles*. Each funiculus forms an enlargement—*processus clavatus*—at its upper end, and is then lost in the corpus testiforme.

FUNICULI SILIQUÆ. Longitudinal fibres seen in the groove which separates the corpus olivare from the corpus pyramidale and corpus testiforme. They enclose the base of the corpus olivare,—those which lie on its inner side forming the *funiculus internus*; and those on its outer side the *funiculus externus*.

FUNICULUS, (dim. of *funis*,) Cord—f. Anterior (of the spinal marrow), see Nerve—f. Externus, see *Funiculi siliquæ*—f. Internus, see *Funiculi siliquæ*—f. Posterior (of the spinal marrow), see Nerve—f. *Spermaticus*, *Spermatic cord*—f. *Tympani*, *Chorda tympani*.

FUNICULUS UMBILICALIS, *Funis umbilicalis*, (dim. of *funis*, 'a cord,') *Intestinulum*, *Vinculum umbilicale*, *Omphaloneuron*, *Umbilical cord*, *Navelstring*. (F.) *Cordon ombilicale*. A cord-like substance, which extends from the placenta to the umbilicus of the fœtus. It is composed of the amnion, an albuminous secretion called the *Jelly of the Cord*, cellular substance, an umbilical vein, and two umbilical arteries. The former conveys the blood from the placenta to the fœtus—the latter return it. All these parts are surrounded by a sheath—*Investitura seu Vagina funiculi umbilicalis*. Its usual length is from 16 to 22 inches.

FUNICULUS VARICOSUS, *Cirsocoele*.

FUNIQUE, *Funic*.

FUNIS, Cord, *Laqueus*—f. *Argentæus*, *Medulla spinalis*—f. *Hippoceratis*, *Achillis tendo*—f. *Umbilicalis*, *Funiculus umbilicalis*.

FUNK, (from *fungus* [?]) *Boletus ignarius*.

FUNNEL, see *Infundibulum*.

FUNNYBONE. When the ulnar nerve is suddenly and forcibly pressed upon above the elbow, it gives occasion to a tingling in the fingers to which it is distributed. In familiar language this is said to be produced by pressing on the "*funny-bone*."

FUR, (low (L.) *furra*,) *Enduit*.

FURCELLA, *Furcula*; (dim. of *furca*, 'a fork,') The upper part of the sternum; the clavicle. The *Fourchette*.

FURCELLA INFERIOR, Xiphoid cartilage.

FURCHMÜHL, MINERAL WATERS OF. These Bavarian springs contain carbonic acid, sulphuretted hydrogen, carbonates of lime and soda; chlorides of lime and magnesium, oxides of iron and magnesium, &c.

FURCIFER, (*furca*, 'a fork,' and *ferre*, 'to carry,') Penis.

FURCILLA, (dim. of *furca*, 'a fork,') *Fourchette*.

FURCULA, (dim. of *furca*, 'a fork,') *Furcella*, *Clavicle*, *Fourchette*.

FUREUR UTERINE, ('uterine furor,') Nymphomania.

FUR'FUR, (from *far*, 'grain,') *Bras*, *Pit'yrion*, *Ach'yrion*, *Apobras'ma*, *Len'ma*, *Canta'brum*, (F.) *Son*. The decoction is sometimes employed as an emollient.

FURFURA, *Scarf*.

FURFURACEOUS, *Scurf'y, Conica'ceous*, *Pityr'ous*, *Pityr'oides*, *Pityr'o'ides*, (F.) *Furfuracé*. Resembling bran. A name given to eruptions, in which the epidermis is detached in small scales resembling bran. Also, a bran-like sediment observed at times in the urine;—*Urin'a furfuracea*, *Sedimentum Urinæ pityr'oides*.

FURFURAMIDE, see Furfurine.

FURFURATIO, Desquamation, Porrigo, Pityriasis.

FURFURIN, *Furfuri'na*. By the action of dilute sulphuric acid on corn meal or bran an oil is obtained—*furfuro'l* or *furfurole*. By the action of ammonia on this, *furfuramide* or *furfuro-lamide* results; and by the action of dilute potassa on this the alkaloid *furfurine* is obtained. This was found by Professor Simpson, of Edinburgh, to possess tonic, if not antiperiodic properties.

FURFURISCA, Pityriasis.

FURFUROL, see Furfurine.

FURFUROLAMIDE, see Furfurine.

FURIA INFERNALIS. A kind of vermiform insect, scarcely two lines long, common in Sweden, which flies about and stings both man and animals, exciting the most excruciating torture.

FURIBUNDUS, (*furere*, 'to be mad,') Manioides.

FURIOSUS, Manioides.

FURIOSUS, Manioides.

FURMAGE, ([F.] *fromage*,) Cheese.

FURMENTY, Frument.

FURNAS, MINERAL WATERS OF. A thermal chalybeate water in St. Michael's, Azores, which contains carbonic acid, and carbonate of iron.

FURONCLE, Furunculus.

FURONCLE GUÉPIER, *Wasp's nest furuncle* or *boil*. A malignant boil, which generally attacks the nape and region of the neck, and rarely others than old people. Hence it has been called *Old People's boil*.

FURONCULEUX, Furuncular.

FUROR, Mania—f. Brevis, Rage—f. Mania, Mania—f. Uterinus, Nymphomania.

FURRED, see *Enduit*.

FURROW, MENTOLABIAL, see Mentolabial furrow.

FURUN'CLAR, *Furun'culous*, *Furunculo'sus*, (F.) *Furunculeux*, (*furere*, 'to rage,' [?] or from *fur*, 'a thief'—see *Felon*.) Relating or appertaining to furunculus,—as 'a *furuncular* epidemic.'

FURUNCLE, Furunculus.

FURUNCULI ATONICI, Ecthyma—f. *Ventriculus*, see Furunculus.

FURUN'CULOID, *Furunculoï'des*, (*furunculus*, and *αἶος*, 'resemblance,') Resembling furunculus,—as 'a *furunculoïd* epidemic.'

FURUN'CLUS, *Chi'adus*, *Chi'oli*, *Doth'ien*, *Doth'ion*, *Furunculus suppurato'rius*, F. *Ve'rus*, F. *benig'nus*, *Phy'na furun'culus*, *Abscessus nucleatus*, *Furuncle*, *Boil*, *Bile*, (Sc.) *Bruick*, *Bruck*, (Prov.) *Pim'ceal*, (F.) *Furuncle*, *Froncle*, *Clou*. A small phlegmon, which appears under the form of a conical, hard, circumscribed tumour, having its seat in the dermal texture. At the end of an uncertain period, it becomes pointed, white or yellow, and gives exit to pus mixed with blood. When it breaks, a small, grayish, fibrous mass sometimes appears, which consists of dead areolar tissue. This is called the *Core*, *Set'fast*, *Sit'fast*, *Ventriculus* seu *Nucleus Furun'culi*, (F.) *Bourbillon*. The abscess does not heal until after its separation. The indications of treatment are,—to discuss by the application of leeches and warm fomentations;—or, if this cannot be done, to encourage suppuration by warm, emollient cataplasms. When suppuration is entirely established, the part may be opened or suffered to break, according to circumstances.

The term *blind boil* is given to an indolent, imperfectly suppurating, phlegmonous tumour, of the kind described above, which is often seated in a sebaceous follicle, as in *Acne indurata*.

FURUNCULUS GANGRENEOSUS, Anthrax—f. *Malignus*, Anthrax.

FUSARIA VERMICULARIS, (from *fusus*, 'a spindle,') *Ascaris vermicularis*—f. *Visceralis et renal*, *Strongylus gigas*.

FUSÉE PURLENTE (F.), (from *fusus*, 'a spindle,') The long and sinuous route which pus takes, in certain cases, in making its way to the surface. These *Fusées* almost always form beneath the skin between the muscles; or along aponeuroses, bones, tendons, &c.

FUSEL OIL, ([G.] *Fusel*, 'bad brandy,') see Oil, fusel.

FUSIBILITY, see Fusion.

FUSIBLE, see Fusion.

FUSION, *Fusio*, *Melt'ing*, *Liquefaction*; (*fundere*, *fusum*, 'to melt,') In chymistry, the transition of a solid body into a liquid by the aid of heat. Substances capable of such transition are said to be *fusi'ble*; or to be possessed of *fusi'bility*.

FUSTIC TREE, (from (F.) *fustet*.) *Cladastrix tinctoria*.

FUTUTIO, (*futuere*, 'to have sexual intercourse,') Coition.

FUTUTRIX, Tribas.

G.

THE Greek G, Γ, with the ancient Greek physicians, signified an ounce.

GAB, ([Da.] *gab*,) Mouth.

GABALLA, Cabal.

GABIR'EA, (*γαβίρσα*,) A fatty kind of myrrh, mentioned by Dioscorides.

GADUS ÆGLIFINUS, see *Oleum jecoris aselli*—g. *Callarias*, see *Oleum jecoris aselli*—g. *Carbonarius*, see *Oleum jecoris aselli*—g. *Lota*, see *Oleum jecoris aselli*—g. *Merluccius*, see *Oleum jecoris aselli*—g. *Morrhua*, see *Oleum jecoris aselli*—g. *Pollachius*, see *Oleum jecoris aselli*.

GÆEPHAGIA, Geophagism.

GÆEPHAGUS, Geophagist.

GÆOPHAGIA, (*γαια*, 'earth,' and *φαγειν*, 'to eat,') Geophagism.

GÆOPHAGUS, Geophagist.

GAGEL, Myrica gale.

GAHET, *Cigut*.

GAÏAC, Guaiacum.

GAILLET ACCROCHANT, *Galium aparine*—g. *Crochant*, *Galium aparine*—g. *Jaune*, *Galium verum*—g. *Vrai*, *Galium verum*.

GAÏNE, Vagina or sheath—g. *de l'Apophyse styloïde*, Vaginal process of the temporal bone—g. *de la Veine porte*, Vagina or sheath of the vena porta.

GAILTING, (dim. of (Sc.) *gat*, 'a child,') Infans.

GALA, (*γαλα*,) Milk.

GALACTACRASIA, (*galacto*, *a*, and *κρasis*, 'mixture.') A morbid mixture or constitution of the milk.

GALACTACRATIA, (*galacto*, *a*, and *κρατος*, 'power.') Galactia.

GALACTÆMIA, (*galacto*, and *αἷμα*, 'blood.') A condition of the blood in which it contains milk.

GALACTAGOGA, (*galacto*, and *αγω*, 'to drive.') Galactopœtica.

GALACTIAPOSTEMA, gen. *Galactapostematis*, (*galacto*, and *ἀποστήμα*, 'abscess.') Mastodynia apostematosa.

GALACTIA, *Galactorrhæa*, *Galactorrhæa*, *Lactis redundantia*, *Polygalactia*, *Galactozæmia*. A redundant flow of milk, either in a female who is suckling, or in one who is not. It may occur without being provoked by suckling. When to a great extent, it sometimes causes wasting; *Tibbes lactea*, *T. nutritum*. Dr. Good uses *Galactia*, in his Nosology, for 'morbid flow or deficiency of milk,' *Galactacratia*.

GALACTICUS, (*γαλακτικός*) Lactic.

GALACTIDROSIS, (*galacto*, and *ἰδρωσις*, 'sweating.') Sweating of milk.

GALACTIFER, (*galacto*, and *ferro*, 'to carry.') Galactophorous.

GALACTINE, Casein.

GALACTINUS, Lactic.

GALACTIRRHŒA, (*galacto*, and *ῥέω*, 'to flow.') Galactia.

GALACTIS, (*γαλακτίς*) Galaxias.

GALACTISCHESIS, *Galactoschesis*, *Lactis retentio*, (*galacto*, and *σχεῖν*, 'to restrain.') Retention or suppression of milk.

GALACTITES, (*γαλακτίτις*) Galaxias.

GALACTO, (*γαλα*, gen. *γαλακτός*) in composition, milk.

GALACTOCATARACTA, Cataract, milky.

GALACTOCELĒ, *Galactocèle*, *Galactoncus*, *Lactocelē*, (*galacto*, and *κῆλη*, 'a tumour.') *Tumour lacteus*. Distension of one or more of the galactophorous sinuses by milk. Lactiferous engorgement. The French use the term *Poil* for such engorgement; a relic of an idea, as old as Aristotle, that a hair swallowed may have obstructed one of the milk tubes, which must be got rid of by sucking.

The name has, also, been given by Vidal de Cassis to a tumour produced by an effusion of a white liquid into the tunica vaginalis, which he considered to be true milk.

GALACTODEN'DRON U'TILĒ, (*galacto*, and *δένδρον*, 'a tree.') *Palo de Vacca*, *Arbol de Leche*, *Cow tree* of South America. Family, *Urticaceæ*. When an incision is made into the trunk, a fluid similar to milk flows out, which is agreeable and nutritious.

GALACTODES, (*γαλακτώδης*, from *galacto*, and *οὐδεις*.) In Hippocrates, the term signifies milky, and likewise a milky colour, as of the urine—*urina galactodes*.

GALACTODIĒ'TA, *Diētā lactea* seu *lactis*, (*galacto*, and *διαιτα*, 'diet.') A milk diet.

GALACTODIARRHŒA, *Celiac Flux*.

GALACTOGANGLION, (*galacto*, and *γαγγλίον*, 'a knot.') Milk knot.

GALACTHÆMIA, *Galacthæmia*, *Galæmia*, (*galacto*, and *αἷμα*, 'blood.') *Lactis sanguinolenti* *Ecretio*. The secretion of bloody or bloodlike milk.

GALACTOID, *Galactoides*, (*γαλακτοειδής*, from *galacto*, and *εἶδος*, 'resemblance.') Resembling milk.

GALACTOMASTOPARTECTOMA, (*galacto*, *μαστος*, 'breast,' and *παρεκτείνειν*, 'to extend.') *Mastodynia apostematosa*.

GALACTOM'ETER, *Lactom'eter*, (*galacto*, and *μετρον*, 'measure.') An instrument for appreciating the quantity of cream in milk. It is a kind

of graduated separatory or *épreuve*—the degrees on the scale indicating the thickness of the layer of cream that forms on the surface of the milk.

GALACTOMYCES, (*galacto*, and *μυκης*, 'a fungus.') See *Encephaloid*.

GALACTONCUS, (*galacto*, and *ογκος*, 'a swelling.') *Galactocèle*.

GALACTOPH'AGOUS, (*γαλακτοφάγος*) *Galactoph'agus*, *Lactif'orus*, sometimes used substantively; (*galacto*, and *φαγω*, 'I eat.') That which feeds on milk. A name given to certain people, with whom milk appears to constitute the chief nourishment.

GALACTOPHORA, (*galacto*, and *φέρω*, 'I bear.') *Galactopœtica*.

GALACTOPHORITIS, (*galactophorous*, (ducts) and *itis*.) Inflammation of the galactophorous ducts. It has been inaccurately used for ulceration of the top of the nipple towards their orifices.

GALACTOPHOROUS, (*galacto*, and *φέρω*, 'to bear.') Lactiferous—g. Ducts, Lactiferous ducts—g. Receptacle, see Lactiferous vessels—g. Sac, see Lactiferous vessels—g. Sinuses, see Lactiferous vessels.

GALACTOPH'ORUS. Some accoucheurs have given this name to an instrument intended to facilitate sucking, when the faulty conformation of the nipple prevents the child from laying hold of it.

GALACTOPHYGUS, (*galacto*, and *φωγῆναι*, 'to shun.') That which arrests or disperses the secretion of milk. Hence *Galactoph'ygga Medicamenta*.

GALACTOPLA'NIA, *Metastasis* seu *Aberratio* seu *Vix extraordinariæ lactis*, *Galactorrhæa erronea*, (*galacto*, and *πλανν*, 'wandering.') Extravasation of milk into the areolar membrane.

Secretion of milk elsewhere than from the breasts.

GALACTOPLERO'SIS, (*galacto*, and *πληρωσις*, 'repletion.') Redundance of milk.

GALACTOPOEA, (*galacto*, and *ποιεω*, 'I make.') *Galactopœtica*.

GALACTOPOESIS, Galactosis.

GALACTOPOE'TICA, (*γαλακτοποιητική*) *Galactoph'ora*, *Galactog'ga*, *Galactopoe'a*. Substances to which has been attributed the property of favouring the secretion of milk and augmenting its quantity.

GALACTOPOIESIS, Galactosis.

GALACTOPO'SIA, (*γαλακτοποσία*, from *galacto*, and *ποσις*, 'drink.') The drinking of milk. Treatment of a disease by means of milk.

GALACTOP'OTES, (*γαλατοποτης*) *Galactop'otus*, *Lactip'otor*. A drinker of milk. One subjected to a milk diet.

GALACTOPYRA, (*galacto*, and *πυρ*, 'fire, fever.') Fever, milk.

GALACTOPYRETUS, (*πυρετος*, 'fever.') Fever, milk.

GALACTORRHŒA, (*galacto*, and *ῥέω*, 'to flow.') Galactia—g. Erronea, Galactoplasia—g. Saccharata, Saccharorrhœa lactea.

GALACTOSACCHARUM, Saccharum lactis.

GALACTOSCHESIS, Galactischesis.

GALACTO'SIS, *Galactopoe'isis*, *Galactopoe'isis*, *Secre'tio lactis*; (*γαλακτωσις*, from *galacto*, and *osis*.) The secretion or formation of milk.

GALACTOSPONGUS, (*galacto*, and *σπογγος*, 'a sponge, a fungus.') See *Encephaloid*.

GALACTOT'ROPHĒ, *Galactotroph'ia*, (*γαλακτοτροφία*, from *galacto*, and *τροφή*, 'nourishment.') Nourishment by means of milk.

GALACTOZE'MIA, (*galacto*, and *ζημια*, 'loss.') Loss of milk. Also, Galactia.

GALACTU'CHOS, *γαλακτουχος*, from *galactos*, and *χειν*, 'to have.') Suckling. Giving milk.

GALACTURIA, (*galacto*, and *ουρον*, 'urine.') Chyluria.

GALÆMIA, (*gala* and *aima*, 'blood.') Galactohæmia.

GALANGA, *Maranta galanga*.

GALANGAL, *Maranta galanga*.

GALANGALE, *Cyperus longus*.

GALARIPS, *Alliameda*.

GALARRHÆUS LATHYRIS, (*gala*, and *lew*, 'to flow.') *Euphorbia lathyris*—*g. Palustris*, *Euphorbia palustris*.

GALAX APHYLLA, (from *gala*.) *G. rotundifolia*, *Erythrorhiza rotundifolia*; *Car'penter's leaf*, *Beetweed*; indigenous; Order, *Ericaceæ*. The root is astringent, and the leaves, in domestic practice, are applied to cuts and wounds.

GALAXIA, (from *gala*.) Thoracic duct.

GALAX'IAS, (*γαλαξίας*.) *Galactites*, *Galac'tis*. A milk stone. A stone supposed to be capable of promoting the secretion of milk.

GALBANUM, (*γαλβαν*.) see *Bubon galbanum*—*g. long-leaved*, *Bubon galbanum*.

GAL'BULUS, (*galbus*, 'yellow.') A kind of congenital jaundice, in which the yellow colour continues through life. It is rather a defect in colour than a disease.

GALE, *Myrica gale*—*g. Scotch*, *Myrica gale*—*g. Sweet*, *Myrica gale*.

GALE, (from *galla*, 'a gallnut,' produced by the puncture of an insect. (?) *Psora*—*g. Canine*, *Psoriasis*—*g. Épidémique*, *Eczema*—*g. Miliàire*, *Psoriasis*—*g. Sèche*, *Lichen*, *Psoriasis*.

GALÉ ODORANT, *Myrica gale*.

GA'LEA. A helmet, (*γαλεα*, 'a cat,' of the skin of which it was formerly made.) A name given to the *amnios*, and also to the *bandage of Galen*. In *Pathology*, it indicates a headach affecting the whole head. See, also, *Caui*.

GALEA APONEUROTICA CAP'ITIS, *Ga'lea tendinea Santorini*, *Ga'lea cap'itis*, *Membrana epicrania*. The tendinous expansion which unites the frontal and occipital portions of the occipitofrontalis muscle.

GALEAMAUROSIS, (*γαλεα*, 'a cat,') *Amaurotic cat's eye*.

GALEAN'CON, *Galian'con*, (*γαλεα*, 'a cat,' 'a weasel,' and *αγκων*, 'an elbow.') *Mustela'neus*. One who has two short arms.

GALEAN'THROPY, *Galeanthropia*, (*γαλεα*, 'a cat,' and *ανθρωπος*, 'a man,') A variety of melancholy in which the patient believes himself changed into a cat. An affection similar to lycanthropy and cynanthropy.

GALE'GA, (from *gala*, which it increases (?) *G. officinalis* seu *vulgaris* seu *Persica*, *Ruta caprarua*, *Goat's Rue*, (F.) *Rue de chèvre*, *Faux Indigo*. Ord. *Leguminosæ*. It is slightly aromatic, and was once used as a sudorific and alexiterial in malignant fevers, &c.

GALEGA APOLLINEA, *Tephrosia apollinea*—*g. Persica*, *Galega*—*g. Purpurea*, *Tephrosia apollinea*—*g. Toxicaria*, *Tephrosia toxicaria*.

GALEGA VIRGINIA'NA, *Tephrosia virginiana*, *Turnkey Pea*, *Hoar'y Pea*, *Devil's shoe'strings*, *Virginia Goat's rue* or *catgut*, is used in some parts of the United States as an anthelmintic. The decoction of the root is given.

GALEGA VULGARIS, *Galega*.

GALENE, *Graphites*.

GALENEA, *Graphites*.

GALEN'IC, *Galen'ical*, *Gale'nicus*, *Gale'nus*, (F.) *Galénique*, (after *Galen*.) That which relates to the doctrine of Galen or to Galenism. Used, substantively, for drugs that are not chymical.

GALENIC MEDICINE, *Galenism*.

GA'LENISM, *Galen'ic medicine*. The doctrine of Galen.

GA'LENIST, *Galenis'ta*, *Galenis'tes*. A follower of the doctrine of Galen.

GALENIUS, *Galenic*.

GALEOB'DOLON, (*γαλεόβολον*, from *γαλεα*, 'a weasel,' and *βύλλω*, 'to emit a stench.') *Galeopsis*.

GALEOP'DOLON, *Galeopsis*.

GALEOP'SIDE, *Galeopsis grandiflora*.

GALEOPSIS, *Lamium album*.

GALEOP'SIS, *Galiop'sis*, *Galeob'dolon*, *Galeop'dolon*, *La'mium rubrum*, *Urtica in'ers magna fetidis'sima*, *Sta'chys fetida*, *Hedgenettle*, (F.) *Ortie morte des bois*. Ord. *Labiatae*. (*γαλεα*, 'a weasel,' and *οψις*, 'appearance,' the corolla having been supposed to resemble the head of the weasel.) This plant was formerly reckoned a vulnerary and anodyne.

GALEOPSIS ANGUSTIFOLIA, *G. grandiflora*—*g. Dubia*, *G. grandiflora*.

GALEOPSIS GRANDIFLO'RA, *G. Ochroleuca* seu *la'danum* seu *angustifolia* seu *du'bia* seu *prostrata* seu *villo'sa*, *Tetrahit longiflorum*, *G. Seg'letum*, *Her'ba Sideritidis*, (F.) *Galeopside*, *Chanvre bâlard*. This plant is regarded in Germany as a 'bitter resolvent.' It is the basis, also, of a celebrated nostrum, the *Blankenheimer Tea*, called likewise *Lieber's pectoral* and *phthisical herbs* (*Liebersche Brust oder Auszehrungs-Kräuter*), which has enjoyed great repute in pectoral complaints. The tops of the plant are given in decoction (3j), boiled in a pint of water for a quarter of an hour.) This quantity to be taken in a day.

GALEOPSIS LADANUM, *G. grandiflora*—*g. Ochroleuca*, *G. grandiflora*—*g. Prostrata*, *G. grandiflora*—*g. Segetum*, *G. grandiflora*.

GALEOPSIS VERSICOLOR, is possessed of the same virtues.

GALEOPSIS VILLOSA, *G. grandiflora*.

GALER'OPIA, (*γαλεος*, 'bright,' and *οψις*, 'vision,') *Oxyopia*.

GALEX, (from *galle*, 'itch,') *Psoric*.

GA'LIA, (from *galle*, 'galls,') An ancient composition, in which galls were an ingredient,—the *Ga'lia pu'ra*. There was also a *Ga'lia aromat'ica*, *moscha'ta* seu *musca'ta*, which consisted of a mixture of several perfumes, such as musk.

GALIANCON, (*γαλιαγκων*.) *Galeancon*.

GALIOPSIS, *Galeopsis*.

GALIPEA FEBRIFUGA, (*G.* is its name in Guiana.) *Cusparia febrifuga*—*g. officinalis*, see *Cusparia febrifuga*.

GALIPOT, see *Pinus sylvestris*.

GA'LIIUM, *Gall'ium*, (*γαλιον*, from *γαλα*, 'milk,' because some species curdle milk.) *G. verum*—*g. Album*, *G. Mollugo*.

GALIUM APARINOIDES, *G. aparine*.

GALIUM APARI'NE, *G. infestum* seu *aparinoides* seu *brachycarp'on* seu *scaber'rimum*, *Valantia aparine*, *Aparine*, *A. his'pida*, *Lap'pa*, *Philanthrop'us*, *Ampelocar'pus*, *Omphalocar'pus*, *Asphari'ne*, *Asper'ula*, *Goosegrass*, *Cleaver's bees*, *Cleaver's*, *Gooses'hare*, *Robin run-the-hedge*, *Hay-riff*, (Old Eng.) *Clith'ren*, (Prov.) *Clider*, *Oliver*; Order, *Rubiaceæ*; *Sex. Syst.* *Tetrandria Monogynia*. (F.) *Gailliet acrochant*, *G. crochant*, *Gat'teron*. The expressed juice has been given as an aperient diuretic in incipient dropsies; also in cancer; and the decoction in *Lepra* and *Psoriasis*.

GALIUM ASPRELL'UM, *Rough bedstraw*, *Rough ladies' bedstraw*; indigenous; has the diuretic properties of most of its genus.

GALIUM BRACHYCARPON, *G. aparine*—*g. Caucasicum*, *G. verum*.

GALIUM CIRCE'ZANS, *Wild Liquorice*, *Master of the Woods*. An indigenous plant, which flowers from June to August. It is demulcent and diuretic, and is a popular domestic remedy.

GALIUM INFESTE, *G. aparine*—*g. Luteum*, *G. verum*.

GALIUM MOLLU'GO, *Gall'ium album* seu *Tyrolen'se*, *Alyssum Plin'tii*, *Great'or ladies' bedstraw*, (F.) *Caillélat blanc*. The herb and flowers have been used medicinally in epilepsy.

GALIUM ODORATUM, *Asperula odorata*—g. *Scaeverrimum*, *G. aparine*.

GALIUM TINCTO'RUM, an American species, closely allied in properties to *G. verum*.

GALIUM TUBERCULATUM, *G. verum*—g. *Tyrolense*, *G. mollugo*.

GALIUM VE'RUM, *Gal'ium*, *G. lu'teum* seu *Caucas'icum* seu *tuberculatum*, *La'dies'* bedstraw, *Cheeserennet*, *Bedstraw*, *Cleave'wort*, *Goosegrass*, *Savoyan*, *Clubbergrass*, *Milk'sweet*, *Poor Rob'in*, *Grav'el-grass*, (F.) *Gaillet jaune ou vrai*, *Vrai Caillelait*. The tops were used in the cure of epilepsy. The leaves and flowers possess the property of surling milk.

GALL, (Sax. *zēala*.) Bile, see Chafing, and Eczema impetiginodes, *Quercus infectoria*, and Vitrum—g. of the Earth, *Prenanthes*, *P. alba*—g. Nut, see *Quercus infectoria*—g. of the Ox, see Bile—g. of the Skin, Chafing.

GALL, TURKEY, see *Quercus infectoria*.

GALLBLADDER, *Vesic'ula fellis*, *Chol'ecyst*, *Cholecyst'is*, *Follie'ulus fell'is*, *Cyst'is fel'lea* seu *choled'ochus*, *Vesi'ca fel'lea* seu *bili'aria*, *Vesic'ula bil'is*, *Follie'ulus fel'leus*, (F.) *Vésicule du fiel* ou *biliaire*, *Réservoir de la bile*. A membranous, pyriform reservoir, lodged in a superficial depression at the inferior surface of the right lobe of the liver, *Fos'sa of the Gallbladder*, (F.) *Fosse de la Vésicule biliaire*. It receives, by the hepatic and cystic ducts, a portion of the bile secreted by the liver, when the stomach is empty, which becomes in it more acrid, bitter, and thick. It receives an artery, called the *cystic*. Its veins empty into the vena porta. Its nerves come from the hepatic plexus, and its lymphatic vessels join those of the liver.

GALLA, see *Quercus infectoria*—g. *Maxima Orbiculata*, see *Quercus infectoria*.

GALLÆ QUERCÛS, ('galls of the oak,') see *Quercus infectoria*—g. *Tinctoriæ*, see *Quercus infectoria*—g. *Turciæ*, see *Quercus infectoria*.

GALLATURA, (from *gallus*, 'a cock,') Mole-cule.

GALLE DE CHÊNE, ('gall of the oak,') see *Quercus infectoria*.

GALLI GALLINACEI CAPUT, *Gallinaginis caput*.

GALLINAG'INIS CA'PUT, *Gal'li gallina'cei* *Ca'put*, *Ca'put gallina'ceum*, *Collie'ulus semina'lis*, *Carunc'ula semina'lis*, *Ve'ru mont'uum*, *Cris'ta urethra'lis* seu *ure'thræ*, *Crête uréthrale*—(Ch.,) (*gallinago*, gen. *gallinaginis*, 'a woodcock,') An oblong, rounded projection, formed by the mucous membrane in the prostatic portion of the urethra, on the sides of which the ejaculatory ducts open.

GAL'LIPOT, (Prov.) *Glumpet*. (*gala*, 'finery.' [?]) A pot painted and glazed, or merely glazed, and commonly used to hold medicines.

GALLITRICHUM, (*Callitrichum*,) *Salvia sclarea*.

GALLIUM, *Galium*.

GALLS, see *Quercus infectoria*—g. *Bussorah*, see *Quercus infectoria*—g. *Mecca*, see *Quercus infectoria*.

GALLSICKNESS, Fever, Walcheren.

GALLSTONES, *Calculi*, biliary.

GALLUS, (*Galli*, priests of Cybele, who castrated themselves.) Eunuch.

GALREDA, ((G.) *Gallerte*, 'jelly,' [?]) Gelatin.

GALVANIA VELLOZII, (after Galvao, a Portuguese naturalist,) *Palicourea Mægrega* v. *ii*.

GALVANIC CAUTERY, see Moxa.

GALVANISATION, *Galvanization*.

GALVANISM, *Galvani's'mus*, *Electric'itas anim'alis* seu *Galvan'ica* seu *metal'lica*, *Irritamen'tum metallo'rum* seu *metal'licum*, *Vol'taism*, *Vol'ta'ic* or *Chem'ical* or *Con'tact Electricity*. (After Galvani.) A series of phenomena, consisting in

sensible movements, executed by animal parts, which are endowed with irritability, when placed in connexion with two metallic plates of different nature, between which a communication is established by direct contact or by means of a metallic wire. Galvanism has been employed medicinally in the same cases as electricity, and especially in neuralgic affections. It has been applied in the form of plates—'Mans'ford's plates.' In asthma, for example, a small blister, the size of a dollar, may be placed on the neck over the course of the phrenic and pneumogastric nerves, and another on the side, in the region of the diaphragm. One metal is placed mediately or immediately over the vesicated surface on the neck, and another over that in the side. They are then connected by means of a wire. The new nervous impression, in this way induced, is often signally beneficial.

Galvanic chains have been devised for the same purpose. The one most in use, called, after its inventor, 'Pulvermacher's Hydroelectric Chain Battery,' is a modification of the voltaic pile, capable of being employed topically.

GALVANIZATION, *Galvanisatio*, (F.) *Galvanisation*. The medical application of the current of galvanic electricity. The act of affecting with galvanism.

GALVANIZATION, LO'CALIZED, *Galvanism, lo'calized*, (F.) *Galvanismus localisēe*. A mode of employing galvanism, proposed by Duchenne, of Boulogne, by which, he conceives, the electric influence may be arrested, at will, in the skin: or, without any incision or puncture, the skin may be traversed, and the electric influence be limited to the organs which it covers—to the nerves, muscles, and even the bones. See *Electrization*, *localized*.

GALVANOPUNCTURE, *Electropuncture*.

GAMBA, *Patella*.

GAMBARUS, (*καμπαρος*,) Crab.

GAMBIER, see *Catechu* and *Nauclea gambir*.

GAMBIR, see *Catechu* and *Nauclea gambir*.

GAMBOGIA, *Cambogia*.

GAMBOIDIA, *Cambogia*.

GAMMARUS, (*καμαρα*, 'an arch,') Crab.

GAM'MATA FERRAMENTA, *Cauteries*, having the shape of the Greek letter *Γ*, *gamma*, which were used for cauterizing hernie.

GAMMAUT, *Gammot*. The Italians, according to Scultetus, gave this name to a kind of crooked bistouri, used for opening abscesses.

GAMMISMUS, *Psammismus*.

GAMPHE, (*γαμφη*,) Gena, Maxillary Bone.

GAMPHELE, (*γαμφήλη*,) Gena, Maxillary Bone.

GANCARDE, *MIN'ERAL WA'TERS OF*. Simple sulphurous springs in the department of Landes, France. The season is from the 1st of May to the 15th of September.

GANGAME, (*γαγγαμή*, 'a drag net,') Epiploon.

GANGAMUM, (*γαγγαμιν*,) Epiploon.

GANGLIA CEREBRI POSTICA, (pl. of *ganglion*,) ('posterior ganglia of the brain,') *Thalami nervorum optico'rum*—g. Formative, see *Ganglion*—g. Hemispherical, *Hemispheres of the brain*—g. of Increase, see *Ganglion*—g. Nervorum, *Ganglions*, nervous, see *Ganglion*—g. Respiratory, see *Encephalon*—g. Sensory, see *Sensory ganglia*—g. Stomato-gastric, see *Encephalon*.

GANGLIAR, *Ganglionic*.

GANGLIATED, *Ganglionic*.

GANG'LIFORM, (*ganglion*, and *forma*, 'form,') *Gang'liform*, *Gangliform'is*. Having the shape of a ganglion.

GANGLIITIS, (*ganglion* and *itis*,) *Ganglionitis*.

GANGLI'OLUM, (dim. of *ganglion*.) A small ganglion.

GANGLIO'MA, (*ganglion* and *oma*.) A tumour

of the glands or of the lymphatic ganglions; and especially epithelioma of them.

GANG'LION, (*γᾱγγλῖον*, 'a knot,') *Gang'lion*. In anatomy, a name generally given to a knot-like enlargement in the course of a nerve. It is applied, however, to organs differing considerably from each other in size, colour, texture, functions, &c. They are divided into *gland'iform*, *lymphatic*, and *ner'vous*. 1. *Gland'iform gang'lions*, called also *adenoid*, *vas'cular*, and *sanguineous gang'lions*; blind, *apor'ic*, *duct'less*, *blood*, and *vas'cular glands*.—*gland'ulæ vasculosæ*; *gland'ulæ spur'ias*, &c., are organs of whose functions we are, in general, ignorant; and which have the appearance of glands. They are formed of agglomerated globules, pervaded by blood-vessels surrounded by areolar membrane, and contain a milky or yellowish fluid. To this class belong the spleen, thymus, thyroid, and supra-renal glands. They are, doubtless, concerned in sanguification. 2. *Lymphatic gang'lions*. See Conglobate. 3. *Ner'vous gang'lions*, *Gang'lia* seu *Ganglio'nes* seu *No'di* seu *No'duli* *Nervo'rum*, *Tumores* seu *Plexus gangli'iform'es*, *Plexus glandi'form'es*, *Tuber'cula nodosa Nervo'rum*, *Divertic'ula spirituum anima'lium*, *Gang'lia of increase*, *Form'ative gang'lia*. Enlargements or knots in the course of a nerve. They belong, in general, to the system of the great sympathetic. One exists on the posterior root of every spinal nerve, and on one cerebral,—the 5th. Bichat regarded them as so many small brains, or centres of nervous action, independent of the encephalon, and intended exclusively for organic life. Being formed by the union of the cerebral and spinal nerves, they may send out the influence of both these nervous centres to the parts to which the nerves proceeding from them are distributed. Ganglia are chiefly composed of vesicular neurine; and appear to be concerned in the formation and dispensation of nerve power.

GANGLION, *Emphy'ma encyst'is gang'lion*. A globular, hard, indolent tumour, without change in the colour of the skin; of a size varying from that of a pea, to that of an egg, and always situate in the course of a tendon. The tumour is formed of a viscid, albuminous fluid, contained in a cyst of greater or less thickness. The cyst is sometimes loose; but in the majority of cases it communicates, by a narrow footstalk, with the sheath of a tendon, or even with the synovial capsule of a neighbouring articulation. The causes are generally unknown. The treatment consists in compression, percussion, the use of discutients, extirpation, or incision.

GANGLION ABDOMINALE, *G. semilunare*—*G. Adenoid*, *G. glandiform*—*G. of Andersch*, Petrous ganglion—*G. Annular*, see Ciliary ligament—*G. of Arnold*, Oticum ganglion—*G. Auricular*, Oticum *G.*—*G. Azygous*, see Trisplanchnic nerve—*G. Cardiac*, Cardiac ganglion—*G. Carotic* or *Carotid*, see Carotid or Carotic nerve—*G. Cerebelli*, Corpus dentatum—*G. Cerebral*, anterior, Corpora striata—*G. Cerebral, inf'erieur*, *grand*, Thalami nervorum optico'rum—*G. Cerebral*, posterior, Thalami nervorum optico'rum—*G. Cerebri Anterior*, Corpora striata—*G. du Cervelet*, Corpus dentatum—*G. Ciliare*, Corpus dentatum—*G. Ciliary*, Ophthalmic ganglion—*G. Corpuseles*, seu *Neurine*.

GANGLION OF EHRENNITTER, *Gang'lion ner'vi glossopharynge'i super'ius*, *G. jugula'rè super'ius* seu *Ehrenritteri* seu *Mulleri*. A reddish-gray mass on the glossopharyngeal nerve in the foramen lacerum, above the ganglion of Andersch.

GANGLION OF THE FIFTH NERVE, *G. of Gasser*.

GANGLION OF GAS'SER, *Gan'glium* seu *Gan'glio* *Gasseri* seu *Gasserianum* seu *semilunare*, *Moles gangli'form'is*, *Intumescen'tia gangli'form'is* seu *semilunaris*, *Ten'ia nervosa Halleri*, *Gang'lion of the fifth nerve*. A semicircular knot on the

5th pair of nerves, before its division into three branches.

GANGLION, **GLANDIFORM**, see Ganglion—*G. Globules*, see Neurine—*G. Impar*, see Trisplanchnic nerve—*G. Jugulare superius*, *G. of Ehrenritter*—*G. Laitenx*, Milk-knot—*G. of Laumonier*, see Carotid or Carotic nerve—*G. Lenticular*, Ophthalmic *G.*—*G. Lymphatic*, Conglobate gland—*G. Muscili'tympanique*, Oticum *G.*—*G. of Meckel*, Sphenopalatine *G.*—*G. Mulleri*, *G. of Ehrenritter*—*G. Nasopalatine*, see Nasopalatine ganglion—*G. Nervi glossopharynge'i superius*, *G. of Ehrenritter*—*G. Nervous*, *Neuroma*—*G. Ophthalmic*, see Ophthalmic ganglion—*G. Optic*, Quadrigemina tubercula—*G. Orbital*, *G. ophthalmic*—*G. Oticum*, *Otic G.*—*G. Petrosal*, see Petrous ganglion.

GANGLION OF THE PNEUMOGASTRIC. A ganglionic structure in the pneumogastric as it passes through the foramen lacerum posterius.

GANGLION OF RIBES. A nervous ganglion upon the anterior communicating artery of the brain, and to be found at the point of junction of the right and left trunks of the sympathetic.

GANGLION, SANGUINEOUS, *G. glandiform*—*G. Semilunare*, *G. of Gasser*, Ophthalmic ganglion—*G. Sensory*, see Sensory ganglia—*G. Solare*, *G. semilunare*—*G. Sphenoidal*, Sphenopalatine ganglion—*G. Spinal*, see Encephalon—*G. Splanchnicum*, *G. semilunare*—*G. Submaxillary*, see Submaxillary—*G. Sup'erieur du cerveau (grand)*, Corpora striata—*G. of the Superior Laryngeal Branch*, see Pneumogastric nerves—*G. Surr'inal*, *G. semilunare*—*G. Thyroid*, see Trisplanchnic nerve—*G. Transversum*, *G. semilunare*—*G. Vascular*, *G. glandiform*—*G. Vertebral*, see Trisplanchnic nerve—*G. de Vieussens*, Celiac plexus.

GANGLIONARY, Ganglionic.

GANGLIONES NERVORUM, Ganglions, nervous.

GANGLION'IC, *Ganglion'icus*, *Gang'lionary*, *Gang'liar*, *Gang'liated*, from *ganglion*, (*F.*) *Ganglionnaire*. Relating to ganglia. Nerves are so called in the course of which ganglions are met with; as the greater part of the branches of the great sympathetic or trisplanchnic, the posterior roots of the spinal nerves, &c. *Ganglion'ics*, according to Dr. Pereira, are agents, which affect the ganglionic or great sympathetic system of nerves (?),—as stimulants and sedatives.

GANGLIONIC NERVE, Trisplanchnic nerve—*G. Nervous System*, see Trisplanchnic nerve.

GANGLIONI'TIS, *Ganglii'tis*, (*ganglion*, and *itis*.) Inflammation of a nervous ganglion. Sometimes used for inflammation of a lymphatic ganglion.

GANGLIONITIS PERIPHERICA et MEDULLARIS, Cholera.

GANGLIONNAIRE, Ganglionic.

GANGLIUM, Ganglion—*G. Gasseri*, Ganglion of Gasser.

GANGRÆNA, (*γᾱγγραινα*, from *γᾱω*, *γᾱρω*, 'to gnaw, to eat,') Gangrene—*G. Alopecia*, Alopecia—*G. Caries*, Caries—*G. Nosocomialis*, Hospital gangrene—*G. Nosocomiorum*, Hospital gangrene—*G. Oris*, Cancer aquaticus—*G. Ossis*, Spina ventosa—*G. Ossium*, Caries—*G. Pottii*, see Gangrene—*G. Pulmonum*, Necropneumonia—*G. Senilis*, Gangrene of old people, see Gangrene—*G. Sphacelus*, Sphacelus—*G. Tonsillarum*, Cynanche maligna—*G. Ustilaginea*, Ergotism—*G. Vaginae*, Colpoceae.

GANGRÆNESCENTIA, Gangrenosis.

GANGRÆNICUS, (*γᾱγγραινός*.) Gangrenous.

GANGRÆNODES, (*gangræna*, and *odes*.) Gangrenous.

GANGRÆNOP'SIS, (*gangræna*, and *osis*, 'the countenance,') Cancer aquaticus; also, gangrenous inflammation of the eyelids; *Blephari'tis gangrænosa*.

GANGRÆNO'SIS, (*γᾱγγραινός*, from *gangræna*,

and *osis*, *Gangraenosis*. The state of being gangrenous or of becoming gangrenous.

GANGRENIOSUS, Gangrenous.

GANGRENE, *Gangra'na*, *Gangra'na*, *Hot mortification*, (F.) *Gangrène*, *G. Chaud*, *Asphyxie des parties*. Privation of life or partial death of an organ. Authors have generally distinguished mortification into two stages; naming the first *incipient mortification* or *gangrene*. It is attended with a sudden diminution of feeling in the part affected; livid discoloration; detachment of the cuticle, under which a turbid fluid is effused; with crepitation, owing to the disengagement of air into the areolar texture. When the part has become quite black, and incapable of all feeling, circulation, and life, it constitutes the *second stage*, or *mortification*, and is called *phacelus*. Gangrene, however, is frequently used synonymously with mortification.—*local asphyxia* being the term employed for that condition, in which the parts are in a state of suspended animation, and, consequently, susceptible of resuscitation. When the part is filled with fluid entering into putrefaction, the affection is called *humid gangrene*, (F.) *Gangrène humide*; on the other hand, when it is dry and shrivelled, it constitutes *dry gangrene*; (F.) *Gangrène sèche*. To this class belongs the *gangra'na senilis*, *G. Poltii*, *Presby-phacelus*, or *spontaneous gangrene* of old people, which rarely admits of cure. Whatever may be the kind of gangrene, it may be caused by violent inflammation, contusion, a burn, congelation, the ligature of a large arterial trunk, or by some inappreciable internal cause.

The treatment, both of external and internal gangrene, varies according to the causes which produce it. Gangrene from excessive inflammation is obviated by antiphlogistics; and that from intense cold by cautiously restoring the circulation by cold frictions, &c. When the gangrene has become developed, the separation of the eschars must be encouraged by emollient applications, if there be considerable reaction; or by tonics and stimulants, if the reaction be insufficient.

GANGRENE, DRY, see Gangrene—g. Hospital, see Hospital gangrene—g. Humid, see Gangrene—g. of the Lungs, Necropneumonia—g. Spontaneous, see Gangrene.

GANGRÈNE DE LA BOUCHE, Cancer aquaticus—g. *Chaud*, Gangrene—g. *Froide*, *Phacelus*—g. *Hôpital*, Hospital gangrene—g. *Humide*, see Gangrene—g. *du Poupon*, Necropneumonia—g. *Sèche*, see Gangrene—g. *des Solénois*, Ergotism.

GANGRÉNEUX, Gangrenous.

GANGRENOUS, *Gangra'nicus*, *Gangra'neus*, *Gangra'neus*, (F.) *Gangréneux*. Affected with or relating to gangrene.

GANGRENOUS EROSION OF THE CHEEK, Cancer aquaticus.

GANJAH, Gunjah.

GANNAL'S SOLUTION, see Aluminae Acetas.

GANTELET (F.), *Chirotheca*, *Fascia digitalis*, *Gant'let*; (from (F.) *gant*, 'a glove.') A sort of bandage which envelops the hand and fingers like a glove. It is made with a long roller, about an inch broad; and is applied so that the fingers are covered to their tips, when it is called *Gantelet entier* or *complet*. The *Demigantelet* includes only the hand and base of the fingers. Both bandages are used in fractures and luxations of the fingers, burns of the hand, &c. See *Chirotheca*.

GANTS DES DAMES, ('gloves of the ladies,') Condom—g. *de Notre Dame*, Digitalis.

GAOL CACHEXIA, see Cachexia, gaol.

GAPING, (Sax. *gæpan*, 'to gape.') Yawning.

GARANÇE, Rubia.

GARCINIA CAMBOGIA, *G. seu Cambogia* *gutta*, *Mangostina Cambogia*. Ord. Guttiferae. (after Dr. L. Garcin, an English traveller, who

accurately described it.) A tree of Ceylon, Family, Guttifera, which affords a concrete juice similar to Gamboge.

GARCINIA GUTTA, G. Cambogia.

GARCINIA MANGOSTA'NA, *Mangostina Gascintia*. The systematic name of the *Mangostin*, *Mangostan*, *Mangostine* or *Mangosteen* tree, *Mangostana*. It grows in great abundance in Java and the Molucca islands. The fruit, which is about the size of an orange, is delicious, and is eaten in almost every disorder. The dried bark is used medicinally in dysentery and tenesmus; and a strong decoction has been much esteemed in ulcerated sore throat.

GARCINIA PURPUREA. From this species a concrete oil of *Mangosteen* is obtained, which is called, in India, *Ko'kum butter*, *Kokum* being the Indian name of Mangosteen. The oil is extracted from the fruit by boiling. It is used in India in cases of chaps.

GARÇON, old (F.) *gars*, 'a lad.' Boy.

GARDE-MALADE, (*garder*, 'to take care of,' and *malade*, 'a sick person.') Nurse.

GARDENIA DUMETORUM, (after Dr. Alex. Garden, of Carolina.) *Randia dumetorum*—g. *Spinosa*, *G. dumetorum*.

GARDEROBE, *Artemisia abrotanu* u.

GARDEROBES, *Favos*.

GARDINER'S ALIMENTARY PREPARATION, see Oryza.

GARDOUCHES, *Vesiculæ seminales*.

GARETUM, (low [L.]) *Poples*.

GARGALISMUS, *Gargalus*.

GARGALUS, (*γάργας*, from *γάργαιος*, 'to tickle,') *Gargalis*, *Gargalis'mus*, *Titillatio*, *Irritatio*, *Pruritus*. Titillation, irritation, itching. Also, masturbation; and, rarely, animal magnetism.

GARGAREON, (*γάργαραν*) *Uvula*.

GARGARISATIO, *Gargarism*.

GARGARISM, *Gargaris'mus*, *Gargaris'ma*, *Anagargalis'ton*, *Gargaris'mum*, *Collut'io*, *Dialys'ma*, *Anagargaris'ton*, *Anagargaris'mus*, *Anaconchylis'mus*, *Anaconchylis'mus*, *Titillat'io*, *Irrit'io*, (from *γάργαιος*, 'I wash the mouth,' 'I gargle, formed by onomatopœia.) A gargle. Any liquid medicine intended to be retained in the mouth, for a certain time, and to be thrown in contact with the uvula, velum pendulum, tonsils, &c. For this purpose, the liquid is agitated by the air issuing from the larynx, the head being thrown back. Gargles are employed in cynanche tonsillar and other diseases of the fauces, and are made of stimulants, sedatives, astringents, refrigerants, &c., according to circumstances.

The process is termed *garg'ling*, *gargarisatio*. The term *collut'io* or *collut'io* *oris* is generally restricted to a wash for the mouth.

GARGET, *Phytolacca decandra*.

GARGLE, *Gargarism*.

GARGLING, see *Gargarism*.

GARGUILLEMENT, (formed, like *Gargarism*, by onomatopœia.) Borborygms, Gurgling. See *Râle muqueux*.

GARLIC, (Sax. *garleac*.) *Allium*—g. *Hedge*, *Alliaria*—g. *Meadow*, *Allium Canadense*.

GARLOCK, *Sinapis arvensis*.

GAROSMUM, *Chenopodium vulvaria*.

GAROU BARK, *Daphne gnidium*.

GARRETUM, *Poples*.

GARROPHYLLUS, *Eugenia caryophyllata*.

GARROT, (F.) *Garrot*, (*garrotter*, 'to tie fast,') A compressing bandage, tightened by twisting a small cylinder of wood, by which the arteries of a limb are compressed, for the purpose of suspending the flow of blood in cases of hemorrhage, aneurism, amputation, &c.

GARROTILL, Cynanche maligna.

GARRULITAS, (*garric*, 'to chatter.') Loquacity.

GARULEUM BIPINNA'TUM, (corr. from *cœruleum*, 'blue'—the flowers.) A South African plant, *Ord. Composite*; known under the name *Snake-root*, from its reputed effects as an antidote to the bites of venomous serpents. The root is a great favourite with the Boers in chest diseases—as asthma—and in affections in which a free secretion from the mucous membrane of the bronchia is indicated. It has diaphoretic properties, and acts as a diuretic in gout and dropsy. It is given in decoction or tincture. Dr. Pappe thinks the root ought to have a place in the *Materia Medica*.

GARUM, (*γαρον*.) *Licui'men*, *Liqua'men*. The ancient Romans gave this name to a kind of pickle made by collecting the liquor which flowed from a salted and half-putrefied fish, (*γapos*.) It was used as a condiment.

GARVANCE, *Cicer arietinum*.

GAS, see *Gaz*—*g.* Ammoniacale, Ammonia—*g.* Animale sanguinis, *Gaz sanguinis*—*g.* Azoticum, Azote—*g.* Azoticum oxygenatum, Nitrogen, gaseous oxide of—*g.* Hepaticum, Hydrogen, sulphuretted—*g.* Hydrogenium sulphuretum, Hydrogen, sulphuretted—*g.* Intoxicating, Nitrogen, gaseous oxide of—*g.* Laughing, Nitrogen, gaseous oxide of—*g.* of the Lungs, *Gaz*, pulmonary—*g.* Nitrous, dephlogisticated, Nitrogen, gaseous oxide of—*g.* Oxygenated muriatic acid, Chlorine—*g.* Oxymuriatic acid, Chlorine—*g.* Palustre, Miasm, marsh—*g.* Paradise, Nitrogen, gaseous oxide of—*g.* Sulphuris, Sulphurous acid.

GASCARILLA, *Croton cascarella*.

GASEOUS, see *Gazeous*.

GASPING, (same etymon as *gaping* [?]), *Anhelatio*.

GASTEIN, MIN'ERAL WATERS OF. Gastein or Gasteiner Wildbad is in the Noric Alps, Austria. The waters are thermal. Temp. 106° to 118° Fah. They contain sulphate of soda, chloride of sodium, chloride of potassium, carbonate of soda, carbonate of lime, magnesia, manganese, iron, &c.

GAS'TER, (*ἡ στερῖς*, 'belly.') The abdomen. At times, but rarely, the uterus. Also, the stomach, in particular.

GAster MUSCULI, see *Muscle*.

GASTERALGIA, *Gastralgia*.

GASTERANAX, (*gaster*, and *αἰαξ*, 'a king.') A name given by Doleus to a hypothetical vital principle, corresponding to the Archæus of Van Helmont, the seat of which he placed in the lower belly. See *Bithnimalca*.

GASTERANGEMPHRAXIS, *Gasterangemphraxis*.

GASTERANGIEMPHRAX'IS, *Gasterangemphrax'is*, *Gasteremphrax'is*. Obstruction of the pylorus; (*gaster*, and *αἶσχος*, 'I strangle,' and *εμφρασις*, 'I obstruct.') Also, and properly, obstruction or congestion of the vessels of the stomach.

GASTÉRASE, (from *gaster*.) *Pepsin*.

GASTERASTHENIA, *Imbecill'itas*, seu *Aton'ia* seu *Lax'itas ventriculi*; (*gaster*, and *ασθενεια*, 'debility.') Debility of the stomach.

GASTERATAXIA, (*gaster*, and *αραξια*, 'irregularity.') *Embarras gastrique*.

GASTRECHE'MA, *gen. Gastreche'matis*, *Gastreche'ma*, *Son'itus stomachicus*; (*gaster*, and *ἤχημα*, 'sound.') Sound heard on auscultating or percussing the region of the stomach, and referred to that organ.

GASTEREMPHRAXIS, (*gaster*, and *εμφρασις*, obstruction.) *Gasterangemphraxis*.

GASTERHYSTEROTOMY, (*gaster*, *ὑστερα*, 'uterus,' and *τομή*, 'section.') *Cæsarean section*.

GASTRÆMIA, (*gastro*, and *αἷμα*, 'blood.') *Hyperæmia* of the stomach.

GASTRALGIA, (*gastro*, and *αλγος*, 'pain,') *Cardialgia*.

GASTRANEURYSMA, (*gastro*, and *ανευρυσμ'ος*, 'dilatation,') *Gastrectasis*.

GASTRATROPH'IA, (*gastro*, and *ατροφια*, 'wasting,') *Atrophy* of the stomach.

GASTRECHEMA, *Gastrectchema*.

GASTRELC'TASIS, *Gastrectasia*, *Gastraneurysma*, *Dilatatio ventriculi*; (*gastro*, and *εκτασις*, 'dilatation,') *Dilatation* of the stomach.

GASTRELCOBROSIS, (*gastro*, *ἔλκος*, 'ulcer,' and *βρωσις*, 'eating,') *Gastrobrosis ulcerosa*.

GASTRELCO'SIS, *Ulceratio seu Erculceratio seu Ulcera ventriculi*, (*gastro*, ad *ἔλκος*, 'ulceration,') *Ulceration* of the stomach. When accompanied with hectic, it constitutes *Gastrophth'isis*, *Gastrophthoë*.

GASTRELYTROTOMIA, see *Cæsarean Section*.

GASTRENCEPHALO'MA, *Gastromyelo'ma*, *Gastromyelo'sis*, *Gastroencephalo'sis*, (*gastro*, *εγκέφαλος*, 'the brain,' and *ομα*.) *Encephaloid* of the stomach.

GASTRENCHYTA, (*gastro*, *εν*, and *χυνω*, 'to pour,') *Stomach pump*.

GASTRENERITIC, *Gastroentericus*.

GASTRENERITIC, *Gastroenteritic*.

GASTRENERITIS, *Gastroenteritis*.

GASTRENTEROMALA'CIA, *Gastrenteromalacia'sis*, (*gastro*, *εντέρον*, 'intestine,' and *μαλακια*, 'softening,') *Softening* of the stomach and intestines.

GASTREPATICUS, *Gastrohepatic*.

GASTREPAT'ITIS, *Inflammatio ventriculi et hepatis*; (*gastro*, and *hepatitis*.) *Inflammation* of the stomach and liver.

GASTREPIPLOICUS, *Gastroëpiloic*.

GASTRERETHIS'IA; (*gastro*, and *ερεθίζω*, 'I irritate,') *Irritation* of the stomach.

GAST'RIC, *Gast'ricus*; (from *gaster*.) *Belonging* or relating to the stomach.

GASTRIC ACIDS, see *Gastric juice*.

GASTRIC ARTERIES, are three in number, *Arteria gastrica-epiploica dextra*, *A. gastrica-epiploica sinistra*, and *A. coronaria ventriculi*.

GASTRIC JUICE, *Sucus gast'ricus*, *Men'strum seu Fermentum ventriculi*, (F.) *Suc Gastrique*, *Gast'ric Acid*. A fluid, secreted from the mucous membrane of the stomach. As met with, it is a mixture of the fluids secreted by that organ with those of the supra-diaphragmatic portion of the alimentary canal. Owing to such admixture, the most contrary properties have been assigned to it. That such a fluid is secreted, which concurs powerfully in digestion, is evident from many considerations, and has been positively proved by the author and numerous others. It was found by him to contain, in man, chlorhydric and acetic acids, which are associated with a ferment, *pepsin*. The gastric fluid in cases of sudden death sometimes corrodes and perforates the stomach; giving rise to interesting questions in medical jurisprudence.

GASTRIC NERVES. The two cords by which the pneumogastric nerves terminate, and which descend on the two surfaces of the stomach; as well as the filaments of the great sympathetic, which accompany the gastric vessels.

GASTRIC PLEX'US, *Plexus coronarius ventriculi*. A nervous net-work, formed by the solar plexus. It accompanies the *Arteria coronaria ventriculi*, and passes along the lesser curvature of the stomach, to which it gives branches.

GASTRIC VEINS follow the same distribution as the arteries, and open into the *Vena porta abdominalis*.

GASTRICISM, *Gastricis'mus*, (from *gaster*.) A name by which is designated the medical theory, that refers all, or almost all, diseases to an accumulation of saburræ in the digestive passages.

GASTRICITE, *Embarras gastrique*.

GASTRILOQUE, (*gastro*, and *loquor*, 'I speak,') *Engastrimyth*.

GASTRILOQUIST, *Engastrimyth*.

GASTRILOQUUS, *Engastrimyth*.

GASTRIMARGUS, (γαστρίμαργος, from *gastro*, and *μαργω*, 'I rage,') Glutton.

GASTRINUM, Potash

GASTRISMUS, (γαστρίσμος, γς,) Gluttony, see *Saburra*.

GASTRIT'IC, *Gastrit'icus*, (*gastro*, and *itis*.) Relating to gastritis.

GASTRITIS, *Ventriculi inflammatio*, *Caum'a seu Empyema seu Cardialgia inflammato'ria*, *Febris stomachica inflammato'ria*, *Inflammatio gastritis*, *I. ventriculi seu stom'achi*, *Phlegmoné ventriculi*, *Inflammation of the stomach*, (F.) *Inflammation de l'Estomac*, *Gastrite*, *Catarrhe gastrique*. A disease, characterized by pyrexia; great anxiety; heat and pain in the epigastrium, increased by taking any thing into the stomach; vomiting and hiccup. Gastritis may either be seated in the peritoneal or mucous coat. It is most frequently in the latter—*Esogastritis*, *Endogastritis*, *Gastromycoenteritis*, *Gastritis mucosa*—being excited directly by acid ingesta. It requires the most active treatment;—bleeding, blistering, fomentations, diluents, &c. Some degree of inflammation of the mucous coat of the stomach was considered by the followers of Broussais to be present in almost all fevers; and the various forms of dyspepsia have been supposed by some to be nothing more than *chronic endogastritis*.

GASTRITIS ARTHRITICA, *Cœliagra*.

GASTRO, (from *gaster*,) in composition, the stomach, or belly.

GASTROARTHRITIS, Gout—g. *Ataxia*, *Dyspepsia*—g. *Atonia*, *Dyspepsia*—g. *Bronchitis*, *Fever*, *adenomenigeal*—g. *Enterite intense*, *Synocha*—g. *Enterite*, with nervous affection of the brain, see *Typhus*—g. *Enteritis*, follicular, *Dothinenteritis*: see *Typhus*.

GASTROBROSIS, *Perforatio ventriculi*, *Gastrorrhœxis*, (F.) *Perforation de l'Estomac*; (*gastro*, and *βρωσις*, 'the act of gnawing,') Corrosion and perforation of the stomach.

GASTROBROSIS ULCEROSA, *Gastrecobros'is*. Destruction and perforation of the coats of the stomach by ulceration.

GASTROCELE, *Gastrocele*, (*gastro*, and *κλη*, 'a tumour,') *Hernia of the stomach*, *Hernia ventriculi*, (F.) *Hernie de l'Estomac*. Hernia formed by the stomach through the upper part of the linea alba: a disease, the existence of which has been doubted by many. See *Epigastrocele*.

GASTROCEPHALITIS, (*gastro*, *κεφαλή*, 'head,' and *itis*.) Inflammation of the stomach and head, —a not uncommon concomitant of certain malignant fevers.

GASTROCHOLOSIS, (*gastro*, *χολή*, 'bile,' and *osis*.) Fever, *gastic*.

GASTROCNEME, (*gastro*, and *κνήμη*, 'the leg,') *Sura*.

GASTROCNEMIA, (γαστροκνημία,) *Sura*.

GASTROCNEMIL. The name of the two fleshy masses which occupy the posterior and superficial part of the leg, *Gemelli*; (F.) *Gastrocnémiens*, *Jumeaux de la jambe*, *Bifémoro-calcanéus*—(Ch. ;) the two constituting the *Gastrocnemius externus*, of English anatomists. These muscles are distinguished into *internal* and *external*, which are distinct above, but united at their inferior extremity. They are long, flat, and thick; and arise—the former from the posterior part of the outer condyle of the femur; the latter, from the posterior part of the inner condyle of the same bone. The aponeurosis, which unites these muscles below, joins with that of the *Solaris*, and forms with it, a large tendon, which, under the name *Tendo Achillis*, is inserted at the posterior part of the calcaneum. These muscles extend the foot on the leg, and the leg on the foot. They can, also, bend the leg and the thigh reciprocally on each other. For the *Gastrocnemius internus*, see *Soleus*.

GASTROCNEMIUM, *Sura*.

GASTROCELIACUS, *Cœliac*.

GASTROCELICUS, *Cœliac*.

GASTROCOLIC, see *Epiploon*, *gastrocolic*.

GASTROCOLICA, *Cardialgia*.

GASTROCOLITIS, (*gastro*, *κώλον*, 'colon,' and *itis*.) Inflammation of the stomach and colon.

GASTROCOLPOTOMIA, (*gastro*, *κολπος*, 'vagina,' and *τομή*, 'section,') see *Cæsarean section*.

GASTRODIDYMUS, *Gastrod'ymus*, *Did'ymus Symphyogast'rius*, *Psod'ymus*; (*gastro*, and *δίδυμος*, 'a twin,') A monstrosity in which twins are united by the abdomen.

GASTRODUODENAL, *Gastroduodena'lis*. Relating to the stomach and duodenum.

GASTRODUODENITIS, (F.) *Gastro-duodénite*. Inflammation of the stomach and duodenum, an affection, which figured largely in the system of Broussais. See *Gastro-enteritis*.

GASTRODUODENOPYRA, (*gastro*, *duodenum*, and *πυρ*, 'fire or fever,') Fever, *adenomenigeal*.

GASTRODYNE, (*gastro*, and *δύνη*, 'pain,') *Cardialgia*.

GASTRODYNIA, *Cardialgia*—g. *Flatulenta*. *Colica flatulenta*.

GASTROËLYTROTOMIA, (*gastro*, *ελτρον*, 'vagina,' and *τομή*, 'section,') see *Cæsarean section*.

GASTROËNTERAL'GIA, (*gastro*, *εντερον*, 'in intestine,' and *αλγος*, 'pain,') Pain in the stomach and intestine.

GASTROËNTER'IC, *Gastroënter'icus*, *Gastroënter'ic*, *Gastroënter'icus*, *Gastroëntes'tinal*, (*gastro*, and *εντερον*, 'intestine,') Relating to the stomach and intestine.

GASTROËNTERITE, *Gastroenteritis*.

GASTROËNTERITIC, *Gastroënterit'icus*, *Gastroënterit'ic*, *Gastroënterit'icus*, (*gastro*, *εντερον*, 'intestine,' and *itis*.) Relating to gastroenteritis.

GASTROËNTERITIS, *Gastroënteritis*, *Inflammatio ventriculi et Intestini'rum*, (F.) *Gastroënterite*, *Gastroenteric disease*. Inflammation of the stomach and small intestine. According to Broussais, the essential fevers of authors are *gastro-enteritis*, simple, or complicated.

GASTROËPIPLOIC, *Gastroëpipl'oicus*, *Gastroëpipl'oicus*, (*gastro*, and *επιπλοον*, 'the epiploon,') That which relates to the stomach and epiploon.

GASTROËPIPLOIC ARTERIES, *Gast'ric inf'erior ar'teries*, are two in number, and distinguished into *right* and *left*. The *right*, also called *Gastrohepatic*, *Gast'rica inf'erior dex'tra*, *Gastroëpipl'oica dex'tra*, is furnished by the hepatic artery. It descends behind the pylorus, and passes from right to left, along the great curvature of the stomach. It gives branches to the pancreas, duodenum, stomach, omentum majus, and terminates by anastomosing with the *Gastroëpipl'oica sinis'tra*, *Gast'rica sinis'tra* seu *inf'erior sinis'tra*. This—the *left*—arises from the splenic artery. It is of considerable magnitude, and passes from left to right, along the great curvature of the stomach, distributing its branches more particularly to the stomach and omentum majus. It terminates by joining the right gastroëpipl'oic.

GASTROËPIPLOIC GANG'LIONS are the lymphatic ganglions or glands, situate towards the great curvature of the stomach, between the two anterior laminae of the omentum majus.

GASTROËPIPLOIC VEINS are distinguished, like the arteries, into *right* and *left*. They empty themselves;—the former, into the superior mesenteric; the latter, into the splenic vein.

GASTROHÆMORRHAGIA, *Hæmatemesis*.

GASTROHEPAT'IC, *Gastrohepat'icus*, *He'pato-gast'ricus*, *Gastrepat'icus*, (*gastro*, and *ήπαρ*, gen. *ήπατος*, 'the liver,') Relating to the stomach and liver. This name has been given to several organs. See *Epiploon*, *gastrohepatic*, &c.

GASTROHEPATITIS, (*gastro*, *ήπαρ*, 'hepatos,

'liver,' and *itis*.) Simultaneous inflammation of the stomach and liver.

GASTROHYSTEROTOMY, (*gastro*, 'stomach,' and *hyster*, 'section,') Caesarean section.

GASTROINTESTINAL, Gastroenteric.

GASTROLIENALIS, (*gastro*, and *lien*, 'the spleen,') Gastrosplenicus.

GASTROLITHIASIS, (*gastro*, and *lithiasis*.) The formation of concretions, *gastrolithi*, in the stomach.

GASTROMALACIA, Gastromalaxia.

GASTROMALAXIA, *Gastromalacia*, *Gastromalacia*, *Malacogaster*, *Malaxia ventriculi*, *Dissolutio ventriculi*, *Emollitio ventriculi*, *Pseudophlogosis ventriculi resolutiva et colligativa*, *Metamorphosis ventriculi gelatinifurvis*, *Ensisio et perforatio spontanea ventriculi*, *Resolutio et diuresis ventriculi*, (F.) *Ramollissement de l'Estomac*, (*gastro*, and *palaxis*, 'softening,') Softening of the stomach, induced at times by the gastric secretions after death,—*Resolutio ventriculi antequam*.

GASTROMANTIS, (*gastro*, and *manis*, 'a diviner,') Engastrimyth.

GASTROMELES, (*gastro*, and *melos*, 'a limb,') A monster with one or more supernumerary limbs on the abdomen.

GASTROMENIA, (*gastro*, and *men*, 'a month,') Vicarious menstruation from the stomach.

GASTROMENINGITIS, (*gastro*, and *meningitis*, 'inflammation of the membranes of the brain,') Fever, gastric.

GASTROMETROTOMY, (*gastro*, *metra*, 'the uterus,' and *tom*, 'section,') Caesarean section.

GASTROMYCODERIS, (*gastro*, *mucos*, 'mucus,' and *deris*, 'skin,') See Stomach.

GASTROMYCODERITIS, (*itis*, 'inflammation,') See Gastritis.

GASTROMYELOMA, (*gastro*, *myelos*, 'medulla, marrow,' and the suffix *oma*.) Gastroencephaloma.

GASTRONOSUS, (*gastro*, and *nosus*, 'a disease,') Gastropathy.

GASTROPARALYSIS, *Gastroparalia*, *Paralysis ventriculi*, (*gastro*, and *paralysis*.) Paralysis of the stomach.

GASTROPATHY, *Gastropathia*, *Gastronosis*, *Gastronosis*, (*gastro*, and *pathos*, 'disease,') Disease of the stomach.

GASTROPERIODYDIA, (*gastro*, *peri*, 'around,' and *odyn*, 'pain,') Sool (India). A violent periodical neuralgic pain at the pit of the stomach, not uncommon in Hindoostan.

GASTROPHRENIC, *Gastrophrenicus*, (*gastro*, and *phren*, 'the diaphragm,') Belonging to the stomach and diaphragm.

GASTROPHRENIC LIGAMENT is a reflection of the peritoneum, which descends from the inferior surface of the diaphragm to the cardia.

GASTROPTHISIS, (*gastro*, and *pthisis*, 'I waste away,') Gastroleosis.

GASTROPTHOE, Gastroleosis.

GASTROPLEGIA, (*gastro*, and *plegia*, 'a stroke,') Gastr paralysis.

GASTROPYRA, (*gastro*, and *pyr*, 'fire, fever,') Fever, gastric.

GASTROPYRETUS, Fever, gastric.

GASTROPYRAPHY, *Gastropathy*, *Gastrorrhaphia*, *Gastrorrhaphy*, *Sutura abdominalis*, (*gastro*, and *pyra*, 'a suture,') The suture used for uniting wounds penetrating the abdomen, when they are too extensive or too unequal to be kept in contact by position, adhesive plaster, or appropriate bandages. The interrupted and quilted sutures are those chiefly employed.

GASTRORRHAGIA, (*gastro*, and *rrhagein*, 'to break forth,') Hematemesis.

GASTRORRHESIS, (*gastro*, and *rrhis*, 'rupture,') Gastrohrosis.

GASTRORRHEA, (*gastro*, and *rho*, 'I flow,')

Blennorrhoea seu Fluxus ventriculi, (F.) *Fluxus sanguinis de l'estomac*, *Catarhe stomachi*. A morbid condition of the stomach, which consists in the secretion of an excessive quantity of mucus from the lining membrane. Also, *Cœliac flux*.

GASTROSCIRRHUS, *Induratio ventriculi scirrhusa*, *Scirrhus seu Carcinoma ventriculi*. Scirrhus induration or cancer of the stomach.

GASTROSCOPIA, (*gastro*, and *σκοπεω*, 'I examine,') Abdominoscopy.

GASTROSES, (from *gaster*, and *osis*.) A generic name for diseases which are seated in the stomach.

GASTROSPLENIC, *Gastrosplenicus*, *Gastrosplenialis*, (*gastro*, and *σπλην*, 'the spleen,') Relating to stomach and spleen.

GASTROSTENOSIS, (*gastro*, *στυος*, 'contracted,' and *osis*.) Stricture of the stomach.

GASTROSTENOSIS CARDIACA ET PYLORICA. Narrowness of the cardiac and pyloric orifices of the stomach from cancer of that organ.

GASTROTOMY, (*gastro*, and *τομή*, 'section,') Several different operations have been so called.

1. The Caesarean Section. 2. An incision made into the abdomen for the purpose of removing some internal strangulation or volvulus, or to reduce hernia, *Laparotomy*; and, 3. The opening made in the stomach to remove a foreign body which has passed into it through the œsophagus.

GASTRYPERNERIA, (*gastro*, *πνερν*, 'in excess,' and *πνερν*, 'a nerve,') Morbidly increased activity of the nerves of the stomach.

GÂTEAU FÉBRILE, ('fever cake,') Ague cake.

GATEUX, ([F.] from *gater*, 'to spoil,') A name given in the insane hospitals, and in the hospitals for incurables and the aged in France, to persons who are incapable of attending to themselves, and have lost all sense of cleanliness.

GATHERING, Abscess, Suppuration.

GATILIER, Vitex.

GAUDIA FEDA, ('foul enjoyments,') Masturbation.

GAULTHERIA, *Gaultheria procumbens*, *Gaultheria repens* (?), *Mountain Tea*, *Partridge Berry*, *Berried Tea*, *Grouseberry*, *Deerberry*, *Spiceberry*, *Tea berry*, *Red berry*, *Wintergreen*, *Red berry Tea*, *Groundberry*, *Ground ivy*, *Ground holly*, *Hill berry*, *Box berry*, *Chequer berry*, (F.) *Gaulthérie*, *Pulmonier*, *Thé du Canada*, (after Dr. Gaulthier, of Quebec.) Ord. Ericaceae. An American plant, the leaves of which—*Gaultheria* (Ph. U. S.)—are one of the principal articles of the materia medica of some Indian tribes. The infusion of the leaves is stimulant and anodyne, and is said to have been used with advantage in asthma. The oil—*Oilum Gaultheriae*, Ph. U. S.—is used, chiefly on account of its pleasant flavour, to cover the taste of other medicines.

GAUNTLET, (from *gant*, 'a glove,') *Gantelet*.

GAUQUAVA, Smilax China.

GAUTIERA REPENS, *Gaultheria*.

GAY FEATHER, *Liatris spicata*.

GAYAC, *Guaiacum*.

GAYLUSSACIA RESINOSA, *Vaccinium resinum*, *Black Huckleberry*. Order, Ericaceae. (after Gay-Lussac.) An indigenous plant, whose fruit is sweet and agreeable. The leaves and root, as well as those of *GAYLUSSACIA DUMOSA*, *Dwarf huckleberry*, and of *G. FRONDOSA*, *Blue tangle*, indigenous, are used in decoction as astringents in diarrhoea, sore throat, &c.

GAZ, *Gas*, (Sax. *gas*, [G.] Geist, 'spirit,') (?) *Air*. Van Helmont first designated by this name—the etymology of which is not certain—the carbonic acid developed in the vinous fermentation. Afterwards, the term was appropriated to every permanently elastic fluid; that is, which preserves its æriform state at all temperatures; and ulti-

mately it was extended to all aëiform bodies; which were divided into *permanent* and *non-permanent* gases. The latter are generally termed vapours; they return to the liquid state, when a portion of their caloric is abstracted. The permanent gases, or *gazes* properly so called, are numerous, and may be divided into three sections with regard to their effects on the animal economy.

- | | |
|----------------------------------|---|
| 1. Irrespirable gases. | { Ammoniacal gaz, chlorohydric acid gas, deutoxide of nitrogen, nitrous acid gaz, and chlorine. |
| 2. Negatively deleterious gases. | { Hydrogen, nitrogen, carbonic acid. |
| 3. Positively deleterious gases. | { Oxygen, protoxide of nitrogen, carburetted hydrogen, carbonic oxide, sulphuretted hydrogen, and arseniuretted hydrogen. |

It is proper to remark that the term *respirable* has been very differently employed by different writers. Sometimes it has meant the power of supporting life when applied to the blood in the lungs. At others, all gases have been deemed irrespirable which are incapable of being introduced into the lungs by voluntary efforts,—without any relation to their power of maintaining vitality; and this is perhaps the best sense. The gases were at one time employed in medicine under great expectations, especially by the enthusiastic Beddoes; but they are now scarcely ever had recourse to. They differ considerably in their effects on the animal economy. Some, as oxygen, are exciting; others, as azote, depressing; whilst others, again, as the *Protoxide of nitrogen* or *laughing gas*, produce the most singular effects.

GAZ DES BALLONS, ('gaz of the balloons,') Hydrogen.

GAZ, PULMONARY, *Gaz of the lungs*. A name given to the expired air, which contains, besides common air, an increase of carbonic acid, water, and some animal matter.

GAZ SAN'GUINIS, ('gas of blood,') *Gaz animalis sanguinis*, *Hal'itus seu Aur'a seu Va'por sanguinis*, *Spir'itus roriferus*, *Hæmat'mus*. The halitus, or vapour, given off by freshly drawn blood.

GAZEL, *Ribes nigrum*.

GAZELLE, *Antilopus*.

GAZE'OUS or **GA'ZE'OUS**, *Gaseous*, *Gazeo'sus*, (F.) *Gazeux*. Relating or appertaining to, or having the form or character of, gas.

GAZ'OGENE, (*gas*, and *γενω*, 'I generate.')

A name given to a portable apparatus for preparing aerated or carbonated water,—as the *soda* water of the shops.

GAZEUX, *Gazeous*.

GE'ANT, Giant.

GE'ANTISME, see Giant.

GE'ARPULVER, ('labor-powder,') Ergot.

GEDE'OLA. The convex part of the liver, according to Du Cange.

GEILNAU, MIN'ERAL WA'TERS OF. Geilnau is a village in the grand duchy of Nassau, at no great distance from Frankfort. The waters contain carbonic acid, carbonate, sulphate, and phosphate of soda, chloride of sodium, and carbonates of lime, magnesia, and iron.

GEISMAR, MIN'ERAL WA'TERS OF. The mineral waters of Geismar, in Bavaria, are acidulous chalybeates.

GEISO'MA, (*γεισωμα*, like *γεισων*, 'the eaves of a house.')

Geisum.

GEIS'UM, (*γεισων*,) *Geis'on*. The part of the frontal bone over the eyes.

GELAPPIUM, (from *Jalapa* or *Xalapa*, in Mexico.) *Convolvulus jalapa*.

GELASINI DENTES, (*γελαστοι*, from *γελω*, 'I laugh,')

Incisive teeth.

GELASMUS, (from *γελω*, 'I laugh,') Canine laugh.

GELATIG'ENOUS, (*gelatin*, and *γενω* or *γενωω*, 'I engender.')

That which forms gelatinous structure. Dr. Carpenter restricts the term to substances yielding gelatin. The propriety of this may well be questioned.

GEL'ATIN, *Gel'atine*, *Gelati'na*, *Gabre'da*, (F.) *Gélatine*, *Gélée*, 'gelly or jelly,' (from *gelare*, 'to freeze, congeal.')

An immediate animal principle. It is semitransparent, insipid, inodorous, insoluble in cold water, very soluble in hot, which it thickens, and transforms into jelly on cooling. Gelatin is a nutritious substance; and, when dissolved in a considerable quantity of water, forms an emollient fluid, much used in therapeutics, but not the most easy of digestion. Two forms of gelatin are admitted—the one *glutin* or gelatin proper; the other *chondrin*.

GELATIN OF WHAR'TON, *Gelati'na Whartoniana*, *Jelly of the Cord*. A soft, dense, fluid, gelatinous substance, which envelops the umbilical cord, and is conceived by some to be inservient to the nutrition of the fetus.

GELATINA AQUATICA, ('aquatic gelatin,') *Brasenia hydropeltis*.

GÉLATINE DES CARILLAGES, ('gelatin of cartilages,') *Chondrin*.

GELATINIFORM MATTER OF THE INTES'TINE, *Albuminose*.

GELAT'INOUS, *Gelatino'sus*, (F.) *Gélatineux*. Relating or appertaining to or having the character or appearance of gelatin or jelly. Viscid.

GELATINOUS NERVE-FIBRE, see Nerve fibres.

GELATIO, (*gelare*, *gelatum*, 'to freeze, congeal.')

Congelation.

GELDING, ([G.] *gelten*, 'to castrate.')

Castration, Eunuch.

GELÉE, Gelly—*g. contre le Goître*, Balsam, ioduretted.

GEL'LY, *Jelly*, *Jus gela'tum seu coagula'tum*, *Ge'lu*, (F.) *Gélée*, (from *gelare*, 'to congeal.')

A substance of a soft consistence, tremulous, and transparent, which is obtained by an appropriate treatment from animal or vegetable matters; hence the distinction into *animal* and *vegetable jelly*. The former is merely a concentrated solution of gelatin, left to cool.

Vegetable Jelly is found in the juice of the currant, mulberry, and of almost all acid fruits, when ripe. It is of itself colourless, but almost always retains a little colouring matter of the fruit which has furnished it. It has an agreeable taste; is scarcely soluble in cold water, but boiling water dissolves it readily: the jelly is, however, almost all deposited on cooling. If this aqueous solution be boiled for a long time, it becomes analogous to mucilage, and loses the property of being jellied on cooling.

GELSEM'INUM or **GELSE'MIUM NIT'IDUM**, *G. Sempervi'rens*, *Bigno'nia sempervi'rens*, *Yellow Jessamine*. The flowers, root, &c., of this shrub of the Southern States, *Sex. Syst.* Pentandria Digynia, *Nat. Ord.* Apocynæ, are narcotic, and the effluvia from the former are said sometimes to induce stupor.

GELU, Gelly.

GELUS, (*γελως*,) Risus.

GEMELLI, ('twins,') *Gastrocnemii*, *Ischio-trochanterianus*, *Testicles*.

GEMEL'LUS, *Gem'inus*, *Did'y'mus*, (F.) *Jumeau*, *Jumelle*, *Besson*, *Bessonne*. One of two children, twins, born at the same accouchement, or gestation. Also, relating to twins, as a 'twin conception.'

GEMELLUS MUS'culus. Cowper applies this name to the long portion of the triceps brachialis united to the inner portion.

GEMINI, ('twins,') Ischiotrochanterianus, Testicles.

GEMINUM CENTRUM SEMICIRCULARE, Tania semicircularis.

GEMINUS, Gemellus.

GEMIPOMA, Mamma.

GÉMISSEMENT, (from *gémir*, 'to groan or moan,') See Moaning.

GEMMA, ('a bud,') Granulation—g. Oculi, Crystalline.

GEMMATION, see Generation—g. Generation by, see Generation.

GEMMIPARITÉ, (*gemma*, 'a bud,' and *pario*, 'to bring forth,') See Generation.

GEMMIPAROUS, see Generation.

GEMUR'SA, a corn; also, a name given by the ancients to a disease seated between the toes;—the nature of which is unknown to us.

GEN, (from *yevaw*, 'I generate,') in composition, generating.

GENA, (*yavus*.) The Cheek, Wang, *Ge'nys*, *Pare'ia*, *Ganph'e*, *Ganphe'le*, *Gna'thos*, *Gnath'mus*, *Ma'la*, (Sc.) *Wonge*, (F.) *Joue*. The cheeks form the lateral parts of the mouth. Externally, they have no precise limits; they are continuous, above, with the lower eyelid; below, they descend as far as the base of the jaw; before, they terminate at the *ala nasi*, and at the commissures of the lips; and behind, at the ear. Their thickness varies according to the degree of fatness of the individual. They are formed of three layers;—one dermoid, another muscular, and the third mucous.

GENCIVES, *Gingivæ*.

GENEANTHROPY, (*gen*, and *ανθρωπος*, 'man,') Anthropogenia.

GENE'AS, (*yevcas*, from *yev*, 'the chin,') *Lani'go pri'ma*, *Probar'bium*. The downy hairs which first cover the cheek. Also, a bandage which passes under the chin.

GENEION, (*yevciou*.) Beard, Mentum.

GENERAL ANATOMY, see Anatomy—g. Practitioner, Surgeonapothecary.

GÉNERALE, Influenza.

GENERATIO, gen. *Generatio'nis*, Generation—g. *Æquivoca*, see Generation—g. *Calculi*, *Lithia*—g. *Homogenea*, see Generation—g. *Primigena*, see Generation—g. *Primitiva*, see Generation—g. *Originaria*, see Generation—g. *Spontanea*, see Generation.

GENERA'TION, *Genera'tio*, *Gen'esia*, *Genne'sis*, *Go'n'e*, *Go'nus*, *Procrea'tio*, *Procrea'tion*, *Breed'ing*, ([L.] *genere*, *genitum*, *yavw*, or *yevaw* 'I engender,') Under this name physiologists comprehend the aggregate of functions, which concur, in organized beings, towards the production of their kind. The act of generation means the union of the sexes. See Coition. The writers of antiquity believed, that all organized bodies are produced, either by what is termed *univocal* or *regular* generation, *Homogen'esia*, *Genera'tio homogen'ea*, *Propaga'tio*, which applies to the upper classes of animals and vegetables, or by *spontaneous* generation, *Autogen'ia*, *Heterogen'esia*, *Heterogen'ia*, *Generatio heterogen'ea* seu *æquivoca* seu *primiti'va* seu *primig'ena* seu *originaria* seu *spontanea*, *Spontéparité* (Dugès), which they considered applicable to the very lowest classes only, as the mushroom, the worm, the frog, &c. There are still many distinguished naturalists who consider that beings, low in the scale of animality, are produced in the latter way. Spontaneous generation and *equivocal* generation have been regarded by many to be synonymous. Others, however, mean by spontaneous generation, the production of a new being from the mere combination of inorganic elements; whilst by equivocal generation they understand the evolution of a new being from organized beings dissimilar to themselves, through some irregu-

larity in their functions, or through the incipient decay or degeneration of their tissues. As to the mode in which regular generation is accomplished, there have been many views. According to the doctrine of Hippocrates, and of the ancient philosophers, the ovaries of the female furnish a prolific fluid, similar to that of the male; and the fœtus results from the mixture of the two seeds in copulation. Steno and others conceived, that the ovaries contain ova, which are not developed until vivified by the male sperm. Bonnet and Spallanzani believed in the pre-existence of germs, created since the origin of the world, but *encased* in each other, and becoming developed in succession; whence it would follow that the ovary of the first female must have contained the germs of all subsequent generations: and that the number of these germs must go on always diminishing, until ultimately extinct. This was the system of *evolution of germs*. According to Leeuwenhoek, the ovaries do not contain eggs, but vesicles destined to receive animalcules; which, in his view, live in the sperm. Thousands of these animalcules are thrown into the uterus during copulation, and the most expeditious and vigorous reaches the ovary, after having scattered and destroyed its competitors. Buffon—admitting the hypothesis of the two seeds—supposed that they were formed of molecules proceeding from every part of the body of each parent; and that, by a kind of elective affinity, those which were furnished by the head, the trunk, or the extremities of the male parent, could only unite with those proceeding from the same parts of the female. Before him, Maupertuis, admitting, with many of the ancient philosophers, the system of *Epigen'esia*, and adopting, as regarded the composition of the sperm, a theory analogous to that of Buffon, had supposed that the molecules, capable of being organized, were attracted towards a centre; that the nose attracts the two eyes; the body, the arms; the arms, the hands, &c., nearly as the particles of a salt, dissolved in a liquid, arrange themselves in regular crystals around the same nucleus. These and various other systems have been successively proposed and abandoned, and the mystery of generation remains impenetrable.

The simplest kind of reproduction does not require sexual organs. The animal separates into several fragments, which form so many new individuals. This is *Fissip'arous* generation, *Fissip'arism*, *G. from fis'sion*, (F.) *Fissiparité*, *Scissiparité*, *Scission*, *Cloisonnement*, *Reproduction ou Multiplication méristématique*. *Gemmip'arous* generation, (F.) *Gemmiparité*, *Génération par Gemmation*, *Sarculation* ou *Bourgeonnement*, consists in the formation of buds, sporules or germs on some part of the body, which at a particular period drop off and form as many new individuals. These kinds of reproduction require but one parent—*Monog'eny*. In *Ovip'arous* generation, (F.) *Oviparité*, the egg is hatched out of the body. In *Ovivivip'arous* generation, the new being is hatched in the excretory passages. In *Vivip'arous* generation, the new individual is born under its appropriate form; and in *marsupial* or *marsupiate* generation, the young being, born at a very early stage of development, is received and nourished in a *marsupium* or pouch. In *alter'nate* generation, (F.) *Génération alternante*, the young not only do not resemble the parent at birth, but remain dissimilar during their whole life, so that their relationship is not apparent until a succeeding generation. Thus, the cercaria undergoes a change into the distoma. See Metagenesis. All these kinds of reproduction require the union of sexes—*Dig'eny*.

All the acts comprising the function of generation in man may be referred to five great heads. 1. *Copulation*. 2. *Conception* or *fecundation*. 3. *Ges-*

tation or Pregnancy. 4. Delivery or Accouchement; and, 5. Lactation.

GENERATION, ACT OF, see Generation—g. Alternate, see Generation, and Metagenesis—g. Equivocal, see Generation—g. by Fission, see Generation—g. Fissiparous, see Generation—g. by Gemination, see Generation—g. Gemiparous, see Generation—g. Marsupial, see Generation—g. Organs of, female, see Vulva—g. Oviparous, see Generation—g. Regular, see Generation—g. Spontaneous, see Generation—g. Univocal, see Generation—g. Viviparous, see Generation.

GEN'EROUS, *Genero'sus*, (*genus*, *gen. generis*, 'a race;' *generosus*, 'of a noble race, noble.') A name given to wines which contain a great quantity of alcohol.

GÉNÈSE, Genesis.

GENESIAL, Genital.

GENESIOLOG'Y, *Genesiologia*, (*genesis*, 'generation,' and *logos*, 'a discourse.') The doctrine of generation.

GÉNÉSIQUE, Genital.

GÉNÉSIQUE, LE, Appetite, venereal.

GEN'ESIS, (*genesis*,) Generation. Also, origin or first formation—as the *genesis*, (F.) *Génèse*, of the teeth.

GENÉT, (old [F.] *Genet*: from *genista*,) Spartium scoparium—g. à Balai, Spartium scoparium—g. des Teinturiers, *Genista tinctoria*.

GENETHLIACUS, (*γενεθλιος*, 'natal,' 'pertaining to nativity,') A name given by the ancients to certain astrologers, who, from the state of the heavens at the time of the birth of an individual, predicted his future character and the events of his life.

GENET'ICA, (*genesis*, 'generation,') Diseases of the sexual functions; the 5th class in Good's Nomenclature. Also, agents that act on the sexual organs.

GENETICUS, (*γεννητικος*,) Genital.

GÉNÉTIQUE, Genital.

GENEVA, (from *genièvre*, 'juniper, gin,') Gin, see Spirit.

GENÉVRIER, *Juniperus communis*—g. *Oxycedre*, *Juniperus oxycedrus*.

GENGIVITE, (*gingivæ*, 'the gums,' and *itis*,) Ulitis.

GÉNI, Genian.

GEN'IAN, *Geni'anus*, *Gen'ial*, (*γενιον*, 'the chin,') The Genian apoph'ysis or process, (F.) *Apophyse génienne* ou *géné*, is situate at the posterior part of the symphysis menti, and is formed of four small tubercles.

GENICULATUM, (from *geniculum*, 'a small knee or joint,' from *genu*, 'the knee,') see Corpora geniculata.

GÉNIE, *Gen'ius*. The French sometimes apply this term to diseases nearly synonymously with nature; as *Génie inflammatoire*, *G. bilieux*, *G. adynamique*. Some use it in the same sense as type; *Génie intermittent*. The unwonted preponderance of any mental faculty is also so called.

GÉNIEN, Genian.

GENIÈVRE, (from *juniperus*,) Gin, *Juniperus communis* (the berry.)

GEN'I'O, (from *γενιον*, 'chin,') in composition, the chin.

GEN'IOGLOSSUS, *Geniohyoglossus*, (*genio*, and *γλωσσα*, 'the tongue,') *Mesoglossus*, *Mesoglossus*, *Nonus linguae musculus*. The name of a flat, triangular muscle, which extends from the genian apophysis to the inferior surface of the os hyoides and tongue, which last it carries forward.

GENIOHYODES, *Geniohyoideus*.

GENIOHYOGLOSSUS, *Genioglossus*.

GENIOHYODES, *Geniohyoideus*.

GENIOHYOÏDEUS, *Geniohyo'des*, *Geniohyoï-*

des, *Mentohyoideus*, (*genio*, and *hyois*, 'the os hyoides,') A muscle which arises from the genian apophysis, and is inserted at the anterior part of the body of the os hyoides. Its use is to raise the os hyoides, and carry it forwards. It may, also, contribute to depress the lower jaw in contracting towards the os hyoides.

GENIOPHARYNGE'US, (*genio*, and *φαρυγξ*, 'the pharynx,') A name given, by some anatomists, to a bundle of fibres which passes from the lower jaw to the sides of the pharynx, and forms part of the constrictor pharyngis superior.

GENIOPLASTY, *Genioplastie*, (*genio*, and *πλασσω*, 'I form,') The operation for restoring the chin. At times used instead of *Genioplasty* for the operation for restoring the cheek.

GEN'IPA OBLONGIFOLIA, *Huito*. A plant of Peru, with the juice of which the Indians paint their legs, to protect them against the stings of insects.

GENIPI and G. ALBUM, *Achillea moschata*, *Artemisia rupestris*—g. *Blanc*, *Artemisia rupestris*—g. *Verum*, *Achillea atrata*.

GENISTA, *Spartium scoparium*—g. *Canariensis*, see *Rhodium lignum*—g. *Hirsuta*, *Spartium scoparium*—g. *Scoparia*, *Spartium scoparium*.

GENISTA SPINO'SA IN'DICA, *Bukel Schulli*. *Ord.* Leguminosae. An oriental tree, a decoction of the roots of which is diuretic. The leaves boiled in vinegar have the same effect.

GENISTA TINCTORIA, *Genisto'des tinctoria*, *Spartium tinctorium*, *Dyers' broom* or *weed*, *Green weed*, *Wood wazen*, *Woodwee*, (F.) *Genêt des Teinturiers*. A shrub cultivated in this country and in Europe. The flowering tops and seed have been used in medicine. It has the same properties as *Spartium scoparium*.

GENISTOIDES TINCTORIA, (*genista*, and *αἶς*, 'resemblance,') *Genista tinctoria*.

GEN'ITAL, (from *geno*, for *gigno*, 'to generate,') *Genitalis*, *Genet'icus*, *Generatio'ni inserviens*, *Gen'ial*, *Genet'icus*, (F.) *Génital*, *Généti-*

que, *Généti-*. That which belongs or relates to generation.

GENITAL OR'GANS, *Sex'ual Or'gans*, *Puden'da*, *Natura'lia*, *Natu'ra*, *Edo'a*, *Me'ra*, *Me'sa*, *Mo'rion*: *Gen'ital*, *Noble*, *Natural* or *Private parts*; *Priv'ities*, *Priv'y parts*, *Priv'y Mem'bers*, *the Parts*, *Pars*, *Pars corporis seu obsc'ra*, *Partes*, *Partes genitalis seu generatio'ni inservientes seu obsc'rae*, *Me'dea*, *Vere'da*, *In'guen*, *Genito'lia*, *Genet'ica*, *Gym'na*, *Mem'bra puden'da*, *Or'ganæ generatio'ni inservientia*, *O. sex'ualia seu genitalia*, (F.) *Organes génitaux*, *Parties génitales* ou *honteuses* ou *génitoires* ou *nobles* ou *sexuelles* ou *naturelles*, *Les Parties Génitoires*. The parts that are inservient to the reproduction of the species. These are very different in the male and female. In man, they are numerous; some secreting the sperm, as the testicles and their appendages; others retaining it, as the vesiculae seminales; and another for carrying it into the organs of the female,—the penis. In the female, the parts of generation form an apparatus, perhaps more complicated than that of the male. Some are inservient to copulation, as the vulva, vagina, &c., others to conception and the preservation of the product for a determinate time,—as the uterus and its appendages; whilst others concur in the alimentation of the infant after birth, as the mammae.

GENITALE, Sperm—g. *Caput*, *Glans*.

GENITALIA, Genital organs—g. *Viri*, *Pudibilia*.

GENITOCRURAL NERVE, *Nervus genitocruralis*, *Subpu'bial* or *Inter'nal inguinal nerve*. A branch of the second lumbar nerve, which passes through the psoas muscle, and approaching the femoral arch divides into two branches,—an internal, *scrotal*, or *gen'ital*, *nervus spermaticus seu puden'dus ext'rius*, and an external or *femoral cutaneous branch*, *lum'bo-inguinalis*.

GENITOÛRINARY, see Urogenital.

GÉNITOIRES, Genital Organs.

GENITÛRA. That which is fecundated or engendered in the maternal womb. This word has been used synonymously with *embryo*, *fœtus*, and infant. Also, the sperm; and the penis. See GONE.

GENIUM, (γενιον,) Mentum.

GENNESIS, (γενεσις,) Generation.

GENNETICA, (γεννητικα,) Genital Organs.

GENNE'TICŌI, (Morbi,) Aidoici, Aëdæci, in the classification of Dr. Wm. Farr are genital diseases.

GENNETICOCNES'MUS, (gennetica, and κνησμος, 'itching,') Itching of the genital organs.

GENNETICON'OSI, Genneticonn'ei, (gennetica, and νόσος, 'disease,') Disease of the genitals.

GENNETICUS, (γεννητικός,) Genital.

GEN'ŌA, CLIMATE OF. The climate of this Italian city and its vicinity has been often selected as favourable for the phthisical valetudinarian during the winter; but it is now admitted to be decidedly improper for pulmonary affections, being subject to frequent and rapid changes of temperature, and to dry cold winds from the north, alternating with warm moist winds from the south-east.

GENONU'SI, Morbi sexūs, (γενος, 'sex,' and νόσος, 'disease,') Sexual diseases.

GENOPLASTIC, (γενν, 'the jaw,' and πλασσω, 'I form,') Meloplastic.

GENOU, Genu.

GÉNOUILLÉS, (Corps,) (from geniculum, 'a small knee or joint,') Corpora geniculata.

GENOUILLET, Convallaria polygonatum.

GENRE, Genus.

GENSANG, Panax quinquefolium.

GENTIA, Gentianina.

GENTIAN, (after Gentius, King of Illyria, who used some species medicinally,) Gentiana lutea, Triosteum—g. Blue, Gentiana catesbei—g. Catesbian, Gentiana catesbei—g. Horse, Triosteum—g. Southern, Gentiana catesbei—g. White, Laserpitium latifolium, Triosteum—g. Yellow, Gentiana lutea, see Calumba.

GENTIANA, G. lutea—g. Alba, Laserpitium latifolium—g. Amaryllodes, G. quinqueflora—g. Cachenlaguen, Chironia Chilensis—g. Centaurium, Chironia centaurium.

GENTIANA CATESBE'I, G. Catesbia'na, Blue Gentian, Catesbian or Southern Gentian, Blue-bells, Bit'terroot. Ord. Gentianaceæ. It is a pure and very bitter, and the root may be used wherever that of the *Gentiana lutea* is proper.

GENTIA'NA CHIRAY'TA, G. Chirayi'ta, Henrice'a Pharmacear'cha, Suer'tia, Agatho'tes chirayi'ta, Ophelia' chirata, Chire'ta, (Ph. U. S.), Chira'ta (Ph. B. 1864), Chirayi'ta, Chirae'ta. A native of India, the herb and root of which has been much employed in that country, in dyspepsia, and as an antiperiodic in intermittents. It is preferred by some to sarsaparilla, where the latter is considered to be indicated,—as after large quantities of mercury have been taken, or where profuse salivation has been induced. It has also been advised in atonic leucorrhœa. It yields its virtues to alcohol and water.

GENTIANA CRINITA, see G. quinqueflora—g. Gerardi, Chironia centaurium.

GENTIANA LU'TEA. The systematic name of the official gentian; Gentia'na, G. ma'jor seu vef'erum seu ru'bra, Suer'tia lu'tea, Aste'rias lu'tea, Yel'low Gentian, Gent'ian, Fel'wort. Ord. Gentianeæ. (F.) *Gentiane jaune*. This is a plant common in the mountains of Europe. The root is almost inodorous, extremely bitter, and yields its virtues to ether, alcohol, and water. It is tonic and stomachic; and in large doses, aperient.

Dose, gr. x to ℥ij. It is most frequently, however, used in infusion or tincture.

GENTIANA MAJOR, G. lutea—g. Ochroleuca, see G. saponaria—g. Peruviana, Chironia Chilensis.

GENTIANA QUINQUEFLO'RA, G. Amaryllôides, Five-flowered Gent'ian; and GENTIA'NA CRINI'TA, Fring'ed Gentian, indigenous, are possessed of like virtues.

GENTIANA RUBRA, G. lutea.

GENTIANA SAPONA'RIA, Soap'wort Gent'ian, and G. ochroleuca, Sam'pson's Snake'root, indigenous, have the properties of *Gentiana lutea*; and are said also to be diaphoretic.

GENTIANA VETERUM, G. lutea.

GENTIANE JAUNE, Gentiana lutea.

GENTIANI'NA, Gent'ianine, Gent'ianin, Gentia'nia, Gent'ia. A supposed neutral substance, obtained from gentian, and, by some, presumed to be its active principle. It is not so.

GENTILITIUS, Hereditary.

GENU, Go'ny, (γενν,) (F.) Genou. The articulation of the leg with the thigh;—the Fem'oro-tib'ial or knee-joint, articula'tio ge'nu, the knee. It is the most complicated in the body, and is formed by the inferior extremity of the femur, the superior extremity of the tibia, and the rotula. The articular surfaces of the bones are covered by layers of cartilage—more or less thick—and by the synovial membrane of the articulation. The soft parts of this joint are,—the ligamentum patellæ, two lateral ligaments, distinguished by the names internal and external; a posterior ligament, Ligamen'tum post'icum Winslow'ii; two crucial ligaments,—the one anterior, and the other posterior; two interarticular fibro-cartilages; some albugineous fibres, which form an imperfect capsule, &c. The knee receives its arteries from the femoral and popliteal. They bear the name articular. Its veins have the same distribution as the arteries, and discharge their blood into the saphena and crural. Its nerves are furnished by the sciatic, popliteal, and crural. The joint is protected by the tendons and muscles which surround it.

The French use the term *Articulation en genou* for a joint, in which the head of a bone is received into a bony cavity of another, where it rolls and moves in all directions.

GENU CORPORIS CALLOSUS, see Corpus callosum—g. Varum, Entogonyancon.

GENUGRA, (genu, and αἴψα, 'seizure,') Gonagra.

GENUINUS, (from genu,) Legitimate.

GE'NUS, gen. Gen'eris, (F.) Genre, (γενος, 'race,') A collection or group of species, analogous to each other, and which can be united by common characters. When a species cannot be referred to a known genus, it constitutes a distinct one.

GENUS CURATIONIS, Ratio medendi.

GENY, (γενν,) Genys.

GENYANTRAL'GIA, (geny, αντρον, 'the antrum,' and αλγος, 'pain,') Pain in the antrum of Highmore.

GENYANTRI'TIS, (geny, αντρον, 'the antrum, and itis,) Inflammation of the antrum of Highmore.

GENYANTRUM, (geny, and αντρον, 'a cave,') Antrum of Highmore.

GENYOCYNANCHE, (geny, and cynanche,) Cynanche parotidæa.

GENYOPLASTY, (geny, and πλασσω, 'I form,') see Genioplasty.

GENYS, (γενος,) Genu, 'the jaw,' also the chin and the beard. See Genæa.

GEOFFRÆA INER'MIS, Geoffræ'a, G. race mo'sa seu Piso'nis seu Jamaicæ'nis, Vouacap'oua America'na, Geoffroy'a, Cabbag'ium, And'ra inermis seu racemo'sa, Cab'bage Tree, Cab'bage Bark Tree, Wormbark Tree. Ord. Leguminosæ

Sex. Syst. Diadelphia Decandria, (after Dr. E. F. Geoffroy, a French naturalist.) The odour of the bark is very unpleasant. It is anthelmintic and cathartic. Dose of the powder, $\mathfrak{H}$ to $\mathfrak{Hj}$.

GEOFFRÆA JAMAICENSIS, G. inermis—g. Pisonis. G. inermis—g. Racemosa, G. inermis.

GEOFFRÆA SURINAMENSIS, Andri'a Surinamensis, has similar properties.

Hutteneschmidt has separated the active principles of G. Jamaicensis and G. Surinamensis, to which he has given the names *Jamaicine* and *Surinamine*. They are alkaline.

GEOFFRÆA VERMIFUGA, Andri'a ibai, Arriba, *Skulemura* Fernambucoensis; a South American plant, has a fruit, the almond of which, called *Angéline*, has a reputation at Rio Janeiro as a vermifuge. Dose, a grain to fifteen; or it may be given in infusion.

GEOFFROYA, Geoffræa inermis—g. Jamaicensis, G. inermis.

GEOGRAPHY, MEDICAL, *Geographia Medica*, ($\gamma\eta$, 'the earth,' and $\gamma\alpha\phi\alpha$, 'I describe,') *Nosogeography*, *Nosogeographia*, *Nosochthonography*, *Nosochthonologia*. The description of the surface of the globe as regards the influence of situation on the health, vital functions, and diseases of its inhabitants, vegetable and animal; but principally on those of man.

GEOGRAPHISM, *Geophag'ia*, *Geophag'ia*, *Geophag'ia*, *Geophag'ia*, *Earth-eating*, *Clay-eating*, *Dirt-eating*, ($\gamma\eta$, 'earth,' and $\gamma\alpha\omega$, 'I eat,') The act or practice of eating earth. See *Chthonophagia*.

GEOGRAPHIST, *Geophagus*, *Geophagus*, *Geophagus*, *Geophagus*, *Earth-eater*, *Clay-eater*. One who eats earth.

GEORGIA, MINERAL WATERS OF. The water of *Indian Springs*, in Butts County, is sulphureous. The *Warm Springs*, in Meriwether County, have a temperature of 90° Fah. *Madison's Springs*, in Madison County, are chalybeate; and so are *Rowland's Springs*, in Cass County. All these are frequented by invalids.

GEREBOLOG'IA, properly *Geratolog'ia*, ($\gamma\epsilon\pi\alpha\varsigma$, 'old age,' and $\lambda\omicron\gamma\omicron\varsigma$, 'a discourse,') The doctrine, or a description, of old age.

GERAFLOURIS, *Dianthus caryophyllus*.

GERANIS, ($\gamma\epsilon\pi\alpha\varsigma$, 'a crane,' which it resembled.) A bandage, used by the ancients in cases of fractured clavicle. Some authors attribute it to Hippocrates; others, to Perigenes.

GERANIUM, (from $\gamma\epsilon\pi\alpha\varsigma$, 'a crane,' because its pistil is long, like the bill of the crane.) *Ord.* Geraniaceæ. The *Crane's Bill*.

GERANIUM CAROLINIANUM, see G. Robertsonium—g. Fœtidum, G. Robertsonium—g. Maculatum, see Geranium, and G. Moschatum.

GERANIUM MOSCHATUM, *Erodium Moschatum*, (F.) *Bec de grue musqué*, A European plant, esteemed to be excitant and diaphoretic.

GERANIUM NOVEBORACENSE, G. Maculatum—g. Purpureum, G. Robertsonium.

GERANIUM ROBERTIANUM, *G. fœtidum* seu *purpureum*, *Stinking Crane's Bill*, *Herb Robert*, (F.) *Herbe à Robert*, *Bec de Grue Robertin*. This plant was, at one time, used as an antispasmodic and slight stimulant, as well as for an external application in various painful sores and inflammations.

Geranium Carolinianum, has similar properties. Most of the species of geranium have been used as astringents. In some of the northwestern parts of the United States, the rhizoma of *Geranium maculatum*—*Geranium*, (Ph. U. S.), *G. Noveboracense*, *Spot'ted Crane's Bill*, *Crane's foot*, *All'un root*, *Tarmen'til*, *Stork's bill*—is called *Racine à Bequet*, after a person of that name. It is highly extolled by the Western Indians as an antispasmodic.

GERAS, ($\gamma\epsilon\pi\alpha\varsigma$,) *Senectus*.

GERAT'ICI (*Morbi*), ($\gamma\epsilon\pi\alpha\varsigma$, gen. $\gamma\epsilon\pi\alpha\varsigma$, old age.) Developmental diseases of old people, in the classification of Dr. W. Farr.

GERATOLOGIA, Geræologia.

GERÇURE, Fissure.

GERM, *Ger'men*, *Blas'te*, *Blas'te'ma*, (F.) *Germe*. The rudiment of a new being, not yet developed, or which is still adherent to the mother.

GERM FORCE, Plastic force.

GERMANDER, COMMON, *Teucrium chamaedrys*—g. Creeping, *Teucrium chamaedrys*—g. Marum, *Teucrium marum*—g. Small, *Teucrium chamaedrys*—g. Water, *Teucrium scordium*.

GERMANDRÉE AQUATIQUE, *Teucrium scordium*—g. Maritime, *Teucrium marum*—g. Officinale, *Teucrium chamaedrys*—g. *Scorodone*, *Teucrium scordium*.

GERME, Germ—g. *Faux*, Conception, false.

GERMEN, gen. *Ger'minis*, (from *gerere*, 'to produce,') Germ, Sperm—g. Dentale, Dental Pulp—g. Falsum, Mole—g. Spurious, Conception, false.

GERMINAL, *Germinative*, *Germinalis*, *Germinativus*, (F.) *Germinativ*; (from *germen*,) Relating or belonging to a germ; as '*germinal cell*.'

GERMINAL CELL, see Cytoblast—g. Membrane and Vesicle, see Molecule—g. Nucleus, see Molecule—g. Spot, see Molecule—g. Vesicle, see Molecule.

GERMS, DISSEMINATION OF, *Panspermia*—g. Pre-existence of, see Preformation.

GERO, GERONTO, ($\gamma\epsilon\tau\alpha\omega$, gen. $\gamma\epsilon\tau\alpha\omega\varsigma$, 'an aged person,') in composition the same.

GEROBOS'CIA, *Gerontobosc'ia*, (*gero*, and $\beta\omicron\sigma\kappa\eta$, 'food,') Nourishment or maintenance proper for the aged.

GEROCOMEUM, *Gerocomeum*.

GEROCOM'IA, *Gerocom'icæ*, *Gerontocom'icæ*, (*gero*, 'an aged person,' and $\kappa\omicron\mu\omicron\varsigma$, 'to take care of,') The part of medicine whose object is the preservation of the health of the aged:—the hygiene of old people.

GEROCOMICÆ, *Gerocomia*.

GEROCOMI'UM, *Gerocom'um*, *Preshyodoch'ium*, *Gerontocom'ium*, *Gerotrophe'um*. An hospital for the aged.

GÉROFLE, see *Eugenia caryophyllata*.

GÉROFLÉE JAUNE, *Cheiranthus cheiri*.

GERONTATROPHIA, (*geronto* and *atrophia*,) *Marasmus senilis*.

GERONTO, in composition, see *Gero*.

GERONTOBOSCIA, *Geroboscia*.

GERONTOCOMICE, *Gerocomia*.

GERONTOCOMIUM, *Gerocomium*.

GERONTOPIA, (*geronto*, and $\omega\psi$, 'the eye,') Weakness of sight of the aged.

GERONTOTOXON, *Gerotoxon*.

GERONTOXON, *Gerotoxon*.

GEROTOX'ON, *Gerontox'on*, *Gerontotox'on*, *Leuco'ma Gerontotox'on*, *Mac'ula cor'neæ arcuata*, *Ar'cus senilis*; (*gero*, and $\tau\omicron\beta\omicron\varsigma$, 'a bow,') A bow shaped opacity at the under margin of the cornea, common to old people, and supposed, by some, to be an indication of the existence of fatty degeneration of the heart.

GEROTROPHÆUM, (*gero*, and $\tau\tau\epsilon\pi\epsilon\iota\nu$, 'to nourish,') *Gerocomium*.

GEROTROPHIA, (*gero*, and $\tau\tau\epsilon\pi\epsilon\iota\nu$, 'to nourish,') *Geroboscia*.

GERSA, *Plumbi subcarbonas*—g. *Serpentariæ*, see *Arum maculatum*.

GÉSIER, ('gizzard,') *Ventriculus callosus*.

GES'TA, 'things done,' (*gerere*, *gestum*, 'to do,') *Acta*. A Latin term, introduced by Hallé into medical language, to designate, among the objects which belong to hygiene, the functions which consist in the voluntary movements of muscles and organs. In the class *Gesta* are found *sleep*, the *walking state*, movements or locomotion, and *rest*.

GESTA'TION, *Gestatio*, *Pho'ra*, (*gestare*, *ges-*

tatum, 'to carry.') The time during which a female who has conceived carries the embryo in her uterus. See Pregnancy. Gestation, likewise, signifies the bearing or carrying of an individual; a kind of exercise easier than that in which he moves by virtue of his own powers. Thus, we speak of *gestation on horseback*, in a carriage, &c.

GESTATION. PROTRACT'ED. Pregnancy protracted beyond the usual period. See Pregnancy.

GESTICULATIO*n*, *Gesticulatio*, (*gestus*, from *gerere*, *gestum*, 'to carry,' 'the carriage of the body.') The act of making many gestures; a symptom in disease, which indicates great encephalic erethism. It is met with in numerous affections.

GETHYL'LIS SPIRA'LIS. A South African plant, *Nat. Ord.* Amaryllidæ, the orange-coloured fruit of which has a peculiar fragrance. An infusion in spirit or tincture is used in flatulence and colic.

GEUM, *G. rivale*—*G. Caryophyllum*, *G. urbanum*—*G. Nutans*, *G. rivale*—*G. Palustre*, *G. rivale*.

GE'UM RIVA'LÊ, *G. palustre* seu *nutans*, *Caryophylla* *ta aquatica* seu *nutans*, *Benedicta sylvestris*, *Wal'ter* or *Pur'ple a'vens*, *Choc'olate root*, (F.) *Benoîte aquatique ou des Ruissaux*. *Ord.* Rosaceæ. *Sex. Syst.* Icosandria Polygynia. The root—*Geum* (Ph. U. S.)—is astringent. It has been much extolled in the cure of intermittents, diarrhœa, hemorrhage, &c.

GEUM URBA'NUM, *G. caryophyllum*, *Caryophylla* *ta*, *C. vulg'aris* seu *urbana*, *Caryophyllus vulg'aris*, *Samama'da*, *Lagophthalmus*, *Caryophylla*, *Janamun'da*, *Her'ba benedicta*, *Common a'vens*, *Herb Ben'net*, *Yellow a'vens*, *Star of the North*, *Bless'ed herb*, (F.) *Benoîte*. The root of this plant has a smell not unlike that of cloves. Taste bitterish, austere; virtues yielded to water and alcohol. It has been used in intermittents, dysentery, chronic diarrhœa, debility, &c. Dose, 5ss to 3j of the powder.

GEUM VER'NUM, *West'ern early a'vens*, has the same properties as

GEUM VIRGINIA'NUM, *White a'vens*, *A'vens*, *Ev'an root*, *Choc'olate root*, *Ben'net*, *Throatroot*, *Cure-all*, (F.) *Benoîte de Virginie*, common from Maine to Carolina and Kentucky, flowering in June and July. It has the same medical properties as *Geum rivale*.

GEUMA, gen. *Geum'atis*, (*γευμα*.) Taste.

GEUSION'OSI, (*gensis*, and *oosis*, 'disease.') *Geusiosis*. Diseases of the organ or sense of taste.

GEUSIS, (*γευσις*.) Taste. Rarely, the root of the tongue—*Rudix lingue*.

GEUTHMOS, (*γευθμος*.) Taste.

GEZIB, see *Pastinaca opoponax*.

GEZIT, *Cagot*.

GEZITAIN, *Cagot*.

GHERKIN, ([G.] Gurke, 'a cucumber,') see *Cucumis sativus*.

GHITTA JECOMO, *Cambogia*.

GIALAPPA, (from *Jalapa* or *Xalapa*, in Mexico.) *Convolvulus Jalapa*.

GI'ANT, *Gi'gas*, (F.) *Géant*, (from *γίγας*, gen. *γίγαντος*.) One much above the ordinary stature or of excessive growth—*Macroso'mia*, *Macroso'natio*, *Magnitudo gigante'a*, (F.) *Géantisme*.

GIBBA, *Hump*.

GIBBER, *Hump*.

GIBBEROSIT'AS, *Gibbositas*.

GIBBOSITAS, *Gibberositas*, *Cyphosis*, *Rhachio'cyphosis*, (*gibbus*, *gibbous*, 'something arched or vaulted; prominent,') *Gibbosities*, *Gib'bousness*, *Curvature of the spine*, *Hybo'ma*. A phenomenon which occurs in different diseases, particularly in rickets and caries of the vertebræ. See *Hump*.

GIBBOSITAS CARIOSA, *Vertebral disease*.

GIBBOUSNESS, *Gibbositas*.

GIBBUS, *Hump*—*G. Pottii*, *Vertebral disease*.

GIB'LETS. (According to Minshew, from *gobbet*, 'a mouthful' [?]; but, according to Junius, from (F.) *gibier*, 'game.') The word seems to be the old (F.) *Gibelez*, *Gibélet*, &c., i. e. *Gibier*, (L.) *Cibarium*, 'food,' (F.) *Abatis*. It means, generally, the parts which are cut off from a goose before it is roasted. Also, the extremities of fowls, such as the head, wings, feet; to which are sometimes added the liver, gizzard, &c. Soup, made from these, is moderately nutritious.

GIDDINESS, *Vertigo*.

GIDDY, (Sax. *gidig*.) *Vertiginous*.

GIFTS. A name given to white specks on the finger-nails, which have been superstitiously supposed to portend gifts.

GIGANTESQUE, *Gigantic*.

GIGANTEUS, (*γίγαντας*, and *γίγαντιος*, from *gigas*.) *Gigantic*.

GIGAN'TIC, *Gigante'us*, *Giganto'des*, (F.) *Gigantesque*. Relating to one much above the ordinary stature.

GIGANTODES, (*γίγαντων*, from *gigas*, and *odes*.) *Gigantic*.

GIGARTINA HELMINTHOCORTON, *Coralina Corsicana*—*G. Lichenoides*, *Fucus amylaceus*.

GIGARUM SERPENTARIA, (*gigas* and *arum*.) *Arum dracuncul*.

GIGAS, gen. *gigan'tis*, (*γίγας*, gen. *γίγαντος*.) *Giant*.

GILARUM, *Thymus serpyllum*.

GILEAD, BALM OF, see *Amyris opobalsamum*.

GILET DE FORCE, *Waistcoat*, *strait*.

GILL, *Glechoma hederacea*.

GILL-GO-BY-GROUND, *Glechoma hederacea*.

GILLA THEOPHRASTI, *Zinci sulphas*—*G. Vitrioli*, *Zinci sulphas*.

GILLE'NIA STIPULA'CEA, (after *Arnoldus Gilenius*, a German physician, [?]) *Small-flowered Ind'ian Phys'ic*, has the same properties as

GILLE'NIA TRIFOLIA'TA, *Spiræ'a trifoliola*, *Common Gille'nia*, *Ind'ian Phys'ic*, *West'ern Drop'root*, *Ind'ian Hip'po*, *Ip'ecac*, *Bow'man's root*, *Mendocino sweet*, *Beaumont root*, (F.) *Gillénie*. *Ord.* Rosaceæ. *Sex. Syst.* Icosandria Pentagynia. The root of this shrub, and of *G. stipulacea*—*Gille'nia*, (Ph. U. S.)—which grows plentifully in the United States, is a safe and efficacious emetic, in the dose of about 30 grains. It resembles *ipeacuanha* in its properties.

GILLÉ'NIE, *Gillenlia trifoliata*.

GILLFLOWER, (*July flower*, or from *Caryophyllum*, [?]) *Dianthus caryophyllus*.

GILLON, *Viscum album*.

GIMBERNAT'S LIGAMENT, (after *Gimbernatus*, a Spanish surgical writer.) A fibrous, triangular expansion, which is detached from the posterior and inner part of the crural arch, and is inserted into the crest of the pubis. This ligament forms the inner part of the superior aperture of the crural canal. It is one of the most frequent causes of strangulation in crural hernia.

GIN, *Hol'lands*, *Gene'ra*, (F.) *Genèvre*. This spirit, which is distilled from corn and juniper berries, or from some substitute for them, is largely used in Great Britain, by the lower classes particularly, and is extremely detrimental to them. It possesses the properties of other spirituous liquors, but is diuretic, and, in popular medicine, is more used than other varieties, in cases of colic or intestinal pain of any kind. *Schiedam Schnapps* is understood to be *Hollands* of good quality. See *Spirit*.

GIN DRINKER'S LIVER, *Liver*, *nutmeg*—*G. Liver*, *Liver*, *nutmeg*.

GINGEMBRE, *Amomum zingiber*.

GINGER, (from *ζγγίβερ*, *zingiber*.) *Amomum zingiber*.

GINGER BEER POWDER, see *Amomum zingiber*.

GINGER, GARDEN, Capsicum—g. Indian, *Asarum Canadense*—g. Jamaica, concentrated essence of, see *Anomum zingiber*—g. Preserved, see *Anomum zingiber*—g. Wild, *Asarum Canadense*.

GINGIBRA'CIUM, (*gingivæ*, 'the gums,' and *brachium*.) A name given to the scurvy, because the gums and arms are chiefly affected by it. It has also been called *Gingipe'dium*, because the lower limbs are in many cases the seat of scorbutic spots. See *Porphyræ nautica*.

GINGIPEDIUM, (*gingivæ*, 'the gums,' and *pes*, gen. *pedis*, 'the foot,') see *Gingibrachium*, *Porphyræ nautica*.

GINGIVÆ, (*gignere*, 'to beget,' because the teeth are, as it were, begotten in them. [?]) The gums, *U'la*, sing. *U'lon*; *U'um*, Carnic'ula, (F.) *Gencivæ*. The portion of the mucous membrane of the mouth which covers the maxillary bones to the level of the alveolar arches. The gums are formed of a red tissue, more or less solid, and of a fibro-mucous nature, which adheres strongly to the necks of the teeth, and transmits, between the roots and their alveoli, a very thin expansion,—the *alveolodental periosteum*. The gums fix the teeth and contribute greatly to their stability. In the aged, after the loss of the teeth, they become fibrous and very solid, and are inservient to mastication.

GINGIVALIS, *Ulet'icus*, *U'licus*. Relating to the gums.

GINGLYMOID, *Ginglymoïdeus*, *Ginglymo'des*, (*γινγλυμος*, 'a ginglymus or hinge,' and *eidos*, 'resemblance,') Resembling a ginglymus or hinge. An epithet applied to joints which resemble a hinge, as a *ginglymoid joint*.

GINGLYMUS, (*γινγλυμος*.) *Cardinamen'tum*, *Car'do*, 'a hinge,' (F.) *Charni'ère*, *Articulation en charnière*. A species of diarthrodial articulation, which only admits of motion in two directions, like a hinge,—as the knee-joint or elbow-joint.

GINSENG, *Panax quinquefolium*—g. Asiatic, *Panax Schinseng*—g. Blue, *Caulophyllum thalictroides*—g. Yellow, *Caulophyllum thalictroides*—g. Horse, *Triosteum*—g. White, *Triosteum*.

GIRAFFE, *Dengue*.

GIRARD-ROUSSIN, *Asarum*.

GIRDLE, ([Sax.] *gýrþel*), *Cingulum*.

GIRL. This seems, formerly, to have been an appellation common to both sexes. Many etymologists deduce the word from the Su. G., Karl, 'a man.' [?] It means a young female, (L.) *Fil'ia*, from *φίλειν*, 'to love'; *Pu'era*, *Pu'ella*, *Pusa*, *Pu'ell'a*; (F.) *Fille*.

GIROFLE, (from *caryophyllum*, [?]) see *Eugenia caryophyllata*.

GIROFLÉE JAUNE, *Cheiranthus cheiri*.

GIROLE, *Sium sisarum*.

GIRON, *Arum maculatum*.

GIRSLE, (*gristle*), *Cartilage*.

GISN, *Anhelatio*.

GISSA, *Cynanche parotidæa*.

GIZZARD, (F.) *Gévier*, *Ventriculus callosus*.

GIZZEN, see *Parturition*.

GLABELLA, (*glaber*, 'smooth,') *Mesophryon*.

GLABELLAD, see *Glabella*.

GLABEL'LAR. An epithet by Dr. Barclay for an aspect towards the glabella. *Glabbellad* is used adverbially to signify 'towards the glabellar aspect.'

GLABRITIES, (*glaber*, 'smooth,') *Calvities*.

GLACE, (from *glacies*), *Ice*.

GLAC'IAL, *Glacia'lis*, (from *glacies*), *Ice*; having the appearance of ice; as '*glacial phosphoric acid*.'

GLACIALE, *Mesembryanthemum crystallinum*.

GLACIALIS, *Glacial*—g. Humor, *Crystalline*.

GLACIES, (from *gelu*, [?]) *Ice*.

GLADI'OLUS, (dim. of *gladius*, 'a sword,'

from the sword-like shape of its leaf,) *Gladiolus communis*—g. *Ceruleus*, *Iris Germanica*.

GLADI'OLUS COMMUNIS, *G. vulg'aris*, *Victoria'lis rotund'a seu femine'a*, *Corn'flag*, (F.) *Glai-eul*. Ord. *Iridiæ*. The root of this plant has been considered aphrodisiac. Applied in cataplasms, it has been extolled against serofulous tumours.

GLADIOLUS LUTETS, *Iris pseudacorus*—g. *Vulgaris*, *G. communis*.

GLADIUS, ('a sword,') *Penis*.

GLAIR, *Albumen*.

GLAIRES, see *Parturition*.

GLAMA, (*γλαμα*) *Lippitudo*, *Chassie*.

GLAND, *Glands*, dim. *Glan'dula*, 'an acorn, a kernel,' *Aden*. The ancient anatomists gave this name to a number of organs of a texture generally soft, and a shape more or less globular, but differing greatly in their nature and functions. They applied it, for instance, 1. To those organs which separate from the blood, any fluid whatever. When such organs were composed of several lobules, united by common vessels, they received the name *conglomerate glands*, as the parotid, pancreas, &c. 2. To the reddish and spongy, knot-like bodies, which are met with in the course of the lymphatics. These they called *conglobate glands*;—see *Ganglion*, (*lymphatic*;) and 3dly and lastly, to various other organs, whose intimate texture and functions are still unknown, as the *Pineal gland*, *Pituitary gland*, *Glands of Pacchioni*, *Thyroid gland*, *Thymus gland*, *Supra-renal glands*, &c. Chaussier restricts the word *gland* to those softish, granular, lobated organs, composed of vessels, and a particular texture, of which there are in the human body, the *lachrymal, salivary, and mammary*, the *testicles*, the *liver*, *pancreas*, and *kidneys*. These *permanent glands*, or *glands with permanent ducts*, are all destined to draw from the blood the molecules necessary for the formation of new fluids, and to convey these fluids externally, by means of one or more excretory ducts. Several glands, besides their excretory ducts, have special reservoirs, in which the fluids, secreted by them, collect, remain for a greater or less space of time, and undergo slight modifications before being evacuated;—such are, the gall-bladder for the liver, the urinary bladder for the kidneys, &c. Each gland has an organization peculiar to it, but we know not the intimate nature of the glandular texture. Malpighi believed that the vessels terminate in small, solid masses, to which he gave the name—*glan'dular grains* or *ac'tini*. In these, he considered, the excretory ducts originate. Ruysch thought that the glands are entirely vascular, and that the excretory ducts are immediately continuous with the *vasa afferentia*, &c. The most accepted view, perhaps, is, that the secreting vessel is distributed on the animal membrane, which lines the minute excretory ducts, and that the secretion is effected from the liquor sanguinis by means of cells.

The term *Glande* (F.) is sometimes appropriated to the tumour formed by inflammation or engorgement of a lymphatic ganglion.

GLAND, *Glands*—g. Accessory, of the Parotid, see *Parotid*—g. Anteprostatic, see *Cowper's glands*—g. Blood, see *Ganglion*—g. Globate, *Conglobate gland*—g. Prostate, *Prostate*—g. Salivary, *abdominal*, *Pancreas*.

GLANDAGE, *Adenophyma*.

GLANDE, *Gland*—g. de Litte, see *Urethra*—g. Pituitaire, *Pituitary gland*—g. Thyroïde, see *Thyroid gland*—g. Vulvaire conglomerée, *Cowper's glands in the female*—g. Vulvo-vaginale, *Cowper's glands in the female*.

GLANDERS, (from *gland*), *Equinia*—g. Farey, see *Equinia*.

GLANDES BRONCHIIQUES, *Bronchial glands*—g. *Céruminéuses*, *Ceruminous glands*—g.

Conglobées, Conglobate glands—*g. de Cowper*, Cowper's glands in the female—*g. de Croissance*, Waxing kernels—*g. Inguinales*, Inguinal glands—*g. de Litte*, see Urethra—*g. de Meibomius*, Meibomius, glands of—*g. de Méry*, Cowper's glands in the female—*g. de Morgagni*, see Urethra—*g. de Pacchioni*, Pacchioni, glands of.

GLANDIFORM CORPUSCLE, Acinus—*g. ganglion*, see Ganglion.

GLANDIUM, Thymus.

GLANDS, see *Quercus Alba*—*g. Accessory*, Cowper's glands—*g. Aggregate*, Peyer's glandulæ—*g. Agminated*, Peyer's glands—*g. Aporic*, Ganglions, glandiform—*g. of Bartholinus*, Cowper's glands in the female—*g. Blind*, Ganglions, glandiform—*g. Blood*, see Ganglion—*g. Brunner's*, see Brunner's glands—*g. Cowper's*, see Cowper's glands—*g. Diaprogenous*, see Perspiration—*g. Ductless*, see Ganglion—*g. of Duverney*, Cowper's glands in the female—*g. Havers's*, Synovial glands—*g. Lenticular*, Lenticulares glandulæ—*g. Lieberkühn's*, Lieberkühn's glands, see Intestine—*g. of Litte*, see Urethra—*g. Military*, Sebaceous glands—*g. of Morgagni*, see Urethra—*g. of Naboth*, Nabothi Glandulæ—*g. Oil*, Sebaceous glands—*g. of Pacchioni*, Glandulæ Pacchioni—*g. Permanent*, see Gland—*g. Peyer's*, Peyer's glandulæ—*g. Renal*, Capsules, renal—*g. Sebaceous*, see Sebaceous glands—*g. Solitary*, Brunner's glands—*g. Sudoriparous*, see Perspiration—*g. Sweat*, see Perspiration.

GLANDS, TEMPORARY, *Glands without permanent orifices*, Glands, that consist of a single primary vesicle or sacculus, which, having elaborated a secretion in its interior, bursts, discharges it, and disappears. The Graafian vesicles and the gland cells afford examples of these.

GLANDS TORRÉFIÉS, ('roasted acorns,') see *Quercus alba*.

GLANDS OF TYSON, Sebaceous glands of Tyson—*g. Vascular*, see Ganglion—*g. of Vesalius*, Bronchial glands—*g. Vulvo-vaginal*, Cowper's glands in the female—*g. of Willis*, Albicantia corpora, Mammillary tubercles.

GLANDULA, (dim. of *glands*, gen. *glandiæ*) Gland—*g. ad Aures*, Parotid—*g. Avicennæ*, Encystis—*g. Bartholiniana*, Sublingual gland—*g. Basilaris*, Pituitary gland—*g. Colli*, Tonsil—*g. Coccygea*, Coccygeal gland—*g. Innominata*, Galeni, Lachrymal gland—*g. Lacrymalis*, Lachrymal gland—*g. Lactifera*, Mammary gland—*g. Mucosa*, Conglobate gland—*g. Pinealis*, Pineal gland—*g. Pituitosa*, Pituitary gland—*g. Riviniana*, Sublingual gland—*g. Salivaris* abdominis, Pancreas—*g. Socia* Parotidis, see Parotid—*g. Thymus*, Thymus—*g. Thyreoidea*, Thyroid gland.

GLANDULÆ ACINOSÆ, see Acinus—*g. Agminated*, Peyer's glandulæ—*g. Articulares*, Synovial glands—*g. Assistentes*, Prostate—*g. Brunneri*, Brunner's glands—*g. Ceruminosæ*, Ceruminous glands—*g. Cervicis* uteri, Nabothi glandulæ—*g. Coronæ* penis, see Sebaceous glands—*g. Cowper's*, Cowper's glands—*g. Duræ* matris, Pacchioni, glands of—*g. Duræ* meningis, Pacchioni, glands of—*g. Inguinales*, Inguinal glands—*g. Intestinales*, Peyer's glandulæ—*g. in agmen* congregatæ Intestinorum, Peyer's glandulæ—*g. Lieberkühni-anae*, Lieberkühn's glands—*g. Littrianæ*, see Urethra—*g. Meibomianæ*, Meibomius, glands of—*g. Muciparæ* racematim congestæ intestinorum, Peyer's glandulæ—*g. Mucosæ* coagminatæ intestinorum, Peyer's glandulæ—*g. Myrtiformes*, Caruncule myrtiformes—*g. Odoriferæ* glandis, see Sebaceous glands—*g. Odoriferæ* penis, see Sebaceous glands—*g. Odoriferæ* Tysoni, see Sebaceous glands—*g. Pacchioni*, Pacchioni, glands of—*g. Peyerianæ*, Peyer's glandulæ—*g. Plexiformes*, Peyer's G.—*g. Prostatæ* mulierum, see Corpus glandulosum mulierum—*g. Sebaceæ*

ciliares, Meibomius, glands of—*g. Sebaceæ* glandis, see Sebaceous glands—*g. Solitaria*, Brunner's glands—*g. Spuriæ*, Peyer's glandulæ, Ganglions, glandiform—*g. Sudoriferæ*, see Perspiration—*g. Suprarenales*, Capsules, renal—*g. Tubulosæ*, see Tubular—*g. Tysoni*, Sebaceous glands of Tyson—*g. Uterinæ*, Utricular glands—*g. Utriculares*, Utricular Glands—*g. Vasculosæ*, see Ganglion—*g. Vesalianæ*, Bronchial glands.

GLANDULAIRE, Glandular.

GLAND'ULAR, *Glandularis*, *Glandulosus*, (F.) *Glandulaire*, *Glanduleux*. Having the appearance, form, or texture of Glands; as a glandular body, a glandular texture, &c.

GLANDULAR SUBSTANCE OF THE KIDNEY, see Kidney.

GLANDULE OF MALPIGHI, see Corpora Malpighiana.

GLANDULEUX, Glandular.

GLANDULO'SOCAR'NEUS. Ruysch gives this epithet to fleshy excrescences which he found in the bladder.

GLANDULOSUS, Glandular.

GLANS, gen. *Glandis*, ('an acorn,') *Balanus*, *Cyltoros*, *Cus'pis*, *Ca'put*, *C. Penis*, *Genita'lis ca'put*, (F.) *Gland*. The extremity of the penis and of the clitoris. The *glans penis* is of a conical, slightly flattened shape. It is continuous with the urethra, which opens at its apex; and is circumscribed by a projecting edge, called the *Coro'na glandis*, behind which is a depression. *Col'um seu Sul'cus retroglandula'ris*, *Fos'sa glandis*. It is covered by a thin mucous membrane; is furnished, at its base, with sebaceous follicles, called *glandulæ odorif'ere Tysoni*, the secretion from which is termed *Smegma prepu'tii*; and can, almost always, be covered by the reflection of the skin, called the *prepuce*. Lastly, it is formed of a spongy texture, susceptible of being thrown into erection.

GLANS, Brnchocele, Gland, Pessary, Suppository.

GLANS CLITO'RIDIS is smaller. It is imperforate, and covered with a sort of prepuce formed by the mucous membrane of the vulva.

GLANS JOVIS THEOPHRASTI, *Fagus castanea*—*g. Pituitum* excipiens, Pituitary gland—*g. Ulmæ*, *Olecranon*—*g. Unguentaria*, *Guilandina moringa*, *Myrobalanus*.

GLAREA, Gravel.

GLASS, (Sax. *glær*,) Vitrum.

GLAUBER'S SALTS, (after R. Glauber, a German chymist,) Soda, sulphate of.

GLAUCEO'DO, gen. *Glaucodinis*, *Glaucoma*.

GLAUCO'MA, gen. *Glaucodinis*, (γλαυκωμα, from γλαυκος, 'sea-green,' and oma.) *Glaucodis*, *Glauc'edo*, *Cataracta* glauc'ea, *Oculus* glauc'ius, *Cæ'cius*, *Phthar'ma* glauc'o'ma, *Parop'sis* glauc'o'sis, *Apoglauc'o'sis*. Amongst the older pathologists, this word was used synonymously with cataract. It is now ordinarily applied to opacity of the vitreous humour or of the tunica hyaloidea, which manifests itself by a grayish, or greenish spot, apparent through the pupil. The diagnosis is generally difficult; and the disease is almost always incurable.

GLAUCOMA WOULHOESI, Cataract.

GLAUCOSIS, (γλαυκωσις,) *Glaucoma*.

GLAYEUL, *Gladiolus communis*—*g. Puant*, Iris foetidissima.

GLECHO'MA HEDERA'CEA, (after γληχων, 'pennyroyal' [?]) *G. hirsu'ta*, *Nep'eta glecho'ma*, *Heder'ula*, *Chamele'ma*, *C. hederacea*, *Calamin'tha hederacea*, *Calamin'tha humil'ior*, *Chamecis'sus*, *Hed'era terre'stris*, *Nep'eta glecho'ma*, *Panace'a pect'oris*, *Ground-ivy*, *Gill*, *Gill-go-by-ground*, *Alch'hoof*, *Robin runaway*, (F.) *Lierre terrestre*, *Terrette*, *Rondote*, *Herbe Saint-Jean* This plant—

Ord. Labiate—has a strong smell, and a bitterish, somewhat aromatic taste. It has been considered expectorant and tonic.

GLECHOMA HIRSUTA, G. hederacea.

GLECHON. (γλεχων.) *Mentha pulegium*.

GLECHONITES, (γλεχωνίτης.) Wine, impregnated with Gledion.

GLEETING (Scl.) Strabismus.

GLEET, (Sax. gleson, 'to glide.') See Gonorrhœa. Sometimes applied to a chronic discharge from any mucous membrane.

GLEME, (γλημε.) *Chuscia*, Lippitudo.

GLENE, (γληνε.) The pupil. The anterior part of the eye. The eyeball. The eye. According to some, the crystalline lens. Also, a glenoid cavity.

GLENGORE (Sc.), Syphilis.

GLENGOUR (Sc.), Syphilis.

GLENITIS, (glene, 'the crystalline lens,' and itis, 'inflammation.') Phacitis.

GLENN'S SPRINGS, see South Carolina, Mineral waters of.

GLENOID, Glene.

GLENOID, *Glenoid'al*, *Glenoid'al*, *Glenoid'al*, *Glenoid'al*, *Glenoid'al*, *Glenoid'al*, *glene*, and *oides*, 'resemblance,' (F.) *Glenoides*, *Glenoidale*, *Glenoidian*. Any shallow, articular cavity, γλην, *gle'ne*, which receives the head of a bone: such as, 1. The *glenoid cavity* or *fossa* of the scapula, *Fossa glenoides*, *Ossoid'gle*, *Acetabulum humeri*, *Cavitas humeri glenoides*, *Entypo'sia*, situate at the anterior angle of the scapula; and articulated with the head of the humerus. 2. The glenoid cavity or fossa of the temporal bone. It is seated between the two roots of the zygomatic process, and receives the condyle of the lower jaw.

GLENOID LIGAMENT, (F.) *Ligament Glenoidien*, is a fibro-cartilaginous ring or *bourrelet*, which seems formed by the expansion of the tendon of the long head of the biceps brachialis, and surrounds the glenoid cavity of the scapula, the depth of which it increases.

GLENOIDALE, Glenoid.

GLIEDY (Scl.) see Strabismus.

GLISCHRASMA, gen. *Glischrasmatia*, *Glischrotes*, (from γλίσχος, (γλα, 'glue,') 'gluey,') Lento, viscidit.

GLISCHROCHOLUS, (γλίσχος, 'gluey,' and χολη, 'bile,') An epithet for excrement which is glutinous and bilious.

GLISCHROTES, (γλίσχροτης.) Glischrasma.

GLISOMARGO, Creta.

GLISEMENT DU LAMBEAU, ('sliding of the flap,') See Flap.

GLISTER, Clyster.

GLOBE, (from *globus*.) Bandage (head)—g. of the Eye, see Eye—g. Flower, *Cephalanthus occidentalis*, *Trollius luteus*.

GLOBI MARTIALES, ('martial balls,') see Ferrum tartarizatum.

GLOBLAIRE PURGATIVE, Globularia alypum.

GLOBULARIA ALYPUM, *Globularia*, *Montpellier Turbith*, (F.) *Globulaire purgative*, *Turbitic blanc*, *Ord.* Globularia. The leaves of this plant are bitter, and have been used in intermittents, and in constipation. See Alypon and Calcifraga.

GLOBULE, *Glob'ulus*, *Sphæ'ron*, *Sphæroid'ion*, *Sphæ'cula*, (dim. of *globus*, 'a ball,') A small globe.

GLOBULES, RED, OF THE BLOOD, *Blood glob'ules*, *Blood corpuscles*, *Blood disks*, *Blood vesicles*, *Globuli* seu *Vesiculae* seu *Sphæ'culæ* seu *Falliculi* seu *Corpuscula* seu *Cellulae* seu *Particulae* seu *Granula sanguinis*, (F.) *Globules* ou *Cellules rouges du sang*, *Disques sanguins*, *Hématies* of Robin, are small bodies, which are particularly observable when the transparent parts of cold-blooded ani-

mals are examined by the aid of the microscope; and are met with in the blood of all animals. They are circular in the mammalia, and elliptical in birds and cold-blooded animals; are flat in all animals, and generally composed of a central nucleus enclosed in a membranous sac. Chemically, they consist of hæmatin and globulin—*hæmato-glob'ulin*. The chemical composition of the nucleus, where it exists,—it does not in man,—has been by some regarded as fibrin; by others, as fat; and by others, again, as a species of horn, to which the name *ne'lein* has been given. See Globulin. Their average long diameter in man is about $\frac{1}{3000}$ th part of an inch.

SIZE OF THE GLOBULES, ACCORDING TO

Sir E. Home and Bauer, with colouring matter.....	1.1700th part of an inch.
Eller.....	1.1930
Sir E. Home and Bauer, without colouring matter.....	1.2000
Müller.....	1.2000 to 1.3500
Marsili.....	1.2025 to 1.3150
Hedekin. Lister, and Rudolphi.....	1.3000
Sprengel.....	1.3000 to 1.3500
Cavallo.....	1.3000 to 1.4000
Donné.....	1.3150 to 1.3250
Jurin and Gulliver.....	1.3240
Blumenbach and Senac.....	1.3350
Tabor.....	1.3600
Milne Edwards.....	1.3600
Wagner.....	1.4000
Kater.....	1.4000 to 1.6000
Prevost and Dumas.....	1.4056
Haller, Wellaston, and Weber.....	1.5000
Young.....	1.6000

Besides these, the blood contains *white* or *pale globules*, (F.) *Globules blancs*, *Leucocytes*, which are probably lymph and chyle corpuscles in process of development into red corpuscles.

Dried blood globules of the ox have been given internally in cases of anæmia.

The minute *pellets* of the homœopathist are called *globules*.

GLOBULE D'ARANTUS, see Sigmoid valves—g. du Sang, Globule of the blood.

GLOBULES BLANCS, see Globules of the blood—g. Chyle, see Chyle—g. Ganglion, see Neurine—g. Lymph, see Lymph—g. Milk, see Milk—g. Mucous, see Mucus—g. Pale, see Globules of the Blood—g. Pus, see Pus—g. Proid, see Pus—g. White, see Globules of the Blood.

GLOBULI ARTERIARUM TERMINI, see Acinus—g. Mammillares, see Mammillary eminence—g. Sanguinis, Globules of the blood—g. Tartari martiales, Ferrum tartarizatum—g. Tartaris ferri et lixivie, Ferrum tartarizatum.

GLOBULIN, *Glob'uline*, *Blood co'sein*. The colourless substance that remains after the abstraction of the colouring matter of the blood-globule. It is a peculiar albuminous principle. The globulin of Berzelius consists of the envelopes of the blood globules, and of the part of their contents that remains after the extraction of the hæmatin. Lecanu regards it as identical with albumen; and, according to Mulder, it belongs to the combinations of protein. Robin and Verdel consider it to be albuminous.

The term *globulin* is likewise given by M. Donné to small granulations appertaining to the chyle, which are observable in the blood with the microscope. They are small, white, roundish, isolated or irregularly agglomerated grains; of about the 1-300 of a millimètre in diameter, and are regarded by M. Donné as the first elements of the blood-globules. They are the *white granulated corpuscles* of Mandl.

GLOBULISM, (from *globule*.) Homœopathy.

GLOBULUS, (dim. of *globus*.) Globule—g. Arantii, see Sigmoid valves—g. Hysterici, Angone—g. Nasi, see Nasus—g. Sanguineus, Punctum saliens—g. Stapedius Ossis, Os orbiculare.

GLOBUS. A ball or globe.

GLOBUS HYSTERICUS, Angene—g. Major, see Epithymis—g. Minor, see Epithymis.

GLOBUS UTERINUS. A term applied by accoucheurs to the round tumour, formed by the uterus in the lower part of the abdomen immediately after delivery.

GLOMERATE, (from *glomus*, gen. *glomeris*, 'a ball.') Conglomerate.

GLOMERULE. *Glomerulus*; (dim. of *glomus*, 'a ball.') A small ball or clew, formed by an agglomeration of vessels; as *Glomerule* of Malpighi; see *Corpora Malpighiana*.

GLOMERULI RENALES, *Corpora Malpighiana*.

GLOXOÏN, *Gloxoin*, *Gloxoinum*, *Nitroglycerin*. A substance prepared by adding glycerin to a mixture of sulphuric acid and fuming nitric acid; pouring it into water and washing upon a filter. It is an oleaginous substance, of a sweet taste, little soluble in water, but readily in alcohol and ether. A drop, brought in contact with the lips, or even the vapour, is said to have caused the most distressing headache; hence it has been used in headache by the homeopaths.

GLOSSA, (γλῶσσα,) *Glotta*, 'the tongue.' The power of speech. Speech.

GLOSSAGRA, (*glossa*, and *αγρα*, 'a seizure.') Glossalgia.

GLOSSAL, *Glossalis*, (from *glossa*.) Relating or appertaining to the tongue.

GLOSSALGIA, *Glossagra*, *Glottalgia*, *Glottagra*, (*glossa*, and *αλγος*, 'pain.') Pain in the tongue.

GLOSSANISCHUM, *Glossanochus*, *Glossanochum*, (*glossa*, and *αρχη*, 'to hold up.') An instrument for holding up the tongue.

GLOSSANOCHUS, *Glossanischum*.

GLOSSANOCHUM, *Glossanischum*.

GLOSSANTHRAX, *Pestis glossanthrax*, (F.) *Bonsode*, (*glossa*, and *ασπς*, 'a carbuncle.') Carbuncle of the Tongue. A disease more common in cattle than in man.

GLOSSEPIGLOTTIC, *Glossepigloticus*. Relating to the tongue and epiglottis, as *Ligamentum glossepigloticum*.

GLOSSIANUS, Lingual muscle.

GLOSSINA MORITANS, *Tsetse*. An insect of South Africa, not much larger than the common housefly, the bite of which, according to Dr. Livingstone, is death to the ox, horse and dog; whilst it is harmless to man and wild animals, and even to calves so long as they suck.

GLOTTIS, (*glossa*, and *itis*.) *Glossitis* inflammatorius, *Angina lingua'ria* seu *lingualis*, *Inflammatio Linguae*, Inflammation of the tongue. (F.) *Inflammation de la Langue*. When confined to the mucous membrane, it is of slight importance. That which affects the whole of the tongue is a serious disease, and requires the vigorous use of antiphlogistics. It is rare.

GLOSSO, (γλῶσσα, 'the tongue.') In composition, relating or appertaining to the tongue.

GLOSSOACE, (*glossa*, and *κακος*, 'evil.') Ulceration of the tongue, with symptoms of adynamic fever.

GLOSSOCARCINOMA, gen. *Glossocarcinomatidis*, *Glossocarcinoma*, *Carcinoma Linguae*, (*glossa*, and *καρκινωμα*, 'cancer.') Cancer of the tongue.

GLOSSOCATACHUS, *Glossocatichê*, (γλῶσσα κατὰ χεῖρα,) *Cataglossum*, *Cataglossum*, (*glossa*, and *κατεχεσθαι*, 'I arrest.') *Linguae Detentor*, *Tongue-holder*, (F.) *Absaisseur de la langue*. An instrument, the invention of which is attributed to Paulus of Aegina, and which was employed to depress the tongue, in order to examine diseases of the fauces. It was composed of two branches; one of which had, at its extremity, a plate for depressing the tongue; whilst the other, shaped like a horse-shoe, was applied under the chin. The

finger, or the handle of a spoon, or a spatula, is generally used in similar cases.

GLOSSOELE, (*glossa*, and *ελη*, 'hernia,' 'tumour.') *Hernia of the Tongue*, *Paraglossa*, *Protopharyngeal*, *Glossomegaly*, *Glossoplasia*. Projection of the tongue from the mouth. It depends, generally, on an inflammatory swelling of the organ. At times, however, a chronic *glossocoele*, or sort of oedematous engorgement, is met with; which proceeds to a great length, and deforms the dental arches, the lips, &c. Inflammatory *glossocoele* must be combated by antiphlogistics. In the oedematous kind, such as is sometimes caused by excessive salivation, the infiltrated fluid may be pressed back by the hand of the practitioner, to get the tongue behind the teeth; and it may be kept there by a piece of gauze tied over the mouth. The chronic, elongated kind sometimes requires amputation of a portion of the organ.

GLOSSOCOMA, (*glossa*, and *coma*.) *Glossospasmus*.

GLOSSOCOMON, *Glossocomum*, *Glossocomion*, (γλωσσόκομοιον, from *glossa*, and *κομω*, 'I guard.') The ancients gave this name to a small case for holding the tongues of their wind instruments. By extension, it was applied to the box or cradle in which fractured limbs were kept. We find in the old writers, a *Glossocomon* of Hippocrates, of Nymphodorus, Galen, &c.

GLOSSODESMUS, (*glossa*, and *δεσμος*, 'a bandage.') *Frænum lingue*.

GLOSSEPIGLOTTIC, *Glossepigloticus*. That which belongs to the tongue and epiglottis. Some anatomists have so denominated certain fleshy fibres, which pass from the base of the tongue towards the epiglottis. These muscles are more evident in some of the mammalia than in man; and their use seems to be,—to raise the epiglottis, and to remove it farther from the glottis. Santorini, who described them after Eustachius, calls them *Retractor Epiglottidis*.

GLOSSEPIGLOTTIC LIGAMENT, *Frænum* seu *Frænulum epiglottidis*, *Ligamentum membranaceum*, *Velamentum linguae*. Three folds of mucous membrane, which connect the anterior surface of the epiglottis with the root of the tongue.

GLOSSOGRAPHY, *Glossographia*, (*glossa*, and *γραφη*, 'a description.') An anatomical description of the tongue.

GLOSSOHYAL. A name given, by Geoffroy Saint-Hilaire, to the posterior cornua of the os hyoides.

GLOSSOLOGY, *Glossologia*, (*glossa*, and *λογος*, 'a treatise,' 'a discourse.') A treatise on the tongue. Nomenclature.

GLOSSOLYSIS, *Glossolytic*, (*glossa*, and *λυσις*, 'solution.') Paralysis of the tongue.

GLOSSOMANTIA, *Prognosis ex lingua*, (*glossa*, and *μαντεία*, 'divination.') Prognosis from the state of the tongue.

GLOSSOMEGISTUS, (*glossa*, and *μεγιστος*, 'the greatest,') *Glossocoele*, *Paraglossa*.

GLOSSONCUS, (*glossa*, and *ογκος*, 'tumour.') *Eumecosis linguae*. Swelling of the tongue.

GLOSSONCUS INFLAMMATORIUS, *Glossitis*.

GLOSSOPALATINUS, *Glossostaphylinus*.

GLOSSOPHARYNGEAL, *Glossopharyngeus*, *Pharyngo-glossal*—g. p. Nerve, *Pharyngo-glossal nerve*.

GLOSSOPHARYNGEUS, *Glossopharyngeal*. Belonging to the tongue and pharynx. Some anatomists thus designate certain fleshy bundles, which arise from the lateral parts of the base of the tongue, and are inserted into the parietes of the pharynx. They form part of the constrictor pharyngis superior.

GLOSSOPLÉGIA, (*glossa*, and *πληγη*, 'a stroke,') *Glossolysis*.

GLOSSOPTOSIS, (γλωσσο, and πωσις, 'a falling,') Glossocele.

GLOSSOSCIRRHUS, Glosso-carcinoma.

GLOSSOSCOPIA, (glossos, and σκοπεω, 'I view,') Inspection of the tongue as an index of disease.

GLOSSOSEMIOTICÆ, (glossos, and semeiotice, 'semeiology,') The signs of disease taken from the tongue.

GLOSSOPAS'MUS, Glosso-pas'ma, Spasmius lin'gue, (glossos and σπασμος, 'spasim,') Cramp or spasm of the tongue.

GLOSSOSPAT'HA, Spat'ula pro o'rē, (glossos, and σπαθη, 'spatula,') A spatula for pressing down the tongue to enable the fauces to be examined.

GLOSSOSTAPHYL'NUS, (glossos, and σταφυλη, 'the uvula,') Glossopalati'nus, Pala'toglossus, Constrictor Isthmi Fau'cium. A small, thin, narrow, and long muscle, which arises from the base of the tongue, and is inserted at the inferior and lateral part of the velum palati, in the anterior pillar of which it is situate. Its use is to contract the isthmus faucium, by depressing the velum palati, and raising the base of the tongue.

GLOSSOSTERE'SIS, Lingue Extirpa'tio, (glossos, and στερησις, 'privation,') Extirpation of the tongue.

GLOSSOSTROPH'IA, (glossos, and στρεφω, 'I turn,') Doubling of the point of the tongue upwards and backwards,—said to have been a mode of suicide. [?]

GLOSSOT'OMY, Glossotom'ia, (glossos, and τεμνειν, 'to cut,') Dissection of the tongue. Amputation of the tongue.

GLOSSYPERTROPH'IA, (glossos, υπερ, 'over, and τρεφειν, 'to nourish,') Hypertrophy or super-nutrition of the tongue.

GLOTTA, (γλωττα,) Glossa, Tongue.

GLOTTAGRA, (glotta, and αγρα, 'a seizure,') Glossalgia.

GLOTTALGIA, (glotta, and αλγος, 'pain,') Glossalgia.

GLOTTICUS, (γλωττικος,) Pertaining to the tongue.

GLOTTIS, gen. Glott'idis, (γλωττις, 'the mouth-piece of a flute,') Lig'ula, Lin'gula. A small oblong aperture in the larynx, comprised between the chordæ vocales. It is narrow, anteriorly; wider, posteriorly; and is capable of being modified by muscular contraction, as may be required by the voice. It is by the chordæ vocales, that voice is produced. The glottis is nearly an inch long in the adult male; less in the female and child. Rima Glottidis being the aperture in which the voice is produced, and therefore synonymous with Glottis, as given above.

Glottis is, by some, used synonymously with ventricle of the larynx: with others, it includes the whole of the larynx.

GLOTTIS, LIPS OF THE, Thyreo-arytenoid ligaments—g. Spasm of the, Asthma thymicum.

GLOUGLOU D'UNE BOUTEILLE, (formed by onomatopoeia,) Gurgling.

GLOUTERON, Arcium Lappa—g. Petit, Xanthium.

GLOUTIUS, (from γλουτος, 'the breech,') Gluteal—g. Maximus et extimus, Gluteus maximus—g. Secundus et medius, Gluteus medius—g. Tertius et intimus, Gluteus minimus.

GLOUTON, Glutton.

GLOWWORM, Cicindela.

GLUANT, Glutinous.

GLUCÆMIA, (γλυκς, 'sweet, and αίμα, 'blood,') Glycæmia.

GLUCOGENESIS, Glycoëny.

GLUCOGENIA, Glycoëny.

GLUCOHÆMIA, Glycæmia.

GLUCOSE, better Gly'cose, (γλυκς, 'sweet,') Grape, Fruit, Starch, and Honey sugar. A variety of sugar, that occurs naturally in many vegetable juices, and in honey. Compared with cane sugar,

it is much less soluble in water, and less disposed to crystallize: and, when injected into the blood-vessels, does not pass off to the like extent by the kidneys. Diabetic, Urinary, and Hepatic sugar appear to have the same chemical composition as Glucose.

GLUCOSIQUE, Saccharine.

GLUCOSURIA, (γλυκς, 'sweet, and ουρον, 'urine,') Diabetes mellitus.

GLUEBONE, Osteocolla.

GLUE, FISH, Ichthyocolla.

GLUMPOT, Gallipot.

GLUTEUS MAGNUS, (γλουτος, 'the buttock, 'the breech,') G. maximus—g. Major, G. maximus.

GLUTEUS MAX'IMUS, G. Major seu magnus, Maximus et ex'timus Glut'ius, (F.) Musculus grand fessier, Ilio-sacro-fémoral; Sacro-fémoral, (Ch.)

This muscle is situate at the posterior part of the pelvis, and at the upper and posterior part of the thigh. It is large, thick, and quadrilateral; and is attached, above, to the posterior part of the crista ili, to the part of the ilium comprised between the crista and the upper curved line, to the posterior surface of the sacrum, coccyx, and great sacro-sciatic ligament; and below, it terminates by a broad aponeurosis, which is inserted into the rugged surface that descends from the trochanter major to the linea aspera of the femur. This muscle extends the thigh on the pelvis, and rotates the thigh outwards. It is greatly concerned in station and progression.

GLUTEUS ME'DIUS, Glut'ius Secun'dus et Mé'dius, (F.) Ilio-trochantérien: Grand Ilio-trochantérien, (Ch.,) Moyen Fessier. This muscle is situate in part beneath the preceding; it is broad, very thick, radiated, and triangular; attached, above, to the crista ili, and to the part of the outer surface of that bone comprised between the three anterior fourths of its crista, its upper curved line, and its lower; and below, it ends by a tendon, inserted at the upper edge of the great trochanter. It is an abductor of the thigh; but can turn the thigh outwards or inwards, according as its posterior or inferior fibres are thrown separately into contraction.

GLUTEUS MIN'IMUS, G. mi'nor, Ter'tius et In'timus Glut'ius, (F.) Petit Fessier:—Ilio-ischii-trochantérien, Petit Ilio-trochantérien (Ch.) This muscle, which is situate beneath the preceding, is flat, triangular, and with radiated fibres. It is attached, above, to the external surface of the os ili, from the inferior curved line to the acetabulum; and, below, is inserted into the anterior part of the great trochanter. It has the same uses as the preceding.

GLUTEUS MINOR, G. minimus.

GLUTEAL, or GLUTEAL, Glut'ius, Glute'us, (F.) Fessier. That which belongs or relates to the nates. This name has been given to many parts which compose the nates.

GLUTEAL APONEURO'SIS. The upper and back part of the femoral fascia. In it is a remarkable opening, called the gluteal arch, for the passage of the gluteal vessels and nerves.

GLUTEAL ARCH, see Gluteal aponeurosis.

GLUTEAL ARTERY, Poste'rior Iliac Artery, (F.) Artère fessière, is one of the largest branches of the hypogastric. It makes its exit from the pelvis at the upper part of the superior sciatic foramen; gains the posterior part of the pelvis, and divides into two branches;—the one superficial, the other deep-seated. The last subdivides into three secondary branches, whose ramifications are distributed particularly to the Glutei, Longissimus Dorsi, Sacro-lumbalis, &c., and anastomose with the sciatic and internal circumflex arteries.

GLUTEAL NERVE, (F.) Nerf Fessier, is a large branch, furnished by the 5th pair of lumbar nerves. It is chiefly distributed to the glutæi muscles.

GLUTEAL VEIN, (F.) *Veine fessière*, follows the same march as the artery of the same name.

GLUTEN, gen. *Glutinis*, ('glue, paste,') *Glutinium*, *Lentor*, *Veg'etabile Gluten*, *Veg'etabile Casein*. An immediate principle of vegetables. It is soft, of a grayish white, viscid consistence, and very elastic. Exposed to the air, it becomes hard, brown, and fragile; and, in moist air, putrefies. Water and alcohol do not dissolve it. It is soluble in vegetable, and in weak mineral, acids, at a high temperature. The farinæ, in which it is found, are those preferred for the preparation of bread; on account of the property it has of making the paste rise. It is a compound of protein, and hence has been ranged amongst the "*proteinaceous alimentary principles*" by Dr. Pereira. By washing wheaten dough with a stream of water, the gum, sugar, starch, and vegetable albumen are removed; the ductile, tenacious, elastic, gray mass left is the gluten, *common gluten*, *Beccari's gluten*. Pure gluten is the soluble portion on boiling common gluten in alcohol. *Granulated gluten*, *Gluten granule*, is a paste made by adding wheat gluten to ordinary wheat. It is said to form an agreeable and nutritious aliment.

GLUTEN ARTICULORUM, *Synovia*—g. *Beccaria's*, see Gluten—g. Bread, see Bread, gluten—g. Common, see Gluten—g. Granulated, see Gluten—g. Granule, see Gluten—g. Pure, see Gluten.

GLUTI, (γλουτοι,) Nates.

GLUTIA, (γλουτια,) Nates, *Quadrigenina corpora*.

GLUTIN, see Gelatin.

GLUTINANS, (*glutinare*, *glutinatum*, 'to glue,') Agglutinant.

GLUTINATIF, Agglutinant.

GLUTINATIO, Agglutination.

GLUTINEUX, Glutinous.

GLUTINOUS, *Glutinosus*, *Colloides*, (*gluten*, 'paste, glue,') (F.) *Glutineux*, *Gluant*. An epithet given to substances taken from the animal or vegetable kingdom, and endowed with unusual viscosity. The decoctions of marsh-mallows, and figs, and the jelly of hartshorn, are said to be glutinous.

GLUTINUM, Gluten.

GLUTITIS, (γλουτις, 'the nates,' and *itis*.) Inflammation of the nates.

GLUTOI, (γλουτοι,) Nates.

GLUTOPYOSIS, (γλουτοι, 'the nates,' and *πυωσις*, 'suppuration,') Suppuration of the nates.

GLUTTON, *Gluto*, *Gluto*, (*glutire*, 'to gulp down,') *Gastrimar'gus*, *Gna'thon*, *Hel'uo*, *Man'do*, *Gulo'sus*, *Lur'co*, *Bellygod*, (F.) *Glouton*, *Gourmand*, *Goulon*. An excessive eater.

GLUTTONY, *Limo'sis*, *Helluo'nium*, *Gastris'mus*, (F.) *Gourmandise*. Excessive appetite, owing often to habitual indulgence.

GLUTTUPATENS, (*gluttus*, and *patere*, 'to extend,') Stomach.

GLUTTUS, *Œsophagus*.

GLUTUS, (sing. of *gluti*, 'the nates,') Trochanter major.

GLYCÆMIA, Glycœmia.

GLYCAMIL, (from *glycerin*, and *amylum*), Plasma.

GLYCANSIS, (γλυκανσις, from γλυκός, 'sweet,') Edulcoration.

GLYCAS'MA, gen. *Glycasm'atie*, (γλυκός, 'sweet,') A sweet wine, prepared from must.

GLYCERATON, (γλυκερος, 'sweet,') Glycyrrhiza.

GLYCERIN, (γλυκός, 'sweet,') see Fat—g. Butyrate of, see Butter.

GLYCEROLE, (F.) *Glycérolé*. A combination of glycerin with some other remedial agent. See Plasma.

GLYCINE APIOS, (from *glyco*,) *Apios tuberosa*.

GLYCIPICROS, (*glyco*, and *πικρός*, bitter.) *Solanum dulcamara*.

GLYCO, GLYCY, (from γλυκός, 'sweet,') in composition, sweet.

GLYCOENE, Glycogenic matter.

GLYCOGEN, Glycogenic matter.

GLYCOGEN'IC, *Glycogen'icus*. Relating or appertaining to the formation of sugar.

GLYCOGENIC MATTER, *Gly'cogen*, *Gly'cocene*, *Hep'atine*, *Am'ylo'n*, *Am'yloid*, *Liverstarch*. The peculiar substance in the liver, or elsewhere, which is converted into sugar. In chemical qualities it is said to be the intermediate between starch and dextrin, and to pass readily into sugar in the presence of any animal ferment.

GLYCOG'ENY, *Glycogen'in*, *Glycogen'exis*, *Glycogen'ia*, *Glycogen'exis*, (F.) *Glycogénie*, (*glyco*, and *γενεσις*, 'generation,') Generation or formation of sugar,—as in the liver.

GLYCOHÆMIA, *Glycœhæ'mia*, *Glycæ'mia*, *Glycæ'mia*, (*glyco*, and *hæma*, 'blood,') A saccharine condition of the blood.

GLYCOSE, Glucose.

GLYCOSIQUE, (from *glyco*,) Saccharine.

GLYCOSURIA, (*glyco*, and *ουρον*, 'urine,') *Diabetes mellitus*.

GLYCY, in composition, see Glyco.

GLYCYPHYTON, (*glycy*, and *φυτον*, 'a plant,') Glycyrrhiza.

GLYCYRRHI'ZA, (*glycy*, and *ρίζα*, 'a root,') *G. Gla'bra* seu *Læ'vis*, *Liquorit'ia*, *Scyth'ica*, *Glycera'ton*, *Glycyph'yton*, *Adip'os*, *Al'imos*, *Liq'norice*, *Lic'orice*, (F.) *Régliasse*, *Bois doux*. *Ord. Leguminosæ. Sez. Syst. Diadelphia Decandria*. The root of this southern European plant is inodorous; has a sweet taste; is mucilaginous; and leaves, when unpeeled, a degree of bitterness in the mouth. It is used as a demulcent, and chiefly in catarrh. The extract, made from it and sold in the shops, is known under the name *Spanish Liquorice* or *Liquorice Juice*, (F.) *Jus de Régliasse*. The sweet principle has been called *Soc'charo-gum'mita*.

Pectoral Balsam of Liquorice—a quack preparation—is said by Dr. Paris to consist chiefly of *Paregoric Elixir*, strongly impregnated with *Oil of Aniseed*.

GLYCYRRHÆ'A, (*glycy*, and *ῥέω*, 'I flow,') A discharge of saccharine fluid from the system.

GLYCYRRHÆA URINOSA, *Diabetes mellitus*.

GLYSTER, Clyster.

GLYSTER HERBS, *Her'bæ pro Enem'atē*. The herbs ordinarily sold by the English apothecary under this title, are:—*mallow leaves*, one part; *chamomile flowers*, one part. (℞iss of these to Oj of water.)

GNAMPSIS, Curvature.

GNAPHALUM DIOICUM, (γναφαλον, 'a lock of wool,' owing to the floccose down of the leaves.) *Antennaria dioica*.

GNAPHALUM MARGARITÆCEUM, *Antennaria Margaritæ'cea*, *Cud'weed*, *Pearly Life everlasting*. An indigenous plant, growing in woods and fields, and flowering in August. Its virtues are not defined, and the same may be said of

GNAPHALUM PLANTAGINIFOLIUM seu *Plantagineum*, *Antennaria plantaginifolia*; and

GNAPHALUM POLYCEPH'ALUM, *Sweet-scented Life everlasting*, *Catfoot*. They appear to be mild astringents.

GNATHALGIA, (*gnatho*, and *αλγος*, 'pain,') *Neuralgia maxillaris*.

GNATHANCYLO'SIS, (*gnatho*, and *αγκυλωσις*, 'stiffness of joint,') *Ankylosis* of the lower jaw.

GNATHANTROPOL'YPUS, (*gnathantrum*, 'the antrum of Highmore,' and *poly'nus*,) *Polypos* of the antrum of Highmore.

GNATHANTRUM, (*gnatho*, and *αντρον*, 'antrum,') *Antrum* of Highmore.

GNATHITIS, *Inflamma'tio ge'næ*, (*gnatho*, and *itis*,) *Inflammation* of the cheek or upper jaw.

GNATHMUS, (γνάθος,) Gnatbus.

GNATHO, (γνάθος,) in composition, the jaw or cheek.

GNATHOCEPHALUS, (gnatho, and κεφαλή, 'head.') A monster which has no head visible externally, but exhibits voluminous jaws.

GNATHOCYNANCHE, Cynanche parotidæa.

GNATHON, (γνάθος,) Glutton.

GNATHONEURALGIA, Neuralgia maxillaris.

GNATHOPLASTY, *Gnathoplasticæ*, (gnatho, and πλαστικός, 'formative.') The formation of an artificial cheek.

GNATHOPLEGIA, *Gnathoparalysis*, (gnatho, and πλῆγη, 'a stroke.') Paralysis of the cheek. *Gnathoparalysis* is employed by Fuchs to signify paralysis of the lesser portion of the trifacial nerve, which supplies the muscles of mastication.

GNATHORRHAGIA, (gnatho, and ῥέω, 'to burst forth.') Hemorrhage from the internal surface of the cheeks.

GNATHOSPASMUS, Trismus.

GNATHUS, (γνάθος,) *Gnathmus*, (gnaw, 'I scrape, rub.') The cheek, the jaw. Also, the part of the jaws in which the teeth are fixed. See Bucca, Genæ, and Maxillary Bone.

GNESIUS, (γνήσιος,) Legitimate.

GNOME, (γνώμη,) Intellect.

GNOSIS, (γνώσις,) 'knowledge.' A common suffix, as in *Diagnosis*, *Prognosis*, &c.

GOACONAX, see *Toluifera balsamum*.

GOADBY'S SOLUTION, see *Solution*, *Goadby's*.

GOATS' BEARD, COMMON, *Tragopogon*.

GOATS' MILK, see *Milk*, *goats'*—g. Milk, artificial, see *Milk*, *goats'*—g. Thorn, *Astragalus verus*.

GOB, Mouth.

GOBELET EMÉTIQUE, Goblet, emetic.

GOBLET, EMETIC, *Poculum emeticum seu vomitorium*, *Calix vomitorius*, (F.) *Gobelet émetique*. A vessel, made by pouring melted antimony into a mould. By putting wine into this and allowing it to stand some time, it acquires the property of producing vomiting. This kind of emetic has been long rejected, as the practitioner could never be certain of the dose he exhibited.

GOD'S MARK. A mark, formerly placed on houses, in England, as a sign that the plague was there.

GOD'BOLD'S VEG'ETABLE BAL'SAM. A quack medicine, said to consist chiefly of simple oxymel.

GODESBERG, MIN'ERAL WATERS OF. These waters, at Godesberg, a German mile from Bonn, are an efficacious, acidulous chalybeate, formerly known by the name, *Draitsch Water*. They contain chloride of sodium, carbonate of soda, carbonate of lime, carbonate of magnesia, and carbonate of iron.

GODFREY'S COR'DIAL. A celebrated nostrum, for which Dr. Paris has given the following formula: Infuse $\frac{3}{4}$ of *sassafras*, and of the seeds of *carragee*, *coriander*, and *anise*, each $\frac{3}{4}$, in six pints of water. Simmer the mixture until reduced to fiv; then add lbvj of *treacle*, and boil the whole for a few minutes. When cold, add $\frac{3}{4}$ of *tincture of opium*. The following form is recommended by a committee of the Philadelphia college of Pharmacy: *Tinct. Opii Oiss; Syrupi Nigri Oxxj; Alcoholis Oij; Aquæ Oxxvj; Carbonatis Potassæ ʒijss; Olei Sassafras fʒiv. M.* It is anodyne.

GODRONNÉ (CANAL), *Canalis Petitiæ nua seu Petiti, Couronne citraire, Canal de PETIT ou godronné, Canal or Bul'ular Canal of Petit*. Petit gave this name (from [F.] *godron*, 'a plait or fold,') to the semicircular canal, formed by the tunica hyaloidea around the edge of the crys-

talline; because it appears plaited or festooned as it were.

GOËTRE, (from guttur, 'throat,' [?]) *Bronchocele*.

GOGGLE-EYED, *Exophthalmus*.

GOITRE, (from guttur, 'throat,' [?]) *Bronchocele*—g. *Exophthalmic*, see *Exophthalmia*—g. Leaf, see *Laminaria*.

GOITRE STICKS. In South America the stems of a seaweed are so called, because, according to Dr. Royle, they are chewed by the inhabitants where goitre prevails.

GOITROUS, (F.) *Goitreux*. Relating or appertaining to goitre. One affected by goitre or Bronchocele, *Goitred*.

GOLD, (Sax. *golo*.) *Aurum*, *Chryseus*, *Sol*, *Rex metallo-rum*, (F.) *Or*. A solid, yellow, very brilliant, hard, very ductile, malleable, tenacious, and heavy metal; found in nature either in its native state or combined with a little silver, copper, or iron; s. g. 19.25.

Muriate or *Chloride of Gold*, *Aur'i Chloridum* seu *Terchloridum* seu *Murius* seu *Chloruretum*, *Aurum muriat'icum* seu *Chlorat'um* seu *Oxydulatum muriat'icum* seu *Solutum*, (F.) *Chlorure ou Muriate ou Hydro-chlorate d'or* has been admitted into the Pharmacopœia of the United States, and into that of Paris, &c. The formulæ, however, differ. That of the United States is a muriate with two bases; and is prepared, according to the form of Dr. Christien, by dissolving the gold in a mixture of nitric and muriatic acids, and adding chloride of sodium to the residuum after evaporation; then redissolving and evaporating slowly to dryness. The Parisian formula for the *Muriate d'or*, *Murius* seu *Chloruretum Aur'i*, consists in simply dissolving the gold in the acids, and evaporating to dryness. It has been recommended as an antisypilitic in old, rebellious, venereal affections, exostoses, and in venereal, scrofulous, or cancerous glandular enlargements. Dose, gr. 1-8th to gr. ss, rubbed on the tongue or gums. Internally, one-sixteenth of a grain, in pills.

Various other preparations, as the *Cy'anide* or *Tercy'anide*, (*Aur'i Cyanidum* seu *Cyanuretum* seu *Tercyanidum*, [F.] *Cyanure d'or*;) the metallic gold in a state of division, (*Aurum metallicum*, *Pulvis Aur'i*, [F.] *Or divisé ou métallique*;) obtained by amalgamating gold with mercury, and driving the latter off by heat, and in the form of filings (*Aurum limatum*); the *Chloride of Gold* and *Sodium*, (*Aurum muriat'icum natronatum* seu *muriat'icum* seu *chloratum natronatum*, *Chloruretum Aur'icum Chloro-to natron*, *Murius Aur'ico-natricum*, *Chloruretum auri et natrii*, *Sol'dii auro-terchloridum*, *Hydrochlorate ou muriate of Gold and So'da*, [F.] *Chlorure d'or et de Sodium*, *Hydrochlorate ou muriate d'or et de Soude*); the *Nitromuriate of Gold*, (*Aurum Nitro-muriat'icum*, *Auri nitromuriæ*, [F.] *Nitromuriate d'or*); the *Oxide of Gold*, (*Aur'i Oxidum*, *Aurum Oxidatum*, *Auri teroxidum*, *Peroxi'de of gold*, *Auric acid*, [F.] *Oxide d'or*, and the *Iodide of Gold*, (*Auri Iodidum* seu *Ioduretum*, [F.] *Iodure d'or*), have been employed in the like affections, and with similar results.

GOLDBEATERS' SKIN. The intestina recta of the ox, which have been beaten quite smooth for the manufacture of gold leaf. Used as a defensive dressing for slight cuts, &c.

GOLD, CHLORIDE OF, see *Gold*—g. *Cyanide* of, see *Gold*—g. *Hydrochlorate* of, see *Gold*—g. *Iodide* of, see *Gold*.

GOLD LEAF, *Gold Foil*, *Aurum foliatum*, *Aurum in libellis*. Used to gild pills and to plug carious teeth. The gold leaf or foil used by the dentist for filling teeth is much thicker than ordinary gold leaf.

GOLD, MURIATE OF, see *Gold*—g. *Nitro-muriate*

of, see Gold—g. Oxide of, see Gold—g. Peroxide of, see Gold—g. and Soda, hydrochlorate of, see Gold—g. and Soda, muriate of, see Gold—g. and Sodium, Chloride of oxide of, see Gold—g. Tereyanide of, see Gold.

GOLD THREAD, Coptis.

GOLDEN BUG, *Coccinella septempunctata*—g. Knop, *Coccinella septempunctata*—g. Rod, *Solidago virgaurea*—g. Rod, fragrant, *Solidago odora*—g. Rod, rigid, *Solidago rigida*—g. Rod, sweet-scented, *Solidago odora*—g. Seal, see Calumba, *Hydrastis Canadensis*.

GOLDENS, *Chrysanthemum leucanthemum*.

GOLDING, *Calendula officinalis*.

GOLDSHRUB, *Palicourea speciosa*.

GOLDWASSER, ('goldwater,') see Spirit.

GOLFE, (κολος, afterwards κολος, 'a gulf,') Sinus—g. de la Veine jugulaire, see Jugular veins.

GOLUNCHEA, *Menispermum cordifolium*.

GOMBO, *Hibiscus esculentus*.

GOMME, Gumma, Gummi—g. *Adragant*, *Tragacanth*—g. *Ammoniaque*, *Ammoniac* gum—g. *Arabique*, *Acacie* gummi—g. *Astringent* of *Gambie*, see *Butea frondosa*—g. *Caragne*, *Caranna*—g. *Carane*, *Caranna*—g. *Élastique*, *Caoutchouc*—g. de *Guinée*, *Guaiacum*—g. de *Guinée*, *Sandarac*—g. *Guttaria*, *Gutta Percha*—g. *Gutte*, *Cambogia*—g. de *Kino*, *Kino*—g. *Laque*, *Lacca*—g. de *Lierre*, see *Hedera helix*—g. du *Pays*, *Gummi nostras*—g. *Sérapique*, *Sagapenum*.

GOMPHIASIS, (γομφιασις,) *Gomphiasis*—g. (γομφος, 'a nail,') A disease of the teeth, and particularly of the molars; looseness of the teeth in their sockets; *Agomphiasis*. Pain in the teeth, *Odontalgia*.

GOMPHIASMUS, (γομφιασμος,) *Gomphiasis*.

GOMPHIOL, (γομφίολοι, pl. of γομφίος,) Molar teeth.

GOMPHOCARPUS CRISPUS, (γομφος, 'a nail,' and καρπος, 'fruit,') A South African plant, the root of which, formerly known to the Dutch apothecaries as *Radix Asclepi'adis crispae*, is extremely bitter and acrid; and, on account of its diuretic virtues, a decoction or infusion of it has been advised in various kinds of dropsy. A tincture of it is said to be valuable in colic.

GOMPHOMA, gen. *Gomphomatis*, (γομφωμα,) *Gomphosis*.

GOMPHOSIS, (γομφωσις,) *Cardinamentum*, *Clavatio*, *Gomphoma*, *Conjunctio*, *Inclavatio*, *Injunctio*, *Engomphosis*. An immovable articulation, in which one bone is received into another, like a nail or peg into its hole. *Gomphosis* is only met with in the articulations of the teeth with the alveoli. It is also called *Articulation par implantation*.

GOMPHUS, (γομφος,) *Clavus*.

GOMUTO PALM, see Toddy.

GOMUTOS GOMUTO, see Toddy.

GONACRASIA, (gono, a, and κρασις, 'mixture,') *Spermatorrhœa*.

GONACRATIA, (gono, a, and κρατος, 'power,') *Spermatorrhœa*.

GON'AGRA, *Gon'yagra*, (gony, and αγρα, 'a prey,') 'That which attacks the knees.' Gout or rheumatism in the knees. *Gonyorheum* a. *Paracelsus* calls it *Gen'agra*.

GONAL'GIA, (gony, and αλγος, 'pain,') *Pain in the knee*. *Gonyal'gia*. This is almost always produced by gout. It may, however, depend on some other disease, either of the knee or of another part, particularly of the hip-joint.

GONARTHRITIS, (gony, αρθρον, 'joint,' and ιτις,) Inflammation of the knee-joint.

GONARTHROCACE, (gony, αρθρον, 'a joint,' and κακος, 'bad,') *Gonoeace*.

GONAURA, (gono, and αερα, 'an air,') See *Sperm*.

GONDOLE OCULAIRE, *Scaphium oculare*.

GONDRET'S AMMONIACAL CAUSTIC,

Pommade de Gondret—g. Counter-irritant, *Pommade de Gondret*.

GO'NĒ, (γονη,) *Go'nos*, *Genit'ra*. The semen (hence gonorrhœa), the uterus, offspring. See *Epigone* and *Generation*.

GONECYSTIDES, (gono, and κυστις, 'a bladder,') *Vesiculæ seminales*.

GONECYSTITIS, *Inflammatio vesiculærum seminalium*, (gono, κυστις, 'bladder,' and ιτις,) Inflammation of the vesiculæ seminales.

GONEPOETICUS, *Spermatorpœus*.

GONEPEUS, (gono, and ποιω, 'I make,') *Spermatorpœus*.

GONFLEMENT, (from *conflare*, 'to blow up,') Swelling.

GONGRONA, (γογγρωνη,) *Bronchocele*.

GONGROPHTHISIS, (γογγρος, 'an excrescence on a tree,' and phthisis,) *Phthisis pulmonalis*.

GONGYLIDIUM, (γογγυλίδιον, from γογγυλος, 'round,') *Pilula*.

GONGYLION, (γογγυλιον,) *Pilula*.

GONGYLIS, (γογγυλις,) *Pilula*.

GONIOCATH'ETER, (γωνια, 'an angle,' and καθητηρ, 'a catheter,') An angular or crooked catheter.

GONIOME'LE, (γωνια, 'an angle,' and μελην, 'a probe,') An angular or crooked probe or sound.

GONIOM'ETER, *Goniometrum*, (γωνια, 'an angle,' and μετρον, 'a measure,') An instrument for measuring angles.

A 'FACIAL GONIOMETER' has been invented by Mr. Turnpenny, of Philadelphia, which is well adapted for measuring the facial angle.

GONO, (from γονος, 'sperm,') in composition, *Sperm*.

GONOBOLIA, (gono, and βολη, from βαλλειν, 'to throw,') *Ejaculation*, *spermatic*.

GONOC'ACE, *Gonarthroc'acæ*, (gony, and κακος, 'evil,') *Tumor genu albus*. White swelling of the knee. *Hydrarthrus*.

GONOC'ELE, (gono, and κηλη, 'a tumor,') *Spermatocele*.

GONOF'DES, (gono, and εἶδος, 'appearance,') *Genit'ra sim'ilis*. Similar to sperm. *Sperm'atoid*, *Spermato'idēs*, *Spermato'des*. A term appropriated to any substance which resembles sperm.

GONOL'OBUS MACROPHYL'LUS, (γωνια, 'an angle,' and λοβος, 'a pod,') indigenous. *Nat. Ord.* *Asclepiadaceæ*. The root is cathartic. It is said to furnish a juice with which the Indians poison their arrows [?].

GONOP'ETICUS, (gono, and ποιω, 'I make,') *Spermatorpœus*.

GONORRHÉE BÂTARDE, *Gonorrhœa spuria*.

GONORRHOBLEPHARRHÆA, *Ophthalmia*, *gonorrhœal*.

GONORRHOPROSTATITIS, see *Prostatitis*.

GONORRHŒA, (erroneously called from gono, and ρευ, 'I flow,' because the older writers believed it to be a flux of semen;) *Blennorrhag'ia*, *Blennorrhœa*, *B. urethral'is*, *Blennorrhœa*, *Phallorrhœa*, *Medorrhœa*, *M. viri'lis*, *Catar'chus gonorrhœa* seu *ure'thræ*, *Urethri'tis*, *Inflammatio ure'thræ*, *Urethral'gia*, *Proflu'vium mucosum ure'thræ*, *Catarrhus urethral'is*, (F.) *Écoulement*, *Ure'thrite*, *Ure'trite*. An inflammatory discharge of mucus from the membrane of the urethra in both sexes; and from that of the prepuce in man, and the vagina in woman. It may be excited spontaneously, or by irritants applied directly to the membrane; but is usually produced by impure connexion. Two great varieties have been generally reckoned.—1. *GONORRHŒA PURA* seu *BENIGNA*. That which does not follow an impure connexion; (F.) *Échauffement*, *Blennorrhag'ia benigna*, *Gonorrhœa benigna*, *Catarrchus Ure'thræ*, *Gonor-*

rhea catarrhalis seu noncontagiosa; and, 2. GONORRHEA IMPURA, *malig'na, contagiosa, syphilitica, et venerea*; *Fluxus albus malig'us, Fluxus venereus, Blennorrhœa ludæ, Bopyrrha'a, Mycic'ia, Lux gonorrhœica, Seroph'ula gonorrhœica, Tubercula gonorrhœica, Clap.* (Old E.) *Chondypis*, (F.) *Chaud-pisse*; that which is the result of impure commerce. The French also distinguish the *Chaud-pisse sèche*, or that unaccompanied with discharge, and the *Chaud-pisse coctée*, *Gonorrhœa coct'ia*, *Chondurethrit'is*, or that accompanied with chordee, and which of course occurs only in the male. See Chordee. It is the kind that most frequently engages the attention of the practitioner, and is characterized by mucous discharge from the urethra or vagina, intermixed with specific matter, and accompanied with burning pain on micturition. It is decidedly infectious. It is, however, a distinct disease from syphilis, and never produces it. Its duration is various, but the inflammatory symptoms usually subside in four or five weeks, generally leaving behind more or less of the gonorrhœa mucosa or gleet. Gonorrhœa of every kind, attended with any inflammatory symptoms, is best treated by the antiphlogistic regimen, avoiding every kind of irritation, and keeping the body cool by small doses of salts, and the urine diluted by the mildest fluids. After the inflammatory symptoms have subsided, cubebæ, or the balsam of copaiba, exhibited in the dose of ʒss to ʒi three times a day, will be found effectual: indeed, during the existence of the inflammatory symptoms, it often affords decided relief. Injections are rarely required.

Sometimes, gonorrhœa affects the glans: when it is called *Gonorrhœa Spuria* seu *Bal'ani*, *Balanoblenorrhœa*, *Balanorrhœa*, *Balanit'is*, *Blennorrhag'ia spuria* seu *no'tha*, (F.) *Blennorrhagie du gland*, *Gonorrhœe bâtarde*, *Fausse Blennorrhagie*. It requires only cleanliness and cooling lotions. This is the simple form of balanitis. It may, however, be of the impure kind — *Balanorrhag'ia*, (F.) *Balanite contagieuse*. When implicating both glans and prepuce, it is *Bal'ano-posthitis*.

Some other varieties of gonorrhœa have been enumerated, but they are of little moment.

In consequence of repeated attacks of gonorrhœa, or of the debility induced by a single attack, it is not unfrequently happens, as already remarked, that a constant, small discharge occurs, or remains behind, after all danger of infection is removed. The great difference between it and gonorrhœa is, that it is uninfectious. The discharge consists of globular particles, contained in a slimy mucus, and is generally devoid of that yellow colour which characterizes the discharge of gonorrhœa virulenta. It is unattended with pain, swelling, &c. To this state the names *Gleet*, *Gonorrhœa mucosa*, *Blennorrhœa chron'ica*, *Blennorrhœa*, &c., have been given. It is commonly a disease of some duration, and demands the use of the copaiba, astringent injections; and, if obstinate, the introduction of the bougie.

GONORRHEA BALANI, G. spuria — g. Benigna, Leucorrhœa — g. Catarrhalis, see Gonorrhœa — g. Chordata, see Gonorrhœa — g. Contagiosa, see Gonorrhœa.

GONORRHEA DORMIENTIUM, G. *One'rag'nos*. The seminal discharge which occurs during sleep, and is occasioned by libidinous dreams. See Pollution.

GONORRHEA IMPURA, see Gonorrhœa.

GONORRHEA LAXORUM, G. *libidinosa*, *Spermatorrhœa aton'ica*, consists of a pellucid discharge from the urethra, whilst awake, without erection of the penis, but with venereal thoughts.

GONORRHEA LIBIDINOSA, G. *laxorum*, see Pollution — g. Maligna, see Gonorrhœa — g. Mucosa

(gleet), see Gonorrhœa — g. Noncontagiosa, G. pura — g. Notha inveterata, Leucorrhœa — g. Oneirogonos, G. dormientium, Pollution — g. Spuria, see Gonorrhœa — g. Syphilitica, see Gonorrhœa — g. Vera, Pollution, Spermatorrhœa — g. Virulenta, see Gonorrhœa.

GONOS, (γῶνος,) Gone.

GONOSTROMA, gen. *Gonostroma'tis*, γῶνος, 'offspring,' and στρώμα, 'bed,' Proligerous disc.

GONOEZEMIA, (γονυ, and ζῆμια, 'loss,') Spermatorrhœa.

GONY, (γόνυ,) *Genu*, 'the knee.'

GONYAGRA, (γονυ, (and ἀγρᾱ, 'a seizure,') Gonagra.

GONYALGIA, (γονυ, and ἀλγος, 'pain,') Gonalgia.

GONYAN'CON, *Gonycamp'sis*, (γονυ, and ἀγκων, 'a curvature,') A curvature or bending of the knee. A bent knee.

GONYCAMPISIS, (γονυ, and ἀγκυλ'is, 'curvature,') Gonyancan.

GONYCROTUS, (γονυ, and κροτω, 'I strike,') One who is knock-kneed, or in-kneed. See Entogonyancan.

GONYON'CUS, (γονυ, and ογκος, 'a tumour,') A swelling of the knee.

GONYORRHEUMA, Gonagra.

GOOD KING HENRY, *Chenopodium bonus Henrius*.

GOODYE'RA PUBES'CENS, (after Goodyer, an English botanist,) *Tussa'ca reticulata*, *Sider'ium*, *Ned'tia*, *Rattlesnake leaf*, *Rattlesnake Plantain*, *Networt*, *Net'leaf*, *Serof'ula weed*, *Cancer weed*, Ord. *Orchidaceæ*. An indigenous plant, used empirically in scrofula — the fresh leaves being applied to the sores. It is employed by the Indians.

GOOSEBERRY, *Ribes uva crispa*.

GOOSEFOOT, *Chenopodium anthelminticum* — g. Angular-leaved, *Chenopodium bonus Henrius* — g. Mercury, *Chenopodium bonus Henrius* — g. Stinking, *Chenopodium vulvaria*.

GOOSEGRASS, *Galium aparine*, G. *verum*.

GOOSESHARE, *Galium aparine*.

GOOSESKIN, *Horrida cutis*.

GORBELLY, *Physconia*.

GORDIUS MEDINENSIS, *Dracunculus*.

GORGE, (from γούργε, 'gullet,') *Ingluvies*, *Pharynx*, *Throat* — g. *Grosse*, *Bronchocele* — g. *Mal de Cynanche*.

GÖRGERET, *Goiget* — g. *Lithotome*, *Goiget*, *lithot'my* — g. *à Repousseir*, see *Goiget*.

GORG'ET, ([F.] *gorge*, 'the throat,') *Cann'lis canaliculatus*, *Ductor canaliculatus*, (F.) *Gorgeret*. An instrument representing a long gutter, in the shape of a throat, which is especially employed in the operations of lithotomy and fistula in ano.

GORGET, CUTTING, see *Goiget*, and *Lithotomy*.

GORGET FOR FISTULA IN ANO consists of a semicylindrical wooden staff, four inches long, without including the handle, and furnished with a wide groove. This is introduced into the rectum, to prevent the point of the bistoury from injuring the intestine, when the internal orifice of the fistula is deeply situate, and it is desirable to perform the operation by incision. This instrument, invented by Marchetti, has been modified by Percy, Runge, &c.

Desault invented an instrument for conducting the wire by the anus, in the operation for fistula by ligature. He called it *Gorgeret à repousseir*.

GORGET, LITHOT'OMY, (F.) *Gorgeret Lithotome*, *Cutting Gorget*, is the one used in the operation for the stone, for the purpose of dividing the prostate and the neck of the bladder, so as to enable the surgeon to introduce the forceps and extract the stone. At the end of this gorget is a crest or beak, which fits the groove of the staff, and admits of the gorget being passed along it

into the bladder. Besides *cutting*, there are also *blunt* gorgets, intended to be introduced into the wound—their concavity serving as a guide for the forceps into the bladder.

The chief modifications in the gorget have been made by Andouillet, Bell, Blicke, Bromfield, Cline, Desault, Foubert, Hawkins, Larrey, Lefèvre, Michalich, Thomas, &c.

GORGONEI FON'TES, (from the fabulous Gorgons, who had the power of turning all who beheld them to stone.) Fountains described by Libavius as containing water which possessed a petrifying property; probably, water holding in solution supercarbonate of lime.

GORGOSSET, (from *gorge*, 'throat,' [?]) Pyrosis.

GOSIER, Pharynx, Throat.

GOSSUM, Bronchocele.

GOSSYP'IIUM, *Gossyp'ium Herba'ceum*, *Gossip'ion*, *Bon'bax*, *Cotton*, (F.) *Coton*. Family, Malvaceæ. *Sex. Syst.* Monadelphia Polyandria. The seeds of the *Cotton Tree*, *Gossyp'ium arbor'ceum*, have been administered in coughs, on account of the mucilage they contain, and a decoction of a pint of the seeds to a quart of water, has been given, in the Southern States, in intermittents. A decoction of the root, *Gossyp'ii ra'dix*, (Ph. U. S.), has been prescribed as an emmenagogue and parturifacient. The *Cotton wool*, *Cotton*, *Raw cotton*, *Gossypium*, (Ph. U. S.), is used in medicine for making moxas, &c., and is applied in the treatment of recent burns and scalds.

GOSSYP'IIUM FULMINANS, see *Collodion*.

GOTCHBELLY, (Prov. *gotch*, 'a large pitcher,')

Physconia.

GOTIUM, Bronchocele.

GOTO, Bronchocele.

GOUDRON, see *Pinus sylvestris*.

GOUET, *Arum maculatum*.

GOUËTRE, (from *guttur*, 'throat,') Bronchocele.

GOULARD, (after the proposer,) *Liquor plumbi subacetatis*, L. p. s. dilutus.

GOULARD'S EXTRACT, *Liquor plumbi subacetatis*—g. Lotion, *Liquor plumbi subacetatis dilutus*—g. Water, *Liquor plumbi subacetatis dilutus*.

GOULU, (from *gula*,) Glutton.

GOUND OF THE EYE, (Prov.) *Chassie*.

GOURD, (abbreviated from *gougourde*, itself from *cucurbita*, [?]) *Cucurbita*—g. Bitter, *Cucumis colocythis*—g. Bottle, *Cucurbita pepo*.

GOURD WORM, *Distoma hepaticum*.

GOURDE, *Cucurbita*.

GOURMAND, Glutton.

GOURMANDISE, Gluttony.

GOURME, *Porrigio larvalis*. Vulgarly, in France, any cutaneous eruption.

GOUSSE, Legumen.

GOURMET (F.) A connoisseur in wines or eating.

GOUT, *Arthritis*, *Arth'ragra*, *Arthral'gia*, *Mor'bus domino'rum seu articula'ris*, *Mal'lum articulo'rum*, *Gut'ta*, *Arthros'ia Pod'agra*, *Podal'gia*, *Pod'agra*, *Arthriticus ve'rus*, *Arthritis Podagra*, *Pod'agra Arthritis*, *Flux'io arthritic'a*, *Fe'bris arthritic'a seu Podag'rica*, *Arthrodyn'ia podag'rica*, *Cum'la podag'ricum*, *Flerecin*, *Gastroarthrit'is*, *Misopto'chos*, (Sc.) *Gut*, (F.) *Goutte*, *Agrie* of Piorry. The gout was formerly regarded as a catarrh, and received its name from (F.) *goutte*, (L.) *gutta*, 'a drop,' because it was believed to be produced by a liquid, which distilled, *goutte à goutte*, 'drop by drop,' on the diseased part. This name, which seems to have been first used about the year 1270, has been admitted into the different languages of Europe. Gout is an inflammation of the fibrous and ligamentous parts of the joints. It almost always attacks, first, the great toe; whence it passes to the other smaller joints, after having produced, or been attended with, various sympathetic phenomena, particularly in the diges-

tive organs: after this, it may attack the greater articulations. It is an affection which is extremely fugitive, and variable in its recurrence. It may be acquired, or hereditary. In the former case, it rarely appears before the age of thirty-five; in the latter, it is frequently observed earlier. It is often difficult to distinguish it from rheumatism. A combination is, indeed, supposed to exist sometimes; hence called *Rheumatic gout*. During the paroxysm or fit, a burning, lancinating pain is experienced in the affected joint, attended with tumefaction, tension, and redness. One or more joints may be attacked, either at the same time or in succession; and, in either case, the attack terminates by resolution in a few days. This is the *Arthritis acuta seu inflammato'ria seu regula'ris*, *Regular gout*, *Arthros'ia pod'agra regula'ris*, *Arth'ragra legit'ima seu vera seu genu'na seu norma'tis*, (F.) *Goutte régulière chaude*. At other times, pains in the joints exist, of more or less acute character; the swelling being without redness. These pains persist, augment, and diminish irregularly, without exhibiting intermission, and, consequently, without having distinct paroxysms. The disease is then called *atonic*, *asthen'ic*, *imperfect* or *irregular gout*, *Chronic G.*, *Arthritis atonic'a seu asthen'ica*, *Arthros'ia Pod'agra larva'ta*, *Dysarthrit'is*, *Arth'ragra*. It is commonly called in France *Goutte froide*, *Goutte blanche*. It may appear primarily, or succeed attacks of regular gout.

Gout does not always confine itself to the joints. It may attack the internal organs: when it is called *Arthritis aber'rans seu errat'ica seu planet'ica seu vaga*, *Planar'thragra*, *Arth'ragra anom'ala*, *Pod'agra aber'rans*, *Vare'ni*, *Wandering*, *misplaced*, or *anomalous gout*, (F.) *Goutte vague*.

Ret'rograde gout, *Arthritis retro'grada*, *Pod'agra retro'grada*, *Arthros'ia Podagra complica'ta*, *Rece'dent* or *misplac'ed gout*, (F.) *Goutte remontée*, *G. malplacée*, *G. rentrée*, is when it leaves the joints suddenly and attacks some internal organ, as the stomach, intestines, lungs, brain, &c.

Gout is also called, according to the part it may affect, *Podagra*, *Gonagra*, *Chiroagra*, &c. It may be acute or chronic, and may give rise to concretions, which are chiefly composed of urate of soda. See *Calculus*, (arthritic.) It may, also, give occasion to nodosities, when it is called *Arthritis nodos'a*, (F.) *Goutte nouée*.

The treatment of active gout is antiphlogistic, and the local disorder should be but little interfered with. Colchicum seems to have great power over the disease. It forms the basis of the *Eau médicinale d'Husson*, a celebrated French gout-remedy. The author—a long sufferer—relies wholly on the *Vinum colechici radicis*. The bowels must be kept regular by rhubarb and magnesia; and a recurrence of the disease be prevented by abstemious habits.

GOUT, (from *gustus*,) Taste.

GOUT, DIAPHRAGMATIC, *Angina pectoris*.

GOUT PAPER, so called, *Charta antiarthrit'ica*, *Charta antirheumat'ica*, is made by spreading a very thin layer of a mixture of an ethereal or spirituous extract of the bark of mezereon root, with wax, spermaceti, and oil, over the surface of paper.

GOUT, RHEUMATIC, see *Rheumatism*, acute—g. Weed, *Ligusticum podagraria*—g. Stones, *Calculi*, arthritic.

GOUTTE, Gout, *Gutta*—g. *Blanche*, Gout (atonic)—g. *Caducue*, Epilepsy—g. *Froide*, Gout (atonic)—g. *Malplacée*, Gout (retrograde)—g. *Nouée*, Gout (with nodosities)—g. *Régulière chaude*, Gout (regular)—g. *Remontée*, Gout (retrograde)—g. *Rentrée*, Gout (retrograde)—g. *Rose*, *Gutta rosea*—g. *Sciaticue*, *Neuralgia femoro-poplitea*—g. *Sereine*, *Amaurosis*—g. *Vague*, Gout (wandering.)

GOUTTES NOIRES, *Guttæ nigræ*.

GOUTTEUX, Arthritic, Podagra.

GOUTTIÈRE, (F.) *Collic'ie*, (from *gutta*, 'a drop,' or from Sax. *geotian*, to pour.) A gutter in a bone, like that used for carrying off rain. Some of these cavities are intended to facilitate the sliding of tendons, such as the *Gouttière Bicépitale* or *Bicipital groove*. Others, as the *Gouttière sagittale* or *Sagittal groove*, lodge blood-vessels and especially veins. Others, again, are merely intended for the support of certain organs; as the *Gouttière basilaire* or *Basillary fossa*, which supports the medulla oblongata.

GOUTTIÈRE BASILAIRE, see *Gouttière*—*g. Bicipitale*, Bicipital groove—*g. Lacrymale*, Lachrymal groove—*g. Sacré*, Sacral groove—*g. Sagittale*, see *Gouttière*—*g. de la Veine cave*, see Liver.

GOUTY, (from *gout*,) Arthritic, Podagric—*g. Rheumatism*, see Rheumatism, acute.

GOVERNAIL DU TESTICULE, Gubernaculum testis.

GOWAN, EWE, (Gael. *gugan*,) Bellis—*g. Yellow*, Ranunculus.

GOWLAND'S LOTION, see Lotion, Gowland's.

GOWN, RED, (from its covering like a gown,) Strophulus—*g. Yellow*, Icterus infantum.

GOWNE OF THE EYE (Prov.), *Chassie*.

GRACE-WIFE, (Prov.) Midwife.

GRACILARIA LICHENOÏDES, (from *gracilis*, 'slender,') *Fucus amylaceus*.

GRAC'ILIS, *Mac'er*, *Macilen'tus*, 'slender, lean.' Also, the slender *Rectus interior femoris* seu *Grac'ilis interior*, (F.) *Sous-pubio-crèti-tibial*, *Sous-pubio-prétibial* (Ch.), *Droit ou grêle interne le la cuisse*. This muscle is situate at the inside of the thigh. It is thin and very long; and arises from the descending ramus of the pubis, to be inserted at the inner and inferior part of the tuberosity of the tibia. It bends the leg and causes adduction of the thigh.

GRACILIS ANTERIOR, *Rectus femoris*.

GRAD'UATE, *Gradu'atus*, (*gradus*, 'a step,' 'a degree,') In medicine, one who has attained a degree,—usually, the degree of doctor.

GRÆA, (*γραια*,) The pellicle which forms on milk. The folds of skin round the umbilicus. An old woman.

GRAFT, *Graff*, *Inser'erè*, (F.) *Greffer*, (from Sax. *græfan*, 'to engrave,') To unite a part to another, so that it may receive support from it; to engraft. The operation is called *grafting* or *engrafting*, *Insit'tio*.

GRAIN, *Granium*; the 60th part of a Troy, and the 72d part of a *Poids de marc* drachm.

GRAIN OIL, Oil, fusel—*g. Oily*, Sesamum orientale.

GRAINE D'ASPIG, see *Phalaris Canariensis*—*g. de Catepnee*, see *Euphorbia lathyris*—*g. d'Écarlate*, *Coccus cacti*—*g. d'Épurgée*, see *Euphorbia lathyris*—*g. de Moluques*, see *Croton tiglium*—*g. Musc*, *Hibiscus abelmoschus*—*g. de Paradis*, *Amomum granum paradisi*—*g. de Perroquet*, *Carthamus tinctorius* (seed)—*g. Royale Mineure*, see *Euphorbia lathyris*—*g. de Santé*, see *Pilulæ aloes et kinæ kinæ*—*g. de Tilly*, see *Croton tiglium*—*g. de Turquie*, *Zea mays*.

GRAINS, GUINEA, see Pepper, Malagueta.

GRAINS OF PARADISE, *Amomum granum Paradisi*.

GRAISSE, (from *crassus*, 'gross' [?]), *Pinguedo*—*g. de Mouton*, *Sevum*—*g. d'Oie*, *Adeps anserina*—*g. Oxygénée*, *Unguentum acidi nitrosi*—*g. de Porc*, *Adeps præparatus*.

GRAMEN, gen. *Graminis*, 'grass,' *Triticum repens*—*g. Egyptiacum*, *G. Crucis cyperioïdis*—*g. Caninum*, *Triticum repens*.

GRAMEN CRUCIS CYPERIOÏDIS, *Gramen Ægypti'acum*, *Ægyptian Cock's foot grass*. Ord. Gramineæ. The roots and plants possess the virtues of the *Triticum repens*, and have been recom-

mended in the earlier stages of dropsy. They were, formerly, considered to possess many other properties.

GRAMEN DIOSCORIDIS, *Triticum repens*—*g. Major*, *Sarsaparilla Germanica*—*g. Orientale*, *Juncus odoratus*—*g. Repens*, *Triticum repens*—*g. Rubrum*, *Sarsaparilla Germanica*.

GRAMIA, *Chassie*, Lippitudo.

GRAMINIV'OROUS, *Graminiv'orus*, (*gramen*, gen. *graminis*, 'grass,' and *voro*, 'I eat,') Feeding or subsisting on grass.

GRAMMARIUM, *Gramme*, *Seruple*.

GRAMME, (*γραμμ*,) *Gramma'rium*. An ancient weight, equivalent to the 24th part of an ounce, or to 24 grains or a scruple, avoirdupois. At the present day, the gramme is equal in weight to a cubed centimètre of water; or to 18 grains, poids de marc—15.434 grains, Troy.

GRAMME, (*γραμμ*,) *Iris*, Line.

GRAMMITIS AUREA, *Asplenium ceterach*—*g. Ceterach*, *Asplenium ceterach*.

GRANA, (abbreviation of *Migrana* or *Hemicrania* [?]), *Hemicrania*—*g.* (pl. of *granum*,) *Guineensis*, *Pepper*, *malagueta*—*g. Molucca*, *Croton tiglium*—*g. Moschi*, *Hibiscus abelmoschus*—*g. Orientis*, see *Menispermum cocculeus*—*g. Paradisi*, *Amomum granum Paradisi*—*g. Tiglii seu Tiglia*, see *Croton tiglium*—*g. Tiliï*, *Croton tiglium*—*g. Triticea*, see *Hyoides os*.

GRANADILLA, APPLE-SHAPED, (dim. of [S.] *granada*, 'the pomegranate,') *Passiflora maliformis*.

GRANATIFRUCTÛS CORTEX, (from *granum*, because full of grains or seeds,) see *Punica granatum*—*g. Radicis cortex*, see *Punica granatum*.

GRANATUM, *Punica Granatum*.

GRAND DORSAL, *Latissimus dorsi*.

GRANDEB'ALÆ. The hair in the arm-pits.

GRANDGORE, ([Sc.]) Syphilis.

GRANDINOSUM OS, (from *grando*, 'hail,') Cuboid.

GRANDO, 'hail,' *Chalaza*.

GRANIV'OROUS, *Gramiv'orus*, (*granum*, 'a grain,' and *voro*, 'I eat,') Feeding or subsisting on grain or seeds.

GRANTRISTUM, Anthrax.

GRANULA SANGUINIS, (pl. of *Granulum*, and dim. of *Grana*,) Globules of the blood—*g. Seminis*, see *Sperm*.

GRANULAR, *Granul'ous*, *Granulo'sus*, (F.) *Granulé*, *Granuleux*, (from *granule*,) Composed of or containing granules or small particles,—as 'granular eyelid,' 'membrana granulosa,' &c., &c.

GRANULAR CONJUNCTIVA, *Trachoma*—*g. Degeneration or Disorganization of the Kidney*, *Kidney*, *Bright's disease of the*—*g. Eyelid*, *Trachoma*—*g. Layer*, *Membrana granulosa*—*g. Liver*, *Cirrhosis*—*g. Tin*, see *Tin*.

GRANULATED LIVER, *Cirrhosis*.

GRANULATION, *Granula'tio*, (from *granule*,) *Gem'ma*, (F.) *Bourgeon*, *B. charn.* Granulations are the reddish, conical, flesh-like shoots, which form at the surface of suppurating wounds and ulcers. They are the product of inflammatory excitement, and may be produced in indolent ulcers, by exciting the parts by proper stimulants. They form the basis of the cicatrix.

GRANULATION is, likewise, a name given by the modern French physicians to an organic lesion, consisting in the formation of small, round, firm, shining, semi-transparent tumours, of the size and shape of millet-seed, or of a pea; which are met with in the lungs particularly, and in considerable quantity; often without materially interfering with their functions.

In pharmacy, *granulation* is a process by which a metal is reduced to fine grains, by melting it, and causing it, whilst liquid, to pass through a kind of sieve into a vessel of water,—as in the

making of shot:— or by shaking or rubbing the melted metal in an appropriate box or vessel,— as in the formation of granular tin or granulated zinc.

GRANULATIONS OÛRÉBRALES, Glandulæ Paccionii.

GRANULATIONS, MIL'IARY, *Miliary tubercles*, are the small, transparent grains, of variable size, from that of a millet-seed to that of a grain of hemp which are presumed to be the primitive state of tubercles.

GRAN'ULE, *Gran'ulum*; (dim. of *granum*, 'a grain.')

A small grain; a small compact particle; a cytoblast. Used, in French pharmacy, for a sugar-coated pill, or any small *dragée*.

GRANULES SEM'INAL, *Gran'ula sem'inis*. Minute, rounded, granulated bodies, observable in the semen, which are, in all cases, much less numerous than the spermatozooids. See *Sperm*.

GRANULEUX, Granular.

GRANULUM, Granule.

GRANULOUS, Granular.

GRANUM, Grain.

GRANVILLE'S LOTION, see *Lotion*, *Granville's counter-irritant*.

GRAPE, ([F.] *grappe*, 'a cluster, a bunch,') see *Vitis vinifera*—*g.* Sea-side, see *Kino*.

GRAPECURE, (F.) *Cure de raisin*, (G.) *Traubenkur*, *Weintraubenkur*. A medication in Germany, which consists in the use of the grape for both meat and drink; nothing more at the farthest being allowed than a piece of dry bread. This diet is continued for weeks. Its effects are altogether revellent, and resemble in many respects those of hydropathy.

GRAPES, DRIED, *Uvæ passæ*.

GRAPHIDOIDES, (*γραφίδος*, gen. *γραφίδος*, 'a style,' and *εἶδος*, 'resemblance,') *Styloid*.

GRAPHIOIDES, *Styloid*.

GRAPHIOIDES, *Styloid*.

GRAPHIS'CUS, (*γραφίσκος*,) *Graphis'cus Di'ocletis*. An instrument invented by Diocles for extracting darts [?]. Said, in many of the Dictionaries, but erroneously, to be described by Celsus. He speaks only of the *Cyathis'cus Di'ocletis*, *Διακλῆτος κυαθίσκος*. See *Cyathiscus*.

GRAPHI'TES, (from *γραφῆς*, 'to write,') *Plumbago*, *Superaccharatum seu Carburetum Ferri*, *Ferri Carbonatum seu Superaccharatum*, *Car'bo mineralis*, *Galena*, *Gal'nè*, *Plumbum nigrum*, *Cerulea nigra*, *Grat'white*, *Carburet of iron*, *Black lead*, *Wad*, (F.) *Crayon noir*, *Plombagine*, *Charbon minéral*, *Carbure de fer*. This substance has been esteemed slightly astringent and desiccative. It has been advised by Weinhold in the cure of herpes.

GRAPHOÏDES, (*γραφῆς*, 'a style,' and *εἶδος*, 'resemblance,') *Styloid*.

GRAPHOSPASMUS, (*γραφῶ*, 'I write,' and *σπασμός*, 'spasm,') Cramp, writers'.

GRAS DES CADAVERES, 'fat of dead bodies,' ([L.] *crassus*, 'thick, fat,') *Adipocire*—*g. des Cimetières*, *Adipocire*.

GRAS FONDURE (F.), *Diarrhœa adiposa*, literally, 'molten grease.' A species of diarrhœa, referred to by old writers; accompanied with great emaciation, and in which the evacuations contain fat-like matter. According to Sauvages, the *Grasfondure* differs from colliquative diarrhœa in not being attended with hectic fever.

GRAS DE JAMBE, *Sura*.

GRASS, (Sax. *græs*,) *Asparagus*, *Triticum repens*—*g.* Bear's, *Yucca filamentosa*—*g.* Bermuda, *Cynodon dactylon*—*g.* Bitter, *Aletris farinosa*—*g.* Blue-eyed, *Sisyrinchium Bermudianum*—*g.* Brome, *Bromus ciliatus*—*g.* Brome, soft, *Bromus ciliatus*—*g.* Canary, cultivated, *Phalaris Canariensis*—*g.* Couch, *Triticum repens*—*g.* Dog, *Triticum repens*—*g.* Eel, *Pila marina*—*g.* Egyptian cock's foot, *Tranen erula cyperoides*—*g.* Goat's, *Scorzonera*

—*g.* Knot, *Polygonum aviculare*—*g.* Lily, *Sisyrinchium Bermudianum*.

GRASS OIL OF NEMAU, *Oleum graminis Indici*.

GRASS, PHYSIC, *Sisyrinchium Bermudianum*—*g.* Rib, *Plantago lanceolata*—*g.* Scurvy, *Sisyrinchium Bermudianum*—*g.* Silk, *Yucca filamentosa*—*g.* Sweet, *Acorus Calamus*.

GRASS TREE, *Xanthorrhœa*.

GRASS, VIPERS', *Scorzonera*—*g.* Yellow-eyed, *Xyris bulbosa*.

GRASSET (F.), (from *crassus*, 'thick' [?]). The anterior region of the thigh, bounded below by the patella.

GRASSEYEMENT (F.), 'speaking thick.' (from *gras*, 'thick,') *So'nus blæ'sus*, *Rotacis'mus*. According to Sauvages, a vicious pronunciation of the letter *r*. They who speak thick, like the inhabitants of Newcastle, in England, or of Havre, in France, have difficulty in pronouncing the *r*, and they frequently substitute for it the letter *l*; but this does not properly constitute *Grasseyement*. It consists in this: that, in words in which the letter *r* is joined to another consonant, a sort of *burring* or guttural rolling is heard, nearly like that produced by gargling. See *Rotacism*.

GRASUS, (*γρασος*,) *Cinabra*, *Hircismus*.

GRATIA DEI, ('grace of God,') *Gratiola officinalis*.

GRATIOLA CENTAURIROIDES, *G. officinalis*.

GRATIOLA OFFICINA' LIS, *Digitalis min'ima*, *Grat'ia Dei*, *Gratiola Centaurioides*, *Hedge hyssop*, *Herb of Grace*, (*gratia*, 'grace or favour,') owing to its presumed excellent medical virtues.) It is a native of the South of Europe. (F.) *Gratiola*, *Séné des prés*, *Petite digitale*, *Herbe au pauvre homme*. The plant is inodorous; taste strong, bitter, nauseous. It is possessed of anthelmintic, purgative, emetic, and diuretic properties. Dose, ten grains.

GRATIOLE, *Gratiola officinalis*.

GRATELLE, (from *gratter*, 'to scratch,') *Psoriasis*.

GRATTERON, *Galium sparine*.

GRATTOIR, *Rasatorium*.

GRAVATIF, (from [L.] *gravis*,) Heavy.

GRAVE, (from *gravis*,) Serious—*g.* Plant, *Datura sanguinea*.

GRAVEDO, gen. *Grave'dinis*, (from *gravis*,) *Catarrh*, *Coryza*—*g.* Neonatorum, *Snuffles*.

GRAVEL, (dim. of old [F.] *grave*, 'a pebble, itself, perhaps, from [Teut.] *graben*, 'to dig out,') *Lith'ia renal'ia arenosa*, *Lith'ia nephrit'ica seu renal'is*, (F.) *Gravelle*. A disease occasioned by small concretions, similar to sand or gravel, *Gla'rea*, (F.) *Gravier*, which forms in the kidneys, passes along the ureters to the bladder, and is expelled with the urine. These concretions, which are commonly composed of urates and an animal matter, are deposited at the bottom of the vessel, immediately after the excretion of the urine; and, by their hardness and resistance under the finger, differ considerably from the ordinary sediment of that liquid. A vegetable diet and alkaline drinks are the best prophylactics. See *Calculi*, *urinary*. A fit of the *Gravel*, *Nephral'gia calculosa seu arenosa*, *Col'ica nephrit'ica*, is the excruciating suffering induced by the passage of gravel from the kidney to the bladder. It can only be relieved by anæsthetics, opiates, the warm bath, &c.

When the deposit is in fine particles, it is termed *Sand*, *Are'na*, *Are'nula*, *Psam'ma*, *Psammus*.

GRAVEL GRASS, *Galium verum*.

GRAVEL, PY'LEUS or HAIR'Y, (F.) *Gravelle pil'ense*. A species of gravel containing hairs, phosphate of lime, ammoniaco-magnesian phosphate, and a little uric acid.

GRAVEL PLANT, *Epigæa repens*—*g.* Root, *Eupatorium purpureum*.

GRAVELEUX, Calculous.
GRAVELLE, (see Gravel,) Chalaza, Gravel—*g. Pileuse*, Gravel, pileous.
GRAVEOLENS, (*gravis*, and *oleo*, 'I smell,') Fetid.
GRAVID, (from *gravis*,) Pregnant.
GRAVIDIN, (from *gravid*,) A sediment in the urine of pregnant women, which, according to Stark, by its decomposition gives rise to the pellicle kystein. It differs from albumen, casein and gelatin.
GRAVIDITAS, Fecundation, Pregnancy—*g. Abdominalis*, Pregnancy, abdominal—*g. Extra-uterina*, see Pregnancy—*g. Extra-uterina in Ovario*, Pregnancy, ovarian—*g. Extra-uterina Secundaria*, Metacyesis—*g. Interstitialis*, Pregnancy, interstitial—*g. Molaris*, Mole—*g. Ovarii*, Pregnancy, ovarian—*g. Spuria*, Pregnancy, false—*g. Tubaria*, Pregnancy, tubal, Salpingocystis—*g. Uteri substantia*, Pregnancy, interstitial—*g. Uterina*, Pregnancy.
GRAVIER, Gravel.
GRAVIMETER, (*gravis*, and *μετρον*, 'a measure,') Areometer.
GRAVIS, Heavy.
GRAVITY, SPECIFIC, *Gravitas specifica*, (F.) *Pesanteur spécifique*. The relation between the weight of a body and its bulk; thus, supposing four bodies to be of the same size, but to weigh, one four, another three, another two, and the fourth one; the specific gravity of the first will be four times greater than that of the last. The specific gravities of different bodies are, therefore, as the weights, bulk for bulk. For solids, and liquids, water is taken as the unit; atmospheric air for the gases. Thus, water is 1.000; mercury, at the common temperature, 13.58. Whence, we conclude mercury is between thirteen and fourteen times heavier than water.
When the specific gravity of a substance is given, its temperature is assumed to be at 60° Fah.
GRAYHAIRD, see Poliosis.
GRAY MATTER OF THE BRAIN, Cortex cerebri, see Neurine.
GRAYMILL, Lithospermum officinale.
GREASE, ([F.] *Groisse*, 'fat.') A specific inflammation, affecting the skin of the heels of the horse, which is especially interesting from the circumstance, that the matter, if inserted under the cuticle of an unprotected individual, may give rise to an affection—*grease-pox*, *variola equina*—which preserves the person from smallpox. [?]
GREASE, BARROW'S, *Adeps suillus*—*g. Goose*, *Adeps anserinus*—*g. Molten*, *Gras-fondure*.
GREASE POX, see Grease.
GREENHEART, see Bebeerru.
GREENHOW'S TINCTURE FOR THE TEETH, *Spiritus armoracae compositus*.
GREEN SICKNESS, Chlorosis.
GREENWEED, *Genista tinctoria*.
GREFFER, Graft.
GRÊLE, old (F.) *Gresle*, *Grac'illa*, 'long and thin.' This epithet is given by the French to various parts, as the
Apophyse Grêle du Marteau, the slender apophysis or process of the malleus. A long process situate at the anterior part of the neck of the malleus, which passes out by the fissure of Glaserius. It is also called the *Apophysis of Rau*, although it was already known to Fabricius ab Aquapendente and to Cæcilius Follius.
GRÊLE, (from *gres*, 'a stone' [?]) Chalaza—*g. Interne de la Cuisse*, *Gracilis*.
GRÊMIL OFFICINAL, *Lithospermum officinale*.
GREMIUM, Vulva.
GRENADÉ, Influenza.
GRENADIER, (from *granatum*,) *Punica granatum*.

GRENADIN, see *Punica granatum*.
GRENIERS, (from *granarium*, 'a granary,') *Vesiculae seminales*.
GRENOUILLE, ('a frog,') *Rana esculenta*.
GRENOUILLETTE, (dim. of *grenouille*, 'a frog,') *Ranula*.
GRÉOULX, MIN'ERAL WA'TERS OF. Sulphuretted springs in the department of Basses-Alpes, France. Temperature, about 100° Fah. The season is from the first of May to the first of October.
GRESSURA, Perineum.
GRESSUS, gen. *Gressûs*, (*gradior*, *gressus*, 'to step,') Walking.
GREVEURÉ, (from *greuer*, 'to aggrrieve, to burden' [?]) Hernia.
GRÎAS CAULIFLO'RA; (from *γρᾶειν*, 'to eat,') *Anchovy Pear*. The fruit of a tree of the *Nat. Ord.* *Onagraceæ*, which grows in the West Indies. It is large, has a stone, and is eaten.
GRIELUM, *Apium petroselinum*, *Smyrniacum olusatrum*.
GRIFF, see Mulatto.
GRIFFO, see Mulatto.
GRINCEMENT DES DENTS, 'grinding the teeth,' (by onomatopœia [?]), *Brygnus*.
GRINCOMES, (old [E.]), Syphilis.
GRINDER. One who grinds or polishes, see Quizzer.
GRINDERS, Molar teeth—*g. Asthma*, see Asthma, grinders'—*g. Disease*, see Asthma, grinders'—*g. Rot*, see Asthma, grinders'.
GRINDING MILL, see Pulverization.
GRIPPE, CUTTING ON THE, (from (F.) *gripper*, Sax. *gripan*, 'to clutch,' 'to seize,') See Lithotomy.
GRIPES, (same etymon,) Tormina, Colic.
GRIPES, WA'TERY. A popular name for a dangerous disease of infancy, common in England, which does not differ essentially from the cholera infantum of this country.
GRIPHOSIS, (*γριπος*, 'bent,') *Onychogryphosis*.
GRIPPE (F.), (*gripper*, 'to clutch,' 'to seize,') A vulgar name for several catarrhal diseases, which have reigned epidemically; as the influenza.
GRIPPE, Pinched.
GRISOU, see Hydrogen carburetted.
GRISTLE, (Sax. *griple*,) Cartilage.
GRITS, Groats.
GRIT GRU'EL, *Water gruel*. This is made as follows:—Take three ounces of *grits*; wash them well in cold water, and, having poured off the fluid, put them into four pints of fresh water, and boil slowly until the water is reduced one-half; then strain through a sieve. It is a good demulcent, and is employed also as a vehicle for clysters.
Cornmeal Gruel may be made in like manner—two large spoonfuls to a pint of water.
Milk may be added to both of these.
GROAN, (from Sax. *groman*, 'to groan,' by onomatopœia.) See Susspirium.
GROATS, (from Sax. *gryt*,) *Gru'tum*, *Avena excorticata*; (F.) *Gruan*, Oatmeal, (Yorkshire.) Oats, hulled, but unground, (*Lancashire*.) Hulled oats, half ground. Oats that have the hulls taken off; *Grits*, (Prov.) *Cutlins*. When crushed, they are termed *Emb'den groats*. In America, fine hominy is called *Grits*, and wheat prepared in the same way is likewise so designated. It is also called *wheat'en hom'iny*. See Avena.
GROATS, CRACOW, Semolina.
GROCERS' ITCH, see Eczema Impetiginodes, Psoriasis.
GROGBLOSSOMS, *Gutta rosea*.
GROGROSES, *Gutta rosea*.
GROIN, Inguen.
GROMWELL, *Lithospermum officinale*.

GROMYL, *Lithospermum officinale*.

GROOVE, *Fur'rou, Sul'cus*, (F.) *Rainure*, (Icelandic, *grifa*, Sax. *grapan*, 'to dig.') A channel or gutter in a bone or surgical instrument. See *Coulisse*.

GROOVE, *PRIM'ITIVE*, *Primitive streak or trace*, *Nota primitiva*. A bright streak in the long axis of the pellucid part of the area germinativa after it presents a central pellucid and a peripheral opaque part, and passes from the round to the pear shape.

GROOV'ED. *Sulca'tus, Stria'tus, Canalicula'tus*, (F.) *Cannelé ou Canelé ou Canaliculé*. Canaliculated. Having a small channel or gutter, as a grooved probe.—*Aulacome'lé, Caelome'lé, Specill'um Sulca'tum*.

GROS, Drachm.

GROS COU, 'large neck,' (from *crassus*, 'thick.') Brachocoelæ—*g. Retombet*, *Passerina tartonraira*.

GROSEILLER NOIR, *Ribes nigrum*—*g. Rouge, Ribes rubrum*.

GROSEILLES, (from *grossulus*, dim. of *grossus*, 'a small fig.') *Ribes nigrum*—*g. Rouge, Ribes rubrum*.

GROSSE GORGE, ('large throat,') Brachocoelæ.

GR OSSESSE, (from *gros, grosse*, 'large,') Pregnancy—*g. Abdominale*, Pregnancy, abdominal—*g. Afetale*, Pregnancy, afetal—*g. Bigéminale*, Pregnancy, bigeminal—*g. Complexe*, Pregnancy, complex—*g. Composée*, Pregnancy, compound—*g. Contre-nature*, Pregnancy, extra-uterine—*g. Fausse ou apparente*, Pregnancy, false—*g. Fétale*, Pregnancy, foetal—*g. Gazo-hystérique*, Pregnancy, gazo-hysterical—*g. Gémellaire*, Pregnancy, bigeminal—*g. Hémato-hystérique*, Pregnancy, hemato-hysterical—*g. Hydro-hystérique*, Pregnancy, hydro-hysterical—*g. Inter-extra-utérine*, Pregnancy, complex—*g. Ovarienne*, Pregnancy, ovarian—*g. Sarcofétale*, Pregnancy, sarcofoetal—*g. Sarcophystérique*, Pregnancy, sarco-hysterical—*g. Simple*, Pregnancy, solitary—*g. Solitaire*, Pregnancy, solitary—*g. Trigéminal*, Pregnancy, trigeminal—*g. Triple*, Pregnancy, trigeminal—*g. Tubaire*, Pregnancy, tubal—*g. Utéro-abdominale*, Pregnancy, utero-abdominal—*g. Utéro-ovarienne*, Pregnancy, utero-ovarian—*g. Utéro-tubaire*, Pregnancy, utero-tubal.

GROSSULARIA, *Ribes uva crispa*—*g. Nigra, Ribes nigrum*—*g. Non spinosa, Ribes nigrum*—*g. Rubra, Ribes rubrum*.

GROTTO DEI SERPI (L.) 'Grotto of serpents.' A grotto near Braccano, in Italy, filled with warm water, to which invalids, affected with skin diseases, resort.

GROUILLEMENT D'ENTRAILLES, ('rumbling of the bowels,') Borborygmus.

GROUND BERRY, *Gaultheria*—*g. Holly, Gaultheria*—*g. Ivy, Gaultheria*, *Glechoma hederacea*—*g. Nut, Apios tuberosa, Arachis hypogaea*, *Pignut*—*g. Pine, Teucrium chamaepitys*—*g. Pine, French, Teucrium iva*.

GROUNDIESWALLOW (Sc.), *Senecio*.

GROUNDSEL, ([Sax.] *grunne-epfelge*,) *Senecio*—*g. Tree, Baccharis halimifolia*.

GROUSEBERRY, *Gaultheria*.

GROWTH, (Sax. *growan*, 'to grow.') *Incrementum, Crescen'tia, Anoplo'sis, Anaptyx'is*, (F.) *Croissance, Crue*. The development of the body; particularly in the direction of its height. Also, any adventitious tissue; thus, we speak of a morbid growth or formation. See *Development*.

GROZET-EYED, (Prov.) *Exophthalmus*.

GRUAU, see *Groats*.

GRUB, Larve, see *Ectozaa*.

GRUBS, *Acne punctata*.

GRUEL, GRIT, see *Grits*—*g. Water*, see *Avena*, and *Grits*.

GRUFF, (Teutonic *ge*, and *ruh*, 'rough.') In pharmacy, the coarse residue, which will not pass through the sieve in pulverization.

GRUNEAU, (from *grunus*,) *Coagulum*.

GRU'MOUS, *Grunu'sus*, (F.) *Grunelé, Grumeleux*, (*grunus*, 'a clot,') Clotted.

GRUMUS, *Coagulum*.

GRUTUM. 'Groats.' *Grutum Mil'ium, Mil'ium*. A hard white tubercle of the skin, resembling, in size and appearance, a millet-seed, found on the eyelids, the root of the nose, the scrotum and ear. See, also, *Groats*.

GRYPHIUS PES. The Griffon's foot, (F.) *Pied de Griffon*. An instrument of which Ambrose Paré speaks, which was used for extracting moles from the uterus.

GRYPHOSIS, *Oncogryphosis*.

GRYPHOSIS, (*γρυψος*, 'bent,' and *osis*,) *Incurvatio*. Curvature or crookedness in general. Crookedness or incurvation of the nails. See *Oncogryphosis*.

GRYPOTES, (*γρυπτός*,) see *Grypus*.

GRYPUS, (from *γρυψος*, 'bent,') One who has a crooked or aquiline nose. The condition is termed *Grypotes*.

GUA'CO, *Hua'co*. The name of a plant, *Eupatorium Guaco*, described by Humboldt and Bonpland under the name *Mikania Guaco*, *Ord.* *Compositæ*, which grows in the valleys of Madalena, Rio Cauca, &c., in South America. The negroes use the juice against the bites of poisonous reptiles;—both in the way of prevention and cure. It has been, of late, brought forward as a remedy in cholera.

GUA'IAC, see *Guaiacum*.

GUA'IACI LIGNUM, see *Guaiacum*—*g. Resina*, see *Guaiacum*.

GUA'IACINE, see *Guaiacum*.

GUA'IACUM, *G. Officinale* seu *America'num*, *Lignum vitæ* seu *sanc'tum* seu *benedictum* seu *Ind'icum*, *Pa'lus sanc'tus*, *Hagio'ylum*, (F.) *Gayac, Gaïac*. *Ord.* *Zygophyllæ*. The resin—*Guai'aci Res'ina*, *Guai'ac*, (F.) *Résine ou Gomme de Gaïac*—and the wood—*Guai'aci lignum*—are both used in medicine. Their odour is slightly fragrant; taste warm and bitter, of the resin more so than of the wood. The resin is concrete, brittle; colour, externally, greenish; internally grayish. Water dissolves about one-tenth; alcohol 95 parts. It is soluble, also, in *liquor potassæ* 15 parts, *liquor ammoniæ* 38 parts. The powder is whitish, but changes to green in the air. The base of the guaiacum is a peculiar resin, called *Guai'acine*.

Guaiacum is stimulant and diaphoretic; and in large doses, purgative. It is administered in chronic rheumatism, gout, cutaneous diseases, and the sequelæ of syphilis. Dose of resin, gr. v to xx:—to purge, gr. xx to xl.

Guai'acum sanc'tum, and *G. angustifolium* have the same virtues as *G. officinale*.

GUALTHERIA, *Gaultheria*.

GUANO,—according to Tschudi, properly *Huano*, 'dung'—is formed of the excrements of different kinds of marine birds—mews, divers, sheerbreaks, &c., but especially of the *Sula variagata*. It is found in enormous layers in the South American islands of the Pacific, and is used as manure.

GUARANA, *Paullinia*.

GUARANHEM, *Monesia*.

GUARANIN, see *Coffea Arabica*.

GUARAPO. A fermented liquor made, in Peru, of sugarcane pulp and water. It is a very favourite beverage of the negroes.

GUARD (for a bed), *Alèse*.

GUARERBA ORBA, *Momordica elaterium*.

GUAVA APPLE, *Psidium pomiferum*.

GUAYAVA, *Psidium pomiferum*.

GUAZA, see *Bangue*.

GUBERNACULUM DEN'TIS, ('rudder of the tooth,' from *gubernare*, 'to govern,') A cord which passes from the follicle of the permanent tooth along a small long canal behind the alveo-

lus of the milk tooth, and becomes continuous with the gum. The gubernaculum has been supposed to direct the permanent tooth outwards. The canal has been termed *Pter dentis*.

GUBERNACULUM TESTIS, *Gubernaculum testis Hunteri*, *G. Hunteri*, *Hunteri Ligamentum suspensorium Testis*, (F.) *Gouvernail du testicule*. A triangular, fibro-cellular cord; which, in the fetus, arises from the ramus of the ischium and the skin of the scrotum, and proceeds to the posterior part of the testicle, before this organ issues from the abdomen. It has been supposed to be a continuation of the fascia superficialis with muscular fibres from the internal oblique muscle, which pass upwards to the testis when in the abdomen; and by their contraction draw the testis down, and ultimately form the cremaster muscle.

GUÊPE, (old (E.) *guespe*, from (L.) *vespa*), Wasp.

GUÉRISON, (from (L.) *curare*, 'to cure.') Cure.

GUÉRISABLE, Curable.

GUI, *Viscum album*—*g. de Chêne*, *Viscum album*.

GUIDER, ('leader.') Tendon.

GUILANDINA MORINGA, (after Melchior Guilandinus (Wieland), a professor at Padua,) *Hyperanthe'ra moringa*. A plant, *Ord. Leguminosæ*, which affords the *Ben nut*, and the *lignum nephriticum*. It is also called *Moringa Oleifera* seu *Zeylanica* seu *Nux ben*. Under the name *Guilandina moringa*, Linnæus is said to have confounded *Moringa pterygosperm'a* and *M. ap'tera*. *Hyperanthe'ra moringa* is a synonyme of the former species. The nut *Ben*, *Ben nux*, *Glans unguenta'ria*, *Balanus Myrep'sica*, *Sandalum cæruleum*, *Oily Acorn* or *Ben nut*, is a West India nut which furnishes an oil, *O'leum Balanitum*, *Oil of Ben*, that does not become rancid by age, and is hence used by perfumers. It is purgative.

The wood of the *Guilandina* is called *Lignum Nephriticum*, and has been used in decoction, in affections of the urinary organs.

GUILLOTINE TONSIL, (after Guillotin, the reviver of the maiden, a Scotch instrument of decapitation,) *Amygdatome*.

GUILTY CUPS, (Prov.) *Ranunculus acris*.

GUIMAUVE, *Althæa*—*g. Veloutée*, *Hibiscus abelmoschus*.

GULA, (akin to *collum* [?]) Œsophagus, Pharynx.

GULÆ IMBECILLITAS, ('weakness of the throat,') Pharyngoplegia—*g. Principium*, Pharynx.

GULLET, (from *gula*), Œsophagus.

GULLION (Prov.), Colic.

GULOSUS, (from *gula*, 'the gullet,') Glutton.

GULSCHÖCH, (Sc. from Belg. *Geelzucht*, 'yellow disease,') Icterus.

GUM ACAROID, (from *gummi*), see *Xanthorrhæa*—*g. Anime*, *Anime*—*g. Arabica*, *Acaciæ gummi*—*g. Artificial*, *Dextrin*—*g. Bassora*, *Bassora gum*—*g. Blackboy*, see *Zanthorrhæa*.

GUM BOIL, Parulis.

GUM, BOTANY BAY, see *Xanthorrhæa*—*g. British*, *Dextrin*—*g. Butea*, see *Butea frondosa*—*g. Caranna*, *Caranna*—*g. Dragon*, *Tragacantha*.

GUM, EAST INDIA, see *Feronia elephantum*—*g. Elastic*, *Coatucheou*.

GUM OF THE EYE, *Chassie*.

GUM, FALLING AWAY OF THE, *Ulatrophia*—*g. Hemlock*, see *Pinus Canadensis*—*g. Indigenous*, *Gummi nostras*—*g. Ivy*, see *Hedera helix*—*g. Juniper*, *Sandarac*.

GUM LANCET, *Dentiscal'pinum Odontoglyphon*, (F.) *Déchaussoir*. An instrument for separating the gum from the cervix of the tooth, prior to extraction. It is formed much like a fleam. The operation itself is called *Lancing the gums*, *Dentium sculptura*, (F.) *Déchaussement*.

GUM MESQUITE, see *Mesquite*—*g. Orenburg*, see *Pinus larix*.

GUM RASH, *Strophulus*.

GUM, RED, *Strophulus*.

GUM RESIN, *Gummi resina*.

GUM, SANDARACH, *Sandaracæ*—*g. Sassa*, *Sassa*—*g. Seneca*, *Acaciæ gummi*—*g. Senega*, *Acaciæ gummi*—*g. Shrinking of the*, *Ulatrophia*—*g. Sweet liquidambar styraciflua*—*g. Tragacanth*, *Tragacanth*.

GUM TREE, *Brown*, see *Kino*.

GUM, WHITE, *Liquidambar styraciflua*, *Strophulus*—*g. Yellow*, *Icterus infantum*.

GUMMA, gen. *Gumma'tis*, (F.) *Gomme*. An elastic tumour formed in the periosteum occupying particularly the cranium and sternum, and produced by the syphilitic virus, when it has been long in the constitution. It is so called, because, when opened, it contains a matter like gum. These tumours—*gumma'ta*—are likewise termed *Tuber'cula gumma'ta* and *Tumores gumma'ti*, (F.) *Tumeurs gommeuses*.

GUMMI, (symp.) *Commi*, (F.) *Gomme*. An immediate principle of vegetables. It is a solid, uncrystallizable, inodorous substance, of a mawkish taste, unchangeable in the air, insoluble in alcohol, but soluble in water, with which it forms a mucilage. It is obtained from various species of *mimosa* and *prunus*; and consequently there are many varieties of gum. They are used in medicine as demulcents, emollients, and relaxants, particularly in catarrh, intestinal irritations, &c.; and in Pharmacy, are employed in the formation of emulsions, pills, &c.

GUMMI ACACIÆ ARABICA, *Acaciæ gummi*—*g. Acanthinum*, *Acaciæ gummi*—*g. Acaroides*, see *Xanthorrhæa*—*g. Adstringens* *Fothergilli*, *Kino*—*g. Ammoniacum*, *Ammoniacæ*—*g. Anime*, *Anime*—*g. Arabicum*, *Acaciæ gummi*—*g. Astragali* *Tragacantha*, *Tragacantha*—*g. Bogia*, *Cambogia*—*g. Brellis*, *Caranna*—*g. Copallinum*, *Copal*—*g. Elasticum*, *Coatucheou*—*g. Euphorbia*, see *Euphorbia officinarum*—*g. Gamandra*, *Cambogia*—*g. Gambiense*, *Kino*—*g. de Goa*, *Cambogia*—*g. Guttæ*, *Cambogia*—*g. Hedera*, see *Hedera helix*—*g. de Jemu*, *Cambogia*—*g. Juniperi*, *Sandaracæ*—*g. Lacææ*, *Lacææ*—*g. Ladanum*, see *Cistus creticus*—*g. Lamac*, *Acaciæ gummi*—*g. Laricis*, see *Pinus larix*—*g. Leucum*, *Acaciæ gummi*—*g. Mimosa*, *Acaciæ gummi*.

GUMMI NOS'TRAS, (F.) *Gomme du Pays*, *Indig'enous Gum*. These generic names are given to several species of gum, which flow spontaneously from certain indigenous fruit trees,—such as the almond, cherry, peach, apricot, &c. The indigenous gums have nearly the same properties as gum Arabic; but they are inferior to it.

GUMMI ORENBURGENSE, see *Pinus larix*—*g. Pannacis*, see *Pastinaca opoponax*—*g. ad Podagram*, *Cambogia*.

GUMMIRESI'NA, *Gumresin*. A milky juice, obtained by making incisions into the branches, stalks and roots of certain vegetables. Gum-resins are compounds of resin, gum, often with essential oil, and different other vegetable matters. They are solid, opaque, brittle, of a strong odour, acid taste, variable colour, and are heavier than water. Water dissolves a part of them, and alcohol another; hence proof spirit is the proper menstruum. The generality of the gum-resins are powerful stimulants to the whole or to parts of the economy. The chief are *asafoetida*, *gum-ammoniac*, *euphorbium*, *gamboge*, *gummyrr*, *olibanum*, *opoponax*, *scammony*, *aloes*, &c.

GUMMI RUBRUM ADSTRINGENS GAMBIENSE, see *Butea Frondosa*, *Kino*—*g. Seneca*, *Acaciæ gummi*, *Senegal*, *gum*—*g. Senega*, *Senegal*, *gum*—*g. Senegalense*, *Senegal*, *gum*—*g. Senica*, *Senegal*, *gum*—*g. Serapionis*, *Acaciæ gummi*—*g. Thebaicum*,

Acaciæ gummi—g. Tragacantha, Tragacantha—g. Uralense, see Pinus larix.

GUMMIODES. (from *gummi*.) Mucilaginous.

GUMMIODES, Mucilaginous.

GUMMOSUS, Mucilaginous.

GUMS. (Sax. *goma*, 'the gum,') Gingivæ.

GUNCOTTON, see Collodion—g. c. Ethereal solution of, Collodion.

GUNJAH, see Bangué.

GUNNERA PERPEN'SA, (after J. Ernst Gunner, a botanist and bishop of Drontheim.) A South African plant, *Nat. Ord.* Urticaceæ; the decoction of which is taken as a domestic remedy by the farmers, as a tonic in dyspepsia. A tincture has been used in gravel. An infusion of the leaves is demulcent, and is employed in pulmonary affections. The leaves are applied, fresh, to wounds and ulcers.

GURGES, gen. *Gur'gitis*, Pharynx.

GURGITELLO, MINERAL WATERS OF. A thermal spring in the isle of Ischia. Temp. at its source, 176° Fah. It contains carbonic acid, carbonates of lime, magnesia, iron and soda, sulphates of lime and soda, chloride of sodium and silica.

GURG'LING. 'Gushing with noise,' as water from a bottle. (Same etymon as *gargling*.) (F.) *Gargouillement*. The rhonchus or *râle* heard on auscultation when there is a cavity in the lungs containing pus. It is the 'cavernous rattle or *rhonchus*,' *Râle caverneux*. The size of the bubbles heard varies, and hence the rhonchus has been called *cavernous* and *cavernulous*, (F.) *Râle cavernuleux*. If the cavern be large, this *râle* will nearly resemble the gurgling of a bottle (*goulglou d'une bouteille*); if, on the contrary, the cavern be small, it will not differ from the *râle muqueux*.

GURGULIO, (like *curculio*.) Penis, Uvula.

GUSTATIF (Nerf.) (from *gustare*, *gustatum*, 'to taste,') see Lingual Nerve.

GUSTATION, Degustation, Taste.

GUSTATORY NERVE, see Lingual Nerve.

GUSTUS, Taste—g. Depravatus, Paragustia.

GUT, Gont, Intestine—g. Blind, Cæcum.

GUTS, SLIPPERINESS OF THE, Lientery.

GUTTA, ('a drop,') Apoplexy, Cambogia, Gout.

GUTTA. A Drop, *Stalagma*, *Stranz*, *Alun'sel*, *Stilla*, (F.) *Goutte*. A quantity of liquid, generally valued, in pharmacy, at the weight of a grain. The weight, however, varies, according to different circumstances, as the degree of tenacity of the fluid, and the extent of moist surface to which the suspended drop is attached before it falls; and it was found by Mr. Alsop to be influenced by the size of the bottle, and the angle of inclination at which it was held during the operation of dropping. The following are some of his results as to the number of drops required to measure a fluidrachm, when dropped from a large and a small bottle.

	From a large bottle.	From a small bottle.
(53) Diluted sulphuric acid . . .	24 drops	84 drops
Schœnle's hydrocyanic acid . . .	35 . . .	70
Distilled water . . .	31 . . .	54
Solution of ammonia . . .	40 . . .	48
Tincture of opium . . .	84 . . .	135
Rectified spirit . . .	100 . . .	130
Tincture of chloride of iron . . .	100 . . .	150

Prof. Wm. Procter, of Philadelphia, has given a table of the number of drops to a fluidrachm of essential oils, when dropped from the bottles from which they are commonly dispensed, and from a minim measure; and Mr. Edward Parrish one of different liquids dropped, under like circumstances, from pint and half pint tincture bottles, and from a minim measure. The following is the number of drops of different liquids equivalent to a fluidrachm—the result of the observations of Mr. Luzzard, of Philadelphia.

One fluidrachm of	Drops.
Acid. acetic, (crystallizable) . . .	contains 120
— hydrocyanic, (medicinal) . . .	" 45
— muriatic, . . .	" 54
— nitric, . . .	" 84
— diluted, (1 to 7,) . . .	" 51
— sulphuric, . . .	" 90
— aromatic, . . .	" 120
— diluted, (1 to 7,) . . .	" 51
Alcohol, (rectified spirit,) . . .	" 138
— diluted, (proof spirit,) . . .	" 127
Arsenite of potassa, solution of, . . .	" 67
Ether sulphuric, . . .	" 150
Oils of aniseed, cinnamon, cloves, pepper, mint, sweet almonds, and olives, } . . .	" 120
Solution of ammonia, (strong,) . . .	" 54
— (weak,) . . .	" 45
Tinctures of assafœtida, foxglove, guaiac, and opium, } . . .	" 120
Tincture of chloride of iron, . . .	" 132
Vinegar, distilled, . . .	" 78
— of colchicum, . . .	" 78
— of opium, (black drop,) . . .	" 78
— of squill, . . .	" 78
Water, distilled, . . .	" 45
Wine, (Teneriffe,) . . .	" 78
— antimonial, . . .	" 72
— of colchicum, . . .	" 75
— of opium, . . .	" 78

By the plural—*guttæ*—is meant a pharmaceutical mixture—*Mistura contracta*—to be given in drops. *Gutta* and *Guttæ*, also, signify certain cutaneous spots or eruptions.

GUTTA GAMANDRÆ, Cambogia—g. Gamba, Cambogia—g. Opaca; Cataract.

GUTTA PER'CHA, *Gutta tin'ban*, *Gomme getta'nia*. The concrete juice of a tree—*Isanan'dra gutta*—which is indigenous in Singapore and its vicinity, and belongs to the *Natural Order* Sapotaceæ. Plunged in boiling water it softens, when it may be moulded like caoutchouc to any form, which it retains on cooling. Splints and other instruments have been made of it.

A solution of *gutta percha*, in chloroform—*Tran'maticine*—has been used with advantage, topically, in various cutaneous affections, serofulous and indolent ulcers, and as an ectrotic in small-pox.

GUTTA ROSACEA SYPHILITICA, Crown of Venus. GUTTA RO'SEA, *G. Rosa'cea*, *Ion'thus corymbifer*, *Crus'ta serpiginosa* seu *pruriginosa*, *Acnè rosa'cea*, *Rose'ola acnè'sa*, *Thylactitia*, *Bac'chia*, *Butiga*, *Cur'buncled Face*, *Ro'sy Drop* or *Welk*, *Coppernose*, *Bottlenose*, *Grogglossoms*, *Groggrokes*, *Brandyface*, *Rumbud*, (F.) *Copperose*, *Goutte Rose*, *Bourgeois*, *Dartre pustulense copperose*. An eruption of small, suppurating tubercles, with shining redness, and an irregular granular appearance of the skin of the part of the face which is affected. The redness commonly appears first at the end of the nose, and then spreads on both sides. It is often produced by hard drinking. Its cure must be attempted by regular regimen, and cooling means internally; weak spirituous or saturnine lotions externally. The affection is usually very obstinate.

GUTTA SERENA, Amaurosis—g. Tuban, *Gutta percha*.

GUTTÆ, pl. of Gutta—g. Abbatiss Rousseau, *Laudanum abbatiss Rousseau*—g. Acidæ tonicæ, *Elixir Acidum Halleri*—g. Ammoniaci, see *Ammoniac gum*—g. Nervine, *Alcohol sulfurico-æthereus ferri*.

GUTTÆ NIGRÆ, 'black drops,' *Acetum o'pii*, *Common Black Drop*, (F.) *Gouttes noires*. (Opii, 3 viij, Aceti distillat. ℥ij. Infuse.)

The celebrated *Black Drop*, *Lan'caster* or *Quaker's Black Drop*, may be made as follows. Take half a pound of opium, sliced; three pints of good verjuice, (juice of the wild crab,) one and a half ounce of nutmegs, and half an ounce of saffron. Boil to a proper thickness, and add a quarter of a pound of sugar, and two spoonfuls of yeast. Set the whole in a warm place, near the fire, for six

or eight weeks; then place it in the open air until it becomes a syrup. Lastly, decant, filter, and bottle it up, adding a little sugar to each bottle. One drop is equal to three of laudanum; and it is nearly devoid of all the unpleasant exciting effects of the latter. An analogous formula is contained in the Pharmacopœia of the United States under the name *Ace'tum opii, Vinegar of Opium*. In the last edition, (1863,) a more easy mode of preparation is given. It consists in the entire extraction of the opium and aromatics by diluted acetic acid; the addition of sugar; and the dilution of the whole, so that a fluidounce shall contain 75 grains of opium. A grain of opium is represented by about $6\frac{1}{2}$ minims.

GUTTERIA, (from *guttur*), Bronchocele.

GUTTUR, gen. *Gutturis*. The throat; the larynx; the trachea.

GUTTUR GLOBOSUM, ('globous throat,') Bronchocele—g. Tumidum, Bronchocele.

GUTTURAL, *Gutturālis*, (from *guttur*.) Relating or belonging to the throat.

The Superior Thyroideal Artery is sometimes called *Gutturāl Artery*.

A *Gutturāl Cough* is one occasioned by irritation of the larynx or trachea.

A *Gutturāl Sound* is one produced, as it were, in the throat.

GUTTURALIS CARTILAGO, Arytenoid Cartilage.

GUTTURIS OS, ('bone of the throat,') Hyoides os.

GUTTURNIA, Arytenoid cartilages.

GYMNA, (from *γυμνος*, 'naked,') Genital Organs.

GYMNASION. (*γυμνασιον*.) Exercise.

GYMNASIUM, (from *γυμνασιον*.) *Palestra*. An establishment amongst the ancients, intended for bodily exercises, as wrestling, running, &c.; a term now used, in Germany more especially, for an academy or higher school.

GYMNAST, *Gymnas'tes*, (*γυμναστής*.) *Gymnas'ta*. The manager of a gymnasium. One, whose profession it is to prevent or cure diseases by gymnastics.

GYMNASTICS, *Gymnas'tica*, (*γυμναστική*, (*τεχνη*),) *Medicīna gymnas'tica seu eue'tica*, *Somascel'tica*, (F.) *Gymnastique*, *Somascel'tique*. That part of hygienic medicine which treats of bodily exercises. It is called *Medical Gymnastics*. The ancients had also *Athletic Gymnastics*, and *Military Gymnastics*. Herodotus, of Selivraea, first proposed gymnastics for the cure of disease.

GYMNASTIQUE, Gymnastics.

GYMNEURA LACTIFERUM, *Ceylon cow-tree*, *Kiriaghuma*. A tree, of the Family Apocynaceae, the juice of which is used by the Cingalese as milk.

GYMNOCLADUS CANADENSIS, *Cof'fee Tree*, *Kentuck'y Cof'fee-bean Tree*, *Mahog'any*, *Nick'ar Tree*, *Bon'due*. Order, Leguminosae; (*γυμνος*, 'naked,' and *κλαδος*, 'a branch,') An indigenous tree, which grows from Ohio to Louisiana. The leaves are cathartic, and said to contain cystin. The seeds are a good substitute for coffee.

GYMNOGRAMME CETERACH, *Asplenium ceterach*.

GYMNOPTERIS CETERACH, (*γυμνος*, 'naked,' and *πτερις*, *ptēris*, 'fern,') *Asplenium ceterach*.

GYMNOSIS, (*γυμνωσις*, from *γυμνος*, 'naked,') Denudation.

GYNECANER, (*gynæco*, and *ανηρ*, 'a man,') *Vir effemina'tus*. An effeminate man.

GYNECANTHE, (*gynæco*, and *ανθος*, 'a flower,') *Tamus communis*.

GYNECATOP'TRON, (*gynæco*, and *κατοπτρον*, 'a speculum,') A speculum for investigating the female organs of generation.

GYNECEA, *Gynæceia*.

GYNECEIA, (*γυναικεια*.) *Gynæci'a*, *Gynæce'a*.)

The catamenia;—the lochia. The diseases of women in general.

GYNÆCEUM, Antimonium, Vulva.

GYNÆCEUS, (*γυναικειος*,) Belonging to women. Female. Feminine.

GYNÆCIA, (*γυναικεια*.) *Gynæceia*, *Menses*.

GYNÆCO, (from *γυνη*, gen. *γυναικος*, 'a woman,') In composition, woman.

GYNÆCOLOG'IA, *Gynæcol'ogy*, (*gynæco*, and *λογος*, 'a description,') The doctrine of the nature, diseases, &c. of women.

GYNÆCOMA'NIA, (*gynæco*, and *μανια*, 'mania,' 'rage,') That species of insanity, which arises from love for women. Some have used the word synonymously with nymphomania.

GYNÆCOMAS'TUS, *Gynæcomas'thus*, (*gynæco*, and *μαστος*, 'a breast,') A man whose breasts are as large as those of a woman. The condition is called *Gynæcoma'zia*.

A considerable enlargement of the breasts of a female was formerly called *Gynæcomas'ton*.

GYNÆCOMA'ZIA, see *Gynæcomastus*.

GYNÆCOMYST'AX, gen. *Gynæcomyst'acis*, (*gynæco*, and *μυσταξ*, 'the beard,') The hair on the pubes of women.

GYNÆCOPHO'NUS, (*gynæco*, and *φωνη*, 'voice,') A man who has an effeminate voice.

GYNANDRIA, (*gyne*, and *ανηρ*, gen. *ανδρος*, 'a man,') Hermaphroditism.

GYNANDROP'SIS PENTAPHYL'LA, *Cleo'nē pentaphylla*; Ord. Cappari'daceae; an indigenous plant, is possessed of diaphoretic properties. The juice alone, or mixed with oil, has been used in earache.

GYNANDRUS, (*gyne*, and *ανηρ*, gen. *ανδρος*, 'a man,') *Gynanthropus*.

GYNANTHRO'PUS, *Gynan'drus*, (*gyne*, and *ανθρωπος*, 'a man,') An hermaphrodite who belongs more to the male than to the female sex.

GYNATRE'SIA, (*gyne*, and *ατρησις*, 'imperforate,') Closure or imperforation of the external parts of generation of the female.

GYNE, (*γυνη*.) Female. Woman.

GYNECOLOGY, *Gynæcologia*.

GYNIACI, (MORBI,) (from *gynē*.) Developmental diseases of women, as chlorosis; those attending the parturient state and amenorrhœa, according to the classification of Dr. W. Farr.

GYNIDA, Hermaphrodite.

GYNOARIUM, (*gyne*, and *ωαριον*, 'ovarium,') Ovarium.

GYNOCARDIA ODORATA, (*gyne*, and *καρδια*, 'a heart,') *Chaulmoogra*.

GYNOPLAS'TIC, *Gynoplas'ticus*, (*gyne*, and *πλασσω*, 'I form,') The gynoplastic operation is employed for opening or dilating the closed or contracted genital openings of the female.

GYPSYWEED, *Lycopus sinuatus*, and *L. Virginicus*.

GYRI CEREBRI, (pl. of *gyrus*.) Convolutions (cerebral).—g. Cochleæ, *Scalæ* of the Cochlea—g. Intestinales, see *Anse*—g. Operi, see *Insula Cerebri*.

GYROMIA VIRGINICA, *Medeola Virginica*.

GYROPHLE, (from *caryophyllum*,) see *Eugenia caryophyllata*.

GYROPHORA, see *Trips de Roche*.

GYRUS, (*γυρος*, 'a circle,') Convolution.

GYRUS INTESTINALIS, *Anse intestinale*.

GYRUS FORNICA'TUS, 'arched circle or convolution.' *Fil'let of the cor'pus callo'sum*. A large convolution of the brain, which lies horizontally on the corpus callosum, and may be traced forwards and backwards to the base of the brain, terminating by each extremity at the fissure of Sylvius. The surface of the hemisphere, where it comes in contact with the corpus callosum, is bounded by it.

GYTLING, (dim. of *Se. get*, 'a child,') Infans.

H.

HABBERING (Sc.), (from Belg. *haperen*, 'to stutter.') Balbuties.

HABEL-KALIMBAT, *Pistacia terebinthus*.

HABBI, *Hagenia Abyssinica*.

HABBLING (Sc.), (from Su. G. *happila*, 'to stutter.') Balbuties.

HABENA, ('a bridle,' from *habeo*, 'to have or hold,') *Tel'amon*. A bandage for uniting the lips of wounds; which, of old, in many instances, replaced the suture.

HABENÆ OF THE PINEAL GLAND, see Pineal gland.

HABENULÆ OF THE PINEAL GLAND, (dim. of *habena*,) see Pineal gland.

HABIL'LA DE CARTHAGE'NA, (dim. of *haba*, (L.) *faba*, 'a bean,') *Bejuco*, *Carthage'na Bean*, (F.) *Fève de Carthagène*. A kind of bean of South America, famed as an effectual antidote for the poison of all serpents, if a small quantity be eaten immediately.

HABIT, *Habitu'do*, (*habere*, *habitu*, 'to have or to hold,') *Assuetu'do*, *Mos*, *U'ens*, *Consuetu'do*, *Ethos*, *Hexis*, *Ethemos'ynē*, (F.) *Habitude*, *Accoutumance*. Habit is the aptitude for repeating certain acts:—or, a facility, which results from the frequent repetition of the same act. It is, according to vulgar expression, 'a second nature.' Habit may predispose to certain diseases, or it may protect against them. It ought not to be lost sight of, in attending to the progress of disease, or its treatment.

HABIT OF BODY, *Constitu'tio*, *Hab'itus*. *H. Cor'poris*, *Catas'tasis*, *Hexis*, *Epiphani'a*, (F.) *Habitude ext'rieure*, *Habitude du Corps*. The aggregate of the physical qualities of the human body.

HABITUDE. Habit—*h. du Corps*, Habit of body—*h. Ext'rieure*, Habit of body.

HABITUDO, (*habere*, 'to have or to hold,') Habit.

HABITUS. Habit of body—*h. Apoplecticus*, Apoplectic habit—*h. Corporis*, Habit of body—*h. Quadratus*, see Apoplectic—*h. Torosus*, see Apoplectic.

HABROMA'NIA, (*ἄβρος*, 'gay,' and *μανία*, 'mania,') Insanity in which the delusions are of a gay character.

HACHICH, see Bangué.

HACHISCH, Bangué.

HACHE (Sc.), Pain.

HACKBERRY, *Celtis occidentalis*, *Prunus padus*.

HADDOCK, see Oleum jecoris aselli.

HÆMA, gen. *Hæ'matis*, (*ἷμα*, gen. *ἱματός*,) 'blood.'

HÆMACHROINE, (*hæma*, and *χρῶς*, 'to colour,') Hæmateine.

HÆMACHRO'SES, ([G.] *Hämachrosen*, from *hæma*, and *χρῶσις*, 'coloration,') A family of diseases in which the blood has its colour different from usual, as in purpura and cyanosis.

HÆMACTUS, (*ἱμακτός*,) Bloody.

HÆMACY'ANIN, (*hæma*, and *κῶνος*, 'blue,') A blue colouring matter, detected by Sanson in healthy blood, and in bile by some chemists, but not by others.

HÆMADON'OSUS, (*hæma*, and *ὄσος*, 'a disease,') *Hæmatangion'osus*, *Hæmatangiom'us*, *Hæmatangio'sis*. Disease of the bloodvessels.

HÆMADOSTO'SIS, (*hæma*, and *ὄσσις*, 'a bony tumour,') Ossification of the bloodvessels.

HÆMADROMOMETER, Hæmodromometer.

HÆMADYNAMETER, Hæmadynamometer.

HÆMADYNAMICS, (F.) *Hémadynamique*,

(*hæma*, and *δυναμῖς*, 'power,') The mechanism of the circulation of the blood.

HÆMADYNAMOMETER, *Hæmodynamom'eter*, *Hæmatodynamom'eter*, *Hæmodynami'eter*, *Hæmatom'eter*, (F.) *Hémodynamomètre*, *Hæmadynamètre*, *Hémomètre*, (*hæma*, *δυναμῖς*, 'power,' and *μετρον*, 'a measure,') An instrument for measuring the force of the blood in the vessels. It consists of a bent glass tube, the lower bent part of which is filled with mercury. A brass head is fitted into the artery, and a solution of carbonate of soda is interposed between the mercury and the blood, which is allowed to enter the tube for the purpose of preventing its coagulation. The pressure of the blood on the mercury in the descending portion of the bent tube causes the metal to rise in the ascending portion; and the degree to which it rises indicates the pressure under which the blood moves. A modification of this instrument has been called *Cardiom'eter*, (F.) *Cardiomètre*.

HÆMAGASTER, Hæmatogaster.

HÆMAGASTRIC or HÆMOGASTRIC PESTILENCE, Fever, yellow.

HÆMAGOGUM, *Pæonia*.

HÆMAGO'GUS, (*hæma*, and *αγω*, 'I drive off,') A medicine which promotes the menstrual and hemorrhoidal discharges.

HÆMAL, (from *hæma*,) Relating to the blood or bloodvessels.

HÆMAL ARCH. The arch formed by the projections anteriorly of the ribs and sternum for the bodies of the vertebrae. It encloses the great bloodvessels.

HÆMAL AXIS, Aorta—*h. Spine*, Sternum.

HÆMALOPIA, (*hæma*, and *ὤψ*, 'the eye,') Hemophthalmia.

HÆMALOPIS, Hæmophthalmia.

HÆMALOPS, (*ἱμαλωψ*,) Hemophthalmia.

HÆMAN'THUS COCCIN'EUS, (*hæma*, and *ανθος*, 'a flower,') A beautiful South African plant. *Nat. Ord.* Amaryllidæe, the bulb of which is employed as a diuretic. It is given as an oxymel in asthma and dropsy. The fresh leaves are antiseptic, and applied to foul, flabby ulcers, and in anthrax.

HÆMANTLI'A, (*hæma*, and *αντλεω*, 'I pump out,') An exhausting syringe, to which a reservoir is attached, containing a needle. It is used for obtaining blood for hæmatological investigations.

HÆMAPERITONIRRHAG'IA, (F.) *Hæmoperitonirrhagie*, (*hæma*, *περιτοναϊον*, 'peritoneum,' and *ῥαγν*, 'a rupture,') An exhalation of blood into the peritoneum.

HÆMAPHÆ'IN, *Hæmatophæ'um*, *Hæmophæ'um*, (*hæma*, and *φαῖος*, 'of a dusky colour,') A term applied by Simon to the brown colouring matter of the blood, supposed by some to be nothing more than hæmatin modified by an alkali. In the urine it is called *Urobæ'matin*.

HÆMAPHOBUS, *Hæmophob'us*, (*hæma*, and *φοβος*, 'dread,') One who has a dread of blood:—who cannot look at it without fainting.

HÆMAPOPHYSES, (*hæma*, and *αποφω*, 'I shoot forth,') Costal cartilages, see Costa.

HÆMAPORIA, *Hæmatopor'ia*, *Hæmatopor'ia*, *Oligophæ'mia*, *Oligæ'mia*, (*hæma*, and *πορος*, 'poor,') Paucity of blood. See Anæmia.

HÆMPTYSIS, (*ἱμαπτυσίς*,) Hæmoptysis.

HÆMAS, (*ἱμας*, gen. *ἱματός*,) A bloodvessel.

HÆMASTATICA, Hæmatostatica.

HÆMASTATICS, *Hæmatostat'icæ*, (*hæma*, and *στασις*, 'statics,') The statics of the blood and bloodvessels; *Hæmatat'icæ*.

HÆMAT'ALLOSCOPY. *Hæmat'oscop'ia*, (*hæmato*, αλλοιος, 'different,' and σκοπεῖν, 'to view.') The mode of distinguishing different kinds of blood from each other.

HÆMATANAGOGĒ, (*hæmato*, ἀνα, and ἀγω, 'I drive off,') *Hæmatemesis*.

HÆMATANGIONOSUS, (*hæmato*, αγγειον, 'a vessel,' and νόσος, 'a disease,') *Hæmadonosus*.

HÆMATANGIOSIS, *Hæmadonosus*.

HÆMATAPORIA, *Hæmaporia*.

HÆMATAPORRHOSIS, (*hæmato*, ἀπο, and ῥησις, 'serum,' and ὁσις, 'Cholera.

HÆMATAPOSTE'MA, gen. *Hæmataposte'matis*, (*hæmato*, and ἀποστήμα, 'an abscess,') An abscess which contains blood.

HÆMATAULICA, (*hæmato*, and αὐλος, 'a tube,') Vascular system.

HÆMATEOLYSIS, *Hæmateol'ia*, *Hæmateol'is*, *Hæmateol'yis*, *Hæmateosol'is*, *Saŋ'guinis dissolutio*, (*hæmato* and ἐκλύσις, 'loosening,') Dissolution of the blood.

HÆMATELĒUM, (*hæmato*, and ελαῖον, 'oil,') see Blood.

HÆMATEMESIS, (*hæmato*, and ἔμω, 'I vomit,') *Hæmorrah'gia*, *Hæmatem'esis*, *Vom'itus cruen'tus*, *Vom'itus seu Vomit'io Saŋ'guinis*, *Gastrorrhag'ia*, *Gastro-hæmorrah'gia*, *Hæmorrah'gia ventric'uli*, *Hæmorrah'gia ventric'uli*, *Hæmorrah'gia ventric'uli*, *Vomiting of Blood*, (F.) *Hématisémèse*, *Vomissement de sang*. *Hæmatemesis* is generally preceded by a feeling of oppression, weight, and dull or pungent pain in the epigastric and in the hypochondriac regions: by anxiety, and, occasionally, by syncope. Blood is then passed by vomiting, and sometimes, also, by stool,—the blood being generally of a gummy aspect. *Hæmatemesis* may be active or passive, acute or chronic. The blood effused proceeds from rupture of vessel, or from a sanguineous exhalation at the surface of the mucous membrane of the stomach. It is often observed in females whose menstrual secretion is irregularly performed. It is not of much danger, except when connected with disease of some of the solid viscera of the abdomen. On dissection of those who have died from protracted hæmatemesis,—for the acute kind is comparatively devoid of danger,—the mucous membrane of the stomach is found red and inflamed, or black, and the vessels considerably dilated. Complete abstinence from food; rest; the horizontal posture; bleeding, if the hæmorrhage be active; cold, acidulous drinks, &c., constitute the usual treatment.

HÆMATENCEPHALUM, (*hæmato*, and ἐνκεφαλον, 'Apoplexy.

HÆMATEPAGOGĒ, (*hæmato*, ἐπι, and ἀγω, 'I drive,') Congestion.

HÆMATERUS, (*hæmateros*; from αἷμα, 'blood,') Sanguine.

HÆMATERYTHRUM, (*hæmato*, and ἐρυθρος, 'red,') *Hæmatin*.

HÆMATEXOOSTOSIS, (*hæmato*, and ὀστωσις, 'Osteosarcoma.

HÆMATHIDROSIS, (*hæmato*, and ἰδρωσις, 'sweating,') *Sudor cruentus*.

HÆMATHORAX, *Hæmatothorax*.

HÆMATIA, (from hæma,) *Jusculum nigrum* *Lacedæmoniurum*.

HÆMATIASIS, (from hæma,) *Hæmatonosus*.

HÆMATIC, (from hæma,) Sanguine—h. Crystals, *Hæmatoidin*.

HÆMATICA, (from hæma,) Diseases of the sanguineous function:—the third class in the nosology of Good. Also, medicines that act on the blood, (Pereira.)

HÆMATICA DOCTRINA, *Hæmatology*.

HÆMATICUS, (*hæmaticos*; from hæma,) Bloody, Sanguine.

HÆMATIDROSIS, (*hæmato*, and ἰδρωσις, 'sweating,') *Sudor cruentus*.

HÆMATIN, *Hæmat'ina*, *Hem'atine*, *Hem'itia*, *Hæmatosin*, *Hem'atosine*, *Hæmater'ylthrum*, *Hæmater'ylthrum*, *Zoöhæmat'in*, *Hæmoch'roin*, *Hæmatech'roin*, *Phæ'uadine*, *Phæ'uicine*, *Hæmatech'roite*, *Cru'orin*, *Rubrin*, *Glob'ulin* of some; *Rouge de Sang*; (from hæma.) The red colouring matter of the blood. It resides in distinct particles or globules, and, in the opinion of some observers, in the envelope of the globules. It appears to be of a peculiar character, and one that has not yet been determined by the chemist. That the colour of the blood is not owing to the peroxide of iron which it contains is shown by the fact mentioned by Scherer, that he removed the iron by acids, and yet a deep red tincture was formed when alcohol was added to the residuum.

HÆMATIN'IC, *Hæmatin'ic*, *Hæmatin'icous*, (from hæmatin.) An agent that augments the number of red corpuscles or the hæmatin of the blood.

HÆMATINIC CRYSTALS, *Hæmatoidin*.

HÆMATINUS, (from hæma,) Sanguine.

HÆMATISCHESIS, *Hæmis'chesis*, *Hæmocr'iph'ia*, (*hæma*, and ἰσχύω, 'to suppress,') The retention or suppression of a natural or artificial flow of blood.

HÆMATISMUS, (from hæma,) Epistaxis, *Hæmorragia*.

HÆMATISTH'MUS, (*hæmato*, and ὁσμος, 'pharynx,') Hemorrhage into the pharynx.

HÆMATITES, *Hæmat'itis*, (*hæmatites*.) Resembling blood. *Lapis Hæmat'ites*. A beautiful ore of iron, called also, *Bloodstone*, *Ochrea rubra*, *Oxydum fer'ricum crystallizatum nativum*, *Ferum Hæmatites*. When finely levigated, and freed from the grosser parts, by frequent washings with water, it has been long recommended in hæmorrhage, fluxes, uterine obstructions, &c., in doses of from one scruple to three or four. Also, a vessel that contains blood, *hæmatitis phleg.*

HÆMATITIS, *Hæmitis*.

HÆMATMUS, (*hæma*, and σμός, 'vapour,') *Gaz sanguinis*.

HÆMATO, (from αἷμα, gen. αἵματος, 'blood,') in composition, blood.

HÆMATOCATHAR'TICA, (*hæmato*, and καθάρσις, 'purification or purgation,') Remedies for purifying the blood.

HÆMATOCELE, (*hæmato*, and κλη, 'tumour,') A tumour formed by blood. By some, this term has been applied to a tumour formed by blood, effused into the areolar texture of the serotum. Others have used it for tumours arising from effusion of blood into the tunica vaginalis:—*hæmatocèle*, according to them, differing from hydrocèle, only in the character of the effusion. (Heister.) Others, again, have applied it to effusions of blood into the interior of the tunica albuginea itself. (Richter.) The first is the usual acceptance. It is most commonly caused by wounds or contusions: and requires the use of antiphlogistics, discutients, &c. Sometimes it is necessary to evacuate the effused blood.

HÆMATOCELE ARTERIOSA, *Aneurism*.

HÆMATOCEPH'ALUS, (*hæmato*, and κεφαλή, 'head,') (F.) *Hæmatoceph'ale*. A monster in which an effusion of blood in the head has given occasion to more or less singular deformities.

HÆMATOCEREBRIFORM DISEASE, (*hæmato*, cerebrum, 'brain,' and forma, 'form,') *Encephaloid*.

HÆMATOCHEZ'IA, (*hæmato*, and χεῖω, 'I go to stool,') *Sedes cruen'ta*, *Cutarrhex'is recta*, *Enterorrhag'ia simp'lex*, *Diarrhæ'a cruenta seu sanguinolenta*, *Hæmato-diarrhæ'a*, *Hæmorrah'gia intestino'rum*, *Enter'o-hæmorrah'gia*, (F.) *Hæmorrah'gie des intestins*, *Écoulement de sang par l'Intestin*. Discharge of blood by stool. See *Melæna*.

HÆMATOCHROITE, (*hæmato*, and χροα, 'colour,') *Hæmatin*.

HÆMATOCHYSIS, (*hæmato* and *χω*, 'to pour out.') Hæmorrhagia.

HÆMATOCE'LLIA, (*hæmato*, and *κελία*, 'the cavity of the abdomen.') Effusion of blood into the abdomen.

HÆMATOCOL'PUS, (*hæmato*, and *κολπος*, 'vagina.') Effusion of blood into the vagina. Accumulation of blood owing to occlusion of the vagina.

HÆMATOCRYSTALLIN, (*hæmato*, and *κρυσταλλος*, 'a crystal,') Hæmatoidin.

HÆMATOCYANOPATHIA, (*hæmato*, *κυανος*, 'blue,' *παθος*, 'affection,') Cyanopathy.

HÆMATOCYANOSIS, (*hæmato*, and *cyanosis*, Cyanopathy.

HÆMATOCYSTE, (*hæmato*, and *κυστις*, 'a bladder,') Hæmatoma saccatum.

HÆMATOCYST'IS, (*hæmato*, and *κυστις*, 'bladder,') Hemorrhage into the bladder.

HÆMATO'DES, *Hæmatoïdes*, *Sanguin'eus*, *Hæmatoid*, *Hem'atoid*, ('*αἱματώδης*, from *hæmato*, and *odes*.) That which contains blood, or has the character of blood.

HÆMATODES FUS'GUS, *Hæmatom'yces*, *Fungus canero'sus hæmato'des*, *Mele'na fungo'sa carci'noïdes*, *Angidiospon'gus*, *Angiom'yces*, *Hæmatospon'gus*, *Tu'mor fungo'sus sanguin'eus seu anom'alus*, *Carcino'ma hæmato'des*, *Carcino'ma Spongio'sum*, *Spon'goid inflammation*, *Pulp'y* or *Med'ullary Sarco'ma*, *Bleed'ing Fungus*, *Soft Can'cer*, (F.) *Carcinôme sanglant*, *Hématoncie fongôide*. An extremely alarming carcinomatous affection, which was first described, with accuracy, by Mr. John Burns, of Glasgow. It consists in the development of cancerous tumours, with violent heat and pain, and with fungus and bleeding excrescences. Even when the diseased part is extirpated at a very early period, recovery rarely follows; other organs being generally implicated at the same time. Fungus hæmatodes was the term first applied to the disease by Mr. Hey of Leeds. Mr. J. Burns called it *Spongoid inflammation*, from the spongy, elastic feel, which peculiarly characterizes it, and continues even after ulceration has taken place. The disease has, most frequently, been met with in the eyeball, the upper and lower extremities, testicle and mamma; but it occurs in the uterus, ovary, liver, spleen, brain, lungs, thyroid gland, and in the hip and shoulder joints.

Some French surgeons designate, by this name, those tumours which were formerly termed *anormales*, *caverneuses*, *variqueuses*, called *Erectiles* by Dupuytren, *Hématoncies*, by Alibert, and *Telan-giectasie* by Græfe.

HÆMATODIARRHŒA, Hæmatochezia.

HÆMATODYNAMOMETER, Hæmadynamometer.

HÆMATOGASTER, (*hæmato*, and *γαστρον*, 'stomach,') Effusion of blood into the stomach.

HÆMATOGENESIS, Hæmatosis.

HÆMATOGENETICA, (*hæmato*, and *γενεσις*, 'generation,') Hæmatopoetica.

HÆMATOGLOBULIN, see Globules of the blood.

HÆMATOGRA'PHIA, *Hæmatog'raphy*, (*hæmato*, and *γραφία*, 'a description,') A description of the blood.

HÆMATOID, (*hæmato*, and *ειδος*, 'resemblance,') Resembling blood. Hæmatodes.

HÆMATO'IDIN, *Hæmatocrys'tallin*, *Hæmatin'ic*, *hæmat'ic*, *hæmin* or *blood crystals*, (F.) *Hématôidine*, *Crystaux d'Hématine*. A substance found in the midst, or in the vicinity, of effusions of blood, which is either in amorphous masses, or, more frequently, in minute rhomboidal crystals, or in fine needles. It is considered to be hæmatin separated from the globules of the blood, and passed from the state of a non-crystallizable organic substance to one of a crystalline chemical compound.

HÆMATOL'OGY, *Hematol'ogy*, *Hæmatolog'ia*,

Hæmat'ica doctri'na, (F.) *Hématologie*, (*hæmatr*, and *λογος*, 'a discourse,') That part of medicine which treats of the blood.

HÆMATOLOGY, PATHOLOG'ICAL, (F.) *Hématologie pathologique*. Observation of the blood to detect its varying characters in disease.

HÆMATOLY'SES, (G.) *Hæmatolysen*, (*hæmato*, and *λυσις*, 'solution,') An order of diseases in which there is diminished coagulability of the blood.

HÆMATOLYSIS, Hæmateclysis.

HÆMATOLYTICUM, Spanamic.

HÆMATO'MA, (from *hæmato*, and *μα*,) *Thrombus*, *Tumor sanguin'eus*. A bloody tumour, especially of the scalp of the newborn.

HÆMATO'MA SACCA'TUM. An encysted tumour containing blood—*Hæmatocys'tis*.

HÆMATOMANTI'A, (*hæmato*, and *μαντεία*, 'divination,') Judgment of disease from the appearance of the blood.

HÆMATOMATRA, (*hæmato*, and *ματρα*, 'uterus,') Metrorrhagia.

HÆMATOMETACHYSIS, (*hæmato*, *meta*, and *χω*, 'I pour,') Transfusio sanguinis.

HÆMATOMETRA, (*hæmato*, and *μητρα*, 'the uterus,') Metrorrhagia.

HÆMATOMMA, gen. *Hæmatom'matis*, (*hæmato*, and *omma*, 'the eye,') Hæmophthalmia.

HÆMATOPHALOCE'LE, *Hæmatom'phalum*, *Hæmatom'phalus*, (*hæmato*, *ομφαλος*, 'the navel,' and *κηλη*, 'a tumour,') Umbilical hernia, the sac of which encloses a bloody serum, or which has, at its surface, a number of varicose veins; constituting *Varicom'phalus*.

HÆMATOPHALUM, (*hæmato*, and *ομφαλος*, 'the navel,') Hæmatomphaloele.

HÆMATOMYCES, (*hæmato*, and *μυκης*, 'a fungus,') Hæmatodes fungus.

HÆMATO'CUS, (F.) *Hématoncie*, (*hæmato*, and *οκος*, 'a tumour,') Alibert has given this name to *Nævi mater'ni*, *Varicose tumours*. He admits three varieties:—the *H. fongôide*, *H. fram-boisée*, and *H. tubéreuse*.

HÆMATO'OSUS, *Hæmat'osis*, *Hæmatopa-thi'a*, *Hæmopathi'a*, *Nosohæ'mia*, (*hæmato*, and *νοσος*, 'disease,') A disease of the blood. A hæmic or blood disease. (F.) *Nosohémie*.

HÆMATOPATHIA, (*hæmato*, and *παθος*, 'an affection,') Hæmatonosus.

HÆMATOPEDESIS, (*hæmato*, and *πῶσις*, 'I spring,') see Diapedesis.

HÆMATOPERICAR'DIUM, (*hæmato*, and *pericardium*,) Effusion of blood into the pericardium.

HÆMATOPHÆUM, Hæmaphavin.

HÆMATOPHIL'IA, *Hæmophil'ia*, *Hæmorrhophil'ia*, *Idiosyncra'sia hæmorrhag'ica*, (F.) *Hæmophilie*, (*hæmato*, and *φιλειω*, 'I love,') A hemorrhagic diathesis.

HÆMATOPHLEBOS'TASIS. Sudden suppression of a hemorrhage:—from *στασις αἱματος φλεβων*, 'suppression of the blood of the veins.'

HÆMATOPHOB'IA, *Hæmaphob'ia*, (F.) *Hæmophobia*, (*hæmato*, and *φοβος*, 'dread,') Dread or horror at the sight of blood, producing syncope, &c. Of old, they who were afraid of, or opposed to, bloodletting, were called *hæmaph'oboi*, *blood-funkers*.

HÆMATOPHTHALMIA, Hæmophthalmia.

HÆMATOPH'THORES, (G.) *Hæmatophthoren*; (*hæmato*, and *θωρα*, 'corruption,') An order of diseases in the classification of Fuchs, in which the blood is materially altered in composition, as in typhus.

HÆMATOP'ISIS, (F.) *Hématopisie*. M. Capuron, of Paris, has applied the term *Hématopisie utérine*, from analogy with *Hydropisie utérine*, to a collection of blood which sometimes takes place in the uterus, when, owing to faulty conformation, the exit of the menstrual flux is prevented.

HÆMATOPLA'NIA. *Hæmatoplane'sis*, *Hæmopla'nia*, (*hæmato*, and *πλαῖν*, 'wandering,') (F.) *Hæmoplanie*. A vicarious hemorrhage.

HÆMATOPLANIA MENSTRUALIS, Menstruation. *ricarius*.

HÆMATOPLASMA, gen. *Hæmatoplasm'atis*, *Liquor sanguinis*.

HÆMATOPLETHORA, *Plethora*.

HÆMATOPŒA, (*hæmato*, and *ποιω*, 'I make,')

Hæmatopoetica.

HÆMATOPOESIS, *Hæmatosis*.

HÆMATOPOËTICA, (*ἀματοποιητικά*,) *Hæmatopœ'a*, *Hæmatogenesi'ca*, 'bloodmakers.' Agents that favour hæmatosis. Iron was called by the ancients *metal'lum hæmatopœst'icum*.

HÆMATOPOIESIS, *Hæmatosis*.

HÆMATOPŒRIA, *Hæmaporia*.

HÆMATOPS, (*hæmato*, and *ὤψ*, 'the eye,')

Hæmophthalmia.

HÆMATOPSIA, *Hæmophthalmia*.

HÆMATOPTYSIS, *Hæmoptysis*.

HÆMATOPTYSIS, *Hæmoptysis*.

HÆMATORRHACHIS, (*hæmato*, and *ῥαχis*, 'the spine,') *Apoplexia myelitica*.

HÆMATORRHAGIA, *Hæmorrhagia*.

HÆMATORRHŒA, (*hæmato*, and *ῥέω*, 'to flow,') *Hæmorrhagia*.

HÆMATORRHOSIS, (*hæmato*, and *ῥοσ*, 'serum,') *Cholera*.

HÆMATORRHHYSIS, (*hæmato*, and *ῥέω*, 'to flow,') *Hæmorrhagia*.

HÆMATOSCHEOCE'LE, *Hæmoscheoce'le*, *Hæmatos'cheum*, (*hæmato*, *σχεον*, 'serotum,' and *κηλη*, 'tumour,') Effusion of blood into the serotum.

HÆMATOSCHEUM, (*hæmato*, and *σχεον*, 'serotum,') *Hæmoscheocele*.

HÆMATOSCOPIA, *Hæmoscop'ia*, (F.) *Hæmorhoscopia*, (*hæmato*, and *σκοπέω*, 'I view,') An examination of blood drawn.

HÆMATOSEPSIS, (*hæmato*, and *σψis*, 'putrefaction,') *Hæmateclisis*.

HÆMATOSIN, (from *hæma*,) *Hæmatin*.

HÆMATO'SIS, (*ἀματωσις*, from *hæmato*, and *οσις*,) *Exæmatosi'sis*, *Enhæmatosi'sis*, *Hæmatopoie'sis*, *Hæmatopœ'sis*, *Hæmatogen'esis*, *Procrea'tio san'guinis*.

Decarbiniza'tion or *Atmospheriza'tion* of the blood, *Sanguifica'tion*. The transformation of the venous blood and chyle into arterial blood by respiration. Called, also, *Aëra'tion*, and *Arterializa'tion*, of the blood. Formation of blood in general.

HÆMATOSPILIA, (*hæmato*, and *σπιλος*, 'a spot,') *Purpura hæmorrhagica*.

HÆMATOSPONGUS, (*hæmato*, and *σπυγγος*, 'a sponge, a fungus,') *Hæmatodes fungus*.

HÆMATOSTATICA, (*hæmato*, and *στατική*, 'statics,') *Ischæ'mia*, *Hæmostat'ica*, *Hæmostat'ica*, *Hæmostat'ics*, *Hæmostat'ics*, *Hæmostat'ics*, *Enæ'ma*, (F.) *Hæmostatistiques*, *Hæmostatiques*, *Hæmostat'iques*. The doctrine of the motion of the blood in living bodies. Also, remedies for stopping blood. *Styptics*; (from *hæmato*, and *στατικός*, 'restraining,')

HÆMATOSTEON, (*hæmato*, and *οστέον*, 'a bone,') Effusion of blood into the bones or joints. **HÆMATOSYMPHORESIS**, (*hæmato*, and *συμφορησις*, 'an accumulation,') *Congestion*.

HÆMATOSYNAGOGÉ, (*hæmato*, and *συναγωγή*, 'a collection,') *Congestion*.

HÆMATOTELANGIOSIS, (*hæmato*, *τελος*, 'end,' *αγγειον*, 'a vessel,' and *οσις*,) *Telangiectasia*.

HÆMATOTHORAX, gen. *Hæmatothora'cis*, *Hæmatho'raz*, *Hæmotho'raz*, *Hæmop'tysis inter'na*, *Pleurorrhæ'a sanguin'ea*, *Pleurorrhag'ia*, *Pneumorrhag'ia inter'na*, *Hæmorrhag'ic Pleurisy*, *Pleur'al hæmorrhage*, (from *hæmato*, and *θώραξ*, 'the chest,') *Extravasation* of blood into the chest.

HÆMATOTOXIC, *Toxæmic*.

HÆMATOXIC, *Toxæmic*.

HÆMATOXYLON, *H. Campechianum*.

HÆMATON'YLON CAMPECHIA'NUM, (*hæmato*, and *ῥυλον*, 'wood,') *Acacia Zeylan'ica*, *Logwood*. The part of the tree used in medicine is the wood, *Hæmatos'yli Liq'num*, *Lignum Campeche'n'se* seu *Campechi'num* seu *Campeesa'num* seu *Indicum* seu *Sappan* seu *Brasil'ianum* *rubrum* seu *cæru'leum*, *Hæmatos'ylium*, *Hæmatos'ylon* (Ph. U. S.), (F.) *Bois de Campêche*. Family, *Leguminosæ*. Ser. *Sypt. Decandria Monogynia*. Logwood is almost inodorous; of a sweetish, subastringent taste; and deep red colour. Its virtues are extracted both by water and alcohol. It is astringent and tonic, and is used in the protracted stages of diarrhœa and dysentery.

HÆMATOZO'A, pl. of *Hæmatozo'on*, (F.) *Hæmatozoaires*, (*hæmato*, and *ζωον*, 'animal,') (F.) *Vers sanguina*. Entozoa in the blood.

HÆMATURESIS, *Hæmaturia*.

HÆMATURIA, *Hæmatu'ria*, (*hæmato*, and *ουρεω*, 'I make urine,') *Voiding* of blood by urine. *Hæmorrhag'ia hæmatu'ria*, *H. ex vi'is urinari'is*, *Hæmure'sis*, *Hæm'ria*, *Mic'tio Cruen'ta* seu *Sanguin'ea*, *Mic'tus Cruen'tus* seu *Sanguin'is*, *Hæmorrh'a vi'a'rum urinari'arum*, *San'guis in Uri'na*, *Bloody urine*, (F.) *Pissement de Sang*, *Hæmatu'rie*, *Hæmure'te*. Hemorrhage from the mucous membrane of the urinary passages. Like other hemorrhages, it may be *active* or *passive*. It may proceed from the kidneys, *Nephreorrhag'ia*, bladder or urethra. The essential symptoms are:—blood, evacuated by the urethra; preceded by pain in the region of the bladder or kidneys, and accompanied by faintness. Whencesoever it proceeds, hæmaturia is usually ascribed to exhalation; rupture of vessels not being thought common in the mucous membranes. *Active hæmaturia* requires general or local blood-letting; diluent and cooling drinks; absolute rest; and the horizontal posture. The chronic kind is more troublesome. It requires acidulated or aluminous drinks; chalybeates and tonics in general. When hæmaturia is excessive, cold injections may be thrown into the rectum or into the vagina of women, and topical applications be made to the perinæum.

HÆMATURIA CYSTICA, *Cystorrhagia*—h. Ejaculatoria, *Spermatoecystidorrhagia*—h. Seminalis, *Spermatoecystidorrhagia*—h. Stillatitia, *Urethrorrhagia*.

HÆMAXIS, (*ἀμασσω*, 'to let blood,') *Blood-letting*.

HÆMENCEPHALUS, (*hæma*, and *εγκεφαλος*, 'encephalon,') *Apoplexy*.

HÆMENDOCAR'DIUM, (F.) *Hæmendocar'de*, (*hæma*, and 'endocardium,') *Inflammation* of the lining membrane of the heart.

HÆMERYTHRUM, (*hæma*, and *ερυθρός*, 'red,') *Hæmatin*.

HÆ'MIC, *Hemic*, *Hæ'micus*, (from *hæma*,) *Relating* or *appertaining* to blood; as a "hæmic disease,"—a blood disease.

HÆMIDROSIS, (*hæma*, and *ἰδρωσις*, 'sweating,') see *Diaporesis*.

HÆMIN CRYSTALS, see *Hæmatoidin*.

HÆMISCHESIS, *Hæmischesis*.

HÆMITIS, *Hæmit'is*, (*hæma*, and *itis*,) *Inflammation* of the blood. (F.) *Hémite*. The alteration of the blood that occurs in inflammatory diseases. (Piorry.) *Inflammatory blood*, *Phlogœ'mia*.

HÆMOARTHRITIS, *Rheumatism* (acute).

HÆMOCARDORRHAG'IA, *Hæmocar'dium*, *Apoplexia cor'dis*, *Ap'oplexy of the heart*, (F.) *Hémorrhagie ou Apoplexie du Cœur*, *Hémicar'de*, (*hæma*, *καρδία*, 'the heart,' and *ῥήγνυμι*, 'I break forth,') *Effusion* of blood into the substance of the heart.

HÆMOCARDIUM, (*hæma*, and *καρδία*, 'heart,')

Hæmocardiorrhagia.

HÆMOCERCHNOS. This term has received two acceptations, owing to the different senses in

which *κερως* is employed; signifying, sometimes, *hissing*; at others, *dry*. Consequently, the Hellenists have translated the compound word, at times, by spitting of blood, with hissing in the throat; at others, by evacuation of dry matters.

HÆMOCRYPHIA, (*hæma*, and *κρυπτεν*, 'to conceal,') *Hæmatischesia*.

HÆMODES, (*αἱμῶδης*, from *hæma*, and *odes*), Sanguine.

HÆMODIA, *Hæmodias'mus*, *Hæmo'dia*, *Hebētū* seu *Stū'por Den'tium*, (*ἀιμῶδω*, (from *hæma*, 'blood,' and *ὀδῶς*, 'tooth,') 'I have pain in the teeth,') Great sensibility of the teeth, and especially *Agacement* or the setting on edge of those bodies by acid or acerb substances. It is also called *Odontia Stuporis*, *Odont'gia hæmo'dia*, *Do'tor den'tium d' strido'rē*, *Cataplex'ia*, *Odontamblyg'mus*, *Odonthyperæsthes'is*, *Toothedge*.

HÆMODIASMUS, *Hæmodia*.

HÆMODROMOMETER, *Hæmadromom'eter*, (*hæma*, *δρομος*, 'course,' and *μετρον*, 'a measure,') An instrument for measuring the rate of the blood's movement. It consists of a glass tube 52 inches long, containing water, bent into the form of a hairpin, which is substituted for a segment of the bloodvessel, the velocity of the blood in which it is required to measure. The column of blood, which comes from the heart, pushes the column of water before it, and thus enables the velocity of the blood to be estimated.

HÆMODYNAMOMETER, *Hæmadynamometer*.

HÆMOGASTRIC, *Hæmagastrie*.

HÆMOIDES, (*hæma*, and *εἶδος*, 'resemblance,') Sanguine.

HÆMOMETER, *Hæmadynamometer*.

HÆMOMETRECTASIA, (*hæma*, *μητρα*, 'uterus,' and *εκτασις*, 'dilatation,') Dilatation of the womb from blood.

HÆMOPATHIA, (*hæma*, and *πάθος*, 'affection,') *Hæmatonosis*.

HÆMOPERICARDIUM, *Pericard'itis exsudato'ria*, (F.) *Hémopéricarde*, (*hæmo*, and *pericardium*.) Effusion of blood in the pericardium.

HÆMOPERITONÆUM, (*hæmo*, and *περιτοναϊον*, 'peritoneum,') Effusion of blood into the peritoneum.

HÆMOPEX'ILÆ, (G.) *Hæmopexien*, (*hæmo*, and *πηξις*, 'coagulation,') An order of diseases in which there is increased coagulability of the blood.—*Hyperinosis sanguinis*, of Fuchs.

HÆMOPHEUM, *Hæmaphein*.

HÆMOPHILIA, *Hematophilia*.

HÆMOPHOBIA, *Hæmatophobia*.

HÆMOPHOBOL, see *Hæmatophobia*.

HÆMOPHTHAL'MIA, *Hæmatophthal'mia*, *Hæmatop'ia*, (*hæma*, and *ὀφθαλμος*, 'eye,') *Hæmalopia*, *Hæmalop'ia*, *Hæmalops*, *Hype'nia* seu *Apoplex'ia oculi*, (F.) *Hémophthalmie*. Effusion of blood into the eye. When the extravasation is external, it is called *Hæmophthal'mia exter'na*, *Hypophag'ma* and *Hæmalops externus*; when internal, *Hæmophthal'mia inter'na*, *Hydrophthal'mus cruentus*, *Hypoch'yris hæmato'des*, *Hæmatom'ma*, and *Hæmalops inter'na*. A bloodshot eye.

HÆMOPLANIA, *Hæmatoplania*—h. *Menstrualis*, *Menstruation*, vicarious.

HÆMOPLETHORA, *Plethora*.

HÆMOPROCTIA, (*hæma*, and *πρωκτος*, 'anus,') *Hæmorrhois*.

HÆMOPTOË, *Hæmoptysis*—h. *Laryngea* et *trachealis*, *Tracheorrhagia*.

HÆMOPTOÏC, *Hæmopto'icus*, (F.) *Hémoptoïque*, *Hémoptique*, *Hæmoptysique*. Relating or appertaining to hæmoptysis. One affected with hæmoptysis.

HÆMOPOTOSIS, *Hæmoptysis*.

HÆMOP'TYSIS, (*hæmo*, and *πτωω*, 'I spit,') *Hæmorrho'ia*, *Hæmop'tysis*, *Hæmap'tysis*, *Hæmoptys'ia*, *Hæmatop'tysis*, *Emp'toë*, *Emop'toë*, *Emp-*

to'ica pas'sio, *Hæmoptys'mus*, *Hæmopto'sis*, *Spu'tum san'guinis* seu *cruentum*, *San'guinis flu'or pulmo'nium*, *Emopto'ica pas'sio*, *Hæmorrho'ia pulmo'nalis*, *Cruent'a expu'tio*, *Hæmorrag'h'ia pulmo'nis* seu *bron'chica*, *Pas'sio hæmopto'ica*, *Rejec'tio san'guinis d' pulmo'nibus*, *Expectora'tio san'guinis*, *Pneumorrhag'ia*, *Pneumorrhag'h'ia*, *Pneumorrhæ'a*, *Branchorrhag'h'ia*, *Emp'tysis*, *Hæmoptoë*, *Spitting of blood*, (F.) *Hémoptysie*, *Crachement de sang*, *Expectoration de sang*, *Angiarrhémie*, *Angiarrhagie*, *Hémangiarrhagie*, of *Piorry*. Hemorrhage from the mucous membrane of the lungs; characterized by the expectoration of more or less florid and frothy blood. It is generally preceded by cough; dyspnoea; sense of heat in the chest, &c. It is important to discriminate between hæmoptysis produced by some accidental cause acting irregularly or periodically on the lungs; and that which is, as it were, constitutional, and dependent on organic affection of the lungs or heart, or some faulty conformation of the chest. These two varieties differ as much in their prognosis and method of treatment as in their causes. Constitutional hæmoptysis is a serious disease, almost always announcing phthisis pulmonalis. The accidental variety is chiefly dangerous by frequent recurrence, and consequent great loss of blood.

The general causes of hæmoptysis are the same as those of other kinds of hemorrhage. It has, besides, particular causes; such as too great exercise of the lungs; loud speaking; playing on wind instruments, breathing acrid vapours, &c. It usually occurs between puberty and the age of 35. A sudden and terrific kind of hæmoptysis is sometimes met with; consisting in a great afflux of blood to the lungs. This has been called *Pulmonary Apoplexy*, *Apoplex'ia pulmo'nalis* seu *pulmonum*, *Pneumorrhag'ia*, *Infarctus hæmorrag'icus pulmonum*, I. *hæmopto'icus Lénec'ii*, (F.) *Apoplexie pulmonaire*, *Hémoptysie foudroyante*, *Hémorrhagie interstitielle du Poupon*. Infiltration of blood into the air-cells may occur without any spitting of blood. Any extravasation of blood into the tissue of the lungs has been termed by M. Hendrin *pneumohémorrhagie*.

Physical signs. Percussion may not always aid us in hæmoptysis, but generally a circumscribed dullness will be perceived. The inspiratory murmur, on auscultation, is feeble or absent, locally; and is replaced by bronchial respiration and bronchophony. A fine liquid crepitus is detected around the affected part; and in the larger tubes, near the spine, a liquid bubbling rhonchus is usually heard. The value of these signs is determined by the nature of the expectoration. The treatment of hæmoptysis must be like that of internal hemorrhage in general.

HÆMOP'TYSIS INTERNA, *Hæmatothorax*—h. *Laryngea* et *trachealis*, *Tracheorrhagia*—h. *Phthisis*, *Phthisis pulmonalis*.

HÆMOTYSMUS, *Hæmoptysis*.

HÆMORMESIS, (*hæma*, and *ορμωω*, 'to drive on,') Congestion, Hyperæmia.

HÆMORRHACHIS, (*hæma*, and *ῥαχis*, 'the spine,') *Apoplexia myelica*.

HÆMORRHAGE, *Hæmorrhagia*.

HÆMORRHAG'IA, (*hæma*, and *ῥήγνυμι*, 'I break forth,') *San'guinis profu'vium copio'sum*, *Sanguifluxus*, *Hæmatoch'yris*, *Aimorrhæ'a*, *Hæmorrho'ia*, *Hæmatorrhag'h'ia*, *Hæmatismus*, *Hæmatorrhæ'a*, *Hæmatorrhæ'ia*, *Hæmatorrhæ'ia*, *Proflu'vium* seu *Prorup'tio seu Erup'tio seu Profu'sio san'guinis*, *Hemorrhage*, *Hæmorrhage*, *Bleeding*, *Loss of blood*, *Rupturing*, *bursting*, or *breaking of a bloodvessel*, (Prov.) *Floating*, (F.) *Hémorrhagie* ou *Hémorrhagie*, *Perte de sang*, *Flux de sang*. Any discharge of blood from vessels destined to contain it; with or without rupture of their coats. Hemorrhages may be spontaneous or

Traumatic: the first belong to the domain of medicine, the latter to that of surgery. They may, also, be *internal or external*; *general*—as in scurvy—or *local*. The hemorrhages by *exhalation*—those which chiefly interest the physician—have been classed, by Pinel, as follows:—1. *Hemorrhage of the Mucous Membranes*; Epistaxis, Hæmoptysis, Hæmatemesis, Hæmorrhoids, Hæmaturia, Uterine Hemorrhage. 2. *Hemorrhage of the Tissues*; Cutaneous, Cellular, Serous, Synovial.

Hemorrhages have generally been distinguished into *active and passive*: in other words into those dependent upon augmentation of the organic actions, and those dependent upon debility. According to Broussais, no spontaneous hemorrhage is passive; all are active,—that is, produced by increased action and excess of irritation of the blood-vessels: they may occur *with* debility, but not *from* debility. He calls those only *passive hemorrhages*, which are owing to an external lesion of the vessels. Hemorrhages have been, by some, divided into *constitutional*, or those depending on original conformation;—*accidental*, or those produced by some adventitious cause; *supplementary*, or those which succeed others; *symptomatic*, *critical*, &c. A *secondary hemorrhage* is one that occurs some time after wounds or operations.

When hemorrhage takes place into any tissue, or is *interstitial*, it receives the name, with many, of *apoplexy*.

Active Hemorrhage, Hemorrhag'ia acti'va seu arterio'sa, Causa hæmorrhag'icæ, Angiorrhag'ia, occurs chiefly in the young and plethoric. Good living; the use of fermented liquor, excessive exercise, or too sedentary a life, may perhaps be ranked as predisponent causes. It is commonly preceded by heaviness and pulsation in the part,—owing to the afflux of blood and consequent hyperæmia,—and by coldness of the extremities. The blood, evacuated, is generally of a florid red. In such active hemorrhages, the great indications of treatment will be, to diminish plethora where it exists, and to lessen the heart's action. Bleeding, purgatives, and cold, with dry diet, will be the chief agents.

Passive Hemorrhage, Hæmorrhag'ia passi'va seu veno'sa, Profusio, P. hæmorrhag'ica, occurs in those of weak constitution; or who have been debilitated by protracted disease, poor diet, long watching, excessive evacuations, &c.

The direct causes may be; previous active hemorrhage; scorbutus, or any thing capable of inducing atony or asthenic hyperæmia of the small vessels. These hemorrhages are not preceded by excitement or by any signs of local determination. They are usually accompanied by paleness of the countenance; feeble pulse; fainting, &c. The indications of treatment will be:—to restore the action of the small vessels and the general tone of the system: hence the utility of styptics and cold externally; and of tonics and astringents, creasote, mineral acids, &c., internally. Hemorrhage also occurs from mechanical hyperæmia, as when hæmoptysis is produced by tubercles in the lungs; apoplexy by softening of the encephalic neurme; hæmatemesis by disease of some of the solid viscera of the abdomen, &c.

In *Traumatic Hemorrhages*, or those which are the consequences of wounds of arterial or venous trunks, the blood is of a florid red colour, and issues by jets and pulses, if it proceed from an artery; whilst it is of a deeper red, issues slowly and by a continuous flow, if from a vein. If the capillary vessels be alone divided, the blood is merely effused at the surface of the wound. Of the means used for arresting these traumatic hemorrhages, some act mechanically as *absorbents, ligature, and compression*; others chymically, as *fire, caustic, creasote, astringents*, &c.

HEMORRHAGIA ACTIVA NARIUM, Epistaxis—

h. Bronchica, Hæmoptysis—h. Cerebri, Apoplexy—h. per Cutem, Sudor cruentus—h. per Diæresin, see Diæresis—h. per Diapedesin, Diapedesum—h. Faucium, Stomatorrhagia—h. Gingivarum, Ulorrhagia—h. Hæmatemesis, Hæmatemesis—h. Hæmaturia, Hæmaturia—h. Hæmoptysis, Hæmoptysis—h. Hepaticæ, Hepathæmorrhagia—h. Hepatis, Hepatorrhagia—h. Intestinorum, Hæmatocœzia—h. Mucosa, see Hæmorrhoids—h. Nabothi, see Parturition—h. Narinae, Epistaxis—h. Narium, Epistaxis—h. Oris, Stomatorrhagia—h. Penis, Stimatosis—h. Proctica, Hæmorrhoids—h. Pulmonis, Hæmoptysis—h. Renum, Nephorrhagia—h. Universalis, Purpura hæmorrhagica—h. Uterina, Metrorrhagia—h. Ventriculi, Hæmatemesis—h. Vesicæ, Cystorrhagia.

HÆMORRHAGICUS, Hæmorrhagic.

HÆMORRHINIA, (hæma, and ῥιν, 'the nose,') Epistaxis.

HÆMORRHINORRHAGIA, (hæma, ῥιν, 'the nose,' and ῥύωμι, 'to break forth,') Epistaxis.

HÆMORRHŒA, (hæmo, and ῥέω, 'I flow,') Hæmorrhag'ia, Loss of blood, (F.) Hémorrhée. Some writers have proposed to restrict this name to passive hemorrhages.

HÆMORRHŒA PETECHIALIS, Purpura hæmorrhagica—h. Pulmonalis, Hæmoptysis—h. Uterina, Metrorrhagia—h. Vasorum hæmorrhoidalium, Hæmorrhoids—h. Ventriculi, Hæmatemesis—h. Viarum urinarium, Hæmaturia.

HÆMORRHOID'AL, Hemorrhoid'al, Hæmorrhoid'al'is, Hæmorrhoid'eus. Relating to hemorrhoids; as *hemorrhoidal flux*, *hemorrhoidal tumours*, &c.

HEMORRHOIDAL ARTERIES have been distinguished into *superior, middle, and inferior*. The first is the termination of the inferior mesenteric artery, which assumes the name *superior hemorrhoidal*, when it reaches the upper and posterior part of the rectum. The *middle hemorrhoidal* is furnished by the hypogastric or internal pudic. It ramifies on the inferior and anterior part of the rectum. The *inferior hemorrhoidal* arteries are branches of the internal pudic, furnished to the inferior part of the rectum and to the muscles of the anus.

HEMORRHOIDAL NERVES. These emanate from the sciatic and hypogastric plexuses; and cover the rectum with their numerous filaments.

HEMORRHOIDAL VEINS follow the same distribution, and generally empty themselves into the lesser mesenteric. Some of them assist in the formation of the hypogastric vein.

HEMORRHOIDAL VESSELS, *Vasa sed'al'ia*, are those vessels which are distributed to the rectum—the seat of hemorrhoids.

HÆMORRHODIALES NODI, Hæmorrhoids.

HÆMORRHOIDES, see Hæmorrhoids—h. Cæcæ, see Hæmorrhoids—h. Fluentes, Hæmorrhoids—h. Furentes, see Hæmorrhoids—h. Maris-cosæ, Hæmorrhoids—h. non Fluentes, Hæmorrhoids—h. Oris, Stomatorrhagia.

HÆMORRHŒDROSIS, (hemorrhœis, and ῥέωσις, 'sweating,') Sweating of hemorrhoidal blood. (?) (Bock.)

HÆMORRHOÏS, Aïmor'rhoïs, Asclepias'mus, (hæmo, and ῥέω, 'I flow,') Aïmor'rhoïs, Proct'ica Maris'ca, Maris'ca, Proct'al'gia Hæmorrhoid'al'is, Morbus Hæmorrhoidalis, Hemorrhoids, Piles, Em'eroïds, Em'eroïds, Em'eraudes, (F.) Hémorrhoides. The essential symptoms of this affection are:—Livid and painful tubercles or excrescences, (Hemorrhoid'al Tumours,) usually attended with a discharge of mucus or blood, (Hemorrhoidal flux, Hemorrhag'ia proctica, Proctor'rhoïs, Hemoproctica, Hemorrhœa vaso'rum hæmorrhoid'alium, (F.) Hémoproctie, Hémoproctie. The most common causes of piles are a sedentary life; accumulation of feces in the rectum; violent efforts at stool; pregnancy, &c. The precursory

symptoms are:—pains in the loins; stupor of the lower limbs; and uneasiness in the abdomen and rectum, with more or less gastric, cerebral, and indeed general disorder;—constituting the *Diathæsis Hæmorrhoidalis*, *Molitus* seu *Turba Hæmorrhoidalis*, and *Molimen Hæmorrhoidale*, of most of the writers of Continental Europe. To these symptoms follow one or more round, smooth, renitent, painful, pulsating, and erectile tumours, around the margin of the anus, or within the anus; some pouring out blood occasionally. After having remained, for a time, tense and painful, they gradually shrink and disappear. The chief symptoms, occasioned by hæmorrhoidal tumours, when much inflamed, are;—constant pain, liable, however, to exacerbations, and to become augmented by the least pressure, or by the passage of the fæces, and obliging the patient to preserve the horizontal posture.

Hæmorrhoids have generally been distinguished into—*Hæmorrhoides Fluentes* seu *Mariæcæ*, *Proctica Mariæca cruenta*, *Bleeding* or *Open Piles*; and into *H. non fluentes*, *Proctica Mariæca cæca*, *Hæmorrhoides cæcæ seu fœnites*, *Hæmorrhoidales* *no di*, *Shut* or *blind piles*. They have, also, been divided into *internal* or *occuli*, and *external*, according to their situation; and into *accidental* or *constitutional*.

Hæmorrhoidal Tumours are extremely troublesome, by their disposition to frequent recurrence; and they are apt to induce fistula; otherwise, they are devoid of danger. When anatomically examined, they are found not to consist in a varicose dilatation of the veins of the rectum; but to be formed of a very close, spongy, texture; similar to that which surrounds the orifice of the vagina; and to be erectile, like it. They are surrounded by a delicate membrane, and have no internal cavity. The treatment, in mild cases of hæmorrhoidal tumours, is simple. Rest; the horizontal posture; the use of mild laxatives, as sulphur, castor oil, and emollient glysters, will be sufficient. If they be much inflamed, leeches may be applied; and warm cataplasms or cold lotions, according to circumstances, be prescribed, with abstinence, and cooling drinks. Afterwards, an ointment, composed of powdered galls and opium, may afford relief. It is in the relaxed kind, that such ointment, and the internal use of *Ward's Paste*, can alone be expected to afford much benefit. If, after repeated attacks, the tumours remain hard and painful, and threaten fistula, they may be removed.

By *Hæmorrhoidal Flux*, *Fluxus hæmorrhoidalis*, *Proctorrhag'ia*, is meant the hæmorrhage which takes place from the rectum, owing to hæmorrhoids. It is a common affection. The quantity of blood discharged is various; at times, it is very trifling; at others, sufficient to induce great debility, and even death. It is announced and accompanied by the same symptoms as precede and attend hæmorrhoidal tumours. Like other hæmorrhages, it may be *active* or *passive*, *accidental* or *constitutional*. The prognosis is rarely unfavourable. The affection may, almost always, be relieved by properly adapted means. These resemble such as are necessary in hæmorrhages in general. Perfect quietude, mental and corporeal; light diet; cooling drinks; bleeding, if the symptoms indicate it; astringents, if the disease be protracted and passive, such as the *Tinctura Ferri Chloridi*; aspersions of cold water on the anus; astringent injections; plugging, and compression. Such will be the principal remedial agents. When the hæmorrhage has become habitual, or is vicarious, some caution may be required in checking it; and, if inconvenience arise from a sudden suppression, its return may be solicited by the semicupium, sitting over warm water, aloetic purgatives, glysters, irritating sup-

positories, &c.; or leeches may be applied to the anus.

To the internal bleeding pile, a soft, red, strawberrylike elevation of the mucous membrane, Dr. Houston, of Dublin, gives the name *vascular tumour*. For its removal he recommends the application of nitric acid, so as to produce sloughing of its surface.

Some authors have described a species of *Leucorrhœa Analis*, or whitish discharge from the anus, which often attends ordinary hæmorrhoids. This they have called *Proctica mariæca mucosa*, *Hæmor'rhois al'ba*, *Hæmorrhag'ia mucosa*, *Leucor'rhois*. It requires no special mention.

The term *Hæmorrhoid* is applied at times to varicose veins developed elsewhere, as at the orifice of the uterus. See *Metarrhœoides*.

HÆMORRHOIS CRUENTA, see *Hæmorrhoids*—h. ab Exaniâ, *Proctocele*—h. *Procedens*, *Proctocele*.

HÆMORRHOIS/CHESIS, (*hæmorrhôis*, *ῥέω*, 'I flow,' and *ἰσχω*, 'I restrain.') *Reten'tio fluxus hæmorrhoidalis*. Suppression or retention of the hæmorrhoidal flux.

HÆMORRHOPHESIS, *Absorptio sanguinis*, (*hæma*, and *ῥοφῶ*, 'I sip up.') Absorption of blood.

HÆMORRHOPHILIA, (*hæma*, *ῥέω*, 'to flow,' and *φιλέω*, 'to love.') *Hæmatophilia*.

HÆMORRHOSCOPIA, (*hæma*, *ῥέω*, 'to flow,' and *σκοπέω*, 'to examine.') *Hæmatoscopy*.

HÆMORRHYSIS, *Hæmorrhagia*.

HÆMOSCHEOCELE, *Hæmatoscheocele*.

HÆMOSCOPIA, *Hæmatoscopy*.

HÆMOSPASIA, see *Hæmospastic*.

HÆMOSPASTIC, (F.) *Hémospastique*, (*hæma*, and *σπᾶω*, 'I draw.') An agent which draws or attracts blood to a part; as a cupping-glass. The operation is termed *Hæmospasia*, (F.) *Hémospasie*. It is generally applied to a process by which the air is exhausted over a considerable surface, as over one or more of the extremities, by an appropriate pneumatic apparatus.

HÆMOSTASIA, *Hæmostasis*, *Epidæsis*, *Sanguinis stagnatio*, (F.) *Hémostasie*, (*hæma*, and *σταίς*, 'stagnation.') Stagnation of blood. This name has also been given to any operation, the object of which is to arrest the flow of blood, as by compression or ligation of the great vessels, (F.) *Perstriction*.

HÆMOSTATIC, **PAGLIA'RIS**, *Pagliari's Styptic*, *A'qua Pagliari*, (F.) *Eau de Pagliari*. A styptic liquor, said to be of great power, discovered by M. Pagliari, a pharmacien at Rome. It is composed as follows.—Eight ounces of tincture of benzoin, one pound of alum, and ten pounds of water, are boiled together for six hours in a glazed earthen vessel, the vaporized water being constantly replaced by hot water, so as not to interrupt the ebullition, and the resinous mass being kept stirred round. The fluid is then filtered, and kept in stoppered bottles. A drop of the fluid, poured into a glass containing human blood, produces an instantaneous magma.

HÆMOSTATICS, *Hæmostaticæ*.

HÆMOTACHOM'ETRY, *Hæmotachomet'ria*, (*hæma*, *ταχος*, 'swiftness,' and *μετρέω*, 'I measure.') The measurement of the velocity of the blood. An instrument, devised for this purpose by Viërrordt, is called by him *Hæmotachom'eter*.

HÆMOTELANGIOSIS, *Telangiectasia*.

HÆMOTEXIA, *Hæmatoclysis*.

HÆMOTEXIS, (*hæma*, and *τείξις*, 'solution,') *Hæmatoclysis*.

HÆMOTHORAX, *Hæmothorax*.

HÆMOTOXIC, *Toxæmic*.

HÆMOTROPHY or **HÆMOT'ROPHY**, *Hæmotroph'ia*, (*hæma*, and *τροφή*, 'nourishment.') Excess of sanguineous nourishment.

HÆMOTUS, (*hæma*, and *οὖς*, gen. *ωτός*, 'the ear.') Effusion of blood into the ear.

HÆMURESIS. Hæmaturia.
HÆMURIA. Hæmaturia.
HÆMYDOR, (*hæma*, and *ὕδωρ*, 'water.') Serum of the blood.
HÆMYDRIA. Hydroæmia.
HEREDITARIUS, (*hæres*, gen. *hære'dis*, 'an heir.') Hereditary.
HESITATIO, (*hæsitare*, *hæsitatum*, 'to stick fast.') Balbuties.
HÆVEA GUIANENSIS, see *Caoutchouc*.
HAG (Prov.), Abdomen.
HAGARD, Haggard.
HAGBERRY, *Prunus padus*.
HAGENIA ABYSSINICA, *Brayera anthelmintica*, *Bank'sia Abyssinica*, (after Prof. K. G. Hagen, of Königsberg.) An Abyssinian tree; Order, Rosaceæ; *Sexual System*, Icosandria Digynia, which the natives plant round their habitations as an ornament. The infusion or decoction of its flowers, *Flores cosso*, is employed by them as an anthelmintic, especially in cases of tape-worm. It is called there *Cus'so* or *Cous'so*, *Kos'so*, *Kous'so*, *Koo'so*, *Kus'so*, *Kwo'so*, *Cotz*, *Cabutz*, and *Hubbi*.
The flowers and unripe fruit—*Brayera, Kosso*—are official in the Pharmacopœia of the United States. An active resinous principle has been obtained from the flowers, to which the names *Tœnin* and *Koussin* have been given. The dose of this is from gr. xx to xl.
HAGGARD, ([G.] *hagard*, 'lean,' [F.] *Hagard*, The French use the term *Air hagard*, *Œil hagard*, 'Haggard air,' 'Haggard eye,' for a physiognomy in which there is at once an expression of madness and terror.
HAGIOSPERMUM, (*ἅγιος*, 'holy,' and *σπέρμα*, 'seed.') *Artemisia santonica*.
HAGIOXYLUM, (*ἅγιος*, 'holy,' and *ξύλον*, 'wood.') *Guaiaecum*.
HAHNEMANNISM, (after Dr. S. C. F. Hahnemann, the founder,) Homeopathy.
HAIL, (Sax. *hægel*.) Chalaza.
HAIR, (Sax. *hær*.) *Cri'nis*. *Pi'lus*, *Thrix*, *Pile*, (F.) *Poil*. A conical, corneous substance, the free portion or *shaft*, *scal'pus*, of which issues to a greater or less distance from the skin, to the tissue of which it adheres by a *bulb* or *button*, *bulbus pi'li*, seated in a *hair follicle*—*follic'ulus pili*—made by an inversion of the integument, the epidermis of which forms a "root sheath," *vagi'na pili*. At the base of the hair follicle there is a small papilla, well supplied with bloodvessels and nerves, *Papil'la pili*; at times called, but improperly, *Pulpa* seu *Blaste'na pili*. The hair receives various names in different parts—as *Beard*, *Cilia*, *Eye-browe*, *Hair of the head* (*Capilli*), &c. See *Capillus*.
HAIR DYE, TWIGGS'S. A preparation much used to give colour to the hair. It may be formed of *precipitated sulphur* and *acetate of lead*, each $\mathfrak{z}\mathfrak{i}$; *rose water*, $\mathfrak{f}\mathfrak{z}\mathfrak{i}\mathfrak{v}$; triturated together in a mortar. It should be applied twice a day until the desired effect is produced.
HAIR, FALLING OFF OF THE, Alopecia.
HAIR FOLLICLE, see *Hair*.
HAIR, MATTED, Plica—h. Plaited, Plica—h. Trichomatose, Plica.
HAIR WORM, *Seta equina*, see *Ectozaa*—h. Worm, muscular, *Dracunculus*.
HAIRY, Pileous. Also, covered with hair.
HAKE, see *Oleum jecoris aselli*.
HAL, (*ἅλς*, *ἅλος*.) Salt.
HALATION, *Hala'tium*, (dim. of *hal*.) A pungent remedy with the ancients, which contained salts.
HALCHEMIA. The alchemists so called the art of fusing salts; (*hal*, and *χεω*, 'I pour out.')
HALCYON, (*ἄλκυων*.) Alcyon.
HALCYONIUM ROTUNDUM, (*ἄλκυονεον*.) *Pila marina*.

HALEINE, (from *halare*, 'to emit a vapour,') Breath—h. *Courte*, Dyspnœa.
HALELÆ'ON, *Halelæ'um*, (*hal*, and *λαίον*, 'oil.') *Oleum Sæle mixtum*. A mixture of oil and salt, for removing swellings of the joints.—
HALÉTERATION, see *Alteration*.
HALFCASTE, *Halfcast*, see *Caste*. A term applied in India to the offspring of a Hindoo and a European: since, extended to the offspring of mixed races. The subjoined list from Tschudi shows the parentage of the different varieties of halfcastes, and also the proper designations of the latter, as observed in South America.

PARENTS.	CHILDREN.
White Father and Negro Mother,	Mulatto.
White Father and Indian Mother,	Meztizo.
Indian Father and Negro Mother,	Chino.
White Father and Mulatta Mother,	Cuarteron.
	{ Creole, (only distinguished from the white by a pale brownish complexion.
	{ Chino-blanco.
White Father and Meztiza Mother,	{ Quintero.
	{ White.
White Father and China Mother....	{ Zambo-negro.
White Father and Cuarterona Mother	{ Mulatto oscuro.
White Father and Quintera Mother	{ Zambo chino.
Negro Father and Mulatta Mother,	{ Zambo-negro, (perfectly black.)
Negro Father and Meztiza Mother,	{ Mulatto, (rather dark.)
Negro Father and China Mother,	{ China-oscur.
Negro Father and Zamba Mother,	{ Meztizo-claro, (frequently very beautiful.)
Negro Father and Cuarterona or Quintera Mother.....	{ Chino-cholo.
Indian Father and Mulatta Mother	{ Zambo-claro.
Indian Father and Meztiza Mother,	{ Indian, (with rather short frizzy hair.)
Indian Father and China Mother.....	{ Meztizo, (rather brown.)
Indian Father and Zamba Mother	{ Zambó, (a miserable race.)
Mulatto Father and Meztiza Mother.....	{ Chino, (of rather clear complexion.)
Mulatto Father and China Mother,	{ Chino, (rather dark.)

For the designation of different admixtures in the United States, see *Mulatto*.

HALICA, *Alica*.
HALICACABUM, (*ἅλς*, 'the sea,' and *κακὰς*, 'nightshade.') *Physalis*.

HALICES, *Pandiculusa*.

HALIDRYS VESICULOSA (*ἅλς*, 'the sea,' and *δρῦς*, 'the oak,') *Fucus vesiculosus*.

HALINA'TRUM, *Halini'trum*, *Halon'i'trum*: (*ἅλς*, 'salt,' and *natrum* or *natron*.) A name given by the ancients to subcarbonate of soda containing a little subcarbonate of ammoniac, which is found, ready-formed, on the plaster of damp walls, in places inhabited by man or animals. Also, *Potassæ nitras*.

HALINITRUM, see *Halinatrum*.

HALINUS, (*ἅλιος*, from *ἅλς*, 'salt,') *Saline*.

HALITUOUS, or **HALIT'UOUS,** *Halituo'us*, (*halitus*, 'vapour,') (F.) *Halitueux*. The skin is said to be *halitueuse* (F.), when covered with a gentle moisture. The vapour, exhaled in all the cavities of the body, so long as the blood is warm, is called *Hal'itus*. The odorous vapour, exhaled by the blood itself whilst warm, is called *Hal'itus San'guinis*.

HALITUS, ('vapour,') Breath, see *Halituous*—h. *Oris Fetidus*, Breath, offensive—h. *Sanguinis*, *Gaz sanguinis*. See *Halituous*.

HALL, MINERAL WATERS OF. The springs of Hall, in Upper Austria, contain iodide.

HALL'S, (DR. MARSHALL,) READY METHOD, see *Ready Method*.

HALLUCINATIO, (*hallucinari*, [or *hallucinari*] *hallucinatus*, 'to be deceived.') Hallucination—h. Hypochondriasis, Hypochondriasis—h. Vertigo, Vertigo.

HALLUCINATION, *Hallucina'tio*, *Hallucina'tio*, *Falla'cia*, *Illda'io*, *I. sensûs*, *Soco's'dia*, *Alu'sia*, *Alucinatio*, *Parora'sis*, *Somnia'tio in statu rig'i'ti*, *Idol'um*, *Waking dream*, *Phan'tasm*. A morbid error, in one or more of the senses. Perception of objects which do not in fact exert any impression on the external senses. Hallucination or *delusion* almost always, if not always, depends on disorder of the brain, but is not an index of insanity, unless the patient believes in the existence of the subject of the hallucination.

HALLUS, Pollex pedis.

HALLUX, Pollex pedis.

HALME, ('*αλμη*, 'saltwater.') Muria.

HALMYRIS, ('*αλμυρις*, 'saltwater.') See Muria.

HALMYRODES, *Salungio'sus*, *Salsu'ginous*, ('*αλμυρος*, '*αλμη*, 'seawater,' '*αλς*, 'salt,') 'saltish,' and *odes*.) An epithet given to any affection in which the heat feels pungent under the finger of the physician.

HALMYRUS, ('*αλμυρος*.) Saline.

HALO, gen. *Halo'nis*, ('*αλως*, 'a circle,') Areola.

HA'LO SIGNA'TUS. The impression made by the ciliary processes on the anterior surface of the vitreous humour. So called from its consisting of a circle of indentations. Called by Haller, *Stri'æ ret'inæ subject'æ ligamen'to cilia'ri*; by Winslow, *Sulci cilii' res*.

HALODERÆUM, (*hal*, and *τεραον*, 'a collar,' from *αμη*, 'neck,') *Colla'rium sali'num*. A collar of salt applied to the neck, as in cases of croup.

HALOGENE, (*hal*, and *γεννω*, 'to generate,') Chlorine.

HALOIDUM OXYGENATUM, (*hal*, and *ειδος*, 'resemblance,') Potassæ murias hyperoxygenatus.

HALONITRUM, Halinatrum.

HALOPE'GÆ, (*hal*, and *πηγη*, 'a spring,') Mineral waters whose chief ingredient is common salt—chloride of sodium.

HALOS, ('*αλως*.) Areola.

HALOT'YLE, *Cul'cita sali'na*, (*hal*, and *τυλη*, 'a bolster,') A bolster or pillow of salt, recommended in croup.

HALSBANE [Sc.], (Sax. *halp*, 'the neck,' and *ban*, 'bone,') Clavicle.

HALSE, (Sax. *halp*.) Collum, Throat.

HALTE'RES, ('*αλτηρες*, from '*αλλομαι*, 'to leap,') Pieces of lead held in the hands by the ancients, to assist them in leaping.

HALLUCINATIO, Hallucination.

HALYCODES, ('*αλυκωδης*.) Saline.

HALYSIS LATA, ('*αλυσος*, 'a chain,') Bothriocephalus latus—h. Solium, Tænia solium.

HAM, (Sax. *ham*.) Poples.

HAMAME'LIS VIRGINIA'NA, ('*αμα*, 'like to,' and *μηδης*, 'an apple-tree,') *H. Virginica* seu *macrophy'lla*, *Witch hazel*, *Winter witch hazel*, *Shrapping hazelnut*, *Winter bloom*. Ord. Hamame-laceæ. The bark of this tree, which is a native of the United States, is somewhat bitter and sensibly astringent; but it has not been much used. A cataplasm of the inner rind of the bark is said to have been found efficacious in painful inflammation of the eyes.

HAMARTHRI'TIS, *Holarthrit'is*, *Catholarthrit'is*, *Arthrit'is universa'lis*, (*αμα*, 'at once,' and *arthrit'is*.) Gout in all the joints. Universal gout.

HAMCH (Prov.), Coxofemoral articulation.

HAM'MA, gen. *Hamm'atis*, (*αμμα*), *Nodus*, 'a tie,' A knot, used for retaining bandages on any part. A truss.

HAMMERING (Prov.), Balbuties.

HAMPSTEAD, MIN'ERAL WA'TERS OF These waters, situate in the neighbourhood of London, are a good chalybeate.

HAMSTRING, see Hamstrings.

HAM'STRINGS, (F.) *Jure'tiera*. The strings or tendons of the ham.

The **HAMSTRING MUS'LES** are the biceps femoris, whose tendon forms the *outer hamstring*; and the semimembranosus, semitendinosus, gracilis and sartorius, whose tendons form the *inner ham string*.

'*To hamstring*' means to cut the strings or tendons of the ham.

HAMULAR PROCESS, see Hamulus.

HAM'ULUS, (dim. of *hamus*, 'a hook,') A hook or crook: *Anci's'tron*. Also, any hook-like process; as the *hamulus* or *hamular process* of the pterygoid process of the sphenoid bone, which projects from the termination of the internal pterygoid plate.

HAMULUS LAMINÆ SPIRALIS, see Lamina spiralis.

HAMUS, 'a hook,' see Hamulus.

HANCHE, Haunch.

HANCLETH (Sc.), Malleolus.

HAND, (Sax. *hanc*, *hond*.) Manus.

HANDS, DROPPED, *Hand-drop*, *Wrist-drop*. A popular term for the paralysis of the hand induced by the action of lead.

HANDWOMAN, Midwife.

HANDWRIST (Prov.), Carpus.

HANGERS, Crusta genu equinæ.

HANG'NAIL, (old Eng.) *Agnail*, (Prov.) *Ang nail*, (Sax. *angnægl*) from *ange*, 'troublesome, and *nægl*, 'a nail,' *Stepmother's blessing*, *Wirt-springs*, *Nail-spring*, *Nang-nail*, *Friend-back*, (F.) *Énvie*. A portion of epidermis, detached so as to tear the integument in the vicinity of the finger nails.

HANNEBANE, ('henbane,') Hyoscyamus.

HAPANTIS'MUS, (from *nas*, gen. *pan'tos*, or *άπας*, gen. *άπαντος*, 'all,') *Oblitera'tio comple'ta*. The matting or growing together of organic parts.

HAPHÊ, (*αφη*, 'feeling, touch,') Hence:

HAPHON'OSI, *Haphon'u'si*, (*haphê*, *άπλος*, 'simple,' and *ωσος*, 'disease,') *Morbi tactûs*. Diseases of the sense of touch.

HAPLOACNE, (*άπλος*, 'simple,' and *acne*.) Acne simplex.

HAPLOPATH'IA, (*άπλος*, 'simple,' and *παθος*, 'disease,') *Morbus simplex*. A simple or uncomplicated affection.

HAPLOTOM'IA, *Simplex sec'tio*, *Incis'io simplex*, *Opera'tio simplex*, (*άπλος*, 'simple,' and *τομη*, 'incision,') A simple incision.

HAPSIS, (*άψις*.) Touch.

HAPTODYSPHOR'IA, *Tactus dolorif'iculus*; (*άπτος*, 'touched,' and *δυσφορος*, 'difficult to be borne,') Painful to the touch.

HAPTOGEN, Haptogenous.

HAPTOGENE, Haptogenous.

HAPTOG'ENOUS, *Haptog'enus*, *Hapt'ogen*, (F.) *Haptogène*, (*άπτω*, 'I attach,' and *γεννω*, 'I engender,') Ascherson calls the pellicle, which forms around fatty matter when albumen is brought in contact with it, *Haptogen membrane*, (F.) *Membrane haptogène*.

HAPTOTICA, *Haptot'icê*, ('*άπτω*, 'to touch,') The doctrine of the phenomena of touch.

HARD, (Sax. *hearp*.) *Du'rus*, *Scl'e'ros*, (F.) *Dur*. That which offers much resistance. In anatomy, the *hard parts* are those which compose the osseous basis of the body; in other words, the skeleton. See Pulse, hard.

HARD'S FARINACEOUS FOOD, see Farina-ceous.

HARDENING, Induration.

HARDESIA, Hibernicus lapis.

HARDHACK, *Spiraea tomentosa*.

HARDS, see Falsifications, table of.

HARD TACK. Army and navy bread, baked without salt and kiln dried.

HARELIP, *Lagochil'us*, *Lagos'toma*, *Lagen'to-mum*, *La'bium Lepori'num*, *Olopho'nia Labii Lo-ba'ta*, *Lepori'num ros'trum*, (Sc.) *Hare'shawe*, (Prov.) *Hay'scale*, (F.) *Bec de Lièvre*. A fissure or perpendicular division of one or both lips. It has been so called, in consequence of the upper lip of the hare being thus divided. Harelip is generally congenital; at other times it is accidental, or produced by a wound, the edges of which have not been brought into contact, and have healed separately. It is *simple*, when there is but one division; *double*, when there are two; and *complicated*, when there is, at the same time, a division or cleft of the superior maxillary bone and of the palate—*Cleft or fissured palate*, *Wolf's jaw*, *Palatum fissum*, *Rictus lupinus*; or a projection of the teeth into the separation of the lip.

In the *Harelip operation*, there are two indications to be fulfilled. First, to pare, with the knife or scissors, the edges of the cleft, and, afterwards, to preserve them in contact; in order to cause adhesion. This last object is accomplished by means of pins, passed through the edges of the division; in other words, by the twisted suture. The projecting teeth must, of course, be previously removed. If there be separation of the palate, it will become less and less after the union of the lip, or the operation of staphylorrhaphy may be performed upon it.

HARE'S EAR, *Bupleurum rotundifolium*—h. Eye, *Lagophthalmia*.

HARESHAW (Sc.), Harelip.

HARGNE, *Hernia*—h. *Anévrysmales*, *Cirsomphalus*.

HARGNEUX, see *Hernial*.

HARICOT, *Phaseolus vulgaris*—h. *Grand de Pérou*, *Jatropha curcas*.

HARMALIA, (*ἀρμαλία*), *Aliment*.

HAR'MONY, *Harmo'nia*, *Harmonus*, (primarily from *ἄνω*, 'I adjust.') Anatomists have called *Suture by Harmony*, or simply *Harmony*, *False* or *superficial suture*, *Sutura os'sium spu'ria*, *Comp'ages os'sium per lin'eam simp'licem*, an immovable articulation, in which the depressions and eminences, presented by the bony surfaces, are but slightly marked; so that it might be presumed that the junction of the bones took place by simple apposition of their surfaces. An instance of harmony occurs in the union of the superior maxillary bones with each other.

HARMUS, (*ἄρμος*, from *ἄνω*, 'I adjust.') Articulation, *Harmony*.

HARNPAN, ([G.] *Hirn*, 'the brain.') *Cranium*.

HARRODSBURG SPRINGS, see *Kentucky*, *Mineral Waters of*.

HARROWGATE, MINERAL WATERS OF. The villages of High and Low Harrogate are situate in the centre of the county of York, near Knaresborough, twenty miles from York, and fifteen from Leeds. Here are several valuable sulphureous and chalybeate springs. The *sulphureous springs* contain chloride of sodium, chloride of calcium, chloride of magnesium, bicarbonate of soda, sulphohydric acid, carbonic acid, carburetted hydrogen, and azote. The *chalybeate springs* contain protoxide of iron, chloride of sodium, sulphate of soda, chloride of calcium, chloride of magnesium, carbonic acid, azote, and oxygen. For the former, Harrogate is celebrated and frequented.

HARROWGATE SALTS, ARTIFICIAL, are much employed, and not unfrequently by those who drink the genuine water, for the purpose of increasing its aperient power. They may be made as follows:—*Potass. Sulph. cum Sulph.* $\mathfrak{z}\mathfrak{v}$; *Potass. bitart.* $\mathfrak{z}\mathfrak{j}$; *Magnes. Sulph. in pulv.* $\mathfrak{z}\mathfrak{v}\mathfrak{j}$; *M.*

The usual dose is a teaspoonful, in a small tumblér of tepid water, early in the morning.

HARROWGATE WATER, ARTIFICIAL, may be formed of *common salt*, $\mathfrak{z}\mathfrak{v}$; *water*, *Oij*; impregnated with the gas from *sulphuret of potassa* and *sulphuric acid*, $\mathfrak{a}\mathfrak{a}$ $\mathfrak{z}\mathfrak{iv}$. The following form has also been recommended. *R. Potass. Sulphat. cum sulph.* (Ph. Ed.) $\mathfrak{z}\mathfrak{j}$; *Potass. bitart.* $\mathfrak{z}\mathfrak{ss}$; *Magnes. sulphat.* $\mathfrak{z}\mathfrak{v}\mathfrak{j}$; *Aquæ destillat.* *Oij*. One-half to be taken for a dose.

HARTFELL, MINERAL WATERS OF. Hartfell Spa is about five miles from Moffat, in Scotland. The water is a chalybeate, and is much used.

HARTSHORN, *Cervus*, *Cornu cervi*, *Liquor cornu cervi*—h. and Oil, *Linimentum Ammoniac carbonatis*—h. Red, *Spiritus lavandulæ compositus*.

HARTS' TONGUE, *Asplenium scolopendrium*.

HARTS' TRUFFLES, *Elaphomyces granulatus*.

HARTWORT OF MARSEILLES, *Seseli tortuosum*.

HASCHISCH, see *Bangue*.

HASCHISCHIN, see *Bangue*.

HASTA, 'a spear,' *Penis*—h. *Nuptialis*, *Penis*—h. *Virilis*, *Penis*.

HASTECLA, (dim. of *hasta*), *Splint*.

HASTINGS, CLIMATE OF. This place has the reputation of being one of the mildest and most sheltered winter residences on the south coast of England. Owing to its low situation, and the height of the neighbouring cliffs, it is protected in a great degree from all northerly winds; and hence is found a favourable residence generally for invalids labouring under diseases of the chest.

HASTULA REGIS, (dim. of *hasta*), *Asphodelus ramosus*.

HATFIELD'S TINCTURE, see *Tinctura Guaiaci ammoniata*.

HATTREL (Sc.), ([G.] *Eiter*, 'pus.') *Abscess*.

HAUNCH, *Coxa*, *Coxen'dix*, of the Latins; *αἶσχος*, or *αἰσχος*, of the Greeks; *Ancha*, *Ischion*, *Omphys*, *Hip*, *Huckle*, *Cossa*, (Prov.) *Hubbon*, *Shoup* (F.) *Hanche*. The region of the trunk which is formed by the lateral parts of the pelvis and the hip-joint, including the soft parts. In women, on account of the greater width of the pelvis, the haunches are more marked and prominent than in men.

HAUNCHBONE, *Ilium*.

HAUSE, (Sax. *half*.) *Collum*, *Throat*.

HAUST, (Sax. *hōrta*, (G.) *Husten*.) *Tussis*.

HAUSTELLATE, *Haustellatus*, *haurio*, *haustum*, 'I drink up.') An epithet for the structure of mouth, which is adapted for drinking or pumping up liquids. Insects, which possess that kind of mouth, are so named. Ehrenberg reiers the fancied spermatozoa to the haustellate entozoa.

HAUSTRA COLI, (*haustrum*, 'a bucket.') See *Colon*.

HAUSTUS, gen. *Haustus*, (*haurire*, *haustum*, 'to drink.') *Potio*, dim. *Potivcula*. A Draught. A liquid medicine, which can be taken at a draught.

HAUSTUS NIGER, ('black draught,') see *Infusum sennæ*.

HAUT MAL, ('great disease;' from (L.) *altus*, 'high, great,' and *malum*, 'disease.') *Epilepsy*.

HAUTLÉ. An'inal bread of the Mexicans. A bread made of three species of hemipterous insects belonging to the group of water bugs.

HAVAN'A or HAVAN'NAH, CLIMATE OF. The climate of Cuba is often selected for the phthisical invalid during the winter months, and so far as regards elevation and comparative equality of temperature, it is more favourable for those of weak lungs than parts of the United States. The mean annual temperature is high (78°), but the difference between the mean temperature of the warmest and coldest months is twice as great as at Madeira.

HAVER, ([G.] Hafer,) Avena.
HAVERSIAN CANALS, (after Dr. Clopton Havers,) Canals, nutritive, of Bones.
HAW (Prov.), Hungry.
HAW, BLACK, (Sax. hæg,) Viburnum prunifolium.

HAWKNUT, Bunium bulbocastanum.
HAWKWEED, VEINY, Hieracium venosum.
HAWSE, (Sax. halp,) Throat.
HAWTHORN, WHITE, Mespilus oxyacantha.
HAY ASTHMA, Fever, Hay.
HAY, CAMEL'S, Juncus odoratus.
HAYRIFF, Galium aparine.
HAYSCALE (Prov.), Harelip.
HAZEL, BEAKED, (Sax. hærl,) Corylus rostrata.

HAZEL CROTTLES, Lichen pulmonarius—h. Nut, snapping, Hamamelis Virginiana—h. Nut tree, Corylus avellana.

HAZEL, WITCH, Hamamelis Virginiana, Sorbus aucuparia.

HAZELRAW, Lichen pulmonarius.

HEAD, (Sax. heapod, heapod, *heaved*;) the past participle of heapan, 'to heave up.' *Caput, Cū'ra, Cū're, Cep'hale, Cy'bē, Poll, Palet,* (Prov.) *Hede, Poir, Scop,* (F.) *Tête*. The head forms the upper extremity of the body, and tops the skeleton. It consists of the cranium and face. The first, which comprises all the superior and posterior part, has the encephalon in its cavity: the latter forms only the anterior part, and serves as the receptacle for the greater part of the organs of the senses.

HEAD, WATER IN THE, Hydrocephalus.

HEAD ACHE, Cephalæa, Cephalalgia—h. Ache, intermittent, Cephalalgia periodica—h. Ache, sick, Cephalæa spasmodica.

HEAD-MOULD-SHOT. An old name for the condition of the skull, in which the bones ride, or are *shot*, over each other at the sutures.

HEADSWOMAN (Prov.), Midwife.

HEAD'Y. Affecting the head. (F.) *Capiteux*. That which inebriates readily. An epithet, applied to wines which possess this quality.

HEAL, (Sax. hæl,) Sanitas.

HEALALL, Collinsonia Canadensis, Prunella vulgaris, Scrophularia nodosa.

HEALING, Curative—h. Art, Medicine—h. Piece, see Touch, royal.

HEALTH, (Sax. hæl,) Sanitas.

HEALTHY, Salutory. Also, whole; not diseased.

HEARING, HARDNESS OF, (Sax. hepan, 'to hear.'). Deafness—h. Perverse, Paracausis perversa.

HEART, (Sax. heort, (G.) Herz,) *Cor, Cear, Cer, Car'dia*, (F.) *Cœur*. An azgyous muscle, of an irregularly pyramidal shape; situate obliquely and a little to the left side, in the chest;—resting on the diaphragm by one of its surfaces;—suspended by its base from the great vessels; free and movable in the rest of its extent, and surrounded by the pericardium. The right side of the body of the heart is thin and sharp, and is called *Margo acutus*: the left is thick and round, and termed *Margo obtusus*. It is hollow within, and contains four cavities; two of which, with thinner and less fleshy walls, receive the blood from the lungs and the rest of the body; and pour it into two others, with thick and very fleshy parietes, which send it to the lungs and to every part of the body. Of these cavities, the former are called *auricles*, the latter *ventricles*. The right auricle and right ventricle form the *Pulmon'ic* or *right or anterior heart, Cor venosum seu pulmona'le*, (F.) *Cœur du poulmon, C. du sang noir, C. droit, C. antérieur*: and the left auricle and ventricle, the *system'ic, corporeal, left, or aortic heart, Cor arteriosum seu aorticum*, (F.) *Cœur du corps, C. gauche, C. cortique ou C. rouge*. In the adult, these are totally distinct from each other, being

separated by a partition;—the *septum cordis*. Into the right auricle, the *venæ cavæ*,—superior and inferior,—and the coronary vein, open. The pulmonary artery arises from the right ventricle (see *Conus arteriosus*): the four pulmonary veins open into the left auricle, and the aorta arises from the left ventricle.

The mean weight of the heart, in the adult, from the twenty-fifth to the sixtieth year, is, according to Bouillaud, from eight to nine ounces. The dimensions, according to Lobstein and Bouillaud, are as follows:—Length, from base to apex, five inches six lines; breadth, at the base, three inches; thickness of the walls of the left ventricle, seven lines; at a finger's breadth above the apex, four lines; thickness of the walls of the right ventricle, two and a quarter lines; at the apex, half a line; thickness of right auricle, one line; of the left auricle, half a line. The heart is covered, externally, by a very thin, membranous reflection from the pericardium. The muscular structure of which it is constituted is much thicker in the parietes of the ventricle than in those of the auricles. Its cavities are lined by a very delicate membrane, the *endocardidium*, which is continuous with the inner membrane of the arteries, as regards the left cavities, and with that of the veins, as regards the right. Its arteries—the *coronary*—arise from the commencement of the aorta. Its nerves proceed, chiefly, from the pneumogastric and the cervical ganglions of the great sympathetic. The heart is the great agent in the circulation. By its contraction, the blood is sent over every part of the body. Its action does not seem to be *directly* owing to nervous influence received from the brain or spinal marrow, or from both. The circulation may, indeed, be kept up for some time, if both brain and spinal marrow be destroyed.

When the ear is applied to the chest, a dull, lengthened sound is heard, which is synchronous with the arterial pulse. This is instantly succeeded by a sharp, quick sound, like that of the valve of a bellows or the lapping of a dog, and this is followed by a period of repose. The first sound appears to be mainly produced by the contraction of the ventricles; the second, by the reflux of the blood against the semilunar valves. These are what are called the *Sounds of the Heart, Cardiache'mata, Son'itus cordis seu cardi'aci*. (F.) *Bruits du Cœur*. Dr. C. J. B. Williams suggests the word *lubb-dup* as conveying a notion of the two sounds. *The Beating or Impulse of the heart, Heart-stroke, Apex beat, Pulsus seu Ictus cordis*, (F.) *Impulsion du Cœur*, against the parietes of the chest, is mainly caused by the systole of the heart, which tends to project it forwards. It is doubted, however, by some, whether the impulsion is produced by the dilatation or the contraction of the ventricles.

The following table exhibits the different actions of the heart, and their coincidence with its sounds and impulse. It presumes, that the period from the commencement of one pulsation to that of another is divided into eight parts; and if the case of a person, whose pulse beats sixty times in a minute, be taken, each of these parts will represent the eighth of a second.

EIGHTHS OF A SECOND.

Last part of the pause.....1...	Auricles contracting; ventricles distended.
First sound and impulse..4...	Ventricles contracting; auricles dilating.
Second sound.....2...	Ventricles dilating; auricles dilating.
Pause.....1...	Ventricles dilating; auricles distended.

The heart is subject to different organic diseases: the chief of which are *anæmia*, *contraction of the upper part*, and *expansion of its parietes*.

HEART, ATROPHY OF THE. *Atrophie du seu Aridra seu Phthisis Cordis, Cardiacophthisis, Anardiophthisis.* A condition of the organ in which there is diminution in the thickness of the parietes of the whole organ.

HEART, BEATING OF THE, see Heart.

HEART CLOT, see Polypus.

HEART, DILATATION OF THE, see Anæmism—h. Displacement of the, *Ectopia Cordis*—h. Fatty, *Stenosis cordis*.

HEART, HYPERTROPHY OF THE. *Hypertrophie Cordis, Hypercardia, Hypercardia Cordis, Hypertrophie Cordis, Cardiacæ, Anæmia cordis, Cardiacæ (Corrisart), Cardiacæ seu Cardiacæ, Cardiacæ Cordis.* (F.) *Hypertrophie du Cœur.* Supernutrition of the muscular parietes of the heart, which are thicker than usual; the cavities being generally diminished. The physical signs which indicate it are the following. In cases of long standing, the præcordial region is generally prominent; the pulsations of the heart are visible over a greater extent than natural; and a marked vibration is communicated to the hand when placed on the cardiac region. The dull sound on percussion is more extensive than natural; and on auscultation there is a permanent increase of the force and extent of the heart's action: there is no increase, however, of frequency, and the rhythm is regular. The pulse is generally strong, full, and hard.

HEART, CONCENTRIC HYPERTROPHY OF THE. (F.) *Hypertrophie concentrique, Hypertrophie concentrique du Cœur, Rétrécissement Cardiaque.* is when the parietes augment at the expense of the cavities.

HEART, HYPERTROPHY WITH DILATATION OF THE. *Atrophie anæmism, Excessive hypertrophy.* In this affection, the pulsations can be seen and felt over a larger space, and the apex is more to the left and lower down than natural. The impulse is less steady, but at times more violent than that which accompanies simple hypertrophy. Percussion gives more distinct evidence of the enlargement, the sound being more extensively dull. On auscultation, the impulse is often violent, but irregular; in extreme cases, it produces the sensation of a large mass of flesh rolling or revolving beneath the ear. The pulse is strong, full, and vibratory. The shock of the heart's action is often transmitted to the whole person, and to the bed on which the patient is lying.

HEART, IMPULSE OF THE, see Heart—h. Neuralgia of the, *Angina Pectoris*—h. Rapture of the, *Cardiorhexis*—h. Sounds of the, see Heart—h. Stroke, see Heart.

HEARTAXES, (Sax. *heaptæces*.) *Ardor ventriculi.*

HEARTBURN, *Ardor ventriculi, Cardialgia.*

HEART'S EASE, *Polygonum persicaria, Viola tricolor.*

HEARTS, LYMPH. *Lymphatic hearts—h. Lymphatic, Lymphatic hearts.*

HEARTSCALD, *Ardor ventriculi.*

HEARTSCIRTS (Prov.) *Diaphragm.*

HEARTWORT, *Laserpium album.*

HEAT, (past participle of Sax. *hefan*, 'to make warm') *Calor, Therma.* (F.) *Chaleur.* The material cause, which produces the sensation—or the particular sensation itself—produced by a body of an elevated temperature, on our organs, especially on the organs of touch. Modern chymists have given the name *Caloric* to the principle, whatever may be its nature, which is the cause of heat. When we touch a body of a temperature superior to our own, a portion of caloric passes from the body to the hand, and produces the sensation of heat. If, on the contrary, we touch a

body of a temperature inferior to our own, we communicate a portion of our caloric to it, and experience the sensation of cold. Our own sensations are but imperfect indexes of temperature. Two men meeting at the middle of a mountain,—the one ascending, the other descending,—will experience different sensations. The one ascending, passes from a warmer to a colder atmosphere;—the one descending from a colder to a warmer.

The chief instrument for measuring heat, used in medicine, is the thermometer. Of this there are three kinds:—that of Fahrenheit, that of Réaumur, and that of Celsius or the Centigrade.

To reduce Centigrade degrees to those of Fahrenheit, multiply by 9, divide by 5, and add 32: thus, $40^{\circ} \text{ Cent.} \times 9 \div 5 + 32 = 104^{\circ} \text{ Fahr.}$ *To reduce Fahrenheit's degrees to those of Centigrade,* subtract 32, multiply by 5, and divide by 9: thus, $104^{\circ} \text{ Fahr.} - 32 \times 5 \div 9 = 40^{\circ} \text{ Cent.}$ *To reduce Réaumur's degrees to those of Fahrenheit,* multiply by 9, divide by 4, and add 32: thus, $32^{\circ} \text{ Réaumur.} \times 9 \div 4 + 32 = 104^{\circ} \text{ Fahr.}$ *To reduce Fahrenheit's degrees to those of Réaumur,* subtract 32, multiply by 4, and divide by 9: thus, $104^{\circ} \text{ Fahr.} - 32 \times 4 \div 9 = 32^{\circ} \text{ Réaumur.}$ *To reduce Réaumur's degrees to those of Centigrade,* multiply by 5, and divide by 4: thus, $32^{\circ} \text{ Réaumur.} \times 5 \div 4 = 40^{\circ} \text{ Cent.}$ and, lastly, *to reduce Centigrade degrees to those of Réaumur,* multiply by 4, and divide by 5; thus, $40^{\circ} \text{ Cent.} \times 4 \div 5 = 32^{\circ} \text{ Réaumur.}$

By the term *gentle heat*, Ph. U. S., 1863, is meant any temperature between 90° and 100° .

The normal temperature of the body may be regarded as on the average, from 98° to 100° or upwards. The blood of the right side of the heart is slightly hotter than that of the left; and the highest temperature observed is where the suprahepatic veins empty their blood into the vena cava ascendens. See Animal heat.

The human body can bear a high degree of heat diffused in the atmosphere. There are cases on record, where air of 400° and upwards, of Fahrenheit's scale, has been breathed with impunity for a short time. It can likewise withstand very severe cold. In the expedition of Capt. Back to the Arctic regions, the thermometer was as low as -70° of Fahr., and Sir George Simpson affirms, that it has fallen in Siberia to -83° , or 113° below the freezing point. Dr. Kane, in the Arctic regions, in 1856-57, observed the range of eleven spirit thermometers, selected as standards, to vary from -60° to -73° . The mean annual temperature was $3^{\circ} 2$, the lowest ever registered.

Excessive heat disposes the body to gastric and intestinal diseases, and particularly to inflammation and enlargement of the liver; hence, the frequency of such affections within the torrid zone. Heat is often used therapeutically; the actual cautery, at a white heat, disorganizes the parts to which it is applied: a lesser degree occasions violent inflammation. Heat higher than that of the human body is excitant: of a lower degree, sedative. Excessive cold acts as a powerful sedative—in tusing sleep, the tendency to which, after long exposure, becomes irresistible. Many of the topical applications—as cataplasms—act wholly by virtue of their warmth and moisture.

HEAT, Artificial, Provoked. (F.) *Chaleur, Chaleur des Animaux, Folia, Rut.* is the periodical sexual desire experienced by animals,—*Estrotrium*. It is supposed by some to be owing to the periodical maturation and discharge of ova,—*Ovulation*.

HEAT, ACRID. see Acid—h. Animal, see Animal Heat.

HEAT APOPLEXY, see *Coup-de-soliel*.

HEAT, INTERNAL. (F.) *Chaleur interne,* is a sensation of heat felt by the patient, but not sensible to the touch. *External heat,* (F.) *Chaleur extérieure,* that which can be felt by others. Heat

A'dor, is called *moist*, (F.) *halitense*, when accompanied with moisture, like that felt after bathing; *dry*, (F.) *sèche*, when the skin has not its ordinary softness. It is called *acrid* and *pungent*, *Calor mordax*, (F.) *âcre et mordicante*, when it conveys a disagreeable tingling to the fingers. The French employ the terms *ner'vous* heat and *errat'ic* heat, *Chaleur nerveuse* and *Chaleur erratique*, for that which comes by flushes, alternating with chills, and which moves rapidly from one part to another. M. Double has used the term *septic heat*, (F.) *Chaleur septique*, for that which produces a pungent sensation on the hand, similar to that of acrid heat, but milder and more uniform, and which is accompanied with feebleness and frequency of pulse, &c.

HEAT, PRICKLY, Lichen tropicus.

HEATH COMMON, (Sax. hæð, Erica vulgaris.

HEATHER (Sc.), Erica vulgaris.

HEATHER-CLU, (Sc.) Malleolus.

HEAUTOPHONICS, (ἑαυτον, 'of himself,' and φωνη, 'voice,') see Autophonia.

HEAVINESS, Somnolency.

HEAVING, (from Sax. heapan, 'to heave, to raise,') Vomiturition.

HEAVY, *Gra'vis*, (F.) *Gravatif*, (Sax. hefig, hæfig.) An epithet given to any condition which consists in a sensation of weight or heaviness, or is accompanied by such sensation.

HEBDOMADAL CYCLE, (ἑβδομας, 'a week,' from ἑπτα, 'seven,') see Cycle.

HEBDOMADARIA, Octana.

HE'BÈ, (ἡβη,) *Juven'ta*, *Juven'tus*, *Juven'tus*, *Ho'ra*. In antiquity, the goddess of puberty. The word has been employed to designate, 1. The first hair that grows on the pubes, 2. The pubic region, and 3. Puberty.

HEBETES, (ἡβητης,) Adolescens.

HEBETER, (ἡβητηρ,) Adolescens.

HEBETUDO ANIMI, (*hebes*, 'dull,') Imbecility — h. Dentium, *Hæmodia* — h. Visus, *Ambyopia*, *Asthenopia*, *Caligo*.

HEBRADENDRON CAMBOGIODES, see Cambogia.

HE'BREWS, MED'ICINE OF THE. Medicine seems to have been at a very low ebb with the ancient Hebrews. Of anatomy they knew nothing. Their physiology was imperfect and filled with superstitions; and their therapeutics unsatisfactory. Hygiene appears to have been most attended to. Of the other departments of medicine we cannot judge of their knowledge.

HECATORIA PALUSTRIS, (from ἑκατον, 'a hundred,' from its numerous flowers,) *Ranunculus sceleratus*.

HECATOPHYLLA, (ἑκατον, 'a hundred,' and φυλλον, 'a leaf,') *Rosa centifolia*.

HECH, (by onomatopœia,) Anhelatio.

HECKBERRY (Prov.), *Prunus padus*.

HEC'TEUS, (ἑκτευσ,) A Greek measure, containing about 72 chopines or pints.

HECTIC, *Hec'ticus*, (from ἔκτισ, 'habit,') (F.) *Hectique*, *Étique*. Relating or appertaining to the constitution or habit.

HECTIC FEVER, *Fe'bris hec'tica* seu *phthis'ica* seu *marasmo'des* seu *tab'ida* seu *len'ta* seu *amphimer'ina* *hec'tica* seu *amato'ria*, *Hectico'py'ra*, *Hectico'py'etos*, *Hec'tica*, *Amphimer'ina* *hec'tica*, *Synteco'py'ra*, *Syntectico'py'ra*, *Marasmo's'py'ra*, *Leuco'py'ra*, *Epan'etus* *hec'ticus*, *Chloro's'is* *amato'ria*, *É'thick*, (Sc.) *Ethick*, (F.) *Fie'vre Hectique* ou *Étique*, (ἔκτισ, 'habit of body,' because in this disease every part of the body is emaciated; or, perhaps, from ἐκτενω, 'I consume,' 'I am exhausted,') The name of a slow, continued, or remittent fever, which generally accompanies the end of organic affections, and has been esteemed idiopathic, although it is probably always symptomatic. It is the fever of irritation and debility;

and is characterized by progressive emaciation, frequent pulse, hot skin,—especially of the palms of the hands and soles of the feet,—and, towards the end, colliquative sweats and diarrhoea. Being symptomatic, it can only be removed by getting rid of the original affection. This is generally difficult, and almost hopeless in the disease which it most commonly accompanies,—consumption.

HECTICA, Hectic Fever—h. Infantilis, Fever, infantile remittent.

HECTICOPYRA, (ἑκτικός, 'hectic,' and πυρ, 'fire or fever,') Hectic fever.

HECTICOPYRETOS, (ἑκτικός, 'hectic,' and πυρετος, 'fever,') Hectic fever.

HECTIQUE, (ἑκτικός,) Hectic.

HECTIQUE, Hectic.

HEC'TISIS, (F.) *Étisie*. The state of those who have hectic fever.

HEC'TOGRAMME, *Hectogram'ma*, (ἑκατον, 'a hundred,' and γραμμα,) A measure of 100 grammes, i. e., 3 ounces, 1 drachm, and 44 grains, Troy.

HECTOLITRE. A measure containing 100 litres or 26.42 wine pints.

HECUSIUS, (ἑκουσις, and ἔων,) Voluntary.

HEDE (Sc.), Head.

HEDEO'MA, *Hedeo'ma pulegioid'es*, (ἡδον, 'agreeable,') *Cumi'la pulegioid'es*, *Melis'sa pulegioid'es*, *Penny'royal*, *Tick'weed*, *Stink'ing Balm*, *Squaw'mint*. *Sez. Syst.* Diandria Monogynia; *Nat. Ord.* Labiatæ. An indigenous plant, common in all parts of the United States, and, which, where it is abundant, perfumes the air for a considerable distance. It is employed in the same cases as the mints and the English pennyroyal. In popular practice it is used as an emmenagogue.

The *O'leum Hedeo'mæ* (Ph. U. S.) or *Oil of Pennyroyal* is used as a stimulating carminative, dropped on sugar. Dose, 2 to 6 drops.

HERERA ARBorea, H. Helix.

HE'D'ERA HE'LIX, *Hed'era arbo'rea*, *Bac'chica*, *Cis'aus*, *Cit'tos*, *Ed'era*, *Corymbe'thra*, *Corym'bos*, *Ivy*, (Sc.) *Bind'wood*, (F.) *Lierre*, *L. commun* ou *grim'pant*. *Ord.* Araliaceæ. The taste of *ivy* leaves is bitter, styptic, and nauseous. They are not used in medicine. According to Haller, they were recommended in Germany against atrophy of children; and the common people of England sometimes apply them to running sores and to keep issues open. The berries were supposed, by the ancients, to have an emetic and purgative quality; and a watery extract was made from them, called by Quercetanus *Extractum purgans*. From the stalk of the tree a resinous juice exudes, in warm climates, called *Gummi Hed'erae*, *Ivy Gum*, (F.) *Gomme de lierre*, *Résine de lierre*. It is possessed of tonic and astringent properties, but is not used.

HERERA TERRESTRIS, *Glechoma hederacea*.

HERERULA, (dim of *hedera*), *Glechoma hederacea*.

HEDEVERK, ([Sax.] heapod-pæpæne,) Cephalalgia.

HEDEWARK, Cephalalgia.

HE'D'RA, (ἡδρα, ἑδρη, 'a vestige,' 'a seat,') A seat. A fracture of the bones of the cranium, in which the trace of the fracturing instrument is still perceptible. It was, also, used by the ancients, for the anus, the breech, excrement, a privy, a night-chair, and for the bottom of an abscess.

HEDROCELE, (*hedra*, and κηλη, 'a tumour,') Proctocoele.

HEDY, in composition (from ἡδύς,) sweet, agreeable.

HEDYCH'ROUM, (*hedy*, and χροα, 'colour,') A remedy of a pleasant colour. Applied to certain trochs, the chief constituent of which was theriac.

HEDYPHONIA, (*hedy*, and *φωνη*, 'voice.') Sweetness of voice.

HEDYPNEUS'TUS, *Hedyp'nous*, (*hedy*, and *πνέω*, 'I breathe.') Breathing sweetly or softly. Smelling sweetly.

HEDYPOIS TARAXACUM, (*hedy*, and *πνέω*, 'to breathe,' from its scenting the breath.) *Leon-todon taraxacum*.

HEDYSARUM ALHAGI, (*ἡδύσαρον*, from *hedy*, and *αρωμα*.) *Agv*

HEDYSMA, gen. *Hedysm'atis*, (*ἡδύσμα*.) Conditioned.

HEEL, (Sax. *hel*.) see *Calcaneum*—h. Clubfoot, see *Kyllosis*—h. String, *Achillis tendo*.

HEGH, (by onomatopœia.) *Anhelatio*.

HEIL, (Sax. *hæl*.) *Sanitas*.

HELCENTERITIS, (*helco*, and *enteritis*.) *Dochthenteritis*.

HELCO, in composition, from *ἔλκος*, 'an ulcer.'

HELCO'ACE, *Cachelo'ma*, (*helco*, and *κακη*, 'badness.') A malignant ulcer.

HELCODES, (*ἑλκωδης*, from *helco*, and *odes*.) Ulcerated, Ulcerous.

HELCOLOGY, *Helcolog'ia*, (*helco*, and *λογος*, 'a discourse.') The doctrine of, or a treatise on, ulcers.

HELCOMA, gen. *Helco'matis*, (*ἑλκωμα*.) Ulcer.

HELCOMENIA, (*helco*, and *μηνες*, 'menses,') *Menelcosis*.

HELCOPTHAL'MIA, *Ophthalm'ia ulcero'sa*. *Ophthalmia* with ulceration.

HEL'COPLASTY, *Hel'koplasty*, *El'koplasty*, *Elo'oplasty*; (*helco*, and *πλασσω*, 'I form.') An operation which consists in grafting on an ulcer a piece of skin from the opposite limb, or from the limb of another person, when the destruction of skin is too great to permit the healing process.

HELCOISIS, (*ἑλκωσις*, from *helco* and *osis*.) *Elcosis*, Ulceration—h. *Cerebri*, *Encephalopyosis*—h. *Laryngis*, *Phthisis laryngea*—h. *Pulmonalis*, *Phthisis pulmonalis*—h. *Renalis*, *Nephrelecosis*—h. *Uteri*, see *Metrophthisis*—h. *Vesicæ*, *Cystophthisis*.

HELCOSTAPHYLO'MA, gen. *Helcostaphylo'matis*, (*helco*, and *σταφυλωμα*, 'staphyloma.') *Staphyloma* ending in ulceration.

HELCOXERO'SIS, (*helco*, and *ξηρωσις*, 'dryness.') The drying of an ulcer.

HELTICA, (from *ἔλκω*, 'to draw,') *Epispas-tics*.

HELCUS, (*ἑλκος*.) Ulcer.

HELCOYDRION, *Helcyd'rion*, *ἑλκυδριον*, (dim. of *Helcus*.) *Ulcus culum*. A small ulcer, a superficial ulceration of the cornea.

HELCYSTER, (*ἑλκυστηρ*, from *ἔλκω*, 'I draw.') An iron hook or erotchet for extracting the fœtus. See *Crotchet*.

HELENIUM, (after *Helena* [?]), *Inula helenium*.

HELEN'IUM AUTUMNA'LE, *False Sunflower*, *Sneeze'wort*, *Sneeze'weed*, *Swamp Sunflower*, *Yellow star*, *Ox'eye*. *Ord. Compositæ*. An indigenous herb, with large goldenyellow compound flowers, which appear in August. All its parts are bitter and somewhat acrid, and when snuffed up the nostrils in powder are powerful sternutatories.

HELIAN'THEMUM CANADEN'SÊ, (*ἡλιος*, 'sun,' and *ἄνθεμον*, 'flower.') *Helianthemum* (Ph. U. S.), *Cistus Canaden'sis*, *Frost'wort*, *Frost'weed*, *Frostplant*, *Rockrose*. *Ord. Cistaceæ*. An herbaceous plant, having large yellow flowers, which grows in all parts of the United States, and flowers, in the Middle States, in June. It has an astringent, slightly aromatic, and bitterish taste. It has been prescribed in scrofula, but probably is nothing more than an aromatic tonic.

HELIAN'THEMUM CORYMBO'SUM, *Rockrose*, an indigenous plant, is used in the same cases.

HELIASIS, (*ἡλιασις*, from *ἡλιος*, 'the sun,') *Astrabolismus*, *Insolation*.

HELIC'RYSUM NUDIFOLIUM, (*ἡλιος*, 'sun,' and *χρυσος*, 'gold,') *Caffer tea*. A South African plant, *Nat. Ord. Compositæ*, which is demulcent, and, in the form of infusion, is recommended in catarrh, phthisis, and other pulmonary affections.

HELICRYSUM SERPYLLIFO'LIUM, *Hot'tentots' tea*, and **HELICBYSUM AURICULA'TUM** have similar virtues.

HELICIA, (*ἡλίκια*.) *Age*.

HEL'ICINE, *Helic'innus*, *Helic'oides*, *Helico'des*, (from *ἑλκx*, gen. *ἑλίκος*, 'the tendril of the vine,') Resembling in form the tendril of the vine.

HELICINE ARTERIES of the penis, as described by J. Müller, are short vessels given off from the larger branches, as well as from the finest twigs of the artery of the organ: most of those come off at a right angle, and project into the cavity of the spongy substance, either terminating abruptly or swelling out into a club-like process without again subdividing. Almost all these vessels are bent like a horn, so that the end describes half a circle or somewhat more. They have a great resemblance to the tendrils of the vine, whence their name. A minute examination of them, either with the lens or the microscope, shows that, although they at all times project into the venous cavities of the corpora cavernosa, they are not entirely naked, but are covered with a delicate membrane, which, under the microscope, appears granular.

HEL'ICIS MA'JOR, (gen. of *helix*.) A muscle of the ear, which originates from the anterior, acute part of the helix, upon which it ascends and is inserted into the helix. It pulls the part into which it is inserted a little downwards and forwards.

HELICIS MI'NOR. This muscle originates from the under and fore part of the helix, and is inserted into the helix, near the fissure in the cartilage, opposite the concha. Its use is to contract the fissure.

HELICOIDES, (*εἰλιξ*, and *εἶδος*, 'resemblance,') *Helicine*.

HELICOTRE'MA, gen. *Helicotre'matis*, (*εἰλιξ*, 'helix,' and *τρημα*, 'a foramen,') The hole by which the two scalæ of the cochlea communicate at the apex.

HELIENCEPHALITIS, (*ἡλιος*, 'the sun,' and *encephalitis*.) see *Coup de Soleil*.

HELIGMUS, (*ἡλιγμος*, from *ἐλίσσω*, 'to turn round,') *Convolution*.

HELIKIA, (*ἡλίκια*.) *Age*.

HELIOMYELITIS, (*ἡλιος*, 'the sun,' *μυελος*, 'the marrow,' and *itis*.) *Myelitis* from exposure to the sun.

HELIONOSIS, (*ἡλιος*, 'the sun,') *Insolation*.

HELIOPSYDRA'CIUM, (*ἡλιος*, 'the sun,' and *psydracium*.) A *psydracium* or pustule from exposure to the sun.

HELIOSCIADIUM NODIFLORUM, (*ἡλιος*, 'the sun,' and *σκιαδιον*, 'a parasol,') *Sium*.

HELIOSIS, (*ἡλιωσις*.) *Astrabolismus*, *Insolation*.

HELIOTROPE, (*ἡλιος*, 'the sun,' and *τροπεω*, 'I turn,') *Heliotropium Europæum*.

HELIOTROPION, *Cichorium intybus*.

HELIOTROP'PIUM EUROPE'UM, *H. erectum* seu *canescens* seu *eupyr'ium*, *Verruca'ria*. The *Heliotrope*, (F.) *Tournefort*, *Herbe aux verrues*. *Ord. Boraginaceæ*. This plant is considered to possess aperient properties; and to be capable of destroying cutaneous excrescences; hence one of its names.

HELIOTROP'PIUM IN'DICUM, an East India plant, naturalized in some of the Western and Southern States, probably has similar virtues.

HELIX, gen. *Hel'icis*, (*εἰλιξ*, gen. *ἑλίκος*, 'the

tendrill of the vine,' from *ελεεινός*, 'to envelop,' 'to roll.') *Copeus olus*. The fold is thus called, which forms the outer circumference or ring of the external ear.

HELIX, *Limax*.

HELIX POMATIA, (F.) *Limacon*, *Colimaçon*, *Escargot*. A large kind of snail, transported from the south of Europe to England by Sir Kenelm Digby, for his lady when in a decline. It was considered highly restorative.

HELLA, *Inebus*.

HELLEBORASTER, (from *ελληβορος*), *Helleborus foetidus*—h. *Fœtidus*, *Helleborus foetidus*.

HELLEBORE, *Dracontium foetidum*—h. American, *Veratrum viride*—h. black, *Helleborus niger*—h. *Blanc*, *Veratrum album*—h. Green, *Helleborus viridis*—h. *Noir*, *Helleborus niger*—h. Stinking, *Helleborus foetidus*—h. Swamp, *Veratrum viride*—h. White, *Veratrum album*.

HELLEBORISMUS, (from *ελληβορος*), *Elleborismus*. The method of treating disease, amongst the ancients, by hellebore. This comprised not only the choice, preparation, and administration of the medicine, but, likewise, the knowledge and employment of preliminary precautions and remedies proper for aiding its action, and preventing the pernicious effects which it might occasion.

HELLEBORUS, (*ελληβορος*), H. *niger*—h. *Albus*, *Veratrum album*.

HELLEBORUS FŒTIDUS, *Helleborus foetidus*, *Elleboraster*, *Stinking Hellebore* or *Beaver's foot*, *Saturnwort*, (F.) *Hellébore* ou *Ellebor foétide*, *Pied de Griffon*, *Pas de lion*. *Ord.* Ranunculaceæ. The leaves of this plant are said to be antihelmintic. The smell of the fresh plant is extremely fetid, and the taste bitter and acrid. It usually acts as a cathartic.

HELLEBORUS GRANDIFLORUS, H. *niger*.

HELLEBORUS NIGER, H. *grandiflorus*, *Elleborus niger*, *Melampodium*, *Melamorchium*, *Ecetonum*, *Black Hellebore*, *Melampyre*, *Christmas Rose*, *Closetongue*, (F.) *Hellébore noir*. The root of this European plant—*Helleborus*, (Ph. U. S.)—has a disagreeable odour, and bitter, acrid taste. It is possessed of cathartic properties, and has been recommended as an emmenagogue. It has been given in mania, melancholia, dropsy, suppressed menses, &c. Dose, gr. x to ℥j. as a cathartic.

HELLEBORUS TRIFOLIUS, *Coptis*.

HELLEBORUS VIRIDIS, *Green Hellebore*. A European plant, naturalized on Long Island, has similar properties to the last.

HELLECBRA, *Sedum*.

HELLENIA GRANDIFLORA, (after C. N. Hellenius, professor at Abo,) *Costus*.

HELMETFLOWER, YELLOW, *Aconitum Anthora*.

HELMET POD, *Jeffersonia Bartoni*.

HELMINS, gen. *Helminthis*, plur. *Helminthes* seu *Elminthes*, (from *ελμιν*, gen. *ελμινος*, 'a worm,') A worm; an entozoon.

HELMINTHOGOGUE, (*helmintho*, and *αγω*, 'to drive,') Anthelmintic.

HELMINTHI, Worms.

HELMINTHIA, Helminthiasis—h. *Alvi*, Worms.

HELMINTHIA ERRATICA, Worms, introduced by accident and without finding a proper habitation in the stomach or intestines; producing spasmodic colic, with severe tormina, and occasionally vomiting or dejections of blood; the 12th genus of the order *Enterica*, class *Celiaca*, of Good.

HELMINTHIA PODICIS, Worms.

HELMINTHIASIS, *Helminthogenes*. A generic name for the condition which gives occasion to the presence of intestinal worms. It is, also, called *Helminthia*, *Verminalio*, *Morbus verminosus*, *Status verminosus*, *Saburra verminos*, &c.

Scoleciasis, *Scoleciosis*, *Entozoiogenes*, *Parasitismus intestinalis*, Worm disease, *Inverminalio*, *Verminalio*. See Worms.

HELMINTHIC, Anthelmintic.

HELMINTHO, (*ελμιν*, gen. *ελμινος*, 'a worm,') in composition, worm.

HELMINTHOCHORTUM, (*helmintho*, and *χортос*, 'grass,') *Corallina Corsicana*.

HELMINTHOCOLICA, (*helmintho*, and *colica*), *Colica verminosa*.

HELMINTHOCORTON, (*helmintho*, and *χортос*, 'grass,') *Corallina Corsicana*.

HELMINTHOGENESIS, (*helmintho*, and *γενεσις*, 'generation,') Helminthiasis.

HELMINTHOL'OGY, *Helmintholog'ia*, *Scolecolog'ia*, (*helmintho*, and *λογος*, 'a description,') A treatise on worms.

HELMINTHONCUS, (*helmintho*, and *ογκος*, 'a tumour,') *Malis*—h. *Medicensis*, *Dracunculus*.

HELMINTHOP'YRA, *Helminthopyretus*, *Febbris verminosa*, (*helmintho*, and *πυρ*, 'a fever,') Fever occasioned by worms. See Fever, verminous.

HELMINTHOPYRETUS, (*helmintho*, and *πυρετος*, 'fever,') *Helminthopyra*.

HELMINTHUS GORDII, *Seta equina*.

HELODES, (*ελος*, 'a marsh,' and *οδης*), *Elodes*.

HELONIAS DIOICA, (from *ελος*, 'a swamp,') *Chamaerium luteum*—h. *Erythrosperma*, *Amianthum muscætoxiænum*—h. *Lutea*, *Chamaerium luteum*—h. *Officinalis*, see *Veratrina*.

HELOPYRA, (*ελος*, 'a marsh,' and *πυρ*, 'fever,') *Elodes* (*febris*).

HELOPYRETUS, (*ελος*, 'a marsh,' and *πυρετος*, 'fever,') *Elodes* (*febris*).

HELO'SIS, *Helot'is*, (*ειλω*, 'I turn,') Eversion of the eyelids, and convulsions of the muscles of the eyes. *Plica Polonica*. *Strabismus*.

HELOTIS, *Helosis*, *Plica*.

HELUS, marsh, (*ελος*); *Clavus*, (*ηλος*).

HELVELLA ESCULENTA, *Morchella esculenta*.

HELVINE, (from *ελω*, 'to draw, from its adherent properties,') *Parietaria*.

HEMA, (F.) *Héma*, (*αιμα*, gen. *αιματος*, 'blood,') in composition, 'blood.'

HÉMACELINEOSE, *Purpura hæmorrhagica*.

HEMACHROIN, (*hema*, and *χρος*, 'colour,') *Hæmatin*.

HÉMADYNAMÈTRE, *Hæmadynamometer*.

HÉMADYNAMIQUE, *Hæmadynamics*.

HÉMAPÉRITONIRRHAGIE, *Hæmaperitonirrhag'ia*, (*hema*, *peritonæum*, and *ρηνεμ*, 'I break forth,') Hemorrhage into the peritoneum.

HÉMAPROCTIE, (*hema*, and *πρωκτος*, 'anus,') Hemorrhoids.

HÉMASTATIQUES, *Hæmatostatica*.

HÉMATIDROSE, (*hema*, and *ιδρος*, 'sweat,') *Sudor cruentus*.

HÉMATIES, Globules, red, of the blood.

HEMATIN, *Hæmatin*.

HEMATO, in composition, see *Hema*.

HÉMATO-ENCÉPHALIE, *Apoplexy*.

HEMATOID, *Hæmatodes*.

HÉMATOIDINE, *Hæmatoïdin*.

HÉMATOLOGIE, *Hæmatology*.

HEMATOLOGY, *Hæmatology*.

HÉMATOMYÉLIE, (*hema*, and *μυελος*, 'marrow,') *Hæmorrhagie de la Moëlle Épineière*.

HÉMATONCIE, *Hæmatoncus*, see *Hæmatodes fungus*—h. *Fonguïde*, *Hæmatodes fungus*.

HÉMATOPISIE, *Hæmatops*.

HÉMATORRHACHIS, *Apoplexia myelitica*.

HEMATOSIN, *Hæmatin*.

HEMATOSTATICS, *Hæmatostatica*.

HÉMATOSTATIQUES, *Hæmatostatica*.

HÉMATOZOAIRES, *Hæmatozoa*.

HÆMATURIE. Hæmaturia.

HÆMENDOCARDE, Hæmendocardium.

HEMERA, ('*ἡμερα*, 'a day.') Dies. Hence:

HEMERALOPIA, see Nyctalopia.

HEMERALOPIA, (*hemera*, and *σκοποι*, 'I see.')

Hæmeralopia, *Dysopia* seu *Culi'go tenebra'rum*, *Parop'ia* *Noctif'uga*, *Vi'sus diurn'us*, *Nyctalopia* (*of some*), *Nyctotroph'ia*, *Amblyopia crepuscula'ris*, *Cæcitas crepuscul'ris* seu *nocturna*, *A'cia diurna*, *Daysight*, *Dayvision*, *Henblind'ness*, *Nightblind'ness*, *Nocturnal Amauro'sis*, (F.) *Héméralopie*, *Vue diurne*, *Augement de Nuit*, *Cécité nocturne*. A disease, in which the eyes enjoy the faculty of seeing, whilst the sun is above the horizon, but are incapable of seeing by the aid of artificial light. Its causes are not evident. The eye, when carefully examined, presents no alteration, either in its membranes or humours.

HEMERALOPS, gen. *Hemeralop'is*, *Do'rea*. One labouring under hemeralopia. Also, Hemeralopia.

HEMERATYPHLOSIS, (*hemera*, *τυφλος*, 'blind,' and *osis*.) Nyctalopia.

HEMERODROMA, gen. *Hemerodrom'atis*, (*hemera*, and *δρομος*, 'a course,') Ephemera.

HEMEROPATHIA; (*hemera*, and *παθος*, 'an affection,') A disease, which continues only a day; or, which is only observed during the day.

HEMEROTYPHLOSIS, (*hemera*, *τυφλος*, 'blind,' and *osis*.) Nyctalopia.

HEMI, ('*ἡμι*, '*ἡμισυς*,' 'half,' 'semi.' Hence:

HEMIACEPHALUS, (F.) *Hémiacéphale*, (*hemi*, *a*, and *κεφαλη*, 'head,') A monster whose head is represented by a shapeless tumour, with some appendages or cutaneous folds anteriorly, the thoracic extremities existing.

HEMIÄMAUROSIS, Hemipia.

HEMIÄNDRUS, ('*ἡμιάνδρος*, from *hemi*, and *ανδρ*, gen. *άνδρος*, 'a man,') Eunuch.

HEMIÄNOR, (*hemi*, and *ανηρ*, 'a man,') Eunuch.

HEMIÄNTHROPIA, ('*ἡμιανθρωπια*, from *hemi*, and *ανθρωπος*, 'a man,') Mania.

HEMIÄNTHROPUS, Eunuch, Maniac.

HEMIÄZYGA (Vena), see Azygos vein.

HEMIC, Hæmic.

HEMICEPHALÆA, (*hemi*, and *κεφαλη*, 'head,') Hemierania.

HEMICEPHALÆUM, Sinciput.

HEMICEPHALIUM, Sinciput.

HEMICEPHALUM, Sinciput.

HEMICEPH'ALUS, *Hemiceph'alus*, *Semi-ceph'alus*. One who has half a head.

HEMICRANIA, ('*ἡμικρανία*.) *Hemicephalæ'a*, *Migra'na*, *Gra'na*, *Hemip'gia*, *Hemipe'gia*, *Heterocrania*, *Monop'gia*, *Menop'gia*, *Cephale'a* *Hemicra'nia*, *Hemipath'i'a*, *Neural'gia cerebralis*, *Hem'icrany*, *Me'grim*, (*hemi*, and *κρανιον*, 'cranium,') (F.) *Migraine*. Pain, confined to one-half the head. It is almost always of an intermittent character;—at times, continuing only as long as the sun is above the horizon; and hence sometimes called *Sun pain*,—and is cured by cinchona, arsenic, and the remedies adapted for intermittents.

HEMICRANIA IDIOPATHICA, Neuralgia, facial.

HEMICRANY, Hemierania.

HEMIDESMUS IN'DICUS, (*hemi*, and *δεσμος*, 'a tie,' in allusion to the filaments,) *Perip'loca In'dica*, *Asclepias Pseudosar'sa*, *Sarsaparil'la In'dica*, *Nannari*, *In'dian Sarsaparil'la*. Ord. Asclepiadaceæ. A Hindoostane plant, the root of which, *Hemidesm'us* (P. B.), has a peculiar aromatic odour, and a bitterish taste. It is used in India as a substitute for sarsaparilla.

HEMIDIALYSIS, (*hemi*, and *διαλυσις*, 'dissolution,') Hemiplegia.

HEMIDIAPHORE'SIS, *Transpira'tio Uni-*

lu'era; (*hemi*, and *διαφορησις*, 'perspiration,') Perspiration of one-half the body.

HEMIECTON, ('*ἡμiekτον*.) *Hemiec'teon*, *Semiec'tum*. A vessel capable of containing 36 chopines or pints, and in which fumigations were formerly made in diseases of the uterus and vagina.

HEMIENCEPH'ALUS, (F.) *Hémiencéphale*, (*hemi*, and *encephalos*, 'the encephalon,') A monster, which has no trace of the organs of the senses, and yet has a brain nearly natural. See Hemicephalus.

HEMIG'YNUS, ('*ἡμιγυνος*, from *hemi*, and *γυνη*, 'a woman,') One who is effeminate, or half a woman.

HEMIM'ELES, (F.) *Hémimèle*, (*hemi*, and *μελος*, 'a limb,') A genus of monsters, in which the upper or lower extremities are very defective—mere stumps,—and the fingers and toes are entirely wanting or very imperfect.

HEMIME'RION, (*hemi*, and *μερα*, 'a part,') 'One half part.' Also, half a drachm.

HEM'INA, ('*ἡμινα*.) A Greek measure, answering to the *Cotyle*, *κότυλη*, i. e. one half the sextarius, or about half a pint, English.

HEMIOBOL'ION, ('*ἡμιβολιον*.) *Hemiob'olon*. Half the obolus. A weight of about five grains.

HEMIOL'IOS, ('*ἡμιολιος*.) A weight of 12 drachms, or oz. 1½. See Sescuncia.

HEMIOPIA, *Hemiop'ia*, *Hemiop'iasis*, *Suf-fu'sio dimid'iana*, *Vi'sus dimidia'tus*, *Marmor'ygê Hippocr'atis*, *Hemiamau'ro'sis*, *Amauro'sis dimidia'ta*, (F.) *Hémiopie*, *Hémiopie*; (*hemi*, and *σκοποι*, 'I see,') Depraved vision, in which the person sees only one half of an object.

HEMIOPIASIS, Hemipia.

HÉMIOPSIE, Hemipia.

HEMIOPSIS, Hemipia.

HEMIPA'GES, *Dicor'ygna's dihyogas'trius*, *Octo'pus synapheceph'alus*, (*hemi*, and *πυγνυμι*, 'I fasten,') A monstrosity, in which twins are united from the navel to the vertex.

HEMIPAGIA, Hemierania.

HEMIPATHIA, (*hemi*, and *παθος*, 'affection,') Hemierania.

HEMIPEGIA, Hemierania.

HEMIPHO'NIA, (*hemi*, and *φωνη*, 'voice,') Great weakness of voice. The husky voice of those affected with cholera.

HEMIPLE'GIA, *Hemiplex'ia*, *Hemidia'l'ysis*, *Para'l'ysis Hemiple'gia*, *Semiple'gia*, *Semisidera'tio*, (*hemi*, and *πλησσω*, or *πληγω*, 'I strike,') Paralysis of one side of the body. See Paralysis. One so palsied is said to be *hemipleg'ic*, *semi-sidera'tus*.

HEMIPLEGIA FACIALIS, Paralysis, Bell's.

HEMIPLEXIA, ('*ἡμικληξία*.) Hemiplegia.

HEMIPROSOPLÉGIA, (*hemi*, *προσωπον*, 'face, and *πληγω*, 'a stroke,') Paralysis, Bell's.

HEMISPHERÆ CEREBRI, Hemispheres of the brain.

HEMISPHERE, *Hemisphæ'ra*, *Hemisphæ'rium*, (*hemi*, and *σφαίρα*, 'a sphere,') One half of a sphere or of a body having a spheroidal shape.

HEMISPHERES OF THE BRAIN, *Hemisphæ'ræ cer'ebrî*, *Hemispher'ical gan'glia*, are the upper spheroidal portions of the brain, separated from each other by the falx cerebri.

HEMISPHERICAL GANGLIA, Hemispheres of the brain.

HÉMITE, Hæmitis.

HEMITRITÆA (FE'BRIS), ('*ἡμιτριταιος* (πυρετος), *Fe'bris semitertiana* seu *sesquial'tera*, (F.) *Hémitritée*, *Demitierce*, *Fèvre demitierce*, (*hemi*, and *τριταος*, 'tertian,') A semitertian fever, so called because it seems to possess both the characters of the tertian and quotidian intermittent.

HEMIUNCIA, (*hemi*, and *ουνκία*, 'an ounce,')
Hemiuncion.

HEMIUNCION, *Hemion'cia*, *Semion'cia*, *Semion'cia* Half an ounce.

HEMLOCK, *Pinus Canadensis*, *Conium maculatum*—*h. American*, *Cicuta maculata*—*h. Bastard*, *Oberophyllum sylvestre*—*h. Common*, *Conium maculatum*.

HEMLOCK DROPWORT, *Oenanthe*.

HEMLOCK, GROUND, *Taxus Canadensis*.

HEMLOCK GUM, see *Pinus Canadensis*—*h. Pitch*, see *Pinus Canadensis*—*h. Poison*, *Conium*—*h. Spruce*, *Pinus Canadensis*.

HEMLOCK, WATER, AMERICAN, *Cicuta maculata*, *Cicuta aquatica*—*h. Water*, fine-leaved, *Phellandrium aquaticum*—*h. Wild*, *Cicuta maculata*.

HÉMOANGIAIRRHAGIE, (*hemo*, *αγγειον*, 'a vessel,' *ang*, 'air,' and *ῥαγναι*, 'to break forth,')
Hæmoptysis.

HÉMOCARDE, (*hemo*, and *καρδία*, 'heart,')
Hæmocardiorrhagia.

HÉMOCARDIOPLASTIES, (*hemo*, *καρδία*, 'heart,' and *πλασσω*, 'to form,') see *Polypus*.

HEMODIA, *Hæmodia*.

HEMO, in composition, see *Hema*.

HÉMODYNAMETRE, *Hæmadynamometer*.

HÉMODYNAMOMÈTRE, *Hæmadynamometer*.

HÉMOENCÉPHALORRHAGIE, (*hemo*, *encephalon*, and *ῥαγναι*, 'to break forth,') *Apoplexy*.

HEMOHÉPATORRHAGIE, *Hepatorrhagia*.

HÉMOMETRE, *Hæmadynamometer*.

HÉMOYÉLORRHAGIE, (*hemo*, *μυελος*, 'marrow,' and *ῥαγναι*, 'to break forth,') *Apoplexy*, spinal.

HÉMOPÉRICARDE, *Hæmopericardium*.

HÉMOPHILIE, *Hæmatophilia*.

HÉMOPHOBIE, *Hæmatophobia*.

HÉMOPHTHALMIE, *Hæmophthalmia*.

HÉMOPLANIE, *Hæmatoplasia*.

HÉMOPROCTIE, see *Hæmorrhoids*.

HÉMOPTOÏQUE, *Hæmoptoic*.

HÉMOPTYIQUE, *Hæmoptoic*.

HÉMOPTYISIE, *Hæmoptysis*—*h. Foudroyante*, see *Hæmoptysis*.

HÉMOPTYISQUE, *Hæmoptoic*.

HEMORRHAGE, *Hæmorrhagia*—*h. Accidental*, see *Hæmorrhagia*—*h. Active*, see *Hæmorrhagia*—*h. Ante partum*, see *Ante partum*—*h. from the Bladder*, *Cystirrhagia*—*h. Constitutional*, see *Hæmorrhagia*—*h. Critical*, see *Hæmorrhagia*—*h. by Exhalation*, see *Hæmorrhagia*—*h. External*, see *Hæmorrhagia*—*h. General*, see *Hæmorrhagia*—*h. Internal*, see *Hæmorrhagia*—*h. Interstitial*, *Apoplexy*—*h. from the Intestines*, *Melæna*—*h. Local*, see *Hæmorrhagia*—*h. of the Mucous membranes*, see *Hæmorrhagia*—*h. Passive*, see *Hæmorrhagia*—*h. from the Pituitary membrane*, *Epistaxis*—*h. Pleural*, *Hæmatothorax*—*h. Post partum*, see *Post partum*—*h. Secondary*, see *Hæmorrhagia*—*h. from the Skin*, *Sudor cruentus*—*h. Spinal*, *Apoplexy*, spinal—*h. Spontaneous*, see *Hæmorrhagia*—*h. Supplementary*, see *Hæmorrhagia*—*h. Symptomatic*, see *Hæmorrhagia*—*h. of the Tissues*, see *Hæmorrhagia*—*h. Traumatic*, see *Hæmorrhagia*—*h. Uterine*, *Metrorrhagia*.

HEMORRHAG'IC, *Hæmorrhag'icus*, (F.) *Hæmorrhagique*. Relating or appertaining to hæmorrhage.

HEMORRHAGIC PLEURISY, *Hæmatothorax*.

HÉMORRHAGIE BUCCALE, *Stomatorrhagia*—*h. Cérébrale*, *Apoplexy*—*h. du Cœur*, *Hæmocardiorrhagia*—*h. du Foie*, *Hepatorrhagia*—*h. Intérielle*, *Apoplexy*—*h. Intérielle du Poupon*, *Hæmoptysis*—*h. des Intestins*, *Hæmatœchezia*—*h. de la Matrice*, *Metrorrhagia*—*h. Mningée*, *Apoplexy*—*h. de la Moëlle épinière*, *Apoplexy*.

plexia myelitica—*h. Nasale*, *Epistaxis*—*h. de la Vessie*, *Cystirrhagia*.

HEMORRHAGIP'AROUS, (F.) *Hæmorrhagipare*, (*hæmorrhagia*, and *parire*, 'to bring forth,') That which gives occasion to hæmorrhage: thus, softening of the neurine may be *hæmorrhagiparous*.

HÉMORRHÉE, *Hæmorrhœa*.

HÉMORRHINIE, (*hemo*, and *ῥιν*, 'the nose,')
Epistaxis.

HÉMORRHOÏDAIRE (F.). One who is subject to hæmorrhoids.

HEMORRHOÏDAL, *Hæmorrhoidal*.

HÉMORRHOÏDES, *Hæmorrhoids*—*h. Aveugles*, *Cæcæ hæmorrhoides*. See *Hæmorrhoids*.

HEMORRHOIDS, see *Hæmorrhoids*.

HÉMORRHOSCOPIE, (*hemo*, *ῥέω*, 'to flow,' and *σκοπεω*, 'to examine,') *Hæmatoscopy*.

HÉMOSPASIE, *Hæmospasia*.

HÉMOSPASIQUE, *Hæmospastic*.

HÉMOSTASIE, *Hæmostasia*.

HEMOSTATICS, *Hæmostatica*.

HÉMOSTATIQUES, *Hæmostatica*.

HEMP, INDIAN, (Sax. *hænep*.) *Apocynum cannabinum*, *Bangue*—*h. Seed*, see *Cannabis sativa*—*h. Wild*, *Ambrosia trifida*.

HEMPWEED, AROMATIC, *Eupatorium aromaticum*—*h. Roundleaved*, *Eupatorium rotundifolium*.

HEMURÉSIE, *Hæmaturia*.

HENBANE, *Hyoscyamus*.

HENBIT, *Lamium amplexicaule*.

HENBLINDNESS, *Hemeralopia*.

HENNA, *Lawsonia inermis*.

HENRICEA PHARMACEARCHA, (after R. S. Henrici, of Copenhagen,) *Gentiana chirayita*.

HENRICUS RUBENS, *Colcothar*.

HEP, (Sax. *heap*,) see *Rosa canina*.

HEP'AR, gen. *Hep'atis*, (*ἥπαρ*, gen. *ἥπατος*, 'liver,') A name for substances resembling liver in appearance. The ancient name for the liver of sulphur, *Hepar sulphuris*: which is sometimes a compound of sulphur and potassium; at others, of sulphur and potassa. See *Potassæ sulphuretum*.

HEPAR ADIPOSUM, *Adiposis hepatica*—*h. Adulterinum*, *Spleen*.

HEPAR ANTIMONIA'TUM is a compound of a sulphuret of antimony and an alkali. See *Oxydum stibii sulphuretum*.

HEPAR KALINUM, *Potassii sulphuretum*.

HEPAR MARTIA'LE, A compound of sulphuret of potassa and an oxide of iron.

HEPAR SANGUINIS, see *Blood*—*h. Sinistrum*, *Spleen*.

HEPAR SUCCENTRIA'TUM. An additional liver or liver lobe; a case of malformation.

HEPAR SULPHURIS, *Potassii sulphuretum*—*h. Sulphuris calcæreum*, *Calcis sulphuretum*—*h. Sulphuris volatile*, *Ammonis sulphuretum*—*h. Uterinum*, *Placenta*—*h. Variegatum*, *Liver nutmeg*.

HEPATAL'GIA, *Hepatodyn'ia*, *Neural'gia hep'atis*, *Colica hepatic*, (*hepato*, and *algos*, 'pain,') (F.) *Hepatalgie*. Pain in the liver. *Neuralgie of the liver*, (F.) *Néuralgie du Foie*.

HEPATALGIA CALCULOSA, *Colica hepatica*—*h. Pettitiana*, *Turgescencia vesiculæ felleæ*—*h. Phlegmonoides*, *Hepatitis*.

HEPATAPOSTEMA, gen. *Hepatapostematis*, (*hepato*, and *αστρομη*, 'an abscess,') Abscess of the liver.

HEPATATROPH'IA, *Atroph'ia* seu *Aridit'ra hep'atis*, (*hepato*, and *atrophia*, 'atrophy,') Atrophy of the liver:—a general concomitant of *Cirrho'ia hep'atis*.

HEPATAUX'Ë, *Hypertroph'ia hep'atis*, (*hepato*, and *αυξή*, 'increase,') (F.) *Hyperhépatistrophie*, *Hypertrophie du foie*. Hypertrophy of the liver.

HEPATECHEMA, gen. *Hepateche'omatis*, *Sonitus hepaticus*, (*hepato*, and *ηχημα*, 'sound,') Sound rendered by the liver on percussion.

HEPATOCE'LE, (*hepato*, and *ετλη*, 'a tumour,')

Hep'ia hepatica: *Hep'ia of the liver*. (F.) *Hépatie du foie*. The liver has never been found entirely out of the abdominal cavity. Increase of its bulk, or injuries of the parietes of the abdomen, have been the sole cause of the protrusions which have been occasionally met with, especially in infants, in whom the upper part of the linea alba is very weak, and indeed scarcely seems to exist. Sauvages has distinguished two species of hepatocèle:—the *ventral* (in the linea alba), and the *umbilical* or *hepatomphalum*.

HEPATOCHOLORRHEA, (*hepato*, *χολη*, 'bile,' and *ρῆω*, 'to flow,') *Hepatirrhœa*.

HEPATOCOLICUM. A ligament of the liver, described by Haller, as passing from the gall-bladder and contiguous sinus portarum, across the duodenum, to the colon. Another, termed *Hepatorenal*, descends from the root of the liver to the kidney. They are both peritoneal.

HEPATOCYSTIC, *Hepatocysticus*, *Cystidepaticus*. (F.) *Hépatocystique*, (*hepato*, and *κυστις*, 'bladder,') Relating to the liver and gall-bladder.

HEPATOCYSTIC DUCT, *Ductus hepatocysticus*. The choledoch duct.

HEPATODES, *Hepatic*.

HEPATODYME, *Hepatodymus*, (*hepato*, and *δύμος*, 'a twin,') A monster united at the liver, and double above.

HEPATODYNIA, (*hepato*, and *δύνη*, 'pain,') *Hepatalgia*.

HEPATODYSENTERIA, *Hepatirrhœa*.

HEPATOGASTRIC, *Gastrohepatic*.

HEPATOGASTROCHOLOSIS, (*hepato*, *γαστήρ*, 'the stomach,' and *χολη*, 'bile,') Fever, bilious, Fever, gastric.

HEPATOGRAPHY, *Hepatographia*, (*hepato*, and *γραφη*, 'a description,') The part of anatomy which describes the liver.

HEPATOHEMIA, *Hyperhemia hepatis*, *Hepatic Engorgement*, (F.) *Hypérémie du Foie*, (*hepato*, and *αἷμα*, 'blood,') Sanguineous congestion of the liver.

HEPATOIDES, ('*ηπατοειδής*, from *hepato*, and *εἶδος*, 'form,') *Hepatic*.

HEPATOLITHIASIS, (*hepato*, and *λίθιασις*, 'formation of stone,') The formation of concretions, *Hepatolithi*, in the liver.

HEPATOLOGY, *Hepatolog'ia*, (*hepato*, and *λογος*, 'a discourse,' 'treatise,') A treatise on the liver.

HEPATOMALACIA, (*hepato*, and *μαλακία*, 'softness,') *Malacosis hepatis*, (F.) *Ramollissement du Foie*. Softening of the liver.

HEPATOMPHALUS, (*hepato*, and *ομφαλος*, 'navel,') Hernia of the liver at the umbilicus.

HEPATOMYELOMA, gen. *Hepatomyelomatus*, (*hepato*, and *μυελος*, 'marrow,') *Hepatencephaloma*.

HEPATONCUS, (*hepato*, and *ογκος*, 'a tumour,') Tumefaction of the liver.

HEPATOPARECTAMA, gen. *Hepatoparectamatus*, (*hepato*, and *παρεκταμα*, 'considerable extension,') Excessive enlargement of the liver.

HEPATOPATHIA, (*hepato*, and *παθος*, 'affection,') *Liver disease*. Disease of the liver.

HEPATOPATHIE CANCÉREUSE, *Hepatoscirrhosis*—h. *Tuberculeuse*, *Hépatostromosie*.

HEPATOPHLEGMON, (*hepato*, and *φλεγμων*, 'inflammation,') *Hepatitis*.

HEPATOPHTHISIS, *Phthisis hepatica*, (*hepato*, and *φθις*, 'I consume,') Consumption from suppuration of the liver.

HEPATORRHAGIA, (*hepato*, and *ρῆξις*, 'rupture,') Rupture of the liver.

HEPATORRHŒA, *Hepatirrhœa*.

HÉPATOSARCOMIE, (*hepato*, and *sarcoma*,) *Hepatoscirrhosis*.

HEPATOSCIRRHUS, (*hepato*, and *σκίρρος*, 'cancerous induration,') *Scirrhosis hepatis*, *Enccephalosis* or *Carcinoma of the liver*, (F.) *Hépatoscarcome*, *Hépatopathie cancéreuse*, *Cancer du Foie*, *Scirrhosis* or *Cancer of the liver*.

HEPATOSTROMOSIE, *Tubercula hepatis*, (F.) *Tubercules du Foie*, *Hépatopathie tuberculeuse*; (*hepato*, and *stroma*, 'a tumour,' 'a scrofulous tumour,') Tubercle of the liver.

HEPATOTOMY, (*hepato*, and *τεμνω*, 'I cut,') Dissection of the liver.

HEPIALOS, (*ήπιαλος*,) *Epialos*.

HEPS, (pl. of *hep*,) see *Rosa canina*.

HEPSEMA, gen. *Hepsematis*, (*ήψημα*,) Decoction.

HEPSESIS, (*ήσις*,) Decoction.

HEPTAL CYCLE, see *Cycle*.

HEPTALION GRAVEOLENS, *Hogwort*, *Bears' freight*. An indigenous plant, which has a fetid porcine smell; and is said to be used by the Indians as a diaphoretic, cathartic, &c.

HEPTAPHARMACUM, (*ήπτα*, 'seven,' and *φάρμακον*, 'a remedy,') A medicine composed of seven substances; ceruse, litharge, pitch, wax, colophony, frankincense, and bullock's fat. It was regarded as laxative, suppurating and healing.

HEPTAPHYLLUM, (*ήπτα*, 'seven,' and *φύλλον*, 'a leaf,') *Tormentilla*.

HEPTAPLEURON, (*ήπτα*, 'seven,' and *πλευρον*, 'a rib,') *Plantago major*.

HERACLEUM, (dedicated to *Ηρακλεις*, 'Hercules,') see *H. lanatum*—h. *Branca*, *H. spondylium*.

HERACLEUM SPONDYLIVM, *H. Branca*, *Branca ursina* seu *Germanica*, *Spondylium*, *Sphondylium*, *Cow Parsnep*, *All'heal*, *Cowmumble*, (Sc.) *Buneward*, (Prov.) *Cushia*, (F.) *Berce*, *Braucersine bâtarde*, *Fausse Acanthe*. Order, *Umbelliferae*. Sex. Syst. *Pentandria Digynia*. The root of this plant has a strong, rank smell; and a pungent, almost caustic, taste. It has been given as a tonic, stomachic and carminative; both in powder and in decoction. The Russians, Lithuanians, and Poles obtain from its seeds and leaves, by fermentation, a very intoxicating spirituous liquor, which they call *Parst*.

The root of *Heracleum Lanatum*, *Masterwort*, *Cow parsnep*,—*Heracleum* (Ph. U. S.)—is in the secondary list of the Pharmacopœia of the United States.

HERACLEUS MORBUS, (*ήρακλειος*, 'herculean,') *Epilepsy*.

HERB, *Herba*, *Bot'anē*. A nonligneous plant, which dies down to the ground at least, in winter.

HERB, BLESSED, *Geum urbanum*—h. *Christopher*, *Actæa spicata*.

HERB-DOCTOR, *Botanical doctor* or *physic'ian*, *Root doctor*. One who treats diseases altogether by herbs, as the—so called—Thomsonian.

HERB OF GRACE, *Gratiola officinalis*—h. *Mastich*, common, *Thymus mastichina*—h. *Mastich*, Syrian, *Teucrium marum*—h. *Peter*, *Primula veris*—h. *Robert*, *Geranium Robertianum*—h. of Saint Cuneconde, *Eupatorium cannabinum*—h. *Sophia*, *Sisymbrium sophia*—h. *Trinity*, *Hepatica triloba*.

HERBA, *Herb*—h. *Alexandrina*, *Smyrnum olusatrum*—h. *Altheæ*, *Pelargonium cucullatum*—h. *Anthos*, *Rosmarinus*—h. *Benedicta*, *Geum urbanum*—h. *Britannica*, *Rumex hydrolapathum*—h. *Canni*, *Artemisia santonica*—h. *Cardiaca*, *Leonurus cardiaca*—h. *Cardui veneris*, *Dipsacus fullonum*—h. *Dorea*, *Solidago virgaurea*—h. *Felis*, *Nepeta*—h. *Genipi veri*, *Achillea atrata*—h. *Ignis*, *Lichen pyxidatus*—h. *Melancholifuga*, *Fumaria*—h. *Militaris*, *Achillea millefolium*—h. *Pa-*

pillaris, Lapsana—*h. Paralyseos*, Primula veris—*h. Paralytica*, Primula veris—*h. Patæ lapinæ*, Leonurus cardiaca—*h. Pulcaris*, Plantago psyllium—*h. Quercini*, Lichen plicatus—*h. Sacra*, Verbena officinalis—*h. Salvaris*, Anthemis pyrethrum—*h. Sardonica*, Ranunculus sceleratus—*h. Scelerata*, Ranunculus sceleratus—*h. Sideritidis*, Galeopsis grandiflora—*h. Tabaci*, Nicotiana tabacum—*h. Trinitatis*, Anemone hepatica, Viola tricolor—*h. Veneris*, Adiantum pedatum—*h. Ventis*, Anemone pulsatilla—*h. Vitri*, see Salsola kali—*h. Zazar-bendi*, Origanum.

HERBÆ PRO ENEMATE, Glyster herbs—*h. pro fotu*, Fomentation herbs.

HERBALIST, Herborist.

HERBARIUM, (from *herba*.) (F.) *Herbier*. A collection of plants. Generally applied to a collection of dried plants—*Hortus siccus*. In Pharmacy, a plant that is used entire.

HERBARIST, Herborist.

HERBARIUS, Herborist.

HERBE, Herb—*h. aux Abeilles*, Spiræa ulmaria—*h. au Cancer*, Plumbago Europæa—*h. au Centaure*, Chironia centaurium—*h. aux Chantres*, Erysimum—*h. aux Charpentiers*, Achillea millefolium, Justitia pectoralis, Sedum telephium—*h. aux Chats*, Nepeta, Teucrium marum—*h. à Chiron*, Chironia centaurium—*h. au Coq*, Tanacetum balsamita—*h. à la Conpure*, Sedum telephium—*h. au Cuillers*, Cochlearia officinalis—*h. aux Écouelles*, Scrophularia nodosa—*h. aux Écus*, Lysimachia nummularia—*h. à Éternuer*, Achillea ptarmica—*h. aux Femmes battues*, Tamus communis—*h. à la Fièvre*, Chironia centaurium—*h. de Grace*, Ruta—*h. aux Guex*, Clematis vitalba—*h. aux Hemorrhoides*, Ranunculus ficaria—*h. à l'Hirondelle*, Chelidonium majus—*h. à la Houette*, Asclepias Syriaca—*h. d'Iroque*, Lolium temulentum—*h. des Juifs*, Solidago virgaurea—*h. aux Ladres*, Veronica—*h. aux Mamelles*, Lapsana—*h. aux Mouches*, Conyza squarrosa—*h. aux Murailles*, Parietaria—*h. des Nones*, Parietaria—*h. aux Oies*, Potentilla reptans—*h. à la Paralyse*, Primula veris—*h. à Paris*, Paris—*h. à Pauvre homme*, Gratiola officinalis—*h. aux Perles*, Lithospermum officinale—*h. à Pisser*, Pyrola umbellata—*h. à la Poudre de Chypre*, Hibiscus abelmoschus—*h. aux Poux*, Delphinium staphisagria—*h. à la Puce*, see Apocynum hypericifolium—*h. aux Puces commune*, Plantago psyllium—*h. à Robert*, Geranium Robertianum—*h. à la Rosée*, Drosera rotundifolia—*h. de Sainte Barbe*, Erysimum barbarea—*h. Sainte Christophe*, Actæa spicata—*h. de Saint Étienne*, Cirsæa lutetiana—*h. Saint Fiacre*, Verbascum thapsus—*h. de Saint Jean*, Artemisia vulgaris—*h. Saint Jean*, Glechoma hederacea—*h. de Saint Quirin*, Tussilago—*h. de Saint Roch*, Inula dysenterica—*h. Sainte Rose*, Pæonia—*h. aux Sorciers*, Cirsæa lutetiana, Datua stramonium—*h. aux Teigneux*, Aretium lappa, Tussilago petasites—*h. à Tous les maux*, Verbena officinalis—*h. Turque*, Herniaria glabra—*h. aux Verrues*, Heliotropium Europæum—*h. aux Vers*, Tanacetum vulgare.

HERBIER, Herbarium, Ingluvies.

HERBIVOROUS, *Herbivorus*, (F.) *Herbivore*, (*herba*, and *voro*, 'I eat.') An epithet applied to animals which feed on herbs.

HERBORIST, *Herbarius*, (from *herba*.) One who deals in useful plants. An *Herbalist* or *Herbarist*.

HERBORIZATION, *Herbarium inquisit'*io. An excursion, made with the view of collecting plants. Such excursions are directed by the Apothecaries' Company of London, for the use of their apprentices, &c.

HERBS, FIVE CAPILLARY, *Quin'que her'bæ capillares*, were, anciently, hart's tongue; black, white, and golden maidenhair, and spleenwort.

HERBS, FIVE EMOLIENT, *Quin'que her'bæ emollientes*, were, anciently, beet, mallow, marshmallow, French mercury, and violet.

HERCULES ALLHEAL, *Pastinaca opoponax*. HERCULEUS MORBUS, (ἡρκλησος νοσος,) Epilepsy.

HERÉDITAIRE, Hereditary.

HEREDITARY, *Hæreditarius*, *Hæreditarius*, *Gentilitius*, *Syn'phytos*, *Syn'genes*, (F.) *Héréditaire*, (*hæres*, gen. *hære'dis*, 'an heir.') An epithet given to diseases, communicated from progenitors. This communication is termed, by the French, *Hérédité*. Such diseases may exist at birth; or they may supervene at a more or less advanced period of existence. *Hereditary diseases*, *Morbi hereditarii*, (F.) *Maladies héréditaires*, often prevail amongst several members of a family, or are family diseases or complaints.

HERÉDITÉ, see Hereditary.

HERMAPHRODITE, *Hermaphroditus*, *Hermaphroditus*, *Hermaphroditus*, *Fab'rica androgyn'a*, *Androgyn'ia*, *Gynand'ria*, *Hermaphroditism*, *Hermaphroditism*; (from *Ἑρμης*, 'Mercury,' and *Ἀφροδίτη*, 'Venus.') Appertaining to Mercury and Venus. Union of the two sexes in the same individual.

HERMAPHRODISIA, Hermaphroditia.

HERMAPHRODITE, *Hermaphroditus*, *Gyn'ida*, *Androg'ynus*, (Sc.) *Searcht*, (Prov.) *Serat*. One who possesses the attributes of male and female: who unites in himself the two sexes. A term, applied to an animal or plant which is, at the same time, both male and female. True hermaphrodites are only met with in the lower degrees of the animal scale, amongst the zoophytes, mollusca, or gasteropoda. The individuals of the human species, regarded as hermaphrodites, owe this appearance to a vicious conformation of the genital organs; a kind of monstrosity, which renders them unfit for generation, although an attentive examination may exhibit the true sex. Hermaphrodites have, likewise, been described, which, instead of uniting the attributes of both sexes, cannot be considered male or female. These have been called *neutral hermaphrodites*.

HERMAPHRODITIC, *Hermaphroditicus*, *Androg'ynus*, (F.) *Hermaphroditique*. Relating or appertaining to an hermaphrodite.

HERMAPHRODITISMUS, Hermaphroditia.

HERMAPHRODITUS, Hermaphrodite.

HERMETICA DOCTRINA, *Hermética ars*, (F.) *Hermétique*. The doctrine of Hermes, a celebrated Egyptian philosopher, who is considered the father of alchemy. That part of chymistry, whose object was the pretended transmutation of the metals.

HERMODACTYLUS, *Dactyle'tus*, *An'ima articulo'rum*, ('Ἑρμης, 'Mercury,' and *δακτυλος*, 'a finger;' or rather from *Hermus*, a river in Asia, upon whose banks it grows, and *δακτυλος*, 'a date;' or from *Ἑρμης*, 'Mercury,' and *δακτυλος*, 'a date;') (F.) *Hermodactyle*, *Hermodacte* on *Hermodate*. The root of the *Hermodactyl* was formerly used as a cathartic. By some, it is supposed to be identical with the *Iris tuberosa*. The best testimony seems to be in favour of its being a variety of the colchicum, — *Colchicum Illyricum*. Under the name *Hermodactyla*, roots or bulbs of an unknown plant, growing in the countries about the eastern extremity of the Mediterranean, are sold in the shops of Europe.

HERMODATE, Hermodactylus.

HERMOPHILUS, ('Ἑρμης, 'Mercury,' and *φίλος*, 'a lover.') One who is fond of mercury as a medicine.

HERNIA, (from *ἥρως*, a sprout, 'a shoot,' [?], *Ru'mex*, *Rup'tura*, *R. herni'osa*, *Crepas'tra*, *Ectop'ia herni'osa*, *Cel'se*, *Rup'ture* *Buxa*, (F.) *Hernie*, *Hargue*, *Descente*, *Effort*, *Greveure*,

Rompure. Any tumour, formed by the displacement of a viscus or a portion of a viscus, which has escaped from its natural cavity by some aperture, and projects externally. Herniæ have been divided into,—1. Hernia of the Brain; Encephalocele; 2. Hernia of the Thorax; Pneumocele; 3. Hernia of the Abdomen.

Abdominal Herniæ are remarkable for their frequency, variety, and the danger attending them. They are produced by the protrusion of the viscera, contained in the abdomen, through the natural or accidental apertures in the parietes of that cavity. The organs, which form them most frequently, are the intestines and the epiploon. These herniæ have been divided, according to the apertures by which they escape, into:

1. *Inguinal or Suprapubic Herniæ*. These issue by the inguinal canal: they are called *Bubonocèle*, when small; and *Sero'tal Hernia* or *Osciocele*, in man, when they descend into the serotum:—*Vulvar Hernia* or *Pudendal* or *Labial Hernia*, *Episiocele*, in women, when they extend to the labia majora. 2. *Crural or Femoral Hernia*, *Mérocèle*, when they issue by the crural canal. 3. *Infrapubic Hernia*, (*F.*) *Hernie sous-pubienne*, *Ooëcocele*, *Opeocèle*, *Opodocèle*, *Her'nia foran'tis ova'lis seu obturatoria*, when the viscera escape through the opening, which gives passage to the infra-pubic vessels. 4. *Ischiatic* or *Sciatic Hernia*; when it takes place through the sacro-sciatic notch. 5. *Umbilical Hernia*, *Exomphalos*, *Omphalocèle*; when it occurs at the umbilicus or near it. 6. *Epigastric Hernia*;—occurring through the linea alba, above the umbilicus. 7. *Hypogastric* or *Infräumbilical Hernia*, *Cæliocèle*, *Hypogastrocèle*,—when it occurs through the linea alba below the umbilicus. 8. *Perineal Hernia*, *Mesocelocèle*, *Hernia perinæ*, *Perineocèle*, *Perineocèle*,—when it takes place through the levator ani, and appears at the perineum. 9. *Vag'inal Hernia*, *Coleocèle* seu *Elytrocèle*—through the parietes of the vagina. 10. *Diaphragmatic Hernia*, *Diaphragmatocèle*; when it passes through the diaphragm.

Herniæ are likewise distinguished,—according to the viscera forming them,—into *Enterocèle*, *Epiplocèle*, *Entéroépiplocèle*, *Gastrocèle*, *Cystocèle*, *Hepatocèle*, *Splenocèle*, &c.

When a hernia can be restored to its natural cavity, by the aid of pressure, &c., properly applied, it is said to be *reducible*. It is, on the contrary, *irreducible*, when adhesion, bulk, &c., oppose its return. When the aperture, which has given passage to the hernia, occasions more or less constriction of the protruded portion, the hernia is said to be *incarcerated* or *strangulated*; and, if the constriction be not removed, constipation, hiccough, vomiting, and all the signs of violent inflammation, followed by gangrene, supervene, with alteration of the features, small pulse, cold extremities, and death.

The therapeutical indications are,—1. *As regards reducible hernia*:—to replace the viscera in the abdomen by the taxis; and to retain them there by the use of a *truss*, which, if properly adapted, may effect a radical cure. 2. *As regards irreducible hernia*:—to support the tumour by an appropriate suspensory bandage. 3. *As regards strangulated hernia*:—to have recourse to the taxis; bloodletting; warm bath; tobacco glysters; ice to the tumour; and, if these should not succeed, to perform an operation, which consists in dividing the covering of the hernia, and cutting the aponeurotic ring, which causes the strangulation;—reducing the displaced viscera, unless their diseased condition should require them to be retained without;—dressing the wound appropriately:—restoring the course of the fæces by means of gentle glysters—preventing or combating inflammation of the abdominal viscera;—conduct-

ing the wound to cicatrization, by appropriate means; and afterwards supporting the cicatrix by a bandage.

The word *hernia* was also used, of old, for the serotum, and, not unfrequently, for the testicle.

HERNIA, ANEURISMAL, Cirsomphalus—*h.* Arteriarum, see Aneurism—*h.* of the Bladder, Cystocele—*h.* Bronchialis, Bronchocele—*h.* Carnosa, Sarcocèle—*h.* of the Cerebellum, Parencephalocele—*h.* Cerebri, Encephalocele.

HERNIA, CONGENITAL, *Her'nia congen'ita*, is a protrusion of some of the contents of the abdomen into the *Tunica vaginalis testis*, owing to a want of adhesion between its sides, after the descent of the testicle.

HERNIA CORNEÆ, Ceratocèle—*h.* Crural, Merocele—*h.* Epiploic, Epiplocèle—*h.* Femoral, Merocele—*h.* Foraminis ovalis, see Hernia—*h.* Gutturis, Bronchocele—*h.* Hepatica, Hepatocèle.

HERNIA HUMORALIS, *Empres'ma Orchitis*, *Didymitis*, *Orchitis*, *Orchiditis*, *Testitis*, *Inflammatio testium*, *Inflammation of the Testicle*, *Swelled Testicle*, *Her'nia Ven'eri*, *Orchiocele*, *Orchidocèle*, *Orchidon'cus*, (*F.*) *Inflammation du testicule*, *Orchite*. Swelling and inflammation of the testicle is a common symptom of gonorrhœa, (*F.*) *Chandepisse tombée dans les Bourses*; but it may arise from external injuries, or from other causes. It is a disease which cannot be mistaken, and the treatment must obviously be strongly antiphlogistic, supporting the testicle during the treatment, and for some time afterwards; methodical compression has also been found useful. The disease is not generally of a serious character, going off as suddenly as it comes on. As it affects the epididymis more especially, when supervening on gonorrhœa or blennorrhœa, it is sometimes termed *blennorrhag'ic epididymitis*.

HERNIA INCARCERATA, Enteropristole—*h.* Inguinalis, Bubonocèle—*h.* Intestinalis, Enterocèle—*h.* Intestinalis omental, Enteropileocèle—*h.* Iridis, Ptosis Iridis—*h.* Ischiatica, Ischiocele—*h.* Lienalis, Splenocèle.

HERNIA LITT'ERICA, *H. e Divertic'ulo Intesti'ni*. Hernia first described by Littre, in which the intestinal canal proper is not included in the hernial sac, the protruded portion of intestine consisting of a digital prolongation of the ileum, which Littre concluded was formed by the gradual extension of a knuckle of the bowel, that had been engaged in the inguinal canal.

HERNIA OF THE LIVER, Hepatocèle—*h.* of the Lung, Pneumocele.

HERNIA, MESOCOLIC, *Her'nia Mesocolica*. Hernia, caused by the bowels gliding between the layers of the mesocolon.

HERNIA OBTURATORIA, Hernia infra-pubic—*h.* Omental, Epiplocèle—*h.* Parorchidoenterica, Parorchido-enterocèle—*h.* Perinæ, see Hernia—*h.* Pharyngis, Pharyngocele—*h.* Phrenic, Diaphragmatic hernia—*h.* Pinguedinosæ scroti, Liparocèle—*h.* of the Pleura, Pleurocele—*h.* Pleurica et pulmonalis, Pleurocele—*h.* Pudendal, Pudendal hernia—*h.* Sacci lacrymalis, Lacrymal hernia, Mucocele—*h.* Sciatic, see Hernia—*h.* Scrotalis, Scrotocèle—*h.* Seminalis scroti, Spermatocèle—*h.* of the Stomach, Gastrocele—*h.* Suprapubic, *H.* inguinal—*h.* of the Tongue, Glossocèle—*h.* Umbilicalis, Epiplophalon, Exomphalos—*h.* Umbilici aquosa, Hydromphalum—*h.* Urachi, Uromphalus—*h.* Uteri, Hysterocele—*h.* Varicosa, Varicocele—*h.* Varicose, Cirsocèle—*h.* Venarum, Varix—*h.* Veneris, Hernia humoralis—*h.* Ventosa, Physocèle—*h.* Ventral, see Ventral—*h.* Ventriculi, Gastrocele—*h.* Vesicæ urinariæ, Cystocèle—*h.* Zirbalis, Epiplocèle.

HERNIAIRE, Hernial, Herniaria glabra.

HERNIAL, *Hernia'rius*, *Her'nious*, (*F.*) *Herniaire*. Belonging to, or concerning hernia:—as *Hernial Bandage*, *Hernial Sac*, &c.

(F.) *Hernië*. *Herniated*, is applied to a part enveloped in a hernial sac.

(F.) *Hernieux*, *Hernieux*, *Celeste*, *Ruptured*, *Durst*, *Burston*, means one affected with hernia; and *Chirurgien herniaire*, one who devotes himself to the treatment of hernia.

HERNIA'RIA GLA'BRA. *H. vulga'ris* seu *hirsu'ta* seu *alpes'tris* seu *an'ua* seu *cine'rea* seu *frutico'sa*. *Miliogra'na*, *Em'petrum*, *Rup'turewort*, from *hernia*, which it was supposed to cure.)

(F.) *Herniaria*, *Turquette*, *Herniole*, *Herbe Turque*. *Ord. Paronychiæ*. This plant, which as its name imports, was formerly considered efficacious in the cure of hernia, seems destitute of all virtues.

HERNIARIUS, Hernial.

HERNIATED, see Hernial.

HERNIE. Hernia — *h. Charnæ*, *Sarcocele* — *h. de l'Épiploûm*, *Epiplocele* — *h. de l'Estomac*, *Gastrocele* — *h. du Foie*, *Hepatocele* — *h. Inguinale*, *Bubonocèle* — *h. du Nomb'il*, *Exomphalos* — *h. Om'bilicale*, *Exomphalos* — *h. du Poymon*, *Pneumocèle* — *h. de l'Utérus*, *Hysterocele* — *h. de la Vessie*, *Cystocèle*.

HERNIÉ, Hernial.

HERNIEMPHRAG'MUS, *Herniemp'hrax'is*, (*hernia*, and *ἐμπραγος*, 'obstruction.') The mechanical obstruction of a hernial canal for the radical cure of the hernia.

HERNIEUX, see Hernial.

HERNIOLE, Herniaria glabra.

HERNIOTOMY, (*hernia*, and *τομή*, 'to cut,') Celotomia.

HERNIOUS, Hernial.

HERO'IC, *Hero'icus*, *Hero'ius*, *Hero'us*. (*hōs*, 'a hero.') An epithet applied to remedies or practice of a violent character.

HERPEDON. (*ἑρπεδον*.) Herpes.

HERPEN. (*ἑρπην*.) Herpes.

HERPES, gen. *Herp'etis*, (*ἑρπης*, genitive, *ἑρπητος*.) *E'rpēs*, *Herpe'don*, *Her'pen*, *Ser'pens*. (*ἑρπω*. 'I creep,' because it creeps and spreads about the skin; *Ec'phlysis* *Her'pes*, *Cytis'ia* *Her'pes*, *Omph'lysis*, *Tet'ter*, *Fret*, (F.) *Herpēs*, *Dartre*. A vesicular disease, which, in most of its forms, passes through a regular course of increase, maturation, decline, and termination, in from 10 to 14 days. The vesicles arise in distinct, but irregular clusters, which commonly appear in quick succession, and near together, on an inflamed base; generally attended with heat, pain, and considerable constitutional disorder. The term, like most others which refer to cutaneous diseases, has not been accurately defined. The ancients had three varieties: the *mili'ary*, *xy-chiaios*; *vesic'ular*, *φλυκταίνων*, and *ero'ding*, *εξδιω-mus*. Bateman has the following varieties: 1. HERPES PHLYCTENO'DES, *Her'pes milia'ria*, *Ec'phlysis* *Her'pes milia'ria*, *Nirles*, (F.) *Dartre phlyctenoïde*, in which the vesicles are millet-sized; pellucid; clusters commencing on an uncertain part of the body, and being progressively strewed over the rest of the surface; succeeded by fresh crops. 2. HERPES ZO'STER, *Zos'ter*, *Zo'na* *ig'nea* seu *serpigin'o'sa*, *Ig'nis* *Per'sicus*, *Cinzill'a*, *Cin'gulum*, *Sa'cer* *ig'nis*, *Ec'phlysis* *Her'pes zos'ter*, *Her'pes peris'celia*, *Ery'sip'elas* *zos'ter* seu *phlycta-noï'des* seu *pustulo'sa*, *Zo'na*, *Cir'cinus*, *Perizo'na*. (F.) *Ceinture*, *C. dartreuse*, *Feu* *Pereique*, *Ery-sipèle* *pustuleux*, *Shingles*, (Prov.) *Cruela*, in which the vesicles are pearl-sized; the clusters spreading round the body like a girdle; at times confluent, and occasionally preceded by constitutional irritation. 3. HERPES CIRCINA'TUS, *Formi'ca* *ambu-lato'ria*, *An'ulus* *re'pens*, *Her'pes* *Serpi'go*, *Serpi'go*, *Ec'phlysis* *Her'pes* *Circina'tus*, *Ring'worm*, *Vesic'ular* *Ring'worm*, (F.) *Herpès* *circiné*, consisting of vesicles with a reddish base, uniting in rings; the area of the rings slightly discoloured,

often followed by fresh crops. 4. HERPES LABIA'LIS, and 5. HERPES PRÆPUTIA'LIS, *Ap'h'is* *præputi'i*, *Ulcus'cula* *præputi'i*, appearing, respectively, on the lips and prepuce. 6. HERPES I'RIS, *Pris*, *Ruin'ous* *worm*, occurring in small circular patches, each of which is composed of concentric rings of different colours. To these may be added, HERPES EX'EDENS, *H. esthiom'enus* seu *depas'cens* seu *fe'rus* seu *esthiom'enus* seu *fe'rox*, *Ec'phlysis* *Her'pes* *ex'edens*, *Dart'a* *excoriati'ca* seu *malig'na*, *Lu'pus* *non* *ex'edens*, *Vitili'go* (?), *Leucè* (?), *Formi'ca* *corro'siva*, *For'mix*, *Pap'ula* *fe'ra*, *Ulc'erative* *Ring'worm*, *A'gria*, (F.) *Dartre* *rongeante*, in which the vesicles are hard; clusters thronged; fluid dense, yellow or reddish, hot, acrid, corroding the subjacent skin, and spreading in serpentine trails.

All the varieties demand simply an antiphlogistic treatment, when attended with febrile irritation. The *herpes circinatus*, alone, requires the use of astringent applications which have the power of repressing the eruption.

HERPES CIRCINATUS, see Herpes — *h. Depascens*, *H. exedens*, see Herpes — *h. Esthiomenus*, *H. exedens*, see Herpes — *h. Esthiomenus*, *H. exedens*, see Herpes — *h. Exedens*, see Herpes — *h. Farinosus*, *Pityriasis* — *h. Ferox*, *H. exedens*, see Herpes — *h. Ferus*, *H. exedens*, see Herpes — *h. Furfuraceus*, *Pityriasis* — *h. Furfuraceus* *circinatus*, *Lep'ra*, *H. exedens* — *h. Guttural*, *Pharyngitis*, *diphtheritic* — *h. Iris*, see Herpes — *h. Labialis*, see Herpes — *h. Miliaris*, *H. phlyctænodes*, see Herpes — *h. Pericælis*, *H. zoster* — *h. Phlyctænodes*, see Herpes — *h. Præputialis*, see Herpes — *h. Serpigo*, *H. circinatus* — *h. Tonsurans*, *Porrig'o* *decalvans* — *h. Zoster*, see Herpes.

HERPET'IC, *Herp'et'icus*, *Her'patus*, (*ἑρπετος*.) (F.) *Herpétique*, *Dartreux*. Possessing the nature of herpes.

HERPETOGRAPHY, *Herpetogra'phia*, (*herpes*, and *γραφία*, 'a description.') A description of the different forms of herpes, as *Herpetology*, *Herpetolog'ia*, is a treatise on the same; and also on reptiles, (from *ἑρπετος*, 'a reptile'.)

HERPETON, (*ἑρπετον*.) *Herpet'icon*, (*ἑρπετις*, 'to creep.') A creeping eruption or ulcer.

HERPYLOS, (*ἑρπελος*.) *Thymus* *serpyllum*.

HERRENSCHWAND'S SPECIFIC, see Specific of Herrenchwand.

HERRIN (Prov.), Urine.

HERSKET, (Heartscauld,) Ardor ventriculi.

HESPERIS ALLIARIA, (*ἑσπερίς*, 'evening,') Alliaria.

HETERADELPH'IA, *Heterodidym'ia*, *Heterodyn'm'ia*, (*hetero*, and *ἀδελφός*, 'a brother.') A double monstrosity, in which the components of the double bodies are very unequal, and of which one portion may be regarded as the stem or trunk, to which another organized part, or even a whole body, less developed than itself, is affixed as a parasite.

HETERADEN'IC, *Heteraden'icus*, (F.) *Hétéradénique*, (*hetero*, and *ἀδην*, 'a gland'.) An accidental tissue is so termed, which has a glandular structure, but occurs in parts devoid of glands.

HETERO, (*ἕτερος*, 'other,') in composition, other.

HETEROCHRON'IA, *Aberra'tio* *tem'poris*, (*hetero*, and *χρῶς*, 'time'.) A deviation as to time in the formation or action of parts.

HETEROCHRONICUS, *Heteroch'ronus*, (*hetero*, and *χρῶς*, 'time'.) Relating to difference of time.

Pulsus heterochron'icus. A pulse of varying rhythm. An irregular or intermittent pulse.

HETEROCHYMEUSIS, (*hetero*, and *χυμῶσις*, 'mixture'.) A state of the blood in which it contains other matters than in health, as urea, bile, &c.

HETEROCLITE, (*ἑτεροκλίτης*, from *hetero*, and *κλίω*, 'I bend,') see Homology, and Tissues.

HETEROCRANIA, (*hetero*, and *cranium*,) Hemierania.

HETEROCRIN'IA, (*hetero*, and *κρίνω*, 'I separate,') Modification in the situation of secretions.

HETERODIDYMA, (*hetero*, and *δίδυμος*, 'a twin,') Heteradelphia.

HETERODYMIA, Heteradelphia.

HETEROGENESIS, (*hetero*, and *γενεσις*, 'generation,') see Generation. Also, any anomaly in organic formation.

HETEROGLA'LIA, (*hetero*, and *λαλέω*, 'to speak,') Heterophonia.

HETEROLOGOUS TISSUES, (*hetero*, and *λογος*, 'a condition,') see Tissues.

HETEROMET'RIA, (*hetero*, and *μετρον*, 'a measure,') A variation in the quantity of parts.

HETEROMORPHISM, *Heteromorphis'mus*, (*hetero*, and *μορφή*, 'shape,') A deviation from the natural shape of parts.

HETEROMORPHOUS, see Homology.

HETEROPA'GIA, (*hetero*, and *παγω*, 'I unite together,') A double monstrosity, in which the accessory subject, which is very small and imperfect, but provided with a distinct head, and at least rudimental lower limbs, has its body implanted in the anterior surface of the body of the principal subject.

HETEROPATHIC, (*hetero*, and *πάθος*, 'affection,') Allopathic.

HETEROPHO'NIA, *hetero*, and *φωνή*, 'voice,') A cracked or broken voice. A change of the voice or speech—*Heterola'lia*.

HETEROPHTHAL'MIA, (*hetero*, and *ὄφθαλμος*, 'eye,') A difference in the two eyes,—as when one squints, or is of a different colour.

HETEROPLA'SIA, (F.) *Hétéroplasie*, (*hetero*, and *πλασσω*, 'to form,') Same etymon as the next. The formation of a heteroplasm.

HETEROPLASM, *Heteroplasm'us*. A morbid formation foreign to the economy, and which is composed of elements different from those that are found in it in the normal condition.

HETEROPLAS'TY, *Heteroplast'icē*, *Heteroplas'tia*. Irregular plastic or formative actions or operations, that do not admit of exact classification.

HETEROPROSO'PUS, (*hetero*, and *πρῶσπον*, 'countenance,') A monster having two faces.

HETEROPUS, (*hetero*, and *πους*, 'foot,') One who has one foot different from the other.

HETEROREXIA, (*hetero*, and *ορέξις*, 'appetite,') Malacia.

HETERORRHYTH'MUS, (*hetero*, and *ῥυθμός*, 'rhythm,') Having another rhythm. An epithet given to the pulse, when it is such, in any individual, as is usually felt at a different age.

HETEROSARCO'SES, (*hetero*, *σαρξ*, 'flesh, and *οσίς*,) A class of diseases which consist in the formation of accidental tissues.

HETEROTAX'IA, (F.) *Hétérotaxie*, (*hetero*, and *τάξις*, 'order,') A malformation, which consists in the general transposition of organs. A change in the relation of organs.

HETEROTOP'IA, *Aheterotio lo'ci*, (F.) *Hétérotopie*, (*hetero*, and *τοπος*, 'place,') A deviation from the natural position of parts. This has been called by M. Gueniot, *objet'ine heterotopia* in contradistinction to *subject'ine heterotopia*, which he applies to a sensation of displacement or approximation; such as is experienced in the stump of an amputated limb.

HÉTRE, *Fagus sylvatica*.

HEUCHERA, (after Heucher, a professor and botanist at Wittenberg,) see *H. cortusa*—*h. Acerifolia*, *H. cortusa*.

HEUCHERA CORTU'SA, *H. America'na* seu *Acerifolia* seu *Visc'ida*, *Cortu'sa America'na*, *Al'um*

Root, American Santicle, Groundmaple, Cliffweed, Splitrock, Heuchère d'Amérique. Ord. Saxifragaceæ. The root—*Heuchera* (Ph. U. S.)—is a powerful astringent, and is the basis of a *cancer powder*. The American Indians apply the powdered root to wounds, ulcers, and cancers. It is said to have been sold for colchicum.

HEUCHERA RICHARDSON'II, which abounds on the rocks of Churchill River, is used medicinally by the Cree Indians, who chew its astringent root, and apply it as a vulnerary to wounds and sores.

HEUCHERA VISCIDA, *H. cortusa*.

HEUCHÈRE, *Heuchera*.

HEUCK-BANE, (Sc.) *Ischion*.

HEUDELOTIA AFRICANA, (after Heudelot, a garden director on the Senegal,) see *Bdellium*.

HEUKS (Prov.), (by onomatopoeia,) *Singultus*.

HEVEA GUIANENSIS, see *Caoutchouc*.

HEXAGIUM, (*ἑξαγων*, from *ἑξ*, 'six,') *Sextula*.

HEXAGONE ARTÉRIELLE, (*ἑξ*, 'six, and *γωνία*, 'an angle,') See *Circulus Willisii*.

HEXATHYRIDIIUM VENARUM, (*ἑξ*, 'six, and *θύρα*, 'a door, *θυρίδιον*, 'a little door,') See *Worms*.

HEXIS, (*ἕξις*,) Habit, habit of body, constitution. Hence, hectic, cachectic, &c.

HEYLE (Sc.), ([Sax.] *hæl*,) *Sanitas*.

HHABB, see *Bouton d'Alep*.

HIA TSAO TONG TCHONG, *Sphæria Sinensis*.

HIA'TUS, (*hiare*, *hiatum*, 'to gape, 'to open,') A foramen or aperture. Mouth. The vulva. Also, yawning.

HIATUS CANALIS FALLOPII, see *Aquæductus Fallopii*.

HIATUS DIAPHRAG'MATIS AOR'TICUS, *Semicirculus excuscul'tus*. The opening in the diaphragm for the passage of the abdominal aorta.

HIATUS FALLOPII, see *Aquæductus Fallopii*—*h. Occipito-pétreus*, *Lacerum posterius foramen*—*h. Sphéno-pétreus*, *Lacerum anterius foramen*.

HIATUS OF WIN'SLOW, *Fora'men Winslow'ii*, *Fora'men of Winslow*. An opening—situate behind the lesser omentum, and behind the vessels and nerves of the liver—which forms a communication between the peritoneal cavity and that of the omenta.

HIBERNATION, *Hiberna'tio*, *Wintersleep*, (F.) *Sommeil hivernal*, (from *hibernare*, 'to winter,') The state of torpidity in which certain animals pass the winter;—the organic actions being reduced so low that animation seems to be suspended.

HIBERNICUS LA'PIS, *Teg'ula seu Arde'sia* ([F.] *Ardoise*, 'slate,') *Hibernica*, *Harde'sia Irish State*. A kind of slate or very hard stone, found in different parts of Ireland, in masses of a bluish-black colour, which stains the hands. It has been taken, powdered, in spruce-beer, against inward contusions.

HIBISCUS, (*ἱβίσκος*,) *Althæa*.

HIBIS'CUS ABELMOS'CHUS, *Abelmos'chus*, *A. moscha'tus*. Ord. Malvaceæ. The name of the plant, whose seeds are called *Gra'na Mosch'i* or *Musk'-seed*. It is the *Belmus'chus*, *Abelmos'chus*, *Gra'nium Mos'chi*, *Ket'mia Ægypti'aca*; *Mosch'us Arabum*, *Ægypt'ia moscha'ta*, *Ba'mia moscha'ta*, *Al'cea*, *A. Ind'ica seu Ægypti'aca Vill'o'sa*, *Abelmosch*, *Abelmusk*, *Muskmulloo*, (F.) *Graine de Musc*, *Herbe à la poudre de Chypre*, *Ambrette*, *Gnimave reloutée*. It is indigenous in Egypt and the Indies. The seeds are chiefly used as perfumes; and especially in the formation of *Cyprus Powder*.

HIBISCUS ESCULENT'US, *O'kra*, *O'kro*, *Gom'bo*. A plant, the green pods of which abound in mucilage, and are much used in the West Indies for soups. In Constantinople, and other portions of

the East, the root is employed as a demulcent like *Aitha*.

HIBISCUS POPULE'US, *Balimbaga*. A small Molucca tree. The fruit is full of a juice similar to camboge. The root is emetic. It is used in chronic diarrhoea, colic, dyspepsia, &c.

HIBRIDA, Hybrid.

HICCOUGH, ([G.] Hick, and Hickkop, by onomatopœia.) Singultus.

HICCUP, Singultus.

HICKET, Singultus.

HICK'ORY. The name of several American trees of the genus *Carya*; Order, Juglandaceæ. The leaves are usually aromatic; and are reputed to be antispasmodic [?]. The bark of those species that have bitter nuts, as *Carya amara* and *C. porcin*a, is somewhat astringent. Some of them bear fruit that is much esteemed, as *Carya aliformis*, *Pecan* or *Pecan*' nut, and *C. sulcata*, and *C. alba*, *Shellbark* or *Shagbark*. The *Hickory nut* is called, also, *Kiskitomas nut*.

HICKOT, Singultus.

HICQUET, Singultus.

HIDDEN SEIZURES. An expression employed by Dr. Marshall Hall for obscure encephalic and spinal attacks, of an epileptoid character for example, which may be immediately owing to trachelismus.

HIDEBOUND, Induration of the cellular tissue.

HIDRISCHESES, Hidroschesis.

HIDROA, (*ιδρωα*, from *hydros*), Desudatio, *Échauboulures*, Hydroa, Sudamina.

HIDRON'OSUS, *Hidronus*'us, (*hidro*, and *nosos*, 'a disease.') A disease accompanied by violent sweats. Sudor Anglicus.

HIDRONUSUS, *Hidronosus*.

HIDROPEDESIS, (*hidros*, and *πηδω*, 'to bound.') Ephidrosis.

HIDROPHORUS, (*hidros*, and *φέρω*, 'to carry.') Sudoriferous.

HIDROPOETICUM, (*hidros*, and *ποιεω*, 'to make.') Sudorific.

HIDROPYRÁ, (*hidros*, and *πυρ*, 'fever.') Sudor Anglicus.

HIDROPYRETUS, (*hidros*, and *πυρεος*, 'fever.') Sudor Anglicus.

HIDORRRHŒA, (*hidros*, and *ρῶω*, 'to flow.') Ephidrosis.

HIDROS, (*ιδῶος*, gen. *ιδῶρος*.) Sudor, 'sweat.'

HIDROS'CHESIS, *Hidris*'chesis, *Retentio sudoris*, (*hidros*, and *ἔχεις*, 'retention.') Suppression of perspiration.

HIDROSIS, (*ιδῶσις*, from *hidros*.) Ephidrosis.

HIDROTTERION, (from *hidros*.) Achicolum.

HIDROTTERIUM, Sudorific.

HIDROTICUM, Sudorific.

HIDROTIQUE, Sudorific.

HIDRO'TIUM, (dim. of *hidros*.) A gentle sweat or perspiration.

HIDROTOPEUM, (*hidros*, and *ποιεω*, 'to make.') Sudorific.

HIEBLE, (from *ebulus*, 'the elder.') Sambucus ebulus.

HIERA OF COLOCYNTH, *H. Diacolocynthidos*.

HIERA DIACOLOCYN'THIDOS, (*hieros*, 'holy.') *Hiera* of *Colocynth*. An electuary, composed of 10 parts of colocynth, as much agaric, germander, white horehound, stœchas:—5 parts of *opopanax*, as much *sagapenum*, parsley, round birthroot root, and white pepper:—4 parts of spikenard, cinnamon, myrrh, and saffron; and 3 pounds, 3 ounces, and 5 drachms of honey.

HIERA LOGADII, *Hiera* piera—*h. Nosos*, (*hieros*, 'holy disease,') Epilepsy.

HIERA PICRA, (*hieros*, 'holy,' and *πικρος*, 'bitter.') *Holy bitter*. *Pulvis aloëticus*, formerly called *Hiera logadii* when made into an electuary with

honey. It is now kept in the form of dry powder;—prepared by mixing *socotrine aloes*, one pound, with 3 ounces of *canella alba*. See *Pulvis aloës cum canella*.

HIERA SYRINX, (*ἱέρα συρίγξ*, 'holy tube,') Vertebral column.

HIERACI'TES, (*hieracins*, from *ἱέραξ*, gen. *ἱέρακος*, 'a hawk,') *La'pis Accipitrum*. The ancient name of a precious stone, believed to be capable of arresting the hemorrhoidal flux.

HIERACIUM LACHENALII, (*hieracium*, 'hawkweed,' from *ἱέραξ*, gen. *ἱέρακος*, 'a hawk,') *H. murorum*.

HIERACIUM MURO'RUM, *H. Lachena'lii*, *Pulmonaria Gallica*, *Auric'ula mu'ris ma'ior*, (F.) *Épervière des murailles*, *Pulmonaire des Français*. Ord. Composite. A European plant, which is a slight tonic.

HIERACIUM OLERACEUM, *Sonchus oleraceus*.

HIERACIUM PILOSEL'LA. The systematic name of the *Auric'ula Mu'ris*, *Pilosel'la*, *R. Alpi'na*, *Myoso'tis*, *Mousse-ear*, (F.) *Piloselle*, *Oreille de Souris*. This plant contains a bitter, lactescent juice, which has a slight degree of astringency. The roots are more powerful than the leaves.

HIERACIUM VENO'SEM, *Rattelnakeveel*, *Veiny Hawkweed*, *Bloodwort*, indigenous, has similar properties.

HIERANOS'US, (*hieros*, 'holy disease,') Chorea, Convulsion, Epilepsy.

HIERAX, (*ἱέραξ*, gen. *ἱέρακος*, 'the sparrowhawk,') Accipiter.

HIEROBOTANE, (*ἱεροβοτανν*, from *hieros*, 'holy,' and *βοτανν*, 'herb,') *Verbena officinalis*.

HIEROGLYPH'ICA, (*ἱερογλυφικά*, from *hieros*, 'holy,' and *γλυφω*, 'I carve,') A name given to the signs employed in medicine; and, also, to the folds in the hands, feet, and forehead, which afford chiromancy its pretended oracles.

HIEROPYR, (*hieros*, 'holy,' and *πυρ*, 'fever,') Erysipelas.

HIKRY PIKRY, (*Hiera picra*), *Pulvis aloës et Canella*.

HILE DU FOIE, see Liver.

HILL'S BALSAM OF HONEY, see Balsam of Honey, Hill's—*h. Essence of Bardana*, *Tinctura guaiaci*.

HILLBERRY, *Gaultheria*.

HILL FEVER, see Fever, hill.

HILON, (F.) *Hilum*, improperly *Hilus*, 'the black spot on the top of a bean.' A name given by some writers to a small blackish tumour, formed by the protrusion of the iris through an opening in the transparent cornea, so called from its comparison with the *hile* or black mark presented by the vicia faba at one of its extremities. In Botany, the *Hile* or *Hilum* is the cicatrix of a seed, which indicates the place by which it was attached in the cavity of the pericarp. It is the umbilicus of the seed. The fissure of the spleen, kidney, lung, ovary, &c., is, also, called *Hilus*. See Porta.

HILUM, *Hilon*.

HILUS, *Hilon*—*h. Hepatis*, see Liver.

HILUS LIENALIS, *Incisura lienalis*. The concave part of the spleen.

HILUS RENALIS, *Incisura renalis*. *Hilus* of the kidney. The concave part of the kidney.

HILWORT, *Mentha pulegium*.

HIMANTOMA, see Himas.

HIMANTOSIS, Himas.

HIMAS, gen. *Himantos*, (*himas*, gen. *himantos*, 'a thong of leather,') The uvula; likewise elongation, and extenuation of the uvula. It is also called *Himantosis*, *Himantoma*. See *Staphylodema*.

HIMEROS, (*ἡμιρος*, 'amorous desire,') *Uvido*.

HINDBERRIES, see *Rubus idæus*.

HINDHEEL, *Tanacetum*.

HIP. ([Sax.] *hipe*.) Haunch, Hypochondriasis—*h.* bone, *Ilion*—*h.* Disease, Coxarum morbus—*h.* Joint, Coxofemoral articulation—*h.* Tree, *Rosa canina*.

HIPPACĒ. (*ἵππακῆ*, from *ἵππος*, 'a horse.') *Culsus Equinus*. A cheese prepared from mare's milk.

HIPANTHROPIA, (*hippo*, and *άνθρωπος*, 'a man.') A variety of melancholy, in which the patient believes himself changed to a horse. The Greek word *ἵππανθρωπος*, means the fabulous Centaur.

HIPPASIA, (*ἵππασία*), Equitation.

HIPPED, Hypochondriac.

HIPPEIA, (*ἵππεα*), Equitation.

HIPPEUSIS, (*ἵππευσις*), Equitation.

HIPPIATER, *Hippiatrus*, *Medicus equarius*, (*hippo*, and *ιατρος*, 'a physician.') A farrier. A horse doctor. A horse leech. Used also for one who treats the diseases of other domestic animals; *Veterinarus*, *Mulomedicus*, *Medicus Veterinaris*, *Zoia'trus*.

HIPPIATRI' A. *Hippiatrica*, *Hippiatricæ*, *Medicina equaria*, (F.) *Hippiatrique*. A science, whose object is the knowledge of the diseases of the horse. It is sometimes made to include other domestic animals. See Veterinary art.

HIPPIATRIQUE, Hippiatria.

HIPPIATRUS, Hippiater.

HIPPO, (*ἵππος*, 'a horse,') in composition, 'horse.' It often expresses 'greatness, excess,' like *Bou*, *Bu*, *Horse*, *Ox*, &c.

HIPPO, *Ipecacuanha*. A corruption, in Ireland, of *Ipecacuanha*; and hence applied, at times, in the United States, to the American plant, *Euphorbia Corollata*, which is, also, called *Ipecacuanha*.

HIPPO, INDIAN, *Gillenia trifoliata*.

HIPPOCAMPE GRANDE, (*ἵπποκαμπος*, 'the sea horse,') *Cornu ammonis*.

HIPPOCAMPUS MAJOR, *Cornu ammonis*.

HIPPOCAM'PUS M'NOR, *Pes hippocampi m'nor*, *Un'guis*, *U. A'vis seu Halleri*, *O'crea*, *Colliculus caecæ posterioris ventriculi*, *rum laterali*, *l'ium*, *Collicar a'vis*, *Un'coiform Em'innence*, (F.) *Ergot*, *Eperon*. A medullary tubercle or projection, observed in the posterior cornu of the lateral ventricle of the brain.

HIPPOCENTAUREA CENTAURIUM, *Chironia centaureum*.

HIPPOCRAS, (from being strained through Hippocrates' sleeve [?]), see *Claret*.

HIPPOCRATES, CAP OF, *Bonnet d'Xippocrate*—*h.* Sleeve, *Chausse*.

HIPPOCRATIC, *Hippocraticus*, (F.) *Hippocraticque*. Relating to Hippocrates, or concerning his doctrine,—as *Hippocratic doctrine*, *Hippocratic face*, &c.

HIPPOCRATISM, *Hippocratism'us*. The doctrine of Hippocrates, which consisted in observing and following nature, as indicated by her presumed efforts, and the crises set up by her.

HIPPOCRATIST. A partisan of the Hippocratic doctrine.

HIPPOGLOSSUM, *Ruscus hypoglossum*.

HIPPOGONYOLEPUS, (*hippo*, *γονν*, 'the knee,' and *λεπος*, 'a scale.') *Crusta genu equine*.

HIPPOLAPATHUM, (*ἵππολαπαθον*, from *hippo*, and *λαπαθον*.) *Rumex patientia*.

HIPPOLITHUS, (*hippo*, and *λιθος*, 'a stone,') Bezoard of the horse.

HIPPOMANE MANCINEL'LA, (*hippo*, and *μανη*, 'madness,' 'horse,' or violent madness.) *Manchineel*, (F.) *Mancenillier*. Family, *Euphorbiaceæ*. See *Syst. Monœcia Monadelphia*. A poisonous tree of South America, with the milky juice of which the Indians poison their arrows. The fruit, also, is highly poisonous.

HIPPOMANES (*ἵππομανης*). Applied to a mare in the state of heat or rutting. Also the secrete-

tion from the vagina of a mare in that condition. Used, of old, in the preparation of philters.

HIPPOMARATHRUM, (*ἵππομαραθρον*, from *hippo*, and *μαραθρον*.) *Puccellanum silaus*.

HIPPOPATHOL'OGY, *Hippopatholog'ia*, (*hippo*, *παθος*, 'a disease,' and *λογος*, 'a discourse,') The science of the diseases of the horse. Pathology of the horse.

HIPPOPHAGOUS, (*hippo*, and *φαγω*, 'I eat,') Equivorous.

HIPPOPUS, (*hippo*, and *πους*, 'a foot,') See *Kyllosis*.

HIPPOS, (*ἵππος*), *Equus*.

HIPPOSELINUM, (*ἵπποσελινον*), *Smyrniolum olusatrum*.

HIPPOSTEOL'OGY, *Hipposteolog'ia*, (*hippo*, *οστέον*, 'a bone,' and *λογος*, 'a discourse,') Osteology of the horse.

HIPPOT'OMY, *Hippotom'ia*, (*hippo*, and *τεμνω*, 'to cut,') Anatomy of the horse.

HIPPU'RIA, (*hippo*, and *ουρον*, 'urine,' because the urine contains hippuric acid, which is found in the urine of the horse.) A pathological condition in which there is an excess of hippuric acid in the urine.

HIPPURIC ACID, see *Acid*, hippuric.

HIPPURIS VULGARIS, (*hippo*, and *ουρα*, 'a tail,') The systematic name of the *Horse's Tail*, *Mare's Tail*, *Equisetum minus*, *Equisetum*, (F.) *Prêle*, *Preele*, *Asprêlle*, *Queue de cheval*. It is an astringent, and frequently used, as tea, by the vulgar, in diarrhoea and hemorrhage. The same virtues are attributed to the *Equisetum arven'se*, *E. fluviat'ilè*, *E. limosum*, &c.

HIP'PUS, (*ἵππος*), *E'quis*, *Nictat'io*. A disease of the eyes, in which from birth they perpetually twinkle, like those of a man on horseback. [?] Also, a tremulous condition of the iris, which occasions repeated alternations of contraction and dilatation of the pupil; *Iridot'romus*, *Iridodone'sis*, *Iridopla'nio*, *Tremor Pridis*.

HIPS, (Sax. *heopa*.) The fruit of the dog-rose, *Rosa canina*: chiefly used as a confection.—See *Confectio Rosæ Caninæ*.

HIRCIS'MUS, *Hircus*, *Hir'quis*, *Gra'sus*, (*hircus*, 'a goat,') Stinking like a goat; applied especially to the odour of the secretions of the axilla.

HIRCUS, ('a goat,') *Canthus* (greater), *Hircismus*, *Tragus*—*h.* *Alarum*, *Cinabra*.

HIRPLING (Prov. and Sc.), *Claudication*.

HIRQUITAL'ITAS, (from *hircus*, 'a goat,') *Parapho'nia pu'berum*. Goat's voice. See *Egophony*.

HIRQUUS, ('a goat,') *Hircus*, *Canthus* (greater), *Hircismus*, *Tragus*.

HIRSUTIES, (*hirtus*, 'hairy,') *Das'yma*, *Das'ytes*, *Tricho'sis*, *Hirsut'ies*, *Hair'iness*. Growth of hairs on extraneous parts, or superfluous growth on parts; as in cases of bearded women. When congenital, it is termed *Hirsut'ies adna'ta*.

HIRU'DO, gen. *Hiru'dinis*. The Leech, *Sanguin'ga*, *Bdel'la*, (F.) *Sanguis*, (*haurio*, 'I draw out,' or from *hæreo*, 'I stick,') [?] In medicine, *Hiru'do seu Sanguin'ga Medicinal'is*, *Bdel'la Medicinal'is* or *Medic'inal Leech*, and *Hirudo Sanguisuga officinal'is* are generally employed. In the United States, *H. dec'ora* is used. The leech lives in fresh water, and feeds on the blood of animals, which it sucks, after having pierced the skin with its three sharp teeth. This habit has been taken advantage of to produce local bloodletting. In applying the leech, the part must be wiped dry; and if there be difficulty in making it suck, a little milk or cream may be applied. When satiated, it will drop off, and by applying a little salt or vinegar to its head it will disgorge the blood. A good English leech will take about half an ounce of blood, including that which flows by fomenting the part subsequently. The American takes less.

HIRUDO ARTIFICIALIS. *Antlia sanguisuga*—h. *Decorata*. see *Hirudo*—h. *Medicinalis*, *Hirudo*.

HIRUNDINARIA. (from *Hirundo*.) *Asclepias vineetoxicum*, *Lysimachia nummularia*.

HIRUNDO. gen. *Hirundo*, *Chelidon*, (from *hirundo*, 'by sticking,' because it sticks its nests against walls. [?]) *The Swallow*. The nests of the swallow were once employed as rutefacients, boiled in vinegar. The edible birds' nests of the *Hirundo asiatica* of the Indian Archipelago, are highly prized, especially by the Chinese, as an article of diet. They have not been accurately analysed, but seem to be of a mucilaginous character. They are formed by the bird vomiting the food which it has taken. See *Aleyon*.

HISPANIUM VIRIDE. ('Spanish green,') *Cypripedium*.

HISPIIDITAS. (*hispidus*, 'hairy,') *Dystrochia*, *Phalangosis*.

HISPIIDULA. *Antennaria dioicum*.

HISSING RESPIRATION. see *Adle sibilant*.

HISTIOLOGICAL. *Histological*.

HISTIOLOGY. (from, dim. of *hios*, 'a tissue,' and *logos*, 'a description,') *Histology*.

HISTOCHEMIA. see *Chymistry*.

HISTOCHEMISTRY. see *Chymistry*.

HISTOCHYMY. see *Chymistry*.

HISTODIALYSIS. (*hios*, and *diastasis*, 'dissolution,') A morbid dissolution of the tissues.

HISTOGENETIC. *Histogeneticus*, 'Tissue making.' (Same etymon as the next.) Relating or appertaining to *histogeny* or the formation and development of the organic textures.

HISTOGENETIC ELEMENTS OF NUTRITION. see *Plastic Element*.

HISTOGENY. *Histogenia*, *Histogenesis*. (F.) *Histogenic*, *Tissue formation*, (*hios*, and *genesis*, 'generation,') The formation and development of the organic textures.

HISTOLOGIA. (*hios*, and *logos*, 'a description,') *Histology*.

HISTOLOGICAL. *Histologicus*, *Histologicus*, *Histologicus*. Relating to histology. Applied, also, at times, to the natural transformations that occur in the tissues in the embryo, in contradistinction to *morphological*, which applies to the alterations in the form of the several parts of the embryo.

HISTOLOGY. *Histologia*, *Histology*, *Histology*, (*hios*, and *logos*, 'a description,') *Anatomy* (general). The term is also more particularly appropriated to the minute anatomy of the tissues. See *Anatomy*.

Pathologic Histology, *Microscopic pathology*, *Anatomy*, is the morbid anatomy of the tissues.

HISTOLYSIS. (*hios*, and *lysis*, 'solution,') The decay and dissolution of the organic tissues and of the blood. It includes the various forms of retrograde metamorphosis and degeneration.

HISTONOMY. *Histonomia*, (*hios*, and *nomos*, 'law,') The aggregate of laws which preside over the formation and arrangement of the organic tissues.

HISTORY, MEDICAL. *Historia Medicæ*. A narration of the chief circumstances, and the persons connected with them, in the progress of medicine.

HISTOS. (*hios*, 'a web or tissue,') The organic texture, *Textura seu Tela organica*.

HISTOTOMY. *Histotomia*, (*hios*, and *tomos*, 'incision,') Dissection of the tissues.

HISTOTRITEUR. (*hios*, and *tere*, 'to rub,') *Écraseur*.

HISTOTRITIE. see *Écraseur*.

HIVE SYRUP. *Syrupus scillæ compositus*.

HIVES. *Cynanche trachealis*, *Urticaria Vari-cella*, (from *Sax.* heapan, 'to raise,') [?] In Scotland, according to Dr. Jameson, *Hives* or *Hyves* means any eruption of the skin, proceeding

from an internal cause; and, in Lothian, it is used to denote both the red and the yellow gum. In the United States it is vaguely employed; most frequently, perhaps, for *Urticaria*.

HIVES. *Bulla*, *Cynanche trachealis*, *Urticaria*—h. *Eating*, *Rupia escharotica*.

HOARHOOUND. *Marrubium*.

HOARSENESS. (from *Sax.* hay, by onomatopoeia.) *Rauco*.

HOAST. (*Sax.* hōsta, [G.] *Huster*,) *Tussis*.

HOBBIL. *Idiot*.

HOBNAIL LIVER. *Cirrhosis of the liver*. *Liver*, *nutmeg*.

HOC. *Alcea rosea*.

HOCK. (*Sax.* hōc, [G.] *Poples*).

HOCKS. *Alcea rosea*.

HOD. (Prov.), *Crick in the neck*.

HOFF. *Poples*.

HOG-LICE. *Onisci aselli*.

HOGWORT. *Heptadon graveolens*.

HOIST. ([Sc.] *Sax.* hōsta, [G.] *Husten*,) *Tussis*.

HOLANENCEPHALIA. (*hōl*, 'the whole,' and *anencephalia*,) Entire absence of brain,—the same as *Anencephalia*.

HOLARTHITIS. (*hōl*, 'the whole,' and *arthrit*,) *Hamarthrit*.

HOLCÉ. (*hōl*,) *A drachm*.

HOLCIMUS. (*hōl*, from *hōl*, 'a weight,') *A tumour of the liver*.

HOLCUS SORGHUM. *Panicum Italicum*.

HOLENE. (Sc.) (*Sax.* hōlen, [G.] *Holen*,) *Helen*.

HOLERA. *Cholera*.

HOLES. *Burnt*, *Rupia escharotica*.

HOLLANDS. *Gin*.

HOLLEN. ([Prov.] *Sax.* hōlen, [G.] *Holen*,) *Helen*.

HOLLY, AMERICAN. (*Sax.* hōlen, [G.] *Holen*,) *Helen*—h. *Common*, *Helen aquifolium*—h. *Daboon*, *Helen vomitoria*—h. *Ground*, *Pyrola maculata*, *Pyrola umbellata*—h. *Sea*, *Eryngium maritimum*.

HOLLYHOCK, COMMON. *Alcea rosea*.

HOLMES WEED. *Scrophularia nodosa*.

HOLMISCU. (*hōl*,) *Alveolus*.

HOLMUS. (*hōl*,) *Mortar*.

HOLM. (from *hōl*, 'whole,') in composition, the whole.

HOLOCYRON. *Tenerium chamæpitys*.

HOLOGANGLIITIS. (*hōl*, *ganglion*, and *itis*,) *Cholera*.

HOLONARCOSIS. (*hōl*, and *narcosis*, 'stupor,') *Narcosis of the whole body*. *Torpefactio universalis*.

HOLOPATHY. *Holopathia*, (F.) *Holopathie*, (*hōl*, 'whole,' and *pathos*, 'affection,') A doctrine which teaches that diseases, as they present themselves, are but phases or 'episodes' of a general affection of the organism—such affection or diathesis producing the phases, when circumstances favor their manifestation.

HOLOPHYCTIDES. *Phlyctena*.

HOLOSTEUM ALSINE. *Alsine media*.

HOLOSTEUS. (*hōl*, and *osteo*, 'a bone,') *Osteocolla*.

HOLOTETANUS. (*hōl*, and *tetanus*,) *See Tetanus*.

HOLOTONIA. (*hōl*, and *tonos*, 'I stretch,') *Holotonicus*, *Tetanus*.

HOLOTONICUS. *Holotonia*. A spasm of the whole body—a variety of tetanus.

HOLYN. (Sc.) (*Sax.* hōlyn, [G.] *Holen*,) *Helen*.

HOLYWELL. *MINERAL WATERS* (F.) *Holywell* is a town in Wales, and takes its name from the famous well of St. Winifred. It is a simple cold water, remarkable for its purity.

HOMAGRA. *Omagra*.

HOMOPATHY. *Hemecopathy*.

HOMERDA. (*hōmo*, 'man,' and *merda*, *excrement*,) *Stereus humanum*.

HOMERIA COLLINA, (after Homer,) *Cape Tulip*. A poisonous South African plant, *Nat. Ord.* Iridææ, the bulb of which acts as a violent acro-narcotic, producing fatal results very speedily.

HOMESICKNESS, Nostalgia.

HOMICIDAL, *Truicidat'us*, *F'rus*, (homo, 'man,' and *cido*, 'I kill.') Pertaining or relating to homicide or the killing of man.

HOMICIDAL INSANITY, *Homicidal Monomania*, (F.) *Monomanie homicide*. Insanity, with an irresistible impulse to destroy life.

HOMILIA, (*ὁμιλία*,) Coition.

HOMINY, (of Indian derivation.) Maize or Indian corn hulled and coarsely broken. It is prepared for food by being mixed with water and boiled.

HOMINY, WHEATEN, see Groats.

HOMIOSIS, Homoiosis.

HOMME, Homo.

HOMO, gen. *Hom'inis*, (F.) *Homme*, Man,—the chief and most perfect of the mammalia; in Greek, *ἄνθρωπος*, *Anthro'pos*, from *ana*, 'upwards,' and *τρεπω*, 'I turn;' because man, alone, of all animals, possesses the natural power of standing erect. He is also the only animal whose incisor teeth, wedged in a projecting jaw, are absolutely vertical. Man is especially distinguished from other mammalia by the faculty which he possesses of classing his ideas; comparing them with each other; and connecting, representing, and transmitting them by signs and articulate sounds. He possesses in the highest degree all the attributes of intelligence—memory, judgment, and imagination. He inhabits all countries—the burning regions of the torrid zone, and the chilling atmosphere of the polar climes. In different situations he presents in his figure, colour and stature, differences which have caused mankind to be divided by naturalists into races or varieties. The number of such races can only be approximated. Blumenbach admits five—the *Caucasian*, *Ethiopian*, *Mongolian*, *Malay*, and *American*. Every division must necessarily be arbitrary, and the individuals composing each variety are far from being alike.

HOMO ALATUS, ('winged man,') see Alatus.

HOMO CAUDA'TUS, 'tailed man.' A fabulous, tailed variety of the human species, 'incola orbis antarctici,' admitted by Linnaeus, although he is uncertain whether to rank them with men or apes!

HOMO FATUUS, Idiot.

HOMOED'RUS, (*ὅμον*, 'together,' and *ἔδρα*, 'seat,') Having the same seat. *Morbi homoedri*: diseases that have the same seat.

HOMEO, (from *ὅμοιος*, 'like,') in composition, like.

HOMEO-MORPHOUS, (*homæo*, and *μορφή*, 'shape,') See Homology.

HOMEO-PATH, Homœopathist.

HOMEO-PATH'IC, *Homœopath'icus*, *Homœopath'ices*, (F.) *Homœopathique*. Relating to homœopathy.

HOMEO-PATHIE, Homœopathy.

HOMEO-PATHIQUE, Homœopathic.

HOMEO-PATHIST, *Homœopath*, *Homœopath'icus*, *Homœopathis'ta*, *Homœopathis'tes*. One who believes in homœopathy.

HOMEO-P'ATHY, *Homœopath'ia*, *Homœopath'ice*, *Ars homœopath'ica*, *Homœopath'ia*, *Hahnemannism*, *Dynamopath'ia*, (F.) *Homœopathie*, *Homœothérapeutique*, *Homœosympathie*, *Doctrine homœo-organique*, (*homæo*, and *πάθος*, 'affection,')

A fanciful doctrine, which maintains that disordered actions in the human body are to be cured by inducing other disordered actions of a like kind [*similia similibus*], and this to be accomplished by infinitesimally small doses, often of apparently inert agents: the decillionth part of a

grain of charcoal, for example, is an authorized dose. It has also been called *Glob'ulism*.

So long ago as in the writings of Gregory the Great, there is the following remark: "Mos medicinæ est ut aliquando similia similibus, aliquando contraria contrariis curet. Nam sæpe calida calidis, frigida frigidis, sæpe autem frigida calidis, calida frigida sanare consuevit."

HOMEOPLASIA, (from *homæo*, and *πλασσω*, 'I form,') The formation of accidental tissues similar to the normal; the opposite to heteroplasia.

HOMEOSIS, Homoiosis.

HOMEOSYMPATHIE, Homœopathy.

HOMEO-THÉRAPEUTIQUE, Homœopathy.

HOMEOZ'YGY, (*homæo*, and *ζυγω*, 'I join together,') (F.) *Soi-pour-soi*. The law of association of organs, by which like parts adhere to like parts.

HOMOETHNIA, (*ὁμοεθνία*,) Sympathy.

HOMOGENESIS, (*homo*, and *γενεσις*, 'generation,') See Generation.

HOMOIO-PATHIA, (*ὁμοιοπάθεια*, from *homæo*, and *πάθος*, 'affection,') Sympathy.

HOMOIO'SIS, *Homœo'sis*, *Homio'sis*, (*ὁμοιοσις*, from *ὁμοιοω*, 'I assimilate,') An elaboration of the nutritious juice, by which it becomes proper for assimilation.

HOMOLINON, (*ὁμος*, 'raw,' and *λεων*, 'flax,') See Apolinosis.

HOMOLOG'ICAL, *Homolog'icus*; (same etymon as Homology.) Relating or appertaining to homology, as *homological anatomy*.

HOMOLOGOUS, (*ὁμος*, 'like,' and *λογος*, 'condition,') See Homology.

HOMOLOGUE, *Homol'ogus*. A term applied to the same organ in different animals under every variety of form and function. (Owen.)

HOMOL'OGY. The doctrine of similarity of structure. Thus, the two sides of the body are said to be 'homologous.'

Homol'ogous, *homomorph'ous*, or *homœomorph'ous* tissues are those that resemble others; in opposition to *heterol'ogous*, *heteroclitic*, or *heteromorph'ous*, which are new formations. Homology seems now to be accepted as the designation of the doctrine or study, the subject of which is the relations of the parts of animal bodies.

HOMOMORPHOUS, (*ὁμος*, 'like,' and *μορφή*, 'shape,') See Homology.

HOMONOPAGIA, Cephalalgia.

HOMOPHAGUS, Omophagus.

HOMOPLAS'TY, (*ὁμος*, 'like,' and *πλασσω*, 'I form,') The formation of homologous tissues.

HOMOPLATA, (*ὁμος*, 'shoulder,' and *πλατυς*, 'broad,') Scapula.

HOMOT'ONUS, (*ὁμοτονος*, from *ὁμος*, 'like,' and *τονος*, 'tone,') *Æqualis*. That which has the same tone. A continued fever, whose symptoms have an equal intensity during the whole course of the disease, has been so called. See Acmaestios and Synocha.

HOMOTYPE, (from *homo*, and *τυπος*, 'type,') A repetition of the same essential part,—in the series of vertebral segments, for example. (Owen.)

HONE, (Prov. *hone*, 'to swell,') A kind of swelling in the cheek.

HONESTY, Lunaria rediviva.

HONEWORT, FIELD, (used in the treatment of hone. [?]) Sison amomum.

HONEY, (Sax. *hunig*,) Mel—h. Balsam of Hill's, see Mel—h. Bloom, Apocynum androsæmifolium—h. of Borate of Soda, Mel boracis—h. of Borax, Mel boracis—h. Clarified, Mel despumatum—h. Prepared, Mel præparatum—h. of Roses, Mel rosæ—h. of Squill, compound, Syrupus scillæ compositus.

HONEYCOMB BAG, Reticulum.

HONEY-SUCKLE, *Lonicera periclymenum*—H.

Bush, *Diervilla trifida*—h. Scarlet, *Lonicera sempervirens*—h. Trumpet, *Lonicera sempervirens*.

HONEY SUGAR, *Glucose*.

HONGLANE (Chinese), *Coptis teeta*.

HONOR CAPITIS, 'honor of the head,' see *Capillus*.

HONORAIRE, *Sostrium*.

HONORARIUM, (*honor*, gen. *honoris*, 'honor,') *Sostrium*.

HONTEUX, 'ashamed,' *Pudic*.

HOODWORT, *Scutellaria lateriflora*.

HOOK, (Sax. *hoec*, *hoec*), *Uncus*, *Uncinus*, *Anchora*, (F.) *Érigne*, *Airigne* on *Érine*. An instrument, consisting of a steel wire, flattened at the middle, and having the extremities crooked and pointed. Some hooks are furnished with a handle at one extremity—the other having one or two hooks;—constituting the *single* or *double hook*. The hook is used by anatomists and surgeons to lay hold of, and raise up, certain parts, the dissection of which is delicate, or which would slip from the fingers.

The *Tenaculum* is a variety of the hook.

HOOK, BLUNT, *Ich'thya*, *Ich'thyē*, *O'nyx*, *Un'guis ferreus*. An instrument which is passed over the flexures of the joints to assist in bringing down the foetus in parturition.

HOOP TREE, *Melia azedarach*.

HOOPER'S PILLS, *Pilulæ Aloes et Myrrhæ*.

HOOPINGCOUGH, (from the attendant *hoop*), *Pertussis*—h. Roche's Embrocation for, see *Roche*.

HOP, ([D.] *hop*), *Humulus lupulus*—h. Pillow, see *Humulus lupulus*—h. Plant, *Humulus lupulus*—h. Tree, *Ptelea trifoliata*.

HÔPITAL, Hospital—h. *des Fous*, *Morotrophiūm*.

HOPLOCHRIS'MA, gen. *Hoplochris'matis*, (*ὅπλον*, 'a weapon,' and *χρῆμα*, 'salve,') *Unguentum arma'rium*, *Arm'atory* or *Sympathetic Unguent*. A salve which was supposed to cure wounds by sympathy.—the instrument with which the wound was inflicted being anointed with it.

HOPLOMOCH'LION, (*ὅπλομαχίον*, from *ὅπλον*, 'a weapon,' and *μαχίον*, 'a lever,') The name of an iron machine or apparatus, which embraced the whole body like armour. A figure of it is given by Fabricius ab Aquapendente.

HOQUET, (by onomatopœia, like *Hiccup*), *Singultus*.

HORA, (*ώρα*), *Hebe*.

HORÆA, (*ώραία*), *Menses*.

HORÆOTES, (*ὥραςτης*), *Maturity*.

HORDEI MALTUM, 'malt of barley,' *Malt*.

HORDEOLUM, (dim. of *hordeum*), *Ordeolum*, *Posthia*, *Sclerophthal'mia*, *Cri'thē*, *Cri'thid'ion*, *Stye*, *Sty'an*, *Stine*, (F.) *Orgelet*, *Orgeolet*. A small, inflammatory tumour, of the nature of a boil, which exhibits itself near the free edge of the eyelids, particularly near the inner angle of the eye.

HOR'DEUM, *Or'deum*, *Cri'thē*. The seeds of *Hor'deum vulgare* or *Scotch Bar'ley*, *Or'd*. *Graminæ*, are ranked amongst the *Cerealia*. (F.) *Orge*. They afford a mucilaginous decoction, which is employed as a diluent and antiphlogistic. The seeds of the *Hor'deum distichon* seu *æsti'vum*, *Zooc'riton distichum*, and *H. heas'tichon* possess similar properties. Barley is freed from its shells in mills, forming the *Hor'deum mundatum* seu *decortica'tum* seu *exco'tica'tum*, *Hor'deum* of the Pharmacopœias; and, at times, is rubbed into small, round grains, somewhat like pearls, when it is called *Hor'deum perla'tum*, *Pearl Bar'ley*, (F.) *Orge perlé*, and forms the *Hor'deum denuda'tum* seu *perla'tum*, the *Hor'dei sem'ina tu'nica nudata* of some Pharmacopœias, — *Hor'deum* (Ph. U. S. and Ph. B.)

HORDEUM CAUSTICUM, *Veratrum sabadilla*—h.

Decortica'tum, see *Hordeum*—h. *Denudatum*, see *Hordeum*—h. *Exco'tica'tum*, see *Hordeum*—h. *Galactium*, *Oryza*—h. *Mundatum*, see *Hordeum*—h. *Perlatum*, see *Hordeum*.

HOREHOUND, *Marrubium*—h. Black, *Ballota foetida*—h. German, *leaved*, *Eupatorium teucri'folium*—h. Stinking, *Ballota foetida*—h. Water, *Lycopus sinuatus*, *Lycopus Virgin'icus*—h. Wild, *Eupatorium rotundifolium*, *E. teucri'folium*.

HORME, (*ὁρμη*), *Instinct*.

HORMINUM, (from *ὁρμαιν*, 'to incite,' as to *venery*'), *Salvia horminum*—h. *Coloratum*, *Salvia horminum*—h. *Sativum*, *Salvia horminum*.

HORMON, (*ὁρμων*), *Enormon*.

HORN, (Sax. *horn*), *Cornu*.

HORN POX, see *Varicella*.

HORNSEED, *Ergot*.

HORNY, *Corneous*.

HORN Y EXCRESCENCES, *Lepido'sis*, *Ichthy'asis cornig'era*, *Cor'nu cuta'nea*, (F.) *Cornes de la Peau*. Certain excrescences, which occasionally form on some part of the skin, and resemble, in shape, the horns of an animal.

HORN Y SUBSTANCE, see *Tooth*.

HORRENTIA, (*horre*, 'to shudder,') *Horripilation*.

HORRIDA CU'TIS, *Cu'tis seu Ca'ro anseri'na* seu *Gallina'cea*, *Dermatospas'mus*, *Spas'mus peripher'icus*, *Gooseskin*, (F.) *Chair de poule*. A state of the skin produced by cold or fear, and accompanying the rigor of an intermittent. It is produced, essentially, by the action of microscopic muscular fibres—the *arrectores pilo'rum*.

HORRIPILATION, *Horripilatio*, *Horror*, *Horren'tia*, *Phr'i'cē*, *Phric'as'mus*, *Phric'asis*, *Phr'i'cia*, (*horre*, 'to bristle,' and *pilus*, 'hair,') (F.) *Horripilation*, *Frissonnement*. General chilliness, preceding fever, and accompanied with bristling of the hairs over the body.

HORROR, (*horre*, 'to shudder,') *Horripilation*—h. *Ciborum*, *Disgust for food*.

HORRORS, THE, see *Delirium tremens*.

HORSE, in composition, often means great—h. *Balm*, *Collinsonia Canadensis*—h. *Cane*, *Ambrosia trifida*—h. *Chestnut*, *Æsculus hippocastanum*—h. *Crust*, *Crusta genu equinæ*.

HORSEFLY WEED, *Sophora tinctoria*.

HORSEFOOT, see *Kyllosis*.

HORSELEECH, *Hippiater*.

HORSEMINT, *Ambrosia trifida*, *Monarda coc'cinea*, *M. fistulosa*, and *M. punctata*—h. *Sweet*, *Cunila mariana*.

HORSE NETTLE, *Solanum Carolinense*.

HORSE RADISH, *Cochlearia armoracia*.

HORSESHOE HEAD. An old name for the condition of the skull in children, in which the sutures are too open,—the coronal suture presenting the form of a horseshoe.

HORSE'S TAIL, *Hippuris vulgaris*.

HORSE WEED, *Ambrosia trifida*, *Collinsonia Canadensis*, *Erigeron Canadense*.

HORTULUS CUPIDINIS, (dim. of *hortus*), *Vulva*.

HORTUS, ('a garden,') *Vulva*—h. *Siccus*, *Herbarium*.

HOSE (Prov.), (from [Sax.] *halp*), *Collum*, *Throat*.

HOSEFISH, *Sepia*.

HOSPITAL, (primarily from *hospes*, gen. *hosp'itis*, 'a guest,') *Nosocom'ium*, *Adynatocom'ium*, *Adynatodoch'ium*, *Xenodoc'ium*, *Xenodoch'ium*, *Infirmari'um*, *Infirmat'orium*, *Nosodoch'ium*, *Valitudinari'um*, *Infirm'ary*, (Old. Eng.) *Farm'icy*, *Spit'al*, (F.) *Hôpital*, *Infirmierie*, contracted in old [F.] to *hostel*, and *hôtel*. An establishment for the reception of the sick, in which they are maintained and treated medically. Hospitals were first instituted about the end of the 4th century a period at which the word *νοσοκομειον* was employed, for the first time, by St. Jerome. They may be

general, receiving all cases; or *special*, admitting only the subjects of certain diseases. See Nosocomial.

HOSPITAL GANGRENE, *Phagedæna gangrenosa*, *Putrid* or *Malignant Ulcer*, *Gangræna Nosocomialis* seu *Nosocomia* *læs* seu *Contagio* *sa*, *Sphacellus nosocomialis*, *Hospital Sore*, *Putredo*, (F.) *Pourriture* or *Gangrène d'hôpital*. Gangrene, occurring in wounds or ulcers, in hospitals, the air of which has been vitiated by the accumulation of patients, or some other circumstance. Hospital Gangrene—many different varieties of which are met with, and always accompanied or preceded by fever—commonly commences with suppression of the suppuration of the wound, which becomes covered with a grayish and tenacious sanies. The gangrene then manifests itself. It extends from the centre of the ulcerated surface towards the edges; these become swollen, painful, and everted; and the patient dies with all the signs of typhus. The treatment must be varied according to circumstances. Sometimes, it requires the use of stimulating, acid, caustic, and antiseptic applications; with, occasionally, the actual cautery, aided by the exhibition of tonics, internally:—at others, the antiphlogistic regimen and emollient applications may be necessary.

HOSPITAL, INSANE, *Morotrophium*—*h. Leper, Ladrerie*.

HOST (Sc.), (Sax. *hōpsta*), Tussis.

HOSTEL, Hospital.

HOT SPRINGS, see Virginia, mineral waters of.

HOTEL, Hospital.

HOTHOUSES, see *Acicholum*.

HOUBLON, (corr. from *lupulus*), *Humulus lupulus*.

HOUGH, (Sax. *hoh*.) Poples; (Prov.), Sickly.

HOUNDS' TONGUE, *Cynoglossum*.

HOUPPE DU MENTON, Depressor labii inferioris, Levator labii inferioris—*h. Nervense*, see *Papilla*.

HOURLASS CONTRACTION OF THE UTERUS, see *Chaton*.

HOUSELEEK, *Sempervivum tectorum*—*h. Small*, *Sedum*.

HOUSEMAID'S KNEE, see *Knee*, housemaid's.

HOUSEPUPIL, see *Housesurgeon*.

HOUSESURGEON, *Resident Surg'con*. Usually a senior housepupil or graduate, who resides in an hospital, and attends to every accident and disease, in the absence of the attending physician or surgeon. It answers, in the British hospitals, to the *Élève interne* or *Interne* of the French. The *Élève externe* or *Externe* is a less advanced pupil; from whom *Internes* are chosen. In ordinary schools, *Élève externe* means a day scholar; whilst *Élève interne* means a boarder.

HOUX, *Ilex aquifolium*—*h. Apalachine*, *Ilex vomitoria*—*h. Petit*, *Ruscus*.

HOWDY (Sc.), *Midwife*.

HOWDYWIFE (Sc.), *Midwife*.

HUACACACHU, *Datura sanguinea*.

HUACO, *Guano*.

HUANU, *Guano*.

HUBBON (Prov.), *Haunch*.

HUCKLE, (from [G.] *Hocker*, 'a protuberance,') *Hip*.

HUCKLEBERRY, see *Gaylussacia*, and *Vaccinium*—*h. Dwarf*, *Gaylussacia humosa*.

HUCKLEBONE, *Ischion*.

HUCKSHOULDERED, (from [G.] *Hocker*, 'a protuberance,') see *Hump*.

HUDSON'S PRESERVATIVE FOR THE TEETH AND GUMS, see *Tinctura Myrrhæ*.

HUILE, (from [L.] *oleum*.) Oil—*h. d'Absinthe*, *Artemisia absinthium* (oil of)—*h. d'Acajou*, see *Anacardium occidentale*.

HUILE ACOUSTIQUE (F.), *Oleum acusticum*, *Acoustic Oil*. An oil for deafness, prepared

of olive oil, $\frac{3}{4}$ ij; garlic, oxgall, and bayleaves, each $\frac{3}{4}$ i; boiled for a quarter of an hour, and strained.

HUILE D'AMANDES, *Oleum amygdalarum*—*h. d'Aneth*, see *Anethum graveolens*—*h. Animale*, *Oleum animale*—*h. Animale de Dippel*, *Oleum animale Dippelii*—*h. Animalisé par infusion*, *Oleum animalizatum per infusionem*—*h. d'Anis*, see *Pimpinella anisum*—*h. Anthelmintique de Chabert*, *Oleum animale empyreumaticum Chaberti*—*h. Aromatique*, *Oleum animalizatum per infusionem*—*h. d'Aspic*, see *Lavandula spica*—*h. d'Aurone*, *Artemisia abrotanum* (oil of)—*h. de Cacao*, *Butter of Cacao*—*h. de Cade*, see *Juniperus oxycedrus*—*h. de Carvi*, *Carum* (oil)—*h. de Cédral*, *Oleum cedrinum*—*h. de petite Chiens*, *Oleum animalizatum per infusionem*—*h. de Corne de Cerf*, *Oleum animale Dippelii*—*h. d'Épurga*, *Euphorbia lathyris*—*h. de Foie de Morue*, *Oleum jecoris aselli*—*h. de Gabian*, *Petroleum*—*h. de Gérosle*, see *Eugenia caryophyllata*—*h. de Laurier*, *Unguentum laurinum*—*h. de Lin*, see *Linum usitatissimum*—*h. de Lis*, see *Lilium candidum*—*h. de Morelle*, see *Solanum*—*h. de Morue*, *Oleum jecoris aselli*—*h. de Noir*, see *Juglans cinerea*—*h. d'Œillette*, *Papaver* (oil)—*h. de Papier*, *Pyrothionide*—*h. de Pieds de Bœuf*, *Oleum bubulum*—*h. de Ricin*, see *Ricinus communis*—*h. de Spic*, see *Lavandula spica*—*h. de Succin*, see *Succinum*—*h. de Tartre par défaillance*, *Liquor potasse subcarbonatis*—*h. Verte*, *Balsam*, green, of Metz—*h. de Vin douce*, *Oleum Æthereum*.

HUILES ANIMALES, *Olea animalia*—*h. Distillées*, *Olea volatilia*—*h. Empyreumatices*, *Olea empyreumatica*—*h. Essentielles*, *Olea volatilia*—*h. Éthérées*, *Olea volatilia*—*h. Fixes* ou *Grosses*, *Olea fixa*—*h. Fugaces*, *Olea fugacia*—*h. Médicinales*, *Olea medicinalia*—*h. Pyrogénées*, *Olea empyreumatica*—*h. Volatiles*, *Olea volatilia*.

HUIT DE CHIFFRE. 'Figure of 8.' A bandage of which the turns are crossed as in the figure 8. Such is the bandage used after bleeding from the arm.

HUIT DU COUP-DE-PIED, *Étrier*.

HUITO, *Genipa oblongifolia*.

HUITRE, (old [F.] *huistre*), *Ostrea*.

HUKEBANE (Sc.), *Ischion*.

HULGIE-BACKED, see *Hump*.

HULLUPING (Prov.), *Vomiting*.

HUM, *VENOUS*, *Bruit de Diable*.

HUMBLE, (from *humilis*), *Rectus inferior oculi*.

HUMECTANTIA, (*humectare*, 'to moisten.')

A name formerly given to drinks, which appeared to possess the property of augmenting the fluidity of the blood.

HUMERAL, *Humeralis*. That which belongs to, or is connected with, the arm or humerus.

HUMERAL ARTERY, *Arteria humeralis*, see *Brachial artery*.

HUMÉRO-CUBITAL, *Brachialis anterior*—*h. Sus-métacarpien*, see *Radialis*—*h. Sus-radial*, *Supinator radii longus*.

HUMERUS, *Scapula*, *O'mos*, *Ar'mus*, (F.) *Épaule*, *Should'er*, (Prov.) *Spauld*. The most elevated part of the arm. The bones, which concur in forming it, are—the scapula, head of the humerus, and the clavicle, united together by strong ligaments, and covered by numerous muscles.

HUMERUS, *Os humeri* seu *brachii* seu *adjutorium* seu *brachiale*, *Brachium*, *Lacer'tus*, *Shoulderbone*, (Prov.) *Spaulbone*, *Spaulbone*, *Shullbone*, (F.) *Os du bras*, is the cylindrical irregular bone of the arm; the upper extremity of which has a hemispherical head connected with the scapula; and two tuberosities or tubercles, a greater and a lesser, for the attachment of muscles, between which is the *Bicipital groove* or *Fossa*. At the inferior extremity may be remarked—the *inner condyle*, the *outer condyle*; the *small head*, which

is articulated with the radius: the trochlea articulated with the ulna, &c. The humerus is developed by seven points of ossification:—one for the body; one for the head; one for the greater tuberosity; one for the trochlea; one for the epitrochlea; one for the epicondyle; and another for the lesser head.

HUMERUS SUMMUS, 'top of the humerus,' Acromion.

HUMEUR, Humour—*h. Aqueuse*, Aqueous humour—*h. de Cotugno*, Cotunnus, liquor of—*h. Crystalline*, Crystalline—*h. Hyaloïde*, Corpus vitreum—*h. de Morgagni*, Morgagni, humour of—*h. Vitree*, Vitreine auditive.

HUMEURS FROIDES, 'cold humors,' Scrofula.

HUMIDE RADICALE, (*humeo*, 'to be moist,') Humidum radicale.

HUMIDUM NATIVUM, *H. radicale*—*h. Nativum* Articulorum, Synovia—*h. Primigenium*, *H. radicale*.

HUMIDUM RADICALÉ, *Humidum primigenium* seu *nativum* seu *seminale*, *Radical Moisture*, (F.) *Humide radicale*. Names formerly given to the liquid which was conceived to give flexibility and proper consistence to the different organic textures.

HUMIDUM SEMINALE, *H. radicale*.

HUMILIS, 'humble,' (from *humus*, 'the ground,') Rectus inferior oculi.

HUMILUS, Humulus.

HUMLOIR (Sc.), Conium.

HUMMUS, see Achicolum.

HUMOR, Humour—*h. Albigneus*, Aqueous

humour—*h. Articularis*, Synovia—*h. Ceruminous*,

Cerumen—*h. Doridis*, Water, sea—*h. Genitalis*,

Sperm—*h. Glacialis*, Crystalline, Corpus vitreum

—*h. Hyalinus* seu *Hyaloïdes*, Corpus vitreum—*h.*

Lacrymalis, Tear—*h. Lacteus*, Milk—*h. of Meibomius*, see Meibomius, glands of—*h. Melancholicus*, see Mercurialis—*h. Mercurialis*, see Mercurialis—*h. Morgagnianus*, Morgagni, humor of—*h. Ovatus*, Aqueous humour—*h. Oviformis*, Aqueous

humour—*h. Pericardii*, see Pericardium—*h. Purgulentus*, Pus—*h. Seminalis*, Sperm—*h. Venericus*,

Sperm—*h. Vitreus*, Corpus vitreum.

HUMORAL, *Humoralis*. Proceeding from, or

connected with, the humours.

HUMORES INQUILINI, 'alien humors,'

Recremental Secretions.

HUMORIQUE, see *Bruit humorique*.

HUMORISM, *Humoral Pathology*, *Pathology*

logia humoralis. A medical theory, founded exclusively on the parts which the humours were considered to play in the production of disease. Although traces of this system may be found in the most remote antiquity, the creation, or, at all events, the arrangement of it may be attributed to Galen, who enveloped it in metaphysical subtleties relating to the union between the elements and the four cardinal humours.

HUMORISTS. The Galenic and other physicians, who attributed all diseases to the depraved state of the humours, or to vicious juices collected in the body.

HUMOUR, *Humor*, (from *humeo*, 'to be moist,')

Hygroïsis, *Hygroïdon*, *Hygrum*, (F.) *Humour*,

Every fluid substance of an organized body;—as

the blood, chyle, lymph, &c. The *Humours*,

Chymi, *Humores*, differ considerably as to number

and quality in the different species of organized

beings; and even in the same species, according

to the state of health or disease. The ancients

reduced them to four; which they called

cardinal humours:—the blood, phlegm, yellow

bile, and atrabillis or black bile. It is not easy to

give a satisfactory classification of the humours.

HUMP. (Perhaps from *umbo*, 'the boss of a

buckler,') *Hunch*, *Bunch*, *Boss*, *Gilber*, *Gilbus*,

Gibba, *Tuber*, (F.) *Bosse*. A prominence, formed

by a deviation of the bones of the trunk. Com-

monly, it is formed by the spine or sternum, and is seated at the posterior or anterior part of the trunk. It may, also, be produced by deviation of the ribs or pelvis. The spine may be curved in three principal directions. 1. *Backwards*, the most common case: this the ancients called *cyphosis*, *Cyphosis*, *Gibbositas*, *Rachiocephosis*. 2. *Forwards*, *Lordosis*, *Recurvatio*; and 3. *Laterally*, *Scoliosis*, *Obstipatio*. Most curvatures occur at a very early age, and are caused by scrofula, rickets, &c.; and, not unfrequently, they are accompanied by caries of the vertebrae. See Vertebral disease. One who is affected with a hump, or is *hump-backed*, *Hunch-backed*, *Bunch-backed*, *Humpy*, *Huckshoultered*, *Shrumpshoultered*, (Sc.) *Hulgiebacked*, *Trudge-backed*, (Prov.) *Croop-backed*, is called, by the French, *bossu*.

HUMPACKED, see Hump.

HUMPTY, see Hump.

HUMULUS, (from *humus*, 'the ground,' on which it creeps,) Humulus lupulus.

HUMULUS LUPULUS, *Lu'pulus*, *L. scandens* seu *communis* seu *salictarius*, *Hu'mulus*, *Hu'milus*, *Com'bulnus* *perennis*, the Hoppant, *Nat. Ord.* Urticaceæ. (F.) *Houblon*, *Vigne du nord*. Its cones or strobiles, *Hu'muli strob'ili*, *Hu'mulus* (Ph. U. S.), have a fragrant odour; and a bitter, aromatic taste, depending on a peculiar principle, named *Lu'pulin*, extractive and essential oil, which may be extracted, equally, by water and spirit, from the dried strobiles. The hop is employed as a tonic and hypnotic, and enters into the composition of ale and beer.

The *Hop pillow*, *Pulv'nar Hu'muli*, has long been used for producing sleep.

HUNCH, Hump—*h. Backed*, see Hump.

HUNGARICA FEBRIS, Fever, Hungary.

HUNGER, (Sax. *hungr*,) *Faim*, *Li'mus*, *Pei'né*, *Pei'na*, *Eau'ries*, *Jeju'nium*, *Jeju'nitas*, *Esuri'tio*, *Esuri'go*, (F.) *Faim*. The necessity for taking food. Hunger is an internal sensation, which some authors have attributed to the friction between the sides of the stomach in its empty state; others, to the compression of the nerves, when the organ is contracted; others, to the action of the gastric juice, &c. It is dictated by the wants of the system:—farther we know not. See Appetite.

HUNGERCURE, Limotherapeia.

HUNGRY, *Famel'icus*, *Li'micus*, *Limo'des*; (Prov.) *Haw*, (vulg.) *Peck'ish*. Affected with hunger.

HUNTINGDON, MINERAL WATERS OF. Huntingdon Springs, called "Warm Springs," are seated five miles north of the town of Huntingdon, in Pennsylvania. They are resorted to, but have not been well examined.

HUNTSMAN'S CUP, *Sarracenia purpurea*.

HU'RA BRASILIENSIS, *Assacu*, *Assacu*, *Ussacu*. A Brazilian tree, of the *Nat. Ord.* Euphorbiaceæ, which, in form of the extract of the bark, is esteemed a specific in leprosy. It is, also, given in elephantiasis, and as an anthelmintic.

HURDIE (Sc.), Nates.

HURTLEBERRY, *Vaccinium myrtillus*.

HYA-HYA, *Tabernamontana utilis*.

HYACINTH, WILD, (*ὕακινθος*,) *Scilla esculenta*.

HYACINTHUS MUSCARI, *Bulbus vomitorius*.

HYÆNAN'CHÈ GLOBO'SA, (*hæna*, *hyæna*, and *ayæu*, 'to strangle,') An arborescent shrub of South Africa, *Nat. Ord.* Euphorbiaceæ, the fruit of which, pounded, is used to destroy hyænas and other beasts of prey, and seems to contain strychnia.

HYALEUS, (from *haleos*, 'glass,') Hyaline.

HY'ALINE, *Hyali'nus*, *Hyale'us*, *Vit'reus*, *Vit'reous*, (F.) *Vitreus*, *Vitré*. Glassy. Resembling glass.

HYALINE SUBSTANCE, Cytoblastema.

HYALITIS, *Inflammati'o tunicæ hyaloïdem*, *Hyaloï'tis*, *Vit'reo capsul'itis*. (F.) *Hyalite* (*hyalo*, and *itis*.) Inflammation of the hyaloid membrane of the eye.

HYALO, (from *υαλος*, 'glass,') in composition, vitreous.

HYALOID, *Hyalo'des*, *Hyaloï'des*, (F.) *Hyaloides*, (*hyalo*, and *oides*, 'resemblance,') Vitreiform; resembling glass. Also relating or appertaining to the hyaloid body, membrane, &c. (F.) *Hyaloïdien*.

HYALOID CANAL, see Hyaloid membrane.

HYALOID FOS'SA, *Fos'sa Hyaloïdea*, is a cup-like excavation in the vitreous humour, in which the crystalline is imbedded.

HYALOID MEMBRANE, *Hyalomem'bra*, *Tu'nica Hyaloïdea* seu *Vit'rea*, *Membr'na Arachnoïdea*, is the extremely delicate membrane, which forms the exterior covering of the vitreous humour, and transmits within it prolongations, which divide it into cells. Fallopius discovered this membrane, and gave it the name *Hy'aloid*. On a level with the entrance of the optic nerve into the eye, the hyaloid membrane has been described as forming, by reflection, a cylindrical canal, which pierces the vitreous humour from behind to before, as far as the posterior part of the crystalline. See Canal, hyaloid. At its anterior part is a depression—*Fossa patella'ris* seu *lenticula'ris*—for the reception of the crystalline lens.

HYALOÏDIEN, see Hyaloid.

HYALOIDITIS, Hyalitis.

HY'ALOMENINX, (*hyalo*, and *μνιγξ*, 'a membrane,') Hyaloid membrane.

HYALONIXIS, (*hyalo*, and *νωσω*, 'to push,') see Cataract.

HYALONYXIS, see Cataract.

HYALOS, (*υαλος*,) Vitrum.

HYANCHE, (*βαγχη*, from *βς*, 'a swine,' and *αγχω*, 'I choke,') Cynanche tonsillaritis.

HYBOMA, gen. *Hybo'matis*, (*ὑβωμα*, from *ὑβωω*, 'to make convex,') Gibbositas.

HYBRID, *Hy'brida*, *Hy'brida*, (F.) *Hybride*, (*ὑβρις*, gen. *ὑβριδος*, 'mongrel or mungrel,') (F.) *Métis*, *Mulet*. A being born of two different species,—as the mule. The term is applied to plants as well as to animals. The result is termed *Hybrid'ity* or *Hy'bridism*. Hybrid is often, also, used to designate words which are formed from two different languages—as *uteritis*, for inflammation of the uterus, in place of *metritis*.

HYBRIDISM, see Hybrid.

HYBRIDITY, see Hybrid.

HYDALEUS, (*ὑδαλεος*,) Hydropic.

HYDARTHROSIS, Hydrarthrus.

HYDARTHROS, Hydrarthrus.

HYDAS, (*ὑδας*,) Hydor.

HYDATID, *Hy'datis*, *Bulla*, *Aqu'ula*, *Hydro'a*, *Hydrocys'tis*, *Hygrocys'tis*, *Tæ'nia hydatig'ena*, *Echinococ'cus huma'nus*, (*ὑδωρ*, gen. *ὑδατος*, 'water,') This name was long given to every encysted tumour which contained an aqueous and transparent fluid. Many pathologists subsequently applied it to vesicles, so that both the tissue of membranes, more or less transparent, which are developed within organs, but without adhering to their tissues. It is by no means clear that these formations are really entozoa. They have been found in various parts of the body; sometimes in the uterus, occasioning signs nearly similar to those of pregnancy, but being sooner or later expelled. The expulsion is generally attended with more or less hemorrhage. See Acephalocyst.

Hy'datis, *Aqu'ula*, *Phlyctæ'nula*, *Verru'ca Palpebrarum*, *Mil'ium*, also, meant a small, transparent tumour of the eyelids.

HYDATIDES (pl. of *hydatis*) CERVICIS I'TERI, Nabothi glandulæ.

HYDATIDIFORM, Hydatiform.

HYDATIDOCE'LE, *Hydatocel'le*, (*hydatid*, and

κελη, 'a tumour,') Osechocele containing hydatids; the *Osechocel'le hydatido'sa*, *Hydatidoscheocel'le* of Sauvages.

HYDATIDOÏDES, Hydatoid.

HYDATIDO'MA, (*hydatid* and *oma*,) A tumour caused by hydatids.

HYDATIDOSCHEOCELE, (*hydatid*, *σχημα*, 'scrotum,' and *κελη*, 'a tumour,') Hydatidocele.

HYDATIFORM, *Hydati'form'is*, *Hydatid'i-form*, *Hydatid'i'form'is*, (*hydatid*, and *forma*, 'form,') Resembling an hydatid.

HYDATINUS, (*ὑδωρ*, gen. *ὑδατος*, 'water,') Hydatoid.

HYDATIS, gen. *Hydat'idis*, Hydatid—h. Finna, Cysticercus cellulose.

HYDATIS'MUS. The noise caused by the fluctuation of pus contained in an abscess.

HYDATO, (*ὑδωρ*, gen. *ὑδατος*, or *ὑδωρ*, gen. *ὑδατος*, 'water,') In composition, water.

HYDATOCELE, Hydatidocele, Hydrocele.

HYDATOCH'OLEOS, (*hydato*, and *χολη*, 'bile,') *Aquo'so-bil'ious*. An epithet given to evacuated matters when mixed with water and bile.

HYDATODES, (*ὑδατωδης*, from *hydato*, and *odes*,) Aqueous.

HY'DATOID, *Hydatoi'des*, *A'queous*, *Aquo'us*, *Hydato'des*, *Hydat'inus*, *Hydatido'des*, *Hydatidoïdes*, (*hydato*, and *oides*, 'resemblance,') Watery. Resembling water. This name has been given to the membrane of the aqueous humour; and, also, to the aqueous humour itself. *Vinum hydato'des*; wine and water.

HYDATONCUS, (*hydato*, and *ογκος*, 'a tumour,') Anasarca, Edema.

HYDATOPO'SIA, (*ὑδατοποσια*, from *hydato*, and *pois*, 'drinking,') Water-drinking;—hence

HYDATOP'OTES, (*ὑδατοποτης*,) A water-drinker.

HYDATOTHERAPEIA, (*hydato*, and *therapeia*,) Hydrosudotherapeia.

HYDERIASIS, (*ὑδεριασις*,) Dropsy.

HYDERICUS, (from *hyderus*,) Hydropic.

HYDERODES, (*ὑδερωδης*, from *hyderus* and *odes*,) Hydropic.

HYDERONCUS, Anasarca, Edema.

HYDERUS, (*ὑδερως*,) Anasarca, Hydrops.

HYDRACHNIS, (*hydro*, and *αχη*, 'down,' or *sordes*,) see Varicella.

HYDRADEN, (*hydro*, and *αδην*, 'a gland,') Conglobate gland.

HYDRADEN'TIS, *Inflammati'o glandula'rum lymphatica'rum*, (*hydraden*, and *itis*,) Inflammation of a lymphatic gland.

HYDRÆDUS, (*hydro*, and *αἰδωα*, 'genital organs,') Edema of the female organs.

HYDRÆMIA, (*hydro*, and *αἷμα*, 'blood,') Hydroæmia.

HYDRAGOGA, Hydragogues.

HYDRAGOGIA, Hydragogues.

HYDRAGOGICA, Hydragogues.

HYDRAGOGUES, *Hydrago'ga*, *Hydrago'gia*, *Hydrago'gica*, *Hydrop'ica*, *Hydro'ica*, *Aquidu'ca*, (*hydro*, and *αγω*, 'I expel,') Medicines, which, by causing watery evacuations, are believed to be capable of expelling serum effused into any part of the body. These are generally cathartics or diuretics.

HYDRAGOGUM BOYLEI, Argenti nitras.

HYDRALLANTE, False Waters.

HYDRAL'LAS, (*hydro*, and *αλλας*, 'a sausage,') see Allantois. A morbid increase in the quantity of the fluid of the allantois.

HYDRALMÆ, (*hydro*, and *αλην*, 'saltness,') Waters, mineral (saline).

HYDRAM'NIOS, (*hydro*, and 'amnios,') An excessive quantity of the liquor amnii.

HYDRANGÆA or HYDRAN'GÆA ARBORIS'CENS, (*hydro*, and *αγγων*, 'a vessel,') *H. vulga'ris*, *Sevenbark*, *Wild Hydran'gea*, *Bis'sium*. Ord. Hydrangææ. An indigenous plant, which

flowers in July. The leaves are said to be tonic, sialogogue, cathartic, diuretic, and antilithic.

HYDRANGEITIS, (*hydrangia*, and *itis*.) Angioleucitis.

HYDRANGIA, (*hydro*, and *αγγιον*, 'a vessel,') Lymphatic vessels.

HYDRANGIOGRAPHIA, (*hydrangia*, and *γραφία*, 'a description,') Angiohydrography.

HYDRANGIOTOMIA, (*hydrangia*, and *τομή*, 'to cut,') Angiohydrotomy.

HYDRARGYRANATRIPSIS, *Hydrargyren-trip'sis*, (*hydrargyrum*, and *ατραπέσις*, 'rubbing in,') The rubbing in of a preparation of quicksilver, — *Hydrargyrotrip'sis*.

HYDRARGYRENTRIPSIS, *Hydrargyranat-ripsis*.

HYDRARGYRI ACETAS, *Hydrargyrus acetatus*—h. Bichloridum, H. oxyurias—h. Bicyanidum, H. cyanuretum—h. Biniodidum, H. iodidum rubrum—h. Bin-oxydum, H. oxydum rubrum—h. Bisulphuretum, H. sulphuretum rubrum—h. Borassias, H. cyanuretum—h. Bromidum, see Bromine—h. Calx alba, *Hydrargyrum præcipitatum*—h. Chloridum, H. suburias—h. Chloridum corrosivum, H. Oxyurias—h. Chloridum mite, H. suburias.

HYDRARGYRI CYANURETUM, *H. Cyan'idum seu Borus'ias seu Bicyan'idum, Hydrargyrum Cyanogenat'um seu Hydrocyan'icum seu Borus'icum seu Zoot'icum, Prus'ias Hydrarg'gri, Cyan'uret or Prus'iate of Mer'cury*, (F.) *Cyanure de Mercury*. (Made by boiling ferrocyanide of iron with red oxide of mercury, filtering, evaporating, and crystallizing.) A modification of this process is contained in the Ph. U. S. 1863. This preparation has been strongly recommended as a powerful antisyphilitic, and is admitted into the Parisian codex. Twelve to twenty-four grains may be dissolved in a quart of distilled water, and three or four spoonfuls of the solution be taken daily in a glass of any appropriate liquid.

HYDRARGYRI DEUTOIODIDUM, H. Iodidum rubrum—h. Deuto-ioduretum, see Iodine—h. Deutonitras, see H. nitras—h. Dipernitras, see H. nitras—h. Hyperoxides, *Hydrargyri nitrico-oxydum*.

HYDRARGYRI IODIDUM, *H. Iodidum viridè*, (Ph. U. S. 1863, and Ph. B. 1864,) *H. Protoiod'idum seu Proto-ioduretum seu Subiod'idum, Hydrargyrum Ioda'tum flat'um seu Iodidula'tum, Iod'idum seu Iodure'tum hydrargyro'sum, Protoiodure'tum mercu'rii, Hydrarg'yrus ioda'tus seu ioda'tus flavus, Mercu'rius oxydula'tus hydr'id'icus, Podide, Proti'odide or Green Iodide of Mercury*, (F.) *Protoiodure ou Proti'odure de Mercure*, (*Hydrarg. 3j*, *Iodin. 3v*, *Alcohol q. s.* Rub the mercury and iodine together, adding sufficient alcohol to form a soft paste, and continue the trituration till the globules disappear. Dry the iodide in the dark, with a gentle heat, and keep it in a well-stopped bottle, the light excluded.—Ph. U. S.) Dose, one grain, gradually raised to three or four.

HYDRARGYRI IODIDUM CHLORIDI, *Mercury, iodide of chloride of*.

HYDRARGYRI IODIDUM RUBRUM, *H. Biniod'idum seu Deuto-iod'idum seu Deuto-iodure'tum seu Period'idum, Hydrarg'yrum ioda'tum rubrum seu Bi'ioda'tum seu Perioda'tum, Iode'tum seu Iod'idum Hydrarg'gri'icum, Deutoiodure'tum mercu'rii, Hydrarg'yrus perioda'tus seu bi'ioda'tus seu ioda'tus ruber, Red Iodide, Bini'odide, Deuti'odide and Peri'odide of Mercury*, (F.) *Deuti'odure ou Bi'iodure ou Peri'odure de Mercure*, (*Hydrarg. corros. chlorid. 3j*, *Potassii iodid. 5x*, *Aque destillat. Oij*. Dissolve the chloride in a pint and a half, and the iodide of potassium in half a pint of distilled water, and mix the solutions. Collect the precipitate on a filter, and, having washed it with distilled water, dry it with a moderate heat, and keep it in a well-stopped bottle.—Ph. U. S.) Dose, onesixteenth to one-fourth of a grain.

HYDRARGYRI IODIDUM VIRIDE, *H. Iodidum—h. Murias basi Oxydi imperfecti, H. oxyurias—h. Murias corrosivus, H. oxyurias—h. Murias dulcis sublimatus, H. suburias—h. Murias oxygenatus, H. Oxyurias—h. Murias spirituosus liquidus, Liquor hydrargyri oxyuriatus—h. Murias suboxygenatus præcipitatione paratus, Hydrargyrum præcipitatum*.

HYDRARGYRI NITRAS, *Nit'ras Hydrar'gryi in crystall'os concere'tus, Hydrar'gyrum nit'ricum, Mercu'rius nitro'sus fri'gore para'tus, Nit'ras hydrar'gyro'sus, Azo'tas hydrargyro'sus, Nit'rate of Mer'cury*, (F.) *Azotate de mercure Protoazotate de mercure, Nitrate mercurieux*. It is employed in syphilis; and, externally, in fungous, obstinate ulcers.

It is used in the formation of the *Soluble Mercury* of Hahnemann.

An *ac'id nitrate of mer'cury, Hydrar'gryi deuto-nitras seu dipernitras seu supernitras, Mercu'rius nitro'sus call'idè para'tus, Nit'ras hydrarg'gri'icus, Azo'tas hydrarg'gri'icus liq'u'idus, Li'quor Hydrar'gryi supernitrat'is, Hydrargyri pernitrat'is Li'quor* (Ph. D.), *Solution of supernitrate of mer'cury, Solution of supernitrate of deutoz'idè of mer'cury*, (F.) *Deutoazotate acide liquide de mercure, Nitrate acide de mercure*, made by dissolving four parts of mercury in eight of nitric acid, and evaporating the solution to nine parts, has been used as a caustic in malignant ulcerations and cancerous affections. *Li'quor Hydrar'gryi Nitrat'is, Solution of Nitrate of Mer'cury*, is in the Ph. U. S., 1863; and *Hydrar'gryi Nitrat'is Li'quor ac'idus* in the Ph. B. (1864.)

Ward's White Drops,—a once celebrated antiscorbutic nostrum,—were prepared by dissolving mercury in nitric acid, and adding a solution of carbonate of ammonia; or, frequently, they consisted of a solution of sublimate with carbonate of ammonia.

HYDRARGYRI NITRICOÏXYDUM, *Hydrar'gryi oxidum rubrum* (Ph. U. S.), *Hydrarg'yrus nitrat'us ruber, Mercu'rius corrosivus ruber seu præcipita'tus corrosivus seu præcipita'tus ruber seu coralli'nus, Arcadum coralli'num, Pulvis prin'cipis, Præcipita'tus ruber, Oxydum hydrar'gryi completum seu hydrarg'gri'icum, Panace'a mercurii rubra, Pulvis Joân'nis de Vigo, Ozo'ides hydrargyri rubrum, Hyeroco'ides hydrargyri, Oxydum hydrar'gryi nit'ricum seu hydrargyri rubrum per ac'idum nit'ricum, Nit'ric ox'idè of mer'cury, Red præcipitate*, (F.) *Oxide nitrique de mercure, Précipité rouge ou per se, Mercure corallin, Nitrate de mercure rouge*, (*Hydrarg. 3xxxvj*; *Acid. nitric. f3xxvij*; *Aque Oij*. Dissolve the mercury with a gentle heat, in the acid and water previously mixed, and evaporate to dryness. Rub into powder, and heat in a very shallow vessel till red vapours cease to rise.—Ph. U. S.) It is a stimulant and escharotic, and used as such in foul ulcers, being sprinkled on the part in fine powder, or united with lard into an ointment.

HYDRARGYRI OXIDES RUBRUM, *Hydrargyri nitricooxydum—h. Oxydi murias ammoniacalis, Hydrargyrum præcipitatum—h. Oxydulum nigrum, H. Oxydum cinereum*.

HYDRARGYRI OXYDUM CINE'REUM, *Æ'thiops persè, Mercu'rius nig'er Moscati seu præcipita'tus nig'er, Oxydum hydrar'gryi nig'rum seu hydrargyro'sum seu hydrarg'gri'icum præcipitatum, Oxydulum hydrar'gryi nig'rum, Pulvis mercuria'tis cine'reus, Mercu'rius cine'reus, Turpe'thum nig'rum, Gray or Black oxide of Mer'cury*, (F.) *Oxide de mercure cendrè, Oxide gris ou noir de mercure, Protoxide de mercure*. This oxide is made in various ways. It may be formed by boiling submuriate of mercury in lime water. The dose of this *Pulvis Hydrar'gryi cine'reus* is from two to ten grains. There are four other preparations of it in estimation, viz:—*Plenck's solution*, made by

rubbing mercury with mucilage. 2. By rubbing equal parts of sugar and mercury together. 3. A compound of honey or liquorice and purified mercury. 4. The blue pill, and blue ointment. All these possess the usual properties of mercury.

The *Hydrargyri Oxidum Nigrum* of the Ph. U. S. is made as follows:—*Hydrarg.* chlorid. mit., Potassa, 3i, Aqua Oj. Dissolve the potassa in the water, allow the dregs to subside, and pour off the clear solution. To this add the chloride, and stir constantly till the black oxide is formed. Pour off the supernatant liquor, wash the black oxide with distilled water, and dry with a gentle heat. Dose, one or two grains.

The *Mercurius solubilis* of Hahnemann is formed from a black oxide of mercury. It is the *Mercurius solubilis Hahnemannii*, *Oxydum hydrargyri nigri mediant ammoniâ et protonitrato hydrargyri præcipitatum*, *Azotus hydrargyro-azotum ammoniacus*, (F.) *Azotate de mercure et d'ammoniaque*, *Oxyde gris ou noir de Mercure*, *Precipité noir*, *Turbith noir*, which has been esteemed, in Germany and France, as an antisyphilitic. Dose gr. 1-5th to gr. i. It is used in the same cases as the *Hydrargyri oxydum cinereum*.

HYDRARGYRI OXYDUM NIGRUM, *H. oxydum cinereum*—*h. Oxydum nigrum mediant ammoniâ et protonitrato hydrargyri præcipitatum*, see *H. oxydum cinereum*—*h. Oxydum nitratum*, *Hydrargyri nitricum-oxydum*.

HYDRARGYRI OXYDUM RUBRUM, *H. Binorum oxydum*, *Mercurius calcinatus*, *Hydrargyri calcinatus*, (F.) *Oxide de Mercure rouge*, *Red oxide of mercury*. (Made by precipitation from a solution of bichloride of mercury by solution of potassa.) See *Hydrargyri nitricum-oxydum*.

It is stimulant and escharotic; and in large doses, emetic. Owing to the violence of its operation, it is seldom given internally.

HYDRARGYRI OXYDUM RUBRUM PER ACIDUM NITRICUM, *Hydrargyri nitricum-oxydum*—*h. Oxydum saccharatum*, *Hydrargyrum saccharatum*—*h. Oxydum sulphuratum*, *Hydrargyrum vitriolatum*.

HYDRARGYRI OXYMURIAS, *H. Chloridum Corrosivum* (Ph. U. S.) seu *Bichloridum* seu *Permurium*, *Hydrargyri murias*, *Murias hydrargyri corrosivus* seu *oxygenatus* seu *hydrargyri halidi oxydi imperfecti* seu *hydrargyri corrosivus*, *Sublimatus corrosivus*, *Sublimatum corrosivum*, *Hydrargyrum corrosivum sublimatum*, *Mercurius corrosivus* seu *corrosivus sublimatus*, *Supermurias hydrargyri*, (F.) *Dentochlorure ou Bichlorure de mercure*, *Muriate oxygéné de mercure*, *Sublimé corrosif*; *Bichloride or Oxygmuriate or Corrosive muriate of mercury*, *Corrosive sublimate*. (*Hydrarg.* ʒij, *Acid sulphur.* ʒiij, *Sodii chlorid.* ʒiiss. Boil the mercury with the sulphuric acid until the sulphate of mercury is left dry. Rub this, when cold, with the chloride of sodium, in an earthenware mortar; then sublime with a gradually increasing heat.—Ph. U. S.) It is used as an antisyphilitic eutrophic in venereal complaints, old cutaneous affections, &c. Gr. iij to Oj of water is a good gargle in venereal sore-throat, or as an injection in gonorrhœa. Externally, it is applied in cases of tetter, and to destroy fungus, or stimulate old ulcers. Dose, gr. 1-16 to gr. 1-8, in pill, once in twenty-four hours. White of egg is the best antidote to it, when taken in an overdose.

The *YEL'LOW WASH*, *Aqua phagedenica*, *Lo'tio Hydrargyri flava*, *Phagedenic Wash*, is made by adding *corrosive sublimate* to *lime-water*, usually in the proportion of fifteen grains to half a pint. It is a common application to syphilitic sores.

HYDRARGYRI PERIODIDUM, *H. Iodidum rubrum*—*h. Permurias*, *Hydrargyri oxyurias*—*h. Proto-iodidum*, *H. Iodidum*—*h. Proto-ioduretum*, *H. Iodidum*—*h. Prototartarus*, *H. tartarus*—*h. Prusias*, *H. cyanuretum*—*h. Saccharum vermifugum*,

Hydrargyrum saccharatum—*h. Subchloridum*, *H. Submurias*—*h. Subiodidum*, *H. Iodidum*.

HYDRARGYRI SUBMURIAS, *H. Chloridum seu Subchloridum seu Chloridum mitè* (Ph. U. S.), *Calomelas*, (Ph. B. 1864), *Calomelans Turqueti*, *Calomeli*, *Hydrargyrum muriaticum mitè*, *Hydrargyrum muriaticum seu Chloratus mitis*, *Hydrargyri submurias*, *Dra'co mitigatus*, *Submurias hydrargyri mitis*, *Submurias Hydrargyri sublimatum*, *Mercurius dulcis* seu *dulcis sublimatus*; when precipitated, *M. dulcis præcipitatus*, *M. sublimatus dulcis*, *M. Zoticus Hartmanni*.—*Panacea Mercurialis* (when nine times sublimed), *Murias hydrargyri dulcis sublimatus*, *Aquila*, *Man'na Metallo'rum*, *Panchymagogum minérale*, *P. Quercetanus*, *mild chloride*, *protobichloride*, *submuriate*, *subchloride*, or *mild muriate of Mercury*, *Calomel*. (F.) *Mercur doux*, *Protobichlorure de mercure*. Mild chloride of mercury is thus directed to be prepared in the Pharmacopœia of the United States:—(*Mercury*, ʒiij; *Sulphuric Acid*, ʒiij; *Chloride of Sodium*, ʒiiss; *Distilled water*, a sufficient quantity. Boil two pounds of the mercury with the sulphuric acid, until the sulphate of mercury is left dry. Rub this, when cold, with the remainder of the mercury, in an earthenware mortar, until they are thoroughly mixed. Then add the chloride of sodium, and rub it with the other ingredients till all the globules disappear: afterwards sublime. Reduce the sublimed matter to a very fine powder, and wash it frequently with boiling distilled water, till the washings afford no precipitate upon the addition of liquid ammonia; then dry it.) *Properties*:—antisyphilitic and sialagogue: in large doses, purgative. Dose: one or two grains given at night gradually excite ptyalism. Gr. v. to xx. purge. Children bear larger doses than adults.

The *BLACK WASH*, *Lo'tio Hydrargyri nigra*, is formed of *calomel*, ʒij; *Lime-water*, Oj. Used for syphilitic sores.

HYDRARGYRI SUBMURIAS AMMONIATUM, *Hydrargyrum præcipitatum*—*h. Subsulphas flavus*, *Hydrargyrum vitriolatum*—*h. Subsulphas pectinati*, *Hydrargyrum vitriolatum*—*h. Sulphas*, *Hydrargyrum vitriolatum*.

HYDRARGYRI SULPHURETUM NIGRUM, *H. sulphuretum cum sulphure*, *Hydrargyrum seu Mercurius cum sulphure*, *H. e sulphure*, *H. sulphuratus niger*, *Æthiops mineralis seu mercurialis seu narcoticus*, *Pulvis hyponitricus*, *Calomelas*, (F.) *Sulfure de mercure noir*, *Black sulphuret of mercury*, *Sulphuret of mercury with sulphur*, *Æthiops mineral*. (*Hydrarg.*, *Sulphur*, ʒiij. Rub together till the globules disappear, Ph. U. S.) Used chiefly in scrofulous and cutaneous affections. Dose, gr. x to ʒss.

HYDRARGYRI SULPHURETUM RUBRUM, *H. Bisulphuretum*, *Hydrargyrum sulphuratum rubrum*, *Minium purum seu Græcum*, *Magma cinnabaris*, *Am'mion*, *Purpurium*, *Mercurius Cinnabarinus*, *Cinnabaris*, *Cinabaris*, *Cinnabarium*, *Bisulphuret or Red Sulphuret of Mercury*, *Cinnabaris*, *Cinquer*, *Vermilion*, (F.) *Sulfure de Mercure rouge*, *Cinabre*, *Cinnabre*. (*Hydrarg.* ʒxl; *Sulphur*, ʒviij. Mix the mercury with the sulphur melted over the fire: and as soon as the mass begins to swell remove the vessel from the fire, and cover it with considerable force to prevent combustion. Rub the mass into powder and sublime. (Ph. U. S.)

It is an antisyphilitic, but chiefly used in fumigation against venereal ulcers of the nose, mouth, and throat;—ʒss being thrown on a red-hot iron. This preparation is the basis of a nostrum, called *Buerhaave's Red Pill*.

HYDRARGYRI SUPERMURIAS, *H. oxyurias*—*h. Supernitras*, see *Hydrargyri nitras*.

HYDRARGYRI TARTRAS, *H. Prototartarus*, *Tartrate of Mercury*. Antisyphilitic. Dose, one or two grains twice a day.

HYDRARGYRI ET ARSENICI IODIDUM, Arsenic and Mercury, iodide of.

HYDRARGYRI ET QUINÆ PROTOCHLORIDUM, *Protochloride of Mercury and Quinia*. A combination of mild chloride of mercury and quinia, administered in obstinate cutaneous diseases.

HYDRARGYRIA, Eczema mercuriale.

HYDRARGYRIASIS, *Hydrargyriasis*. Disease induced by the use of mercury: *Morbus Mercurialis*, *Mercurial disease*, (F.) *Maladie Mercurielle*. Poisoning by mercury. Eczema mercuriale. Also, the phenomena produced by mercury on the organism. *Mercurialism*, *Mercurialis'mus*.

HYDRARGYRICUM, Mercurial.

HYDRARGYRIUM, Mercurial.

HYDRARGYROSIS, Eczema mercuriale, Hydrargyriasis.

HYDRARGYROSTOMATITIS, see Salivation, mercurial, and Stomatitis, mercurial.

HYDRARGYROTRIPSIS, (*hydrargyrum*, and *tripsis*, 'rubbing,') Hydrargyranatripsis.

HYDRARGYRUM, *Hydrargyrum*, (*hydro*, and *argyros*, 'silver,') *Mercurius*, *Argentum vivum* seu *mobile* seu *fulsum* seu *fugitivum* seu *liquidum*, *Miscadum*, *Fumus albus*, *Lolrus*, *Arlea arseno-lum*, *Dal'dalus*, *Mater metallorum*, *Mercury*, *Quick'silver*, (F.) *Mercury*, *M. cou*, *Vif-Arget*, formerly *Argent-vif*. A fluid, brilliant metal; of a slightly bluish white colour; fluid above—39° of Fah. and under 656°. S. g., when liquid, 13.568 (Cavendish); easily oxydized. Metallic quicksilver does not act on the body, when taken into the stomach; but if oxydized, and combined with acids, it acts powerfully. It has been exhibited in cases of constriction of the bowels and in intussusception, from a notion that it must certainly pass through the bowels by its gravity. The water, in which mercury has been boiled, has been recommended as a vermifuge; but probably enjoys no such property, as chemical tests do not exhibit the presence of the metal. When the crude metal is distilled in an iron retort, it forms the *Hydrargyrum purificatum*.

HYDRARGYRUM AMMONIATOFRIATICUM, H. præcipitatum—h. Ammoniatum. H. præcipitatum—h. Biiodatum, Hydrargyri iodium rubrum—h. Biiodatum cum kalio iodato, Potassii hydrargyriodidum—h. Borussicum, Hydrargyri cyanuretum—h. Corrosivum sublimatum, Hydrargyri oxymurias.

HYDRARGYRUM CUM CRETÂ, *Hydrargyrum cum cretâ*, *Mercurius alkalisatus*, *Mercury with chalk*, *Æthiops alcalisatus*, *Gray powder*, (F.) *Mercury avec la craie*, (*Hydrarg.* 3ij; *Creta præparat.* 3v. Rub them together till the globules disappear. Ph. U. S.) A protoxide of mercury, formed by trituration with carbonate of lime. It is somewhat uncertain; and consequently not much employed as a mercurial. It possesses the properties of the black oxide of mercury, and may be advantageously exhibited in cases of diarrhoea in children, dependent upon acidity and vitiated secretions. Dose, gr. v to 3ss, twice a day, in any viscid substance.

HYDRARGYRUM CUM MAGNE'SIÂ of the Dublin Pharmacopœia resembles it in properties.

HYDRARGYRUM CYANOGENATUM, Hydrargyri cyanuretum.

HYDRARGYRUM HYDROCYANICUM, Hydrargyri cyanuretum—h. Iodatum, Hydrargyri iodium—h. Iodatum cum chlorido mercurii, Mercury, iodide of chloride of—h. Iodatum flavum, Hydrargyri iodium—h. Iodatum rubrum, Hydrargyri iodium rubrum—h. Iodidatum, Hydrargyri iodium—h. Muriatricum Mite, Hydrargyri submurias—h. Nitricum, Hydrargyri nitras—h. Periodatum, Hydrargyri iodium rubrum.

HYDRARGYRUM PRÆCIPITATUM, H. ammoniatum (Ph. U. S.), H. ammonia'tomuriat'icum, *Hydrargyri ammonio-nichloridum*, *Mercurius cœncl-*

icus, *Mur'ias ox'idi hydrar'gyri ammonio-nichlor'is*, *M. hydrar'gyri suboxygensatus præcipitatus* nè *paratus*, *Submurias ammonio-achlorargyricus*, S. *Hydrar'gyri ammoniatum seu Præcipitatum*, *Calc Hydrar'gyri al'ba*, *Præcipitatum album*, *Ammonio-nichloride of Mercury*, *Ammoniated Mercury*, *White precipitate of Mercury*, *White precipitate*, *Calcinated white Potash*, (F.) *Sousmuriate de mercure précipité*, *Précipité blanc*, (*Hydrarg.* chlorid. corros. 3vj; *aqua destillat. coag.*, *Liquor ammoniac f3viij*. Dissolve the chloride in the water, with the aid of heat, and to the solution, when cold, add the solution of ammonia, frequently stirring. Wash the precipitate till it is tasteless, and dry it.—Ph. U. S.)

A peroxide, combined with muriatic acid and ammonia, forming a triple salt. It is used in powder, to destroy vermin; and, united with lard, for the same purpose, as well as in scabies and some other cutaneous affections.

HYDRARGYRUM SACCHARATUM, *Æthiops saccharatus*, *Mercurius saccharatus*, *Oxidum hydrargyri saccharatum*, *Saccharum hydrargyri vermifugum*. A mild mercurial formula in several of the Pharmacopœias of continental Europe; formed by trituring one part of mercury with two of white sugar. It is used in the venereal affections of children.

HYDRARGYRUM ZOOTICUM, Hydrargyri cyanuretum.

HYDRARGYRUS, Hydrargyrum.

HYDRARGYRUS ACETATUS, *Sperma mercurii*, *Terra foliata mercurii*, *Mercurius acetatus*, *Hydrargyri Acetatus*, *Acetatus seu Protoacetatus Hydrargyri*, *Acetate of mercury*. This was the basis of *Keyser's pills*, and was once much celebrated in the cure of the venereal disease. The dose is from three to five grains, but it is not much used.

The formula for *Keyser's antivenereal pills* was as follows:—*Hydrarg. Acet.* 3iv; *Manna* 3xxx; *Amyli* 3ij; *Muc. G. Trag.* q. s. To be made into pills of gr. vj each. Dose, two pills.

HYDRARGYRUS BIODATUS, Hydrargyri iodium rubrum—h. Calcinatus, Hydrargyri oxydum rubrum—h. Chloratus mitis, Hydrargyri submurias—h. cum Cretâ, Hydrargyrum cum cretâ—h. Iodatus, Hydrargyri iodium—h. Iodatus flavus, Hydrargyri iodium—h. Iodatus ruber, Hydrargyri iodium—h. Muriatricus mitis, Hydrargyri submurias—h. Muricatus, Hydrargyri oxymurias—h. Nitratu ruber, Hydrargyri nitricooxydum—h. Periodatus, Hydrargyri iodium rubrum.

HYDRARGYRUS PHOSPHORATUS, *Phosphuretted mercury*. This preparation has been recommended in cases of inveterate venereal ulcers, but is now scarcely used.

HYDRARGYRUS CUM SULPHURE, Hydrargyri sulphuretum nigrum—h. Sulphuretus ruber, Hydrargyri sulphuretum rubrum.

HYDRARGYRUS VITRIOLATUS, *Turpethum minérale*, *Mercurius emeticus flavus* seu *causticus flavus* seu *luteus*, *Hydrargyri sulphas* seu *sulphas flavus* seu *sulphas flava* (Ph. U. S., 1863). *Subsulphas Hydrargyri flavus*, *Oxydum hydrargyri sulphuricum*, *Subsulphas hydrargyri peroxidati*, *Calc mercurii vitriolata*, *Turpeth mineral*, (F.) *Sous-sulfate de mercure ou turbith minéral* (*Hydrarg.* 3iv; *Acid. Sulph.* 3vi. Mix in a glass vessel, and boil in a sand-bath till a dry, white mass remains. Rub this into powder, and throw it into boiling water. Pour off the liquor, and wash the yellow precipitated powder repeatedly with hot water; then dry it.—Ph. U. S.) Two grains of this mercurial act on the stomach violently. It is sometimes recommended as an emetic in amaurosis.

HYDRARTHROS, Hydrarthrus.

HYDRARTHROS, Hydrarthrus.

HYDRARTHROSIS, Hydrarthrus.

HYDRARTHROSIS, *Hydrarthros*, *H. synovialis*, *Hydrarthron*, *Hydrarthros*, *Spina ventosa* of Rhazes and Avicenna, *Arthritis Hydrarthros*, *Hydrarthrosis*, *Hydrarthrosia*, *Arthrodema*, *Emmagma articulare*, *Tumour albus*, *White Swelling*, (*hydro*, and *αρθρον*, 'a joint.')

(F.) *Tumeur blanche*, *T. lymphatique des articulations*. Hydrarthrus is now generally restricted to dropsy of the articulations, *Hydrops articulationum*.

White swelling is a formidable strumous disease. It may attack any one of the joints; but is most commonly met with in the knee, the haunch, the foot, the elbow, and generally occurs in scrofulous children. It consists, at times, in tumefaction, and softening of the soft parts and ligaments, which surround the joints; at others, in swelling and caries of the articular extremities of bones; or both these states may exist at the same time. The treatment consists in the employment of counter-irritants; the use of iodine internally and externally, &c. Also, Synovia.

HYDRAS'PIS EXPAN'SA, (*hydro* and *ασπίς*, 'a round shield.')

Great freshwater Tortoise. On the sandy banks of rivers in Peru this animal buries its eggs, from which the Indians extract oil. Its flesh supplies well-flavoured food.

HYDRASTIS, (*hydro*, and *δραω*, 'I am active,' from its active juice, [?] *H. Canadensis*).

HYDRAS'TIS CANADEN'SIS, *Hydras'tis*, *Warneria Canadensis*, *Hydrophyl'um ve'r'um*, *Canad'ian Hydras'tis*, *Yel'low Root*, *Orange Root*, *Yel'low Puccoon*, *Ground Raspberry*, *Yel'low Paint*, *Golden Seal*, *Indian Paint*, *Eye'balm*. *Nat. Ord.* Ranunculaceæ. The root, *Hydras'tis* (Ph. U. S.) is used, in infusion, as a tonic; and as a 'mouth water,' and an outward application in wounds and local inflammations.

HYDRATION, *Hydra'tio*, (*ὑδωρ*, 'water.')

Impregnation or moistening with water; as *Hydraté* (F.) means impregnated or moistened with water.

HYDREDEMA, (*hydro*, and *οίδημα*, 'a swelling.')

Edema.

HYDRELE'ON, *Hydrole'um*, (*ὑδρελαίον*, from *hydro*, and *ελαίον*, 'oil.')

A mixture of water and oil.

HYDRELYTRON, (*hydro*, and *ελυτρον*, 'a sheath,')

see *Hydrocele*.

HYDRÉMIE, *Hydroæmia*.

HYDRENCEPHALITIS, (*hydro*, *encephalon*, and *itis*.)

Hydrocephalus internus.

HYDRENCEPHALIUM, (*hydro*, and *encephalon*.)

Hydrocephalus internus.

HYDRENCEPHALOC'E'LĒ, *Hydrocephaloc'e'lē*, (*hydro*, *encephalon*, and *κλῆλη*, 'rupture, protrusion.')

A monstrosity in which there is a fissure of the cranium, the integument of the head being present, and forming a hernial sac in which the brain lies outside the skull,—the sac containing a large quantity of serous fluid. Also, *Hydrocephalus chronicus*.

HYDRENCEPH'ALOID, (*hydro*, *encephalon*, and *ειδός*, 'resemblance.')

Resembling *hydroencephalus*. *Hydroencephaloid disease*, *Spurious hydroceph'alus*, *Pseudoencephali'tis*. Disorders of the bowels, and exhaustion in children are at times attended with *hydroencephaloid* symptoms.

HYDRENCEPHALON, see *Hydrocephalus chronicus*.

HYDRENCEPHALUS, (*hydro*, and *εγκεφαλος*, 'encephalon,')

Hydrocephalus internus.

HYDRENTEROCE'LE, (*hydro*, *εντέρον*, 'intestine,' and *κλῆλη*, 'a tumour.')

Intestinal hernia, the sac of which encloses fluid.

HYDRETEROMPHALOC'E'LE, *Hydrenteromphalus*, (*hydro*, *εντέρον*, 'intestine,' *μφαλος*, 'umbilicus,' and *κλῆλη*, 'rupture.')

Umbilical hernia with intestine and water in the sac.

HYDRETEROMPHALUS, *Hydrenteromphaloc'ele*.

HYDREPIGASTRIUM, see *Ascites*.

HYDREPIPOCE'LE, (*hydro*, *επιπλοον*, 'omentum,' and *κλῆλη*, 'rupture.')

Omental hernia, with water in the sac.

HYDREPILOMPHALUS, *Hydrepilomphaloc'le*, (*hydro*, *επιπλοον*, 'omentum,' and *μφαλος*, 'umbilicus.')

Umbilical hernia, with omentum and water in the sac.

HYDRETRUM, (*hydro*, and *πτερον*, 'the lower belly,')

Ascites.

HYDRIASIS, (*ὑδωρ*, 'water,')

Hydrosudothe-rapeia.

HYDRIATER, (*hydro*, and *ιατρος*, 'a physician,')

see *Hydrophatic*.

HYDRIATRICA, *Hydrosudotherapeia*.

HYDRIATRICA ARS, *Hydrosudotherapeia*.

HYDRIATRICUS, see *Hydrophatic*.

HYDRIATRUS, see *Hydrophatic*.

HYDRIODAS KALICUS, see *Potassæ hydriodas*.

HYDRIODIC ACID, see *Acid, hydriodic*.

HYDRO, (from *ὑδωρ*, gen. *ὑδατος*, 'water,')

in composition, water.

HYDRO'A, *Hidro'a*, *Aqu'ula*, *Bo'a*, *Plan'ta noct'is*, (*ὑδωρ*, 'water.')

An affection, which consists in an accumulation of water or serous fluid under the epidermis. Some have used *hydroa* synonymously with *sudamina*; others with *pemphigus*. In the first case, it has generally, however, been written *hidro'a*, from *ἵδρος*, 'sweat,' and in the latter *hydra'a*. See *Hydatid*.

HYDROË'MIA, *Hydræ'mia*, *Hæmydria*; (F.) *Hydrémie*, (*hydro*, and *αἷμα*, 'blood.')

Anæmia. The state of the blood in which the watery constituents are in excess.

HYDROA'ËRIC, *Hydroaë'ricus*, (F.) *Hydroaérique*, (*hydro*, and *αἶρ*, 'air.')

An epithet for the sound rendered on percussing or ausculting cavities in which there are water and air.

HYDROAËROPLEURIE, (*hydro*, *αἶρ*, 'air,' and *πλευρα*, 'side.')

Hydropneumothorax.

HYDROA'ION, *Hydroôph'oron*, *Hydroôphor'ia*, *Hydroôph'ion*, *Hydroôph'rium*, *Ôari'y'drops*, *Hy'drops oca'r'ii*, *Ascit'es oca'r'ii*. *A. eucac'tus*, (F.) *Hydropisie de l'ovaire*, (*hydro*, and *ωαριον*, 'ovarium.')

Dropsy of the ovarium.

HYDROATA, *Sudamina*.

HYDROBLEPH'ARON, *Hy'drops Pal'pebræ*, *Blepharœ'dema aqu'um*, *Edæ'ma palpebr'arum*. An edema or watery swelling of the eyelids; from *hydro*, and *βλεφαρον*, 'eyelid.'

HYDROCACHEXIA, *Leucophlegmatia*.

HYDROCARDIA, (*hydro*, and *καρδία*, 'the heart,')

Hydropericardium, see *Pericardium*.

HYDROCATTARRHOPHE'SIS, *Hydrocattarrhop'h'ia*, (*ὑδωρ*, 'water,' and *καταρροφειν*, 'to sip up.')

Absorption of water from without.

HYDROCE'LE, *Hy'drops Scro'ti seu testicul'um*, *Hydroscelenc'e'lē*, *Hydatu'c'le*, *Hydras'cheum*, *Hydror'chis*, *Hydrosc'hœmie*, (Alibert.) *Hy'drocele*, (*ὑδρωκλῆλη*, from *hydro*, and *κλῆλη*, 'a tumour.')

A term generally applied to a collection of serous fluid in the areolar texture of the scrotum or in some of the coverings, either of the testicle or spermatic cord. To the first of these varieties the names — *External Hy'drocele*, *H. œdematodes*, (F.) *H. par infiltration* have been given; and to the second, those of *Hydroc'e'lē inter'na*, *H. tu'nica vaginal'is test'is*, *Hydr'elytron*, (F.) *H. par épanchement*. When the collection occurs in the envelope of the testicle, it is called *H. of the tu'nica vaginal'is*; and the epithet *congen'ital* is added, when the interior of the membrane, in which it is situate, still communicates freely with the cavity of the abdomen. When it exists in the spermatic chord, it is called *encyst'ed*, or *diffused Hy'drocele of the spermatic cord*, as the case may be. The tumour of the distended scrotum is oblong;

greater below than above; indolent and semi-transparent. When it becomes inconveniently large, the fluid may be evacuated by puncturing with a trocar, but, as it collects again, this operation can only be considered palliative. The radical cure consists, usually, in injecting some irritating liquid through the canula of the trocar, which has been left in after puncturing. This is kept in the tunica vaginalis for a few minutes, and then withdrawn. The coat inflames; adhesion takes place, and the cavity is obliterated.

HYDROCELE OF THE NECK. A tumour, filled with a watery fluid, occupying some portion of the neck.

HYDROCELE PERITONÆI, Ascites—h. Spinalis, Hydrothorax.

HYDROCE'LIC, Hydroce'licus, (ὕδροκελικός.) Relating or appertaining to hydrocele.

HYDROCENO'SIS, (*hydro*, and *κένωσις*, 'evacuation.') The evacuation of water morbidly accumulated in the body.

HYDROCEPHALE, Hydrocephalus.

HYDROCÉPHALE AIGÜE, Hydrocephalus internus.

HYDROCEPHALITIS, (*hydrocephalus*, and *itis*.) Hydrocephalus internus.

HYDROCEPHALUM, Hydrocephalus.

HYDROCEPHALOCELE, (*hydrocephalus*, and *κῆλη*, 'a rupture or protrusion,') Hydrocephalocele.

HYDROCÉPHALOËCTASIE, (*hydrocephalus*, and *εκτασις*, 'extension,') Hydrocephalus chronicus.

HYDROCEPH'ALUS, (ὕδροκεφαλος.) *Hydroceph'alum, Hydrocra'nia, Hydrocra'nium, Hydroceph'alium, Hydroceph'alê, Hy'drops Cap'itis seu Cer'e'brî, (hydro, and κεφαλή, 'the head,') Water of the head, Drop'sy of the head, Drop'sy of the brain.* A collection of water within the head. It may be internal or external.

HYDROCEPHALUS ACUTUS, H. internus—h. Acutus senum, Apoplexy, serous—h. Adnatus, see H. chronicus.

HYDROCEPHALUS CHRON'ICUS, Hydrecephaloce'lê, Hy'drops Cap'itis, seu arachnoïdæ, Hydrocephalus exter'nus, (F.) *Hydrocéphale, Hydrocéphaloëctasie* of some, may exist at birth. *Hydreceph'alon, Hydroceph'alus congen'itus* seu *adnatus* commonly commences at an early period of existence, and the accumulation of fluid gradually produces distension of the brain, and of the skull, with separation of the sutures. It commonly proves fatal before puberty.

HYDROCEPHALUS CONGENITUS, see H. chronicus.

HYDROCEPHALUS EXTERNUS, Ede'ma cap'itis, Cephalœde'ma of some, is a mere infiltration into the subcutaneous cellular tissue of the cranium. See Hydrocephalus chronicus.

HYDROCEPHALUS INTER'NUS, Hydroceph'alus acut'us seu meningæus, Hy'drops cer'e'brî, Enceph'al'gia hydrop'ica, Encephali'tis exsudato'ria, Encephali'tis seu Meningi'tis Infan'tum seu hydroceph'al'ica, Mor'bus cerebra'lis Whyt'tii, Enteroccephal'og'ra Infan'tum, Hydrophlogo'sis Ventriculo'rum cer'e'brî, Encephaloch'y'sis, Phrenic'ula hydroceph'al'ica, Hydreceph'alus, Hydrecephali'tis, Hydreceph'al'ium, Hydrocephali'tis, Hydromeningi'tis, Fe'bris seu Apoplex'ia hydroceph'al'ica, Car'us hydroceph'alus, Water Brain Fe'ver, Whyt't's Disease, (F.) *Hydrocéphale aiguë, Fièvre cérébrale des Enfants,* is generally seated, according to modern observers, in the meninges and surface of the encephalon, and is a tubercular meningi'tis, *Meningi'tis seu Menin'gia tuberculo'sa, Encephalotuberculo'sis, Gran'ular or Scroph'ulous meningi'tis,* (F.) *Méningite tuberculeuse ou granuleuse, Pæden-céphalie* of Piorry. It is observed particularly in childhood. Its march is extremely acute and often very rapid; admitting, generally, however,

of division into three stages. The symptoms of the first stage are those of general febrile irritation, with headache, intolerance of light and sound, delirium, &c. Those of the second, which generally denote that the inflammation has ended in effusion, are, great slowness of pulse, crying out as if in distress, moaning, dilated pupil, squinting, &c.; and lastly, in the third stage—profound stupor, paralysis, convulsions, involuntary evacuations, quick pulse, and frequently death. The disease is of uncertain duration; sometimes, destroying in two or three days; at others, extending to two or three weeks. The prognosis is unfavourable. The treatment must be active during the stage of excitement,—precisely that which is necessary in phrenitis. In the second stage, the indication is;—to promote the absorption of the effused fluid. This must be done by counter-irritants, and mercury, chiefly. On dissection, water is generally found in the ventricles, or at the base of the brain; or there are evidences of previous vascular excitement, as effusions of coagulable lymph, &c.

When the effusion occurs suddenly, either primarily, or secondarily in the course of some other disease, it has been termed, by the Germans, *Wasserschlag, Water Stroke.*

HYDROCEPHALUS MENINGEUS, H. internus—h. Spurius, Hydrocephaloid, (disease.)

HYDROCHAMAIMELUM, (*hydro*, and *chamæmelum*, 'chamomile,') *Infusum anthemidis.*

HYDROCHEZIA, (*hydro*, and *χεζειν*, 'to go to stool,') *Diarrhoea serosa.*

HYDROCHLORATE D'OR, see Gold.

HYDROCHLORINAS NATRICUS, Soda, muriate of.

HYDROCHOLECYSTIS, (*hydro*, *χολη*, 'bile,' and *κυστις*, 'bladder,') *Turgescencia vesiculæ felleæ.*

HYDROCH'YSES, (G.) *Hydrochysen,* (*hydro*, and *χυσις*, 'effusion,') A family of diseases, according to the classification of Fuchs, in which there is a sudden effusion of serous fluid, as in serous apoplexy, hydrocephalus, &c.

HYDROCIRSOCE'LE, Hygrocirsoce'lê, Hygroce'lê, (*hydro*, *κυστις*, 'varix,' and *κῆλη*, 'tumour,') A tumour, formed by the varicose distension of the veins of the spermatic cord, and by the accumulation of serous fluid in the areolar texture of the scrotum.

HYDROCELIA, (*hydro*, and *κοιλία*, 'the abdomen,') *Ascites.*

HYDROCONIUM, (*hydro*, and *κονιος*, 'dusty,') Bath, shower.

HYDROCOT'YLE ASIAT'ICA, (*hydro*, and *κωτυλη*, 'a cup,' from growing in moist places, and the leaves being like cups.) *Codagam, Pes equi'nus, Asiat'ic Pen'nywort.* Order, Umbelliferae. A plant of India, Cape of Good Hope, and Jamaica, where it is supposed to be possessed of astringent and diuretic properties. The leaves are applied to bruised parts to prevent inflammation. The Hydrocotyle has been extolled, in India, as a remedy for lepra.

HYDROCOT'YLÊ CENTE'LA. A South African plant, the roots and stalks of which are astringent; and used in diarrhoea and dysentery.

HYDROCOTYLE UMBELLATUM, Aca'icoba.

HYDROCRANIA, (*hydro*, and *κρανιον*, 'skull,') Hydrocephalus.

HYDROCRANIUM, Hydrocephalus.

HYDROCRITHE, (*hydro*, and *κριθην*, 'barley,') Decoction Hordei.

HYDROCYAN'IC ACID, Ac'idum Hydrocyan'icum: (*hydro*, and *κυανος*, 'blue,') *Pruss'ie Ac'id, Ac'idum Pruss'icum* seu *Boruss'icum* seu *Zoët'icum* seu *Zoët'in'icum, Cyanohy'dric* or *Cyanhy'dric Ac'id,* (F.) *Acide Hydrocyanique* ou *Prus-*

siqne. This acid exists in a great variety of native combinations in the vegetable kingdom, and imparts to them certain properties, which have been long known and esteemed; as in the bitter almond, cherry laurel, leaves of the peach tree, kernels of fruit, pips of apples, &c. When concentrated, it is liquid, colourless, of a strong smell and taste, at first cool, afterwards burning. Its s. g. at 7° Centigrade, is 0.7058. It is very volatile, and enters into ebullition at 80° Fah. It slightly undergoes decomposition, sometimes in less than an hour, and consists of a peculiar gaseous and highly-inflammable compound of carbon and nitrogen, to which the name *Cyanogen* has been assigned; and of hydrogen, which acts as the acidifying principle: hence its name *Hydrocyanic acid*. In the Pharmacopœia of the United States, (1851 and 1863,) two formulæ for the preparation of the *Ac'idum hydrocyanicum dilutum*—*Ac'idum hydrocyanicum*, Ph. U. S. of 1842—are given; the one from the *Ferrocyanide of Potassium*; the other from the *Cyanide of Silver*. Both of these as well as the *Ac'idum hydrocyanicum dilutum* of the British Pharmacopœia (1863), contain 2 per cent. of the anhydrous acid. According to Magendie, the acid, prepared after Scheele's method—the one in common use—is of irregular medicinal power: he, therefore, recommends Gay Lussac's acid, diluted with six times its volume, or 8.5 times its weight of distilled water, for medicinal purposes, and this he calls *Medic'inal Prussic Acid*. Dr. Ure has proposed, that the specific gravity should indicate that which is proper for medicinal exhibition; and, after experiments of the specific gravity of the acids obtained by different processes, he states, that the acid, usually prescribed, is of s. g. 0.996 or 0.997. Great caution is, however, necessary. One drop of pure prussic acid may instantly destroy, and the animal show hardly any traces of irritability, a few moments after death. It has been advised in laryngeal phthisis, in pulmonary phthisis, pulmonary inflammation and irritation, dyspepsia, uterine affections, hectic cough, cancer, chronic rheumatism, and mania, and as a local remedy in impetiginous affections; but, although possessed of powerful sedative properties, it is so unmanageable and the preparation so uncertain, that it is not much used. The *Dose* of Scheele's Acid, or of the *Medicinal Prussic Acid*, is from a quarter of a drop to two drops.

HYDROCYSTIS, (*hydro*, and *κυστις*, 'a bladder.') A cyst containing a watery or serous fluid. A hydatis. Also, saccated ascites.

HYDRODERMA, gen. *Hydroderm'atis*, (*hydro*, and *δερμα*, 'skin.') Anasæra.

HYDRODES, (*ὕδρωδης*.) Aqueous.

HYDRODIARRHŒA, Diarrhœa, serous.

HYDROEDEMA, gen. *Hydrœd'ematis*, Œdema.

HYDRŒNCÉPHALORRHŒE, (*hydro*, 'encephalon,' and *ῥέω*, 'to flow.') Apoplexy, serous.

HYDROËNTEROËPIPOCLE/LĒ, *Hydro'epi-ploënteroce'lē*; (*hydro*, *εντερον*, 'intestine,' and *επιπλοον*, 'omentum.') Enteropiepocele, the sac of which contains a serous fluid.

HYDROËNTEROËPILOMPH'ALUM, (*hydro*, *εντερον*, 'an intestine,' *επιπλοον*, 'the caul,' and *ομφαλος*, 'the navel.') Umbilical hernia, the sac of which contains intestine, epiploon, and serum.

HYDROËNTEROM'PHALUM, *Hydrenterom'phalum*; (*hydro*, *εντερον*, 'an intestine,' and *ομφαλος*, 'the umbilicus.') Hernia umbilicalis, the sac of which contains intestine and serum.

HYDROËPIPOCLE/LĒ, (*hydro*, *επιπλοον*, 'omentum,' and *κληρα*, 'a tumour.') Hernia, formed by omentum, the sac of which contains serum.

HYDROËPIPOËNTEROCELE, *Hydroen-tero-epiplocele*.

HYDROËPILOMPH'ALUM, (*hydro*, *επιπλοον*,

'omentum,' and *ομφαλος*, 'the umbilicus.') Umbilical hernia, the sac of which contains epiploon and serum.

HYDROG'ALA, gen. *Hydrogalu'tis*, (*hydro*, and *γαλα*, 'milk.') A mixture of water and milk.

HYDROGASTER, (*ὕδρογαστηρ*, from *hydro*, and *γαστηρ*, 'the stomach, the belly.') Ascites.

HY'DROGEN, *Hydrogen'ium*, *Inflam'mable air*, *Phlogis'ton*, *Princ'ip'ium hydrogenet'icum* seu *hydro'ticum*, *Meph'itis inflammab'ilis*, (F.) *Hydrogène*, *Phlogogène*, *Air inflammable*, *Gaz des ballons*, (*hydro*, and *γεννω*, 'I produce.') This gas, when breathed, proves fatal from containing no oxygen. When diluted with two-thirds of atmospheric air, it occasions some diminution of muscular power and sensibility, and a reduction of the force of the circulation. It has been respired in catarrh, hæmoptysis, and phthisis.

HYDROGEN, BINOX'IDE or PEROX'IDE OF, *Ox'ygenated water*, is a combination of water with a mixing weight of oxygen. It is liquid, and very prone to decomposition. It is a powerful oxidizing agent, and has been given as an excitant, in asphyxia, cholera and narcotic poisoning; and in cachectic affections,—cancer, diabetes, &c.

HYDROGEN, CARBURET'TED, *Inflam'mable air*, *Firedamp*, of miners, (F.) *Hydrogène carboné ou proto-carboné*, *Grisou*—obtained by passing the vapour of water over charcoal, at the temperature of ignition, in an iron tube—has been found possessed of properties like hydrogen, and has been used, diluted, in like cases. After an explosion of fire damp in mines, there remains the *after damp*, *choke damp*, or *black damp*, consisting of carbonic acid and nitrogen, which occasions a greater loss of life, perhaps, than the explosion.

HYDROGEN, PEROXIDE OF, Hydrogen, binoxide of.

HYDROGEN, PROTOXIDE OF, Water.

HYDROGEN, SULPH'URETTED, *Hydrosulph'uric*, *Sulphhy'dric*, *Sulphohy'dric* or *Hydrothion'ic ac'id*, *Hydrothi'on*, *Gaz hepat'icum* seu *hydrogen'ium sulphura'tum*, *Meph'itis hepat'ica*, (F.) *Hydrogène sulfuré*, *Acide hydrosulfurique*, *Air puant ou hépatique*, may be disengaged from any of the sulphurets by the addition of a strong acid. It is a violent poison, but has been recommended to be inhaled, diluted, to allay the increased irritability which occasionally exists after diseases of the lungs. See Hydrosulphuretted Water.

HYDROGENATION, see *Hydrogénésées*.

HYDROGÈNE, Hydrogen—*h. Carboné* ou *protocarboné*, Hydrogen, carburetted—*h. Sulfuré*, Hydrogen, sulphuretted.

HYDROGÈNÈSES. Baumes gives this name to diseases which he fancifully considers to depend upon disturbed hydrogenation. In it he includes intermittent and remittent fevers.

HYDROGENOSULPHURETUM AMMONIACÆ LIQUIDUM, Ammonie sulphuretum.

HYDROGLOSSA, (*hydro*, and *γλωσσα*, 'tongue.') Ranula.

HYDROGRAPHY, (*hydro*, and *γραφω*, 'to describe.') See Hydrology.

HYDROHÉMIE, (*hydro*, and *αἷμα*, 'blood.') Anæmia, Hydroemæmia.

HYDROHYMENT'IS, *Orrhohymen'itis*, *Orrhymen'itis*, (*hydro*, *ὑμνη*, 'a membrane,' and *itis*.) Inflammation of a serous membrane.

HYDROHYSTERA, (*hydro*, and *ὑστερα*, 'uterus.') Hydrometra.

HYDROLÆUM, *Hydreleoon*.

HYDROLAPATHUM, (*hydro*, and *λαπαθον*, 'the dock.') *Rumex hydrolapathum*.

HYDROLATA, (*ὕδωρ*, 'water.') *Aquæ destillatæ*.

HYDROLATS, *Aquæ destillatæ*, Water, distilled.

HYDROLÉ, Decoction, Fomentation, Lotion, Mucilage, Potion, Tisane. See *Hydroleica*.

HYDROLICA, (F.) *Hydroliques*, ('ὕδωρ, 'water.'). Watery solutions of the active principles of medicinal agents. Those prepared by solution or admixture are termed, by some of the modern French *pharmaciens*, *Hydrolés*; those by distillation, *Hydrolats*.

HYDROLIQUES, *Hydroleica*.

HYDROLITE, Decoction, Mucilage, Potion.

HYDROLOGY, *Hydrolog'ia*, (*hydro*, and *logos*, 'a discourse.'). A treatise on water. By the term *Medical Hydrology* is meant that part of physics, whose object is the study of water, considered as it respects medicine; and, consequently, embracing that of mineral waters. *Medical Hydrography* comprises the study of the influence exerted by the sea or by navigation on the health of man.

HYDROLOTIF, (*hydro*, and *lavare*, *lotum*, 'to wash.'). Fomentation, Lotion.

HYDROMANIA, (*hydro*, and *μανία*, 'mania.'). A name given by Strambi to pellagra, in which the patient has a strong propensity to drown himself.

HYDROMEDIASTINUM, *Hy'drops medias-tini*. Effusion of serous fluid into the mediastinum.

HY'DROMEL, *Hydrom'eli*, (ὕδρομελι, from *hydro*, and *μελι*, 'honey.'). *Aqua mul'sa*, *Meliti'tis*, *Mul'sum*, *Melic'ratum*, *Mellie'ratum*, *Brag'gart*, *Me'do*. A liquid medicine, prepared with an ounce and a half of honey and a pint of tepid water. It is used as a demulcent and laxative, and is generally known under the names *Simple hy'dromel*, *Vinous hy'dromel*, *Mead*, *Hydrom'eli vino'sum*, *Ce'nomel*, is a drink made by fermenting honey and water, and is much used in some countries. As a common drink, mead is often made of honey and some alcoholic liquor well beaten together. See *Mahogany*.

HYDROMELLÉ. A preparation formed like the syrups, using hydromel in place of sugar.

HYDROMENINGITIS, *Aquomembranitis*, *Hydrocephalus internus*.

HYDROMETER, (*hydro*, and *μετρον*, 'a measure.'). *Areometer*.

HYDROMETRA, *Hydrom'e'tria*, *Hydrohys'tera*, *Hydrometrecta'ia*, *Hysteri'tes*, *Hystera'de'ma*, *Hy'drops u'teri*, (*hydro*, and *μητρα*, 'the womb.'). *Drop'sy of the womb*. A disease characterized by circumscribed protuberance in the hypogastrium,—with obscure fluctuation, progressively enlarging, without ischury or pregnancy. If it ever occur, it must be a rare disease.

HYDROMETREXTASIA, (*hydro*, *μητρα*, 'uterus,' and *εκτασις*, 'extension.'). *Hydrometra*.

HYDROM'PHALUS, *Hy'drops umbilica'lis*, *Exom'phalus aquo'sus*, *Her'nia umbil'i ci aquo'sa*, (*hydro*, and *ομφαλος*, 'the navel.'). A tumour, formed by the accumulation of serum in the sac of umbilical hernia; or simply by distension of the navel in cases of ascites.

HYDROMYRIN'GA, *Hydromy'rinx*, *Hydrops tym'pani*, (*hydro*, and *myringa* or *myrinx*, 'the membrana tympani.'). *Drop'sy of the drum of the ear*; giving rise to difficulty of hearing.—*Dysac'a hydrop'ica*.

HYDROMYRINX, *Hydromy'ringa*.

HYDRONCUS, (*hydro*, and *ογκος*, 'a tumour.'). *Anasarca*, *Œdema*.

HYDRONEPHROSIS, (F.) *Hydronephrose*, *Hydror'énale distension*, (*hydro*, νεφρος, 'kidney,' and *osis*.). An accumulation in the kidney, owing to the obstruction of the tubes of the papillæ.

HYDRONOSUS, (*hydro*, and *νοσος*, 'a disease.'). *Hydrops*.

HYDRONUSUS, *Hydrops*.

HYDROOARION, *Hydroarion*.

HYDROPHORON, (*hydro*, ωοφορος, 'egg bearing.'). *Hydroarion*.

HYDROOVARIVM, *Hydroarion*.

HYDROPATHIC, *Hydropath'icus*, *Hydriat'ricus*, (*hydro*, and *παθος*, 'affection.'). Relating to hydrophaty or the water-cure,—as a hydrophathic physician, *Hydriü'ter*, *Hydriü'trus*, *Hydrop'athist*.

HYDROPATHIST, see *Hydrophathic*.

HYDROPATHY, *Hydrosudotherapeia*.

HYDROPEDE'SIS, (F.) *Hydropédè'se*, (from *hydro*, and *πηδω*, 'I spring out.'). *Ephidro'sis*. Excessive sweating.

HYDROPEGE, (*hydro*, and *πηγη*, 'a spring.'). Water, spring.

HYDROPELTIS PURPUREA, (*hydro*, and *πελτη*, 'a buckler,' a waterplant with leaves like buckler), *Brasenia hydropeltis*.

HYDROPERICARDE, *Hydropericardium*.

HYDROPERICARDIA, *Hydropericardium*.

HYDROPERICARDITIS, *Hydropericardium*.

HYDROPERICARDIUM, *Hydropericar'dia*,

Hydropericardi'tis, (*hydro*, and *pericardium*.), *Hy'drops Pericar'dii*, *Hydrocar'dia*, *Drop'sy of the pericar'dium*, (F.) *Hydropéricarde*, *Hy'dropsie du Péricarde*. This is not a common disease. Palpitations; irregular or intermitting pulse; excessive dyspnoea, amounting often to orthopnoea, and dulness over a large space on percussion, will cause the pericardium to be suspected. The treatment is that of dropsies in general. It is, usually, however, of the active kind.

HYDROPERIONE, (*hydro*, περι, 'around,' and *ων*, 'an egg, or ovum.'). The seroalbuminous substance, secreted by the lining of the uterus prior to the arrival of the impregnated ovum in that cavity.

HYDROPERITONEUM, *Ascites*.

HYDROPERITONIE, *Ascites*.

HYDROPHAL'LUS, (*hydro*, and *φαλλος*, 'the male organ.'). *Œdema of the male organ*

HYDROPHIMOSIS, *Phimosis œdematodes*.

HYDROPHLEGMASIA TEXTÛS CELLULARIS, *Phlegmasia alba*.

HYDROPHLOGOSIS VENTRICULORUM

CEREBRI, (*hydro*, and *φλογωσις*, 'inflammation.'). *Hydrocephalus internus*.

HYDROPHOB'IA, ('ὕδροφοβία, from *hydro*, and *φοβος*, 'dread.'). *Paraphob'ia*, *Parophob'ia*, *Phobodip'son*, *Pheu'gydron*, *Phengophob'ia*, *Cynolys'sa*, *Cynolys'sum*, *Mor'bus hydrophob'icus seu hydroph'o-bus*, *Lycan'chæ*, *Lycan'chia*, *Ærophob'ia*, *Phre-ni'tis la'trans*, *Lyl'tia*, *Lys'sa*, *L. cani'na*, *Pantophob'ia*, *Ra'bies cani'na*, *Erethi'smus hydropho-bia*, *Clo'nos hydrophobia*, *Hygrophobia*, *A'que me'tus*, *Canine' mad'ness*, *Wa'terfright*, (F.) *Rage*; *Mal de Saint Hubert*.

The term *Ra'bies* is more appropriate for the aggregate of symptoms resulting from the bite of rabid animals. *Hydrophobia* literally signifies a 'dread of water;' and consequently, ought to be applied to one of the symptoms of rabies, rather than to the disease itself. It is a symptom which appears occasionally in other nervous affections. Rabies is susceptible of spontaneous development in the dog, wolf, cat, and fox, which can thence transmit it to other quadrupeds or to man; but it has not been proved that it can supervene,—without their having been previously bitten,—in animals of other species; or that the latter can, when bitten, communicate it to others. Many facts induce the belief, that the saliva and bronchial mucus are the sole vehicles of the rabid virus; the effects of which upon the economy sometimes appear almost immediately after the bite, and are, at others, apparently dormant for a considerable period. The chief symptoms are— a sense of dryness and constriction of the throat; excessive thirst; difficult deglutition; aversion for, and horror at, the sight of liquids as well as

of brilliant objects; red, animated countenance; great nervous irritability; frothy saliva; grinding of the teeth, &c. Death most commonly happens before the fifth day. Hydrophobia has hitherto resisted all therapeutical means. Those which allay irritation are obviously most called for. In the way of prevention, the bitten part should always be excised, where practicable; and cauterized.

In some cases, symptoms like those which follow the bite of a rabid animal are said to have come on spontaneously. This affection has been termed *nervous* or *spontaneous hydrophobia*.

HYDROPHOBIA, STOX'S MEDICINE FOR, see *Anagallis*.

HYDROPHOBUS, (ὕδροφοβος,) *Lyssodæctus*.

HYDROPTHAL'MIA, (*hydro*, and ὀφθαλμος, 'the eye.') *Hydrophthal'mus*, *Dropsy of the eye*, *Hy'drops Oculi*, *Buphtal'mus*, *Zoophthal'mus*, *Oculus Bovinus* seu *Bubulus* seu *Elephantinus*, *Ophthalmoptosis*, *Paropsis* *Staphylo'ma simplex*, (F.) *Hydrophthal'mie*, *Hydropisie de l'œil*. This affection is caused, at times, by an increase in the quantity of the aqueous, at others, of the vitreous, humour. In the former case, the iris is concave anteriorly, and pushed backwards:—in the latter, it is convex, and pushed forwards. Most commonly, the disease seems to depend on both humours at the same time. *Hydrophthal'mia* sometimes affects both eyes; at others, only one. Children are more exposed to it than adults or old persons. The treatment must vary according to the cause; its longer or shorter duration; greater or less extent, &c. Hence, according to circumstances, hydragogue medicines, purgatives, general and local blood-letting, blisters, setons, moxa, cupping-glasses, fomentations, collyria, and fumigations of different kinds, have been employed. When all means fail, and the disease continues to make progress, the fluid may be evacuated by a puncture made with a cataract needle at the lower part of the transparent cornea.

HYDROPTHAL'MIA, CONICAL, *Staphylo'ma of the cornea*.

HYDROPTHAL'MION. An oedematous swelling of the conjunctiva in hydroptic persons.

HYDROPTHAL'MUS, *Hydrophthal'mia*—h. *Cruentus*, *Hæmophthal'mia*.

HYDROPHYLLUM VERUM, (*hydro*, and φύλλον, 'a leaf,') *Hydrastis Canadensis*.

HYDROPHYSOCELE, (*Hydropneumatocele*, (*hydro*, φυσσα, 'wind,' and κηλη, 'a tumour,') *Hernia*, which contains a serous fluid and gas.

HYDROPHYSOME'TRA, (*hydro*, φυσσα, 'I inflate,' and μητρα, 'the womb,') A morbid condition of the womb, in which both fluid and air are contained in it.

HYDROPI'C, (ὕδρωπικος, from ὕδρωψ, 'dropsy,') *Hydrop'icus*, *Hy'phydros*, *Hyder'icus*, *Hydero'des*, *Hyda'leus*, *Hydrop'sical*, (Sc.) *Edroppit*, (F.) *Hydro'pique*. One labouring under dropsy. *Dropsied*. Relating to dropsy. Also, an antihydroptic, and a hydragogue.

HYDROPIPER, (ὕδροπιπερι, from *hydro*, and πιπερι, 'pepper,') *Polygonum hydro'piper*.

HYDROPSIA, *Hydrops*.

HYDROPSIAE, *Hydrops*—h. *du Bas-ventre*, *Ascites*—h. *Cardiaque*, *Dropsy, cardiac*—h. *Cérébrale suraiguë*, *Apoplexy, serous*—h. *Hépatique*, *Dropsy, hepatic*—h. *de l'Œil*, *Hydrophthal'mia*—h. *de l'Ovaire*, *Hydroo'varion*—h. *du Péricarde*, *Hydropericardium*—h. *du Péritoine*, *Ascites*—h. *des Plèvres*, *Hydrothorax*—h. *de Poitrine*, *Hydrothorax*—h. *Rénale*, *Dropsy, renal*—h. *de la Vésicule du Fiel*, *Turgescentia vesicæ fellæ*.

HYDROPSIS, *Hydrops*—h. *Vera*, *Anasarca*.

HYDROPSMUS, *Hydrops*.

HYDROPLEURIE, (*hydro*, and πλευρα, 'the side,') *Hydrothorax*.

HYDROPNEMATOCELE, (*hydro*, πνευμα, 'air,' and κηλη, 'rupture,') *Hydrophysocèle*.

HYDROPNEMON, *Hydropneumonia*.

HYDROPNEMON'IA, (*Hydropneumonia*, *Œde'ma pulmo'num chron'icum*, *Hy'drops pulmo'num* seu *pulmo'num cellulo'sus*, *Anasar'ca pulmo'num*, (*hydro*, and πνευμων, 'the lung,') *Dropsical infiltration of the lungs*. See *Œdema of the lungs*.

HYDROPNEMOSAR'CA, (*hydro*, πνευμα, 'wind, air,' and σαρξ, 'flesh,') An abscess, containing water, air, and matters similar to flesh.

HYDROPNEMOTHOR'AX, gen. *Hydropneumothorac'is*, *Hydrocœdœopleuric*, (*hydro*, πνευμων, 'the lung,' and θωραξ, 'the chest,') *Pneumothorax* with effusion of blood into the chest. See *Pneumothorax*.

HYDROPOIDES, (ὕδρωποιδης, from *hydro*, and αἶς, 'likeness,') An epithet for watery excretions, such as sometimes take place in hydrotics.

HYDROPO'TA, *Hydropotes*, (ὕδωρος, from *hydro*, and ποτις, 'a drinker,') *Pota'tor Aquæ*, *A water drinker*. One who drinks only water, or drinks it in an extraordinary quantity.

HY'DROPS, gen. *Hydro'pis*, (ὕδρωψ, from ὕδωρ, 'water,') *Phlegma'tia*, *Hy'deros*, *Hyderi'asis*, *Affectus hydero'des*, *Hydrop'is*, *Hydrop'sia*, *Hydrop'simus*, *Hydrop'sia*, *Hydrorrhœ'a*, *Hydror'hœ*, *Hydror'hœius*, *Polyhy'dria*, *Hydron'o'mus*, *Hydroni'us*, *Dropsy*, *Hy'dropsy*, (Old Eng.) *Dro'p'ik*, (F.) *Hydropisie*. A preternatural collection of a serous fluid in any cavity of the body, or in the areolar texture. When the areolar texture of the whole body is more or less filled with fluid, the disease is called *Anasar'ca* or *Leucophlegma'tia*;—and when this variety is local or partial, it is called *Œde'ma*. The chief dropsies, designated from their seat, are:—*Anasar'ca*, *Hydrocephalus*, *Hydrorachitis*, *Hydrothorax*, *Hydropericardium*, *Ascites*, *Hydrometra*, *Hydrocele*, &c.

Encysted Dropsy, *Hy'drops sacca'tus*, *inaccerat'us* seu *cyst'icus*, is that variety in which the fluid is enclosed in a sac or cyst: so that it has no communication with the surrounding parts. *Dropsy of the Ovarium*, *Hydrops* seu *Ascites Ovari*, *Ascites sacca'tus*, is an instance of this variety.

Dropsy may be active or passive. The former consists in an increased action of the exhalants, so that those vessels pour out much more fluid than is absorbed: the latter arises from a state of atony of the absorbent vessels, which allows of an accumulation of fluid. It may also be mechanical, or produced by obstructions to the circulation, as in cases of diseased liver. Active dropsy, occurring accidentally in a sound individual, generally ends favourably. That which supervenes on other diseases, or is symptomatic of some internal affection, is rarely curable.

The treatment consists in the use of all those remedies which act on the various secretions: so that, the demand being increased, the supply will have to be increased accordingly; and in this manner some of the collected fluid may be taken up by the absorbents. To this end bleeding if the dropsy be very active; purgatives, diuretics, sudorifics, sialagogues, &c., are the remedies chiefly depended upon.

HYDROPS ABDOMINIS, *Ascites*—h. *Abdominis acereus*, *Tympanites*—h. *Abdominis sacca'tus*, see *Ascites*—h. *Anasarca*, *Anasar'ca*—h. *Anasarca acuta*, see *Anasarca*—h. *Arachnoideæ*, *Hydrocephalus chronicus*—h. *Articulorum*, *Hydrarthrus*—h. *Ascites*, *Ascites*—h. *Capitis*, *Hydrocephalus*, *Hydrocephalus chronicus*—h. *Cardiacus*, *Dropsy, cardiac*—h. *Cavitatis columnæ vertebralis*, *Hydrorachis*—h. *Cellularis artuum*, *Œdema*—h. *Cellularis totius corporis*, *Anasar'ca*—h. *Cellulosus*, *Anasar'ca*—h. *Cerebri*, *Hydrocephalus*, *Hydrocephalus internus*—h. *Cu-*

taneus, Anasarca—h. Cysticus, see Ascites, and Hydrops—h. Glottidis, Œdema of the Glottis—h. Hepaticus, Dropsy, hepatic—h. Incarceratus, see Hydrops—h. Intercus, Anasarca—h. Leucophlegmatia, Leucophlegmatia—h. Matellæ, Diabetes—h. ad Matulam, Diabetes—h. Mediastini, Hydro-mediastinum—h. Medullæ spinalis, Hydrorachis—h. Oculi, Hydrophthalmia—h. Ovarii, Hydro-sarion, see Hydrops—h. Palpebræ, Hydroblepharon—h. Pectoris, Hydrothorax—h. Pericardii, Hydro-pericardium—h. Pleuræ, Hydrothorax—h. Pulmonis, Hydrothorax—h. Pulmonum, Hydro-pneumonia, Œdema of the Lungs—h. Pulmonum cellulosus, Hydro-pneumonia—h. Renalis, Dropsy, renal—h. Saccatus, see Hydrops—h. Sacci lacrymalis, Fistula lacrymalis—h. Seroti, Hydrocele—h. Siccus et flatulentus, Tympanites—h. Spinæ, Hydrorachis—h. Spinæ vertebralis, Hydrorachis—h. Subcutaneus, Anasarca—h. Telæ cellulose, Anasarca—h. Testiculorum, Hydrocele—h. Thoracis, Hydrothorax—h. Tubarum Fallopii, Hydrosalpinx—h. Tympani, Hydromyringa—h. Tympanites, Tympanites—h. Umbilicalis, Hydrophalmarum—h. Uteri, Hydrometra—h. Vesicæ fellæ, Turgescencia vesiculæ fellæ.

HYDROPSIA, Hydrops.

HYDROPSY, Hydrops.

HYDROPYR'ETOS, (*hydro*, and *πυρετος*, 'fever;') *Febris sudatoria*. Fever with sweating. *Hydropyretos* would be more proper; from *ἰδρω*, 'sweat.' See *Sudor Anglicus*.

HYDRORA'CHIS, (*hydro*, and *ραχis*, 'the spine;') *Hydrorrhachis*, *Hydrorachitis*, *Hydrorrhachia*, *Myeloch'ysis*, *Hy'drops Cavita'tis Colum'næ Vertebr'al'is* seu *Sp'inx Vertebr'al'is* seu *Medullæ spin'al'is* seu *Sp'inx*, *Hydrocel'æ spin'al'is*, *Hydrorach'itis spin'o'sa*. An effusion of serum, often owing to inflammation of the spinal membranes—*myel'itis exsudat'iva*—and forming a soft, frequently transparent, tumour, constituted of the membranes of the spinal marrow, which are distended and projecting backwards from the vertebral canal, the posterior paries of which, when the affection is congenital, is wanting to a certain extent—*Sp'inx bif'ida*, *Atelorachid'ia*, *Hydrora'chis dehiscens seu congen'ita*, *Schistorrhach'is*, *Sp'inxola*. The disease is often accompanied with paralysis of the lower extremities. It is congenital, and situate in the lumbar or sacral regions. It is almost always fatal:—the tumour rupturing, and death occurring instantaneously. On dissection, a simple separation or complete absence of the spinous processes of the vertebrae is perceived, with, at times, destruction or absence of spinal marrow. The treatment is the same as in *hydrocephalus chronicus*; and, as in it, advantage seems occasionally to have been derived by puncturing with a fine needle.

HYDRORACHIS DEHISCENS, see Hydrorachis.

HYDRORACHITIS, gen. *Hydrorachitidis*, Hydrorachis.

HYDRORCHIS, gen. *Hydror'chidis*, (from *hydro*, and *ορχis*, 'a testicle;') Hydrocele.

HYDRORE'NALE DISTENSION, Hydro-nephrosis.

HYDRORRHACHIA, Hydrorachis.

HYDRORRHACHIS, Hydrorachis—h. Congenita, see Hydrorachis—h. Dehiscens, see Hydrorachis.

HYDRORRHAGIE, (*hydro*, and *ῥυγμι*, 'to break forth;') Apoplexy, serous.

HYDRORRHŒ, Hydrops.

HYDRORRHŒA, (*hydro*, and *ῥεω*, 'to flow;') Hydrops. Also, a copious watery discharge,—as from the vagina.

HYDRORRHŒUS, Hydrops.

HYDRORTHOPNŒ'A, (*hydro*, and *ορθοπνοια*, 'difficulty of breathing; except in the erect posture;') Orthopnoea, owing to a collection of water in the chest.

HYDROSAC'CHARUM, *A'qua sacchar'ata*, (F.) *Eau sucrée*. Sugared water. Syrupus simplex.

HYDROSAL'PINX, gen. *Hydrosalpin'gis*, *Hy'drops tubarum Fallo'pii*; (*hydro*, and *σάπις*, 'a tube;') Dropsy of the Fallopian tube.

HYDROSAR'CA, (*hydro*, and *σαρξ*, 'flesh;') A tumour containing a fluid, as well as portions of flesh. Also, Anasarca.

HYDROSARCOCE'LE, (*hydro*, *σαρξ*, 'flesh; and *κλη*, 'a tumour;') *Sarcocydrocele*. A tumour, formed by a sarcocele, complicated with dropsy of the tunica vaginalis.

HYDROSCŒOCE'LE, (*hydro*, *σχεο*, 'the scrotum; and *κλη*, 'rupture;') A collection of water in the scrotum. Hydrocele.

HYDROSCŒONIE, Hydrocele.

HYDROSCEUM, (*hydro*, and *σχεον*, 'the scrotum;') Hydrocele.

HYDROSIS, Hidrosis.

HYDROSTATIC BED, Water bed—h. Test of Infantieide, see Docimasia.

HYDROSUDOPATHY, (*hydro*, *sudo*, 'I sweat; and *παθος*, 'affection;') Hydrosudotherapeia.

HYDROSUDOTHERAPEIA, *Hydrop'athy*, *Hydropath'ia*, *Hydrosudop'athy*, *Hydr'asis*, *As hydriat'rica*, *Hydriat'ria*, *Hydatotherape'ia*, *Hydrotherape'ia*, *Watercure*, [G.] *Wasserkur*, (*hydro*, *sudo*, 'I sweat; and *θεραπεω*, 'I cure;') A badly compounded word, formed to express the mode of treating diseases systematically by cold water, sweating, &c.

HYDROSULPH'URETTED WATER, *A'qua hydrosulphur'ata simplex*, *A. hepatic'a*, (F.) *Eau hydrosulphurée simple*. (Sulphuret of iron 1000 parts, sulphuric acid 2000 parts, distilled water 4000 parts; add the water to the acid, and put the sulphuret of iron into a retort, to which a Wolff's apparatus of five or six vessels is adapted; the last containing about an ounce of potassa, dissolved in a quart of water. Pour the diluted acid gradually on the sulphuret, and ultimately throw away the water in the last vessel. *Ph. P.*) It is stimulant, diaphoretic, and deobstruent [?], and is used in rheumatism, diseases of the skin, &c.

It has been also called *Ac'idum Hydrothion'icum bi'uidum*.

HYDROSULPHURETUM AMMONIACALE AQUOSUM, Ammonia sulphuretum—h. Ammoniacum. Ammonia sulphuretum.

HYDROTHERAPEIA, (*hydro*, and *θεραπεω*, 'I cure;') Hydrosudotherapeia.

HYDROTHION, (*hydro*, and *θειον*, 'sulphur;') Hydrogen, sulphuretted.

HYDROTHO'RAX, gen. *Hydrothorac'is*, (*hydro*, and *θωραξ*, 'the chest;') *Hy'drops Thora'cis* seu *pector'is seu pulmo'nis seu pleur'æ*, *Stethoch'ysis*, *Pleurorrhœ'a lymphat'ica* seu *sero'sa*, *Dyspnœ'a* et *Orthopnœ'a hydrothorac'ica*, (F.) *Hydropisie de Poitrine* ou des *Plevres*, *Dropsy of the Chest*. Idiopathic hydrothorax, termed by Laënnec, *Hydropisie des plevres*, *Dropsy of the Pleuræ*—by Piorry, *Hydropleurie*—is a rare disease, and not easy of diagnosis. It generally exists only on one side, which, if the fluid effused be considerable, projects more than the other. Dyspnœa, and fluctuation perceptible to the ear, are characteristic symptoms. When the chest is examined with the stethoscope, respiration is found to be wanting every where except at the root of the lung. The sound is also dull on percussion.

Effusion into the chest, as a result of inflammation of some thoracic viscera, is as common as the other is rare. It is usually a fatal symptom. It has been called *symptomatic hydrothorax*.

In hydrothorax, the course of treatment proper in dropsics in general must be adopted. Diure-

may occasion a rupture or destruction of this membrane. It is often, indeed, found absent in children soon after birth; whilst it may remain entire after copulation. Hence the presence of the hymen does not absolutely prove virginity; nor does its absence prove incontinence, although its presence would be *primâ facie* evidence of continence.

HYMEN, (ὑμην,) Membrane—h. Diaphragmaton, Mediastinum.

HYMENÆA, (after the god Hymen,) see Copal—h. Courbaril, see Anime.

HYMEN'ICA AMENORRHÆ'A. Amenorrhœa occasioned by closure of the hymen.

HYMEN'ICUS, *Hymenodes*. Relating to the hymen. Also, membranous.

HYMEN'ITIS, (*hymen*, and *itis*.) Membranous inflammation. Inflammation of an internal membrane.

HYMEN'IUM, (ὑμενιον, dim. of *hymen*.) *Membranula*. A fine, delicate membrane.

HYMENODES, (ὑμενώδης,) Hymenicus.

HYMENOGANGLIITIS, Cholera.

HYMENOG'ENY, *Hymenogenia*, (F.) *Hyménogénie*, (*hymen*, and γεννα, 'I engender.') The production of membranes by the effect of simple contact of two liquids, as albumen and fat, when the former gives a coating to the globules of the latter. (Ascherson.)

HYMENOG'RAPHY, *Hymenographia*, (*hymen*, and γραφο, 'I describe.') That part of anatomy whose object is the description of the different membranes.

HYMENOL'OGY, *Hymenolog'ia*, (*hymen*, and λογος, 'a description.') A treatise on the membranes.

HYMENOPHTHALMIA, Ophthalmia membranarum.

HYMENOR'RAPHY, *Hymenorrhaphia*, (*hymen*, and ραφη, 'a suture.') A form of elytrorrhaphy, in which the operation is performed in the natural situation of the hymen.

HYMENOT'OMY, *Hymenotomia*, (*hymen*, and τεμνω, 'I cut, I dissect.') The part of anatomy which treats of the dissection of the membranes. The term has also been applied to the incision of the hymen, practised in certain cases of imperforation of the vagina, in order to give exit to the blood retained and accumulated in the cavity of the uterus.

HYMNIUM, (contracted from *hymenium*.) Amnios.

HYO, in composition, an abridgment of *Hyoides os*.

HYOBASIOGLOSSUS, Basioglossus.

HYOCHONDROGLOSSUS, Hyoglossus.

HYODEOGLOSSUS, Hyoglossus.

HYODEOTHYREODES, Thyreohyoides.

HYODEPIGLOTTICUS, Hyoepiglotticus.

HYODES, Hyoides.

HYOPIGLOT'TICUS, *Hyodepiglot'ticus*. Belonging to the os hyoides and epiglottis. Some anatomists have given the name *Hyo-epiglottic ligament* to a bundle of condensed areolar tissue, which passes from the posterior part of the body of the hyoid bone to the base of the epiglottic fibro-cartilage.

HYOGLOSSO-BASI-PHARYNGIEN, Constrictor pharyngis.

HYOGLOS'SUS, *Hyodeoglossus*, *Hyochondroglossus*, *Hyopiglossus*, *Ceratoglossus* of Douglass and Cowper: *Basioceratochondroglossus*. A large, thin, quadrilateral muscle, situate at the anterior and superior part of the neck. Its insertions at three different points of the os hyoides permit it to be divided into three portions:—the first (*Ceratoglossus* of Albinus), is attached to the great cornu of the os hyoides; the second (*Basioglossus* of Albinus), arises from the superior part of the body of the same bone; and the third (*Chondro-*

glossus of Albinus), arises from the lesser cornu and the cartilage situate between the body and the greater cornu. The fibres of these three bundles are inserted into the lateral and inferior parts of the tongue. This muscle depresses the base of the tongue, or raises the os hyoides, when the tongue is fixed.

HYOTHYREOÏDEUS, Thyreohyoides.

HYOTHYROID, Thyreohyoid.

HYOID BONE, Hyoides, os.

HYOÏDES OS, (ὑοειδής, the Greek ὑ and εἶδος, 'shape.') *Os Bicornè* seu *hypseloï'des* seu *Lambdoi'des* seu *Gut'turis* seu *Gutturālē* seu *Lingualē* seu *Lingual'tē*, *Ypsiloï'des*, *Ypsiloïdes*. The *Hyoid Bone*, *Hyoides*, *Hyo'des*. The hyoid bone is a very movable osseous arch; of a parabolic shape, convex before, and suspended horizontally in the substance of the soft parts of the neck, between the base of the tongue and the larynx. This bone, separated entirely from the rest of the skeleton, is composed of five distinct portions, susceptible of motion on each other. The first and most central is the body of the hyoid, *Ossiculum me'dium* *Hyoï'dis*, which affords attachment to several muscles; the two others are lateral, and bear the name of *branches* or *greater cornua*. The last two are smaller, situate above the other, and are known under the name *lesser cornua* and *styloid cornua*, the *Os'sa pisiformia lingualia* of Sömmerring, *Os'sa seu Gra'n'a seu Ossic'ula* seu *Cor'pora* seu *Corpus'cula trit'ica*. The os hyoides is ossified from five points.

HYOÏDES PRIMUS, Sterno-hyoides.

HYOÏDEUS, Hyoides.

HYOÏDIS QUARTUS MUSCULUS, Omohyoides.

HYOSCY'AMUS, (ἵς, 'a swine,' and κωμος, 'a bean,' eaten by swine.) *Fa'ba suil'la*, *Ben'gi*, *Jusqui'amus*, *Hyoscyamus* *u'ger* seu *agres'tis*, *Apollina'ris*, *Alter'cum*, *Ag'onē*, *Altercan'genon*, *Hen'bane*, *Poison Tobac'co*, *Stink'ing night'shade*. (F.) *Jusquiamè*, *Fève a Cochon*, *Hannebane*, *Potée*, *Porcelet*. Ord. Solanaceæ. The leaves, *Hyoscy'ami Fo'lium*, and seeds, *Hyoscy'ami Semen*, Ph. U. S. 1863, are the parts used in medicine. Their odour is narcotic and peculiar; taste insipid and mucilaginous. The virtues are yielded to proof spirit. *Hyoscyamus* is narcotic, anodyne, antispasmodic, and slightly stimulant. It is used as a substitute for opium, where the latter disagrees; and is applied, externally, as a cataplasm in cancer and glandular swellings. Dose, gr. iij to x of the powder. An alkaloid was obtained from it by Brandes, *Hyosciamine*, *Jusquiamine*.

HYOSCYAMUS AGRESTIS, *Hyoscyamus*.

HYOSCYAMUS ALBUS, *White Hen'bane*, possesses similar virtues.

HYOSCYAMUS LUTEUS, *Nicotiana rustica*—h. Niger, *Hyoscyamus*—h. Peruvianus, *Nicotiana tabacum*.

HYOSCYAMUS SCOPO'LIA, *Scopoli'na atropoï'des*. The herb and root of this plant, which grows in Illyria, Hungary, Croatia, Gallizia, and Bavaria, are used in the same cases as belladonna. Dose of the powder, half a grain.

HYPACTICUS, (ὑπακτικός, from *hypo*, and αγω, 'I drive,') Cathartic.

HYPE'MIA, (*hypo*, and αἷμα, 'blood,') *Hypoæ'mia*, *Hypohæ'mia*, *Hypohæ'mia*, *Oligæ'mia*, *Oligohæ'mia*, *Anæ'mia*, (F.) *Hyphémie*. Deficiency of blood. Also, extravasation of blood.

HYPEREMIA OCULI, Hæmophthalmia.

HYPA'GOGE, (ὑπαγωγῆ, from *hypo*, and αγω, 'I drive,') Dejection.

HYPA'GOGUS, (ὑπαγωγός,) Laxative.

HYPALEIM'MA, gen. *Hypaleim'matis*, (*hypo*, and αλειψω, 'I anoint,') *Hypalim'ma*, *Hypaleip'tum*, *Hypalip'tum*. An ointment or liniment to be rubbed or spread on a part.

HYPALEIPTRIS, (ὑπαλειπτρίς.) Hypaleiptrum. HYPALEIP'TRON, (ὑπαλειπτρον.) *Hypaleip'trum*, *Hypaleiptris*, *Hypalip'trum*, *Specil'lum*, *Spatha*. A sort of spatula for spreading ointments.

HYPALEIPTRUM, *Hypaleiptron*.

HYPALEIPTUM, *Hypaleimma*.

HYPALIMMA, *Hypaleimma*.

HYPALIPTRUM, *Hypaleiptron*.

HYPAMAURO'SIS, (*hypo*, and *amaurosis*;) *Amauro'sis imperfecta*. Imperfect amaurosis; *Μεταμαυρο'sis*.

HYPANCONIUM, (*hypo*, and *αγκον*, 'the elbow.') A pillow for supporting the elbow in surgical cases.

HYAPOPLEX'IA, (*hypo*, and *apoplexia*.)

An incomplete attack of apoplexy.

HYPATISMUS, *Fumigation*.

HYPATMUS, (ὑπατμος, from *hypo*, and *ατμος*, 'vapor.') *Fumigation*.

HYPAUCHE'NIUM, (*hypo*, and *αυχνη*, 'the neck.') A pillow or cushion for the neck.

HYPECCHORESIS, (ὑπεκχορσις, from *hypo*, *ec*, and *χωρεω*, 'to make vacant.') *Dejection*.

HYPECCHORETICUS, *Laxative*.

HYPELATUS, (*hypo*, and *ελαυνω*, 'to drive,') *Cathartic*, *Laxative*.

HYPENTIOMIA, (*hypo*, and *εναντιος*, 'opposite.') *Allopathy*.

HYPENTIOSIS, *Allopathy*.

HYPE'NE, (ὑπηνη.) The beard which grows under the chin, according to some. Also, the upper lip. See *Mystax*.

HYPENETES, (ὑπηνητης.) *Adolescents*.

HYPER, ὑπερ, 'above,' 'in excess.'

HYPERAUC'SIS, *Oxyecō'a*, *Hyperac'oë*, *Plithogodypho'ria*, (F.) *Hyperacusie*, (*hyper*, and *ακη*, 'audition.') Excessive sensibility of the organ of hearing.

HYPERADENO'SIS, *Hyperadeno'ma*, *Hyper'troph'ia glandula'rum*, (*hyper*, *αδην*, 'a gland,' and *osis*.) *Hypertrophy* of a gland.

HYPERÆ'MIA, *Hyperhæ'mia*, *Hyperæmo'sis*, *Hæmormæ'sis*, (F.) *Hyperémie*, *Hyperhémie*, *Angiohémie*, (*hyper*, and *αἷμα*, 'blood.') *Preternatural accumulation* of blood in the capillary vessels, more especially local *plethora*—*pletho'ra partialis*; congestion. Various forms of hyperæmia are admitted by pathologists,—for example, the *active* or *sthenic*; the *asthenic* or *passive*; the *cadaveric*, or that which forms immediately before or after death; the *hypostatic*, which occurs in depending parts, and the *mechanical*, which is produced by some mechanical impediment.

HYPERÆMIA ABDOMINIS, *Cœliemia*—h. *Activa*, *Inflammation*—h. of the Brain, *Cephalæmia*—h. *Capitis*, *Cephalæmia*—h. *Cerebri*, *Cephalæmia*—h. *Pectoris*, *Stethæmia*—h. *Hepatic*, *Hepatohæmia*—h. *Pulmonum*, *Stethæmia*—h. of the Lungs, *Stethæmia*.

HYPERÆMIC, *Hyperæmicus*, *Hyperæ'mial*, *Congest'ed*, *Blood'shot*, (F.) *Hyperémie*. Affected with hyperæmia.

HYPERÆMOSIS, *Hyperæmia*.

HYPERÆSTHESIA, (*hyper*, and *αισθησις*, 'the faculty of feeling.') *Hyperæsthesis*—h. *Linguae*, *Hypergæstia*—h. *Olfactoria*, *Hyperosphresia*—h. *Plexus cardiaci*, *Angina pectoris*—h. *Psychica*, *Hypochondriasis*.

HYPERÆSTHES'IS, *Hyperæsthes'ia*, *Oxyæsthes'ia*, (F.) *Hyperæsthésie*. Excessive sensibility, impressibility, or passibility.

HYPERÆSTHET'ICA. Agents that are conceived to augment general sensibility—as *strychnia*, *brucia*, &c.

HYPERANARRHOPHE'SIS, (*hyper*, and *αναρροησις*, 'absorption.') Excessive absorption.

HYPERANTHERA MORINGA, (*hyper*, 'in excess,' and *anthera*, 'anther,') *Guilandina moringa*.

HYPERANTHRAXIS, (*hyper*, and *ανδραξ*, 'a coal,') see *Cholera*.

HYPERAPH'IA, *Oxyaph'ē*, *Oxyaph'ia*, (*hyper*, and *αφη*, 'touch.') Excessive acuteness of touch.

HYPERAPHRODIS'IA, (*hyper*, and *αφροδιτη*, 'Venus.') Excessive venereal desire.

HYPERASTHEN'IA, (*hyper*, and *ασθενεια*, 'weakness.') Excessive debility.

HYPERAUXE'SIS, (*hyper*, and *αυξησις*, 'augmentation.') *Hyperepid'osis*. Excessive increase or enlargement of a part, as *Hyperauxe'sis I'ridis*, an excessive enlargement of the iris, so as to stop up the pupil.

HYPERBOL'IC (*attitude*), (ὑπερβολικος, from *hyper*, and *βαλλω*, 'I throw;' 'excessive.') *Galen*, by this term, designates certain extraordinary attitudes, in which the limbs and vertebral column are in a state of complete extension or flexion.

HYPERBO'REAN, *Hyperbor'eus*, (ὑπερβορεος, from *hyper*, and *βορεας*, 'the north wind.') A race of men found at the extreme north of the two continents, in the vicinity of the polar circle. It includes the Thibetans, Ostiaks, Kamtschadales, Laplanders, Samoiedes, Esquimaux, &c.

HYPERBU'LIA, (*hyper*, and *βουλη*, 'will.') Ungovernable will or volition.

HYPERCARDIA, (*hyper*, and *καρδια*, 'heart,') *Heart*, *hypertrophy* of the.

HYPERCARDIOTROPHIA, (*hyper*, *καρδια*, 'heart,' and *τροφη*, 'nourishment.') *Heart*, *hypertrophy* of the.

HYPERCATAPINO'SIS, (*hyper*, *καταπινειν*, 'to sip up,' and *osis*.) Excessive activity of absorption.

HYPERCATHAR'SIS, *Hyperine'sis*, *Hyper'inos*, *Superpurgat'io*, *Ejffrenitatio*, (*hyper*, and *καθαρος*, 'purgation.') *Superpurgation*.

HYPERCENO'SIS, (*hyper*, and *κνωσις*, 'evacuation.') Excessive evacuation, as of blood, bile, &c.

HYPERCERASIS, (*hyper*, and *κερας*, 'the cornea,') *Staphyloma* of the cornea.

HYPERCERATOSIS, *Staphyloma* of the cornea.

HYPERCHOLIA, (*hyper*, and *χολη*, 'bile,') *Polycholia*.

HYPERCHRO'MA, gen. *Hyperchro'matis*, (*hyper*, and *χρωμα*, 'colour,') A red fleshy excrescence at the inner angle of the eye, near the *caruncula lacrymalis*, which sometimes grows very large, so as to separate the eyelids and interfere with their movements.

HYPERCHROMATOPS'IA, *Hyperchromatop'sy*, (*hyper*, *χρωμα*, gen. *χρωματος*, 'colour,' and *οψις*, 'vision.') A defect of vision, owing to which ideas of colour are attached to objects, which convey no such coloured impressions to a healthy eye. It is the antithesis to *achromatopsia*.

HYPERCINE'SIA, *Hypercine'sis*, (*hyper*, and *κινω*, 'I move,') 'motion.') Excessive motion.' Under the term *hypercinesis*, *Romberg* includes the spasmodic neuroses.

HYPERCINESIA HYSTERIA, *Hysteria*—h. *Nervosa*, see *Irritable*—h. *Uterina*, *Hysteria*.

HYPERCINESIS GASTRICA, *Hypochondriasis*.

HYPERCONJUNCTIVITE, see *Ophthalmia*.

HYPERCORYPHO'SIS, (*hyper*, and *κορυφη*, 'the vertex;' 'the extreme point of anything,') The lobes of the liver and lungs.

HYPERCRIN'IA, *Hyperdia'crisis*, *Chymoze'mia*, (*hyper*, and *κρινω*, 'I separate,') A morbid increase in the quantity of the secretions.

HYPER'CRISIS, *Superæcre'tio*, *Superævacua'tio*, *Superævacua'tio*. An excessive crisis, or evacuation; a flux.—A very violent, critical effort, or too copious critical evacuations.

HYPERCYEMA, gen. *Hypercy'e'matis*, (*hyper*, and *κλειν*, 'to conceive,') *Superfoetation*.

HYPERCYESIS, Superfoetation.

HYPERDERMATO'SIS, *Hyperdermato'ma*, *Hyperdermo'sis*, *Hyperderma'ma*, (*hyper*, *δύω*, 'skin,' and *osis*.) Hypertrophy of the skin.

HYPERDIACRISIS, (*hyper*, *δια*, and *κρίνω*, 'I separate.') Hypercrinia.

HYPERDIURESIS, *Hyperdiuresis*, Diabetes.

HYPERDYNAMIA, (*hyper*, and *δύναμις*, strength,) *Hypersthenia*.

HYPERDYNAMIC, *Hyperdynamicus*. Appertaining to or having the characters of hyperdynamia, or excessive strength—of the vital powers more especially.

HYPERCHEMA, gen. *Hyperche'matis*, (*hyper*, and *ἤκω*, 'to sound,') See Exaggerated.

HYPERCHESIS, See Exaggerated.

HYPEREMESIS, *Hyperemes'ia*, *Vom'itus profusus*, (*hyper*, and *εἶρω*, 'I vomit,') Excessive vomiting.

HYPERÉMIE, *Hyperæmia*—*h. Cérébrale*, *Cephalæmia*—*h. du Cerveau*, *Cephalæmia*—*h. du Foie*, *Hepatohæmia*—*h. Hypostatique*, see *Hypostatic*—*h. de la Moëlle épinière*, *Hypermyelohæmia*—*h. des Poumons*, *Stethæmia*.

HYPERÉMIE, *Hyperæmie*.

HYPERENCÉPHALOTROPHIE; (*hyper*, *εγκεφαλον*, 'the encephalon,' and *τροφή*, 'nourishment,') Hypertrophy of the encephalon of Piorry.

HYPERENCEPHALUS, (F.) *Hyperencéphale*, (*hyper*, and *encephalon*.) A monster whose excessive brain is situate in the skull.

HYPERENDOSMOSE, Inflammation.

HYPERENERGIA, (*hyper*, and *ενέργεια*, 'activity,') Excessive activity, as of the nervous system.

HYPEREPHIDROSIS, *Ephidrosis*.

HYPEREPIDROSIS, (*hyper*, and *επιδροσις*, 'increase,') *Hyperauxesis*.

HYPERERETHIS'IA, (*hyper*, and *ερεθίζω*, 'I excite,') Excessive irritability.

HYPERESTHÉSIE, *Hyperæsthesis*.

HYPERFIBRINATION, *Hyperinosis*.

HYPERGENESIS, (*hyper*, and *γενεσις*, 'generation,') (F.) *Hypergénésie*, *Hyperplastie*. The excess of formative power, which gives occasion to monstrosities by excess of parts. See *Monster*.

HYPERGEUS'TIA, *Hypergeus'is*, *Hyperæsthésia ling'ue*, *Oxygeus'ia*, (*hyper*, and *γεύω*, 'taste,') Excessive sensibility of the organ of taste.

HYPERHÆMATOSIS, Inflammation.

HYPERHÆMIA, *Hyperæmia*.

HYPERHÉMIE, *Hyperæmia*.

HYPERHÉPATOTROPHIE, (*hyper*, *ἥπαρ*, gen. *ἥπατος*, 'liver,' and *τροφή*, 'nourishment,') *Hepatauxes*.

HYPERHIDROSIS, (*hyper*, and *ἰδρώσις*, 'sweating,') *Epidrosis*.

HYPERHORA, (*hyper*, and *ῥα*, 'manhood,') Premature development of the body, or of some part.

HYPERICUM, or HYPERICUM BACCIFERUM, (*ὑπερικόν*), *Arbutuscula gummi'fera Braziliensis*, *Caa-opia*. *Ord.* *Hypericaceæ*. A Brazilian tree, whose bark emits a juice, when wounded, which resembles gamboge. *Hypericum* is considered by Linnæus to be derived from *ὑπερ*, 'over,' and *εἶκω*, 'an image or spectre;' from its having been fancied to possess the power of driving away evil spirits.

HYPERICUM OFFICINALE, *H. perforatum*—*h. officinarum*, *H. perforatum*.

HYPERICUM PERFORATUM, *Hypericum*, *H. officinale* seu *officina'rum* seu *vulga'rè* seu *Virgin'icum*, *Fu'ga De'monum*, *Andros'mum*, *Co'rion*, *Perforated* or *Common St. John's Wort*, (F.) *Milepertuis ordinaire*. It is aromatic and astringent, and enters into a number of aromatic preparations; and, amongst others, into the *Falltranks*. It was, at one time, much used in dysentery, hem-

orrhages, &c. The juice is acrid. The *Oil of St. John's Wort*, *Red oil*, *Linimentum seu O'leum hyperici*, *Balsamum hyperici simplex*, is made by infusing $\frac{3}{4}$ lb of the flowers in a quart of olive oil. It is applied to recent bruises.

HYPERICUM SAROT'HRA, *O'angegrass*, *Pineverew*, indigenous, is applied in domestic practice, boiled, to contusions and sprains.

HYPERICUM VIRGINICUM, *H. perforatum*—*h. Vulgare*, *H. perforatum*,

HYPERIDROSIS, *Ephidrosis*.

HYPERINESIS, (*ὑπερῖσσις*, from *hyper*, and *ἵκω*, 'to purge,') *Hypercatharsis*.

HYPERINOS, (*ὑπερινός*), *Hypercatharsis*.

HYPERINO'SIS, *Ino'sis*, *Hyperplas'ma*, (*hyper*, *ἵς*, gen. *ῖνος*, 'fibre,' and *osis*.) The condition of the blood in which it contains an increase in the proportion of fibrin, *Hyperfibrin'ation*, *Superfibrin'ation*, a decrease of the ratio of red corpuscles, with excess of fibrin, and an increase of the fat,—as in inflammation. In proportion to the increase of the fibrin and fat, and the decrease of the corpuscles, the whole solid residue will be diminished. Also, morbidly increased muscular activity.

HYPERINOT'IC, *Hyperinot'icus*. Appertaining or relating to hyperinosis.

HYPERLYMPH'IA, (F.) *Hyperlymphie*, (*hyper*, and *lymph'a*.) Excessive formation or accumulation of lymph.

HYPERMÉTROHÉMIE, *Metrohæmia*.

HYPERMETROP'IA, (*hyper*, *μετρον*, 'a measure,' and *οὐρα*, 'vision,') *Presbyopia*.

HYPERMNE'SIA, (from *hyper*, and *μνασται*, 'I recollect,') Excessive memory.

HYPERMYELOHÆ'MIA, (F.) *Hyperémie de la Moëlle Épinère*, *Congestion sanguine rachidienne*, (*hyper*, *μυελος*, 'marrow,' and *αἷμα*, 'blood,') *Hyperæmia* of the spinal marrow.

HYPERNÉPHROTROPHIE, (*hyper*, *νέφρος*, 'kidney,' and *τροφή*, 'nourishment,') Hypertrophy of the kidney.

HYPERNEUR'IA, (*hyper*, and *νευρον*, 'a nerve,') Excessive nervous activity.

HYPERNEURO'MA. Morbid development of the neurine or nervous masses.

HYPEROPHARYNGEUS, *Palatopharyngeus*.

HYPERO'A, (*ὑπερωα*, from *hyper*, and *ωον*, 'a high place,') The palatine arch,—the base of the cranium; the palate.

HYPEROÏ'TIS, *Inflammat'io pala'ti*, *Anji'na Palati'na*, (F.) *Inflammation du Palais*; (*hyperoia*, and *itis*.) Inflammation of the velum palati.

HYPEROÏCHAS'MA, gen. *Hyperoïchas'matis*, (*hyperoia*, and *χάσμα*, 'an opening,') *Lycost'oma*, *Palat'um fissum*. Fissure of the palate.

HYPEROPSIA, (*hyper*, and *οψίς*, 'vision,') *Oxyopia*.

HYPEROS, (*ὑπερος*), *Pilum*.

HYPEROSMIA, (*hyper*, and *οσμή*, 'smell,') *Hyperosphresia*.

HYPEROSPHERE'SIA, *Hyperos'mia*, *Hyperæsthésia olfacto'ria*, *Olfac'tus acut'us*, (*hyper*, and *οσφρησις*, 'smell,') Excessive acuteness of the smell.

HYPEROSTOSIS, (*hyper*, *οστέον*, 'a bone,' and *osis*.) *Exostosis*.

HYPERPATH'IA, (*hyper*, and *πάθος*, 'affection,') Excessive sensibility in disease.

HYPERPHLEBO'SIS, (*hyper*, *φλέβη*, gen. *φλέβος*, 'a vein,' and *osis*.) Too great development of the venous system; predominant venosity.

HYPERPHLEGMASIA, *Hyperphlogosis*.

HYPERPHLOGO'SIS, *Epiphlogo'sis*, *Hyperphlegma'cia*, *Inflammat'io peracuta*, (*hyper*, and *φλογωσις*, 'inflammation,') A high degree of inflammation.

HYPERPHRÉNIE, (*hyper*, and *φρην*, 'the mind,') *Mania*.

HYPERPIMELE, (*hyper*, and *πυμλη*, 'fat,') see Polysarcia.

HYPERPLASMA, gen. *Hyperplasm'atis*, (*hyper*, and *πλασσω*, 'to form,') Hyperinosis.

HYPERPLASIA, (*hyper*, and *πλασσω*, 'to form,') Excess of formative action.

HYPERPLASTIE, Hypergenesis.

HYPERPLEXIE, (*hyper*, and *πλησσω*, 'to strike,') Ecstasis.

HYPERPNEUSTIA, (*hyper*, and *πνεω*, 'to blow,') Flatulence.

HYPERPRESBYTIA, Presbytia.

HYPERSARCO'MA, gen. *Hypersarco'matis*, *Hypersarco'sis*, *Hypersarx'is*, (*hyper*, and *sarco-ma*.) A soft fungous excrescence, especially such as appears upon ulcerated parts:—a fungosity.

HYPERSARCOSIS, (*hyper*, and *sarcosis*), Excrescence, Fungosity—h. Cordis, Heart, hypertrophy of the.

HYPERSARXIS, (*hyper*, and *σαρξ*, 'flesh,') Hypersarcoma.

HYPERSPADIAS, (*hyper*, and *σπασω*, 'to draw,') Epispadias.

HYPERSPASMIA, (*hyper*, and *σπασμος*, 'spasm,') Convulsion.

HYPERSPLENOTROPHIE, (*hyper*, *splen*, and *τροφή*, 'nourishment,') Splenoncus.

HYPERSPONGIA, (*hyper*, *σπογγος*, 'a sponge, a fungus,') Spina ventosa.

HYPERSTHENIA, *Hyperdyna'mia*, *Hyperzōdyna'mia*, *Stai'tus inflammato'rius ve'trus*, (*hyper*, and *σθενος*, 'strength,') Super-excitement. A morbid condition, characterized by over-excitement of all the vital phenomena.

HYPERSTHENIC, Stimulant.

HYPERSTHENICUS, see Active, Stimulant.

HYPERTON'IA, (*hyper*, and *τονος*, 'tone,') Excess of tone in parts. It is opposed to atony.

HYPERTRICHOSIS, (*hyper*, and *τριχ*, gen. *τριχος*, 'hair,') Excessive production of hair.

HYPERTROPHÆ'MIA, (*hyper*, *τροφή*, 'nourishment, and *αίμα*, 'blood,') A state in which the plastic powers of the blood are increased.

HYPERTROPHIA, (*hyper*, and *τροφή*, 'nourishment,') Hypertrophy—h. Cerebri, Phrenauxe—h. Cordis, Heart, hypertrophy of the—h. Glandularum, Hyperadenosis—h. Hepatis, Hepatauxe—h. Intestinorum, Enterhypertrophia—h. Lienis, Splenoncus—h. Splenis, Splenoncus—h. Unguini, Onychauxe—h. Uteri, Metrauxe—h. Vesicæ urinariæ, Cystauxe.

HYPERTROPHIÆ, see Hypertrophy.

HYPERTROPHIE, Hypertrophy—h. du Cœur, Heart, hypertrophy of the—h. Centripète du Cœur, Heart, hypertrophy of the—h. Concentrique du Cœur, Heart, hypertrophy of the—h. du Foie, Hepatauxe—h. de la Rate, Splenoncus—h. du Corps Thyroïde, Bronchocèle.

HYPERTROPHIED, see Hypertrophy.

HYPERTROPHIEN, see Hypertrophy.

HYPERTROPHOUS, see Hypertrophy.

HYPERTROPHY, or HYPERTROPHY, *Hypertroph'ia*, *Supernutri'tio*, (F.) *Hypertrophie*, *Supernutrition*. The state of a part in which the nutrition is performed with greater activity; and which, on that account, at length acquires unusual bulk. The part thus affected is said to be *hyper'trophied* or *hyper'trophous*. *Hypertrophie* is used by Neligan, as *Hypertrophien* is by Simon, for an order of cutaneous diseases, characterized by hypertrophy of the derma, or epidermis, or both, or of the hair follicles.

HYPERURESIS, (*hyper*, and *ουρον*, 'urine,') Enuresis—h. Aquosa, Diabetes.

HYPERURORRHÉE, (*hyper*, *ουρον*, 'urine, and *ρρω*, 'to flow,') Diabetes—h. Saccharine, Diabetes (mellitus.)

HYPERZOODYNAMIA, (*hyper*, *ζωον*, 'animal, and *δυναμις*, 'power,') Hypersthenia.

HYPEX'ODUS, (*ὑπεξοδος*, from *hypo*, and *εξοδος*, 'an exit,') An alvine discharge or flux.

HYPEZOCUS, (*hypo*, and *ζωννυμι*, 'to gird,') Diaphragm, Pleura. See Hypozoma.

HYPHA, (*ὑψη*), Texture.

HYPIIÆMA, gen. *Hypihæ'matis*, (*hypo*, and *αίμα*, 'blood,') Ecchymoma.

HYPIÆMIA, Hyæmia.

HYPHÆMATOSIS, *Hypohæmato'sis*, (*hypo*, and *αίματωσις*, 'sanguification,') Morbidly diminished hæmotosis.—*Sanguificatio debili'ior*.

HYPHÆMOS, (*hypo*, and *αίμα*, 'blood,') Subcurrentus.

HYPHE, (*ὑψη*), Texture.

HYPHÉMIE, Hyæmia.

HYPHYDROS, (*hypo*, and *ὑδωρ*, 'water,') Hydropic.

HYPINO'SIS, *Hypoplas'ma*, (*hypo*, and *ισ*, *ινος*, 'fibre,') The condition of the blood in which the quantity of fibrin is frequently less than in health, or if it amounts to the usual quantity, its proportion to the blood corpuscles is less than in health: the quantity of corpuscles is either absolutely increased or their proportion to the fibrin is larger than in healthy blood; the quantity of solid constituents is also frequently larger than in health. Such is the condition of the blood in fevers, hemorrhages, and polyæmia.

HYPNOTICAL, *Hypnotic'us*. Relating or appertaining to hypnosis.

HYPNÆSTHESIS, (*hypnos*, and *αἰσθησις*, 'feeling,') Somnolency.

HYPNAGOG'IC, *Hypnago'gius*, (*hypnos*, and *αγωγος*, 'a leader,') Leading to sleep.

Hypnagogic hallucinations are those visions which occur when we are half asleep.

HYPNIA'TER, (F.) *Hypniatre*, (*hypnos*, and *ιατρος*, 'a physician,') A name given to deluded or designing persons who have affirmed that they were able, during their 'magnetic sleep,' to diagnose disease and its appropriate treatment.

HYPNIC, *Hypnic'us*, (*ὑπνικος*, from *hypnos*.) An agent that affects sleep.

HYPNOBADES, (*hypnos*, and *βαίνω*, 'to move,') Somnambulist.

HYPNOBADISIS, Somnambulism.

HYPNOBASIS, Somnambulism.

HYPNOBATAISIS, Somnambulism.

HYPNOBATES, Somnambulist.

HYPNOBATESIS, Somnambulism.

HYPNOBATA, Somnambulism.

HYPNODES, (*ὑπνωδης*, from *ὑπνος*, 'sleep, and *οδης*.) One in a state of slumber or somnolency.

HYPNODIA, (*ὑπνωδία*), Somnolency.

HYPNOLOG'ICA, *Hypnolog'icæ*, (*hypnos*, and *λογος*, 'a discourse,') The part of hygiene which treats of sleep.

HYPNOL'OGY, *Hypnolog'ia*, (F.) *Hypnologicæ*, (*hypnos*, and *λογος*, 'a discourse,') A treatise on sleep. The doctrine of sleep.

HYPNONERGIA, (*hypnos*, and *ενεργεια*, 'action,') Somnambulism.

HYPNOPHOB'IA, (*hypnos*, and *φοβος*, 'dread,') Dread of sleep. Incubus.

HYPNOPEÛS, (*ὑπνοποιος*, from *hypnos*, and *ποιω*, 'to make,') Somniferous.

HYPNOS, (*ὑπνος*), Sleep.

HYPNOSIS, (*hypnos*, and *osis*.) The super-vention or production of sleep.

HYPNOSIS BIOMAGNETICA, Sleep, magnetic.

HYPNOTIC, Somniferous.

HYPNOTIQUE, Somniferous.

HYPNOTISM, (from *hypnos*,) see Magnetism, animal.

HYPNOTIZED, see Mesmerized.

HYPO, (*ὑπο*, 'sub,') Under, in composition.

HYPO, Hypochondriasis.

HYPOÆMA, gen. *Hypoæ'matis*, (*hypo*, and *αίμα*, 'blood,') Ecchymoma.

HYPOÆMIA, Echy-moma, Hypæmia.

HYPOBLEPHARUM, (*hypò*, and *blepharòs*, eyelid.) Tumefaction under one or both eyelids. Also, an artificial eye, placed under the eyelids.

HYPOCAPNISMA, gen. *Hypocapnismatiss*, (*ὑποκαπνισμα*, from *hypo*, and *καπνός*, 'smoke,') Fumigation, Suffummentum.

HYPOCAPNISMUS, (*ὑποκαπνισμός*,) Fumigation.

HYPOCARO'DES, *Sulapara'tiss*, (*ὑποκαρώδης*, from *hypo*, *καρρός*, 'a heavy sleep,' and *odes*,) One who is in a state approaching carus.

HYPOCATHAR'SIS, (*ὑποκαθάρσις*, from *hypo*, and *καθάρσις*, 'purgation,') Feeble purgation. Opposed to hypercatharsis.

HYPOCATHARTICUS, Laxative.

HYPOCAUSTUM, (*ὑποκαύστης*, from *hypo*, and *καίω*, 'I burn,') A name given to a stove, or any such contrivance, to sweat in. Also, a furnace in any subterraneous place, used for heating baths:—*ὑποκαυστήριον*, *Balneo'rium*, *Vapora'rium*.

HYPOCEPHALÆ'UM, (*ὑποκεφαλαίον*, from *hypo*, and *κεφαλή*, 'head,') A pillow for the head.

HYPOCERCHALÆON, (*ὑποκερχαλίον*, from *hypo*, and *κερχαλίος*, 'hoarse,') Roughness of the fauces affecting the voice.

HYPOCHLORETUM SULPHUROSUM, Sulphur, chloride of.

HYPOCHLOROMELAS, *sub pallide nigricans*, (*hypo*, *χλωρός*, 'green,' and *μέλας*, 'black,') A term applied to one whose skin is pale, with a blackish hue.

HYPOCHONDRE, *Hypochondrium*, *Subcartilagin'eum*, *Re'gio Hypochondri'aca*, *Hypochondria'ca Re'gion*, (*ὑποχόνδριον*, from *hypo*, and *χόνδρος*, 'a cartilage,') Each lateral and superior region of the abdomen is so called, because it is bounded by the cartilaginous margin of the false ribs, which forms the base of the chest. There is a right and a left hypochondrium.

HYPOCHONDRIAC, (*ὑποχονδριακός*,) *Hypochondri'acus*, *Hypochondri'acal*, *Vaporous*, *Vaporous*, (F.) *Hypochondriacque*. Belonging to hypochondriasis. One labouring under hypochondriasis: hippel.

HYPOCHONDRIACISMUS, Hypochondriasis.

HYPOCHONDRIALGIA, (*hypochondrie*, and *άλγος*, 'pain,') Hypochondriasis.

HYPOCHONDRIQUE, Hypochondriac.

HYPOCHONDRIASIS, *Alu'cia seu Hallucina'tio hypochondriasis*, *Mor'bus hypochondri'acus seu resicent'ria seu rustic'ana*, *Mor'bus hypochondri'acum*, *Hypochondriaci'smus*, *Hypochondriaci'smus*, *Dyspeps'ia hypochondriaca*, *Pari'sis seu Af'fec'tio hypochondri'aca*, *Hallucina'tio hypochondri'aca*, *Ana'sthesi'sis*, *Hypocri'sis*, *Hyper'æsthes'ia seu ch'ica*, *Sp'ies*, *Melanchol'ia ner'cea seu flatu's seu hypochondri'aca*, *Suffoca'tio hypochondri'aca*, *Mor'bus flatu's seu flatulent'is seu erudit'um*, *For'ces ventriculi*, *Mirach'ia*, (from Arab. *mirach*, 'the belly,') *Morbus mirach'ia*, *Hypochondri'acum*, *Hypo*, *Hip*, *Spleen*, *Vapors*, *English Melancholy*, *Low Sp'ies*, (F.) *Hypochondria*, *Maladie imaginatione*, on *Anglais*, *Affection Vaporeuse*, *Vapours*, *V. de Rate*. This disease is probably so called, from the circumstance of some hypochondriacs having felt an uneasy sensation in the hypochondriac regions. The disease seems really to be, as Pinel has classed it, a species of neurosis, or of mental alienation, which is observed in persons who in other respects are of sound judgment, but who reason erroneously on whatever concerns their own health. Hypochondriasis is characterized by disordered digestion, without fever or local lesion; Estulence; borborygmi; extreme increase of sensibility; palpitations; illusions of the senses: a succession of morbid feelings, which appear to simulate the greater part of diseases: panics; exaggerated uneasiness of various kinds: chiefly in what regards the health, &c. Indiges-

tion has usually been considered the cause of hypochondriasis. They are, unquestionably, much connected with each other: but there is every reason to believe, that the seat of the affection is really, though functionally, in the brain. The disease almost always appears at the adult age, most commonly in irritable individuals; and in those exhausted, or rather in the habit of being exhausted, by mental labour, overwhelmed with domestic or public affairs, &c.

The treatment is almost entirely moral. The condition of the digestive function must, however, be accurately attended to.

HYPOCHONDRICISMUS, Hypochondriasis.

HYPOCHONDRISM, Hypochondriasis.

HYPOCHONDRUM, Hypochondre.

HYPOCHOREMA, gen. *Hypochorematis*, (*ὑποχόρημα*, from *hypo*, and *χόρηω*, 'to make vacant,') Excrement.

HYPOCHORESIS, (*ὑποχόρησις*,) Dejection, Defecation, Excrement.

HYPOCHORETICUS, Cathartic.

HYPOCHYMA, gen. *Hypochymatiss*, (*ὑποχυμα*, from *hypo*, and *χυω*, 'to pour,') Cataract.

HYPOCHYSIS HÆMATODES, (*ὑποχυσίς*, from *hypo*, and *χυω*, 'to pour,') Hæmophthalmia.

HYPOCISTIS, (*ὑποκίστις*,) Cytinus.

HYPOCLEPTICUM, (*ὑποκλέψω*, and *κλέω*, 'I steal,') A chymical vessel, formerly used for separating oil from water.

HYPOCLYSIS, *Hypoclysis*, (*ὑποκλυσίς*, and *ὑποκλύω*, from *hypo*, and *κλύω*, 'to wash out,') as by a clyster. Cleansing by a clyster.

HYPOCELIS, (*ὑποκέλη*,) Palpebra inferior.

HYPOCELIUM, (*ὑποκέλιον*, from *hypo*, and *κέλη*, 'abdomen,') Abdomen.

HYPOCELUM, Hypocoilon, Palpebra inferior.

HYPOCOILON, *Hypocoilon*, *Hypocoilon*, (*ὑποκοίλον*, and *κεῖλον*, 'a cavity,') A cavity situate under the lower eyelid. The lower eyelid itself.

HYPOCOPHOSIS, (*ὑποκοπός*, and *κωφός*, 'deaf,') Barroquia: Deafness.

HYPOCRANIUM (APOSTEMA), (*hypo*, and *κρανίον*, 'the cranium,') A collection of pus between the cranium and dura mater.

HYPOCRATES, (*hypo*, and *κρατος*, 'strength,') Infirm.

HYPOCYLUM, Hypocoilon, Palpebra inferior.

HYPOCYPHUS, (*ὑποκυφός*, from *hypo*, and *κυφός*, 'gibbous,') Slightly humpedback.

HYPOCYSTOTOMIA, (*hypo*, *κύστις*, 'bladder,' and *τομή*, 'section,') see Lithotomy.

HYPODERIS, (*ὑποδερίς*, from *hypo*, and *δέρμις*, 'skin,') Epideris.

HYPODERMATOMY, *Hypodermatom'ia*, (*hypo*, *δέρμα*, 'the skin,' and *τομή*, 'incision,') The section of subcutaneous parts, as of tendons and muscles.

HYPODERMIC, *Hypodermic'us*, (F.) *Hypodermique*, (*hypo*, and *δέρμα*, 'skin,') That which is under the skin.—as *hypodermic medication*, (F.) *Médication hypodermique*; that which consists in the application of remedies—as morphia—under the skin.

HYPODERMIS, Clitoris, Epideris.

HYPODERRHIS, (see *Hypodermis*,) Epideris. Also, the lower and fore part of the neck, (from *hypo*, and *δέρμα*, 'neck,')

HYPODESIS, or HYPODESIS, (*ὑποδεδίς*, or *ὑποδεδίς*, from *hypo*, and *δέω*, 'to bind,') see Bandage.

HYPODESMA, gen. *Hypodesmatiss*, (*ὑποδέσμα*, from *hypo*, and *δέω*, 'a bandage,') see Bandage.

HYPODYNAMIC, (*hypo*, and *δυναμις*, 'power,') Adynamic.

HYPODYN'IA, (*hypo*, and *δύω*, 'pain,') *Doler m'ia seu l'ia*. A slight pain.

HYPOGALA, (*hypo*, and *γάλα*, 'milk,') Hypopyon.

HYPOGASTRAL'GIA, (*hypogastrium*, and *άλγος*, 'pain,') Pain in the hypogastrium.

HYPOGAS'TRIC, *Hypogas'tricus*, *Hypogas'trius*, (ὕπογαστρος,) (F.) *Hypogastrique*. Relating or belonging to the hypogastrium.

HYPOGASTRIC ARTERY, *A. Ili'aca inter'na*, *A. Ili'aca post'erior*, (F.) *Artère hypogastrique*, *A. pelvienne*—(Ch.) is the more internal of the two branches into which the primary iliac divides. It descends into the cavity of the pelvis, and gives off a considerable number of branches, which arise, at times, separately; at others, by common trunks. These branches are, 1. The *posterior*, i.e. the ilio-lumbar arteries, lateral, sacral, and gluteal. 2. The *anterior*, i.e. the umbilical, vesical, and obturator. 3. The *interval*,—the middle hemorrhoidal arteries, uterine, and vaginal in women. 4. The *inferior*, i.e. the *ischiatric arteries*, and internal pudic.

HYPOGASTRIC GAN'GLION. A large nervous ganglion, described by Dr. Robt. Lee as seated on each side of the cervix uteri, immediately behind the ureter; which receives the greater number of the nerves of the hypogastric and sacral plexuses, and distributes branches to the uterus, vagina, bladder, and rectum.

HYPOGASTRIC OPERATION OF LITHOT'OMY, (F.) *Traille hypogastrique*. The high operation, or that practised above the pubes.

HYPOGASTRIC PLEX'US, *Plexus hypogastricus* or *sous-mésentérique* of Winslow, is situate at the lateral and posterior parts of the rectum and the *bas fond* of the bladder. It is formed by the sacral nerves and the inferior mesenteric plexus, and gives off numerous filaments, which accompany the arteries that pass to the rectum and genital organs.

HYPOGASTRIC VEIN, (F.) *Veine hypogastrique*, furnishes nearly the same branches.

HYPOGASTRION, (ὕπογαστριον,) Abdomen, Hypogastrium.

HYPOGASTRIUM, *Hypogas'trion*, (*hypo*, and *γαστηρ*, 'the stomach or belly;') *Etron*, *Are'a*, *Stachys* [?], *Ven'ter i'mus*, *V. par'vus*, *Aqualic'ulus*, *Su'men*, *Ru'men*; the lower part of the abdomen. The *Hypogastric re'gion*, *Re'gio hypogas'trica* seu *hypogas'tria*, which extends as high as three fingers' breadth beneath the umbilicus, is divided into three secondary regions—one *middle* or *pubic*, and two *lateral* or *inguinal*.

HYPOGASTRIUS, Hypogastric.

HYPOGASTROCE'LE, (*hypogastrum*, and *κηλη*, 'a tumour.') Hernia in the hypogastric region, occurring through the separated fibres of the lower part of the linea alba. See *Hernia*, hypogastric.

HYPOGASTRODID'YMUS, *Did'yimus Sym'phyohypogas'trius*, *Ischiopa'ges*, (*hypogastrum*, and *διώμος*, 'a twin.') A monstrosity in which twins are united by the hypogastrium.

HYPOGASTRORRHEXIS, (*hypogastrum*, and *ῥήξις*, 'rupture,') Eventration.

HYPOGLOSSA, Hypoglossides.

HYPOGLOSSAL NERVE, Hypoglossus.

HYPOGLOSSIA, Hypoglossides.

HYPOGLOSSADENITIS, (*hypo*, *γλωσσα*, 'tongue,' *ἀδην*, 'a gland,' and *itis*.) *Inflammat'io glandula'rum sublingua'lium*. Inflammation of the sublingual gland.

HYPOGLOSSIDIA, Hypoglossides.

HYPOGLOSSIS, (ὕπογλωσσις,) *Hypogloss'ium*, *Hypoglot'tia*, *Hypoglot'tia*, (*hypo*, and *γλωσσα*, 'tongue,') The under surface of the tongue; anything under the tongue; and, hence, *Ranula*.

HYPOGLOSSIUM, Hypoglossis, *Ranula*.

HYPOGLOSSUM, (ὕπογλωσσον,) *Ruscus hypoglossum*, *Ranula*.

HYPOGLOS'SUS, (ὕπογλωσσος,) That which is under the tongue.

HYPOGLOSSUS, *Lingua'lis Me'dius*, *Hypogloss'al* or *Ling'al* or *Gus'tatory Nerve*. (F.) *Nerv' Hypogloss'e* ou *Grand Hypogloss'e*, *Hypogloss'ien* (Ch.), is the *ninth pair* of nerves, *Par' no'um*, *Ner'vus*

hypogloss'us seu *lo'quens*, of many anatomists. It arises by ten or twelve very fine filaments from the grooves, which separate the corpora pyramidalia from the corpora olivaria; issues from the cranium by the foramen condyloideum anterius; and divides, near the angle of the jaw, into two branches; the one, the *cervica'lis descendens* or *descen'dens noni*, which forms, with the cervical plexus, a large anastomotic arch, and furnishes branches to several of the muscles of the neck. The other, the *lingual branch*, is the continuation of the principal trunk, and gives its numerous filaments to the muscles of the tongue and pharynx. The ninth pair communicates motion to the muscles to which it is distributed.

HYPOGLOTTA, Hypoglossis, Hypoglossides.

HYPOGLOTTIDES (PIL'ULÆ,) *Hypogloss'ia*, *Hypoglot'tia*, *Hypogloss'ia*, *Hypoglot'tia*, *Hypogloss'idia*, *Pil'ulæ sublingua'les*. Pills placed under the tongue to dissolve there.

HYPOGLOTTIS, (ὕπογλωττις,) Hypoglossis, *Ranula*.

HYPOGLU'TIS, (ὕπογλωστις,) (*hypo*, and *γλουτος*, 'the breech,') The lower and projecting part of the nates.

HYPOGNATHADEN, (*hypo*, *γναθος*, 'the jaw,' and *ἀδην*, 'a gland,') Submaxillary gland.

HYPOGNATHADENITIS, *Hypoglossadenitis*, (*hypognathaden*, and *itis*.) Inflammation of the submaxillary gland.

HYPOHE'MA, gen. *Hypohe'matis*, (*hypo*, and *αἷμα*, 'blood,') Effusion of blood into the chambers of the eye. It is generally absorbed in time.

HYPOHEMATOSIS, Hypæmatis.

HYPOHE'MIA, Hypæmia.

HYPOLEPSIOMANIA, (*hypo*, *ληψις*, 'a seizure,' and *mania*,) Melancholy.

HYPO'MIA, (*hypo*, and *ωμος*, 'the shoulder,') That which lies under the shoulder, as the axilla.

HYPOMO'RIA, (*hypo*, and *μωρια*, 'folly,') Slight imbecility of mind, or slight delirium.

HYPONARTHÉCIE, *Hyponarth'e'cia*, (*hypo*, and *ναρθήξ*, 'a splint,') A term used by M. Mayor for his mode of treating fractures by position only,—the limb resting upon a properly cushioned board or splint.

HYPONEUR'IA, (*hypo*, and *νευρον*, 'a nerve,') Morbidly diminished nervous energy.

HYPON'OMUS, (ὕπονους, from *hypo*, and *νεμος*, 'I feed,') *Ulcus subtile depas'cens*. A deep fistula or ulcer.

HYPON'YCHON, (*hypo*, and *ονυξ*, gen. *ονυχος*, 'a nail,') *Eechymo'ma Hyponychon*. Effusion of blood under a nail.

HYPOPATH'IA, *Subaffec'tio*, (*hypo*, and *παθος*, 'affection,') A disease of a slight character.

HYPOPATUS, (*hypo*, and *πατω*, 'to go,') Dejection.

HYPOPE'DIUM, (*hypo*, and *pes*, 'the foot,') A cataplasm for the sole of the foot.

HYPOPH'ASIS, (*hypo*, and *φαίνω*, 'I appear,') The state of the eyes in which the white only is seen through the opening of the eyelids.

HYPOPHLEGMASIA, Subinflammatio.

HYPOPH'ORA, (*hypo*, and *φέρω*, 'I carry;') *Ulcus sinu'osum* seu *fistulo'sum*. A fistulous ulcer, also, a dejection.

HYPOPHOSPHITES, see Calcis hypophosphis.

HYPOPTHALMIA, Hypopyon.

HYPOPTHAL'MION, (*hypo*, and *ὀφθαλμος*, 'the eye,') That part under the eye where oedema generally commences in chronic diseases and in cachexia.

HYPOPHYSIS, (*hypo*, and *φωω*, 'to bring forth,') Cataract, Epigone—h. Cerebri, Pituitary gland.

HYPOPION, see Hypopyon.

HYPOPITYS LANUGINO'SA, (*hypo*, and *πitys*, 'a pine tree,') *American Pine-sap*, *Falsæ*

Beech-drops, Birds' Nest. Indigenous: flowering from June to August. *Order*, Ericaceae. Used as a nerve in the form of the powdered root.

HYPOPIUM, (ὑποπιον,) (*hyppo*, and *ωψ*, 'eye,') The part of the face under the eye:—a black eye. *HYPOPIUM OS*, Mala os.

HYPOPLASMA, gen. *Hypoplasma'tis*, (*hyppo*, and *πλασμα*, 'to form,') Hypinosis.

HYPOPLASTEMA, (F.) *Hypoplastic*, (*hyppo*, *πλαστικός*, 'formative,' and *αἷμα*, 'blood,') Diminished plasticity of the blood.

HYPOPLEURIUS, (*hyppo*, and *πλευρα*, 'a rib,') Pleura.

HYPOPO'DIA, *Supplanta'tia*, *Suppeda'nea*, (*hyppo*, and *πους*, 'the foot,') Remedies, as sinapisms, which are applied under the foot.

HYPOPYON, *Hypopyum*, *Hypophthal'mia*, *Pyophthal'mia*, *Empye'sis* seu *Diapye'sis* seu *Abcessus Oculi*, *Oculus purulentus*, *Lunel'la*, *Hypog'ala*, *Hypopyum lacteum*, *Pyo'sis*, *Parop'sis* *Staphylo'ma purulentum*; (*hyppo*, and *πυς*, 'pus,' because the pus is under the cornea.) This name has been given to small abscesses between the laminae of the cornea, as well as to different purulent collections in the chambers of the eye; hence, some pathologists have distinguished *Hypopyon of the chambers* from *Hypopyon of the cornea*. In abscesses of the chambers, the purulent matter is mixed with the aqueous humour, which it renders turbid; and is deposited particularly at the lower part of the eye; forming behind the cornea a kind of whitish crescent, that rises more or less before the pupil, and closes it entirely or in part. It requires the use of local and general antiphlogistics, and sorbafacients. At times, it is necessary to puncture the cornea and evacuate the pus.

Hypopyon was used by Galen to signify "an effusion of blood under the eyelids and tunica conjunctiva, or under the lower eyelid." (Blancard.)

HYPOPYUM LACTEUM, *Hypopyon*.

HYPORINION, *Hyporrhinium*, (ὑπορίνιον, from *hyppo*, and *ῖον*, 'the nose,') That part of the beard which grows beneath the nose. The mustaches. Also, the upper lip.

HYPORRHA'CHIS, (ὑπορραχίς, from *hyppo*, and *ραχίς*, 'the spine, the back,') The hollow in the small of the back.

HYPORRHHYSIS, (*hyppo*, and *ῖω*, 'to flow,') Defluxion, Prolapsus.

HYPOSA'PRUS, *Sub'putria*, (*hyppo*, and *σάπρος*, 'putrid,') *Putres'cens*. Growing putrid; slightly putrid. Game, of various kinds, (F.) *Viandes fisan'des*, is, at times, preferred in an incipient state of decomposition, which certainly renders it more digestible.

HYPOSA'CA, (*hyppo*, and *σαξ*, 'flesh,') *Hypocarcid'us*, *Anasarca*. In the Nosologies of Linnæus and Cullen, it is synonymous with *Phycocoin*.

HYPOSARCIDIUS, *Anasarca*, *Hyposarca*.

HYPOSARCO'SIS, (*hyppo*, and *σαρκωσις*, 'a fleshy growth,') A small, fleshy growth:—a wart.

HYPOSIAGON, (*hyppo*, and *σιγαν*, 'the jaw-bone,' 'lower jawbone,') Maxillary bone, inferior.

HYPOSIAGONARTHRI'TIS, (*hyposiagon*, *αρθρον*, 'a joint,' and *itis*,) *Inflamma'tio articuli maxillæ inferioris*. Inflammation of the joint of the lower jaw.

HYPOSIALADEN, (*hyppo*, *σιαλον*, 'saliva,' and *αδην*, 'a gland,') Submaxillary gland.

HYPOSIALADENITIS, (*hyposialaden*, and *itis*,) *Hypognathadenitis*.

HYPOSPA'DIA, *Hypospadi'asis*, (*hyppo*, and *σπασω*, 'I draw,') A malformation, in which the canal of the urethra, instead of opening at the apex of the glans, terminates at its base, or beneath the penis, at a greater or less distance from

the symphysis pubis. When the orifice of the urethra is very near the root of the penis, the scrotum is divided, as it were, into two great labia; and this malformation has often been taken for a case of hermaphroditism. *Hypospadia* is ordinarily incurable; and an idea has been entertained that it is the cause of impotence. It is not exactly so; but it renders impregnation less probable.

HYOSPA'DIAS, *Hypospadi'asis*, *Hypospadi'icus*, *Hypospadi'an*, (F.) *Hypospade*. One affected with hypospadia.

HYOSPADIASIS, *Hypospadia*.

HYOSPATHIS'MUS, (ὑποσπαθισμός, from *hyppo*, and *σπαθιον*, 'a ladle or spatula,') A surgical operation, with the ancients, in cases of severe 'defluxion to the eyes,' which consisted in incisions made in the scalp down to the bone, and the insertion of an instrument called a *spath'ion* or *hypospathis'ter*, to enable the intermediate parts to be divided. The precise character of the operation is not well described, and it is no longer known.

HYOSPATHISTER, see *Hypospathismus*.

HYOSPHAG'MA, gen. *Hyposphag'matis*, (*hyppo*, and *σφαζω*, 'I kill,') The coagulated blood, which is collected when an animal is killed and used for food. Also, an effusion of blood, especially under the conjunctiva.

HYOSPHAGMA, *Hæmophthalmia*.

HYPOSTAPHYLE, (*hyppo*, and *σταφυλη*, 'uvula,') *Staphylodema*.

HYPOSTAPHYLITIS, (*hypostaphyle*, and *itis*,) see *Staphylitis*.

HYPOSTASIS, (ὑποστασις, from *hyppo*, and *στασις*, 'the act of placing,') A morbid deposition in the body. Sediment.

HYPOSTAT'IC, *Hypostat'icus*, (F.) *Hypostat'ique*, (*hyppo*, and *στασις*, 'stagnation,') Relating to hypostases, sediments or depositions.

HYPOSTATIC HYPEREMIA, *Hyperémie* ou *Congestion hypostatique*. A congestion of blood in the vessels of a part, caused by its depending position.

HYPOSTEMA, gen. *Hypostematis*, (ὑποστέμα,) *Sediment*.

HYPOSTHENIA, (*hyppo*, and *σθενος*, 'strength,') Debility.

HYPOSTHENIC, *Centrostimulant*.

HYOSTHÉNIQUE, *Centrostimulant*.

HYOSTROPHE, (ὑποστροφή, from *hyppo*, and *στροφή*, 'I turn,') Act of a patient in turning himself. Also, a relapse or return of a disease.

HYOSTYPT'IC, *Hypostyp'ticus*, (*hyppo*, and *στυψίς*, 'constriction,') A feeble astrigent.

HYOSYPH'ILIS, (*hyppo*, and *syphilis*,) A mild form of syphilis.

HYPOTHEMA, (ὑποδεμα, from *hyppo*, and *τίθημι*, 'to place,') Suppository.

HYPOTHENAR, *Sub'vola*, (ὑποθηναρ, from *hyppo*, and *θεναρ*, 'the palm of the hand or sole of the foot,') *Hypothenar Em'inence*. The fleshy projection of the palmar surface of the hand, which corresponds with the little finger, and is supported by the fifth metacarpal bone. This eminence is formed of four muscles: the *Palmaris brevis*, *Adductor* seu *Flexor brevis minimi digiti*, and *Opponens minimi digiti*. The name *Hypothenar* has also been given to different muscles of the hand. The *Hypothenar minimi digiti* of Riolan comprehended the *Adductor*, *Flexor brevis* and *Opponens minimi digiti*; and his muscle, *Hypothenar pollicis*, corresponded to the *Adductor* and a portion of the *Flexor brevis pollicis*. Winslow called the *Adductor minimi digiti*, *muscle petit hypothenar* ou *hypothenar du petit doigt*.

HYPOTHENAR MINIMI DIGITI, *Flexor parvus minimi digiti*—h. Minor metacarpus, *Adductor minimi digiti*—h. Rioli, *Flexor parvus minimi digiti*.

HYPOTHETON, (*ὑπο*, and *τιθεμι*, 'to place,') Suppository.

HYPOTHYMIAMA, gen. *Hypothymia'matis*, Fumigation.

HYPOTHYMIASIS, (*ὑποθυμίασις*, from *hypo*, and *θυμιαω*, 'to fumigate,') Fumigation.

HYPOTIUM (*Emplastum*;) (*hypo*, and *ous*, gen. *ωτος*, 'the ear,') *Emplastrum auricula'rē*; a plaster applied behind or under the ear.

HYPOTROPE, (*ὑποτροπή*, from *hypo*, and *τροπω*, 'to turn,') Relapse.

HYPOTROPH'IA, *Hy'trophia*, (*hypo*, and *τροφη*, 'nourishment,') Scanty nourishment or nutrition. Atrophy.

HYPOTROPIASMUS, (*ὑποτροπισμός*, from *hypo*, and *τροπω*, 'to turn,') Relapse.

HYPOTRYGUS, (*ὑποτρύγος*, from *hypo*, and *τρύξ*, gen. *τρύγος*, 'lees,') Feculent.

HYPOUCANNA, *Ipecacuanha*.

HYPOXÉMIE, (*hypo*, *oxygen*, and *αίμα*, 'blood,') Asphyxia.

HYPOX'YS EREC'TA, *Star'grass*; (*hypo*, and *ξύς*, 'sharp,' because the pod is acute at the base [?]); indigenous; *Order*, *Amaryllidaceæ*. The root is eaten, and has been used as a vulnerary, and in chronic ulcers; and in ague.

HYPOZO'MA, gen. *Hypo'zo'matis*, (*ὑποζωμα*, from *hypo*, and *ζωωμι*, 'I bind round,') *Membrana succin'gens*. A membrane or septum, as the mediastinum, diaphragm, &c.

HYPOCRAS, (see *Hippocras*.) Claret.

HYPSELOGLOSSUS, *Basioglossus*.

HYPSILODES, OS, (*υ*, the Greek letter *u psi*-ion,) *Hyoides os*.

HYPSILOGLOSSUS, *Hyoglossus*.

HYPSONOSUS, (*ὑψος*, 'height,' and *νσος*, 'disease,') Puna.

HYP SOPH'ONUS, (*ὑψος*, 'height,' and *φωνη*, 'voice,') One who has a clear high voice.

HYP SISIS, (*ὑψωσις*, 'lifting on high,') Sublimation.

HYPTIASMA, (*ὑπτιάζω*, 'I lie with the face upwards,') Supination.

HYPTIASMUS, (*ὑπτιασμός*.) Lying in a supine posture. Also, inversion of the stomach, as in nausea, regurgitation, or vomiting.

HYP U'LUS, (*ὑποῦλος*, from *hypo*, and *ουλη*, 'cicatriz,') Imperfectly cicatrized.

Ulcera hypu'la. Ulcers healed at the top, but not at the bottom.

HYRA'CEUM. A substance found in the Cape Colony, which Thunberg and other travellers mistook for a kind of bitumen, *Stone sweat*; but, according to Dr. Pappe, it is obtained from the urine of the *Klipdas* or *Hy'rax Capen'sis*, which, when passed, is thick and of a glutinous nature. The animal is in the habit of evacuating the urine at one spot, where its aqueous parts evaporate in the sun—the more tenacious adhering to the rock and hardening.

In smell, and medical properties, it most resembles castor, which, according to Dr. Pappe, it may replace. It is used by the Cape farmers in nervous and spasmodic affections.

HYRAX CAPENSIS, gen. *Hy'racis*, see *Hy'raceum*.

HYRTOCHEILIDES, (*χειλος*, 'a lip,') *Labia pudendi*. See *Lips*.

HYSOPE, *Hyssopus*.

HYSSOP, *Hyssopus*—h. Hedge, *Gratiola officinalis*.

HYSSOPT'ES, (*ὑσσωπεις*). Ancient name of a wine, of which hyssop was the chief ingredient, and which Dioscorides extolled in chronic inflammation of the chest. It was regarded as diuretic and emmenagogue.

HYSSO'PUS (*ὑσσωπος*), from the Hebr. *Azob*, *Qasbi'ala*, *Hyssopus officina'lis*, Common *hyssop*,

Ysaope, (F.) *Hysaope*. It has been chiefly used as an aromatic, stimulant and pectoral, in the form of infusion.

HYSTERA, (*ὑστέρα*.) Secundines, Uterus.

HYSTERAL'GIA, *Hysterodynia*, (F.) *Fortraiture*, (*hystero*, and *αλγος*, 'pain,') Pain in the uterus. See *Metralgia*. *Irritable uterus*, (F.) *Névrologie de l'utérus*. Hippocrates uses the epithet *Hyster'al'ges*, *ὑστέραλγος*, for any thing that excites uterine pain; and, especially, [?] for vinegar.

HYSTERALGIA CATARRHALIS, *Metrorrhœuma*—h. *Galactica*, *Phlegmatia alba dolens*—h. *Lochialis*, *Dyslochia*—h. *Rheumatica*, *Metrorrhœuma*.

HYSTERAPOPNIXIS, (*hystero*, *apo*, and *πνιγω*, 'to strangle,') Angone.

HYSTERATRE'SIA, *Atretome'tria*, (*hystero*, and *ατρεσις*, 'imperforate,') Imperforation of the os uteri.

HYSTERELCO'SIS, (*hystero*, *ελκος*, 'an ulcer,' and *osis*.) *Uteri exulceratio*, *Uteri ul'cus*. Ulceration of the uterus.

HYSTEREOSIS, (*hystero*, and *ελεω*, 'to turn,') *Hysteroloxia*.

HYSTEREMPHYSEMA, gen. *Hysteremphyse'matis*, *Physometra*.

HYSTERIA, (*ὑστέρα*, 'the uterus,') *H. va'ga*, *Hysterici'mus*, *Hysteri'mus*, *Hyste'rias*, *Hysteri'asis*, *Hysteropathi'a*, *Hypercin'e'cia uteri'na seu hyste'ria*, *Uteri adscen'sus*, *Suffoca'tio hyste'rica seu uteri'na seu mul'erum*, *Asth'ma u'teri*, *Præfoca'tio mat'ricis seu uteri'na*, *Sypsa'cia hyste'ria*, *Ma'lum hyste'ricum seu hyste'ricol'gypochoudri'acum*, *Vapo'res uteri'ni*, *Affec'tio seu Pas'sio hyste'rica*, *Mor'bus hyste'ricus*, *Strangula'tio uteri'na seu Vul'væ*, *Va'pors*, *Hyste'ria*, *Hyste'ric fit*, *Mo'ther*, (F.) *Hystérie*, *Mal de Mère*, *Maladie imaginaire*, *Pulsion hystérique*, *Suffocation utérine*, *ou de matrice*, *Ascension de la matrice*, *Névrose utérine*, *Etrangement*, *Épilepsie utérine*, *Vapeurs*, *Maux ou Attaques de Nerfs*. A species of neurosis, classed amongst the spasmi by Sauvages and Cullen, and in the *Névroses de la génération*, by Pinel. It received the name *hysteria*, because it was reputed to have its seat in the uterus. It generally occurs in paroxysms; the principal characters of which consist in alternate fits of laughing and crying, with a sensation as if a ball ascends from the hypogastrium towards the stomach, chest, and neck, producing a sense of strangulation. If the attack be violent, there is, sometimes, loss of consciousness (although the presence of consciousness generally distinguishes hysteria from epilepsy) and convulsions. The duration of the attacks is very variable. It appears to be dependent upon irregularity of nervous distribution in very impressible persons, and is not confined to the female; for well-marked cases of hysteria are occasionally met with in men. During the fit,—dashing cold water on the face, stimulants applied to the nose, or exhibited internally, and antispasmodics, form the therapeutical agents. Exercise, tranquillity of mind, amusing and agreeable occupations constitute the prophylactics. See *Mania*, dancing.

HYSTERIA CATALEPTICA, *Catalepsy*—h. *Vaga*, *Hysteria*.

HYSTERIAS, *Hysteria*.

HYSTERIASIS, *Hysteria*.

HYSTERICA, see *Hysterical*.

HYSTERICAL, *Hyste'ricus*, *Va'pourish*, *Va'pouring*, (F.) *Hystérique*. Relating to or affected with hysteria.

The word *hysterica* was used by Martial for a female affected with nymphomania or with strong sexual desires.

HYSTERICISMUS, *Hysteria*.

HYSTERICUS, *Hysteria*.

HYSTERICUS, (*ὑστέριος*, from *ὑστέρα*, 'uterus,') 'Hysterical.

HYSTÉRIE, Hysteria.
HYSTERISMUS, Hysteria.
HYSTÉRITE, Metritis.
HYSTERITES, Hydrometra.
HYSTERITIS, (*hystero*, and *itis*.) Metritis.
HYSTERO, (from *ὑστέρα*, *ὑστέρα*, 'the uterus,')
in composition, uterus.
HYSTEROCARCINOMA, Metrocarcinoma.
HYSTEROCELĒ, (*hystero*, and *celē*, 'hernia,')
Metrocēlē, *Ἡστέλια* *ὑτῆρι*, *Hernia of the womb*, (F.)
Hernie de l'Utérus. This is a rare disease. The
womb may protrude through the inguinal or the
crural canal, or through the lower part of the
linea alba.
HYSTEROCELE NUDA, Prolapsus uteri.
HYSTEROGNESMUS, (*hystero*, and *σπασμος*,
'itching,') Pruritus of the uterus or genitals.
HYSTEROCOLIA, Ootica uterina.
HYSTEROCRANIUM, (*ὑστέρος*, 'behind,' and
κράνιον, 'cranium,') Oeciput.
HYSTEROCYESIS, (*hystero*, and *κυειν*, 'to be
pregnant,') Pregnancy.
HYSTEROCYSTIC, *Hystero-cysticus*, (F.) *Hys-
téro-cystique*, (*hystero*, and *κύστις*, 'the bladder,')
Relating to the uterus and bladder. Some au-
thors have called *Hystero-cystic Retention of urine*,
that which is caused by the compression of the
bladder by the uterus, during pregnancy.
HYSTEROCYSTOCELĒ, (*hystero*, *κύστις*, 'the
bladder,' and *celē*, 'a tumour,') *Hernia* of the
uterus complicated with displacement of the
bladder.
HYSTERODYNIA, (*hystero*, and *δύνη*, 'pain,')
Hysteralgia.
HYSTERODEMA, gen. *Hystero-de'matis*, (*hys-
tero*, and *οίδημα*, 'a swelling,') Hydrometra.
HYSTEROID, *Hystero'idēs*, *Hystero'idēs*, *Hys-
teroid'is*, *Hystero'id'is*, (*hysteria*, and *ειδός*, 're-
semblance,') Resembling hysteria;—as a hys-
teroid disease, symptom, &c.
HYSTEROLOGŶ, *Hystero-log'ia*, (*hystero*, and
λόγος, 'a description,') A treatise on the sound
and morbid uterine.
HYSTEROLOXIA, *Hystero-lox'is*, *Obliq'uitas*
seu Si'us obli'quis uteri, *Flex'io seu Ver'sio in-
complete* *seu Inclina'tio uteri*, *Metrolox'ia*, *Metro-
camp'is*, *Metrocaxio'ma*, *Uterus obli'quis seu in-
clina'tus*; (*hystero*, and *λόξος*, 'oblique,') An
oblique position of the uterus, occurring during
pregnancy. *Anter'e'sion of the uterus*, *Hystero-
lox'ia ante'rior*, *Anter'e'sio seu Prona'tio uteri*,
Ven'ter prepa'ndens; and *Retrover'sion of the uter-
us*, *Hystero'lox'ia poste'rior*, *Retrover'sio uteri*, *Re-
flex'io uteri comp'ita*, are varieties.
HYSTEROMALACIA, *Hystermalac'o'is* *Hys-
teromalacu'ma*, *Metromalac'o'is*, *Metromalacu'ma*,
Malac'o'is uteri, *Emolli'ties uteri morbo'sa*, *Pu-
trescent'ia uteri gra'vidi*, (*hystero*, and *μαλακία*,
'softness,') Softness of the uterus during pre-
gnancy, which renders it liable to rupture in la-
bour.
HYSTEROMALACOMA, Hysteromalacia.
HYSTEROMANIA, Nymphomania.
HYSTEROMETER, (*hystero*, and *μετρον*,
'measure,') See Sound, uterine.
HYSTÉROMÈTRE, Sound, uterine.
HYSTEROMOCHLIUM, (*hystero*, and *μοχλιον*,
'a lever,') Lever.
HYSTERON'CUS, (*hystero*, and *ογκος*, 'tumour,')
Metron'cus, *Hystero-ph'ia*, *Metraph'ia*, *Tu'mor*
ὑτῆρι. A tumour of the uterus.
HYSTEROPARALYSIS, *Hystero-par'e'gia*, (*hys-
tero*, and *παράλυσις*, 'paralysis,') Paralysis of the
uterus.
HYSTEROPATHIA, (*hystero*, and *πάθος*, 'af-
fection,') Disease or suffering in the uterus.
Also, hysteria, and deutero-pathia.

HYSTEROPHYMA, (*hystero*, and *ψαμα*, 'a
swelling,') Hysteroneus.
HYSTEROPHYSE, Hystero-physis, Physome-
tra.
HYSTEROPHY'SIS, *Hystero-ph'ia*, (*hystero*,
and *πνεν*, 'wind,') Distension of the uterus with
air: *Emph'ysē'ma ὑτῆρι*.
HYSTEROPLEGIA, (*hystero*, and *πληγη*, 'a
stroke,') Hyster-paralysis.
HYSTEROPNIX, (*hystero*, and *πνιξ*, 'suffoca-
tion,') Angone.
HYSTEROPOLYPUS, Metro-polypus.
HYSTEROPSEPHIA, (*hystero*, and *ψοφος*, 'a
sound,') Physometra.
HYSTEROPTOSIS, (*hystero*, and *πτωσις*, 'fall,')
Prolap'sus uteri. Also, *Inver'sio uteri*, *Anatrophē*
uteri. In a general sense, a protrusion of any
part of the genital organs or of excrecences from
them into the genital passages; *E'dopto'sis*.
HYSTEROPTOSIS, *Prolap'sus uteri*—h. Vaginae,
Prolapsus vaginae.
HYSTERORRHAGIA SANGUINEA, (*hystero*,
and *ῥιγναι*, 'to break forth,') Metrorrhagia.
HYSTERORRHESIS, (*hystero*, and *ρῆσις*,
'rupture,') Uterus, rupture of the.
HYSTERORRHŒA, (*hystero*, and *ῥεω*, 'to
flow,') Metrorrhagia—h. Mucosa, Leucorrhœa.
HYSTEROSALPINX, gen. *Hysterosalpin'gis*,
(*hystero*, and *σαλπιγξ*, 'a trumpet,') Tube, Fallo-
pian.
HYSTEROSCIRRHUS, Metrocirrhus.
HYS'TEROSCOPE, (*hystero*, and *σκοπεω*, 'I
view,') A metallic mirror, used by Colombat de
l'Isère, in inspecting the state of the os uteri, to
throw the rays of a taper to the bottom of the
speculum uteri.
HYSTEROSPASM'US, (*hystero*, and *σπασμος*,
'spasm,') Spasm of the uterus. Also, hysterical
spasm.
HYSTEROSTOMA, gen. *Hystero-stom'atis*, (*hys-
tero*, and *στόμα*, 'mouth,') Os uteri.
HYSTEROSTOMAT'OMUS, (F.) *Hystéro-sto-
matome*, (*hystero*, and *τομή*, 'to cut,') An
instrument invented by Couleury for dividing the
os uteri, when it is important to deliver immedi-
ately, as in cases of convulsions.
HYSTEROSTOMIUM, (*hystero*, and *στόμα*,
'mouth,') Os uteri.
HYSTERO-SYNZESIS, (*hystero*, and *συνζῆσις*,
'sitting together,') Metrosynzesis.
HYSTERTOKOTOMIA, (*hystero*, *τοκος*,
'parturition,' and *τομή*, 'section,') Cæsarean
section.
HYSTERTOME, (*hystero*, and *τομή*, 'section,')
Hysterotomus.
HYSTERTOM'IA, *Hysterot'omy*, (F.) *Hysté-
rotomie*. Cæsarean section. Also, dissection of
the uterus.
HYSTERTOMOTOCIA, (*hystero*, *τομή*, 'sec-
tion,' and *τοκος*, 'parturition,') Cæsarean section.
HYSTERTOMUS, *Hystero*, *Metrot'omus*,
Me'trotome, (*hystero*, and *τομή*, 'section,') An in-
strument for dividing the womb through the
vagina. It is a kind of *Bistouri ca. té*, and is
intended to divide the cervix and os uteri.—*Hys-
terostomat'omus*.
HYSTERTRISM'US, (*hystero*, and *τρίβω*, 'I
gnash,') Spasmodic contraction of the uterus,
especially of its mouth.
HYSTRIASIS, Hystriciasis.
HYSTRIC'ASIS, *Hystri'asis*, *Hys'tricæ*, *Hys-
trici'mus*, *Cerostro'sis*, *Tricho'sis seto'sa*, (*ὑστρίξ*,
gen. *ὑστρίξ*, 'a hedgehog or porcupine,') A dis-
ease of the hairs, in which they stand erect like
the quills of the p. r. p.
HYSTRICISMUS, Hystriciasis.
HYSTRIN, gen. *Hys'tricis*, (*ὑστρίξ*,) Hystric-
iasis.
HYVES, Hives.

I.

IAMA. (*ιαμα*, gen. *ιαματος*.) Medicament.
IAMATOLOGY, (*ιαμα*, and *λογος*, 'a discourse.')
Materia medica.

IAMATOSYNTAXIOLOGIA. (*ιαμα*, *συNTAXIS*,
'arrangement,' and *λογος*, 'a discourse.') See
Prescription.

IAMATOSYNTAXIS, see Composition.
IAMATOTAXIOLOGIA, see Prescription.

IASIMUS, (*ιασιμος*.) Curable.

IASIS, (*ιασις*.) Curation.

IASPIS, (*ιασις*.) Jaspis.

IATERIA (*ιατρηια*), Medicina.

IATERIUS (*ιατρηιος*), Medicinal.

IATRALEIP'TICE, (*ιατραλειπτικη*, (*τεχνη*),
Iatralipticē (Ars), *Iatraliptic* or *Iatralip'tic*
meth'od; (*ιατρο*, and *αλειψω*, 'I anoint.') The
method of treating diseases adopted by the Iatra-
leptes,—that is, by friction chiefly, *Perfricatio*.

IATRALEPTES (*ιατραλειπτης*), *Iatraliptes*,
Iatralip'ta, *Iatroleiptes*, *Med'icus Unquentarius*.
One who treats diseases by unguents, frictions,
and by external means generally. Prodicus, a
disciple of Æsculapius, was the chief of the Iatra-
leptes.

IATRALIPTES, Iatroleptes.

IATRALIPTICE, Iatralipticē.

IATREUSIOLOG'IA, (*ιατρευσις*, 'the exercise
of the art of healing,' and *λογος*, 'a description.')
The doctrine of the exercise of the healing art.

IATREUSIS (*ιατρευσις*). The exercise of the
healing art.

IATRIA (*ιατρηια*), Medicina.

IATRICE (*ιατρικη*), Medicina.

IATRICUS (*ιατρικος*), Medical; see *Digitus*.

IATRI'NĒ (*ιατρινη*), Medicina. Also, a female
practitioner of medicine, *Med'ica*, *Mu'lier med'-*
ica. A doctress. A midwife.

IATRI'ON, (*ιατριον*), *Iū'tron*, *Ietri'on*, *Medica-*
tri'na. The house or office of a physician or sur-
geon. Also, the physician's fee or *honorarium*.

IATRIQUE, Medical.

IATROCHEMIA, (*ιατρο*, and *chemia*.) Chymi-
atria.

IATROCHYMIA, Chymiatría.

IATROGNOM'ICA, (*ιατρογνωμ'ικη*, (*ιατρο*, and
γινωσκω, 'I know.') A knowledge of medical ob-
jects.

IATROLEIPTES, Iatroleptes.

IATROLOG'IA, *Iatrol'ogy*, (*ιατρο*, and *λογος*,
'a description.') A treatise on physis and physi-
cians.

IATROMAN'TIS, (*ιατρο*, and *μαντεια*, 'divina-
tion.') A physician who is a soothsayer.

IATROMATHEMATICAL PHYSIC'IANs,
Iatromathematici, *Iatromechanici*. Mechanical physi-
cians. A sect which flourished about the middle
of the seventeenth century, and of which Borelli
was the founder.

IATRON, (*ιατρον*.) Iatron.

IATROPHYSICS, Physics, medical.

IATROS, (*ιατρος*.) Iatrus.

IATROSPHIS'TA, (*ιατρος*, and *σοφιστης*, 'one
skilled in an art or science.') A learned, or thor-
oughly educated physician.

IATROTECH'NA, *Iatrotech'nes*, (*ιατροτεχνες*,
from *ιατρο*, and *τεχνη*, 'art.') A practical physi-
cian or surgeon.

IATROTECHNICE, Medicina, Therapeutics.

IATRUS, (*ιατρος*.) Physician; also, a surgeon.

IBERIS, (*ιβηρις*), *Lepidium iberis*.

IBE'RI'S AMA'RA, *Bitter can'dyust*, (F.) *Passe-*
rage, from *Iberia*, where it is indigenous. A small
herbaceous plant, *Ord.* Crucifere, indigenous in
Europe, which was employed, of old, in gout,

rheumatism, &c. The seeds have been used, in the
dose of one to three grains, in asthma, bronchitis,
dropsy, and hypertrophy of the heart. It is said
to possess acro-narcotic properties; but it is not
much used.

IBERIS BURSA PASTORIS, *Thlaspi bursa*—i.
Campestris, *Thlaspi campestre*—i. *Sophia*, *Carda-*
mine pratensis.

IBICUIBA, *Beeniba*, *Beeniba nux*. A species
of nut from Brazil, the emulsive kernel of which
is ranked amongst balsamic remedies.

I'BIS, (*ιβις*.) A bird held sacred by the Egyp-
tians. When sick, it is asserted that it was wont
to inject the water of the Nile into its fundament:
whence, according to Langius, was learned the
use of gylsters.

IBISCHA MISMALVA, *Althæa*.

IBISCUS, (*ιβισκος*), *Althæa*.

IBIXUMA, *Saponaria*.

ICE, (Sax. *ir*), *Gla'cies*, *Fro'zen water*, (F.)
Glacé. Iced water is much used internally, as the
best refrigerant in fever. It is, also, applied ex-
ternally, in cases of external inflammation, as well
as in phrenitic and hernial affections, &c., in the
form of the *ice cap* and *ice cataplasms* or *ice poult-*
ices, formed by half filling a bladder with pounded
ice.

ICE CAP, see Ice.

ICE COLD, *Icy cold*, (F.) *Froid glacial*. A
very strong morbid sensation of cold, compared
by the patient to that which would be produced
by the application of pieces of ice.

ICE POULTICE, see Ice.

ICELAND, MIN'ERAL WA'TERS OF. Hot
springs are found in every part of Iceland. The
most noted of these is one called Geyser, two days'
journey from Hecla, and near Skalholt. The
diameter of the basin is 59 feet; and the height
to which the water is thrown is often more than
100. The heat of the water is 212°.

ICEPLANT, *Mesembryanthemum* *crystalli-*
num, *Monotropa uniflora*.

ICH'NOS, (*ιχως*.) The foot. The sole of the foot.
The heel.

I'CHOR, (*ιχωρ*), *I'chos*, *Sa'nies*, *Sordes*, *Virus*,
Pus malignum, *Tubum*. The serum of the blood.
A thin, aqueous, and acrid discharge. A cor-
rupted humor.

ICHOROÏDES, (*ιχωρ*, and *ειδος*, 'resemblance.')
Ichorous.

ICH'OROUS, *Ichorous*, *Ichoroï'des*, *Sanio'us*,
Sa'nious, (F.) *Ichoreux*, *Sanieux*. Belonging to
or resembling ichor.

ICHORRHÆ'MIA, (*ichor*, and *αιμα*, 'blood.')
A diseased condition induced, according to Vir-
chow, by the presence of corrupted or vitiated
humors in the blood.

ICHOS, (*ιχως*.) *Ichor*.

ICHTHYA, (*ιχθυα*, 'a fishhook.') Hook, blunt.

ICHTHYO, (*ιχθυς*, gen. *ιχθυς*, 'a fish.') In
composition, fish.

ICHTHYOCOL'LA, (*ichthyo*, and *κolla*, 'glue.')
Col'la Pis'cium, *Pis'glass*, *Fishglue*, (F.) *Ichthy-*
colle, *Colle de Poisson*. A name given to the dried
swimming bladder or *sacculus* of *Acipenser hu'so*,
and other species of fish, which is almost wholly
composed of gelatin, and is employed in medicine
in the formation of nutritive jellies. It is, also,
occasionally used in anatomical injections. The
English Court Plaster is made with it. In Russia,
the *Acipenser* is not the only genus from which
isinglass is procured; it is also obtained from *Si-*
lu'ria Glanis, which Dr. Royle suggests to be the
source of the *Samory* or *Somory Isinglass* of com-
merce. The sounds of the cod, ling, hake, &c.

yield it. From these it is made in New England; and, in England, is termed *New York Isinglass*.

ICHTHYOPHAGISTS, *Ichthyoph'agi*, (ἰχθυοφαγοί, from *ichthyo*, and φαγω, 'I eat.') (F.) *Ichthyophages*. People who feed habitually on fish;—generally the most uncivilized of mankind.

ICHTHYOPHAGOUS, (ἰχθυοφάγος,) Piscivorous.

ICHTHYOSE, *Ichthyosis*.

ICHTHYOSIS, (ἰχθύς, 'a fish,') from the resemblance of the scales to those of a fish. *Lepido'sis Ichthy'asia*, *L. Ichthy'o'sis*, *Lepra Ichthy'o'sis*, *Alveas ni'gra* (Arab.), *Impeti'go excorticat'iva*, (F.) *Ichthyose*; *Fishskin*, *Porcupine Disease*. A permanently harsh, dry, scaly, and, in some cases, almost horny texture of the integuments of the body, unconnected with internal disorder. Willan and Bateman have two varieties, *I. simplex* and *I. cornen*. Alibert has three, the *I. nacreæ* or *pearly*, the *I. cornée*, and the *I. pellagre* or *Pellagra*.

ICHTHYOSIS PELLAGRA, *Pellagra*.

ICHTHYOSIS SEBA'CEA, *Sebaceous Ichthy'o'sis*.

A morbid incrustation of a concrete subaceous substance upon the surface of the epidermis, confounded, according to Mr. E. Wilson, with *ichthyosis*, to which it bears a close resemblance. In many cases there is neither redness nor heat, nor is the affection often accompanied by constitutional symptoms.

ICHTHYOTOX'ICUM, (*ichthyo*, and τοξικον, 'a poison,') *Ich'thyus*, fish poison.

ICHTHYIUS, *Ichthyotoxicum*.

ICICA ARACOUCHINI, (name in Guyana.) *Aracouchini* is a balsam, extracted by incision from this tree in Guyana. The Galibis use it for healing wounds.

ICICA ICICARIBA, see *Amyris elemifera*.

ICTAR, (ἰκταρ,) *Vulva*.

ICTÈRE, *Icterus*—*i. Bleu*. Cyanopathy—*i. des Nouveau-nés*, *Icterus infantum*—*i. Noire*, *Melana*.

ICTERIC, *Ictericus*.

ICTÉRICIE, *Icterus*.

ICTERICUS, (ἰκτερός,) *Antiicteric*. Also, relating to, or resembling *Icterus*; *Icteric*, *Icteroïdes*, *Icteroïdes*, (F.) *Ictérique*.

ICTERITIA ALBA, *Chlorosis*—*i. Flava*, *Icterus*—*i. Rubra*, *Erysipelas*—*i. Rubra*, *Erysipelas*.

ICTERODES, (ἰκτερώδης, from *icterus* and *odes*), *Ictericus*.

ICTERUS, *I. fla'vus* seu *ve'rus*, (ἰκτερος, from ἰκτίς, a species of weasel, whose eyes are yellow [?]. *Icterus* meant, with the ancients, a yellow bird, perhaps the Oriole.) *Morbis arcua'tus* seu *arquatus*, *Leseli morbus* (Paracelsus), *Auri'go*, *Auri'go*, *Mor'bus re'gius* seu *lute'olus*, *Cholelith'ia ic'terus*, *Cholihæ'mia*, *Cholæ'mia*, *Cholopla'nia*, *Cholo'sis*, *Dermatocho'lo'sis*, *Suff'a'sio aurigno'a* seu *fel'tis* seu *bi'lis*, *Icteri'tia* seu *icteric'ia fla'va*, *I'cus fla'vus* seu *ictericoides*, *Canche'ia icter'ica*, *Fel'tis suff'a'sio seu obstruct'io seu superfu'sio*, *Bil'ious Dys'crasy*, *Jau'n'dice*, *Yel'l'ice*, *Jan'ders*, *Jau'n'ders*, (Sc.) *Gul'schoch*, (Prov.) *Jo'nas*, *Yellot*, (F.) *Ictère*, *Ictéricie*, *Jau'nisse*, *Bile répandue*. A disease, the principal symptom of which is yellowness of the skin and eyes, with white faeces and high-coloured urine. It admits of various causes; in fact, anything which can directly or indirectly obstruct the course of the bile, so that it is taken into the mass of blood, produces the yellowness of surface; the bile being separated by the kidneys, causes yellowness of urine, and its being prevented from reaching the intestine occasions the pale-coloured faeces. The prognosis, in ordinary cases, is favourable;—when complicated with hepatic disease, unfavorable. The treatment is simple:—an emetic or purgative, given occasionally so as to

solicit the return of the bile to its ordinary channels; light tonics; unirritating diet; cheerful company, &c.

ICTERUS ALBUS, *Chlorosis*—*i. Cæruleus*, *Cyanopathy*.

ICTERUS INFAN'TUM, *I. Neonato'rum*, *Neotoc'iet'erus*, *Pædi'et'erus*, *Auri'go neophyto'mum*, *Yel'l'ice gum*, *Yell'ice gomen*, (F.) *Ictère des nouveau-nés*, is a common affection, and frequently dependent upon obstruction of the choledoch duct by the meconium. It requires time; and castor oil, occasionally.

ICTERUS MELAS, *Melæna*—*i. Neonatorum*, *I. infantum*—*i. Niger*, *Melæna*.

ICTERUS SATURNI'NUS, *Lead jaundice*. The earthy-yellow hue in saturnine cachexy.

ICTERUS VIRIDIS, *Green jaundice*. *Icterus* when the skin is of a green or olive hue.

ICTODES FETIDUS, (from ἰκτίς, 'a weasel,') *Dracontium fetidum*.

ICTUS, gen. *Ictūs*: 'a stroke or blow;' (from ἰκο, *ictum*, 'to strike,') *Ple'gē*, *Pla'ga*. *Ictus so'lis*, a stroke of the sun.—See *Coup de soleil*. *Ictus*, also, means the pulsation of an artery, and the sting of a bee or other insect.

ICTUS, *Blow*—*i. Cordis*, see *Heart*—*i. Cordis et arteriarum*, *Pulse*—*i. Sanguinis*, *Apoplexy*—*i. Solis*, *Coup de soleil*.

IDEA, *Idea*, (*idea*), *Ido'lum*, *Ideach* [? Paracelsus], (F.) *Idée*, (from ιδω, 'I see,') The image or representation of an object in the mind.

IDEA, *FIX'ED* or *PREDOMINANT*, *Idea fix'a*. *Ideopeg'ma*. Tension of the mind on one notion; often observed in insanity.

IDEA MORBI. Knowledge or idea of a disease.

IDEACH, *Idea*.

IDEAGENOUS, (*idea*, and γένεω, 'to generate,') *Sentient*.

IDEAL, *Idea'lis*. Mental, notional, fancied.

Mor'bi idea'les. Ideal diseases. Diseases of the imagination.

IDEA'TION, *Idea'tio*. The establishment of a distinct mental representation or idea of an object.

IDÉE, *Idea*.

IDEN'TITY (PERSONAL), *Iden'titas*, (ἰδ.,) *Identité*, (*idem*, 'the same,') *Same'ness*. It is sometimes a question in legal medicine to decide upon personal identity: that is, whether an individual be the same he represents himself to be.

Physical marks form the great criteria.

IDEOL'OGY, *Ideolog'ia*, (*idea*, and λόγος, 'a discourse,') The science of ideas. Intellectual philosophy.

IDEOMO'TOR. A name given by Dr. Carpenter to muscular movements, the result of complete engrossment by an idea, which he regards as automatic, although originating in the cerebrum.

IDEOPEGMA, gen. *Ideopeg'matis*, (*idea*, and πηγνυ, 'to fasten,') *Idea*, fixed.

IDÉOSYNCHYSIE, (*idea*, and συγχύω, 'confusion,') *Delirium*.

IDIAPOCAUSIS, (*idio*, apo, and καίω, 'I burn,') *Combustion*, human.

IDIEMPRESIS, (*idio*, and εμπρησος, 'burning,') *Combustion*, human.

IDIO, (from ἰδιος,) in composition, *proper*, peculiar.

IDIOCRASIS, (*idio*, and κρasis, 'mixture,') *Idiosyncrasy*.

IDIOCTONIA, (*idio*, and κτονω, 'murder,') *Suicide*.

IDIOCY, *Idiotism*.

IDIOMIASMATA, see *Miasm*.

IDIOPATHIA, *Idiopath'ia*, *Protopath'ia*, *Pro'prius affect'us*, *Mor'bus idiopath'icus* seu *pro'prius seu prima'rius seu protopath'icus seu origina'lis*, *Ma'lum prima'rium*, (*idio*, and παθος

affection.') A primary disease; one not depending on any other.

IDIOPATH'IC, *Pro'prio hab'itu seu constitutio'nē pro'pria pen'dens*, *Idiopath'icus*, *Idiopath'es*. (F.) *Idiopathique*. Primary affections and their symptoms are so denominated.

IDIOPATCY, (*idio*, and *opsis*, 'vision,') *Achromatopsia*.

IDIOPTS, see *Achromatopsia*.

IDIOSYNCRASIA, *Idiosyncrasy*—i. *Hæmorrhagia*, *Hæmatophilia*—i. *Olfactoria*, *Parosmia*.

IDIOSYNCRASY, *Idiocrasy*, *Idiosyncrasis*, *Idiosyncrastia*, *Idiotropia*, *Idiocrasis*, *Idiosyncrastia*, (F.) *Idiosyncrasie*, (*idio*, *syn*, 'with,' and *crasis*, 'mixture,') A peculiarity of constitution, in which one person is affected by an agent, which, in numerous others, would produce no effect. Thus, shell-fish, bitter almonds, produce urticaria in some, by virtue of their idiosyncrasies. Others faint at the sight of blood, &c.

ID' IOT, *Idiotia*, (*ἰδιωτης*, 'an ignorant person,') Now used for one who is fatuous, or who does not possess sufficient intellectual faculties for the social condition, and for preserving himself from danger,—*Ho'mo fat'uus*, (Prov.) *Dal'cop*, *Fond'ling*, *Hob'bil*. In law, one who has been without understanding from his birth, and whom the law presumes to be never likely to attain any.

IDIOTIA, *Idiotism*—i. *Endermica*, *Cretinism*.

IDIOTIE, *Idiotism*.

ID' IOTISM, *Idiotismus*, *Idiotia*, *Imbecillitas mentis*; *Mo'ria de'mens*, *Ana'a*, *Mo'ria*, *Moro'sis*, *Mei'o'sis*, *Fatu'itas*, *Amen'tia*, *Demen'tia natura'lis*, *Stupid'itas*, *Vecor'dia*, *Imbecill'itas Ingen'ii*, *Id'iotcy*, *Id'iotcy*, *Fatu'ity*. (F.) *Démence innée*, *Idiotisme*, *Idiotie*, *Apyschisme* of *Piorry*. A species of unsound mind, characterized by more or less complete obliteration of the intellectual and moral faculties. It may supervene on mania and melancholia, when it is termed *Demen'tia*, but more commonly it depends upon original conformation. It may also, be symptomatic of organic disease of the brain, which has come on after birth. *Idiotism* exists in various degrees. Some idiots are mere automata, exhibiting scarcely any sensibility to external impressions; others are capable of articulating a few words, and possess certain mental emotions to a limited extent. The physiognomy is usually vacant, step unsteady, and articulation imperfect or broken. The affection is almost always incurable; but it may often be palliated.

ID' IOTISME, *Idiotism*.

ID' IOTISMUS, *Idiotism*—i. *Endemicus*, *Cretinism*.

ID' IOTROPIA, (*idio*, and *τροπω*, 'to turn,') *Idiosyncrasy*.

IDOLUM, (*εἰδωλον*,) *Hallucination*, *Idea*.

IDROSIS, (*ἰδρωσις*, 'sweat,') *Ephidrosis*.

IETRION, (*ιητριον*, *ιητριον*,) *Iatrimon*.

IF, *Taxus baccata*.

IFFIDES, *Plumbi subcarbonas*.

IGDE, (*ιγδρ*,) *Mortar*.

IGDIS, (*ιγδς*,) *Mortar*.

IGNA'TIA AMA'RA, ('bitterignatia,') *Strych'nos Ignat'ii*, *Ignatia'na Philippin'ica*. *Ord.* *Apocynaceæ*. The systematic name of the plant which affords *St. Ignatius's Bean*, *Fa'ba In'dica seu Sancti Ignat'ii seu febrif'uga*, (F.) *Ignatie*, *Fève de Saint Ignace*, *Noix d'Igasur*. The seeds, *Ignatia'na* (Ph. U. S.), are bitter and poisonous, containing strychnia, and only appear to differ from *nux vomica* in containing a larger proportion of that alkaloid.

IGNATIE, *Ignatia amara*.

IGNAVIA, ('laziness,' from *in*, priv., and *gravis*, 'industrious,') *Inertia*—i. seu *Ignavitas partium genitalium*, *Impotence*.

IG'NIS, 'fire.' Hence:

IGNIS ACTUALIS, *Cauterium*—i. *Animalis*, *Animal heat*.

IGNIS CAL'IDUS. 'A hot fire.' A violent inflammation, about to degenerate into gangrene.

IGNIS COLUMELLE, *Erysipelas*.

IGNIS FRIG'IDUS. 'A cold fire.' *Sphacelus*.

IGNIS NATURALIS, *Animal heat*—i. *Persicus*, *Anthrax*, see *Herpes zoster*—i. *Philosophicus*, *Phosphorus*—i. *Potentialis*, see *Cauterium*—i. *Sacer*, *Erysipelas*, *Herpes zoster*—i. *Sancti Antonii*, *Erysipelas*.

IGNIS HÆSIEN'TUM. The ancient name for the heat of horses' dung.

IGNIS SYLVAT'ICUS, *I. sylvest'ris* seu *volat'icus* seu *vola'grius*, *Stroph'ulus sylvest'ris* seu *volat'icus*, *Wild fire rash*, (F.) *Feu sauvage ou volage*, *Flamboise*. Probably, the *Porri'go larva'lis* or *Crusta lac'tea* of infants. Also, a transient redness of the face and neck, sometimes observed in hysterical and chlorotic females.

IGNIS SYLVESTRIS, *I. sylvaticus*—i. *Vitalis*, *Animal heat*—i. *Volagrarius*, *I. sylvaticus*—i. *Volaticus*, *I. sylvaticus*.

IGNIVOROUS, (*ignis*, and *voro*, 'I devour,') *Pyrophagus*.

IGNYE, (*γννα*,) *Poples*.

IGNYS, (*γνys*,) *Poples*.

IL, as a prefix, see *Im*.

ILAPHIS, *Arctium lappa*.

ILEA, *Flanks*.

ILEAC PASSION, *Neus*.

ILEA DELPH'US, (*ileum*, and *ἀδελφος*, 'brother,') A double monster, united at the pelvis, and below.

I'LECH, *Y'lech*, *Nei'as*, *Ilia'ster*, *Ylia'ter*, *Eli-as'ter*, *Ilia'strum*, *Ilia'dos*, *Nei'dos*, *Ilia'dum*, *Ilia'dus*. Terms used by Paracelsus to designate the first matter:—the beginning of every thing.

ILEIADOS, *Ilech*.

ILEIADUM, *Ilech*.

ILEIAS, *Ilech*.

ILEIDOS, *Ilech*.

ILEIT'IS, (F.) *Iléite*, (*ileo*, and *itis*,) *Inflammation of the ileum*.

ILEO, in composition, *ileum* or *ilium*.

ILEOCE'CAL, *Ileocec'alis*. Relating to the ileum and cæcum,—as the *ileocec'al valve*.

ILEOCHOLOSIS, (*ileo*, and *χολη*, 'bile,') *Diarrhoea*, *bilious*.

ILEOCOL'IC, *Ileocol'icus*, (F.) *Ileo-colique*. Relating to the ileum and colon:—as the *ileo-colic valve* or valve of Bauhin.

ILEOCOLITIS, see *Enteritis*.

ILEODICLIDITE, (*ileo*, *δικλς*, gen. *δικλιδος*, 'a valve,' and *itis*,) see *Typhus*.

ILÉOLOMBAIRE, *Ileolumbar*.

ILEOLUM'BAR, *Ileolumba'lis*, *Iliolumba'ris*, (F.) *Réolombaire*. Belonging to the ilium and lumbar region.

ILEOLUMBAR AR'TERY, *Iliolumbar artery*, *Il'aco-muscular*, (Ch.) is given off by the hypogastric, opposite the base of the sacrum. It ascends behind the psoas muscle, and divides into two branches;—an ascending and a transverse, which give off numerous ramifications to the neighbouring parts.

ILEOLUMBAR LIG'AMENT, *Iliolumbar* or *Iliolumbover'tebral ligament*, *Vertebro-iliac ligament*, is a broad, membraniform, triangular ligament, extending horizontally from the transverse process of the 5th lumbar vertebra to the upper and posterior part of the iliac crest. It unites the vertebral column with the pævis.

IL'EON, *I'l'eum*, *É'l'eon*, *I'l'ium*, *Intesti'num circumvolu'tum*, (*ειλεω*, 'to turn,' 'to twist,') Anatomists have given this name to the longest portion of the small intestine, which extends from the jejunum to the cæcum. It was so called from its forming a considerable number of convolutions.

ILEOPYRA, (*ileo*, and *pyr*, 'fire, fever;') see Typhus.

ILEOSIS, Ileus.

ILEOTYPHUS, see Typhus.

ILEN, Flanks.

ILEUM, Ileon.

ILEUS, *Ἐίλεος*, (*εἰλος*, from *εἰλω*, 'I twist,')

Colica Ileus seu *spasmodica*, *Enteroleisia*, *Enteroleisia*, *Ileus spasmodicus*, *Chordapneus*, *Passio Iliaca*, *Iliac* or *Ileac Passion*, *Volvulus*, *Ileo'sis*, *Miserere mei*, *Convolutus*, *Tormentum*, *Interceptio Intestinalis*, *Enterostenosis volvulus*, (F.) *Colique de Miséricorde*, *C. de Misérère*, *Passion iliaque*. A disease, characterized by deep-seated pain in the abdomen, stercoraceous vomiting, and obstinate constipation. It is occasioned by hernia or other obstruction to the passage of the feces through a part of the intestinal canal, *Enterostenosis*. The term Ileus has been applied to various affections—to simple nervous colic, intussusception, and to strangulation of the small intestine, &c. Various remedies have been employed;—the majority for the purpose of forcing alvine evacuations,—as purgatives, in draught, pill, or glyster; suppositories, tobacco glysters; pure mercury; leaden bullets; antispasmodics and narcotics; blisters to the epigastrium; ice by the mouth, or injected into the rectum. Some of these have, occasionally, succeeded,—especially the tobacco glyster, and ice to the tumour, where the disease has been occasioned by strangulated hernia. It is very dangerous.

ILEUS FLAVUS, Icterus—i. Icteroides, Icterus—i. Inflammatorius, Enteritis.

ILEX AQUIFOLIUM, gen. *Ilicis*. The systematic name of the Common *Holly*, *Aquifolium*, *Agriifolium*, *πρινος* of the ancients [?], (Prov.) *Hollen*, (Sc.) *Hollyn*, *Hol'ene*, (F.) *Houx*, *H. commun* ou *Chêne vert*. Ord. Aquifoliaceæ. The leaves of this plant have been recommended as tonic, astringent, and antiseptic, and have been prescribed in atonic gout, intermittents, dyspepsia, &c. *Ilicine*, the active principle, has also been advised. It is obtained by dissolving the alcoholic extract of the leaves of the holly in water, and successively treating it with the subacetate of lead, sulphuric acid, and carbonate of lime. The filtered and evaporated product is then dissolved in alcohol; and the mixture filtered and evaporated in shallow vessels.

ILEX CASSINE, *Ilex vomitoria*.

ILEX MAJOR. From the berries of this tree, called by the Spaniards *Bellotas*, a juice may be expressed, which forms a slightly astringent emulsion with water, and has been recommended by some Spanish physicians in humid cough, hæmoptysis, &c.

ILEX MATE, *Ilex paraguensis*.

ILEX OPA'CA, *American Holly*, grows throughout the Atlantic portion of the United States. It is said to possess the same properties as the European variety.

ILEX PARAGUENSIS, *Viburnum lævigatum* seu *Cassinoïdes*, *Cassine* *Pera'gua* seu *Caroliniana*, *I. Maté*; a native of Paraguay. It affords the celebrated *Mattee* or *Maté*, *Folia Apalachines* seu *Pera'gue*, which is drunk in place of the Chinese tea by the people of Paraguay. By M. Lenoble, of Montevideo, the mate is referred to *Psora lea glandulosa*.

ILEX VOMITORIA, *I. Cassine*, *Cassina*, *Dahoon*, *holly*, *Apalachine*, *Gal'lis*, *Southsea Tea*, *Evergreen Cassine*, *Cassé'na*, *Yaupon*, *Yopon*, (F.) *Thé des Apalaches*, *Houx Apalachine*, *Apalachine*, *Thé de la Mer du Sud*. A tree, indigenous in the southern parts of the United States. The leaves, when dried, are aromatic, stimulant, stomachic, and expectorant, and are used as a tea. When fresh, they are emetic and cathartic.

ILIA, Flanks.

ILIAC, *Iliacus*, *Ili'al*, (F.) *Iliaque*, (from *ilia*.) Relating to, or connected with, the flanks.

ILIAC ARTERIES, *Arterie Iliacæ*. This term has been given to several arteries. The *Primary Iliacs*, (F.) *A. Iliacæ primitives*,—*A. Pelviculares*, (Ch.) arise from the bifurcation of the aorta, opposite the body of the 4th lumbar vertebra, and descend, in a divergent manner; until, opposite the sacro-iliac symphysis, they divide into two considerable trunks,—the *internal Iliac* or hypogastric (which see), and the *external Iliac*—*Portion Iliac de la crurale*, (Ch.) This proceeds from the sacro-iliac juncture as far as the crural arch, when it assumes the name *Femoral Artery*. Before passing under the arch it gives off two pretty considerable branches,—the *Epigastric*, and the *Circumflexa Ilii*. This last, which is called, also, *anterior Iliac* by some anatomists,—*Circumflexa de l'Ilium* (Ch.)—leaves the external iliac at the crural arch. It ascends along the outer edge of the iliacus muscle, and divides into two branches—an *internal* and an *external*.

The *Iliac Interna Minor* seu *Iliolumbalis* is a small artery, which sometimes arises from the hypogastric; at others, from the beginning of the gluteal. It is sent, chiefly, to the psoas and iliacus internus muscles.

ILIAC CREST, *Crista Ilii*, is the upper margin of the ilium. (F.) *Crête Iliaque*. It is very thick, and curved like the Italic *S*. It affords attachment to the broad muscles of the abdomen.

ILIAC FOS'Æ are two in number; the *internal*, the depression presented by the ilium on its interior, and at the upper part in which the *Iliacus internus* is lodged:—and the *external*, an excavation on the outer surface of the same bone, occupied by the *Glutæi* muscles.

ILIAC MUSCLE, INTER'NAL, *Iliacus internus* Muscle, *Iliacus*, *Iliaco-trochanterien* (Ch.); *Femur moventium septimus*, *Iliac muscle*, is situated in the fossa iliaea, and at the anterior and superior part of the thigh. It is broad, radiated, triangular; and is attached, *above*, to the two upper thirds of the fossa iliaea, and to the internal part of the iliac crest:—*below*, it terminates by a tendon, which is common to it and the psoas magnus, and is fixed into the lesser trochanter. When this muscle contracts, it bends the pelvis on the thigh outwards, and prevents the trunk from falling backwards.

ILIAC PASSION, Ileus.

ILIAC RE'GIONS, *Regiones iliacæ*, *Inan'ia*, are the sides of the abdomen between the ribs and the hips.

ILIAC SPINES, *Spi'nous Proc'esses of the Ilium*, are four in number. They are distinguished into *anterior* and *posterior*. Of the *anterior*, one is *superior*. It bounds the crista ilii anteriorly, and affords origin to the sartorius muscle. The other is *inferior*, and receives the tendinous origin of the rectus femoris.

The two *Posterior Iliac Spines* are divided, like the *anterior*, into *superior* and *inferior*. They afford insertion to strong ligaments, which unite the ilium with the sacrum.

ILIACO-TROCHANTÉRIEN, *Iliacus internus*.

ILIACUS, *Iliac muscle*, internal—i. *Externus* *Pyramidalis*.

ILIADUS, Ilech.

ILIAL, Iliac.

ILIASTER, Ilech.

ILIASTRUM, Ilech.

ILINGOS, (*ιλιγγος*;) Vertigo.

ILIO. In composition, relating or appertaining to the ilion or ileon.

ILIO-ABDOMINAL, Obliquus internus abdominis—*i. Apophyseos-femoralis*, Fascia lata muscle—*i. Costal*, Quadratus lumborum—*i. Cresti-tibial*, Sartorius—*i. Femoral*, Coxofemoral—*i. Hypogastricus* (nervus), Musculocutaneous nerve, superior—*i. Inguinal*, Musculocutaneous nerve, inferior—*i. Ischii-trochantérien*, Gluteus minimus—*i. Lombi-costal*, Quadratus lumborum—*i. Lombo-costo-abdominal*, Obliquus internus abdominis—*i. Lumbalis*, Iliolumbar, see Iliac arteries—*i. Lumbovertebral* ligament, Iliolumbar-ligament.

ILIOPECTINEA, *Reopectinea*, (F.) *Ilio-pectiné*, Ilio-pectineal. Belonging to the ilium and to the pecten or pubis.

ILIOPECTINEAL EMINENCE is so called from being formed by the junction of the ramus of the ilium with that of the pubis. It affords attachment to the psoas minor.

ILIOPECTINEA LINEA, *Linea innomina'ta*, is the projecting line or ridge of the ilium and pubis, which forms part of the brim of the pelvis.

ILIOPRÉTIBIAL, Sartorius.

ILIOPSOITIS, (*ilio*, *φωγ*, 'the loins, and *itis*.) Inflammation of the ilium and psoas muscule.

ILIO-PUBO-COSTO-ABDOMINAL, Obliquus externus abdominis—*i. Rotulien*, Rectus femoris—*i. Sacral*, Sacro-iliac—*i. Sacral* articulation, Sacro-iliac articulation—*i. Sacro-femoral*, Glutæus major—*i. Scrotal* nerve, Musculo-cutaneous, superior—*i. Trochantérien*, Glutæus medius—*i. Trochantérien* petit, Glutæus minimus.

IL'ION, *Il'ium*, *Il'eum*, *Haunch* or *Hip bone*. The largest of the three bones which constitute the os innominatum in the fœtus and child. (It was probably so called from its seeming to support the intestine ilion; or, perhaps, because its crest is curved upon itself, from *εἰλω*, 'I twist,' 'I roll.') This portion of the os innominatum is usually so called in the adult. The posterior surface is called *Dorsum*, the internal *Venter*. The upper semicircular edge is the *Crista* or *Spine*, at the anterior part of which is the *anterior* and *superior spinous process*; and, below, the *anterior* and *inferior spinous process*. At the back part of the spine are two *spinous processes*, the *posterior* and *superior*, and *posterior* and *inferior*. See Iliac spines.

ILITHY'IA, (*Εἰλιθυία*), *Luci'na*, *Juno Lucina*. The goddess who presided over parturient females, with the Greeks and Romans.

ILIUM, Ileum, Ilium.

IL'KESTON, MINERAL WATERS OF. Ilkeston is in Derbyshire, England, about eight miles from Nottingham. The water contains carbonate of lime and soda, chloride of calcium, sulphate of magnesia, sulphate of soda, carbonic acid, sulphohydric acid, and a little iron.

ILL, Sick.

ILLACRYMATIO, (*il*, and *lacryma*, 'a tear,') Epiphora.

ILLECEBRA, (from *illicere*, 'to allure,') Sedum—*i. Major*, Sedum telephium.

ILLECEBRUM POLYGOÏDES, Achyranthes repens—*i. Vermiculare*, Sedum.

ILLEGITIMATE, *Illegit'imus*, (F.) *Illegitime*, (*il*, and *legitimus*, (*lec*, *legis*, 'law,') 'legitimate,') *Bastard*. That which is contrary to law: which has not the conditions required by law, as an illegitimate birth—one out of wedlock; *base-born*, *love-begot*, *love-child*.

ILLEGITIMATE or **BASTARD** is also applied to diseases—as to fevers, whose progress is anomalous. See False.

ILLI, (*ἄλλοι*,) see Strabismus.

ILLIC'IUM ANISAT'UM, (*illicere*, 'to allure,') Yellowflowered Anise, Aniseed Tree, *Star Anise*, *Ani'sum stella'tum* seu *Sinen'se*, *Selen Ba'dian*, (F.) *Anis de la Chine*, *A. étoilé*, *Ba'diane*. Ord. Magnoliaceæ. Ser. Syst. Polyandria Polygynia. The seeds are used like the ani-

seed. The same tree is supposed to furnish the aromatic bark called *Cortex An'isi Stella'ti*, *Cortex Lau'cola*.

Illicium Anisatum is said to furnish much of the so-called *Oil of An'ise*, used in the United States.

ILLIC'UM FLORIDA'NUM, *Florida An'ise Tree*, *Star Anise*, *Sweet Lau'rel*. An evergreen shrub, the bark, leaves, &c., of which have a spicy odour like anise, and might be used for it.

ILLIC'UM PARVIFLO'RUM, a shrub of the hilly regions of Georgia and Carolina, has a flavour closely resembling that of sassafras root.

ILLIDES, (*ἄλλος*, gen. *ἄλλιδος*, 'one who squints,') See Strabismus.

ILLINITIO, (*il*, and *lino*, *litum*, 'to anoint,') Inunction.

ILLISIO, (*il*, and *lædere*, 'to strike violently,') Enthlasis.

ILLITIO, (*il*, and *lino*, *litum*, 'to anoint,') Inunction.

ILLITUS, Inunction.

ILLNESS, Disease—*i. Monthly*, see Menses.

ILLOS, (*ἄλλος*,) Eye, Strabismus.

ILLO'SIS, *Ilo'is*, (from *illos*, and *osis*.) Distortion of the eyes. Strabismus.

ILLUSIO, (*il*, and *ludere*, *lucum*, 'to play,')

Illusion—*i. Sensus*, Hallucination.

ILLUSION, Hallucination.

ILLUTAMENTUM, see Illutatio.

ILLUTA'TIO, (*il*, and *lutum*, 'mud,') A word used by the ancients for the act of covering any part of the body with mud—*illutamen'tum*—with therapeutical views.

ILLOSIS, Illosis.

ILOTS DU FOIE, ('Islets of the Liver,') see Liver.

IM, **IN**, **IR**. In composition, generally mean *in*; occasionally, however, they have the privative sense of *un*.

IMAGINA'TION, *Imaginat'io*, *Figurat'io*, *Phanta'sia*, *Dianoë'ma*, (*imago*, gen. *imag'inis*, 'image,') The faculty of forming in the mind an assemblage of images and combinations of ideas which are not always in direct connection with external objects.

IMBALSAMATION, Embalming.

IMBECIL'ITY, *Imbecil'itas*, *Stupid'itas*, *Hebetudo an'imi*, *Oligopsych'ia*, (F.) *Imbecillité*. Weakness, especially of the intellect; incoherency; *Imbecil'itas mentis*.

IMBECILLIS, Infirm.

IMBECILLITAS, Imbecility—*i. Ingenii*, Idiostism, Imbecility—*i. Mentis*, Idiotism, Imbecility—*i. Ventriculi*, Gasterasthenia.

IMBELLIS AD VENEREM, ('powerless for venery,') Frigid.

IMBER'BIS, *Imber'bus*, *Agenei'os*, *Apo'gon*, (*im*, and *barba*, 'beard,') Devoid of beard, beardless.

IMBIBITIO, (*imbibere* (*in*, and *bibere*, 'to drink, to imbibe,') Absorption, Imbibition.

IMBIBIT'ION, *Imbibit'io*, *Emp'osis*, *Aspira'tion*. The action by which a body becomes penetrated by a liquid. See Endosmose. The reverse action has been called *Exbibition*. Many of the phenomena of absorption are owing to imbibition.

IMBIBITION, DOUBLE, Endosmose.

IMBOWELLING, Exenterismus.

IMBEX ('a tile') **NARIUM**, Septum narium.

IMBRICARIA SAXATILIS, Lichen saxatilis.

IM'BRICATED, *Imbrica'tus*, (F.) *Imbriqué*; (*imbrex*, 'a tile,') Placed over each other like the tiles of a roof.

IMMERSUS, (*im*, and *mergere*, *mersum*, 'to dip, because hidden, as it were?') Subscapularis muscle.

IM'MINENCE, *Imminen'tia*, (*immineo* (*im*, and

minco, ('to hang over,') Staying over or upon; impending. Some authors have designated by this term the period which precedes the invasion of a disease, when certain indications foretell its approach.

IMMINENCE MORBIDE, Diathesis, Predisposition.

IMMISSIO CATHETERIS, (*im*, and *mittere*, *missum*, 'to send,') Catheterismus.

IMMISSOR, Catheter.

IMMOBILITAS, (*im*, *movere*, 'to move,' and *habilis*, 'able,') Achesia—i. Pupillæ, Amaurosis.

IMMODERANTIA, (*im*, and *moderari*, 'to be moderate,') Intemperance.

IMMODERATIO, Intemperance.

IMMOVABLE APPARATUS, see *Apparatus*, immovable.

IMMUTANS, (*im*, and *mutare*, 'to change,') Alterative.

IMPACTED, (*impingere* (*im*, and *pangere*, *pac-tum*,) 'to drive in,' 'to strike against,') Wedged.

IMPAC'TION, *Impac'tio*. A collision. A fracture of the cranium, ribs, or sternum, with depression of some of the fragments and projection of others externally. See *Epiepsia*.

IMPATIENS BALSAMINA, *Bal'sam weed*, *Touch me not*, (called from the sudden bursting of the pods when touched.) *Ord.* Balsaminaceæ. This probably resembles the other species in its properties.

IMPATIENS FUL'VA and *I. PAL'LIDA*, *Touch me not*, *Jew'el weed*, *Bal'sam weed*, *Slip'pers*, *Cell-andine*, *Quick in the hand*, *Weath'ercocks*, *Snapp-weed*. Indigenous plants, having tender, juicy, almost transparent stems, and yellow flowers, which appear in July and August. They are found in low, moist ground, in every part of the Union. The properties are probably the same as those of

IMPATIENS NOLI ME TAN'GERE of Europe, which has an acrid taste, and acts as an emetic, cathartic, and diuretic.

IMPATIENTIA VESICÆ, (*im*, and *patior*, *passus*, 'to suffer,') Cysterethismus.

IMPEDIMENTUM, ('obstruction,') Embryagma.

IMPÉRATOIRE, *Imperatoria*.

IMPERATORIA, (from *imperare*, *imperatum*, 'to command;' from its commanding virtues.) *I. Ostruthium*, *Seli'nium ostruthium* seu *Imperatoria*, *Angel'ica officina'lis*, *Astruthium*, *Ostruthium*, *Astran'tia*, *Magistran'tia*, *Mast'hercort*; (old F.) *Austruche*, *Impératoire*. *Ord.* Umbelliferae. The roots of this plant were formerly considered *divi-num reme'dium*. They are merely aromatic and bitter, and are not used.

IMPERATORIA SYLVESTRIS, *Angelica sylvestris*.

IMPERFORATE, (*im*, *per*, 'through,' and *forare*, 'to bore,') *Atretus*.

IMPERFORATIO, Imperforation—i. Ani, *Atresia ani adnata*, *Proctatresia*—i. Pupillæ, *Synezisis*—i. Uteri, *Metratresia*.

IMPERFORATION, *Imperforatio*, *Atre'sia*, *Atr'etis'mus*. Absence of a natural aperture; as of the mouth, anus, vulva, nostrils, &c. It is congenital, being dependent upon faulty conformation.

IMPERFORATION DE L'ANUS, *Atresia ani adnata*.

IMPERIAL. A pleasant, cooling drink, formed of bitartrate of potassa $\frac{3}{4}$ ss; one lemon, cut into slices; white sugar lbs; and water Oijj. Let the mixture stand for half an hour, and strain.

IMPERIUM, ('an order,') Prescription.

IMPETIGINOSITAS, *Impetigo*.

IMPETIG'INOUS, *Impetigino'sus*, (F.) *Impé-tigineux*. Having the nature of, or relating to, *impetigo*.

IMPETIGO, gen. *Impetig'inis*, (*impeto*, 'I in-

vade,') *Impetiginos'itas*, *Dar'ta*, *Der'bia*, *Im'petus*, *Peti'go*, *Ecpye'sis* seu *Phly'sis irpeti'go*, *Le'pra Squamo'sa*, *Melita'gria*, *Ru'n'ning Scall* or *Tetter*, *Crust'ed*, *Pustular* or *Hum'id Tetter* or *Scall*, *Coc'rap*, (F.) *Dartre*, *D. crustaceæ* on *croû-teuse*, *Lèpre humide*. A word used in various acceptions. With some writers it is synonymous with itch. In others, it means a variety of herpes. Sauvages employs it as a generic term, under which he comprises syphilis, scorbutus, rachitis, elephantiasis, the itch, tinea, scrofula, &c. It forms, also, a genus in the class *Cachecticæ* of Cullen. In Bateman, it is the first genus of the 5th order, *Pustule*, and is defined—the *humid or running tetter*; consisting of small pustules denominated *Psydra'cia*. It is unaccompanied by fever; not contagious, or communicable by inoculation. He has five species:—the *I. figura'ta*, *I. spar'sa*, *I. erysipelato'des*, *I. scab'ida*, and *I. ro'dens*. See *Psoriasis*.

IMPETIGO EXCORTICATIVA, *Ichthyosis*—i. *Figurata*, *Porrigo lupinosa*—i. *Pellagra*, *Pellagra*—i. *Ulcerata*, *Zerna*.

IMPETUM FACIENS, ('impulse making,') Enormon.

IMPETUS, *Impetigo*, *Paroxysm*.

IMPLICENTAL, see *Placenta*.

IMPLICATIONES RETICULARES NERVORUM, ('reticular entwinings of nerves,') *Plexus nervorum*.

IMPLIC'ITI MORBI, *Implica'ti* seu *complica'ti Morbi*, (*im*, and *plico*, 'to fold,') Diseases which exist in an organ, and produce, concurrently, disorders in other organs.

IMPLUVIUM, (*im*, and *pluere*, 'to rain,') Embrocation; Bath, shower.

IMPOSTEM, (corr. from *apostem*, [?]) Abscess.

IMPOSTHUME, Abscess.

IMPOTENCE, *Acrati'a*, *Impotent'ia*, *I. generan'di*, *Agene'sia*, *Agenne'sis*, *Ignav'ia* seu *Ignav'itas par'tium genita'lium*, (F.) *Impuissance*; (*im*, and *potens*, 'able,') Loss of power over one or more of the members. Commonly, it means want of sexual vigor; incapacity for copulation, and chiefly on the part of the male. *Astys'ia*, *Astyp'h'ia*, *Ayno'dia*, *Adyna'mia viri'lis*. It has also been used synonymously with sterility. Impotence may be absolute or relative, constitutional or local, direct or indirect, permanent or temporary.

IMPOTENTIA, *Adynamia*, *Impotence*—i. *Generandi*, *Impotence*.

IMPOV'ERISHED, *Effe'tus*, *Depau'peratus*, (F.) *Appau'ri*. ('Having become poor,') originally from (L.) *pauper*, (F.) *pauvre*, 'poor,') The Humorists applied this epithet to a humour deprived of a part of its constituents, and particularly to the blood. This fluid was considered to be impoverished when it was pale, without the proper consistence, and abounding in serum. It was, on the contrary, rich, in their opinion, when of a scarlet colour; possessing due consistence; when it coagulated promptly, and the quantity of serum, compared with that of the clot, was by no means considerable.

IMPRÆGNATIO, *Fecundation*, *Pregnancy*.

IMPREGNATION, *Fecundation*, *Pregnancy*.

IMPRESSIBILITY, (*impressio*, and *habilis*, 'able,') See *Sensibility*—i. *Great*, *Hyperæsthesis*.

IMPRES'SION, *Impres'sio* (*im*, and *premere*, *pressum*, 'to press,') *Pros'bolê*, (F.) *Empr'inte*. A more or less deep indentation which certain organs seem to make on others. Inequalities observable on the bones, which appear to be made by the subjacent organs.

IMPRESSIONS DIG'ITAL, *Impressio'nes digita'tæ cran'ii*, *Fos'sæ digita'tes*, are the depressions of various forms, observable at the inner surface of

the cranium, which look at first view as if they were made with the fingers.

IMPRESSIONABILITÉ, see Sensibility.

IMPUBER, *Impu'bes, Impu'bis, Ane'bus, Capilla'tus*, (F.) *Impubère*, (in, and *puber*, 'at the age of puberty.') One who has not attained the age of puberty.

IMPUBES, Impuber.

IMPUDICUS, (in, and *pudicus*, 'modest.') See *Digitus*.

IMPUISSANCE, (in, and *puissance*, 'power,' from *posse*, 'to be able.') Impotence.

IMPULSE DIASTOLIC, (in, and *pellere, pulsare*, 'to drive.') *Impul'sus diastol'icus*, *Back stroke of the heart*. A jog or stroke which has been termed the 'back stroke,' felt at the end of each pulsation, and which would seem to be owing to the refilling of the ventricles.

IMPULSE OF THE HEART, see Heart.

IMPULSION DU CŒUR, see Heart.

IMPURITATES ALVINÆ, (in, and *purus*, 'pure.') Fæces.

IMPURITIES, Acatharsia.

IN, in composition, see Im.

INANIA, (*inania*, 'empty.') Iliac regions.

INANIMATE, *Inanima'tus*, (F.) *Inanimé*. Not animated or possessed of life.

INANITIATED, (from *inanire*, *inanimus*, 'to empty.') See Inanitation.

INANTIA'TION, *Inantia'tio*. The act of being exhausted for want of nourishment. One so exhausted is said to be *inani't'iated*.

INANIT'ION, *Inanit'io, Inan'itas, Ceno'sis*. Exhaustion for want of nourishment. To die from inanition is to die from exhaustion.

INAPPETENTIA, (in, and *appeto*, 'to desire.') Anorexia, Dysorexia.

INARTICULATIO, (in, and *articulus*, 'a joint.') Enarthrosis.

INAURA'TION, *Inaura'tio*, (in, and *aurum*, 'gold.') A mode of curing diseases by charms, &c., defended by Paracelsus, Van Helmont, and others.

INCANTAMENTUM, (in, and *cantare, cantatum*, 'to sing,'—for example, a magical song.) Charm.

INCANTA'TION, *Incanta'tio, Incantamen'tum, Epo'de*. A mode of curing diseases by charms, &c., defended by Paracelsus, Van Helmont, and others.

INCARCERA'TIO, (in, 'in,' and *carcer*, 'prison.') *Chatonnement, Incarceration*—i. Intestinorum, Enteropneustole—i. Intestinorum interna, Enteremphaxis.

INCARCERATION, *Incarcera'tio*. Hernia is said to be 'incarcerated,' (F.) *Incarcéré, Hernia incarcerationis*, when, owing to constriction about the neck of the hernial sac or elsewhere, it cannot be reduced with facility. Incarceration is sometimes used in the same sense as strangulation.

INCARNANS, *Incarnat'vus, Sarco'ticus, Plerot'icus, Anaplerot'icus, Sarco'tic*, (F.) *Incarnatif, Plérétique*, (in, and *caro, gen. carnis*, 'flesh.') Medicines which were fancied to promote the regeneration of the flesh. Certain bandages and sutures have also been so called.

INCARNATIF, Incarnans.

INCARNATIO, Incarnation—i. Unguim, Onychogryphosis.

INCARNA'TION, *Incarna'tio*. Growth of flesh or granulations.

INCARNATUS, Carneous.

INCENDIUM, (*incendere, incensum* (in, and *candere*, 'to burn.') *Pyrex'a, Cau'sis, Phlogo'sis*. A burning fever or any burning heat. *Incend'ium febr'ile, Incen'sio*. Inflammation, Phlegmon.

INCENDIUM SPONTANEUM, Combustion, human.

INCENSIO, Incendium.

INCENTIVUM, Stimulant.

INCERA'TION, *Incera'tio, Encero'sis*, (in, and *cera*, 'wax.') The act of incorporating wax with

some other body; or, rather, the operation whose object is to communicate to a dry substance the consistence of wax.

INCERNICULUM, (*incernere* (in, and *cernere*, 'to sift.') A strainer or sieve.

INCESSUS, (*incedo, incessum*, 'to go.') Walking.

INCIDENT'IA, (*incidere, incisum* (in, and *cedere*, 'to cut.') (F.) *Incisifs*. This name was formerly given to medicines to which was attributed the property of cutting thick or coagulated humours. The fixed alkalies, hydrosulphurets of alkalies, sulphurous water, &c., were considered to belong to this class.

INCINCTA, (in, and *cingere, cinctum*, 'to gird.') Pregnant.

INCINERA'TION, *Incinera'tio, Cinera'tion, Cinefac'tio*, (in, and *cinis, gen. cineris*, 'ashes.') In pharmacy, a process by which animal or vegetable substances are reduced to ashes.

INCISIFS, Incidentia, Incisive.

INCISIO, (*incidere, incisum*, 'to cut.') Cut, Incision—i. Simplex, Haplotomia.

INCIS'ION, *Incisi'o, En'copé, Incisu'ra, Inci'sus, To'mé*, (F.) *Taillade*. A methodical division of soft parts with a cutting instrument.

INCISION SÈCHE, ('dry incision.') See *Écraseur*.

INCISIVE, *Incisi'vus, Inciso'rius*, (F.) *Incisif*. Relating or appertaining to the incisor teeth.

Fos'sa incisi'va seu myrtifor'mis is a depression in the superior maxillary bone, above the incisor teeth.

INCISIVE BONE, Intermaxillary bone—i. Canal, see Palatine canals.

INCISIVE NERVE, *Nervus incisi'vus*. A branch of the inferior dental nerve, which supplies the incisor teeth.

INCISIVE TEETH, *Incis'or Teeth, Den'tes incisor'es seu tom'ici seu ct'nes seu dichaste'res seu gelas'ini seu primo'res seu raso'rii*. The teeth which occupy the anterior part of the upper and lower jaws are so called, because they are used for cutting the food in the manner of cutting instruments. The two middle incisors of the upper jaw are, at times, called *Butter teeth*.

INCISIVUS (MUSCULUS), Levator labii superioris proprius—i. Inferior, Levator labii inferioris—i. Lateralis et pyramidalis, Levator labii superioris alæque nasi—i. Medius, Depressor alæ nasi.

INCISOR TEETH, Incisive teeth.

INCISOR'RIUM, *Tom'ion*. An operating table. A scalpel.

INCISORIUS, Incisive.

INCISURA, *Échancreure*, Incision—i. Crucialis, Crucial Incision—i. Ischiadica Major, Sciatic notch, greater—i. Ischiadica Minor, Sciatic notch, lesser—i. Lienalis, Hilus lienalis—i. Renalis, Hilus renalis—i. Scapularis, Notch, scapular—i. Septi, see Ventricles of the brain.

INCISURA TRAG'ICA, *Notch of the Con'cha*. A wide, deep and rounded notch, which separates the tragus from the antitragus.

INCISURES DE SANTORINI, Santorini, fissures of.

INCISUS, Incision.

INCITABILITY, (*incitare*, 'to put in quick motion,' and *habilis*, 'able,') Irritability.

INCITAMENTUM, Stimulus.

INCITANS, Stimulant.

INCITATIO, Stimulation.

INCITO or INCITO MOTOR, (F.) *Incito-moteur*. An epithet, applied to an action the reverse of excitomotor; as in the case of muscular motion, which commences in the nervous centres and excites the muscles to contraction.

INCLAVATIO, (in, and *clavus*, 'a nail,') Gomphosis, see Wedged.

INCLINATIO, (*inclinare, inclinatum*, 'to bend down.') Decline.—i. *Caeli*. Climate.

INCLUSION, *MCNSTROSTY BY*, (*in*, and *cludere, clusum*, 'to shut.') *Emboîtement*.

INCLUSUS, see *Wedged*.

INCOCTUS, (*in*, and *coquere, coctum*, 'to cook,') see *Crudity*.

INCOHERENCE, (*incohérence*, (F.) *Anacolutie, Révasserie*; (*in*, *co*, and *hærrere*, 'to stick.') Want of dependence of one part on another. The condition of the mental manifestations in dementia, &c.

INCOMPATIBLE, (*in*, *competere*, 'to agree,' and *habilis*, 'able.') A substance, which cannot be prescribed with another, without interfering with its chemical composition or medicinal activity.

INCONTINENTIA, (*in*, and *contineo*, 'I contain,') *Anepisthesis*. 'Incontinence.' Inability to retain the natural evacuations. Abuse of the pleasures of love.

INCONTINENTIA ALVI, *Diarrhœa*—i. *Urinæ*, *Enuresis*, *Stercorærasia*.

INCORPORATIO, *Incorporatio*, *Corporatio*, *Ensomato'sis*, *Metensomato'sis* (*in*, and *corpus*, gen. *corporis*, 'a body,') An operation, by which medicines are mixed with soft or liquid bodies, in order to give them a certain consistence. Also, the thorough admixture of various substances.

INCRASSANTIA, *Incrassantia*, *Inviscantia*, *Spissantia*, *Epepyctica*, *Condensantia*, *Pachynctica*, *Psychotica*, *Pyrenimasctica*, (*in*, and *crassus*, 'thick,') Medicines which were formerly believed to possess the power of thickening the humours, when too thin. All mucilaginous substances were so regarded.

INCRASSATIVA, *Incrassantia*.

INCREASE, *Incrementum*, *Aux'sis*, *Auxes'sis*, (*in*, and *creascere*, 'to grow,') (F.) *Accroissement*. Augmentation of the size or weight of a body, by the application of new molecules around those which already exist.

INCREMENTUM, Augmentation, Growth, Increase.

INCRUCIATIO, (*in*, and *cruciare, cruciatur*, 'to cross,') Decussation.

INCRUSTATION, *Incrustatio*, (*in*, and *crusta*, 'a crust,') *Conit'asia*. The act of forming a crust on the surface of a body, as well as the crust itself. Also, the calcareous deposits or cartilaginous plates, which sometimes form in organs.

INCUBATION, *Incubatio*, *Incubit'io*, *Incubitus*, (*incubare, incubitum*, (*in*, and *cubare*, 'to lie upon,') This word, which is used in natural history for the sitting of birds upon their eggs, is employed, figuratively, in medicine, for the period that elapses between the introduction of a morbid principle into the animal economy and the invasion of the disease; and, likewise, for the maintenance of a temperature of warm air—say of 98° of Fahrenheit—around a diseased part.

INCUBES, see *Incubus*.

INCUBITIO, Incubation.

INCUBITUS, Incubation.

INCUBO, Incubus.

INCUBUS. *In'cubus*, *Epial'tes*, *Epial'tes*, *I. nocturnus*, *Hypnophob'ia*, *Epilep'sia nocturna*, *Asth'ma nocturnum*, *Pnigal'ion*, *Suc'cubus*, *Onei'rodyn'ia gra'vans*, *Erethis'mus onei'rodyn'ia*, *Nocturna oppressio*, *Epib'ole*, *Babuzicarius*, *Nightmare*, *Wild'mare*, *Ri'dinghag*, (Prov.) *Bitch-daughter*, *Hella*, (F.) *Cauchemar*, *Cauchevieille*, *Cochemar*, *Onei'rodynie gravative*. A sensation of a distressing weight at the epigastrium during sleep, and of impossibility of motion, speech, or respiration; the patient at length awaking in terror, after extreme anxiety. Nightmare is often the effect of difficult digestion or of an uneasy position of the body. At other times, it occurs in

consequence of severe emotions. The sensation of suffocation was formerly ascribed to the person's being possessed, and the male spirits were called incubes—the female succubæ. The disease requires no particular treatment. The causes must be avoided.

INCUBUS VIGILANT'ium, *Epial'tes vigilan'tium*, *E. hypochondri'aca*, *Daymare*. This is produced during wakefulness; the sense of pressure being severe and extending over the abdomen; respiration frequent, laborious, and constricted; eyes fixed; sighing, deep and violent; intellect, undisturbed.

INCUNABULA, (*in*, and *cunabula* [pl.], 'a cradle,') Swathing clothes.

INCUNEATIO, (*in*, and *cuneus*, 'a wedge,') Gomphosis, see *Wedged*.

INCURABLE, *Incurabil'is*, *Rem'ediless*, *Irrem'ediable*, *Insonabil'is*, *Anal'thes*, *Anaces'tos*, *Atherapeut'us*, (F.) *Inguérissable*, (*in*, and *cura* 'cure,') Not susceptible of cure;—applied to both patients and diseases.

INCURIA, (*in*, and *cura*, 'care,') Acedia.

INCURSUS ARTERIARUM, (*in*, and *currere*, *currere*, 'to run,') Pulse.

INCURVATIO, (*in*, and *curvare, curvatum*, 'to crook,') Gryposis.

INCUS, gen. *Incu'dis*, (*in*, and *cadere, cecum*, 'to hammer,') 'an anvil.' *Ac'mon*, *Os incu'di sim'ile*, *Ossic'ulum Incu'di seu mola'ri den'ti compar'atum*, (F.) *Enclume*. One of the small bones of the ear, so called from its fancied resemblance to a smith's anvil. It is situate in the cavity of the tympanum, between the malleus and orbicular. Its body is articulated with the malleus; its horizontal *ra'mus*, which is the shortest, corresponds with the entrance of the mastoid cells; and its vertical *ra'mus* is articulated with the os orbicular. The incus is composed of a compact tissue, and is developed from a single point of ossification.

INDEX, gen. *In'dicis*, *Dig'itus index* seu *secun'dus* seu *salutato'rius*, *Demonstra'tor*, *Indica'tor*, *Indicato'rius*, *Li'chanos*, *Deicticus*, (*indicare*, 'to point out,') The forefinger, index finger: See *Digits*.

INDIAN ARROW WOOD, see *Euonymus*—i. *Bread*, see *Lycoperdon tuber*—i. *Cup*, *Sarracenia purpurea*—i. *Loaf*, see *Lycoperdon tuber*—i. *Mallow*, *Abutilon Avicennæ*—i. *Paint*, *Hydrastis Canadensis*—i. *Physic*, *Gillenia trifoliata*—i. *Physic*, small-flowered, *Gillenia stipulacea*.

INDIAN SPRINGS, see *Georgia*, *Mineral waters of*.

INDIA'NA, MINERAL WATERS OF. A medicinal spring, near Jeffersonville, is much frequented. Its waters are strongly impregnated with sulphur and iron.

INDICANT, *In'dicans*, *Endeicticus*, (F.) *Indicatif*, (from *indicare*, 'to point out,') Any thing which, in the course of a disease, or in what precedes or accompanies it, concurs in pointing out the means to be used for its cure.

INDICATIF, Indicant.

INDICATING DAYS, *Di'es In'dices* seu *Indicato'rii* seu *Indican'tes* seu *Contemplan'tes*, (F.) *Jours indicateurs*. Hippocrates and others have thus called the middle day of the septenary; because they imagined, that indications were sometimes given then of the crisis which would occur in the last day of the septenary.

INDICATION, *Indicatio*, *Accusatio*, *Dela'tio*, *Endeiz'is*, *Endez'is*, *Boethemat'icum Seme'ion*. The object proposed in the employment of any means which are had recourse to for the cure of disease. It may also be defined—the manifestation afforded by the disease itself of what is proper to be done for its removal.

INDICATOR, *Extensor proprius indicis*, see *Digits*, and *Index*.

INDICUM, (from *India*.) Indigo.

INDICUS COLOR, Indigo.

INDIGENCE, (*indigeo*, (from *in*, and *eegeo*), 'to be needy,') Want.

INDIG'ENOUS, *Indig'ena*, 'Inborn,' (*indu*, for 'in,' and *geno*, for *gigno*, 'I generate,') Whatever is native in a country, in opposition to exotic; as an indigenous remedy, indigenous disease, &c.

INDIGESTION, (*in*, and *digerere*, *digestum*, 'to set in order,') Dyspepsia—i. Alkaline, see Dyspepsia—i. Neutral, see Dyspepsia.

INDIGITATIO, (*in*, and *digitus*, 'a finger,') Intussusceptio.

INDIGNABUNDUS, (*indignari*, 'to be indignant,') Rectus externus oculi.

INDIGNATORIUS, Rectus externus oculi.

INDIGO, *In'digum*, *In'dicum*, *Indicus color*, *Pigmentum Indicum*, (from *India*.) A dye-stuff, in small solid masses, of a deep azure blue colour, and devoid of smell and taste. It is obtained, by a fermentative process, from *Indigofera a'n'il*, (*I. suffruticosa*), *I. argentea*, (*I. articulata* seu *tinctoria* seu *glauca* seu *colorata*), and *I. tinctoria* seu *Indica*, and is sometimes used by the apothecary for colouring certain preparations. It has likewise been administered internally in spasmodic diseases, especially in epilepsy. The dose may be at first grains, but it may be elevated to drachms.

INDIGO, FALSE, (TALL WHITE,) see Sophora tinctoria—i. *Indigo Faux*, *Galega officinalis*—i. Weed, *Sophora tinctoria*—i. Wild, *Sophora tinctoria*—i. Yellow, *Sophora tinctoria*.

INDIGOFERA, (*indigo*, and *fero*, 'to bear,') see Indigo, *Sophora tinctoria*.

INDIGO, Indigo.

INDISPOSIT'ION, *Ma'la disposit'io*, *Dyspno'ria*, (F.) *Malaise*, (*in*, and *disponere*, *dispositum*, 'to put in order,') The state of being unwell. A slight functional disturbance, which may scarcely bear the name of disease. A feeling of sickness—*sensus ægritudinis*; vulgarly an all-overish feeling.

INDOLENT, *In'dolens*, (*in*, and *dolere*, 'to be in pain,') Exhibiting little or no pain. An epithet particularly applied to certain tumours.

INDOLENTIA, Anodynia.

IND'OLES, (*in*, and *olescere*, 'to grow,') A natural disposition or character.

INDOLES AN'IMI. The natural disposition or character of mind.

INDOLES MORBI. The nature or character of a disease.

INDOSYNCLONUS, (*Ind*, 'India,' and *συνκλινος*, 'tremor,') Beriberi.

INDUC'TIO, *Apogogē*, (*inducere*, (*in*, and *ducere*, *ductum*), 'to lead in,') A word used especially for the action of extending a plaster upon linen.

INDUCTION, ELECTRICITY OF. Faradization.

INDUCULA, (*inducere*, 'to put on,') Waistcoat, strait.

INDUMENTUM CORDIS, (*inducere*, 'to put on, to cover,') Pericardium—i. Nervorum, *Neurilemma*—i. Ventriculorum, *Ependyma ventriculorum*.

INDUR'ANS, (*in*, and *durus*, 'hard,') *Scleroticus*, *Sclerunticus*, *Scleryn'ticus*. A medicine which hardens the parts to which it is applied.

INDURATIO, Induration—i. Intestinorum, *Enteropathia cancerosa*—i. Maligna, *Scirrhus*—i. Renum, *Nephroscleria*—i. Telæ cellulossæ neonatorum, see Induration—i. Ventriculi *scirrhosa*, *Gastroscirrhus*.

INDURAT'ION, *Indura'tio*, *Sclerys'ma*, *Sclerys'mus*, *Callos'itas*, *Indurescent'ia*, *Poro'na*, *Poro'sis*, *Hard'ening*, (F.) *Endurcissement*. The hardness which supervenes, occasionally, in an inflamed part. It is one of the terminations of in-

flammation, and is owing to a change in the nutrition of the part.

INDURATION OF THE AREOLAR OR CEL'LULAR TISSUE, *Indura'tio cellulosa*, *Induratio* seu *Opilatio* te'læ cellulossæ neonatorum, *Ethmophytosis*, *Æde'ma neonatorum*, *Compact' Æde'ma* of *Infants*, *Cat'ochus Infantum*, *Scleremia*, *Sclerema*, *Sclero'ma*, *Scleroder'mia*, *Cul'tis ten'sa chronica*, *Dermatopericlerys'is*, *Dermatopericlerysmus*, *Phlegma'ia cellularia*, *Scirrhus'æa neonatorum*, *Scler'asis neonatorum*, *Sclero'sis*, *Stipatio* te'læ cellulossæ *Infantum*, *Skin'bound Disease*, *Skin'bind*, *Skin'tight*, *Hide'bound*, (F.) *Endurcissement du tissu cellulaire*; *Ædème du tissu cellulaire des nouveau-nés*, *Asphyxie lente des nouveau-nés*, *Squirrosarque*, is a disease which attacks infants a few days after birth, and which Chaussier proposed to call *Sclérème*, from *σκληρος*, 'hard.'

The Induration of the areolar tissue of the adult, *Sclerosteno'sis euta'nea*, has been regarded as the result of inflammation of the corium, *Chorion'tis*.

INDURATION OF THE BRAIN, *Sclerencephalia*—i. *Grise*, see Hepatization.

INDURATUS, (F.) *Induré*. Affected with induration—as 'an indurated chancre.'

INDURESCENTIA, Induration.

INDU'SIUM, (*inducere*, 'to put on,') *Chitonis'cus*. Strictly, 'a shirt,' 'a small tunic.' Some have so called the amnion.

INE'BRIANT, *Inebrians*, *Phantas'ticus*; (*in*, and *ebrio*, 'I intoxicate,') Intoxicating. An agent that intoxicates.

INEBRIATION, Temulentia.

INEBRIATION QUINIQUE, Quinism.

INEDIA, (*in*, and *edere*, 'to eat,') Fasting.

INERT'IA, (*in*, and *ars*, gen. *artis*, 'art,') *Ignavia*, *Ener'vitas*. Sluggishness, inactivity.

INERTIA OF THE WOMB, (F.) *Inertie de la matrice*. The diminution and even total cessation of the contractions of the uterus during labour; as well as the species of languor into which it sometimes falls after the expulsion of the foetus.

INERTIE PAR ÉPUISEMENT, ('inertia by exhaustion,') see Parturition—i. *de la Matrice*, *Inertia of the Womb*.

INESIS, (*ινησις*.) Cenosis.

INETHMUS, (*ινεθμος*.) Cenosis.

IN EXTREMIS, ('at the extremity or point of death,') Psychorages.

INFANCY, *Infan'tia*, (*in*, and *fans*, from *fari*, 'to speak,') Early childhood. Childhood; *Nepiotes*, *Paid'a*, (F.) *Enfance*. It generally includes the age from birth till the seventh year. See Age.

INFANS, gen. *Infantis*, *Pæ'dion*, *Pa'dion*, *Nepios*, *Puer*, (Sc.) *Gait'ling*, *Gyt'ling*, (Old Eng. and Prov.) *Barn*; (Sc.) *Bairn*, (F.) *Enfant*. An infant, a baby, a child, one in infancy. In law, one who has not attained the age of legal capacity; which is, in general, fixed at twenty-one years.

INFANS RECENS NATUS, *Nouveau-né*.

INFANTIA, Infancy.

INFANTICIDE, *Infanticidium*, *Tecnoctonia*, *Childmurder*, (*infans*, *infantis*, 'a child,' and *cædere*, 'to kill,') The murder of a child newly born, or on the point of being born. It may be perpetrated by the mother, or by some other person, either by commission, that is, in consequence of a direct voluntary act;—or by omission of those cares which are necessary for the preservation of the new-born.

INFARCTION, (from *in*, and *farcire*, *farcum*, 'to stuff,') Emphraxis.

INFARCTUS, Emphraxis—i. Hæmoptoicus *Laennecii*, see Hæmoptysis—i. Intestinorum, *Enteremphraxis*—i. Lactei extremitatum, *Phlegmatia dolens*—i. Lienis, *Splenoncus*.

INFARCTUS MAM'MÆ LACTEUS, *Lac'tis concretio'nes*, *No'di* seu *Throm'bi lac'tei*, *Trich'asis lac'*

ten. Knotty tumours of the female mammæ, owing to the accumulation and arrest of milk in the galactophorous ducts.

INFARCTUS UTERI, Metremphraxia.

INFECTANT, *Infectans*, (*infectere*, *infectum*, [*in*, and *facere*, 'to make']; 'to taint.') That which can taint or infect.

INFECTIEUX, see Contagious.

INFECTIO PURULENTA, see Pyohæmia.

INFECTION, see Contagion—i. Putrid, Septæmia.

INFECTIOUS, see Contagious.

INFECUNDITY, (*in*, and *fecundus*, 'fruitful.')

INFECUNDUS, Sterile.

INFERENT, see Affluent.

INFERNAL, *Infer'nus*; ('relating to hell.')

A name applied to caustic—*Lapis Infernalis*—on account of its strong burning properties.

INFIBULATIO, *Fibulatio*, *Infibulatio*, *Ancerias'mus*; (*in*, and *fibula*, 'a clasp.')

(F.) *Bowelement*. An operation, formerly practised, which consisted in passing a ring through the prepuce, after having drawn it over the glans;—in order to prevent coition. The ancients employed infibulation with their gladiators to preserve all their strength by depriving them of venery. In the women, to preserve their chastity, the ring was passed through the labia majora.

INFILTRATED, *Infiltratus*, (F.) *Infiltré*. Affected with infiltration.

INFILTRATION, *Infiltratio*, (*in*, and *filtrare*, *filtratum*, 'to filter.')

Effusion. The passage or effusion of a fluid into the areolæ of any texture, and particularly of the areolar membrane. The fluid effused is ordinarily the *Liquor sanguinis*, sound or altered—sometimes blood or pus, fæces, or urine. When infiltration of a serous fluid is general, it constitutes *anasarca*; when local, *œdema*.

INFILTRATION PURULENTE, see Hepatization.

INFINITESIMAL. *Infinitesimus*, (*in*, and *finitus*, 'limited.')

Unlimited. An epithet employed by the homeopaths for their endless subdivisions of drugs:—the decillionth of a grain, for example, is "an infinitesimal dose."

INFINITIVIST, see Emboîtement.

INFIRM, *Infirmus*, *Imbecillitas*, *Debilitas*, *Asthene*, *Hypocrætes*, (*in*, and *firmus*, 'firm.')

Not firm or sound; weak; feeble.

INFIRMARIUM, Hospital.

INFIRMARY, Hospital.

INFIRMATORIUM, Hospital.

INFIRMERIE, Hospital.

INFIRMIER (F.), *Infirmarius*, *Nosocomus*, *Cura'tor infirmorum*. One employed in a hospital or infirmary to take care of the sick.

INFIRMITY, *Infirmitas*, *Asthenia*, *Arrhositia*, *Invaletudo*. Any disease which has become habitual, either owing to its chronic character, or its numerous relapses.

INFIRMUS, *Abebasos*, *Infirm*.

INFLAMED, (Old Eng.) *Flamed*, *Inflammat'us*, (F.) *Enflammé*, *Phlogosé*,—*Phlogosé*, according to some,—(*in*, and *flamma*, 'flame, fire.')

Affected with inflammation. A sore, when inflamed, is said to look *irritated*, *angry*.

INFLAMMABLE AIR, Hydrogen, carburetted.

INFLAMMATIO, Inflammation, Phlegmon—i. Abdominalis, Encœlitis—i. Articuli Maxillæ Inferioris, Hyposiagonarthritidis—i. Auris, Otitis—i. Bronchiorum, Bronchitis—i. Cæci, Typhloenteritis—i. Capsulæ lentis, Phacohymenitis—i. Cerebelli, Cerebellitis—i. Clitoridis, Nymphitis—i. Conjunctivæ, see Ophthalmia—i. Corneæ, Ceratitis—i. Coxæ, Osphytitis—i. Crouposa, Croupy inflammation—i. Cystidis fellæ, Cholecystitis—i. Debilis, see Inflammation—i. Epiglottidis, Epiglottitis—i. Erysipelatosa, Erysipelatous inflammation—i. Faucium, Cynanche, Isthmitis—i.

Gastritis, Gastritis—i. Genæ, Gnathitis—i. Genæ, Melitis—i. Gingivæ, Ulitis—i. Glandularum lymphaticarum, Hydradenitis—i. Glandularum lymphaticarum, Lymphadenitis—i. Glandularum sublingualium, Hypoglossiadenitis—i. Gulæ, Œsophagitis—i. Hepatis, Hepatitis—i. Hepatis lenta, Hepatitis (chronic)—i. Interna, Empresma—i. Intestinorum, Enteritis—i. Iridis, Iritis—i. Jecoris, Hepatitis—i. Laryngis, Laryngitis—i. Lienis, Splenitis—i. Ligamentorum, Syndesmitis—i. Lingue, Glossitis—i. Mediastini, Mesodermis—i. Medullæ Spinalis, Myelitis—i. Musculi psoas, Psositis—i. Musculorum, Myositis—i. Musculorum abdominalium, Myocœlitis—i. Nervorum, Neuritis—i. Oculorum, Ophthalmia—i. Œsophagi, Œsophagitis—i. Omenti, Epiploitis—i. Ossis, Osteitis—i. Ovari, Oaritis—i. Palati, Hyperoitis—i. Pancreatis, Pancreatitis—i. Parenchymatica, Parenchymatitis—i. Parotidum, Parotitis, Cynanche parotidæa—i. Pectoris acuta, Pneumonia—i. Pericardii, Pericarditis—i. Perioste, Periostitis—i. Perioste orbitæ, Periorbitis—i. Peritonæi, Peritonitis—i. Pharyngis, Cynanche pharyngæa—i. Phrenitis, Phrenitis—i. Pleuræ, Pleuritis—i. Pneumonica, Pneumonia—i. Pulmonum, Pneumonia—i. Renum, Nephritis—i. Renum succenturiatorum, Paraneuritis—i. Retinæ, Dietyiti. Retinitis—i. Scleroticæ, Sclerotitis—i. Scroti, Oschitis—i. Septi transversi, Diaphragmitis—i. Sinuum frontalem, Metopantritis—i. Stomachi, Gastritis—i. Superficie internæ cordis, Endocarditis—i. Telæ cellulose, Ethmyphytis—i. Telæ fibrosæ, Inohymenitis—i. Testium, Hernia humoralis—i. Tonsillarum, Cynanche tonsillarum—i. Tunicæ hyaloideæ, Hyalitis—i. Tympani, Tympanitis—i. Urethræ, Gonorrhœa—i. Uteri, Metritis—i. Uteri catarrhalis, see Metritis—i. Uteri et peritonæi, Metropertitonitis—i. Uvulæ, Uvulitis—i. Vasorum, Angetitis—i. Vasorum lymphaticorum, Angioleucitis—i. Ventriculi, Gastritis—i. Ventriculi et intestinorum, Gastroenteritis—i. Vesicæ, Cystitis—i. Vesicæ fellæ, Cholecystitis.

INFLAMMATION, *Inflammat'io*, so called in consequence of the acute or burning pain felt in a part affected with it. *Phlegmoné*, *Phlog'osis*, *Ecphlog'osis*, *Epiphleg'ia*, *Exap'osis*, *Phlegma'cia*, *Causo'ma*, *Empres'ma*, *Phlog'ia*, *Hyperendosmose* (Dutrochet), *Inced'dium*, *Hyperhæmat'osis*, *Hyperæ'mia act'iva*, *Phleboptero'sis ecphrac'tica*; (F.) *Inflammation*, improperly *Angiite*. Altered nutrition in a part is occasioned by some agency;—owing to which the blood flows into the capillary vessels in greater abundance than natural, and those vessels become over-dilated and enfeebled; whence result pain, redness, heat, tension, and swelling; symptoms which appear in greater or less severity, according to the structure, vital properties, and functions of the part affected, and its connexion with other parts, as well as according to the constitution of the individual.

Simple accumulation of blood in the capillaries of the part is *hyperæmia*, not inflammation. The latter requires, in addition, altered nutrition of the vessels themselves, and the surrounding tissues.

The inflammations of the areolar and serous membranes greatly agree;—and those of the mucous and skin; the former being more active, and constituting the *phlegmonous* variety;—the latter, the *erythematic* or *erysipelatous*. Of this variety is the *diffusive inflammation* produced by morbid poisons,—as during dissection, where solutions of continuity exist in the fingers of the operator. It is seen, too, in workers in bone, and hence has been called *bone fever*. Preparatory to the turning of bones, it is customary to macerate them in water. The fluid soon becomes putrid, and if the hands be kept in it diffusive inflammation results.

Inflammation may end by resolution, suppuration, gangrene, adhesion, effusion, or induration. Each of the inflammations of internal organs has received a name according to the organ affected;—as, *gastritis, cephalitis, enteritis, hepatitis, &c.* Besides the above inflammations, it has been considered, that there is an instinctive kind established for the union of parts which have been divided, whether the union takes place immediately, or by the aid of an intermediate body. This is the *adhesive inflammation* of some; but it may well be questioned whether inflammation is ever desirable in such cases. See Adhesion, Callus, Cicatrix. Broussais considered that the term *inflammation* should include every local exaltation of the organic movements which is sufficiently great to disturb the harmony of the functions, and disorganize the texture in which it is situated. He farther extended the name *inflammation* to irritations which do not induce disorganization of the textures, and which had been previously, and are still, called *fevers*.

Examination of the blood drawn always exhibits an increase of the fibrinous element—the average proportion of which, in healthy blood, is about three in the thousand. In inflammation, it at times rises as high as ten. In fevers unaccompanied with inflammation, the proportion is natural, or below the average; but whenever inflammation supervenes, it immediately rises.

External inflammation is easily detected by the characters already mentioned:—*internal*, by disturbance of function and pain upon pressure; but the last sign is often not available. Both forms require the removal of all irritation, and the reduction of vascular excitement and nervous irritability; hence, blood-letting—local and general—sedatives, refrigerants, and counter-irritants become valuable remedies in almost all cases of inflammation.

Inflammation is not always of the active kind. There is a form, *Inflammatio debilis*, which occurs in low conditions of the system, and in broken down constitutions, which, like many forms of chronic inflammation, requires a stimulant, rather than a sedative, mode of treatment.

INFLAMMATION DES AMYGDALES, Cynanche tonsillaris—*i. des Arrières*, Arteritis—*i. of the Bladder*, Cystitis—*i. de la Bouche*, Stomatitis—*i. of the Brain*, see Encephalitis, and Phrenitis—*i. des Bronches*, Bronchitis—*i. du Cæcum*, Typhloenteritis—*i. of the Cæcum*, Typhlo-enteritis—*i. du Cerveau et du Cervelet*, Phrenitis—*i. du Cœur*, Carditis—*i. du Colon*, Colitis—*i. de la Conjonctive*, see Ophthalmia—*i. of the Diaphragm*, Diaphragmatitis—*i. Diffusive*, see Inflammation—*i. Diphtheritic*, Diphtheritis—*i. of the Ear*, Otitis—*i. de l'Épiglotte*, Epiglottitis—*i. de l'Estomac*, Gastritis—*i. of the Eye*, Ophthalmia—*i. du Foie*, Hepatitis—*i. des Gencives*, Ulitis—*i. General*, Synocha—*i. of the Internal Membrane of the Heart*, Endocarditis—*i. Internal*, Empresma—*i. des Intestins*, Enteritis—*i. of the Iris*, Iritis—*i. of the Kidney*, Nephritis—*i. de la Langue*, Glossitis—*i. of the Larynx*, Laryngitis—*i. of the Liver*, Hepatitis—*i. de la Luette*, Uvulitis—*i. of the Lungs*, Pneumonia—*i. of the Malpighian bodies*, Kidney, Bright's disease of the—*i. des Mamelles*, Mastitis—*i. de la Matrice*, Metritis—*i. de la Membrane alvéolo-dentaire*, Periodontitis—*i. de la Membrane séreuse céphalo-rachidienne*, Meningitis—*i. of the Mesentery*, Mesenteritis—*i. de la Moëlle épinière ou rachidienne*, Myelitis—*i. of the Mouth*, pseudomembranous, Stomatitis, pseudo-membranous—*i. of the Mouth*, pultaceous, Aphthæ—*i. des Muscles*, Myositis—*i. des Nerfs*, Neuritis—*i. de l'Œil*, Ophthalmia—*i. de l'Oreille*, Otitis—*i. de l'Ovaire*, Ovaritis—*i. du Palais*, Hyperoitis—*i. du Parenchyme pulmonaire*, Pneumonia—*i. de la Parotide*,

Cynanche parotidea—*i. Pellicular*, Diphtheritis—*i. du Péricarde*, Pericarditis—*i. of the Pericardium*, Pericarditis—*i. du Péritoine*, Peritonitis—*i. of the Peritonæum*, Peritonitis—*i. of the Pleura*, Pleuritis—*i. de la Plèvre*, Pleuritis—*i. des Poupmons*, Pneumonia—*i. du muscle Psoas*, Psositis—*i. de la Rate*, Splenitis—*i. des Reins*, Nephritis—*i. Spongoid*, Hamatodes Fungus, see also Encephaloid—*i. of the Stomach*, Gastritis—*i. of the Testicle*, Hernia humoralis—*i. de la Testicule*, Hernia humoralis—*i. des Tissus blancs*, Angioleucitis—*i. of the Tongue*, Glossitis—*i. des Vaisseaux lymphatiques*, Angioleucitis—*i. des Veines*, Phlebitis—*i. de la Vésicule du Fiel*, Cholecystitis—*i. de la Vessie*, Cystitis—*i. of the Womb*, Metritis.

INFLAMMATIUN' CULÀ, (dim. of *inflammatio*.) *Subinflammatio*. A superficial and often insignificant inflammation of the skin, as in many cutaneous affections.

INFLAMMATOIRE, Inflammatory.

INFLAMMATORY, *Inflammato'rius*; *Phlog'icus*, *Causo'des*, *Phlogo'des*, *Phlogisticus*, *Phlogis'tic*, (F.) *Inflammatoire*, *Phlegmasique*, belonging to inflammation;—as, *inflammatory tumour*, *inflammatory fever*, &c. The blood is said to be *inflammatory* when cupped or buffed. See *Corium phlogisticum*.

INFLAMMATORY BLUSH, Erythema.

INFLAMMATUS, Inflamed.

INFLATIO, (in, and *flare*, *flatum*, 'to blow.') Emphysema, Puffiness, Colica flatulenta—*i. Parotidum*, Cynanche parotidea—*i. Uteri*, Physometra.

INFLEXIO, (in, and *flectere*, *flectum*, 'to bend.') Camptis.

INFLUENCE, Influenza.

INFLUENTIA, Influenza.

INFLUEN'ZA. The Italian for 'Influence.' *Influenza Europa'a*, *Influent'ia*, *Catar'rhus epidem'icus*, *C. a contagio*, *Fébris catarrh'alis epidem'ica*, *Rheum'a epidem'icum*, *Morbus Vervecinus seu Catarrah'alis seu Ari'etis*, *Syn'ochus seu Deflux'io catarrh'alis*, *Cephalal'gia contagiosa* (epidemics of the 16th and 17th centuries): (F.) *Tac*, *Quinte*, *Florion*, *Coqueluche*, *Baraquette*, *Générale*, *Grippe*, *Follette*, *Grenade*, *Coquette*, *Cocote*, *Petite Poste*, *Petit Courier*, *Allure*, *Lodendo*, *Fièvre catarrhale épidémique*; *Influence*, *Epidemic catarrh*. A severe form of catarrh occurring epidemically, and generally affecting a number of persons in a community. See Catarrh, epidemic. Gluge, from his investigations, considers that the following is the chronological order of the return of the influenza:—14th century, 1323, 1326—15th century, 1410, 1411, 1414—16th century, 1510, 1557, 1562, 1574, 1580, and 1593—17th century, 1658, 1669, 1675, 1693—18th century, 1708, 1712, 1729, 1732, 1733, 1742, 1743, 1761, 1762, and 1775—19th century, 1800, 1803, 1831, and 1833. To these may be added 1837, and 1843.

INFLUENZA EUROPEA, Influenza.

INFLUEN'ZOID, *Influenzoï'des*, (influenza, and *aidos*, 'resemblance.') Resembling influenza.

INFORMITAS, (in, and *forma*, 'form,') Deformation.

INFORTUNIUM, 'misfortune,' (in, and *fortuna*, 'fortune,') Contrastsura.

INFRA, as a prefix, beneath.

INFRAATLOIDÆUS, Sub-atloidæus.

INFRAAXOIDÆUS, Sub-axoidæus.

INFRACONSTALES, see Intercoastal muscles.

INFRADIAPHRAGMATIC, *Infradiaphragmaticus*, (F.) *Sousdiaphragmatique*. Situate under the diaphragm.

INFRAMAXILLARIS, Sub-maxillary.

INFRAMAXILLOSTERNODYMIA, Cephalosomatodymia.

INFRAORBITAR, Suborbitar.

INFRAPUBIAN. *Infra-pubianus*, (F.) *Sous-pubien*. That which is situate beneath the pubis. See *Subpubis*.

INFRAPUBIAN LIGAMENT, Triangular ligament.

INFRASCAPULARIS, Subscapularis.

INFRASPINALIS, Infra-spinatus.

INFRASPINATUS, *Infra-spinatus*, (infra, and *spina*, 'spine.') Situate beneath the spine of the scapula.

INFRASPINATA FOSSA, (F.) *Fosse sous-épineuse*. A large excavation on the posterior surface of the scapula, beneath its spine. It is filled by the

INFRASPINATUS (*Muscle*, *Grand Scapulo-trachetérien*, *Suprascapularis inferior* (Ch.), (F.) *Sous-épineux*, which is broad, flat, and triangular. It is attached, by its base, to the three inner quarters of the fossa: and is inserted, by a long tendon, into the middle part of the great tuberosity of the os humeri (*Trochiter*.) It turns the arm outwards, and, when the arm is elevated, carries it backwards.

INFRICTION, (in, and *fricare*, *frictum*, 'to rub,') Inunction.

INFRINGENS, (in, and *frangere*, *fractum*, 'to break,') Corrigent.

INFUNDIBULUM, (in, and *fundere*, *fusum*, 'to pour,') A Funnel, *Tritorium*, *Chonch*, *Chonch*, (F.) *Entonnoir*,—of which various kinds are employed in pharmaceutical operations. A name, given to many parts which, more or less, resemble a funnel. See *Proventriculus*, and *Pulmo*. It is particularly appropriated to the following organs:—

INFUNDIBULUM OF THE BRAIN, *Infundibulum* seu *Pellis* seu *Choncha* seu *Cylindrus* seu *Seyphus* seu *Concha* seu *Lacuna* seu *Embolus* seu *Aqueductus* seu *Lacrum* seu *Concavitas conchularis* seu *Processus orbicularis Cerebri*, (F.) *Entonnoir du ventricule moyen du cerveau*, *Tige Pituitaire*, *Tige sub-sphénoïdale*. A depression in the inferior paries of the middle ventricle, above the pituitary gland. It was, anciently, regarded as a canal by which the fluid, collected in the ventricles of the brain, was evacuated, and poured into the nasal fossæ.

INFUNDIBULUM CEREBRI, I. of the Brain.

INFUNDIBULUM OF THE COCHLEA, I. *Cochlea*, *Seyphus* *Vicusenii* seu *auditorium*, *Cucullus*, *Cannalis* *Scala*rum *communis*. This, with the modiolus, forms the nucleus around which the gyri of the cochlea pass. It is an imperfect funnel, the apex of which is common with that of the modiolus: and the base is covered with the apex of the cochlea, termed *Cupula*.

In *Surgery*, infundibula or funnels are used to direct steam or vapours; to conduct the actual cautery to certain morbid parts, &c.

INFUNDIBULUM OF THE ETHMOID BONE, or OF THE NASAL FOSSÆ. One of the anterior cells of that bone, which is broad and expanded above, and narrow below; opening, above, into the frontal sinus; below, into the anterior part of the middle meatus of the nasal fossæ.

INFUNDIBULUM OF THE HEART, *Conus arteriosus*—I. of the Kidney, see *Calix*—I. Lacrymale, Lacrymal Sac—I. Tubarum Fallopii, see *Tube*, *Fallopian*—I. Ventriculi, *Œsophagus*.

INFUSE, Infusion.

INFUSION, *Infusio*, *Enchyseis*, (*infundere* (in, and *fundere*, *fusum*), 'to pour in, 'to introduce,') A pharmaceutical operation, which consists in pouring a hot or cold fluid upon a substance whose medical virtues it is desired to extract. *Infusion* is, also, used for the product of this operation. *Infusum*, *Enchyasma*, (F.) *Infusion*, *Infusé*. In *Surgery*, *Infusion*—*Chirurgia infusoria*, *Art clysmatica nova*—is the act of introducing into the veins medicinal substances, by aid of an instrument called *Infusor*. See *Infusoir*. This mode

of introducing medicines was called *Art infusoria*.

INFUSION OF ANGUSTURA, *Infusum Cuspariæ*—I. of Bark, *Infusum cinchonæ*—I. of Buchu, *Infusum Diœmæ*—I. of Calumba, *Infusum calumbæ*—I. of Cascarella, *Infusum cascariellæ*—I. of Catechu, *Infusum catechu compositum*—I. of Cayenne pepper, *Infusum capsici*—I. of Chamæmelis, *Infusum anthemidis*—I. of Chiretta, *Infusum Chiretæ*—I. of Cichæna, *Infusum cichonæ*—I. of Cloves, *Infusum Caryophyllorum*—I. of Cusparia, *Infusum cuspariæ*—I. of Dandelion, *Infusum Taraxaci*—I. of Ergot, *Infusum Ergotæ*—I. of Foxglove, *Infusum digitalis*—I. of Gentian compound, *Infusum gentianæ compositum*—I. of Ginger, *Infusum Zingiberis*—I. of Hops, *Infusum humuli*—I. of Horseradish, compound, *Infusum armoraciæ compositum*—I. of Kousso, *Infusum Cusso*—I. of Linseed, *Infusum lini compositum*—I. of Matioe, *Infusum matico*—I. of Mint, compound, *Infusum menthæ compositum*—I. of Orange-peel, compound, *Infusum aurantii compositum*—I. of Pareira Brava, *Infusum Pareiræ*—I. of Pinkroot, *Infusum spigeliæ*—I. of Quassia, *Infusum quassiæ*—I. of Rhubarb, *Infusum Krameriæ*—I. of Rhubarb, *Infusum rhei*—I. of the Rose, *Infusum rosæ compositum*—I. of Roses, acid, *Infusum Rosæ acidum*—I. of Sage, *Infusum salviæ*—I. of Sarsaparilla, *Infusum sarsaparillæ*—I. of Sassafras pith, *Infusum sassafrasæ medullæ*—I. of Seneka, *Infusum Senegæ*—I. of Senna, *Infusum sennæ compositum*—I. of Simaruba, *Infusum simarubæ*—I. of Slippery elm, *Infusum ulmi*—I. of Thoroughwort, *Infusum eupatorii*—I. of Tobacco, *Infusum tabaci*—I. of Valerian, *Infusum valerianæ*—I. of Virginia snake-root, *Infusum serpentariæ*—I. of Wild cherry, *Infusum pruni Virginianæ*.

INFUSOIR (F.), *Infusoir*. An instrument for injecting medicinal substances into the veins. It was a kind of funnel, the elongated apex of which was stopped by a metallic rod, which could be withdrawn when the apex was introduced into a vein.

INFUSUM, Infusion.

INFUSUM ACACIÆ CATECH, I. catechu compositum—I. Amarus vinosus, Vinum gentianæ compositum—I. Angusturæ, I. cuspariæ.

INFUSUM ANTHEMIDIS, *Hydrochamæmelum*, *Infusio* of Chamæmelis. *Anthemid. flor.* $\frac{3}{ss}$: *Aq. bullient.* Oj. Macerate for ten minutes in a covered vessel, and strain. Dose, $\frac{f\frac{3}{ij}}$ to $\frac{f\frac{3}{iss}}$.

INFUSUM ARMORACIÆ COMPOSITUM, I. *Armoraciæ* (Ph. U. S.) Compound Infusion of Horseradish. (*Armorac. concis.*, *Sinapis* cont., sing. $\frac{3}{j}$; *Aq. bullient.* Oj. Macerate for two hours, and strain.) Dose, $\frac{f\frac{3}{ij}}$ to $\frac{f\frac{3}{ij}}$.

INFUSUM AFRANTH COMPOSITUM, Compound Infusion of Orange Peel. (*Aurant. cort. sicæ.* $\frac{5}{ij}$; *Limon. cort. recent.* $\frac{3}{j}$; *Caryoph. cont.* $\frac{3}{ss}$: *Aq. fervent.* Oss. Macerate for fifteen minutes, and strain.—Ph. L.) Dose, $\frac{f\frac{3}{iss}}$ to $\frac{f\frac{3}{ij}}$.

INFUSUM BRASHI, Wort—I. Buchu, *Infusum Diœmæ*—I. Byres, Wort.

INFUSUM CALUMBÆ, I. *Calumbæ* seu *Calumbæ*, *Infusio* of Calumbæ. (*Calumb. rad. concis.* $\frac{3}{ss}$: *Aq. fervent.* Oj. Macerate for two hours, and strain. Made also by displacement.—Ph. U. S.) Dose, $\frac{f\frac{3}{iss}}$ to $\frac{3}{ij}$.

INFUSUM CAPSICI, *Infusio* of Cayenne Pepper. (*Capsic. in. pulv. crass.* $\frac{3}{ss}$: *Aq. bullient.* Oj. Macerate for two hours, and strain.—Ph. U. S.) Dose, $\frac{f\frac{3}{iss}}$.

INFUSUM CARYOPHYLLI, *Infusio* of Cloves. (*Caryoph. contus.* $\frac{3}{j}$; *Aq. bullient.* Oj. Macerate for two hours, and strain.) Dose, $\frac{f\frac{3}{iss}}$ to $\frac{f\frac{3}{ij}}$.

INFUSUM CASCARIÆ, *Infusio* of Cascarella. (*Cascariell. contus.* $\frac{3}{j}$; *Aq. bullient.* Oj. Macerate for two hours, and strain.) Made also by displacement, Ph. U. S. Dose, $\frac{f\frac{3}{iss}}$ to $\frac{f\frac{3}{ij}}$.

INFUSUM CASSIÆ SENNÆ, *I. sennæ compositum*.

INFUSUM CAT'ECU COMPOS'ITUM, *I. Catechu seu Aca'eia Catechu, Infusion of Cat'echu*. (Catechu, in pulv. $\frac{3}{4}$ ss; Cinnamon. cont. $\frac{3}{4}$ j; Ag. bullient. Oj. Macerate for an hour, and strain. Ph. U. S.) Dose, f $\frac{3}{4}$ iss to f $\frac{3}{4}$ ij.

INFUSUM CHIRET'Æ, *I. Chiraitæ* (Ph. B. 1864.) *Infusion of Chiretta*. (Chiretta, $\frac{3}{4}$ v; Ag. fervent. Oj. (Imp. meas.) Infuse for two hours, and strain. Dose, as a tonic, f $\frac{3}{4}$ j to f $\frac{3}{4}$ ij.

INFUSUM CINCHONÆ, *I. Cinchonæ lancifoliae, Infusion of Cincho'na*. (Cinch. contus. $\frac{3}{4}$ j; Ag. fervent. Oj. Macerate for two hours, and strain.) This infusion may also be made from the same quantity of bark in coarse powder by the process of displacement with hot or cold water. Dose, f $\frac{3}{4}$ j to f $\frac{3}{4}$ ij. The Pharmacopœia of the United States has an *Infusum Cincho'ne flavae, Infusion of yellow cinchona*, and an *Infusum Cinchonæ rubrae, Infusion of red cinchona*—*Infusum Cincho'ne Compositum* (Ph. U. S. 1851)—which are made as follows: (Cinchon. flav. vel rubr. in pulv. $\frac{3}{4}$ j; Acid. sulph. aromat. f $\frac{3}{4}$ j; aquæ. Oj.) Made by displacement. Dose, same as the last.

INFUSUM CINCHONÆ COMPOSITUM, see *Infusum Cinchonæ*—*i. Cinchonæ flavæ*, see *Infusum Cinchonæ*—*i. Cinchonæ lancifoliae*, *Infusum Cinchonæ*—*i. Cinchonæ rubrae*, see *Infusum Cinchonæ*—*i. Columbae*, *I. Calumbæ*.

INFUSUM CUSPARIÆ, *I. Angusturæ, Infusion of Cusparia* (Cuspar. cort. contus. $\frac{3}{4}$ ss; Ag. bullient. Oj. Macerate for two hours, and strain. Made also by displacement.) Dose, f $\frac{3}{4}$ j to f $\frac{3}{4}$ ij.

INFUSUM CUS'SO, *Infusion of Kou'sso*, (Kou'sso, in coarse powder, $\frac{3}{4}$ l; boiling Distilled water, f $\frac{3}{4}$ iv. Infuse in a covered vessel for fifteen minutes, without straining. Ph. B. 1864.) For a dose.

INFUSUM DIGITALIS, *I. Digitalis purpurea, Infusion of Fox'glove*. (Digital. $\frac{3}{4}$ j; Ag. bullient. Oss; Tinct. cinnamomi, f $\frac{3}{4}$ j. Macerate for two hours, and add sp. cinnam. f $\frac{3}{4}$ j.) Dose, f $\frac{3}{4}$ j.

INFUSUM DIOS'MÆ, (Ph. U. S., 1842.) *I. Buch'u*, (Ph. U. S., 1851.) *I. Buch'co*, (Ph. B. 1864.) *Infusion of Buch'u*. (Diosm. $\frac{3}{4}$ j; Ag. bullient. Oj. Macerate for two hours in a covered vessel, and strain.—Ph. U. S.) Dose, f $\frac{3}{4}$ iss.

INFUSUM ER'GOTÆ, *Infusion of Ergot*. (Ergot. in pulv. crass. $\frac{3}{4}$ j; Ag. bullient. f $\frac{3}{4}$ ix.) Infuse for an hour, in a covered vessel, and strain. Ph. D.) Dose, as a parturifacient, f $\frac{3}{4}$ ij.

INFUSUM EUPATORI, *Infusion of Tho'rough-wort*. (Eupator. $\frac{3}{4}$ j; Ag. bullient. Oj. Macerate for two hours, and strain.—Ph. U. S.) Dose, f $\frac{3}{4}$ ij.

INFUSUM GENTIANÆ COMPOS'ITUM, *Compound Infusion of Gentian*. (Gentian. cont. $\frac{3}{4}$ ss; Aurant. amar. cort., Coriandr. contus. $\frac{3}{4}$ ss; Alcohol. dilut. f $\frac{3}{4}$ iv; Aquæ, f $\frac{3}{4}$ xij. First pour on the alcohol, and, three hours afterwards, the water; then macerate for twelve hours, and strain. Prepared also by displacement.—Ph. U. S.) Dose, f $\frac{3}{4}$ j to f $\frac{3}{4}$ ij.

Mars'den's Anticorbutic Drops, an empirical preparation, consist of a solution of corrosive sublimate in an infusion of gentian.

INFUSUM HUMULI, *I. Lupuli, Infusion of Hops*. (Humul. $\frac{3}{4}$ ss; Ag. bullient. Oj. Macerate for two hours, and strain.—Ph. U. S.) Dose, f $\frac{3}{4}$ iss to f $\frac{3}{4}$ ij.

INFUSUM JUNIP'ERI, *Infusion of Ju'niper*. The common mode of making this infusion is by an ounce of Juniper Berries to a pint of boiling water. The whole may be taken in the twenty-four hours.

INFUSUM KRAMERIÆ, *Infusion of Rhat'any*, (Kramer. contus. $\frac{3}{4}$ j; Ag. bullient. Oj. Macerate for four hours, and strain. Made also by displacement.—Ph. U. S.) Dose, f $\frac{3}{4}$ iss.

INFUSUM LINI, *I. L. compositum*.

INFUSUM LINI COMPOS'ITUM (Ph. U. S., 1851), (*I. Lini*, 1842), *I. Lini usitatis'simi, Infusion of Lin'seed, Flax'seed Tea*. (Lini sem. cont. $\frac{3}{4}$ ss; Glycyrrh. rad. cont. $\frac{3}{4}$ j; Aquæ bullient. Oj. Macerate for two hours, and strain.) Dose, a teacupful, ad libitum.

INFUSUM LUPULI, *Infusum Humuli*—*i. Malti, Wort*.

INFUSUM MATI'CO, *I. Mati'ce, Infusion of Mati'co*. (Matico, $\frac{3}{4}$ ss (avoirdupois); Ag. bullient. Oss (Imp. meas.) Infuse for an hour, and strain.—Ph. D.) Dose, f $\frac{3}{4}$ iss to f $\frac{3}{4}$ ij.

INFUSUM MEN'THÆ VIRIDIS, *Infusion of Spear'mint, Mint Tea*. (Mentha virid. $\frac{3}{4}$ j; Aquæ bullient. Oss. Infuse for an hour, and strain.—Ph. D.) Dose, f $\frac{3}{4}$ j to f $\frac{3}{4}$ ij.

INFUSUM PAREIRÆ, *Infusion of Pareira Brava*. (Pareir. $\frac{3}{4}$ j; Ag. bullient. Oj.) Infuse for two hours, and strain. (Ph. U. S. 1863.) Dose, f $\frac{3}{4}$ j or f $\frac{3}{4}$ ij.

INFUSUM PICIS EMPYREUMATICÆ LIQUIDÆ, see *Pinus sylvestris*—*i. Picis liquidæ*, see *Pinus sylvestris*—*i. Polygalæ*, *Infusum Senegæ*.

INFUSUM PRUNI VIRGINIÆ, *Infusion of Wild Cherry Bark*. (Prun. Virginian. cont. $\frac{3}{4}$ ss; Aquæ, Oj. Macerate for 24 hours, and strain. Prepared, also, by displacement. Ph. U. S. 1863.) Dose, f $\frac{3}{4}$ j to f $\frac{3}{4}$ ij.

INFUSUM QUAS'SIÆ, *I. Quassie excell'æ, Infusion of Quassia*. (Quassia liqn. cont. $\frac{3}{4}$ j; Ag. (frigid.) Oj. Macerate for twelve hours, and strain.—Ph. U. S.) Dose, f $\frac{3}{4}$ j to f $\frac{3}{4}$ iv.

INFUSUM RHEI, *An'ima Rhei, Infusion of Rhubarb*. (Rhei, cont. $\frac{3}{4}$ j; Ag. ferv. Oss. Macerate for two hours, and strain.—Ph. U. S.) Dose, f $\frac{3}{4}$ j to f $\frac{3}{4}$ iv.

INFUSUM RO'SÆ AC'IDUM, *Acid Infusion of Ro'ses*. (Ros. Gallic. $\frac{3}{4}$ l; Acid. Sulph. d. f $\frac{3}{4}$ j; Ag. destillat. bullient. f $\frac{3}{4}$ x. Add the acid to the water; infuse the petals in the mixture for half an hour, and strain. Ph. B. 1864.)

INFUSUM RO'SÆ COMPOS'ITUM, *I. Rosæ Gallicæ, Infusion of the Rose*. (Ros. Gallic. $\frac{3}{4}$ v; Ag. bullient. Oiss; Acid. sulph. d. f $\frac{3}{4}$ ij; Sacch. purif. $\frac{3}{4}$ ss. Add the water, and afterwards the acid;—macerate for half an hour; strain, and add the sugar.—Ph. U. S.) Dose, f $\frac{3}{4}$ iss to Oss.

INFUSUM SALV'Æ, *Infusion of Sage, Sage Tea*. (Salv. $\frac{3}{4}$ ss; Ag. bullient. Oj; macerate for half an hour, and strain.)

INFUSUM SARSAPARILLÆ, *Infusion of Sarsaparilla*. (Sarsaparill. contus. $\frac{3}{4}$ j; Aquæ bullient. Oj. Digest for two hours in a covered vessel, and strain.—Ph. U. S.) It may also be prepared by displacement. Dose, f $\frac{3}{4}$ ij to f $\frac{3}{4}$ iv.

INFUSUM SASS'AFRAS, *I. Sassafras medullæ, Mucila'go Sassa'fras, Infusion of Sassa'fras Pith, Mucilage of Sassa'fras Pith*. (Sassafras medull. $\frac{3}{4}$ j; Aquæ (frigid.) Oj. Macerate for three hours, and strain.—Ph. U. S.) An emollient collyrium, and a demulcent drink.

INFUSUM SEN'EGÆ, *Infusum Polygalæ, Infusion of Sen'eka*. (Seneg. $\frac{3}{4}$ x; Ag. bullient. (Imp. meas.) Infuse for four hours, and strain.—Ph. E.) Dose, f $\frac{3}{4}$ j to f $\frac{3}{4}$ ij.

INFUSUM SENNÆ, *I. S. Compositum, I. Cas'sie Sennæ seu Sennæ simplex, Aqua laxati'va Vienne'nsis, Infusion of Senna*. (Senna, $\frac{3}{4}$ j; Coriandr. cont. $\frac{3}{4}$ j; Ag. bullient. Oj. Macerate for an hour, and strain.—Ph. U. S.) Dose, f $\frac{3}{4}$ j to f $\frac{3}{4}$ iv, as a cathartic.

The Black Draught, Black Dose, Black bottle, Haus'tus ni'ger, is usually formed by this infusion. It may be made of *Infus. sennæ*, f $\frac{3}{4}$ v; Ag. cinnam. f $\frac{3}{4}$ j; Mannæ, $\frac{3}{4}$ v; Magnæ. sulph. $\frac{3}{4}$ vj. Dose, a wineglassful.

Sel'way's Prepared Essence of Senna is a concentrated infusion of the leaves in combination with an alkali.

INFUSUM SENNE COMPOSITUM, *I. sennæ.*

INFUSUM SERPENTARIE, *Infusion of Virgin's Snake-root.* (Serpentar. $\frac{3}{4}$ ss; Aq. bullient. Oj. Macerate for two hours, and strain. Prepared, also, by displacement.—Ph. U. S.) Dose, f $\frac{3}{4}$ ss.

INFUSUM SIMARUBÆ, *Infusion of Simaruba.* (Simarub. cort. cont. $\frac{3}{4}$ ij; Aq. fervent. Oj. Macerate for two hours, and strain.—Ph. E.) Dose, f $\frac{3}{4}$ ij to f $\frac{3}{4}$ ij.

INFUSUM SPIGELLÆ, *Infusion of Pinkroot.* (Spigel. $\frac{3}{4}$ ss; Aq. bullient. Oj. Macerate for two hours, and strain.—Ph. U. S.) Dose, f $\frac{3}{4}$ iv to Oss.

INFUSUM TABACI, *Infusion of Tobacco.* Tabac. fol. $\frac{3}{4}$ j; Aq. ferv. Oj. Macerate for an hour, and strain.—Ph. U. S.) Half of this per anum in strangulated hernia.

INFUSUM TARAXACI, *Infusion of Dandelion.* (Tarazac. contus. $\frac{3}{4}$ ij; Aq. bullient. Oj. Macerate for two hours, and strain.—Ph. U. S.) Dose, f $\frac{3}{4}$ ss, as a diuretic, &c.

INFUSUM ULMI, *Mucilago Ulmi, Infusion of Slippery Elm, Slippery Elm Tea, Mucilage of Slippery Elm bark.* This preparation, in the Pharmacopœia of the United States, is made by infusing one ounce of slippery elm bark in a pint of boiling water.

INFUSUM VALERIANÆ, *Infusion of Valerian.* (Rad. valerian. in pulv. crass. $\frac{3}{4}$ ss; Aq. bullient. Oj. Macerate for an hour, and strain. Prepared, also, by displacement.—Ph. U. S.) Dose, f $\frac{3}{4}$ ss to $\frac{3}{4}$ ij.

INFUSUM ZINGIBERIS, *Infusion of Ginger, Ginger Tea.* (Zingib. contus. $\frac{3}{4}$ ss; Aq. bullient. Oj. Macerate for two hours, and strain.—Ph. U. S.) Dose, f $\frac{3}{4}$ ss to f $\frac{3}{4}$ ij, as a carminative.

ING'A. A genus of plants, *Ord. Leguminosæ*, belonging to South America, many of the species of which are possessed of astringent properties. *Inga bark* is used in South America as a tonic astringent in diarrhœa, gonorrhœa, &c. An extract, prepared from it, possesses all the properties of rhatany.

INGENIUM (Morbi), (*in*, and *geno*, for *gigno*, 'to generate.') The nature or genius of a disease. This word is employed, especially by the French, synonymously with *nature*. They speak, for instance, of *Génie inflammatoire, bilieux*, &c. Some, also, use *génie* in the place of *type* of an intermittent.

INGESTA, (*in*, and *gerere, gestum*, 'to bear or carry.') Substances, introduced into the body by the digestive passages; as food, condiments, drinks, &c.

INGLUVIES, *Aples'tia, Vic'tus intemperantia*: Gluttony, Insatiableness; also, the *Crop* or *Craw* or *Gorge* of Birds, (F.) *Jabot*; and the *first stomach* or *paunch* of ruminant animals—*Pen'ula, Ru'men, Pan'ties, Ven'ter magnus*, (F.) *Panse, Herbie, Double*. Also, the Pharynx.

INGRASCIAS, APOPHYSES OF. The lesser alæ of the sphenoid bone.

INGRAVIDATION, (*in*, and *gravidus*, 'pregnant,') Fecundation, Pregnancy.

INGRESSUS SUPERIOR, ('superior entrance,') Cardia.

INGUEN, *gen. In'guinis, Æde'on, Bu'bon, Bu'bo, Groin*, (Prov.) *Lank, Last, Lisk, Lesk*; (F.) *Aine*. The oblique fold or depression which separates the abdomen from the thigh. It is only, properly speaking, a line that extends from the anterior and superior spinous process of the ilium to the middle part of the horizontal ramus of the pubis. Also, the genital organs.

INGUÉRISABLE, (F.) (*in*, and *guérir*, 'to cure,') Incurable.

INGUINAL, *Inguinal'is*, (from *inguen*.) Belonging or relating to the groin. This epithet has been given to various parts met with in the region of the groin or inguinal region.

INGUINAL AR'TERY is that portion of the femoral artery situate immediately beneath the crural arch in the inguinal region.

INGUINAL CANAL' is a canal, about two inches in length, proceeding obliquely downwards, inwards and forwards at the lower part of the abdomen; through which passes the spermatic cord, in men, and the round ligament of the uterus in women. This canal is formed, inferiorly and anteriorly, by the aponeurosis of the greater oblique muscle; posteriorly by the *fascia transversalis*, which is joined to the preceding aponeurosis, and forms with it a deep channel, into which are received the lesser oblique and transversalis muscles. The inguinal canal has two apertures; the one, the *lower* and *inner*, is called the *inguinal* or *abdominal ring*. It is bounded by two strong tendinous pillars which fix it—the innermost to the symphysis, the outermost to the spine of the pubis. The *upper* and *outer* aperture is formed by the *fascia transversalis*. From the edge of this aperture arises a membranous funnel,—a prolongation of the *fascia transversalis*,—which receives the spermatic vessels; forms their proper sheath, and accompanies them as far as the testicle. On its inside lies the epigastric artery. Above it, is the lower edge of the transversalis muscle: and, below, it is bounded by the channel of the greater oblique. By following the oblique direction of this canal, and passing, consequently, on the outside of the epigastric artery, the viscera are displaced, so as to constitute internal inguinal hernia.

INGUINAL GLANDS, *Glan'dule inguinal'es*, (F.) *Glandes inguinales*, are the lymphatic glands or ganglions of the groin. There are two sets, one *superficial*; the other *deep-seated*; the former are larger, and their number varies, but on the average there are eight or ten which are placed irregularly about Poupart's ligament. The latter, or deep-seated, are situate behind the others around the femoral artery and vein.

INGUINAL RE'GION, *Re'gio inguinal'is, Bubo, Inguen*. The region of the groin.

INGUINAL RING, *Abdom'inal Ring, An'nulus abdom'inis*, (F.) *Anneau Inguinal*, is the inferior aperture of the inguinal canal.

INGUINOCUTA'NEUS. A name given by Professor Chaussier to the middle ramus of the anterior branch of the first lumbar nerve, because it sends its numerous filaments to the groin, serotum, and to the skin of the superior part of the thigh.

INHÆRENS, *Inherent*.

INHALATIO, (*in*, and *halare*, 'to breathe,') Absorption, Inhalation—i. Cutis, see Absorption.

INHALA'TION, *Inhala'tio*. The act of drawing air or vapour into the lungs—*Inhala'tio pulmona'lis*. Also, absorption.

INHA'LER. An apparatus for inhalation. *Mudge's Inhaler* is an apparatus for inhaling the steam of hot water, in affections of the air-passages. It consists of a pewter tankard provided with a lid, into which a flexible tube is inserted. Through this the vapour is inhaled.

INHE'RENT, *Inhæ'rens*, (*in*, and *hæ'rere*, 'to stick,') That which adheres, or which is joined or united to any thing.

INHERENT CAUT'ERY, (F.) *Cautère inhérent*, is the actual cautery, left in contact with a part until it is reduced to the state of a deep eschar.

INHIB'ITORY, *Inhibito'rius*, (*inhibeo*, (*in*, and *habeo*,) 'I hold in,') 'I restrain.') A term which has been applied to nerves, which, when methodically excited, seem to repress the action of muscles to which they are distributed.

INHUMA'TION, *Inhumat'io*, (*inhumare, inhumatum*, (*in*, and *humus*,) 'I put in the ground,') The sepulture of the dead. This belongs to the

subject of medical police. Also, digestion of substances in earth or dung. See *Arenatio*.

INIAD, see *Inial*.

IN'IAL, (from *inion*.) An epithet, proposed by Dr. Barclay for an aspect towards the plane of the ridge of the occiput. *Iniad* is employed by him adverbially to signify 'towards the inial aspect.'

INIENCEPH'ALUS, (F.) *Iniencephale*, (*inion*, and *εγκεφαλος*, 'encephalon.') A monster whose encephalon is in great part in the cranium, and in part out of it, behind, and a little beneath the cranium, which is open in its occipital portion.—(I. G. St. Hilaire.)

INIODYMUS, (*ινον*, and *δυειν*, 'to mingle,') *Diprosopus*.

IN'ION, (*ινον*.) Some of the Greek physicians give this name to the occiput, or the ridge of the occiput; others to the back part of the neck, and the muscles of the occiput. Blancard says it is the commencement of the spinal marrow.

IN'IOPS, *In'iope*, (*inion*, and *ωψ*, 'the eye, face,') A monster, double above the umbilicus, which has two ears very near each other, or only one median ear, and above it, at a greater or less distance, a median orbit and eye, more or less imperfect. The rudimental face appearing to be directly opposite the principal face suggested the name to I. G. St. Hilaire.

INI'TIS, *Inchymen'tis*, (*is*, gen. *ινος*, 'a fibre,') and *itis*.) Fibrous inflammation.

INITIUM, (*in*, and *ire*, 'to go,') 'a beginning,' *Arche*—i. *Asperæ arteriæ*, *Larynx*—i. *Extuberans coli*, *Cæcum*.

INIUM, (*ινιον*.) The nucha. Also, the muscles at the back of the neck. See *Inion*.

INJACULA'TIO, (*in*, and *jacular*, *jaculatus*, 'to shoot,') A term employed by Van Helmont to designate an acute pain of the stomach, with rigidity and immobility of the body.

INJEC'TED, *Injec'tus*, (*injicere*, *injectionem*, (*in*, and *jacere*,) 'to throw into,') The face and other parts are said to be *injected*, when the accumulation of blood in the capillary vessels gives them an evident red colour. A subject or part of a subject is also said to be *injected*, when its vessels have been filled, by the anatomist, with an appropriate composition.

INJEC'TION, *Injec'tio*, *Eis'bolē*. The act of introducing, by means of a syringe or other instrument, a liquid into a cavity of the body. The liquid injected is also called an *injection*. Anatomists use injections—*Injec'tio'nes anatomicæ*—for filling the cavities of vessels, in order that they may be rendered more apparent, and their dissection be facilitated. For this purpose, they employ syringes of different dimensions, and various materials. The most common injections are made of soot, wax, and turpentine, coloured with lamp-black, vermilion, &c. There are three kinds chiefly used by anatomists,—the *coarse*, the *fine*, and the *minute*. The finest injections for histological researches are made with ether as a menstruum.

INJECTION, Clyster—i. Matthews's, see *Tinctura cantharidis*.

INJECTIONES ANATOMICÆ, see *Injection*.

IN-KNEED, *Entogonyneon*.

INK, ([D.] *inkt*), *Atramentum*.

INKBERRY, *Prinos glaber*.

IN'NATE, (*in*, and *nascor*, *natus*, 'I am born,') (F.) *Inné*. *Inborn*.

INNATE DISEASES, *Mor'bi conna'ti seu congen'iti*, (F.) *Maladies innées*. Diseases with which the infant is born. They are not always hereditary, as hereditary diseases are not always innate.

INNERLEITH'EN, MIN'ERAL WA'TERS OF These springs, situate near the Tweed, and

supposed to be the scene of "St. Ronan's Well," contain chlorides of sodium and calcium, and carbonate of magnesia.

INNERVA'TION, *Innerva'tio*, *Enneuro'sis*, (*in*, and *nervus*, 'a nerve,') By this term is meant—the nervous influence, necessary for the maintenance of life and the functions of the various organs;—an influence of whose character and source we are ignorant. It seems to resemble the galvanic or electric agencies. See *Nerves*.

INNOMINA'TUS, *Anon'yms*, (*in*, and *nomen*, 'a name,') (F.) *Innominé*, *Anonyme*. Having no name. See *Anonymous*.

INNOMINATA ARTE'RIA, *Brachiocephalic ar'tery*, *Arte'ria anon'yma*, *Right Subcla'vian*, (F.) *Artère innominée*, *Artère brachio-céphalique* (Ch.), is the trunk common to the right primitive carotid and to the subclavian. It arises from the anterior part of the arch of the aorta, ascends obliquely to the right, along the trachea; and, after a course of about an inch in length, divides into two trunks, which go off at right angles. The one is *external*—the *right subclavian proper*; the other *superior*—the *primitive carotid* of the same side.

INNOMINATA CARTILAGO, *Cricoid*.

INNOMINATA CAV'ITAS. A cavity of the outer ear, between the *helix* and *anthelix*.

INNOMINATA FOSSA, see *Fossa*—i. *Linea*, see *Ilio-pectinea Linea*.

INNOMINATA MINO'RA OS'sa, *Lesser Ossa innominata*, *Ossic'ula innominata*. Some anatomists have given this name to the three cuneiform bones of the tarsus.

INNOMINATA OSSICULA, *Innominata minora ossa*—i. *Tunica Oculi*, *Sclerotic*.

INNOMINATÆ VE'NÆ, OF VIEU'SSENS. *Vieussens* has given this name to two or three veins, which arise on the anterior surface and right margin of the heart, and open into the auricle towards its right margin. The term *VENÆ INNOMINATÆ* OF MECKEL is given to the *brachiocephalic* veins, which are generally included in the description of the subclavian vein, and correspond to the *arteria innominata*, being formed by the union of the internal jugular vein and the subclavian properly so called, which correspond to the common carotid and subclavian arteries.

INNOMINATI seu ANONYMI NERVI. Some anatomists have thus called the nerves of the fifth pair.

INNOMINATUM FORA'MEN. A foramen, near the middle of the anterior surface of the pars petrosa of the temporal bone, leading backwards for the passage of the Vidian nerve, reflected from the 2d branch of the 5th to the portio dura of the 7th pair.

INNOMINATUM OS, *Os Coxen'dicis seu Cox'æ seu anon'ymsu seu pel'vis latera'le*, *Ache-bone*, *Aitch-bone*, (F.) *Os innominé*, *Os Cozal*, *Os anonyme*. A very large, flat bone, which forms the anterior and lateral paries of the pelvis. It is curved upon itself in two opposite directions. In the first periods of life, it is composed of three portions;—the *ilium*, *ischium*, and *pubis*, which join each other in the acetabulum. It is articulated *before* with its fellow, — *behind* with the sacrum; and *laterally* with the femur.

INNOMINÉ, *Innominatus*.

INNUTRITIO OSSIIUM, *Rachitis*.

INO, (*is*, *ινος*, 'a fibre,') in composition, *fibre*.

INOC'ULABLE, (*inoculare* (*in*, and *oc'ulus*, 'an eye'), 'to ingraft,') Capable of inoculation, or of being communicated by inoculation.

INOCULA'TION, *Inocula'tio*, *Insit'io*, *Insit'io variola'rum*. Any operation by which small-pox, for example, may be artificially communicated, by introducing the virus of the particular disease into the economy, by means of a puncture or scratch made in the skin. When the word *inocu-*

lation is used alone, it usually means that for the small-pox.—*Variolatio*. See Syphilization.

INOCULATION, COWPOX, Vaccination—i. Jennerian. Vaccination.

INOCULATOR, *In'sitor*. One who practises inoculation.

INODULAIRE, (from *nodus*, 'fibrous,') see Tissue, inodular.

INODULAR TISSUE, see Tissue, inodular.

INODULE, Tissue, inodular.

INOGENESIS, (*inu*, and *genesis*, 'generation,') Inosis.

INOHYLOMA, gen. *Inohylo'matis*, (*ino*, 'υλη,' 'matter,' and *oma*.) Tumour, fibrous.

INOHYMENITIS, (*ino*, 'υμνη,' 'a membrane,' and *itis*.) *Inflammat'io telæ fibro'se*. Inflammation of the fibrous tissue.

INOMA, (from *is*, gen. *inos*, 'a fibre,') Scirrhus, Tumour, fibrous.

INONCTION, Inunction.

INOPEX'IA, (*ino*, and *πηξις*, 'coagulation,')

Increased coagulability of fibrin.

INOPOLYPUS, (*ino*, and *polypus*,) see Polypus.

INORGANIC, *Inorgan'icus*, *Unor'ganized*, (F.) *Inorgan'ique*, (*in*, and *organon*, 'an organ,')

A term applied to bodies which have no organs;—such as minerals. At the present day, naturalists admit but two classes of bodies,—the *organized* and *inorganic*. Parts of the body which, like the epidermis, are devoid of blood vessels and nerves, have been called *anorgan'ic*.

INOSCLERO'MA, gen. *Inosclero'matis*, (*ino*, and *σκληρωμα*, 'induration,') Induration of the fibrous tissue.

INOSCLATIO, (*in*, and *osculum*, dim. of *os*, 'a mouth,') Anastomosis.

INOSIS, (*is*, gen. *inos*, 'a fibre,') Hyperinosis. Also, fibre formation, *Inogen'esis*.

IN'OSITE, *In'osit*, *Muscle sugar*, (from *is*, gen. *inos*, 'a fibre,') An unfermentable sugar, found only in muscle.

INOSTEATO'MA, (*ino*, and *steatoma*.) A tumour of a mixed fibrous and steatomatous character.

INQUIES, Inquietude.

INQUIETATIO, Inquietude.

INQUIETUDE, (*Inquietu'do*, *In'quies*, *Inquieta'tio*, *Jactita'tio*, (*in*, and *quies*, *quie'tis*, 'rest,') Agitation or trouble, caused by indisposition. Restlessness.

INQUINAMENTUM, (*inquinare*, 'to pollute,') Miasm.

INQUISITIO MEDICOLEGALIS, (*in*, and *quære*, *quæsitum*, 'to search,') see Medicolegal.

INSALIVA'TION, *Insaliva'tio*, (*in*, and *saliva*.) The mixture of the food with the saliva, and other secretions of the mouth.

INSALU'BRIOUS, *Insalu'bris*, (*in*, and *salu'bris*, 'healthy,') That which is unhealthy,—which injures the health.

INSANABILIS, (*in*, *sanare*, 'to cure,' and *habilis*, 'able,') Incurable.

INSANE, *Insanus*; (*in*, and *sanus*, 'sound,') *Aliena'tus*, *Cra'zy*, *Mad*, *Nonsane*, *Dement'ed*, *De-ranged*, *Bed'lamite*, (Sc.) *Daft*, *Wod*, (F.) *Aliéné*, *Fou*, *Insensé*. One affected with mental aberration, or of unsound mind.

INSANIA, Mania—i. *Cadiva*, *Epilepsy*—i. *Lupina*, *Lycanthropia*—i. *Puerperarum*, *Mania*, *puerperal*.

INSANITAS. Insanity.

INSAN'ITY, *Insania*, *Insan'itas*. *Mental aliena'tion*, *Abaliena'tio* seu *Alienatio Men'tis*, *Arrep'tio*, *Unsound Mind*, *Derangement*, *Deranged intellect*, *Cra'ziness*, *Madness*, *Phrensy*, *Frenzy*, (Sc.) *Daff'in*, *Daff'ing*, *Wod'nes*; *Aphros'yne*, *Ephro'nia*, *Ephros'yné*, *Para'llage*, *Parallax'is*, *Delir'ia*, *Vesa'nix*, *Delir'ium* (Crichton), *Insipien'tia*, (F.) *Folie*, *Egarement d'Esprit*, *Paraphré-nie*, *Cérébrie*. This term includes all the varieties

of unsound mind,—*Mania*, *Melancholia*, *Moral Insanity*, *Dementia*, and *Idiocy*. A slight degree of insanity is sometimes popularly called "*a kink in the head*;" in Scotland, "*a bee in the bonnet*."

INSANITY, HOMICIDAL, see Homicidal—i. *Incoherent*, *Dementia*—i. *Moral*, *Pathomania*—i. *Puerperal*, *Mania*, *puerperal*—i. *Religious*, *Sebastomania*—i. *Senile*, *Delirium senile*—i. *Suicidal*, see *Suicide*.

INSANUS, Insane.

INSCRIPTION, (*in*, and *scribere*, *scriptum*, 'to write,') see *Matriculation*.

INSCRIPTIONES TENDIN'EÆ MUSCULO'RUM, *Intersectio'nes* seu *Enervatio'nes tendin'eæ musculo'rum*, *Interne'dia ligamenta'tia seu ner'vae*. The tendinous portions which cross several muscles, and especially the straight muscles of the abdomen.

INSEC'TICIDE, (*insecum*, 'an insect, and *cædere*, 'to kill,') A substance that destroys insects.

INSEMINATION, (*in*, and *semen*, gen. *seminis*, 'seed,') see *Ejaculation*.

INSENESCENCIA, (*in*, and *senescere*, 'to grow old,') *Agerasia*.

INSENSE, Insane.

INSENSIBILITY, *Anæsth'esia*, *Insensibil'itas*, (*in*, *sensus*, 'feeling,' and *habilis*, 'able,') Loss or absence of sensibility. It is very common in cerebral affections, and may extend to every part, or be limited to one or more. Some organs are much more sensible than others. The bones, cartilages, ligaments, &c., are insensible in health, but acutely sensible in disease.

INSEN'SIBLE, *Sensibilita'tis ca'rens*. That which is devoid of sensibility. This word is applied, also, to phenomena which cease to be appreciable to the senses. Thus, we say, the pulse becomes *insensible*.

INSERTION, *Inser'tio*, *Symph'ysis*, *Eno'sis*, (*inserere*, *insertum* (*in*, and *serere*, 'to join or knit,') 'to ingraft,') (F.) *Attache*. The attachment of one part to another. Insertions occur chiefly in bones, cartilages, and fibrous organs; thus, we speak of the insertion of muscular fibres into a tendon or aponeurosis; the insertion of a tendon, aponeurosis, or ligament, into a cartilage or bone. The word *insertion* has likewise been used by pathologists for the act of inoculating or introducing a virus into the body.

INSEN'SIO, (*insidere*, *insessum* (*in*, and *sedere*, 'to sit in,') This term is, sometimes, applied to a vapour bath, the person being seated in a perforated chair, beneath which a vessel, filled with hot water, or the hot decoction of some plant, is placed. See *Bath*, *half*, and *Semicupium*.

INSESSUS, Bath, half.

INSICIUM, (*inseco*, 'to cut up,') *Isicium*.

INSIDEN'TIA, (*insideo* (*in*, and *sedere*, 'to sit in or upon,') *Epi'stasis*. Any things which swim on or in—the urine, for example. It is opposed to the *Hypo'stases* or *subside'ntia*.

INSIPIEN'TIA, (*in*, and *sapiens*, 'wise,') *Insanity*.

INSITIO, (*inserere*, *insitum*, 'to engraft,') *Grafting*, *Inoculation*—i. *Ciliorum*, *Blepharoplastice*—i. *Dentis*, *Transplantatio Dentis*—i. *Variolærum*, *Inoculation*.

INSOLATION, *Insola'tio* (*in*, and *sol*, 'the sun,') *Solica'tio*, *Aprica'tio*, *Helio'sis*, *Heli'osis*, *Heliono'sis*, *Siri'asis*. Exposure to the sun. Exposure of a patient to the rays of the sun is, sometimes, had recourse to, with the view of rousing the vital forces when languishing, or of producing irritation of the skin. *Insolation* is occasionally used in the same sense as *coup de soleil*.

In *Pharmacy*, *insolation* means the drying of chemical and pharmaceutical substances.

INSOLATIONE DE PRIMIVERA. ('spring insulation.') Pellagra.

INSOMNIA. *Insomnitas*, Sahara, Zuara, Per-rigil'ium, Pernoc'tio, App'nia, Agryp'nia, Ahyp'nia, Anyp'nia, Typhoma'nia, Sleep'lessness, Vig'ilance, (in, negative, and *somnus*, 'sleep,' absence of sleep.) This may exist alone, and constitute a true disease; or it may be connected with another affection. It is an unequivocal sign of suffering in some organ; even when the patient experiences no pain.

INSOMNIUM. (in, and *somnus*, 'sleep,') Som-nium.

INSPECTIO MEDICOLEGALIS. (*inspicere*, inspectum, 'to look into,') see *Medicolegal*.

INSPIRATEUR. Inspiratory.

INSPIRATION. *Inspira'tio*, *Empneumato'sis*, *Eisp'mo*, *Adspira'tio*, *Aspira'tio*. (in, and *spirare*, *spiratum*, 'to breathe,') The action by which the air penetrates into the interior of the lungs. A movement opposed to that of expiration. As regards the average quantity of air received into the lungs at each inspiration, there is much discrepancy amongst observers. The following table sufficiently exhibits this:—

	Cubic inches at each inspiration.
Reil	42 to 100
Menzies,	
Sanvages,	
Hales,	
Haller,	
Ellis,	40
Spengel,	
Simmering,	
Thomson,	
Berstock,	
Jurin	25 to 38
Fontana	35
Richerand	30 to 40
Dalton (John)	30
Jedreys	26
Herbst	24 to 30
Herholdt	20 to 29
Jurine and Coathupe	20
Allen and Pepys	16½
J. Borrelli	15 to 40
Goodwyn	14
Sir H. Davy	13 to 17
Abernethy and Mojon	12
Keutsch	6 to 12

From 20 to 25 cubic inches may be regarded as a fair average.

INSPIRATION, CROWING, see *Asthma thymicum*—i. Muscles of, see *Inspiratory*.

INSPIRATION OF VENOUS BLOOD. By this is meant the aspiration of blood towards the heart, occasioned by the approach to a vacuum produced by the dilatation of the thorax during inspiration.

INSPIRATORY. *Inspiratio'ni inser'vians*, (F.) *Inspirator*. A name given to muscles, which, by their contraction, augment the size of the chest, and thus produce inspiration. The diaphragm and intercostal muscles are the chief agents of inspiration. In cases where deep inspirations are necessary, the action of these muscles is aided by the contraction of the pectoralis major and pectoralis minor, subclavius, serratus major anticus, scaleni, serratus posticus superior, &c. Most of these muscles become inspiratory, by taking their fixed point at the part which they ordinarily move, acting inversely and elevating the ribs.

INSPISSANT. *Inspissans*, (in, and *spissare*, *spissatum*, 'to thicken,') Any remedial agent that renders the blood thicker, directly, or indirectly. The opposite of diluent.

INSPISSATIO. *Pyeno'sis*. The act of rendering thick; as in the formation of an extract—*Succus Inspissatus*.

INSTEP. *Col'lum pe'dis*. Neck of the foot. (F.) *Conte-pied*, *Cont-de-pied*. The projection at the upper part of the foot, near its articulation with the leg—the *tarsus*.

INSTILLATION. *Enstilla'io*, *Instilla'tio*, (in,

'in,' and *stilla*, 'a drop,') The act of pouring a liquid drop by drop.

INSTINCT. (L.) *Instine'tus*, (*instinguere*, *instinctum*, 'to impel,') 'inwardly moved.' *Brutia*, *Hor'me*. The action of the living principle, whenever manifestly directing its operations to the health, preservation, or reproduction of a living frame or any part of such frame—*Moll'men natu'ræ salutarium*. The law of instinct is, consequently, the law of the living principle, and instinctive actions are the actions of the living principle. Instinct is natural. Reason is acquired.

INSTINCTIVE. *Instine'tus*, (F.) *Instinctif*. Relating to or caused by instinct; as *instinctive* actions. See *Emotional*, and *Instinct*. Those instinctive actions of animals which are owing to impressions made on the sensory ganglia, exciting respondent motor influences that are propagated to the various muscles of the body, are termed *consensual*. Consensual movements which accompany voluntary efforts without our consciousness are termed *associated*, (F.) *Associés*.

INSTITUTES OF MEDICINE. (*institutum*, 'a principle,' from *instituere*, (in, and *statuere*,) 'to establish,') see *Theory of medicine*.

INSTITUTUM ORTHOPÆDICUM. Ortho-pædic institution.

INSTRUMENT. *Instrumen'tum*, (from *instruere*, (in, and *struere*,) 'to put together,') *Organum*. A tool, an agent.

INSTRUMENTA. Pudibilia.

INSTRUMENTUM. Instrument—i. *Chirurgicum*. A surgical tool or instrument.

INSTRUMENTUM DIGESTIVUM. The digestive apparatus.

INSTRUMENTUM INSTRUMENTORUM. Manus.

INSUFFICIENCY. (F.) *Insuffisance*; (in, and *sufficere*, 'to suffice,') Inadequateness to any end or purpose,—as *Insufficiency of the valves of the heart*, (F.) *Insuffisance des valvules du Cœur*;—a condition in which they are not adapted, as in health, to properly close the apertures.

INSUFFISANCE. Insufficiency—i. *des Valvules du Cœur*, Insufficiency of the valves of the heart.

INSUFFLATION. *Insufla'tio*, (in, *inf*, and *flare*, 'to blow,') The act of blowing a gas or vapour into some cavity of the body; as when tobacco smoke is injected into the rectum; or when air is blown into the mouths of newborn children to excite the respiratory function.

INSULA. ('an island,') *Insula cerebri*.

INSULA CEREBRI. *Island* or *In'sula* of Reil. The intermediate lobe of the brain, *Lobus intermedius cerebri*. A remarkable group of convolutions—*Gy'ri oper'ti*—within the fissure of Sylvius. It is called, by Cruveilhier, *Lob'ule of the fissure of Syl'vius*, *Lobule of the corpus striatum*. The 'island' of Reil, with the substantia perforata, forms the base of the corpus striatum.

INSULA SANGUINIS. see *Blood*.

INSULÆ. *Maculæ albæ*—i. *Pulmonales*, see *Pulmo*.

INSULTUS. (*insilio*, *insultum*, 'to attack,') Attack. Paroxysm.

INSURANCE OF LIFE. see *Life Insurance*.

INTEGRITAS. (*integer*, 'whole,') *Sanitas*, *Virginity*.

INTEGUMENT. *Integumen'tum*, *Tegumen'tum*, *Teg'men*, *Teg'men*, *Involu'crum* seu *Velamen'tum corporis commu'ne*, *Vela'men* seu *Velamentum nat'urum*, (in, and *tegere*, *tectum*, 'to cover,') (F.) *Tégument*. Any thing which serves to cover, to envelop. The *skin*, including the cuticle, rete mucosum, and cutis vera, is the common integument or tegument of the body.

INTEGUMENTA FÆTUS. Membranes of the fetus.

INTEGUMENTUM. Integument—i. *Com-mune*, *Integumentum*.

INTELLECT, *Intellectus*, *Nous*, *Mens*, *Gno'mē*, *Noos*, *Nūs*, (intelligere, intellectum, (inter, and legere, 'to choose,') 'to understand,' 'conceive,' 'know,') (F.) *Entendement*, *Intelligence*. The aggregate of the intellectual faculties—perception, formation of ideas, memory, and judgment. Their exercise constitutes *intellection*, *Noē'sis*, *Syn'esis*.

INTELLECT, DERANGED, Insanity.

INTELECTION, see *Intellect*.

INTELLIGENCE, *Intellect*.

INTEMPERANCE, *Intemperant'ia*, *Immoderant'ia*, *Immoderatio*, *Acras'ia*, *Acolas'ia*, *Plesmonē*, *Amet'ria*, *Aples'tia*, (in, and temperare, 'to temper,') Immoderate use of food and drink, especially the latter;—a fruitful source of disease.

INTEMPERIES, *Dyscras'ia*. Derangement in the constitution of the atmosphere and of the seasons; bad constitution; derangement or disorder in the humours of the body.

INTENSIVUS, (in, and tendere, *tensum*, 'to stretch,') *Entaticus*.

INTENTION, *Intentio*, (in, and tendere, *tensum* or *tentum*, 'to stretch,') *Propositum*. The object which one proposes. In surgery, a wound is said to heal by the first intention, *Reu'nio per primam intentio'nem*, when cicatrization occurs without suppuration; union by the second intention, *Reu'nio per secundam intentio'nem*, being that which does not occur until the surfaces have suppurated. To obtain union by the first intention, the edges of a recent wound must be brought in apposition and kept together by means of adhesive plasters and a proper bandage. Delppech has substituted for those expressions, *Réunion primitive*, and *Réunion secondaire*.

INTER, in composition, between.

INTERANEA, (*intra*, 'within,') *Entrails*.

INTERARTICULAR, *Interarticula'ris*; (F.) *Interarticulaire*, (inter, and articulus, 'a joint,') Parts situate between the articulations are so called; as *interarticular cartilages*, *interarticular ligaments*, &c.

INTERCADENCE, *Intercaden'tia*, *Interciden'tia*, (in, and cadere, 'to fall,') Disorder of the arterial pulsations, so that, every now and then, a supernumerary pulsation occurs. The pulse, in such case, is said to be *intercurrent*.

INTERCALARY, *Intercala'ris*, *Intercidens*, *Embol'imos*, *Interpolatus*, *Provocato'rius*, (F.) *Intercalare*, (*intercalare*, 'to insert,') The days which occur between those that are critical. The term has, also, been applied to the days of apyrexia in intermittent fevers.

INTERCELLULAR, *Intercellula'ris*, (F.) *Intercellulaire*; (inter, and cellula, 'a small cell, a cellule,') That which is between cells or cellules.

INTERCELLULAR FLUID, *Liquor Sanguinis*.

INTERCELLULAR PASSAGES. A term given by Mr. Rainey to irregular passages through the substance of the lung, which form the terminations of the bronchial tubes, are clustered with aircells, and not lined by mucous membrane. [?]

INTERCELLULAR SUBSTANCE, see *Cytoblastema*.

INTERCEPTIO, (from *intercipere*, *interceptum*, inter, and capere, 'to intercept,') A bandage by the aid of which the ancients proposed to arrest the progress of the material cause of gout and rheumatism; and which consisted in covering the affected limbs with carded wool; surrounding them, afterwards, with broad bandages, applied from the fingers to the axilla, or from the toes to the groin.

INTERCEPTIO INTESTINORUM, *Ileus*.

INTERCERVICAUX, (inter, and cervix, gen. *cervicis*, 'neck,') *Interspinales colli*.

INTERCIDENS, (inter, and cadere, 'to fall,') *Intercalary*.

INTERCIDENTIA, *Intercadence*.

INTERCILIUM, (inter, and cilium, 'eyelid,') *Mezophryon*.

INTERCLAVICULAR, *Interclavicula'ris*, (F.) *Interclaviculaire*, (inter, and clavicula, 'a clavicle,') That which is placed between the clavicles.

INTERCLAVICULAR LIGAMENT is a fibrous bundle, placed transversely above the extremity of the sternum, between the heads of the two clavicles. This ligament is flat. Its fibres, which are always longer above than below, are separated by small apertures, which are traversed by vessels. It prevents the separation of the two clavicles in the forced depression of the shoulder.

INTERCOSTAL, *Intercosta'lis*, (inter, and costa, 'a rib,') *Mesopleu'rus*, *Mesopleu'rius*. That which is situate between the ribs.

INTERCOSTAL ARTERIES vary in number. There is constantly, however, a *superior*, *Arte'ria Intercosta'lis superior* seu *Intercosta'lis subclavia*, which is given off from the posterior part of the subclavian, and which sends branches into the first two or three intercostal spaces; and, generally, eight or nine inferior or aortic *intercostals*. These arise from the lateral and posterior parts of the pectoral aorta, and ascend obliquely in front of the vertebral column, to gain the intercostal spaces, where they divide into a *dorsal* branch and an *intercostal*, properly so called.

INTERCOSTAL MUSCLES are distinguished into *internal*, — *inter-plévrocostaux* of Dumas, — and *external*. The former are inserted into the inner lip, the latter into the outer lip of the edge of the ribs. The fibres of the *external intercostals* are directed obliquely downwards and forwards; and those of the *internal* downwards and backwards. Both are inspiratory or expiratory muscles, according as they take their origin on the upper or lower rib. Some small, fleshy fibres, seen occasionally at the inner surface of the thorax, descending obliquely from one rib to another, have been called *Infracostales*, (F.) *Soucostaux*.

INTERCOSTAL NERVE, *Trisplanchnic nerve*.

INTERCOSTAL NERVES, *Branches souscostales* (Ch.), *Costal* or *Dorsal nerves*, proceed from the anterior branches of the dorsal nerves. They are twelve in number, and are distributed especially to the muscles of the parietes of the chest and abdomen.

INTERCOSTAL SPACE, *Interval'lum* seu *Interstit'ium intercosta'le*, *Mesopleu'rum*, *Mesopleu'rium*, is the interval which separates one rib from that immediately above or below it.

INTERCOSTAL VEINS are distinguished like the arteries. The *right superior intercostal vein* is often wanting. When it exists, it opens into the back part of the subclavian. The same vein of the left side is very large. It communicates with the *demi-azygos*, receives the left bronchial vein, and opens into the corresponding subclavian. The *right inferior intercostal veins* open into the *vena azygos*; and those of the left into the *demiazygos*.

INTERCOSTO HUMERAL NERVES. So called from their origin and distribution. They are the cutaneous branches of the second and third intercostal nerves.

INTERCURRENT, *Intercur'rens*, (inter, and currere, 'to run,') Diseases are so called which supervene at different seasons of the year, or which cannot be considered as belonging to any particular season. A disease is likewise so termed which occurs in the course of another disease, as *Intercurrent Pneumonia*.

INTERCUTANEOUS, *Subcutaneous*.

INTERDENTIUM, (inter, and dens, gen. *dentis*, 'a tooth,') The interval between teeth of the same order.

INTERDIGITAIRE, *Interdigital*.

INTERDIGITAL, *Interdigita'lis*, (F.) *Interdigitaire*; (inter, and dig'iti, 'a finger,') That which relates to the spaces between the fingers. *Interdigital space*, (F.) *Espace interdigitaire*, is

used, also, for the commissure between the fingers.

INTERDIGIT'IUM. A corn or wart, which grows between the fingers and toes, especially between the latter.

INTERÉPINEUX, Interspinal—i. *Cervicæux,* Interspinalis colli.

INTERFEMINEUM, (*inter*, 'between,' and *femen*, gen. *feminis*, the inner part of the thigh,) Perineum. Vulva.

INTERFUNGIONARIUM, (*inter*, and *funis*, 'a boundary,') Septum narium.

INTERFEMINEUM, Perineum, Vulva.

INTERFORAMINEUM, Perineum.

INTERGANGLION'IC, *Interganglion'icus*, (*inter*, and *ganglion*, 'a knot,') An epithet for nervous cords, placed between ganglia, which they connect together.

INTERITUS, (*intereo*, *interitum*, 'to perish,') Death.

INTERLOBAR, *Interloba'ris*, (F.) *Interlobaire*, (*inter*, and *lobus*, 'a lobe,') That which is between the lobes,—as of the lungs.

INTERLOBULAR, *Interlobula'ris*, (F.) *Interlobulaire*, *Interlobulinaire*, (*inter*, and *lobulus*, 'a small lobe,') That which is between lobules,—as of the lungs.

INTERLOBULAR FIS'SURES. The intervals between the lobules of the liver. The narrower spaces are called *Fissu'ræ interlobula'res*, the wider *Spa'tia interlobula'ria*. See Liver.

INTERLOBULAR PLEXUS OF THE BILIARY DUCTS. See Liver.

INTERLOBULAR SPACES. The angular interstices formed in the liver by the apposition of several lobules.

INTERLOBULAR TIS'SUE, *Tex'tum interlobula'rè*, *Ligamen'ta interlobula'ria pulmo'nium*. The cellular tissue between the pulmonary lobules.

INTERLOBULAR VEINS, see Liver.

INTERLOBULINAIRE, Interlobular.

INTERMAXILLARY, *Intermaxilla'ris*, (F.) *Intermaxillaire*, (*inter*, and *maxilla*, 'a jaw,') That which is situate between the maxillary bones.

INTERMAXILLARY BONE, *Inci'sive*, *Pal'atine*, or *Labial bone*, is a bony portion, wedged in between the two superior maxillary bones, which supports the upper incisors. This bone is found in the mammalia; and, also, in the human fœtus.

INTERMÈDE, (*inter*, and *medius*, 'middle,') Excipient.

INTERMEDIA LIGAMENTALIA seu **NERVEA**, *Inscriptiones tendinæ musculorum*.

INTERMÉDIAIRE DE WRISBERG, Portio Wrisbergii.

INTERMEDIATE VASCULAR SYSTEM, Capillary system.

INTERMENT, PREMATURE, (*in*, and *terra*, 'earth,') Zoothapsis.

INTERMIS'SION, *Intermis'sio*, *Dialeip'sis*, *Dialip'sis*, *Tempus intercala'rè*, *Interval'lum*, *Interval*, (*inter*, and *mittere*, *missum*, 'to put or send,') (F.) *Intermittence*. The interval which occurs between two paroxysms of an intermittent or other disease—during which the patient is almost in his natural state. There is said to be *intermission* of the pulse, when, in a given number of pulsations, one or more may be wanting.

INTERMITTENCE, *Intermission*.

INTERMITTENT, *Intermit'tens*. That which has intermissions.

INTERMITTENT FE'VER, *Fe'bris intermit'tens*, *Dialeip'ya*, *Intermittens*, *An'teus*, *Pyretotypo'sis*, *A'gue*, *Aque* and *Fe'ber*, *Fever* and *Aque*, *Palu'dal* or *Period'ic Fever*, (Prov.) *Agah*, *Coldchill*, *Shaking*, *Shakes*, (Old Eng.) *Ak'sis*, *Auggere*, *Axes*; (F.) *Fèvre intermittente*, *F. d'Accès*, *F. des marais*, *F. Périodique*, is a fever consisting of paroxysms, with a complete state of apyrexia in the

intervals. The chief types are *Quotid'ian*, *Tertian*, and *Quart'an*. The symptoms of intermittents are those of a decided and completely marked *cold stage*, or *Stage of Concentration*, (F.) *Stade de froid*, attended with paleness; collapse; impaired sensibility; and coldness, more or less diffused, followed by general rigors. After this occurs the *hot stage*, or *stage of expansion*, (F.) *Stade de la chaleur*, the heat returning partially and irregularly, and at length becoming universal, and much above the standard of health. The pulse is now hard and strong; tongue white; urine, high-coloured; thirst, considerable. At length, the *sweating stage*, or *stage of termination*, (F.) *Stade de sueur*, makes its appearance; the moisture usually beginning on the forehead, face, and neck, and soon extending universally; the heat abating; thirst ceasing; the urine throwing down a sediment, and the functions being gradually restored to their wonted state. The tertian type is the most common, and the quartan the most severe. The quotidian more readily changes into a remittent and continued fever. The quartan has, generally, the longest cold stage, the tertian the longest hot. The chief exciting cause is marsh miasmata, the malaria of the Italians. Ague, also, occurs in districts where there are no marshy emanations. Such districts are, generally, of a volcanic nature;—further we know not. When the disease has once attacked an individual, it is apt to recur. The *prognosis* is, in general, favourable, as far as regards life; but long protracted intermittents are apt to induce visceral obstructions, and engorgements, which may end in dropsy. In some countries, the disease is of a very pernicious character. The indications of treatment are, 1. To abate the violence of the paroxysm; and, 2. To prevent its return. The first indication demands the adoption of the general principles and remedies required in ordinary fever. The *second* is the most important. The period of apyrexia is that for action. The means for fulfilling this indication are:—the use of emetics, purgatives, antiperiodics, as cinchona, quinia, &c., arsenic, and forcible impressions made on the mind of the patient.

INTERMUS'ULAR, *Intermuscula'ris*, (F.) *Intermusculaire*, (*inter*, and *musculus*, 'a muscle,') That which is placed between muscles.

INTERMUSCULAR APONEURO'SES are aponeurotic laminae or septa, situate between muscles, to which they often give attachment.

INTERNAL, *In'ter*, *Inter'nus*. That which is placed on the inside. This epithet is given to parts that look towards an imaginary central plane, which divides the body into two equal and symmetrical portions, as well as to those which correspond with a cavity. Thus, we say,—the *internal* or *inner surface* of the arm or thigh, skull, &c.

INTERNAL DISEASES, *Morbi inter'ni*, are those which occupy the inner parts of the body. Their investigation belongs to the physician; *external* diseases falling under the management of the surgeon.

INTERNAT, (F.) The post or office of an interne.

INTERNE, see Houseurgeon.

INTERNODIA DIGITORUM MANUS, (*inter*, and *nodus*, 'a knot,') Phalanges of the fingers—i. *Digitorum pedis*, Phalanges of the toes.

INTERNO'DIUM. The part of the fingers between the joints. A phalanx.

INTERNUN'CIAL, *Internuncia'lis*, (*inter*, and *nuncia*, 'a messenger,') Relating or belonging to a messenger between parties. A term applied by Mr. Hunter to the functions of the nervous system.

INTERNUNTII DIES, Critical days.

INTERNUS AURIS, ('internal of the ear,') Tensor tympani—i. Mallei, Tensor tympani.

INTEROSSEOUS, *Interosseus*, (F.) *Interosseus*, (*inter*, and *os*, 'a bone,') That which is situated between the bones.

INTEROSSEOUS ARTERIES OF THE FOREARM AND HAND. Of these there are several. The common *interosseous artery* arises from the posterior part of the ulnar, a little below the bicipital tuberosity of the radius; passes backwards and divides into two branches: the one called *anterior interosseous* descends vertically, anterior to the interosseous ligament; the other, called *posterior interosseous*, passes above the ligament, appears at its posterior part and divides into two great branches, — the *posterior recurrent radial* and the *posterior interosseous*, properly so called. In the hand, — 1. The *dorsal metacarpal interosseous arteries* are given off by the *dorsalis carpi*, a division of the radial artery. 2. The *palmar interosseous arteries*, which arise from the convexity of the deep palmar arch, and give off the *middle interosseous arteries*. 3. The *dorsal interosseous of the index*, proceeding directly from the radial artery. In the foot, are distinguished: — 1. The *dorsal interosseous arteries*, three in number, which arise from the artery of the metatarsus, a branch of the *dorsalis tarsi*. 2. The *plantar interosseous arteries*, which are, also, three in number, and arise from the plantar arch.

INTEROSSEOUS LIGAMENTS. Ligaments seated between certain bones, which they unite; such are the ligaments between the radius and ulna, and between the tibia and fibula.

INTEROSSEOUS MUSCLES. These occupy the spaces between the bones of the metacarpus and metatarsus; and, consequently, belong, — some to the hand, others to the foot.

INTEROSSEOUS NERVE is a branch, given off by the median nerve, which descends before the interosseous ligament, accompanying the artery of the same name.

INTEROSSEOUS VEINS have the same arrangement as the arteries.

INTEROSSEUS MANUS, (F.) *Muscle interosseus de la main*, — *Métacarpo-phalangien-latéral suspalmaire et métacarpo-phalangien latéral*, (Ch.) These muscles are seven in number; two for each of the three middle fingers, and one for the little finger. Four are situated on the back of the hand, and three only in the palm. They are inserted into the metacarpal bones, and send a tendon to the tendon of the extensor communis. According to their office, they are, to each finger, an *adductor* and an *abductor*. The *index* has a dorsal abductor and a palmar one. The *middle finger* has two dorsal muscles for adductor and abductor; the *ring finger* has a dorsal adductor and a palmar abductor; and the *little finger* has only one interosseous abductor, which is palmar. These muscles produce abduction and adduction of the fingers, which they can also extend, owing to their connexion with the extensor tendons.

INTEROSSEUS PEDIS, *Métatarso-phalangien-latéral*, (Ch.) The number, arrangement, shape, and uses of these are the same as in the case of the preceding muscles. Four are *dorsal*, and three *plantar*: six belong to the three middle toes and one to the little toe. The great toe is devoid of them. As in the hand, they are distinguished, in each toe, into abductor and adductor.

INTERNUS QUARTUS, Prior annularis—i. Secundus, Prior medii digiti.

INTEROSSEUX DE LA MAIN, *Interosseus manus*.

INTERPARIETAL BONE, *Os Interparietale*. A bone found in the skulls of the children of the Peruvian races. It lies in the situation of the upper angle of the occipital bone, where the parietal bones separate from each other; and is the

analogue of the interparietal bone of ruminants and carnivora.

INTERPEDUNCULAR, *Interpeduncularis*; (*inter*, and *pedunculi*, 'peduncles,') A term, applied to that which is placed between the peduncles of the encephalon, — as *Interpeduncular space*; a lozenges shaped interval, behind the diverging peduncles or crura of the cerebrum, in which are found the posterior perforated space, corpora albicantia, tuber cinereum, infundibulum, and pituitary body.

INTERPELLATUS (*Morbus*,) (*interpello*, [*inter*, and *pello*,] 'I interrupt,') A term, by which Paracelsus designated those diseases whose progress is unequal, and paroxysms irregular.

INTERPLÉVRO-COSTAUX, ('interpleurocostals,') Intercoastal muscles.

INTERPOLATUS, (*interpolaris*, *interpolarum*, 'to give a new form to,') Intercalary.

INTERRUPTIO MENSTRUATIONIS, Amenorrhoea.

INTERSCAPULAR, *Interscapularis*, (F.) *Interscapulaire*, (*inter*, and *scapula*,) A term, which is between the shoulders; as the *interscapular region*. The *Interscapular cavities*, *Cavitates scapulae*, are the depressions between the scapulae and the spinous processes of the vertebrae.

INTERSCAPULIUM. The spine of the scapula.

INTERSECTIO, (*inter*, and *seco*, *sectum*, 'to cut,') Decussation, Intersection.

INTERSECTION, *Intersectio*, *Diaopê*. The point where two lines meet and cut each other. The name *aponeurotic intersection* is given to fibrous bands, which certain muscles present in their length, and by which they seem interrupted. Aponeurotic intersections are found in the recti muscles of the abdomen; in the semimembranosus, complexus, sterno-thyroideus, &c.

INTERSECTIONS TENDINEÆ MUSCULORUM, *Inscriptiones tendineæ musculorum*.

INTERSEPIMENTUM THORACIS, (*intersepicio*, 'to fence about,') Mediastinum.

INTERSEPTA HORIZONTALIA PACCHIONI, Tentorium.

INTERSEPTUM, (*inter*, and *septum*, 'a partition,') The uvula. The septum narium. The diaphragm.

INTERSEPTUM NARIUM, Septum narium—i. Virginalis, Hymen.

INTERSPINAL, *Interspinalis*, (*inter*, and *spina*, 'the spine,') (F.) *Intéripéneux*. That which is seated between the spinous processes.

INTERSPINAL LIGAMENT, *Membra'na interspinalis*, (F.) *Ligament intéripéneux*. These occupy the intervals between the spinous processes in the back and loins. In the neck, they are replaced by the muscles of the same name. They prevent the too great flexion of the spine, and keep the spinous processes *in situ*. The outermost thickened portions of these are, sometimes, termed *Ligamenta a'picipum*.

INTERSPINALES COLLI, *Spinales Colli minores*, *Spinali*, (F.) *Intéripéneux-cervicaux*, *Intercervicaux*, (Ch.) These are twelve in number, and occupy, in two parallel rows, the intervals between the spinous processes of the cervical vertebrae, from that of the atlas and vertebra dentata, to that between the last cervical and first dorsal vertebra. They are flat, thin, and quadrilateral. These muscles contribute to the extension of the neck and to throwing the head backwards.

INTERSPINALES DORSI ET LUMBORUM; portions of the *Transversaire-épineux* of Boyer, and the *sacro-spinal* of Chaussier. These muscles are of two kinds. The one (*Muscle grand épineux du dos*, of Winslow) representing fleshy bundles of different lengths, applied upon the lateral surfaces

of the spinous processes, from the third dorsal vertebra to the second lumbar. The other (*Muscle petit épineux du dos*, Winslow) covered by the preceding. They are situate on each side of the interspinal ligament, in the form of small, short, flat bundles, which pass from one spinous process to the second, third or fourth above it. These muscles aid in extending the vertebral column; and incline it a little to one side, when they act on one side only.

INTERSTICE, *Interstit'ium*, *Aræo'ma*, (*inter*, and *sistere*, 'to place.') *Diaste'ma*, *Interval'lum*. Anatomists have given this name to the intervals between organs. The *iliac crest*, *crista ili*, for example, has two *lips* and an interstice between them, which affords attachment to the lesser oblique muscle of the abdomen.

INTERSTICE, Pore.

INTERSTIT'IAL, *Interstitia'lis*, (F.) *Interstitiel*. Applied to that which occurs in the interstices of an organ,—as 'interstitial absorption,' 'interstitial pregnancy,' &c.

INTERSTITIUM, Interstice—i. Ciliare, Ciliary ligament—i. Intercostale, Intercostal space—i. Jugulare, Throat.

INTERTINCTUS, (*inter*, and *tingere*, *tinctum*, 'to dye.') Discrete.

INTERTRACHÉLIENS, (*inter*, and *τραχηλος*, 'the neck.') Intertransversales colli.

INTERTRANSVERSAIRE, *Intertransversalis*—i. *des Lombes*, *Intertransversales lumborum*.

INTERTRANSVERSA'LIS, *Intertransversa'rius*, (F.) *Intertransversaire*. That which is placed between the transverse processes of the vertebrae.

INTERTRANSVERSALES COL'LI, (F.) *Intertransversaires cervicaux*, *Intertrachéliens*, (Ch.) These are small, muscular bundles; quadrilateral; thin and flat; situate, in pairs, in the intervals between the transverse processes of the neck, except between the first and second, where there is only one. They are distinguished into *anterior* and *posterior*. The former are six in number; the latter five. These muscles bring the transverse processes of the neck nearer each other, and contribute to the lateral flexion of the neck.

INTERTRANSVERSALES LUMBO'RUM, *Musculi intertransversari lumborum*, (F.) *Intertransversaires des lombes*. These are almost entirely fleshy, and ten in number; five on each side. They are similar to the preceding in general arrangement; except that they are more marked, and not placed in two rows. Each intertransverse space contains only one. They are quadrilateral, and flat. The first occupies the space between the transverse processes of the first lumbar and the last dorsal vertebrae; and the last is between that of the fourth and fifth lumbar vertebrae. These muscles incline the lumbar region laterally; and straighten it when inclined to one side.

INTERTRIGO, gen. *Intertrig'inis*, (*inter*, and *terere*, 'to rub.') Chafing—i. *Podicis*, Chafing.

INTERVAL, (*inter*, and *vallum*, 'a rampart.') Intermission—i. *Lucid*, see *Lucid*.

INTERVALLUM, (same etymon.) Intermission, Intervall—i. *Intercostale*, Intercostal space.

INTERVENIUM, (*inter*, and *vena*, 'a vein.') Mesoplebium.

INTERVERTEBRAL, *Intervertebra'lis*, (*inter*, and *vertebra*.) That which is situate between the vertebrae.

INTERVERTEBRAL CARTILAGES, *Fibrocartilages* or *discs*, *Ligamenta intervertebra'lia*, *Fibrocartilagin'ines intervertebra'les*. These organs are of a fibrocartilaginous nature; sections of a cylinder; flexible; whitish; resisting; and situate between the bodies of the vertebrae,—from the space between the second and third as far as that

between the last vertebra and sacrum. Their form is accommodated to that of the vertebrae with which they are in connexion; so that, in the neck and loins, they are oval, whilst in the dorsal region they are nearly circular.

INTERVERTEBRAL DISCS, Intervertebral cartilages—i. *Fibrocartilages*, Intervertebral cartilages.

INTESTABILIS, (*in*, and *testis*, 'a testicle.') Castratus.

INTESTATUS, Castratus.

INTESTIN, Intestine.

INTEST'INAL, *Intestina'lis*, *Enter'icus*, *Enter'ic*, (F.) *Entérique*, (*intus*, 'within.') That which belongs to the intestines,—as *intestinal canal*, &c. Bordeu uses the term *Pouls intestinal* for a pulse, which he conceived to announce an approaching crisis by the intestines.

INTESTINAL JUICE, *Succus entericus*.

INTESTINAL TUBE or **TRACT**, *Canalis seu Tractus intestino'rum*. The canal formed by the intestines from the pyloric orifice of the stomach to the anus.

INTES'TINE, *Intesti'num*, *En'teron*, *Chor'da*, *Gut*, *Ned'ya* (pl.), *Pan'tices* (pl.), *Boel'li* (pl.), *Bowel*, (F.) *Intestin*, *Bogau*. A musculomembranous canal, variously convoluted, which extends from the stomach to the anus, and is situate in the abdominal cavity; the greater part of which it fills. In man, its length is six or eight times that of the body. It is divided into two principal portions, called *small intestine* and *large intestine*. The former, *Intesti'num ten'ue*, *I. grac'ile*, *Aræ'a*, (F.) *Intestin grêle*, constituting nearly four-fifths of the whole length, begins at the stomach and terminates in the right iliac region. It is divided into *duodenum*, *jejunum* and *ileum*. Some anatomists give the name *small intestine* to the last two only; which are kept in place by the mesentery, and form a large *paquet*, occupying the umbilical and hypogastric regions, a part of the flanks, of the iliac regions, and of the cavity of the pelvis. It is composed of, 1. A serous membrane, which is peritoneal. 2. Of a muscular coat, whose fibres are very pale, and are placed, in part, longitudinally; but the greater part transversely. 3. Of a whitish, mucous membrane; villous, and forming folds or valves—*valvule conniventes*—at its inner surface, and furnished with a number of mucous follicles, called *glands* of Lieberkühn, (*Crypts* or *Follicles* of Lieberkühn,) and in the duodenum with the *glands* of Brunner. The arteries of the small intestine proceed from the superior mesenteric; its veins open into the vena porta. Its nerves proceed from the superior mesenteric plexus. The large intestine, *Intesti'num cras'sum*, *Megalocœ'lia*, (F.) *Gros intestin*, forms a sequence to the small. It is much shorter, and is solidly attached in the regions of the abdomen which it occupies. It begins in the right iliac region; ascends along the right flank, till beneath the liver, when it crosses the upper part of the abdomen, descends into the left iliac fossa, and plunges into the pelvic cavity, to gain the anus. The great intestine is usually divided into three portions,—the *cæcum*, *colon* and *rectum*. It receives its arteries from the superior and inferior mesenteries. Its veins open into the vena porta. Its nerves are furnished by the mesenteric plexuses. Its lymphatic vessels, which are much less numerous than those of the small intestine, pass into the ganglions or glands seated between the different reflections of the peritoneum, which fix it to the abdominal parietes. The use of the intestines is,—in the upper part, to effect the chylification of the food and the absorption of the chyle;—in the lower, to serve as a reservoir, where the excrementitious portion of the food collects; and, also, as an excretory duct, which effects its expulsion.

INTESTINORUM LÆVITAS, Lientery.

INTESTINULA CEREBRI, Convolutions (cerebral)—i. Meibomii, Meibomius, glands of.

INTESTINULUM, (dim. of *intestinum*.) Funiculus umbilicalis.

INTESTINUM, Intestine—i. Cellulatum, Colon—i. Circumvolutum, Ileon—i. Crassum, Colon. see Intestine—i. Gracile, see Intestine—i. Grande, Colon—i. Laxum, Colon—i. Majus, Colon—i. Medium, Mesentery—i. Plenum, Colon—i. Rectum, Rectum—i. Tenne, see Intestine.

INTIMUM UNGUIS, (*intimus*, 'inmost.') Nail, root of the.

INTONATIO INTESTINALIS, (*intonare*, *intonatum*, 'to make a noise.') Borborygmus.

INTORSIO, (*in*, and *torquere*, *tortum*, 'to wrench.') Distortion.

INTOXICATIO, (*in*, and *τοξικον*, 'a poison.') Poisoning—i. Arsenicalis, Arsenicisms—i. Opiaea, Mec-nismus—i. Saturnina, Saturnism.

INTOXICATION, Poisoning, Temulencia—i. Antimonialis, Stibialism—i. des Marais, see Miasm.

INTOXICATION SATURNINE. Lead poisoning; saturnine cachexy. The aggregate of symptoms which present themselves prior to an attack of lead colic.

INTRA (L.), *Intro*, in composition, within.

INTRAARTICULAR, *Intraarticula'ris*; (*intra*, 'within,' and *articulus*, 'a joint.') Relating or appertaining to the interior of a joint—as *intra-articular* suppuration.

INTRALINGUAL SALIVARY GLANDS, see Salivary glands.

INTRALOBULAR, *Intralobula'ris*, (F.) *Intralobulaire*, *Intralobulaire*, (*intra*, and *lobulus*, 'a lobule.') Relating to the space within a lobule:—as the "*intralobular veins*," *venae seu venulae intralobulares seu centrales* of the liver:—veins which communicate with the interlobular veins, and are the radicles of the hepatic veins.

INTRAOCULAR, *Intraocula'ris*, (from *intra*, and *oculus*, 'the eye.') That which is within the eye.

INTRA-PELVIO TROCHANTÉRIEN. (*intra*, *pelvis*, and *trochanter*.) Obturator internus.

INTRAUTERINE, *Intrauteri'nis*, (*intra*, and *uterus*.) That which takes place within the womb—as *intrauterine* life.

INTRAVAGINAL, *Intravagina'lis*, (*intra*, and *vagina*.) Relating or appertaining to the interior of the vagina.

INTRAVASATION, *Intravasa'tio*, (*intra*, and *vas*, 'a vessel.') The entrance into vessels of matters formed outside of them or in their parietes.

INTRICATURA, (*in*, and *tricolor*, *tricus*, 'to make difficulties.') Decussation.

INTRINSIC, *Intrin'secus*, (F.) *Intrin'sèque*, (*intra*, and *secus*, 'towards.') Applied to the internal muscles of certain organs, as to those of the ear, tongue, and larynx. Linnæus gave the name *Intrin'seci* to internal diseases.

INTRO (L.), in composition, see *Intra*.

INTROITUS PELVIS, (*intro*, and *ire*, *itum*, 'to go.') See Pelvis—i. Vagina, Rima vulvæ, see Vagina.

INTROMISSION, *Intromis'sio*, (*intro*, and *mittere*, *missum*, 'to send.') The act of introducing one body into another—as *Intromis'sio Pe'nis*, (F.) *Intromission de la Vergé*.

INTROPELVIMETER, (F.) *Intropelvimètre*. An instrument, proposed by Madame Boivin, to measure the diameters of the pelvis. It differs from the pelvimeter of Contouly, in one of its branches being intended to be introduced into the rectum.

INTROSUSCEPTIO, (*intro*, and *suscipio*, *suscipitum*, 'to receive.') Intussusceptio—i. Entropion, Entropion.

INTUBUM, (*αὐτὸν*.) Cichorium endivia—i. Erraticum, Cichorium intybus.

INTUMESCENCE, *Intumescen'tia*, *Onco'sis*, *Dionco'sis*, (*intumescere* (*in*, and *tumescere*), 'to swell.') Augmentation of size in a part or in the whole of the body. Sauvages uses the word *Intumescen'tia* for an order of diseases, in which he comprehends polyseria, pneumatoses, anasarca, oedema, phlebotomy, and pregnancy.

INTUMESCENTIA, Intumescence—i. Gangli-formis, Ganglion of Gasser, see Facial nerve—i. Lactea Mammarum, Sparganosis—i. Lienis, Splenoncus—i. Semilunaris, Ganglion of Gasser.

INTUS INVERSUS, (*intus*, 'within,' and *inversus*, 'inverted.') Transposition of the viscera.

INTUSSUSCEPTIO, (*intus*, 'within,' and *suscipio*, *suscipitum*, 'I receive.') *Intussusceptio*. In physiology—the mode of increase peculiar to organized bodies. In pathology, like *Unvolvulus*, *Volvulus* seu *Susceptio* seu *Parasago'gē* *intestini*, *Chordap'us*, *Enteroparasio'gē*, *Indigita'tio*, *Tormen'tum*, and *Invagina'tio*, it means the introduction of one part of the intestinal canal—*intussusceptum*—into another, which serves it as a sort of vagina or sheath. Generally, it is the upper part of the small intestine which is received into the lower—*intussusceptiens*—when the intussusception is said to be *progressive*. At times, however, it is *retrograde*. As the disease cannot be positively detected by the symptoms, it must be treated upon general principles. At times, the invaginated portion has separated and been voided per anum,—the patient recovering. The disease is, however, of a very dangerous character.

INTUSSUSCEPTUM, see Intussusceptio.

INTUSSUSCIPiens, see Intussusceptio.

INTYBUS, Cichorium endivia.

INTYBUS HORTENSIS, Cichorium endivia.

INULA, (corr. from *Helenium*.) [?] See Inula helenium—i. Britannica, I. dysenterica—i. Common, I. helenium—i. Conyzæa, I. dysenterica.

INULA DYSENTERICA. The systematic name of the Lesser Inula, I. Britannica seu Conyzæa, *Aster dysentericus* seu *undulatus*, *Pulicaria dysenterica*, *Conyza*, *C. media*, *Ar'nica Sueden'sica* seu *epu'ria*, *Cuni'la mas'cula* seu *me'dia*, *Cunila'go*, (Prov.) *Mare's fat*, (F.) *Aunée* ou *Aulnée antidysentérique*, *Herbe de Saint Roch*. Nat. Ord. Compositæ. This plant was once considered to possess great antidysenteric virtues. The whole plant is acrid and somewhat aromatic.

INULA HELENIUM. The systematic name of the *El'ecampane*, *En'ula*, *E. camp'dana*, *Elen'ium*, *Necta'rium*, *Helen'ium*, *Aster helen'ium* seu *officina'lis*, *Corvisaria'lia helen'ium*, *Common In'ula*, (F.) *Aunée* ou *Aulnée*, *Aunée* ou *Aulnée officinale*. The root *In'ula* (Ph. U. S.) was formerly in high esteem in dyspepsia, cachexia, pulmonary affections, &c. It is now scarcely used.

INULA, LESSER, Inula dysenterica—i. Squarrosa, Conyza squarrosa.

INUNCTIO, (*inungere*, *inunctum* (*in*, and *ungere*, *unctum*), 'to anoint.') Inunction, Liniment.

INUNCTION, *Inunctio*, *Perunctio*, *Illit'io*, *Illit'us*, *Illinit'io*, *Oblinit'io*, *Chri'sis*, *Catach'risis*, *Diach'risis*, *Epich'risis*, *Unctio*. The act of rubbing in an ointment or a liniment, *Unctio*, *Infri'etion*, (F.) *Oncion*, *Inonction*. Inunction of the whole body with a piece of bacon, lard, or with oil, has been recommended in scarlatina, measles, and in scrofulous or tuberculous cachexia.

INUNDATIO, (*inundare*, *inundatum* (*in*, and *undare*), 'to flow over.') Depuration.

INUSTIO, (*in*, and *urere*, *ustum*, 'to burn.') Cauterization, Cauterium.

INUSTORIUM, Cauterium.

INVAGINATED, *Invagina'tus*, (*in*, and *-a*).

ginn, 'a sheath.') Applied to a part which is received into another, as into a sheath.

The *invaginated* or *slit and tail bandage* is one in which strips or tails pass through appropriate slits or button-holes.

INVAGINATIO, Invagination, Intussusceptio.

INVAGINATION, *Invagina'tio*, *Encoleo'sis*. Intussusceptio. Also, an operation which consists in introducing one end of a divided intestine into the other, with the view of restoring the continuity of the intestinal canal.

The name *Invagination* is likewise given to an operation for the radical cure of crural hernia, which consists in introducing into the crural canal a thick tent of charpie, the effect of which is, by compression and adhesive inflammation, to obliterate the neck of the hernial sac.

INVALETUDO, (*in*, and *validudo*, 'health.') Infirmary.

INVALID', (*in*, and *validus*, 'strong.') Valetudinary.

INVALIDING, (F.) *Réforme*. The removal from the army of a soldier who has become, from infirmity, unfit for service. Such was the meaning of the term in the service generally; but an Invalid Corps is now established in the United States, to which lighter and special duties are assigned, in hospitals and elsewhere.

INVALIDUS, Valetudinary.

INVASIO, (*in*, and *vadere*, *vasum*, 'to go.') Arche, Attack, Paroxysm.

INVERMINATION, (*in*, and *verminare*, *verminatum*, 'to have worms,'—*vermes*). Helminthiasis.

INVERSIO PALPEBRARUM, (*in*, and *vertere*, *versum*, 'to turn.') Entropion—i. Uteri, Hysteroptosis, Uterus, inversion of the.

INVERSION DU LAMBEAU, see Flap—i. de la Matrice, Uterus, inversion of the.

INVERTENTIA, Absorbents, Antacids.

INVESTITURA FUNICULI UMBILICALIS, (*in*, and *vestis*, 'a garment,') see Funiculus umbilicalis.

INVENTERATUS, (*in*, and *vetus*, gen. *veteris*, 'old,') Chronic.

INVISCONTIA, (*in*, and *viscum*, 'bird lime,') Incrassantia.

INVISCA'TIO OC'ULLI. A morbid adhesion of the eyelids to each other, or to the globe of the eye. Also, gluing together of the eyelids by a viscid secretion.

INVOLONTAIRE, Involuntary.

INVOLUCRUM, *Involumen'tum*, (*involvere* (*in*, and *volvere*, *volutum*), 'to fold in,') A covering: hence *Involu'era cer'e'br*i, the membranes of the brain. *Involu'era nervo'rum*, the sheaths of the nerves.

INVOLUCRUM CORDIS, Pericardium—i. Corporis commune, Integument—i. Lingue, see Tongue—i. Membranaceum, Decidua reflexa—i. Nervorum, Neurilemma—i. Reti comparatum, Retina.

INVOLUTUM, Involverum.

INVOLUNTARY, *Involunta'rius*, (F.) *Involontaire*, (*in*, and *voluntas*, 'will,') Not under the influence of the will.

INVOLVENTIA, Demulcents.

IODATED, Iodized.

IODE, (*iōds*, 'violet colored,') Iodine.

IODÉ, Iodized.

IODES, (*ios*, 'verdigris,') Æruginous.

IODIUM HYDRARGYRICUM, Hydrargyri iodidum rubrum.

IODHYDRARGYRITE DE CHLORURE MERCUREUX, Mercury, iodide of chloride of.

IOD'IC or IOD'IC, *Iod'icus*, (F.) *Iodique*. Containing iodine.

IODIDIUM HYDRARGYRI CHLORIDI, Mercury, iodide of chloride of—i. Hydrargyricum, Hydrargyri iodidum rubrum—i. Hydrargyrosus, Hydrargyri iodidum.

I'ODINE, *Iod'ina*, *Iod'ium* (Ph. U. S., 1842', *Iodin'ium* (Ph. U. S., 1851), *Iodin'eum*, *Iod'on* (Ph. B., 1864), *Iod'ium*, *Ion'ium*, *Ion'um*, (F.) *Iode*, (*iōds*, *violat'ceus*, 'of a violet colour;') so called from the violet vapour it exhales when volatilized.) It is contained in the mother waters of certain fuci, and is obtained by pouring an excess of concentrated sulphuric acid on the water obtained by burning different fuci, lixiviating the ashes and concentrating the liquor. The mixture is placed in a retort to which a receiver is attached, and is boiled. The iodine passes over and is condensed. It is solid, in the form of plates; of a bluish gray colour, of a metallic brightness, and smell similar to that of the chloride of sulphur. Its s. g. is 4.946. When heated, it becomes volatilized, and affords the vapour which characterizes it. With oxygen it forms *Iod'ic* acid, and with hydrogen *Hydriod'ic acid*. The tincture of iodine and the iodides have been employed with great success in the treatment of goitre and of some scrofulous affections. It must be administered in a very small dose and for a long period. It is said to be apt, however, to induce cholera morbus, signs of great nervous irritability, *Iodom'ethê*, (F.) *Ivresse iodique*, and emaciation of the mammae. When these symptoms, collectively termed *Iodism*, *Iodo'sis*, and *Iodin'ia*, (F.) *Cachexie iodée*, are urgent, the dose may be diminished, or it may be wholly discontinued, and afterwards resumed. When exposed to the air in a room, it has been found an effective antitubercle.

Various preparations of iodine are employed in medicine.

IODINED, Iodized,

IODINIA, see Iodine.

IODINIUM, Iodine.

IODINUM, Iodine.

IODIQUE, Iodic.

IODISM, see Iodine.

IODIUM, Iodine.

I'ODIZED, *Iodined*, *I'odated*, *Ioda'tus*, *Iodisa'tus*, (F.) *Iodé*. Impregnated or affected with iodine or its preparations.

I'ODOFORM, *Iodofo'r'mum*, (F.) *Iodoforme*, *Iodure de formyle*; (so called from its analogy to chloroform.) It is obtained by the reaction on each other of iodine, bicarbonate of potassa, water, and alcohol. It is in crystalline plates, of a beautiful citrine colour; and may be given in the same cases as iodine and the iodides. Dose, one grain, three times a day. It is the *sesquiodide of carbon*. See Carbonis sesquiodidum.

IOGNOGOSIE, Iodognosis.

IOGNOGOSIS, (F.) *Iodognosie*; (*iōdum*, and *γνωσις*, 'knowledge,') A knowledge of iodine in its various relations.

IODOHYDRARGYRATE OF POTASSIUM, see Iodine.

IODOMETHE, (*iodine*, and *μεθν*, 'drunkenness,') See Iodine.

IODOPE'GÆ, (*iōdum*, and *πνν*, 'a spring,') Mineral waters containing iodine.

IODOPHTHISIE, (*iōdum*, and *φθισις*, 'phthisis,') Loss of flesh and strength produced by iodine.

IODOSIS, see Iodine.

IODUM, Iodine.

IODURE FERREUX, Ferri iodidum.

IODURE DE FORMYLE, Iodoform—i. de Mercure, Hydrargyri iodidum—i. de Plomb, Plumbi iodidum—i. Plombique, Plumbi iodidum—i. de Potassium, Potassæ hydriodas—i. de Quinine, Quinæ hydriodas—i. de Soufre, Sulphuris iodidum—i. de Zinc, Zinci iodidum.

IODURETUM AMMONIÆ, see Iodine—i.

Amyli, see Iodine.—i. Hydrargyrosus, Hydrargyri iodidum.—i. Sulphuris, see Iodine.

ION. (ἰων.) Viola.

IONIA, *Tenacium chamæpitrys*.

IONID' IUM MARCUCCI, (dim. of *ion*, 'the violet,' or from *ion* and *eidos*, 'resemblance,') *Cinchichnechuli*—a South American plant. *Sec. Syst.* Pentandria Monogynia; *Nat. Ord.* Violariæ; said to be extremely efficacious in the Mal de San Lazaro, of Colombia.

IONIUM, (*ion*, 'a violet,') Iodine.

IONTA, (*ἰοντα*) see Excretion.

IONTHUS, *Va'rus*, *Violet Eruption*, (*ἰωνθος*, from *ion*, 'the violet,' and *athos*, 'a flower,' or *athos*, 'foulness' [?].) An unsuppurative, tubercular tumour; stationary; chiefly on the face. A genus in the class *Hæmatica*, order *Phlogotica* of Good.

IONTHUS CORYMBIFER, *Gutta rosea*—i. *Varus*, *Acne*.

IONUM, (*ion*, 'a violet,') Iodine.

IOTACIS'MUS, (from the Greek letter *Iota*.) Defective articulation,—the patient not being able to pronounce the palatals *j* and *g* soft.

IOULUS, (*ἰουλος*.) *Julus*.

IPECAC. *Apocynum androsæmifolium*, *Euphorbia corollata*, *Gillenla trifoliata*, *Ipeacuanha*, *Triosteum*—i. American, *Apocynum androsæmifolium*—i. Wild, *Euphorbia ipeacuanha*.

IPECACUAN'HA. In common parlance, often abridged to *Ipecac*, and in Ireland to *Hipp'o*. The pharmacopœial name of the root of *Cepha'elis seu Cuthico'ca ipeacuan'ha*, *Ipeacuan'ha*, *Ipeacuan*, *Cogoa'anga*, *Hypocuan'na*, *Hypocuan'na*, (F.) *Ipeacuanha améle*. *Nat. Ord.* Cinchonaceæ. It is also obtained from the *Psycho'tria emet'ica* of Peru. The odour of the root, *Ipeacuan'ha* root, *Ipeacuan'ha ra'dix*, *Ra'dix Brazilian'is*, *Ipeacuanha* (Ph. U. S.), (F.) *Racine Brésillienne ou de Brésil*,—is faint and peculiar; taste bitter, subacid, and mucilaginous; both water and alcohol extract its virtues, which depend on a peculiar principle, called *Emet'ica*. It is emetic in large doses; sudorific in smaller. *Dose*, as an emetic, gr. xx to xxx,—alone, or united with gr. i to ij of tartarized antimony.

IPECACUANHA, *Euphorbia corollata*, *Pedilanthus tithymaloides*—i. *Améle*, *Ipeacuanha*—i. Bastard, *Asclepias curassavica*, *Triosteum perfoliatum*—i. *Blanc de l'Île de France*, *Tylophora asthmatica*—i. *Blanc de St. Domingue*, *Asclepias curassavica*—i. of the Isle of France, *Tylophora asthmatica*.

IPECACUANHA, WHITE, is obtained from different species of *Richardsonia* and *Ionid'ium*. *Vit'ola ipeacuan'ha* seu *So'lea ipeacuan'ha* seu *Pom'ba'lea ipeacuan'ha* also affords it. It is weaker than the gray.

IPO. Upas—i. Toxicaria, see Upas.

IPOMŒA BATATOÏDES, (*ἰψ*, 'an insect that infests the vine,' and *ἰμοος*, 'like,' from its twisting round other plants), see *Convolvulus jalapa*—i. *Jalapa*, *Convolvulus jalapa*—i. *Macrorrhiza*, *Convolvulus jalapa*—i. *Mestitanica*, see *Convolvulus jalapa*—i. *Nil*, *Convolvulus nil*—i. *Operculata*, see *Batata de purga*—i. *Orizabensis*, see *Convolvulus jalapa*—i. *Purga*, see *Convolvulus jalapa*—i. *Schiedeana*, see *Convolvulus jalapa*.

IR, as a prefix, has the signification of Im.

IRA, Rage.

IRACUNDUS, (from *ira*.) *Rectus externus oculi*.

IRAL'GIA, *Ireal'gia*, *Iridal'gia*, (*iris*, and *αλγος*, 'pain,') Pain in the iris.

IRASCIBLE, (*ira*, 'anger,') *Rectus externus oculi*.

IREALGIA, *Iralgia*.

IREONCION, (*iris*, and *ογκος*, 'a hook,') *Iriancistron*.

I'RIAN, *Iria'nus*, (F.) *Irien*. Belonging or relating to the iris.

IRIANCIS'TRON, *Iridancist'ron*, *Ireonc'cion*, (*iris*, and *αγκιστρον*, 'a hook,') A hooked instrument, used by Schlagintweit in the formation of an artificial pupil.

IRIDE'ÆA EDU'LIS. One of the *Algæ*, eaten in Scotland and the Southwest of England.

IRIDALGIA, (*irido*, and *αλγος*, 'pain,') *Iralgia*.

IRIDANCISTRON, *Iriancistron*.

IRIDAREO'SIS, (*irido*, and *αραιωσις*, 'diminution,') Atrophy of the iris.

IRIDAUXE'SIS, (*irido*, and *αυξησις*, 'augmentation,') *Staphylo'ma Iridis*, *S. U'veæ*, *Iridomeo'is*, *Lymphon'eus Iridis*. Exudation of fibrin into the tissue of the iris.

IRIDECTOMEDIALYSIS, (*irido*, *εκτομη*, 'excision,' and *διαλυσις*, 'separation,') see *Coretomia*.

IRIDECTOMIA, (*irido*, and *εκτομη*, 'excision,') see *Coretomia*.

IRIDENCELEISIS, (*irido*, *en*, and *κλειω*, 'to close,') see *Coretomia*.

IRIDENCEISMUS, see *Coretomia*.

IRIDERE'MIA, (*irido*, and *ερημος*, 'deprived of,') *Iridostere'sis*. Absence of iris, either apparent or real.

IRID'ESIS, (*iris* and *δew*, 'I bind,') An operation for artificial pupil, which consists in opening the cornea so as to admit the canula forceps, and, by it, drawing out a portion of the iris near its ciliary attachment, and strangulating a piece of it by means of a thread of moistened silk.

IRIDITIS, (*irido*, and *itis*.) *Iritis*.

I'RIDO, (from *iris*, gen. *iridos*), in composition, *iris*.

IRIDOCELE, (*irido*, and *κηλη*, 'a tumour,') *Pto'sis iridis*.

IRIDOCOLOBOMA, *Coloboma iridis*.

IRIDODIALYSIS, (*irido*, and *διαλυσις*, 'separation,') *Coretomia*.

IRIDODONE'SIS, (*irido*, and *δονew*, 'I shake,') Tremulousness of the iris. See *Hippus*.

IRIDOMALA'CIA, (*irido*, and *μαλακια*, 'softness,') Mollescence or softening of the iris.

IRIDOMELANO'MA, gen. *Iridomelano'matis*, *Iridomelano'sis*, (*irido*, and *μελανew*, 'to colour black,') The deposition of black matter on the tissue of the iris.

IRIDONCO'SIS, *Iridon'cus*, *Ironco'sis*, *Hyperronco'sis Irid'is*, (*irido*, and *ογκος*, 'a swelling,') Tumefaction or thickening of the iris. Also *Iridauxesis*.

IRIDONCUS, *Iridoncosis*.

IRIDOPERIPHAC'TIS, *Iridoperiphac'itis*, (*irido*, *περι*, 'around,' and *phacitis*, inflammation of the lens.) Inflammation of the anterior hemisphere of the capsule of the lens.

IRIDOPLANIA, (*irido*, and *πλανη*, 'wandering,') *Hippus*.

IRIDOPTOSIS, *Pto'sis iridis*.

IRIDORRHAGAS, gen. *Iridorrhag'adia*, *Iridorrhage*, *Iridor'rhox*, *Iridoschis'ma*, *Fissu'ra Irid'is*, (*irido*, and *ρῆγας*, 'a fissure,') Fissure of the iris. See *Coloboma*.

IRIDORRHOGE, (*irido*, and *ρῆγῃ*, 'a fissure,') *Iridorrhagas*.

IRIDORRHOX, (*irido*, and *ρῆξ*, 'a fissure,') *Iridorrhagas*.

IRIDOSCHISMA, gen. *Iridoschism'atia*, (*irido*, and *σχισμα*, 'a fissure,') *Coloboma Iridis*, *Iridorrhagas*.

IRIDOSTERESIS, (*irido*, and *στερωσις*, 'privation,') *Irideremia*.

IRIDOTOMENCEISIS, (*irido*, *τομη*, 'section,' and *εγκλειω*, 'to include,') *Coretomia*.

IRIDOTOMEDIALYSIS, (*irido*, *τομη*, 'section,' and *διαλυσις*, 'separation,') see *Coretomia*.

IRIDOTOMIA, (*irido*, and *τομη*, 'section,') see *Coretomia*.

IRIDOTROMUS, (*irido*, and *τρεμος*, 'trembling,') *Hippus*.

IRIEN, Irian.

IRIS, gen. *Iridis*, (iris, gen. *iridis*.) Gram'me. So called from its resembling the rainbow in a variety of colours. A membrane, stretched vertically at the anterior part of the eye, in the midst of the aqueous humour, in which it forms a kind of circular, flat partition, separating the anterior from the posterior chamber. It is perforated by a circular opening called the *pupil*, which is constantly varying its dimensions, owing to the varying contractions of the fibres of the iris. Its posterior surface has been called *uvea*, from the thick, black varnish which covers it. The pigmentary stratum, on its free surface, appears to be bounded by a delicate, but sharply defined, line, which, has been described as a special membrane—*Membrana pigmenti seu limbitans seu limbitans Paci'ni seu Jacobi Arnold*. The greater circumference of the iris is adherent to the ciliary processes and circle. It has an external plane of radiated fibres and an internal one of circular fibres, which serve—the one to dilate, the other to contract the aperture of the pupil. The iris receives the irian nerves. Its arteries are furnished by the long ciliary arteries, which form two circles by their anastomoses; the one very broad, near the great circumference; the other, smaller, and seated around the circumference of the pupil. The veins of the iris empty themselves into the *Vasa vorticosa*, and into the long ciliary veins. The use of the iris seems to be,—to regulate by its dilatation or contraction, the quantity of luminous rays necessary for distinct vision. The different colours of the iris occasion the variety in the colours of the human eye.

IRIS, see Herpes Iris—i. Common, I. Germanica—i. Commun, I. Germanica—i. de Florence, I. Florentina.

IRIS FLORENTINA, *Flor'entine i'ris* or *or'ris*, (F.) *Iris de Florence*. Order, Iridaceæ. The rhizoma of this plant is extremely acrid in its recent state; and, when chewed, excites a pungent heat in the mouth, that continues for several hours. When dried, the acrimony is lost, or nearly so, and the smell is very agreeable. It is indebted to its agreeable flavour for its retention in the pharmacopœias, although it is ranked as an expectorant. The *orris* root of the shops is said to consist of the rhizomes of three species—*I. Florentina*, *I. pallida*, and *I. Germanica*.

IRIS FÆTIDIS'SIMA, *I. fœtida*, *Spathulæ fœtida*, *Xy'ris*, *Stinking i'ris*, (F.) *Iris puant*, *Gayeluy puant*. The root has been esteemed antispasmodic and narcotic.

IRIS GERMANICA. The systematic name of the *Flower-de-Luce*, *Iris nos'tras* seu *vulgaris*, *Aier'sa*, *Common i'ris* or *or'ris*, *Gladi'olus cœruleus*, (F.) *Iris commun*, *Flambe*. The fresh roots have a disagreeable smell, and an acrid, nauseous taste. They are powerfully cathartic, and are given in dropsies, where such remedies are indicated.

IRIS HEXAGONA, I. versicolor.

IRIS LACUS'TRIS, *Dwarf Iris*, *Dwarf lake iris*; has the properties of I. versicolor.

IRIS LAKE, DWARF, I. lacustris—i. Lutea, I. pseudacorus—i. des *Marais*, I. pseudacorus—i. Nostras, I. Germanica—i. Palustris, I. pseudacorus.

IRIS PSEUDACORUS. The systematic name of the *Yellow waterflag*, *Iris palustris* seu *lutæa*, *Gladi'olus luteus*, *Ac'orus vulgaris* seu *adulterinus* seu *palustris*, *Pseudac'orus*, *But'omon*, (F.) *Iris des marais*, *Fauc acore*, *Flambe bâtarde*, *Acore bâtarde*. The root has an acrid, styptic taste. It is an emmenagogue, and acrid astringent. The expressed juice is diuretic, and said to be a useful application to serpiginous and other cutaneous affections. Rubbed on the gums, or chewed, it is said to cure toothache.

IRIS PUANT, I. fœtidissima—i. Stinking, I. fœtidissima.

IRIS VERSICOLOR, *I. hexago'na*, *Blue flag*. The rhizoma of this is an active cathartic, and has been much used, as such, by the American Indians. It is reputed to be diuretic.

IRIS VIRGINICA, *Slender Blue flag*; and *Iris verna*, are said to have virtues like those of I. versicolor.

IRIS VULGARIS, I. Germanica.

IRISITIS, Iritis.

IRITIS, *Inflammation of the i'ris*, *Inflammatio i'ridis*, *Iris'i'tis*, *Iridi'tis*, (*iris*, and *itis*.) *Uve'i'tis* of some. The chief symptoms are;—change in the colour of the iris; fibres less movable; tooth-like processes shooting into the pupil; pupil irregularly contracted, with the ordinary signs of inflammation of the eye. If the inflammation does not yield, suppuration takes place; and although the matter may be absorbed, the iris remains immovable. It is often caused by syphilis. The general principles of treatment are, to deplete largely and exhibit mercury freely; along with attention to other means advisable in ophthalmia. The free use of quinia is sometimes serviceable. When the inflammation is seated in the serous covering of the iris, it is termed *Iritis serosa*.

IRON, (Sax. *Iren* or *Ipen*.) Ferrum—i. Acetate of, Ferri acetas—i. Ammoniated, Ferrum ammoniatum—i. Ammonio chloride of, Ferrum ammoniatum—i. Ammonio-citrate of, Ferri ammonio-citras—i. Ammonio-tartrate of, Ferri ammonio-tartaras—i. and Alumina, sulphate of, Ferri et aluminis sulphas—i. and Ammonia, citrate of, Ferri ammonio-citras.

IRON BARK, see Kino.

IRON, BLACK OXIDE OF, *Oxydum ferri nigrum*—i. Bromide of, see Bromine—i. Carbonate of, precipitated, Ferri subcarbonas—i. Carbonate of, saccharine, Ferri proto-carbonas—i. Carburet of, Graphites—i. Chloride of, Ferri chloridum—i. Citrate of, Ferri citras—i. Solution of Citrate of, *Liquor ferri citratis*—i. Citrate of the magnetic oxide of, see Ferri ammonio-citras—i. and Ammonia, citrate of, Ferri et ammoniæ citras—i. Ferrocyanuret of, Prussian blue—i. Ferropussiate of, Prussian blue.

IRON FILINGS, PURIFIED, Ferri limatura purificata.

IRON HYDRATED OXIDE OF, *Ferrum oxydatum hydratum*—i. Hydrated peroxide of, *Ferrum oxydatum hydratum*—i. Hydrated tritoxide of, *Ferrum oxydatum hydratum*—i. Hydriodate of, Ferri iodidum—i. Hydrobromate of, see Bromine—i. Hydrobromide of, *Ferrum oxydatum hydratum*—i. Hypophosphite of, see Calcei hypophosphis—i. Iodide of, see Iodine, and Ferri iodidum—i. Ioduret of, Ferri iodidum—i. Lactate of, Ferri lactas—i. *Liquor*, Ferri acetas—i. and Magnesia, citrate of, Ferri et Magnesiæ citras—i. Magnetic Oxide of, *Oxydum ferri nigrum*—i. Muriate of, Ferri chloridum—i. Nitrate of, solution of, see *Tinctura ferri muriatis*—i. Perchloride of, Ferri chloridum—i. Peroxide of, Ferri subcarbonas—i. Persulphate of, see Ferri sulphas—i. Persulphate of, solution of, see Ferri sulphas—i. Phosphate of, Ferri phosphas—i. Pills of, compound, *Pilulæ ferri compositæ*—i. and Potass, Tartrate of, *Ferrum tartarizatum*—i. Potassio-citrate of, see Ferri ammonio-citras—i. Protocarbonate of, Ferri protocarbonas—i. Protoxide of, lactate of, Ferri lactas.

IRON PYRITES, Ferri sulphuretum.

IRON, PYROPHOSPHATE OF, see Ferri phosphas—i. Quevenne's Ferri pulvis.

IRON AND QUINIA, citrate of, Ferri et Quiniæ citras—i. and Quinia, hydriodate of, see Quinia, iodide of Iodhydra of.

IRON REDUCED, *Ferri pulvis*.

IRON, SESQUICHLORIDE OF, *Ferri chloridum*—i. Sesquinitrate of, solution of, see *Tinctura ferri muriatis*—i. Sesquioxide of, *Ferri subcarbonas*—i. Sodicitrate of, see *Ferri ammonioicitras*—i. Subcarbonate of, *Ferri subcarbonas*—i. Subsulphate of, solution of, see *Ferri sulphas*—i. Sulphate of, *Ferri sulphas*—i. Sulphuret of, *Ferri sulphuretum*—i. Tannate of, *Ferri tannas*—i. Tartarized, *Ferum tartarizatum*—i. Ternitrate of, solution of, see *Tinctura ferri muriatis*—i. Trithydroferrocyanate of, Prussian blue—i. Valerianate of, *Ferri valerianas*.

IRON WEED, *Vernonia Novboracensis*.

IRONCOSIS, *Iridoncosis*.

IRRADIATION, *Irradiatio*, (*ir*, and *radius*, 'a ray.') Shooting or proceeding from a centre—as an 'irradiating pain.'

IRREDUCIBLE, (*F*.) *Irreductible*. An epithet given to fractures, luxations, herniæ, &c., when they cannot be reduced.

IRREDUCTIBLE, *Irreducibile*.

IRREGULAR, *Irregularis*, *Anomalous*, (*ir*, and *regula*, 'a rule.') A term chiefly applied to the types of a disease; and also, to the pulse when its beats are separated by unequal intervals.

IRREGULAR PRACTITIONER. One who does not practise his profession according to rules sanctioned by law or custom.

IRREMEDIABLE, *Incurable*.

IRREPTIO, (*ir*, and *repo*, *reptum*, 'to creep,') Attack.

IRRIGATION, *Irrigatio*, *Embrocatio*, *Irroratio*: 'the act of watering or moistening,' (*ir*, and *rigare*, *rigatum*, 'to water.') The methodical application of water to an affected part, to keep it constantly wet.

IRRITABILITAS, (*irritare*, 'to excite,') Irritability—i. Halleriana, Irritability—i. Morbosa, see *Subsultus tendinum*—i. Vesicæ, *Cysterethismus*.

IRRITABILITY, *Irritabilitas*, (*F*.) *Irritabilité*; *Vie irritabilité*, *V. Vie's*, *V. insita* of Haller *V. vitalis* of Gorter, *Oscillatio* of Boerhaave, *Tonic power* of Stahl, *Vita propria*, *Irritabilitas Halleriana*, *Inherent power*, *Contractility*, *Excitability*, &c. (*F*.) *Irritabilité*, *I. Hallérienne*. A power, possessed by all living, organized bodies of being acted upon by certain stimuli, and of moving responsive to such stimuli. It is the ultimate vital property.

IRRITABLE, *Irritabilis*, (*irritare*, 'to excite,' and *habilis*, 'capable.') That which is endowed with irritability. Every living organized tissue is irritable; that is, capable of feeling an appropriate stimulus, and of moving responsive to such stimulus. Irritable is often used in the same sense as *impressible*, as when we speak of an irritable person, or habit, or temper. This last condition has been variously termed—*Debilitas nervosa*, *D. Erethisica*, *Hypercinesia nervosa*, *Neurasthenia*, *Neurosthenia*, *Sensibilitas anomala*, *S. Morbosa*, *S. anxia*.

IRRITAMEN, Irritant, Stimulus.

IRRITAMENTUM, Erethism, Irritant, Stimulus—i. Metallicum, Galvanism.

IRRITANT, *Irritans*, *Irrita'men*, *Irritamentum*. That which causes irritation or pain, heat and tension; either mechanically, as puncture, acupuncture, or scarification; chemically, as the alkalies and acids; or in a specific manner, as cantharides.

Irritants—Irritantia, Excitantia—are of great use in the treatment of disease.

IRRITATED, see Inflamed.

IRRITATING POISON, see Poison.

IRRITATIO, Erethism, Gargale, Irritation.

IRRITATION, *Irritatio*, *Erethismus*. The state of a tissue or organ, in which there is excess of vital movement; commonly manifested by

increase of the circulation and sensibility. Broussais defines irritation to be;—the condition of an organ, the excitation of which is carried to so high a degree, that the equilibrium resulting from the balance of all the functions is broken. In this signification, he also uses the word *surirritation*, which he considered as a higher degree, and as the essential cause of fever. Irritation is the common precursor of inflammation.

IRRITATION, MORBID, *Constitutional irritation*, *Ir'ritative fever*, *Pseudofever*, is that excitement which occurs after injuries done to the body, or to any part thereof;—constituting cases of *diseased sympathy*.

IRRITATIVE FEVER, Irritation, morbid.

IRRORATIO, (*ir*, and *ros*, gen. *roris*, 'dew,') Irrigation.

IS, (*is*, gen. *vōs*, &c.) Fibre.

ISADELPHIA, (*isos*, 'equal,' and *adelphos*, 'brother.') A double monstrosity, in which both bodies are equally and perfectly developed, and possessed of independent vital organs; and which are united by parts of little significance.

ISATIS TINCTORIA, (*isatis*,) *Wood*, *Pos'tel*. *Ord.* Capparidaceæ. A European plant, whose leaves have a fugitive pungent smell, and an acrid durable taste. They are not used, however, in medicine at the present day, but are the source of the dye-stuff, *wood*.

ISCA, (*ισκν*,) *Boletus ignarius*.

ISCHÆMIA, (*ischæ*, and *αἷμα*, 'blood,') Morbid retention or suppression of an habitual flux of blood, as of the hemorrhoidal or menstrual flux or of epistaxis. See *Hæmatostatica*. Also local arterial anæmia, in which the blood is prevented from flowing into the parts for which it is destined. (*Virchow*.)

ISCHÆMON, *Ischæmum*. A medicine which restrains or stops bleeding.

ISCHAS, (*ισχος*, gen. *ισχados*, 'a dried fig,') see *Ficus*.

ISCHESIS, (from *ισχω*, 'to restrain,') Suppression or retention of a discharge or secretion.

ISCHIA, MINERAL WATERS OF. In this volcanic isle, five miles from Naples, there are several thermal waters, one of which, that of Gurgitello—raises the thermometer of Fah. to 167°.

ISCHIACUS, (*ισχιακος*,) *Ischiatic*.

ISCHIADELPHIUS, (*ischion*, and *adelphos*, 'a brother,') *Ischiopages*.

ISCHIADICUS MORBUS, (from *ischias*,) *Neuralgia femoro-poplitea*—i. *Nervus Sciatic nerve*.

ISCHIADOCELE, *Ischiocoele*.

ISCHIAGRA, or ISCHIA'GRA, *Is'chias*, *Is'chal'gia*, (*ischion*, and *αἶμα*, 'a seizure,') A name given to ischiatic gout. *Femoro-popliteal neuralgia* has, also, been so called.

ISCHIAL, *Ischia'lis*. Relating or belonging to the ischion.

ISCHIALGIA, (*ischion*, and *αλγος*, 'pain,') *Ischiagra*, *Neuralgia femoro-poplitea*.

ISCHIAS, gen. *Ischi'adis*, (*ισχιας*, gen. *ισχιαδος*,) *Ischiagra*—i. *Nervosa antica*, *Neuralgia femoro-poplitea*—i. *Nervosa Cutunui*, *Neuralgia femoro-poplitea*—i. *Nervosa digitalis*, *Neuralgia cubito-digitalis*—i. *Nervosa postica*, *Neuralgia femoro-poplitea*—i. *Rheumatica*, *Neuralgia femoro-poplitea*—i. à *Sparganosi*, *Phlegmatia dolens*.

ISCHIATIC, *Ischiad'ic*, *Ischiad'icus*, *Ischiad'icus*, *Ischi'acus*, *Sciatic*, (*F*.) *Ischiatique*, (from *ischion*,) whence the word *Sciatic*. An epithet given to parts connected with the ischium. The SCIATIC NOTCHES, (*F*.) *Echancrures Ischiatiques*, are formed by this bone; the ISCHIATIC SPINE belongs to it, and gives attachment to the small sacro-sciatic ligament. The TUBEROSITY OF THE ISCHIUM, *Os sedentarium*. *Tuber Ischi'*, (*F*.) *Tubérosité ischiatique*, is formed by it. It receives the insertions of different muscles of the thigh,

and forms the projection on which the body rests when seated.

ISCHIATIC ARTERY, *Sciatic Artery*, (F.) *Artère Fémoro-Poplitée* (Ch.) arises singly from the hypogastric, or with the gluteal; and seems to be really a continuation of the trunk of the hypogastric. It issues from the pelvis, at the lower part of the great sciatic notch; and, afterwards, divides into a considerable number of branches, which are distributed particularly to the posterior and superior region of the thigh.

ISCHIATIC REGION, *Sciatic region*; *Regio ischiadica*. The region of the hip.

ISCHIATIC VEIN presents the same arrangement as the artery.

ISCHIATICUS, (ισχιακός, from *ischias*.) Sciatic.

ISCHIATOCELE, Ischiocele.

ISCHIDROSIS, *Sudoris suppressio*, *Oligidria*, (*ischio*, and *ἰδρωσ*, 'sweat.') Suppression of perspiration.

ISCHIOCAVERNOSUS. Belonging to the ischium and corpus cavernosum.

ISCHIOCAVERNOSUS MUSCLE, *Collateralis*, *Director* seu *Erector* seu *Sustentator Penis*, (F.) *Érecteur du Pénis*, *Ischio-uréthral*, (Ch.) is a small, long, flat muscle, which surrounds the origin of the corpus cavernosum. It is fixed below to the inner side of the tuberosity of the ischium, and above to the root of the penis, where it is confounded with the fibrous membrane of the corpus cavernosum. It draws the root of the penis downwards and backwards.

ISCHIOCAVERNOSUS OF THE FEMALE, *Erector* seu *Sustentator* seu *Superior rotundus clitoridis*, (F.) *Ischio-clitoriden*, *Ischio sous-clitorien*, (Ch.), is arranged nearly as in the male, but is less bulky. It arises, by aponeurosis, from the tuberosity of the ischium, and terminates by embracing the corpus cavernosum of the clitoris, to the erection of which it appears to contribute.

ISCHIOCE'LE, *Ischiatocele*, *Ischiadocele*, *Her'nia ischiadica*; (*ischion*, and *κληη*, 'tumour') *Ischiatic hernia*. One in which the viscera issue by the great sciatic notch. It is a rare disease. The protruded parts must be reduced—the patient being placed in a favourable position, and they must be retained by a bandage.

ISCHIOCLITORIANUS. That which is connected with the ischium and clitoris.

ISCHIOCLITORIAN ARTERY of Chaussier is a division of the internal pudic, which furnishes the two arteries of the clitoris—the superficial and deep-seated.

ISCHIOCLITORIAN NERVE, of the same professor, is the superior branch of the pudic nerve, which is distributed to the clitoris.

ISCHIOCLITORIDIEN, *Ischiocavernosus*—i. *Coccygeus*, *Coccygeus*—i. *Créti-tibial*, *Semitendinosus*—i. *Femoralis*, *Adductor magnus*—i. *Fémoro-péronier*, *Biceps flexor cruris*.

ISCHIODYMIA, (*ischion*, and *δυειν*, 'to mingle,' or *διδυμος*, 'a twin.') See *Somatodymia*.

IS'CHION, (ισχον,) *Is'chium*, *Os is'chii*, *Os coxen'dicis*, *Hucklebone*, *Seatbone*, (Sc.) *Heuckbane*, *Hukebane*, *Lungiebane*, (Prov.) *Chancebone*, *Pinbone*, (F.) *Os de l'assiette*, (from *ισχω*, 'I support,' or from *ισχω*, 'I arrest,' 'I retain;'—because that bone serves as a base or support for the trunk, when we are seated, see *Ischiatic*.) The lowermost of the three portions which compose the os innominatum in the foetus and young individual. The inferior region of the same bone has, also, been called *is'chium*, in the adult. According to Hesychius, the ancients designated by the word *ischion* the capsular ligament of the coxo-femoral articulation, as well as the articulation itself.

ISCHION, Haunch.

ISCHIONEURALGIA, *Neuralgia femoro-poplitea*.

ISCHIOPA'GES, *Ischiadelphus*, (*ischion*, and *παγω*, 'I fasten.') A monstrous union of two foetuses, in which they are attached to each other by the ischia. See *Hypogastrodidymus*.

ISCHIO-PENIEN, see *Pudic Nerve*.

ISCHIO-PÉRINÉAL, *Transversus perinæi*.

ISCHIOPTHYSIS, *Tabes coxaria* seu *ischiadica*, *Phthisis ischiadica*, (*ischion*, and *phthisis*.) Phthisis in consequence of suppuration of the hip-joint. See *Coxarum morbus*.

ISCHIO-POPLITI-FÉMORAL, *Semi-membranosus*—i. *Popliti-tibial*, *Semi-membranosus*—i. *Prétibial*, *Semitendinosus*.

ISCHIOPROSTATICUS. Winslow, Sancto-rini, Albinus, and Sömmering, have given this name to the fibres of the transversus perinæi muscle, which go towards the prostate.

ISCHIORETAL FOSSA, *Perineal fossa*.

ISCHIO-SOUS-CLITORIEN, *Ischio-cavernosus*—i. *Sous-trochantérien*, *Quadratus femoris*—i. *Spini-trochantérien*, *Ischiotrochanterianus*—i. *Uréthral*, *Ischiocavernosus*.

ISCHIOSIS, (from *ischion*.) *Neuralgia femoropoplitea*.

ISCHIOTROCHANTERIA'NUS, *Gem'ini*, *Gemelli*. Part of the *Maraspia* of Cowper; *Cur-neum Maraspium*, *Ischio-spini-trochantérien*, *Secundus et tertius quadrigem'inus*, (F.) *Muscle cannelé*, *Accessoire de l'obturateur interne*. Two small, fleshy bundles, long and flat, which arise—the superior on the outside of the sciatic spine; the inferior behind the tuberosity of the ischium. Both pass horizontally outwards, and are attached to the tendon of the obturator internus, which they accompany into the fossa of the trochanter. These muscles are rotators of the lower limb outwards. They can, also, turn the pelvis on the femur, in standing on one foot.

ISCHIUM, *Ischion*.

ISCHL MINERAL WATERS OF. These springs, which are in Upper Austria, contain iodine and bromine.

ISCHNOPHON'IA, (ισχνος, 'slender,' and *φωνη*, 'voice,') *Slenderness of voice*.

ISCHNOTES, (ισχνωσις,) *Emaciation*.

ISCHO, (ισχω, 'to restrain,') in composition, arrest, retention.

ISCHOBLENNIA, (*ischio*, and *βλεννα*, 'mucus,') The suppression of a mucous discharge.

ISCHOCENO'SIS, (*ischio*, and *κωσις*, 'evacuation,') Retention or suppression of a natural evacuation—as of the menses.

ISCHOCOL'IA, (*ischio*, and *χολη*, 'bile,') *Retention bilis*. Retention or suppression of the biliary secretion.

ISCHOCOILIA, (*ischio*, and *κοιλια*, 'the bowels,') Constipation.

ISCHOCOPRIA, (*ischio*, and *κοπρος*, 'excrement,') Constipation.

ISCHOGALAC'TIA, *Retention lactis*, (*ischio*, and *γαλα*, gen. *γαλακτος*, 'milk,') Want of milk in the mamma.

ISCHOLOCH'IA or *Ischolocho'ia*, *Lochioschesis*, *Suppression lochio'rum*, (*ischio*, and *λοχεια*, 'the lochial discharge,') *Retention lochio'rum*. Suppression of the lochial discharge.

ISCHOMENIA, (*ischio*, and *μηνες*, 'the menses,') *Amenorrhœa*.

ISCHOPHONIA, (ισχοφωνια, from *ischio*, and *φωνη*, 'voice,') *Balbuties*.

ISCHOSPER'MIA, (*ischio*, and *σπερμα*, 'sperm,') *Seminis retention*. Retention or suppression of the spermatic secretion.

ISCHURET'IC, *Ischuret'icum*, (F.) *Ischurétique*, (*ischio*, and *ουρον*, 'urine,') A medicine for relieving suppression of urine. A diuretic.

ISCHURIA, (ἰσχυρία) *Urinæ suppressio*, *Stoppage of urine*, *Suppression lo'tii*, *Paruria retentio'nis*. Retention of, or impossibility of discharging, the urine, *Anurorrhée* of Piorry. Often restricted to suppression of the secretion, *Renal Ischuria*, *Anuria*, *Anure'sis*, *Nephranuria*, *Anu-voerinie* of Piorry, *Isch'ury*.

Ischuria Vera is that in which the urine, having accumulated in the bladder, the patient is unable to pass it, notwithstanding the inclination which constantly distresses him. In *false ischuria*, *Paruria in'ope*, *P. retentio'nis renalis*, *Ischuria no'tha seu spu'ria*, of some, (F.) *Suppression d'Urine*, owing to some disease of the kidney or ureters, the urine cannot reach the bladder.

Ischuria has likewise received various other names, according to the seat and character of the obstacle which opposes the exit of the urine: hence the expressions—*Renal*, *Ureter'ic*, *Ves'ical*, *Ure'thral*, and *Cal'culous Ischuria*. *Vesical Ischuria* is generally used synonymously with *Retention of Urine*, which see.

ISCHURIA CYSTOPLEGMATICA, *Cystirrhœa*—i. *Phimosica*, see *Phimosius*—i. *Spasmodica*, *Cystospasmus*—i. *Spuria*, see *Ischuria*—i. *Urethralis à phimosi*, see *Phimosius*—i. *Vera*, see *Ischuria*—i. *Vesicalis*, *Retention of urine*.

ISIC'IUM, (ἰσικιον) *Psicos*, *Insic'ium*, *Tomaculum*, *Tom'aculum*, *Tomac'inum*. A kind of sausage, formed of mixed meat and spices.

ISINGLASS, ([G.] Hausen, 'a sturgeon,' and Blase, 'a bladder.') *Ichthyocolla*.

I'SINGLASS, JAP'ANESE. A substance, obtained from seaweed, in China, which is used as a substitute in the formation of jellies, under the name *seaweed jelly*. It is sent from Singapore, sweetened, flavoured, and ready for use. In this state it may be kept for several years.

ISIR, Elixir.

ISIS NOBILIS, see *Coral*.

ISLAND OF REIL, *Insula cerebri*.

ISLE OF WIGHT, *CLIMATE OF*. This beautiful island is a favourable summer retreat for invalids. Undercliff is the situation chosen as a winter residence for phthisical valetudinarians. Cowes, Niton, Sandown, Shanklin, and Ryde, are delightful summer residences.

ISNARDIA PALUSTRIS, (after D'Isnard, a French botanist,) *Ludwigia palustris*.

ISO, (ἴσος, 'equal,') in composition, equal.

ISOCH'RONOUS, *Isch'ronus*, *Isch'ron'ius*, *Isod'romus*, (F.) *Isochrome*, (iso, and χρόνος, 'time,') That which takes place in the same time, or in equal times. The pulsations of the arteries, in various parts of the body, are nearly isochronous.

ISOCH'RYSON, *Auro compar*, (iso, and χρυσός, 'gold,') A collyrium, described by Galen as worth its weight in gold. Libavius has also given this name to an amalgam, made with equal parts of antimony and mercury.

ISOC'RATES, (iso, and κρατος, 'strength,' or κεραννυμι, 'to mix,') A mixture of equal parts of wine and water.

ISODROMUS, (iso, and δρομος, 'a course,') *Isochronous*.

ISOLUSIN, (iso, and λωω, 'to wash,' because washed in water and rectified spirit,) see *Polygala senega*.

ISOM'ERIC, *Isomer'icus*, *Isom'ernus*, *Isom'e'rus*, *Isom'orus*, (iso, and μέρος, 'part,') An epithet applied to different bodies which agree in composition, but differ in properties. The condition is termed *Isom'erism*.

ISOMERISM, see *Isomeric*.

ISOMORPHISM, (iso, and μορφη, 'form,') see *Isomorphus*.

ISOMORPHOUS, *Isomor'phus*, *Isomor'phicus*. An epithet applied to different bodies which have

the same crystalline form. The condition is called *Isomorph'ism*.

ISONANDRA GUTTA, see *Gutta percha*.

ISOPATHIA, *Isop'athy*, (iso, and πάθος, 'disease,') This term has been used by some of the German writers to signify the cure of diseases by the disease itself, or its products;—under the hypothesis, that every contagious disease contains in its contagious matter the means for its cure;—thus, that variola may be cured by homœopathic doses of viruliferous matter; syphilis, by venereal matter, &c., &c. Others have given to isopathy another form;—maintaining, that every diseased organ has its remedy in the same organ,—that eating liver, for example, will remove disease of the liver!

Dr. J. M. B. Harden, of Georgia, has used the term "*Parallexism of Diseases*;" for "the disposition of diseases to 'anastomose with each other,' or to wear each other's livery."

ISSUE, (from exeo, (ex, and eo,) 'to go out,') *Fonticulus*—i. *Pias*, *Aurantia curassaventa*, see *Fonticulus*.

ISTHME DU GOSIER, *Isthmus of the fauces*.

ISTHMION, (ἰσθμιον) *Isthmus*, *Pharynx*.

ISTHMITIS, *Inf'ammatio Fau'cium seu Pa'lati*, *Angi'na sim'plex seu mit'is seu fau'cium*, *Ang'or fau'cium*, *Cyan'chê sim'plex*, *Fau'citis* (F.) *Angine simple*, (ictimus, and itia.) Inflammation of the isthmus faucium, and of the fauces. See *Cynanche pharyngea*, and *Angina*.

ISTHMUM, (ἰσθμους) *Isth'us*.

ISTHMOCATARRHUS, (*isthmus*, and καταρροή, 'catarrh,') *Catarrh of the fauces*.

ISTHMODYN'IA, (*isthmus*, and δύσιν, 'pain,') *Pain in the fauces*.

ISTHMOPL'E'GIA; (*isthmus*, and πλῆγην, 'stroke,') *Paralysis of the fauces*.

ISTHMOS, (ἰσθμς,) *Isthmus*, *Pharynx*.

ISTHMYTYPHUS, *Cynanche maligna*.

ISTHMUS, (ἰσθμς,) *Isth'mion*, *Isth'micus*, (F.) *Isthme*. A tongue of land joining a peninsula to a continent, or which separates two seas; a narrow pass. Anatomists have given the name *Isth'mus of the Fauces*, i. *Fau'cium*, *Claust'rum Gut'turis*, (F.) *Isthme du gosier*, to the strait which separates the mouth from the pharynx. It is formed above by the velum palati and uvula; at the sides, by the pillars of the fauces and the tonsils; and below, by the base of the tongue. Also, the Fauces.

ISTHMUS OF THE ENCEPH'ALON. The narrowed and constricted portion of the encephalic mass, situate between the cerebrum, the cerebellum and the medulla oblongata, which corresponds to the free margin of the tentorium cerebelli, and comprises the pons varolii, and middle peduncles of the cerebellum, the peduncles of the cerebrum, the tubercula quadrigemina, the superior peduncles of the cerebellum, and the valve of Vieussens.

ISTHMUS OF THE FOS'SA OVA'LIS: *Isthmus Vieussen'ii*, *Stri'ga cartilagine'sa cor'dis*, is the prominent arch formed above the fossa ovalis by the union of the two pillars which bound the cavity.

ISTHMUS HEPATIS, see *Lobulus anonymus*.

ISTHMUS OF THE THYROID GLAND is a narrow band that unites the two chief lobules composing the thyroid gland.

ISTHMUS URETHRÆ, see *Urethra*—i. *Uteri*, see *Uterus*—i. *Vieussenii*, *Isthmus of the fossa ovalis*, see *Ovalis fossa*.

ITALIAN JUICE, *Extractum glycyrrhizæ*.

ITALICUS MORBUS, ('Italian disease,') *Syphilis*.

ITCH, (ἱσχη,) *Psora*—i. *Animalcule*, see *Psora*—i. of *Animals*, *Scabies ferina*.

ITCH, BAKERS', see *Psoriasis*—i. *Barbers'*, *Sycosis*—i. *Bricklayers'*, *Eczema impetiginodes*.

ITCH, COW'POX. A cutaneous eruption, observed and described by Göllis, which appeared

after the fourteenth day from vaccination, and consisted of isolated vesicles, often filled with a puriform fluid.

ITCH, DRUNKARDS', *Psora ebriorum*—i. Gro-
vers', see Psoriasis.

ITCH INSECT, see PSORA—i. Mite, see Psora.

ITCH, VENEREAL, Scherlievo.

ITCH WEED, *Veratrum viride*.

ITCH'ING, *Parap'sia pruritus*, *Autal'gia pruriginosa*, *Pruritus*, *Pruri'go*, *Cnes'mos*, *Cnes'ma*, (Sc.) *Yuck*, *Yuke*, (F.) *Prurit*, *Démangeaison*. A sensation, more inconvenient than painful, seated especially at the surface of the body, which provokes the patient to scratch the part. It may be either an external or an internal sensation: that is, produced by an external body, or by some modification in the organic actions of the part to which it is referred.

ITEA, (1764,) *Salix*.

ITER ('a way or passage,' gen. *Itin'eria*) A TERTIO AD QUARTUM VENTRICULUM, *Aquæductus Sylvii*—i. Dentis, see Gubernaculum dentis—i. Femineum, *Perinæum*—i. ad Infundibulum, *Vulva*—i. a Palato ad Arem, *Eustachian tube*—i. ad Quartum ventriculum, *Aquæductus Sylvii*—i. ad Tertium ventriculum, *Vulva*—i. Urinæ, *Urethra*—i. Urinarium, *Urethra*.

ITHYPHAL'LUS, (ἰθυφαλλος, from ἰθυσ, 'straight,' and φαλλος, 'penis'.) An amulet, in the form of an erect penis, anciently worn round the neck, to which were attributed alexiterial properties.

ITINERARIUM, (iter, gen. *itin'eria*, 'a way or journey,') *Conductor*, *Director*.

ITIS, (ἰτις, 'bold,' 'rash,' [?].) A suffix denoting inflammation;—as *encephalitis*, inflammation of the encephalon;—*Pleuritis*, inflammation of the pleura. According to some, as Kraus, *encephalitis*, for example, was, with the ancients, the adjective agreeing with νοσος, 'a disease,' understood to express a disease of the encephalon, and it was only in modern times, that the suffix was used to denote inflammation.

IVA, *Teucrium iva*—i. *Arthritica*, *Teucrium*

chamaepitys—i. *Moschata Monspelienisium*, *Teucrium iva*—i. *Pecanga*, *Smilax sarsaparilla*.

IVETTE MUSQUÉE, (dim. of *Iva*), *Teucrium iva*—i. *Petite*, *Teucrium chamaepitys*.

Ivoire, Ivory—i. *Végétal*. *Phytelephas macrocarpa*.

I'VORY, *E'bur*, *El'ephant*, (F.) *Ivoire*. The tusk of the elephant. It is chiefly composed of phosphate of lime, and is used for the fabrication of pessaries, artificial teeth, handles of instruments, &c. Formerly, when calcined to whiteness, it entered into some pharmaceutical preparations. It was regarded as astringent and anthelmintic, and was called *Spo'dium*.

The dentine is the ivory of the human tooth. See Tooth.

IVORY BLACK, see *Carbo animalis*.

IVORY, VEGETABLE, *Phytelephas macrocarpa*.

IVRAIE, (*ebrius*, 'drunk,') *Lolium temulentum*.

IVRESSE, (same etymon), *Temulentia*—i. *Iodique*, see Iodine—i. *Quinique*, *Quinism*.

IVROGNERIE, see *Temulentia*.

IVROIE, (from *ebrius*, 'drunk,') *Lolium temulentum*.

IVY, (Sax. *1p1x*), *Hedera helix*, *Kalmia angustifolia*—i. American, *Ampelopsis quinquefolia*—i. Big, *Kalmia latifolia*—i. Fine-leaved, *Ampelopsis quinquefolia*—i. Ground, *Glechoma hederacea*.

IXIA, (ἴξις), *Varix*, *Viscum album*.

IXINE, (ἴξις, 'birdlime,') *Atractylus gummifera*.

IXIR, Elixir.

IXO'DES RIC'INUS, (ἰξωδης, 'tenacious,') *Tick*, *Woodtick*, (F.) *Tigue*. An animal parasite, which immerses its proboscis in the skin, and often excites great irritation there. It may be made to detach itself by rubbing it with oil.

IXUS, (ἴξος), *Viscum album*.

IXOMYELITIS, (ἰξος, μυελος, 'marrow,' and *itis*.) Inflammation of the spinal marrow in the lumbar region.

IX'YS, (ἴξυς), *Ix'ya*, *Ix'yē*. Used by different authors for the ilia, flanks, and loins; most frequently for the last.

J.

JABOT, (from *gibber*, 'a bunch, or swelling,' [?]) *Ingluvies*.

JACA INDICA, *Thymus mastichina*.

JACEA TRICOLOR, *Viola Tricolor*—j. Ramo-
tissima, *Centaurea calcitrapa*.

JACENS, (*jacere*, 'to lie,') *Sick*.

JACK OF THE BUTTERY, *Sedum acre*—j.
n the Hedge, *Alliaria*.

JACKET BARK, (from (F.) *Jaque*, *Jaquette*.) A jacket of flannel or cotton in which powdered bark is quilted. It is occasionally worn as an antiperiodic and tonic.

JACKSON'S BATHING SPIRITS, *Linimentum saponis compositum*.

JACOBÆA, (after St. James, [?]) *Senecio Jacobæa*.

JACOBÉE. *Senecio Jacobæa*.

JACOB'S COAT, (after Dr. Jacob, of Dublin,) *Tunica Jacobi*—j.'s Ladder, false, *Polemonium reptans*—j.'s Membrane, *Tunica Jacobi*—j.'s Ulcer, *Lupus*.

JACOBSON, VENOUS SYSTEM OF, (after Prof. Jacobson, of Copenhagen,) see Venous system.

JACOBSON'S ANASTOMOSIS, see Petrosal ganglion.

JACOLATT. Chocolate.

JACTATION, *Jactitation*, *Jacta'tio*, *Jacta'tus*, *Rhaptas'mos*, *Jactita'tio*, (*jacare*, *jactatum*, 'to toss about,') Extreme anxiety; excessive restlessness;—a symptom observed in serious diseases.

JACTITATIO, *Inquietude*, *Jactation*.

JADE NÉPHRITE ou ORIENTALE, *Nephreticus lapis*.

JAGGERY, Spirit (Arrack), see Toddy.

JAGRE, see Tari.

JALAP, (*Chala'pa* or *Xala'pa*, its native place,) *Convolvulus jalapa*, *Phytolacca decandra*—j. Blanc, *Convolvulus mechoacan*—j. Cancer root, *Phytolacca decandra*—j. Fusiform, light male, see *Convolvulus jalapa*—j. Wild, *Convolvulus panduratus*.

JALAPA, *Convolvulus jalapa*.

JALAPINE, see *Convolvulus jalapa*.

JALAPIUM, see *Convolvulus jalapa*.

JALAPPA, *Convolvulus jalapa*—j. Alba, *Convolvulus mechoacan*.

JALEYRAC, MINERAL WATERS OF. These waters are situate two leagues from Mauriac, in France, on the road from Clermont in Auvergne. The waters contain lime and carbo-

nate of soda, and are esteemed to be tonic, aperient, &c.

JALOPA, *Convolvulus jalapa*.

JAMAICA, see West Indies—j. Bark tree, *Bursera gummitera*.

JAMAICINE, see *Geoffræa inermis*.

JAMBES DE BARBADE, (pl. of [F.] *jambe*, 'leg,') see Elephantiasis—j. du Cervelet, Corpora testiformia.

JAMBIER, (from *jambe*, 'leg,') Tibial—j. Antérieur, Tibialis anticus—j. Grêle, Plantar muscle—j. Postérieur, Tibialis posticus.

JAMES'S ANALEPTIC PILLS, see Analeptica.

JAMESTOWN WEED, *Datura stramonium*.

JAMAMUNDA, *Geum urbanum*.

JANDERS, (Prov. for jaundice,) Icterus.

JANICEPS, (*Janus*, a divinity with two faces, and *caput*, 'head,') A double monster, the two bodies intimately united above a common umbilicus; and a double head, with the faces looking in opposite directions.

JANIPHA MANIHOT, *Jatropha manihot*.

JANITOR, (*janua*, 'a door,') Pylorus.

JANITRIX, (*janua*, 'a door,') Porta vena.

JANON-TARENTISME, Tarantismus.

JANUS, (the two-headed god,) Dicephalus.

JAQUIER, *Artocarpus*.

JAKRET, (low [L.] *garretum*,) Poples.

JARRETIER, *Popliteus muscle*, Hamstring.

JARRETIÈRE (F.), *Periæcelis*, a garter.

(from low Latin, *garretum* or *garretum*, 'the ham,') A kind of furfuraceous herpes, which occupies the part of the leg where the garter is worn.

JASMIN, *Jasminum officinale*.

JASMINUM ARABICUM, *Coffea Arabum*.

JASMINUM OFFICINALE, *Jasminum*, *Jesem'inum*, the *Jes'amine*, (F.) *Jasmin*. Ord. *Jasminææ*. The flowers of this beautiful plant have a very fragrant smell, and a bitter taste. They afford, by distillation, an essential oil; which is much esteemed in Italy, for rubbing paralytic limbs, and in the cure of rheumatic pains.

JASPER, *Jaspis*.

JAS'PIS, *lās'pis*, (αἰσπῆς,) *Jas'per*. A precious stone, supposed by the ancients to be capable of arresting hemorrhage when worn as an amulet.

JATAMANSI, *Sumbul*.

JATROPHA CURCASA, (αἰσπῆς, 'I heal,' and ρεσπῆς, 'I nourish,') *Ricinus major*, *Ricinus' des*, *Pineus* seu *Curcas pur'gans*—*Nat. Order*, *Euphorbiacææ*. The seed of this plant, *Pinho'nes Indici*, *Sem'ina Ricini majores*, *Fa'ba cathartica*, *Avella'na cathartica*, *Nux cathartica America'na* seu *Barbade'nsis*, *Physic' or Barba'does Nut*, (F.) *Pignon d'Inde*, *Médecinier cathartique*, *Grand haricot de Pérou*, *Pignon de Barbarie ou des Barbades*, *Noix cathartique*, *N. Américaine ou des Barbades*, affords a quantity of oil, *Oleum infernale*, *O. Ricini majores*—given, in many places, like the castor oil; to which, indeed, it is nearly allied. They contain a peculiar acid, the *Jatroph'ic* or *Igasur'ic*.

The seeds of *JATROPHA MULTIP'IDA* are called *Purging Nuts*, and give out a similar oil.

JATROPHA ELASTICA, see *Caoutchouc*.

JATROPHA MANIHOT, *Jun'ipha Man'ihot*, *Man'ihot utilis'ima*, *Bit'ter Cassa'va*: the plant affording the *Cassa'ndu* or *Cassa'va Root*, *Caca'vi*, *Cassa'ré*, *Cuzabi*, *Pain de Madagascar*, *Ric'imus Minor*, *Man'ivoc*, *Mandioc*, *Magnoc*, *Man'iot*, *Yuc'ca*, *Mani'ibac*, *Aipi*, *Aipima cozera*, *Aipipoca*, *Jau'ipha*. The juice of the root is extremely acrid and poisonous. What remains, after expressing it, is made into cakes or meal; of which the cassava or cassava bread is formed. This bread constitutes a principal food of the inhabitants of those parts where it grows.

The fecula of the root forms *Tapio'ca*, *Cipipa*, *Fec'ula Tapio'ka*, *Am'yllum manihot'icium*, which

is very nutritious. It may be prepared in the same manner as sago.

A factitious Tapioca is met with in the shops, which is in very small, smooth, spherical grains, and is supposed to be prepared from potato starch. It is sold under the name *pearl tapioca*.

JATROPHA U'RENS. A most deadly plant. The touch of its five bristly stings had nearly cost Mr. Smith, curator of Kew gardens, his life.

JAUNDERS, (Prov. for jaundice,) Icterus.

JAUNDICE, (from *jaune*,) Icterus—j. Black, Melæna—j. Blue, Cyanopathy—j. Green, Icterus viridis—j. Lead, Icterus saturninus—j. Red, Phenigmus.

JAUNE, Yellow—j. d'Œuf, see Ovum.

JAUNISSE, (from *jaune*,) Icterus.

JAW BONE, ([F.] *joue*, 'the cheek,') Maxillary Bone.

JAW DISEASE, PHOSPHORUS, see Phosphorus.

JEAN-DE-GLAINES, ST., MINERAL WATERS OF. These waters are situate two leagues from Billom in Auvergne. They contain chloride of calcium, and carbonate of magnesia; dissolved in an excess of carbonic acid.

JEANNETTE, *Narcissus pseudonarcissus*.

JECINUS, gen. *Jecin'oris*, Liver.

JÉCORAIRE, (from *jecur*,) Hepatic.

JECORAL, Hepatic.

JECORARIA, *Marchantia polymorpha*.

JECORARIUS, Hepateros, Hepatic.

JECORIS VOMICA, see Hepatitis.

JECTIGAT'IO, (*jectura*, [from *jacio*,] 'a throwing,') A word used by Van Helmont for a species of epilepsy or convulsion.

JECUR, (gen. *Jec'oris*, seu *Jecin'oris*,) Liver—j. Uterinum, Placenta.

JEFFERSONIA BARTONI, (after President Jefferson,) *J. Diphyl'la*, *Com'mon Twin'leaf*, *Yellow Root*, *Rheum'atum Root*, *Hel'met Peel*, *Ground Squi'rel Pea*. An indigenous plant, belonging to *Nat. Ord.* *Berberideæ*, *Sex. Syst.* *Octandria Monogynia*; which possesses medical properties analogous to those of hydrastis.

JEJU'NAL, *Jejuna'lis*, (from *jejunum*,) Relating or appertaining to the jejunum.

JEJUNITAS, (from *jejunus*, 'empty,') Hunger.

JEJUNT'IS, (*jejunum*, and *itis*,) Inflammation of the jejunum.

JEJUNIUM, Hunger, Fasting.

JEJUNUM, *Nestia*. The part of the small intestine comprised between the duodenum and ileum. It has been so called, because it is almost always found empty in the dead body.

JELLY, Gelly—j. of the Cord, Gelatin of Wharton—j. Rice, see Oryza—j. Seaweed, see Icinglass—j. Water, *Brasenia hydropeltis*.

JENEPERE, *Juniperus communis*.

JENKINSONIA ANTIDYSENTERICA, (after Jenkinson, an English botanist,) *Pelargonium antidy'sentericum*.

JERCON, see Arrowpoison.

JERKING RESPIRATION, Respiration, jerking.

JERKS, see Mania, dancing.

JEROFFLERIS, (July flower, or from *caryo phyllum*,) *Dianthus caryophyllus*.

JERSEY, CLIMATE OF. Jersey is the largest of the islands of the British Channel, and is most frequented by invalids. Its climate closely resembles that of the southwest coast of England, and especially of Penzance, and it is adapted to the same class of invalids.

JERUSALEM OAK OF AMERICA, *Chenopodium anthelminticum*.

JESEMINUM, *Jasminum officinale*.

JESSAMINE, *Jasminum officinale*—j. Yellow, *Gelsemium nitidum*.

JETICUCU, *Batata de purga*.

JEUÑE, (from *jejunus*, 'empty.') Fasting.

JEUNESSE, (from *juvenis*, 'a young person.')

Adolescence.

JEWBUSH, *Pedilanthus tithymaloides*.

JEWEL WEED, *Impatiens balsamina*.

JEW'S EAR, *Peziza auricula*—*j.* Harp, *Trilium latifolium*.

JIGGER, *Chique*.

JIMPSON WEED, (corr. from Jamestown,) *Datura stramonium*.

JIMSTON WEED, *Datura stramonium*.

JIZZEN BED, (Sc. from old *F. gésine*, 'in childbed.')

See Parturition.

JOANNESIA PRINCEPS, *Anda*.

JOANNETTE, MINERAL WATERS OF.

Several springs are found at Joannette, about five leagues from Angers, in France. The waters are both cold and warm, and contain sulphate of lime; subcarbonate of soda; chloride of sodium; some iron; chloride of calcium, and, sometimes, a kind of saponaceous matter. They are chiefly used as a tonic in chlorosis, leucorrhœa, &c.

JOE PYE, *Eupatorium perfoliatum*, and *E. purpureum*.

JOHNE, MINERAL WATERS OF. *Johne* is a village near Dol in Franche-Comté, France, where is a mineral spring, which contains subcarbonate of soda, and subcarbonates of magnesia and iron. It is used chiefly as a bath in certain diseases of the skin, and in atonic affections.

JOHNNY JUMPER, *Viola tricolor*.

JOINT, ([*F.*] *joint*, 'joined,' from *jungere*, *junctum*, 'to join.')

Articulation—*j.* Ball and Socket,

Enarthrosis—*j.* Dovetail, Suture—*j.* Stiff, Ankylosis.

JOINTURE, Articulation.

JONAS, ([*Prov.*] corr. from *jaunisse*,) *Icterus*.

Jo'NAS, MINERAL WATERS OF. This spring is situate to the south-west of Bourbon-l'Archambault, in France. The waters contain chloride of calcium and sulphate of lime; chloride of sodium, and sulphate of soda; carbonate of iron, and carbonic acid gas. They are tonic and aperient.

JORDAN'S WHITE SULPH'UR SPRINGS. These are in Virginia, five or six miles from Winchester. They are used in the same cases as sulphureous springs in general.

JOT-GUT (*Prov.*), Rectum.

JOUAN, ST. MINERAL WATERS OF. *St. Jouan* is a village, near Saint-Malo, in France. The water is a cold chalybeate.

JOUBARBE ACRE, (*Jovis barba*, 'beard of Jupiter.')

Sedum—*j. des Toits*, *Sempervivum tectorum*—*j. des Vignes*, *Sedum telephium*.

JOUE, *Gena*.

JOULUS, (*ιουλος*,) *Julus*.

JOUR, (*dies*, *diurnus*, [*I.*] *giorno*,) *Dies*.

JOURS CANICULAIRES, Dog days—*j. Critiques*, Critical days—*j. Indicateurs*, Indicating days—*j. Paroxysmiques*, see *Paroxysmie*.

JOVIS FLOS ('flower of Jove'), *Crocus*—*j.* Glans, *Juglans*.

JUCATO CALLELOE, *Phytolacca decandra*.

JUDÆ'US, *Judaicus* seu *Phœniciæ La'pis*, *Tecol'ithos*, (*F.*) *Pierre Judæique*. A stone, found in Judæa, Palestine, &c. Called, also, *Lapis Syriacus*. It was formerly esteemed to be diuretic and lithontriptic.

JUDG'MENT, *Judic'ium*, (*F.*) *Jugement*, (from *judicare*, 'to judge.')

The faculty of the intellect,

by which ideas are compared with each other, and their relative worth appreciated.

JUDICATORII DIES, Critical days.

JUDICIUM, (from *judex*, gen. *ju'dicis*, 'a judge.')

Judgment.

JUGA ALVEOLARIA, (pl. of *jugum*, 'a yoke,')

see Alveolar arches.

JU'GA CEREBRA'LIA. Prominences on the inner

surface of the bones of the cranium, between the digital impressions.

JUGAL, *Zygomatic*—*j.* Process, *Zygomatic* process.

JU'GAL RE'GION, *Re'gio Juga'lis*. The region of the cheek-bone.

JUGALE OS, *Malæ os*.

JUGALIS, *Zygomatic*.

JUGA' LIS SUTU'RA. The suture which unites the *Os Malæ* or *Os Juga'le* with the superior maxillary bone. Also, the sagittal suture.

JUGEMENT, Judgment.

JUGEOLINE, *Sesamum orientale*.

JUGLANS, gen. *Juglan'dis*, (*Ju-glans*, quasi *Jovis glans*,) *J. cinerea*, *J. regia*—*j.* Black, *Juglans nigra*.

JU'GLANS CINER'E'A, *But'ter* or *Oil Nut*, *White Wal'nut*. *Ord.* *Juglandacæ*. The inner bark of the root of this tree, *Juglans* (*Ph. U. S.*), which is abundant in the United States, is used in medicine, in the form of extract, as an efficacious and mild laxative, in doses of from 10 to 20 grains. It is in the secondary list of the Pharmacopœia of the United States.

JUGLANS NI'GRA, *Black Juglans*, *Black Walnut*; indigenous. The rind of the unripe fruit is said to remove ringworm and tetter. A decoction has been given as an anthelmintic.

JUGLANS RE'GIA, *Juglans*, *Nux Juglans*, *Ca'rya basil'ica*, *Carya*. The *Wal'nut*, *French nut*, (*F.*) *Noyer*, the tree; *Noir*, the nut, *Car'yon*. The unripe fruit, in the state in which it is pickled, was formerly esteemed to be anthelmintic. The *putamen* or green rind of the walnut (*F.*) *Brou de Noir*, has been celebrated as a powerful antisyphilitic, and used as a sort of diet drink. The *kernel* is an agreeable article of dessert; but, like all nuts, is difficult of digestion. The expressed oil, *Huile de Noir*, is used in France as an aliment, and, like other fixed oils, is laxative.

Various preparations of the leaves have been recommended in scrofulous affections.

JUG'ULAR, *Jugula'ris*, (*F.*) *Jugulaire*, (*jugulum*, (dim. of *jugum*, because the 'yoke' was attached there,)) 'the throat.')

JUGULAR FOS'SA, *Fos'sa Jugula'ris*, is a cavity in the *petrocephical* suture. It is formed by the petrous portion of the temporal bone, and by the occipital bone, and lodges the origin of the internal jugular vein. It is, sometimes, called *thimble-like cavity*.

JUGULAR VEINS, *Ve'næ Jugula'res* seu *sphagit'ides* seu *apoplecticæ* seu *sopora'les*. These are situate at the lateral and anterior parts of the neck. They are two on each side; one *external*, the other *internal*. 1. The *Extern'al Jugular Vein*, *Trachêlo-sous-cutané*, (*Ch.*) is of less size than the internal. It descends, almost vertically, along the anterior and lateral part of the neck, from the cervix of the lower jaw to the subclavian vein, into which it opens, a little above the internal jugular. It is formed by the *internal maxillary*, *superficial temporal*, and *posterior auricular veins*. It is this vein which is commonly opened in bleeding in the neck. 2. The *Intern'al Jugular Vein*, *V. Céphalique* (*Ch.*), *Vena apoplectica*, is much larger, and more deeply seated than the preceding. It descends, vertically, along the anterior and lateral part of the neck, from the posterior part of the foramen lacerum posterius as far as the subclavian vein. It commences at the *sinus* of the jugular vein, (*F.*) *Golfé de la veine jugulaire*, and receives the blood, which returns by the *sinus* of the *dura mater*, and that of the *facial*, *lingual*, *pharyngeal*, *superior thyroid*, *occipital*, and *diploic veins*.

JUGULATION, *Jugula'tio*, (from *jugulare*, *jugulatum*, 'to strangle.')

The sudden cutting short of a disease by remedial agents.

JUGULUM, (dim. of *jugum*, 'a yoke.') Clavicle, Throat—*j.* Uteri, Collum uteri.

JUGUM PENIS, ('yoke of the penis,') *Prepuce* *urethre*.

JUICE, EXPRESSED, see *Succus*—*j.* Musclee, see *Muscle juice*.

JUJUBE, *Ju'juba*, *Zizyphum*, *Bac'ca ju'jubæ*, *B. Zizyphi*. (Of Arabic origin.) The fruit of *Rhamnus Zizyphus* seu *Zizyphus vulgaria*. *Ord.* *Rhamnæ*. A native of the south of Europe. It was formerly ranked amongst the pectoral fruits. It has an agreeable sweet taste. The fruits of two other species of *Zizyphus*—*Z. jujuba*, *Rhamnus jujuba*, *Mons'na arbo'rea*, a native of the East Indies, and *Z. lotus*, *Rhamnus lotus*, *Zizyphus nitida* seu *sativ'a* seu *sylnes'tris*, growing in North Africa, possess properties similar to those of *Zizyphus vulgaris*.

JUJUBE Paste, Paste of jujubes.

JULAPIUM, *Julep*—*j.* Camphoræ, *Mistura Camphoræ*.

JULEB, *Julep*.

JULEP, *Jula'pium*, *Jule'pus*, *Zula'pium*, *Juleb* of the Persians. A sweet drink. A demulcent, acidulous or mucilaginous mixture.

JULEP, CAMPHOR, *Mistura camphoræ*—*j.* *Camphoræ*, *Mistura camphoræ*—*j.* Chalk, *Mistura cretæ*—*j.* Mint, see *Mint julep*.

JULUS (*ουλος*), *Iulus*. The first down that appears upon the chin.

JUM, (*Prov.*) *Lolium temulentum*.

JUMBLE BEADS, see *Abrus precatorius*.

JUMEAU, ('a twin,') *Gemellus*.

JUMEAUX DE LA JAMBE, *Gastrocnemii*.

JUMELLE, ('a twin,') *Gemellus*.

JUMENTOUS URINE, (from *jumentum*, 'a beast of burthen,') *Urine jumentense*.

JUMPERS, see *Convulsionnaire*.

JUNCTURA, (*jungere*, *junctum*, 'to join,') *Articulation*.

JUN'CUS ODORAT'US, (*jungere*, *junctum*, 'to join,' in allusion to the use of rushes for bands. [?]) *Andropogon schænan'thus* seu *bicor'nis* seu *citrat'us* seu *citriodor'us*, *Cymbopogon schænan'thus*, *Fœnum camelo'rum*, *Junc'us aromat'icus*, *Schænan'thus*, *Schænan'thum*, *Cal'amus odorat'us*, *Squinan'thus*, *Gra'men orienta'le*, *Cam'el's hay*, *Sweet rush*, (F.) *Junc odorant*, *Foin de chameau*. *Ord.* *Graminæ*. The dried plant, which is generally procured from Turkey and Arabia, has an agreeable smell, and a warm, bitterish taste. It was formerly used as a stomachic and deobstruent.

JUNGLE FEVER. A variety of remittent occurring in the jungle districts of India.

JUNIPER TREE, *Juniperus communis*.

JUNIPERUM VINUM. Wine impregnated with juniper berries.

JUNIPERUS COMMUNIS, *J. Suec'ica*, *Arcent'hus*, *Ju'niper tree*. (Sc.) *Jenepere*. Family, *Conifere*. *Sex. Syst.* *Dioecia Monadelphia*. The berries or fruit, *Juniper'us* (Ph. U. S.), *Akat'alis*, *Akat'era*, (Sc.) *Eat'in ber'ries*, *El'nagh ber'ries*, are ordered in the pharmacopœias. Their odour is strong, but not unpleasant. Taste warm, pungent. Properties dependent upon essential oil, which they yield to both water and alcohol. Dose, ʒj to ʒss, rubbed with sugar. In Holland, juniper berries are used for flavouring gin. The oil, *O'leum juniperi*, possesses the virtues of the plant. It is called by *Ruland*, *Targac*. (F.) *Genévrier* (the plant), *Genèvre* (the berry).

JUNIPERUS LYCIA. This plant—*Thu'rea*, *Th. vir'ga*, *Ar'bor thurif'era*—has been supposed to afford the true *Frank'incense*, *Cedrus*, *Olib'anum*, *O. Arab'icum*, *Thus Libani'tas*, *Libano'tum*, *Lib'anos*, *Thus mas'culum*, *Thus ve'rum*, (F.) *Encens*. By some, however, it is supposed to be the produce of an *Am'yris*, and by others of *Boswell'ia floribunda*, *B. Papyrif'era*, *Ploes'sea floribunda*. The odour of *olibanum* is peculiar and aromatic;

taste bitterish and slightly pungent; partly soluble in alcohol, and forming a milky emulsion when triturated with water. It was formerly used in dysentery and hæmoptysis, but is now never employed except as a perfume in a sick room.

Boswell'ia serra'ta, *B. thurif'era* of India, Persia and Arabia, is said to yield *East India Olib'anum*, *O. Indicum*.

JUNIPERUS OXYCEDRUS, (F.) *Genévrier Oxycedre*, *Cade*, *Oricède*, *Oxycedre*, grows in the south of Europe, Siberia, &c. By combustion of the wood, a liquid tar, *O'leum ca'dinum*, *Juniperi o'leum empyreumat'icum*, *Oil of Cade*, (F.) *Huile de Cade ou de Genévrier*, is obtained, which is employed externally in various chronic cutaneous and other diseases.

JUNIPERUS SABI'NA, *Sabi'na*, *S. ster'ilis*, *Savi'na*, *Brut'a*, *Cedrus baccif'era*, *Com'mon or bar'ren savin*; *Brat'hu*, *Bar'rathron*, *Bo'rathron*, (F.) *Sabine*, *Savinier*. The odour of *savin tops*, *Sabi'na* of the pharmacopœias, is strong and disagreeable. Taste hot, acrid, and bitter, depending on an essential oil. Their operation is stimulant, emmenagogue, anthelmintic, and abortive; externally irritant. Dose, internally, gr. v to x of the powder. As a local irritant, they are applied in powder or formed into a cerate. The essential oil, *O'leum Sabi'næ* (Ph. U. S.), has the virtues of the *savine*. Dose, two to five drops.

JUNIPERUS SUECICA, *J. communis*.

JUNIPERUS VIRGINIA'NA, *Red ced'ar*. This tree is known throughout the United States by the name of *sav'ine*, and is often used for the same purposes. The tops are official in the Pharmacopœia of the United States.

JUNK, (from *juncus*, 'a bulrush,') *Pad*.

JUNO LUCINA, (Goddess of childbirth,) *Ilithyia*.

JUNO'S TEARS, *Verbena officinalis*.

JUPITER, (*Jus*, and *pater*, 'father,') *Tin*.

JUPITER'S BEARD, *Sempervivum tectorum*.

JURIBALI, *Euribali*. A tree in the forests of Pomeronia. Fam. *Meliaceæ*. Class, *Octandria*. Nat. *Ord.* *Monogynia*. The bark is febrifuge, and may be given in powder or infusion. (ʒj) ad aq. bullient. Oij.)

JURISPRUDENCE, MEDICAL, (*jus*, gen. *juris*, 'law,' and *prudencia*, 'knowledge,') *Jurisprudent'ia Med'ica*, (F.) *Jurisprudence Médicale*. This word is often used synonymously with *Legal Medicine*. It is at times employed for the embodied laws and regulations that relate to the teaching and practice of medicine.

JURY OF MATRONS. A jury formed of women empanelled under a writ of *ventre inspiciendo*, to try the question, whether a woman is with child or not.

JUS, gen. *Juris*, *Zomos*, *Zomid'ium*, *Sorbit'io*, *Sorbit'um*. Soup, broth, *Jusculum* or *bouillon*. See *Broth*.

JUS, Succus.

JES BOVINUM, *Beeftea*—*j.* *Coagulum*, *Gelly*.

JUS D'HERBES (F.). The juice of certain vegetables administered as depuratives; as that of fumitory, burdock, water trefoil, &c.

JUS JELATUM, *Gelly*—*j.* de *Régliose*, see *Glycyrrhiza*—*j.* de *Viande*, see *Bouillon*—*j.* de *Viande Crue*, *Juice of raw meat*.

JUSCULUM, (dim. of *jus*,) *Broth*, see *Jus*.

JUS'CULUM NIGRUM LACEDÆMONI'UM, *Me'las Zo'mos*, (*μελας ζωπος*,) *Hama'tia*, *Black broth of the Lacedæmonians*. A celebrated mess in antiquity, the ingredients of which are not known. It was supposed to be made of blood—hog's blood,—treated in some way, and, therefore, resembling perhaps—as suggested by Warner—the modern black pudding.

JUSÉE (F.) An aqueous *maceratum* of tan, coarsely powdered, in which the skins of animals have been steeped for a time, in the preparation of leather. An extract has been prepared from this, which has been prescribed in syrup and in pill; each pill containing about a grain and a half of the extract; and two being taken daily in phthisis, and in general debility connected with scrofula and rickets.

JUSQUIAMINE, Hyoscyamus.

JUSQUIAMUS, Hyoscyamus.

JUSQUITAME, Hyoscyamus.

JUSTICIA ADHATODA, (after J. Justice, a Scotch botanist,) Adhatoda—j. Bivalvis, Adulasso.

JUSTICEA ECBOLITUM, *Carim curini*. A Malabar plant, the root of which, and the leaves, in decoction, are considered in the country to be lithontriptic,

JUSTICIA PECTORALIS. A West India plant, which is slightly astringent. (F.) *Carmentine*, *C. Pectorale*, *Herbz aux Charpentiers*.

JUVANS, pl. *Juvan'tia*, (*juvare*, 'to aid,') *Auxilium*, *Remedium*. A medicine or substance of any kind, which relieves a disorder. An Adjuvant.

JUVENIS, Adolescens.

JUVENTA, Adolescence, Hebe.

JUVENTAS, Adolescence, Hebe.

JUVENTUS, Adolescence, Hebe.

JUNTAPOSIT'ION, *Junctaposit'io*, (*juncta*, 'near to,' and *ponere*, *positum*, 'to place,') The mode of increase proper to minerals; which consists in the successive application of new molecules upon those that form the primitive nucleus. It is opposed to *intussusception*.

K.

Most of the terms under K, derived from the Greek, are found under the letter C.

K. This letter was formerly used to designate a compound of gold.

KAATH, Catechu.

KAAWY. Ancient name of an Indian drink, prepared from maize.

KABALA, Cabal.

KÆMPFERIA GALANGA, (after Kæmpfer, the German naturalist,) see *Maranta galanga*.

KÆMPFERIA ROTUNDA. *Zedoaria*, *Z. rotunda*, *Amomum zedoaria*, *Colchicum Zeylanicum*, *Curcuma aromatica*, *Zedoary*, (F.) *Zédoaire rond*. Ord. Zingiberaceæ. The roots of this Ceylonese plant have a fragrant smell, and are stimulant and carminative. Dose, ʒj to ʒj of the powder.

According to some, the round *zedoary* is furnished by *Curcuma zerbumbet*; the long, *Zedoaria longa*, by *Curcuma zedoaria*.

KAHINCE RADIX, *Caincæ radix*.

KAIB, *Kayl*. A word employed by the alchemists for sour and coagulated milk.

KAJEPUT, Cajeput.

KAKERLAKISM, (D.) *Kakkerlak*, 'a cockroach,' which comes out only in the dark.) See Albino.

KAKOCHYMIA, Cacoehymia.

KALI, Potash, Potassa—k. Acetas, Potassæ acetas—k. Aeratum, Potassæ carbonas—k. Bichromicum, Potassæ bichromas—k. Causticum, Potassa fusa—k. Causticum cum calce, Potassa cum calce—k. Chloricum, Potassæ murias hyperoxygenatus—k. Chromicum acidum, Potassæ bichromas—k. Chromicum flavum, Potassæ bichromas—k. Chromicum neutrale, Potassæ chromas—k. Chromicum rubrum, Potassæ bichromas—k. Hydriodicum, see Potassæ hydriodas—k. Hypermanganicum, Potassæ permanganas—k. Inermis, see Salsola kali—k. Nitricum, Potassæ nitras—k. Oxalicum acidulum, Potassa, oxalate of—k. Præparatum, Potassæ subcarbonas—k. Præparatum e tartaro, see Potash—k. Purum, Potassa rusa—k. Soda, see Salsola kali—k. Spinosum cochleatum, Salsola kali—k. Subcarbonas, Potassæ subcarbonas—k. Sulphas, Potassæ sulphas—k. Sulphuricum, Potassæ sulphas—k. Sulphuretum, Potassæ sulphuretum—k. Tartarizatum, Potassæ 'atras—k. e Tartaro, see Potash—k. Vitriolatum, Potassæ sulphas.

KALICUM HYDRAS, Potassa fusa.

KALIUM IODATUM, see Potassæ hydriodas—k. Iodatum Hydrargyrum, Potassii hydrargyro-iodidum.

KAL'MIA ANGUSTIFOLIA, (after Peter Kalm, the Swedish botanist.) *Pey*, *Narrowleaved Kalmia* or *Laur'el*, *Dwarf Laurel*, *Sheep Laurel*. Ord. Ericaceæ. This plant has the same virtues as *K. latifolia*. So also have *KAL'MIA CUNEATA* and *KAL'MIA HIRSU'TA*.

KAL'MIA, BROAD LEAVED, *K. latifolia*.

KAL'MIA GLAUCA, *Sicamp Laurel*, *Pale Laurel*. **KAL'MIA LATIFOLIA**, *Broadleaved Kalmia*, *Calico bush*, *Laurel*, *Mountain or Rose* or *Broadleaved Laurel*, *Big Pey*, *Spoonwood*, *Catskill*, *Lamb'kill*, *Sheep-poison*. (F.) *Laurier des Montagnes*. This plant kills sheep and other animals. The Indians use it as a poison. The powdered leaves have been applied successfully in tinea capitis: and a decoction of it has been used for the itch. The powder, mixed with lard, has been applied in herpes.

KAL'MIA, NARROWLEAVED, *K. angustifolia*.

KALO, Arum esculentum.

KAMAS ROOT, Camassia esculenta.

KAMEELA, see *Rottleria tinctoria*.

KAMPHUR, Camphor.

KANKROID, Canceroid.

KAPHUR, Camphor.

KARABE, Asphaltum.

KARABITUS, Phrenitis.

KARAPAT, Ricinus communis, (oil of.)

KARCINOSEN, Carcinoses.

KARENA, Carena.

KARKINOMA, Carcinoma.

KARSON-KHAYI, Swietenia Senegalensis.

KASSADER, Convolvulus panluratus.

KASSAUDE, Convolvulus panduratus.

KATASARCA, (*cata*, and *σαρξ*, gen. *σαρκος*, 'flesh,') Anasarca.

KATASTALATICA, (*κατασταλτικά*, from *cata*, and *στέλλειν*, 'to repress,') Astringents.

KAVA, Ava.

KAVIAC, Caviare.

KAWING, Anhelatio.

KAYL, Kaib.

KECKCORN (Prov.), Trachea.

KECKER, Trachea.

KEENBITE, see Congelation.

KEER, *Sorbus aucuparia*.
 KEITA, *Monsonia ovata*.
 KEKUNE OIL, see *Aleurites triloba*.
 KELLS, (properly *Chelis*, from *χελη*, 'the claw of a crab.') See *Canceroid*.
 KELING (Prov.), (by onomatopœia,) *Eru-tation*.
 KELOID, (properly *Cheloid*, from *χελη*, 'the claw of a crab,' and *ειδος*, 'resemblance,') *Canceroid*.
 KELOTOMIA, *Celotomia*.
 KELP, *Varec*. The impure mineral alkali obtained by burning certain marine plants. See *Soda*.
 KEME, see *Cucurbita citrullus*.
 KENCH (Prov.), *Sprain*.
 KENGIVE, see *Cucurbita citrullus*.
 KENNELWORT, *Scrophularia nodosa*.
 KENTUCK'Y, MINERAL WATERS OF. The *Olympian Springs*, *Mudlick*, in this State, are near the sources of Licking river. There are three different kinds, in the space of half a mile. One of them is saline, impregnated with sulphur;—another is chalybeate, and a third a sulphureous "ring. In various parts of Kentucky, there are saline waters, which are frequented by invalids. The *Salines*' at Bigbone, formerly employed in the manufacture of salt, are resorted to. A spring, near Harrodsburg, in Mercer County, is strongly impregnated with sulphate of magnesia, and is much visited. At the *Blue Licks*, on both banks of the Licking river, there are sulphureous saline waters.
 KERATIASIS, *Ceratiasis*.
 KERATITIS, *Ceratitis*.
 KERATOCELE, *Ceratocele*.
 KERATODEITIS, *Ceratitis*.
 KERATOICRICOID, *Ceratoicricoid*.
 KERATOGLLOSSUS, *Ceratoglossus*.
 KERATOIRITIS, see *Aquomembranitis*.
 KERATOMALACIA, *Cerato-malacia*.
 KERATONYXIS, *Ceratonyxis*.
 KERATOPHARYNGEUS, *Ceratopharyngeus*.
 KERATOPLASTY, *Ceratoplastice*.
 KERATOSTAPHYLINUS, *Ceratostaphylinus*.
 KERATOTOMUS, *Ceratotomus*, Knife, cataract.
 KERECTOMY, *Ceratotomy*.
 KERION, (*κηριον*, 'a honeycomb,') see *Favosus*.
 KER'MES, *Cher'mes*, *Alker'mes*. One of the species of the genus *kermes* lives on a green oak, and is called *Coc'cus ilicis*, *Kermes an'imal*, *Coc'ci gra'num*, *Coc'cum*, *C. baph'icum seu infec'torium seu tinct'orum seu scarlat'um*, &c. The oak, to which allusion has been made, is known by botanists under the name *Quercus coccif'era*, and grows abundantly in the uncultivated lands of southern France, Spain, and in the islands of the Grecian Archipelago. The *kermes* inhabiting it has the appearance of a small, spherical, inanimate shell. Its colour is reddish-brown, and it is covered with a slightly ashecoloured dust. This is the *kermes* of the shops. It is now only used in dyeing; but was formerly reputed to possess aphrodisiac, analeptic, antiabortive, and other virtues.
 KERMES MINERAL, *Antimonii sulphuretum præcipitatum* and *A. sulphuretum rubrum*.
 KERNEL. (Sax. *kerne*l,) *Tuberele*.
 KERNELS, WAXING, see Waxing kernels.
 KERNMILK, ((Se. and Prov.) 'churnmilk,') Buttermilk.
 KER'OSOLENE. A liquid product of the destructive distillation of coal, of very low specific gravity, say '6346. It has anæsthetic properties, but is not easily managed.
 KERVA, *Ricinus communis*.
 KESLOP, (Prov. from Sax. *cepe*lob, 'curdled milk,') *Rennet*.
 KETCH'UP, *Catch'up*, *Cat'sup*, (the *Kitjap* of the Chinese [?]). A pickle prepared from the liquor of the mushroom, walnut, tomato, &c.

KETMIA ÆGYPTIACA, *Hibiscus abelmoschus*.

KEY, (Sax. *cæg*,) *Clav'is*, *Cleis*, *Clav'is Anglica*, (F.) *Clef*; *Clef de Garangeot*. An instrument, used for extracting teeth. It consists of a firm handle, with a claw at right angles to it, and moving upon a pivot. This claw embraces the tooth. It has undergone several modifications, and hence various instruments are used under this denomination. The French have the *Clef à pompe*, *Clef à pivot*, and *Clef à noix*.

KEYRI CHEIRI, *Cheiranthus cheiri*.

KHALA MIMUC, Bit noben.

KHAYA SENEGALENSIS, *Swietenia Senegalensis*.

KHUS-KHUS, *Andropogon muricatus*.

KIASTER, properly *Chias'ter*, (*χιασιν*, 'to cross,') A species of bandage, having the form of the Greek letter *χ*, which the ancients used for approximating, and maintaining in contact, the fragments of the patella, in cases of fracture of that bone. It was applied in the form of the figure 8.

KIBE, *Chilblain*.

KIBISITOME, (*κιβισις*, 'a sac or wallet,' and *τεμνω*, 'to cut,') *Cystitome*.

KIDNEY, (etymology uncertain.) *Serenius* derives it fancifully from *Su. Goth.* *qued*, the belly; and *nigh*, (quasi, *quidney*.) *Ren*, *Neph'ros*, (Se.) *Neir*, *Nere*, (F.) *Rein*. The kidneys or *reins* are the secretory organs of the urine. They are two glands, situate deeply,—the one on the right, and the other on the left side—in the hypochondres: at the sides of the lumbar vertebra; behind the peritoneum; and in the midst of an abundant, fatty areolar tissue, *Tu'nica seu Cap'sula adip'osa*. The kidney is of a reddishbrown colour; oval form; and flattened on two surfaces. It has, at its internal margin, a deep fissure, by which the renal vessels and nerves enter or quit the organ, and the ureter issues. It resembles, pretty accurately, the haricot or kidney-bean. Two substances are readily distinguishable in it;—the outer, *secern'ing*, *cortical*, *gland'ular* or *vascular*, *Substan'tia cortical'is seu glandulo'sa*, which secretes the urine; and the inner, *tub'ular*, *med'ullary*, *urinif'erous*, *conoid'al* or *rad'iated*, *Substan'tia Medulla'ris seu Tubulo'sa seu Fibro'sa*, which appears under the form of small cones or unequal *papillæ* or *mammillæ*, each resulting from the union of small capillary tubes, adherent by one of their extremities to the cortical substance; and opening, by the other, at the summit of the cone, into *calices*, a species of membranous tubes, more or less numerous, which transmit the urine of the *papillæ* to the *pelvis*. By the *pelvis* is meant a small, membranous sac, of an irregularly oval shape, at the base of which are the orifices of the calices, and the other extremity of which is continuous with the ureter. The kidney is surrounded by a fibrous membrane proper to it, *Perineph'rus*, *Tu'nica pro'pria seu albugin'ea*. It has been shown by Mr. Bowman and others that the renal artery is distributed to the corpora Malpighiana, forming a pellet of convolutions, which is received into a flask-like dilatation of the ureter—*Bow-man's Cap'sule*, and through which—it is conceived—the watery portion of the urine is separated. The blood then becomes venous, and is distributed by different veins—*portal veins* of the kidney—to the convoluted tubes through which the proper urine is secreted. Hence the blood passes into the renal vein. The intermediate vessels between the Malpighian bodies and the convoluted tubes have been termed the *Por'tal Sys'tem of the Kid'ney*.

KIDNEYBEAN, *Phaseolus vulgaris*.

KIDNEY, BRIGHT'S DISEASE OF. The *Mor'bua Bright'ii seu albumin'en'sis seu granulo'sus Re-num*, *Nephri'tis albumino'sa seu sociat'a*, *Neph'*

ria, *Nephrochylazo'sis*, *Nephrococco'sis*, *Uroze'mia albuminosa*, *Cachectic* or *Associated nephritis*, *Inflammation of the Malpighian corpuscles*, *Disease of Bright*, *Granular Degeneration* or *Disorganization of the Kidney*, *Granular Kidney of Bright*, *Cachectic nephritis*, *Chalazonephritis*, (F.) *Maladie de Bright*, *Néphrite albumineuse*, *État Granuleux* ou *Dégénérescence granulée du Rein*, *Albuminorrhée*. A granular disease of the cortical part of the kidney, characterized by the secretion of urine that contains albumen, and is of less specific gravity than natural, and which destroys by inducing other diseases. It was first described by Dr. Bright, of London.

KIDNEY, GRANULAR, OF BRIGHT, Kidney, Bright's disease of the.

KIDNEY, MOVABLE. A condition of the kidney, usually congenital, in which the renal vessels are so elongated as to permit the kidney to be moved in certain directions. The tumour, formed by it, and felt on pressure, may readily be mistaken for disease of other parts.

KI'ESTEÏNE, properly *Ky'esteïne*, *Ki'esteïn*, *Ky'estin* or *Ky'estein*, *Kystein*, *Ki'estin*, *Ki'estine*, *Cy'esteine* or *Cy'estein*, *Kiesthein*, (F.) *Kyestéine*, (*κίαιν*, 'to be pregnant,' and *εσθς*, 'a garment or pellice.') A peculiar albuminoid pellicle, which forms on the urine of a pregnant female when it is allowed to stand for a few days. It is whitish, opalescent, slightly granular, and may be compared to the fatty substance that swims on the surface of soups, after they have been allowed to cool. When taken in conjunction with other phenomena, it is a valuable aid in the diagnosis of early pregnancy.

KIESTIN'IC, *Kiestin'icns*. Relating or appertaining to kiestine; as, 'a kiestinic urine.'

KIL'BURN, MIN'ERAL WA'TERS OF. These springs contain carbonic acid, sulphohydric acid; carbonates of lime, magnesia, and iron; sulphates of soda, lime, and magnesia, and chloride of sodium.

KILE, Ulcer.

KILL LAMB, *Andromeda mariana*.

KIL'GRAMME, *Chiliogram'ma*, (*χίλια*, 'a thousand,' and *γραμμα*, 'a gramme.') The weight of a thousand grammes;—two pounds, eight ounces, one drachm, and twenty-four grains, Troy.

KILOLITRE, (*χίλια*, 'a thousand,' and *λίτρα*, 'a litre.') A measure containing a thousand litres.

KINA KINA (Peruvian), *Cinchona*.

KINANCIE, *Cynanche*.

KINCOUGH, (by onomatopœia, from *kin*, Teut. *kink*.) Pertussis.

KINDCOUGH, Pertussis.

KINESIP'ATHY, *Kinesitherapi'a*, *Cinesip'athy*, Movement cure, *Mo'torpathy*, *Medici'na mechanica*, (*κίνησις*, 'motion,' and *παθος*, 'disease.') A mode of treating disease by gymnastics or appropriate movements, proposed by Ling, a Swede, and hence called *Ling'iam*.

KINETIC, (*κινεω*, 'to move,') Motory.

KINGCURE, *Pyrola maculata*.

KING'S EVIL, Scrofula.

KINGSHOOD, Reticulum.

KINICI ACETAS, Quinia, acetate of.

KININUM, Quinine.

KINIUM, Quinine.

KINK, (Teut. and So., by onomatopœia.) *Quinte*.

KINK IN THE HEAD, (Teut. *Kink*, 'a twist in a rope,') see *Insanity*.

KINKHAUST, ([G.] *Kink* and *Husten*, 'a cough,') Pertussis.

KINKHOST, Pertussis.

KINK'INA (Peruv. *K'ina*), *Cinchona*.

KINNIKIN'NICK. An Indian term for a

mixture of tobacco, leaves of the sumach and twigs of the willow, which is used as tobacco.

KINO, (Ind.) *Gum'mi Gambien'sè* seu *rubrum adstringens Gambien'sè* seu *adstringens Pothergill'i*, *African*, *East India* or *Ambony'na kino*, (F.) *Gomme*, *suc* ou *extrait de kino*. The trees, whence one variety of this resin is obtained, are not botanically ascertained. The London college ascribes it to *Pterocarpus erinacea*; the Edinburgh to *Eucalyptus resinifera* seu *gummifera*, *Trou bark*; and the Dublin to *Butea frondosa*. The Pharmacopœia of the United States, (1842), defines it to be "an extract obtained from an uncertain plant." The later editions, and the British Pharmacopœia, (1864,) state it to be the inspissated juice of *Pterocarpus marsupium* (De Candolle) and of other plants. On wounding the bark, the kino flows drop by drop. A West India variety is said to be derived from *Coccoloba uvifera* or *Seaside Grape*; and a *Bo'tany Bay kino* is said to be the concrete juice of *Eucalyptus resinifera* or *broken gumtree* of New Holland. *Sex. Syst.* *Icosandria Monogynia*. *Nat. Ord.* *Myrtaceæ*. Kino consists chiefly of tannic and gallic acids, oxide of iron, and colouring matter. It is inodorous; the taste a sweetish bitter; and it is sometimes gritty between the teeth. It comes to us in fragments of a dark ruby red colour, and is easily pulverized. Its properties are powerfully astringent. Dose, gr. x to gr. xx in powder.

KIONORRHAPHIA, (*κίον*, 'uvula, and *ρᾶψη*, 'suture,') *Staphylorrhaphy*.

KI'OTOME, *Ki'o'tomis*, (*κίον*, 'a pillar,' 'a support,' and *τεμνω*, 'to cut,') (F.) *Coupebride*. An instrument invented by Desault to cut any accidental brides or filaments in the rectum and bladder; and which he afterwards used for the removal of the tonsils. It is composed of a flat, silver sheath, open at one edge. This sheath is provided with a cutting blade, which can be forced through the opening, and thus all the parts can be divided with which it comes in contact.

KIPPERNUT, *Bunium bulbocastanum*.

KIRATE. A weight of four grains, according to Blancard.

KIRIAGHUMA, *Gymneura lactiferum*.

KIRKLAND'S NEUTRAL CERATE, see *Cerate*, *Kirkland's Neutral*.

KIRNMILK, ([S. and Prov.] 'churnmilk,') *Buttermilk*.

KIRRONOSE, *Cirrhosis*.

KIRRHONOSIS, (from *κίρρος*, 'yellow,') *Cirrhosis*.

KIRRHOSIS, *Cirrhosis*.

KIRSCHWASSER, ([G.] *Kirsch* and *Wasser*, 'Cherry water,') *Kirsch*, *Kirschwasser*. An alcoholic liquor, obtained from cherries bruised with their stones, by subjecting them to distillation, after having caused them to ferment. See *Spirit*.

KISKITOMAS NUT, *Hickory nut*.

KISS'INGEN, MIN'ERAL WA'TERS OF. These Bavarian springs have been long frequented. There are three,—two chalybeate, and one alkaline and acidulous.

KIST. A weight of 14 grains.

KITILLING (Prov.), Tickling.

KITTLING (Prov.), Tickling.

KLAPROTHII SULPHAS, (after M. H. Klaproth, the German chemist,) *Cadmii sulphas*.

KLAPROTHIUM SULPHURICUM, see *Cadmii sulphas*.

KLEPTOMA'NIA, *Cleptoma'nia*, *Klopema'nia*, *Clopemania*; (*κλεπτω*, 'I steal,' and *mania*.) *Insanity*, with an irresistible propensity to steal.

KLIÑOCEPHALIA, *Clinoccephalia*.

KLIPDAS, see *Hyrax Capensis*.

KLOPEMANIA, (*κλοπη*, 'theft,' and *mania*;) *Kleptomania*.

KNAP-KNEED. (Prov.) see Entogonyancan.

KNARESBOURGH, MINERAL WATERS OF. see Harrowgate, mineral waters of.

KNEADING, (from Sax. *cnæsan*, 'to kneed,') Shampooing.

KNEE, (Sax. *cnæp*,) Genu—k. Encircled hoof, Crusta genu equinæ.

KNEE, HOUSEMAID'S. An inflammation of the bursa, which in most individuals is in front of the patella—(F.) *Bursæ præpatellæ*.—and is apt to inflame and enlarge from effusion in those in whom it is subjected to much pressure. Hence its name. It is a form of capsular rheumatism.

KNEEHOLLY, Ruscus.

KNEEJOINT, Genu.

KNEE SCAB, Crusta genu equinæ.

KNIFE, (Sax. *cnif*, (F.) *Cuivif*.) *Cuiv'ter*, *Smil'la*, *Cuiv'tell'us*, *Tome'son*, *Tome'son*, *Machæ'ra*, *Machæ'ria*, (F.) *Couteau*. A cutting instrument, used in surgery to divide the soft parts, and which only differs from the bistouri or scalpel in being usually larger.

The most common knives are the following:

KNIFE, AMPUTATION, (F.) *Couteau à amputation*, *C. droit*, *Cuiv'ter rectus*. This is the largest of the knives used in surgery. Formerly, they were curved; now they are straight, and provided with one or two edges.

KNIFE, CATARACT, *Cerat'omus*, *Kerat'omus*, (F.) *Couteau à cataracte*. The cataract knives of Richter, Wenzel, Ward, and others, being intended to perform the section of the transparent cornea, are so shaped as to exactly fill the small wound made by them; and thus to prevent the discharge of the aqueous humour, until the section is completed. The blade of the knives of Wenzel and Ward resembles a very narrow lancet, blunt in the posterior five-sixths of one of its edges. The blade of that of Richter is pyramidal, cutting through the whole length of its inferior edge, and also blunt in the five-sixths of the upper.

KNIFE, CHES'ELDEN'S. A knife with a fixed handle; very convex on its edge, concave on the back, which was used by Cheselden in *lithotomy*.

KNIFE, CROOK'ED, *Cuiv'ter falcatus*, *Cuiv'ter curvus*, (F.) *Couteau courbe*. A knife, which is crooked and concave on its cutting edge. It was formerly employed in amputation of the limbs.

KNIFE, CROTCH'ET, *Cuiv'tell'us un'cus*, (F.) *Couteau à crochet*. A steel instrument, composed of a round staff, furnished with a handle at one extremity, and at the other with a curved knife. It was formerly used to cut to pieces monstrous fœtuses in utero, and to open the head when necessary.

KNIFE, DOUBLEEDGE'D, *An'ceps cuiv'tell'us*, *Amphis'mela*, *Amphis'milè*, *Cuiv'ting*, (F.) *Couteau à deux tranchans ou désarticulateur ou interosseux*. A knife, the blade of which is straight and sharp on both sides. It is used for disarticulating bones; and for cutting the soft parts situate between the bones, in amputation of the leg and forearm.

KNIFE EN SERPETTE (F.) *Couteau en serpette*. A sort of knife, of the shape of a *serpette* or pruningknife, invented by Desault for dividing the bony paries of the maxillary sinus, for the purpose of extracting fungi from it.

KNIFE, LENTICULAR, *Cuiv'ter lenticularis*, (F.) *Couteau lenticulaire*. An instrument, used in the operation of trepanning, for removing inequalities in the inner table of the skull, which may have been left by the crown of the trephine around the opening made by it. It is formed of a lenticular button, fixed at the extremity of an iron staff, which is convex on one side, flat on the other; sharp at both edges, and mounted on an ebony handle.

KNIFE, LITHOT'OMY, (F.) *Couteau lithotome*. A

name, given by Foubert to a large knife, the narrow blade of which, four and a half inches in length, was sharp in its whole extent, and made an obtuse angle with the handle. He used it in the lateral operation.

KNIFE, POLYPSY, Polypotome.

KNIFE, ROOTCUTTING. In Pharmacy, a knife moving on a joint at its pointed extremity, by which roots and other ligneous matters are divided in pharmaceutical processes.

KNIFESMAN, see Surgery, salivation.

KNITBACK, (Prov.) Symphytum.

KNOBWEED, *Collinsonia*.

KNOCK-KNEED, Entogonyancan.

KNOT, (Sax. *cnotta*.) Tubercle.

KNOT, PACKER'S, (F.) *Nœud d'emballeur*. A compressive bandage, used for arresting hemorrhage from the temporal artery or its branches. The *nœud d'emballleur* is made with a double-headed roller, five ells long. A graduated compress is placed over the opening in the artery, and the bandage is applied over it; the balls of the roller being carried horizontally round to the opposite temple, where they are crossed obliquely and carried back to the part where the compress is situate. The hands are then changed with the rollers, crossing them so as to form a knot, and taking one above the head; the other beneath the chin. They are then crossed again, so as to form several knots, one above the other. This bandage is called, by some surgeons, *so'lar* or *oblique chevestre* or *capistrum*.

KNOT, SURGEON'S, *Nœ'dus chirurgicus*, (F.) *Nœud du chirurgien*. A double knot made by passing the thread twice through the same noose. This knot is used frequently in the ligature of arteries, the umbilical cord, &c.

KNOTBERRIES, *Rubus chamaemorus*.

KNOTGRASS, *Polygonum aviculare*.

KNOTROOT, *Collinsonia Canadensis*.

KNOTWEED, *Collinsonia Canadensis*, *Polygonum aviculare*.

KNOWLTONIA VESICATORIA, (after Thomas Knowlton, of the botanic garden at Eltham.) *Nat. Order*, Ranunculaceæ. An acrid plant of South Africa, used by the Cape colonists as a blister in rheumatism.

KOHL SALAT, ('cabbage salad,') Cold slaw.

KOINOMIASMATA, (*κοινος*, 'common,' and *miasmata*.) see Miasm.

KOKUM, *Garcinia purpurea*—k. Butter, see *Garcinia purpurea*.

KOLERUS, a name given by Paracelsus to a dry ulcer.

KOLPORRAPHY, (*κολπος*, 'vagina,' and *ραφειν*, 'suture,') Elytrorrhaphy.

KOLTO, Plica.

KOOCHLA TREE, *Strychnos nux vomica*.

KOOSSO, *Hagenia Abyssinica*.

KOOTAHROOT, see Valeriana.

KOPYOPIA, Copyopia.

KORIS, (*κορις*,) Cimex.

KOSSO, *Hagenia Abyssinica*.

KOUMIS, Kumys.

KOUS, see Valeriana.

KOUSSIN, see *Hagenia Abyssinica*.

KOUSSO, *Hagenia Abyssinica*.

KRAME'RIA, (after the Kramers, German botanists,) *Krameria trian'dra*, *Krame'ria*, *Ratan'hia*, *Rhatan'ia*, *Rat'anhy*, (F.) *Ratanhie*.—*Sex. Syst.* Tetrandria Monogynia. *Nat. Ord.* Polygalæe.—is a native of Java. The root, *Krame'ria* (Ph. U. S.), has a bitter taste; and is astringent, diuretic, and detergent. Dose, ʒi to ʒi.

KRAME'RIA IXTNA or *Ratanhy* of the *Astilles* has similar virtues.

KRANIA, (*κρανια*.) see Cornus mas.

KREASOTON, Creasote.

KREATIC NAUSEA, see Nausea.

KREATINE. Creatine.

KREATININE, see Creatine.

KREATOPHAGOUS. (κρεας, gen. κρεατος, 'flesh,' and φάειν, 'to eat,') Carnivorous.

KREOSOTON, Creosote.

KRETINISMUS, Cretinism.

KRIEBELKRANKEIT (G), (kriebeln, 'to creep,' and krankheit, 'disease,') Convulsio cerebri.

KRONE, Crown.

KRYSTALLINE, see Crystalline.

KUA KAH, Curcuma longa.

KUKUI OIL, see Aleurites triloba.

KUMBECEPHALIC. *Cymbecephalic*, (κυμβη, 'a canoe,' and κεφαλή, 'head,') According to Professor Daniel Wilson, having a boat-shaped head.

KUMYSS. *Koumyss*, *Koumiss*. A beverage used in families of the people of Tartary. It resembles sour buttermilk, without being greasy. According to Sir George Simpson, it is prepared in a very simple way from mare's milk, which is merely allowed to stand for some days in a leathern churn till it becomes sour. It is then bottled for use. This drink is rather nutritious than exhilarating; but from the same material the Burats and the Kirghez prepare an intoxicating spirit in which they indulge to excess. A similar preparation is said to be in use in the Orkney and Shetland islands.

KUNAREE. *Nerium odorum*.

KUSSAUDE, *Convolvulus panduratus*.

KUTKULEJA, *Casalpinia bonducella*.

KUTU KURUNJA, *Casalpinia bonducella*.

KUTUBUTH. An Arabic name for a species of melancholy in which the patient is never quiet at any one place, but wanders about here and there, see Cuttubuth. Also, the name of an insect, which lives at the surface of stagnant waters, and is in a constant state of agitation. Some lexicographers imagine that it is on account of this last circumstance, that the name of the insect has been given to the disease.

KWOSO, *Hagenia Abyssinica*.

KYANOSIS, Cyanopathy.

KYAPUTY, Caleput.

KYESTEINE, Kiesteine.

KYESTINIC, Kiestinic.

KYLLOPODIA, (κυλλος, 'crooked,' and πους, gen. ποδός, 'a foot,') Kyllosis.

KYLLO'SIS, (κυλλος, 'crooked,' 'lame,') *Cyllo'sis*, *Kyllopodia*. Professor Chaussier so calls congenital distortion of the feet, *Club-foot*, *Polt-foot*, *Stump-foot*, (Prov.) *Pommelfoot*, *Pummelfoot*, *Tal'ipes*, (F.) *Pied bot*. Of this there are many varieties. In one, the foot, instead of resting on the soil, by the whole plantar surface, touches it only with the metatarsophalangean articulations. It seems as if turned backwards and broken upon the leg. (*Pes seu Tal'ipes Equinus*, *Hippopus*, *Oxyppodia*, *Tipfoot*, *Horse-foot*.) In other cases the foot is twisted inwards, (*Val'rus*, *Tal'ipes varus*, *Blesop'odes*, *Blw'opus*, *Cyllo'epus*, *Cyllop'oda*, *Cyllopod'ion*, *Lo'ripes*, *Cross-foot*, (F.) *Cig-neux*,) so that it rests only on the ground on its outer edge; or it may be twisted outwards, (*Val'*

gus, *Tal'ipes Valgus*, *Out'lore foot*,) or rest only on its inner edge, or on the back part of the heel, *Talipes calcaneus*, *Heel club-foot*. In the *flat-foot* or *splay-foot*, *Duck-foot*, *Splat-foot*, *Sw'opus*, *Plat'ypus*, *Platypod'ia*, (F.) *Pied plat*, the plantar surface of the foot is flattened instead of being concave.

These deformities are rarely accidental. They are almost always congenital, and may be rectified, at an early period, by proper mechanical means to strengthen the foot gradually, and restore it to its proper shape and direction; and if these means fail, the tendons and muscles concerned in the deformity may be divided. A similar affection of the hand is called, by the French, *Main bot*.

KYMOGRAPH'ION, *Cymograph'ion*; (κυμα, 'wave,' and γραφω, 'I describe,') A wave describer or measurer. An instrument invented by Ludwig, which is self-registering, and exhibits the relation between the waves of the pulse and the undulations produced by respiration.

KYNA, *Pastinaca opponanax*.

KYPHOSIS, Cyphosis—*k. Inflammatoria*, Vertebral disease—*k. Paralytica*, Vertebral disease.

KYPHOSKOLIOSIS, (κυφός, 'gibbous,' and σκολιος, 'crooked,') Cyphoma.

KYST, *Cyst*, *Kys'tis*, *Cys'tis*, *Kys'tē*, *Cys'tē*, (κυστις, *κυστη*, 'a bladder,' 'pouch,') (F.) *Kyste*. This term is generally applied to a pouch or sac, —*Cysto'ma*—without opening, and commonly of a membranous nature, which is accidentally developed in one of the natural cavities, or in the substance of organs. Many theories have been successively emitted to explain the formation of cysts, but none are entirely satisfactory. Some are formed by a thin, translucent membrane, having scarcely the thickness of the arachnoid; others of a whitish, fibro-cellular membrane, more or less thick. Some contain cartilaginous or bony flakes. The greater part have but one cavity; others, on the contrary, have several, separated by incomplete or imperfect septa, as is frequently seen in those developed in the ovaries. The matter contained in cysts is sometimes limpid, serous, yellowish white, reddish, and, at others, more or less thick, albuminous, adipous, or caseous. The tumour, formed by them, is called Encysted.

KYSTE, Kyst—*k. Anévrysmal*, Aneurismal sac.

KYSTEIN, Kiesteine.

KYSTH'ITIS, *Cysth'itis*, (κυσθος, 'the vagina,' and *itis*.) Inflammation of the vulva and of the mucous membrane of the vagina.

KYSTIQUE, Cystic.

KYSTHOPTO'SIS, (κυσθος, 'the vagina,' and *πτωω*, 'I fall,') Prolapsus or inversion of the vagina.

KYSTITOME, *Cystitome*.

KYSTOTOMY, Cystotomy.

KYTE (Sc. and Prov.), Abdomen, Stomach.

KYTARRHAG'IA, (κυτταριον, 'an alveolus,' and *ῥαγω* or *ῥήγνυμ*, 'I break forth,') Discharge of blood from an alveolus.

L.

LA VENO BENO, see Falsifications, table of.

LABA'RIMUM, (*labi*, 'to fall,') Looseness of the teeth.

LABDACISMUS, (λαβδαισμος,) Lallation.

LABDAMEN, *Cistus creticus*.

LABDANUM, see *Cistus creticus*.

LA'BE, (λαβη, from λαβω, λαβανω, 'I seize,' 'I take,' 'the act of grasping,') Invasion. Also, employed to denote the first paroxysm of fevers.

LABEO, (from *labium*,) Chilon.

LABES, Chilon, Macula.

LABIA CUNNI, (pl. of *labium*,) Lips of the

valva.—1. Internus see Minora. Nymphae.—1. Majora. Lips of the valva.—1. Palatum. Lips of the valva.—1. Palatum minora. Nymphae.

LABIAL, Labialis, (from *labium*, 'a lip.') Relating or appertaining to the lips.

LABIAL, Labialis, Orticularis oris.

LABIAL ARTERY, Haller and Sabatier call thus the facial artery of the majority of anatomists. The labial arteries, properly so called, containing arteries of the lips, (F.) *Arteriae faciales*, are two in number. The superior arises from the facial, above, and very close to, the commissure of the lips. It is large and tortuous, and is distributed to the upper lip. The lower arises from the facial, at a considerable distance from the commissure, and proceeds, in a serpentine course, into the substance of the lower lip, to which it is distributed.

LABIAL GLANDS. This name is given to a multitude of mucous follicles, of some size, round, prominent, and separate from each other, which are found on the inner surface of the lips, below the mucous membrane.

LABIAL VEINS are distinguished, like the arteries, into *superior* and *inferior*. They open into the facial vein (—a division of the internal jugular).

LABIALIS, Orticularis oris.

LABIDOMETER, Labimeter.

LABIMETER, (F.) *Labimeter* on *Labidometer*, (from *labu*, gen. *labubo*, 'forceps,' and *metron*, 'measure,') A scale adapted to the handles of the forceps, which indicates the distance of the blades from each other, when applied to the head of the child in *caesare*.

LABIS, gen. *Labidis*, (Labu, from *labuere*, 'to seize,') Forceps.

LABIUM, (Labium, 'to seize,') Lip.—1. Leporinum, Harelip.—1. Uteri, Amphideum.

LABLAB, Dolichos lablab.

LABORANS, (from *labor*, 'disease,') Sick.

LABORATOIRE, Laboratory.

LABORATORIUM, Laboratory.—1. Chymicum seu pharmacotheca, see Pharmacopoeia.

LABORATORY, *Laboratorium*, *Ergasterium*, (F.) *Laboratorium*, (*laborans*, *laboratum*, 'to work,') A workshop. A place for preparing chemical or pharmaceutical products, &c.

LABORIOSUS, Laborious, Sick.

LABORIOUS, *Laboriosus*, (F.) *Laboriosa*, (from *labor*, 'difficulty,') Delivery is said to be laborious, *Partus laboriosus*, *Magnus*, (F.) *Amplius laboriosa*, when attended with more difficulty and suffering than usual. With some, *laboriosa labor* means one that requires the use of instruments. See Dystocia, and Parturition.

LABOUR, (*labor*) Parturition.

LABOUR CHAIR, *Obstetric chair*. A chair, in which a parturient woman is placed during delivery.

LABOUR, COMPLICATED, see Parturition.—1. Difficult, Dystocia.—1. Dry, Partus siccus.—1. Forced, see Parturition.—1. Impassible, see Parturition.—1. Laborious, see Laborious.—1. Morbid, Dystocia.

LABOUR PAINS, see Pains.

LABOUR, POWERLESS, see Parturition.—1. Premature, Parturition, (premature)—1. Preternatural, Metastocia, see Parturition.

LABOUR SHOW, see Parturition.

LABRISULCIUM, (*labrum*, and *ensula*, 'a furrow,') Chelicoene, Stomacoe.

LABRUM, Lip.—1. Cerebri, Infundibulum of the brain.

LABRUSCA, Bryonia alba.

LABYRINTH, (*labyrinthos*,) *Labyrinthus*, *Antrum tympanicum*. A place full of turnings, the exit of which is not easily discoverable. Anatomists have given this name to the aggregate of parts, constituting the internal ear, *Labyrinthus auris internae*, *Intra pars operculi et cilia*, *Labyrinthi cavity of the ear*. The Labyrinth is situated between the tympanum and meatus auditorius

internus. It is composed of several varieties, which communicate with each other in the dried bone; as the *membrana vestibularis*, *membrana cochlearis*, &c. It is lined by periosteum, and also by a pulpy substance, constituting the *membrana labyrinthi*, in which the auditory nerve is regularly dispersed. This membrane forms two sacs in the vestibule, called *utricle* and *sacculus*, and *utricle*, respectively, which resemble in shape that of the honey cavities containing them. Each sac contains calcareous matter, constituting the *Crystals* and *Otoliths*. When the sac is laid open, upon the upper and outer part, a partition appears, partaking of the nature of the sac, and called by Meckel, *Sepalum cristae utriculo-membranaceum*.

Labyrinth is also applied to the lateral portions of the ethmoid bone, which are formed of large irregular cells.

LABYRINTH, MEMBRANOUS, see Labyrinth.

LABYRINTHIC, *Labyrinthicus*, *Labyrinthicus*, (F.) *Labyrinthique*. Relating or appertaining to the labyrinth.

LABYRINTHIC CAVITY OF THE EAR, Labyrinth.

LABYRINTHODES, (*Labyrinthos*, from *labyrinthos*, and *odes*,) Labyrinthos.

LABYRINTHUS, *Labyrinthos*, Labyrinth.—1. Auris internae, see Labyrinth.

LAC, *Lacca*, *Lacquer*. Milk.—1. Ammoniac, Mixture ammoniac.—1. Amygdala, Emulsion amygdala.—1. Asaferida, Mixture asaferida.—1. Aris, Albumen ovi.—1. Guaiac, Mixture Guaiac.

LAC, GUM. Lacca.

LAC LUXA, Marga candida.—1. Maris, Spermi.—1. Pressum, Urd.—1. Primam Puergera, Colostrum.

LAC, SEED, Lacca.—1. Shell, Lacca.—1. Stick Lacca.

LAC SULPHUREUS, Sulphur præcipitatum.—1. Terre, Magnesie carbonas.—1. Virginia, Virgin's milk.

LACAPHTHON. Presumed to be the bark of some fir or other tree. It entered into the composition of the *Cypri*, used in fumigations. Supposed to be the *Naraphothum* or *Nemaphothum*.

LACCA, (Lakk, Arab.,) Gum of lac, Stick Gum, Seed, or Shell Lac, (F.) *Lacca*, *Gummi Laccae*. Lac is a substance formed by an insect, and deposited on different species of trees, chiefly in the East Indies, (for example, on *Croton tigliifer*, and two species of *Ficus*,—*Ficus religiosa*, and *F. indica*). The various kinds, distinguished in commerce, are *stick-lac*, which is the substance in its natural state, investing the small twigs of the tree; and *shell-lac*, which is the same broken off. When melted it is called *shell-lac*.

Lac was, at one time, used in the form of tincture, as a tonic and astringent; and it still forms part of particular dentifices.

LACERATION, *Laceratio*, (*lacerare*, *laceratum*, 'to tear,') *Ruptura*, *Rheuma*, *Rheuma*, *Ruptura*, *Ruptura*, *Dislocation*, *Spargium*, (F.) *Arachnoid*, *Dischizement*, *Dislocation*, *Arachnoid*. The act of tearing or rending. The breach made by tearing or rending; as a *lacerated wound*, *Traus ciliatus laceratum*, *Laceratum*, *Fallus laceratum*.

LACERATURA, see Laceration.

LACERTA, Lizard.—1. Aquatica, see Ectocera.

LACERTI CORDIS, Columnæ carnea.—1. Muscularum, see Muscular fibre.

LACERTOSUS, Muscular.

LACERTULI CORDIS, (dim. of *Laceratus*,) Columnæ carnea.

LACERTUS, Brachium, Muscle, see Muscular fibre.

LACERUM, (*lacer*, 'to tear,') Any thing torn, or appearing as if torn. (F.) *Lacerat*.

LACERUM FORAMEN ANTERIUS, (F.) *Traus ciliatus anterior*, *Huic aperturæ pte*, (Ch.) is an

irregular opening, formed by the sphenoid, occipital, and petrous portion of the temporal bone. It is closed by cartilage, and forms a kind of fontanelle, situated at the termination of the petro-occipital suture.

LACERUM FORAMEN POSTERIUS, *Foramen jugulârè*, *F. lacerum* in *Ba'si Cra'nii*, (F.) *Trou déchiré postérieur*, *Hiatus occipito-pétreux*, (Ch.) is formed by the occipital bone, and the inferior edge of the petrous portion of the temporal bone. Through it, the internal jugular vein, the eighth pair of nerves, and accessory nerve pass out of the cranium.

LACERUM FORAMEN SUPERIUS, Sphenoidal fissure.

LACH'ANON, (λαχανον, from λαχανω, 'I dig.') A pot herb; *Oler*, pl. *Olera*, a vegetable used as food. (F.) *Potager*.

LACHESIS PICTA, see Arrow-poison.

LACHESIS RHOMBEA'TA, *Flammon*. A poisonous serpent common in the lower forests of Peru.

LACHNANTH'ES TINCTO'RIA, (λαχνη, 'wool,' and ανθος, 'a flower,' owing to its woolly flowers.) *Dilat'ris tincto'ria*, *Red'root*: indigenous. *Ord.* *Hæmodoracææ*. The root is astringent and tonic.

LACHRYMA, (δακρυμα,) Tear; see *Vitis vinifera*.

LACHRYMAL, *Lac'rymal*, *Laeryma'lis*, (from *laeryma*.) Belonging to the tears. This epithet is given to various parts.

LACHRYMAL APPARATUS, see *Lachrymal Passages*.

LACHRYMAL AR'TERY proceeds from the ophthalmic; and distributes its principal branches to the lachrymal gland.

LACHRYMAL BONE, *Unguis os*.

LACHRYMAL CANAL' or DUCT, *Na'sal Canal' or duct*, *Dacryoso'len*, *Canal'is laeryma'lis* seu *or'bite nasa'lis* seu *Sac'ci laeryma'lis*, *Duct'us ad Na'sum* seu *Nasa'lis or'bite*, is formed by the superior maxillary bone, os unguis, and os turbinatum inferius; and is seated in the outer paries of the nasal fossæ. It is lined by a prolongation of the mucous membrane of the lachrymal sac; and its inferior orifice is furnished with a valvular duplication. This duct transmits the tears, which have been absorbed at the great angle of the eye, by the puncta laerymalia, into the nasal fossæ.

LACHRYMAL CARUNCLE, see *Caruncula laerymalis*—1. *Cyst*, *Dacryops*—1. *Duct*, *Lachrymal Canal*.

LACHRYMAL FOS'SA, *Fossa* seu *Fo'vea laeryma'lis*, is a slight depression at the upper part of the orbit, which lodges the lachrymal gland.

LACHRYMAL GLAND, *Glandula laeryma'lis* seu *innomina'ta Gale'ni*, is seated in a depression of the frontal bone at the upper, anterior, and outer part of the orbit. It is of about the size of an almond; and of an oval shape, flattened above and below:—its great diameter being the antero-posterior. It is composed of several small lobules, united by areolar tissue, and separated by it as well as by vessels and nerves which creep in the intervals. This gland has seven or eight excretory ducts, which open behind the upper eyelid. Its use is to secrete the tears, and pour them on the globe of the eye by the excretory ducts.

LACHRYMAL GROOVE, (F.) *Gouttière laerymale*, is the bony channel, which lodges the lachrymal sac. It is seated at the anterior and inner part of the orbit and is formed by the os unguis and the ascending process of the upper jaw bone.

LACHRYMAL HER'NIA, *Lachrymal Tu'mour*, *Her'nia Sac'ci Laeryma'lis*, is when the tears enter the puncta, but cannot pass to the nose, and accumulate. By *Anel*, this was called *Dropsy of the Lachrymal sac*.

LACHRYMAL MUSCLE, *Tensor tarsi*.

LACHRYMAL NERVE is the smallest of the three branches formed by the ophthalmic nerve. It is distributed, particularly, to the lachrymal gland and to the upper eyelid. In its course it gives off a *sphenomaxillary* and a *malar* filament.

LACHRYMAL PAPILLA, see *Lachrymal Puncta*.

LACHRYMAL PASS'AGES, *Viæ laeryma'les*, (F.) *Voies laerymales*. The organs concerned in the secretion of tears, in spreading them over the eye, and taking them up again to transmit them into the nasal fossæ. The lachrymal passages are composed of the *lachrymal gland*, *caruncle*, *puncta*, *ducts*, *lachrymal sac*, and *nasal duct*, which, together, form the *lachrymal appara'tus*, (F.) *Appareil laerymal*.

LACHRYMAL PUNCTA, *Tear puncta*, *Puncta Laeryma'lia*, *Spiram'ina Palpebra'rum*, (F.) *Points laerymaux*, are two small, round, and contractile openings, situate in the centre of a tubercle or papilla, *Papilla laeryma'lis*, *Tuber'culum laeryma'le*, about a line and a half distant from the inner commissure of the eyelids, and continuous with the lachrymal ducts. These ducts, *Laeryma'les Canalic'uli*, *Canal'es* seu *Duct'us laeryma'les*, *Duct'us laerymales latera'les* seu *puncto'rum laeryma'lium*, *Cor'nua laeryma'lia* seu *lima'cum*, *Collic'is puncto'rum laerymalium*, *Canalic'uli lima'cum*, (F.) *Conduits laerymaux*, are two in number—a *superior* and an *inferior*—which extend from the puncta to the lachrymal sac. They seem formed by a very delicate prolongation of the conjunctiva, which is continuous with the mucous membrane of the lachrymal sac, (F.) *Reservoir des larmes*:—the *Sac'cus* seu *Si'nus* seu *La'eus laeryma'lis*, *Dacryocys'tis*, *Infundib'ulum laeryma'le*.

LACHRYMAL SAC, see *Lachrymal puncta*.

LACHRYMAL TUBERCLE, see *Lachrymal puncta*.

LACHRYMAL VEINS accompany the artery of the same name, and open into the ophthalmic and palpebral veins.

LACHRYMATION, *Laerymatio*—1. *Sanguineous*, *Dacryhæmorrhysis*.

LACIN'IA, 'a hem or fringe;' hence

LACINLÆ TUBARUM FALLOPII, see *Tube*, *Fallopian*.

LACIN'IATED, *Lacinia'tus*, (F.) *Lacinié*, (from *lacinia*.) *Fringed*,—as the lacinated extremities of the Fallopian tubes.

LACIS, (F.) (from *lacer*, 'to lace,') *Plexus*.

LACMUS, *Lichen roccella*—1. *Tinctorius*, *Lichen roccella*.

LACONICUM, (used by the Λακωνες, *Lacedæmonians*.) *Vaporarium*, see *Stove*.

LACQ, *Laqueus*.

LACRIMA, *Tear*.

LACRUMA, *Tear*.

LACRYMA, *Tear*.

LACRYMAL, *Lachrymal*.

LACHRYMALIS, *Lachrymal*.

LACRYMA'TIO, *Lachryma'tion*, (from *laeryma*.) *Epiphora*. Also, a profuse secretion of tears from any cause,—weeping, *Fle'tus*, *Plora'tio*, *Plora'tus*.

LACRYMINE, see *Tears*.

LACTALBUMEN, (*lac*, gen. *lactis*, 'milk,' and *albumen*.) *Casein*.

LACTANS, (*lactare*, 'to give milk,') *Nurse*.

LACTA'RIMUM, *Lacticin'ium*, (from *lac*.) Food prepared from milk. Milk diet or food.

LACTATE DE FER, *Ferri lactas*—1. *de Quinine*, *Quinîe lactas*.

LACTA'TION, *Lacta'tio*, *Thela'sis*, *Thelas'mus*, (*lactare*, *lactatum*, (*lac*, 'milk,') 'to suckle,' 'to give milk.') *Suck'ling*:—the giving of *suck*, (F.) *Allaitement*. The French make four varieties of lactation. 1. *Allaitement maternel*—*Maternal Lactation*, when the mother suckles the child. 2. *A. étranger mercenaire*,—when another

suckles it. 3. *A. artificial*, when the child is brought up by hand. 4. *A. animal*, when the child is suckled by an animal.

LACTÉ, Lactic.

LACTEAL, Lactic—1. Sac, see Lactiferous vessels—1. Swelling, Lactiferous swelling.

LACTEALS, Chyliferous vessels.

LACTEIN, Milk, solidified.

LACTENS, (*lacteo*, 'to suck,') Sucking child.

LACTES, Mesentery, Pancreas.

LACTEUS, Lactic.

LACTIC, *Lacteus*, *Galacticus*, *Galactinus*, *Lactéal*, *Milky*, (F.) *Laitue*, *Laitoux*. Appertaining to, or resembling milk.

LACTIC ACID, *Ac'idium Lactéum* seu *Lactis* seu *Lacticum*, Acid of milk, *Galactic*, *Nan'ceic* or *Ca'seic Ac'id*, (F.) *Acide Lactique*. This has been recommended as a therapeutical agent in atonic dyspepsia, owing to its being presumed to be one of the gastric acids secreted in health. It is given either in the form of lemonade or of lozenge. The acid is obtained either from milk or from the juice of the red beet. S. g. 1.212.

LACTICA (Arab.), Typhoid.

LACTICANS, (from *lac*, 'milk,') Nurse.

LACTICINIUM, Lactarium.

LACTIFÈRE, Lactiferous.

LACTIFEROUS, *Galactoph'orous*, *Lactifer*, (F.) *Lactifère*, (*lac*, and *fero*, 'I carry,') Milk-conveying.

LACTIFEROUS or **LACTEAL SWELLING**. A tumefaction of the breast, supposed by Sir Astley Cooper to arise from a large collection of milk in one of the lactiferous tubes, the result of chronic inflammation of the tube near the nipple, with closure of its aperture, and obliteration of the canal for an inch or more. The tube requires to be punctured.

LACTIFEROUS VESSELS, *Lactif'erous* or *Galactoph'orous Ducts*, *Tu'buli lactif'eri*, *Ductus lactif'eri* seu *lac'tei*, (F.) *Vaisseaux ou conduits lactifères*, are the excretory ducts of the mammary gland, each of which, running towards the nipple, dilates beneath the areola into an elongated sacculus—the *lactéal* or *galactophorous sac*, *receptacle* or *sinus*, *ac'culus* seu *si'nus lactiferus*.

LACTIFUGA, *Lactifuge*, (*lac*, and *fugo*, 'I drive away,') Medicines which dispel milk.

LACTIGO, gen. *Lactig'inis*, (from *lac*, 'milk,') Porrigo larvalis.

LACTIN, Sugar of milk.

LACTINATED, *Lactinatus* (from *lactin*.) Relating or appertaining to lactin. Lactin, or sugar of milk, being hard, it is used for the division and dilution of substances; and powders thus formed have been called *lactinated powders*.

LACTIN'IA, Food prepared with milk.

LACTIPOTOR, (*lac*, and *potare*, 'to drink,') Galactopotes.

LACTIS CONCRETIONES, Infarctus Mam-mæ lacteus.

LACTIS REDUNDANTIA, Galactia—1. Retentio, Galactischesis—1. Sanguinolenti Excretio, Galactohæmia.

LACTISUGIUM, (*lac*, and *sugere*, 'to suck,') Anthia lactea.

LACTIVORUS, (*lac*, and *vorare*, 'to devour,') Galactophagous.

LACTOCELE, (*lac*, and *κηλη*, 'a tumour,') Galactocèle.

LACTOLINE, Milk, solidified.

LACTOMETER, (*lac*, and *μετρον*, 'a measure,') Galactometer.

LACTOSE, (from *lac*, 'milk,') Sugar of milk.

LACTUCA, (from *lac*, 'milk,') so called, from its milky juice.) *Lactuca Sat'iva*, *Let'tuce*, *Går-den Lettuce*, *Eunu'chium Meco'nis*, *Thri'dax*, *Thri-dar'inè*, *Cher'bas*, (F.) *Laitue ordinaire*; Ord. Com-

positæ, is used as a wholesome salad. The seeds possess a quantity of oil, which, when formed into an emulsion, has been advised in ardor urinae, &c.

The inspissated juice, *Lactuca'rium*, *Thrid'acè*, *Lettuce opium*, resembles, in odour and appearance, that of opium, and is, like it, narcotic, but uncertain. Dose, gr. j to x, and more.

LACTUCA ELONGATA, see *L. virosa*—1. *Floridana*, *Mulgedium Floridanum*—1. *Graveolens*, *L. virosa*—1. *Marina*, *Fucus vesiculosus*.

LACTUCA SCAR'OLA, *L. Sylvest'ris*, *Scari'ola*, (F.) *Laitue Scariole* ou *Sauvage*, possesses a stronger degree of bitterness than *L. sativa*. It has similar virtues with *L. virosa*.

LACTUCA SYLVESTRIS, *L. scariola*—1. *Villosa*, *Mulgedium acuminatum*.

LACTUCA VIRO'SA, *L. grave'olens*, *Strong-scented Lettuce*, (F.) *Laitue vireuse*. The odour of this plant, the leaves of which are used in medicine, is strongly narcotic, and the taste bitter. They are narcotic, diuretic, and aperient; and have been used in dropsies. *Lactuca elong'ata*, *Wild Lettuce* of the United States, has been employed for *L. virosa*.

LACTUCARIUM, see *Lactuca*.

LACTUCIMEN, (from *lac*, 'milk,') Aphthæ.

LACTUCIMINA, Aphthæ.

LACTUMEN, (from *lac*, 'milk,') Porrigo larvalis.

LACTUMINA, Aphthæ.

LACUNA, *Canalic'ulus*, (dim. of *lacus*, 'a lake, or deep ditch, or cistern,' *λακος*, and *λακος*.) *A Fos'sa* or *Ditch*. A small cavity in a mucous membrane, the parietes of which secrete a viscid humour. It is used synonymously with *crypt*. Also, the external opening of such cavity.

LACUNA, *Crypta*, *Fontanella*—1. *Cerebri*, *Infundibulum* of the brain, *Pituitary gland*.

LACUNA seu **SULCUS** seu **SULCULUS** **LABII SUPERIORIS**, *Amab'ilis*, *Amato'rium*, *Phil'trum*. The hollow of the upper lip under the nose.

LACUNA MAGNA, see *Urethra*.

LACUNE, (pl. of *Lacuna*), see *Urethra*.

LACUNE OF BONE. Certain dark stellate spots, with threadlike lines radiating from them, seen under a high magnifying power. These were at first believed to be solid os'seous or bone cor'puscles or cells, *Corpus'cula os'sium*, *Cor'puscles of Purkinje*, (F.) *Cellules des os ou osseuses*, *Corpuscules des os ou noirs ou ramifiés ou calcaires*, *Ostéoplastes*; but are now regarded as excavations in the bone—*Sac'culi chalcoph'ori*—with minute tubes or *canaliculi* proceeding from them, and communicating with the Haversian canals. The lacune and canaliculi are fibres concerned in the transit of nutrient fluid through the osseous tissue.

LACUNE GRAAFIA'NÆ, *L. mucosæ vulvæ*. The mucous follicles of the vagina.

LACUNE MUCOSÆ VULVÆ, *L. Graafianæ*—1. *Palpebrarum*, *Meibomius*, glands of.

LACUNAR, *La'quear*, 'an arched roof or ceiling.' The roof of a chamber. Hence,

LACUNAR OR'BITÆ. The roof of the orbit.

LACUNAR VENTRICULI QUARTI SUPERIOR, *Vavula Vieussenii*.

LACUNE DE LA LANGUE, *Cæcam lora-men*—*L. de Morgagni*, see *Urethra*.

LACUS LACRYMALIS, (*lacus*, 'a lake,' or cistern, *λακος*.) *Lachrymal sac*.

LACUS LACRYMA'RUM. A small space in the inner angle of the eye between the lids, towards which the tears flow, and at which the triangular canal formed between the closed lids terminates.

LADA, *Piper nigrum*.

LADANUM, see *Cistus creticus*.

LADENDO, *Influenza*.

LADIES' MANTLE, *Alchemilla*.

LADIES' SLIPPER, *SHOWY*, *Cypripedium*

spectabile—*l. Slipper*, stemless, *Cypripedium acaule*—*l. Slipper*, yellow, *Cypripedium luteum*.

LADIES' SMOCK, *Cardamine pratensis*.

LADRE, (from *Laz'arus* [?]), Leprous.

LADRIÈRE (F.), (from *ladre*.) (F.) *Léproserie*, *Maladerie*. A vulgar name for elephantiasis, or lepra. Also, an hospital for the reception of the leprous, *Leprosarium*, *Do'mus leprosa'ria*, *Lepor hospital*.

LADYBIRD, *Coccinella septempunctata*.

LADYBUG, *Coccinella septempunctata*.

LADYCOW, *Coccinella septempunctata*.

LADY CRESPIGNY'S PILLS, see *Pilulæ Aloes et Kinæ Kinæ*.

LADY HESKETH'S PILLS, see *Pilulæ Aloes et Kinæ Kinæ*.

LADY WEBSTER'S PILLS, see *Pilulæ Aloes et Kinæ Kinæ*.

LADY'S THUMB, *Polygonum persicaria*.

LÆDENS, pl. *Læden'tia*, (*lædo*, 'I hurt.') Any thing hurtful or injurious, as to the economy.

LÆMOPARALYSIS, (*læmus*, and *paralysis*.) Cæphaloplegia.

LÆMOSCIRRHUS, *Cancer pharyng'is et œsoph'agi*, *Læmosten'ois* seu *Dyspha'gia scirr'hosa*; (from *læmus*, and *scir'pos*, 'scirrhus.') Cancer of the pharynx or œsophagus.

LÆMOSTENOSIS, (*læmus*, and *stenosis*, 'narrowness,') *Dysphagia constricta*—*l. Scirr'hosa*, *Læmoscirrhus*.

LÆMUS, (*læmos*.) Pharynx.

LÆSIO, (*lædere*, *læsum*, 'to hurt.') Lesion—*l. Continui*, see *Continuity*, solution of.

LÆTIFICANTIA, (*lætifico* (*lætus*, and *facio*.) 'I make glad.') Medicines formerly used as cordials, in depression of spirits, &c.

LÆVIGATIO, (*lævis*, 'light, smooth.') Levigation.

LÆVITAS INTESTINORUM, (*lævis*, 'smooth,') Lientery.

LAG TEETH, Molar teeth.

LAGENTOMUM, (*λαγος*, 'a hare,' *en*, and *τομη*, 'section,') Harelip.

LAGERBEER, see *Cerevisia*.

LAGNEA, (*λαγνεία*) Coition, Satyriasis, Sperm.

LAGNEIA, (*λαγνεία*) Coition, Satyriasis, Sperm.

LAGNESIS, (from *λαγνός*, 'lustful.') Furor Femininus, Nymphomania—*l. Furor masculinus*, *Satyriasis*—*l. Salacitas*, *Satyriasis*.

LAGNEUMA, (*λαγνευμα*) Coition, Sperm.

LAGNIA, (*λαγνεία*) Coition, Satyriasis, Sperm.

LAGNOSIS, (from *λαγνός*, 'lustful.') Satyriasis.

LAGOCHELILUS, (*λαγος*, 'hare,' and *χειλος*, 'lip,') Harelip.

LAGONES, (*λαγones*.) Flanks.

LAGONOPONOS, (*lagones*, and *πονος*, 'suffering,') Pleurodynia.

LAGOPHTHALMIA, (*λαγος*, 'a hare,' and *ὀφθαλμος*, 'an eye,') *Lagophthal'mus*, *Lepori'nus ocul'us*, *Hare's Eye*, (F.) *Œil de Lièvre*. A vicious arrangement of the upper eyelid, which is so retracted that it cannot cover the globe of the eye during sleep. It has been asserted that this condition of the eye is natural in the hare when asleep.

LAGOPHTHALMUS, (*λαγωφθαλμος*.) Lagophthalmia, Geum urbanum.

LAGOSTOMA, gen. *Lagostom'atis*, (*λαγος*, 'hare,' and *στομα*, 'mouth,') Harelip.

LAÏ, Aï.

LAÏCHE, Sarsaparilla Germanica—*l. des Sables*, Sarsaparilla Germanica.

LAIT, (from *lac*, gen. *lactis*, 'milk,') Milk—*l. Adoucissant*, Looch ex ovo—*l. d'Amandes*, Emulsion amygdalæ—*l. d'Anesse*, Milk, asses'—*l. d'Asafetida*, Mistura asafetidæ—*l. de Beurre*, Butter-milk—*l. de Brebis*, Milk, ewes'—*l. Caillé*, Curds—*l. de Chèvre*, Milk, goats'—*l. Épanché*, *L. ré-*

pandu—*l. de Femme*, Milk, human—*l. de Jument*, Milk, mares'—*l. de Poule*, see *Ovum*.

LAIT RÉPANDU, ('milk diffused,') (F.) *L. épanché*. A popular expression in France, under which is comprehended every kind of disease—and particularly vague pains—occurring after delivery; all being ascribed to diffusion or deposition of milk.

LAIT DE TERRE, ('milk of earth,') Magnesie carbonas—*l. de Vache*, Milk of the cow—*l. Virginal*, Virgin's milk.

LAITERON DOUX, (from *lait*, from its milky juice.) Sonchus oleraceus.

LAITEUX, Lactic.

LAITAT, ([F.] from *lait*.) Sour whey in which different wild fruits have been macerated. Said to be much used in the Jura as a refreshing drink.

LAITUE ORDINAIRE, (from *lait*, from its milky juice.) Lettuce—*l. Sauvage*, *Lactuca scariola*—*l. Scariote*, *Lactuca scariola*—*l. Vireuse*, *Lactuca virosa*.

LA'ITY, La'ici, (*λαῖ*, 'the people,') Formerly, and generally at the present day, the people as distinguished from the clergy. Frequently, however, employed in the sense of the people as distinguished from those of the other learned professions. A *lay'man* is one of the laity.

LAKEWEED, *Polygonum hydropiper*.

LALIA, (*λαλία*, 'speech,') Voice, articulated.

LALLA'TION, *Lalla'tio*, *Lambdacis'mus*, *Lambdacis'mus*, *Lul'laby speech*. Sauvages uses this term for a vicious pronunciation, in which the letter L is improperly doubled or softened or substituted for R.

LAMAC, *Acaciæ gummi*.

LAMBACISMUS, (from the Greek letter Λ, *λμβάα*) Lallation.

LAMB'DOID, *Lambdoid'al*, *Lambdō'id'ees*, *Lambdo'd'es*, *Lambdō'id'ees*, (Λ, (*λμβάα*), and *είδος*, 'shape, resemblance,') Anatomists have given the name LAMBDOID'AL SUTURE, *Sutur'a Lambdoid'a'lis* seu *lambdō'id'ees* seu *lambdō'idea* seu *Pro'ra*, to the suture, formed by the parietal bones and the occipital, because it resembles the letter Λ (*lambda*) of the Greeks. It is the *Occip'ito-pari'etal suture*, *Suture occipitale*, (Ch.) In this suture, the ossa Wormiana are most frequently met with; and the denticulations are most distinctly marked.

LAMBEAU, Flap.

LAMBITIVUM, (*lambere*, 'to lick,') Electos.

LAMBKILL, *Kalmia latifolia*.

LAMBKILLER, *Andromeda mariana*.

LAME, *Lamina*—*l. Cornée*, *Tænia semicircularis*—*l. Ruyschienne*, *Ruyschiana tunica*—*l. Vitrée*, see *Table*.

LAMELLA, *Lamina*—*l. Plana*, *Planum os*.

LAMEL'LAR, *Lamello'sus*, *Laminated*, (F.) *Lamel'le*, *Lamelleux*, *Lamineux*, composed of thin laminae or leaves—as the *Lamellar* or *laminated tissue*; i. e. the areolar tissue.

LAMELLÉ, Lamellar.

LAMELLEUX, Lamellar.

LAMENESS, Claudication.

LAMIER, *Lamium album*.

LAM'INA, *Lamel'la*, ('a plate,') (F.) *Lame*. A thin, flat part of a bone, a plate or table, as the cribriform lamina or plate of the ethmoid bone. *Lamina* and *Lamella* are generally used synonymously; although the latter is properly a diminutive of the former.

LAMINA CILIARIS, Ciliary zone.

LAMINA CINER'EÆ. A thin layer of gray substance which forms the anterior part of the inferior boundary of the third ventricle of the brain.

LAMINA COR'NEÆ, *Tænia Tarsi*. A yellowish band or a thickening of the lining membrane of the ventricle, by which the vena corporis

striati is overlaid in the lateral ventricle of the brain.

LAMINA CRIBROSA, Cribriform lamella—*l.* Cribrosa ossis ethmoidalis, see *Cribell*—*l.* Fusca, see *Sclerotie*—*l.* Fusca of the Choroid, see *Choroides tunica*—*l.* Medullaris triangularis cerebri, *Lyra*—*l.* Papyracea, *Planum os*—*l.* Perforata anterior, *Loeus perforatus anticus*—*l.* Perforata posterior, see *Tarini pons*.

LAMINA PERITONEI EXTERNA. The outer lamina or fold of the peritonæum.

LAMINA PRIMA CUTIS, Epidermis.

LAMINA SPIRALIS, *Septum seu lœ seu coch'leæ auditivæ*. A partition between the scalæ of the cochlea. The largest part of this, next the modiolus, is formed of bone. The remainder, or that part next the opposite side of the scalæ, is composed of a cartilaginous membrane, called by Val-salva, *Zo'na seu Zo'mula Coch'leæ*. By some anatomists the lamina is divided into a *Zona os'sea* and *Z. mollis*. By others it is considered to consist of four laminae, when examined with a strong glass: a *Zona os'sea* next to the modiolus—a *Zona coria'cea* on the outer side of this; a *Zona vesicul'ris*—and a *Zona membrana'cea*, which is, perhaps, the lining membrane of the cochlea. At the apex of the cochlea, the lamina spiralis terminates by a pointed hook-shaped process, *ham'ulus lam'inae spiralis*.

LAMINA OF A VERTEBRA, see *Vertebra*.

LAMINE DORSAL'ES, *Dorsal lam'inae*. Two oval masses on each side of the primitive groove of the embryo, which approach so as to form a groove, in which are lodged the future brain and spinal marrow.

LAMINE SPONGIOSÆ NASI, Turbinate bones.

LAMINE VENTRAL'ES, *L. viscerales*. Thickened prolongations of the serous layer of the germinal membrane, which, by their union, form the anterior wall of the trunk of the new being.

LAMINARIA DIGITATA, *Tan'gle, Sea tan'gle*. One of the Algæ eaten in Scotland, and hawked about the streets with the Pepperdulse. It has been proposed to make tents of it.

The leaf of a seaweed—a species of *Laminaria*—is employed in the Himalayas under the name of *goitre leaf*; so called because chewed by the inhabitants where goitre prevails.

LAMINATED, Lamellar.

LAMINEUX, Lamellar.

LAMIUM ALBUM, (from *lama*, 'a ditch or bog,' because often found there, [?]) *L. foliosum*, *Urtica mor'tua*, *Galeop'sis*, *Archangel'ica*, *Dead or White Archangel Nettle* (F.) *Ortie blanche ou morte*, (Sc.) *Day'nettle*, *Lamier*. Ord. Labiatæ. Infusions of this plant have been recommended in uterine hemorrhage and leucorrhœa. It is not used.

LAMIUM AMPLEXICAUL'È, *Dead Nettle*, *Henbit*; naturalized; flowering from May to October; is regarded as tonic, diaphoretic, and laxative.

LAMIUM FOLIOSUM, *L. album*—*l.* *Montanum*, *Melittis melissophyllum*—*l.* *Plinii*. *Melittis melissophyllum*—*l.* *Rubrum*, *Galeopsis*.

LAMOTTE, MINERAL WATERS OF. These thermal springs are in the department of Isère, France. Temperature, 184°.

LAMPOURDE, *Xanthium*.

LAMPROPH'NUS, (*λαμροφ'ωνος*, from *λαμρος*, 'clear,' and *φωνη*, 'voice.' One who has a clear voice.

LAMPSONA, (*λαψων*.) *Lapsana*.

LAMPYRIS NOCTILUCA, (*λαμυρις*, from *λαμπος*, 'to shine,' and *ουρα*, 'a tail.') *Cicindela*.

LANA PHILOSOPHORUM, ('wool of philosophers'.) *Zinci oxydum*.

LANARIA, (from *lana*, 'wool.') *Saponaria*, *Verbascum nigrum*.

LANCE DE MAURICEAU, (F.) An instru-

ment invented by Mauriceau for perforating the head of the fetus. A perforator.

LANCELET, *Lancet*.

LANCEOLA, *Lancet*.

LANCET, *Lance'ola*, *Lancet'ta*, *Schaste'ron*, *Scal'pum chirur'gicum*, *Lancetel*, (Prov.) *Fame*, (F.) *Lancette*, dim. of *lancea*, 'a lance or spear.' A surgical instrument used in the operation of phlebotomy. It is composed of two parts—the handle, (F.) *Chasse*, and the blade, (F.) *Lame*. The former is made of two small plates of ivory, bone, or shell, movable on the blade, for whose preservation they are intended. The blade is formed of well-polished steel. Lancets are made of different shapes; some being broad-shouldered—others, *spear-pointed*. The French distinguished three kinds: 1. *Lancette à grain d'orge*, which, on account of the almost oval shape of its point, makes a large opening. 2. The *L. à langue de serpent*, which is very narrow towards the point; and 3. *L. à grain d'avoine*, which holds a medium station between the two former, and is generally preferred.

The *Abscess Lancet* is merely a large lancet for the purpose of opening abscesses.

The employment of a lancet for opening abscesses, &c., is called *Lancing*.

LANCETTA, *Lancet*.

LANCETTE, *Lancet*.

LANCETTIER, (F.) A lancet case.

LANCINANT, *Lancinating*.

LANCINATING, *Lancinans*, (*lancinase*, *lancinatus*, (from *lancea*, 'a lance,' 'to strike or thrust through.') (F.) *Lancinant*, (substantive, *Élancement*.) A species of pain which consists in lancements or shootings, similar to those that would be produced by the introduction of a sharp instrument into the suffering part. It is especially in cancer that this kind of pain is felt.

LANCING, see *Lancet*—*l.* the Gums, *Déchaussement*, see *Gum lancet*.

LAND'S END, CLIMATE OF. The climate of the Land's End, in England, resembles that of the south of Devonshire, but is more relaxing. It is considered to be most likely to prove beneficial in consumptive cases in which the disease is accompanied by an irritated state of the pulmonary mucous membrane, producing a dry cough. Where the system is relaxed, and the secretion from the lungs considerable, the climate, it is conceived, will generally prove injurious. As a brumal retreat, the southern coast of Devonshire would seem to be preferable to it.

LANGDEBEF, ('ox tongue.') *Anehusa officinalis*.

LANGCEAL, MINERAL WATERS OF. Acidulous mineral waters at Langeac, in the department of Haute Loire, France. They are employed as refrigerant, aperient, and diuretic. They contain carbonic acid, carbonates of soda and magnesia, and a little iron.

LANGENSCHWALBACH, *Schwalbach*.

LANGHII A'QUA EPILEPTICA, *Epilep'tic Water of Lan'gins*, formerly employed against epilepsy. It was composed of the flowers of *concallaria* and lavender, *Spanish wine*, cinnamon, nutmeg, mastic, penny and dittany roots, long pepper, cubebæ, and rosemary flowers.

LANGUE, (from *lingua*.) *Tongue*—*l.* *Abaisseur de la Glossoepatochus*—*l.* de *Carpe*, *Fish-tongue*, see *Lever*—*l.* de *Cœf*, *Asplenium scolopendrium*—*l.* de *Chenal*, *Ruscus hypoglossum*—*l.* de *Chien*, *Cynoglossum*—*l.* de *Serpent*, *Ophioglossum vulgatum*.

LANGUEUR, *Languor*.

LANGUOR, *Apl'esis*, *Faint'ness*, (F.) *Languueur*. A species of atony, depression or debility, which generally comes on slowly.

LANGWORT (Prov.), *Veratrum album*.

LANIARI DENT'ES, (F.) *Dents Laniaræ*, (from *laniare*, 'to tear.') The canine teeth.

LANK (Prov.). Inguen.

LANTANA, *Sage'twee*, *Blue'berry*, (F.) *Civil-teau*. The leaves of this indigenous plant form a finescented tea, like *L. Cam'ara* or *Baha'ma Tea*, and *L. Pseudothea* or *Brazil' Tea*. The tea is used as a diaphoretic.

LANUGO, gen. *Lanuginis*, *Pi'li cu'tis*, *Plu'ma*, (*lana*, 'wool.') The soft, fine hair on different parts of the body, especially in the young.

LANUGO PRIMA, Genies—1. Pudendum, Pubes—1. Siliquæ hirsute, see *Dolichos puriens*.

LANUVIUM, (from *lana*, 'wool.') Vulva.

LAONICA CURATIO seu CU'RA, (*λαος*, 'the people.') A popular mode of treating the gout in the middle ages, "which consisted in the employment of local applications proper for evacuating the morbid matter." [2]

LAOS, Tin.

LAPACTICUS, (*λαπαζειν*, 'to evacuate.') Cathartic, Laxative.

LAPAGMA, (*λαπαγμα*.) An evacuation.

LAPARA, (*λαπαρά*.) Abdomen, Flanks, Lumbi.

LAPARENTEROTOMIA, Laparoenterotomy.

LAPAROCE'LE, *La'paroccele*, (*lapara*, and *κηλη*, 'rupture.') *Lum'bar Her'nia*, through a separation of the fibres of the quadratus lumborum, and a protrusion of the aponeurosis of the transverse muscle on the outside of the mass common to the sacrolumbalis and longissimus dorsi.

LAPAROCYSTOTOMIA, (*lapara*, *κυστις*, 'bladder,' and *τομη*, 'section.') See Lithotomy.

LAPAROELYTROTOMIA, (*lapara*, *ελυτρον*, 'vagina,' and *τομη*, 'section.') Cæsarean section.

LAPARENTEROTOMY, *Lap'aroenterotom'ia*, *Laparenterotom'ia*, (*lapara*, *εντρον*, 'intestine,' and *τομη*, 'incision.') The operation of opening the abdomen and intestinal canal for the removal of disease.

LAPAROSCOPIA, (*lapara*, and *σκοπεω*, 'to examine.') Abdominoscopia.

LAPAROTOMY, *Laparotom'ia*, (*lapara*, and *τομη*, 'section.') Gastrotomy, especially when executed in the lumbar region. See Gastrotomy.

LAPATHUS, (*λαπαθος*.) *Rumex acutus*.

LAPATHUM, (*λαπαθον*.) *L. acutum*, *Rumex acutus*—1. Aquaticum, *Rumex hydrolapathum*—1. Chinense, Rheum—1. Hortense, *Rumex patientia*—1. Orientale, Rheum—1. Pratense, *Rumex acetosum*—1. Sanguineum, *Rumex sanguineus*—1. Scutatum, *Rumex scutatus*—1. Sylvestre, *Rumex acutus*, *R. obtusifolius*—1. Unctuosum, *Chenopodium bonus Henrius*.

LAPAXIS, (*λαπαξις*.) Evacuation.

LAPE, (*λαπη*.) Mucus.

LAPIDILUM, (*lapis*, gen. *lapidis*, 'a stone.') Blasius has given this name to a kind of scoop used for extracting stone from the bladder.

LAPILLATIO, (from *lapillus*.) Lithia.

LAPILLI GLANDULÆ PINEALIS, see Pineal gland.

LAPIL'LUS, (dim. of *lapis*, 'a stone.') A small stone; gravel; a grain of sand.

LAPIS, gen. *Lap'idis*, Calculus—1. Accipitrum, Hieracites—1. Ærosus, Calamina—1. Ammoniaci, see Ammoniac gum—1. Animalis, Blood—1. Armenius, Melochites—1. Aureus, Urine—1. Bezoardicus, Bezoar—1. Cæruleus, Copri sulphas—1. Calaminaris, Calamina—1. Carneolus, Cornelian—1. Causticus, Potassa cum calce, Potassa fusa—1. Collymus, Ætites—1. Contraryervæ, Pulvis contraryervæ compositus—1. Cyanus, *L. lazuli*.

LAPIS DIVI'NUS, *L. Ophthalmicus* seu *Ophthalmicus Sti. Ivesii*. *Cuprum alumina'tum*, (F.) *Pierre divine*, (*Cupri sulphat.*, *Alumin.*, *Potass. nitrat.* 3j. Melt together, adding at the end *Camphor.* ʒss.) Employed to make an eye-water, ʒij ad aquæ ʒiv.

LAPIS FULMINEUS, Coraunion—1. Hematites, Hæmatites—1. Heracleus, Magnet—1. Infernalis,

Argenti nitras—1. *Infernalis alkalinus*, *Potassa fusa*—1. *Infernalis sive septicus*, *Potassa cum calce*—1. *Judaicus*, *Judæus* (*lapis*).

LAPIS LA'ZULI, *L. Cy'anus*, *Asulci*, *Laz'ulite*, (F.) *Pierre d'azur*. A stone, of a beautiful blue colour; opaque; and close-grained; fracture, dull. It is composed of silice; alumina; carbonate and sulphate of lime; oxide of iron, and water. It was formerly looked upon as a purgative and emetic, and given in epilepsy.

LAPIS MALUCENSIS, Bezoard of the Indian porcupine—1. Nauticus, Magnet—1. Ophthalmicus, *L. Divinus*—1. Ophthalmicus *St. Ivesii*, *L. Divinus*—1. Phœnicites, *Judæus* (*lapis*)—1. Porcinus, Bezoard of the Indian porcupine—1. *Prunellæ*, *Potassæ nitras fusus sulphatis paucillo mixtus*—1. *Pumicis*, *Pumex*—1. *Sardius*, *Cornelian*—1. *Septicus*, *Causticum commune*, *Potassa fusa*—1. *Specularis*, *S. lucidum*—1. *Syderitis*, Magnet—1. *Syriacus*, *Judæus* (*lapis*)—1. *Vini*, *Potassæ supertartas impurus*.

LAPPA, (*λαβειν*, 'to lay hold of.') *Arctium lappa*, *Galium aparine*, *Lippitudo*—1. Minor, *Xanthium*.

LAPPULA HEPATICA, Agrimony.

LAP' SANA, (*λαψανη*.) *Lamp'sana*, *Napi'um*, *Papilla'ris her'ba*, *Dockcresses*, *Nippleswort*, (Sc.) *Bo'lgan leaves*, (F.) *Lamp'sane*, *Herbe aux Mamelles*. This plant is a lactescens bitter. It has been chiefly employed, however, as an external application to sore nipples, &c.

LAPSUS PILORUM, ('falling of hairs.') Alopecia.

LAQ, Laqueus.

LAQUE, Lacca.

LAQUEAR, ('an arched roof or ceiling.') Lacunar.

LA'QUEAR VAGI'NÆ, *Fun'dus Vagi'næ* (*laquear*, 'the inward roof of a house'). The part of the vagina in which the cervix uteri terminates.

LA'QUEUS, ('a noose, a snare,') *Fu'nis*. A cord, ligature or bandage, with running knots;—a Noose, a loop. A fillet, *Bro'chos*, *Pach'etos*, (F.) *Lac*, *Laq* on *Lacq*. The term is applied to a bandage or fillet of any kind, attached by means of a loop upon any part, with the view of fixing it; as in certain cases of labour, where a hand or foot presents; or to facilitate extension in luxations and fractures.

Also:—A prominent band in the brain, *Lemnis'cus*, behind the brachium posterius of the corpora quadrigemina, which marks the course of the superior division of the fasciculus olivaris.

LAQUEUS GUT'TURIS, 'Noose of the throat.' Violent inflammation of the tonsils, in which the patient appears as if suffocated by a noose. According to some, gangrenous cynanche.

LARCH, *Pinus larix*.

LARD, (from *lardum*.) Adeps—1. Benzoated, Unguentum Benzoini.

LARDAOE, Lardaceous.

LARDA'CEOUS, *Larda'ceus*, *Lard'ceus*, *Lard'iform*, *Lardifor'mis*, (F.) *Lardacæ*; (from *lardum*.) An epithet given to certain organic alterations in the textures, whose aspect and consistence resemble lard. (F.) *Tissus lardacæ*. They constitute what has been termed the *Larda'ceous* or *Chol'esterinic Disease*. From their waxy appearance the pathological condition has been called *Cero'ma*.

LARDEUS, Lardaceous.

LARDIFORM, Lardaceous.

LARDUM, Adeps.

LARGE, (from *largus*.) Broad.

LARDUM, Adeps.

LARIX, gen. *La'ricis*, (*λαριξ*, gen. *λαρικος*.) *Pinus larix*—1. *Communis*, *Pinus larix*—1. *Decidua*, *Pinus larix*—1. *Europæa*, *Pinus larix*—1. *Pyramidalis*, *Pinus larix*.

LARKSPUR, BRANCHING, *Delphinium consolida*.

LARKE (from Journal) Tour.

LARMOIEMENT, from *larmoyer*, 'to shed tears.' *Exilire*.

LARNAUDÈS, DISINFECTING LIQUID
OF, see ZINC sulphat.

LAROCHE POSAY, MINERAL WATERS
OF. Simple sulphurous waters in the department
of Vienne, France. Season from the 1st of June
to the 15th of September.

LARREA MEXICANA, (after J. A. de Larrea, a Spanish savant.) *L. glutinosum*, *Ononis* plant. Order, Zygophyllaceae. A shrub growing in Arkansas, Southern Texas, New Mexico, and Northern Mexico, which abounds in a strong-scented resinous matter; and is used, externally and internally, in rheumatism, and in syphilitic, and other pains.

LARVA. Mask. Also the larva, grub, or vermiform condition of an insect: the first change it experiences after leaving the ovum. Larvae of insects are occasionally developed in the intestinal canal from one swallowed. See Potosoa.

LARVE, Mosk.

LARFÉ. Masked.

LARYNGÉ, Larvageal

LARYNGE'AL or *Laryng'eal*, *Laryng'eus*, (F. *Laryngé*, *Laryngien*, (from *larynx*.) That which belongs to the larynx.

LARYNGEAL ARTERIES are given off from the thyroid arteries.

LARYNGEAL NERVES. [F.] *Nerfs Laryngaux*, are two in number:—a *superior* and an *inferior*. The *superior laryngeal nerve* is given off from the trunk of the pneumogastric, at the upper and deep part of the neck. It passes downwards and inwards, behind the internal carotid artery, and divides into two secondary branches; the one, *external*, which distributes its filaments, on the outside of the larynx, to the sterni-thyroid, hyo-thyroid, constrictor inferior, crico-thyroid muscles, &c. The other, the *internal*, which crosses the thyro-hyoid membrane, and gives filaments to the epiglottis, the mucous membranes of the pharynx and larynx, to the arytenoid gland, the arytenoid and crico-thyroid muscles, and ultimately intermingles with the inferior laryngeal nerve. The *inferior laryngeal nerves* or *recurrents*, *Nerfs r  versifs*, *Rameaux Trach  aux* (Ch.), arise from the trunk of the pneumogastric within the thorax. They ascend in the furrow, which separates the trachea from the oesophagus, to be distributed on the neck, after having been reflected:—the left around the arch of the aorta; the right, around the corresponding subclavian. They send off filaments to the cardiac plexusses; to the plexus of the oesophagus, and trachea; to the thyroid gland; to the inferior constrictor of the pharynx; the posterior and lateral crico-arytenoid and thyro-arytenoid muscles; and to the mucous membrane of the pharynx and larynx.

LARYNGEAL PHTHISIS. Phtthisis, laryngeal.

LARYNGEAL POUCHES. Small *cysts-à-cave*, which lead from the anterior part of each ventricle of the larynx upwards, for the space of half an inch, between the superior ligaments on the inner side, and the thyroid cartilage on the outer, reaching as high as the upper border of that cartilage at the sides of the epiglottis. Numerous small glands open into them.

LARYNGEAL SOUND. Laryngeeche.

LARYNGEAL VEINS open into the internal jugular. Winslow gives the name *laryngeal* to the superior thyroid artery.

LARYNGEAL VOICE, see Voice.

LARYNGE'CHÉ. (*larynx*, and *syn.* or *son.* 'sound.') The *laryngeal* sound heard by the stethoscope during breathing and speaking.

LARYNGEMPHRAXIS, (*laryngo*, and *emphrax*).

'obstruction.') Obstruction of the larynx from
 SEX CHASE.

LARYNGIEN. Larværeal.

LARYNGISM US. Spasm of glottis, giving occasion to contraction or closure of the opening.

LARYNGISMUS STRIDULUS. Asthma chy-
micum.

LARYNGITE, Laryngitis.—1. *Mucosa*, Laryngitis (simple).—1. *Edematosa*, Edema of the glottis.—1. *Edematosa et adhaerentia*, Edema of the glottis.—1. *avec Production de Fusses membraneuses*, Cynanche trachealis.—1. *Pseudo-membraneuse*, Cynanche trachealis.—1. *avec Stricture de Poi*, Phiditis laryngea.—1. *Sous-mucosa*, Edema of the glottis.—1. *Stridulense*, Asthma thymicum.—1. *Syn-glottica*, Edema of the glottis.

LARYNGITIS. (*Laryngitis*, and *itis*, *Inflam-*
atio Laryngitis, *Quæritur* the sen *Angina laryngis*,
Inflammation of the Larynx, (F.) *Laryngitis*, *C.*
Larynge *laryngien*, *Angine laryngé*. This disease,
in some measure, resembles croup; but is usually
dieroid of that peculiar sonorous inspiration, which
attends the latter. There is, also, pain upon
pressing the larynx: and, whilst laryngitis is a
disease of more advanced life, croup attacks
children. The membraniform exudation is, also,
absent; probably, because the inflammation, being
seated above the glottis, the *conspicible* lymph is
readily expectorated. It requires the most active
treatment.

Simple Laryngitis is called by some authors *Laryngitis*, *Laryngitis acuta* *submucosa* *acuta*, (F.) *Laryngitis muquense*, to distinguish it from *submucous Laryngitis* or *Edema of the glottis*.

Chronic Laryngitis is generally regarded as synonymous with laryngeal phthisis; but it may exist independently.

LARYNGITIS ACUTA, Laryngitis—1. Chronic, see Phthisis Laryngea—1. Exudative, Cynanche trachealis—1. Membranacea, Cynanche trachealis—1. Mucosa acuta, Laryngitis—1. Mucous, Laryngitis (simple)—1. Edematous, Edema of the glottis—1. Pterygosa, Cynanche trachealis—1. Seropurulent, Edema of the glottis—1. Submucous, Edema of the glottis—1. et Tracheitis chronica, see Phthisis laryngea—1. et Tracheitis infantilis, Cynanche trachealis.

LARYNGO (from λαρύξ, gen. λαρύγος), in composition, larvnx.

LARYNGOACE, (*laryngo*, and *ace*, 'bad,')
Cynanche trachealis.

LARYNGOCATARRHUS, *Catarrhus Laryngæus et trachealis*. Catarrh affecting the larynx and trachea more especially, as indicated by alteration of the voice — hoarseness — itching and sensation of burning in those parts; short cough and expectoration, &c.

LARYNGO-ET-TRACHEO-PHTHISIS,
Phtisis laryngea.

LARYNGOGRAPHY. *Laryngographia*: (la-ryngo, and grapho, 'a description.') An anatomical description of the larynx.

LARYNGOLOGY, (*larynx*, and *logos*, 'description.') A treatise on the larynx.

LARYNGOPARALYSIS, see Aphonia.

LARYNGOPHONY, *Laryngophonia*, *Tracheophony*, *Laryngeal voice*, *Tracheal voice*; (*larynx*, and *phon*, 'voice'). The sound heard in health, when the stethoscope is placed over the larynx or trachea at the time a person speaks. The voice appears to pass immediately up to the ear of the auscultator. A similar physical sign exists when there is a cavity in the lungs, and the instrument is placed over it whilst the patient speaks. See *Pneusthony*.

LARYNGOPHTHISIS. *Phthisis laryngea.*

LARYNGOSCOPY. *Laryngoskopia*. (Laryngo, and *scopos*, 'I view.') Inspection of the larynx.

An instrument, on the same principle as the ophthalmoscope, which, by means of a mirror, en-

bles the larynx to be inspected, is termed a *laryngoscope*, *larynx speculum*.

LARYNGOSPASMUS, Asthma thymicum.

LARYNGOSTASIS, (*laryngo*, and *stasis*, 'a settling, or stagnation,') Cynanche trachealis.

LARYNGOSTENOSIS, (*laryngo*, and *stenosis*, 'contraction,') Contraction or narrowness of the larynx.

LARYNGOTOMY, *Laryngotomia*, (*laryngo*, and *tomia*, 'to cut,') A surgical operation, which consists in opening the larynx, either to extract a foreign body, or to remedy an obstruction of the glottis. The operation is, sometimes, erroneously called *Bronchotomy*, and *Tracheotomy*.

LARYNGOTRACHEITIS, Cynanche trachealis—1. Tracheitis with Diphtheritic exudation, Cynanche trachealis.

LARYNX, gen. *Laryngis*, (*λάρυγξ*, gen. *λάρυγγος*), *Caput seu Operculum seu Intimum seu Finis superior seu Terminus superior seu Pars prima asperæ arteriæ*. The apparatus of voice is situate at the superior and anterior part of the neck; and at the top of the trachea, with which it communicates. It is composed of four cartilages,—the thyroid, cricoid, and two arytenoid; is moved by a number of muscles, and lined by a mucous membrane, having certain membranous reflections, constituting the superior and the inferior ligaments of the glottis, &c.

The larynx is destined to give passage to the air, in the act of respiration, and to impress upon it certain modifications, which constitute voice. Its dimensions vary in different individuals. In men, it is always larger, and situate lower, than in women.

LARYNX, PELLICULAR or PLASTIC INFLAMMATION OF THE, Cynanche trachealis.

LARYNX SPECULUM, Laryngoscope.

LAS'ANUM, (*λασάνη*), *Aph'edra*, *Aph'edron*, *Sella familiar'ica*, (F.) *Chaise percée*. A close stool.

LASCIVIA, (from *lascivus*), Satyriasis.

LASCIVITAS, Satyriasis.

LASCIVUS, Libidinous.

LASER, *Laserpitium*—1. *à Grandes feuilles*, *Laserpitium latifolium*.

LASERPIT'IUM, *La'ser*, *Sil'phium*, *σός αλφειον*. A term applied, anciently, both to a plant and its juice, regarding the nature of neither of which we possess any precise information. Bentley, Laurence, Geoffroi, &c., regard it to have been the same as asafetida:—Theophrastus, Dioscorides, and the ancient scholiast of Aristophanes, however, by assigning a sweet and agreeable flavour to the laserpitium, discountenance the idea. From whatever plant obtained, it was so rare, and consequently so costly, that the Romans deposited it in the public treasury. It was obtained from Cyrene—*Succus Cyrena'icus*—and likewise from Persia—the latter being the most valuable. The Laserpitium is called by Avicenna, *Althit*.

LASERPITIUM ASPERUM, L. latifolium.

LASERPITIUM LATI'FOLIUM, L. *asperum*, *Gentiana alba*, *White Gentian*, *Cervaria alba*; (F.) *Laser à larges feuilles*. The root of this plant is bitter and tonic.

LASERPITIUM MONTANUM, L. *siler*.

LASERPITIUM SIL'ER, L. *trifoliatum* seu *montanum*, *Ses'eli*, *Seselis*, *Sil'ler montanum* seu *lancifolium*, *Heart'wort*, *Sermountain*. The seeds and roots are possessed of aromatic properties.

LASERPITIUM TRIFOLIATUM, L. *siler*.

LASK, Diarrhoea.

LASSITUDE OCULAIRE, Copyopia.

LASSITUDO, (*lassus*, 'tired,') Copos.

LAST, Inguen.

LATENS IN ORE, ('lying hid in the mouth,') *Pterygoideus internaus*.

LATENT, *La'tens*, (*latere*, 'to lie hid,') 'lying

hid,' 'concealed.' An epithet applied to certain diseases or states of disease, in which the symptoms are so concealed and obscure, *morbi occulti*, as to escape the observation of the physician. Thus, we say *latent inflammation*, *latent period of smallpox*.

LAT'ERAD, (*latus*, gen. *lateralis*, 'the side,') A term used averbally by Dr. Barclay to signify 'towards the lateral aspect.'

LATERAL ASPECT, see Mesial.

LATERIT'IOUS, *Laterit'ius*, *Lateric'ius*, (F.) *Briqueté*, (*later*, gen. *lateralis*, 'a brick,') An epithet applied to the bricklike sediment, occasionally deposited in the urine of people afflicted with fever; *Sedimentum urinae lateric'ium*;—(F.) *Urine briqueté*.

LAT'EROFLEX'ION, *Lat'eroflex'io*, (*latus*, gen. *lateralis*, and *flexio*, 'a bending,') A bending to one side,—as of the uterus.

LATÉRO-VERSION, (F.) *Lat'erover'sio*. A turning or version to one side,—as of the uterus.

LATESCENTIS CHORDÆ MUSCULUS, ('muscle of the hidden cord,') *Palmaris longus*.

LATEX NIVEUS, ('snowy liquor,') Milk.

LATHYRIS, Euphorbia lathyris.

LATIB'ULUM, (from *lateo*, 'I lie hid,') The foyer of a febrile poison; whence it spreads to every part to induce a paroxysm. See *Clavier*.

LAT'ICA, (from *lateo*, 'I lie hid,') *Amphimer'ina lat'ica* of Sauvages. A species of quotidian remittent, whose paroxysms are very long, and which is accompanied with latent or slightly developed fever,—whence its name.

LATICES LACTEI, Receptaculum chyli.

LATISSIMUS COLLI, (superlative of *latus*, 'broad,') *Platysma myoides*.

LATIS'SIMUS DORSI, *Aniscalp'tor*, *Bra'chium movens quartus*, (F.) *Lombo-huméral* (Ch.), *Dorsilombo-sacro-huméral*, *Muscle grand dorsal*, M. *très large du dos*. A flat muscle; broad, especially below; thin; quadrilateral; and situate at the posterior, lateral, and inferior region of the trunk. It is attached to the posterior half of the outer lip of the crest of the ilium; to the posterior surface of the sacrum; to the spinous processes of the six or seven last dorsal vertebrae, to all those of the loins, and to the last four false ribs, and is inserted by a strong tendon at the posterior edge of the bicipital groove of the humerus. Its upper fibres are almost horizontal; the middle very long and oblique upwards and outwards; and the anterior almost vertical. This muscle carries the arm backwards, depressing it, and making it turn on its axis. It also draws backwards and downwards the prominence of the shoulder. When, suspended by an arm, we make an effort to raise ourselves, it draws the trunk towards the arm. It can, also, raise the ribs by assuming its fixed point on the humerus, and become an inspiratory muscle.

LATITUDO HUMERI, (from *latus*, 'broad,') Scapula.

LATTICE WORK, (from (F.) *latte*, 'a lath,') Cancelli.

LATUS, Broad—1. Ani, Levator ani.

LAU, see Spirit.

LAUCA'NTA, (*λαυκάνη*), *Leuca'nia*, *Lau'chané*. The fauces and oesophagus. Also, the chin.

LAUDANISÉ, see Opiated.

LAUD'ANUM or LAUDA'NUM. (Perhaps, from *laus*, gen. *laudis*, 'praise; lauda'tum, 'praise-worthy,') Every preparation of opium, solid or liquid, but more particularly the extract and tincture, and especially the latter.

LAUDANUM ABBA'TIS ROUSSEAU, *Gut'tæ Abba'tis Rousseau*, *Vinum opiatum fermentatio'nē paratū*, *Abbe Rousseau's Drops*. (*Mel. Narbonne*, 3xij; *Aque calidæ*, Oij. Set in a warm place, and, as soon as the mixture ferments, add *Opium*, 3iv, dissolved in *Water*, f 3xij. Let it ferment for

a month, and evaporate to $\frac{1}{3}$ x: strain, and add Rectified spirit of wine, $\frac{1}{3}$ vss.

LAUDANUM. FORD'S. Vinum opii—1. Liquid, Tinctura opii—1. Liquidum Hoffmanni. Vinum opii—1. Liquidum Sydenhami. Vinum opii—1. Opium. Extractum opii—1. Simplex, Extractum opii—1. Sydenhami. Vinum opii.

LAUGH. (Sax. hlahan, by onomatopœia,) Rhus—1. Sardonis. Canine laugh.

LAUGHING. Rhus.

LAUGHTER. Rhus.

LAUREL, *Kalmia latifolia*, *Laurus*, *Magnolia macrophylla*—1. Broadleaved, *Kalmia latifolia*—1. Cherry, *Prunus laurocerasus*—1. Common, *Prunus laurocerasus*—1. Dwarf, *Kalmia angustifolia*—1. Great, *Rhododendron maximum*—1. Ground. *Epigaea repens*—1. Mountain, *Kalmia latifolia*, *Rhododendron*—1. Narrow-leaved, *Kalmia angustifolia*—1. Pale, *Kalmia glauca*—1. Poison. *Prunus laurocerasus*—1. Rose, *Kalmia latifolia*—1. Sheep. *Kalmia angustifolia*—1. Swamp, *Kalmia glauca*—1. Sweet, *Illicium Floridanum*—1. Water. See *Prunus laurocerasus*—1. White, *Magnolia glauca*.

LAURENT. SAINT, MIN'ERAL WATERS OF. A thermal spring, five leagues from Joyeuse in France. Temp. 127° Fahr.

LAURENTIA PINNATIFIDA, *Pepperdulse*. One of the Algae, eaten in Scotland, and hawked about the streets of Edinburgh along with *Laminaria digitata* or Tangle.

LAUREOLA, (dim. of *laurus*, 'laurel,') *Daphne laureola*.

LAURIER, *Laurus*—1. *Alexandrin*, *Ruscus hypoglossum*—1. *Amandier*, *Prunus laurocerasus*—1. *d'Apollon*, *Laurus*—1. *Cerise*, *Prunus laurocerasus*—1. *de Ceylon*, *Laurus cinnamomum*—1. *Commun*, *Laurus*—1. *des Cuisines*, *Laurus*—1. *des Montagnes*, *Kalmia latifolia*—1. *Noble*, *Laurus*—1. *Rose*, *Nerium oleander*—1. *de Trébizonde*, *Prunus laurocerasus*.

LAUROCERASUS, (*laurus* and *cerasus*, 'the cherry tree,') *Prunus laurocerasus*.

LACROSE, (from *laurus*.) *Nerium oleander*.

LAURUS, *Laurus nobilis*, *Daphne*, *Laur'el*, *Sweet Bay*. *Nat. Ord.* Lauraceæ. (F.) *Laurier*, *L. noble* ou *des cuisines* ou *commun* ou *d'Apollon*. The leaves and berries, *Laurus* (Ph. L.), have a sweet, fragrant smell, and an aromatic, astringent taste. Sweet bay has been advised as a stomachic and carminative, but is now rarely used. It is, sometimes, employed as a fomentation and in gylsters; and the berries are an ingredient in the *Emplastrum Cumin'i*.

LAURUS ÆSTIVALIS, *L. benzoïn*—1. *Alexandrina angustifolia*, *Ruscus hypoglossum*.

LAURUS BENZOÏN, *Benzoïn odoriferum*, *Laurus Pseudobenzoïn* seu *æstivalis*, *Spice wood*, *Spice bush*, *Ben'jamin bush*, *All'spice bush*, *Wild all'spice*, *Spice berry*, *Fever wood*, *Fever bush*. An indigenous shrub, growing in moist, shady places, in all parts of the United States; flowering early in spring. All parts of the shrub have a spicy, agreeable flavour, which is strongest in the bark and berries. An infusion or decoction of the small branches is used in popular practice as a vermifuge, and an agreeable drink in low fevers. The bark has been used in intermittents; the berries, dried and powdered, for allspice. The oil of the berries is used as an excitant.

LAURUS CAMPHORA, see *Camphor*—1. *Canella*, *L. Cassia*.

LAURUS CAS'SIA, *L. Canel'la*, *Per'eca cassia*. The species of *laurus* which yields the *Ca'sia*, *Can'cia*, *Canel'la*, *Cassia lignea* seu *cinnamomea*, *C. lignea Malabar'ica*, *Cortex Canel'la Malabar'ica*, *Cortex coras'ior*, *Xyl'oca'sia*, *Canel'la Malabar'ica* et *Javan'ica*, *C. Cubana*, *Ar'bor Juda'ica*, *Cannell'era Malabar'ica*, *Cinnamo'mum Malabar'icum* seu *Ind'icum* seu *Sinen'se*, *Culihacha canel'*

la, *Wild Cin'namon*, *Malabar' Cinnamon*, *Cassia*, (F.) *Cannelle de Malabar* ou *de Java* ou *de la Chine* ou *des Indes* ou *de Caramandel*, *C. javanica*, *C. malle*, *Casse en bois* ou *aromatique*. The bark and leaves abound with the flavour of cinnamon, for which they may be substituted; but they are much weaker. The unopened flowerbuds are used in the same manner. *AGUA CAS'SIA*, *Water of Cassia*, prepared by distillation, is official in the Edinburgh Pharmacopœia.

LAURUS CINNAMOMOIDES, *Nectandra cinnamomoides*.

LAURUS CINNAMOMUM, *Per'eca cinnamomum*, *Cinnamo'mum*, *C. Zeylan'icum*, *Dar'seni*, *Darsi'ni*, *Cin'namon*, *Xyl'ocinnamo'mum*, *Cu'rdo Cannel'la*, (Sc.) *Cannel*, (F.) *Cannelle*, *Baume de Cannelle*, *Cannelle officinal* ou *de Ceylon*. Cinnamon bark, which is obtained, also, from *Cinnamo'mum aromaticum*, is stimulant and carminative, and is employed, chiefly, as a grateful aromatic, to cover the taste of nauseous remedies. Dose, gr. x to $\frac{3j$. The Flowers, called *Cassia Flores* in the shops, possess aromatic and astringent virtues, and may be used wherever cinnamon is required. The volatile oil of the bark—*Oleum Cinnamo'i*—*Oil of Cinnamon*, is official in the Pharm. U. S.

LAURUS CUBEBA, *Piper cubeba*—1. *Culilaban*, *L. Culilawan*.

LAURUS CULILAWAN, *L. Culilaban* seu *Caryophyllus*, *Cinnamo'mum Culilawan*. The tree that affords the *Cortex Culilawan* seu *Culilaban*, *Culilawan*, *Culitawan*, *Cortex caryophyllodes* seu *Caryophylloides Amboinensis*. This bark resembles the sassafras in appearance and properties, and is used in Java as a condiment.

LAURUS MALABATHRUM, see *Malabathrum*—1. *Nobilis*, *Laurus*—1. *Persea*, *Persea gratissima*—1. *Pseudobenzoïn*, *L. Benzoïn*.

LAURUS SAS'SAFRAS, *Per'sea sas'sifras*, *Sassafras*, *S. officinalis*, *Cor'nus mas odora'ta*, *Ambiba*, *Aguafree*, *Aque-tree*, (F.) *Pavanne*. Indigenous in the United States. Sassafras wood and root, and especially the bark of the root, *Sassafras Radicis Cortex* (Ph. U. S.), have been considered stimulant, sudorific, and diuretic. The virtues depend upon essential oil, *Oleum Sas'safras*, the odour of which is not unlike that of fennel. It has been used in cutaneous diseases, chronic rheumatism, &c., in the form of infusion, which has also been taken as tea. See *Saloop*.

The pith of the stems, *Sassafras medulla*, (Ph. U. S.), abounds in gummy matter, which it readily imparts to water, forming a limpid mucilage, which is much employed as a collyrium in ophthalmia, and as a drink in dysentery, catarrh, &c. (one drachm of the pith to a pint of boiling water).

LAUTISSIMA VIN'A. Superlative of *laurus*, 'elegant.' Wines were formerly so called, which were strongly impregnated with myrrh.

LAVACRUM OPHTHALMICUM, (from *lavare*, 'to wash,') see *Collyrium*.

LAVAMENTUM, (from *lavare*, 'to wash,') *Clyster*.

LAVANDE, *Lavandula*—1. *de Mer*, *Statice limonium*—1. *Triste*, *Statice limonium*.

LAVAN'DULA, *Laven'dula*, (from *lavare*, 'to wash,' from being used in baths.) *Lavan'dula sp'ica* (Linnaeus) seu *angustifolia* seu *officinalis* seu *vera*, *Com'mon Lav'ender*, (F.) *Lavandula*, *Ord.* Labiatæ. The odour of lavender flowers, *Lavandula* (Ph. U. S.), is fragrant and agreeable; taste warm and bitterish—depending upon an essential oil. It has been used as a stimulant; particularly in the form of the oil,—*Oleum Lavan'dula*. The dried leaves have been employed as an emmenagogue.

The French use, also, *LAVANDULA STOE'CRAS*, *Stoe'chas* et *Stoe'chas Arab'ica*, *French Lav'ender*, of which they have a compound syrup, *Syrupus de stoe'chaud compos'itus*; given as a pectoral.

LAVANDULA LATIFOLIA, Lavandula, L. Spica.
LAVANDULA SPI'CA (Linnaeus), *L. latifolia*, Spica, French *Lav'ender*, (F.) *Lavande*, *Aspic*, Spic, yields the Oil of Spike, *O'leum spi'ce*, (F.) *Huile d'Aspic* ou de Spic, which is used to make varnishes, and as a liniment in paralysis.

LAVEMENT, (from [L.] *lavare*, 'to wash,') Clyster, Enema—*L. Antispasmodique*, Enema foetidum.

LAVENDER, COMMON, Lavandula—*L. French*, Lavandula spica—*L. Sea*, Statice Caroliniana, Statice limonium,

LAVENDULA, Lavandula.

LAVER, Ulva latissima—*L. Broad green*, Ulva latissima—*L. Germanicum*, Veronica beccabunga—*L. Laciniated purple*, Ulva umbilicalis—*L. Odoratum*, Sisymbrium nasturtium.

LAVIPEDIUM, (*lavare*, 'to wash,' and *pes*, gen. *pedis*, 'foot,') Pediluvium.

LAW MEDICINE, Medicine, legal.

LAWSONIA ALBA, (after Dr. J. Lawson, author of a Voyage to Carolina,) *L. Inermis*.

LAWSONIA INER'MIS, *L. alba*, *Alcan'a ve'ra* seu *Oriental'is*, *Cyp'e'rus antiquo'rum*, *Liquis'trum Egyptiacum*, Smooth *Lavso'nia*, *Henna*, (F.) *Henné*, *Alhaune*. An East Indian and African plant, the root of which is slightly astringent.

In India, the root of the *Lavso'nia spinosa* is employed in lepra and other cutaneous affections.

LAX, (from *laxus*, 'loose,') Diarrhoea.

LAXANS, (*laxare*, *laxatum*, 'to loosen,') Laxative, Relaxant.

LAXATIF, Laxative.

LAX'ATIVE, *Laxati'vus*, *Lax'ans*, *Le'niens*, *Minorati'vus*, *Soluti'vus*, *Alvidi'vus*, *Hypochore'ticus*, *Hypog'o'gus*, *Hypel'atus*, *Hypocathar'ticus*, *Lapac'ticus*, (F.) *Laxatif*, *Minoratif*. A medicine which gently opens the bowels; such as tamarinds, manna, &c.

LAXATIVUS INDICUS, Cambogia.

LAXATOR AURIS INTERNUS, *L. tympani*.

LAXATOR TYM'PANI, *L. ma'jor tym'pani*, *Exter'nus* seu *Anterior mall'ei*, *Obli'quus* seu *Externus aur'is*, *Laxator aur'is inter'nus*, *Eusta'chii mus'culus*, (F.) *Antérieur du marteau*, *Aconstico-malléen*, *Sphéno-salpingo-malléen*. A muscle which arises from the spine of the sphenoid bone and from the cartilage of the Eustachian tube, and is inserted, by a tendon, into the apophysis of Rau. It relaxes the membrana tympani.

LAXATOR TYM'PANI MI'NOR. A very small muscle which extends from the upper part of the external auditory canal, and is inserted at the inferior part of the process of the handle of the malleus. Its existence is denied by most anatomists.

LAXITAS, Atony, Laxity—*L. Alvi*, Diarrhoea—*L. Ingestorum*, Lientery—*L. Intestinorum*, Lientery—*L. Scroti*, Rachosis—*L. Ventriculi*, Gasterasthenia.

LAX'ITY, *Lax'itas*, *Lax'ness*. Condition of a tissue when loose or relaxed; or of one which wants tone. We say *laxity of fibre*, *laxity of skin*, to express that those parts have lost some of the senseness proper to them.

LAY'ER, AN'IMAL, ('that which is laid,') see *Tache embryonnaire*—*L. Mucous*, see *Tache embryonnaire*—*L. Serous*, see *Tache embryonnaire*—*L. Vascular*, see *Tache embryonnaire*—*L. Vegetative*, see *Tache embryonnaire*.

LAYERS OF THE BLASTODERMA, see *Tache embryonnaire*.

LAYMAN, see Laity.

LAZARET'TO, *Laz'aret*, *Laz'zarhouse*, ([I.] *lazzo*, 'a leper,' from *Lazarus*.) A solitary edifice in most seaports of magnitude, intended for the disinfection of men and goods proceeding from places where communicable diseases are prevailing.

LAZULITE, Lapis lazuli.

LEACH, ([Sax.] *læce*, [Da.] *læge*.) Physician.

LEACHCRAFT, Medicina.

LEACHMAN, Physician.

LEAD. (Sax. *læc*.) Plumbum—*L. Black*, Graphites—*L. Chloride* of, Plumbi chloridum.

LEAD COLIC, see Colica metallica.

LEAD, IODIDE of, Plumbi iodidum—*L. Nitrate* of, Plumbinitras—*L. Oxyd* of, semivitrified, Plumbi oxydum semivitreum.

LEAD PARALYSIS, see Palsy, lead.

LEAD POISONING, *Molybdo'sis*, *Molybdo'nosus*, *Mor'bus plumb'eus*, *Cucochym'ia plumbea*, (F.) *Intoxication saturnine*. Morbid phenomena induced by lead received into the system.

LEAD, RED, Plumbi oxydum rubrum.

LEAD RHEU'MATISM, *Lead Neural'gia*, *Arthral'gia* of M. Tanquerel. The neuralgic and spasmodic pains caused by the poison of lead.

LEAD, SUBCARBONATE of, Plumbi superacetatis—*L. Tannate* of, see Tannin.

LEAD WATER, Liquor Plumbi subacetatis dilutus.

LEAD, WHITE, Plumbi subcarbonas.

LEADER, ('that which leads,') Tendon.

LEADWORT, Plumbago Europæa.

LEAF, SOUR, Andromeda arborea.

LEAM'INGTON, MIN'ERAL WA'TERS OF. Saline waters at Leamington, about two miles east of Warwick, England, which contain chloride of sodium, sulphate of soda, and chlorides of calcium and magnesium.

LEANNESS, (from Sax. *hlæne*, 'wanting in flesh,') Emaciation.

LEAP, (Sax. *hlæpan*.) *Sal'tus*, *Salit'io*, *Exal'sis*, (F.) *Saut*; Bound, Jump,—the act of leaping. Muscular movement or movements, by which the body is detached from the soil by the forcible and sudden extension of the lower limbs, previously flexed upon the pelvis.

LEAPERY, Lepra.

LEAPING A'GUE. This disease is said by the Scotch writers to be characterized by increased efficiency, but depraved direction, of the will, producing an irresistible propensity to dance, tumble, and move about in a fantastic manner, and often with far more than the natural vigour, activity, and precision. See Mania, dancing.

LEATHER FLOWER, Clematis viorna—*L. Wood*, *Direa palustris*.

LEB'ANON, MIN'ERAL WA'TERS OF. The spring at Lebanon, 26 miles east of Albany, New York, is a simple thermal; temperature, 73° Fah.

LECANE, (*λεκανη*, 'a bowl,') Pelvis.

LEC'HE, (Sax. *læce*.) Physician.

LECHENEION, (*λεχηνειον*) Torcular Herophili.

LECHEROUS, Libidinous.

LECHO, (*λεχω*.) Puerpera.

LECHOPYRA, (*lecho*, and *pyr*, 'fire or fever,') Puerperal fever.

LECITHODES, (*lecithos* and *odes*.) Vitelline.

LECITHOS, (*λειδος*, 'yolk of egg,') see Ovum.

LECO, Penis.

LECON'TIA VIRGINICA, (after Major Leconte, U. S.) *Peltandra Virginica*.

LE CRAN, Cochlearia armoracia.

LECTISTER'NIUM, (*lectus*, 'a bed,' and *sternere*, 'to spread,') The arrangement of a bed so as to adapt it to a particular disease. Also, a supplication, with the Romans, in times of public danger, when beds or couches were spread for the gods, as if they were about to feast, and their images were taken down from their pedestals and placed upon these couches around the altars. The lectisterium was first introduced in the time of a pestilence.

LECTUA'LIS, (*lectus*, 'a bed,') An epithet applied to a protracted disease.

LECTULUS MEDICATUS, (dim. of *lectus*, 'a bed,') see Fomentation—*L. Stramineus*, *Fanon*.

LEDE SAUVAGE, *Ledum sylvestre*.

LEDOYEN'S DISINFECTING LIQ'UID or FLU'ID. A solution of nitrate of lead, (*Plumb.*

nitrat. $\frac{5}{16}$ ad aqua $\frac{1}{5}$ l. used as an antiseptic and antihæmorrhagic.

LEDUM LATIFOLIUM, *Labrador's Tea*, *Ord.* Ericaceæ, grows in damp places, in Canada and the United States. The leaves have a pleasant odour and taste, and have been used as tea. They have also been esteemed pectoral and tonic.

LEDUM PALUSTRE, *Rosmarinus* seu *Anthon's eyebright*, *Marsh Tea*, (F.) *Lêde* ou *Rosmarin sauvage*, *Oliver de Bohême*, *Nat. Ord.* Ericaceæ. *Sec. Syst.* Decandria Monogynia. This plant has a bitter, substringent taste, and was formerly used in Switzerland in place of hops: the virtues are equivocal. The leaves are chewed by the Cree Indians, and applied to burns, which are said to heal rapidly under their influence. The cake of chewed leaves is left adhering to the sore until it falls off.

LEE, ([F.] *lie*, from *limus*, 'mud,' [?]) Feculence, Lixivium, Lye, Urine.

LEECH, *Hirudo*, see Ectozoa, Physician.

LEECHCRAFT, *Medicina*.

LEECH, HORSE, *Hippiater*.

LEEK, (Sax. *leac*.) *Allium porrum*.

LEES, SOAP, *Liquor potassæ*.

LEG, ([from [Sax.] *legan*, 'to place,' [Da.] *Læg*, *Cons.* *Soc'us*, *Gn'më*. The portion of the lower extremity, which extends from the knee to the foot. It consists of three bones, *Tibia*, *Fibula*, and *Patella*, and also of a great number of muscles, vessels, and nerves. The projection, formed by the muscles at the back part of the leg, has received the name of *Calf of the leg*. The leg is the special attribute of man, and proves that he is destined to be biped.

LEG ASLEEP. A familiar name for the obtunding of sensation produced by pressure on the great nerve of the leg. As the nerve fibres gradually recover from the effect of the pressure, a pricking sensation is experienced, familiarly known under the name *pins and needles*.

LEG, COCHIN, see Elephantiasis—*L. Swelled*, *Phlegmatia dolens*.

LEGITIMATE, *Legitimus*, (F.) *Légitime*, (*lex*, gen. *legis*, 'law,' *Gn'minus*, *Gn'esinus*. An epithet applied to things which are according to rule. *A legitimate child*, (F.) *Enfant légitime*, is one conceived or born during marriage. *Legitimate diseases*, (F.) *Maladies légitimes*, are those which follow a regular march.

LEG'NA, (*λεγω*, *λεγων*, 'a fringed edge,' 'the border of a garment,') The orifice of the pudendum mulieb're, or of the uterus.

LÉGUME, *Legumen*.

LEGUMEN, gen. *Leguminis*, (*lego*, 'I gather' [?], akin to *λεγω*, 'pulse,') (F.) *Légume*, *Gousse*. So called because it is usually gathered by the hand, instead of being reaped. All kinds of pulses, as peas, beans, &c., are thus termed.

LEGUMIN, see Casein.

LEICHEN, (*λεγω*,) *Lichen*.

LEIMANTHIUM VIRGINICUM, (*λεωω*, 'a meadow,' and *ανθος*, 'a flower,') *Melanthium Virginicum*.

LEIOPUS, *Λέωπος*, *Plan'cus*, *Plan'tus*, *Plo'tus*, (*λεω*, 'smooth,' and *πους*, 'a foot,') One who is affected with flat footedness, *splay footedness*, *Leopodia*, *Liopodia*. One, the soles of whose feet are flat, instead of having the concavity which they commonly present.

LEIOSIS, (*λεω*, 'ground,') *Comminution*.

LEIPHÆMIA, *Liphæmia*, (*leipo*, and *αἷμα*, 'blood,') Poverty or paucity of blood.

LEIPHÆMI, *Achroi*.

LEIPHÆMOS, *Liphæmos*, (F.) *Leiphème*. A word sometimes used adjectively; at others, substantively, either for a vicious state of the blood—or rather for a sort of anæmia—or for the pa-

tient who labours under this condition. See *Achroi*.

LEIPHÈME, *Leiphæmos*.

LEIPO, (from *λεω*, 'I fail, I want,') in composition, want, defect.

LEIPODERMIA, (*leipo*, and *δέρμα*, 'skin,') *Aposthia*.

LEIPODERMUS, *Lipoder'mus*, *Leipoder'matus*. One who wants a part of his skin. It is especially applied to one who wants the prepuce. See *Apella*, and *Aposthia*.

LEIPOMERIA, *Lipomeria*.

LEIPOPSYCHIA, (*leipo*, and *ψυχη*, 'breath of life,') *Syncope*.

LEIPOTHYMIA, (*leipo*, and *δυναμις*, 'mind,') *Syncope*.

LEIPYRIAS, (*leipo*, and *πυρ*, 'fire,' or 'heat,')

A species of continued fever, referred to by the Greek physicians, in which there is burning heat of the internal parts and coldness of the extremities. *Avicenna* described, under this name, a kind of hemitritæa.

LEIRION, (*λεῖριον*,) *Lilium candidum*.

LEMA, (*λεμα*,) *Chassie*.

LEME, (*λεμα*,) *Lippitudo*.

LEMITHOCORTON, *Corallina Corsicana*.

LEMMA, gen. *Lemmatis*, (*λεμμα*,) *Cortex*, *Feculence*, *Purfur*, *Sediment*.

LEMNISCUS, (*λεμνισκος*,) *Pessary*, *Tent*. See *Laqueus*.

LEMON, see *Citrus medica*—*L. Ground*, *Podophyllum montanum*—*L. Juice*, see *Citrus medica*—*L. Juice*, artificial, see *Citrus medica*—*L. Peel*, see *Citrus medica*—*L. Tree*, *Citrus medica*.

LEMON, WILD, *Podophyllum montanum*.

LEMONADE, *Limona'da*, (from *lemon*,) (F.) *Limonade*, *Citronade*. *Lemon juice* diluted with water and sweetened. See *Citrus medica*.

LEMONADE, DRY, (F.) *Limonsade sèche*. *Citric* or *tartaric acid* reduced to powder and mixed with sugar. *Lemonade powders* may be made as follows:—Pound $\frac{3}{4}$ of *citric acid* with a few drops of essence of *lemon-peel* and $\frac{5}{8}$ of more of lump sugar. Divide into six papers, each of which will make a glass of lemonade. See *Citric acid*.

Limonsade Gazeuse (F.), is an agreeable drink prepared by adding syrup of lemons, raspberry, &c., to water saturated with carbonic acid.

LEMONADE, MAGNESIAN, *Magnesiæ citras*.

LEMONADE, NITRIC. *Nitric acid* considerably diluted with water, and sweetened.

LEMONADE, SULPHURIC, and **LEMONADE, TARTARIC**, are made with the sulphuric and tartaric acids.

LEMOSITAS, (from *lema*,) *Chassie*, *Lippitudo*.

LENDE, (Sax. *leno*,) *Lumbi*.

LENDIS, *Lumbi*, *Nates*.

LENIENS, (*lenio*, *lenitum*, 'to assuage,') *Laxative*, *Lenitive*.

LENIS, ('gentle,') *Lenitive*.

LENITIVE, *Lenitivus*, *Len'ia*, *Len'iens*, *Mit'igans*, (F.) *Lenitif*. A medicine which allays irritation or palliates disease: also, a laxative medicine. A *lenitive electuary* is one that purges gently.

LENOS, (*λενος*,) *Torcular Herophili*.

LENS, gen. *Lenitis*, 'a lentil,' *Ervum lens*—*L. Crystalline*, *Crystalline*—*L. Esculenta*, *Ervum lens*.

LENT ROSE, *Narcissus pseudonarcissus*.

LENTICULA, (dim. of *lens*,) A freckle. Also, the eruption of lenticular fever. See *Ephelides*.

LENTICULAIRE, *Lenticular*.

LENTICULAR, *Lenticular'is*, (F.) *Lenticulaire*. Having the shape of a lens or lentil.

LENTICULAR GANGLION, *Ophthalmic G.*

LENTICULARES GLANDULÆ, *Lenticula' glands*. Mucous follicles, having the shape of a

lentic, which are observed especially towards the base of the tongue.

LENTIGO, gen. *Lentig'ina*, (from *lens*), Ephe-lides.

LENTIL, *Ervum lens*.

LENTILLE, *Ervum lens*—*l. Crystalline*, Crystalline—*l. Ers. Ervum ervilia*.

LENTIS/CINUM VINUM, (from *Lentiscus*, 'the mastich tree.') Wine impregnated with mastich.

LENTISCUS VULGARIS, *Pistacia lentiscus*.

LENTITIA, *Lentor*.

LENTITIS, (*lens*, gen. *lensis*, and *itis*), Phacitis.

LENTITUDO, *Lentor*.

LENTOR, *Lentit'ia*, *Lentitu'do*, (*lentus*, 'clammy,') A viscosity or siziness of any fluid. See Gluten.

LENUM, (*ἄλως*, 'a pool or lake,') Torcular Herophili.

LEON, (*ἰών*, gen. *λεωνος*, 'a lion,') in composition, lion.

LEONOTIS LEONURUS, (*leon*, and *ovs*, gen. *ωτος*, 'ear,') *Ord. Labiata*. A South African plant, which has a peculiar smell and nauseous taste, and is said to produce narcotic effects if incautiously used. It is employed in decoction in chronic cutaneous diseases. The Hottentots smoke it like tobacco, and take a decoction of its leaves as a strong cathartic. It is also given as an emmenagogue. In the eastern districts of the Cape Colony, *Leonotis ovata*, (Prov.) *Crowpar-nep*, *Male*, is used for the same purpose.

LEONOTIS OVATA, see *Leonotis leonurus*.

LEONTIASIS, *Leontion*, (*λεων*, 'a lion,') A name given to lepra of the face, from some fancied resemblance between the countenance of those labouring under it and that of the lion. To this kind of lepra the epithets *le'onine* and *le'ontine* have been given.

LEONTICE THALICTROIDES, *Caulophyllum thalictroides*.

LEONTION, *Leontiasis*.

LEON'TODON TARAX'ACUM, (*leon*, and *odous*, gen. *odōros*, 'a tooth,') *L. officinalē* seu *vulga'rē*, *Tarax'acum officinalē* seu *Dens Leonis*, *Dens leonia*, *Hedyp'nois tarax'acum*, *Urina'ria*, *Caput Mon'achi*, *Dan'delion*, *Piss-a-bed*, (Prov.) *Shitabed*, *Puffball*, (Sc.) *Dentiloun*, (F.) *Pissenlit*, *Liondent*, *Dent de lion*, *Couronne de Moine*. *Order*, *Compositæ*. The young leaves are sometimes eaten as salad. The roots are, also, roasted and used as a substitute for coffee. The root, *Tarax'acum*, (Ph. U. S.) is, moreover, reputed to be aperient and diuretic; hence its vulgar names. Its efficacy is doubtful.

LEONTOPODIUM, (*leon*, and *πους*, gen. *πόδος*, 'a foot,') *Alchemilla*.

LEONURUS, (*leon*, and *oupa*, 'a tail,') *Lion's Tail*.

LEONURUS CARDI'ACA, *Agripal'ma Gall'is*, *Marrubium*, *Cardi'aca crispa* seu *triloba'ta* seu *vulga'ria*, *Her'ba cardi'aca* seu *Pat'e lupinæ*, *Motherwort*, *Throatwort*, (F.) *Agripaume*. *Ord. Labiata*. Its properties are those of a nauseous bitter; and hence it has been used in hysteria and other nervous affections.

LEONURUS LANATES, *Ballota lanata*.

LEOPARD'S BANE, ROMAN, *Arnica montana*, *Doronicum pardalanches*.

LEPER, see Leprous—*l. Hospital*, *Ladriere*.

LEPIA CAMPESTRIS, *Thlaspi campestre*—*l. Sativa*, *Lepidium Iberis*.

LEPID'IUM, (from *λεπς*, gen. *λεπιδος*, 'a scale,') so called from its supposed usefulness in cleansing the skin from scales and impurities.) *Pepperwort*. *Ord. Crucifere*.

LEPIDIUM CAMPESTRE, *Thlaspi campestre*.

LEPID'IUM IBE'RIIS, *Ibe'ria*, *Cardaman'tica*, *Scitrica cretensis*. This plant possesses a warm,

penetrating, pungent taste, like other cresses, and is recommended as an antiscorbutic, antiseptic, and stomachic.

LEPIDIUM SATIVUM, *Lep'ia sativa*, *Thlaspi nasturtium* seu *sativum*, *Nasturtium hortensū* seu *sativum*, *Gar'den cress*, *Ditt'ander*, *Green mustard*, (F.) *Cresson alénois*, *Cresson des Jardins*. This plant possesses warm, stimulating properties, and is used like the last.

LEPIDIUM SCUMATUM, *Cochlearia coronopus*.

LEPIDO, (from *λεπς*, gen. *λεπιδος*, 'a scale,') in composition, scale.

LEPIDODES, (*lepidō*, and *odes*), Squamous.

LEPIDOIDES, (*lepidō*, and *ειδός*, 'resemblance,') Squamous.

LEPIDOÏDE, Squamous.

LEPIDOSARCO'MA, gen. *Lepidosarco'matis*, *Tumor squamiformis carnosus*. A fleshy tumour, covered with scales; (*lepidō*, and *σάρκωμα*, 'a fleshy tumour,') Marcus Aurelius Severinus describes tumours of this kind in the interior of the mouth.

LEPIDOSIS, (*lepidō*, and *osis*), Scaly disease.

LEPIDOSIS ICHTHYIASIS, Ichthyosis—*l. Ichthyiasis cornigera*, Horny excrescences—*l. Lepriasis*, *Lepra*—*l. Pityriasis*, *Pityriasis*—*l. Psoriasis*, *Psoriasis*.

LEPIDOTIS CLAVATA, *Lycopodium clavatum*.

LEPIRA, *Lepira*.

LEPIS, (*λεπς*, gen. *λεπιδος*), Scale.

LEPISMA, gen. *Lepis'matis*, Squamous portion.

LEPORINUM LABIUM, (*lepus*, gen. *lep'oris*, 'a hare,') *Harelip*—*l. Rostrum*, *Harelip*.

LEPORINUS OCLUS, *Lagophthalmia*.

LEPOSTOPHYTON: (*λεπς*, *στέων*, 'a bone, and *φυτον*, 'a plant or growth,') A thin scale of bone, the result of a morbid growth.

LE'PRA, *Lep'ira*, (*λεπρος*, 'scaly,') *Lepido'sis Lepri'asis*, *Lepro'sis*, *Lepro'sitas*, *Vitil'go*, *Lep'ra Græcorum*, *Her'pes furfuraceus circina'tus*, *Lep'rosy*, *Lep'ery*, (Sc.) *Lepyr*, *Lipper*, (F.) *Lèpre*, *Mal Saint-Main*, *Dartre furfuracée arrondie*. This term has been applied to various affections, very different in character. 1. To the *Lep'rosy* of the Jews, *Leu'cē*, *Lepido'sis Lepri'asis canes'cers*, *Lep'ra Mosa'ica* seu *Hæbræorum*,—a variety of the *Al'phos* or *Lep'ra al'phoides*. The leuce was, generally, not scaly, but consisted of smooth, shining patches, on which the hair turned white and silky, and the skin, with the muscular flesh, lost its sensibility. It was incurable. 2. To the *Elephantiasis* or *Lep'ra of the A'rabs*, see *Elephantiasis*; and 3. To the *Lep'ra of the Greeks*, which includes all the varieties met with at the present day. It is characterized by scaly patches of different sizes, but having always nearly a circular form. Bateman and Willan describe three chief varieties of this lepra.

1. *Lep'ra al'phoides*, *Lepido'sis Lepri'asis al'bida*, *Al'phos*, *Morphe'a al'ba*, *Vitil'go al'phus*, *Al'baras al'ba*, *Al'bares*, *Al'baros*, *Lèpre écaillée* of Alibert, *White lep'rosy*. An affection, characterized by white patches, surrounded by a rose-coloured areola, which appears here and there on the surface; depressed in the middle.

2. *Lep'ra nigricans* seu *me'las* seu *maculo'sa nigra*, *Vitil'go Me'las* seu *nigra*, *Morphe'a* seu *Al'baras nigra*, *Me'las*, *Lepido'sis lepri'asis nigricans*, *Black lep'rosy*; in which the scales are livid; the size of half a dollar; and diffused over the body, but less widely than in the *Al'phoides*. The French pathologists usually admit three species of lepra, to which they give the epithets *scaly*, (*squamens*, *crustaceus*, (*crustacée*), and *tubercular*, (*tuberculeuse*,) according as the skin may be covered with scales, crusts, or tubercles.

3. *Lep'ra vulga'ris*, *Lepido'sis lepri'asis vulga'ris*, *Dartre furfuracée arrondie*, of Alibert, characterized as follows: scales glabrous, whitish

size of a crown piece; preceded by smaller reddish and glossy elevations of the skin, encircled by a dry, red, and slightly elevated border; often confluent; sometimes covering the whole of the body except the face.

Lepra appears to be endemic in Egypt, in Java, and certain parts of Norway and Sweden. Imperfect and faulty nutriment seems to contribute to its development. The means, best adapted for its removal, are:—a mild, unirritating diet, emollient fomentations—sulphureous baths, fumigations, &c.; but, often, all remedial agents will be found ineffectual.

LEPRA ARABUM, Elephantiasis Arabica—*l. As-turica*, see Pellagra—*l. Borealis*, Radzyge—*l. Fungifera*, Frambœsia—*l. Gangrænosa*, Ngerengere—*l. Græcorum*, Lepra—*l. Hebræorum*, see Lepra—*l. Ichthyosis*, Ichthyosis—*l. Lombardica*, Pellagra—*l. Maculosa nigra*, *l. Nigricans*—*l. Mediolanensis*, Pellagra—*l. Melas*, Lepra nigricans—*l. Mercurial*, Eczema mercuriale—*l. Mo-saica*, see Lepra—*l. Norvegica*, Radzyge—*l. Squa-mosa*, Impetigo—*l. Taurica*, *Mal de Crimée*.

LÉPRE, Lepra—*l. des Cosaques*, *Mal de Crimée*—*l. Écailleuse*, Lepra alphoides—*l. Hu-mide*, Impetigo.

LÉPREUX, Leprous.

LEPRICUS, Leprous.

LEPRODES, (λεπρός, from *lepra*, and *odes*.) Leprous.

LEPROSARIUM, (from *leprosus*.) Ladrerie.

LÉPROSERIE, Ladrerie.

LEPROSIS, Lepra.

LEPROSITAS, Lepra.

LEPROSUS, Leprous.

LEPROSY, Lepra—*l. Black*, Lepra nigricans—*l. Norwegian*, see Radzyge—*l. White*, Lepra alphoides.

LEPROUS, *Lepro'sus*, *Leprot'icus*, *Lep'ricus*, *Lepro'des*, (Sc.) *Lip'per*, (F.) *Lépreux*, *Ladre*. Relating to or resembling or affected with leprosy; a *lep'er*.

LEPSIS, (λεψις.) Attack.

LEPTAN'DRA VIRGIN'ICA, (*lepto*, and *ανθη*, 'an anther,') *Vero'nica* seu *Padero'ta Virgin'ica*, *Cul'ver's phys'ic*. Ord. Scrophulariaceæ. An indigenous plant, which grows throughout the United States, and flowers in August. The flowers are white, and terminate the stem in a long spike. A variety with purple flowers has been described, as

LEPTANDRA PURPU'REA, *Phys'ic* root, *Black* root, *Whorl'wort*, *Cul'ver's* root, *Brin'ton* root, *Bow-man* root. The root is bitter and nauseous, and when fresh is emetic and cathartic. In the dried state it is more uncertain. Dose of the powder, gr. xx to 3j.

LEPTO, (from λεπτός, in composition, thin, light.

LEPTOCEPHALIA, (*lepto*, and κεφαλή, 'head,') Monstrosity, in which the head is abnormally small.

LEPTOCHROA, *Lep'tochros*, (*lepto*, and χροα, *χρoα*, 'the colour of the skin: the skin,') Finesness, thinness of skin.

LEPTOCHYMIA, (*lepto*, and χυμος, 'a juice,') Morbid thinness of the juices.

LEPTOHYME'NIA, (*lepto*, and ὑμην, 'a membrane,') Thinness, delicacy of membrane.

LEPTOMER'IA, (*lepto*, and μέρος, 'a part,') Finesness, delicacy of bodily formation.

LEPTONTIQUES, (λεπυνω, 'to make thin,') Attenuants.

LEPTOPHONIA, (*lepto*, and φωνη, 'voice,') Oxyphonia.

LEPTOSPERMUM LEUCADENDRUM, (*lepto*, and σπερμα, 'a seed,') *Melaleuca cajaputi*.

LEPTOTHRIX, (*lepto*, and θριξ, gen. τριχος, 'hair,') Leptotrichus.

LEPTOTRICH'IA. Fineness of hair.

LEPTOTRICHUS, *Lep'tothrix*. One who has fine hair.

LEPTOTROPH'IA, *Microtroph'ia*, *Microtra-pez'ia*, (*lepto*, and τροφή, 'nourishment,') Light nutrition.

LEPTYNSIS, λεπυνσις, from λεπυνω, 'to make thin,') Emaciation.

LEPTYNTICA, Attenuants.

LEPTYSMUS, (λεπυσμος,) Emaciation.

LEPYR, (Sc.) Lepra.

LEREMA, gen. *Lere'matis*, (ληρημα,) see Dementia.

LERESIS, (ληρησις,) see Dementia.

LERUS, (ληρος,) Delirium, see Dementia.

LESEOLI MORBUS, Icterus.

LESE'OLUS. According to Paracelsus, a perfectly transparent salt, which cured jaundice; hence Lescoli morbus.

LE'SION, *Læ'sio*, (lædere, læsum, 'to injure,') Derangement, disorder; any morbid change, either in the exercise of functions or in the texture of organs. *Organ'ic* lesion is synonymous with *organic* disease.

LESION OF CONTINUITY, see Continuity, solution of.

LESK (Prov.), Inguen.

LESSIVE, Lixivium.

LESSIVE DES SAVONNIERS, ('lye of soapboilers,') Liquor potassæ. Also, a solution of caustic soda in water, containing about 3 parts of soda to 8 of water.

LESSIVE DE TARTRE, ('lye of tartar,') Liquor potassæ subcarbonatis.

LETALITAS, (letum, 'death,') see Mortal.

LETHALIS, (lethum, 'death,') Lethiferous, Mortal.

LETHALITAS, see Mortal.

LETHALITÉ, see Mortal.

LETHAR'GIC, *Lethar'gicus*, *Veterno'sus*, (F.) *Léthargique*, (ληθνη, 'oblivion, and ἀργος, 'inactive,') Relating to lethargy; affected with lethargy.

LÉTHARGIQUE, Lethargic.

LETH'ARGY, *Lethar'gia*, *Lethar'gus*, *Ca'rus* *lethargus*, *Veter'nus*, *Obliv'io ners*. A constant state of stupor from which it is almost impossible to arouse the individual; and, if aroused, he speedily relapses into his former condition.

LETHEA, (ληθη, 'oblivion,') Papaver.

LE'THEON, (ληθη, 'oblivion,') A name at first given to sulphuric ether, when inhaled as an anæsthetic agent.

LETHIF'EROUS, *Le'thifer*, *Letha lis*, *Lethif'icus*, *Mort'ifer*, *Mortifer'ous*, (F.) *Léthifère*, (ληθη, 'oblivion, and φερω, 'I bear,') Deathbearing; deadly.

LETHUM, Death.

LETTING BLOOD, Bloodletting.

LETTUCE, Lactuca—*l. Blue*, *Mulgedium acu-minatum*—*l. False*, *Mulgedium Floridanum*—*l. Indian*, see Calumba.

LETTUCE OPIUM, see Lactuca.

LETTUCE, STRONGSCENTED, Lactuca virosa—*l. White*, *Nabalus albus*—*l. Wild*, *Lactuca elongata*.

LETUM, Death.

LEUCADES, (λευκος, 'white,') see Sclerotic.

LEUCE'MIA, *Leukæ'mia*, *Leuchæ'mia*, (*leuco*, and αἷμα, 'blood,') (F.) *Leucémie*. A condition of the blood in which it is deficient in colouring matter. See Leucoerythæmia.

LEUCÆTHIOPIA, (*leuco*, and Αἰθωπ, 'an Ethiopian,') see Albino.

LEUCÆTHIOPS, Albino.

LEUCANIA, (λευκανια,) Laucania.

LEUCANTHEMUM, (*leuco*, and ανθερον, 'a flower,') *Anthemis nobilis*, *Matricaria chamomilla*—*l. Vulgare*, *Chrysanthemum leucanthemum*.

LEUCE, (λευκη, from λευκος, 'white,') Herpes exedens, Lepra (of the Jews.)

LEUCELECTRUM, (*leuco*, and λεκτρον, 'amber,') Succinum (album.)

LEUCÉMIE, Leucæmia.

LEUCÉTHIOPIE, see Albino.

LEUCHÆMIA, Leucæmia.

LEUCITIS, (*leuco*, and ιτις,) Sclerotitis.

LEUCO, (from λευκος, 'white,') in composition, white.

LEUCOCYTE, (*leuco*, and κυτος, 'a cell,') see Cytoid, and Globules of the blood.

LEUCOCYTHÆMIA, 'white cell blood,' (*leuco*, κυτος, 'cell, and 'αἷμα, 'blood,') A condition of the blood, which consists in a superabundant development of the white corpuscles.—*Polyleucocythæmia*—a disease which has been observed at times to be accompanied by enlargement of the spleen and liver, and at others by increased size of the lymphatic glands, organs which are concerned in hæmatisis. It is the *Leucocyto'sis* of Virchow. See Leucæmia.

LEUCOCYTOSIS, (*leuco*, and κυτος, 'cell,') Leucocythæmia.

LEUCOCYTOTIC, *Leucocyto'ticus*. Relating or appertaining to Leucocytosis.

LEUCODENDRON, (*leuco*, and δένδρον, 'tree,') Melaleuca cajuputi.

LEUCÆNUS, (*leuco*, and οἶνος, 'wine,') see Wine.

LEUCOÏUM, (*leuco*, and ἰον, 'the violet,') Lunaria rediviva—*L. Luteum*, Cheiranthus cheiri.

LEUCOL, Leukoleinum.

LEUCOLACHANUM, (*leuco*, and λαχανον, 'herb,') Valeriana.

LEUCOLEIN, Leukoleinum.

LEUCOMA, gen. *Leuco'matis*, (from λευκος, 'white,') Oculo'rum albu'go, *Leuco'm'ma*, *Leuco'sis*, *Albu'go*, *Al'bula*, *Ceratoleuco'ma*, *Ephel'otes*. *Leucoma* and *Albugo* are often used synonymously to denote a white opacity of the cornea. Both are essentially different from nebula of the cornea; nebula being usually the result of chronic ophthalmia and an effusion of a milky serum into the texture of the delicate continuation of the conjunctiva over the cornea;—the others are the result of violent, acute ophthalmia. In this state, a thick, coagulable lymph is extravasated from the arteries, sometimes superficially, at other times deeply into the substance of the cornea. On other occasions, the disease consists of a firm, callous cicatrix on this membrane,—the effect of a wound or ulcer with loss of substance. The affection is more difficult of cure in proportion to its duration and to the age of the individual; the activity of the absorbents being greater in youth. If inflammation still exist, antiphlogistics must be persevered in, and, afterwards, gentle stimulants be used to excite absorption; along with the internal use of mercury or iodine.

LEUCOMA, Albumen—*L. Gerontotoxon*, Gerontoxon—*L. Margaritaceum*, see Margaritaceus.

LEUCOMMA, gen. *Leuco'm'matis*, Leucoma.

LEUCOMORIA, (λευκος, 'feeble,' and μωρια, 'fatuity,') Cuttubuth.

LEUCONECRO'SIS, *leuco*, and νεκρωσις, 'death,') A form of dry gangrene,—the opposite in appearance to anthraconecrosis.

LEUCONYMPHÆA, Nymphæa alba.

LEUCOPATHIA, (*leuco*, and παθος, 'affection,') see Albino, Chlorosis—*L. Partialis* acquista, Achroma.

LEUCOPHAGIUM, (*leuco*, and φάγω, 'to eat,') *Blauemanger*.

LEUCOPHLEGMASTIA, Leucophlegmatia—*L. Dolens* puerperarum, Phlegmatia dolens.

LEUCOPHLEGMA'TIA, *Leucophlegma'tia*, *Hy'drops leucophlegma'tia*, *Tumescen'tia pituitosa*, (*leuco*, and φλεγμα, gen. φλεγματος, 'phlegm,') A dropsical habit. *Hydrocachex'ia*. Some writers

use the word synonymously with *anasarca* and *œdema*; others with *emphysema*.

LEUCOPHLEGMASTIA ÆTHIOPUM, Chthonophagia.

LEUCOPIPER, (*leuco*, and πιπρι, 'pepper,') Piper album.

LEUCOPYRIA, (*leuco*, and πυρ, 'fire, fever,') Hectic fever.

LEUCORRHÉE, Leucorrhœa.

LEUCORRHÆIC, *Leucorrhœ'ic*, *Leucorrhœ'icus*, (F.) *Leucorrhéique*, (same etymon as *Leucorrhœa*.) Relating or appertaining to leucorrhœa.

LEUCORRHÆA, (*leuco*, and ρέω, 'I flow,') *Flux'us* seu *Flu'or* multi'ebri's, *Proflu'vium* multi'ebri's, *Cur'us* matric'is, *Flux'us* matric'is, *Elytrit'is*, *Coleosit'is*, *Colpit'is*, *Destilla'tio* uteri, *Flux'ic* al'ba seu vul'væ, *Ul'us* uteri, *Catar'rhus* genita'lium, *Hysterorrhœ'a* mucosa, *Catame'nia* al'ba, *Men'ses* al'bi, *Men'strua* al'ba, *Menorrhag'ia* al'ba, *Flu'or* multi'ebri's non Gall'icus, *Blennely'tria*, *Gonorrhœ'a* benigna no'tha invetera'ta, *Purgatio* multi'ebri's al'ba, *Al'ba* purgamenta, *Cachex'ia* uteri'na, *Rheu'ma* uteri, *Uteri* Cory'za, *Menorrhœ'a* femina'rum in'sons, *Menorrhœ'a* seu *Blennorrhag'ia* genita'lium, *Ædœoblennorrhœ'a* seu *Medoblennorrhœ'a* femina'rum; *The whites*; (F.) *Flours* ou *Flueurs* blanches, *Pertes* blanches, *Écoulement blanc*, *Catarrhe* utérin, *Perte* utérine blanche. A more or less abundant discharge of a white, yellowish, (yellows) or greenish mucus; resulting from acute or chronic inflammation, or from irritation of the membrane lining the genital organs of the female. Vag'inal Leucorrhœa has been termed *Blennorrhœ'a* seu *Flu'or* al'bus vag'inæ, *Leucorrhœ'a*, *Medorrhœ'a* vaginæ, *Vaginit'is*, *Elytrobrennorrhœ'a*, *Colporrhœ'a*.—Uterine Leucorrhœa has received the names *Flu'or* al'bus uteri, *Leucorrhœ'a* seu *Medorrhœ'a* uteri, *Metrobrennorrhœ'a*, *Blennometri'tis*, *Blennorrhœ'a*, *Metrobrennozen'ia*, *Metrocata'rhus*, *Metrolencorrhœ'a*, (F.) *Leucorrhée* utérine, *Catarrhe* utérin. It is often attended with pain and a sense of heaviness in the loins, abdomen and thighs; disordered digestive functions, &c., so that, at times, the health suffers largely, although there are few females who are not occasionally subject to moderate leucorrhœa. Attention to the general health, change of air, keeping up a perspirable state of the surface by flannel worn next the skin, the horizontal posture, &c., do more than pharmaceutical agents; which are almost entirely confined to astringent injections. These may be employed, when the discharge is so great as to require them.

LEUCORRHÆA ANALIS, see Hæmorrhœis—*L. Nabothi*, see Parturition.

LEUCORRHOIS, (*leuco*, and ρέω, 'to flow,') See Hæmorrhœis.

LEUCO'SES, (from λευκος, 'white,' and σις.) In the nosology of M. Alibert, all the diseases of the lymphatic apparatus. The 7th family in his *Nosologie*. In that of Fuchs it is a family of diseases, (G.) Leukosen, which includes the various forms of anæmia.

LEUCOSIS, (λευκωσις,) Leucoma, Paleness. See Albino.

LEUK or LOCHE, MIN'ERAL WA'TERS OF. Saline chalybeate waters, about six leagues distant from Sion. They contain chloride of sodium, with a little sulphate of magnesia; sulphate of lime, carbonate of magnesia, carbonic acid, and protoxide of iron.

LEUKÆMIA, Leucæmia.

LEUKOLEIN'UM, *Chinolein'um*, *Leu'coleine*, *Leu'colein*, *Chi'nolein*, *Chi'nolin*, *Qui'nolin*, (from λευκο, and ελαιον, 'oil,') (?) This substance is the product of the dry distillation of coal; mixed with picolin, anilin and other substances, in mineral tar. It is procured, also, by heating quinia, cinchonia, and strychnia, with as concen-

trated a ley of *potassa* as can be made. Its specific gravity is 1.081; and it is slightly soluble in water, and miscible in all proportions with alcohol, ether, and essential oils.

LEVAIN. (*levere*, 'to raise.') Ferment.

LEVANDIÈRE, Midwife.

LEVATOR, (*levere*, 'to raise.') Elevator,

LEVATOR ANGULI O'RIS, *Abducens labiorum*, *Eleva'tor labiorum commu'nis*, *Caninus*. (F.) *Submaxillo-labialis*, *Petit sus-maxillo-labialis* (Ch.), *Muscle canin*. A small, flat, long, quadrilateral muscle, which arises from the fossa canina, and is inserted at the commissure of the lips, where it is confounded with the triangularis. It raises the corner of the mouth, and draws it towards the ear.

LEVATOR A'NI, *Levator magnus seu inter'nus*, *Lat'us a'ni*, *Eleva'tor a'ni*, *Se'dem attol'lens*, *Diaphragma pel'vis*, (F.) *Pubio-coccygien annulaire*, *Sous-pubio-coccygien* (Ch.), *Releveur de l'anus*. A muscle situate at the lower part of the pelvis. It is broad, flat, quadrilateral, and broader above than below. It represents a kind of membranous partition, which closes the outlet of the pelvis, and the upper concavity of which is opposed to that of the diaphragm. It is attached, above, to the posterior surface of the body of the pubis, to the upper part of the obturator foramen, and to the spine of the ischium, and is inserted into the coccyx, into an aponeurotic line common to it and its fellow, and into the lateral parts of the rectum. This muscle supports the rectum; raises it, and carries it upwards during the evacuation of the excrement. It can, also, compress the bladder and vesiculae seminales, and thus favour the expulsion of the urine and sperm.

LEVATOR ANI PARVUS, *Transversus perinaei*—*l. Auris*, *Attollens aurem*—*l. Coccygis*, *Coccygeus*.

LEVATOR GLAN'DULÆ THYROIDÆÆ. A muscle occasionally found connected with the upper border or isthmus of the thyroid gland; and attached superiorly to the body of the os hyoides, or to the thyroid cartilage.

LEVATOR LA'BII INFERIO'RIS, *L. menti*, *Eleva'tor labii inferior'is seu labii inferior'is proprius*, *Incisivus inferior*, *Musculus penicillatus*, (F.) *Houpe du menton*, *Releveur de la lèvre inférieure ou du menton*. A portion of the *mento-labialis* of Chaussier. A small muscle situate before the symphysis menti. It is thick, conical, and attached by its apex to a fossette at the side of the symphysis in the inferior maxillary bone. Its fibres proceed, diverging and vanishing in the manner of a tuft, (F.) *Houpe*, on the skin of the chin. This muscle raises the chin, and pushes upwards the lower lip.

LEVATOR LA'BII SUPERIO'RIS A'LEQUE NA'SI, *Incisivus lateralis et pyramidalis*, (F.) *Grand sus-maxillo-labialis* (Ch.), *Élévateur commun de l'aile du nez et de la lèvre supérieure*. This muscle is a fleshy, thin, triangular bundle, situate at the sides of the nose. It arises from the ascending process of the superior maxillary bone; thence its fibres descend in a diverging manner, a part being inserted into the ala of the nose, and a part losing themselves in the upper lip. This muscle raises the upper lip and ala nasi, which it draws a little outwards.

LEVATOR LA'BII SUPERIO'RIS PRO'PRIUS, *Musculus incisivus*, *Eleva'tor labii superior'is proprius*, (F.) *Moyen sus-maxillo-labialis*, (Ch.), *Orbitomaxillo-labialis*. This thin, flat, quadrilateral muscle is situate at the middle and inner part of the face. It arises from the os male and the os maxillare superius, and is inserted into the upper lip, which it raises, at the same time carrying it a little outwards.

LEVATOR MENTI, *Levator labii inferioris*.

LEVATOR OCULI, *Rectus superior oculi*.

LEVATOR PALA'TI, *L. Pala'ti Mol'lis*, *Petrosalpin'gostaphyli'nus*, *Salpin'gostaphyli'nus inter'nus*, *Salpin'gostaphyli'nus*, *Pter'ygostaphyli'nus externus*, *Sphe'nostaphyli'nus*, *Sphe'nopalati'nus*, *Peristaphyli'nus inter'nus superior*, *Péto-staphylin*, (Ch.). This muscle is long, narrow, and almost round above; broader and flatter inferiorly. It arises from the lower surface of the *pars petrosa*; and from the cartilage of the Eustachian tube; and is inserted into the substance of the velum palati. Its use is to raise the velum palati.

LEVATOR PAL'PEBRÆ SUPERIO'RIS, *Pal'pebræ superior'is primus*, *Aper'tor Oc'uli*, *Ape'riens Palpebra'rum rectus*, *Recl'u'sor palpebra'rum*, (F.) *Orbito-palpébral* (Ch.), *Orbito-sus-palpébral*, *Élévateur de la paupière supérieur*. A long, small, thin muscle, situate at the upper part of the orbital cavity. By its posterior extremity it is inserted into the little ala of the sphenoid bone, immediately in front of the foramen opticum, and by its anterior extremity, which is expanded, it terminates at the upper margin of the tarsal cartilage of the eyelid. This muscle raises the upper eyelid, draws it backwards, and sinks it into the orbit.

LEVATOR PROS'TATÆ, (F.) *Releveur de la prostate*. Santorini has given this name to the anterior fibres of the levator ani, which embrace the prostate.

LEVATOR SCAP'ULÆ, *Eleva'tor scap'ulæ*, *E. seu Musculus Patient'is*, *Angula'ris*, *Levator proprius scap'ulæ*, (F.) *Trachélo-scapulaire*, (Ch.), *Angulaire de l'omoplate*, *Releveur de l'omoplate*. This muscle is so called because it is attached to the upper and internal angle of the scapula. It is situate at the posterior and lateral part of the neck, and at the upper part of the back. It is long, flat, and broader above than below. In this latter direction, it is inserted into the superior internal angle of the scapula; and in the former, into the tops of the transverse processes of the first four cervical vertebrae. It depresses the prominence of the shoulder by raising the posterior angle of the scapula, on which it impresses a kind of rotary motion. It can, also, draw the head to one side.

LEVATOR PROPRIUS SCAPULÆ, *Levator scapulæ*.

LEVATOR URE'THRE, (F.) *Releveur de l'Urèthre*. Santorini describes under this name a portion of the transversus perinaei muscle.

LEVATORES COSTARUM, *Supracostales*.

LEVEN, (from *levere*, 'to raise.') Ferment.

LE'VER, (from *levere*, 'to lift up.') *Vect'is*, *Moch'lus*, *Porre'ctum*. An inflexible rod, turning round a fixed point, and used for moving bodies, bearing burdens, or raising them. The point on which the lever moves is called the *Fulcrum Hypomoch'lion*. The force which moves the lever is called the *power*; and the weight to be moved the *resistance*. There are three kinds of levers. A lever of the first kind has the fulcrum between the power and resistance. A lever of the second kind has the resistance between the fulcrum and power; whilst a lever of the third kind has the power between the fulcrum and resistance. In the locomotive system of the human body, we have examples of all the three kinds. The bones represent levers; the muscles of locomotion are powers; the weight of parts to be moved constitutes the resistance. The fulera are, at times, the joints; at others, the ground, &c. The head moves on the neck as a lever of the first kind; the first cervical vertebra forming the fulcrum. We rise on tiptoe by a lever of the second kind, the fulcrum being the ground under the toes; and we have examples of a lever of the third kind

in the flexion of the forearm on the arm, in the elevation of the arm, &c.

LEVIER, *Hystero-moch'lum*, *Mochlis'eus*, *Vec'tis obstetric'i'us*, *Vec'tis*, (F.) *Lévier*, is an instrument curved at the extremity, and having a fenestra. It is used to assist the extraction of the child's head when instrumental aid is necessary, either as a lever proper or as a tractor. Levers are also used by the dentist for extracting stumps, &c. The *Lévier de l'Écluse*, *Langue de Carpe*, *Trevelin*, *Pied de biche*, *Punch*, is employed for extracting the molar teeth.

LEVY'ATHAN PE'NIS, *Pria'pus Cel'ti*, *Bale'nas*. The penis of the whale. This singular medicine was at one time given in powder in cases of dysentery and leucorrhœa.

LEVIER, Lever—*l. de l'Écluse*, see Lever,

LEVIGAT'ION, *Læviga'tio*, (*lævigare*, *lævigatum* (*lævis*, 'smooth,') 'to bruise small,') *Porphy-riza'tion*. An operation by which bodies are reduced to very fine powder. It is performed by putting substances already pulverized into water; the coarser parts are not long in being deposited, whilst the finer molecules remain suspended in the water. The liquor is decanted into another vessel, and suffered to remain at rest until the fine particles are collected at the bottom. The fluid part is then separated by decantation.

LEVISTICUM, *Ligusticum levisticum*.

LÈVRE, (from *labrum*,) Lip.

LÈVRES GRANDES, *Labia pudendi*—*l. Petites*, *Nymphæ*—*l. de la Vulve*, *Labia pudendi*.

LEVÛRE, (from *levare*, 'to raise,') Yest—*l. de la Bière*, Yest.

LEXIPHARMACUS, (*λεγειν*, 'to make cease,' and *φαρμακον*, 'a poison,') *Alexipharmic*.

LEXIPYRETICUS, (*λεγειν*, 'to make cease,' and *πυρ*, 'fire, fever,') *Febrifuge*.

LEY, *Lixivium*, Lye—*l. Soap*, *Liquor potasse*.

LEYSERA GNAPHAL'IDES, (after Fr. W. von Leyser, of Halle.) A South African plant, *Nat. Ord.* Compositæ, which is emollient, and highly recommended at the Cape in catarrh, cough, and even in phthisis.

LÉZARD, Lizard.

LIANE À RÉGLISSE, *Abrus precatorius*.

LIA'TRIS SPICA'TA, *Gay's feather*, *Button snakeroot*. *Ord.* Compositæ. An indigenous plant, growing in meadows and moist grounds in the Middle and Southern States. Its beautiful purple compound flowers are in a spike. They appear in August. The root has been considered diuretic.

LIATRIS SCARIO'SA and **L. SQUARRO'SA**, *Throatwort*, *Sow'wort*, *Back'ache root*, *Dev'il'sbite*, *Blazing Star*, *Prairie pines*, *Rough root*, are called from their reputed powers in bites from the rattlesnake, *Rattle'snake's master*. The roots, bruised, are applied to the wound; and the decoction, in milk, is given internally.

LIBANOTIS ANNUA, (from *λίβανος*, 'the frankincense tree,') *Athamanta cretensis*—*l. Coronaria*, *Rosmarinus*—*l. Cretensis*, *Athamanta cretensis*—*l. Hirsuta*, *Athamanta cretensis*.

LIBANOTUS, see *Juniperus lycia*.

LIBANUS, (*λίβανος*,) *Juniperus lycia*.

LIBER PHARMACEUTICUS, ('pharmaceutical book,') *Dispensatory*.

LIBIDINIS SEDES, ('seat of lust,') *Clitoris*.

LIBIDINOSUS, (*libido*, gen. *libid'inis*, 'lust,') *Libidinous*.

LIBID'INOUS, *Libidino'sus*, *Lasci'vus*, *Veneriv'agus*, *Lech'erous*, *Lick'orous*. Lewd; lustful.

LIB'IDO, gen. *Libid'inis*. Desire, necessity. Authors speak of *Libi'do uri'næ*, *Libi'do intesti'ni*. Some employ it synonymously with *Pruri'go*; others, with *Salac'i'tas*, *Lubi'do*, *H'i'meros*.

LIBOS, (*λίβος*, from *λειβω*, 'to drop,') A delusion from the eyes.

LIBRA, (*λίτρα*,) Pound.

LICENT'IATE, (*licet*, 'it is permitted,') *Permis'sus*, (F.) *Licencié*. One licensed to practise his profession.

LICHANOS, (*λίχανος*,) Index, see *Digitus*.

LICHE, Body.

LICHEN, gen. *Liche'nis*, (*λεχην* or *λεχην*, pronounced *L'i'ken*.) *Exor'mia Li'chen*, *Le'i'chen*, *Serpi'go*, *Volat'ica*, *Pap'ula*, *P. sic'ca*, *Peti'go*, *Pustula sic'ca*, *Scu'bica sic'ca seu a'gria*, *Lichen'asis adulto'rum*, *Li'chenous rash*, (F.) *Gale sèche*, *Dartre furfuracée volante*, *Poussée*. The cutaneous affection described under this name by the Greek writers is not clearly defined. Some have believed it to be *Impetigo*, but this is doubtful. The name is now generally applied to a diffuse eruption of red pimples, accompanied by a troublesome sense of tingling or pricking. Drs. Willan and Bateman define it—"an extensive eruption of papule affecting adults, connected with internal disorder, usually terminating in scurf; recurrent, not contagious." One of their varieties, however, the *Lichen Tropicus*, does not accord well with this definition; for it affects children as well as adults, and is unconnected with internal disorder.

LICHEN A'GRIUS, *L. A'gria*, *Exor'mia Li'chen fe'rus*, *Li'chen fe'rus*, *Pap'ula a'gria*, *Wild Li'chen*, is distinguished by pimples in clusters or patches, surrounded by a red halo; the cuticle growing gradually harsh, thickened, and chappy, often preceded by general irritation. In addition to antiphlogistics, a cooling ointment may be used, to allay itching.

LICHEN BARBATUS PLICATUS, *L. plicatus*.

LICHEN CIRCUMSCRIP'TUS is characterized by clusters or patches of papule; having a well defined margin, and an irregularly circular form; continuing for six or eight weeks. These varieties require but little medical treatment. The antiphlogistic plan is all that is necessary.

LICHEN FERUS, *Lichen agrius*.

LICHEN LIV'IDUS. The papule have a dark red or livid hue, without any fever. They are more permanent in this variety. It requires the mineral acids and bark.

LICHEN PILA'RIS, *Exor'mia Li'chen pila'ris* is merely a modification of the preceding; the papule appearing at the roots of the hair.

LICHEN SIM'PLEX, *Exor'mia Li'chen sim'plex*, consists of an eruption of red papule, appearing first on the face or arms, and afterwards extending over the body; preceded for a few days by slight febrile irritation, which usually ceases when the eruption appears,—with an unpleasant sense of tingling during the night. It generally dies away in ten days or a fortnight.

LICHEN TROP'ICUS, *Exormia Li'chen trop'icus*, *Eczema*, *Es'sera* [?], *Sudam'ina* [?], *Prickly Heat*, *Summer Rash*. The pimples are bright red, and of the size of a small pin's head; with heat, itching, and pricking, as if by needles. It is local; produced by excessive heat; and disappears when the weather becomes cooler, or the individual is injured to the climate.

LICHEN URTICA'TUS, *L. Urtico'sus*, *Exor'mia Li'chen urtico'sus*, is another variety. The *Nettle Li'chen* consists of papule, accompanied by wheals like those of nettle rash. A variety of this is said to be called, in America, *blights*.

LICHEN, *WILD*, *Lichen agrius*.

LICHEN APHTHO'SUS, *Pelti'dea aphtho'sa*; *Ord.* *Lichenaceæ*; is said to be cathartic and anthelmintic.

LICHEN ARBORUM, *Lichen pulmonarius*—*l. Barbatus plicatus*, *Lichen plicatus*—*l. Blanc de Neige*, *L. caninus*—*l. Bôttier*, *L. pyxidatus*.

LICHEN CANI'NUS seu *epu'rius seu terres'tris seu veno'sus seu ciner'eus terres'tris*, *Mus'eus cani'nus*, *Phys'cia nive'lis*, *Peltig'era cani'na*, *Pelti-*

aea canina seu *leucorrhiza* seu *mala'cea* seu *amplissima* seu *spuria*, Ashcoloured Ground Liverwort, (F.) *Lichen contre-rage*, *Lichen blanc de neige*. This cryptogamous plant was, for a long time, considered capable of preventing and curing *Rabies canina*. It has, also, been used in mania and in spasmodic asthma.

LICHEN CARRAGEEN, *Fucus crispus*—1. *Cinereus terrestris*, *L. caninus*—1. *Cocciferus*, *L. pyxidatus*—1. *Entomoir*, *L. pyxidatus*—1. *Eryngifolius*, *L. islandicus*—1. *Floridus hirtus*, *L. plicatus*—1. *Hirtus*, *L. plicatus*.

LICHEN ISLANDICUS, *L. eryngifolius*, *Lichen des Islandicum*, *Lobelia Islandica*, *Muscus Islandicus* seu *catharticus*, *Cladonia* seu *Physcia* seu *Cetraria* seu *Parmelia* *islandica*, *Iceland Lichen* or *Liverwort* or *Moss*, (F.) *Lichen d'Islande*, *Mousse marine perlée*, *Mousse d'Islande*. This plant is inodorous, with a bitter and mucilaginous taste. It is esteemed to be tonic, demulcent, and nutrient. Dose, $\mathfrak{z}$ i to $\mathfrak{z}$ iv, being first steeped in water holding in solution some carbonate of potassa to extract the bitter, and then boiled in milk. A bitter principle has been extracted from it, termed *Cetrarin*, *Cetrarinum*, *Cetrarium*, which has been given in intermittents.

LICHEN LACINIATUS, *L. saxatilis*—1. *Parietinus*, *Parmelia parietina*.

LICHEN Plicatus, *L. hirtus* seu *barbatus plicatus* seu *Floridus hirtus*, *Parmelia plicata*, *Usnea plicata*, *U. hirta* seu *Florida hirta*, *Muscus arborescens* seu *albus* seu *quercinus*, *Quercini Herba*. This plant is applied, by the Laplanders, as an astringent, to bleeding vessels; and to parts which are excoriated after long journeys.

LICHEN PULMONARIUS, *L. arborum* seu *reticulatus*, *Parmelia pulmonacea*, *Reticularia officinalis*, *Muscus pulmonarius quercinus*, *Pulmonaria arborescens*, *Lobelia pulmonaria*, *Sticta pulmonacea*, *Oak Lungs*, *Tree Lungwort*, *Ha'zel Cro'bles*, *Ha'zelraac*, (F.) *Lichen pulmonaire*, *Pulmonaire en arbre ou de chêne*. This plant is subastringent, and rather acid. It was once in high repute for curing diseases of the lungs.

LICHEN PYXIDATUS, *Muscus Pyxidatus*, *Seyphophorus pyxidatus*, *Muscus pyxoides terrestris*, *Lichen pyxidatus major* seu *cocciferus*, *Herba l'ymis*, *Cup Moss*, (F.) *Lichen entomoir*, *Lichen Bôtier*, *L. Pyxidé*. This plant is subastringent, and has been used in decoction in hooping-cough.

LICHEN CONTRE RAGE, *L. caninus*.

LICHEN RETICULATUS, *L. pulmonarius*.

LICHEN ROCCEL'LA, *Parmelia roccella*, *Roccella*, *R. tinctoria*, *Scleria roccella*, *Litmus*, *Lacmus*, *L. tinctorius*, *Orchill*, *Canary Archell*, *Chinny Weed*, *Herb Archell*, (F.) *Orseille*. The chief use of this plant is as a blue dye. It has been employed to allay cough in phthisis, &c.

The blue pigment—*Litmus*—is much used as a delicate test for acid—in the urine for example. *Blue Litmus paper* is unsized paper steeped in tincture of litmus—which is made of *Litmus powder* $\mathfrak{z}$ i, *proof spirit*, $\mathfrak{f}$ 3x, and dried. It is reddened by acids. *Red litmus paper* is unsized paper steeped in tincture of litmus, which has been previously reddened by the addition of a very small quantity of sulphuric acid, and dried. The blue colour is restored by alkalis.

LICHEN ROCK, *Tripe de roche*.

LICHEN SAXATILIS, *L. tinctorius* seu *lacinia'tus*, *Parmelia saxatilis*, *Imbricaria saxatilis*, *Lobelia saxatilis*, *Muscus* seu *Usnea* *Cra'nii humani*. This moss, when found growing on a human skull, was formerly in high estimation against head affections, &c.

LICHEN SPURIUS, *L. caninus*—1. *Stellatus*, *Marchantia polymorpha*—1. *Tinctorius*, *L. Saxatilis*—1. *Wall*, yellow, *Parmelia parietina*.

LICHENIASIS ADULTORUM, *Lichen*—1. *Strophulus*, *Strophulus*.

LICHENOID, *Lichenoides*, (*lichen*, and *eidos*, 'resemblance,') Resembling lichen.

LICHENOIDES, *Lichenoid*—1. *Islandicum Lichen Islandicus*.

LICINIUM, (from *linum*, 'thread' [?]), Tent.

LICKOROUS, *Licibidinus*.

LICORICE, *Glycyrrhiza*.

LICULMEN, *Garum*.

LIEBENSTEIN, MINERAL WATERS OF These waters, situated in the duchy of Saxe-Meiningen, are amongst the strongest acidulous chalybeates in Germany. They contain sulphate of soda, chloride of sodium, sulphate of lime, chloride of calcium, carbonate of lime, chloride of magnesium, and carbonate of iron.

LIEBERKÜHN'S GLANDS or FOLLICLES, *Glandulae seu Cryptae Lieberkühniæ*, so called from their first describer, are fine, columnar, blind sacs, the openings of which are from 1-20th to 1-30th of a line in diameter, so closely placed over the whole of the small intestine as to give the mucous membrane a general sieve-like or perforated appearance. They secrete the *succus entericus*.

LIEBERSCHES KRAUTER, ('Lieber's consumption herbs,') see *Galeopsis grandiflora*—1. *Brustkraut*, see *Galeopsis grandiflora*.

LIEBWERDA, MINERAL WATERS OF. Liebwerda is a Bohemian village, near the Silesian frontier. The springs are much frequented. There are four, all of which are rich in carbonic acid, and contain but little saline matter.

LIÈGE, (from *leger*, 'light,') Suber.

LIEN (F.) (from *lier*, [L.] *ligure*, 'to bind,') *Vinculum*. A band, strap or garter, used in certain operations; as to tie patients during the operation of lithotomy; to fix the apparatus in fracture, &c.

LIEN, Spleen—1. *Accessorius*, *Lienculus*—1. *Ingens*, *Splenoneus*—1. *Succenturiatus*, *Lienculus*.

LIEN'CULUS, (dim. of *lien*, 'spleen,') *Lien succenturiatus* seu *accessorius*. A supernumerary spleen.

LIENITIS, (*lien*, and *itis*,) *Splenitis*.

LIENOSUS, *Splenice*.

LIENTERIC, *Lientericus*, (F.) *Lientérique*. Relating or appertaining to lientery.

LIENTÉRIE, *Lientery*.

LIENTERY, *Lienteria*, (*λαος*, *laeis*, 'smooth,' 'slippery,' and *εντερον*, 'intestine,') *Læritus* seu *Læritus intestinum*, *Diarrhœa lienteria*, *Fluxus Lientericus*, *Chymorrhœa* seu *Læritus ingestorum*, *Bromatœcrosis*, *Diarrhœa Dyspeptica* seu *cum Apepsiâ* seu *Ingestorum*, (F.) *Lientérie*, *Slip'periness of the Guts*. Frequent liquid evacuations, the food only half digested. This condition is always symptomatic of great irritation in the intestinal canal, the sensibility of which is so much augmented that it cannot bear the sojourn of the food in it.

LIERRE, *Hedera helix*—1. *Grimpant*, *Hedera helix*—1. *Terrestre*, *Glechoma hederacea*.

LIFE, (Sax. *lyr*, *lyr*,) *Vita*, *Bîas*, *Bîoté*, *Bîotos*, *Zoë*, *Pneum'a*, *Spir'itus*, (F.) *Vie*. The state of organized beings, during which, owing to the union of an unknown force with matter, they are capable of performing functions different from those that regulate other natural bodies; all of which functions, however numerous and diversified, work to one end. Life has only a limited duration; beyond which,—the organic functions ceasing to be executed,—the body is given up to the agency of chemical affinity. Hence Bichat has defined life to be—the aggregate of the functions which resist death. On account of the difference that exists among the vital functions, be

applied the term *Organic Life* to the functions, inservient to composition and decomposition;—as digestion, respiration, circulation, calorification, absorption, secretion, and nutrition; and *Animal life*, to the functions which connect man and animals with external bodies; as the understanding, sensations, locomotion and voice.

LIFEBLOOD. A term commonly used for a twitching or quivering of the eyelids—a kind of involuntary or muscular tic. See Cillo.

LIFE, DURATION OF, see Longevity.

LIFE EVERLASTING, *Gnaphalium margaritaceum*—l. Everlasting, plaitain, *Antennaria plantaginifolia*—l. Everlasting, sweetscented, *Gnaphalium polyccephalum*.

LIFE, EXPECTATION OF, see Longevity.

LIFE INSURANCE. A contract entered into, usually by an *insurance company*, to pay a certain sum of money on a person's death, on the condition of his paying an annual premium during his life. The medical practitioner may be applied to, to certify that the life of the insured is one that is insurable according to the rules of the company.

LIFE ROOT, *Senecio aureus*.

LIFE, VALUE OF, see Longevity, and Mortality.

LIFE'S BLOOD, Lifeblood, Cillo.

LIGAMENT, *Ligamentum*, (*ligare, ligatum*, 'to bind;') *Desmus, Syndes'mus, Colliga'men, Cop'ula*. A name given to fibrous structures, which serve to unite bones, and to form articulations; hence the division into *interosseous* and *articular* ligaments. They are of a white, close texture; are but little extensible, and difficult to break. The name *ligament* has, also, been given to any membranous fold, which retains an organ in its situation.

TABLE OF THE PRINCIPAL LIGAMENTS.

1. <i>Of the Lower Jaw.</i>	- Capsular ligament. - Suspensory ligament of the stylo-glossus. - Lateral ligament.		- Two transverse,—one superior and one inferior. - Iliosacral. - Capsular of the sacroiliac synchondrosis.
2. <i>Connecting the Head with the first and second Vertebrae, and these with each other.</i>	- Two capsular ligaments between atlas and head. - Circular ligament. - Two capsular between atlas and axis. - Perpendicular ligament. - Two lateral or moderator ligaments. - Transverse ligament and its appendices.	5. <i>Of the Pelvis.....</i>	- Two sacrospinatic. — posterior and anterior, with the superior and inferior appendices. - Longitudinal of os coccygis. - Inguinal ligament. - Capsular of symphysis pubis. - Ligament of foramen thyroideum.
3. <i>Of the other Vertebrae.</i>	- Anterior common. - Crucial intervertebral. - Ligaments running from the edge of the bony arch and spinous process of one vertebra to that of the next. - Interspinous ligament. - Ligamentum nuchæ. - Interspinous. - Capsular. - Posterior or internal common.	6. <i>Of the Clavicle.....</i>	- Radiated ligament. - Capsular. - Interclavicular. - Ligamentum rhomboideum. - Claviculoacromial. - Conoid. - Trapezoid.
4. <i>Of the Ribs, Sternum, &c.</i>	- Capsular of the head of ribs. - Capsular of the tubercles. - Ligamenta transversaria interna. - Ligamenta transversaria externa. - Ligamenta cervicis costarum externa. - Ligamentous fibres running from the margins of the extremities of the ribs to the corresponding cartilages. - Radiated ligaments from cartilage of ribs to the sternum. - Capsular ligaments of the cartilages of the ribs. - Proper membrane of sternum. L. of cartilago ensiformis. - Tendinous expansions over the intercostales, &c.	7. <i>Of the Scapula.....</i> 8. <i>Of the Shoulder Joint.</i> 9. <i>Of the Elbowjoint..</i>	- Anterior triangular. - Proper posterior. - Capsular. - Capsular. - Brachioulnar. - Brachioradial. - Coronary of the radius. - Anterior and posterior accessory. - Intermuscular of the os humeri.
		10. <i>Carpal Extremity of Radius and Ulna, and between those bones.</i> 11. <i>Between Fore-arm and Wrist.</i> 12. <i>Of the Carpus.....</i> 13. <i>Between Carpal and Metacarpal bones.</i> 14. <i>Between the extremities of the Metacarpal bones.</i> 15. <i>At the base of the Metacarpal bone of the Thumb, and at the first joint of the Fingers.</i> 16. <i>Of the first and second joints of the Thumb, and second and third joints of the Fingers.</i> 17. <i>Retaining the Tendons of the Muscles of the Hand and Fingers in situ.</i> 18. <i>Connecting the Os femoris with the Os innominatum.</i> 19. <i>Of the Kneejoint.</i> 20. <i>Connecting Fibula and Tibia.</i> 21. <i>Connecting the Tarsal with the Leg Bones.</i>	- Interosseous ligament. - Oblique or chorda transversalis cubiti. - Capsular. - Capsular. - Two lateral. - Mucous. - Annular. - Capsular. - Short ligaments, — oblique, transverse, capsular, and proper. - Articular lateral. - Straight, perpendicular, &c. - Interosseous, at the bases and heads, — dorsal, lateral, palmar. - Capsular. - Lateral. - Capsular. - Lateral. - Annular. - Vaginal or flexor tendons. - Vaginal or crucial of the phalanges. - Accessory of the flexor tendons. - Posterior annular. - Vaginal of extensors. - Transverse of extensors. - Capsular and accessory slips. - Round or teres ligament. - Cartilaginous ligament. - Double cartilaginous ligament. - Ligamenta mucosa. - Lateral, — internal and external. - External short lateral ligament. - Posterior ligament. - Ligament of the patella. - Capsular. - Ligamentum alare, — majus et minus. - Ligamentum mucosum. - Two crucial, — anterior and posterior. - Transverse. - Capsular. - Interosseous. - Anterior superior. - Posterior superior. - Anterior ligament of the fibula. - Posterior of fibula. - Deltoides of tibia. - Capsular.

- Capsular.
Short ligaments.
Capsular, broad superior, and lateral ligaments, connecting astragalus and navicular.
Superior, lateral, and inferior, fixing os calcis to os cuboides.
Long, oblique, and rhomboid, forming the inferior ligaments.
Superior superficial, interosseous and inferior transverse ligaments, fixing the os naviculare and os cuboides.
Superior lateral, and plantar, which fix the os naviculare and cuneiform.
Superior-superficial and plantar, connecting the os cuboides, and os cuneiforme externum.
Dorsal and plantar, uniting the ossa cuneiformia.
The proper capsular of each bone.
22. *Of the Tarsus*.....
Capsular.
Dorsal, plantar, lateral, straight, oblique, and transverse.
24. *Connecting the Metatarsal Bones.*
Dorsal, plantar, and lateral, connecting the metatarsal bones.
Transverse ligaments.
25. *Of the Phalanges of the Toes.*
Capsular.
Lateral.
Annular.
Vaginal of the tendons of the peronei.
Laciniated.
Vaginal of the tendon of the flexor longus pollicis.
Vaginal and crucial of the tendons of the flexors of the toes.
Accessory of the flexor tendons of the toes.
Transverse of the extensor tendons.
26. *Retaining the Tendons of the Muscles of the Foot and Toes in situ.*
Vaginal of the tendon of the flexor longus pollicis.
Vaginal and crucial of the tendons of the flexors of the toes.
Accessory of the flexor tendons of the toes.
Transverse of the extensor tendons.

LIGAMENT, ANTERIOR, OF THE BLADDER. A name given by the older anatomists to a portion of the superior pelvic aponeuroses, which becomes attached to the front of the neck of the bladder.

LIGAMENT ARTÉRIEL, Arterial ligament—*l.* Camper's, Perineal fascia—*l.* Cervical supraspinal, see Supraspinosa ligamenta—*l.* Ciliaire, Ciliary ligament.

LIGAMENT, CORONARY, OF THE LIVER, is a reflection formed by the peritoneum, between the posterior margin of the liver, and the lower surface of the diaphragm. See Falx.

LIGAMENT, COSTOXIPHOID, Xiphoid ligament—*l.* Dentelé, Denticulatum ligamentum—*l.* Denticulé, Denticulatum ligamentum—*l.* Dorsolumbo-supraspinal, see Supraspinosa ligamenta—*l.* of Fallopius, Crural arch—*l.* Gimbernati's, see Gimbernati's ligament—*l.* Glossepiglottic, see Glossepiglottic—*l.* Infrapubian, Triangular ligament—*l.* Intérépineux, Interspinal ligament—*l.* of the Ovary, see Ovarium—*l.* Palmar inferior, Metacarpal ligament—*l.* Poupert's, Crural arch—*l.* Rond, Teres ligamentum—*l.* Sur-épineux cervical, see Supraspinosa ligamenta—*l.* Sur-épineux-dorsolombaire, see Supraspinosa ligamenta—*l.* Suspensory, of the liver, see Suspensory—*l.* Triangular, Perineal fascia.

LIGAMENT OF ZINN, *Ten'don of Zinn, Ten'do oculi*. A tendon from which the inferior rectus muscle of the eye arises, in common with the rectus internus and rectus externus.

LIGAMENTA ALARIA, Odontoid ligaments—*l.* Apicum, see Interspinal ligament—*l.* Coli, see, Colon—*l.* Interlobularia pulmonum, Interlobular tissue—*l.* Intervertebralia, Intervertebral cartilages—*l.* Lata uteri, see Uterus—*l.* Maucharti, Odontoid ligaments—*l.* Rotunda uteri, Round ligaments of the uterus—*l.* Tarsæ lata, see Tarsæ lata (ligament'a).

LIGAMENTARIS, Ligamentous.

LIGAMENTEUX, Ligamentous.

LIGAMENTOSUS, Ligamentous.

LIGAMENTOUS, *Ligamento'sus, Ligamenta'ris, Ligamentarius, Des'mous, Desmo'sus, (F.) Ligamenteux, Desmeux*. Having the character of, or relating to, a ligament.

LIGAMENTS, BROAD, OF THE UTERUS, see Uterus—*l.* Croisés, Crucial ligaments—*l.* Cruciformes, Crucial ligaments—*l.* Jaunes, Yellow ligaments—*l.* Grandes de l'utérus, see Uterus—*l.* of the Larynx, inferior, Thyreoarytenoid ligaments—*Ronds de l'utérus*, Round ligaments of the uterus—*l.* Thyrohyoid, see Thyrohyoid membrane.

LIGAMENTUM, Ligament—*l.* Arteriosum, see Arterial duct—*l.* Capsulare, Capsular ligament—*l.* Ciliare, Ciliary ligament—*l.* Conoïdes, Conoid ligament—*l.* Dentatum, Denticulatum ligamentum—*l.* Fallopii, Crural arch—*l.* Iridis, Ciliary ligament—*l.* Latum, Suspensory ligament of the liver—*l.* Membranaceum, Glosso-epiglottic ligament—*l.* Nuchæ, Cervical ligament—*l.* Posticum Winslowii, see Genu—*l.* Suspensorium dentis, see Odontoid ligaments—*l.* Suspensorium hepatis, Suspensory ligament of the liver—*l.* Suspensorium testis, Gubernaculum testis—*l.* Teres, Teres ligamentum—*l.* Trapezoides, Trapezoid ligament—*l.* Triangulare, Triangular ligament.

LIGATE, see Ligature.

LIGATIO, Fascia, Ligature—*l.* Lingue, Aphonia.

LIGATION, see Ligature.

LIGATURA, Ligature, Fascia—*l.* Glandis, Phimosis.

LIG'ATURE, *Ligatu'ra, Liga'tio, Alligatu'ra, Deliga'tio, Vincitu'ra, Fas'cia, Epid'esis*. This word has various acceptations. It means, 1. The thread with which an artery or vein is tied, to prevent or arrest hemorrhage. 2. The cord, or thread, or wire, used for removing tumours, &c. 3. The bandage used for phlebotomy. *Ligature* is, also, sometimes applied to the act of tying an artery or considerable vessel—*Liga'tion, Deliga'tion*; and occasionally we meet with the terms 'to *ligature*' and 'to *ligate*,' used unnecessarily in this sense, in place of to take up or tie a vessel. When the artery alone is tied, the ligature is said to be *immediate*; when any of the surrounding parts are included, it is said to be *mediate*. The ligature occasions obliteration or adhesion of the arterial parietes, by cutting through the middle and internal coats; the adhesion being favoured by the formation of a coagulum, which acts, in some degree, as a barrier against the impulse of the blood, and subsequently disappears by absorption.

LIGATURE ARTICULÉE, *Écraseur*.

LIGHT, (Sax. leoh't, lht,) (G.) Licht, *Lus, Lu'men, Phos, (F.) Lumière*. An extremely rare fluid; diffused over the universe; emanating from the sun and fixed stars; traversing more than four millions of leagues in a minute; passing through transparent bodies, which refract it according to their density and combustibility; and arrested by opaque bodies, by which it is reflected at an angle equal to the angle of incidence. It is the cause of colour in all bodies, being entirely reflected by white surfaces and absorbed by black. It is decomposed in passing through a transparent prism into seven rays—red, orange, yellow, green, blue, purple, and violet.

Light acts upon the body as a gentle and salutary stimulus. It urges to exercise, whilst privation of it induces sleep and inactivity, and disposes to obesity. Hence it is, that, in rural economy, animals which are undergoing the process of fattening are kept in obscurity. When vegetables are deprived of light, their nutrition is interfered with, and they become *etiolated*. To a certain extent this applies to animals, and there

is every reason to believe, that want of light prevents the due development of organized bodies. It has been found that, when tadpoles were deprived of light, they did not undergo the perfect metamorphosis into the frog, but that monstrosities from arrest of development were induced.

LIGHTNING, (from *light*.) *Astrape*.

LIGHTS, (from *lht*, 'light,' from their little specific gravity.) Pulmo—*l*. Rising of the, Cynanche trachealis.

LIGNE, Line—*l*. *Apré*, Linea aspera—*l*. *Blanche*, Linea alba—*l*. *Courbe*, Curved line—*l*. *Médiane* de l'abdomen, Linea alba—*l*. *Semilunaire* de Spiegel, Linea semilunaris Spigelii—*l*. *Sou-trochantérienne*, Linea aspera.

LIGNEOUS, *Ligno'sus*, (*lignum*, 'wood,') (F.) *Ligneux*. Woody. Having the character or appearance of wood; as '*Ligneous* Scirrhus,' or S. of the mamma, which has for its principal characteristics, hardness, the inextensibility of wood, and no defined limits. (Velpeau.) A form of this, in plates, or diffused over the mammary integuments, is termed, by M. Velpeau, *Squirre en cuirasse*.

LIGNIN, (from *lignum*, 'wood,') Ligneous or woody fibre; the fibrous structure of vegetable substances.

LIGNUM ALOËS, ('wood of aloes,') *Agallochum*—*l*. *Aspalathi*, *Agallochum*—*l*. *Benedictum*, *Guaiaicum*—*l*. *Brasilianum rubrum*, *Hæmatoxylo-n campechianum*—*l*. *Cæruleum*, *Hæmatoxylo-n campechianum*—*l*. *Campechense*, *Hæmatoxylo-n campechianum*—*l*. *Campechianum*, *Hæmatoxylo-n campechianum*—*l*. *Campescanum*, *Hæmatoxylo-n campechianum*—*l*. *Colubrinum*, see *Strychnos*—*l*. *Febrifugum*, *Quassia*—*l*. *Hæmatoxyli*, *Hæmatoxylo-n campechianum*—*l*. *Indicum*, *Guaiaicum*, *Hæmatoxylo-n campechianum*—*l*. *Infelix*, *Sambucus*—*l*. *Nephriticum*, *Guilandina Moringa* (the wood)—*l*. *Pavana*, *Croton tiglium*—*l*. *Quassia*, see *Quassia*—*l*. *Sanctum*, *Guaiaicum*—*l*. *Sandalinum*, see *Pterocarpus santalinus*—*l*. *Sappan*, *Hæmatoxylo-n campechianum*—*l*. *Serpentum*, *Ophioxylum serpentinum*—*l*. *Vitæ*, *Guaiaicum*.

LIGNYODES, (*λινυοδές*, from *lignys*, and *odes*.) Fuliginous.

LIGNYS, (*λινυς*.) Fuligo.

LIG'ULA, *Lin'gula*, 'a small tongue, a tongue of land; the tongue of a shoe.' The clavicle; also, the giottis and epiglottis. A measure containing 3 drachms and a scruple, or about half an ounce. Also, a species of bandage or ligature.

LIGUSTICUM, (from *λινυστικός*, appertaining to *Liguria*.) *L. Levisticum*.

LIGUSTICUM ACTÆIFO'LIUM, *Thapsium actæifo-lium*, *Angel'ica lū'cida*, *Angel'ica Nen'do*, *White root*, indigenous; *Ord.* Umbelliferae. The root is aromatic and carminative.

LIGUSTICUM CAPILLACUM, *Æthusa meum*—*l*. *Carvi*, *Carum*—*l*. *Fœniculum*, *Anethum*.

LIGUSTICUM LEVISTICUM, *Levisticum*, *Laserpitium Germanicum*, *Ligusticum*, *Angel'ica levis'ticum* seu *paludapifolia*, *Lov'age*, (F.) *Livèche*, *Ache des montagnes*. *Ord.* Umbelliferae. The properties of this plant are said to be stimulant, carminative, emmenagogue, &c.

LIGUSTICUM MEUM, *Æthusa meum*—*l*. *Phellandrium*, *Phellandrium aquaticum*.

LIGUSTICUM PODAGRA'RIA, *Podagra'ria ægopo'dium*, *Ægopo'dium* seu *Sil'son podagra'ria*, *Pimpinella angelicifolia*, *Sil'um vulga're*, *Tragoselinum angel'ica*, *Ses'eli ægopo'dium*, *Angel'ica sylvestris*, *Goutweed*. A British plant, once considered useful in cases of gout.

LIGUSTICUM SILAUS, *Peucedanum silaus*.

LIGUSTRUM ÆGYPTIACUM, (*ligare*, 'to bind,' because used for bands.) *Lawsonia inermis*.

LIGUSTRUM VULGA'RE, *Privet*, *Priv'y*, *Prim*, (F.) *Troëne*. *Ord.* Oleaceæ. A shrub, which grows wild both in Europe and in the United

States, usually in hedges. The leaves are astrigent and bitter; and the flowers, which are snow-white, and of an agreeable odour, have been employed in decoction, in sore throat, and ulcerous stomatitis. The berries are said to be cathartic.

LILAC, COMMON, *Syringa vulgaris*—*l*. *Vulgaris*, *Syringa vulgaris*.

LILI, *Lilium* Paracelsi.

LILIA'GO, (dim. of *lilium*, 'the lily,') *Spiderwort*; *Lilias'trum*. This plant was formerly said to be alexipharmic and carminative.

LILIASTRUM, *Liliago*.

LIL'LIUM CAN'DIDUM, ('white lily,') *L. album*, *Lei'ron*, *Cri'non*. The white lily. (F.) *Lis blanc*. *Ord.* Liliaceæ. The infusion of the flowers of the lily in olive oil is emollient, and often applied externally, under the name of *Lily oil*, (F.) *Huile de lis*. The scales of the bulb, roasted, are sometimes employed as maturatives.

LILIUM CONVALLIUM, *Convallaria maialis*.

LILIUM PARACEL'SI, *Tinctura Metallorum*, *Lili*. A medicine employed by Paracelsus. It was prepared by melting in a crucible four ounces of each of the following alloys; *Antimony and iron*, *antimony and tin*, *antimony and copper*, previously mixed with eighteen ounces of nitrate of potassa and as much salt of tartar. The melted mass, when pulverized, was treated with rectified alcohol, which really dissolved only the potassa set at liberty by the decomposition experienced by the nitre and salt of tartar. The *Lilium Paracelsi* was used as a cordial. It entered into the composition of the theriacal elixir, &c.

LILY, *Lilium*—*l*. *Ground*, *Trillium latifolium*—*l*. *May*, *Convallaria maialis*—*l*. *Pond*, *Nelumbium luteum*—*l*. *Pond*, white, *Nymphæa odorata*—*l*. *Pond*, yellow, *Nuphar advena*—*l*. *Toad*, *Nymphæa odorata*—*l*. of the Valley, *Convallaria maialis*—*l*. *Water*, little, *Brasenia hydropeltis*—*l*. *Water*, sweet, *Nymphæa odorata*—*l*. *Water*, sweet-scented, *Nymphæa odorata*—*l*. *Water*, white, *Nymphæa alba*—*l*. *Water*, yellow, *Nymphæa lutea*—*l*. *White*, *Lilium candidum*, *Nelumbium luteum*.

LIMA, CLIMATE OF. The climate of Peru does not appear to be favourable to the generation of consumption; and Lima would seem to be a good residence for the phthisical valetudinarian. Many have been benefited by a residence there; but when they have gone farther south, as to Chili, the effect, according to Dr. M. Burroughs, has generally been fatal.

LIMA DENTARIA, ('dental file,') *Scalprum dentarium*.

LIMACIEN (F.), (from *limaçon*, [*limax*, gen. *lima'cis*, 'the snail,'] 'the cochlea of the ear,') A branch of the acoustic or labyrinthic nerve sent to the cochlea.

LIMACIEN, *Cochlea*, *Helix pomatia*, *Limax*—*l*. *Rampes du*, see *Cochlea*.

LIMACUM CORNUA, ('snails' horns,') see *Lachrymal puncta*.

LIMANCHIA, (*λινανχία*, from *limus*, and *αγχεν*, 'to torture,') Abstinence.

LIMA'TIO, (*limare*, *limatum*, 'to file,') 'Filing.' An operation employed by the dentist more especially to prevent immediate contact of the teeth with each other.

LIMATURA FERRI, *Ferri Limatura*.

LIMAX, gen. *Lima'cis*, (*λίμαξ*, gen. *λίμακος*.) *Hel'ix*, *Slug* or *Snail*, (F.) *Limaçon*, *Colimaçon*, *Escarbot*. A syrup has been prepared from these animals, which has been given in phthisis, &c. The raw snails have also been taken in consumption. They have been used as food.

LIMB, (Sax. *lim*.) *Membrum*.

LIMBUS ('a border') **ALVEOLARIS**, Alveolar border—*l*. *Fossæ Ovalis*, see *Ovalis fossa*—*l*. *Luteus retinae*, see *Foramen centrale*—*l*. *Posterior corporis striati*, *Tænia semicircularis*.

LIME, (from *limon*.) A fruit like a small

lemon, the juice of which is strongly acid, and is much used for making punch. It is, also, used in long voyages as an antiscorbutic, &c. It is a species of lemon, the fruit of *Citrus ac'ida*, *Citrus lim'etta*.

LIME, (Sax. lim.) Calx—1. Carbonate of, Creta—1. Chloride of, Calcis chloridum—1. Chlorite of, Calcis chloridum—1. Chloruret of, Calcis chloridum—1. Hydrate of, see Calx—1. Hydrosulphate of, Calcis sulphuretum—1. Hypochlorite of, Calcis chloridum—1. Hypophosphite of, Calcis hypophosphis—1. Muriate of, solution of, see Calcis murias—1. Oxymuriate of, Calcis chloridum—1. Phosphate of, see Cornu ustum—1. Bone Phosphate of, see Cornu ustum—1. Phosphate of, precipitated, see Cornu cervi—1. Slaked, see Calx—1. Sulphuret of, Calcis sulphuretum.

LIME TREE, Tilia—1. Water, Liquor calcis—1. Water, compound, Liquor calcis compositus.

LIMELLIER, see Citrus mella rosa.

LIMICUS, (from *limus*.) Hungry.

LIMITROPHES, (*lines*, 'a boundary,' and *τροφειν*, 'to nourish,') see Trisplanchnic nerve.

LIMNE, (λίμνη,) Marsh.

LIMNE'MIC, *Limnæ'mic*, *Limnhe'mic*, *Limnæ'micus*, *Limnhe'micus*, (λίμνη, 'a marsh,' and *αἷμα*, 'blood,') An epithet given to affections induced by paludal emanations; (F.) *Affectiones limn-hémiques*.

LIMNOMEPHITIS, (*limne*, and *mephitis*,) see Miasm.

LIOMOCTONIA, (λιμοκτονία, from *limus*, and *κτονος*, 'death,') Abstinence to death. Death from hunger. Suicide by hunger.

LIMODES, (λιμώδης, *limus*, and *odes*,) Hungry.

LIMON, see Citrus medica.

LIMONADA, Lemonade.

LIMONADE, Lemonade—1. *Gazeuse*, see Lemonade—1. *Sèche*, Lemonade, dry; see Citric acid.

LIMONIUM, Statice limonium—1. *Malum*, see Citrus medica.

LIMONUM BACCA, see Citrus medica.

LIMOPHTHISIS, (from *limus*, and *φθίσις*, 'wasting,') Wasting from privation of food.

LIMO'SIS, (from *limus*,) *Stomach disease*, *Morbid appetite*. A genus in the class *Caliciaca*, order *Enterica*, of Good.

LIMOSIS CARDIALGIA MORDENS. Cardialgia—1. Cardialgia sputatoria, Pyrosis—1. Dyspepsia, Dyspepsia—1. Expers, Anorexia—1. Expers protracta, Fasting—1. Flatus, Flatulence—1. Heliunum, Gluttony—1. Pica, Malacia.

LIMOTHERAPEIA, *Nestitherapeia*, *Nestotherapeia*, *Nestiatr'i'a*, *Peinotherapeia*, *Cura'tio mor'bi per in'e'diam*, (*limus* and *Σεπάρεια*, 'treatment,') Hunger-cure. Cure by fasting.

LIMUS, (λίμος,) Hunger.

LIN, *Linum usitatissimum*—1. *Graines de*, see *Linum usitatissimum*—1. *Purgatif*, *Linum catharticum*.

LINAIRE. Antirrhinum linaria.

LINAMENTUM, (from *linum*,) Lintum.

LINARIA, (from *linum*, owing to the similarity of the leaves,) Antirrhinum linaria—1. Cymbalaria, Antirrhinum linaria—1. Elatine, Antirrhinum elatine—1. Vulgaris, Antirrhinum linaria.

LINCTUARIUM, (from *lingere*, *linctum*, 'to lick,') Electos.

LINCTUS, gen. *Linctus*, Electos—1. Albus, Looch album—1. Amygdalinus, Looch album—1. ad Aphthas, Mel boracis—1. de Borace, Mel boracis—1. Communis, Looch album.

LIND, (Sax. lind,) Tilia.

LINDEN TREE, Tilia, Tilia Americana.

LINE, *Lin'ea*, *Gram'me*, (F.) *Ligne*. Extent in length, considered without regard to breadth or thickness. As a measure, it means the 12th part of an inch; with some, however, the 10th part of an inch.

MEDIAN LINE OF THE BODY is an imaginary line supposed to set out from the top of the head and to fall between the feet, so as to divide the body vertically into two equal and symmetrical parts.

LINE, (from *linum*,) Lintum, *Linum usitatissimum*.

LINÆA, Line.

LIN'EA AL'BA, *Linea alba Abdom'inis*, (F.) *Ligne blanche ou médiane de l'abdomen* (Ch.), *Cand'id'ula abdom'inis lin'ea*, *Lin'ea cent'ral'is*. A tendinous, strong, and highly resisting cord; extending from the ensiform cartilage of the sternum to the symphysis pubis, with the umbilicus near its middle. The linea alba is formed by the decussation of the aponeurosis of the abdominal muscles; and its use is to limit the movement of the chest backwards; to prevent it from separating too far from the pelvis, and to furnish a fixed point for the muscles of the abdomen in their contraction.

LINÆA AS'PERA, *Rug'ged ridge*, (F.) *Ligne âpre*, *Ligne sous-trochantérienne*, (Ch.) A rough projection at the posterior surface of the femur, which gives attachment to muscles.

LINÆA CANDIDULA ABDOMINIS, L. alba—1. Centralis. L. alba—1. Innominata. Heopectinea Linea.

LINÆA ALBICANTES. (pl. of *linea*, 'a line,') The white lines left where the skin, as of the abdomen or mammae, has been greatly stretched.

LINÆA ARCUATÆ, Curved lines—1. Semicirculares externæ, Curved lines.

LINÆA SEMILUN'ARES are the lines which bound the outer margin of the recti muscles of the abdomen. They are formed by the union of the abdominal tendons. The lines which cross these muscles are called *Linæa transvers'æ*.

LINÆA SEMILUN'ARIS SPIGE'LII, (F.) *Ligne semi-lunaire de Spiegel*. The semilunar notched inner edge of the fleshy portion of the transversalis abdominis, at the point where the muscular fibres are in connexion with the tendinous or aponeurotic fibres, which serve for the insertion of the muscle.

LINÆA TIB'Æ, *An'guli tib'ia*. Sharp lines on the tibia.

LINÆA TRANSVERSÆ, see L. semilunares, and Processus teretes.

LINÉAIRE, Linear.

LIN'EAMENT, *Lineamen'tum*, (*linea*, 'a line,' *lineare*, 'to mark by lines,') A delicate trait observed on the countenance, which constitutes its special character, enables us to preserve its image, and is the cause of resemblance to others. A feature.

Bonnet gave the name *Linéament* to the first traces of organization in the embryo of man and animals.

LIN'EAR, *Linea'ris*, *Linea'rius*, (from *linea*, 'a line,') (F.) *Linéaire*. Pathologists apply the epithet *linear* to fractures which are very narrow, and in which the fragments are scarcely separated.

LIN'EOLA, (dim. of *linea*,) A small line.

LINEOLE MAMMA'RUM. The white lines on the breasts.

LINGISM, (after Ling, of Sweden, the proposer,) Kinesipathy.

LINGUA, (from *lingere*, 'to lick,') Tongue—1. Avis, see Fraxinus excelsior—1. Bovis, Anchusa officinalis—1. Canina, Cynoglossum—1. Cervina, Asplenium scolopendrium—1. Exigua, Epiglottis—1. Prognosis ex, Glossomantia—1. Serpentaria, Ophioglossum vulgatum.

LINGUADENT'AL, *Linguodental* *Linguadenta'lis*, *Linguodenta'lis*, (from *lingua*, 'the tongue,' and *dens*, gen. *dentis*, 'a tooth,') Relating or appertaining to the tongue and teeth.

LINGUE DETENTOR, Glossocatochus—*l. Exoncosis*, Glossoncus—*l. Scalpium*, *Cure-langue*.

LINGUAL, *Lingualis*, (from *lingua*.) Relating or belonging to the tongue.

LINGUAL ARTERY, *Arteria lingualis*, arises from the external carotid; and, after several tortuosities, reaches the base of the tongue, becomes horizontal, and, under the name *Ravine*, advances to the tip, where it anastomoses with its fellow. In its course, it gives off the *Dorsalis lingue* and *sublingual*.

LINGUAL MUSCLE, *Lingualis*, *Basilio-glossus* (Cowper), *Glossianus*, is a small, long, fasciculus of fibres, hidden beneath the sides of the tongue, between the hyoglossus and styloglossus muscles, which are on the outside, and the genioglossus, within. This muscle passes from the base to the tip of the tongue; and, at its sides, is confounded with the muscles just referred to. It shortens the tongue, depresses its point, and can carry it to the right or left side.

LINGUAL NERVE is a name which has been given to the ninth pair or hypoglossus. It is, generally applied to a branch given off from the *Inferior maxillary*, or third branch of the fifth pair. Near its origin it runs close to, and anastomoses with, the chorda tympani. It afterwards gives off a considerable number of filaments which are distributed to the tongue, and some of which are said to have been even traced as far as the papillæ;—a distribution which has occasioned it to be regarded as the *Gustatory nerve*, (*F.*) *Nervus gustatif*. It is doubtless the nerve of general sensibility to the tongue.

LINGUAL VEIN follows nearly the same distribution as the artery. It opens into the internal jugular.

LINGUALIS, *Lingual* muscle.

LINGUETTA LAMINOSA, (dim. of *lingua*), *Lingula*.

LINGULA, (dim. of *lingua*), *Ligula*.

LINGULA, *Linguetta lamino'sa*. A thin, transversely grooved lobule of gray substance, derived from the anterior border of the cerebellum, which, for a short distance, lies over the velum medullare anterius. A depression in the cerebellum in front of the lingula is called (*F.*) *Niche*, *Recessus*.

LINGULA FISTULE, Epiglottitis.

LINGUODENTAL, *Linguadental*.

LINI FARINA, see *Linum usitatissimum*—*l. Usitatissimum semina*, see *Linum usitatissimum*.

LINIMENT, *Linimentum*, *Lin'tus*, *Enchris'ton*, *Catachris'ton*, *Perich'risis*, *Perichris'ton*, *Aleiph'a*, *Aleim'na*, *Enchris'ta*, *Fric'tum*, *Fricato'rium*, *Inunctio*, (from *linire*, *linitum*, 'to anoint.') An unctuous medicine, containing usually oil or lard, which is used externally in the form of friction.

LINIMENT OF ACONITE, *Linimentum Aconiti*—*l. of Ammonia*, Compound, see *Lotion*, *Granville's*.

LINIMENT AMMONIACAL, *Linimentum ammoniæ*—*l. de carbonate d'Ammoniaque*, *Linimentum ammoniæ carbonatis*.

LINIMENT, ANODYNE, *Linimentum opii*.

LINIMENT ANTHÉMORRHŒDALE DE SIEUR ANDRY, (*F.*) This is formed of Narbonne honey, Olive oil, and Turpentine.

LINIMENT ANTIPARALYTIQUE, (*F.*) Composed of subcarbonate of ammonia alcoholized, oil, black soap, and oil of rosemary.

LINIMENT, ANTISCROFULOUS, of HUFELAND. Composed of fresh ox-gall, White soap, Unguentum althææ, Volatile oil of petroleum, Carbonate of ammonia, and Camphor.

LINIMENT OF BELLADONNA, *Linimentum Belladonnæ*.

LINIMENT CALCAIRE, *Linimentum aquæ calcis*—*l. Camphor*, *Linimentum camphoræ*—*l. Camphré*, *Linimentum camphoræ*—*l. of Cantharides*, camphorated, *Linimentum e cantharidibus*

camphoratum—*l. Cantharide camphré*, *Linimentum e cantharidibus camphoratum*—*l. of Chloroform*, *Linimentum chloroformi*—*l. Croton oil*, *Linimentum crotonis*—*l. d'Eau de chaux*, *Linimentum aquæ calcis*—*l. of Iodine*, *Linimentum Iodii*—*l. of Limewater*, *Linimentum aquæ calcis*—*l. de Mercure*, *Linimentum hydrargyri*—*l. Mercurial*, *Linimentum hydrargyri*—*l. Oléo-calcaire*, *Linimentum aquæ calcis*—*l. of Opium*, *Linimentum opii*.

LINIMENT RÉSOLUTIF DE POTT, composed of oil of turpentine and muriatic acid. Used in rheumatism, swellings, &c.

LINIMENT, SAINT JOHN LONG'S. A liniment used by a celebrated empiric at the commencement of the second quarter of the 19th century. It is said to have consisted of oil of turpentine and acetic acid, held in suspension by yolk of egg. It was a powerful counterirritant.

LINIMENT DE SAVON, *Linimentum saponis compositum*—*l. de Savon opiacé*, *Linimentum saponis et opii*.

LINIMENT SAVONNEUX HYDROSULFURÉ DE JADELLOT. Composed of sulphuret of potass, ʒij; white soap, lbij; oil of poppy, lbiv; oil of thyme, ʒij. Used in itch.

LINIMENT, SIMPLE, *Linimentum simplex*—*l. Soap*, *Linimentum saponis*—*l. Soap and opium*, *Linimentum saponis et opii*—*l. Soap*, compound, *Linimentum saponis compos.*—*l. Turpentine*, *Linimentum terebinthinæ*—*l. of Turpentine and Acetic Acid*, *Linimentum Terebinthinæ aceticum*—*l. of Verdigris*, *Linimentum æruginis*—*l. de Vert-de-gris*, *Linimentum æruginis*—*l. Volatile*, *Linimentum ammoniæ*.

LINIMENTUM, *Liniment*.

LINIMENTUM ACONITI, *Liniment of Aconite*. (*Aconite root*, in powder, ʒxx; Camphor, ʒj; Rectified spirit, fʒxxx or q. s. Moisten the aconite with a portion of the spirit and macerate for seven days; then percolate into a receiver containing the camphor, until one pint is procured. Ph. B. 1864.) In neuralgic and other pains,—℥xx or xxx being rubbed on the part.

LINIMENTUM ÆRUGINIS, *Oxymel Æruginis*, *Unguentum Ægyptiacum*, *Mellitum de aceto'isæ cu'pri*, *Liniment of verd'gris*, (*F.*) *Liniment de Vert-de-gris*, *Miel d'acétate de cuivre*. (*Ærugin*, cont. ʒj; Aet. ʒvij; Mellis despum. pond. ʒxiv. Liquefied, strained, and inspissated by boiling.—Ph. L.) It is used as an escharotic and detergent;—diluted, as a gargle in venereal ulcerations and in foul ulcers.

LINIMENTUM ALBUM, *Ceratum cetacei*, *Unguentum cetacei*—*l. ad Ambustiones*, *L. aquæ calcis*.

LINIMENTUM AMMONIÆ, *Sa'po ammoniacal'is*, *Linimentum ammoniatum seu ammoniacum seu Anglica'rum*, *Sa'po ammoniæ olea'ceus*, *Unguentum album resolutens*, *O'leum ammoniatum*, *Strong liniment of ammo'nia*, *Volatile liniment*, (*F.*) *Liniment volatil ou ammoniacal*, *Savon ammoniacal*. (*Aq. ammon.* fʒj; *Olei olivæ*, ʒij. Mix.—Ph. U. S.) A stimulating and rubefacient soap.

LINIMENTUM AMMONIÆ CARBONATIS, *Liniment of subcarb'onate of ammo'nia*, *Linimentum ammoniæ sesquicarbonat'is*, *L. volat'ilè*, *Harts'horn and oil*, (*F.*) *Liniment de carbonate d'ammoniaque*. (*Solut. subcarb. ammon.* fʒj; *Olei oliv.* fʒij. Shake till they unite.—Ph. L.) A stimulating liniment, mostly used to relieve rheumatic pains, bruises, &c.

LINIMENTUM AMMONIÆ COMPOSITUM, see *Granville's Lotion*—*l. Ammoniæ Sesquicarbonatis*, *Linimentum Ammoniæ Carbonatis*—*l. Ammoniatum seu Ammoniacum*, *L. Ammoniæ fortis*—*l. Anglicanum*, *L. Ammoniæ fortis*—*l. ad Aphthas*, *Mel boracis*.

LINIMENTUM AQUÆ CALCIS, *L. Calcis* (*Ph. U. S.*), *L. ad ambustio'nes*, *O'leum li'ni cum cal'cè*, *Sa'*

po calca'rius, Lin'iment of lime'water, Car'ron oil, (because much used in the Carron iron works in Scotland), (F.) *Liniment d'eau de chaux*, *Savon calcaire*, *Liniment calcaire ou oléo-calcaire*. (*Olei lini*, 3vij; *Aquæ calcis*, f3vij. *Misce.*) A cooling and emollient application to burns and scalds.

LINIMENTUM ARCEI, Unguentum elemi composi-
tum.

LINIMENTUM BELLADONNÆ, Lin'iment of Bel-
ladonna. (*Belladonna root*, in powder, 3xx; *Camphor*, 3j; *Rectified spirit*, f3xxx, or a sufficiency. Prepared like Linimentum Aconiti, so that a pint is produced.—Ph. B. 1864.) Applied like Extractum Aconiti.

LINIMENTUM DE BORACE, Mel boracis—*L. Calcis*, *L. aquæ calcis*.

LINIMENTUM CAMPHORÆ, *O'leum Camphoratum*, *Solutio camphoræ oleosa*, *Camphor lin'iment*, (F.) *Liniment Camphré*, (*Camphoræ*, 3ij; *Olei olivæ*, 3xij. Dissolve.—Ph. U. S.) It is used as a stimulant and discutient.

LINIMENTUM CAMPHORÆ COMPOSITUM, *L. saponato-ammoniato-camphoratum*, *Compound camphor liniment*, *Ward's essence for the head-ache*. (*Camphor*, 3iiss; *Ol. Lavand.* f3j; *Sp. rectificat.* f3xvij; *Liquor ammon. fortior* f3ij. Dissolve the camphor and oil in the spirit; add the ammonia, and shake till they are mixed. Ph. L.) It is stimulant and discutient.

LINIMENTUM CANTHARIDIS, Lin'iment of Span-
ish Flies. (*Cantharid.* in pulv. 3j; *Ol. terebinth.* Oss. Digest for three hours by means of a water-bath, and strain.—Ph. U. S.) Used as an excitant liniment in adynamic cases.

LINIMENTUM E CANTHARIDIBUS CAMPHORATUM, *Camph'orated lin'iment of cantharides*, (F.) *L. cantharide camphré*. (*Tinct. cantharid.* 3ss; *Ol. amygd. dulc.* 3iv; *Sapon. amygd.* 3j; *Camphor.* 3ss. Dissolve the camphor in the oil, and add the mixture to the tincture and soap.—Ph. P.) Rubefacient, and discutient.

LINIMENTUM CHLOROFORMI, Lin'iment of Chlo-
roform. (*Chloroform. purif.* 3ij; *Ol. Oliv.* 3iv. M.—Ph. U. S. 1863.) f3ss to f3j to be applied.

LINIMENTUM CROTONIS, *Cro'ton Oil Lin'iment*, (*Ol. Croton.* f3j; *Ol. Terebinth.* f3vij. M. Ph. D. *Croton oil*, f3ss; *Olive oil*, f3iiss, Mix, Ph. B. 1864.) Rubefacient, causing an eruption when continued.

LINIMENTUM HYDRARGYRI, *L. Hydrargyri composi'tum*, *Mercurial lin'iment*, (F.) *L. de Mer-
cure*. (*Unq. Hyd. fort.* *Adip. præp.* 3iiv; *Camphoræ*, 3j; *Sp. rect.* 3t. xv; *Liquor ammon.* f3iv. Rub the camphor with the spirit; add the ointment and lard; and, lastly, gradually add the solution.—Ph. L.) It is used as a stimulant and discutient to venereal swellings, &c.

LINIMENTUM HYPERICI, see Hypericum perfor-
atum.

LINIMENTUM IODI, Lin'iment of Iodine. (*Iod.* 3i; *Potass. iodid.* 3ss; *Sp. rectific.* f3v. Dis-
solve. Ph. B. 1864.) To paint the part.

LINIMENTUM OPII, Lin'iment of Opium, *An'o-
dyne Lin'iment*. (*Sapon.* 3vj; *Opii.* 3iiss; *Camphor.* 3ij; *Ol. Rosmarin.* f3vj; *Spirit rectificat.* Oij. (*Imp. meas.*) Macerate the soap and opium in the spirit for three hours; filter, and add the oil and camphor.—Ph. L.) Anodyne and slightly rubefacient.

LINIMENTUM PLUMBATUM, Unguentum plumbi
superacetatis—*L. Saponaceum opiatum*, *L. Sapo-
nis et opii*.

LINIMENTUM SAPONA'CEUM HYDROSULPHURATUM. (*Common Soap*, 500 p. Liquefy in a water
bath in an earthen vessel; and add white poppy
oil, 250 p. Mix intimately, and add dry sulphuret
of potass, 100 p. Beat together, and add oil of
poppy seed, 750 p.—Ph. P.) In cutaneous affec-
tions, as psora, herpes, &c.

**LINIMENTUM SAPONATO-AMMONIATO-CAMPORA-
TUM**, *L. Camphoræ compositum*—*L. Saponato-camphoratum*, *L. Saponis camphoratum*—*L. Saponis*,
L. S. camphoratum.

LINIMENTUM SAPONIS CAMPHORATUM, *L. S.*
composi'tum, *L. Saponato-camphoratum*, *Bal'sa-
mum opodel'doc*, *Tinctura saponis camphorata*
(Ph. U. S.), *Linimentum Saponis* (Ph. U. S. 1863),
Opodel'doch, *Steer's Opodel'doch*, *Soap Liniment*,
Compound Soap Liniment, (F.) *L. de Savon.*
(*Sapon. rasur.* 3iv; *Camphor.* 3ij; *Ol. Rosmarini*
f3ss; *Aquæ* f3iv; *Alcohol.* Oij. Mix the alcohol
and water: digest the soap in the mixture by
means of a water-bath, until it is dissolved; filter,
and add the camphor and oil.—Stimulant and
anodyne; in bruises, local pains, &c.

Freeman's Ba'thing Spir'its consist of *Lin. sapon.*
camphoratum, coloured with *Daffy's elixir*.

Jackson's Ba'thing Spir'its differ from *Freeman's*
in the addition of some essential oils.

LINIMENTUM SAPONIS COMPOSITUM, *L. S. camphoratum*.

LINIMENTUM SAPONIS ET OPII, *L. saponaceum*
opiatum, *Soap and opium lin'iment*, *Bates's an'o-
dyne bal'sam*, *Bal'samum anodynum*, *Tinctura sa-
ponis et opii*, (F.) *L. de Savon opiacé*. (*Saponis*
duri. 3iv; *Opii.* 3j; *Camphoræ*, 3ij; *Olei roris-
marin.* f3ss; *Alcohol.* Oij.—Ph. E.) Anodyne;
in chronic rheumatism and local pains in general.

LINIMENTUM SIMPLEX, *Simple Lin'iment*. (*Ol.*
oliv. 4 p.; *Ceræ albæ*, 1 p. fiat linimentum.—Ph.
E.) Emollient; used in chaps, &c.

LINIMENTUM SIMPLEX, Unguentum ceræ.

LINIMENTUM TEREBINTHINÆ, *L. Terebinthina'tum*,
Tur'pentine lin'iment. (*Cerat. resinæ*, 3ij; *Ol. tere-
binth.* Oss. Melt the cerate and stir in the oil.)
A stimulant; applied to burns, &c.

LINIMENTUM TEREBINTHINÆ ACETICUM, Lin'-
iment of Tur'pentine and Acetic Ae'id. (*Ol.*
Tereb. f3i; *Acid. Acet.* f3i; *Linim. Camphor.*
f3i; M. Ph. B. 1864.) Said to resemble St.
John Long's liniment.

LINIMENTUM VOLATILE, *L. Ammoniae carbonatis*.
LINN. (*Sax. linn.*) *Tilia Americana*.

LINNÆA, *L. Borealis*. *Ord. Caprifoliaceæ*,
(after Linnaeus.) This plant has a bitter, sub-
astringent taste; and is used, in some places, in
the form of fomentation, in rheumatic pains. An
infusion in milk is used in Switzerland for the
cure of sciatica.

LINOSPERMUM, (*λινος*, 'flax,' and *σπέρμα*,
'seed.') See *Linum usitatissimum*.

LINSEED, see *Linum usitatissimum*.

LINT, Carbasus, Lintum.

LINTEAMEN, Lintum, Pledget.

LINTEUM, 'a linen cloth,' (from *linum*, 'flax,
linen,') *L. carp'tum* seu *ra'sum*, *Linteamen*, *Oth'-
oné*, *Othon'ion*, *Carbasus*, *Lin'basu*, *Car'pia*, *Mo'-
tos*, *Mo'te*, *Mo'tum*, *Linamen'tum*, *Til'ma*, *Xys'tos*,
Xys'ma, *Ach'né*, *Lint*, *Line*, (F.) *Charpie*. A soft,
flocculent substance, made by scraping old linen
cloth, (F.) *Charpie râpée*, or by unravelling old
linen cut into small pieces—(F.) *Charpie brute*;
and employed in surgery as a dressing to wounds,
ulcers, &c., either simply or covered with oint-
ment. *Picked oak'um* has been recommended as
a substitute for lints in dressings.

LINTEUM CARPTUM, see Lintum—*L. Rasum*,
see Lintum—*L. Scissum*, *Splitcloth* (bandage.)

LINUM (*λινος*), see Lintum, and *Linum usita-*
tissimum—*L. Arvense*, *L. usitatissimum*.

LINUM CATHARTICUM, *L. min'imum*, *Chamæli-*
num, *Pur'ging flax*, *Mill moun'tain*, (F.) *Lin pur-*
gatif. *Ord. Linacææ*. This plant is possessed of
cathartic properties, and has a bitterish, disagree-
able taste. Dose, 3j, in substance.

LINUM CRUDUM, see Apolinosia—*L. Minimum*
L. Catharticum.

LINUM USITATISSIMUM, *L. arven* 3ē, *Com'mon*

flax, Lin. (F.) *Lin.* The seed, *Linum*, (Ph. U. S.), *Sem'ina li'ni usitatis'simi*, *Linosp'er'mum*, *Lin-seed*, *Flax'seed*, (F.) *Grains de lin*, are inodorous, and almost tasteless; yielding mucilage to warm water, and oil by expression. They are demulcent and emollient. Linseed, when ground into powder, *Li'ni Far'i'na*, *Flax'seed meal*, (Ph. U. S.) forms a good emollient poultice. It is only necessary to stir the powder into boiling water. The oil, *O'leum li'ni*, *Flax'seed* or *Lin'seed oil*, (F.) *Huile de Lin*, is emollient and demulcent.

LIONDENT, ('lion's tooth') *Leontodon taraxacum*.

LION'S FOOT, *Nabalus albus*, *Prenanthes*—l's. Tail, *Leonurus*.

LIOPODIA, see *Leiopus*.

LIOPUS, *Leiopus*.

LIP, (Sax. and Germ. *Lippe*.) *Chei'los*, *La'bium*, *La'brum*, (F.) *Lèvre*. [In Entomology, *labium* means the lower lip,—*labrum*, the upper.] The lips are composed of different muscular fasciuli, nerves, and vessels, covered by the skin and mucous membrane of the mouth. They circumscribe the anterior aperture of that cavity; and are inservient to mastication, pronunciation, &c. They are distinguished into *upper* and *lower*—*Anoche'il'on*; and *Catoche'il'on*, (Prov.) *Fipple*—and are placed in front of each jaw, forming between them the anterior aperture of the mouth. They unite at each side, and form what are called the *angles* or *commissures* of the mouth—*Chal'ini*. Their free edge is covered with a mucous membrane, of a more or less livid red, according to the individual. They receive their arteries from the external carotid. Their veins open into the two jugulars. Their lymphatic vessels descend into the ganglions situate beneath the chin. Their nerves are derived from the infraorbital, mental, and facial.

LIPS, *La'bia*, (F.) *Lèvres de la vulve*, are folds belonging to the genital organs of the female, and distinguished into—1. *La'bia puden'di* seu *Cun'ni seu majo'ra*, *Epist'a*, *A'tæ puden'di muliebris*, *Hyrtoche'ilides*, *Ru'pes*, *A'tæ majo'res*, *Crem'noi*, *Piniac'ula*, (F.) *Grandes Lèvres*. These are two membranous folds, thicker above than below, which limit the vulva laterally, and extend from the inferior part of the mons veneris to the perinæum. They unite anteriorly and posteriorly, forming commissures; the posterior of which is called *Fourchette*. Their outer surface is convex; formed of skin and covered with hair. The inner surface is white, and covered by a mucous membrane, continuous with that lining the other parts of the vulva. The space between the skin and mucous membrane is filled with a fatty tissue and fibrous bands, some fibres of the constrictor vaginae muscle, vessels, and nerves. 2. *Labia puden'di mino'ra*, see *Nymphæ*.

We speak, also, of the *lips* of a wound, ulcer, &c., when alluding to the edges of these solutions of continuity.

LIPÆMIA, *Piærhæmia*.

LIPARA, (from *liparus*.) Plasters, containing much oil or fat.

LIPARIA, (from *liparus*.) see *Polysarcia*.

LIPAROC'E'LE, *Lipoc'e'lê*, (*liparus*, and *κηλη*, 'tumour.') *Lipo'ma*, or fatty tumour of the scrotum, *Her'nia pinguedino'sa scroti*.

LIPARODES, (from *liparus*, and *odes*.) Fatty.

LIPAROIDÊ, *Cerate*.

LIPAROLÊ, *Pomatum*, *Pommade*.

LIPAROTES (*λιπαροτης*), see *Polysarcia*.

LIPAROTRICH'IA, (*liparus*, and *τριχος*, gen. 'hair.') Too great oiliness of the hair.

LIPARUS (*λιπαρος*), Fatty.

LIPASMA (*λιπασμα*), see *Polysarcia*.

LIPE (*λίπε*), Fat.

LIPHÆMIA, *Leiphæmia*.

LIPHÆMOS, *Leiphæmos*.

LIPOCELE, (*lipe*, and *κηλη*, 'tumour.') *Liparoc'ele*.

LIPODERMIA, (*leipo*, and *δερμα*, 'skin.') *Aposthia*.

LIPODERMUS, *Leipodermus*.

LIPO'MA, gen. *Lipo'matis*, (F.) *Lipôme*, (from *lipe*, and *oma*.) A fatty tumour of an encysted or other character.

LIPOMATODES, (*lipoma*, and *odes*.) *Lipomatous*.

LIPOM'ATOUS, *Lipomato'sus*, *Lipomato'des*, (F.) *Lipomatœuz*. Having the nature of lipoma—as a '*lipomatous* mass.'

LIPÔME, *Lipoma*.

LIPOM'E'RIA, *Leipom'e'ria*, (*leipo*, and *μερος*, a 'part.') Monstrosity from arrest of development, or from defect. Deficiency of one or more of the parts of the body; for example, in one who has only four fingers.

LIPOPSYCHIA, (*λειποψυχια*, from *leipo*, and *ψυχη*, 'breath.') *Syncope*.

LIPOSIS, (from *lipus*, and *osis*.) see *Polysarcia*.

LIOSPHYXIA, (*leipo*, and *σφυξις*, 'pulse.') *Syncope*.

LIPOTHY'MIA, (*λειποθυμια*, from *leipo*, and *θυμος*, 'mind.') *Syncope*.

LIPPA, *Chassie*.

LIPPER, *Lepra*, *Leprous*.

LIPP'IA, *Adali*.

LIPPITU'DO, gen. *Lippitu'dinis*, *Ophthal'mia chron'ica* of some, *Lip'pitude*, *Bleat'eye*, *Bleat-eyedness*, *Bleat'edness*, *Lap'pa*, *Lip'pa*, *Xerophthal'mia*, *Le'ma*, *Le'mê*, *Lemos'itas*, *Gle'mê*, *Gla'ma*, *Gra'mia*. A copious secretion of the sebaceous humour of the eyelids which renders them gummy. It is owing to a state of chronic inflammation of the tarsal margins; the eyelids being generally red, tumefied, and painful.

LIPPITUDO NEONATORUM, see *Ophthalmia*.

LIPPUS, one affected with bleat eye; *bleat-eyed*; (F.) *Chassieus*.

LIPSIS (*λειψις*) *ANIMI*, ('a forsaking of mind.') *Syncope*.

LIPOTRICH'IA, (*λειψις*, 'a forsaking,' and *τριχ*, gen. *τριχος*, 'hair.') *Alopecia*, *Calvities*.

LIPUS (*λίπος*), Fat.

LIQUAMEN, gen. *Liqua'minis*, ('a liquid mixture,' from *liquare*, 'to dissolve.') *Garum*.

LIQUAMU'MIA, *Adeps huma'nus*. Human fat.

LIQUARIUM, (*liquare*, *liquidum*, 'to melt or dissolve.') *Syrupus simplex*.

LIQUATIO, *Liquefaction*.

LIQUEFA'CIENT, *Liquefa'ciens*, (*liquere*, 'to be liquid,' and *facere*, 'to make.') A medicinal agent, which seems to have the power of liquefying solid depositions. To this class mercury, iodine, &c., have been referred by some.

LIQUEFA'C'TION, *Liqua'tio*, *Liquefac'tio*. Transformation of a solid substance into a liquid. It is used particularly in speaking of metals and fatty bodies, see *Fusion*.

LIQUEUR, *Liquor*—l. *d'Alumine Composee*, *Liquor aluminis compos.*—l. *de l'Amnios*, *Liquor amnii*—l. *Anodine d'Hoffman*, *Spiritus ætheris sulphurici compositus*—l. *Anodine nitreus*, *Spiritus ætheris nitrici*—l. *Arsénicale*, *Liquor arsenicalis*—l. *Fumante de Boyle*, *Ammonia sulphuretum*, *Liquor fumans Boylii*—l. *de Lampadius*, *Carbonis sulphuretum*—l. *volatile de Corne de cerf*, *Liquor volatilis cornu cervi*—l. *de Cuivre ammoniacal*, *Liquor cupri ammoniaci*—l. *de Fer alcaline*, *Liquor ferri alkalini*—l. *de Labarraque*, *Liquor sodæ chlorinata*—l. *de Sous-acétate de plomb*, *Liquor plumbi subacetatis*—l. *de Potasse*, *Liquor potassæ*—l. *du Sang*, *Liquor sanguinis*—l. *de Sous-carbonate de potasse*, *Liquor potassæ subcarbonatis*.

LIQUID, ADHESIVE, MAYNARD'S. Collo-dion—*l.* Disinfecting, Burnett's, see Burnett's disinfecting liquid—*l.* Disinfecting, Labarraque's, Liquor sodæ chlorinatæ—*l.* Disinfecting, of Larnaude's, see Zinci sulphas—*l.* Disinfecting, Ledeyen's, see Ledeyen's disinfecting liquid.

LIQUIDAMBAR ASPLENIFOLIA. *Comptonia asplenifolia*—*l.* Officialis, see *Syrax*—*l.* *Peregri-na*, *Comptonia asplenifolia*.

LIQUIDAMBAR SYRACIFLUA. *Liquidambar*, *Sweet gum*, *White gum*. *Ord.* Balsamineæ. The name of the tree which affords the *Liq'id am'ber*, and *Sto'raz liq'uida*, *Syr'raz liq'uida* or *Liq'id sto'raz*. Liquid amber is a resinous juice of a yellow colour, inclining to red; at first of about the consistence of turpentine; by age, hardening into a solid, brittle mass. It is not used medicinally.

Syr'raz Liq'uida is obtained from this plant by boiling. There are two sorts,—one more pure than the other. It is used occasionally, as a stomachic, in the form of plaster.

LIQUIDE AMNIOTIQUE. *Liquor amnii*—*l.* *Céphalo-rachidienne*, *Cephalo-spinal fluid*.

LIQUIDUM NERVEUM. *Nervous fluid*.

LIQUIRITIA. *Glycyrrhiza*.

LIQUOR. *gen.* *Liquor'is*, (from *liqueo*, 'to be liquid,') *Liqua'men*, *Hy'gron*, *Hy'grotes*, (F.) *Liqueur*. A name given to many compound liquids, and especially to those the bases of which are water and alcohol.

LIQUOR ACIDUS HALLERI. *Elixir acidum Halleri*—*l.* *Ethereus*, *Ether*—*l.* *Ethereus sulphuricus*, *Spiritus ætheris sulphurici*.

LIQUOR ALUMINIS COMPOSITUS. *A'qua alu'mi-nis compos'ita seu alumin'o'sa Bate'na*, *Bate's alu'm wa'ter*, *Com'pound solution of Al'um*, (F.) *Liqueur d'alumine composée*. (*Aluminis*, *Zinci sulphat*, *sing.* $\mathfrak{z}$ ss, *Aque ferrent*, *Oij.* Dissolve and filter through paper.) Detergent and stimulant. Used as a collyrium, when properly diluted, in ophthalmia; as an injection in gleet, leucorrhœa, &c.

LIQUOR EX AMMONIÆ ET OLEO SUCCINI. *Spiritus ammoniæ succinatus*.

LIQUOR AMMONIÆ. *L. ammoniæ pu'ra*, *Al'cali ammoniacum flu'idum*, *Ammo'nia liq'uida seu can'stica liq'uida seu pu'ra liq'uida*, *Lixiv'ium ammoniacale*, *Spiritus sal'is ammoni'aci aqu'o'sus*, *A'qua ammo'nia seu ammo'nia can'stica*, *Liq'id ammo'nia*, *Solutio of ammo'nia*, (F.) *Ammoni-aque liq'uide*. Directed to be made in the Ph. U. S. from *Muriate of Ammonia*, in fine powder; *Lime*, and *Water*. Heat is applied so as to drive off the ammonia, which is made to pass into the distilled water. *Liquor Ammonia* may also be made by mixing one part of liquor ammonia fortior with two parts of distilled water. Its s. g. is 0.860. **LIQUOR AMMONIÆ FORTIOR.** *Stronger solution of ammonia*, *Aqua Ammonia fortior*, *Stronger water of ammonia*, (Ph. U. S. 1864.) is an aqueous solution of ammonia of the s. g. 0.900; that of Ph. B. 1864, is 0.891. *Liquor ammonia* is stimulant, antacid, and rubefacient. Dose, gr. x to xx, in water or milk.

LIQUOR AMMONIÆ ACETATIS. *Al'cali ammoni'acum aceta'tum*, *A. volat'ile aceta'tum*, *A'qua aceta'tis ammoniæ seu ammo'niæ aceta'tis*, *Solutio of ac'etate of ammo'nia*, *Ac'etus ammo'niæ*, *Sal ammoniacum vegetabile*, *Spiritus ophthal'micus seu Spiritus seu Liquor Mindere'ri*, *Sal aceto's ammoniacalis*, *Ammonium aceti'cum liq'u'idum*, *Spirit of Mindere'ri*, (*Acid. acetic. dilut.* *Oij.* *Ammon. carbonat.* q. s. Add the salt to the acid until it is saturated. Ph. U. S. 1851.) A sudorific; externally, cooling. Dose, f $\mathfrak{z}$ j to f $\mathfrak{z}$ iss.

LIQUOR AMMONIÆ CITRATIS. *Solution of Cit'rate of Ammo'nia*. This preparation is official in the Dublin Pharmacopœia. It is made by saturating a solution of citric acid with sesqui-

carbonate of ammonia. It is given in cases of adynamic fever.

LIQUOR AMMONIÆ SESQUICARBONATIS. *Liquor Ammonia Subcarbonatis*.

LIQUOR AMMONIÆ SUBCARBONATIS. *Liquor Ammonia Sesquicarbonatis*, *Solutio subcarbonatis ammoniæ*, *A'qua carbonatis ammoniæ*, *Solutio of subcarbonate of ammo'nia*, *Wa'ter of carbonate of ammo'nia*. (*Ammonia carbon.* $\mathfrak{z}$ iv, *Aque destillat.* *Oj.* Dissolve and filter through paper.) Use;—the same as that of the carbonate of ammonia.

LIQUOR AMMONIÆ ETHEREUS. *Spiritus ammonia aromaticus*—*l.* *Ammonii aromaticus*, *Spiritus ammonia aromaticus*—*l.* *Ammonii hydrothiodis*, *Ammonia sulphuretum*—*l.* *Ammonii oleoso-aromaticus*, *Spiritus Ammonia aromaticus*—*l.* *Ammonii succinei*, *Spiritus ammonia succinatus*—*l.* *Ammonii vinosus*, *Spiritus ammonia*.

LIQUOR AM'NI. *A'qua am'ni*. The liquor of the amnios, (F.) *Eau ou Liqueur de l'amnios*, *Liquide amniotique*. The fluid exhaled by the amnios, and which envelops the fœtus during the whole period of uterogestation. It is often simply called the *waters*, (F.) *Les Eaux*. Its relative quantity diminishes as pregnancy advances, although its absolute quantity continues to increase till the period of delivery. In some women, only five or six ounces are met with; in others, it amounts to pints. It is limpid, yellowish, or whitish; exhales a faint smell, and has a slightly saline taste. It contains water in considerable quantity; albumen; chloride of sodium; phosphate of lime; and alkaline substance; and a particular acid. It facilitates the dilatation of the uterus, and aids delivery by acting as a soft wedge enclosed in its membranes. *Bag of waters*, *Poche des Eaux*, &c. It is probably inservient to useful purposes in the nutrition of the fœtus.

LIQUOR AMNI, FALSE. The fluid contained between the amnion and chorion in the early periods of fetal existence.

LIQUOR ANODYNE, HOFFMANN'S. *Spiritus ætheris sulphurici compositus*—*l.* *Anodynus martialis*, *Alcohol sulphuricoæthereus ferri*.

LIQUOR ANODYNUS TEREBINTHINATUS. A formula prescribed by Rademacher in cases of gallstone, and of obstructions and indurations of the liver and spleen. It was composed of *Hoffmann's anodyne liquor* $\mathfrak{z}$ j; *Rectified oil of turpentine* $\mathfrak{z}$ ij. Dose, 5 to 10 drops. It resembles the *Remède de Durand*, *Mixture lithonatriptique de Durand*, *M. de Whitt*, *Éthéré d'essence de térébenthine*, which is formed of *Sulphuric ether*, 30 parts; *Essence of turpentine*, 15 parts. It is used in the same cases.

LIQUOR ANTIMONII TARTARIZATI. *L. Tartari emetici*, *Solutio of Tartarized Antimony*, (*Antim. et potass. tartrat.* gr. liiv, *Aq. destillat.* *Oj.* Solve et adde *Sp. Vini rectif.* f $\mathfrak{z}$ vij.) Emetic, like the *Vinum antimonii*; and in the same dose.

LIQUOR ANTIMONII TERCHLORIDI. see *Antimonium muriatum*.

LIQUOR ARSENICALIS. *L. Potas'se arseni'tris* (Ph. U. S.), *Solutio arsenicalis seu arsenicata seu arsenitis kali'æ*, *Arsenical solution*, *Min'eral sol'vent*, *Ar'senic potas'se liq'u'idus seu potas'se aqu'o'sus*, *Fowler's solution of arsenic*, *Solvens mineral'is*, *Italian poison*, *Aqua seu Aqua Toffania*, *Acqua della Toffana*, *A. di Napoli*, *Acquet'a* [?], *Tasteless aque drops*, (F.) *Liquor arsenicalis*. (*Acid. Arsenios*, in frustulis, *Potas'se carbonatis pur.*, *sing.* gr. lxiv, *Aque destillat.* q. s. Boil together the arsenious acid and carbonate of potassa with twelve fluidounces of distilled water, in a glass vessel, until the arsenic is dissolved. When the solution is cold, add *Spirit. lavand.* c. f $\mathfrak{z}$ iv, and as much distilled water as will make the whole one pint. Ph. U. S. 1851 contains gr. ss of the arsenious acid. Dose, gr. x.

LIQUOR ARSENICALIS BIETII, see Arseniate of ammonia.—I. Arsenicalis Pearsonii, see Arseniate of soda.

LIQUOR ARSENICI CHLO'RIDI, *Solution of Chloride of Arsenic*, of the London Pharmacopœia of 1851, is made by boiling *Arsenious acid* $\mathfrak{Zss}$, in *Hydrochloric acid* $\mathfrak{f}\mathfrak{zss}$ (Imp. meas.), mixed with a fluidounce of distilled water, until it is dissolved: then adding water to make the quantity a pint (Imp. meas.) Each fluidounce contains a grain and a half of arsenious acid, or an equivalent quantity of terchloride. Properties the same as those of Liquor Arsenicalis. It is a substitute for, and imitation of, *De Valangin's Arsenical Solution*, *Solutio solven'tis minera'alis*, much used by many British practitioners. Dose, grt. v to x.

LIQUOR ARSENICI ET HYDRARGYRI IODIDI, see Arsenic and Mercury, iodide of.

LIQUOR ATROP'LE, *Solution of Atropia*, (*Atropia*, in crystals gr. iv, *Rectified Spirit* $\mathfrak{f}\mathfrak{z}\mathfrak{j}$; *Distilled water*, $\mathfrak{f}\mathfrak{z}\mathfrak{vi}\mathfrak{j}$. Dissolve the atropia in the mixed fluid, Ph. B. 1864.) Used by the oculist to dilate the pupil, a drop being placed on the eye.

LIQUOR BARI CHLORIDI, Baryta, muriate of, solution of.—I. Bellostii, L. Hydrargyri nitrici.—I. Calcei chloridi, see Calceis murias.

LIQUOR CAL'cis, *Solutio cal'cis*, *Aqua cal'cis seu benedicta seu Calca'rie us'tæ*, *Calca'ria pu'ra liquida*, *Solution of Lime*, *Lime-water*, (F.) *Eau de chaux*. (Calceis $\mathfrak{Z}\mathfrak{iv}$, Aq. destill. cong. Pour the water on the lime, and stir. Let it stand in a covered vessel three hours; bottle the lime and water in stopped bottles, and use the clear solution.) It is astringent, tonic, and antacid; and is used in diarrhœa, diabetes, heartburn, &c., and as a lotion to foul and cancerous ulcers, &c. Dose, $\mathfrak{f}\mathfrak{z}\mathfrak{i}\mathfrak{j}$ to Oss, in milk.

The British Pharmacopœia has a *Liquor Cal'cis saccharatus*, *Saccharated Solution of Lime*, which is made by mixing *slaked lime* $\mathfrak{Z}\mathfrak{j}$, with refined sugar, in powder, $\mathfrak{z}\mathfrak{i}\mathfrak{j}$; putting them in a bottle containing a pint of distilled water; shaking occasionally for a few hours and separating the clear solution. Dose, $\mathfrak{f}\mathfrak{zss}$ to $\mathfrak{f}\mathfrak{z}\mathfrak{i}$.

LIQUOR CALCEIS CHLORINATÆ, L. C. Chlorinatæ, *Solution of Chlorinated Lime*, is made by triturating half a pound of chlorinated lime with half a gallon of water; shaking the mixture; filtering, and keeping in a well-stopped bottle. Used in the same cases as chlorinated lime. Dose, $\mathfrak{f}\mathfrak{zss}$ to $\mathfrak{f}\mathfrak{z}\mathfrak{j}$.

LIQUOR CALCEIS COMPOSITUS. *Aqua Cal'cis composita seu benedicta composita*, *Compound Lime-water*. (F.) *Eau de chaux composé*. (Lign. quaiac. ras. lss. Rad. glycyrrh. $\mathfrak{z}\mathfrak{j}$, Cort. cassiafras, $\mathfrak{zss}$; Semin. coriand. $\mathfrak{z}\mathfrak{i}\mathfrak{j}$. *Liquor. calceis*, Ovj. Macerate for two days, and filter.) It is stimulant, diaphoretic, and astringent, and is used in cutaneous affections.

LIQUOR CALCEIS MURIATIS, see Calceis murias.—I. Cerebrospinalis, Cephalospinal fluid.—I. Cerebris, Cerevisia.—I. Chloreti natri, L. sodæ chlorinatæ.—I. Chlorini, see Chlorine.—I. Chlorureti natri, L. Sodæ chlorinatæ.—I. Chlorureti sodæ, L. sodæ chlorinatæ.—I. Chyli, see chyle.—I. Cornu cervi, *Liquor Volatilis cornu cervi*—I. of Cotugno or Cotunnus, Cotunnus, *Liquor of*.

LIQUOR CUPRI AMMONIAT'II, *Liquor Cupri ammoniosulphatis*, *Aqua seu solutio cupri ammoniati seu sappharina*, *Blue eye-water*, *Solution of ammoniated copper*. (F.) *Liquor on Eau de cuivre ammoniacal*. (Cupri ammoniat. $\mathfrak{z}\mathfrak{i}$. *Aque destill.* Oj. Dissolve and filter the solution through paper. Ph. L.) Corrosive and detergent. Used externally to foul ulcers; and diluted with an equal part of distilled water, it is applied by means of a hair pencil to specks and films on the eye.

LIQUOR CUPRI AMMONIOSULPHATIS, *Liquor Cupri ammoniati*.

LIQUOR CUPRI SULPHATIS COMPOSITUS, *Aqua*

cupri vitriola'ti composita. (Cupri sulphat., Alumin. sulphat. $\mathfrak{aa}$ $\mathfrak{z}\mathfrak{i}\mathfrak{j}$. *Aque puræ* Oij., *Acid. sulph.* $\mathfrak{z}\mathfrak{i}\mathfrak{j}$. Boil the salts in the water until they are dissolved; then filter the liquor, and add the acid.) Used as an astringent in epistaxis, &c. It was also called *Aqua Styptica*.

LIQUOR CYRENIACUS, Benjamin.—I. Entericus, Succus entericus.—I. Donovanii, see Arsenic and Mercury, iodide of.—I. Excitans, Spiritus ammoniaci succinatus.

LIQUOR FERRI ALKALI'NI, *Solution of Alkaline Iron*, (F.) *Liquor de fer alkaline*. (Ferri $\mathfrak{z}\mathfrak{i}\mathfrak{ss}$, *Acid. nitricæ*. $\mathfrak{z}\mathfrak{i}\mathfrak{j}$, *Aque destillat.* $\mathfrak{f}\mathfrak{z}\mathfrak{vi}\mathfrak{j}$, *Liq. potass. subcarb.* $\mathfrak{f}\mathfrak{z}\mathfrak{v}\mathfrak{j}$. To the acid and water mixed, add the iron; and, after the effervescence, add the clear solution, gradually, to the liq. potassæ subcarb.; shaking it occasionally till it assumes a deep browned colour, and the effervescence stops. After six hours' settling, pour off the clear solution. Ph. L.) It is tonic like other preparations of iron. Dose, $\mathfrak{f}\mathfrak{zss}$ to $\mathfrak{f}\mathfrak{z}\mathfrak{i}\mathfrak{ss}$.

LIQUOR FERRI CITRATIS, *Solution of Citrate of Iron*, (Ph. U. S. 1863,) (made by saturating with citric acid the hydrated oxide of iron formed by decomposing the solution of tersulphate of iron by water of ammonia; and reducing the liquid by evaporation to the proper measure.) Used to prepare the Ferri citras.

LIQUOR FERRI IODIDI, *Solution of Iodide of Iron*, *Syrupus Ferri iodidi*, (Ph. U. S. 1863,) *Syrup of Iodide of Iron*. (Iodin. $\mathfrak{z}\mathfrak{i}\mathfrak{j}$, Ferri rament. $\mathfrak{z}\mathfrak{i}$, Sacchar. pulv. $\mathfrak{z}\mathfrak{x}\mathfrak{i}\mathfrak{j}$, *Aque destillat.* q. s. Mix the iodine with $\mathfrak{f}\mathfrak{z}\mathfrak{x}$ of the distilled water, in a porcelain or glass vessel, and gradually add the iron filings, constantly stirring. Heat the mixture gently until the liquor acquires a light greenish colour; then, having added the sugar, continue the heat a short time, and filter. Lastly, pour distilled water upon the filter, and allow it to pass until the whole of the filtered liquor measures twenty fluidounces. Keep the solution in closely stopped bottles.—Ph. U. S. 1851.) Dose, 20 or 30 drops.

LIQUOR FERRI MURIATIS, *Tinctura ferri muriatis*—I. Ferri muriatæ oxydati, see Ferri chloridum.

LIQUOR FERRI NITRATIS, L. Ferri Sesquinitrat'is seu ternitrat'is seu pernitrat'is, *Solutio Ferri nitrat'is*. *Solution of nitrate, of ternitrate of sesquioxide, or of sesquinitrate, of iron*, has been recommended in chronic diarrhœa and dysentery. Its virtues exactly resemble those of chloride of iron. It is prepared as follows:—Ferri fili incis. $\mathfrak{z}\mathfrak{i}\mathfrak{ss}$, *Acid. nitricæ*. $\mathfrak{f}\mathfrak{z}\mathfrak{v}$, *Aq. destillat.* q. s. Mix the acid and a pint of distilled water, until the gas ceases to be given off; filter, and add distilled water to make $\mathfrak{f}\mathfrak{z}\mathfrak{xxxvi}$ —Ph. U. S.) Dose, 10 to 20 drops.

LIQUOR FERRI PERCHLORIDI, see Ferri chloridium.—I. Ferri sesquichlorati, see Ferri chloridum.—I. Ferri subsulphatis, see Ferri sulphas.

LIQUOR FERRI TERsulphATIS, *Solution of Tersulphate of Iron*. (Ferri sulph. in pulv. crass. $\mathfrak{z}\mathfrak{x}\mathfrak{i}\mathfrak{j}$; *Acid. sulphur.* $\mathfrak{z}\mathfrak{i}\mathfrak{j}$ and gr. lx; *Acid. nitricæ*. $\mathfrak{z}\mathfrak{i}\mathfrak{j}$ and gr. cccx; *Aq. q. s.* Mix the acid with half a pint of the water; heat the mixture to the boiling point, and add the sulphate of iron gradually until effervescence ceases. Continue the heat until the solution has a reddishbrown colour, and is free from nitrous odour. Add water, when nearly cold, to make a pint and a half. Ph. U. S., 1863.) Used in the formation of different chalybeate preparations.

LIQUOR FOLLICULI, see Folliculi Graafiani.

LIQUOR FUMANS BOYL'II, *Sulphuretum Ammoniaci Hydrogenatum*, *Hydrosulphuretum Ammoniaci*, *Tinctura Sulphuris Volatilis*, *Aqua Sulphureti Ammoniaci*, Boyle's Fulminating liquor, (F.) *Liquor fumante de Boyle*. It is possessed of nauseating and emetic properties, and has been

given in diabetes and diseases of excitement as a deoxygenuiser!

LIQUOR FUMING, BOYLE'S, Liquor fumans Boylei—l. Fuming, of Libavius. Tin, muriate of—l. Genital, Speru—l. of Hartshorn, volatile, Liquor volatilis cornu cervi—l. Hydrargyri bichloridi, Liquor hydrargyri oxy muriatis—l. Hydrargyri chloridi corrosivi, l. hydrargyri oxy muriatis—l. Hydrargyri et Arsenici iodidi, see Arsenic and Mercury, iodide of.

LIQUOR GUTTA-PER'CHÆ, *Solution of Gutta-per'cha*, (Ph. U. S. 1863), *Traumaticine*. (A solution of gutta-percha in chloroform in the proportion of 3iiss of the former to 3xvij of the latter.) Applied by means of a camel's-hair pencil in cutaneous affections, as erysipelas and eczema, and to indolent ulcers.

LIQUOR HYDRARGYRI NITRATIS ACIDUS, see Hydrargyri Nitrates.

LIQUOR HYDRARGYRI NI'TRICTI, *L. Bellost'ii*, (F.) *Eau mercurielle ou contre la gangrène, Remède du Duc d'Antin, R. du Capucin*, (Hydrarg. 120 p., Acid. nitr. (33°) 150 p. Dissolve, and add to the solution distilled water, 900 p. Ph. P.) Dose, two or three drops in a glass of water. Not much used.

LIQUOR HYDRARGYRI OXYMURIA'TIS, *L. Hydrargyri Chlo'ridi corrosivi seu Hydrargyri bichlo'ridi, Solution of Oxy muriate of Mercury, L'iquor Suietensis seu Syphiliticus Turneri, Mu'rias hydrargyri spirituosus liq'u'idus, Solutio Muria'tis hydrargyri oxygena'ti, Liquor or Solution of Corro'sive Sublimate or of Van Swiet'en*, (Hyd. oxyg., Ammon. muriat. 55 gr. x. Aquæ destill. Oj., (Imp. meas.) Dissolve in the water and add the spirit. Ph. P.) Dose, f3ss, or f3i.

Nor'ton's Drops, Green's Drops, and Solomon's Antimptig'ones—all nostrums—seen, to be disguised solutions of Corrosive Sublimate.

LIQUOR HYDRARGYRI PERNITRATIS, see Hydrargyri nitrates—l. Hydrargyri Supernitratis, see Hydrargyri nitrates—l. Hydriodatis Arsenici et Hydrargyri, Arsenic and mercury, iodide of—l. Intestinalis, Succus entericus.

LIQUOR IO'DINI COMPOS'ITUS, (Ph. U. S. 1842,) *L. Iodin'ii compos'itus*, (Ph. U. S. 1851.) *Solutio Potas'sii Iodidi Iodure'ta, Compound Solution of Potass. Iodide, Lugol's Solution*, (Iodin. 3vj, Potass. iodid. 3iiss, Aquæ destillat. Oj. Dissolve the iodine and iodide of potassium in the water.—Ph. U. S.) Dose, gtt. vj, ad xij, in sugared water.

LIQUOR LACTIS, see Milk—l. Lithargyri Subacetatis, L. plumbi subacetatis—l. Lithargyri subacetatis compositus, Liquor plumbi subacetatis dilutus—l. Lymphæ, see Lymph.

LIQUOR MAGNESIÆ CITRATIS, see Magnesiae citras—l. Mindereri, Liquor ammoniæ acetatis.

LIQUOR MONRO', *Solution of Monro'*. A solution, used by Monro for the preservation of anatomical preparations. It was composed of alcohol at 22° or 24°, with a drachm of nitric acid to each pint.

LIQUOR MORGAGNII, Morgagni, humour of.

LIQUOR MORPHIÆ ACETATIS, *Solution of Acetate of Morph'ia, Anodyne Drops*, (Acetate of morph'ia, 3iv, Distilled water, Oj. (Imp. meas.), Spirit. tenuior. Oss, (Imp. meas.) Ph. L. Acid. acetic. ℥xv. Dose, from six to twenty-four drops.

LIQUOR MORPHIÆ MURIA'TIS, *Solution of Muriate of Morph'ia*, (Morphiæ muriat. 3iv, Aq. destillat. Oj. (Imp. meas.), Sp. tenuior. Oss, (Imp. meas.) Ph. L.) Dose, from 6 to 20 minims.

LIQUOR MORPHIÆ HYDROCHLORA'TIS, *Solution of Hydrochlo'rate of Morph'ia*, Ph. B. 1864, consists of hydrochlorate of morph'ia, gr. iv. dilute hydrochloric acid ℥viij, rectified spirit f3ij, distilled water, f3vj. Dose, ℞x to f3ss.

LIQUOR MORPHIÆ SULPHA'TIS, *Solution of Sulphate of Morph'ia*, (Morphiæ sul'phat. gr. viij, Aquæ destillat. Oss. Dissolve the sulphate of morphia in the water.—Ph. U. S.) Dose, f3i to f3ij—containing from an eighth to a quarter of a grain.

LIQUOR MUCI, see Mucus—l. Natri Oxymuriat'ici, L. Sodæ chlorinatæ—l. Nervinus Bangii, Tinctura ætherea camphorata—l. Oleosus Sylvi, Spiritus ammoniæ aromaticus—l. Opii sedativus, (Haden's), see Tinctura opii.

LIQUOR OPII SEDATI'VUS. An empirical preparation by a London druggist of the name of Battley. It is said to be an aqueous solution of opium, evaporated to dryness to get rid of the acid resin, redissolved in water, and a small portion of alcohol added to give it permanence. (Redwood.) It is devoid of many of the narcotic effects of opium.

LIQUOR OVI ALBUS, Albumen ovi—l. Pancreaticus, see Pancreas—l. Pericardii, see Pericardium—l. Plumbi acetatis, L. P. subacetatis—l. Plumbi diacetatis, Liquor Plumbi subacetatis.

LIQUOR PLUMBI SUBACETA'TIS, *L. Subacetatis Lithar'gyri seu Plum'bi Aceta'tis seu Plum'bi Diaceta'tis, Goulard's Extractum Satur'ni, Lithar'gyri Acetum, Acetum plum'bicum seu satur'num, Solution of Subac'etate of Lead, Goulard's extract, Goulard's extract of lead, Goulard*, (F.) *Liquor de sous-acétate de Plomb*, (Plumb. acet. 3xvj, Plumb. oxid. semivit. in pulv. subtil. 3ixss, Aq. destillat. Oiv. Boil together in a glass or porcelain vessel for half an hour, occasionally adding distilled water, so as to preserve the measure. Filter through paper, and keep the solution in closely-stopped bottles.—Ph. U. S.) It is used externally, as a cooling astringent, and dis-cutient, when diluted with distilled water.

LIQUOR PLUMBI SUBACETA'TIS DILU'TUS, *L. Subacetatis Lithar'gyri Compos'itus, L. Plum'bi aceta'tis dilu'tus, L. Plumbi Diaceta'tis dilu'tus, Acetas Plumbi dilu'tum alcohol'icum, Diluted Solution of Subac'etate of Lead, A'qua Satur'ni seu veg'etomineral'is seu Plum'bi seu Goulard'ii seu Lithar'gyri Aceta'ti compos'ita, Tinctura plumb'osa, Lead water, Goulard' water, Goulard's lotion, Goulard*, (F.) *Eau blanche, Eau de Goulard, Eau végété-minérale, White Wash, Royal Preventive*, (Liq. plumbi subacetat. 3ij. Aquæ destillat. Oj. Ph. U. S.) Properties the same as the last, but feebler.

LIQUOR POTAS'SÆ, *A'qua Potas'sæ seu Kali Caust'ici seu Kali pur'i, Solution of Potash or of Potassa, Lixiv'ium magistra'te seu Saponar'ium, Soap Lees, Soap Ley, Lixiv'ium caust'icum, Potas'sa liq'u'ida*, (F.) *Eau, solution ou liqueur de Potasse, Potasse liquide, Lessive des Savonniers*, (Potassæ carb. ℔j, calcei. ℔ss, Aquæ destill. fervent. congium. Dissolve the alkali in Oij of the water, and add the remainder to the lime. Mix the whole: set aside in a close vessel, and, when cold, filter through calico. It may, also, be obtained by dissolving potassa, 3j, in distilled water, Oj.) It is antilithic in cases of uric acid calculi, and antacid. Externally, stimulant and escharotic. Dose, gtt. x to xx.

LIQUOR POTASSÆ ARSENITIS, L. arsenicalis—l. Potassæ Carbonatis, L. P. Subcarbonatis.

LIQUOR POTASSÆ CITRA'TIS, *Mistura Potas'sæ Citra'tis, Mixture of Citrate of Potassa, Solution of Citrate of Potas'sa, Neutral Mix'ture, Saline Mix'ture*, (Succ. Limon. Oss, Potass. Bicarbonat. q. s.) saturate and filter. This is the *Mistura Potassæ citra'tis*, Ph. U. S. 1863. Or, take Acid. Citric. 3ss; Ol. Limon. M. ij; Aquæ Oss. Potassæ Bicarbonat. q. s.; dissolve, saturate by the carbonate of potassa, and filter. This is the *Liquor Potassæ citra'tis*, Ph. U. S. 1863. Used in fever, but probably of little or no efficacy.

LIQUOR POTASSÆ PERMANGANATIS, see Potassæ permanganas.

LIQUOR POTASSÆ SUBCARBONATIS, *L. Potassæ Carbonatis* (Ph. U. S.), *A'qua Kal'i seu Kal'i preparati seu Subcarbonatis Kal'i*, *O'leum Tartari per deliquium*, *Liriv'ium Tartari*, Oil of Tartar, *Saline oil of Tartar*, *Solution of Subcarbonate of Potassæ*, (F.) *Liqueur de sous-carbonate de Potasse*, *Lessive de Tartre*, *Huile de Tartre par défaillance*. (Potass. subcarb. ℞j, *Aque destillat.* Oj. Dissolve and filter. Ph. U. S.) Dose, grt. x to xxx.

LIQUOR POTASSII IO'DIDI COMPOSITUS. *Compound Solution of Iodide of Potassium*. (Potassii iodid. gr. x, Iodin. gr. v, *Aque destillat.* Oj. (Imp. meas.) Ph. L.) Dose, ℥j.

LIQUOR PROPAGATORY, *Sperm—*1. Prostaticus, Prostatis liquor, see Prostate—1. Puris, see Pus.

LIQUOR SANG'UINIS. *Blood liquor*, (F.) *Liquen du Sang*. A term given by Dr. B. Babington to one of the constituents of the blood, the other being the red particles. It is the *Hæmatoplasma*, the *Plasma*, of Schultz, *Lymph*, *Coagulable* or *plastic Lymph*, the *Mucin* or *Mucilage* of Harvey, and others, and the *intercellular fluid* of Lehmann, (F.) *Lymphy plastique ou coagulable*, *Suc nourricier*, and is the effused material from which the cells obtain the constituents of the different tissues and secretions. See Blood.

LIQUOR OF SCARPA, *Vitrine auditive*.

LIQUOR SEMINIS, *Spermatic liquor*. The homogeneous transparent fluid, in which the spermatozooids and seminal granules are contained. See Sperm.

LIQUOR SODÆ, *L. Sodæ Caustica*, *Solution of Soda*, *Solution of Cautic Soda*. (Prepared, like Liquor Potassæ, from Carbonate of Soda.) It has the same properties as Liquor Potassæ.

LIQUOR SODÆ ARSENIATIS, see Arseniate of Soda.

LIQUOR SODÆ CARBONATIS, *Solution of Carbonate of Soda*. (*Sodæ carbon.* ℥iss, (avoidrupois); *Aque destillat.* Oj. (Imp. meas.) Ph. D.) Dose, as an antacid, one to two tablespoonfuls, diluted with water.

LIQUOR SODÆ CHLORIDI, *L. sodæ chlorinatæ*.

LIQUOR SODÆ CHLORINATÆ, *L. Sodæ chloratæ seu sodæ chloridi seu sodæ oxymuriaticæ seu chloreti nat'ri seu chlorureti nat'ri seu chloreti sodæ seu chlorureti sodæ seu nat'ri oxymuriat'ici, Nat'rium chloratum liquidum*, *A'qua nat'ri oxymuriat'ici*, *Labarraque's Disinfecting Liquid*, *Solution of Chlorinated Soda*, (F.) *Liquen de Labarraque*. (*Calcis Chlorinat.* ℞j, *Sodæ Carbonat.* ℞ij; *Aque cong. iss.* Dissolve the carbonate of soda in three parts of the water, with the aid of heat. To the remainder of the water add, by small portions at a time, the chlorinated lime, previously well triturated, stirring the mixture after each addition. Set the mixture by for several hours, that the dregs may subside; decant the clear liquid, and mix it with the solution of carbonate of soda. Lastly, decant the clear liquor from the precipitated carbonate of lime, pass it through a linen cloth, and keep it in bottles secluded from the light.—Ph. U. S.) Used in the same cases as the chloride of lime. Internally, 10 drops to a fluidrachm, for a dose. Diluted with water, it is an excitant and disinfectant in various morbi externi.

LIQUOR SODÆ EFFERVESCENS, Acidulous water, simple—1. Sodæ Oxymuriaticæ, *L. sodæ chlorinatæ*—1. Strychniæ, see Strychniæ murias—1. Stypticus Loofii, see Ferri chloridum—1. Stypticus Ruspini, Stypticæ Ruspini's—1. Sulphurico-æthereus Constringens, Colloidum—1. Sulphuricus Alcolisatus, Spiritus ætheris sulphurici—1. Swietenii, *L. hydrargyri oxymuriatis*—1. Syphiliticus Turneri. *L. hydrargyri oxymuriatis*—1. of Van

Swieten, *L. hydrargyri oxymuriatis*—1. Tartari emetici, *L. antimonii tartarizati*—1. Vesicans, Colloidum, cantharidal.

LIQUOR VOLATILIS CORNU CERVI, *L. cornu cervi*, *L. volatilis Cornu Cervini*, *L. volatilis ossium*, *Volatile Liquor of Harts'horn*, *Spiritus Lumbricorum seu Millepedarum seu Cornu Cervi*, *Harts'horn*, *Spirit of Harts'horn*, *Bone Spirit*, (F.) *Liqueur volatile de Corne de cerf*. This is a solution of subcarbonate of ammonia, impregnated with empyreumatic oil. It possesses the same virtues as the subcarbonate of ammonia. It is in common use to smell at, in faintings, &c.

LIQUOR VOLATILIS OSSIUM, *L. volatilis cornu cervi*—1. Zincæ Chloridi, see Burnett's Disinfecting Liquid.

LIQUOR ZINCI SULPHATIS cum CAMPHORÆ, *A'qua Zinci vitriolata cum Camphorâ seu vitriolica camphorata seu ophthalmica*, *Compound Eye-water*. (*Zinci sulph.* ℥ss, *Camphor.* ℥ij, *Aq. bullient.* Oij; dissolve and filter.) Used as a lotion for ulcers; or, diluted with water, as a collyrium.

LIQUORICE, Glycyrrhiza—1. Bush, *Abrus precatorius*—1. Juice, see Glycyrrhiza.

LIQUORICE, REFINED, *Extractum glycyrrhizæ*—1. Spanish, see Glycyrrhiza—1. Wild, *Aralia nudicaulis*, *Galium circeæans*.

LIQUORITIA, Glycyrrhiza.

LIRIODENDRON, (*Λιριον*, 'lily,' and *δένδρον*, 'tree,') *L. Tulipifera*, *Tulipifera Liriodendron*, *Old wife's shirt*, *Tulip or Poplar tree*, *Tulip-bearing or American Poplar*, *White Wood*, *Cypress Tree*, (New England.) (F.) *Tulipier*. The bark—*Liriodendron* (Ph. U. S.)—especially of the root, of this noble forest tree, which is indigenous in the United States, is a strong aromatic bitter, and has been employed advantageously as a tonic. An active principle was separated from it by Professor J. P. Emmet of the University of Virginia, and has been called *Liriodendrin*. It is not used in medicine.

LIS ASPHODELE, ([F.] *lis*, 'a lily,') *Asphodelus ramosus*—1. *Blanc*, *Lilium candidum*—1. *des Étangs*, *Nymphæa alba*.

LISERON, GRAND, (from *lis*,) *Convolvulus sepium*—1. *des Haies*, *Convolvulus sepium*—1. *Mechameck*, *Convolvulus panduratus*.

LISK, (Se. and Prov.), Inguen.

LISTS, (Prov.) Flanks.

LISTON'S ISINGLASS PLASTER, see Sparadrapum adhesivum.

LITE, (*λίτη*), A plaster, formerly made of verdigris, wax, and resin.

LITHAGOGECTASIA, (*lithagogum*, and *εκτασία*, 'dilatation,') Lithectasy.

LITHAGO'GUM, (*lithos*, and *αγω*, 'I expel,') A remedy which was supposed to possess the power of expelling calculi. Also, a lithotomy forceps.

LITHANTHRAX, (*λίθο*, and *ανθράξ*, 'a coal,') *Carbo fossilis*.

LITHANTHRAKOKALI, *Anthrakokali*.

LITHARGE, (*λίθαργος*, from *λίθος*, and *αργός*, 'silver,') *Plumbi oxydum semivitreum*—1. of Gold, see *Plumbi oxydum semivitreum*—1. of Silver, see *Plumbi oxydum semivitreum*.

LITHARGYRI ACETUM, *Liquor plumbi subacetatis*.

LITHARGYRUM, *Plumbi oxydum semivitreum*.

LITHARGE, *Plumbi oxydum semivitreum*.

LITHAS, gen. *Litha'tis*, Urate.

LITHATE, Urate—1. of Soda, Urate of soda.

LITHECTASY, *Lithectasia*, *Lithagogectasia*, *Lithodectasia*, (*lithos*, and *εκτασία*, 'dilatation,') *Cystectasy*. An operation which consists in extracting stone from the bladder by dilating the neck of the organ, after making an incision in

the perineum, and opening the membranous portion of the urethra.

LITHIA, (from *lithos*), *Lithi'asis*, *Lithogen'ia*, *Uri'asis*, *Urolithi'asis*, *Cachex'ia calculo'sa*, *Cal'culi Mor'bus*, *Lopilla'tio*, *Genera'tio cal'culi*. The formation of stone, gravel, or concretions in the human body.

Also, an affection in which the eyelids are edged with small, hard, and stone-like concretions.

Lithia is also the oxide of *lith'ium*, *lith'ion* or *lithon*.

LITHIA CARBONATE OF, *Lith'ie Carbo'nas*, *Lith'ium Carbo'nicum*, (F.) *Carbonate de Lithine*. A salt found in certain mineral waters, which have been serviceable in lithuria by its power of dissolving uric acid and the urates. Hence, it has been suggested in that morbid condition. It may be obtained by decomposing chloride of *lithium* by means of carbonate of ammonia. Dose, gr. v to x, three or four times a day. It has, also, been injected into the bladder, in cases of deposition of lithuria and oxaluria.

Cit'rate of Lith'ia, *Lith'ie cit'ras* made by the action of citric acid on carbonate of *lithia* is official in the Ph. B. for 1864. It has been given in gout. Dose, gr. x to gr. xx.

LITHIA RENALIS ARENOSA, Gravel—l. Renalis, Nephrolithiasis—l. Vesicalis, Calculi, vesical.

LITHIÆ CARBONAS, *Lithia*, carbonate of—l. *Citras*, see *Lithiæ carbonas*.

LITHIASIS, (*Λιθιασις*), *Lithia*—l. *Cystica*, *Cal'culi*, vesical—l. *Nephritica*, Gravel, Nephrolithiasis.

LITHIASIS PULMONUM, *Pulmo'nes tartariza'ti*. The formation of concretions in the lungs, occasioning at times *Lithophthi'sis*, *Phthisis calculo'sa*, *Phthisis calculeuse*, of Bayle.

LITHIASIS RENALIS ARENOSA, Gravel—l. Renalis, Nephrolithiasis—l. Vesicalis, Calculi, vesical.

LITHIC, *Lith'icus*. Belonging to lithic or uric acid, or to stone: hence *Lithic Diathe'sis*. Also, an antilithic.

LITHIC ACID, Uric acid—l. Acid diathesis, Lithuria—l. Diathesis, Lithuria—l. Sediments, see Lithuria.

LITHINE CARBONATE DE, *Lithia*, carbonate of.

LITHION, see *Lithia*.

LITHIUM, see *Lithia*—l. Carbonicum, *Lithia*, carbonate of.

LITHIURIA, *Lithuria*.

LITHOBEXIS, (*lithos*, and βῆξ, 'a cough.') Cough with calculous expectoration.

LITHOCENOSIS, (*lithos*, and κενώσεις, 'evacuation.') Lithotrity.

LITHOCLAST, (*lithos*, and κλω, 'to break,') see Lithotrity.

LITHOCLASTIA, Lithotrity.

LITHOCYSTOTOMY, (*lithos*, κυστις, 'bladder,' and τομή, 'incision,') Lithotomy.

LITHODECTASIA, (*lithos*, ὁδός, 'a way,' and εκτασις, 'dilatation,') Lithectasy.

LITHODIALYSIS, *Lithol'y'sis*, (*lithos*, and διαλυω, 'to dissolve,') The dissolving of a stone in the bladder.

LITHODRASIS, *Lithodras'sicus*, (F.) *Lithodras'sique*, (*lithos*, and δρασσειν, 'to seize hold of,') An epithet given to a form of stone forceps—*Pince lithodras'sique*—used in the operation of lithotrity, by MM. Meirieu and Tanchou.

LITHOGENIA, (*lithos*, and γεννωω, 'to generate,') *Lithia*.

LITHOID, *Litho'ides*, *Lithoi'des*, (*lithos*, and εἶδος, 'resemblance,') Of the nature of stone, or resembling stone: as

LITHOIDES OS, see Temporal bone.

LITHOLABE (F.), *Lithol'abum*, (*lithos*, and λαβω, λαμβανω, 'I seize hold of,') An instrument,

employed for laying hold of a stone in the bladder, and keeping it fixed, so that lithotritic instruments can act upon it.

LITHOLABON, Forceps, (lithotomy.)

LITHOLABUM. An instrument concerned in extracting stone from the bladder. It had various shapes.

LITHOLYSIS, (*lithos*, and λυω, 'I dissolve,') Lithodialysis.

LITHOLYTE. An instrument, of the catheter kind, for conveying solvents of calculi into the bladder.

LITHOLYTIC, Lithontripctic.

LITHOMETRA, (*lithos*, and μετρα, 'the uterus,') Osseous, or other concretions of the uterus.

LITHOMYLIE, (*lithos*, and μυλη, 'a mill,') Lithotrity.

LITHON, see *Lithia*.

LITHONEPHRITIS, see Nephritis.

LITHONLYTIC, (*lithos*, and λυω, 'I dissolve,') Lithontripctic.

LITHONTHRYPTIC, (*lithos*, and θρυπτω, 'I break in pieces,') Lithontripctic.

LITHONTRIP'TIC, *Lithontrip'ticus*, *Lithon-thryp'tic*, *Litholyt'ic*, *Lithonlyt'ic*, *Calculif'ragus*, *Saxif'ragus*, (*lithos*, and τριβω, 'I rub,') A remedy believed to be capable of dissolving calculi in the urinary passages. There is not much reliance to be placed upon such remedies. By *antilithics* exhibited according to the chemical character of the calculus, (see Calculi, urinary,) the disease may be prevented from increasing; but most of the vaunted lithontripitics for dissolving the calculus already formed have been found unworthy of the high encomiums which have accompanied their introduction.

LITHOPEDION, *Infans lapid'eus*, *Osteopa'dion*, (*lithos*, and παις, gen. παῖδος, 'a child,') A foetus, petrified in the body of the mother.

LITHOPHAGUS, (*lithos*, and φάγω, 'I eat,') A stone-eater. Cases are on record of persons who have swallowed stones for purposes of exhibition.

LITHOPHTHISIS, see Calculi, pulmonary, and Lithiasis pulmonum.

LITHOPRINIE, (*lithos*, and πρινω, 'to saw,') Lithotrity.

LITHOPRIONE, (*lithos*, and πριων, 'a saw,') An instrument proposed by M. Leroy for preventing the fragments of a calculus, when subjected to lithotrity, from falling into the bladder. It is a variety of *litholabe*.

LITHOPRISIE, (*lithos*, and πρινω, 'to saw,') Lithotrity.

LITHORINEUR, (*lithos*, and ῥινειν, 'to file,') An instrument, proposed by MM. Meirieu and Tanchou for filing down calculi in the bladder.

LITHOS, (*Λιθος*), Calculus.

LITHOSPERMUM OFFICINALE, (*Λιθoσπepμoν*, from *litho*, and σπέρμα, 'seed,') *Mil'ium So'lis*, *Ægon'yehon*, *Grom'well*, *Grom'il*, *Grom'yl*, *Graymill*, *Gray millet*, *Bas'tard Al'kanet*, *Little-wale*, (F.) *Grémil officinal*, *Herbe aux Perles*. Ord. Boraginæ. The seeds of this plant were formerly supposed, from their stony hardness, (*lithos*, and σπέρμα, 'seed,') to be efficacious in calculous affections. They have, also, been considered diuretic.

LITHOSPERMUM VILLOSUM, *Anehusa tinctoria*.

LITHOTECNION, (*lithos*, and τεκνω, 'a child,') Lithopædion.

LITHOTERE'THRUM, (*lithos*, and τερειν, 'to rub,') A lithotritor.

LITHOTHLIBIE (F.), (*lithos*, and θλιβω, 'I crush,') The crushing of a friable calculus between a finger, passed by the rectum or vagina as far as the *bas-fond* of the bladder, and a catheter introduced into the bladder.

LITHOTHRYPsis, (*lithos*, and θρυπτω, 'I break in pieces,') Lithotrity.

LITHOTHRYPATORS, see Lithotrixy.

LITHOTOME, *Lithot'omus*, (*lithos*, and *tomē*, 'an incision.') This name has been given to a number of instruments of different shapes and sizes which are used in the operation for the stone, to cut the neck or body of the bladder. They ought with more propriety to be called *Cystotomes*.

The *Lithotome Caché* of Frère Côme is the most known, and is still occasionally used. It is composed of a handle and a flattened sheath, slightly curved: in this there is a cutting blade, which can be forced out, by pressing upon a *bas-cule* or lever, to any extent that may be wished by the operator.

A *Double Lithotome* was used by Dupuytren in his bilateral operation. See Lithotomy.

LITHOTOMIST, *Lithot'omus*. One who devotes himself entirely to operating for the stone. One who practises lithotomy.

LITHOTOMY, *Lithotom'ia*, *Cystotom'ia*, *Urolithotom'ia*, *Sec'tio vesica'lis*, *Lithocystot'omy*, (F.) *Lithotomie*, *Taille*. The operation by which a stone is extracted from the bladder. The methods according to which this operation may be practised are reducible to five principal, each of which has experienced numerous modifications.

1. The *Meth'od of Celsus*, *Meth'odus Celsiana*, *Cystotom'ia cum appa'atu par'vo*, *Appa'atus Minor*, *Cutting on the Gripe*. This consisted in cutting upon the stone after having made it project at the perinæum by means of the fingers introduced into the rectum. The method was attended with several inconveniences; such as the difficulty of dividing the parts neatly, injury done to the bladder, as well as the impossibility of drawing down the stone in many persons. It is sometimes also called *Meth'odus Guytoniana*; from Guy de Chauliac having endeavored to remove from it the discredit into which it had fallen in his time. It was termed *Apparatus Minor*, (F.) *Le petit appareil*, from the small number of instruments required in it.

2. *Appa'atus Major*. This method was invented in 1520 by John de Romani, a surgeon of Cremona, and communicated by him to Mariano Santo di Barletta, whence it was long called *Maria'no's Meth'od*, *Sec'tio Maria'na*. It was called, also, *Appa'atus Major*, and *Cystotom'ia seu Meth'odus cum appa'at'umag'no*, (F.) *Le grand appareil*, from the number of instruments required in it. An incision was made on the median line; but the neck of the bladder was not comprehended in it. It was merely dilated. The greater apparatus was liable to many inconveniences; such as ecchymoses; contusion; inflammation of the neck of the bladder; abscesses; urinary fistulæ; incontinence of urine, impotence, &c.

3. The *High Operation*, *Appa'atus altus*, *Cystotom'ia cum appa'atu alto*, *C. Hypogastrica*, *Epi-cystotom'ia*, *Laparo-cystotom'ia*, *Sec'tio seu Meth'odus Franconiana*, *S. Hypogastrica seu alt'a*, (F.) *Haut appareil*, *Taille Hypogastrique*, *Taille sus-pubienne*, was first practised by Peter Franco, about the middle of the 16th century. It consisted in pushing the stone above the pubis by the fingers introduced into the rectum. Rousset afterwards proposed to make the bladder rise above the pubis by injecting it. The method had fallen into discredit, when Frère Côme revived it. It is used when the calculus is very large. It was practised by opening first the membranous part of the urethra upon the catheter passed into the canal. Through this incision the *Sonde à dard*—a species of catheter, having a spear-pointed stilet—was introduced into the bladder. An incision was then made into the linea alba, above the symphysis pubis, of about four or five fingers' breadth, and the peritoneum detached

to avoid wounding it. The stilet was pushed through the bladder, and used as a director for the knife, with which the bladder was divided anteriorly, as far as the neck; and the stone extracted. It was performed in England by Douglass in 1719, and since by others, with various modifications.

4. The *Lat'eral Operation*, *Hypocystotom'ia*, *Cystotom'ia latera'lis*, *Cystauchenotom'ia*, *Cystotrachelotom'ia*, *Urethrocystauchenotom'ia*, *Urethrocystotrachelotom'ia*, *Sec'tio latera'lis*, *Appa'atus latera'lis*, (F.) *Appareil latéralisé*, so named from the prostate gland and neck of the bladder being cut laterally, was probably invented by Peter Franco. It was introduced into France by Frère Jacques de Beaulieu. He performed it with rude instruments invented by himself, and improved by the suggestions of some of the Parisian surgeons. In England, it received its earliest and most important improvements from the celebrated Cheselden. It is the method practised at the present day, according to different modes of procedure. In this method, the patient is placed upon a table; his legs and thighs are bent and separated; the hands being tied to the feet. The perinæum is then shaved, and a staff is introduced into the bladder, the handle being turned towards the right groin of the patient. An oblique incision is now made from the raphe to the middle of a line drawn from the anus to the tuberosity of the ischium of the left side; and, taking the staff for a guide, the integuments, areolar tissue of the perinæum, membranous portion of the urethra, transversus perinæi muscle, bulbo-cavernosus, some fibres of the levator ani, the prostate and neck of the bladder, are successively divided. For this latter part of the operation, the knife, the beaked bistoury, *Bistouri ou Lithotome Caché*, cutting gorget, &c., is used, according to the particular preference. The forceps are now introduced into the bladder, and the stone extracted. In the operation care must be taken not to injure the rectum or the great arterial vessels distributed to the perinæum.

A variety of the *Lat'eral Appa'atus*, called by the French *Appareil latéral*, consisted in cutting into the *bas fond* of the bladder, without touching the neck of that organ; but it was soon abandoned, on account of its inconveniences.

The method of Le Cat and of Pajola—*Urethrocystoënu'rysmatotom'ia*—consists in dividing the prostate in part only,—the enlargement of the wound being effected by a peculiar dilator.

The *Bilat'eral Operation* is founded on that of Celsus. It consists in making an incision posterior to the bulb of the urethra, and anterior to the anus, involving both sides of the perinæum by crossing the raphe at right angles: an incision is then made through the membranous part of the urethra, and the prostate may be cut bilaterally, either with the double lithotome of Dupuytren, or the prostatic bisector of Dr. Stevens, of New York.

5. *Lithotomy by the Rectum*, *Proctocystotom'ia*, *Sec'tio rectovesica'lis*, (F.) *Taille par la Rectum*, *Taille postérieure*, *T. Recto-vésicale*. This was proposed by Vegetius in the 16th century; but it was never noticed until M. Sanson, in the year 1817, attracted attention to it; since which time it has been successfully performed in many instances. It consists in penetrating the bladder through the paries corresponding with the rectum by first cutting the sphincter ani and rectum abor; the root of the penis, and penetrating the bladder by the neck of that organ, dividing the prostate, or by its *bas-fond*.

Lithotomy in women, from the shortness of the urethra, is a comparatively insignificant operation.

LIVER, ALBU'MINOUS, *L. amyloid*, 'arda'ceous.

scrofulous, var'y. Altered nutrition of the liver, in which its substance is altered so as to have those epithets applied to it.

LIVER, AMYLOID, L. albuminous.

LIVER DISEASE, Hepatopathia—l. Fatty, Adiposus hepatica—l. Gin, L. nutmeg—l. Gin-drinkers', L. nutmeg—l. Granulated, Cirrhosis of the liver.

LIVERGROWN, *Tu'mido jec'orè præ'ditus.* Having a large liver.

LIVER, HOBNAIL, Cirrhosis of the liver—l. Lardaceous, Liver albuminous—l. Mammillated, Cirrhosis of the liver.

LIVER, NUTMEG, *He'par variegatum, Tu'beriform liv'er.* An appearance of the liver when cut across, resembling that of the section of a nutmeg, *Myristicatio hepatis*, supposed by some to be the result of intemperance in the use of alcoholic drinks; but occurring under other causes. The terms *whis'ky liv'er, gin-drinkers' liv'er*, and *gin liv'er*, occasionally applied to it, are, consequently, not distinctive. Under the name *Pigment liv'er*, Frerichs describes the liver as it sometimes appears in those who die from the effects of malarious fevers; when it is of a steel gray or blackish, or, not unfrequently, chocolate colour; brown insulated figures on a dark ground—a condition, which "is produced by an accumulation of pigment matter in the vascular apparatus of the gland."

LIVER PIGMENT, see Liver, nutmeg.

LIVER, SCROFULOUS, Liver, albuminous.

LIVER SPOT, Chloasma—l. Starch, Glycogenic matter—l. Sugar, see Saccharum.

LIVER, TUBERCULATED, Cirrhosis of the liver—l. Tuberiform, L. nutmeg—l. Waxy, Liver, albuminous.

LIVER WEED, Hepatica triloba.

LIVER, WHISKY, Liver, nutmeg.

LIVER WORT, Hepatica triloba, Marchantia polymorpha—l. Wort, ground, ash-coloured, Lichen caninus—l. Wort, Iceland, Lichen Islandicus—l. Wort, noble, Hepatica triloba.

LIVIDUS MUSCULUS, Pectinalis.

LIVOR, ('a blue or livid spot,') Suggillation—l. Emortualis, see Suggillation—l. Sanguineus, see Eechymoma.

LIVRE, (from *libra*,) Pound.

LIVRET, (dim. of (F.) *livre*, (L.) *liber*, 'a book,' from its arrangement,) Omasum.

LIXIVIA TARTARIZATA, (from *lix*, 'ashes,') Potassæ tartras—l. Vitriolata, Potassæ sulphas—l. Vitriolata sulphurea, Potassæ sulphas cum sulphure.

LIXIVIE seu KALICUM ACETAS, Potassæ acetas.

LIXIV'IAL, *Lixivio'sus*, (F.) *Lixiviel, Lixivien*. An ancient term for salts obtained by washing vegetable ashes,—as the fixed alkalis.

LIXIVIA'TION, *Elxiviation, Lixivia'tio*. An operation which consists in washing wood-ashes with water, so as to dissolve the soluble parts. The filtered liquor is the *ley*.

LIXIVIEL, Lixivial.

LIXIV'IUM, *Lixiv'ia, Com'ia, Lee, Ley, Lye*, (F.) *Lessive*. Any solution containing potassa or soda—*Sal Lixivio'sum*—in excess. In domestic practice a ley is made from wood-ashes, and mixed with the ordinary basis of a poultice, to form a *ley cataplasm* or *poultice*, which is applied to whitlow and similar local inflammations.

LIXIV'IVUM AMMONIACALE, Liquor ammoniæ—l. Ammoniacale aromaticum, Spiritus ammoniæ aromaticus—l. Causticum, Liquor potassæ—l. Magistrale, Liquor potassæ—l. Saponarium, Liquor potassæ—l. Tartari, Liquor potassæ subcarbonatis.

LIXIVUS CINIS, ('lye ash,') Potash of commerce.

LYZ'ARD, *Lacer'ta, Lacer'tus*, (said to be so called in consequence of its limbs resembling the

arms (*lacerti*) of man [?].) *Sau'ra, Sau'ros*, (F.) *Lézard*. Lizards were formerly employed in medicine as sudorifics; and were, at one time, extolled in syphilis, cutaneous affections, and in cancer.

LIZARD'S TAIL, *Saururus cernuus*.

LOADED, see Enduit.

LOADSTONE, (from *lead* and *stone*,) Magnet.

LOATHING, (Sax. *laðian*, 'to loathe,') Disgust.

LOBAIRE, Lobar.

LO'BAR, *Loba'ria*, (F.) *Lobaire*. Relating or appertaining to a lobe or lobes.

LOBARIA ISLANDICA, Lichen Islandicus—l. Pulmonaria, Lichen pulmonarius—l. Saxatilis, Lichen Saxatilis.

LO'BATED, *Loba'tus*, (F.) *Lobé*, (from *lobe*.) Divided into lobes; as *Lob'ulated, Lobala'tus*, (F.) *Lobulé*, means divided into lobules.

LOBE, (*Λοβος*,) *Lo'bua*. A round, projecting part of an organ. The liver, lungs, and brain, for example, have lobes.

The under surface of the brain is divided into two anterior, two lateral, two posterior, and two intermediate lobes or processes. These Chaussier calls *lobules* of the brain: the cerebral hemispheres he terms *lobes*.

LOBE, BIVEN'TRAL. A wedgeshaped lobe of the cerebellum, situate behind the amygdala.

LOBE DOUDÉNAL, Lobulus Spigelii.

LOBE OF THE EAR, *Lob'ule of the Ear*, is a soft, rounded prominence, which terminates the circumference of the pavilion inferiorly, and is pierced in those who wear rings.

LOBE PANCRÉATIQUE, Lobulus Spigelii—l. *Petit du foie*, Lobulus Spigelii—l. *de Spigel*, Lobulus Spigelii.

LOBÉ, Lobated.

LOBELIA, (after Lobel, a Flemish botanist,) L. inflata—l. Blue, L. syphilitica.

LOBE'LIA CARDINA'LIS, *Lobe'lia coccin'ea, Trache'tium America'num, Car'dinal Plant, Car'dinal Flow'er, Scar'let Lobe'lia*. Ord. Lobeliaceæ. This species is indigenous in the United States. It blooms in autumn, having beautiful carmine flowers. The root is a reputed anthelmintic with the Indians.

LOBELIA COCCINEA, L. Cardinalis.

LOBELIA INFLA'TA, *In'dian or Wild Tobac'co, Puke Weed, Asthma Weed, Eye'bright, Emet'ic Weed, Lobe'lia* (Ph. U. S.), (F.) *Lobélie, L. enflée*. The prominent virtues of this American plant are those of an emetic. In smaller doses it is sedative, and has been given as a pectoral in croup, asthma, &c. It is, also, sudorific and cathartic, and is an acro-narcotic poison. Twenty grains acts as an emetic.

LOBELIA, PALESPIKED, L. spicata.

LOBELIA, PINIFOLIA. A South African plant, the root of which is excitant and diaphoretic. A decoction of it is sometimes used in the Cape Colony as a domestic remedy in cutaneous affections, chronic rheumatism, and gout.

LOBELIA, SCARLET, L. Cardinalis.

LOBELIA SPICA'TA, *Palespiked Lobelia*; indigenously: is a diuretic.

LOBELIA SYPHILIT'ICA, *L. reflex'a, Ranunculus Virginia'us, Ranu'tium Syphilit'icum, Blue Lobe'lia, Blue Car'dinal Flow'er*. The root of this plant, which is indigenous in the United States, is an emetic and drastic cathartic. It has been used in syphilis; hence its name. The mode of preparing it is to boil $\frac{3}{4}$ ss of the dried root in Oij of water, until the fluid is reduced to Oviij. Dose, Oss.

LOBÉLIE, Lobelia—l. *Enflée, Lobelia inflata*.

LOBES BIJUMEAUX, Quadrigena tubercula—l. Cerebral, see Lobe.

LOBES OF THE LIVER, *Pin'ne* seu *Lo'bi* seu *Pin'nae* *He'patis*. See Lobule.

LOBES, OPTIC, *Quadrigenina tubercula*—*l. Optiques*, *Quadrigenina tubercula*.

LOBI (pl. of *lobus*.) HEPATIS, Lobes of the liver—*l. Pulmonum*, see Pulmo.

LOBLOLLY, see Avena.

LOBLOLLY BOY, ('water gruel boy.') An attendant and aid to the surgeon on board ship.

LOBULAR, *Lobula'ris*, (F.) *Lobulaire*. Relating to or belonging to a lobule:—as *lobular pneumonia*, (F.) *Pneumonie lobulaire*, *P. mamelonnée*, *P. disséminée*. Pneumonia anatomically characterized by nuclei of red or gray hepatization disseminated in variable numbers in one or both lungs.

LOBULAR BIL'IARY PLEX'US. The plexus formed of lobular hepatic ducts, which are derived chiefly from the interlobular. This plexus forms the principal part of the substance of the lobule.

LOBULAR VE'NOUS PLEX'US. The plexus interposed between the interlobular portal veins, and the intralobular hepatic vein.

LOBULATED, see Lobated.

LOBULE, *Lobulus*, (dim. of *lobus*.) Mr. Kierman uses the term *lobule* for the acinus of the liver of many anatomists.

LOBULE OF THE CORPUS STRIATUM, *Insula cerebri*—*l. Carré*, *Lobulus anonymus*—*l. of the Ear*, *Lobe of the Ear*—*l. of the Fissure of Sylvius*, *Insula cerebri*—*l. du Foie*, *Lobulus Spigelii*—*l. antérieur du Foie*, *Lobulus anonymus*—*l. of the Liver*, see Liver—*l. of the Lung*, see Pulmo—*l. Pneumogastric*, *Flocculus*.

LOBULÉ, see Lobated.

LOBULES HEPATIQUES, see Liver.

LOBULI, (pl. of *lobulus*.) see Lobulus—*l. Pulmonales*, see Pulmo.

LOBULINS, see Liver.

LOBULUS, *Lobule*—*l. Accessorius anterior quadratus*, *L. anonymus*.

LOBULUS seu LO'BUS ANON'YMUS, *L. accessorius anterior quadratus*, *L. quadratus*, (F.) *Lobule carré*, *L. antérieur du foie*, *Éminence porte antérieure*. This is situate in the liver between the passage for the round ligament and the gallbladder, and is less prominent, but broader, than the *Lobulus caudatus*. From the lobulus anonymus a bridge runs across the passage for the round ligament. It is called *Pons seu Isth'mus he'patis*.

LOBULUS seu LO'BUS CAUDA'TUS, *Processus caudatus*. This is merely the root or one of the angles of the lobulus Spigelii, advancing towards the middle of the lower side of the great lobe, and representing a kind of tail. Also, the termination of the helix and anthelix of the ear, which is separated from the concha by an extensive fissure.

LOBULUS CENTRA'LIS. A small lobule or prominence of the superior vermiform process of the cerebellum situate in the incisura anterior.

LOBULUS NASI, see Nasus—*l. Pneumogastricus*, *Flocculus*—*l. Posterior*, *L. Spigelii*—*l. Posticus papillatus*, *L. Spigelii*—*l. Quadratus*, *L. anonymus*—*l. Renum*, *Ferrein*, pyramid of.

LOBULUS seu LO'BUS SPICE'LI, *L. posterior seu posticus papillatus*, (F.) *Éminence porte postérieure*, *Lobule ou Petit lobe du foie*, *Lobe de Spigel* ou *duodénal* ou *pancréatique*, is situate near the spine, upon the left side of the great lobe of the liver, and is of a pyramidal shape, projecting like a nipple, between the cardia and vena cava, at the small curvature of the stomach.

LOBUS, (*Λοβος*.) Lobe, see Lobulus—*l. Anonymus*, *Lobulus anonymus*—*l. Quadrangularis*, see Cerebellum—*l. Semilunaris*, see Cerebellum—*l. Superior anterior*, see Cerebellum—*l. Superior posterior*, see Cerebellum.

LO'CAL, *Loca'lis*, (from *locus*, 'a place,') *Topi-*

icus, *Mer'icus*, *Partia'lis*, *Topical*, (F.) *Locale*, *Topique*. Affections are called local—*Morbi locales*—when confined to a part, without implicating the general system; or, at all events, only secondarily. *Local* is thus opposed to *general*. In the classification of Dr. Wm. Farr, *Local diseases*, *Amorphanici* (*Morbi*), include sporadic diseases in which the functions of particular organs and systems are disturbed or obliterated, with or without inflammation; sometimes hereditary.

A local or topical application is one used externally. See Topic.

LOCH, *Loch*.

LOCHI'A or LO'CHIA, (*λοχεια*, *λοχια*.) *Purgament'a Puere'rii* seu *U'teri*, *Purga'tio puere'rii*, *Ly'ma*, (*λογος*, 'a woman in childbed,' *λοχια*, 'parturition,') (F.) *Lochies*, *Suites de couches*, *Vidanges*. The *clannings*. A serosanguineous discharge following delivery. During the first two or three days it is bloody; but afterwards becomes green-coloured,—*green waters*—and exhales a disagreeable and peculiar odour. The lochia, in different stages, have received the names *Lochia cruenta*, *L. serosa* and *L. alba* seu *muco'sa* seu *lacte'a*. The duration, quantity, and character of the discharge vary according to numerous circumstances. It flows from the part of the uterus which formed a medium of communication between the mother and foetus, and continues, usually, from 14 to 21 days. See Parturition.

LOCHIOCELIITIS, (*λοχεια*, 'childbirth,' *κελια*, 'abdomen,' and *itis*.) Puerperal fever.

LOCHIODOCHUM, *Lochodochium*.

LOCHIOPYRA, (*λοχεια*, 'childbirth,' and *πυρ*, 'fever,') Puerperal fever.

LOCHIORRHAG'IA, (*λοχια*, and *ῥήγνυμι*, 'to break forth,') An immoderate flow of the lochia. Hemorrhage from the uterus in the childbed state.

LOCHIORRHE'A, (*lochias*, and *ῥέω*, 'I flow,') (F.) *Lochiorrhée*. Discharge or immoderate discharge of the lochia.

LOCHIORUS RETENTIO, *Ischlochias*.

LOCHIOSCHESIS, (*lochias*, and *σχίζω*, 'to restrain,') *Ischoiochia*.

LOCHIOCH, *Looch*.

LOCHODOCHI'UM, *Lochiodochi'um*, (*lochias*, and *δέχομαι*, 'I receive,') An institution for the reception of pregnant and childbed females. A *Lying-in-hospital*.

LOCHOMETRITIS, (*lochias*, *μητρα*, 'uterus,' and *itis*.) see Metritis.

LOCHOMETROPHLEBITIS, (*lochias*, *μητρα*, 'uterus,' and *phlebitis*.) see Phlebitis.

LOCHOOPHORITIS, (*locho*, *οἰφωρον*, 'ovary,' and *itis*.) see Oaritis.

LOCHOPERITONITIS, see Peritonitis.

LOCHOS, (*λοχος*.) Puerpera.

LOCI, (pl. of *locus*, 'a place,') Uterus—*l. Muhières*, *Uterus*, *Vulva*.

LOCK HOSPITAL. A term of uncertain origin, generally given to venereal hospitals in Great Britain.

LOCKED JAW, Trismus.

LOCKING Coxofemoral articulation.

LOCOMOTEUR, Locomotive.

LOCOMOTILITY, see Locomotion.

LOCOMO'TION, *Locomoti'o*, (*locus*, 'a place,' and *movere*, *motum*, 'to move,') An action peculiar to animal bodies, by which they transport themselves from place to place. It, as well as *musculati'o*, has also been used for the function of animal movements. The faculty is sometimes called *Locomotive'ity* and *Locomotiv'ity*.

LOCOMOTION OF AN AR'TERY, is the movement produced in vessels having a curvature, by the impulse of the blood sent from the heart, which tends to straighten the artery, and causes the movement in question.

LOCOMO'TIVE, *Locomoto'ry*, *Locomoto'rius*, (F.) *Locomoteur*: (same etymon.) Relating or appertaining to locomotion—as 'the *locomotive* or *locomotory* apparatus,' (F.) *Appareil locomoteur*.

LOCOMOTIVITY, see Locomotion.

LOCULAMENTA COLI, (pl. of *loculamentum*, (from *locus*), 'a compartment, a cell,') see Colon.

LOCULAIRE, Locular.

LOC'ULAR, *Locula'ris*, (F.) *Loculaire*, *Loculeux*, (*loculus*, dim. of *locus*, 'a place,' 'a small place,') Provided with or divided into small spaces or compartments.

LOCULEUX, Locular.

LOCUS CINEREUS, *Substantia ferruginea*.

LOCUS NIGER, see Peduncles of the Brain.

LOCUS PERFORATUS ANTI'CUS seu ANTE'RIOR.

A triangular flat surface of the brain, which corresponds to the posterior extremity of each olfactory process in front of the optic tract, and near the entrance of the fissure of Sylvius: *Substantia seu Lam'ina perforata anterior*.

LOCUS PERFORATUS POSTICUS, see Tarini pons.

LOCUST, BLACK, *Robinia Pseudo-acacia*.

LOCUST EATERS, *Acridophagi*—l. Plant, *Cassia Marilandica*—l. Tree, *Robinia Pseudo-acacia*.

LOCUST, YELLOW, *Cladrastis tinctoria*.

LODGED, Wedged.

LODOICEA, see Coco of the Maldives.

LOECHE, MINERAL WATERS OF, Leuk, mineral waters of.

LEME, (*λοιμη*,) Plague.

LEMIA, (*λοιμη*,) Plague.

LEMICUM, (from *λοιμικος*, 'pertaining to the plague,') see Læmology.

LEMOCHOLOSIS, (*læmos*, and *χολη*, 'bile,') Fever, yellow.

LEMOCOMI'UM, (*læme*, and *κομω*, 'I take care of,') A hospital for those affected with plague.

LEMOGRAPHY, Loimography.

LEMOLOGIUM, see Læmology.

LEMOL'OGY, *Læmolog'ia*, (*læmos*, and *λογος*, 'a description,') The doctrine of plague and pestilential diseases. A treatise on the same,—*Læmicium, Læmolog'ium*.

LEMOPHTHALMIA, see Ophthalmia.

LEMOPYRA, (*læmos*, and *πυρ*, 'fire, fever,') Plague.

LEMOS, (*λοιμος*,) Plague.

LOGADES, (*λογας*, pl. *λογαδες*, 'white of the eye,') Sclerotic.

LOGADITIS, (*logades*, and *itis*,) Sclerotitis.

LOGIATROS, *Logia'ter*, (*logos*, and *ιατρος*, 'a physician,') In the bad sense, a position without experience; a mere theorist. In the good sense, a rational physician; one who treats disease according to theoretical or scientific principles.

LOGONEURO'SES, (*logos*, and *νευρον*, 'a nerve,') Diseases of the mental activity.

LOGOS, (*λογος*,) Reason.

LOGWOOD, *Hæmatoxylon Campechianum*.

LOHOCH, Looch.

LOIMOCHOLOSIS, (*loimos*, and *χολη*, 'bile,') Fever, yellow.

LOIMOGRAPHY, *Loimograph'ia*, *Læmog'raphy*, (*læmo*, and *γραφω*, 'I describe,') A description of the plague and pestilential diseases.

LOIMOLOGY, (*loimos*, and *λογος*, 'a discourse,') Læmology.

LOIMOS, (*λοιμος*,) Plague.

LOINS, ([W.] *lœyn*,) Lumbi.

LOISELEURIA PROCUMBENS, (after Loiseleur Deslongchamps, the botanist,) *Azalea procumbens*; indigenous; Order, Ericaceæ; has some reputation as an astringent.

LOLIACEUM RADICE REPENTE, (after Solium, Triticum repens.

LOLIUM ANNUUM, *L. temulentum*.

LOLIUM TEMULENT'UM, *L. annuum, Crapa'lia seu Crapa'lia temulen'ta, Bromus temuler'tus, Dar'*

nel, (Prov.) *Cock'grass*, *Jum*, (F.) *Herbe d'Irrogne, Ieraie, Ieraie*. A species of the genus *Lolium*; Ord. Graminææ; *Sax. Syst.* Triandria digynia; which has decidedly poisonous properties; occasioning, when mixed in bread or beer, intoxication, vertigo, nausea, and vomiting.

LOMBAIRE, Lumbar.

LOMBES, Lumbi.

LOMBO-ABDOMINAL, *Transversalis abdominis*—l. *Costal*, *Serratus posticus inferior*—l. *Costo-trachélien*, *Sacrolumbalis*—l. *Dorso-spinal*, *Transversalis dorsi*—l. *Dorso-trachélien*, *Longissimus dorsi*—l. *Huméral*, *Latissimus dorsi*—l. *Sacré*, *Lumbosacral*.

LOMBRIC, *Ascaris lumbricoides*.

LOMBRICAUX, Lumbricales.

LOMBRICOIDE, *Ascaris lumbricoides*.

LONGHADES, (for *Logades*,) see Sclerotic.

LONGHADITIS, (for *Logaditis*,) Sclerotitis.

LONGCHITIS, (*λογχον*, 'the head of a lance,') *Polypodium filix mas*.

LONG, *Lon'gus*, ([Sax.] *lang*, *long*,) *Ma'eros*. That which is much greater in length than in breadth; as the *long bones*. The epithet is, also, applied to several muscles, to distinguish them from others of similar function, when the latter are shorter. We say, for instance, *long flexors*, and *long extensors*, in opposition to *short flexors*, and *short extensors*.

LONG DU COU, *Longus colli*—l. *du Dos*, *Longissimus dorsi*.

LONGÆVUS, (*longus*, 'long,' and *ævum*, 'age,') *Macrobiotic*.

LONGANON, *Rectum*.

LONGAON, *Rectum*.

LONGAS, *Rectum*.

LONGEV'ITY, *Longæ'vitas*, (F.) *Longévité*, (*longus*, 'long,' and *ævum*, 'age,') *Macrobiosis*, *Macrobio'tes*. The prolongation of existence to an advanced age. Haller collected examples of more than one thousand centenarians. He had knowledge of sixty-two persons aged from 110 to 120 years; of twenty-nine, from 120 to 130 years; and of fifteen, who had attained from 130 to 140 years. Beyond this advanced age, examples of longevity are much more rare and less sufficiently attested.

Longevity also means *length or dura'tion of life*, (F.) *Durée de la vie*. The mean age at death, (F.) *Vie moyenne*, of different classes and professions enables an estimate to be formed of the expectation or value of life in each.

LONGING, *Malacia*.

LONGISSIMUS, (superlative of *longus*,) A term applied to parts, which are very long in their relation to others.

LONGIS'SIMUS DOR'SI, *Semispina'tus*, (F.) *Lombo-dorso-trachélien*, *Portion costo-trachélienne du sacro-spinal* (Ch.), *Long dorsal*, *Long du dos*, is situate vertically at the posterior part of the trunk, and fills, in a great measure, the vertebral furrows. It is thick and almost square below; thin and pointed above. It is attached to the posterior surface of the sacrum, to the transverse processes of all the lumbar and dorsal vertebræ, and to the inferior margin of the last 7 or 8 ribs. It maintains the vertebral column in a straight position; straightens it when bent forwards, and can even carry it back. It also assists in the rotatory motion of the trunk.

LONGISSIMUS FEMORIS, *Sartorius*—l. *Oculi*, *Obliquus superior oculi*.

LONGSIGHTEDNESS, *Presbytia*.

LONGUS, *Long*.

LON'GUS COL'LI, ('long [muscle] of the neck,') (F.) *Pré-dorso-cervical*, *Pré-dorso-alloïdien*, (Ch.), *Long du cou*. This muscle is situate at the anterior and superior part of the vertebral column. It is long, flat, and broader at its middle

than at its extremities, which are pointed. It is attached to the anterior surface of the bodies of the first three dorsal and last six cervical vertebrae; to the intervertebral ligaments; to the anterior edge of the transverse processes of the last five cervical vertebrae; and to the tubercle on the anterior arch of the first. This muscle bends the cervical vertebrae upon each other and upon the dorsal vertebrae. If the upper portion acts on one side only, it occasions the rotation of the atlas on the vertebra dentata; and, consequently, of the head on the neck.

LONICE'RA BRACHYP'ODA, (called after *Lonicera*, a German botanist.) *Ord.* Caprifoliaceae. A Japanese plant. An infusion or decoction of the leaves is possessed of diuretic properties.

LONICERA DIERVILLA, *Diervilla trifida*—*l.* Germanica, *l.* Periclymenum—*l.* Marilandica, *Spigelia Marilandica*.

LONICE'RA PERICLY'MENUM, *L. Germanica*, *Periclymenum*, *P. vulgare*, *Caprifolium*, *C. Periclymenum* seu *syneal'icum* seu *distinctum*, *Common Woodbine*, *Hon'ey-suckle*, (Prov.) *Suck'ling*, (F.) *Chèvre-feuille*. This common plant is slightly astringent and tonic, and was formerly much used in gargles.

LONICERA SEMPERVIRENS, *Caprifolium sempervirens*, *Scar'let* or *Trump'et Hon'ey-suckle*. Indigenous. A syrup of the leaves is said to have been given in asthma and in amygdalitis. [?]

LONK, Coxofemoral articulation.

LO'OCH, *Lo'hoch*, *Loch*, *Loch'och*, (of Arabic extraction.) A linctus. See *Eclegma* and *Eclectos*.

LOOCH ABS'QUE EMULSIO'NÈ PARA'TUM, *Look préparé sans émulsion*; *Lo'och préparé* without emulsion. (*Puls. g. trag.* gr. xvj—gr. xxx, *Ol. amygd. dulc.* 3ss., *Sacchar.* 3j. *Aque* 3iij, *Aque flor. aurant.* 3ij. Mix by rubbing in a marble mortar. Ph. P.) Demulcent.

LOOCH AL'BUM, *L. amygdalinum*, *Linctus albus* seu *amygdalinus* seu *communis*. *Eclegma album* seu *gummo'soileum*, (F.) *Looch blanc*, *L. b. amygdalin*, *L. b. pectoral*. (*Amygd. dulc.* 3ss., *Amygd. amar.* No. ij, *Sacchar. alb.* 3iv. Make an emulsion by gradually adding 3iv of *Water*. Then take *Puls. tragacanth.* gr. xvj. *Ol. amygd. dulc. recent.* 3ss., *Sacch.* 3ij. Add the almond milk gradually to this, and afterwards *Aq. flor. aurant.* 3ij. Ph. P.) It is demulcent and pectoral.

LOOCH AMYGDALINUM, *L. album*—*l. Blanc*, *L. album*—*l. cum Croco* et *pistaciis*, *L. viride*—*l. of Egg*, *L. ex Ovo*.

LOOCH EX O'VO, *L. pectoralè le'niens*, *Po'tio* seu *emul'sio* seu *mistu'ra le'niens* seu *oleo'sa*, *Looch of Egg*, (F.) *Look d'œuf*, *Looch rouge*, *Émulsion huileuse*, *Mixture calmante*, *Potion pectorale*, *Lait adoucissant*. (*Vitell. ovi. recent.* 3ss., *Ol. amygd. dulc.* 3iss., *Syrup. althææ*, 3j. Rub in a mortar, and add by degrees, *Aq. flor. aurant.* 3j, *Aq. papav. rhæad.* 3ij. Ph. P.) Virtues the same as the preceding.

LOOCH ROUGE, *L. ex Ovo*—*l. Vert.*, *L. viride*.

LOOCH VIRIDÈ, *L. cum cro'co* et *pista'ciis*, (F.) *Looch vert.* (*Syrup. Violar.* 3j, *Tinct. croci* gtt. xx., *Aque*, 3iv. Mix, and add *Pistacia semin. sic.* 3vj. Ph. P.) Virtues like the last.

LOOK, *Looch*—*l. d'œuf*, *Looch ex ovo*—*l. Préparé sans émulsion*, *Looch absque emulsione paratum*.

LOOSESTRIFE, CREEPING, *Lysimachia nummularia*—*l.* Fourleaved, *Lysimachia quadrifolia*.

LOOSENESS OF THE BOWELS, *Diarrhoea*—*l.* of the Teeth, *Odontoseisis*.

LO'PEZ RA'DIX, *Ra'dix lopezia'na* seu *In'dica lopezia'na*. The root of an unknown Indian tree, not possessed of any remarkable smell or

taste, or of any appearance of resinous matter. It has been extolled, notwithstanding, in cases of colliquative diarrhoea. Gaubius compares its action to that of simarouba, but thinks it more efficacious.

LOPHADIA, *Lophia*.

LOPHIA (λοφία), *Lophadi'a*, *Lophos*. The upper part of the neck. The first vertebra of the back—*Vertebra dor'si prima*.

LOPHOS (λοφος), *Lophia*.

LOPIMA, *Fagus castanea*.

LOQUAC'ITY, *Garrulitas*; (from *loqui*, 'to speak,') *Loquacitè*. (F.) *Babillement*. The act of speaking with volubility. It is sometimes a symptom of disease, and is observable in hysteria, &c.

LOQUELA, (from *loqui*, 'to speak,') Voice, articulated—*l.* Abolita, *Aphonia*—*l.* Blæsa, *Balbuties*—*l.* Impedita, *Baryphonia*.

LORDO'SIS, *Lordo'ma*, (from *lordos*, 'curved,' 'bent,') *Repa'ndatio*, *Repa'nditas*. A name given to curvatures of the bones in general; and particularly to that of the vertebral column forwards; *Spina dor'si intror'sum flex'a*, *Emprostocyrto'ma*. This gives rise to the projection of the sternum called *chickenbreasted* or *pigeon-breasted*; the *Pectus carina'tum*, *Pig'con breast*, *Narrow chest*.

LORICA'TIO, (*loricare*, 'to plaster,' 'to encrust,') In Pharmacy, the application of a coating of clay, or other substance, to vessels which are to be exposed to fire.

LORIND MATRICOIS, *Epilepsia uteri'na*, *Convulsivus uteri morbus*. A barbarous name given to a pretended epilepsy of the womb.

LORIPES, gen. *Lorip'edis*, (*lorum*, 'a thong,' and *pes*, 'a foot,') See *Kyllosis*.

LORUS, *Hydrargyrum*.

LOSANGE, *Labella*.

LOT, *Urine*.

LOTA MOLVA, see *Oleum jecoris aselli*—*l.* *Vulgaris*, see *Oleum jecoris aselli*.

LOTIO, (*lavare*, *lotum*, 'to wash,') *Clyster*, *Lotion*—*l.* *Cupri sulphatis camphorata*, *Lotion*, camphorated, of *Bates*—*l.* *Rubra*, *Lotion*, camphorated, of *Bates*.

LOTIO RU'BRA, *Red wash*, (*Hydrarg. chlorid. corrosiv.* gr. xviii; *H. sulphureti rubr.* gr. ix; *Creasot.* ℥vi; *Aque f3viij.*) May be used in the proportion of f3j; to from f3i to f3iij of water, like the yellow wash.

LOTIO SAPONACEA, see *Sapo*.

LO'TION, *Lo'tio*, *Lotu'ra*, *Collu'tion*, (F.) *Hydrolotif*, *Hydrolé*. A fluid external application. Lotions are ordinarily applied by wetting linen in them and keeping it on the part affected.

LOTION, *BAR'LOW'S*, *Lotion of Sulph'uret of potass'ium*. (*Potassii sulphur.* 3iij; *Sapon.* 3iss; *Aq. Calcis*, f3viiss; *Alcohol. dilut.* f3ij. M.) Used in various chronic cutaneous diseases.

LOTION, CAMPHORATED, OF BATES, *Bates's camphorated water*, *Lo'tio cupri sulphatis camphorata*, *Lo'tio ru'bra*, *Red wash*. (*Cupri sulphat.* 3ij; *Bol. rubr.* 3ij; *Camphor.* 3ss; *Aq. bullient.* Oij. Strain through linen. f3ij of this, added to f3iv of water, is *Bates's Collyrium*, *Collyrium Batesanum*, which may also be made extemporaneously as follows:—*Cupri sulphat.* gr. viij; *Boli*, gr. viij; *Camphor.* gr. ij; *Aq. bullient.* f3viij.) Mix and filter. See *Lotio rubra*.

LOTION, FEBURE'S. A cancer remedy, which consisted of *Arsenic*, gr. x; *Water*, Oj; *Extract of hemlock*, 3j; *Liquor Plumbi subacetatis*, f3iij; *Laudanum*, f3j.

LOTION, GOULARD'S, *Liquor plumbi subacetatis dilutus*.

LOTION, GOW'LAND'S. An empirical preparation. (*Bitter almonds*, 3j; *Sugar*, 3ij; *Distilled water*, ℥ij. Grind together, strain, and add *Cor-*

rosive sublimate, *℞ij*, previously ground with *Sp. vini rect.* *℞ij*.) Used in obstinate eruptions.

LOTION, GRANVILLE'S COUNTERIRRITANT, *Granville's Lotion*, *Granville's antidyneous lotion*. Of this lotion, Dr. Granville gives two forms—a milder, and a stronger. The milder is made as follows: *Liq. ammon. fort.* *℥ij*; *Sp. Rosmarin.* *℥vj*; *Tinct. camphor.* *℥ij*, *M*.

The stronger is made as follows: *Liq. ammon. fort.* *℥x*; *Spir. Rosmar.* *℥ss*; *Tinct. camphor.* *℥ij*, *M*.

The stronger lotion vesicates rapidly. A piece of cotton or linen folded six or seven times, or a piece of thick flannel, may be imbued with it, and laid for a few minutes on the part to be irritated.

The Dublin Pharmacopœia has a *Linimentum Ammoniac Compositum*, Compound Liniment of Ammonia, which is a close imitation of *Granville's Lotion*.

LOTION, HAN'NAY'S Preventive wash. This famous nostrum for the prevention of venereal infection was nothing more than a solution of caustic potassa.

LOTION, HYDROCYAN'IC, *Lo'tio Ac'idi Hydrocyan'ici*. (*Hydrocyanic acid*, *℥iv*; *Rectified spirit of wine*, *℥ij*; *Distilled water*, *℥vss*.) Used with much success in impetigo, &c.

LOTION, STRUVE'S, FOR HOOP'ING-COUGH. (*Antim. et Potass. tart.* *℥j*; *Aque*, *℥ij*. Add *Tinct. cantharid.* *℥j*.) A counterirritant to the chest.

LOTIUM, Urine.

LOTURA, (*lavare, lotum*, 'to wash.') Lotion.—*L. Carnium*. The washings of flesh.

LOTUS SYLVESTRIS, (*λωτος*), *Trifolium melilotus*—*L. Virginiana*, *Diospyros Virginiana*.

LOUCHEMENT, (from *Lucus*,) Strabismus.

LOUCHES, see Strabismus.

LOUPE, Lupus.

LOUPE, (from *λοφος*, 'a lobe' [?],) Wen.

LOUSE, (*Sax. luf*), Pediculus.

LOUSINESS, Phtheiriasis.

LOUSY DISEASE, Phtheiriasis.

LOUTRON (*λουτρον*), Bath.

LOVAGE, *Ligusticum levisticum*.

LOVE, *E'ros*, *A'mor*, (from *Sax. lufran*, 'to love.') (*F.*) *Amour*. A tender and elevated feeling, which attracts one sex to the other. Love is occasionally a cause of disease, especially of insanity.

LOVE APPLE PLANT, *Solanum lycopersicum*—*L. Begot*, Illegitimate—*L. Child*, Illegitimate—*L. Pea*, *Abrus precatorius*.

LOW SPIRITS, Hypochondriasis.

LOW'ER, TU'BERCLE OF, *Tuber'culum Low'eri*, (after Dr. Richard Lower). Anatomists have given this name to a small projection, the existence of which is by no means constant, and which is found in the sinus venosus between the superior and inferior cava.

LOXAR'THRUS, *Loxar'thrum*, (*λοξος*, 'oblique,' and *αρθρον*, 'articulation.') (*F.*) *Lor'w'ithe*, *Per'v'ersion de la t'ete des os et des muscles*. A vicious deviation or direction of the joints, without spasm or luxation,—as in *clubfoot*.

LOXIA, (*λοξος*, 'oblique,') Torticollis.

LOXOPHTHALMUS, (*λοξος*, 'oblique,' and *οφθαλμος*, 'eye,') Strabismus.

LOZANGIA, Tabella.

LOZENGE, Tabella.

LOZENGES, BARK, *Tabellæ cinchonæ*—*L. Bismuth*, *Trochisci Bismuthi*—*L. Catechu*, *Trochisci catechu*—*L. Catechu* and *magnesia*, *Tabellæ antimoniales Kunckelii*—*L. Faustinus's Faustini pastilli*—*L. for the Heart-burn*, *Trochisci carbonatis calcis*—*L. Magnesia*, *Tabellæ de magnesia*—*L. of Marshmallows*, *Tabellæ de althæa*—*L. Opium*, see *Trochisci Glycyrrhizæ cum Opio*—*L. of Oxalic acid*, *Tabellæ acidæ oxalici*—*L. Pectoral*, *lack*, *Trochisci glycyrrhizæ glabræ*—*L. Pectoral*,

of *emetine*, *Trochisci emetinæ pectorales*—*L. Rhu-barb*, *Tabellæ de rheo*—*L. of Scammony and senna*, compound, *Tabellæ de scammonio et sennâ*—*L. Spitta's*, see *Trochisci glycyrrhizæ cum opio*—*L. Steel*, *Tabellæ de ferro*—*L. Steel*, aromatic, see *Tabellæ de ferro*—*L. Sulphur*, compound, *Tabellæ de sulphure compositæ*—*L. Sulphur*, simple, *Tabellæ de sulphure simplicis*—*L. of Sulphuret of antimony*, *Tabellæ antimoniales Kunckelii*—*L. Tannin*, *Trochisci acidi tannici*—*L. Wistar's*, *Trochisci glycyrrhizæ cum opio*—*L. Worm*, *Ching's*, see *Worm lozenges*, *Ching's*.

LUBIDO, gen. *Lubi'dinis*, Libido—*L. Intestini*, *Voluntas desedendi*.

LUBRICANTIA, (*lubricare*, 'to make slippery,') Demulcents, Emollients.

LUBRICUM CAPUT, ('slippery or smooth head,') Penis.

LUC'CA, MIN'ERAL WATERS AND CLIMATE OF. The baths and waters, near this ancient Italian city, have been long celebrated. They are thermal, and resemble, in properties, those of *Plombières* in France, and of *Bath*, England. They contain carbonic acid, sulphates of alumina, soda, magnesia, and iron. There are ten different sources, the temperature of which varies from 94° to 130° Fah.

Luca is much frequented in summer; partly on account of its mineral waters, but more on account of the coolness of the situation.

LUCHON, Bagnères-de-Luchon.

LU'CID, *Lu'cidus*, (*luceo*, 'I shine.') In medicine, the word *lucid* is particularly applied to the intervals, *Inter'v'ella lu'cida*, of apparent reason, which occur in mental alienation.

LUCIFERMATCHMAKERS' DISEASE, see Phosphorus.

LUCIF'UGUS, (*lux*, gen. *lucis*, 'light,' and *fugere*, 'to shun,') *Photoph'obus*, *Photophob'icus*. Dreading or avoiding the light.

LUCINA, (from *lux*, gen. *lucis*, 'light,' because she brought to light; a surname of Juno and Diana,) *lithyia*.

LUCKYHOOD, (because supposed to give luck,) *Caul*.

LUCOMANIA, (*λυκος*, 'a wolf,' and *mania*,) *Lycanthropia*.

LUCTUEUX (*F.*), *Luctuo'sus*, 'mournful,' *Plaintive*. An epithet applied to the respiration—(*F.*) *Respiration luctueuse*—when accompanied by moaning.

LUCUMA, (*S.*) A fruit which grows in the southern provinces of the coast of Peru, and the north of Chili. It is round; and the gray-brown husk encloses a fibrous, dry, yellow-coloured fruit with its kernel.

LUCUMORIA'NUS, (probably from *lux*, gen. *lucis*, 'light,' and *morari*, 'to tarry,') Continuing for several days: hence, *Dormit'io leucomoria'na*. A morbid sleep persisting for several days.

LU'DUS HELMONTII, *L. Paracel'si*, *Fel ter'æ*. A calcareous stone, the precise nature not known, which was used by the ancients in calculous affections. The term was also applied to every species of calculous concretion occurring in the animal body.

LUDUS PARACELSI, *L. Helmontii*.

LUDWIG'IA PALUSTRIS, (after Ludwig, a botanist and physician of Leipzig,) *L. nit'ida*, *Inar'dia palus'tris*, *Walter Pur'slane*, *Phth'i'sisweed*. Order, *Onagraceæ*; indigenous. An infusion of the dried plant has been used as a pectoral in asthma and chronic cough.

LUES, (*lucere*, 'to dissolve, melt away,') Disease, Plague, Syphilis—*L. Deifica*, Epilepsy—*L. Divina*, Epilepsy—*L. Dysenterica*, Dysentery—*L. Gonorrhœica*, *Gonorrhœica impura*—*L. Gutturis epidemica*, *Cynanche maligna*—*L. Indica*, *Frambœsia*—*L. Inguinaria*, Plague—*L. Pannoniæ*, Fever, Hungary—*L. Polonica*, Plica—*L. Sarmatica*, Plica—*L.*

Scorbutica, see Purpura.—1. Syphilis, Syphilis.—1. Syphilodes, Syphilis pseudo-syphilis.—1. Trichomatia, Plica.—1. Venerea, Syphilis.

LUETTE, Uvula.—1. *Vésicale*, see Urinary bladder.

LUG, Ear.

LUGDUS, Erysipelas.

LUJULA, (corruption of *Hallelujah*), Oxalis acetosella.

LULLABY SPEECH, Lallation.

LUMBA'GO, gen. *Lumbag'ginis*, (from *lumbi*.) *Arthro'sia lumborum*, *Lumba'go seu Nephral'gia rheumat'ica*, *Rachir'rheum'a*, *Rachior'rheum'a*, *Rheumatism'us dors'al'is*, *Osp'h'yrrheum'a*. Rheumatism affecting the lumbar region.

Lumba'go psöä'd'ica, *L. apostemato'sa*, *L. ab arthro'acë*. Pain in the loins from abscess. See Psöitis.

LUMBAGO A NISU, (*a nisu*, 'from effort,') see Effort.

LUM'BAR, *Lumba'ris* seu *Lumba'lis*, (F.) *Lombaire*, (from *lumbi*.) Belonging or having reference to the loins.

LUMBAR AB'SCESS, *Pso'a's ab'scess*, *Apostema psöä'ticum*, *Abscess'us lumborum*, *Mor'bus psöä'd'icus*. This abscess is so called from the matter being found on the side of the *psaos* muscle, or betwixt that and the *iliacus internus*. Between these muscles is a quantity of loose, areolar substance; and when an abscess takes place there, it can find no outlet except by a distant course. Generally, it descends along the *psaos* muscle, forming a swelling immediately beneath *Poupart's* ligament; at times, however, it extends down the thigh under the fascia. Severe hectic follows the bursting of the abscess, and often death. Its causes are,—scrofula, injury to the loins, &c.

When, from the pain of the back continuing for some time, with other symptoms, the disease is suspected, caustics, or the moxa, applied opposite the transverse processes of the lumbar vertebrae, may be recommended; and, when the abscess is ready for opening, it may be emptied by repeated discharges of the matter, through a small opening, made with a lancet, or small, lancetpointed trocar. The medical treatment consists in supporting the system, under the great reparative efforts required of it.

LUMBAR ARTERIES are four or five in number on each side. They arise from the sides of the abdominal aorta, and pass behind the muscles situate in front of the lumbar portion of the spine, to gain the broad muscles of the abdomen. They give off, 1. Spinal branches. 2. Anterior, posterior, and external muscular branches.

LUMBAR NERVES are five in number, and issue from the vertebral column by the spinal foramina of the loins. The first lumbar nerve gives off three branches:—the *external or ilioacral*; the *middle or inguinoventaneous*, and the *internal or infrapubian*. Along with the three pairs below it, it forms the lumbar plexus.

LUMBAR PLEX'US, *Portio lombaire du plexus crural* (Ch.), *Plexus Lombo-abdominal*. This plexus is formed by the union of the *Rami communican'tes* of the anterior branches of the first four lumbar nerves. It is situate behind the *psaos* muscle, and before the transverse processes of the lumbar vertebrae. It furnishes, besides the branches which proceed from the first pair, several filaments, that are distributed to the *psaos* muscle, to the *iliacus*, the integuments, and glands of the groin; and three great branches,—the *crural*, *obturator*, and *lumbosacral*.

LUMBAR REGION, *Lumbi*.

LUMBAR VEINS have an arrangement analogous to that of the arteries of the same name. They communicate with the vertebral sinuses, *azygos* veins, &c., and pass into the *vena cava inferior*.

LUMBARIS EXTERNUS, *Quadratus lumborum*—1. *Internus*, *Psoas magnus*.

LUM'BI, (pl. of *lumbus*.) *The Loins or Loins*, *Lumbar' region*, (Old. Eng.) *Lende*, (Sc.) *Lendis*; *Re'gio lum'balis*, *Lap'ara*, *Pso'a*, plural *Pso'a's*, *Os'phys*, *Reins*, (F.) *Lombes*. The posterior regions of the abdomen, comprised between the base of the chest and the pelvis. The parts which enter into the formation of the lumbar region are,—the skin; a considerable quantity of areolar texture; broad and strong aponeuroses;—the *Latissimus Dorsi*, *Obliquus externus*, and *Obliquus internus abdominis*, *Transversalis abdominis*, *Quadratus lumborum*, and the mass common to the *Sacro-lumbalis*, *Longissimus dorsi*, and *Multifidus spinæ*. These muscles surround the lumbar region of the vertebral column. The vessels, nerves, &c. of the loins, are called *lumbar*.

LUMBO, in composition, relating or belonging to the loins or lumbar vertebrae.

LUMBOCOSTALIS, *Sacrospinalis*.

LUMBOSACRAL. Belonging to the lumbar and sacral regions. (F.) *Lombo-sacré*. *Bichat* calls this a very large nerve, given off from the anterior branch of the fifth lumbar pair, which descends into the pelvis before the sacrum to join the sciatic plexus.

LUMBRICA'LES, *Vermicula'ris*. Resembling a *lumbricus* or 'earthworm,' *Lumbricoïdes*. A name given to small muscles, (F.) *Lombricux*, met with in the palm of the hand and sole of the foot.

LUMBRICA'LES MA'NÛS, *Fidicina'les*, *Flexor primi internodii digitorum manûs seu perforatus lumbrical'is*, (F.) *Annul-tendino-phalangiens*, *Palui-phalangiens*. Four small, fleshy, thin, round, long, fusiform fasciculi, situate in the palm of the hand, and distinguished into first, second, third, and fourth, counting from without to within. They arise from the tendons of the flexor communis digitorum, and are inserted at the outer and posterior side of the superior extremity of the first phalanges of the last four fingers. These muscles bend the fingers on the metacarpus, and fix the tendons of the flexor digitorum communis.

LUMBRICALES PE'DIS, (F.) *Planti-tendino-phalangiens*, *Planti-sous-phalangiens*, (Ch.) They are analogous to those of the hand in form, number, and arrangement. They increase the flexion of the toes, and draw them inwards.

LUMBRICUS, ('earthworm,') *Ascaris lumbricoïdes*—1. *Latus*, *Bothriocephalus latus*, *Tænia solium*—1. *Teres hominis*, *Ascaris lumbricoïdes*.

LUMBUS VENERIS, ('loin of Venus,') *Achillea millefolium*.

LUMEN, gen. *Lu'minis*, Light, Pupil—1. *Conjunctans*, *Phosphorus*.

LUMIÈRE, (from *lumen*, 'a light,') Light.

LUMINOSITY, *Phosphorescence*.

LUNA, ('the moon,') *Argentum*, Moon—1. *Albini*, *Sciatic notch*, lesser—1. *Imperfecta*, *Bismuth*—1. *Potabilis*, *Argenti nitras*.

LUNACY, see Lunatic.

LUN'AR, *Luna'ris*, (from *luna*, 'the moon, silver,') Relating or appertaining to the moon. See Lunatic;—or to silver, as '*Lunar caustic*.'

LUNA'RE OS. *Os semilunare*. The second bone in the upper row of the carpus.

LUNARIA, (*luna*, 'the moon,') *Menses*.

LUNARIA REDIV'VA. *Leucoïum*, *Bulbonach*, *Sat'in*, *How'city*. *Ord. Cruciferae*. Said, by Ray, to be a warm diuretic.

LUNARIS, Lunatic.

LUN'ATIC, *Lunaticus*, *Luna'ris*, *Luna'rius*, *Lu'nar*, (F.) *Lunatique*, (from *luna*.) Relating to the moon. An epithet given to diseases—as epilepsy and insanity—which were supposed to appear at certain phases of the moon; or to those

who were affected by them; *Lunatic'i*, *Seleni'aci*. The term *lunatic* (*Moon'ed*, *Moon' struck*, *Seleno-ble'tus*), is now restricted to one labouring under lunacy or mental alienation, *Dement'ia accident'a-lis* seu *adventit'ia*, that is, under any form of unsoundness of mind, except idiocy. In law, a *lunatic* is one who has had an understanding, but by disease, grief, or other accident, has lost the use of his reason, and become *non compos mentis* or *non compos*.

LUNATICUS, Epileptic, Lunatic, Somnambulist.

LUNATISMUS, Somnambulism.

LUNE, (from *luna*, 'the moon,') A fit of lunacy or insanity; a mad freak.

LUNE, Moon — *l. d'Eau*, Nymphæa alba.

LUNELLA, (dim. of *luna*, 'the moon,') Hyppoon.

LUNES, (pl. of *lune*, 'a moon or month,') Menses.

LUNETTES ORDINAIRES, (dim. of *luna*, 'moon,' from their shape,) Spectacles.

LUNG, (Sax. *lunge*.) Pulmo — *l.* Black, of coal miners, Anthracosis — *l.* Cancer of the, Phthisis, cancerous — *l.* Collapse of the, see Atelectasis pulmonum.

LUNG FEVER, Catarrh, Pneumonia.

LUNG, PERFORATING ABSCESS OF THE. A purulent collection, which forms exterior to the lung, and afterwards perforates its tissue, so that it is evacuated through the bronchial tubes.

LUNG PROOF, Docimasia pulmonum — *l.* Vesicles, see Pulmo — *l.* Wort, Pulmonaria — *l.* Wort, cow's, Verbascum nigrum — *l.* Wort, tree, Lichen pulmonarius.

LUNULA, (dim. of *luna*, 'the moon,') Onyx — *l.* Unguim, see Nail — *l.* Scapulæ, Notch, scapular.

LUNYIE (Sc.), Lumbi.

LUNYIEBANE, Ischion.

LUPIA, (from *λυπω*, 'to harass,') Encystis, Wen — *l.* Junctura, Spina ventosa.

LUPINUS. Under this term the *white lupin*, *Lupinus albus* seu *sativus*, (F.) *Lupin blanc*, *Ord.* Leguminosæ, is often meant. The seeds, which were much eaten in the days of Pliny and of Galen, are now neglected. The meal is, however, occasionally used as an anthelmintic, and as a cataplasm.

LUPOID, *Lupoïdes*, (*lupus*, and *ειδος*, 'resemblance,') Resembling, or having the character of, lupus.

LUPULINA, *Lu'pulin*, *Lupulin'ic glands*. A substance which exists in *Humulus lupulus* or hop. It is in the form of small, shining, yellowish grains, which cover the base of the scales of the hop; is pulverulent, and of an aromatic odour. When analyzed, it is found to contain resin, volatile oil in small quantity, and a bitter principle — *Lu'pulite*. It is aromatic and tonic, and — according to some — narcotic.

LUPULINIC GLANDS, *Lupulina*.

LUPULITE, see *Lupulina*.

LUPULUS, *Humulus lupulus* — *l.* Communis, *Humulus lupulus* — *l.* Salictarius, *Humulus lupulus* — *l.* Scandens, *Humulus lupulus*.

LUPUS, 'a wolf.' (So named from its rapacity.) *Lu'pus exedens* seu *vox'ra* seu *dev'orans*, *Ja'cob's Ul'cer*, *Ul'cus Tuberculo'sum*, *Can'cer* seu *Phymato'sis lu'pus*, *No'li me tang'ere*. (F.) *Loup*. Tubercular excrescences, with ragged, spreading ulcerations, chiefly about the *ala nasi*, where they destroy the skin, &c., for some depth. Sometimes they appear in the cheek, circularly, or in the shape of a sort of ringworm, destroying the substance, and leaving a deep and deformed cicatrix. The knife or caustic should be used to separate the sound from the morbid parts. Arsenic has been given internally with advantage. See, also, *Herpes exedens*, and *Lycoides*.

LUPUS CANCRO'SUS, Cancer — *l.* Devorans, see

Lupus — *l.* Exedens, *Lupus* — *l.* Varicosus, *Nævus* — *l.* Vorax, *Herpes exedens*, see *Lupus*.

LURCO, Glutton.

LURID, *Lu'idus*. Pale-yellow, sallow, — applied to the complexion.

LURKEYDISH, *Mentha pulegium*.

LUROR, (*λαρος*, 'pale,') Paleness.

LUSCIOTAS, *Luscitas*, *Myopia*.

LUSCIOSUS, *Borgne*, *Myops*.

LUSCITAS, *Luscio'sitas*, *Luscit'ies*, *Luscit'ion*, (*Lusciosis*, 'dim-sighted,') Strabismus. The term has also been given to any case of obliquity in which the eye is fixed in an unnatural position. The obliquity termed *Cockeye* is a form of *Luscitas*.

LUSCITIES, *Luscitas*.

LUSCUS, ('one eyed,') *Borgne*.

LUSS (Sc.), *Pityriasis*.

LUSTRAMENTUM, (*lustrare*, 'to purify,') Cathartico.

LUT, *Lute*.

LUTE, *Lu'tum*, 'mud,' *Cement'um*. (F.) *Lut*. A composition employed either for closing apertures in a pharmaceutical instrument, or for covering the surface of retorts, tubes, &c., which have to support a great degree of heat. Lutes are composed differently, according to the object to be accomplished. Commonly, they are made of linseed meal and starch. The *fat lute* is formed of clay and drying oil. Sometimes, the white of egg and lime are used; and that which is employed for covering vessels, intended to be strongly heated, is made of clay, sifted sand, and water.

LUTEOLA, (dim. of *luteus*,) *Reseda luteola*.

LUTEUS, Yellow.

LUTRON, (*λουτρον*, from *λουειν*, 'to wash,') A bath. Also, an ophthalmic medicine.

LUX, gen. *Lu'cis*, Light.

LUXATIO, *Luxation* — *l.* Imperfecta, Sprain.

LUXATION, *Lu'xat'ion*, (*luxare*, *luxatum*, 'to put out of place,') *Disloca'tion*, *Olisthe'ma*, *Luza'tio*, *Disloca'tio*, *Ec'clisis*, *Strem'ma*, *Luza'tura*, *Ectop'ia*, *Ecpto'sis*, *Elap'sus*, *Lygis'mus*, *Emo'tio*, *Ec'stasis*, *Exarthre'ma*, *Paracine'ma*, *E. luza'tio*, *Exarthro'ma*, *Exarthro'sis*, *Exo'sis*, *Exo'thesis*, *Paratop'ia*, *Streblo'sis*, *Deloca'tio*, *Exarticula'tio*, (F.) *Luxation*, *Dislocation*, *Déboîtement*. A displacement of a part from its proper situation. A putting out of joint. A displacement of two or more bones, whose articular surfaces have lost, wholly, or in part, their natural connexion; either owing to external violence, (*accident'al luxation*), or to disease of some of the parts about the joint (*spontaneous luxation*.) *Luxation* is *complete* when the bones have entirely lost their natural connexion; *incomplete*, — *Luxatio imperfecta*, *Sub-luxat'ion* — when they partly preserve it; and *compound*, when a wound communicates with the luxated joint. The general indications of treatment, are; — 1. To reduce the protruded bone to its original place. 2. To retain it *in situ*. 3. To obviate any attendant or consequent symptoms.

To reduce requires extension, counterextension, and coaptation.

LUXEUIL, MINERAL WATERS OF. Saline waters, at the town of Luxeuil, at the foot of the Vosges, in the department of Haute Saône. Five springs are thermal, and two cold. They seem to contain carbonic acid, carbonates of iron and lime, and chloride of sodium; and are employed as aperient, tonic, and stimulant.

LYCAN'CHE, *Lycan'chis*, (*lyco*, and *αγκη*, 'I strangle,') *Wolf quin'sy*, *Wolf choak*; — as *Cynanche* means *Dog choak*. Also, hydrophobia. See *Lycoides*.

LYCANCHIS, *Lycanche*.

LYCANTHROPE, see *Lycanthropia*.

LYCANTHROPIA, (*lyco*, and *ανθρωπος*, 'a man,') *Lyca'um*, *Insa'nia lupi'na*, *Lycoma'nia*. A

variety of melancholy, in which the person believes himself to be changed into a wolf, and imitates the voice and habits of that animal. One so circumstanced is called a *Ly'canthrope*, *Lycanthro'pus*.

LYCANTHROPUS, see *Lycanthropia*.

LYCAUM, *Lycanthropia*.

LYCHNIDIUM, (λυχνιδιον, dim. of λυχνος, 'a light,') *Biolychnium*.

LYCHNIS GITHA'GO, (λυχνις,) *Agrostem'ma githa'go*, *Corn cockle*, *Rose cam'pion*. A naturalized European plant. *Ord.* Caryophyllaceæ. The seeds are somewhat acrid, but not poisonous. They were, at one time, employed in cutaneous diseases, and in hemorrhage, and as a topical application to ulcers and fistule.

LYCHNIS OFFICINALIS, *Saponaria*—*L. Sylvestris*, *Saponaria*.

LYCHNIUM, (from λυχνος, 'a light,') *Biolychnium*.

LY'CIUM, (Λυκιον.) A medicinal substance, anciently obtained from the roots and branches of a thorny shrub growing in Lycia. It was astringent, and used in dysentery, ulcers of the gums, &c. By some, it has been considered to be the catechu.

LYCO, (from λυκος, 'a wolf,') in composition, wolf.

LYCODONTES, (*lyco*, and οδους, 'a tooth,') Canine teeth.

LYCOÏ'DES, *Lyco'des*, (*lyco*, and ειδος, 'form,') *Lupo sim'ilis*. An epithet for a species of cynanche, called, also, *Ly'pus* and *Strangulator*, and absurdly attributed to excessive retention of sperm in its reservoirs, and its passage into the blood [?].

LYCOMANIA, (*lyco* and mania,) *Lycanthropia*.

LYCOPE DE VIRGINIE, *Lycopus Virginicus*.

LYCOPER'DON, (*lyco*, and περδω, 'I break wind,') *L. Bovis'ta* seu *Arrhi'zon* seu *Globo'sum*, *Bovis'ta*, *Crep'itus Lupi*, *Fun'gus Chirurgo'rum*, *Puffball*, *Bullfists*, *Mol'tipuffs*, (Prov.) *Puff*, (Sc.) *Deil's Snuffbox*, (F.) *Vesseloop*. *Ord.* Fungi. The puff-ball dries into a very fine, light brownish dust, which is sometimes used as a mechanical styptic to arrest hemorrhage.

LYCOPERDON ARRIZON, *Lycoperdon*—*L. Cervinum*, *Elaphomyces granulatus*—*L. Globosum*, *Lycoperdon*—*L. Gulosorum*, *L. Tuber*.

LYCOPERDON PRO'TEUS, *Common puffball*. The fumes of this substance have been found to produce anesthetization in animals.

LYCOPERDON TU'BER, *L. Gulosorum*, *Tu'ber Agriocae'tanum* seu *Ciba'rium* seu *Guloso'rum* seu *Ni'grum*, *Tu'bera Ter'ræ*, *Truba*, *Truffle* (pronounced *troofle*), (F.) *Truffe*. A fleshy, firm body, unknown in its mode of reproduction, which is found under ground, especially in different parts of France, Italy, &c., and is much esteemed as an aliment. It seems to belong to the champignons, and has a particular perfume. Aphrodisiac virtues have been ascribed to it.

A Virginia truffle is called *Tuckaho'*. It is the *Sclero'tium gigante'um*, and is sometimes named *Indian bread* and *Indian loaf*.

LYCOPERSICIUM ESCULENTUM, *Solanum lycopersicum*—*L. Pomum amoris*, *Solanum lycopersicum*—*L. Tuberosum*, *Solanum tuberosum*.

LYCOPEDE, *Lycopodium*.

LYCOPO'DIUM, (*lyco*, and ποος, gen. ποδος, 'a foot,') *L. Clavatum* seu *Officina'le*, *Lepido'tis clavata*, *Cingula'ria*, *Mus'cus clavatus*, *Clubmoss*, *Wolf'sclaw*, *Witchmeal*. *Ord.* Lycopodiaceæ. This is the largest of the European mosses. The dust, or spores, *Lycopodium*, (Ph. U. S.), which fills the capsules of its spikes, is very inflammable, and hence has been called *Veg'etabile Sul'phur*;—

and, also, *Fari'na* seu *Pul'vis* seu *Se'men* seu *Sul'phur Lycopo'dii*, (F.) *Lycopode*, *Sonfre végétal*. It is used in France and in the United States to roll pills and boluses in; and in medicine is a desiccative in the excoriations to which infants are liable. It is collected chiefly in Switzerland and Germany for commerce.

LYCOPODIUM CLAVATUM, *Lycopodium*.

LYCOPODIUM COMPLANATUM, *Common Club-moss*, *Ground Pine*; indigenous; has the same properties as *Lycopodium selago*.

LYCOPODIUM OFFICINALE, *L. clavatum*—*L. Recurvum*, *L. Selago*.

LYCOPODIUM SELA'GO, *L. Recurvum*, *Plananthus fastigatus* seu *sela'go*, *Mus'cus erectus* seu *Cathar'ticus*, *Sela'go*, *Up'right Clubmoss*, *Fir Clubmoss*, *Up'right Firmoss*. A decoction of this plant acts violently as an emetic and purgative; and was formerly, on this account, used as an abortive. It is also employed to kill vermin.

LYCOPUS, *L. Virginicus* seu *uniflorus* seu *pumila*, *Bu'gleweed*, *Wa'ter Bu'gle*, *Wa'ter Horehound*, *Gypsyweed*, *Paul's Be'tony*, (F.) *Lycopoe de Virginie*, (*lyco*, and ποος, 'a foot,' from the likeness of the leaves.) This indigenous herb, *Sc. Syst.* *Diandria Monogynia*, *Nat. Ord.* Labiatae, grows throughout the greater part of the United States. The whole herb is said to be slightly narcotic. It is given in infusion, (*Lycop. 3j*); *Aq. fervent. Oj.* Dose, Oss to Oj.

LYCOPUS PUMILA, *Lycopus*.

LYCOPUS SINUA'TUS, *Wa'ter Horehound*, *Gypsy Weed*, *Paul's Be'tony*, indigenous, has similar properties.

LYCOPUS UNIFLORUS, *L. pumila*—*L. Virginicus*, *Lycopus*.

LYCOREXIS, (*lyco*, and ορεξις, 'appetite,') *Fames lupina*.

LYCOSA TARENTULA, see *Tarantula*.

LYCOSTOMA, (*lyco*, and στομα, 'mouth,') *Hyporöchasma*.

LYE, ([Sax.] leah, læs,) *Lixivium*, *Ley*—*L.* Dyspeptic, see *Fuligokali*—*L.* Medical, see *Fuligokali*—*L.* Tea, *Physick's*, see *Fuligokali*.

LYGISMUS, (λυγισμος, from λυγιζω, 'to bend,') *Luxation*.

LYGODES, (*lygnus*, and οδες,) *Singultus*.

LYGMUS, (λυγμος,) *Singultus*.

LYGODES, (λυγδης,) *Singultus*.

LYING-DOWN STATE, *Parturient state*.

LYING-IN STATE, *Parturient state*.

LYMA, *Lochia*, (λυω, 'to wash,') also, filth or sordes, (to be washed away,) *Ly'më*.

LYME, (λυμα,) *Lyma*.

LYMPH, *Lymph'a*, (akin to νυμφη, 'water,' by changing ν into λ.) *White blood*, *Lymph'a Nutrit'ia*, (F.) *Lymphe*. A name given to the fluid contained in the lymphatic vessels and thoracic duct of animals, which have been made to fast for 24 hours. According to Chevreul, the lymph of the dog contains water, fibrin, albumen, common salt, subcarbonate of soda, phosphates of lime and magnesia, and carbonate of lime. The properties and composition of lymph vary somewhat according to the part whence the lymphatic vessels obtain it. Generally, it is under the form of a transparent, slightly alkaline fluid, sometimes of a madder-red or yellowish colour,—of a spermiac odour, and saline taste; soluble in water,—the solution becoming turbid, when mixed with alcohol. When left to itself, it coagulates. The clot or solid portion becomes of a scarlet red, if put in contact with oxygen; and of a purple red, if placed in carbonic acid.

Like the blood, the lymph consists of a fluid—*Liquor lymph'a*—in which *lymph corpuscles* or *globules* are suspended.

The lymph is probably the product of internal absorption in different parts of the body: it then flows along the lymphatic vessels, uniting with

the chyle, and is poured with it into the veins; thus becoming one of the materials of the blood. According to others, the blood, when it reaches the arterial radicles, is divided into two portions,—the one red, which is carried to the heart,—the other serous or white, which is absorbed by the lymphatic vessels, and constitutes the lymph. By others, again, the lymphatics are considered to be the vessels of return for the white blood sent to certain tissues. White blood, however, has probably no existence.

The word *lymph* is sometimes used unhappily by the surgeon to signify liquor sanguinis.

LYMPH, COAGULABLE, Fibrin, Liquor sanguinis.

LYMPH CORPUSCLES, see Lymph—I. Globules, see Lymph—I. Hearts, Lymphatic hearts.

LYMPH, PLASTIC, Liquor sanguinis.

LYMPHA, Lymph—I. Arborum, see Sap—I. Muculentarium, Nasal mucus—I. Nutritia, Lymph—I. Pancreatis, see Pancreas—I. Pericardii, see Pericardium—I. Plastica, Fibrin.

LYMPHADENITIS, (*lymph* and *adenitis*.) *Inflammatio glandularum lymphaticarum*, *Adenitis lymphatica*, (F.) *Adénite lymphatique*. Inflammation of a lymphatic gland or ganglion.

LYMPHÆDUCTUS, (*lymp̄ha*, and *ductus*, 'a duct,') Lymphatic vessels.

LYMPHANGELITIS, (*lymphangia* and *itis*.) Angioloecitis.

LYMPHANGIA, (*lymp̄ha*, and *αγγειον*, 'a vessel,') Lymphatic vessels.

LYMPHANGIECTASIS, *Lymphenrysis*, (*lymphangia*, and *ectasis*, 'dilatation,') Dilatation of lymphatic vessels.

LYMPHANGIOGRAPHY, *Lymphangiographia*, (*lymphangia*, and *γραφία*, 'a description,') A description of the lymphatic vessels.

LAMPHANGIOITIS, (*lymphangia*, and *itis*.) Angioloecitis.

LYMPHANGIOLOGY, (*lymphangia*, and *λογος*, 'a description,') A treatise on the lymphatics.

LYMPHANGION, sing. of Lymphangia.

LYMPHANGIOTOMY, *Lymphangiotomy*, (from *lymphangia*, and *τομή*, 'incision,') The anatomy or dissection of the lymphatic vessels.

LYMPHANGITIS, (*lymphangia*, and *itis*.) Angioloecitis.

LYMPHANGIONCUS, *Lymphoncus*, (*lymp̄ha*, *αγγειον*, 'a vessel,' and *ογκος*, 'a tumour,') Tumefaction of the lymphatics.

LYMPHATIC, *Lymphaticus*, (F.) *Lymphatique*. That which relates to lymph; also, abounding in lymph. See Phlegmatic.

LYMPHATIC HEARTS, *Lymphhearts*. The frog and several other animals are provided with large receptacles for the lymph immediately underneath the skin, which exhibit distinct and regular pulsations like the sanguiferous heart. Their use appears to be to propel the lymph.

LYMPHATIC SYSTEM is that particular system of organs which is inservient to the formation and circulation of lymph. These organs are:

1. LYMPHATIC GANGLIONS OR GLANDS, see Conglobate.

2. LYMPHATIC VESSELS, *Lymphæductus*, *Lymphangia*, *Lymphangia*, *Venæ lymphaticæ*, *Ductus serosi*, *Vasæ lymphatica seu resorbentia seu hydragoga*, *Hydragoga*. These are very numerous. Arising at the surface of membranes and in the tissue of the organs, they carry into the veins the lymph from those parts. Lymphatic vessels are found in almost every part. Wherever they are met with, however, they form two orders,—one superficial, the other deep-seated;—the two orders frequently communicating with each other. Lymphatic vessels are generally smaller than arteries and veins. They are very thin, diaphanous, and cylindrical; but present, here and there, more or less considerable dilatations, caused by valves in their interior. They are slightly tortuous in

their course; their anastomoses are very numerous, and they often cross each other, forming successive plexuses. Of the arrangement of the extreme radicles we are ignorant. All the branches, before joining the principal trunks, pass through lymphatic ganglions, in which they are convoluted, or subdivided almost *ad infinitum*. They are formed of an outer areolar membrane and an internal coat, similar to that of the veins; of the latter, the valves are formed. All the lymphatics of the body ultimately discharge themselves at the junction of the subclavian and internal jugular veins. Two of these trunks are considerably larger than the others,—the *thoracic duct*, and the great trunk of the right side, (F.) *La grande veine lymphatique droite*. The former receives the lymphatics of the abdomen, of the lower extremities, the left side of the thorax, the left upper extremity, and the corresponding side of the head and neck; the latter receives those of the right upper extremity, and of the right side of the head, neck and thorax.

LYMPHATITIS, (*lymp̄ha*, and *itis*.) Angioloecitis.

LYMPHE, Lymph—I. Coagulable, Liquor sanguinis—I. de Cotugno, Cotunnus, liquor of—I. Plastica, Liquor sanguinis.

LYMPHEURYSMA, (*lymp̄ha*, and *συσμα*, 'dilatation,') Lymphangiectasis.

LYMPHITIS, (*lymp̄ha*, and *itis*.) Angioloecitis.

LYMPHIZATION. A term used by Professor Gross to signify effusion of coagulable lymph. It is not a happy word, inasmuch as lymph has another meaning, whilst the term 'coagulable lymph' is now almost abandoned.

LYMPHOCHEZIA, (*lymp̄ha*, and *χέζω*, 'I go to stool,') Serous diarrhoea.

LYMPHONCUS, (*lymp̄ha*, and *ογκος*, 'a tumour,') Lymphangione—I. Iridis, Iridaexesis.

LYMPHOPYRA, (*lymp̄ha*, and *πυρ*, 'fire, fever,') Fever, adenomenigeal.

LYMPHORRHAGIA, (*lymp̄ha*, and *ῥήγνμι*, 'I break forth,') Discharge of lymph from a lymphatic vessel, owing to a lesion of its coats.

LYMPHOSIS, (F.) *Lymphose*. The preparation or elaboration of lymph.

LYMPHOTOMY, (*lymp̄ha*, and *τέμνω*, 'I cut,') Dissection of the lymphatics.

LYNCH'S EMBROCA'TION. An emollient nostrum, formed of olive oil impregnated with bergamot and other essences, and coloured with alkannet root.

LYNGODES, (*Λυγγώδης*), Singultous—I. Febris, Fever, singultous.

LYNN WAHOO, *Ulmus alata*.

LYNX, (*Λυξ*, gen. *λυκκος*), Singultus.

LYPE, (*λυπη*, 'sadness,') Athymia.

LYPEMANIA, (*lype*, and *mania*,) Melancholy.

LYPERIA CROCEA, (from *λυπρος*, 'sad,' from the dull color of the flowers.) A South African plant, *Nat. Ord.* Scrophulariaceæ, the flowers of which closely resemble saffron in smell and taste, and possess similar medicinal virtues.

LYPEROPHRENIE, (*λυπρος*, 'sad,' and *φρεν*, 'the mind,') Melancholy.

LYPOTHY'MIA, (*lype*, and *θυμος*, 'the mind,') Very great sadness or despondency.

LYRA, (*λύρα*, 'the lyre,') *Psalter*, *Psalterium*, *Corpus Psalterii*, *Lyra Davidica*, *Lam'ina medullaris triangularis cerebri*, *Spatium trigonum*. The under surface of the posterior part of the body of the fornix is impressed with numerous transverse and oblique lines, which have been so called from some resemblance they bear to the ancient lyre.

LYSIA, Lysis.

LYSIMACHIA NUMMULARIA, (*λυσίμαχια*, from *λυσις*, 'solution,' and *μαχη*, 'strife,') *L. nemorum*, *Nummularia*, *Hirundinaria*, *Centimorbia*,

Concepting Loosestrife, Moneywort. (F. *Herb. auc. éus.* Ord. Primulaceæ. This plant was formerly accounted vulnerary. It has been considered to possess antiseptic and astringent qualities, and has been used in leucorrhœa.

LYSIMACHIA PTYCHÆA, *Lythrum salicaria*.

LYSIMACHIA QUADRIFOLIA, *Four-leaved Loosestrife, Crosswort*, an indigenous plant, is astringent; and has been used as a stomachic and antiperiodic.

LYSIS, (λύσις, from λύνω, 'I dissolve,') *Ly'sis*. Solution. A common suffix; also, a name given to a solution or termination of disease, which is operated insensibly; that is, gradually and without critical symptoms.

LYSSA, (λύσσα,) *Hydrophobia*—*L. Canina*, *Hydrophobia*.

LYSSAS, (λύσας,) *Maniodes*.

LYSSETER, (λύσσητρο, *Maniodes*.

LYSSODECTUS, (λύσσοδεκτος, from λύνω, and δέκω, or δάκνω, 'I bite,') *Hydrophobus; Cané rabido mor'sus*. One who has been bitten by a mad dog, or is actually labouring under hydrophobia.

LYSSODEG'MA, gen. *Lyssodeg'matis*, *Lyssodeg'ma*, *Lyssodeg'ma*. The bite of a mad dog.

LYSSODEXIS, *Lyssodegma*.

LYTERIOS, (λύτηριος, from λύνω, 'I dissolve,') 'solving.' An epithet given to those signs which announce the solution of a very violent disease.

LYTHRUM ALATUM, (from λυθρον, 'blood, and dust,' on account of the colour of the flowers of some species, see *Lythrum salicaria*—*L. Hyssopifolia*, *Lythrum salicaria*.

LYTHRUM SALICARIA, *Lythra'chia purpurea*, *Salica'ria vulgaris* seu *spicata*; — the *Common* or *Purple Willow Herb*. Ord. *Lythraceæ*. The herb, root, and flowers possess a considerable degree of astringency, and are used, occasionally, in the cure of diarrhœa and dysentery, leucorrhœa, hæmorrhæsis, &c.

Lythrum Hyssopifolia, and *L. Alatum*; indigenous, have similar properties.

LYTTA CERULEA, (from λυτταω, 'to be mad,' from its irritant properties [?], see *Cantharis*—*L. Gigas mas*, see *Cantharis*—*L. Ruficeps*, see *Cantharis*.

LYTTA RUPIFES. Class, *Insecta*; Ord. *Coleoptera*. A variety of *Lytta*, peculiar to Chili, which is more active as a vesicant than the *cantharis*, or the *Lytta vittata*.

LYTTA SEGETUM, see *Cantharis*—*L. Vesicatoria*, *Cantharis*.

LYTTA VITTA'TA, *Cantharis vittata*, *Pota'to Fly*, (F.) *Cantharide tachetie*. Four species of mead that blister are found in the United States. The *lytta vittata* feeds principally upon the potato plant, and, at the proper season of the year, may be collected in immense numbers. The potato fly resembles the *cantharides* in every property, and is fully equal to them.

M.

M. This letter signifies, in prescriptions, *manipulus*, 'a handful.' Also, *Misce*, 'mix.' See Abbreviation.

MAAGONI, *Swietenia Mahogani*.

MACAHALEF, see *Calaf*.

MACAPATLL, *Smilax sarsaparilla*.

MACARONI. An alimentary paste, moulded of a cylindrical shape, and formed of rice or wheat flour. It is eaten—when boiled—in soup; prepared with cheese, &c. See *Turundæ Italice*.

Also, a name formerly given to a pulverulent compound of sugar and glass of antimony, introduced into France by the Italian monks, and employed, at the hospital *La Charité*, in the treatment of painters' colic.

MACARONS, Biscuits, medicinal.

MACÉ, see *Myristica moschata*—*m. Reed*, *Typha latifolia*.

MACEDON'ICUM EMPLASTRUM, (μακεδονικόν, from *Macedonia*, *Macedonia*.) An ancient plaster referred to by *Aëtius*.

MACEDONISUM, (from *Macedonia*), *Smyrnum olusatrum*.

MACER, ('lean,') *Gracilis*.

MACERATION, *Macera'tio*, (*macerare*, *maceratum*, 'to soften by water,') An operation which consists in infusing, usually without heat, a solid substance in a liquid, so as to extract its virtues.

MACERA'TUM, (F.) *Macéré*. The product of maceration.

MACERONA, *Smyrnum olusatrum*.

MACESCERE, *Emaciate*.

MACHERA, (μαχαίρα,) *Culter*, *Knife*, *Penis*.

MECHERIDION, *Machæridion*.

MACHÆRION, *Machæ'ria*, *Machærid'ion*, *Gladi'olus*, (μαχαίρα, 'a sword; μάχη, 'a fight,') A knife. An amputating knife. *Rufus* of *Ephesus*

asserts, that the *Aruspices* gave this name to a part of the liver of animals.

MACHÆRIS, (μαχαίρις,) *Knife*, *Novacula*.

MACHA'ON, gen. *Machao'nis*, (Μαχάων.) The son of *Æsculapius*, and a celebrated physician.

MACHAÖ'NIA seu MACHAÖ'NICA ARS, (called after *Machaon*,) *Medicine*. The *Healing Art*.

MACH'INAL, *Mechan'icæ*, (μαχινά, 'a contrivance,') This epithet is added especially by French writers to the word *movement*, to express that the will takes no part in it.

MACHINE', *Mach'ina*, *Machinamen'tum*, *Mech'ani*, *Mechan'e'ma*, (same etymon.) A more or less compound instrument or arrangement for putting a body in motion, or for producing any action whatever. Physiologists sometimes use it for the *animal body*;—as the *machine* or *animal machine*.

MACHLOSINE, (μαχλοσίνη, from μαχλος, 'unchaste,') *Nymphomania*.

MACHOIRE, from [L.] *masticare*, 'to chew,') *Maxillary bone*—*m. Dinocraniæ*, *Maxillary bone*, lower—*m. Sphenocraniæ*, *Maxillary bone*, superior.

MACIES, *Atrophy*, *Emaciation*—*m. Infantum*, *Tabes mesenterica*.

MACILENTUS, (*macere*, 'to be lean,') *Gracilis*.

MACIS, (μακίς,) see *Myristica moschata*.

MACRAU'CHEN, gen. *Macrau'chenis*, (μακραυχόν, from *macro*, and *αυχόν*, neck,') *Longo collo præditus*. One who has a long neck.

MACRE FLOTTANTE, *Trapa natans*.

MACRITUDO, (*macere*, 'to be lean,') *Emaciation*.

MACRO, (from μακρός, 'long,') in composition, long.

MACROBIOSIS, (μακροβίωσις, from *macro*, and βίος, 'life,') Longevity.

MACROBIOTES, (μακροβίους,) Longevity.

MACROBIOTIC, *Macrobiot'icus*, *Macrobi'otus*, *Macro'bius*, *Long'eus*, *Macobiotic*. That which lives a long time. The *macrobiotic art*, *Macrobiot'ics*, (F.) *Macrobiotique*, is the art of living a long time.

MACROBIOTUS, (μακροβίος,) *Macriobiotic*.

MACROBIUS, (μακροβίος,) *Macrobiotic*.

MACROCEPHALIA, (*macro*, and κεφαλή, 'head,') see *Capitones*.

MACROCEPHALUS, (μακροκεφαλός,) *Macroceph'alus*. (F.) *Macrocéphale*; *Qui magnūm caput habet*. 'One who has a large head.' This epithet is given to children born with heads so large that they seem to be hydrocephalic; but in which the unusual development is owing to a large size of the brain. Such are supposed to be more than ordinary liable to convulsions. [?] The term has also been applied by Hippocrates to certain Asiatics who had long heads. See *Capitones*.

MACROCHIRIA, (*macro*, and χείρ, 'hand,') Monstrosity, characterized by excessive development of the hands.

MACROCO'LLIA, (μακροκόλλια, from *macro*, and κόλλω, 'a limb,') Great length of limbs in general, and of the lower limbs in particular.

MACROCOSM, *Macrocosm'us*, (*macro*, and κόσμος, 'world,') The universe, in contradistinction to *Microcosm*, which means man.

MACRODACTYLUS, (μακροδάκτυλος, from *macro*, and δάκτυλος, 'a finger,') Having long fingers.

MACROGASTER PLATYPUS, (*macro*, and γαστήρ, 'belly,') *Acarus folliculorum*.

MACROGLOSSA, (*macro*, and γλῶσσα, 'tongue,') The condition of a *Macroglossus*.

MACROGLOS'SUS. One who has a very large or prolapsed tongue. See *Paraglossa*.

MACROMEL'IA, (*macro*, and μέλος, 'limb,') Monstrosity, which consists in the excessive size of a limb.

MACRONOSIE, (μακρονοσία, from *macro*, and νόσος, 'disease,') Chronic diseases.

MACROPHAL'LUS, (*macro*, and φάλλος, 'the male organ,') A large size of the male organ.

MACROPHO'NUS, (*macro*, and φωνή, 'voice,') One who has a strong voice.

MACROPHTHAL'MUS, (*macro*, and οφθαλμος, 'eye,') Having large eyes.

MACROPHYSOCEPH'ALUS, (*macro*, φύσα, 'air,' and κεφαλή, 'head,') A word used by Ambrose Paré to designate an augmentation of the head of the fetus, produced by a sort of emphysema [?], which retards delivery.

MACROPIPER, (*macro*, and πείρι, 'pepper,') *Piper longum*.

MACROPNE'A, (μακροπνοία, from *macro*, and πνέω, 'I breathe,') A long and deep respiration.

MACROP'NUS, *Macrop'noius*, (μακροπνους, and μακροπνοός,) Long-breathed. One who breathes slowly:—a word met with in some authors.

MACROP'ODUS, *Mac'ropus*, *Me'pal'opus*, (μακροπους, from *macro*, and πους, gen. ποός, 'foot,') One who has a large foot.

MACROPROS'PIA, (*macro*, and προσωπον, 'face,') Monstrosity, characterized by excessive size of face.

MACROPUS, *Macropodus*.

MACROR, *Emaciation*.

MACRORRHIS, gen. *Macrorrh'nis*, (μακρορρις, from *macro*, and ρις, or ριν, 'nose,') One who has a long nose.

MACROS'CELES, *Orura longa habens*; (μακροσκελής, from *macro*, and σκελος, 'the leg,') One who has long legs.

MACROSIÆ, (μακρος, 'long,') Chronic diseases.

MACROSOMATIA, (*macro*, and σωμα, gen. σώματος, 'body,') see *Giant*.

MACROSOMIA, see *Giant*.

MACRO'TES, (*macro*, and οὖς, ὠτος, 'an ear,') One who has long ears.

MACROTYS RACEMOSA, (same etymon, from its capsules,) see *Actæa racemosa*.

MAC'ULA. A spot. *Dyschra'a*, *Ce'lis*, *La'bes*, (F.) *Tache*. A permanent discoloration of some portion of the skin, often with a change of its texture. *Ephelis*, *Nævus*, *Spilus*, &c., belong to *Macule*.

MACULA, *Molecule*—*m. Aurea*, see *Foramen centrale retinæ*—*m. Cornæ*, *Caligo*—*m. Cornæ arcuata*, *Gerotoxon*—*m. Cornæ margaritacea*, *Paralampsis*—*m. Cribrosa*, see *Auditory canal*, internal—*m. Emortalis*, see *Suggillation*—*m. Flava*, see *Foramen centrale retinæ*—*m. Fusca*, *Ephelides*—*m. Germinativa*, see *Molecule*—*m. Hepatica*, *Chloasma materna*, *Nævus*—*m. Lenticularis*, *Ephelides*—*m. Lutea retinæ*, see *Foramen centrale*—*m. Matricalis*, *Nævus*—*m. Matricis*, *Nævus*—*m. Nigra*, see *Optic nerve*—*m. Solaris*, *Ephelides*.

MAC'ULÆ AL'BÆ, (pl. of *macula*,) *M. Lac'tes*, *In'sule*, *Mil'kapots*. White spots, seen on serous membranes, especially on that investing the heart, and which appear to be the result of previous inflammatory action.

MACULÆ ANTE OCULOS VOLITANTES, *Metamorphopsia*—*m. Lactea*, *M. Albæ*.

MACULO'SUS, *Mac'ulose*, *Maculous*, *Mac'ulated*, (F) *Maculeux*. Spotted. Containing *maculæ* or spots.

MAD, ([Sax.] *gemæb*,) *Insane*—*m. Apple plant*, *Solanum melongena*—*m. Doctor*, *Psychiater*.

MADAR, *Mudar*.

MADARO'SIS, (μαδαρωσις, from μαδάω, 'to be bald,') *Madaro'ma*, *Made'sis*, *Madar'otes*, *Made'ma*, *Mad'isis*, *Deplumatio*, *Mil'phæ*, *Milpho'sis*, *Pti'los*, *Ptilo'sis*, *Calvit'ies*. Loss of the hair, particularly of the eyelashes.

MADAROTES, (μαδαρωσις,) *Madaros*.

MADDER, *DYERS*, ([Sax. *maðdere*,) *Rubia*.

MAD'IRA, *CLIMATE OF*. This island is much frequented by pulmonary invalids, on account of the mildness and equability of its climate. Owing, indeed, to the mildness of the winter, and the coolness of the summer, together with the remarkable equality of the temperature during day and night, as well as throughout the year, it has been considered that the climate of Madeira is the finest in the northern hemisphere. Sir James Clark is of opinion that there is no place on the continent of Europe, with which he is acquainted, where the pulmonary invalid could reside with so much advantage, during the whole year, as in Madeira.

MADELCON, (μαδέλκων,) *Bdellium*.

MADEMA, gen. *Made'matis*, (μαδέμμα, from μαδάω, 'to be bald,') *Madarosis*.

MADESIS, (μαδσις, from μαδάω, 'to be bald,') *Depilation*, *Madarosis*.

MADISIS (*madisis*), *Depilation*, *Madarosis*.

MADISON SPRINGS, see *Georgia*, *Mineral waters of*.

MADISTE'RIMUM, (μαδιστηριον, from μαδίζω, 'to make bald,') *Trichola'bium*, *Trichol'abis*, *Vol'sella*. Tweezers. An instrument for extracting hairs.

MADJOUND, *Madjoun*. A mixture of powdered hemp and honey, used as a narcotic agent, like tobacco, by the Algerines.

MADNESS, *Insanity*—*m. Canine*, *Hydrophobia*—*m. Raving or furious*, *Mania*.

MA'DOR, *Moisture*. A cold sweat.

MADORIUS, *Mudar*.

MADREPORA OCLATA, see *Coral*.

MADWEED, *Scutellaria lateriflora*.

MÆA (μαία), *Midwife*.

MÆEIA (*μαία*), Obstetrics.

MÆEUMA (*μαίωμα*), see Parturition.

MÆUTICA ARS. Obstetrics.

MÆIA (*μαία*), Obstetrics.

MAESA PICTA, Saoria.

MÆUSIS (*μαῖσις*), Parturition.

MÆUTRIA (*μαευρία*), Midwife.

MAGDAL'IA, (*μαγδαλία*, from *μαστόν*, 'to knead.') *Magdal'oon*, *Mag'dalis*. 'Crumb of bread.' Any medicine, as a pill, formed of crumb of bread. A roll of plaster.

MAGEIRICE, (*μαγειρικὴ τέχνη*.) Culinary art.

MAGGOT PIMPLE, see Acne.

MA'GIS (*μαῖς*, 'a cake.') A sort of cake, composed of cloves, garlic and cheese, beaten together.

MAGISTERIUM, Magistery—m. Bismuthi, Bismuth, subnitrate of—m. Jalapæ, Resin of Jalap—m. Marcasite, Bismuth, subnitrate of—m. Plumbi, Plumbi subcarbonas—m. Sulphuris, Sulphur lotum—m. Tartari purgans, Potassæ acetas.

MAG'ISTERY, *Magiste'rium*, (*magister*, 'a master.') Certain precipitates from saline solutions were formerly so called; as well as other medicines, the preparation of which was kept secret.

MAGISTERY OF BISMUTH, Bismuth, subnitrate of.

MAG'ISTRAL or MAGIS'TRAL, *Magistra'lis*. *Extempora'neous*. A prescription is so called—*Form'ula magistra'lis seu extempora'nea*—which is prepared extemporaneously; an *offic'inal* prescription—*Form'ula officina'lis*—being such as has been prepared for some time before it is prescribed.

MAGISTRANTIA, (*magistrare*, 'to rule.') Imperatoria.

MAG'MA, gen. *Mag'matis*, (*μαγμα*.) (F.) *Marc*. The thick residuum, obtained after expressing certain substances to extract the fluid parts from them. The grounds which remain after treating a substance with water, alcohol, or any other menstruum. Also, a salve of a certain consistence.

MAGMA OF MARC OF OLIVES is the residuum after the greatest possible quantity of oil has been extracted from olives by making them ferment. It was formerly employed as a stimulant, under the form of a *bath*—to which the name *Bain de Marc* was given by the French.

MAGMA OF MARC OF GRAPES, *Bry'tia*, was once employed for the same purposes.

MAGMA RÉTICULÉ, 'reticulated magma.' The gelatiniform substance found between the chorion and amnion in the early period of embryonic existence.

MAGNES, gen. *Magne'tis*, (*μαγνης*.) *Magnet*.

MAGNES ARSENICA'LIS. (*Sulphur*, *White arsenic*, and *Common antimony*, of each equal parts. Mix by fusion.) It is corrosive. See *Magnetic plaster*.

MAGNES EPILEPSIÆ, Hydrargyri sulphuretum rubrum.

MAGNE'SIA, *Abarna'as*, *Cham'bar*, *Ter'ra ama'ra*, *Magnesia ter'ra*, *Talc earth*; (from *magnes*, or from *Magnesia* in *Lydia*.) An alkaline earth whose metallic base is *magnesium*. The term *Magnesia* was employed, of old, for any substance that seemed to have the power of attracting some principle from the air. At present it is restricted to—

MAGNESIA. *M. usta seu calcina'ta*, *Cal'cined Magnesia*, *Ox'ide of magnesi'um*, (F.) *Magnésie brûlée*, *Magnésie*, *M. Caustique*. (This is obtained by exposing carbonate of *magnesia* to a strong heat, to drive off the carbonic acid.) It is inodorous taste very slightly bitter; in the form of a

white, light, spongy, soft powder. S. g. 2-3; requiring 2000 times its weight of water for its solution. It is antacid, and laxative when it meets with acid in the stomach. Dose, gr. x to ʒi in water or milk.

Henry's, *Husband's*, and *Ellis's Magnesia*, are dense forms of this preparation.

A light *Magnesia*, *Magne'sia le'vis*, differing from the above only in its greater levity, is officinal in the British Pharmacopœia (1864).

MAGNESIA, ACETATE OF, *Magnesia acetas*—m. *Aerata*, *Magnesia carbonas*—m. *Alba*, *M. carbonas*—m. *Calcinata*, *M. usta*—m. *Citrate* of, *Magnesia citras*—m. *Dinneford's*, see *Magnesia*, fluid—m. *Edinburgensis*, *M. carbonas*.

MAGNESIA, EFFERVES'CENT, *Mox'on's*. (*Magnes. carb.*; *M. Sulphat.*; *Soda bicarbon.*, *Acid tartaric*. *aa partes æquales*; to be pulverized, well dried, mixed, and enclosed in bottles hermetically sealed.) Dose, a teaspoonful in half a tumbler of water, drunk in a state of effervescence.

MAGNESIA, ELLIS's, see *Magnesia*.

MAGNESIA, FLU'ID. Under this name a preparation is designated, which consists of a solution of carbonate of *magnesia* in carbonated water. It is also termed *carbonated magnesia water*, *aerated magnesia water*, and *condensed solution of magnesia*, (F.) *Eau magnésienne*. *Dinneford's Magnesia* is a solution of this kind.

MAGNESIA, HENRY'S, *Magnesia*—m. *Husband's* see *Magnesia*—m. *Mitis*, *Magnesia carbonas*—m. *Muriate* of, *Magnesi chloridum*—m. *Nigra*, *Manganese*, black oxide of.

MAGNESIA OPAL'INA. A name given by Léméry to a mixture of equal parts of *antimony*, *nitrate of potassa*, and *chloride of sodium*, decrepitated. It has emetic properties, but is not used.

MAGNESIA SALIS AMARI, *Magnesia carbonas*—m. *Salis Ebsdamensis*, *Magnesia carbonas*—m. *Saturni*, *Antimonium*—m. *Solution* of, *condensed*, *M. fluid*—m. *Subcarbonate* of, *Magnesia carbonas*—m. *Subcarbonate* of, *hydrated*, *Magnesia carbonas*—m. *Terra*, *Magnesia*—m. *Vitriolata*, *Magnesia sulphas*—m. *Usta*, *Magnesia*—m. and *Soda*, *sulphate* of, see *Soda*, *sulphate* of.

MAGNE'SIÆ ACET'AS, *Ac'etate of Magnesia*. (This salt is prepared by saturating carbonate of *magnesia* with *acetic acid*.) It is of syrupy consistence, and about four ounces is a dose. It acts like citrate of *magnesia*.

MAGNE'SIÆ CARBO'NAS seu *Subcarbo'nas*, *Magne'sia* (Dublin), *M. aëra'ta seu carbo'nica seu al'ba*, *M. Subcarbo'nica seu Mit'is seu Edinburgen'sis seu Sa'lis Ebsdamen'sis seu Salis ama'ri*, *Carbo'nas magnesi'um*, *Lac ter're*, *Hypocarb'onas magnesi'æ*, *Terra ama'ra aëra'ta seu absor'bens mine'ra'lis seu talco'sa oxyanthrac'o'des*, *Subcar'bonate of Magnesia*, *Hy'drated Subcar'bonate of Magnesia* (F.) *Sous-carbonate ou Carbonate de Magnésie*, *Magnésie aérée ou blanche ou crayeuse ou douce ou effervescente ou moyenne ou Anglaise*, *Cràie ou Terre magnésienne*, *Lait de Terre*, *Panacée Anglaise*, *Poudre de Sentinelli ou de Valentini ou du Comte de Palme ou de Zwinger*. (Prepared from sulphate of *magnesia* by subcarbonate of potassa.) It is inodorous; insipid; light; white; spongy; opaque; effervescing with acids; insoluble in water. Properties the same as the last; but the carbonic acid, when set free, sometimes causes unpleasant gastric distension.

A heavy and a light carbonate of *magnesia*—*Magne'siæ Carbo'nas* (*M. C. Ponderosum*, Ph. D.) and *Magne'siæ carbo'nas le'vis* are officinal in the British Pharmacopœia (1864).

MAGNE'SIÆ CITRAS, *Cit'rate of Magnesia*, (F.) *Citrate de Magnésie*. A saline preparation, formed by saturating a solution of *citric acid* with either *magnesia* or its carbonate. Dose, an ounce. It

is devoid of the bitter taste of the magnesian salts.

A solution in water, or in mineral water, sweetened with syrup, and acidulated with citric acid, makes an agreeable purgative. A simple solution in water has been called *magnesian lemonade*. In the effervescent state, it is the *effervescent magnesian lemonade*. The Pharmacopœia of the United States (1851) has a form for the *Liquor Magnesiæ Citratæ*, *Solution of Citrate of Magnesia*. (Carbonate of magnesia, $\mathfrak{z}\mathfrak{v}$; Citric acid, $\mathfrak{z}\mathfrak{vi}\mathfrak{ss}$; Syrup of citric acid, $\mathfrak{f}\mathfrak{z}\mathfrak{i}\mathfrak{j}$; Water, a sufficient quantity. Dissolve the citric acid in $\mathfrak{f}\mathfrak{z}\mathfrak{i}\mathfrak{v}$ of water, and add $\mathfrak{z}\mathfrak{i}\mathfrak{v}$ of the carbonate of magnesia, previously rubbed with $\mathfrak{f}\mathfrak{z}\mathfrak{i}\mathfrak{j}$ of water. When the reaction has ceased, filter into a strong $\mathfrak{f}\mathfrak{z}\mathfrak{x}\mathfrak{i}\mathfrak{j}$ glass bottle, into which the syrup of citric acid has been previously introduced. Rub the remaining carbonate of magnesia with $\mathfrak{f}\mathfrak{z}\mathfrak{i}\mathfrak{j}$ of water, and pour the mixture into the bottle, which must be well corked, and secured with twine; and shake the mixture occasionally until it becomes transparent. A modified form is introduced into the Ph. U. S. for 1863.) As a cathartic the whole of this may be taken.

MAGNESIÆ HYPOCARBONAS, *M. carbonas*—*m. Subcarbonas*, *M. carbonas*.

MAGNESIÆ SULPHAS, *Sulphas Magnesiæ purificata*, *Magnesia vitriolata*, *Sal catharticus amarus* seu *catharticum amarum* seu *amarum* seu *Anglicanum* seu *An'glicum* seu *Epsomensis* seu *catharticus Anglicanus* seu *Sedlitzensis* seu *Ebsdamsen'se* seu *Seydschützen'se*, *Ten'ra amara sulphurica*, *Vitriolicum Magnesiæ*, *Sulphate of Magnesia*, *Ep'som Salt*, *Bit'ter pur'g'ing Salt*, (F.) *Sulfate de magnésie*, *Sel admirable de Lénéry*, *Sel d'Égra*. (Generally obtained from seawater.) Its taste is bitter and disagreeable. It is soluble in an equal quantity of water at 60°. It is purgative and diuretic. Dose, as a cathartic, $\mathfrak{z}\mathfrak{ss}$ to $\mathfrak{z}\mathfrak{i}\mathfrak{j}$.

MAGNESIÆ VITROLICUM, *Magnesiæ sulphas*.

MAGNESIAN LEMONADE, *Magnesiæ citras*.

MAGNÉSIE AÉRÉE, *Magnesiæ carbonas*—*m. Anglaise*, *Magnesiæ carbonas*—*m. Blanche*, *Magnesiæ carbonas*—*m. Brûlée*, *Magnesia usta*—*m. Carbonate de*, *Magnesiæ carbonas*—*m. Cautique*, *Magnesia usta*—*m. Citrate de*, *Magnesiæ citras*—*m. Crayeuse*, *Magnesiæ carbonas*—*m. Douce*, *Magnesiæ carbonas*—*m. Effervescente*, *Magnesiæ carbonas*—*m. Moyenne*, *Magnesiæ carbonas*—*m. Souscarbonate de*, *Magnesiæ carbonas*—*m. Sulfate de*, *Magnesiæ sulphas*.

MAGNE'SII CHLO'RIDUM, *Chlo'ride of Magnesium*, *Mu'riate of Magnesia*. This bitter deliquescent salt has been given as a mild and effective cholagogue cathartic, in the dose of half an ounce to the adult. Being deliquescent, it may be kept dissolved in its weight of water.

MAGNESIUM, see *Magnesia*—*m. Chloride of*, *Magnesium chloridum*—*m. Oxide of*, *Magnesia usta*.

MAGNET, *Magn'es*, *Magne'tes*, *Ma'gn'etis*, *Fer'rum Magn'es attractivum*, *Sider'ites*, *Sider'itis*, *La'pis heracle'us seu Syder'itis seu nauticus*, *Ad'aman't*, (from *Magnes*, its discoverer, or from *Magnesia*, where it is said to have been first found.) (F.) *Aimant*, *Pierre d'Aimant*. The magnet or loadstone. An amorphous, oxydulated ore of iron, which exerts an attraction on unmagnetized iron, and has the property of exhibiting poles; that is, of pointing by one of its extremities to the north. This ore, by constant or long rubbing, communicates its properties to iron; and thus artificial magnets are formed. Magnetic ore is found in many countries, and particularly in the island of Elba. The magnet is sometimes used to extract spicula of iron from the eye or from wounds. It has been employed as an antispasmodic; but acts only through the imagination. The powder has been given as a tonic. In

Pharmacy, it is used to purify iron filings. It attracts the iron, and the impurities remain behind. It formerly entered, as an ingredient, into several plasters, to draw bullets and heads of arrows from the body—as the *Emplast'rum divinum Nicola'i*, the *Emplast'rum nigrum* of Augsburg, the *Opodel'doch*, and *Attractivum* of Paracelsus, &c.

MAGNETES (μαγνητις), Magnet.

MAGNET'IC, *Magnet'icus*, (F.) *Magnétique*. That which belongs or relates to magnetism;—mineral or animal.

MAGNETIC FLU'ID. A name given to the imponderable fluid to which the magnet owes its virtues. By analogy it is applied to a particular principle, supposed to be the source of organic actions, which affects, it is conceived, the nervous system principally, and is susceptible of being transmitted from one living body to another, by contact or simple approximation, and especially under the influence of fixed volition. See *Magnetism*, *Animal*.

MAGNETIC PLAS'TER. A plaster, at present, not used. It has for its base a mixture, called *Magn'es arsenical'is*; formed of equal parts of antimony, sulphur and arsenic melted together in a glass cucurbit. The name *Magnetic plaster* was, likewise, given to one that contained powdered magnet.

MAGNETINUS, *Potassæ supertartras impurus*.

MAGNÉTISEUR, Magnetizer.

MAGNETISM, AN'IMAL, *Mes'merism*, *Path'etism*, *Psy'cheism*, *Neuroga'mia*, *Bioga'mia*, *Bio-magnetis'mus*, *Zoömagnetis'mus*, *Exon'eurism* (proposed by Mr. H. Mayo), *Anthropomagnetis'mus*, *Gar'gate*, *Gargalis'mus*, *Gar'galus*. Properties attributed to the influence of a particular principle, which has been compared to that which characterizes the magnet. It is supposed to be transmitted from one person to another, and to impress peculiar modifications on organic action, especially on that of the nerves. The discussions, to which this strange belief has given rise, are by no means terminated. There is no evidence whatever of the existence of such a fluid. Highly impressible persons can be thrown into a kind of hysterical or 'magnetic' sleep and somnambulism, designated by Mr. Braid, *hyp'notism*, *neurohyp'notism*, and *nervous sleep*, and at times called, after him, *Braid'ism*; but farther than this, the efforts of the magnetizer cannot reach. It is a mode of action upon the nerves through the medium of the senses.

MAGNETIZATION, Mesmerization.

MAGNETIZED, Mesmerized.

MAGNETIZER, Mesmerizer.

MAGNETICOÉLECTRICITY, see *Electro-magnetism*.

MAGNETIS (μαγνητις), Magnet.

MAGNITUDO CORPORIS, ('size of the body,')

Stature—*m. Gigantea*, see *Giant*.

MAGNOC, *Jatropha manihot*.

MAGNOLIA FRAGRANS, (after P. Magnol, a Professor of Botany at Montpellier,) *M. glauca*.

MAGNO'LIA GLAU'CA, *Magnolia*, *M. fra'grans*, *Small Magnolia*, *Swamp Sas'safras*, *Elk Bark*, *Ind'ian Bark*, *White Lau'rel*, *Sweet Bay*, *Beav'er Wood*, *Beav'er Tree*, *White Bay*, *Cincho'na of Virginia*, *Cas'ton Bark*, *Sweet or Lau'rel magnolia*, (F.) *Magnolier glauque*, *Ord. Magnoliaceæ*. The bark is possessed of tonic properties, resembling those of casarilla, canella, &c. The same may be said of the *Magnolia tripet'ala* seu *Umbrel'la*, *Umbrel'la tree*; the *M. acuminata* or *Cucumber tree*, the *M. grandiflora* and *M. macrophylla*, *Lau'rel*, *Elkwood*, *Silverleaf*, *Big leaf*, *White Bay*, *Beavertree*, *Elk Bark*, *Big bloom*.

Magnolia (Ph. U. S.), is the bark of *Magnolia glauca*, *M. acuminata*, and *M. tripet'ala*.

MAGNOLIER GLAUQUE, *Magnolia glauca*.

MAGNUM DEI DONUM, ('great gift of God,') *Cinchona*.

MAGNUM OS, ('great bone,') The third bone of the lower row of the carpus, reckoning from the thumb. It is the largest bone of the carpus; and is, also, called *Os capitulum*; (F.) *Grand Os*.

MAG'RUMS. A popular name in the State of New York for a singular convulsive affection, which resembles chorea. It rarely, however, occurs before the adult age; never ceases spontaneously, and when fully developed, is devoid of any paroxysmal character.

MAGUEY, *Agave Americana*.

MAHAGONI, *Swietenia mahogani*.

MAHMOUDY, *Convolvulus scammonia*.

MAHOGAN-FEBRIFUGE, *Swietenia febrifuga*.

MAHOGANY, *Gymnocladus Canadensis*, *Swietenia mahogani*. Also, a drink in use among the Cornish fishermen of England, which is made of two parts of gin and one part treacle, well beaten together. An analogous drink—a mixture of whisky and honey, (see Mead, under Hydromel,) is known in the Highlands of Scotland by the name *Ath'ol por'ridge*.

MAHOGANY, MOUNTAIN, *Betula lenta*.

MAHOGANY TREE, *Swietenia mahogani*.

MAHURA, *Covolum*.

MAIANTHEMUM, (*Maius*, May, and *ανθεμον*, 'a flower,') *Convallaria maialis*.

MAIDENHAIR, *Adiantum capillus veneris*—*m. American*, *Adiantum pedatum*—*m. Canada*, *Adiantum pedatum*—*m. Common*, *Asplenium trichomanoides*—*m. Golden*, *Polytrichum*—*m. White*, *Asplenium ruta muraria*.

MAIDENHEAD, *Virginity*.

MAIDENHOOD, *Virginity*.

MAIEIA, (*μαία*,) *Obstetrics*.

MAIEUMA, gen. *Μαίευματ'is*, (*μαίευμα*,) see Parturition.

MAIEUSIS, (*μαίευσις*,) Parturition.

MAIEUTER, (*μαίευτρος*,) *Accoucheur*.

MAIEUTES, (*μαίευτρος*,) *Accoucheur*.

MAIEUTICA ARS, (*μαίευτική τέχνη*, 'art of midwifery,') *Obstetrics*.

MAIEUTICAL, *Obstetrical*.

MAIGREUR, (from [L.] *macere*, 'lean,') *Ema-ciation*.

MAIHEM, *Mayhem*.

MAILLET, (from *malleus*,) *Mallet*.

MAILLOT, *Swathing clothes*.

MAIMING, *Mayhem*.

MA'N, *Manus*—*m. Bot*, see *Kyllosis*.

MAINS. An instrument introduced by Palfin into the art of Obstetrics, and which is considered to have led to the invention of the forceps. It consisted of two spoons (*cuillères*) without fenestræ, mounted on wooden handles, and the branches of which were united by means of a steel *bride*.

MAIRANIA UVA URSI, (*μαίρω*, 'to shine,')—the leaves,) *Arbutus urva ursi*.

MA'IS, *Zea mays*.

MAISON PETITE, ('small house,') (from [L.] *manere*, *mansum*, 'to remain,') *Morotrophum*.

MAISON-DE-SANTE, ('health house,') A private establishment, usually founded by a physician, into which cases of disease are received for treatment.

MAIZENA, see *Zea mays*.

MAJOR HEL'ICIS, ('larger muscle of the helix,') A narrow band of muscular fibres situate upon the anterior border of the helix of the ear, just above the tragus.

MAJORANA, *Origanum majorana*—*m. Hortensis*, *Origanum majorana*—*m. Syriaca*, *Teu-erium marum*.

MAKROCEPHALUS, *Macrocephalus*.

MAL (F.), *Malum*, ('an evil, injury,') Any

pain or suffering, disease or infirmity, which interferences with the healthy play of the functions.

MAL D'AMOUR, *Odontalgia*—*m. Anglais*, *Mal de la Baie de Saint Paul*.

MAL DES ARDENS, *Mal d'Enfer*, *Feu sacré*. A name given to a species of pestilential erysipelas or *Saint Anthony's fire*, which reigned epidemically in France, in 1130.

MAL D'AVENTURE, *Paronychia*.

MAL DE LA BAIE DE SAINT PAUL, ('disease of the Bay of St. Paul,') *Mal Anglais*, *Mal de Chien*, *Mal de Chicot*, *Mal des Éboulements*, *O'toway disease*. A syphilitic affection, which prevailed in Canada from 1760 to 1780 and upwards.

MAL DE BASSINE, *Mal de Vers*—*m. Caduc*, *Epilepsy*—*m. de Cerf*, *Tetanus*—*m. de Chicot*, *Mal de la Baie de Saint Paul*—*m. de Chien*, *Mal de la Baie de Saint Paul*.

MAL DE CRIMÉE (F.), *Lèpre des Cosaques*, *Lepra Tau'rica*. A variety of lepra in the Crimea.

MAL DE DENT, *Odontalgia*—*m. Divin*, *Epilepsy*—*m. des Éboulements*, *Mal de la Baie de Saint Paul*—*m. d'Enfant*, *Pain (Labour)*—*m. d'Enfer*, *Mal des Ardens*—*m. d'Estomac*, *Chthonophagia*—*m. Française*, *Syphilis*—*m. de Gorge*, *Cynanche*—*m. Grand*, see *Epilepsy*—*m. Haut*, *Epilepsy*.

MAL DE LAIRA, *Bark'ing disease*. An epidemic hysteroid affection, a form of theomania, which prevailed in several convents of Germany and elsewhere, in the seventeenth century.

MAL DE MACHOIRE, *Trismus*—*m. de Mer*, *Nausea marina*—*m. de Mère*, *Hysteria*—*m. de Misère*, *Pellagra*—*m. de Montagne*, *Puna*—*m. de Naples*, *Syphilis*—*m. Petit*, see *Epilepsy*—*m. des Pieds et des mains*, *Acrodynia*—*m. Régionale*, *Endemy*—*m. du Roi*, *Scrofala*—*m. de la Rosa*, *Pellagra*—*m. Rouge de Cayenne*, *Elephantiasis of Cayenne*—*m. Saint Antoine*, *Erysipelas*—*m. de Saint Eutrope*, *Hydrops*—*m. de Saint Hubert*, *Hydrophobia*—*m. Saint Jean*, *Epilepsy*—*m. Saint Main*, *Lepra*, *Psora*—*m. de Saint Médard*, *Odontalgia*—*m. de Siam*, *Fever*, *yellow*.

MAL DE SAN LAZARO, *Cocobay*. A leprous disease, common in Colombia, S. America.

MAL DE SEPT JOURS, ('disease of seven days,') see *Trismus*—*m. de Sologne*, *Ergotism*—*m. de Terre*, *Epilepsy*—*m. à Tête*, *Cephalalgia*—*m. del Valle*, *Proctocæce*—*m. Vat*, see *Anthrax*.

MAL DE VERS, ('disease of worms,') *Mal de Bassine*. A vesico-pustular eruption on the fingers, and on the back and palm of the hand, observed in women engaged in the filature of silk.

MALA, (*malum*, 'an apple,') *Gena*, *Melum*—*m. Aurea*, see *Citrus aurantium*.

MALABATH'RINUM, (*μαλαβαθρινον*,) Ancient name of an ointment and a wine, into which the *malobathrum* entered.

MALABA'THRUM, (*μαλαβαθρον*,) *Malabathrum*, *Cadeji-Indi*. The leaves of a tree of the East Indies. These leaves entered into the theriac, mithridate, and other ancient electuaries. They are believed to be from a species of laurel—*Laurus Cassia*; but, according to others, from *Laurus Malabathrum*. The *O'leum Malaba'thri* is obtained from it.

MALACCÆ RADIX, ('Malacca root,') *Sagittarium alexipharmacum*.

MALACHE, (*μαλαχία*,) *Malva rotundifolia*.

MALA'CIA, (from *μαλακός*, 'effeminate,') A depravation of taste, in which an almost universal loathing is combined with an exclusive longing for some particular article of food. If the patient desires substances that are noxious or not eatable, it constitutes *Pi'ca*, *Heterorexia*, *Heterorrhexia*, *Cis'sa*, *Citto'sis*, *Cit'ta*, *Limo'sis*, *Pice*,

Allotriophagia, *Dopra'vèd ap'petite*, (F.) *Envie*. These symptoms accompany several nervous affections, — those of females in particular. In pregnancy it is common, and is termed *Long'ing*.

MALACIA AFRICANORUM, *Chthonophagia* — m. Cordis, *Cardiomalacia*.

MALACISMUS, (from *μαλακός*, 'soft,') Mollities. MALACO, (from *μαλακός*, 'soft,') in composition, soft.

MALACOGASTER, (*malaco*, and *γαστήρ*, 'stomach,') *Gastromalacia*.

MALACOMA, (from *μαλακός*, 'soft,' and the suffix *oma*,) Mollities.

MALACOPHONUS, (*malaco*, and *φωνή*, 'voice,') One who has a soft voice.

MALACOPŒA, (*malaco*, and *ποιεῖν*, 'to make,') Emollients.

MALACOSARCO'SIS, (*malaco*, and *σαρξ*, *carne*, 'flesh,') State of softness of the muscular system.

MALACOSIS, (*μαλακός*, 'soft,') Mollities — m. Cerebri, Mollities cerebri — m. Cordis, *Cardiomalacia* — m. Hepatis, *Hepatomalacia* — m. Uteri, *Hysteromalacia*.

MALACOSTEON, (*malaco*, and *οστέον*, 'a bone,') Mollities ossium.

MALACOSTEOSIS, Mollities ossium.

MALACTICA, (*μαλακτικά*, from *μαλακός*, 'soft,') Emollients, Relaxants.

MALADE, (from [L.] *malum*, [F.] *mal*,) Sick.

MALADIE, Disease — m. Anglæ, Hypochondriasis — m. de Basedow, see *Exophthalmia* — m. Bleue, Cyanopathy — m. de Bright, Kidney, Bright's disease of the — m. Bronzé, see Capsule, renal — m. Bruyante, *Streptopus morbus* — m. Bulleuse, see Bullous, Pemphigus — m. de Cruveilhier, see Brash, weaning — m. Cuculaire, Pertussis — m. Glandulaire, Elephantiasis Arabica — m. Imaginaire, Hypochondriasis, Hysteria — m. Mercurielle, Hydrargyriasis.

MALADIE DES MINEURS, Anæmia occurring in the workers in mines.

MALADIE NOIRE, Melæna — m. du Pays, Nostalgia — m. Pédiculaire, Phtheiriæsis — m. de Plomb, Saturnismus — m. de Pott, Vertebral disease — m. Typhoïde, see Typhus — m. Vermineuse, Syphilis — m. de Vénus, Syphilis — m. Vénérienne, Helminthiasis — m. de Werlhof, Purpura hæmorrhagica.

MALADIES ACQUISES, Acquired diseases — m. Annuelles, Annual diseases — m. Chroniques, Chronic diseases — m. Connées, Connate diseases — m. Dissimulées, Feigned diseases — m. Extérieures, External diseases — m. Feintes, Feigned diseases — m. Hérititaires, Hereditary diseases — m. Intimes, Innate diseases — m. Légitimes, see Legitimate — m. Nerveuses, Nervous diseases — m. Religieuses, Religious diseases — m. Simulées, Feigned diseases — m. Sporadiques, see Sporadic — m. Supposées, Feigned diseases — m. Ventesuses, Pneumatosis.

MALADIF, Sickly.

MALADRERIE, (from *mal*, and *ladrerie*,) *Ladverie*.

MALADY, ENGLISH, Hypochondriasis.

MALE' OS, (*malum*, 'an apple,' so called from its roundness.) *Zygoma*, *Os mala'rè* seu *Juga'tè* seu *Jugamentum* seu *Genæ* seu *Zygomaticum* seu *Hypopium* seu *Suboccul'tè* seu *Pudi'cum*, (F.) *Os Maleire* ou *Zygmatique* ou *de la Pommette*. The cheek or malar bone. This bone is situate at the lateral and superior part of the face; and constitutes the zygomatic region of the cheek. It is irregularly quadrilateral. Its outer surface is convex, covered by muscles and skin, and pierced with canals, called *malar*, through which vessels and nerves pass. Its upper surface is concave, and forms part of the orbit. Its posterior surface is concave, and enters into the composition of the temporal fossa. This bone is thick and cellular. It is articulated with the frontal, temporal, sphenoid, and superior maxillary bones, and is developed by a single point of ossification.

The part of the face made prominent by it, the French call *Pommette*.

MALAG'MA, gen. *Malag'matis*, (*μαλαγμα*, from *μαλακω*, 'I soften,') An emollient cataplasm, and, in general, every local application which enjoys the property of softening organic tissues.

MALAIRE, Malar.

MALAISE, (*mal*, 'ill,' and *aïse*, 'ease,') Indisposition.

MALAKIEN, (from *μαλακός*, 'soft,') Mollities.

MALAMBO BARK, Matias.

MALANDERS, Malandria.

MALANDRIA, (of old.) A species of lepra or elephantiasis. *Mal'anders* or *Mal'lenders*, is a disease of the horse, which consists of a scurfy eruption on the inside of the hock, or a little below it, as well as at the bend of the knee. It is called *Mallenders* in the foreleg, and *Sa'llenders* in the hind leg.

MALANDRIO'SUS, *Lep'rous*. Affected with malandria.

MA'LAR, *Mala'ris*, (F.) *Malaire*, (from *mala*,)

Belonging to the cheek,—as the *malar* bone.

MALAR PROCESS, Zygomatic process.

MALARE OS, Mala' os.

MALARIA, ([I.] 'bad air,') see Miasm.

MALARIOUS, *Mala'rial*. Owning to, or connected with, malaria,—as a *malarious soil*, *malarious disease*, &c.

MALASSIMILATION, *Malassimila'tio*, (*mala*, 'bad,' and *assimilitio*,) Imperfect or morbid assimilation or nutrition.

MALAXATE, *Moll'rè*, *Subig'erè*, *Malacissarè*, (F.) *Malacer*, (*μαλακω*, 'I soften,') To produce softening of drugs, by kneading them. The process is called *Malacea'tion*, *Malacea'tio*.

MALAXATION, see Malaxate.

MALAXIA VENTRICULI, (*μαλακωσις*, 'to soften,') *Gastromalacia*.

MALAXIS, (*μαλαξίς*,) Mollities—m. Cordis, *Cardiomalacia*—m. Hepatica, *Adiposis hepatica*.

MALAY, see Homo.

MALAZISSA'TUS, (*μαλακωσις*, 'to soften,') *Malacissatus*. One in whom the testicles have not descended. It has, also, been used synonymously with *emascu'latus* and *muliera'tus*.

MALCE, (*μαλκή*, 'torpor from cold,') Chilblain.

MALE, *Mas*, *Mas'culus*. Of the sex that begets young. Not female. What belongs to the male sex: as the *male organs of generation*.

MALE, (*μαλη*,) Axilla.

MALE ORGAN, Penis.

MALE ([I.] disease) *DI BRENDO*, see Scherlievo — m. di *Fiume*, Scherlievo — m. *Scabioso venereo*, Scherlievo — m. di *Scaurliero*, Scherlievo — m. di *Scherliero*, Scherlievo — m. del *Sole*, Pellagra.

MALEFICIUM, (*malus*, 'bad,' and *facere*, 'to make,') Poisoning.

MALFORMATION, *Mal'forma'tio*, *Malconforma'tio*; (*malus*, 'bad,' and *forma*, 'form,') A wrong formation; or irregularity in the structure of parts. See Monster.

MALHERBE, *Plumbago Europæa*.

MALIA, (*μαλη*,) see Equinia.

MALIASMUS, (*μαλιασμος*,) Malis, see Equinia.

MALICE, *Althæa*.

MALICHORIUM, see *Punica granatum*.

MALICORIUM, (*malum*, 'apple,' and *corium*, 'skin,') see *Punica granatum*.

MALIE, (*μαλη*,) Equinia.

MALIG'NANT, *Malig'nus*, (*malus*, 'bad,') (F.) *Malin*. A term applied to any disease whose symptoms are so aggravated as to threaten the destruction of the patient. A disease of a very serious character, although it may be mild in appearance; — *Mor'bus malig'nus*.

MALIN, Malignant.

MALING'ERER. ([F.] *malinge*, 'sickly.') A simulator. (F.) *simulateur*, of disease, so termed in the British military service. See Abraham man.

MAL'IS, (*malis*.) *Malias'mus*, *Cutaneous vermination*, *Helminthion'cus*. The cuticle or skin infected with animalcules. — *Phtheiri'asis*, *Parasiti's-mus superfi'ciei*. In Persia, this affection is produced by the Guinea worm; in South America, by the Chigre; and in Europe, occasionally by the Louse. See Equinia, and Phtheiri'asis.

MALIS DRACUNCULUS, *Dracunculus*—*m. Gordii*, *Dracunculus*—*m. Pediculi*, *Phtheiri'asis*.

MALLENDERS, Malandria.

MALLE'OLAR, *Malleola'ris*, (from *malleolus*.) (F.) *Malléolaire*. Belonging or relating to the ankles.

MALLEOLAR ARTERIES are two branches furnished by the *anterior tibial* about the instep; the one—the *internal*—passes transversely behind the tendon of the tibialis anticus, to be distributed in the vicinity of the malleolus internus;—the other—the *external*—glides behind the tendons of the *extensor communis digitorum pedis* and the *peroneus brevis*, and sends its branches to the parts which surround the outer ankle, as well as to the outer region of the tarsus.

MALLE'OLUS, (dim. of *malleus*, 'a mallet.') *Racet'a*, *Racet'ta*, *Ras'cha*, *Raset'ta*, *Ras'e'ta*, *Sphy'ra*, *Sphyron*, *Ta'lus*, *Diab'ebos*, *Tale'olus*, *Pe'za*, *An'kle*, (Prov.) *An'cler*, *An'clet*, *An'cliff*, *An'cline*, *An'kley*, *Rick*. (Sc.) *Cute*, *Coot*, *Hancloth*, *Heatherolu*, (F.) *Malléole*, *Chevillon du Pied*. The two projections formed by the bones of the leg at their inferior part. The *inner* belongs to the tibia; the *outer* to the fibula. The ankles afford attachment to ligaments; and each has a sort of gutter, in which certain tendons slide. See Malleus.

MAL'LET, *Malle'olus*, (F.) *Maillet*. A kind of hammer, used with a gouge for removing or cutting bones, in certain surgical and anatomical operations.

MAL'LEUS, ('a mallet,') *Malle'olus*, *Ossic'ulum malle'olo assimila'tum*, (F.) *Marteau*. The longest and outermost of the four small bones of the ear. It is situate at the outer part of the tympanum, and is united to the *membrana tympani*. It has, 1. An oval head, which is articulated behind with the *incus*, and is supported by a narrow part called the neck: this has, anteriorly, a *long apophysis*, which is engaged in the glenoid fissure, and is called the *Apophysis* or *Process*, *Process'us grac'ilis*, of *Rau*. It affords attachment to the *anterior* *seu externus mallei* muscle. 2. A handle, which forms an obtuse angle with the neck, and corresponds to the membrane of the tympanum, which it seems to draw inwards. It is furnished at its upper extremity with a process—the *process'us bre'vis*, to which the *inter'nus mal'lei* is attached. This bone is developed by a single point of ossification.

MALLEUS, Equinia—*M. farciminosus*, see Equinia—*m.* Slender Process of the *Grêle apophyse du marteau*.

MALLOW, COMMON, *Malva*—*m.* Compound decoction of, *Decoctum malvæ compositum*—*m.* Indian, *Abutilon Avicennæ*—*m.* Yellow, *Abutilon Avicennæ*.

MALLS, *Rubeola*.

MAL'MEDY, MIN'ERAL WA'TERS OF. Malmedy is a town in Rhenish Prussia, between Spa and Coblenz. In its immediate vicinity are several acidulous chalybeate springs.

MALOBATHRUM, (*μαλοβαθρον*.) *Malabathrum*.

MALOGRANATUM, (*malum*, 'an apple,' and *granum*, 'a grain,'—having grainlike seeds.) *Punica granatum*.

MALPIGHI, ACINI OF, *Corpora Malpighiana*.

MALPIG'HIA MOUREL'LA (after Malpighi), (F.) *Mourellier*, *Samarouba fauce*. The bark of this shrub—a native of Cayenne—is reputed to be febrifuge, and useful in diarrhoea.

MALPIGHIAN BODIES. *Corpora Malpighiana*—*m. b.* Inflammation of the Kidney, Bright's disease of the—*m. b. of the Spleen*, see Spleen—*m.* Vesicles, see Pulmo.

MALPRAX'IS, *Mal'a praxis*, *Malum reg'imen*. Bad management or treatment.

MALT, (Sax. *mealt*.) (*μαλατω*, 'I soften.' [?]) *By'nz*, *Mal'um*, *Hor'dei mal'tum*, *Bra'sium*, (F.) *Drêche*. Barley made to germinate, for the purpose of forming beer. It has been recommended in medicine as antiscorbutic, antiscrofulous, &c.

MALT SPIRIT. A spirit distilled from malt. It is the basis of most of the spirituous cordials.

MAL'TA, CLI'MATE OF. The climate of Malta is pretty equable, the range of temperature in the twenty-four hours being rarely more than 6°. The air is almost always dry and clear. The most disagreeable wind is the sirocco, which is the source of more or less suffering to the pulmonary invalid. The winter climate is favourable. Dr. Liddell thinks that no place which he has seen in the south of Europe can compete with Malta for a mild, dry, bracing air in November, December, and part of January; and during the other winter and spring months he thinks it is equal to any of them.

MAL'THA, (*μαλθα*.) *Mal'thē*, (*μαλαττω*, 'I soften.') Wax, particularly soft wax.

MALTHAXIS, (*μαλθαξις*.) Mollities.

MALTHE, (*μαλθη*.) Maltha.

MALTUM, Malt.

MALUM, (*malus*, 'bad,') Disease, *Me'tum*, ('an apple,') see *Pyrus Malus*—*m.* Articuli, Gout—*m.* Caducum, Epilepsy—*m.* Caducum pulmonum, Asthma—*m.* Canum, see *Pyrus cydonia*—*m.* Coenae, Averrhoa carambola—*m.* Cotoneum, see *Pyrus cydonia*—*m.* Cotunni, Neuralgia femoro-poplitea—*m.* Hypochondriacum, Hypochondriasis—*m.* Hysterium, Hysteria—*m.* Insanum, see *Solanum Melongena*—*m.* Ischialicum, Neuralgia femoro-poplitea—*m.* Lazari, Elephantiasis—*m.* Lycopersicum, *Solanum lycopersicum*.

MALUM MOR'TUUM. A species of lepra, in which the affected portions of skin seem to be struck with death.

MALUM PILARE, Trichosis—*m.* Pottii, Vertebral disease—*m.* Primarium, Idiopathia—*m.* Regimen, Malpraxis—*m.* Spinosum, *Datura stramonium*—*m.* Terrestris, *Atropa Mandragora*—*m.* Venereum, Syphilis.

MALUS, ('an apple tree,') *Pyrus malus*—*m.* Aurantia major, *Citrus aurantium*—*m.* Communis, *Pyrus malus*—*m.* Dasyphylla, *Pyrus malus*.

MAL'US IN'DICA, *Bilambi biting-bing* of Bon-tius. *Ord.* Rosaceæ. The juice of this East India tree is cooling, and is drunk as a cure for fevers. The leaves, boiled and made into a cataplasm with rice, are famed in all sorts of tumours. The juice, mixed with arrack, is drunk for the cure of diarrhoea. The ripe fruit is eaten as a delicacy, and the unripe is made into a pickle for the use of the table.

MALUS LIMONIA ACIDA, see *Citrus medica*—*m.* Medica, see *Citrus medica*—*m.* Sylvestris, *Pyrus malus*.

MAL'VA, (from *μαλας*, 'soft,') *Mal'vus sylves'tris seu cultiva'ris*, *Com'mon Mal'low*, (F.) *Mauve sauvage*. *Ord.* Malvaceæ. The leaves and flowers are chiefly used in fomentations, cataplasms, and emollient enemata. Its properties are demulcent.

Malva rotundifolia, *Mal'achē*, *Mal'ochē*, has like virtues, as well as the other varieties.

MALVAT DE LANGUEDOC, see Anthrax.

MALVAVISCUM, (*malva*, and *viscum*, 'bird lime,') Althæa.

MALVERN, WATERS OF. The village of Great Malvern (pronounced *Mac'vern*), in Worcestershire, England, has for many years been celebrated for a spring of remarkable purity, which has acquired the name of the *Holy well*. It is a carbonated water, containing carbonates of soda and iron, sulphate of soda, and chloride of sodium; and is chiefly used externally in cutaneous affections.

MAMA PIAN. (F.) *Maman pian*. An ulcer of a bad aspect, which is the commencement of the pian, and which, after having destroyed the flesh, extends to the bones. It is also called the *Mother of Pians*, (F.) *La mère des pians*.

MAM'EI, Mam'oe, Mom'in or Tod'dy tree. From incisions made in the branches of this West Indian tree, a copious discharge of pellucid liquor occurs, which is called *Momin* or *Toddy wine*. It is diuretic, and is esteemed to be a good antilithic and lithontriptic. See *Mammea Americana*.

MAMELLE, (dim. of *mamma*,) see *Mamma*.

MAMELON, Nipple.

MAMELONNE, Mammillated.

MAMELONS DU REIN, Papillæ of the kidney.

MAMILLA, Mammula, (dim. of *mamma*,) see *Mamma*.

MAMILLAIRE, Mammillary.

MAMILLARY, Mammillary.

MAM'MA, (*mamma*, one of the earliest cries of the infant, ascribed to a desire for food.) *Mas'thos, Mas'tus, Maz'os, The'le, Ti't'hé, Ti't'his, Ti't'hos, Ti't'y, Ru'ma, Runis, U'ber, Nu'trix, Gemi-po'ma*. The *fé'male breast, Pap, Pappé*, (*Mam'milla, Mam'illa* being the male breast), (F.) *Mamelle*. A glandular organ proper to a class of animals—the *mammalia*—and intended for the secretion of milk. The *mammæ* exist in both sexes, but they acquire a much greater size in the female, especially during pregnancy and lactation. In women, before the age of puberty, the breasts are but little developed. At this period, however, towards the central part of each breast, the skin suddenly changes colour, and assumes a rosy tint. It is of a reddish brown in women who have suckled several children. This circle has a rugous appearance, owing to the presence of sebaceous glands, and is called *Are'ola* or *Aure'ola*. These glands—*Tu'bercles of the Are'ola* of Sir Astley Cooper—furnish an unctuous fluid for defending the nipple from the action of the mouth of the sucking infant. In the midst of the aureola is the nipple, a conoidal eminence, of a rosy tint, susceptible of erection, and at the surface of which the galactophorous ducts open. Besides the skin covering them, the breasts are also composed of a layer of fatty areolar tissue, more or less thick; of a large gland; excretory ducts; vessels, nerves, &c. See *Mammary*.

In the animal the *mamma* is called the *U'd' der, U'ber, the Bay*, (Prov.) *Yever*.

The breasts plus the space between are called the *bo'som, si'nus, col'pus*. At times these terms are restricted to the space between the breasts. (F.) *Sein*.

Mamma also means a nurse.

MAMMAIRE, (from *mamma*,) *Mammary*.

MAMMAL, pl. *Mamma'tia, Mam'mifer, Mam'miferous animal*, (from *mamma*.) An animal that suckles its young.

MAM'MARY, Mam'marius, (F.) *Mammaire*. Relating to the breasts.

MAMMARY ABSCESS, *Mastodynia apostematosa*.

MAMMARY ARTERIES are three in number. They are distinguished into—1. The *Internal Mam'mary, Arte'ria sterna'talis, A. Sous-sterna'l* (Ch.), *Internal thora'cic*. It arises from the subclavian, and descends obliquely inwards, from its origin to the cartilage of the third rib. Below

the diaphragm it divides into two branches; the one *external*, the other *internal*. From its origin until its bifurcation, it gives branches to the muscles and glands of the neck, to the thymus, mediastinum, pericardium, and oesophagus. In each intercostal space it gives off *internal* and *external musculo-cutaneous* branches, and also on each side the *superior diaphragmatic*. Its two ultimate branches are distributed on the parietes of the abdomen, and anastomose with the external mammary, intercostal, lumbar, circumflexa ilii, and epigastric arteries. 2. The *External Mam'mary Arteries* are two in number, and are distinguished into *superior* and *inferior*. The *superior external mam'mary, First of the thora'cics* (Ch.), *Superior external thora'cic, Superior thora'cic*, is furnished by the axillary artery. It descends obliquely forwards between the pectoralis major and pectoralis minor, to which it is distributed by a considerable number of branches. The *inferior external mam'mary, the second of the thora'cics* (Ch.), *Long or inferior thora'cic*, arises from the axillary artery, a little below the preceding. It descends vertically over the lateral part of the thorax; curves afterwards inwards; becomes subcutaneous, and divides into a number of branches, which surround the breast. It gives branches to the pectoralis major, serratus major anticus, the intercostal muscles, the glands of the axilla, and the integuments of the breast.

MAMMARY GLAND, Glandula lactifera, Mastaden, is the secretory organ of the milk. It is situate in the substance of the breast, to which it gives shape. The tissue of this gland results from the assemblage of lobes of different size, united intimately by a dense areolar tissue. Each of these is composed of several lobules, formed of round granulations, of a rosy white colour, and of the size of a poppy seed. The glandular grains give rise to the radicles of the excretory canals of the *mamma*, which are called *galactophorons* or *lactiferous*. These excretory vessels unite in ramusculi, rami, and in trunks of greater or less size; collect towards the centre of the gland; are tortuous, very extensible, and semi-transparent. All terminate in sinuses, situate near the base of the nipple, which are commonly from 15 to 18 in number. These sinuses are very short, conical, and united by areolar tissue. From their summits a fasciculus of new ducts sets out, which occupy the centre of the nipple and open separately at its surface. The arteries of the mammary gland come from the thoracic, axillary, intercostal, and internal mammary. The veins accompany the arteries; the nerves are furnished by the intercostals and brachial plexus; the lymphatic vessels are very numerous, and form two layers. They communicate with those of the thorax, and pass into the axillary ganglions.

MAMMARY SARCO'MA, Mas'toid sarco'ma of Abernethy, *Emphy'ma sarco'ma mamma'rum*. A tumour, of the colour and texture of the mammary gland; dense and whitish; sometimes softer and brownish; often producing on extirpation a malignant ulcer with indurated edges. Found in various parts of the body and limbs.

MAMMARY VEINS follow the same course as the arteries, and have received the same denominations. The *inter'nal mam'mary vein* of the right side opens into the superior cava; that of the left into the corresponding subclavian vein. The *exter'nal mam'mary veins* open into the axillary vein.

MAM'MEA AMERICA'NA, (F.) *Abbricot sauvage, Ord. Guttifera*. The systematic name of the tree on which the *mamme* or *mamei* fruit grows. This fruit has a delightful flavour when ripe, and is much cultivated in Jamaica, where it

is generally sold in the markets as one of the best fruits of the island. See *Mammea Americana*.

MAMMEA'TA, *Mamma'sa*, (from *mamma*.) One who has large breasts.

MAMMELLA, (dim. of *mamma*.) Nipple.

MAMMIFER, (*mamma*, and *fero*, 'to bear.') Mammal.

MAMMIFEROUS ANIMAL, Mammal.

MAMMIFORM, (*mamma*, and *forma*, 'form.') Mastoid.

MAMMILLA, (dim. of *mamma*.) Mamma

(male), Nipple.

MAMMILLÆ MEDULLARES, see *Mammillary*.

MAMMILLAIRE, Mammillary.

MAMMILLARIS, (from *mammilla*.) Mastoid.

MAM'MILLARY, *Mammilla'ris*, (from *mammilla*, (F.) *Mammillaire*, *Manillaire*.) Having the shape of a nipple. See *Mastoid*.

MAMMILLARY EM'INENCE is a name given, 1. To more or less marked prominences on the inner surface of the bones of the cranium, which correspond to the anfractuositities of the cranium. 2. To white, round, medullary tubercles of the size of a pea, situate at the base of the brain, behind the gray substance from which the *Tige pituitaire* of the French anatomists arises. These *Mammillary Tubercles*, *Corpora albican'tia* seu *candican'tia* seu *mammilla'ria* seu *risifor'mia*, *Bulbi for'nici*, *Mammill'æ* seu *Globuli medulla' res*, *Prominentiæ albican'tes*, *Processus mammilla' res cer'e'ri*, *Protensio'nes glandula' res*, *Eminent' iæ candican'tes*, *Priorum crurum for'niciis bulbi*, *Willis's Glands*, (F.) *Bulbes de la voûte à trois piliers*, *Tubercles pisiformes* (Ch.), are united to each other by a small grayish band, which corresponds with the third ventricle. They receive the anterior prolongations of the fornix.

Some ancient anatomists, taking the nervous trunks, to which Willis first gave the name of *olfactory nerves*, for simple appendages of the brain, called them on account of their shape, *Carunculae mammilla' res*. Vesalius, Fallopius, Columbus, and several others, termed them *Processus mammilla' res cer'e'ri ad na' res*. They have also been called *Trigona olfacto'ria*.

MAM'MILLATED, (F.) *Mamelonné*. That which has mammiform projections on its surface.

MAMMILLATED LIVER, Cirrhosis.

MAMMITE, Mastitis.

MAMMITIS, (*mamma*, and *itis*), Mastitis.

MAMMOSA, Mammeata.

MAM'MULA, (dim. of *mamma*.) *Mamill'a*. A small breast.

MAMOE, Mamei.

MAN, (Sax. *man* or *mon*.) *Aner*, *Anthropos*, *Homo*.

MANAKIN, (dim. of *man*.) *Fantom*.

MAN-IN-THE-GROUND, *Convolvulus panduratus*—m. of the Earth, *Convolvulus panduratus*.

MANCENILLIER, *Hippomane mancinella*.

MANCHE D'HIPPOCRATE, ([L.] *manica*, 'a sleeve.') *Chaussée*.

MANCHINEEL, (from (S.) *mangana*, 'apple,') *Hippomane mancinella*.

MANCURANA, *Origanum*.

MANDIBULA, (*mandere*, 'to chew') *Maxillary bone*.

MANDIBULARIS MUSCULUS, *Masseter*.

MANDIO, *Jatropha manihot*.

MANDO, (from *mandere*, 'to chew.') *Glutton*.

MANDRAGORA, (*μανδραγόρας*.) *Atropa mandragora*—m. *Acaulis*, *Atropa mandragora*—m. *Officinalis*, *Atropa mandragora*—m. *Vernalis*, *Atropa mandragora*.

MANDRAGORI'TES, (*μανδραγορίτης*.) (from *mandragora*.) Wine in which the roots of mandrake have been infused.

MANDRAKE, *Atropa mandragora*, *Podophyllum peltatum*, *P. montanum*.

MANDRIN À VIRGULE, see *Brise-coque*.

MANDUCATIO, (*manducare*, *manducatum*, 'to chew,') *Mastication*—m. *Difficilis*, *Bradymasesis*.

MANDUCATORY, *Manducato'rius*, (from *manducatio*, 'mastication,') Appertaining or relating to mastication;—as,

MANDUCATORY NERVE, see *Trigemini*.

MANE, *Arachis hypogæa*.

MANGANESE, *Mangane'sium*, *Manga'nium*, *Man'ganum*, ([G.] *Manganerz*, 'Mangan earth,') (?). A metal, which is pretty abundant in nature in an oxidized state. It is of a grayish-white colour, resembling some varieties of cast iron, is hard and brittle, and destitute of magnetic properties. Specific gravity about 8.

MANGANESE, BLACK OXIDE OF, *Mangane'sii oxidum* seu *oxidum nigrum* seu *Binox'idum*, *Tetrox'idum* of *manganese*, *Magne'sia nigra*, *Mangane'sium vitrario'rum* seu *oxydatum nativum* seu *nigrum* seu *ochraceum nigrum* seu *oxydatum nativum*, *Man'ganum oxydatum nativum*, *Molybdaenum magne'sii*, *Oxodes man'gani nativæ*, *Perox'ydum mangane'sii nigrum nativum*, *Superox'ydum mangani'cum*, (F.) *Oxyde noir de manganese*. This oxide is not much used in medicine. It has been advised to dust the affected parts, in tinea capitis, with the powder. It was officinal in the Edinburgh pharmacopœia, and is used in preparing chlorine, solution of chlorine, and solution of chlorinated soda.

MANGANESE, SALTS OF, see *Manganese, sulphate of*.

MANGANESE OXYDE NOIR DE, *Manganese, black oxide of*.

MANGANESE, SULPHATE OF, *Mangane'sii Sulphas* seu *Protox'idi Sulphas*, *Sulphate of Protox'ide of manganese*. A rosecoloured and very soluble salt, isomorphous with sulphate of magnesia. It is prepared on a large scale for the use of the dyer, by heating, in a close vessel, peroxide of manganese and coal, and dissolving the impure protoxide, thus obtained, in sulphuric acid, with the addition of a little chlorohydric acid towards the end of the process. The solution is evaporated to dryness, and again exposed to a red heat, by which the persulphate of iron is decomposed. Water then dissolves the pure sulphate of manganese, leaving the oxide of iron behind.

This salt has been recommended as a cholagogue, in doses of a drachm or two.

The salts of manganese have been prescribed in chlorosis and amenorrhœa, and as substitutes for chalybeates generally. The subcarbonate and the oxide have been chiefly employed. The chloride is the basis of Mr. James Young's disinfectant.

MANGANESE, TETROXIDE OF, *M. Black oxide of*.

MANGANESII BINOXIDUM, *Manganese, black oxide of*—m. *Oxidum*, *Manganese, black oxide of*—m. *Perox'ydum nigrum nativum*, *Manganese, black oxide of*—m. *Protoxidi sulphas*, *Manganese, sulphate of*—m. *Sulphas*, *Manganese, sulphate of*.

MANGANESEIUM, *Manganese*.

MANGANEIUM OCHRACEUM NIGRUM, *Manganese, black oxide of*—m. *Oxydatum nativum* seu *nigrum*, *Manganese, black oxide of*—m. *Vitrarium*, *Manganese, black oxide of*.

MANGANI OXODES NATIVA, *Manganese, black oxide of*.

MANGANIC ACID, see *Potassæ permanganæ*.

MANGANICUM SUPEROXIDUM, *Manganese, black oxide of*.

MANGANUM, *Manganese*.

MANGANUM, *Manganese*—m. *Oxydatum nativum*, *Manganese, black oxide of*.

MANGE, ([F.] *démangeaison*, 'itching,') *Scabies ferina*.

MANGIFERA INDICA, *M. domestica*, (*man-go*, 'the fruit,' and *fero*, 'I bear.')

The *Man'go tree*, (F.) *Manguier*. *Ord.* Terebinthaceæ. A tree cultivated over Asia, and in South America. Mangos, when ripe, are juicy, of a good flavour, and so fragrant as to perfume the air to a considerable distance. They are eaten, either raw or preserved with sugar. From the expressed juice a wine is prepared; and the remainder of the kernel can be reduced to an excellent flour for bread.

MANGO TREE, *Mangifera Indica*.

MANGONISATIO, (from *παγῶσις*, 'a deception,') Falsification.

MANGONIUM, Falsification.

MANGOSTAN, *Garcinia mangostana*.

MANGOSTANA, *Garcinia mangostana*—*m.* Cambogia, *Garcinia cambogia*—*m.* *Garcinia*, *Garcinia mangostana*.

MANGOSTAN, *Garcinia mangostana*.

MANGOSTEEN, *Garcinia mangostana*.

MANGUIER, *Mangifera Indica*.

MANHOOD, Adult age.

MANIA, (*μᾶνία*), *Fu'ror*, *Hemianthro'pia*, *Fu'ror* seu *Ephro'nia* seu *Delir'ium ma'nia*, *Insa'nia*, *Delir'ium mani'acum* seu *furi'osum*, *Ma'nia uni-versa'lis*, *Ves'a'nia ma'nia*, (F.) *Manie*, *Hyperphrénie*, *Ra'ving* or *fu'rious mad'ness*, (Sc.) *Wod'nes*, (from *παύωμαι*, 'I am furious.')

With some, it means *insanity*. Disorder of the intellect, in which there is erroneous judgment or hallucination, which impels to acts of fury. If the raving be not directed to a single object, it is mania properly so called; if to one object, it constitutes *monomania*, which term is, however, usually given to melancholy. Mania attacks adults chiefly; and women more frequently than men. The prognosis is unfavourable. About one-third never recover; and they who do are apt to relapse. *Separation* is one of the most effective means of treatment, with attention to the corporeal condition and every thing that can add to the mental comfort of the patient, and turn his thoughts away from the subjects of his delusion. In the violence of the paroxysms, recourse must be had to the strait waistcoat, the shower bath, &c. Separation should be continued for some weeks during convalescence, with the view of preventing a relapse.

MANIA, DANC'ING. *Danc'ing plague*. A form of convulsion, which has appeared, at various times, epidemically under the form of St. Vitus's dance, St. John's dance, Tarantism, Hysteria, Tigretier (in Abyssinia), and diseased sympathy; and which has been fully described by Hecker in his 'Epidemics of the Middle Ages.' See *Convulsionnaire*.

MANIA EPILEPTICA, see Epilepsy—*m.* Erotica, *Erotomania*—*m.* Lactea, *M. puerperal*—*m.* Melancholica, *Melancholy*—*m.* a Pathemate, *Empathema*—*m.* Pellagria, *Pellagra*—*m.* a Potu, *Delirium tremens*.

MANIA, PUERPERAL, *M. puerperarum acuta* seu *puerpera'lis* seu *lac'tea*, *Insa'nia puerperarum*, *Euphthalpathi'a puerpera'lis*, *Puerperal Insan'ity*. Mania which supervenes in the childbed state.

MANIA PUERPERAL ACUTA, *M. puerperal*—*m.* sine *Delirio*, *Pathomania*—*m.* a Temulentiâ, *Delirium tremens*.

MANIAC, see Maniodes.

MANIACAL, Maniodes.

MANIACUS, Maniodes.

MANICA HIPPOCRATIS, ('a sleeve,' from *manus*, 'the hand,') *Chausse*.

MANICOCOMIUM, (*manicus*, and *κομῶν*, 'to take care of,') *Morotrophium*.

MANICUS, (*μανικός*), Maniodes.

MANIE, Mania—*m.* sans *Délire*, *Empathema*.

MANIGUETTA, *Amomum grana paradisi*.

MANIHOT UTILISSIMA, *Jatropha manihot*.

MANIIBAR, *Jatropha manihot*.

MANIKIN, (dim. of *Man*,) *Fantom*.

MANILUVIUM, *Maniluvium*, (*manus*, and *lavo*, 'I wash,') A bath for the hands. It may be rendered stimulating, by means of muriatic acid, mustard, &c.

MANIOC, *Jatropha manihot*.

MANIO'DES, (*mania*, and *odes*,) *Mani'acus*, *Ma'nicus*, *Mani'acal*. (F.) *Maniaque*. One labouring under mania; *Ma'niac*, *Hemianthro'pus*, *Furi'osus*, *Furibundus*, *Lys'sas*, *Lyss'eter*.

MANIPLUS, *Manipulus*.

MANIPULATION, (from *manus*,) Mode of working, as in pharmacy.

MANIP'ULUS, *Manipulus*, (from *manus*, and *pleo*, 'to fill,') 'a handful,' (F.) *Poignée*. The quantity of a substance capable of filling the hand. A handful. See *Fasciculus*.

MANMIDWIFE, *Aecoucheur*.

MANNA, see *Fraxinus ornus*—*m.* *Briançon*, see *Pinus larix*.

MANNA BRIGANTINA, see *Pinus larix*—*m.* *Calabrina*, see *Fraxinus ornus*—*m.* *Croup*, *Semolina*—*m.* *Laricea*, see *Pinus larix*—*m.* *Metallorum*, *Hydrargyri submurias*.

MANNEQUIN, (dim. of *Man*,) *Fantom*.

MANNIKIN, *Fantom*.

MANNIN, see *Fraxinus ornus*.

MANNITE, see *Fraxinus ornus*.

MANŒUVRE (F.), anglicized, and pronounced *maneu'vre*, (*main*, 'the hand,' and *œuvre*, 'work,') A dexterous movement. Applied in France to the practice of surgical or obstetrical operations on the dead body or phantom; *Opera'tio chirur'gica* seu *obstet'rica*.

MANSANA ARBOREA, see *Jujube*.

MANSFORD'S PLATES, see *Galvanism*.

MANSORIUS, (*mandere*, *manum*, 'to chew,') *Buccinator*.

MANSTUPRA'TIO, (*manus*, 'the hand,' and *stuprare*, *stupratorum*, 'to ravish,') *Masturbation*.

MANTELE, ('a napkin,') *Bandage* (body).

MANTIA, *Man'tica*, *Man'tice*, in English, *mancy*; a common suffix, from *μανεια*, denoting 'divination;' *Divina'tio*, *Prædivinatio*, *Præsig'ium*.

MANTILE, ('a napkin,') *Bandage* (body).

MANTLE, (*mantelum*, 'a cloak,') *Panniculus carnosus*.

MANUBRIUM, (from *manus*,) The handle of any thing:—as *manu'brium mallei*, *pet'iolus mallei*, 'the handle of the malleus.'

MANUBRIUM, *Chasse*—*m.* *Manus*, *Radius*.

MANUBRIUM STERNI. The uppermost broad part of the sternum.

MANULUVIUM, (*manus*, 'the hand, and *lavo*, 'I wash,') Bath, hand, *Maniluvium*.

MA'NUS, gen. *Manus*, *Cheir*, *Chir*, *Instrumentum instrumento'rum*, *Hand*, *Paw*, (F.) *Main*. The part which terminates the upper extremity in man, and which is inservient to prehension and touch. It extends from the fold of the wrist to the extremity of the fingers. The hand is sustained by a bony skeleton, composed of a number of pieces, movable on each other; of muscles, tendons, cartilages, ligaments, vessels, nerves, &c. It is divided into three parts—the *carpus* or wrist, the *metacarpus*, and *fingers*. Its concave surface is called the *palm*; the convex surface the *back of the hand*. The facility of being able to oppose the thumb to the fingers in order to seize objects forms one of the distinctive characters of the human hand.

MANUS CHRIST'I PERLA'TA. A name anciently given to troches, prepared of pearls and sugar of roses. They were called *Manus Christi's simplices*, when pearls were not employed.

MANUS DE'I. An ancient plaster, prepared of

rax, myrrh, frankincense, mustich, gum ammoniac, galbanum, oil, &c. See Opium.

MANUS HEPATIS, see Liver—*m.* Jecoris, see Liver—*m.* Parva majori adjuvix, see Digitus.

MANUSTUPRATIO, (*manus*, 'the hand,' and *stuprare*, 'to ravish,') Masturbation.

MANUSTUPRATOR, Masturbator.

MANYPLIES, ('many plies or folds,') Omasum.

MAPLE, (*Sax. mapulber,*) Acer saccharinum—*m.* Ground, Heuchera cortusa—*m.* Red, Acer rubrum—*m.* Striped, Acer Pennsylvanicum—*m.* Sugar, Acer saccharinum, see Saccharum acernum.

MARAI, (*Sax. meppe,*) Marsh.

MARANSIS, (*μαρανσις,*) Atrophy.

MARANTA, (after Maranta, a Venetian physician,) Arrowroot—*m.* Arundinacea, Arrowroot.

MARANT'GA GALAN'GA, *Alpinia galan'ga, Amomum galan'ga, Galan'ga.* Ord. Zingiberaceae. The smaller galan'gal. Two kinds of galangal are mentioned in the pharmacopœias; the greater, obtained from *Kempferia galan'ga*, and the smaller, from the root of *Marant'ga galan'ga*. The dried root is brought from China, in pieces from one to two inches in length, but scarcely half as thick; branched; full of knots and joints, with several circular rings, of a reddish brown colour, on the outside, and brownish within. It was formerly much used as a warm stomachic bitter, and generally ordered in bitter infusions.

MARANT'IC, *Marant'icus*, (F.) Marantique, (from *maransis*, 'atrophy,') Of, or belonging to, *maransis* or atrophy.

MARASCHINO, see Spirit.

MARASMO'YRA, *Febris marasmo'des*, (*marasmus*, and *rup*, 'fever,') Fever of emaciation general. Hectic fever.

MARASMUS, (*μαρασμος,*) Atrophy—*m.* Lactantium, Pedatrophia—*m.* Phthisis, Phthisis pulmonalis.

MARASMUS SENI'LIS, *Tu'bes senum, Gerontoph'ia.* Progressive atrophy of the aged.

MARASMUS TABES, *Tabes*—*m.* Tabes dorsalis, Tabes dorsalis.

MARASQUINO, see Spirit.

MARATHRI'TES, (*μαραθριτης*, from *μαραθρον*, 'fennel,') Wine impregnated with fennel.

MARATHROPHYLLUM, (*marathrum*, and *φυλλον*, 'a leaf,') Peucedanum.

MARATHRUM, (*μαραθρον*,) Anethum—*m.* Sylvestre, Peucedanum.

MARAUGIA, (*μαραυγω*, 'to dazzle,') Metamorphopsia.

MARBLE, Marmor.

MARBRE, Marmor.

MARC, Magma.

MARCASITA, Bismuth—*m.* Alba, Bismuth, subnitrate of—*m.* Plumbea, Antimonium.

MARCASITE MAGISTERIUM, Bismuth, subnitrate of.

MARCHAN'TIA POLYMER'PHA, (after Marchant, a Parisian botanist,) *M. stella'ta seu umbella'ta, Hepat'ica fonta'na, Li'chen stella'tus, Jecora'ria, Liv'erwort.* (F.) *Hépatique des fontaines.* Ord. Hepaticæ. This plant is mildly pungent and bitter. It is recommended as aperient, resolvent, and antiscorbutic; and is used in diseased liver, &c.

MARCHE (LA), Walking.

MARCHIO'NIS PUL'VIS, *Pow'der of the Mar'quis.* A powder formerly considered to be antiepileptic; and composed of *Male peony root, Mistletoe, Ivory shavings, Horn of the hoof of the stag, Spodium, Tooth of the monodon, coral, &c.*

MARCORES, (*marcere*, 'to wither,') Atrophy, Emaciation.

MARCORY, Stillingia.

MARÉCAGEUX, (from *marais*, 'a marsh,') Elodes.

MAREMMATIQUE, (from *Maremma*, a mala-

rious region of Italy near the sea [*mare*]). See Elodes.

MAREO, Puna.

MARE'S FAT, *Inula dysenterica*—*m.*s. Tail, *Hippuris vulgaris*.

MARGA CAN'DIDA, *Lac lu'næ.* An ancient name for a variety of spongy, white, friable marl, which was employed as an astringent and refrigerant.

MARGARETIZZA, *Scherlievo.*

MARGARIN, (from *μαργαρον*, 'a pearl,') see Fat.

MARGARITA, (*μαργαριτης*,) Pearl.

MARGARITA'CEOUS, *Margarita'ceus, Na'creous*, (F.) *Nucré*, (from *margarita*, 'pearl, mother of pearl,') Resembling, or of the nature of, mother of pearl,—as *Leuco'ma Margarita'ceum*. Pearl-like leucoma.

MARGARON, (*μαργαρον*,) Pearl.

MARGELIS, (*μαργηλις*,) Pearl.

MARGELLUM, Pearl.

MARGINI-SUS-SCAPULO-TROCHITÉRIEN. Teres minor.

MARGO, gen, *Mar'ginis, Bord*—*m.* Dentatus, see Retina—*m.* Orbitalis, see Orbit.

MARGOSA TREE, see Melia azedarach.

MARGUERITE PETITE, (from *margarita*,) Bellis—*m.* des Près, grande, Chrysanthemum leucanthemum.

MARIENBAD, MIN'ERAL WA'TERS OF. Celebrated springs in Bohemia. The Kreuzbrunn contains sulphate of soda, carbonate of iron, and carbonic acid. Season, from the 1st of June to the 15th of September.

MARIGOLD, DIAMOND FIG, Mesembryanthemum crystallinum—*m.* Garden, Calendula officinalis—*m.* Marsh, *Caltha palustris*—*m.* Single, Calendula officinalis—*m.* Wild, Calendula arvensis.

MARIOTTE, EXPERIMENT OF, (F.) *Expérience de Mariotte.* A celebrated experiment of the Abbé Mariotte, which consists in placing two small round spots on a wall at some distance from each other, standing opposite the left-hand object, and looking at it with the right eye, the left being closed. By walking backwards, until the distance from the object is about five times as great as the distance between the two objects, the latter will be found to disappear. Mariotte and, after him, many ophthalmologists, inferred that the optic nerve, on which the ray doubtless falls in this experiment, is insensible; and hence that the choroid may be the seat of vision,—not the retina. The inference is illogical; for it doubtless falls on the part of the optic nerve without doubt, where the central artery enters, and the central vein leaves the eyeball, and where there is necessarily no neurine.

MARIS, (*μαρις*,) Ancient name of a measure, containing 83 pints and 4 ounces.

MARISCA, ('a large fig,') Ficus, Hæmorrhoids.

MARJOLAINE, ([low L.] *majorana*,) Origanum majorana.

MARJORAM, COMMON, Origanum—*m.* Sweet, Origanum majorana—*m.* Wild, Origanum.

MARMALADE, Marmelade.

MARMALET, Marmalade.

MARMAROS, (*μαρμαρος*, from *μαρμαριζω*, 'to shine,') Marmor.

MARMARYGE, (*μαρμαρυγη*, 'splendor,') *ερα* Metamorphopsia.

MARMARYGO'DES, (*μαρμαρυγιδης*, from *μαρμαρυγη*, 'splendor,' and *odes*,) An epithet, joined particularly to the word *οφθαλμος*, to indicate a brilliant eye, a flashing eye. An eye which transmits the sense of imaginary objects. See Metamorphopsia.

MAR'MELADE, *Marmela'da, Marmela'ta, Marmalade, Marmalet, Mica*, ([S.] *marmelosa*, 'a quince,') Parts of quinces, or other fruits, connected with sugar, and reduced to a pulsatious consistence.

MARMELADE OF APRICOTS. A marmelade, prepared with two parts of ripe apricots deprived of their stones, and one part of white sugar.

MARMELADE OF FERNEL, M. of Tronchin.

MARMELADE OF TRONCHIN or of FERNEL. A kind of thick looch, of an agreeable taste, prepared with two ounces of oil of sweet almonds, as much syrup of violets, manna in tears, very fresh pulp of cassia, 16 grains of gum tragacanth, and two drachms of orange flower water. It is used as a laxative, demulcent, and pectoral.

MARMELATA, Marmelade.

MAR'MOR, gen. Mar'moris, (μαρμαρος, from μαρμαίρω, 'to shine,') Marble, Cal'cis Carbo'nas du'rus, (F.) Marbre native. White granular carbonate of lime. Used in pharmacy for the preparation of a pure lime, and the disengagement of carbonic acid.

MARMORATA AURIUM, Cerumen.

MARMORYGE, (μαρμαρυγή, 'splendor,') see Metamorphopsia—m. Hippocratis, Hemipopia.

MARUTE, Anthemis cotula.

MARRIABLE, Nubile.

MARRIAGEABLE, Nubile.

MAR'RIOT, DRY VOMIT OF. This once celebrated emetic, called dry, from its being exhibited without drink, consisted of equal portions of tartarized antimony, and sulphate of copper.

MARRONIER D'INDE, Æsculus Hippocastanum.

MAR'ROW, (Sax. meap̃, My'elos, Osteomy'elon, Medul'la, M. os'sium, Med'ullary Juice, Azin'gia de Muntia, Muntia de Medul'lis, (F.) Moëlle, Suc médullaire. The oily, inflammable, whitish or yellowish juice, which fills the medullary canal of the long bones, the cancellated structure at the extremities of those bones, the diploë of flat bones, and the interior of short bones. The marrow is furnished by the exhalation of the medullary membrane. It is fluid during life, and appears under the form of small points or brilliant grains after death. It is enveloped in the medullary membrane.

In the bony canals it is called *Oil of Bones*.

MARROW, SPINAL, Medulla spinalis—m. Vertebral, Medulla spinalis.

MARRUBE BLANC, Marrubium—m. Noir, Ballota foetida.

MARRUBIUM, M. vulgare seu al'bum seu Germanicum, Pra'sium, Phra'sum, Horehound, Horehound, (F.) Marrube blanc. Ord. Labiatae. The leaves have a moderately strong, aromatic smell; and a very bitter, penetrating, diffusive, and durable taste. It has often been given in coughs and asthmas, united with sugar. Dose. ʒss to ʒj, in infusion; dose of extract, gr. x to ʒss.

FORD'S BALSAM OF HOREHOUND, is made as follows:—*horehound, liquorice root, aa* ℥ij and ʒviij; *water, q. s.* to strain ℥vj. Infuse. To the strained liquor add:—*proof spirit or brandy, ℥l2; camphor, ʒj and ʒij; opium and benjamin, aa ʒj; dried squills, ʒij; oil of aniseed, ʒj; honey, ℥ij and ʒviij.* It is pectoral.

MARRUBIUM, Leonurus cardiaca—m. Album, Marrubium—m. Germanicum, Marrubium—m. Nigrum, Ballota foetida—m. Vulgare, Marrubium.

MARS, ('the God of War,') Ferrum—m. Solubilis, Ferrum tartarizatum.

MARSEILLES, CLIMATE OF. The remarks made upon the climate of Montpellier apply even in greater force to that of Marseilles. It possesses all the objectionable qualities of the climate of southeastern France.

MARSH, (Sax. meppe, Lim'nē, He'lus, Pal'us, (F.) Marais. Marshy districts give off emanations, which are the fruitful source of disease and the cause of great insalubrity in many countries. The chief disease, occasioned by the malaria or miasm, is intermittent fever. Hence it be-

comes important to drain such regions, if practicable. Some marshy countries are not so liable to phthisis pulmonalis, and it has been found, that where intermittents have been got rid of by draining, consumption has, at times, become frequent. The most unhealthy periods for residence in a marshy district are during the existence of the summer and autumnal heats; at which times the water becomes evaporated, and the marshy bottom is more or less exposed to the sun's rays. This postulatam seems necessary for the production of the miasmata: for whilst the marsh is well covered with water, no miasm is given off.

MARSH MALLOW, Althæa—m. Poison, Miasm (marsh)—m. Root, Statice Caroliniana—m. Tea, Ledum palustre.

MARSHALL'S CERATE, see Cerate, Marshall's.

MARSHALL HALL'S METHOD, Ready method.

MARSHY, Elodes.

MARSUPIUM, (μαρσπιον,) Marsupion.

MAR'SUM, Mar'sium, Mar'sicum (vinum.) An ancient wine of Marsia, in Italy, which was used as an astringent in certain diseases of the mouth.

MARSUPIAL, see Marsupion.

MARSUPIALIS, Ischiotrochanterianus, Ob-turator internus.

MARSUPIATE, see Marsupion.

MARSU'PION, (μαρσπιον, 'a small sac, a pouch,') Marsyp'ion, Marsip'pon, Marsu'pium, Marsip'ium, Sac'culus. A sack or bag, with which any part is fomented. Also, the abdominal pouch in the kangaroo, opossum, &c., into which the young, born at a very early stage of development, are received and nourished with milk secreted from glands which open into the pouches. Such animals are termed *Marsu'pial, Marsu'piale, Marsu'pialia*. See Generation.

MARSU'PIUM, Marsupion, Scrotum—m. Musculosum, Dartos.

MARSYPION, Marsupion.

MARTEAU, Malleus.

MARTIAL, (from Mars,) Chalybeate.

MARTIALIS, Chalybeate.

MARTIA'TUM UNGUEN'TUM, (F.) Onguent de Soldat, Soldier's ointment. This was composed of bay berries, rue, marjoram, mint, sage, wormwood, basil, olive oil, yellow wax, and Malaga wine. It was invented by Martian; and was employed by soldiers as a preservative against injuries from cold.

MARTIS LIMATURA, Ferri limatura.

MARTUM CORTUSI, Teucrium marum—m. Creticum, Teucrium marum—m. Syriacum, Teucrium marum—m. Verum, Teucrium marum—m. Vulgare, Thymus mastichina.

MARUTA COTULA, Anthemis cotula.

MAS, gen. Ma'ris, Male, Modiolus.

MASCARPIO, (mas, and carpere, 'to enjoy,') Masturbator.

MASCHALADENITIS, (μασχλη, 'the axilla,' and adenitis, 'inflammation of a gland,') Inflammation of the glands of the axilla.

MASCHALE, (μασχλη,) Axilla.

MASCHALIÆUS, Axillary.

MASCHALIATRI'A, (μασχλη, 'the axilla,' and ιατρεία, 'healing,') Treatment of disease by applications made to the axilla.

MASCHALIS, (μασχλη,) Axilla.

MASCHALISTER, (μασχλιστηρ,) Axis.

MASCHALON'CUS, Maschalopa'nus, (μασχλη, 'the axilla,' and ογκος, 'a tumour,') A tumour or bubo or swelling in the axilla.

MASCHALOPANUS, (maschale, and panus, 'a tumour,') Maschalonus.

MASCULA, ('imitating the male,') Tribas.

MASCULUS, (dim. of Mas, 'a male,') Male.

MASESIS, (μασησις,) Mastication.

MASETER, (μασητηρ,) Masseter.

MASHUA (S.). A tuberous root of a flat, pyramidal shape, which is cultivated and cooked like the potato by the Serranos of Peru. It is watery and insipid, but nevertheless is much eaten by them. The Indians use the mashua as a medicine in dropsy, dyspepsia, and dysentery: The plant is unknown to botanists.

MASK, *Lar'va*, (F.) *Masque, Larve*. A bandage applied over the face, as a sort of mask, in cases of burns, scalds, or erysipelas. It serves to preserve the parts from the contact of air, and to retain topical applications *in situ*. It is made of a piece of linen, of the size of the face, in which apertures are made corresponding to the eyes, nose, and mouth, and which is fixed by means of strings stitched to the four angles.

MASK'ED, *Larva'tus*, (F.) *Larvé*. Disguised, —as a "masked fever," or one which is not clearly developed. See *Fever, masked*.

MAS'LACH, *Mos'lich, Amph'lon, Anfi'on*. A medicine much used by the Turks, and into the composition of which opium enters. It is excitant.

MASQUE, Mask.

MASS, *Ma'za, Mas'sa*, (F.) *Masse*, (μασσω, 'I knead.') The compound, from which pills have to be formed.

MASSA, Mass — m. *Carnea Jacobi Sylvi*, see *Flexor longus digitorum pedis profundus perforans* — m. de *Dactylis*, Paste, date — m. de *Extracto glycyrrhizæ*, Pasta glycyrrhizæ, &c. — m. de *Gummi Arabico*, Paste, marshmallow — m. de *Zizyphorum fructu*, Paste of jujubes.

MASSAGE, (from μασσω, 'to knead,') Sham-pooing.

MASSE, Mass — m. *d'Eau, Typha latifolia*.

MASSEMA, (μασημα,) Mastication.

MASSEMENT, (from μασσω, 'to knead,') Sham-pooing.

MASSEPAINS, (μασσω, 'to knead, and pain, 'bread,') Biscuits, medicinal.

MASSES APOPHYSAIRES, ('apophysar masses,') see *Vertebrae*.

MASESIS, (μασησις,) Mastication.

MASSE'TER, *Mase'ter*, (μασσηταιρ, 'I chew.') *Musculus mandibularis*, (F.) *Zygomatico-maxillaire*. A muscle situate at the posterior part of the cheek, and lying upon the ramus of the lower jaw-bone. It is long, quadrilateral, and is attached, *above*, to the inferior edge and to the inner surface of the zygomatic arch; *below*, it terminates at the angle of the jaw, and at the outer surface and inferior margin of the ramus of that bone. It is composed of an intermixture of fleshy and aponeurotic fibres. Its office is to raise the lower jaw, and to act in mastication.

MASSETER INTERNUS, *Pterygoideus internus*.

MASSETER'IC, *Mas'seterine, Masseter'icus, Masseter'inus*. (F.) *Massétérin, Massétérique*. Relating or belonging to the masseter muscle.

MASSETERINE, *Masseteric*.

MASSETERINE ARTERY arises from the trunk of the internal maxillary or temporalis profunda posterior, and is distributed to the masseter muscle, after having passed, horizontally, through the sigmoid notch of the lower jaw-bone.

MASSETERINE NERVE, is given off from the inferior maxillary branch of the fifth pair. It passes through the sigmoid notch, and is distributed on the inner surface of the masseter. In luxation of the lower jaw, this nerve is strongly stretched, and considerable pain, consequently, produced.

MASSETERINE VEIN has the same distribution as the artery. It opens into the internal maxillary vein.

MASSETERINUS, *Masseteric*.

MASSETTE, *Typha latifolia*.

MASSICOT, *Plumbi oxydum semivitreum*.

MASSING, (from μασσω, 'to knead,') Sham-pooing.

MASSULA, (dim. of *massa*,) Molecule.

MASTADEN, (*mastos*, and *adēn*, 'a gland,') see *Mamma*.

MASTADENITIS, (*mastaden*, and *itis*,) Mastitis.

MASTALGIA, (*mastos*, and *αλγος*, 'pain,') Mastodynia.

MASTAX, (μασταξ,) *Mystax*.

MASTEMA, Masticatory.

MASTER OF THE WOODS, *Galium circæ-zans*.

MASTERWORT, *Angelica atropurpurea*, *Hecræolum lanatum*, *Imperatoria*.

MASTHELICOSIS, (*mastos*, and *ἑλκος*, 'an ulcer,') Mastodynia apostematosa.

MASTHOS, (μασθος,) *Mamma*.

MASTIC, see *Pistacia lentiscus*.

MASTICA'TION, *Mastica'tio, Mase'sis, Mase'sis, Masse'ma, Manduca'tio, Commanduca'tio, Manduca'tion, Chew'ing, Chaw'ing, (masticare, masticatum, 'to chew,')* The action of bruising food in the mouth, to prepare it for the digestion it has to undergo in the stomach. This is executed by the joint action of the tongue, cheeks, and lips, which push the alimentary substance between the teeth; and by the motions of the lower jaw it is cut, torn, or bruised.

MASTICATORY, *Masticato'rium, Maste'ma, Diamaste'ma*. (F.) *Masticatoire*. Chewing. Relating or appertaining to mastication or chewing. Also, a substance, chewed with the intention of exciting the secretion of saliva.

MASTICATORY NERVE, see *Trigemini* — m. Spasm of the face, *Trismus*.

MASTICH, (μαστιχη,) see *Pistacia lentiscus* — m. Herb, common, *Thymus mastichina* — m. Herb, Syrian, *Teucrium marum* — m. Tree, *Pistacia lentiscus*.

MASTICHE, see *Pistacia lentiscus*.

MASTICHELEON, (*mastich*, and *ελαιον*, 'oil,') see *Pistacia lentiscus*.

MASTICHINA GALLORUM, (after *mastich*,) *Thymus mastichina*.

MASTIGODES HOMINIS, *Trichocephalus*.

MASTIGO'SIS, (μαστιγωσις, from *μαστιγωω*, 'I whip,') *Flagella'tio, Flagella'tion, Scourg'ing*. Employed by the ancients as a remedy in many diseases.

The *Flagellation* or *Scourging Mania*, was a form of theomania, which prevailed in Europe, in the 13th and 14th centuries. They who were affected with it were termed *Flagell'ants*.

MASTITIS, *Mastoï'tis, Mammi'tis, Mazoï'tis, Inflamma'tio Mam'mæ, Mastadeni'tis*, (F.) *Inflammation des Mamelles, Mammité, (mastos, and itis,)* Inflammation of the breast. Inflammation of the mammary gland of the pregnant or parturient female is vulgarly called a *weid* or *weed*, and a *weed in the breast*.

MASTITIS APOSTEMATOSA, Mastodynia apostematosa.

MASTITIS PUERPERA'LIS. Inflammation of the mamma in the childbed woman.

MASTIX, (after *mastich*,) see *Pistacia lentiscus*.

MASTOCARCINO'MA, (*mastos* and *carcinoma*,) Carcinoma or cancer of the breast.

MASTOCHONDRO'SIS, (*mastos*, *χονδρος*, 'a cartilage, and *osis*,) A cartilaginous formation in the breast.

MASTODES, (μαστωδης,) Mastoid.

MASTODYN'IA, *Mastal'gia, Mazodyn'ia, (mas-tos, and δυνν, 'pain,')* Pain in the breasts; a form of neuralgia. See *Neuralgia mammae*.

MASTODYNIA APOSTEMATOSA, *Phleg'moné Mam'mæ, Masth'co'sis, Masti'tis apostematosa, Galac-*

lapete'ma, Galactomastoparecto'ma, Absces'sus Mam'mæ seu lac'teus, Phleg'monë Mastodyn'ia, Ab'scess of the Breast, Mam'mary Ab'scess, Milk-abscess. Phlegmonous inflammation of the breast, running on to suppuration, generally in the child-bed female. It is one of the best examples of acute phlegmonous inflammation, and requires the active use of appropriate treatment. A 'broken breast,' in common parlance, means an abscess of the breast, which has suppurated, and opened spontaneously.

MASTODYNIA POLYGALA, Sparganosis—m. Polygala, Mastospargosis.

MAS'TOID, (*μαστοειδής*) *Mastoi'des, Masto'des, Mastoïdeus, Mammilla'ris, Papil'li seu mammilliformis, Mammiform'is, Mam'miform, (F.) Mastoïde, Mastoïdien, (mastos, and idos, 'resemblance.')* Having the form of a nipple. Also, that which relates to the mastoid process, *Mastoïdeus*.

MASTOID AP'ERTURE. The opening of communication between the cavity of the tympanum and the mastoid cells.

MASTOID AR'TERY, *Arte'ria mastoïdea*. A posterior branch of the external carotid artery.

MASTOID CAN'CEr. A kind of firm carcinomatous growth, the section of which is thought to resemble the boiled udder of the cow.

MASTOID CELLS, *Cell'ulæ Mastoïdes, An'trum mastoïdeum, Mas'toid Si'nuses*. These are situate in the mastoid process; communicate with each other, and open into the cavity of the tympanum. Their use seems to be to increase the intensity of sound.

MASTOID FOR'AMEN is situate behind the mastoid process, and gives passage to a small artery of the dura mater, as well as to a vein which opens into the lateral sinus.

MASTOID Fos'sa, (F.) *Gouttière mastoïdienne*, is a depression at the inner surface of the mastoid portion of the temporal bone, which forms part of the lateral sinus.

MASTOID OR DIGAS'TRIC GROOVE, (F.) *Rainure mastoïdienne ou digastrique*, is a groove, situate at the inner side of the mastoid process, which affords attachment to the posterior belly of the digastric muscle.

MASTOID MUSCLE, POSTERIOR, *Splenius*.

MASTOID OR MAM'MIFORM OR MAM'MILLARY PROC'ESS, *Proces'sus mastoïdeus, Apoph'ysis mammilla'ris, Pars mastoïdea*, is situate at the inferior and posterior part of the temporal bone, and gives attachment to the digastric and mastoid muscles.

MASTOÏDE, Mastoid.

MASTOÏDEUS, *Sternocleidomastoïdeus—m. Lateralis, Complexus minor*.

MASTOÏDOCONCHINIEN, *Retrahens auris—m. Génien, Digastricus—m. Hyogénien, Digastricus—m. Oriculaire, Retrahens auris*.

MASTOME'NIA, (*mastos, and μην, 'a month.'*) Vicarious menstruation from the mamma.

MASTON'CUS, *Thelon'cus, (mastos, and ογκος, 'a tumour.')* A tumefaction of the nipple, or of the breast itself. When the tumour is of an adipous character it is termed *Mastopion'cus* or *Mastopimelon'cus*.

MASTONCUS POLYGALACTICUS, *Mastospargosis*.

MASTOPAREC'TAMA, gen. *Mastoparectem'a-tis, (mastos, and παραρτα, 'excessive distension.')* Immoderate distension of the mamme.

MASTOPATH'IA, (*mastos, and παθος, 'disease.'*) An affection of the breast.

MASTOPIMELONCUS, (*mastos, πιμελην, 'fat,' and ογκος, 'tumour,')* see Mastoncus.

MASTOPIONCUS, (*mastos, πινω, 'fat,' and ογκος, 'tumour,')* see Mastoncus.

MASTORRHAG'IA, (*mastos, and ρηνυσι, 'to break forth.')* An unusual flow of milk.

MASTOS, (*μαστος*;) Mamma.

MASTOSCIR'RHUS, (*mastos, and σκίρρως, 'a hard tumour.')* Scirrhus, or hard cancer of the breast.

MASTOPARGO'SIS, *Mastodyn'ia polygala Maston'cus polygalacticus, (mastos, and παραρτα, 'I am full to bursting.')* Fulness of the breasts with milk, so that they are ready to burst.

MASTOSYRINX, gen. *Mastosyrin'gis, Fis'tula mamme, (mastos, and σφύρις, 'a pipe,')* Fistula of the mamma.

MASTURBA'TION, (*μανος, 'hand,' and stupro, 'I ravish,')* *Cinæ'dia, Gar'galz, Gar'galus, Gargalis'mus, Mastupra'tio, Manustupra'tio, Manustupra'tio, Ona'nia, On'anism, Edæogargaris'mus, Edæogargalus, Anaplas'mus, Chiroma'nia, Gau'dia fo'da, Duct'io præputii, Vol'untary Pollutio, Self Pollution, Selfabuse, Secret vice, (F.) Abus de soimême, Attouchement.* Excitement of the genital organs by the hand.

MASTURBA'TOR, *Mastupra'tor, Manustupra'tor, Mascarp'pio*. One given to masturbation.

MASTUS, (*μαστος*;) Mamma.

MAT'SON (F.) A dull or flat sound. The obscure noise, (F.) *Matité*, afforded in certain diseases when any part, as the chest, is percussed. It has been compared to that produced when the thigh is struck. It is opposed to the *Son clair*, or 'clear sound.'

MATCHES DISEASE, see Phosphorus.

MATE, see Ilex Paraguensis.

MATECO, Matico.

MATELLA, (like *Matula*;) Urinal.

MATER, gen. *Matris, (μήτηρ, 'mother,')* Uterus—m. Dura, Dura mater—m. Herbarum, *Artemisia vulgaris—m. Metallorum, Hydrargyrum—m. Mollis, Pia mater—m. Perlum, see Pearl—m. Pia, Pia mater—m. Secalis, Ergot—m. Tenuis, Pia mater*.

MATERIA, (from *mater, 'mother,'* [?]) Matter—m. Fibrosa, Fibrin.

MATE'RIA MED'ICA, *Pharmacolog'ia, Pharmacology, Acol'ogy or Akol'ogy or Acecol'ogy, Hylé Iat'ricé, Iamatolog'ia, Iamatol'ogy, (F.) Matière Médicale*. The division of medical science which treats of the knowledge of medicines; their action on the animal economy, and mode of administration. The study of the *Materia Medica* is one of great importance;—it is a study of the tools with which the practitioner has to work in the cure of disease. Much labour has been spent in contriving classifications of the *Materia Medica*. Some have arranged the articles according to their natural resemblances: others according to their real or presumed virtues: others, according to their active constituent principles. The *Pharmacopœias* place them alphabetically. Perhaps the best classification would be one founded on the agency exerted by the articles on the different tissues; but this arrangement, in the present state of science, is by no means easy; and, moreover, ideas in regard to the action of medicines are so associated with certain terms,—as narcotics, tonics, sedatives, &c., employed to denote certain operations, which they are esteemed capable of producing, that, to abandon them, would be to throw obstacles in the way of the student, without the ultimate advantage accruing to him of possessing a better knowledge of the *modus operandi* of medicines than when a classification, somewhat resembling those usually embraced, is adopted.

The following is the classification given by the Author, in his "General Therapeutics and *Materia Medica*," 6th edit., Philad., 1857:—

- | | | |
|---|---|----------------|
| 1. Agents that affect prominently the alimentary canal or its contents. | { | Emetics, |
| | | Cathartics, |
| | | Anthelmintics. |
| 2. Agents that affect prominently the respiratory organs, | { | Expectorants. |
| | | |

3 Agents that affect prominently the follicular or glandular organs,

4. Agents that affect prominently the nervous system,

5. Agents that affect prominently the organs of reproduction,

6. Agents that affect various organs,

7. Agents whose action is prominently chemical,

8. Agents whose action is prominently mechanical,

Errhines.
Sialogogues,
Diuretics,
Antitithics,
Diaphoretics,
Narcotics,
Tetanus,
Antispasmodics.
Emmenagogues,
Parturificants.

Excitants,
Tonics,
Astringents,
Sedatives,
Refrigerants,
Revellents,
Eutrophics.
Antacids,
Antalkalics,
Disinfectants,
Demulcents,
Diluents.

Of old, the *Materia Medica* consisted of more articles than at present. The tendency, indeed, is, and must be, to diminish it still further; to get rid of those articles which possess no advantages over others equally common, or whose properties are doubtful. In a dictionary, it becomes necessary to insert all that have been reputed to possess virtues; but the majority are unnecessary. The catalogue might be largely reduced with impunity.

MATERIA MEDICA, DYNAMICAL, Pharmacodynamics.

MATERIA MORBO'SA, *M. Pec'cana*, *Mate'ries Mor'bi*, 'Morbid matter.' The matter or material which is the cause of disease.

MATERIA OSSEA, *Terra Ossea*—*m. Peccans*, *M. Morbosa*—*m. Scytdopephica*, *Tannin*—*m. Sialina*, see *Saliva*—*m. Testacea dentium*, see *Tooth*—*m. Urinosa*, *Urea*.

MATERIA LISTA, (from *materia*.) Druggist.

MATÉRIAUX IMMÉDIATS, (from *materia*.) Principles, immediate.

MATERIES, Matter—*m. Morbi*, *Materia morbo'sa*.

MATERNITÉ (F.) (*maternus*, (*mater*, 'mother,' 'maternal.') Formerly, in France, a house for the reception of foundlings; at the present day, an establishment for the reception of pregnant women—a lying-in establishment.

MATHEW'S PILLS, *Pilulæ ex helleboro et myrrha*.

MATHIEU'S VERMIFUGE, see *Vermifuge*.

MATIAS. The bark of a South American tree not yet determined. It is used in its native country in intermittents, and as a tonic generally. Its principal characteristic constituent is a bitter resinous matter. It is probably the same as the *Malambo bark*. The latter has been referred to *Croton Malambo*.

MATICO, (pronounced *mate's'co*.) *Mateco*, *Matica*, *Yerba del Soldado*, *Soldier's weed* or herb. A South American herb.—*Piper angustifolium* seu *longatum*, *Stephens'ia elongata*, *Artan'thè elongata*, *Ord*. *Piperaceæ*—which is possessed of astringent virtues, and is used both internally and externally. It is given in infusion made of one ounce of the leaves—*Matico*, (Ph. U. S.)—to a pint of boiling water, of which the dose is f ʒiiss, or in tincture, made of ʒiiss of the leaves to a pint of dilute alcohol.

MATIERE, Matter—*m. Caséuse*, *Casein*—*m. Cérébriforme*, *Encephaloid*—*m. Extractive du Bouillon*, *Osmazome*—*m. Fécale*, *Fæces*—*m. Fibreuse du Sang*, *Fibrin*—*m. Médicale*, *Materia Medica*—*m. de la Mélanose*, see *Choroidæ tunica*—*m. Noire des Pommans*, *Anthraco'sis*—*m. Perlée de Kerkring*, *Antimonium diaphoreticum*—*m. Pulmonaire noire*, see *Pulmo*.

MATITÉ, see *Mat em*.

MAT'LOCK, MINERAL WATERS OF. *Matlock* is a village in Derbyshire, England, at which there is a mineral spring of the acidulous class. Temperature 66°. It differs but little, except in temperature, from good spring water. It is generally used as a tepid bath.

MATONIA CARDAMOMUM, (after Dr. Wm. Maton, of London,) *Amomum cardamomum*.

MATORIUM, *Ammoniac*, gum.

MATRACIUM, *Matrass*, *Urinal*.

MAT'ASS, *Matracium*. A glass vessel with a long neck; and a round, and sometimes oval, body. It may be furnished with tubulures, or not. It is used in *Pharmacy* for distillation, digestion, &c.

MATRES CEREBRI, *Meninges*.

MATRICAIRE, *Matricaria*.

MATRICA'LIS, *Matricarius*, from *matrix*. Relating to the uterus.

Matricaria are remedies for diseases of the uterus.

MATRICA'RIA, (*matrix*, 'the womb,' so called from its reputed virtues in affections of that organ.) *M. Parthenium*, *Parthenium febrifugum*, *Py'rethrum* seu *Chrysanthemum parthenium*, *Argyrochæ'ta*, *Feverfew*, *Featherfew*, *Motherwort*, *Couthwort*, (Sc.) *Fever-fouille*, (F.) *Matricaire*. *Nat. Ord.* *Compositæ*. The flowers—*Matricaria*, (Ph. U. S.)—resemble, in their properties, chamomile and tansy; and, like them, have been esteemed tonic, stomachic, resolvent, emmenagogue, vermifuge, &c. They are not much used.

MATRICARIA, see *Matricaria chamomilla*.

MATRICARIA CHAMOMIL'LA, *M. Suave'olens*, *An'themis vulg'aris*, *Chamæ'mum Vulg'are*, *Chamomil'la nostras*, *Leucanthemum* of *Dioscorides*, *Wild Corn*, *Dog's Cham'follie*, *German Chamomile*, (F.) *Camomille vulgaire*. The flowers, *Matricaria* (Ph. U. S.), resemble *Matricaria* in properties.

MATRICARIA GLABRA'TA. A South African plant, known at the Cape as *Wild Cham'omile*, has the same properties as the other species of *matricaria*.

MATRICARIA LEUCANTHEMUM, *Chysanthemum* *Leucanthemum*—*m. Suaveolens*, *M. Chamomilla*.

MATRICE, ([L.] *matrix*, gen. *matric'is*.) *Uterus*—*m. Col de la*, *Collum uteri*.

MATRICE DE L'ONGLE, see *Nail*.

MATRIC'ULATE, (*matricula*, diminutive of *matrix*, gen. *matric'is*, 'a roll,' originally 'an army roll or register.') One who is admitted into a university or college, by enrolling or having his name enrolled on the register of the institution. In France, *prendre inscription* means 'to matriculate'; and, in the university regulations of that country, it is required, that the inscription shall be made every three months, until the termination of the prescribed period of study; the student having to take his inscription within the first fortnight of each trimestre or of every three months, and to present himself within the last fortnight of the trimestre to establish the fact of his attendance.

In this country, it is only necessary to matriculate at the commencement of each session.

One who has thus enrolled himself in an institution is called a *Matric'ulate*.

MATRISYLVA, (*mater*, 'mother,' and *sylva*, 'wood,') *Asperula odorata*.

MATRIX, gen. *Matric'is*, (from *μῆρ*, *mater*, 'mother,' 'uterus,') *Cytoblastema*, *Uterus*—*m. Unguis*, see *Nail*.

MATRONA, (from *mater*, 'mother,') *Midwife*.

MATRONALIS, *Viola*.

MATTEE, see *Ilex Paraguensis*.

MATTER, *Materia*, *Hy'le*, *Mate'ries*, (F.) *Matière*. Any substance which enters into the composition of a body. In *Medicine*, it is sometimes applied to the substance of evacuations; and is

also used synonymously with pus. 'To matter' is to suppurate or discharge pus or matter.

MATTING OF PARTS, Hapantismus.

MATULA, Urinal.

MATURATIF, Maturative.

MATURATION, *Maturatio*, (*maturare*, *maturatum*, 'to ripen.') *Pepas'mus*, *Pepan'ria*. Progression of an abscess towards maturity. The state of maturity. Coction.

MAT'URATIVE, *Maturans*, *Pepan'ticos*, (F.) *Maturatif*. A medicine which favours the maturation of an inflammatory tumour.

MATURATUS, Concocted.

MATU'RITY, *Maturitas*, *Hor'e'otes*, (F.) *Maturité*. The state of fruits and seeds, when comparatively developed. State of an abscess, in which the pus is completely formed.

MATURITY, PRECOCIOUS, Precocity.

MAUDLIN, Achillea ageratum—m. Tansey, Achillea ageratum—m. Wort, Chrysanthemum leucanthemum.

MAULES, Rubeola.

MAUVE SAUVAGE, Malva.

MAUX (pl. of *mal*) DE NERFS, Hysteria.

MAW, (Sax. *maga*), Stomach—m. Bound, Constipated—m. Worm, Ascaris.

MAXILLA, (from *mala*, 'the cheek.' [?]) Maxillary Bone.

MAX'ILLARY, *Maxilla'ris*, (F.) *Maxillaire*, (from *maxilla*, 'a jaw,' *maxilla*, 'I chew.') Relating or belonging to the jaws.

MAXILLARY ARTERIES are three in number. 1. *External maxillary*. See Facial. 2. *Internal maxillary*—A. *Guturo-maxillaire* (Ch.). This arises from the external carotid with the temporal. It is remarkable for its complex course, and for the number of branches which it transmits to the deep-seated parts of the face. Immediately after its origin, it buries itself under the neck of the lower jaw, curving inwards and downwards. It then advances directly inwards; proceeding in the space between the two pterygoid muscles towards the *maxillary tuberosity*. It turns again, becomes vertical, and ascends into the bottom of the zygomatic fossa, until, having arrived at the floor of the orbit, it takes a horizontal and transverse direction; enters the sphenomaxillary fossa, and divides into several branches. See Artery, (table.)

MAXILLARY BONE, *Maxil'la*, *Mandib'ula*, *Gam'phē*, *Gam'phelē*, *Gna'thus*, *Mo'la*, *S'i'agon*, *Jaw'bone*, (Prov.) *Chaff'bone*, *Chall*, *Chaille*, *Chavel*, *Chaw'bone*, *Gill*, (F.) *Mâchoire*. A name given to two bones, which support the teeth, and, by means of them, are inservient to the cutting, bruising, and tearing of alimentary substances.

MAXILLARY BONE, INFE'RIOR, *Lower jawbone*, *Os Maxilla'rē infe'rius*, *Maxil'la infe'rior*, *Hypo'si'agon*, (F.) *Os maxillaire* (Ch.), *Mâchoire dia-cranienne*, is a symmetrical, nearly parabolic bone, the middle portion of which is horizontal, and called the *body*; and the posterior is vertical, the angular portions being termed *Rami* or *branches*. These have behind a *parotidian edge*, which forms—by uniting with the base—the *angle of the jaw*. The branches terminate above by two processes, separated by the *sigmoid notch* or *fossa*; the anterior of which is called the *coronoid*; the posterior, the *condyloid process* or *maxillary condyle*, supported by a *Cer'vix*, *Collum*, or *Neck*. The chief parts observed in the lower jaw are—1. *Externally*,—the *Sym'physis men'ti*, *Apoph'ysis men'ti*, *Men'tal fora'men*, and the *external oblique line*. 2. *Internally*,—the *Genian-apoph'ysis*, the *internal* or *My'loid oblique lines*, and the *entrance of the dental canal*. The lower jaw has, also, an *alveolar edge*, which contains alveoli for the reception of the teeth. The lower jawbone is developed by two points of ossification, which unite at the sym-

physis of the chin. It is articulated with the temporal bone and with the teeth.

MAXILLARY BONE, SUPE'RIOR, *Upper jawbone*, *Maxil'la supe'rior* seu *synera'nia*, *Os maxilla'rē supe'rius*, (F.) *Os sus-maxillaire* (Ch.), *Mâchoire syncranienne*. The upper jawbones are to the face what the sphenoid bone is to the cranium. They are articulated with all the portions composing it. They determine, almost alone, the shape of the face, and give it solidity. Their size is considerable; form unequal. They occupy the middle and anterior part of the face; and enter into the composition of the nasal fossæ, orbit, and mouth. The chief parts observable in the upper jaw are—1. *Externally*,—the *Nasal proc'ess*, *Processus nasal'is* seu *frontal'is* seu *ascen'dens*, (F.) *Apophyse montante*, the *superior* or *orbital surface*, *Superfic'ies orbital'is*, *Planum orbitalē*, which forms almost the entire floor of the orbit, the *Fora'men infra-orbital'ium*, *Zygoma'tic proc'ess*, *Canine* or *My'rtiform fossa*. 2. *Internally*,—the *Palatine proc'ess*, *Ante'rior palatine canal*, and the *Antrum of Highmore*. Its circumference is very unequal, and it has behind a round, unequal prominence, called the *Maxillary tuberosity*, which is pierced by the *posterior dental canal*. Anteriorly, there is a notch, which forms part of the anterior aperture of the nasal fossæ, and beneath, an eminence, called the *anterior nasal spine*. The lower part of this circumference forms the *alveolar marg'in*. Each superior maxillary bone is articulated with the ethmoid, frontal, nasal, lachrymal, palatine, inferior spongy, vomer, its fellow, the teeth of the upper jaw, and sometimes the sphenoid bone. It is developed by four or five points of ossification.

MAXILLARY NERVES are two in number, and formed of the second and third branches of the fifth pair. The *Superior Maxillary Nerve*, *Nervus maxillaire* (Ch.), arises from the middle of the gangliform enlargement of the fifth pair; passes forwards, and issues from the cranium through the foramen rotundum of the sphenoid bone; enters the sphenomaxillary fossa, which it crosses horizontally; passes into the infraorbital canal, which it traverses; and makes its exit to vanish on the cheek. It gives off the following branches,—the *orbital*, a branch which goes to the *sphenopalatine ganglion*; *posterior dental branches*; the *anterior dental*,—and terminates in the *infraorbital* nerves, which are divided into *superior*, *inferior*, and *internal*. The *inferior maxillary nerve*, *Nervus maxillaire* (Ch.), is the largest of the three branches furnished by the fifth pair. It issues from the cranium by the foramen ovale of the sphenoid. Having reached the zygomatic fossa, it divides into two trunks; the one *superior* and *external*, which gives off the *temporales profundi*, *masseterine*, *buccal*, and *pterygoids*;—the other—*inferior* and *internal*—the larger of the two, which furnishes the *inferior dental*, *lingual*, and *auricular*.

MAXILLARY VEINS present the same arrangement as the arteries they accompany.

MAXILLO-ALVEOLI-NASAL. Depressor alæ nasi—m. *Labial*, Depressor anguli oris—m. *Narinal*, Compressor naris—m. *Palpebralis*, Orbicularis palpebrarum—m. *Scleroticus*, Obliquus inferior oculi.

MAY APPLE, *Podophyllum peltatum*—m. a. Mountain, *Podophyllum montanum*—m. Flower, *Anthemis cotula*—m. Weed, *Anthemis cotula*.

MAYELLA, *Curcuma longa*.

MAYHEM, *Maïhem*, *Maiming*, (low Latin *ma-hemium*, 'maim or loss of a limb,' old French, *mahem* or *mahe*.) The act of depriving a person of a limb necessary for his defence.

MAYNARD'S ADHESIVE LIQUID, Collo-dion.

MAYS, Zea mays—m. Americana, Zea mays—m. Zea, Zea mays.

MAYWORM, TRUE, *Meloe maialis*.

MAYZ, Zea mays.

MAZA (*μαζα*), Mass, Placenta.

MAZALYSIS, Mazolysis.

MA'ZICUS, (from *maza*.) Relating to the placenta.

MAZISCH'ESIS, (*maza*, and *σχέσις*, 'holding,' 'retention.') Retention of the placenta.

MAZODYNIA, (*μαζος*, 'a breast,' and *δύνη*, 'pain,') Mastodynia.

MAZOITIS, (*μαζος*, 'breast,' and *ίτις*.) Mastitis.

MAZOLYSIS, *Mazal'ysis*, (*maza*, and *λυσις*, 'solution,') Separation of the placenta.

MAZOPATHY'A, (*maza*, and *πάθος*, 'disease,') A disease of the placenta. One originating from the placenta.

MAZOS (*μαζος*), Mamma.

MEAD, ([Sax.] *meo*, *meou*.) Hydromeli, Melizomum.

MEADOW BLOOM, *Ranunculus acris*—m. Fern, *Comptonia asplenifolia*—m. Pride, see *Calumba*—m. Rue, *Thalictrum*—m. Saffron, *Colchicum autumnale*—m. Sweet, *Gillenia trifoliata*, *Spiræa ulmaria*—m. Sweet, American, *Spiræa lobata*—m. Sweet, red, *Spiræa tomentosa*.

MEALY TREE, *Viburnum dentatum*.

MEASLES, ([G.] *Mase*, 'a spot,' *masern*, 'spotted,') *Rubeola*. Also, a diseased condition of pork—*measly pork*—which has been ascribed to the presence of cysticercus cellulose; and may be owing to trichina; see *Trichiniasis*.

MEASLES, BLACK, see *Rubeola*—m. Camp, see *Rubeola*—m. False, *Roseola*—m. French, *Roseola*.

MEASLINGS, *Rubeola*.

MEASLY, Morbillous—m. Pork, see *Measles*.

MEASURE, GRADUATED. A glass measure, used in pharmacy, which is so divided as to indicate accurately the quantities of liquids in dispensing medicines.

MEASUREMENT, Mensuration.

MEASURES, see *Weights and Measures*.

MEASURING, MEDICAL, Mensuration.

MEAT BISCUIT, see *Biscuit*, meat.

MEAT, EXTRACTIVE OF, *Osmazome*.

MÉAT, Meatus.

MEA'TUS, gen. *Meat'us*, (*meo*, *meatum*, 'to go,') *Duct'us*, *Canal'is*, *Po'ros*, *Po'rus*, (F.) *Méat*. A passage or canal.

MEATUS AUDITORIUS EXTERNUS, Auditory canal, external—m. Auditory internus, Auditory canal, internal—m. Cæcus, Eustachian tube—m. Cuticularis, Pori cutanei—m. Cysticus, Cystic duct—m. Narium, see *Nasal fossæ*—m. Seminales uteri, *Tubæ Fallopianæ*—m. Seminars, Corpus Highmori—m. Urinarius, Urethra.

MECHAMECK, *Convolvulus panduratus*—m. Bindweed, *Convolvulus panduratus*.

MECHANE (*μηχανή*), Machine.

MECHANEMA (*μηχανημα*), Machine.

MECHAN'ICAL, *Mechan'icus*, (*μηχανικός*, from *mechane*.) An epithet given to irritating bodies, which do not act chymically—as a *mechanical irritant*.

MECHANICAL or IATROMATHEMAT'ICAL PHYSICIANS, *Iat'romechan'ici* seu *Iat'romathemat'ici*, are such as refer every function, healthy or morbid, to mechanical or mathematical principles;—*Medic'na mechan'ica*.

MECHAN'ICS, AN'IMAL. That part of physiology whose objects are to investigate the laws of equilibrium and motion of the animal body.

MECH'ANISM, *Mechanism'us*. The structure of a body; the collection or aggregate of the parts of a machine; the mode in which forces produce any effect, &c.

MECHANOL'OGY, *Mechanolog'ia*, (*mechane*, and *λογος*, 'a description,') A description of, or treatise on, machines or apparatus used in medicine or surgery.

MÈCHE, (from *myxa*, 'the wick of a lamp or candle,') Tent. This term is usually applied in French surgery to a collection of threads of charpie, cotton or raw silk, united together, which are used for detaching sinuous or fistulous ulcers; or to keep open or enlarge natural or artificial apertures. They are generally applied by means of an instrument called *Porte-mèche*.

MÉCHOACAN DU CANADA, *Phytolacca decandra*.

MECHOACANA NIGRA, (from *Mechoacan*, in Mexico,) *Convolvulus jalapa*—m. Nigricans, *Convolvulus jalapa*.

MECHOACANÆ RADIX, *Convolvulus mechoacan*.

MECON'ETER, (F.) *Mécomètre*, (*μικρος*, 'length,' and *μετρον*, 'measure,') A kind of graduated compass,—*compas de proportion*,—used at the Hospice de Maternité of Paris, to measure the length of newborn infants.

MECON (*μηκων*), Meconium, Opium, Papaver.

MECONICUM, (*μηκων*, 'the poppy,') Opiate.

MECONIS'MUS, *Intoxica'tio opia'ca*; Poisoning by opium.

MECO'NIUM, (*μηκωνιον*, from *μηκων*, 'the poppy,' the inspissated juice of the poppy;) *Pop'py juice*, *Papaver'culum*; Opium. Also, the excrement, *Purgamen'ta Infant'is*, *Me'con*, passed by the infant a short time after birth, which had accumulated in the intestines during pregnancy. It is of a greenish or deep black colour, and very viscid. It seems formed of the mucous secretions of the intestines mixed with bile.

MECONOLOG'IA, *Opiolog'ia*; (*μηκων*, 'opium,' and *λογος*, 'a description,') A treatise on opium.

MECONOPSIS DIPHYLLA, (*μηκων*, 'poppy,' and *σφίς*, 'appearance,') *Stylophorum diphyllum*.

MEDCINARE, Physician.

MEDEA, Genital organs.

MÉDECIN, (from *medeor*, 'to heal,') Physician—m. *Consultant*, Consulting physician—m. *de Fous*, see *Psychiatria*—m. *Piqueur*, see *Piqueur*.

MÉDECINE, *Medicina*—m. *Expectante*, Expectation—m. *Légale*, Medicine, legal—m. *Militaire*, Military medicine—m. *Opératoire*, Surgery, operative—m. *Perturbatrice*, *Perturbatrix* (*Medicina*)—m. *Pneumatique*, *Atmiatria*.

MEDEL'TIS, (*medea*, and *ίτις*.) Inflammation of the genital organs.

MEDELA, (*medeor*, 'to heal,') Curation.

MEDE'NI. A term used by Paracelsus for cutaneous ulcerations like those of boils.

MEDEOLA VERTICILLIFOLIA, (after *Medea*, the sorceress,) *M. Virginica*.

MEDE'OLA VIRGIN'ICA, *M. verticillifolia*, *Gyro'mia Virgin'ica*, *Ind'ian cu'cumber*. Order, *Smilacæ*. An indigenous herb, growing in every part of the United States, the root of which resembles a small cucumber. It has been thought to be diuretic.

ME'DIAN, *Media'mus*; (*medium*, 'the middle,') That which is situate in the middle.

MEDIAN LINE. A vertical line, supposed to divide a body longitudinally into two equal parts; the one right, the other left. Chaussier calls the *linea alba* the *ligne médiane* of the abdomen.

MEDIAN NERVE, *Médian digital*, (Ch.) This nerve arises chiefly from the anterior branches of the last two cervical nerves and first dorsal. The fifth and sixth cervical pairs also send it a branch, which separates from the musculocutaneous nerve. The median nerve descends the inner part of the arm along the biceps muscle. Opposite the elbow

joint it buries itself behind the aponeurosis of that muscle, and engages itself between the two fasciculi of the pronator teres. Lower down, it is situate between the flexors—sublimis and profundus—and passes, with their tendons, under the anterior annular ligament; it then divides into five branches, which are distributed to the muscles of the thenar eminence, to the lumbricales, the integuments of the thumb, the index, middle finger, and outer part of the ring finger. This nerve gives no branches to the arm. In the forearm, it furnishes filaments to all the pronator and flexor muscles; and one of them accompanies the anterior interosseous artery. It also gives off a filament to the integuments of the palm of the hand.

MEDIAN VEINS. Three of the superficial veins of the forearm are so called. The *median basilic*—the *median cephalic* and *common median* or *ful-nis bra'chii*. See Basilic and Cephalic.

MEDIANUM, Mediastium, Mesentery.

MEDIAS'TINAL, Mediastina'lis, (F.) Mé-diastin. Relating to the mediastinum.

MEDIASTINAL ARTERIES are very delicate arterial branches, distributed in the areolar texture of the mediastinum. They are distinguished, according to their situation, into anterior and posterior.

MEDIASTINITIS, Mesodmitis.

MEDIASTI'NUM, Mediasti'nus, (quasi, in *medio stans*, as being in the middle;) *Mesod'mē, Media'nium, Mesotē'chium, Mesotē'chum, Sep'tum thorac'is, Hy'men Diaphrat'ton, Membra'na Diaphrat'ton* seu *thorac'em intersep'iens, Intersep'iēntium thorac'is, Divitō'rium, Dissipien'tium thorac'is, (F.) Médiastin. A membranous space formed by the approximation of the pleurae, dividing the chest into two parts, the one right, the other left. The mediastinum, formed by a double reflection of the pleura, extends from the spine to the posterior surface of the sternum. Its anterior part, called *Anterior mediastinum, Mediastinum pectora'le*, lodges, at its upper part, the thymus gland in the fœtus, and is filled below with fatty areolar tissue. Its posterior part, parallel to the spine, is occupied by the œsophagus, vena azygos, thoracic duct, the lower part of the windpipe, the origin of the bronchia and a number of lymphatic glands. This part is called the *posterior mediastinum—Mediastinum dorsa'le*.*

By some anatomists, Mediastinum is defined to be the pleural walls of the membranous space, whilst the space itself is termed *Cavum mediasti'ni*.

MEDIASINUM AURIS, see Tympanum—*m. Cerebri, Falx cerebri, Septum lucidum—m. Testis, Corpus Highmori.*

MEDIASTINUS, Mediastinum,

MEDICABILIS, (medicor, medicatus, 'to cure or heal,' itself from μέδος, 'care,' and habilis, 'capable,') Curable.

MEDICABLE, Curable.

MED'ICÆ. Female practitioners of old, see Iatrinæ. Also, sworn midwives, whose duty it was to inspect women in cases of suspected pregnancy.

MED'ICAL, Med'icus, Medica'lis, Iat'ricus, Pœo'nus, Pœo'nicus, Phys'ical (of old), *Escula'pian, (F.) Médical, Iatrinque.* Appertaining or relating to medicine or to medicines. See Medicinal.

MEDICAL JURISPRUDENCE, Medicine, legal—m. Man, Surgeonapothecary.

MEDICALIS, Medical.

MED'ICAMENT, Medicamen'tum, Remed'ium, Ace'sis, Aces'sa, Aces'sium, Aces'sys, Al'thos, Hy-gias'ma, Ia'ma, A'cos, Rhy'ma, Medica'men, Phar'macon, Phar'macum, Pharmaceu'ma, Alkar, Auxil'ium, Boëthe'ma, Med'icine, (old E.) Mēdsine; Phys'ic, Rem'edy, Rem'etic, (F.) Remède. A medicine. Any substance exhibited with the view

of curing or allaying morbid action. Medicines are obtained from the three kingdoms of nature, and are divided into *internal* and *external*, according as they are administered internally or applied externally.

In common language, medicine is often called *Stuff, Doctors' stuff*, and, in the Southern States, *Truch*.

MEDICAMENTAL, Medicinal.

MÉDICAMENTER, see Physician.

MÉDICAMENTEUX, Medicinal.

MEDICAMENTO'SUS LA'PIS, Medic'inal stone, (F.) Pierre médicamenteuse. A name formerly given to a mixture of *peroxyd of iron, litharge, alum, nitre, sal ammoniac, and vinegar*; evaporated and calcined at a red heat for an hour. The product was regarded as eminently astringent.

MEDICAMENTUM, Drug, Medicament—m. ex Palmulis, Diaphenicon.

MEDICAS'TER, Medicastra. An ignorant practitioner. A charlatan.

MEDICA'TION, Medica'tio. The change in the animal economy produced by the operation of remedies. Treatment by medicine. The act of impregnating with some medicine.

MÉDICATION HYPODERMIQUE, see Hypodermic—*m. Substitutive,* see Substitutive.

MEDICATRINA, Iatrin.

MEDICAT'RIX, gen. Medicat'ricis. 'Having the power of healing.' A female practitioner.

MEDIC'NA, Ars med'ica seu *Machaō'nica* seu *Machaō'nica, Res med'ica, Iat'rice, Iat'ria, Iat'rica, Iat'rinē, Pœo'nis, Pœo'syne, Pœo'nicē, Iatrotech'nicē, Acesto'ria, Scien'tia med'ica* seu *med'en'di, Sote'ria doct'rina, Med'icine, (old E.) Med'sine, The healing art, Leechcraft or Leachcraft, Phys'ic, (F.) Médecine.* A science, the object of which is the cure of disease and the preservation of health. Occasionally, it is used to comprehend all the branches of the healing art; at others to comprise one great division, in contradistinction to *surgery and obstetrics*. Medicine, in this sense, includes many branches;—the chief of which are, Anatomy, Physiology, Pathology, Therapeutics, Hygiene, Materia Medica, and Pharmacy.

MEDICINA CONSERVATIVA, Hygiene—m. Diæ-tetica, Dietetics—m. Diasostica, Hygiene—m. Eclectica, see Eclectic—m. Efficax, Surgery—m. Equaria, Hippiat'ria—m. Euectica, Gymnastics—m. Forensis, Medicine, legal—m. Gymnastica, Gymnastics—m. Hermetica, see Spagyrista—m. Judiciaria, Medicine, legal—m. Magnetica, Sym-pathy, cure by—m. Mechanica, Kinesipathy, see Mechanical—m. Methodica, see Methodists—m. Militaris, Military medicine—m. Operativa, Surgery—m. Paracelsistica, see Spagyrista—m. Perturbatrix, see Perturbatrix—m. Politica, Police, medical—m. Politicoforensis, Medicine, state—m. Publica, Medicine, state—m. Sinica, Chinese medicine—m. Spagyrica, Chymiat'ria; see Spagyrista—m. State, Police, medical—m. Statica, Statica medicina—m. Sympathetica, Sympathy, cure by—m. Tristitia, Crocus—m. Veterinaria, Veterinary art.

MEDICINABLE, Medicinal.

MEDIC'INAL, Medicina'lis, Medicament'al, Medic'inable, Med'ical, Iat'ricus, Rem'edial, Rem'ediate, (F.) Médicinal, Médicamenteux. Having a remedial power;—as *medicinal plants, &c.* Relating to medicine, as

MEDICINAL DAYS, Medicina'les Di'es. Days on which the ancients considered that remedies might be administered; and especially evacuants. Such days were not esteemed critical.

MEDICINAL HOURS. Times at which—the stomach being freed from its digestive actions—medicine, it was thought, could be most advantageously administered, as in the morning fast-

ing, an hour before dinner, some hours after it, and at bedtime.

MEDICINER, Physician.

MEDICINE, Medicina.

MEDICINE is, also, used in the same sense as Medicament, and for a purging potion.

To Medicine was formerly used for "to restore or cure by medicine."

MEDICINE CHEST, Pharmacotheca.

MEDICINE, CLINICAL, see Clinical.

MEDICINE, CONSERVATIVE, *Medic'na Conservativa*. A term applied by Dr. Flint to that mode of treating disease, which tends to the preservation, development, and support of the vital powers.

MEDICINE, ECLECTIC, see Eclectic—m. Empirical, Arcanum—m. Galenic, Galenism.

MEDICINE, LEGAL, *Med'ical jurisprudence* of some, *Law or Forensic medicine*, *Medic'na forensis* seu *judicia'ria*, *Diatri'a*, (F.) *Médecine légale*. The application of medical knowledge to the solution of every question connected with the administration of justice.

MEDICINE, PATENT, see Patent medicine—m. Political, Police, medical—m. Psychical or Psychological, see Psychical—m. Quack, Arcanum.

MEDICINE SEAL, *Cachet*—m. Stamp, *Cachet*.

MEDICINE, STATE, *Medic'na publica* seu *politicoforensis*. The medical knowledge, brought to bear on state objects, as on public hygiene, and matters appertaining to medical jurisprudence.

MÉDICINIER CATHARTIQUE, *Jatropha curcas*.

MEDICOCHIRURGICAL, *Med'icochirurg'gicus*. Relating or appertaining to medicine and surgery;—as 'medicochirurgical society.'

MEDICOCHIRURGICAL ANATOMY, see Anatomy.

MEDICOLEGAL, *Med'icolegalis*. Relating to legal medicine; as 'a medicolegal inquiry, *Inquisit'io medicolegalis*;'—'a medicolegal inspection,' *Inspect'io med'icolegalis*.

MEDICOMANIA, (from *medicus*, and *mania*.) A mania for the science of medicine without the necessary study.

MEDICOSTATISTICAL, see Statistics, medical.

MEDICUS, (from *medeor*, 'to heal,') Doctor, Medical, Physician—m. Equarius, Hippiator—m. Ocularis, Oculist—m. Ophthalmicus, Oculist—m. Unguentarius, Iatralaptes—m. Veterinarius, Hippiator—m. Vulnerarius, see Deligation—m. Venerum, Surgeon.

MEDIMNUS, (*μεδιμνος*.) An ancient measure, capable of containing about 4 pecks and 6 pints.

MEDINOGRDIUS, *Draconculus*.

MEDITULLUM, (*medium*, 'middle,') Diploë.

MEDIUM MUSCULI, see Muscle—m. Ventris, Umbilicus.

MEDIUS, Middle, Median.

ME'DIUS DIGITUS, *Ver'pus*, *Mid'dle Fin'ger*, (F.) *Doigt du Milieu*, is between the index and ring finger.

MEDLAR, (Sax. *mæb*.) *Mespilus*.

MEDO (Sax.) Hydromeli.

MEDOBLENNORRHŒA, (*medea*, and *blennorrhœa*.) *Leucorrhœa*.

MEDORRHŒA, (*medea*, and *ρρω*, 'to flow,') Gonorrhœa—m. Feminarius insons, *Leucorrhœa*—m. Virilis, Gonorrhœa.

MEDSINE (old E.), Medicament, Medicina.

MEDULLA, (from *medius*, 'in the middle,') Marrow—m. Cerebri, see Cerebrum—m. Dorsalis, M. Spinalis—m. Dorsualis, M. Spinalis—m. Nervorum, Neurine.

MEDULLA OBLONGATA, *Oblong medulla*, *Cerebral protuberance*, *Nervous system of the senses*, (Gall), *Cerebrum elongatum*, *Mesocephalon*, (F.) *Mésocéphale*, *Moëlle allongée*, *Mésencéphale*, (Ch.)

The medullary substance that lies within the cranium upon the basiliary process of the occipital bone. The anterior surface which rests in the basiliary groove, is impressed by the basiliary artery. At the upper extremity, and on its posterior surface are the *Tubercula quadrigemina*. At the same extremity, the medulla gives rise to two prolongations, the *Peduncles of the brain*, separated from each other by the mammillary eminences, and becoming lost in the optic thalami. The lower extremity is called the *tail* or *Rachidian bulb*, *Bulbus medullæ spinalis* seu *rachidicus* (F.) *Bulbe rachidien*, and is continuous with the medulla spinalis. It is to this part only that some anatomists apply the name *Medulla oblongata*. From the posterior angles two other prolongations arise, called *Peduncles* of the cerebellum. The medulla oblongata has several eminences—the *Pons varolii*, *Corpora pyramidalia*, *C. olivaria*, &c.—and it has a longitudinal fissure before, and another behind, called the *anterior* and the *posterior median fissure*. The vesicular neurine in the centre of the medulla is the nervous centre of respiration and deglutition: hence it has been called *centrum vitale*.

MEDULLA OSSIIUM, Marrow—m. Spinae, M. spinalis.

MEDULLA SPINALIS seu *Spinae seu dorsualis* seu *vertebralis* seu *dorsalis*, *Cor'da spinalis*, *Fu'nis argenteus*, *My'elus*, *M. rhachites* seu *notius*—*us* seu *diache'nus* seu *pro'tes*, *My'elon*, *Notomy'elus*, *Rhachites*, *Æ'on*, *Processus rachidianus*, *Sac'ra fistula*, *Spinal prolongation*, *Vertebral marrow*, *Spinal cord* or *mar'row*, *Nervous system of voluntary motion* and *tactile impression* (Gall), (F.) *Moëlle épinière*, *Moëlle vertébrale*, called by Ch., *Prolongement rachidien* of the encephalon, is the continuation of the medulla oblongata. It commences at the foramen magnum of the occipital bone, and descends in the vertebral canal as low as the 2d lumbar vertebra, without filling it; presenting, in its course, several evident enlargements. It is grooved on both its anterior and posterior surfaces by a furrow, which divides it, in its whole length, into two great nervous cords, intimately united with each other. It terminates by an oval tubercle, whence a number of nerves set out called *Cauda equina*, (F.) *Queue de cheval*, from its resemblance to a horse's tail. The spinal marrow has no analogy, as regards its structure, with the marrow of long bones. It is formed of two substances; one white, the other gray, presenting an inverse arrangement to that which they have in the brain, the white being external, the gray at the centre, *Substantia grisea centralis*. It is enveloped by a yellowish, fibrous membrane, very resisting, which seems to be continued insensibly from the pia mater, and by two other membranes, which are merely prolongations of the arachnoid and dura mater. See Nerve.

MEDULLA VERTEBRALIS, M. spinalis.

MEDULLÆ (pl. of medulla,) Sperm.

MEDULLARY, *Medullaris*, (F.) *Médullaire*, (from medulla.) Relating to the marrow, or analogous to marrow.

MEDULLARY ARTERIES. The arteries, which enter bones and pass to the marrow.

MEDULLARY CANAL, see Canal, medullary—m. Cells, see Medullary Membrane—m. Juice, Marrow—m. Matter, see Nerve-fibres—m. Matter of the brain, see Cerebrum.

MEDULLARY MEMBRANE, *Periosteum Inter'nun*, *Endosteum*, *Perymy'elis*. A vascular, areolar web of extreme tenuity, which envelops the marrow and lines the inner surface of the medullary canal of the long bones. This membrane has been considered as a species of internal periosteum of those bones. It has numerous vessels, which bury themselves in the thickness

of the marrow; and others which nourish the innermost plates of the bone. The cells formed by it are termed *Cellule seu Sacculi medullares*.

MEDULLARY NEURINE, see Neurine — m. Sarcoma, Encephaloid, Hæmatodes fungus — m. Sheath, see Nervefibres.

MEDULLARY SUBSTANCE OF THE KIDNEY, see Kidney — m. s. of Schwann, see Nerve-fibre.

MEDULLARY SYSTEM. Bichat gives this name to the marrow and its membranes. He distinguishes two species of medullary systems; the one occupies the cellular tissue at the extremities of the long bones and that of the flat and short bones; the other is found merely in the central canal of long bones.

MEDULLARY TUMOUR OF THE LUNGS, Phthisis, cancerous.

MEDULLITIS. (*medulla* and *itis*.) Myelitis. M. Gerdy uses *Medullite* for inflammation of the marrow of bones, *Osteomyelitis*; — *Myelitis* being usually appropriated to inflammation of the spinal marrow.

MEERAN, *Daucus carota*.

MEGALANTHROPOGENESIS, (*megalo*, *av-Spas*, 'man,' and *genesis*, 'birth.') A term used by a French physician, named Robert, to designate the art of procreating great men; men of mind; men of genius. He considered that they may be perpetuated by always taking care to have talented men united to clever women!

MEGALO, (from *μεγας*, *μεγαλη*, 'great.') In composition, great.

MEGALOCARDIA, (*megalo*, and *καρδια*, 'heart.') The state of having a very large heart.

MEGALOCÆLIA, (*megalo*, and *καλια*, 'abdomen,') Intestine, great, Megalosplanchnia.

MEGALOMELIA, (*megalo*, and *μελος*, 'a limb.') A monstrosity characterized by very large limbs.

MEGALOPHONIA, (*megalo*, and *φωνη*, 'voice.') The condition of having a full, strong voice.

MEGALOPHTHALMUS. (*megalo*, and *οφθαλμος*, 'eye.') A congenital deformity, in which the eye is inordinately large.

MEGALOPUS, (*megalo*, and *πους*, 'a foot,') Macro-podius.

MEGALOSPLANCHNUS, (*megalo*, and *σπλγχχνον*, 'a viscus.') *Megalocælia*. Hippocrates applies the epithet to those in whom a viscus, or the viscera, are tumid. Some have used the substantive *Megalosplanchnia* for the tumour itself.

MEGALOSPLENIA, (*megalo*, and *σπλην*, 'the spleen,') Splenonecus.

MEGETHOS, (*μεγεθος*, from *μεγας*, 'great,') Stature.

MEGRIM, (from [F.] *migraine*, itself abridged from *hemicrania*.) Hemicrania.

MEIBOMIUS, GLANDS OF, *Follicles of M.*, *Palpebral* or *Ciliary Follicles*, *Glandulæ sebaceæ ciliares* seu *Meibomia-næ*, *Intestinalia Meibomia-næ*, *Lacrimæ palpebrarum* (F.) *Follicles palpebræ* ou *ciliares*, *Glandes de Meibomius*. Small sebaceous follicles, called after Meibomius, although known long before his description of them. They are situate in special grooves in the tarsal cartilages; and are ranged by the side of each other, in the form of yellowish, parallel, and vertical lines; sometimes straight, at others, tortuous. They may be seen at the inner surface of the eyelids, through the conjunctiva, and secrete a sebaceous humour, called *Humor of Meibomius*, *Lip-piti'do*, *Gum*, (F.) *Chassie*.

MEIOSIS, (*meiosis*, from *μειωω*, 'I lessen,') *Meo-sis*, *Mio-sis*, *Minu-tio*, *Minutho-sis*, *Minutho-sis*. The period of a disease, in which the symptoms begin to diminish; — the decline. Remission. Also, idiotism.

MEIURUS, (*μειουρος*, from *μειωω*, 'to lessen,' and *ουρα* 'a tail,') Decurtatus, Myurus.

MEKILWORT. *Atropa belladonna*.

MEL. gen. *Mellis*, (*μελι*.) *Meli*, *Nili-acum*, *Honey*, (F.) *Miel*. A liquid of a mucosaccharine nature, prepared by the *Apis mellifica* or common bee, which collects the materials from the nectaries of flowers. Honey is employed as aliment, condiment, and medicine. It is demulcent and aperient; and is prescribed as an adjunct to gargles in cynanche tonsillaris, &c. It is, at times, used as a detergent to foul ulcers. *Virg'in Honey*, *Mel virginum* seu *virginæum*, (F.) *Miel vierge*, is that which flows from the wax spontaneously.

HILL'S BAL'SAM OF HONEY is formed of *balsam of tolu*, ℞j, *honey*, ℞j, *rectified spirit*, one gallon. It was long a celebrated empirical pectoral.

HONEY WATER is a mixture of essences coloured with saffron. A little honey is added to communicate a clamminess; the effect of which is to make it retain the scent the longer. It is used as a scent.

MEL ACETATUM, Oxymel — m. *Ægyptiacum*, *Ægyptiacum* — m. *Aërium*, *Fraxinus ornus* — m. *Arundinaceum*, *Saccharum*.

MEL BORACIS, *M. subboracis* seu *boraxatum* seu *Sodæ Boracis*, (Ph. U. S. 1863.) *Colluto'rium adstrin-gens*, *Limimentum de Boracæ* seu *ad aph-thas*. *Linctus de Boracæ* seu *ad apthas*, *Honey of Boracæ* or of *Borate of Soda*, (F.) *Mellite de borax*, *Miel boratè*. (*Boracis contrit.* ʒj, *Mellis depumati*, ʒj.) Detergent. Applied to the mouth in aphthous affections.

MEL CANNE. *Saccharum* (non purificatum) — m. *Coctum*, *Pelicide*.

MEL DESPUMATUM, *M. Depura'tum*, *Clar'ified Honey*, *Anaphrom'eli*, (F.) *Miel clarifié* ou *despumé* ou *dépuré*, *Mellite simple*, *Sirap de miel*. (Melt the honey in a waterbath, and remove the scum.) Uses the same as honey.

MEL ROSÆ, *M. rosatum*, *Mellitum de Rosæ*, *Rhodome'li*, *Rhodostacton*, *Honey of Rosæ*, (F.) *Miel rosat*, *Mellite de Rosæ*. (*Rosæ gallicæ* in pulv. ʒij, *Mellis depum.* fʒxxv, *Alcohol*, dilut. q. s. Gradually pour the diluted alcohol on the powder in a percolator until six fluidrachms have passed. Set this aside, and continue the percolation until half a pint more has passed. Evaporate to ten fluidrachms; add the liquid set aside, and mix the whole with the clarified honey. Ph. U. S. 1863.) Astringent and detergent. Used chiefly in gargles and washes in apthæ, &c.

MEL SCILLÆ, Oxymel scillæ — m. *Scillæ compositum*, *Syrupus scillæ compositus* — m. *Sodæ boracis*, *Mel boracis* — m. *Vinosum*, *Onomel* — m. *Virginum*, see Mel.

MELENA, (*μελαινα*, 'black,') *Melanorrhag'ia*, *Melenorrhag'ia*, *Melenodiarrhœa*, *Icterus nig'er*, *Melancho'rus*, *Melanicterus*, *Mélas Icterus*, *Melanism'us*, *Morbus nig'er* seu *nig'er Hippoc'ratis*, *Enterorrhœa*, *Black Jaun'dice*, (F.) *Mélena*, *Mé-lène*, *Mélanictère*, *Maladie noire*, *Méline*, *Ictère noir*. A name given to vomiting of black matter, ordinarily succeeded by evacuations of the same character. It seems to be often a variety of hæm-temesis. The *Black Vomit* in yellow fever is owing to a morbid secretion from the lining membrane of the stomach and small intestines mixed with blood. Melena also signifies hemorrhage from the intestines; *Fluxus sple'nicus*, *Dysenter'ia sple'nica*, *Dejectio'nes nig'æ*, *Seces'sus nig'er*.

MELÆNA FUNGOSA CARCINODES, Hæmatodes fungus.

MELÆNODIARRHŒA, Melæna.

MELÆNAGO'GUS, (*μελαινα*, 'black,' and *αγω*, 'I drive away,') A medicine believed to be capable of expelling atrabilis or 'black bile.'

MELÆNORRHAGIA, (*μελαινα*, 'black,' and *ῥηγνμι*, 'to break forth,') Melæna.

MELAINÉ, (from *μελαινα*, 'black,') see Choroidea tinea.

MELALEUCA CAJUPUTI, (μελας, 'black,' and λευκος, 'white,' from the trunk being black and the branches white.) *M. Myrtor* seu *Leucaeden'dron*, *Myrtus Leucaeden'dron*, *Leptospermum Leucaeden'drum*. Ord. Myrtaceæ. This plant affords the *Cajuput Oil*, (see *Cajuput*.) The leaves are esteemed diuretic, stomachic, and emmenagogue.

MELALEUCA LEUCADENDRON, *M. Cajuputi*—m. Minor. *M. Cajuputi*.

MELAMPONUS, (μελας, 'black,' and φωνη, 'voice,') Having a hoarse or indistinct voice.

MELAMPHYLLUM, (μελαμφύλλον, from μελας, 'black,' and φύλλον, 'leaf,') *Acanthus mollis*.

MELAMPODE, *Helleborus niger*.

MELAMPIDIUM, μελαμπίδιον, after the shepherd Melampus, or from μελας, 'black,' and πους, gen. ποδος, 'foot,' in allusion to the seeds,) *Helleborus niger*.

MELANÆMA, (melano, and αἷμα, 'blood,') Suffusion.

MELANÆMIA, Venosity. A morbid condition in which the vessels contain an unusual quantity of pigment.

MELANAGOGUE, *Melanagógus*, (melano, and αγω, 'I expel,') A medicine which the ancients believed adapted for expelling black bile or melancholy.

MELANCHLO'RUS, (melano, and χλωρος, 'green,') The ancients gave this name to certain darkcoloured topical remedies. See *Melæna*.

MELANCHOLE, *Atrabilis*.

MELANCHOLIA, (μελαγχολία, from melano, and χολη, 'bile,) Melancholy—m. Autochirica, Suicide—m. Errabunda, Cuttubuth, Kutubuth—m. Flutuosa, Hypochondriasis—m. Hypochondriaca, Hypochondriasis—m. Nervea, Hypochondriasis—m. Pleonectica, see *Pleonectica*—m. Sultans, Chorea—m. Suicidium, Suicide—m. Uterina, Nymphomania—m. Zoanthropia, Melancholy.

MELANCHOLIE, Melancholy.

MELANCHOLIC, (μελαγχολικός,) *Melanchol'icus*, *Melancholus*, *Melanchol'des*, (Sc.) *Dram*, *Drum*, (F.) *Mélancolique*, *Mélancolique*. One labouring under melancholy. That which belongs or relates to melancholy. In popular language, one of a gloomy disposition.

MELANCHOLINESS, Melancholy.

MELANCHOLIQUE, Melancholic.

MELANCHOLODES, Melancholic.

MELANCHOLUS, (μελαγχολος,) Melancholic.

MEL'ANCHOLY, *Melanchol'ia*, *Lypemania*, *Ephro'nia* *Melanchol'ia*, *Ma'nia* *Melanchol'ica* seu *Melanchol'ia*, *Tristema'nia*, *Barythym'ia*, *Hypoleptoma'nia*, *Anoë'ia* adstricta, *Fixa'tio monaca'ia*, *Mel'ancholiness*, (Sc.) *Dismal*; (F.) *Mélancolie*, *Mélancolie*, *Lyperophrénie*.

A disease supposed, by the ancients, to be caused by black bile. A variety of mental alienation, characterized by excessive gloom, mistrust, and depression, generally, with insanity on one particular subject or train of ideas, *Monoma'nia*, *Monoma'ria*, or on a few subjects, *Oligoma'nia*, *Panophro'bia*; *Demonoma'nia*, *Erotoma'nia*, *Nostal'gia*, &c., may be referred to this head.

Melancholy is also used for unusual gloominess of disposition.

MELANCHOLY, EROTIC, Erotomania—m. Love, Erotomania.

MELANCHRUS, *Melan'ochrus*, (melano, and χρος, 'a colour,') One attacked with black jaundice. Supposed, by some, to be an abbreviation of *Melanchlorus*.

MÉLANCOLIE, Melancholy.

MÉLANCOLIQUE, Melancholic.

MELANÉ, Melanic.

MELANÉMIE, Venosity.

MELANEPHIDROSIS, (melano, and ephidrosis,) Sudor Anglicus niger.

MÉLANGE, *Mistura*—m. *Pectoral*, *Mistura acidi hydrocyanici*.

MELANIC, *Melan'icus*, *Melanot'ic*, *Melanot'icus*, (F.) *Mélané*, *Mélanique*, *Mélanotique*; (from μελας, gen. μελανος, 'black,') Of or belonging to Melanosis;—as *Melan'ic deposit*, a black colouring matter deposited from the blood under special circumstances:—see *Melanosis*.

MELANIC DEPOSIT, Melanosis.

MELANICTERUS, (melano, and icterus, 'jaundice,') *Melæna*.

MÉLANIENNES (*Taches*), *Nævus*.

MÉLANINE, see *Choroidea tunica*.

MÉLANIQUE, Melanic.

MÉLANISME, Melanismus.

MELANISMUS, *Melæna*. M. I. G. St. Hilaire employs the term *Melanisme*, for the change of coloration of the skin in which the pigment is of a deeper hue and in greater quantity than natural.

MEL'ANO, (from μελας gen. μελανος, 'black,') in composition, black, obscure.

MELANOCARCINOMA, Cancer, melanotic.

MELANOCHRUS, Melanchrus.

MELANOMA, (melano, and oma,) Melanosis—m. *Pulmonum*, see Melanosis.

MELANOMYCES, (melano, and μυκης, 'a fungus,) *Melanospongus*.

MELANOPATHIA, (melano, and παθος, 'affection,') *N'igritism*, *Nigrit'ies*. A disease of the skin, which consists in augmentation of black pigment; generally in patches.

MELANOPHYMA, gen. *Melanophy'matis*, (melano, and φυμα, 'a swelling,') *Ergot*.

MELANOPIPER, (melano, and πιπερι, 'pepper,') *Piper nigrum*.

MELANOPNEUMON, (melano, and πνευμων, 'lung,') see Melanosis.

MELANORRHAGIA, (melano, and ραγη, 'a breaking forth,') *Melæna*.

MELANORRHIZUM, (melano, and ριζα, 'a root,') *Helleborus niger*.

MÉLANOSE, Melanosis.

MELANO'SIS, *Melan'is*, *Melano'ma*, *Nigritu'do*, (F.) *Dégénérescence noire*, *Mélanose*; (μελας, gen. μελανος, 'black,' and οσις,) An organic affection, in which the tissue of the parts is converted, owing to a melan'ic deposit, into a black, hard, homogeneous substance, near which ulcers or cavities may form;—owing to the softening, either of the substance itself, or of some other morbid tissue,—of tubercles especially, This morbid change affects the lungs particularly; when it is called *Melano'sis* seu *Melano'ma* seu *Anthraco'sis Pulmo'nium*, *Melanopneumon*, *Nigritu'do melano'ica*, *Melan'is Pulmo'nium*, *Pneumonmelano'sis*, *Black tu'bercle*, (F.) *Mélanose Pulmonaire*. It is, also, met with in the liver and areolar texture. Its causes are very obscure. Melanosis of the lungs constitutes one of the species of phthisis of Bayle; but it is impossible to distinguish it from the other species during life.

In the classification of Fuch, *Melanosen* (G.) forms a family of diseases.

The deposition of melanic matter in the tissues is not necessarily connected with malignant disease. See Cancer, melanotic.

MELANOSIS PULMONUM, see Melanosis—m. *Universalis*, Venosity.

MELANOSMEGMA, (melano, and σμυγμα, 'the act of cleansing,') see Sapo.

MELANOSPONGUS, *Melanomy'ces*, (melano, and σπγγος, 'sponge,') The tuberiform variety of melanosis of the lungs.

MELANOTHANATUS, (melano, and θανατος, 'death,') Death, black.

MEL'ANOTHRIX, *Atricapillus*, (F.) *Mé'yno*.

trigue, (*melano*, and *σπῆξ*, 'hair.') Having black hair.

MELANOTIC, *Melanic*—*m.* Cancer, Cancer, melanotic.

MELANSIS, *Melanosis*—*m.* Pulmonum, *Melanosis pulmonum*.

MELANTHIUM, (*melano*, and *ανθος*, 'flower,') *Nigella*, *M. Virginicum*—*m.* Muscætoxicum, *Amianthium muscætoxicum*.

MELANTHIUM VIRGINICUM, *Melanthium*, (*melano*, and *ανθος*, 'flower,') *Leimanthium Virginicum*, *Quas'adil*; indigenous; Order, *Melanthaceæ*, flowering in July, is said to be a sure but severe remedy for itch.

MELAS, (*μελας*, 'black,') *Lepra nigricans*—*m.* Icterus, *Melena*—*m.* Zomos, *Jusculeum nigrum* *Lacedæmoniorum*.

MELAS'MA, *gen. Melas'matis*, (*μελασμα*,) *Echymoma* *Melas'ma*, *Nig'ror*. A black spot or ecchymosis, occurring on the lower extremities of old people especially. Also, a cutaneous affection analogous to chloasma, differing from it only in the dark colour of the morbid pigment. It is also called *Pityriasis nigra*.

MELASMA SUPRARENALÉ, see *Capsule*, renal.

MELAS'SES, *Molas'ses*, *Melus'tum*, *Theriaca communis*, *Syrupus empyreumaticus* seu *niger* seu *fuscus*, *Sac'chari Fæx*, *Tre'acle*, (Old Eng.) *Triacle*; (F.) *Mélasse*, (from *mel ustum*, 'burnt honey,' or from *μελας*, 'black,') The syrup, which remains after the juice of the sugar-cane has been subjected to all the operations for extracting sugar from it. In the United States, the syrup, made in the process of forming common sugar, is called *melasses*; that remaining after the refinement of sugar being termed *sugarhouse melasses* or *treacle*.

MELASSES SPIRIT, Rum.

MELATROPH'IA, (*μελος*, 'a limb,' and *ατροφία*, 'wasting,') Wasting of the limbs.

MELCA, (*μελκα*,) Food made of milk, coagulated by boiling vinegar.

MELE, (*μηλη*,) *Specillum*.

MELEA, (*μηλις*, 'the apple-tree,') *Pyrus malus*.

MELEGUETTA, *Amonium grana paradisi*.

MELEI'OS, (*μηλειος*,) *Meli'na*, *Meli'sa*. A species of alum found in the island of Melos.

MÉLÉ'NA, *Melæna*.

MÉLÈNE, *Melæna*.

MÉLÈZE, *Pinus larix*.

MELI, (*μελι*,) *Mel*.

MELIA, (*μελία*,) *Fraxinus ornus*.

MEL'IA AZED'ARACH, *Azedara'cha amœ'na*, *Poinberry Tree*, *Azed'arach*, *Pride of India* or of *China*, *Pridetree*, *Hoptree*, *Bendtree*. *Nat. Ord.* *Meliaceæ*; *Ser. Syst.* *Decandria Monogynia*. The bark of the root—*Azedarach*, (Ph. U. S.)—is usually given in decoction, in the proportion of three or four ounces of the bark of the fresh root to a quart of water, boiled down to a pint. The dose is one or two tablespoonfuls, every two or three hours, till purging is induced. It is given in this manner as an anthelmintic. It is sometimes formed into an ointment, and used in tinea capitis. The tree is a native of Syria, Persia, and the north of India, and is abundant in the southern states of the Union.

The bark of *Melia azadirach'ta*, *Azadirach'ta Indica*, *Neem Tree*, *Margo'sa Tree*, is bitter and astringent, and is prescribed, in India, as a tonic and antiperiodic. The seeds, and oil obtained from the pericarp, have been regarded as anthelmintic; the bark of the root as an emmenagogue; the gum, which exudes from it, as an aphrodisiac; and the leaves have been applied in the form of poultice to ulcers, and cutaneous affections.

MELIA TERRA, (from the island Melos,) *Creta*.

MELIANTHUS MAJOR, (*meli*, and *ανθος*, 'a flower,') *Nat. Ord.* *Zygophyllæ*. A south Afri-

can plant, a decoction of whose leaves is a good external remedy in tinea capitis, crusta serpigiosa, necrosis and foul ulcers. It is also useful as a gargle and lotion in sore throat and diseases of the gums. The bruised leaves applied to ulcers promote granulation.

MELICERA, (*meli*, and *κeros*, 'wax,') *Porrigio favosa*.

MELICERIA, *Porrigio favosa*.

MELICE'RIS, *Melice'rium*, *Emphy'ma encystis melice'ris*. An encysted tumour filled with a substance resembling honey. Also, *Porrigio favosa*.

MELIC'EROUS, *Melicer'itos*. Having the characters of meliceris,—as a *melicerous* tumour.

MELICERUM, *Porrigio favosa*.

MELICHEI'UM, (*meli*, and *χεω*, 'I pour out,') A honeylike discharge from an ulcer.

MELICRATUM, (*μελικρατον*, from *meli*, and *κραω*, 'to mix,') *Hydromeli*.

MELIFAVIUM, (*meli*, and *favus*, 'a honeycomb,') *Meliceris*, *Porrigio favosa*.

MELIGE'I'ON, (*μελι*, 'honey,') A fetid humour of the consistence of honey, discharged from ulcers, accompanied with caries of a bone.

MÉLILOT, *Trifolium melilotus*.

MELLILOTUS, (*μελιλωτον*,) *Trifolium melilotus*.

MELIME'LUM, (*meli*, and *μηλον*, 'an apple,') A name given to two compounds:—one of honey and quince, the other of honey and apples.

MÉLINE, *Melæna*.

MELINI SULPHAS, (from the island Melos,) *Cadmii sulphas*.

MEL'NUM, (*μηλων*,) The ancient name of an ointment, and of several plasters, described by Dioscorides and Galen; so called from their resembling the *μηλον* or quince in colour.

MELINUM SULPHURICUM, *Cadmii sulphas*.

MELINUS, *Meleios*.

MÉLIOLE, *Mellitum*.

MELIPHYLLUM, (*meli*, and *φυλλον*, 'a leaf,') *Melissa*.

MELIS'SA, (*μελισσα*, 'a bee,' because bees gather honey from it) *M. officinalis* seu *Roma'na* seu *hirsuta* seu *citrina* seu *citra'ta* seu *hortensis*, *Citra'go*, *Citra'ria*, *Melitei'a*, *Melissob'otos*, *Melite'na*, *Melissobot'anum*, *Melissophyll'um*, *Meliphyl'um*, *Men'tha citra'ta*, *Melitt'ia*, *Cedronel'la*, *Apia'strum*, *Balm*, (F.) *Mélisse*, *La Citronelle*, *Céline*; *Nat. Ord.* *Labiatae*. The leaves of balm, *Melis'sa* (Ph. U. S.), have a pleasant odour; and an austere and aromatic taste. Balm was formerly much used in nervous diseases, but is now only employed when made into tea, as a diluent, in febrile affections. It is the basis of a celebrated preparation,—the *Eau des Carmes*, *Aqua Carmelit'na*, *Car'melite water*.

MELISSA CALAMIN'THA, *Calamin'tha*, *C. vulgaris* seu *officinalis* seu *grandiflora*, *Thymus calamin'tha*, *Cal'amin't*, *Nespite*, (F.) *Calament*. This plant smells like wild mint, though more agreeably. It is used, popularly, as a tea in dyspepsia; flatulent colic; hysteria; uterine obstructions, &c.

MELISSA CANARIENSIS, *Dracocephalum Canariense*—*m.* *Citrata*, *Melissa*—*m.* *Citrina*, *Melissa*—*m.* *Clinopodium*, *Clinopodium vulgare*.

MELISSA GRANDIFLO'RA, *Thymus glandiflorus*, *Calamin'tha mag'no flo're*, *C. mont'na*, *Mount'ain Cal'amin't*. This plant resembles the last in virtues.

MELISSA HIR'SUTA, *Melissa*—*m.* *Hortensis*, *Melissa*—*m.* *Humilis*, *Melittis melissophyllum*.

MELISSA NEP'ETA, *Thymus nep'eta* seu *multiflora*, *Calamin'tha Ang'lica* seu *Pule'gii odorata* seu *nep'eta* seu *parviflora* seu *trichot'oma*, *Nep'eta agrest'is*, *Field Cal'amin't*, *Spot'ted Cal'amin't*. Formerly used as an aromatic.

MELISSA OFFICINALIS, *Melissa*—*m.* *Pulegioides*, *Hedeoma*—*m.* *Pennana*, *Melissa*—*m.* *Tragi*,

Melittis melissophyllum — m. Turcica, Dracocephalum Canariense.

MELISSE DES BOIS, Melittis melissophyllum — m. de Moldavie, Dracocephalum Canariense.

MELISSOBOTANUM, (melissa, and βοτανή, 'a plant,') Melissa.

MELISSOBOTOS, (melissoβotos, from melissa, 'a bee,' and βoskw, 'to feed,') Melissa.

MELISSOPHYLLUM, (melissa, and φύλλον, 'a leaf,') Melissa, Melittis melissophyllum.

MÉLISSOT, Melittis melissophyllum.

MELITÆMIA, Melithæ'mia, Mélithémie, (meli, and αίμα, 'blood,') A condition of the circulating fluid, in which it contains an unusual quantity of saccharine matter.

MELITÆNA, (μελιτανα,) Melissa.

MELITAGRIA, (meli, and αγρα, 'a seizure,') Impetigo.

MELITEIA, Melissa.

MELITHÆMIA, Melitæmia.

MELITHYPERURIA, (meli, υπερ, 'in excess,' and ουρον, 'urine,') Diabetes.

MELI'TIS, Inflammat'io ge'næ, (μυλον, 'the cheek,' and itis.) Inflammation of the cheek.

MELITITES, (μελι, 'honey,') Hydromeli.

MELITOPTYA LISM, Mel'itoptyalis'mus, Ptyalis'mus mell'itus, (from meli, and ptyalism.) A copious secretion of sweet tasted saliva.

MELITTA, (μελιττα,) Bee.

MELITTIS, Melissa.

MELI'TIS MELISSOPHYLLUM, La'mium montanum, Meli'sa hu'milis seu sylvest'ris seu Tra'gi, La'mium Plin'ii, Melissophyll'um, Bostard Balm, (F.) Mélisse des Bois, Méli'sot. Ord. Labiate. This plant was formerly employed like the balm in uterine obstructions.

MELITTIS SYLVESTRIS, M. Melissophyllum.

MELITURIA, (meli, and ουρον, 'urine,') Urine, diabetic, see Diabetes.

MELI'TUS, Mell'itus, (F.) Miellé, (from meli.) Appertaining to honey. Of the nature of, or sweetened with, honey.

MELIZOMUM, (meli, and ζωμος, 'broth,') Mead, Mell'ina, Metheglin, (Prov.) Mullin. A drink prepared with honey.

MELLA'GO, (from meli.) Any medicine having the consistence of honey. A fluid extract.

MELLÉOLE (F.), Mellitum.

MELLICRATUM, (meli, and κραω, 'to mix,') Hydromeli.

MELLIFAVIUM, (meli, and favus, 'a honey-comb,') Porrigo favosa.

MELLINA, (from meli,) Melizomum.

MELLITE, Mellitum — m. de Borac, Mel boracis — m. de Roses, Mel rosæ — m. Simple, Mel depumatum.

MELLI'TUM, (from mel,) (F.) Miel médicinal, Mellite, Hydromellé, Saccharolé liquide, Melliolé, Melliolé. A prepared honey: — a medicated honey.

MELLITUM DE ACETATE CUPRI, Linimentum Æruginis.

MELLITUM DE MERCURIA'LI COMPOS'ITUM, (F.) Miel de mercuriale composé, Syrop de longue vie. (Succ. purif. mercurialis lbij, Boraginis, Anchusæ officinalis, aa lbss, Iris pseudocori radicea recent. ℥ij. Rad. sicc. gentiane ℥j, Mellis lbij, Vini albi ℥xiiij. Macerate the bruised roots in the wine for 24 hours; strain; mix the juices and honey; boil slightly, and filter; then add the two liquors, and boil to the consistence of syrup. Ph. P.) Cathartic, stomachic, &c.

MELLITUM DE ROSIS, Mel rosæ.

MELLITUM SIM'PLEX, Syr'upus de Mell'le, Syr'up of Honey. (Mellis lbvj, Aquæ Biss, Carbonat. calcis preparat. ℥ij. Mix the honey, water, and carbonate of lime in a silver vessel: boil the mixture, stirring, at the same time, for 2 or 3 minutes:

then add prepared animal charcoal ℥vj, two whites of eggs, mixed in a pint of water. Mix all, and boil to the consistence of syrup: remove from the fire; let the syrup rest for 15 minutes, and pass through a cloth. Ph. P.) Demulcent.

MELLITUS, Melitus.

MELÔË'MIA, (μελος, 'black,' and αίμα, 'blood,') A state of blood, characterized by its incoagulability, black colour, and septic properties.

MELOCHI'TIS, Malachi'tes, Lapis Armē'nus, Armeni'tes, Armē'nian Stone. A variety of blue carbonate of copper. It is found in Germany, the Tyrol, and especially in Armenia. It was formerly employed as a cardiac, and as proper for purging away melancholy.

MELODID'YMUS, (μελος, 'a limb,' and διδυμος, 'double,') A double member or limb.

MELÔE MAIALIS, see Cantharis.

MELÔE N'GER, M. Pennsylvan'icus; Class, Insecta; Ord. Coleoptera. A blistering fly, native of the United States. It feeds upon Prunell'a vulga'ris or Selfheal, and Ambrosia trif'ida or Stickweed. These flies resemble the Spanish flies in properties.

MELÔE PENNSYLVANICUS, M. niger.

MELÔE PROSCARABÆ'US, Cantarel'us, was anciently used as a diuretic and antihydrophobic. See Cantharis.

MELÔE VESICATORIUS, Cantharis.

MELOM'ELES, (melos, and μελεος, 'unprofitable,') A monster with supernumerary limbs.

MEL'ON, (μυλον, 'an apple,') Mel'um. A disorder of the eye, in which it protrudes out of the socket. See Exophthalmia.

MELON, Cucumis melo — m. d'Eau, Cucurbita citrullus — m. Plant, Cucumis melo — m. Water, Cucurbita citrullus.

MELONGENA, (from μυλον, 'an apple,') Solanum melongena.

MELOPLACUNTIUM, Meloplaeus.

MELOP'LACUS, Meloplacun'tium, (μηλοπλακος, from μυλον, 'a quince,' and πλακος, 'a sake,') A compound obtained by boiling wine, honey, quince, pepper, &c., together.

MELOPLAS'TIC, Meloplas'ticus, Genoplas'tic, (μυλον, 'the cheek,' and πλασσω, 'I form,') (F.) Meloplastic, an epithet for the operation for forming a new cheek when any part of it has been lost.

MEL'OS, (μελος,) Mem'brum. A member. A limb. An organized part, composed of other parts.

MELÔ'SIS, (μυλην, 'a probe,') Catheteris'mus, Ctenemate'sis. The act of probing a wound, ulcer, &c. Catheterism.

MELOTHRUM, (μυλωθρον,) Bryonia alba.

MELÔ'TIS, (μυλωτις, from μυλην, 'a probe,' and οσ, gen. ωος, 'the ear,') Melo'tris, Specil'um minus. A small probe, — a probe for the ear. See Apyromelo.

MELOTRIS, (μυλωτρις,) Melotis.

MELT, (Sax. milc.) Spleen. Also, the sperm of fish, — Milt.

MELTING, Fusion.

MEL'UM, (μυλον,) Mel'on, Ma'lum. An apple. Fruit in general. Also, a round, firm female breast. The cheek, mala. The apple eye or melon eye is an apple-formed projection of the eye from the orbit. See Melon.

MELUM ARMENIACUM, Prunus armeniaca — m. Cydonium, see Pyrus cydonia.

MELUSTUM, ('burnt honey,') Melasses.

MEMBER, Membrium.

MEMBRA PUDEENDA, Genital organs — m. Villia. Pudibilia.

MEMBRANA, (quia membra tegit, 'because it covers limbs,') Membrane — m. Abdominis, Peritonæum — m. Adiposa, Cellular membrano — m. Adventitia, Decidua reflexa — m. Agnina, Amnios, — m. Amphiblestroides, Retina — m. Arachnoidea,

Hyaloid membrane—*m.* Capsularis testis, Albuginea—*m.* Carnosa, Dartos—*m.* Cellulosa, Cellular membrane—*m.* Cellulosa, Decidua—*m.* Cerebri mollis, Pia mater—*m.* Cerebri propria, Pia mater—*m.* Cerebri tenuis, Pia mater—*m.* Chorior-capillaris, see Choroidæ tunica—*m.* Circumossalis, Pericæteum—*m.* Communis, Decidua—*m.* Cor circumplexa, Pericardium—*m.* Cordis, Pericardium—*m.* Costalis, Pleura—*m.* Costas succingens, Pleura—*m.* Crassa, Decidua reflexa—*m.* Cribrosa, Decidua—*m.* Cuticularis, Dura mater—*m.* Decidua serotina, see Decidua—*m.* Demuriana, see Aqueous humour, and Cornea—*m.* Dentata, Denticulatum ligamentum—*m.* Dermatodes, Dura mater—*m.* Descemetii, see Aqueous humour, and Cornea—*m.* Diaphratton, Mediastinum—*m.* Dura cerebrum ambiens, Dura mater—*m.* Elastica, Membrane, elastic—*m.* Epicrania, Galea aponeurotica capitis—*m.* Externa dentium, Tapetum alveoli—*m.* Farcinialis, Allantois—*m.* Fenestræ rotundæ, see Fenestra—*m.* Filamentosa, Decidua reflexa—*m.* Flocculenta, Decidua—*m.* Foetum involvens, Amnios—*m.* Fusca, see Sclerotic.

MEMBRANA GRANULOSA, *Granular membrane* or *lay'er.* (F.) *Membrane granuleuse, Couche celluleuse* (Coste). A layer of yellow, granular matter, which lines the inner layer of the Graafian follicle, and is regarded by Kölliker as epithelium.

MEMBRANA HUMORIS AQUEI, see Aqueous humour—*m.* Interspinalis, Interspinal ligament—*m.* Intestinalis, Allantois—*m.* Jacobi, Tunica Jacobi—*m.* Jacobi Arnold, see Iris—*m.* Limitans, see Iris—*m.* Limitans Pacini, see Iris—*m.* Lingue, see Tongue—*m.* Macilentior, Epiploon, gastrohepatie—*m.* Mucosa, Decidua, see Membrane—*m.* Nervorum, Neurilemma—*m.* Nictitans, Valvula semilunaris—*m.* Olfactoria, Pituitary membrane—*m.* Ossis, Periosteum—*m.* Ovi materna, Decidua—*m.* Ovi corticalis, Ovoculymma—*m.* Pellucida, Amnios—*m.* Pigmenti, see Iris—*m.* Pinguedinosæ, see Cellular membrane—*m.* Pinguis intestinorum, Mesentery—*m.* Pituitaria, Pituitary membrane, see Membrane—*m.* Pituitosa, see Membrane—*m.* Pleuritica, Pleura—*m.* Præexistens, Decidua—*m.* Propria, Basement membrane, and Folliculi Graafiani, see Cortex ovi—*m.* Pupillaris, Pupillary membrane.

MEMBRANA PUTAMINIS, (*putamen*, 'a shell or husk.') *Pellucula O'vi.* The membrane adherent to the inner surface of the eggshell.—*Putamen* seu *Testa O'vi.*

MEMBRANA RETIFORMIS CHORII, Decidua reflexa—*m.* Ruyschiana, Choroid (inner layer.)

MEMBRANA SACCIFORMIS. A separate synovial membrane for the inferior radiocubital articulation, which covers the upper surface of the triangular ligament, and the sort of incomplete ring which circumscribes the head of the ulna.

MEMBRANA SEMIPELLUCIDA, Caligo—*m.* Serosa, Membrane (serous)—*m.* Sinuosa, Decidua—*m.* Subcostalis, Pleura—*m.* Succingens, Diaphragm, Pleura.

MEMBRANA TESTÆ, *Shell mem'brane.* The membrane that lines the shell of the bird's egg:—a simple membrane.

MEMBRANA THORACEM INTERSEPIENS, Mediastinum—*m.* Tympani, see Tympanum—*m.* Tympani secundaria, see Fenestra—*m.* Urinaria, Allantois—*m.* Uteri interna evoluta, Decidua—*m.* Uvea, Uvea—*m.* Vasculosa cerebri, Pia mater—*m.* Vasorum Communis, Endangium—*m.* Verriularis, Retina.

MEMBRANA VERSICOLOR. A peculiar membrane—according to Mr. Fielding, of Hull—situate immediately behind the retina, and in connexion with it. It is separable into distinct layers from the choroid, and is supplied with bloodvessels.

MEMBRANA WACHENDORFIANA, Pupillary membrane.

MEMBRANACEOUS, Membranous.

MEMBRANE, *Membrana, Hy'men, Me'ninx, Epen'dyma, Pan'iculus, Pan'nicle.* A name given to different thin organs, representing a species of supple, more or less elastic, webs; varying in their structure and vital properties, and intended, in general, to absorb or secrete certain fluids; and to separate, envelop, and form other organs. Bichat has divided the membranes into simple and compound.

Simple membranes comprise three orders. 1. *Mucous mem'brane, Membra'na muco'sa* seu *pituitosa* seu *pituitaria, Phleg'my'men, Blen'nymen, Phlegmat'hymen, Myco'derma*, (F.) *Muqueuse, Membrane muqueuse, Membrane folliculeuse* (Ch.), is so called, on account of the mucous fluid by which it is constantly lubricated. Mucous membranes line the canals, cavities, and hollow organs, which communicate externally by different apertures on the skin. Bichat refers them to two great divisions—the *gastropulmonary*, and the *genitourinary*. The mucous membranes have a striking analogy with the cutaneous tissue, in organization, functions, and diseases. They are composed of corium, papillæ, and epidermis; and are furnished with a multitude of follicles, which secrete a viscid humour,—mucus. They receive a quantity of arterial vessels, veins, lymphatics, and nerves. 2. The *Serous mem'branes, Membra'næ sero'sæ, Orrhohym'enes, Orrhyn'enes, M. villieuses simples* (Ch.), are transparent, thin, and composed of one lamina. One surface adheres to other textures; the other is smooth, polished, and moistened by a serous fluid. They are arranged—in the form of sacs without apertures—as great, intermediate reservoirs for the exhalant and absorbent systems, in which the serous fluid, in passing from one system, tarries some time before it enters the other. The serous membranes resemble the areolar membrane in structure and diseases. They facilitate the motion on each other of the organs which they envelop. They may be divided into (a) *Serous membranes*, properly so called; as the arachnoid, pleura, peritoneum, and tunica vaginalis. (b) *Synovial membranes or capsules*; which belong to joints, tendons, aponeuroses, &c. These membranes—mucous and serous—are constituted of similar layers—epithelium, basement membrane, condensed areolar tissue; and a looser form of areolar tissue, termed *submucous* in one case,—*subserous* in the other. 3. *Fibrous mem'branes, Membranes albugineuses* (Ch.) These are almost all continuous, and terminate at the periosteum—their common centre. They have been divided into two sections. 1. *Enveloping aponeuro'ses, Aponeuro'ses of insertion, Fibrous capsules of the joints, and Fibrous sheaths of tendons.* 2. *The Perios'teum, Du'ra ma'ter, Sclero'tica, Tu'nica albugin'ea testis, Fibrous mem'brane of the spleen, &c.* The fibrous membranes are not free or moistened by any particular fluid. They adhere by both surfaces to the neighbouring parts; are firm, resisting, but slightly elastic, and of a white colour; sometimes pearly and glistening. Their vessels are numerous, in some, as in the dura mater and periosteum; in others, scarcely perceptible, as in the aponeuroses. The presence of nerves has never been proved, although several circumstances, regarding their sensibility, render their existence probable. The fibrous membranes serve, in general, to augment the solidity of the organs which they envelop; to retain the muscles in their respective positions; to favour the motion of the limbs, and that of the muscles and skin; to form canals and rings for the passage of different organs, &c.

Bichat admits three species of *Compound Membranes*. 1. The *Serofibrous*, formed of a serous

and fibrous lamina, intimately adherent to each other;—as the pericardium, dura mater, and tunica albuginea. 2. The *Sc'rom'e'ous*, formed of a serous and mucous lamina;—as the gall-bladder at its lower part. 3. The *F'i'bromucous*, constituted of the union of a fibrous and mucous membrane; as the mucous membrane of the nasal fossæ, gums, &c. Chaussier admits six kinds of membranes. 1. The *laminated*. See Cellular Tissue. 2. The *serous or simple villous*. 3. The *follicular or complicated villous*. 4. The *mucular or fleshy*. 5. The *albugineous*. 6. The *albuminous*.

MEMBRANE, ACCIDENTAL, *M. false*—*m. Accidentelle*, *M. false*—*m. Adnæe*, Conjunctiva—*m. Adventitious*, *M. false*—*m. Anhistous*, Decidua—*m. of Arnold*, see Iris.

MEMBRANE, BASEMENT, *Primary or lim'itary membrane*, *Membr'a pro'pria*. A delicate structureless lamella of membrane found beneath the epidermis or epithelium, on all the free surfaces of the body.

MEMBRANE CADUQUE, Decidua—*m. Caduque réfléchie*, Decidua reflexa—*m. de Demours*, see Aqueous Humour, and Cornea—*m. de Descent*, see Aqueous Humour, and Cornea.

MEMBRANE, ELASTIC, *Membr'a elastica*. Membrane formed of the elastic areolar tissue, and which occurs in the coats of vessels, especially of the arteries, in the trachea and bronchi, &c.

MEMBRANE, FALSE, *Accidental or Adventitious membrane*, *Pseudomembrane*, *Pseudymen*, *Pseudomennæ*, (*F. Fausse membrane*, *M. accidentelle*. Membranous productions, which form on all the free natural surfaces, and on every free accidental surface, are so called. They are, in general, produced by the exudation of a fibrinous matter, susceptible of organization, which takes place in consequence of inflammation of the various tissues. These accidental membranes occur on the skin after the application of a blister; on mucous surfaces, as in croup; on the parietes of inflamed veins and arteries, &c. The cicatrices of wounds are formed of them.

By some it has been proposed to give the name *pseudo-membrane* to fibrinous exudations, as in diphtheritis, which are devoid of organization, whilst *neo-membrane* may be applied to such fresh productions, as from serous membranes, as have become supplied with vessels and nerves.

MEMBRANE FAUSSE, Membrane, false—*m. Fibrous*, see Membrane—*m. Germinal*, see Molecule—*m. Granular*, *Membrana granulosa*—*m. Granulæuse*, *Membrana granulosa*—*m. Hapto-gen*, see Haptogenous—*m. Hyaloid*, see Hyaloid membrane—*m. Jacob's*, *Tunica Jacobi*—*m. Lim'itary*, Membrane, basement—*m. Mucous*, see Membrane—*m. Mucousæ*, see Membrane—*m. Nasale*, Pituitary membrane—*m. Nasmyth's*, see Enamel—*m. Olfactive*, Pituitary membrane—*m. Pituitaire*, Pituitary membrane—*m. Primary*, Membrane, basement—*m. Pseudo*, *M. false*—*m. Ruyschienne*, Choroid (inner layer), *Ruyschiana tunica*—*m. de Schneider*, Pituitary membrane—*m. Schneiderian*, Pituitary membrane—*m. Serous*, see Membrane—*m. Shell*, *Membrana testæ*—*m. Simple*, see Membrane—*m. Synovial*, see Membrane—*m. du Tympan*, see Tympanum—*m. Tympanal*, see Tympanum—*m. Utero-épichorale*, Decidua—*m. Virginal*, Hymen—*m. Vitellary*, *Zona pellucida*—*m. Vitelline*, *Zona pellucida*.

MEMBRANES ALBUGINEUSES, Membranes, fibrous, see Albuginea—*m. Folliculæuses*, Membranes, mucous—*m. Veloutées*, Villous membranes—*m. Villæuses simples*, Membranes, serous.

MEMBRANES OF THE FÆTUS, *Membr'ne fætum involv'ntes*, *Velamenta fœtalis*, *Integumenta fætalis*. The membranes. The membranes which immediately envelop the foetus in the cavity of the uterus, and the rupture of which gives rise to

the discharge of the liquor amnii. These membranes are the *decidua*, *chorion*, and *amnion*.

MEMBRANES, THE, Membranes of the Fœtus.

MEMBRANEUX, Membranous.

MEMBRANIFORM or MEMBRANIFORM, *Membraniformis*, (*membrana* and *forma*;)—resembling a membrane. A name given to thin and flat parts, which resemble membranes.

MEMBRANIFORM MUSCLES. Very broad and thin muscles, as the *platysma myoides*, obliquus abdominis, &c.

MEMBRANOSUS, (from *membrana*;) Fascia lata muscle, Membranous.

MEMBRANOUS, *Membranosus*, *Membranoceus*, *Hymenodes*, (*F. Membraneux*. Having the nature of membrane. Formed of membrane.

MEMBRANULA, (dim. of *membrana*;) Hymenium, Membranule—*m. Coronæ ciliaris*, see Ciliary—*m. Lunata vaginæ*, Hymen—*m. Nervorum*, Neurilemma—*m. Semilunaris conjunctivæ*, Valvula semilunaris.

MEMBRANULE, *Membran'ula*. A small membrane.

MEMBRE, Membrum—*m. Viril*, Penis.

MEMBROSUS, *Membr'sior*, *Mentulatus*, *Mutonia'tus*, *Nasatus*, *valdè Mentulatus*, *Fascinosus*, *Pso'lon*, (*F. Membra*, (from *membrum*, 'the male organ'.) One whose penis is very large.

MEMBRU, Membrus.

MEMBRUM, *Ar'tus*, *M'los*, *Co'l'on*, a limb, a member, (*F. Membre*. The limbs or extremities or members of animals are certain parts exterior to the body, which are more or less apparent, long, and movable. It is by means of their limbs, that animals transport themselves from one place to another; and that they defend themselves, or attack others. The limbs are arranged on each side of the trunk, with which they are articulated. In man, they are four in number:—two upper or thoracic, and two lower, pelvic or abdominal. Also, the male organ.

MEMBRUM GENITALE VIRORUM, Penis—*m. Mulieris*, Clitoris—*m. Seminale*, Penis—*m. Virile*, Penis.

MÉMOIRE, Memory—*m. Perte de*, Amnesia.

MEMORIA, Memory—*m. Deleta*, Amnesia.

MEM'ORY, *Memo'ria*, *Mne'mè*, *Recordatio*, *Recordatus*, (*F. Mémoire*. The cerebral faculty, by virtue of which past impressions are recalled to the mind.

MEMPHITES LA'PIS. A sort of stone, found anciently in the environs of Memphis; which was esteemed narcotic.

MENAGOGUES, (*menes*, and *αγω*, 'to drive.')

Emmenagogues.

MENDESION, *Ægyptiacum*.

MENE (*μην*), Moon.

MENECRATIS ACCIPITER, Accipiter.

MENELCO'SIS, *Hæmorrhæmia*, (*menes*, and *ἐλκος*, 'ulceration.')

Vicarious menstruation from an ulcer.

MENES (*μηνς*), Menses.

MÉNESPAUSIE, (*menes*, and *παυσι*, 'to cause to cease.')

See Menses.

MENFRIGE, see Pistacia lentiscus.

MENIDRO'SIS, (*menes*, and *ῥοις*, 'sweating.')

A sweating of menstrual blood.

MENINGARTHROACÉ, *Inflammati'o men-*

brancarum articulatio'nis, *Arthromeningi'tis*, (*με-*

νιγγο, *αρθρον*, 'a joint,' and *κακος*, 'disease.')

Inflammation of the membranes of a joint.

MÉNINGE, (from *meninæ*;) Dura mater.

MENINGÉ, Meningeal.

MENINGEAL, or MENIN'GEAL, *Menin'geus*, (*F. Ménin'gé*. Relating to the meninges, or merely to the dura mater.

MENINGEAL APOPLEXY, see Apoplexy.

MENINGEAL ARTERY, MID'DLE, *Arter'ia menin-*

gea me'dia seu Sphenospino'sa seu Du'ræ Ma'

tris me'dia max'ima seu Sphenospina'lis, Spi'nal

Ar'tery, (F.) *Artère méningée moyenne* ou *Sphéno-épineuse*, ou *Épineuse*, is the largest of the branches given off by the internal maxillary artery. It enters the cranium by the foramen spinale, and distributes its branches chiefly to the dura mater.

MENINGEAL VESSELS, *Vas'a meningeae*. The vessels of the membranes of the brain.

MENINGES, (*μνινγες*, pl. of *meninx*.) *Éilam-ides*, *Ματ'ρες* seu *Ινβολ'ερα* *κερ'εβρι*, *Ομεν'τα* seu *Τελομεν'τα* *κερ'εβρι'α*. The three membranes which envelop the brain—*Dura mater*, *Arachnoid*, and *Pia mater*.

MENINGETTE, *Pia mater*.

MENINGIA TUBERCULOSA, see *Hydrocephalus internus*.

MENINGITIS, *Meningitis*, and, especially, *Inflammation of the meningium or arachnoid membrane*.

MENINGINA, (F.) *Méningine*. A name given by Chausser to the pia mater, united to the cerebral layer of the arachnoid.

MENINGINA, INNER LAMINA OF THE, *Pia mater*.

MENINGINE, *Meningina*.

MÉNINGITE, *CÉRÉBRO-SPINALE*, *Meningitis*, *cerebro-spinal*.

MENINGINITIS, (*meningina*, and *itis*.) *Inflammation of the meningina*.

MENINGION, (*μνινγιον*, dim. of *μνινξ*, 'a membrane.') *Arachnoid membrane*.

MENINGIS CUSTOS, *Meningophylax*.

MENINGITE, *Meningitis*—*m. Granulose*, *Hydrocephalus internus*—*m. Rachidienne*, *Meningitis*, *spinal*—*m. Tuberculose*, *Hydrocephalus internus*.

MENINGITIS, *Meningit'is*, *Encephalit'is* *peripher'ica* seu *membrano'sa*, (F.) *Inflammation de la membrane séreuse céphalo-rachidienne*, *Méningite*, (*meninges*, and *itis*.) *Inflammation of a membrane and especially of the meninges or membranes of the brain*. See *Phrenitis*.

MENINGITIS ARTHRITICA, *Cephalagra*.

MENINGITIS CEREBROSPINAL, *Myelit'is petechial'is*, *Cerebral or cerebrospinal typhus*, (F.) *Méningite cérébro-spinale*, *Fievre cérébro-spinale*, *Typhus cérébro-spinal*. An alarmingly fatal form of epidemic fever, which prevailed in different countries of Europe, and, also, in certain portions of the United States, near the middle of the 19th century. It is considered, by Boudin, to constitute the cerebro-spinal form of the nosological group of typhus.

MENINGITIS GRANULAR, *Hydrocephalus internus*—*m. Hydrocephalica*, *Hydrocephalus internus*—*m. Infantum*, *Hydrocephalus internus*—*m. Membranosa*, *M. spinal*.

MENINGITIS MESENCEPHALICA. *Inflammation of the meninges of the medulla oblongata and pons varolii more especially*.

MENINGITIS PERIPHERICA, *M. spinal*—*m. Phantasmaphora*, *Delirium tremens*—*m. Potatorum*, *Delirium tremens*—*m. Rachidian*, *M. spinal*—*m. Scrophulous*, *Hydrocephalus internus*.

MENINGITIS SPINAL, *M. Rachidian*, *M. Spinal'is* seu *peripher'ica* seu *membrano'sa*, *Perimyelit'is*, *My'elomeningit'is*, (F.) *Méningite rachidienne*. *Inflammation of the meninges of the spinal marrow*.

MENINGITIS, TUBERCULAR, see *Hydrocephalus internus*—*m. Tuberculosa*, *Hydrocephalus internus*.

MENINGIUM, (*μνινγιον*, dim. of *μνινξ*, 'a membrane.') A fine, delicate membrane. The tunica arachnoidea of the brain.

MENINGO, (*μνινξ*, gen. *μνινγος*, 'membrane,') in composition, *gen.*

MENINGOCEPHALITIS, *Meningoencephalit'is*, *Ceph'alomeningit'is*, *Enceph'alomeningit'is*, (from

meningo, κεφαλη, 'head,' and *itis*.) *Inflammation of the membranes and brain*.

MENINGOGASTRIC FEVER, *Fever, gastric*.

MENINGOGASTRITIS, *Fever, gastric*.

MENINGOMALACIA, (*meningo*, and *μαλακία*, 'softening,') *Softening of membranes in general*.

MENINGOPHYLAX, *Cust'os menin'gis*, *Dépresso'rium*, (F.) *Dépressoire*, (from *meningo*, and *φλάξ*, 'a preserver,') *An instrument for depressing the dura mater, and guarding it from injury, whilst the bone is cut or rasped*.

MENINGORRHŒA, (*meningo*, and *ῥεω*, 'flow,') *Effusion of blood upon or between the membranes of the brain*.

MENINGOSIS. The union of bones by means of membrane; e. g. the articulation of the bones of the cranium in the fœtus. *Meningosis* is a variety of *Syndesmosis*.

MENINGOSYMPHYSIS, *Meningosyzyg'ia*, (*meningo*, and *συνψυσις*, 'growing together,') *Adhesion of or by membrane*. *Adhesion of the membranes of the brain to each other, or to the brain*.

MENINGOSYZYGIA, (*meningo*, and *συνζυγία*, 'a conjunction,') *Meningosymphysis*.

MENINGURIA; (*meningo*, and *ουρον*, 'urine,') *The passage of urine containing membranous shreds*.

MENINX, gen. *Menin'gis*, (*μνινξ*, gen. *μνινγος*.) *Membrane*, (especially one that invests the brain) *m. Choroides*, *Pia mater*—*m. Crassa*, *Dura mater*—*m. Dura*, *Dura mater*—*m. Exterior*, *Dura mater*—*m. Interior*, *Pia mater*—*m. Media*, *Arachnoid membrane*—*m. Pacheia*, *Dura mater*—*m. Sclera*, *Dura mater*—*m. Serosa*, *Arachnoid membrane*—*m. Vasculosa*, *Pia mater*.

MENISCHEISIS, (*menes*, and *ισχειν*, 'to restrain,') see *Amenorrhœa*.

MENISCI, (*μηνισκος*, dim. of *μηνη*, 'moon,') *Cartilages, interarticular*.

MENISPERMUM ANGULATUM, (*μηνη*, 'the moon,' and *σπέρμα*, 'seed,' in allusion to the crescentic shape of the seed.) *M. Canadense*.

MENISPERMUM CANADENSE, *M. angulatum*, *Ord. Menispermaceæ*. A climbing plant, growing in various parts of the United States. Said to be used in Virginia as a substitute for sarsaparilla in scrofula, and to be called, in the West, *Sarsaparilla*. It is an excitant tonic.

MENISPERMUM COCCULUS, *M. glaucum*, *Anamir'ta Cocculus* seu *paniculata*, *Cocculus suberosus*. The systematic name of the plant, the dried fruit of which are well known by the name of *Cocculus Indicus* seu *suberosus*, *Indian Berries*, *Bac'cæ Piscator'icæ*, *Indian Cock'les*, *Cocculus Indicus*, *Cocculi officinarum*, *Cocculi orientales*, *Gra'na Orient'is*, (F.) *Coque du Levant*, *C. levant*. These berries are remarkable for their inebriating and destructive quality to fish. The deleterious principle appears to reside in the kernel, and in the active principle called *Picrotox'ine*. *Cocculus Indicus* has been used in decoction to kill vermin. It has, at times, been added to beer, by fraudulent tradesmen, to render it more inebriating.

MENISPERMUM GLAUCUM, *M. cocculus*—*m. Palmatum*, *Calumba*.

MENISPERMUM TUBERCULATUM, *M. verrucosum*, *Cocculus crispus*. A Chinese plant, an extract from the root of which is tonic to the stomach and bowels, and possesses some astringency. Dose, 5 to 10 grains.

The same may be said of *Menispermum Cordifolium*—the *Goluncha*, of Bengal.

MENISPERMUM VERRUCOSUM, *M. tuberculatum*. **MENOCRYPTIA**, (*menes*, and *κρυπτος*, 'concealed,') *Amenorrhœa*.

MENOLIPSIS, (*menes*, and *λαψις*, 'want,') see *Menses*.

MENOPAUSIS, (*menes*, and *παυειν*, 'to cause to cease,') see *Menses*.

MENOPHANIA, (*menses*, and φαῖνω, 'to appear,') see *Menses*.

MENOPLANIA, (*menses*, and πλανη, 'a wandering,') Menstruation, vicarious.

MENORRHAG'IA, (*menses*, and ἰσχυρῶς, 'to flow fiercely,') Flow of the menses. Frequently, the word is used synonymously with uterine hemorrhage or *metrorrhagia*, or for immoderate flow of the menses—*profuse menstruation*, *Paramenia superflua seu profusa*, *Menorrhag'ia rubra*, *Catameniorum fluxus immodicus*, *Menstrua immodica seu superflua*, *Menorrha'a*.

MENORRHAGIA, Menstruation. *Metrorrhagia*—*m. Alba*, *Leucorrhœa*—*m. Difficilis*, *Dysmenorrhœa*—*m. Erronea*, Menstruation, erroneous—*m. Stillititia*, *Dysmenorrhœa*.

MENORRHŒA, (*menses*, and ῥέω, 'I flow,') The ordinary flow of the menses. Also, too long continuing, or too often returning, menstruation. *Menorrhagia*.

MENOSCHESIS, (*menses*, and ὀχεω, 'to hold,') see *Amenorrhœa*.

MENOSTASIA, (*menses*, and στας, 'standing,') *Amenorrhœa*.

MENOS'TASIS. This word, according to some, signifies the retention of the menses and their accumulation in the uterus. According to others, it means the acute pain which, in some females, precedes each appearance of the menses: a pain which has been presumed to proceed from the stasis of blood in the capillary vessels of the uterus.

MENOXEN'IA, *Menoxeno'is*, (*menses*, and ξενος, 'foreign,') Irregular menstruation.

MENOXENOSIS, *Menoxenia*.

MENS, *Anima*, *Intellect*.

MEN'SA, *Trapa'za*, 'a table.' The upper superficial part or corona of the jaw teeth.

MEN'SES, pl. of *Mensis*, from *mens*, 'a month,') *M. Muliebres*, *Mulieb'ris*, *Men'es*, *Fluxus menstrualis* seu *lunaris*, *Profluvium mulieb'rè seu genit'alè mul'ebre*, *Consectu'do menstrualis*, *Catamenia*, *Mens'trua*, *Eumenia*, *Tributum lunat'rè seu menstrualium*, *Purgatio'es*, *P. menstrualis*, *Gynaecol'ia*, *Aphedria*, *Cut'sus menstrualis*, *Hor'a'a*, *Luna'ria*, *Courses*, *Men'strial flux*, *Monthly courses* or *periods*, *Flow'ers*, *Turns*, *Terms*, *the Reds*, *Troubles*, (F.) *Mois*, *Règles*, *Lunes*, *Flucurs*, *Affaires*, *Époques*, *Ordinaires*, *Purgations*, *Fleurs*, *Menstrues*. The sanguinous evacuation from the uterus; *Sanguis menstrualis*, *Men'strial blood*, the monthly occurrence of which constitutes menstruation or monthly illness. The first appearance of the menses—*menophania*—is usually preceded by the discharge of a fluid whitish matter from the vagina: by nervous excitement, and by vague pains and heaviness in the loins and thighs; numbness of the limbs, tumefaction and hardness of the breasts, &c. More or less indisposition and irritability also precede each successive recurrence of the *menstrual flux*. In temperate climates, each period, *men'strial period*, *peri'odus lunaris*, ordinarily continues from three to six days; and the quantity lost varies from four to eight ounces. The menses continue to flow from the period of puberty till the age of 45 or 50. At the term of its natural cessation, *Menop'is*, *Ménopause* (Gardanne), *Menopaus'is*, the flux becomes irregular; and this irregularity is occasionally accompanied with symptoms of dropsy, glandular tumours, &c., constituting the *Paramenia cessatio'nis* of Good, and what is called the *critical time* or *turn* or *change of life*; yet it does not appear that the mortality is increased by it. With the immediate causes of menstruation we are unacquainted. We express only our ignorance, when we assert it to depend upon periodicity; the discharge comes from the vessels of the uterus and vagina, and differs from ordinary blood by its peculiar odour, and by its not coagu-

lating. It is evidently connected with the condition of the ovaries, and—it is generally supposed—with the periodical discharge of ova from them. It is arrested, as a general principle, during pregnancy and lactation. In warm climates, women usually begin to menstruate early, and cease sooner than in the temperate regions. The quantity lost is also greater. In the colder regions, the reverse of this holds as a general rule.

Dried menstrual blood, especially of a woman in her first menstruation—*Sanguis vir'ginis primum menstrual'is*—was formerly given internally as an anti-epileptic.

MENSES ALBI, *Leucorrhœa*—*m. Anomali*, *Paramenia*—*m. Devii*, Menstruation, vicarious—*m. Dolorifici*, *Dysmenorrhœa*—*m. Retention* of the, see *Amenorrhœa*—*m. Suppression* of the, see *Amenorrhœa*.

MENSIUM RETENTIO, ('retention of the menses,') see *Amenorrhœa*.

MENSTRU'A, *Menses*—*m. Alba*, *Leucorrhœa*—*m. Difficilia*, *Dysmenorrhœa*—*m. Dolorosa*, *Dysmenorrhœa*—*m. Immodica*, *Menorrhagia*—*m. Superflua*, *Menorrhagia*.

MENSTRUAL, *Catamenial*—*m. Flux*, *Menses*—*m. Period*, see *Menses*.

MENSTRUANT, *Men'struant*, *Menstruat'us*, (from *menstrua*.) One subject to the catamenia.

MENSTRUATIO, Menstruation—*m. Anomala*, *Paramenia*—*m. Difficilis*, *Dysmenorrhœa*—*m. Dolorifica*, *Dysmenorrhœa*—*m. Impedita*, see *Amenorrhœa*—*m. Per insolitas Vias*, Menstruation, vicarious.

MENSTRUATIO PRÆCOX, *Premature menstruation*. Menstruation occurring before the usual age or time.

MENSTRUATIO RECIDIVA. Menstruation, when protracted beyond the usual age.

MENSTRUATIO RETENTA, *Amenorrhœa*, *Paramenia*—*m. Suppressa*, *Amenorrhœa*, *Paramenia*.

MENSTRUATION, *Menstruat'io*, *Chronogyn'ia*, *Menorrhag'ia*, *Fluxus menstrual'is*. The flow of the menses.

MENSTRUATION DIFFICILE, *Dysmenorrhœa*—*m. Difficult*, *Dysmenorrhœa*—*m. Laborious*, *Dysmenorrhœa*—*m. Premature*, *Menstruatio præcox*—*m. Profuse*, *Menorrhagia*.

MENSTRUATION, VICA'RIOUS, is that which occurs from other parts than the uterus. It is called *Paramenia error'is*, *Menorrhag'ia error'is*, *Men'strum per alie'na loc'a excretio*, *Menses devii*, *Aberratio mensium seu menstruorum*, *Menstruat'io per insol'itas vi'as*, *Hæmatopla'nia seu Hæmatopla'nia menstrual'is*, *Menopla'nia*, (F.) *Déviation des Règles*, *Règles déviées*. At times, the discharged fluid has all the characters of the menstrual; at others, it appears to be mere blood.

MENSTRUATUS, *Menstruat*.

MENSTRUE, *Menstruum*.

MENSTRUUEL, *Catamenial*.

MENSTRUËS, *Menses*.

MENSTRUOUS, *Catamenial*. Also, one affected with the menses or catamenia.

MENSTRUUM, (F.) *Menstrue*, (from *mensis*, 'a month,')—the menstruum being, of old, usually continued in action for a month.) The monthly discharge of the female. The plural, *menstrua*, is now, however, always used for the menses. The name was formerly given to every substance which possesses the property of dissolving others slowly and with the aid of heat. At present, it is used synonymously with *solvent*. Thus—water, alcohol, ether, acids, oils, &c., are *menstrua*.

MENSTRUUM AURI, *Nitromuriatic acid*—*m. Ventriculi*, *Gastric juice*.

MENSURA MEDICA. A measure of 48 ounces.

MENSURATION, *Meta'ria*, (from *mensura*,

'measure.') *Meas'urement*, *Med'ical meas'uring*. One of the means used for exploring the state of the thoracic and other cavities. It consists in a comparative measurement of each side of the chest, by means of a ribband extended from the median line of the sternum to the spine. When effusion exists on one side, the measurement is usually greater than on the other. When, on the other hand, a portion of a lung has ceased, for some time, to exert its respiratory functions, the corresponding side of the chest becomes smaller, in consequence of the contraction of the pleura, retraction of the pulmonary tissue, and greater approximation of the ribs. Instruments have been devised for this purpose. See *Stethometer*.

MENTA, *Mentha*, *Penis*.

MENTAGRA, (*mentum*, 'the chin,' and *aypa*, 'a seizure,') *Sycosis*—*m. Infantum*, *Porrigo lupinosa*.

MEN'TAGRAPHYTE, *Men'tagrophyte*, *Microsporion mentagrophytes*, from *mentagra*, and *αἴσρον*, 'a plant.' A name proposed by M. Gruby, of Vienna, for a cryptogamous plant, which he considers to produce *mentagra*.

MENTAL, *Mentalis*, (*mens*, gen. *mentis*,) belonging or relating to the mind, as *mental phenomenon*, *mental diseases* (*Mor'bi mentales*), &c.

In Anatomy, mental, *Mentalis*, (F.) *Mentonier*, signifies that which relates to the chin; (from *mentum*.)

MENTAL AR'TERY, (F.) *Artère mentonnière*. This is given off by the inferior dental, and issues at the mental foramen to be distributed on the lower lip.

MENTAL FORA'MEN, *Anterior maxillary F.*, (F.) *Trou mentonnier*, is the outer orifice of the inferior dental canal. It is situated on the outer surface of the lower jaw-bone, opposite the second incisor, or the canine tooth, and gives passage to the vessels and nerves.

MENTAL FOSSA, (see *Fossa*, *mental*).

MENTAL NERVE, (F.) *Nerf mentonnier*, is furnished by the inferior dental nerve. It issues by the mental foramen, and is distributed, in numerous filaments, to the muscles of the lower lip.

MENTAL REGION, *Re'gio mentalis*. The region of the chin.

MEN'THA seu MEN'TA AQUAT'ICA, (*μνθη*, *μνθος*, 'mint,') *M. palustris* seu *hirsuta* seu *rotundifolia palustris*, *Menthastrium*, *Sisymbrium menthastrium*, *Balsamum palustris*, *Watermint*, (F.) *Menthe aquatique*, *M. rouge*. *Nat. Ord. Labiateæ*. This is less agreeable than the *Mentha viridis*, and more bitter and pungent. It is used like *spearmint*.

MENTHA BALSAMEA, *M. piperita*.

MENTHA CAPEN'SIS, which grows at the Cape of Good Hope, possesses the medical properties of the *mints*.

MENTHA CERVI'NA, *Hart's pen'nyroyal*, *Hys'sop-leaved mint*, *Pule'gium cervinum*, (F.) *Menthe cervine*. Possesses the properties of pennyroyal, but is very unpleasant.

MENTHA CITRATA, *Melissa*.

MENTHA CRIS'PA, *M. hercynica*, *Colymbifera minor*, *Curled Mint*, (F.) *Menthe frisée* ou *crêpe*, *Baume d'eau à feuilles ridées*. Possesses the properties of peppermint.

MENTHA GENTILIS, *M. viridis*—*m. Hercynica*, *M. crispa*—*m. Hirsuta*, *M. Aquatica*—*m. Lævigata*, *M. viridis*—*m. Officinalis*, *M. piperita*—*m. Palustris*, *M. Aquatica*.

MENTHA PIPERITA, *M. Piperitis* seu *officinalis* seu *balsamea*, *Pep'permint*, (F.) *Menthe poivrée*. The odour of this variety is strong and agreeable. Taste, pungent, aromatic, and producing a sensation of coldness in the mouth. Virtues depend upon essential oil and camphor. It is stom'achic and carminative. It is chiefly

used in the form of essential oil—the *oleum menthae piperitæ*.

Ess'ence of *Pep'permint* consists of *ol. menthae pip.* 3ij, *sp. vin. rectif.*, coloured with *spinach leaves*, Oij. See *Tinctura olei menthae piperitæ*.

MEN'THA PUL'E'GIUM, *Pule'gium*, *P. regale* seu *latifolium*, *Gle'chon*, *Pen'nyroyal*, *Pud'dinggrass*, (Prov.) *Hil'wort*, *Lur'key-dish*, *Organ*, (F.) *Pouliot*. The oil—*Oil'um pule'gii*, *Oil of pen'nyroyal*,—possesses the virtues of the plant. It resembles the mints in general in properties. Amongst the vulgar, it is esteemed an emmenagogue. *A'qua PUL'E'GI*, *A'qua Men'thae pule'gii*, *Pen'nyroyal Wa'ter*, was official in the Edinburgh and Dublin Pharmacopœias.

MEN'THA ROMANA, *Balsamita suaveolens*, *Tanacetum balsamita*—*m. Rotundifolia palustris*, *M. aquatica*—*m. Saracenicæ*, *Balsamita suaveolens*, *Tanacetum balsamita*—*m. Sativa*, *M. viridis*—*m. Spicata*, *M. viridis*.

MEN'THA VIR'IDIS, *M. Sativa* seu *vulgaris* seu *gentilis* seu *spicata* seu *lævigata*, *Spear'mint*, (F.) *Menthe sauvage* ou *verte* ou *des jardins*, *Baume des jardins*. Odour strong, aromatic; taste warm, austere, bitterish. Virtues the same as those of the peppermint. The *Oleum menthae viridis* is obtained from it.

MEN'THA VIRGARIS, *M. viridis*.

MEN'THASTRUM, *Mentha aquatica*.

MEN'THA AQUATIQUE, *Mentha aquatica*—*m. Cervine*, *Mentha cervina*—*m. Coq*, *Tanacetum balsamita*—*m. Crêpe*, *Mentha crispa*—*m. Frisée*, *Mentha crispa*—*m. des Jardins*, *Mentha viridis*—*m. Poivrée*, *Mentha piperita*—*m. Romaine*, *Balsamita suaveolens*—*m. Rouge*, *Mentha aquatica*—*m. Sauvage*, *Mentha viridis*—*m. Verte*, *Mentha viridis*.

MENTIGO, (from *mentum*, 'chin,') *Porrigo larvalis*, *Sycosis*.

MENTISME, (from *mens*, gen. *mentis*, 'mind,') Any irregular movement of the mind, whether from emotion or a vivid imagination.

MEN'TOBICORNEUS, (*mentum*, 'chin,' and *bicornis*, 'two horned,') *Geniohyoides*.

MEN'TOLA'BIAL, *Mentolabialis*, (from *mentum*, 'the chin,' and *labium*, 'a lip,') Under this name *Chaussier* has united the muscles *Levator labii inferioris*, and *Quadratus*. See *Depressor labii inferioris*.

MEN'TOLABIAL FUR'ROW: a transverse depression situate between the lower lip and the chin, which is remarkable for the perpendicular direction of the hairs growing upon it.

MENTON, *Mentum*.

MENTON'NE (I.), (F.) *Menton*—now in France. A small village on the Mediterranean, in a sheltered bay bounded by Capes St. Martin and Murtola, which is much frequented by valetudinarians, especially those laboring under pulmonary affections.

MENTONNIER, *Mental*—*m. Labial*, *Depressor labii inferioris*.

MENTONNIERE, see *Funda*.

MENTULA, (like *menta*,) *Penis*—*m. Facititia*, *Dildo*—*m. Muliebris*, *Cercosis*, *Clitoris*—*m. Parva*, *Pippina*.

MEN'TULAGRA, (*mentula*, 'the penis,' and *aypa*, 'a seizure,') A convulsive erection of the *mentula* or penis; such as is said to be sometimes observed in eunuchs. See *Pudendagra*.

MEN'TULATUS, *Membrosus*.

MEN'TUM, *Geni'on*, *Geni'um*, *Anthe'reon*, *Chin*. (F.) *Menton*. The inferior and middle part of the face, situate below the lower lip. The chin is formed of skin, areolar tissue, muscles, vessels, nerves, and the os maxillare inferius.

MENYAN'THES TRIFOLIA'TA, *Menyan'thes*, *Minyan'thes*, (*men*, and *ανθος*, 'a flower,' because it keeps in flower about a month [?].) *Trifolium paludosum* seu *aquat'icum* seu *palustris* seu *fibri'*.

num, Wa'ter tref'oil, Buck'bean, Bog'bean. Nat. Ord. Gentianaceæ. (F.) Trêfle d'eau ou des Marais ou de Castor. The taste is intensely bitter, and is extracted by water. It is tonic, anthelminthic, diuretic, cathartic, and, in large doses, emetic. In some countries, it is used as a substitute for hops in making beer.

MENYANTHES VER'NA, *Amer'ican buck'bean, Marsh tref'oil, Wa'ter sham'rock, Bit'ter root*, has similar properties.

MEOSIS, *Meiosis*.

MEPHIT'IC, *Mephit'icus, (F.) Méphitique; (from mephitis.)* Any thing possessed of an unwholesome property; chiefly applied to exhalations.

MEPHITIC ACID, Carbonic acid—*m. Air, Carbonic acid.*

MEPHITIS, see Mephitism—*m. Hepatica, Hydrogen, sulphuretted—m. Inflamabilis, Hydrogen—m. Urinosa, Ammonia.*

MEPHITISM, *Mephitism'us, (F.) Méphitisme.* Any pernicious exhalation. *Mephitis* and *Mophe'ta, (F.) Mofette* ou *Monfette*, are old names for nitrogen; and, in general, for all exhalations and gases that are unfit for respiration.

MERACUS, *Merus*.

MERAMAUROSIS, (*mero*, and *amaurosis*.) *Hypamaurosis*.

MERANÆSTHÆSIS, *Meranæsthæ'sia, (mero*, *av*, 'privative,' and *αισθησις*, 'sensation.') Insensibility of a part of the body.

MERATROPH'IA, (*mero*, and *ατροφία*, 'want of nourishment.') Atrophy of some part of the body.

MERCURE, (after the god *Mercury*.) *Hydrargyrum—m. Bichlorure de, Hydrargyri oxyurias—m. Biiodure de, Hydrargyri iodium rubrum—m. Bromures de, see Bromine—m. Coralli, Hydrargyri nitrico-oxydum—m. avec la Craie, Hydrargyrum cum cretâ—m. Cru, Hydrargyrum—m. Cyanure de, Hydrargyri cyanuretum—m. Deutochlorure de, Hydrargyri iodium rubrum—m. Deutochlorure de, Hydrargyri oxyurias—m. Doux, Hydrargyri suburias—m. Gommeux de Plench, Syrupus de mercurio mediante gummi—m. Muriate oxygéné de, Hydrargyri oxyurias—m. Oxide de, cendré, Hydrargyri oxydum cinereum—m. Oxide gris de, Hydrargyri oxidum cinereum—m. Oxide nitrique de, Hydrargyri nitrico-oxydum—m. Oxide noir de, Hydrargyri oxydum cinereum—m. Oxide rouge de, Hydrargyri oxydum rubrum—m. Périodure de, Hydrargyri iodium rubrum—m. Protiodure de, Hydrargyri iodium—m. Protochlorure de, Hydrargyri suburias—m. Protoxide de, Hydrargyri oxydum cinereum—m. Sous-muriate de, précipité, Hydrargyrum præcipitatum—m. Sous-sulfate de, Hydrargyrum vitriolatus—m. Sulfure de, noir, Hydrargyri sulphuretum nigrum—m. Sulfure de, rouge, Hydrargyri sulphuretum rubrum.*

MERCUR'IAL, *Mercuria'lis, (F.) Mercuriel.* That which contains or relates to mercury. A preparation of mercury—*Hydrargyri'cum, Hydrargyri'um.*

MERCURIAL DISEASE, *Hydrargyriasis—m. Erethism, see Erethism—m. Rash, Eczema mercuriale.*

MERCURIALE ANNUELLE, *Mercurialis—m. des Bois, Mercurialis perennis—m. de Chien, Mercurialis perennis—m. Vivace, Mercurialis perennis.*

MERCURIA'LIS, (after *Mercury*.) *Mercuria'lis an'na, French mer'cury, (F.) Mercuriale annuelle, Foïrolle, Ramberge, Vignoble; Class, Dicoecia; Order, Eneandria.* It has been esteemed cathartic, hypnotic and cosmetic, and has been chiefly used by way of *laxament*. See *Chenopodium Bonus Henricus*.

MERCURIALIS HU'MOR, *Hu'mor melanchol'icus.* A supposititious humour with the older physicians,

out of which they presumed that *morbi melancholici seu mercuriales* arose.

MERCURIALIS PEREN'NIS, *Cynocram'bè, Cani'na bras'sica, M. mont'ana, M. Sylvest'ris, Dog's mer'cury, (F.) Mercuriale vivace, M. de chien, M. des bois.* This plant is possessed of acronarcotic properties.

MERCURIALISMUS, *Hydrargyriasis.*

MERCURIALIST, *Mercurialis'ta.* One inordinately addicted to prescribing mercury.

MERCURIALIZA'TION. The state of being affected by mercury. One under the influence of mercury is said to be *mercurialized*.

MERCURII CALX VITRIOLATA, *Hydrargyrum vitriolatus—m. Deutoioduretum, Hydrargyri iodium rubrum—m. Proto-ioduretum, Hydrargyri iodium—m. Sperma, Hydrargyrum acetatus—m. Terra foliata, Hydrargyrum acetatus.*

MERCURIUS, (after the god,) *Hydrargyrum—m. Acetatus, Hydrargyrum acetatus—m. Alkalisatus, Hydrargyrum cum cretâ—m. Calcinatus, Hydrargyri oxydum rubrum—m. Causticus flavus, Hydrargyrum vitriolatus—m. Cinereus, Hydrargyri oxydum cinereum—m. Cinnabarinus, Hydrargyri sulphuretum rubrum—m. Corallinus, Hydrargyri nitrico-oxydum—m. Corrosivus, Hydrargyri oxyurias—m. Corrosivus ruber, Hydrargyri nitrico-oxydum—m. Corrosivus sublimatus, Hydrargyri oxyurias—m. Cosmeticus, Hydrargyrum præcipitatum—m. Duleis, Hydrargyri suburias—m. Emeticus flavus, Hydrargyrum vitriolatus—m. Gummosus Plenchii, Syrupus de mercurio mediante gummi—m. Luteus, Hydrargyrum vitriolatus—m. Mortis, Algaroth—m. Niger Moseati, Hydrargyri oxydum cinereum—m. Nitrosus calide paratus, see Hydrargyri nitras—m. Nitrosus frigore paratus, Hydrargyri nitras—m. Oxydulatus hydriodicus, Hydrargyri iodium—m. Præcipitatus corrosivus, Hydrargyri nitrico-oxydum—m. Præcipitatus niger, Hydrargyri oxydum cinereum—m. Saccharatus, Hydrargyrum saccharatum—m. Solubilis of Hahnemann, see Hydrargyri oxydum cinereum—m. Sublimatus duleis, Hydrargyri suburias—m. cum Sulphure, Hydrargyri sulphuretum nigrum—m. Terrestris, Polygonum Hydropiper—m. Vitæ, Algaroth—m. Zoticus Hartmanni, Hydrargyri suburias.*

MERCURY, (after the god,) *Hydrargyrum—m. Acetate of, Hydrargyrum acetatus—m. Acid nitrate of, see Hydrargyri nitras—m. Ammoniated, Hydrargyrum præcipitatum—m. Ammoniochloride of, Hydrargyrum præcipitatum—m. Bichloride of, Hydrargyri oxyurias—m. Biniodide of, Hydrargyri iodium rubrum, see Iodine—m. Bisulphuret of, Hydrargyri sulphuretum rubrum—m. Borate of, see Borax—m. Bromide of, see Bromine—m. with Chalk, Hydrargyrum cum cretâ—m. Chloride of, mild, Hydrargyri suburias—m. Compound Cerate of, Ceratum Hydrargyri compositum—m. Cyanide of, Hydrargyri cyanuretum—m. Cyanuret of, Hydrargyri cyanuretum—m. Deutoiodide of, see Iodine—m. Dog's, Mercurialis perennis—m. English, Chenopodium bonus Henricus—m. French, Mercurial—m. Iodide of, Hydrargyri Iodium—m. Iodide of, red, Hydrargyri Iodium rubrum.*

MERCURY, IODIDE OF CHLO'RIDE OF, Tod'idum Hydrargyri chlo'ridi, Hydrargyrum iodatum cum chlo'rido mercur'ii, (F.) Iodhydrargyrite de chinure mercurieux, Sol de Boutigny. (Made by the reaction of iodine on mild chloride of mercury.) Two iodides may be prepared—the iodide and the biniodide. Both are violent irritants; the biniodide, especially, is a powerful caustic. Both have been given in scrofula. The biniodide is only used externally like nitrate of silver in scrofulous and certain syphilitic ulcerations. An ointment of the iodide (gr. xv ad adipis ʒij) is rubbed on scrofulous tumefactions.

MERCURY, IODOARSENITE OF, Arsenic and Mercury, iodide of.—*m.* Muriate of, corrosive, Hydrargyri oxyurias.—*m.* Muriate of, mild, Hydrargyri suburias.—*m.* Nitrate of, Hydrargyri nitras.—*m.* Nitrate of, solution of, see Hydrargyri nitras.—*m.* Nitric oxyd of, Hydrargyri nitrico-oxydum.—*m.* Oxide of, black or gray, Hydrargyri oxydum cinereum.—*m.* Oxide of, red, Hydrargyri oxydum rubrum.—*m.* Oxy muriate of, Hydrargyri oxyurias.—*m.* Periodide of, Hydrargyri iodidum rubrum.—*m.* Phosphuretted, Hydrargyri phosphoratus.—*m.* Precipitate of, white, Hydrargyrum præcipitatum.—*m.* Protochloride of, Hydrargyri suburias.—*m.* Protoiodide of, see Hydrargyri iodidum, and Iodine.—*m.* Prussiate of, Hydrargyri cyanuretum.—*m.* Soluble, Hydrargyri nitras.—*m.* Subchloride of, Hydrargyri suburias.—*m.* Submuriate of, Hydrargyri suburias.—*m.* Sulphuret of, black, Hydrargyri sulphuretum nigrum.—*m.* Sulphuret of, red, Hydrargyri sulphuretum rubrum, Realgar.—*m.* Sulphuret of, with sulphur, Hydrargyri Sulphuretum nigrum.—*m.* Supernitrate of, solution of, see Hydrargyri nitras.—*m.* Tartrate of, Hydrargyri tartras.—*m.* Three-seeded, Acalypha Virginica.

MERCURY AND ARSENIC, IODIDE OF, Arsenic and Mercury, Iodide of.—*m.* and Quinia, Protochloride of, Hydrargyri et Quiniæ Protochloridum.

MERDA, Excrement.

MERDIVOMUS, (*merda*, and *vomo*, 'I vomit,') Copremetus.

MERDUS, Excrement.

MÈRE DES PIAIS (F.), ('mother of the pians,' from [L.] *mater*, 'mother,') Pian.—*m.* *de Seigle*, Ergot.

MEREMPHRAX'IS, (*mero*, and *ἐμφραξις*, 'obstruction,') Obstruction or infarction of an organ or part.

MERICUS, (μερικός,) Local.

MERIDROSIS, *Su'dor partialis seu localis*; (*mero*, and *ἰδρωσις*, 'sweating,') A partial perspiration.

MERIMNA, (μεριμνα,) Cura.

MERIMNE, (μεριμνη,) Cura.

MERISIER, (from *cerise*, [?]) *Prunus avium*.

MERISMAT'IC, *Merismat'icus*, (F.) *Mérismat'ique*; (μερισμα, gen. μερισματος, 'division,') What takes place by division;—as 'merismatic multiplication or reproduction,' that which occurs by the splitting or division of cells or of whole beings.

MERISMOPÆDIA VENTRICULI, Sarcina ventriculi.

MERKEL'S MUSCLE, Cerato-cricoid.

MERO, (from μέρος, 'a part,') in composition, a part.

MEROBALANEUM, (*mero*, and βαλανειον, 'a bath,') Bath, partial.

MEROBALNEUM, Bath, partial.

MEROCE'LE, *Me'rocele*, *Miroce'le*, *Merorrhex'is*, (*meros*, and κληη, 'tumour,') *Her'nia cruralis seu femoralis*, *Fem'oral* or *crural* *her'nia*, (F.) *Hernie crurale* ou *fémorale*. In this hernia, the viscera issue from the abdomen through the crural canal; or through an opening immediately on the outer side of Gimbernat's ligament, which gives passage to lymphatic vessels. This affection is more frequent in the female than in the male, and especially in those who have had children. The tumour, formed by merocele, is generally small, round, and more or less difficult of reduction. In other respects, it possesses all the characters of hernia. The neck of the sac has, close on the outside, the epigastric artery; above, the spermatic cord and spermatic artery in the male—the round ligament in the female; on the inside, Gimbernat's ligament; and, below, the pubes. When the obturator artery arises from the epigastric, it generally passes on the outside of and below the orifice of the sac; sometimes, however,

takes a turn above, and then to the inside of the opening. J. Cloquet asserts, that of 134 cases, in one only did he find the epigastric artery on the inside of the orifice of the sac.

The operation, required in strangulation of this variety, may be practised on the aponeurotic opening, by cutting downwards and inwards on the side of Gimbernat's ligament.

MEROCO'XALGIA, (*meros*, *coxa*, 'the hip,' and *algos*, 'pain,') Coxalgia.

MEROL'OGY, *Merolog'ia*, (F.) *Mérologie*, (from *mero*, and *logos*, 'a discourse,') The branch of anatomy, which treats of the elementary parts or constituents of the tissues and humours of the organism.

MERO'PIA, (*mero*, and *opsis*, 'vision,') A partial obscurity of vision.

MERO'RHEU'MA, gen. *Merorrhœumatis*, *Rheumatismus partialis seu localis seu Membro'rum singulo'rum*; (*mero*, and *ῥευμα*, 'defluxion, rheumatism,') Rheumatism affecting a part. Topical or local rheumatism.

MERORRHÆXIS, (*meros*, and *ῥηξις*, 'a rupture,') Merocele.

MEROS (μῦρος), Femur, Thigh.

MERRYAN'DREW. An itinerant quack, who exposes his nostrums for sale at fairs and markets. See CHARLATAN.—So called from Dr. Andrew Boorde, who lived in the reigns of Henry VIII., Edward VI., and Queen Mary, and who was in the habit of frequenting fairs and markets, at which he harangued the populace.

MERTEN'SIA VIRGIN'ICA, (after C. H. Mertens, botanist of Bremen,) *Pulmonaria Virgin'ica*, *Virgin'ian Cowslip*, *Smooth Lungwort*; indigenous. *Family*, Boraginaceæ. The plant is mucilaginous, and may be used as a demulcent.

MERULIUS, *Morchella Esculenta*—*m.* Auricula, Peziza auricula.

MERUS, *Merac'us*, *A'ceratos*, 'pure, genuine;' as *Vinum merum*, 'unmixed wine.'

MERUS (μῦρος), Femur, Thigh.

MERVINE, MALE, *Cypripedium luteum*.

MERYCISMUS (μυρκισμος), Rumination.

MÉRYCOLE, see Rumination.

MERYCOLOG'IA, (μυρκαω, 'I ruminate,' and *logos*, 'a description,') Any work on rumination may be so termed.

MERYPERTROPH'IA, (*mero*, and *hypertrophia*, 'supernutrition,') Hypertrophy of a part.

MESA, (μεσος, 'middle,') Genital organs.

MESARÆUM, (μεσαραιον, from *meso*, and *arai*, 'small intestines,') Mesentery.

MESARÆICUS, Mesenteric.

MESARAÏC, Mesenteric.

MESARTERII'TIS, (*meso*, and *arteriitis*, 'inflammation of an artery,') Inflammation of the middle layer of an artery.

MESEMAR, Mismar.

MESEMBRYANTHEMUM COPTICUM, see Soda—*m.* *Acinaciforme*, see *M. Edule*.

MESEMBRYANTHEMUM CRYSTALL'INUM, *M. ficoi'des*, (μεσημβρια, 'noon,' because its flowers expand at midday.) *Di'amon'd fig-marygold*, *Iceplant*, (F.) *Glaciale*, *Crystalline*. *Nat. Ord.* Ficoides: a plant common in the neighborhood of Cape Town. It has been recommended in dysuria, ischuria, and some other affections of the urinary organs.

MESEMBRYANTHEMUM EDU'LE, and *M. acinaciform'e*, South African plants, are much used as domestic remedies at the Cape. The expressed juice of the leaves acts as an astringent in dysentery, and as a mild diuretic. It is also used as an antiseptic gargle in malignant sore throat, violent salivation and aphthæ, and as a lotion to burns and scalds.

MESEMBRYANTHEMUM FICOIDES, *m.* *Crystallinum*.

MESEMBRYANTHEMUM TORTUO'SUM, also a South

African plant, is said to possess narcotic properties.

MÉSENCEPHALE, (*meso*, and *ἐγκεφαλος*, 'encephalon,') Medulla oblongata, Pons Varolii.

MESENTERIC, *Mesara'ic*, *Mesentericus*, *Mesara'icus*, *Mesary'icus*, (F.) *Mésentérique*, *Mésaraïque*. That which relates or belongs to the mesentery.

MESENTERIC ARTERIES are two in number, and distinguished into, 1. The *Superior mesenter'ic*, which arises from the anterior part of the abdominal aorta, below the celiac. It immediately descends to the left, forwards, behind the pancreas, and in front of the third portion of the duodenum,—to gain the superior extremity of the mesentery, where it makes a long curvature, the convexity of which is turned forwards and to the left. Towards the termination of the ileum, it ends by anastomosing with a branch of the *A. colica dextra inferior*. At its concavity, it gives off the three *Arteriae colicæ dextræ*, which belong to the great intestine; and, at its convexity, it sends off fifteen or twenty branches, which are distributed to the small intestines, after having formed numerous arches by their anastomoses.

2. The *Inferior Mesenter'ic Artery* arises from the anterior part of the abdominal aorta, an inch and a half before its termination. It descends, on the left side, behind the peritoneum; engages itself in the substance of the iliac mesocolon, forming a considerable curvature, whose convexity faces the left side. When it reaches the brim of the pelvis, it passes along the posterior separation of the mesorectum, and attains the neighbourhood of the anus, under the name *Superior hemorrhoidal artery*. It gives off no branch at its concave part; but, from the convex, the three *Arteriae colicæ sinistræ* arise.

MESENTERIC DISEASE, *Tabes mesenterica*.

MESENTERIC GLANDS are the lymphatic ganglions of the mesentery. Through them, the chyliferous vessels pass to the thoracic duct. Their uses are unknown. When diseased, nutrition is interfered with, and atrophy produced.

MESENTERIC HERNIA. If one of the layers of the mesentery be torn by a blow, whilst the other remains in its natural state, the intestines may insinuate themselves into the aperture, and form a kind of hernia. It is not known during life.

MESENTERIC PLEX'USES are furnished by the solar plexus, and have been distinguished into *superior* and *inferior*; like the mesenteric arteries which they accompany.

MESENTERIC OR MESARA'IC VEINS are two in number, and belong to the vena porta. They are distinguished into, 1. The *Superior mesenteric* or *mesara'ic* or *great mesara'ic*. This trunk receives, from above to below and on the right side, the three *venæ colicæ dextræ* and the *Gastroepiploica dextra*. Into its left side, the veins of the small intestine open. It passes in front of the transverse portion of the duodenum; and, behind the pancreas, unites with the splenic vein to concur in the formation of the vena porta.

The *inferior* or *lesser mesenteric vein* corresponds to the artery of the same name, and opens into the splenic, near the union of that vein with the superior mesenteric, and behind the pancreas.

MESENTERITIS, *Mesenteritis*.

MÉSENTERIQUE, *Mesenteric*.

MESENTERITIS, *Mesenterit'is*, *Empres'ma periton'is mesenterica*, *Enterit'is mesenterica*, *Inflammation of the mesentery*, (F.) *Mésentérite*; (*mesentery*, and *itis*.) The pain is here deeper seated and more immediately in the mesenteric region. The external tenderness is less than in some of the other varieties of peritonitis.

MESENTERIUM (*μυστερῖον*), *Mesentery*—m. *Crassum*, *Mesocolon*.

MESENTERON, *Mesentery*.

MESENTEROPHTHISIS, *Tabes mesenterica*.

MES'ENTERY, *Mesenter'ium*, *Mesara'ion*, *Mesary'um*, *Medi'um*, *Membra'na pin'guis intestinor'um*, *Luct'es*, *Mesenteron*, *Medi'um intesti'um*, *Epichor'dis*; (*μυεντεριον*, from *meso*, and *εντερον*, 'intestine,') (F.) *Mésentère*. A term in anatomy, applied to several duplicatures of the peritoneum, which maintain the different portions of the intestinal canal in their respective situations; allowing, however, more or less motion. They are formed of two laminae, between which are contained the corresponding portion of intestine and the vessels that pass to it. One only of these duplicatures has received the name *mesentery*, properly so called. This belongs to the small intestine, which it suspends and retains *in situ*. Its posterior margin, which is the smallest, is straight, and descends obliquely from the left side of the body of the second lumbar vertebra to the right iliac fossa. Its anterior margin is curved, undulating, plaited, and corresponds to the whole length of the small intestine. The mesentery contains, between the two laminae which form it, a number of lymphatic ganglions; the trunks and branches of the mesenteric vessels; the nervous plexuses accompanying them, and many lacteals and lymphatics.

MESERA, *Tutia*.

ME'SIAL, (*μεσος*, 'middle,') Relating or appertaining to the middle.

MESIAL PLANE, an imaginary plane, dividing the head, neck, and trunk into similar halves, towards right and left. Every aspect towards this plane is *mesial*; and every aspect towards right or left is *lateral*; every lateral aspect being *dextral* or *sinistral*.

MESIANUM, *Aniceton*.

MESMER'IC or **MES'MERIC**, *Mesmer'icus*; (from Anthony Mesmer, the great promulgator of animal magnetism.) Relating to mesmerism or animal magnetism:—as the 'mesmeric state,' 'mesmeric sleep,' &c. &c.

MESMERISM, *Magnetism*, animal.

MES'MERIST. A practitioner of, or believer in, mesmerism.

MESMERIZA'TION, *Magnetiza'tion*. The act of mesmerizing. The state of being mesmerized.

MES'MERIZED, *Mesmeriza'tus*, *Mag'netized*. Affected with mesmerism or animal magnetism. When the person is in a state of 'magnetic sleep,' he is said to be *hypnotized*.

MES'MERIZER, *Animal Magneti'zer*, *Mag'netizer*, (F.) *Magnétiseur*. One who practises mesmerism.

MESMEROPHRENOLOGY, *Phrenomagnetism*.

MESO, (from *μεσος*, 'middle,') in composition, middle.

MESOARIUM, (*meso*, and *οαριον*,) see *Ovary*.

MESOBlast, (*meso*, and *βλαστος*, 'germ,') See *Cytoblast* and *Molecule*.

MESOBREGMATE, (*meso*, and *bregma*, 'the top of the head,') See *Cranium*.

MESOCÆ'CUM, (*meso*, and *cæcum*). A name given to a duplicature of the peritoneum, (in some persons only,) at the posterior part of the cæcum.

MÉSOCÉPHALE, (*μεσο*, and *κεφαλη*, 'head,') Medulla oblongata, Pons Varolii.

MESOC'HON'DRIAC, *Mesochondri'acus*; (*meso*, and *χονδρος*, 'cartilage,') A name given by Boerhaave to fleshy fibres situate between the cartilaginous rings of the trachea.

MESOCOL'IC, *Mesocoli'cus*, (*meso*, and *κολον*, 'the colon,') Belonging or relating to the mesocolon.

MESOCO'LON (*μυσοκωλον*), *Mesoco'lium*, *Mesenter'ium cras'sum*. A name given to the duplicatures of the peritoneum, which fix the different parts of the colon to the abdominal parietes. It

has received different names, according to its situation. The *right lumbar mesocolon* fixes the ascending colon to the corresponding lumbar region. The *transverse mesocolon* arises from the concave arch of the colon, and forms a septum between the epigastric and umbilical regions. Its inferior portion is continuous with the mesentery. The *left lumbar mesocolon*, which contains the ascending colon, is continuous below with the *iliac mesocolon*. The last includes between its layers the sigmoid flexure of the colon, and ends in the mesorectum. Under the right kidney, it is narrow and firm, and forms the *right ligament of the colon*: at the under end of the left kidney, it forms the *left ligament of the colon*.

MESOCRANIUM, (*meso*, and *κρανιον*, 'skull.') Sinciput.

MESOCRANIUM, (*μεσοκρανιον*, from *meso*, and *κρανιον*, 'skull.') Sinciput.

MESODERMUM, (*meso*, and *δερμα*, 'skin.') Corpus mucosum.

MESODME, (*μεσοδμη*, 'a crossbeam,' from *meso*, and *δομος*, 'a house.') Mediastinum.

MESODMITIS, (*mesodme*, and *itis*.) *Mesotæchi'tis*, *Mediastini'tis*, *Inflammati'o mediasti'ni*. Inflammation of the mediastinum.

MESOËPIDID'YMIS, (*meso*, and *ἐπιδιδυμις*, 'epididymis.') A double serous fold, like the mesocolon, formed by the tunica vaginalis passing from the body of the epididymis, and uniting it to the testicle.

MESOGASTRIUM, (*meso*, and *γαστηρ*, 'belly.') Umbilical region.

MESOGASTRIUM MÜLLERI. A portion of mesentery, which passes from the vertebral column to the left curvature of the stomach.

MESOGLOSSUS, (*meso*, and *γλωσσα*, 'tongue.') Genioglossus.

MESOGLOTTUS, (*meso*, and *γλωττα*, 'tongue.') Genioglossus.

MESOGONIUM, (*meso*, and *γονυ*, 'the knee.') Internodium.

MESOLO'BAR. Belonging or relating to the *Mesolobe* or *Cor'pus callo'sum*.

MESOLOBAR ARTERIES, *Arte'riæ mesolob'icæ*, *A. cor'poris callo'si cer'ebrî*, are the arteries of the corpus callosum.

MESOLOBE, *Corpus callosum*.

MESLOBUS, (*meso*, and *λοβος*, 'a lobe.') Corpus callosum.

MESOMËTRE, *Mesome'trium*; (*meso*, and *μητρα*, 'uterus.') The fold of the peritoneum, which unites the uterus to the abdominal parietes.

MESOME'RIA, (*μησομηρια*, from *meso*, and *μηρος*, 'the thigh.') The parts of the body situate between the thighs and hips.

MESOMERION, *Perinæum*.

MESOMPHALUM, (*meso*, and *ομφαλος*, 'navel.') Umbilicus.

MESOPHLEBITIS, (*meso*, and *phlebitis*.) Inflammation of a vein. Inflammation of the middle layer of a vein.

MESOPHLEBIUM, (*meso*, and *φλεβις*, gen. *φλεβος*, 'a vein;') *Interve'nium*. The space between two veins.

MESOPH'RYON (*μεσοφρυον*), *Glabb'la*, *Gabb'la*, *Intercil'ium*, *Metop'ium*, *Na'sal em'innence*, (*meso*, and *οφρυα*, 'the eyebrows.') The part between the eyebrows.

MESOPLEURIUM, (*meso*, and *πλευρον*, 'a rib.') Intercostal space.

MESOPLEURIUS, *Intercostal*.

MESOR'CHIUM, (*meso*, and *ορχις*, 'a testicle.') A duplicature of the peritoneum, which supports the testicle in its passage from the abdomen into the scrotum.

MESOREC'TUM, (*meso*, and *rectum*, 'the intestine rectum.') A triangular reflection, formed by the peritoneum, between the posterior surface

of the rectum and the anterior surface of the sacrum. Between the two layers of which the mesorectum is composed are found much areolar tissue, and the termination of the inferior mesenteric vessels.

MESOSCELO, (*meso*, and *σκελος*, 'leg.') *Perinæum*.

MESOSCELOCEL'E, (*mesoscelon*, and *κηλη*, 'a rupture.') *Hernia*, *perineal*.

MESOSCELOPHY'MA, (*mesoscelon*, and *φυμα*, 'swelling.') A tumour of the perinæum.

MESOTH'ENAR, (*meso*, and *θεναρ*, 'the thenar,' 'the palm of the hand.') A muscle, which carries the thumb towards the palm. Winslow applied the term to the abductor pollicis, united to the deep-seated portion of the flexor brevis pollicis.

MESOT'ICA, (from *μεσος*, 'middle.') Diseases affecting the parenchyma. Pravity in the quantity or quality of the intermediate or connecting substance of organs; without inflammation, fever, or other derangement of the general health. The first Order, class *Eccritica* of Good.

MESOT'ECHITIS, (*mesotæchum*, and *itis*.) *Mesodmitis*.

MESOT'ECHIUM, *Mediastinum*.

MESOT'ECHIUM, (*μεσοτοιχον*, 'a partition wall,' from *meso*, and *τοιχος*, 'a wall.') *Mediastinum*.

MES'PILUS, (*μεσπιλη*, from *meso*, and *πιλος*, 'a ball,' from the resemblance of the fruit to half a ball.) *Mes'pilus German'ica* seu *domest'ica*. *Ord. Rosaceæ*. *The med'lar*, (F.) *Néflier*. The fruit (F.) *Nêfle*, and seeds of the medlar, have both been used medicinally:—the immature fruit as an astringent, and the seeds in nephritic diseases. See *Cratægus Aria* and *Sorbus aucuparia*.

MESPILUS ARIA, *Cratægus aria*—m. *Domestica*, *Mespilus*, *Sorbus domestica*—m. *Germanica*, *Mespilus*—m. *Intermedia*, *Mespilus oxyacantha*—m. *Lævigata*, *Mespilus oxyacantha*.

MESPILUS OXYACAN'THA, *M. Oxyacanthoï'des* seu *interme'dia* seu *læviga'ta*, *Oxyacantha*, *Spi'na al'ba*, *White Haw'thorn*. (F.) *Aubépine*. The flowers of this uncommon European plant are sometimes used in infusion as a pectoral.

MESPILUS OXYACANTHOIDES, *Mespilus oxyacantha*.

MESQUITE, GUM. A gum, obtained in New Mexico from a tree of the same family as the *Acacia*. It has the same virtues as gum Arabic.

META, *μετα*, *Meth'*, *μὲθ'*, 'with,' 'together with,' 'after,' 'change of form and place.' A common prefix to terms.

METAB'ASIS, *Trans'itus*, *Metab'oliz*, *Metabol'ia*, *Metab'lage*, *Metallur'gic*, (*μεταβασις*, from *μεταβαινω*, (*meta*, and *βαινω*, 'I go,' 'I digress.') A change of remedy, practice, &c.

METABOLE, (*μεταβολη*, from *μεταβαλλειν*, 'to transpose,') *Metabasis*, *Transformation*.

METABOLEL'OGY, *Metabolelog'ia*; (*μεταβολη*, (*meta*, and *βαλλω*), 'change,' and *λογος*, 'a description.') A description of the changes which supervene in the course of a disease.

METABOLIA, (*μεταβολια*.) *Metabasis*.

METABOL'IC, (*μεταβολικος*.) *Metabol'icus*. Appertaining to change or transformation.

METABOL'IC FORCE. *Via metabol'ica*. A term employed by Schwann for the power possessed by living cells of changing the character of the substances brought in contact with them.

METACARP'AL, *Metacarpia'nus*, (F.) *Métacarpien*, (from *metacarpus*.) Relating or belonging to the metacarpus.

METACARPAL ARTERY, *Arte'ria dorsa'lis metacar'pi*, arises from the radial, at the moment it engages itself in the upper extremity of the abductor indicis. It descends obliquely upon the back of the hand. Its branches are distributed to the abductor indicis, and the integuments: some communicate with the dorsalis carpi.

METACARPAL ARTICULA'TIONS are those by

which the last four metacarpal bones are united together at their upper extremity.

METACARPAL BONES, or bones which compose the metacarpus, are five in number; and distinguished into *first, second, third, &c.*, beginning from the outer or radial side. They are articulated by their superior extremity with the bones of the second range of the carpus; and by the lower with the first phalanges.

METACARPAL LIG'AMENT, *Inferior palmar Lig'ament*, is a fibrous band, stretched transversely before the inferior extremities of the last four metacarpal bones, which it keeps in their respective positions.

METACARPAL PHALAN'GES are the first phalanges of the fingers; so called, because they are articulated with the bones of the metacarpus.

METACARPAL RANGE or ROW of the carpal bones, (F.) *Rangée métacarpienne du carpe*, is the lower row of carpal bones; so called because they are articulated with the bones of the metacarpus. It is composed of the trapezium, trapezoides, magnum, and unciniforme.

METACARPEUS, Abductor metacarpi minimi digiti.

MÉTACARPIEN, Metacarpal—*m. du Petit doigt*, Opponens minimi digiti—*m. du Pouce*, Opponens pollicis.

METACARPION, (μετακάρπιον,) Metacarpus.

METACARPOPHALANGÆUS POLLICIS, Abductor pollicis manus.

METACAR'POPHALAN'GIAN, *Metacarpophalangianus*, *Metacarpophal'angal*, (F.) *Métacarpophalangien*. That which belongs to the metacarpus and phalanges.

METACARPOPHALANGIAN or METACARPOPHALANGAL ARTICULA'TIONS are formed by the bones of the metacarpus and the corresponding phalanges. In these articulations, the bony surfaces are incrustated with cartilage, covered by a synovial membrane, and kept in connection by means of an anterior and two lateral ligaments.

MÉTACARPOPHALANGIEN LATÉRAL SUS-PALMAIRE, ET MÉTACARPO-PHALANGIEN LATÉRAL, Interosseus manus.

METACARP'US, (meta, and καρπος, 'wrist,') *Metacarp'ion*, *Postbrachia'te*, *Postcar'pium*, *Tor'sus ma'nus*, (F.) *Métacarpe*. The part of the hand comprised between the carpus and fingers. It is composed of five parallel bones, called *metacarpal*; forming the back of the hand, externally, and the palm internally.

METACERASMA, gen. *Metacerasum'atis*, (μετακέρasma, from meta, and κεραννμι, 'to mix,') *Cerasma*.

METACHEIRIXIS, (meta, and χειρ, 'the hand,') *Surgery*.

METACHIRISIS, *Surgery*.

METACHIRISMUS, *Surgery*.

METACHORESIS, (μεταχώρησις, from meta, and χωρεω, 'to go,') *Metastasis*.

METACHYSIS, (meta, and χεω, 'to pour,') *Transfusion*.

METACINEMA, gen. *Metacine'matis*, (μετακίνημα, from meta, and κινω, 'to move,') *Metastasis*.

METACINESIS, (μετακίνησις,) *Metastasis*.

METACONDYLE, see Phalanx.

METACYE'SIS, (meta, and κυνσις, 'pregnancy,') *Gravid'itas extraiter'ina secunda'ria*. Extrauterine pregnancy in which the foetus is at first in the uterus, but subsequently in some other place.

METADERMATO'SIS, (meta, and δερμα, gen. δερματος, 'skin,') A morbid development of the epidermis or epithelium.

METAGEN'ESIS, (F.) *Metagénèse*, *Altern'ate generation*, (meta, and γέναις, 'generation,') A term, proposed by Professor Owen, to express the changes of form, which one animal species undergoes in a series of successively produced

individuals, extending from the one developed from the ovum to that which becomes the image or last perfected individual. This term he employs in contradistinction to *metamorphosis*, which he restricts to the changes of form undergone by one and the same individual. See *Generation*, *alternate*.

METAL, (μεταλλον,) *Metal'lum*. A class of simple, bodies; in general, distinguished from others by considerable specific gravity; a particular splendour; almost total opacity; insolubility in water; and the property they have of ringing when struck. Metals have no effect, except of a mechanical nature, when taken into the stomach; unless they have already undergone, or undergo in the stomach, oxidation or union with an acid; when, at times, deleterious compounds may be formed. Copper cents; half-pence; quicksilver; lead, have frequently been swallowed in the metallic state with impunity. Tin and mercury are the only metals prescribed for a mechanical effect; the former as an anthelmintic, —the latter, idly enough, in cases of fancied intussusception.

MÉTAL DES CLOCHES, Bell-metal.

METALLAGE, (μετάλλαγη, from meta, and αλασσω, 'I exchange,') *Metabasis*.

METALLAXIS, (μετάλλαξις,) *Metallage*.

METALLIC VOICE, *Tintement métallique*.

METALLODYN'IA, (μετάλλων, 'a metal,' and δυννη, 'pain,') Pain owing to the injurious influence of a metal—as lead, quicksilver, &c.

METALLUM, (μετάλλον,) *Metal*—*m. Hamato poeticum*, *Ferrum*.

METAMORPH'ICI, (Morbi,) (μεταμορφωσις, (meta, and μορφη, 'form,') *I transform.*) Developmental diseases; special diseases, the incidental result of the formative reproductive and nutritive powers. (W. Farr.)

METAMORPHO'SIA, *Phantasmascopia*, *Phantasmatoscop'ia*, (μεταμορφωσις, (from meta, and μορφη, 'form,') 'I transform,' and οψις, 'vision,') *Suffusio*, *S. Oculorum seu metamorphops'ia seu myo'des*, *Imagination* (Maitre-Jean), *Crup'sia*, *Marmar'ygge*, *Marmor'ygge*, *Mac'ula an'te oculos volitan'tes*, *Maraug'ia*, *Oculi marmar'ygges*, *Vis'io phantas'matum*, *Vis'io defigur'atus*, *False sight*, (F.) *Berlue*. Aberration of the sense of sight, which transmits the image of imaginary objects. This affection sometimes depends on a slight opacity in the transparent parts of the eye: the cornea, crystalline, or vitreous humour, when it is symptomatic and of no consequence. At other times, it appears to be idiopathic; and occurs particularly in those who have been in the habit of constantly fixing their eyes on very brilliant or small bodies. The objects fancied to be seen are various. They are sometimes circular, perpetually moving; or shining, or black spots, cobwebs, insects, or pieces of wool—when they are often termed *Mus'cæ seu Flo'ci volitan'tes*, (F.) *Mouches volantes*, and the condition *Vis'io musca'rum*, *Scotom'ata*, *Suffusio Myo'des*, *Myodeop'sia*, *Myi'odop'sia*, *Myodeop'sia*, *Myodesop'sia*, *Visus musca'rum*. These appearances sometimes continue for a few days; being dependent on the state of the nerves of the individual at the time; or they may exist for life, and ultimately impair the sight. Certain of them change their position, and appear to be seated in the humours of the eye, and—it has been supposed—in the vitreous humour more especially: hence the term *ento-hyaloid muscæ* applied to them.

If the affection be symptomatic, it is of but little moment. If idiopathic, and connected with any excitement of the brain, which is not often the case, attention will have to be paid to that organ.

METAMORPHOSIS, (μεταμορφωσις, from meta, and μορφη, 'to shape,') see *Metagenesis*.

METAMORPHOSIS VENTRICULI GELATINIFORMIS, Gastromalacia.

METANGISMUS, (μεταγγισμός, from *meta*, and ἀγγίον, 'a vessel,') Decantation.

METAPEDIUM, (*meta*, and ποδίου, dim. of πούς, gen. ποδός, 'a foot,') Metatarsus.

METAPHRENON, (μεταφρενον, from *meta*, and φρενός, 'the diaphragm,') Dorsum.

METAPHYTEIA, (μεταφυτεω, from *meta*, and φυω, 'to produce,') 'to transplant,') Transplantation.

METAPODIUM, (*meta*, and ποδίου, dim. of πούς, gen. ποδός, 'a foot,') Metatarsus.

METAPOROPŒA, *Metaporopœsis*, (*meta*, πορος, 'a pore, and ποιεῖν, 'to make,') The change produced in the minute pores, in the capillary extremities of vessels,—when they pass from the morbid to the healthy condition. (?)

METAPTO'SIS, (μεταπτωσις, *Metaschematism'us*, *Metaschemat'isis*. Mutation, change; from μεταπτω, (*meta*, and πτω, 'I fall,') 'I digress,') Any change in the form or seat of a disease. Transformation.

METARRHŒA, (μεταρρῳα, *Metar'rhy'sis*, (*meta*, and ῥεω, 'I flow,') Reflux. The transfer of a disease from without to within, or from one part to another.

METARRHYSIS, (μεταρρῳις, *Metarrhœa*.

METASCHEMATISIS, (μετασχηματισμός, from *meta*, and σχημα, 'form,') Metaptosis.

METASCHEMATISMUS, (μετασχηματισμός, *Metaptosis*.

METAS'TASIS, (μεταστασις, *Metachore'sis*, *Metacine'ma*, *Metacine'sis*, (*meta*, and ἵστημι, 'to place,') *Displace'ment*, *Translation*. A change in the seat of a disease; attributed, by the Humorists, to the translation of the morbid matter to a part different from that which it had previously occupied; and by the Solidists, to the displacement of the irritation. It has also been used in the same extensive sense as Metaptosis. Disputes have often been indulged, whether a case of metastasis ought not rather to be esteemed one of extension of the disease. The phenomena of gout and acute rheumatism are in favour of metastasis occasionally supervening.

METASTASIS LACTIS, Galactoplasia, Phlegmatia dolens.

METASTAT'IC, (μεταστατικός, *Metastat'icus*, (F.) *Métastatique*. Belonging or relating to metastasis. A metastatic crisis is one produced by metastasis:—a metastatic affection, one caused by metastasis, &c. See Abscess, metastatic.

METASYN'CRISIS, (μετασυνκρίσις, from *meta*, and συνκρανω, 'I join,') *Recomposition'ion*. A word employed by some disciples of Asclepiades. This physician supposed, that animals are formed by the union or assemblage of atoms. He designated all bodies by the word συγκρίματα, which signifies 'assemblage.' Συγκρίνεται, 'to be assembled,' was, with him, synonymous with 'to exist,' and διακρίνεται, 'to separate,' was, with him, synonymous with 'to dissolve,' 'to cease to exist.' The word *Metasyncretisis* was invented to express the recomposition of bodies after their momentary dissolution. Some have rendered the word by the Latin *Recorporatio*.

METASYNCRIT'IC, (μετασυνκριτικός, *Metasyncret'icus*, *Recorporat'ivus*. Belonging or relating to metasyncretisis. A name formerly given to medicines to which was attributed the virtue of producing the metasyncretisis or regeneration of the body, or of some of its parts.

METASYNCRITIC CYCLE meant a determinate series of remedies employed for this purpose.

METATARSAL, *Metatar'seus*, (F.) *Métatarsien*. Relating or belonging to the metatarsus.

METATARSAL ARTERY, *Artère sus-métatarsienne* of Chaussier, arises from the *Arteria dorsalis Tar'si*; passes transversely over the back of the

foot; and furnishes, at its convexity, which is anterior, three branches, called *Arteriæ interos'seæ dorsales Pedis*. These branches are distributed in the interosseous spaces.

METATARSAL ARTICULA'TIONS are those resulting from the junction of the metatarsal bones with each other. They are strengthened, behind, by dorsal and plantar ligaments; and before, by a transverse metatarsal ligament, which is plantar, and has the greatest analogy to the inferior transverse metacarpal ligament.

METATARSAL BONES, (F.) *Os métatarsiens*, are five in number, and distinguished by their number; first, second, third, &c., reckoning from the outer side.

METATARSAL PHALANG'ES are the first phalanges of the toes; so called because they are united to the metatarsus.

METATARSAL ROW—(F.) Rangée métatarsienne—of the bones of the tarsus, is the second row, or that contiguous to the metatarsus; comprehending the cuboides and three cuneiform bones: some add the scaphoides.

METATARSEUS, see Abductor minimi digiti pedis.

MÉTATARSIEN, *Metatarsal*.

MÉTATARSIIUM, *Metatarsus*.

METATARSOPHALANG'IAN, *Metatarsophalangianus*, *Metatarso-phalangal*. Relating to the metatarsus and phalanges.

METATARSOPHALANGIAN or **METATARSOPHALANGAL ARTICULATIONS** are formed by the bones of the metatarsus and the corresponding phalanges. They bear the greatest analogy to the metacarpophalangeal articulations.

MÉTATARSÓ-PHALANGIEN-LATÉRAL, see *Interossei pedis*—*m. Sous-phalangien transversal du premier orteil*, *Transversus pedis*.

METATARS'US, *Metatar'sium*, *Metape'dium*, *Metapod'ium*, *Præcor'dium* seu *Sol'um ped'is*, *Vestig'ium ped'is*, (F.) *Métatars*, (*meta*, 'after,' and τάρπος, 'the tarsus,') That part of the foot which is situate between the tarsus and toes, corresponding with the metacarpus. It is composed of five parallel bones; one to each toe. Like the metacarpal bones, they are developed from two points of ossification.

METATH'ESIS, (μεταθεσις, from μετα-ἵστημι, (*meta*, and ἵστημι, 'to place,') 'I change place,') *Transposit'io*, *Transposit'ion*. An operation, by which a morbid agent is removed from one place to another, where it may produce less disturbance in the exercise of the functions:—as, for example, in the operation of depressing cataract, or when calculus in the urethra is pushed back into the bladder. Also, Derivation.

METATIO, (*metor*, *metatus*, 'to measure,') Mensuration.

METATOC'IA, (*meta*, and τοκος, 'birth,') Parturition in a preternatural manner. Preternatural labour.

METAPTOSIS, (μεταπτωσις, from μεταπτω, 'to change place,') *Diadexis*.

METENSOMATOSIS, (μετενσωματωσις, from *meta*, εν, and σωμα, 'body,') *Incorporation*.

METEORISMUS, (μετεωρισμός, from μετεωρίζω, 'to elevate,') *Sublimation*, *Tympanites*.

METEOROL'OGY, (μετεωρολογία, *Meteorology'ia*, (*μετεωρος*, (*meta*, and αἰωρεω, 'I lift up,') 'a meteor,' and λόγος, 'a discourse,') The science, whose object is the knowledge of the origin, formation, appearance, &c., of meteors. The state of the atmosphere has a most important bearing upon the health of animals. Every epidemic affection has its causes seated there. Meteorological conditions have, hence, ever been attended to by the physician for the purpose of detecting the precise character of any particular epidemic influence. The barometer, thermometer, and hygrometer are the instruments used with this intent—to de-

fect, as well the varying weight or pressure, and the temperature, as the moisture. Perhaps, of the three conditions, the last exerts more influence in the production of disease than either of the others. Our knowledge, however, of this part of physics is extremely limited and unsatisfactory.

METER, (μετρον, 'a measure,') A suffix to words denoting 'a measure,' as in *Barometer*, *Pleximeter*, &c. Also uterus, (from μετρον.)

METH. A prefix to terms. See *Meta*.

METHÆMACHYMIA, (meth, αἷμα, 'blood,' and χέω, 'to pour,') Transfusion.

METHÆMATA, (meth, and αἷμα, 'blood,') The capillary or intermediate system of vessels in which the blood undergoes the change from venous to arterial, and conversely. See *Capillary vessels*.

METHE, (μεθη,) *Temulentia*.

METHEGLIN, ([W.] meddylglynn,) *Melizomum*.

METHEMERINUS, (μεθημερινος, 'daily,' from meth, and ημερα, 'a day,') *Quotidian*.

METHOD, (μεθοδος,) *Meth'odus*, (meth, and ὁδος, 'way,') This word has different acceptations in the sciences. In medicine, *curative method*, *meth'odus medendi*, is the methodical treatment of disease.

METHOD OF COUNTING, *Method*, numerical—*m. Mariano's*, see *Lithotomy*.

METHOD, NUMERICAL, *Method of Observation* or of *Counting*, of Louis, (F.) *Méthode numérique*, consists in observing every case and every symptom of a case numerically, so as to ensure, as far as practicable, accuracy of observation; and to enable us, by the analysis and collation of such facts, to deduce general laws and conclusions. Also, the application of numbers to the study of disease. See *Statistics*, medical.

METHOD, READY, of Dr. Marshall Hall, see *Ready Method*—*m. Sylvester's*, see *Sylvester's method*.

MÉTHODE NUMÉRIQUE, *Method*, numerical, *Statistical*, medical—*m. Perturbatrice*, *Perturbatrix* (*Medicina*)—*m. Stibio-dermique*, see *Unquenum antimonii*.

METHODICAL COMPRESSION, see *Compression*.

METHODICS, *Methodists*.

MÉTHODIQUES, *Methodists*.

METHODISTS, *Methodical sect*, *Method'ics*, (F.) *Methodistes*, *Methodiques*. A sect of physicians whose doctrine was in vogue after that of the Empirics and Dogmatists, towards the end of the first century. According to the Methodists, of whom *Themison* was the chief, almost every disease is dependent on contraction or relaxation—*strictum* or *laxum*. To these two causes, they added a third—*mixed* or *compound*—to include those affections which partook of the two characters. The doctrine, *Medic'ina method'ica*, *Methodism'us*, *Meth'odism*, resembled, in some respects, that of *Brown*.

METHODOLOGY (MEDICAL), *Methodolog'ia Med'ica*, (μεθοδος, 'method,' and λογος, 'a discourse,') A word used, by the French more especially, to signify *method* applied to the study of any science. *Medical Methodology*, consequently, means *method* applied to the study of medicine.

METHODUS, (μεθοδος,) *Method*—*m. Cataleptica*, see *Cataleptic*—*m. Celsiana*, see *Lithotomy*—*m. Curatoria*, *Therapeutics*.

METH'ODUS DERIVAT'RIA. The derivative or revellent system of treatment.

METHODUS ENDERMATICA seu ENDERMICA, see *Endermic*—*m. Franconiana*, see *Lithotomy*—*m. Guytoniana*, see *Lithotomy*—*m. Medendi*, *Method of cure*, *Therapeutics*.

METHOMA'NIA, (μεθη, 'drunkenness,' and μανα, 'mania,') An irresistible desire for intoxicating substances. *Temulentia*.

METHU, (μεθυ,) *Wine*. See *Temulentia*.

METHYLE, HYDRATED OXIDE OF, *Spiritus pyroxylicus*.

METHYLENE, BIHYDRATE OF, *Spiritus pyroxylicus*.

METHYSIS, (μεθυσις,) *Temulentia*.

METHYSMUS, (from μεθη, 'intoxication,') *Temulentia*.

METHYS'TICUS, (μεθυστικός,) That which causes, or pertains to, drunkenness.

METHYSTOPHYLLUM GLAUC'UM, (μεθυστικός, 'intoxicating,' and φυλλον, 'a leaf,') A South African plant, *Nat. Ord. Amyridaceæ*, an infusion of whose leaves is pleasant to the taste, and is used in bronchitis, asthma, and other thoracic diseases. With the Bushmen and others, it is a favourite beverage, and is called by them *Bosch-jeesmanthee*, 'Bushman's tea.' It is, also, chewed.

MÉTIS, (from mictus, 'mixed,') see *Hybrid*.

MÉTISSAGE, *Crossing*.

METODONT'IASIS, (meta, and ὀδοντίαςις, 'dentition,') Faulty development of the teeth.

METOPAGES, (μετωπον, 'front,' and πηννυμι, 'to fix,') *Symphcephalus*.

METOPANTRA, (metopon, and αντρον, 'a cave,') *Frontal sinuses*.

METOPANTRAL'GIA, (metopantra, and αλγος, 'pain,') *Pain* in the frontal sinuses.

METOPANTRI'TIS, *Inflammat'io sinuum frontali'um*, (metopantra, and itis,) *Inflammation* of the frontal sinuses.

METOPION, *Bubon galbanum*.

METOPIMUM, (μετωπιος, 'pertaining to the forehead,') *Mesophryon*.

METOPODYNIA, (metopon, and ὀδυνη, 'pain,') *Neuralgia frontalis*.

METOPON, (μετωπον, from meta, and ψ, 'the eye,') *Front*, *Frontal bone*.

MET'OPOSCOPY, *Metoposcop'ia*, (μετοσκοπια, from metopon, and σκοπειν, 'to view,') The art of knowing the temperament and character of a person by inspecting the traits of his forehead or face. See *Physiognomy*.

METRA, *Uterus*.

METRAE, (μετραι, pl. of μετρα, 'the womb,') *Secundines*.

METREMORRHAGIA, (metra, and hæmorrhagia,) *Metrorrhagia*.

METREMORRHOÏDES, (metra, and αἱμορροιδες, 'hemorrhoids,') *Hemorrhoids* or varicose veins at the orifice of the uterus.

METRAL'GIA, *Metrodyn'ia*, *Uteral'gia*, (metra, and αλγος, 'pain,') *Pain* in the uterus. See *Hysteralgia*.

METRANÆ'MIA, (metra, and anæmia,) An anæmic condition of the uterus.

METRANASTROPHE, (metra, and αναστροφή, 'inversion,') *Uterus*, inversion of the.

METRANEURYSMUS, (metra, and ανευρυσμος, 'dilatation,') *Metreuryisma*.

METRATRE'SIA, *Imperforat'io u'teri*, (metra, and αντροσις, 'imperforation,') An unnatural closure of the uterus.

METRATROPH'IA, (metra, α, 'privative,' and τροφη, 'nourishment,') *Atrophy* or want of development of the uterus.

METRAUX'Ë, *Hypertroph'ia seu Sarco'sis u'teri*, (metra, and αυξη, 'increase,') *Hypertrophy* of the uterus.

METRECTA'SIA, (metra, and εκτασις, 'dilatation,') *Dilatation* of the womb.

METRELCO'SIS, *Metrelco'sis*, (metra, and ἔλκος, 'an ulcer,') *Ulceration* of the uterus.

METREMPHRAX'IS, (metra, and εμπρασσω, 'I obstruct,') *Obstruction* of the womb or of the vessels of the womb—*Infarc'tus u'teri*. A name under which some authors have confounded chronic inflammation of that viscus, and the different degenerations to which it is exposed.

METREMPHYSEMA, Physometra.

METREN'CHYTA, *Metren'chytes*, (*metra*, and *εἰχυν*, 'I inject.') Injection of the uterus. Substances injected into the uterus are called *Metren'chyta*.

METREURYS'MA, gen. *Metreury'smatis*, *Metreury'smus*, (*metra*, and *εὐρύσμα*, 'dilatation.') A morbid dilatation of the womb.

METRHELICOSIS, *Metrelcosis*.

METRHYMENITIS, (*metra*, *ὑμην*, 'membrane,' and *itis*), see *Metritis*.

METRIA, (from *metra*), Puerperal fever.

METRIOPATH'IA, (*μετρία*, 'tempered,' and *πάθος*, 'affection.') State of an individual whose passions are temperate.

METRI'TIS, (*metra*, and *itis*), *Fe'bris uteri'na*, *Hysteri'tis*, *Empres'ma Hysteri'tis*, *Inflammati'o Uteri*, *Metrophlogo'sis*, *Inflammation of the Uterus* or *Womb*, (F.) *Inflammation de la matrice*, *Hysterite*. The characteristic symptoms of this affection are:—pain, swelling, and tenderness in the hypogastric region; with heat, pain, and tenderness of the os uteri; vomiting, smallness, and frequency of pulse. It occurs most frequently after delivery, *Lochometri'tis*, when there is generally suppression of the lochial discharge. The treatment must be vigorous,—bleeding early, so as to make a decided impression; followed by a full dose of opium, fomentations, blisters, &c.

Acute inflammation of the womb, eated in its internal membrane, *Endometri'tis*, *Metrhymenit'is*, *Enmetri'tis*, has been called *Inflammati'o catarrh'al'is u'teri* or *Acute' catarrh'*, (F.) *Catarrhe aigu*, of that viscus. It is known by the discharge of a clear, stringy fluid *per vaginam*; preceded by pains, which, from the hypogastric region, shoot to the thighs, groins, &c., with more or less fever. It requires the antiphlogistic treatment. *Chronic metritis* sometimes succeeds the acute. To it must be referred the indurations, observed in the uterus, and many of the leucorrhœal discharges to which females are subject.

METRITIS RHEUMATICA, *Metrorrhœma*—m. *Septica*, *Metrocæcæ*—m. *Venosa*, see *Phlebitis*.

METROBLENNORRHŒA, *Leucorrhœa uteri*.

METROBLENNOEZEMIA, (*metra*, *βλιννα*, 'mucus,' and *ζημία*, 'loss,' *Leucorrhœa uteri*.

METROBOTRYTES, (*metra*, and *βωτρης*, 'a cluster of grapes,') Cauliflower excrecence of the uterus.

METROC'ACE, *Metri'tis sep'tica*, *Cacome'tra*, *Cacome'tria*, (*metra*, and *κακος*, 'evil,') Putrescency or gangrene of the uterus.

METROCAMPSIS, (*metra*, and *καμψις*, 'a bending,') *Hysteroloxia*.

METROCARCINO'MA, gen. *Metrocarcino'matis*, *Hystero'carcino'ma*, *U'teri carcino'ma*, *Canc'er u'teri*, (*metra*, and *καρκινωμα*, 'cancer,') Cancer of the uterus.

METROCATARRHUS, (*metra*, and *κατάρρως*, 'catarrh,') *Leucorrhœa uteri*.

METROCELE, (*metra*, and *κηλη*, 'a rupture,') *Hysterocele*.

METROCELIDES, (*μητηρ*, 'mother,' and *κελιδ*, 'a mark,') *Nævus*.

METROCHOLÓ'SIS, (*metra*, and *χολος*, 'bile,') *Fe'bris puerper'al'is bilio'sa*.

METRODYNIA, (*metra*, and *δυνη*, 'pain,') *Metralgia*.

METROHÆMIA, (F.) *Hypermétrohémie*, (*metra*, and *αἷμα*, 'blood,') *Hyperæmia* or congestion of blood in the uterus.

METROHEMORRHAGE, *Metrorrhagia*.

METROLEUCORRHŒA, *Leucorrhœa uteri*.

METROLOXIA, (*metra*, and *λοξος*, 'oblique,') *Hysteroloxia*.

METROMALACOMA, gen. *Metromalaco'mutis*, '*metra*, and *μαλακος*, 'soft,') *Hysteromalacia*.

METRCMALACOSIS, *Hysteromalacia*.

METROMANIA, *Nymphomania*.

METROMY'ZEUS, (*metra*, and *μυζω*, 'I suck,') An exhausting syringe, to be applied to the os uteri for exhausting the womb in cases of amenorrhœa.

METRONCUS, (*metra*, and *ογκος*, 'a tumour,') *Hysteroncus*.

METROPARALYSIS, (*metro*, and *παράλυσις*, 'paralysis,') Paralysis of the uterus.

METROPATH'IA, (*metro*, and *πάθος*, 'affection,') An affection of the womb.

MÉTRO-PÉRITONITE, *Metropéritonitis*—m. *Puerpérale*, see *Peritonitis*.

METROPÉRITONITIS, (*metra*, and *peritonitis*), *Inflammati'o u'teri et peritonæ'i*, (F.) *Métropéritonite*. Inflammation of the uterus and peritoneum. Puerperal Fever. See *Peritonitis*.

METROPHLEBITIS, *Phlebometri'tis*, (*metra*, *φλεψ*, gen. *φλεβος*, 'a vein,' and *itis*.) Inflammation of the veins of the uterus. See *Phlebitis*.

METROPHLEBITIS PUERPERALIS, see *Phlebitis*.

METROPHLOGOSIS, (*metra*, and *φλογωσις*, 'inflammation,') *Metritis*.

METROPHTHISIS, *Phthi'sis uteri'na*, (*metra*, and *φθιω*, 'I consume,') Consumption from ulceration of the uterus—*U'tera seu Helco'sis u'teri*.

METROPHYMA, gen. *Metrophy'matis*, (*metra*, and *φύμα*, 'a swelling,') *Hysteroncus*.

METROPHYSETERION, (*metra*, and *φυσσω*, 'I inflate,') An instrument for applying gasiform agents to the uterus.

METROPLETHO'RA, *Metrypene'mia*, (*metra*, and *πληθωρα*, 'fulness,') Congestion or hyperæmia of the uterus.

METROPOL'YPUS, *Hys'teropol'ypus*, *Polypus U'teri*, (*metra*, and *πολύπος*.) *Polypus of the uterus*.

METROPROPTOSIS, (*metra*, *pro*, and *πτωσις*, 'a falling,') *Prolapsus uteri*.

METROPTOSIS, *Prolapsus uteri*.

METRORRHAG'IA, *Metrorrhag'ia*, *Hæmorrhagia uteri'na* seu *u'teri*, *Sanguifluxus uteri'nus*, *Sanguinis stillicid'ium ab U'tero*, *Fluor uteri'ni san'guinis*, *Profluvium San'guinis ex u'tero*, *Hysterorrhag'ia sanguinea*, *Hystero'rrhœa*, *Hæmorrhœa uteri'na* (of some), *Menorrhag'ia* (of some), *Hæmatome'tra*, *Hæmatoma'tra*, *U'terine Hem'orrhage*, *Hem'orrhage from the Womb*, *Flooding*, *Metrohem'orrhage*, (F.) *Hémorrhagie de la Matrice*, *Pertes*, *Pertes utérines rouges*, *P. de sang*, (*metra*, and *ρήγνυμι*, 'I break forth,') An effusion of blood from the inner surface of the uterus, either at the menstrual or other period; but in a greater quantity than proper. Uterine hemorrhage may be caused by those influences which produce hemorrhage in general. It happens, however, more frequently during pregnancy, and during or after delivery, when the vascular system of the uterus is so circumstanced as to favour its occurrence more than at other times. The termination of metrorrhagia is usually favourable. Should it, however, be very copious, or frequently recur; or should it happen to a great extent after delivery, death may take place very speedily; and, in some cases, without the discharge being perceptible; constituting *internal hemorrhage*.

Uterine hemorrhage may be *active* or *passive*; requiring obviously a different treatment. The main management is similar to that of hemorrhage in general;—the horizontal posture; acid drinks; free admission of cool air; cold applications to the loins, thighs, and abdomen; injection of cold water, even of iced water, into the vagina; plugging the vagina, so as to prevent the discharge *per vaginam*, and thus induce a coagulum in the mouths of the bleeding vessels. Such will be the *special* plan adopted where the hemorrhage has occurred in one not recently delivered. In uterine hemorrhage after delivery, the same cooling plan must be followed; but, as the flow of

blood is owing to the uterus not contracting so as to constrict its vessels, pressure must be made on the abdomen to aid this; and, if necessary, the hand must be introduced into the uterus to stimulate it to contraction. Should the female be excessively reduced, so as to render the accoucheur apprehensive that she may expire from loss of blood, excitants may be exhibited. The profuse exhibition of opium in such cases is, at least, a doubtful plan.

Transfusion has, at times, been practised as a last resource.

METORRHEU'MA, gen. *Metorrhœum'atis*, *Rheumatism'us Uteri*, *Hysteral'gia rheumat'ica* seu *catarrh'al'is*, *Metrit'is rheumat'ica*, (F.) *Rheumatisme de l'utérus*; (metra, and *ῥευμα*, 'defluxion,' 'rheumatism.') Rheumatism of the uterus.

METORRHEXIS, (metra, and *ῥηξις*, 'rupture.') Uterus, rupture of the.

METORRHE'IA, (metra, and *ῥευω*, 'I flow.') A protracted discharge of any fluid from the uterus.

METROSALPINGITIS, (*metrosalpinx*, and *itis*, 'inflammation.') Inflammation of the Fallopian tube.

METROSALPINGORRHEXIS, (*metrosalpinx*, and *ῥηξις*, 'rupture.') Rupture of a Fallopian tube.

METROSALPINX, gen. *Metrosalpin'gis*, (metra, and *σαλπιγξ*, 'a trumpet.') Tube, Fallopian.

METROSCIRRHUS, *Hysteroscir'rhus*, *Scir'rhus* seu *Carcin'oma uteri*, (metra, and *σκίρρως*, 'scirrhous.') Scirrhous of the uterus.

METROSCOLIOMA, (metra, and *σκολιος*, 'oblique.') Hysteroxolia.

ME'TROSCOPE, *Metroscop'ium*, (metra, and *σκοπεω*, 'I view.') An instrument, invented by M. Nauche, for listening to the sounds of the heart of the foetus in utero gestation, when the sounds and movements are imperceptible through the parietes of the abdomen. The extremity of the instrument—the first notion of which was given to M. Nauche by the stethoscope of Laënnec—is introduced into the vagina and applied against the neck of the uterus.

METROSIDEROS GUMMIFERA, see Kino.

METROSTERE'SIS, (metra, and *στερησις*, 'privation.') Extirpation of the uterus. Want of uterus.

METROSYNIZE'SIS, *Hysterosynize'sis*, (metra, and *συνιζις*, 'running together.') Morbid union of the uterus to adjacent parts.

MÉTROTOME, Hysterotomus.

METROTOMIA, (metra, and *τομή*, 'section.') Cæsarean section.

METRYPERÆMIA, (metra, and *hyperæmia*), Metroplethora.

METRYPERÆSTHESIS, (metra, and *hyperæsthesia*.) See Uterus, irritable.

MEU, *Æthusa meum*.

MEULIÈRE, (from *mola*, 'a mill.') Molar.

MEUM, (from *μειον*, 'small,' in allusion to the leaves.) *Æthusa meum*—*m.* *Anethifolium*, *Æthusa meum*—*m.* *Athamantium*, *Æthusa meum*.

NEURTRISSION, (from (L.) *mors*, gen. *mortis*, 'death.') Contusion.

MEVIUM, Syphilis.

MEZEA (*μίζα*), Genital organs.

MEZEREON, *Daphne mezereum*.

MEZEREUM, *Daphne mezereum*—*m.* *Acetatum*, see *Daphne mezereum*.

MIAMMA gen. *Miamm'atis*, (*μιαμμα*), Miasma.

MIANSIS (*μιανσις*), Miasma.

MIARIA (*μιαρα*), Miasma.

MIAS'MA, pl. *Mias'mata*, *Mi'asm*, (*μιασμα*, 'a stain,' from *μιανω*, 'I contaminate.') *Miam'ma*, *Mian'sis*, *Mia'ria*, *Mias'mus*, *Inquinamentum*, *Moly'sis*, *Contagium*. The word *miasm* has, by some, been employed synonymously with contagion. It is now used more definitely for any

emanation, either from the bodies of the sick, or from animal and vegetable substances, or from the earth, especially in marshy districts, (*Marsh poison*, *Efflu'vium palus'træ*, *Gas palus'træ*, *Limnomeneph'itis*, (F.) *Intoxication des Marais*), which may exert a morbid influence on those who are exposed to its action. To these terrestrial emanations—the *Koinomias'mata* of Dr. E. Miller, of New York—the Italians give the name *aria cattiva*, but, more commonly, *malaria* ('bad air'); a word which has been adopted into those languages. The deleterious effluvia, originating from the decomposition of matter derived from the human body, have been called by Dr. Miller, *Idiomias'mata*; the epithets *Koino* and *Idio* being derived respectively from *κοινος*, 'common, and *ιδιος*, 'personal.' Of the miasms which arise either from the animal body or from the most unhealthy situations, we know, chemically, nothing. All that we do know is, that, under such circumstances, emanations take place, capable of causing disease in many of those who are exposed to their action.

MIASMAT'IC, *Miasmatic'us*. Belonging or relating to miasmata. Sauvages, in his classification of diseases according to their etiology, has a class under the name *Morbi miasmatic'i*, *miasmatic' diseases*; and *Miasmatici (morbi)* is an order of the class *Zymotici* of Dr. Wm. Farr. A fever that arises from marshy miasms, is styled *malaria*, *paludal*, *Helop'gra*, *Helopyr'etous*, &c.

MIASMUS, Miasma.

MI'ALULANTES, (*mi'auler*, 'to mew,' formed by onomatopœia.) See *Convulsionnaire*.

MICÆ PANIS, ('crumbs of bread,') see *Triticum*.

MICATIO CORDIS, (*micare*, *micatum*, 'to move quickly,') Systole.

MICHE'LIA CHAM'PACA, (after P. A. Micheli, a Florentine botanist.) *Champaca*, *Michelia Tojampaca*. Ord. Magnoliaceæ. An Oriental tree, much prized for the odour and beauty of its flowers. The oily infusion of the flowers is employed in the Moluccas in headache.

MICLE'TA. A medicine used by Mesue for arresting hemorrhage; perhaps, according to Siebenhaar, from the Arabic, *micnata*, "proved by experience."

MICRANATOMIA, (*micro*, and *anatomia*), see Anatomy.

MICRANGIA, (*micro*, and *αγγιον*, 'a vessel,') Capillary vessels.

MICRENCEPHALIUM, (*micro*, and *εγκεφαλος*, 'encephalon,') Cerebellum.

MICRENCEPHALON, Cretinism.

MICRENCEPHALUM, Cerebellum.

MICRO, (*μικρος*, 'small,') in composition, smah **MICROCEPHALIA**, (*micro*, and *κεφαλη*, 'head,') See Microcephalus.

MICROCEPH'ALUS, *Mikroceph'alus*, *Micro cran'ius*. One who has a small head. A monster having a small imperfect head, or a small imperfect cranium. The condition is called *Microceph'al'ia*, *Craniostenosis*, (F.) *Microcéphalie*: with some it means Idiocy.

MICROCORIA, (*micro*, and *κορη*, 'pupil,') Myosis.

MI'CROCOSM, *Microcos'mus*, (*μικροκοσμος*), (*micro*, and *κοσμος*, 'world,') A little world. Some philosophers have given this appellation to man, whom they consider as the epitome of all that is admirable in the world. The world they call *Ma'crucosm*, (from *μακρος*, 'great,')

MICROCOSMETOR, (*micro*, and *κοσμητωρ*, (from *κοσμεω*,) 'a regulator,') Vital principle.

MICROCOSM'ICA MACH'INA. The organism of man.

MICROCOSMICA SCIENTIA, Physiology.

MICROCOSMICUM SAL, with the ancients meant the salts of the urine.—*Sal urinæ*.

MICROCOSMOGRAPHIA, (*microcosmus*, and *γραφία*, 'a description.') Physiology.

MICROCOSMUS, *Microcosm*.

MICROCOUSTIC, *Microcous'ticus*, (*micro*, and *ακουω*, 'I hear.') This word, as well as *Microph'onus*, (*μικρος*, 'small,' and *φωνη*, 'voice,') means any thing that contributes to increase the intensity of sound—as the speaking trumpet—by collecting the sonorous rays.

MICROCRANIUS, (*micro*, and *κρανιον*, 'cranium.') *Microcephalus*.

MICROGLOS'SIA, (*micro*, and *γλωσσα*, 'tongue.') Original smallness of tongue.

MICROLEUCONYMPHÆA, (from *micro*, λευκος, 'white,' and *nymphaea*.) *Nymphæa alba*.

MICROLOGY, (*μικρολογία*, from *micro*, and *λογος*, 'a discourse,') See *Microscopy*.

MICROM'ELUS, *Hemim'eles*, (*micro*, and *μελος*, 'a limb.') A monster having imperfectly developed extremities.

MICROMMATUS, (*μικροματος*, from *micro*, and *ομμα*, gen. *ομματος*, 'the eye.') *Microphthalmus*.

MICROPHONOUS, (*micro*, and *φωνη*, 'voice,') See *Microcoustic*.

MICROPTHAL'MUS, *Microm'matus*, (*micro*, and *οφθαλμος*, 'eye,') One who has small eyes. A monster with two small, or imperfectly developed eyes. A small eye,—*Ophthalmidium*, *Ophthalmium*.

MICROPODIA, *Micropus*.

MICROPROS'PUS, *Aproso'pus*, (*micro*, and *προσωπον*, 'face,') A monster in which a part of the face is absent.

MIC'ROPUS, (*micro*, and *πους*, gen. *ποδος*, 'foot,') One who has small feet. The condition is called *Micropod'ia*.

MIC'ROPYLE, *Microp'yla*, (*micro*, and *πυλη*, 'gate,') An opening in the epispem of seeds, through which the fecundating pollen enters the ovule to fecundate it. It has been supposed, that such a micropyle exists in the human ovule.

MICRO'RCHIDES, (*micro*, and *ρχις*, gen. *ρχιδος*, 'a testicle,') They who have very small testicles.

MIC'ROSCOPY or **MICROS'COPY**, *Microscop'ia*; (*micro*, and *σκοπη*, 'a view,') Observation by the microscope; an important agency in the examination of the healthy and morbid tissues. A description of the results of such observation is *Micrography*. *Micrology* is often used with the same signification, and for a treatise on the microscope and the results of microscopy.

MICROSOMATIA, (*micro*, and *σωμα*, gen. *σωματος*, 'body,') *Nanosomia*.

MICROSOMIA, (*micro*, and *σωμα*, 'body,') *Nanosomia*.

MICROSPHYC'TUS, (*micro*, and *σφυγμος*, 'pulse,') Smallness or weakness of pulse.

MICROSPORON AUDOUIN, (*micro*, and *σπορον*, 'a seed, a spore,') See *Fungus*, and *Porrigo decalyans*—*m. Furfur*, see *Chloasma* and *Fungus*—*m. Mentagrophytes*, *Mentagrophyte*, see *Fungus*.

MICROS'TOMUS, (*micro*, and *στομα*, 'mouth,') One who has a small mouth.

MICROTE'SIA, (*μικροτης*, 'smallness,') *Par'tium organica'rum par'vitas morbo'sa*. Morbid smallness of organic parts.

MICROTRAPEZIA, (*micro*, and *τραπεζα*, 'a table,') *Leptotrophia*.

MICROTRICH'IA, (*micro*, and *τριχ*, gen. *τριχος*, 'hair,') Fineness or shortness of hair.

MICROTROPHIA, (*micro*, and *τροφή*, 'nourishment,') *Leptotrophia*.

MICTIO, (*mingere*, *mictum*, 'to pass the urine,') Micturition—*m. Cruenta*, *Hæmaturia*—*m. Inop'ortuna*, *Enuresis*—*m. Involuntaria*, *Enuresis*—*m. Sanguinea*, *Hæmaturia*.

MICTION, Micturition.

MICTERIT'ION, *Mictio*, *Mictus*, *Ure'sis*, *Uri'asis*, *Omiche'sis*, (F.) *Miction*, *Action de pisser*, *Pissement*, (*micturio*, (*mingere*, *mictum*,) 'I make water,') The act of making water—*Urina'tion*. Also, morbid frequency of passing the water.

MICTUS, Micturition—*m. Cruentus*, *Hæma'turia*.

MIDRIFF, (from Sax. *mid*, 'middle,' and *hrif*, 'the bowels,') Diaphragm.

MID'WIFE, (*mid*, 'with,' and *þig*, 'wife,' or *með*, 'meed,' 'recompense,' and *þig*, 'wife,') *Mat'rona*, *Obstet'rix*, *Mul'lier obstetrix*, *Mæ'a*, *Mæ'as*, *Mau'tria*, *Aces'toris*, *Aces'tris*, *Aces'tria*, *Saga*, (Sc.) *How'dy*, (Prov.) *Gravewife*, *Hau'dwoman*, *Heads'woman*, *How'dywife*, (F.) *Sage femme*, *Accoucheuse*, *Ventrière*, *Matrone*, *Levandière*. A female who practises obstetrics.

MIDWIFERY, *Obstetrics*.

MIEL, *Mel*—*m. d'Acéte de Cuivre*, *Liniment æruginis*—*m. Boraté*, *Mel boracis*—*m. Dépuré*, *Mel despumatum*—*m. Despumé*, *Mel despumatum*—*m. Médicinal*, *Mellitum*—*m. Mercuriale composé*, *Mellitum de mercuriali compositum*.

MIELLÉ, *Melitus*.

MIGMA (*μυγμα*), *Mistura*.

MIGRAINE, (abridged from) *Hemicrania*.

MIGRANA, *Hemicrania*.

MIGRATION DU LAMBEAU, ('migration of the flap,') See *Flap*.

MIKANTIA GUACO, (after Prof. J. Mikan, botanist of Prague,) *Guaco*.

MICROCEPHALUS, *Microcephalus*.

MIL, (from *milium*, 'millet,') *Panicum milia-ceum*.

MILFOIL, (*mille*, 'a thousand,' and *folium*, 'a leaf,') *Achillea millefolium*.

MILIAIRE, *Miliary*, *Miliary fever*.

MILIARIA, (*milium*, 'millet,') *Miliary fever*—*m. Sudans*, *Miliary fever*—*m. Sudatoria*, *Miliary fever*.

MILIARIS, *Miliary*—*m. Sudatoria*, *Sudor Anglieus*, *S. Picardicus*.

MILI'ARY, *Mil'aria*, (F.) *Mil'aire*. Resembling millet seed,—hence, *miliary fever*, *miliary glands*, &c.

MILI'ARY FE'VER, *Em'phlysis milia'ria*, *Milia'ris*, *Mil'aria*, *M. sudans seu sudato'ria*, *Su'dor*, *S. milia'ris*, *Pap'ula milia'ris*, *Ecanthe'ma milia'ria*, *Spy'ochus milia'ria*, *Aspre'do milia'cea*, *Fel'bris milia'ris seu essero'sa seu puncticula'ris seu cuticula'ris seu vesicula'ris seu lenticula'ris seu purpura'te rub'ra et al'ba milia'ris*, *Pur'pura puerpera'rum*, *P. milia'ris*, *Pap'ule sudo'ris*, *Mil'let seed rash*, (F.) *Mil'aire*, *Millet*, *Pourpre blanc*, *Millet*. It is so called from the eruption resembling the seed of the *milium* or *millet*. Fever, accompanied by an eruption of small, red, isolated pimples, rarely confluent, but almost always very numerous, slightly raised above the skin, and presenting, at the end of 24 hours, a small vesicle filled with a white transparent fluid, which quickly dries up, and separates in the form of scales. Miliary fever is now rare, in consequence of the cooling practice in fevers and other states of the system. It is almost always brought on by external heat, and hence the prevention is obvious. The treatment is simple. It requires merely the antiphlogistic regimen, in ordinary cases.

MILIARY GLANDS, *Sebaceous glands*—*m. Granulations* or *Tubercles*, see *Granulation*.

MILII SEMINA, ('seeds of millet,') *Panicum Italicum*.

MILI'OLUM, (dim. of *milium*, 'millet,') *Cenchr'id'ion*. A small tumour on the eyelids, resembling, in size, a millet-seed.

MILITARY HERBA, *Achillea millefolium*.

MIL'ITARY, *Mil'aria*, (*miles*, *mil'itis*, 'a sol-

dier.') (F.) *Militaire*. Pertaining to a soldier or an army—as '*Military medicine*,' *Medicina militaris*, (F.) *Médecine militaire*; and *Military surgery*, *Chirurgia militaris*, (F.) *Chirurgie militaire*. Medicine and surgery applicable to the soldier.

MILKUM, *Grutum*, *Panicum miliaceum*, see Hydatid—*m. Esculentum*, *Panicum miliaceum*—*m. Indicum*, *Panicum Italicum*—*m. Panicum*, *Panicum miliaceum*—*m. Solis*, *Lithospermum officinale*.

MILK, (Sax. *melc*.) *Lac*, *Ga'la*, *Hu'mor lac'teus*, *La'tex ni'veus*, (F.) *Lait*. A fluid secreted by the mammary glands of the females of the mammalia. The skimmed milk of the cow, (F.) *Lait de Vache*, contains water, caseous matter, traces of butter, sugar of milk, chloride of sodium, phosphate, and acetate of potassa, lactic acid, lactate of iron, and earthy phosphates. The cream is formed of butter, casein, and whey, in which there is sugar of milk and salts. When examined by the microscope, milk is seen to contain a large number of particles, of irregular size and shape, varying from $\frac{1}{32000}$ th to $\frac{1}{30000}$ th of an inch in diameter, floating in a colorless fluid—*liquor lactis*. They consist of oily matter, surrounded by a delicate pellicle, and are the *milk globules*.

Cow's milk is employed for the preparation of cream, butter, cheese, whey, sugar of milk, and frangipane. It is useful in a number of cases of poisoning; either by acting as a demulcent, or by decomposing certain poisons, or by combining with others so as to neutralize them. It is constantly employed as aliment, and may be regarded as an emulsion in which butter and casein are found in suspension. When taken into the stomach, it is coagulated by the gastric fluids, and the coagulum is digested like any other solid. The watery parts are absorbed.

Between milk, flour, and blood, there is great similarity of composition. The following table is given by Dr. Robert Dundas Thomson:

MILK.	FLOUR.	BLOOD.
Curd or Casein	Fibrin,	Fibrin,
	Albumen,	Albumen,
	Casein,	Casein,
	Gluten,	Colouring matter.
Butter . . .	Oil.	Fat.
Sugar . . .	Sugar,	Sugar.
	Starch,	
Chloride of potassium,		
—sodium,		
Phosphate of soda,		
—lime,	Do.	Do.
—magnesia,		
—iron,		

* From a considerable number of experiments, Messrs. Deyeux and Parmentier class the six kinds of milk, which they examined, according to the following table, as regards the relative quantity of materials they contain.

CASEIN.	BUTTER.	SUGAR OF MILK.	SERUM.
Goat	Sheep	Woman	Ass
Sheep	Cow	Ass	Woman
Cow	Goat	Mare	Mare
Ass	Woman	Cow	Cow
Woman	Ass	Goat	Goat
Mare	Mare	Sheep	Sheep

MILK ABSCESS, *Mastodynia apostematosa*.

MILK, **ALMOND**, *Emulsio amygdalæ*.

MILK AND SODA WATER. An agreeable mode

of taking milk in cases where it lies heavily on the stomach. Heat, nearly to boiling, a teacupful of milk; dissolve in it a teaspoonful of sugar, put it into a large tumbler, and pour over it two-thirds of a bottle of soda water.

MILK, AS'SES', (F.) *Lait d'Anesse*, considerably resembles human milk, of which it has the consistence, smell, and taste; but it contains a little less cream, and more soft, caseous matter. It is often used by those labouring under pulmonary affections.

Artificial Asses' Milk may be made by taking gelatin $\frac{3}{4}$ ss; dissolving it, by the aid of heat, in barley water Oij; adding refined sugar $\frac{3}{4}$ j; pouring into the mixture new milk Oj; and beating the whole with a whisk.

It may also be prepared by dissolving sugar of milk $\frac{3}{4}$ j in tepid skimmed cow's milk Oj.

MILK BLOTCH, *Porrigo larvalis*—*m. Crust*, *Porrigo larvalis*.

MILK, EWES, (F.) *Lait de Brebis*. It affords more cream than cows' milk; but the butter is softer. The caseum, on the contrary, is fatter, and more viscid. It contains less serum than cow's milk. The Roquefort cheeses are made from it.

MILK FEVER, see Fever, milk—*m. Glass*, *Breast glass*—*m. Globules*, see Milk.

MILK, GOATS', (F.) *Lait de Chèvre*, resembles cows' milk: the butyrateous matter, however, which enters into its composition, is more solid than that of the cow.

Artificial Goat's Milk may be made by taking fresh suet $\frac{3}{4}$ j; cutting it into small pieces; tying it in a muslin bag, large enough to leave the morsels free from compression; and boiling in a quart of cow's milk, sweetened with a quarter of an ounce of white sugar candy.

Used as a diet in scrofulous cases, and also in phthisis.

MILK, GREEN, *Colostrum*.

MILK, HU'MAN, (F.) *Lait de femme*,—contains more sugar of milk, and cream, and less caseum, than cows' milk. Its composition differs according to the distance of time from delivery.

MILK KNOT, *Galactogan'gion*; (F.) *Ganglion lacteum*. The knots often observed in the breast after inflammation of the organ, or for some time after the suppression of the secretion. They generally end by resolution.

MILK LEG, *Phlegmatia dolens*.

MILK, MALES', *Sperm*.

MILK, MARES', (F.) *Lait de Jument*,—contains only a small quantity of fluid butyrateous matter; a little caseum, softer than that of cows' milk, and more serum.

MILK SCALL, *Porrigo larvalis*.

MILK SICK'NESS, *Sick stom'ach*, *Swamp sick'ness*, *Tires*, *Slows*, *Stiff joints*, *Pu'king fe'ver*, *Riv'er sick'ness*. A disease occasionally observed in the States of Alabama, Indiana, and Kentucky, which affects both man and cattle, but chiefly the latter. It is attributed in cattle to something eaten or drunk by them; and in man to the eating of the flesh of animals labouring under the disease. Owing to the tremors that characterize it in animals, it is called the *Trem'bles*. It is endemic.

The symptoms of the disease are such as are produced by the acronarcotic class of poisons—vomiting, purging, extreme nervous agitation, &c.: and the approved indications of treatment appear to be—gentle emetics and laxatives, with quiet, and mucilaginous drinks.

MILK, SNAKES', *Euphorbia corollata*.

MILK, SOLIDIFIED. Milk, to which little less than an equal weight of white sugar has been added, evaporated to dryness. When redissolved in water, it has all the qualities of fresh milk, with the additional quantity of sugar.

To milk, evaporated to dryness, *M. Griman*2

and Galais have given the names *Lactoline* and *Lacteine*.

MILK SPOTS, *Maculæ albæ*, *Strophulus*—m. Stone, *Morochthus*.

MILK, SUGAR OF, Sugar of milk.

MILK OF SULPHUR, Sulphur præcipitatum—m. Sweet, Galium verum—m. Teeth, see Dentition—m. Thrush, Aphthæ—m. Vetch, Astragalus verus—m. Vetch, stemless, Astragalus exscapus—m. Weed, Apocynum androsæmifolium, Asclepias Syriaca, Euphorbia corollata—m. Weed, long-leaved, green, Acerates longifolia—m. Weed, smooth, Asclepias Sullivantii—m. Wort, bitter, Polygala amara—m. Wort, common, Polygala—m. Wort, dwarf, Polygala paucifolia—m. Wort, rattlesnake, Polygala.

MILKWOMAN, Nurse, wet.

MILKY, Lactic.

MILL MOUNTAIN, Linum catharticum.

MILLEFEUILLE, ((L.) mille, 'a thousand,' and folium, 'a leaf,') Achillea millefolium—m. Aquatique, Phellandrium aquaticum.

MILLEFEUILLET, Omaseum.

MILLEFOLIUM, Achillea millefolium.

MILLEMORBIA SCROPHULARIA, (mille, 'a thousand,' and morbus, 'a disease') Scrophularia nodosa.

MILLEPEDES, (mille, 'a thousand,' and pedes, 'feet,') Onisci aselli.

MILLEPERTUIS ORDINAIRE, (from mille, 'a thousand,' and pertuis, 'a hole,' because the leaf, when held to the light, seems pierced with holes,') Hypericum perforatum.

MILLET, (from milium, 'millet,') Aphthæ, Miliary fever—m. Barbadoes, Panicum Italicum—m. Common, Panicum miliaceum—m. Gray, Lithospermum officinale—m. Indian, Panicum Italicum—m. des Oiseaux, Panicum Italicum—m. Ordinaire, Panicum miliaceum.

MILLETSEED RASH, Miliary fever.

MIL/LIGRAMME, (mille, 'a thousand,' and gramme, 'a gramme,') The thousandth part of a gramme:—about 0.0154 Troy grain.

MILLIGRANA, Herniaria glabra.

MILLIMÈTRE, Millim'eter, (mille, 'a thousand,' and μέτρον, 'measure,') The thousandth part of the metre:—equal to 0.03937 English inch, or about twofifths of a line.

MILLOT, Miliary fever.

MILPHÆ, (μυλφαί), Madarosis.

MILPHOSIS, (μυλφωσις), Madarosis.

MILT (Sax.), Melt, Spleen.

MILTLIKE. An epithet for a form of cancer, characterized by a soft pale growth, containing scarcely any blood;—so called from its resemblance to the milt of a fish.

MILTLIKE TUMOUR, Encephaloid.

MILTUS, Plumbi oxydum rubrum.

MILTWASTE, Asplenium.

MIMOSA CATECHU, (from μῖμος, 'a mimic,' the leaves of many species mimicking sensibility,') Catechu—m. Cochliacarpa, see Cortex adstringens Brasiliensis—m. Leucophea, see Spirit (Arrack)—m. Nilotica, see Acacia gummi—m. Scandens, Cachang-parang—m. Senegal, see Senegal, gum—m. Virginalis, see Cortex adstringens Brasiliensis.

MIND, ABSENCE OF, Aphelxia socors—m. Abstraction of, Aphelxia intentia—m. Unsound, Insanity.

MINERALIUM, Antimonium.

MIN'ERS EL'BOW. An enlargement of a bursa over the olecranon, occurring in such as are in the habit of leaning much upon it; and, therefore, often seen in those who work on the side in low-roofed mines.

MIN'IM, Min'imum, 'the least part or portion.' The 60th part of a fluidrachm. This measure was introduced by the London College of Physi-

cians, in consequence of the uncertainty of the size of the drop, (see Gutta.) The subdivision of the wine pint has, accordingly, been extended to the 60th part of the fluidrachm; and glass measures, called "*minim measures*," have been adopted by the London College. The proportion between the minims and the drops of various fluid preparations is given under Gutta.

MINIM'ETER, (minimium, and μέτρον, 'measure,') An instrument for measuring minims, invented by Mr. Alsop. It consists of a glass tube, graduated from the conical point into minims; and having a piston, by the elevation of which, fluid may be drawn into the tube, and by its depression be forced from it.

MINIM MEASURE. A measure for the divisions of a fluidrachm.

MINIMUM, Minim.

MINISTER GYMNASTÆ, Pædotribes.

MINITHOSIS, (from μινῶω, 'to diminish,') Meiosis.

MINIUM, Plumbi oxydum rubrum—m. Græcorum, Hydrargyri sulphuretum rubrum—m. Purum, Hydrargyri sulphuretum rubrum.

MINOR HELICIS, ('smaller muscle of the helix,') A muscle situate on the posterior border of the helix of the ear, at its commencement in the fossa of the concha.

MINORATIF, Laxative.

MINORATIVUS, (from minor, 'less,') Laxative.

MINT, Mentha—m. Cat, Nepeta—m. Cock, Tanacetum balsamita—m. Curled, Mentha crispa—m. Horse, Monarda coccinea and M. punctata—m. Horse, hairy, Blephilia hirsuta—m. Horse, Ohio, Blephilia hirsuta—m. Horse, sweet, Cunila Mariana—m. Hyssop-leaved, Mentha cervina.

MINT JU'LEP. A drink, consisting of brandy, sugar, and pounded ice, flavoured by sprigs of mint. It is an agreeable alcoholic excitant.

MINT, MOUNTAIN, Monarda coccinea—m. Mountain, common, Pycnanthemum incanum—m. Pepper, Mentha piperita—m. Spear, Mentha viridis—m. Squaw, Hedeoma—m. Stone, Cunila Mariana.

MINT TEA, Infusum Menthæ viridis.

MINT, WATER, Mentha aquatica.

MINTHA, (μνθῆ), Mint.

MINUTHESES, (from μινῶω, 'to diminish,') Meiosis.

MINU'TTO MON'ACHI, ('restriction of the monk,') A term said to have been used in the middle ages for any treatment adapted for 'lessening' the sexual energy.

MINYANTHES, Menyanthes.

MIOSIS, Meiosis.

MIRACH, Abdomen.

MIRE (old F.), A physician or surgeon.

MIRLES, Rubeola.

MIROCELE, Meroccele.

MIRROT, Daucus Carota.

MISADIR, Ammonia murias.

MISANTHRO'PIA, (μισανθρία,) *Misan'thropy*, *Exanthro'pia*, *Phyganthro'pia*, (μῖσος, 'hatred,' and ἀνθρωπος, 'man,') Aversion to man and society;—a symptom of melancholy, and hypochondriasis.

MISCARRIAGE, Abortion.

MISCEGENATION, (miscere, 'to mix,' and genus, 'a race,') A mixture or amalgamation of races, as of the black with the white.

MISE-BAS, [F.] ('laid down,') Parturition.

MISERERE MEI, ('have pity on me,') Iteus.

MISERY (from miser, 'suffering,') see Pain.

MISHME TEETA, Coptis teeta.

MIS'MAR, *Mecenar*. A name given by Avicenna to a kind of nodus, which forms on the

toes as the sequel of contusion or inflammation of those parts.

MISOG'YNY, *Misogyn'ia*, (μῖσος, 'hatred,' and γυνή, 'a woman.')

 Hatred of women. Repulsion for sexual intercourse.

MISOPTO'CHUS, (μῖσος, 'hatred,' and πτωχος, 'poor.')

 That which has hatred for the poor. The gout has been so called by some, because it commonly affects the rich.

MISSADAN, Hydrargyrum.

MISSIO SANGUINIS, Bloodletting.

MISSISSIP'PI, MIN'ERAL WA'TERS OF. *Cooper's Well*, in Hinds County, four miles from Raymond, and thirteen from Jackson, is a sulphureous saline, the chief constituents being the sulphates of soda and magnesia. Of the solid ingredients upwards of 3 per cent. are of peroxide of iron. The waters are chiefly used in chronic gastroenteric affections. *Ocean Springs*, in Jackson County, is an acidulous sulphureous chalybeate.

MISTIO, (from *miscere*, *mistum*, 'to mix,') Mistura.

MISTLETOE, ([Sax.] *mýrtel-tan*.) *Viscum album*—m. Yellowish, *Viscum flavesens*.

MISTU'RA, *Mis'tis*, *Propolis'ma*, *Mig'ma*, *Mis'tio*, *Miz'tio*, *Miz'tura*, *Mix'ture*; (F.) *Potion*, *Mé-lange*. A mingled compound, in which different ingredients are contained in the fluid state, or suspended or not by means of mucilaginous or saccharine matter. In this sense, it is synonymous with the French *Potion*. In France, however, the word *mixture* is more frequently understood to mean a liquid medicine, which contains very active substances, and can only be administered by drops. A mixture, in other words, in the French sense, may be regarded as a *potion* deprived of watery vehicle. When it consists of materials, that require to be shaken up when taken, it is sometimes termed *Mistu'ra m'edia*.

MISTURA, (S.) A fragrant yellowcoloured water, used as a perfume by the ladies of Peru. It is prepared from gillyflower, jasmine, and *flor de mistela* (*Talinum umbellatum*). See *Campomanesia lineatifolia*.

MISTURA ACALÆ, *Mucilago Acaciæ*.

MISTURA AC'IDI HYDROCYAN'ICI, *Mix'ture of Prussic acid*, *Mélange pectoral* (Magendie). (*Medicinal Prussic acid*, $\mathfrak{z}$ j, *Distilled water*, $\mathfrak{z}$ xiv, *Pure sugar*, $\mathfrak{z}$ iss. A dessertspoonful every morning and evening, as a pectoral, &c.

MISTURA AMMONI'ACI, *Lac ammoniaci*, *Mix'ture of ammo'niac*. (*Ammoniac*. $\mathfrak{z}$ ij, *aquæ Oss.* Rub the ammoniacum: adding the water gradually until they are perfectly mixed. Ph. U. S.) Dose, $\mathfrak{f}$ 3ss to $\mathfrak{z}$ j.

MISTURA AMYGDALÆ, *Emulsio amygdalæ*.

MISTURA ASAFOETIDÆ, *Lac asafetidæ*, *Emulsio antihysterica*, *Mix'ture of asafetida*. (F.) *Lait d'asafetida*. (*Asafetide* $\mathfrak{z}$ ij, *Aquæ Oss.* Rub together, adding the water by degrees. Ph. U. S.) Used where pills cannot be swallowed, and as a glyster in irritations during dentition, and in ascariæ. Dose, $\mathfrak{f}$ 3ss to $\mathfrak{f}$ 3iiss.

MISTURA ASAFOETIDÆ PRO CLYSMATE, *Enema fœtidum*.

MISTURA CAM'PHORÆ, *A'qua cam'phoræ* (Ph. U. S.), *Cam'phor mix'ture* or *julep*, *Mistu'ra camphora'ta*, *Julepium e cam'phorâ seu camphora'tum*, (F.) *Julep camphrê*. (*Camphor*. $\mathfrak{z}$ ij, *Alcohol*, $\mathfrak{m}$ xl, *Magnes. carb.* $\mathfrak{z}$ iv, *Aquæ Oij*. Rub the camphor with the alcohol, and afterwards with the magnesia; add the water, and filter. Ph. U. S.) Virtues like those of camphor. See *Emulsio camphoræ*.

MISTURA CHLOROFOR'MI, *Mix'ture of Chloroform*. (*Chloroform. purif.* $\mathfrak{z}$ ss. *Camphor. gr. lx*, *Vitell. ovi. i*, *Aquæ*, $\mathfrak{f}$ 3vj. Rub the yolk with the camphor previously dissolved in the chloroform, and gradually add the water. Ph. U. S., 1863.) Dose, $\mathfrak{f}$ 3ij to $\mathfrak{z}$ ss.

MISTURA CONTRACTA, ('a small mixture,') see *Gutta*.

MISTURA COR'NU USTI, *Decoc'tum cor'nu cervini seu al'buni*, *Decoc'tion of burnt harts'horn*, *Mix'ture of burnt harts'horn*, *White Decoc'tion*, (F.) *Decoc'tion blanche*. (*Cornuum. ust.* $\mathfrak{z}$ ij, *Acaciæ gum. cont.* $\mathfrak{z}$ j, *Aquæ Oijj*. Boil to Oij constantly stirring, and strain. Ph. L.)

MISTURA CREASO'TI, *Creasote mix'ture*. (*Creasot.*, *Acid. acetic.* $\mathfrak{a}\mathfrak{a}$ $\mathfrak{m}$ xvj, *Sp. Junip. comp.*, *Syrup.*, $\mathfrak{a}\mathfrak{a}$ $\mathfrak{f}$ 3j; *Aquæ* $\mathfrak{f}$ 3xiv. Mix the creasote and the acid; add gradually the water, and then the syrup and spirit. Ph. E.) A fluidounce contains a minim of creasote.

MISTURA CRE'TÆ, *Po'tio carbonat'is cal'cis*, *Mistu'ra creta'cea*, *Mucilo'go cre'tica*, *Chalk or Creta'ceous mix'ture*, *Chalk julep*, (F.) *Mixture de Craie*. (*Crete* pp. $\mathfrak{z}$ ss, *Sacchar.*, *Acaciæ gum. in pulv.*, $\mathfrak{a}\mathfrak{a}$ $\mathfrak{z}$ ij, *Aquæ cinnam.*, *Aquæ*, $\mathfrak{a}\mathfrak{a}$ $\mathfrak{f}$ 3iv. *Misce. Ph. U. S.*) Antacid and absorbent. Dose, $\mathfrak{f}$ 3j to $\mathfrak{f}$ 3ij.

MISTURA DIABOL'ICA. A mixture under this name is kept in many hospitals, for malingerers. It is made of sundry nauseous ingredients, as aloes, asafetida, castor, &c., and is given so as to keep up a disagreeable impression on the gustatory nerves.

MISTURA EMETI'NÆ VOMITO'RIA, *Emetic mix'ture of emetine*. (*Emetine* 4 gr., *Orange flower water* $\mathfrak{z}$ ij, *Syrup* $\mathfrak{z}$ ss; M.) A dessert-spoonful every half hour till it acts.

MISTURA FERRI COMPOS'ITA, *Com'pound mix'ture of iron*, (F.) *Mixture de fer composé*, *Griffith's mixture*. (*Myrrh. cont.*, $\mathfrak{z}$ j, *Potassæ carb. gr. xxv*, *Aquæ rosæ* $\mathfrak{f}$ 3viiss, *Ferri sulph. in pulv.* $\mathfrak{z}$ j, *Sp. lavand.* $\mathfrak{z}$ ss, *Sacchar.* $\mathfrak{z}$ j. Rub together the myrrh, subcarbonate of potassa, and sugar; then add, while triturating, the rosewater, spirit of nutmeg; and, lastly, the sulphate of iron. Pour the mixture directly into a glass bottle, and stop it close. Ph. U. S.) It is tonic and emmenagogue, and is useful wherever iron is indicated.

MISTURA FERROSALINA, *Mixture, ferrosaline*.

MISTURA FUSCA, ('brown mixture,') *M. Glycyrrhizæ composita*.

MISTURA GLYC'YRRHI'ZÆ COMPOS'ITA, *Mistu'ra fusca*, *Com'pound mix'ture of Liquorice*, *Brown Mix'ture*, (*Extract. Glycyrrhiz. pulv.*, *Acaciæ pulv.*, *Sacchar.*, $\mathfrak{a}\mathfrak{a}$ $\mathfrak{z}$ ss; *Tinct. opii camphorat.* $\mathfrak{f}$ 3ij; *Vin. antimon.* $\mathfrak{f}$ 3j; *Spirit. æther. nitric.* $\mathfrak{f}$ 3ss; *Aquæ* $\mathfrak{f}$ 3xij. Rub the liquorice, gum arabic, and sugar with the water, gradually poured upon them; add the other ingredients and mix. Ph. U. S.) A popular cough medicine, but not deserving of being made officinal. Dose, a table-spoonful or $\mathfrak{f}$ 3ss.

MISTURA GUA'I'ACI, *Mix'ture of guai'ac*, *Lac guai'aci*, *Emul'sio guai'aci'na*, *Mistu'ra guai'aci gummosa seu gum'mi guai'aci Bergeri*, *Solutio guai'aci gummosa seu resinæ guai'aci aquosa*, (F.) *Mixture de Gayac*. (*Guaiac. gum resin.* $\mathfrak{z}$ iss, *Sacch. pur.* $\mathfrak{z}$ ij, *Muc. acaciæ gum.* $\mathfrak{z}$ ij, *Aq. cinnam.* $\mathfrak{f}$ 3viij. Rub the guaiacum with the sugar; then with the mucilage. Add, gradually, the cinnamon water. Ph. L.) Dose, $\mathfrak{f}$ 3ss to $\mathfrak{f}$ 3j.

MISTURA GUMMI GUA'IACI BERGERI, *Mistura Guai'aci*—m. Hordei, *Decoc'tum*, *Hordei compositum*—m. Leniens, *Looch ex ovo*—m. Media, see *Mistura*.

MISTURA MOS'CHI, *M. moscha'ta*, *Musk mix'ture*. (*Moschi*, *Acaciæ gum. contus.*, *Sacch. purif.*, *sing.* $\mathfrak{z}$ j, *Aq. rosæ* $\mathfrak{f}$ 3vj. Rub the musk with the sugar; add the gum, and, by degrees, the rosewater. Ph. L.) Dose, $\mathfrak{f}$ 3ss to $\mathfrak{f}$ 3ij.

MISTURA POTASSÆ CITRATIS, *Liquor Potassæ Citratiss*.

MISTURA SCAMMO'NIJ, *Scam'mony Mix'ture*. *Scammon. gr. vij*, *Lactis* $\mathfrak{f}$ 3ij. Triturate to form an emulsion. Ph. E.) The whole is a cathartic dose.

MISTURA SPIR'ITUS V'I'NI GALL'IC, *Bran'dy*

Mixture. (*Sp. Vin. Gallic.*, *Aq. cinnam.* āā f ʒiv; *Vitel. ori.* ij; *Sacchar.* ʒss; *Ol. cinnam.* ℥ ij. *M. Ph. L.*) An excitant draught for the low stages of fever.

MISTURA STRYCHNINÆ, *Mixture of Strychnia.* (*Distilled water,* ʒij, very pure *Strychnia*, 1 gr., *White sugar,* ʒij; *M.*) Dose, a dessertspoonful.

MISTURA SULPHURICO-ACIDA, *Elixir acidum Halleri*—*m. Vulneraria acida, Argemone, Eucd.*

MITCHELLIA REPENS, (after Dr. John Mitchell, of Virginia, a botanist.) *Partridge berry.* A pretty little indigenous trailing evergreen, of the *Cinchona family*—*Cinchonaceæ*,—which flowers from June to July. It has been considered an expectorant, emmenagogue, and diuretic; and has been prescribed in dropsy and gout.

MITELLA, (dim. of *μῆτρα*, *mitra*, 'a bandage,') Sling.

MITHRIDATE, *Mithridatium, Mithridatium medicamentum, Antidotum Mithridatium, Dioscoridi antidotum, Confectio Damocratis.* A very compound electuary, into which entered—*Myrrh of Arabia, Saffron, Agaric, Ginger, Cinnamon, Frankincense, Garlic, Mustard, Birthwort, Galbanum, Castor, Long pepper, Opoponax, Bdelium, Gum Arabic, Opium, Gentian, Orris, Sappanum, Valerian, Acacia, Hypericum, Canary wine, Honey, &c.* It was invented by Mithridates, king of Pontus and Bithynia, and was formerly regarded as alexipharmic. It is little used at the present day, and, from its heterogeneous nature, should be wholly abandoned.

MITHRIDATICUM MEDICAMENTUM, *Mithridate.*

MITHRIDATIUM, *Mithridate.*

MITIGANS, (*mitigare*, 'to assuage,') Lenitive.

MITRA HIPPOCRATICA, (*μῆτρα*, 'a mitre,' 'a bandage,' 'a head dress,') *Bonnet d'Hippocrate.*

MITRAL, *Mitra'lis*. Having the form of a mitre; resembling a bishop's mitre.

MITRAL REGURGITATION, see *Regurgitation*.

MITRAL VALVES, *Valvule mitra'les* seu *Cor'dis mitra'les* seu *episcopa'les* seu *bicuspidata'les* seu *bicuspidatæ*. Two triangular valves at the opening of communication between the left auricle of the heart and the corresponding ventricle. These valves are formed by the inner membrane of the left cavities of the heart; and are retained on the side of the ventricle by tendinous cords, proceeding from the columnæ carneæ. They form a species of valve, which permits the blood to pass from the auricle into the ventricle and opposes its return.

MITRIUM, (*μῆτριον*, dim of *μῆτρα*, 'a bandage,') Sling.

MITTE (F.), *Effluviū latrina'rium*. An emanation exhaled from privies, which strongly irritates the eyes. It consists of ammonia, united to the carbonic and hydrosulphuric acids.

MIUAMARU, see *Arrow poison*.

MIVA, *Marmelade*.

MIXIS, (*μιξίς*,) Coition, *Mistura*.

MIXTIO, (*miscere, mixtum*, 'to mix,') *Mistura*.

MIXTURA, *Mistura*.

MIXTURE, *Mistura*—*m. Almond, Emulsio Amygdalæ*—*m. of Ammoniac, Mistura ammoniaci*—*m. of Asafetida, Mistura asafetidæ*.

MIXTURE, BRANDY, *Mistura spiritus vini Gallici*—*m. Brown, Mistura glycyrrhizæ composita*—*m. Calmante, Looch ex ovo*.

MIXTURE, CAMPHOR, *Mistura camphoræ*—*m. Chalk, Mistura cretæ*—*m. of Citrate of Potassa, Liquor potassæ citratis*.

MIXTURE DE CRAIE, *Mistura cretæ*—*m. of Creasote, Mistura creasoti*.

MIXTURE, CRETACEOUS, *Mistura cretæ*.

MIXTURE OF EMETINE, (*emetic*,) *Mistura emetinæ vomitoria*—*m. de Fer composée, Mistura ferri composita*.

MIXTURE, FERROSALINE, *Mistura ferrosali'na*.

Under this name, the author has long prescribed, in public and in private, the following combination in torpor of the colon. *Magnes. sulphat.* ʒi, *Potassæ bitart.* ʒi, *Ferri sulph.* gr. x; for a quart of water. A wineglassful every morning, or rising, until the torpor is removed.

MIXTURE DE GAYAC, *Mistura Guaiaci*.

MIXTURE, GRIFFITH'S, *Mistura ferri composita*.

MIXTURE OF GUAIAC, *Mistura Guaiaci*—*m. of Gum Arabic, see Emulsio Acaciæ*—*m. of Harts-horn, burnt, Mistura cornuusti*.

MIXTURE, HOPE'S. A mixture recommended by a Mr. Hope in diarrhoea and dysentery. It is essentially a nitrate of morphia. Although the proposer employed the acidum nitrosus of the Edinburgh Pharmacopœia, nitric acid is generally used. (*Acid. nitros.* ʒj; *Aquæ camphor.* f ʒviij; *Tinct. opii* gtt. xl; *M.*) Dose, a fourth part every three or four hours.

MIXTURE OF IRON, COMPOUND, *Mistura ferri composita*.

MIXTURE OF LIQUORICE, COMPOUND, *Mistura glycyrrhizæ composita*—*m. Lithontriptique de Durande*, see *Liquor anodynus terebinthinatus*—*m. of Marsh mallows, Decoetum althææ*.

MIXTURE, MUSK, *Mistura moschi*—*m. Neutral, Liquor potassæ citratis*.

MIXTURE OF PRUSSIC ACID, *Mistura acidi hydrocyanici*—*m. of Scammony, Mistura scammonii*.

MIXTURE, SCUDAMORE'S. A mixture recommended by Sir C. Scudamore in gouty and rheumatic affections, and much prescribed in the United States. *Magnes. sulph.* ʒi—ʒij; *Aquæ menthæ* f ʒx; *Aceti colchic.* f ʒj—f ʒiiss; *Syrup. croci* f ʒj; *Magnes.* ʒviij, *M.* Dose, one, two, or three tablespoonfuls, repeated every two hours in a paroxysm of gout, until from four to six evacuations are produced in the twenty-four hours.

MIXTURE OF STRYCHNIA, *Mistura strychniæ*—*m. de Whitt*, see *Liquor anodynus terebinthinatus*.

MNEME, (*μνήμη*,) *Memory*.

MNEMECEPHALICUM BAL'SAMUM. A very compound medicine, into which entered *Balm, Lily, Rosemary, Lavender, Borage, Broom, Roses, Violet, Saffron, Thyme, Storax, Galbanum, &c.* Supposed to have great mnemonic virtues.

MNEMONETICE, *Mnemonics*.

MNEMONICS, *Mnemon'icæ, Mnemoneuticæ, Mnemotechnæ*, (*μνήμη*, 'memory,' *μνημονικός*, 'appertaining to memory,' from *μνᾶμαι*, 'I recollect,') The art of aiding the memory; and of forming, in some sort, an artificial memory.

MNEMOTECHNE, (*μνήμη*, 'memory,' and *τεχνή*, 'art,') *Mnemonics*.

MOAN'ING, (*Sax. mænan*, 'to grieve,' by onomatopœia,) *Respiratio luctuosæ, Mycthis'mos*. A plaintive respiration, in which the patient utters audible groans,—*moans*, (F.) *Gémissements*.

MOAP-EYE, *Myopia*.

MOBILE, see *Saint Augustine*.

MOBILITY, *Mobil'itas*, (*mobilis*, contraction of *movibilis*, 'that can be moved,' from *movere*, 'to move,' and *habilis*, 'capable,') The power of being moved. In *physiology*, great nervous susceptibility, often joined to a disposition to convulsion. Greatly developed excitability.

MOCCASIN FLOWER, *Cypripedium acaule, C. luteum*—*m. Plant, Cypripedium luteum*.

MOCHLIA, (*μολχία*, from *μολχεω*, 'to move,') Reduction of a luxated bone; (from *μολχος*, 'a lever,')

MOCHLICUS, (*μολχικός*,) Relating or appertaining to *mochlia*. Also, a purgative; (from *μολχεω*, 'to set in motion,' as with a lever.)

MOCHLISCUS, (dim. of *μολχος*, 'a lever,') Lever.

MOCHLUS, (*μολχος*,) *Lever*.

MOCHTHUS, (*μολθος*,) *Agony*.

MOCK-KNEES, *Crusta genu equinæ*.

MODELLING PROCESS. A term proposed

by Dr. Macartney, of Dublin, to signify the mode in which wounds are healed without inflammation or suppuration, by a deposit of plastic matter from the surface of the wound, by which the gap is more rapidly filled, — portion being laid upon portion, without waste, after the manner of clay in the hands of the sculptor;—hence the term.

MODIOLUS, ('the nave of a wheel.') *Pyrænia* seu *Az'is* seu *Columel'la coch'lex*. A hollow cone in the cochlea of the ear, forming a nucleus, axis, or central pillar, round which the gyri of the cochlea pass. The modiolus forms the inner and larger portion of the central pillar, and is the cavity seen at the bottom of the meatus auditorius internus. It lodges a branch of the auditory nerve. The central portion of the modiolus contains a number of minute canals, and is called in consequence *Tractus spiralis foraminulosus*. Into these the nerves of the cochlea enter, and pass out at right angles between the bony plates forming the zona ossea of the lamina spiralis, to be expanded on the membranous portion of the lamina.

MODIOLUS, *Mas*, also means the crown of the trepan.

MODIRA CANIRAM, *Strychnos colubrina*.

MO'DIUS. A dry measure, the third of an amphora, equal to the bushel.

MO'DUS OPERAN'DI, ('mode of operating.') This term is applied to the general principles upon which remedies act in morbid states of the body;—a subject of much interest, although involved in considerable obscurity.

The following classification will convey some idea of the ways in which different organs may be excited into action.

1. By actual or immediate contact of the remedy, and by absorption or mediate contact.
2. By an impulse conveyed by the nerves, through an impression made on the stomach or elsewhere, or by remote sympathy.
3. By contiguous or continuous sympathy, or by mere proximity or continuity of parts.

MOËLLE, (from *medulla*;) according to some, from *μυελος*, 'marrow,') Marrow—*m. Allongée*, Medulla oblongata—*m. Épinid'le*, Medulla spinalis—*m. Vertébrale*, Medulla spinalis.

MŒNIA DENTIUM, ('walls of the teeth,') Alveoli dentium.

MËROR, ('sorrow,') *Athymia*.

MOFETTE, Azote, see Mephitism.

MOFFAT, MINERAL WATERS OF. Moffat is a village situate about 56 miles S. W. of Edinburgh. The water is a cold sulphureous: containing, in a wine gallon, 36 grains of chloride of sodium, 5 cubic inches of carbonic acid, 4 of nitrogen, and 10 of sulphohydric acid. It resembles Harrogate water in its properties.

A strong chalybeate spring was discovered there about the year 1828, which contains sesquioxide of peroxide of iron, sulphate of alumina, and uncombined sulphuric acid.

MOGIGRAPHIA, (*μωγισ*, 'with difficulty,') and *γραφα*, 'I write,') Cramp, writers'.

MOGILALIA, *Paralalia*, (*μωγισ*, 'with difficulty,') and *λαλεω*, 'to speak,') This word is inaccurately written by Paul Zacheias,—*Mola'lia*, and *Molila'lia*. Difficult or defective articulation. Impediment of speech. See Balbuties.

MOGOSTOCIA, (*μωγος*, 'difficulty,') and *τοκος*, 'parturition,') Dystocia, Laborious labour.

MOGUS, (*μωγος*,) Agony.

MOHAWK TASSEL, *Eupatorium purpureum*.

MOHR'IA THURIF'RAGA, (after Prof. D. M. H. Mohr, a botanist of Kiel.) A South African fern, *Nat. Ord.* Filices, the dry leaves of which, pulverized and made into an ointment with fat, are serviceable in burns and scalds.

MOIGNON, (from Celt. *Moign*, 'naimed,') Stump.

MOIS, ('month,') Menses.

MOISTURE, *Ma'dor*, (F.) *Moiteur*. Simple humidity of the skin. The skin is said to be moist or in a state of moisture when there is slight perspiration.

MOÏTEUR, (old F. *moisteur*,) Moisture.

MOLA, ('a mill,') Maxillary bone, Molar tooth, Mole, Patella.

MOLAGOCODI, *Piper nigrum*.

MOLAIRE, Molar.

MOLALIA, Mogilalia.

MO'LAR, *Mola'ris*, *My'licus*, (F.) *Molaire*, *Moulière*. That which bruises or grinds; (from *mola'ris*, 'a grindstone,') or *mola*, 'a mill,')

MOLAR GLANDS. Two small bodies formed by a collection of mucous crypts; seated in the substance of the cheeks, between the masseter and buccinator muscles, and whose excretory ducts open into the mouth opposite the last molar teeth.

MOLAR TEETH, *Grind'ers*, *Jaw Teeth*, *Mola'ris*, *permanen'tes den'tes*, *Den'tes mola'res* seu *maxilla'res* seu *gom'phii* seu *clava'les*, *Mo'la*, *Gom'phioi*, *My'lacri*, *My'lodon'tes*, *My'lodi*, *Momis'ci*, *Lag'teeth*, *Wall teeth*, (Prov.) *Az'le teeth*, *Az'le teeth*, *Cheek teeth*, *Wang teeth*, (F.) *Dents molaires* ou *mâchelières*, occupy the farther part of each alveolar arch. Their coronæ are broader than they are high: unequal; tuberculated; and the roots are more or less subdivided. They are 20 in number, 10 in each jaw. The first two pairs of molar teeth in each jaw have been called *lesser molares*, *premo'lar* or *bicus'pid*, *Dentes bicuspidati* seu *bucca'les*. The coronæ have two tubercles; the outer being more prominent and larger than the inner. The root is commonly simple; sometimes bifurcated. The other three pairs have been termed, *Great molares*—*Dents multicuspidées* (Ch.). Their coronæ are cubical, rounded, and have, at the upper surface, four or five tubercles, separated by deep furrows. The root is divided into two, three, four, and five fangs, which are more or less curved. The third great molaris appears a long time after the others, and hence has been called *Dens sapient'ie* or *Wisdom tooth*.

MOLASSES, Melasses.

MOLE, *Mo'la*, (*moles*, 'mass,') *My'le*, *Zephyr'ius* *foetus*, *Ger'men fal'cum*, *Pseudocye'sis mola'ris*, *Gravid'itas mola'ris*, *Moon calf*, old French, *Fardean*. A fleshy, insensible, at times, soft—at others, hard—mass; of variable and determinate size, which forms in the uterus, and is slightly united by vessels to that organ; from which its feeble vitality is derived. It has been conceived by some to be always owing to imperfect conception; but moles may form in the undoubted virgin. They seem to be owing to a morbid process; and certainly are generally connected with conception. At times, they contain parts of the foetus; but commonly do not. At very different periods, in different women, the diseased mass is expelled from the uterus, with ordinary symptoms of abortion; and the case requires similar management. See, also, *Nævus*, (Sax. mal,) and Conception, false.

MOLE PLANT, *Euphorbia lathyris*.

MOLECULAR, *Molecula'ris*, (F.) *Moléculaire*. Of or belonging to molecules or minute portions of any thing. Hence *molecular death*, in contradistinction to *somatic death*.

MOLECULE, *Molec'ula*, *Mass'ula*: (F.) *Molé-cule*, (dim. of *moles*, 'a mass,') A minute portion of any body. Also, the *cicatri'cula*, *mac'ula*, *gallat'ura*, *gelatinous molec'ule*, tread of the cock, or embryo part of the impregnated ovum, observable by the microscope before the ovum has left the ovarium of the hen. It lies under the epidermic coats of the yolk, and upon its proper coat. If the ovum, according to Valentin, be lacerated and its contents minutely examined, the *cicatri'cula* is found like a grayish white disk, which in

its whole periphery is dense, granulous, and opaque; but in the centre presents a clear, non-granulous, and perfectly diaphanous point. Purkinje found, that when he removed the dark granulous mass by suction with a small tube, there remained a perfectly transparent vesicle filled with a pellucid lymph, which had a decidedly spherical form, but, being extremely delicate, was easily lacerated, and then its fluid escaped. As he found this, which later naturalists have named—after its discoverer—the *Purkinjean vesicle*, in the ova of the ovary, but could not see it in ova, which had already entered the oviduct, he gave it the name *germinal vesicle*, or *germ cell*, *vesicula prolifera* seu *germinativa*, *Blastocystina*, (F.) *Vesicula germinativa*. Besides a perfectly colourless fluid, this contains one or more dark corpuscles, which appear as a nucleus through the including membrane in the shape of opaque spots—the *germinal spot*, *macula germinativa* seu *germinans*, *nucleus germinativus*, *Blastocardia*, *Corriculum germinis*, *Blastocellia*, *Wagnerian spot* or *vesicle*, (F.) *Tache germinative*. The small dots, observed by Valentin in the Wagnerian vesicles and arising in them have been termed *Valentinian Vesicles*. The granulous membrane—its thickened portion, the so-called '*ciatricula*,'—and the germinal vesicle, constitute those parts of the ovum which pass immediately into the original foundation of the embryo, the *blastodermis* or *germinal membrane*, *vesicula blastodermatica* seu *blastodermica*, *blastodermic vesicle*.

In the nomenclature of Professor Agassiz, the yolk membrane is an *ectoblast*, the '*Purkinjean Vesicle*,' a *mesoblast*; the '*Wagnerian Vesicle*' an *entoblast*, and the '*Valentinian Vesicle*,' an *entostoblast*. See *Cytoblast*.

MOLECULE, GELATINOUS, Molecule.

MOLÈNE, *Verbascum nigrum*.

MOLES GANGLIFORMIS, ('gangliform mass,') Ganglion of Gasser.

MOLIBDUS, (μολιβδος,) Plumbum.

MOLILALIA, *Mogilalia*.

MOLIMEN, pl. MOLIMINA, (*moliōr*, 'to strive,') An attempt, a struggle. Hence:

MOLIMEN CRITICUM. An impulsion towards a sudden solution or crisis of a disease.

MOLIMEN HÆMORRHAGICUM. The hemorrhagic diathesis or impulsion.

MOLIMEN HÆMORRHOIDALÈ. The hemorrhoidal diathesis or impulsion. See *Hæmorrhoids*.

MOLIMEN MENSTRUALÈ. The menstrual diathesis or impulsion.

MOLIMEN NATURE SALUTARIUM, Instinct.

MOLKENKUR, *Whey cure*.

MOLLESCENCE, (*mollis*, 'soft,') Mollities.

MOLLET, (dim. of (F.) *mou*, *mol*, 'soft,') *Sura*.

MOLLIPUFFS, *Lycoperdon*.

MOLLIS MATER, ('tender mother,') *Pia mater*.

MOLLIT'IES, (from *mollis*, 'soft,') *Mollescence*, *Malacismus*, *Mollitia*, *Mollitudo*, *Emollit'ies*, *E. morbo'sa*, *Mollit'io*, *Malax'is*, *Malaco'ma*, *Malaco'sis*, *Malthac'is*, (F.) *Ramollissement*; (G.) *Malakien* of Fuchs. Preternatural softness of an organ or part of an organ.

MOLLITIES CEREBRI, *Malaco'sis Cerebri*, *Encephalomalacia*, *Encephalomalax'is*, *Encephalomalaco'sis* *Neurencephalus* (W. Farr), *Cerebromalacia*, *Encephaloseps'is*, *Mollescence* or softening or a kind of liquefaction of the cerebral substance; the remainder preserving its ordinary consistency. (F.) *Ramollissement du cerveau*. The neurine often contains small clots of blood, giving rise to what has been termed *Apoplexie capillaire*, (Cruveilhier.) The symptoms denoting it are equivocal.

MOLLITIES CORDIS, *Cardiomalacia*.

MOLLITIES MEDULLÆ SPINALIS, *Myelomala-*

cia, (F.) *Ramollissement de la Moëlle Épinrière*. Softening of the spinal marrow.

MOLLITIES OS'SIUM, *Malacosteosis*, *Malacosteon*, *Emollit'ies os'sium*, *Parostia flexilis*, *Osteomalaco'sis*, *Osteomala'cia*, *Rachit'is* seu *Rachit'is adultorum*, *Spina vento'sa* (of some), *Softening of the bones*, (F.) *Osteo-malacia*, *Osteomalacie*, *Osteo-malakie*, *Ramollissement des os*. A rare affection, in which the bones are deprived of their salts, particularly of the phosphate of lime, and consist only, or mainly, of gelatin; hence they acquire a degree of suppleness which renders them unfit for the performance of their functions. The disease generally affects all the bones; but it is especially remarkable during life in the long bones, which assume any curvature that may be wished. Very violent pain is often experienced in them; and the urine frequently contains an enormous proportion of calcareous phosphate. The patient is compelled to remain in the horizontal posture; the bones no longer being fixed points for the muscles to act upon. The disease, has, hitherto, always terminated fatally, and dissection has exhibited the gelatinous nature of the bones; which, by desiccation, have become transparent, as if they had been macerated in acid.

Experience has afforded no means of relief in this dreadful affliction. The alkaline salts, earthy phosphates, &c. are of little or no use; the disease being in the cells of nutrition.

MOLLITIO, Mollities.

MOLLITUDO, Mollities.

MOLLUS'CUM, *Atheroma*, *Ochthi'asis*, ([L.] *Molluscum*, 'a knotty growth on the bark of the maple,') A cutaneous affection, which consists of numerous tumours, varying in size from that of a pea to that of a pigeon's egg, filled with an atheromatous matter, which are developed in the substance of the derma, and are of various shapes, some having a large base,—others adherent by means of a pedicle.

Molluscum contagiosum, (F.) *Acné molluscoides*, a singular variety, is characterized by the presence of hard, round, tubercles, which are smooth and transparent, and, when pressed, pour out from an orifice in their summits a little opaque or milky fluid.

The disease is probably seated in the sebaceous follicles. If internal treatment be adopted at all, it must consist of eutrophics, as arsenic, iodine, &c. The external treatment consists in the employment of measures calculated to excite the tubercles to inflammation, as by touching them with potassa.

MOLOCHE, (μολοχη,) *Malva rotundifolia*.

MOLOPES, (μολωψ, 'a weal,') *Vibices*.

MOLYBDÆNUM MAGNESII, (μολυβδος, 'lead,') Manganese, black oxide of.

MOLYBDOCACHEXIA, (μολυβδος, 'lead,' and *cachexia*,) *Saturnismus*.

MOLYBDOCOLICA, (μολυβδος, 'lead,' and *colica*,) *Colica metallica*.

MOLYBDONOSUS, (μολυβδος, 'lead,' and *nosos*, 'disease,') *Leadpoisoning*.

MOLYBDOPARESIS, (μολυβδος, 'lead,' and *paresis*, 'paralysis,') *Palsy*, *lead*.

MOLYBDUS, (μολυβδος,) *Plumbum*.

MOLYBDOSIS, (from *molybdus*, and *osis*,) *Leadpoisoning*.

MOLYNE, (μολυνη,) *Anus*.

MOLYNIE, (μολυνι,) *Anus*.

MOLYNSIS, (μολυνσις, from *μολυνω*, 'to taint,') *Miasma*.

MOMIE, *Mummy*.

MOMIN, *Mamei*.

MOMISCI, *Molar teeth*. Also the parts of the molar teeth connected with the gums.

MOMORDICA, *M. Balsam'ina*, (*mordico*, *mordidi*, *morsum*, 'to bite,') *Balsam'ina*, *Neurosp'er-*

ma cuspidata, Balmapple, Balsam apple, (F.) *Balsamine*, *Pomme de merveille*. Ord. Cucurbitaceæ. The fruit of this oriental tree, *Pomum mirabile* seu *hierosolymitanum*, was formerly considered vulnerary.

MOMORDICA ASPERA, *M. Elaterium*.

MOMORDICA ELATERIUM: *M. aspera*, *Elaterium*, *Eboalia Elaterium*, *Eboalum officinarum* seu *agrestis* seu *cordifolium*, *Cucumis agrestis* seu *asininus* seu *sylvestris*, *Charantia*, *Guarea* or *ba*, *Eboalum elaterium*, *Wild*, *Squirtling*, or *Asses' cucumber*, (F.) *Concombre Sauvage*, *Momordique*. Nat. Order, Cucurbitaceæ. The dried sediment from the juice is the elaterium of the shops. (See *Extractum Elaterii*.) It is a most powerful cathartic, and, as such, has been used in dropsies. Its active principle is the *Elatin*, *Elaterin* or *Ecballin*. Dose, gr. $\frac{1}{2}$ to gr. j. until it operates.

MOMORDIQUE, *Momordica elaterium*.

MO'NAD, ($\mu\omega\nu\alpha\varsigma$, gen. $\mu\omega\nu\alpha\delta\omicron\varsigma$, 'a unit.'). A simple particle, or atom, or unit. The smallest of all visible animalcules. A primary cell or germ.

MONÆ, Nates.

MONARDA, (after N. Monardes, a Spanish botanist.), *M. didyma* seu *coccinea* seu *purpurea* seu *Kalmia*, *Scarlet Rose balm*, *Monntain mint*, *Ousego Tea*, *Monntain Balm*, *Horsemint*, *Square stalk*, *Red balm*; a beautiful indigenous plant, *Sex. Syst.* Diandria Monogynia. Nat. Ord. Labiata, has the excitant properties of *Monarda punctata*.

MONARDA ALLOPHYL'LA, *M. fistulosa*—m. *Clinopodia*, *M. fistulosa*—m. *Coccinea*, *Monarda*—m. *Didyma*, *Monarda*.

MONARDA FISTULO'SA, *M. rugosa* and *oblongata*; *M. clinopodia* seu *alloyhylla*, *Wild bergamot*, *Horsemint*. The leaves have a fragrant smell; and an aromatic, somewhat bitter taste. They are reputed to be nerve, stomachic, and deobstruent [?].

MONARDA KALMIANA, *Monarda*—m. *Oblongata*, *M. fistulosa*.

MONARDA PUNCT'ATA, *Horsemint*, *Monarda*, (Ph. U. S.) Indigenous in the United States. Stimulant and carminative. The *Oleum monardæ* is official in the United States.

MONARDA PURPUREA, *Monarda*—m. *Rugosa*, *M. fistulosa*.

MONDER (F.), (from [L.] *mundare*, 'to clean.'). To render clean or pure. In *Pharmacy*, it means to separate any substance from its impure or useless portion. In surgery, *monder* ou *mondifier une plaie*, is to clean or deterge a wound.

MONDIFICATIFS, ([L.] *mundus*, 'clean,' and *facere*, 'to make,') Detergents.

MONDIFIER UNE PLAIE, ('to cleanse a wound,') see *Monder*.

MONE'MERON, *Monœ'meron*, *Monohe'meron*; (*mono*, and $\mu\eta\epsilon\rho\alpha$, 'a day.'). A name given to several collyria, which were supposed to be capable of curing diseases of the eyes in a day.

MONENTERUM, (*mono*, and $\epsilon\pi\epsilon\rho\omega\nu$, 'intestine,') Colon.

MONE'SIA. A vegetable extract imported into Europe from South America, in hard, thick cakes. It is prepared from the bark of a tree, whose botanical name is uncertain—probably *Chrysophyllum glycyphæum* seu *Buranhem*, *Buranhem*, *Guaranhem*; Ord. Sapotaceæ. It is very friable, and its fracture very much resembles that of a well-torrefied cocoanut. It is wholly soluble in water; and its taste—which is at first sweet like liquorice—sometimes becomes astringent. It is on account of its astringent properties that it has been prescribed in chronic bronchitis, hæmoptysis, diarrhœa, leucorrhœa, uterine hemorrhage, &c. It has been applied locally, in the form of ointment, (*Mones. p. 1, Adipis, p. vij.*)

The dose of *Monesia* is from 12 to 40 grains a day.

MONETWORT, *Lysimachia nummularia*.

MONGO'LIAN. Anthropologists give the name *Mongolian race* to a variety of the human species, spread over a great part of the north of Asia, in China, India, Thibet, the Moluccas, &c. The individuals composing it have the skin of a brown red, forehead flat, nose broad, cheeks prominent, and lips large. See *Homo*.

MONGREL, ([Sax.] *mengan*, or *mængan*, 'to mix,') Hybrid.

MONPLIES, [Sc.] ('many plies or folds,') *Omasum*.

MO'NISM, *Monismus*, ($\mu\omega\nu\alpha\varsigma$, 'unity.'). The doctrine, which declares matter and mind to be identical.

MONK PHYSIC' IANS AND SUR'GEONS. A class of practitioners of whom Frère Cosme and Jacques Beaulieu in France, and Pravetz in Germany, were the most distinguished.

MONKSHOOD, *Aconitum*—m. Common, *Aconitum napellus*—m. Salutory, *Aconitum anthora*—m. Wild, *Aconitum uncinatum*.

MONNI'NA POLYSTA'CHIA, (after Monnino, Count de Florida Blanca, a Spanish promoter of botany.) A beautiful South American plant. Nat. Ord. Polygalæ. *Sex. Syst.* Diadelphia Octandria. The bark of the root is a powerful astringent, and much used in South America in diseases of the bowels.

MONNYPLIES, [Sc.] ('many plies or folds,') *Omasum*.

MONO, (from $\mu\omega\nu\alpha\varsigma$, 'one alone,') in composition, one alone, only one.

MONOBLEP'SIS, (*mono*, and $\beta\lambda\epsilon\psi\iota\varsigma$, 'sight.'). An affection in which vision is confused, imperfect, and indistinct, when both eyes are employed; but perfect or nearly so, when either eye is used singly.

MONOCAR'PÆ, (*mono*, and $\kappa\alpha\rho\pi\omicron\varsigma$, 'fruit.'). A division of cutaneous affections, of the family *Eczematosen*, of Fuchs, which includes *strophulus*, *psyracria*, and *eethyma*.

MONOCEPH'ALUS, ($\mu\omega\nu\omicron\kappa\epsilon\phi\alpha\lambda\omicron\varsigma$, from *mono*, and $\kappa\epsilon\phi\alpha\lambda\eta$, 'head.'). A compound monster having two bodies with a single head.

MONOCLE, *Monoculus*.

MONOCOLON, *Cæcum*, *Rectum*.

MONOCRA'NUS, (*mono*, and $\kappa\rho\alpha\nu\omicron\nu$, $\kappa\rho\alpha\nu\iota\sigma$, 'cranium.'). A monster with one cranium, but with the face in part double.

MONOCULUM, *Cæcum*.

MONOC'ULUS, *Mon'ocle*, *Monophthal'mus*, *Unioculus*, *Unoc'ulus*, (F.) *Monocle*, (*mono*, and *oculus*, 'an eye.'). A bandage employed to maintain topical applications over one of the eyes. This bandage, called by some surgeons *Simplex oculus*, (F.) *Ëil simple*, is made of a roller three or four ells long, rolled into a single ball. See *Cyclops*.

MONODIPLO'PIA, (*mono*, $\delta\iota\pi\lambda\omicron\sigma$, 'double,' and $\omega\psi$, 'eye.'). Double vision with one eye.

MONOMERON, *Monemerion*.

MONOGAS'TRIC, *Monogast'ricus*, (*mono*, and $\gamma\alpha\sigma\tau\rho$, 'stomach.'). That which has but one stomach. Man is *monogastric*; ruminating animals, on the contrary, are *polygastric*.

MONOGENY, (*mono*, and $\gamma\epsilon\nu\alpha\varsigma$, 'kind,') see *Generation*.

MONO'GRAPH, MED'ICAL, *Monograph'ia medica*, (*mono*, and $\gamma\rho\alpha\phi\eta$, 'description.'). An *ex professo* treatise on a single class of diseases, or on a single disease or medical subject.

MONOHEMERON, *Monemerion*.

MONOMACHON, *Cæcum*.

MONOMACUM, *Cæcum*.

MONOMANIA, see *Melancholy*—m. *Homicidal*, *Homicidal insanity*—m. *Incendiary*, *Pyromania*.

MONOMANIAC, *Monomaniacus*, (F.) *Monomane*, *Monomaniacque*. One who labours under monomania, or insanity chiefly on one subject.

MONOMANIE. Melancholy — *m. Homicide*, Homicidal insanity.

MONOMMATOS, (*μονοματος*, from *mono*, and *μμα*, 'eye,') Cyclops.

MONOMORIA, (*mono*, and *μωρια*, 'silliness,') see Melancholy.

MONOMPHALUS, (*mono*, and *μφαλος*, 'a navel,') A monster produced by the union of two bodies having a common umbilicus. Such a double monster is called by I. G. St. Hilaire *Monomphalian*, (F.) *Monomphalien*.

MONONEA, (*mono*, and *νοος*, 'thought,') Thought on one subject, as in monomania.

MONOPAGIA, (*mono*, and *παγιος*, 'fixed,') Clavus hystericus, Hemieria.

MONOPATHY, *Monopathia*, (*mono*, and *παθος*, 'disorder,') A state in which one organ or function is disordered. Applied to melancholy or monomania, which is said to be a *monopathic* affection.

MONOPEGIA, (*mono*, and *πηνυμι*, 'to fix,') Clavus hystericus, Hemieria.

MONOPHTHALMUS, (*mono*, and *οφθαλμος*, 'eye,') Cyclops, Monoculus.

MONOPLASTIC, *Monoplasticus*, (*mono*, and *πλασσω*, 'I form,') That which has one form. A *monoplastic element*, in histology, is one which retains its primary form.

MONOPODIA, (*mono*, and *πους*, gen. *ποδος*, 'a foot,') Symphodia.

MONOPS, (*μονωψ*, from *mono*, and *ωψ*, 'eye,') Cyclops.

MONOPSIA, (*mono*, and *οψις*, 'vision,') Cyclopia.

MONOPUS, (*mono*, and *πους*, 'a foot,') Having one foot. See Symmeles.

MONORCHIS, gen. *Monorchididis*, (*μονορχις*, from *mono*, and *ορχις*, gen. *ορχιδος*, 'testicle,') (F.) *Monorchide*. One who has only one testicle. The condition is *Monorchidia*.

MONORGANICI, see Local.

MONOSCELUS, (*μονοσκελης*,) (*mono*, and *σκελος*, 'a leg,') One legged.

MONOSITIA, (*μονοσιτια*, from *mono*, and *σιτος*, 'food, 'repast,') The habit of taking only one meal in the day.

MONOSOMIAN, (F.) *Monosomien*, (*mono*, and *σωμα*, 'a body,') An epithet applied to a double monster, which seems, in the first instance, to have had but one body.

MONOTROPA UNIFLORA, (*mono*, and *τροπος*, 'a turn, — the solitary flower turned to one side,') *Indian pipe*, *Iceplant*, *Pipe-plant*, *Nest root*, *Fit root*. Ord. Ericaceæ. An indigenous plant, whose juice, mixed with water, has been extolled by the Indians in ophthalmia. The dried root in powder has been given in epilepsy and nervous diseases.

MONS VENERIS, ('mount of Venus,') *Monticulus Veneris*, *Ephæbum*, (F.) *Mont de Venus*, *Pénit*, *Motte*, *Éminence sus-pubienne*. The more or less projecting eminence, situate at the base of the hypogastrium, above the vulva and in front of the os pubis. At the period of puberty it becomes covered with hair, formerly called *Tressoria*.

At times, *mons veneris* is applied to the corresponding part in the male.

MONSEL'S SALT, see Ferri sulphas — *m. Solution*, see Ferri sulphas.

MONSONIA OVA'TA, (after Lady Ann Monsen, aid to Lee in his Introduction to Botany,) *Keita*, of the Hotentots. A plant of the *Nat. Ord.* Geraniaceæ, which grows at the Cape of Good Hope. The root and herb are very astringent, and are used successfully in dysentery.

MON'STER, *Mon'strum*, ('anything strange,') *Portentum*, *Terror*, *Pe'lor*, *Pelo'ria*, *Pelo'rium*, *Pelo'rum*, (F.) *Monstre*. Any organized being, having an extraordinary vice of conformation, or a preternatural perversion of every part, or of certain parts only.

The following classification embraces the main varieties of malformations — *Viti'a primæ conformatio'nis*. — 1. Those in which certain parts of the normal body are absent or defective — *mon'stra deficient'ia*, constituting *Ectrog'eny* of Serres, — *Monstros'ity by defect*, (F.) *Anomalie ou Monstruosité par défaut*, *Agénésie*. 2. Those produced by fusion or coalition of organs — *coalit'io partium*, *syn'physis*. 3. Those, in which parts, united in the normal state, are separated from each other — *clefts*, *fissures*. 4. Those in which normal openings are occluded — *atre'sia*. 5. Those by excess, or in which certain parts have a disproportionate size — *mon'stra abundan'tia*, *Monstros'ity by excess*, (F.) *Monstruosité par excès*, *Hypergénésie*. 6. Those, in which one or more parts have an abnormal position — *si'tus muta'tus*. 7. Those affecting the sexual organs — *hermaph'roditism*; and to these "true malformations" Vogel adds 8. Diseases of the foetus, and abnormal states of its envelopes.

Amongst the numerous hypotheses entertained on the origin or cause of monsters, or of monstrosity — *terato'gia*, *monstros'itas*, (F.) *monstruosité*, as the state has been called — three only are worth mentioning. They have been attributed, 1. To the influence of the maternal imagination on the foetus in utero. 2. To accidental changes, experienced by the foetus at some period of its uterine existence: and 3. To a primitive defect in the germs. The second seems to be the only one that is philosophical.

Cases of *supernumerary fingers and toes*, which fall under the 5th division of the above classification, are, probably, owing to an excess of formative action in the part; and it may be a question, whether many of the cases of diplogensis are not owing to this rather than to a fusion of two germs.

As a medicolegal question, monsters, if capable of action as individuals, have the same rights as other persons.

MONSTERS, COMPOUND, see *Unitaire* — *m. Double*, see *Diplogensis* — *m. Triplet*, see *Triplet*.

MONSTRA ABUNDANTIA, see *Monster* — *m. Deficientia*, see *Monster* — *m. Trigemina*, see *Triplet* — *m. Triplica*, see *Triplet*.

MONSTRE, *Monster* — *m. Composé*, see *Unitaire* — *m. Unitaire*, see *Unitaire*.

MONSTROSITAS, see *Monster*.

MONSTROSITY, see *Monster* — *m. by Defect*, see *Monster* — *m. by Excess*, see *Monster* — *m. by Inclusion*, see *Emboitement*.

MONSTRUM, *Monster*.

MONSTRUOSITÉ, see *Monster* — *m. par Défaut*, see *Monster* — *m. par Excès*, see *Monster* — *m. par Inclusion*, see *Emboitement*.

MONT DE VENUS, *Mons Veneris*.

MONTBRI'SON, MIN'ERAL WATERS OF. Montbri'son is a town in France, in the department of the Loire, fifteen leagues from Lyons. There are three cold springs which are highly aculeous, and used as refrigerants and aperients.

MONT-DE-MAR'SAN, MIN'ERAL WATERS OF. Mont-de-Marsan is a small French town, 10 leagues from Dax: where there is a chalybeate spring, somewhat frequented.

MONT-D'OR, MIN'ERAL WATERS OF. Mont-d'Or is a village seven leagues from Clermont, (Puy-de-Dôme,) at which are several mineral springs, cold and thermal. Those of the Fountain *La Madelaine* and of the *Great Bath*, are thermal, temp. 112° Fahr., and contain carbonic

acid, subcarbonate and sulphate of soda, chloride of sodium, subcarbonate of lime and magnesia, alumina and oxide of iron. Those of *St. Marguerite* are cold, and contain much carbonic acid. Season, from the 15th of June to the 14th of September.

MONTECATINI, MINERAL WATERS OF. This town is situate near Borgo-Buggiano, in Tuscany. The springs are thermal; ranging from 78° to 118° Fahrenheit.

MONTHLY COURSES, Menses — *m.* Illness, see *Menses* — *m.* Periods, *Menses*.

MONTICULUS CEREBELLI, (dim. of *mons*, *gen. montis*, 'a mountain,') see Vermiform processes of the Cerebellum — *m.* Veneris, *Mons veneris*.

MONTPELLIER, CLIMATE OF. This seat of a celebrated medical school in the S. E. of France was at one time renowned for its climate, and especially in regard to its adaptation to consumptive cases. This is now exploded, and it is at present esteemed an unfit residence for them. The climate possesses the general characters of that of southeastern France—being dry, hot, and irritating; subject to frequent vicissitudes, and especially to blasts of keen, cold, northerly winds. It is, indeed, most injurious in pulmonary diseases.

MOON, Lu'na, Sele'nē, Me'nē, (Sax. *Mona*, [G.] *Mond*.) (F.) *Lune*. The moon has been supposed to exert considerable influence over the human body, in health and disease. Such influence has been grossly exaggerated. Not many years ago, it would have been heretical to doubt the exacerbation of mania at the full of the moon; yet it is now satisfactorily shown, that if the light be excluded at this period, the *lunatic* is not excited more than ordinarily.

MOONCALF, Mole.

MOONED, Lunatic.

MOONSTROKE, see *Coup de soleil*.

MOONSTRUCK, Lunatic.

MOORBERRY, Vaccinium oxycoccos.

MOORGRASS, Potentilla anserina.

MOORSHEAD, Alembic.

MOORWORT, BROADLEAVED, Andromeda mariana.

MOOSEWOOD, Dirca palustris.

MOPE-EYE, Myopia.

MOPHETA, see *Mephitism*.

MOPSICAL, Myopic.

MORBEUX, Morbid.

MORBI, pl. of *Morbus* — *m.* *Acquisiti*, Acquired diseases — *m.* *Acuti*, see *Acute* — *m.* *Adventitii*, Acquired diseases — *m.* *Anniversarii*, Annual diseases — *m.* *Annu*, Annual diseases — *m.* *Asthenici*, Adynamia — *m.* *Celati*, Feigned diseases — *m.* *Chronici*, Chronic diseases — *m.* *Cognati*, Innate diseases — *m.* *Complicati*, see *Complication*, *Impliciti morbi* — *m.* *Congeniti*, Innate diseases — *m.* *Connati*, Connate diseases — *m.* *Connexi*, *Conjuncti morbi* — *m.* *Consequentes*, *Conjuncti morbi* — *m.* *Constrictorii*, see *Spasmodicus* — *m.* *Dermatodes*, see *Dermatoid* — *m.* *Dispersi*, Sporadic diseases — *m.* *Dissimulati*, Feigned diseases — *m.* *Epicteti*, Acquired diseases — *m.* *Evolutionis*, see *Development* — *m.* *Hereditarii*, Hereditary diseases — *m.* *Homœdri*, see *Homœdus* — *m.* *Infiltrati*, Feigned diseases — *m.* *Interni*, Internal diseases — *m.* *Melancholici*, see *Mercurial* — *m.* *Mercuriales*, see *Mercurial* — *m.* *Nervosi*, Nervous diseases — *m.* *Occulti*, see *Latent* — *m.* *Olfactūs*, *Osmonosi* — *m.* *Perplexi*, see *Complication* — *m.* *Pseudalei*, Feigned diseases — *m.* *Recidiva*, *Relapse* — *m.* *Rekursus*, *Palindromia* — *m.* *Sexuales*, see *Sexual* — *m.* *Sexūs*, *Genonosi* — *m.* *Simulati*, Feigned diseases — *m.* *Sparsi*, Sporadic diseases — *m.* *Spasmodici*, see *Spasmodicus* — *m.* *Subacute*,

see *Acute* — *m.* *Tactūs*, *Haphonosi* — *m.* *Tropici*, *Troponidi*.

MORBID, Mor'bidus, Morbo'sus, Path'ic, Path'ical, Path'icus, (F.) *Morbeux, Morbide*, (from *morbus*.) Diseased, or relating to disease.

MORBID ANATOMY, see *Anatomy*.

MORBIF'IC, Morbif'icus, (F.) *Morbifique*, (*morbus*, and *facere*, 'to make,') Causing or producing disease—as *morbific emanations*.

MORBILITY, Disease.

MORBILLEUX, Morbillous.

MORBILLI, (dim. of *morbus*.) *Rubeola* — *m.* *Confluentes*, *Scarlatina* — *m.* *Ignē*, *Scarlatina*. **MORBILLO'SUS, Morbillous.**

MORBIL'LOUS, Morbillo'sus, Measly, (F.) *Morbilleux*, (from *morbilli*.) Affected or connected with measles.

MORBOSUS, (from *morbus*.) Sickly.

MORBULENT, (from *morbus*.) Sickly.

MORBUS, Disease, see *Morbi* — *m.* *Ampullaceus*, *Pemphigus* — *m.* *Anglicus*, *Phthisis*, *Rachitis* — *m.* *Aphrodisius*, *Syphilis* — *m.* *Arcuatus*, *Icterus* — *m.* *Arietis*, *Influenza* — *m.* *Arquatus*, *Icterus* — *m.* *Articularis*, *Gout* — *m.* *Astralis*, *Epilepsy* — *m.* *Attonitus*, *Apoplexy* — *m.* *a. Celsi*, *Catalepsy* — *m.* *Brightii*, *Kidney*, *Bright's disease* of the — *m.* *Bullosus*, *Pemphigus* — *m.* *Caducus*, *Epilepsy* — *m.* *Ceruleus*, *Cyanopathy* — *m.* *Calculi*, *Lithia* — *m.* *Cardiacus*, *Cardialgia* — *m.* *Catarrhalis*, *Influenza* — *m.* *Catoxys*, see *Catoxys* — *m.* *Cerealis*, *Ergotism* — *m.* *Cerebralis* *Whyttii*, *Hydrocephalus Internus* — *m.* *Cirrhorum*, *Plica* — *m.* *Cœliacus*, *Cœlia flux* — *m.* *Comitialis*, *Epilepsy* — *m.* *Cordis*, *Cardionosus* — *m.* *Costalis*, *Pleuritis* — *m.* *Coxarius*, *Coxarum morbus* — *m.* *Croatus*, *Scherlievo* — *m.* *Cucullaris*, *Pertussis* — *m.* *Cucullus*, *Pertussis* — *m.* *Cystophlegmaticus*, *Cystirrhœa* — *m.* *Dæmoniæus*, *Epilepsy* — *m.* *Dæmonius*, *Epilepsy* — *m.* *Deificus*, *Epilepsy* — *m.* *Dissolutus*, *Dysentery* — *m.* *Divinus*, *Epilepsy* — *m.* *Dominorum*, *Gout* — *m.* *Endemius*, *Endemie* — *m.* *Epidemicus*, *Epidemy* — *m.* *Epidemicus gutturis* *Foresti*, *Scarlatina* — *m.* *Eruditorum*, *Hypochondriasis* — *m.* *Extrinsecus*, *External disease* — *m.* *Farcinosus*, see *Equinia* — *m.* *Febrilis*, *Pyreticosis* — *m.* *Fellidus*, *Cholera* — *m.* *Flatulentus*, *Hypochondriasis* — *m.* *Flatuosus*, *Hypochondriasis* — *m.* *Fluminienis*, *Scherlievo* — *m.* *Fœdus*, *Epilepsy* — *m.* *Frigidus*, *Atrophy* — *m.* *Gallicus*, *Syphilis* — *m.* *Gesticulatorius*, *Chorea* — *m.* *Granulosus* *renum*, *Kidney*, *Bright's disease* of the — *m.* *Gravis*, see *Serious* — *m.* *Hæmorrhoidalis*, *Hæmorrhoids* — *m.* *Heracleus*, *Epilepsy* — *m.* *Herculeus*, *Epilepsy* — *m.* *Hispanicus*, *Syphilis* — *m.* *Hydrophobicus*, *Hydrophobia* — *m.* *Hypochondriacus*, *Hypochondriasis* — *m.* *Hystericus*, *Hysteria* — *m.* *Idiopathicus*, *Idiopathia* — *m.* *Indicus*, *Syphilis* — *m.* *Interlunus*, *Epilepsy* — *m.* *Ischiadicus*, *Neuralgia femoro poplitea* — *m.* *Italicus*, *Syphilis* — *m.* *Jecinatoris* *Hepatitis* — *m.* *Lascivus*, *Chorea* — *m.* *Lateralis*, *Pleuritis* — *m.* *Lazari*, *Elephantiasis* — *m.* *Leseeli*, *Icterus* — *m.* *Lunaticus*, *Epilepsy* — *m.* *Luteolus*, *Icterus* — *m.* *Maculosus hæmorrhagicus* *Werlhofii*, *Purpura hæmorrhagica* — *m.* *Magnus*, *Epilepsy* — *m.* *Major*, *Epilepsy* — *m.* *Mensalis*, *Epilepsy* — *m.* *Mercurialis*, *Eczema mercuriale*, *Hydrargyriasis* — *m.* *Metallicus*, *Colic*, *metallie* — *m.* *Mirachialis*, *Hypochondriasis* — *m.* *Mucosus*, *Fever*, *adenomeningeal* — *m.* *Nauticus*, *Nausea marina* — *m.* *Neapolitanus*, *Syphilis* — *m.* *Niger*, *Melena* — *m.* *Œsophagi*, *Œsophagopathia* — *m.* *Originalis*, *Idiopathia* — *m.* *Oryzeus*, *Cholera* — *m.* *Pallidus*, *Chlorosis* — *m.* *Pancœnus*, *Pandemie* — *m.* *Pandemius*, *Pandemie* — *m.* *Parthenius*, *Chlorosis* — *m.* *Pedicularis*, *Phtheiriæsis* — *m.* *Percutus*, see *Catoxys* — *m.* *Phœnicus*, *Elephantiasis* — *m.* *Pilaris*, *Trichiasis* — *m.* *Pleuriticus*, *Pleuritis* — *m.* *Plumbeus*, *Lead-poisoning* — *m.* *Popularis*, *Epilepsy* — *m.* *Porcinus*, *Essera* — *m.* *Primarius*, *Idiopathia* — *m.* *Proprius*, *Idiopathia* — *m.* *Proteiformis*, see

Proteiformis—*m. Protopathicus, Idiopathia*—*m. Psodæicus, Lumbar abscess*—*m. Publicus, Epidemy*—*m. Puerilis, Epilepsy*—*m. Pulicularis, Typhus gravior*—*m. Pustulosus Finnicus*, see **Anthrax**—*m. Recidivus, Palindromia, Relapse*—*m. Regionalis, Endemic*—*m. Regius, Icterus*—*m. Resiccatorius, Hypochondriasis*—*m. Ruetuosus, Hypochondriasis*—*m. Ruetuosus*, see **Eruetation**—*m. Sacer, Epilepsy*—*m. Saltatorius, Choreia*—*m. Sancti Joannis, Epilepsy*—*m. Sancti Valentini, Epilepsy*—*m. Scarlatinus, Scarlatina*—*m. Seclustus, Epilepsy*—*m. Scrophulosus, Scrofula*—*m. Secundarius, Deuteropathia*—*m. Seleniacus, Epilepsy*—*m. Sideratus, Epilepsy*—*m. Silesiacus, Raphania*—*m. Simplex, Haplopathia*—*m. Solstitialis, Coup-de-Soleil*—*m. Sonticus, Epilepsy*—*m. Spasmodicus malignus seu popularis, Raphania*—*m. Strangulatorius, Cynanche trachealis*—*m. Sudatorius, Sudor Anglicus*—*m. Suffocans, Cynanche maligna*—*m. Suffocativus, Cynanche maligna*—*m. Truculentus infans, Cynanche trachealis*—*m. Tuberculosus*, see **Tubercle** and **Tubercular cachexia**—*m. Tuberculosis pedis, Podalcoma*—*m. Ungaricus, Fever, Hungaric*—*m. Verminosus, Helminthiasis*—*m. Vernaculus, Endemic*—*m. Vervecinus, Influenza*—*m. Vesicularis, Pemphigus*—*m. Virgineus, Chlorosis*—*m. Viridellus, Epilepsy*—*m. Vitriolatus, Epilepsy*—*m. Vocis, Phononcus*—*m. Vulpis, Alopecia*.

MORCEAU DU DIABLE, ('devil's bit,' from [L.] *mordere, morsum*, 'to bite,') see **Tube, Fallopi**—*m. Frangé*, see **Tube, Fallopi**.

MORCELLEMENT (F.), (from [L.] *mordere, morsum*, 'to bite,') 'Dividing into pieces.' A mode adopted by M. Maisonneuve for removing successfully a large fibrous tumour of the neck.

MORCHEL/LA ESCULEN'TA, (from *Morchel*, the German name,) *Phallus esculentus*, *Helvella esculenta*, *Fungus fagino'sus*, *Mercutius*, *Com'mon Morell*, (F.) *Morille*. *Nat. Order, Fungi*. A fungus, which is esteemed a luxury, entering into ragouts and other dishes, or cooked itself between slices of bacon. It has been considered to be aphrodisiac.

MORDEX, gen. *Mor'dicis*, (from *mordere*, 'to bite,') **Tooth**.

MOR'DICANT, *Mor'dicans*, *Ca'lor mor'dicans*, (*mordeo*, 'I bite,') A morbid heat, causing a disagreeable pungent sensation in the fingers of the physician.

MORDICES, (pl. of *mordex*,) **Teeth**: see **Tooth**.

MOREA (*μωρεα*,) **Morus nigra**.

MORELL, **COMMON**, *Morchella esculenta*.

MORELLE, *Boletus esculentus*—*m. à Fruit noir*, *Solanum*—*m. Furieuse*, *Atropa belladonna*—*m. en Grappes*, *Phytolacca decandra*—*m. Grim-pante*, *Solanum dulcamara*.

MORETARIUM, **Mortar**.

MORETUM. A kind of salad, with the ancient Romans, made of garlic, parsley, vinegar, oil, &c.

MORETUS, (*morum*, 'the mulberry,') Ancient name of a cordial julep, into the composition of which the syrup of mulberries entered.

MORGAGN'I, HU'MOUR OF, Hu'mor Morgagnianus, (F.) *Humeur de Morgagni*. A peculiar, transparent, slightly viscid fluid, found between the crystalline and its capsule.

MORGELENE, (from [L.] *morsus gallinae*, 'hen's bite' [?]), *Alsine media*.

MORGUE (F.). A *deathhouse*, in France, wherein persons, found dead, are exposed with the view of being recognized by their friends—*Neerocomium*, *Neerobutichium*, *Neerostoa*.

MORI SUCCUS, see **Morus nigra**.

MORIA, (*μωρια*,) **Idiotism**—*m. Demens*, **Idiotism**, **Dementia**—*m. Demens lerema*, see **Dementia**—*m. Imbecilis amnesia*, **Amnesia**.

MORIBUND, (from *moriors*, 'to die,') **Psychorages**.

MORILLE, *Morchella esculenta*.

MORINGA APTERA, (*Moringa*, in Malabar,) see **Guilandina moringa**—*m. Nux*, **Ben**, **Guilandina moringa**—*m. Oleifera*, **Guilandina moringa**—*m. Pterygosperma*, **Guilandina moringa**—*m. Zeylanica*, **Guilandina moringa**.

MORION, (*μοριον*, 'a part, a member,') **Genital organ, Penis**.

MORIOPLAS'TICĒ, *Chirur'gia curto'rum seu anaplastica*, *Restitutio organica seu partium deperdita'rum*, *Transplantatio*, *Autoplas'ticē*, *Chirur'gia plas'tica*, *Anaplas'tic Surg'ery*, *Autoplas'ty*, *Autoplas'tic or Plastic Surg'ery*, (*μοριον*, 'a part,' and *πλαστικός*, 'forming,') The restoration of lost parts. The operations for this purpose have various names, according to the part concerned, as *Cheiloplasticæ*, *Ceratoplasticæ*, *Rhinoplasticæ*, &c.

MORNING GLORY, BLUE, *Convolvulus nil*.

MO'RO, (*morum*, 'a mulberry,') **Morum**, (F.) *Mûre*. A small abscess resembling a mulberry. A small tumour, of a similar appearance, particularly on the genital organs after impure coition.

MOROCUTES, **Morochthus**.

MOROC'THUS, *Moroch'tus*, *Moroch'i'tes*, *Moroch'i'tes*, *Moroc'tes*, *Galax'ia*, *Leucograph'ia*, (F.) *Pierre au lait*, *Milkstone*. An Egyptian stone, used by the ancients as an application to the intertrigo of children: probably a variety of fuller's earth, inasmuch as it appears to have been used by fullers.

MOROC'THUS, **Morochthus**.

MOROCOMIUM, (*μωρια*, 'folly, idiocy,' and *κομω*, 'to care for,') **Morotrophium**.

MORODOCHIUM, (*μωρια*, 'folly, idiocy,' and *δεχομαι*, 'to receive,') **Morotrophium**.

MOROSIS, (*μωρωσις*,) **Fatuitas, Idiotism**.

MOROSITA'TES, (*morosus*, 'morose,' 'peevish,' 'fretful,' from *μωρια* and *μωρωσις*, 'stupidity,')

A generic name given by Linnæus to an order of Vesania, in which he included pica, bulimia, polydipsia, antipathia, nostalgia, panophobia, satyriasis, nymphomania, tarentismus, and hydrophobia.

MOROTROPHIUM, *Morocomi'um*, *Morodochi'um*, *Manicocomi'um*, (*μωρος*, 'fatuous,' and *τροφη*, 'support,') An insane establishment, *lunatic asylum*, *mad'house*, *Bed'lam*, (F.) *Hôpital des fous*, *Maison petite*.

MORPHÆA, **Morphew**—*m. Alba*, **Lepra al-phoides**—*m. Nigra*, **Lepra nigricans**.

MORPHEE, **Morphew**.

MORPHEW, (low L.) *Morphæ'a*, (F.) *Morphée*. A term vaguely applied to scurfy eruptions on the face. *Morphæa* was formerly applied to squamous diseases in general.

MORPHIA, (*μωρεος*, *Morpheus*, the 'god of sleep,') **Morphina**—*m. Acetate of*, **Morphinæ acetat**—*m. Citrate of*, see **Morphinæ sulphas**—*m. Lactate of*, see **Morphinæ sulphas**—*m. Muriate of*, see **Morphinæ sulphas**—*m. Sulphate of*, **Morphinæ sulphas**—*m. and Zinc, double iodide of*, **Zinc and Morphina, double iodide of**.

MORPHIÆ ACETAS, **Morphinæ acetat**—*m. Citras*, see **Morphinæ sulphas**—*m. Hydrochloras*, **Morphinæ murias**—*m. Murias*, see **Morphinæ sulphas**—*m. Sulphas*, **Morphinæ sulphas**.

MORPHIC, **Morphique**.

MORPHIATED, *Morphia'tus*, (F.) *Morphiné*. Impregnated with morphia or with one of its salts.

MOR'PHICA, (see *morpho*,) **Monstrosities of birth**. The 3d order, class *Typhica* of Good.

MORPHI'NA, *Mor'phia*, **Morphium**, *Morphi'nē*, **Morphine**. A solid, colourless alkali; crystallizable in beautiful pyramids, truncated and transparent; soluble in alcohol, and slightly so in boiling water. It exists, combined with meconic acid, in opium. It may be obtained by decomposing, by means of ammonia or calcined

magnesia, an aqueous solution of opium made in the cold. The following is the process of the Pharm. U. S. *Opium*, sliced, ℥j; *Distilled water*, Alcohol, *Animal charcoal*, each a sufficient quantity; *Solution of ammonia*, f ʒvj. Macerate the opium with four pints of distilled water for 24 hours, and having worked it with the hand, digest for 24 hours, and strain. Macerate the residue twice successively with distilled water, and strain. Mix the infusions: evaporate to six pints, and filter; then add first five pints of alcohol, and afterwards three fluidounces of the solution of ammonia, previously mixed with half a pint of alcohol. After 24 hours, pour in the remainder of the solution of ammonia, mixed as before with half a pint of alcohol; and set the liquor aside for 24 hours, that crystals may form. To purify these, boil them with two pints of alcohol till they are dissolved; filter the solution, while hot, through animal charcoal, and set it aside to crystallize.

Morphia dissolves perfectly in the acids, which it saturates; and with which it forms very crystallizable salts, with the exception of the acetate. All these salts have a bitter taste, and act upon the animal economy in the same manner as opium, but more powerfully. The acetate is the salt commonly employed in medicine. It was, at one time, supposed that Morphia is the purely sedative part of opium, divested of its irritating properties; but experience has not confirmed this. On the contrary, it will generally disagree where opium does. Onesixth of a grain has been considered equal to a grain of opium.

MORPHINE ACETAS, *Morphine Acetas*, *Morphium aceticum*, *Acetate of Morphine*, *Acetate of Morphia*. A salt formed by saturating morphia with acetic acid. (*Morphia*, in powder, freed from narcotina by boiling with sulphuric ether, ʒj; *Distilled water*, Oss; *Acetic acid*, a sufficient quantity. Mix the morphia with the water; then carefully drop in the acid, constantly stirring until the morphia is saturated and dissolved. Evaporate the solution in a waterbath to the consistence of syrup. Lastly, dry the acetate with a gentle heat and rub it into powder.—Ph. U. S.) Dose, from ʒth of a grain to half a grain.

MORPHINE SULPHAS, *Morphine Sulphas*, *Sulphate of Morphia*. A salt, formed like the sulphate, by saturating morphia with sulphuric acid, evaporating and crystallizing. Dose, the same as of the acetate.

Lactate, *Muriate*, and *Citrate of Morphia*, *Morphine citras*, have likewise been used; but they possess little or no advantages over the other salts. The sulphate and muriate are official in the Ph. U. S. The muriate—*Morphine Muriat*, *M. Hydrochloras*, (Ph. B., 1864.)—is directed to be prepared as follows. (*Morphia*, in pulv., ʒj; *Aque destillate*, Oss; *Acid. muriat*, q. s. Mix the morphia with the water; carefully drop in the acid, till the morphia is saturated and dissolved. Evaporate and crystallize.) Dose from ʒth of a grain to half a grain.

MORPHINE, Morphiated.

MORPHIQUE, *Morphicus*, *Morphic*, (from *morphia*. Applied in pharmacy to any composition of morphia.

MORPHIUM, *Morphina*—m. *Aceticum*, *Morphina acetas*.

MORPHO, (*μορφη*, 'shape or form,') in composition, the same.

MORPHOLOGICAL, *Morphologicus*, (*morpho*, and *logos*, 'a description,') That which has relation to the anatomical conformation of parts. Applied, at times, to the alterations in the form of the several parts of the embryo, in contradistinction to *histological*, which is applied to the

transformation by which the tissues are gradually generated. In comparative anatomy, it is applied to the history of the modifications of forms, which the same organ undergoes in different animals, and, more generally, to the doctrine of the forms of organized bodies.

MORPHOLOGY, Anatomy, see Morphological.

MORPHOLYSIS, (*morpho*, and *lysis*, 'solution,') Destruction of organization. An agent that occasions such destruction is called a *morpholytic*.

MORPHOLYTIC, see Morpholysis.

MORPHONOMY, *Morphonomia*, (*morpho*, and *nomos*, 'a law,') The laws of organic formation. The department of anatomical science which teaches the laws of organic configuration.

MORPHOTIC, *Morphoticus*, (*μορφη*, 'shape,') A term applied to organic or anatomical elements in contradistinction to the inorganic.

MORPHOTOMY, (*morpho*, and *τεμνω*, 'to cut,') see Anatomy.

MORPIO, see Pediculus.

MORPION, Crab-louse, see Pediculus.

MORRHUA AMERICANA, see *Oleum jecoris aselli*.

MORS, gen. *Mortis*, Death—m. Apparens, Asphyxia—m. Putativa, Asphyxia—m. Repentina, Demotivus lapsus.

MORSELLUS; (dim. of *morsus*, 'a bite,') Tabella.

MORSULI ANTIMONIALES, (pl. of *morsulus*,) *Tabellæ antimoniales Kunckelii*—m. *Stibii*, *Tabellæ antimoniales Kunckelii*.

MORSULUS, (dim. of *morsus*, 'a bite,') Tabella.

MORSURE, Degmos—m. *du Diable* ('devil's bite'), see Tube, Fallopien.

MORSUS, gen. *Morsus*, (*mordeo*, *morsum*, 'to bite,') Degmos—m. Diaboli, Scabiosa succisa, see Tube, Fallopien—m. Gallinae, Alsine succisa—m. Stomachi, Cardialgia—m. Ventriculi, Cardialgia.

MORT, Adeps.

MORT, (from *mors*, gen. *mortis*, 'death,') Dead, Death—m. Apparente, Asphyxia—m. *du Chien*, Cholera—m. *aux Chiens*, Colicium autumnale—m. *du Diable*, Scabiosa succisa—m. *Né*, Still-born—m. *aux Panthères*, *Doronium pardalianches*.

MORTA, Pemphigus.

MORTAL, *Mortalis*, (from *mors*,) *Lethalis*, *Letales*, *Thanatodes*, *Thanasimus*, (F.) *Mortel*. That which is subject to death;—that which causes death:—as, *man is mortal*; a disease is mortal. Of old, it was the custom to have a division of mortal or deadly wounds, which gave rise to many errors in medico-legal investigations, as the mortality, *Lethalitas*, *Letalitas*, (F.) *Léthalité*, of wounds depends upon various circumstances; and it is often a matter of extreme difficulty to pronounce whether or not a wound is necessarily mortal.

MORTALITY, *Mortalitas*, (F.) *Mortalité*. This word, taken in an extended sense, expresses the condition of all organized bodies,—of being subject to the cessation of life. In the sense in which it is most frequently employed, it signifies, the death rate,—the proportional quantity of individuals who, in a certain population, die in a given time. If we assume the population of the earth to be one thousand millions, and a generation to last thirty-three years; in that space of time, the one thousand millions must all die, and, consequently, the number of deaths will be, by approximation,

Each year.....	30,000,000
Each day	82,109
Each hour	3,421
Each minute.....	57
Each second	1 nearly.

If, on the other hand, as has been supposed, the number of deaths is to that of the births as TEN to TWELVE: there will be born,

Each year.....	36,000,000
Each day.....	98,356
Each hour.....	4,098
Each minute.....	68
Each second.....	1 & 2-15ths.

It would not be proper to regard the *value of life* in different countries, or in different periods in the same country, to be indicated by the average mortality; inasmuch as in one case a greater mortality may occur amongst children, and in another amongst adults.

MORTALITY, BILLS OF, *Bills of Necrology, Mortuary Registers, (F.) Tables de Mortalité.* Tables, instituted for the purpose of exhibiting the number of deaths, &c., in a given time. Well-kept bills of mortality are of great use to the physician and political economist. Those of London were proverbial for their inaccuracy and insufficiency; especially as regards the complaints of which the persons had died. They were formerly made out by the parish clerks, and the information was conveyed to them by two old women, who were appointed in each parish, and were called *Searchers*;—their duty being to see that the deceased had died a natural death. This defective system has, however, been abolished; and the Registrar-General's annual reports of births, deaths, and marriages, exhibit how admirably vital statistics are now collected in Great Britain.

MORTAR, *Mortarium, Moretarium, Mortarium, P'so, Ac'onē, P'la, Hol'mus, Ig'dē, Ig'dus, (F.) Mortier.* A vessel for reducing to powder different solid substances, and for making certain mixtures. Various substances are employed in the construction of mortars;—iron, marble, glass, wedgewood ware, &c.

MORTARIOLUM, (dim. of *mortarium*, 'a mortar,' or cavity resembling a mortar.) Alveolus.

MORTARIUM, Mortar.

MORTELL, Mortal.

MORTIER, Mortar.

MORTIFER, (*mors*, gen. *mortis*, 'death,' and *fero*, 'I carry.') Lethiferous.

MORTIFICATIO, (*mors*, gen. *mortis*, 'death,' and *fio*, 'I become.') Mortification—*m. Pulmonum, Neoropneumonia.*

MORTIFICATION, *Mortificatio, Necrosis, Sphacelatio.* The loss of vitality in a part of the body. The incipient stage of mortification, when the case is still recoverable, is called *Gangrene*; when totally dead, *Sphacelus*. Mortification of a bone is called *Necrosis*.

MORTIFICATION, COLD, Sphacelus—m. Ergot, Ergotism—m. Hot, Gangrene—m. Mildew, Ergotism.

MORTOOTH, Cupri sulphas.

MORTUARY REGISTERS, Mortality, bills of.

MORTUUS, (from *mors*,) Dead.

MORULA, (from *morum*, 'a mulberry.') But-ton scurvy.

MORUM (*морум*), see *Morus nigra—m. Palpebræ internæ, Trachoma carunculosum.*

MORUS NIGRA, *M. lacinia'ta, Mul'berry Tree, Mor'ea, Sycami'nus, Sycamin'ea. (F.) Mûrier Noir. Ord. Urticacæ.* The fruit of the mulberry tree, *Morum, Sycami'num, (F.) Mûre*, is pleasant, sweet, subacid, and abounds with a deep violet-coloured juice, *Mo'ri suc'cus* (Ph. L.). A syrup is directed in the London Pharmacopœia. The bark of the root has been regarded as an antelmintic.

MORUS RU'BRA, Red Mul'berry, Wild Mul'

berry; indigenous. The root has been used to destroy tania.

MORVE, (from *morbus*, 'a disease.') Mucus, Nasal mucus; see *Equinia—m. Farcinæus, see Equinia.*

MOS, gen. *Mo'ris*, Habit.

MOS MORBI BENIG'NUS seu MALIG'NUS. The benign, or malignant character of a disease.

MOSCHARDI'NA, *Muscer'da, (μοσχος, 'musk.')* A lozenge for the mouth, composed of musk, ambergris, and other aromatics.

MOSCHELÆ'ON, (*μοσχος*, 'musk,' and *ελαιον*, 'oil.') *Moscola'a.* A compound aromatic oil, containing musk.

MOSCHUS, (*μοσχος*,) Musk—*m. Arabum, Hibiscus abelmoschus—m. Facitius, Musk, artificial.*

MOSCOLÆA, Moschelæon.

MOSLICH, Maslach.

MOSQUITA, see Mosquito.

MOSQUITO (S.), (dim. of [S.] *mosca*, 'a fly.') *Musquito, Cu'lex mosqui'to.* A very troublesome insect in warm, moist situations. Curtains, called *Mosquito curtains or bars*, are used by way of protection. The entrance of mosquitos into the bed-chamber may also be prevented, to a certain extent, by keeping the doors and windows closed until the lights are extinguished. The pain and itching from the bites are alleviated by washing them with hartshorn.

MOSQUITA is a name given, by the Portuguese, to small, red pimples on the skin, resembling the bites of the mosquito.

MOSS BERRY, *Vaccinium oxycoccus—m. Ceylon, Fucus amylaceus—m. Club, Lycopodium—m. Cup, Lichen pyxidatus—m. Edible, Fucus amylaceus—m. Haircap, Polytrichum Juniperinum—m. Iceland, Lichen Islandicus—m. Irish, Fucus crispus—m. Jaffna, Fucus amylaceus—m. Marine, Fucus amylaceus—m. Pearl, Chondrus crispus—m. Stone, Tripe de roche—m. Wall, common yellow, Parmelia parietina.*

MOTAMEN, (*movere, motum*, 'to move.') Motion.

MOTATIO, Motion.

MOTE (*μωτη*), Carbasus, Lintum.

MOTEUR, Motory.

MOTHER, ([Sax.] *moðon*,) Hysteria, Phlegm.

MOTHER'S MARKS, Nævus—m. Spots, Nævus.

MOTHERWORT, *Leonurus cardiaca, Matricaria.*

MOTIL'ITY, *Motil'itas*, (from *movere, motum*, 'to move.') (F.) *Motilité, Motricité.* Faculty of moving; moving power; contractility.

MOTION, *Mo'tus, Mo'tio, Mota'tio, Mota'men, Commo'tio, Permo'tio, Cine'sis, Movement, (F.) Mouvement.* The act of changing place. The various motions may be divided into, — *First*, the *voluntary*, or those that are executed under the influence of the brain. *Secondly*, the *involuntary*, which may be subdivided into, 1. Those of the *reflex* kind, as the closure of the larynx on the contact of acrid vapours, of the pharynx on that of the food,—a function of the spinal marrow; and, 2. Those that are executed under the organic and other nerves of involuntary function. It is probable, too, that every living tissue is capable of moving responsive to its appropriate irritant. See Irritability.

MOTION, CILIARY, see Cilia—m. Involuntary, see Automatic—m. Vibratory, see Ciliary—m. Voluntary, Autocinesis.

MOTIVE, Motory.

MOTOR, Motory—m. Externus, Motor oculi externus—m. Oculi, Motores oculorum.

MOTOR OCULI EXTER'NUS, *Motor externus, Ner'vus tim'idus seu abdu'cens, (F.) Nerf oculo-musculaire externe* (Ch.), *Nerf moteur oculaire ex-*

terne, *Sixth pair of nerves*, arises from the furrow which separates the pons Varolii from the medulla oblongata. It enters the cavernous sinus; anastomoses on the outside of the carotid artery with two filaments furnished by the superior cervical ganglion; enters the orbit by the sphenoidal fissure, and is lost on the rectus externus oculi. This pair is, also, called *Abducens*.

MOTORES OCULO'RUM, *Moto'rii Oculo'rurum, Oculares communes, Ophthalmici ext'eri, Motor Oculi, M. O. communis, Oculo-moto'rius, Common Oculo-muscular Nerves*, (F.) *Nerfs moteurs oculaires communs* (Ch.), The third pair of nerves, *Par tert'ium*. This nerve arises from a depression at the inner side of the peduncles of the brain, between the tuber annulare and corpora mamillaria. It enters the outer part of the cavernous sinus, and passes into the orbital cavity. Behind the sphenoidal fissure, it divides into two branches; the one *superior* and the other *inferior*. The former is lost on the rectus superior oculi and the levator palpebræ superioris; the latter—the larger of the two—divides into three branches, which pass to the rectus internus, rectus inferior, and lesser oblique. The last furnishes a filament, which passes to the ophthalmic ganglion.

MOTORIAL, Motory.

MOTORPATHY, (*motor*, and *πάθος*, 'affection,') Kinesipathy.

MOTORY, *Mo'tive, Mo'tor, Moto'rial, Mo'trix, Kinet'ic or Cinet'ic*, (F.) *Moteur*. That which moves or causes movement, as *motor nerves* in contradistinction to *sensory*.

MOTOS, (*μωτος*), Lintum, Tent.

MOTO'SIS, The application of lint to a wound.

MOTRICITÉ, Motility.

MOTRIX, (*movere, motum*, 'to move,') Motory.

MOTTE, Mons veneris.

MOTUM, (*μωτον*), Carbasus, Lintum.

MOTUS, gen. *Mo'tus*, (*movere, motum*, 'to move,')

Motion—*m. Assimilationis*, Plastic force—*m. Automaticus*, see *Automatic*—*m. Compressorius*, *Peristole*—*m. Cordis diastalticus*, *Diastole*—*m. Hæmorrhoidalis*, *Hæmorrhoids*—*m. Intestinorum*, *Borborygmus*—*m. Involuntarius*, see *Autocinesis*—*m. Peristalticus*, *Peristole*—*m. Testudineus*, *Peristole*—*m. Vermicularis*, *Peristole*—*m. Voluntarius*, *Autocinesis*.

MOUCHE, (from *musca*, 'a fly,') *Cantharis*—*m. d'Espagne*, *Cantharis*.

MOUCHER L'ACTION DE, (from (L.) *mucus*) Munctio.

MOUCHES, see *Pains, labour*—*m. Volantes*, see *Metamorphopsia*.

MOUCHETURE, see *Scarification*.

MOUCHOIR EN TRIANGLE, *Couvrechef*, see *Bandeau*.

MOUFETTE, see *Mephitism*.

MOULD, (from *modulus*, dim. of *modus*, 'a measure, or standard of measure,') *Fontanella*.

MOULE, (from *musculus*, [?]) *Mytilus edulis*.

MOULEUR, (from *molaris*, 'a millstone,') Muller.

MOULTING, (from *mutare*, 'to change,') Desquamation.

MOUNTAIN ASH, *Sorbus aucuparia*—*m. Ash*, *American*, *Sorbus Americana*—*m. Service*, *Sorbus aucuparia*—*m. Sickness*, *Puna*.

MOUREILLER, *Malpighia mourella*.

MOURON DES OISEAUX, (from *murus*, 'a wall,') *Alsine media*—*m. Rouge*, *Anagallis*.

MOUSE EAR, *Alsine media*, *Hieracium pilosella*.

MOUSESIGHT, *Myopia*.

MOUSSE DE CORSE, (from *muscus*, 'moss') *Coralina Corsicana*—*m. d'Islande*, *Lichen Islandicus*—*m. d'Irlande*, *Fucus crispus*—*is. Marine*

perlée, *Lichen Islandicus*—*m. Perlée*, *Fucus crispus*.

MOUSSERON, see *Agaric*.

MOUSSEUX, Frothy.

MOUSTACHE, Mystax.

MOÛT DE LA BIERE, ([old F.] *moust*, from (L.) *mustus*, 'new,') Wort.

MOUTARDE, (*Mustard*,) *Sinapis*—*m. des Allemands*, *Cochlearia armoracia*—*m. des Moines*, *Cochlearia armoracia*.

MOUTH, Sax. *muð*, *Os*, *Ca'vum O'ris*, *Sto'ma*, *Gab*, (Sc.) *Moue*, (Prov.) *Gob*, (F.) *Bouche*. This word sometimes signifies the cavity situate between the jaws, and containing the tongue, &c.;—at others, the outer orifice of that cavity, *Peristom'ium*. The mouth, in the first acception, *Ca'vitas seu Spa'tium O'ris*, is the cavity bounded, above, by the palatine arch; below, by the tongue; before, by the lips; and behind, by the velum palati and pharynx. The sides of the mouth and the organs it contains are lined by a mucous membrane. The anterior aperture of the mouth is, sometimes, called *facial*—the posterior, *pharyngeal*. In the mouth are the teeth, gums, alveolar margins, tongue; the excretory ducts of the salivary glands, and those of a number of mucous follicles, &c. It is in this cavity that the food is cut, torn, or bruised by the teeth; is impregnated with saliva, and formed into a mass or *bolus*, which is then subjected to the act of deglutition. The mouth contains the organs of taste; and it serves in respiration, articulation, expectoration, suction, &c.

The condition of the mouth requires to be attended to in *Semeiology*. The state of the mucous membrane, like that of the tongue, indicates the condition of the membrane of the alimentary canal generally.

MOUTH, Os, *Apertu'ra*, *Orific'ium*, *Hia'tus*, *Peristom'ium*, is, also, applied to the open extremities of vessels or other canals.

MOUTH ROOT, *Coptis*.

MOUTH, SORE, *Aphthæ*.

MOUTH WATERING, see *Salivation*.

MOUVEMENT, Motion—*m. Péristaltique*, see *Peristole*.

MOVEMENT, (*movere, motum*, 'to move,') *Dejection*, *Motion*—*m. Cure*, *Kinesipathy*.

MOW (from [F.] *moue*,) Mouth.

MOX'A, *Moxibur'ium*. A word by which the Chinese and Japanese designate a cottony substance, which they prepare by beating the dried leaves of the *Artemisia moxa*,—a kind of mugwort. With this down they form a cone, which is placed upon the part intended to be cauterized, and is set fire to at the top. The heat and pain gradually increase, in proportion as the combustion proceeds downwards, until ultimately an eschar may be formed. In Europe and this country, the same operation is usually practised with a cylinder of cottonwool, or with one formed from the pith of the greater sunflower. This mode of cauterization is employed as a powerful counter-irritant; which it assuredly is. Sloughing may be in some measure prevented, according to Baron Larrey, by the application of liquid ammonia to the burnt part.

The term *moxa* has been extended to any substance, which by gradual combustion on or near the skin is employed as a counterirritant,—as the *electric* or *galvanic moxa* or *caut'ery*, in which voltaic electricity is employed as the cauterizing agent.

MOXIBURIUM, (*moxa*, and *bu'ro*, for *uro*, *ustum*, 'to burn,') *Moxa*.

MOXIBUSTION, *Moxibus'tio*, *Moxocaut'is*, *Encaut'is*. Mode of cauterization by means of *moxa*.

MOXOCAUSIS, (*moxa*, and *kauais*, 'burning,') Moxibustion.

MOXON'S EFFERVESCING MAGNESIA, see Magnesia, effervescing, Moxon's.

MUCAGO, (from *mucus*.) Mucilage.

MUCARUM, Mucharum.

MUCINIUM, (from *mucus*.) Mucilage.

MUCÉDINÉES, (*μυκός*, 'fungus,') see Fungus.

MUCEDO, (from *mucus*.) Mucilage.

MUCHA'RUM, *Muca'rum*. A barbarous term, formerly used for an aqueous infusion of roses, sweetened, and evaporated to the consistence of syrup.

MUCIFIC, (*mucus*, and *facere*, 'to make,') Blenno-genous.

MUCIFORM, *Muciform'is*, *Myxo'des*, *Bleno'i'des*, *Blenno'i'deus*, (*mucus*, and *forma*, 'form,') Resembling mucus in character or appearance.

MUCILAGE, *Mucila'go*, *Muca'go*, *Muce'do*, *Mucin'ium*, (from *mucus*.) A mixture of gum and a small quantity of matter analogous to mucus, which is found in abundance in linseed, quince-seed, &c. It is obtained by heating in water the parts, or products, of plants which contain it. It is much used in the preparation of emollient cataplasms and the greater part of the demulcent tisanes, (F.) *Hydrolés*, *Hydrolites*.

MUCILAGE, see *Liquor sanguinis*—*m. Adragant*, *Mucilago astragali tragacanthæ*—*m. de gomme Adragante*, *Mucilago astragali tragacanthæ*—*m. Arabique*, *Mucilago acaciæ*—*m. de gomme Arabique*, *Mucilago acaciæ*.

MUCILAGE, ANIMAL, Mucus.

MUCILAGE OF BARLEY, *Mucilago hordei*—*m. of Quince-seeds*, *Decoctum cydoniæ*—*m. of Rice*, see *Oryza*—*m. of Sassafras pith*, *Infusum sassafras medullæ*.

MUCILAGE OF SLIPPERY ELM BARK, *Infusum ulmi*.

MUCILAGE, STARCH, *Mucilago amyli*.

MUCILAGINEUX, Mucilaginous.

MUCILAGINOUS, *Mucilagino'sus*, *Gummo'sus*, *Gummi'o'des*, *Gummi'des*, (F.) *Mucilagineux*. Gummy. Resembling gum. That which possesses the character of mucilage.

MUCILAGO, gen. *Mucilag'inis*, Mucilage, *Mucilago Acaciæ*, *Synovia*.

MUCILAGO ACA'CLE, *M. Acaciæ Arab'icæ* seu *Gum'mi Arab'icæ*, *Mucila'go*, *Mistu'ria Aca'ciæ*, *Mucilage of aca'cia or of gum Arab'ic*, (F.) *Mucilage Arabique ou de gomme Arabique*. (*Acaciæ gum. contus. ʒiv; Aq. bullient. Oss. Rub the gum with the water gradually added.*) Demulcent. To allay cough; but, chiefly, to suspend insoluble matters in water.

MUCILAGO AM'YLI, *Starch mu'cilage*, *Decoc'tum Am'yl'i*. (*Amyli, ʒij; Aque, Oj. Rub the starch with the water gradually added; then boil till it forms a mucilage.*—Ph. L.) Demulcent. Generally given *per anum*.

MUCILAGO ARTICULORUM seu **JUNCTURARUM**, *Synovia*.

MUCILAGO ASTRAG'ALI TRAGACAN'THÆ, *M. tragacanthæ* (Ph. U. S.) seu *gum'mi tragacanthæ*, *Mu'cilage of trag'acanth*, (F.) *Mucilage Adragant ou de gomme Adragante*. (*Gummi tragacanth. ʒj; Aque bullient. Oj. Macerate for 24 hours; triturate till the gum is dissolved, and press through linen.*) Used for pharmaceutical purposes.

MUCILAGO CRETICA, *Misturæ cretæ*—*m. Gummi Arabici*, *M. acaciæ*.

MUCILAGO HOR'DEI, *Mu'cilage of Bar'ley*. (*Hor'dei, ʒss, (avoirdupois), Aque, f ʒxvi. Triturate well; and boil. Ph D.*) A vehicle for substances given in enemata.

MUCILAG SASSAFRAS, *Infusum sassafras medullæ*.

MUCILAGO SEMINIS CYDONII MALLI, *Decoc'tum cydoniæ*—*m. Gummi Tragacanthæ*, *M. astragali*

tragacanthæ—*m. Tragacanthæ*, *M. astragali tragacanthæ*—*m. Ulmi*, *Infusum ulmi*.

MUCIN, Mucus.

MUCIPAROUS, *Mucip'arus*, (F.) *Mucipare*, (*mucus*, and *pario*, 'I bring forth,') Mucus-producing; as *muciparous glands* or follicles.

MUCITIS, *Mycodermi'tis*, *Blennophlogo'sis*, *Blennymeni'tis*, (*mucus* and *itis*.) Inflammation of a mucous membrane.

MUCKSWEAT, (Sc. *muck*, 'dirt,') Desudatio.

MUCOCE'LE, *Hernia sacci lacryma'lis*, (*mucus*, and *κηλη*, 'rupture,') An enlargement or protrusion of the mucous membrane of the lachrymal passages, giving occasion to fistula lacrymalis. Also, dropsy of the lachrymal sac.

MUCOENTERITIS, see Enteritis.

MUCOPURULENT, *Mucopurulen'tus*, *Pyoblen'nicus*, (from *mucus*, and *pus*.) Having the character or appearance of mucus and pus.

MUCOR, Euros, Mucus—*m. Mucedo*, see Fungus—*m. Narium*, Nasal mucus.

MUCORINÉES, see Fungus.

MUCOSIN, see Mucus.

MUCOSITAS, Mucosity, Mucus.

MUCOSITY, *Mucosi'tas*, (F.) *Mucosité*. A fluid, which resembles mucus, or contains a certain quantity of it.

MUCOUS, *Muco'sus*, *Blenno'des*, *Myxo'des*, (from *mucus*.) (F.) *Muqueux*. An epithet for all bodies containing mucilage or mucus. It is, also, sometimes used synonymously with gummy. In pathology, it is occasionally employed to express the seat of a disease, as *mucous disease*, *mucous phlegmasia*; that is, having its seat in a mucous membrane.

MUCOUS CORPUSCLES, see Mucus—*m. Layer*, see *Tache embryonnaire*—*m. Membranes*, see *Membranes*—*m. Web*, *Corpus mucosum*.

MUCRO, Apex.

MUCRO seu *Cor'yp'hæ* seu *Fun'dus* seu *Ver'tez Cor'dis*. The apex or point of the heart.

MUCRO HUMERI, Acromion.

MUCRONATA CARTILAGO, (from *mucro*,) Xiphoid cartilage.

MUCRONATUM OS, Xiphoid cartilage.

MUCULENT, *Muculen'tus*, (from *mucus*.) Slimy, viscid.

MUCUNA, see *Dolichos pruriens*—*m. Pruriens*, see *Dolichos pruriens*.

MUCUS, (*μυκός*) *Mu'cor*, *Muco'sus hu'mor*, *Mucosi'tas*, *Myx'a*, *My'cus*, *La'pæ*, *Pitu'ita*, *Blen'na*, *Zoömy'cus*, *Zoömyx'a*, *Pit'uite*, *An'imal mu'cus*, *An'imal mu'cilage*, (F.) *Muqueux animal*, *Morre*, *Mucilage animal*, (presumed to be from *munger*, 'to wipe the nose,') A substance, analogous to vegetable mucilage; from which, however, it differs by affording subcarbonate of ammonia on distillation. Mucus exudes through the skin, in a state of combination with a peculiar oily matter; and, drying, forms the epidermis. It constitutes, in part, the different epidermic productions, as the hair, nails, wool, and horn of animals, feathers of birds, and scales of fish. It is found at the surface of the mucous membranes, and presents some difference in its composition and properties, according to the particular mucous membrane from which it is obtained. Its chief organic constituent is an albuminoid compound,—*mu'cin*, *mu'cosin*.

Mucus preserves the membranes moist, and in a state best fitted for the performance of their functions. The French give the term *glair* to the thick, stringy mucus, secreted by the mucous membranes when in a state of disease.

When mucus is examined with the microscope it is found to contain numerous epithelial scales or flattened cells; together with round, granular bodies, commonly termed *mucous corpuscles* or *mucous glob'ules*, which are contained in a fluid portion—*li'quor mu'ci*.

MUCUS CARPHODES, see *Carphodes*.—*m. Catharticus*, *Lichen Islandicus*.—*m. Malpighii*, *Corpus mucoosum*.—*m. Narium*, *Nasal mucus*.

MUDAR, *Madar*. The *Calo'tropis muda'rii* seu *mada'rii*, *C. gigante'a*, *Asclepias gigante'a*, *Mado'rius*. *Ord. Asclepiadaceæ*. In the Hindoo practice of physic, the bark of the root, as well as the concrete juice of the plant, enters into various compound formulæ for the cure of elephantiasis and many other disorders. Experiments instituted by Dr. Duncan, Jr., exhibit its properties to be like those of *ipecacuanha*. Mixed with oil it is used in gout in Malabar.

MUDGE'S INHALER, see *Inhaler*.

MUDLICK, see *Kentucky*, mineral waters of.

MUET, (from *mutus*), *Mute*.

MUGUET, (from *muscus*, 'moss,' [?]) *Aphthæ*, *Convallaria maialis*.—*m. des Bois*, *Asperula odorata*.—*m. de Mai*, *Convallaria maialis*.

MUGWORT, *Artemisia vulgaris*.

MULATRE, *Mulatto*.

MULAT'TO, (F.) *Mulâtre*, (S.) *Mulata*. An individual of the human species engendered of a white and black race, (from *mulus*, 'a mule.')

The following table, according to Mr. Lawrence, after Ulloa, exhibits the proportion of white blood in the various castes,—arising from the hybridous admixture of white and black,—according to the principles sanctioned by usage.

PARENTS.	OFFSPRING.	DEGREE OF MIXTURE.		
Negro and White.....	Mulatto.....	1-2	White,	1-2 Black.
White and Mulatto.....	Terceron.....	3-4	1-4	1-4
Negro and Mulatto.....	{ Griffo, Griff, or Zambo, or Black Terceron, }	1-4	—	3-4 —
White and Terceron.....	Quarteron or Quadroon.....	7-8	—	1-8 —
Negro and Terceron.....	Black Quarteron or Quadroon.....	1-8	—	7-8 —
White and Quarteron.....	Quinteron.....	15-16	—	1-16 —
Negro and Black Quarteron.....	Black Quinteron.....	1-16	—	15-16 —

The last two were considered to be respectively white and black, in the British West India Islands; and the former, prior to modern changes, were white by law, and consequently free.

Terceron is now scarcely used; and Quarteron or Quadroon takes its place. See *Half Caste*. The Octoroon is the old Quadroon. In other words, at the present day, a Quadroon is the offspring of a white and mulatto; an Octoroon, of the white and quadroon. No other term is used for farther dilutions.

MULATTO JACK, Fever, yellow.

MULBERRY CALCULUS, ([G.] *Maulbeere*), see *Calculi*, urinary and mural.

MUL'BERRY MASS. The mulberry-like appearance, presented by the embryo at a very early stage, when the different cells have drawn around them a portion of the vitellus or yolk.

MULBERRY RASH, see *Typhus*.—*m. Red*, *Morus rubra*.—*m. Tree*, *Morus nigra*.—*m. Wild*, *Morus rubra*.

MULES (F.), *Mulæ*. Chilblains on the heels.

MULET, (from *mulus*, 'a mule,') Hybrid.

MULGEDIUM ACUMINATUM, (*mulgere*, *mulsum*, 'to milk,') *Lactu'ca villosa*, *Blue Let'tuce*; and

MULGEDIUM FLORIDANUM, *Lactu'ca Florida'na*, *Son'chus Florida'na*, *False Let'tuce*; indigenous plants; *Order*, *Compositæ*; have the reputation of curing the bites of rattlesnakes.

MULIEBRIA, (from *mulier*,) *Menses*, *Vulva*.

MULIEBRITY, *Mulieritas*.

MULIEBROSITAS, (*mulieb'ris*, appertaining to women,') *Philogyn'ia*. Fondness for women. Hence, *Mulieb'rosus*, *Muliera'rius*, and *Muliero'rus*; one who is fond of women.

MULIEBROSUS, see *Muliebrositas*.

MULIER, gen. *Mul'ieris*. Woman.—*m. Medica*, see *Iatrine*.—*m. Obstetric*, *Midwife*.

MULIERARIUS, see *Muliebrositas*.

MULIERITAS, *Mulieb'ritiy*, *Womanhood*. The state of puberty in the female.

MULIEROSUS, see *Muliebrositas*.

MULIER PLENA, ('a full woman,') *Pregnant*.

MULLED WINE, (from *mollire*, 'to make milder,') see *Wine*.

MULLEIN, **BLACK**, *Verbascum nigrum*.—*m. Broad-leaved*, great, *Verbascum nigrum*.—*m. Yellow*, *Verbascum nigrum*.

MUL'LER, (from *molaris*, 'a millstone,' *mola*, 'a mill,') (F.) *Mouleur*. A moulder. A stone held

in the hand, with which any powder is ground upon a horizontal slab. It is sometimes called *mullet*, (F.) *Mollette*.

MULLIGRUBS, *Tormina*, *Colic*.

MULLIN, *Melizomum*.

MULOMEDICINA, (*mulus*, 'a mule,' and *medicina*,) *Veterinary art*.

MULOMEDICUS, see *Hippiater*.

MULSA ACIDA, (*mulsus*, 'mixed with honey') *Oxyglyeus*.

MULSUM, *Hydromeli*.—*m. Vinosum*, *Enomel*.

MULTIFIDUS SPINÆ, (*multus*, 'many,' and *findere*, 'to divide,') *Transversalis dorsi*.

MULTIFETA'TION, *Multifeta'tio*, (*multus*, 'many,' and *fetus*,) *Pregnancy* with more than two fetuses.

MULTILOC'ULAR, *Multilocula'ris*, (F.) *Multiloculaire*, (*multus*, 'many,' and *loculus*, 'a small place, a cell,') Having many cells or small cavities.

MULTIMAM'MÆ, (F.) *Multimammæ*, (*multus*, 'many,' and *mamma*, 'a breast,') A variety of hypergenesis, in which there are supernumerary mammae.

MULTIP'AROUS, *Multip'arus*, (F.) *Multipare*, (*multus*, 'many,' and *parire*, 'to bring forth,') One that brings forth several young at the same time. Such a birth is called *plural*.

MULTIPLICATION MERISMATIQUE, see *Generation*.

MULTITESTIS, (*multus*, and *testis*, 'a testicle,') *Polyorchis*.

MULTITUDO, gen. *Multitu'dinis*, (*multus*, 'many,') *Plethora*.

MULTIVORANTIA, (*multus*, and *vorare*, 'to devour,') *Polyphagia*.

MULTIVOROUS, *Polyphagous*.

MUMIA, *Mummy*.—*m. de Medullis*, *Marrow*.

MUMISATIO, *Mummification*.

MUMMIFICATION, *Mumisa'tio*, (*mummy*, and *fio*, 'to be made,') *Sceletu'sis*, *Skeleton'sis*. The mode of preparing a mummy.

MUM'Y, *Mum'ia*, (Arab. *Moumya*, from *mum*, 'wax,') *Rebolea*, *Rebona*, *Sceletum'a*, (F.) *Momie*. A dead body simply dried, or dried after having been embalmed. The latter acceptation is the most common. Formerly, the Egyptian mummy was extolled as useful in contusions. It was presumed, also, to have healing, tonic, and resolvent properties. It is now only regarded as an archæological curiosity. See *Skeleton*.

MUMPS, *Cynanche parotidea*.

MUNCHATOE, *Mystax*.

MUNC'TIO, (F.) *l'Action de moucher*, (mungere, munc'tum, 'to blow the nose.') The act of blowing the nose.

MUND'IA SPINO'SA, (from *mundus*, 'clean, neat,' from its appearance.) A plant of the *Nat. Ord.* Polygalæ, which grows in Southern Africa, and a decoction of whose branches is used in atrophy, phthisis, &c.

MUNDIFICANTIA, (*mundus*, 'clean,' and *facere*, 'to make,') Detergents.

MUNDIFICATIVA, Detergents.

MUNGOS RADIX, Ophiorrhiza mungos.

MUNGREL, (see Mongrel.) Hybrid.

MUNIMENTUM CASTITATIS, ('muniment of chastity,') Hymen.

MU'QUEUX, Mucous—*m. Animal*, Mucus.

MUQUEUSE, Membrane, (Mucous.)

MUR, Murr, see Coryza.

MURAL (F.), *Muralis*, ([F.] *Mûre*, 'a mulberry,') Mulberry. Vesical calculi are so called, when rugous and covered with tubercles or asperities. They are composed of oxalate of lime.

MÛRE, Moro, see *Morus nigra*.

MURFLES, Ephelides, Papule.

MURIA, *Hal'mé*, *Salsila'o*, *Salse'do*, *Salsitu'do*, *Salsu'go*, Brine, (F.) *Saumure*, ('admixis, 'salt water,') This was formerly used as a glyster in dysentery, in certain inveterate neuralgias, &c. Muria was also a kind of pickle or sauce made by the Romans from the tunny fish.

MURIA, Soda, muriate of.

MURIAS AURICONATRICUM, see Gold.

MURIATE D'AMMONIAQUE, Ammoniac murias—*m. d'Ammoniaque et de fer*, Ferrum ammoniatum—*m. d'Or*, Gold, muriate of, see Gold.

MURIATIC AC'ID, (from *muria*; *Ac'idum sa'lis seu mari'num concentra'tum seu sa'lis culina'ris seu sa'lis mari'ni*, *Spir'itus sa'lis mari'ni seu sa'lis Glauberi seu sa'lis ac'idus seu fu'mans*, *Ac'idum hydrochloricum*, *A. muriaticum* (Ph. U. S.), *Spir'it of salt*, (F.) *Acide hydrochlorique ou muriatique*. An aqueous solution of chlorohydric acid gas of S. g. 1.16. The odour of muriatic acid is suffocating; taste very acid and caustic. It is nearly colourless when pure; but commonly of a pale yellow; volatile; the fumes visible. Muriatic acid is possessed of tonic and antiseptic properties. It is used in typhus; cutaneous eruptions, in gargles for inflammatory and putrid sore throats, (zett. xxx to f 3vj of water.) &c. When added to a pediluvium, it renders it stimulating.

ACIDUM MURIATICUM DILU'TUM of the Ph. U. S. contain four troycounces of muriatic acid, to a pint of distilled water.

MURIDE, (after *Muria*.) Bromine.

MÛRIER NOIRE, *Morus nigra*.

MURIFORM, *Muriformis*, ([F.] *mûre*, 'a mulberry,' and [L.] *forma*, 'shape,') Mulberry shaped, as *Muriform mass* or *body*, (F.) *Amas* or *Corpus mûriforme*. The collection of vitelline globes proceeding from the segmentation of the vitellus.

MURIGENE, (*muria* and *gennao*, 'to generate,') Chlorine.

MURINA, (after *Muria*.) Bromine.

MUR'MUR, (F.) *Murmure*. (A word existing both in the Romanic and Teutonic languages, and probably a variety of onomatopœia;—*mur mur* expressing the kind of sound which the word indicates.) A low continued or continuously repeated sound, as that of flame, or of a stream running over a stony bottom.

MURMUR AURIUM, Tinnitus Aurium—*m. Intestinalis seu Intestinarum seu Ventris*, Borborygmus.

MURMUR, RESPIRATORY, (F.) *Bruit respiratoire on vésiculaire*, *Murmure vésiculaire*. The noise heard during inspiration and expiration, especially the former. It is produced by the passage of the air through the bronchial tubes and into the air-cells. It has been also called *mur'*

mur of the expansion of the lungs, (F.) *Bruit d'expansion pulmonaire*; and, when distinctly vesicular, *Respiration of the cells or vesicular respiration*, (F.) *Respiration on murmure vésiculaire*. Vesicular respiration is of course absent when the cells of the lungs have been obliterated from any cause. We may then have the *Respiration nulle*, *Absence du bruit respiratoire*, *Silence*, and *Respiration silencieuse* of the French writers.

At times, it is rude during inspiration or expiration, or both—the *Respiration rude* or *R. râpeuse* of the French. At others, there is a blowing sound, (F.) *Souffle*, *Respiration soufflante*, as if some one were blowing into the auscultator's ear through a tube. This is heard in the healthy state over the larynx, trachea, (F.) *Souffle trachéal*, and about the bifurcation of the bronchia, (F.) *Souffle bronchique*; but when it proceeds from the lungs it denotes disease. It may be tubular or diffused. In the former, the whiffing murmurs appear to occur in a space limited to the immediate neighbourhood of the part examined. In the latter, they are produced with but moderate intensity, and sometimes at a distance from the ear, over a tolerably extended space.

The respiration, perceived over the trachea and bronchia in health, is called *tracheal* or *bronchial* or *tubal*, (F.) *Respiration bronchique*, *Souffle tubaire*, according to the situation in which it is heard.

MURMUR, UTERINE, *Bruit placentaire*—*m.* Utero-placental. *Bruit placentaire*—*m.* Vesicular, Murmur, respiratory.

MURMUR, WHIFFING, see Murmur, respiratory.

MURR, *Mur-rain*, (Sax. *mýrphan*, 'to destroy,' or from (L.) *mori*, 'to die,') (F.) *Claveau*, *Clavelle*, *Pestilent'ial Fe'ver*. An epizootic, perhaps contagious, disease, having some resemblance to smallpox, which affects cattle—especially sheep; and is said to have been transferred to man.

MURRAIN, Murr, see Coryza.

MURREN, Murr, Coryza.

MURTON, (*improv.*) Clitoris.

MURTUM, Clitoris.

MUSA, (after Antonius Musa, physician to Augustus,) *M. Paradisiaca*—*m.* Cliffortiana, *M. Paradisiaca*—*m.* Mensaria, *M. Paradisiaca*.

MU'SA PARADISI'ACA, *Mu'sa*, *M. Mensaria* seu *Cliffortiana*, *Amu'sa*, *Pal'ma hu'milis*, *Ficus Indica*, *Ba'ta*, *Platanus*, *Plan'tain tree*, (F.) *Plaintain-arbre*, *Platanier*, *Bananier commun*. Family, Musaceæ. *Sex. Syst.* Hexandria Monogynia. It grows in many parts of India and South America. The fruit, which is largely eaten for bread, consists of a mealy substance, *Plan'tain meal*, *Cinquin-tay*. It is clammy; has a sweetish taste, and will dissolve in the mouth without being chewed. The spike of the fruit often weighs forty or fifty pounds. The leaves of the tree serve the Indians for tablecloths and napkins. Being smooth and soft, they are also employed as dressings for blisters. The water from the soft trunk is astringent, and sometimes used in diarrhæa.

MUSA SAPIEN'TUM. The *Bana'na tree*, *Baco'ba*, (F.) *Figuier-bananier*, *Bananier*, *Bacove*. This differs somewhat from the last. The fruit, *Bana'na*, (S.) *Platano*, is shorter, straighter, and rounder; the pulp softer and of a more luscious taste. When ripe, it is very agreeable; it is eaten like the plantain, and relished by all ranks of people in the West Indies. Both varieties are natives of Guinea.

MUSCO, (from *μωσχος*.) Musk.

MUSCA HISPANICA, ('Spanish fly,') *Cantharis*.

MUSCÆ, ENTOHYALOID, see Metamorphopsia—*m.* Volitantes, see Metamorphopsia.

MUSCADE, see Myristica moschata.

MUSCADIER, Myristica moschata.

MUSCERDA, Moschardina. Also, the dung

of mice, (*mus*, 'a mouse,' and *merda*, 'dung,') formerly given in epilepsy.

MUSCLE, (old Eng.) *Bracon*, *Musculus*, *Totulus*, *Laceratus*, *Mys*, *My'on*, (from *mys*, 'a mouse or rat,' because, say some etymologists, the ancients compared the muscles to flayed mice or rats. According to Diemerbroeck, Douglass, Chaussier, &c., *μύω*, comes rather from *μύω*, 'to close,' 'to move,' &c., a function proper to muscles. This etymon is the more probable.) Muscles have been divided into those of *Animal life* or *of the life of relation*—*voluntary muscles*—which execute movements under the influence of the will; as the muscles of the limbs, head, trunk, &c., and into those of *organic life*—*involuntary muscles*—which contract under the influence of certain special stimuli; as the heart, fleshy fibres of the stomach, &c. *Mixed muscles* are those which belong partly to each of these divisions;—as the muscles of respiration, and the sphincters. Muscles that act in opposition to each other are called *antagonists*; thus, every extensor has a flexor for an antagonist, and conversely. Muscles that concur in the same action are termed *congen'erous*. The muscles present numerous varieties in form, size, situation, use, &c., and have been divided, by some, into *long*, *broad*, and

short. Each of these divisions comprises *simple* and *compound* muscles. *Simple* or *rectilinear muscles* have all their fibres in a similar direction, and only one body—as the *Sartorius*, *Pronator quadratus*, &c. *Compound muscles* are those which have only one belly and several tendons, as the flexors of the fingers and toes; or several bellies and several tendons,—as the biceps flexor cubiti, *sacrolumbalis*, &c. To the compound muscles belong, also, the *radiated muscles*. Their fibres set out from a common centre, and are arranged like the radii of a circle;—such are the diaphragm, *iliacus*, *temporal*, &c. *Pen'nated* or *Pen'niform Muscles*: their fibres are arranged in two rows, which are united at a median line, at greater or less angles; nearly as the feathers are inserted into a quill. The *palmaris longus* is one of these. *Sem'ipenniform muscles*: their fibres are oblique, as in the last case; but they are inserted only on one side of the tendon. *Hollow Muscles* are,—the heart, intestines, urinary bladder, &c.

Much difference has existed in the enumeration of muscles. Some authors reckon them at upwards of 400. Chaussier admits only 368. The greater part of them are in *pairs*. Very few are *az'ygous*.

TABLE OF THE MUSCLES, ARRANGED AFTER THE MANNER OF DR. BARCLAY, ACCORDING TO THEIR ACTIONS.

THE HEAD IS MOVED

Forwards by

Platysma myoides,
Sternomastoideus,
Rectus anticus major,
" " minor,

Assisted (when the lower jaw is fixed) by

Mylohyoideus,
Geniohyoideus,
Geniohyoglossus,
Digastric.

Backwards by

Part of trapezius,
Splenius capitis,
Complexus,
Trachelomastoidens,
Rectus posticus major,
" " minor,
Obliquus capitis superior.

To either side by

Platysma myoides,
Sternomastoideus,
Part of trapezius,
Splenius capitis,
" colli,
Trachelomastoideus,
Complexus.

THE NECK IS MOVED

Forwards by

Platysma myoides,
Sternomastoideus,
Digastricus,
Mylohyoideus,
Geniohyoideus,
Geniohyoglossus
Omohyoidei,
Sternohyoidei,
Thyrohyoidei,
Rectus anticus minor,
Longus colli.

Backwards by

Part of trapezius,
Rhomboides minor,
Serratus posticus superior,
Splenius capitis,
" colli,
Complexus,
Trachelomastoidens,
Transversalis colli,
Interspinales colli,
Semispinales colli,
Rectus posticus major,
" " minor,
Obliquus capitis superior,
" " inferior,
Scaleni postici,
Levator scapulae.

Laterally by

Various combinations of those muscles which separately move it forwards and backwards, assisted by the scaleni, intertransversales, and recti laterales.

THE TRUNK IS MOVED

Forwards by

Rectus abdominis,
Pyramidalis,
Obliquus externus abdominis,
Obliquus internus,
Psoas magnus,
" parvus,

Assisted (when the arms are carried forwards) by

Pectoralis major,
" minor,
Serratus magnus.

Backwards by

Trapezius,
Rhomboides major,
Latissimus dorsi,
Serratus posticus superior,
" " inferior,
Sacrolumbalis,
Longissimus dorsi,
Spinales dorsi,
Semispinales dorsi,
Multifidus spinæ,
Intertransversales dorsi et lumborum.

Laterally by

Obliquus externus,
" internus,
Quadratus lumborum,
Longissimus dorsi,
Sacrolumbalis,
Serrati postici,
Latissimus dorsi.

THE SCAPULA IS MOVED.

<i>Upwards by</i>	<i>Downwards by</i>	<i>Forwards by</i>	<i>Backwards by</i>
Trapezius, Levator scapulæ, Rhomboides.	Lower part of trapezius, Latissimus dorsi, Pectoralis minor.	Pectoralis minor, Serratus magnus,	Part of trapezius, Rhomboides, Latissimus dorsi.

THE HUMERUS IS MOVED

<i>Forwards by</i>	<i>Backwards by</i>	<i>Inwards by</i>	<i>Rotated inwards by</i>
Part of deltoid, Part of pectoralis major, <i>Assisted in some circumstances by</i> Biceps, Coracobrachialis.	Part of deltoid, Teres major, " minor, Long head of triceps, Latissimus dorsi.	Part of pectoralis major, Latissimus dorsi.	Subscapularis, <i>Assisted occasionally by</i> Pectoralis major, Latissimus and teres major.
			<i>Outwards by</i> Supraspinatus, Infraspinatus, Teres minor.

THE FOREARM IS MOVED

<i>Forwards by</i>	<i>Backwards by</i>	<i>Rotated inwards by</i>
Biceps, Brachialis anticus, Pronator teres, <i>Assisted by</i> Flexor carpi radialis, " sublimis, " ulnaris, Supinator longus.	Triceps, Anconeus.	Pronator teres, Flexor carpi radialis, Palmaris longus, Flexor sublimis, Pronator quadratus.
		<i>Outwards by</i> Biceps, Supinator brevis, Extensor secundi internodii.

THE CARPUS IS MOVED

<i>Forwards by</i>	<i>Backwards by</i>	<i>Outwards by</i>	<i>Inwards by</i>
Flexor carpi radialis, Palmaris longus, Flexor sublimis, " carpi ulnaris, " profundus, " longus pollicis.	Extensor carpi radialis longior, Extensor carpi radialis brevior, Extensor secundi internodii, Indicator, Extensor communis digitorum, Extensor proprius pollicis.	Flexor carpi radialis, Extensor carpi radialis longior, Extensor carpi radialis brevior, Extensor ossis metacarpi, Extensor primi internodii.	Flexor sublimis, " carpi ulnaris, " profundus, Extensor communis digitorum, Extensor minimi digiti, Extensor carpi ulnaris.

THE THUMB IS MOVED

<i>Inwards and forwards, across the palm, by</i>	<i>Outwards and backwards by</i>	<i>Upwards and forwards, away from the other fingers, by</i>	<i>Backwards and inwards, to the other finger, by</i>
Opponens pollicis, Flexor brevis, " longus.	Extensor ossis metacarpi pollicis, Extensor primi internodii, Extensor secundi internodii.	Abductor, <i>Assisted by part of the</i> Flexor brevis.	Adductor, Extensor primi internodii, Extensor secundi internodii.

THE FINGERS ARE MOVED

<i>Forwards, or flexed, by</i>	<i>Backwards, or extended, by</i>	<i>Outwards, to radial border, by</i>	<i>Inwards by</i>
Flexor sublimis, " profundus, Lumbricales, Interossei, Flexor brevis digiti minimi, Abductor digiti minimi.	Extensor communis, " minimi digiti, Indicator.	Abductor indicis, " digiti minimi, Interossei.	Abductor digiti minimi, Interossei.

THE THIGH IS MOVED

<i>Forwards by</i>	<i>Backwards by</i>	<i>Inwards by</i>	<i>Outwards by</i>
Psoas magnus, Iliacus, Tensor vaginæ femoris Pectineus, Adductor longus, " brevis.	Gluteus maximus, Part of gluteus medius, Pyriformis, Obturator internus, Part of adductor magnus, Long head of biceps, Semitendinosus, Semimembranosus.	Psoas magnus, Iliacus, Pectineus, Gracilis, Adductor longus, " brevis, " magnus, Obturator externus, Quadratus femoris.	Tensor vaginæ femoris Gluteus maximus, " medius, " minimus, Pyriformis.

THE THIGH IS ROTATED

<i>Inwards by</i>	<i>Outwards by</i>
Tensor vaginæ femoris, Part of gluteus medius, <i>And, when the leg is extended,</i> <i>by</i> Sartorius, Semitendinosus.	Gluteus maximus, Part of gluteus medius, Pyriformis, Gemellus superior, Obturator internus, Gemellus inferior, Quadratus femoris, Obturator externus, Psoas magnus, Iliacus, Adductor longus, " brevis, " magnus, Biceps cruris, slightly.

THE LEG IS MOVED

<i>Backwards, or flexed, by</i>	<i>Extended by</i>
Semitendinosus, Biceps, Semi-membranosus, Gracilis, Sartorius, Popliteus.	Rectus, Crureus, Vastus externus, " internus.

THE FOOT IS MOVED

<i>Forwards, or flexed, by</i>	<i>Backwards, or extended, by</i>	<i>Inclined inwards by</i>	<i>Outwards by</i>
Tibialis anticus, Extensor proprius pollicis, Extensor longus digito- rum. Peroneus tertius.	Gastrocnemius, Plantaris, Soleus, Flexor longus digitocum, " longus pollicis, Tibialis posticus, Peroneus longus, " brevis.	Extensor proprius pollicis, Flexor longus digitorum, " longus pollicis, Tibialis posticus.	Peroneus longus, " brevis, Extensor longus digito- rum, Peroneus tertius.

THE TOES ARE MOVED

<i>Backwards, or flexed, by</i>	<i>Forwards, or extended, by</i>	<i>Inclined inwards by</i>	<i>Outwards by</i>
Abductor pollicis, Flexor brevis digitorum, Abductor minimi digiti, Flexor longus pollicis, " digitorum, " accessorius, Lumbricales, Flexor brevis pollicis, Adductor pollicis, Flexor brevis minimi di- giti, Interossei.	Extensor longus digito- rum, Extensor proprius pollicis, " brevis digitorum.	Abductor pollicis, Interossei.	Abductor pollicis, " digiti minimi, Interossei.

Muscles have been variously named. 1. *According to their uses*, as diaphragm, buccinator, extensors, flexors, adductors, abductors, levators, depressors, &c. 2. *According to their position*, as interspinales, interossei, subclavius, popliteus, anconeus, cubitalis, iliacus, temporalis, &c. 3. *According to their shape*, as trapezius, splenius, lumbricalis, serratus, digastric, deltoid, scalenus, rhomboides, &c. 4. *According to their dimensions*, as pectoralis major, rectus capitis anticus major, pectoralis minor, glutæus maximus, medius, and minimus. 5. *According to their direction*, as obliquus abdominis, transversalis abdominis, rectus femoris, rectus abdominis, &c. 6. *According to their composition*, as semimembranosus, semitendinosus, complexus, &c. 7. *According to their attachments*, or the different points of the skeleton with which they are connected by means of tendons or aponeuroses; as sternocleidomastoideus, sternohyoideus, &c. On this is grounded the nomenclature of M. Dumas, and that of Chaussier, which is full of meaning.

The end of the muscle, which adheres to the most fixed part, is usually called the *origin* or *head*, *caput*, (F.) *Tête*; and that which adheres to the more movable part, the *insertion* or *tail*, *cauda*, (F.) *Queue*; the intervening part or *body* of the muscle being called the *venter*, *gaster*, or *belly*, *Venter musculi*, *Me'dium musculi*, (F.) *Ventre*: hence the names *gastrocnemii*, *digastricus*, *biceps*, and *triceps*; according as they have two bellies, two or three heads, &c.

Muscles are formed,—1. Essentially of the *muscular* or *fleshy fibre*, (see Muscular Fibre.) 2. Of *Areolar tissue*, which unites together the fibres. This areolar tissue is not very visible between the fine and loose fibres; but becomes more so, when they unite in more considerable fasciculi. It forms, moreover, to each muscle, an external envelope, which unites it to the neighbouring parts, and admits of its motion. This envelope was formerly called *Tu'nica pro'pria musculo'rum*. 3. Of *Ar'teries*. These proceed from neighbouring trunks, and are, generally, very large. Their size and number are always in proportion to the bulk of the muscle. With the exception of some viscera, as the lungs and the kidneys, there are few organs that receive as much blood as the muscles. 4. Of *Veins*. They follow the same course in the muscles as the arteries. Bichat asserts that they are generally devoid of valves. 5. Of *Lymphat'ics*. Of these we know little, and cannot easily follow them between the fleshy fibres. 6. Of *Nerves*. These are numerous, and of different sizes. They, almost all, proceed from the encephalon; some, however, issue from ganglions, and accompany the arteries. In general, they penetrate the fleshy tissue along with the vessels, with which they are closely united. After they have entered the muscles, they divide and subdivide until they are lost sight of.

MUSCLE CANIN, Levator anguli oris — *m. Canellé*, Ischiotrochanterianus.

MUSCLE OF GAVARD. The oblique muscular fibres of the stomach.

MUSCLE GRAND FESSIER, Glutæus major — *m.* of Guthrie, Compressor urethræ — *m.* of Horner, Tensor tarsi — *m.* of Houston, Compressor venæ dorsalis penis.

MUSCLE JUICE. A fluid surrounding the fibres of striated muscle and the fibre-cells of smooth muscles, which differs from the liquor sanguinis of the blood; containing a considerable amount of casein. It seems to be produced within the sarcolemma, through which it exudes.

MUSCLE TRÈS LARGE DU DOS, Latissimus dorsi — *m.* Merkel's, ceratocricoid — *m.* of Wilson, Compressor urethræ.

MUSCULAR, *Muscula'ris*, *Musculo'eus*, *Toro'*-

us, (F.) *Musculaire*, *Musculeux*. That which belongs or relates to the muscles. Well furnished with muscles, *Lacerta'sus*.

MUSCULAR ARTERIES. Arteries that are distributed to the muscles. The name *Muscular Arteries* of the eye has been especially given to two branches sent off by the ophthalmic artery:—the one, *inferior*, which furnishes branches to the rectus inferior oculi, lachrymal sac, &c., and some of the anterior ciliary arteries: the other—the *superior*, which is sometimes wanting; but, when it exists, gives branches to the rectus superior oculi and levator palpebræ superioris.

MUSCULAR CONTRACTION. The exertion of the power, possessed by muscles, of shortening themselves, or of contracting to produce motion:—*muscular motion* being the change in the situation and relation of organs, induced by muscular contraction. When a muscle contracts, its fibres assume more the zigzag direction, and the extremities approximate; but the bulk of the whole muscle is not augmented. This contraction takes place, at times, with extreme velocity: a single thrill, in the letter R, can be pronounced in the 1-30,000th part of a minute. The force of contraction, *Myody'namis*, depends upon the healthy physical condition of the muscle, combined with due energy of the brain. The duration, in voluntary motion, is for a certain time dependent upon the will:—contractions, excited involuntarily, cannot be so long maintained.

MUSCULAR CUR'RENT. A current which, according to Matteucci, exists in every living muscle, whose tendinous extremities are not equally disposed—one, for example, forming a cord; the other broad and ribbonlike—and which circulates in the muscle from the tendinous extremity to the muscular surface.

MUSCULAR FIBRE, *Fleshy fibre*, *Filum muscula're*, *Fibril'la muscula'ris*. A name given to the filaments, which, by their union, form the muscles. This fibre is flat, soft, downy, linear, but little elastic, and more or less red. It is firmer in adults than in the young or the aged; of the same size in the great and small muscles; and runs its course without bifurcation or ramification. It is but slightly resisting in the dead body, tearing readily; but during life it supports very great efforts without laceration. It is, itself, composed of a considerable number of fibrils, similar to each other, and subdividing almost *ad infinitum*. The ultimate filaments, into which the fibre can be decomposed by mechanical means, seem to be hollow or tubular. The fibre exists under two forms, the *striated* or *striped*, (F.) *Fibre striée*, and the *non-striated*, *smooth*, or *unstriated*, (F.) *Fibre lisse*; the muscles composed of the former ministering, as a general rule, to the *animal functions*,—the latter always, perhaps, to the *organic*. The colour of the muscular fibres is red in man, and white in several animals. A greater or less number,—united in fasciculi, *Fasciculi seu Lacer'ti musculo'rum*, approximated to each other, and forming a distinct mass, of very variable size and shape, the extremities being attached to bones by means of tendons,—constitutes a muscle. In this are included areolar membrane, vessels and nerves. See Muscle.

MUSCULAR NERVES. The nerves distributed to the muscles are so named. Winslow calls the 4th pair of nerves—*Nerf musculaire oblique supérieur*.

MUSCULAR SYSTEM. A term given to the aggregate of the muscles of the body.

MUSCULAR VEINS. These bring back the blood, which has been carried to the muscles by the muscular arteries.

MUSCULARIS, 'stout,' 'strong,' Corroborant.

MUSCULATION, See Locomotion.

MUSCULATURE (F.) The condition of the

muscular system:—as *une bonne musculature*, 'a good condition of the muscular system.'

MUSCULEUX, Muscular.

MUSCULI ACCESSORII AD SACROLUMBALEM, see *Sacrolumbalis*—*m. Papillares*, *Columnæ carnae*—*m. Pectinati*, Pectinated muscles.

MUSCULIN, see *Fibrin*.

MUSCULOCUTANEOUS, *Musculocuta'neus*, (from *musculus*, and *cutis*, 'skin.') That which appertains to muscles and skin.

MUSCULOCUTANEOUS NERVE. This name is especially given to two nerves;—the *one*, the *External cutaneous nerve*, furnished by the brachial plexus (see *Cutaneous*):—the *other*, given off from the popliteus externus, (F.) *Nerv. Sciatique*, *Poplite externe*, *Prétibio digital* (Ch.). It descends on the anterior and outer part of the leg,—at first, hid among the muscles; becomes superficial about the middle of the leg; and divides into two branches, which pass superficially on the back of the foot. Two musculocutaneous nerves—superior and inferior—proceed from the first lumbar. The *superior musculocutaneous*—*il'ioscerotal*, *il'iohy-pogast'ricus*, which divides into two branches,—abdominal and scrotal. The *inferior musculocutaneous* is smaller than the superior, and is distributed as its name, *il'ioin'guinal*, indicates.

MUSCULORACHIDÆUS. Belonging or relating to muscles and to the spine. A name given to branches, furnished behind by the intercostal, lumbar, and sacral arteries, which are distributed to the spine and to the muscles of the neighbourhood.

MUSCULOSPIRAL NERVE, Radial nerve.

MUSCULOSA EXPANSIO, ('muscular expansion,') *Platysma myoides*.

MUSCULOSUS, Muscular.

MUSCULUS, Muscle—*m. Aclivis*, *Obliquus internus abdominis*.

MUSCULUS ANOMALUS. A muscular fasciculus, attached only to the superior maxillary bone, with which the Levator labii superioris alaeque nasi muscle is related.

MUSCULUS AUXILIARIUS, *Pyramidalis abdominis*—*m. Constrictorius*, *Sphincter*—*m. Cutaneus*, *Platysma myoides*—*m. Eustachii*, *Laxator tympani*—*m. Fallopii*, *Pyramidalis abdominis*—*m. Femoris membranousus*, *Fascia lata*—*m. Patientiæ*, *Levator scapulae*—*m. Penicillatus*, *Levator labii inferioris*—*m. Risorius* *Santorini*, see *Platysma myoides*—*m. Scandularius*, *Parathenar*—*m. Subcutaneus*, *Platysma myoides*—*m. Succenturiatus*, *Pyramidalis abdominis*—*m. Suceingens*, *Diaphragm*—*m. Supercilli*, *Corrugator supercillii*—*m. Testicodius*, *Cremaster*—*m. Testis*, *Cremaster*—*m. Tubæ novæ*, *Circumflexus*.

MUSCUS ARBOREUS, ('tree moss,') *Lichen plicatus*—*m. Caninus*, *Lichen caninus*—*m. Catharticus*, *Lycopodium selago*—*m. Clavatus*, *Lycopodium*—*m. Corallinus*, *Corallina Corsicana*—*m. Crani humani*, *Lichen saxatilis*—*m. Erectus*, *Lycopodium selago*—*m. Helminthocortos*, *Corallina Corsicana*—*m. Islandicus*, *Lichen islandicus*—*m. Marinus*, *Corallina Corsicana*—*m. Maritimus*, *Corallina*—*m. Pulmonarius quercinus*, *Lichen pulmonarius*—*m. Pyxidatus*, *Lichen pyxidatus*—*m. Pyxoides terrestris*, *Lichen pyxidatus*—*m. Villosus*, *Peristroma*.

MUSEAUDE TANCHE, ('the tench's mouth or snout,') *Os uteri*.

MUSENA BARK, see *Rottlera Schimperii*.

MUSEUM ANATOMIUM, *Supellex anatomica*. An anatomical museum.

MUSHROOM, ([F.] *Mousseron*,) *Fungus*.

MUSIC, *Musica*, (F.) *Musique*, (*музыка*, *musa*, 'a song,') The art of producing harmonious and cadenced sounds; an art, which has, at times, been beneficially used in diseases, particularly in those of the mind; or on which the mind could act in a salutary manner.

MUSICAL SOUND, *Sifflement modulé*.

MUSICOMANIA, *Musomania*, (*music*, and *mania*.) A variety of monomania in which the passion for music is carried to such an extent as to derange the intellectual faculties.

MUSING, *LISTLESS*, (from *musa*, 'study,') *Aphelexia otiosa*.

MUSIQUE, Music.

MUSK, *Mos'chus*, *Mosch*, (Arab.), (F.) *Musc*. A peculiar concrete animal substance, of a very diffusible odour, bitter taste, and a deep brown colour: solid, and enclosed in the preputial follicles of *Mos'chus moschiferus*, *Musk Deer*, (F.) *Chevrotain porte-musc*. It is possessed of stimulant and antispasmodic properties. Dose, gr. v to ʒj in bolus.

MUSK, **ARTIFICIAL**, *Mos'chus factit'ius*, *Resina Succini*. (*Ol. succini rectific.* 1 part, *Acid. nitric.* 4 parts. Digest;—a black matter will be deposited, which must be well washed with water.)

MUSKGRAPE FLOWER, *Bulbus vomitorius*.

MUSKMALLOW, *Hibiscus abelmoschus*.

MUSKROOT, *Sumbul*.

MUSKSEED, see *Hibiscus abelmoschus*.

MUSKWOOD, *Thymiana*.

MUSOMANIA, *Musicomania*.

MUSQUITO, *Mosquito*.

MUSSËN'DA FRONDO'SA, *Belil'la*, *Beleson*, A decoction of this Ceylonese plant has been esteemed refrigerant.

MUSSEL, *Mytilus edulis*.

MUSSELLED, see *Mytilus edulis*.

MUSSITA'TIO, *Mussitatio*, (from *mussitare*, *mussitatum*, itself from *mussare*, 'to murmur,') A condition, in which the tongue and lips move, as in the act of speaking, but without sounds being produced. This sort of murmuring is an unfavourable sign in disease, as indicating great cerebral debility.

MUSTA'CEUM, (from *Mustum*, 'must,' used in making it.) A kind of wedding-cake with the ancients, which consisted of meal, aniseed, cummin, and several other aromatics; its object—it has been conceived—being to prevent or remove the indigestion occasioned by too great indulgence at the marriage feast. *Laureolam in mustaceo quærere*—"to seek for glory in pitiful performances." (Cicero.)

MUSTACHE, *Mystax*.

MUSTARD, **BLACK**, [F.] *mountarde*, (from *moût*, 'must,' because the seeds were bruised with vinegar or must—hence, according to some, from *mustum ardens*, [?]), *Sinapis*—*m. Clammy*, *Polanisia graveolens*—*m. Essence of*, *Whitehead's*, see *Sinapis*—*m. False*, *Polanisia graveolens*—*m. Green*, *Lepidium sativum*—*m. Hedge*, *Erysimum*—*m. Mithridate*, *Thlaspi campestre*—*m. Stinking hedge*, *Alliaria*—*m. Treacle*, *Thlaspi arvense*—*m. White*, *Sinapis alba*—*m. Wild*, *Sinapis arvensis*.

MUSTELANEUS, (from *mustela*, 'a weasel,') *Galeancon*.

MUSTUM, ('juice of the grape,') *Wort*.

MUTACIS'MUS, *Mutacisim*, (*μωτακισμ*, 'I use the letter *μ*, *μν*, too frequently,') A vicious pronunciation; consisting, according to Sauvages, in the frequent repetition of the letters *μ*, *ν*, and *ρ*, which are substituted for others.

MUTA'TIO SEX'US. 'A change or conversion of sex.' A notion prevailed in antiquity that such a conversion was possible.

MUTE, *Mu'tus*, *Dumb*, (F.) *Muet*. Affected with mutitas.

MUTEOS'IS, (*mutus*, 'dumb,') (F.) *Mutéose*. Under this name is comprehended every phenomenon of expression, voluntary and involuntary, which impresses the sight or touch.

MUTILA'TION, *Mutila'tio*, *Maim'ing*, (*mutitus*, 'broken,') The removing or privation of a limb, or of some other external part of the body. In all

countries, this crime has been punished with severity.

MUTINUM, Penis.

MUTISME, Mutitas.

MUTITAS, (from *mutus*, 'dumb,') *Ala'lia*, *Ana'd'ia*, *Dumb'ness*, *Speech'lessness*, *Mut'ism*, (F.) *Mutisme*, *Mutité*. Impossibility of articulating sounds; although they can be elicited. Dumbness is often congenital, and united with deafness, of which it is an effect. In some cases it is accidental.

MUTITAS SURDO'RUM, *Apho'nia Surdorum*, *Surdomut'itas*, *Deaf'lumbness*, *Deaf'mutism*, (F.) *Surdimutité*. Speechlessness from deafness; congenital, or produced during infancy. The subjects of this affection are called *deafdumb*, (F.) *Sourdsmuts*.

MUTITÉ, Mutitas.

MUTO, Penis.

MUTONIATUS, (from *muto*,) *Membrosus*.

MUTTERKORN, Ergot.

MUTTON TEA. Prepared from a pound of mutton, freed from the fat and cut into thin slices; and a pint and a half of boiling soft water poured over them, as in the case of beef tea; except that it requires to be boiled for half an hour after the maceration, before it is strained through a sieve.

MUTUNNUS, (from *muto*,) *Priapus*.

MUTUS, Mute.

MYACANTHA, *Ruscus*.

MYALGIA, (*myon*, and *algos*, 'pain,') Cramp.

MYASTHENIA, (*myon*, and *astheneia*, 'debility,') Muscular debility.

MYCES, (*μυκης*,) Fungus.

MYCETO'MA, (from *myces*, and *oma*,) Fungus disease. An affection of the hands and feet which prevails extensively in India, and is ascribed to the growth of a fungus,—*Chio'nyphe Carteri*, (after Dr. H. V. Carter.)

MYCHMUS, (*μυχμος*,) *Suspirium*.

MYCHTHISMUS, (*μυχθισμος*,) Moaning, *Suspirium*.

MYCODERMA, gen. *Mycoderm'otis*, *My'coderm*, (*myces*, and *derma*, 'skin,') A cryptogamous growth, which constitutes the crusts of favus. See *Porrigo favosa*. Also, a mucous membrane, see *Membrane*.

MYCODERMA CEREVISIÆ, *Torula cerevisiæ*.

MYCODERMA PLI'CAE, *Tricomaph'ytion*. A cryptogamous growth, said to be the essence of *Plica*.

MYCODERMITIS, (*mycoderma*, and *itis*,) *Mucitis*.

MYCODES, (from *myces*, and *odes*,) Fungoid.

MYCIDIARRHŒA, see *Diarrhœa*.

MYCODYSENTERIA, see *Dysentery*.

MYCOPHTHISIS, see *Bronchorrhœa*.

MYCORTHOPNŒA. *Orthopnoea* from excessive secretion of mucus in the air passages—*Orthopnoea pituitosa*.

MYCOSIS, (*myces*, and *osis*,) Fungus.

MYCTER, (*μυκτηρ*,) *Nasus*.

MYCTERES, (*μυκτηρες*,) *Myxæ*, *Nares*.

MYCTEROPHONIA, (*mycter*, and *phōnē*, 'voice,') *Rhinophonia*.

MYCTEROXEROTES, *Mycterozero'sis*, (*mycter*, and *ξηρότης*, 'dryness,') *Narium sic'citas*. Dryness of the interior of the nose.

MYCUS, (*μυκης*,) *Mucus*.

MYDE'SIS, (*μυδαι*, 'I abound with moisture, I am putrid,') *Moisture*. In its most general sense it means *corruption*. By some it is used for a mucous discharge from inflamed eyelids.

MY'DON, (*μυδων*,) Fungous or putrid flesh in certain fistulous ulcers. Also, flesh putrid from mouldiness.

MYDRIASIS, *Platyco'ria*, *Platyco'ria'sis*, *Mydri'asia*. A name given by several writers to morbid dilatation of the pupil—*Coramydri'asia* and by others to weakness of sight produced by

hydropthalmia, or superabundance of humours, (from *μυδος*, 'moisture,')

MYDRIATICUS, *Mydriat'ic*. Relating to or causing dilatation of the pupil. Certain narcotics, as opium, contract the pupil in poisonous doses; whilst others, as belladonna, dilate it. Owing to this antagonism, the mydriatic belladonna has been given in poisoning from opium.

MYELAL'GIA, (*myelus*, and *algos*, 'pain,') Pain in the spinal marrow.

MYELANALOSIS, (*myelus*, and *analōsis*, 'wasting,') *Tabes dorsalis*.

MYELAPOPLEXIA, *Apoplexia myelitica*.

MYELATEL'IA, (*myelus*, and *areleia*, (a, privative, and *telos*, 'end,') 'want of end or finish,') Incompleteness of the spinal marrow.

MYELATROPHIA, (*myelus*, and *atrophia*, 'atrophy,') *Tabes dorsalis*.

MYELIN, (from *myelus*,) see *Nerve fibres*.

MYÉLITE, *Myelitis*.

MYELITIS, *M. parenchymato'sa seu spina'lis*, *Inflammat'io medullæ spina'lis*, *Spini'tis*, *Spino-dorsi'tis*, *Notomyeli'tis*, *Rachiomylei'tis*, *Rachial-gi'tis* (of some), (F.) *Myélite*, *Inflammation de la moëlle épinière*, *I. rachidième*, (*myelus*, and *itis*,) Inflammation of the spinal marrow or its membranes;—indicated by deep-seated burning pain in the spine, with various nervous and vascular irregularities of function. It is not common. Dr. Marshall Hall proposes to call inflammation of the membranes of the brain *Meningi'tis*; that of the substance of the brain *Myeli'tis*.

MYELITIS CONVULSIVA, *Chorea*, electric—*m. Parenchymatosa*, *Myelitis*—*m. Petechialis*, *Meningitis*, *cerebrospinal*—*m. Spinalis*, *Myelitis*—*m. Exsudativa*, *Hydrorachis*.

MYELOCHYSIS, (*myelus*, and *χω*, 'to pour out,') *Hydrorachis*.

MYELOCON, (F.) (*myelus*, and *κονια*, 'dust,') A mixture of fatty matters found in the substance of the brain. (Couverbe.)

MYELOCYTES, (*myelus*, and *κυτος*, 'a mass, body, or cell,') Granules of the brain. Nuclei of the cells of the gray substance.

MYELOGANGLIITIS, (*myelus*, *γαγγλιον*, 'a ganglion,') and *itis*,) *Cholera* (sporadic).

MYELOID, *Myelo'id'es*, *Myelo'd'es*, (*myelus*, and *ειδς*, 'resemblance,') Marrowlike.

MYELOID TU'MOUR, *Fibroplast'ic tumour*. A sarcomatous tumour, the more characteristic constituents of which are—peculiar many-nucleated corpuscles, which have been recognized as constituents of the marrow and diploe of bones, especially in the fœtus and in early life.

MYELOMA, gen. *Myelo'matis*, (from *myelus*,) See *Encephaloid*.

MYELOMALACIA, (*myelus*, and *μαλακία*, 'softness,') *Mollities medullæ spinalis*.

MYELOMENINGITIS, (*myelus*, *μηνιγξ*, 'a membrane,') and *itis*,) *Meningitis spinal*.

MYELOMYCES, (*myelus*, and *myces*,) see *Encephaloid*.

MYELON, (from *myelus*,) *Medulla spinalis*.

MYELOPARALYSIS, (*myelus*, and *paralysis*,) *Paraplegia*.

MYELOPHTHISIS, *Tabes dorsalis*—*m. Sicca*, *Tabes mesenterica*.

MYELOPHY'MATA, *Tuber'cula medullæ spina'lis*, (*myelus*, and *φυμα*, 'a tumor,') Tubercles of the spinal marrow.

MYÉLOPLAXE, (*myelus*, and *πλαξ*, 'a plate or lamella,') Plates or lamellæ with numerous nuclei in the marrow of bones. (Robin.)

MYELORRHAGIA, (*myelus*, and *ῥηγνυμι*, 'to break forth,') *Apoplexia myelitica*.

MYELOSPONGUS, (*myelus*, and *σπογγος*, 'a sponge, a fungus,') See *Encephaloid*.

MYELUS, (*μυελος*,) *Marrow*, *Medulla spinalis*—*m. Diauchenius*, *Medulla spinalis*—*m. Dorsites*,

Medulla spinalis.—m. Notieus, Medulla spinalis—m. Psoties, Medulla spinalis.

MYENERGIA, (*myon*, and *energia*.) 'Muscular strength.'

MYGMUS, (*μυγμός*.) 'Suspirium.'

MYIOCEPHALUM, (*μυια*, 'a fly,' and *κεφαλή*, 'head,' from its resemblance.) Ptoxis iridis.

MYIODEOPSIA, (*μυιωδής*, 'appertaining to flies,' and *ὄψις*, 'vision,') see Metamorphopsia.

MYIODOPSIA, Metamorphopsia.

MYITIS, (*myo*, and *itis*.) Myositis, Rheumatism, acute.

MYLABRIS CICHORII, (*μυλαβρις*.) see Cantharis.—m. Punctata, see Cantharis.—m. Pustulata, see Cantharis.

MYLACEPHALUS, (*μυλή*, 'a mole,' a, privative, and *κεφαλή*, 'head,') A monster with only traces of viscera, and without head or extremities.

MYLACRI, (*μυλακροι*, from *μυλή*, 'a millstone,')

Molar teeth.

MYLACRIS, Patella.

MYLE, (*μυλή*, 'a millstone,') Patella, Mole.

MYLICUS, Molar.

MYLO, (from *μυλή*, 'a mill,') In composition, the jaw, especially the lower jaw.

MYLODI, (*μυλή*, 'a mill,' and *ὄδον*, 'a tooth,')

Molar teeth.

MYLODONTES, Molar teeth.

MYLOGLOSSUS. Winslow has given this name to muscular fibres which pass from the posterior part of the myloid line of the lower jaw, and from the sides of the base of the tongue to the parietes of the pharynx. These fibres belong to the constrictor superior pharyngis.

MYLOHYOID FURROW or GROOVE. A furrow which passes from the superior orifice of the inferior dental canal of the lower jaw in the same direction as the canal, and lodges the mylohyoid nerve, a branch of the inferior-dental.

MYLOHYOID LINE, see Mylohyoideus.

MYLOHYOID NERVE, see Mylohyoid furrow.

MYLOHYOIDEUS, *Mylojides*, *Diaphragma oris*, (F.) *Mylo-hyoidien*. The *Mylopharyngeus* of Morgagni and Santorini. This muscle is situate at the upper and anterior part of the neck, behind the lower jaw. It is broad, flat, and has the form of a truncated triangle. It arises from the internal oblique, internal maxillary, or mylohyoid line of the lower jaw bone, and is inserted at the fore part of the body of the os hyoideus. Its innermost fibres unite with those of the opposite side to form a raphe on the median line. The mylohyoideus raises the os hyoideus and carries it forward, or it depresses the lower jaw.

MYLOPHARYNGEUS, Constrictor pharyngis, Mylo-hyoideus.

MYLOÏDES, Mylohyoideus.

MYOCARDITIS, (*myon*, and *carditis*.) Carditis.

MYOCLE, *Myoclele*, (*myon*, and *κλήη*, 'a tumor,')

A muscular tumor.

MYOCEPHALITIS, *Myotitis cephalica*, (*myon*, *κεφαλή*, 'head,' and *itis*.) Inflammation of the muscles of the head.

MYOCEPHALON, (*μυια*, 'a fly,' and *κεφαλή*, 'head,' from its resemblance.) Ptoxis iridis.

MYOCELIALGIA, (*myon*, *κοιλία*, 'abdomen,' and *αλγος*, 'I suffer,') Pain in the muscles of the abdomen.

MYOCELITIS, *Myocælititis*, *Inflammatio musculo-rum abdominalium*, (*myon*, *κοιλία*, 'lower belly,' and *itis*.) Inflammation of the muscles of the abdomen.

MYODEMIA, (*myon*, and *ἔμπος*, 'fat,') Fatty degeneration of muscle.

MYODEOPSIA, (*μυιωδής*, 'appertaining to flies,' and *ὄψις*, 'vision,') Metamorphopsia.

MYODES, (*myon*, and *odes*.) Like unto muscle. Muscular.

MYODESOPIA, (*μυιωδής*, 'appertaining to flies,' and *ὄψις*, 'vision,') Metamorphopsia.

MYODESOPSIA, Metamorphopsia.

MYODYNAMIA, *Vis muscularis*, (*myon*, and *δυναμις*, 'power,') The force or power of a muscle.

MYODYNAMICS, see Muscular contraction.

MYODYNAMIGETER, (*myodynamia*, and *μετрон*, 'measure,') Dynamometer.

MYODYNAMOMETER, Dynamometer.

MYODYN'IA, (*myon*, and *ὄδυνη*, 'pain,') Pain in the muscles,—rheumatismal or other. Rheumatism.

MYODYNIA INFLAMMATORIA, Myositis.

MYOGRAPHY, *Myographia*, (*myon*, and *γραφειν*, 'to describe,') An anatomical description of the muscles.

MY'OID, *Myoides*, (*myon*, and *εἶδος*, 'resemblance,') An epithet given to tumours composed of fibre cells or muscular fibres of organic life.

MYOLEMMA, gen. *Myolem'matis*, *Myole'ma*, (*myon*, and *λεμμα*, 'a coat,') The membranous tube of each muscular fibre. It has also the same signification as *Sarcolemma*.

MY'OLIN, (from *myon*.) A name given by Mr. Erasmus Wilson to a transparent substance that fills the cells, which, he conceives, by their juxtaposition form the ultimate muscular fibril.

MYOL'OGY, *Myolog'ia*, *Sarcolog'ia*, *Myoiolog'ia*, (*myon*, and *λογος*, 'a discourse,') That part of anatomy which treats of the muscles.

MYOMALACIA, (*myon*, and *μαλακία*, 'softness,') Softening of muscle.

MYON, (*μυων*.) Muscle.

MYONARCO'SIS, (*myon*, and *ναρκη*, 'stupor,') Numbness of the muscles.

MYONITIS, (*myon*, and *itis*.) Myositis.

MYON'OSUS, *Myon'us*, *Myopathia*, (*myon*, and *ὄσος*, 'disease,') A disease of the muscles.

MYOPALMUS, (*myon*, and *παλμος*, 'agitation,') Subsultus tendinum.

MYOPARALYSIS, see Paralysis.

MYOPATHIA, (*myon*, and *πάθος*, 'affection,') Myonosis.

MYOPE, Myops.

MYOPHON'IA, (*myon*, and *φωνή*, 'voice,') The sound of muscular contraction,—as that of the ventricles of the heart during the systole of that organ.

MYO'PIA, *Myopi'asis*, *Lusciō'itas*, *Myo'sis*, *Parop'sis propin'qua*, *Ambly'opia dissito-rum*, *Dyso'pia dissito-rum*, *Dyspho'tia*, *Visu' juvenum*, (*μυς*, 'a mouse,' and *ὤψ*, 'the eye,') *My'opy*, *Short-sightedness*, *Nearsightedness*, *Mousesight*, *Farblindness*, (Old E.) *Mo-pe-eye*, *Moap-eye*, (F.) *Vue courte*, *V. basse*. The state of those who can only see objects very near. The defect is owing to the too great convexity or depth of the eye, or to too great density of the humours, and is palliated by wearing concave glasses.

MYO'PIC, *Myop'icus*, *Mo-pe-eyed*, *Mop'sical*, *Shortsighted*, *Por'blind*, *Pur'blind*, *Spur'blind*. Relating or appertaining to myopia.

MYOPODIO'RTHO'TICON, (*myopia*, and *διορτωτικον*, 'having power to correct,') An apparatus for the cure of shortsightedness.

MY'O'PS, *Lusciō'sus*, *Myo'pus*, *My'ope*. One affected with myopia or shortsightedness.

MYOPY, Myopia.

MYORRHEN'IS, (*myon*, and *ρηξις*, 'rupture,')

Rupture of a muscle.

MYOSARCO'MA, gen. *Myosarco'matis*, (*myon*, and *sarcoma*.) A sarcomatous tumour in which muscular fibres are found.

MYO'SIS, (*μυω*, 'I close the eyes,') *Microcor'ia*. Smallness of the pupil. *Phthi'sis pupillari'a*. Permanent contraction of the pupil. It is usually caused by iritis, and is extremely difficult to cure. When it exists to such an extent as to obliterate the pupil, it is called *Synize'is*. Also, *Myopia*.

MYOSITE, *Myositis*.

MYOSITIC, *Myositicus*, from *Myosis*. Causing contraction of the pupil,—as opium.

MYOSITIS, *Myitis*, *Mysitis*, *Myonitis*, *Sarcitis*, *Rheumatismus phlegmonodes*, *Myodynitis inflammatoria*, *Phlegmatis myoica*, *Inflammatio seu Phlegmonè Musculorum*, (F.) *Myosite*, *Inflammation des muscles*, (*myon*, and *itis*.) A name proposed by Sagar for inflammation of the muscles. Also, Rheumatism.

MYOSOTIS, (μυσοῦσις, from μυς, gen. ὠς, 'a mouse,' and οὖς, gen. ὠς, 'ear.') *Hieracium pilosella*, Rheumatism, Rheumatism, acute; (from *myon*.)

MYOSPASMUS, Cramp.

MYOSITICI (*Morbi*), (*myon*, and ὀστέον, 'a bone.') Diseases of the bones and muscles, in the classification of Dr. Wm. Farr.

MYOTILITY, *Myotilitas*, (F.) *Myotilité*, (from *myon*.) Muscular contractility.

MYOTOMY, *Myotomia*, (*myon*, and τέμνειν, 'to cut'.) The part of practical anatomy which treats of the dissection of the muscles. Also, the surgical operation of the division of muscles to remove deformity.

MYOTYRBE, (*myon*, and τυρβη, 'disorder.') Chorea.

MYRACOPUM, (μυρον, 'an ointment,' a, privative, and κόπος, 'fatigue'.) An ointment used by the ancients in cases of fatigue.

MYRCIA ACRIS, see *Spiritus myrciæ*.

MYREPSUS, (μυρπος, from μυρον, 'an ointment'.) Unguentarius.

MYRIAGRAMME, (μυρία, '10,000,' and γραμμα, 'gramme'.) A weight equal to 10,000 grammes, or to 26 pounds, 9 ounces, and 6 drachms Troy.

MYRIAMÈTRE. A measure of 10,000 metres; equal to 16 miles, 1 furlong, 156 yards, and 14 inches.

MYRICA ASPLENIFOLIA, (after μυρική, 'the tannarisk'.) *Comptonia asplenifolia*—m. *Cenifera*, see *Cera flava et alba*, and Wax, myrtle.

MYRICA GAL'E, *M. palustris*, *Myrtus Brabantica* seu *Anglica*, *Myrtifolia Belgica*, *Gal'e*, *Ga'gel*, *Rhus sylvestris* seu *Belgica*, *Ac'aron*, *Elæagnus*, *E. Cor'do*, *Chamaelæagnus*, *Dodonæa*, *Dutch myrtle*, *Sweet gale*, *Sweet willow*, (Sc.) *Scotch gale*, (F.) *Piment Royal*, *Thé de Simon Pauli*, *Gale odorant*. Ord. Myricaceæ. Sex. Syst. Dicoecia Tetandria. The leaves, flowers, and seeds have a strong, fragrant smell, and a bitter taste. They are used for destroying moths and cutaneous insects. The infusion is given internally as a stomachic and vermifuge.

MYRICA PALUSTRIS, M. Gale.

MYRINGA, see Tympanum.

MYRINGITIS, *Inflammatio tympani*, (*myrin'ga*, 'the membrana tympani,' and *itis*.) Inflammation of the membrana tympani; and, also, of the tympanum.

MYRINX, gen. *Myrin'gis*, see Tympanum.

MYRIS, (μυρίς, *Myrothe'ce*, *Narthe'cia*, *Narthe'cium*, *Nar'theux*, (μυρον, 'a perfumed oil or ointment'.) A perfumed oil jar or ointment box.

MYRIS'TICA, (μυρον, 'a liquid perfume,' from μυρον, 'to flow,') *M. Moscha'ta* seu *aromat'ica* seu *officinalis* seu *fra'grans*, *Comacon*. Nat. Ord. Myristicæ. The tree which produces the nutmeg and mace. (F.) *Muscadier*. The *Nutmeg*, *Myristica*, (Ph. U. S.), *Myristicæ* seu *Myristicæ moscha'tæ nucleus*, *Nux moscha'ta* seu *unguentaria*, *Nucis'ta*, *Nux myristica* seu *aromat'ica*, *Chrysobalanus Gale'ni*, *Unguentaria*, *As'sala*, (F.) *Muscade*, is the kernel of the fruit. It has a fragrant, aromatic odour; an agreeable pungent taste, and is much used for culinary purposes. Alcohol extracts its active matter. It has the properties of aromatics in general; being stimulant and stomachic. The volatile oil—*O'leum myristicæ* seu *nucis'tæ ether'eum*—possesses the

virtues of the nutmeg. Dose of the nutmeg, gr. v to ℥j;—of the oil, gtt. iij to gtt. vj.

Mace, *Ma'cis*, *Flores macis* seu *ma'cidis*, (F.) *Fleurs de muscade*, is the arillus of the fruit. It is membranous, with the odour and taste of the nutmeg, and is possessed of similar qualities. The *O'leum macis* seu *ma'cidis* is a fragrant, sebaceous substance, expressed in the East Indies. It is only used externally. It is the *Adeps myristicæ*, *Myristicæ o'leum express'um*, *O'leum* seu *Balsamum nucis'tæ*, *But'ter of nut'megs*, *Express'ed oil of Mace*.

MYRISTICÆ NUCLEUS, see *Myristica moschata*.

MYRISTICATIO HEPATIS, Liver, nutmeg.

MYRMECIA, (μυρμηκία,) Formica.

MYRMECIASIS, (μυρμηκiasis,) Convulsio cerealis, Formication.

MYRMECIASMUS, (μυρμηκiasmos,) Convulsio cerealis, Formication.

MYRMECISMUS, Convulsio cerealis, Formication.

MYRMECIZON, (μυρμηκίζων,) Formicant.

MYRMECOSIS, Convulsio cerealis, Formication.

MYRMEX, (μυρμηξ,) Formica.

MYROBALANUS, (*myron*, and βαλανος, 'a nut'; so called, because formerly used in ointments;) *Angeloc'acos*, *Myro'bolan*, *Glans unguentaria*, *Pal'ma unguentario'rum*. A dried Indian fruit, of the plum kind, of different species of *Terminalia*. Ord. Combretaceæ. Of this there are several varieties—the *M. Bellir'ica* or *Bellir'ic myro'balan*, *Bellegu*, *Bellerigi*, *Belmleg*—the *M. cheb'ula* or *Chebule myro'balan*; the *M. cit'rina* or *Yell'ow myrobalan*, *Ara'ra*; the *M. Emb'lica* or *em'blic myrobalan*, and the *M. In'dica*, *As'nar*, *In'dian* or *Black myro'balan*. All the myrobals have an unpleasant, bitterish, very austere taste; and strike an inky blackness with a solution of steel. They are said to possess laxative as well as astringent properties.

MYROLES, (*myron* and *oleum*,) see *Olea volatilia*.

MYRON, (μυρον,) Myrum, Unguentum.

MYROPISSOCE'RON. A topical application in alopecia, referred to by Galen; (*myron*, πισσα, 'pitch,' and κηρος, 'wax'.)

MYROPÆUS, (*myron*, and ποιω, 'to make,') Unguentarius.

MYROPOLES, (μυροπωλης, from *myron*, and πωλω, 'to sell,') Unguentarius.

MYROSPERMUM FRUTESCENS, (*myron*, and σπέρμα, 'seed,') *Myroxylum Peruiferum*—m. *Pereire*, see *Myroxylon Peruiferum*—m. *Peruiferum*, see *Myroxylon Peruiferum*—m. of *Sonsonate*, see *Myroxylon Peruiferum*—m. *Toluiferum*, see *Toluifera Balsamum*.

MYROXYLON, (*myron*, and ξύλον, 'wood,') see *M. Peruiferum*.

MYROX'YLON PERUIF'ERUM, *Myrosperm'um frutescens* seu *Peruif'erum*, *Quinquino*, *Caburei'ba*. Nat. Ord. Leguminosæ. Supposed to be the tree which affords the *Peruvian bal'sam*, *Bal'samum Peruvianum* (Ph. U. S.) seu *Peruvianum nigrum* seu *Peruanum* seu *In'dicum*, *Putzochill*, *Myrox'yl' Peruif'eri bal'samum*, *Myrox'ylon*, *Cabureiciba*, *In'dian*, *Mex'ican*, or *American bal'sam*, (F.) *Baume de Pérou ou des Indes*. This balsam consists of benzoic acid, resin, and essential oil. Its odour is fragrant and aromatic; taste hot and bitter: it is soluble in alcohol, and miscible in water by the aid of mucilage. It is stimulant and tonic, and considered to be expectorant: as such, it has been employed in paralysis, chronic asthma, chronic bronchitis, rheumatism, gleet, leucorrhœa, &c., and externally, for cleansing and stimulating foul indolent ulcers. Dose, gtt. v to gtt. xxx.

White Bal'sam of Peru', *Nat'ural bal'sam*, *Bal'-*

samum al'ba, *Sty'rax al'ba*, *Balsamel'on*, is obtained by incision from *Myrospermum peruiferum*.

The Myroxylon of Sonsonate in Central America—*Myrospermum Peréire* of Dr. Royle, according to the Ph. B. (1864), furnishes balsam of Peru.

MYROXYLON OF SONSONATE, see *M. Peruiferum*—*m. Toluiferum*, *Toluifera Balsamum*.

MYRRHA, (*μυρρά*), *Bo'la*, *Stac'te*, *Myrrh*, *Calo'nia*, *Smyr'na*, *My'rha ru'bra*, (F.) *Myrrhe*. The exudation (*μυρον*, from *μυρνω*, 'to flow') of a plant of Abyssinia or Arabia Felix, the *Balsamodendron myrrha*. This gum-resin has a fragrant, peculiar odour; and bitter aromatic taste. It is in reddishyellow, light, brittle, irregular tears; partially soluble in distilled water when aided by friction. S. g. 1.360. It is stimulant; and has been used in cachectic affections, humoral asthma, chronic bronchitis, &c. Dose, gr. x. to ʒi.

MYRRHA IMPERFECTA, *Bellium*.

MYRRHINE, (*μυρρην*), *Myrtus*.

MYRRHIS ANNUA, (*μυρρίς*), *Athamanta Crentensis*—*m.* Major, *Cherophyllum odoratum*—*m.* *Odorata*, *Cherophyllum odoratum*.

MYRSINA AFRICANA, *Tatzé*.

MYRSINE, (*μυρσιν*), *Myrtus*.

MYRSINELÆ'ON, (*μυρσιν*, 'myrtle,' and *λαῖον*, 'oil,') Oil of myrtle.

MYRSINTES, (*μυρσιντες*.) Wine in which branches of myrtle have been macerated.

MYRTACANTHA, (*myrtus*, and *ακανθα*, 'a thorn,') *Ruscus*.

MYRTE COMMUN, *Myrtus*.

MYRTID'ANON, (*μυρτιδανον*, from *myrtus*, 'myrtle,') An excrescence growing on the trunk of the myrtle, and used as an astringent. With Hippocrates, either the *myrtus caryophyllata* or *myrtus pimenta*. [?] Also, a wine—*Vinum Myrtid'anum*—made from wild myrtle berries.

MYRTIFOLIA BELGICA, (*myrtus*, and *folium*, 'a leaf,') *Myrica gale*.

MYRTIFORM, *Myrtiformis*, *Myrtodes*, (*myrtus*, and *forma*, 'shape,') Having the shape of a leaf of myrtle. A name given to a muscle (depressor *ale nasi*), to the fossa incisiva; to certain caruncles, &c.

MYRTIFORMIS, *Compressor naris*, *Depressor ale nasi*.

MYRTITES, (*μυρτιτης*, from *μυρτον*, 'the myrtle berry,') A name given to a medicine prepared with honey and myrtle berries.

MYRTLE, *Myrtus*—*m.* Berry, *Vaccinium myrtillus*—*m.* Candleberry, see *Wax*, myrtle—*m.* Dutch, *Myrica gale*—*m.* Wax, see *Wax*, myrtle—*m.* Wild, *Ruscus*.

MYRTOCHEILIDES, (*myrton*, and *χειλις*, 'a small lip,') *Nymphæ*.

MYRTOCHILA, (*myrton*, and *χειλος*, 'a lip,') *Nymphæ*.

MYRTODES, (*myrtus*, and *odes*), *Myrtiform*.

MYRTON, (*μυρτον*), *Clitoris*.

MYRTUS, (*μυρτος*), *Myrtus communis* seu *communis Italica*, *Myrsinè*, *Myrrhinè*, *The Myrtle*, (F.) *Myrte Commun*. Ord. *Myrtacæ*. The berries of this plant have been recommended in alvine and uterine fluxes and other disorders of relaxation and debility. They are moderately astringent and somewhat aromatic.

MYRTUS ACIS, see *Spiritus myrciæ*—*m.* Anglica, *Myrica gale*—*m.* Brabantica, *Myrica gale*—*m.* *Caryophyllata*, see *Spiritus myriciæ*.

MYRTUS CARYOPHYLLATA, *Calyptanthus caryophyllata*, *Cassia* seu *Canel'la caryophyllata*, (F.) *Capulet*, *Cannelle giroflée*. The tree which is considered to afford the *Clove bark*; *Cortex caryophyllatus* seu *caryophyllata*. This bark is

a warm aromatic; resembling clove with an admixture of cinnamon. It may be used with the same views as cloves or cinnamon.

MYRTUS CARYOPHYLLUS, *Eugenia caryophyllata*—*m.* *Leucodendron*, *Melaleuca cajuputi*.

MYRTUS PIMENTA, *Eugenia Pimenta*. The tree which bears the *Jamaica pepper*, *Pimentæ bac'ca*, *Pimentæ ber'ries*, *Pimentæ*, *Pi'per caryophyllatum* seu *chia'pæ* seu *odoratum Jamaicensè* seu *Jamaicensè* seu *tabas'cum*, *Coc'culi Ind'i aromatici*, *Amomum pimenta*, *Fructus pimentæ*, *Carive*, *Caryophyllus Americanus* seu *Pimenta*, *Allspice*, (F.) *Poivre de Jamaïque*, *Toute épice*, *Assouvi*. The unripe berries—*Pimenta*, *Pimentæ* (Ph. U. S.)—have an aromatic odour; resembling a mixture of cinnamon, nutmeg, and cloves; the taste is pungent, but mixed, like the odour. Like other peppers, this is stimulant and carminative. The oil—*Oleum Pimentæ*—possesses the virtues of the berries. The powdered fruit has been called *Quatre épices* or *Four épices*. Dose, gr. v to ʒij.

Aqua Pimentæ, *Pimentæ water*, is official in the London, Edinburgh, and Dublin Pharmacopœias. It is carminative. Dose, f ʒij or f ʒij.

MYRUM, (*μυρον*), *My'ron*. A perfumed oil or ointment. A liquid perfume. The spontaneously exuding juice of many plants, especially of that from which myrrh is obtained.

MYS, (*μῦς*.) *Muscle*.

MYSIOLOGIA, (*mys*, and *λογος*, 'a discourse,') *Myology*.

MYSITIS, (*mys*, and *itis*, 'inflammation,') *Myositis*.

MYS'TAX, gen. *Mys'taxis*, *Mastax*, (*μυσταξ*, gen. *μυσταξες*.) The hair growing on each side of the upper lip in men. The *mustache* or *mustachio*, (Old. Eng.) *Munchat'oe*, (F.) *Moustache*. Also, the upper lip.

MYSTERION, (*μυστηριον*, 'a mystery,') An arcanum, nostrum, or secret preparation in general. Also, an antidote, referred to by Galen and others.

MYSTRON, (*μυστρον*.) A Greek measure, which held about three drachms.

MYTILUS EDULIS, (*μυtilος*, from *mys*.) The common mussel, (F.) *Moule*. A bivalve, the flesh of which, when at all in a state of decomposition, is highly poisonous. When fresh, it is an agreeable, but not very digestible article of diet, producing, in some persons, all the symptoms of urticaria. One affected with such phenomena is said, occasionally, to be *muscalled*.

MYU'RUS, *Mei'rus*. A pulse is so called when it sinks progressively and becomes smaller and smaller like a mouse's tail; (from *μῦς*, 'a mouse,' and *οὐρα*, 'tail,') *Pulsus myurus reciprocus*, a pulse, which, after having become gradually weaker, resumes, by degrees, its former character.

MYXA, (*μύξα*), *Mucus*, *Sebestina*.

MYXÆ, (*μύξαι*), *Nares*.

MYXIOSIS, (from *μύξα*, 'mucus, and *osis*,') *Gonorrhœa impura*.

MYXOIDES, (from *myxa*, 'mucus,' and *ειδος*, 'resemblance,') *Muciform*, *Mucous*.

MYXOR'HOOS, (*μυχορροος*, from *μύξα*, 'mucus,' and *ῥεω*, 'I flow,') One who is subject to mucous discharge. Applied to an infant that discharges a considerable quantity of mucus and saliva.

MYXOSARCO'MA, gen. *Myxosarco'matis*, (*μύξα*, 'mucus,' and *σάρξ*, gen. *σαρκος*, 'flesh,') A tumour which is *mu'cocar'neous*; partly mucous and partly fleshy.

MYXOTER, (*μύξητηρ*, from *μύξα*, 'mucus,') *Nasus*.

MYZESIS, (*μύζω*, 'to suck,') *Sucking*.

N.

N. This letter, in prescriptions, is an abridgment of *numero*, 'by number.'

NABULUS ALBUS, (from *ναββα*, 'a harp,' owing to the lyrate leaves,) *Prenanthes alba* seu *serpentaria*, White Lettuce, Lion's Foot, Rattlesnake's Master, Rattlesnake root, Gall of the Earth. An indigenous plant, of the order *Compositæ*, which, with several other species of the genus, is reputed to possess the power of curing the bites of serpents. The root has been used in dysentery.

NABULUS FRASERI has similar properties.

NABOT, *Saccharum candidum*.

NABOTHI GLANDULÆ, *Ovula* seu *Ova Nabothi* seu *Nabothia'na*, *Ovaria Nabothi*, *Folliculi rotundi et oblongi*, *Vesiculae* seu *Bullae rotundae cervicis uteri*, *V. seminales mulierum*, *Corpora* seu *Corpuscula globosa*, *Glandulae* seu *Hydatides cervicis uteri*, *Glands* of *Naboth*, (F.) *Oufs* ou *Vésicules* de *Naboth*. Small, mucous crypts or follicles situate in the interstices of the duplicatures of the lining membrane of the cervix uteri. Naboth, from noticing them in a morbid condition, mistook them for ova; and hence they received the name *Ovula Nabothi*.

NACRA, *Nakra*, *Nasa*. A kind of influenza common in the East Indies.

NACRÆ, ([F.] *naere*, 'mother of pearl,') *Margaritaceus*.

NACREOUS, *Margaritaceus*.

NÆVI, see *Nævus*—n. Pigmentary, see *Nævus*.

NÆVOID, *Nævoides*, (*nævus*, and *ειδος*, 'resemblance,') Resembling *nævus*.

NÆVUS, pl. *Nævi*; *Nævus mater-nus* seu *Sigillum*, *No'ta mater-ni*, *Nævus*, *Macula matricis* seu *mater-næ* seu *matric'al'es*, *No'tæ infantum*, *Lupus varicosus*, *Stigmata*, *Metrocel'ides*, *Congen'itæ no'tæ*, *Mother's* or *Fairy* marks, *Mother's spots*, (F.) *Envie*, *Tache* de naissance ou pigmentaire, *Signe*. Spots of various kinds on the skin of children when born, which have been attributed to the influence of the maternal imagination on the fœtus in utero. They are of various appearances, some much better supplied with blood than others. Some are merely superficial or stain-like spots, *Pigmentary nævi*; others are prominent, *Vascular tumours*, and often have long, irregular hairs growing from them. These have usually been called *Moles*, *Spili*, *Spilomata*, *Epichor'æes spili*, &c. The red, rosy, violet or bluish spots are called by I. G. St. Hilaire, *Taches sanguines*;—those of an intermediate hue, between the normal colour and black, *Taches melaniennes*. See *Telangiectasia*.

When *nævi* are superficial, without any disposition to enlarge or spread, they need not be meddled with; but all those that partake of the character of aneurism by anastomosis had better be removed, where practicable.

NÆVUS SIGILLUM, *Nævus*.

NAFDA, *Naphtha*.

NAFTA, *Naphtha*.

NAIL. (Sax. *nægl*.) *Un'guis*, *O'nyx*, (F.) *Ongle*.

A whitish transparent substance, similar to horn, which covers the dorsal extremity of the fingers. Three portions are distinguished in it; the *extremity*, which is free, at the end of the finger;—the *body* or middle portion adherent by its inner surface; and the *root*, *O'nychorrh'iza*, *Rhizonych'ia*, *Rad'ix* seu *Ma'trix un'guis*, *In'timum un'guis*, (F.) *Matrice de l'ongle*. The last presents two distinct parts; the one, terminated by a thin, serrated edge, is buried in a duplicature of the skin, *Vall'ecula un'guis*, which acts the part of a *naill'icicle*; the other, called *Lu'nula* seu *Semi-lu'nula* seu *Sel'ē* seu *Ar'cus* seu *Exor'tus* seu

Anat'olē seu *Albe'do un'guum*, is whitish and of a semilunar shape, and is situate above the part where the epidermis terminates. The nails are composed of a horny tissue, of the same nature as that which forms the hoofs, horns, and scales of different animals. When the nail has been torn off, the papillæ of the skin become covered by a soft, whitish lamina, whose consistence gradually augments. New laminae are then formed underneath, and give the nail the thickness it ought to possess. The corneous substance, being thus constantly produced at the extremity of each of these laminae, the whole of the nail is pushed forwards, and it would grow indefinitely, were it not cut or worn by friction.

The nails protect and support the extremities of the fingers against the impression of hard bodies. They are, also, useful in laying hold of small bodies; and dividing those that have but little consistence.

NAILFOLLICLE, see *Nail*.

NAILSPRING, *Hangnail*.

NAIN, *Nanus*.

NAISSANCE, (from *nascor*, *natus*, 'to be born,') see *Parturition*.

NAJA TRIPUDIANS, *Cobra di Capello*.

NAKRA, *Nacra*.

NANGNAIL, *Hangnail*.

NANISME, see *Nanus*.

NANNARI, *Hemidesmus Indicus*.

NANNYBERRY, *Viburnum lentago*.

NANOCEPHALIA, (*nanus*, and *κεφαλή*, 'head,') see *Nanocephalus*.

NANOCEPHALUS. One of a monstrous state—*Nanocephalia*—in which the whole head or certain of its parts are too small, whilst the trunk and extremities are normal.

NANOCORMIA, see *Nanocormus*.

NANOCORMUS, (*nanus*, and *κορμος*, 'a trunk,') One of a monstrous condition—*Nanocormia*—in which the trunk is too small, whilst the head possesses its normal size.

NANOMELIA, see *Nanomelus*.

NANOMELUS, (*nanus*, and *μελος*, 'a limb,') One of a monstrous condition—*Nanomelia*—in which some part of an extremity is too small, and the whole limb too short.

NANOSOMIA, (*nanus*, and *σωμα*, 'body,') *Microsomnia*, *Microsoma'tia*, *Dwarf'ishness*, (F.) *Nanisme*. A state in which the entire body with all its parts is smaller than common.

NANOSOMUS, *Nanus*.

NA'NUS, (*νανος*;) *Pumilio*, *Pumilio*, *Pusill'us*, 'a dwarf.' *Nanosom'us*, (Prov.) *Crut*, *Rope*, *Spin'nick*, (Sc.) *Croil*, *Crok*, *Dearch*, *Dersch*, *Droich*, *Wrat'ack*, (*Pumilio*, *Pumilio*, 'dwarfish,') (F.) *Nain*. One who is much below the usual stature.

The term is applicable to all organized beings from man to the vegetable. Trees have their dwarfs, as the human species have theirs. Some curious authenticated instances of human dwarfs are on record. Bebe, the dwarf of Stanislaus, king of Poland, was 33 inches (French) long, and well proportioned. The Polish nobleman, Borwaski, who was well made, clever, and a good linguist, measured 28 Paris inches. He had a brother, 34 inches high, and a sister 21. A Friesland peasant, at 26 years of age, had attained 29 Amsterdam inches. C. H. Stöberin, of Nürnberg, was under three feet high at 20, yet he was well proportioned and possessed of talents. General Tom Thumb, so called, was seen by the Author in 1847. He was said to be 15 years old; measured 28 inches in height, and when weighed at the mint was found to weigh

20 pounds and 2 ounces. Two remarkable Indian dwarfs were exhibited in the United States and in Europe, about the middle of the 19th century, under the name of 'Aztec children,'—the boy, from 7 to 8 years of age, was 33½ inches in height, and weighed 20½ pounds;—the girl, from 4 to 6 years of age, was 29½ inches high, and weighed 17 pounds. They were very deficient in intellect. See Pygmy.

NAPE OF THE NECK, *Nucha*.

NAPELLUS VERUS, (dim. of *napus*, 'a turnip,') *Aconitum napellus*.

NAPHÆ FLORES, see *Citrus aurantium*.

NAPHTHA, *Acetone*.

NAPHTHA, (*ναφθα*,) *Naf'da*, *Naf'ta*, *Nap'ta*, *Nap'ta'lis*, *O'leum pet'ree al'bum*; (*ἡ*) *Naphthe*; from a Chaldaic and Syriac word signifying bitumen. A bituminous substance, found in Persia, Calabria, Sicily, &c. It is liquid, limpid, of a yellowishwhite colour, a smell slightly resembling that of oil of turpentine, and lighter than water. It resembles petroleum in its properties, and has been chiefly used as an external application; although occasionally, as an anthelmintic, and in inhalation in phthisis pulmonalis. Also, *Æther*.

NAPHTHA ACETI, *Æther aceticus*—*n*. Coal Tar, light, see *Anæsthetic*—*n*. Sulphuricæ, *Æther sulphuricus*—*n*. Vitrioli, *Æther sulphuricus*—*n*. Vitrioli camphorata, *Tinctura atheræ camphorata*—*n*. Vitrioli martialis, *Tinctura seu Alcohol sulphuricoæthereus ferri*—*n*. Wood, *Acetone*.

NAPH'THALIN, *Naphthalina*. When coal tar is subjected to distillation, naphthaline passes over after coal naphtha. It is a white, shining, concrete, crystalline substance, fusible at 176° and boiling at 423°. It is soluble in alcohol, ether, naphtha and the oils, but insoluble in water. It has been used as an excitant expectorant, in the dose of 8 to 30 grains, in emulsion or syrup, and repeated. It has also been used as an anthelmintic; and, when made into an ointment, in psoriasis, lepra vulgaris, &c.

NA'PIFORM, *Napiformis*, *Bul'noid*, *Buniôides*, (*napus*, 'a turnip,' and *forma*, 'shape,') An epithet for a form of scirrhus tumour whose cut surface presents concentric lines resembling those seen in the interior of a turnip.

NAPIUM, *Lapsana*, *Sinapis*.

NA'PLES, CLIMATE OF. The climate of Naples in its general characters resembles that of Nice, but it is more changeable: the sirocco too, which is little known in Nice, is severely felt at Naples. It is not a good residence for the phthisical invalid: Sir James Clark, indeed, considers it altogether unsuitable.

NAPLES, MIN'ERAL WA'TERS OF. In the Quarter Santa Lucia, near the coast, is a cold spring, rich in sulphuretted hydrogen and carbonic acid. It is much used as an aperient tonic, and in cutaneous affections.

NAPLES WA'TER, FACTITIOUS, *A'qua Neapolitana seu acid'ula hydrosulphurata*, (*F.*) *Eau de Naples*. (*Acidulous water*, containing four times its bulk of carbonic acid, 3xv, and 3ij; *Hydro-sulphuretted water*, 3ix; *Carbonate of soda*, gr. viij; *Carbonate of magnesia*, gr. x. *M. Ph. P.*)

NAPTA, *Naphtha*, *Nata*.

NAPUS, (*ναψ*, 'mustard,') *Sinapis*—*n*. *Leucosinapis*, *Sinapis alba*—*n*. *Sylvestris*, *Brassica napus*.

NAPY, (*ναψ*,) *Sinapis*.

NARCA, (*ναρκ*, 'stupor,') *Narcosis*.

NARCAPHTHE, *Narcaphthon*.

NARCAPH'THON, (*ναρκαφθον*,) *Narcaph'thon*. The bark of an aromatic tree formerly brought from India. By some, supposed to be that of the tree which affords the oilbanum. It was used in fumigation, in diseases of the lungs. See *Lacaphthon* and *Thymiana*.

NARCE, (*ναρκ*, 'stupor,') *Narcosis*.

NARCEMA, (*ναρκημα*,) *Narcosis*.

NARCESIS, (*ναρκεις*,) *Narcosis*.

NARCISSE DAUTOMNE, *Colchicum autumnale*—*n*. *Flour*, *Narcissus pseudonarcissus*—*n*. *des Prés*, *Narcissus pseudonarcissus*—*n*. *Sauvage*, *Narcissus pseudonarcissus*.

NARCIS'SUS PSEUDONARCIS'SUS, (*ναρκισσος*, from *ναρκ*, 'stupor,') *N. festalis seu glaucus seu grandiflorus seu hispanicus seu majus seu serratus seu sylvestris*, *Bulbocodium*, *Pseudonarcissus*, *Daf'fodil*, (old Eng.) *Aff'adil*, *Belle-Blume*, (*Prov.*) *Affodile*, *Bell'flower*, *Butter and Eggs*, *Lent rose*. *Ord.* *Amoryllidaceæ*. (*F.*) *Narcisse sauvage*, *Narcisse des prés*, *Faux narcissé*, *Porillon*, *Ailaut*, *Zouzinette*, *Jeannette*. The root is emetic and cathartic, in the dose of ʒij. The flowers are antispasmodic. Dose, 24 grains.

NARCODES, (*ναρκωδες*,) *Narcotized*.

NARCO'SIS, (*ναρκωσις*,) *Narcotism*, *Nar'cē*, *Nar'ca*, *Narce'ma*, *Narce'sis*, *Torpe'do*, *Tor'por*, *Stu'por*, *Stupefactio*. The aggregate effects produced by narcotic substances. At times, narcotism is confined to a state of more or less profound stupor; and constitutes, in certain cases, a useful remedial condition; at others, it is a true poisoning, characterized by vertigo, nausea, a state of intoxication, or apoplexy, constant delirium, convulsive motions, &c. Emetics in strong doses, and not much diluted with water; purgatives and glysters are the first means to be used in this condition. The stupor may afterwards be combated by the use of exciting and stimulating drinks.

The production of narcosis or the effect of narcotics has been called *narcotization*.

NARCOSIS FOLLICULO'RUM. A state of the scalp, which Mr. Erasmus Wilson conceive to be dependent upon torpid action of the hair follicles, and in which the scalp and hair are found covered with a yellowish, dirtylooking powder, composed of an admixture of granular particles and furfuraceous scales.

NARCOSPASM 'US, (*narce*, and *σπασμος*, 'spasm,') Stupor combined with spasm.

NARCOTIA, *Narcotin*.

NARCOTIC POISON, see *Poison*.

NARCOTICOACRID, see *Poison*.

NARCOTICS, *Narcotica*, *Carotica*, *Obstupefacientia*, *Stupefacientia*, *Stupeficients*, (*F.*) *Stup'factifs*, *Stup'fians*. Substances, which have the property of stupefying;—as opium, stramonium, and hyoscyamus, (from *Narce*.) They are used in medicine as soothing agents; exerting their special influence on the brain and tubular matter of the spinal marrow. In small doses, as a general rule, narcotics stimulate; in large, they act as sedatives. The following is a list of the chief narcotics:—*Aconitum*, *Ætherea*, *Belladonna*, *Cannabis*, *Conium*, *Digitalis*, *Humulus*, *Hyoscyamus*, *Lactucarium*, *Opium* and *Morphia*, *Stramonii Folia*, *Stramonii Semina*; *Mental Narcotics*, (*Appropriate Music*, *Monotonous sounds*, or any succession of monotonous impressions.)

NAR'COTIN, *Narcoti'na*, *Narco'tia*, *Anarco'tina*, *Opia'mum*; (same etymon,) (*F.*) *Sel de Déroïne ou d'opium*, *Opiane*, *Principe cristallisable de Déroïne*. A solid, white, inodorous, and insipid substance; by some considered to be alkaloid; by others neuter; crystallizable in straight prisms with a rhomboidal base; fusible like the fats; soluble in boiling alcohol and ether, and scarcely soluble in water. *Narcotin* produces all the unpleasant effects of opium; but at the same time, throws the animal into a state of stupor. It is not used in medicine, on account of these objections. The salts, which are very bitter, have been used successfully in India for the cure of intermittents.

NARCOTIZING, see *Narcosis*.

NARCOTISM, *Narcosis*.

NARCOTIZED, *Torpidus*, *Narco'des*. Affected with stupor, as from the use of a narcotic.

NARD, CELTIC, *Valeriana Celtica*—n. India, *Nardus Indica*—n. *Indien*, *Nardus Indica*—n. *Indique*, *Nardus Indica*—n. *Petit*, *Aralia nudicaulis*—n. *Saurage*, *Asarum*.

NARDOSTACHYS JATAMANSI, *Nardus Indica*—n. *Patrinia*, *Nardus Indica*.

NARDUM GALLICUM, *Valeriana Celtica*.

NARDUS AMERICANUS, *Aralia nudicaulis*—n. *Celtica*, *Valeriana Celtica*—n. *Celtica altera*, *Arnica montana*.

NARDUS INDICA, (*vapōs*), *Spica nardi* seu *Indica*, *Andropogon nardus* seu *citridorus*. Ord. Gramineæ [?]; or *Nardostachys* seu *Valeriana* seu *Patrinia Jatamansi*, *Valeriana spica*; Order, *Valerianaceæ*; *Indian nard*, *Spikenard*. The root of this plant is one of the ingredients in the mithridate and theriaca. It is moderately warm and pungent, and has a flavour by no means disagreeable. It is used by the Orientals as a spice:—(F.) *Nard Indien* ou *Indique*. An ointment was formerly used, called *Unguentum nardium*. It was prepared of *Nard*, *Malabathrum leaves*, *Oil of worms*, *Costus*, *Anonum*, *Myrrh*, &c., and was used as a detergent.

NARDUS MONTANA, *Asarum*—n. *Rustica*, *Asarum*.

NARES, (pl. of *naris*), *Rhînes*, *Cavitas* seu *Cavum* seu *Ca'væ* seu *Caver'na* na'rîum, *Na'vus* *inter'nus*, *Mycteres*, *Myxæ*, *Na'rîum ad'itus*, *Ocheteum'ata*, *The nostrils*, (Prov.) *Suir'rels*, *Suures*, (F.) *Narines*. Two elliptical apertures, situate beneath the nose, and separated from each other by the cartilaginous *septum* seu *colum'na nasi*. These apertures are continually open, and give passage to the air we breathe, and to the mucous fluids secreted in the nasal fossæ.

NARES INTERNÆ, Nasal fossæ.

NARES, POSTERIOR, *Na'res postre'mæ*, *Extre'mæ* *na'res*, *Ostia posteriora* seu *Choanæ* na'rîum, *Na'rîum Ex'itus*, (F.) *Arrière-narines*, *Ouvertures postérieures des fosses nasales*, are the posterior apertures of the nasal cavities, which establish a communication between those cavities and the pharynx. They are bounded, *above*, by the body of the sphenoid bone; *below*, by the palate bone, and the base of the velum pendulum; and, on the *outside*, by the internal ala of the pterygoid process. They are separated from each other by a septum, of which the vomer is the bony part.

NARIFUSORIA, (*nares*, 'the nostrils,' and *funder, fusus*, 'to pour.') Medicines, which are dropt or put into the nostrils.

NARINES, *Nares*.

NARIUM ADITUS, ('entrance or passage of the nostrils,') *Nares*—n. *Siccitas*, *Mycterorotes*.

NARROW CHEST, see *Lordosis*.

NARTHECIA, *Formulary*, *Myris*.

NARTHECIUM, (*ναρθηκίον*, dim. of *ναρθηξ*, 'a case of ointments,') *Formulary*, *Myris*.

NARTHEX, (*ναρθηξ*), *Formulary*, *Myris*, *Splint*—n. *Asafœtida*, see *Asafœtida*.

NASA, *Nacra*, *Nata*.

NASAL, *Nasa'lis*, (from *nasus*, 'the nose,') That which relates to the nose.

NASAL ARTERY. This is the largest of the two branches in which the ophthalmic artery terminates. It issues from the orbit, above the tendon of the orbicularis palpebrarum, passes above the side of the root of the nose, and anastomoses with the last extremity of the facial. Haller gave the name *nasal* to the sphenopalatine. He also called the dorsales nasi, furnished by the external maxillary, *Nasa'les laterales*.

NASAL BONES, *Os'na* na'vi seu *nasa'lia*, *Ossicula* na'vi, (F.) *Os nasæ*, *Os propriæ du nez*. These bones are situate beneath the nasal notch

of the os frontis, and occupy the space between the nasal or angular processes of the superior maxillary bone. Their shape is nearly quadrilateral. They have an *external* or *cutaneous* surface, an *internal* or *nasal*, and four margins. Each is articulated with its fellow, with the os frontis, ethmoid, and superior maxillary bones. They ossify from a single point.

NASAL CARTILAGE, *Cartilago triangul'aris* *Na'si*. A cartilage formed of three portions, which unite at the dorsum nasi, and are distinguished into the *cartilage of the septum*, and the *lateral cartilages*. This cartilage is continuous, *above*, with the ossa nasi, and, *inferiorly*, with membranous fibrocartilages, which form the supple and movable part of the nostrils. They are two in number—the one before, the other behind.

NASAL DUCT, Lachrymal duct—n. *Eminence*, *Mesophryon*.

NASAL FOS'SÆ, *Ca'vi* na'rîum, *Na'res inter'næ*. Two large, anfractuous cavities, situate between the orbits below the cranium, and lined by the pituitary or Schneiderian membrane. These cavities have no communication with each other; but the various sinuses in the neighbouring bones—the ethmoidal, sphenoidal, superior maxillary, &c.—all communicate with them. The general cavity of each nostril is divided by the ossa spongiosa into three *meatus* or *passages*, which run from before backwards. 1. The *Meatus na'rîum super'ior*, placed at the upper, inner, and back part of the superior spongy bone. 2. The *Meatus me'dius*, situate between the superior and inferior spongy bones; and 3. The *Meatus inf'erior*, situate between the inferior spongy bone and the bottom of the nose. The Schneiderian membrane receives the first pair of nerves, and various branches from the fifth pair. The arteries are furnished by the branches of the *internal maxillary*, known under the names of sphenopalatine, infraorbital, superior alveolar, palatine, pterygopalatine; by the supraorbital, and ethmoidal branches of the ophthalmic artery; by the internal carotid, superior labial, and dorsales nasi. Its veins are little known, and generally follow the course of the arteries. The lymphatics are almost unknown. The nasal fossæ are the seat of smell; they aid, also, in respiration and phonation.

NASAL MEATUS, see *Nasal fossæ*.

NASAL MUCUS, *Mucus* na'rîum, *Phlegma* na'rîum *cras'sum*, *Mucor* seu *Pituita* seu *Blen'na* seu *Lym'pha muculen'ta* na'rîum, vulgarly called *Snot*, *Snob*, *Snoth*, (F.) *Morve*, is the mucus secreted by the Schneiderian membrane.

NASAL NERVE, *Naso-palpébral* (Ch.), *Nasocula'ris* (Sömmering), *Nasocilia'ria*. One of the three branches of the ophthalmic nerve of Willis. It enters the orbit by the sphenoidal fissure, passes along the inner paries of that cavity, and divides, opposite the internal and anterior orbital foramen, into two branches. 1. The *internal* and *posterior*, which passes into the *Foramen orbitarium internum anteri'us*, enters the cranium beneath the dura mater, and passes into the nasal fossæ, through an aperture at the side of the *Crista galli*. It afterwards divides into several filaments: one of them—the *Nasolobar*, of Chaussier—is very small, and descends on the posterior surface of the os nasi, and ramifies on the integuments of the ala nasi. A second terminates near the septum; others descend along the outer paries of the nasal fossæ. 2. The other is *external* and *anterior*, and is called the *external nasal nerve*. It is distributed to the outside of the orbit. Before dividing, the nasal nerve communicates with the ophthalmic ganglion, and gives off two or three ciliary nerves. Sömmering has given the name *Posterior Nasal Nerves* to the nerves which arise from the internal part of the sphenopalatine ganglion.

NASAL NOTCH, (F.) *Échancrure nasale*. A semi-

circular notch, situate between the nasal prominence of the frontal bone, and articulated with the nasal bones and the nasal processes of the superior maxillary bones.

NASAL PROCESS, Maxillary bone, superior.

NASAL PROM'INENCE, (F.) *Bosse nasale*. A prominence, situate on the median line, at the anterior surface of the os frontis, between the two superciliary arches.

NASAL REGION, *Re'gio nasa'lis*. The region of the nose.

NASAL SPINES. These are three in number.

1. The *Superior nasal spine* of the os frontis, occupying the middle of its nasal notch, and articulated before with the nasal bones, behind with the ethmoid. 2. The *inferior and anterior nasal spine*, situate at the inferior part of the anterior opening of the nasal fossæ. It is formed by the two superior maxillary bones; and 3. The *inferior and posterior nasal spine*, *Épine gutturale* (Ch.), a process, formed on the mesial line by the two palate bones at the posterior part of the palatine arch.

NASALE, (from *nasus*.) Errhine.

NASALIS, Compressor naris.

NASA'LIS LA'BII SUPERIO'RIS. A small muscular slip, which runs up from the middle of the orbicularis and the lip to the tip of the nose. It lies exactly in the furrow, and is occasionally a levator of the upper lip, or a depressor of the tip of the nose.

NASAS, Nata.

NASATUS, ('long nosed,' from *nasus*:—the size of the nose being supposed an index of that of the male organ.) Membrosus.

NAS'CALE. A kind of pessary, made of wool, or cotton, which was formerly introduced into the vagina, after being impregnated with oil, ointment, or some other proper medicament.

NASCAPHTHON, Narcaphthon.

NASDA, Nata.

NASI, see Oryza.

NASITAS, (from *nasus*.) Rhinophonia.

NASITIS, (*nasus*, and *itis*.) Rhinitis—n. Postica, Angina nasalis.

NA'SO, *Nasu'tus*, (from *nasus*.) One who has a long nose. In composition, the nose.

NASOCILIARIS (Nervus), Nasal nerve.

NA'SOLA'BIAL, *Na'solabia'lis*; (from *naso*, and *labium*, 'a lip.') Relating to the nose and lip.

NASOLABIAL LINE. A line or furrow, which separates the lip from the cheek, and commences at the ala nasi.

NASOLABIALIS. A muscular fasciculus described by Albinus, which arises from the anterior extremity of the septum nasi, and terminates in the orbicularis oris.

NA'SOPAL'ATINE, *Na'so-palati'nis*. That which belongs to the nose and velum palati.

NASOPALATINE GAN'GLION is situate in the foramen palatinum anterior. Its greater extremity receives the two naso-palatine branches; whilst the smaller gives off two or three filaments, which reach the palatine vault, where they ramify on the membrane of the same name, anastomosing with filaments of the great palatine nerve.

NASOPALATINE NERVE, *Nerve of Cotu'ninus*, is furnished by the sphenopalatine, which proceeds from the ganglion of Meckel. It traverses the vault of the nasal fossæ, and proceeds upon the septum between the two layers of the pituitary membrane. It enters the anterior palatine canal, and terminates at the superior angles of the naso-palatine ganglion, without attaining the mouth.

NASOPALPÉBRAL, Orbicularis palpebrarum.

NASTA, Nata.

NASTURTIUM AMPHIBIUM, (*nasus*, and *torquere*, 'to twist,' owing to the effect of its pun-

gency,) see *Sisymbrium*—n. *Aquaticum*, *Cardamine pratensis*, *Sisymbrium nasturtium*—n. *Bursa pastoris*, *Thlaspi bursa*—n. *Hortense*, *Lepidium sativum*—n. *Indicum*, *Tropæolum majus*—n. *Officinale*, *Sisymbrium nasturtium*—n. *Palustre*, see *Sisymbrium*—n. *Peruvianum*, *Tropæolum majus*—n. *Pratense*, *Cardamine pratensis*—n. *Sativum*, *Lepidium sativum*.

NASUM DILATANS, ('dilating the nose,') *Pyramidalis nasi*.

NA'SUS, *Na'sum*, *Nose*, ([Sax.] *nære*, *ne're*.) *Rhin*, *Rhis*, *Or'ganon olfactûs seu odoratûs seu olfacto'rium*, *Promonto'rium fa'ciei*, *Myxo'ter*, *Emuncto'rium cer'e'brî*, *Snout*, (Sc.) *Neis*, *Nes*, (F.) *Nez*. The nose is a pyramidal eminence, situate above the anterior apertures of the nasal fossæ, which it covers; and, consequently, occupying the middle and upper part of the face, between the forehead and upper lip, the orbits and the cheeks. Its lateral surfaces form, by uniting angularly, a more or less prominent line, called *Dorsum seu Rho'dchis seu Spi'na Na'si*, (F.) *Dos du nez*. This line terminates, anteriorly, by the *lobe*,—*Lob'ulus*. The sides are called *Al'e Na'si*, *Pin'næ Na'ris*. (F.) *Ailes du nez*. The *columna* is the inferior part of the partition. Its apex or tip has been called *Glob'ulus nasi*.

The chief varieties of the nose are the *aquiline*, the *flat nose*, (F.) *Nez camarat ou épaté*, and the *snub nose*, (F.) *Nez retroussé*. The nose is formed, besides its bones, of fibro-cartilage, cartilage, muscles, vessels, and nerves; and its use seems to be to direct odours to the upper part of the nasal fossæ.

NASUS INTERNUS, Nares.

NASUTUS, Naso.

NA'TA, *Na'ta*, *Na'sa*, *Nas'da*, *Nas'ta*, *Na'sas*, *Nap'ta*. A fleshy, indolent excrescence, having the shape of the nates.

NATARON, Natron.

NATA'TION, *Nata'tio*, (*nature*, *natatum*, (itself from *nare*, 'to swim,') *Swim'ming*. The action of swimming, or of supporting one's self, or moving upon the water. Swimming resembles the horizontal leap in its physiology—the medium being water instead of air. The difference between the specific gravity of the human body and that of water is not great: so that but little exertion is required to keep a part of the body above water. Swimming is a healthy gymnastic exercise, combining the advantages of bathing.

NA'TES, sing. *Na'tis*, 'a buttock,' *Ephre'ra*, *Po'dex*, *Clu'nes*, *Glu'toi*, *Glu'ti*, *Glu'tia*, *Sedi'tia*, *Py'gæ*, *Mo'næ*, *Bot'tom*, *Back'side*, *Poste'riors*, *But'tocks*, (Prov.) *Tout*, (Sc.) *Dowp*, *Dowp*, *Dolp*, *Drod'dum*, *Hur'dies*, *Len'dis*, &c., (F.) *Fesses*. Two round projections, at the inferior and posterior part of the trunk, on which we sit, the *seat*, (F.) *Siège*. Amongst the mammalia, man alone has the nates prominent and round. They are formed chiefly by the skin, and a thick layer of areolar tissue, which covers the three glutæi muscles.

The cleft between the nates is the *Ri'ma clu'nium*.

NATES CEREBRI, Quadrigemina corpora—n. et Testes, Quadrigemina corpora.

NATIS, see Nates.

NATRIUM, Sodium.

NATRIX, ('a water serpent,') Penis.

NATROCRENÆ, (*natron*, and *κρηνη*, 'a well,') *Natropegæ*.

NA'TRON, (*νιτρον*.) *Na'trum*, *Na'taron*, *Ni'trum*, *Ana'tron*, *Aphroni'trum*, (from *Natron*, a lake in Judæa.) A saline compound, very abundant in Egypt, which is almost wholly formed of subcarbonate of soda.

NATRONIUM, Sodium.

NATROPE'GÆ, *Natrocre'næ*, (*natron*, and *πηγή*, 'a spring,') Soda springs.

NATRUM, Natron—*n.* Arsenicicum, Arseniate of soda—*n.* Biberacium, Borax—*n.* Chloratum liquidum, Liquor sodæ chlorinatæ—*n.* Muraticum, Soda, muriate of—*n.* Muriatum, Soda, muriate of—*n.* Vitriolatum, Soda, Sulphate of—*n.* Nitricum, Soda, nitrate of—*n.* Oxymuriaticum, Soda, chloride of—*n.* Oxyphosphorodes, Soda, phosphate of—*n.* Subboracicum, Borax—*n.* Sulphuricum, Soda, sulphate of—*n.* Præparatum, Soda, subcarbonate of—*n.* Tartarizatum, Soda, tartrate of.

NATTA, Nata.

NATULÆ, (dim. of *nates*.) Quadrigemina tubercula.

NATU'RA, (*nasci*, *natus*, 'to be born or arise.')

Phy'sis. Nature. Also, the genital organs.

NATURA MORBI. The essence or condition of a disease.

NATURAL PARTS, Genital organs.

NATURALIA, Genital organs.

NAT'URISM, *Nat'uralism*. A view which attributes every thing to nature, as a sage, prescient, and sanative entity. See Expectation, and *Vis Medicatrix Naturæ*.

NAT'URIST. A physician who scrupulously investigates, interprets, and follows the indications presented by nature in the treatment of disease.

NATUS MORTUUS, Still-born.

NAU'CLEA GAM'BIR, (*navis*, 'a ship,' and *κλειω*, 'to enclose,'—the half capsule being hull-shaped.) *Unca'ria gam'biri*. A plant of the *Nat. Order*, Rubiaceæ, native of the Malayan Peninsula and Indian Archipelago, which yields large quantities of the kind of Catechu known by the names *Terra Japonica* and *Square Catechu*, and which, in Indian commerce, is called *Gambier* or *Gambir*. It is a powerful astringent, much used in tanning, and in medicine. See Catechu.

NAUPATHIA, (*navis*, 'a ship,' and *παθος*, 'affection,') *Nausea marina*.

NAU'SEA, (*ναυσία*.) *Nau'sia*, *Nau'si'asis*, *Nau'sio'sis*, *Naut'ia*, *Queas'iness*, *Squeas'iness*, (F.) *Nausée*, *Envie de vomir*; (from *navis*, (L.) *navis*, 'a ship,' because those unaccustomed to sailing are so affected.) Sickness. Inclination to vomit.

NAUSEA, *KREAT'IC*, (*κρεας*, gen. *κρεατος*, 'flesh.')

The sickness and vomiting, excited, in some nervous patients, by the smallest portion of animal food.

NAUSEA MARI'NA, *Mor'bus nau'ticus*, *Vom'itus navigan'tium seu mari'nus*, *Naupath'ia*, *Seasick'ness*, (F.) *Mal de mer*. The sickness, vomiting, &c., experienced at sea by those unaccustomed to a seafaring; and from which those who are accustomed are not always exempt. It generally ceases when the person becomes habituated to the motion of the vessel, and not till then.

NAUSEABOND, Nauseant.

NAUSEABUNDUS, (from *nausea*.) Nauseant.

NAU'SEANT, *Nau'seans*, *Nauseabun'dus*, (F.) *Nauséabond*, *Nauséant*. An agent that excites nausea, which is a state of diminished action. Nauseants are, hence, valuable remedies in diseases of excitement.

NAUSÉE, Nausea.

NAUSIA, (*ναυσία*.) Nausea.

NAUSIASIS, Nausea.

NAUSIO'SIS, (*nausia* and *osis*.) This word, besides being synonymous with nausea, has been used to express the state of venous hemorrhage, when blood is discharged by jets.

NAUTIA, (*ναυτία*.) Nausea.

NAVEL, ([Sax.] *napela*.) Umbilicus—*n.* String, Funiculus umbilicalis—*n.* Wort, Cotyledon umbilicus—*n.* Wort, Venus's, Cotyledon umbilicus.

NAVET, (from *napus*, 'the turnip,') *Brassica rapa*—*n.* du *Diable*, *Bryonia alba*—*n.* *Galant*, *Bryonia alba*.

NAVETTE, *Brassica rapa*.

NAVIC'ULAR, *Navicula'ris*, *Naviform'is*, (F.) *Naviculaire*, (from *navicula*, 'a little ship.')

NAVICULAR FOS'SA, *Fos'sa Navicularis seu Scaphoi'des*, *Navic'ula*, *Scaph'ula*. See Scaphoid. This name has been given, 1. To a small depression between the entrance of the vagina and the posterior commissure of the labia major or fourchette. 2. To a perceptible dilatation, presented by the urethra in man, near the base of the glans. 3. To the superficial depression which separates the two roots of the helix. This is also called *Scaph'a*.

NAVICULARE OS, *Scaphoides os*.

NAVIFORMIS, (*navis*, 'a ship,' and *forma*, 'form,') *Navicular*.

NAVIS, ('a ship,') *Vulva*.

NEARSIGHTEDNESS, *Myopia*.

NEARTHRO'SIS, (*νεος*, 'young or new,' and *arthron*, 'a joint,') A new joint.

NEBLE, Nipple.

NEB'ULA, (from *νεφελη*, 'a cloud,') *Nu'bes*, *Nubec'ula*, *Ne'phus*, *Neph'elē*, *Nephe'lion*, *Vi'sus nebulosus*, (F.) *Nuage*, *Ombra'ge*. A slight speck on the cornea. A mist or cloud suspended in the urine. See *Caligo*.

NEBULOUS, *Nepheloid*.

NEC'ESSARY, *Necessari'us*, (*ne*, and *cessare*, 'to cease,') The *Necessaries of Life*, *Vitæ necessitates*, (F.) *Besoins de la vie*, include every thing requisite for the maintenance of life, and particularly food.

NECK, ([Sax.] *hnecca*, *necca*.) *Collum*—*n.* *Derbyshire*, *Bronchocele*.

NECK OF THE FOOT, *Instep*.

NECK, STIFF, *Torticollis*—*n.* Swelled, *Bronchocele*—*n.* *Wry*, *Torticollis*.

NECK'LACE, *AN'ODYNE*, (F.) *Collier anodin*. These are formed of the roots of *hyoscyamus*, *Job's tears*, allspice steeped in brandy, or the seeds of the wild liquorice vine, to suit the fancy of the prescriber. They are employed to facilitate dentition in children, and to procure sleep in fever!

NECKWEED, *Veronica beccabunga*, *V. peregrina*.

NECRÆ'MIA, (*necro*, and *αἷμα*, 'blood,') Death of the blood. Death beginning with the blood.

NECRENCEPHALUS, (*necros*, and *εγκεφαλος*, 'encephalon,') *Mollities cerebri*.

NECROBIOSIS, see Degeneration.

NECROBIOT'IC, *Necrobioti'cus*, (*necros*, and *βιος*, 'life,') Relating or appertaining to necrobiosis or morbid metamorphosis.

NECROCEDIA, (*necros*, and *κηδος*, 'funereal attentions,') *Embalming*.

NECROCŌMIUM, (*necros*, and *κομεω*, 'to care for,') *Morgue*.

NECRODES, (*νεκρωδης*, from *necros*, and *odes*.) *Cadaverous*.

NECRODOCHIUM, (*νεκροδοχειον*, from *necros*, and *δοχμα*, 'to receive,') *Morgue*.

NECROLOGY, *BILLS OF*, (*necros*, and *λογος*, 'a discourse,') *Mortality*, *bills of*.

NEC'ROMANCY, *Necromanti'a*, *Necyomanti'a*, (*νεκρομαντεια*, from *necros*, and *μαντεια*, 'divination,') *Divination by the dead*. Conjurat'ion; the black art, *Nigromanti'a*, *Negromanti'a*, (*niger*, 'black,' and *μαντεια*, 'divination,')

NECROMANTIA, *Necromancy*.

NECRONARCEMA, (*necros*, and *ναρκεμα*, 'stiffness,') *Rigor mortis*.

NECROPHOB'IA, (*necros*, and *φοβος*, 'fear,') Exaggerated fear of death. This symptom occurs in patients where the disease is not mortal; as in hypochondriasis. In fevers, it is not a good symptom.

NECROPNEUMO'NIA, *Pneumo'nia gangræna'sa seu typh'o'sa*, *Gangræ'na seu Mortificatio'seui An'thrax seu Carbun'culus Pulmo'nium*, *Pneumosep'sis*, *Pneumoc'acē*, (F.) *Gangrène du Pouton*,

Pneumonecrosis of Piorry; (*necros*, and *pneumonia*.) Gangrenous inflammation of the lungs. This may be *diffused* or *circumscribed*. The only pathognomonic symptom is the extraordinary and repulsive odour of the breath and expectoration. The treatment consists in the use of the chlorides internally, or of chlorine by inhalation; allaying irritation by opium, and supporting the patient by winewhey and nourishing diet.

NECROPSIA, (*necros*, and *opsis*, 'seeing,') Autopsia cadaverica.

NECROPSY, Autopsia cadaverica.

NECROS, (*νεκρος*,) Cadaver.

NECROSCOPIA, (*necros*, and *σκοπεω*, 'to examine,') Autopsia cadaverica.

NECROSCOPIC, *Necroscopicus*, (F.) *Necroscopique*, (*necros*, and *σκοπεω*, 'I view,') Relating to necroscopy or examination after death.

NECROSCOPY, Autopsia cadaverica.

NECROSED, see *Necrosis*.

NECROSEMIOTICÉ, (*necros*, and *σημειον*, 'a sign,') The doctrine of the signs of death.

NECROSIAL, *Necrosialis*, *Necrotic*, *Necroticus*. Relating or appertaining to death or mortification, or to necrosis,—as '*necrosial fever*;' the fever that accompanies necrosis.

NECROSIS, (F.) *Nécrose*, (*νεκρωσις*, from *νεκρω*, 'I kill,') Death, mortification. Especially, the state of a bone or of a portion of a bone deprived of life. *Osteogangræna*, *Osteonecrosis*. The bone, in such case, is said to be *necrosed*. Necrosis may take place without the surrounding soft parts being struck with gangrene. It is to the bones what gangrene is to the soft parts. The part of the bone affected with necrosis becomes a foreign body, similar to the gangrenous eschar, and its separation must be accomplished by the efforts of nature, or by art. When necrosis occurs in the centre of long bones, it never extends to their articular extremities. The exterior layers of bone form a canal round the dead portion or *sequestrum*; between these swollen layers and the sequestrum suppuration takes place; the matter presses against the bony canal; perforates it, and is discharged by apertures, which become fistulous.

In the treatment, the exit of the sequestrum must be facilitated by proper incisions, by the application of the trephine to the bone, &c.

NECROSIS CEREALIS, Ergotism—*n. Dentium*, Dental gangrene—*n. Ustilaginea*, Ergotism.

NECROSTOA, (*necros*, and *στοα*, 'a gallery,') *Morque*.

NECROTIC, *Necrosial*.

NECROTOCION, (*νεκροτοκιον*, from *necros*, and *τοκος*, 'a child,') A dead or stillborn child.

NECROTOMY, (*necros*, and *τομη*, 'section,') Dissection.

NECTAN'DRA CINNAMOMOI'DES, *Laurus cinnamomoides*, *Cinnamomum sylvestre America-num*: Order, Lauracæ; native of New Grenada, where it is called *Canela*; has the odour and taste of cinnamon, and is used as such.

NECTANDRA CYMBAR'UM, *Ocotea cymbarum* seu *amara*, grows in Oronoko, where it is called *Sassafra*s. The bark is aromatic, bitter, and stomachic; and Von Martius thinks it is an ingredient in the curare poison.

NECTANDRA PUCHURY, see *Pichurim* beans—*n. Rodiei*, see *Bebeery*.

NECTAR, (*νεκταρ*, from *νη*, a particle of negation, and *τραω*, 'I kill,') A pleasant liquor, feigned by the poets to have been the drink of the gods, and to have rendered immortal those who partook of it. A name given, by the ancients, to many drinks; and particularly to one made with wine, evaporated and sweetened with honey.

NECTARIUM, *Inula helenium*.

NECUSIA, (*νεκος*, 'to kill,') see *Wound*.

NECYOMANTIA, (*νεκος*, 'a dead body,' and *μαντια*, 'divination,') Necromancy.

NEDYIA, (*νηδεια*,) Intestines.

NEDYS, (*νηδυσ*,) The belly, abdomen, stomach, uterus.

NEE'DLE, (Sax. *nædl*, *nædl*, from Teut. *neten*, 'to sew,') *A'cus*, *Bel'oné*, *Ra'phis*, *Ra'phion*, *Rha'phis*, *Rha'phion*, *Aces'tra*, (F.) *Aiguille*. A steel instrument, used in many professions. In *Surgery*, a steel, gold, silver, or platinum instrument, that may be round, flat, or triangular, straight, or curved, supported or not by a handle, but having almost always a point, by means of which it can penetrate the textures; and either near the point, or more commonly, near the other extremity, an aperture or eye for the reception of a thread or tape, which it introduces into the parts. The *Aiguille à appareil* of the French is the ordinary sewing needle used in the making of bandages, &c.

NEEDLE, *A'cupuncture*, (F.) *Aiguille à Acupuncture*. An inflexible gold or silver needle; conical, very delicate, four inches long, furnished with a handle, and, at times, with a canula shorter than it by about half an inch. An ordinary needle, waxed at the head, will answer as a substitute for this. See *Acupuncture*.

NEEDLE, *Cat'aract*, *A'cus ophthalmica*, (F.) *Aiguille à cataracte*. Needles of gold, silver, and steel have been used; the latter, alone, at the present lay. The cataract needle is employed to depress or tear the crystalline when opaque. This needle is usually made from 15 to 24 lines long; and is attached to a fine handle. The extremity may be, as in Scarpa's and Langenbeck's, pointed, prismatic, triangular and curved; in Dupuytren's and Walther's, flat, curved, and sharp-edged; in Hey's, flat, with a semicircular and sharp end; or, as in Beer's, Siebold's, Schmidt's, Himly's, Von Gräfe's, &c., straight and spearpointed. A mark is generally placed upon the handle to inform the operator,—when the instrument is engaged in the eye,—what side corresponds to the crystalline.

NEEDLE FOR A COUNT'ROPENING, *A'cus invaginatus*, (F.) *Aiguille à Contre-ouverture ou en gaine* ou *à gaine*,—a long, narrow instrument of steel; the point of which is fine and sharp on both sides;—the heel (*talon*) having an aperture to receive a thread, tape, &c., provided with a flat, silver sheath, shorter than the blade, the point of which it covers when passing through parts that have to be respected.

NEEDLE, *Deschamps's*, *Paupé's needle*, (F.) *Aiguille de Deschamps*, *A. à manche*, &c., is the last described needle fixed to a handle. The eye is placed near the point. It is employed in the ligation of deep-seated arteries.

NEEDLE, *Fis'tula*, (F.) *Aiguille à Fistule*. A long, flat, flexible, silver instrument; having an aperture near one extremity; blunt at the other. This was formerly used for passing a seton into fistulous ulcers. On one of its sides was a groove for guiding a bistouri in case of necessity.

Also, a long, steel instrument, terminated by a point like that of a trocar, which Desault employed for penetrating the rectum, when operating for fistula that had no internal aperture.

NEEDLE, *Harelip*, (F.) *Aiguille à Bec de Lièvre*. A small, silver canula, to which is attached a spearpoint, that can be readily withdrawn. This needle, armed with the point, is introduced at one side of the fissure in the lip, and through the other. The twisted suture is then applied, and the pin withdrawn.

NEEDLE, *Lig'ature*, (F.) *Aiguille à ligature*. A long, steel instrument, sharp towards one extremity, with an eye near the other, which was formerly used for suspending the circulation of blood prior to amputation, by being passed through

the limb, so as to include the principal artery and a part of the muscles and integuments.

Also, a steel instrument of various dimensions, round towards one of its extremities, which was straight, and furnished with an aperture; curved and flat towards the opposite, which was pointed, and had a slight ridge on its concave side. At the commencement of the last century, this instrument was used to pass ligatures around vessels.

Also, a steel instrument of various dimensions, flat, regularly curved in the form of a semicircle, with a sharp or lance point, and a long eye, used with advantage in place of the last.

NEEDLE, PAUPE's, Needle, Deschamps'.

NEEDLE, SE'TON, (F.) *Aiguille à Séton*. A long, narrow, steel blade; pointed and sharp at one extremity; pierced at the other by an aperture. The *Aiguille à contre-ouverture* may be used for the same purpose.

NEEDLE, SU'TURE, (F.) *Aiguille à Suture*. For the twisted suture, the hare-lip needle is used; for the others, the straight or curved needle: the straight needle is preferable for stitching up the abdomen, &c., in dissection. In the suture of the tendons, a curved needle has been used; flat on both sides, and cutting only at the concave edge, in order that the instrument may pass between the tendinous fibres without dividing them.

NEEDLEBEARER, *Porte-aiguille*.

NEEDLECARRIER, *Porte-aiguille*.

NEM, see *Melia azedarach*.

NEESING, Sneezing.

NÉFLE, Mespilus (the fruit).

NÉFLIER, Mespilus.

NEFREN'DES, properly, 'sucking pigs.' *No'-doi, Eden'tuli*. Persons devoid of teeth. Young children, for instance, who have not cut them; or aged persons, who have lost them. This state is called *Nefren'dis, Odon'tia eden'tula*.

NEGOTIUM PARTURITIONIS, ('affair of parturition,') Parturition.

NÈGRE, Negro—*n. Blanc* ('white negro,') Albino.

NEGRETIA PRURIENS, *Dolichos pruriens*.

NE'GRO, (S. and P. *negro*, from (L.) *niger*, 'black,') *Nègrita*, (F.) *Nègre*. One of the Æthiopian race. See *Homo* and *Mulatto*.

NEGRO, PIED, see *Achroma*.

NEGROMANTIA, (*niger*, 'black,' and *μαντεία*, 'divination,') see *Neeromancy*.

NEI'ERA, (*νεῖρα*), Abdomen.

NEIGE, Snow.

NEIR, (Sc. *niere*,) Kidney.

NEIRA, (*νεῖρα*), Abdomen.

NEIS, Nasis.

NELUMBIUM LU'TEUM, *Yel'low ne'lumbo, Yel'low wa'ter lil'y, Pond lily, Water shield, Water nuts, Water chin'capin, Rat'tle nut, Sa'cred bean*. A beautiful water plant, common in the United States, and belonging to *Nat. Ord. Nymphaeaceæ; Sex. Syst. Polyandria Polygynia*. The leaves are cooling and emollient when applied to the surface. The roots, leaves, and nuts are eaten. The last are called by the Indians and others *water chin'capins*.

NELUMBO, YELLOW, *Nelumbium luteum*.

NENDO, *Angelica lucida*.

NENNDORF, MIN'ERAL WA'TERS OF. Nenndorf is a village three and a half German miles from Hanover. Its cold sulphureous spring is much celebrated.

NENUPIAR, *Nymphaea alba*—*n. Blanc*, *Nymphaea alba*—*n. Jaune*, *Nymphaea lutea*—*n. Lutea*, *Nymphaea lutea*—*n. Odorant*, *Nymphaea odorata*.

NEO, (from *νῆς*, 'new,') in composition, new.

NEOARTHRO'SIS, (*neo*, and *arthron*, 'a joint,') A new joint; an artificial joint.

NEOG'ALA, (*neo*, and *γάλα*, 'milk,') Milk se-

creted immediately after the colostrum. Also, the colostrum.

NEOGENES, (*νεογενής*, from *neo*, and *γενω*, 'to be born,') *Nouveau-né*.

NEOGILUS, (*νεογίλος*), *Nouveau-né*.

NEOGNUS, (*νεογνός*), *Nouveau-né*.

NEOGONUS, (*νεογονός*), *Nouveau-né*.

NEOMEMBRANE, see *Membrane*, false.

NEONATUS, (*neo*, and *natus*, 'born,') *Nouveau-né*.

NE'OPLASM, *Neoplasm'a*, gen. *Neoplasm'atis*, (F.) *Néoplasme*, (*neo*, and *πλασσω*, *πλάττω*, 'I form,') A new formation or tissue; the product of morbid action.

NE'OPLASTY, *Neoplas'ticē*. An operative process for the formation of new parts. It includes autoplasty, cicatrization of wounds, and the formation of adhesions.

NEOSCYTECHUS, (*neo*, *σκυτος*, 'leather,' and *ηχος*, 'a sound,') see *Craquement de Cuir neuf*.

NEOTOCITERUS, (*neo*, *τοκος*, 'one born,' and *icterus*,) *Icterus infantum*.

NEOTIA, (*νεοτία*, 'a nest,'—from the arrangement of the roots,) *Goodyera pubescens*.

NEP, *Nepeta*.

NEPEN'THA DESTILLATO'RIA, *Bandu'ra*. A Ceylonese plant, the root of which is astringent.

NEPEN'THES, (*νεπενθής*, from *νη*, 'neg,' and *πενθος*, 'grief,') A remedy much extolled by the ancients against sadness and melancholy.

The women of Thebes, according to Diodorus Siciliotes, alone possessed the secret of its composition; and, according to Homer, Helen introduced it from Egypt. Some suppose it to have been opium.

NEPENTHES, Bangu—*n. Opiatum*, *Pilulæ opiatæ*.

NEP'ETA, (from *Nepete*, a town in Italy [?]), *N. Cata'ria seu vulga'ris, Cata'ria vulga'ris, Her'ba fe'lis, Nep. Cat'nepe, Cat'nip, Cat'mint*, (F.) *Herbe aux Chats, Cataire*. *Ord. Labiateæ*. The leaves, *Cata'ria*, (Ph. U. S.), have a smell and taste like those of an admixture of spearmint and pennyroyal. It has been recommended in uterine disorders, dyspepsia, flatulency, &c., like pennyroyal; and is much used on the American continent in domestic medicine, in flatulencies, &c., of children.

NEPETA AGRESTIS, *Melissa nepeta*—*n. Glechoma, Glechoma hederacea*—*n. Vulgaris, Nepeta*.

NEPHALIOTES, (*νεφθαλιότης*, from *νηφω*, 'I abstain,') Temperance.

NEPHELE, (*νεφελή*, 'a cloud,') *Enœorema, Nebula*.

NEPHELION, (*νεφελιον*, 'a little cloud,') *Nebula*.

NEPH'ELOID, *Nepheloi'des, Nephelo'des, Neb'ulous, Nubilo'sus*, (*νεφελειοειδής*, from *νεφελή*, 'a cloud,' and *εἶδος*, 'resemblance,') An epithet applied to urine when it is cloudy—*Uri'na nephelo'des*.

NEPHOS, (*νεφός*, 'a cloud,') *Nebula*.

NEPHREMO'RHAGIA, (*nephritis*, and *hæmorrhagia*,) see *Hæmaturia*.

NEPHRAL'GIA, *Do'tor Nephret'icus, Neural'gia Re'num*, (F.) *Néuralgie des Reins, Dysnéphro-nervie*, (*nephritis*, and *αλγος*, 'pain,') Pain and neuralgia in the kidney.

NEPHRALGIA ARENOSA, see *Gravel*—*n. Calculosa*, see *Gravel*—*n. Rheumatica, Lumbago*.

NEPHRANURIA, (*nephritis*, *a*, priv. and *ουρον*, 'urine,') *Ichuria renalis*.

NEPHRAPOS'TASIS, *Nephropy'o'is, Abces'sus ren'al'is*, (*nephritis*, and *αποστήσις*, 'abscess,') Renal abscess.

NEPHRATON'IA, (*nephritis*, and *ατονία*, 'want of tone,') *Re'num aton'ia seu paral'yo'is*. Atony of the kidney.

NEPHRAUX'E, (*nephritis*, and *αυξω*, 'to increase,') Enlargement of the kidney.

NEPHRELCOISIS, *Nephrogonia'sis*, *Hælo'sis rena'lis*, (*nephrus*, and *ἄλωσις*, 'ulceration.') Ulceration of the kidney.

NEPHRELMINTIC, *Nephrelminticus*, (*nephrus*, and *ἔμυς*, gen. *ἑμύβος*, 'a worm.') That which is owing to the presence of worms in the kidney.

NEPHREMPHRAX'IS, (*nephrus*, and *εμψρασσω*, 'I obstruct.') A name given by Plouquet to obstruction of the kidneys.

NEPHRETIC, *Nephrid'ius*, *Nephrit'ic*, *Nephrit'icus*, (from *nephrus*.) That which relates to the kidney. Applied, especially, to pain, &c., seated in the kidney.

NEPHRETICUM, in *Materia Medica*, means a medicine employed for the cure of diseases of the kidney.

NEPHRETICUM LIGNUM, *Guilandina moringa*.

NEPHRITICUS LA'PIS, *Tal'cum nephrit'icum*, (F.) *Pierre néphrétique*, *Jade néphrite* ou *oriental*. A green, fatty kind of stone, — once used as an amulet against calculus and epilepsy, — an absurd name, as there can be no such remedy.

NEPHRIA, (*νεφρος*, 'kidney,') Kidney, Bright's disease of the.

NEPHRID'ION, *Pingue'do rena'lis*, (*νεφρίδιον*, 'pertaining to the kidney,') The fat which surrounds the kidney.

NEPHRIDIDIUM, Capsule. renal.

NEPHRIDIVS, (*νεφρίδιος*.) Nephretic.

NEPHRINE, (from *nephrus*;) *Oxyl*, cystic, *Urea*.
NÉPHRITE, Nephritis—*n. Albumineuse*, Kidney, Bright's disease of the.

NEPHRITES, (from *nephrus*;) *Asphaltites*.

NEPHRITIC, Nephretic.

NEPHRITICI (*morbi*), in the classification of Dr. Wm. Farr, renal diseases.

NEPHRITICUM LIGNUM, *Guilandina moringa*.

NEPHRITIS, (from *nephrus*;) *Asphaltites*.

NEPHRITIS, *Empres'na Nephrit'is*, (*nephrus*, and *itis*.) *Renum inflammatio*, *Nephrophleg'monē*. — *Inflammation of the Kidneys*, (F.) *Néphrite*, *Inflammation des Reins*, is characterized by acute pain; burning heat, and a sensation of weight in the region of one or both kidneys; suppression or diminution of urine; fever; dysuria; ischuria; constipation, more or less obstinate; retraction of the testicle, and numbness of the thigh of the same side. It may be distinguished into *simple*, and *calculus nephritis*—*Lithonephrit'is*. In the latter, the urine often contains small particles of uric acid or of urate of ammonia. The most common causes of nephritis are,—excess in irritating and alcoholic drinks; abuse of diuretics; blows or falls on the region of the kidneys; the presence of renal calculi, &c. It may be distinguished from lumbago by the pain which attends the latter on the slightest motion, &c. It usually terminates by resolution in from one week to two or three. It may, however, end in suppuration—*pyonephritis*;—or may become chronic—*chrononephritis*. In the treatment, antiphlogistics, as bleeding, baths, &c., are required to the full extent; with the use of diluents, opiates, &c.

NEPHRITIS ALBUMINOSA, Kidney, Bright's disease of the—*n. Albuminosa*, Kidney, Bright's disease of the—*n. Cachectic*, Kidney, Bright's disease of the—*n. Calculous*, see Nephritis.

NEPHRITIS DES'QUAMATIVE. A condition of Bright's disease, in which epithelial cells are seen by the microscope in the urine.

NEPHRITIS SOCIATA, Kidney, Bright's disease of the.

NEPHROCHALAZOSIS, (*nephrus*, and *χάλαζα*, 'hail,' 'granulation,') Kidney, Bright's disease of the.

NEPHROCOCCOSIS, (*nephrus*, and *κοκκος*, 'a grain or berry,') Kidney, Bright's disease of the.

NEPHROCOLICA, *Colica nephretica*.

NEPHROCYSTOSIS, (*nephrus*, and *κύστις*, 'a cyst or bladder,' and *osis*.) A morbid condition, which consists in the formation of cysts in the bladder.

NEPHRODES, (*νεφροειδής*.) Nephroid.

NEPHRODIUM FILIX MAS, (from *nephrus*;) Polypodium filix mas.

NEPHROGRAPHY, (*nephrus*, and *γραφη*, 'a description,') An anatomical description of the kidney.

NEPHROHÆMIA, (*nephrus*, and *αἷμα*, 'blood,') Hyperæmia or congestion of the kidney.

NEPHROID, *Nephroï'des*, *Nephroï'des*, *Reniform'is*, (*nephrus*, and *ειδος*, 'form, resemblance,') Reniform. Having a resemblance to a kidney—as 'nephroid cancer,' so called because the morbid growth resembles the kidney in structure.

NEPHROLITHIASIS, (*nephrus*, and *λίθος*, 'a stone,') The disease of calculus in the kidney; *Lith'ia rena'lis*, *Lith'iasis nephrit'ica*, *L. Rena'lis*. See Gravel.

NEPHROLITHIC. Belonging to calculi in the kidneys. This epithet has been applied to ischuria, occasioned by calculi formed in the kidneys.

NEPHROLITHOTOMY, Nephrotomy.

NEPHROLOGY, *Nephrolog'ia*, (*nephrus*, and *λογος*, 'a treatise,') A treatise on the kidneys. Dissertation on the kidneys and their functions.

NEPHROMALACIA, (*nephrus*, and *μαλακία*, 'softness,') Softening of the kidney.

NEPHRONCUS, *Tumor rena'lis*, (*nephrus*, and *ογκος*, 'a tumour,') A tumefaction of the kidney.

NEPHROPARALYSIS, *Nephropar'esis*, *Nephrophle'gia*, *Paral'ysis renum* seu *rena'lis*. Paralysis of the kidneys.

NEPHROPARESIS, (*nephrus*, and *παρεσις*, 'loss of tone,') Nephroparalysis.

NEPHROPHLEGMATIC, *Nephrophlegmat'icus*, (*nephrus*, and *φλεγμα*, gen. *φλεγματος*, 'phlegm,') A name given, by some authors, to ischuria produced by mucus contained in the urine.

NEPHROPHLEGMONE, (*nephrus*, and *φλεγμονη*, 'inflammation,') Nephritis.

NEPHROPTHISIS, (*nephrus*, and *φθισις*, 'consumption,') Phthisis from suppuration of the kidney.

NEPHROPLEGIA, (*nephrus*, and *πληγη*, 'a stroke,') Nephroparalysis.

NEPHROPLETHORIC, *Nephroplethor'icus*, (*nephrus*, and *πληθωρα*, 'plethora,') Belonging to plethora of the kidneys. An epithet given to ischuria dependent upon this cause.

NEPHROPYIC, *Nephropy'icus*, (*nephrus*, and *πυον*, 'pus,') Belonging to suppuration of the kidney.

NEPHROPYOSIS, Nephropostasis, Nephrelcosis.

NEPHRORRHAGIA, *Hæmatu'ria rena'lis*, *Hæmorrhag'ia re'num*, (*nephrus*, and *ρῆξις*, 'rupture,') *Proflu'vium sanguinis e re'nibus*. Hemorrhage from the kidney.

NEPHROS, (*νεφρος*.) Kidney.

NEPHROSCLERIA, (*nephrus*, and *σκληρία*, 'hardness,') *Induratio re'num*. Induration of the kidneys.

NEPHROSPASTIC, *Nephrospas'ticus*, (*nephrus*, and *σπασμ*, 'I draw,') That which depends upon spasm of the kidney. An epithet given to a variety of ischuria.

NEPHROTHROMBOD, *Nephrothromboï'des*, (*nephrus*, and *θρομβος*, 'a clot,') That which depends upon clots of blood, contained in the kidneys or their ducts. An epithet for a species of ischuria.

NEPHROTOMY, *Nephrolithot'omy*, *Nephrotom'ia*, *Nephrolithotom'ia*, *Sec'tio rena'lis*. *S. re-*

nis, (*nephros*, and *ρᾶνναι*, 'to out.') Dissection of the kidney. Also, an operation proposed with the view of extracting calculi formed in the kidney, by means of an incision into the tissue of that organ.

NEPHRUS, (*νεφρος*,) Kidney.

NEPIOS, (*νηπιος*,) Infans.

NEPIOTES, (*νηπιότης*,) Infancy.

NEPTA, Asphaltum.

NERA, Toddy.

NERE, (Sc.) ([G.] Niere,) Kidney.

NERF, Nerve—*n. Circonflexe*, Axillary nerve—*n. Dentaire*, Dental nerve—*n. Fessier*, Gluteal nerve—*n. Glosso-pharyngien*, Pharyngoglossal nerve—*n. Gustatif*, see Lingual nerve—*n. Gustatif innominé*, Trigemini—*n. Gutturale*, Palatine (middle) nerve—*n. Gutturo-palatin*, Palatine nerve—*n. Honteux*, Pudic nerve—*n. Irien*, Ciliary nerve—*n. Ischio-clitorien*, Pudic nerve—*n. Ischiopénien*, Pudic nerve—*n. Labyrinthin*, Auditory nerve—*n. Mentonnier*, Mental nerve—*n. Moteur oculaire externe*, Motor oculi externus—*n. Oculomusculaire externe*, Motor oculi externus—*n. Oculo-musculaire interne*, Patheticus nervus—*n. Pathétique*, Patheticus nervus—*n. Péronière branche*, Popliteal nerve, external—*n. Pharyngoglossien*, Pharyngoglossal nerve—*n. Pneumogastrique*, Pneumogastric nerve—*n. Première paire trachélienne*, Occipital nerve—*n. Prétibio-digital*, Musculocutaneous nerve, see Popliteal nerve—*n. Prétibio-susphalangéaire*, see Popliteal nerve—*n. Prétibio susplantaire*, Tibial nerve, anterior—*n. Radio-digital*, Radial nerve—*n. Sciatique grand*, Sciatic nerve, great—*n. Sciatique petit*, Sciatic nerve, lesser—*n. Sciatique poplitée externe*, Musculocutaneous nerve—*n. Sous-occipital*, Occipital nerve—*n. Sous-pubio-fémoral*, Obturator nerve—*n. Spino-cranio-trapézien*, Spinal nerve—*n. Susmaxillaire*, Maxillary superior nerve—*n. Suspucien*, Suprapubic nerve—*n. Sympathique*, Trisplanchnic nerve—*n. grande Sympathique*, Trisplanchnic nerve—*n. Tibiale branche*, Popliteal internal—*n. Trachéal*, Laryngeal inferior nerve—*n. Trachélo-dorsal*, Spinal nerve—*n. Trijumeau*, Trigemini—*n. à Trois cordes*, Trigemini—*n. Tympanique*, Chorda tympani—*n. Vague*, Pneumogastric nerve—*n. Vertébro-digital*, Musculocutaneous nerve.

NERFES BRONCHIQUES, Bronchial nerves—*n. Ciliaires*, Ciliary nerves—*n. Craniens*, Cerebral nerves—*n. Moteurs oculaires communs*, Motores oculorum—*n. Rachidiens*, Vertebral nerves—*n. Sacrés*, Sacral nerves—*n. Sous-costales*, Intercostal nerves.

NÉRION, Nerium oleander.

NÉRIS, MINERAL WATERS OF. Nérís is on the high road from Moulins to Limoges, eighty leagues from Paris. There are four springs, the water of which is clear, inodorous, tasteless, and has an unctuous feel. It contains carbonic acid gas, carbonate of soda, and sulphate of soda, chloride of sodium, silicic acid, and an animal matter to which its 'oleaginous' property is probably owing. The waters are generally used in the form of thermal baths. Season, from the 20th of May to the 15th of October.

NERIUM, ('humid,' because growing in moist places.) *N. oleander*.

NERIUM ANTIDYSENTERICUM, *Wrightia antidysenterica*. The tree which affords the *Codagapala Bark*, *Comesici cortex*, *Tillischer's cortex*, *Cortex profluvi*. Family, Apocynaceæ. Sex. Syst. Pentandria Monogynia. (F.) *Codagapala*, *Cropala*. The bark of this Malabar tree is of a black colour, externally, and is generally covered with a white moss or scurf. It has an austere, bitter taste, and has been recommended as an astringent in diarrhoea, dysentery, &c.

NERIUM ODO'RUM, *Kunaree*, of India, is a well known poison, and extensively used as such.

NERIUM OLEAN'DER, *Nerium*, *Rhododaphnis*, *Rosa'go*, (F.) *Laurier rose*, *Laurrose*, *Nérion Rouge*, *Rhododaphné*. The leaves are reputed to be narcotic. The infusion is employed internally, in herpetic affections, and the powder, incorporated with lard, is used in the itch.

NEROLI, see Citrus aurantium.

NERONIA'NA seu NERO'NICA PHLEBOTOM'IA. An epithet given, for some cause, to phlebotomy, when more than one vein was opened on the same day, or to bleeding *coup sur coup*. Termed *Neroniana*, because the philosopher Seneca was led to death under Nero [?].

NERPRUN PURGATIF, (or *Noirprun*, 'prunus nigra,') *Rhamnus*.

NERVE, (*νεῦρον*, 'a string.') *Nervus*, *New'ron*, (F.) *Nerf*. (*Neuron* and *Nervus* meant also, with the ancients, a tendon and a ligament, *Partes nervosæ*; and hence the different acceptations of 'nervous':—'a man of nerve—a strong, nervous man; and a weak, nervous woman.') The nerves are tubular cords of the same substance as that which composes the encephalon and spinal marrow. They extend from one or other of the nervous centres to every part of the body, communicating, frequently, with each other; forming *plexuses*, and, occasionally, *ganglions*; and being, at length, lost in the parenchyma of organs. There are 42 pairs, and, according to their origin, they are termed *Cranial* or *Encephalic*, and *Spinal*. Each nerve is composed of several filaments or cords placed alongside each other, and is surrounded by a neurilemma. The encephalic nerves, in general, have only one root in the brain, whilst the spinal arise from the marrow by two roots: the one from an anterior fasciculus of filaments, the other from a posterior, separated from each other by the *Ligamentum denticulatum*; uniting outside this ligament, and presenting, near the intervertebral foramen, a ganglion formed only by the posterior root. The two roots make, afterwards, but one nerve; and, like the encephalic nerves, proceed to their destination, subdividing into rami and ramusculi, until they are finally lost in the texture of the organs. The trunks first formed are commonly round, and proceed alone, or accompany the great vessels, being placed in the areolar spaces which separate the organs, and are thus protected from injury. Their manner of termination we are not acquainted with; whether the nervous pulp, for instance, as is more probably the fact, is distributed or lost in a membrane, as seems to be the case with the nerves of sight, hearing, and smell,—or is looped;—see *Corpuscles*, *Pacinian*, and *Corpuscula tactils*. Certain it is, that there is considerable difference in the organs, as respects the quantity of nerves that terminate in them; and the particular arrangement of the nervous extremities. Some organs have numerous nerves; others seem to have none: a circumstance which influences considerably the sensibility of parts. Observation has shown, that certain motor fibres cross the median line and affect muscles on the side opposite to that from which they originate; but the influence is very limited. See *Entrecroisement*.

The *Encephalic nerves* arise from the encephalon, or are inserted into it, (according as we consider the brain the origin or termination of the nerves;) and make their exit by foramina at the base of the skull. They are 12 in number. The spinal nerves are 31 in number, 8 *cervical*, 12 *dorsal*, 5 *lumbar*, and 6 *sacral*: the four inferior cervical being much larger than the superior, because they furnish the nerves of the upper extremities.

SYNOPTICAL TABLE OF THE NERVES.

I. Cranial or Encephalic Nerves.

1. <i>Olfactory</i> (1st pair.)	Divided into internal, external, and middle branches, which are distributed on the Schneiderian membrane.	6. <i>Abducentes</i> (6th pair.)	To the rectus externus oculi.
2. <i>Optic</i> (2d pair.)	Terminate in the retina.		
3. <i>Motores oculorum</i> ... (3d pair.)	<i>Superior Branch.</i> To the rectus superior oculi and levator palpebrae superioris. <i>Inferior Branch.</i> To the rectus internus, rectus inferior and lesser oblique muscles; a filament which goes to the ophthalmic ganglion.	7. <i>Facial</i> (Portio dura of the 7th pair.)	1. At its exit from the cranium, the posterior auricular, stylohyoid, and infra-mastoid, to the pavilion of the ear, the mastoid process, the digastricus, and the muscles attached to the styloid process. 2. Near the parotid gland, the temporal, malar, buccal, supra-maxillary, and infra-maxillary to the whole superficies of the face.
4. <i>Pathetici</i> (4th pair.)	To the greater oblique muscle of the eye. <i>Ophthalmic Branch.</i> Divided into three branches. 1. The lachrymal branch, to the lachrymal gland and upper eyelid. 2. <i>Frontal branch</i> , to the forehead and upper eyelid. 3. <i>Nasal branch</i> , to the eyelids, nasal fossae, and nose. <i>Superior Maxillary Branch.</i> 1. The orbital branch, to the orbit. 2. The posterior and superior dental, to the last three molar teeth and gums. 3. The anterior dental, to the incisor, canine, and two lesser molares. 4. <i>Infraorbital</i> , to the upper lip, cheek, and nose. <i>Inferior Maxillary Branch.</i> 1. Temporal profound branches, to the temporal muscle. 2. <i>Masseterine branch</i> , to the masseter muscle. 3. <i>Buccal</i> , to the inner surface of the cheek. 4. <i>Pterygoid</i> , to the internal pterygoid muscle. 5. <i>Lingual</i> , to the mucous membrane of the tongue. 6. <i>Inferior dental</i> , to the teeth of the lower jaw, and to the lower lip. 7. <i>Auricular branch</i> , to the pavilion of the ear and forehead.	8. <i>Auditory</i> (Portio mollis of the 7th pair.) 9. <i>Glossopharyngeal</i> ... (Portion of the 9th pair.)	To the vestibule, semicircular canals, and cochlea. To the base of the tongue and pharynx.
5. <i>Par Trigeminum</i> ... (5th pair.)		10. <i>Par Vagus</i> (8th pair.)	1. <i>IN THE NECK.</i> — a pharyngeal branch to the pharynx; superior laryngeal branch to the larynx, and to some muscles of the inferior hyoid region. Cardiac branches to the cardiac plexus. 2. <i>IN THE CHEST.</i> — the inferior laryngeal branch to the larynx pulmonary branches, which form the plexus of the same name; oesophageal branches to the oesophagus. 3. <i>IN THE ABDOMEN.</i> — gastric branches to the plexus of the stomach; filaments which go to the neighbouring plexuses. At its exit from the cranium — the accessory of the pneumogastric nerve anastomosing with this nerve. In the neck — filaments to the trapezius muscles. <i>Cervicalis descendens</i> , to the muscles of the inferior hyoid region and to the cervical nerves. Filaments to the muscles of the tongue.
		11. <i>Spinal or Accessory of Willis</i>	
		12. <i>Hypoglossal</i> (9th pair.)	

II. Spinal or Vertebral Nerves.

1. <i>Cervical Nerves.</i>			
1st Cervical Pair.....	<i>Anterior Branch.</i> Anastomosing par arcade with the second pair.	Cervical Plexus.....	1st. to the parotid gland and pavilion of the ear. <i>Superficial Cervical Branches.</i> (2.) To the platysma myoides, digastricus, the integuments of the neck, &c. <i>Anterior Branches.</i> Assisting in the formation of the brachial plexus.
2d Cervical Pair.....	<i>Anterior Branch.</i> A branch anastomosing with the first pair; a branch which goes to the cervical plexus.		<i>Posterior Branches.</i> To the muscles and integuments of the posterior part of the neck and upper part of the back.
3d and 4th Cervical Pairs.....	<i>Anterior Branch.</i> Concurring in the cervical plexus.	5th, 6th, and 7th Cervical Pairs.....	1. <i>Thoracic Branches.</i> To the anterior and lateral parts of the chest. 2. <i>Suprascapular</i> branch to the muscles of the posterior scapular region. 3. <i>Infra-scapular</i> branches (3.) to the subscapularis, teres major, teres minor, and latissimus dorsi. 4. The brachial internal cutaneous, to the integuments of the palmar and dorsal surfaces of the forearm, near the ulnar margin. 5. The brachial external cutaneous, principally to the integuments of the palmar and dorsal surfaces of the forearm, near the radial margin. 6. The median nerve, to the
The Posterior Branch of these four pairs ramifies on the occiput and muscles of the superficial and deep-seated cervicocephalal regions.			
	DESCENDING BRANCHES. — 1. <i>Internal descending</i> , anastomosing with a branch of the hypoglossus. 2. <i>Parenchymal branch</i> , to the diaphragm. 3. <i>External descending branches</i> , dividing into supra-clavicular, supra-acromial, infra-clavicular, and cervicalae profundae, to the muscles and integuments of the upper part of the chest and shoulder, to the trapezius, levator scapulae, rhomboidaeus, &c.	Brachial Plexus.....	
Cervical Plexus.....	ASCENDING BRANCHES. — 1. <i>Mastoid</i> , to the posterior and lateral part of the head and the inner surface of the pavilion of the ear. 2. <i>Auricu-</i>		

Brachial Plexus..... { forearm, hand, and palmar surface of all the fingers. 7. The *cubital nerve*, to the last two fingers. 8. The *radial*, to the first three. 9. The *axillary*, around the shoulderjoint, and to the neighbouring muscles.

2. Dorsal Nerves, (*Nervi dorsales*.)

1st Dorsal Pair..... { *Anterior Branch*. To the brachial plexus.

2d and 3d Dorsal Pairs..... { *Anterior Branches*. An intercostal and brachial branch.

4th, 5th, 6th, and 7th Dorsal Pairs..... { *Anterior Branches*. *Internal branches*, to the intercostals, triangularis sterni, pectoralis major, and to the skin. *External branches* to the integuments of the chest, the obliquus externus abdominis, and the skin of the abdomen.

8th, 9th, 10th, and 11th Dorsal Pairs... { *Anterior Branches*. *Internal branches*, to the transversalis, obliquus internus, and rectus muscles, and to the skin of the abdomen. *External branches*, to the integuments of the chest, and to the muscles and skin of the abdomen.

12th Dorsal Pair..... { *Anterior Branch*. To the first lumbar nerve, and to the muscles and skin of the abdomen as far as the iliac crest.

The *Posterior Branches* of the dorsal nerves are distributed to the muscles and integuments of the back and loins.

3. Lumbar Nerves.

1st, 2d, 3d, and 4th Lumbar Pairs..... { *Anterior Branches*. Concurring to form the lumbar plexus.

5th Pair..... { *Anterior Branch*. Aiding in forming the sciatic plexus.

The *Posterior Branches* of the lumbar nerves are distributed to the loins, sacrum, and nates.

Lumbar Plexus.....

1. *Musculocutaneous Branches*, to the number of three. One *superior*, to the muscles of the abdomen, to the fold of the groin, and the scrotum: a *middle*, to the integuments and muscles of the abdomen: and an *inferior*, to the skin of the thigh. 2. A *genitocrural* branch to the integuments of the scrotum, the groin, and the thigh. 3. The *crural nerve*, to the integuments and muscles of the thigh, skin of the leg, and foot. 4. The *obturator nerve*, to the muscles at the inner part of the thigh. 5. The *lumbosacral*, to the sciatic plexus. It gives off the *gluteal nerve* to the glutei muscles.

4. Sacral Nerves.

1st, 2d, 3d, and 4th Sacral Pairs..... { *Anterior Branches*. They form by their union the sciatic plexus.

5th and 6th Pairs..... { *Anterior Branches*. To the parts in the vicinity of the coccyx.

The *Posterior Branches* ramify on the muscles and integuments of the nates.

1. *Hemorrhoidal nerves*, to the rectum. 2. *Vesical*, to the bladder. 3. *Uterine and vaginal*, to the vagina and uterus. 4. *Inferior gluteal*, to the gluteal muscles, perineum, and integuments of the posterior part of the thigh. 5. *Pudic*, to the perineum, penis, or vulva. 6. *Sciatic*, divided into the *external popliteal*, which ramifies on the integuments and muscles of the external side of the leg, on the dorsum of the foot, and the dorsal surface of the toes; and into the *internal popliteal*, distributed on the dorsal surface of the two last toes, to the muscles of the foot, and on the plantar surface of all the toes.

Sciatic Plexus.....

Classifications of the nerves have been recommended according to their uses, in preference to the ordinary anatomical arrangement. It has been remarked that the encephalic nerves have generally one root; the spinal two. Experiments and pathological facts have proved that the anterior column of the marrow—*funiculus anterior*—and the anterior roots of the spinal nerves are inservient to volition or voluntary motion; and that the posterior column—*funiculus posterior*—and roots are destined for sensibility. Hence the spinal nerves, which have two roots, must be the conductors both of motion and feeling: whilst the encephalic, which, with but few exceptions, have but one, can possess but one of these properties:—they must be either *sensitive* or *motor*, according as they arise from the posterior or anterior column of the medulla; and, consequently, three classes of nerves may be distinguished.

1. *Sensory*..... { Arising, by a single root, from the posterior column of the medulla oblongata or spinal marrow.
2. *Motor*. { Arising, by a single root, from the anterior column of the same parts.
3. *Sensory and Motor*..... { Which have two roots: one from the anterior, and one from the posterior column.

The tubules of all the afferent or sensory nerves cross to the opposite side of the spinal marrow, and an impression received by them is conveyed along the vesicular matter to the sensory ganglia of the encephalon; whilst the efferent or motor fibres cross only at the anterior pyramids of the medulla oblongata.

According to Sir Charles Bell, the medulla oblongata is composed of three columns on each side,—an *anterior*, a *middle*, and a *posterior*. Whilst the anterior and posterior columns produce the nerves of motion and sensation respectively, the middle, according to Sir Charles, gives rise to a third set of nerves—the *respiratory*. To this order belong:—

1. The accessory nerve of Willis, or *superior respiratory*.
2. The par vagum.
3. The glossopharyngeal.
4. The facial or *respiratory of the face*.
5. The phrenic.
6. A nerve which has the same origin as the phrenic;—the *external respiratory*.

When a horse has been hardridden, every one of these nerves is in action.

This division is now, however, generally abandoned, and there does not seem to be a third column, especially destined for the movements of respiration.

Sir C. Bell, again, reduced the system of nerves to two great classes. 1. Those that are regular, primitive, symmetrical, and common to all animals, from the worm to man, which have double roots, and preside over sensibility and motion; and, 2. The irregular or superadded, which are added to the preceding, in proportion as the organization of animals offers new or more complicated organs. To the first class belong all the spinal nerves and at least one encephalic—the 5th pair;—to the second, the rest of the nervous system.

Dr. Marshall Hall proposed a division of the nervous system, which is calculated to explain many of the anomalous circumstances so frequently witnessed. He proposed to divide all the nerves into, 1. The cerebral or the sentient and voluntary. 2. The true spinal or excitomotory. 3. The ganglionic or cycloganglionic, the nutrient and secretory. If the sentient and voluntary functions be destroyed by a blow upon the head, the sphincter muscles will still contract when irritated, because the irritation is conveyed to the spine, and the reflex action takes place to the muscle so as to throw it into contraction. But if the spinal marrow be now destroyed, the sphincters remain entirely motionless, because the centre of the system is destroyed. Dr. Hall maintained that a peculiar set of nerves constitutes, with the vesicular centre of the marrow as their axis, the second subdivision of the nervous system; and as those of the first subdivision are distinguished into sentient and voluntary, these may be distinguished into the *excitor* and *motor*. The first, or the *excitor nerves*, pursue their course principally from internal surfaces, characterized by peculiar excitabilities, to the centre of the medulla oblongata and m. spinalis; the second or the *motor nerves* pursue a reflex course from the medulla to the muscles, having peculiar actions concerned principally in ingestion and egestion. The motions connected with the first or cerebral subdivision are sometimes, indeed frequently, *spontaneous*; those connected with the true spinal are, he believes, *always excited*. He thought, too, that there is good reason for viewing the fifth and posterior spinal nerves as constituting an external ganglionic system for the nutrition of the external organs; and he proposed to divide the ganglionic subdivision of the nervous system into, 1. The *internal ganglionic*, which includes that usually denominated the sympathetic, and probably filaments of the pneumogastric; and, 2. The *external ganglionic*, embracing the fifth and posterior spinal nerves. To the cerebral system he assigned all diseases of sensation, perception, judgment, and volition—therefore all painful, mental, and comatose, and some paralytic diseases. To the true spinal, excitomotory, reflex, or distal nervous system, he referred all spasmodic and certain paralytic diseases. He properly added that these two parts of the nervous system influence each other both in health and disease, as they both influence the ganglionic system.

The main views of Dr. Hall on the excitomotory or reflex system have been generally embraced. Much, however, has still to be accomplished, notwithstanding the activity of modern observers, before the precise functions of the different nervous centres can be regarded as definitely determined.

The nerves are sheathed and are united to the neighbouring parts, by an abundant layer of fatty areolar texture, which sends inwards prolongations that separate the nervous cords and filaments from each other. The arterial trunks distributed to them transmit branches into their interior. The veins follow the same course as the arteries. Absorbents are not easily traced even on the greatest trunks.

NERVE, ACCESSORY, OF WILLIS, Spinal nerve—n. Articular, Axillary nerve—n. Buccinator, Buccal nerve.

NERVE CELLS, see Neurine—n. of Cotunnus, Nasopalatine nerve.

NERVE, FACIAL, see Facial nerve—n. Femoral Cutaneous, see Genitoocrural nerve—n. Femoropopliteal, lesser, Sciatic nerve, lesser.

NERVE FIBRES, *Fibra nervosa*, *Fibra nervosa*. In the nerves of the higher animals two forms of nerve fibre exist—the *tubular*, and the *gelatinous*. In the minute fibrils or tubules, *Tubuli nervosi*, full of nervous matter, which constitute the former, there is a difference between the central and peripheral portion;—the former has been called the *axis cylinder* and *primitive* and *flattened band*; the latter, the *medullary* or *white substance* of Schwann—*nerve medulla*, *medullary sheath* or *matter*, *myelin*—being that to which the peculiarly white aspect of cerebro-spinal nerves is principally due. In the gelatinous fibre no tubular envelope can be distinguished, and the white substance of Schwann seems to be wanting. It contains numerous cell-nuclei. See Neurine.

NERVE, GENITAL, see Genitoocrural nerve—n. Genitoocrural, see Genitoocrural nerve—n. Gluteal of the Sacral plexus, Sciatic nerve, lesser—n. Gustatory, Hypoglossal nerve—n. Inguinal, internal, Genitoocrural nerve—n. of Jacobson, see Petrosal ganglion—n. Lateral nasal, Sphenopalatine nerve—n. Lingual, Hypoglossal nerve.

NERVE MEDULLA, see Nerve fibres.

NERVE, NASOOCULARIS, Nasal nerve—n. Nasopalatine, see Nasopalatine—n. Nasopalpebral, Nasal nerve—n. Ocular, Optic nerve—n. Orbitofrontal, Ophthalmic nerve—n. of Organic life, Trisplanchnic nerve—n. Palpebrofrontal, Frontal nerve—n. Pang, Neuralgia—n. Phrenic, Diaphragmatic nerve—n. Popliteal internal, Tibial nerve, posterior.

NERVE POWER, Nervous power.

NERVE, RESPIRATORY, EXTERNAL, see Thoracic nerves—n. Respiratory, of the Face, Facial nerve—n. Respiratory, superior, Spinal nerve—n. Scapulohumeral, Axillary nerve—n. Scrotal, see Genitoocrural nerve—n. Spiral, Radial nerve—n. Spiral, muscular, Radial nerve—n. Suboccipital, Occipital nerve—n. Subpubian, Genitoocrural nerve—n. Superficial, temporal, Auricular nerve—n. Sympathetic, Trisplanchnic nerve—n. Temporal cutaneous, Auricular nerve—n. Ulnar, Cubital nerve—n. Vidian, Pterygoid nerve—n. Vocal, see Spinal nerve—n. of Wrisberg, see Cutaneous nerves—n. Zygomaticauricular, Auricular nerve.

NERVES, COSTAL. Intercoastal nerves—n. Dorsal, Intercoastal nerves—n. Eighth pair of, Pneumogastric—n. Encephalic, first pair, Olfactory nerves—n. Encephalic, eleventh pair, Spinal nerve—n. Ethmoidal, Olfactory nerves—n. Fifth pair, Trigemini—n. Fourth pair of, Patheticus nervus—n. Infraorbital, Suborbitar nerves—n. Ninth pair of, Hypoglossal nerve—n. Oculomuscular, common, *Motores oculorum*—n. Phrenic, Diaphragmatic nerves—n. Recurrent, Laryngeal inferior nerves—n. Sixth pair of, *Motor oculi externus*—n. Spinal, Vertebral nerves—n. Tenth pair of, Pneumogastric—n. Third pair of, *Motores oculorum*.

NERVEUX, Nervous.

NERVI ACROMIALES, Acromial nerves—n. Anonymi, Innominati nervi—n. Claviculares, Clavicular nerves—n. Cutanei clunium inferiores, see Sciatic nerve, lesser—n. Divisi, Trigemini—n. Entobiontes, Afferent nerves—n. Gustatorii, Trigemini.

NERVI MOLLES: 'soft nerves.' The anterior branches of the superior cervical ganglion of the great sympathetic, which accompany the carotid artery and its branches, around which they form intricate plexuses, and here and there small gan-

glia. They are called *molles* from their softness, and *subrufi* from their reddish hue.

NERVI ODORATORII, Olfactory nerves—*n. Reversivi*, Laryngeal inferior nerves—*n. Subrufi*, Nervi molles—*n. Vulneratio*, Neurotrosis.

NERVIMOT'ION, *Nervimotio*, (*nerve*, and *motio*, 'motion.') A name given by M. Dutrochet, to the movement which is caused in the organs of the senses by external agents, and transmitted to the muscles by the nerves.

NERVIMOT'OR, (*F.*) *Nervimoteur*. An epithet given by M. Dutrochet to external agents which are capable of causing nervimotion.

NER'VINE, *Nervinus*, *Neurotic*, *Neuritic*, *Antineurotic*, *Antineuropathic*, (*F.*) *Nervin*, *Névritique*, *Névrolique*. A medicine which acts on the nervous system.

NERVOPATHIC, Neuropathic.

NER'VOSISM. The doctrine which maintains that all morbid phenomena are owing to variations in the nerve force.

NER'VOUS, *Nervosus*, *Neurodes*, *Nervens*, *Neuricus*, (*F.*) *Nerveux*. (Same etymon.) Relating or belonging to the nerves. Strong. Also, weak, irritable.

NERVOUS ATTACK', (*F.*) *Attaque des nerfs*, is an affection accompanied with spasm, pain, and different nervous symptoms, to which impressible individuals are liable.

NERVOUS CENTRES, see Centres, nervous.

NERVOUS DIATHESIS, termed *Ataxia spirituum*, *Anomalia nervorum*, *Status nervosus* seu *erethiticus*, *Diathesis* seu *Constitutio nervosa*, *Nervousness*, (*F.*) *Névroisme*. Medically, *nervousness* means unusual impressibility of the nervous system. Formerly, it signified strength, force, and vigour. Recently, it has been applied to a hypochondriacal condition verging upon insanity, occasionally occurring in those in whom the brain has been unduly tasked;—a condition termed by some *cerebrop'athy*; by others, *brain fog*.

NERVOUS DISEASES, *Morbi Nervosi*, (*F.*) *Maladies nerveuses*. Affections seated in the nervous system. To purely functional disease of the nerves Dr. Laycock has given the name *neuræmia*, and to the class of diseases he applies the epithet *neuræmic*.

NERVOUS FLU'ID, *Nervous principle*, *Fluidum* seu *Liquidum nervinum*, *Sucus nervus*, *Spiritus vitalis*. The fluid which is supposed to circulate through the nerves, and which has been regarded as the agent of sensation and motion. Of this fluid we know nothing, except that it resembles, in many respects, the electric or galvanic. It was formerly called *Animal spirits*, *Spiritus animalis*, (*F.*) *Esprits animaux*.

NERVOUS PAIN, Neuralgia.

NERVOUS POW'ER, *Nerve power*, *Via nervosa* seu *nervorum*, *Neurodynamia*, *Neurodynamis*. The power of the nerves as exhibited in the living organism.

NERVOUS PRINCIPLE, Nervous fluid.

NERVOUS SYSTEM. The nerves and nervous centres considered collectively. See Nerves.

NERVOUS SYSTEM OF THE AUTOMATIC FUNCTIONS, Trisplanchnic nerve—*n. System*, ganglionic, see Trisplanchnic nerve—*n. System*, organic, see Trisplanchnic nerve—*n. System* of the senses, *Medulla oblongata*—*n. System* of voluntary motion, &c., *Medulla spinalis*.

NERVOUSNESS, Nervous diathesis.

NERVUS, see Nerve, Penis—*n. Acusticus*, Auditory nerve—*n. Ambulatorius*, Pneumogastric nerve—*n. Anonymus*, Trigemini—*n. Auditorius*, Auditory nerve—*n. Caroticus*, Carotid nerve—*n. Cervicofacialis*, Cervicofacial nerve—*n. Communicans faciei*, Facial nerve—*n. Cutaneus internus*, see Saphenous nerves—*n. Cutaneus internus longus*, see Saphenous nerves—*n. Decimus*, Pneumo-

gastric—*n. Ganglionicus*, Trisplanchnic nerve—*n. Genitoceuralis*, Genitoceural nerve—*n. Hypoglossus*, Hypoglossus.

NERVUS IMPAR. A prolongation of the neurilemma below the lower extremity of the spinal cord, as a fibrous filament, which is inserted into the base of the coccyx. It was formerly regarded as a nerve, hence its name.

NERVUS INCISIVUS, Incisive nerve—*n. Innominatus*, Trigemini—*n. Ischiadicus*, Sciatic nerve—*n. Juvenilis*, Penis—*n. Latus*, Achillis tendo—*n. Lingualis lateralis*, Pharyngoglossal nerve—*n. Lingualis medius*, Hypoglossal nerve—*n. Loquens*, Hypoglossus—*n. Lumboinguinalis*, Genitoceural nerve—*n. Mixtus*, Trigemini—*n. Musculi pterygoidei*, Pterygoid nerve—*n. Nasociliaris*, Nasal nerve—*n. Opticus*, Optic nerve—*n. Pneumogastricus*, Pneumogastric—*n. Pudendus externus*, see Genitoceural nerve—*n. Quintus*, Trigemini—*n. Spermatikus*, see Genitoceural nerve—*n. Subcutaneus male*, Orbital nerve—*n. Sympatheticus magnus*, Trisplanchnic nerve—*n. Sympatheticus medius*, Trigemini, Pneumogastric nerve—*n. Sympatheticus minor*, Facial nerve—*n. Sympathicus*, Trisplanchnic nerve—*n. Sympathicus medius*, Trigemini—*n. Temporofacialis*, Temporofacial nerve—*n. Timidus*, Motor oculi externus—*n. Tremellus*, Trigemini—*n. Trisplanchnicus*, Trisplanchnic nerve—*n. Trochlearis*, Patheticus nervus—*n. Vagus*, Pneumogastric nerve—*n. Vagus cum accessorio*, Spinal nerve—*n. Vasomotorius*, Trisplanchnic nerve.

NES, Nasus.

NESPIE, Melissa calamintha.

NESTIA, (*νηστεία*,) Fasting.

NESTIATRIA, (from *nestia*, and *ιατρία*, 'medical treatment,') Limotherapeia.

NESTIS, (*νηστις*, 'a faster,') Jejunum.

NESTITHERAPEIA, Limotherapeia.

NESTOTHERAPEIA, Limotherapeia.

NESTROOT, *Monotropa uniflora*.

NETLEAF, *Goodyera pubescens*.

NETTING, Urine.

NETTLE, ARCHANGEL, WHITE, (*Sax. netel*), *Lamium album*—*n. Day*, *Lamium album*—*n. Dead*, *Lamium album*, *L. amplexicaule*—*n. Dwarf*, *Urtica urens*—*n. Hedge*, *Galeopsis*—*n. Horse*, *Solanum Carolinense*—*n. Pillbearing*, *Urticaria pilulifera*.

NETTLE RASH, Urticaria—*n. Springe*, Urticaria.

NETTLE, STINGING, COMMON, Urtica.

NETTLE TREE, *Celtis occidentalis*.

NETWORT, *Goodyera pubescens*.

NEURADYNAMIA, (*neuron*, *a*, and *δυναμις*, 'power,') Neurasthenia.

NEURÆMIA, (*νευρον*, 'a nerve,' and *αἷμα*, 'blood,') See Nervous diseases.

NEURÆMIC, see Nervous diseases.

NEU'RAL, (from *neuron*.) Relating to a nerve or to the nervous system or centres.

NEURAL ARCH. The arch formed by the posterior projections connected with the body of the vertebra, which protect the medulla.

NEURAL AXIS, see Eucephalon—*n. Spines*, Spinous processes.

NEURAL'GIA, (*neuron*, and *αλγος*, 'pain,') *Rheumaticus epuricus nervosus*, *Neurodynia*, *Nervous pain* or *pang*, (*F.*) *Néuralgie*, *Névrodynie*. A generic name for a number of diseases, the chief symptom of which is a very acute pain, exacerbating or intermitting, which follows the course of a nervous branch, extends to its ramifications, and seems, therefore, to be seated in the nerve. The principal neuralgias have been distinguished by the names *facial* (of which the *infraorbital*, *maxillary*, and *frontal* are but divisions)—the *iliocrotal*, *femoropopliteal*, *femoropretibial*, *plantar*, and *cubito-digital*. A division of anomalous neuralgia has likewise been admitted.

All varieties of neuralgia are obstinate, and the greatest diversity of means has been made use of:—bleeding, general and local,—emetics, purgatives, rubefacients, vesicants, actual cautery, narcotics, mercurial frictions, electricity, destruction of a portion of the nerve, &c. The most successful remedy, perhaps, is the carbonate of iron, given in doses of some magnitude; as, for instance, $\mathfrak{ss}$ or $\mathfrak{ij}$ twice or thrice a day, in molasses. This plan of treatment, continued for a month or two, will often relieve, and ultimately remove this much dreaded affection. The mode in which it acts is by no means clear; but it is almost as certain as any other remedy used in disease in producing its salutary effects. The bowels must be kept free, and all inflammatory symptoms removed during its administration.

NEURALGIA ANOMALOUS. Under this name Chaussier has included different neuroses, some of which are characterized by acute pains circumscribed within a short compass, or extending by irradiations, but not having their seat in the course of a nerve; and others which are occasioned by tumours in the course of a nerve, or which succeed contusions or incomplete divisions of nerves.

NEURALGIA BRACHIALIS, Brachialgia—*n.* Brachiothoracica, Angina pectoris—*n.* Cerebralis, Hemiplegia—*n.* Celiacæ, Neuralgia of the solar plexus (Romberg)—*n.* Cruralis, *N. femorotibialis.*

NEURALGIA CUBITODIGITALIS, Is'chias nervo'sa digitalis. In this variety the pain extends from the part where the nerve passes under the inner condyle to the back of the hand and to its cubital edge.

NEURALGIA DENTALIS, Odontalgia nervosa.

NEURALGIA, FA'cial, Neuralgia fac'ciei, Tris'mus maxilla'ris seu *Dolorificus* seu *Clon'icus, Opsial'gia, Do'l'or cru'cians* fac'ciei seu *faciei* period'icus, *Hemicra'nia idiopath'ica, Autal'gia dolorosa, Tic douloureux, Do'l'or fac'ciei* seu *faciei* typ'ico caracte'rè seu *faciei* Fothergill'i, *Rheumatis'mus cancro'sus, Rheumatis'mus larva'us, Prosopop'alia, Prosopodyn'ia, Fe'bris top'ica, Ophthalmodyn'ia period'ica, Tortu'ra O'ris, Affe'ctus spasmodicoconvulsi'vus Labio'rum, Face ague,* is characterized by acute lancinating pains, returning at intervals; and by twinges in certain parts of the face, producing convulsive twitches in the corresponding muscles. It may be seated in the frontal nerve, in the infraorbital, or in the maxillary branch of the fifth pair.

Metopodyn'ia, Brocague, Neural'gia fronta'lis, Ophthalmodyn'ia, Tic douloureux, Do'l'or period'icus, &c., commences at the superciliary foramen and extends along the ramifications of the frontal nerve that are distributed on the forehead, upper eyelid, caruncula lacrymalis, and nasal angle of the eyelids. Sometimes it is felt particularly in the orbit.

Neuralgia infraorbital'ia, Do'l'or fac'ciei a'trox, Rheumatis'mus cancro'sus, Tris'mus clon'icus, Prosopop'alia, Fe'bris top'ica, Odontalgia remittens et intermittens, Hemicra'nia se'ra, Infraorbital' neuralgia, Tic douloureux, is chiefly felt in the infraorbital foramen, whence it passes to the cheek, upper lip, ala nasi, lower eyelid, &c.

In *Neuralgia maxilla'ris, Gnathalgia, Gnathoneural'gia, Tic douloureux, Tris'mus cataracta'lis maxilla'ris,* the pain usually sets out from the mental foramen and passes to the chin, lips, temple, teeth, and tongue.

NEURALGIA, FALSE. A term assigned to pains along a nerve or its ramifications, produced by some body compressing it,—those pains terminating with the removal of the compressing cause.

NEURALGIA FEMORO-POPITE'A, N. Ischiad'ica,

Sciati'ca, Sci'as, Mal'um ischiad'icum, Mar'bus ischiad'icus, Pas'sio ischiad'ica, Ischi'agra, Ischiad'igia, Is'chias, Ischio'sis, Ischioneural'gia, Coxal'gia, Do'l'or Ischiad'icus nervo'sus, Is'chias, I. nervo'sa Cotu'g'ni seu Cotumiti, Malum Cotu'g'ni, I. nervo'sa post'ica seu rheumat'ica, Neurisch'ias, Cox'agra, Ischiad'ica, (Prov. Devonshire) Bone-shave, (F.) Sciatique, Goutte sciaticque. This is characterized by pain following the great sciatic nerve from the ischiatic notch to the ham, and along the peroneal surface of the leg to the sole of the foot. It is often considered to be a form of rheumatism—*rheumatism of the hip.*

NEURALGIA FEMOROPRETIBIALIS, N. cru'ral, Is'chias nervo'sa anti'ca, Scelalgia anti'ca. In this the pain, setting out from the groin, spreads along the fore part of the thigh, and passes down, chiefly on the inner side of the leg, to the inner ankle and back of the foot.

NEURALGIA FRONTALIS, see *N. Facial*—*n.* of the Heart, Angina pectoris.

NEURALGIA HEPATICA, Hepatalgia.

NEURALGIA ILIOSCROTAL has been rarely observed. It is characterized by a very acute pain in the course of the branches of the first lumbar pair; this pain follows the crista ilii and accompanies the spermatic vessels to the testicle, which is often painfully retracted.

NEURALGIA INFRAORBITAR, see *N. Facial*—*n.* Ischiadica, *N. Femoropopliteal*—*n.* Lead, see *Lead rheumatism*—*n.* Lienis, Splenalgia.

NEURALGIA MAM'ME, Irritable breast. An exceedingly painful affection of the female mamma, unaccompanied by inflammation.

NEURALGIA MAXILLARIS, see *N. Facial*

NEURALGIA PLANTARIS. This is rare, and the pain is confined to the course of the plantar nerves.

NEURALGIA RENUM, Nephralgia—*n.* Spinalis, Spinal irritation—*n.* Testis, Orchidalgia.

NEURANAGENNESIS, Neur'anaph'ysis, (neuron, and αναγεννω, 'to regenerate.') Regeneration or renewal of a nerve.

NEURANAPHYSIS, (neuron, ανα, 'again,' and γεννω, 'to grow.') Neur'anagenesis.

NEURAPOPHYSIS, see *Vertebræ.*

NEURASTHENIA, Neuradyn'ia, Debilitas nervo'sa. Debility or impaired activity of the nerves, (*neuron*, and ἀσθενεια, 'debility.')

NEURECTOMIA, (neuron, and εκτομη, 'excision.') Excision of a nerve.

NEURICUS, (from neuron,) Nervous.

NEURILEMMA, gen. Neurilemm'atis, Neurile'ma, Neurily'ma, Neurymen, Perineur'ion, Fis'tula seu Tu'bulus seu Cap'sula seu Involucrum nervo'rum, Membra'na seu Membran'ula seu Tu'nica seu Indumen'tum nervo'rum, Vagina nervo'rum, (F.) Nérilème, Périnèvre, (neuron, and λεμμα, 'a coat.') The fine transparent, and apparently fibrous membrane that surrounds the nerves, to every filament of which it forms a true canal.

Robin restricts the term *perinèvre* to the structure that invests the primitive bundles of nerves; and every nervous filament. It differs from the neurilemma in its ordinary acception, which forms a coat of some thickness in the larger nerves, and supports the nutrient vessels, whilst the *perinèvre* exhibits no capillaries.

NEURILEMMATITIS, Neurilemmitis.

NEURILEMMITIS, Neurilemmat'is, Neurolemmat'is, Neurymenit'is, Neurilymit'is, (F.) Nérilémite, (neurilemma, and itis.) Inflammation of the neurilemma.

NEURILITY, Neuril'itas, (from neuron.) The property, possessed by nerve-fibre, of exciting contraction in a muscle, secretion in a gland, and sensation in a ganglionic centre.

NEURILYMA, (neuron, and ελυμα, 'a covering,') Neurilemma.

NEURINE, (from *neuron*.) *Medulla nervorum*, (F.) *Névrine*, *Albumine cérébrale* ou du *cerveau*. The substance of which the nervous system is composed. It consists chiefly of albumen and a peculiar fatty matter, associated with phosphorus. There are two kinds of neurine—the one *vesicular* or consisting essentially of *nerve vesicles* or *nerve cells* or *corpuscles*, or, as they are sometimes called, from their prevailing in the ganglia—*ganglion corpuscles* and *ganglion globules*; the other, the *tubular*, formed—as the word imports—of *tubules*. The former is the *cineritious* or *cortical* nervous matter of the older anatomists;—the latter, the *white* or *medullary*. The *vesicular* neurine has appeared to be restricted to the production and distribution of nervepower; the *tubular*, to its conduction; yet the experiments of Dr. Brown-Séquard show, that the former conveys sensitive impressions from the posterior spinal roots to the brain. Most of the nerve cells are furnished with one process, hence termed *unipolar*; others have two processes—the *bi'polar*; whilst others are devoid of processes—the *a'polar*.

NEURISCHIAS, Neuralgia femoropoplitæa.

NEURITIC, Nervine.

NEURITIS, *Inflammat'io nervorum*, *Neurophlogo'sis*, *Neuropheg'monê*, (F.) *Inflammation des Nerf*, *Nérite*, (*neuron* and *itis*), Inflammation of a nerve.

NEUROBLACTIA, (*neuron*, and *βλακεια*, 'stupor.') Insensibility of the nerves.

NEUROCHONDRO'DES, (*neuron*, *χονδρος*, 'a cartilage,' and *odes*.) A hard substance between a sinew and a cartilage.

NEURODES, (*νευρώδης*, from *neuron* and *odes*), Nervous.

NEURODYNAMIA, (*neuron*, and *δυναμις*, 'power,') Nervous power.

NEURODYNAMIS, Nervous power.

NEURODYNIA, (*neuron*, and *δύνη*, 'pain,') Neuralgia.

NEUROGAMIA, (*neuron*, and *γάμος*, 'marriage,') Magnetism, animal.

NEUROGLIA, (*neuron*, and *γλτα*, 'glue.') A form of connective tissue, which binds the nervous elements together. (Virchow.)

NEUROGRAPHY, *Neurograph'ia*, (F.) *Névrographie*, (*neuron*, and *γραφη*, 'a description.') The part of anatomy which describes the nerves.

NEUROHYPNOLOGY, *Neurohypnology'ia*, (*neuron*, *υπνος*, 'sleep,' and *λογος*, 'a treatise.') A name given by Mr. Braid to his theory of magnetic sleep. See Magnetism, animal.

NEUROHYPNOTISM, see Magnetism, animal.

NEUROLEMMATITIS, Neurilemmitis.

NEUROLOGY, *Neurolog'ia*, (F.) *Névrologie*, (*neuron*, and *λογος*, 'a discourse.') The part of anatomy which treats of the nerves.

NEUROMA, gen. *Neuro'matis*, (F.) *Névrôme*, (from *neuron*.) A morbid enlargement of a nerve. Applied to subcutaneous, circumscribed, and highly painful tumours formed on the tissue of the nerves; and likewise to a small, hard, grayish tumour, *Neuroma'tion*, *Nervous ganglion*, *painful tubercle*, of the size of a pea, which is observed in the course of nerves, and appears to be formed from the neurilemma.

NEUROMALACIA, (*neuron*, and *μαλακία*, 'softening,') Softening of nerves.

NEUROMATION, Neuroma.

NEUROMATOID, *Neuromato'id*es, (*neuroma*, and *ειδος*, 'resemblance,') Resembling neuroma; as a '*neuromatoid* tumor.'

NEUROMETRÆ, (*νευρομετραί*), Psœ.

NEUROMYELITIS, (*neuron*, *μυελος*, 'marrow,' and *itis*.) Inflammation of the medullary matter of the nerves.

NEURON, (*νευρον*), see Nerve.

NEURONOSOS, *Neuronu'sos*, (*neuron*, and *νοσος*, 'a disease,') A disease of the nerves.

NEURONYG'MÊ, *Neuronyg'mus*, *Neuronyg'is*, (*neuron*, and *νυγμα*, 'puncture,') *Punctio ner'vi*. Puncture of a nerve.

NEUROPATHIC, *Neuropath'icus*, *Nervopath'ic*, (F.) *Névropathique*, (*neuron*, and *παθος*, 'affection,') Belonging to disease of the nerves, *Neuropath'ia*, *Neuron'osis*, *Neurop'athy*, (F.) *Névropathie*: see *Neuroses*.

NEUROPHLEGMONÊ, (*neuron*, and *φλεγμονη*, 'inflammation,') Neuritis.

NEUROPHLOGOSIS, (*neuron*, and *φλογωσις*, 'inflammation,') Neuritis.

NEUROPLAS'TY, *Neuroplas'tice*, (F.) *Névroplastique*, (*neuron*, and *πλαστικός*, 'forming,') M. Serres applies this term to a ganglionic alteration of the peripheral nerves.

NEUROPYRA, (*neuron*, and *πυρ*, 'fire, fever,') Fever, nervous, Typhus mitior.

NEUROPYRETUS, Fever, nervous, Typhus mitior.

NEURO'SES, (F.) *Névroses*, (*neuron*, and *osis*.) A generic name for diseases supposed to have their seat in the nervous system, and which are indicated by disordered sensation, volition, or mental manifestation; without any evident lesion in the structure of the parts, and without any material agent producing them. Such is the usual definition. Broussais attributes them to a state of irritation of the brain and spinal marrow.

NEUROSKELETON, see Skeleton.

NEUROSPASMI, see Spasm.

NEUROSTHEN'IA, (F.) *Névrosthénie*, (*neuron*, and *σθενία*, 'strength,') Excess of nervous irritation. Nervous irritation.

NEUROTHELÊ, (F.) *Névrothèle*, (*neuron*, and *θηλη*, 'a nipple,') A nervous papilla.

NEUROTHELIC APPARATUS, Corpus papillare.

NEUROTIC, Nervine, see Neurotica.

NEUROTICA. Diseases of the nervous function. The 4th class of Good's Nosology. *Neurotics*, in the classification of Dr. Taylor, are poisons, which belong to the narcotic and narcotico-acrid classes of Orfila. They are divided by him into the *Cerebral*, as morphia; the *Spinal*, as strychnia; and the *Cerebrospinal*, as conia and aconitia.

NEUROTOME, *Névroto'me*.

NEUROTOMY, *Neurotom'ia*, (*neuron*, and *τομή*, 'incision,') (F.) *Névrotomie*. Dissection of nerves. Also, an incised wound and section of a nerve.

NEUROTROSIS, *Neurotros'mus*, (*neuron*, and *τροσις*, 'wounding,') *Nervi vulneratio*. Wound or wounding of a nerve.

NEUROTROSMUS, Neurotrosis.

NEURYMEN, (*neuron*, and *μυρον*, 'a membrane,') Neurilemma.

NEURYMENITIS, Neurilemmitis.

NEUTA, *Neutha*. A membrane, according to Paracelsus, which covers the eyes and ears of the fœtus in utero.

NEUTRAL, (*neuter*, 'neither,') Belonging to neither in particular.

NEUTRAL SALTS, *Sec'ondary salts*, (F.) *Sels neutres*. Salts, composed of two primitive saline substances, combined together, and not possessing the characters of acid or of base.

NÉVIS, MIN'ERAL WATERS OF. Thermal springs in the department of Allier, France. Temperature, 136° to 148° Fahrenheit.

NÉVRAGMIE (F.), (*névro*, and *αγμας*, 'fracture,') The bruising or section of a nervous cord.

NÉVRALGIE, Neuralgia—*n. du Foie*, *Hepatalgia*—*n. des Reins*, *Nephralgia*—*n. de l'Utérus*, *Hysteralgia*.

NÉVRILÈME, Neurilemma.

NÉVRILÉMITE, Neurilemmitis.

NÉVRINE, Neurine.

NÉVRITE, Neuritis.

NÉVRITIQUE, Nervine.

NÉVRO (F.), (from *neuron*,) in composition, 'nerve.'

NÉVRODYNIE, Neuralgia.

NÉVROGRAPHIE, Neurography.

NÉVROLOGIE, Neurology.

NÉVRÔME, Neuroma.

NÉVROPATHIE, Neuropathy.

NÉVROPATHIQUE, Neuropathic.

NÉVROPLASTIE, Neuroplasty.

NÉVROSE DU CŒUR, Angina pectoris—n. Utérine, Hysteria.

NÉVROSES, Neuroses.

NÉVROSISME, Nervousness.

NÉVROSTHÉNIE, Neurosthenia.

NÉVROTHÈLE, Neurothela.

APPAREIL NÉVROTHÈLE. The papillary apparatus, or arrangement of the skin in the aggregate.

NÉVROTIQUE, Nervine.

NÉVROTOME, *Neurotome*, *Neurot'omus*; (from *neuro*, and *temno*, 'I cut,' 'I dissect.') An instrument used by anatomists to dissect the nerves. It is a long and very narrow scalpel, having two edges.

NÉVROTOMIE, Neurotomy.

NEVUS, Nævus.

NEW JERSEY, MIN'ERAL WATERS OF. In the upper part of Morris county and in the county of Hunterdon, near the top of Musconetcong mountain, there are chalybeate springs, which are resorted to. See Schooley's Mountain.

NEW YORK, MIN'ERAL WATERS OF. The chief waters are those of Saratoga, Ballston, Sharon, and Avon, which see. A spring at Catskill contains sulphates of magnesia, soda, and lime; carbonates of lime, magnesia, and iron, and chlorides of sodium, calcium and magnesium. The Stockport and Columbia springs contain sulphates of magnesia, lime, and soda; chlorides of magnesium and calcium; carbonates of magnesia and lime; organic matter and carbonic acid.

NEXUS STAMINEUS OCULI, ('stamineous connection of the eye,') Ciliary body.

NEZ, Nasus.

NGERENGERE. A singular disease, with the New Zealanders, which Dr. Arthur S. Thomson considers to be a variety of such diseases as the Elephantiasis Græcorum, Lepra of the Cossacks, &c., and which he proposes to call *Lepra gangræno'sa*.

NICANDRA ANOMALA, (after Nicander, a physician of Colophon,) *Anisodus luridus*.

NICARAGUA WOOD, see *Cæsalpinia*.

NIC'COLI SALES, *Salts of Nick'el*, (F.) *Sels de Nickel*. These salts are gentle tonics, and have been given with advantage in periodic headache, and in chlorosis and amenorrhœa. The dose of the sulphate is gr. ss or gr. j, three times a day.

NICCOLUM, Nickel.

NICE, CLIMATE OF. The climate of Nice possesses some advantages over the neighbouring climates of Provence and Italy, being free from the sirocco of the latter, and protected from the mistral of the former. Spring is the most unfavourable season. The climate is very dry. It has been a great winter retreat for the consumptive; but does not deserve the encomiums that have been passed upon it.

NICHE, see *Lingula*.

NICKAR TREE, *Gymnocladus Canadensis*.

NICK'EL, *Nic'olum*, (from [G.] *Nickel*, 'a prostitute,') A term of detraction, applied by the old German writers to what was regarded as a kind of false copper ore.) A white malleable metal, s. g. 8.8; less oxidizable than iron. The sulphate has been given as a tonic in the dose of half a grain to a grain.

NICKEL, SALTS OF, Niccoli sales.

NICODEMI O'LEUM, *Oil of Nicodemus*. An oil made by digesting, for some time, in a mixture of white wine and olive oil,—old turpentine, litharge, aloes, saffron, Oxyd of zinc, &c.

NICOTIANA, (after Jean Nicot, who first introduced it into Europe about 1560,) *N. tabacum*—n. Minor, *N. rustica*.

NICOTIANA RUS'TICA, *N. mi'nor*, *Priape'ia*, *Hyoscy'amus luteus*, *English* or *Wild Tobacco*, *Ord. Solanaceæ*. The leaves possess the properties of tobacco, but are milder.

NICOTIANA TABACUM or TAB'ACUM, *Nicotia'na*, *Her'ba taba'ci*, *Taba'cum*, *Pe'tum*, *Pe'tum*, *Hyoscy'amus Peruvia'nus*, *Tabac'co*, *Virgin'ia tobacco*, *Drunk'wort*, (F.) *Tabac*, *Nicotiane*. Tobacco is a violent acronarcotic; its properties seeming to depend upon a peculiar principle, *Nicotin* or *Nicotin'amin*. It is narcotic, sedative, diuretic, emetic, cathartic, and errhine. In incarcerated hernia, it is injected, in the form of smoke or infusion, but requires great caution. It is extensively and habitually used as an errhine and sialogogue. The infusion is used to kill vermin, and in some cutaneous eruptions.

NICOTIANE, *Nicotiana tabacum*.

NICOTIANIN, see *Nicotiana tabacum*.

NICOTIN, see *Nicotiana tabacum*.

NICTATIO, *Hippus*, *Nictation*.

NICTA'TION, *Nicta'tio*, *Nictita'tio*. (*nictare*, *nictatum*, 'to wink,') 'winking.' *Clo'nus nictita'tio*, *Blepharism'us*, *Palpebra'tio*, *Treink'ling of the eye*, (F.) *Clignotement*, *Cillement*. A rapid and repeated movement of the eyelids, which open and shut alternately. As occasionally performed, it is physiological; if repeatedly, a disease. It seems to be executed chiefly by the motor 7th pair of nerves; but it is necessary that the excitator 5th pair should likewise be in a state of integrity.

NIDAMENTUM, (*nidus*, 'a nest,') *Decidua*.

NIDDICK, *Nucha*.

NIDOREUX, *Nidorous*.

NIDOROUS, *Nidoro'sus*, (F.) *Nidoreux*, from *Ni'dor*, *Criv'sa*, 'the smell of any thing burnt,' &c. Having the smell of burnt or corrupt animal matter.

NIDUS, ('a nest,') *Nidus hirundinis*.

NI'DUS HIRUN'DINIS, *Ni'dus*, *Swa'llow's Nest*. A deep fossa in the cerebellum—so called from its fancied resemblance—situate between the velum medullare posterius in front, and the nodulus and uvula behind.

NIEDERBRONN, MIN'ERAL WATERS OF. Saline waters in the department of Bas-Rhin, France, which contain chloride of sodium, sulphate of magnesia, and carbonic acid.

NIELLE, *Nigella*.

NIEPA BARK, see *Samadera Indica*.

NIGELLA, *Coptis*.

NIGEL'LA, *N. Suti'va*; *Ord. Ranunculaceæ*; (*nigrella*, (from *niger*, 'black,') in allusion to the seed.) *Melan'thium*, *Cumi'num Ni'grum*, *Fen'nel flower*, *Nut'meg flower*, *Dev'il in a bush*, (F.) *Nigelle*, *Nielle*, *Faux cumin*, *Fleur Sainte-Catherine*. This small southern European and Syrian plant was formerly used medicinally as an expectorant and deobstruent, errhine, sialogogue, &c.

NIGELLE, *Nigella*.

NIGHTBLINDNESS, *Hemeralopia*.

NIGHT EYES, *Crusta genu equinæ*—n. *Mare Incubus*.

NIGHTSHADE, AMERICAN, *Phytolacca decandra*—*n.* Bittersweet, *Solanum dulcamara*—*n.* Common, *Solanum*—*n.* Deadly, *Atropa belladonna*—*n.* Enchanter's, *Circæa Lutetiana*—*n.* Garden, *Solanum*—*n.* Palestine, *Solanum sanctum*—*n.* Red, *Phytolacca decandra*—*n.* Stinking, *Hyoscyamus niger*—*n.* Vine, *Solanum dulcamara*—*n.* Woody, *Solanum dulcamara*.

NIGREDO À SOLE, ('blackness from the sun,') Ephemelids.

NIGRITA, Negro.

NIGRITIES, ('blackness,') *Melanopathia*—*n.* Ossium, Caries.

NIGRITISM, *Melanopathia*.

NIGRITUDO, gen. *Nigritudinis*, ('blackness,') *Melanosis*—*n.* Pulmonum, *Melanosis pulmonum*.

NIGROMANTIA, (*niger*, 'black,' and *μαντεία*, 'divination,') *Neeromancy*.

NIGROR, ('blackness,') *Melasma*.

NIGRUM OCULI, ('black of the eye,') Pupil.

NIHIL ALBUM, *Zinci oxydum*—*n.* *Griseum*, *Zinci oxydum*.

NILIACUM, (from Nilus, 'the Nile,') Mel.

NIMA QUASSIOIDES, *Simu'ba quassioïdes*, a native of Nepal and the Himalaya mountains, is as bitter as quassia, and is used like it.

NINDSIN, *Sium ninsi*.

NINE BARK, *Spiræa opulifolia*.

NINEDAY FITS, *Trismus nascentium*.

NINSI, *Sium ninsi*.

NINSING, *Panax quinquefolium*.

NINZIN, *Sium ninsi*.

NIPABLEPSIA, (*νίφα*, 'snow,' and *αβλεψία*, 'blindness,') *Snowblindness*.

NIPPLE, (Sax. *nýpele*.) The *Teat*, *Tit*, *Te'e*, *Dug*, *Pap*, (Old Eng.) *Nebble*, *Tit'ty*; *Acromas'tium*, *Uber*, *Staphyl'itis*, *Staphyl'ium*, *Bubo'na*, *Mam'milla*, *Mamm'ella*, *Papill'la*, *P. mam'mæ*, *Uberis apex*, *The'le*, *Tit'th'e*, *Tit'thos*, *Tit'thion*, *Tit'this*, (F.) *Mamelon*, *Bout de sein*. The conical tubercle, situate at the centre of the breast. Towards the central part of each breast the skin changes colour, and assumes a rosy tint in young females, or a reddish brown in those who have suckled several children. The circle is called the *Are'ola* or *Aure'ola* of the nipple. The nipple is capable of erection on being excited.

NIPPLE, ARTIFICIAL, *Bout de Sein*.

NIPPLEWORT, *Lapsana*.

NIRLES, *Herpes phlyctænodes*.

NIRMALLEES, *Strychnos potatorum*.

NIRMILLIES, *Strychnos potatorum*.

NI'SUS, *Nix'us*, *Pei'ra*, *Strain'ing*, (*nitor*, *nisus*, and *nixus*, 'to endeavour,') A voluntary retention of the breath, so as to force down the diaphragm; the abdominal muscles being at the same time contracted forcibly. In this manner the contents of the abdomen are compressed; and the evacuation of the fæces, urine, &c., is effected.

NISUS, Effort—*n.* *Formativus*, *Plastic force*.

NISUS PARTURIEN'TIUM. The efforts or forcing during parturition.

NITEDULA, (*niteo*, 'to shine,') *Cicindela*.

NITON, MIN'ERAL WA'TERS OF. These springs in the Isle of Wight contain iron, and sulphate of alumina and potassa.

NITRAS, gen. *Nitra'tis*, (from *nitre*,) Nitrate—*n.* *Hydrargyricus*, see *Hydrargyri nitras*—*n.* *Hydrargyrosus*, *Hydrargyri nitras*—*n.* *Kalicus*, *Potassæ nitras*—*n.* *Lixiviæ*, *Potassæ nitras*—*n.* *Natrium*, *Soda*, *nitrate* of—*n.* *Plumbicus*, *Plumbi nitras*—*n.* *Potassæ cum sulphure fusus*, *Potassæ nitras fusus sulphatis paucillo mixtus*—*n.* *Subbismuthicum*, *Bismuth*, *sub-nitrate* of.

NITRATE, *Ni'tras*, *Azo'tas*, *A'zotate*. A salt, formed of a base and nitric acid. Several nitrates are employed in medicine.

NITRATE ACIDE DE MERCURE, see *Hydrargyri nitras*—*n.* *d'Ammoniaque*, *Ammonias nitras*—*n.* *d'Argent*, *Argenti nitras*—*n.* *Mercureux*, *Hydrargyri nitras*.

NITRE, *Potassæ nitras*—*n.* *Cubic*, *Soda*, *nitrate* of.

NITRIC, *Ni'tricus*, (F.) *Nitrique*, (from *nitrum*, 'nitre,') Appertaining or relating to nitre.

NITRIC AC'ID, *Ac'idum ni'tricum* seu *azo'ticum* seu *sept'icum* seu *ni'tri*, *A'qua for'tis*, *Ni'trous ac'id*, *Spir'itus ni'tri du'plex* seu *ni'tri fu'mans* seu *ni'tri Glauberi* seu *ni'tri ac'idus*; *Azot'ic acid*, *Dephlogis'ticated nitrous acid*, *Oxysepton'ic acid*, (F.) *Acide nitrique* ou *azotique*. Nitric acid is obtained from nitre—*Ni'trate of potas'sa*. Its odour is suffocating; taste very acid and caustic. It is corrosive, liquid, colourless, and transparent. S. g. 1.42. Ph. U. S.

Strong nitric acid is rarely used except as an application to foul, indolent ulcers, or to warts. When given internally, it is in the form of the *Acidum ni'tricum dilu'tum*, *Spir'itus ni'tri simplex* seu *ni'tri vulga'ris*—*Dilu'ted nitric acid*, which, in the Pharmacopœia of the United States, (1863,) consists of *nitric acid*, three Troy ounces, *water*, a pint. Diluted largely with water, it is used, as a drink, in fevers of the typhoid kind; in chronic affections of the liver, syphilis, &c.; but in the latter affections, it is not to be depended upon.

NITROGEN, (*nitre*, and *γεννω*, 'I generate,') *Azote*.

NITROGEN, GAS'EUS OX'IDE OF, *Ni'trous ox'ide*, *Protox'ide of nitrogen* or of *azote*, *Par'adise*, *In'tox'icating* or *Laughing gas*, *Dephlogis'ticated ni'trous gas*, *Gas azo'ticum oxygenatum*. This gas, when respired, produces singular effects; great mental and corporeal excitement; and, generally, so much exhilaration as to render the appellation, "laughing gas" by no means inappropriate. It has not been much used in medicine, although recommended in paralysis. Its effects are too violent and too transient to render it a valuable remedial agent; and, in the delicate, it has been productive of unpleasant effects, inducing palpitation, fainting, and convulsions.

NITROGEN, PROTOXIDE OF, Nitrogen, gaseous oxide of.

NITROGÈNE, *Azote*.

NITROGENIUM, *Azote*.

NITROGENIZED, *A'zoted*, *A'zotized*. (F.) *Azoté*. Containing nitrogen or azote;—as a *nitrogenized*, *azoted*, or *azotized* aliment, animal food, for example,—in contradistinction to the *nonnitrogenized*, *nonazoted*, or *nonazotized*, (F.) *Nonazotés*, *anazotiques*, or those that do not contain nitrogen;—saccharine and oleaginous aliments, for example.

NITROGLYCERIN, *Glonoin*.

NITROMURIATIC AC'ID, *Ac'idum nitro-muriaticum* seu *muriaticum ni'troso'cygena'tum*, *Mens'trium au'ri*, *Nitrohydrochlor'ic* or *Hydrochloroni'tric acid*, *Aqua re'gia* seu *styg'ia* seu *re'gia*, *Chrysæalea*, (F.) *Eau régale*. A mixture of the nitric and muriatic acids has been used in diseases of the liver, in the form of a bath for the feet and legs, made sharply acidulous. It has, also, been employed, and with more uniform results, in cutaneous affections. Whatever advantages it may possess in internal diseases—and these advantages are doubtful—they are probably dependent upon the chlorine formed by the mixture, or upon the properties possessed by the mineral acids in general. See *Chlorine*.

The *Acidum nitromuriaticum* of the Pharmacopœia of the United States (1863) is formed by mixing three Troy ounces of *nitric acid* with five of *muriatic acid*.

Acidum Nitromuriaticum dilu'tum, *Dilu'ted Ni-*

nitromuriatic Acid is made of *nitric acid*, a Troy ounce and a half; *muriatic acid*, $\frac{3}{4}$ üss; *Distilled water*, Oj.

NITROUS OXIDE, Nitrogen, gaseous oxide of.

NITRUM, (*νιτρον*.) *Natron*, *Potassæ nitras*—*Antiquorum*, *Soda*—*n. Cubicum*, *Soda*, *nitrate of*—*n. Factitium*, *Borax*—*n. Flammans*, *Ammoniæ nitras*—*n. Lunare*, *Argentii nitras*—*n. Prismaticum*, *Potassæ nitras*—*n. Rhomboidale*, *Soda*, *nitrate of*—*n. Saturninum*, *Plumbi nitras*—*n. Tabulatum*, *Potassæ nitras fusus sulphatis paucillo mixtus*—*n. Vitriolatum*, *Potassæ sulphas*—*n. Vitriolatum Schroederi*, *Potassæ sulphas*.

NIX, gen. *Nivis*, *Snow*—*n. Fumans*, *Calx viva*—*n. Zinei*, *Zinci oxydum*.

NIXUS, *Nisus*—*n. Parturientium*, *Nisus parturientium*.

NOAH'S ARK, *Cypripedium luteum*.

NO'BLE, *No'bilis*. Principal, Essential.

NO'BLE PARTS, *Par'tes no'biles seu essential'es*.

Some anatomists have given this name to parts, without which life cannot exist; such as the heart, liver, lungs, brain, &c. The organs of generation have, likewise, been so called.

NO'CAR, (*νωκαρ*.) *Tor'por Soporif'icus*. *Lethargie torpor*.

NOCTAMBULATIO, (*nox, noctis*, 'night,' and *ambulare*, *ambulatum*, 'to walk,') *Somnambulism*.

NOCTAMBULISMUS, *Somnambulism*.

NOCTAMBULUS, *Somnambulist*.

NOCTISURGIUM, (*nox, noctis*, 'night,' and *urgere*, 'to arise,') *Somnambulism*.

NOCTU'NI OC'ULI, ('owl's eyes,' from *noctua*, 'an owl,') *Gray or blue eyes*.

NOCTURNA BELLA, ('nocturnal wars,') *Coitus*.

NOCTUR'NAL, *Noctur'nus*, *Nycter'inus*, (from *nox*, gen. *noctis*, 'night,') Relating to night, as *Febris noctur'na seu nycter'ina*. A fever occurring in the night.

NODDING, *Annuitio*, *Nutatio*.

NODDLEPOX, *Syphilomania*.

NODE, *No'dus*, *Emphy'ma exosto'sis perios'tea*, (F.) *Nodosité*. *Nœud*. A hard concretion or incrustation, which forms around joints attacked with rheumatism or gout. Some include, under this name, exostoses, articular calculi, ganglions, and even the chronic swellings of the joints known under the name of white swellings.

NODES, (*νωδός*, from *νη*, priv. and *οδός*, 'a tooth,') *Edentulus*.

NODI DIGITORUM MANUS, *Phalanges of the fingers*—*n. Hæmorrhoidales*. see *Hæmorrhoids*—*n. Lactea*, *Infarctus mamæ lacteus*—*n. Nervorum*, *Ganglions*, *nervous*.

NODOI, (*νωδοί*, sing. *νωδός*, from *νη*, priv. and *οδός*, 'a tooth,') *Nefrendes*.

NODO'SIS, (from *nodus*, and *osis*.) The formation or existence of nodes.

NODOSITÉ, *Node*.

NOD'ULI ARAN'TII, (pl. of *nodulus*.) *Noduli Morgagnii*, *Cor'pora Aran'tii*. The small sessoid bodies situate on the periphery of the semilunar valves of the aorta and pulmonary artery for the better occlusion of the artery.

NODULI MORGAGNII, *Noduli Arantii*—*n. Nervorum*, *Ganglions*, *nervous*.

NOD'ULUS, (dim. of *nodus*, 'a knot,') A small knot. A small prominence or lobule in the portion of the cerebellum, which forms the posterior boundary of the fourth ventricle. The nodulus is on the median line, and before the uvula. See *Vermiform process*, inferior.

NODULUS UTERINI, *Pessary*.

NO'DUM RELIGA'RE, ('to fasten a knot or tie,') (F.) *Nouer l'Aiguillette*. To prevent by enchantment the venereal act; or to render impotent!

NODUS, ('a knot or tie,') *Articulation*, *Edentulus*, *Encystis*, *Hæmma*—*n. Cerebri*, *Pons Varolii*—*n. Chirurgicus*, *Knot*, *surgeon's*—*n. Encephali*, *Nœud de l'Encéphale*, *Pons Varolii*—*n. Hystericus*, *Angone*—*n. Vitæ*, *Centrum vitale*.

NOEMA, (*νοημα*, from *νοεω*, 'to think,') *Thought*.

NOESIS, (*νοησις*.) see *Intellect*.

NŒUD, ('a knot,') *Node*—*n. du Chirurgien*, *Knot*, *surgeon's*—*n. d'Emballeur*, *Knot*, *packer's*.

NŒUD DE L'ENCEPHALE (F.), *Nodus Enceph'ali*. *M. Cruveilhier*, under this name, includes the pons Varolii, peduncles of the cerebrum and cerebellum, and the tubercula quadrigemina.

NŒUD VITAL, ('vital knot,') *Centrum vitale*.

NŒUDS, (pl. of *nœud*), *Calculi*, *arthritic*.

NOIR (from [L.] *niger*, 'black,') **ANIMAL**, ('animal black,') *Carbo animalis*—*n. d'Os*, *Carbo animalis*.

NOISETIER, *Corylus avellana*.

NOISETTE, (dim. of *noix*.) *Corylus avellana* (nut.)

NOIX, ('a walnut,' from *nux*, gen. *nucis*, 'a nut,') *Juglans regia* (*nux*)—*n. Américaine*, *Jatropha curcas*—*n. des Barbades*, *Jatropha curcas*—*n. Cathartique*, *Jatropha curcas*—*n. d'Eon*, *Trapa natans*—*n. de Galle*, see *Quercus infectoria*—*n. d'Igaur*, *Ignatia amara*—*n. de Serpent*, see *Thevetia Ahouai*—*n. Vomique*, see *Strychnos nux vomica*.

NOLI ME TANGERE, ('do not touch me,') *Lupus*, see *Chancreux* (*Bouton*).

NOMA, (*νομα*, from *ναιω*, 'to corrode,') *Cancer aquaticus*—*n. Pudendorum*, *Colpoceae infantilis*.

NO'MAD, *Nomade*, *No'mas*, (*νομας*, gen. *νομάδης*, from *νομα*, 'pasturage,') An epithet given to people who have no fixed habitation, and who travel, with their flocks, from country to country, for pasturage. Such are the Tartars. By analogy, the word *Nomad'ic* has been applied to a spreading ulcer.

NOMBRIL, *Umbilicus*—*n. de Venus*, *Cotyledon umbilicus*.

NOME, (*νομη*), *Cancer aquaticus*, *Phagedænia ulcer*.

NO'MENCLATURE, *Nomenclat'io*, *Nomenclatura*, *Onomatolog'ia*, *Onomatocle'sis*, *Terminolog'ia*, *Terminolog'ia*, *Glossolog'ia*, *Orismatolog'ia*, (*νομα*, 'name,' and *καλεω*, 'I call,') A collection of terms or words peculiar to a science or art. In all sciences, nomenclature is an object of importance; and each term should convey to the student a definite meaning. The Lavoisierian nomenclature was a valuable gift to chemistry; and anatomy has derived advantage from the labours of Barclay, Dumas, and Chaussier, who have given names to parts indicative of their situation. See *Muscle*. The nomenclature of pathology has required the greatest attention; and although repeated attempts have been made to improve it, the barbarous terms that disgrace it are still frequently adopted. It consists of Hebrew and Arabic terms; Greek and Latin, French, Italian, Spanish, German, English, and even Indian, African, and Mexican; often barbarously and illegitimately compounded. A want of principle in forming the technical terms of medicine is everywhere observable. They have been formed for example, — 1. From colour; as *Melanæ*, *Melas*, *Atrabilis*, *Leuce*, *Alphos*, *Chlorosis*, *Rubeola*, *Scarlatina*, *Purpura*, &c. 2. From duration; as *ephemeral*, *quidians*, *tertians*, and *quartans*, *continued*, and *intermittent*, &c. 3. From Birds, Beasts, Fishes, Insects, and Plants; as *Rabies canina*, *Cynarcho*, *Boulimia*, *Pica*, *Hippanthropia*, *Elephantiasis*, *Urticaria*, *Lichen*, *Ichthyosis*, &c. 4. From Persons or Places; as *Morbus Heruleus*, *Favien Hippocratica*, *Lepra Arabum*, *Plica Polonica*, *Sudor Anglicus*, *Morbus Gallicus*, *Ignis Sancti Antonii*, *Chorea Sancti Viti*, &c.

NOMUS, (νομος, from νέμω, 'to corrode,') Cancer aquaticus.

NONAZOTED, see Nitrogenized.

NONCOMPOS, Lunatic.

NONCOMPOS MENTIS, ('not possessing mind,') Lunatic.

NONNAT'URALS, *Nonnaturalia*. Under this term the ancient physicians comprehended air, meat, and drink, sleep and watching, motion and rest, the retentions and excretions, and the affections of the mind. They were so called, because they affect man without entering into his composition, or constituting his nature; but yet are so necessary that he cannot live without them.

NONNITROGENIZED, see Nitrogenized.

NONPLACENTAL, see Placental.

NONSANE, Insane.

NONSTRATED MUSCULAR FIBRE, see Muscular fibre.

NONVIABLE (F.) (from *non*, and *viable*.)

Applied to a new-born child not capable of continued existence. See Viable.

NONUS HUMERI PLACENTINI, (*nomus*, 'the ninth,') Teres minor—n. Linguae musculus, Genioglossus—n. Vesalii, Peronæus tertius.

NOODLES, see Turundæ Italicæ.

NOOS, (νοος,) Intellect.

NORICE, ([F.] *Nourrice*.) Nurse.

NORMA VERTICALIS, ('vertical rule,') see Normal.

NORMAL, *Norma*'lis, (*norma*, 'a perpendicular,') 'a rule,') According to rule; perpendicular.

The *normal line*, *norma verticalis*, of Blumenbach, is a vertical line let fall from the prominence of the frontal bone and shaving the superior maxillary, so as to mark the projection of the latter bone beyond the arch of the forehead.

NORRIS'S DROPS, see Antimonium tartarizatum.

NORTH CAROLINA, MIN'ERAL WA'TERS OF. The chief springs are the *Warm* and *Hot Springs* of Buncombe. Temperatures varying from 94° to 104° Fah. They have not much mineral impregnation, and have the properties of the thermal waters in general.

NORTON'S DROPS, Liquor hydrargyri oxy-muriatis.

NORYSS, ([F.] *Nourrice*.) Nurse.

NOSACERUS, (νοσάκερος,) Sickly.

NOSE, Nasus,—n. Bleed, Epistaxis.

NOSE, RUNNING AT THE, Coryza.

NOSELIA, (νοσηλία,) *Nosocomie*'sis, *Nosocomi*'a. Care of the sick.

NOSEMA, (νοσημα,) gen. *Nose*'matis, Disease.

NOSENCEPH'ALUS, (F.) *Nosencephale*, (*nosos*, and *εγκεφαλος*, 'brain,') A monster whose skull is open only on the frontal and parietal regions, the posterior fontanelle being distinctly present.

NOSERUS, (νοσηρος,) Insalubrious, sick.

NOSEUMA, gen. *Noseu*'matis, (νοσηυμα,) Disease.

NOSOCHTHONOGRAPHIA, (*nosos*, χθων, 'the earth,' and γραφή, 'to describe,') Geography, medical.

NOSOCHTHONOLOGIA, (*nosos*, χθων, 'the earth,' and λογος, 'a description,') Geography, medical.

NOSOCOMESIS, (νοσοκομσις, from *nosos*, and *κομω*, 'to care for,') Noselia.

NOSOCOMIA, Noselia.

NOSOCO'MIAL, *Nosocomia*'lis, (from *Nosocomium*.) Relating to an hospital,—as '*nosocomial* or hospital fever.'

NOSOCOMIUM, (νοσοκομειον,) Hospital.

NOSOCOMUS, (νοσοκομος,) *Infirmier*.

NOSOCRATIC, (*nosos*, and *κρατος*, 'power,') Specific.

NOSOCRINIES, (*nosos*, and *κρνω*, 'I separate,') Diacrisis.

NOSODES, (νοσώδης, from *nosos*, and *odes*,) Insalubrious, Sick, Sickly.

NOSODOCHIUM, (νοσοδοχιον, from *nosos*, and *δεχομαι*, 'to receive,') Hospital.

NOSOGENESIS. (*nosos*, and *γενεσις*, 'generation,') Pathogeny.

NOSOGENIA, (*nosos*, and *γεννω*, 'to engender,') Pathogeny.

NOSOGENY, Pathogeny.

NOSOGEOGRAPHY, Geography, medical.

NOSOG'RAPHY, *Nosograph*'ia, (F.) *Nosographie*, (*nosos*, and *γραφω*, 'I describe,') A description of diseases.

NOSOHÆMIA, (*nosos*, and *αίμα*, 'blood,') *Hæmatonosus*.

NOSOHÉMIE, *Hæmatonosus*.

NOSOL'OGISM, *Nosologis*'mus. The abuse of nosological arrangements in pathology.

NOSOL'OGY, *Nosolog*'ia, (F.) *Nosologie*, (*nosos*, and *λογος*, 'a discourse,') A name given to that part of medicine whose object is the classification of diseases. The most celebrated nosological systems have been those of Sauvages (1763), Linnæus (1763), Vogel (1764), Sagar (1776), Macbride (1772), Cullen (1772), Darwin (1796), Selle, Crichton (1804), Parr (1809), Swediaur (1812), Pinel (1813), Young (1813), Good (1817), Hosack (1818), &c. Besides these general nosographies, others have been published on *Surgery* exclusively, none of which are particularly worthy of enumeration amongst nosological systems. Nosological arrangements have, also, been formed of single families or groups of diseases. Plenck, of Baden, is the author of two different treatises of this kind: the one, a methodical arrangement of the diseases of the eyes, and the other, of cutaneous diseases. Dr. Willan published an arrangement of cutaneous diseases, which was completed by Dr. Bateman, and adopted into the *Nosology* of Dr. Hosack. Mr. Abernethy, also, published a methodical classification of tumours, and many other partial nosological classifications might be enumerated. Also, *Pathology*.

NOSON'OMY, *Nosonom*'ia, (*nosos*, and *ονμα*, 'name,') The nomenclature of diseases.

NOSOPHYTA, (*nosos*, and *φυτον*, 'a plant,') Diseases supposed to be produced by, or to consist in the development of parasitic plants—as porrigio, mentagra, &c.

NOSOPHYTODERM'ATA, (from *nosophyta*, and *derma*, 'skin,') *Nosophyta* affecting the skin. Mr. Erasmus Wilson reckons five of these—*Favus*, *Trichosis*, *Mentagra*, *Alopecia areata*, and *Chloasma*.

NOSOPH'US, *Nosopoiēt*'icus, (F.) *Nosopoiétique*, (*nosos*, and *ποιω*, 'I make,') That which causes disease.

NOSORGANIES, (*nosos*, and *οργανον*, 'an organ,') Organic diseases.

NOSOS, (νοσος,) Disease.

NOSOTAXY, *Nosotax*'ia, (*nosos*, and *ταξις*, 'arrangement,') The distribution and classification of diseases.

NOSOTHEO'RIA, (*nosos*, and *θεωρια*, 'doctrine,') The doctrine or theory of disease.

NOSTAL'GIA, (νοστος, 'a journey home,' and *αλγος*, 'pain,') *Nostoma*'nia, *Nostras*'sia, *Apodemial*'gia, *Pathopatridal*'gia, *Pothopatridal*'gia, *Philopatridal*'gia, *Philopatridomania*, *Ademon*'ia, *Ademos*'yne, *Homesickness*, (F.) *Nostalgie*, *Maladie du pays*. An affection produced by the desire of returning to one's country. It is commonly attended by slow wasting, and sometimes by hectic, which may speedily induce death. M. Pinel properly regards it as a variety of melancholy.

NOSTOMANIA, (νοστος, 'a journey home,' and *μανια*, 'mania,') *Nostalgie*.

NOSTRASSIA, (*nostras*, 'of our country,') Nostalgia.

NOSTRILS, ([Sax.] *nær-ðýnl*, 'nosehole,') Nares.

NOSTRUM, (*nostrer*, 'our own,') Arcanum.

NOSTRUM, CHITTICK'S. An empirical remedy for stone in the bladder, said to be a solution of alkali in veal-broth.

NOTA MATERNA, ('mother's mark,') *Nævus*—n. Primitiva, Groove, primitive.

NOTÆ INFANTUM, ('infants' marks') *Nævi*.

NOTAL, (from *notus*,) Dorsal.

NOTAL'GIA, *Notal'gia*, (*Notalgia*, improperly, of Köchlin and others,) (*νωτος*, 'back,' and *αλγος*, 'pain,') Pain in the back. Spinal irritation.

NOTANENCEPHALIA, (*νωτος*, and *anencephalia*,) see *Notencephalus*.

NOTCH, (Teut. *No che*, Ital. *Nocchia*,) *Emarginatio*, *Emarginatu'ra*, (F.) *Échancrure*. A depression or indentation of different shape and size, observed on the circumference or edges of certain bones.

NOTCH OF THE CONCHA, *Incisura tragica*.

NOTCH, ETHMOID'AL, (F.) *Échancrure ethmoïdale*, is situate on the frontal bone, and joins the ethmoid.

NOTCHES, ISCHIAT'IC, (F.) *Échancrures Ischiatiques*, are two in number:—the greater and the less. The former is large, situate at the inferior part of the pelvis, and formed by the sacrum and ilium. It gives passage to the sciatic nerve, pyramidalis muscle, and to the superior gluteal vessels and nerves. The latter is much smaller than the other, from which it is separated by the sciatic spine. It gives passage to the tendon of the obturator internus, and to the internal pudic vessels and nerves.

NOTCH, PAROT'ID, (F.) *Échancrure parotidienne*, is the triangular space comprised between the parotid edge of the inferior maxillary bone and the mastoid process; so called, because it lodges the parotid gland. The notches in soft parts are generally called *Fissures*.

NOTCH, SCAP'ULAR, *Incisura scapularis*, *Lu'nula scap'ular*, (F.) *Échancrure scapulaire*. The notch on the superior edge or costa of the scapula, which is converted into a foramen by means of a ligament, and gives passage to the suprascapular nerve.

NOTCH, SEMILUNAR OF THE STERNUM, *Fourchette*.

NOTENCEPHALIA, see *Notencephalus*.

NOTENCEPH'ALUS, (*νωτος*, 'the back,' and *εγκεφαλος*, 'the brain,') A monster whose head, with the brain, is on the back. The condition is termed *Notencephalia* and *Notanencephalia*.

NOTHEUSIS, (*νοθευσις*,) Degeneration.

NOTHIA, (*νοθεια*,) Degeneration.

NOTHROTES, (*νωθροτες*,) Torpor.

NOTHUS, (*νοθος*,) False.

NOTIÆUS, (*νωτιας*,) Dorsal—n. *Myelus*, *Medulla spinalis*.

NOTIALGIA, *Notalgia*.

NOTOMÈLE, (*notus*, and *μελος*, 'unprofitable,') A monster with superfluous flesh on the back.

NOTOMYELITIS, (*notos*, and *myelitis*,) *Myelitis*.

NOTOMYELUS, (*notos*, and *μυελος*, 'marrow,') *Medulla spinalis*.

NOTUS, (*νωτος*,) Dorsum, Vertebral column.

NOUË, (*Bandage*) (F.); 'knotted,' (from *nœud*, (L.) *nodus*, 'a knot,') A bandage which has a considerable number of knots placed over each other. It is made with a roller, 6 or 7 ells long, rolled into two balls, and is used to compress the

parotid region, after the extirpation of the parotid gland.

Also, an epithet applied to children in whom the disease of rickets has swollen the articulations.

It is, likewise, applied to the gout when it has caused nodes on the joints.

NOUER L'AIGUILLETTE, ('to knot the aiguillette,') *Nodum religare*.

NOUËT (F.), *Nod'ulus*. A bag filled with medicinal substances, infused in a liquid to communicate their properties to it.

NOUFFER'S, MADAME, REMEDY, *Polypodium filix mas*.

NOURRICE, Nurse.

NOURRICIER, Nutritious.

NOURISSON, Nursling.

NOURRITURE, Aliment.

NOUS, (*vous*,) Intellect.

NOUËRE, (from *nouer*, 'to knot,') *Rachitis*.

NOUVEAU-NÉ (F.), 'newborn,' *Neonatus*, *nuperrimè Natus*, *Neog'enes*, *Neog'iles*, *Neog'ilus*, *Neog'onus*, *Neog'nus*, *Infans recens natus*. That which has been just born. A newborn infant.

NOVACULA, Razor.

NOW ([Sc.], from [Sax.] *enolle*, 'the top of a hill,') Vertex.

NOYAU, (from *noix*, 'a nut,') see *Cytoblast*—n. *Cancéreux*, see *Cancer cell*.

NOYAU CENTRAL DES PÉDONCULES DU CERVELET, *Corpus dentatum*.

NOYER, (from *noix*, 'a nut, a walnut,') *Juglans regia*—n. *de Ceylon*, *Adhatoda*.

NOYRIS, Nurse.

NUAGE, (from [L.] *nubes*, 'a cloud,') *Nebula*—n. *de la Cornée*, *Caligo*.

NUBECULA, (dim. of *nubes*,) *Enæorema*, *Nebula*.

NUBES, ('a cloud,') *Enæorema*, *Nebula*.

NU'BILE, *Nu'bilis*, *Marriageable*, *Marriageable*, 'fit to marry.' Generally, the period of puberty is considered to be the age at which both sexes are *nubile*. They are truly *nubile*, however, only when they are fitted to procreate healthy and vigorous children, and are competent to discharge their duties as parents.

NUBILITY, *Nubil'itas*, (F.) *Nubilité*. The state of being *nubile* or marriageable.

NUBILOUS, (*nubes*, 'a cloud,') *Nepheloid*.

NUCES (pl. of *nux*, 'a nut,') AQUATICÆ, ('water nuts,') see *Trapa natans*—n. *Quercus*, see *Quercus alba*.

NU'CHA, *Nu'cium*, *Nucha cap'itis*, (F.) *Nuque*. The nuke, hinder part or nape of the neck, (Prov.) *Niddick*. The part where the spinal marrow begins.

Ligamentum Nuchæ. A strong ligament from the neck, proceeding from one spinous process to another, and inserted into the occipital bone. It is very strong in quadrupeds. It is called in them *Poxyceax*, *Paxwax*, *Packwax*, formerly, *Faxwax*, and *Taxwax*.

NU'CHAL, *Nucha'lis*, (from *nucha*,) Relating or appertaining to the nucha or nape of the neck.

NUCHAL RE'GION, *Re'gio nu'chæ seu nucha'lis seu occipita'lis infe'rior*. The region of the nucha or nape of the neck.

NUCISTA, (from *nux*, gen. *nucis*, 'a nut,') see *Myristica moschata*.

NUCK, CANAL OF. A small prolongation of the peritoneum often sent into the inguinal canal of the female fœtus. So called, from Nuck, who first described it.

NU'CLEATED, *Nucleatus*, (F.) *Nucléé*, (*nucleus*, 'a kernel,') Having a nucleus or central particle. Applied to the elementary cells of organized tissues; the vital properties of which are seated in the nucleus. See *Cytoblast*.

NUCLEATED CELL, see Cytoblast.

NUCLEI CEMBRIÆ, see Pinus Cembra — n. Ossei, Ossification, points of — n. Pineæ, see Pinus pinea.

NUCLEIN, see Globules of the blood.

NUCLÉOLE, see Cytoblast.

NUCLÉOLULE, see Cytoblast.

NUCLEOLUS, (dim. of *nucleus*), see Cytoblast.

NUCLEUS, 'a kernel,' see Cytoblast — n. Blastodermatis, *Tache embryonnaire* — n. Centralis, Corpus dentatum — n. Cicatriculæ, *Tache embryonnaire* — n. Dentatus, Corpus dentatum — n. Dentis, Dental pulp — n. Encased, Cytoblast — n. Fimbriatus, Corpus dentatum — n. Furunculi, see Furunculus — n. Germinal, see Molecule — n. Germinativus, see Molecule — n. Olivæ, Corpus dentatum — n. Ossificationis, Ossification, point of — n. Rhomboidalis, see Corpus dentatum.

NUCULA TERRESTRIS, (dim. of *nux*, gen. *nucis*, 'a nut,') Bunium bulbocastanum.

NUKE, Nucha.

NULE-KNEED, Entogonyancœon.

NUMERICAL METHOD, see Method, numerical.

NUM'MULAR, *Nummularis*, *Num'mulated*, (from *nummus*, 'money,') Relating to money. An epithet applied to the sputa in phthisis, when they flatten at the bottom of the vessel, like a piece of money; and to the mode of aggregation of the red corpuscles of the blood — '*nummular* rolls.'

NUMMULARIA, *Lysimachia nummularia*.

NUMPOST, Abscess.

NUPHAR AD'VENA, (Arab.) *Nymphaea ad'vena*, *Yellow Pondlily*, *Spatterdock*. Order, *Nymphaeaceæ*; indigenous in Canada and the United States. The root is a popular tonic, and the fresh root bruised is applied as a cataplasm.

NUPHAR LUTEUM, *Nymphaea lutea*.

NUQUE, Nucha.

NURICE, Nurse.

NURSE, ([Sax.] *noþice*), *Nutrix*, (from *nutrio*), *Titheñe*, *Lactans*, *Lacticans*, *Nutriciens*, *Thelactria*, *Mam'ma*, (I.) *Nutrice*, (F.) *Nourrice*, (Sc.) *Noyria*, *Noryss*, *Nurice*, *Norice*. One who suckles her own child or another's. Also, one that has the care of a sick person, *Caretaker*, (F.) *Garde-malade*.

NURSE, DRY. One who gives every care to a child, but does not suckle it.

NURSE, MONTH'LY. One who attends a woman during the month of her parturient state.

NURSE, WET, (Sc.) *Milkwoman*. A female, who suckles the child of another.

To 'nurse artificially,' is to bring up a child by the hand.

NURSING, (from *nurse*), (F.) *Nourrison*. An infant nursed by its mother or by another.

NUS, (*vous*), Intellect.

NUSUS, (*νοσος*), Disease.

NUT ([Sax.] *hnut*, [L.] *nutr.*) BARBADOES, see *Jatropha curcas* — n. Brazilian, see *Pichurim beans* — n. Butter, *Juglans cinerea* — n. Cembros, see *Pinus cembra* — n. Clearing, *Strychnos potatorum* — n. French, *Juglans regia* — n. Ground, *Apios tuberosa*, *Arachis hypogea* — n. Para, see *Pichurim beans* — n. Physic, *Jatropha curcas* — n. Pine, see *Pinus picea* — n. Pistachio, see *Pistacia vera* — n. Poison, see *Strychnos nux vomica* — n. Purg- ing, see *Jatropha* and *Croton tiglium* — n. Rattle, *Nelumbium luteum* — n. Sassafras, see *Pichurim beans* — n. Soap, see *Sapindus saponaria*.

NUT TREE, MALABAR, *Adhatoda*.

NUT, VOMIC, see *Strychnos nux vomica* — n. Zibel, see *Pinus picea*.

NUTATION, *Nutatio*, (*nutare*, *nutatum*, 'to nod,') Nodding. Also, a constant oscillation of the head, by which it moves involuntarily in one or more directions.

NUTATOR CAPITIS, ('nodder of the head,') *Sternocleidomastoideus*.

NUTMEG, (*nux moscata*), see *Myristica moscata* — n. Butter, see *Myristica* — n. Flower, *Nigella* — n. Liver, *Liver*, *nutmeg*.

NUTMEGS, *Testes*.

NUTRICATIO, (*nutrio*, 'to nourish,') *Nutri-*tion.

NUTRICIER, *Nutritious*.

NUTRICIUS, *Nutritious*.

NUTRIENS, *Aliment*, *Nurse*.

NUTRIMEN, *Aliment*.

NUTRIMENTAL, *Nutrimen'talis*, *Aliment'al*, *Alimen'tary*, *Alimen'talis*, (F.) *Alimenteux*. Having the qualities of food or nutriment.

NUTRIMENTUM, *Aliment*, *Pabulum*.

NUTRITIF, *Nutritive*.

NUTRITION, *Nutrit'io*, *Nutrica'tio*, *Nutri-tus*, *Alitu'ra*, *Threp'sis*, (*nutrire*, *nutritum*, 'to nourish,') Nutrition is that function by which the nutritive matter already elaborated by the various organic actions, loses its own nature, and assumes that of the different living tissues, — to repair their losses and maintain their strength. Sometimes the word is used in a more extended signification, to express the whole series of actions by which the two constant movements of composition and decomposition are accomplished, in organized bodies. Nutrition, then, would comprehend digestion, absorption, respiration, circulation, and assimilation; the latter being *nutrition*, properly so called, and being effected in the intermediate system over the whole of the body, — the cells of the tissues attracting from the blood the elements necessary for their repa-ration.

NUTRITION, FORCE OF, *Plastic force*.

NUTRIT'IONAL, *Nutritionalis*. Relating or appertaining to nutrition.

NUTRIT'IOUS, *Nutric'ius*, *Nutritive*, *Alib'ilis*, *Troph'imus*, *Tropho'des*, (F.) *Nourricier*, *Nutricier*, *Alibile*. Having the quality of nourishing: as nutritious food, nutritious lymph, &c.

NUTRITIOUS or NUTRITIVE ARTERIES, (F.) *Ar-tères nutritives*. Arterial branches which enter the *foramina nutricia* of long bones, and pene-trate to the medullary membrane.

NUTRITIVE, *Nutritious*, (F.) *Nutritif*. Also, relating to nutrition: hence the 'nutritive func-tions,' or those that are concerned in nutrition.

NUTRITUS, *Aliment*, *Nutrition*.

NUTRIX, gen. *Nutric'is*, *Mamma*, *Nurse*.

NUTS, WATER, *Nelumbium luteum*.

NUX (gen. *Nucis*, 'a nut') AROMATICA, see *Myristica moscata* — n. *Avellana*, *Corylus avellana* (nut) — n. *Barbadensis*, *Jatropha curcas* — n. *Becuba*, *Ibicuba* — n. *Cathartica Americana*, *Jatropha curcas* — n. *Gallæ*, see *Quercus cerris* — n. *Juglans*, *Juglans regia* — n. *Medica*, *Coco* of the *Maldives* — n. *Metella*, *Strychnos nux vomica* — n. *Nethel*, *Datura stramonium* — n. *Moscata*, see *Myristica moscata* — n. *Myristica*, see *Myristica moscata* — n. *Pistacia*, see *Pistacia vera* — n. *Unguenteria*, see *Myristica moscata* — n. *Vomica*, *Strychnos nux vomica*.

NYCTALOPE, see *Nyctalopia*.

NYCTALOPEX, (*νυκταλωπής*), see *Nyctalopia*.

NYCTALOP'IA, (*νυκταλωπία*, from *nycto*, and *οπτομαι*, 'I see,') *Parop'sis*, *Lucif'uga*, *Nyctalopi'-asis*, *Cæcitas diurna*, *Vit'us noctur'nus*, *Oxyo'pia*, *Hemeral'opia* (moderns), *Amblyo'pia meridiana*, *Hemerat'ophlo'sis*, *Hemerot'ophlo'sis*, *Photophob'ia*, *Dyso'pia tu'nnis*, *Vit'us a'crior*, *Nyc'talopy*, *Day-blindness*, *Diur'nal amauro'sis*, (F.) *Vue nocturne*, *Avenglement de Jour*. The faculty of seeing during the night, with privation of the faculty during the day. It affects both eyes at once, when idio-pathic. Its duration is uncertain, and treatment very obscure. It is, however, a disease of nervous irritability, and one of excitement of the visual

nerve in particular. The indications of cure will consequently be—to allay direct irritation in every way; to excite counterirritation by blisters; and to gradually accustom the eye to the impression of light.

One labouring under this affection is called a *Nyctalope*, *Nyctalops*, *Nyctalopex*.

Nyctalope has also been used for night blindness. See Hemeralopia.

NYCTALOPS, gen. *Nyctalop'is*, (νυκταλωψ,) see *Nyctalopia*.

NYCTALOPY, *Nyctalopia*.

NYCTERINUS, (νυκτερινος,) Nocturnal.

NYCTHEMERON, see *Dies*.

NYCTHEMERUM, (*nycto*, and *hēmera*, 'day,')

The space of 24 hours, or of a day and night. Certain complaints continue only so long.

NYCTO, (from νύξ, gen. νυκτος, 'night,') in composition, night.

NYCTOBADIA, (*nycto*, and *βαδω*, 'to walk or mount,') *Somnambulism*.

NYCTOBASIS, (*nycto*, and *basis*, 'walking,') *Somnambulism*.

NYCTOBATESIS, (*nycto*, and *βαδω*, 'to walk or mount,') *Somnambulism*.

NYCTOBATIA, *Somnambulism*.

NYCTOTYPHLOSIS, (*nycto*, and *τυφλος*, 'blind,') *Hemeralopia*.

NYGMA, gen. *Nyg'matis*, (νύγμα, 'a prick,' from νύσσειν, 'to pierce,') Wound, punctured.

NYGMATICUM EMPLASTRUM, ('wound plaster,') *Emplastrum resinæ*.

NYMPHÆ, (νύμφα,) *Clitoris*, see *Nymphæ*.

NYMPHÆ, (νύμφαι,) *Alæ mino'res seu inter'næ mino'res clitor'idis seu mul'ebres mino'res*, *Carunculæ cuticulæ'res*, *Cris'tæ clitor'idis*, *Collie'uli vaginæ*, *Myrtoschi'la*, *Myrtosche'i'lideæ*, *Lab'ia mino'ra seu inter'na seu puden'di mino'ra*, (F.) *Nymphæ*, *Petites lèvres*. Two membranous folds, which arise from the lateral parts of the prepuce of the clitoris, and descend on the inner surface of the labia majora; terminating, by becoming gradually thinner, about the middle of the circumference of the orifice of the vagina. They are formed each of two folds of the mucous membrane of the vulva; and contain, in their substance, a thin layer of spongy, erectile tissue. Their use seems to be,—not, as was once supposed, to direct the course of the urine, which notion gave rise to their name, (after the fabulous *Water nymphæ*,) but to favour the elongation and dilatation of the vagina in pregnancy and labour.

The word *νύμφη*, *Nymphê*, has also been used synonymously with clitoris by Oribasius, Aëtius, &c.

NYMPHÆA ADVENA, (νυμφαία,) *Nuphar advena*.

NYMPHÆA AL'BA, *Leuconymphæ'a*, *Nen'uphar*, *Microleuconymphæ'a*, *Casta'lia specios'a*, *White Waterlily*, (Sc.) *Cam'biateaf*, (F.) *Nénuphar* ou *Némufar blanc*, *Lis des étangs*, *Lune* ou *Volant d'Eau*, *Nat. Ord. Ranunculaceæ*. *Sec. Syst. Polyandria Monogynia*. Formerly employed as a demulcent, antaphrodisiac, emollient, and slightly anodyne remedy.

NYMPHÆA INDICA, *N. nelumbo*—*n. Major lutea*, *N. lutea*.

NYMPHÆA LU'TEA, *N. ma'jor lu'tea seu umbili-*

ca'lis, *Nu'phar lu'teum*, *Nen'uphar lu'tea*, *Yellow Waterlily*, (Sc.) *Cam'biateaf*, (F.) *Nénuphar jaune*. Used for the same purposes as *Nymphæa alba*. The flowers are called in Norfolk, England, *Brandybottles*.

NYMPHÆA NELUM'BO, *N. In'dica*, *Fa'ba Ægypti'aca*, *Cy'anus Ægypti'acus*, *Pon'tic* or *Ægypti'an Bean*. The fruit of this is eaten raw in Egypt and some of the neighbouring countries; and is considered to be tonic and astringent.

NYMPHÆA ODORA'TA, *Sweet-scented Waterlily*, *Sweet water* or *White Pond* or *Toad Lily*, *Cow* or *Water Cab'bage*, (F.) *Nénuphar odorant*. An indigenous plant, growing in most parts of the United States in fresh water ponds, and on the borders of streams, and having large white, beautiful, sweet-scented flowers. The root is very astringent and bitter. It is sometimes made into a poultice and used as a discutient.

NYMPHÆA UMBILICALIS, *N. lutea*.

NYMPHE, (νύμφη,) *Clitoris*, see *Nymphæ*.

NYMPHES, *Nymphæ*.

NYMPHITIS, (*nymphæ*, and *itis*,) *Epiderrh'i'tis*, *Inflammat'io Clitor'idis*. Inflammation of the clitoris.

NYMPHOMA'NIA, (*nymphæ*, and *μανία*, 'fury,') *Fu'ror uteri'nis*, *Uteroma'nia*, *Lagne'sis fu'ror femini'nis*, *Metroma'nia*, *Ædæogar'galus*, *Ædæogargaris'mus*, *Thelygon'ia*, *Erotoma'nia*, *Ædema'nia*, *Aidoioma'nia*, *Cytheroma'nia*, *Melanchol'ia uteri'na*, *Sympto'ma turpitu'dinis*, *Androma'nia*, *Gynæcoma'nia*, *Machlos'ynê*, *Tenti'go vene'rea*, *Hysteroma'nia*, *Sulac'tias vulvæ*, *Uteri prur'itus*, *Brachu'na*, *Ara'tum*, *Æstroma'nia*, (F.) *Nymphomanie*, *Fureur utérine*. An irresistible and insatiable desire, in females, for the venereal act. It occurs in those particularly, who possess a nervous temperament, and vivid imagination, especially if excited by improper language, masturbation, &c. Its course, as described, is as follows. In the commencement, the sufferer is a prey to perpetual contests between feelings of modesty and impetuous desires. At an after period, she abandons herself to the latter, seeking no longer to restrain them. In the last stage the obscenity is disgusting; and the mental alienation, for such it is, becomes complete. The treatment consists in the use of the same means as are required in the satyriasis of man. When the mental alienation is complete, solitude is indispensable.

NYMPHON'CUS, (*nymphæ*, and *ογκος*, 'a tumour,') A morbid tumefaction of the nymphæ.

NYMPHOTOMY, *Nymphotom'ia*, *Nympharum Sect'io*, (F.) *Nymphotomie*, (*nymphæ*, and *τομή*, 'to cut,') An operation, known and practised for a long time, which consists in the excision of the nymphæ. The operation is had recourse to, when they are attacked with scirrhus, cancer, fungus, or gangrene; or when they are so large as to interfere with walking or coition. *Nymphotomy* is the circumcision of the female. It is practised in some countries.

Some authors have used the term *Nymphotomy* for amputation of the clitoris.

NYSTAG'MUS. A partial rotatory movement of the eyeball from side to side. (F.) *Sou'ris*. Also, *Coma vigil*, (*νυσταγμος*, 'sleep,')

NYXIS, (νύσσειν, 'to pierce,') *Puncture*.

O.

OAK, BLACK. (Sax. *ac*, *æc*.) *Quercus tinctoria* —o. Common, *Quercus robur* —o. Jerusalem, *Che-nopodium botrys*.

OAK LUNGS, *Lichen pulmonarius*.

OAK ORCHARD AC'ID SPRINGS. These springs — eight in number — are in Genesee county, New York, eight miles from Lockport. They are acid salines, consisting of free sulphuric acid, sulphate of protoxide of iron, and sulphates of alumina, magnesia, and lime. They are powerfully astringent and tonic.

OAK, POISON, *Rhus toxicodendron* — o. Red, *Quercus rubra montana* — o. Sea, *Fucus vesiculosus* — o. Spanish, *Quercus rubra montana* — o. White, *Quercus alba*.

OAKUM, see *Linteum*.

OÄR'IC, *Oär'icus*, (F.) *Oarique*, (from *oarion*.)

Relating or appertaining to the ovary.

OÄRIOCYESIS, (*oarion*, and *κρησις*, 'pregnancy,') Pregnancy, ovarian.

OÄRION, (*ωαριον*, from *ωον*, *ovum*, 'an egg,') Ovary.

OÄRION'CUS, *Oariophy'ma*, *Ova'rium tu'mi-dum*, *Tu'mor Ova'rrii*, (*oarion*, and *ογκος*, 'swelling,') Ovarian tumour.

OÄRIOPAREC'TAMA, gen. *Oarioparectem'atis*, *Oophoraua'is*, (*oarion*, and *παρεκτειναι*, 'to extend,') Enlargement of the ovary.

OÄRIOP'ATHY, *Oariopathi'a*, *Oophor'opathi'a*, (*oarion*, and *παθος*, 'affection,') Disease of the ovary.

OÄRIOPHYMA, (*oarion*, and *φυμα*, 'a swelling,') Oarioncus.

OÄRIORRHEX'IS, (*oarion*, and *ρηξις*, 'rupture,') *Ruptu'ra Ova'rrii*. Rupture of the ovary.

OÄRIOSCI'R'HUS, (*oarion*, and *scirrhus*.) Scirrhus of the ovary.

OÄRIOSTERESIS, (*oarion*, and *στερησις*, 'privation,') Oariotomy.

OÄRIOT'OMY, *Oariotom'ia*, *Ovariotom'ia*, *Oäriostere'sis*, *Ovariot'omy*, (*oarion*, and *τομή*, 'incision,') The operation for removing the ovary.

OÄR'ITIS, *Oöphori'tis*, *Oori'tis*, *Inflamma'tio Ova'rrii*, *Ovari'tis*, (F.) *Inflammation de l'Ovaire*, (*oarion*, and *itis*.) Inflammation of the ovarium. When occurring in childhood, it is the *Lochoöphoritis* of some.

OÄRIULE, (*oarion*, and *ἐλη*, 'cicatrix,') *Corpus luteum*.

OÄRIUM, (*ωαριον*.) Ovary.

OÄRIYDROPS, (*oarion*, and *ἰδρωψ*, 'dropsy,') Hydroarion.

OÄRTHROC'ACË, (formed by contraction from *ομοärthrocæce*.) Said to be used by Rust to designate inflammation of the scapulohumeral articular surfaces.

OATMEAL, see *Avena*.

OATS, (Sax. *æten*.) *Avena* — o. Water, *Zizania aquatica*.

OB (L.), in composition, 'on, before, over, against, towards.' At times, it is only intensive; and, at others, it means inversely. Before *c* and *f*, it is often changed into *oc*, *of*; and before *p*, always into *op*.

OBAUDITIO, (*ob*, and *audire*, *auditus*, 'to hear,') Baryecoa.

OBAUDITUS, Baryecoa.

OBÆCATIO, (*ob*, and *cæcare*, *cæcatum*, 'to blind,') Cæcitas.

OBDO'RMIT'IO, (*ob*, and *dormio*, *dormitum*, 'I sleep,') The state of being asleep; also, of the limbs being asleep; — *Stu'por ar'tuum*.

OBDUCTIO LEGALIS, Autopsia cadaverica legalis.

OBELÆA RAPHE, (*οβελος*, 'a spit or arrow,') Sagittal suture.

OBESITAS, ('fatness,') Polysarcia — o. Colli, *Struma adiposa* — o. Nimia, *Pimelosis* — o. Viscerum, *Physconia adiposa*.

OBESITË, Polysarcia.

OBSITY, Polysarcia.

OBESUS, ('fat,') Corpulent.

OBFUSCATIO, (*ob*, and *fuscare*, *fuscatum*, 'to make dark,') Amaurosis.

OBITUS, (*ob*, and *ire*, 'to go,') Death.

OBJECTIVE CONE. The cone of light proceeding from an object, the apex of which is on the object, and the base on the cornea.

OBJECTIVE SENSATIONS, see Sensation.

OBLINITIO, (*ob*, and *linere*, 'to besmear,') Inunction.

OBLIQUE', *Obli'quus*. Any thing inclined, or which deviates from the vertical line. Anatomists have given this name to certain muscles, which have an oblique direction as regards the plane that divides the body into two equal and symmetrical halves. These are:—

OBLIQUE MUSCLES OF THE ABDOMEN. They are two in number, and distinguished into: 1. *Obliquus Exter'nus Abdom'inis seu descen'dens exter'nus seu descendens seu ma'jor*, (F.) *Ilio-pubo costo-abdom'inal*, *Costo-abdom'inal* (Ch.), *Grand oblique*, *Oblique externe*. One of the broadest muscles of the body. It is situate at the lateral and anterior part of the abdomen; and is flat and quadrilateral. It is attached, *above*, to the outer surface and lower edge of the 7 or 8 last ribs; *below*, to the anterior third of the external lip of the crista illi; *before*, it terminates at the linea alba by means of a broad and strong aponeurosis, which covers the rectus, and presents towards its inferior part two very solid fibrous fasciuli, which are inserted, — the one at the symphysis, the other at the spine of the pubis, — under the name of *Pillars of the Abdominal Ring*. These pillars leave between them an opening, which forms the inferior orifice of the inguinal canal. The obliquus externus abdominis depresses the ribs, and carries them backwards during a strong expiration. It impresses on the chest a movement of rotation, and bends the thorax upon the pelvis, and conversely. It contracts, also, the abdominal cavity. 2. *Obliquus Inter'nus Abdom'inis seu ascen'dens seu mi'nor seu inter'nus seu ascendens internus*, *Musculus accl'i'vis*, (F.) *Ilio-lombo-costo-abdom'inal*, *Ilio-abdom'inal* (Ch.), *Muscle petit oblique ou oblique interne*, is broad, especially before; thin, and irregularly quadrilateral, like the preceding, beneath which it is situate. It is attached, *above*, to the inferior edge of the cartilages of the 5th, 4th, 3d, and 2d false ribs; *below*, to the anterior two-thirds of the interstice of the crista illi, to the posterior part of the crural arch, and to the pubis; *behind*, to the spinous processes of the last two lumbar vertebrae, and to those of the first two portions of the sacrum; *before*, to the linea alba. Its upper fibres run obliquely upwards and forwards; the middle are horizontal; and the lower pass obliquely downwards and forwards.

These last, in the male, are dragged down through the inguinal ring, when the testicle descends, and form the two fasciuli of the cremaster.

The obliquus internus resembles the O. externus in function.

OBLIQUE MUSCLES OF THE EYE, *Amato'rrii*, *Circumagen'tes*, *Rotato'res Oe'uli*, are two in number. They are distinguished into: 1. *Obliquus Super'ior Oe'uli*, *Amato'rius musculus*, *Trochlea'ris*,

Trochlear, *Obliquus major*, *Circumductio* *ophtice*, *Longissimus oculi*, (F.) *Optico-trachlei sclerotici*, *Grand trochléateur* (Ch.), *Grand oblique de l'œil*, *Oblique supérieur de l'œil*, *Amoureux* (Muscle), is situate at the inner and upper part of the orbit. It is small, round, fusiform, and reflected upon itself in the middle of its course. Behind, it is attached to the inside of the foramen opticum; and when it arrives opposite the internal orbital process, it becomes a small, round tendon, which slides in a cartilaginous pulley fixed to the os frontis, and is reflected, at an acute angle, to proceed downwards and outwards, and to attach itself to the outer and back part of the globe of the eye: This muscle carries the globe of the eye forwards and inwards; making it experience a movement of rotation, which directs the pupil downwards and inwards. This is conceived to be an involuntary muscle as well as the next. In sleep, according to Sir C. Bell, when the power over the straight or voluntary muscles of the organ is nearly lost, the eye is given up to the oblique muscles, which lodge the transparent cornea under the upper eyelid. At the approach of death, the same thing is observable; hence, the turning up of the eye, at such time, is not an evidence of agony or suffering, but of insensibility. 2. *Obliquus Inferior* seu *mi'nor oculi*, (F.) *Maxillo-sclerotici*, *Petit Trochléateur* (Ch.), *Petit oblique ou oblique inférieur de l'œil*, is situate at the anterior and inferior part of the orbit. It is flat and attached to the inner and anterior part of the orbital surface of the superior maxillary bone, on the outside of the lachrymal gutter; from thence it passes outwards and backwards, and terminates by an aponeurosis, at the posterior and inner part of the globe of the eye. It carries the globe of the eye inwards and forwards; and directs the pupil upwards and outwards.

OBLIQUE MUSCLES OF THE HEAD. These are two in number. 1. *Obliquus Superior* seu *mi'nor Cap'itis*, (F.) *Trachélo-atloïdo-occipital*, *Atloïdo-sous-mastôïdien* (Ch.), *Muscle oblique supérieur ou petit oblique de la tête*. This muscle is situate at the sides of, and behind, the articulation of the head: it is flat and attached, on the one hand, to the top of the transverse process of the atlas; and, on the other, terminates at the occipital bone, beneath the inferior curved line, and sometimes, also, at the mastoid region of the temporal bone. It extends the head,—inclining it to one side. 2. *Obliquus Inferior Cap'itis*, *Obliquus major*, (F.) *Spino-axoïdo-trachéi-atloïdien*, *Axoïdo-atloïdien* (Ch.), *Oblique inférieur ou grand oblique de la tête*, is situate at the posterior part of the neck and head. It is round, fusiform; attached to the spinous process of the axis, and proceeds to terminate behind and below the summit of the transverse process of the atlas. It impresses, on the first vertebra and the head, a movement of rotation, which turns the face to one side.

OBLIQUE PROCESSES, see *Vertebrae*.

OBLIQUE EXTERNE, *Obliquus externus abdominis*—o. *Grand*, *Obliquus externus abdominis*—o. *Grand de l'œil*, *Obliquus superior oculi*—o. *Grand de la tête*, *Obliquus inferior capitis*—o. *Inferieur de l'œil*, *Obliquus inferior colli*—o. *Inferieur de la tête*, *Obliquus inferior capitis*—o. *Interne*, *Obliquus internus abdominis*—o. *Petit*, *Obliquus internus abdominis*—o. *Petit de l'œil*, *Obliquus inferior oculi*—o. *Petit de la tête*, *Obliquus superior oculi*—o. *Supérieur de l'œil*, *Obliquus superior oculi*—o. *Supérieur de la tête*, *Obliquus superior capitis*.

OBLIQUITÉ DE LA MATRICE, *Retroversio uteri*.

OBLIQUUS, (*ob*, and *liquis*, 'slanting,') *Oblique*—o. *Ascendens*, *O. Internus abdominis*—o. *Auris*, *Laxator tympani*—o. *Descendens externus*, *O. externus abdominis*—o. *Externus abdominis*, see *Ob-*

lique muscles of the abdomen—o. *Internus abdominis*, see *Oblique* muscles of the abdomen—o. *Major*, *O. externus abdominis*, *O. inferior capitis*, *O. superior oculi*—o. *Minor*, *O. internus abdominis*—o. *Minor capitis*, *O. superior capitis*—o. *Minor oculi*, *O. inferior oculi*.

OBLITERATED, *Obliteratus*: (*obliterare*, (*ob*, and *litera*,) 'to efface,') A vessel or duct is said to be obliterated, when its parietes have approximated and contracted such an adhesion to each other that the cavity has completely disappeared.

OBLITERATIO COMPLETA, *Hapantismus*.

OBLIVIO, ('forgetfulness,') *Amnesia*—o. *Iners*, *Lethargy*.

OBLÖBIUM, (*ob*, and *λοβος*, 'a lobe,') *Antilobium*.

OBNUBILATION, (F.) *Obnubilatio*, (*ob*, and *nubilare*, *nubilatum*, 'to be cloudy,') *Giddiness*, *dazzling*.

OB'OLUS, *Onolosat*. A weight of 9 or 10 grains.

OBSCENÆ PARTES, ('obscene parts,') *Genital organs*.

OBSCURISSEMENT DE LA VUE, (from *obscurus*, 'obscure,') *Caligo*.

OBSERVATION, *Observatio*, (*ob*, and *servare*, *servatum*, 'to keep,' e.g. in sight.) *Tere'sis*, *Symparater'sis*. Act of examining a thing by means of the external senses. This word is employed in several acceptations. It expresses—1. The action of observing—2. The aggregate of knowledge, afforded by observation. In French—but not in English—it means the case or history of the phenomena presented by a patient in the course of a disease.

OBSERVATION, Case—o. Method of, *Numerical method*.

OBSTETRIC, *Obstet'ric*, *Obstet'ricus*, *Maien'tical*, *Maien'ticus*, (*obstare*, (*ob*, and *stare*,) 'to stand before,') Relating or appertaining to *Obstetrics*,—as 'obstetric auscultation,' 'obstetric exploration,' &c.

OBSTETRIC CHAIR, *Labour chair*.

OBSTETRICANS, *Accoucheur*.

OBSTÉTRICIE, *Obstetrics*.

OBSTETRICS. *Tokol'ogy*, *Tocol'ogy*, *Maie'ia*, *Maei'a*, *Maei'a*, *Maeen'tica ars*, *Ars obstetric'ia*, *Obstet'ricy*, (F.) *Obstétrique*, *Obstétricie*. The art of midwifery. Midwifery in general.

OBSTETRICY, *Obstetrics*.

OBSTÉTRIQUE, *Obstetrics*.

OBSTETRIST, *Accoucheur*.

OBSTETRIX, *Midwife*.

OBSTIPATIO, (*ob*, and *stipare*, *stipatum*, 'to cram close,') *Constipation*—o. *Alvina*, *Constipation*—o. *Tenesmus*, *Tenesmus*.

OBSTIPATIO, see *Hump*.

OBSTIPITAS CAPITIS seu **COLLI,** (*obstipo*, 'to lean to one side,') *Torticollis*.

OBSTIPITÉ, *Torticollis*.

OBSTRUCTION, (*obstruo*, (*ob*, and *struere*, *structum* 'to build,') 'I stop up by building against,') *Emphraxis*, *Stegnosis*—o. *Alvi*, *Constipation*—o. *Ductus alimentarii*, *Constipation*—o. *Ductus Stenoniani*, *Stenostenosis*—o. *Intestinalis*, *Constipation*—o. *Pulmonum pituitosa febrilis*, *Peripneumonia notha*—o. *Recti Spastica*, *Stricture of the Rectum*, *spasmodic*.

OBSTRUCTION OF THE INTESTINES, *Enteroemphraxis*.

OB'STRUENS, *Oppilati'vus*. A medicine which closes the orifices of ducts or vessels.

OBSTUPEFACIENTIA, (*ob*, and *stupefacere*, 'to benumb,') *Narcotics*.

OBTONDANT, *Obtundens*.

OBTURCATIO, (*ob*, and *truncare*, *truncatum*, 'to cut off,') *Decollation*.

OBTUNDENS, (F.) *Obtondant*, (*obtundere*, (*ob*, and *tundere*, 'to beat,') 'to beat against,' and

therefore to blunt the edge.) An epithet applied to remedies that were supposed, according to an erroneous theory, to be possessed of the power of blunting the acrimony of the humours. A demulcent is one of these.

OBTURAMENTUM, (*obturare, obturatum, 'to stop up,'*) Emphragma.

OBTURATEUR, Obturator—*o. du Palais*, Palate, artificial.

OBTURATIO, Emphraxis.

OBTURATION DES DENTS, Plugging.

OBTURATOR, *Obturato'rius, (F.) Obturateur*. An instrument adapted for closing an opening caused by a wound or disease, as '*Obturator of the palate*,' see Palate, artificial. Also a name given in anatomy to several parts connected with the Obturator foramen.

OBTURATOR ARTERY, *Arte'ria obturato'ria seu obtura'trix, (F.) Sous-pubio fémorelle (Ch.), Artère obturatrice*, arises, most commonly, from the hypogastric. It is, however, frequently given off from the epigastric; a matter of importance to be determined in cases of femoral hernia. Of 500 obturator arteries examined by M. J. Cloquet, 340 were furnished by the hypogastric, and 152 by the epigastric or crural. When it arises from the hypogastric, it passes forwards and outwards, and then turns horizontally into the cavity of the pelvis, to issue from this cavity by the opening left at the upper part of the obturator membrane. When, on the contrary, the obturator artery arises from the epigastric or the crural, it descends obliquely inwards, behind the os pubis, to the obturator foramen. At its exit from the pelvis, the artery divides into two branches, a *posterior* and an *anterior*, which are distributed to the muscles of the anterior and superior part of the thigh.

OBTURATOR FORAMEN, *Fora'men Obturato'rium seu obtura'tum seu infrapubia'num seu ova'lè seu thyroïdeum seu Thyroï'des seu An'plum Pelvis, (F.) Trou sous-pubien*. A large opening, of an oval or triangular form, in the anterior part of the os innominatum, on the outside of the symphysis pubis and beneath the horizontal ramus of the os pubis. This foramen is closed by a membranous ligament.

OBTURATOR LIGAMENT OR MEMBRANE, *Sub-pub'ic mem'brane*, is a fibrous membrane, fixed to the whole circumference of the obturator foramen, except above, where an opening remains for the passage of the vessels and nerves of the same name.

OBTURATOR MUSCLES, *Obturato'res, Rotato'res fem'oria*. These are two in number. They are divided into

a. *Obturator Exter'nus, Extra-pelvio-pubi-trochan'térien, Sous-pubio-trochan'térien externe (Ch.)*. A muscle, situate at the anterior and inner part of the thigh. It is broad, flat, and triangular; and is attached, on the one hand, to the anterior surface of the os pubis, to that of the ischium, and to the anterior surface of the obturator ligament. Its fleshy fibres converge to the tendon, which proceeds to be inserted at the inferior part of the cavity of the great trochanter. This muscle rotates the thigh outwards.

b. *Obturator Inter'nus, Marsupia'lis, Bursa'lis, Intra-pelvio-trochan'térien, Sous-pubio-trochan'térien interne (Ch.)*, is seated, almost entirely, in the pelvis. It arises from the inner surface of the obturator ligament, and from the posterior part of the circumference of the obturator foramen, and is inserted, by means of a strong tendon, running between the two portions of the gemini, into the cavity at the root of the great trochanter; after having turned upon the ischium, which forms for it a kind of pulley. This muscle also rotates the thigh outwards.

OBTURATOR NERVE, *Sous-pubio-fémorale, (Ch.)*, proceeds principally from the 2d and 3d lumbar

nerves. It descends into the pelvis; gains the obturator foramen; gives branches to the obturator muscles, and divides, behind the abductor primus and pectinalis, into two branches; one *anterior*, whose branches are distributed to the first two adductors, gracilis, and integuments; the other, *posterior*, distributing its ramifications to the obturator externus and third adductor.

OBTURATOR VEIN has, ordinarily, the same arrangement as the artery. It is common, however, to find it arising from the epigastric; whilst the corresponding artery proceeds from the hypogastric, and conversely.

OBTURATORIES, Obturator muscles.

OBVOLVENTIA, (*obvolvere, 'to cover over,'*) Demulcents.

OC, as a prefix, see Ob.

OCA, Oxalis tuberosa.

OCCECATIO, (*oc, and cæcus, 'blind,'*) Cæcitas.

OCCIP'ITAL, *Occipita'lis, (from Occiput, gen. Occip'itis.)* That which belongs to the occiput.

OCCIPITAL ARTERY. This arises from the posterior part of the external carotid, beneath the parotid. It proceeds backwards, passes between the mastoid process and the transverse process of the atlas; reaches the occipital bone and divides into two branches, which may be called ascending posterior and anterior, and are distributed to the neighbouring muscles and ligaments.

OCCIP'ITAL BONE, *Os occip'itis seu occip'itii seu occipita'le seu sph'e'nobasila'rè seu memo'ria' seu nervo'sum seu basila'rè seu pro'ria' seu pyr'idis seu sea'tum cra'nii seu lambdoï'des seu lamb'dæ seu pup'pis seu nervo'sum seu fibro'sum seu pelvi-cephal'icum, (F.) Os occipitalis*, is situate at the posterior and inferior part of the cranium, which it assists in forming. It is flat, symmetrical, and curved upon itself. It presents, 1. An *occip'ital or posterior surface*, which is convex, and has, upon the median line, the *bas'iliary surface*, the *fora'men mag'num*, through which passes the spinal marrow with its membranes and vessels,—the *external occipital crest*, the *external occipital protuberance*; and, at the sides, the *upper curved line*, *large rough arched ridge or transverse arch* or *lin'ea semicircula'ris*, the *lower curved line*, the *posterior condyloid fossæ*, the condyles for the articulation of this bone with the atlas; and the *anterior condyloid fossæ*, pierced by a foramen for the passage of the ninth pair of nerves. 2. A *cer'ebral or anterior surface*. On the median line are: the *bas'iliary fossæ*, the *inner orifice* of the foramen magnum, the *internal occip'ital crest*, the *internal occip'ital protuberance*, the *cru'ciform spine*; a *channel*, which lodges the termination of the straight sinus, and, on each side, the *occip'ital fossæ*, distinguished into *superior or cer'ebral*, and *inferior or cerebel'lous*, and separated by a groove which lodges the lateral sinus. 3. The surfaces of the occipital bone are separated by four ridges and four angles. The two superior edges are articulated with the parietal bones; the two lower join the temporal; and the anterior angle, under the name *bas'iliary process*, is united to the sphenoid.

The occipital bone is developed from four points of ossification; and sometimes from a greater number.

OCCIPITAL MUSCLE, *Occipita'lis*. Many anatomists have given this name to the posterior fasciculus of the occipitofrontalis.

OCCIPITAL NERVE, *Suboccip'ital nerve, (F.) Première paire trachélienne (Ch.), Nerf occipital ou sous-occipital*. It arises from the upper part of the spinal marrow by eight or ten filaments, united into two fasciculi. Thus formed, it passes between the foramen magnum and the posterior arch of the atlas; and, at this place, forms a long ganglion, afterwards dividing into two branches. Of

these, the *anterior*, which is long and small, makes a turn above the transverse process of the atlas and forms an anastomotic noose with a branch of the second cervical nerve. The *posterior* branch, larger and shorter, divides into seven or eight branches, which are distributed to the muscles of the upper and back part of the neck.

OCCIPITAL REGION, Occiput.

OCCIPITAL VEIN. Its roots follow exactly the course of the branches of the artery, and unite into a single trunk, which opens into the internal jugular vein, and sometimes into the external.

OCCIPITUM, Occiput.

OCCIPITOATLOID, *Occipitoatloïdeus*, (F.) *Occipitoatloïdien*. That which has reference to the occiput and atlas.

OCCIPITOATLOID ARTICULATION is the articulation of the condyles of the occipital bone with the superior articular cavities of the atlas. It is strengthened by two ligaments—the one *anterior*, the other *posterior*, called *occipitoatloïd ligaments*: the one extends from the anterior, the other from the posterior arch of the atlas to the corresponding portion of the circumference of the foramen magnum.

OCCIPITOAXOID, *Occipitoaxoïdeus*, (F.) *Occipito-axoïdien*. That which relates to the occipital bone and the axis or second vertebra.

OCCIPITOAXOID ARTICULATION is the connexion of the occipital bone with the axis or second vertebra, although these bones are not really articulated, but are merely retained in apposition by three strong ligaments, the posterior of which is called the *occipito-axoid*, and the two others *odontoid*.

OCCIPITOFRONTALIS, *Digastricus cranii*, *Epicranialis*, *Frontalis et occipitalis*. The majority of anatomists call by this name the whole of the fleshy plane, with the epicranial or coronal aponeurosis (see *Calotte*), which covers the head from the occiput to the forehead. It is attached, by its posterior fasciculus, to the two outer thirds of the upper curved line of the occipital bone, and to the outer surface of the mastoid portion of the temporal; and, by its anterior fasciculus, it terminates at the eyebrow, where it becomes confounded with the superciliaris, pyramidalis nasi, and orbicularis palpebrarum.

The occipitofrontalis, by the contraction of its anterior fasciculus, draws forward a part of the integuments of the cranium. It wrinkles the skin of the forehead transversely, and may also contribute to open the eye by its decussation with the orbicularis palpebrarum. The posterior fasciculus of the muscle draws backwards a part of the skin of the cranium, and assists in stretching the common aponeurosis.

OCCIPITOMENINGEAL. That which belongs to the occipital bone and to the meningeal or dura mater.

OCCIPITOMENINGEAL ARTERY, in Chaussier's nomenclature, is a branch of the vertebral, given off to the dura mater at its entrance into the cranium.

OCCIPUT, gen. *Occip'itis*, *Occipit'ium*, (*oc*, and *caput*, 'the head.') *Regio occipitalis*, *In'ion*, *Opisthocranium*, *Opisthocephalon*, *Hysteroecranium*, *Pro'ra*. The back part of the head, formed by the occipital bone.

OCCIPUT, SORT, Craniotabes.

OCCCLUSION, *Occlusio*, (*occludere*, *occlusum*, 'to shut up.') Sometimes this word signifies, simply, the transient approximation of the edges of a natural opening—the *occlusion of the eyelids*, for example; at others, it is synonymous with imperforation, as *occlusion of the pupil*, *vagina*, &c.

OCCULT DISEASES, see Latent.

OCEAN SPRINGS, see Mississippi, mineral waters of.

OCHEMA, gen. *Oche'matis*, (*οχημα*.) Vehicle.

OCHETEUMATA, (pl. of *οχητευμα*, 'a canal.') Nares.

OCHELOS, (*οχερος*.) Canal.

OCHEUMA, (*οχηυμα*.) Coition, Fecundation.

OCHEUS, (*οχευς*, from *οχεω*, 'to hold.') Scrotum.

OCHLE'SIS, (*οχλος*, 'a crowd.') *Crowded condition*. A morbid condition induced by the crowding together of sick persons under one roof.

OCHREA (from *οχρος*, 'pale') RUBRA, ('red ochre.') Hematites.

OCHRIASIS, (*οχριασις*.) Paleness.

OCHROMA, (*οχρος*, 'pale.') Paleness.

OCHROPYRA, (*οχρος*, 'yellow,' and *πυρ*, 'fever.') Fever, yellow.

OCHROTES, (*οχροτης*.) Paleness.

OCHROTYPHUS, (*οχρος*, 'yellow,' and *typhus*.) Fever, yellow.

OCHTHIASIS, (*οχθος*, 'a hill, a protuberance.') Molluscum.

OCHTHODES, (*οχθος*, 'the hard edge of an ulcer.') Callous.

OCIMUM ADSCENDENS, (*ωκιμον*, from *ωκυς*, 'swift,' owing to its swift growth.) O. Basilicum.

OCIMUM BASILICUM, *O. adscendens* seu *pinissum* seu *racemosum* seu *citratum*, *Beren'daros*, *Basilicum*, *B. majus* seu *citratum*, *Com'mon* or *Cit'ron basil'*, (F.) *Basilic commun.* Nat. Ord. Labiate. *See Syst. Didynamia* Gymnospermia. This herb has a fragrant odour and aromatic taste. It is used as a condiment, and has been supposed to possess nervine properties.

OCIMUM CARYOPHYLLATUM, *O. min'imum*, *Small* or *Bush basil*, possesses properties similar to the former. It is sometimes used as snuff.

OCIMUM CITRATUM, O. Basilicum—o. Pilosum, O. Basilicum—o. Racemosum, O. Basilicum.

OCOTEA AMARA, *Nectandra cymbarum*—o. Cymbarum, *Nectandra cymbarum*—o. Pichurim, see Pichurim beans—o. Puchury, see Pichurim beans.

OCREA, *Hippocampus minor*, Shin.

OCTA'NA, *Hebdomada'ria*, *Fe'bris hebdomada'ria*, (*octo*, 'eight.') A fever whose paroxysms recur every week;—a supposititious case.

OCTA'RIOUS, (from *octo*, 'eight.') A pint. The eighth part of a wine-gallon. Same etymon. It contains sixteen fluidounces (Ph. U. S.)—20 fluidounces imperial measure. See Weights.

OCTOPUS SYNAPHOCEPHALUS, Hemipages.

OCTOROON, (from *octo*, 'eight,') see Mulatto.

OCTUNX, (*octo*, 'eight,' and *uncia*, 'an ounce.') A weight of eight ounces.

OCULAR, *Ocularis*, (F.) *Oculaire*, (from *oculus*.) Of or belonging to the eye.

OCULAR CONE. The cone formed within the eye by a pencil of rays proceeding from an object; the base of the cone being on the cornea, the apex on the retina.

OCULARES COMMUNES, *Motores oculorum*—o. Dentes, Canine teeth.

OCULARIA, (from *oculus*.) *Euphrasia officinalis*.

OCULI MARMARYGODES, *Metamorphopsia*—o. Palpebrarum scabies pruriginosa, see Ophthalmia tarsi.

OCULIST, *Oculis'ta*, *Ophthalmia'ter*, *Med'icus ocularius* seu *ophthal'micus*, (from *oculus*.) One who occupies himself chiefly with the management of diseases of the eye.

OCULISTIQUE, Ophthalmology.

OCULOMOTORIUS, Motor oculi.

OCULOMUSCULAR NERVES, COMMON, *Motores oculorum*.

OCULUM MOVENS PRIMUS, *Rectus internus oculi*—o. *Movens quartus*, *Rectus inferior oculi*—o. *Movens secundus*, *Rectus externus oculi*—o. *Movens tertius*, *Rectus superior oculi*.

OCULUS, (*οκος*.) Eye—o. Bovinus, *Hydrophthalmia*—o. Bovis, *Chrysanthemum leucanth-*

rum—o. Bubulus, Hydrophthalmia—o. Cæsius, Glaucoma—o. Duplex, Binoculus—o. Elephantinus, Hydrophthalmia—o. Genu, Patella—o. Lærymans, Epiphora—o. Leporinus, Lagophthalmia—o. Purulentus, Hypopyon—o. Simplex, see Monoculus—o. Typhonis, Scilla.

OCY, (ὤκως, 'quick.') In composition, quick.

OCYUM, see Ocimum—o. Sylvestre, Clino-podium vulgare.

OCYODINIC, (ocy, and ὤδν, 'a labor pain.') Ocytoic.

OCYPH'ONUS, (ocy, and φόνος, 'murder.') An agent that kills speedily.

OCYTOCEUS, (ὠκυτοκεύς,) Ocytoic.

OCYTOCIC, *Ocyto'cic*, *Ocytoce'us*, *Ocyto'cius*, *Ocyodinic*, *Odyagopus*, (ocy, and τὸκος, 'labor.') Any thing that expedites parturition.

OCYTOCIUS, (ὠκυτοκεύς,) Ocytoic.

OCYT'OCUS, (ὠκυτοκεύς,) A quick or easy labor.

OD. A term employed by Reichenbach to designate an "effective force," which, according to him, not only manifests itself in contact with man, as in the case of magnets, crystals, the hands, &c., but also at distances, as from the sun, moon, and stars; indeed, from all matter. Such a force is termed by him *od'ic*.

ODAXIS'MUS, *Odaxes'mus*, (ὀδᾶξιμος, from ὀδᾶω, 'I bite.') *Odontocene'sis*, *Odontocnes'mus*. A sensation of biting or pricking, as that in the gums which precedes the appearance of the teeth.

ODES, (ὠδός.) A suffix, at times having the signification of *oid*, but generally indicating, like *osis* in Latin terms, 'consisting of,' or 'full of.'

ODEUR, Odour.

ODIC, see Od.

ODIN, (ὠδν,) Pain (labour).

ODINAGOGUS, (odin, and αγω, 'I lead.') See Ocytoic.

ODINOL'YSIS, (odin, and λυσις, 'solution.') Mitigation of labour pains.

ODINOPE'IA, (odin, and ποιεω, 'I make.') Agents that excite or encourage labour pains.

ODIS, (ὠδισ,) Pain (labour).

ODME, (ὀδμ,) Odour.

ODONTAGOGON, (ὀδονταγωγόν, *odonto*, and αγω, 'I draw.') Dentagra.

ODONTAGRA, (ὀδονταγρα, from *odonto*, and αγρα, 'a seizure.') A rheumatic or gouty pain in the teeth. Also, Dentagra.

ODONTAL'GIA, (ὀδονταλγία, from *odonto*, and αλγος, 'pain.') *Dent'ium dol'lor*, *Tooth'ache*, *Odont'ia*, *O. dolorosa*, *Gomphi'asis*, *Gomphias'mus*, *Odontodyn'ia*, (F.) *Odontalgie*, *Douleur des dents*, *Fluxion sur les dents*, *Mal de dent* ou *d'amour* ou *de Saint Médard*. A disease dependent upon a variety of causes affecting the cavity of the tooth, but generally owing to caries, which exposes the cavity to the action of the air, and to extraneous matters in general. Hence the treatment consists in plugging the tooth or destroying the sensibility of the nerve by powerful stimulants; and, if these means fail, in extracting the tooth.

ODONTALGIA CARIOSA, Dental gangrene—o. Dentitionis, Dentition—o. Hæmodia, Hæmodia.

ODONTALGIA NERVO'SA, *Neural'gia Dent'alis*. Neuralgia of the teeth. Characterized by periodical pain, shooting with the utmost violence along the branches of the fifth pair distributed to the affected jaw.

ODONTALGIA REMITTENS ET INTERMITTENS; Neuralgia infraorbitalia.

ODONTALGIC, Antiödontalgic.

ODONTALGIE, Odontalgia.

ODONTALGIQUE, Antiödontalgic.

ODONTAMBLYOGMUS, (*odonto*, and αμβλυν, 'blunt.') Hæmodia.

ODONTATROPH'IA, *Atroph'ia Dentis*, (F.) *Atrophie des Dents*, (*odonto*, and *atrophia*, 'atrophy.') Atrophy of the teeth.

ODONTHEMODIA, Hæmodia.

ODONTHARPA, (*odonto*, and ἀρπάζειν, 'to lay hold of.') Dentagra.

ODONTHYALOPH'THORA, (*odonthyalus*, 'enamel,' and φθορά, 'corruption.') Destruction of the enamel of the teeth.

ODONTHYALUS, (*odonto*, and βάλες, 'glass.') Enamel of the teeth.

ODONTHYPERÆSTHESIS, Hæmodia.

ODONTIA, Odontalgia.

ODONT'IA DEFOR'MIS. Deformity of the teeth from error of shape, position, or number.

ODONTIA DENTITIONIS LACTANTII, see Dentition (first)—o. Dentitionis puerilis, see Dentition (second)—o. Edentula, see Nefrendes—o. Excrescens, Epulis—o. Incrustans, Odontolithus—o. Stuporis, Hæmodia.

ODONTIASIS, (ὀδοντίασις,) Dentition.

ODONTIATER, (*odonto*, and ιατρος, 'physician.') Dentist.

ODONTIATRIA, Dentistry.

ODONTIC, Antiödontalgic.

ODONTIT'IS, *Odontophleg'monē*, (*odonto*, and *itis*.) Inflammation of the teeth.

ODONTO, (ὀδόντος, gen. ὀδόντος, 'tooth.') In composition, tooth.

ODONTOBOTHRY'TIS, *Odontophatni'tis*, *Inflamma'tio alveolo'rum*, (*odontobothrium*, and *itis*.) Inflammation of the alveoli.

ODONTOBOTHRIUM, (*odonto*, and βόθριον, 'a small pit.') Alveolus.

ODONTOCERAM'IC, *Odontoceram'icus*, (*odonto*, and κεραμικός, 'potters' earth.') Relating or appertaining to porcelain teeth;—as the '*odontoceram'ic art*,' *Odontoc'er'amotech'ny*, the art of making porcelain teeth.

ODONTOCERAMOTECHNY, see Odontoceram'ic.

ODONTOCLA'SIS, (*odonto*, and κλάσις, 'fracture.') *Fractu'ra dent'is*. Fracture of a tooth.

ODONTOCNE'SIS, (*odonto*, and κνησις, 'itching.') Odaxismus.

ODONTOCNE'SMUS, Odaxismus.

ODONTODES, (ὀδοντωδής,) Odontoid.

ODONTODYNIA, (*odonto*, and ὀδυνή, 'pain.') Odontalgia.

ODONTOG'ENY, *Odontogen'ia*, *Odonto'sis*, (*odonto*, and γενεσις, 'generation.') Generation or mode of development of the teeth.

ODONTOGLYPHON, (ὀδοντογλυφόν, from *odonto*, and γλύφειν, 'to hollow.') Denticulipium, Gum lancet.

ODONTOGLYPHUM, Denticulipium, Gum lancet.

ODONTOG'RAPHY, *Odontogra'phia*, (*odonto*, and γραφή, 'a description.') A description of the teeth.

ODONT'OID, (ὀδοντοειδής,) *Odontoï'des*, *Odonto'des*, *Dentiform'is*, *Denta'lis*, *Dentoidēs*, *Pyrenoidēs*, (*odonto*, and εἶδος, 'shape, resemblance.') An epithet given to the *Processus dentatus* of the second vertebra or axis.

ODONTOID LIG'AMENTS, *Ligamen'ta ala'ria* seu *Maucha'r'ti*, (F.) *Ligaments odontoidiens*, are two strong and short conical fasciculi, whose truncated summits embrace the sides of the odontoid process, and whose bases are fixed in the fossæ at the inner side of the condyles of the occipital bone. Their direction is obliquely outwards and slightly upwards. They enter into the composition of the occipito-axoid articulation, strengthen the junction of the head with the vertebral column, and limit the movement of rotation of the atlas on the axis.

Other ligamentous fibres pass from the highest portion of the processus dentatus to the anterior margin of the foramen magnum of the occipital bone, constituting the *Ligamentum suspensorium dent'is*.

ODONTOID PROCESS, see Axis.

ODONTOLITHUS, (F.) *Odontolithe*. (*odonto*, and *λίθος*, 'a stone.') A sort of incrustation of a yellowish colour, which forms at the coronæ of the teeth, and is called *Tar'tar*, *Tar'tarus Dentium*, *Odontia incrustans*, *Calculus dentalis*. (F.) *Tartre des dents*, *T. dentaire*. It consists of seventy-nine parts of phosphate of lime, twelve and a half of mucus, one of a particular salivary matter, and seven and a half of animal substance, soluble in chlorohydric acid. A species of infusoria, *Denticula humilis*, has been found in it.

ODONTOLGY, *Odontolog'ia*, (*odonto*, and *λογος*, 'a discourse.') A treatise on the teeth.

ODONTOLOXIA, (*odonto*, and *λῶξις*, 'oblique.') Odontoparallaxis.

ODONTONECROSIS. Dental gangrene.

ODONTOPARALLAX'IS, *Odontolox'ia*, (*odonto*, and *παράλλιξις*, 'deviation.') Irregularity and obliquity of the teeth, (Prov.) *Snaggle teeth*.

ODONTOPHATNE, (*odonto*, and *φατή*, 'a crib, a stall.') Alveolus.

ODONTOPHATNITIS, (*odontophatne*, and *itis*.) Odontobothritis.

ODONTOPHYIA, (*odonto*, from *odonto*, and *φυω*, 'to grow.') Dentition.

ODONTOPLEROSIS, (*odonto*, and *πληρω*, 'I fill.') See Filling.

ODONTOPRISIS, (*odonto*, and *πρίσις*, 'gnashing of teeth.') Brygmus, Stridor dentium.

ODONTORRHIZA, (*odonto*, and *ρίζα*, 'root.') See Tooth.

ODONTOSEY'SIS, *Odontoseis'mus*, *Dentium vacillan'tia*, (*odonto*, and *σεισις*, 'shaking.') Loose-ness of the teeth.

ODONTOSISMUS, *Odontoseisis*.

ODONTOSIS, Dentition, Odontogeny.

ODONTOSMEGMA, (*odontosmegma*, from *odonto*, and *σμεγμα*, 'a detersive substance.') Dentifrice.

ODONTOSPHACELISIS, (*odonto*, and *σφακελις*, 'gangrene.') Dental gangrene.

ODONTOSPHACELISMUS, Dental gangrene.

ODONTOSTERE'SIS, (*odonto*, and *στερησις*, 'privation.') Loss of the teeth.

ODONTOSTYNERISMUS, (*odonto*, and *συνεπι-ζω*, 'to strike together.') (F.) *Claquement*. Chattering of the teeth.

ODONTOTECHNY, (*odonto*, and *τεχνη*, 'art.') Dentistry.

ODONTOTHECA, (*odonto*, and *θήκη*, 'a sheath.') Dental follicle.

ODONTOTHÈQUE, Dental follicle.

ODONTOTHERAPIA, (*odonto*, and *θεραπεία*, 'healing.') Dentistry.

ODONTOTHRYP'SIS, *Odontotripsis*.

ODONTOTRIMMA, gen. *Odontotrim'matis*, (*odonto*, and *τριβω*, 'to rub.') Dentifrice.

ODONTOTRIP'SIS, *Odontotryp'sis*, *Abra'sio den'tium*, (F.) *Usure des Dents*, (*odonto*, and *τριβις*, 'friction.') Abrasion or wearing away of the teeth.

ODONTOTRYPY, *Odonttrypy*.

ODONTTRYPY, *Odontotryp'y*, *Odontotryp'ia*, (*odonto*, and *πυρω*, 'I perforate.') The operation of perforating a tooth to evacuate purulent matter confined in the cavity of the pulp.

ODOR, gen. *Odo'ris*, Odour.

ODORAMENTUM, (from *odor*.) Odoriferum.

ODORANT, see Odour.

ODORAT, Olfaction.

ODORATIO, Olfaction.

ODORATUS, Olfaction — o. Deperditus, Anos-mia.

ODORIFEROUS GLANDS, see Sebaceous glands.

ODORIFERUM, (*odor*, and *fero*, 'I carry;') *Odoramen'tum*. A medicine that gives odour or flavour. A scent. A perfume.

ODOROUS, see Odour.

ODOS, (ὁδός,) Way.

O'DOUR. O'dor, Od'mē, Os'mē, (F.) *Odeur*. A smell. Odours are subtle particles, constantly escaping from the surface of certain bodies hence called *odorons*, (F.) *Odorant*. They act, in some manner, by actual contact with the nerves of the Schneiderian membrane, and give occasion to the sense of smell or olfaction.

ODOUS, (ὁδός, gen. ὁδός,) see Teeth.

OD'YNE, (*odyn*, 'pain;') *Do'lor*, *Odyne'ma*. A very common suffix to words: as in *Pseudodyn-*.

ODYNEMA, gen. *Odyne'matis*. (ὀδύνημα, / Odyne.

ODYNEPHO'BIA, (*odyne*, and *φοβία*, 'I fear,') Fear or dread of pain.

ODYNODES, (ὀδυνοδός,) Painful.

ODYNOPOETIC, *Odynopoe'ticus*, *Dolorif'ic*us, *Dolorif'ic*, (F.) *Odynopoe'ticus*, *Dolorif'ique*, (*odyne*, and *ποιω*, 'to make,') Causing pain.

ŒCIOMANIA, *Œciomani'a*, (ὀκνέσις, 'belonging to a household or family,' and *mania*.) A form of moral insanity, characterized by domestic perversity.

ŒCONOMIA, (ὀικονομία, from οἶκος, 'a house,' and *νομος*, 'law,') Economy — œ. Animalis, Physiology.

ŒDALICUS, (ὀιδῶν, 'to swell,') Œdematous.

ŒDALIUS, (ὀιδῶλος,) Œdematous.

ŒDE'MA, gen. *Œde'matis*, (ἄδῆμα, 'a swelling,')

Hydrone'cus, *Œdemat'ia*, *Hydron'cus*, *Hydaton'cus*, *Hydrocœ'dema*, *Hyd'cœ'dema*, *Hy'drops cellula'ris ar'tium*, *Phlegmat'ia*, *Leucophlegmat'ia*, *Eo-phy'ma œdemat'icum*, (F.) *Œdème*, *Œdématis*, *Enflure*. Swelling produced by the accumulation of a serous fluid in the interstices of the areolar texture. This swelling is soft; yields under the finger; preserves the impression for some time, and is pale and without pain. It presents the same characters as anasarca, which is general œdema. Its etiology and treatment are also the same. See Anasarca.

ŒDEMA ACUTUM, see Anasarca.

ŒDEMA ARSENICALIS. The swelling of the eyelids and face, induced by continued use of the preparations of arsenic.

ŒDEMA CALIDUM, see Anasarca — œ. Capitis Hydrocephalus externus.

ŒDEMA CEREBRI, (F.) *Œdème du cerveau*.

Œdema of the brain. A condition of the cerebral pulp, in which there is an infiltration of serous fluid into it, so that it appears more moist or watery than common; and, when sliced or pressed, small drops of water are seen to ooze out.

ŒDEMA, COMPACT, Induration of the cellular tissue — œ. Cruentum, Sanguillation — œ. Febrile, see Anasarca — œ. Fugax, Anathymiasis.

ŒDEMA OF THE GLOTTIS, (*Œ. glottidis*, *Œdem'atous Laryng'itis* or *ang'ina*, *Laryng'itis œdemat'ica* seu *œdemat'ica* seu *seropurulent'a*, *Hy'drops glottidis*, *Ang'ina aquo'sa* seu *laryng'e'a œdemat'ica*, *Submucosa Laryng'itis*, (F.) *Laryngite œdémateuse*, *L. œdémateuse et séro-purulente* ou *ang-glottique* ou *sub-muqueuse*, *Angine laryngée œdémateuse*, *Œdème de la glotte*. A disease consisting of serous or seropurulent infiltration into the submucous tissue of the glottis. The symptoms resemble those of croup; but the disease attacks the adult rather than the child. The age is, indeed, a principal means of diagnosis between the two affections.

The disease is almost always fatal. The treatment has to vary according to the accompanying general symptoms.

ŒDEMA HYSTERICUM, Anathymiasis.

ŒDEMA LACTEUM, Phlegmatia dolens.

ŒDEMA OF THE LUNGS, (*Œ. pulm'onum*, *Pneumock'ysie*, *Pneumonœ'dema*, *Hy'drops* seu *anasar'ea* *Pulmo'num*, *Hydropneumo'nia*, (F.) *Œdème du Pommou*. Laënnec has so called the infiltration of serum into the tissue of the lung, carried to such an extent as to diminish its permeability

to air. It is not an uncommon sequela of pneumonia, and the major exanthemata. The respiration is laborious; the respiratory murmur scarcely perceptible, although the thorax is largely expanded, and there is a slight *râle crépitant*, particularly at the base and inferior part of the lung. The sound on percussion is clear, and on both sides equally so. The cough is attended with aqueous expectoration. In some cases the respiration becomes purile in a small portion of the summit of the lung.

ŒDEMA NEONATORUM, Induration of the cellular tissue—œ. Palpebrarum, Hydroblepharon—œ. Puerperarum, Phlegmatia dolens—œ. Pulmonum, Œdema of the lungs—œ. Pulmonum chronicum, Hydropneumonia—œ. Scroti aquosum, Oschydroedema—œ. Scroti cruentum, Oschæmatœdema—œ. Scroti purulentum, Oscheopyœdema—œ. Scroti urinosum, Urocele—œ. Spasticum, Anathymiasis—œ. Uvulae, Staphylœdema.

ŒDEMATIA, Œdema.

ŒDÉMATIE, Œdema.

ŒDEMATOSARCA, Œdemosarca.

ŒDEMATOSCHEOCELE, Œdemoscheocèle, Oscheocèle Œdématic. Oscheocèle with œdema of the serotum.

ŒDEMATOUS, Œdemato'sus, Œdématicus, Œdemato'des, Œdal'ius, Œdal'icus, Œdem'atous, (F.) Œdémateux. Affected with œdema, (F.) Œdématic; of the nature of œdema.

ŒDÈME, Œdema—œ. du Cerveau, Œdema cerebri—œ. Douloureux des femmes en couche, Phlegmatia dolens—œ. de la Glotte, Angina œdematosa, Œdema of the glottis—œ. des Nouvelles accouchées, Phlegmatia dolens—œ. actif des Nouvelles accouchées, Phlegmatia dolens—œ. du Tissu cellulaire des nouveau-nés, Induration of the cellular tissue.

ŒDEMOSAR'CA, (œdema, and σαρκ, gen. σαρκος, 'flesh,') Œdematosar'ca. A species of tumour mentioned by M. A. Severinus, which holds a middle place between œdema and sarcoma.

ŒDEMOSCHEOCELE, Œdematoscheocèle.

ŒIL, (oculus,) Eye—œ. de Bœuf, Anthemistinctoria—œ. Double, Binoculus—œ. de Lièvre, Lagophthalmus—œ. Simple, see Monoculus.

ŒILLET GIROFLÉE, Dianthus caryophyllus.

ŒILLIÈRE, (from œil,) Scaphium oculare.

ŒNANTHE, Œ. crocata'ta seu cherophyll'i fo'liis, Hem'lock drop'wort, Hem'lock water drop'wort. Nat. Ord. Umbelliferae. Sec. Syst. Pentandria Digynia, (œnos, and andes, 'a flower,' its flowers smelling like the vine.) A violent poison of the acro-narcotic class. Its juice has been recommended in certain cutaneous diseases; but it is scarcely ever used. It is employed in fomentations.

ŒNANTHE AQUATICA, Phellandrium aquaticum—œ. Phellandrium, Phellandrium aquaticum—œ. Striata rigida, Seseli tortuosum.

ŒNELÆ'ON, (œnos, and ελαιον, 'oil,') A mixture of oil and wine. Also, rectified spirit or alcohol.

ŒNODES, (οινωδης, from œnos, and odes,) Vinous.

ŒNOG'ALA, gen. Œnogalac'tis, (οινοναλα, from œnos, and γαλα, 'milk,')—'a mixture of wine and milk.' According to some, wine as warm as new milk.

ŒNOG'ARON, Garum vino mistum. Wine mixed with garum.

ŒNOIDES, (οινωειδης, from œnos, and ειδος, 'resemblance,') Vinous, Winelike.

ŒNOLATURE, (from œnos,) see Vinum.

ŒNOLE, see Vinum.

ŒNOLOTIF (F.). A name given by M. Bérat to a medicine intended for external use, whose excipient is wine.

ŒNOMA'NIA, Oinoma'nia, (œnos, and mania,) Delirium tremens. Properly, wine-mania. An insatiable desire for intoxicating liquors.

Œ'NOMEL, Œnom'eli, (œnos, and μελι, 'honey,') Mel vino'sum, Mul'sum Vino'sum. Honey wine. Wine mead. See Hydromel.

ŒNOPHLYGIA, (αινοφλυγια, from œnos, and φλυα, 'to overflow,') Temulentia.

ŒNOPHLYXIS, (αινοφλυξις,) Temulentia

ŒNOS, (οινος,) Wine.

ŒNOSTAGMA, gen. Œnostag'matis, (œnos, and σταζω, 'to distil,') Spiritus vini rectificatas.

ŒNOTHE'RA BIEN'NIS, (œnos, and θηρ, 'wild,') [?] Œ. Mollis'sima seu murica'ta seu gau'roi'des, On'agra, Eve'ning Prim'rose, Tree Prim'rose, Prim'rose tree, Cureall, Sea'bish. An indigenous plant, common on the borders of fields, and in natural hedges. Ord. Onograceæ. Sec. Syst.

Oetandria Monogynia. Its properties are mucilaginous and slightly acid. A decoction has been used in cases of infantile and other eruptions.

ŒNOTHERA GAUROIDES, Œ. biennis—œ. Muricata, Œ. biennis.

ŒSOPHAGE, Œsophagus.

ŒSOPHAGE'AL, or ŒSOPHAG'EAL, Œsophag'eus, (from œsophagus,) (F.) Œsophagien. Relating or belonging to the œsophagus.

ŒSOPHAGEAL AP'ERTURE OF THE DI'APHRAGM, For'a'men œsophag'eum, (F.) Ouverture œsophagienne du diaphragme. An opening in the diaphragm for the passage of the œsophagus.

ŒSOPHAGEAL AP'ERTURE OF THE STOM'ACH. A name given to the superior or cardiac orifice of the stomach, to distinguish it from the inferior or pyloric. See Cardia.

ŒSOPHAGEAL MUS'CLE, Œsophag'eus. Some anatomists have given this name to the transverse muscular fibres which surround the œsophagus at its upper extremity.

ŒSOPHAGEAL TUBE, see Tube, œsophageal.

ŒSOPHAGEURYS'MA, gen. Œsophageurys'matis, (Œsoph'agus succenturiat'us, (œsophagus, and ευρυσμος, 'dilatation,') Dilatation of the œsophagus.

ŒSOPHAGIALGIA, (œsophagus, and αλγος, 'pain,') Œsophagismus.

ŒSOPHAGIARTICIA, (œsophagus, and arctare, 'to make narrow,') Dysphagia constricta.

ŒSOPHAGIEN, Œsophageal—œ. (muscle), Constrictor Œsophagi.

ŒSOPHAGIS'MUS, Œsophagis'mum, Spasm of the œsophagus, Dysph'agia Spasmod'ica seu Spas'tica seu Nerv'osa, Strict'ura œsoph'agi spasmod'ica, Œsophagial'gia, Œsophagospas'mus, Tenes'mus Gu'læ, (F.) Ténèsmè de l'œsophage. By some used synonymously with œsophagitis, dysphagia, and contraction of the œsophagus.

ŒSOPHAGITIS, Angi'na seu Cynan'chè œsophag'e'a, Dysph'agia inflammato'ria, Inflammato' œsoph'agi seu Gu'læ, (F.) Œsophagite, Angine œsophagienne, (œsophagus, and itis,) Inflammation of the œsophagus.

ŒSOPHAGODYN'IA, (œsophagus, and οδυνη, 'pain,') Pain in the œsophagus.

ŒSOPHAGOPATHI'A, Mor'bus Œsoph'agi, (œsophagus, and παθος, 'affection,') A morbid condition of the œsophagus.

ŒSOPHAGOPLE'GIA, Dysph'agia paralyt'ica, Lætoparaly'sis, (œsophagus, and πλεγη, 'a stroke,') Paralysis of the œsophagus.

ŒSOPHAGORRHAG'IA, (œsophagus, and ραγη, 'a rupture,') Hemorrhage from the œsophagus.

ŒSOPHAGORRHŒA, (œsophagus, and ρευω, 'I flow,') Discharge of blood or mucus from the œsophagus.

ŒSOPHAGOSPAS'MUS, (œsophagus, and σπασμος, 'spasm,') Spasm of the œsophagus.

ŒSOPHAGOSTENOMA, (œsophagus,

στενός, 'narrow,' and oma.) Dysphagia constricta.

ŒSOPHAGOTOMY, (Œsophagotom'ia) (F.) Œsophagotomie, (œsophagus, and τέμνω, 'to cut.') An incision made into the œsophagus for the purpose of extracting foreign bodies from it.

ŒSOPHAGUS, (œsôphagos, from œw, future œσω; I carry,' and φαγω, 'I eat;') *Gu'ta, Fis'tula cibalis, Fis'tula seu Infundib'ulum Ventriculi, Vi'a stom'achi et ven'tris, Glu'tis*, (F.) Œsophage. The Gullet. A musculomembranous canal, cylindrical, and depressed from before to behind, which extends from the inferior extremity of the pharynx to the upper orifice of the stomach. At its origin, it is situate in the median line; but beneath the larynx, it deviates to the left, and in the chest experiences different reflections. In its cervical portion, it corresponds, behind, with the spine; before, with the larynx and trachea; and, at the sides, it is close to the primitive carotids, internal jugular veins, par vagum, recurrent nerves, &c. In its inferior or thoracic portion, the œsophagus is entirely contained in the posterior mediastinum; and enters the abdomen through the œsophageal aperture, *Fora'men œsophage'um*, of the diaphragm. The œsophagus is composed of a very strong muscular layer, sometimes called *Tu'nica vagina'lis gu'tis*; formed, itself, of two sets of fibres, the external being generally longitudinal, the internal transverse or annular; and of a mucous membrane which is soft, fine, thin, and white, especially at its lower part. It is continuous, above, with the mucous membrane of the pharynx. The mucous follicles, found beneath it, are not numerous, and have been called *Œsophage'al glands*. The arteries of the œsophagus proceed, in the neck, from the thyroid; in the chest, from the bronchial arteries and directly from the aorta;—in the abdomen, from the inferior phrenic, and coronaria ventriculi. Its veins empty themselves into the inferior thyroid, the vena cava superior, the internal mammary, azygos, bronchial, phrenic, and coronaria ventriculi. Its lymphatics pass into the ganglia surrounding it. Its nerves are afforded by the pharyngeal and pulmonary plexuses; by the cardiac nerves; the thoracic ganglia, and, especially, by the pneumogastrics and their recurrent branches.

ŒSOPHAGUS SUCCENTURIATUS, Pharyngocele. ŒSTROMANIA, (œstros, 'imperious erotic impulse,' and mania,) Nymphomania.

ŒSTRUAL, *Œstru'al'is*, (from *œstrum*.) Appertaining or relating to orgasm, especially venereal orgasm.

ŒSTRUATION, see Heat.

ŒSTRUM, Clitoris, Orgasm.

ŒSTRUM VEN'ERIS, *Œ. vene'reum, Œ'strus vene'reus*; (œstros, œstrus, 'imperious erotic impulse;') A vehement desire for sexual intercourse. Orgasm. With some, *œstrum* signifies Clitoris.

ŒSTRUS, (œstros,) Clitoris, Orgasm, see Ectozoa.—œ. Veneres, Œstrum venereum.

ŒSYSPUS, (œsytos, from œs, 'a sheep;') The greasy matter of unwashed wool; formerly employed in friction in diseased joints.

ŒUF, Ovum, *Œ. de De Graaf*, Folliculi Graafiani—*œ. de Naboth*, Nabothi glandulæ.

OFFICE, PHYSICIAN'S or SURGEON'S, Iatron.

OFFICINA, for *Opificina*, (from *opifex*, (*opus*, and *facio*), 'a workman;') 'a shop,' Pharmacopolium.

OFFIC'INAL, *Officina'lis*. An epithet for medicines found in the shop of the apothecary, ready prepared—*usu'al'ia*; in opposition to *magistral* or *extemporaneous*,—those prepared after the prescription of the physician.

OFFIUM, Affion, Opium.

OFFSPRING, Epigone.

OFFUSCATIO, (*of*, for *ob*, and *fuscare*, 'to make dark;') Amaurosis.

OGLA, Oogala.

OGNON or OIGNON, (from *unio*), Allium cepa, a Corn—o. *Marin*, Scilla.

OH'IO, MIN'ERAL WA'TERS OF. *Yel'low Spring* is a chalybeate, situate in Greene county, 64 miles from Cincinnati. It is somewhat frequented.

OID, *Oi'des*, (οἶδος, from εἶδος, 'form, resemblance;') A suffix, denoting 'resemblance,' as in ethmoid, sphenoid, &c.

ODIUM ABORTIFACIENS, (*ovum*, 'an egg,' and *oidos*, 'resemblance;') see Ergot—o. Albicans, Aphthaphyte—o. Schoenleini, see Porrigo favosa.

OIE, Sorbus domestica.

OIGNON, Ognon.

OIKEIOMANIA, Œciomania.

OIKIOMIASM'ATA, *Oikeiomiasm'ata*, (from οἶκος, belonging to a household and *miasmata*.) Morbific emanations from domestic sources.

OIL, ([Sax.] *el*), *O'leum, Elæ'on*, (F.) *Huile*; (from *elaia*, 'the olive;') A collective name, under which two classes of fluids are included, very different from each other: those belonging to the one class, are viscid, mawkish, or almost insipid; those of the other are nearly devoid of viscosity, and are caustic and very volatile. The former are called *fat or fixed oils*; the latter *volatile or essential oils or essences*.

OIL, see Ricinus communis.

OIL OF ALMONDS, *Oleum amygdalarum*—o. of bitter Almonds, *Oleum amygdalæ amaræ*—o. of Amber, rectified, see *Succinum*—o. of Andæ, see Andæ—o. Animal, *Oleum animale*—o. Animal, of Dippel, *Oleum animale Dippelii*—o. of Bay, *Daphnelæon*—o. of Bays, *Oleum laurinum*—o. and Beeswax, *Unguentum ceræ*—o. of Ben, see *Guilandina moringa*—o. of Benjamin or Benzoin, *Oleum benzoini*—o. Benne, see *Sesamum orientale*—o. of Bergamot, see Bergamot.

OIL OF BONES, see Marrow.

OIL, BRITISH. An empirical preparation often used in cases of sprains. A committee of the Philadelphia College of Pharmacy recommended the following form for its preparation.—*Ol. Terebinth*, *Ol. Lini usitatis*. aa f fij , *Ol. Succini*, *Ol. Juniper*. aa f fij , *Petrol. Barbadaens*. f fij , *Petrol. American*. (*Seneca Oil*), f fij . M.

OIL OF CACAO, Butter of Cacao—o. of Cade, see *Juniperus Oxycedrus*—o. Camphor, see Camphor—o. of Canada fleabane, *Oleum Erigerontis Canadensis*—o. Carron, *Linimentum aquæ calis*—o. of the Cashew nut, see *Anacardium occidentale*—o. Castor, see *Ricinus communis*—o. of Chabert, *Oleum animale empyreumaticum Chaberti*—o. of Cinnamon, see *Laurus cinnamomum*—o. of Cloves, *Eugenia caryophyllata*—o. Cod, o. *Jecoris aselli*—o. Codliver, o. *Jecoris aselli*—o. of Copaiba, *Oleum Copaibæ*—o. Cornspirit, Oil, fusel—o. of Cubebs, see *Piper cubeba*—o. of Cypress, *Cyprium oleum*—o. of Dill, see *Anethum graveolens*—o. Dippel's, *Oleum animale Dippelii*—o. of Egg, see *Ovum*—o. Ethereal, *Oleum Æthereum*—o. of Fennel, see *Anethum*—o. of Fern (male), see *Polypodium filix mas*—o. Flaxseed, see *Linum usitatissimum*.

OIL, FUSEL, *Al'cohol amylicum* (Ph. U. S., 1863), *Fermento'leum Sola'ni*, *Fow'sel oil, Potat'o oil, Grain oil, Corn'spirit oil, Potat'o spir'it oil, Oil of potat'oes, Amylic al'co'hol, Hy'drated ox'ide of am'yle*, ([G.] Fusel, an epithet for poor brandy.) An acrid volatile oil, formed in the manufacture of potato or grain brandy, which is not easily separable from it, and is obtained by continuing the process after the ordinary spirit has ceased to come over. Its chemical constitution is analogous to that of alcohol. It exhales a powerful and peculiarly suffocating odour.

S. g. 818. In small doses it is highly stimulating,—acting like narcotics in general. In large doses it destroys the mucous membrane of the stomach. It is used by the British Pharmacopœia in the preparation of *Sodæ valerianæ*.

OIL OF GERANIUM, *Oleum graminis Indici*—o. Gingergrass of Nemauro, *Oleum graminis Indici*—o. Grain, Oil, fusel—o. Grass of Nemauro, *Oleum graminis Indici*.

OIL, HÆR'LEM. An empirical preparation supposed to consist chiefly of petroleum, turpentine, and balsam of sulphur. Used internally in renal and rheumatic affections.

OIL, KÉKUNÉ, see *Aleurites triloba*—o. Krumholz, see *Pinus mughos*—o. Kukin, see *Aleurites triloba*—o. of Lemons, see *Citrus medica*—o. Lily, see *Lilium candidum*—o. Linseed, see *Linum usitatissimum*—o. of Mace, expressed, see *Myristica*—o. of Mastich, see *Pistacia Lentiscus*—o. of Mucilages, *Oleum e mucilagibus*—o. Neats-foot, *Oleum bubulum*—o. of Nicodemus, *Nicodemium oleum*—o. Nut, *Juglans cinerea*—o. of Nutmegs, see *Myristica moschata*—o. Olive, *Oleum olivæ*—o. Palm, see *Cocos butyracea*—o. Paper, *Pyrothonide*—o. Parsley, *Apiol*—o. Pennyroyal, see *Hedeoma pulegioides*—o. of Pike, *Oleum lucii piscis*—o. Poppy, see *Papaver*—o. of Potatoes, Oil, fusel—o. Potato, Oil, fusel—o. Potato spirit, Oil, fusel—o. Rag, *Pyrothonide*—o. Red, see *Hypericum perforatum*—o. Rock, Petroleum—o. Roosa, *oleum graminis Indici*—o. of Rose Geranium, *Oleum graminis Indici*—o. of Roses, see *Rosa centifolia*—o. of Rue, *Peganelæon*, see *Ruta*—o. Salad, *Oleum olivæ*—o. of Scorpion, see *Scorpion*—o. Seneca, Petroleum, New York—o. Skatelliver, see *Oleum Jecoris aselli*—o. of Spike, *Oleum terebinthinæ*—o. of Spike, see *Lavandula spica*—o. of Spikenard, *Oleum graminis Indici*—o. Sulphuretted, *Balsamum sulphuris simplex*—o. of Tartar, *Liquor potassæ subcarbonatis*—o. of Theobroma, *Butter of Cacao*—o. of Thyme, see *Thymus*—o. of Tobacco, *Oleum tabaci*—o. of Turpentine, *Oleum terebinthinæ*—o. of Turpentine, rectified, *Oleum terebinthinæ rectificatum*—o. of Valerian, *Oleum valerianæ*—o. of Vitriol, *Sulphuric acid*—o. Walnut, Belgium, see *Aleurites triloba*—o. Walnut, Spanish, see *Aleurites triloba*—o. of Wine, *Oleum aethereum*.

OILS, ANIMAL, *Olea animalia*—o. Distilled, *Olea volatilia*—o. Empyreumatic, *Olea empyreumatica*—o. Essential, *Olea volatilia*—o. Ethereal, *Olea volatilia*—o. Expressed, *Olea fixa*—o. Fatty, *Olea fixa*—o. Fixed, *Olea fixa*—o. Fugacious, *Olea fugacia*—o. Medicinal, *Olea medicinalia*—o. Volatile, *Olea volatilia*.

OILY, *Oleaginous*—o. Grain, *Sesamum orientale*.

OLINOMANIA, *Enomania*.

OLINOS KRITHES, (*ολινος κριθης*, 'barley wine,') *Cerevisia*.

OINTMENT, *Unguentum*—o. of Aconitia, *Unguentum Aconitiæ*—o. Antimonial, *Unguentum antimonii*—o. of Antimony, tartarized, *Unguentum antimonii*.

OINTMENT, ARSENICAL, (of *Sir Astley Cooper*.) This is made of *Arsenious acid*, 3j; *Sulphur*, 3i; *Spermacei cerate*, 3i. It is spread on lint, and applied to cancerous sores.

OINTMENT OF ATROPIA, *Unguentum atropiæ*—o. Banyer's, *Unguentum Calomelanos compositum*—o. Basilicon, *Unguentum resinæ*—o. Bay, *Unguentum laurinum*—o. of Belladonna, *Unguentum belladonnæ*—o. of Benzoin, *Unguentum Benzoini*—o. Blister, *Cerate of cantharides*—o. Blistering, green, *Unguentum lyttæ medicatum*—o. Blistering, milder, *Unguentum lyttæ*—o. Blistering, yellow, *Unguentum lyttæ medicatum aliud*—o. Blue, *Unguentum hydrargyri*—o. of Calomel, *Unguentum calomelanos*—o. of Calomel, compound, *Unguentum calomelanos compositum*—o.

Citrine, *Unguentum hydrargyri nitratis*—... of *Cocculus Indicus*, *Unguentum cocculi*—o. of *Creasote*, *Unguentum creasoti*—o. *Cucumber*, see *Cucumis sativus*—o. *Cyrillo's*, *Unguentum muratis hydrargyri oxygenati medicatum*—o. Digestive, simple, *Unguentum digestivum simplex*—o. Edinburgh, see *Unguentum veratri*—o. Elder, *Unguentum sambuci*—o. of Elemi, *Unguentum elemi*—o. for the Eyes, (*Smellome's*,) see *Ceratum resinæ*—o. of Galls, *Unguentum gallæ*—o. of Galls, compound, *Unguentum gallæ compositum*—o. Garlic, see *Allium*—o. of Glycerin, see *Fat*—o. Golden, *Unguentum hydrargyri nitrico-oxidi*—o. Golden, Singleton's, see Singleton's golden ointment—o. Goulard's, *Ceratum plumbi compositum*—o. Green, *Unguentum sambuci*—o. Heliebore, white, *Unguentum veratri*—o. Helmerich's, *Unguentum sulphuratum alcalinum ad scabiem*—o. of Hemlock, *Unguentum conii*—o. of Hydriodate of Potass, *Unguentum potassæ hydriodatis*—o. of Iodide of Lead, *Unguentum plumbi iodidi*—o. of Iodide of Mercury, *Unguentum hydrargyri iodidi*—o. of red Iodide of Mercury, *Unguentum hydrargyri iodidi rubri*—o. of Iodide of Sulphur, *Unguentum sulphuris iodidi*—o. of Iodine, *Unguentum Iodini*—o. of Iodine, compound, *Unguentum iodini compositum*—o. Issue, Dr. Physick's, see *Unguentum lyttæ medicatum aliud*—o. Itch, *Unguentum sulphuris compositum*—o. Itch, Bailey's, see *Unguentum sulphuratum ad scabiem*—o. Itch, Bateman's, see *Unguentum sulphuratum alcalinum ad scabiem*—o. Itch, Helmerich's, *Unguentum sulphuratum alcalinum ad scabiem*—o. Itch, Jackson's, *Unguentum sulphuris compositum*—o. of Lard, *Unguentum ceræ*—o. Lead, compound, *Unguentum plumbi compositum*—o. of white oxide of Lead, *Unguentum oxidi plumbi albi*—o. of Lydia, *Bacaris*—o. Marshmallow, *Unguentum de althæâ*—o. Mercurial, *Unguentum hydrargyri*—o. Mercurial, milder, see *Unguentum hydrargyri*—o. Mercurial, strong, see *Unguentum hydrargyri*—o. of gray oxide of Mercury, *Unguentum oxidi hydrargyri cinerei*—o. of nitrate of Mercury, *Unguentum hydrargyri nitratis*—o. of nitric oxyd of Mercury, *Unguentum hydrargyri nitrico-oxidi*—o. of Meze-reon, *Unguentum mezerii*—o. of Nitrous acid, *Unguentum acidi nitrosi*—o. of Pitch, *Unguentum resinæ nigre*—o. of Pitch, black, *Unguentum resinæ nigre*—o. Plunkett's, for cancer, *Plunkett's ointment*—o. red Precipitate, *Unguentum hydrargyri nitrico-oxidi*—o. of white Precipitate, *Unguentum hydrargyri præcipitati albi*—o. Resin, *Ceratum resinæ flavæ*—o. Resin, black, *Unguentum resinæ nigre*—o. of Rosewater, *Unguentum aquæ rosæ*—o. Saturnine, *Unguentum plumbi superacetatis*—o. Savine, *Ceratum Sabinæ*—o. Soldier's, *Martiatum unguentum*—o. of Spanish flies, *Cerate of Cantharides*, *Unguentum lyttæ*—o. Spermacei, *Unguentum cetacei*—o. of Stramonium, *Unguentum stramonii*—o. of Subacetate of copper, *Unguentum subacetatis cupri*—o. of Sugar of lead, *Unguentum plumbi superacetatis*—o. Sulphur, *Unguentum sulphuris*—o. Sulphur, compound, *Unguentum sulphuris compositum*—o. Sympathetic, *Hoplochrisma*—o. Tar, *Unguentum picis liquidæ*—o. Tartar emetic, *Unguentum antimonii*—o. Tobacco, *Unguentum Tabaci*—o. Tutty, *Unguentum oxidi zinci impuri*—o. of Veratria, *Unguentum veratriæ*—o. Verdigris, *Unguentum subacetatis cupri*—o. Wax, *Unguentum ceræ*—o. White, *Unguentum oxidi plumbi albi*—o. Zinc, *Unguentum zinci*—o. of impure oxide of Zinc, *Unguentum oxidi zinci impuri*.

OIOCALYMN'MA, gen. *Oiocalymn'atis*, *Oiocalypt'rum*, (*ωιω, ωω, 'egg,' and καλυπται, 'to cover,'*) *Membrana O'vuli cortica'lis*. The membrane of the eggshell.

OKEN'S BODY, *Corpus Wolffianum*.

OKRA, *Hibiscus esculentus*.

OLD AGE, *Senectus*.

OLD MAN, *Artemisia abrotanum*.

OLD WIFE'S SHIRT, *Liriodendron*.

O'LEA ANIMALIA, *Animal oils*. (F.) *Huiles animales*. A name given to fixed oils holding in solution the mucilaginous and gelatinous principles of certain animals, as the *Oil of frogs*, *Oil of scorpions*, *Oil of spiders*, &c. Sometimes, also, the term *animal oils* is given to empyreumatic oils, produced during the decomposition of animal substances by heat.

OLEA DESTILLATA, O. Volatilia.

OLEA EMPYREUMATICA, *Empyreumatic oils*. (F.) *Huiles empyreumatiques ou pyrolytiques*, *Pyrolytiques*. Oils which have an empyreumatic or burnt smell. They are obtained by treating vegetable or animal matters by heat, in close vessels. They do not exist in organized bodies, but are formed during their decomposition by fire. The animal oil of Dippel is an empyreumatic oil.

OLEA, O. *Europea*, O. *orientalis* seu *lunifolia* seu *polymorpha* seu *Gallica*, *Olivea*, *Olive tree*, *Elia*, (F.) *Oliver*; (Fruit) *Oliva*. *Nat. Ord.* *Jasminaceae*. *Sex. Syst.* *Diandria Monogynia*. The leaves of the olive are bitter, and an extract prepared from them, and a substance called *oliva*, the bitter principle of the leaves, have been given in Italy as antiperiodics. The fruit, when pickled, is extremely grateful to some stomachs. Olives, as met with in the shops, are prepared from the green, unripe fruit, repeatedly steeped in water. To this, some quicklime or alkaline salt is added, and, afterwards, they are washed and preserved in a pickle of common salt and water. From this fruit is prepared the *Olive oil* or *Salad oil* of the Pharmacopœias, which is obtained by grinding and pressing the olives, when thoroughly ripe. The finer and purer oil issues first by gentle pressure, and the inferior sort on heating what is left, and pressing it more strongly. See *Oleum Olivæ*. In Calabria, an odorous resin exudes from its trunk, which is employed as a perfume by the Neapolitans.

A gum flows from certain wild olives, in warm countries, which consists, according to Pelletier, of a resin, a little benzoic acid, and a peculiar substance, called *Olivette*. It is in yellow masses, of a slightly acrid taste, and of a vanilla smell. It is called *O'lea gummi*, O. *resina*, and O. *balsamum*, (F.) *Gomme olivier*, and is esteemed astringent and detensive.

O'LEA FIXA seu PIN'GUA, *Expressed*, *Fixed* or *Fatty oils*. (F.) *Huiles fixes ou grasses*, *Oléa*. All the oils obtained from the seeds or pericarps of vegetables, without distillation, and which are viscid, but slightly odorous and rapid; lighter than water, and insoluble in alcohol. The rancidity of oils depends on the absorption of oxygen, and therefore they should be kept in bulk as much as possible, and in narrow-necked bottles, so that a very small surface only can be exposed to the air. All the fixed oils are emollient, and, in a certain dose, they act as purgatives and emetics. They are prepared by expressing the fruit or seed containing them.

OLEA FUGACIA, *Fugacious oils*. (F.) *Huiles fugaces*. A name given to oils which are so volatile that, in order to obtain them, recourse must be had to a different process from that employed for other essential oils. Such are the oils of *jessamine*, *lily*, *violet*, &c.

OLEA GALICA, O. *Europea*—O. *Lancifolia*, O. *Europea*.

OLEA MEDICINALIA, *Medicinal oils*. (F.) *Huiles Médicinales*, *Éléa*, *Solutions parles huiles*. A name given to oils prepared by macerating, infusing or boiling medicinal substances in olive or any other fixed oil. These oils may then be

regarded as oily solutions of certain medicinal substances; whence they can never be simple. They have, however, been divided into *simple* and *compound medicinal oils*. To the former belong the *Oils of St. John's wort*, of the *Solanum nigrum*, &c.; to the other—which have often been called *Oily Balsams*, (F.) *Baumes huileux*—the *Balsams of Fioravanti*, *Metc.*, &c. Medicinal oils are, almost always, employed externally.

OLEA POLYMORPHA, O. *Europea*—O. *Sativa*, O. *Europea*.

OLEA VOLATILIA, O. *destillata*, *Ethereal*, *Volatile*, *Essential*, or *Distilled Oils*, *Essences*, (F.) *Huiles volatiles ou essentielles ou distillées ou échantées, volatiles, oléales*. Oils found in aromatic vegetables, and in every part of them, except in the interior of the seeds. The majority are obtained by distillation; but some by expression. They possess unctuousity, inflammability, and viscosity, like the fixed oils; but they are generally odoriferous, pungent, and acrid. The greater part are lighter than water; but some are heavier, and congeal at a moderate temperature. They dissolve, in small quantity, in distilled water, by simple agitation. Almost all are soluble in alcohol. The odour and taste are the usual tests of their goodness. To preserve them, they should be kept in a cool place, in small bottles, quite full and well corked. Volatile oils are possessed of the aromatic properties of the plants whence they are obtained. They are all, when applied externally, stimulant and rubefacient. Pharmaceutical preparations, in which substances are dissolved in volatile oils, (F.) *Solutions par les huiles volatiles*, are termed by Bérard, *Myrades*.

OLEAGINOUS, *Oleaginous*, *Oleoous*. (F.) *Oléagineux*; (from *oleum*.) Oily; containing oil,—as '*an oleaginous or oily mixture*.'

OLEA MEN, gen. *Oleamina*, *Oleamenium*, (from *oleum*.) Any soft ointment prepared of oil.

OLEANDER, (from *olea*, 'the olive,') *Rhododendron chrysanthemum*.

OLECRANARTHRI'TIS, (*olecranon*, *αρθρον*, 'joint,' and *itis*.) Inflammation of the elbow joint.

OLECRANARTHROCA'CE, (*olecranon*, *αρθρον*, 'articulation,' and *κατα*, 'vice or disease,') Inflammation of the articular surfaces of the elbow.

OLECRANIAL, (F.) *Olécranien*. Relating or belonging to the olecranon.

OLECRANOID CAVITY, see *Ulna*.

OLECRANON, *Olecranon*, *Olecranium*, (*ωλεκράνον*.) *Olecranon mobile*, (F.) *Olécrane*, (*ωλεκη*, 'the elbow,' and *κρανον*, 'the head,') *Acromion*, *Additamentum humeri*, *Ancon*, *Processus anconæus* seu *uncinatus*, *Glan* seu *Coro'na* seu *Coro'na post'rior* seu *Additamentum uncatum Ulnæ*, *Ver'tex Cu'biti*, *Patell'a fix'a*, *Rostrum externum* seu *post'erior*, *Top of the cubit*. Head or protection of the elbow. A large process at the upper extremity of the ulna, on which we lean. When this process is fractured, it is apt to be drawn up by the triceps, and much care is required to keep the parts in apposition.

OLEFIANT GAS, CHLORIDE OF, see *Anæsthetic*.

OLEIN, (from *oleum*.) see *Fat*—O. *Codliver*, see *Oleum jecoris aselli*.

OLENE, (*ωλενη*.) *Ulna*.

OLEOCERATUM, *Cerate*—O. *Aqua subacutum*, *Ceratum Galeni*.

OLÉO-CÉROLÉ, *Cerate*.

OLÉOLATS, *Olea volatilia*.

OLÉOLÉS, *Olea volatilia*.

OLÉOLS, *Olea fixa*.

O'LEORESIN, *O'leo-resina*. A resin dissolved in volatile oil. The oleo-resins are made by passing ether through the powdered drug by displace-

ment, and evaporating the ether, until the liquid is of an oily consistence. They not only contain the resin but the essential oil of the drug.

OLEORESIN OF CAPSICUM, *Oleoresina Capsici*—o. of Cubebæ, *Extractum Cubebæ fluidum*—o. of Ginger, *Oleoresina Zingiberis*—o. of Lupulin, *Oleoresina Lupulinæ*—o. of Pepper, *Extractum Piperis fluidum*.

OLEORESINA, *Oleoresin*.

OLEORESINA CAPSICI, *Oleoresin of Capsicum*. (Prepared by the action of ether on capsicum powder, by displacement, and evaporating the ether.)

OLEORESINA CUBEÆ, *Extractum Cubebæ fluidum*.

OLEORESINA LUPULINÆ, *Oleoresin of Lupulin*. (Prepared like *Oleoresina Capsici*.)

OLEORESINA PIPERIS, *Extractum Piperis fluidum*.

OLEORESINA ZINGIBERIS, *Oleoresin of Piperoid of Ginger*. (Prepared by the action of ether and alcohol on ginger powder, by displacement, and evaporating the volatile part.) Used in the formation of ginger-drops chiefly.

OLÉOSACCHAROLE, *Elæosaccharum*.

OLEOSACCHARUM, *Elæosaccharum*.

OLEOSUS, *Oleaginous*.

OLERA, (pl. of *olus*, 'a potherb,') see *Lachnon*.

OLETTE, MINERAL WATERS OF. At Olette, in the department Pyrénées Orientales, is a thermal spring, which raises the thermometer to 190° F^{ah}. It is the hottest in France.

OLEULES, *Olea volatilia*.

OLEUM, (from *olea*, 'the olive,') Oil—o. Abietis, see *Pinus picea*—o. Absinthii, *Artemisia absinthium*, (oil of)—o. Acusticum, *Huile acoustique*.

OLEUM ÆTHE'REUM, *Spiritus ætheris oleosus*, *Æthereal oil* (formed in the distillation of ether), *O'leum vini*, *Oil of wine*, *Heart'y oil of wine*, *Sweet oil of vit'riol*, (F.) *Huile douce de vin*. A peculiar oleaginous matter, obtained by continuing the distillation, after the whole of the sulphuric ether has passed over in the process for the preparation of the latter. It is used only as an ingredient in the compound spirit of ether. It is official in the Ph. U. S.

OLEUM AMMONIATUM, *Linimentum ammoniæ fortius*.

OLEUM AMYGDALÆ AMA'RÆ, *Oil of Bit'ter Almonds*. An oil, obtained by distilling with water the kernels or the fruit of *Amygdalus communis*, variety *Amara*. Its action is like that of hydrocyanic acid—four drops having destroyed a dog. Dose, a quarter of a drop to a drop. A drop to a fluidounce of water forms a wash, which has been used in Prurigo senilis, and other forms of itching.

OLEUM AMYGDALA'RUM, *Oleum amygdalæ*, *O. A. dulcis*, (Ph. U. S. 1863.) *Oil of Almonds*, (F.) *Huile d'amandes*. (Expressed from sweet almonds, of *Amygdalus communis*. $\mathfrak{Z}$ xvj of almonds yield $\mathfrak{z}$ v of oil.) It is inodorous, insipid, and of a pale straw colour, and is employed as a demulcent and emollient.

OLEUM ANACARDII, see *Anacardium occidentale*.

OLEUM ANIMALĒ, *Animal oil*, (F.) *Huile animale*. An oil obtained by the decomposition of the immediate principles of animals, subjected to the action of heat. It is fetid, and always contains a certain quantity of carbonate of ammonia. See *Olea empyreumatica*. The name *animal oil* is sometimes also given to the fat contained in the fatty vesicles. The composition of this fat does not, indeed, differ from that of the fixed oils.

OLEUM ANIMALĒ DIPPE'LII, *O. Cor'ni Cer'vi*, *O.*

C. C. rectificatum, *O. pyroanimalē depuratum* seu *animalē æthereum*, *Pyro'leum os'ium rectificatum*, *Animal oil*, *Animal oil of Dip'pel*, (F.) *Huile animale de Dippel* ou de *corne de cerf*, is obtained by distilling animal matters, especially hartshorn, on the naked fire. The carbonate of ammonia, which it contains, renders it partly soluble in water, and communicates to it the stimulant properties for which it is used in medicine. It is employed as an antispasmodic.

OLEUM ANIMALĒ EMPYREUMATICUM CHABERTI, *O. Emphyreumaticum* seu *anthelmin'ticum* seu *contra téniam Chabert'i*, *Emphyreumatic oil of Chab'ert*, *Oil of Chab'ert*, (F.) *Huile anthelmin'tique de Chabert*, is made by adding one part of animal oil to three parts of oil of turpentine, leaving them to combine for four days, and then distilling three parts. An effective anthelmintic. Dose, a teaspoonful three times a day.

OLEUM ANIMALIZATUM PER INFUSIO'NEM, (F.) *Huile animalisée par infusion*, *H. aromatique ou de petits chiens*. A preparation, formerly esteemed tonic and cephalic. It was obtained by boiling new-born puppies in oil, first depriving them of their blood, skin, and intestines. When the decoction was cold, origanum, thyme, pennyroyal, St. John's wort, and marjoram were added.

OLEUM ANISI, see *Pimpinella anisy*—o. Anthelmin'ticum Chaberti, *Oleum animale empyreumaticum Chaberti*—o. Anthemidis, see *Anthemis nobilis*—o. Aurantii, see *Citrus aurantium*—o. Balaninum, *Guilandina moringa* (oleum)—o. Balsami, see *Amyris opobalsamum*.

OLEUM BENZO'INI, *Oil of Ben'zoin* or *Ben'jamin*. An oil obtained by heating, in a sandbath, the matter which remains after benzoic acid has been separated from benzoin by the aid of heat. It has been regarded as balsamic and sudorific.

OLEUM BERGAMII, see *Bergamot*.

OLEUM BU'BULUM, *Neat'sfoot oil*, (F.) *Huile de pieds de bœuf*. It is obtained by boiling in water for a long time the feet or bones of the beef—*Bos domesticus*—previously deprived of the hoof. It is introduced into the official list of the Ph. U. S. as an ingredient of the ointment of nitrate of mercury.

OLEUM CACAO SPISSATUM, *Butter of Cacao*—o. Cadinum, see *Juniperus oxycedrus*—o. Cajuputi, *Caieput* (oil)—o. Camphoræ, see *Camphor*—o. Camphoratum, *Linimentum camphoræ*—o. Carapæ, see *Carapa Guianensis*—o. Cari seu Carui, see *Carum*—o. Caryophylli, see *Eugenia caryophyllata*.

OLEUM CED'RINUM, *Essentia de cedra*, (F.) *Huile de cédrat*. The oil of the peel of citrons, obtained in Italy in a particular manner, without distillation.

OLEUM CHABERTI, *O. animale empyreumaticum Chaberti*—o. Chenopodii, see *Chenopodium anthelmin'ticum*—o. Cicinum, see *Ricinus communis*—o. Cinnamomi, see *Laurus cinnamomum*—o. Contra Téniam Chaberti, *Oleum animale empyreumaticum Chaberti*.

OLEUM COPAI'Æ, *Oil of Copai'ba*, (*Copaib.* $\mathfrak{Z}$ xij; *Aque*, $\mathfrak{Oxvj}$. Distil three gallons; separate the oil; return the water to the copai'ba, and again distil three gallons. Separate the oil, and add it to the other. Ph. U. S.) Dose, gtt. x to xxx.

OLEUM CORNU CERVI, *O. animale Dippelii*—o. Cubebæ, see *Piper cubeba*.

OLEUM ERIGERON'TIS CANADEN'SIS, *Oil of Canada Fleabane*. (By distillation from the herb.)

OLEUM EUPHORBIE LATHYRIDIS, see *Euphorbia lathyris*—o. Fixum nucum cocos butyracæ, see *Cocos butyracæ*—o. Fœniculi, see *Anethum*—o. Gabianum, *Petroleum*—o. Gallinæ, *Alkalæ*—o. Gaultheriæ, see *Gaultheria*—o. Hedeomæ, see *Hedeoma pulegioides*—o. Hyperici, see *Hyperic-*

eum perforatum — o. Infernale, Jatropha curcas
(oil)

OLEUM GRAMINIS INDICI, *Grass oil of Nemaour*.
Gingergrass oil, *Roussa oil*; called, by the perfumers, oil of *geranium*, and oil of *rose geranium*, and occasionally sold as oil of *spikenard*, is obtained from *Andropogon calamus aromaticus* of India. It is used as a perfume, and like the essential oils in general.

OLEUM JECINORIS ASELLI, Oleum jecoris aselli.

OLEUM JECORIS ASELI. *O. Morrhua* (Ph. U. S.) seu *Jecoris Morrhua* seu *Jecoris Aselli*, *Arctia* *Gad'i*, *A. Piscina marina*, *Codliver oil*, *Cod oil*, *F. Huile de morue ou de Foie de morue*. The animal oil, which appears under this name in commerce, is obtained from several of the species belonging to the genus *Gadus*. The *Cod*, *Gadus morrhua* (Ph. B. 1864); the *Dorse* or *Dorch*, *Gadus Callarias*, *Morrhua Americana*; the *Coalfish*, *Gadus carbonarius*; the *Burbot*, *Gadus lota*, *Lota vulgaris*; the *Pollock*, *Gadus pollachius*; the *Ling*, *Lota mullus*; the *Torsk*, *Borpinus vulgaris*. There is reason also to believe, that on the American coast it is obtained, too, from the *Hake*, *Gadus merluccius*, and the *Haddock*, *Gadus aeglefinus*. The clearest sorts are generally used. An excellent cataplectic, in tuberculous and strumous affections, rheumatism, chronic cutaneous diseases, tumours of the mammae, &c. The dose for an adult is from ʒij to ʒiiss.

Codliver olein, prepared by subjecting codliver oil to a low temperature and separating the olein by pressure, has been used in the same cases as the oil.

Skateliner oil, *O'leum ra'ig*, obtained from *Ra'tia*
batis and *R. clava'ta*, is used in the same cases
and doses.

OLEUM JECORIS MORRHUE. *Oleum jecoris aselli*—o. *Juniperi*, see *Juniperis communis*—o. *Juniperi empyreumaticum*, see *Juniperus oxycedrus*—o. de Kervâ, see *Ricinus communis*—o. *Kervinum*, see *Ricinus communis*.

OLEUM LAURINUM, *O. laur'i*, *Daphnolæon*, Oil of baye. An oil obtained from bayberries, and sometimes used in sprains and bruises, unattended with inflammation. See *Unguentum laurinum*.

OLEUM LAVANDULÆ, see Lavandula—o. Lentiscinum, Schinella—o. Limonis, see Citrus medica—o. Lini, see Linum usitatissimum—o. Lini cum calce, Linimentum aquæ calcis.

OLEUM LU'ON PISCEIS, Oil of Pike. From the liver of the *Esox lucius* an oil is spontaneously separated, which is used in some countries to remove specks on the cornea.

OLEUM MACIDIS, see Myristica.

OLEUM MALABATHRI, see Malabathrum.—*o. Maria*, see *Fagara octandra*.—*o. Martis*, see *Ferri chloridum*.—*o. Mastichinum*, see *Pistacia lentiscus*.—*o. Melaleuce leucodendri*, Caigut (oil).—*o. Menthae piperitæ*, see *Mentha piperita*.—*o. Mentha viridis*, see *Mentha viridis*.—*o. Monarda*, *Monarda punctata*.—*o. Morrhuæ*, *O. jecoris aselli*.

OLEUM E MUCILAGINIBUS. *Oil of Mucilages.*
(*Rad. althææ* rec. 1858; *Sem. lini, Sem. frægraci*,
33 ̄ijj; *Aqua, Oij.* Boil for half an hour; add
Oil. olive, Oiv. continue the boiling till the water
is nearly consumed, and pour off the oil.) Emul-
sient.

OLEUM MYRISTICÆ, see *Myristica moschata*—o.
 Myristicæ expressum, see *Myristica*—o. *Nardi*, see
Citrus aurantium—o. *Nucistæ*, see *Myristica*—o.
 Nucistæ æthereum, see *Myristica*.

OLEUM OLIVÆ, O. *Olivæ* rum, Olive or Sal'ad oil. An inodorous, insipid, and transparent oil; obtained by expression from the olive, when ripe. It is demulcent and emollient, — possessing the qualities of the fixed oils in general.

An inferior kind, obtained by boiling olives in

water, and skimming the oil from the surface, is also used in Pharmacy. See *Olea Europæa*.

OLEUM OPHACINUM, Omatribes.—o. Origan, see Origanum.—o. Oxidi cupri viride, Balsam, green, of Metz.—o. de Palma Christi, see Ricinus communis.—o. Palmae, see Cocos butyracea.—o. Palmae liquidum, see Ricinus communis.—o. Petrae, Petroleum.—o. Petrae album, Naphtha.—o. Picinum, Brutia, Pisselæum.—o. Pimentæ, see Myrs pimenta.—o. Pini purissimum, O. terebinthinæ rectificatum.—o. Pulegii, see Mentha pulegium.—o. Pyro-animale, O. animale Disspelli.—o. Ruta, see Oleum jeorris aselli.—o. Ricini, see Ricinus communis.—o. Rosæ, see Rosa centifolia.—o. Rosarum, Rhodolæon.—o. Rosmarini, see Rosmarinus.—o. Ruta, Peganelæon.—o. Ruta, see Ruta.—o. Sabina, see Juniperus Sabina.—o. Sassafras, see Laurus sassafras.—o. Sesami, see Sesamum orientale.—o. Sinapis, see Sinapis.—o. Spicæ, see Lavandula spica.—o. Spicæ vulgaris, O. terebinthinæ.—o. Succini, see Succinum.—o. Succinum rectificatum, see Succinum.

OLEUM SULPHUREATUM, *Bol'samum sulphuris im'plex.* Sulph'uretted oil. (Sulphur. *lot.* 3ij; Ol. olivæ. ℞j. Heat the oil in a large iron pot, and throw in the sulphur by degrees; stirring the mixture after each addition till they unite.) It is stimulating, and was formerly much used in coughs, asthma, &c. and, externally, to foment ulcers.

OLEUM TABACI, Oil of Tobacco. In the form of ointment, (gtt. xx to ℥j of lard,) applied to indolent tumours, ulcers, &c. (An empyreumatic oil, obtained from coarsely-powdered tobacco by heating the retort to dull redness.—Ph. U. S.)

OLEUM TARTARI PER DELIQUITUM, Liqueur potassæ subcarbonatis — o. Templinum, see Pinus mugho.

OLEUM TEREBIN'THINE, *Oil of Tur'pentine*. The volatile oil distilled from the turpentine of *pinus palustris*, and other species of *pinus*.

OLEUM TEREBINTHINÆ RECTIFICATUM, O. pini
purificatum. O. T. *purificatum* seu *terebinthinæ*
altheæ resin. Rectified oil of turpentine, Sp. of tur-
 pentine. Common oil of turpentine is also called
Common oil of Spike, O. leunipæ ex vulgaria, Olei
terebinth. Oij. Aqua, Civ. Distil over the oil. It
 is stimulant, diuretic, anthelmintic, and rubefa-
 cient. Dose, gr. x to ℥i.

Guaiaconian Embrocation for rheumatism consists of *Ol. terebinth.*, *Ol. oliv.* āā f ℥ iss; *Acid. sulph. dil.* f ℥ ij.

OLEUM THEOBROMÆ CACAO EXPRESSUM. Butter of cacao—o. Thymi, see Thymus—o. Tiglini, see Croton tiglium.

OLEUM VALERIA'NÆ. *Oil of Valerian*. The distilled oil of the root of *Valeriana officinalis*. — Ph. U. S. Dose, four or five drops.

OLEUM VINI, O. athereum — o. Vitrioli, Sulphuric acid — o. Vitrioli dulce, Æther sulphuricus.

OLFACTORY. *Olfactory.*
OLFACTION. *Olfacere, [olfacere, olfactum,*
(olere, 'to smell,' and facere, 'to make, 'to smelt;'
Ophre'ois, Ophre'ois, Ophre'ois, Ome'ois, Odu-
rat'io, Odu'rat'io, Sen'sus communi'ricus. Sense of
smell, Smelling. (F.) Olorat. The sense by which
 we perceive the impressions made on the olfactory
 nerves by the odor-us particles suspended in the
 atmosphere. The olfactory nerve or first pair has
 usually been considered the great nerve of smell;
 and it is probably the nerve of special sensibility,
 general sensibility being communicated by the
 branches of the fifth pair, distributed on the
 pituitary membrane of the nose and sinuses.

OLFACTORY, *Olfactory*.
 OLFACTORY, *Olfactive*, *Olfact'ive*, *Olfact'o-
 rius*, *Olfactantius*, *Olfactant'ius*, *Olfact'ivus*,
 (F.) *Olfactif*. That which belongs or relates
 to the sense of smell).

OLFACTORY BULB, see Olfactory nerves.
OLFACTORY FORAM'INA, (F.) *Trous olfactifs*.

are the holes in the cribriform plate of the ethmoid bone, through which the olfactory nerve passes.

OLFACTORY GANGLION. see Olfactory nerves.—o. Lobe, see Olfactory nerves.

OLFACTORY NERVES, *Ethmoidal nerves*, *Par prium Nervo'rum cerebri*, *Nervi odoratorii*, *Proces'sus mamilla'res seu papilla'rum seu mamillares cerebri ad na'res seu papilla'ras*, *Cir'cu'le' manilla'ras*,—the first pair of encephalic nerves. This nerve, which probably arises from the medulla oblongata, is observed to leave the brain, opposite the inner part of the fissure of Sylvius, by three roots; which, by their union, form a triangular knot or expansion. When it reaches the ethmoid fossa it expands and forms a triangular ganglion, or grayish, soft bulb,—*Bul'bus olfacto'rius*, *Olfac'tory bulb*, *gang'lion*, *tuber'cle*, or *lobe*,—which furnishes, from its inferior surface, the branches that have to be distributed to the nasal fossæ. These filaments are numerous; they pass through the foramina in the cribriform plate and enter the nasal fossæ. They are distinguished into the *internal*, *external*, and *middle*. The former are distributed over the mucous membrane, covering the outer paries of the nasal fossæ; the second descend upon the septum, and the third are lost, almost immediately, on the portion of the pituitary membrane that lines the vault of the fossæ.

OLFACTORY TUBERCLE. see Olfactory nerves.

OLFACTUS, gen. *Olfactûs*, Olfaction.—o. Acutus. Hyposphresia.—o. Amissio, Anosmia.—o. Deficiens, Anosmia.—o. Depravatus, Parosmia.

OLIBANUM, (from *λίβανος*, 'the frankincense tree' [?]), see *Juniperus lycia*—o. Arabicum, see *Juniperus lycia*—o. Indicum, see *Juniperus lycia*—o. Sylvestre, see *Pinus abies*—o. Vulgare, see *Pinus abies*.

OLIGEMIA, (*oligo*, and *αἷμα*, 'blood,') *Anæmia*, *Hæmaphoria*, *Hypæmia*.

OLIGIDRIA, (*oligo*, and *ἔρως*, 'sweat,') *Ischidrosis*.

OLIGO, (from *ολιγός*, 'few, little,') in composition has these significations.

OLIGOBLÉN'NIA, (*oligo*, and *βλεννα*, 'mucus,') A deficiency of mucus.

OLIGOCHOL'IA, (*oligo*, and *χολή*, 'bile,') Paucity of bile.

OLIGOCH'YLUS, (*oligo*, and *χυλός*, 'juice, 'chyle,') An epithet for food which is but little nutritive; which furnishes little chyle.

OLIGOCOP'RIA, (*oligo*, and *κόπος*, 'excrement,') Scantiness of alvine evacuations.

OLIGOCYTHÆ'MIA, (*oligo*, *κύτος*, 'a cell, 'and *αἷμα*, 'blood,') A condition of the blood in which there is paucity of red corpuscles.

OLIGODAC'RYA, (*oligo*, and *δακρυ*, 'a tear,') Paucity of the lachrymal secretion.

OLIGOGALACTIA, (*oligo*, and *γαλαξ*, gen. *γαλακτός*, 'milk,') *Agalactia*.

OLIGOGALIA, *Agalactia*.

OLIGOHÆMIA, (*oligo*, and *αἷμα*, 'blood,') *Anæmia*, *Hæmaphoria*, *Hypæmia*.

OLIGOMANIA, (*oligo*, and *mania*,) see *Me-lancholy*.

OLIGOPO'SIA, (*oligo*, and *ποσις*, 'drink,') Diminution in the quantity of drinks.

OLIGOPSYCHIA, (*oligo*, and *ψυχή*, 'mind,') Imbecility.

OLIGOSIA'LIA, (*oligr* and *σάλων*, 'saliva,') Paucity of saliva.

OLIGOSPER'MIA, (*oligo*, and *σπερμα*, 'sperm,') Paucity of spermatie secretion.

OLIGOSPOND'YLUS, (*oligo*, and *σπονδυλος*, 'a vertebra,') A monster with defective vertebrae.

OLIGOTRICH'IA, (*oligo*, and *τριχός*, gen. *τριχός*, 'hair,') Want of hair. Paucity of hair.

OLIGOTROPH'IA, *Par'cior nutrit'io*, (*oligo*, and *τροφή*, 'I nourish,') Deficient nourishment.

OLIGURE'SIA, *Oligure'sis*, *Oligoure'sis*, *Oligoure'sia*, *Oligu'ria*, (*oligo*, and *ουρῶν*, 'I pass urine,') Morbidity diminished urinary secretion.

OLIGURIA, *Oliguresia*.

OLISTHEMA, (*ολισθημα*, from *ολισθαίνω*, 'to slip,') Luxation.

OLIVA, ('the olive,') see *Olea*.

OLIVÆ, Corpora olivaria.

OLIVAIRE, Oliveshaped.

Caute're Olivaire, of the French surgeons, is a cautery whose extremity is terminated by a button having the shape of an olive.

OLIVARIS, Oliveshaped.

OLIVARY, Oliveshaped—o. Bodies, Corpora olivaria.

OLIVARY PROC'ESS, *Proces'sus Oliva'ris*, is a small ridge, running transversely between, and a little behind, the roots of the anterior clinoid processes of the sphenoid bone, and by some considered as the fourth clinoid process.

OLIVE, see *Olea*—o. Tree, *Olea Europæa*.

OL'IVESHAPED, *Oliva'ris*, *Olivifor'mia*, *Ol'ivary*, (F.) *Olivaire*. Resembling an olive. See *Corpora olivaria*.

OLIVES, Corpora olivaria—o. Pickled, Colymbades.

OLIVIER, *Olea Europæa*—o. *de Bohême*, *Ledum palustre*.

OLIVIFORMIS, (*oliva*, and *forma*, 'shape,') Oliveshaped.

OLIVINA, see *Olea Europæa*.

OLLA CAPITIS, (*olla*, 'a pot,') Cranium.

OLMITELLO, MIN'ERAL WATERS OF. A thermal spring in the isle of Ischia. Temperature, 100° Fahrenheit.

OLOPHLYC'TIS, gen. *Olophlyc'tidis*, (*ολοφλυκτις*, from *ὄλος*, 'entire, 'and *φλυκτις*, 'a vesicle,') Herpes.

OLOPHO'NIA, (*ὄλλω*, *ὀλλω*, 'I lose, 'and *φωνή*, 'voice,') Congenital misconstruction of the vocal organs.

OLOPHONIA LABII LOBATA, Harelip.—o. *Lingua frænata*, see *Ankyloglossum*.

OLSNITIUM, *Peucedanum palustre*.

OLUS, (pl. of *olera*, 'a pot herb,') Lachanon.

OLYMPIAN SPRINGS, see Kentucky, mineral waters of.

OMA, (*ωμα*,) in the termination of many pathological words, signifies morbid condition, as in *sarcoma*, *atheroma*, &c.

OMACEPH'ALUS, (F.) *Omacéphale*, (*omo*, and *κεφαλος*, 'devoid of head,') A genus of paracephalous monsters, without any neck—a slight depression indicating the limits of the head and body.

O'MAGRA, *Ho'magra*, (*omo*, and *αγρα*, 'a seizure,') Gout in the shoulder.

OMAL'GIA, (*omo*, and *αλγος*, 'pain,') Pain in the shoulder.

OMARTHRI'TIS, (*omo*, and *αρθρον*, 'a joint, 'and *itis*,) Inflammation of the shoulder-joint.

OMARTHROC'ACĒ, (*omo*, *αρθρον*, 'a joint, 'and *κακός*, 'bad,') Arthroceae of the shoulder-joint. Caries or suppuration of the shoulder-joint.

OMA'SUM, *Oma'sus*, *Psalt'erium*, *Ma'nuplies*, (Sc.) *Mun'plies*, *Mun'plies*, (F.) *Feuillet*, *Mille-feuillet*, *Livret*, *Psautier*. The third stomach of ruminant animals.

OMBILIC, Umbilicus.

OMBILICAL, Umbilical.

OMBILIQUE, Umbilicated.

OMBRAGE, (from *umbra*,) 'a shade,') *Nebula*.

OMENTA, (pl. of *omentum*,) *Meninges*.

OMENTESIS, *Epiploitis*.

OMENTITIS, (*omentum*, and *itis*,) *Epiploitis*.

OMENTULA. (pl. of *omentum*, dim. of *omentum*.) Appendicula pinguedinosa.

OMENTUM, Epiploon—o. Calicium, Epiploon, colic—o. Gastrocolicum, Epiploon, gastrocolic—o. Gastrosplenic, Epiploon, gastrosplenic—o. Great, Epiploon, gastrocolic—o. Hepatogastrium, Epiploon, gastrohepatic—o. Lesser, Epiploon, gastrohepatic—o. Majus, Epiploon, gastrocolic—o. Micus, Epiploon, gastrohepatic—o. Osium, Peristomium—o. Sac of the, see Epiploon, gastrohepatic—o. Third, Epiploon, colic.

OMICHESES, (*ομικτω*, 'to pass urine,') Micturition.

OMICHTMA, (*ομικτμα*.) Urine.

OMITIS, (*ομις*, and *ιτις*.) Inflammation in or about the shoulder-joint.

OMMA, gen. *Ommatias*, (*ομμα*.) Eye. Vision.

OMNIPHAGUS, (*ομνις*, 'all,' and *φαγω*, 'to eat,') Omnivorous.

OMNIVORE, Omnivorous.

OMNIVOROUS, *Omnivorus*, *Omniphagus*, *Pantophagus*, (F.) *Omnivore*, (*ομνις*, 'all,' and *εσθω*, 'I eat,') An epithet for animals which eat every kind of food, animal or vegetable—*Pantophagus*.

OMO, (from *ωμος*, 'shoulder,') in the composition of anatomical terms, means the scapula or shoulderblade; in other cases, the shoulder.

OMOCALVICULAR, see Coracoclavicular.

OMOCOTYLE, see Glene.

OMOHYOIDEUS, (*ομοχυοειδης*, *Hyoidis Quartus Musculus*, (F.) *Scapula-hyoidien*, *Omo-hyoidien*, *Omo-plat-hyoidien*.) This muscle is situate obliquely at the sides and front of the neck. It is slender, long, and flat. It arises from the superior costa of the scapula, near the semilunar notch, and from the ligament that runs across it, and is inserted at the sides of the inferior margin of the body of the os hyoides. It consists of two fasciculi, united by a common tendon, and is a true digastric muscle. It depresses the os hyoides, carrying it a little backwards, and to one side, except when it acts with its fellow, when the bone is depressed, and drawn obliquely backwards.

OMOPHAGIA, (*ομοφαγια*.) *Omositia*, (*ωμος*, 'raw,' and *φαγω*, 'I eat,') Fondness for raw food. Eating of raw food.

OMOPHAGUS, *Omo-phagus* (improperly.) One who eats raw flesh.

OMOPLAT-HYOIDIEN, *Omo-hyoideus*.

OMOPLATE, (*ομο*, and *πλατος*, 'broad, flat,') Scapula—o. *Col de l*, Collum scapulae.

OMOS, (*ωμος*.) Crude, Humerus.

OMOSITIA, (*ωμοσιτια*, from *ωμος*, 'raw,' and *εσθω*, 'food,') *Omo-phagia*.

OMOTARICHOS, (*ομοταριχος*, from *ομο*, and *ταριχος*, 'pickled,') *Salsamentum crudum*; *Salted Tunny Fish*, in particular. Properly, the neck or shoulder-piece of a salted animal. Once much recommended against the bites of vipers, and in hydrophobia.

OMOTES, (*ωμοτες*.) Crudity.

OMOTOCIA, (*ωμοτοκια*, from *ωμος*, 'unripe,' and *τοκος*, 'birth,') Abortion.

OMOTRIBES, (*ωμοτριβες*, from *ωμος*, 'unripe,' and *τριβω*, 'I bruise,') *Omphacium Oleum*. Oil expressed from unripe olives.

OMPHACE, (*ομφακ*, from *ομφαλ*, 'an unripe grape,') Verjuice.

OMPHACION, (*ομφακτιον*, from *ομφαλ*, 'an unripe grape,')—*Succus uvae acer'is*, *Omphacium*. The juice of unripe grapes. Also, Verjuice.

OMPHACIS, (*ομφακς*, from *ομφαλ*, 'unripe,') An unripe gallnut.

OMPHACITES (VINUM), (*ομφακτις*.) A name given to wine prepared from the unripe grape.

OMPHACIUM, (*ομφακτιον*.) *Omphacium*, Verjuice.

OMPHACOMELLI, (*ομφαλ*, 'a 1 unripe grape,')

and *μελι*, 'honey,') A sort of oxymel, made of the juice of unripe grapes and honey.

OMPHALELCO SIS, (*ομφαλος*, and *ελεος*, 'an ulcer,') Ulceration of the navel.

OMPHALEXOCHE, (*ομφαλος*, and *εξωχ*, 'a standing out,') Exomphalus.

OMPHALITIS, (*ομφαλος*, and *ιτις*.) Inflammation of the umbilicus.

OMPHALO, (*ομφαλος*, 'navel,') in composition, navel.

OMPHALOCARPUS, (*ομφαλος*, and *καρπος*, 'fruit,') *Galium sparine*.

OMPHALOCLELE, (*ομφαλος*, and *κλη*, 'rupture,') Exomphalos.

OMPHALOCRA-NIODIDYMUS, (from *ομφαλος*, *κρανιον*, 'the cranium,' and *διδυμος*, 'a twin,') A double monster, in which the umbilical cord or the rudiment of one fetus is rooted in the cranium of another.

OMPHALOMANTIA, *Omphalomancy*, (*ομφαλος*, and *μαντις*, 'prophecy,') A species of divination, practised by credulous matrons, who pretend to be able to know the number of children a female will have, by the number of knots in the navel string of the child.

OMPHALOMESENTERIC, *Omphalomesentericus*, *Omphalomesentericus*, (*ομφαλος*, and *mesenterium*, 'the mesentery,') Appertaining or relating to the umbilicus and mesentery.

OMPHALOMESENTERIC VESSELS, *Vitelomesenterica*, *Vitelline vessels*. Haller gave this name to two very fine vessels, which spread their ramifications on the parietes of the umbilical vesicle. There is an *omphalo-mesenteric artery and vein*. The *omphalo-mesenteric artery* is a branch of the superior mesenteric;—the vein empties itself into the trunk, or into one of the branches of the superior mesenteric. Velpeau affirms, that they inosculate with a branch of the second or third order of those great vessels, with those in particular that are distributed to the cæcum, and he regards them to be the vessels of nutrition of the umbilical vesicle. They are occasionally met with in the fetus, at the full period, under the form of whitish filaments, which extend from the mesenteric vessels to the umbilicus.

OMPHALOMESERICA, *Omphalomesenteric*.

OMPHALONCUS, (*ομφαλος*, and *ογκος*, 'a swelling,') *Omphalephyma*.

OMPHALONEURON, (*ομφαλος*, and *νευρον*, 'a sinew, a ligament,') *Funiculus umbilicalis*.

OMPHALONEURORRHEXIS, (*ομφαλονευρον*, and *ρhexis*, 'rupture,') *Omphalorrhaxis*.

OMPHALOPHLEBITIS, (*ομφαλος*, and *phlebitis*, 'inflammation of a vein,') Inflammation of the umbilical vein.

OMPHALOPHYMA, gen. *Omphalophymatis*, *Omphaloncus*, (*ομφαλος*, and *φυμα*, 'tumour,') A tumefaction of the navel.

OMPHALOPROPTOSIS, (*ομφαλος*, and *προπτωσις*, 'a falling forward,') *Exomphalus*.

OMPHALORRHAGIA, (*ομφαλος*, and *ρραγη*, 'I break out,') Hemorrhage from the umbilicus, in the new-born in particular.

OMPHALORRHENTS, (*ομφαλος*, and *ρρηξις*, 'rupture,') *Omphalorrhæctis*. Rupture of the navel string.

OMPHALUS, (*ομφαλος*.) Umbilicus.

OMPHALOSITE, (*ομφαλος*, and *σιτος*, 'nourishment,') A monster that possesses an imperfect kind of life, which ceases when the umbilical cord is divided.

OMPHALOTOMY, (*ομφαλοτομια*.) *Omphalotomy*, (*ομφαλος*, and *τομη*, 'I cut,') The division of the navel string.

OMPHALUS, (*ομφαλος*.) Umbilicus.

OMUS, (*ωμος*.) Crude, Humerus.

ONAGRA, *Oenothera biennis*.

ONANIA, (after *Onan*.) Masturbation.

ONANISM, Masturbation.

O'NAR, (*ovap*.) Somnium.

ONCOME. A Scotch term, according to Sir Walter Scott, for "mysterious diseases, which baffle the regular physician."

ONCOS, (*ovkos*.) Tumour.

ONCOSES, (*oncos*, and *oais*.) Tumours, as diseases.

ONCOSIS, (*ovkωsis*.) Intumescence.

ONCOT'OMY, *Oncotomia*, *Onkotomy*, (*oncos*, and *tomē*, 'incision'.) The opening of an abscess with a cutting instrument, or the excision of a tumour.

ONCTION, (from *ungere*, *unctum*, 'to anoint'.) Inunction.

ONCUS, (*ovkos*.) Swelling, Tumour.

ONDULATION, (from *undū*, 'a wave'.) Fluctuation.

ONEBERRY, Paris.

ONEIROCRITICUS, *Oneirocriticus*, (*oneiros*, and *kritēs*, 'judgment'.) One who judges according to dreams.

ONEIRODYN'IA, (*oneiros*, and *odynē*, 'pain'.) A painful dream. *Incubus* and *Somnambulism* are oneirodynia.

ONEIRODYNIA ACTIVA, Paroniria, Somnambulism—o. Gravans, Incubus.

ONEIRODYNIE GRAVATIVE, Incubus.

ONEIROG'MUS, (*oviroγμος*.) *Oneirogynē*. A lascivious dream; pollution; nocturnal pollution.

ONEIROGONORRHŒA, Pollution, nocturnal.

POLLUTION. ONEIROGONOS, (*oneiros*, and *gonos*, 'seed'.) Pollution.

ONEIROLOGY, *Oneirologia*, (*oneiros*, and *logos*, 'a description'.) The doctrine of dreams.

ONEIROMANTIA, (*oneiros*, and *μαντεία*, 'divination'.) The art of divining by dreams; or of interpreting dreams.

ONEIRON, (*ovειρον*.) Somnium.

ONEIRON'OSUS, (*oneiros*, and *ovos*, 'a disease'.) *Somnia'tio morbo'sa*. Morbid, uneasy dreaming.

ONEIROPOLEIS, (*ovειροπολεις*.) Somnium.

ONEIROPHILIA, (*ovειροφιλια*.) Somnium.

ONEIROS, (*ovειρος*.) Somnium.

ONERA ALVI, ('load of the bowels'.) Excrement—o. Ventris, ('load of the belly'.) Excrement.

ONGLADE, (from *ungula*, dim. of *unguis*, 'a nail'.) See Onychogryphosis.

ONGLE, (from [L.] *unguis*.) Nail—o. *Incarné*, Onychogryphosis.

ONGLÉE (F.), *Digitorum stupor à gelu*. Painful numbness at the extremities of the fingers, caused by cold.

ONGLET, (after *Ongle*.) Pterygion.

ONGUENT, Unguentum—o. *d'Althéa*, Unguentum de Althæa—o. *Blanc*, Ceratum cetacei—o. *de Blanc de baleine*, Unguentum cetacei—o. *Blanc de Rhazas*, Unguentum plumbi subcarbonatis—o. *de Céruse*, Unguentum plumbi subcarbonatis—o. *Citrin centre la gale*, Unguentum hydrargyri nitratis—o. *Digestif simple*, Unguentum digestivum simplex—o. *Gris*, Unguentum oxidi hydrargyri cinereum—o. *de Guimauve*, Unguentum de althæa—o. *de Laurier*, Unguentum laurinum—o. *Mercuriel*, Unguentum hydrargyri—o. *Napolitain*, Unguentum hydrargyri—o. *de Nitrate de mercure*, Unguentum hydrargyri nitratis—o. *de Poix et de cire*, Basilicum—o. *de Soldat*, Martiatum unguentum—o. *de Sureau*, Unguentum sambuci—o. *de Torumira*, Unguentum plumbi subcarbonatis.

ONION. COMMON, ([L.] *unio*.) *Allium cepa*—o. *Sea*, Scilla.

ONIS'CUS, *Asellus*, *Cu'tio*, *Porcellio*, *Porcellus*, *Por'culus*, (dim. of *ovos*, 'an ass'.) A genus

of insects. *Class*. Crustacea, very common in cellars and dark and moist places.

ONISCI ASEL'LI, *Millep'edes*, *Asell'li*, *Millep'eda*, *Fa'ba*, *Wood'lice*, *Sla'ters*, *Hog'lice*, (F.) *Chopartes ordinaires*, had, at one time, a place in the pharmacopœias. They were considered stimulant and diuretic, and useful in jaundice.

ONITIS, (*ovitis*, gen. *ovitidos*.) *Origanum dictamnus*.

ONKOTOMY, Oncotomy.

ONLOSAT (Arab.), Obolus.

ONOMATOCLESIS, (*ovoma*, gen. *ovomatos*, 'a name', and *κλέω*, 'to call'.) Nomenclature.

ONOMATOLOGIA, (*ovomatoλογία*, from *ovoma*, gen. *ovomatos*, 'a name', and *logos*, 'a discourse'.) Nomenclature.

ONONIS ANTIQUORUM, (*ovonis*.) *O. spinosa*.

ONONIS ARVEN'SIS, (F.) *Bugrane des champs*, has properties like the next.

ONONIS SPINOSA, *Asu'nis*, *Res'ta* seu *Arres'ta bo'vis*, *Ren'ora Ara'tei*, *Ono'nis antiquorum*, *Rest harrouc*, (F.) *Arrête-beauf*, *Bugrande épineuse*, *Bugrane*, *Nat. Ord. Leguminosæ*. *Sec. Syst. Diadelphia Decandria*. The root of this plant was once used as a diuretic.

ONOPORDON ACANTHIUM, (*ovos*, 'the ass', and *πρόω*, 'to break wind'.) *Onopordum acanthium*, *Carduus tomentosus*, *Acum'thium*, *Ac'annus*, *A. spi'na*, *Sp'i'na al'ba*, *Cot'ton Thistle*, (F.) *Chardon aux Aues*. *Ord. Composite*. *Sec. Syst. Syngenesia Polygamia equalis*. The expressed juice has been recommended as a cure for cancer, applied externally.

ONOPORDUM ACANTHIUM, *Onopordon acanthium*.

ONTHUS, (*ovθος*.) Fimus.

ONTOL'OGY, *Ontolog'ia*, (F.) *Ontologie*, (from *ov*, gen. *ovos*, 'a being', and *logos*, 'a discourse'.) A term sometimes used for metaphysics generally; but, more frequently, for that portion of it which deals in generalities; and which embraces, therefore, the speculative and chimerical. It was in this last sense, that Broussais spoke of *medical ontology*, (F.) *Ontologie médicale*.

ONYCHA, see *Bellium*, and *Cistus creticus*.

ONYCHATROPH'IA, *Atroph'ia un'guium*, (*onycho*, and *atrophia*, 'want of nourishment'.) Atrophy of the nails.

ONYCHAUX'Ë, *Hypertroph'ia un'guium*, (from *onycho*, and *αύξη*, 'increase'.) Unusual increase of the size and thickness of the nails.

ONYCHEXALLAX'IS, (*onycho*, and *εξάλλαξις*, 'change'.) A morbid condition of the nails.

ONYCHIA, (from *ovv*, gen. *ovvchos*, 'a nail'.) Paronychia.

ONYCHITIS, (*onycho*, and *itis*.) Inflammation of a nail. See Paronychia.

ONYCHO, (*ovv*, gen. *ovvchos*, 'a nail'.) in coin position, a nail.

ONYCHOCAMPE, (*onycho*, and *καμπή*, 'curvature'.) Onychogryphosis.

ONYCHOCCL'ASIS, (*onycho*, and *κλασις*, 'fracture'.) Fracture of a nail.

ONYCHOCLIN'Ë, *Onychostro'ma*, (*onycho*, and *κλινή*, 'a bed'.) The sensitive bed or stroma under the nail.

ONYCHOCRYPTOSIS, (*onycho*, and *κρυπτος*, 'concealed'.) Onychogryphosis.

ONYCHOGRYPH'OSIS, *Onychogryphosis*, *Onychogrypto'sis*, *Onychocam'pē*, *Gripho'sis*, *Grypho'sis*, (*onycho*, and *gryposis*, 'crookedness'.) Curvature of the nails; such as occurs in hectic individuals. Also, growing in of the nails; *Onychocrypto'sis*, *Incar'atio* seu *Adunca'tio* seu *Arctura un'guium*, *Ony'is*, (F.) *Ongle incarné*, *Onglade*.

ONYCHOMANTIA, (*onycho*, and *μαντεία*, 'divination'.) Divination from the appearance of the nails.

ONYCHON'OSI, *Onychon'si*, (*onycho*, and *oos*, 'a disease.') Disease of the nails.

ONYCOPH THORA, *Onychophthoria*, *Onychophthoria*, (*onycho*, and *phtho*, 'degeneration,') A degenerate condition or destruction of the nails.

ONYCOPHY MA, gen. *Onychophy'matis*, (*onycho*, and *pus*, 'a tumour,') A painful degeneration of the nails, which become thick, rough, and crooked. It occurs as a symptom of syphilis and of lepra.

ONYCHOPTO'SIS, *Ptyntonychia*, (*Dia'resis Un'gula*, *onycho*, and *ptosis*, 'falling,') The falling off of the nails.

ONYCOPHTHORA, *Onychophthoria*.

ONYCHORRHIZA, (*onycho*, and *rhiza*, 'a root,')

See Nail.

ONYCHOSTROMA, (*onycho*, and *stroma*, 'a bed,') Onychocline.

O'NIX, gen. *O'nychie*, (from *οὐξ*, gen. *οὐχος*,) Hook (blunt), Nail, Pterygion. Also, a collection of purulent matter between the laminae of the cornea, having the shape of a nail.

ONYXIS, see Onychogryphosis.

ONYXITIS, (*onyx*, and *itis*,) see Paronychia.

OOCYESIS, (*oos*, and *cyosis*, 'pregnancy,')

Pregnancy, ovarian.

OODEOCELE, see Hernia (foraminis ovalis.)

OODES, (*οὐδης*, from *οον*, 'an egg,' and *oides*,)

Aqueous humour, Oval.

OÜG'ALA, (*oos*, and *gala*, 'milk,') *Oüg'la*, *Og'la*, *Pula* ex *Oc'ia* et *Lacté*. Milk of eggs. A preparation of eggs and milk.

OÜGLA, Oogala.

OÜIDES, (*οὐιδης*, from *oos*, and *uides*, 'resembling,') Aqueous humour, Oval.

OULEMMA PELLUCIDUM, gen. *Oulemm'atis*, (*oos*, and *lemma*, 'a sheath,') Zona pellucida.

OLEUCON, (*oos*, and *leucos*, 'white,') Albumen ovi.

OÜLOG'IA, *Ovolog'ia*, *Ovul'ogy*, (*oos*, and *logos*, 'a discourse,') A description of the ovum.

OON, (*οον*,) Ovum.

OONINE, (from *oos*,) Albumen.

OÜPH'AGOUS, *Oüph'agus*, (*oos*, and *phayw*, 'I eat,') Feeding on eggs.

OÜPHORAÜXE, (*oosphorum*, and *αὐξη*, 'increase,') Oarioparectama.

OÜPHORITIS, (*oosphorum*, and *itis*,) Oaritis.

OÜPHOROPATHIA, (*oosphorum*, and *πάθος*, 'afection,') Oariopathy.

OÜPHÖRUM, (*oos*, and *φέρω*, 'to bear,') Ovary.

OÖRITIS, Oaritis.

OP, as a prefix, see Ob.

OPAC'ITY, *Opac'itas*, *Opakeness*, (*opacus*, 'obscure,') Quality of that which is opaque. The property possessed by some bodies of not allowing the light to traverse them. It is opposed to transparency. *Opacity* of the cornea constitutes *albugo* or *leucoma*; — opacity of the crystalline causes cataract.

OPAKE, *Opac'ens*, *Opac'ous*. An epithet given to bodies, which do not permit the passage of rays of light. The *opaque cornea* is the sclerotica, in contradistinction to the transparent cornea or true cornea.

OPAKENESS, Opacity.

OPALESCENT, Opaline.

OP'ALINE, *Opalescent*. That which resembles the opal. A fluid is said to be opaline when milky, and when it presents an appearance more or less like that of the opal.

OPE, (*οπη*,) Foramen.

OPEOCELE, (*ope*, and *εηλη*, 'rupture,') see Hernia.

OPERATIO, (*opus*, gen. *operis*, 'work,') Action. Operation—o. Chirurgical, see Operation—o. Chymica seu pharmaceuticala, Operation, chymical, &c.—o. Simplex, Haplotomia.

OPERATION, *Operatio*. The application of instruments to the human body with the view of removing disease. The object of an operation, *operatio chirurgica*, is generally to divide or reunite parts, to extract extraneous or noxious bodies, and to replace organs that are wanting, by different instruments or artificial means. The principal operative methods have been called *Syn'thesis*, *Dia'resis*, *Eva'resis*, and *Proth'esis*. Frequently the most difficult subject connected with an operation is to decide when it is absolutely called for or advisable, and when improper.

OPERATION, BILATERAL, see Lithotomy—o. *Cæsarienne*, Cæsarean section.

OPERATION, CHYM'ICAL OR PHARMACEUTICAL, *Operatio chym'ica* seu *pharmaceutica*, is any process whose object is the preparation of medicines; their combinations with each other, analysis, and decomposition;—such are, *solutio*, *distillatio*, *sublimatio*, *evaporatio*, *digestio*, *maceratio*, *infusio*, *decoctio*, *calcinatio*, &c. Some of these operations are mechanical; others, really chymical.

OPERATION, HIGH, see Lithotomy—o. *à Lambeau*, see Flap—o. Lateral, see Lithotomy—o. Sigaultian, Symphysectomy.

OPERATIONS, Capital, see Surgery, operative—o. Major, see Surgery, operative—o. Minor, see Surgery, operative.

OP'ERATOR. A surgeon who is in the habit of practising the greater operations. One who performs any operation. To be a good operator, the surgeon must be well acquainted with anatomy, and be possessed of strong nerve. See Surgeon.

OP'ERCULUM, *Operot'rium*, *Operimen'tum*, (from *operire*, 'to cover,') A cover or lid. In fishes, the gill cover.

OP'ERCULUM ASPERÆ ARTERIÆ, LARYNX—o. *Cutis*, *Epidermis*—o. *Hel*, *Bauhin*, valve of—o. *Laryngis*, *Epiglottis*—o. *Oculi*, Palpebra.

OP'ERCULUM PAPILLÆRUM. A shield for the nipple.

OP'ERIMENTUM, (from *operire*, 'to cover,') Operculum—o. *Intestinerum*, *Epiploon*—o. *Præ-tensum* abdominis, *Peritoneum*.

OP'ERTORIUM, Operculum.

OPHELIA CHIRATA, *Gentiana chirayta*.

OPHIASIS, (*οφιαις*) Calvities, Porrigio decal vans.

OPHIOGLOS'SUM VULGA'TUM, *O. ova'tum* seu *echinogloss'um*, (*ophis*, and *γλῶσσα*, 'tongue,') *Lia'gua* *serpenta'ria*, *Ad'der's* *Tongue*, (*F.*) *Langue* *de* *Serpent*. *Ord.* Filices. *Secz.* Syst. Cryptogamia. This plant was formerly considered to be vulnerary.

OPHIOPH'AGOUS, *Ophioph'agus*, (*ophis* and *phayw*, 'I eat,') An epithet for animals which feed on serpents. Pliny has called certain African tribes by this name.

OPHIORRHIZ'A, *O. mun'gos*, (*ophis*, and *rhiza*, 'root,') *Family*, *Gentianaceæ*. *Secz.* Syst. *Pentandria* *Digynia*. The name of the plant whose root has been called *Ra'dix* *Serpent'um*, *Mun'gos* *Ra'dix*. The bitter root is much esteemed in Java, Sumatra, &c., for preventing the effects that usually follow the bite of the *naja*, a venomous serpent: with which view it is eaten by the natives. It is, also, a reputed anthelmintic.

OPHIOTAPHYLON, (*ophis*, and *σταφύλη*, 'a grape,') *Bryonia* *alba*.

OPHIÖXYLON, *Ophiöxylum*.

OPHIÖXYLUM, (*ophis*, and *ξύλον*, 'wood,') *Ophiöxylum* seu *Ophiöxyton* *serpent'ium* seu *trifolia'tum*, *Acace'ria*, *Lignum* *serpen'tinum*. *Ord.* *Apocynaceæ*. This is recommended in the bites of serpents, and in intermittents. It is said to be very bitter.

OPHIS, (*οφας*,) Serpent.

OPHITES, (*οφιτης*, from *ophis*,) *Serpenti'nus*,

Serpentine or *Black Porphyry*. This rock was formerly worn as an amulet, to cure diseases of the head.

OPHRYS, (ὀφρυς,) Supercilium.

OPHTHALMAGRA, (ophthalmos, and αἶμα, 'a seizure.') A sudden gouty or other pain of the eye.

OPHTHALMAL'GIA, (ophthalmos, and αλγος, 'pain.') Pain in the eye, Neuralgia of the eye.

OPHTHALMEMPASM, gen. *Ophthalmempasmat'is*, (ophthalmos, and ἐμπασσεν, 'to strew upon,') Collyrium siccum.

OPHTHAL'MIA, (from ophthalmos,) *Ophthalmi'tis*, *Oculo'rum inflammatio*, *Caui'ma ophthalmi'tis*, *Ophthalmopo'nia*, *Inflammation of the Eye*, *Ophthalm'ny*, (F.) *Ophthalmie*, *Ophthalmite*, *Inflammation de l'œil*, *Catarrhe oculaire*. Three great varieties of ophthalmia, independently of the *acute* and *chronic* (which conditions occur in all inflammations,) may be reckoned:—the *Ophthalmia membrana'rum*, *O. purulen'ta*, and *O. tar'si*. The first is characterized by the general symptoms of ophthalmia;—pain and redness of the eye or its appendages; with intolerance of light, and unusual flow of tears; the inflammation being seated chiefly in the coats of the eyeball. It is the *Hymenophthal'mia*, *Ophthalmia tarax'is* seu *hu'mida* seu *te'ra* seu *chemo'sis*, *Conjunctivi'tis*, *Inflammatio conjuncti'væ*, *Symphymen'i'tis*, *Syndesmi'tis*, *Catarr'h'al ophthalmia*, *Ophthalmococonjunctivi'tis*, *Ophthalmodesmi'tis*, *Epiph'ora* (Galen), *Conjuncti'val* or *Lach'rymose Ophthalm'ia*, *Tarax'is* (Paulus of Ægina), *Chemo'sis* (Aëtius), (F.) *Conjunctivite*, *Inflammation de la conjonctive*. In the second,—*Ophthalm'ia purulen'ta* seu *puriform'is* *Blennophthal'mia*, *Bleph'aroblenorrhæ'a*, *Ophthalmoblenorrhæ'a*, *Blennorrhæa ocu'li*—the internal surface of the palpebræ associates in the inflammation of the eyeball; and there is a copious secretion of a purulent fluid. An epidemic and contagious [?] variety of this is the *Ægyptian Ophthalmia*, *O. epidem'ica* seu *purulen'ta epidem'ica* seu *contagio'sa* seu *catarrha'lis bel'lica* seu *bel'lica* seu *Asia'tica* seu *purulen'ta contagio'sa*, *Blennorrhæ'a ocu'li* *Ægypti'aca*, *Læmophthal'mia*, *Conjunctivi'tis puromuco'sa contagio'sa* seu *Ægypti'aca*, *Blepharo'tis glandula'ris contagio'sa*, *Ade-*

n'i'tis palpebra'rum contagio'sa, *Sup'purative ophthalmia*, *Epidem'ic contagio'sus ophthalm'ia*—called *Ægyptian*, from its prevalence in Egypt during the British Expedition under Sir Ralph Abercrombie. The inflammation is rapid and destructive; granulations shoot from the tunica conjunctiva, and occasionally there is intolerable pain, often succeeded by delirium. In newborn children, a species of purulent ophthalmia, *O. purulen'ta infan'tum* seu *neonata'rum*, *Blepharophthal'mia* seu *Lippitu'do* seu *Blennorrhæ'a ocu'li* seu *Ophthalm'oblenorrhæ'a* seu *Blepharoblenorrhæ'a* seu *Blepharopyorrhæ'a* seu *Pyophthal'mia*, *Psorophthal'mia neonato'rum*, *Adenoseychiton'i'tis*, *Adenologadi'tis*, (F.) *Ophthalmie puriforme des nouveau-nés*, in which the palpebræ are florid and peculiarly tumid, is by no means uncommon. It seems to be caused by acrid discharges from the mother, applied to the eye of the infant during its exit; or to the stimulus of the light, when the child first opens its eyes. A severe form of purulent ophthalmia—*Hyperconjunctivite* (Piorry)—is produced by the application of gonorrhœal matter to the eye. It is the *Ophthalmia gonorrhœ'ica*, *Blennorrhæ'a ocu'li gonorrhœ'ica*, *Gonorrhoblepharhæ'a*, *Conjunctivi'tis blennorrhag'ica* seu *gonorrhœ'ica*, *Gonorrhæ'al Ophthalm'ia*, (F.) *Ophthalmie blennorrhagique*, *Conjunctivite blennorrhagique*.

Ophthalmia is likewise modified by the condition of the constitution; and hence we have *strumous*, *variolous*, and other inflammations of the conjunctiva.

The third variety—the *Ophthalmia Tar'si* seu *glutino'sa*, *Blepharophthal'mia*, *Blepharo'tis*, *Blepharoti'tis*, *Blephari'tis*, *Blepharadeni'tis*, *Adenophthal'mia*, is seated chiefly in the tarsus; the sebaceous crypts secreting a viscid and acrid fluid, that glues and ulcerates its edges, and irritates the eye.

When accompanied with ulceration and much itching, it constitutes *Psorophthal'mia*, *Ocu'li palpebra'rum Scab'ies pruriginosa*, *Bleph'arophthal'mia ulcero'sa*, *Ophthalm'ia psor'ica*, *Tin'ea tarsi* seu *cilia'ris*, *Tinea of the Eyelids*.

The different forms of inflammation of the conjunctiva are thus classed by M. Desmarres:—

Conjunctivitis.....	{	1. Pure or phlegmonous.	{	{	{			
		2. Pustular.				Contagious.-	Erysipelatous.	
		3. Granular or catarrhal.				Noncontagious.		Variolous.
						Miasmatic.		Morbillous.
Exanthematic.....	Scarlatinous.							
		{	{	{				
					4. Purulent.....	Of new-born infants.	Erysipelatous.	
Gonorrhœal.	Variolous.							
Egyptian.	Morbillous.							

Conjunctivitis, when slight, requires little treatment: the antiphlogistic regimen—and particularly avoiding exposure to light—being all that is necessary. When more severe, along with this, general and local bloodletting, especially by leeches to the side of the septum narium, must be employed, so as to make a decided impression on the system; with nauseating doses of antimony, purgatives, blisters (at some distance from the organ), refrigerants, and astringents, cold or warm, according to circumstances, to the eye. In *Purulent Ophthalm'ny* these measures must be still more actively employed, otherwise disorganization will speedily follow;—the granulations which form on the adnata of the eyelids may be removed with the scissors, and the sore be touched with a solution of alum. *Ophthalm'ia Tar'si* must be treated on general principles, when severe. It usually, however, requires

merely the antiphlogistic regimen, with the occasional use, especially at night, of a slightly stimulating ointment, such as the unguentum hydrargyri nitricooxydi, considerably reduced with lard. The ointment prevents the eyelids from being glued together during the night, and thus obviates the irritation caused by attempts at separating them in the morning.

OPHTHALMIA, ÆGYPTIAN, see Ophthalmia—o. Asiatica, see Ophthalmia—o. Bellica, see Ophthalmia—o. Biliosa, Ophthalmocholosis—o. Catarrhal, see Ophthalmia—o. Catarrhalis bellica, see Ophthalmia—o. Chronica, Lippitudo—o. Conjunctival, see Ophthalmia—o. Contagiosa, see Ophthalmia—o. Epidemica, see Ophthalmia—o. Glutinoso, see Ophthalmia—o. Gonorrhœal, see Ophthalmia—o. Gonorrhœica, see Ophthalmia—o. Granular, Trachoma—o. Humida, see Ophthalmia—o. Lachrymose, see Ophthalmia—o. Membrana-

rum. see Ophthalmia.—o. Neonatorum, see Ophthalmia.—o. Phlyctenular, see Ophthalmia.—o. Psorica, see Ophthalmia tarsi.—o. Puriformis, see Ophthalmia.—o. Purulenta, see Ophthalmia.—o. Purulenta contagiosa, see Ophthalmia.—o. Purulenta epidemica, see Ophthalmia.—o. Purulenta infantum, see Ophthalmia.—o. Rheumatic, Sclerodermatitis.—o. Sicca, Xerophthalmia.—o. Suppurative, see Ophthalmia.—o. Taraxis, see Ophthalmia.—o. Tarsi, see Ophthalmia.—o. Ulcerosa, Helicophthalmia.—o. Varicose, Cirsophthalmia.—o. Vera, see Ophthalmia.

OPHTHALMIATER, (*ophthalmos*, and *iater*, 'a physician.') Oculist.

OPHTHALMIATRICA, *Ophthalmiatrotechnica*, *Ophthalmotherapia*, (F.) *Oculistique*. The art of the oculist. Treatment of diseases of the eye.

OPHTHALMIATROTECHNICA, (*ophthalmiater*, and *τεχνη*, 'art.') Ophthalmiatry.

OPHTHALMIC, (*ophthalmicus*, (F.) *Ophthalmique*, (from *ophthalmos*.) That which relates or belongs to the eye.

OPHTHALMIC ARTERY, *A. orbitaire*, (Ch.) arises from the internal carotid, and issues from the cranium by the foramen opticum. At first it is situate at the outer and lower side of the optic nerve, but ascends above this nerve and passes towards the inner angle of the orbit. It furnishes in its course the *lacrimal*, *centralis retinae*, *infraorbitalis*, *ciliaries*, *anterior ciliaries*, *inferior and superior muscular*, *anterior and posterior ethmoidals*, and *superior and inferior palpebrals*. After having given off these, it divides into two branches—the *frontal* and *nasal*, which furnish a number of ramifications—some superficial, others deep-seated.

OPHTHALMIC GANGLION, *Lenticular or Ciliary ganglion*, *Ganglion ophthalmicum*, seu *semilunare seu ciliare*, *G. orbitaire*, (Ch.) is one of the smallest ganglions of the body, and formed by the 5th and 3d pairs of encephalic nerves. It is situate at the outer side of the optic nerve near the bottom of the orbit; is irregularly quadrilateral and flat; its colour of a reddish gray. Behind, it communicates by its posterior angles—by means of nervous filaments—with the nasal nerve of the ophthalmic, the motor oculi, and offsets from the sympathetic. Each of its anterior angles furnishes a fasciculus of small nerves. These are the ciliary nerves.

OPHTHALMIC NERVE, *Ophthalmic Nerve of Willis*, *Orbito-frontal*, (Ch.) is the smallest and first of the three branches given off by the 5th pair. It proceeds along the external paries of the cavernous sinus, and enters the orbit by the sphenoidal fissure. It divides into three branches, one, *external*—the *lacrimal nerve*; another, *superior*—*frontal nerve*; and the last, *internal*—the *nasal nerve*. See *Lacrimal*, *Frontal*, *Nasal*.

OPHTHALMIC or OPTIC REMEDY, *Ophthalmicum* seu *Opticum*, (F.) *Topique ophthalmique*, is a medicine employed locally in ophthalmia.

OPHTHALMIC SINUS, see *Cavernous Sinus*.

OPHTHALMIC VEIN has the same arrangement as the artery, which it accompanies in all its divisions. It issues from the orbit at the inner part of the sphenoidal fissure, and discharges its blood into the cavernous sinus.

The name *Facial Ophthalmic* has been given to the branch by which the facial vein communicates with the ophthalmic.

OPHTHALMICA, *Euphrasia officinalis*.

OPHTHALMICI EXTERNI, *Motores oculorum*.

OPHTHALMICUM, *Ophthalmic remedy*.

OPHTHALMIDIUM, (dim. of *ophthalmos*.) Microphthalmus.

OPHTHALMIE BLENNORRHAGIQUE, see

Ophthalmia.—o. *Puriforme des Nouveau-nés*, see Ophthalmia.

OPHTHALMITE, Ophthalmia.

OPHTHALMITIS, (*ophthalmos*, and *itis*.) Ophthalmia. Occasionally restricted to inflammation of the globe of the eye, in which both the external and internal structures are involved.

OPHTHALMIUM, (dim. of *ophthalmos*.) Microphthalmus.

OPHTHALMO, (from *ὀφθαλμος*, 'eye.') In composition, eye.

OPHTHALMOBLENNORRHEA, Ophthalmia (purulent)—o. Neonatorum, see Ophthalmia (purulent) infantum.

OPHTHALMOCARCINOMA, pen. *Ophthalmocarcinoma*, cancer of the eye.

OPHTHALMOCELE, (*ophthalmos*, and *κηλη*, 'tumor.') Exophthalmia.

OPHTHALMOCHOLOSIS, (*ophthalmos*, and *χολη*, 'bile.') *Ophthalmia biliosa*. Ophthalmia from biliary derangement. [?]

OPHTHALMOCHROITES, (*ophthalmos*, and *χρῶσις*, 'to colour.') See *Choroida tunica*.

OPHTHALMOCONJUNCTIVITIS, see Ophthalmia.

OPHTHALMOCOPIA, (*ophthalmos*, and *κοπος*, 'fatigue.') Copyopia.

OPHTHALMODESMITIS, (*ophthalmodesmum*, and *itis*.) See Ophthalmia.

OPHTHALMODESMUM, (*ophthalmos*, and *δεσμος*, 'a band.') *Conjunctiva tunica*.

OPHTHALMODULIA, (*ophthalmos*, and *ὄσλη*, 'servitude.') The title of a book by Bartisch on diseases of the eye.

OPHTHALMODYNIA, (*ophthalmos*, and *δύνη*, 'pain.') Pain, and especially rheumatic pain of the eye. Plenck has given this name to neuralgia of the frontal nerve, in which the pain radiates particularly towards the bottom of the orbit. See *Neuralgia*, *frontal*.

OPHTHALMODYNIA PERIODICA, *Neuralgia*, *facial*.

OPHTHALMŒDEMA, gen. *Ophthalmœdematis*, (*ophthalmos*, and *οἴδημα*, 'œdema.') Œdema of the conjunctiva.

OPHTHALMOGRAPHY, (*ophthalmos*, and *γραφη*, 'a description.') The part of anatomy which gives a description of the eye. An anatomical description of the eye.

OPHTHALMOLOGY, *Ophthalmologia*, (*ophthalmos*, and *λογος*, 'a discourse.') The part of anatomy which treats of the eye. An anatomical treatise on the eye. A description of the eye in health and disease.

OPHTHALMOMELANOMA, *Ophthalmomelanosis*, (*ophthalmos*, and *melanoma*.) Melanoma of the eyeball.

OPHTHALMOMETER, (*ophthalmos*, and *μετρον*, 'measure.') An instrument of the nature of compasses, invented by F. Petit, for measuring the capacity of the anterior and posterior chambers of the eye in anatomical experiments.

OPHTHALMOMYITIS, *Ophthalmomyositis*, (*ophthalmos*, *μῦς*, 'a muscle,' and *itis*.) Inflammation of the muscles of the eye.

OPHTHALMOPARALYSIS, *Ophthalmoplegia*.

OPHTHALMOPHLEBOTOMIA, (*ophthalmos*, *φλεψ*, gen. *φλεβος*, 'a vein,' and *τομή*, 'incision.') Bleeding from the vessels of the conjunctiva.

OPHTHALMOPHTHARISIS, *Ophthalmophtharisis*, (*ophthalmos*, and *φθορα*, 'corruption.') Destruction of the eyeball.

OPHTHALMOPHTHISIS, *Ophthalmophtharsis*.

OPHTHALMOPLEGIA, *Ophthalmoplegia*, (*ophthalmos*, and *πληγω*, 'I strike.') Paralysis of one or more of the muscles of the eye.

OPHTHALMOPONIA, (*ophthalmo*, and *πονος*, 'suffering,') Ophthalmia.

OPHTHALMOPTOMA, Ophthalmoptosis.

OPHTHALMOPTOSIS, (*ophthalmopto*'ma, (*ophthalmo*, and *πτωσις*, 'a prolapse,') A word employed by some authors in the sense of exophthalmia, by others in that of hydropthalmia.

OPHTHALMORRHAG'IA, (*ophthalmo*, and *ῥαγναι*, 'I break forth,') Hemorrhage from the tunica conjunctiva. It is rare.

OPHTHALMOS, (*οφθαλμος*,) Eye.

OPHTHALMOSCOPE, see Ophthalmoscopia.

OPHTHALMOSCOPIA, (*ophthalmoscopy*, (*ophthalmo*, and *σκοπεω*, 'I regard attentively,') The art of judging of the temper, &c., of a person by examining his eyes. The art of judging of health or disease by inspection of the eyes. Exploration of the eyes in order to a diagnosis.

Ophthalmoscope, *ophthalmoscopium*, *ophthalmoscopium*, *Speculum oculi*. An instrument which, by reflecting the light on the retina, enables the condition of the interior of the eye to be appreciated. To a form of the ophthalmoscope the name *Anagnostakis* has been given, after a young Greek physician, who, it was thought, invented it.

OPHTHALMOSTA'TUM, (*Ophthalmostat*, (*ophthalmo*, and *στασις*, 'station,') *Speculum oculi*. An instrument for fixing the eye. The action of fixing the eye is termed *Ophthalmos'tasis*.

OPHTHALMOTHERAPIA, (*ophthalmo*, and *θεραπευα*, 'treatment,') Ophthalmiatria.

OPHTHALMOT'OMY, (*Ophthalmotomia*, (*ophthalmo*, and *τεμνω*, 'I cut,') The part of anatomy which treats of the dissection of the eye. It has also been applied to extirpation of the eye.

OPHTHALMOXEROSIS, (*ophthalmo*, and *ξηρωσις*, 'dryness,') Xerophthalmia.

OPHTHALMOX'YSIS, (*ophthalmos*, and *ζωω*, 'I scrape,') A name given to the scarification sometimes practised on the conjunctiva in cases of ophthalmia.

OPHTHALMOXYS'TRUM, (*Ophthalmoxyster*, *Ophthalmoxystrium*, *Xystrium ophthalmicum*. An instrument for scraping the eye. Name given to a small brush, with barbs like an ear of barley or rye, intended to scarify the eyelids in certain cases of ophthalmia.

OPHTHALMOZO'A, (F.) *Ophthalmozaires*, (*ophthalmo*, and *ζωον*, 'an animal,') Entozoa or parasites in the globe of the eye and its appendages.

OPHTHALMUS, (*οφθαλμος*,) Eye.

OPHTHALMYMENITIS, (*Ophthalmohymenitis*, (*ophthalmo*, *ὑμην*, 'membrane, and *ιτις*,') Inflammation of one or more of the membranes of the eye.

OPIACÉ, Opiate.

OPIACUM, Opiatum.

OPIANE, Narcotine.

O'PIANINE, (*Opiantina*, *Opi'ania*, (from *opium*,) An alkaloid discovered in opium, which resembles morphia in its properties and doses.

OPIANUM, Narcotine.

OPIAT, Opiate—o. *Dentifrice*, Dentifrice.

O'PIATE, (*Opiatum*, *Opiacum*, *Papavera'ceum*, *Meco'nicum*, (F.) *Opiat*, *Opiacé*, (from *οπιον*, (*opos*, 'juice,') 'opium,') A medicine containing opium. A medicine that procures sleep. An electuary;—formerly, an electuary which contained opium.

O'PIATED, (*Opiatus*, (F.) *Opiaté*, *Laudanisé*. Impregnated with opium or laudanum. Affected by opium or laudanum.

OPIATUM, Opiate.

OPIA'TUM FEBRIS'CEUM, (*Electuarium de K'ina K'ina*; *Electuary of Cincho'na*. (*Pulv. cinch. 3viii*, *Ammon. muriat. 3j*, *Mellis*, *Syrup. absinth. aa 3ij*. Make into an electuary. *Ph. P.*) Given when cinchona is indicated.

OPIATUM MESENTER'ICUM, (*Electuarium de Al'oë*,

Muria'te Hydrar'gyri et Fer'ro. (*Gum. ammon. 3ss*, *Sennæ 3vj*, *Hydrargyri submuriat.*, *Rad. Ari.*, *Aloës socotrin. aa 3ij*, *Pulv. scammon. comp.* (vulg. *de tribus*), *Rhei rad. aa 3ij*, *Ferri limatur. porphyrisat. 3ss*. Bruise and mix together, add of Compound syrup of apples double the weight of the other matters, and make into an electuary. *Ph. P.*) Dose, 3ss to 3ij, in obstructions of the liver, mesentery, &c.

OPIATUS, Opiated.

OPION, (*οπιον*, dim. of *opos*, 'juice,') Opium

O'PIOPHIL, (*Opiophilos*, *Opioph'ilus*; (*οπιον*, *opium*, and *φιλεω*, 'I love,') A lover of opium. There was in Paris, in 1853, a club of opium smokers, who called themselves *O'piophils*.

OPISMA, (*οπισμα*,) Succus.

OPISTHENAR, (*οπισθεναρ*,) *Dorsum manus*, (*opistho*, and *θεναρ*, 'the flat of the hand,') The back of the hand.

OPISTHO, (*οπισθε*, 'behind, backwards,') in composition the same.

OPISTHOCEPHALON, (*opistho*, and *κεφαλη*, 'the head,') Occiput.

OPISTHOCRANIUM, (*opistho*, and *κρανιον*, 'the skull,') Occiput.

OPISTHOCYPHOSIS, (*οπισθοκυφωσις*, from *opistho*, and *cyphosis*,) Cyphoma.

OPISTHOLOB'IUM, (*Opiethotium*, (*opistho*, and *λοβιον*, 'the lobe of the ear,') Any agent applied behind the ear.

OPISTHOTUM, (*opistho*, and *ους*, gen. *ωτος*, 'the ear,') Opisthobolium.

OPISTHOT'ONUS, (*οπισθοτονος*,) *Raptus posterga'neus*, *Tet'anus dorsa'lis seu post'i'eus seu posterga'neus*, (*opistho*, and *τενω*, 'I stretch,') A species of tetanus, in which the body is bent backwards.

OPIUM, (*οπιον*, dim. of *opos*, 'juice,') see Papaver—o. Colatum, Extractum opii—o. Depuratum, Extractum opii.

OPIUM EATERS, Theriaki.

OPIUM, LETTUCE, see Lactuca.

OPOBALSAMUM, (*οπος*, 'juice, and *βαλσαμον*, 'a balsam,') see Amyris opobalsamum.

OPOCARPASON, Carpasium (juice.)

OPOCEPH'ALUS, (*ωψ*, 'the face, and *κεφαλη*, 'head,') (F.) *Opocephale*. A genus of monsters having the two ears approximated or united under the head; the jaws atrophied; and no mouth or nose.

OPODELDOCH, Opodeldoch.

OPODEL'DOCH, (*Opodel'toch*, *Opodel'doc*. An unmeaning term, frequently used by Paracelsus. Formerly, it signified a plaster for all external injuries, and was occasionally used, generically, like *Saponula*, for the union of soap with one or more volatile oils: now, it is applied to the Linimentum Saponis Compositum.

OPODELDOCH, STEER'S. A liniment, called after the inventor. There are many formulæ for its preparation. The following is one. *Sop. alb. ʒij*, *Camphor. ʒij*, *Ol. rosmarini ʒiv*, *Spiritus vini rectificati Oij*. See Linimentum Saponis Compositum.

OPODECELE, (*οπη*, 'a foramen,') *εἶδος*, 'resemblance, and *κηλη*, 'a rupture,') see Hernia.

OPODIDYMUS or OPODYMIUS, (*ωψ*, 'the face, and *διδυμος*, 'double,') Diprosopus.

OPOIDIA GALBANIFERA, (from *opos*, 'juice,')

οπος, 'juicy,') see Bubon Galbanum.

OPOLÉ, (*οπος*, 'juice,') Succus.

OPOLIDE, Succus.

OPOPANACUM, Pastinaca opopanax.

OPOPANAX, (*οποπαναξ*; from *opos*, 'juice, and

panax,) see Pastinaca opopanax—o. Chironium, Pastinaca opopanax.

OPOPANAXWORT, Pastanaca opopanax.

OPOPIOS, (*οπωπιος*,) Optic.

OPOPONAX, Pastinaca opopanax.

OPORA, (*οπορα*,) see Autumn.

OPORICE, (ὄπωρα, 'autumn.') A medicine, composed of several autumnal fruits, particularly quinces, pomegranates, &c., and wine. It was formerly administered in dysentery, diseases of the stomach, &c.

OPPOS. (ὀπός,) Succus expressus—o. Silphion, *Laserpitium*.

OPPIDULUM, ('a little town.') Vulva.

OPPILATIO, (*oppillare, oppilatum*, 'to shut up.') *Emphraxis*—o. *Tela Cellulosa*, Induration of the cellular tissue.

OPPILATIVUS, *Obstruens*.

OPPO'NENS, (F.) *Opponant, (op, and ponere, 'to place.')* That which faces or is put in opposition to something. The name has been given to two muscles of the hand.

OPPO'NENS MINIMI DIG'ITI, (F.) *Opposant du petit doigt, Carpo-métacarpien du petit doigt* (Ch.), *Métacarpien du petit doigt*. This muscle has the same shape and arrangement as the preceding, but is of less size. It is situate in the hypothenar eminence. Its fibres are inserted into the anterior annular ligament of the carpus, and terminate on the whole length of the inner edge of the 5th metacarpal bone. This muscle carries the 5th metacarpal bone forwards and outwards, and thus augments the concavity of the palm of the hand.

OPPO'NENS POL'ICIS, *Flexor ossis metacarpi pollicis, F. primi internodii* (Douglas), *Opponens pollicis manus, Antith'enor sive semi-interosseus pollicis*, (F.) *Carpo-métacarpien du pouce* (Ch.), *Métacarpien du pouce, Opposant du Pouce*. A small, flat, triangular muscle, situate in the substance of the thenar eminence. It is attached, on the one hand, to the anterior annular ligament of the carpus and to the trapezium; and on the other, to the whole of the outer margin of the first metacarpal bone. This muscle impresses on the first bone of the metacarpus a movement of rotation, which opposes the thumb to the other fingers.

OPPOSANT, *Opponens*—o. *du Petit doigt, Opponens minimi digiti*—o. *du Pouce, Opponens pollicis*.

OPPRESSIO, (from *opprimere*, (*op*, and *premere, pressum*.) 'to press against.') *Catalepsy, Oppression*—o. *Nocturna, Incubus*.

OPPRESSION, *Oppress'io, Thlip'sis, Cata-thlips'is*. A state, in which the patient experiences a sensation of weight in the part affected. When employed abstractly, it means, particularly, *Oppression of the chest, Oppress'io, Pect'oria*, (F.) *O. de Poitrine*.

OPPRESSION, *Oppress'io vi'rium*, (F.) *Oppression des forces*, is, also, used for that condition, at the commencement of fevers, &c., in which the system is opposed rather than debilitated, and where the vascular action arises, as the obstruction to free circulation is relieved by bleeding, purging, &c.

OPS, (ὄψ,) Eye.

OPSEONUSI, *Opsionusi*.

OPSIALGIA, (ὄψις, 'the countenance,' and *algos*, 'pain,') *Neuralgia, facial*.

OPSIGONUS (*DENS*), (ὀψιγόνος, from *ὄψις*, 'late,' and *γενω*, 'to be born,') see *Dentition*.

OPSIODOUS, (ὄψις, 'late,' and *ὄδους*, 'a tooth,') see *Dentition*.

OPSIONUSI, *Opseonusi*, (ὄψις, 'vision,' and *νοσος*, 'a disease,') *Mor'bi vis'us*. Diseases of vision.

OPSIOTOC'IA, (ὄψις, 'late,' and *τοκος*, 'birth,') *Partus serotinus*. Parturition after the usual period. See *Pregnancy*.

OPSI, (ὄψις, 'countenance,') Pupil, Vision.

OPSOMA'NIAC, *Opsom'anes*, (F.) *Opsomane*, (ὄψωρ, 'aliment,' and *pavia*,) One who loves some particular aliment to madness.

OPTESIS, (ὀπτησις, 'roasting or broiling,') *Asatio*.

OPTIC, (ὀπτικός, *Opticus, Opo'pico*, (F.) *Optique*, (from *ὀπταί*, 'I see,') That which relates to vision.

OPTIC CEN'TRE, *Centrum opticum*. The optic centre of the crystalline is the point at which the various rays proceeding from an object cross in their way to the retina.

OPTIC COMMISSURE. See *Chiasm*.

OPTIC FORA'MEN, *Fora'men opticum*, (F.) *Trou optique*. A round opening in the sphenoid bone, near the base of its lesser ala, through which the optic nerve passes.

OPTIC GANGLIA, *Quadrigenina tubercula*.

OPTIC GROOVE. A transverse groove on the superior surface of the sphenoid bone, on which the commissure of the optic nerve rests, and which is continuous on each side with the optic foramen.

OPTIC LOBES, *Quadrigenina tubercula*.

OPTIC NERVE, *Nervus opticus seu vis'us seu vis'orius, Par secundum seu opticum seu vis'orium, Nerve of the 2d pair, Ocular nerve of Chaussier*. The optic nerves are remarkable for their size; for their running a longer course within than without, the cranium; and for their furnishing no branch from their origin to their termination. They do not seem to arise, as was long supposed, from the optic thalami, but from the tubercula quadrigenina. Immediately after their origin, the optic tracts, *Tractus optici*, proceed forwards; are, at first, broad and flat; but afterwards become narrower and round. In front of the fossa pituitaria, they unite and decussate, each nerve proceeding through the optic foramen of the sphenoid bone with the ophthalmic artery, and through the *Fora'men opticum choroideæ*, to form the retina. The nerve passes to the back part of the globe of the eye, becomes narrower, and enters that organ at a flattened circular papilla—*Collie'ulus seu Papilla nervi optici*—to give rise to the nervous expansion called the retina. Between the colliculus and the yellow spot of Sömmerring, a small projection or fold of the retina passes—*plica central'is ret'inae*. Besides its neurilemma, the optic nerve is surrounded by a sheath, furnished by the dura mater. This accompanies it as far as the eye.

The optic nerve is the nerve of special sensibility of the eye. In the centre of the nerve, there is a *punctum cal'cum, macula nigra*, where the central artery enters the eye, and the corresponding vein passes out. If a ray of light falls on this spot it is not appreciated.

OPTIC REMEDY, see *Ophthalmic remedy*—o. *Thalami, Thalami nervorum opticorum*—o. *Traets*, see *Optic nerve*.

OPTICO-TROCHLÉI-SCLÉROTICIEN, *Oblique, superior, of the eye*.

OPTICUM, *Ophthalmic remedy*.

OPTICUS, (ὀπτικός,) *Optic*.

OPTILOS, (ὀπτικός,) *Eye*.

OPTIQUE, *Optic*.

OPTOMETER, (ὀπτοματ, 'I see,' and *μετρον*, 'a measure,') An instrument for measuring the distance of distinct vision.

OPUNTIA, (from *ὄπους*, *Opus*, gen. *Opuntis*, 'a town of Locris,') *Cactus opuntia*—o. *Vulgaris, Cactus opuntia*.

OR. (from *aurum*,) *Gold*—o. *Blanc, Platinum*—o. *Cyanure d'*, see *Gold*—o. *Divisé*, see *Gold*—o. *Hydrochlorate d'*, see *Gold*—o. *Iodure d'*, see *Gold*—o. *Métallique*, see *Gold*—o. *Nitromuriate d'*, see *Gold*—o. *Oxide d'*, see *Gold*—o. *et de Sodium chlorure d'*, see *Gold*—o. *et de Soude, Hydrochlorate d'*, see *Gold*—o. *et de Soude, Muriate d'*, see *Gold*.

ORA SERRATA, ('serrated edge,') see *Clivary (Body)*.

ORACHE, *SPREADING, Atriplex patula*—o. *Stinking, Chenopodium vulvaria*.

O'RAL, *Vocal*, (F.) *Oral*, (from *os*, gen. *oris*, 'a mouth.') Relating to the mouth or to speech.

ORANGE, (from modern Lat. *aurantium*, itself from *aurum*, 'gold'—the colour of the fruit,) *Citrus aurantium*—o. Bitter, see *Citrus aurantium*—o. Flowers of the, see *Citrus aurantium*.

ORANGE GRASS, *Hypericum sarothra*—o. Root, *Hydrastis Canadensis*—o. Tree, *Citrus aurantium*.

OR'ANGEADE, (from orange.) A drink, made with orange juice diluted with water. It is antiphlogistic, and often recommended in acute diseases.

ORANGES, see *Citrus aurantium*—o. *Curassoa*, *Aurantia curassavatica*.

ORBES CARTILAGINOSI TRACHEÆ, (*orbis*, 'circle,') see *Trachea*.

ORBICULAIRE DES LÈVRES, *Orbicularis oris*—o. des *Paupières*, *Orbicularis palpebrarum*.

ORBICULAR, *Orbicularis*, (F.) *Orbiculaire*, (from *orbis*, 'a circle,') Spherical, Circular.

ORBICULAR BONE, *Os orbicularē seu lenticularē seu Syll'vii seu orbicularē Syll'vii*, *Globulus stap'edis os'sis*, *Epiph'ysis crur'is longioris incud'is*, *Ossiculum orbicularē seu squamosum seu cochlearē seu quar'tum*, is the smallest of the four bones of the ear. It is scarcely perceptible, round, convex on two surfaces, and situate between the long ramus of the incus and the head of the stapes.

ORBICULAR MUSCLES are muscles with circular fibres surrounding some natural opening of the body. 1. *Orbicularis* seu *Constrictor O'ris*, *Sphincter Labiorum*, *Sem'i seu Demi-orbicularis*, *Subpræsemi-orbicularis*, *Labialis*, *Osculato'rius*, *Basia'tor*, (F.) *Labial*, *Orbiculaire des lèvres*. A muscle situate in the substance of the lips, and extending from one commissure to the other. It is formed of two very distinct portions, of a semi-oval shape; one belonging to the upper lip, the other to the lower. Their extremities cross at the commissures, and are confounded with the other muscles of the parts. The use of this fleshy muscle is to bring the lips together, and to close the aperture of the mouth, by making it represent a sort of *bouffet* with radiated wrinkles. It is an antagonist to every other muscle of the lips. 2. *Orbicularis Palpebrarum* seu *oculi seu palpebrarum ciliaris*, *Palpebrarum du'o musc'uli*, *Palpebralis*, *Constrictor seu Sphincter Palpebrarum seu Oculi*, (F.) *Naso-palpébral* (Ch.); *Maxillo-palpébral*, *Palpébral*, *Orbiculaire des paupières*. A muscle common to both the eyelids, and seated in their substance. It is broad, thin, transversely oval, and cleft in its great diameter. It is attached to the nasal process of the superior maxillary bone; to the internal angular process of the frontal bone, and to the orbital process of the superior maxillary bone. From these origins the muscle passes outwards, under the skin of the eyelids, surrounding the orbit in a circular manner, extending somewhat beyond it, and covering the upper part of the cheek. It is inserted into the skin of the eyelids, its upper and inner edge being intimately connected with the frontal and corrugator muscles. Its use is to close the eye, by bringing the eyelids together. The part of the orbicularis which covers the cartilages of the eyelids, and which is remarkably thin, is the *Musculus Ciliaris* of some authors.

ORBICULARIS OCULI, *Orbicularis palpebrarum*, see *Orbicularis muscles*—o. *Oris*, see *Orbicular muscles*—o. *Palpebrarum*, see *Orbicular muscles*—o. *Palpebrarum ciliaris*, see *Orbicular muscles*—o. *Recti*, *Sphincter ani externus*.

ORBICULUS CILIARIS, (dim. of *orbis*, 'a circle,') Ciliary ligament, Ciliary zone.

ORBIS GENU, *Patella*.

OR'BIT, *Or'bita*, *O. oculi*, *Troch'ia*, *Con'chus seu Cav'itas seu Fo'vea seu Pelvic'ula oculi*, (F.)

Orbite, (from *orbis*, 'a circle,') The circular cavities are so called, which lodge the organs of sight. The *orbis* or *orbital fossæ* or *cavities*, *conchi*, *κογχoi*, are situate at the upper part of the face, and are composed of seven bones, viz.,—the *frontal*, above; the *palate* and *superior maxillary*, below; the *sphenoid* and *malar*, externally; and the *ethmoid* and *lachrymal*, internally. The orbit is filled by the globe of the eye, with its different accessory parts—muscles, nerves, vessels, the lachrymal gland, &c. Its margin is termed *margo orbita'lis*.

ORBITA, *Orbit*—o. *Oculi*, *Orbit*.

ORBITAL, *Orbital*—o. *Fissure*, inferior, *Sphenoidal fissure*—o. *Fissure*, superior, *Maxillary fissure*.

OR'BITA R, *Orbita'rius*, *Orbita'lis*, *Or'bital*, (F.) *Orbitaire*. Relating or belonging to the orbit.

ORBITAR APOPH'YSES or PROC'ESSES terminate the two extremities of the orbital arch. The external is much more prominent, and is articulated with the cheekbone; the internal is thinner, and joins the os unguis.

ORBITAR ARCH, (F.) *Arcade orbitaire*, is the projecting, round margin, which separates the *frontal* and *orbital* surfaces of the os frontis, and forms a part of the circumference of the orbit.

OR'BITAR FIS'SURES, (F.) *Fentes orbitaires*. These are situate in the orbit. They are two in number: one *superior*, the *Sphenoidal*; the other *inferior*, the *Sphenomaxillary*.

ORBITAR FORAM'INA. Of these there are several.—1. The *Foram'ena orbitaria super'ius*, (F.) *Trou orbitaire supérieur*, which is situate at the junction of the inner third of the orbital arch with its two outer thirds, and gives passage to the frontal nerve. 2. The *Foram'ina orbitaria interiora seu inter'na*, *Foram'ina Ethmoïdea*. These are situate at the upper part of the internal paries of the orbit, are formed by the union of two notches of the os frontis with two similar ones of the ethmoid; and are distinguished into *anterior* and *posterior*. The former gives passage to the ethmoidal filament of the nasal nerve and to a small artery.

ORBITAR NERVE, *Nervus orbita'lis seu orbita'rius seu subcutaneus malar'is*, is a branch given off from the superior maxillary. It enters the orbit by the sphenomaxillary fissure, and divides into two filaments; the one—*malar*—which passes over the cheekbone, is distributed to the orbicularis palpebrarum, and anastomoses with the facial nerve; the other—*tem'poral*—which passes to the temporal fossæ, and anastomoses with the inferior maxillary and facial, to become afterwards subcutaneous.

ORBITAR SURFACE, see *Maxillary bone* (superior).

ORBITO-EXTUS-SCLÉROTICIEN, *Rectus externus oculi*—o. *Intus-Scléroticien*, *Rectus internus oculi*—o. *Maxillo-labial*, *Levator labii superioris proprius*—o. *Palpébral*, *Levator palpebræ superioris*—o. *Sus-palpébral*, *Levator palpebræ superioris*.

ORCHANET, *Anchusa tinctoria*.

ORCANETTE, *Anchusa tinctoria*.

ORCHALGIA, *Orchidalgia*.

ORCHEA, (*ορχέα*,) *Serotum*.

ORCHET'IS, *Osci'tis*, *Oschei'tis*, *Inflammatio scro'ti*, (*orchæa*, and *itis*.) Inflammation of the serotum, also *Hernia humoralis*.

ORCHEOCATABASIS, *Orchidocatabasis*.

ORCHEOCELE, *Orchiocèle*.

ORCHEOTOMY, (*orchis*, and *τεμνω*, 'to cut,') *Castration*.

ORCHESTROMANIA, (*ορχηστρη*, 'a dancer,' and *mania*.) *Chorea*.

ORCHIDA, (from *orchis*,) *Spirocolon*.

ORCHIDAL'GIA, *Orchial'gia*, *Orchidodyn'ia*, *Orchiodyn'ia*, (*orchis*, and *αλγος*, 'pain.') Pain of the testicle. Neuralgia of the testicle, Irritable testicle; a form of neuralgia.

ORCHIDATROPH'IA, *Atroph'ia testicul'i*, *orchis*, and *atrophia*, 'atrophy.') Atrophy or wasting of the testicle.

ORCHIDITIS, (*orchis*, and *itis*.) Hernia humoralis.

ORCHIDOCARPUM ARIETINUM, (*orchis*, and *καρπος*, 'fruit,') see *Carica papaya*.

ORCHIDOCATAB'ASIS, *Orcheocatab'asis*, *Descensus testiculo'rum*, (*orchis*, and *καταβασις*, 'descent.') The descent of the testes into the scrotum.

ORCHIDOCELE, (*orchis*, and *κηλη*, 'tumor,') Hernia humoralis.

ORCHIDODYNIA, (*orchis*, and *δυνη*, 'pain,') Orchidalgia.

ORCHIDOMYELO'MA, (*orchis*, *μυελος*, 'marrow,' and *ωμα*.) Medullary, fungus of the testicle.

ORCHIDONCUS, (*orchis*, and *ογκος*, 'a swelling,') Hernia humoralis.

ORCHIDOSCIRRHUS, see *Orchiocoele*.

ORCHIDOTOMY, (*orchis*, and *τεμνειν*, 'to cut,') Castration.

ORCHILL, *Lichen roccella*.

ORCHIOCE'LE, *Orcheoce'le*, (*orchis*, or *orchea*, and *κηλη*, 'a rupture,') A tumour of the testicle. Scrotal hernia. Hernia humoralis. A name given to several diseases of the testicle and its envelopes, essentially differing from each other.

ORCHIOCE'LE SCIRRHOS'A, *Scirrheoce'le*, *Orchidoscirrhus*, *Orchiocirrhus*. A scirrhous tumor of the testis.

ORCHIODYNIA, (*orchis*, and *δυνη*, 'pain,') Orchidalgia.

ORCHIOSCIRRHUS, see *Orchiocoele*.

ORCHIOTOMIA, (*orchis*, and *τεμνειν*, 'to cut,') Castration.

ORCHIS, gen. *Or'chidis* or *Orchis*, (*ορχις*, gen. *ορχιδος*, or *ορχως*.) Testicle—o. *Bifolia*, see *Salep*—o. *Butterfly*, *Orchis bifolia*—o. *Latifolia*, see *Salep*—o. *maculata*, see *Salep*.

ORCHIS MAS'CU'LA, *Saty'rion*, *Testiculus caninus*, *Cynosor'chis*, *Dogs' stones*, *Male or'chis*, *Ballcockgrass*, *Male Fools' stones*, *Twinroot*. *Ord.* *Orchidaceæ*. This root had a place in some pharmacopœias, on account of the glutinous, slimy juice it contains. The root of the *Orchis bifolia* or *Butterfly orchis* has likewise been collected. It has been used as a demulcent. Its fancied aphrodisiac virtues seem to be owing to its resemblance to *ορχις*, 'a testicle,' whence its name.

Salep is obtained from this variety of the orchis, and from the *Orchis morio*, &c. See *Salep*.

ORCHIS MILITARIS, see *Salep*—o. *Round-leaved*, *Large*, *Platanthera orbiculata*.

ORCHITE, Hernia humoralis.

ORCHITIS, (*orchis*, and *itis*, 'inflammation,') Hernia humoralis.

ORCHITOMIA, (*orchis*, and *τεμνειν*, 'to cut,') Castration.

ORCHOI, (*ορχος*, 'a row,') see *Tarsus*.

ORCHOTOMIA, (*orchis*, and *τεμνειν*, 'to cut,') Castration. Also, removal of the tarsi.

ORDEAL NUT, *Calabar' Bean*; (F.) *Fève du Calabar*, *Fève épreuve du Calabar*; called *Esere* by the natives of Calabar, is from a plant of the natural family, *Leguminosæ*—the *Physostig'ma venenosum*—closely resembling a *Dolichos*. It is possessed of virulent acro-narcotic properties. One or two drops of a solution of an extract of the bean, introduced under the eyelids, rapidly contracts the pupil, and may be used where myositis are indicated. Its active alkaloid is termed *Physostig'mine*.

ORDEOLUM, *Hordeolum*.

ORDEUM, *Hordeum*.

ORDINAIRES, (from (L.) *ordo*, gen. *ordinis*, 'an established process,') *Menses*.

ORDO MEDICINALIS, *Police*, medical.

ORDONNANCE, (from *ordonner*, 'to order,') Formula, Prescription.

OREILLE, (from (L.) *auris*.) Ear—o. *d'Âne*, Symphylum—o. *d'Homme*, *Asarum*—o. *d'Ours*, *Primula veris*—o. *de Souris*, *Hieracium pilosella*. **OREILLETTE**, (dim. of *oreille*, (L.) *auricula*, 'a small ear,') *Asarum*, *Auricle* of the heart.

OREILLONS, (from *oreille*.) *Parotis*, *Cynanche parotidæa*.

ORELIA GRANDIFLORA, *Allamanda*.

ORELLANA, see *Terra Orleana*.

OREOSELINUM, (*ορος*, 'a mountain,' and *σελινον*, 'parsley,') *Athamanta aureoselinum*—o. *Africanum*, *Bubon galbanum*—o. *Legitimum*, *Athamanta aureoselinum*—o. *Nigrum*, *Athamanta aureoselinum*.

OREXIS, (*ορεξις*.) Appetite, Pyrosis—o. *Cynodes*, *Boulimia*.

ORGAN, *Or'ganum*, *Or'ganon*, (*οργανον*, from *εργον*, 'work,' 'an instrument,') Part of an organized being, destined to exercise some particular function. Thus, the eyes are organs of sight; the muscles are organs of motion, &c. The collection of organs, which concur in any function, is called *apparatus*.

ORGAN, *Mentha pulegium*.

ORGAN OF GIRALDÈS, *Corps innommé de Giralde*. A small linear body at the upper end of the testicle in the spermatic cord, regarded by Giralde as the remains of the Wolffian body, and analogous to the 'appendage to the ovary' of the female.

ORGAN OF ROSENMULLER, *Parovarium*.

ORGANA GENERATIONI INSERVIENTIA, Genital organs—o. *Genitalia*, Genital organs—o. *Sexualia*, Genital organs—o. *Sudoripara*, see *Perspiration*—o. *Urogenitalia*, Urogenital organs—o. *Uropœtica*, Urinary organs.

ORGANES GÉNITAUX, Genital organs.

ORGAN'IC, (*οργανικος*.) *Organ'icus*, (F.) *Organique*. Relating to an organ or organs, and to beings possessed of organs. Hence, **ORGANIC FUNCTIONS** are those possessed by both animals and vegetables. It, as well as *teleorgan'ic*, is also used synonymously with *vital*, in contradistinction to *physical*.

ORGANIC CHEMISTRY, see *Chymistry*.

ORGANIC DISEASES or LE'SIONS—*Nosorganica* of Bouchut—are such as concern the organ itself, in contradistinction to *functional*, *rational*, or *dynam'ic*, which merely concern the function. Of course, *organic diseases* of a part are always more serious than *functional*.

ORGANIC NERVOUS SYSTEM, see *Trisplanchnic nerve*.

ORGANICISM, or ORGAN'ICISM, *Organicism'us*, (from *organ*.) The doctrine of the localization of disease, or which refers it always to a material lesion of an organ. *Organicism* is the antithesis to *vitalism*. The great principle of the supporters of the former doctrine, (F.) *École organicienne*, is, that life is the result of organization.

ORGANISATIO, *Organization*.

ORGANISATUS, *Organized*.

ORGANISÉ, *Organized*.

ORGANISM. The living economy. Many physiologists have used this word synonymously with organization; but it more especially designates the aggregate of vital actions in organized beings, animal as well as vegetable. The collection of parts composing an organized body, and the laws which govern it.

ORGANITE, (from *organ*.) According to Dr. Milne Edwards, an elementary organ,—a blood-corpuscle, for example.

ORGANIZABILITY, (*organisatio*, and *ἡαδι*

lis, 'suitable.') Capability of organization, or of being formed into living tissue. Substances possessed of such capability—as fibrin—are said to be organizable.

ORGANIZABLE, see Organizability.

ORGANIZATION, *Organisat'io*. The manner of structure proper to different beings. It is also employed in the sense of the structure of a part; as the organization of the heart, &c.

ORGANIZATION, COMPOUNDS OF, Principles, immediate.

ORGANIZED, *Organis instructus*, *Organisatus*, (F.) *Organisé*. That which is composed of organs.

ORGANIZED BOD'IES, (F.) *Corps organisés*, are those endowed with life; i. e. animals and vegetables: in contradistinction to *inorganic* bodies, which comprise the mineral kingdom. See *Corpus*.

ORGANO-CHEMIA, Chymistry, organic.

ORGANOGEN'Y, *Organogen'ia*, *Organogen'esis*, (organ, and *γενναις*, 'generation.') The doctrine of the formation of different organs.

ORGANOGENO'SIA, (organ, and *γνωσις*, 'knowledge.') The knowledge and discrimination of organized bodies.

ORGANOGRAPHY, *Organograph'ia*, (organ, and *γραφη*, 'a description.') A description of the organs of a living body.

ORGANOLEPTIC, (organ, and *ληψις*, 'action of receiving,') Plastic.

ORGANOLOGY, (organ, and *λογος*, 'a description,') Anatomy, Craniology.

ORGANON, (*organon*), Organ—o. *Olfactūs*, *Nasus*.

ORGANONOM'IA, (organ, and *νομος*, 'a law.') The doctrine of the laws of organic life.

ORGANOPATHISM, *Organopathol'ogy*, (organ, and *παθος*, 'an affection.') The doctrine of organopathies, or of 'organopathic states' of Piorry, which consists, essentially, in studying the pathological condition of the organs in each individual case.

ORGANOPLASTIES, (organ, and *πλασσειν*, 'to form,') see Training.

ORGANOSCOPY, (organ, and *σκοπεω*, 'I examine,') see Craniology.

ORGANOZOONOMIA, (organ, *ζωον*, 'an animal, and *νομος*, 'a law,') Zoonomia.

ORGANUM, (*organon*), Instrument, Organ.

ORGASM, (*orgasmos*), *Orgas'mus*, *Orgas'is*, (*Ἔστρος*, *Ἔστρον*, *οργασμ*, 'I desire ardently, 'I am excited,') A strong impulse or desire for something. State of excitement and turgescence of any organ whatever. Erethism. Applied particularly to the height of venereal excitement in sexual intercourse.

ORGASMUS, Orgasm, Turgescence.

ORGAS'TICA. The second order of the class *Genetica* of Good. Defined to be diseases affecting the orgasm; organic or constitutional infirmity disordering the power or the desire of procreating.

ORGE, (*οργη*), Rage.

ORGE, Hordeum—o. *Perlé*, see Hordeum.

ORGEAT, (from *orge*, because formerly made with barley water,) Syrupus amygdale.

ORGELET, (dim. of *orge*), Hordeolum.

ORGEOLET, (dim. of *orge*), Hordeolum.

ORGOSIS, Orgasm.

ORICALCUM, (auricalcolum, or from *oros*, 'a mountain, and *χαλκος*, 'brass,') Brass.

ORICULAR, Auricular.

ORICULE, Auricle, Pavilion of the ear.

ORIFICIUM, (*os*, gen. *oris*, 'a mouth, and *facere*, 'to make,') Mouth.—o. *Dextrum Ventriculi*, *Pylorus*—o. *Uteri*, *Os uteri*—o. *Uterinum*, see *Uterus*—o. *Vaginale*, see *Uterus*—o. *Ventriculi sinistrum*, *Cardia*.

ORIGANUM, *O. vulgare* seu *heracleoticum*, *origanum*, from *oros*, 'a mountain, and *γενος*, 'delight,') *Cimicifuga bulbosa*, *Majorana mancarina*,

Zuzarhen'di her'ba, Wild or Common *mar'joram*, *Mancur'na*, (F.) *Pied de lit*. *Ord. Labiatae*. *Sec. Syst* *Didynamia Gymnospermia*. This plant resembles *marjoram* in smell; and has a pungent taste, much like thyme. It is aromatic and stimulant; and its oil possesses the properties of the essential oils in general. The oil, *Oleum origani* (Ph. U. S.), is very pungent, and is rarely given internally. It is used in toothache.

ORIGANUM AQUATICUM, *Eupatorium canadense*—o. *Creticum*, *O. dictamnus*.

ORIGANUM DICTAM'NUS, *O. Creticum*, *Dictamnus Creticus*, *Amaracus tomentosus*, *Onitis*, *Dittany of Crete*, (F.) *Dictanne de Crète*. The leaves of this plant were once recommended as emmenagogue and alexipharmic.

ORIGANUM HERACLEOTICUM, *Origanum*.

ORIGANUM MAJORA'NA, *O. majoranoides*, *Majorana*, *M. hortensis*, *Sweet mar'joram*, *Samp'sychem*, *Samp'sychem*, *Agriorig'annum*, *Amaracus*, (F.) *Marjolaine*. Its odour is strong and fragrant; taste aromatic and bitterish. Its properties are those of a stomachic and errhine. It is chiefly used, however, for culinary purposes, and as a snuff in headache.

ORIGANUM SYRIACUM, *Teucrium marum*.

ORIGINAIRE (F.), (*origo*, gen. *originis*, 'origin,') This term has been used, by French writers, in the same sense as *congenital*. It is particularly applied to affections depending upon faulty conformation, or to faulty conformation itself.

ORIGO, (*orior*, 'to arise,') *Arche*.

ORISMOLOGY, (*ορισμος*, 'a definition, and *λογος*, 'a discourse,') Terminology.

ORLEANA, see *Terra Orleana*.

ORME, (*ορμη*), Appetite.

ORME, Ulmus—o. *Champêtre*, *Ulmus*—o. *Pyramidalis*, *Ulmus*.

ORMEAU, Ulmus.

ORMS'KIRK MED'ICINE. A celebrated antilyssic remedy, supposed to consist of *Pulv. Crætæ*, ʒss; *Bol. Armen.* ʒijj; *Alumin. gr. x*; *Pulv. Iulmæ*, ʒj; *O. Anisi*, gtt. vj.

This is taken for a dose every morning for six times, in a glass of water, with a small proportion of fresh milk.

ORNAMENTUM FOLIACEUM, see *Tube*, *Fallopian*.

ORNITHOG'ALUM ALTIS'SIMUM. (*ορνιθoγαλον*, a fanciful name, from *ornis*, gen. *ornidis*, 'a bird, and *γαλα*, 'milk,') A South African plant, *Nat. Ord.* *Asphodeleæ*, the fleshy bulb of which is diuretic. An oxymel is prepared of it, which is given in catarrh, asthma, phthisis, and hydrothorax. It resembles squill in its medical properties.

ORNITHOGALUM MARITIMUM, *Scilla*—o. *Squilla*, *Scilla*.

ORNITHOGLOSSA, (*ornis*, gen. *ornidis*, 'a bird, and *γλωσσα*, 'a tongue,') see *Fraxinus excelsior*.

ORNUS EUROPEA, see *Fraxinus ornus*—o. *Mannifera*, *Fraxinus ornus*—o. *Rotundifolia*, *Fraxinus ornus*—o. *Sylvestris*, *Fraxinus excelsior*.

OROBANCH'E VIRGINIA'NA, (*orobus*, and *αγχω*, 'I strangle,') *Epiphe'gus seu Epifa'gus Americana* seu *Virginiana*, *Virgin'ia* broomrape, *Beechdrops*, *Cancerroot*. Family, *Orobanchoides*. This parasitic plant is astringent, and a peculiar and extremely nauseous bitter. It is most powerful when fresh. It has been used in dysentery; and, externally to obstinate ulcers.

OROBANCH'E AMERICA'NA, and *O. UNIFLO'RA*, have the same properties, and are likewise, called *Cancerroot*, *Earthclub*, and *Claywort*.

OROB'E, (*οροβος*), *Ervum erilia*.

OROBION, see *Ervum erilia*.

OROBOT'DES, (*orobus*, and *ειδος*, 'form,') *Oroboides*. Having the shape of the orobus. A name given to urine, *Urina oroboides seu oroboi-*

des, when depositing a substance of a fawn colour, like the meal of the orobus.

OROBUS, (οροβος,) *Ervum ervilia*—o. *Faba*, *Vicia faba*.

ORANGE, (from *orange*, its color,) see *Agaric*.

ORONOSUS, (ορος, 'a mountain,' and νοσος, 'a disease,') *Puna*.

OROPHALL'US, (ορος or οροφος, 'the rump,' and φαλλος, 'the male organ,') A monster having a second male organ originating from the rump.

OROS, (ορος,) *Croupion*, Serum.

ORPIMENT, *Auripigmentum*, *Or'pin*, *Rissigallum*, *Rissigallum*, *Clo'ans*, *Ela'mē*, *S'ira*, *Lempnia*, *Spec'ulum cil'rimum*, *Y'rides*, *Y'ride*. The native yellow sulphuret of arsenic. It is poisonous, but less so than the oxide. It is, also, called *Adari'go*, *Adar'neck*, *Ahu'sal*, *Althana'cha*, *Azar'net*. The Turks prepare a depilatory with it, which they call *Rus'ma*.

ORPIN, Orpiment—o. *Brulant*, Sedum—o. *Grand*, Sedum telephium—o. *Reprise*, Sedum telephium.

ORPINE, Sedum telephium.

ORRHOCHEZIA, (ορρος, 'serum,' and χεζω, 'to go to stool,') *Diarrhœa*, serous.

ORRHOCYST'IS, (ορρος, 'serum,' and κυστις, 'a bladder,') An encysted tumour containing a serous fluid.

ORRHOMYMENES, (ορρος, 'serum,' and ὑμνη, 'a membrane,') see *Membrane* (serous).

ORRHOMYMENTIS, (ορρος, 'serum,' ὑμνη, 'a membrane,' and itis,) *Hydrohymenitis*.

ORRHOPYG'ION, (ορρος, 'extremity,' and πυγν, 'the nates,') The inferior extremity of the vertebral column. The os coccygis. Also, the raphe, extending from the penis to the anus, and separating the scrotum into two parts.

ORRHORRHŒ'A, (ορρος, 'serum,' and ρευ, 'I flow,') A morbidly increased secretion of a serous fluid.

ORRHOS, (ορρος,) *Croupion*, Serum.

ORRHOMENES, (ορρος, 'serum,' and ὑμνη, 'a membrane,') see *Membrane* (serous).

ORRHIMENTIS, (orrhymenes, and itis,) *Hydrohymenitis*.

ORRIS, *Iris Florentina*—o. Common, *Iris Germanica*.

ORSEILLE, *Lichen roccella*.

ORTEIL, (from *articulus*, 'a joint,') *Digitus pedis*.

ORTHO, (from ορθος,) straight, right, upright, in composition.

ORTHOC'LUM, (ortho, and κωλον, 'limb,') Stiffness of the limb so that it is kept quite straight.

ORTHODON'TIA, *Orthodon'ticē*, *Orthodontosie*, (ortho, and οδους, gen. οδοντος, 'a tooth,') The part of dentistry which is concerned in the treatment of irregularity of the teeth.

ORTHODORON, (ορθοδωρον, from ortho, and δωρον, 'a palm,') A measure with the Greeks, equal to eleven finger-breadths.

ORTHOGNATHOUS, (ortho, and γναθος, 'jaw,') Having a vertical jaw. A term applied to the form of head in which the facial angle approaches the right angle; *Orthognathus*.

ORTHOMORPHIA, (ortho, and μορφη, 'shape,') *Orthopædia*.

ORTHOMORPHOSIS, *Orthopædia*.

ORTHONTROPIA, (ortho, and εντροπειν, 'to convert,') *Orthopædia*.

ORTHOPÆD'IA, *Orthopæ'dicē*, *Orthomor'phia*, *Orthomorpho'sia*, *Orthontrop'ia*, *Orthopæ'dies*, (ortho, and παις, gen. παιδος, 'a child,') The part of medicine whose object is to prevent and correct deformity in the bodies of children. Often used, however, with a more extensive signification, to embrace the correction or prevention of deformities at all ages. *Orthosomat'ics*, *Orthosomat'icē*, (from ορθος, 'right,' and σωμα, 'body,') has been proposed as a preferable term.

ORTHOPE'DIC, *Orthopæ'dicus*. Relating to orthopædia,—as *Orthopæ'dic Institution*, *Institutum orthopæ'dicum* seu *orthopæ'dium*: an institution for the correction or prevention of deformities in children.

ORTHOPEDICS, *Orthopædia*.

ORTHOPE'DIST, *Ortho'ter*, *Orthopæ'dicus*. One who practises orthopædia.

ORTHOPHON'IA, (ortho, and φωνη, 'voice,') A good pronunciation.

ORTHOPHREN'IC, *Orthophren'icus*, (ortho, and φρην, 'mind,') Relating to sound mind, and to the cure of insanity.

ORTHOPHTHAL'MIC, *Orthophthal'micus*, (ortho, and οφθαλμος, 'eye,') Relating or belonging to the correction of deviations of the eyeball,—as of strabismus.

ORTHOPNŒ'A, *Suffocatio*, *Strangulatio*, *Profocatio*, *Phia*, *Phia'tis*, *Phig'os*, *Phig'ma*, *Euthyp'noē*, *Euthypnœ'a*, *An'gor*, *Apnœ'a* (ortho, and πνœα, 'I respire,') Impracticability of breathing in the horizontal posture. Necessity of being in the erect posture, in order to respire.

ORTHOPNŒA CARDIACA, *Angina Pectoris*—o. *Convulsiva*, *Asthma*—o. *Cynanchica*, *Cynanche trachealis*—o. *Hydrothoracica*, *Hydrothorax*—o. *Hysterica*, *Angone*—o. *Pituitosa*, *Mycorthopnœa*—o. *Tussiculosa*, *Pertussis*.

ORTHOSOMATICE, (ortho, and σωμα, 'body,') *Orthopædia*.

ORTHOSOMATIQUE, *Orthopædia*.

ORTHOTER, (ορθωτηρ,) *Orthopedist*.

ORTHYSTEROPTOMA, gen. *Orthysteropto'matis*, (ortho, ὄσθηα, 'uterus,' and πτωμα, 'a fall,') *Prolapsus uteri*.

ORTHYSTEROPTOSIS, *Prolapsus uteri*.

ORTIE, *Urtica*—o. *Blanche*, *Lamium album*—o. *Brûlante*, *Urtica urens*—o. *Morte*, *Lamium album*—o. *Morte des Bois*, *Galeopsis*—o. *Romaine*, *Urtica pilulifera*.

ORTILLUS, (from *articulus*, 'a joint,') *Digitus pedis*.

ORTYX, (ορτυξ,) *Tetrao coturnix*.

ORUS, (ορος,) *Croupion*, Serum.

ORVALE. *Salvia sclarea*.

ORVIETANUM. (From the Italian quack, Hieronymus Ferrantes, of Orvieto.) An electuary regarded as an invaluable antidote, and composed of *Old theriac*, *Dried vipers*, *Scorzonera*, *Carlina*, *Imperatorium*, *Angelica*, *Bistort*, *Rosemary*, *Juiper*, *Cinnamon*, *Cloves*, *Mace*, *Honey*, and a crowd of other substances. Its medical properties resembled, in some respects, those of the theriac. (It was so called, according to some, because invented by Orvietano, a celebrated charlatan; or, according to others, from Orvieto, a town in Italy.)

ORY'ZA, (ορυζα,) *Ory'za sat'iva*, *Ory'zum*, *Rice*, (F.) *Riz*. Family, Gramineæ. *Sex. Syst.* Hexandria Monogynia. A plant, almost aquatic, indigenous in China, one of the richest productions of Egypt, and which prospers in the warmer regions of the four quarters of the globe. The grain inclosed in the husk is called by the Malays *Pad'dy*, *Padi*, or *Pad'die*; when deprived of the husk, *Bras*, and when boiled, *Nasi*. Its amylaceous seeds, *Ory'za*, *Ory'zum*, *Hor'dæum galactiticum*, nourish more men than those of wheat and rye together. Besides being alimentary, they are demulcent and emollient. *Gard'ner's alimentary preparation* is very finely ground rice meal.

Mix'clage of Rice, *Rice water*, may be formed by taking one ounce of rice, and, having washed it, macerating it for three hours in a quart of tepid soft water, in a pan placed upon a stove; then boiling the whole slowly for another hour, and straining through a sieve.

It may be sweetened and acidulated, or be prepared with milk, like arrowroot, and be used as a demulcent.

Rice Jelly is made by macerating a sufficient quantity of clean *rice* in as much *water* as will cover it; boiling slowly, adding *water* as it evaporates, until the *rice* is reduced to a pap, sweetening and flavouring with *lemonjuice* or *vanilla*, and passing through a fine sieve into a form or mould. It may be eaten alone or with milk; and is a good dietetic preparation for those of weak digestive powers.

Ground rice. Take a tablespoonful of *ground rice*, a pint and a half of milk, and half an ounce of *can'died lem'oupeel*. Rub the *rice* smooth with the milk: then add the *lemonpeel* cut into small pieces; boil for half an hour, and strain whilst the milk is hot. It is used in the same cases as *rice jelly*.

ORYZUM, (ορυζον,) Oryza.

OS, gen. *Os'sis*, Bone, dim. *Ossiculum*: OS, gen. *O'ria*, Mouth — o. Acromii, Acromion — o. Adjutorium, Humeri os — o. Alaeforme, Sphenoid — o. Alagus, Coccyx, Sacrum — o. Alatum, Sphenoid bone — o. Amphideon, Os uteri — o. Anonyma, Innominatum os — o. Anonymum, Innominatum os — o. Anticum, *Avant bouche* — o. Arcuale, Temporal bone — o. Armale, Temporal bone — o. Asser, Sternum — o. *de l'Assiette*, Ischion — o. Azygos, Sphenoid — o. Ballistæ, Astragalus — o. Basilare, Occipital bone, Palate bone, Sacrum, Sphenoid — o. Baxillare, Sphenoid — o. Bicornæ, Hyoides os — o. Brachiale, Humeri os — o. Brachii, Humeri os — o. *du Bras*, Humerus — o. Calcis, Calcaneum — o. Capitatum, Magnum os — o. Caudæ, Occipital bone — o. Clunium, Sacrum — o. Coccygis, Coccyx — o. Colatorii, Sphenoid bone — o. Colatorium, Ethmoid bone — o. Coliforme, Ethmoid — o. Convolutum superius, Turbinate bone, middle — o. Coronale, Frontal bone — o. Coxæ, Innominatum os — o. *Coxal*, Innominatum os, Ischion — o. *Cribleux*, Ethmoid — o. Cribiforme, Ethmoid — o. Cribrosum, Ethmoid — o. Cristatum, Ethmoid — o. *Crochu*, Os unciniforme.

OS CROTAPHALÆ. A name given by M. Bèclard to a bone sometimes met with at the anterior and inferior angle of the parietal bone; from κροτάφος, 'the temple.'

OS CROTAPHITICUM, Temporal bone — o. Cubicum, Ethmoid bone — o. ad Cubitale, Radius — o. Cubiti Inferius, Ulna — o. *de la Cuisse*, Femur — o. Cuneiforme, Sphenoid — o. Cuneo comparatum, Sphenoid — o. Cymbiforme, Scaphoid bone — o. Disciforme, Patella — o. Durum, Temporal bone — o. Ensiforme, Sternum.

OS EPACTALÆ, (from επακτος, 'superadded.') A name given to one of the ossa Wormiana; particularly to one met with in the posterior fontanelle, and which has been described by G. Fischer, under the name *Os Epactal'le* seu *Goëthia'num*.

OS ÉPACTAUX, Wormiana ossa.

OS EXTER'NUM. The entrance into the vagina, in contradistinction to the *Os inter'num* or *Os u'teri*.

OS EXTRA ORDINEM CARPI, Os pisiforme — o. Femoris, Femur — o. Fibrosus, Occipital bone — o. Foraminulentum, Ethmoid — o. Frontis, Frontal bone — o. Genæ, Malæ os — o. Gladioli, Sternum — o. *Grand*, Magnum os — o. *Grand, col du*, Collum ossis magni — o. Grandinosus, Cuboid — o. Gutturis, Hyoides os — o. Hamatum, Os unciniforme — o. *Hors du rang*, Os pisiforme — o. Hyoides, Hyoides os — o. Hypopium, Malæ os — o. Hypseloides, Hyoides os — o. Inconjugatum, Sphenoid — o. Incudi similis, Incus — o. Inferius ventriculi, Pylorus — o. *Innominé*, Innominatum os — o. *Intercale*, Wormiana ossa — o. Internum, see Uterus — o. Interparietale, Interparietal bone — o. Inverecundum, Frontal bone — o. Ischii, Ischion — o. Jugale, Malæ os — o. Jugamentum, Malæ os — o. Juguli, Clavicla — o. Lacrymale, Unguis os — o. Lambdæ, Occipital bone — o. Lambdoides, Hyoides

os, Occipital bone — o. Lapideum, Temporal bone — o. Lapidosum, Temporal bone — o. Latum, Sacrum — o. Latum humeri, Scapula — o. Lenticulare, Os orbiculare, Os pisiforme — o. Lepidoides, Temporal bone — o. Linguae, Hyoides os — o. Linguale, Hyoides os — o. Lithoides, Temporal bone — o. Magnum, neck of the, Collum ossis magni — o. *Malair*, Malæ os — o. Malare, Malæ os — o. Maxillæ superioris undecimum, Vomer — o. *Maxillaire inférieure*, Collum mandibulæ — o. Memento mori, Temporal bone — o. Memoria, Occipital bone — o. Mendosum, Temporal bone — o. Morsus Adami, Thyroid cartilage — o. Mucronatum, Xiphoid cartilage — o. Multangulum majus, Trapezium os — o. Multangulum minus, Trapezoides os — o. Multiforme, Ethmoid, Sphenoid — o. *Nasauz*, Nasal bones — o. Naviculare, Os scaphoides — o. Nervale, Temporal bone — o. Nervosum, Occipital bone — o. Occipiti, Occipital bone — o. Orbiculare, Os pisiforme — o. *du Palais*, Palate bone — o. Palati, Palate bone — o. *Palatin*, Palate bone — o. Papillare, Sphenoid — o. Parietale inferius, Temporal bone — o. Paxillare, Sphenoid — o. Pectinis, Pubis, os — o. Pectoris, Sternum — o. Pelvicephalicum, Occipital bone — o. Pelvis laterale, Innominatum os — o. Perone, Fibula — o. Petrosus, Temporal bone — o. Polygonum majus, Trapezium os — o. Polygonum minus, Trapezoides os — o. Polymorphon, Sphenoid — o. *de la Pommette*, Malæ os — o. Posterum, Pharynx — o. Præruptæ rupi assimilatam, Temporal bone — o. Procutibale, Ulna — o. *Propres du nez*, Nasal bones — o. Proræ, Frontal bone, Occipital bone — o. Pterygoideum, Sphenoid — o. Pubis, see Pubis, os — o. Pudicum, Malæ os — o. Puppis, Frontal bone, Occipital bone — o. Pyramidale carpi, Trapezoides os — o. Pyxidis, Occipital bone — o. Rationis, Frontal bone — o. *du Rayon*, Radius — o. Sacrum, Sacrum — o. Saxeum, Temporal bone — o. Scutiforme, Patella — o. *de Sèche*, Sepia — o. Sedentarium, see Ischiatic — o. Semilunare, Lunare os — o. Sextum cranii, Occipital bone — o. *du Sinciput*, Parietal bones, see Sinciput — o. *Sons ethmoidal*, Turbinate bone, inferior — o. Sphecoideum, Sphenoid — o. Sphænobasilare, Occipital bone — o. Sphenoidale, Sphenoid — o. Sphenoides, Sphenoid — o. Spongiosum, Ethmoid — o. Spongiosum inferius, Turbinate bone, inferior — o. Spongiosum medium, Turbinate bone, middle — o. Spongiosum superius, Turbinate bone, middle, Turbinate bone, superior — o. Spongoides, Ethmoid — o. Squamosus, Temporal bone — o. Suboculare, Malæ os — o. *Surnuméraires*, Wormiana ossa — o. *Susmaxillaire*, Maxillary (superior) bone — o. Sylvii, Os orbiculare — o. Synepitidis, Frontal bone — o. Temporis, Temporal bone — o. Tessera, Astragalus, Cuboid — o. Thyreoides, Patella — o. Tibiæ minus, Fibula — o. Tineæ, Os uteri — o. *Triangulaires*, Wormiana ossa — o. Turbinatum, Turbinate bone, middle — o. Turbinatum inferius, Turbinate bone, inferior — o. Uncinatum, Os unciniforme — o. Unguis, see Unguis os — o. Upsiloides, Hyoides os.

OS U'TERI, *Os tin'ce* seu *inter'num* seu *amphid'leon* seu *amphid'eum*, *Protos'porus*, *Os'chion*, *Hysteros'toma*, *Os'culum* seu *orific'ium u'teri*, *Hysteros'tom'ium*, (F.) *Musæum de Tanche*. The mouth of the womb, see Uterus.

OS UTERI EXTERNUM, see Uterus — o. Uteri internum, see Uterus — o. Varium, Cuboid — o. Vespiforme, Sphenoid — o. Vomeris, Vomer — o. *Wormiens*, Wormiana ossa — o. Xiphoides, Sternum — o. Ypseloides, Hyoides os — o. Zygomaticum, Malæ os.

OSANORE, see Artificial teeth.

OSCEDO, (os, 'the mouth,' and *ciere, citum*, to move,) Yawning.

OSCHÆMATÆDE'MA, gen. *Oschemate'dematia*, *Oschemede'ma*, *Ed'e'ma scroti cruent'um*, (osche, 'aqua,' 'blood,' and *oichma*, 'oedema.') Effusion of blood into the scrotum.

OSCHÆMEDEMA, *Oschæmatœdema*.

OSCHE, (ὄσχη,) Scrotum.

OSCHEITIS, (*osche*, and *itis*,) Orchitis, Oschitis.

OSCHEOCARCINOMA, Cancer, chimney-sweepers'.

OSCHEOCE'LE, *Oschoece'lē*, *Serotoce'lē*, (*oscheon*, and *κλήη*, 'tumour,') *Sero'tal her'nia*, see Bubonocœle. Sauvages uses *oscheocœle* in the sense of *Drop'sy of the scrotum*, see Hydrocœle.

OSCHEOCELE AQUOSA, *Hydroscheocœle*—o. Flutula, *Physocœle*—o. *Hydatidosa*, *Hydatidocœle*—o. *Œdematica*, *Œdematoscheocœle*—o. *Seminalis*, *Spermatoœle*—o. *Urnalis*, *Urocœle*—o. *Vari-cosa*, *Varicoœle*.

OSCHEOCHALASIE, (*oscheon*, and *χαλασις*, 'relaxation,') A tumour resulting from hypertrophy of the areolar tissue of the scrotum, and the excessive extension of the skin of the part,—an affection which has been confounded with sarco-cœle.

OSCHEON, (ὄσχειν,) Scrotum.

OSCHEON'CUS, *Oschon'cus*, *Oschœophy'ma*, *Oschœophy'ma*, (*oscheon*, and *ὄγκος*, 'a tumour,') A preternatural swelling of the scrotum.

OSCHEOPHYMA, (*oscheon*, and *φύμα*, 'a swelling,') *Oscheoncus*.

OSCHEOPLAS'TIC, *Oscheoplast'icus*, (*oscheon*, and *πλασσω*, 'I form,') An epithet given to the operation for curing the scrotum when lost. *Os'cheoplasty*.

OSCHEOPYŒDE'MA, gen. *Oscheopyœde'matis*, *Oschopyœde'ma*, *Œde'ma scroti purulen'tum*, (*oscheon*, *πύον*, 'pus,' and *œdema*.) Purulent tumefaction of the scrotum.

OSCHEUS, (ὄσχεος,) Scrotum.

OS'CHION, (ὄσχιον.) The raised margin of the os uteri. According to some, an excrescence from the os uteri.

OSCHI'TIS, *Orchei'tis*, *Oschei'tis*, *Inflamma'tio scro'ti*, (*osche*, and *itis*.) Inflammation of the scrotum.

OSCHOCARCINOMA, gen. *Oschocarcino'matis*, Cancer, chimney-sweepers'.

OSCHOCELE, *Oscheocœle*, *Serotoœle*.

OSCHONCUS, *Oscheoncus*.

OSCHOPHYMA, gen. *Oschœophy'matis*, (*oschus*, and *φύμα*, 'a swelling,') *Oscheoncus*.

OSCHOPYŒDEMA, *Oscheopyœdema*.

OSCHURŒDE'MA, gen. *Oschurœde'matis*, (*oschus*, *ουρον*, 'urine,' and *ὄδημα*, 'œdema,') *Œdema* from effusion of urine into the scrotum.

OSCHUS, (ὄσχος,) Scrotum.

OSCHYDRŒDE'MA, (*oschus*, *ὕδωρ*, 'water,' and *ὄδημα*, 'œdema,') *Œdema scroti aquo'sum*. Watery œdema of the scrotum.

OSCILLA'TIO, Irritability, Oscillation.

OSCILLA'TION, *Oscilla'tio*, (from *oscillare*, *oscillatum*, 'to swing,') Vibration, *Vibra'men*, *Vibra'tio*, *Vibra'tus*. Also, a partial rotatory movement of the eyeball to and from its antero-posterior axis.

OS'CITANT (FEVER), *Fe'bris os'citans*, (F.) *Fèvre oscitante*, (*oscitare*, 'to yawn,' from *os*, 'the mouth,' and *cierē*, *citum*, 'to move,') A fever, in which the patient is continually yawning.

OSCITATIO, Yawning.

OSCU'LATORIUS, (from *osculum*, dim. of *os*, 'a mouth,') Orbicular oris.

OSCU'LU M UTERI, (*osculum*, dim. of *os*, 'a mouth,') Os uteri.

OSCUS, (ὄσχος,) Scrotum.

OSEILLE, (from *oxalis*,) *Rumex acetosus*—o. *Bouche*, *Rumex scutatus*—o. *Petite*, *Rumex scutatus*—o. *Ronde*, *Rumex scutatus*—o. *Rouge*, *Rumex sanguineus*.

OSIER, (from *osier*,) *Salix alba*—o. Red, *Cornus sericea*.

OSIS, (ωσις.) A suffix in many terms, which

signifies the condition in general; and, in pathological terms, the morbid condition.

OS'MAZOME, *Os'mozome*, *Osmazo'ma*, *Osmo-zo'ma*, *Os'mazone*, *Zonod'mum*, (F.) *Matière extractive du Bouillon*, *Extrait de viande*; *Extrac-tive* or *Suppona'ceous ex'tract of meat*; (*osme*, and *ζωμος*, 'soup,') M. Thénard gave this name to an extractive matter, contained in muscular flesh and in the blood of animals, which he considers of a peculiar nature. It has an agreeable smell and taste, and is found in *Bouillons* of meat, in the proportion of one part to seven of gelatin. Vauquelin discovered it in some fungi. It is the substance which gives the flavour of meat to soups, and hence its name. It is a mixture of different crystallizable and uncrystallizable principles with empyreumatic products.

Osmazome has been prescribed in the dose of ʒss to ʒj, in broth or in powder, with aromatics, as a stimulant to the digestive actions and a restorer of appetite in convalescence. An *Osmazome choc'olate*, *Chocola'ta cum osmazo'ma*, has been introduced as an anæsthetic article of diet. It consists of chocolate lbj to osmazome ʒj.

OSME, (ὀσμη,) Odour.

OSMESIS, (ὀσμησις,) Olfaction.

OSMIDRO'SIS, (*osme*, and *ἰδρωσις*, 'sweat,') Per-spiration of an unusual odour.

OSMITOP'SIS ASTERISCOIDES, (from *osmē*, 'an agreeable odour,' (?) *Bel'lis*. A plant, of the Cape of Good Hope, which, from its smell and taste, seems to contain camphor; hence its anti-spasmodic virtues. In infusion, it is said to be beneficially employed in cough, hoarseness, and diseases of the chest generally, and is said to be serviceable in flatulent colic. Infused in spirit, *spir'itus bell'idis*, it has been used successfully as an external remedy in paralysis.

OSMOMETRICUS SENSUS, (*osme*, and *μετρον*, 'measure,') Olfaction.

OSMON'OSI, *Osmōnu'si*, *Mor'bi olfac'tūs*, (*osme*, and *νόσος*, 'a disease,') Diseases of olfaction.

OSMONOSOLOG'IA, (*osme*, *νόσος*, 'disease,' and *λόγος*, 'description,') The doctrine of, or a treatise on, the diseases of the sense of smell.

OSMORRHIZA LONGISTYLIS, (*osme*, and *ρίζα*, 'a root,') *Sweet Cic'ely*. An indigenous plant, Order, Umbelliferae, which flowers in May and June. It is an agreeable aromatic, having much of the flavour of anise.

OSMOSE, *Osmo'sis*, (ὀσμησις, 'impulse,') The force by which fluids are impelled through moist membrane and other porous septa, in endosmotic and exosmotic actions.

OSMOT'IC, *Osmot'icus*. Relating or belonging to osmose: as 'osmotic force,' the force which induces osmose.

OSMOZOME, Osmazome.

OSMUND ROYAL, *Osmunda regalis*.

OSMUN'DA CINNAMOM'EA, (from *Osmunder*, one of the names of Thor, a Scandinavian deity,) *Cin'namon Fern*. Indigenous; Order, Filices; is regarded to be demulcent, subastringent, and tonic. Boiled in milk, it yields a fine mucilage, which is useful in diarrhoea.

OSMUN'DA REGA'LIS, *O. spectab'ilis*, *Filix Flor'ida*, *Flow'ering Fern*, *Osmund-royal*. This plant was once thought to possess astringent and emmenagogue virtues.

OSMUNDA SPECTABILIS, *O. regalis*.

OSOROR, Opium.

OSPHRANTERIUS, (ὀσφραντήριος,) Olfactory.

OSPHRANTICOS, (ὀσφραντικός,) Olfactory.

OSPHRASIA, (ὀσφρασια,) Olfaction.

OSPHEREIOL'OGY, *Osphresiology'ia*, (*osphre-sis*, and *λόγος*, 'a discourse,') A treatise on olfac-tion and odours. *Osmol'ogy*, *Osmolog'ia*.

OSPHERESIS, (ὀσφρησις,) Olfaction.

OSPHRETICUS, (ὀσφρητικός,) Olfactory.

OSPHRISTICÆ, (from *osphresis*.) The doctrine of the phenomena of odours.

OSPHYALGEMA, gen. *Osphyalg'e'matis*, (*osphy's*, and *αλγῆμα*, 'pain,') Coxalgia.

OSPHYALGIA, (*osphy'algia*, from *osphy's*, and *αλγος*, 'pain,') Coxalgia.

OSPHYARTHROSIS, (*osphy's*, and *arthrosis*), Osphyritis.

OSPHYARTHROCE, (*osphy's*, and *arthroce'ae*), Coxarum morbus.

OSPHYITIS, *Osphy'itis*, *Osphyarthri'tis*, *Inflamma'tio cox'æ*, (*osphy's*, and *itis*.) Inflammation of the parts about the hip.

OSPHYRRHEUMA, (*osphy's*, and *ῥευμα*, 'rheumatism,') Lumbago.

OSPHY'S, (*osphy's*), Haunch, Lumbi.

OSPHYTIS, Osphyitis.

OSSA BREGMATIS, Parietal bones—o. *Convoluta inferiora*, Turbinated bones, inferior—o. *Digitorum manus*, Phalanges of the fingers—o. *Digitorum pedis*, Phalanges of the toes—o. *Epactalia*, Wormiana ossa—o. *Nasalia*, Nasal bones—o. *Nasi*, Nasal bones—o. *Nervalia*, Parietal bones—o. *Pisiformia lingualia*, see *Hyoides*, os—o. *Raphogeminantia*, Wormiana ossa—o. *Sesamoidæa*, Sesamoid bones—o. *Sincipitis*, Parietal bones—o. *Spongiosa*, Turbinated bones—o. *Suturarium*, Wormiana ossa—o. *Tertia maxillæ superioris*, Turbinated bones—o. *Tetragona*, Parietal bones—o. *Triangularia*, Sphenoidalia cornua—o. *Triangularia Blasii*, Wormiana ossa—o. *Triquetra*, Wormiana ossa—o. *Triticea*, see *Hyoides*, os—o. *Turbinata*, Turbinated bones—o. *Verticalia*, Parietal bones.

OSSEINE, (from *os*, 'a bone,') Ostein.

OSSELET, (dim. of *os*, 'a bone,') Ossiculum.

OSSELETS DE BERTIN, Sphenoidalia cornua—o. *de l'Oreille*, Ossicula auditûs—o. *de l'Ouïe*, Ossicula auditûs—o. *du Tympan*, Ossicula auditûs.

OS'SEOUS, *Os'se'ous*, *Osteo'des*, *Osto'des*, *Ostoï'des*, *Os'toïd*, (F.) *Osseux*, (*os*, 'a bone,') Bony. Resembling bone.

OSSEUX, Osseous.

OSSICLES, Ossicula, see Ossiculum.

OSSICULA, pl. of Ossiculum.

OSSICULA AUDITÛS, *O. Aur'is*, (F.) *Osselets de l'oreille ou de l'ouïe ou du tympan*. The small bones or ossicles situate in the cavity of the tympanum, and forming an uninterrupted chain from the membrane of the tympanum to that of the fenestra ovalis. They are four in number; and their series, from without to within, is as follows: *Malleus*, *Incus*, *Os orbiculare*, *Stapes*.

OSSICULA BERTINI, Sphenoidalia cornua—o. *Innominata*, *Innominata minora* ossa—o. *Nasi*, Nasal bones—o. *Sesamoidæa*, Sesamoid bones—o. *Triticea*, see *Hyoides*, os.

OSSICULUM, pl. *Ossic'ula*, *Osta'rion*, (F.) *Ossicule*, *Osselet*, (dim. of *os*, 'a bone,') A small bone.

OSSICULUM COCHLEARE, Orbicular bone—o. *Includi Comparatum*, *Incus*—o. *Malleolo assimilatam*, *Malleus*—o. *Molari denti comparatum*, *Incus*—o. *Orbiculare*, Orbicular bone—o. *Quantum*, Orbicular bone—o. *Squamosum*, Orbicular bone.

OSSIFICATION, *Ossific'a'tio*, *Ossifac'tio*, (*os*, 'a bone,' and *facere*, 'to make,') *Osteogen'ia*, *Osteogen'es'is*, *Osteo'sis*, *Osto'sis*. Formation of bone. Development or increase of the osseous system. Ossification takes place in the same manner as the nutrition of other parts. The bones are, at first, mucous, and afterwards cartilaginous; the cartilage at length receives the phosphate of lime, and is, at the same time, replaced by a gelatinous parenchyma, when the bone has acquired the whole of its development.

OSSIFICATION, POINTS OF, *Punc'ta seu Nu'clei ossificatio'nis*, *Nu'clei os'sei*, are the points where the ossification of a bone commences, whence it

extends to other parts. Almost all the bones present a greater or less number of points of ossification. Besides the natural ossification, which we observe in the fœtus and in the first periods of life, there are also *acciden'tal ossifications*, such as those frequently remarked, after the inflammation of serous membranes, in the parietes of arteries, and to which the terms *Osteo'thes'ia*, *Incrusta'tion*, and *Petrifac'tion* have sometimes been applied.

OSSIFRAGA, (*os*, 'a bone,' and *frangere*, 'to break,') Osteocolla.

OSSIS SACRI ACUMEN, ('point of the sacrum bone,') Coccyx.

OSSISANA, (*os*, 'a bone,' and *sanare*, 'to heal,') Osteocolla.

OSSIV'OROUS, (*os*, 'a bone,' and *vorare*, 'I devour,') A species of tumour, mentioned by Ruysch, which destroys a bone.

O'STAGRA, *Oste'agra*, (*osteon*, and *αγρα*, 'seizure,') A forceps to cut or remove portions of bone.

OSTALGIA, (*osteon*, and *αλγος*, 'pain,') Osteocopus.

OSTARIUM, (*οσταριον*, dim. of *οστέον*, 'a bone,') Ossiculum.

OSTARTHROSIS, Arthrophlogosis.

OSTARTHROCE, *Spina ventosa*.

OSTEAGRA, *Ostagra*.

OSTEALGIA, Osteocopus.

OSTEIDE, Osteoid.

OSTEIN, *Os'teine*, *Os'seine*, (from *osteon*.) The animal matter of bone.

OSTEITES, (from *osteon*), Osteocolla.

OSTETIS, *Osti'tis*, (*osteon*, and *itis*), *Inflamma'tio Os'is*. Inflammation of bone; characterized by violent shooting and lancinating pains in a bone.

OSTEMBRYON, (*osteon*, and *εμβρυον*, 'embryo,') Lithopædion.

OSTEMPY'SIS, *Osteëmpye'sis*, *Ostempyo'sis*, (*osteon*, and *εμπυσις* (*ev*, and *puv*), 'effusion of pus,') An abscess in the interior of a bone.

OSTEMPYOSIS, Osteomyelitis.

OSTEOANABROSIS, Osteonabrosis.

OSTEOCE'LE, (*osteon*, and *κηλη*, 'a rupture,')

A hernia in which the sac is cartilaginous and bony:—a rare case. Also, osseous induration of one or both testicles.

OSTEOCHONDROPHYTE, (*osteo*, *χονδρος*, 'a cartilage,' and *φυω*, 'to grow,') Chondroma.

OSTEOCLASIS, (*osteo*, and *κλασις*, 'a fracture,') Fracture.

OSTEOCOL'LA, (*osteon*, and *κόλλα*, 'glue,') *Gluebone*, *Stone or Bone binder*; *Ossif'raga*, *Hollos'teus*, *Ossisa'na*, *Ostei'tes*, *Amos'teus*, *Osteol'i'thos*, *Stelochi'tes*. A name given to petrified carbonate of lime, found in some parts of Germany and other portions of the globe; so called, because it has been supposed to possess the power of favouring the formation of *callus* in fractures. It is the *Chaux carbonatée concrétionnée incrustante* of Haüy. Not used.

OSTEOCOPUS, (*οστεοκοπος*), *Osteal'gia*, *Ostal'gia*, *Osteodyn'ia*, *Os'teoco'pe*, (*osteon*, and *κοπος*, 'fatigue,') *Boneache*, *Bone-ague*. Pain in the bones,—as in syphilis.

OSTEOCYST'OID, *Osteocystoï'des*, (*osteon*, *κυστις*, 'a sack or cyst,' and *ειδος*, 'appearance,') A morbid growth in bone, consisting of the formation of cysts filled with puriform or other fluid.

OSTEODENTINE, see Tooth.

OSTEODES, (from *osteon*), Osseous.

OSTEODYNIA, Osteocopus.

OSTEOANGRÆNA, Necrosis.

OSTEOGENIA, (*osteon*, and *γενω*, 'to beget,') Ossification.

OSTEOG'RAPHY, *Osteogra'phia*, (*osteon*, and *γραφειν*, 'to describe,') Description of the bones. The part of anatomy which describes the bones,

OSTEOHELICOSIS, (*osteon*, and *ἑλκος*, 'an ulcer,') Caries.

OSTEOID, *Osteoï'des*, (F.) *Ostéide*, (*osteon*, and *εἶδος*, 'appearance,') An accidental bony production, or more frequently a morbid production, which is a calcareous incrustation—(F.) *Concrétion ossiforme*—of the normal tissue, or of fibrous tumours.

OSTEOLITHOS, (*osteon*, and *λίθος*, 'a stone,') Osteocolla.

OSTEOLQGY *Osteolog'ia*, (*οστεολογία*, from *osteon*, and *λογος*, 'a discourse,') The part of anatomy which treats of bones.

OSTEOLYSIS, *Osteol'ysis*, (*osteon*, and *λυω*, 'I dissolve,') Necrosis or death of the bone, as in hospital gangrene, in which the dead parts are of soft consistence.

OSTEOMA, gen. *Osteo'matis*, (from *osteon*,) Exostosis.

OSTEOMALACIA, (*osteon*, and *μαλακία*, 'softness,') Mollities ossium—o. Infantum, Rachitis.

OSTEOMALACOSIS, Mollities ossium.

OSTEOMYELITIS, (*osteomyelon*, and *itis*,) Medullitis.

OSTEOMYELON, (*osteon*, and *μυελος*, 'marrow,') Marrow.

OSTEON, (*οστέον*,) Bone—o. Hieron, Sacrum.

OSTEONABROSIS, *Osteonabro'sis*, (*osteon*, a, priv., and *βρωσις*, 'nutriment,') Atrophy of the osseous texture marked by diminution of volume.

OSTEONCOSIS, (*osteon*, and *ογκος*, 'a tumour,') see Exostosis.

OSTEON'CUS, (*osteon*, and *ογκος*, 'a tumour,') A bony tumour.

OSTEONECROSIS, Necrosis.

OSTEON'OSI, *Osteon'osi*, (*osteon*, and *νοσος*, 'a disease,') *Mor'bi os'sium*. Diseases of the bones.

OSTEOPÆDION, (*osteon*, and *παιδίον*, 'a small child,') Lithopædion.

OSTEOPALINCLASIS, (*osteon*, *παλιν*, 'again,' and *κλασις*, 'fracture,') The breaking again of a bone which has united unfavourably.

OSTEOPHY'MA, gen. *Osteophy'matis*, (*osteon*, and *φύμα*, 'a swelling,') Osteoncus. Sometimes used in the sense of Exostosis. By Lobstein applied to a bony production originating in the deep-seated layers of the periosteum, in the vicinity of a carious bone, and which seems to be intended to obviate the weakness resulting from the destruction of the bone.

OSTEOPHYTE, (*osteon*, and *φύω*, 'to grow,') Exostosis, Osteophyton.

OSTEOPHYTON, *Os'teophyte*. Under this name, Rokitsky describes a deposition of bony matter, found by him on the inner surface of the parietes of the skull of pregnant women.

OSTEOPLASTIC, see Osteoplasty.

OSTEOPLASTY, (F.) *Ostéoplastie*, (*osteon*, and *πλασσειν*, 'to form,') An operation by which the total or partial loss of a bone is remedied. The filling of the cavity of a tooth with some appropriate substance; hence termed *Osteoplas'tic*.

OSTEOPLEU'RA, *Pleurasto'sis*, (*osteon*, and *πλευρον*, 'a rib,') Ossification of the cartilages of the ribs.

OSTEOPORO'SIS, *Osteosclero'sis*, (*osteon*, and *πώρασις*, 'induration,') Induration of a bone, from the deposition of too much bony matter. Also, a bony degeneration, which consists in too great porosity with hardening. [?]

OSTEOSATHYROSIS, (*osteon*, and *ἄσθυρος*, 'friable,') *Fragilitas ossium*.

OSTEOSARCO'MA, gen. *Osteosarco'matis*, *Osteosarco'sis*, *Sarcosto'sis*, *Hæmateoscto'sis*, (*osteon*, *σαρξ*, gen. *σαρκος*, 'flesh,' and *ομα*,) Disease of the bony tissue, which consists in softening of its laminae, and their transformation into a fleshy substance, analogous to that of cancer; accompanied by general symptoms of cancerous

affection. The word has often, also, been used synonymously with *spina ventosa*.

OSTEOSARCOSIS, (*osteon*, *σαρξ*, gen. *σαρκος*, 'flesh,' and *οσις*,) Osteosarcoma.

OSTEOSCLEROSIS, (*osteon*, and *σκληρος*, 'hard,') Osteoporosis.

OSTEOSIS, (from *osteon*,) Ossification.

OSTEOSPONGIO'MA, gen. *Osteospongio'matis*, *Osteospongio'sis*, (*osteon*, and *σπγγος*, 'a sponge or fungus,' and *ομα*,) A fungous condition of bone, *Spina ventosa*.

OSTEOSTEATO'MA, gen. *Osteosteato'matis*, *Exosto'sis steatomato'des*, (*osteon*, and *στέαρ*, gen. *στέατος*, 'suet or fat,') A name given to bony tumours or degenerations, which sometimes belong to osteosarcoma; at others, to *spina ventosa*, exostosis, &c.

OSTÉOTOME; same etymon as Osteotomy. A bone saw, which has the form of a chain. Also any bone saw, *Osteot'omus*, *Osteotom'ium*, *Osteotom'icum*.

OSTEOTOMICUM, Osteotome.

OSTEOTOMIST, *Osteotomis'ta*. An instrument, invented by Dr. D. Davis, for cutting the bones of the foetal cranium, where it becomes necessary to greatly reduce the size of the head in parturition.

OSTEOTOMIUM, Osteotome.

OSTEOTOMUS, Osteotome.

OSTEOT'OMY, *Osteotom'ia*, (*osteon*, and *τεμνειν*, 'to cut,' (*τομή*, 'an incision,')) The part of practical anatomy whose object is the dissection of bones.

OSTEOTOPHUS, Tophus.

OSTEOTYLUS, (*osteon*, and *τυλος*, 'callus,') Callus.

OSTEULCUM, (*οστέυλος*, from *osteon*, and *ελκειν*, 'to draw,') Bonenippers.

OSTEUM, (*οστέον*,) Bone.

OSTHELCUS, (*osteon*, and *ἑλκος*, 'an ulcer,') Caries.

OSTHEXIA, (*osteon*, and *ἔξις*, 'habit,') see Ossification.

OSTIARIUS, ('a doorkeeper,' from *ostium*, 'a door,') Pylorus.

OSTIOLUM, (dim. of *ostium*,) A small door or gate.

OSTIOLA CORDIS, *Valvula cordis*, *Pellieu'la cordis*. The valves of the heart.

OSTITIS, Osteitis.

OSTIUM, (from *os*, 'a mouth,') *Orific'ium*, a door or gate,—of the heart, for example. The opening—*Ostium a'trioventriculārē seu ventric'uli veno'sum*—between the auricle and ventricle of each side. *Ostium ventric'uli arterio'sum*, is the opening at the origin of the pulmonary artery and aorta.

OSTIUM ABDOMINALE, see Tube, Fallopian—o. Arteriosum, see Ostium—o. Atrioventriculare, see Ostium—o. Cutaneum, see Urethra—o. Duodenale ventriculi, Pylorus—o. Internum, see Ostium uteri—o. *Cephagæum*, Cardia.

OSTIUM U'TERI. The vulva; the vagina. The term *Ostium inter'num* has been applied to the inferior opening of the uterus, which establishes a free communication between the cavities of the body and neck, and which is often obliterated in old women. The neck of the uterus; the mouth of the uterus. See Uterus.

OSTIUM UTERINUM, see Tube, Fallopian—o. Vaginale urethrae, see Urethra—o. Vesicale, see Urethra—o. Venosum, see Ostium—o. Ventriculi Duodenale seu Pyloricum seu Dextrum seu Inferius, Pylorus.

OSTODES, (*οστώδης*, from *osteon*, and *οδης*,) Osseous.

OSTOIDEA SUBSTANTIA, see Tooth.

OSTOMA, (*osteon*, and *ομα*,) Exostosis.

OSTOSIS, (from *osteon*, and *osis*.) Ossification.

OSTREA, *Ostreum*, *Ostrea edulis*, (F.) *Huitre*. Class, Acephala. The oyster. The oyster is a wholesome article of diet, often advised as anaplectic and easy of digestion.

The shells, *Tes'tæ Ostreum*, *Tes'tæ* (Ph. U. S.), *Conchæ*, (F.) *Écailles d'huitres*, have been received into the Pharmacopœias, and used medicinally. Their virtues are similar to those of the carbonate of lime. See *Creta*.

OSTREUM, (ὀστρεον,) *Ostrea*.

OSTREION, (ὀστρειον,) *Ostrea*.

OSTRUTHIUM, *Imperatoria*.

OSWEGO TEA, *Monarda coccinea*.

OSYRIS, (ὀσυρίς,) *Antirrhinum linaria*.

OTACOSTIC, *Otacus'ticus*, (F.) *Otacus-tique*, (*oto*, and *ακουω*, 'I listen.') Any instrument is so called which improves the sense of hearing, —as the different species of trumpets.

OTAGRA, (ὠταγρα, from *oto*, and *αγρα*, 'a seizure,') *Otalgia*.

OTAL'GIA, (ὠταλγία,) *O'tagra*, *Otod'ymē*, *Otal'gy*, (F.) *Otalgie*, (*oto*, and *αλγος*, 'pain,') Pain in the ear; *Do'lor au'rium*, *Spas'mus au'rium*, *Ear-ache*. See *Otitis*.

OTAL'GIC, *Otal'gicus*, (F.) *Otalgique*. An epithet for remedies used in *otalgia*.

OTECUS, (*oto*, and *ηχος*, 'sound,') *Tinnitus aurium*.

OTEMPLAS'TRUM, *Otoplas'ta*, *Otoplas'trum*, (*oto*, and *εμπλαστρον*, 'a plaster,') A plaster put behind the ears.

OTEN'CHYTES, (ὠτεγκυτῆς,) *Oten'chyta*, (*oto*, *en*, 'into,' and *χωω*, 'I pour,') *Si'pho auricula'ris*. A species of syringe, used for injecting the ear.

OTHELICOSIS, (*oto*, and *ἐλκωσις*, 'ulceration,') Ulceration or suppuration of the ear.

OTHONE, (ὀθονη,) *Lintenum*.

OTHONION, (ὀθονιον,) *Lintenum*.

OTHYGROTES, (*oto*, and *ὕγροτης*, 'moisture,') *Otitirrhœa*.

OTIATER, (*oto*, and *ιατρος*, 'physician,') *Aurist*.

OTIATRI'A, *Otiat'ricæ*, *Otiatri'a*, (*oto*, and *ιατρεία*, 'medicine,') The business of the aurist. The management of diseases of the ear; *Aural Medicine and Surgery*, *Ear Surgery*.

OTIATRICE, *Otiatria*.

OTIATRUS, *Aurist*.

O'TIC, *O'ticus*, (F.) *Otique*, (from *ovs*, gen. *ωτος*, 'ear,') Belonging or relating to the ear. A medicine employed in diseases of the ear.

OTIC GAN'GLION, *Otogan'glium*, *Gan'glium auricula'rē* seu *oticum*, *Auric'ular gan'glium*, *Gan'glium of Ar'nold*, (F.) *Ganglion maxillo-tympanique*. This is a small ganglion, although more than double the size of the ophthalmic; ash-coloured and pulpy like the ganglions of the sympathetic. It is situate in advance of the ganglion of Gasser, on the lower surface of the inferior maxillary nerve, at the inner margin of the foramen ovale of the sphenoid bone. See *Petrus ganglion*.

OTIRRHŒ'A, *Otorrhœ'a*, *Othy'grotæ*, *Blenno-rhœ'a*, (F.) *Catarhe de l'Oreille*, (*oto*, and *ῥεω*, 'I flow,') Discharge of a purulent or puriform liquid, from the meatus auditorius externus, *Pyorrhœ'a au'rium*, *Otopyorrhœ'a*, *Pyoōtorrhœ'a*, *Otorrhœ'a purulen'ta*. It is a symptom of chronic otitis. Some authors, have, indeed, used the term synonymously with chronic otitis. When it extends to the brain, it is termed *cer'eb'al otorrhœ'a*; and when from the middle or internal ear, it is termed *Internal otorrhœa*, *Otorrhœ'a inter'na*, and *Entotorrhœ'a*.

OTIRRHŒA, CEREBRAL, see *Otitirrhœa*.

OTITES, (ὠτιτις, 'appertaining or relating to the ear,') see *Digitus*.

OTI'TIS, *Otiophleg'monē*, *Empres'ma oti'tis*, *Inflammatio au'ris*, *Inflammation of the ear*, (F.) *Oüte*, *Inflammation de l'oreille*, (*oto*, and *ἰτις*.)

Inflammation of the mucous membrane of the ear, characterized by excruciating pain; intolerable humming in the ear, with a discharge of mucus generally from the meatus externus or from the Eustachian tube. When the inflammation is restricted to the lining membrane of the meatus, it is termed *otitis catarrhalis*. Otitis is divided into *external* and *internal*, according as it affects the meatus auditorius, or the cavity of the tympanum or internal ear. It may be *acute* or *chronic*. *Acute otitis* commonly terminates in a few days in a favourable manner, with the aid of antiphlogistics. *Chronic otitis*, *otirr'hœ'a*, is almost always accompanied by a purulent discharge from the meatus auditorius. It is often incurable, and may ultimately give occasion to disease of the internal ear, and even of the brain.

OTITIS CATARRHALIS, see *Otitis*.

O'TIUM, (ὠτιον, dim. of *ovs*, gen. *ωτος*, 'ear,') *Oti'rium*. A small ear. The auricle or pavilion of the ear.

OTO, (from *ovs*, gen. *ωτος*, 'the ear,') in composition, ear.

OTOCEPH'ALUS, *Mono'tia*, (*oto*, and *κεφαλη*, 'head,') A monster, whose ears are in contact, or united into one.

OTOCLEI'SIS, (*oto*, and *κλειω*, 'I close,') Closure or obstruction of the ear.

OTOCONIA, (*oto*, and *κονία*, 'dust,') see *Otoliths*.

OTODYNE, (*oto*, and *δύνη*, 'pain,') *Otalgia*.

OTOGANGLIUM, *Otic ganglion*.

OTOLYPTIS, (ὠτολυψίς, from *oto*, and *λυψειν*, 'to hollow,') Earpick.

OTOLYPHUM, (ὠτολυφον,) Earpick.

OTORAPHY, *Otograf'phia*, (*oto*, and *γραφη*, 'a description,') The part of anatomy which describes the ear.

OTOIATRIA, *Otiatria*.

O'TOLITHS, *Otol'ithi*, *O'tolites*, (F.) *Otolithes*, (from *oto*, and *λίθος*, 'a stone,') White calcareous substances contained in the membranous vestibule, and seeming to be suspended in the fluid of the vestibule by means of a number of nervous filaments proceeding from the auditory nerve. Their universal presence in the ear of the mammalia would seem to show that they are inservient to audition. When the substance is of a looser consistence, it is called *Otocon'ia*, (F.) *Otoconie*, *Ear powder*, (from *oto*, and *conia*.)

OTOL'OGY, *Otolog'ia*, (*oto*, and *λογος*, 'a discourse,') The part of anatomy which treats of the ear. An anatomical treatise on the ear.

OTOPATHEMA, *Otopathy*.

OTOPATHY, *Otopath'i'a*, *Otopathe'ma*, (*oto*, and *πάθος*, 'affection,') A diseased condition of the ear.

OTOPHLEGMON, *Otitis*.

OTOPHONE, (*oto*, and *φωνη*, 'sound,') Ear trumpet.

OTOPLASTA, (*oto*, and *πλασσειν*, 'to cover over,') *Otemplastrum*.

OTOPLAS'TICŒ, *O'toplasty*, (*oto*, and *πλαστικός*, 'forming,') The operation for restoring a lost ear.

OTOPLASTRUM, *Otemplastrum*.

OTOPYORRHŒA, (*oto*, *πυον*, 'pus,' and *ῥεω*, 'to flow,') *Otitirrhœa*.

OTORRHAG'IA, (*oto*, and *ῥαγη*, 'rupture,') Bleeding from the ear or ears.

OTORRHEUMATIS'MUS, (*oto*, and *ρευματισμός*, 'rheumatism,') *Otal'gia rheumat'ica*. Rheumatic earache.

OTORRHŒA, *Otitirrhœa*—o. Cerebral, see *Otitirrhœa*—o. Interna, see *Otitirrhœa*—o. Purulenta, *Otitirrhœa*.

O'TOSCOPE, *Otoscop'ium*, (*oto*, and *σκοπη*, 'examination,') Any instrument for examining the condition of the ear. An elastic stethoscope, the

ends of which are tipped with ivory—one to be inserted into the meatus of the patient, and the other applied to the ear of the examiner has been so called; as well as one formed on the principle of the ophthalmoscope, laryngoscope, &c.

OTOT'OMY, *Ototom'ia*, (oto, and *τεμνειν*, 'to cut.') The part of practical anatomy which teaches the mode of dissecting and preparing the ear.

OTTO OF ROSES, see *Rosa centifolia*.

OTTOWAY DISEASE, *Mal de la Baie de Saint Paul*.

OUACHITA SPRINGS, see Arkansas, mineral waters of.

OUÏE, (from *audire*, 'to hear,') Audition.

OULA, (ουλα,) Gingivæ.

OULE, (ουλη,) Cicatrix.

OULITIS, (ουλα, and *itis*,) Ulitis.

OULORRHAGIA, (ουλα, and *ρην*, 'a breaking forth,') Ulorrhagia.

OUNCE, *Un'cia*. A weight equal to the 16th part of a pound avoirdupois, the 12th of a pound troy. It is composed of 8 drachms. See *Weights and Measures*.

OURAQUE, Urachus.

OURARI, Curare.

OUREMA, gen *Oure'matis*, (ουρημα,) Urine.

OURLES, (from *oreille*,) Palotitis, Cynanche parotidæa.

OUROLOGY, (ouron, and *λογος*, 'a discourse,') Uronology.

OURON, (ουρον,) Urine.

OUROSCOPY, (ouron, and *σκοπεω*, 'I examine,') see *Uromantia*.

OUS, gen. *O'tos*, (ous, gen. *ωτος*,) Ear.

OUTHAR, (ουθαρ,) see *Mamma*.

OUVERTURE CADAVÉRIQUE, Autopsia cadaverica.

OVA GRAAFIANA, Folliculi Graafiani—o. Nabothi, Nabothi glandulæ.

OVAIRE, Ovarium.

O'VAL, *Ova'lis*, *Oô'des*, *Oûi'des*, *Ovn'tus*, (F.) *Ovale*, *Océ*, (from *ovum*,) Egg-shaped; elliptic. That which is round and oblong like an egg.

VAL FORAMEN, *Fora'men ova'le*, *Foramen oû'des*, (F.) *Trou ovale*. The foramen obturatorium; also, the foramen in the inner paries of the middle ear, which opens into the vestibule; the foramen in the sphenoid bone, through which the third branch of the fifth pair issues from the cranium; and the aperture, in the fœtus, between the auricles of the heart. See *Botal Foramen*.

VAL FOS'SA, *Fossa ova'lis*, *Fo'vea ova'lis*, *Valvula* seu *Vestig'ium Foram'inis ova'lis*, is a depression observed in the right auricle of the heart, in the septum, which separates it from the left auricle. It has thick and strong edges, at its upper and fore part, called *Colum'næ foram'inis ova'lis*, *Isth'mus* seu *An'nulus Vieussæ'nii*, *An'nulus seu Lim'bus fo'veæ ova'lis*: in the fœtus called *An'nulus foram'inis seu fossæ ova'lis*. The fossa ovalis, in the adult, replaces the inter-auricular aperture in the septum auricularum of the fœtus.

VALBUMEN, Albumen ovi.

OVARIAN, *Ova'rial*, *Ovaria'nus*, (F.) *Ovarien*, *Ovarique*; (from *ovarium*,) Relating or belonging to the ovary—as *Ovarian pregnancy*.

OVARIAN ARTERY, *Spermatic artery*—o. Dropsy, Hydroo'rtion—o. Nerves, Spermatic plexus of nerves—o. Pregnancy, see *Pregnancy*, ovarian—o. Veins, Spermatic veins—o. Vesicles, Folliculi Graafiani.

OVARIOTOMY, (*ovarium*, and *τεμνειν*, 'to cut,') Oariotomy.

OVARIQUE, Ovarian.

OVARISTS, *O'vists*. Those physiologists who think that the phenomena of generation, in the human species and in every species of animal, result from the developmen't of the ova or ovula

of the female, merely incited by the male. Not many of the physiologists of the present day can be classed among the ovarists. The doctrine is called *O'varism* and *O'vism*.

OVARITIS, (*ovarium*, and *itis*,) Oaritis.

OVARIULE, (*ovarium*, and *ύλη*, 'cicatrix,') Corpus luteum.

OVARIUM, (*ωαριον*, dim. *ωον*, *ovum*,) The *Ovary*, *Eggbed*; *Testis muliebri* seu *femineus*, *Oa'rtion*, *Ooph'orum*, *Gynoa'rium*, *Vesicarium*, (F.) *Ovaire*. The ovaries are the organs in which the ova are formed in oviparous animals. By analogy, the name has been given to the organs which the ancients called the *testicles* of the female. They are two ovoid bodies, almost as large as the testicles in man, placed on each side of the uterus, between the Fallopian tube and round ligament, and in the substance of the broad ligament. Their outer extremity gives attachment to one of the projections of the fimbriated extremity of the Fallopian tube; and the internal is fixed to the uterus by a small ligamentous cord, called *Ligamentum rotundum ova'rii*, *Ligament of the Ovary*, *Mesoa'rium*. The ovaries are composed of a very close, spongy texture—*stroma*,—and of small vesicles—*Folliculi Graafia'ni*,—filled with a clear fluid: these vesicles contain ovules, which detach themselves from the ovary before and after fecundation, and are carried into the cavity of the uterus by the Fallopian tube.

OVARIUM NABOTHI, Nabothi glandulæ—o. Tumidum, Oarioncus.

OVATION, see *Parturition*.

OVATUS, (from *ovum*,) Oval.

OVÊ, Oval.

OVEREATING, COLIC FROM, Colica erapulo-sa.

OVERO BUTUA, Pareira brava.

OVICAPSULE, (*ovum*, and *capsule*,) see *Folliculi Graafiani*.

OVIDUCT, (*ovum*, and *ductus*, 'duct,') Tube, Fallopian.

OVIDUCTUS MULIEBRIS, Tuba Fallopiana.

OVIG'EROUS, *O'viger*, *Ovig'erus*; (*ovum*, and *gero*, 'I bear,') Containing or producing ova or eggs.

OVIPARITÉ, see *Oviparous*.

OVIPAROUS, *Ovip'arus*, (F.) *Ovipare*, (*ovum*, and *pario*, 'I bring forth,') An epithet applied to animals which are hatched from eggs outside of the body. The condition may be termed *Ovip'arousness*, (F.) *Oviparité*. See *Generation*.

OVIPAROUSNESS, see *Oviparous*.

O'VIPONT, (F.) *Oviponte*, ([L.] *ovum*, [F.] *œuf*, and *ponte*,) 'the laying of an egg.' A term applied to the escape of the ovum from the ovary; whilst *ovulation*, includes, also, the process of its maturation.

OVISAC, (*ovum*, and *sac*,) see *Folliculi Graafiani*—o. Tunic of the, see *Folliculi Graafiani*.

OVISM, see *Ovarists*.

OVISTS, Ovarists.

OVULOLOGY, (*ovum*, and *λογος*, 'a discourse,') Ologia.

OVOVIVIPAROUS, (F.) *Ovo-vivipare*, (*ovum*, *vivus*, 'living, and *pario*, 'I bring forth,') Oviparous animals, whose ova are hatched within the mother,—vipers, for example. See *Generation*.

OVULA (pl. of *Ovulum*) **GRAAFIANA**, 'ovules of De Graaf,' Folliculi Graafiani—o. Nabothi, Nabothi glandulæ.

O'VULAR, *Ovula'ris*; (from *ovulum*,) Relating or belonging to the ovule.

OVULATION. The formation of ova in the ovary, and the discharge of the same. Also, the development of the embryo from the egg. See *Ovipont*.

SPONTANEOUS OVULATION, (F.) *Ovulation spontané*, is the spontaneous formation and discharge

of ova which takes place in the female of the mammalia. See Heat, and Menses.

OVULE, Ovulum.

OVULINE, (from *ovulum*.) Decidua reflexa.

O'VULUM, *O'cule*, (dim. of *ovum*.) A small egg. That which has the appearance of a small egg. See Ovum.

O'VUM, *Oön*, (F.) *Ouf*, *Egg*. The egg of *Phasianus Gallus*, the common hen, is chiefly used as food. The different parts are, likewise, employed in medicine. The shell, *Auranum*, *Auran'tum*, (F.) *Coque* ou *Coquille d'œuf*, calcined, is esteemed an absorbent. The Oil of the egg is emollient, and is used, externally, to burns and chaps. The Yolk or Yelk of the egg—*Vitel'um* seu *Vitel'us* o'vi, *Lec'ithos*—renders oils, &c., miscible with water. Eggs, when lightboiled, are easy of digestion; but, when very hardboiled or fried, they are more rebellious.

Egg Brandy is made by taking of Brandy $\frac{1}{2}$ iv; Cinnamon water, $\frac{1}{2}$ iv; the yolks of two eggs; Sugar, $\frac{1}{2}$ ss; and Oil of cinnamon, gr. ij. Mix the yolks of the eggs first with the water, the oil and the sugar, stirring constantly. Then add the brandy little and little, until a smooth fluid is formed. It is an agreeable mode of giving brandy in adynamic states.

Egg'nog, is a drink made of eggs, rum or brandy, milk and sugar beaten up together.

In Pharmacy, white of egg, which consists chiefly of albumen, is used for clarifying syrups, &c. Yolk of egg, (F.) *Jaune d'œuf*, beaten up with warm water and sugar, and aromatized with orangeflower water, forms an emulsion, which is emollient, and known under the name *Lait de poule* or *chicken's milk*.

Anatomists give the name *Ova*, *O'vula*, *O'vules*, (F.) *Œufs*, to round vesicles, containing a humour similar to the yolk of egg, which are situate in the ovaries of the female, and, when fecundated, constitute the rudiments of the fœtus. During gestation, the embryo and its enveloping membranes retain the name *Ovum*, (F.) *Œuf*. The changes induced in the mammalia after impregnation greatly resemble those in the bird.

OVUM, Testicle—o. Hystericum, see Clavus hystericus.

OX, in composition, see Oxy.

O'XALAS SUBPOTASSICUS, Potass, oxalate cf.

O'XALATIC, *Oxalat'icus*. Relating to oxalates: hence *Oxalat'ic* or *Oxal'ic acid Diathesis*; the habit of body which favours the formation and deposition of oxalates from the urine.

O'XALIC AC'ID, (*oxalis*, 'sorrel,') *Ac'idum oxal'icum* seu *sacchar'icum* seu *hydrocarbo'nicum* seu *acetosel'æ* seu *oxal'icum* seu *sac'chari*, *Ac'id* of sugar or of sor'el, *Carbonous acid*. An acid found in sorrel. It crystallizes in long, colourless, quadrangular prisms, and is generally found, 'a nature, united with lime or potassa. It has been proposed, in a diluted state, as a refrigerant, but it is unnecessary; and, besides, in quantity, it is a virulently acrid poison. It was once largely employed for cleaning boottops; and, as it strongly resembles the sulphate of magnesia, has often been taken for it. Several fatal cases are, indeed, on record. For its antidotes, &c., see Poison.

O'XALIC ACID DIATHESIS, see Oxalatic.

O'XALIDE, *Oxalis acetosella*.

O'XALIS ACETOSA, gen. *Oxal'idis*, (from *oxus*, 'acid,' owing to the sharpness of its juice,) *Rumex acetosus*.

O'XALIS ACETOSEL'IA, *O. America'na*, *Ox'ys al'ba*. The systematic name of the Woodsorrel, *Cuckowbread*, *Sour Tref'le*, *White sor'el*, *Mountain Sor'el*, *Trifo'l'ium acetosum*, *Oxytriphyl'ium*, *Lul'ula*, *Allelu'ja*, *Acetosel'ia*, *Oxyphyl'ion*, *Oxytriphyl'ion*, *Pa'nis cuculi*, (F.) *Surelle*, *Oxalide*, *Pain a Coucou*, *Alleluia*. Ord. Oxalidaceæ. Sex.

Syst. Decandria Decagynia. This plant has a gratefully acid taste, and is used in salads. It has been esteemed refrigerant, antiscorbutic, and diuretic.

O'XALIS AMERICANA, *O. Acetosella*.

O'XALIS STRIC'TA, *Yel'low woodsorrel*; indigenous, has similar virtues to *O. Acetosella*.

O'XALIS TUBERO'SA, *O'ca*. A tuberous plant, whose ovalshaped root is used as food by the Peruvians. It is watery, has a sweetish taste, and is much liked.

O'XALUM, Potass, oxalate of.

O'XAL'ME, (*oxalum*, from *oxus*, and *âls*, 'salt,') A mixture of vinegar and salt.

O'XALURIA, Urine, oxalic.

O'XELE'ON, (*oxelaion*, from *oxy*, and *elaion*, 'oil,') A mixture of vinegar and oil, used as a condiment.

O'XÉOLÉ, (from *oxus*.) *Acétolè*.

O'X EYE, *Helenium autumnale*—o. Gall, Bile of the ox.

O'XICÉDRE, (*oxycedrus*;) *Juniperus oxycedrus*.

O'XID, Oxyd.

O'XIDATION, Oxydation.

O'XIDE, Oxyd.

O'XIDUM, Oxyd, see Oxydum.

O'XINES, (*oxinus*, [*oxus*], from *oxus*, 'sharp.' Wine already sour, but not yet vinegar.

O'XITRIPH'YLLUM, (*oxy*, and *τριφυλλον*, 'trefoil,') *Oxalis acetosella*.

O'XODE, Oxyd.

O'XODES, (*oxodys*;) Acidulous.

O'XOIDES, (from *oxy*, and *ειδος*, 'resemblance,') Acidulous.

O'XTER, Axilla.

O'XURE, Oxyd.

O'XUS, (*oxus*;) Aceticum acidum, *Ac'ium*.

O'XY, (from *oxus*, 'acute,') in composition, acute, quick, pointed, sharp, acid. In the composition of chemical terms, *oxy* and *ox* mean oxygen.

O'XYA, (*oxya*;) *Fagus sylvatica*.

O'XYACANTHA, (*oxycantha*, from *oxy*, and *ακανθα*, 'a thorn,') *Mespilus oxyacantha*.

O'XYCAN'THA GALE'NI, *Berberis*, *B. vulgaris*, *Sp'ina ac'id'a*, *Crespi'nus*, (F.) *Épine vinette*, *Vinettier*. Family, *Berberideæ*. Sex. *Syst.* Hexandria Monogynia. The *Barberry*. The fruit of the berries, *Barberries*, is gratefully acid, and moderately astringent. By preparing an alcoholic extract of the root *Berberis*, (Ph. U. S.) and adding water to it, a pulverulent brown substance is thrown down, which, when dissolved in alcohol and evaporated, yields the bitter principle, *Berberin*, which is an excellent stomachic in the dose of 2, to 10 grains.

The American *Barberry*, *Berberis Canadensis*, *Pip'peridge Bush*, *Sow'berry*, differs slightly from the European. It flourishes on mountains and hilly districts from Canada to Virginia.

O'XYÆSTHESIA, (*oxy*, and *æsthesia*;) Hyperæsthesia.

O'XYAP'HE, (*oxy*, and *αψη*, 'touch,') Hyperaphia.

O'XYAPHIA, Hyperaphia.

O'XYBAPHION, (*oxybaphion*, from *oxus*, 'acid,' or *oxos*, 'vinegar,' and *βαπτειν*, 'to dip,') see *Acetabulum*.

O'XYBLEP'SIA, (*oxyblepsia*, from *oxy*, and *βλεπω*, 'to see,') Acuteness of sight.

O'XYBOL'IA, (*oxy*, and *βολη*, 'ejection,') Too rapid emission of sperm.

O'XYCÉDRE, (*oxycedrus*;) *Juniperus oxycedrus*.

O'XYCEPHA'LIA, 'Sugar-loaf head,' (*oxy*, and *κεφαλη*, 'head,') Monstrosity in which the head is peaked like a sugar-loaf.

O'XYCOCCOS, (*oxycoccus*, from *oxys*, 'berry,') *Vaccinium oxycoccus*—o. *Macrocarpus*, see *Vaccinium oxycoccus*—o. *Palustris*, *Vaccinium oxycoccus*.

O'XYCRATE, *Oxyera'tum*, (*oxykra'ton* from *oxy*,

and *ωρα*, 'I mix.') *Pos'ea*, *Plus'ea*. A mixture of vinegar and water; frequently used in medicine as a refrigerant and antiseptic.

OXYCROC'EUM EMPLASTRUM, (*oxy*, and *κροκος*, 'saffron.') A plaster, composed of *saffron*, *pitch*, *colophony*, *yellow wax*, *turpentine*, *gallum*, *gum ammoniac*, *myrrh*, *olibanum*, and *mastic*. It was employed as a strengthening plaster.

OXYCUM, Oxygen.

OXYD, *Ox'id*, *Ox'ide*, *Ox'yde*, *Ox'ydim*, *Ox'idum*, (F.) *Oxyde*, *Oxide*, *Oxure*. A combination, not acid, of a simple body with oxygen.

OXYD, **CALCULOUS**, *Oxyd*, *cystic*.

OXYD, *Cryst'ic*, *Cal'culous* or *Ves'ical oxy'd*, *Cyst'ine*, *Neph'rine*, *Sorodod'mine*. A species of urinary calculus, having the shape of confused, yellowish, semitransparent, insipid, very hard crystals. When distilled, it furnishes subcarbonate of ammonia, like all nitrogenized matters. It dissolves in the nitric sulphuric, phosphoric, and oxalic acids.

OXYD, **VESICAL**, *Oxyd*, *cystic*.

OXYDA'TION, *Oxyda'tio*, *Oxida'tion*, *Oxygenat'ion*. The action of oxidizing a body; that is, of combining it with oxygen, a combination from which results an oxyd.

OXYDE, *Oxyd*.

OXYDE D'ANTIMOINE SULFURÉ, *Oxydum stibii sulphuratum* — o. *Gris ou noir de Mercure*, see *Hydrargyri oxydum cinereum*.

OXYDER'CIA, (*οξύεα*.) *Oxyder'cia*, *Vis'us a'cies*, (*oxy*, and *ἰσος*, 'I see.') Sharpness of vision.

OXYDER'CICUS, *Oxyder'ces*. A medicine which sharpens the sight.

OXYDORCIA, (*οξύδωρεια*.) *Oxydereia*.

OXYDUM, *Oxid* — o. *Antimonii cum phosphate calcis*, *Antimonial powder*.

OXYDUM FER'RI NI'GRI, *Fer'ri ox'idum ni'grum*, *Black oxy'd of iron*. The scales, which fall from iron, when heated, consist of iron combined with oxygen. They have been used like the chalybeates in general. The Edinburgh Pharmacopœia has a *Fer'ri ox'idum ni'grum*, made by precipitating, by ammonia, a solution of the mixed sulphate of protoxide and sesquioxide of iron. An analogous preparation is in the Dublin Pharmacopœia, under the name *Fer'ri oxy'dum magnet'icum*, *Magnet'ic oxy'de of iron*, *Martial E'thiops*.

OXYDUM FERRICUM CRYSTALLIZATUM NATIVUM, *Hæmatites* — o. *Hydrargyri completum*, *Hydrargyri nitrico-oxydum* — o. *Hydrargyricum*, *Hydrargyri nitrico-oxydum* — o. *Hydrargyricum præparatum*, *Hydrargyri oxydum cinereum* — o. *Hydrargyrosus*, *Hydrargyri oxydum cinereum*.

OXYDUM STIBII SULPHURATUM, *Hæ'par antimoni'i*, *Cro'cus metallo'rum seu antimoni'i*, (F.) *Oxyde d'antimoine sulfuré*, *Poic d'Antimoine*. This was formerly exhibited in the cure of fevers, but it is now rarely employed.

OXYDUM URICUM, *Uric oxide*.

OXYECOIA, (*οξύεωια*, from *oxy*, and *αων*, 'hearing,') *Hyperacusis*.

OXYG'ALA, (*οξύγαλα*, from *oxy*, and *γαλα*, 'milk,') *Sour milk*. The *Oxygala equinum*, *Curcæus mos*, is reckoned amongst the delicacies of the Tartars.

OXYG'ARON, (*οξύγαρον*, from *oxy*, and *γαρον*, 'garum,') A composition of garum and vinegar.

OXYGEN, *Oxygen'ium*, *Empy'real*, *Pure*, *Vital* or *dephlogisticated Air*, *Oxyg'enes*, *Ox'igenum*, *Princip'ium oxyg'ennans seu oxygenet'ium seu acidif'icans seu ac'idum*, *Elementum acidif'icum*, (F.) *Oxygène*, *Air du feu*; (*οξύς*, 'acid,' and *γενω*, 'I engender,' that is, *generator of acids*; and such it was believed to be exclusively, at the period when the name was given to it. This is now known not to be the case.) Oxygen is an elementary principle, largely distributed in nature. It exists in the air, in water, in seve-

ral acids, in all the oxyds, and in vegetable and animal substances. It is obtained by decomposing the peroxyd of manganese or the chlorate of potassa by heat in close vessels. Although oxygen, in the state of admixture in which it is found in the atmosphere, is of vital importance, it cannot be desired in a pure state with impunity. Animals die in it long before the whole of the oxygen is consumed. The properties of oxygen seem to be stimulant. It increases the force and velocity of the pulse, and has, accordingly, been used in cases of chronic debility, chlorosis, asthma, serofula, dropsy, paralysis, &c. It requires to be diluted with from 10 to 20 parts of atmospheric air; one to two quarts being used during the day.

OXYGENATED WATER, *Hydrogen*, *peroxide* of.

OXYGENATION, *Oxydation*.

OXYGÈNE, *Oxygen*.

OXYGÈNÈSES. A name given by Baumes to diseases which he attributed to disordered oxygenation.

OXYGEUSIA, (*oxy*, and *γεωσις*, 'taste,') *Hypergeusia*.

OXYGLICUS, *Oxyg'lyces*, (*οξύγλυκες*, from *oxy*, and *γλυκός*, 'sweet,') *Mul'sa ac'id'a*, *Acetum melleum dul'ce*. Prepared by macerating and boiling honeycomb (from which the greater part of the honey has been taken) in water, and adding vinegar. An *Oxymel*.

OXYHYDROCARBURETUM EX OLEO PYROXYLICO PARATUM, *Creasote*.

OXYLAPATHUM, (*oxy*, and *λαπάθος*, 'dock,') *Rumex acetos.*

OXYMEL, *Apom'eli*, *Mel aceta'tum*, (F.) *Acetometel*, (*οξύμελι*, from *oxy*, and *μελι*, 'honey,') *Honey* and *vinegar* boiled to a syrupy consistence. (*Mel. despumat.* Bv, *Acid. acetic.* ʒvij, *Aq. destillat.* fʒvij. Add the acid to the water and mix them with the honey previously heated.—Ph. L.) It is cooling;—externally detergent.

OXYMEL ERUGINIS, *Linimentum Eruginis*.

OXYMEL COLCHICI, *O. of mead'ow saffron*, (F.) *Oxymel colchique*. (*Rad. Colch. recent. in laminas tenues secta* ʒi, *Aceti destillati* Bj, *Mellis despumat.* pond. Bij. Macerate in a gentle heat for 48 hours. Press and boil the liquor with the honey to the thickness of a syrup; stirring with a wooden spoon. Ph. D.) It is expectorant and diuretic. Dose, fʒi, in gruel.

OXYMEL COLCHIQUE, *O. colchici* — o. of *Meadow saffron*, *O. colchici*.

OXYMEL SCILLÆ, *O. of squills*, *O. Scillicit'icum*, *Mel scill'æ*. (*Mellis despumat.* Oiss, *Aceti scillæ*, Oij. Boil in a glass vessel over a gentle fire to a proper consistence. Ph. U. S.) It is reputed to be expectorant and diuretic. Dose, fʒi to fʒij.

OXYMEL OF SCILLÆ, *O. scillæ*.

OXYMYRRHINE, (*oxy*, and *μύρρινη*, 'myrtle,') *Ruscus*.

OXYMYRSINE, (*οξύμυρσινη*.) *Ruscus*.

OXYNGIUM, *Adeps suillus*.

OXYNITRON, (*oxy*, and *νίτρον*, 'nitre,') Name of a plaster, composed chiefly of vinegar and nitre, and recommended by Aëtius.

OXYNOSEMA, gen. *Oxygnose'matis*, (*oxy*, and *νοσημα*, 'disease,') *Acute disease*.

OXYNOSOS, (*oxy*, and *νοσος*, 'disease,') *Acute disease*.

OXYNUSOS, (*oxy*, and *νοσος*, 'disease,') *Acute disease*.

OXYOPIA, *Galero'pia*, *Hyperop'ia*, *Suffu'sio accla'rana*, *A'cies visus*, (*oxy*, and *ωψ*, 'sight,') Excessive acuteness of the sense of sight. Instances have occurred where persons could see the stars in the day time. The cause is seated in great sensibility of the retina. The term has been used synonymously with *nyctalopia*.

OXYOSPHERESIA, *Oxyosphra'sia*, (oxy, and σφῆραι, 'smell.') Excessive acuteness of the sense of smell.

OXYPHLEGMA'SIA, οξύφλεγμασία, (from oxy, and φλεγμασία,) *Acu'ta et ve'hemens inflammatio*. A violent inflammation.

OXYPHENICON, (oxy, 'acid,' and φοινίκος, 'deep red.' [?]) *Tamarindus*.

OXYPHONIA, *Vox acu'ta*; (οξύφωνια, from oxy, and φωνή, 'voice;') *Parapho'nia clan'gens seu ulcero'sa*, *Clan'gor*, *Vox clango'sa*, *Leptopho'nia*. A shrill and squalling voice,—frequently, a symptom of morbid affection of the larynx.

OXYPHYLLON, (oxy, and φύλλον, 'a leaf') *Oxalis acetosella*.

OXYPODIA, (oxy, and πους, gen. ποδος, 'foot,') see *Kyllosis*.

OXYREG'MIA, (οξύρεγμα, from oxy, and ερεγγω, 'I belch.') *Rue'tus ac'idus*, Acid eructation.

OXYRIA RENIFORMIS, (οξύς, 'sharp, acid,') *Ru'mex dig'yinus*, *Bo'real sour'dock*, *Mountain Sor'rel*, *Welsh sor'rel*. A plant, *Ord. Polygana-cee*, *Sex. Syst. Diandria Digynia*, which grows in the northern parts of America and Europe; blossoming in the spring. It possesses the same properties as *Oxalis acetosella*.

OXYRRHOD'INON, *Acetum rosa'tum*, (oxy, and ῥόδον, 'a rose,') A composition of vinegar and roses. (*Ol. rosat. 3ij, acet. rosat. 3j.*) Used as a liniment in herpes and erysipelas.

OXYS, (οξύς,) Acid, Acute—o. Alba, *Oxalis acetosella*.

OXYSAC'CHARUM, (oxy, and σακχαρον, 'sugar,') A mixture of sugar and vinegar; termed *Oxysac'charum vomiti'vum*, when it held in solution the glass of antimony; and *Oxysac'charum scillit'icum*, when it contained squills.

OXYSAI DIAPHORETICUM AN'GELI SALÆ. A preparation of Angelo Sala. Acetate of potass exposed to the atmosphere; and consequently fallen into a state of deliquium.

OXYTARTARUS, *Potassæ acetas*.

OXYTES, (οξύτης,) Acidity.

OXYTOC'IA, (oxy, and τιτω, 'I bring forth,') Rapidity of parturition.

OXYTOC'IC, *Oxy'tocus*. A medicine which promotes delivery.

OXYTRIPHYLLON, (oxy, and τριφυλλον, 'trefoil,') *Oxalis acetosella*.

OXYURE VERMICULAIRE, *Ascaris vermicularis*.

OXYURIS VERMICULARIS, (oxy, and ουρα, 'a tail,') *Ascaris vermicularis*.

OYSTER GREEN, *Ulva latissima*—o. Plant, *Tragopogon*—o. Root, *Tragopogon*—o. Vegetable, *Tragopogon*.

OYSTER SHELLS, *Ostreæ testæ*—o. s. Prepared, *Testa præparata*.

OZOSTOMIA, (οζή, 'a stench,' and στομα, 'mouth,') Breath, offensive.

OZÆ'NA, (οζαίνα,) *Cory'za enton'ica seu purulen'ta seu ozæno'sa seu ulcero'sa seu virulen'ta seu malig'na*, *Pyorrhœ'a nas'al'is*, *Entozæ'na*, *Rhini'tis ulcero'sa*, *Ulcus na'r'ium fœ'tens*, (οζή, 'a fetid smell,') (F.) *Ozène*, *Punaisie*. An affection of the pituitary membrane, which gives occasion to a disagreeable odour similar to that of a crushed bed-bug; hence the name *Punais*, by which the French designate one labouring under ozæna. It is, sometimes, owing to caries of the bones; but is, perhaps, most frequently dependent upon syphilitic ulceration of the pituitary membrane, with or without caries of the bones of the nose.

OZÆNA BENIGNA, *Coryza*, chronic.

OZE, (οζή,) Breath, offensive.

OZEMUN, *Albumen*.

OZÈNE, *Ozæna*.

O'ZON, **O'ZONE**; (οζω, 'I smell of something,') The powerfully odorous matter produced when a current of ordinary electricity passes from pointed bodies into the air. It generally is presumed to be a peculiar modification of oxygen; and in varying quantity in the atmosphere is supposed to affect the health of man; but knowledge of this matter is exceedingly vague.

Air containing ozone is said to be ο'zonizèd, (F.) *Ozonisé*, *Ozoné*.

P.

P. A contraction of *Pugillus*.

P. Æ., A contraction of *Partes æquales*. } See Abbreviation.

P. P., A contraction of *Pulvis patrum*.

PAB'ULUM, *Tro'phê*, *Phor'bê*, *Alimen'tum*, *Nutrimen'tum*. Food, aliment. The animal heat and animal spirits—unmeaning terms—were formerly considered the *Pabulum vite*.

PACAY, *Prosopis dulcis*.

PACCHIONI, GLANDS or CORPUSCLES OF, *Pacchio'nian Bod'ies*, *Glandulæ Du'ræ Ma'tris seu du'ræ Menin'gis*, *Tuber'cula par'va seu Cor'pora glandifor'mia du'ræ ma'tris*, (F.) *Glandes de Pacchioni*, *Granulations cérébrales*, (after Pacchioni, an Italian anatomist.) Small, whitish, or yellowish bodies, sometimes separate, at others united like a bunch of grapes, which are observed in several parts of the dura and pia mater. They receive vessels, but apparently no nerves. Their texture and uses are unknown. A considerable number is observed in the longitudinal sinus, covered by the internal membrane.

PACHÆ'MIA, *Pachyæ'mia*, *Pachæ'ma*, *Pachyæ'ma*, (pachy, and αἷμα, 'blood,') Too great thickness of the blood.

PACHEABLEPHARO'SIS, *Pacheableph'ara*,

Pachybleph'aron, *Pachyblepharo'sis*, *Pach'ytes*, (pachy, and βλέφαρον, 'an eyelid,') A name given, by some authors, to a thickening of the eyelids, produced by tubercles or excrescences formed on their margins. It is the same affection as that which Sauvages designates *Coli'go à Pacheablepharo'si*.

PACHETOS, (παχῆτος,) see *Laqueus*.

PACHULOSIS, (παχυλός, 'thick,') *Pachylosis*.

PACHY, (παχύς, 'thick,') in composition, thick.

PACHYÆ'MIA, *Pachæmia*.

PACHYBLEPHARON, *Pacheablepharosis*.

PACHYCHOL'IA, (pachy, and χολή, 'bile,') Morbid thickness of the bile.

PACHYDERMATOCE'LE, (pachy, δερμα, gen. δερματος, 'skin,' and κηλη, 'a tumour,') A disease of the dermoid tissue, apparently congenital, which increases, at times from a small mole or spot to the size and appearance of a relaxed and very emaciated mamma.

PACHYDERMIA, *Elephantiasis*.

PACHYLO'SIS, *Pachulo'sis*, (παχυλός, 'thick,') An inordinate production of the epidermis dependent upon hypertrophy of the papillæ of the skin.

PACHYMENINGITIS, *Per'imeningi'tis*, (pa

chymeninx, 'dura mater,' and *itis*, 'inflammation.'). Inflammation of the dura mater.

PACHYMENINX, (*pachy*, and *μηνίς*, gen. *μηνίγος*, 'a membrane,') Dura mater.

PACHYNSIS, (*παχύσις*, from *παχύς*, 'to make thick,') Polysarcia adiposa.

PACHYNTICA, (*παχύντικα*,) Incrassantia.

PACHYSMUS, (*παχύσμος*, from *παχύς*, 'thick,') Polysarcia adiposa.

PACHYTES, (*παχύτης*, 'thickness,') Pacheeablepharosis.

PACINIAN CORPUSCLES, (after Pacini,) see Corpuscles.

PACKING, OF THE HYDROPATHISTS; *Émaillotage*.

PACKWAX, see Nucha.

PAD, ([S.] *Pajado*, from *paja*, 'straw' [?];) *Plumaceolus*, *Pulvil'lus*, *Pulvin'ulus*, *Par'vus pulvin'ar*;—a cushion, a little cushion; a junk;

(F.) *Coussinet*. A small bag, filled with feathers, wool, hair, bran, chaff, &c. It is used to compress or support parts. In the treatment of certain fractures, soft pads or cushions, called by the French *Remplissages*, are placed between the splints and the fractured limb, in order that the prominent parts may not be chafed by the splints.

PADDIE, see Oryza.

PADDY, see Oryza.

PADNOON, Bitnoben.

PAD'UA, MIN'ERAL WATERS OF. Near Padua, in Italy, there are several thermal sulphureous springs, the hottest of which is 188° Fah. The chief establishments are Abano, Battaglia, and Montegrotto.

PADUS, (*παδός*,) *Prunus padus*—P. Avium, *Prunus padus*—p. *Laurocerasus*, *Prunus laurocerasus*.

PÆDAN'CHONĒ, (*pædo*, and *αἴχω*, 'I strangle,') *Angi'na Sic'ca*. Name of a fatal species of cynanche, mentioned by M. A. Severinus; probably the *Cynanche Trachealis*, or *Cynanche Maligna*.

PÆDARTHROCEAE, (*pædo*, and *arthroceae*,) *Spina ventosa*.

PÆDATROPH'IA, *Atrophia infant'ilis*, *Marasmus lactant'ium*, (*pædo*, a, priv., and *τροφή*, 'nourishment,') Atrophy of children. *Tabes mesenterica*.

PÆDATROPHIA GLANDULOSA, *Scrofula*.

PÆDENCÉPHALIE, (*pædo*, and *εγκεφαλός*, 'encephalon,') *Hydrocephalus internus*.

PÆDERAST'IA, (*παιδερασία*,) *Pædast'ia*, *Cinnam'dia*, (F.) *Pédérastie*, (*pædo*, and *ἔρως*, 'love,') The love of boys. A disgraceful passion, common in ancient times, but now unknown. These terms are likewise applied to the crime of sodomy.

PÆDERAST'IC, *Pæderast'icus*. Relating or appertaining to pæderastia.

PÆDEROTA VIRGINICA, *Leptandra virginica*.

PÆD'IA, (*παιδεία*,) *Educat'io infant'um*. The bringing up or education—physical as well as moral—of children.

PÆDIATRI'A, *Pædiat'rica*, *Pædiat'rics*, (*pædo*, and *ιατρία*, 'medical management,') Treatment of the diseases of children.

PÆDICATIO, (*pædicare*, *pædicatum*, 'to commit pæderastia,') *Pæderastia*.

PÆDICI, *Pai'dici* (*morbi*,) Developmental affections, as malformations and teething.—(Dr. W. Farr.)

PÆDICTERUS, (*pædo*, and *icterus*, 'jaundice,') *Icterus infantum*.

PÆDION, (*παιδιον*,) *Infans*.

PÆDO, (*παις*, gen. *παῖδος*, 'a child,') in composition, child, boy.

PÆDOBAROMACROMETER, (*pædo*, *βαρος*, 'weight,')

μακρός, 'large,' and *μετρον*, 'measure,') *Baromacrometer*.

PÆDOMETER, (*pædo*, and *μετρον*, 'measure,') *Baromacrometer*.

PÆDON'OSUS, (*pædo*, and *νόσος*, 'disease,') A disease of childhood.

PÆDOPHLEBOTOM'IA, (from *pædo*, *φλέψ*, 'a vein,' and *τομή*, 'incision,') *Venæsectio puero'rum*. Bleeding performed on children.

PÆDOT'RIBES, *Min'ister Gymnas'tæ*, (*pædo*, and *τρίβω*, 'I train,') In ancient gymnastics, the officer acquainted with all the prescribed exercises, and who carried them into effect, according to the direction of the superior.

PÆDOT'ROPHY, *Pædotroph'ia*, (F.) *Pédotrophie*, (*pædo*, and *τροφή*, 'nourishment,') The part of hygiene which concerns the nourishment of children.

PÆEOSYNĒ, (*παιρσυνή*,) *Medicine*.

PÆON, (*Παιών*, 'the god of physic,') *Physician*.

PÆO'NIA, *P. officin'alis seu mas'cula seu nemora'lis seu fen'ina seu loba'ta*, *Hæmago'gum*, *Pe'ony*, or *Pi'ony*, male and female. (F.) *Pivoine*, *Pione*, *Herbe Sainte-Rose*, (from *Παιών*, 'the god of physic,') The root, flowers, and seeds were long esteemed powerful medicines. They are feebly astrigent, and have been considered to be narcotic and antispasmodic. They have been given, particularly, in intermittents and epilepsy, but are not now used. The flowers are called *Rosæ benedict'æ*, *R. re'gine*.

PÆONICE, (*παιωνική*, [*τεχνη*,] 'medical,') [*art.*] *Medicina*.

PÆONICUS, (*παιωνικός*,) *Medical*.

PÆONIS, (*παιωνίς*,) *Medicina*.

PÆONIUS, (*παιωνίος*,) *Medical*.

PAGAPOPLEX'IA, (*παγος*, 'icy coldness,' and *αποπληξία*, 'apoplexy,') Apoplexy from cold.

PAGOS, (*παγός*,) *Peritonæum*.

PAIDIA, (*παιδεία*,) *Infancy*.

PAIDICI, *Pædici*.

PAIDION, (*παιδιον*,) *Infans*.

PAIGIL, *Primula veris*.

PAIN, ([old F.] *painer*, 'to torment,') *Sax. piman*, 'to torture,') *Do'lor*, *Al'gos*, *Alge'ma*, *Alge'do*, *Alge'ia*, *Od'ynē*, *Ache*, *Aitch*, (Sc.) *Hæschæ*, (Prov.) *Crazy*, *Wark*; *Mis'ery*, (especially a constant annoying pain or suffering,) (F.) *Douleur*. A disagreeable sensation, which scarcely admits of definition. It is generally symptomatic, and is called *acute*, (F.) *aiguë*, when sharp and violent; when short and sudden a *twinge*, and, when more violent, a *pang*; *pungent*, (F.) *pungitive*, when it resembles that which would be produced by a sharp instrument run into the part: *heavy*, (F.) *gravative*, when attended with a sensation of weight; *tensive*, when the part seems distended: *lan'cinating*, when occurring in shoots: *lar'earing* or *tearing*, when the part seems to be tearing: *burning*, (F.) *brûlante*, when resembling that produced by a burn, &c.

PAIN, *SUN*, *Hemicrania*.

PAIN, (from *panis*, 'bread,') see *Triticum*—p. à *Coucou*, *Oxalis acetosella*—p. *de Madagascar*, *Jatropha manihot*—p. *de Porcean*, *Cyclamen*.

PAINFUL, *Odyno'des*, (F.) *Douloureux*. Full of pain.

PAINS, *La'bour pains*, *Throes*, *Throes*, *Parturient'is dolor'es*, *O'din*, *O'dis*, *Dolo'res ad par'tum*, *Tor'mina parturient'ium*, (F.) *Mal d'Enfant*, *Douleurs*. The uneasy sensations that accompany labor, and are owing to uterine contraction. The pains that precede actual delivery, and are simply employed in dilating the os uteri, are called *grinding*: those which extrude the child, *forcing*. Those which take place in the first days after delivery, and are employed in forcing away coagula, &c., are termed *Afterpains*, *Dolo'res seu*

Tor'mina post par'tum, D. puerpera'rum, (F.)
Tranchées utérines.

"To take a pain,"—"To try a pain,"—is to make an examination *per vaginam*, during labour, to discover its progress, character, &c.

The French term the pains, which precede and announce labour, *mouches*; those which accompany it, *douleurs*; and those which occur immediately afterwards, *tranchées ou coliques*.

PAINS, AFTER, see P. labour.

PAINS, GNAW'ING, *Dolo'res roden'tes*. Pains resembling what might be produced by the gnawing of an animal.

PAINT, (*pingere, pinctum*, 'to paint,' [L.] *pinto, (F.) peint*;) *Pigmentum, Fulcus, (F.) Fard*. A name given to different substances, employed for embellishing the complexion or rendering the skin softer. The substances most commonly used for this purpose are:—the *Subn'trate of Bis'muth, (F.) Blanc de fard*. A spirituous solution of benzoin, precipitated by water, and forming *Vir'gin's milk, (F.) Lait virginal*; the red or *rouge of the earthanus; Spanish white*; the oxides of lead, tin, and mercury, vermilion, red saunders, &c. All paints may be said to be noxious. They injure the skin; obstruct perspiration; and, thus, frequently lay the foundation for cutaneous affections. See Cosmetic.

PAINT, Indian, *Hydrastis Canadensis, Sanguinaria Canadensis*—p. Yellow, *Hydrastis Canadensis*.

PAIS, (*παῖς, gen. παῖδος*;) Boy.

PALAD, Vertex.

PALÆSTRA, (*παλαιστρα*, 'a place for wrestling,' from *παλν*, 'wrestling,') Gymnasium.

PALAIS, Palace.

PALAME, (*παλαμῶν*;) Palm.

PALATAL, Palatine.

PAL'ATE, *Palat'um, Hypero'a, Hypero'um, Uvulit'cus, Uvulus, Roof of the mouth, Fos'sa Palati'na, Palat'um stabile seu du'r'um seu os'seum, Hard Pal'ate, (F.) Palais*. The upper part of the cavity of the mouth; a kind of parabolic vault formed by the two superior maxillary bones and the two palate bones, covered with a thick and dense mucous membrane; bounded, before and at the sides, by the superior dental arch; and behind by the velum palati. A whitish line runs along the middle, from before to behind, at the anterior extremity of which is a tubercle corresponding to the inferior orifice of the anterior palatine canal. The arteries of the palate and corresponding gums proceed from the palatine, alveolar, infraorbital, labial, and buccal branches. The veins follow a similar course. The nerves are given off from the palatine, facial, infraorbital, superior dental, and from the nasopalatine ganglion.

PALATE ARTIFIC'IAL, *Obturator for the pal'ate, (F.) Obturateur du Palais*. A plate of tortoiseshell, silver, vulcanite, &c., used to close a deficiency of the palate.

PALATE BONE, *Os Palati seu basila'rē, (F.) Os Palatin'um du Palais*, is a small bone of an irregular shape, situate at the posterior part of the nasal fossæ and palate. It seems formed by the union of two bony plates joined at right angles to each other; so that one is inferior and horizontal;—the other, superior and vertical. The horizontal or palatine portion, Arch of the palate, is quadrilateral. Its upper surface forms part of the nasal fossæ; the lower forms part of the palatine arch, and contains the inferior orifice of the posterior palatine canal. Its anterior edge is articulated with the palatine process of the superior maxillary bone; the posterior is free and sharp. It gives attachment to the velum palati, and presents an eminence, which, when joined with one on the opposite side, forms the posterior nasal spine. Its inner edge is articu-

lated with the corresponding bone; its outer is confounded with the vertical portion. The vertical or ascending portion has the form of a long square. The inner surface forms part of the outer paries of the nasal fossæ. Its outer surface is unequal, and is articulated with the superior maxillary bone; behind, it presents a vertical groove, which forms part of the posterior palatine canal; and, above, a small free surface, which looks into the zygomatic fossa. Its anterior edge is united with the superior maxillary bone; the posterior is unequal, and rests on the internal ala of the pterygoid process. The angle, which it forms by uniting with the posterior edge of the horizontal portion, presents a very prominent pyramidal eminence, called the *tuberosity of the palate bone*. Its upper margin is surmounted by two eminences;—the one, anterior, called the *orbital process*;—the other, posterior, the *sphenoid process*. These two processes are separated by an almost circular notch, which the sphenoid bone forms into a foramen, called the *sphenopalatine*.

This bone appears to be developed from a single point of ossification. It is articulated with the sphenoid, ethmoid, the cornua sphenoidal, superior maxillary bone, the inferior spongy bone, the vomer, and its fellow.

PALATE, CLEFT, see Harelip—p. Falling down of the, Uvulitis—p. Fissured, see Harelip—p. Hard, Palate—p. Soft, Velum pendulum palati.

PAL'ATINE, *Pal'atal, Palati'num, Pal'atic, (from palatium)* Relating or belonging to the palate.

PAL'ATINE or PAL'ATAL ARTERIES are two in number. They are distinguished into, 1. The superior palatine or palatine, *Palati'na descendens* or *Pal'atomaxillary*, which leaves the internal maxillary behind the top of the orbit; descends into the pterygomaxillary fissure, and enters the posterior maxillary canal. Its branches are distributed to the velum palati and the nasal fossæ. 2. The inferior or ascending palatine or palatal, which is very small, and separates from the facial near its origin. It passes along the lateral and superior part of the pharynx, between the pillars of the velum palati, and divides into a number of ramusculi, which are distributed to the pharynx, the corresponding tonsil, the Eustachian tube, and the velum palati.

PALATINE CANALS or DUCTS. These are two in number. The anterior, *Ductus incisivus* seu *Naso'lo-palati'nis, Incisive canal*, is situate at the anterior part of the palatine arch, and is formed by the two superior maxillary bones. It is single beneath, but opens above by two apertures, one into each nasal fossa. The foramen is called *Fora'men incisivum* seu *Palati'num anterie'us* seu *Alveola'rē anterie'us* seu *cæ'cum os'sis maxilla'ris superio'ris*. The posterior palatine canal is situate at the posterior part of the palate. It is formed by the palate and superior maxillary bones, ascends between these two bones, and in its course gives rise to two or three small accessory palatine canals, which open above the tuberosity of the palate bone. It transmits the nerves and vessels which bear its name. The foramen is called *F. Palati'num poste'rius* seu *Palatomaxilla'rē* seu *alveola'rē poste'rius*.

PALATINE MEMBRANE is the fibro-mucous membrane that lines the palate or roof of the mouth.

PALATINE or PALATAL NERVES, *Gutturopalatins* (Ch.), are three in number, and arise from the inferior part of the sphenopalatine ganglion. 1. The great Palatine, *Palatin inférie'ur* (Ch.), is situate anteriorly to the two others, and descends into the posterior palatine canal; gives a nasal filament and a *staphyline* filament, and is distributed on the palate. 2. The middle palatine, *Gut'tural* (Ch.), which arises behind the preceding,

enters one of the accessory palatine canals, and is distributed to the tonsils and velum palati. 3. The *little palatine nerve*, situate still more backward than the last, which loses itself by several filaments in the uvula, tonsils, and follicles of the palatine membrane.

PALATITE, (*palatum*, and *itis*.) *Uranisconitis*.

PALATITIS, *Uranisconitis*.

PALATOGLOSSUS, *Glossostaphylinus*.

PALATOPHARYNGEUS, *Pharyngostaphylinus*, *Staphylinopharyngeus*, *Therapharyngostaphylinus*, part of the *Hyperpharyngeus*, (F.) *Palato-pharyngien*, *Stylo-pharyngien* (Ch.). A membranous muscle, broader at its middle than at its extremities, and situate vertically in the lateral paries of the pharynx and velum palati. It includes the peristaphylopharyngeus, pharyngostaphylinus, and thyrostaphylinus of Winslow. When the palatopharyngei contract together, they depress the velum. They elevate and shorten the pharynx, and act principally in deglutition.

PALATO-PHARYNGIEN, *Palatopharyngeus*—p. *Salpingeus*, *Circumflexus*—p. *Salpingien*, *Circumflexus*.

PALATOSTAPHYLINI, see *Azygos* muscle.

PALATUM, *Palate*—p. *Durum*, see *Palate*—p. *Fissum*, *Hyperoëchasma*, see *Harelip*—p. *Mobile*, *Velum pendulum palati*—p. *Molle*, *Velum pendulum palati*—p. *Osseum*, see *Palate*—p. *Pendulum*, *Velum pendulum palati*.

PALE, (*παλῆ*, 'dust.') *Pollen*.

PA'LEA CABOT'II. A fern of Java, the delicate filaments of which, derived from the stipes, act as mechanical styptics.

PALE'NESS, *Pall'or*, *Och'rotæ*, *Ochro'ma*, *Ochro'asia*, *Leuco'sia*, *Dentib'ria*, *Lu'ror*, (F.) *Pâleur*. Whiteness of complexion. It appears to be owing to a diminution or alteration of the blood circulating in the capillary vessels, and is often a sign of disease.

PALES COULEURS, ('pale colors.') *Chlorosis*.

PALET, (from [F.] *pelote*,—'itself,' from *pila*, 'a ball.') *Head*.

PALETTE (F.) 'a battle-dore.' *Pal'mula*, *Fer'ula*. An instrument of percussion. It has the shape of a spatula with a long handle; is made of very light, white wood, and has been proposed by M. Percy for what is called the *Massage*, a kind of pressure or kneading, exerted by the hand on the body and limbs, to excite the tone of the skin and subjacent tissues.

Palette (from *patella*) also means a cup for bleeding; and the xiphoid cartilage [?].

PALETTE DE CABANIS. An instrument, invented by Cabanis, of Geneva, to draw out the extremity of the stylet when passed into the nose, in the operation for fistula lacrymalis. It is composed of two plates of silver, pierced with several holes, and movable on each other. To use it, the instrument must be carried into the nose, so that the holes of the two plates correspond. When the end of the stylet is received into one of these holes, it is pinched, by sliding one of the plates on the other, and is easily withdrawn.

PALETTE À PANSEMENT. A small splint of thin, scooped wood, having the shape of the hand, and used to support the hand in certain cases,—when burnt, for example,—in order to prevent the fingers from being deformed by the cicatrices.

PÂLEUR, ([L.] *pallor*.) *Paleness*.

PALI PLAQUE, see *Plague*.

PALICOUR'EA CRO'CEA, *Psychot'ria crocea*, *Ordi'or*, *Cinchonaceæ*, a native of the West Indies, is possessed, according to Von Martius, of emetic properties.

PALICOUR'EA MARGGRAAT'II, *Erva do rato*, *Gal'ena'ia Velloz'ii*, a native of Brazil, is poisonous, and employed there to kill rats and mice.

PALICOUR'EA OFFICINA' LIS, a native of Brazil, is reported to be a powerful diuretic in small doses.

PALICOUR'EA SPECIO'SA, *Goldahrub*, a native of New Granada and Brazil. The leaves are considered to be antisyphilitic. The decoction in large doses is poisonous.

PALILLO, *Compomanesia lineatifolia*.

PALIMPISSA, (*παλιμπίσσα*, 'pitch reheated,' from *παλιν*, 'again,' and *πίσσα*, 'pitch.') *Pinus sylvestris*.

PALINCOTESIS, (*παλινκωσις*, from *παλιν*, 'again,' and *κωσις*, 'animosity,') *Recrudescencia*.

PALINDROME, *Palindromia*.

PALINDROM'IA, *Palin'drom'ia*, (*παλιν*, 'again,' and *δρομή*, 'a course,') This word signifies with some a reflux of fluids from the exterior towards the interior; and, according to others, the relapse of an affection—the recurrence—*Recur'sio*, *Repetit'io*,—of a paroxysm or disease,—*Mor'bus recidi'vus*, *Mor'bi recur'sus*.

PALINGENESIA, (*παλιγενεσία*, from *παλιν*, 'again,' and *γενεσις*, 'generation,') *Regeneration*.

PALLA, (*πάλλα*,) *Pila*.

PALLAT, *Vertex*.

PALLIATION, *Palliatio*, (*palliare*, *palliatum*, 'to cover, mask,' from *pallium*, 'a mantle,') The act of palliating; that is, of causing the symptoms of a disease to mitigate, without curing it. The medical art can frequently only moderate the symptoms of a disease, prevent it from making progress, prolong the days of the patient, and diminish his sufferings. This is *palliative treatment*, (F.) *Traitement palliatif*.

PALLIATIVE, *Palliati'vus*, *Antipath'ic*, *Evan'tiopath'ic*, (F.) *Palliati'f*. A remedy or mode of treatment which only relieves a disease, without curing it. The *palliative treatment*,—*Cu'ra palliati'va*, is placed in antithesis to the radical,—*Cu'ra radica'lis*: see *Radical*.

PALLIDULUS, (dim. of *pallidus*, 'pale,') *Blafard*.

PALLIDUS MORBUS, *Chlorosis*.

PALLOR, *Paleness*—p. *Virginum*, *Chlorosis*.

PALM, *Pal'ma*, *Pal'amé*, *Vo'la*, *V. ma'nús*, *Plat'e'a*, *The'mar*, *A'pes*, *Apul'tus*, *Poc'ulum*, *Pat'ora*, *Sup'plex* seu *Sup'plex'tile*, *Diag'n'osis*, (F.) *Pau'me* ou *Cœur de la main*. The hollow or inside of the hand. It is limited *without* by the thenar eminence, which answers to the thumb; *within*, by the hypothenar eminence, which answers to the little finger; *above*, by the wrist-joint; and *below* by the four fingers.

PALM, *GOM'UTO*, see *Toddy*.

PALMA, *Palm*—p. *Christi*, ('palm of Christ,')

Ricinus communis—p. *Cocos*, *Cocos nucifera*—p. *Dactylifera*, *Date tree*—p. *Humilis*, *Musa paradisiaca*—p. *Indica*, *vinaria* *secunda*, see *Toddy*—p. *Unguentarium*, *Myrobalanus*.

PALMÆ UTERI PLICATÆ, (pl. of *palmæ*, 'the branch of a tree,') *Ar'bor ut'ri uter'ina*, *A. ut'ri ric'ijana*. The arborescent striæ in the interior of the cervix uteri.

PALMAIRE, *Palmar*—p. *Cutané*, *Palmeris brevis*—p. *Grand*, *Palmaris magnus*—p. *Grêle*, *Palmaris longus*—p. *Long*, *Palmaris longus*.

PALMAR, *Palma'ris*, (F.) *Palmaire*, (from *palmæ*.) That which concerns the palm of the hand. The palm has sometimes been divided into three regions, called *palmar*. 1. The *external palmar region* corresponds to the thenar eminence. 2. The *internal palmar region* is formed by the hypothenar eminence; and 3. The *middle palmar region* is the hollow of the hand, (F.) *Cœur de la main*.

PALMAR APONEURO'SIS, *Palmar fac'ia*. A strong triangular aponeurosis, which arises from

the inferior tendon of the palmaris brevis and the anterior annular ligament of the carpus. It covers the whole palm of the hand, and adheres, somewhat strongly, to the skin of the part. The *palmar ligaments* are small, very numerous, fibrous fasciæ, which unite the different bones of the carpus and metacarpus.

PALMAR ARCH'ES, (F.) Arcades ou Croisces palmaires. The curved extremities of the radial and cubital arteries in the palm of the hand. There are two palmar arches:—the one, *cubital* or *superficial*,—*grande Arcade palmaire* of Chaussier, furnishes at its convexity towards the fingers the collateral arteries of the fingers, and the internal collateral of the thumb; the other, the *radial*, *profund*, or *deep-seated*,—*petite Arcade palmaire* of Chaussier, is deep in the palm of the hand beneath the tendons of the flexor muscles of the fingers. It gives some small branches only to the surrounding parts. The palmar veins accompany the corresponding arteries.

PALMAR ARTERIES, see *Palmar arches*—p. Fascia, P. aponeurosis.

PALMAR MUS'LES. These are three in number. 1. *Palmaris Magnus, Flexor carpi radialis, Radialis internus*, (F.) *Épitrochlo-métacarpien* (Ch.), *Palmaire grand, Radial antérieur*. A long fusiform muscle, which passes obliquely from the inner tuberosity of the humerus to the upper extremity of the second metacarpal bone. It bends the hand. 2. *Palmaris Longus, Latescens chor'dæ musculis, Ulnaris gracilis*, (F.) *Épitrochlo-carpi-palmaire* (Ch.), *Palmaire long, P. grêle*. This muscle is situate on the inner side of the last. It is very slender, and passes from the inner tuberosity of the humerus to the anterior annular ligament of the carpus, and to the palmar aponeurosis, which it seems to form by the vanishing of its fibres. It stretches the palmar aponeurosis, and bends the hand. 3. *Palmaris Brevis, Carpi quadrata, Carpiæus*, (F.) *Palmaire cutané*. A small, thin, quadrilateral muscle; situate before the thenar eminence. It passes from the annular ligament of the carpus to the integuments of the palm of the hand, which it wrinkles by its contraction.

PALMAR VEINS, see *Palmar arches*.

PALMARIS BREVIS, see *Palmar muscles*—p. Longus, see *Palmar muscles*—p. Magnus, see *Palmar muscles*.

PALMETTO, SAW, *Chamærops serrata*.

PALMI-PHALANGIENS, *Lumbricales manus*.

PALMISTRY, (from *palma*, 'the palm of the hand,') *Chiromancy*.

PALMOSCOPY or PALMOS'COPY, *Palmoscopia*, (παλμος, 'palpitation,' and σκοπεω, 'I view,') Investigation of the beating of the heart and pulse. Prognosis from the beating of the heart and pulse.

PALMULA, (dim. of *palma*,) *Date, Palette*.

PALMUS, (παλμος,) *Palpitation*—p. Cordis, *Palpitation*—p. Plumbarius, *Colic, metallic*—p. Vomitus, *Vomiting*.

PALO DE CALENTURAS, (S.) 'fever wood,' *Calenturas, Cinchona*—p. de Vacca, *Galactodendron utile*—p. de Velas, *Parmentiera cefrifer*.

PALOMMIER, *Gaultheria*.

PALPA'TION, (*palpare, palpation*,) 'to feel,') The sense of touch. It is also used for the mode of exploring diseases by feeling or pressing upon the diseased organ. A delicate and discriminating touch—*tactus eruditus*—is important in the diagnosis of many morbid conditions.

PALPEBRA, Palpebrum, (supposed to be from *palpitare*, 'to palpitate,' owing to its frequent motion. [?]) *Bleph'aron, Operculum seu Ooper'culum oculi, Eyelid*, (Prov.) *Eyebreck*,

(Sc.) *Eewinker, (F.) Paupière*. The eyelids are two movable vela, situate in front of the eye; which by their separation permit the organ to receive the impression of light, or by their greater or less occlusion protect it from the influence of too great light, or from the action of extraneous bodies. They are separated, above, from the forehead, by the eyebrows; and, below, are confounded with the cheeks. They are distinguished into *upper, Palpebra major seu superior, Epical'lis*; and *lower, Palpebra minor seu inferior, Hypocall'is, Hypocallum*. The former is much broader and more movable than the latter. They are separated by a transverse cleft—*Fissura palpebrarum*—and their edges are thick, and furnished with the cilia or eyelashes. Supported by the tarsal cartilages, they unite at the extremities of the transverse diameter of the orbit, forming two angles, the innermost being called the *greater angle of the eye, or inner or greater canthus*; the other, the *lesser, or external angle or canthus*. When brought together, a triangular space is left between their posterior surfaces and the conjunctiva, which has been called *Rivus lacrymalis*. They are formed, 1. Of a thin dermoid texture: 2. Of a muscular membrane,—a portion of the orbicularis: 3. Of a particular fibrous membrane: 4. Of the tarsi: 5. Of a mucous membrane,—a part of the tunica conjunctiva.

Their vessels and nerves are called *Palpebral*.

PALPEBRA INFERIOR EXTRORSUM FLEXA, *Ectropion*—p. Tertia, *Valvula semilunaris*.

PALPEBRÆ SUPERIORIS PRIMUS, ('first [muscle] of the upper eyelid,') *Levator palpebræ superioris*.

PALPEBRAL, Palpebralis, (F.) Palpébral, Blépharique, (from *palpebra*.) That which belongs or relates to the eyelid.

PALPEBRAL ARTERIES are those distributed to the eyelids. They are two in number; an *upper* and a *lower*. The former arises from the ophthalmic, a little beyond the cartilaginous pulley of the greater oblique muscle, and descends towards the upper eyelid, over which it is distributed. The latter arises at the side of, and a little behind, the preceding, and sends its ramifications to the lower eyelid and the neighbouring parts. The palpebral veins have a similar arrangement; except that some of them open into the temporal and labial veins.

PALPEBRAL FLUX, PU'RIFORM. A puriform discharge, furnished by the glands of Meibomius, when inflamed; and to which Scarpa attributes fistula lacrymalis in many cases.

PALPEBRAL LIG'AMENTS are distinguished into *superior* and *inferior*. Some anatomists, with Winslow, have described, under this name, the fibrous layer which is attached,—on the one hand, to the outer part of the circumference of the orbit; on the other, to the tarsal cartilages.

PALPEBRAL NERVES, Nerves of the eyelids, arise, particularly, from the frontal and lachrymal branches; from the ophthalmic nerve of Willis; and from the facial, nasal, and infraorbital branch of the superior maxillary nerve.

PALPEBRALIS, Orbicularis palpebrarum.

PALPEBRARUM APERIENS RECTUS, *Levator palpebræ superioris*—p. Duo musculi, *Orbiculares palpebrarum*.

PALPEBRATIO, (palpebrare, 'to wink,') *Nictation*.

PALPEBRUM, Palpebra.

PALPITATIO, Palpitation—p. Arteriarum, see *Palpitation*—p. Cordis *trepidans, Cardiotromus*.

PALPITATIO MEMBRORUM. A term used by Gaubius for involuntary shaking of the legs.

PALPITA'TION, Palpita'tio, Pal'mus, P. seu Tremor Cor'dis, Palpita'tio seu Palsa'tio Cor'dis,

Vibratio Cordiaca seu *Cor'dis*, *Cardiog'mus*, *Cardiopal'mus*, *Subul'tio*, *Clo'nus Palpita'tio*, (Sc.) *Dant*, *Donat*, (F.) *Palpitation*, *P. du Cœur*, (*pul'sare*, *pulpitatum*, 'to throb.') A name given to the pulsations of the heart, when they become stronger, more extensive, or more sensible to the individual than ordinarily. Palpitations occur in a number of affections. They are, sometimes, owing to organic diseases of the heart—hypertrophy, inflammation, &c.; at others, to a disease in some other viscus; whilst, frequently, they are merely nervous.

PALPITATION is sometimes felt in the arteries, constituting *Palpitatione Arterio'sa*.

PALPITATION DU CŒUR, Palpitation.

PALSIED, Paralytic.

PALSY, Paralysis—p. Bell's, Paralysis, Bell's—p. Histrionic, Paralysis, Bell's.

PALSY, LEAD, *Paral'ysis saturni'na* seu *metal'lica*, *Mul'dipar'esia*, (F.) *Paralysie saturnine*, *Metal'lic Palsy*, *Painters' Palsy*. The paralysis of the upper extremities especially, induced by the poison of lead. Palsy of the intestines, thus induced, is called, by Piorry, *Anetérnerrie saturnine*.

PALSY, METALLIC, see Palsy, lead—p. Painters', P. lead—p. Scribe, Cramp, writers'.

PALSY, SEMP'STRESSES. An affection analogous to writers' cramp, which usually consists in an entire or partial loss of sensation at the end of the right thumb and index finger, with inability to hold the needle.

PALSY, SHAKING, Paralysis agitans—p. Trembling, Paralysis agitans—p. Wasting, *Atrophie musculaire progressive*.

PALSY WORT, *Primula veris*.

PALTA, *Persea gatissima*.

PALUDAL, (*palus*, gen. *palu'dis*, 'a marsh,') Elodes.

PALUDAPIUM, ('marsh apium,') *Apium graveolens*.

PALUDÉEN, Elodes.

PALUDOSUS, Elodes.

PALUS, gen. *Palu'dis*, Marsh, Penis—p. Sanctus, *Guaiaecum*.

PALUSTER, (from *palus*, 'a marsh,') Elodes.

PALUSTRE, Elodes.

PAMBIOMA, gen. *Pambio'matis*, Panbioma.

PAMPOISON, (*pamer*, 'to faint,') Syncope.

PAMPINIFORM PLEXUS, (*pampinus*, 'a tendrill,') and *forma*, 'form,') Corpus pampiniforme.

PAMPINIFORME CORPUS, see Corpus pampiniforme.

PAMPLE'GIA, *Panple'gia*, (*pam*, and *πλεγω*, 'I strike,') General paralysis. Palsy of the whole body.

PAN, PANTO, (*pan*, neuter of *pas*, gen. *παντος*, 'all,') in composition, all.

PANACE'A, (*πανακεια*,) *Cathol'icon*, *Cuthol'icum*, (F.) *Panacée*, (*pan*, and *ακευαι*, 'I cure,') *Universal reme'dium*, *Panchres'ton*. A pretended universal remedy. There is no such thing in existence.

PANACEA DUCIS HOLSATIE, Potassæ sulphas—p. Duplicata, Potassæ sulphas—p. Lapsorum, *Arnica montana*—p. Mercurialis, Hydrargyri submurias—p. Mercurii rubri, Hydrargyri nitrico-oxidum—p. Pectoris, *Glechoma hederacea*—p. Vegetabilis, *Crocus*.

PANACÉE, Panacea—p. *Anglaise*, *Magnesiae carbonas*.

PANA'DA, (dim. of [L.] *pane*, [L.] *panis*, 'bread,') *Pana'do*, *Pana'ta*, *Panatel'la*. Bread, boiled in water to the consistence of pap; sometimes administered to the sick.

SIMPLE BREAD PANADA may be made by putting any quantity of grated stale bread into enough water to form a moderately thick pulp; covering it up, and leaving it to soak for an hour; then beating it up with two tablespoons

of milk, and a small portion of refined sugar, and boiling the whole for ten minutes, stirring all the time.

PANAIS, *Pastinaca sativa*.

PANARIS, Paronychia.

PANARTIUM, Paronychia.

PANARIUM, Paronychia.

PANATA, Panada.

PANATELLA, Panada.

PANAX, gen. *Pan'acis*, (*παναξ*, gen. *πανακος*, from *pan*, and *ακευαι*, 'to cure,') see *Panax quinquefolium*—p. *Costinum*, *Pastinaca opoponax*—p. *Heracleum*, *Pastinaca opoponax*—p. *Pastinacea*, *Pastinaca opoponax*.

PANAX QUINQUEFO'LIUM, *Aurelia'na* seu *Ar'elia Canadensis*, *Gin'seng*, *Nin'sing*, *Gon'sang*, *Sung*, *Red'berry*, *Five fingers*. (Same etymon as *Panacea*.) *Family*, *Araliaceæ*. *Sex. Syst.* *Polyandria Monœcia*. This plant grows in North America. The root—*Pa'na'x* (Ph. U. S.), *Ra'dix gin'seng*—is about the thickness of the little finger; an inch or two in length, often dividing into two branches; of a whitish-yellow colour; wrinkled on the surface, and of a compact, almost horny, texture. It has no smell, but a very sweet taste, combined with a slight degree of aromatic bitterness. It is much esteemed by the Chinese, who pay a high price for it. It is a gentle and agreeable stimulant.

PANAX SCHIN'SENG, *Asiat'ic Ginseng*, a native of Asia, has been usually confounded with the last. The root, *ra'dix nin'si*, has like virtues.

PANBIO'MA, *Pambio'ma*, gen. *Pambio'matis*, (*pan*, and *βιος*, 'life,') The original animating principle. The general principle of life.

PANCHRESTON, *Panchrestum*, (*πανχρηστος*, from *pan*, and *χρηστος*, 'useful,') *Panacea*.

PANCHRESTUM CHINÆ, see *Quinium*.

PANCHYMAGO'GUM, *Pantago'gon*, (*pan*, *χυμος*, 'juice,' and *αγω*, 'I drive away,') A medicine which the ancients believed capable of purging away all sorts of humours.

PANCHYMACOGUM MINERALE, *Hydrargyri submurias*.

PANCENONOSUS, (*pancenus*, and *νοσος*, 'disease,') *Pandemic*.

PANCENOSUS, *Pandemic*.

PANCENUS MORBUS, (*παγκυνος*, from *pan*, and *κυνος*, 'common,') *Pandemic*.

PANCENONUSUS, *Pandemic*.

PANCRATIUM, (*πανκρατιον*, from *pan*, and *κρατος*, 'strength,') An athletic exercise with the Greeks. Applied also to a plant, which was considered all powerful.

PANCREAS, gen. *Pancre'atis*, (*παγκρεας*, gen. *παγκρεατος*, from *pan*, and *κρεας*, 'flesh,') i. e. quite fleshy,') *Callio'reas*, *Pan'creum*, *Lac'tes*, *Pulvi'nar ventric'uli*, *To'tum carno'sum*, *Gland'ula sali'vallis abdom'inis*, *Pancre'nē*, *Sweet'bread*. A gland, deeply seated in the abdomen, and lying transversely on the vertebral column, between the three curvatures of the duodenum, behind the stomach, and to the right of the spleen. Its left extremity is called the *tail* of the pancreas; it is smaller than the right, which is called the *head*. At its right portion, it has, almost always, a greater or less prolongation; called, by Winslow, and some others, the *les'ser pan'creas*, (F.) *Petit pancre'as*, *Pancreas of Aselli*, *Pancreas mi'nus*. The pancreas resembles the salivary glands in structure, and has been called the *Abdom'inal Salivary Gland*. It is composed of lobes and granulated lobules, distinct, and united by areolar tissue. From each of the granulations of these lobes arise the radicles of its excretory duct, which are very delicate, and united like veins. The duct itself, *Duct'us Pancreat'icus* seu *Wirung'ia'nis*, *Pancreat'ic duct*, *Canal* or *Duct* of *Wir'sun* or *Wir'sungs*, (F.) *Canal pancréatique*, proceeds in a serpentine course through the substance of the organ: and when it has reached behind

the second portion of the duodenum, it becomes free, and is of the size of a crow's quill. It opens at an acute angle into the choledoch duct, or proceeds close to it, opening separately into the duodenum. The *Pancreatic arteries* are very numerous and small. They proceed from the cœliac, splenic, superior mesenteric, right gastroepiploic, coronaria ventriculi, and left capsular. Its *veins* open into the radicles of the vena porta; and, particularly, into the splenic and lesser mesenteric. Its *nerves* emanate from the solar plexus, and its lymphatics pass into ganglions, to which it gives its name. The pancreas secretes the *Pancreat'ic juice*, *Suc'cus* seu *Li'quor Pancreat'icus*, *Lym'pha Pancreat'is*, (F.) *Suc pancreatique*, which resembles the saliva. When this juice is mixed with amylaceous matters, it converts them into dextrin and glucose. Its great use appears, however, to be, to emulsify fatty matters by virtue of a peculiar albuminoid principle—*pan'-creatin*, coagulable by heat or alcohol—which it contains.

PANCREAS OF ASELLI, see Pancreas—p. Minus, see Pancreas—p. Petit, see Pancreas—p. Second, Brunner's glands.

PANCREATAGIA, (*pancreas*, and *αλγος*, 'pain.') Pain or disease of the pancreas.

PANCREATEMPHRAXIS, (*pancreas*, and *εμπράσσω*, 'I obstruct.') Obstruction of the pancreas.

PANCREAT'IC, *Pancreat'icus*, (F.) *Pancréatique*. Relating to or affecting the pancreas.

PANCREATIC ARTERIES, see Pancreas—p. Duct, see Pancreas—p. Juice, see Pancreas—p. Nerves, see Pancreas.

PANCREATIC SARCO'MA OF AB'ERNETHY, *Emphy'ma sarco'ma pancreat'icum*. A tumour, in irregular masses, connected by loose fibrous substance, like the irregular masses of the pancreas. Found in the areolar substance; but, more usually, in lymphatic glands, chiefly in the female breast.

PANCREATIC VEINS, see Pancreas.

PANCREATIN, see Pancreas.

PANCREATITIS, *Inflammat'io pancreat'is*, (F.) *Inflammation du Pancréas*, (*pancreas*, and *itis*.) Inflammation of the pancreas;—an affection, the phenomena of which are not well known.

PANCREATOID, (*pancreas*, and *εἶδος*, 'appearance.') *Pancreatoid'es*. A tumour resembling the pancreas in structure.

PANCREATON'CUS, (*pancreas*, and *ογκος*, 'a swelling.') Tumour of the pancreas.

PANCRENE, (*pan*, and *κρηνη*, 'a spring,') Pancreas.

PANCREUM, Pancreas.

PANDA'LEA, *Panda'leum*. A kind of lozenge used by the Arabian physicians.

PANDALEUM, Pandalea.

PANDALITIUM, (corrupted from *panaritium*), Paronychia.

PANDEMIA, (*πανδημία*), Pandemic.

PANDEMI'C, *Pandemy*, *Pandemia*, *Mor'bus Pandem'ius* seu *Pandem'ius* seu *Pancem'us*, *Pancem'osus*, *Pancænon'osus*, *Pancænonu'sus*, (F.) *Pandémie*, (*pan*, and *δῆμος*, 'people.') An epidemic which attacks the whole, or a great part, of the population.

PANDEMI'E, Pandemic.

PANDEMY, Pandemic.

PANDICULA'TION, *Pandiculatio*, *Scordin'es-ma*, *Scordinis'mus*, *Clo'nus Pandiculatio*, (*pandiculari*, (*pandere*, 'to stretch out,') 'to stretch, as one awaking from sleep.') *Stretch'ing*, *Hal'tices*, (pl.) An action, which consists in elevating the upper limbs, and throwing the head and trunk backwards, at the same time that the lower limbs are extended. This series of motions of the extensor muscles is in some measure in-

dependent of the will, and is accompanied by yawning, which is equally involuntary. In the state of health, stretching occurs before and after sleep; especially when we are fatigued. It is also observed at the onset of certain paroxysms of fever, hysteria, &c.

PANDOCE'UM, *Pandochi'um*, *Pandoci'um*, (*pan*, and *δέχομαι*, 'to receive.') A house or hotel that receives all classes. A general hospital, or one that admits all cases.

PANDOCHIUM, (*πανδοχεῖον*), Pandoceum.

PANDOCIUM, (*πανδοκίον*), Pandoceum.

PANG, (like Pain, or from Sax. *pingan*, 'to prick, to torture,') see Pain—p. Breast, suffocative, Angina pectoris.

PANHIDRO'SIS, (*pan*, and *ἰδρῶσις*, 'sweating,') *Sudor universa'lis*. Sweating over the whole body.

PANHYPERÆMIA, (*pan*, and *hyperæmia*), Plethora.

PANIC D'ITALIE, Panicum italicum.

PANIC GRASS, ITALIAN, Panicum italicum.

PANICAUT, *Eryngium campestre*—p. Maritime, *Eryngium maritimum*.

PANICUM DACTYLON, *Cynodon dactylon*.

PAN'ICUM ITAL'ICUM, (from *panis*, 'bread,') *Italian Pan'ic Grass*, *Indian Mil'let*, *Guin'ea Corn*, *Barbadoes Mil'let*, *Mil'ium Indicum*, *Sorghum*, *Hor'ba Sorghum*, *Durra*, (F.) *Millet des oiseaux* ou *à grappes*, *Panic d'Italie*. *Nat. Ord.* Gramineæ. The seed—*Mil'ii Sem'ina*—are much esteemed in Italy, being a constant ingredient in soups, and made into a variety of forms for the table.

PANICUM MILIA'CEUM, *P. mil'ium*, *Common Mil'let*, *Mil'ium*, *M. Panicum* seu *Esculent'um*, *Cen'chron*, *Cen'chrus*, (F.) *Millet ordinaire*, *Mil.* Milletseed are esteemed a nutritious article of diet, and are often made into puddings.

PANICUM MILIUM, *P. miliaceum*.

PANIDRO'SIS, (*pan*, and *ἰδρῶς*, 'sweat,') General or universal perspiration.

PANIS, ('bread,') see *Triticum æstivum*—p. Canicaceus, see Canicæ—p. Cibaricus seu secundarius, *Syncomistos*—p. Cuculi, *Oxalis acetosella*—p. Porcinus, *Cyclamen*.

PANIVOROUS, *Paniv'orus*, (*panis*, 'bread, and *voros*, 'I devour,') Subsisting on bread.

PANNA AFRICA'NA. A tæniæfuge root, introduced into Germany, and supposed to be the male fern.

PANNI, Swathing clothes.

PANNICLE, (dim. of *pannus*, 'a web,') Membrane.

PANNICULE, *Panniculus*—p. *Charnu*, *Panniculus carnosus*.

PANNICULUS, (dim. of *pannus*, 'a web,') (*P. Pannicule*. A membrane. Pterygion.

PANNICULUS ADIPOSUS, Cellular membrane.

PANNICULUS CARNO'SUS, *Tu'nica carno'sa*, *Mus'cular Web*, *Man'tle*, (F.) *Pannicule charnue*, is a general covering found in the quadruped, and formed by a thin, subcutaneous muscle, which serves to move the skin. It exists only in certain parts of the human body,—as in the forehead, where it is formed by the occipito-frontalis; in the neck, by the platysma myoides; and, in the scrotum, by the cremaster muscle.

PANNICULUS CORDIS, Pericardium—p. Hymenæus, Hymen—p. Virginalis, Hymen.

PANNIKELL, Cranium.

PAN'NUS. A piece of cloth. A web. This word has several other meanings, many of which are equivocal. It means a tent for a wound, and also an irregular nevus or mark on the skin. Modern ophthalmologists apply it to an opaque and vascular condition of the anterior membrane of the cornea, almost invariably produced by the continuance of purulent or strumous ophthalmia, or by chronic inflammation of the cornea. This

part becomes nebulous, covered with red vessels, and opaque. At times, the sclerotic conjunctiva undergoes a similar change to that which occurs in the corneal portion.

PANNUS HEPATICUS, Chloasma—p. Lenticularis, Epelides.

PANOPCHIA, Bubo.

PANOPHOBIA, Pantopho'bia, Panpho'bia, Athym'ia, Pa'vor, (F.) Frayeur nocturne, Souci, (from Παν, 'the god Pan,' and φόβος, 'fear,') A panic or fear inspired by the god Pan, according to the opinion of the ancients. Others derive it, and with more probability, from παν, 'all,' and φόβος, 'fear.' This universal and groundless fear is morbid, and is a symptom of melancholy.

PANOPHTHALMITIS, (pan, and ophthalmia.) Inflammation of the whole eye.

PANPHOBIA, Panophobia.

PANPLEGIA, Pamplegia.

PANSCLEROSIS, (pan, and σκληρωσις, 'hardness,') Complete induration of a part.

PANSE, (from παντεξ, pl. pantes,) Ingluvies.

PANSEMENT, Dressing.

PANSIES, Viola tricolor.

PANSPERMIA, (pan, and σπέρμα, 'sperm,') The theory of Dissemination of Germs, according to which, ova or germs are disseminated over all space, undergoing development under favourable circumstances.

PANSY, Viola tricolor.

PANTAGOGON, (panto, and αγω, 'to drive out,') Panchymagogen.

PANTANENCEPHALIA, (panto, and anencephalia.) Entire absence of brain—the same as anencephalia with I. G. St. Hilaire.

PANTATROPHIA, (panto, and τροφία, 'atrophy,') Universal atrophy.

PANTHODIC, Panthod'icus, (panto, and ὁδός, 'a way,') An epithet applied by Dr. Marshall Hall, to a course of nervous action proceeding from one point in all directions to every other.

PANTICES, Ingluvies, Intestines.

PANTING, ([F.] panteler, 'to pant,') Anhelatio.

PANTO, in composition, see Pan.

PANTOGLANGLITIS, (panto, ganglion, and ἰτις,) Cholera.

PANTOPHAGISTS, (panto, and φάγω, 'I eat,') see Omnivorous.

PANTOPHAGUS, Omnivorous.

PANTOPHOBIA, (panto, and φόβος, 'fear,') Hydrophobia, Panophobia.

PANTOZOOTIA, Panzootia.

PANUS, Phygethlon—p. Faucium, Stomatopanus—p. Inguinalis, Bubo.

PANZOOTIA, Pantozootia; (pan, and ζωον, 'animal,') An endemic or epidemic disease affecting animals generally.

PAP, (from papilla,) Mamma, Nipple. Also, a soft food made of bread boiled or softened with water, (Sc.) Breadberry.

PAP OF THE HASS, Uvula—p. of the Throat, Uvula.

PAPAS AMERICANUS, Solanum tuberosum.

PAPAVER, P. somniferum, Me'con, Lethe'a. The White Poppy, Papaver sativum, (F.) Pavot des Jardins. Ord. Papaveraceæ. Sec. Syst. Polyanthia Monogynia. The ripe CAPSULE OF THE WHITE POPPY, Cod'ia, Sca'phion, Papaveris Cap'sula, Papaver, (Ph. U. S.,) Me'con, Pop'pyhead, is used in fomentation as an anodyne and relaxant to inflamed or ulcerated parts. The expressed oil of the seeds—Poppy oil—is called by the French Huile d'Éillette. It is from the poppy that OPIUM—called, also, Opium Theba'icum, Succus Theba'icus, Meconium, Opion, Ma'nus De'i, Affion, Affium, Am'phiam, A'fium, Osoror, Lau'danum, Me'con, So'pio, (Prov.) Pop'py pill, (F.) Opium brut or cru—is procured. It is the concrete juice obtained after

the flowering of the plant, by making longitudinal incisions in the unripe capsules; and is commonly obtained from the Levant or India, where the poppy grows abundantly. Opium is a compound of morphia, narcotina, meconic acid, codein, narcotin, meconin, caoutchouc, fecula, resin, oil, and several other substances. Turkey Opium is of a heavy and slightly fetid odour. Its taste is nauseous, bitter, acrid, and warm: it is in flat pieces; solid and tenacious; of a dark brown colour, and yellowish when powdered; making a light brown, interrupted, streak on paper. In East India Opium the odour is the same, and empyreumatic; the taste is less bitter, but more nauseous, and the colour darker. The varieties of opium known in commerce are the Smyrna, the Constantinople, the Egyptian, the Persian, and the Indian. Opium is stimulant in small doses; sedative in large; narcotic and anodyne; operating through the nervous system. It is a most valuable remedy in all painful affections and inflammations, when given so as to produce its full, sedative effects, especially in peritonitis. In diarrhoea and dysentery it acts as an astringent; and in intermittents it will often prevent the paroxysm. In typhus, in small doses, it acts as a cordial; in larger it allays irritation and produces sleep. When applied externally, its narcotic effects are developed; and, hence, it is a useful application to irritable sores, painful tumours, &c. Dose, gr. $\frac{1}{2}$ to $\frac{1}{4}$ as a stimulant; gr. j to iij as a sedative.

PAPAVER CEREALE, P. rhœas—p. Corniculatum, Chelidonium majus—p. Erraticum, P. rhœas—p. Luteum, Chelidonium majus.

PAPAVER NIGRUM. This is merely a variety of the white poppy, producing black seeds.

PAPAVER RHÆAS. The systematic and pharmacopœial name of the Red corn'poppy, Corn' rose, Papaver erraticum seu cereal'e seu nigrum, Cop'per rose, Red-mailles, (F.) Coquelicot. The heads of this species contain a milky juice of a narcotic quality. A syrup, Syrupus Rhæados, of its flowers—Rhæas—has been directed, which has been esteemed diaphoretic and slightly soothing; but it is more admired for its color than for any other property.

PAPAVERACEUM, (from papaver,) Opiate.

PAPAVERCULUM, (from papaver, the inspissated juice of which it resembles,) Meconium.

PAPAVÉRINE, Papaver'ina, Papaverinum; (from papaver.) An alkaloid contained in opium; Codeine.

PAPAW TREE, Carica papaya.

PAPAYER, Carica papaya.

PAPER, BLISTERING, see Sparadrapum vesicatorium.

PAPIER ÉPISPASTIQUE, ('blistering paper,') see Sparadrapum vesicatorium—p. Médicamenté, see Sparadrapum—p. Vésicant, see Sparadrapum vesicatorium.

PAPILLA, pl. Papillæ, (F.) Papille. The end of the breast, the nipple. See Nipple. An eminence similar to a nipple. A name given to small eminences, more or less prominent, at the surface of several parts, particularly of the skin and mucous membranes, in which latter situation they are also called villi. See Villous. They contain the ultimate expansions of the vessels and nerves, and are susceptible, in some cases, of a kind of erection. They are also called Papillæ nervosæ seu Nervæ, P. Pyramida'les, Nervous papillæ, (F.) Papilles ou Papilles nerveuses, Houppes nerveuses.

PAPILLA LACHRYMALIS, see Lachrymal puncta—p. Mammæ, Nipple—p. Nervi optici, see Optic nerve—p. Pili, see Hair.

PAPILLÆ, see Corpuscles, Pacinian—p. Calyciformes, see Papillæ of the tongue—p. Capitata, see Papillæ of the tongue—p. Circumvallata, see

Papillæ of the tongue—p. Conical, see Papillæ of the tongue—p. Conoidal, see Papillæ of the tongue—p. Filiform, see Papillæ of the tongue—p. Fungiform, see Papillæ of the tongue—p. Fungoid, see Papillæ of the tongue.

PAPILLÆ OF THE KIDNEY, (F.) *Mamelons du rein*, are the projections into the pelvis of the kidney of the apices of the cones—*Pyramides Malpighii*, *Pyramides Malpighianæ*—of the tubular substance of the organ. They are also called *Processus mamillares*, *Caruncula papillares*, and *Papillæ renum*.

PAPILLÆ, LENTICULAR, see Papillæ of the tongue—p. Maximæ, see Papillæ of the tongue—p. Mediæ, see Papillæ of the tongue—p. Minimæ, see Papillæ of the tongue—p. Mucosæ, see Papillæ of the tongue—p. Nervæ, Corpuseles, Pacinian—p. Nervous, see Papilla—p. Pyramidal, see Papillæ of the tongue—p. Pyramidales, see Papilla—p. Renum, see Papillæ of the kidney—p. Semilenticulares, see Papillæ of the tongue.

PAPILLÆ OF THE TONGUE are very numerous and marked. They are observed particularly at the upper surface of the organ, and are distinguished into—1. The *Lenticular papillæ*, *Papillæ mucosæ* seu *maximæ* seu *capitatae* seu *vallatæ* seu *circumvallatæ* seu *truncatæ* seu *calyciformes*. They are arranged at the base of the tongue in two oblique lines, having the shape of the letter V, and united angularly at the foramen cæcum. They have, in general, an ovoid or spherical shape, and have seemed to be mucous follicles similar to those of the palate, lips, &c. 2. The *Fungiform* or *Fungoid Papillæ*, *Papillæ medise* seu *semilenticulares*, are of indeterminate number, but always more numerous than the preceding. They are irregularly disseminated near the edges and the point of the tongue; have a rounded and flattened head, supported by a short and narrow footstalk. Their tint is whitish. 3. The *Conical*, *Conoidal* or *Pyramidal papillæ*, *P. minimæ* seu *villosæ*, are the most numerous of all. Their arrangement is more regular in the forepart of the tongue than behind. They resemble small cones; the base being towards the tongue, and the apex free. The anterior are very small, and have been called *Filiform papillæ*. They appear to be produced by the terminations of the filaments of the lingual nerve, and are surrounded by an evident vascular network.

PAPILLÆ TRUNCATÆ, see Papillæ of the tongue—p. Vallatæ, see Papillæ of the tongue—p. Villosæ, see Papillæ of the tongue.

PAPILLARY, *Papillaris*, *Papillous*, (F.) *Papillaire*, (from *papilla*.) Of or belonging to the nipple or to the papilla.

PAPILLARY BODY, Corpus papillare.

PAPILLI-AUT-MAMILLI-FORMIS, Mastoid.

PAPILLIFORM, *Papilliformis*, (*papilla*, and *forma*, 'form.') Having the shape or appearance of a papilla.

PAPILLOMA, gen. *Papillo'matis*, (*papilla*, and *oma*.) A tumour, whose specific elements are those of the papillæ of the normal skin; and which, consequently, is composed of a stem of fibrous or homogeneous connective tissue; is club-shaped, cylindrical or acuminated, and covered with a layer of epithelial cells, similar to those of the normal papillæ.

PAPILLOUS, Papillary.

PAPOOSE ROOT, Caulophyllum thalioides.

PAPPE, Mamma.

PAPPEA CAPEINIS, (after Dr. Karl W. L. Pappel, a botanist of Leipzig.) A small tree of the *Nat. Ord.* Sapindaceæ, common in Cafferland, the fruit of which is called *Wild Plum*. It furnishes a vinous beverage and excellent vinegar. Its kernel contains an oil, which, although

eatable, is somewhat cathartic, and is recommended as an external application in tinea capitis, alopecia, &c.

PAPPUS, (παππος.) The down on the chin and cheeks.

PAPPUS AMERICANUS, *Solanum tuberosum*.

PAPULA, pl. *Papulae*. Akin to *papilla*. A *pimple*, *Eccor'mia*, (Prov.) *Poucke*, *Puckle*, (Sc.) *Pluke*, *Plouk*, (F.) *Papule*, *Bouton*, *Bourgeon*. A small acuminated elevation of the cuticle, with an inflamed base; very seldom containing a fluid, or suppurating, and commonly terminating in scurf or desquamation. An eruption of pimples is called in Devonshire *Murples*.

Papule is the first order of Willan and Bateman's arrangement of cutaneous diseases.

PAPULA AGRIA, *Lichen agrius*—p. Fera, *Herpex exedens*—p. Miliaris, *Miliary fever*—p. Typhosa, *Spots*, red.

PAPULÆ, *Lichen*—p. Siccæ, *Lichen*—p. Sudorales, *Sudamina*—p. Sudoris, *Miliary fever*.

PAPULAR, Papulose.

PAPULÆ, Papula.

PAPULEUX, Papulose.

PAPULOSE, *Papulosus*, *Papulous*, *Papularis*, *Papularis*, *Pimp'led*, *Papular*, (F.) *Papulæus*. Of or belonging to, resembling, or covered with papule or pimples.

PAQUERETTE VIVACE, *Bellis*.

PAR NERVORUM, *Syzygi nervorum*. A pair of nerves.

PAR NONUM, *Hypoglossus*—p. Opticum nervorum, *Optic nerve*—p. Primum nervorum cerebri, *Olfactory nerve*—p. Quartum, *Pathetic nerve*—p. Quintum nervorum cerebri, *Trigemini*—p. Secundum nervorum, *Optic nerve*—p. Septimum sive faciale, *Facial nerve*—p. Tertium, *Motores oculorum*—p. Trigemini, *Trigeminus*—p. Trium funicularum, *Trigeminus*—p. Vagum, *Pneumogastrie*—p. Visorium, *Optic nerve*.

PARA, (παρα.) A common prefix, 'by, near, contrary to, through,' (*per*); 'above and beyond,' (*ultra*); 'besides,' (*preter*). In certain terms, as in *Paracausis*, *Parapsis*, *Parabysma*, &c., used by Dr. Good, *para* means a defective or morbid condition.

PARA NUT, see *Pichurim beans*.

PARABALANUS, (παραβαλλειν, (from *para*, and βαλλειν, 'to cast,') 'to expose one-self.') A name given by the ancients to a person who devoted himself to the service of the sick in hospitals; and particularly during epidemics and the plague.

PARABYSMA, gen. *Parabys'matis*, (παραβυσμα, from *para*, and βυειν, 'to stuff.') Infarction or engorgement of an organ—as *Parabysma Lienis*, Engorgement of the spleen.

PARABYSMA MESENTERICUM, *Tabes mesenterica*.

PARABYS'TIA, Engorgement or infarction of an organ with blood.

PARACELSIS'TA, *Paracelsist*, *Paracelsian*, (after Paracelsus.) A follower of Paracelsus.

PARACELSISTICA MEDICINA, see *Spagyrist*s.

PARACENTE'SIS, (παρακέντησις, from παρακέντω, (*para*, and κέντω, 'I pierce,') 'I pierce through.') *Perforatio*, *Punctio*, *Punctura*, *Cenembat'esis*, *Cente'sis*, *Compunctio*, (F.) *Paracentese*. Puncturing. The operation of tapping to evacuate the collected fluid in ascites, ovarian dropsy, &c.

PARACENTESIS THORACIS, *Thoracocentesis*.

PARACENTETERION, (παρακέντητριον,) *Trocar*.

PARACENTETERIUM, (παρακέντητριον,) *Woolhouse* has given this appellation to a small trocar employed by Nuck for puncturing the eye in cases of dropsy of that organ.

PARACENTICION, Trocar.

PARACEPHALUS, (F.) *Paracéphale*, (from *para*, and *κεφαλή*, 'head.') A genus of monsters, with the head badly formed, but voluminous; the face distinct, with a mouth; rudimental sensorial organs, and thoracic members.

PARACHROEA, (*para*, and *χρῶς*, 'colour.') Morbid change of colour of the surface, especially of the face.

PARACHROMA, gen. *Parachromatis*, (*para*, and *χρῶμα*, 'colour.') Achromatopsia.

PARACHROSIS, Decoloration.

PARACINEMA, Luxation.

PARACINESES, (*para*, and *κίνησις*, 'motion.') Diseases of the motor nerves; morbid movements of voluntary muscles.

PARACMASIS, (*παράκμασις*,) Decline.

PARACMAS'TIC, *Paracmas'ticus*, (*παράκμαζω*, (*para*, and *αἰσθ*, 'the highest point,') 'I decrease,') Decreasing. An epithet for a distemper which gradually decreases. Also, according to Galen, the period of life when a person is said to grow old, which he places at from 35 to 49, after which he is said to be old—a definition that would not suit us at the present day.

PARACMASTICUS, (*παράκμαστικός*,) see Ac-masticus.

PARACME, (*παράκμη*,) Decline.

PARACOE, (*παράκοη*, from *para*, and *ακοή*, 'hearing,') See Cophosis, Paracusis.

PARAC'OPHE, (*παράκοπη*,) *Paraph'ora*, *Paracu'ois*, (*παράκοπτω*, (*para*, and *κοπτω*, 'I strike,') 'I strike falsely,') Delirium. In Hippocrates, it means the slight delirium which sometimes occurs in febrile diseases.

PARACOUSIA, (*para*, and *ακουω*, 'to hear,') See Cophosis.

PARAC'RISES, (*para*, and *κρίσις*, 'secretion,') Diseases of the secretory function.

PARACRUSIS, (*παράκρουσις*, from *para*, and *κρῶνω*, 'I strike,') Paracoepe.

PARACU'SIS, (*παράκουσις*, from *para*, and *ακουσις*, 'the act of hearing,') Cophosis. Also, perverted audition.

PARACUSIS A'CRIS, *Audi'tus a'crior*, *Paracu'sis oxyco'i'a*. The hearing painfully acute, and intolerant of the lowest sounds; a symptom in otalgia, epilepsy, cephalitis, and febrile affections.

PARACUSIS DUPLICATA, Double hearing—p. Imaginaria, Tinnitus aurium—p. Imperfecta, Double hearing, p. perversa—p. Obtusa, Barycoia—p. Oxycoia, P. acris.

PARACUSIS PERVERSA, P. *Willisia* na seu *imperfecta*, *Perverse* hearing. The ear only sensible to articulate sounds when excited by other and louder sounds intermixed with them.

PARACUSIS SURDITAS, see Cophosis—p. *Willisiana*, Paracusis perversa.

PARACYCLESSES, (*para*, and *κυκλῆσις*, (*κύκλος*, 'a circle,') 'circulation,') Diseases of the blood as regards its distribution and motion.

PARACYESIS, (*para*, and *κύησις*, 'pregnancy,') Pregnancy, morbid, Pregnancy, preternatural—p. Abortus, Abortion—p. Tubaria, Salpingocyesis.

PARACYNAN'CHE, (*para*, and *cynanche*,) Inflammation of the external muscles of the larynx. According to others, a slight cynanche.

PARÆSTHESIA, Paresthesia—p. Olfactoria, Parosmia.

PARÆSTHESIS, Paresthesia—p. Auditus flacca, Proptoma auricularum.

PARÆTHENAR, Parathenar.

PARAGEUS'IS, (*para*, and *γεύσις*, 'taste,') Ageusia. Also, perversion of taste, *Paragen'isia*, *Parageusia*, *Allotriogen'isia*, *Gus'tus deprava'tus*.

PARAGEUSTIA, Parageusis.

PARAGLOS'SA, *Paraglos'sē*, (*para*, and *γλῶσσα*, 'the tongue,') Swelling of the tongue; *Dysphu'*

gia Lingu'o'sa, *Glossomegis'tus*, *Macroglos'sa*, *Pro-lap'sus lin'gue*. Sauvages, under the term *Paraglossa deglutito'ria*, means a pretended doubling of the tongue into the pharynx, (F.) *Renversement de la langue*. See Glossocoele.

PARAGOGE, (*παράγωγη*, from *para*, and *αγω*, 'to lead,') Abduction, Coaptation.

PARAGOMPHO'SIS, *Paragomphosis cap'itis*, (*para*, and *γῶμφωσις*, 'a nailing,') Incomplete wedging of the head of the child in the pelvis during labour.

PARAGUAY-ROUX, see *Spilanthus oleracea*.

PARALALIA, (*para*, and *λαλία*, 'speech,') Mogilalia—p. Nasalis, Rhinophonia.

PARALAMP'SIS, (*παράλαμψις*, from *παράλαμψω*, (*para*, and *λαμψω*, 'I shine,') 'I shine at the side of,') *Mac'ula corneæ margarita'cea*, *Per'ila*. A cicatrix on the transparent part of the eye.

PARALEREMA, (*παράληρημα*, from *para*, and *ληρος*, 'foolish talk,') Delirium.

PARALLAGE, (*παράλλαγη*,) Insanity.

PARALLAGMA, gen. *Parallag'matis*, (*παράλλαγμα*,) *Chevauchement*.

PARALLAXIS, (*παράλλαξις*, 'aberration,') *Chevauchement*, Insanity.

PARALLE'LA. A kind of scurfy, cutaneous eruption, affecting the palms of the hands in parallel lines, and symptomatic of the venereal disease, according to Forestus.

PARALLELISM OF DISEASES, Isopathia.

PARALOGIA, (*παράλογια*, from *para*, and *λογος*, 'a discourse,') Delirium.

PARALOPH'IA, (*para*, and *λοφία*, 'the first vertebra of the back,') The lower and lateral part of the neck, near the vertebrae.

PARALYSE, Paralytic.

PARALYSIE, Paralysis—p. *Musculaire atrophique*, *Atrophie musculaire progressive*—p. *Sarturiana*, Palsy, lead.

PARALYSIS, (*παράλυσις*,) *Par'esis*, *Resolutio membro'rum seu nervo'rum*, *Ca'rus paraly'sis*, *Cata'lysis*, (F.) *Paraly'sie*, *Résolution des membres*; *Ane'u'ria*, *Anev'ria*, *Anerv'isme* of Piorry; *Palsy*, (*παράλυω*, (*para*, and *λυω*, 'I loosen,') 'I relax,') Abolition or great diminution of the voluntary motions, *Myoparaly'sis*, and sometimes of sensation in one or more parts of the body.

Palsy is said to be *local* when it affects only a few muscles. When it extends to half the body, vertically or horizontally, it is called *Hemiple'gia* or *Paraple'gia*. It is commonly symptomatic of disease of the brain or of the nerves themselves, and is a frequent accompaniment, precursor, or successor of apoplexy,—*Phrenoparaly'sis*. The immediate cause is, generally, pressure, either by blood effused, or by serum, or by vascular turgescence. At its onset, therefore, blood-letting may be advisable; but, afterwards, the disease becomes one of nervous debility, and requires stimulants to the paralyzed parts, or to the mucous membranes,—frictions, sinapisms, blisters, the moxa, baths, electricity, galvanism, purgatives, and the nux vomica, which last has the property of exciting movements where the will has no longer power. The disease generally admits only of palliation, and is extremely apt to recur.

PARALYSIS, *Primula veris*.

PARALYSIS AG'ITANS, P. *trem'ula*, *Syn'clonus ballis'mus*, *Scelotyr'bē festi'nans*, *Tre'mor*, *Tro'mos*, *Shak'ing Pal'sy*, *Trem'bling Pal'sy*. A variety of tremor, in which the muscles are in a perpetual alternation of contraction and relaxation.

PARALYSIS, BELL'S, *Bell's Pal'sy*, *Histrion'ic Paraly'sis* or *Pal'sy*, *Paraly'sis facia'lis*, *Proso-poraparaly'sis*, *Tris'timus diastroph'ic Sauvage'sti*, *Dis'troph'ic Gale'ni*, *Hemiple'gia facia'lis*, *O'ria tortu'ra paraly'tica Linne'i*, *Hemiprosopie'gia*. The paralysis of the face induced by a lesion of

the portio dura has been called Bell's Palsy, owing to Sir Charles Bell having been one of the first to investigate it minutely.

PARALYSIS BERIBERIA, Beriberi—p. Extremittatum, Acroparalysis.

PARALYSIS, DIPHTHERITIC, *P. diphtheritica*. A singular consequence, at times, of diphtheria, supervening when the patient is convalescing or apparently well. It may affect only the palate or pharynx, or be more general; and usually gets well, although slowly.

PARALYSIS FACIALIS, Paralysis, Bell's—p. General, of the Insane, Dementia paralytica—p. Hemiplegia, Hemiplegia—p. Histrionic, Paralysis, Bell's—p. of the Insane, Dementia paralytica—p. Intestini recti, Proctoparalysis—p. Intestinorum, Enteroparalysis—p. Linguae, Glossolysis—p. Medullaris, Paraplegia—p. Metallica, Palsy, lead—p. Muscular, progressive, *Atrophie musculaire progressive*—p. Nervi hypoglossi, Glossoplegia—p. Nervi vagi in parte thoracica, Asthma—p. Nervorum thoracis, &c., Stethoparalysis—p. from Pisons, P. Venenata—p. Rachialgia, Colic, metallic.

PARALYSIS, REFLEX. Palsy occasioned by irritation reflected through the excitomotory nervous system to parts in themselves healthy,—as in the case of palsy from dentition, or of an injury of a nerve on one side of the body occasioning palsy on the other. Reflex paralysis would seem, at times, to assume the paraplegic form—*reflex paraplegia*. The cure must depend on the removal of the primary cause.

PARALYSIS RENUM, Nephroparalysis—p. Saturnine, Paraplegia—p. Tremula, Paralysis agitans—p. Palsy, lead—p. Scorbutica, Pellagra—p. Spinalis, Paraplegia—p. Vacillans, Chorea.

PARALYSIS VENENATA, *Paralysis e venenis*, *Paralysis from poisons*. Paralysis, the result of poisonous substances—as of lead, mercury, arsenic, ergot, &c.

PARALYSIS E VENENIS, *P. Venenata*—p. Venetriculi, Gastroparalysis.

PARALYTIC, (*παρλυτικός*) *Paralyticus*, *Paralyzed*, *Pal'sied*, (F.) *Paralysé*, *Paralytique*. Affected with or causing paralysis.

PARALYTIC STROKE, see Stroke, paralytic.

PARALYTICUS, (*παρλυτικός*) *Paralytic*.

PARALYZED, *Paralytic*.

PARAME'NIA, (*para*, and *μην*, 'a month.') *Menstruatio anomala*, *Menses anomali*, *Mis-menstruation*. Difficult menstruation. Disordered menstruation.

PARAMENIA DIFFICILIS, Dysmenorrhœa—p. Erroris, Menstruation, vicarious—p. Obstructionis, see Amenorrhœa—p. Profusa, Menorrhagia—p. Superflua, Menorrhœa.

PARAME'RIA, (*παρμηρία*, *para*, and *μηρος*, 'the thigh.') The internal parts of the thigh.

PARAMESOS, (*παρμεσος*, from *para*, and *μεσος*, 'the middle.') Annular finger, see Digitus.

PARAMO DE RUIZ, MIN'ERAL WATERS OF. A remarkable acid spring in New Grenada. It contains free sulphuric and chlorohydric acids, and has a temperature of 157° Fah.

PARANEPH'RTIS, (*para*, and *nephritis*.) Inflammation of the renal capsules, *Inflammatio re'num eucenturiatorum*.

PARANEPHRUS, (*para*, and *νεφρος*, 'kidney.') Capsule, renal.

PARANCEA, (*παρανεια*, from *para*, and *νοος*, 'understanding.') Delirium, Dementia.

PARANYMPH'US, (*παρυνμφος*, *para*, and *νυμφη*, 'a young spouse, a bride.') A name given by the ancients to him who led the bride to the house of her husband. The term was adopted into the ancient schools of medicine to designate the usual discourse pronounced at the end of the prescribed period of study for licentiate, in which the qualifications and merits of the candidates were described.

PARAPATHIA, (*para*, and *παθος*, 'affection.') Pathomania.

PARAPECHYON, (*παρπαγχιον*, from *para*, and *πηχυς*, 'the elbow.') Radius.

PARAPHIA, (*para*, and *αφή*, 'touch.') Parapsis.

PARAPHIMO'SIS, (*παρφαμωσις*) *Periphimo'sis*, *Circumligatu'ra*, *Phimo'sis circumligata*, *Pe'nis lipoderm'ia*, (*para*, and *φίμωσις*, 'I bridle.') Strangulation of the glans penis, owing to the opening of the prepuce being too narrow. It frequently happens in persons affected with phimosis, when the prepuce has been pushed violently behind the base of the glans, during copulation, &c. In paraphimosis the narrow and resisting aperture of the prepuce forms, behind the corona glandis, a kind of ligature, which strangulates the part, causes it to swell and inflame, and even to become gangrenous. Attempts must be made to bring back the prepuce. These excite considerable pain, yet they must be persevered in for some time, and should they not succeed, the ring formed by the tumefied prepuce must be cut transversely, and the antiphlogistic treatment be adopted.

PARAPHOBIA, (*para*, and *φοβος*, 'dread.') Hydrophobia.

PARAPHON'IA, *παρφωνια*, from *para*, and *φωνη*, 'voice.') Affection of the voice, when its character or timbre is less agreeable.

PARAPHONIA CLANGENS, Oxyphonia—p. Gut-turalis, Asaphia—p. Nasalis, Rhinophonia—p. Puberum, Hirqitalitas—p. Rauca, Raucedo—p. Ulcerosa, Oxyphonia.

PARAPHORA, (*παρφορα*, from *para*, and *φέρω*, 'to carry,') Delirium, Paracope.

PARAPHRENESIS, (*para*, and *φρην*, 'the mind,') Paraphrenitis—p. Diaphragmatica, Diaphragmitis.

PARAPHRÉNIE, Insanity.

PARAPHREN'ITIS, (from *para*, *φρην*, 'the mind,' or *φρενις*, 'the diaphragm,' and *itis*,) *Paraphren'esis*, (F.) *Paraphrénésie*. A name given to inflammation of the diaphragm, which was supposed always to produce delirium:—as well as to the delirium itself accompanying that affection. See Phrenitis.

PARAPHROS'YNE, (*παρφροσυνη*) Delirium.

PARAPLAS'TIC, *Paraplasticus*, (*para*, and *πλαστικός*, 'formative.') Possessed of depraved formative powers,—as Carcinoma, Encephaloma, Melanosis, &c.

PARAPLE'GIA, (*παρπληγια*) *Paraplex'ia*, *Rachio'paral'ysis*, *Rhachio'paral'ysis*, *Rhachio'ple'gia*, *Myelo'paral'ysis*, *Paral'ysis spina'lis seu medulla'ris*, *Cæ'rus paral'ysis paraple'gia*, (F.) *Paraplégie*, (*para*, and *πλησσω*, 'I strike.') Palsy of the lower half of the body, including the bladder and rectum. It is, generally, owing to disease of some part of the cerebrospinal axis; especially of the spinal portion. The ancients sometimes used the term for incomplete apoplexy. Some authors have, also, made a distinction between *paraplegia* and *paraplexia*,—using the former for *partial apoplexy*,—the latter for *general or complete apoplexy*.

PARAPLEGIA, REFLEX, see Paralysis, reflex.

PARAPLEURITIS, Pleurodynia.

PARAPLEXIA, (*παρπληξια*) Paraplegia.

PARAPOPHYSES, see Vertebra.

PARAP'OP'LEXY, *Parapoplex'ia*, (*para*, and *αποπληξια*, 'apoplexy.') False apoplexy; a soporose state, resembling apoplexy; and especially that which occurs in the paroxysms of pernicious intermittents.

PARAP'SIS, *Paraph'ia*, (*para*, and *απτομαι*, 'I touch.') Sense of touch or general feeling vitiated or lost.

PARAP'SIS AC'RI'S TENERITUDO, Soreness—p. Ex.

pers, Torpor, Anæsthesia—p. Illusoria, Pseudæsthesia—p. Pruritus, Itching.

PARARMA, (παράρμα,) Fimbria.

PARARRHYTHMICUS, Pararrhythmus.

PARARRHYTHMUS, (παρὰρρυθμός,) Pararrhythmicus, (para, and ῥυθμός, 'rhythm.') An epithet for the pulse, when it has not the rhythm proper to the age and constitution of the individual.

PARATHRE'MA, gen. Parathre'matis, (παρθρέμμα,) Pararthro'ma, Pararthre'ois, Pararthro'ois, Parec'tasis, Parastre'ois, (para, and ἀρσπον, 'a joint,') Incomplete luxation; subluxation.

PARACEPAS'TRA, (para, and σκεπίζω, 'I cover,') A cap or bandage, which covers the whole head.

PARASCEUE, (παράσκευν, from para, and σκευν, 'equipment,') Apparatus.

PARASCHIDES, (παράσχιδες, from para, and σκίζω, 'I cleave,') Fragments or splinters of a broken bone.

PARASITAIRE, Parasitic.

PAR'ASITE, (παράσιτος,) Parasi'tus, (παράσιτω, (para, and σιτος, 'corn,') 'I eat by the side of,') Parasites are plants which attach themselves to other plants; and animals which live in or on the bodies of other animals,—so as to subsist at their expense. The mistletoe is a parasitic plant;—the louse a parasitic animal. Examples of parasitic animals are the *Pediculus humanus corporis*, *P. capitis*, *P. pubis*, *Pulex irritans*, *P. penicillatus*, the *Acarus Sca'biei*, and the different entozoa.

PARASITIA, (παράσιτια,) Parasitism.

PARASIT'IC, (παράσιτικός) Parasi'ticus, *Parasitoid*, *Parasitoides*. Having the nature of, or resembling a parasite, — as 'parasitic growths,' tubercles, cancer, monster, (F.) *Parasitaire*, &c.

PARASITIC DISEASES, *Parasit'ici* (morbi) are such as are induced by, or connected with animal or vegetable parasites. See *Porrigio favosa*, *Psora*, &c.

PARASITICIDÈ, (parasite, and cedere, 'to kill,') Antiparasitic.

PARASITISM, *Parasitism'us*, *Parasi'tia*. The condition of a parasite, or of an organized structure or body which lives on another organized body.

PARASITISMUS INTESTINALIS, Helminthiasis—p. *Superficieii*, *Malis*.

PARASITOG'ENY, *Parasitogen'ia*, *Parasitogen'esis*, (parasite, and γένω, 'to beget,') The corporeal condition, which favours the development of parasites.

PARASITOID, (parasite, and εἶδος, 'resemblance,') Parasitic.

PARASITUS, (παράσιτος,) Parasite.

PARASPA'DIA, (para, and σπῶω, 'I draw,') A preternatural opening of the urethra at the side of the penis. One whose urethra opens in this manner is called *Paraspadie'us* and *Paraspa'dias*.

PARASPADIÆUS, see *Paraspadia*.

PARASPHAGIS, (παράσφαγς, from para, and σφάγη, 'the throat,') The part of the neck contiguous to the clavicles.

PARAS'TATA, *Paras'tates*, (παρίστημι, (para, and ἵστημι, 'I stand near,') The prostate. In Hippocrates, it means the epididymis. The epididymes were called by Herophilus and Galen *Varico'æ parastate*, to distinguish them from *glandulo'æ parastate* or *prostate*. Rufus of Ephesus called the Tubæ Fallopiæ *Parastate varico'æ*.

PARASTATA ADENOIDES, Prostate—p. *Glandula*, *Prostate*—p. *Glandulosa*, *Prostate*.

PARASTATES, *Parastata*.

PARASTATY'TIS, *Epididymy'tis*; (parastata, 'epididymis,' and itis,) Inflammation of the epididymis.

PARASTREM'MA, gen. *Parastrem'matis*, *Parastrophê*, (παράστρεμμα, from παράστροφω, (para, and στροφή, 'I turn,') 'I distort or pervert,') Distortion. Convulsive distortion, especially of the face, the soft parts being drawn laterally.

PARASTREPSIS, *Parastrema*.

PARASTROPHE, *Parastrema*.

PARASYNANCHE, (παρσυναγχη,) *Cynanche parotidæa*.

PARASYSTOLE, (para, and συστολή, 'contraction,') *Acinesia*.

PARATH'ENAR, *Para'thenar*, (para, and ἔναρ, 'the sole of the foot,') Winslow called *grand parathénar*, *parathénar major*, a portion of the abductor of the little toe, *musculus scandula'rius*; and *petit parathénar*, *parathénar minor*, the short flexor of that toe.

PARATHENAR MAJOR, Abductor minimi digiti pedis—p. *Minor*, *Flexor brevis minimi digiti pedis*.

PARATONIA, (para, and τόνος, 'tension,') see *Spasm*.

PARATOPIA, (para, and τόπος, 'place,') *Luxation*.

PARATOP'IE. Diseases in which organs are removed from their proper position. See *Ectopia*.

PARATRIMMA, gen. *Paratrim'matis*, (παπατρίμμα, from para, and τριβω, 'to rub,') Chafing.

PARATROPE, (παράτροπη, from para, and τρεπω, 'to turn,') *Apotrope*.

PARAT'ROPHY, *Paratrophê*, *Paratroph'ia*, (para, and τροφή, 'nourishment,') *Misnutrition*.

PARBREAKING, ([G.] brechen, 'to vomit,') *Vomiting*.

PARDALIANCHES, (παρδαλιανχες, from παρδαλις, 'a panther, and αἰσχιν, 'to strangle,') *Aconitum*.

PARDALLANCHUM, *Aconitum*.

PARC'TAMA, gen. *Parc'tamatis*, (παρκεταμα,) *Parc'tasis*, (para, εκ, 'out, and ἔεινω, 'I stretch,') *Immoderate extension, distension, or dilatation of a part—as of the heart*.

PARÆTASIS, (παρεκτασις,) *Parathrema*, *Parcetama*.

PAREGOR'IC, (παρηγορικός,) *Paregor'icus*, *Solipiens*, (F.) *Parégorique*, (παρηγορεω (para, and αγορεω), 'I soothe, I calm,') That which soothes or assuages—as *paregoric elixir*. An anodyne.

PAREIA, (παρεία,) *Gena*.

PAREIRA, see *P. brava*.

PAREIRA BRA'VA, *Cissampelos Pareira*, *Purey'ra*, *Bu'tua*, *Abu'ta*, *Albu'tua*, *Boton*, *Caa-peba*, *Overo butua*, *Wild Vine*, *Velvet leaf*. Family, *Menispermæ*. *See. Syst. Dioecia Monadelphia*. The root of this plant—*Pareir'a* (Ph. U. S.)—a native of South America and the West Indies, has a sweet taste, with considerable bitterness and slight roughness. It has been recommended in nephritis and calculous affections; but it is not much used. By A. de St. Hilaire, *Pareira* is assigned to *Cissampelos glaber'ima*.

PARELLE, *Rumex hydrolapathum*, *R. patiens*.

PAREMPTOSIS, (παρεμπτωσις, 'intrusion,') see *Coincidental*.

PARENCEPHALIS, (παρεγκεφαλις, from παρα, and εγκέφαλος, 'the brain,') *Cerebellum*.

PARENCEPHALITIS, (parencephalis, and itis,) *Cerebellitis*.

PARENCEPHALOC'E'LÈ, (parencephalis, and κλη, 'a rupture,') *Her'nia of the cerebel'lum*; a very rare disease. It occurs under the form of a soft, indolent, or slightly painful tumour, without change in the colour of the skin, and is situate in the occipital region. It is constituted of the cerebellum, which escapes through an aperture in the occipital bone. Great attention must be paid not to mistake it for an ordinary tumour, as its extirpation would be attended with fatal consequences.

PARENCEPHALUS, *Cerebellum*.

PARENCHYMA, gen. *Parenchymat'is*, (*παρυσχυμα*.) *Cu'ra parenchymat'ica seu viscerum seu viscerat'is seu viscerosa*, *Substantia parenchymat'osa*, *Prochem'ma*, (F.) *Parenchyme*. The texture of glandular and other organs, composed of agglomerated globules united by areolar tissue, and tearing with more or less facility. Such is the texture of the liver, kidneys, &c. *Παρυσχυμα* signifies, in Greek, effusion, (*para*, *α*, and *χω*, 'I pour,') and it was believed, that this tissue consisted of effused blood or other fluid.

PARENCHYMA TESTIS, *Pulpa testis*.

PARENCHYMAL, *Parenchymatous*, *Parenchymous*, *Parenchymat'is*, *Parenchymat'icus*, (F.) *Parenchymateux*. Belonging or relating to the parenchyma of an organ.

PARENCHYMATITIS, (*parenchyma*, and *itis*.) *Inflammat'io parenchymat'ica*. Inflammation of the substance of an organ.

PARENCHYMATOUS, *Parenchymal*.

PAR'ESIS, (*παρεσις*, from *παρηνμι*, 'I relax,') (F.) *Parésie*. Paralysis. According to some, a slight, incomplete paralysis, affecting motion but not sensation.

PAR'ESTH'ESIS, *Paræsthes'is*, *Paræsthes'ia*, *Sens'us imperfect'us*, (*para*, and *αἰσθησις*, 'sensation,') Congenital misconstruction of the external organs of sense. A genus in the nosology of Good. Also, perversion of sensibility.

PAR'ETICA EXARTHROSIS. Luxation from relaxation of the ligaments of a joint.

PAR'ETICUS, (*παρετος*, 'relaxed,') *Relaxant*.

PAREYRA, *Pareira brava*.

PAR'FUM, (*per*, and *fumus*, 'smoke,') *Suffimentum*.

PARIERA, *Pareira brava*.

PARIES, gen. *Pariet'is*, 'a wall,' *Table*.

PARI'ETAL, *Pariet'al'is*, *Pariet'inus*, (from *paries*.) (F.) *Pari'etal*. Appertaining to the parietes or walls of an organ or to the parietal bones.

PARIETAL ARTERY, see *P. Bones*.

PARIETAL BONES, *Os'sa pariet'al'ia seu vertical'ia seu vertic'is seu sincip'itis seu bregmat'is seu bregmat'ica seu nerc'al'ia seu tetrag'onia*, are two fellow-bones, occupying the lateral and upper parts of the cranium, in the formation of which they assist. They have a quadrilateral shape: their external surface is convex; and presents, at its middle part, an eminence called the *Pari'etal protuberance*, *Tu'ber pariet'al'e*, (F.) *Bosse pariet'ale*. Their inner or cerebral surface, which is concave, has a depression that corresponds with the preceding eminence, and which is called the *Pari'etal fos'sa*, *Fos'sa pariet'al'is*. Above and behind, near the superior edge of these bones, is a foramen, which does not always exist, called the *Pari'etal fora'men*, *Fora'men pariet'al'e*, through which passes a vein—*pari'etal vein*—that opens into the superior longitudinal sinus of the dura mater: and an artery—*pari'etal artery*—a branch of the occipital, which ramifies on the same sinus. Each parietal bone is articulated above, with its fellow; *below*, with the temporal bone; *before*, with the frontal; *behind*, with the occipital, and by the anterior and inferior angle with the great ala of the sphenoid; and is developed from a single point of ossification.

PARIETAL FORAMEN, see *P. Bones*—*p. Fossa*, see *P. Bones*—*p. Protuberance*, see *P. Bones*—*p. Veins*, see *P. Bones*.

PARIETA'RIA, (from *paries*, 'a wall,' because found on old walls.) *P. officinat'is*, *Hel'si'nië*, *Wit-traria*, *Wall pel'titory*, (F.) *Parit'aire commune*, *Perce-muraille*, *Ammor*, *Herbe aux murailles* ou *des Mores*, *Family*, *Urticæ*. See *Syst. Polygamia Monœcia*. This plant contains much nitrate of potassa, and was formerly largely employed as a diuretic, and, also, as an emollient; but it is not such used at the present day.

PARIETARIA PENNSYLVANICA, *American Pel'titory*, a small weed, flowering from June to August, has been used in juice or decoction as a diuretic, 'deobstruent,' and emmenagogue.

PARI'ETES, (pl. of *paries*, 'a wall,') (F.) *Parois*. A name given to parts which form the enclosures or limits of different cavities of the body, as the *Parietes* of the cranium, chest, &c.

PARIETINUS, *Parietal*.

PARIS, gen. *Par'idis*, *P. quadrifo'lia*, *Sola'mum quadrifo'lum*, *U'va inver'sa seu lupi'na seu vulpi'na*, *One'berry*, *True'love*, (F.) *Parisette*, *Raisin de Renard*, *Herbe à Paris*, *Étrangle-loup*. It possesses narcotic properties, and was formerly considered a powerful philter. It is emetic, and is sometimes used externally as a resolvent and anodyne.

PARISAGOGUE INTESTINORUM, (from *παρεισαγω*, 'to introduce improperly,') *Intussusceptio*. *PARISETTE*, *Paris*.

PARISTH'MIA, (*παρισθυια*, from *para*, and *ισθος*, 'the throat,') A word employed to designate the tonsils, and, likewise, inflammation of the tonsils, *Cynanche tonsillar'is*, and *Cynanche* in general.

PARISTHMIOTOMUS, (*paristh'mia*, and *τομω*, 'I cut,') An instrument with which the tonsils were formerly scarified.

PARISTHMITIS, (*paristh'mia*, and *itis*.) *Cynanche*, *C. tonsillar'is*.

PARISTHMIUM, (*παρισθυμιον*) *Tonsil*.

PARISWORT, *Trillium latifolium*.

PARMACITY, (corr. from *spermaceuti*.) *Ceta-ceum*—*p. Poor Man's*, *Thlaspi bursa*.

PARMELIA ISLANDICA, *Lichen Islandicus*.

PARMELIA PARIET'INA, *Lichen pariet'inus*, *Yel'low wall Li'chen*, *Com'mon Yel'low Wall Moss*, which grows abundantly on trees and walls, *Order*, *Lichenaceæ*, is used in intermittent fever on account of its bitterness.

PARMELIA PLICATA, *Lichen plicatus*—*p. Pulmonacea*, *Lichen pulmonarius*—*p. Roccella*, *Lichen roccella*—*p. Saxatilis*, *Lichen Saxatilis*.

PARMENTIERA CERIF'ERA, (after *Parmentier*.) *Palo de velas*, *Can'dle Tree*. A forest tree in the valley of the Chagres,—*Nat. Ord. Crescentiaceæ*,—the long cylindrical fruit of which, resembling candles, serves for food for cattle.

PARMENTIERE, *Solanum tuberosum*.

PAROARION, (*para*, and *ωαριον*, 'ovarium,') *Parovarium*.

PAROCHEUTEUSIS, (*παροχευσις*, from *para*, and *οχευω*, 'to divert water from,') *Derivation*.

PARODINIA, (*para*, and *ωδιν*, 'labour pains,') *Dystocia*—*p. Perversa*, *Presentation*, *Preternatural*.

PARODONTIDES, (*para*, and *οδους*, gen. *οδοντος*, 'a tooth,') *Tumours on the gums*; *parulis*, &c. **PAROIS**, *Parietes*.

PAROLE, (from low Lat. *parabolare*, [F.] *parler*, 'to speak, to parley,') *Voice*, articulated.

PAROMPHALOC'E'L'E, (*para*, *ωμφαλος*, 'the navel,' and *κηλη*, 'rupture,') *Hernia* near the navel.

PARON'IRIA, (*para*, and *ωνειρον*, 'a dream,') *Depraved*, *disturbed*, or *morbid dreaming*; *Onei'rodyn'ia act'iva*, *Ereth'ismus onei'rodyn'ia*. The voluntary organs connected with the passing train of ideas, overpowered by the force of the imagination during dreaming, and involuntarily excited to their natural or accustomed actions, while the other organs remain asleep.

PARONIRIA AMBULANS, *Somnambulism*—*p. Salax*, *Pollution*.

PARONYCH'IA, *Parony'chia*, *Paronychi'tis*, *Pan'aris*, *Redu'via*, *Redi'e'ia*, *Ver'mis seu Pteryg'ium dig'iti*, *Panarit'ium*, *Digit'ium*, *Pandali'tium*, *Pas'sa*, *Dactyl'itis*, *Pan'arium*, *Onych'ia*, (F.) *Onglade*, (*para*, and *ωνυς*, gen. *ωνυς*, 'the

nailed. The ancients gave this name to an inflammatory tumour. — *Oncobolus*, *Oncrytius* (Gross), — seated near the nail, and involving the pulp or matrix. Some of the moderns have extended it to inflammation of the hand, and even of the forearm. It usually means every phlegmonous tumour of the fingers or toes, and especially of the first phalanx. Four kinds of paronychia are commonly pointed out: 1. That seated between the epidermis and skin, called, vulgarly, in this country, *Rod'round* — in France, *Tournaile*, &c. 2. That seated in the subcutaneous areolar tissue. 3. That occupying the sheath of a tendon; and 4. That considered to be seated between the periosteum and bone. The last three are only different degrees of the same disease, constituting *Whitlow*, (Prov.) *Whitlow*, *Tusk*, (F.) *Mal d'ongle*; — the worst kinds, in English, are called *Fellon*, *Cacoparonychia*. The inflammation generally commences in the subcutaneous areolar tissue, and spreads to other parts, occasioning excessive lancinating pain, owing to the parts being put upon the stretch; hence, the necessity of dividing them freely, — even down to the bone, when there is reason to believe that the affection belongs to the fourth division, — making use of emollient applications, the antiphlogistic treatment, and opiates to relieve pain.

PARONYCHIA, *Asplenium ruta muraria* — p. *Digitum*, *Digitaria*.

PARONYCHIS, *Paronychia*.

PARONYCHITIS, *Paronychia*.

PAROPHOBIA, (*para*, and *φοβος*, 'dread,') *Hydrophobia*.

PAROPIA, (*παροπια*, from *para*, and *ωψ*, 'the eye,') *Canthus* (besser.)

PAROPLEXIA, (*para*, and *πλησσω*, 'to strike,') *Paraplegia*.

PAROPSIS, (*para*, and *opsis*, 'sight,') *Dysæsthesia visualis* — p. *Ammaurosis*, *Ammaurosis* — p. *Cataracta*, *Cataract* — p. *Glaucomosis*, *Glaucoma* — p. *Ilusoria*, *Pseudoblepsia* — p. *Lateralis*, *Dysopia lateralis* — p. *Linguisque*, *Presbytia* — p. *Lucifuga*, *Nyctalopia* — p. *Nostalgia*, *Hemeralopia* — p. *Propinqua*, *Myopia* — p. *Staphyloma*, *Staphyloma* — p. *Staphyloma purulentum*, *Hypopyon* — p. *Staphyloma simplex*, *Hydrophthalmia* — p. *Strabismus*, *Strabismus* — p. *Synæsthesia*, *Synæsthesia*.

PAROPTESIS, (*παροπτις*, from *para*, and *πτωω*, 'I roast,') Forcing a sweat by placing a patient before the fire or putting him in a dry bath.

PARORASIS, (*para*, and *οραω*, 'I see,') *Perversion of sight*, which prevents the person from judging accurately of the colour of objects. *Achromatopsia*. *Hallucination*.

PARORCHIDIUM, (*para*, and *ορχις*, gen. *ορχιδος*, 'testicle,') Any position of one or both testicles, different from that which they ought to occupy in the scrotum: — whether they may not have descended through the abdominal ring, or are drawn up to the groin by the cremaster muscle.

PARORCHIDO-ENTEROCELE, *Hernia parorchid-enterica*. Internal hernia complicated with displacement of the testicle.

PAROSMIA, *Anosmia*. Also, perversion of smell; *Parosmia olfactoria*, *Olfactua depravata*, *Idiosyncrasia olfactoria*, (from *para*, and *οσμεω*, 'I smell,')

PAROSTIA, (*para*, and *οστωω*, 'a bone,') Bones untended in their substance, and incapable of affording proper support.

PAROSTA FLEXILIS, *Mollities ossium* — p. *Fragilis*, *Fragilitas ossium*.

PAROTIA, (*παροπια*, from *para*, and *ωψ*, gen. *ωψος*, 'the ear,') *Cynanche parotidea*.

PAROTID, *Glandula parotidea* a sea ad Auris, *Parotis*, *Animalis*, *Alimentis*. The largest of the salivary glands, seated under the ear, and near the angle of the lower jaw. It is composed of many separate lobes, giving rise to excretory

ducts, which unite to form one canal, called the *Parotid duct*, *Stoma* or *Stoma's canal* or *duct*, — the *Ductus superior* or *Superior salivary canal*, of some, see *Ductus salivaris superior*. This duct, after having advanced horizontally into the substance of the cheek, proceeds through an opening in the buccinator muscle, and terminates in the mouth opposite the second upper molaris. About the middle of its course, it sometimes receives the excretory duct of a glandular body, situate in its vicinity, and called the *Accessory Gland of the Parotid*, *Stoma Parotidis*, *Glandula stoma parotidis*. In the substance of the parotid are found — a number of branches of the facial nerve, of the transverse arteries of the face, and the posterior auricular. It receives, also, some filaments from the inferior maxillary nerve, and from the ascending branches of the superficial cervical plexus. Its lymphatic vessels are somewhat numerous, and pass into ganglions situate at its surface or behind the angle of the jaw. The parotid secretes saliva, and pours it copiously into the mouth.

PAROTID, *ACCESSORY GLAND OF THE*, see *Parotid*.

PAROTID APONEUROSIS. A sheath of great thickness, continuous, below, with the cervical fascia. It forms a frame-work to the parotid by means of fibrous prolongations from its deep surface. Its density accounts for the pain of parotitis, and for the difficulty with which pus makes its way to the surface.

PAROTID ARTERIES, see *Parotid* — p. *Duct*, see *Parotid* — p. *Veins*, see *Parotid*.

PAROTIDE, *Parotis*.

PAROTIDEAN, *Parotidea*, (F.) *Parotidien*. Belonging or relating to the parotid.

PAROTIDITIS, (*parotid*, and *itis*,) *Cynanche parotidea*, *Parotis*.

PAROTIDONCUS, *Tumor parotidis*, *T. Glandula parotidea*, (*parotid*, and *ocys*, 'a tumour,') Swelling of the parotid.

PAROTIS, gen. *Parotidis*, (*παροπια*, from *para*, and *ωψ*, gen. *ωψος*, 'ear,') *Phlegmon parotidea*, *Inflammatio parotidis*, *Cynanche parotidea*, *Parotitis*, *Parotiditis*, *Dissectio*, (F.) *Parotide*, is the name of a tumour seated under the ear; which is reddish, hard, and attended with obtuse pain: the progress to suppuration being slow and difficult. Sometimes it is of a malignant character, sloughing, and long protracted. Some French pathologists restrict the term *parotide* to symptomatic parotitis — as to that which occurs in plague; whilst they apply the terms *Oreillons* or *Ourlas*, *Parotomies*, to *Cynanche Parotidea* or *Mumps*. Also, the parotid.

PAROTIS, *Parotid* — p. *Contagiosa*, *Cynanche parotidea* — p. *Epidemica*, *Cynanche parotidea* — p. *Serosa glutine tumens*, *Cynanche parotidea* — p. *Sparis*, *Cynanche parotidea*.

PAROTITIS, (*parotid*, and *itis*,) *Cynanche parotidea*, *Parotis* — p. *Epidemica*, *Cynanche parotidea* — p. *Erysipelatosa*, *Cynanche parotidea* — p. *Polymorpha*, *Cynanche parotidea* — p. *Specific*, *Cynanche parotidea*.

PAROTONCIES, (*parotid*, and *ocys*, 'a tumour,') see *Parotis*.

PAROVA RIUM, properly *Parovarium*, *Corpus seu Corpusculum ovium* *Rosenmülleri*, *Organ* or *Body of Rosenmüller*, *Appendage to the Ovary*, (*para*, and *ovum*, 'the ovary,') A body closely analogous in structure to the epididymis, which is seated between the Fallopian tube and the ovary in the broad ligament. The *Corpus Wolffianum* concurs in its formation.

PAROXYNTIC, (*παροξυντικός*,) *Paroxyntica*, (F.) *Paroxynismus*, *παροξυνω*, 'I excite,') Relating to a paroxysm.

Jours paroxyniques are days on which paroxysms take place.

PAR'OXYSM, (παροξυσμός, *Paroxysmus*, (παροξυνω, from *para*, and *oxynai*, 'to render sharp.' 'I irritate.') *Impetus*, *Inasultus*, *Invasio*, *Exacerbatio*, *Accessio*, *Accessus*, (Old. Eng.) *Access*, *Access*, (F.) *Exacerbation*, *Redoublement*, *Accès*. The augmentation which supervenes at equal or unequal intervals in the symptoms of acute diseases. Also, a periodical exacerbation or fit of a disease.

PAROXYSMAL, *Paroxysmalis*, (F.) *Paroxysmique*. Of or belonging to a paroxysm;—as a *paroxysmal day*, (F.) *Jour paroxysmique*; a day on which a paroxysm of disease occurs.

PAROXYSMUS FEBRILIS, Pyrexia.

PAROXYSTIQUE, Paroxysmal.

PARS, gen. *Partis*, 'a part,' Genital organs—p. *Corporis*, Genital organs—p. *Inferior pedis*, Sole—p. *Intermedia*, see *Corpus spongiosum urethræ*—p. *Mastoidea* of Temporal Bone, Mastoid process—p. *Naturalis medicince*, Physiology—p. *Obscena*, Genital organs—p. *Petrosa*, see Temporal Bone—p. *Prima aspera arteria*, Larynx—p. *Pyramidalis ossis temporis*, see Temporal Bone—p. *Squamosa*, see Temporal Bone—p. *Urethræ membranacea*, see *Urethra*—p. *Vaga*, *Pneumogastrie*—p. *Virilis*, Penis.

PARSLEY, COMMON, (corr. from *petroselinum*, [F.] *persil*.) *Apium petroselinum*—p. Fool's, *Æthusa cynapium*—p. Macedonian, *Bubon macedonicum*—p. Mountain, black, *Athamanta aureo-selinum*—p. Poison, *Conium maculatum*—p. Spotted, *Conium maculatum*—p. Water, *Cicuta maculata*.

PARSNEP, COW, (corr. from *pastinaca* [?]), *Heracleum spondylium*, *Heracleum lanatum*—p. Crow, *Leontodon taraxacum*—p. Garden, *Pastinaca sativa*—p. Meadow, *Thapsium Barbinode*—p. Water, common, *Sium latifolium*.

PART, (from *parere*, *partum*, 'to bring forth,') see Parturition—p. *Légitime* et *illégitime*, see Parturition.

PARTES, pl. of *Pars*, Genital organs—p. Generation inservientes, Genital organs—p. *Genitales*, Genital organs—p. *Laterales nasi*, *Scapulæ nasi*—p. *Nervosæ*, see Nerve—p. *Obscena*, Genital organs.

PARTHENEIA, (παρθενα, *Virginity*.

PARTHENIUM FEBRIFUGUM, (from *παρθενος*, 'a virgin,' used in diseases of virgins,) *Matricaria*—p. *Mas*, *Tanacetum*.

PARTHENIUS (*Morbis*), (παρθενιος, 'relating or appertaining to a virgin.') A disease in a young female. Also, chlorosis.

PARTHENOGENESIS, 'virgin birth,' (παρθενος, 'a virgin,' and *γενεσις*, 'generation.') The successive production of procreating individuals from a single ovum, without any renewal of fertilization.

PARTHENOLOGY, *Parthenology'ia*, (παρθενα, 'virginity,' and *λογος*, 'a description.') A description or consideration of the state of virginity in health or disease.

PARTHENONOSUS, (*parthenus*, and *νοσος*, 'disease,') Chlorosis.

PARTHENUS, (παρθενος, *Virgin*.

PARTIALIS, (*pars*, gen. *partis*, 'a part,') Local.

PARTICLES, PRIMITIVE, OF MUSCLES, Sarcous Elements.

PARTICULÆ SANGUINIS, (dim. of *pars*, gen. *partis*, 'a part,') Globules of the blood.

PARTIES, LES, 'the parts,' Genital organs—p. *Genitales*, Genital organs—p. *Génitoires*, Genital organs—p. *Honteuses*, Genital organs—p. *Naturelles*, Genital organs—p. *Nobles*, Genital organs—p. *Sexuelles*, Genital organs.

PARTIO, (*parere*, *partum*, 'to bring forth,') Parturition.

PARTIDGE-BERRY, *Gaultheria*, *Mitchella repens*.

PARTS, THE, Genital organs.

PARTURA, (*parturire*, *parturitus*, itself from *parere*, 'to bring forth,') Parturition.

PARTURIENT, *Parturiens*. Bringing forth, or about to bring forth, or having recently brought forth young.

The *Parturient*, *Puerperous*, *Child'ing*, *Child'bed* or *Lying-in* or *Lying-down state*, *Confinement*, *Puerperium*, generally comprises the period from parturition to perfect recovery, which is usually a month. In the north of England this is called the *steg month*, requires careful management, and is subject to various diseases.

A *Parturient* or *Parturificient* is a medicine which induces or promotes labour; as ergot, and perhaps borate of soda. See Abortive.

PARTURIFACIENT, (*parturio*, 'to bring forth,' and *facere*, 'to make,') Parturient.

PARTURITION, *Puerperium*, *Partus*, *Partio*, *Parturit'io*, *Partura*, *To'cus*, *To'cetus*, *Tec'is*, *Lochi'a*, *Apocye'is*, *Mæu'is*, *Maieu'is*, *Nego'tium* seu *Ac'tus parturitio'nis*, *Eniz'us* seu *Enic'io* seu *Expul'sio* *fa'ctus*, *Apol'ysis*, *Stat'us parturient'is*, *Deliv'ery*, *La'bour*, *Child'bearing*, *Child'birth*, *Birth*, *Trav'ail*, (Sc.) *Childill*, *Crying*, *Crying out*, (F.) *Accouchement*, *Couche*, *Travail d'Enfant*, *Enfantement*, *Mise-bas*. The act of delivery of the foetus and its appendages; also the state during and immediately after delivery—*Childbed*, (Sc.) *Gizzenbed*, *Gizzen*, see Parturient. Labour is the necessary consequence of conception, pregnancy, and the completion of gestation. It does not take place, at the full time, until after nine months of utero-gestation,—the causes producing it being the contraction of the uterus and abdominal muscles. By different authors, the stages of labour have been variously divided. We may, perhaps, admit of four. The first stage comprises the precursory signs. One, two, or three days before labour, a mucous discharge, streaked with blood, takes place from the vagina, which is called the *Signum*, *Leucorrhæa* *Nabo'thi*, *Hæmorrhag'ia* *Nabo'thi*, *La'bourshow* or *Show*, (F.) *Glaïres*, and along with this, trifling grinding pains, called by the French *Mouches*, are felt in the loins and abdomen. Second stage.—Peculiar pains, which extend from the lumbar region towards the uterus; tension and dilatation of the neck and orifice of that organ; protrusion into the vagina of the envelopes of the foetus, forming the *bag of waters*, (F.) *Poche des eaux*; and rupture or breaking of the waters. Third stage.—Contractions of the uterus and abdominal muscles; forcing pains; the head of the foetus becomes engaged in the pelvis: the occiput being commonly situate above the left acetabulum, passes beneath the arch of the pubis. Fourth stage.—The head now presents at the vulva, the perineum being considerably distended; at length it clears the external parts, and the rest of the body is easily disengaged. The child still continues attached to the mother by the umbilical cord, which has to be divided. In the course of fifteen or twenty minutes, trifling pains occur, and the secundines are expelled. This completes delivery—the *birth*, (F.) *naïssances* of the child. In by far the majority of cases, the head is the presenting part, the occiput directed anteriorly, and the face posteriorly. Of 12,633 children, born at the Hospital *La Maternité*, in Paris, from the 10th of December, 1797, to the 31st of July, 1806, the following were the presentations:

The head, with the face posteriorly...	12,120
The head anteriorly.....	63
The breech.....	198
The feet.....	147
The knees.....	3
Other positions.....	102
	12,633

The following table, drawn up from data furnished by Velpeau, gives an approximate view of the comparative number of presentations, in

1000 cases, according to the experience of the individuals mentioned :

	ACCORDING TO							
	Merri- man.	Bland.	Madame Boivin.	Madame Lacha- pelle.	Naegele.	Lovati.	Hospital of the Faculté.	Boer-
Regular or of the vertex.....	924	944	969	933	933	911	950	
I. <i>Occipitoanterior</i>	908		944	910		895		
a. <i>Occipitocotyloid</i> (left).....			760	717		557		
Do. (right).....			179	209				
b. <i>Occipitopubian</i>			0.29					
II. <i>Occipitoposterior</i>			9.4	9				
a. <i>Frontocotyloid</i> (left).....			5.3	7.3				
b. Do. (right).....			4.4	2.9				
Face presentation.....	2.2	2.6	3.6	4.6				8.8
Mentoiliac (right).....				2.6				
Of the pelvis.....	36	28	29	36	47			29
Of the foot.....	12.7	9.4		14				10.3
Of the knees.....			0.19	0.40				
Of the breech.....	23	13	18	22				19
Of the trunk.....			4.6	5.3	4.8			
Requiring Forceps.....	6.6	4.7	4.6	3.4	36			5.7
“ Turning.....	16	4.7		7.8	7.2			5.9
“ Cephalotomy.....	3.3	5.2	4.77	0.53	2.4			1.5

Labour has received different names, according to the period of pregnancy at which it occurs, and the mode in which it takes place. It is called *miscarriage* or *abortion*, prior to the seventh month; *premature labour*, *Omotocia*, *Partus præcox* seu *præmaturus*, between this and eight months and a half; and *labour at the full period*, (F.) *Accouchement à terme*, when it happens at the end of nine months. Labours have likewise been variously divided by different authors. Perhaps the one into *Natural* and *Preternatural*, is as good as any;—*natural Labour* meaning that which would occur without manual assistance; and *preternatural*, requiring the assistance of art, either by the hand solely, or with the aid of instruments—*instrumental labour*, which M. Capuron calls *Accouchement Mécanique*. Professor Hodge admits the following classification :

EUTOCIA. { Vertex presentations.
 { Pelvic presentations.

DYSTOCIA. { Various states of { Large heads.
 the fetus..... { Malpresentations.
 { Plurality of children.
 { Deformities.

DYSTOCIA. { Conditions of the { Deformed pelvis.
 pelvis or uterus. { Displacements of the
 uterus.
 { Physiological states { Inordinate excite-
 ment.
 { Pathological states { Deficient excitement.
 Local.
 General.

Powerless labour, the *Inertie par épuisement* of Madame Lachapelle, is that in which the muscular powers from exhaustion are unable to accomplish delivery.

Forced labour or *parturition*, (F.) *Accouchement ou Avortement provoqué*, is labour or abortion, induced by art. When accompanied with other phenomena of a deranging character, it is said to be *complicated*. When the child, even when reduced in size, cannot pass, it is termed *impracticable*, &c. See *Dystocia*.

The French use the term *Part*, (L.) *Partus*, at times, synonymously with *delivery*; at others, with the child, *Maicuma*, *Mæu'ma*. *Suppression de part* means concealment of delivery; *Part légitime*, a fetus which comes into the world with the conditions necessary for its enjoying advantages granted by law; and *Part illégitime*, one that does not fall under those conditions.

TABLE OF THE COURSE AND MANAGEMENT OF DIFFERENT KINDS OF LABOURS.

I. NATURAL LABOURS ;

Or those in which Delivery can be effected without Assistance.

HEAD PRESENTING. (Four Positions.)	1. <i>Occiput towards the Left Acetabulum.</i>	The head with the chin bent upon the breast clears the brim of the pelvis. Movement of rotation produced by the inclined planes of the pelvis; the occipital protuberance is carried behind the symphysis pubis, and the face towards the hollow of the sacrum. The head attains the outlet: clears it; the occiput being first disengaged, and the face afterwards, which passes successively along the whole of the hollow of the sacrum. The shoulders enter the brim diagonally; and execute the movement of rotation. The one behind first clears the outlet. The same thing occurs with the nates. The same course is here followed: except that after the rotary movement, the face passes behind the symphysis pubis, and the occiput into the hollow of the sacrum. The occiput is first extruded;—the face next.
	2. <i>Occiput towards the Right Acetabulum.</i>	
	3. <i>Occiput towards the Right Sacroiliac Symphysis.</i>	

NATURAL LABOURS.—CONTINUED.

LOWER EXTREMITY
PRESENTING.
(Four positions of
the Feet, Knees,
and Breech.

- | | |
|---|--|
| <p>1. <i>Occiput towards the Left Sacroiliac Symphysis.</i></p> | <p>When the chin presents and is turned forwards, the face is first expelled; the head is thrown backwards. This presentation is uncommon. If the chin present and be thrown backwards, natural delivery cannot be effected.</p> |
| <p>1. <i>Heel, Tibia, or Sacrum towards the left Acetabulum.</i></p> | <p>The nates enter the brim diagonally; movement of rotation: the one passes behind the symphysis pubis; the other towards the hollow of the sacrum. The one behind is first extruded.</p> |
| <p>2. <i>Heel, Tibia, or Sacrum towards the Right Acetabulum.</i></p> | <p>The same thing happens to the shoulders. The chin is bent upon the chest to clear the brim: movement of rotation: the occiput passes behind the symphysis pubis, and the face towards the hollow of the sacrum. The face clears the outlet first; the occiput last.</p> |
| <p>3. <i>Heel, Tibia, or Sacrum towards the Right Sacroiliac Symphysis.</i></p> | <p>The same progress, except that after the movement of rotation, the face is towards the pubis, and the occiput towards the sacrum. The face is here again delivered first.</p> |
| <p>4. <i>Heel, Tibia, or Sacrum towards the Left Sacroiliac Symphysis.</i></p> | <p></p> |

II. PRETERNATURAL LABOURS.

1. *Those which require Manual Assistance only.*

I. PRESENTATION OF THE
HEAD, UPPER EXTREMITIES,
BACK, ABDOMEN,
&c.

1. When the labour is without any serious complication, and the head is not in a good position at the brim of the pelvis, endeavours are to be made to place it in its natural position: the hand to be introduced with the fingers united, and the long diameter of the head to be placed diagonally in the brim, if possible. Or the lever may be fixed over the occiput, and, by drawing it down, a closer approximation of the chin and chest may be effected; the position of the head to the pelvis being at the same time attended to.
2. If complicated with syncope, convulsions, hemorrhage, want of uterine action, &c., the feet must be laid hold of, and the child *turned*. For this purpose, empty the rectum and bladder; and turn, if possible, prior to the escape of the liquor amnii. The female may be placed on her back or left side, with the breech over the edge of the bed. Use the right or left hand, according as the feet of the child are to the right or left side of the pelvis; always so introducing the hand, that the child shall be in its palm, and the back of the hand opposed to the inner surface of the uterus. The hand and arm to be introduced during an interval of pain; and they must be flattened and tranquil whilst the uterus is contracting forcibly. If both feet cannot be readily brought down, seize one, and tie a fillet around it.
3. If the toes point to the vertebral column, or to the abdomen of the mother, the head, in its descent, will not enter the pelvis: the chin and occiput will be hitched on the pubis and promontory of the sacrum. In such case, grasp the nates as soon as they have passed the os externum; and, during a pain, endeavour to direct the toes towards either sacro-iliac synchondrosis. When the fingers can be passed along the body of the child, and over the shoulders to the bend of the elbows, draw down the arms successively. When the body is expelled, and the head filling up the brim, the termination of labour must be accelerated by passing two fingers over the shoulder of the child; using moderate and steady extracting power, whilst one finger of the other hand, passed into the mouth, depresses the chin on the sternum. In this way, the head may be extracted.

2. *Those requiring the assistance of appropriate Instruments.*

GENERAL OBSERVATIONS.

1. *Short Forceps*.....2. *Long Forceps*.....3. *Lever or Vectis*.....4. *Fillet and Blunt Hook*.....

- If the pains become feeble, or inadequate from any cause;—or if the presenting part be firmly wedged in the pelvis; or the woman be becoming exhausted, instrumental assistance may be required. In all such cases, place the female on the back or left side, empty the bladder and rectum; wait till the os uteri is in a fit state; afford assistance during the pains; introduce the instrument during the intervals between the pains; have the patient on her left side (generally); and extract according to the axes of brim and outlet, as the case may be.
- Apply these to the sides of the head of the child, so that the ears and parietal protuberances may be within the fenestrae. Introduce the left-hand blade first; the index and middle fingers of the right hand guiding it to the ear. With the right hand pass the other in an opposite line, corresponding with the course of the first blade, guided by the fore and middle fingers of the left hand; the third and little fingers being employed to retain the lefthand blade in place; humour the instrument so as to allow the blades to lock. The handles may now be tied gently together. The force used must be from blade to blade, with a gentle, tractile effort.
- Applicable where the head has not entered the brim. The long forceps must be applied, in most cases, over the occiput and face of the child, so that the convex edges of the blades may be towards the hollow of the sacrum. When used, the power may be exerted from side to side, with moderate traction.
- The lever is applicable to the same cases as the forceps. It must not be used strictly as a lever, but as a hook with which to draw downwards. It may be applied to any part of the head, but is generally hitched on the occiput.
- In certain cases of breech and knee presentation, where the hand is sufficient, it may be necessary to pass over the joint a fillet, or the blunt hook, with which to exert some force of traction. The operation is very simple.

PRETERNATURAL LABOURS.—CONTINUED.

Disproportionate size of head, or distorted or deformed pelvis, may require the child to be destroyed, and its bulk lessened; or the Cæsarean section or symphyotomy, or the induction of premature labour may be called for.

The instruments required here, are the *Perforator*, *Crotchet*, and *Craniotomy Forceps*. Make a steady pressure on the abdomen; pass two fingers the left hand up to the head: feel for a suture or fontanel; introduce the perforator and bore through, until the progress of the instrument is arrested by its shoulders. Open the handles, and turn the instrument in different directions, so that the opening may be large enough to admit the perforator, with which the brain must be broken down.

If, after this, delivery cannot be accomplished without further aid, pass up the craniotomy forceps; open the handles slightly, and introduce the blade without touch within the osseum. On sliding the forceps, a hold is obtained; and a traction force can now be exerted during the pains, and the extrusion of the child be effected.

Should great difficulty exist at the brim, the bones at that part of the head may have to be removed, until the base alone remains. The chin must then be brought through first.

The crotchet is often used instead of the craniotomy forceps, being passed into the opening made by the perforator, and hooked upon some bony projection in the interior of the skull; but it is not a good instrument. In presentations of the face demanding perforation of the cranium, the perforator should be introduced just above the nose, in the occipital suture.

When it is necessary to open the head after the lower extremities have been expelled, the perforation must be made behind the ear.

In cases of great narrowness of the pelvis, the operation of *Symphiotomy* has been adopted by French practitioners, rarely by British or American; and, in extreme cases, *Embole* has been had to the Cæsarean Section.

If the body of the foetus has been firmly separated, and the head left in the uterus, it must be laid hold of with one hand, and the forceps be applied; or the head be opened and delivered, as under *embryulcia*. Should the head have been brought away, and the body left behind, the feet must be brought down.

In cases of narrowness of the pelvis, where the foetus, at the full term, cannot be born alive, delivery may be brought about at an earlier period, or as soon as the child is capable of carrying on its functions independently of the mother.

The most advisable plan for accomplishing this object is merely to pass the finger round and round within the os and cervix uteri, so as to stretch the os, the membranes being thus left entire, and the life of the child not so much endangered, as where the membranes are punctured, and the waters evacuated. Parturition usually follows within forty-eight hours.

EMBOLE.

SYMPHYOTOMY AND CÆSAREAN SECTION.

SEPARATION OF THE HEAD OF THE FŒTUS FROM THE BODY.

INDUCTION OF PREMATURE LABOUR.

The laying of eggs by animals—*Ovulation*—is called, in French, *Ponte*. The term has been applied to the periodical discharge—*Ponte périodique*—of ova, (F.) *Ovule des (Œufs)*, which is supposed to take place from the ovaries at each menstrual period.

PARTURITION, DRY, *Partus siccus*.

PARTURIUM VANUM, *Fœtus coeque*.

PARTUS, gen. *Partūs*, (parens, partum, 'to bring forth,') Parturition—*p. Abortus*, Abortion—*p. Abactus*, Abortion—*p. Cæsareus*, Cæsarean section—*p. Difficilis*, Dystocia—*p. Laboriosus*, Dystocia, Laborious labour—*p. Præcox*, Parturition (premature)—*p. Præmaturus*, Parturition (premature)—*p. Serotinus*, Oxytocia.

PARTUS SIC'CUS, *Dry labour* or *parturition*. Labour, which is neither preceded nor attended by a discharge of the liquor amnii.

PARULIS, (parula, from *pava*, and *alvus*, 'the gum,') *Apocynum parulis* seu *paron'ia*, *Phlegmon parulis*, *Gumbail*, *Alveolar abscess*, (F.) *Parulis*. Small abscesses are so called which generally form in the alveoli and involve the gums, sometimes without any known cause, but which often depend upon carious teeth.

PARURIA, (para, and *urina*, 'I pass urine,') Morbid secretion or discharge of urine.

PARURIA INCONTINENS, Enuresis—*p. Incontinens aquosa*, see *Diabetes*—*p. Inaps*, Ischuria, false—*p. Mellita*, Diabetes—*p. Retentio*, Ischuria—*p. Retentionis renalis*, Ischuria—*p. Retentionis vesicalis*, Retention of urine—*p. Stillitia*, Strangury—*p. Stillitia mucosa*, Cystirrhoea.

PARVITAS MORBOSA PARTIUM ORGANICARUM, 'morbid smallness of organic parts,' *Microtesia*.

PAR'YGRON, (paragon, from *pava*, and *typos*, 'bumid,') A liquid or moist preparation for allaying topical inflammation.

PAS, (from [L.] *passus*,) Step—*p. d'Asc*, Tas-

siago, *Speculum oris*—*p. de Lion*, *Helleborus foetidus*.

PASMA, gen. *Pasm'atis*, (casiv, 'to sprinkle,') Catapasma.

PASQUE FLOWER, (from flowering about Easter,) *Anemone pulsatilla*.

PASSA, (patior, *passus*, 'to suffer,') Paronychia. PASSE CORINTHIACÆ, (*passus*, 'dried,') see *Vitis Corinthiaca*.

PASSAGE, Dejection.

PASSAGES, DIGESTIVE, Ways, digestive—*p. Second*, Ways, second.

PASSAMBIO, *Rio vinaigro*.

PASSE-PLEUR, *Anemone pulsatilla*.

PASSE-PIERRE, ('pass stone,') *Crithmum maritimum*.

PASSERAGE, *Iberis amara*—*p. Sauvage*, *Cardamine pratensis*.

PASSERTYNA TARTONRAIRA, (from *passer*, 'a sparrow,' alluding to the beaked seed,) (F.) *Passerina de Tarton-raire*, *Tarton-raire*, *Gras Roussin*, *Trintandie Malherbe*, *Family*, *Daphnoides*. A plant, whose bark is possessed of epispastic properties. An ointment, prepared like that of Mezeron, is said to possess more activity than the latter.

PASSERINE DE TARTON-RAIRE, *Passerina tartonraira*.

PASSIBILITY, GREAT, (patior, *passus*, 'to suffer,') and *habilis*, 'able,') Hyperæsthesia.

PASSIF, Passive.

PASSIFLORA LAURIFOLIA, (*passio*, 'passion,' and *flor*, gen. *floris*, 'a flower,') *Boyleana Passifloræ flor'æ*, *Ord.* *Passifloræ*. A native of Surinam. The fruit has a delicious smell and flavour, and quenches thirst, abates heat of the stomach, increases the appetite, &c.

PASSIFLORA MALIFOR'IS, *Apple-shaped Gra-nadilla*. The fruit of this species is esteemed a delicacy in the West Indies, where it is served up at table in desserts.

Some of the genus are said to possess narcotic and emetic properties.

PASSIO, (*passio*, *passus*, 'to suffer,') Disease, Passion—p. *Æmoptoica*, *Hæmoptysis*—p. *Cadiva*, *Epilepsy*—p. *Cardiaca*, *Cardialgia*—p. *Cholerica*, *Cholera*—p. *Cœliaca*, *Cœliac flux*—p. *Fellidua*, *Cholera*—p. *Hæmoptoica*, *Hæmoptysis*—p. *Hypochondriaca*, *Hypochondriasis*—p. *Hysterica*, *Hysteria*—p. *Iliaca*, *Ileus*—p. *Ischiadica*, *Neuralgia femoropoplitea*—p. *Pleuritica*, *Pleuritis*—p. *Stomachica*, *Dyspepsia*—p. *Ventriculosa*, *Cœliac flux*—p. *Vomiculua*, *Phthisis pulmonalis*.

PASSION, *Pas'sio*, *An'imi Pathe'ma*, *Affectus*, *Emo'tio*. An active affection or emotion of the mind; as rage, terror, love, hatred, &c.

PASSION FLOWER, *BAYLEAVED*, *Passiflora laurifolia*—p. *Hystérique*, *Hysteria*—p. *Iliac*, *Ileus*.

PASSIONES (pl. of *passio*) **ANIMI**, Affections of the mind.

PAS'SIVE, *Passi'vus*, (F.) *Passif*. An epithet for diseases which seem owing to a greater or less diminution of the strength, or which are without apparent reaction. Thus, passive hemorrhages are such as supervene in debilitated individuals, or under conditions in which increased action of the arteries can scarcely be presumed. The term *passive* has, also, been given to aneurisms of the heart with extenuation of the parietes; in opposition to *active* aneurisms, with thickness of the parietes—hypertrophy.

PASSULÆ MAJORES, (dim. of *passus* [*uvæ*],) see *Vitis vinifera*—p. *Minores*, see *Vitis Corinthiaca*.

PASSULA'TUM. A medicine in which the pulp of raisins or currants is a chief ingredient.

PASSUS, (*pandere*, *passum*, 'to extend,') Step.

PAS'SY, *MIN'ERAL WA'TERS* OF. This village is in the vicinity of Paris. There are several springs, containing sulphates of lime, iron, and magnesia, alum, common salt, carbonate of iron, carbonic acid, &c. They are employed as a tonic and astringent.

PASTA, (*παστη*), Paste—p. de *Althæa*, Paste, marshmallow—p. *Amygdalina*, Confection (almond)—p. *Cautica Viennensis*, see Powder, Vienna—p. de *Dactylis*, Paste, date—p. *Emulsiva*, Confection (almond).

PASTA GLYCYRRHIZÆ GUMMA'TA ET ANISA'TA seu *Mas'sa de extracto glycyrrhizæ*. A compound of *Extract of liquorice*, *Gum Senegal*, *Sugar*, *Florence orris-root*, and *Volatile oil of aniseed*. Demulcent.

PASTA DE JUJUBIS, Paste of jujubes—p. *Regia*, Confection (almond).

PASTA ZIN'CI CHLO'RIDI, (F.) *Pâte de Canquoin*, Paste of *Chlo'ride of Zinc*, *Canquoin's Caustic Paste*. This paste is made of three strengths, consisting, respectively, of *Chloride of zinc*, 1 part; *Wheaten flour*, parts 4, 3 and 2; *Water*, sufficient to make a paste. This caustic destroys the part in a direct ratio with the thickness of the layer applied.

PASTE, *Pas'ta*, (F.) *Pâte*. A viscid, tenacious mixture. Also, a compound medicine, like the *pastil*, but less consistent, flexible, less saccharine, and more mucilaginous. Pastes of opium, &c., are used by the dentist in dental caries to relieve pain; and similar preparations of astringents and aromatics are applied to the gums, to give tone to them, and improve the breath. See *Dentifrice*.

A convenient paste for attaching labels to glass, or for other services, is made of *gum tragacanth* and *water*, to which *alum* is added. This the author always keeps by him. Another form is to make a solution of *Gum Arabic*, $\mathfrak{z}\text{ij}$, *Boiling water*, $\text{f}\mathfrak{z}\text{ij}$, and *Glycerin*, $\text{f}\mathfrak{z}\text{ij}$; and another is made by mixing powdered *Gum Arabic*, and powdered *Tragacanth*, of each $\mathfrak{z}\text{ss}$, *Water*, $\text{f}\mathfrak{z}\text{ss}$, *Acetic acid*, mxx .

PASTE, *CAGLIARI*, see *Turundæ Italicæ*.

PASTE OF CHLORIDE OF ZINC, *Pasta Zinci chloridi*.

PASTE, DATE, *Mas'sa seu Pas'ta de dactylis*, Composed of *dates*, *sugar*, *gum Arabic*, *water*, and *orange-flower water*.—Ph. P. Demulcent.

PASTE OF ICE'LAND MOSS, *Pas'ta Lichenis*, (F.) *Pâte de lichen*. A paste made by boiling *lichen Islandicus* to a proper consistence, and adding *gum Arabic* and *sugar*.

The *Pâte de Lichen opiacæ* contains opium in addition; about $\frac{1}{4}$ of a grain to an ounce.

PASTE, ITALIAN, see *Turundæ Italicæ*.

PASTE OF JU'JUBES, *Pas'ta de ju'jubis*, *Mas'sa de zizyphorum fructu*, (F.) *Pâte de Jujube*. Composed of *jujubes*, *gum Senegal*, *sugar*, *water*, and *tincture of lemon-peel*. Demulcent.

PASTE, MARSHMAL'LOW, *Mas'sa de gum'mi Arab'ico*, *Pas'ta de althæa*, (F.) *Pâte de Guimauve*. A compound of *Gum Arabic*, *Sugar*, *White of egg*, and *Orangeflower water*.—Ph. P. It is demulcent.

PASTE, TOOTH, see *Dentifricium*—p. *Vienna*, see *Powder*, *Vienna*—p. *Ward's*, *Confectio piperis nigra*.

PASTEL, (dim. of *pasta*), *Isatis tinctoria*.

PASTÈQUE, *Cucurbita citrullus*.

PASTIL, *Pastil'us*, *Avic'ula Cyp'ria*, *Cande'la fuma'lis*, *Phtho'is*, *Phthois'eus*, (dim. of *pasta*.) A compound of aromatic substances in various shapes; used as a fumigating material when set fire to. See *Fumigation*.

The French use the term *Pastille*, (L.) *Pastil'us*, for a solid, dry, round, oblong, square, or triangular compound, whose base is essential oil, and sugar the excipient. These pastilles are in fact small lozenges, and only differ from the *tablette* in their lesser size. They are termed, also, *Saccharolæ solidæ*.

PASTILLE, see *Pastil*.

PASTILLES DIGESTIVES D'HAUTE-RIVE-VICHY, see *Trochisci Sodæ bicarbonatis*—p. *Fumigating*, see *Fumigation*—p. de *Renaudot*, see *Pilule de Hydrargyro*, *Scammonio et Aloe*—p. de *Vichy*, see *Trochisci Sodæ bicarbonatis*.

PASTILLI DIGESTIVI DARCETII, see *Trochisci Sodæ bicarbonatis*—p. *Emetinæ Pectorales*, *Trochisci emetinæ pectorales*.

PASTILLI DE MENTHÆ PIPERITÆ, *Pep'ermint drops*. (*Sacch. alb.*, *Liquæ menthæ pip.*, *Aq. destillat.* $\mathfrak{z}\mathfrak{z}$ $\mathfrak{z}\text{ij}$.) Boil to the consistence of an electuary. Take of fine *White sugar* in powder, $\mathfrak{z}\text{iv}$; *Volatile oil of peppermint*, $\mathfrak{z}\text{ss}$. Mix, and add the electuary whilst warm. Drop it then on marble, and dry the *drops*.—Ph. P.)

PASTILLUS, *Pastil*.

PASTINACA ALTISSIMA, (dim. of *pasta*, from *pastus*, 'food,') P. *opoponax*—p. *Anethum*, *Anethum graveolens*—p. *Graveolens*, *Anethum graveolens*—p. *Opaca*, P. *sativa*.

PASTINACA OPOF'ANAX. *Ord.* Umbelliferae. The name of the plant which furnishes the *Opoponax*. The plant, whence the *gumresin* is obtained, has, also, been called *Fer'ula opop'onax* seu *opop'anax*, *Pastina'ca altissima*, *Laserpitium* seu *Seli'um chiro'nium*, *Opopana'eum*, *Pa'nax heracle'um* seu *costi'nium*, *Pastina'cea*, *Ky'na*, *Hercules allheal*, *Opop'anaxvoort*.

Opop'anax, *Opopana'eis gummi'resina*, *Opop'onax*, *Ge'zir*, *Pop'anax*, *Gum'mi Pan'acis*, exudes from the roots when wounded. Its odour is strong and peculiar; taste, bitter and acrid. It is met with in the shops, in lumps of a reddish-yellow colour without; white within. It has been recommended as an antispasmodic and emmenagogue; but is seldom used. Dose, gr. x, to $\mathfrak{D}\text{j}$, in powder.

PASTINACA PRATENSIS, P. *sativa*.

PASTINACA SATI'VA, *Flaphoboc'eum*, *Pastina'ca*

syvestris seu *opa'ca* seu *praten'sis*, *Ane'thum* seu *Seli'nium pastina'ca*, *Ban'ica*, *Gar'den pars'-nep*, (F.) *Panais*. The root is sweet and nutritious, and is in esteem as an article of food. It has been used as a diuretic and demulcent in calculous affections, &c.

PASTINACA SYLVESTRIS, P. sativa — p. Sylvestris tenuifolia officinarum, Daucus carota.

PATATE, ([S.] *batata*), *Convolvulus batatas*.

PATHOULLI, *Pogostemon patchouli*.

PÂTE, (old [F.] *paste*), *Paste* — p. *Arsénicale*, Arsenical paste — p. de *Canquoin*, *Pasta Zinci chloridi* — p. *Caustique de Vienne*, Powder, Vienna — p. de *Guimauve*, Paste, marshmallow — p. du *Jujube*, Paste of Jujubes.

PATELLA, *Rot'ula*, *R. ge'mu*, *Epigon'atis*, *Epigon'tis*, *Aeromy'lē*, *Epigou'nis* seu *Epigu'nis* seu *Epig'onis*, *Mo'la* seu *Con'cha* seu *Oe'ulus* seu *Or'bis* seu *Scu'tum ge'mu*, *Oscu'tifor'mē* seu *discifor'mē* seu *thyreoi'des*, *Supergenu'a'lis*, *Caucalo'i'des*, *My'læris*, *My'lē*, *Gam'ba*, *Olecranon mobi'lē*, (Prov.) *Copbone*, (F.) *Rotule*. (Diminutive of *patera*, 'a bowl,' so called from its shape.) A sort of sesamoid, flat, short, thick, rounded bone, situate in front of the knee, and enveloped in the substance of the tendon common to the extensor muscles of the leg. The posterior surface of the patella is divided into two *facettes*, united angularly, which are encrusted with cartilages, and each of which is articulated with one of the condyles of the femur. The patella is developed by a single point of ossification. When the patella is fractured, it is difficult to produce bony union. This can only be accomplished by keeping the fractured extremities closely in apposition.

PATELLA, Cup — p. *Fixa*, *Olecranon*.

PATENCY OF THE FORAMEN OVALE, (from *pateo*, 'to be open,') see *Cyanopathy*.

PATENT MEDICINE, (F.) *Médecine brevetée* ou *patentée*, *Remède breveté* ou *patenté*. A medicinal compound, for the sale of which the proposer obtains from government an exclusive privilege. A specification of his *invention* must be registered in the Patent Office, and, hence, it differs from a secret medicine.

PATER OMNIUM VIVENTIUM, ('father of all living,') *Penis*.

PATERA DIOGENIS, (*patera*, 'a flat dish,') *alm*.

PÂTEUX, (from *pâte*), Clammy.

PATHEMA, (*παθημα*), Affection, Disease.

PATHEMATOLOGY, (*pathema*, gen. *pathe'matis*, and *logos*, 'a discourse,') Pathology.

PATHETIC, *Pathet'icus*, *Pathétique*, (*παθητικός*, from *pathos*), That which moves the passions. A name given to the superior oblique muscle of the eye, and, also, to the 4th pair of nerves.

PATHETIC MUSCLE, Oblique, superior, of the eye.

PATHETIC NERVE, *Nervus Pathet'icus* seu *trochlea'ris* seu *trochleato'rius* seu *trochlea'tor*, the 4th pair of nerves, *Par qu'artum*, (F.) *Oculo-musculaire interne* (Ch.), *Nerv pathétique*, is the smallest encephalic nerve. It arises behind the inferior part of the tubercula quadrigemina (*testes*) at the lateral parts of the valvula Vieussenii. Its roots unite into one thin round cord, which makes a turn upon the peduncles of the brain, and passes into a canal of the dura mater, near the posterior clinoid process. It enters the orbit by the broadest part of the sphenoidal fissure, and terminates in the superior oblique muscle of the eye, by dividing into several filaments.

PATHÉTIQUE, Pathetic.

PATHETISM, Magnetism, animal.

PATHIC, (from *pathos*), Morbid.

PATHICAL, Morbid.

PATHICUS, Morbid.

PATHE, (*παθη*), Alveolus.

PATHOGENETIC, *Pathogenet'icus*, *Pathogen'-*

icus, (*pathos*, and *γενεσις*, 'generation,') Relating to pathogeny.

PATHOGENETIC PHYSIOLOGY, *Physiolog'ia pathogenet'ica*. That part of physiology which relates to the origin of disease.

PATHOGENICUS, Pathogenetic.

PATHOG'ENY, *Pathogen'ia*, *Nosogen'esis*, *Nosogen'ia*. The branch of pathology which relates to the generation, production and development of disease.

PATHOGNOMON'IC, *Pathognomon'icus*, *Sig'n-um conjunctum* seu *characterist'icum* seu *mor'bi essentia'lē*, (*pathos*, and *γνωω*, 'I know,') A characteristic symptom of a disease.

PATHOG'RAPHY, *Pathograph'ia*, (*pathos*, and *γραφω*, 'I describe,') A description of disease.

PATHOLOGIA, Pathology — p. *Humoralis*, Humorism — p. *Solidaris*, Solidism.

PATHOLOG'ICAL, *Patholog'icū*, *Pathol'ogus*, (F.) *Pathologique*. Relating to pathology. Often used in the sense of *pathical*.

PATHOLOGICAL ANATOMY, see *Anatomy*.

PATHOLOGIST, *Pathol'ogus*, (from *pathos*, and *λογος*, 'a discourse,') One versed in pathology.

PATHOL'OGY, *Pathematol'ogy*, *Pathol'og'ia*, *Pathematolog'ia*, *Patholog'icē*, (*παθολογική* (*τεχνη*)). The branch of medicine whose object is the knowledge of disease. It has been defined *diseased physiology*, and *physiology of disease*. It is divided into *general* and *special*. The first considers diseases in common; — the second the particular history of each. It is subdivided into *internal* and *external*, or *medical* and *surgical*.

PATHOLOGY, HUMORAL, Humorism.

PATHOMANIA, *Parapath'ia*, *Mor'al Insan'ity* (*pathos*, and *mania*), *Ma'nia sine Delir'io*. A morbid perversion of the natural feelings, affections, inclinations, temper, habits, moral disposition, and natural impulses, without any remarkable disorder or defect of the intellect or knowing and reasoning faculties, and particularly without any insane hallucination.

PATHOMYOTOM'IA, (*pathos*, *μυς* or *μωμ*, 'muscle,' and *τομή*, 'incision,') The title of a work by Dr. John Bulwer (1649), being "a dissection of the significative muscles of the affections of the mind."

PATHOPTRIDALGIA, (*pathos*, *παις*, 'one's fatherland,' and *αλγος*, 'pain,') Nostalgia.

PATHOPOËTIC, *Pathopoët'icus*, (F.) *Pathopoétique*, (from *pathos*, and *ποιεω*, 'to make,') Producing disease.

PATHOS, (*παθος*) Affection, Disease.

PATIENCE, *Rumex patientia* — p. *d'Eau*, *Rumex hydrolopathum* — p. *Garden*, *Rumex patientia* — p. *Ronge*, *Rumex sanguineus* — p. *Sauvage*, *Rumex acutus*.

PATIENT, (*patior*, 'to suffer,') see *Sick*.

PATIENT OF DEATH, Agony.

PATIENTIA, *Rumex patientia*.

PATIENTIÆ MUSCULUS, Levator scapulae

PATNE, (*παρνη*), Alveolus.

PATOR NA'RIMUM, (*pateo*, 'to be open,') The cavity or the opening of the nostrils.

PATRATIO, (from *patrare*, *patratum*, 'to accomplish,') Coition, Ejaculation, (of sperm.)

PATRINIA JATAMANSI, (after E. L. M. Patrin, a naturalist and traveller of Lyons.) *Nardus Indica*.

PATTE D'OIE (F.), 'Goose's foot.' An aponeurotic expansion, which covers the internal surface of the tibia at its upper part, and consists of the expansion formed by the tendons of the sartorius, gracilis, and semi-tendinosus muscles

PATURSA, Syphilis.

PAU, CLIMATE OF. Pau is the capital of the department of the Lower Pyrenees, and is about 150 miles from Bordeaux, and 50 from Bayonne. The climate corresponds with that of the southwest of France generally; its great

quality is the comparative mildness of its spring, and exemption from cold winds. It is regarded as a favourable winter residence for invalids labouring under chronic affections of the mucous membranes. It is too changeable for the consumption; and to be avoided by the rheumatic.

PAULLINIA, (after S. Pauli, Prof. of Botany at Copenhagen.) This is an extract from a plant of the same name in Brazil. It is prepared by the Indians, and appears to possess excitant powers. In Brazil, and the neighbouring countries, it is given in the form of tisanæ—the powder being mixed with cocoa—in diarrhoea and dysentery. It has been employed also as a tonic. An extract is prepared in Brazil from *Paulinia sorbifolia*, Ord. Sapindaceæ, known there under the name of *Guarana*, which is administered in similar pathological cases.

PAUME DE LA MAIN, Palm.

PAUNCH, (from *pantex*, pl. *pan'tices*), Inguvies.

PAUNTA, (*navau*, 'to allay,') Sedatives.

PAUPIÈRE, Palpebra.

PAUSIMENIA, (*navais*, 'cessation,' and *μνεις*, 'menses,') Cessation of the menses.

PAUSIS, (*navais*,) Sedation.

PAVANA WOOD, *Croton tiglium*.

PAVANNE, *Laurus sassafras*.

PAVILION. The French give the name *pavillon*, (L.) *Papilio*, 'a canopy,' to the expanded extremity of a canal or cavity,—for example, of a catheter, sound, &c.

PAVILION OF THE EAR, *Auricula*, *Pin'na*, *O'tium*, *A'la*, (F.) *Auricule* ou *O'ricule*, *Pavillon de l'oreille*, is seated behind the cheeks, beneath the temple, and anterior to the mastoid process. Its size varies in individuals. It is free above, behind, and below. Anteriorly and within, it is continuous with the neighbouring parts. Its outer surface has several prominences, the *Hel'ix*, *Ant'helix*, *Tr'agus*, *Anti'ragus*, and *Lobe*; and several cavities—the groove of the *Hel'ix*, the *Fos'sa navicula'ris* seu *scaphoïdes*, and the *Con'cha*. The skin which covers the pavilion is very fine, and studded with a number of sebaceous follicles. The auricle is fixed to the head by three fibro-areolar ligaments;—a *superior*, *anterior*, and *posterior*.

PAVILLON DE L'OREILLE, Pavilion of the ear—*p. de la Trompe*, see *Tuba Fallopiiana*.

PAVIMENTUM CEREBRI, ('pavement of the brain,') Base of the brain.

PAVINA, *Æsculus hippocastanum*.

PAVITA'TIO, *Tre'mor*, (*pavitare*, *pavitatum*, 'to tremble from fear,') Trembling from fright.

PAVOR (L.), Dread. Great fright. *Panophobia*.

PAVO'RES NOCTUR'NI seu *DORMIENT'ium*. Fear during sleep.

PAVOT, Papaver.

PAW, ([F.] *patte*, (W.) *pawen*, 'a foot,') Manus.

PAWPAW, see *Carica papaya*.

PAXWAX, see *Nucha*.

PAXYWAXY, see *Nucha*.

PAZHAR, Bezoar.

PEA, ([F.] *pois*,) *Pisum*—*p. Garden*, *Pisum*—*p. Ground Squirrel*, *Jeffersoniana Bartoni*—*p. Hoary*, *Galega Virginiana*—*p. Love*, *Abrus precatorius*—*p. Nut*, *Arachis hypogea*—*p. Partridge*, *Cassia chamaecrista*—*p. Turkey*, *Galega Virginiana*.

PEACH, (from *Pessicum*, corr. from *Persicum*, 'Persian,') see *Amygdalus Persica*—*p. Brandy*, see *Amygdalus Persica*—*p. Tree*, *Amygdalus Persica*—*p. Wood*, see *Cæsalpinia*.

PEAGLE, *Primula veris*.

PEAR, ([Sax.] *pepa*, [L.] *pyrum*,) see *Pyrus malus*—*p. Alligator*, *Persea gratissima*—*p. Anchovy*, *Grias cauliflora*—*p. Avocado*, *Persea gratissima*.

PEARL, ([Sax.] *perla*,) *Per'la*, *Mar'gavan*, *Margari'ta*, *Margell'ium*, *Margell'is*, *Unio*, (F.) *Perle*. A roundish concretion of a silvery white colour, and brilliant polish, found in several shells, and especially in the *Avic'ula margaritif'era*, which has, hence, been called *Ma'ter perla'tum*, *Margari'ta*, and *Mother of pearl*, and is much used in the fabrication of handles for surgical instruments, &c. Pearls were formerly given in powder as astringents and antacids. They consist of lime and an animal matter.

PEARL ASH, Potash of commerce.

PEARL, MOTHER OF, see *Pearl*—*p. White*, *Bismuth*, subnitrate of.

PEAU, (from *pellis*,) *Cutis*—*p. Divine*, *Condom*.

PEAUCLIER, (from *pellis*,) *Platysma myoides*.

PECAN or **PECCAN NUT**, see *Hickory*.

PEC'CANT, *Pec'cans*, (*peccare*, 'to sin,') Morbid, not healthy. An epithet given by the humourists to the humours when erring in quality or quantity. See *Materia Morbosa*.

PECHAN, *Stomach*.

PÊCHE, *Peach*, see *Amygdalus Persica*.

PECHEGUERA. A pulmonary disease affecting children three or four months old in South America, which is promptly fatal.

PÊCHER, *Amygdalus Persica*.

PECHIAGRA, *Pechyagra*.

PECHING, *Anhelatio*.

PECHURIM seu **PICHURIM CORTEX**. A highly aromatic bark, the produce of a species of *laurus*. It is extremely fragrant, like cinnamon, which it greatly resembles in its properties. In Portugal it is used in the cure of dysentery, &c. See *Pichurim beans*.

PECHY'AGRA, *Pechi'agra*, *Ancon'agra*, (*πηχυς*, 'the elbow,' and *αγρα*, 'a seizure,') Gout affecting the elbow.

PECHYS, (*πηχυς*,) Elbow.

PECHYTYBEE, *Porphyra nautica*.

PECKISH, ('disposed to peck,') (Prov.) Hungry.

PECTE, (*πηκτης*,) Cheese.

PECTEN, gen. *Pec'tinis*, ('a border, a row,') Pubes. Also, a comb.

PECTEN DENTIUM, see *Sepes*—*p. Veneris*, *Scandix cerefolium*.

PECTIN, (*πηκτος*, 'coagulated, curdled,') A name given by Braconnot to a principle which forms the basis of vegetable jelly. It is extensively diffused in the juices of pulpy fruits and roots, especially when they are mature; and occasions these juices to coagulate when they are mixed with alcohol or boiled with sugar.

It is the basis of one of the classes of elementary principles of Dr. Pereira—the *pectinaceous*.

PECTINACEOUS, *Pectinosus*.

PECTINA'LIS, (*pecten*, gen. *pec'tinis*, 'the pubes,') *Pectin'eus*, *Mus'culus liv'idus*, *Pectina'tus*, (F.) *Pubio-fémoral*, *Supubio-fémoral* (Ch.), *Pectiné*. This muscle is situate at the inner and upper part of the thigh. It is long, flat, and triangular; is attached, above, to the space which separates the iliopectineal eminence from the spine of the os pubis; and, below, by a flat tendon, to the oblique line which descends from the lesser trochanter to the linea aspera. The pectinalis bends the thigh on the pelvis, and carries it outwards in adduction and rotation. It may also bend the pelvis on the thigh.

PECTINATED, *Pectina'tus*, *Pect'ini-form*, *Pect'inate*, (F.) *Pectiné*, (from *pecten*, gen. *pectinis*, 'a comb,') Having the shape of the teeth of a comb.

PECTINATED MUS'cles, *Pectina'tinus'culi*. The fascicular texture observed in the right auricle of the heart:—*Mus'culi Auric'ulæ Pectina'ti*. Their main use probably is to prevent the over dilatation of the auricles.

PECTINATUS, *Pectinalis*.

PECTINÉ, *Pectinalis*, *Pectinated*.

PECTINEUS, *Pectinalis*.

PECTINIFORM, (*pecten*, 'a comb,' and *forma*, 'shape,') Pectinated.

PECTINOUS, *Pectinaceus*, *Pectinosus*, *Pectinaceus*, (from *pectus*.) Of or belonging to pectin or vegetable jelly,—as a *pectinous* or *pectinaceous* vegetable principle.

PECTORAL, (*pectus*, 'the breast,') Relating to the breast. See *Pectoralis*.

PECTORAL, GRAND, *Pectoralis major*.

PECTORALE, Corset.

PECTORALIS INTERNUS, *Triangularis sterni*.

PECTORALIS MAJOR, *Pectoralis*, *Sternocleidobrachialis*, (F.) *Sterno-costo-clavio-humeral*, *Sterno-humeral* (Ch.), *Grand Pectoral*. A large, flat, triangular muscle, situate at the anterior part of the chest, before the axilla. It is attached, on the one hand, by means of aponeurotic fibres, to the inner half of the anterior edge of the clavicle; to the anterior surface of the sternum, and to the cartilages of the first six true ribs;—on the other, by a strong tendon, which is inserted at the anterior edge of the bicipital groove of the humerus. The use of this muscle is to move the arm in various directions. If it be hanging by the side of the body, it carries it inwards and a little forwards. When the arm is raised, it depresses it and carries it inwards. It can, also, move it in rotation inwards. If the humerus be raised and fixed, the pectoralis major draws the trunk towards the upper extremity.

PECTORALIS MINOR, *Serratus anticus minor*, (F.) *Costo-coracoidien*, (Ch.), *Petit Pectoral*, *Petit dentelé antérieur*. This muscle is situate beneath the preceding, at the anterior and upper part of the chest. It is flat and triangular; is attached by its base, which appears digitated, to the upper edge and outer surface of the 3d, 4th, and 5th true ribs, and is inserted at the anterior part of the coracoid process. The pectoralis minor draws the scapula forwards and downwards, and makes it execute a rotatory motion, by virtue of which its inferior angle is carried backwards, and the anterior depressed. When the shoulder is fixed, it elevates the ribs to which it is attached, and aids in the dilatation of the chest.

PECTORALS, *Pectoralia*, *Thoracica*. Medicines considered proper for relieving or removing affections of the chest.

PECTORAL DROPS, **BATEMAN'S**, consist chiefly of *tincture of castor*, with some *camphor* and *opium*, flavoured by *aniseed*. Gray gives the following formula: *Castor*, ℥i; *Ol. anisi*, ℥i; *Camph.* ℥v; *Cocci*, ℥ss; *Opii*, ℥vj; *Proof spirit*, a gallon.

A form, advised by a committee of the Philadelphia College of Pharmacy, is the following:—*Alcohol. dil. cong.* iv; *Santal. rubr. ras.* ℥i; *digere per horas xxiv*; *cola*, et adde *Pulv. opii*, *Pulv. catechu*, *Camphoræ*, āā ℥ij; *Ol. Anisi*, ℥iv. Digest for ten days.

PECTORILOQUE, (*pectus*, 'the breast,' and *loqui*, 'to speak,') One, according to Laënnec, who presents the phenomenon of *Pectoriloquism*. Also, a stethoscope.

PECTORILOQUIE, *Pectoriloquy*—p. *Chérotante*, *Egophony*.

PECTORILOQUOUS BRONCHOPHONY, see *Pectoriloquy*.

PECTORILOQUY, *Pectoriloquism*, *Cav'ernous voice*, *Pectoriloquia*, (F.) *Pectoriloquie*, *Voix caverneuse*, *V. articulée*. Speech or voice coming from the chest. Laënnec has designated, by this name, the phenomenon often presented by consumptive individuals, when their chests are examined with the stethoscope. The voice seems to issue directly from the chest, and to pass through the central canal of the cylinder,—a phenomenon owing to the voice resounding in the anfractuons cavities, produced in the lungs by the suppurative

or breaking down of tubercles, which constitute abscesses or ulcers of the lungs. By some, it is not separated from bronchophony, having been termed, at times, *strong bronchophony*, and *pectoriloquous bronchophony*. See *Laryngophony*. *Whispering pectoriloquy* is that produced by the whispering of the patient.

PECTOROSUS, (from *pectus*.) One who has a broad chest.

PECTUS, gen. *Pectoris*. Sternum, Thorax—p. *Carinatum*, see *Carina* and *Lordosis*.

PECULIUM, ('private property,') Penis.

PEDAGRA, *Potassæ supertartaris impurus*.

PEDAL, *Pedalis*, *Podalis*, *Podalicus*, (*pes*, gen. *pedis*, 'the foot,') Relating to the foot:—as

PEDAL APONEUROSIS, (F.) *Aponévrose pédiécuse*. A thin, aponeurotic layer, which invests the tendons of the foot from each other.

PEDEBELLY (Prov.), *Physonia*.

PÉDÉRASTIE, *Pæderastia*.

PEDESIS, (*πηδάσας*, from *πηδάω*, 'to leap,') Pulsation.

PEDETHMOS, (*πῆδημος*), Pulsation.

PEDIAL'GIA, *Pedional'gia*, *Pedioneural'gia*, (*pedion*, and *algos*, 'pain,') Pain in the side of the foot. Neuralgia of the foot in general. An affection of this kind, under the name *Pédionalgie*, prevailed in 1762 at Savigliano in Piedmont, and in 1806 amongst the military at Padua.

PEDICEL, *Pedicel*.

PEDICELLATED, *Pediculated*.

PÉDICELLÉ, *Pediculated*.

PED'ICLE, *Ped'icel*, *Pediculus*, (F.) *Pédicule*, (dim. of *pes*, gen. *pedis*, 'a foot,') A slender stalk; also, the narrow portion of a part of a tumor for example,—from which it was developed, and is supported.

PEDICLE, **VITELLINE**, see *Vesicula umbilicalis*.

PEDICULARIA, (from *pediculus*), *Delphinium staphisagria*.

PEDICULATED, *Ped'icellated*, *Pediculatus*, *Pedicellatus*, (F.) *Pédiculé*, *Pédicellé*. Supported by a pedicle.

PEDICULATIO, (from *pediculus*), *Phthieriasis*.

PÉDICULE, *Pedicel*.

PÉDICULÉ, *Pediculated*.

PEDICULUS, *Peduncululus*, *Phtheir*, *Louse*, (Prov.) *Bid'dy*, (F.) *Pou*, (dim. of *pes*, gen. *pedis*, 'a foot,' from its having many feet.) A genus of parasitic insects. The human body is infested by three kinds: the *Bod'ylouse*, or *Clotheslouse*, *Pediculus vestiment'i*, (F.) *Pou de corps*; the *Headlouse*, *Pediculus cap'itis*, (F.) *Pou de la tête*, which lives in the hair; the *Crablouse*, *Mor'pion*, *Fera'lis pediculus*, *Plat'ula*, *Phthi'rius inguinalis*, *Pediculus Pu'bis*, (F.) *Mor'pion*, which infests the hair of the pubes. Infusion of tobacco, or mercurial ointment, or ointment of white precipitate, or of red precipitate of mercury, readily destroys them. The louse occurring in phthieriasis, *pediculus tabescens*, is said to differ from the common louse. Also, a pedicle.

PEDICULUS CAPITIS, see *Pediculus*—p. *Pubis*, *Crab-louse*—p. *Tabescens*, see *Pediculus*—p. *Vestiment'i*, see *Pediculus*.

PÉDIOURE, (*pes*, gen. *pedis*, 'foot,' and *cure*, 'to care for,') Chiropodist.

PEDICUS, (from *pes*, gen. *pedis*), *Extensor brevis digitorum pedis*.

PÉDIEUX, (*muscle*), *Extensor brevis digitorum pedis*.

PEDIGAMBRA, *Ysambra*.

PEDILANTHUS TITHYMALOÏDES, (*πέδιλον*, 'a shoe,' and *ανθος*, 'a flower,') *Euphorbia tithymaloïdes* seu *myrtifolia*, *Jewbrush*, *Order*, *Euphorbiaceæ*, grows in varicose parts of the

West Indies. The practitioners of Curaçoa give a decoction in venereal cases. It is also used in domestic practice, in amenorrhœa. It is, moreover, known and used as *Ipécacuanha*.

PEDILUVIUM, (*pedes*, 'the feet,' and *lavo*, 'I wash.') *Lavipedium*, (F.) *Pédiluvee*, *Bain de pied*. A bath for the feet. See Bath.

PEDIOLUS, (dim. of *pes*, gen. *pedis*, 'a foot,') *Petiulus*.

PEDION, (*πῆδιον*,) Sole.

PEDIONALGIA, Pedialgia.

PEDIONEURALGIA, Pedialgia.

PEDIUM, (*πῆδιον*,) Tarsus.

PÉDONCULE, Peduncle.

PÉDONCULES DU CERVEAU, Peduncles of the brain—*p. du Cervelet*, Corpora restiformia—*p. du Conarium*, Peduncles of the pineal gland—*p. de la Glande pinéale*, Peduncles of the pineal gland.

PEDORA. The sordes of the eyes, ears, and feet.

PÉDOTROPHIE, Pædotrophy.

PED'UNCLE, *Pedunculus*, (F.) *Pédoncule*, (dim. of *pes*, gen. *pedis*, 'a foot,') A flower stalk. This term has been applied to different prolongations or appendices of the encephalon.

PEDUNCLES OF THE BRAIN, *Pedunculû seu Cru'ra seu Can'dex cer'e'brî*, called, also, *Cru'ra antio'ra medull'æ oblongat'æ*, (F.) *Pédoncules du Cerveau*. Two white cords, about $\frac{3}{4}$ ths of an inch in diameter, on the outside of the corpora albicantia. They arise from the medullary substance of the brain, and gradually approach, till they join the tuber annulare. They are formed, internally, of a mixture of cineritious and medullary matter; the former of which, being of a darker colour at one part than in any other part of the brain, has been called *Lo'cus nig'er cru'rûm cer'e'brî*, *Substantia nigra pedunculû*. It separates the main body of peduncular fibres into two parts,—the lower or superficial consists, almost wholly, of white fibres, collected into coarse fasciculi, and is called the *crust or basis or fasciculated portion of the peduncle*. The upper part is named *tegmen'tum*.

PEDUNCLES OF THE CEREBELLUM, *Pedunculû seu Cru'ra cerebell'î*, called, also, *Cru'ra posteriora*, are six in number, three on each side—a superior, a middle, and an inferior. The superior peduncles are generally known as the *Processus cerebell'î ad test'es*, *Cru'ra cerebell'î ad corpora quadrigem'ina*, *Bra'chia copulati'va*: the inferior are the *Processus cerebell'î ad medull'am oblongatam*: and the middle are called, also, *Cerebell'î ped'uncles*—*Processus seu Cru'ra cerebell'î ad pontem*, *Bra'chia pontis*.

PEDUNCLES, INFERIOR, OF THE CEREBELLUM, Corpora restiformia—*p. of the Medulla oblongata*, Corpora restiformia.

PEDUNCLES OF THE PIN'EAL GLAND, *Pedunculû cond'it'î*, (F.) *Pédoncules de la glande pinéale*, *P. du Conarium*, are two medullary bands or strips, which seem to issue from the pineal gland, and proceed on each side upon the upper and inner region of the optic thalami. See Pineal Gland.

PEDUNCULI CEREBELLI, Peduncles of the Cerebellum—*p. Cerebrî*, Peduncles of the brain—*p. Conariî*, Peduncles of the pineal gland, see Pineal gland—*p. Medullæ oblongatæ*, Corpora restiformia.

PEDUNCULUS, Peduncle.

PEGA, (*πηγν*,) Canthus (lesser).

PEGANELE'ON, *O'leum ruta'ceum*, *O'leum rutæ*, (*πεγανον*, 'rue,' and *ελαιον*, 'oil,') Oil of Rue.

PEGANUM, (*πεγανον*,) Ruta.

PEGMIN, (from *πηγνω*, 'to coagulate,') Corium phlogisticum.

PEINE, (*πεινα*,) Hunger.

PEINOTHERAPEIA, (*peine*, and *θεραπεια*, 'treatment,') Limnotherapeia.

PEIRA, (*πειρα*,) Effort, Experience, Nisus.

PELA'DA, (F.) *Pelade*. A word of uncertain origin, applied to alopecia, especially to that resulting from syphilis. See Alopecia and Porrigo decalvans.

PELADE, Pelada—*p. Achromatouse*, Porrigo decalvans—*p. Décalvante*, Porrigo decalvans.

PELAGRA. A kind of scaly erysipelas of the hands—sometimes of the legs; at others, of the face. Pellagra [?].

PELAGRA, Pellagra.

PELARGONIUM AN'CEPS, (*πελαργος*, 'a crane,' from the shape of the capsules,) *Peris'tera an'ceps*. A South African plant, *Nat. Ord. Geraniaceæ*, the decoction of which is used by the Malays in amenorrhœa, and to promote parturition and abortion.

PELARGONIUM ANTIDYSENTER'ICUM, *Jenkinsonia antidynter'ica*. This South African plant is used by the natives in dysentery. It is boiled in milk.

PELARGONIUM CUCULLA'TUM. Common on the Table Mountain, Africa. It has been recommended in decoction as an enema in colic, nephritis, and suppression of urine. It is an excellent emollient, and was formerly exported to Holland under the name *Herba Althææ*.

PELARGONIUM TRIS'TE, *Polyac'tium trist'e*. The root of this plant, which grows at the Cape of Good Hope, is somewhat astringent, and is used there in diarrhœa and dysentery, and as an antelmintic.

PEL'ICAN, *Pelica'nus*, *Peleca'nus*, *Pel'ecan*, *Pel'ecas*, (*πελικαν*, from *πελκαω*, 'I cleave,') An instrument, curved at the end like the beak of a pelican. Its use is to extract teeth.

PELI'CIDE, *Pelu'dê*, *Mel coc'tum*. Boiled honey.

PELICOMETER, (*πελιξ*, gen. *πελικος*, 'a basin,') 'the pelvis,' and *μετρον*, 'a measure,') Pelymeter.

PELIDNOMA, (*πελιδνώμα*, from *πελιδνός*, 'livid,' and *ομα*,) Ecchymoma, Pelioma.

PELIO'MA, gen. *Pelio'matis*, *Pelidno'ma*, (*πeliωμα*, from *πελος*, 'livid,' and *ομα*,) A livid ecchymosis. See Ecchymoma.

PÉLIOSE, Purpura.

PELIOSIS, (*πελιωσις*, from *πελος*, 'livid,' and *osis*,) Purpura hæmorrhagica—*p. Rheumatica*, see Purpura simplex.

PELLA, (*πελλα*, 'skin,') Cutis, Prepuce.

PEL'LAGRA, *Pel'agra*, *Ichthyo'sis seu Tu'ber seu Impet'igo Pel'agra*, *Elephant'iasis Ital'ica*, *Le'pra Mediolanen'sis seu Lombard'ica*, *Scorbut'us alpi'nus*, *Ma'n'ia pella'gia*, *Eryth'ma eudem'icum seu pel'agrum*, *Derm'atagra*, *Erysip'elas per'iod'ica nervo'sa chron'ica*, *Paral'ysis scorbut'ica*, *Insolazione de Primavera*, *Male del Sole*, *Scur'vy of the Alps*, (F.) *Mal de misère*, *Scorbut des Alpes*; (from *πελλα*, [?] 'skin,' and *αγπα*, 'seizure,') A disease, particularly noticed among the Milanese, which consists in the skin becoming covered with wrinkles, and assuming a scaly appearance, especially in the parts exposed to the air. The strength diminishes, the intellectual faculties and sensations become obscure, and cramps, convulsions, and other lesions of muscular contractility supervene. It is a singular endemic affection, and has been supposed to follow everywhere the introduction of Indian corn! Hence the name *Roph'ania maiz'tica*, given to it by some of the Italian physicians.

An affection endemic in the Asturias is said to be a variety of pellagra. It is the *Ro'sa seu Le'pra Astu'rica*, *Mal de la Rosa*.

PEL'LENS, (*pellere*, 'to drive,') Driving; impelling.

PELLEN'TIA MEDICAMEN'TA are medicines that promote the occurrence of the menstrual or hæmorrhoidal flux.

PELLET, (from *pila*, 'a ball,') see Globule.

PELLICLE, Pellicula.
PELLICULA, *Pell'icle*, (dim. of *pellis*, 'a skin.') A thin skin or membrane; a film.

PELLICULA OVI, Membrana putaninis—p. Summa, Epidermis—p. Superior, Epidermis.

PELLICULÆ CORDIS, Ostiola cordis.

PELLIS, Cutis—p. Summa, Epidermis.

PELLITORY, Xanthoxylum fraxineum—p. American, Parietaria Pennsylvanica—p. Bastard, Achillea ptarmica—p. of Spain, Anthemis pyrethrum—p. Wall, Parietaria.

PELMA, (πελαα,) Sole.

PELOHÆMIA, (F.) *Pélohémie*, (πελος, 'dark,' and 'αίμα,' 'blood,') A condition of the blood in which it is thick, syrupy, and of a very dark colour—as in gangrene, &c.

PELOR, (πελωρ,) Monster.

PELORIA, (from πελωρ,) Monster.

PELORIUM, (from πελωρ,) Monster.

PELORUM, (from πελωρ,) Monster.

PELOTE DE MER, (from *pila*, 'a round mass,') *Pila marina*.

PELTALIS seu PELTATUS CARTILAGO, (πελτη, 'a small shield,') Thyroid cartilage, Xiphoid cartilage.

PELTAN'DRA VIRGIN'ICA, (πελτη, 'a small shield,' and *ανθη*, gen. *ανθος*, 'anther,' from the shield-shaped stamens.) *Arum Virgin'icum*, *Calla* seu *Lecon'tia* seu *Russelae'ria Virgin'ica*, *Ar'row* *Ar'um*, *Taraho'*, *Wam'pee*; indigenous; of the Order Araceæ. The fresh roots and seeds are acrid stimulants, like *Arum*.

PELTIDEA AMPLISSIMA, (πελτη, 'a small shield,') Lichen caninus—p. Aphthosa, Lichen aphthosus—p. Canina, Lichen caninus—p. Leucorrhiza, Lichen caninus—p. Malacea, Lichen caninus—p. Spuria, Lichen caninus.

PELTIGERA CANINA, Lichen caninus.

PELUDE, Pelicidae.

PELVIC, *Pel'veius*, *Pelvi'nus*, (F.) *Pelvien*, (from *pelvis*.) Belonging or relating to the pelvis.

PELVIC APONEURO'SIS, (F.) *Απονέυρωσε pelvienne*. A tendinous lamina, given off from the sides of the pelvis, and from the entire circumference of the brim, which passes into and lines the pelvis, and is soon divided into two distinct layers;—one external, the lateral pelvic or obturator fascia, which is continued to line the sides of the pelvis, and covers the obturator internus muscle; the other, internal or superior, which passes inwards upon the side of the prostate, bladder, and rectum in the male, and of the bladder, vagina, and rectum in the female, in order to form the floor of the pelvis.

PELVIC CAV'ITY, (F.) *Cavité pelvienne*. The cavity of the pelvis.

PELVIC MEMBERS. The lower extremities.

PELVIC SURFACE OF THE IL'IUM. That which faces the pelvic cavity.

PELLICULA OCULI, (dim. of *pelvis*, 'a basin,') Orbit.

PELVID'YMUS, (F.) *Pelvidyme*, (*pelvis*, and *δευν*, 'to mingle,') A monster united at the pelvis, and double above.

PELVIAN, Pelvic.

PELVIMETER, *Pelycom'eter*, *Pelicom'eter*, *Pelyom'eter*, *Pylom'eter*, (F.) *Pelvimètre*, (*pelvis*, and *μετρον*, 'a measure,') A barbarous hybrid. This name has been given to different instruments, invented for measuring the diameters of the pelvis, and particularly the antero-posterior or fore-and-aft diameter of the brim:—*Pelvim'etry*, (F.) *Pelvimétrie*. Two have been chiefly employed, especially by French practitioners:—the *Compas d'épaisseur*, and the *Pelvimeter* of Coutouly. 1. The *Compas d'épaisseur* or *Call'ipers* of Baudeloque, is formed like a pair of compasses—with blunt extremities—the branches of which can be moved at will. One of these is applied on the symphysis pubis, and the

other on the sacrum. About three inches must be deducted from this measurement for the thickness of the mons veneris, pubis, and the base of the sacrum. 2. The *Pelvimeter* of Coutouly resembles the instruments used by shoemakers for measuring the length of the foot. The two branches are introduced, in a state of approximation, into the vagina, and then separated, so that one touches the promontory of the sacrum, the other comes behind the os pubis. It is a barbarous contrivance, and its introduction ought never to be attempted on the living subject. It does not, indeed, seem possible to introduce it without mischief.

The finger is, decidedly, the best pelvimeter, and by it we can judge whether the base of the sacrum be unusually prominent. Measurement of the pelvis by the hand has been called *Pelycoch'irometre'sis*; (from *πελός*, 'the pelvis,' *χειρ*, 'the hand,' and *μετρον*, 'a measure,')

PELVIMETRY, see Pelvimeter.

PELV'IS, gen. *Pelvis*, 'a basin,' *Cho'anna*, *Py'elus*, *Lec'anē*, (F.) *Bassin*; (from *πελος*, *πελός*, or *πελός*, 'a basin,' because fancied to be shaped like an ancient basin.) The part of the trunk which bounds the abdomen below. It is a large, bony, irregular, conoidal cavity,—open above and below,—which supports and contains a part of the intestines, and the urinary and genital organs; and serves, at the same time, as a fixed point for the articulation of the lower limbs, the attachment of their muscles, and the execution of their movements. The pelvis supports, behind, the vertebral column, and is sustained, before, by the ossa femorum. It is situate, in the adult, near the middle part of the body, and is composed of four broad, flat, unequally thick bones, differing much in their shape, size, and arrangement, which touch, are articulated at some part of their surface, and intimately united by means of a number of ligamentous fasciæ. Of these bones, two are behind, on the median line,—the sacrum and the coccyx; the two others are before and at the sides,—the ilia. They are fellows, and unite, before, with each other. The most important parts of the pelvis, in an obstetrical point of view, are the brim and the outlet. The BRIM, *Angus'tia abdomina'lis*, *Intro'itus*, *Apert'ura pelvis superior*, *Up'per O'pening or Strait of the Cavity of the Pelvis*, (F.) *Détroit supérieur*, *D. abdominal*, is the narrow part which separates the greater pelvis from the less—the false from the true, *Pelvis v'e'ra* seu *mi'nor*. In the well-formed woman it is elliptical, and slightly inclined forwards. Its antero-posterior, sacro-pubic or conjugate diameter, in a standard pelvis, measures 4½ inches, but with the soft parts, 3½ inches; its transverse or iliac or lateral, 5½ inches, but with the soft parts 4 inches; and its oblique, *Diam'eter of Devent'er*, with the soft parts, 4½ inches. The OUTLET, *Ex'itus*, *Infe'rior o'pening or strait*, *Angus'tia perine'a'lis*, (F.) *Détroit inférieure*, *D. périnéal*, forms the lower aperture of the pelvis. The anteroposterior diameter is here, on account of the mobility of the coccyx, 5 inches: the lateral, 4 inches. The AX'IS OF THE PELVIS is important to be known in obstetrics. The Axis of the Brim is indicated by a straight line drawn from the umbilicus to the apex of the coccyx;—the Axis of the Out'let by a line drawn from the first bone of the sacrum to the entrance of the vagina. An imaginary curved line which indicates the direction of the canal of the pelvis, has occasionally been termed the curve of *Call'rus*, in consequence of its having been pointedly described by that German obstetrician.

PELVIS AURIUM, Cochlea—p. Cerebri, Infundibulum of the brain.

PELVIS OF THE KID'NEY, *Pelvis ren'al's* seu *renum*, *S'anus*, *Ven'ter* seu *Al'veus Re'nium*, (F.) *Bassin*. This is a small, membranous pouch,

occupying the posterior part of the fissure of the kidney. It is placed behind the renal artery and vein; is elongated from above to below; flattened from before to behind; irregularly oval in figure; and, below, contracts considerably, to be continuous with the ureter. It receives the orifices of the infundibula, which pour the urine secreted in the kidney into its cavity.

PELVIS MINOR, Pelvis—p. Ocularis, Scaphium oculare—p. Renalis, Pelvis of the kidney—p. Renum, Pelvis of the kidney—p. Vena, Pelvis.

PELVITROCHANTE'RIAN, *Pelvitrochanterianus*. That which relates to the pelvis and great trochanter. The *Pelvitrochanterian* re'gion is formed by the muscles—*pyramidalis, two obturators, gemini, and quadratus femoris*—which pass from the pelvis to the digital cavity of the great trochanter.

PELYCOCHROMETRESIS, (πελυσ. gen. πελκος, 'pelvis,' χερ, 'hand,' and μετρον, 'measure,') see Pelvimeter.

PELYCOMETER, (πελυσ. gen. πελκος, 'pelvis,' and μετρον, 'a measure,') Pelvimeter.

PELVOMETER, Pelvimeter.

PEM'PHIGAN or PEM'PHIGAN. Meat cured, pounded, and mixed with fat. It has been much used as nutriment on long overland journeys.

PEMPHIGO'DES, (πεμφιγοδής,) *Pemphigoid*, (F.) *Pemphigode*. An epithet given by the ancients to the fever that accompanies pemphigus.

PEM'PHIGUS or PEMPHIGUS, *Emphygis* *Pemphigus*, *Pompholyx*, (from πεμφίς, gen. πεμφγος, 'a blister,') *Febbris bullosa* seu *vesicularis* seu *ampullosa* seu *pemphigodes* seu *pemphigodes*, *Exanthema serosum*, *Morbus*, *Pemphigus mortis* seu *Helveticus* seu *major* seu *minor*, *Morbus bullosus* seu *vesicularis* seu *ampullaceus*, *Pemphix*, *Pemphix*, *Typhus vesicularis*, *Vesicular Fe'ver*, *Bladdery Fe'ver*, *Waterblebs*, (F.) *Fievre bulleuse ou vesiculaire*, *Maladie bulleuse*. A disease, defined to consist of vesicles, scattered over the body; transparent, filbert-sized, with a red, inflamed edge, but without surrounding blush or tumefaction; on breaking, disposed to ulcerate; fluid, pellucid or slightly coloured; fever, adynamic. It is doubtful whether any such idiopathic fever has ever existed: the probability is, that the fever and vesications have been an accidental complication. Cullen thinks the *Pemphigus Helveticus* must have been malignant sore throat. If, however, such a distinct disease did exist, it was probably only as an endemico-epidemic.

PEMPHIGUS GANGRE'NOSUS, *Rupia escharotica*—p. *Hungaricus*, see *Anthrax*—p. *Infantilis*, *Rupia escharotica*—p. *Major*, *Pemphigus*—p. *Minor*, *Pemphigus*—p. *Morta*, *Pemphigus*—p. *Variolodes*, see *Varicella*.

PEMPHIX, gen. *Pemphigis*, (πεμφίς,) *Pemphigus*.

PEMPHYX, (πεμφίς,) *Pemphigus*.

PEMPTÆA FEBRIS, (πεμπτως, 'five,') *Quintan*.

PENÆA MUCRONATA, (after P. Penna, a French botanist,) see *Sarcocolla*—p. *Sarcocolla*, *Sarcocolla*.

PENCHE, Abdomen.

PENCIL LIKE PROCESSES, *Styloid processes*.

PENDULOUS ABDOMEN, *Physeonia*.

PENDULUM PALATI, (from *pendere*, 'to hang,') *Velum pendulum palati*.

PENÉTRANT, Penetrating.

PENÉTRATING, *Penetrans*, (F.) *Péntrant*, (*penetrare*, *penetratum*, (*penitus intrare*,) 'to go into,') A wound is so called which penetrates one of the great splanchnic cavities—*Vulnus penetrans*.

A medicine is also so called, which is supposed to pass through the pores, and stimulate.

PENICILLUM, Compress, *Penicillus*.

PENICILLUS, *Penicillum*, *Peniculus*, ('a painter's brush; dim. of *penis*, 'a tail,') A tent or plectet. A pessary. The secreting glandiform extremities of the *venæ portæ*, (*Ad'ini bilio'si*), have been so termed, as well as the villous textures.

PENICILLUS, Tent.

PENICULUM, Compress.

PENICULUS, *Penicillus*, Tent.

PÉNIDE, *Penidium*.

PENIDES, *Saccharum hordeatum*.

PENID'IUM, *Penidium saccharatum*, (*penis*, 'a tail,' from the shape [?],) (F.) *Pénide*, *Sucre tora*. A kind of clarified sugar made up into rolls. *Barley sugar*. It is demulcent; see *Saccharum hordeatum*.

PÉNIL, *Mons veneris*.

PENIS, gen. *Penis*, 'a tail,' (probably from *pendere*, 'to hang down,') *Caulis*, *Colles*, *Colis*, *Caul'es*, *Cau'los*, *Men'tula*, *Men'ta*, *Phal'lus*, *Pe'los*, *St'e'ma*, *Nat'rix*, *Cap'ulus*, *Rhop'ulum*, *Pos'th'e*, *Pos'thium*, *Pro's'thium*, *Cri'th'e*, *Cor'yn'e*, *Vo'mer*, *Pa'ter on'ium viven'tum*, *Pria'pus*, *Vir'ga*, *V. viri'lis*, *V. genita'lis*, *Ver'etrum*, *Sal'th'e*, *Mem'brum viri'le seu semina'le seu genita'le viro'rum*, *Mo'rion*, *Cy'on*, *Sic'ula*, *Ten'tum*, *Ha'e'ta*, *H. nuptia'lis*, *H. viri'lis*, *Vas*, *Vas'culum*, *Pecu'lium*, *Vir'gula*, *Vir*, *Thyr'sus*, *Ten'sus*, *Ola'vus*, *Cau'da sa'tax*, *Cau'da*, *Fus'cinum*, *P. viri'le*, *Mu'to*, *Ner'veus*, *N. fistulo'sus seu fistula'ris seu juveni'lis*, *Per'tica per se*, *Scap'us*, *Le'co*, *Curcul'io*, *Ve'na*, *Con'tus*, *Tel'm*, *Gurgu'lus*, *Scep'trum*, *Ar'ma*, *Gla'dius*, *Lu'bricum*, *Cap'ut*, *Mu'tinum*, *Pa'lus*, *Pen'sulus*, *Ra'dix*, *Ra'mus*, *Ruta'bulum*, *Ar'ma Ven'tris*, *Colum'na ad'stans inguin'ibus*, *Colum'na*, *Pyra'mis*, *Trabs*, *Sp'i'na*, *Catapul'ta viri'lis*, *Ver'pa*, *Mach'e'ra*, *Tau'rus*, *Pos'te*, *Tur'tur*; the *Yard*, *Male organ*, &c., (F.) *Verge*, *Membre viril*. This organ, the use of which is to carry the seminal fluid into the female organs of generation, is cylindroid, long, and erectile, and situate before and beneath the symphysis pubis. In the ordinary state, it is soft and pendent in front of the scrotum. During erection, it becomes elongated, upright, and assumes a triangular shape. Its upper surface is called the *Dorsum penis*, (E.) *Dos de la verge*; and, at its anterior surface, there is a longitudinal projection formed by the canal of the urethra. The two sides of the penis are round, and its posterior extremity or root is attached to the pelvis. Its anterior extremity is free, and presents the glans, prepuce, and orifice of the urethra. The penis is formed of the corpora cavernosa, the principal seat of erection; the corpus spongiosum of the urethra, for the conveyance of the urine and sperm; and the glans, which terminates the canal. The *Arteries* of the penis are branches of the internal pudic. The *Veins* correspond with the arteries. The *Nerves* are from the internal pudic.

PENIS CEREBRI, Pineal gland—p. *Faetitious*, *Dildo*—p. *Femineus*, *Clitoris*—p. *Lipodermis*, *Paraphimosis*—p. *Muliebris*, *Clitoris*—p. *Succedaneus*, *Dildo*.

PENNATUS, Penniform.

PEN'NIFORM, *Penniformis*, *Penna'tus*, (*penna*, 'a pen,' and *forma*, 'form,') An epithet for muscles whose fleshy fibres are inserted at the side of a middle tendon, like the feathers of a pen on their common stalk. When on one side of the tendon, they are called *simple* or *single penniform*; when on both sides, *double penniform*.

PENNSYLVANIA, MINERAL WATERS OF. The most celebrated are those of Bedford. The Carlisle, Ephrata, Yellow Springs, &c., are also frequented.

PENNYCRESS, *Thlaspi*.

PENNYROYAL, *Mentha pulegium*, *Hedeoma pulegioides*—p. *Hart's*, *Mentha cervina*.

PENNYWORT, ASIATIC. *Hieracotyle Asiatica*.

PENSACOLA, see Saint Augustine.

PENSÉE, *Viola tricolor*—*p. Sauvage*, *Viola tricolor*.

PEN'SILE, *Pen'silis*, (from *pendere*, 'to hang,') Applied to tumours which hang by small roots.

PENSILIA, (*pendere*, 'to hang down,') *Pudibilia*.

PENSY, *Viola tricolor*.

PENTA, (*πεντε*, 'five,') in composition, five.

PENTAM'YRON, (*πεντα*, and *μυρον*, 'ointment,')

An ancient ointment, consisting of five ingredients. These are said to have been—storax, mastich, wax, opobalsam, and unguentum nardium.

PENTAPHAR'MACON, (*πενταφαρμακον*, from *πεντα*, and *φαρμακον*, 'remedy,') Any medicine consisting of five ingredients.

PENTAPHYLLUM, (*πενταφυλλον*, from *πεντα*, and *φυλλον*, 'a leaf,') *Potentilla reptans*.

PENTATEUCH, (**SURG'ICAL**), *Pentateuchus*, (*πεντατευχος*, from *πεντα*, and *τευχος*, 'a book,') which signifies the five books of Moses—Genesis, Exodus, Leviticus, Numbers, and Deuteronomy. By analogy, some surgeons have given the name *Surgical Pentateuch* to the division of external diseases into five classes:—wounds, ulcers, tumours, luxations, and fractures.

PENTATH'ETUM, (*πεντα*, and *τιθημι*, 'to place,') An ancient plaster consisting of five ingredients.

PENULA, ('a cover,') *Ingluvies*.

PENZANCE, **CLIMATE** OF. This is the chief residence of invalids in Cornwall, England, during the winter. It is situate on Mount's Bay, about ten miles from the Land's End. It is $5\frac{1}{2}^{\circ}$ warmer in winter than in London, 2° colder in summer; scarcely 1° warmer in the spring, and only about $2\frac{1}{2}^{\circ}$ warmer in the autumn. It is a very favourable winter residence for the phthisical invalid.

PEONY, *Pæonia*.

PEOS, (*πεις*), *Penis*.

PEOTOMY, *Peotomia*, (from *peos*, and *τομη*, 'incision,') Amputation of the penis.

PEPANISIS, (*πεπασις*, from *πεπαίω*, 'to mature,') Coction, Maturation.

PEPANTICUS, (*πεπαντικός*,) *Maturative*.

PEPASMUS, (*πεπασμος*,) Coction, Maturation.

PEPAS'TIC, *Pepas'ticus*, *Peptic*, *Pep'ticus*, (F.) *Pépastique*, *Peptique*. A medicine supposed to have the power of favouring the concoction of diseases. *Maturative*.

PEPEIRUS, (*πεπεϊρος*,) *Concocted*.

PEPERI, (*πεπερι*,) *Piper*.

PEPINO (S.). A *Cucurbitacea*, which is cultivated in great abundance in the fields of Peru. The pulp or edible part is solid, juicy, and well flavoured; but is apt to disagree.

PEPO, (*πεπων*,) *Cucurbita pepo*—*p. Lagenarius*, *Cucurbita lagenaria*—*p. Vulgaris*, *Cucurbita pepo*.

PEPPER, (*πικερα*,) *Piper*—*p. Black*, *Piper nigrum*—*p. Bush*, sweet, *Clethra alnifolia*—*p. Cayenne*, *Capsicum annuum*—*p. Chilly*, *Capsicum annuum*—*p. Cubeb*, *Piper Cubeba*—*p. Decorticated*, *Fulton's*, see *Piper nigrum*—*p. Guinea*, *Capsicum annuum*—*p. Jamaica*, *Myrtus Pimento*—*p. Japanese*, *Fagara piperita*—*p. Long*, *Piper longum*.

PEP'PER MALAGUET'TA. This term has been applied to the fruit or seeds of several zingiberaceous plants, as well as to the pimento. It is usually considered to be synonymous with *Grana Paradi'si*, *Grains of Par'adise*, and *Grana Guineensis*, *Guinea Grains*.

PEPPER POD, *Capsicum annuum*—*p. Poor-man's*, *Polygonum hydropiper*—*p. Tailed*, *Piper eubeba*—*p. Turnip*, *Arum triphyllum*—*p. Wa't*,

Sedum—*p. Water*, *Polygonum hydropiper*—*p. Water*, of America, *Polygonum punctatum*—*p. White*, *Piper album*.

PEPPERWORT, *Lepidium*.

PEP'SIN, *Peps'ina*, *Pepsin'um*, *Chy'mosin*, *Gastræne*, *Digestive Principle*, (from *πέψις*, 'coction, 'digestion,') A peculiar organic matter, which, in combination with the gastric acid, is considered to form the proper digestive solvent. Its chemical constitution is unknown. It would appear that its presence is necessary to induce changes in the elements of the food, which may enable the gastric acids to exert their proper action on them. One mode of obtaining it is by opening and reversing rennet bags—the fourth stomachs of ruminant animals—and washing them under a thin stream of cold water; scraping off the mucous membrane; reducing it to a pulp, and macerating it in distilled water for twelve hours; filtering, and adding to the liquor a sufficient quantity of acetate of lead; collecting the precipitate, and passing through it a current of sulphuretted hydrogen; filtering again; evaporating at a low temperature, and powdering the dry residue. Pepsin has been given in dyspepsia. A form of it, under the name *Poudre nutritive*, has been much used. A wine of pepsin has been made of calves' rennets 3, sherry Oij, alcohol, Osr. The rennets are washed clean, cut up, and macerated for fourteen days in the wine, shaking often. The alcohol is then added. Dose, a teaspoonful immediately after eating.

PEPSINUM, *Pepsin*.

PEPSIS, (*πέψις*,) *Coction*, *Digestion*.

PEP'TIC, *Pep'ticus*, (*πεπτικός*, 'I digest,') *Pepas'tic*. An agent that promotes digestion, or is digestive. Also, applied adjectively to an article of food that is easy of digestion.

PEPTIC PERSUADER, see *Pilulæ aloës et kinæ kinæ*.

PEP'TONE, *Pep'tonum*, (from *πεπτός*, 'I digest,') A presumed definite combination between the solvent fluid in the stomach, and albuminous matters received into the organ as aliments. The same as *Albuminose*.

PER, 'by or through,' in composition. It often, also, gives intensity to the signification.

PERACEPH'ALUS, (*per*, and *ακεφαλος*, 'acephalus,') (F.) *Péracéphale*. A genus of monsters comprising one of the most extensive groups of the teratological series, characterized by a sufficiently regular form of the body: the absence of upper extremities,—the thorax being more imperfect than in acephali, or even completely null,—the essential conditions, consequently being in their degree of anomaly passing 'beyond' the true type of the acephalous monster.

PERACUTUS, *Catoxys*.

PERARTICULATIO, *Diarthrosis*.

PERATODYNIA, (*πρηπα*, 'a pouch for provisions, a stomach, 'and *ὀδυνή*, 'pain,') *Cardialgia*.

PERCE-CRANE, (*percer*, 'to pierce, 'and *crane*, 'cranium,') *Perforator*.

PERCE-FEUILLE, (*percer*, 'to pierce, 'and *feuille*, 'a leaf,') *Bupleurum rotundifolium*.

PERCE-MOUSE, (*percer*, 'to pierce, 'and *mousse*, 'moss,') *Polytrichum*.

PERCE-MURAILLE, (*percer*, 'to pierce, 'and *muraille*, 'a wall,') *Parietaria*.

PERCE-PIERRE, (*percer*, 'to pierce, 'and *pierre*, 'a stone,') *Cithrum maritimum*.

PERCEPTA, (*percipere*, *perceptum*, (*per*, and *capio*,) 'to perceive, 'receive,') A word used by some writers on hygiene to indicate, in a general manner, the effects of the nervous action on the animal economy: in other words, the sensations, the functions of the mind, and their deterioration or privation.

PERCEPTIBILITY, see *Sensibility*.

PERCEP'TION, *Percep'tio*. The appreciation which the brain has of an impression made upon an organ of sense.

PERCEPTIVITY. The power of perception.

PERCOLATIO, (*percolare*, *percolatum*, (*per*, and *colare*), 'to strain through.') Filtration, Percolation.

PERCOLA'TION, *Percola'tio*. The terms percolation and *displacement* are applied, in pharmacy, to an operation of filtration, which consists in placing any substance, the virtues of which have to be extracted by a menstruum, in a funnel-shaped instrument, having a septum perforated with holes, or its tube stuffed with cotton or tow, and pouring fresh portions of the menstruum upon it until all its virtues have been extracted. The operation is used in the formation of certain infusions, extracts, tinctures, &c. See Ph. U. S. 1863.

An instrument used for this purpose is called a *displacer* or *percolator*.

PERCOLATOR, see Percolation.

PERCUSS'. *Percuterè*, (*per*, and *quaterè*, 'to strike,') (F.) *Peruter*, *Frapper*. To strike upon, with the view of appreciating the resulting sound. To practise percussion.

PERCUS'SION, *Epicrou'sis*, *Percus'sio*. When immediate or direct percussion is made on the chest or abdomen, the more or less perfect resonance is an index of the state of the contained organs; and the physician is thus aided in his diagnosis. For this purpose the part may be struck with the fingers, gathered into a bundle, and their tips placed upon a level. It is better, however, to employ *mediate percussion*, which consists in interposing, between the point of the fingers and the part, the finger of the other hand, or a *pleximeter*, and striking this instead of the naked chest.

PERCUSSION, **AUSCULTATORY**, see Acouphonia.

PERCUTER, *Percuss*.

PERCUTEUR COURBE À MARTEAU (F.), ('curved hammer percussor.') An instrument used by Baron Heurteloup in the operation of lithotripsy, in which a hammer is employed instead of a screw, to force together the blades of the instrument on the foreign body.

PERDO'NIUM. A medicated wine of herbs.

PERETERIUM, (*περητηριον*, from *περαω*, 'to bore,') Trepan.

PERETORIUM, Trepan.

PERFORANS, (*perforare*, (*per*, and *forare*), 'to bore through,') A name given to different muscles, whose tendons pass through intervals between the fibres or tendons of other muscles; hence called *perforated*. See *Flexor Longus Digitorum*, &c. Under the name *Perforating Arteries*, are included, 1. *In the hand*—arterial branches, given off by the profound palmar arch, which traverse the muscles and interosseous spaces. 2. *In the thigh*—three or four arteries, furnished by the profunda, which pass through the openings of the abductor magnus. 3. *In the foot*—the anterior and superior branches of the plantar arch.

PERFORANS CASSERI, see Cutaneous—p. Man's, *Flexor profundus perforans*—p. Profundus, *Flexor longus digitorum pedis profundus perforans*.

PERFORATED SPOT, *Locus perforatus*.

PERFORATING ARTERIES, see Perforans.

PERFORATIO, *Perforation*—p. Cranii, see Perforator—p. Intestinorum, *Enterobrosis*—p. Ventriculi, *Gastrobrosis*.

PERFORA'TION, *Perfora'tio*, (*perforare*, *perforatum*, 'to bore through,') *Anatre'sis*, *Diatre'sis*. An accidental opening in the continuity of organs, either from an external or internal cause.

PERFORATION, *Paracentesis*—p. de *Ŷstomac*,

Gastrobrosis—p. *des Intestins*, *Enterobrosis*—p. Uteri, Uterus, rupture of the.

PERFORATOR, *Perforatorium*, (F.) *Percecrane*. An instrument for opening the head of the fœtus in utero, when it is necessary to diminish its size. The operation is called *Cephalotomia*, *Perfora'tio cranii*. The Perforator, most commonly used, is *Smellie's Scissors*, which present a spear point when the handles are closed; with cutting edges at the sides and exterior, which act when the handles are separated.

PERFORATORIUM, *Perforator*.

PERFORA'TUS, (F.) *Perforé*. That which is pierced. Anatomists have given this name to muscles whose fibres or tendons separate to suffer other parts to pass through them. Such are the *Flexor brevis digitorum pedis*, *F. sublimis perforatus*, and *Coracobrachialis*. Also, to parts that are perforated for any purpose—as the *substantia perforata* of the brain.

PERFORATUS CASSERI, *Coracobrachialis*.

PERFRICATIO, (*per*, and *fricare*, *fricatum*, 'to rub,') see *Iatralcēptice*.

PERFRIC'TIO, *Perfrige'rium*, *Catarrh'is*, *Peripy'is* (*perfrigere*, *perfrictum*, (*per*, and *frigere*), 'to shiver with cold,') Considerable refrigeration. Great sense of cold—shivering.

PERFRIGERATIO, see *Rigor*.

PERFRIGERIUM, *Perfrictio*.

PERFUSIO, (*perfundere*, *perfusum*, (*per*, and *fundere*), 'to pour over,') *Fomentation*—p. *Frigida*, see *Affusion*.

PERI, *περι*, 'about, on all sides, round about, 'also, an augmentative prefix.

PERIÆ/RESIS, (*peri*, and *αιρω*, 'I take away,') A sort of circular incision, which the ancients made in the neighbourhood of large abscesses. The *periæresis* is now only practised in some cases of tumours.

PERIAL'GIA, (*peri*, and *αλγος*, 'pain,') A very violent pain.

PERIAMMA, gen. *Periam'matis*, (*περιαμμα*, from *περι*, and *απτειν*, 'to hang,') *Amuletum*.

PERIANAL, (*peri*, and *ανος*,) *Periproctico*.

PERIAPTON, (*περιαπτος*, 'hung around,') *Amuletum*.

PERIARTERII'TIS, (from *peri*, and *arteriitis*, 'inflammation of an artery,') Inflammation of the sheath of an artery.

PERIBLAST, *Periplast*, (F.) *Périblaste*, (*peri*, and *βλαστος*, 'a germ,') The amorphous matter, which surrounds the *endoblast*, *endoplast*, (F.) *Endoblaste* or cell nucleus, and undergoes segmentation.

PERIBLEMA, gen. *Perible'matis*, (*περιβλημα*, from *peri*, and *βαλλω*, 'to throw,') *Catoblema*.

PERIBLEP'SIS, (*περιβλεψις*, from *peri*, and *βλεπω*, 'I look,') The wild look that accompanies delirium.

PERIB'OLĒ, (*περιβολη*, from *peri*, and *βαλλω*, 'to throw,') The dress of a person. The pericardium. Also, the translation of morbid matters towards the surface of the body.

PERIBRO'SIS, (*περιβρωσις*,) (*περι*, and *βρωσσω*, 'I eat,') 'I eat around,') Ulceration or erosion at the corners of the eyelids.

PERICÆ/CAL, *Pericæc'alis*, (*peri*, and *cæcum*.) That which surrounds the cæcum, hence '*pericæcal abscess*.'

PERICAR'DIAC, *Pericar'diæ*, *Pericar'diul*, *Pericar'diacus*, *Pericar'dicus*, (F.) *Péricardique*. Relating to the pericardium,—as '*pericardial murmur*,' '*pericardial effusion*,' '*pericardial arteries*, veins,' &c. &c.

PERICARDIAL, *Pericardiac*.

PERICARDIQUE, *Pericardiac*.

PERICARDI'TIS, (*pericardium*, and *itis*,) *Inflamma'tio seu Plen'itis pericar'dii*, *Cardi'tis extér'na seu sero'sa seu membrano'sa*, *Exocardi'tis*, *Inflamma'tion of the pericar'dium*, (F.) *Péricardite*,

Inflammation du Péricarde. This is probably the proper appellation for most of those cases which have received the names *Carditis*, *Cardipericarditis*, *Cardopericarditis*, and *Cardiopericarditis*. Along with signs of pyrexia, the local symptoms resemble those of pneumonia. Those which point out that the pericardium is the seat of disease, are the following:—pain, referred to the region of the heart, or the scrobiculus cordis,—sometimes pungent, at others, dull and heavy; palpitation, accompanied with spasmodic twichings in the neighbourhood of the heart, shooting up to the left shoulder; pulsation, and sometimes soreness of the carotids, with tinnitus aurium and vertigo: the breathing is by catches; dyspnoea considerable; pulse jarring, jerking, peculiar; the tongue white, covered with a mucous coat, and the skin often bathed in sweat, as in acute rheumatism. The physical signs during the first period are as follows. The action of the heart is generally evident to the eye, and may be felt by the hand. There is soreness to the touch over the intercostal spaces, and over a small surface in the epigastric region, when the pressure is directed upwards towards the pericardium. Percussion is usually natural, but at times there is dulness. On auscultation, the cardiac movements are found to be frequent, abrupt, jerking, and tumultuous; often irregular and intermittent. The pulse presents corresponding characters. When effusion of lymph has occurred, percussion may be negative, or be but slightly affected. On auscultation, in addition to the preceding signs, there may be one or more of the rubbing or friction sounds resembling the rustling of parchment, or of a sawing or rasping character. In some cases, the sound is like the creaking of new leather. This has been supposed to be pathognomonic of effused lymph. The most important point in the pathology of pericarditis is its connexion with acute rheumatism; and it forms one of the most dangerous occurrences in the latter disease. It may be *acute* or *chronic*: in either case, it is, of course, formidable. The most active depletion must be used; with large doses of opium, counterirritants, and all the means required in the most violent internal inflammations.

PERICARDITIS EXSUDATORIA SANGUINOLENTA, Hæmopericardium.

PERICARDIUM, *Pericar'dion*, (*peri*, and *καρδία*, 'heart.') *Peritolē*, *Membra'na Cor circumplex'a*, *Involucrum* seu *Ar'cula* seu *Cap'sula* seu *Cap'sula* seu *Camer'a* seu *Indumen'tum* seu *Pariet'ulus* seu *Membra'na* seu *The'ca* seu *Sac'cus* seu *Sac'culus* seu *Sero'tum* seu *Vagi'na* seu *Vesi'ca* seu *Ar'ca* seu *Thal'amus* *rega'tis* seu *Amphies'ma* seu *Bur'sa* *Cor'dis*, *Sac* or *Cap'sule* of the Heart, (F.) *Péricarde*. The pericardium is a membranous sac, which envelops the heart, and the arterial and venous trunks that pass from or into it. It is seated in the separation of the mediastinum, above the central aponeurosis of the diaphragm, to which it strongly adheres. Its shape is triangular, like that of the heart, to which it is fitted. The pericardium is composed of two membranes:—the one—*external*—fibrous; the other—*internal*—serous. The latter, having lined the inner surface of the external layer, is reflected over the heart, and covers it entirely; (*Epicar'dium*, (F.) *Épicarde*), without, however, having the heart within it; in which arrangement it resembles other serous membranes. The pericardium envelops the heart; retains it in position, and facilitates its movements by means of the serous fluid contained in its cavity—*Ca'vum pericar'dii*—and secreted by the membrane—*Li'quor* seu *Aqua* seu *Hu'mor* seu *Lym'pha* seu *Uri'na* *Pericar'dii*, *Hydrocar'dia*, which it contains in greater or less quantity.

The arteries of the pericardium are small, and derived from the superior phrenic, anterior mediastinal and bronchial. The veins accompany them, and open into the brachiocephalic. No nerves have been traced to it. The lymphatics enter the lymphatic glands that surround the vena cava superior.

PERICARPIUM, (*περικαρπιον*, from *peri*, and *καρπος*, 'fruit,') *Epicarpium*.

PERICHOLIA, (*peri*, and *χολη*, 'bile,') *Polycholia*.

PERICHONDRI'TIS, *Inflammat'io perichondrii*, (*perichondrium*, and *itis*.) Inflammation of the perichondrium: hence *Perichondri'tis larynge'a*; inflammation of the perichondrium covering the larynx.

PERICHON'DRIUM, (F.) *Périchondre*, (*peri*, and *χονδρος*, 'a cartilage,') A membrane of a fibrous nature, which covers cartilages that are nonarticular, and bears considerable analogy to the periosteum in organization and uses.

PERICHRISIS, (*περιχρσις*, from *peri*, and *χρω*, 'to anoint,') *Circumlitio*, *Liniment*.

PERICHRISTON, *Circumlitio*, *Liniment*.

PERIC'LASIS, (*περικλάσις*,) (*peri*, and *κλαω*, 'I break,') A comminuted fracture, with denudation of the bone.

PERICLYMENUM, (*περικλυμενον*), *Lonicera periclymenum*—p. *Vulgare*, *Lonicera periclymenum*.

PERICNE'MIA, (*peri*, and *κνημη*, 'the leg,') The parts surrounding the tibia:—the tibia itself.

PERICOLONITIS, *Pericolit'is*, (*peri*, and *colonitis*.) Inflammation of the areolar tissue surrounding the colon.

PERICOLPI'TIS, (*peri*, *κολπος*, 'vagina,' and *itis*.) Inflammation of the areolar tissue surrounding the vagina.

PERICRANIEDE'MA, gen. *Pericranie'demat'is*, (*pericranium*, and *οίδημα*, 'a swelling,') *Œdema* of the head, owing to fluid effused under the pericranium.

PERICRA'NIUM, *Pericra'nion*, *Pericra'nia*, (F.) *Péricrane*, (*peri*, and *κρανιον*, 'the cranium,') The periosteum, which covers the cranium externally.

PERICYSTI'TIS, (*peri*, and *cystitis*.) Inflammation of the areolar tissue separating the bladder from the neighbouring parts.

PERIDER'IA, (*peri*, and *ἐρη*, 'the neck,') Twisting of the cord round the neck of the child.

PERIDES'MIC, *Perides'micus*, (F.) *Perides'mique*, (*peri*, and *ἔσμος*, 'a ligament,') Some nosologists have given this name to ischuria caused by a ligature round the penis or by stricture of the urethra.

PERIDES'MIUM. The delicate areolar tissue covering a ligament.

PERIDIASTOLE, see *Diastole*.

PÉRIDI'DYME, (*peri*, and *ἰδύμης*, 'testicle,') See *Testicle*.

PERIDIDYMI'TIS, (F.) *Pérididymite*, (*peri*, *ἰδύμης*, 'testicle,' and *itis*.) Inflammation of the tunica albuginea testis.

PERID'ROMUS, (*πεύδωρος*, from *peri*, and *δωρος*, 'a course,') The extreme circumference of the hairs of the head.

PERIGLOTTIS, (*peri*, and *γλωττα*, 'tongue,') Epiglottic gland. See *Tongue*.

PERIG'RAPHE, (*περιγραφη*, (*περι*, and *γραφω*, 'I write,') 'circumscription,') The aponeurotic intersections of the rectus abdominis are so called.

PERIHEPATITIS, (*peri*, and *hepatitis*.) See *Hepatitis*.

PERILARYNGI'TIS, (*peri*, and *laryngitis*.) Inflammation of the areolar tissue surrounding the larynx, which not unfrequently ends in *peri laryngeal abscess*.

PERILYMPH, Cotunnus, liquor of.
PERIMENINGITIS, (*peri*, and *meningitis*.) Pachymeningitis.

PERIMENIX, (*peri*, and *μηνιξ*, 'a membrane.') Dura mater.

PERIMYELIS, (*peri*, and *μυελος*, 'marrow.') Medullary membrane.

PERIMYELITIS, (*peri*, and *myelitis*.) Meningitis, spinal.

PERIMYSIUM, *Fas'cia*, (*peri*, and *μυς*, 'a muscle.') The areolar membrane or sheath—*vagina muscularis*—that surrounds a muscle, or its fasciculi. The sheath of the whole muscle is termed *Perimysium externum*; of the fasciculi, *Perimysium internum*.

PERIN, *Per'ia*, (*πριν*, *πηρα*, 'a sac or pouch.') With some this means the scrotum; with others, the testicle; with others, the vagina; with others, the peritonæum; and with others, the anus.

PERINEOCELE, (*perineum*, and *κηλη*, 'rupture.') Hernia, perineal.

PERINEORRHAPHIA, *Perinæorrhaphy*, (*perineum*, and *ρᾶψη*, 'suture.') Suture of the perineum, (F.) *Suture du Périnée*, in cases of laceration of the same.

PERINÆUM, (*πριν*, gen. *πρινος*, 'the scrotum' [?].) *Interfeminium*, *Interfeminium*, *Interfeminium*, *Interfemina*, *Pter feminium*, *Pe'ris*, *Pe'rin*, *Perine'um*, *Gressura*, *Plé'chas*, *Am'phiplex*, *Mesos'celon*, *Mesos'celus*, *Perine'on*, *Perinæ'os*, *Perine'os*, *Tra'mē*, *Tra'mis*, *Tau'rus*, *Interforamine'um*, *Cochonē*, *Plé'chos*, *Mesome'ron*, *P'e'men*, *Re'gio*, *Perinæ'i*, (F.) *Périnée*. The space at the inferior region of the trunk, between the ischiatic tuberosities, anus, and genital organs. It is smaller in the female than in the male; has a triangular shape, and is divided into two equal parts by a median line called *Ra'phē*. It is occasionally ruptured in labour. At times it has been made to extend posteriorly as far as the os coccygis. The part between the pudendum and anus is sometimes called *anterior perinæum*, to distinguish it from that which extends from the anus to the coccyx, called *posterior perinæum*.

PERINÆUS, Perineal.

PERINE'AL, *Perinæ'us*, *Perinæ'al'is*, *Perineal'is*. That which relates or belongs to the perineum, as *Perineal artery*, *P. hernia*, &c.

PERINEAL APONEUROSIS, P. Fascia.

PERINEAL ARTERY, *Arteria perinæ'al'is*, *Superficial artery of the Perinæum*. A branch of the internal pudic distributed to the perineum.

PERINEAL CUTANEOUS NERVE, See Sciatic nerve, lesser.

PERINEAL FAS'CIA, *Fas'cia perinæ'i*, *Perine'al aponeurosis*. The fasciæ,—*superficial and deep-seated*,—*Perine'al ligament*—*Cam'per's ligament*, *Triangular ligament*,—which belong to the perineum.

PERINEAL FOS'SA, *Fos'sa perinæ'i*, *Ischiorectal fossa*. A conical fossa, the base of which corresponds with the skin; is formed anteriorly by the transversus perinæi muscle; behind, by the inferior border of the glutæus maximus; internally, by the levator ani; and externally, by the tuber ischii. It is filled up with fat and fibrous strise.

PERINEAL ISCHURIA, *Ischuria Perine'al'is*. A name given by Sauvages to retention of urine, caused by a tumour seated in the perineum.

PERINEAL LIGAMENT, see P. fascia.

PERINEAL NERVE. A branch of the internal pudic; which is mainly distributed to the perineum and scrotum of the male, and to the vulva and perineum of the female.

PERINEAL REGION, *Perinæum*.

PERINEAL SEC'TION, *Sec'tio perinæ'al'is*. An incision into the perineum for the cure of stricture of the urethra.

PÉRINÉE, *Perinæum*.

PERINEORRHAPHY, *Perinæorrhaphia*.

PERINEPHRITIS, (F.) *Phlegmon périnéphrétique*, (*peri*, and *nephritis*.) Inflammation of the external areolar and fibrous membranes of the kidney, or of their investing adipose areolar tissue.

PERINEPHRUS, (*peri*, and *νεφρος*, 'kidney.') See Kidney.

PERINEURION, (*peri*, and *νευρον*, 'a nerve.') Neurilemma.

PÉRINÉVRE, *Neurilemma*.

PERINCYTIDES, (*peri*, and *νυξ*, gen. *νυκτος*, 'night.') An eruption appearing at night and disappearing by day.

PERIOD, (*περιόδος*), *Peri'odus*, *Period'e'ia*, *Periodeu'sis*, *Sta'dium*, *Circu'itus*, (F.) *Période*, (*peri*, and *ὅδος*, 'way.') *Cir'cuit*. *Periods* are the different phases or revolutions of a disease; or the epochs which are distinguishable in the course of it. Three periods are commonly enumerated. 1. The augmentation, increase, or progress, (*Incrementum*;) 2. The acmé or height, (F.) *État*, (*Sta'tus*;) and 3. The decline, (*Decrementum*.) Some authors reckon only the invasion and termination.

Period is sometimes used in describing an intermittent, for the time between the commencement of a paroxysm and that of the next, including the fit as well as interval.

PERIODS, MONTHLY, Menses.

PERIODEIA, (*περιόδια*), *Period*.

PERIODEUSIS, *Period*.

PERIODEUTES, (*περιόδευτης*, from *peri*, and *ὄδευς*, 'to travel.') Charlatan.

PERIODICITY, *Periodic'itas*, *Rhythm*, (F.) *Périodicité*. The aptitude of certain physiological or pathological phenomena, in health or disease, to recur at particular periods, after longer or shorter intervals, during which they cease completely. Diseases thus recurring are called *periodical* or *rhythmical*,—*Typo'ses*.

PERIODOL'OGY, *Periodolog'ia*, (*period*, *ὅδος*, 'period,' and *λογος*, 'a discourse.') The doctrine of periodicity in health and disease.

PERIODONTITIS, (F.) *Périodontite*, *Inflammation de la Membrane alvéolo-dentaire*, (*peri*, *ὄδους*, gen. *ὀδοντος*, 'a tooth,' and *itis*.) Inflammation of the membrane that lines the socket of a tooth.

PERIODONTITIS GINGIVARUM, *Ulitis*.

PERIODOSCOPE, (*period*, and *σκοπεω*, 'I view.') An instrument, proposed by Dr. Tyler Smith, for the ready calculation of the periodical functions of the sex. It consists of a movable circular dial, upon which the months and days are engraved, fixed on a pivot in the centre of a large plate on which are numbered the different conditions of the reproductive system, as conception, abortion, premature labour, hemorrhage, labour, &c. By a knowledge of the date of conception, say November 14th, and fixing the movable plate opposite the point on the fixed plate which indicates conception, the observer is enabled at once to see that, allowing 280 days for gestation, labour may be expected about the 20th of August. The dial is made of card-board, and is affixed to a small volume, in which its uses and applications are explained.

PERIODUS LUNARIS, 'a lunar period.' See Menses—p. Morbi, Type—p. Sanguinis, Circulation.

PERIODYN'IA, (*peri*, and *ὄδυν*, 'pain.') A violent and extensive pain.

PERION, (*peri*, and *ωον*, 'ovum.') Decidua.

PERIORBITA, *Periosteum orbitæ*, (F.) *Orbit'or bite*, (*peri*, and *orbita*, 'the orbit.') An appellation used by some for the periosteum lining the orbit, which is a continuation of the dura mater. The inflammation of this periosteum is termed

Periorbitis or *Periorbititis*, and *Inflammati'o periorbitæ* *or* *orbitæ*.

PERIORBITIS, (*periorbita*, and *itis*.) See *Periorbita*.

PERIORBITITIS, see *Periorbita*.

PERIOSTEITIS, *Periostitis*.

PERIOSTEOPHYTE, (*periosteum*, and *φύω*, 'I grow.') An osseous formation, commencing in the periosteum.

PERIOSTEOTOMY, *Periosteotomia*, (*periosteum*, and *τομή*, 'incision.') The operation of dividing the periosteum.

PERIOSTEUM, (*periosteon*.) *Perios'teon*, *Perios'teos*, *Perios'tium*, *Circumossal'le*, *Membrana oss'is*, *Omentum oss'ium*, *Circumossal'is Membrana*, (F.) *Périoste*, (*peri*, and *οστέον*, 'a bone.') The periosteum is a fibrous, white, resisting medium, which surrounds the bones every where, except the teeth at their coronæ, and the parts of bones that are covered with cartilage. The external surface is united in a more or less intimate manner to the neighbouring parts by areolar tissue. Its inner surface covers the bone, whose depressions it accurately follows. It is united to the bone by small, fibrous prolongations; and especially by a prodigious quantity of vessels, which penetrate their substance. The periosteum unites the bones to the neighbouring parts. It assists in their growth, either by furnishing, at its inner surface, as M. Bécord demonstrated, an albuminous exudation, which becomes cartilaginous, and at length ossifies;—or by supporting the vessels which penetrate them to carry the materials of their nutrition. See *Medullary membrane*.

PERIOSTEUM INTERNUM, *Medullary membrane*—p. *Orbitæ*, *Periorbita*.

PERIOTIC, *Periosticus*, (F.) *Périostique*. Relating or appertaining to the periosteum, or to parts surrounding a bone.

PERIOSTITIS, *Periostitis*, *Inflammati'o periorbitæ*, (*periosteum*, and *itis*.) Inflammation of the periosteum.

PERIOTOMEDULLITIS, (F.) *Périostomédulite*, (*periosteum*, *medulla*, 'marrow,' and *itis*.) Simultaneous inflammation of the periosteum and medulla spinalis.

PERIOTO'SIS, (*peri*, *osteon*, and *osis*), (F.) *Périostose*. Tumour of the periosteum. Tumours are so called which are developed on bones, and formed by their outer membrane. These tumours are more rapidly developed, have a less degree of consistence, and sooner disperse than exostoses, which they resemble greatly.

PERIOTOSTEITIS, (F.) *Périostostéite*, (*periosteum*, *οστέον*, 'a bone,' and *itis*.) Simultaneous inflammation of the periosteum and bone.

PERIPHACUS, (*peri*, and *φακος*, 'lens.') See *Crystalline*.

PERIPHERAD, see *Peripheral aspect*.

PERIPHERAL, *Peripheric* or *Periph'eric*, *Peripher'ical*, (*peri*, and *φέρω*, 'I bear.') Relating or appertaining to the periphery or circumference.

PERIPHERAL ASPECT. An aspect towards the circumference of an organ. *Periph'erad* is used adverbially to signify 'towards the peripheral aspect.'

PERIPHERAL VASCULAR SYSTEM, *Capillary system*.

PERIPHIMOSIS, *Paraphimosis*.

PERIPHLEBITIS, (*peri*, and *phlebitis*, 'inflammation of a vein.') Inflammation of the sheath of a vein.

PERIPLAST, (*peri*, and *πλασσω*, 'to form,') *Periblast*.

PERIPLEUMONIA, (*περιπνευμονία*.) *Peripneumonia*.

PERIPLEURITIS, (*peri*, and *pleuritis*.) In-

flammation of the subserous areolar tissue of the costal pleura.

PERIPLOCA INDICA, (*peri*, and *πλωειν*, 'to twine,') *Hemidesmus Indicus*—p. *Mauritiana*, *Scammony*, *Bourbon*.

PERIP'LYSIS, (*περιπλυνσις*, from *peri*, and *πλυνσις*, 'the act of washing;') *Profluvium*. A copious discharge from any part, especially from the bowels.

PERIPNEUMONIA, *Peripleumonia*, *Peripneumonia*, *True peripneumonia*, (*περιπνευμονία*, from *peri*, and *πνευμων*, 'the lung,') Inflammation of the substance of the lungs. See *Pneumonia*.

PERIPNEUMONIA BILIOSA. Inflammation of the lungs, accompanied with bilious fever.

PERIPNEUMONIA CATARRHALIS. *Bronchitis* or *pulmonary catarrh*, with pain in some part of the chest. *Peripneumonia notha*.

PERIPNEUMONIA NOTHA, *P. spuria* seu *pituitosa* seu *catarrhalis*, *Pneumonia notha*, *Pseudoperipneumonia*, *Pseudopneumonia*, *Pseudopneumonitis*, *Bronchoacé*, *Emphothorax*, *Obstructio pulmonum pituitosa febrilis*, *Bronchitis asthenica*, *False* or *Bastard Peripneumony*, (F.) *Péri-pneumonie bâtarde* ou *fausse*, *Fausse pneumonie*. An inexact name, under which some affections are comprised that resemble pneumonia; and, especially chronic bronchitis with pleurodynia. See *Bronchitis* (chronic).

PERIPNEUMONIA PITUITOSA, *P. notha*.

PERIPNEUMONITIS, *Peripneumonia*.

PERIPNEUMONY, *Pneumonia*—p. *Bastard*, *Peripneumonia notha*.

PERIPNEUMONY, LA'TENT. *Peripneumony* whose symptoms are so obscure as to be recognized with difficulty.

PERIPROC'TIC, *Periprocticus*, (F.) *Périprouctique*, (*peri*, and *πρωκτος*, 'anus,') Applied to organs or lesions around the anus.

PERIPROCTYTIS, (*peri*, *πρωκτος*, 'the anus,' and *itis*.) Inflammation of the areolar tissue surrounding the rectum.

PERIPSYXIS, (*περιψυξις*, from *peri*, and *ψυχω*, 'to become cold,') *Cataplexis*, *Perfrictio*.

PERIPYEMA, (*περιπυημα*, from *peri*, and *πυον*, 'pus,') Suppuration around an organ;—a tooth for example.

PERIRRHOEA, *Perirrhoë*, (*περιρροια*, from *peri*, and *ῥεω*, 'I flow,') Afflux of fluids from every point of the body towards an organ which has to remove them from the economy. Also, *enuresis*.

PERIS, (*περις*.) *Perin*.

PERISCELIS, (*peri*, and *σκελος*, 'leg,') *Jarretière*.

PERISCYPHISMUS, *Periscythismus*.

PERISCYTHIS'MUS, (*περισκυθισμος*.) *Periscythosis*, *Periscythismus*, *Periscythitis*, *Periscyphismus*, *Scalping*, (*περισκυθισμος*, from *peri*, and *Σκυθος*, 'a Scythian,' that is, 'scalping after the manner of the Scythians,') An operation described by Paulus of Aëgina. It consisted of an incision made around the cranium, and was employed in habitual weakness of the eyes, pains of the head, &c.

PERISP'HAL'SIS, *Circumductio*, (from *peri*, and *σφαλλω*, 'I move,') A circular motion impressed on a luxated bone for the purpose of reducing it.

PERISPLENITIS, (*peri*, and *splenitis*.) see *Splenitis*.

PERISSODACTYLUS, (*περισσος*, 'superabundant,' and *δακτυλος*, 'finger,') *Polydactylus*.

PERISTAL'TIC, *Peristalticus*, (*περισταλτικός*, from *peri*, and *σπellaω*, 'to contract,') (F.) *Péristaltique*. Relating or appertaining to peristole,—as 'peristaltic action.' See *Systaltic*.

PERISAPHYLINUS EXTERNUS, (*peri*, and *σφαυλην*, *υγία*,) *Circumflexus musculus*—p.

Inferior, Circumflexus musculus—p. Internus, Levator palati.

PERISTERA ANCEPS, (περιστερα, 'a pigeon,') Pelargonium anceps.

PERISTERIUM, (περιστεριον,) Verbena officinalis.

PERISTER'NA, (peri, and στερον, 'the sternum,') The lateral parts of the chest.

PERIS'TOLĒ, (περιστολη,) *Motus Peristalticus* seu *compressorius seu vermicularis seu testudinæus*, (from *peri*, and στελλω, 'to contract,') The peristaltic action of the intestines, *Motus peristalticus*, (F.) *Mouvement ou Action péristaltique*. It consists of a kind of undulation or vermiform movement, in appearance irregular, but in which the circular fibres of the muscular membrane of the intestine contract successively, from above downwards, in proportion as the chyme advances in the alimentary canal; so that, being compressed above, it is pushed into the next portion of the intestine, whose fibres are in a state of relaxation. When the fibres contract inversely, they occasion an *antiperistaltic action*. The peristaltic action is involuntary, and is not under the immediate influence of either brain or spinal marrow. It continues for some time after death.

PERISTOMIUM, (περιστομιον, from *peri*, and στομα, 'mouth,') See Mouth.

PERISTRO'MA, gen. *Peristro'matis*, *Musculus villosus*, *Crusta membrana'cea seu vermicularis*, (περιστρομα, 'a coverlet, from *peri*, and στρωωμι, 'to spread,') The villous or mucous coat of the intestines.

PERISYS'TOLĒ, (peri, and συστολη, 'contraction,') The interval that exists between the systole and diastole of the heart, which is only perceptible in the dying.

PERITERION, (peri, and τερειν, *terere*, 'to rub, to drill,') Τρέπαν perforatif.

PERITESTIS, (peri, and testis,) Albuginea.

PÉRITOINE, Peritonæum.

PERITOME, (περιτομη, from *peri*, and τεμνω, 'to cut,') Circumcision.

PERITONÆREX'IS, *Peritonæorex'is*, (peritonæum, and ῥηγνυμι, 'to break out,') Hernia formed by the rupture of the peritonæum.

PERITONÆ'UM, (περιτοναϊον, peritonæon, peritonæum,) *Peritonæum*, *Peritonæ'on*, *Peritonæ'ion*, *Peritonæ'os*, *Periton'ion*, *Peritonæ'os*, *Velum'entum abdominalē*, *Tunica præten'sa seu Operimentum præten'sum seu Membra'na abdom'inalis*, *Pé'gion*, *Sil'phac*, *Syl'phac*, *Zé'pach*, *Zi'phac*, (F.) *Péritoine*, (peri, and τεινω, 'I stretch,') A serous membrane, which lines the abdominal cavity, *Peritonæum parietalē*, extends over the greater part of the organs contained there, *Peritonæum intestinalē* seu *viscera'le*, envelops them wholly or in part, and maintains their respective relations by means of different prolongations and ligamentous folds:—the mesentery, epiploon, mesocolon, &c. Like all the serous membranes, the peritonæum, except in the female, is a sort of sac without aperture, which covers the abdominal organs, without containing them within it, and the internal surface of which is smooth, and lubricated by a serous fluid. In the male fœtus, the peritonæum furnishes a prolongation, which accompanies the testicle at the time of its descent, and, in the female fœtus, forms the *Canal' of Nuck*. Below the neck of the gall-bladder, the peritonæum has a triangular opening, called the *Fora'men* or *Hid'rus of Win'slow*, through which it proceeds behind the stomach, to form a sort of secondary cavity, called the *posterior cavity of the peritonæum*, (F.) *Arrière cavité péritonéale ou Cavité des Épiploons*.

PERITONÆUM DUPLICATUM, Epiploon, gastrocolic.

PERITONÆ'AL, *Peritonæus*, (F.) *Péritonéal*.

That which appertains or relates to the peritonæum.

PÉRITONITE, Peritonitis—p. *Puerpérale*, see Peritonitis.

PÉRITONITIS, *Empres'ma seu Caum'a Peritonit'is*, *Peritonæit'is*, *Inflammat'io seu Phleg'mon' Peritonæ'i*, *Phlegma'sia* or *Inflammation of the Peritonæum*, (peritonæum, and itis,) (F.) *Péritonite*, *Inflammation du Péritoine*. The characteristic signs of acute inflammation of the peritonæum are,—violent pain in the abdomen, increased by the slightest pressure, often by the simple weight of the bed-clothes. It generally occurs in the parturient state, *Locho-peritonit'is*, and begins on the second or third day after delivery. At times, a malignant epidemic, and perhaps contagious, variety has made its appearance, and destroyed numbers of females. This has been described under the name *Puer'peral Fever*, *Metropéritonit'is*, *Enc'ia Syn'ochus Puerperarum*, *Erythemat'ic* or *Nonplastic Puer'peral Peritonit'is*, *Typhohæmic Peritonit'is*, (F.) *Péritonite puerpérale*, *Métropéritonite puerpérale*, *Fèvre puerpérale*, *Typhus puerpéral*, *Adynam'ic* or *Malignant Puer'peral Fever*, *Low Fever of Child'bed*. It is, according to some, a malignant inflammation of the peritonæum and uterus; according to others, a *U'terine Phlebi'tis*. In any form it requires active treatment, early employed. The appearances on dissection are such as occur whenever a serous membrane has been inflamed, and such inflammation has produced death:—effusion of fluid with flakes of coagulable lymph in it; appearances of vascularity, &c., in the membrane. The treatment—in *active peritonitis*—must consist of the same course as recommended under Enteritis. In the *chronic* kind, symptoms must be combated as they arise, and it may be advisable to excite a new action in the system by means of mercury. In the epidemic and malignant variety, unless bleeding be freely employed at the very outset, it does harm. If used early, it may be of decided advantage. Everything depends upon arresting the morbid process at the outset.

PÉRITONITIS, ERYTHEMATIC, see Peritonitis—p. Hepatic, see Hepatitis—p. Nonplastic, see Peritonitis—p. Puerperal, see Peritonitis—p. Typhohæmic, see Peritonitis.

PÉRITOMA, (περιτωμα, from περισσος, 'over and above,') Excrement.

PÉRITYPHLITIS, (F.) *Pérityphlitis*, (peri, and typhlitis, 'inflammation of the cæcum,') Inflammation of the areolar substance surrounding the cæcum. See Typhloenteritis.

PÉRIÜTERINE, *Periüteri'nus*, (F.) *Périüترین*, (peri, and uterus,) That which is situated around the uterus; hence "*periüterine inflammation*" or "*periüterine phlegmon*,"—inflammation of the areolar tissue around the uterus.

PERIWINKLE, LESSER, ([L.] *pervinca*, [Sax.] *peynince*), *Vinca minor*.

PERIZO'MA, gen. *Perizo'matis*, (περιζωμα, from *peri*, and ζωννυμι, 'to gird,') A bandage; a girdle. In Fabricius Hildanus, a truss. Herpes zoster. The diaphragm.

PERKINISM. A therapeutical means, first employed by Dr. Elisha Perkins, of Norwich, Connecticut, towards the termination of the last century, and named after him. It consisted in drawing over affected or other parts the pointed extremities of two metallic rods, called *Metallic Tractors*, each made of a different metal. The success obtained was through the influence of the imagination; and, consequently, as a remedial agent, it ranks as high as animal magnetism, but no higher. The operation was, also, called *Tractoration*.

PERKINIST. A believer in, and practiser of, Perkinism.

PERKINIS'TIC. Relating or belonging to Perkinism.

PERLA, Pearl. Paralampsis.

PERLE, Pearl.

PERMISSUS, (*permittere, permisum*, from *per*, and *mittere*, 'to suffer to pass through,') Licentiate.

PERMISTIO, (*per*, and *miscere, mistum* or *mixtum*, 'to mix,') Coition.

PERMIXTIO, Coition.

PERMOTIO, (*per*, and *movere, motum*, 'to move,') Motion.

PERNICIEUX, Deleterious.

PERNICIOSUS, (*per*, and *neco*, 'to destroy,') Deleterious.

PERNIO, Chilblain.

PERNOCTATIO, (*per*, and *nox*, gen. *noctis*, 'night,') Insomnia.

PERO, (*προς*, 'deficient,') in composition, deficient.

PEROCEPH'ALUS, (*pero*, and *κεφαλη*, 'head,') A monster with a defective head.

PEROCOR'MUS, (*pero*, and *κορος*, 'trunk,') A malformation in which the trunk is defective, and too short, from the absence of one or more vertebrae.—the head and limbs being normal.

PERODACTYLEUS, (*perone*, and *δακτυλος*, 'a finger or toe,') Flexor longus digitorum pedis profundus perforans.

PERODYN'IA, (*πρια*, 'a sac, stomach,') and *οδυνη*, 'pain,') see Cardialgia.

PEROMEL'IA, (*pero*, and *μελος*, 'a limb,') Congenital misconstruction, or mutilation of the limbs. A genus in the nosology of Good.

PEROM'ELUS. A monster possessed of the deformity mentioned above.

PERONE, (*περονη*, from *πειρω*, 'to transfix,') Fibula.

PÉRONÉ, Fibula—*p. Col du, Collum fibulae.*

PERONE'AL, *Peronæus, Fibula'ris, Fib'ular,* (F.) *Péronier*, (from *perone*.) Belonging or relating to the fibula.

PERONEAL AR'TERY, *Fib'ular, A.*, arises from the popliteal, and is situate deeply in the posterior and outer part of the leg. It gives off branches of little importance. Near the outer ankle, it divides into two branches. 1. The *posterior peroneal*, which descends behind the lower joint of the fibula, on the outer surface of the calcaneum. 2. The *anterior peroneal*, which crosses the inferior extremity of the interosseous ligament, to be distributed on the dorsal surface of the foot.

PERONEAL MUS'LES are three in number. 1. *Peronæus Brevis* seu *Secundus* seu *Anticus* seu *Medius*, *Semifibularis*, (F.) *Grand-péronéo-sus-métatarsien, Court péronéo-latéral, Moyen Péronier.* A muscle, situate beneath the peronæus longus. It is attached, above, to the outer surface of the fibula, and terminates, below, at the posterior extremity of the 5th metatarsal bone, after having been reflected beneath the malleolus externus. It extends the foot upon the leg, at the same time raising a little the outer edge. It may, also, act upon the leg, and extend it upon the foot. 2. *Peronæus Longus* seu *primus* seu *posticus* seu *maximus* seu *posterior*, (F.) *Tibi-peronéo-tarsien, Long péronier latéral, Péronéo-sus-tarsien*, (Ch.) *Muscle grand péronier.* This muscle is situate at the outer part of the leg. It is long, thick, and triangular above; thin, narrow, and flat below. It is attached above to the outer edge of the upper extremity of the fibula, and to the upper third of the outer surface of that bone. Below, it terminates at the outer part of the posterior extremity of the first metatarsal bone. This muscle extends the foot on the leg, turning its point outward, and raising its outer edge. It acts, also, upon the leg, which it ex-

ten is on the foot. 3. *Peronæus Tertius, Nodus Vasa'lii*, (F.) *Péronier antérieur, Paît péronéo-sus-métatarsien* (Ch.) *Petit Péronier.* A muscle, situate at the anterior, outer, and inferior part of the leg. It is long and flat, and is attached, above, to the inferior third of the anterior margin and inner surface of the fibula; terminating, below, at the posterior extremity of the 5th metatarsal bone. This muscle bends the foot on the leg, by raising its outer edge. It can, also, bend the leg on the foot.

PERONEAL NERVE, see Popliteal nerves.

PERONEAL VEIN follows the same course as the artery.

PERONEDACTYLIUS, (*perone*, and *δακτυλος*, 'a finger or toe,') Flexor longus digitorum pedis profundus perforans.

PÉRONÉO-LATÉRAL COURT, *Peronæus brevis*—*p. Phalangien du gros orteil, Flexor longus pollicis pedis*—*p. Sus-phalangien du pouce, Flexor longus pollicis pedis*—*p. Sus-tarsien, Peronæus longus*—*p. Sus-métatarsien, Peronæus tertius*—*p. Sus-métatarsien, grand, Peronæus brevis*—*p. Sus-phalangien du pouce, Extensor proprius pollicis pedis*—*p. Sus-phalangien du pouce, Extensor proprius pollicis pedis*—*p. Tibi-sus-phalangien commun, Extensor communis digitorum pedis.*

PERONEUM, (from *perone*), Fibula.

PÉRONIER, Peroneal—*p. Grand, Peronæus longus*—*p. Latéral, long, Peronæus longus*—*p. Moyen, Peronæus brevis*—*p. Petit, Peronæus tertius.*

PERONODACTYLIÆUS, (*perone*, and *δακτυλος*, 'a finger or toe,') Flexor longus digitorum pedis profundus perforans.

PERONODACTYLIUS, Flexor longus digitorum pedis profundus perforans.

PEROSOMUS, (*pero*, and *σωμα*, 'body,') A monster whose whole body is imperfectly developed.

PEROSPLANCH'NICA, (*pero*, and *σπλαγχων*, 'a viscus,') Congenital misconstruction of the viscera. A genus in the nosology of Good.

PERPENDICULUM HEPATIS, (*perpendicularum*, 'a plumb line,' from *per*, and *pendere*, 'to hang,') Suspensory ligament of the liver.

PERPERACUTUS, 'Extremely acute.' An epithet applied to very violent and rapid diseases.

PERPLEXI MORBI, (*per*, and *plectere, plexum*, 'to interweave,') see Complication.

PERPLICATION, *Perplicitio*, (*per*, and *plico, plicatum*, 'to fold,') A method of tying arteries, which consists in making a small incision in the side of the artery, near its bleeding orifice, introducing a small pair of forceps, seizing the open extremity, and drawing it backward, through the aperture made in the side of the vessel, so as to form a kind of knot.

PERROSIN, see Pinus abies.

PERRY, see Cider.

PERRY COUNTY WARM SPRINGS. These are in Pennsylvania, eleven miles from Carlisle, and fourteen from Harrisburg. Temperature 67°; impregnation not much. They are used for bathing.

PERSEA CAMFORA, (*περσεα*,) see Camphor—*p. Cassia, Laurus cassia*—*p. Cinnamomum, Laurus cinnamomum*—*p. Cubeba, Piper cubeba.*

PERSEA GRATIS'SIMA, *Laurus Persea, Patta tree, Avocado* or *Al'ligator Pear.* A slender, very high tree of Peru, with a small domelike top, which grows on the eastern declivity of the Andes, and rises, at times, to the height of more than 60 feet. Its fruit—(S.) *Patta*—which is pear-shaped, is generally much liked. It dissolves, like butter, on the tongue, and hence is called, in some of the French colonies, *beurre végétale*. The leaves are reckoned balsamic, pectoral, and vulnerary. The seeds are very astringent.

PERSEA PICHURIM, see Pichurim beans—p. Sassafras, *Laurus sassafras*.

PERSIAN FIRE, see Anthrax.

PERSICA VULGARIS, (*Περσικα*, 'Persian,') *Amygdalus Persica*.

PERSICAIRE ACRE, Polygonum hydropiper —p. *Brillante*, Polygonum hydropiper—p. *Douce*, Persicaria—p. *Poirée*, Polygonum hydropiper.

PERSICARIA, (from *persica*, 'peach-tree,' the blossoms being similar,) Polygonum Persicaria—p. Minor, Polygonum Persicaria—p. Mitis, Polygonum Persicaria—p. Urens, Polygonum hydropiper—p. Wandspiked, Polygonum Virginianum—p. Water, Polygonum amphibium.

PERSIL, (abbreviation of *petrosil*, from *petroselinum*.) *Apium petroselinum*—p. *d'Âne*, Chærophyllym sylvestre—p. *Faux*, *Æthusa cynapium*—p. *de Macédoine*, *Bubon Macedonicum*—p. *de Montagne*, *Athamanta aureoselinum*.

PERSIMMON, *Diospyros Virginiana*.

PERSOLATA, *Arctium lappa*.

PERSOLLATA, *Arctium lappa*.

PERSOLUTA, *Arctium lappa*.

PERSPECILLUM, (*per*, and *specio*, 'to behold,') see Spectacles.

PERSPIRABILE SANCTORIANUM, (after Sanctorius,) Perspiration.

PERSPIRATION, *Perspira'tio*, *Transpira'tio*, *Diffusa'tio*, *Diap'noë*, *Diapnoë'a*, *Diapneu'sis*, *Perspirabi'le Sanctoria'nium*, (*per*, and *spirare*, *spiratum*, 'to breathe, exhale,') The insensible transpiration or exhalation continually going on at the surface of the skin and membranes. Sensible perspiration is called *sweat*. The perspiratory fluid is secreted by an appropriate glandular apparatus, termed by Breschet *diapnog'eous*, (F.) *Appareil diapnogène*; the sudorip'arous glands, *Glan'dulæ sudorip'aræ*, *Organa sudorip'ara*, *sweat glands*, consisting of a convoluted tube, situate beneath the true skin, and of spiral ducts, which open obliquely under the scales of the epidermis. A certain portion of the perspired fluid is thrown off by ordinary physical evaporation.

The *Pulmonary exhalation* or *transpiration* is that which takes place into the bronchia, and is sent out with the expired air.

PERSPIRATORY, *Perspirato'rius*, *Transpi'ratory*, *Transpirato'rius*, (F.) *Perspiratoire*, *Transpiratoire*. Relating or belonging to, or the product of, perspiration.

PERSTRICION, (from *perstringere*, (*per*, and *stringere*, *strictum*,) 'to bind,') see Hæmostasia.

PERSUDATIO, (*per*, and *sudare*, *sudatum*, 'to sweat,') Diaporesis.

PERSULTATIO, (*persultare*, *persultatum*, 'to leap through,' from *per*, and *saltare*, 'to leap,') Diapodesis.

PERTE, (from *perdere*, *perditum*, 'to lose,') Metrorrhagia—p. *d'Appetit*, Anorexia—p. *de la Mémoire*, Amnesia—p. *de l'Odorat*, Anosmia—p. *de Sang*, Hæmorrhagia—p. *de la Voix*, Aphonia—p. *de la Vue*, Cæcitas.

PERTÉRÉBRANT, *Pertér'ebrans*, (*per*, and *terebrare*, 'to bore,') The French give this epithet to an acute pain—*Douleur pertér'ébrante*—which occasions a sensation like that which would be produced by an instrument penetrating and lacerating a part;—the pain from whitlow, for example.

PERTES BLANCHES, (pl. of *perte*,) Leucorrhœa—p. *de Sang*, Metrorrhagia—p. *Séminales*, Spermatorrhœa—p. *Uérines*, Metrorrhagia—p. *Uérin'es blanches*, Leucorrhœa—p. *Uérines rouges*, Metrorrhagia.

PERTICA PER SE, ('a pole,') Penis.

PERTRUSIO MUTUA, (*per*, and *trudere*, *trusum*, 'to thrust,') Endosmose.

PERTURBATIONES ANIMI, (*perturbare*,

perturbatum, (*per* and *turbare*,) 'to disturb greatly,') Affections of the mind.

PERTURBATRIS (*Medici'na*,) *Per'turbating treat'ment*, (F.) *Méthode ou Médecine perturbatrice*; a mode of treatment in which very active means are employed, and such as impress upon diseases a different course from that natural to them. It is the antithesis to the *Medicina expectans*. Much evil has resulted from such treatment, especially in febrile diseases.

PERTUSS'AL, *Pertussa'lis*, Relating or appertaining to pertussis.

PERTUSS'IS, (*per*, and *tussis*, 'a cough,') 'a violent cough.' *Tus'sis feri'na* seu *convulsi'va* seu *pu'eros strang'ulans* seu *quin'ta* seu *stomacha'lis* seu *an'hela clamo'sa* seu *stran'gulans* seu *delas'sans* seu *suf'focans* seu *amphemer'ina* seu *tussicu'lo'sa* seu *spasmod'ica* seu *asin'i'na* seu *can'i'na*, *Bronchi'tis convulsi'ca*, *Bronchocephali'tis*, *Pneu'sis pertus'sis*, *Amphemer'ina*, *Orthopne'a tussicu'lo'sa* seu *clango'sa* seu *suffocati'va* seu *clamo'sa* seu *spasmod'ica*, *Mor'bus cucu'llus* seu *cuculla'ris*, *Cucu'llus*, *Bechorthopne'a*, *Bex convulsi'va* seu *therio'des*, *Whoopingcough*, *Whoopingcough*, *Chin'coughs*, *Kin'cough*, *Kind'cough*, (Germ. *Kind*, 'a child,') (Se.) *Kink-host*, (Prov.) *Kinkhaust*, (F.) *Coqueluche*, *Catarrhe* ou *Bronchite convulsive*, *Toux convulsive* ou *quintense* ou *bleue*, *Maladie cuculla'ire*. A violent convulsive cough, returning by fits (F.) *Quintes*, at longer or shorter intervals; and consisting of several expirations, followed by a sonorous inspiration or *whoop*. The fits of coughing generally recur more frequently during the night, morning, and evening, than in the day. It is esteemed to be contagious, and attacks the young more particularly. It is rare for it to effect an individual for the second time. The duration is various,—six or eight weeks or more. Although the paroxysms are very violent, it is not a dangerous disease. It may, however, give rise to other affections, as convulsions, pneumonia, &c., when the complication is very dangerous, as the cause cannot be removed. Those children suffer the least, who evacuate the contents of the stomach during the fit. In the treatment, all that can be done is to palliate. It must be borne in mind, that the disease will, in time, wear itself out. If there be much tensive pain of the head, or fever, bleeding may be required, but it is seldom necessary. Narcotics occasionally afford relief, but it is temporary. Gentle emetics, given occasionally, when the paroxysms are long and dry, give decided relief, and aid in the expectoration of the morbid secretions. After the disease has continued for some weeks, and persists in part from habit, change of air is essential, and this, even should the change be to an atmosphere that is less pure.

PERU, see Lima.

PERUNCTIO, (*per*, and *ungere*, *unctum*, 'to anoint,') Inunction.

PERVENCHE, ([L.] *pervinea*,) *Vinca minor*.

PERVERSION, *Perver'sio*, (*per*, and *vertere*, *versum*, 'to turn,') *Dias'trophê*, *Diastrem'ma*. One of the four modifications of function in disease: the three others being augmentation, diminution, and abolition. The humorists used this term also to designate disorder or morbid change in the fluids.

PERVERSION DE LA TÊTE DES OS ET DES MUSCLES, *Loxarthus*.

PERVIGILIUM, (*per*, and *vigilare*, 'to watch,') Insomnia, *Coma vigil*.

PES, gen. *Pe'dis*, (*πους*, gen. *ποδος*,) *Ich'nus*, (Se.) *Fit*, (F.) *Pied*, 'the foot.' The inferior extremity of the abdominal member, representing a bony arch, an elastic vault, which transmits the weight of the body to the ground. It is divided into three parts;—the *tarsus*, *metatarsus*, and *toes*.

PES ALEXANDRINUS, *Anthemis pyrethrum*—p.

Anserinus, Chenopodium bonus Henricus, Facial nerve—p. Cati, Antennaria dioica—p. Equinus, Pes equinus, see Kylosis—p. Hippocampi major, Cornu ammonis—p. Hippocampi minor, Hippocampus minor—p. Hippopotami major, Cornu ammonis—p. Hippopotami minor, Cornu ammonis—p. Leonis, Alechemilla.

PESANTEUR, (from [F.] *peser*, [L.] *pendere*, *pensum*, 'to weigh,') Weight—p. *Spécifique*, Gravity, specific.

PÈSE-ACIDE, (*peser*, 'to weigh,' and *acid*,) Acidometer—p. *Liqueur*, Areometer—p. *Sirap*, Saccharometer.

PESSETTE, Cicer arietinum.

PESSAIRE, Pessary—p. *Intra-utérin*, Pessary, intra-uterine.

PES'SARY, *Pessa'rium*, (from *πῆσος*, 'a small stone or other substance for playing at draughts; a pessary,') *Pes'us*, *Pes'sum*, *Pes'sulus*, *Penicillus*, *Talus*, *Nodulus uterinus*, *Suppositorium uterinum*, *Glane*, *Lemni'cus*, (F.) *Pessaire*. A solid instrument, composed of cork, ivory, or elastic gum, which is introduced into the vagina to support the uterus, in cases of prolapsus or relaxation of that organ; to keep vaginal hernia reduced, &c. The ancients made use of *medicated pes'saries*, which they distinguished into emollient, astringent, aperient, &c., and they are still used. The form of pessaries is very variable;—round, oval, &c., according to fancy and circumstances.

PESSARY, EGYPTIAN, *Ægyptius pessus*.

PESSARY, INTRA-UTERINE, *Stem pes'sary*, (F.) *Pessaire intra-utérin*, *Redresseur utérin*. An instrument for rectifying uterine displacements—as anteversion and retroversion, consisting of a stem which is introduced into the uterus; this stem is attached to a flattened ovoid bulb or ball, on which the cervix uteri rests.—Connected with this is an external part or wire frame, which is attached, at one extremity, to a flat tubular portion, passing into the vagina, to be fixed to the intra-uterine portion. The wire frame is made to press upon the pubis, so that the pessary can be kept in position *in utero*. This is essentially the intra-uterine pessary of Professor Simpson of Edinburgh.

A modification of Prof. Simpson's instrument for rectifying malpositions of the uterus has been made by Dr. Sims, of New York. This consists in having a joint or hinge near enough the end to prevent its striking against the fundus uteri, and yet far enough to allow it to enter fully the cavity; making the os uteri the centre of motion. It is called by Dr. Sims the *uterine elevator*.

PESSARY, STEM, Pessary, intra-uterine.

PESSULUS, (dim. of *pessus*,) Pessary, Penis.

PESSUM, Pessary.

PESSUS, Pessary.

PEST, (*pestis*,) Plague.

PESTE, Plague—p. *Antonine*, Plague of Antoninus—p. *d' Athènes*, Plague of Athens—p. *Noire*, Death, Black.

PESTICHIÆ, *Petechiæ*.

PESTICLE, *Petechiæ*.

PES'TILENCE, *Pestilen'tia*, *Pestill'itas*, (from *pestis*, 'a pest,') A malignant spreading disease;—applied especially to plague.

PESTILENCE, CHOLEMIC, Cholera spasmodica—p. *Glandular*, Plague—p. *Hæmagastic*, Fever, yellow—p. *Septic*, Plague.

PESTILENTWORT, Tussilago petasites.

PESTILENTIA, Pestilence, (from *pestis*,) Plague—p. *Hæmagastica*, Fever, yellow.

PESTILENT'IAL, *Pestilentia'lis*, *Pestilentia'rins*, *Pestilentio'sus*, *Pestilen'tus*, (F.) *Pestilential*. Relating or appertaining to a pest or plague. *Pestilential diseases* are those that bear some resemblance to the plague in respect to mode of propagation, symptoms, danger, &c.

PESTILITAS, (from *pestis*,) Plague.

PESTIS, Plague—p. *Acutissima*, Plague—p. *Adeno-septica*, Plague—p. *Americana*, Fever, yellow—p. *Antoniniana*, Plague of Antoninus—p. *Bellica*, Typhus—p. *Britannica*, Sudor Anglicus—p. *Bubonaria orientalis*, Plague—p. *Contagiosa*, Plague—p. *Glandulosa*, Plague—p. *Glossanthrax*, Glossanthrax—p. *Inguinaria*, Plague—p. *Inter-tropica*, Fever, yellow—p. *Nigra*, Death, black—p. *Occidentalis*, Fever, yellow—p. *Orientalis*, Plague—p. *Septica*, Plague—p. *Variolosa*, Variola.

PESTLE, (from *pistillum*,) Pilum—p. Spring, see Pilum.

PET, (from *pedere*,) Faru.

PETALE, Phtheiriasis.

PÉTASITE, Tussilago petasites.

PETASITES, (*πετασῖτις*, from *πετασος*, 'a sun hat,') Tussilago petasites—p. *Hybrida*, Tussilago petasites—p. *Major*, Tussilago petasites—p. *Officinalis*, Tussilago petasites—p. *Rubens*, Tussilago petasites—p. *Vulgaris*, Tussilago petasites.

PETAUR'UM, (*πεταυρον*,) A seat suspended by ropes, in which a person taking exercise sat, and was tossed about by assistants.

PETE'CHIE, *Petic'ule*, *Pestich'ix*, *Pestic'ix*, *Punctic'ula*, *Pur'pura maligna*, *Por'phyra Græcorum*, (F.) *Pétéchies*, (from [L.] *Pedechio*, 'a flea-bite,') Small spots, similar in shape and colour to flea-bites, which occur spontaneously upon the skin, in the course of severe fevers, &c. They are, under such circumstances, signs of great prostration.

PETECHIE SINE FEBRE, *Purpura simplex*.

PETE'CHIAL, *Petechi'alis*. Resembling or accompanied by petechiæ.

PETECHIAL SCURVY, *Purpura simplex*.

PETECHIANOSIS, (*petechiæ*, and *νσος*, 'disease,') *Purpura hæmorrhagica*.

PETER'S PILLS, *Pilulæ Aloes et Cambogiæ*—p. Worm lozenges, see Worm lozenges, Sherman's.

PETICULÆ, *Petechiæ*.

PETIGO, Impetigo.

PETINA, (from *pes*, gen. *pedis*, 'a foot,' or from *peto*, 'to go,') Sole.

PETI'OLUS, *Pedi'olus*, (dim. of *pes*, gen. *pedis*, 'a foot,') A footstalk or leafstalk of a plant. A petiole.

PETIOLUS EPIGLOT'TIDIS. The root of the Epiglottis.

PETIOLUS MALLEI, *Manubrium mallei*.

PETIT ARGENT, ('small silver,') Platinum—p. *Courier*, Influenza—p. *Lait*, Serum lactis—p. *Lait d' Hoffmann*, Serum lactis Hoffmanni—p. *Mal*, see Epilepsy—p. *Sus-mazillo-labial*, Levator anguli oris.

PETITE-POSTE, Influenza—p. *Vérole*, Variola.

PETITS-LAITS, see Serum lactis.

PETRÆLEUM, (*petro*, and *ελαιον*, 'oil,') Petroleum.

PETRÆUM, (*πετραῖος*, 'living among rocks,') Solidago virgaurea.

PETRAPIUM, (*petro*, and *apium*, 'parsley,') Bubon Macedonicum.

PÉTRÉE, Petrous.

PÉTRÉOLE, Petroleum.

PÉTRÉUX, Petrous—p. *Os*, see Temporal bone.

PETRO, (*πετρα* and *πετρος*, 'rock,') in composition, rock; and the petrous or rocky portion of the temporal bone.

PETRO DEL PORCO, ('stone of the hog,') Bezoar of the Indian porcine.

PÉTROLE, Petroleum.

PETROLEUM, *Petrolæum*, *Petræ'leum*, *Fe'træ o'leum*, *Bitu'men Petroleum*, *Pissela'm 'indicum*, *Rock Oil*, *Petroleum Barbadensê*, *Barba'dos Tar*,

(F.) *Pétrole* ou *Pétrôle*, (*petro*, and *ελαιον*, 'oil.') A bituminous substance flowing spontaneously from the earth, in various parts of the world, and abounding on this continent over a large area, embracing a part of Canada West from Lake Ontario to Lake Huron, and portions of Western New York and Western Pennsylvania, the south-eastern half of Ohio, all Northwestern Virginia, and the eastern district of Kentucky. It is of a fetid odour, and bitter, acrid taste. It is semiliquid, tenacious, semi-transparent; of a reddish-brown colour; insoluble in water and alcohol; combines with fixed and essential oils and sulphur; and is partially soluble in ether. The petroleum found at Gabian, near Béziers, in France, has been called *Oleum Gabianum*, (F.) *Huile de Gabian*, *Petroleum rubrum*. It has been given as an excitant expectorant; and, mixed with tincture of assafœtida, in tapeworm. It is, also, used externally in chilblains, chronic rheumatism, affections of the joints and skin, and paralysis, and is a disinfectant and parasiticide.

New York Petroleum, called *Seneca Oil*, is used externally in similar cases.

PETROLEUM BARBADENSE, *Petroleum*.

PETROLEUM SULPHURATUM, *Bul'namum Sulphuris Barbadien'se*,—(Composed of *Petrol. Barbadi.* 3xvj, *Flor. Sulph.* 3iv) has been used as a pectoral; and as a detergent to ulcers. It has been accounted antispasmodic and sudorific. Dose, grt. x to xxx. Externally, it is used as a stimulant and discutient.

PETROÛCCIPITAL, *Petroûccipitalis*. Belonging to the petrous portion of the temporal bone and to the occipital bone.

PETROÛCCIPITAL SUTURE is formed by the junction of the petrous portion of the temporal with the occipital bone. It is a deep groove separating the bones, which have between them a thin layer of cartilage.

PETROSALPIN'GOPHARYNGE'US. A fleshy fasciculus, which extends from the sphenoid bone, from the petrous portion of the temporal bone, and from the Eustachian tube to the upper part of the pharynx. See *Constrictor*.

PETROSALPINGOSTAPHYLINUS, *Circumflexus*—p. *Salpingo-staphylinus*, *Levator palati*.

PETROSPHENOIDAL, *Petrosphenoidalis*. Belonging to the petrous portion of the temporal bone and to the sphenoid bone.

PETROSPHENOIDAL SUTURE—a name given to the small suture which is formed by the anterior edge of the petrous portion of the temporal bone and the posterior edge of the sphenoid.

PETROSAL, *Petrous*.

PETROSELINUM, (*πετροσελινον*, from *petro*, and *σελινον*, 'parsley,') *Apium petroselinum*—p. *Macedonicum*, *Bubon Macedonicum*.

PETROSTAPHYLIN, *Levator palati*.

PETROSUM, *OS*, see *Temporal*.

PETROUS, *Petro'sus*, *Petro'sal*, (F.) *Pétroux*, *Pétré*, *Pierroux*, (*πετρος*, 'a rock.') Resembling stone; having the hardness of stone. Relating or belonging to the petrous portion of the temporal bone.

PETROUS GAN'GLION, *Petro'sal Ganglion*, *Gan'glion of An'dersch*, is formed by the glossopharyngeal nerve, shortly after it emerges from the jugular fossa. From this ganglion, M. Jacobson of Copenhagen traced an anastomosing loop through the cavity of the tympanum, which left branches there, and was thence reflected downwards to join the Vidian nerve.—See *Otic Ganglion*. This has been called the *Nerve of Jacobson*, *Tympanic branch of the glossopharyngeal*; and the *inoculation*, *Jacobson's Anastomosis*.

PETROUS PORTION OF TEMPORAL BONE, see *Temporal Bone*—p. *Process*, see *Temporal Bone*.

PETROUS S'INUSES, *Petro'sal S'inuses*, are two *venosæ* sinuses of the dura mater, connected with

the petrous portion of the temporal bone, and distinguished on each side, into:—1. The *superior petrosal sinus*. It arises from the cavernous sinus; follows the upper edge of the petrous portion which affords it a gutter; passes into the great circumference of the tentorium cerebelli, and opens into the lateral sinus where the latter experiences a bend near the base of the pars petrosa. Its form is triangular. 2. The *inferior petrosal sinus* arises from the cavernous sinus at the same point as the last, with which it communicates at the moment of its origin. It descends between the inferior edge of the pars petrosa and the basilar process, and terminates in the lateral sinus, at the sinus or gulf of the internal jugular vein. It is triangular, and broader at its extremities than at the middle.

PETUM, *Nicotiana tabacum*.

PETUN, *Nicotiana tabacum*.

PEUCE, (*πενκη*) *Pinus sylvestris*.

PEUCEA BALSAMEA, see *Pinus abies*.

PEUCEDANUM, (*πενκη*, 'the pine tree,' its leaves resembling those of the pine;) *P. officinale* seu *absoluticum* seu *altissimum* seu *paniculatum* seu *ternatum*, *Selinum peucedanum*, *Mar'athrum sylv'es'tre*, *Mar'athrophyl'ium*, *Pinastel'ium*, *Fœniculum porc'inum*, *Bo'nus Gén'ius*, *Hoy's Fen'nel*, *Sul'phurwort*, (F.) *Peucedane*, *Queue de Pourcenn*, *Fenouil de Porc*. *Ord.* Umbellifère. *Sec. Syst.* Pentandria Digynia. The root has a strong, fetid smell—somewhat resembling that of sulphureous solutions—and an acrid, unctuous, bitterish taste. It has been recommended as antihysterical and nervine.

PEUCEDANUM ALSATICUM, *Peucedanum*—p. *Altissimum*, *Peucedanum*—p. *Austriacum*, see *P. palustre*—p. *Officinale*, *Peucedanum*—p. *Oreoselinum*, *Athamanta aureoselinum*.

PEUCEDANUM PALUS'TRE, *Thysse'lium* seu *Selinum palus'tre*, *Selinum sylv'es'tre* seu *thysse'lium* seu *lactescens*, *Thysse'lium*, *Olsnii'um*, *Marsh seli'num*, a European plant, has been recommended as an anti-epileptic. In Courland it has been a famous remedy in epilepsy. *Peucedanum Austriacum* is possessed of the same virtues.

PEUCEDANUM PANICULATUM, *Peucedanum*—p. *Pratense*, *P. silaus*.

PEUCEDANUM S'ILAUS, *P. pratense*, *Ses'eli pratense*, *Sil'um* seu *Ligusticum* seu *Onid'ium sil'aüs*, *Saxifraga vulg'aris* seu *Ang'lica*, *Hippomar'athrum*, *Fœniculum errat'icum*, *Eng'lish* or *Meadow Saxifrage*, (F.) *Séséli des prés*. The roots, leaves, and seeds have been recommended as aperients, diuretics, and carminatives. Rarely used.

PEUCEDANUM TERNATUM, *Peucedanum*.

PEUPLIER, *Populus*—p. *Baumier*, *Fagara octandra*—p. *de la Caroline*, *Fagara octandra*—p. *Frane*, *Populus*—p. *Noir*, *Peucedanum*.

PEXIS, (*πηξις*) *Concretion*.

PEYERI GLAND'ULÆ, *Peyeri Plex'us glandula'si*, *Gland'ulæ Peyeria'næ* seu *mucosæ conminatæ* seu *agminatæ* seu *mucip'aræ racemati'nges'tæ intestino'rum* seu *intestinal'es plexifor'mes* seu *spu'riæ*, *Corpus'cula glandula'rum similia*, *Gland'ulæ in ag'men congregatæ intestino'rum*, *Enterad'enes*, *Peyer's Glands*, *Ag'minated* or *ag'gregate glands*. Small glands situate beneath the villous coat of the intestines, (called after Peyer, who well described them.) They are clustered together, (F.) *Plaques de Peyer*, having a honeycomb appearance—*Plaques gaufrées*—and are extremely numerous. They occur most in the lower portion of the ileum, and resemble, in their morphology and functions, the lymphatic ganglia. They are, evidently, concerned in the great function of lymphosis.

PE'ZA, (*πεζα*.) The foot: according to some, the sole of the foot; and to others, the astragalus, the ankle.

PEZIZA seu PEZEZA AURICULA. *Tremella auricula* Ju'dæ, *Auricula* Ju'dæ, *Auricula-ria sambuci* seu *sambucina*, *Exidia auricula* Ju'dæ, *Merulius auricula*, *Fungus sambucinus*, *Agaricus Auriculae formæ*, *Jew's Ear*. Ord. Fungi. This fungus is said to be astringent; but has chiefly been used in decoction or infusion as a gargle in relaxed sore throat.

PFEFFERS or PFAFFERS, MINERAL WATERS OF, *Thermæ Piperi-næ*, *Th. Juba-rigæ*. These springs, which are in the Swiss Alps, are simply thermal: temperature 100° of Fahrenheit.

PHA'CÊ, (φακή, 'a grain of the lentil,') *Errum lens*. Also, chloasma; lens; the crystalline lens; and a flask for the application of heat, so called owing to its flattened lenslike shape.

PHACELLUS, (φακελλος,) Fasciculus. PHACELUS, (φακελος,) Fasciculus. PHACHYMENTIS, Phachymenitis. PHACI, (pl. of *Phacis*), Epelides. PHA'CICUS, *Phaco'tus*, *Phaco'des*, (from *phace*.) Lens-shaped, lenticular.

PHACITIS, *Phacodēit'is*, *Lenti'tis*, *Crystallit'is*. Also, chloasma; *Glenit'is*, (*phace*, and *itis*.) Inflammation of the crystalline lens. Inflammation of the capsule of the lens is termed *Cap-sulit'is*, *Cap-sulit'is*, *Crystallinocapsulit'is*, and *Phacocapsulit'is*.

PHACOCOPIIS, (*phace*, and *κοπίς*, 'a knife,') *Phacopis*.

PHACOCYSTE, (*phace*, and *κύστις*, 'a cyst or bladder,') see Crystalline.

PHACOCYSTECTOMIA, (*phacocyste*, and *εκτομία*, 'cutting out,') An operation for cataract, which consists in cutting out a part of the capsule of the lens.

PHACOCYSTITIS, (*phacocyste*, and *itis*), see Phacitis.

PHACO'DES, (φακώδης, *phace*, and *odes*), *Phacius*.

PHACHYMENTITIS, *Phachymenit'is*, *Phachymenit'is*, *Phachymenit'is*, *Inflammatio capsulae len'tis*; (*phace*, *ὑμν*, 'a membrane,' and *itis*.) Inflammation of the capsule of the crystalline.

PHA'CID, *Phacodēdes*, (from *phace*, and *είδος*, 'resemblance,') Resembling a lens. See Crystalline.

PHACOIDEITIS, (*phacoid*, and *itis*), *Phacitis*.

PHACOMALACIA, (*phace*, and *μαλακός*, 'soft,') Cataract, soft.

PHACOMETACHORE'SIS, (*phace*, and *μετακίρω*, 'to change place,') Dislocation of the crystalline.

PHAC'OPIS, *Phacop'is*, *Phacot'us*, (*phace*, and *κοπίς*, 'a knife,') *Scalpel'ium lenticula'rē*. A lenticular-shaped knife.

PHACOPALINGENESIS, (*phace*, *παλιν*, 'again,' and *γένεσις*, 'generation,') Regeneration of the crystalline.

PHACOPS'IS, (*phace*, and *ὥψις*, 'vision,') The state of one who has chloasma or freckles on the face.

PHACOPTIS'ANA, *Phacit'ium*, (*phace*, and *πίνω*, 'drink,') A drink of lentils. A poultice of lentils.

PHACOS, (φακος,) *Errum lens*.

PHACOSCOTOMA, (*phace*, and *σκοτώμα*, 'darkness,') Cataract.

PHACOSCLERO'MA, (*phace*, and *σκληρός*, 'hard,' and *ωμα*.) Induration of the crystalline.

PHACOTUS, (φακώτος, 'lens-shaped,') *Phacius*, *Phacopis*.

PHACUS, (φακος,) *Errum lens*.

PHACHYMENTIS, Phachymenitis.

PHÆNOMENOLOGIA, (*phænomenon*, and *λογος*, 'a discourse,') Semeiology.

PHÆNOMENON, (φαίνω, 'to manifest,') *Phænomenon*.

PHÆNOMENOSCOPIA, (*phænomenon*, and *σκοπεῖν*, 'to examine,') see Symptomatology.

PHAGÆNA, (φαγᾶνα, from *φαγῶ*, 'to eat,') *Boulimia*.

PHAGEDÆNA, (φαγεῖσθαι, 'a ravenous hunger,' from *φαγῶ*, 'to eat,') *Boulimia*, *Phagedenic ulcer*—p. *Gangrenosa*, *Hospital gangrene*—p. *Sloughing of the mouth*, *Cancer aquaticus*.

PHAGEDEN'IC, *Phagedēnicus*, *Phagedēnicus*, *Depress'ens*, *Ind'ent*, (F.) *Phagédénique*, *Rongeant*. That which rapidly eats away.

PHAGEDENIC CHAN'CRE. A destructive form of primary syphilitic sore. A virulent variety has been long known in the London Hospitals under the name, "*Sivan alley sore*," from the miserable locality inhabited by prostitutes, in which it formerly prevailed.

PHAGEDENIC UL' CER, *Nô'mê*, *Phagedēna*, *Ul'cus depress'ens* seu *obambulans seu fer'num*, (F.) *Ulcère phagédénique ou rongeant*, is one which rapidly eats and corrodes the neighbouring parts.

Phagedēnic, in *Materia Medica*, is a substance used for destroying fungous granulations in ulcers, &c.

PHAGEDENOUS, *Phagedenic*.

PHAKOHYMENTIS, *Phachymenitis*.

PHALACRA, (φαλακρα,) Calvities.

PHALACROMA, (φαλακρομα,) Calvities.

PHALACROSIS, (φαλακρωσις,) Calvities.

PHALACROTES, (φαλακρωτής,) Alopecia.

PHALÆNA PINGUINALIS, (φαλαίνα, 'a moth,') see *Ectozoa*.

PHALATA. A barbarous term, employed by Basil Valentine to designate a universal medicine or panacea.

PHALANGAGRA, (*phalanx*, and *αγρα*, 'a seizure,') *Phalangarthritis*.

PHAL'ANGAL or PHALAN'GAL, *Phalan'gian*, *Phalangia'nus*, (from *phalanx*.) Having relation to a phalanx.

PHALANGARTHRI'TIS, (*phalanx*, *αρθρον*, 'a joint,' and *itis*), *Phalan'gagra*. Gout in the fingers. Inflammation of the joints of the fingers.

PHALANGE, (φαλαγγ,) *Phalanx*.

PHALANGES DES DOIGTS, see *Phalanx*—p. of the Fingers, see *Phalanx*—p. *des Orteils*, see *Phalanx*—p. of the Toes, see *Phalanx*.

PHALANGETTES, see *Phalanx*.

PHALANGETTIEŒ. Professor Chaussier uses this term for anything relating to the third phalanx of the fingers or toes,—*Phalangien*, for what relates to the first,—and *Phalanginiën*, to the second.

PHALANGIAN, *Phalanga*.

PHALANGIEN, see *Phalangettien*.

PHALANGINES, see *Phalanx*.

PHALANGINIEN, see *Phalangettien*.

PHALANGIUM ESCULENTUM, (from *φαλαγγιον*, 'a venomous spider,' whose poison it was supposed to neutralize,) *Scilla esculenta*.

PHALANGO'SIS, *Hispid'itas*, (from *φαλαγξ*, 'a row of soldiers,' and *osis*.) An affection of the eyelids, in which the lashes are arranged in two or three rows. See *Trichiasis*. Also, dropping of the upper eyelid, from paralysis of the levator palpebræ superioris muscle.

PHAL'ANX, gen. *Phalan'gis*, (φαλαγξ, gen. *φαλαγγος*, 'a row of soldiers,') *Phalan'gæ*, *Phalan'goma*. A name given to the small bones which form the fingers and toes, because placed alongside each other like a phalanx. They are distinguished into the *Phalan'ges of the fingers*, *Os sa seu Ag'mina seu A'cies seu Artic'uli seu Scutic'ula seu Scut'ula seu Nodi seu Interno'dia seu Seg'mental'ia seu Con'dyli Digito'rum Ma'nūs*; and the *Phalanges of the Toes*, *Os sa seu Artic'uli seu Interno'dia Digito'rum Pe'dis*. I. *Phalanges of the fingers*, (F.) *Phalanges des doigts*. They are,

in each hand, fourteen.—three to each finger, and two to the thumb; and are situate one above the other. They are distinguished, numerically, reckoning from the base towards the tip of the finger.

1. The *first* or *metacarpal*, *Procon'dyli*, (F.) *Pre-mières phalanges* ou *Phalanges métacarpiennes*.
2. The *second* or *middle* *phalanges*, (F.) *Secondes phalanges* ou *Phalanges moyennes*, *Phalangines*, (Ch.) 3. The *third* *phalanges*, *Metaco'n'dyli*, *Phalanges unguis*, (F.) *Troisièmes phalanges* ou *Phalanges unguéales* ou *unguifères* ou *unguinales*, *Phalangettes* (Ch.), which terminate each finger, and support the nail. II. *Phalan'ges of the toes*, (F.) *Phalanges des Orteils*. They are the same in number as those of the fingers, but much smaller, and of a very different shape.

The phalanges of a finger or toe are united with each other; and the first with the metacarpal or metatarsal bone. They are developed from two points of ossification; one for the superior extremity, and the other for the body and inferior extremity.

PHALARIS CANARIENSIS, *P. ora'ta*, (φα'ρας,—from φαω, 'I shine,'—'bright, shining,' in application to the seed.) *Cultivated Canary Grass*, (F.) *Alpiste*. Ord. Gramineæ. See. Syst. Triandria Digynia. The seed of this grass, *Canary seed*, (F.) *Graines d'aspic*, afford a nourishing meal. The ancients recommended them for allaying pains in the kidney and bladder.

PHALARIS ZIZANOIDES, *Andropogon muricatus*.

PHALLAL'GIA, *Phallosdyn'ia*, (*phallo*, and αλγος, 'pain,') Pain in the male organ.

PHALLANAS'TROPHE, (*phallo*, and αναστροφή, 'a turning back,') Distortion of the male organ backwards.

PHALLANXYLOSIS, (*phallo*, and αγκυλωσις, a bending,') Chordee.

PHALLITIS, *Priap'itis*, (*phallo*, and itis.) Inflammation of the male organ.

PHALLO, (from φαλλος, 'the male organ,') in composition, the male organ.

PHALLOCAMPSIS, (*phallo*, and καμψις, 'bending,') Chordee.

PHALLOCARCINO'MA, gen. *Phallocarcino'matis*, (*phallo*, and καρκινωμα, 'cancer,') Carcinoma or cancer of the penis.

PHALLODYNTIA, (*phallo*, and οδυνη, 'pain,') **PHALLON'CUS**, (*phallo*, and οκος, 'swelling,') Tumefaction of the male organ.

PHALLOPSOPH'IA, (*phallo*, and ψοφος, 'a noise,') Discharge of wind from the male urethra.

PHALLORRHAG'IA, (*phallo*, and ρηγνυμι, 'I break forth,') Hemorrhage from the male urethra. Stomatosis.

PHALLORRHŒA, (*phallo*, and ρεω, 'to flow,') Gonorrhœa.

PHALLORRHOÏS'CHESIS, (*phallo*, ρη, 'discharge,' and ισχειν, 'to withhold,') Sudden arrest of a gonorrhœal discharge.

PHALLUS, (φαλλος,) Dildo, Penis—p. Esculentus. Morehella esculenta.

PHANEROG'AMOUS, *Phanerogam'ic*, *Phanerogam'icus*, (F.) *Phanérogame*, (φανερως, (φαω, 'I shine,') 'evident,' and γαμος, 'marriage,') An epithet used by botanists for plants whose sexual organs are apparent. It is, consequently, the opposite to *cryptogamous*.

PHANTAS'IA, (φαντασια, from φανταζω, (φανω, 'to bring to light,') 'to make appear,') Delirium, Imagination.

PHAN'TASM, *Phantas'ma*, *Phanto'ma*, *Phas'ma*. A phantom, false appearance, apparition. Hallucination. Derangement of the sense of sight or of the intellectual faculties, causing objects to be perceived, which are not before the eyes.

PHANTASMA, gen. *Phantasm'atis*, (φαντασμα, gen. φαντασματις,) **Phantasm**, **Pseudoblepsia**.

PHANTASMATICUS, (from *phantasma*,) **Phantasticus**.

PHANTASMATOSCOPIA, (*phantasma*, and σκοπειω, 'I view,') **Metamorphopsia**.

PHANTASMO'SCOPIA, **Metamorphopsia**.

PHANTAS'TICUS, (φανταστικός,) **Imbriant**. Creating phantoms—*phantasmaticus*.

PHANTOM, (φανω, 'to cause to appear,') **Fantom**—p. Tumour, see **Fantom**.

PHANTOMA, gen. *Phanto'matis*, **Phantasm**—p. Obstetricium. **Fantom**.

PHARMACEIA, (φαρμακεια,) **Pharmacy**, **Poisoning**.

PHARMACEUM, (φαρμακειον,) **Apotheca**.

PHARMACEU'MA, gen. *Pharmaceu'matis*, (φαρμακευμα,) **Medicament**.

PHARMACEUS, (φαρμακευς,) **Apothecary**.

PHARMACEUTA, (φαρμακευτης,) **Apothecary**.

PHARMACEUTIC, *Pharmaceu'ticus*, *Pharmaceu'tical*, *Pharmaceutica'lis*, (F.) *Pharmaceutique*, (from *pharmakon*,) An epithet for everything belonging to pharmacy. Thus, we say, *Pharmaceutical preparations*, &c.

PHARMACEUTICE, (φαρμακευτικη [τεχνη]) **Pharmacy**.

PHARMACEUTIST, (φαρμακευτης,) **Apothecary**.

PHARMACEUTRIA, (φαρμακευτρια,) A woman who prepares or deals in drugs or poisons.

PHARMACIA, (φαρμακια,) **Pharmacy**, **Poisoning**.

PHARMACIEN, **Apothecary**.

PHARMACO, (from φαρμακον, 'a medicine,' 'poison,') in composition, generally medicine; at times, poison.)

PHARMACOCATAGRAPHOLOG'IA, *Catagrapholog'ia*, *Ars formulæ concinnan'di*, (*pharmaco*, κατα, 'beneath,' γραφη, 'a writing,' and λογος, 'a description,') The art of writing prescriptions.

PHARMACOCHYMIA, (*pharmaco*, and chymia,) **Chymistry**, **pharmaceutical**.

PHARMACODYNAM'ICS, (*pharmaco*, and δυναμις, 'power,') *Pharmacodynam'icæ*, *Dynam'ical Mate'ria Med'ica*, *Pharmacodynam'ica doctri'na*, *Pharmacodynamilog'ia*. A division of pharmacology, which considers the effects and uses of medicines.

PHARMACODYNAMIOLOGIA, (same etymon and λογος, 'a discourse,') **Pharmacodynamic**.

PHARMACOGNO'SIA, *Acogno'sia*, *Acogno'eia*, (*pharmaco*, and γνωσις, 'knowledge,') A division of pharmacology, which treats of simples, or unprepared medicines.

PHARMACOLOGIA, (*pharmaco*, and λογος, 'a discourse,') **Materia medica**.

PHARMACOMA'NIA, (*pharmaco*, and mania, 'mania,') A mania for prescribing or for taking or giving medicine, or for drugging.

PHARMACON, (φαρμακον,) **Medicament**, **Poison**.

PHARMACOPŒ'IA, *Pharmacopœ'a*, *Co'dex medicamentarius*, (F.) *Pharmacopée*, (*pharmaco*, and ποιω, 'I make,') A work, containing a collection of formulæ for the preparation, &c., of medicines: generally published by authority. Formerly, a chemical laboratory, *Laborato'rium chym'icum* seu *pharmaceut'icum*.

PHARMACOPOËSE. The preparation of medicines.

PHARMACOPŒUS, **Apothecary**, **Druggist**.

PHARMACOPOLA, (φαρμακωπλη, from *pharmaco*, and πωλω, 'to sell,') **Apothecary**, **Druggist**.

PHARMACOPOLE, **Apothecary**, **Druggist**. **PHARMACOPOLIUM**, *Apothe'ca*, *Offici'na*, *Tuber'na Med'ica*. The shop of the druggist or apothecary, (F.) *Apothicairerie*.

PHARMACOPŒS'IA, (*pharmaco*, and ποσις, 'a potion,') A liquid medicine, especially one that is cathartic.

PHARMACOSA MEDICAMENTA, from

pharmacœon. Medicines that contain poisonous substances.

PHARMACOTHE'CA. (φαρμακοθήκη, *pharmacoe*, and *thea*, 'a sheath.') A case for medicines. A medicine chest.

PHARMACOTHERAPI'A. *Pharmac'is*, (*pharmacœon*, and *θεραπεία*, 'treatment.') Cure of diseases by medicine.

PHARMAC'TER. (φαρμακτερός.) Apothecary.

PHARMACUM. (φαρμακόν.) Medicament, Poison — *p.* Ægyptiacum, Ægyptiacum — *p.* Amatorium, Philter.

PHAR'MACUM AD AU'RES. *Ægyptium Medicamentum ad Aures.* A name given to a medicine composed of bitter almonds, white pepper, saffron, myrrh, opium, frankincense, castor, verjuice, sulphate of iron, vinegar, pomegranate bark, unguentum nardium, &c. It was believed to be useful for cleansing fetid ulcers in the ears.

PHARMACURGICUS. (*pharmacœon*, and *εργειν*, 'to work.') Apothecary.

PHARMACURGUS. Apothecary.

PHAR'MACY. *Pharmac'ia*, *Pharmacœuticē*, *Pharmac'is*, (from *pharmacœon*.) The art which teaches the knowledge, choice, preservation, preparation, and combination of medicines. It was formerly divided into *Galenical* and *Chymical*. The former, called after Galen, embraced the operations to which medicines were subjected without chymistry;—the latter, *Chymical Pharmacy* or *Pharmacœutic Chymistry*, *Pharmacœchym'ia*, comprised the preparation of medicines founded on the chymical action of their principles.

PHARMAXIS. (φαρμαξίς.) Pharmacotherapy, Pharmacy.

PHARNA'CEUM LINEA'RĒ. (called after Pharnaces, King of Pontus.) A South African plant, *Nat. Ord.* Paronychiaceæ; employed at the Cape in pulmonary affections. It is generally given in infusion.

PHARUS. (φάρος.) Pharynx.

PHARYGATHRON. (φάρυγαθρον.) Pharynx.

PHARYGETHRUM. (φάρυγεθρον.) Pharynx.

PHARYNGÆ'UM (SAL.). (from *pharynx*.) A compound of cream of tartar, nitrate of potassa, burnt alum, and distilled vinegar. It was formerly employed, united with honey and plantain-water, to make gargles, which were advised in quinsy.

PHARYNGALGIA. (*pharyngo*, and *αλγος*, 'pain,') Pharyngodynia.

PHARYNGĒ. Pharyngeal.

PHARYNGE'AL or **PHARYNGE'AL.** *Pharyngæus*, *Fau'cial*, (F.) *Pharyngē*, *Pharyngien*, (from *pharynx*.) That which relates to the pharynx or fauces. A name given to the muscles, vessels, and nerves of the pharynx.

PHARYNGEAL ARTERIES. These are two in number. The one—*superior* or *pterygo-palatine*, *Ramus Pharyngæus superior* of Haller—arises from the internal maxillary at the base of the zygomatic fossa, passes through the pterygo-palatine canal, and proceeds to be distributed to the pharynx and the corresponding part of the Eustachian tube. The other—the *inferior*—*Pharyngæa ascendens* of Haller—arises from the external carotid, on a level with the facial. It ascends along the lateral and posterior part of the pharynx, and divides into two branches; the one—the *pharyngeal*, properly so called—is distributed to the pharynx; the other—the *meningeal*—gives its branches particularly to the dura mater.

PHARYNGEAL NERVE. A branch sent by the pneumogastric to the pharynx. It descends behind the internal carotid; communicates with the glosso-pharyngeal; and divides, opposite the middle constrictor of the pharynx, into a great number of filaments, which anastomose, in a multitude of different directions, with filaments of the

glossopharyngeal nerve, of the superior laryngeal branch, and of the first cervical ganglion. These different nervous filaments united constitute the *pharyngeal plexus*, the irregular meshes of which send numerous ramifications to the pharynx.

PHARYNGEAL VOICE. see *Voice*.

PHARYNGEMPHRAX'IS. (*pharyngo*, and *εμφραξις*, 'obstruction.') Obstruction of the pharynx.

PHARYNGETHRON. (φάρυγεθρον.) Pharynx.

PHARYNGEURYSMA. (*pharyngo*, and *ευρυσμα*, 'dilatation,') Pharyngocoele,

PHARYNGIUS. (from *pharynx*.) Pharyngeal

PHARYNGIEN. Pharyngeal.

PHARYNGISMUS. Pharyngosasmus.

PHARYNGITE. Cynanche pharyngea — *p.* *Tonsillaire*, Cynanche tonsillaris.

PHARYNGITIS. (*pharyngo*, and *itis*.) Cynanche pharyngea.

PHARYNGITIS APOSTEMATOSA. Abscess of the pharynx.

PHARYNGITIS, DIPHTHERITIC. *Angina pseudomembranacea seu membrana'cea seu pellicularia seu plastica seu diphtheritica seu maligna seu pseudomembranosa seu pestidentia'lis seu fau'cium exudativa Pseudomembranous Inflammation or Diphtheritis or Diphtheria of the Throat, Hogskin angina*, (F.) *Angine couenneuse ou gutturale couenneuse ou pseudo-membraneuse ou plastique ou diphtheritique, Herpes guttural.* Diphtheritic inflammation of the pharynx. See Diphtheritis.

When *diphtheria* is spoken of, it is usually this affection that is meant; although the local phenomena are rather expressions of the peculiar adynamic condition, termed *diphtheritis*; which may destroy without much or any evidence of the pharyngeal affection. It is this adynamic condition which demands the main attention;—the pharyngeal exudation requiring but little. During convalescence, peculiar paralytic phenomena supervene at times. See Paralysis, diphtheritic. Diphtheritic pharyngitis is not new. It is manifestly the malignant sore-throat, which is described as having been so fatal—in the time of Horace Walpole, for example. See Cynanche maligna.

PHARYNGITIS, FOLLICULAR. *Follicular Inflammation of the Pharynx, Angina folliculosa* and *A. glandulosa* of the *Pharynx*, *Ac'nē of the Throat.* Inflammation and enlargement of the follicles of the pharynx, occasionally extending to the larynx. Sometimes termed *Clergyman's Sorethroat*, *Rhe-torari'chē*, *Tubercles of the Larynx* and *Fauces*, and *Tubercular Sorethroat*.

PHARYNGITIS, GANGRENOUS. Cynanche maligna.

PHARYNGO. (φάρυγξ, gen. φάρυγγος, in composition, the pharynx.)

PHARYNGOCACE. (*pharyngo*, and *κακος*, 'evil,') Cynanche maligna.

PHARYNGOCE'LE. *Pharyngeurysma*, *Diverticulum seu Hernia seu Prolap'sus Œsophagi seu pharyng'is*, *Œsophagus secenturiatus*, (from *pharyngo*, and *κελη*, 'a tumour,') An enlargement at the upper part of the gullet, in which the food sometimes rests in considerable quantity.

PHARYNGODYN'IA. *Pharyngal'gia*, (*pharyngo*, and *δύνη*, 'pain,') Pain in the pharynx.

PHARYNGOGLOSSAL. *Glossopharyngeal*, *Pharyngoglossia'lis*. That which belongs to the pharynx and tongue.

PHARYNGOGLOSSAL NERVE. *Glossopharyngeal Nerve*, *Lingual'is lateralis*, (F.) *Nerv. Glossopharyngien*, *Pharyngoglossien*—a portion of the eighth pair—arises from the superior and lateral parts of the spinal marrow, between the facial and pneumogastric nerves, by several filaments,

plant is stimulant and narcotic. It has been used in intermittents and dyspepsia. It has, also, been given in phthisis pulmonalis, and in cases of scrofulous atonic ulcers.

PHENE, Benzole.

PHENGOPHOBIA, (φειγος, 'light,' and φοβος, 'dread,') Photophobia.

PHENIG'MUS, (φαινιγμος,) *Phenig'mus*, (from φαινι, 'purple or red,') *Red Jaw'dice*. A cutaneous affection, consisting of redness diffused over the skin, without fever. Sauvages makes it a genus of his order *Ictericæ*. Also, the act of irritating a part by a stimulating application.

PHENOL, Acid, phenic.

PHENOMÈNE, Phenomenon.

PHENOM'ENON, (φαινομενον,) *Phænomen'on*, (F.) *Phénomène*, (φαινομαι, 'I appear,') An extraordinary and unexpected event. In *Medicine*, it means any change, appreciable by our senses, which supervenes in an organ or function:—as the *phenomena* of the circulation, respiration, &c. It is used in pathology, synonymously with symptom.

PHENYLE, HYDRURET OF, Benzole.

PHEUGYDRON, (φευγω, 'to flee,' and υδωρ, 'water,') Hydrophobia.

PHIALA, A *Phial* or *Vi'al*, (F.) *Fiole*, (φιαλη, φιαλη, 'a broad drinking cup, a saucer,') A small bottle, in which apothecaries are in the habit of sending out their mixtures, draughts, &c.: hence called *Med'icine Vials*, (F.) *Fioles à Médecine*. They are usually sold in grosses, and assorted into vials of oz. 8; oz. 4; oz. 3; oz. 2; oz. 1½; oz. 1; and oz. ½.

PHILAN'DYNAMICA, (*philo*, *a*, and δυναμις, 'power,') Debilitants.

PHILANTHROPUS, (*philo*, and ανθρωπος, 'man,') Galium aparine.

PHILIA'TER, *Philia'tros*, (φιλιατρος, *philo*, and ιατρικη, 'medicine,') An amateur of medicine. One who studies medicine; *Studio'sus medic'i'næ*.

PHILO, (φιλω, 'I love,') in composition, to love.

PHILOBIO'SIS, (F.) *Philobiosie*, (*philo*, and βιος, 'life,') The love of life.

PHILÆ'NIA, (φιλαινια, from *philo*, and οινος, 'wine,') Love of wine. Addiction to wine.

PHILOGY'NIA, (φιλογυνια, from *philo*, and γυνη, 'a woman,') Muliebrositas.

PHILONIUM LONDINENSE, (after its proposer, Philo,) Confectio opii.

PHILOPATRIDALGIA, (*philo*, πατρις, 'one's country,' and αλγος, 'pain,') Nostalgia.

PHILOPATRIDOMANIA, (*philo*, πατρις, 'one's country,' and mania,) Nostalgia.

PHILOSOPHER'S STONE, see Alchemy.

PHILOSOPHIA CORPORIS VIVI, Physiology—p. Hermetica, Alchemy—p. per Ignem, Chymistry.

PHIL'TER, (φιλτρον, from φιλω, 'to love,') *Philtrum*, *Phar'macum* seu *Poc'ulum amato'rium*, *Amato'rium venefic'ium*, (F.) *Philtre*. A medicine or drink considered capable of causing love.

PHILTRUM, Lacuna labii superioris.

PHILYRA, Tilia.

PHILY'REA LATIFO'LIA, (after *Phily'ra*, 'mother of Chiron,') A shrub of the south of France and Spain, *Ord.* Oleaceæ, whose leaves have been used as an astringent in ulcers of the mouth and throat. The flowers, steeped in vinegar, have been applied to the head in headache. The decoction of the leaves is said to be diuretic.

PHIMOS, (φίμος,) Capistrum.

PHIMO'SICUS. Relating to Phimos. Sauvages calls *Ischu'ria phimo'sica* seu *urethra'lis à phimo'si*, the retention of urine which depends upon phimos.

PHIMO'SIS, improperly *Phymo'sis*, *Capistra'tio*, *Epago'yium*, *Ligatu'ra glan'dis*, *Strictu'ra præ-*

put'ii, *Reductio præput'ii impedita*, (φίμω, 'I bind tight,') A disease which consists in a preternatural narrowness of the opening of the prepuce, so that it cannot be carried behind the corona glandis. This affection is most commonly congenital. Sometimes it is accidental, and dependent upon tumefaction of the glans, or of the prepuce, as occurs in syphilis. To remedy this state, the prepuce may be divided by means of a bistoury and director; or circumcision may be practised.

PHIMOSIS ADNATA seu CINGENTA seu PUERILIS, Congenital phimos—p. Circumligata, Paraphimos—p. Congenita, P. adnata.

PHIMOSIS EDEMATO'DES, *Hydrophimo'sis*. Phimos with cedematous swelling of the prepuce.

PHIMOSIS PUERILIS, Ph. adnata.

PHLASMA, gen. *Phlasm'atis*, (φλασμα, from φλω, 'to bruise,') Contusion.

PHLEBARTERIODIALYSIS, (*phlebo*, αρτηρια, 'artery,' and διαλυσις, 'a loosing or separating,') see Aneurism.

PHLEBECTA'SIA, *Phlebec'tasis*, (*phlebo*, and εκτασις, 'dilatation,') Dilatation of a vein, or of a portion of a vein.

PHLEBECTIARCTIE, (*phlebo*, and αρκτο, 'I contract,') Contraction of a vein.

PHLEBEMPHRAX'IS, (*phlebo*, and εμφραξις, 'obstruction,') Obstruction of a vein.

PHLEBES, pl. of *phlebs*, (φλεβες,) see Artery.

PHLEBEURYSMA, (gen. *Phlebeurysm'atis*, (*phlebo*, and ευρυσμα, 'dilatation,') Varix.

PHLEB'ICUS, (φλεβικος,) Relating or appertaining to a vein.

PHLEB'ION, (φλεβιον,) (dim. of *phlebs*,) A vein. A small vein,—*Venula*.

PHLEBIS'MUS, (from *phlebs*.) A term employed by Dr. Marshall Hall for the turgescence of veins, produced by impeded return of the blood. See Trachelismus.

PHLEBITIS, *Inflammatio venarum*, (F.) *Phlébite*, *Inflammation des veines*, (*phlebo*, and itis.) Inflammation of the inner membrane of a vein, *Endophlebitis*, which sometimes follows blood-letting, and extends from the small wound made in that operation to the neighbouring parts of the venous system. The symptoms are, first, inflammation in the punctured part; and, afterwards, a knotty, tense, painful cord, following the direction of the vessel, and accompanied with more or less fever, according to the extent of the inflammation, the nervous symptoms, &c. Phlebitis, also, sometimes supervenes on the ligature or excision of the varices; after amputation, gun-shot wounds, delivery (*U'terine Phlebitis*, *Lochometrophlebitis*, *Metrophlebitis puerpera'lis*, *Phlebitis uteri'na*, *Metri'tis venosa*), the ligature of the umbilical cord, &c. It may terminate by resolution, suppuration, ulceration, or gangrene.

PHLEBITIS, CRURAL, *Phlegmatia dolens*—p. Uterina, see Peritonitis, and Phlebitis.

PHLEBO, (φλεβ, gen. φλεβος, 'a vein,') In composition, a vein.

PHLEBOG'RAPHY, *Phlebogra'phia*, (*phlebo*, and γραφη, 'a description,') A description of the veins.

PHLEB'OLITE, *Phlebol'ithus*, *Vein stone*, (F.) *Phlébolithe*, (*phlebo*, and λιθος, 'a stone,') A loose concretion, varying in size from that of a currant to that of a pea, occasionally found in the veins. It would seem to be formed in the blood.

PHLEBOLITHUS, Phlebolite.

PHLEBO'OGY, *Phlebolog'ia*, (*phlebo*, and λογος, 'a discourse,') The part of anatomy which treats of the veins.

PHLEBOMETRITIS, (*phlebo*, and metri'tis,) Metrophlebitis.

PHLEBOPA'LIA, (*phlebo*, and παλλω, 'I palpitate,') With the an-ients, *pulsation*, which they

conceived to be produced by the veins. See *Pulsus venarum*.

PHLEBOPHTHALMOTOM'IA, *Ophthalmophlebotom'ia*, (*phlebo*, *ὀφθαλμος*, 'the eye,' and *τομή*, 'I cut.') Blood-letting in the eye.

PHLEBOPLEROSIS ECPHRACTICA, (*phlebo*, and *πληρωσις*, 'fulness,') Inflammation.

PHLEBORRHAG'IA, (*phlebo*, and *ῥήγναι*, from *phlebo*, and *ῥήγναι*, 'I break out.') Rupture of the veins; and, in a more extensive sense, venous hemorrhage.

PHLEBORRHEX'IS, (*phlebo*, and *ῥήξις*, 'rupture,') Rupture of a vein. Also, hemorrhage from a vein, *Phleborrhag'ia*.

PHLEBOSTENO'SIS, (*phlebo*, and *στενωσις*, 'contraction,') Constriction or contraction of a vein.

PHLEBOTHROMBOSIS, (*phlebo*, and *thrombosis*,) Coagulation of blood in a vein.

PHLEBOTOMOMANIA, *Phlebotomom'nia*, (*phlebotomy*, and *mania*,) An extravagant estimate of the value of blood-letting as a curative agent.

PHLEBOTOMIST, (*phlebo*, and *τομή*, 'I cut,') A bleeder. One who practises particularly the art of phlebotomy. See *Blood-letting*.

PHLEBOTOMUM, (*φλεβοτόμιον*,) Fleam.

PHLEBOTOMY, (*φλεβοτομία*,) See *Blood-letting*.

PHLEBS, (*φλεψ*, gen. *φλεβος*,) Vein.

PHLEDONIA, (*φλεδονεία*, from *φλεω*, 'to talk idly,') Delirium.

PHLEGM, (*φlegma*,) *Phleg'ma*, *Pituita*, *Pituite*, *Mucus*, *Mother*, (F.) *Phlegme*, *Pituite*. One of the four natural humours of the ancients, which, according to them, was cold and moist, as atrablis was cold and dry. It predominated, especially, in winter.

Pituita was afterwards applied to every aqueous or excrementitious humour, such as the saliva, nasal and intestinal mucus, serum, &c. The terms *phlegm* and *pituita* are no longer used in physiology,—the different humours having received particular names; but the vulgar still use *phlegm* to designate a stringy mucus, expectorated, or rejected by vomiting. The ancient chymists gave the name 'phlegm' to aqueous, insipid, and inodorous products obtained by subjecting moist vegetable matters to the action of heat.

PHLEGMA, gen. *Phleg'matis*, (*φlegma*, gen. *φlegματος*,) *Phlegm*—p. *Narium Crassum*, Nasal mucus—p. *Vitrioli*, Sulphuric acid (very dilute.)

PHLEGMAGOGUE, *Phlegmagog'us*, (*phlegma*, and *αγω*, 'I expel,') A medicine formerly believed capable of 'cutting' or removing phlegm.

PHLEGMAPYRA, (*phlegma*, and *πυρ*, 'fire, fever,') Fever, adeno-meningeal.

PHLEGMASIA, (*φlegmasia*, from *φλεγω*, 'to inflame,') Inflammation, *Phlegmon*—p. *Adenosa Adenitis*—p. *Alba dolens* puerperarum, Ph. dolens—p. *Cellularis*, Induration of the cellular tissue—p. *Glandulosa*, *Adenitis*—p. *Myoica*, *Myositis*—p. of the Peritoneum, *Peritonitis*.

PHLEGMASIQUE, Inflammatory.

PHLEGMA'TIA, (from *phlegma*,) (F.) *Phlegmatie*. *Edema*, *Anasarca*, *Hydrops*.

PHLEGMATIA DO'LENS, *P. lac'tea* seu *alba dolens* seu *Leucophlegmatia dolens* seu *Spargano'sia* seu *Scelut'gia* seu *Ede'ma puerperarum*, *Scelon'cus anasar'ca sero'sa*, *Ecephy'ma edemat'icum*, *Echymo'ma lymphat'icum*, *Is'chias a spargano'sia*, *Ede'ma lac'teum*, *Galactode'ma*, *Inflar'ctus lac'tei extremitatum*, *Metast'asis lactis*, *Hysteral'gia galactica*, *Hydrophlegma'sia testis cellul'o'si*, *Cru'al Phlebitis*, *White leg*, *Swelled leg*, *Milk leg*, *White swelling* of lying in women, (F.) *Phlegmatie puerpérale*, *Edème des nouvelles accouchées*, *Edème douloureux des femmes en couche*, *Dépôt lactéux sur la cuisse*, *Edème actif des nouvelles accouchées*, *Enflure ou Engorgement des jambes et des*

cuisse de la femme accouchée, *E. lactéux des membranes abdominaux*. It occurs, for the most part, in the second or third week after delivery:—is limited to the lower extremity, and chiefly to one side; exhibiting to the touch a feeling of numerous irregular prominences under the skin. It is hot, white, and unyielding; and is accompanied, sooner or later, with febrile excitement. After a few days, the heat, hardness, and sensibility diminish, and the limb remains oedematous for a longer or shorter period. The disease frequently, if not generally, consists in the obstruction of the iliac or other veins. Owing to the pressure of the gravid uterus—the flow of blood being obstructed—the liquid part of it is thrown out into the cellular membrane of the limb. Sometimes the vein is found completely obliterated.

The treatment consists in the use of antiphlogistics:—the application of leeches or cupping-glasses near the groin, fomentations to the limb, &c.; and, when the active state has subsided, the use of a roller, applied gently at first, but, afterwards, gradually tightened.

PHLEGMATIA LACTEA, P. dolens.

PHLEGMATIÆ, see *Empresma*—p. *Membrano'se et parenchymatose*, *Empresma*.

PHLEGMAT'IC, (*φlegmatikos*,) *Phlegmat'icus*, *Phlegmato'des*, (F.) *Phlegmatique*. Abounding in phlegm,—as the 'phlegmatic or lymphatic temperament.'

PHLEGMATIE, *Phlegmatia*—p. *Puerpérale*, *Phlegmatia dolens*.

PHLEGMATO'DES, (*φlegmatōdes*, from *phlegma*, and *odes*,) Full of inflammation, or phlegm.

PHLEGMATOCHEMY, (*phlegma*, and *χημεία*, 'chemistry,') see *Chymistry*.

PHLEGMATOPYRA, (*phlegma*, and *πυρ*, 'fire, fever,') Fever, adenomeningeal.

PHLEGMATORRHAG'IA, *Phlegmatorrhœ'a*, *Catarrh*, (*phlegma*, and *ῥαγναι*, 'a breaking forth,') Stahl and his partisans gave this name to a copious secretion, by the nostrils, of a limpid and thin mucus, without inflammation. It is the *Coryza Phlegmatorrhag'ia* of Sauvages. See *Coryza*.

PHLEGMATORRHEA, (*phlegma*, and *ῥεω*, 'to flow,') Catarrh.

PHLEGMHYMENITIS, *Phlegmymenitis*.

PHLEG'MON, (*φlegmō*, 'I burn,') *Phlego'sis* *phlegmonē*, *Phleg'monē*, *Phlegma'sia*, *Iscen'dium*, *Inflammati*, (Prov.) *Beal*. Inflammation of the areolar texture accompanied with redness, circumscribed swelling, increased heat and pain; which is, at first, tensile and lancinating; afterwards, pulsatory and heavy. It is apt to terminate in suppuration. It requires the antiphlogistic treatment, until suppuration seems inevitable, when the suppuration must be encouraged as the next favourable termination to resolution.

PHLEGMON PÉRINÉPHRÉTIQUE, *Periphrenitis*—p. *Periuterine*, see *Periuterine*.

PHLEGMONE, (*φlegmonē*,) Inflammation, *Phlegmon*—p. *Mammæ*, *Mastodynia apostematosa*—p. *Mastodynia*, *Mastodynia apostematosa*—p. *Musculorum*, *Myositis*—p. *Parotidæ*, *Parotitis*—p. *Parulis*, *Parulis*—p. *Peritonæi*, *Peritonitis*—p. *Ventriculi*, *Gastritis*.

PHLEGMONEUX, *Phlegmonous*.

PHLEGMONODES, (*φlegmonōdes*,) *Phlegmonous*.

PHLEGMONOID, (*phlegmon*, and *ειδός*, 'resemblance,') Resembling phlegmon.

PHLEG'MONOUS, *Phlegmono'des*, *Phlegmonoid*, (F.) *Phlegmonoux*, (from *phlegmon*,) That which relates or belongs to phlegmon.

PHLEGMONOUS INFLAMMATION is the active inflammation of the areolar membrane; in contradistinction to erysipelatous inflammation, or that of the skin.

PHLEGMOPYRA. (*phlegma*, and *πῦρ*, 'fire, fever,') Fever, adenomeningeal.

PHLEGMORRHAGIA. (*phlegma*, and *ῥαγή*, 'a breaking forth,') Blennorrhœa.

PHLEGMORRHAGIE PULMONAIRE. Bronchorrhœa.

PHLEGMORRHŒA. (*phlegma*, and *ῥέω*, 'to flow,') Blennorrhœa.

PHLEGMYMEINITIS. *Phlegmymentitis*. (*phlegma*, *ἔμνη*, 'a membrane,' and *itis*.) Inflammation of a mucous membrane.

PHLEGMYMEINITIS EXTERNA, see Enteritis.

PHLEPS. (*φλεψ*, gen. *φλεβος*.) Vein.

PHLOGŒMIA. (*phlogia*, and *αἷμα*, 'blood,') see Hæmicitis.

PHLOGIA. (*φλογία*.) Inflammation.

PHLOGICUS. (from *phlogia*.) Inflammatory.

PHLOGISTIC. (from *phlogia*.) Inflammatory.

PHLOGISTICI. Eupresma.

PHLOGISTON. (*φλογίζω*, 'to burn,') Hydrogen.

PHLOGMUS. Verbascum thapsus.

PHLOGODES. (from *phlogia*, and *odes*.) *Inflamed, Inflammatum, Inflammativus*. Inflammatory. An epithet employed particularly to express the red colour of the face.

PHLOGOGENE. (*φλεξ*, gen. *φλογος*, 'flame,' and *γενω*, 'to beget,') Hydrogen.

PHLOGOPYRA. (*φλεξ*, gen. *φλογος*, 'flame,' and *πῦρ*, 'fever,') Synocha.

PHLOGOSÉ, Inflamed.

PHLOGOSÉD, Inflamed.

PHLOGOSIS. (*φλογωσις*.) *Incedium, Inflammation*, (from *phlogia*.) Some writers use this word exclusively for external inflammation—phlegmon; others, for that which is superficial or erysipelatos.

PHLOGOTICA. *Inflammations*. The second order in the class *Hæmatia* of Good.

PHLOIOS. (*φλοιος*.) Cortex.

PHLOMUS. Verbascum thapsus.

PHLOOS. (*φλοιος*.) Cortex.

PHLORIDINE. *Phloridina*. *Phlor'idin*. (from *phloos*, and *ρίζα*, 'root,') The bitter or active principle of the root and trunk of the apple, pear, cherry, and plum tree. It is deposited from the decoction on cooling; crystallizes in silky spicula of a dead white colour when in masses; or in long slender prisms or tables, when crystallized with care. 1000 parts of water, at a temperature of from 32° to 71°, only dissolve about one part; but at from 70° to 212°, it is dissolved in all proportions. It is very soluble in pure alcohol, but only slightly so in ether; e.g. 1.429. It has been administered successfully in intermittents,—10 to 20 grains being generally required to arrest the paroxysm. It may be given either in the form of pill or powder.

PHLOUS. (*φλοιος*.) Cortex.

PHLOX. Verbascum thapsus.

PHLYCTÆNÆ. (*φλυκταινά*, from *φλεω*, 'to bubble up,') *Phlyctænia*, *Phlyctis*, *Phlyctænia*, *Amphulla*, *Holophlyctis*, *Phlyctis*, *Phlyctis*, (F.) *Phlyctène*. A tumour formed by the accumulation of a serous fluid under the epidermis.

PHLYCTENOIDES. (*φλυκταινοειδής*.) *Phlyctenoid*.

PHLYCTENOPHTHALMIA, see *Phlyctenulæ*.

PHLYCTENOSIS. (*φλυκταινωσις*.) The formation of *phlyctenæ*.

PHLYCTENULA. (dim. of *phlyctæna*.) see *Hydatid* and *Phlyctenula*.

PHLYCTÈNE. *Phlyctæna*.

PHLYCTENOID. (*Phlyctenoides*, (*phlyctæna*, and *αἶδος*, 'resemblance,') Resembling *phlyctæna*. An epithet for a variety of herpes.

PHLYCTENULAR. (from *phlyctenula*.) Of or belonging to *phlyctenula*,—as *phlyctenular ophthalmia*; inflammation of the eye, accompanied

with *phlyctenulæ* on the cornea. *phlyctenophthalmia*.

PHLYCTIDES. (pl. of *phlyctis*.) *Phlyctenæ*.

PHLYCTIDIUM. (dim. of *phlyctis*.) A circular or annular spot of inflammation of the corium, encircled by a red ring or zone, within which suppuration takes place. The genuine small-pox pustule, and that produced by the application of tartar emetic ointment, are examples of it.

PHLYCTIS. gen. *Phlyctidis*, (*φλυκτις*.) *Phlyctæna*.

PHLYSIS. (*φλυσ*, from *φλεω*, 'to buldle up,') A cutaneous eruption. An ulcerative subcutaneous tumour: flat, tensile, glabrous: diffused, hot, throbbing; at length fluctuating, with an acrid ichor.

PHLYSIS. *Phlyctæna*—p. *Ecthyma*, *Ecthyma*—p. *Impetigo*, *Impetigo*—p. *Porrigio*, *Porrigio*—p. *Scabies*, *Psora*.

PHLYZACIA. *Ecthyma*.

PHLYZACION. (*φλυσάκιον*, from *φλεω*, 'to buldle up,') *Phlyctæna*.

PHOBODIPSON. (*φοβόδιψον*, from *φοβος*, 'dread,' and *διψα*, 'thirst,') *Hydrophobia*.

PHOCOMELUS. (*φωκ*, 'a seal,' and *μελος*, 'a limb,') A monster, in which the limbs are shortened; the hands and feet appearing to exist alone, and to be inserted immediately on the trunk, as in the seal and the herbivorous cetacea.

PHŒNICINE. (*φαική*, gen. *φαικής*, 'red,') *Hæmatin*.

PHŒNICISMUS. (*φαική*, gen. *φαικής*, 'red,') *Rubella*.

PHŒNIGMUS. (*φαιγμός*.) *Phenigmus*—p. *Pecthialis*, *Purpura simplex*.

PHENIX DACTYLIFERA. (*φαινίξ*.) *Date tree*—p. *Excelsa*, *Date tree*.

PHŒNODINE. (from *φαινός*, 'blood red,') *Hæmatin*.

PHONAS'CIA. (*φωνή*, 'voice,' and *ασκία*, 'to exercise,') The art of exercising the voice.

PHONATION. *Phonatio*. *Phonè'sis*, *Phonè'sis*, (from *phone*.) The physiology of the voice.

PHONE. (*φωνή*.) *Voice*.

PHONESIS. *Phonation*.

PHONETIC. *Phonè'ticus*, *Pho'nic*, *Pho'nicus*, (F.) *Phonétique*, *Phonique*, (*φωνητικός*, from *φωνή*, 'sound,' 'voice,') Relating or appertaining to sound or to the voice.

PHŒNICA. (*φαινικά*, from *phone*.) The first order of the class *Pneumonia*, in Good's Nosology. Defined: diseases affecting the vocal avenues:—the passage of the voice impeded, or its power suppressed or depraved.

PHŒNIQUE. *Phonetic*.

PHONON'OSI. *Phonon'osi*. (*phone*, and *ωσος*, 'a disease,') *Phonopath'ia*, *Phon'ic'osis*. Diseases of the voice.

PHONOSIS. *Phonation*.

PHONOPATHIÆ. (*phone*, and *πάθος*, 'affection,') *Phonosis*.

PHORA. (*φορά*.) *Gestation*.

PHORBE. (*φορβή*.) *Palatum*.

PHORONOMIA. (*φωρ*, 'I move,' and *νομος*, 'law,') The doctrine of muscular motion.

PHOROUS. (*φορος*, from *φωρ*, 'I move or carry,') A suffix denoting conveyance. Hence, *Galactophorous*, &c.

PHOS. (*φως*, gen. *φωτος*.) *Light*.

PHOSPHAS. gen. *Phosphatis*, *Phosphate*. A salt formed by the union of phosphoric acid with a salifiable base.

PHOSPHAS CALCIUM STIBIATUM. *Antimonial powder*—p. *Calci stibiatum*, *Antimonial powder*—p. *Natrium*, *Soda*, *phosphate of*.

PHOSPHATE. *Phosphas*.

PHOSPHATIC. *Phosphat'icus*, (F.) *Phosphatique*, (from *Phosphorus*.) Relating to the phosphates; hence the *Phosphatic Diathesis* is the

habit of body which favours or occasions the deposition of the phosphates from the urine.

PHOSPHÈNE, (*phos*, and *φαίνω*, 'to become manifest,') Photopsia.

PHOSPHORE, Phosphorus.

PHOSPHORÈNÈSES. M. Baumes unites, under this generic name, diseases which he attributes to disordered phosphorization; i. e. to excess or deficiency of calcareous phosphates or its decomposition. Amongst these affections he ranks rachitis, osteomalacia, and gout.

PHOSPHORES'CENCE, *Phosphorescen'tia*, *Luminous'ity*. (from *phosphorus*.) The properties possessed by certain bodies of being luminous in obscurity, or under particular circumstances. This phosphorescence has been occasionally witnessed in man, when suffering from some wasting disease, and near death.

PHOSPHOR'IC AC'ID, *Ac'idum phosphor'icum*, (F.) *Acide phosphorique*. A solid, colourless compound; highly sapid; very soluble in water, and vitrifiable by fire. It exists in bones combined with lime. It has been employed in medicine in the form of lemonade, in passive hemorrhage, typhus, phthisis, marasmus, &c., and in injection in cases of cancer of the uterus. It does not seem preferable to other acids. *Acidum phosphoricum glaciale*, and *Acidum phosphoricum dilutum* are official in the Pharmacopœia of the United States of 1863.

PHOSPHORNECROSIS, (*phosphorus* and *ne-crosis*), see Phosphorus.

PHOSPHORURIA, (*phosphorus*, and *ουρον*, 'urine,') Urine, phosphorescent. Also, urine containing an excess of phosphates. See Urine.

PHOS'PHORUS, (*φωσφορος*, 'a lightbearer,' from *phos*, and *φέρω*, 'to carry,') *Phos'phorum*, *Auto-phos'phorus*, *l'g'nus philosoph'icus*, *Lu'men constans*, (*φως*, 'light,' and *φέρω*, 'I carry,') (F.) *Phosphore*. An elementary, simple, or undecomposed substance; not metallic; solid; transparent or semitransparent; colourless; brilliant; flexible; of a sensible garlic odour; very inflammable, and susceptible of combining with the oxygen of the atmosphere, at even the ordinary temperature. S. g. 1.8. It does not exist pure in nature, but is found frequently united to other bodies. Its fumes in the manufacture of lucifer and Congreve matches have caused, it is said, neurosis of the lower jaw—*Phosphorneuro'sis*, *phosphorus jaw disease*, *Lu'cifermatchmakers' Disease*, *Matches Disease*.

Phosphorus has been used in medicine, dissolved in ether or suspended in the yolk of egg, as a stimulant; but the greatest precautions are necessary in its administration: hence it is rarely employed.

PHOSPHORUS JAW DISEASE, see Phosphorus.

PHOSPHURIA, (*phosphorus*, and *ουρον*, 'urine,') Urine, phosphorescent. Also, urine containing an excess of phosphates. See Urine.

PHOTOCAMPSIS, (*phos*, and *καμψις*, 'bending,') Refraction.

PHOTOMANIA, (*phos*, and *μανία*,) Insane delirium induced by light.

PHOTOPHOBIA, *Phenopho'bia*, (*phos*, and *φοβος*, 'fear,') Aversion to light:—intolerance of light. Nyctalopia.

PHOTOPHOBICUS, *Lucifugus*.

PHOTOPHOBOPHTHAL'MIA, (from *phos*, *φοβος*, 'dread,' and *ophthalmia*,) Ophthalmia with great intolerance of light.

PHOTOPHOBUS, *Lucifugus*.

PHOTOP'SIA, *V'isus lu'cidus*, (*phos*, and *ωψ*, 'the eye,') A false perception of light, as of sparks, flashes of fire, &c., occasioned by certain conditions of the retina and brain, and by pressure on the eyeball. When pressure does not induce the luminous appearance, (F.) *Phosphène*,

the existence of amaurosis may be inferred. See Metamorphopsia.

PHOTURIA, (*phos*, and *ουρον*, 'urine,') A luminous condition of the urine.

PHRAGMUS, (*φραγμος*, 'a fence,' from *φρασσω*, 'I enclose,') A row of teeth.

PHRASIS, (*φρασις*,) Voice, articulated.

PHRASUM, *Marrubium*.

PHREN, gen. *Phren'is*, (*φρην*, 'the mind,' and the organ of the mind.) Also, the diaphragm and scrobiculus cordis.

PHRENATROPH'IA, *Atroph'ia cer'ebr'i*, (*phren*, a, priv, and *τροφή*, 'nourishment,') Wasting or atrophy of the brain.

PHRENAUX'É, *Hypertroph'ia cer'ebr'i*, (*phren*, and *αυξη*, 'increase,') Hypertrophy of the brain.

PHRENES, (*φρενες*,) Diaphragm.

PHRENESIA, (from *phren*,) Phrenitis.

PHRÉNÉSIE, Phrenitis.

PHRENESIS, (*φρενησις*,) Delirium, Phrenitis.

PHRENITIASIS, Delirium, Phrenitis.

PHRENET'IC, (*φρενητικός*,) *Phrene'ticus*, *Phreni'ticus*, *Frene'tic*, (F.) *Phrénétique*. Relating or appertaining to phrenitis; frantic; mad.

PHRENITISMUS, Phrenitis.

PHRENIC, (from *phren*,) Diaphragmatic.

PHRENICA, (from *phren*,) Diseases affecting the intellect; characterized by error, perversion, or debility of one or more of the mental faculties, (*Phrenopath'ia*.) The first order of the class *Neurotica* of Good. Also, remedies that affect the mental faculties—*Phren'ica*.

PHRENICUS, see Phrenica.

PHRENICULA or **PHRENETICULA HYDROCEPHALICA**, *Hydrocephalus internus*.

PHRÉNIQUE, Diaphragmatic.

PHRENISMUS, Phrenitis.

PHRENITICUS, (*φρενητικός*,) Phrenetic.

PHRENITIS, gen. *Phreni'tidis*, (*φρενιτις*, gen. *φρενιτιδος*,) *Phrene'sis*, *Phrene'sia*, *Phreneti'asis*, *Phrenis'mus*, *Phrenetis'mus*, *Cephal'i'tis*, *Encephali'tis*, *Cerebri'tis*, *Cephalalgia inflammato'ria*, *Karabitus* (Arab.), *Sibare*, *Sphacelis'mus cer'ebr'i*, *Empres'ma cephaliti's*, *Siri'asis*, *Sciri'asis*, *Sideratio*, *Inflammat'io seu Caum'a phreni'tis*, *Phren'tey*, *Frenzy*, *Delir'ium*, *Brain Fever*, (F.) *Phrénésie*, *Cérèbrite*, *Inflammation du cerveau et du cercelet*, (from *phren*, and *itis*,) A disease characterized by violent pyrexia, pain in the head, redness of the face and eyes, intolerance of light and sound, watchfulness, and delirium, either furious or muttering. Its causes are the same as those of other inflammations: sometimes, however, it is induced by a *Coup de soleil*: by inebriation; violent mental emotions, &c. Some authors have given the name *Phreni'tis* and *Meningi'tis* to inflammation of the membranes of the brain; and that of *Cephal'i'tis*, *Encephali'tis*, or *Cerebri'tis* to inflammation of the substance of the brain; but there are no pathognomonic symptoms, which can lead us to refer the seat of the disease to either of those parts exclusively. Phrenitis is, of course, a very dangerous affection, from the importance of the organ concerned. The most active treatment must necessarily be pursued. Repeated bleeding; purging; the application of ice to the shaved head; a position in which the head is raised; and the avoiding of irritation of every kind;—form the remedial means during the violent stage. After this has yielded, counterirritants to the nape of the neck or to other parts of the body will be useful adjuvants. When the brain and meninges were affected only symptomatically, as in inflammation of the diaphragm, the disease was formerly called *Paraphreni'tis*.

Hydrocephalus acutus is a variety of phrenitis.

PHRENITIS CALENTURA, *Coup-de-soleil*—p. Latrans, *Hydrophobia*—p. Potatorum, *Delirium tremens*—p. Vermiosa, *Verocoquin*.

quired to rouse the predisposition into action. It is less a disease of the torrid and frigid zones; and in the temperate regions its pathology is often intimately allied with that of scrofula. It consists in the formation of tubercles in the lungs, which sooner or later soften and break down. In such a constitution, ulcerations of the lungs do not readily heal; and hectic fever—the universal attendant upon irritability and debility—is established. This persists, and gradually wears down the patient, till he ultimately expires, exhausted by the irritative fever. The chief symptoms are the following. The patient feels a sense of uneasiness in the chest, with severe fits of coughing, or a short and dry cough with dyspnea. This may be considered the incipient stage. The cough then becomes extremely troublesome, especially in the morning; and dyspnea is increased on the slightest exertion. The expectoration is evidently purulent, with fever in the evening, and circumscribed redness of the cheeks; wasting and colliquative sweats early in the morning. This must be considered the confirmed stage.—*Phthisis consummata*, *Pneumophthisis*. In the third, or last stage, the hectic is constant, with small pulse, pungent heat, colliquative sweats, and diarrhoea. The weakness becomes extreme, and the patient exhausted. *Physical Signs*.—1. In the first stage, whilst there are yet scattered miliary tubercles,—the only appreciable change at the exterior of the chest, and this is not constant, is a want of freedom in the movements under one or both clavicles. Percussion will generally discover a slight diminution of sound corresponding to the summit of one lung. In the same situation, auscultation shows that the inspiratory murmur is either weaker, rougher, or confused. The sound of expiration is more distinct. The voice and cough are rather more resonant, and wheezing or mucous râles are occasionally heard in the same region. 2. When the tubercular deposition has become more abundant, the motion of the chest is more markedly diminished under one or both clavicles. The loss of sound on percussion is more distinct, the inspiration, on auscultation, is bronchial, the expiration louder and more superficial, and the resonance of the voice and cough is increased. The rest of the lung is perhaps healthy, or the respiration may be purile. 3. When the tubercles are undergoing softening,—in addition to the preceding signs, auscultation exhibits a subcrepitant râle around the apex of the lung, decreasing from above downwards. It is most distinct during or after the cough, and at the close of a deep inspiration. As the disease proceeds, the bubbling becomes successively larger, and ultimately gurgling is heard. The inspiration and expiration are cavernous or tracheal. The voice and cough indicate unusual resonance and pectoriloquy. The intercostal spaces are often depressed locally, and percussion may become gradually clearer and even tympanitic as the cavern increases.

It is only in the incipient state of this formidable disease, that advantage can be expected from any plan of treatment; and of all means that can be devised, that of removing to a climate in which air and exercise can be daily enjoyed during the autumnal, winter, and spring months is the best. See Climate. If this be impracticable, the patient must be kept in a regulated temperature, taking care that the chamber is duly ventilated; and gestation in the open air be used, whenever the atmosphere, although cold, is dry. The other indications will be: 1. To moderate inflammatory action: 2. To support the strength; and, 3. To palliate urgent symptoms. The whole treatment indeed is palliative. Counterirritants relieve pain: demulcents and mild

opiates alleviate the cough: the mineral acids, united with opium, relieve the colliquative sweats; and chalk and opium the diarrhoea.

Dr. Wilson Philip has pointed out a particular species of phthisis, which is characterized, in the early stage, by a deranged state of the digestive organs: flatulent, irregular bowels; furred tongue; impaired appetite; unnatural fæces; and the epigastric region more or less full and tender on pressure. The affection of the lungs is here secondary. He calls it *Dyspeptic phthisis*.

PHTHISIS PULMONALIS PURULENTA EXULCERATA, *Phthisis pulmonalis*—p. *Pulmonalis scrofulosa*, *Phthisis pulmonalis*—p. *Pulmonalis tuberculosa*, *Phthisis pulmonalis*—p. *Pulmonalis tuberculo-ulcerata*, *Phthisis pulmonalis*—p. *Pulmonalis vera*, *Phthisis pulmonalis*—p. *Pupillæ*, *Synechisis*—p. *Pupillaris*, *Myosis*—p. *Scrophulosa*, *Phthisis pulmonalis*—p. *Sicca*, *Tubercularis*—p. *Tuberculosa*, *Phthisis pulmonalis*—p. *Uterina*, *Metrophthisis*—p. *Vesicularis*, *Cystophthisis*.

PHTHISIS WEED, *Ludwigia palustris*.

PHTHISURIA, (*phthisis*, and *ουρον*, 'urine,') *Diabetes*.

PHTHISURIA, *Diabetes*—p. *Sucre*, *Diabetes mellitus*.

PHTHOË, (*φθον*,) *Phthisis*.

PHTHOIS, gen. *Phthoidis*, (*φθοις*,) *Pastil*, *Trochiscus*.

PHTHOISCUS, (*φθοισκος*,) *Pastil*, *Trochiscus*.

PHTHONGODYSPHORIA, (*φθονγγος*, 'a cry,' and *δυσφορία*, 'impatience,') *Hyperæmis*.

PHTHORA, (*φθορα*, 'destruction,') *Abortion*, *Corruption*, *Plague*.

PHTHORIUS, (*φθοριος*,) *Abortive*.

PHTHOROCTONUS, (*φθορα*, 'destruction,' and *τεκνον*, 'a child,') *Abortive*.

PHUCAGROSTIS MINOR, *Pila marina*.

PHUSCA, (*φουσκα*,) *Oxyerate*.

PHYGANTHROPIA, (*φανα*, 'to flee,' and *ανθρωπος*, 'man,') *Misanthropia*.

PHYGETH'LON, (*φυγεθλον*,) *Pa'nus*, *Phypella*, (from *φυω*, 'I grow,' [?].) *Inflammation of the superficial lymphatic glands, not ending in suppuration*.

PHYGEDEMA, (*φυγη*, 'flight,' and *oedema*,) *Anathymiasis*.

PHYLACTERION, *Amuletum*.

PHYLACTERIUM, (*φυλακτηριον*, from *φυλασσω*, 'I preserve,') *A Phylactery*, (F.) *Phylactère*. An amulet:—a prophylactic.

PHYLAXIS, (*φυλαξ*,) *Conservation*.

PHYLLANTHUS URINARIA, (*φυλλον*, 'a leaf,' and *ανθος*, 'a flower,') *Binkohumet*. *Ord.* *Euphorbiaceæ*. A plant employed in Ceylon, in decoction, as a diuretic.

PHYLLITIS, (*φυλλον*, 'a leaf,') *Asplenium scolopendrium*—p. *Muraria*, *Asplenium ruta muraria*—p. *Rotundifolia*, *Asplenium trichomanoides*.

PHYMA, gen. *Phymatis*, (*φυμα*, gen. *φυματος*, from *φυω*, 'I grow,') With the ancients, this word was used with great latitude: sometimes, in the sense of phlegmon, for tumours larger than a boil, and perfectly suppurative. In the nosology of Sauvages, *Phymata* is an order of diseases, to which he refers erysipelas, oedema, phlegmon, emphysema, scirrhus, &c. In that of Dr. Good it is a genus, including hordeolum, furunculus, syccosis, and anthrax.

PHYMA ANTHRAX, *Anthrax*—p. *Anthrax terminus*, *Terminus*—p. *Syccosis*, *Syccosis*.

PHYMATODES, (*φυματωδης*,) *Affected with phymata*.

PHYMATOID, *Phymatoides*, (F.) *Phymatoides*, (*phyma*, and *ειδος*, 'resemblance,') A state of cancerous tissue, in which the dull yellow hue and the consistence resemble those of tubercle.

PHYMATOSES, (from *phyma*, and *osis*,) *Tuberculous diseases*.

PHYMATOSIS, *Excrecence*—p. *Cerebri*, *En-*

cephalophymata—p. Elephantiasis, Elephantiasis—p. Lupus, Lupus—p. Pulmonum, see Tubercle—p. Verrucosa, Verruca.

PHYMOSIS, Phimosi.

PHYELLA, Phyethlon.

PHYSA, (*ψαα*, 'wind,') Crepitation, Flatulence.

PHYSAGOGA, (*ψαα*, 'wind,' and *αγω*, 'to carry off,') Carminatives.

PHYSALIS, (*ψααδης*, 'a bladder,' referring to the inflated calyx,) *P. Alkeken'gi* seu *halicac'abum*, *Solanum vesicarium*, *Alchucen'gê*, *Alkeken'gi*, *Halicac'abum*, *Winter Cher'ry*, (F.) *Alkekengi*, *Copurca*. Ord. Solanaceae. *Sex. Syst.* Pentandria Monogynia. The berries are recommended as a diuretic, from 6 to 12 for a dose, in dropsical and calculous cases. It has been affirmed, that the whole plant—twigs, leaves, capsules and berries—possesses the antiperiodic virtues of Cinchona.

PHYSALIS ALKEKENGII, Physalis—p. Halicacabum, Physalis—p. Obscura, Ph. viscosa—p. Pennsylvanica, Ph. viscosa—p. Pubescens, Ph. viscosa—p. Stramonium, Anisodus luridus.

PHYSALIS VISCOSA, *P. obscura* seu *Pennsylvanica* seu *pubescens*, *Ground cherry*; indigenous. The berries are said to be diuretic and sedative.

PHYSCE, (*ψαα*.) Abdomen, Colon.

PHYSICA ISLANDICA, Lichen Islandicus—p. Nivalis, Lichen caninus.

PHYSICONIA, (*ψαακον*, 'the paunch,') *Ventros'itus*, *Euphy'ma physiconia*, *Euphrac'tica*, *Visconia*, *Pathelly*, *Tumbelly*, *Pendulous abdom'en*, *Stucagbilly*, *Gorbelly*, (Prov.) *Gotchbilly*, *Pedbilly*. Physiconia is a generic term, under which Sauvages has compromised every large tumour developed in the abdomen, that is neither fluctuating nor sonorous. Linnæus calls it *Hypoposarca*. Often, these various terms are applied to excessive corpulency which gives to the abdomen the appearance indicated by them.

PHYSICONIA ADIPO'SA, *Obes'itus* seu *Polysar'cia viscerum*. Accumulation of fat in the intestines.

PHYSCONIA BILIOSA, *Turgescencia vesiculæ fellæ*.

PHYSCONIA EMPHYSEMA'TICA. Accumulation of air in the areolar tissue, or between the coats of particular viscera.

PHYSCONIA LIENS, Splenoneus—p. Mesenterica, *Tabes mesenterica*—p. Splenica, *Ague cake*.

PHYSCONIA STRIMO'SA, *Serpy'ula abdomin'alis inter'na*. Tumor abdomen from scrophulous enlargement of the glands.

PHYSCONIE MÉSÉNTÉRIQUE, *Tabes mesenterica*.

PHYSECHEMA, (*ψαα*, 'wind,' and *ηχηα*, 'a sound,') *Bruit de soufflet*.

PHYSEMA, (*ψααα*,) Tympanites, Phyesis.

PHYSESIS, *Physe'ma*, (*ψαασις*, from *ψααω*, 'I inflate,') An emphysematous tumour. Tympanites.

PHYSETER MACROCEPHALUS, (*ψαατηρ*, 'one that blows,') see Cetaceum.

PHYSIATRICE, (*physis*, and *ιατρεα*, 'treatment,') Vis medicatrix nature.

PHYSIAUTOCRATIA, (*physis*, *αυτος*, 'self,' and *κρατος*, 'power,') Vis medicatrix nature.

PHYSIAUTOTHERAPIA, (*physis*, *αυτος*, 'self,' and *θεραπια*, 'treatment,') Vis medicatrix nature.

PHYSIC, (from *physis*,) Medicament, Medicina. To *physic* is to administer medicine, (F.) *Medicament*; and, at times, to purge.

PHYSIC, INDIAN, *Euphorbia corollata*, *Gillenia trifoliata*—p. Root, *Leptandra purpurea*.

PHYSICAL, Medical—p. Signs, see Symptom.

PHYSICIAN, *Med'icus*, *Int'ros*, *Aces'tor*, *Aces'ter*, *All'thus*, *Pæon*, (Old Eng.) *Fesic'ian*, *Ficicion*, *Leche*, *Leech*, *Leach*, *Leachman*, (Sc.)

Mediciner, *Medicinare*, *Medcinare*, *Medicinar*, (*ψαασις*, 'an investigator of nature,' *ψαασις*,) (F.) *Médecin*. Properly, one who has received his degree from an incorporated Institution, as Doctor of Medicine; but often applied in the United States to any one who practises physic. The French formerly used the word *physicien* in the same sense. It is now appropriated by them to the natural philosopher. *Phys'icist* has the same signification.

PHYSICIAN-ACCOUCHEUR, *Accoucheur*—p. Alienist, Psychiater—p. Botanical, Herb doctor.

PHYSICIEN, FAMILY. One regarded as the regular attendant on a family.

PHYSICIAN LEARNED, *Iatrosophista*—p. Practical, *Iatrotechna*—p. Priest, see *Aselepiadæ*.

PHYSICIEN, Physician.

PHYSICIST, (from *physis*,) see Physician.

PHYSICS, MEDICAL, *Iatrophys'ica*. Physics directly applied to medicine,—either to the explanation of the vital phenomena, the preservation of the individual, or the treatment of disease.

PHYSIOAUTOCRATIA, (*ψαασις*, 'nature,' *αυτος*, 'self,' and *κρατος*, 'power,') Vis medicatrix nature.

PHYSIOGNOMIST, *Physiognom'on*, one who busies himself especially with, or pretends to be versed in, physiognomy.

PHYSIOGNOMY, (Old Eng.) *Vis'nomy*, *Physiognom'ia*, *Physiogn'mica*, *Physiogn'micê*, *Physiogn'mia*, *Physiognomon'ia*, *Physiogn'monê*, *Physiognomos'ynê*, *Physiognom'o'sia*, *Anthroposcopia*, *Prosopomanti'a*, *Prosopolog'ia*, *Semiol'icê*, *fac'icis*, *Physiognomon'ica*, *Physiognom'ony*, *Me'toposcopia*, *Prosop'o'sia*. (F.) *Physiognomie*, (*επισκοπωμα*, from *physis*, 'reality,' and *γνωσις*, 'knowledge,') The art of knowing the dispositions of men from their external appearance; especially from the features. Every inference of this kind must necessarily be extremely fallacious, and cannot be reduced to rule, as was attempted by Lavater and others. In infantile diseases, numerous shades of expressions are evident, which experience teaches how to appreciate; and which afford useful guides in understanding the pathology of that period of existence.

PHYSIOLOGIA PATHOGENETICA, see Pathogenetic.

PHYSIOLOGICAL, *Physiolog'icus*, (F.) *Physiologique*. Relating or belonging to physiology.

PHYSIOLOGICAL ANATOMY, see Anatomy—p. Doctrine, Broussaism.

PHYSIOLOGICE, Physiology.

PHYSIOLOGISM, *Physiologis'mus*. Abuse of physiology. The improper application of physiological data to the theory of disease.

PHYSIOLOGIST, *Physiol'ogus*. One who occupies himself specially, or is well acquainted, with physiology.

PHYSIOLOGY, *Physiolog'ia*, *Biol'ogy*, *Bion'omy*, *Biol'ica*, *Biophænomenolog'ia*, *Physiolog'icê*, (*Εκνομ'ia animi'alis*, *Anatom'ia vi'va*, *Anat'omê animi'ata*, *Anthropolog'ia*, *Microcosmog'phia*, *Microcos'mica scien'tia*, *Cognit'io physiolog'ica*, *Pars natura'lis medic'ine*, *Philosoph'ia corporis vi'vi*,) (from *ψαασις*, 'nature,' and *λογος*, 'a description,') Formerly, Physiology meant the same as Physics, in its extensive signification, now does. At the present day, it includes the science which treats of the functions of animals or vegetables; an acquaintance with the phenomena the aggregate of which constitute life. It is the science of life. It is divided into *animal*—*Zoôphysiolog'ia*, or *Zoobiolog'ia*; and *vegetable*—*Phytophysiol'og'ia*, or *Phytobiolog'ia*, according as it considers the life of animals or of vegetables singly. *Comparative physiolog'ogy* comprises both. Physiology is, also, *general* or *special*, according as it treats of life in the abstract or in some particular

species. To the latter belongs the *Physiology of Man*—called, also, *Hygienic Physiology* to distinguish it from *Pathological Physiology* or *Pathology*.

PHYSIOLOGY, PATHOGENETIC, see Pathogenetic.

PHYSIONOMIA, Physiognomy.

PHYSIONOMIE, Physiognomy—p. *Égarée*, see Wild.

PHYSIS, (φύσις,) Natura.

PHYSO, (from φύσα, 'air,') in composition, air.

PHYSOCE'LE, Pneumatocoe'le, Her'nia vento'sa, Oscheocoe'le flatulen'ta, Emphyse'ma Sero'ti, (physo, and κλη, 'a tumour,') An emphysematous tumour of the serotum; probably a case of intestinal hernia, containing much flatus.

PHYSOCEPH'ALUS, (physo, and κεφαλη, 'head,') (F.) *Physocéphale*. Emphysematous tumefaction of the head.

PHYSOCELIA, (physo, and κυλία, 'intestines,') Tympanites.

PHYSO'DES, (φυσώδης, from φύσα, 'air,' and odes,) Full of air; apparently full of air—applied to tumours, &c.

PHYSOME'TRA, Emphyse'ma U'teri seu Uteri'nium, Hysteroph'ysē, Hysteremphyse'ma, Metremphyse'ma, Infla'tio U'teri, Tym'pany of the Womb, U'terine tympani'tes, (from physo, and μτρα, 'the womb,') A light, tense, circumscribed protuberance in the hypogastrium, obscurely sonorous, with wind occasionally discharged through the os uteri with noise, *Hysteropsoph'ia*, *Ædæropsoph'ia uteri'na*.

PHYSONCUS, (physo, and ογκος, 'a swelling,') Emphysema.

PHYSOPSOPH'IA, (from physo, and ψοφος, 'a noise,') A discharge of air with noise.

PHYSOSPASMUS, Colica flatulenta.

PHYSOSTIGMA VENENOSUM, (physo, and stigma, in Botany, the top of the style,) Ordeal nut.

PHYSOSTIGMINE, see Ordeal nut.

PHYSOTHORAX, gen. *Physothora'cis*, Pneumothorax.

PHYTANATOMY, (phyton, and anatomy,) see Anatomy.

PHYTEL'EPHAS MACROCAR'PA, (phyton, and ελεφας, 'an elephant,') A tree of South America, called, by Willdenow, *Elephanta'sia*, the dense albumen of whose nut is the vegetable ivory, (F.) *Ivoire végétal*, which is as durable and nearly as hard as the ivory of the elephant.

PHYTICUS, (φυτικός,) Vegetative.

PHYTIVOROUS, (phyton, and voro, 'I devour,') Phytophagous.

PHYTOALOPECIA, (phyto, and alopecia,) Porrigo decalvans.

PHYTOBIOLOGIA, (phyton, βιος, 'life,' and λογος, 'a discourse,') see Physiology.

PHYTOCHYMISTRY, see Chymistry.

PHYTOLAC'CA DECAN'DRA, *P. vulga'ris*, *Pokeweed*, *American nightshade*, *Cancerroot*, *Poke*, *Shoke*, *Jucato Calleloe* [?], *Ja'vap*, *Pig'leon Ber'ry*, *Jalap can'cer root*, *Red weed*, *Red nightshade*, *Coakum*, *Sola'nium racem'o'sum* *America'nium* seu *nag'nium* *Virginia'nium* *ru'brum*, *Blu'tum* *America'nium*, *Gar'get*, (F.) *Raisin d'Amérique*, *Physolaque à dix étamines*, *Morelle en grappes*, *Méchoucan du Canada*. Ord. Chenopodaceae; Sox. Syst. Decandria Decagynia, (phyton, and lacca, because the colour of the juice of the berries resembles that of lacca.) The berries—*Phytolac'cæ Bac'ca* (Ph. U. S., 1863)—and roots—*Phytolac'cæ Ra'dix*—are employed. The young stems, when boiled, are sometimes eaten at table; but when old, they must be cautiously used, as the plant is emetic and cathartic. It is most celebrated as a remedy in chronic rheumatism, and is given in the form of tincture of the ripe berries. It is, also, used in cutaneous eruptions, tinea capitis, itch, cancerous ulcers, &c.

The root is gathered and dried in the fall.

PHYTOLAQUE À DIX ÉTAMINES, *Phytolacca decandra*.

PHYTOLOGIA, (phyton, and λογος, 'a discourse,') see Vegetable—p. *Medica*, Botany, medical.

PHYTON, (φυτον,) Vegetable.

PHYTOPARASITE, Epiphyte.

PHYTOPATHOLOGY, *Phytopatholog'ia*, (from phyton, and pathology,) Properly, the pathology of vegetables; but occasionally applied to the knowledge of morbid states induced by vegetable growths.

PHYTOPH'AGOUS, *Phytoph'agus*; *Phytiv'orous*, *Phytiv'orus*; (phyton, and φάγω, 'I eat,') Feeding or subsisting upon vegetables.

PHYTOPHYSIOLOGIA, (phyton, and physiologia,) see Physiology.

PHYTORGANIES, (phyton, and οργανον, 'an organ,') A name given by Bouchet to vegetable growths on the living body.

PHYTOSPERM'ATA, (F.) *Phytospermes*, (phyton, and σπέρμα, 'sperm,') Particles similar to the spermatozooids of animals, observed in vegetables, and presumed to possess a similar agency in reproduction.

PHYTOTOMY, (phyton, and τέμνειν, 'to cut,') see Anatomy, (vegetable.)

PHYTOZO'AIRE, (phyton, and ζωον, 'animal,') Zoophyte.

PHYTOZOON, (phyton, and ζωον, 'animal,') Zoophyte.

PI'A MA'TER, 'delicate mother,' *Pia* seu *moll'is* seu *ten'uis* *Mater*, *Pia me'ninx*, *Meninx inte'rior* seu *vasculo'sa* seu *choroi'des*, *In'ner lam'ina of the Méninge*, (Ch.), *Membra'na vasculo'sa Cer'ebr'i*, *M. Cer'ebr'i ten'uis* seu *moll'is* seu *pro'pria*, (F.) *Pie mère*, *Méningette*. The pia mater is a very delicate membrane, which covers the brain immediately, penetrates into its anfractuositities, and envelops, also, the cerebellum and the spinal prolongation, &c. Bichat does not regard it as a membrane, properly so called, but as a network of blood-vessels, united by a lax areolar tissue. The *Pia Mater exterior* is the portion of membrane which covers the exterior of the encephalon, and faces the arachnoid externally. The *Pia Mater interior* is that which lines the cerebral fosse, into which it penetrates by a cleft situate between the posterior part of the corpus callosum and the fornix; and by two other fissures, situate between the corpora fimbriata and thalami nervorum optico'rum. The pia mater contains a great number of granulations similar to the *Glandulae Puccioni*.

PIA MATER TESTIS, Tunica vasculosa—p. *Meninx*, *P. Mater*.

PIAN, *Frambesia*.

PIANS, MOTHER OF, *Mama-pian*.

PIAR, (πιαρ,) Fat.

PIARHÆ'MIA, *Piarrhæ'mia*, *Lipæ'mia*, (*piar*, and αἷμα, 'blood,') Fat in the blood.

PICA, Malacia—p. *Africanorum*, *Chthonophagia*.

PICAC, *Euphorbia corollata*.

PICACISM'US, *Pica'tio*, (*pix*, gen. *picis*, 'pitch,') Depilation by means of a pitch plaster.

PICATIO, Picacismus.

PICEA, (*pix*, gen. *picis*, 'pitch,') *Pinus abies*.

PICERION, (πικριον,) Butter.

PICHU'RIM BEANS. The seed,—*Fa'ba Pichurim*, *F. Pechu'rei*, *Pa'ra nut*, *Sas'safras nut*, *Puciri*, *Brazil'ian nut'meg*,—of a tree, supposed to be *Per'sea pichurim*, *Ocote'a pichu'rim* seu *Puchu'ry*, *Tetranthe'ra pichurim*, *Necton'dra puchu'ry*, *Puchu'ry* or *Puchy'ry* of the Brazilians, Ord. Lauraceae, which grows in Brazil, Guiana, Venezuela, and other parts of South America. They are of an elliptical shape; flat on one side, convex on the other; of an aromatic odour between that of

nutmeg and of saffrafas, and of a spicy, pungent taste. The medical properties are the same as those of other aromatics. For the bark, see *Pichurim cortex*.

PICK, (from *puke*.) Emetic.

PICKERELWEED, SHOVEL, *Unisema deltifolia*.

PICOTE, Variola.

PICOTEMENT, (from *piquer*, 'to prick,') Pricking.

PICOTIN, *Arum maculatum*.

PICRA, (from *πικρος*, 'bitter,') *Pulsis aloës et canellæ*.

PICRÆNA EXCELSA, (*πικραῖνα*, 'to be bitter,') Quassia.

PICRIA, (*πικρία*.) Bitterness.

PICRIS, (*πικρος*, 'bitter,') *Cichorium intybus*.

PICRUM SPICATUM, *Coutoubea alba*.

PICROCHOLOS, (*πικροχολος*, from *picros*, and *χολη*, 'bile,') Bilious.

PI'CROMEL, *Bi'tlin*, (*picros*, and *μελι*, 'honey,') A name given by Thénard to a colourless, soft substance, of an acrid, bitter, saccharine taste, nauseous smell, and very soluble in water, which is considered to be a *bi'tlate* of *so'da*. It is not an immediate principle, but a mixture of different substances. It forms part of the bile, and is generally classed amongst the animal immediate principles.

PICROPE'GÆ, *Picrope'gæ*, (*picros*, and *πηγή*, 'a spring,') Mineral waters bitter from the presence of chloride of magnesium.

PICROS, (*πικρος*, 'bitter,') *Amarus*.

PICROTOX'IN, *Picrotox'ine*, *Picrotox'ine*, (*picros*, and *toxicum*, 'poison,') A vegetable alkali, discovered by M. Boullay in the *Menispermum Cocculus*. It crystallizes in quadrangular, white, brilliant, semitransparent prisms; which are excessively bitter, soluble in three parts of alcohol, and in fifteen of boiling water.

The *Cocculus Indicus* owes its poisonous properties to this principle.

PIE PLANT, *Rheum rhaponticum*—*p. Rhu-barb*, *Rheum rhaponticum*.

PIE MÈRE, *Pia mater*.

PIE-MÉRITE, *Inflamma'tio pi'æ ma'tris*, (from *pie mère*, and *ite*, for *itis*.) Inflammation of the *pia mater*.

PIÈCE, Hepatitis—*p. d'Anatomie Artificielle*, see Artificial.

PIED, *Pes*—*p. d'Alexandre*, *Anthemis pyrethrum*—*p. d'Alouette des champs*, *Delphinium consolida*—*p. de Biche*, see *Lever*—*p. Bot*, see *Kyllosis*—*p. de Chat*, *Antennaria dioica*—*p. de Cheval marin*, *Cornu ammonis*—*p. de Griffon*, *Gryphius pes*, *Helleborus foetidus*—*p. d'Hippocampe*, *Cornu ammonis*—*p. de Lièvre*, *Arum maculatum*—*p. de Lion*, *Alchemilla*—*p. de Lit*, *Origanum*—*p. d'Ours*, *Acanthus mollis*—*p. Plat*, see *Kyllosis*—*p. de Veau*, *Arum maculatum*—*p. de Veau triphyll*, *Arum triphyllum*.

PIEROS, (*πικρος*.) Fatty.

PIERRE, (from *petrus*, *πετρος*, 'a stone,') Calculus—*p. d'Aigle*, *Ætites*—*p. d'Aimant*, *Magnet*—*p. d'Azur*, *Lapis lazuli*—*p. Caluminaire*, *Calamine*—*p. à Caùtère*, *Potassa cum calce*—*p. Divine*, *Lapis divinus*—*p. d'Ècervise*, *Cancerorum chelæ*—*p. Infernale*, *Argenti nitras*—*p. Judaïque*, *Judæus (lapis)*—*p. au Lait*, *Morochthus*—*p. Médicamenteuse*, *Medicamentosus lapis*—*p. Néphrétique*, *Nephreticus lapis*—*p. Ponce*, *Pumex*.

PIERRES CRAYÈSES, 'chalk stones,' *Calculi*, *arthritis*—*p. au Fiel*, *Calculi*, *biliary*—*p. Stercorales*, *Calculus of the Stomach and Intestines*—*p. Urinaires*, *Calculi*, *urinary*.

PIERREUX, *Petrus*.

PIESTER, (*πιεστηρ*.) Press.

PIESTERION, (*πιεστηριον*.) Press.

PIESTRON, (*πιεστρον*, from *πιεζω*, 'I press,')

An instrument recommended by Hippocrates for breaking the bones of the foetal head when too large to be extracted otherwise.

PIGAMON JAUNÂTRE, *Thalictron*.

PIGEON BERRY, *Phytolacca decandra*—*p. Breasted*, see *Lordosis*—*p. Tree*, *Aralia spinosa*.

PIGMENT CELLS, see *Cell*, *pigment*.

PIGMENTAIRE, *Pigmental*.

PIGMENT'AL, *Pig'mentary*, *Pigmenta'rius*, (*F.*) *Pigmentaire*, *Pigmenteux*, (from *pigmentum*.) Having relation to pigment or colouring matter.

PIGMENTAL APPARA'TUS, (*F.*) *Appareil pigmen-tal*. The apparatus concerned in the coloration of the skin of the coloured varieties of man.

PIGMENTARIUS, see *Unguentarius*.

PIGMENTARY, *Pigmental*—*p. Nævi*, see *Nævus*.

PIGMENTEUX, *Pigmental*.

PIGMENTUM, (*pingere*, *pingtum*, 'to paint,') Paint—*p. Indicum*, *Indigo*—*p. Nigrum*, see *Choroidæ tunica*—*p. Uracu*, *Terra Orleana*.

PIGMY, *Pygmy*.

PIGNEROLE, *Cantaurea calcitrapa*.

PIGNON DES BARBADES, *Jatropha curcas*—*p. de Barbarie*, *Jatropha curcas*—*p. d'Inde*, *Jatropha curcas*—*p. d'Inde*, *petit*, *Croton tiglium*.

PIG'NUT, *Earth'nut*, *Ground'nut*. The bulbous root of *Bunium Bulbocastanum*. *Ord. Umbelliferae*. (Called *pignuts*, from the fondness of the pig for them.) They have been considered to possess a styptic quality; and are deemed serviceable in atonic affections of the kidney. See *Bunium bulbocastanum*.

PIG'S FLARE, *Adeps suillus*.

PIKROPEGÆ, *Picropegæ*.

PIKRY, (from *πικρος*, 'bitter,') *Pulsis aloës et canellæ*.

PILA, (*πῖλος*.) *Sphæ'ra*, *Pa'lla*, *Ball*. A gymnastic exercise, much used by the ancients as well as moderns. Also, a mortar and a pestle—*Ac'ônê*.

PILA, Mortar, Pillar—*p. Damarum*, *Ægagropila*.

PILA MARI'NA, *Sphæ'ra Mari'na* seu *Thalas'sia*, *Halcyon'ium rotundum*, (*F.*) *Pelote de Mer*, *Zostère*, *Algue marine*, *A. commune* ou des *Verriers*. Certain light, round, depressed, or oblong masses—inodorous and insipid—in which rushes, hairs, and the débris of shells are met with. These are found on the sea-shores, and have been regarded by Spielmann as consisting chiefly of the roots of the *Zostè'ra Mari'na* seu *Marit'ima*, *Alga Mari'na*, (*Big tangles*, *Ware*, *Wair*, *Sea-ware*, *Sea wrack*, *Eel grass*.) *Phucagros'tis minor*. Iodine is detected in the ashes, which accounts for their success in goitre. They have, also, been given as a vermifuge.

PILA RUPICAPRARUM, *Ægagropila*.

PILAIRE, *Pileous*.

PILAR, *Pileous*.

PILARE MALUM, *Trichosis*.

PILA'TIO, (*pilus*, 'a hair,') Fracture of the cranium in which the fissure is narrow, and has the appearance of a hair—a *capillary fissure*, (*F.*) *Fente capillaire*.

PILE, (*pilus*.) Hair.

PILÆA PU'MILA, (from *pilus*, 'a hair,' *pilæus*, 'hairy,') *Richweed*, *Clearweed*, *Cookweed*; indigenous; *Order*, *Urticacæ*; flowering from July to September; has been used as a wash in poisoning by *Rhus*. It has a strong, grateful smell.

PILEOLUM, (dim of *pilæus*.) *Calotte*.

PILEOLUS, *Caul*, *Pileus*.

PILEOUS, *Pilo'us*, *Pi'lose*, *Pi'lous*, *Pi'lar*, (*F.*) *Pileux*, *Pilaire*, (from *pilus*, 'a hair,') That which relates to the hair. *Hairy*.

PILEOUS SYSTEM of Bichat is that formed by

the hair. It comprises the different species of hair of the human body.

PILES, BLEEDING, see *Hæmorrhoids*—*p.* Blind, *Cæce hæmorrhoides*, see *Hæmorrhoids*—*p.* Open, see *Hæmorrhoids*.

PILEUS, (*πίλος*, 'a hat, cap, or bonnet for the head.') *Pileolus*. A shield. A nipple-like instrument, formed of various materials, and used by females for protecting the nipple during the child's sucking, when the part is painful. Also, a caul.

PILEUS, *Cucupha*—*p.* Hippocraticus, *Bonnet d'Hippocrate*.

PILEUX, *Pileous*.

PILEWORT, *Ranunculus acris*, and *R. ficaria*.

PILI ANALES, (*pl. of pilus*.) see *Anal*—*p.* Capitis, see *Capillus*—*p.* Cutis, *Lanugo*—*p.* Palpebrarum, *Cilia*—*p.* Pudendorum, *Pubes*—*p.* Subaxillares, see *Axilla*.

PILIER, (*[L.] pila*.) *Pillar*.

PILIFEROUS, *Pilifer*, (*F.*) *Pilifère*, (*pili*, 'hairs,' and *fero*, 'I bear.') *Pile* or *hair bearing*,—as a '*piliferous cyst*,'—one containing hair.

PILIFORM, *Piliformis*, (*F.*) *Piliforme*, (from *pilus*, and *forma*, 'form.') Having the form of a hair.

PILIMICTIO, (*pili*, 'hairs,' and *mictio*, 'urination,') see *Trichiasis*.

PILL, *Pilula*—*p.* Abernethy's, see *Pilulæ hydrargyri*—*p.* Barbadoes Aloes, see *Pilulæ aloes*—*p.* of Socotrine Aloes, see *Pilulæ aloes*—*p.* of Aloes and camboge, *Pilulæ aloes et cambogiae*—*p.* of Aloes and cinchona, *Pilulæ aloes et kinæ kinæ*—*p.* of Aloes and colocynth, *Pilulæ aloes et colocynthidis*—*p.* of Aloes and fetid gums, *Pilulæ de aloes et fetidis*—*p.* of Aloes and ginger, *Pilulæ aloes et zingiberis*—*p.* of Aloes and iron, *Pilulæ aloes et ferri*—*p.* of Aloes and mastich, *Pilulæ aloes et mastiches*—*p.* Aloes and myrrh, *Pilulæ aloes et myrrhæ*—*p.* Aloetic, *Pilulæ aloetica*—*p.* Aloetic and assafetida, *Pilulæ aloes et assafetida*—*p.* Aloetic, compound, *Pilulæ aloes compositæ*—*p.* of Ammoniuiret of copper, *Pilulæ ammoniuireti cupri*—*p.* Analeptic, James's, see *Analeptica*—*p.* Anderson's, see *Pilulæ aloes et jalapæ*—*p.* Anodyne, *Pilulæ opiatæ*—*p.* Antibilious, Barclay's, *Pilulæ antibiliosæ*—*p.* Antibilious, Dixon's, *Pilulæ antibiliosæ*—*p.* Asiatic, *Pilulæ Asiaticæ*—*p.* Assafetida, *Pilulæ Assafetida*—*p.* Bacher's, *Pilulæ ex helleboro et myrrhæ*—*p.* Barbarossa's, *Barbarossæ pilulæ*—*p.* Barthez's, see *Pilulæ aloes et myrrhæ*.

PILL, BLAIR'S. An empirical preparation of colchicum, much used in England in cases of gout.

PILL, BLAUD'S. These pills, proposed by M. Blaud as antichlorotic, are formed as follows:—Take of *gun tragacanth*, in powder, six grains; *water*, one drachm. Macerate in a glass or porcelain mortar, until a thick mucilage is formed, and if it be desired to prevent the formation of peroxide of iron, and to make the pill similar to that of Vallet, substitute a drachm of powdered sugar for the mucilage. Add afterwards of *sulphate of iron*, in powder, half an ounce. Beat well until the mixture is homogeneous, and add *eubarbonate of potassa*, half an ounce. Beat until the mass becomes of a deepgreen colour, and of a soft consistence. Divide into 48 pills or boluses. Dose, one a day; gradually increasing to two, and afterwards to three.

PILL, BLUE, *Pilulæ hydrargyri*—*p.* Brandreth's, see *Pilulæ aloes et cambogiae*—*p.* Calomel, *Pilulæ hydrargyri chloridi mitis*—*p.* Calomel, compound, *Pilulæ hydrargyri submuriatis compositæ*—*p.* Calomel and opium, *Pilulæ calomelanos et opii*—*p.* Camboge, *Pilulæ cambogiae compos.*—*p.* Camboge, compound, *Pilulæ cambogiae compositæ*—*p.* of Carbonate of iron, *Pilulæ ferri carbonatis*—

p. Cathartic, compound, *Pilulæ cathartice compositæ*—*p.* Chamberlain's restorative, see Chamberlain's restorative pills—*p.* of Colocynth and henbane, *Pilulæ colocynthidis et hyoscyami*—*p.* of Colocynth, compound, *Pilulæ colocynthidis compositæ*—*p.* Common, *Pilulæ aloes et myrrhæ*—*p.* of Ammoniated copper, *Pilulæ cupri ammoniati*—*p.* of Digitalis and squill, *Pilulæ digitalis et scillæ*—*p.* Dinner, see *Pilulæ aloes et kinæ kinæ*, and *Pilulæ rhei*.

PILL, DINNER, see *Pilulæ aloes et kinæ kinæ*, and *Pilulæ aloes et mastiches*.

PILL, DINNER, of MR. MAYO, is composed of *P. Rhei. gr. iv*; *Sode Carb. gr. j.*

PILL, EVERLASTING, *Pilulæ æternæ*, (*F.*) *Pilule perpetuelle*. A pill once in vogue, and said to be composed of metallic antimony. It was believed to have the property of purging as often as it was swallowed. A lady having swallowed one of these pills, became seriously alarmed at its not passing. "Madam," said her physician, "fear not; it has already passed through a hundred patients without any difficulty."—(*Paris*.)

PILL, FEMALE, *Pilulæ de aloes et fetidis*—*p.* Ferruginous, of Vallet, see *Ferri protocarbonas*—*p.* Fothergill's, see *Pilulæ aloes et colocynthidis*—*p.* Frank's, see *Pilulæ aloes et kinæ kinæ*—*p.* Fuller's, see *Pilulæ de aloes et fetidis*—*p.* Galbanum, compound, *Pilulæ galbani compositæ*—*p.* Griffith's, *Pilulæ ferri compositæ*—*p.* Griffith's, see *Pilulæ rhei compositæ*—*p.* Gum, *Pilulæ galbani compositæ*—*p.* Harvey's, *Pilulæ aloes et colocynthidis*—*p.* of Hellebore and myrrh, *Pilulæ ex helleboro et myrrhæ*—*p.* of Hemlock, compound, *Pilulæ conii compositæ*—*p.* Holloway's, see *Pilulæ aloes et jalapæ*—*p.* Hooper's, see *Pilulæ aloes et myrrhæ*—*p.* of Iodide of iron, *Pilulæ ferri iodidi*—*p.* of Ipecacuanha and opium, *Pilulæ ipecacuanhæ et opii*—*p.* of Ipecacuanha and squill, *Pilulæ ipecacuanhæ cum scillâ*—*p.* of Sulphate of iron, *Pilulæ ferri sulphatis*—*p.* James's Analeptic, see *Analeptica*—*p.* Keyser's, see *Hydrargyrus acetatus*—*p.* Lady Crespigny's, *Pilulæ aloes et kinæ kinæ*—*p.* Lady Hesketh's, see *Pilulæ aloes et kinæ kinæ*—*p.* Lady Webster's, *Pilulæ aloes et kinæ kinæ*, and *Pilulæ aloes et mastiches*.

PILL, LARTIGUE'S, *Pilulæ Colchici et Digitalis*, (*F.*) *Pilules de Lartigue*. (*Ext. Coloc. c. 3ss*; *Ext. Sem. Colchic. alcoholic. Extract. Digital. alcoholic. ãâ gr. iss*; *f. massa in pilul. x. dividend.*) Dose, two, three, or more, in the 24 hours, in gout, until purging is induced.

This is a form that has been employed as a substitute for the *Pilules de Lartigue*, the composition of which is not known. M. Bouchardat recommends for the same purpose a combination of *Extractum colocynthidis compositum*, *Extractum colchici* and *Extractum opii*, which is almost identical with pills prescribed by Sir Henry Holland for the gout; the only difference being, that the latter substitutes *Dover's powder* for the *Extractum opii*. Prof. Wittstein affirms that their composition is *powdered colchicum*, two grains; *sugar and mucilage*, each half a grain.

PILL, OPIATE, OF LEAD, *Pilulæ plumbi opiatæ*. **PILL, LEE'S NEW LONDON**, is composed of *scammony*, *gamboge*, *calomel*, *jalap*, *aloes*, *soap*, and *syrup of buckthorn*.

PILL, LEE'S WINDHAM, consists of *gamboge*, *aloes*, *soap*, and *nitrate of potassa*.

PILL MACHINE, *Pilulier*.

PILL, MATTHEWS'S, see *Pilulæ ex helleboro et myrrhæ*—*p.* Mercurial, *Pilulæ hydrargyri*—*p.* of Mercury, *Pilulæ hydrargyri chloridi mitis*—*p.* Moffat's, *Pilulæ aloes et cambogiae*—*p.* Morrison's, *Pilulæ aloes et cambogiae*, see *Pilulæ cambogiae compositæ*—*p.* Moseley's, *Pilulæ rhei compositæ*—*p.* Night, common, *Pilulæ opiatæ*—*p.* Opiate,

Pilulæ opiatæ—p. Peter's, see *Pilulæ aloes et cambogiæ*—p. Plummer's, *Pilulæ hydrargyri submuriatis compositæ*—p. of sulphate of Quinia, *Pilulæ quiniæ sulphatis*—p. Red, *Pilulæ hydrargyri submuriatis compositæ*—p. Red, Boerhaave's, see *Hydrargyri sulphuretum rubrum*—p. of Rhubarb, *Pilulæ rhei*—p. of Rhubarb, compound, *Pilulæ rhei compositæ*—p. Rudius's, *Pilulæ aloes et colocynthidis*—p. Rufus's, *Pilulæ aloes et myrrhæ*—p. Rush's, see *Pilulæ aloes et cambogiæ*—p. Scotch, *Pilulæ de aloes et foetidis*—p. of Soap and opium, *Pilulæ saponis cum opio*—p. Speediman's, see *Pilulæ aloes et myrrhæ*—p. Squatting, *Pilulæ opiatæ*—p. Squill, compound, *Pilulæ scillæ compositæ*—p. Stahl's, see *Pilulæ de aloes et foetidis*—p. Starkey's, see *Pilulæ ex helleboro et myrrhâ*—p. Stomachic, see *Pilulæ aloes et kinæ kinæ*, and *Pilulæ rhei*—p. Storax, *Pilulæ e styrace*—p. of Strychnia, *Pilulæ strychniæ*—p. of Submuriate of Mercury, compound, *Pilulæ hydrargyri submuriatis compositæ*—p. Tanjore, *Pilulæ Asiaticæ*.

PILL TILE, see *Pilula*.

PILL, TURPENTINE, *Pilulæ de terebinthinâ*—p. Vallet's, see *Ferri protocarbonas*, and *Pilulæ ferri carbonatis*—p. Webster's, *Pilulæ aloes et kinæ kinæ*—p. Whytt's, see *Pilulæ de aloes et foetidis*.

PIL'LAR, P'ila, Colum'na, Columel'la, (F.) *Pilier*. A name given to several parts; as the pillars of the *velum palati*, *diaphragm*, &c.

PILLULA, *Pilula*.

PILNA, MINERAL WATERS OF, see *Pillna*.

PILON, *Pilum*.

PILOSELLA, (from *pilosus*,) *Hieracium pilosella*—p. Alpina, *Hieracium pilosella*.

PILOSUS, (from *pilus*, 'a hair,') *Pileosus*.

PILQUS, *Pileosus*.

PIL'ULA, pl. *Pil'ulæ, Pill'ula*, (dim. of *pila*, 'a ball,') *Coccus, Sphærid'ion, Catapot'ion, Catopotum, Sphæ'rion, Gongyl'ion, Gong'ylia, Gongylid'ium, Sphæ'rula Gale'ni*. A pill. (F.) *Pilule*. A simple or compound medicine, weighing from one to five grains, of a firm consistence and spherical form; intended to be swallowed at once, without chewing. The *Pill* differs from the *Bolus*, the latter being softer and larger.

Medicines are often given in the form of pill, on account of their disagreeable taste and odour, and for the like reason they are now sugar-coated. *Pilula* also means a *Pilular mass*, or the mass formed by the amalgamation of the different articles that enter into the composition of the pill. To form the mass into pills, it is rolled out and divided by the spatula on a *pill tile* or by a *pill machine*.

PILULÆ, see *Pilula*—p. *Æternæ*, *Pill*, everlasting—p. ex *Aloe*, *P. aloes compositæ*—p. de *Aloe et cambogiâ*, *P. aloes et cambogiæ*.

PILULÆ AL'OEÏS, *P. Aloët'icæ seu ex Aloë et Sapo'næ, Aloët'ic Pill*, (F.) *Pilules d'aloës ou aloëtiques*. (*Aloës in pulv.*; *Saponis*, sing. $\mathfrak{z}$ j; *Syr.* q. s. to form 240 pills—Ph. U. S.) Dose, gr. x to xx, in habitual costiveness.

PILULÆ AL'OEÏS BARBADEN'SIS, *Pill of Barbadoes Aloes* (Ph. B. 1864), is made of *Aloës Barbada*, in powder, $\mathfrak{z}$ ij, *Sapo. dur.*, in powder, $\mathfrak{z}$ j; *Ol. carui*, f. $\mathfrak{z}$ j; *Confect. Ros.* q. s. to form a mass.

PILULÆ AL'OEÏS COMPOSITÆ, (F.) *P. ex Aloë, Compound Aloët'ic Pills*. (*Aloës spic. extract. cont. $\mathfrak{z}$ j*; *Ext. gentiânæ*, $\mathfrak{z}$ ss; *Olei carui*, gtt. xl; *Syr.* q. s.—Ph. L.) These are sometimes, also, called *Family pills*, *Antibilious pills*, &c.

PILULÆ AL'OEÏS SOCOTRINÆ, *Pill of Socotrine Aloes*, of the same Pharmacopœia, is made of *Aloës Socotrin*, in powder, $\mathfrak{z}$ ij; *Sapo dur.* in powder, $\mathfrak{z}$ j; *Ol. myrist.* f. $\mathfrak{z}$ j; *Confect. Ros.* $\mathfrak{z}$ j.

PILULÆ AL'OEÏS ET ASSAFÔG'IDÆ, *Aloët'ic and Assafetida Pills*. (*Aloës, Assafetide, Saponis*

sing. $\mathfrak{z}$ ss; *Aquæ*, q. s. to form 180 pills—Ph. U. S.) *Purgative and stomachic*. Dose, gr. x.

PILULÆ AL'OEÏS ET CAMBOG'ICÆ, *P. de Aloë et Cambogiâ seu Hydrago'icæ Bon'tii, Pill of Aloes and Camboge*. (*Aloës, Camboag, Gum ammoniac*, $\mathfrak{a}\mathfrak{a}$ p. $\mathfrak{a}$. Make into a powder. Dissolve in vinegar; pass through a cloth; express the residuum strongly; and afterwards evaporate, in a water-bath, to a pilular consistence—Ph. P.) *Purgative*; used in dropsy. Dose, gr. xij to xxx.

Peter's Pills—a nostrum—are formed of *Aloes, Jalap, Scammony*, and *Camboge*, of each equal parts.

The basis of *Mor'rison's Pills*, often called *Hygiene Pills*, of *Brand'reth's Pills*, and *Moff't's Pills*, is said to be *aloes*, with *Camboge, Scammony*, &c.

Rush's Pills were formed of *Calomel, Aloes*, and *Camboge*, with *Oil of Mint*.

PILULÆ AL'OEÏS ET COLOCYN'THIDIS, *P. ex Colocyn'thidæ cum aloë seu Aloës cum Colocyn'thidæ seu coc'cinæ seu coc'chii seu colocyn'thidis compos'itæ seu coc'cinæ mino'res, Pill of Aloës and Colocyn'th.* (F.) *Pilules panchymagogues ou universelles, Pilules cochées*. (*Pulp. coloc. $\mathfrak{z}$ ss; Aloës hepat.*, *Scammon.* $\mathfrak{a}\mathfrak{a}$ $\mathfrak{z}$ j; *Sapon.* $\mathfrak{z}$ ij; *Ol. caryoph. $\mathfrak{z}$ j*; *Syrup.* q. s.—Ph. D.) *Cathartic*. Dose, gr. v to x or more.

Poth'ergill's Pills consist of *aloes, scammony, colocynth*, and *oxide of antimony*; and *Har'vey's Antibilious Pills*, and *Ru'dius's Pills*, (see *Extractum Rudii*), do not differ much from them.

PILULÆ AL'OEÏS ET FER'RI, *Pills of aloës and Iron*. (*Ferri sulphat.* p. iij; *Aloës Barbada.* p. ij; *Pulv. arom.* p. vj; *Conserv. rosæ*, p. viij—Ph. Ed.) In atonic constipation. Dose, one to three pills.

PILULÆ DE AL'OE ET FÆTIDIS, *P. Benedic'tæ Fulleri seu Ecphrac'ticæ seu Aloës martia'les, Pills of aloës and fet'id gums, Fe'male pills*, (F.) *Pilules bénites de Fuller*. (*Aloës socotr.*, *Sennæ, Assafetide, Galban.*, $\mathfrak{a}\mathfrak{a}$ $\mathfrak{z}$ ij; *Myrrh.* $\mathfrak{z}$ iv; *Croci, Macis*, $\mathfrak{a}\mathfrak{a}$ $\mathfrak{z}$ j; *Ferri sulphat.* $\mathfrak{z}$ iss. Reduce to powder separately, and add *Ol. succin.* gtt. viij; *Syr. artemisiz*, q. s.—Ph. P.) *Purgative*, and given in hysteria. Dose, gr. xv to xx.

Whytt's Pills, given in similar cases, consisted of *Chloride of iron, Aloes, Extract of horehound*, and *Assafetida*.

Stahl's Pills, much used as a tonic aperient in Germany, are made as follows:—*Ferri pulv.* $\mathfrak{z}$ ss; *Aloës pulv.*, $\mathfrak{z}$ ij; *Ext. colocynth.* $\mathfrak{z}$ j; *Mucilag. acaciæ* q. s. ut fiant pil xl.

PILULÆ AL'OEÏS ET JALAP'Æ, commonly called *Anderson's pills, Anderson's Scots' pills, Scotch pills*. (*Aloës Barbada.* lbj; *Rad. helleb. nigr.*, *Rad. jalapæ, Potassæ subcarb.* $\mathfrak{a}\mathfrak{a}$ $\mathfrak{z}$ j; *Ol. anisi*, $\mathfrak{z}$ iv; *Syrup. simpl.* q. s. M.) Dose, gr. x to xxx.

A committee of the Philadelphia College of Pharmacy recommend the following: *Aloës Barbada.* $\mathfrak{z}$ xxiv; *Saponis.* $\mathfrak{z}$ iv; *Colocyn'thid.* $\mathfrak{z}$ j; *Cambogie*, $\mathfrak{z}$ j; *Ol. anis.* f. $\mathfrak{z}$ ss.

Hollownay's Pills are said to be composed as follows: *Aloës* gr. lxij; *Rhei pulv.* gr. xxvj; *Piperis*, gr. viij; *Croci, Sodæ sulphat.* $\mathfrak{a}\mathfrak{a}$ gr. iij, M. et f. pil. exliv.

PILULÆ AL'OEÏS ET KINÆ KINÆ, *P. stomachicæ seu ante ci'bum, Pill of aloës and cincho'na*, (F.) *Pilules gourmandes*. (*Aloës socotr.* $\mathfrak{z}$ vj; *Ext. cinchon.* $\mathfrak{z}$ ij; *Cinnam.* $\mathfrak{z}$ j; *Syrup. absinth.* q. s.—Ph. P.) *Stomachic and laxative*. Dose, gr. x to xx.

La'dy Crespign'y's pills, La'dy Web'ster's pills, La'dy Hesk'eth's pills; the *Pilules de Duchesne* (*Querquetanus*), the *Grains de Vie* or *Pilules de Clérambourg*; the *Pilules Indiennes stomachiques de De la Croix*, the *Pilules de Barbier*, and the *Pilulæ stomachicæ Mesuës*, (Ph. L. 1835,) resemble these. See *Pilulæ Aloës et Mastiches*.

Frank's Pills, much employed, also, as 'dinner

pills, or,—to use the language of the eccentric Kitchener,—as ‘*peptic persuaders*,’ are said to be much used, under the name *Graines de santé*, in various parts of Europe. They consist of *Aloes*, *Jalap*, and *Rhubarb*, with *Syrup of wormwood* as an excipient.

PILULÆ ALOES MARTIALES, P. de aloes et fœtidis.

PILULÆ ALOËS ET MAS'TICHES, *Pills of Aloes and mastich*. (*Aloës socotrin*, in pulv. ʒiiss; *Mustich*, pulv. Ros. Gallie. aa ʒss, M. et divide in pil. cccc.—Ph. U. S. 1864.) The mastich is added to the aloes to prevent the too ready solution of the latter in the gastric and intestinal fluids; so that it is long in operating as a laxative. One is usually taken before dinner as a ‘dinner pill.’ It has been introduced into the Ph. U. S., 1864, as a form for *Lady Webster's pills*.

PILULÆ ALOËS ET MYRRHÆ, P. aloës cum myrrhâ seu de aloë et myrrhâ, P. Ru'fî seu Rufî seu commun'es, Ru'fus' or Ru'f'us' pills. *Common pills, Pills of aloes and myrrh*. (*Aloës*, ʒij; *Croci*, in pulv. ʒss; *Myrrh*, ʒj; *Syr*. q. s. to form 480 pills.—Ph. U. S.) Cathartic and emmenagogue. Dose, gr. x to xx.

Hoop'er's Pills—possessed of similar properties—according to one of Gray's formulæ, are made as follows: *Ferri sulph.*, *Aque*, aa ʒviij. Dissolve. Add *Aloes Barbado.* lbij and ʒviij; *Cannelæ albæ*, ʒvj; *Gum. myrrh.* ʒij; *Opoponacis*, ʒiv.

The committee of the Philadelphia College of Pharmacy recommend the following form: *Aloës Barbadosis*, ʒviij; *Ferri sulphatis exsiccata*, ʒij, and ʒiiss, vel *Ferri sulph. crystall.* ʒiv; *Ext. helleb.* nigr. ʒij; *Myrrh.* ʒij; *Saponis*, ʒij; *Cannelæ*, in pulv. tritæ, ʒj; *Zingib.* in pulv. trit. ʒj.

Speediman's Pills consist of *aloës*, *myrrh*, *rhubarb*, extract of chamomile, and essential oil of chamomile; and the *Pills of Barthez*, of *aloës*, *myrrh*, *musk*, *camphor*, and *balsam of Peru*.

PILULÆ DE ALOE ET SAPONE, P. aloeticæ.

PILULÆ ALOËS CUM ZINGIB'ERÆ, *Pills of aloes and ginger*. (*Aloës hepat.* ʒj; *Rad. zingib.* in pulv. ʒj; *Sapon. hispan.* ʒss; *Ol. ess. menthæ pip.* ʒss.—Ph. D.) In habitual costiveness. Dose, gr. x to xx.

PILULÆ AMMONIURETI CU'PRI, P. Cu'pri, *Pills of Ammoniu'ret of copper*, (F.) *Pilules d'ammonure de cuivre*. (*Ammoniu'r. cupri*, in pulv. ten. trit. gr. xvij; *Miccæ panis*, ʒiv; *Aque carbonatis ammoniæ*, q. s. Beat into a mass, and divide into xxxij pills.—Ph. E.) Tonic and antispasmodic, in epilepsy, &c. Dose, one pill twice daily; gradually increasing the dose.

PILULÆ ANTE CIBUM, ‘pills before-food,’ P. aloes et kinæ kinæ.

PILULÆ ANTIBILIO'SÆ, *Bar'clay's Antibilious pills*. (Extract of colocynth, ʒij; Resin of jalap, ʒj; Almond soap, ʒiiss; Guaiac, ʒij; Emetic tart. gr. viij; Oils of juniper, caraway, and rosemary, aa iv drops; Syrup of buckthorn, q. s. to form 64 pills.) Dose, two or three.

DIC'ON'S Antibilious pills are formed of *Aloes*, *Scammony*, *Rhubarb*, and *Tartarized antimony*. **PILULÆ ANTIBERPETICÆ**, *Pilulæ hydrargyri submuriatis comp.*—p. Antibystericæ, P. galbani composita.

PILULÆ ANTISCROPHULO'SÆ, P. ex ox'ido stib'ii et hydrargyri sulphure'to nig'ro compos'itæ. (P. scammon., Hydrarg. sulphur. nigr. aa ʒij; Antimonii acidi albi, Milleped., Sapon. amygdalin. aa ʒij; Extract. glycyrrh. ʒv. Make into a mass.—Ph. P.) Dose, gr. iv to xx.

PILULÆ ANTISPASTICÆ, P. galbani composita.

PILULÆ ASIAT'ICÆ, *Asiat'ic or Tonjore's pills*. (*Acid. arsenios.* gr. lv; *Piper. nigr.* pulv. ʒix; *Acacie*, q. s. ut fiant pil. 800.) Each pill contains about one-fiftieth of a grain of arsenious acid. These pills are employed in India for the cure of syphilis and elephantiasis

PILULÆ ASSAFETIDÆ, *Assafet'ida pills*. (*Assafetid.* ʒiiss; *Saponis*, ʒss. Beat with water so as to form a mass to be divided into 240 pills.—Ph. U. S.)

PILULÆ ASSAFETIDÆ COMPOSITÆ, P. galbani composita.

PILULÆ BALSAM'ICÆ, P. Docto'ria Mor'ton. (*Milleped.* pulverisat. ʒxviij; *Gum ammon.* ʒxj; *Acid. benzoic.* ʒvj; *Croci.* *Balsami Peruæ*. aa ʒj; *Balsam. sulph. anisat.* ʒvj. Mix.—Ph. P.) Tonic, expectorant, and balsamic.

PILULÆ BARBAROSSÆ, see *Barbarossæ pilulæ*—p. *Benedictæ Fuller*, P. de aloes et fœtidis.

PILULÆ CALOMELANOS COMPOSITÆ, P. Hydrargyri submuriatis composita.

PILULÆ CALOMEL'ANOS ET O'PII, P. Hydrargyri chloridi et o'pii, *Pills of Calomel and Opium*. (*Hydrarg. chlorid.* mit. p. iij; *Opii*, p. j; *Confect. Ros.* q. s. To be divided into pills each containing two grains of calomel.—Ph. Ed.)

PILULÆ CAMBOGIÆ COMPOSITÆ, P. de gut'tâ gaman'drâ seu hydrago'gæ seu gum'mi gut'tæ aloet'icæ, *Gamboge's pills*, *Com'pound gamboge's pills*, (F.) *Pilules de gomme gutte composées*. (*Cambogiz contrite*, *Aloes epicat. extract.* contriti, *Pulv. cinnam.* comp. sing. ʒi; *Saponis*, ʒij.—Ph. L.) Cathartic. Dose, gr. x to xx.

Mor'rixon's Pills are said by Dr. Paris to be nothing more than the above, with the addition of a small portion of Cream of tartar.

PILULÆ CATHARTICÆ COMPOSITÆ, *Com'pound cathartic pills*. (*Ext. colocynth.* comp. in pulv. ʒss; *Ext. jalapæ*, in pulv., *Hydrarg. chlorid.* mit. aa ʒij; *Gambog.* in pulv. ʒij. M. Form into a mass with water, and make into 180 pills.—Ph. U. S.) Three for a dose.

PILULÆ COCHII, P. aloes et colocynthidis—p. *Coccia minores*, P. aloes et colocynthidis—p. *Coccinæ*, P. aloes et colocynthidis—p. *Colehici et digitalis*, P. Lartigue's—p. *Colocynthidis composita*, P. aloes et colocynthidis—p. ex *Colocynthide cum aloes*, P. aloes et colocynthidis.

PILULÆ COLOCYNTH'IDIS ET HYOSC'YAMI, *Pills of Colocynth and Hen'bane*. (*Pil. Colocynth. comp.* p. ij; *Ext. Hyoseyam.* p. i.—Ph. E.) Dose, gr. v to gr. xx.

A much more elaborate formula is contained in the Ph. B. (1864.) (*Colocynth*, in powder, ʒj; *Barbadoes Aloes*, in powder, ʒij; *Scammony*, in powder, ʒij; *Sulphate of potassa*, in powder, ʒi; *Oil of cloves*, f ʒij; *Extract of Hyoseyamus*, ʒij; *Distilled water*, q. s. Beat into a mass.)

PILULÆ COMMUNES, P. aloes et myrrhæ.

PILULÆ CONT' COMPOSITÆ, *Com'pound Pill of Hem'lock*. (*Ext. Conti* ʒv; *Ipecac. pulv.* ʒj; *The'riac.* q. s.—Ph. L.) Dose, gr. v three times a day.

PILULÆ COPAI'BÆ, *Pills of Copai'ba*. (*Copaib.* ʒij; *Magnesia*, recenter præpar. ʒj. Mix, and set aside till it concretes into a pilular mass, to be divided into 200 pills.—Ph. U. S.) Each pill contains nearly five grains of copaiba. Two to six for a dose.

PILULÆ CUPRI, P. ammoniureti cupri.

PILULÆ CUPRI AMMONI'ATI, *Pills of ammoniated Cop'per*. (*Cupri ammon.* in pulv. p. j; *Miccæ panis*, p. vj; *Liq. ammon. carbonat.* q. s.—Ph. E.) Dose, one to six pills.

PILULÆ DE CYNOGLOSSO, P. de extrac'to o'pii. (*Pulv. rad. cynoglossæ*, *Semin. hyoseyam. alb.*, *Extract. opii vinos.* aa ʒiv; *Pulv. myrrh.* ʒvj; *Oliban.* ʒv; *Croci.* *Castor.* aa ʒiiss; *Syrup opii*, q. s.—Ph. P.) Anodyne. In cough, &c. Dose, gr. iv to gr. viij.

PILULÆ DEPURANTES, ‘depurative pills,’ P. hydrargyri submuriatis composita.

PILULÆ DIGITALIS ET SCILLÆ, *Pills of Digitalis and Scillæ*. (*Digital.* *Scillæ*, aa p. j; *Elect. aromat.* p. ij.—Ph. E.) One or two pills for a dose.

PILULÆ ECPHRACTICÆ. P. de aloë et fetidiss.

PILULÆ FER'R'I CARBONAT'IS, *Pills of ferrous carbonate of iron, Valler's pills, Valler's ferruginous pills.* (Ferri sulph. $\frac{3}{4}$ ij; Sodæ carbonat. $\frac{3}{4}$ ix; Mek. depurcat. $\frac{3}{4}$ ij; Sacchar. $\frac{3}{4}$ ij; Aquæ bullientis, Oij; Syrup. q. s. Dissolve the sulphate of iron and carbonate of soda, each in a pint of the water, and to each solution add a fluidounce of syrup; then mix the two solutions in a bottle just large enough to contain them; close it accurately with a stopper, and set it by, that the carbonate of iron may subside. Pour off the liquid, and having washed the precipitate with warm water, sweetened with syrup in the proportion of f $\frac{3}{4}$ ij of the latter to Oj of the former, until the washings no longer have a saline taste, place it upon a flannel cloth, and express as much water as possible; then mix it with the honey and sugar. Lastly, heat the mixture in a water-bath, until it attains a pilular consistence.—Ph. U. S. 1863.) Dose, gr. v to xv. or more.

The formula in the Ph. B. (1864) is—*Saccharatæ Carbonatæ of iron, $\frac{3}{4}$ ij; Confection of Roses, $\frac{3}{4}$ ij. Beat into a mass.*

PILULÆ FER'R'I COMPOS'ITÆ, *P. ferri cum myrrhâ, Compound pills of iron, Griffith's Pills, (F.) Pilulæ de fer compositis.* (Myrrh. in pulv. $\frac{3}{4}$ ij; Sodæ carb. Ferri sulph. $\frac{3}{4}$ ij; Syrup. q. s. Mix, to form 80 pills.—Ph. U. S.) Tonic. Dose, gr. x to xx.

PILULÆ FER'R'I IO'DIDI, *Pills of Iodide of iron.* (Ferri sulphat. $\frac{3}{4}$ ij; Potass. iodid. $\frac{3}{4}$ iv; Tragacanth. pulv. gr. x; Sacchar. pulv. $\frac{3}{4}$ ss. Beat with syrup into a mass, to form 40 pills.—Ph. U. S. 1851.) A modified form is given in the Ph. U. S. for 1863. Dose, one to three where a tonic and euphoric is indicated.

PILULÆ FERRI CUM MYRRHÂ, *P. Ferri composita.*

PILULÆ FERRI SULPHAT'IS, *Pills of Sulphate of Iron.* (Ferri sulph. exsic. p. ij; Ext. Taraxac. p. v; Conserv. Rosæ, p. ij; Glycyrrhiz. pulv. p. iij.—Ph. E.) Dose, gr. v to gr. xx.

PILULÆ FETIDÆ, *P. galbani composita.*

PILULÆ GAL'BANI COMPOS'ITÆ, *P. asafœtida composita seu fa'tida seu gummo'sa seu gum'mi seu antispas'tica seu antihyster'ica seu myrrhæ composita, Compound galbanum pill, Gum pill.* (Galban., Myrrhæ, $\frac{3}{4}$ ij; Asafœtida, $\frac{3}{4}$ ij; Syrup. q. s. to form 240 pills.—Ph. U. S.) Antispasmodic and emmenagogue. Dose, gr. x to xx.

PILULÆ GUMMI, *P. galbani composita*—p. Gummi guttæ aloeticæ, *P. cambogiæ composita*—p. Gummosæ, *P. galbani composita*—p. de Guttâ gamandrá, *P. cambogiæ composita.*

PILULÆ EX HELLEBORO ET MYRRHÂ, *P. tonica, Ba'cher's pills, Tonic pills of Ba'cher, Pills of hell'ebore and myrrh.* (Ext. helleb. nig., Myrrh. $\frac{3}{4}$ ij; Fol. cardui benedicti in pulv. $\frac{3}{4}$ ij. M.—Ph. P.) Tonic and emmenagogue. In dropsical, cutaneous, and verminous affections. Dose, gr. iv to gr. viij.

Matthews's pills, Star'key's pills, are formed of Rad. helleb. nigr., Rad. glycyrrh., Sapon. Castill., Rad. curcumæ, Opii purif., Syrup. croci, $\frac{3}{4}$ ij; Ol. tereb. q. s. ut fiat massa.—(Gray.)

PILULÆ HYDRAGOGÆ, *P. cambogiæ composita*—p. Hydragogæ Bontii, *Pilulæ aloës et cambogiæ.*

PILULÆ HYDRAR'GYRI, *P. mercurial'les, Mercu'rial or Blue pill, Blue mass, Pills of Mer'cury, (F.) Pilulæ de Mercurio.* (Hydrag. purif. $\frac{3}{4}$ ij; Conf. rosæ, $\frac{3}{4}$ ss; Glycyrrh. pulv. $\frac{3}{4}$ ss. Rub the quicksilver with the confection, until the globules disappear; then add the liquorice powder, and beat the whole into a uniform mass, to be divided into 480 pills.—Ph. U. S.) Antisiphilitic; mercurial;—in large doses, purgative.

A purgative pill, much prescribed by Mr. Aber-

nethy, and called *Ab'ernethy's pill*, consisted of Pil. hydrarg. gr. x; Jalap. pulv. gr. xx; Syrup. rhamni, q. s. ut fiant pil. vj. Two were given at night, and the next morning a wineglassful of infusion sennæ.

PILULÆ HYDRARGYRI CHLORIDI ET OPII, *P. Calomelanos et Opii.*

PILULÆ HYDRAR'GYRI CHLO'RIDI MI'TIS, *Pills of mild chlor'ide of mer'cury, Cal'omel pills.* (Hydrag. chlorid. mit. $\frac{3}{4}$ ss; Acaciæ, in pulv. $\frac{3}{4}$ ij; Syrup. q. s. ut fiant pil. 240.—Ph. U. S.) Each pill contains one grain.

PILULÆ HYDRAR'GYRI SUBMURIA'TIS COMPOS'ITÆ, *Pilulæ Antimo'nii Compositæ, (Ph. U. S. 1863.) P. ex stib'io et hydrar'gyro seu antihyper'ticæ seu ex hydrargyro murial'ico mi'ti cum sulphure stib'io auriant'ico seu resolut'es cum mercurio seu depuran'tes seu hydrar'gyri submuriat'is, Pilulæ Hydrar'gyri chlo'ridi composita, Pilulæ calomel'anos composita, Compound Cal'omel pill, Compound pills of submuriate of mer'cury, Cal'omel pills, Plum'mer's pills, Red pill.* (Hydrag. submur., Antimon. sulphur. precipitat. sing. $\frac{3}{4}$ ij; guaiac. resin. pulv., Syrup. fusc. $\frac{3}{4}$ ss. Beat together, and divide into cexl pills. Mercurial, and reputed to be diaphoretic and alterative. Dose, gr. v to x.

Leake's pil'ula saluta'ria resembles these. It is used in syphilis.

PILULÆ HYDRARGYRI SUBMURIATIS ET OPII, *P. Calomelanos et Opii.*

PILULÆ DE HYDRAR'GYRO, SCAMMONIO ET AL'OË, vulgo MERCURIA'LES, (F.) *Pilulæ Mercurielles ou de Mercure, Scammonée et Aloës.* (Hydrag. purif. $\frac{3}{4}$ ij; Mellis, $\frac{3}{4}$ xj. Rub until the globules disappear; then add Aloës Socotr., Scammon. $\frac{3}{4}$ ij; Maciæ, Cinnam. $\frac{3}{4}$ ij. Make into a mass.—Ph. P.) Uses, same as the last, but they are more active. Dose, gr. viij to xxx. These pills are nearly the same as the *Pastilles de Renaudot*.

PILULÆ EX HYDRARGYRO MURIATICO MITI CUM SULPHURE STIBIATO AURANTIACO, *Pilulæ hydrargyri submuriatis composita.*

PILULÆ IPECACUAN'HÆ ET O'PII, *Pills of Ipecacuan'ha and Opium.* (Pulv. Ipecac. et Opii, p. iij; Conserv. Rosæ, p. j.—Ph. E.) One grain of opium in about thirteen grains.

PILULÆ IPECACUAN'HÆ CUM SCILLÂ, *Pills of Ipecacuan'ha with Squill.* (Pulv. Ipecac. comp. $\frac{3}{4}$ ij; Scillæ pulv., Ammoniac. in pulv. $\frac{3}{4}$ ij; Theriac. q. s.—Ph. L.) Dose, gr. v to x.

PILULÆ MERCURIALES, *P. hydrargyri, P. de hydrargyro, scammonio et aloë*—p. Doctoris Morton, *P. balsamicæ*—p. Myrrhæ composita, *P. Galbani composita.*

PILULÆ OPIATÆ, *Pilulæ Opii* (Ph. U. S.). *P. Thebat'icæ, Opiate pills, Con'mon night pills, Anodyne pills, (Prov.) Squatting pills, Nepen'thes opiatum, (F.) Pilulæ opiacées.* (Opii pulv. $\frac{3}{4}$ ij; Sapon. gr. xij; to be made into 60 pills.—Ph. U. S.) Each pill contains a grain of opium.

PILULÆ DE EXTRACTO OPII, *P. de cynoglossa*—p. ex Oxido stibii et hydrargyri sulphureto nigro composita, *Pilulæ antiscrophulosæ.*

PILULÆ PLUMBI OPIATÆ, *Pilula Plumbi cum Opio, (Ph. B. 1864.) Opiate Pills of Lead, Pill of Lead and Opium.* (Plumb. acetat., p. vj; Opii, p. j; Conserv. Rosæ, p. j. To be made into four grain pills.—Ph. E.) Each pill contains three grains of acetate of lead.

PILULÆ QU'INIE SULPHAT'IS, *Pills of sul'phate of qu'inia.* (Quin. sulph. $\frac{3}{4}$ ij; Acaciæ, in pulv. $\frac{3}{4}$ ij; Mel. q. s. ut fiant pilulæ 480.—Ph. U. S.) Each pill contains a grain of the sulphate of quina.

PILULÆ RESOLVENTES CUM MERCURIO, *P. hydrargyri submuriatis composita*—p. de Khababar, *Pilulæ rhei composita.*

The stomachic nostrum, sold under the name

of *Moseley's pills*, consists merely of *rhubarb* and *ginger*.

PILULÆ RHEI, *Pills of Rhubarb*. (*Rhei* pulv. ʒvj; *Saponis*, ʒij. M. et fiant pil. 120.—Ph. U. S.) Each pill contains three grains of rhubarb.

A good *Dinner* or *Stomachic* pill is made of *Rhei* pulv. ʒij; *Capsic*. pulv. ʒi; make into a mass with molasses, and divide into 60 pills. Dose, two or three before dinner.

PILULÆ RHEI COMPOSITÆ, *P. de Rhubarbaro, Compound rhubarb pills*. (*Rhei* pulv. ʒij; *Aloes*, pulv. ʒvj; *Myrrh*. pulv. ʒiv; *Ol. menthæ pip.* f ʒss. Beat into a mass, with *Syrup of orange-peel*, to form 240 pills.—Ph. U. S.) Laxative and stomachic. Dose, gr. x to xx.

PILULÆ RHEI ET FERRI, *Pills of Rhubarb and Iron*. (*Ferri sulph.* exsiccet. p. iv; *Ext. Rhei*, p. x; *Conserv. Rosæ* p. v. Make into five grain pills.) Dose, as a tonic and laxative, two or three pills.

Griffitt's pills, so called after Dr. S. P. Griffitts, of Philadelphia, who prescribed them as a tonic aperient, were formed as follows:—*Rhei* pulv. ʒss; *Ferri sulph.* ʒss; *Saponis*, ʒij; *Aque*, q. s. ut fiat massa in pill. xl dividenda. Dose, three or four at bedtime.

PILULÆ RUFÆ, *P. aloes et myrrhæ*—p. Salutarie, Leake's, see *Pilulæ hydrargyri submuriatis compositæ*—p. *Saponaceæ*, *P. saponis cum opio*—p. *Saponis compositæ*, *P. saponis cum opio*.

PILULÆ SAPONIS CUM OPIO, *P. Saponaceæ seu saponis compositæ*, *Pills of soap and opium*, (F.) *Pilules de savon et d'opium*. (*Opii*, ʒj; *Sapon.* ʒss. M.—Ph. U. S.) Four grains contain one of opium.

The Pharmacopœia of the United States has a form for the *Pilulæ opii*, *Opium pills*, as well as for the *Pilulæ saponis compositæ*; the only difference between them being in the proportion of the ingredients. See *Pilulæ opiatæ*.

PILULÆ SCILLÆ, *P. Scillæ compositæ*.

PILULÆ SCILLÆ COMPOSITÆ, *P. scillitice seu scillæ cum zingibere seu scillitæ, Compound squill pill or Compound Pills of Squill*. (*Scillæ* pulv. ʒj; *Zingib.* pulv., *Ammoniac.* pulv. aa ʒij; *Saponis*, ʒij; *Syrup.* q. s. ut fiant pil. 120.—Ph. U. S.) Dose, two pills. Expectorant and diuretic. Dose, gr. x to xx.

PILULÆ SCILLÆ CUM ZINGIBERE, *P. scillæ compositæ*—p. *Scillitice*, *P. scillæ compositæ*—p. *ex Stibio et hydrargyro*, *P. hydrargyri submur. compos.*—p. *Stomachicæ*, *P. aloes et kinæ kinæ*.

PILULÆ STRYCHNIE, *Pills of Strychnia*. (Very pure *Strychnia*, 2 gr.; *Conserve of roses*, ʒss. Make into 24 pills.) One or two for a dose.

PILULÆ STYRACIS, *P. S. Compositæ, Storax Pills*. (*Styracis pur.* ʒvi; *Opii*, *Croc.*, sing. ʒij.—Ph. E.) Five grains contain one of opium.

PILULÆ SUBLINGUALES, *Hypoglossitides*.

PILULÆ DE TEREBINTHINÆ, *Turpentine Pills*. Made of turpentine boiled in 3 times its weight of water, until, when thrown into cold water, it forms a soft paste, which has the properties of turpentine, and has been administered in gleet, &c.

PILULÆ THEBAICÆ, *P. opiatæ*—p. *Tonicæ*, *P. ex Helleboro et myrrhâ*.

PILULAR, Pilularis, (F.) *Pilulaire*, from *pilula*.) Relating or appertaining to pills,—as 'a pilular mass,' 'pilular form,' &c.

PILULES, *Pilulæ*—p. *Aggregatæ*, see *Aggregate*—p. *d'Aloes ou aloétiques*, *Pilulæ aloeticæ*—p. *d'Ammoniac de cuire*, *Pilulæ ammoniacreti cupri*—p. *de Barbier*, see *Pilulæ Aloes et kinæ kinæ*—p. *Bénites de Fuller*, *Pilulæ de aloes et foetidis*—p. *Cochées*, *Pilulæ aloes et colocynthidis*—p. *de Duchesne*, see *Pilulæ Aloes et kinæ kinæ*—p. *de Ferri compositæ*, *Pilulæ ferri compositæ*—p. *Ferrugineusæ de Vailet*, see *Ferri protocarbo-nas*—p. *Gourmandes*, *Pilulæ aloes et kinæ kinæ*

—p. *Indiennes stomachiques de Delacroix*, see *Pilulæ aloes et kinæ kinæ*—p. *de Lartigue*, *Pill.* *Lartigue*—p. *de Mercure*, *Pilulæ hydrargyri*—p. *de Mercure*, &c., *Pilulæ de hydrargyro*, et *scammonio et aloes*—p. *Mercurielles*, *Pilulæ de hydrargyro*, *scammonio et aloes*—p. *Opiacées*, *Pilulæ opiatæ*—p. *Panchymagogues*, *Pilulæ aloes et colocynthidis*—p. *Perpetuelles*, *Pill.* *everlasting*—p. *de Saron et d'opium*, *Pilulæ saponis cum opio*—p. *Universelles*, *Pilulæ aloes et colocynthidis*.

PILULIER (F.), *Pill Machine*. An instrument for rolling and dividing pills. Also the earthen pot in which pills are preserved.

PILUM, Pistillum, Pistillum, Hyperos, Cercis, (F.) *Pilon*. A pestle. A wooden, iron, bell-metal, marble, agate, porcelain, or glass instrument for beating or dividing substances in a mortar.

A *spring pestle* is one attached above to a spring, so as to take off the weight, and thus lessen the labour of pounding.

PILUS, *Capillus, Caul, Hair*.

PIMELE, (πιδελν.) *Fat*.

PIMELECHYSIS, (*pimele*, and *εχysis*, 'a pouring out,') *Pimelorrhœa*.

PIMELICUS, (from *pimele*), *Fatty*.

PIMELITIS, (*pimele*, and *itis*.) *Inflammation of the adipous tissue*.

PIMELODES, (πιδελος,) *Fatty*.

PIMELOMA, gen. *Pimelomatia*, (*pimele*, and *oma*.) A fatty swelling.

PIMELORRHÆA, *Pimelechysis*, (*pimele*, and *ρρ*, 'I flow.') A morbid discharge or disappearance of fat.

PIMELOTHOPNÆA, *Piorthopnæa*, (*pimele*, *opnos*, 'upright,' and *πναιν*, 'to breathe.') *Dyspnœa*, practicable only in the erect attitude, owing to fat.

PIMELO'SIS, *Obesitas nimia*, (from *pimele*, and *osis*.) Conversion into, or formation of, fat,—as *Pimelosis hepatica*, 'fatty degeneration of the liver.'

PIMELOSIS HEPATICA, *Adiposis hepatica*, see *Pimelosis*—p. *Nimia*, see *Polysarcia*.

PIMÉLOTIQUE, *Corpulent*.

PIMELURIA, *Adiposuria*, (*pimele*, and *ουρον*, 'urine,') *Fatty urine*.

PIMENT, *Capsicum annum*—p. *d'Eau*, *Polygonum hydropiper*—p. *Royal*, *Myrica gale*.

PIMENTA, see *Myrtus pimenta*.

PIMENTÆ BACCÆ, see *Myrtus pimenta*.

PIMENTO, ([S.] *pimienta*, 'pepper,') see *Myrtus pimenta*.

PIMIACULA, *Lips of the Vulva*.

PIMPINEL BLUE, (see *Pimpinel*.) *Scutellaria lateriflora*—p. *Red*, *Anagallis*—p. *Scarlet*, *Anagallis arvensis*—p. *Water*, *Veronica beccabunga*.

PIMPILIM, *Piper longum*.

PIMPINEL, (*bipinnella* or *bipennula*, from the double pennate order of its leaves,) see *Pimpinnella*.

PIMPINEL ITALIAN, *Sanguisorba officinalis*.

PIMPINELLA ALBA, *P. saxifraga*—p. *Angelicefolia*, *Ligusticum podagraria*.

PIMPINEL'LA ANI'SUM, *Anise plant*, *Ani'sum*, *A. officinalis seu vulgare*, *Sison ani'sum*, *A'pium ani'sum*, *Sola'nien intestino'rum*, *Ani'sum*, (F.) *Anis*, *Boucage anis*. *Ord.* *Umbelliferae*. *Sec. Syst.* *Pentandria Digynia*. A native of Egypt. The seeds, *Ani'sum*, (Ph. U. S.) *Semina Ani'si*, *Aniseed*, (Old Eng.) *Aneys*, have an aromatic odour; and a sweetish, warm, grateful taste. They are carminative, and used in dyspepsia and in tormina of infants. Dose, gr. xv to ʒj, bruised. The oil.—*O'LEUM ANI'SI*, (F.) *Huile d'anis*—is official. Dose, gtt. v to xv. Most of the oil used in this country under the name of

Oil of Anise is said to be obtained from *Illic'ium Anisat'um*, in China.

Aqua anisi, *Anise wa'ter*, (Ph. D.), is made by mixing a fluidounce of *essence of anise*, with half a gallon (Imp. meas.) of *distilled water*, and filtering through paper.

PIMPINELLA, GREATER, *P. magna*—*p. Hircina*, *P. Saxifraga*—*p. Laterifolia*, *Sion ammi*.

PIMPINELLA MAG'NA seu *nigra* seu *ma'jor* seu *rubra*; *Great'er Pimpinell'a*; *Tragosel'num mag'num* seu *ma'jus*, *Dan'cus*, *Cyan'opus*, (F.) *Boncuje majeur*. The root has been extolled as diuretic and resolvent, as well as in erysipellatous ulceration, tinea capitis, rheumatism, &c.

PIMPINELLA MAJOR, *P. magna*—*p. Nigra*, *P. magna*—*p. Nostras*, *P. saxifraga*—*p. Officinalis*, *Sanguisorba officinalis*—*p. Rubra*, *P. magna*.

PIMPINELLA SAXIFRAGA, *Saxifraga*, *Bur'net saxifrage*, *Poter'ium sanguisor'ba*, *Sorbastrel'la*, *Tragosel'num*, *T. saxif'raga*, *Pimpinell'a hirci'na* seu *umbellif'era* seu *al'ba* seu *no'tras*, (F.) *Boncuje mineur*, *Petit B.*, *Pimpinelle*, *Rompt-pierre*. The root has an unpleasant smell, and hot, pungent, bitterish taste. It has been recommended as a stomachic; and as a stimulating gargle in paralysis of the tongue.

PIMPINELLA UMBELLIFERA, *P. saxifraga*.

PIMPLE, ([Sax.] *pimpel*.) *Papula*.

PIMPLED, *Papulose*.

PIMPRENELLE, *Pimpinella saxifraga*—*p. Noire*, *Sanguisorba officinalis*.

PIN. (Perhaps from *pennum*, low Latin: or from *spina* [?], or from Gael. *pin*.) *Sp'ina fer'rea*, *Sp'ina'la*, *A'cus capita'ta*, (F.) *Épingle*. An iron or brass instrument, generally of a small size, pointed at one extremity, and having a head at the other. It is used, in surgery, to fix rollers and dressings; and occasionally in sutures.

PIN, *Pinus*—*p. à Pignon*, see *Pinus pinea*.

PIN AND WEB, *Caligo*, *Pterygion*.

PINASTELLUM, (from *pinus*.) *Peucedanum*.

PINASTER, (from *pinus*.) *Pinus sylvestris*.

PINBONE, *Ischion*.

PINCE, (from *pincer*, 'to pinch.') Forceps—*p. à Anneau*, see Forceps—*p. à Dissection*, see Forceps—*p. à Ligature*, see Forceps—*p. Lithodras-sique*, see *Lithodras-sique*—*p. de Museux*, see Forceps—*p. à Pansement*, see Forceps.

PINCÉ, *Pinched*.

PINCÉE, *Pugillus*.

PINCERS, Forceps.

PINCETTE, Forceps.

PINCHED, *Contractus*, *Retrac'tus*, (F.) *Pincé*, *Grippé*. An epithet applied to the face, when the features are contracted or shrunken; as in violent abdominal affections, or during intense pain.

PICKNEYA PU'BENS, *P. Pubes'cens*, *Geor'gia Bark*, *Bitter Bark*, *Flor'ida Bark*, *Fel'ver tree*. This bark has been used in Georgia in the cure of intermittents, and successfully. It is a powerful bitter. Dose of the powder, $\mathfrak{z}$.

PINDARS, *Arachis Hypogea*.

PINE, *Pinus*.

PINE APPLE, *Bromelia ananas*—*p. Apherousli*, *Pinus cembra*—*p. Ground*, *Lycopodium complanatum*—*p. Ground*, stinking, *Camphorosma Monspeliaca*—*p. Loblolly*, *Pinus taeda*—*p. Mountain*, see *Pinus mughos*—*p. Mugho*, see *Pinus mughos*—*p. Prince's*, *Pyrola umbellata*—*p. Sap*, American, *Hypopitys lanuginosa*—*p. Stone*, *Pinus pinea*—*p. Sugar*, see *Arrow Root*—*p. Yellow*, *Pinus palustris*.

PINEA, *Pinus Pinea*.

PINEAL or PINE'AL, *Pinea'lis*, (F.) *Pinéal*, (*pine'a*, 'a pine cone.') That which resembles a pine cone.

PINEAL GLAND, *Glan'dula pinea'lis*, *Cer'eb'ral epiph'ysis*, *Cona'drium*, *Cono'id'es cor'pus*, *Cor'pus*

turbina'tum, *Pe'nis* seu *Vir'ga* seu *Tur'bo cer'eb'ri*, is a small body, of conical shape; pale red or grayish colour; and soft consistence; situate between the fornix and the tubercula quadrigemina. It almost always contains sabulous particles, *Sabulum cona'drii*, *Brain Sand*, *Pinéal Sand*. When these are grouped together over the base of the gland, they form the *Acer'vulus Cer'ebri* of Sömmerring, *Acer'culi* seu *Lapilli Glan'dula pinea'lis*. From its anterior parts arise two medullary striæ, which proceed over the posterior commissure; coast along the optic thalami, and unite at the anterior pillar of the fornix,—the *Habe'næ*, *Haben'ulae* or *reins of the pineal gland*—*Pedunculi cona'drii*, see *Peduncles of the Pineal Gland*.

The uses of the pineal gland are unknown. Descartes supposed it to be the seat of the soul!

PINEAL SAND, see *Pineal gland*.

PINEL, see *Pinus pinea*.

PINEOLI, see *Pinus pinea*.

PINEUS PURGANS, *Jatropha curcas*.

PINEWEED, *Hypericum sarothra*.

PINGUEC'ULA, *Pingue'ula*, *Pteryg'ium pin'guë* seu *lardaceum*, (from *pinguis*, 'fat,' 'fatty.') A small, whitishyellow tumour in the sclerotic conjunctiva and subjacent areolar tissue, close to the margin of the cornea on its nasal or temporal side; so called from its being supposed, but erroneously, to be of a fatty nature.

PINGUEDINOUS, *Fatty*.

PINGUEDO, gen. *Pingue'dinis*, *Fat*—*p. Renalis*, *Nephridion*.

PINGUICULA, *Pinguecula*.

PINGUIC'ULA VULGARIS, (from *pinguis*, 'fat.') *P. alpi'na*, *Sanic'ula monta'na* seu *Eboracen'sis*, *V. ol'ia palustris*, *Dodecat'yon Plin'ii*, *But'terwort*, *York'shire San'icle*. Ord. *Lentibulaceæ*. Sec. Syst. *Diandria Monogynia*. The unctuousity of this plant has caused it be applied to chaps, and as a pomatum to the hair. Decoctions of the leaves, in broths, are used by the common people in Wales as a cathartic.

PINGUID, *Fatty*.

PINGUIN, *Bromelia pinguin*.

PINGUITUDO, *Polysarcia adiposa*.

PINHOLE PUPIL, see *Pupil*, *pinhole*.

PINHONES INDICI, *Jatropha curcas*.

PINK, CAROLINA, *Spigelia Marilandica*.

PINK DYE. (*Stripped Spigelia*, $\mathfrak{z}$), *subcarb. of potass*. gr. xvij. *spirit of wine*, $\mathfrak{z}$ ij; digest for two hours; add *distilled water*, $\mathfrak{z}$ ij; digest for two hours more, and add *distilled vinegar* or *lemon juice* q. s., to reduce it to a fine rosecolour.) Used as a cosmetic.

PINK, GROUND, *Silene Virginica*—*p. Indian*, *Spigelia Marilandica*—*p. Wild*, *Silene Virginica*.

PINKROOT, *Spigelia marilandica*—*p. Demerara*, *Spigelia anthemlia*.

PINNA, 'a wing,' *Ala*, *Pavilion of the ear*—*p. Marina*, see *Byssus*.

PINNACULUM FORNICIS GUTTURALIS, (dim. of *pinnæ*.) *Uvula*.

PINNÆ (pl. of *pinnæ*) HEPATIS, *Lobes of the liver*—*p. Naris*, see *Nasus*.

PINNULA, (dim. of *pinnæ*.) *Aileron*.

PINNULÆ HEPATIS, *Lobes of the liver*.

PINO'LI (S.). A preparation used for subsistence on long journeys in the West. It is parched corn, beaten or ground very fine, and sweetened with sugar, to be used with water, and drunk on the march. It is the *cold flour of the Indians*, and early Western pioneers.

PINONCILLO TREE, *Castigleonia lobata*.

PINS AND NEEDLES, see *Leg asleep*.

PINSWEAL, *Furunculus*.

PINT, ([Sax.] *pýn'*), *Octarius*.

PINTA, (from [S.] *pintar*, 'to paint.') *Blas'tain*: a disease, which is said to prevail in Mexico, and to be a variety of *Pityriasis versicolor*.

PINUS ABIES. *P. excel'sa* seu *pice'a*, *A'bies*, *A. rubra* seu *excel'sa* seu *pice'a*, *Al'tat' thelo'a*, *Pice'a*, *Nor'way Spruce Fir*, *Yewleaved Fir*. *Nat. Ord.* Conifereæ. *Sex. Syst.* Monœcia Monadelphia. The tops are used in making *Spruce Beer*.

Essence of Spruce, *Essentia Abietis*, is a fluid extract, prepared by decoction from the twigs of this species of fir. From this is made *Spruce Beer*.

This fir affords the Burgundy pitch and common frankincense. 1. *Burgundy or White pitch*, *Pix Burgundica* (Ph. U. S. 1863) seu *al'ba* seu *ar'ida* seu *abiet'ina* seu *abietis*, *Resi'na abietis hu'mida*, *R. al'ba hu'mida*, *Pi'ni abietis resi'na spon'te concre'ta*, (F.) *Poix blanche ou grasse ou jaune ou de Bourgogne*. This prepared concrete juice is of a close consistence, rather soft, of a reddishbrown colour, and not unpleasant smell. It is very adhesive to the skin; and, consequently, forms excellent plasters for remaining upon the part for some time;—as in cases of affections of the chest. 2. *Abietis Resi'na* (Ph. L. since 1809), *Com'mon Frank'incense*, *Perrosin*, *Tus*, *Thus*, *Th. femini'nun* seu *vulgar'e*, *Olib'anum vulgar'e* seu *sykes'tr'e*, *Resi'na abietis sic'ca*, *Resi'n of the Spruce Fir*. It is solid, dry, brittle; externally, brownishyellow; internally, whitish. Used in plasters like the last.

PINUS ABIES, see *P. picea*—p. *Alba*, *P. picea*.

PINUS BALSAMEA, *A'bies balsame'a* seu *balsamif'era*, *Peu'cea balsame'a*, *Bal'sam fir*. The tree which affords the *Can'ada Bal'sam*, *Bal'samum Canaden'se* seu *de Can'ada*, *Resi'na strobili'na*, *Terebinth'ina Canaden'sis* seu *Pi'ni balsame'e*, *Can'ada Tur'pentine* or *Bal'sam*, *Bal'sam of Fir*, (F.) *Baume de Canada*. It is one of the purest turpentines, and has the common properties of those substances, and is much employed by microscopic observers.

PINUS CANADEN'SIS, *A'bies Canaden'sis*, *Hem'lock*, *Hemlock Spruce*. A tree, which is abundant in Canada, Nova Scotia, and the more northern parts of New England, and is found in the elevated and mountainous regions of the Middle States. The pitch—*Pix Canaden'sis*, *Can'ada Pitch*, *Hem'lock Pitch*—obtained from it is commonly known under the name *Hemlock Gum*.

PINUS CANDICANS, *P. Picea*.

PINUS CEMBRA, *P. Monta'na*, *Aphernous'li Pine*. It yields an agreeably scented turpentine:—the *Carpathian Bal'sam*, *Bal'samum Carpathic'um* seu *Lib'ani*, *Carpathic'um*, *Bri'aucon Tur'pentine*. The nuts, *Cem'bro nuts*, *Nu'clei Cem'bræ*, have an eatable kernel, and yield oil. The shoots yield *Riga Balsam* by distillation.

PINUS DAMAR'RA, *Ag'athis Damar'ra*, grows in the East India Islands. The juice speedily concretes into a very hard resin,—the *Damar'a turpentine*.

PINUS EXCELSA, *P. abies*—p. *Gallica*, *P. picea*.

PINUS LA'RIX, *A'bies la'rrix*, *La'rrix*, *L. communis* seu *decid'ua* seu *Europæ'a* seu *pyramida'lis*. The *Larch*, (F.) *Mélèze*. From this tree exudes *O'renberg gum*, *Bri'aucon mawna*, *Gum'mi lar'icis* seu *O'renburgen'se* seu *Uralen'se*, *Mon'na Briganti'na* seu *larice'a*. It also yields, by boring, *Com'mon Ven'ice Tur'pentine*, *Resi'na Lar'icis*, *Terebinth'ina Ven'eta* seu *larici'na*, *Pi'ni Lar'icis resi'na liq'uida*, (F.) *Térébenthine de Vénise* ou de *Mélèze*. It is usually thinner than the other sorts of turpentine.

PINUS MU'GHOS. The *Moun'tain* or *Mu'gho Pine*, *Pi'nus Pumil'io*. From the extremities of the branches exudes the *Resi'na Strobili'na*, of the Germans, or *Hunga'rian Bal'sam*. It is also obtained, by expression, from the cones. By distillation, the Hungarian balsam affords the *Krum'holz oil*, *Oleum Templi'nun*.

PINUS PALUSTRIS, see *Pinus sylvestris*.

PINUS PICEA, *P. A'bies*, *A'bies*, *A. pice'a* seu *pectin'ata* seu *Gal'lica* seu *al'ba* seu *can'dicans* seu *vulgar'is* seu *taxifolia*, *Europe'an Sil'ver Fir Tree*, *Al'tat'*, (F.) *Sapin commun*. By piercing the tubercles of the bark of this fir, the *Strasburg Turpentine*, is obtained:—the *Resi'na Abietis* (Ph. L. before 1809), *Oleum Abietis*, *Terebinth'ina Argentoratensis*.

PINUS PINASTER, see *P. sylvestris*.

PINUS PINEA, *Stone Pine*, *Pi'na*, *Pi'nus*, *P. uber'rima* seu *sativa*. The nuts, *Zir'bel nuts*, *Pine nuts*, *Nu'clei Pi'neæ*, *Pi'nei*, *Pine'oli*, (F.) *Pin à pignons*, are eaten raw, or preserved like almonds. They are, also, used in emulsions.

PINUS PUMILIO, see *Pinus Mughos*—p. *Sativa*, *P. pinea*.

PINUS SYLVESTRIS, *Pinas'ter*, *Peu'cè*, *Scotch Fir*. This pine, as well as *P. marit'ima* (*P. Pinaster*), and other species of *Pinus*, affords common turpentine and its oil, resin, tar, and pitch. 1. *Com'mon Tur'pentine of Eu'rope*, *Terebinth'ina*, *T. vulgar'is* seu *commun'is*, *Resi'na pi'ni*, *Bijou*, *Horse Tur'pentine*, *Bordeaux Tur'pentine*, (F.) *Térébenthine de Bordeaux*, *Térébenthine commune*, is obtained by wounding the tree in hot weather. It is used, chiefly, as a dressing for wounds, &c., in horses, and for the distillation of the oil, (see *Oleum Terebinthinæ*.) The *white Tur'pentine* or *common Tur'pentine of America*—*Terebinth'ina*, Ph. U. S.—is produced chiefly from *Pi'nus palustris*, *Yellow pine*, and *P. ta'eda*, *Loblolly pine*, and perhaps from other species inhabiting the Southern States. When the oil is distilled with water, yellow resin, or *Ros'in*, *Resi'na*, Ph. U. S., *Resi'na fla'va*, is left, which is only used in the formation of ointments and plasters. If without the addition of water, the residuum is *common res'in* or *Col'ophony*. 2. When the cold begins to check the exudation of the common turpentine, part of it concretes in the wounds. This is called, in France, *Gall'ipot*, *Barras*; and *White Ros'in*, *Resi'na alba*, when hardened after long exposure to the air. 3. When the old trees are subjected to distillation, in a coarse manner, Tar is obtained—*Hygropis'uos*, *Pis'sa*, *Pix ced'ria* seu *liq'uida*, *Resi'na pi'ni empyreumat'ica liq'uida*, *Terebinth'ina empyreumat'ica*, *Alchit'ram*, *Alchit'ra*, *Ced'ria*, (F.) *Goudron*, *G. végétale*, *Poix liquide*, *Brai liquide*.

Tar water, *Aqua Pice'a* seu *Picia*, *Infusum Picia liq'uida* (Ph. U. S. 1863,) seu *Picia empyreumat'ice liq'uide*, *Pos'tio pice'a*, (F.) *Eau de Goudron*, which is made by mixing two pints of tar with a gallon of water, stirring with a wooden rod for fifteen minutes, and when the tar has subsided, straining and keeping in well stopped bottles, was, at one time, a fashionable remedy in numerous complaints, and its use has been revived, since its virtues have been presumed to be owing to creasote. It is employed chiefly in pulmonary affections, and the vapour has been recommended in phthisis and other diseases of the lungs. It is used externally as a detergent. A *wine or beer of tar*, *Tar'beer*, *Jews' beer*, has been employed in Philadelphia in similar cases. It may be made by adding to tar enough beer to saturate it, mixing, and allowing them to macerate for three days, with occasional agitation. The dose is a table-spoonful. 4. *Com'mon Pitch*, *Pix nigra*, *Black Pitch*, *Bur'nea*, *Bur'nia*, *Stone Pitch*, *Pix sic'ca*, *P. al'ba*, *P. nava'lis*, *Zopis'sa*, *Palimpis'sa*, *P. ar'ida* (Ph. L. before 1809), (F.) *Poix navale*, *P. noire*, is obtained by inspissating tar. It is used only as a resolvent in plasters.

PINUS TAEDA, see *P. sylvestris*—p. *Taxifolia*, *P. picea*—p. *Uber'rima*, *Pinus pinea*—p. *Vulgaris*, *P. picea*.

PIONE, *Pæonia*.

PIONY, *Pæonia*.

PIORTHOPNŒA, (πιωρ, 'fat,' and orthopnœa,) Pimelorthopnœa.

PIOULQUES (F.). A kind of sucking pump, invented by Louis, for extracting water that had entered internal cavities, in cases of drowning. A useless instrument.

PIP, Syphilis.

PIPE PLANT, *Monotropa uniflora*.

PIPER, (πῑπερ.) see *Piper nigrum*.

PIPER ALBUM, *Leucopeper*, *White Pepper*, is black pepper freed from its cuticle.

PIPER ANGUSTIFOLIUM, *Matico*—p. *Aromaticum*, *P. nigrum*—p. *Betel*, *Betel*—p. *Brazilianum*, *Capsicum annuum*—p. *Calecuticum*, *Capsicum annuum*.

PIPER CAPENSE, a South African species, has all the properties of the peppers, and, in appearance and taste, greatly resembles cubebs. It possesses, too, similar virtues.

PIPER CARYOPHYLLATUM, see *Myrtus pimenta*—p. *Caudatum*, *P. cubeba*—p. *Chiapæ*, see *Myrtus pimenta*.

PIPER CUBERÆ, *Cubebæ officinalis*, *Persea seu Lau'rus cubebæ*, *Litsæa cubebæ* seu *piperi'ta*. *Ord. Piperacæ*. A native of Java and Guinea. The odour of these berries—*Cubebæ* (Ph. U. S.), *Com'peper*, *Com'peba*, *Cubal sini*, *Pi'per caudatum*, *Bac'cæ Pi'peris glab'ri*, *Cu'beb* or *Tail'ed Pepper*, *Canna'mus*, (F.) *Poire-à-queue*, *Quabebe*—is aromatic; taste at first cooling, afterwards pungent. The active principle is an essential oil—*O'leum Cubebe*, oil of cubebs—which is official in the Pharm. U. S. The properties of the cubeb are stimulant and purgative. It is used only in gonorrhœa. Dose, from ℥j to ʒj, in powder, three or four times a day; of the volatile oil, 10 or 12 drops.

Tur'key yel'low ber'ries,—the dried fruit of the *Rhamnus Catharticus*—are often substituted for cubebs.

PIPER ELONGATUM, *Matico*.

PIPER GUINEENSE, *Capsicum annuum*—p. *Hispanicum*, *Capsicum*—p. *Indicum*, *Capsicum annuum*—p. *Jamaicense*, see *Myrtus pimenta*.

PIPER LONGUM, *Macrop'iper*, *Acapali*, *Caturip'ali*, *Pon'pili'm*, *Chav'ica Roxburg'hii*, *Long Pep'per*, (F.) *Poire long*. As we receive the berries, they consist of the unripe fruit dried in the sun; and are small, round grains, disposed spirally on a cylindrical receptacle. They are stimulant, carminative, and tonic, like the other peppers. Dose, gr. v to ℥j.

PIPER LUSITANICUM, *Capsicum annuum*—p. *Methysticum*, see *Ava*—p. *Murale*, *Sedum*.

PIPER NIGRUM, *Melanop'iper*, *Molagoco'di*, *La'da*, *Pi'per aromaticum*, *Black Pep'per*, (F.) *Poire noir*. *Black Pepper*, *Piper* (Ph. U. S.), is the unripe fruit dried in the sun. *Virtues*, same as those of the other peppers. When freed from its husk by mechanical trituration it constitutes *Ful'ton's deco'rated pepper*.

PIPER ODORATUM JAMAICENSE, see *Myrtus pimenta*—p. *Tabascum*, see *Myrtus pimenta*—p. *Turcicum*, *Capsicum annuum*.

PIPERINA, *Piperinum*, *Pip'erine*, *Piperin*, (from *piper*.) A resinoid substance, obtained from the *piper nigrum* by M. Erstädt, in which a part of the virtues of the pepper resides. It has been employed, like quinia, in intermittents, and successfully. Dose, gr. v—x.

PIPEROID, *Piperoidum*, *Zingib'erin*, (*piper*, and *eidos*, 'resemblance.') A yellow syrupy mass obtained by exhausting ginger root by ether. It is soluble in ether, alcohol, and oils, and may be given in tincture, sugar, lozenge, or syrup, in cases where ginger is indicated.

PIPEROID OF GINGER, *Oleoresina zingiberis*.

PIPEROPAS'TUS, (*piper*, and *σαστ*, 'to strew.') Strewed with pepper,—socks, &c. ex-

ample, where it is desirable to excite the surface of the foot.

PIPETTE. 'A small pipe.' In Pharmacy, a small tube terminating in a perforated point, which is passed into a liquid to be removed,—from the surface of a precipitate, for example. The liquid is drawn into the tube by applying the mouth to the free extremity.

PIPIN'NA, *Men'tula par'va*. A small penis.

PIPPERIDGE BUSH, see *Oxyacantha Galeni*.

PIPSISSEWA, *Pyrola umbellata*—p. *Spotted*, *Pyrola maculata*—p. *White*, *Pyrola maculata*.

PIPTONYCHIA, (πιπτο, 'to fall,' and *ονyx*, gen. *ονυχος*, 'the nail,') *Onychoptosis*.

PIPTOSTEGIA OPERCULATA, see *Batata de purga*—p. *Pisonis*, *Convulvulus mechoacan*.

PIQUETÉ, (from *piquer*, 'to prick, to lard,') see *Sablé*.

PIQÛRE, *Puncture*, *Wound*, *punctured*.

PIQUEURS MÉDECINS, 'Puncture doctors.' A name at one time given in derision, in the Parisian hospitals, to those physicians who employed acupuncture in almost all diseases.

PIRIFORMIS, *Pyramidalis*.

PIROGOFF'S OPERA'TION. A mode of amputating at the ankle, proposed by Pirogoff, of Russia, which consists in retaining a portion of the calcaneum to impart greater length and roundity to the stump.

PIRTLING, (old E.) *Slaving*.

PIRUS, see *Pyrus malus*.

PISA, CLIMATE OF. Pisa has long been celebrated as one of the most favourable climates in Italy, for the consumptive. It is genial, but somewhat oppressive and damp. It is softer than that of Nice, but not so warm; less soft, but less oppressive than that of Rome.

PISCIARELLI, MIN'ERAL WA'TERS OF. A spring at Monte Secco, in Italy, between Lake Agnano and Solfatara. It is thermal (100° Fah.), sulphureous, and carbonated; and is chiefly used in old ulcers, leucorrhœa, chronic diarrhœa, and in cutaneous affections.

PISCID'IA ERYTHR'NA, (*piscis*, 'a fish,' and *cædere*, 'to kill,') *Jama'ica Dog'wood*. *Ord. Leguminosæ*. A small branching tree, common in most of the West India Islands; *Diadelphia Decandria* of Linnaeus. It is used in the West Indies to poison fish. It appears to be an acro-narcotic, and has been used in the form of a tincture of the bark of the root in toothache.

PISCINA, 'a fish pond,' *Baignoire*.

PISCIV'OROUS, *Pisciv'orus*, *Ichthyoph'agous*, *Ichthyoph'agus*, (*piscis*, 'a fish,' and *voro*, 'I eat,') Feeding or subsisting on fish.

PISIFORM, *Pi'siformis*, (*pisum*, 'a pea,' and *forma*, 'shape,') Pealike, pea-shaped.

PISIFORM'E, *Os*, *Os lenticula'ræ* seu *orbicula'ræ* seu *ex'tra ord'inem car'pi*, (F.) *Os hors du rang*, is the fourth bone of the first row of the carpus. It is round, and gives attachment to the tendon of the cubitalis anterior, and to the anterior transverse ligament of the carpus. It is articulated, behind, with the *os cuneiforme*.

PISO, *Mortar*.

PISSA, (πίσσα,) *Pitch*, see *Pinus sylvestris*.

PISSASPHAL'TUS, *Pissasphal'tum*, *Min'eral Pitch*, (*pisssa*, and *ασφαλ'tος*, 'bitumen,') (F.) *Poiz minérale*. A thicker kind of *Petro'leum* or *Rock oil*; formerly employed as a vulnerary. It is synonymous with *Glu'tinous bitu'men*, *Mal'ta bitu'men*, and *Min'eral tar*.

PISSELE'UM, *O'leum Pic'inum*, (*pisssa*, and *ελαον*, 'oil,') An oily matter obtained from boiling pitch, and used for the same purposes as tar.

PISSELEUM INDICUM, *Petroleum*.

PISSEMENT, *Micturition*—p. *de Pus*, *Puria*—p. *de Sang*, *Hæmaturia*.

PISSENLIT, ('piss a-bed,' *Leontodon taraxacum*).

PISSE-SANG, ('piss blood,' *Fumaria*).

PISSE'ROS, (*πισσῆρος*, 'made of pitch.') Ancient name of a cerate, prepared of *olive oil*, *oil of roses*, and *dried pitch*. It was, also, called *Cera'tum picat'um*, and *C. tetraphyl'um*.

PISSTITES, (*πισσιτης*), *Vinum picat'um*. A wine prepared from tar and the must of the grape. It was regarded to be stomachic.

PISSOSIS, (from *πασα*, 'pitch,') see Depilatory.

PISSWEED, *Anthemis cotula*.

PISTACHIO NUTS, (*πισταχια*), see *Arachis hypogea*, *Pistacia vera*.

PISTACIA CHIA, *P. lentiscus*.

PISTACIA LENTIS'CUS, *P. Chi'a*, *Terebin'thus lentis'cus*, *Lentis'cus vulga'ris*, *Schi'nos*; the *Mas'tich Tree*. *Ord. Terebinthaceæ*. The wood of this tree abounds with the resinous principles of mastich; and a tincture made from it has been used in the cure of dyspeptic affections, gout, and dysentery. It yields, by incision, the resin *Mas'tich*, *Mas'tic*, *Resi'na Lentis'cina*, *Men'frige*, *Mas'tiche* (Ph.U. S.), *Mas'tic*, *Pista'cie Lentis'ci Resi'na*. This resin has an agreeable smell when heated: is almost insipid; in globular, irregular, yellowish, semitransparent masses; soluble in ether; partially so in alcohol. It is stimulant, and has been used in old coughs, and gleet, and is chewed to sweeten the breath. It is sometimes, also, employed to stop holes in carious teeth.

Oil of Mas'tich, *Mastichela'on*, *O'leum Mastichi'num*, is mentioned by Dioscorides.

PISTACIA NARBONENSIS, *P. lentiscus*—*p. Reticulata*, *P. lentiscus*.

PISTACIA TEREBIN'THUS, *P. Narbonen'sis* seu *reticulata* seu *ve'ra*, *Tur'pentine Tree*, *Hab-el-Kalimbut*, *Terebin'thus vulga'ris*, *Termin'thus*. The tree which affords the *Cy'prus Tur'pentine*, *Resi'na Terebin'thi*, *Terebin'thina ve'ra* seu *Chi'a* seu *Cy'pin*, (F.) *Terebinthine de Chio*. This substance is classed amongst the resins. It is procured by wounding the bark of the trunk. The best Chio turpentine is of about the consistence of honey; very tenacious, clear, and almost transparent; of a white colour inclining to yellow, and of a fragrant smell; moderately warm to the taste, but free from acrimony and bitterness. It resembles in virtues the other turpentine.

PISTACIA VE'RA, (*πισταχια*). Name of the tree which affords the *Nux Pista'cia*. *Pista'chio nuts* have a sweetish, unctuous taste, resembling that of sweet almonds. Like these, they afford an oil, and may be formed into an emulsion.

PISTATION, Epistation.

PISTILLUM, (*pastare*, 'to pound,') *Pilum*.

PISTOLOCHIA, (*πιστος*, 'faithful,' and *λοχεια*, 'parturition,') *Aristolochia pistolochia*.

PISTORIENSIS GLADIUS, (from *Pistorium*, now *Pistoia*, in Tuscany,) *Bistouri*.

PI'SUM, (*πισον*), *P. sativum*, *Pea*, *Garden pea*, (F.) *Pois*, *P. cultivé*. The seeds of the pea. *Ord. Leguminosæ*, *Sec. Syst. Diadelphia Decandria*, are much employed as an aliment, both in the fresh and dried state. They are farinaceous, and, when dried, not easy of digestion.

PIT, ([Sax.] *pyt* or *pic*), Pock mark—*p.* of the Stomach, *Fossette du cœur*.

PITA'YA BARK, of South America. The bark of *Cincho'na pita'ya*, *Ord. Cinchonaceæ*, is preferred in Colombia to any other. It contains a new vegetable alkali, to which the name *Pitayine* has been given.

PITAYINE, see *Pitaya bark*.

PITCAITHLEY, MIN'ERAL WATERS OF. These saline springs, at Pitcaithley, about four miles from Perth, Scotland, contain chlorides of sodium and calcium.

PITCH, (from [L.] *pix*), *Colophonia*, *Pissa*—*p.*

Black, see *Pinus sylvestris*—*p. Burgundy*, see *Pinus abies*—*p. Canada*, see *Pinus Canadensis*.

PITCH CAP, see Depilatory.

PITCH, COMMON, see *Pinus sylvestris*—*p. Hemlock*, see *Pinus Canadensis*—*p. Jews'*, *Asphaltum*—*p. Mineral*, *Pissasphaltum*.

PITCH PLASTER, see Depilatory, *Emplastrum picis*.

PITCH, STONE, see *Pinus sylvestris*—*p. White*, see *Pinus abies*.

PITCHER PLANT, *Sarracenia purpurea*.

PITHYRINUS, (from *pityron*, 'bran,') *Acerosus*, *Furfuraceus*.

PITTA'Cium. A pitch plaster, (from *πιττα* 'pitch,')

PITTING, Pock mark.

PITTOSIS, (from *πιττα*, 'pitch,') see Depilatory

PITTO'TA. Medicines in which pitch is the chief ingredient.

PITTS'BURG, MIN'ERAL WATERS OF. A chalybeate mineral spring, situate about four miles south-west of Pittsburg, in Pennsylvania. It contains chloride of sodium, chloride of magnesium, oxide of iron, sulphate of lime, and carbonic acid.

PITUITA, Mucus, Phlegm—*p. Narium*, Nasal mucus.

PITUITAIRE, Pituitary, Pituitary membrane.

PITU'ITARY, *Pituita'rium*, (F.) *Pituitaire*, (from *pituita*.) Concerned in the secretion of mucus or phlegm. A name given to several parts.

PITU'ITARY BODY, *P. Gland*—*p. Fossa*, *Sella turcica*.

PITUITARY GLAND, *Glandula Pituitosa* seu *pituitaria* seu *basilaris*, *Lacuna* seu *Appendicula cerebri*, *Senti'na*, *Hypophysis* seu *Appendix cerebri*, *Glans pituitam excipiens*, *Corpus pituitaire*, *Pituitary body*;—(F.) *Glande ou Corps pituitaire*, *Appendice sus-sphénoïdale du cerveau*, (Ch.) A small, round body; the anterior portion of which is of a cineritious yellow hue, and the posterior, which is smaller, is whitish and pulpy. It occupies the sella turcica of the sphenoid bone. Its functions are unknown.

PITUITARY MEMBRANE, *Schneiderian Membrane*, *Membra'na pituitaria* seu *olfactoria*, (F.) *Membrane nasale ou olfactive ou pituitaire ou de Schneider*, *Pituitaire*. The mucous membrane which lines the nasal fossæ, and extends to the different cavities communicating with the nose. In the nasal fossæ, the membrane appears formed of two layers, intimately united;—the one in contact with the bone is fibrous;—the other is free at one surface. The latter is a mucous membrane, in which papillæ are not easily distinguishable, but which contains them, as well as a number of very small mucous follicles, that open into the cavity of the nares. On penetrating the maxillary, frontal, sphenoidal, and ethmoidal sinuses, the membrane becomes very thin, transparent, less vascular, and seems reduced to its mucous layer. The pituitary membrane receives the impression of odours by means of the expansion of the olfactory nerves upon it. It is the seat of smell.

PITUITE, Mucus.

PITU'ITOUS, *Pituito'sus*, (F.) *Pituiteux*, (from *pituita*.) Consisting of, or resembling, or full of mucus or phlegm.

PITYRE'NUS, (*πιτυρηνος*), *Acerosus*, *Furfuraceus*.

PITYRI'ASIS, (*πιτυρις*, 'bran,') *Lepido'sis Pityri'asis*, *Pityri'sma*, *Her'pes furfuraceus* seu *farinosus*, *Porri'go* (of some), *Tin'ea furfuracea* seu *porriginosa*, *Furfuratio*, *Furfuris'ca*, *Dandruff*, *Dander*, (Sc.) *Luas*, (Prov.) *Room*, *Rummet*, (F.) *Teigne*, *Dartre*, *D. furfuracea volante*;—a very superficial affection; characterized by irregular patches of thin scales, which repeat

edly exfoliate and recur; but which never form crusts, or are accompanied with exoriations. It is not contagious. It occurs under three or four varieties of form.

PITYRIASIS, Porrigo.—p. Nigra, Melasma.—p. Versicolor, Chloasma.

PITYRIASMA, gen. *Pityriasis*, (πυρίσμα,) Pityriasis.

PITYROIDES, (πυροειδής, from πυρον, 'bran,' and εἶδος, 'resemblance,') Furfuraceus.

PITYRON, (πυρον, 'bran,') Furfur.

PIVOINE, Pæonia.

PIVOTING. In dentistry, an operation which consists in fastening a new crown upon the root of a tooth by means of a piece of gold wire, called 'a pivot.'

PIX, gen. *Piceis*, (πίσσα, 'pitch,') see Pinus sylvestris, Sperm.—p. Abietina, see Pinus abies.—p. Abietis, see Pinus abies.—p. Alba, see Pinus abies.—p. Arida, see Pinus abies, and Pinus sylvestris.—p. Atra, see Pinus sylvestris.—p. Brutia, Brutia.—p. Burgundica, see Pinus abies.—p. Canadensis, see Pinus Canadensis.—p. Cedria, see Pinus sylvestris.—p. Græca, Colophonia.—p. Liquida, see Pinus sylvestris.—p. Navalis, see Pinus sylvestris.—p. Nigra, see Pinus sylvestris.—p. Sicca, see Pinus sylvestris.

PLACE'BO, 'I will please,' (from placeo.) A medicine, prescribed rather to satisfy the patient than with any expectation of its effecting a cure.

PLACEN'TA, (πλάκων, gen. πλακωντος, 'a cake,') *P. uteri* seu *uterina* seu *uteri car'nea*, *He'par* seu *Jecur uteri'num*, *Cu'ro* seu *Affu'sio orbicularis*, *Ma'za*. A soft, spongy, vascular body, adherent to the uterus, and connected with the fœtus by the umbilical cord. It is not in existence during the first period of the embryo state, but its formation commences perhaps with the arrival of the embryo in the uterus. It is generally considered to have two portions, one *fœtal*, and the other *maternal*. The fœtal consists of highly vascular villi and tufts, which contain the inosculating loops of the umbilical arteries and umbilical vein of the fœtus. The maternal portion is considered by Dr. Reid to consist, essentially, of a large sac formed by the inner coat of the vascular system of the mother, into which the maternal blood is poured by the curling arteries of the uterus, and from which it is returned by the uteroplacental veins. The tufts and villi of the fœtal placenta are ensheathed by this inner coat of the maternal vascular system; and, according to Prof. Goodsir, two distinct sets of nucleated cells—one maternal and the other fœtal—separate the maternal and fœtal portions where they come in contact, one set of which are probably, he thinks, concerned in the separation of nutritive matter from the maternal blood, and the other destined to convey it into the vessels of the fœtus. The placenta is formed of several lobes or cotyledons, which can be readily distinguished from each other on the uterine surface; but towards the fœtal surface are confounded into one mass. It is composed of the umbilical vessels, areolar tissue, and whitish, resisting filaments, which are vascular ramifications obliterated. Lymphatic vessels have also been presumed to exist in it; and nerves have been traced, proceeding from the great sympathetic of the fœtus.

The umbilical cord usually arises from the centre of the placenta; but occasionally from its edge, when it is called *bat'ledore placenta*.

The main function of the placenta appears to be like that of the lungs in the adult. It admits of the blood of the fœtus being shown to that of the mother, and undergoing requisite changes. It may, also, be an organ for nutritive absorption, as observed above.

Animals, which have no placenta—as the Marsupialia—are called *implacental* and *nonplacental*.

PLACEN'TA, BATTLEDORE, see Placenta.—p. Cruoris, see Blood.—p. Febrilis, Ague cake.—p. Oblata, P. prævia.—p. Obvia, P. Prævia.

PLACENTA PRÆ'VIA, (præ, 'before,' and via, 'way,') *P. ob'via* seu *obla'ta*. Presentation of the placenta; a condition which gives rise to *unavoidable* hemorrhage from the uterus. The safety of the female depends upon speedy delivery.

PLACENTA SANGUINIS, see Blood.—p. Uteri Succenturiata, Decidua.

PLACEN'TAL, *Placenta'lis*, (F.) *Placentaire*. Relating or appertaining to the placenta. Possessed of a placenta, —as 'a placental mammal.'

PLACENTITIS, *Placenti'tis*, *Placoi'tis*, *Inf'lamma'tio placen'tis*, (placenta, and itis.) Inflammation of the placenta.

PLACEN'TULA, 'a little cake,' (dim. of placenta, 'a cake,') A rudimental placenta. The placenta of an abortion.

PLACOTIS, (πλάκων, 'placenta,' and itis,) Placitis.

PLACOUNTITIS, (πλάκων, gen. πλακωντος, 'placenta,' and itis,) Placentitis.

PLADAROMA, gen. *Pladaro'matis*, (πλαδάρωμα,) Pladarosis.

PLADAROSIS, *Pladaro'ma*, *Pladar'otes*, (from πλαδάρων, 'soft,' and osis.) A soft tumour on the eyelid, unaccompanied with redness or pain. Trachoma.

PLADAROTES, (πλαδάρων,) Pladarosis.

PLAFOND DES VENTRICULES DU CERVEAU, ([F.] *plafond*, 'a ceiling,' from plat, 'flat,' and fond, 'ground,') Corpus callosum.

PLAGA, (πλάγη,) Ictus, Wound. Formerly, any kind of wound or disease from violence.

PLAGIOBOL'IA, *Planiobol'ia*, (πλάγος, 'to one side,' and βάλλειν, 'to throw,') Imperfect emission of sperm into the vagina. Emission to one side; sometimes practised to render impregnation less probable.

PLAGUE, (from πλῆγη, *plaga*, 'a stroke,') *Pestis*, *P. Orienta'lis* seu *bubon'ria orientalis* seu *inguina'ria* seu *contagio'ria* seu *acutis'sima* seu *septic'a* seu *glandu'osa* seu *ad'eno-septic'a* seu *inguina'ria*, *Pestil'entia*, *Ephem'era pestilenti'alis*, *E. mortif'era*, *Ta'bun*, *Lu'ës inguina'ria*, *Clu'des glandula'ria*, *Phtho'ra*, *Pestilen'tia*, *Lu'ës*, *Lo'i'mos*, *Fe'bris ad'enuer'osa*, *Læ'mos*, *Lu'm'e*, *Læ'mia*, *Læmop'gra*, *Chaos'da*, *Anthrax'ia Pestis*, *An'thracoty'phus*, *Fe'bris pes'tilens*, *Syn'ochus pestilenti'alis*, *Ty'phus pestis* seu *pestilenti'alis* seu *gravis'simus* seu *anthraci'cus* seu *buboni'cus*, *Ex-anthe'ma pestis*, *Pest*, *Lev'ant plague*, *Sep'ticæ* or *glandular pestilence*, *The sick'ness*, (F.) *Peste*, *Typhus d'Orient*, *Fièvre du Levant*, *F. ad'enuer'veuse*. An eminently malignant disease; endemic in the Levant; frequently epidemic, and destroying at least two-thirds of those it attacks. It is a fever of the most aggravated kind, with affection of the lymphatic glands of the groins or axillæ, and carbuncles. Its miasmata—it has been conceived—adhere to different organic textures, to woollen goods, clothing, and furniture; and may thus be transported to a considerable distance; but this is disputed. The mean duration of the disease is six or seven days; some die in twenty-four hours, others not till ten or twelve days. Pathological anatomy has afforded little light with respect to it. Various means have been used for arresting it, but none have seemed to be pre-eminently distinguished. The great point is, to watch the indications as they develop themselves; and to treat the case, in general, like one of typhus gravior. It is universally agreed that the suppuration of the buboes should be aided as far as practicable. For preventing the importation and spread of the plague, the Quarantine Laws were instituted; and when the disease has actually appeared, a cordon sanitaire

has been drawn round the infected district, so as to prevent all communication.

PLAGUE OF ANTONINUS, *Pes'tis Antoniniana*, (F.) *Peste Antonine*. A severe febrile disease, which prevailed in the Roman empire, during the reign of Antoninus. It began in Asia, and passed westward. It resembled the Plague of Athens.

PLAGUE OF ATHENS, (F.) *Peste d'Athènes*. A febrile disease, which raged at Athens during the Peloponnesian war, in the 5th century before the Christian era; and a description of which is given in Thucydides. It was characterized by a cutaneous eruption, accompanied by small ulcerations, vomiting, affection of the respiratory organs, and diarrhoea. It began in the far East, and, before reaching Athens, ravaged Egypt, and the greater part of the Persian Empire.

PLAGUE, BLACK, Death, Black.

PLAGUE, COLD. A severe form of congestive fever, seen in the Southern States. Bilious pneumonia, in which there is no reaction, has been, also, so called.

PLAGUE CROSS. A cross formerly affixed by the authorities to the door of a house in which there was plague.

PLAGUE, PA'LI. A malignant fever, greatly resembling plague, which prevailed in Rajpootana after July, 1836.

PLAGUE SPOT, Plague token.

PLAGUE TO'KEN, *Plague mark*, *Plague spot*. A mark by which one struck with the plague was known. Perhaps the early stage of carbuncle in unfavourable cases. It is described as a small tubercle—*Bos'sa*—somewhat resembling a wart, callous, and more or less deficient in sensibility, varying in size from that of a millet-seed to that of a bean.

PLAGULA, (dim. of *plaga*,) Compress, Pledget, Splint.

PLAIE, (from *plaga*,) Wound—*p. d'Arme à feu*, Wound, gunshot—*p. d'Arquebuse*, Wound, gunshot—*p. Contuse*, Wound, contused—*p. Envenimée*, Wound, poisoned—*p. par Arrachement*, Wound, lacerated—*p. à Lambeaux*, see Flap.

PLAINTIVE, (from [F.] *plaindre*, 'to complain') (F.) *Luctueux*.

PLANANTHUS FASTIGATUS, *Lycopodium selago*—*p. Selago*, *Lycopodium selago*.

PLANARIA LATIUSCULA, (from *planus*, 'flat,') *Distoma hepaticum*.

PLANARTHRA'GRA, (*πλανη*, 'a wandering,' *αρθρον*, 'a joint,' and *αγχα*, 'a seizure,') Gout (wandering).

PLANCHER (F.), 'floor.' In anatomy, the inferior wall or boundary of a cavity.

PLANCHER DU CERVEAU, Tentorium.

PLANCUS, (*planca*, 'a plane, a table,') *Leiodius*.

PLANE, (*πλανη*, 'a wandering,') *Plexus*.

PLANE, MESIAL, see Mesial.

PLANETSTRUCK, see Sideratio.

PLANE'TES, (*πλανητης*,) *Planeticus*, *Planeticus*, *Errabundus*, *Erratic*, (from *πλανη*, 'a wandering,') An epithet given to diseases whose returns are irregular—especially to intermittent fever; *Planet's Fe'bris*, *F. Erratica*, *F. Erronea*, *F. Vaga*.

PLANETICA ARTHRITIS, Gout, wandering.

PLANICUS, *Planetes*.

PLANIPEDES, (*planus*, 'flat,' and *pes*, gen. *pedis*, 'foot,') *Steganopodes*.

PLANITIES PEDIS, (from *planus*, 'flat,') Sole.

PLANO'BOLIA, (*πλανη*, 'a wandering,' and *βαλλειν*, 'to throw,') *Plagiobolia*.

PLANODIA, (*πλανη*, 'a wandering,' and *ὄδω*, 'a way,') A false passage,—as occasionally results from introducing the catheter.

PLANTA, Sole—*p. Noctis*, *Hydroa*, *Sudamina*—*p. Pedis*, Sole—*p. Prima pedis*, *Tarsus*.

PLANTA'GO, gen. *Plantag'inis*, (*planta*, 'sole,' ... from resemblance of the leaves,) *P. Ma'jor*,

Centiner'via, *Arnoglos'sum*, *Septinsr'via*, *Polyn'e'ron*, *Heptapleu'ron*, *Planta'go latifo'lia* seu *ma'jor* seu *vulgaris*, *Great plantain*, *Way'bread*, (Sc.) *Wa'bran leaves*, (F.) *Plantain commun.* Ord. *Plantaginaceæ*. See. *Syst. Tetrandria Monogynia*. The leaves have been said to be refrigerant, attenuant, subtyptic, and diuretic.

PLANTAGO AQUATICA, *Alisma plantago*.

PLANTAGO CORDA'TA, *Heartleaved plantain*, *P. LANCEOLA'TA*, *Ribgrass*; and *P. VIRGINICA*, *White plantain*, or *Rib'wort*, indigenous, have properties like *Plantago major*.

PLANTA'GO CORON'OPUS, *Coronopod'ium*, *Cor'næ cervi'nium*, *Stell'a ter'ræ*, *Plantago*, *Buckshorn Plantain*. Properties like the last.

PLANTAGO NITIDA, *P. psyllium*.

PLANTAGO PSYLLIUM, *P. nitida*, *Psyllium*, *P. erectum*, *Pulica'ris her'ba*, *Crystal'lion*, *Cynomoi'a*, *Branch'ing Plantain*, *Flea'wort*, (F.) *Herbe aux puces commune*. The seeds have a nauseous, mucilaginous taste, and no remarkable smell. The decoction of the seeds is recommended in hoarseness and roughness of the fauces.

PLANTAGO VULGARIS, *Plantago*.

PLANTAIN DES ALPES, *Arnica montana*—*p. Arbre*, *Musa paradisiaca*—*p. Branching*, *Plantago psyllium*—*p. Buckshorn*, *Plantago coronopus*—*p. Commun*, *Plantago*—*p. d'Eau*, *Alisma plantago*—*p. Great*, *Plantago major*—*p. Heartleaved*, *Plantago cordata*.

PLANTAIN MEAL, see *Musa paradisiaca*.

PLANTAIN, RATTLESNAKE, *Goodyera pubescens*—*p. Robin's*, *Erigeron bellidifolium*.

PLANTAIN TREE, *Musa paradisiaca*.

PLANTAIN, WATER, *Alisma plantago*, *Unisema deltifolia*—*p. White*, *Plantago Virginica*.

PLANTAIRE, *Plantar muscle*—*p. Grêle*, *Plantar muscle*.

PLAN'TAR, (from *planta*,) *Planta'ris*, (F.) *Plantaire*. That which relates or belongs to the sole of the foot. This epithet has been given to several parts.

PLANTAR APONEURO'SIS, *P. fas'cia*. A thick, dense, resisting aponeurosis; of a triangular shape, and superficially situate, which occupies the middle and sides of the sole of the foot, and furnishes insertions to several of the muscles of the region, either directly or through the medium of septa, which it sends between them.

PLANTAR ARCH, see *Plantar arteries*.

PLANTAR ARTERIES. These are two in number, and arise from the extremity of the posterior tibial artery. They are distinguished into:—The *inter'nal plantar ar'tery*, situate above the abductor pollicis pedis. It extends from the termination of the posterior tibial to the trunk of one of the collaterals of that toe.—The *exter'nal plantar ar'tery* is larger than the preceding, and ought to be considered as the termination of the posterior tibial. Situate above the flexor communis digitorum pedis, it extends from beneath the calcaneum to the posterior extremity of the last metatarsal space. When it has reached this part, it turns from without inwards; advances towards the posterior extremity of the first metatarsal bone, and anastomoses with the perforating branch of the *dorsalis tarsi*. In the first part of its course, it furnishes calcaneal, articular, muscular, and adipous branches. The second portion of the artery is called the *Plan'tar Arch* or *Cross*. It gives branches in all directions. The inferior and posterior have received no name. The superior—three in number—are the *poste'rior per'forating ar'teries*. The anterior branches are larger, and there are commonly four. They furnish the *ante'rior per'forating*, and the *collat'e'als* of the last four toes.

PLANTAR CROSS, see *Plantar arteries*—*p. Fas'cia*, *P. aponeurosis*.

PLANTAR LI'G'AMENTS. A name given to the

inferior ligaments, which unite the bones of the tarsus and metatarsus, to distinguish them from the superior, called *dorsal*.

PLANTAR MUSCLE, *Plantaris, P. gracilis, Tibialis gracilis, Extensor tar'si m'ior, (F.) Plantaris, P. Grêle, Jambier grêle, Petit fémoro-calcaneus, (Ch.)* A muscle, situate at the posterior part of the leg. It is long, thin, and narrow. Above, it is attached to the posterior part of the external condyle of the os femoris: below, it terminates by a long tendon, which is attached to the posterior and inner part of the calcaneum. This muscle extends the leg on the foot, and, conversely. It can, also, assist in the flexion of the leg on the thigh.

PLANTAR NERVES. These are two in number, and are furnished by the termination of the posterior tibial nerve. They are distinguished into—1. The *internal plantar nerve*, which extends from the inferior part of the tibial nerve to both sides of the first three toes, and the inner side of the fourth. It furnishes, also, branches to the abductor of the great toe; to the short flexor of the toes; to the accessory; to the short flexor of the great toe, the lumbricals, &c. 2. The *external plantar nerve* proceeds from the termination of the posterior tibial to the posterior extremity of the 5th metatarsal bone, and gives branches to the flexor brevis digitorum pedis, the accessory and the abductor of the little toe. This trunk subsequently divides into two branches;—the one—*profunda*—is distributed to the short flexor of the great toe: the interosseus and abductor pollicis pedis: the other—*superficialis*—sends filaments to the short flexor of the little toe, the two sides of that toe, the outer side of the 4th, and the last lumbricalis.

PLANTAR REGIONS or SURFACES are different regions or surfaces of the sole of the foot. Three are generally reckoned,—the *internal, middle, and external*.

PLANTAR VEINS follow an arrangement similar to that of the arteries.

PLANTARIA, Dengue.

PLANTARIS, Plantar, P. muscle—p. Superficies pedis, Sole—p. Verus, see Flexor longus digitorum pedis profundus perforans (accessorius.)

PLANTARIUM, ('a nursery ground,' from *planta*, 'a plant,') Pubes.

PLANTE DU PIED, Sole.

PLANTI-SOUS-PHALANGIENS, Lumbricales pedis—p. *Tendino-phalangiens, Lumbricales pedis*.

PLANUM APONEUROTICUM DIAPHRAGMATICIS, (*planus*, 'flat, level, smooth.') The tendinous expansion of the diaphragm.

PLANUM ORBITALE, see Maxillary bone (superior.)

PLANUM OS. The papyraceous or orbital plate of the ethmoid bone—*Lamina papyracea, Lamella plana*—was formerly so called.

PLANUM PEDIS, Sole.

PLANUM SUPERIOR LINGUÆ. The upper surface of the tongue.

PLANURIA, (*πλῶν*, 'a wandering,' and *ουρον*, 'urine.') Discharge of urine through unwonted ways.

PLAQUEMINIER D'EUROPE, Diospyros lotus.

PLAQUES GAUFRÉES, 'honeycomb patches,' see Peyer glandulae—p. *Laitueuses*, see *Taches laiteuses*—p. *de Peyer*, see Peyer glandulae.

PLASMA, gen. *Plasmatis*, (*πλάσμα*, 'a thing formed,' from *πλάσσειν*, 'to form,') Liquor sanguinis. In antiquity, a gargle or mouthwash to clear the voice. Also, a pharmaceutical preparation from *glycerin* and *starch* heated together in the proportion of a fluidounce of the former and seventy grains of the latter in powder, gradually

raised to the temperature of about 240°, constantly stirring. It constitutes the basis of different ointments, and keeps better than they do as usually made. Called also *Glycamil*.

PLASMAT'IC, *Plasmaticus*, (*πλαστικόν*, 'to form,') Having the power to give form, plastic.

PLASMEXHIDROSIS, (*plasma*, *ἐξ*, 'out of,' and *ἵδρωσις*, 'perspiration.') Exudation of the plasma of the blood from the vessels.

PLASTER, (*εμπλάστρον*, from *πλάσσειν*, 'to smear, to cover over,') Emplastrum—p. Adhesive, Emplastrum resinae, Sparadrapum adhesivum—p. Adhesive, Baynton's, see Emplastrum resinae—p. Adhesive, of pitch and resins, Emplastrum de pice et resinis glutinans—p. Ammoniacum, Emplastrum ammoniaci—p. Ammoniacum with mercury, Emplastrum ammoniaci cum hydrargyro—p. of Antimony, Emplastrum antimonii—p. of Arnica, Emplastrum arnicæ—p. Aromatic, Emplastrum aromaticum—p. Assafoetida, Emplastrum asafetida—p. of Belladonna, Emplastrum belladonnæ—p. Black, see Emplastrum plumbi—p. Blister, Blister—p. Blistering, Emplastrum lyttæ—p. of Blistering fly, Emplastrum lyttæ—p. Calefacient, Emplastrum calefaciens—p. of Canada Pitch, Emplastrum picis Canadensis—p. of Cantharides, compound, Emplastrum lyttæ compositum—p. Cephalic, Emplastrum picis comp.—p. Corn, Sparadrapum viride—p. Corn, Kennedy's, Sparadrapum viride—p. Court, English, see Ichthyocolla, Sparadrapum adhesivum—p. Cummin, Emplastrum cumini—p. Defensive, Sparadrapum Galteri—p. Diachylon, Emplastrum plumbi—p. Fly, Emplastrum lyttæ—p. Frankincense, Emplastrum thuris—p. Galbanum, Emplastrum galbani—p. Galbanum, compound, Emplastrum galbani compositum—p. Gum, Emplastrum gummosum—p. of Gumresins, Emplastrum cum gumiresinis—p. Hemlock, Emplastrum cicutæ—p. Hemlock Pitch, Emplastrum picis Canadensis—p. Iron, Emplastrum ferri—p. Isinglass, see Sparadrapum adhesivum—p. Issue, Sparadrapum pro fonticulis—p. Lead, Emplastrum plumbi—p. of red Lead, see Emplastrum plumbi—p. Machine, Sparadrapier—p. Mercurial, Emplastrum hydrargyri compositum—p. Mustard, Cataplasma sinapis—p. Nuremberg, Emplastrum Norimbergense—p. Opium, Emplastrum opii—p. Pitch, see Depilatory—p. Pitch, comp., Emplastrum picis compositum—p. of Red oxide of iron, Emplastrum oxidi ferri rubri—p. Resin, Emplastrum resinae—p. Soap, Emplastrum saponis—p. of Spanish flies, compound, Emplastrum lyttæ compositum—p. Spice, see Emplastrum aromaticum—p. Sticking, Emplastrum resinae—p. Strengthening, Emplastrum ferri, Emplastrum oxidi ferri rubri—p. Warm, Emplastrum calefaciens—p. Warming, Emplastrum calefaciens—p. Wax, Emplastrum cereæ.

PLASTIC, *Plasticus, Formativus, Formativus*, (F.) *Plastique*, (*πλαστικός*, from *πλάσσειν*, 'to form,') That which forms, or serves to form.

PLASTIC ELEMENT. One which bears within it the germs of a higher form.—(Gerber.) *Plastic* or *histogenetic* or *organoleptic elements of nutrition* are those elements of food which go to the formation of organized tissue.—See Aliment.

PLASTIC FORCE. The generative or formative power in organized bodies. It is also called *Nisus formativus, Vis seu Virtus plastica seu formativa seu essentialis seu reproductiva, Facultas formativa seu nutritiva seu auctrix seu vegetativa; Blas alterativum, Motus assimilativus, Anima vegetativa, Plasticus, Virtus formativa, Plastic'itas, Plastic'ity, Force of Nutrit'ion, F. of assimilation, F. of Formation, F. of vegetation, Puissance du moule intérieur (Buffon), Bildungstriebe, Germ force, of the Germans.*

PLASTIC LYMPH, Liquor Sanguinis—p. Surgery, Morioplastice.

PLASTICISMUS, (πλασσειν, 'to form,') Plastic Force.

PLASTICITAS, Plastic Force.

PLASTICITY, Plastic Force.

PLASTILYTIC, (πλασσειν, 'to form,' and λυω, 'to loose,') Antiplastic.

PLASTILYTICUM (HÆMATOLYTICUM), Spasmemic.

PLASTIQUE, Plastic.

PLASTODYNAMIA, (πλασσειν, 'to form,' and δυναμις, 'force,') The manifestation of nutritive activity or power.

PLATA, (πλατη, from πλατυς, 'broad,') Scapula.

PLATANO, (platanus,) Musa Sapientum.

PLATANIER, (platanus,) Musa paradisiaca.

PLATANIERA ORBICULATA, (πλατυς, 'wide,' and ωρη, for 'anther,') *Large roundleaved Orchis*; indigenous. Order, Orchidaceæ; flowering in July. The leaves are large, soft and fleshy, and are used, in certain places, for dressing blisters.

PLATANUS, (πλατανος, from πλατυς, 'broad,') Musa paradisiaca.

PLATEA, (πλατυς, 'broad,') Bothriocephalus latus.

PLATEBONE, Scapula.

PLATEIASMA, Platiasmus.

PLATES, MANSFORD'S, see Galvanism.

PLATIASMUS, Plateiasma, (from πλατυς, 'broad,') Excessive development of the lips, causing the articulation to be imperfect.

PLATINA, Platinum.

PLATINUM, Platina, (from [S.] plata, 'silver,') (F.) Platine, Or blanc, Petit argent. The preparations of platinum resemble in their therapeutical properties those of gold. The BICHLORIDE, Platini Bichloridum, made by dissolving platinum in aqua regia, and the DOUBLE CHLORIDE of PLATINUM and SODIUM, Sodii chloroplatinus, Chloroplatinate of Sodium, prepared by dissolving bichloride of platinum and pure chloride of sodium, in proper proportions, in water, evaporating and crystallizing,—are the preparations used. They are not much prescribed.

PLATULA, (πλατυς, 'broad,') see Pediculus.

PLATY, (πλατυς, 'broad,') in composition, broad.

PLATYBREGMATE, (platy, and bregma,) see Cranium.

PLATYCEPHALIC, Platycephalicus, (from platy, and κεφαλη, 'head,') Having a broad skull.

PLATYCORIA, (πλατυκορια, from platy, and κορη, 'pupil,') Mydriasis.

PLATYCORIASIS, Mydriasis.

PLATYELMINS, (platy, and ελμινς, 'worm,') Bothriocephalus.

PLATYNOSIS, (from πλατυς, 'broad,' and οσις, 'Amplification,') Morbid extension of parts.

PLATYOPHTHALMUS, (πλατυοφθαλμος, from platy, and οφθαλμος, 'eye,') One who has broad eyes. Antimony was of old called *Platyophthalmion*, under the idea, that it improved the appearance of the eyes.

PLATYPODIA, (platy, and πους, gen. ποδος, 'a foot,') see Kylosis.

PLATYPUS, (πλατυπους,) see Kylosis.

PLATYSMA, gen. *Platysmatis*, (πλατυσμα, from πλατυς, 'broad,') Anything broadly extended. A spread plaster.

PLATYMA MYOIDES, *P. Myoides*, *Musculus cutaneus subcutaneus* (Winslow), *Quadratus genæ seu latissimus colli*, *Explanatio musculoæ*, *Tetragonus*, *Stomomaxicon*, (F.) *Thoraco-maxillofacial*, *Thoraco-facial* (Ch.), *Peancier*. A muscle situate superficially on the lateral parts of the neck. It is flat, broad, and quadrilateral. Its fibres, which are all parallel to each other, pass obliquely upwards, and from without inwards.

They arise from the areolar tissue, which covers the anterior and superior part of the chest; pass before the clavicle, and proceed to attach themselves at the interior part of the symphysis menti, at the external oblique line of the jaw, and at the commissure of the lips. The fibres of this last insertion form a distinct fasciculus, called by some anatomists *Musculus Risorius Santorini*. The platysma myoides depresses the commissure of the lips, and carries it outwards. It contributes, also, to the depression of the lower jaw. When it contracts, it wrinkles the skin of the neck transversely.

PLATYSTERNUS, (platy, and στερνον, 'the sternum,') One who has a broad, well developed chest.

PLATYSTOMUS, (platy, and στομα, 'mouth,') One who has a broad mouth.

PLAUTUS, Leiopus.

PLECHAS, (πληχας,) Perinaeum.

PLECTANÆ, Cornua uteri.

PLECTANE, *Plica*, *Plegma*, *Plexus*, (πλεκτανη, from πλεκω, 'I twine,') A fold; a plexus. Also, a cornu of the uterus.

PLECTRUM, (πλεκτρον,) The styloid process of the temporal bone. Also, the uvula, and the tongue.

PLEDGET, ([D.] Plagghe,) *Plumaceolus*, *Lintamen*, *Plagula*, (F.) *Plumaceau*, *Plumaseau*. A small compress or gâteau of lint—the filaments arranged parallel to each other—flattened between the hands after the extremities have been cut off or folded down. It is applied over wounds, ulcers, &c., to preserve them from the contact of the air, to retain dressings *in situ*, and to absorb the discharges.

PLEGE, (πληγη,) Blow, Ietus.

PLEGMA, (πληγμα,) Plectane, Plege, Plexus.

PLEGOMETER, (πληγη, 'a stroke,' and μετρον, 'measure,') Pleximeter.

PLEIN (F.), (from plenus, 'full,') The middle part of a roller. See Bandage or Roller.

PLEIOMASTHUS, (πλειων, 'more,' and μασθος, 'a breast,') Polymasthus.

PLEIOMAZIA, (πλειων, 'more,' and μαζος, 'a breast,') see Polymasthus.

PLENA MULIER, 'a full woman,' Pregnant.

PLENCK'S DEPILATORY, see Depilatory, Colley's.

PLENITUDE, (plenus, 'full,') Fulness.

PLENITUDO, Plethora.

PLEOMAZIA, (πλειων, 'more,' and μαζος, 'a breast,') see Polymasthus.

PLEONECTICUS, (πλεονεκτικός, from πλειων, 'more,' and εχειν, 'to have,') Covetous. Desirous to acquire: hence *ma'nia pleonectica*.

PLEONECTICA ATHYMIA seu MELANCHOLIA. *Pleonexia*. A form of insanity characterised by a desire for gain or by covetousness.

PLEROSIS, (πληρωσις, from πληρης, 'full,' and οσις,) *Repletio*, *Refectio*, *Repletio*, (F.) *Plerose*. Restoration of flesh after sickness. Plethora.

PLEROTICUS, (πλερωτικός, from πληρωω, 'to fill up,') Incarnans.

PLÉROTIQUE, Incarnans.

PLESMONÉ, (πλησμονη,) *Plenitudo*, *Satietas*, *Satiety*. Intemperance. A condition the opposite to that of hunger.

PLESSER, (πλησσω, 'I strike,') Plexor.

PLESSIMETER, Pleximeter,

PLESSIMÈTRE, Pleximeter.

PLETHOMERIA, (πληθος, 'a great number,' and μερος, 'a part,') Superabundance of parts, as in the case of six fingers to one hand.

PLETHORA, (πληθωρα,) *Polyæmia*, *Polyhyperæmia*, *Panhypæmia*, *Polychymia sanguinea*, *Angiopleurosis*, *Angiopleurosis*, *Erythræ-*

nia, *Erythro'sis*, *Hæmatopletho'ra*, *Hæmopletho'ra*, *Plenitudo*, *Multitudo*, *Redundantia*, *Repletio*, *Enchyma*, *Completio*, *Plethory*, *Repletio*, (Old Eng.) *Plow'sy*, *Sanguineness*, *Fullness*, (F.) *Abundance*, *Réplondance*. The word *plethora*, which means repletion, *Pletho'sis*, expresses a superabundance of blood in the system, or in some part of it:—hence the division of *plethora* into *general* and *local*; the latter being called, according to its seat, *cerebral*, *pulmonary*, *uterine*, &c. The principal symptoms of *plethora* exist in the circulatory system; such as redness of the surface, swelling of the veins, increase in the fullness of the pulse, in the strength of the heart's pulsations, &c., with spontaneous hemorrhages occasionally. With these are usually united general heaviness, torpor, lassitude, vertigo, tinnitus aurium, flushes of heat, &c. The blood of plethoric persons differs from healthy blood in the smaller ratio of water which it contains, and especially in the augmentation of the red corpuscles.

The tendency to *plethora*, or its actual existence, must be obviated by purgatives, proper diet, exercise, &c.; and, if the danger from it be pressing, by bloodletting. This is, however, a doubtful remedy for general *plethora*, as it increases hæmotosis.

PLETHORA ARTERIO'SA is the form in which the blood is rich in fibrin and red corpuscles—the *arterial constitution*.

PLETHORA, LOCAL, Hyperæmia—p. Partialis, Hyperæmia.

PLETHORIC, (πληθωρικός,) *Plethoricus*, *Polyæmic*, *Polyæmicus*, *Polyæmatus*, *Sanguine*, *Sanguineus*, (F.) *Pléthorique*. Full of blood. Relating to or affected with *plethora*.

PLETHORICUS, *Plethoric*.

PLETHORY, *Plethora*.

PLETHOS, (πληθος,) *Satiety*.

PLEUMON, (πλευμων,) *Pulmo*.

PLEUMONIA, *Pneumonia*.

PLEURA, (πλευρα,) *Pleurum*, *Pleuro'ma*. *Membrana pleurica succin'gens seu cost'as succin'gens*, *M. seu Tunica costalis seu subcostalis*, *Hypopleurios*, *Hypezocus*, (F.) *Plevre* on *Pleure*. The pleurae are two thin, diaphanous, perspirable membranes, which line each side of the chest, and are reflected thence upon each lung. Like other serous membranes—to which class they belong—each represents a sac without aperture. From the junction of the pleura of each side the mediastina are formed. That portion of the pleura which lines the parietes of the chest is called *Pleura Costalis* seu *parietalis*; the portion that covers the lungs, *Pleura Pulmonalis*. The arteries of the pleurae are from the intercostals, internal mammaries, phrenic, inferior thyroid, &c. The veins correspond with the arteries. They are, also, supplied with lymphatics, but nerves are not traceable in them.

Pleura is also used for *Cavity of the Pleura*, *Sac'cus Pleuræ*.

PLEURÆ, (pl. of *pleura*.) The sides of the chest.

PLEURAL, *Pleuralis*, *Pleuricus*, *Pleurites*. That which concerns the pleura;—as a *pleural fistula*, *pleural cavity*, *pleural friction*, *pleural hemorrhage*, &c.

PLEURALGIA, (πλευρα, 'side,' and αλγος, 'pain,') *Pleurodyne*.

PLEURAPOPHYSES, (πλευρα, 'side,' and ἀποφυσες, 'processes,') *Costæ veræ*, see *Costa*.

PLEURAPOSTEMA, (*pleura*, and *apostema*, 'abscess,') *Empyema*.

PLEURARTHRO'ACE, (πλευρα, 'a rib,' and ἄρθρον, 'a joint,' and κακος, 'bad,') *Caries of the ribs*.

PLEURE, *Pleura*.

PLEURESIA, (from *pleura*, *Pleuritis*).

PLEURÉSIE, *Pleurisy*—p. *Fausse*, *Pleurodynia*.

PLEURESIS, (from *pleura*.) *Pleuritis*.

PLEURÉTIQUE, *Pleuritic*.

PLEURICUS, *Pleural*.

PLEURISY, *Pleuritis*.

PLEURISY, BILIOUS, *Pleuritis biliosa*. *Pleurisy* accompanied with bilious symptoms;—the effect of duodenitis or duodenohepatitis.

PLEURISY, FALSE, *Pleurodynia*.

PLEURISY, HEAD. A ridiculous term, occasionally used by the vulgar, in the Southern States especially, for bilious pneumonia before the pneumonic phenomena are developed, and whilst the head is prominently affected.

PLEURISY, HEMORRHAGIC, *Hæmatothorax*—p. *Rheumatic*, *Pleurodynia*—p. *Root*, *Asclepias tuberosa*.

PLEURISY, TY'PHOID, *Pleuritis typho'ides* seu *typho'ides* seu *typho'sa*. *Pleurisy* accompanied by adynamic symptoms.

PLEURITES, (πλευριτης,) *Pleural*.

PLEURITIC, (πλευριτικός,) *Pleuriticus*, *Pleuritical*, (F.) *Pleurétique*; (same etymon as *pleuritis*.) Relating or appertaining to *pleuritis*,—as '*pleuritic pain*'.

PLEURITIS, gen. *Pleuritidis*, (ι, *πleuritis*, from *pleura*, and *itis*, or from *πλευριτης*, (νόσος, 'a disease,' understood,) *Morbus lateralis* seu *pleurit'icus* seu *costalis*, *Empres'ma* seu *Cau'ma* *Pleuritis*, *Pleur'e'sis*, *Pleur'e'ia*, *Pleuritis costalis* seu *ve'ra*, *Pleurit'ica Fe'bris*, *Pas'sio pleurit'ica*, *Inflammat'io Pleure*, *Pleur'isy*, *Inflammat'ion of the Pleu'ra*, (F.) *Pleurésie*, *Fièvre pleurétique*, *Inflammation de la Plevre*, may be acute or chronic. Perhaps the most violent of all internal inflammations is the active form; known by acute, lancinating pain in one side of the chest, increased by inspiration, by coughing, and often by pressure; dyspnoea, the inspirations short and frequent; cough dry, or with little expectoration; and difficulty of lying on the affected side; symptoms, attending the phlegmasia in general, strongly marked.

Physical Signs. There is seldom any perceptible alteration in the exterior of the chest. When effusion has taken place to a great extent, the signs will be similar to those of chronic pleurisy. The motion of the affected side is diminished, and the vibrations of the voice cannot be detected by the hand. There is more or less dulness of sound on percussion, the dulness diminishing or disappearing by change of position. On auscultation, the inspiration is feeble, distant, or inaudible; but change of position modifies it. Egophony is usually present, when the effusion is in moderate quantity, and is best heard over a portion of the thoracic parietes, represented by a band of three inches, running from below the inferior margins of the scapula in the direction of the ribs to the sternum. Simple pleurisy rarely attacks both sides at once: when such is the case, the presence of tubercles may be suspected.

Formerly, a *true pleurisy* meant one seated in the pleura: a *mediastinal pleurisy*, one situated in the anterior mediastinum; and *false pleurisy* or *rheumatism'al pleurisy*, that which occupies the intercostal muscles. The last has received the name *Pleurodynia*; that of *pleurisy* being exclusively appropriated to inflammation of the pleura. Pleurisy has also been distinguished into *dry* and *humid*, according as it is or is not accompanied by expectoration. *Humid* or *moist pleurisy* is now properly regarded as a complication of pleurisy with pulmonary catarrh. It is the *catarrhal pleurisy* of some. The majority of the species of pleurisy, admitted by Sauvages, consist of complications.

The causes of pleurisy are like those of other internal inflammations. It may terminate by resolution, effusion, or suppuration.

CHRONIC PLEURISY, which has been particularly described by recent pathologists, most commonly terminates by a serous or purulent effusion, which sometimes simulates hydrothorax; at others, phthisis pulmonalis. It may form slowly, without evident symptoms; or it may succeed acute pleurisy. In the former case, vague pains are felt in the chest, with small, dry cough, oppression at intervals, shivering, irregular febrile symptoms, and hardness of pulse.

Physical Signs. The affected side is smoother, more rounded and motionless; the intercostal spaces are dilated and filled up, or may even protrude slightly. In very chronic cases, however, the affected side may be smaller. The triangular space above the clavicle and the depression immediately above the sternum are often drawn downwards on the diseased side. No vibration is experienced by the hand when the patient speaks. The parietes of the thorax are sometimes cedematous, and fluctuation may be occasionally felt between the ribs. On percussion the sound is dull, or if clear, only so in the upper portions of the chest. On auscultation, there is an absence of the usual sounds over the affected parts, excepting occasionally transmitted sounds. There are no physical signs that can enable us to know whether pus or serum is effused into the chest. The introduction of a grooved needle through the intercostal space, as recommended by Dr. Thomas Davies, would of course settle the question.

On dissection of those who have died of pleurisy, the pleura is found thickened, red, inflamed, and covered with membranous exudations or false membranes. Sometimes it seems cartilaginous and ossified. Its cavity frequently contains serous or seropurulent effusions.

The acute form of the disease requires most active treatment. General bloodletting, to such an extent as to make a very decided impression on the system, and repeated if necessary. This is the most important agent. After the violence of the disease has been got over, counterirritants will be found highly beneficial; as well as when it has passed into the chronic state. The cough may be relieved by demulcents.

Sauvages has given the name **PLEURITIS VENTOSA**, and Pringle that of **PLEURODYNIA VENTOSA**, to a pain behind the false ribs, attributed to the presence of air in the intestines.

PLEURITIS BILIOSA, Pleurisy, bilious.—p. Bronchialis, Bronchitis.—p. Costalis, Pleuritis.—p. Diaphragmatica, Diaphragmatitis.—p. Humida, Bronchitis.—p. Muscularis, Pleurodynia.—p. Notha, Pleurodynia.—p. Pericardii, Pericarditis.—p. Pulmonis, Pleuropneumonia.—p. Rheumatica, Pleurodynia.—p. Spuria, Pleurodynia.—p. Spuria simulata, Pleurodynia.—p. Typhoides, Pleurisy, typhoid.—p. Vera, Pleuritis.

PLEUROCELE, (*pleura*, and *κηλη*, 'a tumour.')

Hernia of the pleura, *Hernia pleurica et pulmonalis*. This name, employed by Sagar, is inaccurate, as the pleura never protrudes alone. It only forms a hernia, when it serves as an envelope to the lung, to tumours, or to purulent collections, which have protruded from the thoracic parietes.

PLEUROCOLLE/SIS, (*pleura*, and *κολληω*, 'I glue.')

An adhesion between the pleura costalis and pleura pulmonalis.

PLEURODYNE, Pleurodynia.

PLEURODYNIA, *Pleuralgia*, *Pseudopleuritis*, *Pseudopleurisy*, *Pleurodynia*, *Auralgia colorosa*, *Parapleuritis*, *Stethorrhœum*, *Rheumatis-mus pectoris*, *Pleuritis rheumatica seu muscularis seu notha seu epurica seu epurica simulata*, *Pneumonia externa*, *Lagomyonosis*, *Thoracodynia*, *Thoracodynia*, *Thoracomyodynia*, *Dolior Pectoris externa*, Pain in the side, *Stitch in the side*, *False, rheumatic or rheumatismal pleurisy*, (Old

Eng.) *Shott*, (Sc.) *Caik*, (F.) *Point de côté*, *Douleur de côté*, *Pleurésie faussée*; (*πλευρα*, 'side,' and *δύνη*, 'pain.')

A spasmodic or rheumatic affection, generally seated in the muscles of the chest, and, ordinarily, in the intercostals. It is usually augmented by outward pressure, inspiration, coughing, exertion, moving the thorax, and sometimes even the corresponding arm. It is unaccompanied by fever, cough, or dyspnoea, excepting under the form of a *stitch*, when dyspnoea exists. It is, generally, by no means obstinate, and yields promptly to warm applications and to cupping or leeches. If it still resists, a blister will, almost always, remove it.

PLEURODYNIA VENTOSA, Pleuritis ventosa.

PLEURODYNIC, *Pleurodynical*, *Pleurodynicus*, (F.) *Pleurodyrique*. Relating or belonging to pleurodynia.

PLEUROMA, gen. *Pleuro'matis*, (*πλευρωμα*, from *pleura*, and *ομα*,) Costa, Pleura.

PLEUROPATHIA, (*pleura*, and *παθος*, 'a disease.')

A disease of the pleura.

PLEUROPERIPNEUMONY, *Pleuroperipneumonia*, *Pleuropneumonia*, *Pneumopleuritis*, *Pneumopleuritis*, *Pleuritis pulmonis*, *Pneumonia pleuritica*, (*pleura*, and *peripneumonia*.)

Inflammation occupying the pleura and the lung at the same time. Pleurisy and peripneumony existing simultaneously. **Physical signs.** A dull sound is yielded by percussion for a variable extent over the more dependent portions of the affected side. The loss of sound is partially modified by change of position. The respiratory murmur on auscultation is very feeble or absent inferiorly; higher up, the crepitation or bronchial respiration of pneumonia may be detected; round the root of the lung, and near the inferior angle of the scapula, egophony is generally present. There is seldom much enlargement of the affected side, or displacement of the adjacent organs, as in the case of simple chronic pleurisy.

PLEUROPNEUMA, gen. *Pleuropneum'atis*, (*pleura*, and *πνευμα*, 'wind,')

Pneumothorax.

PLEUROPNEUMONIA, Pleuroperipneumony.

PLEUROPNEUMONITIS, Pleuroperipneumony.

PLEUROPYESIS, (*pleura*, and *πυσις*, 'suppuration,')

Empyema.

PLEURORRHAGIA, (*pleura*, and *ῥαγη*, 'a breaking forth,')

Hæmatorrhax.

PLEURORRHŒA, (*pleura*, and *ῥεω*, 'I flow.')

Accumulation of fluid in the pleura.

PLEURORRHŒA CHYLOSA, Chylothorax.—p. Lymphatica, Hydrothorax.—p. Purulenta, Empyema.—p. Sanguinea, Hæmatorrhax.—p. Serosa, Hydrothorax.

PLEURORTHOPNŒA, (*pleura*, *ὀρθος*, 'erect,' and *πνεω*, 'I respire.')

Pain of the side, which does not permit the patient to breathe, except when in the vertical position.

PLEUROSOMUS, (*pleura*, and *σωμα*, 'body.')

A malformation in which the fissure is somewhat lateral, with eventration extending chiefly upon the upper part of the abdomen and upon the chest; the upper extremity of the fissured side being more or less atrophied.

PLEUROSTOSIS, (*pleura*, *ὀστέον*, 'a bone,' and *οσις*.)

Ossification of the pleura, or in the cavity of the pleura. *Osteopleurica*.

PLEUROTETANUS, (*πλευρα*, 'a side,' and *tetanus*,) Pleurothotonus.

PLEUROTHOTONUS, *Pleurotonus*, *Pleurotel'anus*, *Tel'anus lateralis*, from *πλευροσος*, 'laterally,' and *τονος*, 'tension.')

A variety of tetanus, in which the body is curved laterally by the stronger contraction of the muscles of one side of the body.

PLEUROTONUS, (*πλευρα*, 'a side,' and *τονω*, 'to stretch,')

Pleurothotonos.

PLEURUM, (*πλευρον*,) Costa, Pleura.

PLÈVRE, Pleura.

PLEXIM'ETER, *Plexom'eter*, *Plegon'eter*, *Plessim'eter*, *Anticoposcop'ium*, *Anticop'oscope*, (F.) *Plessimètre*, (πλησσω, 'I strike,' and μετρον, 'measure,' 'a measure of percussion.') An ivory plate of a circular or ovoid shape, from an inch and a half to two inches in diameter, and about one-sixth of an inch in thickness. It has either a raised rim or edge, or projecting handles on its upper side, to permit its being held between the finger and thumb of the left hand, whilst it is struck with the right. It is used in percussion of the chest. A piece of metal, a coin, or the finger of the left hand may be used with equal advantage.—See Percussion, mediate.

PLEXOMETER, Pleximeter.

PLEX'OR, (πλησσω, 'I strike.') *Ples'ser*. Any instrument employed in percussion. The ends of the fingers of the right hand brought together are the best *plexor*; as the finger of the left is, perhaps, the best *pleximeter*.

PLEX'US, gen. *Plexūs*, *Pleg'ma*, *Plec'tanē*, *Pla'nē*, *Re'tē*, (πλεω, *plecto*, *plexum*, 'to interlace,' 'to entwine,') (F.) *Lacis*. A network of blood-vessels or of nerves. The nervous plexuses, *Implicatio'nes reticula'res* seu *Tex'tus nervo'rum* belong—some to the system of encephalic nerves—others to that of the great sympathetic; whilst some, as the pharyngeal, seem to be formed of the two sets. The plexuses represent complex networks, with more or less loose meshes, formed by the numerous and diversified anastomoses of the nervous filament; from which proceed other branches, that are distributed to organs or to other plexuses.

PLEXUS BRACHIALIS, Brachial plexus.—p. Cardiac, Cardiac plexus.—p. Carotid, see Carotid nerve.—p. Cavernosus, Bulbus vestibuli.—p. Choroidea, see Choroïd.—p. Choroidea Medius, Choroidea tela.—p. Ciliaris, Ciliary ligament.—p. *Cœliacque*, *Cœliac plexus*—p. Coronariis ventriculi, Gastric plexus.—p. Coronary, see Cardiac plexus, and Coronary plexus.—p. Gangliiformis semilunaris, *Cœliac plexus*—p. Gangliiformes, Ganglions, nervous.—p. Glandiformes, Ganglions, nervous.—p. Glandulosi Peyer, Peyer's glandulæ.—p. Hederaceus, Corpus pampiniforme.—p. Hepaticus, Hepatic plexus.—p. *Hypogastrique*, Hypogastric plexus.—p. Infrathyroidal, see Thyroidal veins.—p. *Lombo-abdominal*, Lumbar plexus.—p. Median, *Cœliac plexus*—p. Mesenterii proprius et maximus, *Cœliac plexus*—p. Nervorum anserinus, Facial nerve.—p. *Opiotogastrique*, *Cœliac plexus*—p. Pampiniformis, Corpus pampiniforme.—p. Pharyngeal, see Pharyngeal nerve.—p. Pneumogastrius, see Pneumogastic nerve.—p. Reticularis, Bulbus vestibuli, Choroidea plexus.—p. Retiformis, Corpus cavernosum vaginæ.—p. Solar, *Cœliac plexus*—p. *Sous-mésentérique*, Hypogastric plexus.—p. *Testiculaire*, Spermatic plexus.—p. *Trachélosous-cutané*, Cervical plexus.—p. Vasculosus funiculi spermatici pampiniformis, Corpus pampiniforme.

PLI (F.), ([L.] *plica*, 'a fold,')—p. *du Coude*, 'fold of the elbow,' see Elbow.—p. *de Douglas*, 'fold of Douglas,' see Uterus.—p. *de Vater*, 'fold of Vater,' *Plica longitudinalis Vateri*.

PLICA, (*plico*, πλεω, 'to fold,') *Tricho'sis pli'ca*, *Tricho'ma*, *Trichono'sis pli'ca*, *Tri'cæ*, *T. Incubo'rum* seu *Serofo'rum*, *Plica Polon'ica* seu *Polon'ica Juda'ica* seu *Cachec'tica* seu *Saxon'ica* seu *ca'put Medu'sæ* seu *longica'u'da latera'tis*, *Affectio Sarmat'ica*, *Lw'es Sarmat'ica* seu *Polon'ica* seu *Trichomat'ica*, *Helo'sis*, *Helo'tis*, *Cir'ragra*, *Cir'rakra*, *C. Polono'rum*, *Ecphy'ma tricho'ma*, *Mor'bus Cirrho'rum*, *Co'ma Cesare'a*, *Kol'to*, *Rhopal'o'sis*, *Plicat'io*, *Plicatu'ra*, *Mott'ed hair*, *Plai'ed hair*, *Trichom'atose hair*, (F.) *Plique*, *P. Polonaise*. A disease endemic in Poland, Lithuania, and other parts of Northern Europe; so

called on account of its being characterized by interlacing, twisting, and agglutination or matting of the hair. By some it has been regarded as a real disease; by others, as the want of attention to cleanliness. However this may be, it generally appears upon the hair of the head, but sometimes on that of other parts, as the beard, the hair on the axilla, pubes, &c. Alibert admits three species of *plica*. 1. *Plica multiform'is*, or *Plica ca'put Medu'sæ*, in which the hairs are mixed and agglutinated in greater or less masses; and this has been again subdivided into two varieties, according as the meshes are almost straight (*P. C. M. lacinia'ta*, [F.] *Plique en lanières*), or much twisted ([F.] *P. en vrilles*). 2. *Plica longica'u'da*, (F.) *Plique solitaire ou à queue*; when the hair is united into a single long mass, and 3. *Plica ces-pito'sa*, (F.) *Plique en masse*, in which the hairs are matted into one large, shapeless mass. Pinel places this disease amongst the *Cutaneous phlegmasiæ*; but it is far from demonstrated, that it is the result of inflammation of the skin. Certain it is, however, that the tangling of the hair is symptomatic of an affection—*sui generis*—of the dermoid system. The idea that it is entirely owing to want of cleanliness, as has been conceived by some, is not tenable. It has been supposed to be produced by a fungous growth—a *mycoder'ma*, *trichomaphyte* or *trichophyte*.

PLICA, 'a fold,' *Plectane*—p. *Cachectica*, *Plica*—p. *Caput Medusæ*, *Plica*—p. *Centralis retinæ*, see *Optic nerve*—p. *Longicauda lateralis*, *Plica*—p. *Lunata*, *Valvula semilunaris*—p. *Polonica*, *Plica*—p. *Polonica Judaica*, *Plica*—p. *Saxonica*, *Plica*—p. *Semilunaris*, *Valvula semilunaris*.

PLICE CONNIVENTES, (pl. of *plica*,) *Valvulæ conniventes*.

PLICA LONGITUDINA'LIS VATERI, *Divertic'ulum Vateri*, (F.) *Plî de Vater*. A vertical projection, about an inch in length, formed at the posterior part of the duodenum by the mucous membrane raised up by the common choledoch duct before it opens into the intestine.

PLICE SEMILUNARES, see *Urinary bladder*, and *Uterus*.

PLICATIO, *Plica*.

PLICATURA, *Plica*.

PLICATURA CEREBRI, *Gyri cerebri*.

PLICHOS, (πλῆκος,) *Perinæum*.

PLIN'THIUM, (πλινθιον.) A machine invented by one Nileus, which was formerly employed in the reduction of fractures and luxations. Scultetus describes it in his *Armamentarium Chirur-gicum*.

PLIQUE, *Plica*—p. *Polonaise*, *Plica*.

PLÔCARIA CANDIDA, *Fucus amylaceus*—p. *Helminthochorton*, *Corallina Corsicana*.

PLOESSLEA FLORIBUNDA, (after Ploessl, an optician of Vienna,) see *Juniperus lycia*.

PLOMB, *Plumbum*. Also, the sulphuretted gas disengaged from privies, which proves fatal at times to the nightmen, (F.) *Vidangeurs*, engaged in emptying them.

PLOMB, ACÉTATE ACIDE DE, *Plumbi superacetas*—p. *Blanc*, *Plumbi subcarbonas*—p. *Blanc* de, *Plumbi subcarbonas*—p. *Carbonate* de, *Plumbi subcarbonas*—p. *Chlorure* de, *Plumbi chloridum*—p. *Iodure* de, *Plumbi iodidum*—p. *Nitrate* de, *Plumbi nitras*—p. *Oxide* de, *fondus*, *Plumbi oxydum semivitreum*—p. *Oxide* rouge de, *Plumbi oxydum rubrum*—p. *Oxide*, *semi-vitreux* de, *Plumbi oxydum semivitreum*.

PLOMBAGE DES DENTS, (from *plomb*, 'lead,') *Plugging*.

PLOMBAGINE, ('*plomb*, 'lead,') *Graphites*.

PLOMBIÈRES, MINERAL WATERS OF. *Fon'tes medica'ti Plumbi'ri*, *Ther'mæ Plumbi'riæ*. *Plombières* is a town in the department of the Vosges, 17 leagues from Nancy, and two from

Remiremont. Here are several sources of thermal water. That which is called *Sappon'ceus*, (F.) *Savonneuse*, is formed of the subcarbonate and sulphate of soda, chloride of sodium, silica, carbonate of lime, and much vegetable matter. This last component, (F.) *Plombiérine*, when held in solution by means of the subcarbonate of soda, gives to the water an unctuous character. This is the origin of its name. The waters of Plombières are employed as solvents in engorgements of the abdominal viscera, and as alteratives in chronic diseases of the skin. Season, from the 15th of May to the 15th of October.

PLOMBIÉRINE, see Plombières.

PLORATIO, (*plorare*, *ploratum*, 'to weep,') *La crymatio*.

PLORATUS, *La crymatio*.

PLOTUS, 'flat,' *Leipous*.

PLOUK, (Gael. *plucan*,) *Papula*.

PLUG'GING, (from [D.] *plug*,) (F.) *Tamponnement*. The act of stopping a hole. The introduction of a plug or dossil of lint or rag—(F.) *Tampon*—into a wound or natural cavity—as the uterus or vagina—to arrest hemorrhage; or of some substance into a carious tooth to prevent toothache, (F.) *Plombage* ou *Obturation des dents*. See *Filing*.

PLUKE, (Gael. *plucan*,) *Papula*.

PLUM, (Sax. *plume*,) *Prunum*, *Prunus domestica*—p. Assyrian, *Sebestina*.

PLUMA, 'down,' *Lanugo*.

PLUMACEAU, (from *pluma*,) *Pledget*.

PLUMACEOLUS, (from *pluma*,) *Pad*, *Pledget*.

PLUMASSEAU, *Pledget*.

PLUMBAGO, (from *plumbum*, the leaves having lead-coloured spots,) *Graphites*, *Pescicaria*.

PLUMBA'GO *EUROPÆA*, *P. lapathifolia* seu *undulata*, *Dentaria*, *Orepanella*, *Squama'ria*, *Dentel'ria*, *Leul'wort*, (F.) *Dentel'rie*, *Herbe au Cancer*, *Malherbe*. Ord. *Plumbaginaceæ*. Sex. Syst. *Pentandria Monogynia*. The root was formerly much esteemed as a cure for toothache, like the *Pellitory of Spain*, which, also, is called *Dentaria*.

PLUMBI ACETAS, *P. superacetas*—p. *Acetas dilutum alcoholicum*, *Liquor plumbi subacetatis dilutus*—p. *Acidulus siccus*, *Plumbi superacetas*—p. *Azotas*, *P. nitras*—p. *Carbonas*, *P. subcarbonas*. *PLUM'BI CHLO'RIDUM*, *Chlo'ride of Lead*, (F.) *Chlorure de Plomb*. This salt of lead has been used in the form of lotion and ointment in cancerous ulcerations.

PLUMBI DEUTOXYDUM RUBRUM, *P. Oxydum rubrum*—p. *Hydriodas*, *P. Iodidum*—p. *Iodhyd'ras*, *P. Iodidum*.

PLUM'BI IO'DIDUM, *P. Iodure'tum* seu *Hydriodas* seu *Hydroi'odas* seu *Iodhy'dras*, *Plum'bium ioda'tum* seu *Hydroi'od'icum*, *Iodide* or *Iod'uret of Lead*, (F.) *Iodure de Plomb*, *I. plumbique*. (This salt is formed by the double decomposition of iodide of potassium and nitrate of lead.) It is used in scrofulous affections internally and externally; and is official in the *Pharmacopœia U. S.* Dose, gr. ss. to gr. iij or iv.

PLUMBI IODURETUM, *P. Iodidum*.

PLUMBI NI'TRAS, *P. Azo'tas*, *Plumbum ni'tricum*, *Ni'tras plumb'icus*, *Ni'trum saturni'num*, *Ni'trate of lead*, (F.) *Nitrate ou Azotate de Plomb*. (A salt formed by the action of diluted nitric acid on litharge.) Nitrate of lead has the same medical action as the other soluble salts of lead, and is a valuable antihemic. It has been used, also, as an excitant antiseptic to wounds, ulcers, &c., and is the basis of Ledoyen's disinfecting liquid.

PLUMBI NITROSAC'CHARAS, *Nitrosac'charate of lead*. (*Suechr.* p. j; *Acid. nitric.* p. ij; *Aquæ* p. x. Beat together in a porcelain dish as long as chemical action continues: dilute the liquor with water; neutralize with chalk, and to the fil-

tered solution add acetate of lead so long as a precipitate is formed. Collect, wash, and dry the precipitate, which is saccharate of lead. Dissolve in dilute nitric acid, (one part acid to nineteen of water,) filter and evaporate to form crystals, which are transparent, of an amber colour, and in regular hexagonal plates or prisms.)

Advised as a solvent of phosphatic calculi.

PLUMBI OXYDUM seu *OXYDUM ALBUM*, *Plumbi subcarbonas*—p. *Oxydum carbonatum per acidum acetosum*, *Plumbi subcarbonas*—p. *Oxydum fuscum*, *Plumbi oxydum semivitreum*.

PLUMBI OXYDUM RUBRUM, *Min'ium*, *Mill'us*, *Dentox'ydum plumb'i rubrum*, *Superox'ydum Plumb'ium*, *Plumb'ium oxyda'tum rubrum* seu *superoxyda'tum*, *San'dix*, *San'dyx*, *Red lead*, (F.) *Oxide de plomb rouge*. It is sometimes used to kill vermin.

PLUMBI OXYDUM SEMIVIT'REUM, *P. Ox'idum* (Ph. U. S. 1863), *Plumbum oxyda'tum*, *Lithar'gyrus*, *Lithar'gyrum*, *Litharag'um*, *Ox'ydum plumb'i fuscum*, *Spu'ma argenti*, *Semivitrified ox'yd of lead* or *lith'arge*, *Calcite'os*, *Cath'mia*, (F.) *Oxide de plomb fondu*, *Oxide de plomb semivitreux*, (a yellow protoxyd of lead, prepared by heat, combined with carbonic acid.) It occurs in scales of a whitish red colour, and semivitrified. It is employed for pharmaceutical purposes, as in the *Liquor plumbi acetatis*, *Emplastrum plumbi*, &c. Formerly, the *Litharge of gold*, *Chrys'itis*, *Mas'sicot*, and *Litharge of silver*, *Almariac*, were distinguished in commerce. These were merely ordinary litharge, resembling gold or silver in colour.

PLUMBI SUBCARBO'NAS, *Carbo'nas plumb'i* seu *plumb'icus*, *Plumbi Carbo'nas* (Ph. U. S.), *Plumbum carbo'nicum*, *P. oxyda'tum carbo'nicum*, *Magnesium plumbi*, *Oxydum plumbi al'bum*, *Cerus'sa*, *Aeru'go plumbi*, *Simmit'ium*, *Paimy'thion*, *Ps'my'thus*, *Blan'ca*, *Ger'sa*, *Iff'ides*, *Subcarbo'nate of lead*, *Cerussa*, *White lead*, *Flake white*, (F.) *Carbonate de plomb*, *Blanc de plomb*, *Plomb blanc*, *Céruse*, *Blanc d'Argent*, *Blanc de Céruse*, *Blanc de Krems*, *B. de Krems*, *B. de Kremnitz*, *Craie de plomb*, (prepared by exposing spiral rolls of lead to the fumes of vinegar, in vessels placed in dung.) It is inodorous; has a sweet taste; is brittle; friable; snow white, and of a minute, scaly texture. It is astringent and sedative, and is sprinkled on parts affected with local inflammation, or to prevent chafing in children. It is also used in the formation of ointments and plasters.

PLUMBI SUPRACET'AS, *Cerus'sa aceta'ta*, *Plumbi acid'ulus sic'cus* *P. ac'etas* (Ph. U. S.), *Sal* seu *Sac'charum Saturni*, *Plumb'um ac'eticum*, *Dulce'do Saturni*, *Ac'etas plumb'icum crystallina'tum*, *Sai plumbi*, *Sugar of lead*, *Supracet'ate of lead*, *Protoac'etate of lead*, (F.) *Acétate acide de plomb*, *Sel ou sucre de Saturne*. (*Plumb. carb. lbj*, *Acid acetic. cong. lʒ*. Boil the lead with the acid till it is saturated; then strain through paper, and evaporate to a pellicle that crystals may be formed. Wash the crystals and dry on blotting-paper. Ph. L. In the Ph. B. (1864), it is directed to be prepared by the action of dilute acetic acid on litharge; filtering, evaporating, and crystallizing.) Acetate of lead has a sweet, styptic taste; a very white colour, and silky lustre. It is astringent; and, in weak solution, cooling and sedative; in strong, stimulant. It is given internally, in visceral and other hemorrhages, combined with opium, and is used externally, in solution, in inflammation, burns, bruises, gonorrhœa, &c. Dose, gr. ss to iss.

PLUMBI TANNAS, see *Tannin*.

PLUMBICUM ACETAS CRYSTALLISATUM, *Plumbi superacetas*.

PLUMBOSUM SUPEROXYDUM, *Plumbi oxydum rubrum*.

PLUMBUM. *Malib'los, Malib'idus, Saturnus, Capricornus, (F.) Plumb.* Lead. A solid metal of a bluish white colour; more malleable than ductile; soft enough to be scratched by the nail; very fusible and oxidizable, when heated in the air, and existing, in nature, under a number of different shapes; the principal of which are those of sulphuret and oxyd. S. g. 11.352.

Its use is to form oxides and salts. In its metallic state, it has no action upon the body; but when oxydized or united with an acid, it acts, in large doses, as a powerfully astringent poison.

PLUMBUM ACETICUM, Plumbi superacetat.—p. Album, Tin—p. Candidum, Tin—p. Carbonicum. Plumbi subcarbonas.—p. Cinereum, Bismuth, subnitrate of.—p. Hydriodicum, Plumbi Iodidum—p. Iodatum, Plumbi Iodidum—p. Nigrum, Graphites—p. Nitricum, Plumbi nitras.—p. Oxydatum, Plumbi oxydum semivitreum.—p. Oxydatum carbonicum, Plumbi subcarbonas.—p. Oxydatum rubrum, Plumbi oxydum rubrum—p. Superoxydatum, Plumbi oxydum semivitreum.

PLUMIERA ALBA, (after Plumier, a French botanist;) *Order, Apocynaceæ*; a native of tropical America, has a milky juice which is cathartic.

PLUMIERA DRASTICA, is given in Brazil in jaundice, and chronic obstructions, mixed with milk of almonds.

PLUMIERA PHAGEDENICA, is in repute in Brazil as an anthelmintic.

PLUMS, SEEDED, see Diospyros Virginiana—p. Winter, see Diospyros Virginiana—p. Yellow, see Diospyros Virginiana.

PLUNGE BATH, see Bath.

PLUNGING SIPHON, see Siphon.

PLUNKETT'S CAUSTIC FOR CANCERS, Plunkett's Ointment.

PLUNKETT'S OINTMENT, Plunkett's Caustic for Can'cers. This empirical cancer remedy consists of *Arsenious acid, Sulphur,* and the powdered flowers of the *Ranunculus flammula* and *Cotula foetida*;—levigated and made into a paste with the white of egg, and applied, on a piece of pig's bladder, to the surface of the cancer.

PLURAL BIRTH, see Multiparous.

PLURIMAMME, (*plures*, 'many,' and *mamma*, 'breasts,') Polymasthus.

PLURISY, (from *plus*, gen. *pluris*, 'more,') Plethora.

PLUTEA. Aricenna gives this name to duplicatures of the dura mater, such as the *Falx cerebri*.

PLUTONIUM, (after *Pluto*,) Barium.

PNEOBIO MANTIA, (*πνευ*, 'to breathe,' *βίος*, 'life,' and *μάντις*, 'divination,') Docimasia pulmonum.

PNEOBIO MANTICA, Docimasia pulmonum.

PNEOMETER, (*πνευ*, 'to breathe,' and *μετρον*, 'measure,') Spirometer.

PNEUMA, gen. *Pneumatis*, (*πνευμα*, gen. *πνευματος*,) Air, Life. See Pneumatic Physicians.

PNEUMAPOSTEMA, gen. *Pneumapostematis*, *Pneumapostema*, *Abcessus pulmonum*, (*pneumon*, and *αποστημα*, 'abscess,') Abscess of the lung.

PNEUMARTHROSIS, (*pneuma*, *αρθρον*, 'articulation,' and *osis*,) Secretion of air into an articular cavity. Seen at times, it is said, after acute rheumatism, but soon disappearing [?].

PNEUMATELECTASIS, (*pneumon*, and *atelectasis*,) Atelectasis pulmonum.

PNEUMATHEMIA, *Pneumatemia*, (*pneuma*, and *ημα*, 'blood,') A morbid condition in which air is developed in the bloodvessels.

PNEUMATHORAX, Pneumothorax.

PNEUMATIC, (*πνευματικός*,) *Pneumaticus*, *Pneumaticeus*, (F.) *Pneumatique*. Relating to air or life.

PNEUMATIC MEDICINE, Atmiatrica.

PNEUMATIC PHYSICIANS, *Pneumatici*, *Pneumatists*. Name given to a sect of physicians, at the head of whom was Athenæus, who made health and disease to consist in the different proportions of an element—which they called *Pneuma*, (*πνευμα*)—to those of the other elementary principles. This *πνευμα* was a fancied spiritual principle, which the Stoics regarded as an element, different from water, air, earth and fire;—the sole elementary principles previously admitted. The doctrine was called *Pneumatism*. At the present day, the epithet *pneumatic* is sometimes applied to those who treat diseases chiefly by the inhalation of gases.

PNEUMATICA, (from *pneuma*,) Appertaining to the breathing. Diseases of the respiratory function;—the second class in Good's system of nosology. Agents that act on the respiratory organs.

PNEUMATICAL, Pneumatic.

PNEUMATOCELE, (*πνευματοκύλη*, from *pneuma*, and *κύλη*, 'a tumour,') *Physocèle*.

PNEUMATODES, (*πνευματώδης*, from *pneuma*, and *οδός*,) One distended with air, or who breathes with difficulty, owing to an accumulation of flatus in the digestive tube; or, according to others, owing to emphysema.

PNEUMATOMETER, (*pneuma*, and *μετρον*, 'measure,') Spirometer.

PNEUMATOMPHALOCÈLE, *Pneumatomphalus*.

PNEUMATOMPHALUS, *Pneumatomphalus*, *celle*, (*pneuma*, and *ομφαλος*, 'umbilicus,') Umbilical intestinal hernia, containing a considerable quantity of flatus.

PNEUMATORRHACHIS, gen. *Pneumatorrhachidis*, (*pneuma*, and *ῥαχίς*, 'spine,') Accumulation of air in the vertebral canal.

PNEUMATOSE, *Pneumatosis*—p. *Péritonéique*, Tympanites.

PNEUMATOSIS, *Aëro'sis*, (from *pneuma*, and *osis*,) *Flat'us*, (F.) *Maladie ventreuse*, *Pneumatose*. Flatulence. Flatulent distension of the stomach or bowels; or both. Sauvages uses the word synonymously with emphysema. It is now generally employed for every abnormal collection of gaseous matters in any part of the body.

PNEUMATOSIS, Emphysema—p. *Abdominis*, Tympanites—p. *Enterica*, *Colica flatulenta*—p. *Pericardii*, *Pneumopericardium*—p. *Pulmonum*, Emphysema of the lungs—p. *Ventriculi et Pneumatosis enterica*, Flatulence.

PNEUMATOTHORAX, *Pneumothorax*.

PNEUMECTASIS, (*pneumon*, and *εκτασις*, 'dilatation,') Emphysema of the Lungs.

PNEUMIC ACID, see Acid, pneumatic.

PNEUMOCACE, (*pneumon*, and *κακός*, 'evil,') Necropneumonia.

PNEUMOCELÈ, *Her'nia pulmonis*, (F.) *Hernie du pmonon*, (*pneumon*, and *κύλη*, 'rupture,') Hernia of the lung through one of the intercostal spaces.

PNEUMOCHOLOSIS, (*pneumon*, and *cholosis*, (*χολή*, 'bile,') 'a biliary disease,') *Pneumonia (bilio)*.

PNEUMOCHYSIS, (*pneumon*, and *χύω*, 'to pour out,') Edema of the lungs.

PNEUMOECTASIE, (*pneumon*, and *εκτασις*, 'dilatation,') Emphysema of the lungs.

PNEUMOGASTRIC, *Pneumogastrius*, (F.) *Pneumogastrique*, (*pneumon*, and *γαστρον*, 'the stomach,') Belonging to the lungs and stomach.

PNEUMOGASTRIC NERVES, *Eighth pair of nerves* (Tenth pair, of some,) *Par vagum, Pars v'ga, V'agus, Nervus vagus seu ambulato'rius seu pneumogastrius seu dec'imus seu sympathicus mel'dius, Mid'dle sympathet'ic*, (F.) *Nervi pneumogastriques ou vagues*; so called because distributed especially to the organs contained in the chest

and abdomen. This nerve arises from behind the eminentia olivaria and near the corpora testiformia by one or two rows of filaments, regularly arranged. These filaments unite to form a flat cord, which issues from the cranium by the foramen lacerum posterius; behind the glossopharyngeal nerve. At its exit from the cranium, the pneumogastric nerve has the appearance of a very compact plexus—*Plexus gangliiformis*, *Ganglion of the superior laryngeal branch*, of Sir Astley Cooper. It then descends along the neck, deeply seated on the outside of the carotid artery and the internal jugular vein. When it has attained the lower part of the neck, it enters the chest, gliding behind the subclavian vein, and passing on the right side, in front of the subclavian artery, and, on the left, before the arch of the aorta. It proceeds, at first, backwards, increasing in size. Afterwards, it passes from behind the bronchia to the oesophagus, which it accompanies through the diaphragm, and terminates on the stomach. In the neck, the pneumogastric nerve anastomoses with the spinal, glossopharyngeal, great hypoglossal, and great sympathetic nerves; and furnishes, 1, a *pharyngeal branch*, 2, a *superior laryngeal*, 3, *cardiac branches*. In the chest, it gives off, 1, the *inferior laryngeal* or *recurrent*, 2, the *pulmonary plexus*, and 3, the *oesophageal branches*. On entering the abdomen, the oesophageal branches, which terminate the pneumogastric, are distributed upon the stomach, and give numerous filaments to that viscus, to the hepatic, coeliac, gastroepiploic, and solar plexuses. Some go, also, to the pancreas, liver, gall-bladder, duodenum, &c.

PNEUMOGRAPHY, *Pneumographia*, (*pneumon*, and *γραφη*, ‘a description.’) The part of anatomy that describes the lungs.

PNEUMO-HEMORRHAGIE, see *Hæmoptysis*.

PNEUMOLITHIASIS, (*pneumon*, and *λιθιασις*, ‘formation of stone.’) A morbid state in which calculi form in the lungs.

PNEUMOLOGY, *Pneumologia*, (*pneumon*, and *λογος*, ‘a discourse.’) An anatomical treatise on the lungs. An anatomical description of the lungs.

PNEUMOMETER, (*pneumon*, and *μετρον*, ‘measure.’) Spirometer.

PNEUMOMETRY, *Pneumometria*. Measurement of the capacity of the lungs for air. See Spirometer.

PNEUMON, (*πνευμων*), Pulmo.

PNEUMONALGIA, (*pneumon*, and *αλγος*, ‘pain.’) Angina pectoris.

PNEUMONATELECTASIS, (*pneumon*, and *atelektasis*), Atelectasis pulmonum.

PNEUMONECROSIS, (*pneumon*, and *νεκρωσις*, ‘death.’) Necropneumonia.

PNEUMONECTASIS, (*pneumon*, and *εκτασις*, ‘dilatation.’) Emphysema pulmonum.

PNEUMONEMPHRAXIS, (*pneumon*, and *εμφραξις*, ‘obstruction.’) Congestion and infarction of the lungs.

PNEUMONIA, (*πνευμονία*, of old, meant a disease of the lungs), *Peripneumonia*, *Empresma pneumonitis*, *Pleumonia*, *Peripneumonia*, *Pneumonitis*, *Pulmonitis*, *Pulmonia*, *Pulmonaria*, *Inflammatio pneumonica*, *I. pectoris acuta seu pulmonum*, *Pneumotaxis*, *Febris pneumonica*, *Pneumony*, *Peripneumony*, *Inflammation of the Lungs*, *Lung-fever* (vulgarly), (F.) *Inflammation des pneumons*, *Fievre pneumonique ou peripneumonique*, *Fluxion de poitrine*, *Inflammation du parenchyme pulmonaire*. (From *pneumon*.) The chief symptoms of pneumonia are:—pyrexia, accompanied by pain, sometimes obtuse, at others pungent,—in some part of the thorax; pulse more or less quick and hard, according to the violence and extent of the local disorder; pain, aggra-

vated by the cough, which, with dyspnoea, exists throughout the disease. At first, the expectoration is difficult and painful; but, in the course of a few days, it becomes free, and the oppression of breathing is mitigated. When the inflammation, instead of going off by resolution, passes on to suppuration, rigors are experienced; the respiration becomes more oppressed, but less painful, and a sense of weight is felt in one of the pleurae. Pneumonia may, also, terminate by gangrene—but this rarely happens,—by induration and by hepatization.

Physical Signs. **FIRST STAGE**, (F.) *Engouement*. Diminution of the sound is generally observed over the affected part on percussion: auscultation exhibits an equal, dry, crepitating *râle*, which is best heard at the close of a deep inspiration, and on coughing. The respiratory murmur is intermingled with the crepitant *râle*, or it may be absent. In the neighbourhood of the affected parts it is natural or puerile. The voice and cough are rather more resonant than usual. **SECOND STAGE**, (F.) *Hépatisation rouge*. When hepatization has occurred, the motion of the affected side is impeded, and immediately above the sternum and in the corresponding triangular space on either side, there is often an evident depression. Percussion is dull over the affected part in every position of the patient. On auscultation, the crepitation is found to be replaced by bronchial respiration. The respiratory murmur is louder in the other portions of the lung, the voice and cough are more resonant, and the heart's action is more distinct. **THIRD STAGE**, (F.) *Hépatisation grise*. When suppuration supervenes, the same signs persist; but usually the voice and cough are less resonant, and a coarse, mucous *râle* is heard. When gangrenous or suppurative cavities are formed, they present the ordinary signs.

The prognosis must be guarded. It is not a disease which we can always subdue. Great extent of inflammation; very considerable oppression; orthopnoea, and difficult expectoration are unfavourable symptoms. The most active treatment is of course necessary. General bleeding forms the sheet anchor,—employed so as to make a decided impression on the system, and repeated, if necessary; local bloodletting; nauseating doses of antimonials; purgatives; and when the violence of the inflammation has been got under,—counterirritants, &c. Chronic pneumonia sometimes succeeds the acute, and, at others, occurs accidentally. It must be managed on general principles. Small bleedings, especially locally; and counterirritants of all kinds are indicated.

PNEUMONIA BILIOSA, P. bilious.

PNEUMONIA, BILIOUS, P. biliosa, *Pneumocholosis*. Inflammation of the lungs, accompanied by gastric fever, and not uncommonly by typhoid symptoms;—*Pneumotiphus*, *Pneumonia typhoides seu typhoides seu typhosa*, *Putrid*, *typhoid*, *adynamic*, or *erysipelatous pneumonia*.

PNEUMONIA, ERSISELATOUS, P. bilious—p. Gangrenosa, Necropneumonia—p. Hypostatica, Pneumonypostasis—p. Intermitiens, Pneumotyposis—p. Lobular, see Lobular—p. Notha, Peripneumonia notha—p. Pleuritis, Pleuropneumonia—p. Putrid, P. bilious—p. Typhodes, P. bilious—p. Typhoid, P. bilious—p. Typhosa, P. bilious, Necropneumonia—p. Vesicular, Bronchitis, vesicular.

PNEUMONIC, (*πνευμονικος*), Pulmonic.

PNEUMONICA, (from *pneumon*.) Diseases affecting the lungs, their membranes or motive power; characterized by irregular, impeded, or painful respiration. The second order of the class Pneumonia of Good.

PNEUMONICI (*morbi*), in the classification of Dr. Wm. Farr, means lung diseases.

PNEUMONIC'ULA, (dim. of *pneumonia*.) A slight inflammation of the lung.

PNEUMONICUS, (πνευμονικός,) Pulmonic.

PNEUMONIE, Pneumonia.

PNEUMONIE DES AGONISANTS, ('pneumonia of the dying,') *Pneumonie hypostatique*. The hyperæmic engorgement, which takes place in the lungs during the last moments of life, has been so termed. It would seem to be hyperæmia rather than inflammation.

PNEUMONIE DISSÉMINÉE. see Lobdular — *p. Lobulaire*, see Lobular — *p. Mamelonnée*, see Lobular.

PNEUMONIT'IC, *Pneumonit'icus*. Of, or belonging to pneumonitis, or inflammation of the lungs.

PNEUMONITIS, (*pneumon*, and *itis*,) Pneumonia.

PNEUMONOCARCINO'MA, gen. *Pneumono-carcinomat'is*, (*pneumon*, and *καρκίνωμα*, 'cancer,') Cancer of the lungs.

PNEUMONOCELĒ, (*pneumon*, and *κηλη*, 'rupture,') Hernia of the lung.

PNEUMONOCE'LE DIAPHRAGMAT'ICA INTER'NA. Hernia of the lung through the diaphragm.

PNEUMONOCELE EXTER'NA seu **THORAC'ICA**. Hernia through the parietes of the chest.

PNEUMONOCELE THORACICA, *P. externa*.

PNEUMONŒDEMA, gen. *Pneumonœdemat'is*, Œdema of the lungs.

PNEUMONOMELANOSIS, see Melanosis.

PNEUMONOMETER, (*pneumon*, and *μετρον*, 'measure,') Spirometer.

PNEUMONOMYCO'SIS, (*pneumon*, *μυκης*, 'a fungus,' and *osis*,) A vegetable parasitic growth in the lungs.

PNEUMONOPHTHISIS, Phthisis pulmonalis.

PNEUMONOPHTHOE, (*pneumon*, and *φθοη*, 'a wasting away,') Phthisis pulmonalis.

PNEUMONOPLEURESIS, Pleuroperipneumony.

PNEUMONOPLEURITIS, Pleuroperipneumony.

PNEUMONORRHAGIA, (*pneumon*, and *ῥαγη*, 'a breaking forth,') Hæmoptysis.

PNEUMONORRHŒA, (*pneumon*, and *ῥεω*, 'to flow,') Hæmoptysis.

PNEUMONOSCIR'RHUS, (*pneumon*, and *κρίσις*, 'induration,') Induration of the lungs.

PNEUMONOSSES, (from *pneumon*, and *osis*,) *Pneumoses*.

PNEUMONY, Pneumonia.

PNEUMONYPOSTASIS, (*pneumon*, and *ὑποστασις*, 'a settling down,') *Pneumo'nia hypostat'ica*. Pneumonia in a depending portion of the lung, caused by lying on the back.

PNEUMOPÉRICARDE, Pneumopericardium.

PNEUMOPERICARD'IUM, *Pneumopericarditis*, *Pneumat'is pericard'ii*, (F.) *Pneumopéricarde*, (*pneumon*, and *pericardium*,) Laënnec designates, under this name, the effusion of air into the cavity of the pericardium.

PNEUMOPHTHISIS, Phthisis pulmonalis.

PNEUMOPHYMATA, (*pneumon*, and *φυμα*, 'a swelling,') Tubercles of the lungs.

PNEUMOPLEURESIS, Pleuroperipneumony.

PNEUMOPHTHOE, (*pneumon*, and *φθοη*, 'a wasting away,') Phthisis pulmonalis.

PNEUMOPYOTHORAX, see Pneumothorax.

PNEUMORRHAGIA, (*pneumon*, and *ῥαγη*, 'a breaking forth,') Hæmoptysis — *p. Interna*, Hæmatothorax.

PNEUMORRHŒA, (*pneumon*, and *ῥεω*, 'to flow,') Bronchorrhœa.

PNEUMOSEPSIS, (*pneumon*, and *σηψις*, 'putrefaction,') Necropneumonia.

PNEUMOSSES, (from *pneumon*, and *osis*,) A term under which Alibert comprises every morbid affection of the lungs.

PNEUMOTHOR'AX, *Pneumothor'ax*, *Pneumatothor'ax*, *Emphyse'ma pectoris*, *Asth'ma æri-um* seu *emphysemat'icum*, *Dyspne'a pneumat'ica* seu *physothorac'ica*, *Physothor'ax*, *Ærothor'ax*, *Æropleurie*, (*pneuma*, and *thorax*,) An accumulation of air in the cavity of the pleura. A complaint generally sudden in its invasion and fatal in its character. In many instances, along with the air, a liquid is effused, consisting of a serous fluid — *Hy'dropneum'othor'ax*, or of one that is purulent — *Pneum'opy'othor'ax*. The disease may be spontaneous and simple, but in a large proportion of cases it is complicated with pleurisy and pulmonary tubercles. The signs vary according as there is, or is not, a communication between the pleura and the bronchia. The affected side gives a hollow, tympanitic sound, even where the thickness of the parietes of the thorax is great. When the effusion is considerable, the affected side is dilated, but there is no rôle of any kind. When a gaseous and a liquid effusion are present at the same time, the sound on percussion is found to be clear at the superior part of the thorax, but dull inferiorly; hence, by varying the position of the patient, and by consequence that of the contained fluids, the seats of the clear and the dull sound will be varied. When the gaseous effusion is owing to a fistulous communication between the pleura and bronchia, the metallic respiration and resonance are heard; and if there be both gaseous and liquid effusion, with a fistulous communication, in addition to these signs there is the *tintement métallique*. The presence of liquid may be ascertained by succussion. See Emphysema of the lungs.

PNEUMOTOMY, *Pneumotom'ia*, (*pneumon*, and *τεμνω*, 'I cut,') Dissection of the lungs.

PNEUMOTYPHUS, Pneumonia, typhoid.

PNEUMOTYPO'SIS, *Pneumo'nia intermit'tens*, (*pneumon*, *τυπος*, 'a stamp, a type,' and *osis*,) Pneumonia characterized by periodicity.

PNEUBIOMANTIA, (πνέω, 'to breathe,' βίω, 'life,' and *μαντεία*, 'divination,') Docimasia pulmonum.

PNEUSIOBIOGNOSIS, (*pneusis*, βίω, 'life,' and *γνωσις*, 'knowledge,') Docimasia pulmonum.

PNEUSIOBOSCOPE, (*pneusis*, βίω, 'life,' and *σκοπεω*, 'to examine,') Docimasia pulmonum.

PNEUSIS, (πνευσις,) Respiration — *p. Pertussis*, Pertussis — *p. Singultus*, Singultus — *p. Tussis*, Tussis.

PNEUSOMETER, (*pneusis*, and *μετρον*, 'measure,') Spirometer.

PNIGALION, (πνιγάλων, from πνιγω, 'to suffocate,') Incubus.

PNIGMA, (πνιγμα,) Orthopnœa, Strangulation.

PNIGOPHOBIA, (*pnigos*, and *φοβος*, 'dread,') Angina pectoris, Suffocation.

PNIGOS, (πνιγος,) Orthopnœa, Strangulation.

PNIX, gen. *Pni'gis*, (πνιξ, gen. *πνιγος*,) Orthopnœa, Strangulation.

PNIXIS, (πνιξις,) Orthopnœa, Strangulation.

PNOE, (πνοη,) Aura, Breath.

PNŒA, (πνοη,) Aura, Breath.

PNŒUM. A nostrum of Dr. Samuel Hahnemann, the founder of the 'Homœopathic System,' which consisted — it is said — of borax only!

PNOOCOLYTICUS, (πνοος, 'breath,' and *κωλυειν*, 'to hinder,') Asthmatic.

PNOUS, (πνοός,) Aura, Breath.

POADMILK, Colostrum.

POCHE DES EAUX, 'bag of waters,' see Liquor amnii, and Parturition.

POCK, (Tent. ποττα, 'a pock or little pouch.') A pustule of smallpox.

POCKARR, Pockmark, see Cicatrix.

POCKBROKEN, see Pockmark.

POCK, KINE, Vaccina.

POCKFRETTE, see Pockmark.

POCKHOLE, Pockmark.

POCK'MARK, *Pock'hole*, (Prov.) *Pockarr*, *Sorob'ulus* seu *Cicatrix Vari'ole*. The mark or pit or pitting left from a smallpox pustule. One pitted with smallpox is said to be *pockmarked*, *pockbroken*, *pockfretten* or *pockfretten*, (Prov.) *Pockurrd*.

POCKY. Infected with, or appertaining to, or resembling, smallpox or syphilis.

POC'ULUM ('a cup') ABSINTHIA'TUM. This was regarded, in remote ages, as a wholesome beverage; the *absinthium*, or 'wormwood,' being supposed to act as an antidote to drunkenness.

POCULUM AMATORIUM, Philter — p. Diogenis, Palm — p. Emeticum, Goblet, emetic — p. Vomitorium, Goblet, emetic.

POD'AGRA, *Podarthrit'is*, *Podorrheu'ma*, *Podal'gia*, (*podō*, and *agrap*, 'seizure.') Pain which attacks the feet. Gout, situate in the articulations of the foot. It has, also, been used in a more extensive signification synonymously with *gout*.

PODAGRA ABERRANS, Gout, (wandering) — p. Arthritis, Gout — p. Retrograda, Gout, (retrograde.)

PODAGRARIA, *Ligusticum podagraria* — p. *Ægopodium*, *Ligusticum podagraria*.

PODAGR'IC, *Podag'ric*, (*ποδαγρικός*,) *Podag'ricus*, *Arthrit'icus*, *Arthrit'ic*, *Gout'y*, (F.) *Goutteux*. Relating or belonging to gout. Affected with, or liable to gout.

PODAGRISM, see Diathesis.

PODALGIA, (*ποδάλγια*, from *podō*, and *algos*, 'pain.') Pain in the foot. Gout, Podagra.

PODALIC, Pedal — p. Version, see Turning.

PODALIRIA TINCTORIA, (after Podalirius, son of Æsculapius,) *Sophora tinctoria*.

PODANENCEPHALIA, see Podencephalus.

PODARTHITIS, Podagra.

PODARTHRO'ACÉ, (*podō*, *arthron*, 'an articulation,' and *κακός*, 'bad.') Caries of the articulation of the foot.

PODELO'MA, *Podelco'ma*, (*podō*, and *ἔλκος*, 'an ulcer.') A peculiar ulcerous affection of the foot, described by Prof. Miller, of Edinburgh, which is but little amenable to treatment; and which has been likewise called *Mor'bus tuberculo'us pe'dis*.

PODENCEPHALIA, see Podencephalus.

PODENCEPH'ALUS, *Podanenceph'alus*, (*podō*, and *κεφαλή*, 'the head.') A monster whose brain is placed outside the skull, and seems to be supported on a pedicle, which traverses the summit of the skull. This state of monstrosity is termed *Podanenceph'al'ia*, or more properly, *Podenceph'al'ia*.

PODEX, (gen. *Po'dicis*,) Anus, Nates.

PODO, (*πους*, gen. *πόδος*, 'a foot,') in composition, foot.

PODODYN'IA, (*podō*, and *δύνη*, 'pain.') Pain in the soles of the feet induced by long standing, as in cutters of garments, &c.

PODŒDE'MA, gen. *Podœde'matis*, (*podō*, and *ἔδημα*, 'a swelling.') Edema of the foot.

PODOL'OGY, *Podolog'ia*, (*podō*, and *λογος*, 'a discourse,') A description of the foot. A treatise on the foot.

PODOPHYLLUM MONTANUM, *Mountain May'apple*, *Man'drake*, *Wild Lem'on*, *Ducks'foot*, *Raccoon's Berry*, *Yellow Ber'ry*, *Ground Lem'on*, Ord. Berberidaceæ, *Sez. Syst.* Polyandria Monogynia, (*podō*, and *φυλλον*, 'a leaf,' in reference to the shape of the leaf,) has the same properties as

PODOPHYLLUM PELTATUM, *Podophyl'lum*, *Ana-podophyl'lum Canadense*, *Mayapple*, *Mandrake*. A common plant throughout North America. The fruit is eatable, and esteemed by many. The leaves are said to be poisonous. The root or rhizoma, *Podophyllum* (Ph. U. S.), is purgative in the dose of 20 grains. It has, also, been used as an anthelmintic. *Podophyl'lin*, obtained by evaporating an alcoholic solution of the root, or by displacement by alcohol, throwing down the precipitate by water, and drying — *Res'ina Podophyl'li* — is an excellent purgative in the dose of 10 grains, and a laxative in the dose of a grain or more. See *Resina Podophylli*.

PODORRHEUMA, (*podō*, and *ῥευμα*, 'a defluxion,') Podagra.

PODOTH'E'CA, (*podō*, and *θηκη*, 'a receptacle, a sheath,') The cuticle of the foot. An anatomical preparation. *Chirothe'ca* has been used for the cuticle of the hand.

POECLIA, (*ποικιλία*, 'a variegated appearance,') see Achroma.

POËLETTE, Cup, Palette.

POEPHAGUS, (*ποη* or *ποα*, 'a plant, an herb,' and *φαγω*, 'I eat,') One who subsists on herbs or vegetables. The act of subsisting or feeding on herbs or vegetables is called *Pœphag'ia*. Hence Strabo calls the Irish *ποηφαγοί*.

POGON, (*πωγων*,) Beard.

POGONI'ASIS, *Pogo'nia*, (from *pogon*,) A female beard. Also, great strength or quantity of beard. A female having a beard — *Vira'go*.

POGO'NIUM, (*πωγωνιον*, dim. of *pogon*,) A weak or small beard.

POGONOL'OGY, *Pogonolog'ia*, (*pogon*, and *λογος*, 'a description,') A description of, or treatise on, the beard.

POGOSTEMON INTERMEDIUS, (*pogon*, and *στημων*, 'a stamen,') P. Patchouli.

POGOSTEMON PATCHOULI, P. *Sua'vis* seu *interme'dius*, *Pucha-Pat*, *Patchouli*. Nat. Ord. Labiate. Native of Penang and the Malayan peninsula. The herb is used as a perfume. The distilled oil is used for the same purpose.

POGOSTEMON SUAVIS, P. Patchouli.

POHON ANTAR, Upas — p. Upas, Upas.

POI. A favorite aliment in the Sandwich Islands. It is a sort of paste made from the root of the Kalo — *Arum esculentum*.

POIDS ET MÉSURES, (from [L.] *pondus*, 'a weight,') Weights and measures.

POIGNÉE, (from [L.] *pugnus*, 'the fist,') Manipulus.

POIGNET, Carpus.

POIL, [L.] (*pilus*, 'a hair,') Galactoele, Hair.

POILETTE, Cup.

POILS, (pl. of *poil*,) see *Typha latifolia*.

POING, [L.] (*pugnus*,) Fist.

POINT, (from *pungere*, *punctum*, 'to prick,') Puncta, Stitch.

POINT, BLIS'TERING. A name given by Dr. Rush to an indeterminate period in the course of a continued fever, intermediate between the stages of high excitement and collapse, in which blisters will — he thought — usually produce unequivocally good effects.

POINT DE COTÉ, 'stitch of the side,' Pleurodynia.

POINT, VITAL, see *Centrum vitale*.

POINTING OF AN ABSCESS, see Suppuration.

POINTS LACRYMAUX, Lacrymalia puncta.

POIRE, see *Pyrus communis*.

POIRÉ, see Cider.

POIREAU, (*porrum*, 'a leek,') *Allium porrum*, Verruca.

POIRIER, ([*L. pyrus*,) *Pyrus communis*.

POIS, *Pisum*—*p. à Cautère*, see *Fonticulus*—*p. Chicke*, *Cicer arietinum*—*p. Cultivé*, *Pisum*—*p. à Gratter*, *Dolichos pruriens*—*p. Patate*, *Dolichos tuberosus*.

POISON, *Phar'macon*, *Phar'macum*, *Tox'icum*, *Vene'num*, *Vi'rus*, *Delete'rium*, *Venena'tion*, (F.) *Poison*. A generic name for all substances which, when introduced into the animal economy, either by cutaneous absorption, respiration, or the digestive canal, act in a noxious manner on the vital properties or the texture of organs. Hence we speak of *fever poison*, *cholera poison*, &c. Poisons exist in the three kingdoms of nature; but those which proceed from animals are often called *venoms*, as the venom of the viper, scor-

pion, tarantula, &c.; whilst those that are the products of disease have the name *virus*. In common parlance, therefore, *poison* is restricted to deleterious articles, furnished by the mineral and vegetable kingdoms. Orfila has divided poisons into four classes. 1. **ACRID**, **IR'ITATING**, **CORRO'SIVE**, or **ESCHAROT'IC**, as the concentrated acids and alkalis; mercurial, arsenical, cupreous, and antimonial compounds, cantharides, &c. 2. **NARCOT'IC**, those that act particularly upon the brain; as hyoscyamus, opium, &c., but without inflaming the organ with which they come in contact. 3. **NARCOT'ICOACRID** or **ACRO-NARCOT'IC**; those that act on the brain, or spinal marrow or both, but, at the same time, irritate the parts to which they are applied; as aconite, belladonna, &c. 4. **SÉP'TIC** or **PUTRES'CENT**; those furnished by the animal kingdom. See *Venom* and *Virus*. Various classifications, of a similar character, have been recommended by different toxicologists; but they are liable to the objection, that they throw substances together whose physiological action on the system is very different. It is, indeed, difficult to avoid unnatural compression of matters into places not properly belonging to them in all such arrangements.

The classification of Dr. A. S. Taylor is as follows:

IRRITANTS	{	MINERAL	{	Non-metallic	{	Acids, Alkalies, and their Salts.		
		VEGETABLE		Metallic		Metalloids.		
		ANIMAL			(Arsenic).			
NEUROTICS	{	CEREBEAL	(Morphia).					
		SPINAL	(Strychnia).					
		CEREBRO-SPINAL	(Conia, Aconitia).					

The accurate investigations of modern chemistry, by separating the active alkaloids of different vegetables, have furnished numerous agents for poisoning; and, hence, deaths from morphia and strychnia have repeatedly become subjects of medicolegal inquiry in courts of justice. Like investigations have facilitated the discovery of many toxical substances after death in the contents of the alimentary canal and elsewhere; although the failure of such discovery may not be a positive evidence that no such poison was administered, as it may have been rejected by vomiting; or, if sufficient time has elapsed,

may have been eliminated in some other manner.

One of the most astonishing of tests is that by spectrum analysis, which, in certain cases, is capable of detecting the most inconceivably minute portions of a metal,—in one instance the 195,000,000 part of a grain. See *Spectrum analysis*.

The following table, which exhibits a *comp d'ais* of the chief poisons, with many of the circumstances of importance connected with them, is not free from the objections above stated

TABLE OF THE CHIEF POISONS

N.B. In all cases the Stomach Pump should be used as soon as possible.

I. INORGANIC POISONS.

POISONS.	SYMPTOMS.	TREATMENT.	TESTS.
ACIDS.	The acids, generally, are strong corrosive poisons. Sour, acid taste, burning in the throat, which is increased by pressure, swallowing, or coughing; eructation, and excruciating pain in the stomach; more or less corrugation of the lining membranes of the mouth and primæ viæ; excoriation about the mouth or such other parts of the skin as the acid may have touched. The matter vomited effervesces with carbonate of lime. The countenance becomes glazed, extremities cold and clammy; convulsions and death. Nitric acid occasions yellow stains, and sulphuric acid, black.	The carbonates of soda, potassa, lime and magnesia, are all antidotes to the acids; calcined magnesia also. They are to be used with the following restrictions: for the acetic, citric, muriatic, sulphuric, and tartaric acids, they may be used indiscriminately. For the nitric and oxalic, carbonates of magnesia and lime can alone be employed with safety. In the case of sulphuric acid, water should not be drunk, on account of the great heat which is produced by their mixture. Subsequent inflammation to be treated by ordinary means.	<i>Sulphuric acid</i> is known by a white precipitate with any salt of baryta, insoluble in nitric acid. <i>Muriatic acid</i> is known by a white precipitate with nitrate of silver, insoluble in nitric acid, which turns black by the action of light. <i>Nitric acid</i> gives rise to orange-coloured fumes, when copper is placed in it, and an ability to dissolve gold when mixed with muriatic acid. <i>Acetic acid</i> exhales the odour of vinegar. <i>Citric acid</i> blackens when heated. <i>Tartaric acid</i> precipitates crystals, with a solution of potassa; and a white precipitate is thrown down by lime-water. <i>Oxalic acid</i> gives a white precipitate with lime-water, which, when heated, is converted into carbonate of lime.
<i>Prussic Acid.</i> — <i>Oil of Bitter Almonds.</i> — <i>Laurel Water.</i>	Is a sedative poison; nausea, giddiness, debility, hurried pulse, weight and pain in the head; eructations having the flavour of the acid; spasms, tetanus, contractile pupil; convulsions, death.	The carbonates of the alkalies and of magnesia and lime have the inconvenience, that a large quantity of gas is extricated in the stomach. Ammonia is an antidote; but it should not be employed in a very concentrated form. Liquid chlorine has also been found efficacious. The cold <i>douche</i> to the head has been recommended.	<i>Prussic acid</i> exhales the odour of peach kernels; forms a white precipitate with nitrate of silver neutralized with an alkali, gives a blue precipitate with the persalts of iron.
ALKALIES and their SALTS. — AMMONIA. <i>Strong Liquor of Water of.</i> — <i>Muriate of Ammonia or Sal Ammoniac.</i> — POTASSA. <i>Caustic Potassa, and Liquor Potassa.</i> — <i>Carbonate of Potassa, or Pearl Ash, and Salt of Tartar.</i> — <i>Nitrate of Potassa or Salpêtre.</i> — <i>Sulphuret of Potassium or Liver of Sulphur.</i> — SODA.	Violent, caustic, acid taste; great heat in the throat, with destruction of its lining membrane; difficult and painful deglutition; vomiting of bloody matter, which turns the yellow of turmeric brown; acute pain in the stomach; cold sweats, weakness, hicough; violent colic pains, with purging of bloody stools and membranous flakes; death.	The vegetable acids, such as vinegar, lemon juice, citric and tartaric acid in solution are antidotes to the alkalies and their carbonates. The fixed oils, such as castor, linseed, almond and olive, form soaps with the free alkalies, and therefore destroy their caustic effects.	The alkalies are known by their action on turmeric paper: restoring the colour of litmus, which has been reddened by an acid. Carbonates are known by their effervescence with an acid; some of them affect the turmeric paper. Sal ammoniac gives out the pungent odour of ammonia, when mixed with quicklime.
<i>Carbonate of Potassa, or Pearl Ash, and Salt of Tartar.</i> — <i>Nitrate of Potassa or Salpêtre.</i> — <i>Sulphuret of Potassium or Liver of Sulphur.</i> — SODA.	The nitrate of potassa will not manifest the effect with turmeric paper. The carbonates, when vomited, will effervesce with acids; and the liver of sulphur will give rise to eructations of sulphuretted hydrogen.	Poisoning by nitrate of potassa to be treated on general antiphlogistic principles; mucilaginous drinks. Liver of sulphur is said to be decomposed and neutralized by common salt. The liquid chloride of soda will also decompose it.	Nitrate of potassa decrepitates and deflagrates when thrown on hot coals. Liver of sulphur emits the odour of sulphuretted hydrogen when dissolved, or when treated with an acid.
EARTHS and COMPOUNDS. — BARYTA. — <i>Carbonate of Baryta.</i> — <i>Chloride of Baryum.</i> — <i>Nitrate of Baryta.</i> — LIME.	Analogous to those of the corrosive metals. Violent burning in the stomach, vomiting, gripes, diarrhœa; excessive muscular debility, headache, convulsions, death. Lime differs from baryta in being a pure irritant.	The sulphates of soda and magnesia are prompt and effective antidotes to all the poisonous salts of baryta. Phosphate of soda will also counteract their effects. Lime may be neutralized by dilute acids. Carbonic acid, in soda-water, effervescing draught or yeast, it is supposed, would answer a good purpose. The fixed oils may be employed either for baryta or lime, when not in a compound state.	Baryta and its salts invariably give a white precipitate, insoluble in acid, with a soluble sulphate. Lime, when dissolved, gives a white precipitate with oxalic acid. Also with carbonic acid or a soluble carbonate. The same of baryta.

POISONS.	SYMPTOMS.	TREATMENT.	TESTS.
ALCOHOL. <i>Brandy, Wines, and all Spirituous Liquors.</i>	Intoxication, and when taken very freely, complete insensibility, with apoplexy or paralysis of one side; the countenance swollen and of a dark-red colour; the breathing difficult, and often stertorous, with a peculiar puffing out of the lips; the breath smells of liquor, which will distinguish the symptoms from those of spontaneous apoplexy.	A powerful emetic of white vitriol, or tartar emetic, should be got into the stomach as soon as possible; and if the person has lost the power of swallowing, a flexible catheter or tube should be the means of conveying it thither. The vomiting should be encouraged as much as possible with warm water; and large and active glysters of salt and water should be thrown up. The patient should be placed erect, and if the countenance and other appearances be not improved after these means have been used, the jugular vein may be opened, and cold wet cloths applied to the head, particularly if the body be hotter than natural. If the extremities become cold, warmth and friction should be perseveringly used.	
VOLATILE OILS. <i>Creosote.</i> <i>Dippel's Animal Oil.</i> <i>Oil of Tar.</i> <i>Oil of Tobacco.</i> <i>Oil of Turpentine.</i> <i>Fusel Oil.</i>	General action, that of irritant poisons. Burning pain, vomiting, pungent taste, purging, &c. The oils of turpentine and tobacco affect the nervous system; the peculiar odour of each oil will be manifested in the matter vomited.	Creosote is immediately coagulated by albumen. Dippel's animal oil may be counteracted with dilute acids and the fixed oils. The other oils have no particular antidotes, and their effects must, therefore, be counteracted upon general principles.	No better mode of recognizing these substances exists, than that derived from their peculiar odour. Dippel's oil has the pungent odour of ammonia; creosote and oil of tar, a peculiar smell of smoke; the odour of tobacco and turpentine are well known; and fusel oil has a peculiar, offensive, suffocating odour.
GASES. <i>Carbonic Acid, or Fixed Air.</i> <i>Carbonic Oxide.</i> <i>Fumes of Burning Charcoal.</i> <i>Chlorine.</i> <i>Sulphuretted Hydrogen.</i>	Chlorine produces, when inhaled, violent irritation of the organs of respiration; cough, bloody expectoration, inflammation of the lungs, and permanent pulmonary disease. The other gases, although producing some effect on the respiratory organs, act as poisons, in consequence of their sedative agency. The symptoms, therefore, are those of apoplexy, or narcotic poisoning.	The antidotes to chlorine are the cautious inhalation of ammonia, or sulphuretted hydrogen. The inflammatory symptoms from chlorine to be treated on general principles. For the other gases, cold affusions to the head, blood-letting, artificial respiration. In poisoning by chlorine, ether may be cautiously inhaled.	Chlorine is recognized by its peculiar odour and bleaching properties. Sulphuretted hydrogen, by its fetid smell, and the rapid blackening of lead. Carbonic acid, by its occasioning turbidness in lime-water placed in its atmosphere. Carbonic oxide, by the blue colour of its flame.
IODINE. <i>Iodide of Potassium.</i>	Irritant symptoms; burning pain in the throat, lacerating pain in the stomach, and fruitless efforts to vomit; suffusion of the eyes; excessive pain and tenderness of the epigastrium.	Iodine combines with starch, and forms an insoluble compound. The prompt administration of starch, wheat flour, or other vegetable matter containing fecula, beat up in water, is recommended. Iodide of potassium has no antidote. Vomiting should be promoted by draughts of warm water, and inflammation be subdued by general treatment.	Iodine is known by its odour, and the formation of a blue precipitate, when brought in contact with a cold solution of starch. Iodide of potassium gives a crystalline precipitate, with tartaric acid in excess. The supernatant fluid will give the blue colour to starch.
METALS. ANTIMONY. <i>Tartar Emetic.</i> <i>Chloride, or Buter of Antimony.</i> <i>Oxide of Antimony.</i>	Vomiting. If vomiting do not occur promptly, violent irritant effects are produced. Burning pain in the pit of the stomach; Purging; colicky pains, sense of tightness in the throat, violent cramps; repeated recurrence of vomiting.	If vomiting have not been produced, it should be brought about by tickling the fauces, and administering copious draughts of warm water. Astringent infusions, such as of galls, oak bark, Peruvian bark, act as antidotes, and should be given promptly; powdered yellow bark may be used, until the infusion is prepared.	Antimony in solution is best distinguished by the peculiar orange-red precipitate, which it forms with sulphuretted hydrogen or solutions of the hydrosulphates. Free alkalis produce white precipitates. The chloride is known by a white precipitate, when it is poured into water. In its solid forms, it may be known by the formation of white fumes, when heated, which reddens litmus. It may also be converted into chloride, and then precipitated by adding water.

POISONS.	SYMPTOMS.	TREATMENT.	TESTS.
ARSENIC. <i>4-venious Acid, or White Arsenic.</i> <i>Orpiment, or Yellow Sulphuret of Arsenic.</i> <i>King's Yellow.</i> <i>Realgar, or Red Sulphuret of Arsenic.</i> <i>Fly Powder.</i> <i>Fowler's Solution.</i> <i>Arsenical Paste.</i> <i>* Arsenical Soap.</i> <i>Arsenite of Copper.</i> <i>Scheele's Green.</i>	Violent burning pain in the region of the stomach and bowels; tenderness on pressure; retching; vomiting; sense of dryness and tightness in the throat; thirst; hoarseness and difficulty of speech; the matter vomited, greenish or yellowish, sometimes streaked with blood; diarrhœa; tenesmus; sometimes excoriation of the anus; urinary organs occasionally affected with violent burning pains and suppression; convulsions and cramps; clammy sweats; lividity of the extremities; countenance collapsed; eyes red and sparkling; delirium; death.	The hydrated peroxide of iron diffused through water; or the precipitated carbonate; or the rubigo ferri, in very fine powder, to be administered every five or ten minutes, until relief is obtained. This is particularly efficacious when the <i>white arsenic</i> has been swallowed. If the arsenic have been taken in the form of Fowler's Solution, lime-water, in copious draughts, may be given. For either of the other forms, emetics of sulphate of zinc; diluents; demulcents, such as flaxseed tea, infusion of slippery elm, &c. Counter-irritants may be used to relieve the spasm, and violent pain in the stomach. Bleeding, as promoting absorption, should not be employed until the stomach is completely evacuated. Magnesia, freshly precipitated or not too highly calcined has been used with advantage in arsenious acid poisoning.	If the suspected substance can be obtained free from organic matter, the presence of arsenic may readily be shown by boiling it in distilled water, filtering, and testing with, 1st. Ammoniacal sulphate of copper. 2d. Ammoniacal nitrate of silver. (<i>Hume's Test</i>) 3d. Sulphuretted hydrogen gas. The first will produce a <i>green</i> precipitate; second, a <i>pale yellow</i> precipitate; third, a <i>bright yellow</i> precipitate, if arsenic be present. A portion of the powder, or one of these precipitates, carefully and thoroughly dried, is then to be mixed with charcoal and exsiccated carbonate of soda, to be placed in a small glass tube, closed at one end, and then to be heated to a red heat in the flame of a lamp. The arsenic will be reduced, and sublime in the upper part of the tube, forming a dark crust, having considerable lustre on its exterior surface, resembling polished steel. By gently heating this crust in a very small flame, crystals of white arsenic will be formed, having a high adamantine lustre. If organic matter be present, it must be entirely destroyed, by heating the substance with nitromuriatic acid. After this has been done, and a clear, transparent solution in distilled water has been obtained, the subsequent process is as above indicated. If arsenious acid be submitted to the action of nascent hydrogen, it is deoxidized, and the metallic arsenic, thereby obtained, combining with hydrogen, forms arsenuretted hydrogen gas. (<i>Marsh's Test</i>.) Add to a suspected solution a few drops of pure chlorohydric acid, and place in it a slip of bright copper; no change occurs until the liquid is brought to the boiling point, when, if arsenic be present, even in small quantity, the copper acquires an iron-gray coating, from the deposit of that metal. Remove the slip of copper; wash it in water, dry it and gradually heat it in a reduction tube, when arsenious acid will be sublimed in minute octohedral crystals. The test succeeds with powdered arsenic, the arsenites, arsenic acid, the arseniates, and orpiment. It will even separate the arsenic from arsenite of copper, and from common lead shot. <i>Reinsch's Test</i>. Bismuth is known by the formation of a white precipitate, when its solution in nitric acid is poured into water; and by the formation of a black precipitate with sulphuretted hydrogen gas, and hydrosulphates in solution.
BISMUTH. <i>Nitrate of Bismuth.</i> <i>Pearl Powder.</i> <i>Oxide of Bismuth.</i> COPPER. <i>Sulphate of Copper</i> <i>Blue Vitriol.</i> <i>Acetate of Copper,</i> <i>Verdigris.</i> <i>Carbonate of Copper.</i> <i>Blue Verditer.</i> <i>Arsenite of Copper</i> <i>Scheele's Green.</i> <i>Food cooked in dirty copper vessels, or pickles made green by copper.</i>	Similar to those produced from other irritant poisons. General inflammation of the whole alimentary canal; suppression of urine; hicough, disagreeable metallic taste; vomiting; cramps; delirium; death. Very similar to those produced by arsenic. Coppery eructations and taste. Fatal cases are generally terminated by convulsions, palsy, insensibility.	Milk, and sweet mucilaginous drinks are recommended. Leeches, general bleeding; glysters; fomentations to be employed on the general principles of treatment for inflammatory symptoms. Albumen to be administered in either of its forms which can be most readily obtained, as milk or whites of eggs. Vinegar should not be given. The inflammatory symptoms are to be treated on general principles, and so of the nervous.	The presence of copper is readily shown in solutions, by ammonia a beautiful bluish precipitate is thrown down, which dissolves in an excess of the precipitant, and the solution has a rich blue colour. A bright iron bar, introduced into solutions of copper, occasions the separation of the copper, in a metallic form, which deposits itself upon the iron. A more delicate test than the ammonia is the ferrocyanide of potassium, which gives a brown precipitate with copper salts.

POISONS.	SYMPTOMS.	TREATMENT.	TESTS.
GOLD. — <i>Chloride of Gold.</i> — <i>Fulminating Gold.</i>	Very analogous to those of other irritant poisons. This substance communicates a pink stain to the flesh, and patches of that colour may be found about the lips, and inside the mouth.	The salts of gold are decomposed by sulphate of iron; and this has, therefore, been recommended as an antidote. Inflammatory symptoms to be treated on general principles.	Chloride of gold is very readily decomposed, and the gold separated in a metallic state. Proto-sulphate of iron, nitrate of silver and protochloride of tin, all answer this purpose. The precipitated powder will be found insoluble, except in nitromuriatic acid.
IRON. — <i>Sulphate of Iron,</i> <i>Copperas,</i> <i>Green Vitriol.</i> — <i>Chloride of Iron.</i>	Symptoms of irritant poisoning; colic pains, constant vomiting and purging. Violent pain in the throat, tension of the epigastrium, coldness of the skin, and feebleness of the pulse.	Carbonate of soda would be a most excellent antidote to either of these substances. Mucilaginous drinks might also be employed; and particular symptoms relieved by general treatment.	Iron is recognised by a bluish-black precipitate, with tincture of galls; rich blue, with ferrocyanide of potassium; and a greenish or reddish precipitate, with the free alkalies or their carbonates.
LEAD. <i>Acetate of Lead.</i> <i>Sugar of Lead.</i> <i>Carbonate of Lead.</i> <i>White Lead.</i> <i>Red Oxide, or Red Lead.</i> <i>Litharge.</i> — <i>Wines sweetened by Lead.</i> <i>Water which has been kept in Leadens vessels.</i> <i>Acid food, cooked or left standing in vessels glazed with Lead.</i>	Irritation of the alimentary canal; spasm; nervous symptoms; paralysis, either partial or complete. When taken for some time, in small quantity, violent and obstinate colic; rigidity of abdominal muscles, cramps; remission of pain; obstinate constipation, urine diminished, saliva increased; countenance anxious and gloomy. If relief be not promptly obtained, giddiness, debility, torpor, coma, convulsions and death. The paralysis affects generally the upper extremities.	Sulphate of magnesia and phosphate of soda are both good antidotes for the soluble salts of lead. For the solid forms, dilute sulphuric acid may be drunk. These are applicable to the irritant forms of poisoning by lead. In the chronic form, or colica pictorum, purgatives and anodynes are resorted to; venesection; and external applications to relieve the cramps. The use of strychnia is recommended for the paralysis, and iodide of potassium for the chronic form generally.	Soluble salts of lead yield a white precipitate with sulphates and carbonates in solution. Salts of chromic acid and iodide of potassium occasion a yellow precipitate. Sulphuretted hydrogen, a black precipitate. If solid, the lead may be converted into a state of solution by nitric acid and the dilute solution be tested as above.
MERCURY. <i>Corrosive Sublimiate.</i> — <i>Cyanide of Mercury.</i> — <i>Nitrate of Mercury.</i> <i>White Precipitate.</i> <i>Red Oxide, or Red Precipitate.</i> — <i>Sulphate, or Turbith Mineral.</i> <i>Vermilion, or Red Sulphuret.</i>	Violent symptoms of irritant poisoning; harsh metallic astringent taste; burning pain in the stomach; vomiting and purging frequently of bloody matter; often irritation of the urinary organs, and sometimes suppression; tightness and burning in the throat, occasionally so great as to prevent speech; countenance not always pale, but sometimes flushed; tendency to doze; stupor, convulsions and death.	Albumen, in some form, must be promptly administered; either white of eggs beaten up with water, milk, or wheat flour beaten up. The inflammatory symptoms to be counteracted by the usual means. Gold, finely mixed in dust, with fine iron filings. The iron in filings or in powder— <i>ferri pulvis</i> —has been given enclosed in gold leaf. In the chronic form iodide of potassium has been given.	The free alkalies differ in their precipitates from solutions of mercury. Potassa gives, with corrosive sublimiate, a yellowish one; ammonia, a white; lime-water, an orange; and sulphuretted hydrogen, a black. If a drop of the solution be placed on a piece of gold, and touched with a knife or key, the mercury will be reduced, which amalgamates with the gold, producing a white spot. Protochloride of tin is the best liquid test. It gives a grayish precipitate. All the solid forms of mercury are volatile, and may be reduced by heating them in a glass tube with charcoal and carbonate of soda, when the metallic mercury will sublime, and may be collected in a globule.
SILVER. <i>Nitrate of Silver, or Lunar Caustic.</i>	Those of other irritant poisons.	Chloride of sodium, or common salt, immediately decomposes this substance, and destroys its activity. Antiphlogistic treatment is to be employed for the inflammatory symptoms.	Nitrate of silver is distinguished by the formation of a white precipitate, insoluble in nitric acid, with chloride of sodium; a gray precipitate with ammonia, which is redissolved in an excess of ammonia; yellow precipitate with phosphate of soda. The precipitate of chloride of silver, obtained by adding chloride of sodium to the nitrate of silver, is readily reduced, and the metallic silver obtained, by mixing it with dilute sulphuric acid and granulated zinc.
TIN. <i>Chloride of Tin.</i> <i>Solution of Tin, used by Dyers.</i> <i>Oxide of Tin, or Putty Powder.</i>	The same as those from other irritant poisons, and a peculiar tanned appearance of the villous coat of the stomach.	Milk to be given copiously, and the subsequent treatment to be regulated by the symptoms.	The chloride precipitates vegetable colouring matter; also albumen and gelatin. The white oxide is precipitated from its solution by free alkalies; with sulphuretted hydrogen gas, a brown precipitate.
ZINC. <i>Sulphate of Zinc.</i> <i>White Vitriol.</i> <i>Acetate of Zinc.</i>	Violent vomiting; astringent taste, burning pain in the stomach; pale countenance; cold extremities; dull eyes; fluttering pulse. Death seldom ensues, in consequence of the emetic effects.	The vomiting may be relieved by copious draughts of warm water. Carbonate of soda, administered in solution, will decompose the sulphate of zinc. Milk and albumen also act as antidotes. General principles to be observed in the subsequent treatment.	A white precipitate is thrown down by the free alkalies from the soluble salts of zinc, which becomes yellow when strongly heated—recovering its white colour on cooling. Sulphuretted hydrogen occasions a white precipitate, but only in neutral or alkaline solutions. Sulphohydrate of ammonia is better.

POISONS.	SYMPTOMS.	TREATMENT.	TESTS.
PHOSPHORUS.	Symptoms of irritant poisoning; pain in the stomach and bowels; vomiting, diarrhoea, tenderness and tension of the abdomen.	An emetic to be promptly administered; copious draughts containing magnesia in suspension; mucilaginous drinks; general treatment for inflammatory symptoms.	Phosphorus is known by its ready inflammability, luminousness at ordinary temperatures, and alliaceous odour.
GLASS or ENAMEL.	If taken in very coarse powder, it produces irritation and inflammation of the bowels.	Large quantities of crumb of bread should be eaten, to envelop the particles. (?) An emetic of sulphate of zinc should then be given, and vomiting be promoted by demulcent drinks.	

II. ORGANIC POISONS.

1. VEGETABLE POISONS.

POISONS.	SYMPTOMS.	TREATMENT.
IRRITANT.		
<i>Anemone pulsatilla</i> . Wind Flower.	The general effects of this class of vegetable poisons are;—an acrid pungent taste, with more or less of bitterness, excessive heat, great dryness of the mouth and throat, with sense of tightness there; violent vomiting, the efforts being continued even after the stomach is emptied; purging, with great pain in the stomach and bowels; pulse strong, frequent, and regular; breathing often quick and difficult; appearances of intoxication; the pupil of the eye frequently dilated, insensibility resembling death; the pulse becomes slow and loses its force, and death closes the scene.	If vomiting have been occasioned by the poison, and the efforts are still continued, they may be rendered easier by large draughts of warm water, or thin gruel; but if symptoms of insensibility have come on without vomiting, it ought to be immediately excited by the sulphate of zinc, or some other active emetic substance, and after its operation a sharp purgative should be given. After as much as possible of the poison is got rid of, a very strong infusion of coffee, or vinegar diluted with water, may be given with advantage. Camphor mixture with ether may be taken frequently, and if insensibility be considerable, warmth, frictions, and blisters, may be employed. If inflammation or other dangerous consequences have been induced, they are to be treated upon general principles.
<i>Arum maculatum</i> . Wake Robin.		
<i>Bryonia dioica</i> . Bryony.	If applied externally, many of them produce violent inflammation of the skin, with vesications or eruptions of pustules.	The fruit of <i>Feuillea Cordifolia</i> has been recommended as a powerful antidote against vegetable poisons; (?) it is to be used in as recent a state as possible.
<i>Caladium seguinum</i> . Dumbcane.		
<i>Calla palustris</i> . Water Arum.	N. B. Bromine, chlorine, and Iodine are said to be antidotes to the alkalis generally, and so are tannic acid and animal charcoal.	
<i>Caltha palustris</i> . Marsh Marigold.		
<i>Chelidonium majus</i> . Celandine.		
<i>Clematis vitalba</i> . Virgin Bower.		
<i>Convolvulus jalapa</i> . Jalap.		
<i>Convolvulus scammonia</i> . Scammony.		
<i>Croton tiglium</i> . Purging Croton.		
<i>Cucumis colocynthis</i> . Colocynth.		
<i>Cyclamen Europeum</i> . Sow Bread.		
<i>Daphne gnidium</i> . Spurge Flax.		
<i>Daphne mezereum</i> . Mezereum.		
<i>Delphinium staphisagria</i> . Stavesacre.		
<i>Dioica palustris</i> . Swamp leather-wood.		
<i>Equisetum hyemale</i> . Scourgrass.		
<i>Euphorbia officinarum</i> . Euphorbium.		
Spurge.		
<i>Gratiola officinalis</i> . Hedge Hyssop.		
<i>Hippomanne mancinella</i> . Manchineel.		
<i>Hura crepitans</i> . Sand Box.		
<i>Hydrocotyle vulgaris</i> . Marsh Pennywort.		
<i>Jatropha curcas</i> . Indian Nut.		
<i>Jatropha manihot</i> . Cassada.		
<i>Juniperus sabina</i> . Savin (oil of).		
<i>Juniperus Virginiana</i> . Red Cedar (oil of).		
<i>Momordica elaterium</i> . Squirtng Cucumber.		
<i>Narcissus pseudo-narcissus</i> . Daffodil.		
<i>Pastinax sativa</i> . Common Parsnep.		
<i>Pedicularis palustris</i> . Marsh Lousewort.		
<i>Phytolacca decandra</i> . Poke.		
<i>Piper cubeba</i> . Cubebs.		
<i>Plumbago Europaea</i> . Toothwort.		
<i>Ranunculus acris</i> , and other species.		
Crowfoot.		
<i>Rhododendron chrysanthemum</i> . Oleander.		
<i>Ricinus communis</i> . Castor Oil Plant.		
<i>Sambucus ebulus</i> . Elder.		
<i>Sedum acre</i> . Stone Crop.		
<i>Stalagmitis cumbogioides</i> . Gamboge.		
<i>Tanacetum vulgare</i> . Tansy (oil of.)		
ACRO-NARCOTIC.		
<i>Æthusa cynapium</i> . Common Fool's Parsley.		
<i>Aconitum napellus</i> . Monkshood.		
<i>Agaricus</i> , five species. Mushrooms (poisonous).		
<i>Amanita muscaria</i> . Truffles (poisonous).		
<i>Anagallis arvensis</i> . Meadow Pimpernel.		
<i>Anda Gomesii</i> .		
<i>Apocynum androsamifolium</i> . Dogsbane.		
<i>Aristolochia clematitis</i> . Birthwort.		
<i>Arnica montana</i> . Leopard's baue.		
<i>Asclepias Syriaca</i> . Swallowwort.		
<i>Atropa belladonna</i> . Deadly Night Shade.		
<i>Æsculus Ohioensis</i> . Buckeye.		
<i>Brucea antidysenterica</i> . False Angustura Bark.		
<i>Cerbera</i> , three species. <i>Cerbera</i> .		
<i>Cherophyllum sylvestre</i> . Bastard Hemlock.		

IRRITANT POISONS.	SYMPTOMS.	TREATMENT.
<i>Chaillotia toxicana</i>. Ratbane. <i>Chenopodium murale</i>. Wormseed. <i>Cicuta maculata</i>. American Hemlock. <i>Cicuta virosa</i>. Water Hemlock. <i>Cissus</i>. Cissus. <i>Cocculus Indicus</i>. Fish berries. <i>Colchicum autumnale</i>. Meadow saffron. <i>Conium maculatum</i>. Hemlock. <i>Coriaria myrtifolia</i>. Myrtle-leaved Sumach. <i>Curare</i>. Indian War Poison. <i>Cynanchum erectum</i>. Cynanchum. <i>Cytisus laburnum</i>. Laburnum. <i>Datura stramonium</i>. Thorn Apple. <i>Digitalis purpurea</i>. Foxglove. <i>Ergot</i>. See <i>Secale</i>. <i>Ervum ervilia</i>. Bitter Vetch. <i>Gaultheria procumbens</i>. Wintergreen (oil of). <i>Hæmanthus toxicarius</i>. <i>Helleborus niger</i>. Black Hellebore. <i>Hypophyllum sanguinaria</i>. Paddock Stool. <i>Ipecacuanha</i>. <i>Ipecacuanha</i>. <i>Lathyrus cicera</i>. Lathyrus. <i>Laurus camphora</i>. Camphor. <i>Lobelia inflata</i>. Indian Tobacco. <i>Lolium temulentum</i>. Darnel. <i>Melia azederach</i>. Pride of China. <i>Mercurialis perennis</i>. Mountain Mercury. <i>Nerum oleander</i>. Common Oleander. <i>Nicotiana tabacum</i>. Tobacco. <i>Oenanthe crocata</i>. Hemlock Dropwort. <i>Passiflora quadrangularis</i>. Barbadoine. <i>Phyostigma venosum</i>. Calabar bean. <i>Fiscidia erythrina</i>. Jamaica Dogwood. <i>Polygala venosa</i> (of Java). <i>Rhus radicans</i>. Poison Vine. <i>Rhus toxicodendron</i>. Poison Oak. <i>Robinia pseudo-acacia</i>. Locust Tree. <i>Ruta graveolens</i>. Rue. <i>Sanguinaria Canadensis</i>. Blood Root. <i>Scilla maritima</i>. Squill. Sea Onion. <i>Secale cornutum</i>. Ergot. Spurred Rye. <i>Sium latifolium</i>. Procumbent Water Parsnep. <i>Spigelia Marilandica</i>. Pink Root. <i>Strychnos Ignatii</i>. St. Ignatius's Bean. <i>Strychnos nux vomica</i>. Nux vomica. <i>Symplocarpus fetida</i>. Skunk Cabbage. <i>Ticunas</i>. Extract of various plants—South America. <i>Tienté</i>. <i>Upas Tienté</i>. Tree of Java. <i>Triticum Hybernum</i>. Wheat (diseased.) <i>Upas antiar</i>. Tree in Java. <i>Veratrum album</i>. White Hellebore. <i>Veratrum viride</i>. American Hellebore. <i>Woorara</i>. War Poison of Guiana. <i>Zea mays</i>. Maize (diseased).	The narcotic vegetable poisons, if taken into the stomach or applied to a wound, occasion the following effects:—stupor; numbness; heaviness in the head; desire to vomit, slight at first, but afterwards insupportable; a sort of intoxication, stupid air, pupil of the eye dilated; furious or lively delirium, sometimes pain; convulsions of different parts of the body, or palsy of the limbs. The pulse is variable, but at first generally strong and full; the breathing is quick, and there is great anxiety and dejection, which, if not speedily relieved, soon ends in death.	The stomach to be effectually evacuated by giving four or five grains of tartar emetic, or from ten to twenty of the sulphate of zinc, repeated every quarter of an hour, till the full effect is produced. These means may be assisted by tickling the throat with a feather or the finger. Large and strong glysters of soap dissolved in water, or of salt and gruel, should be speedily administered, to clear the bowels, and assist in getting rid of the poison; and active purgatives may be given after the vomiting has ceased. When as much as possible of the poison has been expelled, the patient may drink, alternately, a tea-cupful of strong hot infusion of coffee, and vinegar diluted with water. If the drowsiness, which is sometimes extreme, and the insensibility bordering on apoplexy, be not remedied by these means, blood may be taken from the jugular vein, blisters may be applied to the neck and legs, and the attention be roused by every means possible. If the heat declines, warmth and frictions must be perseveringly used. Vegetable acids are on no account to be given before the poison is expelled, and it is desirable that but little fluid of any kind should be administered. N. B. Bromine, chlorine, and iodine are said to be antidotes to the alkaloids generally. Electromagnetism has been found efficacious in the drowsiness and sleep of narcotic poisoning. Where myositic narcotic poisons, as opium, have been taken, mydriatics, as belladonna and stramonium, have been recommended; and conversely.
NARCOTIC. <i>Actæa spicata</i>. Baneberry. <i>Amygdalus communis</i>. Bitter Almond. <i>Amygdalus Persica</i>. Peach. <i>Gelseminum nitidum</i>. Yellow Jessamine. <i>Helonias erythrosperma</i>. Fly Poison. <i>Hyoscyamus albus</i>. White Henbane. <i>Hyoscyamus niger</i>. Black Henbane. <i>Kalmia latifolia</i>. Mountain Ivy. <i>Lactuca virosa</i>. Strong-scented Lettuce. <i>Opium</i>, and its proximate principles. <i>Papaver somniferum</i>. Poppy. <i>Paris quadrifolia</i>. Herb Paris. <i>Prunus Caroliniana</i>. Wild Orange. <i>Prunus lauro-cerasus</i>. Cherry Laurel. <i>Prunus nigra</i>. Black Cherry. <i>Prunus padus</i>. Cluster Cherry. <i>Prunus Virginiana</i>. Wild Cherry. <i>Solanum dulcamara</i>. Bitter-sweet. <i>Sorbus acuparia</i>. Mountain Ash. <i>Taxus baccata</i>. Yew. POISONOUS MUSHROOMS. <i>Agaricus muscarius</i>. Fly Agaric. <i>Agaricus piperatus</i>. Pepper Agaric. <i>Agaricus necator</i>. Deadly Agaric. <i>Agaricus bulbosus</i>. Bulbous Agaric. <i>Agaricus chantarellus</i>. Champignon.	Nausea, heat, and pain in the stomach and bowels, with vomiting and purging, thirst; convulsions, and faintings; pulse small and frequent; delirium; dilated	The stomach and bowels to be cleared by an emetic of tartarized antimony followed by frequent doses of Glauber's or Epsom salt, and large, stimulating glysters. After the poison is evacuated, ether may be administered, with small

POISONS.	SYMPTOMS.	TREATMENT.
	pupil, and stupor; cold sweats, and death. Poisonous mushrooms may be distinguished from the edible by their botanical characters, and by the following criteria. The former grow in wet, shady places, have a nauseous odour; are softer, more open, and porous; have a dirty-looking surface, sometimes a gaudy colour, or many very distinct hues, particularly if they have been covered with an envelope; they have soft, bulbous stalks, grow rapidly, and corrupt very quickly.	quantities of brandy and water; but if inflammatory symptoms manifest themselves, such stimuli should be omitted, and other appropriate means be had recourse to.

2. ANIMAL POISONS.

POISONS.	SYMPTOMS.	TREATMENT.
POISONOUS FISH. <i>Balistes monoceros.</i> Old wife. <i>Cancer astacus.</i> Crawfish. <i>Cancer ruricolus.</i> Land Crab. <i>Clupea thrissa.</i> Yellow-billed Sprat. <i>Coracinus fuscus major.</i> Gray Snapper. <i>Coracinus minor.</i> Hyne. <i>Coryphæna splendens.</i> Dolphin. <i>Mormyra.</i> Blue Parrot Fish. <i>Murena major.</i> Conger Eel. <i>Mytilus edulis.</i> Mussel. <i>Ostracion globellum.</i> Smooth Bottle Fish. <i>Perca major.</i> Barracuda. <i>Perca venenosa.</i> Grooper. <i>Perca venenata.</i> Rock Fish. <i>Physalia.</i> Portuguese Man of War. <i>Scomber caruleus.</i> Spanish Mackerel. <i>Scomber maximus.</i> King Fish. <i>Scomber thynnus.</i> Bonetta. <i>Sparus chrysops.</i> Porgree. <i>Tetrodon sccleratus.</i> Tunny. <i>Tetrodon ocellatus.</i> Blower.	In an hour or two, or often in a much shorter time, after the fish has been eaten, a weight at the stomach comes on, with slight vertigo and headache, sense of heat about the head and eyes, considerable thirst, and often an eruption of the skin (urticaria), and in many cases death. A sharp pain in the wounded part, which soon extends over the limb or body; great swelling, at first hard and pale, then reddish, livid, and gangrenous in appearance; faintings, vomitings, convulsions, and sometimes jaundice; pulse small, frequent, and irregular; breathing difficult, cold sweats, the sight fails, and the intellectual faculties are deranged. Inflammation, and often extensive suppuration and gangrene, followed by death.	An emetic should be speedily administered, or, in the absence of it, vomiting may be excited, by tickling the throat with the finger, and taking large draughts of warm water. After full vomiting, an active purgative should be given, to remove any of the noxious matter that may have found its way into the intestines. Vinegar and water may be drunk after the above remedies have operated, and the body may be sponged with the same. Water, made very sweet with sugar, to which ether may be added, may be drunk freely as a corrective, and a very weak solution of alkali has been recommended to obviate the effects of the poison. If spasm ensue; after evacuations, laudanum, in considerable doses, is necessary. If inflammation should occur, the usual means of removing it must be employed. A cupping-glass to be applied over the wound, or a moderately tight ligature above the bites, and the wound left to bleed after being well washed with warm water; the actual cautery, lunar caustic, or butter of antimony, to be then applied freely to it, and afterwards covered with lint, dipped in equal parts of olive oil and spirit of hartshorn. The ligature to be removed if the inflammation be considerable. Warm, diluting drinks, and small doses of ammonia or hartshorn, to cause perspiration; the patient to be well covered in bed, and a little warm wine given occasionally. If gangrene be threatened, wine may be given more freely, and the bark should be had recourse to. Arsenic, the principal ingredient in the Tanjore Pill, has been strongly recommended, and, also, Bibron's antidote.
POISONOUS SERPENTS. <i>Boa crotaloides.</i> Copperhead. <i>Cenchris mockeson.</i> Mockeson. <i>Cerastes nasicornis.</i> Horned Viper of Western Africa. <i>Coluber berus.</i> Viper. <i>Coluber prester.</i> Black Viper. <i>Crotalus, (5 species.)</i> Rattle Snake. <i>Scytale piscivorus.</i> Water Viper.	Nauseous odour of the breath; acrid taste; burning heat in the throat, stomach, and abdomen; frequent vomitings, often bloody, with copious bloody stools; excruciating pain in the stomach; painful and obstinate priapism, with heat in the bladder, and strangury or retention of urine; frightful convulsions, delirium and death.	Vomiting to be excited by drinking sweet oil, sugar and water, milk, or linseed tea, very freely. Emollient glysters should be administered, and if symptoms of inflammation of the stomach, kidney, or bladder supervene, they must be subdued by appropriate treatment. Camphor dissolved in oil may be rubbed over the belly and the thighs.
CANTHARIS VESICATORIA. Spanish, or Blistering Fly. <i>Lytta vittata.</i> Potato Fly.	In general, the sting of these insects occasions only a slight degree of pain and swelling; but occasionally the symptoms are more violent, and sickness and fever are produced by the intensity of the pain.	Hartshorn and oil may be rubbed on the affected part, and a piece of rag, moistened in the same, or in salt and water, may be kept upon it till the pain is removed. A few drops of hartshorn may be given frequently in a little water, and a glass or two of wine may be taken. The sting may, in general, be removed by making strong pressure around it with the barrel of a small watch-key.
VENOMOUS INSECTS <i>Tarantula.</i> <i>Scorpio.</i> Scorpion. <i>Vespa crabro.</i> Hornet. <i>Vespa vulgaris.</i> Wasp. <i>Apis mellifica.</i> Bee. <i>Culex pipiens.</i> Gnat. <i>Æstrus bovis.</i> Gad-fly.		

	SYMPTOMS.	TREATMENT.
SALIVA OF THE RABID DOG.	At an uncertain interval after the bite, generally, however, between the twentieth day and three or four months, pain or uneasiness occurs in the bitten part, though the wound may have been long healed. Anxiety, uneasiness, languor, spasms, horror, disturbed sleep, difficult respiration succeed, and are soon very much increased; violent convulsions affect the whole body, hideously distorting the muscles of the face; the eyes are red and protruded, the tongue swells and often hangs out, and viscid saliva flows from the mouth; there is pain in the stomach, with bilious vomitings, a horror of fluids, and impossibility of drinking them. All these symptoms are aggravated till the sufferer is relieved by death.	Hydrophobia is more easily prevented than cured; indeed, it is doubted whether it ever has been cured. Mercury, arsenic, opium, musk, camphor, acids, wine, vegetable and mineral alkali, oil, various herbs, and many other remedies whose effects are quite opposite, have been employed, but none can be relied on. Large blood-lettings, the warm and cold bath, and almost every other remedial agent, have been tried without success. The bitten part should be completely cut out, even after it has healed, if the symptoms have not yet come on; the part should then be immersed in warm water, or be washed with it as long as it will bleed, and, after the most persevering ablation, caustic should be applied to every part of the surface, and then the wound covered with a poultice, and suffered to heal by granulation. No milder discipline can ensure safety.

The most energetic poisons are used therapeutically, and often with considerable advantage. They have, of course, to be administered in extremely small doses, to avoid producing true poisoning.

POISON, ARROW, see Arrow poison.

POISON BERRY-TREE, *Melia azedarach*.

POISON DE FLÈCHES, Arrow poison.

POISON, HEMLOCK, *Conium*.

POISON, ITALIAN, *Liquor arsenicalis* — p. Morbid, Virus.

POISON OAK, *Rhus toxicodendron* — p. Root, *Cicuta maculata*.

POISON, SAUSAGE, *Allantotoxium* — p. Vegetable, *Urtica*.

POISON VINE, *Rhus radicans*.

POISONING, *Veneficium*, *Toxicatio*, *Intoxicatio*, *Entoxicismus*, *Entoxicismus*, *Pharmacologia*, *Maleficium*, *Venenatio*, (F.) *Empoisonnement*, *Intoxicacion*. The art of applying a poisonous substance to any of the textures in a dose capable of occasioning more or less serious symptoms. Rigorously speaking, therefore, if a physician, in treating any case, employs a deleterious medicine in a small dose, which, owing to particular circumstances, produces unpleasant symptoms, he may be said to have poisoned the patient. The term is, however, generally restricted to poisoning by design.

Poisoning has also been used, by a few, to denote the series of symptoms resembling those produced by certain poisons, which occasionally follow the use of substances that are generally innoxious. Overloading the stomach with indigestible food, has, for example, given occasion to symptoms resembling those produced by a narcotic poison, and the effect has been termed *poisoning*.

Secret poisoning was, at one time, practised as an art. It consisted in administering medicines, which should act as *slow poisons*, and gradually undermine the strength and life. Its agency was far from being so extensive as was imagined.

POISONING, CROWD, *Ochlesis*.

POISONOUS, *Toxical*, *Noxius*, (F.) *Toxique*, *Vénéneux*. Having the qualities of a poison.

POISONS, ACRO-NARCOTIC, p. narcotico-acrid — p. Corrosive, p. acrid — p. Escharotic, p. acrid — p. Irritating, p. acrid — p. Putrescent, p. septic. See poison.

POITRINAIRE, Phthisicus.

POITRINE, (from [L.] *pectus*, gen. *pectoris*), Thorax.

POIVRE. Piper — p. *d'Eau*, *Polygonum hydro-piper* — p. *d'Espagne*, *Capsicum annuum* — p. *de Guinée*, *Capsicum annuum* — p. *d'Inde*, *Capsicum annuum* — p. *de Jamaïque*, see *Myrtus pimenta* — p. *Long*, *Piper longum* — p. *de Moine*, *Vitex* — p. *Noir*, *Piper Nigrum* — p. *Petit*, *Vitex* — p. *à Queue*, *Piper cubeba* — p. *Sauvage*, *Vitex*.

POIX, (from [L.] *pix*), Pitch, see *Pinus sylvestris* — p. *Blanche*, see *Pinus abies* — p. *De Bourgogne*, see *Pinus abies* — p. *Grasse*, see *Pinus abies* — p. *Jaune*, see *Pinus abies* — p. *Liquide*, see *Pinus sylvestris* — p. *Minérale*, *Pissasphaltum* — p. *Minérale scoriacée*, *Asphaltum* — p. *Navale*, see *Pinus sylvestris* — p. *Noire*, see *Pinus sylvestris*.

POKE, *Phytolacca decandra* — p. Indian, *Veratrum viride* — p. Stink, *Dracontium fœtidum* — p. Weed, *Phytolacca decandra*, *Veratrum viride*.

POLANIS'IA GRAVE'OLENS, *Clammyweed*, *Stinkweed*, *Wormweed*, *Clammy Mustard*, *False mustard*, (πολις, 'many,' and *avicos*, 'unequal,' in reference to its stamens.) An indigenous plant, Ord. Capparidaceæ, *Sex. Syst.* Dodecandria Monogynia, which blossoms in summer, and is found in every part of North America. To the whole plant have been ascribed anthelmintic properties.

POLAR PLANT, *Silphium laciniatum*.

POLECAT WEED, *Dracontium fœtidum*.

POLEMONIUM REP'TANS, *False Jacob's Ladder*, Greek *Valerian*, *Siccatroot*. Order, Polemoniaceæ; indigenous, flowering in May; has been used as a diaphoretic and expectorant.

POLENTA, (*pollen*, 'fine flour,') Alphon.

POLES, (from πολες, 'a pole, axis,') Chalazæ.

POLEY MOUNTAIN OF CANDY, *Teucrium Creticum* — p. Mountain of Montpellier, *Teucrium capitatum*.

POLIANTHES TUBERO'SA, (πολις, 'a town,' and *avros*, 'a flower,') *Amaruacahu*, Ord. Liliaceæ. The root is cut into slips, and laid upon the bites of serpents, by the Peruvian Indians.

POLIA'TER, (πολις, 'a town,' and *iatros*, 'a physician,') A physician exercising his profession in a town, by the appointment of the government.

POLICE', MED'ICAL, (πολιτεια, the administration of a state or town,) *Politica medica*, *Poliobiotherapia*, *Medicina politica*, *Ordo medicinalis*, *Political medicine*, (F.) *Police médicale* ou *sanitaire*. Hygienic rules established, or

which ought to be established, for the prevention of disease, and to contribute to the progress of medicine. The salubrity of districts, especially during the existence of spreading disease, is greatly ensured by an efficient medical police.

POLICLINICA, (πολις, 'a town,' and κλινη, 'a bed.') Treatment of diseases in a town. A city or town clinic or hospital.

POLIO'SIS, *Tricho'sis polio'sis*, *Spilo'sis polio'sis*, *Pol'ioties*, *Canit'ies*, (πολιωσις, from πολιος, 'hoary, gray.') Hair prematurely gray or hoary. The state of becoming gray, *Grayhaired*, *Polio'trichos*.

POLIOTES, (πολιοτης,) Poliosis.

POLIOTRICHOS, (πολιος, 'gray,' and τριξ, gen. τριχος, 'hair,') see Poliosis.

POLITIA MEDICA, Police, medical.

POLITOBIOThERAPIA, (πολιτη, 'a citizen,' βιος, 'life,' and θεραπεια, 'medical treatment,') Police, medical.

POLIUM CAPITATUM, (πολιος, 'gray, whitish,') *Teucrium capitatum*—p. *Creticum*, *Teucrium Creticum*—p. *Montanum*, *Teucrium capitatum*.

POLL, ([D.] Bol.) Head.

POLL'EN, gen. *Pol'linis*, *Pol'lis*, (παλη, 'dust,') The fecundating dust contained in the anthers of flowers, which is dispersed on the stigma. It is conveyed to the ovary through the stigma by the *Pollen tube*, *Tu'bus pollin'icis*, (F.) *Boyan Pollinique*.

POL'LEX, gen. *Pol'licis*, *Ant'icheir*, *Prom'anus*, *Dig'itus primus* seu *mag'nus*, *Al'tus pollex*, *Thumb*, (Prov.) *Thame*, (from *pollere*, 'to have much strength,') (F.) *Pouce*. The first of the fingers in size and strength. It is asserted that the term *poltroon* is derived from *pol'licē trunco'to*—the Romans who were desirous of being exempted from military service being accustomed to cut off the thumb, and thus render themselves unfit to handle arms.

POLLEX, see *Digitus*.

POLLEX PE'DIS, *Hallus*, *Hallux*. The great toe. (Prov.) *Tomtoe*.

POLLINC'TOR, *Pollinc'ter*, (*pollingere*, *pollinctum*, 'to lay out a corpse,') In antiquity, an anointer, dresser, or embalmer of the dead.

POLLINCTURA, Embalming.

POLLIS, Pollen.

POLLOCK, see *Oleum jecoris aselli*.

POLLOD'IC, *Polloed'icus*, (πολος, 'many,' and 'ολος, 'a way,') An epithet proposed by Dr. Marshall Hall for a course of nervous action proceeding from one point, in many directions, to every other.

POLLOM, LITTLE, *Polygala paucifolia*.

POLLUTION, *Pollu'tio*, (*polluo*, *pollutum*, 'to profane,') The excretion of the seminal liquor or sperm, or the production of orgasm in male or female, at other times than during coition. When occasioned by a voluntary act, it is called simply *Pollution* or *Masturba'tion*, (F.) *Souillure*; when excited during sleep, by lascivious dreams, it takes the name *Noctur'nal pollution*, *Ezono'ro'sis*, *Onei'rog'mos*, *Onei'rog'onos*, *Onei'rogonor'ra'a*, *Ezono'rog'mus*, *Spermatoclem'ma*, *Spermatoclep'sis*, *Spermatoclep'sis*, *Spermatolip'sis*, *Gonor'ra'a dormien'tium* seu *onei'rog'onos* seu *er'a* seu *libidino'sa*, *Proflu'vium sem'inis*, *Spermatorr'ha'a*, *Paron'ir'ia ea'la'z*, *Night pollution*, (F.) *Pollution involontaire*.

POLLUTION INVOLONTAIRE. Pollution—p. *Nocturnal*, Pollution—p. *Self*, *Masturbation*—p. *Voluntary*, *Masturbation*.

POLONYCH'IA, *Polyonych'ia*, (poly, and ονχ, gen. ονχος, 'a nail,') The condition of having supernumerary nails,—a rare congenital monstrosity.

POLYFOOT, see *Kylosis*.

POLY, *Poly*, (πολος,) 'many, much, full,' in composition:

POLYACTIUM TRISTE, *Pelargonium triste*.

POLYÆMATUS, (πολυαιματος,) *Plethoric*.

POLYÆMIA, (πολυαιμα, from poly, and αιμα, 'blood,') *Plethora*.

POLYÆMIC, *Plethoric*.

POLYÆMICUS, *Plethoric*.

POLYANHÆMIA, *Anæmia*.

POLYANHÆMIE, *Anæmia*.

POLYBLÉN'NIA, (poly, and βλεννα, 'mucus,') Excessive secretion of mucus.

POLYBRACH'IA, (poly, and βραχιων, 'an arm,') A monstrosity characterized by one or more supernumerary arms.

POLYBREP'HA, (poly, and βρεφος, 'a fœtus, a child,') *Pregnancy*, compound.

POLYCARP'Æ, (poly, and καρπος, 'fruit,') A division of cutaneous diseases of the family *Eczematosen*, of Fuchs, including *Psoriasis*, *Lichen*, *Herpes*, and *Impetigo*.

POLY'CERAS, (poly, and κρας, 'horn,') A monstrous animal having many horns.

POLYCHOL'IA, *Perichol'ia*, *Hyperchol'ia*, (poly, and χολη, 'bile,') *Superabundance of bile*; such as is supposed to exist in bilious fevers.

POLYCHREST'US, (poly, and χρηστος, 'useful,') Having many virtues. An ancient name for certain medicines considered to be useful in many diseases.

POLYCHRONIUS, (poly, and χρονος, 'time,') *Chronic*.

POLYCHY'LLIA, (poly, and χυλος, 'juice,') *Polychy'mia*. Fulness of juices.

POLYCHYMIA, *Polychylia*—p. *Sanguinea*, *Plethora*.

POLYCLIN'ICA, (poly, and κλινη, 'bed,') Provided with many beds;—as an hospital.

POLYCOR'IA, (poly, and κορη, 'pupil,') The state in which there is more than one pupil.

POLYCRINOSES, (poly, and κρινω, 'to separate, and o'is, see *Proflu'vium*).

POLYCYE'SIA, *Polycy'e'sis*, (poly, and κυσις, 'pregnancy,') Frequent or quickly recurring pregnancies: also, compound pregnancy.

POLYCYTHÆ'MIA, (poly, κυτος, 'a cell,' and αιμα, 'blood,') A condition of the blood in which there is an increase of the red corpuscles.

POLYDACTYLUS, *Perissodact'ylus*, (πολυδάκτυλος, from poly, and δακτυλος, 'a finger,') One who has a supernumerary finger or fingers, or toes.

POLYDIP'SIA, (poly, and διψη, 'thirst,') *Sit'is morbo'sa* seu *inten'sa*, *Polydip'sia*, *Dipso'sis a'cens*, *Excessive thirst*, (F.) *Soif excessive*. A constant desire for drinking, with a sense of dryness in the mouth and throat. It is mostly symptomatic of fever, inflammation, dropsy, excessive discharges, &c., and is so much a part of diabetes, that it has been called *Polydipsia*.

POLYG'ALA, (πολυγαλον, from poly, and γαλα, 'milk,') *P. vulga'ris*, *Com'mon Milk'wort*. Ord. *Polygalaceæ*. See *Syst. Diadelphia Octandria*. The leaves are bitter, and are used in the same cases as the *Polygala amara*.

POLYGALA AMÁ'RA, *P. amare'lla* seu *vulga'ris*, *Bitt'er Milk'wort*. A powerful bitter; and, consequently, tonic. Its reputed virtues are balsamic, demulcent, and corroborant. The *Polygala rubell'a*, or *Bitt'er Polyg'ala* of the United States, has the same virtues.

POLYGALA AMARELLA, *P. amara*—p. *Bitter*, see *P. amara*—p. *Grandiflora*, *P. senega*.

POLYGALA PAUCIFOL'IA, *Trickl'oper'ma grandifl'o'ra*, *Dearf' Milk'wort*, *Little Poll'om*, *Ever-green Snake'root*, (F.) *Polygale na'ive*, possesses the same properties as *P. Senega*.

POLYGALA RUBELLA, see *Polygala amara*.

POLYGALA SÉN'EGA, *P. Virginia'na* seu *grandifl'o'ra*, *Sen'e'ka*, *Rattlesnake Milk'wort*, *Sen'ega*, or *Rattlesnake root*, (F.) *Polygale de Virginie* ou

de Senega. The bark is the active part of the root, *Senega* (Ph. U. S.). Its taste is at first acrid, and afterwards very hot and pungent. It was once esteemed to be specific against the poison of the rattlesnake, and in pleurisy, pneumonia, &c., after the inflammatory action had been got under. It has been given also in humoral asthma, chronic rheumatism, dropsy, croup, amenorrhœa, &c. According to Peschier, it contains three principles, *Polygalin* or *Senegin*, *Isol'usin*, and *Polygalic Acid*. Dose, gr. 30 to 40 of the powder.

POLYGALA SERPENTARIA. A shrub of Caffra-ria, the root of which is considered by the natives as a sure antidote against the bites of venomous serpents.

POLYGALA VIRGINIANA, *P. senega*—p. *Vulgaris*, *P. amara*.

POLYGALACTIA, *Polygal'ia*, (*poly*, and *γαλα*, gen. *γαλακτος*, 'milk.') Superabundant secretion of milk; galactia.

POLYGALÉ NAINE. *Polygala paucifolia*—p. *de Virginii*, *Polygala Virginiana*.

POLYGALIA, *Polygalactia*.

POLYGALIN, see *Polygala senega*.

POLYGONATUM, (*πολυγονατον*, from *poly*, and *γωνη*, 'a knee or joint,') *Convallaria polygonatum*—p. *Acneops*, *Convallaria polygonatum*—p. *Caniculatum*, *Convallaria canaliculata*—p. *Latifolium*, *Convallaria canaliculata*—p. *Multiflorum*, *Convallaria multiflora*—p. *Pubescens*, *Convallaria pubescens*—p. *Uniflorum*, *Convallaria polygonatum*—p. *Vulgare*, *Convallaria polygonatum*.

POLYGONIA, (*πολυγονια*, from *poly*, and *γενωαι*, 'to produce,') Fecundity.

POLYGONIUS, (*πολυγωνιος*,) *Polygonus*.

POLYGONUM, (*πολυγονον*,) *P. aviculare*, (from *poly*, and *γωνη*, 'a knee or joint,') *P. centinodium* seu *erectum*, *Centinodia*, *Centumodia*, *Sanguinalis* *Corrig'isla*, *Sanguinalia*, *Knot'grass*, *Knot'-weed*, *Bind'-weed*, *Door'-weed*, (F.) *Renouée*, *Trainasse*, *Centinode*. Ord. *Polygonaceæ*. Sex. Syst. Octandria Trigynia. This grass is said to be a feeble astringent.

POLYGONUM ACRE, *P. hydropiper*.

POLYGONUM AMPHIBIUM, *P. coccineum*, *Water Persicaria*, of Europe and this country. The roots have been used like sarsaparilla, especially in diseases of the skin.

POLYGONUM BISTORTA, *P. bistortoides*, *Bistorta*, *Colubri'na*, *Offic'inal Bistort*, *Great Bistort* or *Snake'-weed*, (F.) *Bistorte*, *Couleuvreine*, *Serpentaire rouge*. The root is a powerful astringent and tonic; and, as such, has been used in internal hemorrhage, atonic diarrhœa, ague, &c. Dose, gr. xv to ʒj.

POLYGONUM BISTORTOIDES, *P. bistorta*—p. *Centinodium*, *P. aviculare*.

POLYGONUM COCCINEUM, *P. amphibium*.

POLYGONUM DIVARICATUM; *East'ern Buck'-wheat plant*. The roots, reduced to a coarse meal, are said to be much used as an article of diet by the Siberians.

POLYGONUM ERECTUM, *P. aviculare*.

POLYGONUM FAGOPYRUM, *Fagopyrum*, *F. esculentum*, *Trit'icum Fagi'num*, *Buck'-wheat*, (F.) *Sar'raisin*, *Blé noir*. The grain of this plant constitutes a principal article of food with the inhabitants of Russia, Germany, and Switzerland. It is much used in the United States.

POLYGONUM HYDROPIPER, *P. acris* seu *hydro-piperoides* (Pursh) seu *punctatum*, *Hydropiper*, *Persicaria urens*, *Mercurius terrestris*, *Poor Man's Pepper*, *Bit'ing Ase'smart*, *Smart'-weed*, *Lake'-weed*, *Water'-pepper*, (F.) *Poivre d'eau*, *Persicaire poivrée* ou *brûlante* ou *âcre*, *Piment d'eau*, *Ourage*, *Renouée âcre*. The leaves of this plant have an acrid, burning taste, and resemble those of the arum. They have been recommended as possessing antiseptic, aperient, and diuretic vir-

tues. The fresh leaves have been applied externally as a stimulating cataplasma.

POLYGONUM HYDROPIPEROIDES, *P. Hydropiper*, *P. punctatum*. *Polygonum hydro-piperoides* of Michaux, *mild water'-pepper*; indigenous; has no acrid properties.

POLYGONUM PERSICARIA, *Persicaria*, *P. minor* seu *nit'is*, *Plumbago*, *Lat'dy's Thumb*, *Heart's ease*, (Prov.) *Red'shank*, (F.) *Persicaire douce*. This plant is said to possess vulnerary and antiseptic qualities. It has been given in wine to arrest gangrene.

POLYGONUM PUNCTATUM, *P. hydro-piperoides*, *Water'-pepper* of America, *Smart'-weed*, possesses similar properties to *Polygonum hydropiper*, and is occasionally used internally in gravel, and externally in chronic ulcers. A fluidrachm of the saturated tincture has been extolled in amenorrhœa. See *Polygonum hydropiper*.

POLYGONUM VIRGINIANUM, *Wand-spiked Persicaria*, *Virgin'ia Bistort*; indigenous; is astringent, diuretic, and tonic.

POLYGONUS, *Polygon'nus*, (*poly*, and *γωνια*, 'an angle,') *Polygonal*; multangular,—hence *Os polygonum*.

POLYGROMA, gen. *Polygro'matis*, (*poly*, and *hygro'ma*,) *Staphylo'ma*.

POLYHIDRIA, *Polyhid'ria*, *Polyhidrosis*, (*poly*, and *ἵδρω*, 'sweat,') Excessive sweating.

POLYHYDRIA, (*πολυῖδρα*, from *poly*, and *ἵδρω*, 'water,') *Hydrops*.

POLYHYPERHEMIA, *Plethora*.

POLYIDRIA, *Polyhidria*.

POLYIDROSIS, *Polyhidria*.

POLYLEUCOCYTHÆMIA, *Leucocythæmia*.

POLYLYMPHIA, *Anasarca*.

POLYMASTHUS, *Pleiomasthus*, (*poly*, and *μασθος*, 'breast,') (F.) *Plurimamme*. A monster having several breasts. The condition has been called *Polymast'ia*, *Pleiomast'ia*, and *Plesma'zia*.

POLYMAZIA, (*poly*, and *μαζος*, 'a breast,') see *Polymasthus*.

POLYMELE'IAN, (F.) *Polymélien*, (*poly*, and *μελος*, 'a limb,') An epithet given to monsters characterized by the insertion, in a well formed subject, of one or more accessory limbs, accompanied, at times, by the rudiments of certain other parts.

POLYMERIA, *Polymerismus*—p. *Promiscua translative*, *Transposition of the Viscera*.

POLYMERISMUS, *Polymer'ia*, (*poly*, and *μερος*, 'a part,') A monstrosity, which consists in the existence of supernumerary organs; as a sixth finger of the hand.

POLYNEURON, *Plantago*.

POLYONYCHIA, *Polonych'ia*.

POLYOP'SIA, *Polyop'sy*, *Polyop'ia*, *Polyop'py*, (*poly*, and *οψει*, 'sight,') Vision is so called, when multiple or manifold.

POLYOPY, *Polyops'ia*.

POLYORCHIS, *Multites'tis*, (*poly*, and *ορχις*, 'a testicle,') One who has more than two testicles.

POLYOREX'IA, (*poly*, and *ορεξις*, 'appetite,') Excessive appetite or hunger. A genus in the *Gastroes* of M. Alibert.

POLYPE'RIA, (*poly*, and *προς*, 'mutilated,') Congenital misconstruction of various parts or organs.

POLYPHA'GIA, *Multivoran'tia*, (*πολυφαγια*, from *poly*, and *φαγω*, 'I eat,') Voracity. Faculty of eating or subsisting on many kinds of food. See *Boulimia*.

POLYPHAGOUS, (*πολυφαγος*,) *Polyph'agus*, *Multivorous*, (F.) *Polyphage*. Relating or appertaining to polyphagia.

POLYPHARMAC'IA, *Polypharm'acy*, (*poly*, and *φάρμακον*, 'a medicine,') The prescribing of too many medicines. A prescription, consisting of a number of medicines; hence the name *Poly*

pharmacus, (F.) *Polypharmaque*, given to one who is in the habit of prescribing a number of medicines, and whose prescriptions are loaded with ingredients. The term is taken in bad part.

POLYPHARMAQUE, see Polypharmacia.

POLYPHYSIA, (*poly*, and *φύσα*, 'wind,') Flatulence.

POLYPIFORM CONCRETIONS, (*polypus*, and *forma*, 'form,') see Polypus.

POLYPIONIA, (*poly*, and *πιον*, 'fat,') see Polysarcia—*p.* Infantum, see Polysarcia.

POLYPIOSIS, (*poly*, *πιον*, 'fat,' and *osis*,) see Polysarcia.

POLYPLASTIC, *Polyplasticius*, (*poly*, and *πλασσω*, 'I form,') That which has many shapes.

POLYPLASTIC ELEMENT, in histology, is one which does not retain its primary shape; but undergoes transformation into others.

POLYPOD'IA, (*poly*, and *πους*, gen. *ποδός*, 'a foot,') A monstrosity in which there are more than two feet.

POLYPODITES, (*πολυποδιτης*,) Wine impregnated with polypody.

POLYPOD'IDIUM, (*πολυποδιον*,) *P. Vulgarē* seu *Virginianum* seu *pinnatifidum*, *Filix a dulcis*, *Common Polypody*, *Fern root*, *Rock Brake* or *Bracken*, *Brake* or *Bracken root*, (F.) *Fougère douce*. Ord. Filices. (From *poly*, and *πους*, gen. *ποδός*, 'a foot,' owing to its many roots.) This fern grows in the clefts of rocks, walls, &c., in both hemispheres. It was formerly regarded as a cholagogue, and was given in head diseases. It is not used now.

POLYPODIUM ADIANTIFORME, see *Calagualæ radix*—*p.* *Ammifolium*, see *Calagualæ radix*—*p.* *Argentum*, see *Calagualæ radix*—*p.* *Calaguala*, *Calagualæ radix*—*p.* *Cetaracium*, *P. incanum*—*p.* *Coriaceum*, see *Calagualæ radix*—*p.* *Dentatum*, *Asplenium filix fœmina*—*p.* *Filix fœmina*, *Asplenium filix fœmina*.

POLYPODIUM FILIX MAS, *Aspid'ium* seu *Athyrium* seu *Nephrodium* seu *Polystichum Filix Mas*, *A. depas'tum* seu *eros'tum*, *Filix pinna'ta*, *Pter'is*, *Alechl'non*, *Polyrrhizon*, *Blanc'non* *Oribas'ii*, *Lonic'itis*, *Filic'ula*, *Filix Mas*, *F. non ramo'sa* *denta'ta*, *Nephrodium crenatum*, *Polystichum callipteris*, *Male Polypody* or *Fern*, (F.) *Fougère mâle*. The rhizoma of this plant, *Filix Mas*, Ph. U. S., *Filix*, Ph. B. (1864), has been much extolled as an anthelmintic, and in cases of tœnia, and has, by many, been placed above all tœniacuges. It was the basis of a once celebrated remedy—*Madame Nouffer's*. Dose, ʒij to ʒiij of the powdered solid part of the root; followed in the morning by a strong cathartic of camboe or jalap. An ethereal extract has been proposed by Dr. Peschier, of Geneva,—*Extractum filicis æthereum*, or *Oleum filicis*, the dose of which is from 12 to 24 grains. *Extractum filicis liq'uidum* is official in the Ph. B. (1864). Dose, ℞xxx to fʒj.

POLYPODIUM INCANUM, *P. cetarac'ium*, *Aeros'tichum polypodioides*, *Polypody*, *Rock Brake* or *Bracken*; indigenous; has been considered demulcent, cathartic and anthelmintic.

POLYPODIUM INCISUM, *Asplenium filix fœmina*—*p.* *Molle*, *Asplenium filix fœmina*—*p.* *Pennatifidum*, *P. vulgare*—*p.* *Politum*, see *Calagualæ radix*—*p.* *Trifidum*, *Asplenium filix fœmina*—*p.* *Virginianum*, *P. vulgare*—*p.* *Vulgare*, *Polypodium*.

POLYPODY, *Polypodium*, *P. incanum*—*p.* *Common*, *Polypodium*—*p.* *Male*, *Polypodium filix mas*.

POLYPOID, *Polypoi'des*, (*polypus*, and *ειδος*, 'resemblance,') Having the shape of, or resembling, a polypus.

POLYPORE DU MÊLÈZE, *Boletus laricis*.

POLYPORUS IGNIARIUS, (*poly*, and *πορος*,

'a pore,') *Boletus ignarius*—*p.* *Officinalis*, *Boletus laricis*.

POLYPO'SIA, (*πολυποσια*, from *poly*, and *ποσις*, 'drink,') Great thirst; desire for drink. *Polydipsia*.

POLYPO'TOME, *Polypot'omus*, *Pol'ypotome*, *Poly'p'tomus*, *Pol'ypus knife*, (*polypus*, and *τμήναι*, 'to cut,') An instrument for the removal of polypus by excision.

POLYPOUS, *Polyppo'eus*. Relating, or belonging to, or resembling polypus.

POLYPTOME, *Polypotome*.

POLYYPUS, (*πολυπους*,) *Pseudometamorpho'sis polypo'sa*, (from *poly*, and *πους*, gen. *ποδός*, 'a foot,') A name given to tumours, which occur in mucous membranes especially; and which have been compared to certain zoophytes. Polypi may form on every mucous membrane. They are most commonly observed in the nasal fossæ, *Rhinopol'ypii*, pharynx, and uterus; more rarely in the stomach, intestines, bronchial tubes, bladder, and meatus auditorius externus. They vary much in size, number, mode of adhesion, and intimate nature. Accordingly, they have received various appellations; *mucous*, *soft*, *vesic'ular*, when their substance has been soft, spongy, vesicular, and, as it were, gorged with fluid. Others have been called *hard*; and these have been distinguished into *fib'rous* or *flesh'y*, and into *scir'rhus* or *cancerous*. *Fib'rous polypi*, *Pol'ypii fibro'si*, *Inopol'ypii*, are of a dense, compact texture, and whitish colour. They contain few vessels, and do not degenerate into cancer. The *scir'rhus* or *carcinom'atous* are true cancerous tumours, painful and bleeding. They present all the pathological changes which belong to cancerous affections, and the prognosis is of course unfavourable.

The mode of treating polypi must differ according to their character. The means used to destroy them have been, 1. To subject them to the action of certain astringent powders or solutions, to obtain their resolution. 2. *Cauterization* or the application of the actual cautery or caustics. 3. *Excision* or ablation with a cutting instrument. 4. *Extirpation* with the fingers or with the polypus forceps. 5. A *ligature*, put around them so as to prevent their nutrition. A seton has, also, been sometimes used to gradually destroy the pedicle.

The term *POL'YPI*,—*Pseudopol'ypii*, *Carditis polypo'sa*, *Polypiform Concretions*, (F.) *Concretions polypeuses ou polypiformes ou sanguines du Cœur*, *Hémocardioplasties*, (Piorry,)—has likewise been applied to collections of blood—fibrinous concretions—found in the heart—*Heart clots*—or great vessels after dissolution. The presence of these concretions during life may be suspected, when in the course of an acute or chronic affection of the heart, or at the close of chronic diseases in general—especially those of the lungs—the heart's action becomes suddenly tumultuous, obscure, and accompanied with a soft bellows' murmur, whilst the general symptoms indicate the effects of obstructed circulation. The right cavities are most frequently affected.

POLYPUS BRONCHIA' LIS, *Bronchlemm'itis*, *Bronchi'tis membrana'cea* seu *exsudati'va seu croupo'sa*, *B. plas'tic*, *Bronch'ial Croup*, *Chron'ic Croup* [?], *Pseudomembranous Bronchitis*, (F.) *Bronchitis pseudo-membranæuse*. A membraniform secretion in the bronchial tubes of a diphtheritic character, having the shape, when expectorated, of the bronchia. It is the result of inflammation of the lining membrane of those tubes.

POLYPUS CARNOSUS, *Sarcoma*—*p.* *Fibrosus*, see *Polypus*—*p.* *Uteri*, *Metropolypus*.

POLYRRHIZON, (*πολυρριζον*, from *poly*, and *ρίζα*, 'a root,') *Polypodium filix mas*.

POLYRRHŒA, (*poly*, and *ῥέω* 'to flow,') *Profluviu'm*.

POLYSAR'CIA, *Polysarcosis*, (πολυσαρκία, from *poly*, and *σάρξ*, gen. *σάρκος*, 'flesh.') *Carnositas*, *Polysomatia*, *Adelipar'ia*, *Corpulentia*, *Steatites*. Excessive corpulency. Authors have distinguished two varieties of this:—**POLYSAR'CIA CARNO'SA**, *Corpulency*, and **P. ADIPO'SA**, *Polypio'nia*, *Polypio'sis*, *Pimelo'sis nim'ia*, *Pachys'mus*, *Pachyn'sis*. *Obes'itas*, *Hyperpim'ele*, *Lipo'ria*, *Adipo'sis*, *Polysar'cia fugino'sa*, *Empyme'ium polysar'cia*, *Baryso'mia*, *Barysomatic'a*, *Lipo'sis*, *Lipar'otes*, *Lipasp'ma*, *Pinguitu'do*, *Pinguitude*, *Obes'ity*, (F.) *Obésité*, *Embonpoint excessif*. A severe regimen and exercise are the best preventives. Children are occasionally met with of unusual fatness, in spite of every care. This morbid disposition has been termed *Polypio'nia infantum* and *Chloro'sis gigante'a*.

POLYSARCIA VISCERUM, *Physconia adiposa*. **POLYSARCOSIS**, *Polysarcia*. **POLYSCEL'IA**, (*poly*, and *σκελος*, 'a leg.') A monstrosity, characterized by the presence of one or more supernumerary legs.

POLYSIA'LIA, (*poly*, and *σialον*, 'saliva.') Excessive discharge of saliva.

POLYSOMAT'IA, (*poly*, and *σωμα*, 'body,') *Polysarcia*.

POLYSO'MIA, (*poly*, and *σωμα*, 'body,') A monstrosity characterized by the existence of more bodies than one.

POLYSPAS'TON, (*poly*, and *σπασ*, 'I draw.') A powerful instrument for reducing luxations.

POLYSPER'MIA, (πολυσπερμια, from *poly*, and *σπερμα*, 'sperm.') Richness or fulness of sperm or seed.

POLYSTHEN'IC, *Polysthen'icus*, (πολυσθενής, from *poly*, and *σθενος*, 'strength,') Having much strength.

POLYSTICUM CALLITERIS, *Polypodium filix mas*—*p. Filix mas*, *Polypodium filix mas*.

POLYSTOMA PINGUICOLA, (*poly*, and *στομα*, 'a mouth,') see *Worms*—*p. Venarum*, see *Worms*.

POLYTOCIA, (πολυτοκία, from *poly*, and *τοκος*, 'a child,') *Fecundity*.

POLYTRIC, *Asplenium trichomanoides*.

POLYTRICHIA, see *Polytrichus*.

POLYTRICHUM, (πολυτριχον, from *poly*, and *τριχος*, gen. *τριχος*, 'hair,') *P. vulga'r'e*, *Polytrichon*, *Adiantum au'reum*, *Gold'en Maid'enhair*, (F.) *Perce-mousse*. *Ord. Musci*. It is possessed of astringent properties, and was formerly given in diseases of the lungs, and in calculous affections.

POLYTRICHUM COMMUNE, *Asplenium trichomanoides*.

POLYTRICHUM JUNIPER'INUM, *Hair'cap moss*, *Rob'bin's Rye*, common in New England, is said to possess powerful diuretic properties.

POLYTRICHUS, (πολυτριχος.) Very hairy. One that has much hair. The condition is termed *Polytrich'ia*; although this term is, at times, employed for the growth of hair on parts not usually having it.

POLYTROPH'IA, (*poly*, and *τροφη*, 'nourishment,') Abundant or excessive nutrition.

POLYURESIS, (*poly*, and *ουρις*, 'urination,') *Diabetes*.

POLYURIA, (*poly*, and *ουρον*, 'urine,') *Diabetes*.

POLYU'RICUS. Sauvages calls *Ischu'ria polyu'rica* a kind of ischuria, caused by accumulation of urine in the bladder.

POMA, gen. *Po'matis*, (πομα.) Drink—*p. Chinen-sia*, (pl. of *pomum*, 'an apple,') see *Citrus aurantium*.

POMACEUM, (*pomum*, 'an apple,') *Cider*.

POMA'TUM, *Pomma'tum*, (from *poma*, 'apples,') which formerly entered into its composition. (F.) *Pommade*, *Liparole*, *Stéarolé*. A pharmaceutical preparation, employed only externally, which differs but little from an ointment, except in hav-

ing generally less consistence. It is usually scented and coloured to render it more agreeable.

Common *Pomatum* is made of *lard*, 12lbs; *suet*, 4lbs; *essence of lemon*, $\frac{3}{4}$ viij.

POMATUM AMMONIACALE GONDRETI, *Pommade de Gondret*—*p. ad Labia Demulcenda*, *Cerate for the lips*—*p. Epispasticum flavum*, *Unguentum lyttæ medicatum aliud*—*p. Epispasticum viride*, &c., *Unguentum lyttæ medicatum*—*p. for the Lips*, *Cerate for the lips*—*p. Nitricum seu oxygenatum*, *Unguentum acidi nitrosi*—*p. Saturni*, *Unguentum plumbi superacetatis*.

POMBALEA IPEACACUANHA, (after the Marquis de Pombal, a celebrated Portuguese statesman and *savant*,) see *Ipeacacuanha*.

POMEGRANATE, *Punica granatum*.

POMMADE, ([F.] *pomme*, [L.] *pomum*, 'an apple,') *Pomatum*—*p. d'Allyon*, *Unguentum acidi nitrosi*—*p. Antipsorique de Helmerich*, *Unguentum sulphuratum alkalinum ad scabiem*—*p. Antipsorique de Jasser*, *Unguentum sulphuris compositum*—*p. d'Autenrieth*, *Unguentum antimonii tartarizati*—*p. de Cyrrillo*, *Unguentum muriatis hydrargyri oxygenato medicatum*—*p. Epispastique jaune*, *Unguentum lyttæ medicatum aliud*—*p. Épispastique verte*, *Unguentum lyttæ medicatum*—*p. de Garou*, *Unguentum epispasticum de daphne gnidio*.

POMMADE DE GONDRET, *P. ammoniacale de Gondret*, *Cuustique ammoniacale*, *Poma'tum seu Unguentum ammoniacal'e Gondreti*, *Gondret's Counterir'itant*, *Gondret's ammoniacal Cuustic*. (*Liq. ammon. p. ij; Sevi, p. j; Ol. amygdal. p. j; M.*) Used to excite a speedy revulsion in cases of chronic affections of the brain, incipient cataract, amaurosis, &c., as well as to cauterize the integuments deeply.

The above formula is that of the French Codex, but, according to M. Trousseau, it is much too hard for use, unless the weather is warm. He proposes two formulæ, one for summer, the other for winter. In the former, he directs three parts of *lard*, one of *suet*, and four of *liquor ammoniz*; in the latter, equal parts of *lard* and *liquor ammoniz*.

POMMADE DE HELMERICH, *Unguentum sulphuratum alkalinum ad scabiem*—*p. Pour les Lèvres*, *Cerate for the lips*—*p. Ophthalmique* (Regent's), *Unguentum ophthalmicum*—*p. Oxygénée*, *Unguentum acidi nitrosi*—*p. Stibiée*, *Unguentum antimonii tartarizati*.

POMME, ([L.] *pomum*, 'an apple,') see *Pyrus malus*—*p. d'Adam*, *Pomum Adami*—*p. d'Amour*, *Solanum lycopersicum*—*p. Blanche*, *Camassia esculenta*, *Psoralea esculenta*—*p. de Prairie*, *Camassia esculenta*, *Psoralea esculenta*—*p. Épineuse*, *Datura stramonium*—*p. de Merveille*, *Momordica balsamica*—*p. Mousseuse*, *Bedegar*—*p. de Terre*, *Solanum tuberosum*.

POMMEL FOOT, 'knobfoot,' *Kyllosis*.

POMMERASSE, *Aristolochia clematitis*.

POMMES HÉMORRHOÏDALES, see *Viscum album*.

POMMETTE, ([F.] *pomme*, 'an apple,') see *Malæ os*.

POMMIER, *Pyrus malus*.

POMPE POUR L'ESTOMAC, *Stomach pump*.

POMPHOLYGE'RON, (πομφολυγηρον, from *πομφολυγω*, 'I raise blisters,') The name of a plaster employed by the ancients.

POMPH'OLYX, gen. *Pompholygis*, (πομφολγῆ, from *πομφος*, 'a blister,') A small vesicle on the skin. An eruption of *bullæ* or *blebs*, appearing without any inflammation around them, and without fever. It is the *Erythymis Pompholyx* of Good. See *Pemphigus*.

POMPHOLYX, Tutia, Zinci oxydum.

POMPHUS, (Πομφος,) Vesicula, Wheal.

POMPION, (Πομπιον,) Vesicula pepo.

POMUM, see *Pyrus malus*.

POMUM ADA'MI, *Eminentia thyroidea*, (F.)

Pomme d'Adam, *Ad'am's App'le*. The projection formed by the thyroid cartilage in the neck—of males particularly. It is so called from a notion that it was caused by the apple sticking in the throat of our first parent.

POMUM AMORIS, 'apple of love,' Testicle—p. Hierosolymitanum, Momordica balsamina—p. Melongena, see *Solanum melongena*—p. Mirabile, Momordica Balsamina—p. Spinosum, *Datura stramonium*.

PONCTION, Puncturing,

POND SHOVEL, Unisema deltifolia.

PONDERA ET MENSURÆ, Weights and measures.

PONDUS PHARMACEUTICUM, see *Weights and measures*.

PONGITIF, Pungent.

PONS, gen. *Pontis*, (F.) *Pont*, 'a bridge.' A name given in anatomy to a part which connects others,—by a bridge, as it were.

PONS CEREBELLI, *Pons Varolii*—p. Hepatis, see *Lobulus anonymus*—p. Sylvii, *Quadrigena Tubercula*—p. Tarini, *Tarini pons*.

PONS VAROLII, P. Cerebelli, *Corpus seu Tubercula annularia*, *Processus annularis*, *Prominentia annularis cerebri*, *Eminentia annularis*, *Nodus encephali seu cerebri*, *Protuberantia basilaris*, *Protuberantia annularis Willisii*, *Varolii's bridge*, *Cerebral protuberance*, (F.) *Protuberance cérébrale* on *annulaire*, *Méencéphale*, *Méocéphale*, *Pont de Varole*, *Annular protuberance*. An eminence at the upper part of the medulla oblongata, first described by Varoli. It is formed by the union of the crura cerebri and crura cerebelli.

PONS ZYGOMATICUS, Zygomatic arch.

PONT, *Pons*—p. de *Tarin*, *Tarini pons*—p. de *Varole*, *Pons Varolii*.

PONTE, (from *pondre*, 'to lay an egg,') see *Parturition*—p. *Périodique*, see *Parturition*.

PONTERIA CORDATA, (from *Guilio Pontederia*, Professor of Botany at Padua,) *Unisema deltifolia*.

PONTICUM ABSINTHIUM, *Artemisia Pontica*.

POOK, Abdomen.

POPLAR, AMERICAN, ([F.] *Peuplier*, [L.] *Populus*,) *Liriodendron*—p. *Balm of Gilead*, *Populus Candicans*—p. *Black*, *Populus*—p. *Tree*, *Liriodendron*—p. *Tulipbearing*, *Liriodendron*.

POPLES, gen. *Poplitis*, *Gare'tum*, *Garretum*, *Ig'nye*, *Ig'nyé*, *Suffra'go*, *Fos'sa ge'nu*, *Fos'sa poplite'a*, the *Ham*, *Hough*, *Hoff*, *Hock*, (F.) *Jarret*, (from [L.] *post*, 'behind,' and *plico*, 'I fold,') The posterior part of the knee joint. The ham forms what is called the *popliteal region* or *space*.

POPLESY, Apoplexy.

POPLITEUS, (from *popes*,) *Popliteal muscle*.

POPLITÉ, *Popliteus muscle*.

POPLITEAD, see *Popliteal aspect*.

POPLITEAL, *Popliteus*, *Poplitic*, (F.) *Poplité*, (from *popes*.) That which relates to the ham. A name given to various parts.

POPLITEAL ARTERY—*Popliteal portion* of the *crural*,—(Ch.), is really a continuation of the femoral, which changes its name on passing through the abductor magnus muscle. It descends a little obliquely outwards into the hollow of the ham, and extends from the commencement of the lower third of the thigh to the end of the upper quarter of the leg, where it terminates by dividing into the *posterior tibial* and *peroneal* arteries. In the ham, it gives off the three *superior articular arteries*; and, at the top of the leg, the arteries of the *gemelli* or *geminii* muscles, *geminii branches*, (F.) *branches jumelles*, the two

inferior articular arteries, and the *tibialis antica*.

POPLITEAL ASPECT. An aspect towards the side on which the *popes* or ham is situated. (Barclay.) *Poplitic'al* is used by the same writer adverbially, and signifies 'towards the popliteal aspect.'

POPLITEAL FOSSA, *Poples*.

POPLITEAL GLANDS, *Glandulæ poplitæ*. Lymphatic glands, four in number, seated in the popliteal region or space.

POPLITEAL MUSCLE, *Poplitæus Mus'cle*, *Subpoplitæus*, (F.) *Muscle poplité*, *Muscle Jarretier*, *Fémoro-popliti-tibial*. (Ch.), is seated in the popliteal region. It is triangular, long, flat; and is inserted, on the one hand, by means of a strong tendon, into the tuberosity at the outer condyle of the femur, terminating below at the upper part of the posterior surface of the tibia. This muscle bends the leg on the thigh, and, reciprocally, the thigh on the leg. When the leg is bent, it causes it to turn, in rotation, inwards.

POPLITEAL NERVES. They result from the bifurcation of the sciatic, and are two in number, viz. 1. The *external popliteal*, *Peroneal nerve*, *Branche péronière*, (Ch.) It is smaller than the internal; turns to the outside of the fibula at the upper part of the leg, and divides into the *musculocutaneous branch*, *nerf prétiéo-digital*, (Ch.) and the *anterior tibial*, *nerf prétiéo-susphalangéaire*, (Ch.) 2. The *internal popliteal nerve*, *Branche tibiale* (Ch.), which is larger than the other, and seems to be the real continuation of the sciatic nerve. It descends vertically into the hollow of the ham; and passes beneath the aponeurotic arch of the solaris muscle, beyond which it takes the name *tibial nerve*. It furnishes the *external saphæna nerve*, and a number of branches, that are distributed to the posterior muscles of the leg. It then passes to the sole of the foot to be divided into two branches, called *Plantar nerves*.

POPLITEAL REGION, *Poples*—p. *Space*, *Poples*.

POPLITEAL VEIN has the same arrangement as the artery behind which it is situate.

POPLITIC, *Popliteal*.

POPONAX, see *Pastinaca opoponax*.

POPPY, ([Sax.] *popig*,) see *Papaver*—p. *Celandine*, *Stylophorum diphyllum*—p. *Corn*, red, *Papaver rhœas*.

POPPY HEADS, see *Papaver*.

POPPY, HORN, *Stylophorum diphyllum*.

POPPY JUICE, *Meconium*—p. *Pill*, *Opium*.

POPPY, PRICKLY, *Argemone Mexicana*—p. *Spatling*, *Cucubalus Behen*—p. *Thorn*, *Argemone Mexicana*—p. *White*, *Papaver*—p. *White*, capsule, see *Papaver*.

POPULUS, P. *ni'gra*, *Ægei'ros*, *Achero'is*, *Black pop'lar*, (F.) *Peuplier*, P. *noir* ou *franc*. The young buds, *oculi* or rudiments of the leaves, which appear in the beginning of spring, were formerly used in an official ointment—the *unquen'tum popule'um*. It was, also, recommended in chronic affections of the lungs.

The bark of *Populus Tremulo'id'es*, *American aspen*; of P. *Trem'ula seu pen'dula*, *Europe'an aspen*; and of *POPULUS CAN'DICANS*, *Balm of Gilead* *Pop'lar*, is antiperiodic, and furnishes *salicin* and *pop'ulin*, in which the properties probably reside.

POPULUS BALSAMIFERA, see *Fagara octandra*—p. *Candicans*, see *Populus*—p. *Pendula*, P. *tremula*—p. *Tacamahaca*, see *Fagara octandra*—p. *Tremula*, see *Populus*—p. *Tremuloides*, see *Populus*.

PORCELAIN, *Essera*, *Urticaria*.

PORCELET, *Hyoscyamus*.

PORCELIA TRILOBA, (after A. Porcel, of Spain, a promoter of botany,) see *Carica papaya*.

PORCELLIO, (dim. of *porcus*, 'a hog,') *Oniscus*.

PORCELLUS, *Oniscus*.

PORCULUS, Oniscus.

PORCUPINE DISEASE, Ichthyosis.

PORC'UPINE MEN, *Echinoder'mi*. Persons have been so called in whom the cuticle is produced in the form of pointed prolongations, as if it had been moulded on the papillæ, like the shorter and blunter quills of the porcupine.

PORCUS, (porcus,) Vulva.

PORDE, (πορδη,) Crepitation, Fart.

PORE, *Por'us*, (poros, 'a passage,') *In'terstice*.

Pores are the excessively minute spaces or interstices which separate the integrant molecules of bodies, and render these bodies permeable. Anatomists formerly gave this name to supposititious openings at the extremities of vessels at the surface of different membranes. Thus EXHALANT PORES have been supposed to exist in the exhalants, to transmit the fluids exhaled.—ABSORBENT PORES to be employed in taking up parts that have to enter the circulation. Pores exist in the cuticle; yet Humboldt, with a powerful magnifying-glass, was unable to observe them. These pores have also been called *spira'cula*.

A body that is possessed of pores is said to be *po'rous*, *poro'sus*, (F.) *Porous*.

POREBLIND, (peerblind [?],) *Myopic*.

PORED MILK, Colostrum.

POREUX, Porous.

PO'RI (pl. of *porus*, 'a passage,') BILIA'RII, *Duct'us bilia'rii seu bilif'eri*, *Canalic'uli bilif'eri*, *Sur'culi fell'e'i*. Ducts which receive the bile from the penicilli of the liver, and convey it to the hepatic duct.

PORI CUTANEI, *P. Exhalan'tes seu Sudorif'eri seu Resorben'tes*, *Mea'tus cuticula'res*. The pores of the skin.

PORI EXHALANTES, *P. cutanei*—*p. Pulmonum*, *Cellule, bronchie*—*p. Resorbentes*, *P. cutanei*—*p. Sudoriferi*, *P. cutanei*.

PORILLON, *Narcissus pseudonarcissus*.

PORNIA, (πορνεία,) Prostitution.

PORNOGRAPHY, *Pornograph'ia*, (πορνεία, 'prostitution,' and *γραφω*, 'I describe,') A description of prostitutes or of prostitution, as a matter of public hygiene.

POROCE'LE, (πωρος, 'callus,' and *κληη*, 'a tumour,') A hard tumour of the testicles or its envelopes.

POROMA, gen. *Poro'matis*, (πωρωμα,) Induration.

POROMPHALOCÉLE, *Poromphalon*.

POROM'PHALON, *Poromph'alus*, *Porompha-loce'lē*, (πωρος, 'callus,' and *ομφαλος*, 'navel,') Umbilical hernia with hard and thickened parietes.

POROS, (πωρος,) Canal, Meatus; (πωρος,) Callosity.

POROSIS, (πωρωσις,) Chalaza, Induration.

POROSUS, (from *poros*, 'pore,') Porous.

POROT'IC, *Porot'icus*, (F.) *Porotique*, (from *πωρος*, 'callus,') A remedy believed to be capable of assisting the formation of callus.

PORPHYRA, (πορφυρα, 'purple,') *Purpura*—*p. Græcorum*, *Petchiæ*—*p. Hæmorrhagica*, *Purpura hæmorrhagica*—*p. Laciniata*, *Ulvæ umbilicalis*—*p. Umbilicalis*, *Ulvæ umbilicalis*.

PORPHYRISMA, (from *πορφυρα*, 'purple,') *Scarlatina*.

PORPHYRISMUS, *Scarlatina*.

PORPHYRIZATION, *Levigation*.

PORPHYROTYPHUS, see *Typhus*.

PORPHYRU'RIA, *Porphyu'ria*, (πορφυρα, 'purple,' and *ουρον*, 'urine,') A state of the urine—*por'puric u'rine*—in which it deposits the remarkable colouring matter—*Pur'purin*, *Uro'sacin* or *Uro'sacin*, *Rosac'ic acid*, *Uroërythrin*, *Uro'rhodin*, *U'r'rhodin*—in association with urate of ammonia. It is considered to be a modification of Urohematin, hæmaphæin.

POR'PHYRY, *Porphyri'tes*, (from *πορφυρα*,

'purple,') A species of very hard, compound rock, susceptible of considerable polish, and used in the fabrication of mortars and slabs for the apothecary.

PORPHYRY, BLACK, *Ophites*.

PORPHYURIA, *Porphyuria*.

PORRACE, *Porraceus*.

PORRA'CEOUS, *Porra'ceus*, *Pra'sinus*, (from *porrum*, 'a leek,') (F.) *Porracé*. Having the green colour of the leek. An epithet given to certain excreted matters,—the sputa, bile and fæces particularly.

PORREAU, *Allium porrum*.

PORRECTUM, (*porrigo*, *porrectum*, 'to stretch,') *Lever*.

PORRET, *Allium porrum*.

PORRIDGE, *ATHOL*, see *Mahogany*.

PORRIGINE, *Porrigio favosa*.

PORRIGINEUX, *Porrigineus*.

PORRIGINOSUS, *Porriginosus*.

PORRIG'INOUS, *Porrigino'sus*, (F.) *Porrigineux*. Appertaining to *porrigo*.

PORRIGO, gen. *Porrig'inis*, 'scurf or scall in the head,' *Tin'ea*, *T. Cap'itis*, *Fa'vus*, *Furfura'tio*, *Furfures*, *Ecpye'sis seu Phly'sis porrigio*, *Trichono'sis*, *Scal'led head* in general, *Scal'lard*; from *porrigere*, 'to spread,' or from *porrum*, 'a leek,' from the odour of the discharge. [?] (F.) *Teigne*, *Rache*. Some of the varieties are communicable. With the ancients a scurfy affection of the head; dandriff. Since often used like tinea for scalled head in general; and by Willan and Bateman for a class of diseases of the hairy scalp, principally characterized by an eruption of the pustules, called favi and achores, unaccompanied by fever.

PORRIGO CONFERTA, *P. scutulata*.

PORRIGO DECAL'VANS, *Tricho'sis a'rea*, *A'rea*, *Alope'cia are'a'ta seu partial'is seu circumsc'ripta*, *Phytoalope'cia*, *Trichono'sis furfura'cea*, *Tyr'ia*, *Tyro'ma*, *Ophi'asis*, *Tin'ea ton'dens*, *Her'pes seu Tin'ea tonsu'rans*, *Vitili'go*, *Squa'ra ton'dens*, *Lim'ited or partial Bald'ness* (F.) *Teigne tondante ou pelade*, *Pelade achromateuse*, *P. décalvante*, *Dartre tonsurante*, consists of patches of baldness without decay or change of colour in the surrounding hair, the exposed parts of the scalp being glabrous, white, and shining. The morbid condition sometimes spreads, so that the patches coalesce, rendering the baldness extensive. To a vegetable growth, forming a tube around the roots of the hair in *Tinea tonsdens*, Gruby gave the name *Micro'sporon seu Micro'sporum Audou'ni*. It is, also, called *Trichoph'yton seu Trichomyces decal'vans*. Another vegetable parasite, called *Trichoph'yton seu Trychomyces tonsu'rans*, (F.) *Teigne tonsurante*, is developed in the root of the hair. The treatment consists in keeping the scalp cleared by constant shaving, and in the steady application of some stimulating liniment or ointment.

PORRIGO FAVO'SA, *Ce'rian*, *Fa'vus*, *Ecpye'sis porri'go favo'sa*, *Tin'ea favo'sa seu ve'ra*, *Scab'ies cap'itis favo'sa*, *Melice'ra*, *Melice'rum*, *Melice'ria*, *Mellifa'vium*, *Porrigoph'ytia*, *Honeycomb Scall* or *Tetter*, (F.) *Teigne favéuse*, *Porrigine*. This consists in an eruption of pustules common to the head, trunk, and extremities, peaisized, flattened at the top, in clusters, often uniting; discharge, fetid; scabs, honeycombed; the cells filled with fluid. It occurs both in early and adult life. Modern researches have satisfied most observers, that the crusts of favus are of vegetable nature, and formed of a parasitic plant, *Fungus porrig'inis*, belonging, according to Gruby, to the obsolete genus *mycoderm'a*. MM. J. Müller and Lebert refer it to the genus *Oidium*, and call it *Oid'ium Schönléin'i*, whilst Remak assigns it to a new genus under the name *Acho'zion Schönléin'ii*. The point is not, however, settled. The treatment of this variety must consist in the exhibition of laxatives and the mildest stimulating ointments,

such as the *Ung. zinci*, *Ung. hydrargyri præcipitat. alb.*, &c. Eczema of the hairy scalp is often confounded with Porrigo.

PORRIGO FURFURANS, *Ecypse'sis porrigo furfuracea*, *Tin'ea seu Tricho'sis furfuracea* seu *porrigino'sa*, *Pityr'iasis*, *Teigne furfuracée* of Alibert, commences with an eruption of small aches, the fluid of which soon concretes and separates in innumerable thin, laminated scabs, or scalelike exfoliations. It is attended with a good deal of itching and some soreness of the scalp, to which the disease is confined. It occurs chiefly in adults. In the treatment, the scalp must be kept shaved. The part must be washed repeatedly with soap and water, and an oiled silk cap worn. The nature of the ointments used must vary according to the period of disease and the irritability of the part affected. *Zinc ointment*; — one made with the *Coccus indicus*; — the *Sulphur ointment*; — *Citrine ointment*; *Ung. hydrargyri nitricooxydi*; *Tar ointment*, and *Nitric acid ointment* have all been found useful. Should inflammation or discharge arise, these applications must of course be intermitted.

PORRIGO LARVALIS, *P. lupinosa*.

PORRIGO LARVA'LIS, *Crusta lac'tea*, *Ecypse'sis Porri'go crusta'cea*, *Menti'go*, *A'chor*, *Tin'ea lac'tea* seu *Fa'ciei*, *Lactu'men*, *Lacti'go*, *Milky scall* or *Tetter*, *Milkblotch*, (F.) *Croûte de lait ou lait-euse*, *Feux volages*. This is, almost exclusively, a disease of infancy. It commonly appears first on the forehead and cheeks, in an eruption of numerous, minute, whitish aches on a red surface. These break and discharge a viscid fluid, which becomes incrustated in thin, yellowish, or greenish scabs. The patches spread, until the whole face becomes, as it were, enveloped in a mask; hence the epithet *larvalis*. The disease is rather a variety of Eczema. Sometimes the eruption appears on the neck and breast; and, occasionally, on the extremities. In general, the health of the child does not suffer, but the eruption is always attended with much itching and irritation. The treatment consists in mild, tepid ablution, and the application of gently stimulating ointments, as the unguentum zinci. The bowels must be kept in a soluble state by small doses of calomel, or by rhubarb and magnesia.

PORRIGO LUPINO'SA, *Tin'ea lupi'na*, *Fa'veus dispersus*, *Scab'ies cap'itis*, *Ecypse'sis porrigo lupino'sa*, *Porri'go larva'lis*, *Impet'i'go figura'ta*, *Tin'ea granulata* seu *muc'o'sa*, *Crusta lac'tea* (of some), *Ment'agra infantum*, (F.) *Teigne, Dartre crustacée flavescente*, *Favus disséminé*, consists of minute pustules in small patches, mostly commencing on the scalp; the patches terminate in dry, delving scabs, resembling lupine seeds; the interstices often covered with a thin, whitish and exfoliating incrustation. It occurs chiefly in early life, and requires the same treatment as the last.

PORRIGO SCUTULA'TA, *P. confer'ta*, *Tin'ea granulata* seu *feco'sa*, *Ecypse'sis porri'go galea'ta*, *Acho'res seu Scab'ies cap'itis*, *Fa'vus confer'tus*, *Ring'worm of the Scalp*, *Scall'd head*, (F.) *Teigne annulaire*, *T. en anneau*, *T. nummulaire*, *Favus en groupes*. It commences with clusters of small, light, yellow pustules, which soon scab off; and, if neglected, become hard by accumulation. If the scabs be removed, the surface is left red and shining, but studded with slightly elevated points. If not attended to, it involves the whole head. It occurs in children three or four years of age, and is very unmanageable. The rules laid down under *Porri'go furfurans* must be here still more perseveringly enforced. The whole tribe of stimulating ointments may be used in succession. No one will always succeed, and hence the necessity of change, provided advantage should not seem to follow any particular application. The

Unguentum ad Scabiem of Banyer sometimes cures, when others fail.

PORRIGOPHY'TA, (*porrigo*, and φυτόν, 'a vegetable,') *Porri'go favo'sa*. *Por'rigophyte* means the plant of favus or of porrigo favosa.

PORRUM, *Allium porrum*, *Thymion*.

PORRUM, *Thymion*.

POR'TA, *Py'læ*, (F.) *Porte*, 'a gate.' The part of the liver or other organ—*hilus*—where its vessels enter as by a gate. See *Liver*. Also, the *Vulva*.

PORTA HEPATIS, see *Liver*—p. *Jecoris*, see *Liver*—p. *Lienis*, see *Spleen*.

PORTA VE'NA, *Ve'na por'tæ*, *V. porta'rum* seu *ad por'tas seu porta'lis seu mag'na seu quæ ad por'tas est seu stelechi'a seu lac'tea seu ostia'ria seu arterio'sa seu arteria'lis*, *Rama'lis ve'næ*, *Jan'itrix*, *Ma'nus jec'oris seu he'patie*.

The *Syst'em of the ve'na por'ta* or the *Abdom'inal ve'nous syst'em* or *Por'tal syst'em*, (F.) *Veine porte*, *Porte veine*, is a vascular apparatus of black blood, situate in the abdomen, and resulting from two distinct orders of vessels, united into one common trunk. 1. One of these trunks, much more extensive than the other, has received the name *Vena porta abdominal'is*, because it arises from all the organs enclosed in the cavity of the abdomen, except the kidneys and bladder; and, in the female, the uterus. The *Abdom'inal ve'na por'ta* has two principal roots, the *splenic vein* and the *superior mesenteric*, which have their ramifications in all the digestive organs. 2. The other trunk, distributed solely to the liver, has been called the *Hepatic ve'na por'ta*. It appears to be a continuation of the abdominal *vena porta*, and proceeds towards the inferior surface of the liver. Opposite the transverse fissure of that organ, it divides, like an artery, into two branches, which, at their union, seem to form a canal or duct, called by some the *Si'nus of the ve'na por'ta* or of the *portal vein*. These branches are distributed in the substance of the liver, whither they are accompanied by a prolongation of the fibrous capsule of that organ,—the capsule of Glisson. The different branches of the *vena porta* are devoid of valves. See *Liver*.

PORTAL, (from *porta*.) Relating to the porta of the liver,—as the *portal system*, or system of the *vena porta*. By extension, the term has been applied to an analogous system of vessels in the kidney. See *Kidney*.

PORTAL FISSURE, see *Liver*—p. *Fossa*, see *Liver*—p. *System*, see *Porta Vena*—p. *System* of the kidney, see *Kidney*—p. *Vein*, *Porta Vena*—p. *Veins* of the kidney, see *Kidney*.

PORTE-AIGUILLE, *Acutenac'ulum*, *Acuduct'or*, *Rhaphidago'gus*, *Needlecarrier*, *Needlebearer*, ([F.] *porter*, 'to carry,' and *aiguille*, 'a needle,') An instrument for accurately laying hold of a needle, and giving it greater length, when it is so fine and small that it cannot be held by the fingers. It is of steel or silver, about two inches long, and throughout the whole, almost, of its length, divided into two branches, so as to form a kind of forceps, capable of being closed by means of a sliding ring. A modification of this instrument was devised by Dr. Physick for taking up deep-seated arteries beyond the reach of a tenaculum or common needle. It consists of a forceps so constructed as to hold in its extremity a needle armed with a ligature. The handles of the forceps are fastened together, temporarily, by a spring or catch; and, when the needle is fairly deposited beneath the vessel, it is disengaged from the forceps and drawn out, leaving the ligature behind, which can be tied without difficulty.

PORTE-BOUGIE (F.) A silver canula similar to that of a trochar, except that it is longer. It is used for conducting bougies into the urethra.

PORTE-CAUSTIQUE, *Porte-pierre*—*p. Fenille*, *Subsopularis*.

PORTE-LAQ. An instrument for carrying the laq or fillet over the limb of the fetus, whilst in utero, when turning is necessary.

PORTE-LIGATURE. (*porter*, 'to carry,' and *ligature*.) An instrument for conveying a ligature to desiccated parts.

PORTE-MÈCHE (F.), (from *porter*, 'to carry,' and *mèche*, 'a tent.') An instrument for carrying tents or *mèches* deeply into fistulous openings, or sinuses, to the bottom of ulcers, &c. It is a wire of silver or steel, four or five inches long, having a notch at one extremity, and a button at the other.

PORTE-MOXA (F.). A small instrument for keeping the moxa applied to a part to be cauterized. It consists of a metallic ring—separated from the skin by three supports of ebony—to which a handle is adapted.

PORTE-NEUD (F.), (from *porter*, 'to carry,' and *neud*, 'a knot.') An instrument for carrying a ligature round the base or pedicle of a tumour.

PORTE-PIERRE (F.), (*porter*, 'to carry,' and *pierre*, 'a stone.') *Porte-caustique*, *Crusticaph'o-rum*, *Caustic bearer*. An instrument similar to a *porte-crayon* for carrying lunar caustic.

PORTE-SONDE (F.). An instrument resembling a *porte-crayon*, employed for holding the style or *sonde*, and inserting it more readily into the nasal duct, during the operation for fistula lacrymalis.

PORTE VEINE. *Porta vena*.

PORTEMUM. *Menster*.

PORTER, (from *beign drunk* by *porters*.) *Cerevisia*.

PORTIO DURA, 'hard portion,' Facial nerve—*p. Inter duram et mollem*. *Portio Wisbergii*—*p. Intermedia Wisbergii*, *Portio Wisbergii*—*p. Media inter portionem duram et portionem mollem*, *Portio Wisbergii*.

PORTIO MOLLIS, 'soft portion,' Auditory nerve.

PORTIO WISBERGII, *P. intermedia Wisbergii* seu *media inter portionem duram et portionem mollem* seu *inter duram et mollem*. (F.) *Fasciculus intermediarius de Wisberg*, *Intermédiaire de Wisberg*. A small, white fasciculus, intermediate between the *portio mollis* and *portio dura* of the seventh pair of nerves, which was first described by Wisberg. The Chorda Tympani is supposed by some to emanate from this nerve.

PORTION COSTO-TRACHÉLIENNE DU SACRO-SPINAL, *Longissimus dorsi*—*p. Écailleuse ou squameuse du temporal*, *Squamous portion of the temporal*.

PORTION GODRONNÉE ('plaited') *DE LA CORNE D'AMMON*. A name given by Vieq d'Azur to a small, denticulated cori. of a compact tissue and granular appearance, which unites the cornu ammonis to the paries of the lateral ventricle.

PORTION LOMBAIRE DU PLEXUS CRURAL, *Lumbar plexus*.

PORIONARIUM, (from *porta*, 'a gate,') *Pylo-rus*, *Duodenum*.

PORTEULACA, *P. oleracea*, *Andrach'nè*, *Al-lum Gall'icum*, *Portulaca*, *Portulaca*. (F.) *Pourpier cultivé*. This plant abounds with a watery and somewhat acid juice; and is often put into soups or pickled with spices. It is regarded as antiseptic and aperient.

PORUS, (*πoros*.) *Callosity*; (*πoros*.) *Canal*, *Meatus*—*p. Acusticus*, *Auditory canal*, *internal*—*p. Acusticus externus*, *Auditory canal*, *external*.

Porus Opticus. An opening in the centre of the cribriform lamella of the sclerotic, through which the central artery of the retina enters the eyeball.

POSCA. *Oxyerate*.

POSE. ([Sax.] *ζεπορε*.) *Coryza*.

POSIMOS. (*ποσιμος*.) *Potable*.

POSITION, FALSE, *False position*.

POSOL'OUY, *Posology*—*in*, *Dosislog'ia*. (*ποσος*, 'quantity,' and *λογος*, 'a discourse.') *Indication of the doses in which the different articles of the materia medica ought to be exhibited*. (F.)

Dosage.

POSQUERIA DUMETORUM, *Randia Dumetorum*.

POSSÉDÉ. *Possessed*.

POSSESS'ED, *Demoni'acus*, (F.) *Possédé*. One affected with epilepsy, or other strange neurotic affection, was formerly supposed to be 'possessed' by a demon. Hence, exorcism and various religious ceremonies, at one time employed to drive off the intruder.

POSS'ET, *Possitum*, (from *posui*, 'sour wine and water' [?].) *Milk curdled with wine, or any acid, or treacle. A preparation of beer and milk; Zythog'ala*.

POSSET SACK, see *Sack*—*p. Sago*, see *Sago*.

POSSETTING. *Regurgitation* (of Infants).

POSTBRACHIALE. (*post*, 'after,' and *brachium*, 'arm,') *Metacarpus*.

POSTCARPIUM, (*post*, 'after,' and *carpus*, 'wrist,') *Metacarpus*.

POSTERIOR ANNULA'RIS. An external, interosseous muscle, which arises from the corresponding sides of the metacarpal bones of the ringfinger and little finger, and is inserted into the inside of the tendon on the back of the ringfinger. Its use is to draw the ringfinger inwards.

POSTERIOR AURIS, *Retrahens auris*.

POSTERIOR INDICIS MA'NŪS. An internal, interosseous muscle, which arises from the inner or ulnar side of the metacarpal bone of the forefinger, and is inserted into the inside of the tendon on the back of the forefinger. Its use is to draw the forefinger inwards.

POSTERIOR INDICIS PE'DIS, *Abduc'tor Indicis Pe'dis*. An external, interosseous muscle, which arises from the contiguous sides of the metatarsal bones of the fore-toe, and second of the small toes, and is inserted into the outside of the root of the first bone of the foretoe. Its use is to pull the foretoe outwards.

POSTERIOR ME'DII DIG'ITI MA'NŪS. An external, interosseous muscle, which arises from the corresponding sides of the metacarpal bones of the middle and ringfingers: and is inserted into the inside of the tendon on the back of the middle finger. Its use is to draw the middle finger inwards.

POSTERIOR ME'DII DIG'ITI PE'DIS, *Abduc'tor me'dii dig'iti pe'dis*. An external, interosseous muscle, which originates from the contiguous sides of the metatarsal bones of the second and third of the small toes, and is inserted into the outside of the root of the first bone of the second of the small toes. Its use is to pull the toe outwards.

POSTERIOR TER'TII DIG'ITI, *Adduc'tor ter'tii dig'iti*. An external, interosseous muscle, which arises from the contiguous sides of the metatarsal bones of the third and fourth of the small toes, and is inserted into the outside of the root of the first bone of the third of the small toes. Its use is to pull the toe outwards.

POSTHE. (*ποσθη*.) *Penis*, *Prepuce*.

POSTHETOMIST, *Posthet'omus*, (*ποσθη*, 'prepuce,' and *τομη*, 'incision,') One who performs the operation of circumcision.

POSTHETOMY, *Circumcision*.

POSTHIA, (*ποσθια*.) *Hordesum*.

POSTHIOPLASTIC, *Posthioplast'icus*, *Posio-plas'tic*, *Posthioplast'icus*, (*posthium*, and *πλασσω*, 'I

form.') An epithet applied to the operation for restoring the prepuce.

POSTHITIS, *Acrobystitis*, *Acroposthitis*, *Epagogistis*, (from *ποσθν*, 'prepuce,' and *itis*.) Inflammation of the prepuce.

POSTHIUM, (*ποσθιον*.) Penis, Prepuce.

POSTHOCALYPTRON, (*ποσθον*, 'penis,' and *αλκντρον*, 'a covering,') Condom.

POSTHONCUS, (*posthe*, and *ονκος*, 'a tumour,') Swelling of the prepuce or penis.

POSTHOPLASTIC, Posthioplastic.

POSTHUMOUS, (*post*, 'after,' and *humus*, 'the ground,') Occurring after death.

POSTHUMOUS CHILD. One born after the death of its father, or taken from the dead body of its mother, as by the Cæsarean section.

POSTIME, (impostume,) Abscess.

POSTMORTEM, 'after death;' as 'a postmortem or neuroscopic examination.'

POSTNATUS. 'Born after.' In the old law writers, the second son.

POSTPARTUM. 'After delivery.' As 'postpartum hemorrhage.'

POSTPOSITIO, (*post*, 'after,' and *ponere*, *positum*, 'to place,') 'a postponement,') Delay in the return of a paroxysm. It is opposed to *anticipatio*, which means the return of a paroxysm earlier than its accustomed hour.

In this sense, we speak of an *anticipating*, and a *postponing*, intermittent (*Febris hystericalis*).

POSTURAL, *Posturalis*, (from *ponere*, *positum*, 'to place,') Relating or appertaining to posture or position—as 'postural treatment,' the treatment by position,—of a fractured limb for example.

POSTURAL RESPIRATION, see Ready method.

POTABLE, *Potabilis*, (*potare*, 'to drink,' and *habilis*, 'able,') *Potimos*, *Potos*, *Posimos*, drinkable. Water is said to be potable, when it contains atmospheric air; is not brackish; and has no mineral impregnation of consequence.

POTAGE, (from (L.) *potus*, 'drink,') see Broth.

POTAGER, Lachanon.

POTAMOPYRA, (*ποταμος*, 'a river,' and *πυρ*, 'fever,') Elodes (febris).

POTASH. So called from the pots or vessels in which it was made;—*Veg'etabile alkali*, *Gastrinum*. Also, Potass, (F.) *Potasse*.

POTASH OF COMMERCE, *Potassa impura*, *Impure potassa*, *Alumen catinum*, *Cineres Rusæci*, *Subcarbonas potassæ impurus*, *Potassæ carbonis impurus* seu *impura*, *Lixivus cinis*, the Pearlash of commerce, is prepared from the ashes of land-plants by calcination, solution in water, filtration, and evaporation. It usually consists of subcarbonate of potassa, sulphate of potassa, chloride of potassium, silex, oxide of iron, &c.

From the ashes of lees of wine, and from vine-twigs, a much purer alkali is obtained. It is called *Cinix infectorius* seu *fulcum*, *Cineres clavellati* seu *gravellati*, *Alumen fæcum*, (F.) *Cendres gravellées ou de sarment*.

Another variety of the alkali is afforded by burning argol in a crucible; afterwards powdering and calcining it, till it is nearly white; then dissolving in water, filtering, and evaporating. It is called *Salt of Tartar*, *Sal Tartari*, *Kali præparatum* e *tartaro*, *K. e tartaro*, *Subcarbonas potassæ purissimus*, *Potassæ subcarbonas e tartaro*. Lastly, when wormwood is burnt to ashes, dissolved in water, filtered, and evaporated to dryness, it affords another variety of vegetable alkali, the *Sal absinthii* or *Salt of wormwood*.

Impure potash is used for preparing the subcarbonate for medical use.

POTASS, Potassa, Potassa fusa, Potash. The protoxide of potassium.

POTASSA, Potass—p. Acetate of, Potassæ acetat—p. Azotate of, Potassæ nitrat—p. Bichromate

of, Potassæ bichromat—p. Bisulphate of, Potassæ supersulphat—p. Bitartrate of, Potassæ supertartras—p. Carbazotate of, Potassæ picras—p. Carbonate of, Potassæ carbonas—p. Carbonate of, pure, see Potassæ subcarbonas—p. Caustica cum calce. Potassa cum calce—p. Chlorate of, Potassæ murias hyperoxygenatus—p. Chromate of, Potassæ chromas—p. Chromate of, neutral, Potassæ chromas—p. Chromate of, yellow, Potassæ chromas.

POTASSA CUM CALCÈ. *P. d'opè cal'cis parata et ignè fusa*, *Potassa caustica cum calcè*, *Pot'assæ with lime*, *Calc cum kali puro*, *Causticum communè fortius*, *La'pis infernalis sive septicus*, *Kali causticum cum calcè*, *La'pis causticus*, (F.) *Potasse préparée avec la chaux*, *P. à la chaux*, *Pierre à caudère*. (*Potassæ*, *Calcis*, *aa 3j*. Rub together, and keep in a wellstopped bottle.—Ph. U. S.) Used for the same purposes as the next, but more manageable; being less deliquescent.

POTASSA, EUCHLORATE OF, Potassæ murias hyperoxygenatus—p. Ferrocyanate of, Potassii ferrocyanuretum—p. Ferropotassate of, Potassæ cyanuretum.

POTASSA FUSA, *Potassa*, *P. Caustica*, *Kali causticum* seu *purum*, *Fu'sed pot'assæ*, *Alkali vegetabile fixum causticum*, *Causticum communè acerimum*, *C. alkalinum* seu *communè* seu *potentiale* seu *Sali'num*, *Hy'dras kalicum*, *H. potassæ puræ fulsæ*, *Potassæ hy'dras*, *La'pis causticus* seu *causticus chirurgorum* seu *infernalis alcalinus* seu *septicus*, *Ruptorium communè*, *Fixèd veg'etabile alkali*, *Hy'drate of Potassa*, *Caustic Pot'ash*, (F.) *Potasse caustique*. (Prepared by evaporating solution of potass to dryness in an iron vessel.) This preparation is generally run into cylindrical moulds, covered with paper, and kept in wellcorked bottles. It is powerfully escharotic, and is used for forming issues, curing strictures, &c.

POTASSA, FUSED, *Potassa fusa*—p. Hydriodate of, Potassæ hydriodas—p. Hydrobromate of, see Bromine—p. Hydrocyanate of, Potassii cyanuretum—p. Hypermanganate of, Potassæ hypermanganas—p. Hyperoxalas, Potassæ, oxalate of—p. Hypertartras, P. supertartras—p. Hypophosphite of, see Calcis hypophosphis—p. Impura, Potash of commerce—p. and Iron, tartrate of, Ferrum tartarizatum—p. with Lime, Potassa cum calce—p. Liquida, Liqueur potassæ—p. Mephite of, Potassæ subcarbonas—p. Monochromate of, Potassæ chromas—p. Muriate of, Potassæ murias—p. Nitrate of, Potassæ nitras—p. Ope calcis parata et igne fusa, Potassa cum calce—p. Oxymuriate of, Potassæ murias hyperoxygenatus—p. Permanganate of, Potassæ permanganas—p. Prussiate of, Potassii ferrocyanuretum—p. and Soda, tartrate of, Soda, tartrate of—p. Subcarbonate of, Potassæ subcarbonas—p. Sulphate of, Potassæ sulphas—p. Sulphate of, acid, Potassæ supersulphas—p. Sulphurata, Potassii sulphuretum—p. Sulphuret of, Potassii sulphuretum—p. Supercarbonate of, Potassæ carbonas—p. Supersulphate of, Potassæ supersulphas—p. Supertartrate of, Potassæ supertartras—p. Tartrate of, Potassæ tartras.

POTASSÆ ACETAS, *Oxytartarus*, *Acetas kali* seu *lixivisæ* seu *kalicæ*, *Alkali tartari aceto saturatum*, *A. vegetabile cum aceto*, *Magisterium tartari purgans*, *Terra foliata Tartari*, *Kali acetalum*, *Sol diureticus* seu *Sennerti*, *Arca'num tartari*, *Ac'etate of pot'assæ*, (F.) *Acétate de potasse*. (*Acid. acetic*. Oj, *Potassæ bicarbon*. q. s. Mix and add, by degrees, enough acetic acid to saturate the alkali; then filter and evaporate till a pellicle is formed, which is to be dried upon bibulous paper; evaporate again in a water-bath, and, removing the pellicles, dry.—Ph. U. S.)

It is cathartic and diuretic. Dose, ʒj to ʒij, or more.

POTASSÆ ET ALUMINÆ SULPHAS, *Alumen*—p.

Azetos, Potassæ nitras—p. Biantimonias, Antimonium diaphoreticum—p. Bicarbonas, Potassæ carbonas.

POTASSÆ BICHRÔ'MAS. *Ka'li chro'micum ru'drum seu ac'idum seu bichro'micum, Bichro'mate of potassæ.* (Procured by adding sulphuric, nitric, or acetic acid to a solution of the neutral chromate, and setting it aside to crystallize.) It is a violent irritant poison; and has been used as a caustic in cases of morbid growths, in the proportion of from half a drachm to a drachm of the salt to from an ounce to an ounce and a half of water.

POTASSÆ BISULPHAS, P. supersulphas—p. Bitartas, P. supertartas.

POTASSÆ CARBO'NAS, *Ka'li aëra'tum, Bicarbo'nas potassæ, Potassæ bicarbonas* [Ph. U. S., L. E., D. and B.], *Denticarbo'nas potassæ, Car'bonate, bicarbonate or supercar'bonate of potassa.* This is not caustic, like the subcarbonate of potass, and hence, has been preferred to relieve acidity. (*Potass. carbon. liv. Aquæ destillat. Ox.* Pass carbonic acid through the solution until it is saturated, and filter. Evaporate at a heat not exceeding 160°, and dry the crystals on bibulous paper. Carbonic acid is obtained from marble by the addition of dilute sulphuric acid.—Ph. U. S.) Dose, gr. x to ʒss. It is used in forming the soda powders, as well as for its antacid properties.

Distillers and brewers prepare this salt with great facility by suspending a solution of the purified carbonate of potassæ in the fermenting tun. The alkaline salt is thus surrounded by an atmosphere of carbonic acid; and by absorbing it, it crystallizes into bicarbonate. The salt in powder is called *sal aëra'tus*. Its composition is between that of a carbonate and of a bicarbonate.

POTASSÆ CARBO'NAS, P. subcarbonas—p. Carbonas e Lixivo Cinere, P. subcarbonas—p. Carbonas purus, Pure carbonate of potassa, see P. subcarbonas—p. Chloras, P. murias hyperoxygenatus—p. Chloruretum oxydatum, P. murias hyperoxygenatus.

POTASSÆ CHRO'MAS, *Ka'li chro'micum fla'tum seu neutra'le, Neu'tral yel'low Chro'mate or Mono-chro'mate of potassa.* (Prepared by igniting a mixture of four parts of native chromite of iron (chrome iron ore) and one part of nitre; dissolving out the chromate of potassa by water, and crystallizing.) It has been used as an emetic, expectorant and diaphoretic. Dose, as an emetic, from two to four grains; as an expectorant, from ʒth to ½ grain. In the proportion of one drachm to 32 ounces of water, it has been used as an antiseptic wash—called by Jacobson *Liquor conservatrix*.

POTASSÆ CITRAS, *Cit'rate of potassæ. (Acid. citric. ʒx, Potass. bicarb. ʒxiv, Aquæ q. s.* Dissolve the acid in the water; and after effervescence has ceased, strain and evaporate to dryness. The salt is deliquescent, and must be kept in wellstopped bottles. Ph. U. S.) It possesses the refrigerant properties of the neutral salts, and may be given in fever, when to do little or nothing is the most advisable course. Dose, gr. x to ʒss.

POTASSÆ ET FERRI TARTRAS, Ferri tartarizatum—p. Hydras, Potassa fusa.

POTASSÆ HYDR'O'DAS, *Potassæ Protohydr'o-das, Potassæ protox'idi hydr'o'das, Ka'li Hydr'idicium seu Hydr'idicum, Hydr'o'das Ka'licus, Hydr'idate of potass, (F.) Hydr'idate de potasse.* If a solution of potass be poured on iodine, an iodate and hydriodate are formed. These may be separated by means of alcohol, which only dissolves the latter of these salts. The hydriodate, or rather the Iodide of Potassium, *Ka'lium Ioda'tum, Potassii Iodure'tum seu I'didum, (F.) Iodure de potassium*, may be obtained by evaporation. When the Iodide of potassium is dissolved in water, it becomes the *Hydr'idate of Potassæ*.

(The following is the formula in the Ph. U. S. *Potass. ʒvj; Iodine, in pulv. ʒxvj; Carbon. liqvi in pulv. subtiliss. ʒij; Ag. ioduret. Oij.* Dissolve the potassa in the water; add the iodine gradually, stirring until the solution becomes colourless, and continue the additions until the liquid remains slightly coloured from excess of iodine. Evaporate to dryness, stirring in the charcoal towards the close. Rub to powder, and heat to dull redness in an iron crucible, keeping up the temperature for 15 minutes. When cool, dissolve out the saline matter with pure water, filter the solution, evaporate and set aside to crystallize. By evaporating and crystallizing as before, an additional quantity of crystals may be obtained.) Virtues,—those of iodine.

POTASSÆ MU'RIAS, *Potassii chlo'ridum, Sal Digestivum seu febrifugum Spl'vii, Spir'itus salis mari'ni coaguatus, Mu'r'ate of Potass, Chloride of potassium.* Bitter and refrigerant.

POTASSÆ MU'RIAS HYPEROXYGENA'TUS, *Potassæ Chlo'ras seu oxygenu'rias, Oxygenu'rias, Hyperoxygenu'rias, Chlo'rate or Euc'hlo'rate of potass, Protochlo'rate of potassium, Al'cali vegetabil'e sal'tis-dephlogisticat'um, Chlorure'tum potassæ oxydat'um, Habi'tum oxygenu'rum, Oxyg'eno-chlorure'tum potassii, Ka'li chlo'ricum, Chlo'ras Ka'licus depuratus, (F.) Chlorate de potasse.* (It may be prepared by passing chlorine gas into a solution of potassa or carbonate of potassa, evaporating and crystallizing. Chloride of potassium is less soluble and remains in the mother water. A more complex form for its preparation is given in Ph. B. 1844.) It has been regarded as anti-neuropathic, and as an alterative purifier of the blood and humours. It is of doubtful efficacy. Dose, ʒ to 6 grains or more.

POTASSÆ NITRAS, P. Azotas, *Nit'rate or Azotate of Potass, Nitrum, N. prismat'um, Sal nitrum, Nit're, Salp'etre, Sal petra, Sal'do mul'cerum, Sump'is'ea, Berenic'ium, Halini'trum, Proto-nit'rate of Potassium, Nit'rate of protox'ide of Potassium, Ka'li nitricum, Nitrus ka'licus seu lixiv'ia, Azotas potassicus, (F.) Nitrate on Azotate de potasse.* (Obtained from the putrefaction of animal matters, in contact with calcareous or alkaline earths, by elixivation; adding, if necessary, woodashes to supply the alkaline basis. It is formed, in an impure state, by nature, in warm climates.)

It is diuretic and refrigerant, and, in large doses, purgative; externally, cooling and detergent. Dose, gr. x to ʒi. In large doses it is an acrid poison.

POTASSÆ NITRAS FU'SUS SULPHA'TIS PATE'LLO MIXTUS, *Sal prunellæ, Crystal'lus mineral'is, Nitras potassæ cum sulphur'e fixo, Nitrum tabula'tum, La'pis Prunellæ, Amalgam' mineral'e, Sorethro'at salt, Mineral' crys'tal, (F.) Sal de Prunelle.* Nitrate of potass, mixed with a little sulphate of potass and run into moulds. Its properties are those of nitre.

POTASSÆ OXALAS, *Ox'alas Acid'ulum Potassæ seu Potassæ acid'ulum seu ka'licus, Superox'alas ka'licum seu Potassæ, Hyperox'alas Potassæ, Sal Acetos'ile, Oxa'tum, Ka'li oxal'icum acid'ulum, Ox'alas subpotassicus, Superox'alate of Protox'ide of Potassium, Ox'alate of Potass, Salt of Sor'rel, (F.) Sel d'oseille, Sel à détacher.* Prepared from the juice of wood sorrel. It is refrigerant, and is used to make lemonade, &c.

POTASSÆ OXALAS ACIDULUM, Potass. oxalate of—p. Oxyurias, P. murias hyperoxygenatus.

POTASSÆ PERMAN'GANAS, *Kali Hypermangan'icum, Permann'ganat'um potassæ, Permann'ganate of potassæ, Hypermanganate of potassa.* (May be made by mixing equal parts of deutoxide of manganese and chlorate of potassa with rather more than an equal part of caustic potassa, dissolving in a little water, evaporating to dryness and ex-

posing to a temperature just short of redness. By hot water a solution is obtained, which on evaporation crystallizes. A form for its preparation is contained in Ph. B. 1864.) This salt has been given in cases of diabetes. Dose, from gr. ij to gr. v, three times a day. It has been prescribed, also, in obstinate dyspepsia; and has been highly recommended as a caustic, less painful and of more intense action than others; for this purpose, the powder is made into a paste with water. As an antibromic and escharotic combined, in the quantity of 20 grains to the pint of water, it, as well as manganic acid, has been used as a wash to cancerous and other ulcers.

Liquor Potassæ Permanganatæ, Solution of Permanganate of Potash, four grains of the permanganate to one fluidounce of water, is official in the British Pharmacopœia of 1864. See Condy's Disinfecting Fluid, which is double the strength of this.

POTASSÆ PICRAS, *P. Carbazo'tas, Pi'crate* or *Carbazotate of Potassa*: (obtained by neutralizing picric acid with potassa.) It has been used as an antiperiodic. Dose two to five grains.

POTASSÆ PROTOHYDRIDAS, *P. hydriodas*—*p. Puræ fusæ hydras, Potassa fusa*—*p. et Sodæ tartars, Soda, tartrate* of.

POTASSÆ SUBCARBO'NAS, *P. Carbo'nas* (Ph. U. S.), *P. carbo'nas e Lixiv'eo Cin'ere, Subcarbo'nas Kid'li, Carbo'nas Potassæ, Kali præparatum, Sal Herbar'um seu Plantarum seu Tartari seu Absin'thii, Calcareous Tartar, Subcarbonate or Meph'ite of Potass, &c.* (See Potash.) (A filtered solution of the impure subcarbonate, *Potassæ carbo'nas impu'ra* (Ph. U. S.), *Pearlash* (see Potash), evaporated to dryness.) This deliquescent salt is given in the same cases as the potassæ carbonas, than which it is more caustic and disagreeable.

The Parisian Codex has a *Subcarbo'nas Potassæ ex tar'taro combus'to*, vulgo *Sal Tar'tari* or *Salt of Tar'tar*:—(it is the *Potassæ carbo'nas pu'rus* seu *pura, Pure Carbonate of Potassa, Ph. U. S.*); a *Subcarbo'nas Potassæ ex deflagrantibus simul tartra'te acid'ulo et Nitra'te Potassæ paratus*, vulgo *Nitrum ex tar'taro fixum*; and a *Subcarbo'nas potassæ ex deflagrantibus cum Carbo'nibus Nitra'te Potassæ paratus*, vulgo *Nitrum fixum*. They are all used like the preceding.

POTASSÆ SUBCARBO'NAS IMPURUS, Potash of commerce.

POTASSÆ SUL'PHAS, *Kali vitriola'tum seu sulphur'icum, Sul'phas Kali, Lixiv'ia vitriola'ta, Nitrum vitriola'tum Schroederi, Sal Sapient'is seu Tartari sulphuratum seu Polychres'tus, Specif'icum Paracel'si, Spiritus vitrioli coagula'tus Mynsich'ti, Al'kali vegeta'bile vitriola'tum, Sal de duobus, Arcanum duplicatum, Tartarum vitriola'tum, Panacea Du'cis Holsat'is seu duplicata, Sul'phate of Potass, (F.) Sulfate de potasse.* (The salt, that remains after the distillation of nitric acid, dissolved; the excess of acid saturated with subcarbonate of potass and crystallized.) Purgative and diuretic. Dose, $\mathfrak{ss}$ to $\mathfrak{zj}$ as a laxative. In a large dose it has proved an irritant poison.

POTASSÆ SUL'PHAS cum SUL'PHURE, *Sul Polychres'tus Glaseri, Lixiv'ia Vitriola'ta Sulphurea.* (Formed by deflagrating nitrate of potass with sublimed sulphur.) Same virtues as the last. Dose, gr. xv to xxx.

POTASSÆ SULPHURETUM, Potassii sulphuretum—*p. Superoxalas, Potassæ oxalas.*

POTASSÆ SUPERSUL'PHAS, *P. Bisul'phas, Sal enia'm seu febrif'igus of Lem'ery, Supersulphate of Potass, Ac'id Sul'phate or Bisul'phate of Potass, Superprotosulphate of Potassium, (F.) Bisulfate de Potasse.* (The salt remaining after the distillation of nitric acid, dissolved, strained, and crystallized.) It is refrigerant and purgative. Dose, $\mathfrak{ss}$ to $\mathfrak{zj}$.

POTASSÆ SUPERTARTRAS, *P. hypertartaras, P. Bitartaras* (Ph. U. S.) seu *tartaras ac'idæ, Crystalli Tartari, Tartaras Acid'ulus Potassæ purus, T. superpotassæicus, T. subpotassæicus, Supertartarate of Prolox'idæ of Potassium, Bitartarate or Supertartarate of Potass, Crystals of Tartar, Crémor Tartari, Cream of Tartar* (when in powder), *Supertartarate of Potass, (F.) Tartrate acide de potasse, Crème de Tartre.* (The tartar of wine purified.) It requires 60 parts of water at 60° to dissolve it. It is mildly purgative, refrigerant and diuretic. Dissolved in water, with a small quantity of white wine, some sugar and lemon-peel, it forms an agreeable beverage in febrile diseases, under the name of *Imperial*. Dose, as a diuretic, $\mathfrak{zj}$ to $\mathfrak{zj}$; as a purgative, $\mathfrak{ss}$ to $\mathfrak{zj}$.

POTASSÆ SUPERTARTRAS IMPURUS, *Tartarus crudus, Tartarum, Lapis vini, Magneti'us, Pedagra, Tartar, (F.) Tartre cru ou brut.* It is called, also, *Tartarum rubrum* and *Red Ar'gol*, when obtained from red wines; *Tartarum album* or *White Ar'gol*, when from white wines. Tartar is the saline deposit from wines as they become aged; which attaches itself to the sides of the casks and bottles in which they are kept. It consists of a considerable quantity of acidulous tartrate of potass, tartrate of lime, silica, alumen, oxyd of iron, manganese, and a red colouring matter, if the wine be red. It is chiefly used for the preparation of cream of tartar.

POTASSÆ TARTRAS, *Kali tartariza'tum, Al'kali fixum tartarisa'tum, Tartarum solu'bile, Tartarus tartariza'tus, Lixiv'ia Tartarisa'ta, Sal Panche's'tum, Tar'tras Kali'us seu lixiv'is, Sal vegeta'bilis, Al'kali vegeta'bile tartariza'tum, Tar'trate of Potass, Soluble Tartar, (F.) Tartrate de Potasse, (Potassæ carbon. $\mathfrak{z}$ xvj; Potassæ Bitartar. in pulv. subtiliss. $\mathfrak{liij}$, vel q. s., *Aque bullientis*, cong.) Dissolve the carbonate of potassa in the water, add the bitartrate to saturation, and boil; filter, evaporate till a pellicle forms, and set aside to crystallize. Pour off the liquid; dry the crystals on bibulous paper, and keep in closelystopped bottles.—Ph. U. S.) A saline cathartic in the dose of $\mathfrak{zj}$ to $\mathfrak{zj}$.*

POTASSÆ TARTRAS ACIDA, *P. supertartaras*—*p. Tartras acidulus purus, P. supertartaras*—*p. Tartras stibialis seu stibiosus, Antimonium tartariza'tum.*

POTASSE, Potash—*p. Acetate de, Potassæ acetas*—*p. Bisulphate de, Potassæ supersulphas*—*p. Caustique, Potassa fusa*—*p. à la Chaux, Potassa cum calce*—*p. Chlorate de, Potassæ murias hyperoxygenatus*—*p. Hydriodate de, Potassæ hydriodas*—*p. Liquide, Liquor potassæ*—*p. Nitrate de, Potassæ nitras*—*p. Préparée avec la chaux, Potassa cum calce*—*p. et de Soude, tartrate de, Soda, tartrate* of—*p. Sulfate de, Potassæ sulphas*—*p. Sulfure de, Potassii sulphuretum*—*p. Tartrate de, Potassæ tartras*—*p. Tartrate acide de, Potassæ supertartaras.*

POTASSII BROMIDUM, *Potassæ hydrobro'mas, Bro'mide of potassium, (F.) Bromure de Potassium.* (Bromin. $\mathfrak{zj}$; Ferri fili, $\mathfrak{zj}$; Potassæ carbon, $\mathfrak{zj}$ and $\mathfrak{zj}$, vel. q. s.; *Ag. destillat., Oiv.* Add the iron filings and bromine to a pint and a half of the water; stirring for half an hour. Apply a gentle heat, and when the liquor assumes a greenish colour, add gradually the carbonate of potassa, dissolved in Oiss of the water, until it ceases to cause a precipitate. Continue the heat for half an hour, and filter. Wash the precipitate with the remaining water boiling hot, and filter. Mix the liquors and crystallize by evaporation.—Ph. U. S.) Virtues:—those of iodide of iron. Dose, gr. v to gr. x. It has been used, also, in the treatment of insomnia and restlessness, accompanied by, and dependent upon, nervous excitement. Large doses are said to occasion drowsiness, and to be antaphrodisiac. See *Bromine*.

POTASSII CHLORIDUM, Potassæ murias—p. Cyanidum, P. Cyanuretum.

POTASSII CYANURETUM, P. Cyanidum (Ph. U. S., 1863), *Cyan'uret of Potassium, Hydrocyanate of Potassa.* (Potassii ferrocyanur., in pulv., $\mathfrak{Z}$ viii; Potass. carbonat. exsicc., $\mathfrak{Z}$ ij. Mix intimately, and throw into a deep iron crucible, heated to redness; keep up the heat till effervescence ceases, and the fused mass concretes, of a pure white colour, upon a warm glass rod dipped into it; then pour out the liquid carefully into a shallow dish to solidify, stopping before the salt becomes contaminated with the precipitated iron. Break up the mass whilst warm, and preserve in well-stopped bottles. — Ph. U. S.) Used in the same cases as the hydrocyanic acid. Dose, a quarter of a grain.

POTASSII DEUTOCARBONAS, Potassæ carbonas.

POTASSII FERROCYANURETUM, P. Ferrocyaniidum, Ferrocy'anide or Ferrocy'anuret of Potassium, Ferrocy'anate or Ferrop'rus'siate or Prus'siate of Potassa, (F.) *Protocyanure jaune de fer et de potassium, Cyanure de fer et de potassium.* The yellow double cyanuret of potassium and iron, the salt from which the cyanuret of potassium is obtained by calcination. It is not much used in medicine. It has been advised as an astringent in the colliquative sweats of phthisis, in leucorrhœa, &c.; but it does not seem to be possessed of much activity. Dose, 10 or 15 grains every four or five hours.

POTASSII HYDRARGYROIDIDUM, Potassæ Iodo-hydrargyras, *Hydrargyrum biiodatum cum Kali'o iodo'to, Kalium ioda'tum hydrargy'ratum, Iodo-hydrargyrate or Hydrargyroi'dide of Potassium.* (This salt is made by the combination of iodide of potassium with red iodide of mercury.) Its properties are similar to those of the iodides of mercury, but in a less degree.

POTASSII IODIDUM, Potassæ hydriodas — p. Ioduretum, Potassæ hydriodas — p. Oxygenochloruretum, Potassæ murias hyperoxygenatus — p. Protoxidi hydriodas, Potassæ hydriodas.

POTASSII SULPHOCYANURETUM, *Sulphocyan'uret of Potassium.* This salt has been prepared as a substitute for hydrocyanic acid, and cyanuret of potassium, on the ground of its possessing their advantages without their inconveniences.

POTASSII SULPHURETUM, Potassæ Sulphure'tum seu tersulphure'tum, Potassa sulphurata, Kali sulphure'tum, He'par Sul'phuris seu Kali'mum, H. sul'phuris salinum seu sulphuris vulga're, Sulphure'tum lixi'cie seu Kali'mum, Li'xer of Sulphur, Sulphuret of Potass, Sulph'urated potass or potash, (F.) *Sulfure de potasse, Sulfure de potassium enlaté, Poie de Soufre.* (Sulphur. $\mathfrak{Z}$ ij, Potassæ carb. $\mathfrak{Z}$ ij. Rub them together, and place the mixture in a covered crucible upon the fire until they unite.) It has been employed in affections of the chest, as an expectorant, in the dose of from $\mathfrak{Z}$ ss to $\mathfrak{Z}$ j, largely diluted; but is chiefly used as a bath or wash in cases of itch, tinea capitis, and other cutaneous affections.)

POTASSII TERSULPHURETUM, Potassii sulphuretum.

POTASSIUM. Same etymon as potash. The metallic base of potassa.

POTASSIUM BROMURE DE, see Bromine.

POTASSIUM, CHLORIDE OF, Potassæ murias—p. Cyanuret of, Potassii cyanuretum — p. Ferrocy'anide of, Potassii ferrocyanuretum — p. Ferrocy'anuret of, Potassii ferrocy'anuretum — p. Iodide of, Potassæ hydriodas — p. Iodo-hydrargyrate of, Potassii hydrargyroididum — p. Iodure de, Potassæ hydriodas — p. Oxide of, Potass — p. Protoarsenite of, Arsenite of protoxide of potassium — p. Protochlorate of, Potassæ murias hyperoxygenatus — p. Protonitrate of, Potassæ nitras — p. Protoxide of, Potass — p. Protoxide of, nitrate of, Potassæ nitras — p. Protoxide of, superoxalate of,

Potass, oxalate of—p. Protoxide of, supertartrate of, Potassæ supertartras — p. Sulfure de, sulfaté, Potassii sulphuretum — p. Sulphuret of, Potassii sulphuretum — p. Superprotosulphate of, Potassæ supersulphas.

POTATO, ([S.] *batata*.) Solanum tuberosum — p. Carolina, Convolvulus batatas.

POTATO FLY, Lytta vittata.

POTATO, HOG, Convolvulus panduratus.

POTATO OIL, see Oil. fusel. By distilling potato oil with anhydrous phosphoric acid, a volatile colourless oily liquid is procured, which is lighter than water, and boils at 102°. This is *Am'ylene* or *Am'ylene*, (F.) *Amylène*. It is best prepared from fusel oil by the action of chloride of zinc. The vapour has been employed as an anæsthetic.

POTATO, SPANISH, Convolvulus batatas — p. Sweet, Convolvulus batatas.

POTATO VINE, WILD, Convolvulus panduratus.

POTATO, WILD, Convolvulus panduratus.

POTBELLY, Physconia.

POTECARY, Apothecary.

POTELÉE, Hyoscyamus.

POTENCY, (*potens*, 'able,') Faculty.

POTENTIA, (*potens*, 'able,') Force.

POTEN'TIA seu POTES'TAS COEUN'DI, Capability of copulation.

POTENTIA GENERAN'DI. The power of procreation.

POTENTIA IRRITANS, Stimulus.

POTENTIAL, *Potentia*, (F.) *Potentiel*; (from *potentia*.) An epithet for remedies which, although energetic, do not act till some time after their application. The caustic alkalies, for instance, are called *Potential cauteries* in contradistinction to the hot iron, which is termed *actual*.

POTENTILLA, (from *potentia*.) P. reptans.

POTENTIL'LA ANSERI'NA, P. argenti'na seu im'polita seu neglecta, Argenti'na vulga'ris, Fraga'ria anseri'na, Ar'gentine, Chenobos'com, Argenta'ria, Anseri'na, Sil'verweed, Wild tansy, Mooi'-grass, (F.) Argentine, Potentille, Herbe aux vries. Ord. Rosaceæ. The leaves are mildly astringent and tonic.

POTENTILLA ARGENTINA, P. anserina — p. Impolita, P. anserina — p. Neglecta, P. anserina.

POTENTILLA NORVEG'ICA, Norway Cinquefoil, an indigenous plant, is possessed of similar properties.

POTENTILLA PALUSTRIS, Comarum palustre.

POTENTILLA REP'TANS, Pentaphyllum, Fraga'ria pentaphyllum, Potentilla, Quinquefoilium, Q. majus, Com'mon cinquefoil, (F.) Potentille rampante, Quinquefeuille. The roots are possessed of astringent properties.

POTENTILLA TORMENTILLA, Tormentilla.

POTENTILLE, Potentilla anserina — p. Rampante, Potentilla reptans.

POTENTIZING, (from *potentia*,) see Dynamic.

POTERIUM SANGUISORBA, (ποριον,) Pimpinella saxifraga.

POTESTAS COEUNDI, ('power of copulating,') Potentia coeundi — p. Generandi, Potentia generandi.

POTEX, gen. *Po'ticis*, (*podex*.) Anus.

POTHOMORP'HA PELTA'TA, Pothomorp'he pelta'ta, (pothos, and ποπή, 'shape,') Caa'peba [?]. A Brazilian plant, Order, Piperacæ, which is possessed of diuretic properties, and has been given in strangury.

POTHOMORP'HA UMBELLA'TA, also called Caa'peba [?], is given in Brazil to modify nutrition, as in glandular affections, in which it is also used externally, in the form of decoction.

POTHOPATRIDA'L'GIA, (ποθος, 'a longing,' πατρις, 'one's country,' and αλγος, 'pain,') Nostalgia.

POTHOS FÆTIDA, Dracontium fœtidum — p. Putorii, Dracon'tium fœtidum.

POTIMOS, (ποτιμος,) Potable.

POTIO, Drink, Haustus, Potion—p. Calcis carbonatis, Mistura cretæ—p. Effervescens antiemetica, Potion of Riverius—p. Leniens, Looch ex ovo—p. Oleosi, Looch ex ovo—p. Picea, see Pinus sylvestris—p. Riverii citrata, Potion of Riverius.

POTION (F.), Potio; Hydrolé, Hydrolite. A liquid medicine, from 4 to 8 ounces or more in quantity, the composition of which varies, and which is administered in divided doses. It corresponds to the English *Mixture*. See *Mistura*.

POTION PECTORALE, Looch ex ovo.

POTION OF RIVERIUS. A mixture of Lemon Juice and Subcarbonate of Potass, generally given in a state of effervescence. It is the *Potio effervescentes antiemetica dicta Riverii* of the Parisian codex, *Potio Riverii citrata*. The name in the Codex indicates one of its uses.

POTIONIS DESIDERIUM, 'desire for drink,' Thirst.

POTIRON, Cucurbita pepo.

POTIUNCULA, (dim. of potio,) Haustus.

POTOMANIA, (ποτος, 'drink,' and mania,) Delirium tremens.

POTOPARANEA, (ποτος, 'drink,' and paranea,) Delirium tremens.

POTOS, (ποτος,) Potable.

POTOTROMOPARANEA, (ποτος, 'drink,' τρομος, 'trembling,' and paranea,) Delirium tremens.

POTTINGAR, Apothecary.

POTT'S CURVATURE, Vertebral disease.

POTT'S DISEASE, Vertebral disease.

POTUS, (ποτος,) Drink—p. Hippocraticus, Claret.

POU, Pediculus.

POUCE, Pollex, see Digitus.

POUCHED, (from [Sax.] pocea, [F.] poche, 'a bag,') Encysted.

POUCHES, LARYNGEAL, see Laryngeal pouches.

POUDRE, Powder, Pulvis—p. Absorbante et anti-acide, Pulvis de magnesiâ compositus—p. d'Alun composée, Pulvis sulphatis aluminæ compositus—p. Antiarthritique amère, Pulvis de amaris compositus—p. Antiasthmaticque, Pulvis de sulphure et scillâ—p. Antimonial composée ou de James, Antimonial powder—p. Aromatique, Pulvis cinnamomi compositus—p. d'Asarum composée dite sternutatoire, Pulvis asari compositus—p. Balsamique, Pulvis myrrhæ compositus—p. de Birckmann, Pulvis de aro compositus—p. de Blanchement, Calois chloridum—p. de Cannelle composée ou aromatique, Pulvis cinnamomi compositus—p. Castillon, see Powders, Castillon—p. Cathartique, Pulvis scammonæ compositus—p. Caustique du frère Cosme, Powder, arsenical, of Côme—p. de Chypre, see Cladonia rangiferina—p. du Compté de Palme, Magnesîæ carbonas—p. Cornachine, Pulvis cornachini—p. de Corne brûlée avec opium, Pulvis cornuusti cum opio—p. Coton, see Collodion—p. de Craie composée, Pulvis cretæ compositus—p. de Craie composée avec opium, see Pulvis cretæ compositus cum opio—p. de Dover, Pulvis ipecacuanhæ compositus—p. du Duc de Portland, Pulvis Ducis Portlandiæ—p. de Dupuytren, Powder of Dupuytren—p. pour Eau de Vichy, see Vichy, mineral waters of—p. Gazifère laxative, Powders, Seidlitz—p. Gazifère simple, Powders, soda—p. de Gomme adragant composée, Pulvis tragacanthæ compositus—p. de Gomme gutte, Pulvis e gummi guttâ—p. Hydragogue, Pulvis e gummi guttâ—p. Incisive, Pulvis de sulphure et scillâ—p. d'Ipecacuanhæ et opium, Pulvis ipecacuanhæ compositus—p. de Jalap composée, Pulvis jalapæ compositus—p. de Kino composée, Pulvis kino compositus—p. de la Mirindole, Pulvis Ducis Portlandiæ—p. Nutrimen-

tive, see Pepsin—p. de Pérard, Pulvis de sennæ, scammonæ et lignis—p. de Quercetanus, Pulvis e gummi guttâ—p. de Rousselot, Powder, arsenical, of Côme—p. de Saint Ange, Pulvis asari compositus—p. de Savon végétale, Pulvis gummosus alkalinus—p. de Scammonée composée, Pulvis scammonæ compositus—p. de Seidlitz, Powders, Seidlitz—p. de Séné composée, Pulvis sennæ comp.—p. de Sentinelli, Magnesîæ carbonas—p. Styptique, Pulvis sulphatis aluminæ compositus.

POUDRE SUBLTIL DE DELCROIX. A fashionable depilatory "for removing superfluous hair in less than ten minutes." It is said to consist of Quicklime and Sulphuret of Arsenic, with some vegetable powder.

POUDRE TEMPÉRANTE DE STAHL, Pulvis de sulphate potassæ compositus—p. de Tennant, Calcis chloridum—p. de Valentini, Magnesîæ carbonas—p. Vermifuge mercurielle, Pulvis e sulphureto hydrargyri nigro et scammonio—p. Vermifuge sans mercure, Pulvis ex helminthocorto compositus.

POUDRE DE ZWINGER, Magnesîæ carbonas.

POUGUES, MINERAL WATERS OF. These alkaline waters are in the department of Nièvre, France. They contain carbonic acid, carbonates of soda, lime, and magnesia; chloride of sodium; alumina, and silica, with protoxide of iron. Season from the 15th of May to the 1st of October.

POUKE, Papula.

POULAIN, Bubo.

POULET, VIN DE (F.) Pullet Wine. A disgusting preparation, made by infusing the excrement of the pullet in white wine. According to Nysten, it is employed by the vulgar in France in a number of diseases, and especially in falls, contusions, &c.

POULIOT, (from *pulegium*,) Mentha pulegium.

POULS, Pulse—p. Abdominal, Pulsus inferior—p. des Adultes, Pulse, natural—p. Agité, Pulsus fortis—p. Bas, Pulse, low—p. Capital, Pulse, nasal—p. Combiné, Pulse, complex—p. Compliqué, Pulse, complex—p. Composé, Pulse, complex—p. Concentré, Pulse, contracted—p. Défaillant, Pulsus deficiens—p. Déprimé, Pulse, depressed—p. Développé, Pulse, developed—p. de la Diarrhée, Pulse, intestinal—p. Dur, Pulse, hard—p. Elevé, Pulse, ardent—p. Embarrassé, Pulsus intricatus—p. de l'Estomac, Pulse, stomachal—p. Étroit, Pulsus strictus—p. de l'Expectoration, Pulse, pectoral—p. Faible, Pulse, feeble—p. du Foie, Pulse, hepatic—p. Grand, Pulse, large—p. Insensible, Pulsus deficiens—p. Intercedent, Pulse, intermittent—p. des Intestina, Pulse, intestinal—p. Languissant, Pulse, languid—p. Lent, Pulse, slow—p. de la Matrice, Pulsus uterinus—p. Modéré, Pulse, natural—p. Mou, Pulse, soft—p. du Nez, Pulse, nasal—p. Ondoyant, Pulse, undulating—p. Ondulant, Pulse, undulating—p. Palpitant, Pulse, tremulous—p. Parfait, Pulse, natural—p. Petit, Pulse, small—p. Plein, Pulse, full—p. Précipité, Pulse, frequent—p. Profond, Pulse, deep—p. Prompt, Pulse, sharp—p. Rare, Pulse, slow—p. Rebondissant, Pulsus dicrotus—p. Redoublé, Pulsus dicrotus—p. des Règles, Pulsus uterinus—p. Roide, Pulse, tense—p. Serratile, Pulsus serratus—p. Serré, Pulse, contracted—p. de la Sueur, Pulsus sudoralis—p. Tardif, Pulse, slow—p. Tendu, Pulse, tense—p. Tremblant, Pulse, tremulous—p. de l'Urine, Pulsus urinalis—p. Véhément, Pulsus fortis—p. Ventral, Pulsus inferior—p. Vide, Pulsus vacuus—p. Vif, Pulse, sharp—p. Vite, Pulse, sharp.

POULTICE, (πολτις; *puls*, gen. *pultis*, 'porridge, pottage, pap,') Cataplasm. For different poultices, see Cataplasm and Cataplasma.

POUNCE, (F.) Ponce, Sandarac.

POUND, *L'bra*, *L'itra*, *As*, *Pon'do*, *Bage'dia* (of twelve oz.), *Barda'dia*, (F.) *Livre*, (from *pondere*, 'to weigh.') A weight, consisting of twelve ounces Troy,—sixteen ounces avoirdupois. See Weights and Measures.

POUPART'S LIGAMENT, Crural arch.

POURD MILK, Colostrum.

POURPIER CULTIVÉ, Portulaca.

POURPRE, *Purpura*—*p. Blanc*, Miliary fever—*p. de Cassius*, Purple of Cassius.

POURRITURE, (from [L.] *putrere*, 'to rot,') Putrefaction—*p. d'Hôpital*, Hospital gangrene.

POUS, (πους,) Pes.

POUSSE, Pursiness.

POUSSEE, Lichen.

POUSSIERE TUBERCULEUSE (F.), 'tuberculous dust.' A name given by MM. Rilliet and Barthez to fine white or yellow minute points of tuberculous matter occurring in myriads, so as to be likened to dust.

POUSSIF, see Pursiness.

POW, (*poll*,) Head.

POWDER, (F.) *poudre*, Pulvis.

POWDER OF ÆTHIOPIA, GUY'S. A once celebrated antelmintic, which consisted of pure *susped tin*, *mercury*, and *sulphur*, triturated together.

POWDER OF ALOES, COMPOUND, Pulvis aloes compositus, Pulvis sulphatis aluminæ compositus—*p. of Aloes with canella*, Pulvis aloes cum canella—*p. Alum*, compound, Pulvis sulphatis aluminæ compositus—*p. Antimonial*, Chenevix's, Antimonial powder—*p. Aromatic*, Pulvis cinnamomi compositus—*p. Aromatic of chalk*, see Confectio Aromatica.

POWDER, ARSENICAL, OF CÔME, *Côme's powder*, *Pulvis Cos'mi*, (F.) *Poudre caustique du frère Cosme ou du Rousselot*—a caustic—is formed of *Arsenious acid*, gr. x; *Red sulphuret of mercury*, gr. xl; powdered *Animal charcoal*, gr. x. It is made into a paste with mucilage.

POWDER, ARSENICAL, OF JUS'TAMOND, is made by melting *sulphuret of antimony* 16 parts; *arsenious acid* 1 part, in a crucible. When the mass is cold it must be reduced to powder, and five parts of *extract of opium* be added.

POWDER OF ARUM, COMPOUND, Pulvis de arum compositus—*p. of Asarabacca*, compound, Pulvis asari compositus—*p. of Bladder-wrack*, yellow, Pulvis quercus marinæ—*p. Bleaching*, Calceis chloridum—*p. Botanical*, Chinese, see Falsifications, table of—*p. Camboge*, Pulvis egummi guttâ—*p. Cardinal del Lugo's*, Cinchona—*p. Castillon*, see Powders, Castillon—*p. of Catechu*, compound, Pulvis Catechu compositus—*p. of Chalk*, compound, Pulvis cretæ compositus—*p. of Chalk with opium*, compound, Pulvis cretæ compositus cum opio—*p. of Cinnamon*, compound, Pulvis cinnamomi compositus—*p. Côme's*, Powder, arsenical of Côme—*p. of Contrayerva*, compound, Pulvis contrayervæ compositus—*p. Countess's*, Cinchona—*p. Countess of Kent's*, Cantianus pulvis—*p. Cyprus*, Hibiscus abelmoschus—*p. Dover's*, Pulvis ipecacuanhæ compositus.

POWDER, DO'SIA. A celebrated secret remedy in Japan, which, for a thousand years, has caused divine honors to be paid to Kobodaisi, its discoverer.

POWDER OF DUPUYTREN, CAUSTIC; (F.) *Poudre de Dupuytren*, formed of 1 part of *Arsenious acid* and 200 parts of *Mild chloride of mercury*. A mild and manageable caustic for causing an eschar on exposed surfaces, and in lupus, &c.

POWDER, EARL OF WARWICK'S, Pulvis cornachini—*p. of Ethiopia*, Powder of Ethiopia.

POWDER OF FAYNARD. A celebrated powder for arresting hemorrhage, said to be nothing more than the charcoal of beech wood finely pulverized.

POWDER, FEVER, SCHWANBERG'S, Antimonial powder.

POWDER FOLDER. In Pharmacy, an apparatus by which papers containing powders are folded of the desired length.

POWDER, GOUT, Duke of Portland's, Pulvis Ducis Portlandiæ—*p. Gray*, Hydrargyrum cum cretâ—*p. of burnt Hartshorn with opium*, Pulvis cornu cervi cum opio—*p. of Ipecacuanha*, compound, Pulvis ipecacuanhæ compositus—*p. of Jalap*, compound, Pulvis jalapæ compositus—*p. James's*, see Antimonial powder—*p. James's factitious*, Antimonial powder—*p. Jesuit's*, Cinchona—*p. of Kino and opium*, Pulvis kino compositus—*p. of Kino*, compound, Pulvis kino compositus.

POWDER, KNOX'S. A disinfecting preparation, composed of eight parts of *Chloride of sodium*, and three parts of *Chlorinated lime*. An ounce, dissolved in a tumblerful of water, forms a solution of proper strength.

POWDER OF LEMONADE, see Lemonade—*p. of Magnesia*, compound, Pulvis de magnesiâ compositus—*p. of the Marquis*, Marchionis pulvis—*p. Mead's*, against the bite of a Mad dog, Pulvis antilyssus—*p. Myrrh*, compound, Pulvis myrrhæ compositus—*p. Portland*, Pulvis Ducis Portlandiæ—*p. of Rhubarb*, compound, Pulvis rhei compositus—*p. Saline*, compound, Pulvis salinus compositus—*p. of the three Sanders*, see Diatrium—*p. Scammony and aloes*, Pulvis scammonii cum aloë—*p. Scammony*, compound, Pulvis scammonæ compositus—*p. of Senna*, compound, Pulvis senæ compositus—*p. Sneezing*, see Veratrum album—*p. Spruce beer*, see Spruce beer—*p. of Sympathy*, Sympathetic powder—*p. Talbor's*, Cinchona—*p. Tennant's*, Calceis chloridum—*p. Tonquin*, Pulvis cobbi—*p. Tooth*, Dentifrice—*p. Tragacanth*, compound, Pulvis tragacanthæ compositus—*p. Vichy water*, see Vichy, mineral waters of.

POWDER, VIEN'NA, *Pulvis causticus Vindobonensis*; (composed of equal parts of *Quicklime* and *Potassa cum calce*.) In using it, it is made into a paste—*Vien'na paste*, *Pasta caustica Viennensis*, (F.) *Caustique de Vienne*, *Pâte Caustique de Vienne*—with spirits of wine, and is applied on the diseased part, previously circumscribed by a hole out in sticking plaster.

To be left on from 10 to 20 minutes, according to the depth of the eschar required. Used in cases of lupous ulceration.

By fusing the potass and lime, and raising the heat so as to cause ebullition, the caustic may be run into cylinders:—when it is called *Caustique-Filhos*, *Caus'ticum Viennensê fu'sum Fil'hos*, from the name of the author of the process.

POWDERS, CASTILLON or CASTILHON, so called from the name of the physician who first proposed them. They are formed of *Sago*, in powder; *Sulep*, in powder; *Tragacanth*, in powder, each, four parts; *Prepared oysters*, one part, *Cochineal*, half a part, divided into powders of one drachm each; one of which is boiled in a pint of milk, and the decoction used *ad libitum*, as a demulcent in diarrhœa and dysentery. Dorvault makes *poudre de Castillon* synonymous with arrow root.

POWDERS, EFFERVESCENT, Powders, Soda.

POWDERS, EFFERVESCENT, APERIENT, Powders, Sedlitz—*p. Ginger-beer*, see Amomum zingiber.

POWDERS, ROCHELLE, see Powders, soda; and Sedlitz, mineral waters of.

POWDERS, SO'DA, *Pulveres effervescentes*, (Ph. U. S. 1863.) seu *aërophori*, *Effervescentes powders*, (F.) *Powders gazifères simples*. (*Soda bicarb.* 3vj. in *tartaric acid*, 3v. Divide each powder into twelve equal parts; and keep part of the bicarbonate and acid in a paper of different

colours. A pleasant, cooling drink; taken during effervescence.

Sedlitz Powders, Rochelle powders, Aperient effervescing powders, Pulveres aperientes effervescentes (Ph. U. S. 1863.) (F.) *Poudres gazeuses laxatives* ou *Poudres de Sedlitz, (Soda bicarb. in pulv. 5j; Potass. et Soda tartret. in pulv. 5iij; Acid Tartar. in pulv. ʒvij. Mix the bicarbonate with the tartrate, and divide into 12 parts. Divide, also, the tartaric acid into 12 parts. Keep the parts of the mixture and of the acid in separate papers of different colours. See Sedlitz, mineral waters of.*

POWER, ([F.] *Pouvoir*, from *posse*, 'to be able,') Faculty—p. Inherent, Irritability—p. Sensoria, see Sensorial—p. Tonic, Irritability.

POX, (from *pocks*), Syphilis, Variola — p. Chicken, Varicella.

Pox Doctor, Syphilidiater.

POX, FRENCH, Syphilis — p. Grease, see Grease — p. Horn, see Varicella — p. Nodde, Syphilomania — p. Small, Variola — p. Swine, see Varicella — p. Water, see Varicella.

POZZUOLI, MINERAL WATERS OF. Pozzuoli is in the neighbourhood of Naples. The springs are, 1. *Acqua della pietra*, Temp. 91° Fah. 2. *Acqua di Cavalanti*, Temp. 100°. *A. di subveni homini*, Temp. 102°; *A. del cantarello*, Temp. 86 to 89°; and that of the Temple of Serapis in Pozzuoli, Temp. 102 to 107°. The last contains carbonic acid, carbonates of lime, magnesia, alumina, and iron, carbonate of soda, chloride of sodium, sulphate of lime, and silica.

PRACTICE OF PHYSIC, Therapeutics.

PRACTITIONER, GENERAL, Surgeon apothecary—p. Irregular, see Irregular—p. Regular, see Regular—p. Routine, Routinist.

PRÆ. In composition, 'before.'

PRÆBIUM, (from *præbeo*, contracted from *præhibeo*, 'to offer, allow,') Dose.

PRÆCAUTIO, (*præ*, and *caveo*, *cautum*, 'to guard against,') Prophylaxis.

PRÆCIPITANS, (*præcipitare*, 'to throw down,') Absorbent—p. Magnum, Sepia.

PRÆCIPITATUM ALBUM, 'white precipitate,' Hydrargyrum præcipitatum—p. Rubrum, 'red precipitate,' Hydrargyri nitricæoxydum.

PRÆCISIO PRÆPUTII, (*præcidere*, *præcisum*, 'to cut off,') Circumcision.

PRÆCOCIA, (*præcox*, 'ripening early,') Prunus Armeniaca.

PRÆCOCITY, Precocity.

PRÆCOGNITIO, (*præ*, and *cognitio*, 'knowledge,') Prognosis.

PRÆCOR'DIA, (*præ*, and *cor*, gen. *cordis*, 'the heart,') The diaphragm; also, the thoracic viscera, and the epigastrium.

PRÆCOR'DIAL or PRÆCOR'DIAL RE'GION, *Re'gio præcordi'aca* seu *præcordia'lis*, is the epigastric region. Also, and more properly, the region of the heart. A line drawn from the inferior margins of the third ribs across the sternum, passes over the valves of the pulmonary artery a little to the left of the mesial line, and those of the aorta are behind them, but about half an inch lower down. A vertical line, coinciding with the left margin of the sternum, has about one-third of the heart, comprising the upper portion of the right ventricle, on the right,—and two-thirds, comprising the lower portion of the right ventricle and the whole of the left, on the left. This is the præcordial region.

PRÆCORDIUM, *Fossette du cœur* — p. Pedis, Metatarsus.

PRÆCOX MATURITAS, Precocity.

PRÆCUSTODIA, (*præ*, and *custodio*, 'to watch,') Prophylaxis.

PRÆDIASTOLICUS, Prediastolic.

PRÆDICTIO, (*præ*, and *dicere*, *dictum*, 'to say,') Prognosis.

PRÆDISPOSITIO Predisposition.

PRÆDIVINATIO, (*præ*, and *divinare*, *divinatum*, 'to divine,') Mantia.

PRÆDOMIN'IUM, (*præ*, and *domo*, *domitum*, 'I subdue,') Predominance of one thing over another.

PRÆDOMINIUM A'QUE. A predominance of watery fluid,—*Tenu'itas aquo'sa*.

PRÆDOMINIUM SAN'GUINIS ARTERIO'SI. A predominance of arterial blood, *Arterios'itas san'guinis*.

PRÆDOMINIUM SAN'GUINIS VENO'SI. *Venos'itas San'guinis*. Predominance of venous blood.

PRÆFOCATIO, (*præfacere*, *præfocatum*, 'to strangle,') Orthopnoea, Suffocation—p. Faucium, Angone — p. Matricis, Angone — p. Uterina, Angone.

PRÆFOCATION, Strangulation — p. Uterine, Angone.

PRÆFORMATIO, Preamformation.

PRÆGNANS, Pregnant.

PRÆGNAS, Pregnant.

PRÆGNATIO, Fecundation, Pregnancy.

PRÆGNATUS, Fecundation, Pregnancy.

PRÆGNAX, Pregnant.

PRÆGUSTATIO, (*præ*, and *gustare*, *gustatum*, 'to taste,') *Avant-gout*.

PRÆHENSIO, Prehensio.

PRÆLING'UA. The anterior part or tip of the tongue.

PRÆLIUM, ('a contest,') Coition.

PRÆLUM, Press, Tourniquet — p. Arteriale, Tourniquet.

PRÆLUMBARIS, Prelumbar.

PRÆMORPHISMUS, Premorphism.

PRÆNOTIO, (*præ*, and *noscere*, *notum*, 'to know,') Prognosis.

PRÆNUNTIIATIO, (*præ*, and *nuntiare*, *nuntiatum*, 'to announce,') Prognosis.

PRÆPARANTES ARTERIÆ, (from being presumed to prepare the sperm.) Spermatic arteries.

PRÆPARATIO, (*præ*, and *parare*, *paratum*, 'to prepare,') Preparation — p. Chyli, Chylification.

PRÆPARATA VENA, see Facial Vein.

PRÆPARATUM, Preparation.

PRÆPOTEN'TIA, (*præ*, and *potens*, 'powerful,') Great power or strength.

PRÆPOTEN'TIA VIRI'LIS. Great, or too great, procreative power on the part of the male.

PRÆPOTIO, (*præ*, and *potio*, 'a drink,') Potipotismus.

PRÆPUTIALIS, Preputial.

PRÆPUTII DUCTIO, Masturbation.

PRÆPUTIUM, Prepuce.

PRÆRECTALIS, *Prærectal*.

PRÆSAGIUM, (*præ*, and *sagio*, 'to perceive quickly,') Mantia.

PRÆSCIENTIA, (*præ*, and *scire*, 'to know,') Prognosis.

PRÆSCITIO, (*præ*, and *scire*, 'to know,') Prognosis.

PRÆSCRIPTIO, Prescription.

PRÆSCRIPTUM, Prescription.

PRÆSEPIOLUM, (dim. of *præsepium*,) Alveolus.

PRÆSEPIUM, (*præ*, and *sepes*, 'an enclosure,') Alveolus.

PRÆSERVATIO, (*præ*, and *servare*, 'to save,') Prophylaxis.

PRÆSERVATIVUM, Amuletum.

PRÆSERVATO'RIA INDICA'TIO. A preservative indication; one proper for preventing the development of disease.

PRÆSERVATORIUS, Prophylactic.

PRÆSPINALIS, Prespinal.

PRÆTIBIALIS, Pretibial.

PRAIRIE BUSH, STINKING, *Ptelea trifoliata*—p. Pines, *Liatris scabiosa*.

PRASINUM VIRIDE, ('leek green,') *Cupressaceas*.

PRASINUS, (πρασινός, 'green as a leek,') *Porraceous*.

PRASITES, (πρασίτης, from *prasinum*.) Wine in which the leaves of horehound have been infused.

PRASIUM, (πρασινόν,) *Marrubium*.

PRASUM, (πρασινόν,) *Allium prasum*.

PRAXIS, (πράξις,) *Action*.

PRAXIS MED'ICA. The practical exercise of the healing art.

PREASER, *Rennet*.

PRECIPITATE, see *Precipitation*—p. Red, *Hydrargyri nitricooxydum*—p. White, *Hydrargyrum præcipitatum*.

PRECIPITATION, *Præcipitatio*, (F.) *Précipité*, (*præcipitare*, *præcipitatum*, (*præ*, and *caput*, 'the head,') 'to throw headlong,') The action by which a body abandons a liquid in which it is dissolved, and becomes deposited at the bottom of the vessel. The matter, so thrown down or precipitated, is called a *precipitate*.

PRÉCIPITATION DE LA MATRICE, *Procidencia uteri*.

PRÉCIPITÉ BLANC, 'white precipitate,' *Hydrargyrum præcipitatum*—p. Noir, see *Hydrargyri oxydum cinereum*—p. per se, *Hydrargyri nitricooxydum*—p. Rouge, *Hydrargyri nitricooxydum*.

PRECOO'ITY, *Preco'ciousness*, *Præ'cox matu'ritas*, *Matu'rity preco'cious*, (F.) *Précocité*, (*præcox*, gen. *præco'cis*, from *præ*, and *coquere*, 'to ripen,') Rapid development before the usual time. See *Præcopia*.

PRECORDIAL, *Præcordial*.

PRECUR'SORY, *Præcursorius*, *Premon'itory*, *Præmonito'rius*, (F.) *Prémonitoire*, *Précurseur*, (*præ*, and *curro*, *cursum*, 'I run,') That which precedes or foretels.

PRECUR'SORY or PREMON'ITORY SIGNS or SYMPTOMS, (F.) *Signes précurseurs ou prémonitoires*, *S. avant-coureurs*, are those which indicate an approaching disease.

PREDIASTOL'IC or PREDIAS'TOLIC, *Prædiastol'icus*, (*præ*, and *diastole*.) That which precedes the diastole of the heart,—as '*prediastolic* friction sound.'

PREDISPO'NENT or PREDISPOSING CAUSES, see *Causes*, *predisponent*.

PREDISPOSIT'ION, *Prædisposit'io*, *Semin'ia morbo'rum* (Gaubius); (*præ*, and *disponere*, *dispositum*, 'to dispose,') (F.) *Prédisposition*, *Imminence morbide*. That constitution or condition of the body, which disposes it to the action of disease under the application of an exciting cause. Thus, in hereditary diseases, the conformation is such, that but a slight exciting cause may be required to produce them. When the disease arises solely from the predisposition, or under the additional influence of an extremely slight exciting cause, it is, by some, termed a *disposition*.

PREDOR'SAL, *Prædorsalis*, (*præ*, and *dorsum*, 'the back,') That which is in front of the back.

PREDORSAL REGION of the vertebral column is the anterior surface of the dorsal region.

PRÉ-DORSO-ATLOIDIEN, *Longus colli*.

PRÉ-DORSO-CERVICAL, *Longus colli*.

PRÆEXISTENCE OF GERMS, see *Preformation*.

PREFORMATION, *Præformatio*, (*præ*, 'before,' and *formatio*, 'forming,') An ancient theory of generation, which maintained that the germs of every part exist, but not visibly, before fecundation, and that fecundation occasions their de-

velopment. It was the doctrine of *præexistence of germs*.

PREGNANCY, *Pregnatio*, *Pregnatus*, *In-pregnatio*, (from *prægnatio*, *præ*, and *genero*, 'to beget,') *Ingravidatio*, *Hysterocye'sis*, *Gravid'itas*, *G. uterina*, *Eucymon*, *Cye'sis*, *Eucye'sis*, *Eulcysis*, *Eucymo'sia*, *Cyopho'ria*, *Gestatio*, *Fœtatio*, *Ingravidatio*, *Utero-gestatio*; (F.) *Grossesse*. The state of a female who has within her a fecundated germ, which gradually becomes developed in or out of the uterus. When placed in the cavity of the uterus,—by virtue of the vital force by which it is animated, it becomes attached to the inner surface of that viscus, and obtains from it indirectly the nutriment necessary for its gradual development, during the space of ten lunar months or two hundred and eighty days—the ordinary term or duration of *gestation*, although the period may be protracted to 300 days or upwards,—at the expiration of which parturition occurs. Pregnancy is commonly limited to a single fetus: at times, the uterus contains two or three.—See *Reproduction*. Occasionally, one or two moles—the degenerated products of conception—alone occupy the uterus:—at other times, a fetus likewise exists. In certain cases, and owing to accidental circumstances, the fecundated ovum remains in the ovarium, or is arrested in the Fallopian tube. Pregnancy has, accordingly, been distinguished into, 1. *Natural* or *uterine*, when the fetus is carried in the cavity of the womb: and, 2. *Preternatural* or *extrauterine*. *Expœtation*, *Conception vitio'sa*, *Paracye'sis* seu *Eecye'sis*, *Echysterocye'sis*, *Echthysterocye'sis*, *Dysto'cia* seu *Dyscye'sis* seu *Gravid'itis extrauterina*, (F.) *Grossesse contre-nature*,—when it occurs in the ovarium, tube, or cavity of the abdomen.

During pregnancy, the female experiences remarkable changes in the condition of her organs and functions. The genital apparatus is, of course, most affected. The uterus changes its shape, situation, size, and texture. The usual signs of pregnancy are,—suppression of the catamenia; the abdomen gradually enlarging, and forming a hard tumour; the umbilicus prominent in the latter months; the breasts increasing in size; the areola round the nipple becoming darker, and the follicles more developed; kyestein in the urine from an early period; a bluish tint of the vagina after the fourth week; about the fifth month, the evidence by *ballotement*, the *placental souffle*, and by the sound of the fetal heart heard on auscultation; the motion of the fetus perceptible, and the neck of the uterus diminishing in length at the advanced periods.

PREGNANCY, ABDOM'INAT, *Eecye'sis abdominalis*, *Caliocye'sis*, *Gravid'itas abdominalis*, (F.) *Grossesse abdominale*. That in which the ovum has escaped into the cavity of the abdomen.

PREGNANCY, AFE'TAL, *Pregnancy without a fœtus*, (F.) *Grossesse afœtale* ou *sans fœtus*. Pregnancy produced by a false or degenerate conception, or any other affection, which has given place to unusual development of the uterus. Madame Boivin includes in this class, *sarcophysteria*, *hydrohysteria*, *gastrohysteria*, and *hematohysteria* pregnancies.

PREGNANCY, BIGEM'INAL, (F.) *Grossesse bigémale* ou *gémellaire*, *Double pregnancy*. Pregnancy in which there are twins in utero.—Madame Boivin.

PREGNANCY, COM'PLEX, (F.) *Grossesse complexe*, *G. inter-extra-utérine*, is when the womb, besides a fetus, contains much water, hydatids, a mole, &c.

PREGNANCY, COM'PC'IND, *Polybreph'ia*, *Polycye'sia*, *Polycye'sis*, (F.) *Grossesse composée*, is when there are two or three fetuses in the cavity of the uterus.

PREGNANCY, DOUBLE, *Pregnancy, bigeminal*—

p. Duration of, see Pregnancy—p. Extrauterine, see Pregnancy.

PREGNANCY, FALSE, Graviditas spuria, (F.) Grossesse fausse ou apparente, Pseudogestis. Authors have thus called different affections of a more or less serious character, which resemble pregnancy; so as, at times, to impose upon the best informed practitioners. These diseases may be seated in the uterus or its dependencies, in the intestines or some other part of the abdomen. They may be;—a mole, hydatids, water, blood, or polypus distending the parietes of the uterus;—chronic enlargement of the uterus; development of tumours, &c., in its substance;—scirrhus; dropsy of the ovaria; ascites; tympanites, &c.

PREGNANCY, FETAL, (F.) Grossesse fœtale. Pregnancy in which there is a fœtus, or more than one, in the uterus or out of it. Madame Boivin puts in this class—the solitary, bigeminal, trigeminal, and sarcofœtal pregnancies, which form the first order; and the tubal, ovarian, and abdominal, which constitutes the second.

PREGNANCY, GAZOXYSTER'IC, (F.) Grossesse Gazo-hystérique. Fœtal pregnancy, caused by development of air or gas in the cavity of the womb.—Madame Boivin.

PREGNANCY, HÆMATOXYSTER'IC, (F.) Grossesse hémato-hystérique. Fœtal pregnancy, caused by accumulation of blood in the uterus.—Madame Boivin.

PREGNANCY, HYDROXYSTER'IC, (F.) Grossesse hydro-hystérique. Fœtal pregnancy, occasioned by the secretion and accumulation of serous fluid in the uterus.

PREGNANCY, INTERSTIT'IAL, Graviditas in uteri substantia seu interstitia. The development of the embryo in the proper substance of the uterus. The case generally terminates fatally at an early period of gestation; the thin external paries of the uterus being ruptured, and the ovum passing into the abdominal cavity.

PREGNANCY, MOR'BID, Paracystis, Dystocia, Dyseystis. The progress of pregnancy disturbed or endangered by the supervention of general or local disorder.

PREGNANCY, OVA'RIAN, Eccyestis ova'ria, Oöriocystis, Oöcyestis, Graviditas uteri seu extrauterina in ova'rio, (F.) Grossesse ovarienne ou de l'ovaire. Pregnancy in which the fœtus is developed in the interior of the ovary.

PREGNANCY, PLEA OF. A plea, which a woman, capitally convicted, may set up in arrest of execution until she is delivered. See Jury of Matrons, and De ventre inspicendo.

PREGNANCY, PRETEND'ED. The criminal act of a female, who, not having been delivered, presents an infant as having been born of her. (F.) *Supposition de part.* This term, in French, likewise includes the case of a female, who, having been delivered of a dead child, substitutes for it a living child which is not hers.

PREGNANCY, SARCOFETAL, (F.) Grossesse sarco-fœtale. Pregnancy in which there are one or more fœtuses, and one or more moles.—Madame Boivin.

PREGNANCY, SARCOXYSTER'IC, (F.) Grossesse sarco-hystérique. Fœtal pregnancy formed by moles, hydatids, or a false conception.—Madame Boivin.

PREGNANCY, SOL'ITARY, (F.) Grossesse solitaire ou simple. Pregnancy in which there is a single fœtus.—Madame Boivin.

PREGNANCY, TRIGEM'INAL, (F.) Grossesse trigéniale ou triple. Pregnancy in which there are three fœtuses in the cavity of the uterus.—Madame Boivin.

PREGNANCY, TU'BAL, (F.) Grossesse tubaire; Eccyestis tubalis, Graviditas tubaria. Pregnancy where the fœtus remains and is developed in the Fallopian tube.

PREGNANCY, UTERINE, see Pregnancy.

PREGNANCY, UTEROABDOMINAL, (F.) Grossesse utero-abdominale. Pregnancy consisting of two fœtuses; one in the uterus—the other in the abdominal cavity.—Madame Boivin.

PREGNANCY, UTEROÖVA'RIAL, (F.) Grossesse utéro-ovarienne. Pregnancy with two fœtuses, one in the womb, the other in the ovary.

PREGNANCY, UTEROTU'BAL, (F.) Grossesse utéro-tubaire. Pregnancy in which there is at the same time a fœtus in the womb and another in the Fallopian tube.

PREGNANT, Prægnans, Prægnas, Prægnax Fetus, Fœtus, Encarpus, Gravida, Plena mulier, Incincta, Gravid, Breed'ing, (Old Eng.) Beal'ing, (F.) Enceinte. One in a state of pregnancy;—one with child.

PREHENSIO, (prehendere, prehensum, 'to lay hold of,') Prehension, Catalepsy, Epilepsy.

PREHENSION, Prehensio. The act of laying hold of.

PREHENSION OF FOOD is the act of seizing and carrying it to the mouth, and introducing it into that cavity. It is, also, called *Assumptio, Proslepsis, Prosphora.*

PRÊLE, ([I.] asparello, 'rough,'—the stalk,) Hippuris vulgaris.

PRÊLOMBAIRE, Prelumbar.

PRÊLOMBO-PUBIEN, Psos parvus—p Trochantin, Psos magnus.

PRELUM, Press—p. Arteriale, Tourniquet.

PRELUM'BAR, Prælumbaris, (F.) Prêlombaire, (præ, and lumbi, 'the loins,') That which is placed before the loins. The prelumbar surface of the spinal column is the anterior surface of the lumbar portion.

PREMIÈRES VOIES, 'first passages,') Primæ viæ.

PREMOLAR, (præ, and molares,) see Molar teeth.

PREMONITORY, (præ, and moneo, 'to admonish,') see Precursory.

PREMORPHISM, Præmorphismus, (præ, and μορφή, 'I shape,') The aggregate of laws by which a new formation is developed from the primordial cell.

PRENAN'THES, (πρηνος, 'bending downward,') and ανθος, 'a flower,') *Gall of the Earth, Devil Snake-root, Lion's Foot.* There are many species of this indigenous plant, whose root and milky juice are very bitter. They are used in popular practice in dysentery; and in the form of cataplasm to bites of serpents.

PRENANTHES ALBA, Nabalus albus—p. Serpentaria, Nabalus albus.

PRENDRE INSCRIPTION, 'to take inscription,') see Matriculate.

PRÉPARATE (VEINE), see Facial vein.

PREPAR'ATION, Præparatio, (præ, and parare, paratum, 'to get ready,') The act of making ready any substance whatever that has to be employed for any purpose.

Also, the product of any pharmaceutical operation;—*Præparatum.*

Any part of a body, preserved for the use of the anatomist, or others, is so called.

PREPUCE, Præputium, Epugo'gium, Pos'thæ, Pelt'a, Pos'thium, Put'tium, Acropos'thia, Acrohy'ntia, Fore'skin, (said to be from præ, and puto, 'to cut off,') or from puteo, 'to smell,') or from præ, and pudendum; but, most probably, from pro, or præ, and ποσιον, 'penis,') The prolongation of the integuments of the penis, which covers the glans. The prepuce is composed of two membranous layers: one external or cutaneous; the other internal or mucous, separated by areolar membrane. The mucous membrane lines the inner surface of the cutaneous layer till beyond the glans, when it is reflected over the latter; forming, behind the

corona, a small *cul-de-sac*, above the corpora cavernosa, which is interrupted by a triangular duplication of the mucous membrane, called the *frænum*, fixed in the furrow at the inferior part of the glans, and terminating at a little distance from the orifice of the urethra. The clitoris is covered by a semilunar fold formed by a continuation of the skin of the labia. It is called *Præputium clitoridis*.

PREPUTIAL, *Præputialis*. Belonging or relating to the prepuce;—as the 'preputial secretion,'—or that which takes place from the lining of the prepuce covering the glans.

PRÆRECTAL, *Prærectalis*, (*præ*, and *rectum*.) applied to that which is in front of the rectum.

PRÉROTULIEN, see *Bourse*.

PRESBYODOCHIUM, (*πρεσβυος*, 'old,' and *δοχμα*, 'to receive,') Gerocoemium.

PRESBYONOSI, (*πρεσβυος*, 'old,' and *νοσος*, 'disease,') Diseases of old age.

PRESBYOPE, *Presbytes*, (F.) *Presbyte*, (*πρεσβυος*, 'old,' and *ωψ*, 'the eye,') One who is long-sighted or affected with presbyopia.

PRESBYOPIA, *Presbytia*.

PRESBYOPIC, *Presbyopicus*, *Presbytic*, *Presbyticus*, (F.) *Presbytique*. Relating or belonging to presbyopia or longsightedness.

PRESBYSPHACELUS, (*πρεσβυος*, 'old,' and *σφακελος*.) Gangrene of old people.

PRESBYTIA, *Presbyopia*, *Telora'sia*, *Amblyopia proximo'rum*, *Visus senilis*, *Hypermetropia*, *Presbyopia*, (F.) *Vue longue*; (from *πρεσβυς*, 'an old person,') A very high degree is termed *Hyperpresbytia*. A condition of vision, common in old persons, which consists in the circumstance of near objects being confusedly seen, whilst those at a greater distance are clearly distinguished. It is supposed to depend generally on diminution of the convexity of the cornea; whence results an alteration in the convergency of the rays, so that they form a focus behind the retina. This defect is remedied by the use of convex glasses, which augment the convergence of the luminous rays. It is, also, called *Long'sightedness*, *Far'sightedness*, *Parop'sia longinqua*, *Dysopia proximo'rum*.

PRESBYTIC, *Presbyopic*.

PRESCRIPTION, *Præscriptio*, (*præ*, and *scribere*, *scriptum*, 'to write,') *Anagraphe*, *Formula*, *F. medica*, *Schedula*, *Imperium*, *Præscriptum*, *Præscriptio medica*, *Receptum*, *Recepta*, *Receptum*, (F.) *Formule*, *Ordonnance*, *Recette*. The formulæ which the physician writes for the composition of medicines adapted to any case of disease. A prescription should be as simple as possible, and should bear upon its face the evidence of the objects to be fulfilled by it. No article should form part of it, unless adapted for serving some useful purpose. A compound prescription has been divided into four parts:—the *basis* or principal ingredient of the prescription; the *adjuvans*, or that which is designed to promote the action of the former; the *corrigens*, or that intended to correct its operation, or obviate any unpleasant symptom which it may be apt to produce; and the *constituens*, *excipiens* or *vehiculum*, *constituent*, *excipient* or *vehicle*, the substance which gives to the other ingredients consistence or form. All these are seen in the following formula for cathartic pills:

R—Aloës..... ʒi *Basis*.
Hydrarg. Chlorid. Mit. gr. x. *Adjuvans*.
Olei Carui..... gtt. v. *Corrigens*.
Syrupi..... q. s. *Constituent*.
M. et fiant pilulæ xxx. See *Symbol*.

It is obvious, however, that most prescriptions are more simple than this. The basis, for exam-

ple, may require neither adjuvant, corrigent, nor constituent.

The doctrine of medicinal combination is termed *Iamatotaxiologia*, *Iamatotaxiologia*, *Artis formulæ medicæ concinnandi*.

The vocabulary in the author's "Medical Student," second edition, Philad., 1844, will aid not only in translating, but in writing, prescriptions more *solito*. See, also, Pereira, *Selecta à Præscriptis*, 14th edition, London, 1864.

PRESENTATION, from (Lit.) *præsens*, (*præ*, 'before,' and *ens*, *entis*, 'being,') The part of a fœtus which is felt *presenting*, on examination *per vaginam*. When the head presents, and especially the vertex, or the feet, knees, or breech, the presentation is said to be *natural*; when any other part, *preternatural*; and the labour is styled *perverse* or *preternatural*, *Parodynia* seu *Dystocia perversa*, *Cross-birth*, (F.) *Accouchement contre nature*. When any part besides the head, feet, knees, or breech presents, the operation of turning becomes necessary.

When a portion of the presenting mass of the fœtus becomes changed for another, in complicated presentations, the case has been termed one of *substituted presentation*.

PRESERVATIVE, (*præ*, and *servare*, 'to save,') *Prophylactic*.

PRESIS, (*πρεσις*.) Swelling.

PRÊSLE, (*Prêle*.) *Hippurus vulgaris*.

PRESMA, (*πρεσμα*.) Swelling.

PRESPI'NAL, *Præspinalis*, (*præ*, and *spina*, 'the spine,') That which is situate before the spine. The prespinal surface of the vertebral column is the anterior surface.

PRESS, *Preclum*, *Prælum*, *Pressorium*, *Pièster*, *Pièsterion*, (F.) *Pressoir*, *Pressé*, (*premere*, *pressum*, 'to squeeze,') An instrument for subjecting matters to considerable pressure, for the purpose of separating the liquid from the solid portion.

PRESS OF HEROPHILUS, *Torcular Herophili*.

PRESSE-ARTÈRE, *Serre-nœud de Deschamps*, *Serre-Artère*. An instrument invented by Deschamps for the immediate compression of arteries. It is composed of a plate six or seven lines long, and three broad, and of a rod or stem two inches long, placed perpendicularly above the plate and riveted. At each extremity of the plate is a hole, through which each of the ends of the ligature, to be placed upon the artery, is passed. The ligature is then to be tightened by drawing the ends, and fixing them to the perpendicular rod. This *presse-artère* or *serre-artère*, surrounded with charpie, remains in the wound, until there is no longer danger of hemorrhage. The ligature is then cut, and the instrument withdrawn. It is an unnecessary instrument.

PRESSE-URÈTHRE, *Jugum Pe'nis*. A kind of forceps of elastic iron, the branches of which, padded, may be approximated or separated by means of a screw. The penis is introduced between the branches, one of them being applied to the urethra, and the screw is tightened so as to exercise a degree of pressure sufficient to bring the parietes of the canal in contact. This instrument is sometimes used in incontinence of urine, to prevent the constant discharge of that fluid.

PRESSIO, (*premere*, *pressum*, 'to press,') *Pressure*.

PRESSIO ABDOMINALE, *Pressure*, abdominal.

PRESSOIR, *Press*—*p. d'Hérophile*, *Torcular Herophili*.

PRESSORIUM, *Press*.

PRESSURE, *Pressio*. The action of pressing.

PRESS'URE, ABDOM'NAL, (F.) *Pressio Abdominale*. A mode of examination by means of which the sensibility, size, suppleness, or hardness of

any morbid abdominal viscus may be compared, in these respects, with its healthy condition. Bichat proposed, with the view of exploring the state of the thoracic organs, to exert a pressure from below upwards on the abdominal organs, so as to crowd them upon the diaphragm, and thus to diminish the dimensions of the cavity of the chest, noticing to what extent respiration was affected by this action. No great information can of course be derived from it.

PRESSURE, *Rennet*.

PRESYSTOLIC or **PRESYSYSTOLIC**, *Præ-systolicus* (*præ*, and *systole*.) That which precedes the contraction of the heart—as ‘*presystolic friction sound*.’

PRETENDED DISEASES, Feigned diseases.

PRETIBIAL, *Prætibialis*, (*præ*, and *tibia*, ‘the tibia.’) That which is situated before the tibia; as the *iliopretibial* and *ischiopretibial* muscles.

PREVENTIVE, (*prævenire*, *præventum*, (*præ*, and *venire*), ‘to come before.’) Prophylactic.

PREVENTIVE, **ROYAL**. An empirical preparation, used as a prophylactic against the venereal virus. It is a solution of *acetate of lead*,—the *Liquor Plumbi Subacetatis dilutus*.

PREVERTEBRAL APONEUROSIS. The aponeurosis which covers the muscles of the prevertebral region.

PREVERTEBRAL ARTERY, (F.) *Artère prévertébrale ou méningienne postérieure*. A ramus of the meningeal branch, and sometimes even of the trunk, of the pharyngeal artery, which is distributed to the prevertebral region.

PREVERTEBRAL MUSCLES, see Cervical region, anterior—p. Region, Cervical region, anterior.

PRIAPEIA, (after Priapus.) *Nicotiana rustica*.

PRIAPIS/CUS, *πριαπικός*.) Name of a piece of wood which forms part of the *scamnum Hippocraticum*, and is like the male organ of generation. Paulus, of Ægina, calls thus a tent of linen shaped like a penis.

PRIAPISM/US, (*πριαπισμός*), *Tenti'go*, *T. Pe'nis*, *Pri'apiem*, *Horncolic* (vulg.), *Enta'sia Priapi's-mus*, *Rhopali's-mus*, *Styl'ma*, *Styl'sis*, (after Priapus.) Constant and distressing erection, with or without any voluptuous idea or desire for venery. It is often symptomatic; and sometimes the immediate result of morbid causes,—of the action of cantharides, for example. The application of cold liquids to the penis, and the internal use of opium and camphor, are the means usually had recourse to for its removal.

PRIAPITIS, (*priapus*, ‘penis,’ and *itis*.) Phallitis.

PRIAPUS, (*Πριαπός*), *Mutun'mus*. In ancient mythology the deity who presided over gardens and the parts of generation, so called *propter deformitatem et membri virilis magnitudinem*.

PRIAPUS, Penis—p. Cervi, see Cervus—p. Ceti, Leviathan penis.

PRICKET, *Sedum acre*.

PRICKING, (from [Sax.] *priccean*, ‘to prick.’) (F.) *Picotement*. A variety of pain, compared to that which pointed bodies would occasion.

PRIDE OF CHINA, *Melia azedarach*—p. of India, *Melia azedarach*—p. Tree, *Melia azedarach*.

PRIDEWEED, *Erigeron Canadense*.

PRIEST PHYSICIANS, see Asclepiadæ.

PRIEST'S PINTLE, *Arum maculatum*.

PRIM, *Ligustrum vulgare*.

PRIMA CELLA COLI, ‘first cell of the colon,’ *Cæcum*.

PRIMÆ VIÆ, (F.) *Premièræ viæ*. ‘The first passages.’ The stomach and intestinal canal:—the latter being the *secundæ viæ* or ‘second passages.’ See Ways, digestive.

PRIMAIRE, Primary.

PRI'MARY, *Primarius*, *Primiti'vus*, (F.)

Primiti'f, *Primaire*, (from *primus*, ‘first.’) First in place, time or character; as *primary symptoms*, in contradistinction to *secondary*.

PRIMARY CELL, see Cell.

PRIMEVÈRE, (from *primum ver* ‘first spring.’) *Primula veris*.

PRIMIP'ARA, *Primipartu'riens*, (F) *Primipare*, (*primus*, ‘first,’ and *parere*, ‘to bring forth.’) A female who brings forth for the first time.

PRIMIPARTURIENS, (*primus*, ‘first,’ and *parturio*, ‘to bring forth,’) *Primipara*.

PRIMISTERNAL, *Primisternal'is*, (*primus*, ‘the first,’ and *sternum*, ‘the sternum.’) M. Béchard has called *os primisternal* on *clavisternal*, the bony portion of the sternum.

PRIMIT'VE, (from *primus*, ‘first.’) The first waters or the waters discharged before the extrusion of the fœtus.

PRIMITIVE, Primary—p. Band, see Nerve fibres—p. Streak, Groove, primitive.

PRIMORDIAL CELL, see Cell.

PRIMORDIUM, (*primus*, and *ordior*, *orais*, ‘to begin,’) *Arche*.

PRIMROSE, **EVENING**, *Oenothera biennis*—p. Tree, *Oenothera biennis*.

PRIMULA OFFICINALIS, (*prim'ulus*, ‘first,’ so called, because it flowers in the beginning of the spring.) P. *veris*—p. *Variabilis*, P. *veris*.

PRIM'ULA VE'NIS, P. *officina'lis* seu *varia'bilis*, *Verbas'culum*, *Paral'ysis*, *Her'ba paralys'eos* seu *paralytica*, the Cow'slip, *Pai'gil* or *Pea'gyle*, *Herb Pe'ter*, *Pal'sywort*, (Prov.) *Cow'striplings*, *Cow'strophle*, *Crow'el*, (F.) *Primevère*, *Herbe à la paralysie*, *Fleur de coucou*, *Oreille d'ours*, *Bravète*; *Ord.* *Primulacæ*. *Ser. Syst.* *Pentandria Monogynia*. The flowers have been considered to be mildly tonic, antispasmodic, and anodyne.

PRIMULA VULGA'NIS. The leaves and roots of this plant have been used as snutatories.

PRINCE ARTHUR'S FARINACEOUS FOOD, see *Farinaceus*.

PRINCIPS, gen. *Prin'cipis*, Rectum.

PRIN'CEPS CERVI'CIS (*Arteria*), *Cer'vical Art'ery*. A branch of the occipital artery, which descends between the splenius and complexus muscles, and may be followed down to the lower part of the neck.

PRINCIPS POL'ICIS (*Arteria*), *Arte'ria magna pol'licis*. When the two collateral arteries of the thumb take origin from a common trunk—the trunk is the *Princeps pollicis*.

PRINCIPE CRYSTALLIZABLE DE DÉ-ROSNE, (from *principium*.) *Narcotine*—p. *de la Mélanose*, see *Choroidea tunica*.

PRINCIPES IMMÉDIATS, Principles, immediate.

PRINCIPIUM, ‘a beginning,’ (from *princeps*, ‘the first,’) *Arche*, Element, Principle—p. Acidificans, Oxygen—p. Adstringens, Tannin—p. Byrsodepiscum, Tannin—p. Coriaceum, Tannin—p. Hydrogeneticum, Hydrogen—p. Hydroticum, Hydrogen—p. Oxygenans, Oxygen—p. Salivale, see Saliva—p. Seytodepiscum, Tannin—p. Urinosum, Urea.

PRINCIPLE, (*princeps*, ‘the first,’) Element, Principium.

PRINCIPLE, COLOURING, OF THE BLOOD. See Hæmaphæin, and Hæmatin. P. Digestive, Pepsin—p. Vital, see Vital principle.

PRIN'CIPLES, **IMME'DIATE**, (F.) *Principes immédiats*, *Matériaux immédiats*. A name given to a considerable number of substances, composed of at least three elements, and which are obtained from animals and vegetables without alteration, by simple processes, and, in some measure, immediately.

PRINCIPLES, IMMEDIATE, OF ANIMALS, *Pro'imate principles*, *Organi'c El'ements*, *Compounds of Organization*. These are divided into acid

principles, fatty principles and principles which are neither fat nor acid. The first include the *uric, rosacic, purpuric, allantoinic, caseic, butyric, &c.* The fatty principles are:—*stearin, elain, cholesterin, &c.*; and the last division includes the most important compounds of organization:—*fibrin, gelatin, albumen, casein, osmazome, &c.*

PRINCIPLES, PROXIMATE, *P.* immediate.

PRINOS (of the ancients), (*πρινος*), *Ilex aquifolium*.

PRI'NOS, *P. verticillatus*, *Alcan'na major latifolia dentata*, *Aquifolium foliis deciduis*, *Pri'nos Gronovii seu padifolius*, *Black Alder*, *Virgin'ia* or *Whor'led Win'terberry*, *Feverbush*, (*F.*) *Apalachine à feuilles de Prunier*. *Ord. Aquifoliaceae*. *Sex. Syst.* Hexandria Monogynia. The bark of this common shrub of the United States—*Prinos*, (*Ph. U. S.*)—is astringent, bitter, and pungent. The berries are, likewise, bitter. The bark has been used as a substitute for the cinchona in intermittents, and as a tonic.

PRINOS GLABER, see *P. laevigatus*—*p.* Gronovii, *Prinos*.

PRINOS LÆVIGATUS, *Smooth Win'terberry*, and *PRINOS GLABER*, *Ink'berry*; indigenous; have the same properties as *Prinos*.

PRINOS PADIFOLIUS, *Prinos*.

PRION, (*πριων*), *Saw*.

PRIONO'DES, (*prion, and odes*), *Serrated*. Applied to the sutures of the head.

PRI'OR, 'first, foremost,' as

PRIOR ANNULA'RIS, *Interos'seus quar'tus* of Winslow. An internal interosseous muscle which arises from the outside of the metacarpal bone of the ringfinger, and is inserted into the outside of the tendon on the back of the ringfinger. Its use is to draw the ringfinger outwards.

PRIOR IN'DICIS, *Extensor ter'tii interno'dii in'dicis*, (*F.*) *Sous-métacarpo-latéri-phalangien*. It arises from the outer or radial side of the metacarpal bone of the forefinger, and is inserted into the outside of the tendon on the back of the forefinger. *Action*:—to draw the finger outwards, towards the thumb.

PRIOR INDICIS PE'DIS, *Abduc'tor In'dicis pe'dis*. An external, interosseous muscle of the foot, which arises from the contiguous sides of the metatarsal bones of the great and fore toes, and is inserted into the inside of the root of the first bone of the foretoe. *Action*:—to pull the foretoe inwards.

PRIOR MEDII, *Prior medii digiti*.

PRIOR ME'DII DIG'ITI, *Prior Me'dii, Sec'ond interos'seus* of Douglas. An external interosseous muscle of the hand, which arises from the corresponding sides of the metacarpal bones of the fore and middle fingers, and is inserted into the outside of the tendon on the back of the middle finger. *Action*:—to draw the middle finger outwards.

PRIOR ME'DII DIG'ITI PE'DIS, *Adduc'tor me'dii dig'iti pe'dis*. An internal, interosseous muscle of the foot, which arises from the inside of the metatarsal bone of the middle toe, and is inserted into the inside of the root of the first bone of the middle toe. *Use*:—to pull the middle toe inwards, or towards the inner side of the foot.

PRIOR MIN'IMI DIG'ITI, *Adduc'tor min'imi dig'iti*. An internal, interosseous muscle of the foot, which originates from the inside of the metatarsal bone of the little toe, and is inserted into the inside of the root of the first bone of the little toe. *Action*:—to pull the little toe inwards.

PRIOR TER'TII DIG'ITI PE'DIS, *Adduc'tor ter'tii dig'iti pe'dis*. An internal, interosseous muscle of the foot, which arises from the inner and under part of the metatarsal bone of the third of the small toes, and is inserted into the side of the root of the first bone of the third of the small

toes. *Action*:—to pull the third of the small toes inwards.

PRISE, Dose.

PRISIS, (*πρισις*), *Brygmus*, *Trepanning*.

PRISMA, gen. *Prism'atis*, (*πρισμα*), *Scobs*.

PRISMUS, (*πριμ*, 'to saw, to grind,') *Brygmus*, *Trepanning*.

PRIVATE PARTS, Genital organs.

PRIVET, *Ligustrum vulgare*.

PRIVITIES, Genital organs.

PRIVY, *Ligustrum vulgare*—*p.* Members, Genital organs—*p.* Parts, Genital organs.

PRO, *προ*, 'for, before.' A common prefix to words.

PROAGOREUSIS, (*προαγορευσις*, from *pro*, and *αγορευω*, 'to announce,') *Prognosis*.

PRO'BANG, *Sponge probang*, *Detrusorium ex bale'na*. (Allied to *probe*.) A long slender rod of whalebone, with a piece of sponge at its extremity, intended to push down extraneous bodies, arrested in the œsophagus, into the stomach. A similar instrument is called by J. L. Petit, *Repoussoir d'arêtes*.

PROBARBIUM, (*pro*, and *barba*, 'beard,') *Geneias*.

PROBASCANIUM, (*προβασκανιον*, from *pro*, and *βασκανειν*, 'to cast a spell upon,') *Amuletum*.

PROBASCANTIUM, *Amuletum*.

PROBE, (*probo*, 'to try,') *Catheterize*, *Specillum*—*p.* Eyed, see *Specillum*.

PROBING, see *Melosis*.

PROBOLE, (*προβολη*, from *pro*, and *βαλλω*, 'to throw,') *Processus*.

PROCARDIUM, (*pro*, and *καρδια*, 'heart,') *Scrobiculus cordis*.

PROCÉDÉ, *Process*.

PROCEDURE, *Process*.

PROCEPHALI, (*προκεφαλοι*, from *pro*, and *κεφαλη*, 'head,') *Capitones*.

PROCE'US ('tall') *NASI*, *Pyramidalis nasi*.

PROCES, *Process*—*p.* *Ciliaires*, *Ciliary processes*.

PROCESS, (*procedere, processum*, 'to go forwards,') *Apophysis*, *Processus*.

PRO'C'ESS, *Ra'tio*, (*F.*) *Procédé*. *Procedure*,—method of performing any operation, chemical, pharmaceutical, or surgical.

PROCESS, CUNEIFORM, *Basilar process*—*p.* *Modelling*, see *Modelling Process*—*p.* of *Rau*, see *Malleus*—*p.* *Vertical superior longitudinal*, *Falx cerebri*.

PROCES'SUS. *Aproc'ess*. *Projectu'ra*, *Prob'ole*, (*F.*) *Proceda*. An apophysis or eminence of a bone; see *Apophysis*. Also, any part which seems prolonged beyond others with which it is in connexion; as the *Ciliary processes*.

PROCESSUS ANCONÆUS, *Olecranon*—*p.* *Annularis*, *Pons Varolii*—*p.* *Ascendens*, see *Maxillary bone*, (*superior*)—*p.* *Belenoides*, see *Belenoid*—*p.* *Blumenbachii*, *Unciform process*—*p.* *Caudatus*, *Lobulus caudatus*—*p.* *Cerebelli ad pontem*, *Peduncles of the cerebellum*—*p.* *Cerebelli ad testes*, *Peduncles of the cerebellum*—*p.* à *Cerebello ad medullam oblongatam*, *Corpora restiformia*—*p.* *Cerebri lateralis*, *Cornu ammonis*—*p.* *Ciliares*, *Ciliary processes*—*p.* *Clavatus*, see *Funiculi graciles*—*p.* *Cochleariformis*, see *Tympanum*—*p.* *Cuneiformis ossis occipitis*, *Basilar process*—*p.* *Dentatus*, see *Axis*—*p.* *Ensiformis*, see *Sphenoid bone*—*p.* *Enteroides cerebri*, *Convulsions (cerebral)*—*p.* *Falciformis cerebelli*, *Falx cerebelli*—*p.* *Falciformis duræ matris*, *Falx cerebri*—*p.* *Frontalis*, see *Maxillary bone*, (*superior*)—*p.* *Gracilis* of *Rau*, see *Malleus*—*p.* *Laterales uteri*, *Tubæ Fallopiæ*—*p.* *Mamillares*, *Papillæ of the kidney*, *Olfactory nerves*—*p.* *Mamillares cerebri*, *Mamillary tubercles*—*p.* *Mastoides*, *Mastoid process*—*p.* *Nasalis*, see *Maxillary bone* (*superior*)—*p.* *Odontoides*, see *Axis*—*p.* *Olivaris*, *Olivary Process*—*p.* *Orbicu-*

PRODUCTION, *Productio*. The act of producing; the thing produced. Also used, especially by the French, synonymously with *production*. The mesentery is, in this sense, a production of the peritoneum.

PRODUCTION ACCIDENTELLE. An accidental or adventitious structure.

PRODUCTIVITAS, Fecundity.

PRODUIT, Product.

PRODUCTUM, Product.

PROËGUMENE CAUSÆ, (*προγενεως*, from *pro*, and *γενεσις*, 'to lead.') Causes, predisponent.

PRÆLUM, Press, Tourniquet.

PRÔËMÎNENT, *Proëminens*, (*pro*, and *eminere*, 'to project.') Some anatomists call the 7th cervical vertebra the *proëminent vertebra*, on account of the length of its spinous process, which passes the level of that of the neighbouring vertebrae.

PROENCEPHALUS, (F.) *Proencéphale*, (*pro*, and *encephalos*, 'encephalon.') A monster, whose encephalon is situated, in a great measure, out of, and in advance of, the cranium, which is open in the frontal region.

PRÔTÏA, *Proïotes*, (*πρωιotes* or *πρωιotes*, from *πρωι*, 'early.') 'Precocity.' *Præcox matu'ritas*, *Præcox matu'ritas*. Premature development of sexual organization or power.

PROFLUVII CORTEX, Nerium antidysentericum.

PROFLUVIUM, *Discharge*, *Fluxus*, *Polyrrhea*, *Flux*; (*profluo*, (*pro*, and *fluo*,) 'I flow forth.') A term under which—in the plural, *Profluvia*, *Polyrrheas*—some nosologists have comprised all morbid discharges or fluxes:—others, increased excretions attended by fever.

PROFLUVIUM ALVI, *Diarrhoea*—p. *Genitale mulieb're*. *Menses*—p. *Mucosum urethrae*, *Gonorrhoea*—p. *Mulieb're*, *Leucorrhoea*. *Menses*—p. *Sanguinis*, *Hæmorrhagia*—p. *Sanguinis è renibus*, *Nephorrhagia*—p. *Sanguinis ex ore*, *Stomatorrhagia*—p. *Sanguinis ex utero*, *Metrorrhagia*—p. *Seminis*, *Pollutio*—p. *Ventris*, *Diarrhoea*.

PROFOND, *Profundus*.

PROFUNDA, fem. of *Profundus*.

PROFUNDA ARTERIA FEMORIS, *A. cruralis posterior seu muscularis femoris*, *Artère-grande musculaire de la cuisse*. (Ch.) This large branch arises commonly from the posterior part of the crural, between the pubis and lesser trochanter. It descends deeply, situate before the adductors; passes through the third adductor above the opening in it for the trunk of the femoral; and terminates in the short portion of the biceps. The *musculus profunda* gives off, 1. The *external circumflex*: 2. The *internal circumflex*: and 3. The *three perforantes*.

PROFUNDA ARTERIA HUMERI, *P. superior*, *Arteria spiritalis*, (F.) *Artère humérale profonde*, *A. collatérale externe*, *Grand musculaire du bras*, (Ch.) Its origin is variable. When it arises from the brachial artery, it is given off opposite the groove of the humerus destined for the radial nerve, and descends backwards between the three portions of the triceps, accompanied by the nerve; giving branches to that muscle and to the humerus. When it reaches the posterior part of the bone, it divides into two branches, which are distributed to the triceps, supinator longus, &c.

PROFUNDA ARTERIA INFERIOR seu MINOR is often a branch of the last; but, commonly, is given off from the brachial, near the middle of the arm. It gives off branches to the muscles, &c., about the inner side of the os humeri. It is, also, called *large communicating ulnar* or *profundoulnar*.

PROFUNDA ARTERIA PE'NIS. Chaussier has given this name to the *cat'ernous artery*, which

arises from the artery of the penis, furnished by the internal pudic.

PROFUNDA ARTERIA SUPERIOR, *Profunda humeri*.

Certain muscles are distinguished by the names *profundus* or *deepseated*, and *superficial*. Thus, we speak of the *superficial* and *deepseated* muscles of the neck, &c.; the *Flexor profundus perforans*, &c.

PROFUNDUS, (*pro*, and *fundus*, 'bottom,' 'having a deep base.') *Alvus*, (F.) *Profundus*. A name given to different parts, which are seated profoundly as regards others.

PROFUSIO, (*pro*, and *fundere*, *fundum*, 'to pour.') A pouring out, a shedding; see *Hæmorrhagia*.

PROFUSIO A'QUÆ. The discharge or breaking of the waters at birth.

PROFUSIO SANGUINIS, *Hæmorrhagia*—p. *Seminis*, *Ejaculation*—p. *Subcutanea*, *Purpura simplex*.

PROGASTOR, (*pro*, and *γαστηρ*, 'belly.') One who has a pendulous belly.

PROGENIES, (*pro*, and *gigno*, 'to bring forth,') *Epigone*.

PROGENY, *Epigone*.

PROGLOS'SIS, (*προγλωσσις*, from *pro*, and *γλωσσα*, 'the tongue;') *A'pex linguae*. The extremity or tip of the tongue.

PROGNATHOUS, *Prognathus*, (*pro*, and *γναθος*, 'the jaw.') Having a projecting jaw. A term applied to the form of the head in which there is a prolongation or forward extension of the jaws, as in the negro.

PROGNOSIS, (*προγνωσις*,) *Proëgoreusis*, *Proërhesis*, *Præcognit'io*, *Prognosis*, *Prænotio*, *Prænuntiatio*, *Præscientia*, *Præsci'tio*, *Prono'a*, *Proph'asia*, *Prædictio*, (F.) *Prognostic*, *Pronostic*, (*pro*, and *γνωσις*, 'knowledge.') A judgment formed by the physician regarding the future progress and termination of any disease.

PROGNOSIS, GEN'ERAL. The opinion formed of a disease in the abstract. Thus, the *General prognosis of Cynanche tonsillaris* is favourable;—of *Phthisis pulmonalis*, unfavourable.

PROGNOSIS, PARTIC'ULAR. The opinion formed of any particular case of disease. The *particular or special prognosis* of one case of typhus may be favourable, whilst that of another may be unfavourable.

PROGNOSIS EX LINGUÂ, *Glossomantia*.

PROGNOSTIC, *Prognosticus*, (F.) *Prognostique*, *Pronostic*. Relating or appertaining to prognosis,—as 'prognostic signs.'

PROGNOSTIC, *Prognosis*.

PROGNOSTICE, (*προγνωστικη*) (*τεχνη*,) *Prognosis*.

PROGNOSTIQUE, *Prognostic*.

PROGRESSIO, (*pro*, and *gradior*, *gressus*, 'to step,') *Augmentation*.

PROGRESSION, *Walking*.

PROGRESSUS, *Augmentation*.

PRÔTÏTES, (*πρωιotes*, or *πρωιotes*,) *Præntia*.

PROJECTURA, (*pro*, and *jacio*, *jectum*, 'to throw,') *Process*.

PROLABIUM, (*pro*, and *labium*, 'lip,') *Procheilon*.

PROLAPSUS, (*prolabor*, *prolapsus*, (*pro*, and *labor*, *lapis*, 'to slide,') 'to slip down,') *Prædent'ia*, *Delapsio*, *Delapsus*, *Hypar'rhysis*, *Proptoma*, *Propt'ia*, *Productio*, *Propendent'ia*, *Pro'minent'ia*, *Pro'sis*, *Ca'sis*, a *protrusion*; a *falling down*; (F.) *Chute*. A genus of diseases in the class *Locales* and order *Ectopie* of Cullen, distinguished by the falling down of a part through the orifice with which it is naturally connected.

PROLAPSUS ANI, *Proctocele*—p. *Bulbi oculi*, *Exophthalmia*—p. *Cornea*, *Staphyloma* or *the cornea*—p. *Iridis*, *Pto'sis iridis*—p. *Lingua*, *Glos-*

socle, Paraglossa — p. Œsophagi, Pharyngocele — p. Palpebræ, Blepharoptosis — p. Pharyngis, Pharyngocele — p. Umbilici, Exomphalus.

PROLAPSUS UTERI, *Exometra*, *Metroppto'sis*, *Metroppto'sis*, *Adopto'sis* seu *Procidencia uteri*, *Hysteropto'sis*, *Hysterocto'te nu'da*, *Falling down of the womb*, (F.) *Abaissement ou Chute ou Descente ou Précipitation de la matrice*. Some use the term 'Procidencia Uteri' for a less degree of the affection than prolapsus; — *Relaxatio uteri* expressing the slightest state of all. A falling down of the uterus, owing to relaxation of the parts about the uterovaginal region. In the treatment, the horizontal posture must be insisted upon; the injection of astringent substances *per vaginam* be advised; and if the affection be not remedied by these means, a pessary may be employed as a palliative, or the operation of *Episiorrhaphy* be performed.

Prolapsus uteri without inversion has been termed *Orthysteroto'ma*, and *Orthysteropto'sis*.

PROLAPSUS UVULÆ, Staphylœdema.

PROLAPSUS VAGINÆ, *Hysteropto'sis* seu *Adopto'sis* *Vaginæ*, *Coleopto'sis*, *Colpopto'sis*, *Elytrop'to'sis*. Protrusion of the upper part of the vagina into the lower. This, like the descent of the uterus, may be *relaxation*, *procidencia*, *prolapsus*, or complete *inversion*.

PROLAPSUS VESICÆ, *Exocyste*.

PROLECTA'TIO; (*prolecture*, *prolectatum*, 'to entice,') Action of separating the finer parts of a body from the grosser.

PROLEPTICS, *Prolepticiæ*, (*προληψις*, 'anticipation,' from *pro*, and *λαμβάνω*, 'I seize hold of,') A term proposed by Dr. Laycock to signify the art and science of predicting in medicine.

PROLEPTICUS, (*προληπτικός*,) Anticipating.

PROLES, (*pro*, and *olere*, 'to grow,') Epigone.

PROLICYDE, *Prolucidum*, (*proles*, 'offspring,' and *cedere*, 'to kill,') Destruction of offspring: a term which includes feticide as well as infanticide.

PROLIFERATION OF CELLS, (*proles*, 'offspring,' and *fero*, 'I carry,') see Cytogeny.

PROLIFIC, *Prolificus*, (F.) *Prolifique*, (*proles*, 'offspring,' and *facere*, 'to make,') That which has the faculty of engendering. Applied to man, and animals, and to their sperm or seed; as well as to remedies, which, by strengthening the genital organs, increase the secretion of sperm.

PROLIG'EROUS, *Pro'liger*, *Prolig'erus*, (F.) *Proligère*, (*proles*, 'offspring,' and *gero*, 'I carry,') That which is connected with carrying the offspring.

PROLIGEROUS DISC or **LAY'ER**, *Discus prolig'erus* seu *vittell'inus*, *Stratum prolig'erum*, *Gonostro'ma*, (F.) *Disque proligère*. A granular layer, situate generally towards the most prominent part of the ovarian vesicle, in the centre of which the true ovum or ovule exists.

PROLONGEMENT RACHIDIEN, (*pro*, and *longus*, 'long,') *Medulla spinalis* — p. *Sous-occipital*, Basillary process.

PROMALACTERIUM, (*pro*, and *μαλασσω*, 'I soften,') The room, in the ancient gymnasium, in which the body was softened and anointed. One of the operations in bathing.

PROMANUS, (*pro*, and *manus*, 'hand,') see *Digitus*, *Pollex*.

PROMETOPIDIA, (*προμετωπίδα*,) Frontal bandages.

PROMETO'PIS, *Prometopid'ion*, (*pro*, and *μετωπον*, 'the forehead,') The skin of the forehead, in which wrinkles are.

PROMINENTIA, (*pro*, and *mineo*, 'to jut,') Prolapsus, Protuberance — p. *Annularis cerebri*, *Pons Varolii* — p. *Corneæ*, *Ceratocoele* — p. *Ossis continua*, *Apophysis*.

PROMINENTIE ALBICANTES, *Mamæillary tubercles* — p. *Semiövales Medullæ Oblongatæ*, *Corpora olivaria*.

PROMONTOIRE, *Promontorium*.

PROMONTORIUM, *Promontory* — p. *Faciei*, *Nasus* — p. *Ossis sacri*, see *Sacrum*.

PROMONTORY, *Promontorium*, (F.) *Promontoire*, (*pro*, and *mons*, gen. *montis*, 'a mountain,') *Tu'ber seu Tuberos'itus Tym'pani*. A small projection at the inner part of the cavity of the tympanum, which corresponds to the external scala of the cochlea, and especially to the outer side of the vestibule.

PROMONTORY OF THE SACRUM, see *Sacrum*.

PROMONATEUR, *Pronator* — p. *Carré*, *Pronator radii quadratus* — p. *Grand ou rond*, *Pronator radii teres* — p. *Petit*, *Pronator radii quadratus*.

PRONA'TION, *Prona'tio*, (*pronus*, 'inclined forwards,') Anatomists understand, by pronation, the motion by which the inferior extremity of the radius passes before the ulna, and thus causes the hand to execute a kind of rotation from without inwards.

PRONA'TOR. That which produces the motion of pronation. (F.) *Pronateur*. This name has been given to two of the muscles of the forearm.

PRONATOR OBLIQUUS, *P. radii teres* — p. *Quadratus*, *P. radii quadratus*.

PRONATOR RA'DII QUADRA'TUS, *P. quadratus* seu *transvers'us* (Winslow) seu *ra'dii bre'vis* seu *ra'dii quadratus* (Cowper), *Quadratus ra'dii*, (F.) *Cubito-radial* (Ch.), *Petit ou carré pronateur*. This muscle is situate at the anterior, inferior, and profound part of the forearm. It is flat and square, and is attached, within, to the inferior quarter of the anterior surface of the ulna; and, without, to the inferior quarter of the anterior surface of the radius. It produces the motion of pronation.

PRONATOR RA'DII TE'RES, *P. te'res* sive *obli'quus* (Winslow), (F.) *Epitrochlo-radial* (Ch.), *Grand ou rond pronateur*. A muscle seated at the upper and anterior part of the forearm. It is long, flat, and larger above than below. It is attached, above, to the inner tuberosity of the humerus and to the coronoid process of the ulna. Thence it passes obliquely downwards and outwards, and terminates at the middle of the outer surface of the radius. It causes the radius to turn on the ulna, to produce the motion of pronation. It can, also, bend the forearm on the arm, and conversely.

PRONATOR TERES, *P. radii teres*.

PRONAUS, (*pro*, and *ναος*, 'a temple,') *Vestibulum*.

PRONERVATIO, (*pro*, and *nervus*, 'a tendon,') *Aponeurosis*, *Tendon*.

PRONŒA, (*προνοια*, from *pro*, and *νοω*, 'to know,') *Prognosis*.

PRONONCIATION, *Pronunciation*.

PRONOSTIC, *Prognosis*.

PRONOSTIQUE, *Prognostic*.

PRONUNCIATION, *Pronuncia'tio*, (F.) *Pronunciation*, (*pro*, and *nunciare*, 'to tell,') The mode of articulating or giving accent or utterance to words.

PROPATHIA, (*προπαθεια*, from *pro*, and *παθος*, 'affection,') *Prodromus*.

PROPENDENTIA, (*pro*, and *pendere*, 'to hang,') *Prolapsus*.

PROPERTIES, PHYSICAL, OF THE TISSUES. These are *flexibility*, *extensibility*, *elasticity*, *racornissement*, *imbibition* and *osmose*.

PROPERTIES, VITAL. Those which depend upon the vital force, as *contractility*.

PROPERTY, *Proprietas*, *Virtus*, (F.) *Propr'ieté*, (*proprius*, 'proper,') That which belongs distinctively to a thing.

PROPHASIS, (*προφασις*, from *προφαίνω*, 'to su-

source.' A cause, especially a remote cause. Also, *Prognosis*.

PROPHYLACTIC, *Prophylacticus*, *Diaphylacticus*, *Diastetic*, *Synsteticus*, *Preservativus*, (Fr.) *Prophylactique*, *Preservatif*, (*pro*, and *phylassein*, 'I defend.') A preservative or preventive.

PROPHYLACTIC MEDICINE, Hygiene.

PROPHYLLAXIS, *Prophyllaxis*, *Prophyllacticē*, *Synstaxis*, *Preservatio*, *Preservatio*, *Preservatio*, *Preservatio*. Preservative or preventive treatment.

PROPOLIS, (*pro*, and *polis*, 'city,' literally, 'that which is before the city.') *Ceraulthemus*, *Ceraulthemus*, *Beebread*. A red, resinous matter, with which the bees cover the bottom of the hive. Its fume has been esteemed antiasthmatic.

PROPOMA, gen. *Propom'atis*, (*pro*, and *oma*, 'a drink.') Ancient name of a medicine composed of seven parts of honey and four of wine. It was taken before a meal.

PROPORTIO, (*pro*, and *partio*,) Symmetry.

PROPOSITUM, (*pro*, and *ponere*, *positum*, 'to place,') Intention.

PROPTISMA, gen. *Proptism'atis*, (*proptis*,) Mixture.

PROPTISMUS, (*proptis*, from *pro*, and *proptis*, 'to give to drink,') *Propotio*. An ancient name for medicines which were given to the patients before purging them, and, in some sort, to prepare them.

PROPRIETAS, Property.

PROPRIÉTÉ, Property.

PROPRIUS AURIS EXTERNÆ, *Retrahens auris*.

PROPTOMA, gen. *Propto'matis*, (*pro*, and *ptoma*, 'to fall,') *Prolapsus*.

PROPTOMA AURICULÆ, *Parasthe'sis auriculae flac'ca*, *Flap Ear*. Lobe of the ear broad, loose, and pendent from birth.

PROPTOMA SCOTI, *Rachosis*.

PROPTUSIS, (*proptus*, *pro*, and *ptusis*, 'a falling down,') *Prolapsus*—p. *Palpebræ*, *Blepharoptosis*—p. *Utriculi*, *Staphylodema*.

PROPTYSIS, (*pro*, and *ptus*, 'to spit,') Expectoration.

PROPYLAMIN, *Propylaminum*, occurs in *chenopodium vulvaria*, in the flowers of *crataegus oxyacantha*; the fruit of *serotina aucuparia*; in *sessile cornutum*, herring pickle, codliver oil, urine, &c. (It may be prepared from herring pickle, by distillation with caustic potassa, neutralizing the distillate with muriatic acid; purifying the salt by dissolving it in strong alcohol or alcoholic ether, and again distilling with potassa.) It has been used in acute and chronic rheumatism, 25 drops being added to six ounces of mint water sweetened; and a tablespoonful given every two hours. Its virtues have been greatly exaggerated. The chloride and iodide have been given in like cases.

PRORA, Occiput.

PRORRHESIS, (*prorrhesis*,) *Prognosis*.

PRORUPTIO SANGUINIS, (*pro*, and *rumpere*, *rumpere*, 'to break,') *Hæmorrhagia*.

PROS, (*pros*,) in composition, generally signifies 'in addition to,' 'over and above,' 'besides.'

PROSARTHROSIS, (*pros*, and *arthron*, 'to attach by joints,') *Diarthrosis*.

PROSBOLÆ, (*pros*, and *ballō*, 'to strike,') *Impression*.

PROSCHYSIS, (*pros*, and *schō*, 'to pour,') *Affusion*.

PROSCOLLESIS, (*proscollēsis*, from *pros*, and *colla*, 'glue,') *Adherence*, *Agglutination*.

PROSECTEUR, *Dissector*.

PROSECTIO, (*pros*, and *secare*, *sectum*, 'to cut,') *Anatomy*.

PROSECTOR, *Dissector*.

PROSLEPSIS, (*proslēsis*, from *pros*, and *lambanō*, 'to seize hold of,') *Prehension*.

PROSODOS, (*prosōdos*, from *pros*, and *ōdos*, 'a way,') *Adjutus*.

PROSOPALGIA, (*prosōpon*, and *algos*, 'pain,') *Neuralgia facialis*.

PROSOPANTRA, (*prosōpon*, and *antron*, 'a cavity,') *Frontal sinuses*.

PROSOPANTREITIS, (*prosōpantra*, and *itis*,) *Inflammation of the frontal sinuses*.

PROSOPIS DULCIS, *Ord. Leguminosæ*. A tree of rather large size, native of Peru. The fruit—*Pacay*—is a pod, from 20 to 24 inches long, containing black seeds imbedded in a white, soft, flaky substance; which last is eaten by the Limes with pleasure.

PROSOPODYNIA, (*prosōpon*, and *odynē*, 'pain,') *Neuralgia facialis*.

PROSOPODYSMORPHIA, (*prosōpon*, *δys*, 'with difficulty, and *morpha*, 'shape,') *Malformation or atrophy of the face*,—as where the process of nutrition suffers from diminished nervous influence.

PROSOPOLOGIA, (*prosōpon*, and *logos*, 'a discourse,') *Physiognomy*.

PROSOPOMANTIA, (*prosōpon*, and *μαντεια*, 'divination,') *Physiognomy*.

PROSOPON, (*prosōpon*,) *Face*.

PROSOPOPARALYSIS, (*prosōpon*, and *paralysis*,) *Palsy*, *Bell's*.

PROSOPORRHEUM'IA, gen. *Prosoporrheum'atis*, *Rheumatismus facialis*, (*prosōpon*, and *ῥευμα*, 'defluxion,') *rheumatism*,) *Rheumatism affecting the face*.

PROSOPOSIS, (*prosōpon*,) *Physiognomy*.

PROSOPOSPASMUS, (*prosōpon*, and *spasmus*,) *Canine laugh*.

PROSOPOSTERNODYMIA, (*prosōpon*, *σθηνον*, 'the sternum, and *dynē*, 'to mingle,') see *Cephalosomatodymia*.

PROSPERITAS VALETUDINIS, (*prosper*, 'favourable,') *Sanitas*.

PROSPHORA, (*prospora*, from *pros*, and *phō*, 'to bring,') *Prehension*.

PROSPHYSIS, (*prosphe'sis*,) *Adhæscen'tia*, *Coalitio*, 'adhesion, adherence, connexion,' (from *pros*, and *phō*, 'I grow,') In a more limited sense, this word means morbid adhesion of the eyelids, either between themselves, or with the globe of the eye: *Ankyloblepharon*.

PROSTASIS, (*prostasis*, from *pro*, and *stasis*, 'station,') *'superiority, 'preponderance.'* A pre-dominance or abundance—as of excrementitious humours.

PROSTATA, *Prostate*—p. *Adenoides*, *Prostate*—p. *Bartholini*, *Cowper's glands in the female*—p. *Glandula*, *Prostate*—p. *Muliebris*, *Cowper's glands in the female*.

PROSTATÆ, *Prostate*—p. *Adenoides*, *Prostate*—p. *Glandulose*, *Prostate*.

PROSTATAL, *Prostatic*.

PROSTATAL'GIA, (*prostate*, and *algos*, 'pain,') *Pain in the prostate*.

PROSTATAUXE, (*prostate*, and *auxē*, 'increase,') *Prostatoparæctasis*.

PROSTATĒ, (*pro*, and *στημι*, 'I stand,') *Prostatata*, *P. glandula*, *Prostatata*, *P. glandulosa seu adenoides*, *Paras'tata adenoides seu glandulosa*, *Ad'ens*, *Paras'tata*, *P. glandula*, *Tes'tis m'ina*, *Cor'pus glandulosa seu glandulosa seu adeniforme seu glanduliforme seu glandulosa seu adenoides*, *Assen'tor glandula*, *Ad'itus glandulosa*; the *Prostate gland*. A glandular, cordiform body, of the size of a chestnut, situate before the neck of the bladder, behind the symphysis pubis, and surrounding the first portion of the urethra. Its excretory outlets, the number of 10 or 12, open into the part of the urethra that traverses it, and pour into it a whitish viscid humour—*Liquor*

prostat'icus—intended to lubricate the interior of the urethra, and to serve as a vehicle for the sperm in its ejaculation.

PROSTATE, ANTERIOR, see Cowper's glands.

PROSTATELCO'SIS, *Prostatelco'sis*, (*prostate*, and *'elcos*, 'an ulcer.') Ulceration of the prostate.

PROSTATES INFÉRIEURES, Cowper's glands—*p. Petites*, Cowper's glands.

PROSTATHELCOSIS, (*prostate*, and *'elcos*, 'an ulcer.') Prostatelcosis.

PROSTAT'IC, *Prostat'icus*, *Prostat'ical*, (F.) *Prostatique*. That which belongs or relates to the prostate. Winslow has named—*Superior Prostatic Muscles*—ligaments which pass from the pubis to the lateral parts of the prostate. He, also, calls—*Inferior prostatic muscles*—the fleshy fibres, which, from the lateral parts of the urethra, near its membranous portion, pass to be inserted into the pubis.

PROSTATIC BISECTOR. An instrument employed by Dr. Stevens, of New York, in the bilateral operation of lithotomy. In form it resembles a large olive, with a beak at the extremity, cutting edges at the sides parallel to its longest axis, and with a straight handle.

PROSTATIC LIQ'UOR, *Li'quor prostat'icus*. The fluid secreted by the prostate, which mixes with sperm during emission.

PROSTATIC PORTION OF THE URETHRA. The part of the canal of the urethra included by the prostate: it is about 15 lines long.

PROSTATIC SINUS, see Sinus, prostatic.

PROSTATICUS SUPERIOR, Compressor prostate.

PROSTATIQUE, Prostatic.

PROSTATITIS, *Inflamma'tio Prostat'icæ*, (*prostate*, and *itis*.) Inflammation of the prostate; when accompanying gonorrhœa, it is the *gonorrhœo-prostat'itis*.

PROSTATOCELE, (*prostate*, and *κηλη*, 'tumour,') Prostatoncus.

PROSTATOLITHI, (*prostate*, and *λιθος*, 'a stone,') see Calculi, urethral.

PROSTATON'CUS, *Prostatocœ'lē*, *Tumor prostate*, (*prostate*, and *o'yxos*, 'a tumour,') Swelling of the prostate.

PROSTATOPAREC'TASIS, (*prostate*, and *παρεκτείνω*, 'to extend,') Enlargement of the prostate; also, induration of the same, *Prostatoscir'rius*, *Prostatoscir'e*.

PROSTATORRHŒA, (F.) *Prostatorrhée*, (*prostate*, and *ῥέω*, 'I flow,') A morbidly increased discharge from the prostate.

PROSTATOSCIRRIUS, see Prostatoparectasis.

PROSTERNID'IUM, (*προστερνιδιον*, from *pro*, and *σενον*, 'the sternum,') Any agent applied to the anterior part of the chest; a plaster to the breast.

PROSTHESIS, (*προσθεσις*,) *Adjunct'io*, *Adjec'tio*, *Appos'itio*, 'addition, application;' (from *pros*, and *τιθέναι*, 'to put,') That part of surgery, whose object is to add to the human body some artificial part, in place of one that may be wanting, as a wooden leg, an artificial eye, &c. Sometimes written *Prothesis*, (F.) *Prothèse*.

PROSTHETA, (*προσθετα*,) Subdita.

PROSTHIUM, (*προσθιον*,) Penis.

PROSTITUTION, *Prostitu'tio*, *Porni'a*, (from *prosto*, (*pro*, 'before,' and *sto*, 'I stand,') 'I stand to be hired,') The condition and practice of a prostitute, (F.) *Fille publique*, *F. de Joie*. In many countries of Europe, prostitution is a matter of public hygiene—the prostitutes being regularly examined medically and licensed. In Great Britain, and in this country, it is not subjected to any regulation.

PROSTOM'IA, *Commissu'ra labio'rum*; (*pro*, and *στομα*, 'mouth,') The commissure of the lips.

PROSTRA TION, *Prostra'tio vi'rium*; (*pro*,

and *sternere*, *stratum*, 'to throw down,') (F.) *Prostration des forces*, *Abattement*. Great depression of strength. Almost total loss of power over the muscles of locomotion.

PROTEA MELLIFERA, (after Proteus.) A South African plant, *Nat. Ord.* Proteaceæ, whose involucre and flowers, at the time of inflorescence, are filled with a sweet watery liquid, which contains a great deal of honey. By inspissation it forms a delicious syrup—*Syrupus Pro'teæ*, which is much used in pulmonary affections.

PROTEA LEPIDOCARPON, and some other Proteæ, supply the same kind of juice.

PROTEAN or PROTE'AN, *Pro'teiform*, *Pro'teiform'is*, (from *Proteus*, who could assume various shapes.) Assuming different shades.

PROTEIC, Proteinous.

PROTEIFORMIS, Protean.

PROTEIFORM'IS MORBUS. A disease which assumes various characters.

PROTEÏN, *Pro'teine*, *Protei'na*, (from *πρω-τινω*, 'I take first rank,') A product of the decomposition of albumen, &c., by potassa. When, according to Mulder, animal albumen, fibrin or casein is dissolved in a moderately strong solution of caustic potassa, and the solution is exposed for some time to a high temperature, these substances are decomposed. The addition of acetic acid causes, in all three, the separation of a gelatinous translucent precipitate which has exactly the same characters and composition. This is protein. The existence of any such radical free from sulphur is denied.

Protein having been regarded as the basis of animal and vegetable fibrin, albumen, casein, and gluten, they have been classed under the "proteinaceous alimentary principle," by Dr. Pereira.

PROTEINACEOUS, Proteinous.

PROTEÏNOUS, *Proteinæ'ous*, *Proteinô'sus*, *Pro'teic*, (F.) *Protéineux*, *Protéique*. Of, or belonging to protein;—as "a proteinous alimentary principle." For proteinous, albuminous is now generally substituted.

PROTEÏNEUX, Proteinous.

PROTEÏQUE, Proteinous.

PROTENSIONES GLANDULARES, (*pro*, and *tendere*, *tensum*, 'to stretch,') Mammillary eminences.

PROTHÈSE, Prosthesis.

PROTHESIS, Prosthesis.

PROTMESIS, (*προτμησις*, from *pro*, and *τεμνω*, 'to cut,') Umbilicus.

PROTO, (*πρωτος*, 'first,') in composition, first.

PROTOAZOTATE DE MERCURE, Hydrargyri nitras.

PROTOGALA, (*proto*, and *γαλα*, 'milk,') Colostrum.

PROTOIATROS, (*proto*, and *ιατρος*, 'physician,') Archiater.

PROTOMEDICUS, Archiater.

PROTOMUS, (*pro*, and *τεμνω*, 'to cut,') Dissector.

PROTONEPHRIUM, (*proto*, and *νεφρος*, 'a kidney,') Corpus Wolfianum.

PROTOORGANISM, *Protoörganism'us*. A being, whose animal or vegetable nature is so difficult of determination, that it might be referred to either one or the other division of organized bodies. See Zoophyte.

PROTOPATHIA, (*proto*, and *παθος*, 'affection,') Idiopathia.

PROTOPHYTE, *Protophy'ton*, (*proto*, and *φυτον*, 'a plant,') A vegetable production at the foot of the scale; as a cryptogamous plant,—a fungus.

PRO'TOPLASM, *Protoplas'ma*, (*proto*, and *πλασσω*, 'to form,') The blastema in the sense of cyblastema. Also, according to Robin, the liquid contained in the cavities of vegetable cells,

or in the embryonic cells, when the embryo has as yet no blood; which liquid is capable, like the liquor sanguinis, of furnishing materials for the origin of other anatomical elements.

PRO'TOPLAST, (*proto*, and πλασσω, 'to form.')

A primary formation. A first formed organized body—the parent of all others.

PROTOSPOROS, (*proto*, and σπειρω, 'to sow,') *Os uteri*.

PROTOZOIDS, (*proto*, ζωον, 'an animal,' and ζωος, 'resemblance,') *Spermatozoa*.

PROTOZO'ON, (F.) *Protozoaire*, (*proto*, and ζωον, 'an animal,') An animal lowest in the scale; —as a monad.

PROTRAHENS LENTIS, (*pro*, and *trahere*, 'to draw,') Ciliary muscle.

PROTRUSION, (*pro*, and *trudere*, *trusum*, 'to thrust,') *Prolapsus*.

PROTRUSOR, *Detrusor urinæ*.

PROTUBERANCE, *Protuberantia*, *Extuberantia*, *Prominentia*, *Ecrescentia*, *Extuberantia*, (*pro*, 'before,' and *tuber*, 'a projection,') *Bosse*, (F.) *Bosse*. A name given to rough, unequal protuberances, seen at the surface of certain organs; the *parietal protuberances*, *occipital protuberances*, *angular protuberances*, &c.

PROTUBERANCE, **ANNULAR**, *Pons Varolii* — *p. Cerebral*, *Medulla oblongata*, *Pons Varolii* — *p. Cylindroide*, *Cornu ammonis*.

PROTUBERANTIA, *Eminence*, *Protuberance* — *p. Annularis Willisii*, *Pons Varolii* — *p. Basilaris*, *Pons Varolii* — *p. Cylindrica*, *Cornu ammonis*.

PROUD FLESH, *Fungosity*.

PROUNE, *Prunus domestica*.

PROVENTRIC'ULUS, *Bulbus glandulosus*, *Infundibulum*, *Sec'ond stomach*, (*pro*, and *ventriculus*, 'the stomach,') A bulbous expansion at the termination of the oesophagus, immediately above the gizzard of birds, the walls of which are thickly studded with a layer of glands or follicles which secrete a digestive fluid.

PROVERB, **MED'ICAL**, *Proverbium medicum*. A proverbial expression frequently contains sound sense in its directions. In medicine, the greatest collection is in the *Regimen of Health of the School of Salerno*, composed in the 11th century, by John of Milan. Yet, although good sense is often inculcated, we frequently meet with the reverse.

Most of the proverbs are hygienic. The following are a few.

1. *Qu' après la soupe un coupe d'excellent vin
Tire un écu de la poche du médecin.*
2. *Surge quintâ, prande nonâ, cæna quintâ, dormi
nonâ, nec est mortî vita prona.*
3. *Nunquam recte corpus exerceri sine animo,
neque animum sine corpore posse.*
4. *Viande bien machée est à demi digérée.*
5. *Optimum condimentum fames.*
Appetite is the best sauce.
6. *Plures occidit gula quam gladius.*
7. Early to bed and early to rise,
Makes a man healthy, wealthy, and wise.
8. One man's meat's another man's poison.
9. An ounce of prevention is better than a pound
of cure.

PROVERSIOS, (*pro*, and *vertere*, *versum*, 'to turn,') *Anteversion*.

PROVINS, **MIN'ERAL WA'TERS OF**. Chalybeate springs in the department of Seine et Marne, France, which contain carbonic acid and iron.

PROVOCATORIUS, (*pro*, and *vocare*, *vocatum*, 'to call,') *Intercalary*.

PROX'IMAD, see *Proximal aspect*.

PROX'IMAL, (*proximus*, 'next,') *Proximate*; nearest; next.

PROXIMAL AS'PECT. An aspect towards the trunk, in the course of an extremity.—*Barclay*. *Prox'imal* is used by the same writer adverbially to signify 'towards the trunk.'

PRUNA, *Anthrax*.

PRUNE, *Prunum*.

PRUNE JUICE EXPECTORA'TION or **Spu'ta**. The sputa resembling prune juice, which occur in the third or purulent stage of pneumonia.

PRUNEAU, *Prunum*.

PRUNELLA, (*dim. of prunum*,) *Angina pectoris*, *Aphthæ*, *Cynanche*, (all from [G.] *Braune*, 'sorethroat,') *Pupil*.

PRUNEL'LA, *P. vulgaris seu officinalis*, *Brunella*, *Consolida mi'nor*, *Symphytum mi'nus*, *Self'heal*, *Healall*, *Bu'gle*, (F.) *Prunelle*, *Brunelle*; *Ord. Labiate*. *Sex. Syst. Didynamia Gymnospermia*. This plant has been recommended as an astringent, in hemorrhages and fluxes; and, also, in gargles against aphthæ, and in inflammation of the fauces; hence its name from ([G.] *Braune*, 'sorethroat,')

PRUNELLÆ SAL, *Potassæ nitras fusus sulphatis pauillo mixtus*.

PRUNELLE, *Prunella*, *Pupil*.

PRUNELLIER, (*dim. of prunus*,) *Prunus spinosa*.

PRUNELLOE, *Prunum Brignolense*.

PRUNEOLA SYLVESTRIS, *Prunus spinosa*.

PRUNES, see *Prunum* — *p. Pulp of, Pruni Pulpæ*.

PRU'NI PUL'PA, *Pulp of prunes*. (Take of *prunes* a sufficient quantity, soften them in the vapour of boiling water, and having separated the stones, beat the remainder in a marble mortar and press through a hair-sieve.—*Ph. U. S.*)

PRUNER, *Prunus domestica* — *p. Sauvage*, *Prunus spinosa*.

PRU'NUM, *Coccyne'lon*. A dried *Plum*, a *Prune*. (F.) *Pruneau*. Three sorts of *Prunes* have been usually ranked amongst the articles of the *matéria medica*. 1. The *Prunum Brignolensê*; *Brignole plum* or *Prunelloe*; from *Brignole* in *Provence*. It is of a reddish-yellow colour; and has a very grateful, sweet, subacid taste. 2. The *Prunum Gal'licum*, *Prunum* (*Ph. U. S.*), the *French Prune*; and 3. The *Prunum Damascenum*, *Dam'son*, formerly *Dam'ascene*, *Brab'ylon*. All these possess the same general qualities as the other summer fruits. They are emollient and laxative.

PRUNUM STELLATUM, *Averrhoa carambola*.

PRUNUS ACACIA, (προυν,) *Prunus padus*.

PRU'NUS ARME'NICA, *Armeni'aca vulgaris seu epirotica*. The *A'pricot*, *A'pricock*. *Bericoc'cæ*, *Præcoc'ia*, (F.) *Abricotier*. *Ord. Rosacæ*. *Sex. Syst. Icosandria Monogynia*. The *apricot*, *Me'lum Armeniacum*, (F.) *Abricot*, when ripe, is easily digested, and considered as a great delicacy.

PRUNUS A'VIUM, *P. cer'asus seu sylvest'ris seu macrophyl'la*, *Cer'asus A'vium seu ni'gra seu dul'cis*, *Black Cher'ry Tree*, (F.) *Merisier*. The fruit is eaten; and a gum exudes from the tree, the properties of which are similar to those of gum Arabic.

PRUNUS CAPULIN, *Capuli*, grows in the open fields of Peru; and is cultivated in gardens in the towns. The fruit is acid, and not often eaten; but on account of its agreeable odour it is used in making *Pucheros de flores*; or with odoriferous flowers to perfume linen.

PRUNUS CER'ASUS, *Cer'asus ac'ida seu hortens'is seu vulgaris seu rubra*. The *Red Cher'ry Tree*; (F.) *Cerisier*. The fruit *Cer'asum*, *Cera'sion*, has a pleasant, acidulous sweet flavour. It is wholesome. See *Prunus avium*.

PRUNUS DOMESTICA. The *Plum* or *Dam'son tree*, *Coccyus'lea*, *Prunae*, (F.) *Pruvier ordinaire*. The damson, when perfectly ripe, affords a wholesome fruit for pies, tarts, &c. It is gently laxative. See *Prunum*.

PRUNUS HORTENSIS, *P. cerasus*.

PRUNUS INSIT'IA. The *Bull'ace Plum Tree*. The fruit of this tree is used like the damson.

PRUNUS LAURO-CERASUS, *Cerasus lauro-cerasus*, *Pa'dus laurocerasus*, *Po'son Lau'rel*, *Cherry Laurel*, *Common Laurel*, *Laurocerasus*, (F.) *Laurier-cerise*, *L. amandier*, *L. de Trébisonde*. The leaves, *Laurocerasus*, (Ph. E., D., and B.), have a bitter, styptic taste, with the flavour of bitter almonds. The flowers have a similar flavour. The powdered leaves, applied to the nostrils, excite sneezing. The flavour of the leaves has given occasion to their being employed in tarts, custards, &c.; but in large quantities they are poisonous. The poisonous principle is the prussic acid. The distilled water, called *Laurelwater*, is poisonous. (See Poisons, Table of.) An *Aqua LAURO-CERASI*, *Cherrylaurel water*, is official in the Edinburgh, Dublin and British (1864) Pharmacopœias. Dose, ℞ xxx to ℥j.

PRUNUS MACROPHYLLA, *P. avium*.

PRUNUS PA'DUS, *Pa'dus*, *P. avium*, *Cerasus Pa'dus* seu *racemosa sylvestris*, *Wild cluster* or *Bird Cherry Tree*, (Sc.) *Hag'berry*, *Hack'berry*, (Prov.) *Heck'berry*, (F.) *Cerisier à grappes*, *Bois puant*. The bark of the tree has a fragrant smell and a bitter, subastringent taste, somewhat similar to that of bitter almonds. It has been recommended in intermittents, &c., in the form of decoction. The berries have been used in dysentery.

PRUNUS SEBESTENA, *Sebestena*.

PRUNUS SPINOSA, The *Sloe Tree*, *P. Sylvestris* seu *Aca'cia*, *Agriococcine'lea*, *Agru'na*, *Agru'ne'la*, (F.) *Prunellier*, *Pruvier sauvage*. The fruit—the *sloe*—is sometimes employed in gargles, and was formerly much used in hemorrhage, owing to its astringent properties. The *Aca'cia nostras* seu *German'ica* seu *vulga'ris*, *Prune'ola sylvestris*, *Succus acaciæ nostrat'is*, *S. Acaciæ Germanicæ inspissat'us*, is obtained from this.

PRUNUS SYLVESTRIS, *P. Avium*, *P. Spinosa*.

PRUNUS VIRGINIANA, *Cerasus acro'tina* seu *Virginia'na*, *Choke Cherry*, *Wild Cherry Tree*, (F.) *Cerisier de Virginie*. The bark of this tree has been found useful in intermittents. The leaves are poisonous to certain animals, and the berries intoxicate different kinds of birds. The Indians use the bark in the cure of syphilis. It is bitter and astringent, and possesses some aromatic warmth, and likewise a sedative quality. It is stimulant and tonic. The leaves contain prussic acid. The distilled oil of the bark is very nearly identical with that of the bitter almond. A strong decoction of the bark is anthelmintic.

PRUNUS VULGARIS, *P. Padus*.

PRURIG'INOUS, *Pruriginosus*, (F.) *Prurigneux*, (from *prurigo*.) Relating, or belonging to, or resembling *prurigo*.

PRURI'GO, gen. *Prurig'inie*, *Pruri'tus*, *Chris'ma*, (prurio, prurimum, 'to itch.') The word is often used synonymously with *itching*. Willan and Bateman use the term for a genus of cutaneous diseases, the characteristic symptoms of which are a severe itching, accompanied by an eruption of papulæ of nearly the same colour as the adjoining cuticle; *Ones'mos*, *Scab'ies populi-form'is*, *Eror'mia pruri'go*. It affects the whole surface of the skin, under three varieties of form—the *P. mil'is*, *P. form'icans*, *P. sen'ilis*, &c., as well as some parts of the body, as the *P. pod'icis*, and *P. pud'en'di mul'i'eb'ris*, *Puden'dagra pruri'ens*, *Pruri'tus vulvæ*, (F.) *Prurit' de la vulve*.

PRURIGO, Heat—*p. Pedicularis*, *Phthieriasis*.

PRURIT, Itching—*p. de la Vulve*, see *Prurigo*.

PRURITUS, (*prurio*, 'to itch.') Gargalus, Heat, Itching, Prurigo—*p. Vulvæ*, see *Prurigo*.

PRUSSIAN BLUE, *Fer'rum Zoat'icum* seu *Borus'sicum* seu *Cyanogenat'um* seu *oxigulat'um hydrocyan'icum*, *Fer'ri Ferrocy'anus* seu *Ferrocy'anuret'um* seu *Ferrocyan'idum* (Ph. U. S. 1863) seu *Cyanuret'um* seu *Percyan'idum*, *Borus'sias* seu *Prus'sias* seu *Hydrocy'anus Fer'ri*, *Ceru'leum Borus'sicum* seu *Berolinens'e*, *Tritohydroferrocy'ane*, *Ferrocyanide*, *Ferrocyanuret* or *Ferroprussiate* of iron, (F.) *Bleu de Prusse*, *B. de Berlin*. *Ferri Ferrocyan'idum*, *F. Ferrocy'anuret'um*, *Pure Prussian Blue*, is directed, in the Ph. U. S., to be made by adding a solution of *ferrocyanide of potassium* to a solution of *ter sulphate of iron*, filtering the liquid, and washing the precipitate until the washings pass nearly tasteless, and drying it. This salt is chiefly used in the preparation of the hydrocyanic acid and the cyanide of mercury. It has been advised in the treatment of intermittents, and in epilepsy and scrophulosis. Externally, it has been applied to ill-conditioned ulcers. (℥i to ℥j of cerate.) Dose, four to six grains.

PSALIDIUM, (ψαλίδιον, dim. of *psalis*.) Fornix.

PSALIS, (ψαλις.) Fornix, Scissors.

PSALTERIUM, (ψαλτηριον, dim. of ψαλτηρ, 'a psalter,') Lyra, Omasum.

PSAMMA, (ψαμμη.) see *Gravel*.

PSAMMIS'MUS, (ψαμμιςμος.) *Ammis'mus*, *Gammis'mus*, (from ψαμμος, 'sand,') *Arenat'io*. The application of sand to any part of the body for the cure of disease.

PSAMMODES, (ψαμμοδης, from ψαμμος, and odes.) Sabulous.

PSAMMUS, (ψαμμος.) see *Gravel*.

PSAUTIER, (from (L.) *psalterium*.) Omasum.

PSELLIS'MUS, (ψελλισμος.) *Psell'otes*, (from ψελλω, 'I stutter,') Imperfect or depraved state of the articulation. A genus in the class *Pneumatica*, order *Phonica*, of Good.

PSELLISMUS NASITAS, *Rhinophonia*—*p. Rho tæcismus*, *Rotacismus*.

PSELLOTES, (ψελλωτης.) *Balbuties*, *Psellismus*.

PSEUDACACIA ODORATA, (*pseudo*, and *acacia*.) *Robinia pseudacacia*.

PSEUDAC'OË, *Pseudacōia*, *Pseudecoia*, (*pseudo*, and *akōn*, 'audition,') *Pseudacous'is*. False hearing.

PSEUDACORUS, (*pseudo*, and *acorus*.) *Iris pseudacorus*.

PSEUDÆSTHÉS'IA, *Sen'sus pover'sus*, (*pseudo*, and *αἰσθάνομαι*, 'I feel, αἰσθesis, 'feeling,') *Parapsis illu'soria*, *Pseud'aphæ*, *Pseudaph'ia*, (F.) *Pseudæsthésie*. Depraved feeling. Imaginary sense of touch or general feeling in organs that have no existence. Common to those who have suffered amputation, and to hypochondriacs and the insane.

PSEUDALEI MORBI, (Ψευδαλει, 'counterfeit,') from ψεύδης, 'false,') Feigned diseases.

PSEUDALEUS TRICHINA, *Trichina spiralis*.

PSEUDANGUSTURINUM, (*pseudo*, and *angustura*, 'false angustura,') *Bruceine*.

PSEUDAPHE, (*pseudo*, and *ἅψη*, 'touch, feeling,') *Pseudæsthesia*.

PSEUDARTHRO'SIA, *Pseudarticula'tio*, *Pseudartic'ulus*, *Articula'tio notha seu artificia'lis*, (*pseudo*, and *αἰθρον*, 'a joint,') A false joint or articulation; *Artic'ulus no'vus* seu *epu'rius* seu *præternatural'is*, (F.) *Articulation surnuméraire*.

PSEUDARTICULATIO, *Pseudarthrosis*.

PSEUDECOIA, *Pseudacoe*.

PSEUDENCEPH'ALUS, *Paraceph'alus*, (*pseudo*, and *εγκεφαλος*, 'the encephalon,') A monster, whose cranium is open in its whole extent, from before to behind, its base supporting a vascular tumour.

PSEUDESTHÉSIE, *Pseudæsthesia*.

PSEUDINOMA, (*pseudo*, *is*, *αἰς*, 'fibre,' and *oma*.) Scirrhus.

PSEUDO, (*ψευδής*.) False—p. Asthma, Dyspnoea.

PSEUDOBLEPH'SIA, *Pseudobleph'sis*, *Pseudo-ra'sis*, *Suffu'sio oculi*, (*pseudo*, and *βλέψις*, 'sight.') *Pseudop'sia*, *Parop'sis illusoria*, *Suffu'sio*, *Phantasma*, False sight. A generic name, used by Cullen for perversion of vision. See *Metamorphopsia*.

PSEUDOBLEPSIS, *Paropsis*.

PSEUDOCARCINOMA LABII, Cancer aquaticus.

PSEUDOCATARACTA, see *Cataract*.

PSEUDOCHROMIA, (*pseudo*, and *χρῶμα*, 'color,') *Achromatopsia*.

PSEUDOCROUP, Asthma thymicum.

PSEUDOCYESIS, (*pseudo*, and *κύσις*, 'pregnancy,') Pregnancy, false—p. *Molaris*, *Mole*.

PSEUDOENCEPHALITIS, (*pseudencephalus*, and *itis*.) Hydronecephaloid disease.

PSEUDOFEVER, Irritation, morbid.

PSEUDOGEUSIA, *Pseudogen'sia*, *Pseudogen'stia*, (*pseudo*, and *γεύσις*, 'taste,') *Hallucination gustiva*. False taste.

PSEUDOGEUSTIA, *Pseudogeusia*.

PSEUDOHELMINTHES, (*pseudo*, and *ἕλμινς*, *gen.* *ἕλμινθς*, 'a worm,') *Ectozoa*.

PSEUDOLI'EN, *Pseudosplen*, (*pseudo*, and *lien*, 'spleen,') Glands situate in the neighbourhood of the spleen; probably, lymphatic glands.

PSEUDOMECHOACANA, *Convolvulus panduratus*.

PSEUDOMEDICUS, (*pseudo*, and *medicus*, 'a physician,') *Charlatan*.

PSEUDOMELANOTIC FORMATION, (*pseudo*, and *melanosis*,) see *Anthraxis*—p. *Meninx*, Membrane, false—p. *Metamorphosis polyposa*, *Polypus*.

PSEUDOMEMBRANA, (*pseudo*, and *membrana*.) Membrane, false.

PSEUDOMEMBRANE, Membrane, false.

PSEUDOMEMBRANOUS, *Pseudomembranous*, (*F.*) *Pseudomembraneux*. Relating or appertaining to false membranes.

PSEUDONARCISSUS, *Narcissus pseudonarcissus*.

PSEUDONOMA'NIA, (*pseudo*, and *mania*.) A form of insanity, which is characterized by a morbid propensity for lying.

PSEUDOPARASITES, *Ectozoa*.

PSEUDOPERIPNEUMONIA, *Peripneumonia notha*.

PSEUDOPHLOGOSIS VENTRICULI RESOLUTIVA ET COLLIQUATIVA, (*pseudo*, and *φλογωσις*, 'inflammation,') *Gastromalacia*.

PSEUDOPHTHISIS. 'False phthisis.' Emaciation occasioned by other cause than organic lesion of the lungs.

PSEUDOPLAS'MATA, (*pseudo*, and *πλασσω*, 'I form,') A term applied to malignant heterologous tumours.

PSEUDOPLEURITIS, *Pleurodynia*—p. *Polypi*, see *Polypi*—p. *Pyrethrum*, *Achillea ptarmica*—p. *Spleen*, *Pseudolien*.

PSEUDOPNEUMONIA, *Peripneumonia notha*.

PSEUDOPNEUMONITIS, *Peripneumonia notha*.

PSEUDOPSIA, (*pseudo*, and *ὄψις*, 'vision,') *Pseudoblepsia*.

PSEUDORASIS, (*pseudo*, and *ὤραω*, 'to see,') *Pseudoblepsia*.

PSEUDOREX'IA, *Pseudorex'ia*, (*pseudo*, and *ἄρεσις*, 'appetite,') False appetite.

PSEUDORHONCHUS, (*pseudo*, and *ῥογχος*, 'rattle,') A sound heard during respiration, which appears to be produced, like the ordinary rhonchi, in the air-tubes, but is exterior to them

—as in the case of *pleural pseudorhonchi*. See *Rattle* and *Râle*.

PSEUDOSCOPI'IC, *Pseudoscop'icus*, (*pseudo*, and *σκοπεῖν*, 'I view,') Conveying false impressions to the eye; as where projecting surfaces are seen depressed, and conversely.

PSEUDOS'MIA, *Pseudosphre'sia*, *Pseudosphre'sia*, (*pseudo*, and *ὀσμη*, 'smell,') False sense of smell.

PSEUDOSPHERIA, (*pseudo*, and *σφῆρις*, 'smell,') *Pseudosmia*.

PSEUDOSYPHILIS, *Syphilis pseudosyphilis*.

PSEUDOTIHANATUS, (*pseudo*, and *θανάτος*, 'death,') *Asphyxia*.

PSEUDOVARIOLÆ, *Variceila*.

PSEUDYMEN, (*pseudo*, and *ὑμην*, 'a membrane,') Membrane, false.

PSIDIUM POMIFERUM. The *Ap'ple Guava*, *Guava*, *Guayava*. *Ord.* *Myrtaceæ*. This plant and *P. Pyrif'erum* bear fruits; those of the former like apples—of the latter like pears. The apple kind has an acid flavour; the other is sweet. Of the inner pulp of either, the natives of the Indies make jellies, and, of the outer rind, tarts, marmalades, &c. They are somewhat astringent: this quality, indeed, exists in every part of the tree, and abundantly in the leaf-buds, which are occasionally boiled with barley and liquorice as an excellent drink in diarrhoea. A simple decoction of the leaves is said to be useful in the itch and most cutaneous eruptions.

PSILOSI, (*ψιλλωσις*, from *ψιλλω*, 'to remove the hair,') *Depilation*.

PSILOTHRUM, (*ψιλλωθρον*, from *ψιλλω*, 'to remove the hair,') *Bryonia alba*, *Depilatory*.

PSIMYTHION, (*ψιμυθιον*,) *Plumbi subcarbonas*.

PSIMYTHUS, (*ψιμυθος*,) *Plumbi subcarbonas*.

PSOA, (*ψοα*,) *Dysodia*.

PSO'Æ, (*ψοα*,) *Psoi'æ*, *Psy'æ*, see *Lumbi*, 'the loins.' The *psœa* muscles; *Alop'eces*, *Neuromel'træ*.

PSO'AS MAG'NUS, (*ψοα*, 'the loins,') *Pso'as* seu *lumba'ris inter'nus*, *Femur noventium sex'tus*, *Pré-lombo-trochantin*, *Pré-lombo-trochantinien* (*Ch.*) A muscle, seated on the lateral parts of the lumbar portion of the vertebral column; and passing down to the superior and anterior part of the thigh. It is long, fusiform, and attached, above, to the body and transverse processes of the first four lumbar vertebrae and last dorsal. Below, it terminates by a very thick tendon, common to it and the iliaceus muscle,—which is inserted into the top of the lesser trochanter. This muscle bends the thigh upon the pelvis, and carries it in rotation outwards. It may, also, bend the pelvis on the thigh, and the loins on the pelvis. It acts considerably in station and progression.

PSOAS PAR'VUS, *Pré-lombo-pubien* (*Ch.*) A muscle situate anteriorly to the last. It is long, thin, flat, and narrow, and is attached, above, to the body of the last dorsal vertebra, and below, to the linea ileopectinea, by means of a slender tendon, which furnishes two expansions—one to the *iliac aponeurosis*, and the other to the *pelvis*. This muscle bends the vertebral column on the pelvis, and, reciprocally, the pelvis on the vertebral column.

PSODYMUS, (*ψοα*, 'the loins,' and *δυω*, 'to mingle,') *Gastrodydymus*.

PSOÆ, *PSOÆ*.

PSOITE, *Psoitis*.

PSOÏTIS, *Inflammatio mus'culi Pso'as*, *Lumbago inflammatoria spasmad'ica*, (*F.*) *Psoïte*, *Inflammation du muscle Psoas*, (*ψοα*, 'the loins,' 'the psœa muscles,' and *itis*.) Inflammation of the psœa muscles and of the parts situate before the lumbar vertebrae; accompanied, from its

commencement, by considerable fever, violent pains in the part affected, and in the thigh of the same side, which remains commonly half bent. Psoritis may terminate by suppuration, and form a lumbar abscess.

PSOÏLE, (ψωλη,) Penis. Also, the glans when uncovered.

PSOLON, (ψωλον,) Membrosus.

PSOLON/CUS, (*psole*, and *oikos*, 'a tumour,') Swelling of the glans, or of the penis generally.

PSO/LUS, (ψωλος, from *psole*.) One who has the glans uncovered. One who has been circumcised.

PSOPHOMETER, (*psophus*, and *μετρον*, 'a measure,') Stethoscope.

PSOPHUS, (ψωφος,) a noise; a sound.

PSORA, (ψωρα, from ψω, 'to scratch,') *Scabies*, *Phlyctis* seu *Eccyptosis* *Scabies*, *Scabiola*, *Itch*, *Scurfy*, *Scotch fiddle*. (Prov.) *Ichet*, *Serat*, *Yecke*. (Sc.) *Reif*, *Yunk*, *Yuke*, *Yecke*. (F.) *Gale*, *Mal Saint-Main*, *Rogne*. A contagious eruption of very minute pimples—pustular, vesicular, papular, intermixed, and alternating, itching intolerably and terminating in scabs. It is found to be connected with an insect of the genus *Acarus*;—the *Ac'arus* seu *Sarcop'tes* seu *Sarcop'tus* seu *Sarcocop'tes* seu *Phthi'rium* seu *Cheyle'tes* *Scabi'ei*, *Ac'arus* *Ci'ro*, *Itch insect*, *Itch animal'cule*, *Itch mite*, (F.) *Ciron*. The itch occupies, particularly, the spaces between the fingers, the back of the hands, wrists, elbows, axillæ, groins, hams, &c., and rarely affects the face. The best applications, for its cure, are,—the *Unguentum Sulphuris* or *Ung. Sulph. comp.* night and morning; but the unpleasant smell of the sulphur has given occasion to the use of other means;—as the *White Hellebore*, *Potass* in deliquescence, *Muriate of Ammonia*, *Sulphuric Acid*, &c. The repugnance to the smell of sulphur ought not, however, to be regarded; especially as, in the course of four or five days, it will, in almost all cases, produce a perfect cure.

PSORA AGRIA, Psoriasis inveterata.

PSORA EBRIO'RUM, *Drunk'ard's Itch*. An eruption, attended by afflicting itching, to which elderly people, who have been much addicted to drinking alcoholic liquors, are liable.

PSORA LEPROSA, Psoriasis—p. *Squamosa*, Psoriasis.

PSORALEA, Psoralea pentaphylla.

PSORA'LEA ESCULEN'TA, *Breadroot*, *Indian turnip*, *Pomme blanche* ou de *Prairie*, (ψωραλεως, 'scurfy or scaly,' in reference to its scurfy glands or dots.) *Oral*. Leguminosæ. A plant, resembling the beet in shape, which is found near the Rocky Mountains. It contains a white pulpy substance, which is sweet and palatable.

PSORA'LEA GLANDULO'SA, *Albaquill'la*. A shrub, used in Chili as a vulnerary. The Paraguay tea is referred to it by M. Lenoble, of Montevideo. See *Ilex Paraguensis*.

PSORALEA PENTAPHYLLA, *Psora'lea*, *Contrayer'va* no'va, *Mex'ican Contrayer'va*. The root is little, if at all, inferior to the contrayer'va in its medical properties; which, by the by, are unimportant.

PSORENTERIA, (*psora*, and *εντερων*, 'intestinal,') Cholera.

PSORIASIS, or PSORIA'SIS, (ψωριασις,) formerly, the state of ψωρα. *Lepido'si psori'asis*, *Serpe'do*, *Serpi'go*, *Impeti'go* (of some), *Scabi'ies sic'ca* seu *feri'na*, *Ps'o'ra lepro'sa* seu *aquamo'sa*, *Scal'y Tet'ter*, *Dry Scale*, *Dry Scall*, (F.) *Dartre squameuse* lichenoidé, *D. écailleuse*, *Grattelle*, *Gale miliaire*, *G. canine* ou *âche*. A cutaneous affection, consisting of patches of rough, amorphous scales; continuous, or of indeterminate outline; skin often chappy. Psoriasis occurs under a considerable variety of forms. The surface, under the scales, is more tender and irritable than in iecra, which psoriasis, in some respects,

resembles. Dr. Willan has given names to eleven varieties of psoriasis: the chief of which are the *Psori'asis gutta'ta*, *P. diffu'sa*, one form of which is the *Bu'ker's Itch*, *Psori'asis pinto'rnum*, and *Grocer's Itch*; and another, the *Washer'woman's Scall*, the *P. gyra'ta*, and *P. invetera'ta* seu *Ps'o'ra a'gria*. The treatment of irritable psoriasis must be antiphlogistic, with the internal use of the fixed alkalies, sulphur, &c. They do better without any local application. The more local and less inflammatory are ameliorated by emollient and soothing applications; or by slightly stimulating means, seabathing, the *Ung. hydrag. nitrico-oxyd.*, *Ung. oxidi zinci*, &c.

PSORTIC, (ψωρικός,) *Psor'icus* (F.) *Psorique*, *Galenx*; (from *psora*.) That which has the nature of itch. A medicine for curing the itch.

PSOROCOMI'UM, (*psora*, and *κομω*, 'I take care of,') An hospital for those affected with itch.

PSORODES, (ψωροδης,) Psorous.

PSO'ROID, *Psoroi'des*; (from *psora*, and *ειδος*, 'resemblance,') Resembling psora.

PSOROPHTHALMIA, (ψωροφθαλμία,) see Ophthalmia tarsi—p. Neonatorum, see Ophthalmia (purulenta infantum).

PSO'ROUS, *Psoro'sus*, *Psoro'des*, (from *psora*.) Appertaining or relating to itch.

PSYÆ, (ψυα,) Psœ.

PSYCHAGOGA, Psychagogica.

PSYCHAGOG'ICA, (ψυχωγωγικά,) *Psychago'ga*, (*psycha*, and *αγω*, 'I lead,') Medicines which resuscitate, when life seems extinct; as in syncope and apoplexy.

PSYCHAL, Psychical.

PSYCHE, (ψυχη,) Anima, Thymus.

PSYCHEISM, (from *psyche*,) Magnetism, animal.

PSYCHIA'TER, *A'lienist*. One who treats diseases of the mind—a *Mad-doctor*, *A'lienist phy-sic'ian*, (F.) *Médecin de fous*, *Aliéniste*.

PSYCHIATRI'A, (*psycho*, and *ιατρεία*, 'healing,') Treatment of diseases of the mind.

PSY'CHICAL, *Psy'chicus*, (ψυχικός, from *psyche*,) *Psy'chal*. That which relates to the mind or mental endowments. *Psycholog'ical* is not unfrequently used with the like signification.

PSYCHICAL or PSYCHOLOGICAL MEDICINE includes the history, pathology, treatment, &c., of mental affections.

PSYCHO, (from ψυχη, anima, 'the mind, the soul,') in composition the same.

PSYCHOLOGICAL, see Psychical.

PSYCHOL'OGY, *Psycholog'ia*, (*psycho*, and *λογος*, 'discourse,') A treatise on the intellectual and moral faculties. Also, the intellectual and moral faculties or sphere.

PSYCHONOSEMA, (*psycho*, and *νοσημα*, 'disease,') see Psychoses.

PSYCHONOSOL'OGY, *Psychonosolog'ia* (*psycho*, *nosos*, 'disease, and *λογος*, 'a description,') The doctrine of, or a treatise on, diseases of the mind.

PSYCHOPATHIA, (*psycho*, and *παθος*, 'affection,') see Psychoses.

PSYCHOPATHIES, Psychoses.

PSYCHOPATHIST, *Psychopathis'ta*. One versed in diseases or affections of the mind.

PSYCHOR'AGES, (ψυχωραγίαι,) *Agoni'zana*, *Moribund'us*, *Mor'ibund*, *Fey*, *Fee*, *Fie*, (*psycho*, and *ραγη*, 'rupture,') Dying; in *Articulo Mortis*, "in the article of death;" in *extre'mis*. *Moribund* means, also, one who is dying.

PSYCHORAGIA, (ψυχωραγία,) Agony, Death.

PSYCHORRHAGIA, (ψυχωρραγία,) Agony, Death.

PSYCHO'SIS, (*psyche* and *osis*.) Neurosis of the intellect. Disease of the mind; *Psychop'a-thy*, *Psychonose'ma*, *Psychopathi'a*.

PSYCHOTHERAPEIA, (*psyche*, and *θεραπεωω*, 'I attend upon the sick,') Treatment of disease through the mind.

PSYCHOTICA, (from ψυχω, 'to animate,')
Analeptica.

PSYCHOTRIA CROCEA, *Palicourea crocea*—
p. Emetica, *Ipecacuanha*.

PSYCHOTROPHUM, (*psycho*, and τρεφειν, 'to
nourish,') Beticana officinalis.

PSYCHROLUSIA, *Psychrolitron*, (Ψυχρολουσία,
from ψυχρος, 'cold,' and λουω, 'I wash,') A cold
bath.

PSYCHROPHOBIA, (Ψυχρος, 'cold,' and φοβος,
'fear,') Dread of cold, especially of cold water.
Impressibility to cold.

PSYCHTICA, (ψυκτικός, 'cooling,') Refrigerants.
PSYCHUS, (ψυχος,) Cold.

PSYDRACIA, *quasi* ψυκρά ὑδράκια, 'cold drops,'
(Ψυδράκια, from ψυδρά, gen. ψυδρακος, 'a small
blister or pimple,') The ancients sometimes used
this term for *pimples*; at others, for *vesicles*.
Willan and Bateman define *Psydracium* to be, a
small pustule, often irregularly circumscribed,
producing but a slight elevation of the cuticle,
and terminating in a laminated scab. Many of
the psydracia usually appear together, and be-
come confluent; and, after the discharge of pus,
they pour out a thin, watery humour; which fre-
quently forms an irregular incrustation. Frank
limits the term to a psoriform eruption, which
differs from itch in not originating from an insect,
and is not contagious;—perhaps the lichen.
Authors have differed regarding its acceptance.

PSYDRACIA ACNE, Acne.

PSYDRACIUM, sing. of Psydracia.

PSYLLIUM, (from ψυλλον, ψυλλα, or ψυλλος,
'a flea,' from the likeness of the seeds,) *Plantago*
psyllium—p. Erectum, *Plantago psyllium*.

PTARMICA, (παρμηκη; from παρμεν, 'to
sneeze,') *Achillea ptarmica*—p. Montana, *Arnica*
montana.

PTARMICUM, (παρμικον,) *Erhrine*, *Sternuta-*
tory.

PTARMUS, (παρμος,) Sneezing.

PTOLEA, (πτελεα,) *Ulmus*.

PTOLEA TRIFOLIATA, *Shrubby trefoil*, *Sicamp*
dogwood, *Stink'ing ash*, *Stink'ing prairie bush*,
Wingseed, *Wufer ash*, *Hop tree*; indigenous.
Order, *Xanthoxylaceae*. Flowering in June. Is
said to be possessed of antiperiodic virtues, and
to be anthelmintic.

PTERIS, (πτερίς, 'a wing,' from the shape of its
leaves.) *Polypodium filix mas*.

PTERIS AQUILINA, *Pteris*, *Thelypteris*, *Filix*
Nymphaea seu fœminina, *Asplenium aquilinum*,
Common Brake or Bracken, *Female Fern*; *Eagle*
Fern, *Bracken* (F.) *Fougère femelle*, *La grande*
Fougère. *Nat. Ord.* Filices. The root is con-
sidered to be anthelmintic; and as efficacious as
that of the male fern in cases of tapeworm.

PTERIS PALUSTRIS, *Asplenium filix fœmina*.

PTERNA, (πτερινα and πτερνιον,) *Calcaneum*.

PTERNIUM, (πτερνιον, dim. of pterna,) *Calca-*
neum.

PTERNOBATES, (πτεροβατης, from pterna, and
βαινω, 'to walk,) *Calcigradus*.

PTEROCARPUS ERINACEA, see *Kino*—p.
Marsupium, see *Kino*—p. *Sandalinus*, *P. Sanda-*
linus.

PTEROCARPUS SANTALINUS, *P. Santalinus*,
(πτερον, 'a wing,' and καρπος, 'fruit,') *Santalum seu*
Santalum Rubrum, *Red Sand'ers* or *Saunders*
Tree, (F.) *Santal rouge*. *Ord.* Leguminosae. *Sex.*
Syst. *Diadelphia Decandria*. *Red Saunders wood*,
Santalum, (Ph. U. S.,) *Pterocarpus*, (Ph. B. 1864,) *Lignum sandalinum*, is used only as a colouring
matter: it gives to rectified spirit a fine deep red.
The juice of this tree, like that of *P. Draco*,
affords a species of *Dragon's Blood*.

PTEROCAULON PYNOSTACHYUM, (πτερον,
'a wing,' and καυλος, 'a stalk;' and πυνος, 'thick,'
and σπας, 'an ear or spike,') *Onyza pynostachya*,
Blackroot; indigenous. *Order*, *Compositae*. The

root is used as an alterative in the form of de-
coction. It is said to be possessed of cathartic
properties.

PTEROSPORA ANDROMEDEA, (πτερον, 'a
wing,' and σπαρα, 'seed,' in allusion to the singular
wing borne by the seeds,) *Scaly Dragonclaw*,
Dragonroot, *Felverroot*, *Abany beech'drop*. An
indigenous plant, *Nat. Ord.* *Monotropææ*, *Sex.*
Syst. *Decandria Monogynia*:—found in the State
of New York, which blossoms in July. The root
has been regarded as an anthelmintic, diuretic,
emmenagogue, &c.: but is not used.

PTERYGION, *Pterygium*, (πτερυγιον, dim. of
pteryx, 'a wing,') *Al'a*, *Web* [?], *Web eye* [?],
Pin and *web* [?], *O'nyx*, *Pyo'sis*, *Un'guis*, (F.)
Drapeau, *Onglet*. A varicose excrescence of the
conjunctiva, of a triangular shape, and commonly
occurring at the inner angle of the eye; whence
it extends over the cornea. Sometimes two or
three pterygia occur on the same eye, and cover
the whole cornea with a thick veil, which totally
prevents vision. According to Scarpa, the ancients
gave the name *Panniculus* to this last complica-
tion. If the pterygion cannot be dispelled by
means of discutient collyria, it must be removed
by the scissors.

PTERYGIUM, *Pterygion*, *Scapula*—p. *Digiti*,
Paronychia—p. *Lardaceum*, *Pinguecula*—p. *Pin-*
gue, *Pinguecula*.

PTERYGO, in the composition of anatomical
terms, means relating, or appertaining, to the
pterygoid processes.

PTERYGO-ANGULI-MAXILLARE, *Ptery-*
goides internus—p. *Colli-maxillare*, *Pterygoides*
externus—p. *Maxillare*, *grand*, *Pterygoides in-*
ternus—p. *Maxillare*, *petit*, *Pterygoides ex-*
ternus—p. *Palatine canal*, *Pterygoid canal*.

PTERYGOPALATINE, *Pterygopalati'nus*.
That which belongs to the pterygoid process and
palate.

PTERYGOPALATINE ARTERY or *Superior pha-*
ryngeal is a very small artery, and is given off by
the internal maxillary at the bottom of the zygomatic
fossa: after having passed through the
pterygopalatine canal, it is distributed to the
superior part of the pharynx, the sphenoid bone,
and the Eustachian tube.

PTERYGOPALATINE CANAL'. A small canal,
formed by the internal ala of the pterygoid pro-
cess, and the palate bone. It gives passage to
the vessels of the same name.

PTERYGOPHARYNGEUS, *Constrictor pha-*
ryngeus—p. *Staphylinus*, *Circumflexus*.

PTERYGOSTAPHYLINUS INTERNUS, *Le-*
vator palati—p. *Syndesmo-staphyli-pharyngien*,
Constrictor pharyngis.

PTERYGODES, (πτερυγώδες, from *pteryx*, and
odes,) *Alatus*.

PTERYGOID, *Pterygoi'des*, *Pterygoï'des*, *Al-*
iformis, (*pteryx*, and εἶδος, 'form,') A name given
to two processes at the inferior surface of the
sphenoid bone; the two laminae which form them
having been compared to wings. These *ala* or
wings or *processes*,—*Processus aliformes*,—have
been distinguished, according to their position,
into *internal* (*entopterygoid*) and *external*.

PTERYGOID ARTERY, *Vid'ian Artery*, *Ramulus*
Ductus Pterygoï'dis, *Arteria pharyngea* a *super-*
ma, arises from the internal maxillary artery, at
the bottom of the zygomatic fossa, and enters the
pterygoid canal with the nerve of the same name,
to proceed to be distributed on the Eustachian
tube and velum palati. The name, *pterygoid*
arteries, is likewise given to the branches, which
the internal maxillary and some of its divisions,
as the meningeal media, and posterior temporalis
profunda, furnish to the pterygoid muscles, be-
hind the neck of the lower jaw.

PTERYGOID BONE, *Sphenoid*.

PTERYGOID CANAL', *Vid'ian canal*', *Ductus pte-*

rygoides. *Canalis Vidianus*, (F.) *Conduit pterygoïdien* ou *Vidian*. *Pterygopalatine canal*, is a narrow channel, which traverses, in a direction from before to behind, the base of the pterygoid process, and gives passage to the pterygoid nerve. The *foramina* of the sphenoid, which terminate the canals, have the same epithets applied to them.

PTERYGOID FOS'SA, *Fos'sa Pterygoïde'a*, (F.) *Fosse pterygoïdienne* ou *pterygoïde*, is the depression which separates the two laminae or alae of the pterygoid process.

PTERYGOID MUS'CLES, *Ala'es mus'culi*, *Aliformes mus'culi*, are two in number. 1. The *Pterygoïdeus externus* seu *minor*, (F.) *Pterygocolli-maxillaire*, *Petit Pterygo-maxillaire* (Ch.) *Petit pterygoïdien*, *Pterygoïdien externe*, is a thick, short muscle, which arises, by short aponeuroses, from the outer surface of the external ala of the pterygoid process, and from the outer surface of the great ala of the sphenoid bone; and is inserted at the anterior part of the neck of the lower jawbone, and the corresponding part of the interarticular fibrocartilage. This muscle draws the condyle of the lower jaw and the interarticular ligament forwards. It carries the chin forwards, and to the opposite side. If the two external pterygoids act together, the jaw is carried directly forwards. 2. The *Pterygoïdeus internus* seu *major*, *Masse'ter internus*, *Latens* in *O'ry*, (F.) *Grand Pterygo-maxillaire*, *Pterygo-anguli-maxillaire*, *Grand Pterygoïdien* ou *Pterygoïdien interne*. A thick, quadrilateral muscle, which is attached by short and very strong aponeuroses; above, to the pterygoid fossa, whence it proceeds downwards, backwards, and outwards, to be inserted into the inner surface of the ramus of the lower jaw, near its angle. This muscle raises the lower jaw, and carries it a little forwards. It also performs the grinding motion.

PTERYGOID NERVES. These are two in number. 1. The *Pterygoid* or *Vidian Nerve*. It arises from the posterior part of the sphenopalatine ganglion; traverses the Vidian canal of the sphenoid; and, beyond, divides into two filaments. 1. A *superior* or *cranial*, *Nervus Superficialis* sive *petrosus*, which ascends into the cranium, and enters the hiatus of Fallopius to unite with the facial nerve. 2. An *inferior* or *carotid*, *Nervus profundus* seu *major*, which traverses the carotid canal, and unites with the ascending branches of the superior cervical ganglion of the trisplanchnic. The *Pterygoid Nerve*, properly so called, *Nervus musculi pterygoïdei*, is given off by the inferior maxillary branch of the 5th pair.

PTERYGOID PROCESSES, see *Pterygoid*.

PTERYGOIDEUS, *Pterygoid*.

PTÉRYGOÏDIEN EXTERNE, *Pterygoideus externus*—p. *grand* ou *interna*, *Pterygoideus internus*—p. *Petit*, *Pterygoideus externus*.

PTERYGO'MA, gen. *Pterygo'matis*, (πτερυγμα, from *pteryx* and *oma*, 'a projecting or pendulous body.') A swelling of the vulva, which prevents coition.

PTERYX, gen. *Pter'ygis*, (πτερυξ, gen. πτερυγος,) *Ala*.

PTILOS, (πτίλος,) *Deplumatio*, *Madarosis*.

PTILOÏSIS, (πτίλωσις,) *Deplumatio*, *Madarosis*.

PTISANA, (πτισανη, 'peeled barley, and a decoction of the same,') *Tisane*—p. *Communis*, *Decoctum hordei compositum*—p. *Hippocratica*, *Decoctum hordei*.

PTISSANA, (πτισσανη,) *Tisane*.

PTOCHIA'TER, *Ptochia'trus*, (πτωχος, 'poor,' and ιατρος, 'a physician,') A physician to the poor, or to an establishment for the poor.

PTOCHIATRIA. Practice amongst the poor. Also, an institution for the treatment of the diseases of the poor.

PTOCHOCOMI'UM, (πτωχος, 'poor,' and κομω,

'I take care of;') *Ptochodochi'um*, *Ptochei'um*, *Ptochotroph'eum*. An almshouse. An hospital for the poor.

PTOCHODOCHIUM, (πτωχος, 'poor,' and δαχμου, 'to receive,') *Ptochocomium*.

PTOCHOTROPHEUM, (πωχοτροφειον, from πτωχος, 'poor,' and τρεφειν, 'to nourish,') *Ptochocomium*.

PTOMA, gen. *Pto'matis*, (πτωμα,) *Cadaver*.

PTOSIS, (πτωσις,) *Prolapsus*—p. *Bulbi oculi*, *Exophthalmia*.

PTOSIS IRIDIS, *Iridoccl'is*, *Her'nia seu Proci'dent'ia seu Prolap'sus seu Staphylo'ma I'ridis*, *Iridopto'sis*. A prolapsus of the iris through a wound of the cornea. It is known by a blackish tubercle, which projects a little from the cornea in various forms, and which, owing to its supposed resemblance to the head of a fly, has been called *Myioceph'alon*, and *Myoceph'alon*, (μυια, 'a fly,' and κεφαλη, 'head,')

PTOSIS PALPEBRE, *Blepharoptosis*.

PTYALOGOGUE, (*ptyalum*, and αγω, 'to lead,') *Sialogogue*.

PTYALIN, (from *ptyalum*,) see *Saliva*.

PTYALISED, *Salivated*.

PTYALISM, *Salivation*.

PTYALISME, (πτυαλισμος,) *Salivation*—p. *Mercuriel*, *Salivation*, *mercurial*.

PTYALISMUS, *Salivation*—p. *Acutus mellitus*, see *Salivation*—p. *Acutus sympatheticus*, see *Salivation*—p. *Hydrargyratus*, *Salivation*, *mercurial*—p. *Iners*, *Slavering*—p. *Mellitus*, *Melitoptyalismus*—p. *Mercurialis*, *Salivation*, *mercurial*—p. *Pyrosicus*, *Pyrosis*.

PTYALOGOGUE, (*ptyalum*, and αγω, 'to drive,') *Sialogogue*.

PTYALUM, (πτυαλον,) *Saliva*.

PTYELUM, (πτυελον,) *Saliva*, *Sputum*.

PTYSIS, (πτωσις,) *Exspuition*.

PTYSMA, gen. *Ptysm'atis*, (πτυσμα,) *Saliva*, *Sputum*.

PTYSMAGOGUE, (*ptysma*, and αγω, 'to drive,') *Sialogogue*.

PTYSMATIS'CHESIS, *Ptysmatos'chesis*, (*ptysma*, and χηω, 'to retain,') *Suppression of pulmonary expectoration*.

PTYSMATOSCHESIS, *Ptysmatischesis*.

PUANTEUR, (from *putere*, 'to smell badly,') *Dysodia*.

PUBENS, 'at the age of puberty,' *Ephebus*.

PUBER, *Ephebus*.

PUBERAL, *Pubera'lis*, *Puber'ic* or *Pub'erie*, *Puber'icus*. Relating or appertaining to the age of puberty—as 'a *puberal* disease.'

PUBÈRE, *Ephebus*.

PUBERIC, *Puberal*.

PUBERTAS, *Ephebia*, *Puberty*.

PUBERTY, *Pu'bertas*, *Ho'te*. *Pu'bes*, *Aphrodis'ia*, *Ephebi'a*, *Ephe'bia*, *Ephe'botes*, *Ephebo'synè*, *Pubes'cence*, (F.) *Puberté*, (from *pubescere*, 'to begin to have hair on the chin and sexual organs,') The period of life at which young people of both sexes are *pubes'cent* or *nubile*. According to the civil law, puberty occurs at 12 years in females, and 14 in males. The period varies somewhat in different climates, and in different individuals of the same climate.

PUBES, *Pu'bis*, *Epis'chion*, *Oleis*. *Pee'ten*: the *Pu'bic region*. The middle part of the hypogastric region; so called, because it is covered with hair, in both sexes, at the period of puberty. The hair is also called *Pu'bes*, *Lanu'go* seu *Pi'ti Pudend'o-rum*, *Planta'r'ium*, *Fem'inal*; and, of old, *Epicten'ion* and *pubertas* signified the same. See *Tressoria*.

PUBES, *Puberty*.

PUBESCENCE, *Puberty*.

PUBESCENT, see *Pub'erty*.

PUBIÆUS, *Pubic*.

PUB'IC, *Pu'bion*, *Pu'bicus*, *Pubi'cus*, (F.)

PUBIEN. That which belongs to or concerns the pubis.

PUBIC ARCH. *Arch. of the Pubis*, (F.) *Arche pubienne*. The arch at the anterior part of the inferior circumference of the pelvis. It is more expanded in the female than in the male, to correspond with the additional functions which she has to execute.

PUBIC ARTICULATION, *Sym'physis Pu'bis*, results from the junction of the ossa pubis. The medium of union is cartilage.

PUBIC LIG'AMENTS are two very solid, fibrous fasciæ, situate before and beneath the symphysis pubis, which they strengthen. The one is called the *anterior pubic ligament*; the other the *inferior or subpubic*.

PUBIC RE'GION, *Re'gio Pu'bis*. The region of the pubes.

PUBIEN, Pubic.

PUBIO-COCCYGIEN-ANNULAIRE.

Pubio-coccygeus annularis. Dumas gave this name to the levator ani and ischio-coccygeus, which he considers a single muscle. See *Levator Ani*.

PUBIO FÉMORAL. *Pectinalis*—p. Femoralis, *Adductor longus femoris*—p. *Ombilical*, *Pyramidalis abdominis*—p. *Prostaticus*, *Compressor prostate*—p. *Sous-ombilical*, *Pyramidalis abdominis*—p. *Sternal*, *Rectus abdominis*—p. *Urethral*, see *Compressor Urethrae*.

PUBIS, Pubes, Pubis os.

PUBIS OS, *Os Pect'linis*, *Episei'on*, *Episi'on*, *Epis'chion*, *Sharebone*, *Barbone*, (Sc.) *Arkbein*. The anterior part of the os innominatum; so called because it corresponds to the genital organs and pubic region. The angle of the pubis is formed by the union of the body and rami of each side.

PUBLICÆ PARTES, 'public parts.' Those organs are so called, the object of whose functions is the nutrition of all the others. Such are the digestive and respiratory organs.

PUCCINIA FAVI, (after Th. Puccini, Professor of Anatomy in Florence), see *Fungus*.

PUCCOON, *Sanguinaria Canadensis*—p. Red, *Sanguinaria Canadensis*—p. Yellow, *Hydrastis Canadensis*.

PUCE, (from *pulex*, gen. *pu'licis*.) *Pulex irritans*—p. *de Bourguigne*, see *Anthrax*—p. *Maligne*, see *Anthrax*—p. *Pénétrante*, *Chique*.

PUCELAGE, (from [L.] *puella*, 'a young girl,') Virginity.

PUCELLE, Virgin.

PUCHA-PAT, *Pogostemon patchouli*.

PUCHURY, see *Pichurim beans*.

PUCHYRY, see *Pichurim beans*.

PUCKLE, *Papula*.

PUDDINGGRASS, *Mentha pulegium*.

PUDDINGPIPE TREE, *Cassia fistula*.

PUDENDA, pl. of *pubendum*, (from *pubere*, 'to be ashamed,') Genital organs—p. *Muliebria*, *Vulva*—p. *Virorum*, *Pubibilia*.

PUDEND'AGRA, (*pubenda*, and *aypa*, 'a seizure,') According to some, this term means pain in the genital parts; *Do'lor pudendo'rum*, *Men'tulagra*, *Edeceodyn'ia*: according to others, it is synonymous with syphilis.

PUDENDAGRA PRURIENS, *Prurigo pudendi muliebris*.

PUDEND'DAL, *Pudenda'lis*. Relating or appertaining to the pudenda.

PUDENDAL HERNIA. Hernia, which descends between the vagina and ramus ischii, and forms an oblong tumour in the labium; traceable in the pelvis as far as the os uteri.

PUDENDALIS, *Pudendal*.

PUDENDALIS LONGUS INFERIOR, see *Sciatic nerve*, lesser.

PUDENDUM, FEMALE, Vulva—p. *Muliebre*, *Vulva*.

PUDENDUS, *Pubic*.

PUDIBIL'IA. The male genital organs; *Genitalia Viri*, *Virilia*, *Pensil'ia*, *Instrumen'ta*, *Puden'da vir'o'rum*, *Vir'ia membra*, *Andro'dia'ia*.

PUD'IC, *Pud'icus*, *Puden'dus*, (F.) *Honteux*, (from *pubere*, 'to be ashamed,') That which causes shame. A name given to the genital organs as well as to different other parts.

PUDIC ARTERIES are distributed to the organs of generation. They are,—1. The *internal Pud'ic*, *Arte'ria Pud'ica*, *A. puden'da commun'is*, (F.) *Artère honteuse interne*, *A. Sous-pubienne* (Ch.), *Genital Artery*. It is furnished by the hypogastric, and divides into two branches: the one—the *inferior or perineal*—which gives off the inferior hemorrhoidal and the artery of the septum; the other—the *superior*, (F.) *Ischio-pénienne*—which furnishes the transverse perineal artery, and divides into two branches; the artery of the *corpus cavernosum*, and the *dorsalis penis*. This latter, in the female, has the name *Dorsalis Clitoridis*. 2. The *external Pud'ic Arteries*, (F.) *Artères scrotales ou vulvaires* (Ch.), *A. honteuses externes*, which are two in number. The one is *superficial or subcutaneous*, and the other *deep-seated or subaponeurotic*. They are given off from the femoral artery, or some one of its branches; and are distributed to the external parts of generation.

PUDIC NERVE, (F.) *Nerf honteux ou ischio-pénien* on *Ischio-clitorien*, is a branch furnished by the sacral plexus, which is distributed to the genital organs; and, especially, to the penis in man, and the clitoris in woman.

PUELLA, Girl.

PUELLUS, (contracted from *puerulus*,) A little boy.

PUER, Boy, Child, Infans.

PUERA, Girl.

PUÉRILE, *Pueri'lis*, (from *puer*, 'a child,') (F.) *Puéril*. Appertaining or relating to childhood.

Respiration puérile ou exagérée ou supplémentaire ou hyperlésiculaire:—a term applied to the respiration when it is heard loud through the stethoscope, as in healthy children.

PUERILITAS, Dementia of the aged.

PUERITIA, see *Age*.

PUER'PERA, (*puer*, and *parere*, 'to bring forth,') *To'kas*, *Lo'chos*, *Enix'a*, *Le'cho*, a *Lying-in woman*, (F.) *Femme en Couche*, *Accouchée*. A female in child-bed. One recently delivered. The condition of a woman, recently delivered, is termed, in France, *Puerpéralité*.

PUER'PERAL, *Puerpera'lis*. Relating to parturition and its consequences.

PUERPERAL CONVULSIONS, *Eclampsia gravidarum et parturientium*.

PUERPERAL FE'VER. Any fever occurring during the puerperal state—*Febria puerpera'rum*, *Lochop'yra*, *Lochica'itis*, *Lochop'yra*, *Lochop'yra*, *Me'tria*, *Child-bed fever*. Most commonly restricted to a malignant variety of peritonitis.

PUERPÉRALITÉ, see *Puerpera*.

PUERPERIUM, Child-bed state.

PUERPEROUS, Parturient.

PUFF, ([D.] *puffen*, 'to blow,') *Lycoperdon*.

PUFFBALL, *Lycoperdon*, *Leontodon taraxacum*.

PUFF VEILED, see *Cavernous respiration*.

PUFFINESS, *Inflatio*, *Insufflatio*, *Tumefactio mol'lis*, (F.) *Bouffissure*, *Bouffissure*. A soft tumescence, without redness, which may be partial or general, and be produced by accumulation of serous fluid, extravasation of blood, effusion of air into the subcutaneous areolar tissue, &c. Puffiness is most frequently seen in the face and legs; and sometimes appears around a wound.

PUGILLUS, (dim. of *pugna*, 'the fist,') *Drac*, *Drac'hion*, *Drac'ma*, *Drac'mis*, (F.) *Pinci-*

The quantity of any substance, which may be taken with the ends of two or three fingers. The eighth part of a handful.

PUGNUS, Manipulus.

PUTEETH, Canine teeth.

PUISSANCE DU MOULE INTÉRIEUR, 'power of the internal mould,' Plastic force.

PUKE, Emetic.

PUKEWEEB, Lobelia inflata.

PUKING, Vomiting, Regurgitation.

PULEGIUM, (from *pulex*, 'a flea,' to which the odour is disagreeable,) *Mentha pulegium*—p. Cervinum, *Mentha cervina*.

PULEX, gen. *Pu'licis*, *P. Irritans*, *P. Vulgaris*. The common flea, (F.) Puce. A parasitic insect found on the bodies of man and animals.

PULEX PENETRANS, *Chique*—p. *Vulgaris*, *P. Irritans*.

PULICARIA DYSENTERICA, (from *pulex*, 'a flea,' to which the odour is disagreeable,) *Inula dysenterica*.

PULICARIS, *Pulicula'ris*, (F.) *Pulicaire*. An epithet given to cutaneous eruptions resembling flea-bites;—and, also, to diseases in which such eruptions occur.

PULICARIS HERBA, *Plantago psyllium*.

PULLEY, Trochlea.

PULL'NA, PIL'NA OR BIL'NA, MIN'ERAL WATERS OF. These well known Bohemian bitter springs contain sulphates of soda and magnesia, and chloride of magnesium.

PULLULAT'ION, *Pullula'tio*, (from *pullulare*, *pullulatum*, 'to bud,' 'budding,') A morbid production, which buds, as it were,—condyloma, for example.

PUL'MO, gen. *Pulmo'nis*; (by transposition of the letter *l* from *pleumon*.) *Pneum'on*, *Pleu'mon*, *Spiramen'tum ani'mæ*, *Ergaste'ri'on spir'itis*, *Fla-bel'lum et ventila'brum cor'dis*, Lights. The essential organ of respiration, which is double, and occupies the two sides of the chest. The lungs, which are of a spongy, soft, flexible, compressible, and dilatable structure, fill exactly the two cavities of the thorax, and are separated from each other by the mediastinum and the heart. The right lung, which is shorter and broader than the left, is divided, by two oblique fissures, into three unequal lobes, *Lobi seu Alæ pulmo'num*. The left has only two lobes, and consequently only one fissure. At the internal surface of those organs, which is slightly concave, there is, about the middle, a pedicle, formed by the bronchia and pulmonary vessels, and called, by anatomists, the *root of the lungs*, (F.) *Racine des Pouxmons*. Essentially, the lungs are composed of prolongations and ramifications of the bronchia, and of the pulmonary arteries and veins,—the divisions of which are supported by a fine, areolar tissue.

When the surface of the lungs is examined in a clear light, we may see, even through the pleura, that their parenchyma is formed by the aggregation of a multitude of small vesicles, of an irregularly spheroid or ovoid shape, full of air, and separated by white and opaque septa, constituting lozenge-shaped spaces, which are called *lob'ules*—*lob'uli seu in'sulæ pulmona'les*—and which are separated by interlobular areolar tissue. These lobules do not communicate with each other. The vesicles are called *Aircells*, *Air or Lung Ves'icles*, *Spiramen'ta seu Spiram'ina seu Cell'ule Pulmo'num*. See Cellules, bronchio. They, who regard the bronchial tubes as terminating in elongated cavities, have termed those cavities *Airaces*, *Infundib'ula*, *Malpigh'ian ves'icles*, *Ter'minal cav'ities*, &c., and the cuplike cavities, observed in these, have been called *alve'oli*. They are the aircells.

Along the partitions or septa is deposited, in greater or less quantity, *black pulmonary matter*, (F.) *Matière pulmonaire noire*, as it has been called,

which seems to be normal. Sometimes it is seen in points; at others, in spots. The colour of the lungs varies according to age and other circumstances. In youth, it is more red; and afterwards, grayish or bluish; often, as if marbled. The pleura pulmonalis is their investing membrane. The air is carried to the lungs by means of the trachea and bronchia. The black venous blood, which requires oxygenation, is conveyed to them from the heart by the pulmonary artery; and when it has undergone this change, it is returned to the heart by the pulmonary veins. The blood-vessels inservient to the nutrition of the lungs are the bronchial arteries. The pulmonary lymphatics are very numerous. Some are superficial; others, deepseated. They pass, for the most part, into the bronchial ganglions or glands. The nerves are furnished by the pulmonary plexus.

PULMOMETER, (*pulmo*, and *μετρον*, 'a measure,') Spirometer.

PULMOMETRY, see Spirometer.

PULMONAIRE, Pulmonaria, Pulmonary—p. *en Arbre*, *Lichen pulmonaria*—p. *de Chêne*, *Lichen pulmonaria*—p. *des Français*, *Hieracium murorum*.

PULMONAL, see Pulmonary.

PULMONARIA, *P. Officinalis* seu *Maculata* seu *Maculo'sa*, *Lung'wort*, *Spot'ted Lung'wort*, *Jeru'salem Cow'slip*, *Jeru'salem Sage*, *Spot'ted Com'frey*, (F.) *Pulmonaire*, *Sauge de Jerusale'm*. Ord. Boraginaceæ. Sex. Syst. Pentandria Monogynia. The leaves of this plant have been used as demulcents and pectorals; partly from their resemblance to lungs—hence the name, from *pulmo*, gen. *pulmonis*, 'the lung'—and partly because they have a slightly mucilaginous taste. The plant is not worthy of notice.

PULMONARIA, Pneumonia—p. *Arborea*, *Lichen pulmonarius*—p. *Gallica*, *Hieracium murorum*—p. *Officinalis*, *Pulmonaria*—p. *Virginica*, *Mertensia Virginica*.

PUL'MONARY, *Pulmona'ris*, *Pulmona'lis*, (F.) *Pulmonaire*, (from *pulmo*.) That which relates or belongs to the lungs.

Son pulmonal (F.), 'pulmonal or pulmonary sound,' is the sound rendered by the lung on percussion. It means, also, a similar sound rendered on percussing other parts.

PULMONARY ARTERY, *Arte'ria pulmona'ris*, *Ve'na arterio'sa*, arises from the upper and left part of the right ventricle of the heart; whence it passes obliquely upwards and to the left side; dividing into two trunks, opposite the second dorsal vertebra—one passing to each lung. At its origin, it is furnished with three sigmoid or semilunar valves. It carries the blood to the lungs for aeration.

PULMONARY MATTER, BLACK, see Pulmo.

PULMONARY PLEX'US, *Plex'us pulmona'ris*. A considerable nervous network, situate behind the bronchia, which is formed by the filaments of the pneumogastric nerve or par vagum, and by other filaments from the inferior cervical ganglion and the first thoracic ganglia. This plexus forms a kind of areolar web, with numerous meshes of different sizes, in which vessels are lodged, surrounded by much areolar tissue. The plexus sends filaments to the bronchia, which they accompany to their extremities.

PULMONARY VEINS, *Ve'næ pulmona'res*, *Arte'riæ veno'sæ*. These arise in the lungs from the final terminations of the arteries of the same name; collect in ramusculi and rami of greater and greater size, and ultimately unite in four trunks, two of which issue from each lung to open into the left auricle of the heart. The pulmonary veins have the same structure as the other veins of the body; from which they differ, in conveying arterial blood, as the pulmonary artery differs

from the others in conveying venous blood. They carry the blood back to the heart after aeration.

PULMONES TARTARIZATI, Lithiasis pulmonum.

PULMONIA, (from *pulmo*.) Phthisis pulmonalis, Pneumonia.

PULMONIC, *Pulmon'icus*, *Pneumon'icus*, *Pulmonarius*, *Pneumon'ic*, (F.) *Pulmonique*, *Pneumonique*. One affected with phthisis—one labouring under disease of the lungs. Any thing relating to or belonging to the lungs. A medicine for the lungs.

PULMONITIS, (*pulmo*, and *itis*.) Pneumonia.

PULP, *Pul'pa*, *Pulpa'men*, *Sarc*, (F.) *Pulpe*, *Pulpolite*. The soft and parenchymatous parts of vegetables reduced to a paste by the operation of pulping. Applied, also, to parts of the human body, which have the characters of, or resemble, a pulp.

PULP OF THE FINGERS, (F.) *Pulpe des doigts*. The fleshy extremity of the fingers.

PULP OF THE TOOTH, Dental pulp.

PULPA, *Pulp*—*p. Cassia*, see *Cassia fistula*—*p. Dentis*, Dental pulp—*p. Pili*, see *Hair*—*p. Pruni*, see *Pruni pulpa*—*p. Prunorum sylvestrium condita*, *Conserva pruni sylvestris*—*p. Tamarindi*, see *Tamarindi pulpa*.

PULPA TESTIS, *Par'en'chyma testis*. The parenchyma of the testis.

PULPAMEN, *Pulp*.

PULPATION, *Pulping*.

PULPE, *Pulp*—*p. Cérébrale*, see *Cerebrum*—*p. Dentaire*, Dental pulp—*p. des Doigts*, *Pulp* of the fingers.

PULPEUX, *Pulpy*.

PULPEZIA, *Apoplexy*.

PULP'ING, *Pulpa'tion*, *Pulpa'tio*, (from *pulp*.)

An operation, the object of which is to reduce certain vegetable substances to a pulp. For this purpose, they are bruised in a mortar, and then passed through a hair sieve, by the aid of a wooden spatula, called a *Pulper*, (F.) *Pulpoire*.

PULPOIRE, see *Pulping*.

PULPOLITE, *Pulp*.

PULP'Y, *Pulpo'sus*, (from *pulpa*.) (F.) *Pulpeux*. Full of pulp.

PULQUE, see *Agave Americana*.

PULS EX OVIS et **LACTE**, 'pap of eggs and milk.' *Oogala*.

PULSATIE, (from *pulsare*, 'to beat,') *Throbbing*.

PULSATILLA NIGRICANS, *Anemone pratensis*—*p. Patens*, *Anemone patens*—*p. Pratensis*, *Anemone pratensis*—*p. Vulgaris*, *Anemone pulsatilla*.

PULSATILLA NOIRE, *Anemone pratensis*—*p. des Prés*, *Anemone pratensis*.

PULSATIO, (*pulsare*, *pulsatum*, 'to agitate,') *Pulsation*—*p. Cordis*, *Palpitation*.

PULSA'TION, *Pulsa'tio*, *Sphyg'mus*, *Sphyx'is*, *Pede'sis*, *Pedeth'mos*, (from *pulsare*, *pulsatum*, 'to beat,') (F.) *Battement*. The beating of the heart and arteries. Also, the beating felt in an inflamed part.

PULSATIVUS, *Throbbing*.

PULSATORY, *Throbbing*.

PULSE, *Pul'sus*, *Pulsa'tio*, *Pul'sus* seu *Ictus cordis et arteriarum*, *Incur'sus arteria'rum*, *Sphyg'mus*, *Sphyg'ma*, *Diasphyx'is*, (F.) *Pouls*, (from *pellere*, *pulsam*, 'to beat,') *Beating of the arteries*, produced by the afflux of the blood propelled by the heart in its contractions. For the sake of convenience, we choose the radial artery at the wrist to detect the precise character of the pulse. These characters relate to the number of pulsations in a given time; to their degrees of quickness; hardness and strength; to the equality or inequality either of the pulsations themselves or of their intervals; to the development or fulness of the artery, and to the different impressions it may produce on the finger.

The following estimates have been made of the pulse at various ages :

AGES.	NUMBER OF PULSATIONS PER MINUTE.		
	HEBERDEN.	SÖMMERING.	MÜLLER.
In the embryo.....			150
At birth.....	130 to 140		Do.
One month.....	120		
One year.....	120 to 108	120	115 to 130
Two years.....	108 to 90	110	100 to 115
Three years.....	90 to 80	90	90 to 100
Seven years.....	72		85 to 90
Twelve years.....	70		
Puberty.....		80	80 to 85
Adult age.....		70	70 to 75
Old age.....		60	50 to 65

It does not appear that the pulse of the aged is less frequent, but the contrary. In general it is more frequent in females and in irritable persons than in males and those of an opposite temperament. In the healthy state, the pulse, besides having the proper number of pulsations, is neither hard nor unusually soft; it is moderately developed; of a medium strength; and perfectly equal, both as regards the pulsations and their intervals. In disease it wanders, more or less, from these physiological conditions. The different characters of the pulse that have been recorded are remarkably numerous. In a dictionary, it is necessary to detail them, although many of them are not now regarded, and some are ridiculous.

PULSE, **AR'DENT**, *Pul'sus ar'dens*, (F.) *Pouls ardens*, *Pouls élevé*. One which seems to raise itself to a point in order to strike the finger.

PULSE BREATH, see *Breath*, *pulse*.

PULSE, **CAP'RIZANT**, *Pul'sus Capri'zans* seu *dorcadis'sans* seu *gazel'lans* seu *fusa'tis*, *Goaleap pulse*, (F.) *Pouls caprisant*. One which seems to leap, like a goat, as it were; an imperfect dilatation of the artery being succeeded by a fuller and a stronger one.

PULSE, **COM'PLEX**, (F.) *Pouls combiné* ou *composé*, *P. compliqué*. A pulse which has the character of several critical pulses united.

PULSE, **CONTRACT'ED**, (F.) *Pouls serré* ou *concentré*. One whose pulsations are narrow, deep, and slightly hard. See *Concentration*.

PULSE, **CONVULSIVE**, *Pul'sus convulsivus*, (F.) *Pouls convulsif*. That which is unequally frequent or unequally hard; in short, which resembles the irregular pulsations of a muscular fibre in a state of convulsion.

PULSE, **CRIT'ICAL**, *Pulsus critic'us*. A pulse,

which, after having possessed the characters of irritation, becomes free, open, soft, &c.

PULSE, DECURTATE, *Pulsus myurus*: see, also, *Myurus*.

PULSE, DEEP, *Pulsus profundus*, (F.) *Pouls profond*. One which cannot be felt without difficulty, and without strong pressure by the finger.

PULSE, DEFICIENT, *Pulsus deficiens seu animum et circumveniens*, P. *Flickering*, (F.) *Pouls défaillant*, P. *insensible*. One which seems, every instant, about to cease.

PULSE, DEPRESSED, *Pulsus depressus*, (F.) *Pouls déprimé*. That which is deep and weak.

PULSE, DEVELOPED, (F.) *Pouls développé*. That which is broad, full, strong, and frequent.

PULSE, DICROTIC, *Pulsus Dicrotus seu bi-crotus seu intestinalis seu bifidus seu malleatus seu inter-crotus seu duplex*, (F.) *Pouls dicrote ou rebondissant ou redoublé*. That in which the finger is struck twice at each pulsation; once lightly, the other time more strongly.

PULSE, EQUAL, *Pulsus equalis*. That in which the pulsations resemble each other, and return at equal intervals.

PULSE, FEBRILE, *Pulsus febrilis*. One very frequent.

PULSE, FEBRILE, P. *Weak*, *Pulsus debilis*, (F.) *Pouls faible*. One which strikes the finger feebly.

PULSE, FILIFORM, P. *thready*. One which resembles a thread slightly vibrating.

PULSE, FLICKERING, Pulse, deficient.

PULSE, FORMICANT, *Pulsus formicans*, (F.) *Pouls formicant*. One which resembles, by being small and scarcely perceptible, the motion of an ant.

PULSE, FREQUENT, (F.) *Pouls agité ou précipité*. One which strikes more frequently than usual in a given time.

PULSE, FULL, *Pulsus plenus*, (F.) *Pouls plein*. One which gives the notion of great repletion in the vessel.

PULSE, GOATLEAP, P. *Caprizant*.

PULSE, GUTTURAL, *Pulsus gutturalis*, (F.) *Pouls guttural*. A pulse which has been supposed to indicate the termination of affections of the throat.

PULSE, HARD, *Pulsus durus*, (F.) *Puls dur*. That in which the pulp of the finger seems to yield to the pulsation.

PULSE, HECTIC, *Pulsus hecticus*. The weak, frequent pulse observed in hectic fever.

PULSE, HEMORRHOIDAL, (F.) *Pouls des Hémorroides*. A critical pulse, announcing the appearance of the hemorrhoidal flux. It is described as unequal;—the pulsations not being alike in force, and still less the intervals.

PULSE, HEPATIC, (F.) *Pouls du Foie*. A pulse imagined to indicate affections of the liver. It is not worthy of description.

PULSE, INFERIOR, (F.) *Pouls inférieur ou abdominal ou ventral*. One supposed to announce evacuations below the diaphragm.

PULSE, INTERCURRENT, *Pulsus intercurrents seu inter-cidens*. One in which a superfluous pulsation seems to occur from time to time.

PULSE, INTERMITTENT, (F.) *Pouls intercadent*. One in which the pulsations fail from time to time.

PULSE, INTES'TINAL, (F.) *Pouls des intestins*, P. *de la diarrhée*. A critical pulse, presumed to announce an evacuation by the intestines.

PULSE, IN'TRICATE, *Pulsus intricatus*, (F.) *Pouls embarrassé*. That which is but little developed and unequally slow.

PULSE, IRREGULAR, *Pulsus anormalis*. One whose pulsations are unequal, and return at unequal intervals.

PULSE OF IRRITATION, *Pulsus ab irritatio[n]e*

seu nervinus seu acriticus. One which is frequent, hard, and contracted.

PULSE, JARRING, P. *Vibrating*—p. *Jerking*, Pulse, sharp.

PULSE, LASS'GUD, (F.) *Pouls languissant*. A slow and feeble pulse.

PULSE, LARGE, *Pulsus crassus seu magnus*, (F.) *Pouls grand*, P. *large*. That which is large and full.

PULSE, LONG. One which strikes the finger to a great extent in length.

PULSE, LOW, *Pulsus humilis*, (F.) *Pouls bas*. One in which the pulsations are scarcely perceptible.

PULSE, NA'SAL, (F.) *Pouls nasal ou capital ou du Nez*. One supposed to announce the end of an irritation in the nasal fossæ.

PULSE, NAT'URAL, (F.) *Pouls naturel des adultes ou parfait ou des adultes ou modéré*. One that is equal and regular in strength and frequency; compressible, and devoid of hardness.

PULSE, ORGANIC, (F.) *Pouls organique*. One which indicates the affected organ, or the organ by which an evacuation will take place.

PULSE, PECTORAL, (F.) *Pouls de l'expectoration*. A critical pulse, announcing the termination of some pulmonary irritation.

PULSE, QUICK. One which strikes sharply, but not forcibly, against the finger.

PULSE, REGULAR, *Pulsus normalis seu regularis*. One whose pulsations are equal, and succeed each other at equal intervals.

PULSE, RESISTING, (F.) *Pouls résistant*. One a little tense, a little hard.

PULSE, SHARP, (F.) *Pouls prompt ou vite ou rif*. One which strikes the finger sharply, rapidly. When to a greater extent, it is called a *jerking* pulse.

PULSE, SHARPTAILED, *Pulsus myurus*: see *Myurus*.

PULSE, SIM'PLE, *Pulsus simplex*. A critical pulse, indicating an evacuation by one organ only.

PULSE, SLOW, *Pulsus tardus seu rarus*, (F.) *Pouls lent ou tardif ou rare*. One which strikes the finger slowly.

PULSE, SMALL, *Pulsus parvus*, (F.) *Pouls petit*. One whose pulsations are slender and weak.

PULSE, SOFT, *Pulsus molis*, (F.) *Pouls mou*. One which yields under the finger.

PULSE, STOM'ACHAL, (F.) *Pouls stomacal ou de l'estomac*. One supposed to announce an evacuation by the stomach.

PULSE, STRONG, *Pulsus fortis*, (F.) *Pouls fort*, P. *véhément*. One which strikes the finger strongly.

PULSE, SUPERIOR, *Pulsus Superior*, (F.) *Pouls supérieur*. A critical pulse, supervening on irritation of organs situate above the diaphragm.

PULSE, SUP'PLE, *Pulsus elasticus*, (F.) *Pouls souple*. One in which the pulsations have strength without hardness.

PULSE OF SWEAT, *Pulsus Sudorialis*, (F.) *Pouls Sudoral ou de la sueur*. A critical pulse, indicating an approaching perspiration.

PULSE, TENSE, P. *Cor'dy*, *Pulsus tensus*, (F.) *Pouls tendu*, P. *roide*. One in which the artery resembles a cord fixed at its two extremities. When resembling a wire, it is called *wiry*.

PULSE, THREADY, P. *filiform*.

PULSE, TREM'ULOUS, *Pulsus tremulus*, (F.) *Pouls tremblant ou palpitant*. One in which each pulsation oscillates.

PULSE, UN'DULATING, *Pulsus fluctuans seu fluens seu incidens Solani*, (F.) *Pouls undulant ou ondoyant*. One which resembles, in its movements, those of the waves.

PULSE, UNEQUAL, *Pulsus inæqualis*, (F.) *Pouls inégal*. One whose pulsations are not alike, or return at unequal intervals.

PULSE, URINAL, *Pulsus urinarius*, (F.) *Pouls*

de l'Urine. A critical pulse, fancied to denote an approaching evacuation of urine.

PULSE, UTERINE. *Pulsus uterinus*, (F.) *Pouls Uterine* on *de la Matrice* on *des Règles*. 'One which indicates the flow of the menses, or the occurrence of metrorrhagia.

PULSE, VAG'INAL. *Pulsus vagina'lis*. The pulsation of the arteria vaginalis, which is increased in pregnancy, during abortion, labour, &c.

PULSE, VENOUS. *Pulsus venarum*.

PULSE, VERMIC'ULAR. That which imitates the movement of a worm.

PULSE, VIBRATING. *Jar'ring Pulse*. One in which the artery seems to vibrate like a musical string.

PULSE, WIRY, see *Pulse*, tense.

PULSELESS, *Asphycticus*, Devoid of pulsation.

PULSELESSNESS, *Asphyxia*.

PULSEY, *Cataplasm*.

PULSIF'IC, *Pulsif'icus*, (*pulsus*, 'pulse,' and *facere*, 'to make,') That which causes or excites pulsation.

PULSILE'GIUM, *Pulsilo'gium*, *Pulsim'eter*, (*pulsus*, and *lego*, 'I read,') *Sphygmole'gium*, *Sphygman'eter*, *Sphygmoscop'ium*, *Sphyg'moscope*. Any instrument for measuring the quickness or force of the pulse.

PULSIMANTA, (*pulsus*, and *μαντα*, 'divination,') The art of predicting or prognosticating by the pulse.

PULSIMETER, (*pulsus*, and *μετρον*, 'measure,') *Pulsilegium*.

PULSUS, *Pulse*—*p. Acrificus*, *Pulse* of irritation—*p. Aequalis*, *Pulse*, equal—*p. Annuens* et *circumnuens*, *Pulse*, deficient—*p. Anormalis*, *Pulse*, irregular—*p. Biorotus*, *Pulse*, dierotic—*p. Biferiens*, *P. dierotic*—*p. Caprizans*, *Pulse*, caprizant—*p. Cordis*, see *Heart*—*p. Crassus*, *Pulse*, large—*p. Debilis*, *Pulse*, feeble—*p. Deficiens*, *Pulse*, deficient—*p. Dierotus*, *Pulse*, dierotic—*p. Doreadissans*, *Pulse*, caprizant—*p. Duplex*, *Pulse*, dierotic—*p. Durus*, *Pulse*, hard—*p. Elasticus*, *Pulse*, supple—*p. Fluctuosus*, *Pulse*, undulating—*p. Fluens*, *Pulse*, undulating—*p. Formicans*, *Pulse*, formicant, and *Pulse*, formicant—*p. Fortis*, *Pulse*, strong—*p. Fusalis*, *Pulse*, caprizant—*p. Gazellans*, *Pulse*, caprizant—*p. Heterochronicus*, see *Heterochronicus*—*p. Humilis*, *Pulse*, low—*p. Incidens* *Solani*, *Pulse*, undulating—*p. Interidens*, *Pulse*, intercurrent—*p. Intercisus*, *Pulse*, dierotic—*p. Intercurrens*, *Pulse*, intercurrent—*p. Intestinalis*, *Pulse*, dierotic—*p. Intricatus*, *Pulse*, intricate—*p. Magnus*, *Pulse*, large—*p. Mallearis*, *Pulse*, dierotic—*p. Mollis*, *Pulse*, soft.

PULSUS MYU'RUS, *Pulse* *decurtate* or *sharp-tailed*, (F.) *Pouls myure*. One which goes on diminishing. See *Myurus*.

PULSUS NERVINUS, *Pulse* of irritation—*p. Normalis*, *Pulse*, regular—*p. Parvus*, *Pulse*, small—*p. Plenus*, *Pulse*, full—*p. Rarus*, *Pulse*, slow.

PULSUS SERRIN'US, (F.) *Pouls serrin* on *serratile*. One which strikes, sometimes strongly, at others feebly, and recalls the idea of a saw.

PULSUS STRIA'TUS, (F.) *Pouls étroit*. One that strikes the finger to a small extent in breadth.

PULSUS SUDORALIS, *Pulse* of sweat—*p. Tardus*, *Pulse*, slow—*p. Urinalis*, *Pulse*, urinal—*p. Uterinus*, *Pulse*, uterine.

PULSUS VAC'UUS, (F.) *Pouls vide*. One which gives the idea of emptiness of the artery.

PULSUS VAGINALIS, *Pulse*, vaginal.

PULSUS VENA'RUM, *Venous Pulse*, *Phlebopaltia*. A pulsation occasionally noticed in the jugular veins. It is owing to some of the blood of the right ventricle reflowing into the right auricle, and communicating an impulse to the venous blood arriving in the auricle through the descending cava. It is often a sign of impediment to the flow of blood along the pulmonary artery.

PULTA'CEOUS, *Pulta'ceus*, (F.) *Pultacé*, (*puls*,

gen *pultis*, 'pap,') Resembling pap. Having a softened, semifluid appearance.

PULTATIO, (*puls*, gen. *pultis*, 'pap,') *Atheroma*. **PULTISE**, *Cataplasm*.

PULTICULA, (dim, of *puls*, gen. *pultis*, 'pap,') *Bouillie*.

PULVERES AEROPHORI, *Powders*, *soda*—*p. Effervescentes*, *Powders*, *soda*.

PULVERIZABLE, *Pul'verable*. (From *pulvis*, gen. *pulveris*, 'powder,' and *habilis*, 'able,') Capable of being reduced to powder.

PULVERIZATION, *Pulverisa'tio*, *Pulveriza'tio*, *Coniosto'is*, (F.) *Pulvérisation*. The operation of dividing substances, so as to reduce them to powder.

Most drugs are reduced to powder by persons who make it a special business, and who are called *drug grinders*. This is done on a large scale in *drug mills*; and the implements used are the *grinding mill* and the *stamping mill*. In the shop of the apothecary, pulverization is effected by the pestle and mortar, and the sieve.

Where substances require the addition of other bodies to facilitate their reduction to powder,—as where camphor needs a few drops of alcohol—the process is called *pulverization by mediation*.

PULVERMACHER'S CHAIN BATTERY, see *Galvanism*.

PULVÉROLÉS, see *Pulvis*.

PULVERULENT, *Pulverulentus*, (from *pulvis*.) Reduced to the state of a fine powder. Covered, as it were, with dust. An epithet applied to the eyes when they appear dirty, and, as it were, sprinkled with dust, by reason of granulations or grayish striae, resulting from thickening of the mucous fluid of the conjunctiva.

PULVILLUS, *Pad*, *Bourdonnet*—*p. e* *Lina-mentis confectus*, *Bourdonnet*—*p. Rotundus*, *Bourdonnet*.

PULVINAR, *Pulvina'rium*. A pillow.

PULVINAR HUMULI, see *Humulus lupulus*.

PULVINAR MEDICATUM seu *MEDICINA'LE*, *Sacculus medicina'lis*. A medicated pillow; as of hops.

PULVINAR PARVUM, *Pad*—*p. Ventriculi*, *Pancreas*.

PULVIS, gen. *Pul'veris*, *Co'nis*, *Por'ider*, (F.) *Poudre*. A substance reduced into small particles by pulverization. *Simple powders* are those of a single substance; *compound*, those which result from a mixture of several different powders. An *impalpable powder* is one which is so fine that it can scarcely be felt under the finger. Compound powders have been called *Spécies*, (F.) *Espèces*, *E. pulvérentes*, *Pulvérolés*.

PULVIS ABSORBENS, *P. de magnesiâ compos.*—*p. Adpersorios*, *Catapasma*—*p. Aerophorus*, *Powders*, *soda*—*p. Alexiterius*, *P. contrayervæ comp.*—*p. Alexiterius*, *P. ipecacuanhæ compositus*—*p. Algarothi*, *Algaroth*.

PULVIS AL'OE'S COMPOSITUS, *P. Aloës cum Guai'aco*, *P. Aloëticus cum Guaiaco*, *Compound powder of Aloës*. (*Aloës*, ʒiiss, *Guaiac. res.* ʒj. *Pule. cinnum. comp.* ʒss. Rub the aloes and guaiacum separately, then mix the whole. *Ph. L.*) Cathartic, stomachic, and sudorific. Dose, gr. x to ʒj.

PULVIS AL'OE'S CUM CANELL'IA, *P. aloës et canellæ*, (*Ph. U. S.*) *P. Aloëticus*, *Powder of Aloës with Canella*, *Spécies Hérois p'era*, *Hérois p'era*, vulg. *Hérois p'ery*, *Hérois p'ery*, *P'era*, and *P'ery*. (*Aloës* ʒij, *Canell.* ʒij. Rub separately into a powder, and mix. *Ph. U. S.*) A warm cathartic. Dose, gr. x to ʒj.

PULVIS AL'OE'S CUM GUAIA'CO, *P. A. compositus*—*p. Aloeticus*, *P. aloës cum canellâ*—*p. Aloeticus*, *Hiera piera*—*p. Aluminis compositus*, *P. sulphatis aluminæ compositus*—*p. Aluminis kinosatus*, see *P. sulphatis aluminæ compositus*—*p. Aluminosus*, *P. sulphatis aluminæ compositus*.

PULVIS DE AMARIS COMPOSITUS. *P. antiarthriticus amarus*, (F.) *Poudre antiarthritique amère*. (Gentian., Aristoloch. rotund., $\bar{a}\bar{a}$ p. 3, *Flor. centaur. min.* p. 4, *Fol. chamædr.*, *Chamæpith.* manip. — *f. pulvis*, *Ph. P.*) In gouty affections, dyspepsia, &c. Dose, gr. xx.

PULVIS AMYGDALÆ COMPOSITUS, *Confectio amygdalarum* — *p. Angelicus*, *Algaroth* — *p. Anodynus*, *P. ipecacuanhæ compositus* — *p. Antacidus*, *P. cretæ compositus* — *p. Antiacidus*, *P. de magnesiâ compositus* — *p. Antiarthriticus amarus*, *P. de amaribus compositus* — *p. Antiarthriticus purgans*, *P. de sennâ, scammonæâ et lignis* — *p. Antiasthmaticus*, *P. de sulphure et scillâ*.

PULVIS ANTILYSÆSUS, *Mead's powder against the bite of a mad dog*. It consisted of ash-coloured liverwort in powder, with an equal quantity of black pepper; a drachm and a half of the two forming a dose for an adult.

PULVIS ANTIMONIALIS, *Antimonial powder* — *p. Antimonii compositus*, *Antimonial powder*.

PULVIS DE AËRO COMPOSITUS, *Compound Powder of Aërum*, (F.) *Poudre de Birkmann*, (*Rad. ari.*, *Acori veri*, *Pimpinell. saxifrag.* $\bar{a}\bar{a}$ 40 p., *Ocul. cancer.* ppt. 12 p., *Cinnam.* 7 p., *Potassæ sulph.* 6 p., *Ammonizæ muriat.* 2 p. — *f. pulv.* *Ph. P.*) *Sternutatory*.

PULVIS AROMATICUS, *p. cinnamomi compositus*.

PULVIS ASARI COMPOSITUS, *P. sternutatorius seu errhinus seu capitalis Sancti Angeli seu cephalicus*, *Cephalic snuff*, *Compound powder of Asarabacca*, (F.) *Poudre d'asarum composée dite sternutatoire*, *P. de Saint-Ange*. (*Asar. Europ.* p. 3, *Origan. majoran.*, *Lavand.*, sing. p. 1. Rub into powder. *Ph. L.*) *Sternutatory*.

PULVIS AURI, see *Gold* — *p. e. Bolo compositus cum opio*, *P. cretæ compositus cum opio* — *p. e. Bolo compositus sine opio*, see *P. Cretæ compositus* — *p. Cambogiæ compositus*, *P. e. gummi guttâ* — *p. Capitalis Sancti Angeli*, *Pulvis asari compositus* — *p. Carbonatis calcis compositus*, *P. cretæ compositus* — *p. Cardiacus*, *P. Contrayervæ comp.* — *p. Carthusianorum*, *Antimonii sulphuretum rubrum*.

PULVIS CATECHU COMPOSITUS, *Compound Powder of Catechu*. (*Catechu, kino*, $\bar{a}\bar{a}$ $\bar{z}$ ij; *Cinnam.*, *Myrist.*, $\bar{a}\bar{a}$ $\bar{z}$ ss.) Dose, as an astringent, gr. xv to xxx.

PULVIS CATHARTICUS, *P. Scammonææ compositus* — *p. Causticus*, *Vindobonensis*, *Powder*, *Vienna* — *p. Cephalicus*, *P. asari compositus*.

PULVIS CINNAMOMI COMPOSITUS, *Compound Powder of Cinnamon*, *Pulvis seu Species Imperatoris*, *Pulvis aromaticus* (*Ph. U. S.*), *P. vitæ Imperatoris seu Diaromatton*, *Species diacinnamomi seu latificantes Rhazis seu Diambræ si nê odoratis seu aromaticæ*, *Tragæa aromatica*, *Aromatic powder*, (F.) *Poudre de canelle composée ou aromatique*, (*Cinnam.*, *Zingib.* $\bar{a}\bar{a}$ $\bar{z}$ ij, *Cardam.*, *Myrist.* $\bar{a}\bar{a}$ $\bar{z}$ j. Rub to a fine powder. *Ph. U. S.*) *Stimulant and carminative*. Dose, gr. x to $\bar{z}$ j.

PULVIS COBBII seu TUNQUINENSIS. The famous *Tonquin powder*, introduced into England by a Mr. Cobb. It was composed of 16 grains of musk, and 48 grains of cinnabar; and was mixed in a gill of arrack. It was formerly regarded as a specific in hydrophobia.

PULVIS COMITIS WARWICENSIS, *P. Cornachini* — *p. Comitissæ*, *Cinchona*.

PULVIS CONTRAYERVÆ COMPOSITUS, *P. cardifolius seu alexiterius seu Mantanus*, *La'pis contrayervæ*, *Contrayerva balls*, *Compound Powder of Contrayerva*, (*Contrayerv. contrit.* $\bar{z}$ v, *Test. præp.* libss. *Mix.* *Ph. L.*) *Stimulant and sudorific*. Dose, gr. xv to $\bar{z}$ ss.

PULVIS CORNACHINI, *P. Oc'yd'i Stib'ii compositus seu Comitissæ Warwicensis seu de Tri'bus seu Scammonii antimonialis*, *Diaceclat'son Porac'lei*, *Earl of Warwicke's Powder*, *Cerberus*

Tri'ceps, (F.) *Poudre Cornachine*. (*Scammon.*, *Potass. superart.*, *Antim. diaph.* $\bar{a}\bar{a}$ p. æ. f. *pulvis*, *Ph. P.*) *Cathartic*. Used chiefly in cutaneous diseases. Dose, $\bar{z}$ ss to $\bar{z}$ j.

PULVIS CORN'US US'TI CUM O'PIO, *P. opiat'us*, *Powder of burnt Hartshorn with opium*, (F.) *Poudre de corne brûlée avec l'opium*, (*Op'ii duri* cont. $\bar{z}$ j. *Corn. ust. præp.* $\bar{z}$ j. *Coccor.* cont. $\bar{z}$ j. *Mix.* *Ph. L.*) Ten grains contain one of opium. *Anodyne*.

PULVIS COSMI, *Powder*, *arsenical*, of *Cômpe* — *p. Cretaceus*, *P. cretæ compositus*.

PULVIS CRETÆ AROMATICUS, see *Confectio aromatica*.

PULVIS CRETÆ COMPOSITUS, *P. Carbonatis Calcis compositus seu cretæceus seu antac'idus seu terrest'ris*, *Species e Scord'io si nê o'pio*, *S. Diacret'ie*, *Compound Powder of Chalk*, *Dioscor'idium*; (with *Bole Armeniæ* in place of chalk — the *Pulvis e Bolo compositus sine opio*;) (F.) *Poudre de craie composée*. (*Cretæ ppt.* libss. *Cinnam.* $\bar{z}$ iv, *Tormentill.*, *Acacis*, sing. $\bar{z}$ ijj, *Pip. long.* $\bar{z}$ iv. Reduce separately to a fine powder, and mix. *Ph. L.*) *Antacid*, *stomachic*, *absorbent*. Dose, gr. xx to $\bar{z}$ j.

PULVIS CRETÆ COMPOSITUS CUM O'PIO, *Compound Powder of Chalk with opium*, *P. e Cretâ compositus cum opio*; (with *Bole Armeniæ* instead of chalk — *P. e Bolo compositus cum opio*;) *Species e Scord'io cum opio*, (F.) *Poudre de craie composée avec l'opium*. (*Pulv. cret. comp.* $\bar{z}$ vjss, *Op'ii duri* cont. $\bar{z}$ iv. *Mix.* *Ph. L.*) Contains one grain of opium in 40. *Anodyne* and absorbent. Dose, $\bar{z}$ j to $\bar{z}$ ij.

The British Pharmacopœia (1864) has a *Pulvis Cretæ aromaticus cum opio*, *Aromatic powder of chalk and opium*, which is composed of aromatic powder of chalk $\bar{z}$ ix $\bar{z}$, opium, in powder, $\bar{z}$ 4. Mix and pass through a fine sieve. Dose, $\bar{z}$ ss to $\bar{z}$ j.

PULVIS DIAPHORETICUS, *P. ipecacuanhæ comp.* — *p. Diapnoicus*, *P. ipecacuanhæ comp.* — *p. Diaromatton*, *P. cinnamomi compositus* — *p. Diasenæ*, *P. sennæ compos.* — *p. Diaturpethi compositus*, *P. jalapæ compositus*.

PULVIS DU'GIS PORTLAND'IE, *Duke of Portland's Gout Powder*, (F.) *Poudre de Duc de Portland ou de la Mirandole*. (*Rad. aristoloch.*, *Gentianæ*, *Summit. chamædr.*, *Centaur. min.* $\bar{a}\bar{a}$ p. æ.) Dose, $\bar{z}$ ss to $\bar{z}$ j.

PULVIS EMMENAGOGUS, *P. myrrhæ compositus* — *p. Epilepticus Marchionum*, *P. de magnesiâ compositus* — *p. Errhinus*, *P. asari compositus* — *p. Febrifugus Jacobi*, *Antimonial powder* — *p. Galactopeus*, *P. de magnesiâ compositus*.

PULVIS E GUMMI GUTTÂ, *P. Cambogiæ compositus seu hydragogus*, *Camboge' Powder*, (F.) *Poudre hydragogue ou de gomme gutte ou de Quercetanus*. (*Rad. jalap.* 24 p., *Rad. mechoacan.*, 12 p., *Cinnam.*, *Rhei*, $\bar{a}\bar{a}$ 8 p., *Gum. Cambog.* 3 p., *Fol. soldanellæ* sic. 6 p., *Semin. anisi* 12 p., fiat pulvis. *Ph. P.*) *A drastic cathartic*. In anasarca and in worms. Dose, gr. xv to xxx.

PULVIS GUNMO'SUS ALKALI'NIS, *Sal'p'o vegetabilis*, (F.) *Poudre de savon végétal*. (*Guaïac. pulv.* 32 p. *Potass. carb.* 4 p. Make into a powder. *Ph. P.*) *Antacid*.

PULVIS EX HELMINTHOCORT'O COMPOSITUS, *P. vermifugus ab'sque mercurio*, (F.) *Poudre vermifuge sans mercure*. (*Fuc. helminthocort.*, *Sem. contra.*, *Summitat.*, *absinth.*, *Tanacet.*, *Fol. scordii*, *Sennæ*, *Rhei*, $\bar{a}\bar{a}$ p. æ, fiat pulvis. *Ph. P.*) Dose, $\bar{z}$ j to $\bar{z}$ j.

PULVIS HYDRAGOGUS, *P. e gummi guttâ* — *p. Hydrargyri cinereus*, see *Hydrargyri oxydum cinereum* — *p. Hypnoticus*, *Hydrargyri sulphuretum nigrum* — *p. Imperatoris*, *P. cinnamomi compositus* — *p. Incisivus*, *P. de sulphure et scillâ* — *p. Pro Infantibus*, *P. de magnesiâ compositus* — *p. Infantum*, *P. de magnesiâ compos.* — *p. Infantum antacidus*, *P. de magnesiâ compositus*.

PULVIS IPECACUANHÆ COMPOSITUS, *P. alexi-*
terius seu anodynus seu diaphoreticus seu diapo-
noicus seu emorificus Doveri seu Ipecacuanhæ
et Opii, Com'pound Powder of Ipecacuanha,
Dover's Pow'der, (F.) *Poudre de Dover ou d'ipe-*
cacuanha et d'opium ou de Dover, (*Ipecacuanha*
pulv., *Opii pulv.* sing. $\mathfrak{z}$ ij. *Potassæ sulphat.* $\mathfrak{z}$ ij.
Mix. Diaphoretic. Dose, gr. v to gr. xij.

PULVIS IPECACUANHÆ ET OPII, *P. ipecacuanhæ*
compositus—p. Jacobi, Antimonial powder.

PULVIS JALAPÆ COMPOSITUS, *P. Diatriptethi*
compositus seu laxans seu purgans seu laxativus
vegetabilis seu laxans cum sale seu Jalapæ salu-
mus seu Jalapæ tartariæ, Spécies Diatriptethi,
Com'pound Powder of Jalap. (F.) *Poudre de*
Jalap composée, (*Pulv. jalap.* $\mathfrak{z}$ ij. *Potassæ bitart.*
 $\mathfrak{z}$ ij. Mix.) Cathartic. Dose, $\mathfrak{z}$ ij to $\mathfrak{z}$ ij.

PULVIS JALAPÆ SALINUS, *P. jalapæ compositus*
—p. Jalapæ tartariæ, *P. jalapæ compositus*—p.
de Jalapæ et scammonio compositus, *P. scam-*
monæ comp.—p. Jamesii, Antimonial powder—p.
Jesuiticus, Cinchona—p. Joannis de Vigo, Hy-
drargyri nitricooxydum.

PULVIS KINO COMPOSITUS, *P. Kino cum o'pio*
(Ph. B. 1864.) *Com'pound Powder of kino*, (F.)
Poudre de kino composée, (*Kino* $\mathfrak{z}$ xv. *Cinnamon*, $\mathfrak{z}$ iv.
Opii duri $\mathfrak{z}$ ij. Rub each to powder, and mix.
Ph. L.) Astringent. Dose, gr. v to xx.

PULVIS LAXANS, *P. jalapæ compositus*—p.
Laxans cum sale, *P. jalapæ compositus*—p. *Lax-*
ativus vegetabilis, *P. jalapæ compositus*.

PULVIS DE MAGNESIÆ COMPOSITUS, *P. Galac-*
topæus seu Nutri'cum seu Infantum seu Infantum
anticid'idus seu pro Infantibus seu anticid'idus seu
absorbens seu epilepticus Marchionis, Com'pound
Pow'der of Magnesia, (F.) *Poudre absorbante et*
anti-acide, (*Magnesia*, *Sacchar*, $\mathfrak{a}\mathfrak{a}$ p. æ. Mix.
Ph. P.)

PULVIS MASTUANS, *P. contraryæ comp.*—
p. Marchionis, Marchionis Pulvis—p. Mercurialis
cinereus, Hydrargyri oxidum cinereum—p. ad
Mucum pulmonalem liberius ciendum, *P. de sul-*
phure et scilla.

PULVIS MYRRHÆ COMPOSITUS, *P. Myrrhæ*
compositus seu ad ulcera tend'ium seu emmena-
gorræ, Com'pound Myrrh Pow'der, (F.) *Poudre*
Balsamique, (*Myrrhæ*, *Sabina*, *Ruta*, *Castor*, *Russ*,
 $\mathfrak{a}\mathfrak{a}$ $\mathfrak{z}$ ij.) A stimulant, antispasmodic, and emme-
nagogue. Dose, gr. xv to xl.

PULVIS NUTRICUM, *P. de magnesiâ compositus*
—p. Opiatus, *P. cornu uti cum opio*—p. Oxydi
stibii compositus, *P. cornachini*—p. ad Partum,
Ergot—p. Parturiens, *Ergot*—p. de Phosphate
calcis et stibii compositus, Antimonial powder—p.
Principis Hydrargyri nitricooxydum—p. Pur-
gans, *P. jalapæ compositus*—p. Quercetanus,
Hydrargyri submuriæ.

PULVIS QUERCUS MARIÆ, *Pow'der of Yell'one*
Blad'derwrack, (The wrack in fruit, dried,
cleaned, exposed to a red heat in a crucible with
a perforated lid, and then reduced to powder.
Ph. L.) Virtues dependent on iodine: used, con-
sequently, in bronchitis, &c. Dose, gr. x to xl.
See *Fucus vesiculosus*.

PULVIS RHEI COMPOSITUS (Ph. U. S. 1863.)
Com'pound Powder of Rhu'barb, (*Magnes*, $\mathfrak{ij}$,
Zingib. pulv. $\mathfrak{z}$ ij. *Rhei pulv.* $\mathfrak{z}$ iv. M.) An antacid
and laxative. Dose, $\mathfrak{z}$ ij to $\mathfrak{z}$ ij.

PULVIS SALINUS COMPOSITUS, *Com'pound Sal-*
ine Pow'der, (*Sodii chlorid.*, *Magnesiæ sulphat.*,
 $\mathfrak{a}\mathfrak{a}$ $\mathfrak{z}$ ij. *Potassæ sulphat.* $\mathfrak{z}$ ij. Ph. E.) Dose, as an
aperient, two or three drachms dissolved in water.

PULVIS SCAMMONIÆ COMPOSITUS, *P. e Scam-*
monio compositus seu Scammonii seu de Jalap'æ et
Scammonio compositus dictus Cathar'ticus, Com-
pound Powder of Scammony, (F.) *Poudre cathar-*
tique ou Poudre de Scammonie composée, (*Scam-*
mon, *Ext. jalap. dur.*, sing., $\mathfrak{z}$ ij, *Zingib.* $\mathfrak{z}$ ss. Rub
separately to powder, and mix. Ph. L.) Cathar-
tic. Dose, gr. viij to gr. xv.

PULVIS SCAMMONII CUM AL'OE, *P. e Scammonio*
cum Al'oe, Scammony and Aloes Pow'der, (*Scam-*
mon, $\mathfrak{z}$ ij. *Ext. jalap. dur.*, *Aloës Sacchar*, $\mathfrak{a}\mathfrak{a}$ $\mathfrak{z}$ iss,
Zing. $\mathfrak{z}$ ss.) Cathartic. Dose, gr. x to xv.

PULVIS SCAMMONII ANTIMONIALIS, *P. corna-*
chini.

PULVIS SEN'NÆ COMPOSITUS, *P. e Sen'na compo-*
situs seu e Sen'na compositus seu Diac'na,
Com'pound Powder of Sen'na, (F.) *Poudre de séné*
composée, (*Senne*, *Potassæ superart.*, sing., $\mathfrak{z}$ ij,
Scammon, $\mathfrak{z}$ ss, *Zingib.* $\mathfrak{z}$ ij. Ph. L.) Cathartic,
hydragogue. Dose, $\mathfrak{z}$ ij to $\mathfrak{z}$ ij.

**PULVIS DE SENNÆ, SCAMMONIÆ ET LIC'NIS COM-
POSITUS dictus ANTIARTHRITICUS PUR'GANS, (F.)
Poudre de Pérad, (*Sen Arab.*, *Potassæ superart.*,
Fol. senne, *China*, $\mathfrak{a}\mathfrak{a}$ 4 p., *Scammon*, *rad.*
Sarsapar., *China*, *Guaic.* *lygn.* $\mathfrak{a}\mathfrak{a}$ 2 p.: fiat pulvis.
Ph. P.) Purgative. Used to obviate goat. Dose,
 $\mathfrak{z}$ ij.**

PULVIS STANNI, see Tin—p. Sternutatorius, *P.*
assri compositus—p. Stibiatu, Antimonial pow-
der—p. Stypticus, *P. sulphatis alumine compo-*
situs—p. Sudorificus, *P. ipecacuanhæ compositus*—
p. Sudorificus Doveri, *P. ipecacuanhæ compo-*
situs.

PULVIS DE SULPHA'TE POTASSÆ COMPOSITUS,
P. temperans Stahl'i, (F.) *Poudre tempérante de*
Stahl, (*Potassæ sulph.*, *P. nitrat.* $\mathfrak{a}\mathfrak{a}$ 9 p., *Hydragr.*
sulph. rubr. 2 p. Mix. Ph. P.) As an anti-
phlogistic, and in acidity of the first passages.
Dose, $\mathfrak{z}$ ij to $\mathfrak{z}$ ij.

PULVIS SULPHATIS ALUMINÆ COMPOSITUS,
P. aluminosus seu aluminis kinosus seu stypti-
cus seu aluminis compositus, Alu'men kinosu-
tum, Com'pound Alu'm Pow'der, (F.) *Poudre stypti-*
que ou d'alun composée, (*Sulphatis alumina*, p.
iv. *Kino*, p. 1. Ph. E.) Astringent. In intestinal
hemorrhage and diarrhoea. Dose, gr. x to xv.

PULVIS DE SULPHURÆ ET SUL'LI, *P. antia'sm-*
maticus seu incisivus seu ad m'cum pulmona'lem
liberius ciend'um, (F.) *Poudre incisive ou anti-*
asthmaticque, (*Sacchar*, p. 3, *Sulphur*, *lot*, p. 2,
Scilla, *siccata*, p. j. Ph. P.) Dose, gr. v to $\mathfrak{z}$ ss.

PULVIS E SULPHURE TO HYDRARG'YRI NIGRO
ET SCAMMONIO, dictus VERMIFUGUS MERCURIA-
LIS, (F.) *Poudre vermifuge mercurielis*, (*Pulv.*
cornachini, *Hydragr. sulphuret. nigr.* *recent. præp.*,
 $\mathfrak{a}\mathfrak{a}$ p. æ. Mix. Ph. P.) Dose, $\mathfrak{z}$ ij to $\mathfrak{z}$ ss.

PULVIS SYMPATHETICUS, Sympathetic powder
—p. Temperans Stahl'i, *P. de sulphate potassæ*
compositus—p. Terrestris, *P. cretæ compositus*—
p. Tonquinensis, *P. Cobbii*.

PULVIS TRAGACANTHÆ COMPOSITUS, *P. e Tra-*
gacanth'â compositus, Spécies Diatriptethi
frig'idæ, Com'pound Powder of Tragacanth, (F.)
Poudre de gomme adragant composée, (*Traga-*
canth. contus., *Acacia cont.*, *Ampli.* sing., $\mathfrak{z}$ iss,
Sacch. purif. $\mathfrak{z}$ ij. Ph. L.) Demulcent. In
coughs, &c. Dose, $\mathfrak{z}$ ss to $\mathfrak{z}$ ij.

PULVIS DE TRIBUS, *P. cornachini*—p. ad Ulcera
tendinum, *P. myrrhæ compositus*, *p. Vermifugus*
absque mercurio, Pulvis ex helminthocorto comp.
—p. Vermifugus mercurialis, *P. e sulphureto hy-*
drargyri nigro et scammonio—p. Vitæ Impera-
toris, *p. cinnamomi compositus*.

PUMEX, *La'pis pumiceus, Pum'ice*, (F.) *Pierre*
Ponce. This was formerly employed as a desicca-
tive and as a dentifrice.

PUMICE, *Pumex*.

PUMILIO, *Nanus*.

PUMILUS, see *Nanus*.

PUMMELFOOT, (*Knobfoot*,) *Kyllosis*.

PUMP, **BREAST**, *Anthia lactea*—p. *stomaen*,
Stomach pump.

PUMPKIN, (*pompion*,) *Cucurbita pepo*.

PUNA, (*S.*) *Veta*, *Sorocce*, *Sorocche*, *Sorocche*,
Marco, *Hypoc'ans*, *Orn'ans*, *Dysp'ia a vomit-*
ina, *Mountain Sickness*, (F.) *Mal de Montagne*.
A sickness common in the elevated districts of
South America and elsewhere, which attacks those

unaccustomed to breathe the rarified atmosphere. It strongly resembles sea-sickness, with a most acute and depressing throbbing in the head. It does not often affect the respiration greatly.

PUNAIS, (perhaps from *puer*, 'to stink,' and *nas*, 'the nose,' see *Ozana*.)

PUNAISE, Cimex.

PUNASIE, *Ozana*.

PUNCH, *Rhizagra*, (*pungere*, *punctum*, 'to prick,') *Repulsorium*, (F.) *Repoussoir*. A surgical instrument used for extracting the stumps of teeth. See *Lever*.

Also, an agreeable alcoholic drink, made of spirit, wine, sugar, lemon juice, and water. Milk is sometimes substituted wholly or in part for the water; the lemon juice being left out and nutmeg added: it then forms *Milk Punch*. *Punch*, as a drink, is said to be of oriental origin, and to be derived from Persian *punji* or Sanscrit *puncha*, 'five,' from the number of ingredients that entered into its composition [?].

PUNCTA, *Punctum*, *Punctus*, (*pungere*, *punctum*, 'to prick, to dot,') a point, a prick, a puncture.

PUNCTA OSSIFICATIONIS, Ossification, points of.

PUNCTÆ MUCOSÆ, see *Acne*.

PUNCTICULA, (dim. of *puncta*, 'a point,') *Petechiæ*.

PUNCTIO, (from *pungere*, *punctum*, 'to puncture,') *Paracentesis*, *Puncture*—p. *Nervi*, *Neurogyme*.

PUNCTUM, ('a point,') *Puncta*, *Puncture*, see *Stitch*.

PUNCTUM AURÆUM, *Punctura auræa*. Formerly, when an intestinal hernia was reduced by an incision made through the integuments down to the upper part of the spermatic vessels, a golden wire was fixed and twisted, so as to prevent the descent of anything down the tunica vaginalis. Hence the term. See, also, *Foramen centrale retinæ*.

PUNCTUM CÆCUM, 'blind spot,' see *Optic nerve*—p. *Lacrymale*, *Lachrymal punctum*.

PUNCTUM SALLIENS, 'leaping point,' *P. sanguineum saliens*, *Globulus sanguineus*, *Punctus ruber saliens*. The first moving point which makes its appearance after the fecundation of the germ. It has been generally, but erroneously, applied to the heart of the embryo.

PUNCTURA, *Paracentesis*, *Puncture*—p. *Aurea*, *Punctum aureum*.

PUNCTURE, *Punctura*, *Punctio*, *Cente'sis*, *Paracente'sis*, *Nyx'sis*, *Tre'sis* *Punctura*, (F.) *Figure*. A wound made by a pointed instrument, as a sword, bayonet, needle, thorn, &c. A puncture is, generally, of trifling consequence; but, sometimes, it becomes important from the parts implicated, and the after consequences.

PUNCTURE DOCTORS, *Piqueurs médecins*.

PUNCTURING, *Compunctio*, *Paracente'sis*, (F.) *Ponction*. The operation of opening, by means of a trocar, lancet, &c., any natural or accidental cavity, with the view of evacuating any fluid, effused or accumulated therein. *Paracente'sis* is a sort of puncturing.

PUNCTUS, 'a point,' *Puncta*—p. *Ruber Salliens*, *Punctum saliens*.

PUN'GENT, (F.) *Pongitif*. Pain is said to be pungent, when it seems as if a pointed instrument was forced into the suffering part.

PUNICA GRANATUM, (*Punica*, 'Punic, Carthaginian,' whose colors were a deep red, *garnet*.) The *Pomegranate*, *Rho'a*, *Rho'a*, *Granatum*, (F.) *Grenadier*: *Ord.* *Myrtacæ*. *Sex. Syst.* *Icosandria Monogynia*. The rind of the fruit, called *Malico'rimum*, *Malico'rimum*, *Granati Fructus Cor'tex* (Ph. U. S.); and the flowers—called *Balaustine flowers*—are used in medicine. They have been employed both internally and externally as astringents. The seeds are called

Cocco'nes. The portion, however, of the plant, that has attained most celebrity, is the bark of the root, *Granati Rad'icis Cor'tex* (Ph. U. S.), which appears to be very efficacious in tænia. It is given in decoction, (*Granat. rad. cort.* ʒij; *Aque Oij*, boiled to Oiss.) Dose, fʒij, every half hour. Three or four doses are usually sufficient to expel the worm.

The bark contains a peculiar matter called *Grenadin*.

PUNK, *Boletus ignarius*.

PUOGENIA, *Pyogenia*.

PUPIL, *Pupilla*, *Pupula*, *Pupillio*, *Vis'io*, *Op'sis*, *Co'rë*, *Co'ra*, *Rhoæ*, *Fenes'tra* seu *Ni'grum* seu *Fora'men Oc'uli*, *Prunell'la*, *Lu'men*, *Gla'zë*, *Sight* or *Ap'ple* or *Ball* of the eye, (Sc.) *Blak* of the *Eie*, (Prov.) *Candle*, (F.) *Pupille*, *Prunelle*, (from *pupa*, 'a babe,' *pupilla*, 'a little babe;' so called from the image seen in the eye.) The aperture of the iris, through which the rays of light pass that have to impress the image of an object on the retina. This aperture can be dilated or contracted so as to allow a greater or smaller quantity of luminous rays to penetrate. The pupil, in man, is round, and by it the anterior and posterior chambers of the eye communicate with each other. In the fetus, during the first six months of gestation, it is closed by the pupillary membrane.

PUPIL CLOSURE OF THE, *Synechisis*.

PUPIL, PIN'HOLE. A condition of the pupil in typhus, in which it is so contracted as to resemble a pinhole. It indicates a dangerous form of brain affection.

PUPILLÆ VELUM, 'curtain of the pupil,' *Pupillary membrane*.

PUPILLARIS HERBA, *Lapsana*.

PUPILLARY, *Pupilla'ris*, (F.) *Pupillaire*. That which belongs to the pupil.

PUPILLARY MEMBRANE, *Membrana Pupilla'ris*, *Pupillæ Velum*, *Membrana Wachendorffiana*. The central aperture of the iris is closed, in the fetus, during a certain period of gestation, by a very thin membrane. It disappears in the seventh month.

PUPPET ROOT, *Veratrum viride*.

PUPPIS OS, 'stern bone,' *Frontal bone*.

PUPULA, *Pupil*.

PUPULÆ, *Ap'ices digito'rum*. The pulps or extremities of the fingers.

PURBLIND, (peebblind [?]) *Myopia*.

PURBLINDNESS, *Myopia*.

PURGAMENT, (from *purgare*, 'to cleanse,') *Purgative*.

PURGAMENTA, *Excrement*—p. *Alba*, *Leucorrhœa*—p. *Infantis*, *Meconium*—p. *Puerperii*, *Lochia*—p. *Uteri*, *Lochia*.

PURGANS MEDICAMENTUM, 'a purging medicine,' *Cathartic*.

PURGATIO, (from *purgare*, *purgatum*, 'to cleanse,') *Catharsis*—p. *Alvi*, *Catharsis*—p. *Muliebris alba*, *Leucorrhœa*—p. *Puerperii*, *Lochia*.

PURGATION, *Catharsis*.

PURGATIONES, [pl. of *purgatio*, 'cleansing,'] *Menses*—p. *Menstruæ*, *Menses*.

PURGATIONS, *Menses*.

PURGATIVE, *Purgati'vus*, *Pur'gans*, *Moch'licus*, *Catoret'icus*, *Catoter'icus*, *Cathar'tic*, *Pur'gament*, *Purge*, (F.) *Purgatif*. A medicine which operates more powerfully on the bowels than a laxative, stimulating the muscular, and exciting increased secretion from the mucous coat.

PURGE, *Purgative*—p. *Root*, *Euphorbia corollata*.

PURGING, *Diarrhœa*.

PURIFICANS, *Purific'us*, (*pnrus*, 'pure,' and *facere*, 'to make,') *Purifying*; hence:—

PURIFICANTIA, *Medicines* that cleanse or purify the blood.

PURIFICUS, *Purificans*.

PURIFORM, *Puriformis*, (*pus*, gen. *puris*, and *forma*, 'form.') Having the appearance of pus.

PURL. A medicated malt liquor, in which wormwood and aromatics are infused. Also, boiled beer having gin, sugar and spice mixed with it.

PUROHEPATITIS, (*pus*, gen. *puris*, and *hepatitis*,) see Hepatitis.

PURMUCOUS, *Purmuco'sus*, (*pus*, gen. *puris*, and *muco'sus*.) Having the character or appearance of pus and mucus.

PURPLE OF CASSIUS, *Pur'pura Mineralis Cassii*, (F.) *Pourpre de Cassius*. A precipitate, obtained by adding the protochloride of tin to a dilute solution of gold; or by dipping rods of tin in a solution of gold in diluted chlorohydric acid. Used in syphilis.

PURPLES, *Purpura hæmorrhagica*.

PURPURA, *Porph'ra*, 'a purple colour.' This word has been employed in many senses. It generally means livid spots on the skin from extravasated blood, with languor and loss of muscular strength, and pains in the limbs. — (F.) *Pourpre*.

In **PURPURA SIMPLEX**, *Porph'ra simplex*, *Petechie si'nè fe'brè*, *Pel'io'sis*, *Phenig'mus Petechia'lis*, *Pro'f'isio subcut'nea*, *Pet'chial Sc'ur'cy*, the spots are numerous, small and fleabite-shaped; chiefly on the breast, arms, and legs; with paleness of countenance. At times, the affection is accompanied by severe pains in the extremities, constituting *Pel'io'sis rheumat'ica*.

In **PURPURA HÆMORRHAGICA**, *Porph'ra hæmorrhag'ica*, *Stomac'acè univers'al'is*, *Hæmorrhag'ia univers'al'is*, *Petechi'na'sis*, *Mor'bus macula'sus hæmorrhag'icus Werthof'is*, *Pel'io'sis hæmorrhag'ica*, *Scorbut'us*, *Hæmorrh'ia petechi'alis*, *Hæmorrh'ia*, *Laud Sc'ur'cy*, *Pur'ples*, (F.) *Scorbut*, *S. de Terre*, *Maladie de Werthof*, *Hémorrhagique*, *Pétiase hæmorrhagique*—the spots are circular, and of different sizes; often in stripes or patches, irregularly scattered over the thighs, arms, and trunk; with occasional hæmorrhage from the mouth, nostrils, or viscera, and great debility and depression of spirits.

In **PURPURA NAUTICA**, *Porph'ra naut'ica*, *Scorbut'us*, *Pechytr'is*, *Gingibra'cium*, *Gingep'dium*, *Seelotgr'bè*, *Stomac'acè*, *Scorbut'us naut'icus*, *Sea sc'ur'cy*, (Sc.) *Serbie*, (F.) *Scorbut*, there are spots of different hues, intermixed with the livid, principally at the roots of the hair; the teeth are loose; the gums are spongy and bleeding; the breath is fetid, and the debility universal and extreme. This occurs chiefly at sea, after exposure to a moist, cold, foul atmosphere; with long use of one kind of food and of stagnant water.

All these varieties of purpura considerably resemble each other. In the first two, however, there seems, at times, to be some excitement, which has led to the employment of purgatives, and even of bleeding, with advantage. They are dependent upon a peculiar diathesis, which has been termed *Scorbut'ic Cachex'ia*, *Lues Sc'orbut'ica*, and *Cacochym'ia Sc'orbut'ica*. Generally speaking, the use of the mineral acids will be indicated; and, in sea scurvy, vegetable acids must be regarded as the best antiscorbutics. Nitre is, also, a good antiscorbutic, and in cases of great prostration it may be united with cinchona or turpentine. Tamarinds and cream of tartar, made into drink, are also serviceable. But the great remedy is change from low to cheerful spirits; from filth to cleanliness; from a cold and damp, to a temperate and dry climate, with sufficiency and change of food, and especially the use of green vegetables.

The general principles of management are the same in sea, as in land scurvy.

PURPURA, *Scarlatina* — p. Alba benigna et maligna, Miliary fever — p. Maligna, *Petechie* — p.

Miliaris, Miliary fever — p. *Mineralis Cassii*, Purple of Cassius — p. *Puerperarum*, Miliary fever — p. *Urticata*, *Urticaria*.

PURPURIC URINE, see *Porphyria*.

PURPURINE, see *Porphyria*.

PURPURISUM, *Hydrargyri sulphuretum rubrum*.

PURRING TREMOR, *Frémissement cataire*.

PURSE, ([L.] *bursa*, [F.] *bourse*.) Scrotum — p. Shepherd's, *Thlaspi bursa*.

PUR'SINNESS, *Dyspno'a pinguetudo'sa*, (F.) *Pousse*. Dyspnoea accompanied with oppressive fatness. The word is, also, applied to dyspnoea of every kind. One so affected is said to be *purey*, or *shortwinded*. Prov. *Short-of-puff*, *Thicklisted*, (F.) *Poussif*, old (F.) *Poussif*.

PURSLAIN, ([L.] *portulaca*, [L.] *porcellana*, old (F.) *pourcelaine*.) *Purslane*.

PURSLANE. *Portulaca* — p. Milk, *Euphorbia corollata* and *E. maculata* — p. Speedwell, *Veronica peregrina* — p. Water, *Ludwigia palustris*, *Veronica beccabunga*.

PURSY, see *Pursinness*.

PURULENCE, (from *pus*, gen. *puris*.) Pus, Suppuration.

PURULENCY, *Pus*, Suppuration.

PURULENT, *Purulent'us*, *Em'pyus*, *Pyo'des*, *Py'icus*. That which has the character of pus or consists of pus; as, *purulent matter*, a *purulent collection*, &c.

PURULENT DEPOSIT, *Pus*.

PURULENTIA, Suppuration.

PURVAIN, *Verbena officinalis*.

PUS, gen. *Pu'ris*, (*πυς* and *πυω*.) *Hu'mor purulent'us*, *Py'on*, *Py'us*, *Mut'ter*, *Pu'ulent deposit*, *Pu'ulence*, *Pu'ulency*, (Sc.) *Woursum*, *Worsum*. A secretion from inflamed textures, and especially from the areolar membrane. It is, almost always, of the same nature, whatever may be the part it proceeds from. Pus of a good quality, — *laud'able pus*, *Pus laud'um seu laudab'ile seu nutritivum seu ve'r'um*, (F.) *Pus louable*, — is of a yellowish-white colour, opaque, inodorous, and of a creamy appearance. Heat, acids, and alcohol coagulate it. When analyzed by Schwilgué, it afforded albumen and water, a particular extractive substance, and a small quantity of soda, phosphate of lime, and other salts. Normal pus consists essentially of two distinct parts, *pus corpuscles* or *pus globules*, *Cy'toid corpuscles* of Henle — and a colourless, aqueous fluid, *li'quor pu'ris*, in which the corpuscles are suspended. A variety of the pus corpuscles is described by M. Lebert under the name *py'oid*. Fully formed pus is aplastic.

PUS BONUM, see *Pus* — p. *Corpuscles*, see *Pus* — p. *Corrosivum*, *Sanies* — p. *Globules*, see *Pus* — p. *Laudable*, see *Pus* — p. *Louable*, see *Pus* — p. *Malignum*, *Ichor*, *Sanies* — p. *Nutritivum*, see *Pus* — p. *Verum*, see *Pus*.

PUSA, *Girl*.

PUSH, *Pustule*.

PUSILLA, (dim. of *pusa*.) *Girl*.

PUSILLATUM, *Pusilla'tum*, (from *pusillus*, 'small' [?].) A coarse powder.

PUSILUS, 'small.' *Nanus*.

PUSKILE, *Pustule*.

PUSTULA, (from *pus*.) *Pustule* — p. *Ardens*, *Eczema* — p. *Gangrenosa*, see *Anthrax* — p. *Oris*, *Apthæ* — p. *Livida* *Esthoniz*, see *Anthrax* — p. *Maligna*, see *Anthrax*.

PUSTULE SACRÆ, 'dry pustules,' *Lichen*.

PUSTULAR, *Pus'tulous*, *Pustula'ria*, *Pustulo'sus*, (F.) *Pustulencus*. Of or belonging to pustules. Affected with pustules.

PUSTULATION, *Pustula'tio*, (*pustulare*, *pustulatum*, 'to cause pustules,') The formation of pustules.

PUSTULE, *Pus'kile*, *Push*, *Pus'tula*, *Ecp'y'sis*, *Ec'thyma*, *Beneth*, *Bubò*, *Epyras'tosis*. An elevation of the cuticle, with an inflamed base;

Willan and Bateman reckon four varieties of pustules, *Phlyctenium*, *Psudacium*, *Achor*, and *Favus*. The varicelous pustule is often called, by the French, *Grain*: English, *Pock*.

PUSTULE MALIGNA, see *Anthrax*.

PUSTULOUS, Pustular.

PUSULATUM, Pusillatum.

PUSUS, Boy.

PUT TO BED, Deliver.

PUTAMEN OVI, ('shell of an egg,' from *pute*, 'to cast off,') see *Membrana putaminis*.

PUTIUM, (πυθιον,) Prepuce.

PUTREDO, gen. *Putredinis*, 'rotteness,' Hospital Gangrene, Putrefaction.

PUTREFACTION, *Putrefactio*, *Putrescentia*, *Sep'is*, *Sap'rotes*, *Se'pedon*, *Putrid* or *Putrefactive fermentation*, *Put'ror*, (*putris*, 'rotten,' and *facere*, 'to make,') A decomposition, experienced by animal substances, when deprived of life, and placed under special circumstances. The presence of water is indispensable; the temperature most favourable to it, is from 60° to 90° of Fahrenheit. The most common products of putrefaction are, water, carbonic acid, acetic acid, ammonia, carburetted hydrogen, and a semi-putrid substance, which is volatilized, and has an infected odour. It has been supposed that something like putrefaction may occur in the living body, as in case of putrid fever. To this condition, Piory has given the name *Typhoe'mia*, conceiving it to be owing to alteration of the blood by putrid animal matters. *Putrefaction* or *Putrescence* properly signifies the state of becoming putrid: whilst *Putridity*, *Putre'do*, *Putriditas*, *Putridness*, (F.) *Putridité*, *Pourriture*, is the state of rottenness.

PUTRESCENT, (from *putrescere*, 'to grow rotten,') Hyposaprus—p. Poison, see *Poison*.

PUTRESCENTIA, Putrefaction—p. Uteri gravidi, Hysteromalacia.

PUTRID, *Putridus*, *Sap'ros*, *Sepedono'des*, (F.) *Putride*, (*putreo*, 'to rot,') An epithet for affections, in which the matters excreted and the transpiration itself exhale a smell of putridity. It is particularly applied to typhus.

PUTRIDITAS, see *Putrefaction*.

PUTRIDITY, see *Putrefaction*.

PUTRILA'GO, gen. *Putrilag'inis*, (*puter*, 'rotten,') (F.) *Putrilage*. The putrescent matter or *slough*, which forms in certain gangrenous and other ulcers, and is thrown off.

PUTRORE, Putrefaction.

PUTZSCHILL, see *Myroxylon Peruiferum*.

PUXIRI, see *Pichurim beans*.

PYÆMIA, Pyohamia.

PYAR, (πυαρ,) Colostrium.

PYARTHRO'SIS, (*pyo*, αρθρον, 'a joint,' and *osis*,) Suppuration of a joint.

PYAR'THRUM, (*pyo*, and αρθρον, 'joint,') A suppurated joint.

PYCNA'THEMUM INCA'NUM, (πυχνος, 'dense,' and ανθερον, 'a blossom,') *Clinopodium incanum*, Common Mountain Mint, Wild Basil. An indigenous plant of the mint family—Ord. Labiate—which flowers in August. It has the aromatic properties of the mints.

PYCNA'THEMUM LINIFOLIUM, *Virginia Thyme*, has like virtues.

PYCNICMA'SIA, *Pycnic'masia*, (πυκνος, 'thick,' and ικμας, and ικμασια, 'moisture,') A thickening of the juices.

PYCNICMASTICA, *Incrassantia*.

PYCNOMETER, (πυκνος, 'thick,' and μετρον, 'measure,') Areometer.

PYCNO'SIS, (πυκνωσις, from πυκνος, 'thick,' and *osis*,) Inspissatio.

PYCNOTICA, (πυκνωτικά,) *Incrassantia*.

PYEC'CHYSIS, (πυσω, 'pus,' and εκχυνσις, 'effusion,') An effusion of pus.

PYELITIS, (F.) *Pyélite*, (*pyelus*, and *itis*,) Inflammation of the pelvis and calices of the kidney.

PYELOMETER, (*pyelus*, and μετρον, 'measure,') Pelvimeter.

PYELONEPHRITIS, (*pyelus*, νεφρος, 'kidney,' and *itis*,) Inflammation of the kidney, and of the pelvis and calices.

PYELUS, (πυελος,) Pelvis.

PYEMESIS, Pyoemesis.

PYESIS, (πυσις,) Suppuration.

PYETIA, (πυετια,) Colostrium, Rennet.

PYGE, (πυγη,) Nates.

PYGMÆUS, (πυγμαιος, from πυγμα, 'pygme,') Pygmy.

PYGMÆ, (πυγμα,) Fist, Forearm.

PYGMÆE, Pygmy.

PYGM'Y, *Pig'my*, *Pygmæus*, (F.) *Pygmée*, (from *pygme*, as big as the fist or forearm,) A dwarf. The Pygmæi were a fabulous nation of dwarfs in the extreme parts of India; or, according to others, in Ethiopia.

PYGODID'YMUUS, *Did'yimus*, *symphyoperinæus*, *Pygop'ges*, (*pyge*, and ιδυμος, 'a twin,') A monstrosity, in which twins are united by the sacrum and coccyx.

PYGO'M'ELES, (F.) *Pygomèle*, (*pyge*, and μελες, 'unprofitable,') A monster with excessive growth on the nates or rump.

PYGOPAGES, (*pyge*, and παγω, 'to fix,') *Pygodidymus*.

PYICUS, (from πυον, 'pus,') Purulent.

PYLE, (πυλη, 'a gate,') Porta.

PYLEMPHRAX'IS, (*pyle*, 'a gate,' and εμφραξις, 'obstruction,') Obstruction of the vena porta.

PYLEPHLEBITIS, (*pyle*, φλεψ, gen. φλεβος, 'a vein,' and *itis*,) Inflammation of the vena porta.

PYLOR'IC, *Pylor'icus*, (F.) *Pylorique*, (from *pylorus*,) That which relates to the pylorus. An epithet given to different parts.

PYLORIC ARTERY, *Arteria Coronaria dextra*, (F.) *Petite artère gastrique droite*, arises from the hepatic, and is distributed to the pylorus and to the lesser curvature of the stomach, anastomosing, particularly, with the A. coronaria ventriculi and A. gastroëpiploica dextra.

PYLORIC VEIN follows the same distribution as the artery.

PYLORIQUE, Pyloric.

PYLO'RUS, (πυλωρος,) Janitor, Portona'rium, (πυλη, 'a gate,' and ορος, 'a guardian,') *Ostium ventriculi duodena'te seu pylo'ricum seu dextra'tum seu inferius*, *Orific'ium dextra'tum seu inferius seu Egres'sus seu Os inferius seu An'nulus seu Sphinc'ter ventriculi*, *Ostia'rius*, (F.) *Pylore*. The lower or right orifice of the stomach—*Orifice intestinal* (Ch.)—is called *Pylorus*, because it closes the entrance into the intestinal canal, and is furnished with a circular, flattened, fibro-mucous ring, which occasions the total closure of the stomach during digestion in that organ. This ring has been called the *Valve of the Pylorus*, *Sphincter Pylori*, *Pylorus propriè sic dictus*, (F.) *Valvule du pylore*. It is a fold of the mucous and muscular membranes of the stomach; and is the *Pyloric muscle* of some authors.

PYO, in composition, means pus, (from πυον, 'pus,')

PYOBLENNICUS, (*pyo*, and βλεννα, 'mucus,') Mucopurulent.

PYOBLENNORRHÆA, (*pyo*, βλεννα, 'mucus,' and ρεω, 'I flow,') A discharge of puriform mucus.

PYOCENO'SIS, (*pyo*, and κενωσις, 'evacuation,') Evacuation of pus.

PYOCH'EZIA, *Diarrhœa purulenta*, (*pyo*, and χεζω, 'I go to stool,') Discharge of pus by stool. *Purulent diarrhœa*.

PYOCÆLIA, *Asci'les purulen'tus*, (*pyo*, and *καῖλα*, 'the belly.') Pus in 'the abdominal cavity.

PYOCYSTIS, (*pyo*, and *κυστίς*, 'a sac,') Vomica.

PYODES, (*πυώδης*,) Purulent.

PYODE'MA, gen. *Pyode'matis*, (*pyo*, and *οἶημα*, 'swelling,') Tumefaction of the surface owing to effusion of pus.

PYOE'MESIS, *Pyem'esia*, *Vom'itus purulen'tus*, (*pyo*, and *εμεσις*, 'the act of vomiting,') Vomiting of pus.

PYOE'MIE, (*pyo*, and *αἷμα*, 'blood,') Pyohæmia.

PYOGENESIS, Pyogenia.

PYOGENETIC, Pyogenic.

PYOGEN'IA, *Puogen'ia*, *Pyogen'esia*, *Pyopoë'sia*, (*pyo*, and *γεννω*, 'to generate,') Generation of pus. The theory or mechanism of the formation of pus. Some have believed that pus is formed by the arterial system, and is deposited, by way of excretion, in the inflamed parts; others, that it is formed by the destruction of solid parts. It seems to be owing to a degeneration of the liquor sanguinis and exudation corpuscles.

PYOGENIA CORROSIVA, Ulceration—p. Simplex, Suppuration.

PYOGEN'IC, *Pyogen'icus*, *Pyogenet'ic*, *Pyogenet'icus*, (F.) *Pyogénique*, *Pyogénétique*. Having relation to the formation of pus.

PYOHÆ'MIA, *Pyæ'mia*, (F.) *Pyohémie*, *Pyhémie*, *Pyôémie*, (*pyo*, and *αἷμα*, 'blood,') Alteration of the blood by pus, giving occasion to the *Diath'esis seu Infect'io purulen'ta*.

PY'OID, (*πυοειδής*, from *pyo*, and *εἶδος*, 'resemblance,') Resembling pus.

PYOID CORPUSCLES or GLOBULES, see Pus.

PYOMA, gen. *Pyo'matis*, Pyonoma.

PYOME'TRA, *P. purulen'ta*, (*pyo*, and *metra*,) A collection of pus in the uterus.

PYOMETRA PURULENTA, Pyometra.

PYON, (*πυόν*,) Pus.

PYONÉPHRITE, (*pyo*, and *nephritis*,) see Nephritis.

PYONO'MA, gen. *Pyono'matis*, *Pyo'ma*, (from *pyon*, and *oma*,) A suppurating sore.

PYOOTORRHŒA, (*pyo*, and *otorrhœa*,) Otorrhœa.

PYOPERICAR'DIA, (*pyo*, and *pericardium*,) *Pyopericar'dium*. A collection of pus in the pericardium.

PYOPHTHALMIA, (*pyo*, and *ophthalmia*,) Hypopyon—p. Neonatorum, see Ophthalmia (purulenta infantum).

PYOPLA'NIA, (*pyo*, and *πλάνος*, 'wandering,') An extension of pus to other parts; purulent infiltration.

PYOPLEURITE, (*pyo*, and *pleuritis*,) Empyema.

PYPOÏESIS, (*pyo*, and *ποιω*, 'to make,') Pyogenia.

PYPOÏETIC, Suppurative.

PYOPTYSIS, (*pyo*, and *πτωω*, 'I spit,') Spitting of pus.

PYORRHAG'IA, (*pyo*, and *ῥαγή*, 'violent rupture,') A sudden discharge of a large collection of matter.

PYORRHŒA, (*πυορροία*, from *pyo*, and *ῥέω*, 'I flow,') A discharge of pus.

PYORRHŒA ALVINA, Pyochezia—p. Aurium, Otorrhœa—p. Nasalis, Ōzena—p. Palpebræ, Blepharopyorrhœa—p. Urinalis, Pyuria—p. Viarum lacrymalium, Dacryopyorrhœa.

PYOS, (*πυός*,) Pus. Also, first milk, colostrum.

PYOSIS, (*πυώσις*,) Hypopyon, Pterygion, Suppuration—p. Pectoris, Empyema.

PYOTHORAX VERUS, (*pyo*, and *thorax*,) Empyema.

PYOTORRHŒE, (*pyo*, and *otorrhœa*,) Otorrhœa.

PYOTURIA, (*pyo*, and *ουρον*, 'urine,') Pyuria.

PYR, (*πυρ*,) Fever.

PYRA, (*πυρα*, from *πυρ*, 'fire,') Anthrax—p. Crustumerina, Crustuminatum—p. Crustumina, Crustuminum.

PYR'AMID, *Pyram'is*, *Eminen'tia pyramida'lis tym'pani*, (*περαμ'is*, gen. *πυραμ'idos*, from *πυρ*, 'flame,' which has a conical shape.) A small bony projection in the cavity of the tympanum, which is excavated, to lodge the muscles of the stapes. The *Centrepin* of the trephine. A small, obtusely pointed eminence of the inferior veruiform process of the cerebellum.

PYRAMID, see Calumba—p. of Malacarne, Vermiform process, inferior.

PYRAMIDAL DE L'ABDOMEN, *Pyramidalis abdominis*—p. de la *Cuisse*, *Pyramidalis*—p. du *Nez*, *Pyramidalis nasi*—p. *Stapédien*, *Stapedius*.

PYRAMIDALE (OS), Cuneiform, (bone.)

PYRAMIDA'LIS, *P. Fem'oris*, *Ili'acus exter'nus*, *Pyriform'is*, *Piriform'is*, *Pri'mus et super'ior quadrang'inus*, (F.) *Sacro-trochantérien* (Ch.), *Pyramidal de la cuisse*. A flat, triangular muscle, situate at the posterior part of the pelvis. By its base, it is attached to the anterior surface of the sacrum; to the corresponding surface of the great sacrosciatic ligament and to the posterior part of the ilium. It terminates by a tendon, which is attached to the inner surface of the great trochanter. This muscle is a rotator of the thigh outwards, or of the pelvis in an opposite direction.

PYRAMIDA'LIS ABDOM'INIS, *Mus'culus Fallo'pii seu succenturiatus seu auxiliiarius*, (F.) *Pubio-ombilical*, *Tubio-sous-ombilical* (Ch.), *Pyramidal de l'abdomen*. A small, fleshy, pyramidal fasciculus, whose existence is not constant; and which is inserted, by its base, into the upper part of the pubis, and terminates, by its summit, at the inferior part of the linea alba. When it contracts, it stretches the linea alba.

PYRAMIDALIS FEMORIS, *Pyramidalis*.

PYRAMIDALIS NA'SI, *Proc'erus na'si*, *Na'sum dilatans*, (F.) *Fronto-nasal* (Ch.), *Pyramidal du nez*. A thin, triangular muscle, which has its summit upwards. Above, it is continuous with the occipitofrontalis. It descends vertically before the root of the nose, on which it terminates, becoming confounded with the transversalis. This muscle wrinkles the skin of the root of the nose transversely, and stretches that which covers the tip.

PYRAMIDES, (pl. of *pyramis*,) *Corpora pyramidalia*.

PYRAMIDES ANTÉRIEURES, *Corpora pyramidalia*—p. Ferreini, Ferrein, pyramids of—p. Malpighiane, *Papillæ* of the kidney—p. *Postérieures du cercelet*, *Corpora testiformia*.

PYRAMIDS OF FERREIN, see Ferrein—p. of Malpighi, *Papillæ* of the kidney—p. Posterior, *Corpora testiformia*—p. of Wistar, *Sphenoidal cornua*.

PYRAMIS, gen. *Pyram'idis*, (*πυραμ'is*, gen. *πυραμ'idos*,) Penis, Pyramid—p. Cochlea, Modiolus—p. Trigona, see Temporal bone.

PYRCLÆ, (*πυρκαϊα*, 'a conflagration,' from *πυρ*, 'fire,' and *καωω*, 'to kindle,') Incendium.

PYREC'TICA, 'febrile,' (*pyreticus*, and *εχω*, 'I have,') -Fever: the first order in the class *Hæmatica* of Good.

PYRÉLAINES, (*πυρ*, 'fire,' and *ελαϊον*, 'oil,') Olea empyreumatica.

PYRENOIDES, (*πυρηνοειδής*, from *πυρην*, 'the kernel, the stone of a fruit,' and *εἶδος*, 'resemblance,') Odontoid.

PYRÉTHRE, *Anthemis pyrethrum*.

PYRETHRUM, (*πυρεθρον*, from *πυρ*, 'fire,') *Anthemis pyrethrum*, *Urticaria*—p. *Parthenium*, *Matricaria parthenium*—p. *Sylvestre*, *Achillea ptarmica*.

PYRETICO'SIS, (from *pyretus*.) *Morbus febrilis*. A febrile affection.

PYRETICUS, (from *pyretus*.) Febrifuge, Febrile.

PYRETOGEN'ESIS, *Pyretogen'ia*, (*pyretus*, and *genesis*, 'generation.') The origin and formation of fever.

PYRETOGENET'IC, (same etymon.) Stimulant.

PYRETOGRAPH'IA, (*pyretus*, and *γραφη*, 'a description.') A description of fever.

PYRETOL'OGY, *Pyretolog'ia*, (*pyretus*, and *λογος*, 'a discourse.') *Pyrol'ogy*. A treatise on fevers. A title sometimes given to a monograph on fever.

PYRETOTYPOSIS, (*pyretus*, *τυπος*, 'type, order,' and *osis*.) Intermittent.

PYRETUS, (*πυρετος*.) Fever.

PYREX'IA, *Pyrex'ia*, (*πυρεξις*, from *pyr*.) Fever. The febrile condition. Also, a paroxysm of fever—*Paroxysmus febrilis*.

PYREX'IE, *Febrile diseases*. The first class of Cullen's Nosology.

PYRI MARTIALES, (pl. of *pyrus*, 'a pile, or pear,') *Ferrum tartarizatum*.

PYRIA, (*πυρία*.) Fomentation, see Stove.

PYRIATERIUM, (*πυριατηριον*.) see Stove.

PYRICASTUM, (*πυρικαστος*, from *pyr*, 'fire,' and *καω*, 'to kindle,') Burn.

PYRIFORMIS, (*pyrus*, 'a pear,' and *forma*, 'shape,') *Pyramidalis*.

PYRIPH'LEGES, (*πυριφλεγες*.) *Pyriphlectus*, (*πυριφλεκτος*, from *pyr*, and *φλεγω*, 'I burn,') One who has a burning fever.

PYRITES, **IRON**, (from *pyr*, 'fire,') *Ferri sulphuretum*.

PYR'MONT, **MIN'ERAL WA'TERS OF**. A celebrated mineral spring on the Weser, in Westphalia, four leagues from Hamelot. The waters are tonic, and contain chlorides of sodium and magnesium, sulphates of soda and magnesia, and carbonates of iron, lime, and magnesia dissolved in carbonic acid, with some resinous (?) principle. It is a highly carbonated chalybeate. Season, from the 10th of June to the 1st of September.

ARTIFIC'IAL PYR'MONT WA'TER may be formed of *Epsom salt*, gr. xv; *Common salt*, gr. v; *Magnesia*, gr. x; *Iron filings*, gr. v; *Water*, Oij; impregnated with the gas from *Marble-powder* and *Sulphuric acid*, aa 3vij.

PYRÆNUS, (*πυρ*, 'fire,' and *οινος*, 'wine,') *Spiritus rectificatus*.

PYROLA, *P. rotundifolia*, *Roundleaved Pyrola* or *Win'tergreen*; *Shinleaf*. Ord. Ericaceæ. (Dim. of *pyrus*, 'a pear tree,' from a fancied resemblance of the foliage.) This plant was once used as a gentle astringent and tonic.

PYROLA MACULA'TA, *Chimaph'ila Macula'ta*, *Spot'ted Pipsis'seava*, *Spot'ted Win'tergreen*, *White'leaf*, *White Pipsis'seava*, *King'cure*, *Ground holly*, *Rheumatism weed*, &c., has similar properties.

PYROLA, **ROUNDLEAVED**, *Pyrola*.

PYROLA UMBELLA'TA, *Chimaph'ila seu Chimoph'ila umbella'ta*, *C. Corymbo'sa*, *Ground holly*, *Pipsis'seava*, *Win'tergreen*, *Rheumatism-weed*, *Prince's pine*, *Herbe a pisser* (Canada). This common American plant is allied to the *uva ursi* in botanical as well as medical qualities. It is tonic and diuretic. The bruised leaves irritate the skin. From the decoction or infusion an agreeable beverage may be formed, by adding sugar, ginger to flavour it, and yeast to produce fermentation. Under the name *Pipsis'seava Beer* it has been used in scrofulous affections.

PYROLEUM OSSIUM RECTIFICATUM, (*πυρ*, 'fire,' and *oleum*, 'oil,') *Oleum animale Dipelii*.

PYROLIG'NEOUS AC'ID, (*πυρ*, 'fire,' and *λίγνιον*, 'wood,') *Ac'idum pyroac'eticum seu pyro-*

λίγνιον seu pyrolig'neum seu pyroxy'licum seu lig'ni pyroileo'num seu ac'eticum empyreumaticum seu lig'neum seu ligno'num empyreumaticum, *Pyrolig'nic ac'id*. (F.) *Acide pyroactique ou pyrolig'nique ou pyrolig'neuse*, *Vinigre de bois*. An acid, obtained by distillation from wood, coal, &c., which was once believed to be of a peculiar character. It is now known to be formed of acetic acid and an empyreumatic oil, from which it may be freed; and, in this state, is sold in commerce for *Acetum distillatum*, *Acetic acid of commerce*, *Ac'idum ac'eticum e Ligno venal'e*. It is used for similar purposes as distilled vinegar, and for preserving animal substances. It has also been used as an antiseptic in gangrene and sphacelus, and to foul ulcers; wherever indeed creasote is indicated. The dose of the impure pyroligneous acid internally is from five to thirty drops, three or four times a day.

PYROMA'NIA, *Empresomoma'nia*, *Incen'diary monoma'nia*, (*πυρ*, 'fire,' and *mania*.) Insanity, with an irresistible desire to destroy by fire.

PYROPH'AGUS, *Igniv'orous*, (*πυρ*, 'fire,' and *φαγω*, 'I eat,') One who eats fire. A juggler, who eats substances in a state of ignition.

PYRUS, (*πυρος*.) *Triticum*.

PYRO'SIS, (from *πυρω*, 'to burn,') *Ptyalis'mus pyros'icus*, *Limo'sis cardial'gia sputato'ria*, *A'dor seu Ebulli'tio stom'achi seu ventric'uli*, *Cardial'gia sputato'ria*, *Pyro'sis Suec'ica*, *Apoceno'sis vom'itus pyro'sis*, *Dyspep'sis pyro'sis*, *Cnismoreg'mia*, *Orex'is* (of some), *So'da*, *Wa'terbrash*, *Wa'terqualm*, *Wa'terping*, *Blackwater*, (F.) *Ardeur de l'estomac*, *Fer chaud*, *Fer rouge*, *Cremaison*, *Gorgoeset*. This affection consists of a hot sensation in the stomach, with eructations of an acrid, burning liquid, that causes a distressing sensation in the parts over which it passes. Attention to diet, and the use of bitter absorbents, will usually relieve, when the disease is functional. Occasionally, it is symptomatic of organic disease of the stomach.

Pyrosis also signified, of old, inflammation, gangrene, and a burn.

PYROSIS STECICA, *Pyrosis*.

PYROSOPHIA, (*πυρ*, 'fire,' and *σοφία*, 'wisdom,') Chymistry.

PYROTECHNIA, (*πυρ*, 'fire,' and *τεχνη*, 'art,') Chymistry.

PYROTECHNY, *Pyrotechn'ia*, *Pyrotech'nia*. Art of making fire-works, &c. Also, chymistry. M. Percy has used the term *Pyrotechnie Chirurgicale* for the art of applying fire or the actual cautery in surgery.

PYROTHONIDE, (*πυρ*, 'fire,' and *οβιον*, 'linen,') *Liquor pyroileo'sus e lin'teo para'tus*. A kind of pyrogenous or empyreumatic oil, produced by the combustion of textures of hemp, linen, or cotton in a copper vessel. The brown product is acid. Its medical properties probably resemble those of creasote. Diluted with three or four times its weight of water, it has been used as a gargle in cynanche tonsillar, but is not now employed.

It is said, also, to be a popular remedy for toothache and skin diseases. When prepared from rags, it is called *Ros oil*; when from paper, *Paper oil*, (F.) *Huile de Papier*.

PYROTICUS, (*πυρωτικός*, from *πυρω*, 'to burn,') Caustic.

PYROXYLIN, (*πυρ*, 'fire,' and *ξύλον*, 'wood,') Gun cotton, see Collodion.

PYRUS, (from *πυρ*, 'fire,' because flame is shaped like the pear, see *Pyrus malus*—p. Americana, *Sorbus Americana*.)

PYRUS CYDO'NIA, *Cydo'nia malifor'mis seu vulg'aris*, *Sor'bus Cydo'nia*, *Cydo'nia*, *Coto'nia*, *Quince Tree*, (F.) *Coignassier*, *Cognassier*. Ord. Rosaceæ. *Sex. Syst.* Icosandria Pentagynia. The fruit is termed *Cydo'nium ma'lum*, *Ma'lum ca'dnum seu*

coto'neum, *Me'lum cydo'nium*, *Quince*, (F.) *Coign*. Quince-seed — *Cydo'nium* (Ph. U. S.) — are sometimes used in decoction as a demulcent. The fruit makes a pleasant preserve.

PYRUS DOMESTICA. *Sorbus domestica*.

PYRUS MALUS. *Malus*, *M. dasycarpa* seu *communis* seu *sylvestris*, *Sorbus malus*, *Me'lea*, the *Ap'ple Tree*, (F.) *Pommier*. The *apple* — *Po'mum*, *Ma'lum*, (F.) *Pomme* — is an agreeable fruit when ripe. When unripe, it disagrees, especially with those of weak digestive powers. The same may be said of the *pear*, (F.) *Poire*; the fruit of *Py'rus communis*, *P. Sorbus* seu *Sat'iva*, *P'rus*, *A'pios*, (F.) *Poirier*.

Ap'ple Brand'y is a spirituous liquor obtained from the juice of the apple, and much used in the United States.

PYTHOGENIC FEVER, (πυθω, 'to putrefy,' and γενναω, 'to generate.') Typhoid fever.

PYTHIA, (πυθια,) *Colostrum*, *Rennet*.

PYTISMA, (πυτισμα,) *Sputum*.

PYUL'CON, *Pyul'cum*, (πυυλκον, from *pyo*, and λαω, 'I extract.') (F.) *Tire-pus*. A surgical instrument for extracting pus from any cavity. Many pyulca have been employed, all of which act as exhausting syringes.

PYURIA, *Pyotu'ria*, *Pyorrhœ'a urina'lis*, (F.) *Pyurie*, *Pissement de Pus*, (*pyo*, and ουρον, 'urine.') Discharge of purulent urine: a symptom which occurs in many diseases; particularly in cases of renal calculi, and in organic affections of the bladder. It receives various epithets, according to the part whence the pus proceeds: — as *Pyuria urethralis*, *P. renal'is*, and *P. vesical'is*.

PYURIA CHYLOSA, *Chyluria* — p. *Lactea*, *Chyluria* — p. *Mucosa*, *Cystirrhœa* — p. *Serosa*, *Cystirrhœa* — p. *Viscida*, *Cystirrhœa*.

PYXIS, gen. *Pyx'idis*, (πυξίς, gen. πυξίδος, made of box-wood, πυξος, *Pyxus*.) A box; a pill-box. Also, an emollient ointment.

PYXIS, *Boîte*, see *Cotyloid*.

Q.

Q. see Abbreviation.

QUABEBE, *Piper cubeba*.

QUACK, *Charlatan*.

QUACK MEDICINE, *Arcanum*.

QUACKSALVER, ([G.] *Quacksalber*, from quaken, 'to quack,' and Salbe, 'salve,') *Charlatan*.

QUADRANS, *Quarta pars li'bræ*. A quarter of a pound. Three ounces (Troy.)

QUADRA'TUS, (F.) *Carré*, (*quadrus*, 'square,') A name for muscles whose figure approaches a square. Depressor Labii inferioris, *Quadratus femoris*.

QUADRATUS DORSI, *Q. lumborum*.

QUADRATUS FEMORIS, *Quadra'tus*, *Quadratus quadrigen'ius quadratus*, (F.) *Tuber-ischio-trochan'térien*, (Ch.) *Ischio-sous-trochan'térien*, *Carré de la Cuisse*. A muscle situate at the posterior and upper part of the thigh. It is thin, flat, and, quadrilateral, and is attached to the outer part of the tuberosity of the ischium; whence it proceeds transversely to the inferior part of the posterior margin of the great trochanter. It rotates the limb, turning the toe outwards.

QUADRATUS GENÆ, *Platysma myoides* — q. *Radii*. Pronator radii quadratus.

QUADRATUS LUMBO'RUM, *Quadra'tus* seu *Lumbaris exter'nus*, *Q. dor'si*, *Flec'tens par lumborum*, (F.) *Ilio-costal*, (Ch.) *Ilio-lombi-costal*, *Carré des lombes*. A flat, somewhat thick, irregularly quadrilateral muscle, situate in the loins, at the sides of the vertebral column. It is attached, below, to the crista illi and ilio-lumbar ligament; and above, to the inferior margin of the last rib; and besides, to the base of the transverse processes of the last four lumbar vertebræ. This muscle inclines the loins to its side; depresses the last rib, and is thus a muscle of respiration. It can also raise the hip.

QUADRATUS MENTI, *Depressor labii inferioris*.

QUADRIGA, (*quatuor*, 'four,' and *jugum*, 'a yoke,') *Cataphractus*.

QUADRIGEMINA TUBER'cula, (*quatuor*, 'four,' and *geminus*, 'double,') *Éminence Quadrigem'ine* seu *Bigem'ine*, *Optic Lobes*, *Optic Ganglia*, *Pons Syl'vii*, *Cor'pora quadrigem'ina* seu *bigem'ina*, *Nates* et *Tes'tes*, (F.) *Tubercules quadrigemeaux* ou *bijumeaux*, *Éminences sigéennes*, *Lobes optiques* ou *bijumeaux*. Four medu-

lary tubercles, situate at the posterior surface of the tuber annulare. They are white, externally; gray, internally; oblong; rounded; connected by pairs, and separated by two grooves, which divide them crucially. Of these tubercles — the superior called *Nates*, *Nat'ulæ*, *Tuber'cula antero'ra*, *Glu'tia*, are larger and more prominent than the lower, called *Tes'tes*, *Tuber'cula postero'ra*. The pineal gland corresponds exactly to the point of intersection of the two furrows that separate them.

They give origin to the optic nerves.

QUADRIGEMINUS PRIMUS ET SUPERIOR, *Pyramidalis* — q. *Quartus quadratus*, *Quadratus femoris* — q. *Secundus et tertius*, *Ischiotrochanterianus*.

QUADRIGEMINA TUBERCULES), *Quadrigenina Tubercula*.

QUADRIMAMMA'TUS, *Tetramas'thus*, (*quatuor*, 'four,' and *mamma*, 'a breast,') *Four breasted*.

QUADROON, (from [L.] *quatuor*, 'four,' a fourth part,) see *Mulatto* — q. *Black*, see *Mulatto*.

QUÆ EX QUIBUS, (*quæ ex quibus*.) A Hippocratic phrase, which comprised the vicissitude, succession, and transmutation of diseases, of which mutation and succession four species were chiefly enumerated, — *epigenesis*, *metaptosis*, *metastasis*, and *apostasis* or abscess. Stephanus Rodericus a Castro wrote a small work under the title "*Quæ ex quibus*," de Morborum Vicissitudinibus.

QUAFADIL, *Melanthium Virginicum*.

QUAHUIL, *Cacao*.

QUAIL, (low L. *qualea*, [F.] *Caille*,) *Tetrao coturnix*.

QUAMASH, *Camassia esculenta* — q. *Eastern*, *Scilla esculenta*.

QUAN'DROS. Ancient name for a precious stone, believed to exist in the brain of the vulture; and to which was attributed the property of augmenting the secretion of milk, and preserving from deadly accidents. No such stone exists.

QUAR'ANTINE. *Quar'antine*, *Quarentaine*, ([L.] *Quaranti'na*, itself from *quaranta*, 'forty,') *Quarantine*. The time during which travellers, returning from a country where the plague or any other pestilential disease exists, are compelled to remain on board ship or in a lazaretto. They, as well as the ship, are said 'to perform quarantine.'

and 'to be quarantined.' Quarantine is sometimes extended beyond forty days, whilst its duration is generally much less. It is sometimes enforced by land: for example, when contagious diseases require the establishment of *cordons sanitaires*, and when persons are desirous of leaving infected districts. During quarantine, no communication is allowed between the travellers and the inhabitants of the country. Quarantine is a great impediment to commerce; and is, doubtless, very frequently enforced when there is but little danger of contagion. The evils, however, that might arise from the introduction of a widely-spreading contagious disease, are so disastrous, as to induce legislators to maintain these laws in perhaps an unnecessary degree of rigour.

QUAR'TAN, *Quarta'nus*, (from *quartus*, 'the fourth,') *Quarta'na Febris*, *Acutus Quarta'nus*, *Tetarta'nus*, *Quarta'tan Agne*, (F.) *Quarte*. An intermittent, the paroxysms of which recur every fourth day, leaving between them two days' interval. We speak also of the *Quar'tan Type*. A *Double Quar'tan*, (F.) *Double quarte*, is one in which one paroxysm occurs on the third and another on the second day, so that there is only one day of apyrexia and two of paroxysms in succession;—the paroxysms of every fourth day resembling each other. A *Triple Quar'tan* is one in which there is a paroxysm every day; and on every three days the paroxysms correspond. A *redoub'ting or repeat'ing quar'tan*, (F.) *Fievre quarte doublee*, is one in which there are two paroxysms every fourth day. See *Intermittent*.

QUARTANUS, *Quartan*—q. Remittens, *Tetartophia*.

QUARTARIUS. An ancient measure, equal to one-fourth of the sextarius. See *Weights and Measures*.

QUARTE, ([L.] *quartus*, 'a fourth,') *Quartan*.

QUARTERON. (from [L.] *quatuor*, 'four,' a fourth part,) see *Mulatto*—q. *Black*, *Mulatto*.

QUAR'ISTERNAL, *Quadrister'nalis*. The fourth osseous portion of the sternum, corresponding to the fourth intercostal space.

QUARTIO, (*quatuor*, 'four, being foursided,') *Astragalus*.

QUASSATIO, (from *quassare*, *quassatum*, 'to shake,') *Conquassation*.

QUASSATURA, *Conquassation*.

QUASS'IA, (Ph. U. S.), (*Lignum Quassiae*, *Febrif'ugum lignum*, *Quassy*, (F.) *Quassie*, *Bois amer*. The wood of *Quassia amara*—and of *Q. excel'sa*, *Picro'na excel'sa*, *Bittera febrif'uga*, *Lofty Bitterwood Tree*, *Bitter ash*, *Simaru'ba excel'sa*, *Ord. Simarubaceae*; West India trees. The former is the *Jamaica Quassia*—the one in common use; the latter the *Surinam' Quassia*. *Quassia*, called after *Quassi*, a Surinam negro, who first employed it in the fevers of the country. It is an intensely durable bitter, without astringency; and has been employed as a tonic and stomachic. It is generally given in the form of infusion. The bitter principle is called *Quassin*, *Quassiin*, *Quas-eit* and *Bitterin*.

QUASSIA SIMAROU'BA, *Simarou'ba*, *Simaru'ba*, *S. officina'lis*, *Euc'onymus*, *Bitter Simaruba*, *Moun'tain Damson*. The bark of the root, *Simaru'ba*, (Ph. U. S.), is bitter, and not unpleasant. Both water and alcohol extract its virtues. It is not astringent. It is exhibited wherever a simple tonic is required.

QUASSIN, see *Quassia*.

QUASSIT, see *Quassia*.

QUATERNIO, (from *quatuor*, 'four,' having four sides,) *Astragalus*.

QUATRE ÉPICES, 'four spices,' see *Myrtus pimenta*.

QUATRIO, (from *quatuor*, 'four,' having four sides,) *Astragalus*.

QUEASINESS, Nausea.

QUEEN OF THE MEADOW, *Spiraea lobata*—q. of the *Prairie*, *Spiraea ulmaria*.

QUEEN'S DELIGHT, *Stillingia*—q. *Root*, *Stillingia*.

QUELLES, Elixir.

QUENOUILLE D'EAU, *Typha latifolia*.

QUERCERA, (from *καραειν*, 'to tremble,') *Epialos*.

QUERCINI HERBA, *Lichen plicatus*.

QUERCULA MINOR, (dim. of *quercus*,) *Teucrium chamædrys*.

QUER'CUS AL'BA, *White Oak*; QUER'CUS RO'BUR, *Com'mon Oak*; Q. TINCTO'RIA, *Black Oak*; Q. RU'BRA MONTA'NA, *Red Oak* or *Span'ish Oak*; Q. PEDUNCULA'TA, *Common British Oak*. *Ord. Cupuliferae*. *Soc. Syst. Monœcia Polyandria*; (F.) *Chêne*. The bark of all these varieties—*Quer'cus*—is, probably, alike in medicinal properties. It is powerfully astringent and tonic, and has been used internally in intermittents, and externally in decoction, as an astringent wash, where such is indicated.

The *A'corn*, (Prov.) *Ackerna*, *Nux Quer'che*, (F.) *Gland*, was at one time much used as food; and a decoction was once recommended in dysentery and diarrhoea, in glandular obstructions, &c. A pessary made of it has been advised in immoderate flow of the menses. It is not much used. *Roasted acorns*, (F. *Glands torréfiés*, have been used in the same cases as the decoction.

QUERCUS COCCIFERA, see *Kermes*.

QUERCUS INFECTO'RIA. A tree of Asia Minor, which affords *Tur'key* or *Nut Galls*, *Gall'æ Tur'cicae* seu *Quer'cinæ* seu *Tincto'riae*, *Nux Gall'æ*, *Galla maxima orbicul'ata*, *Ce'cia*, *Gall'æ*, *Galls*, *Gallnut*, (F.) *Noix de Galle*, *Galle de chêne*, *Cas-seoles*. The *nut-gall*—*Gall'æ* (Ph. U. S.)—is a morbid protuberance or tubercle, produced by the puncture of the *Diplo'lepis* seu *Cynips Gall'æ Tincto'riae*, *Cynips Quer'cûs fo'lii*. It is powerfully astringent and tonic, and has been employed in diarrhoea, intestinal hemorrhage, and intermittents: but is chiefly used, externally, in gargles, and injections. The powder, made into an ointment with lard, is used in piles. The ordinary nutgalls of the shops, are procured from the *Levant*. The large *Me'cea* or *Bus'sorah galls*, at times called *Dead Sea ap'ples*, *Mad ap'ples*, and *ap'ples of Sod'om*, are produced on *Quercus infectoria*.

There are upwards of thirty species of oak in the United States, all of which are probably more or less astringent and tonic.

QUERCUS MARINA, *Fucus vesiculosus*—q. *Suber*, see *Suber*.

QUERQUERA, (from *καραειν*, 'to tremble,') *Epialos*.

QUEUE DE CHEVAL, 'horse's tail,' (from [L.] *cauda*, 'a tail,') *Hippuris vulgaris*, *Cauda equina*, see *Medulla spinalis*—q. de la *Moëlle épinière*, *Cauda equina*—q. de *Pourceau*, *Peucedanum*—q. de *Rat*, *Sium lunare*.

QUICK-IN-THE-HAND, *Impatiens*.

QUICKEN TREE, *Sorbus aucuparia*.

QUICK'ENING, (Sax. *epiecan*, 'to make alive,') The period of gestation when the motion of the fœtus first becomes perceptible. This usually occurs about the eighteenth week, and was at one time erroneously believed to mark the time at which the fœtus becomes alive—*Animatio fœtus*. It need scarcely be said, that it possesses the principle of life from the moment of the union of the materials furnished by the sexes at a fecundating copulation. When the motion is felt, the female is said to be 'quick with child.'

QUICKENS, *Triticum repens*.

QUICKLIME, *Calx viva*.

QUICKSILVER, *Hydrargyrum*.

QUIES, 'rest,' *Acinesia*.

QUINA, (from *quinaquina*.) Quinine—*q. do Campo*, *Strychnos pseudoquina*—*q. Quina*, *Cinchona*.

QUINÆ ACETAS, SULPHAS, &c., see *Quinæ Acetas*, &c.

QUINCE, (from *quinsy* [?].) *Serofula*; see *Pyrus cydonia*, ([*F.*] *Coing*)—*q. Tree*, *Pyrus cydonia*.

QUINIA, (from *quinaquina*.) Quinine—*q. Acetate* of, *Quinæ acetas*—*q. Amorphous*, see *Quinia*, *extract of*, and *Chinioidine*—*q. Arseniate* of, *Quinæ arsenias*—*q. Arsenite* of, *Quinæ arsenis*—*q. Citrate* of, *Quinæ citras*.

QUINIA, EXTRACT OF, *Quinæ sulphas impurus*. A preparation which consists in evaporating the liquor poured off the crystals of sulphate of quinia, to the consistence of a pilular mass.

It has the properties of sulphate of quinia, and its active principle appears to be amorphous quinia or chinioidine. Twenty-four grains will generally arrest ague.

QUINIA, FERROCYANATE OF, *Quinæ ferrocyanas*—*q. Hydriodate* of, *Quinæ hydriodas*—*q. Hypophosphite* of, see *Calcis hypophosphis*—*q. Impure sulphate* of, *Quinia*, *extract of*—*q. Iodhydrate* of, *Quinæ hydriodas*—*q. Iodide* of, *Quinæ hydriodas*—*q. Iodide of iodhydrate* of, see *Quinæ hydriodas*—*q. Ioduret* of, *Quinæ hydriodas*—*q. and Iron*, *Hydriodate* of, see *Quinæ hydriodas*—*q. Lactate* of, *Quinæ lactas*—*q. and Mercury*, *protochloride* of, *Hydrargyri et Quinæ Protochloridum*—*q. Muriate* of, *Quinæ murias*—*q. Nitrate* of, *Quinæ nitras*—*q. Phosphate* of, *Quinæ phosphas*—*q. Sulphate* of, *Quinine*, *sulphate* of.

QUINIA AND CINCHONIA, *Tannate* of, *Quinæ et Cinchonie tannas*.

QUINIA ACETAS, *Chini'num aceticum*, *Acetas Chini'ni seu Chini'ni seu Chini'nicus seu Quinæ seu Quinæ seu Quini'ni seu Kini'ni, Quina aceticæ*, *Ac'etate of Quinia or of Quinine*. (Prepared by saturating *Quinia* with diluted *Acetic acid*.) Has the properties of the other salts of Quinia.

QUINIA ARSENIAS, *Chini'num seu Quinæ Arsenias*, *Chini'num Arsenico'bum seu Arsenic'icum*, *Arseniate of Quinia or of Quinia or of Quinine*, (*F.*) *Arséniate de quinine*. (Formed by the union of *arsenic acid* and *quinia*.) Employed as an antiperiodic in the dose of from three quarters of a grain to a grain and a half in the twenty-four hours.

QUINIA ARSENIS, *Q. Diar'senis*, *Ar'senite of Quinia*. (A salt resulting from the double decomposition of *arsenite of potassa* and *sulphate of quinia*.) It has been used in chronic cutaneous diseases; and as an antiperiodic in ague, neuralgia, &c. Dose, oneithird of a grain, three times a day.

QUINIA CITRAS, *Quinæ seu Chini'ni citras*, *Citras chini'nicus seu quini'nicus*, *Chini'num seu Chini'num Citricum*, *Cit'rate of Quinia or Quina or Quinine*. (Formed from the union of *citric acid* and *quinia*.) It has the same properties as the sulphate.

QUINIA DIARSENIS, *Quinæ Arsenis*.

QUINIA FERROCYANAS, *Chini'num ferrocyanogenatum seu fer'rohydrocyanicum*, *Ferrocyanate or hydrocyanof'er'rate of Quinia or of Quinine*. (This salt is obtained by the decomposition of *sulphate of quinia* by means of a solution of *ferrocyanuret of potassium*.) It has all the properties of sulphate of quinia, and, according to some, in a superior degree.

QUINIA HYDRIODAS, *Chini'num hydriod'icum*, *Hydriodate or Iodhy'drate of Quinia*; called, also, *Iodure'tum seu Iodidum Quinæ*, *Iod'uret* or *Iodide of Quinia*, (*F.*) *Hydriodate ou Iodhydrate ou Iodure de Quinine*. (This is formed by precipitating *sulphate of quinia* by means of *iodide of potassium*.) Given in serofulous affections; half a grain to a grain, three times a day, to a child.

An *iodide* and a *biniodide* have been formed;

the latter made by mixing double the quantity of *iodide of potassium* with *sulphate of quinia*.

AN IODIDE OF IODHY'DRATE OF QUINIA, (*F.*) *Iodure d'Iodhydrate de Quinine*, is prepared by pouring into an acid solution of *quinia* a solution of *iodide of iron*, containing a slight excess of iodine. The precipitate is treated with boiling alcohol: the liquor filtered, and crystals suffered to form. The salt possesses all the properties of the other salts of quinia. Under the name, *Hydriodate of Iron and Quinia*, a preparation was introduced by Mr. Battley. It is possessed of tonic and antiperiodic virtues.

QUINIA IODIDUM, *Quinæ hydriodas*—*q. Iodure'tum*, *Quinæ hydriodas*.

QUINIA LACTAS, *Quinæ lactas*, *Chini'num lacticum*, *Lac'tate of Quinia or of Quinine*, (*F.*) *Lactate de quinine*. (Made by the action of *lactic acid* on *quinia*.) Used in the same cases as the other salts of quinia.

QUINIA MURIAS, *Chini'num muriat'icum seu salitum seu hydrochloricum*, *Murias chini'nicus*, *Muriate of Quinia or of Quinine*. (May be prepared by dissolving *quinia* in dilute *muratic acid*.) Given as an antiperiodic. Dose, half a grain to a grain.

QUINIA NITRAS, *Chini'num seu Chini'num nitricum*, *Nit'rate of Quinia or of Quinine*, (*F.*) *Nitrate ou Azotate de quinine*. (May be formed by the addition of *nitric acid* to *quinia*.) Its properties are those of the sulphate of quinia.

QUINIA PHOSPHAS, *Chini'num seu Chini'num phosphoricum*, *Phosph'ate of Quinia or of Quinine*; (may be prepared by the addition of dilute *phosphoric acid* to *quinia*.) It is thought by many to be next to the sulphate of quinia in its remedial powers.

QUINIA SULPHAS, *Quinine*, *sulphate* of.

QUINIA VALERIA'NAS, *Chini'num seu Chini'num Valerian'icum*, *Vale'rianate of Quinia*, of *Quina*, or of *Quinine*; (may be formed by the union of *valerianic acid* with *quinia*.) It has the properties of the salts of quinia. Dose, gr. j to gr. v.

QUINIA ET CINCHONIA TAN'NAS, *Chini'num seu Chini'num tan'nicum*, *Tan'inate of Quinia and Cinchonia*. This salt has the same properties as the salts of quinia in general.

QUINICINE, *Quinici'na*, *Quinic'ia*. When a salt of quinia is exposed to heat with certain precautions, the alkaloid is changed into another isomeric with it, to which this name has been given.

Quinicine, and its sulphate, possess the same medical properties as quinia and its salts. Dose, gr. v to gr. xv.

QUINICUS, *Cinchonia*.

QUINIDIA, *Quinidi'na*, *Quinidine*, *Chini'dine*, *Cin'chotin*. An alkaloid of the cinchonas—of many of the genuine at least—obtained from them by the same process as that by which quinia is obtained from the barks that yield it. Its sulphate is more soluble however than the sulphate of quinia, and it is left, therefore, in the mother waters.

The salts of quinidia are markedly antiperiodic. From gr. x to gr. xxx will usually arrest a paroxysm.

QUININE', or QUI'NINE, (from *quinaquina*.) *Quinia*, *Quina*, *Kini'num*, *Ki'num*, *Sul'essentiale corticis Peruvia'ni*, *Chini'num*, *Quini'na*. An alkaline, encrystallizable substance; under the form of a porous, whitish mass; almost insoluble in water, soluble in alcohol and ether. It forms, with acids, salts that are in general soluble. It is obtained from different cinchonas, but chiefly from the yellow, and is the active principle of those valuable drugs.

QUININE, ACETATE OF, *Quinæ acetas*—*q. Arseniate* of, *Quinæ arsenias*—*q. Arsenite* of, *Quinæ*

arsenis—q. Azotite of, Quiniae nitras—q. Citrate of, Quiniae citras—q. Ferrocyanate of, Quiniae ferrocyanas—q. *Hydriodate de, Quiniae hydriodas*—q. *Iodhydrate de, Quiniae hydriodas*—q. Iodide of, Quiniae hydriodas—q. *Iodure de, Quiniae hydriodas*—q. *Iodure d'iodhydrate de, Quiniae, iodide of iodhydrate of*—q. Lactate of, Quiniae lactas—q. Muriate of, Quiniae murias—q. Nitrate of, Quiniae nitras—q. Phosphate of, Quiniae phosphas.

QUININE, QU'INA or QU'NIA, SULPHATE OF, *Qu'ine seu Qu'ina seu Qu'ina Sul'phas, Qu'ine Disul'phas, Chini'num seu Chini'num Sulphuricum, (F.) Sulfate de Quinine*—the salt usually employed in medicine—occurs in needles of a pearly and satiny appearance. It is employed with great success in the treatment of intermittents; and is available in many cases, where the bark in substance could not be retained, or would be injurious. Dose, as a tonic, from 3 to 10 grains in the 24 hours. As an antiperiodic it may be carried much farther. Its action, in a large dose, is decidedly sedative. (It is obtained, by treating the yellow bark with sulphuric acid. The following form is adopted in the Pharmacopœia of the United States.—*Cinchon. flav.* in pulv. crass. ℥iv; *Acid. mariat.* f℥ijj; *Calceis.* in pulv. ℥v; *Aqua*, cong. v; *Acid. Sulphur.*, *Alcohol.*, *Carbon animal.* āā q. s. Boil the bark in one-third of the water mixed with one-third of the muriatic acid, and strain through linen. Boil the residue twice successively with the same quantity of acid and water as before, and strain. Mix the decoctions, and, while the liquor is hot, gradually add the lime, previously mixed with two pints of water, stirring constantly, until the quinia is completely precipitated. Wash the precipitate with distilled water, and having pressed and dried it, digest it in boiling alcohol. Pour off the liquor and repeat the digestion several times, until the alcohol is no longer rendered bitter. Mix the liquors, and distil off the alcohol, until a brown viscid mass remains. Upon this substance, removed from the vessel, pour about half a gallon of distilled water, and having heated the mixture to the boiling point, add as much sulphuric acid as may be necessary to dissolve the impure alkali. Then add an ounce and a half of animal charcoal; boil for two minutes; filter the liquor while hot, and set it aside to crystallize. Should the liquor, before filtration, be entirely neutral, acidulate it very slightly with sulphuric acid. Should it, on the contrary, change the colour of litmus paper to a bright red, add more animal charcoal. Separate the crystals from the liquor, dissolve them in boiling water slightly acidulated with sulphuric acid; add a little animal charcoal, filter and set aside to crystallize. Wrap the crystals in bibulous paper, and dry with a gentle heat. The motherwaters may be made to yield an additional quantity of sulphate of quinia by precipitating the quinia with solution of ammonia, and treating the precipitated alkali with water, sulphuric acid, and animal charcoal, as before.)

QUININE, TANNATE OF, *Quiniae et cinchonae tannas*—q. Valerianate of, *Quiniae valerianas*.

QUININISM or QUININISM, *Quininis'mus, Qu'nis'm, Cin'chonism, (F.) Inébriation ou Ivesse quinique*. The aggregate of encephalic or neuropathic phenomena induced by overdoses of quinia.

QUINOIDINE, Chinioidine.

QUINIQUE, Cinchonice.

QUINIMUM. An alcoholic extract, made from

cinchona barks the composition of which is known, and which are mixed together in such proportions that the quinia and the cinchonina exist in the relation of two of the former to one of the latter. It is used in pills or in wine, and is said to represent, faithfully, all the useful principles of cinchona. To a similar preparation containing quinia and cinchonina, in the proportion of 91 of the former to 25 of the latter, M. Labarraque has given the name *Panchres'tum Chi'ne*.

QUINOLIN, (*quina*, and *laon*, 'oil,') Leukoleinum.

QUINOL'OGY, *Quinol'og'ia, Cinchonol'ogy, Cinchonol'og'ia, (quini, and laos, 'a discourse.')* A treatise on the cinchonas, or on quinia.

QUINQUEFOLIUM, (*quinque*, 'five,' and *folium*, 'a leaf,') *Potentilla reptans*—*Quinquefolium majus, Potentilla reptans*.

QUINQUINA, (*Peruvian kina kina*,) *Cinchona*—q. *Aromatique, Croton cascarilla*—q. *Bicolor, Cinchona Caribæa cortex*—q. *Faux de Virginie, Cusparia febrifuga*—q. *Gris de Loza, Cinchona lanceifolia cortex*—q. *Jaune, Cinchona cordifolia cortex*—q. *Piton, Cinchona Caribæa cortex*—q. *Rouge, Cinchona oblongifolia cortex*.

QUINQUINO, *Myroxylon peruvianum*.

QUINSY, *Cynanche, C. tonsillaris*—q. *Nervous, Angone*—q. *Wolf, Lycanche*.

QUINTAN, *Febris quinta'na seu pempt'a seu quin'ta*; (from *quintus*, 'a fifth,') A fever whose paroxysms return only every five days inclusively; and between which there are, consequently, three days of apyrexia.

QUINTE (F.), (formed, like *kink*, by onomatopœia [?].) This word is used synonymously with *paroxysm* or *fit*, when speaking of cough,—as *une Quinte de Toux*—'a fit of coughing,' (Sc.) *Kink*. Also, the influenza.

QUINTEFEUILLE. ([F.] *quinte*, 'a fifth,' and *feuille*, 'a leaf,') *Potentilla reptans*.

QUINTERON, (from *quintus*, 'a fifth,') see *Mulatto*—q. *black*, see *Mulatto*.

QUINTES, see *Quinte*, and *Pertussis*.

QUINTES'SENCE, *Quin'ta Essen'tia*, 'fifth essence.' A name formerly given to alcohol, impregnated with the principles of some drug. It was, consequently, often synonymous with *Alcohol'ic tincture*. The most volatile principles of bodies were, also, called *Quintessences*.

QUINTISTERN'AL. The fifth osseous portion of the sternum. It corresponds to the 5th and 6th intercostal spaces.

QUINUA, *Chenopodium quinoa*.

QUIZ'ZER, *Grinder*. One who examines and prepares students in medical colleges on the subjects taught in them.

QUOTIDIAN, *Quotidia'nus, Ephem'erus, (F.) Quotidien*—itself from *quotidiū*, 'daily,' (*quot*, 'every,' and *dies*, 'day,') That which takes place every day.

QUOTIDIAN FE'VER, *Q. a'gue, Fe'bris quotidiana seu hemere'sia seu hemer'ina seu ephen'era seu methemer'ina seu methemer'ia, An'etus quotidiana'nus, Cathem'erus, Cathemer'inus, Methemer'inus, Amphemer'inus, Amphimer'inus, Amphem'erus, Quotidia'nus, (F.) Fièvre quotidienne*, is an intermittent, the paroxysms of which recur every day.

A *sin'ple, doub'le*, or *trip'le quotidia'n*, is a quotidian, which has one, two, or three paroxysms in the 24 hours.

R.

R or R is placed at the commencement of a prescription as a contraction of *Radicis*. See Abbreviation.) Originally, it was the sign of Jupiter, and was placed at the top of a formula, to propitiate the king of the gods, that the compound might act favourably.

RABDOIDES. (SUTURA.) (ραβδοΐδης, from *ραβδος*, 'a rod,' and *αδός*, 'resemblance,') Sagittal suture.

RABID. *Rabidus*, (F.) *Robique*, *Rabétique*, (from *rabies*, *rabes*, 'I am mad or furious,') 'canine madness,') That which belongs to hydrophobia; as *rabid virus*, &c. *Rabid* also means affected with hydrophobia, (F.) *Enragé*.

RABIÉTIQUE. *Rabid*.

RABIES CANINA. Hydrophobia.

RABIQUE. *Rabid*.

RACAHOUT. *Amylum querneum*. A name given by the Arabs to the starch prepared from an edible acorn obtained from *Quercus Ilex*, but, according to Plagge, the so called *Racahout* of the Arabs, sold in Paris, is a compound of potatoe, starch, chocolate, and aromatics, — as vanilla.

RACCOON BERRY. *Podophyllum montanum*.

RACE. (from *radix*, ablative of *radix*, 'root,') *Breed*. The union or aggregate of individuals whose conformation or particular condition differs perceptibly from those of neighbouring varieties. In the human species, several races are designated. See *Homo*.

RACÉMEUX. *Racemosus*.

RACÉMO'SUS. *Rac'emosus*, *Rac'emo'se*, (F.) *Racemose*, (from *racemus*,) In clusters, like grapes.

RACE'MUS. A cluster—as of grapes.

RACHAMEL'CA. (Hebr. רחם *rechem*,) 'the womb,' and מלך *molech*, 'a king,') *Rechamel'ca*. A peculiar formative principle, supposed, by Doleus, to exist in the uterus.

RACHE. Porrigio.

RACHILÆL. *ῥαχίς*, 'the spine,') *Rhachitæ*.

RACHIALGIA. (*rachis*, and *αλγος*, 'pain,') Colic, metallic, *Rhachialgia*, *Rhachidynia*, *Vertebral disease*—*r. Mesenterica*, *Tabes mesenterica*—*r. Pictoniensium*, *Colic, metallic*—*r. Pictonium*, *Colic, metallic*.

RACHIALGIE MÉSENTÉRIQUE. *Tabes mesenterica*.

RACHIALGITIS. *Rachialgitis*.

RACHIDIAN. (from *rachis*,) *Spinal*—*r. Arteries*, *Spinal arteries*—*r. Bulb*, see *Medulla oblongata*—*r. Canal*, *Vertebral canal*.

RACHIDIEN. *Spinal*.

RACHIOPHYMA. *Rachiphyma*.

RACHIPHY'MA. gen. *Rachiphy'matis*, *Rhachiphy'ma*, *Rachiphy'gy'ma*, *Tu'mor Rach'i*, (*rachis*, and *ορμα*, 'a tumour,') A tumour on the spine or on the back.

RACHIS. gen. *Rach'idis*, (*ραχίς*, gen. *ραχίδος*,) *Vertebral column*.

RACHIS'AGRA. *Rhachis'agra*, *Rhachi'agra*, *Rhache'agra*, *Rhachid'agra*, *Rhachi'tis spina'lis*, (*rachis*, and *αγχα*, 'a seizure,') A gouty or rheumatic affection of the spine.

RACHITÆ. *Rhachitæ*.

RACHIT'IC. (*ῥαχίτις*,) *Rachit'icus*, *Rhachit'icus*, (F.) *Rachitique*. Relating or appertaining to, or affected with rachitis or rickets. Weak, feeble in the joints; defective in development.

RACHIT'IS. *Rhachit'is*. Inflammation of the spine. Also, *Cyrtosis Rad'chia*, *Cyrtosis*, *Morb'us Ang'licus*, *Osteomalacia Infantum*, *Tu'ber pector'ea*; *Immutu'io os'sium*, *Spina nodo'sa*,

Rachitismus, *English disease*, *Rickets*, (from *ῥαχίς*, 'the spine,' *ῥαχίτις*, *raχitis*, 'spine disease,') (F.) *Rachitisme*, *Novum*. A disease not known in England before the first quarter of the seventeenth century, and then termed *Rickets*, (probably from the Sax. *riec* or *hrec*, (Sc. and Prov.) *Rig*, all of which signify the back. Glisson first gave it the scientific name from *Rachis*, which, in Greek, also means the spine.) It is characterized by crookedness of the long bones; swelling of their extremities; crooked spine; prominent abdomen; large head; and often precocity of intellect. It is accompanied by leanness, general debility, indigestion; and frequently induces atrophy and hectic. *Rickets* occurs, particularly, during the first years of existence, in weakly children, born of rickety or scrupulous parents; brought up in damp or confined situations; who are badly nourished, and do not take sufficient exercise. The progress and termination of the disease are very variable. Some children gradually recover their health: others become more and more deformed, and ultimately die of consumption, dropsy, or other organic disease. The treatment is almost wholly hygienic. Pure air; a healthy situation; nourishing diet; exercise; sea or common cold bathing, and tonics, afford the best prospect of success.

RACHITIS ADULTORUM. *Mollities ossium*.

RACHITISME. *Rachitis*.

RACHITOME. (*rachis* and *τεμνω*, 'to cut,') An anatomical instrument for opening the medulla spinalis without injuring the medulla.

RACHO'SIS. *Rhach'o'sis*, *Rach'o'sis* (from *ῥαχίς*, or *ραχίς*, 'I wrinkle,' and *osis*,) A relaxation of the serotum. *Propt'oma seu Lux'io Seroti*, *Scrotum pendulum*. An excoriation of the relaxed pendulum.

RACINE. *Radix*—*r. à Becquet*, *Geranium moschatum*—*r. de Bengal*, *Cassumunar*—*r. Bré'silienne*, *Ipecacuanha*—*r. de Characis*, *Dorstenia contrayerva*—*r. de Chine*, *Sinilax china*—*r. de Dracæ*, *Dorstenia contrayerva*—*r. des Philippines*, *Dorstenia contrayerva*—*r. de Saint Esprit*, *Angeica*—*r. Salsolaire*, *Anthemis pterithrum*—*r. de Turbith*, *Convolvulus turpethum*.

RACINES DU CERVELET. (pl. of *racine*,) *Corpora testiformia*.

RACK. *Arack*.

RACLURES DES BOYAUX. (from *racler*, 'to scrape,') see *Abrasion*. *Ramenta intestinum*.

RACORNISSEMENT (F.), (*racornir*, ([L.] *cornu*, 'horn,') 'to make hard as horn,') *Cor'ni'fication*. A physical property, possessed by animal substances, which consists in a kind of contractility, accompanied with sudden corrugation and curling. It is produced by heat, and by chemical agents, especially by the strong mineral acids.

RACOSIS. (*ῥαχίσις*,) *Rachosis*.

RADCLIFFE'S ELIXIR, see *Tinctura aloes composita*.

RADESYGE. *Radzyge*.

RADEZYGE. *Radzyge*.

RADIAD. see *Radial aspect*.

RA'DIAL. *Radia'lis*, (from *radius*,) That which relates to the radius.

RADIAL ANTERIEUR. *Palmaris magnus*.

RADIAL ARTERY. *A. externa Cubiti*. It arises from the brachial, at the upper and anterior part of the forearm; descends on the outer side as far as the lower part of the radius, and proceeds downwards and backwards towards the superior extremity of the space which separates the first

two metacarpal bones. It then buries itself in the palm of the hand, where it forms the *profund* or *radial palmar arch*. The radial artery gives off the *recurrent radial*, several *radiomuscular* branches, the *transverse palmar radiocarpal artery*, the *external superficial artery of the palm of the hand*, the *external dorsal of the thumb*, the *dorsalis carpi*, *dorsal interosseous artery of the index*, &c.

RADIAL ASPECT. An aspect towards the side on which the radius is situated.—(Barclay.) *Radial* is used by the same writer, adverbially, to signify 'towards the radial aspect.'

RADIAL EXTERNE PREMIER, see *Radialis*—*r. Externe second*, see *Radialis*—*r. Grand*, see *Radialis*.

RADIAL NERVE, *Radio-digital* (Ch.), *Spiral Nerve*, *Spiral muscular N.*, *Musculospiral nerve*, arises from the four inferior branches of the brachial plexus. It is, at first, situate behind the other nerves of the plexus. It then becomes engaged between the three portions of the triceps brachialis, passes behind the humerus, and descends between the supinator longus and brachialis internus, as far as opposite the upper extremity of the radius. In its course it gives numerous filaments to the triceps, supinator longus, extensor carpi radialis longior, and to the integuments. Opposite to the upper extremity of the radius it divides into two branches;—the one *posterior*, which turns backwards into the substance of the supinator brevis, and when it has reached the space between the two layers of muscles on the posterior surface of the forearm, divides into a great number of filaments, which are distributed to those muscles, and to the integuments of the hand. The other branch is *anterior*: it descends before the supinator brevis and the radius; until, near the inferior third of that bone, it engages itself between the tendons of the supinator longus and extensor carpi radialis longior; and, becoming afterwards subcutaneous, divides into two branches, whose filaments are distributed to the integuments of the thumb, index, middle finger, to the outside of the ring-finger, and to the first interosseous muscle of the back of the hand.

RADIAL PETIT, see *Radialis*.

RADIAL VEINS, DEEPSEATED. These follow the same course as the radial artery.

RADIALIS EXTENSOR LONGIOR, *R. externus longior*.

RADIALIS EXTERNUS BREVIOR, *Extensor Carpi Radialis Brevior*, *Radialis secundus*, (F.) *Second radial externe*, *Petit radial*, *Épicondylus-métacarpien*, (Ch.) An extensor muscle of the wrist, situate beneath the extensor carpi radialis longior. It is attached, above, to the external tuberosity of the humerus, and terminates, below, by a long tendon, inserted into the posterior part of the upper extremity of the third bone of the metacarpus. It has the same uses as the next muscle.

RADIALIS EXTERNUS LONGIOR, *Extensor Carpi Radialis Longior*, *Radialis externus primus*, (F.) *Premier radial externe*, *Grand radial*, *Humérus-métacarpien*, (Ch.) It is seated at the outer part of the forearm; is thin, but thicker on the outside than on the inside. It is attached, above, to the inferior part of the outer edge, and to the corresponding tuberosity of the humerus; and terminates, below, by a long tendon, which is attached to the superior extremity of the second bone of the metacarpus. It extends the hand on the forearm.

RADIALIS EXTERNUS PRIMUS, *R. Externus longior*—*r. Internus*, *Palmaris magnus*—*r. Secundus*, *R. Externus longior*.

RADIATED, *Radialis*, (from *radius*, 'a ray,')

(F.) *Rayonné*. That which is arranged in *rays*; in lines, diverging from a common centre. An epithet given to several ligaments, &c.; as to those which unite the ribs to the sternum; those which unite the tibia and fibula at their inferior extremity, &c.

RADIATED SUBSTANCE OF THE KIDNEY, see *Kidney*.

RADICAL, *Radicalis*, (from *radix*.) A *radical cure*, *cura radicalis*, is one in which the disease is destroyed, as it were, from the root. It is used in opposition to *pal/iative cure*.

RADICAL MOISTURE, *Humidum radicale*.

RADICAL VESSELS, *Vascular Radicles*, (F.) *Radicules vasculaires*, are the small vessels that take their origin in the tissues, and by their union form larger vessels.

RADICAL VINEGAR, *Aceticum acidum*.

RADICES HYOIDEI OSSIS, (pl. of *radix*.) *Cornua hyoidei ossis*.

RADICIS/ECA, *Rhizotomus*, *Rhizotomist*, (*radix* and *secare*, 'to cut.') One employed in collecting and preparing roots was formerly so called.

RADICLE, *Radicula*.

RADICULA, *Radicle*, (F.) *Radicule*, (dim. of *radix*, 'a root.') A small root or rootlet; also, *Raphanus hortensis*.

RADICULES VASCULAIRES, *Radical vessels*.

RADII CILIARES, (*radius*, 'a ray,' pl. *radii*.) *Ciliary processes*.

RADII FRONDIS. The folds or wrinkles of the forehead.

RADIO, (from *radius*.) In composition the *radius*.

RADIOCARPAL, *Radiocarpia/nnus*. That which relates to the radius and carpus.

RADIOCARPAL ARTICULATION is the *wrist joint*, or articulation of the os scaphoides, os semilunare, and os cuneiforme of the carpus with the inferior surface of the radius, and the fibro-cartilage, situate below the ulna. It is strengthened by lateral, anterior, and posterior ligaments.

RADIOCARPAL TRANSVERSE PALMAR ARTERY, (F.) *Artère Radio-carpienne-transversale-pal-maire*. A transverse branch, given off by the radial artery, opposite the lower edge of the pronator quadratus, which sends several branches to the anterior or palmar surface of the carpus.

RADIO-CUTANÉ (NERF), see *Cutaneous*—*r. Phalangettien du ponce*, *Flexor longus pollicis manus*.

RADIOLUS, (dim. of *radius*, 'a rod or spoke.') *Sound*.

RADIOMUSCULAR, *Radiomuscula/ris*. A name given to the branches sent off from the radial artery to the muscles of the forearm in the first part of its course; as well as to the nervous filaments which the radial nerve sends to the same muscles.

RADIS, (*radix*.) *Raphanus hortensis*—*r. de Cheral*, *Cochlearia armoracia*.

RADISH, *Raphanus hortensis*—*r. Water*, *Nasturtium amphibium*.

RADIUS, 'a staff,' 'a rod,' 'a spoke:'—so called from its shape. *Cer/cis*, *Foc/ile m/nnus seu sup/erius*, *Con/na m/nnor*, *Os ad cubitallē*, *Additamentum ul'næ*, *Manubrium m/nnis*, *Parap/chyon*, *Arum/do bra/chii m/nnor*, (F.) *Os du Rayon*. A long, prismatic bone, the upper and lesser extremity of which is called the *head*. This is supported by a *cer/vix* or *neck*. At the part where the neck is confounded with the *body* of the bone is the *tu/bercle* or *bicipital tuberosity* or eminence for the insertion of the biceps. The radius is articulated, above, with the os humeri and with the lesser sigmoid cavity of the ulna: below, with the scaphoides, semilunare, and the head of the ulna. Its inferior extremity, which is larger than

the superior, is flattened before and behind; is irregularly quadrilateral; and has, below, a double facette to be articulated with the first two bones of the carpus. On the outer side is the styloid process; and, on the inner, a concave facette, which joins the ulna. Behind, are grooves for the passage of the extensor tendons. The radius is developed from three points of ossification; one for the body, and one for each extremity.

RADIUS, COL DU, Collum radii.

RADIX, gen. *Radi'cis, Rhi'za, (F.) Racine.*

♂ root; also, the male organ.

Radix, Root or fang of a tooth, is the part contained in the alveolus. The *radix* or *root* of the nail is the portion hidden under the skin. Surgeons give the name to prolongations, sent by scirrhus, cancerous, or other tumors into the neighboring parts.

Five Aperient Roots, Quinque radi'ces aperien'tes, were, of old, asparagus, butchers' broom, fennel, parsley, and smallage.

Five Lesser Aperient Roots, Quinque radi'ces aperien'tes mino'res, were, caper, dandelion, eryngo, madder, and restdarrow.

RADIX ASCLEPIADIS CRISPA, see *Gomphocarpus crispus*—*r. Braziliensis, Ipecacuanha.*

RADIX COR'DIS, Ba'sis Cordis. The base of the heart.

RADIX GINSENG, see *Panax quinquefolium*—*r. Indica* Lopeziana, *Lopez radix*—*r. Linguae,* see *Tongue*—*r. Lopeziana, Lopez radix*—*r. Ninsi, Panax schinseng, Sium ninsi*—*r. Rubra, Rubia*—*r. Serpentinum,* see *Ophiorrhiza mungos*—*r. Unguis, Nail, root of*—*r. Ursina, Æthusa meum*—*r. Ventris, Umbilicus.*

RADULA, (from *radere, 'to scrape,')* Raspatorium.

RADZY'GĖ, *Radzygin, Radezyge, Radezyge,* (from obsolete *Da* word *rada, 'malignant,'* and *ayge, 'disease,')* *The'ria, (properly The'ria, from θηρία (thēria), 'a malignant ulcer,')* *Le'pra borea'lis seu Norveg'ica, Norwe'gian Lep'rosy.* A name given, in Norway, to a disease, bearing considerable analogy to the yaws. Some have esteemed it a species of lepra or elephantiasis, and others a syphiloid affection.

Another form—the *Spedalske, or Spedalskhd*—of Norway, appears to be a variety of tubercular lepra. Under the epithelial scales, numerous acari have been found; and the affection has been called *Scab'ies crusta'sa, and Scab'ies Norveg'ica Boeck'ii,* after Boeck, who well described a case of it.

RAFFLE'SIA ARNOLD'I, (after Sir Stamford Raffles.) A plant of Java. Order, *Rafflesiaceæ,* a decoction of which is employed there as an astringent in relaxed conditions of the genito-urinary apparatus.

RAFAÏCHISSANT, (re, and fraîche, 'cool,') from [L.] *refrigerescere, 'to become cool again,')* Refrigerant.

RAGE, I'ra, Fu'ror bre'vis, Or'gê, Thy'mos, (F.) Col'ere, Rage. Violent passion, characterized by contraction of the muscles of the face, violence in every movement, extreme irritation of the nervous system, acceleration of the blood's motion, and, generally, redness and swelling of the face.

RAGE, (payn,) Rhage.

RAGE, (from [L.] rabies,) Hydrophobia, *Rage.*

RAGE DE TÊTE. An epidemic and fatal malady, which prevailed in France in 1481. It was presumed to be epidemic meningitis.

RAGLE (F.), (said to be derived from an Arabic word, which signifies 'to traverse the desert.')

A kind of hallucination observed in travellers in the deserts, which resembles the effect of intoxicating liquors, opium, &c.

RAGWEED, Ambrosia elatior—*r. Golden, Senecio aureus*—*r. Great, Ambrosia trifida.*

RAGWORT, Senecio Jacobæa.

RAIA BATIS, and R. CLAVATA, see *Oleum jecoris aselli.*

RAIDEUR, (from rigidus, 'stiff,') Rigidity—*r. Cadavérique, Rigor mortis.*

RAIFORT, (from raphanus, or from radix fortis, or from rapum forte, [?]) Raphanus hortensis—*r. Savage, Cochlearia armoræica.*

RAINBOW WORM, Herpes iris.

RAINURE, Groove—*r. Digastrique, Mastoid groove.*

RAISIN, 'a grape,' (from *racemus, [?]*) see *Vitis vinifera*—*r. d'Amérique, Phytolacca decandra*—*r. de Bois, Vaccinium myrtillus*—*r. d'Ours, Arbutus uva ursi*—*r. de Renard, Paris.*

RAISING, Expectoration.

RAISINIERE, (F.) (from raisin,) A name given to a small granular, brownish or blackish tumour, which forms occasionally at the surface of the cornea, in consequence of ulcers or accidental wounds of that membrane.

RAISINS, Uvæ passæ, see *Vitis vinifera*—*r. de Corinthe,* see *Vitus Corinthiaca.*

RAISON, ([L.] ratio,) Reason.

RAKASI'RA. An American balsam; of a brownish or brownishred colour; semitransparent; brittle; softening and becoming tenacious by heat; inodorous when cold, but exhaling a very agreeable smell when heated; and possessing a balsamic and slightly bitter taste. The precise vegetable that furnishes it is not known. It has been used chiefly as a balsam in gonorrhœa and urinary affections.

RAKETEETH. A term, used in the north of England for teeth which are wide apart, and similar to those of a rake.

RAKIA, see Spirit.

RÂLE, (F.) *Rat'tle, Rhon'chus, Rhen'chus, Rhén'is, (from [D.] Ratel,)* *Cerc'h'nus, Rhog'mus* (probably an example of onomatopœia). Noise produced by the air in passing through mucus, of which the lungs are unable to free themselves. This condition, *Râle trachéal,* is chiefly observed at the approach of death, and is commonly called "the rattles."

The term *Râle* has been given to different sounds during respiration, caused by the air in connection with fluid contained in the bronchi or areolæ of the pulmonary tissue; which are perceived by the stethoscope. *Râle* has generally been used for a 'moist' sound, heard on auscultation of the chest, whilst *rhonchus* has meant a 'dry' sound.

RÂLE BRONCHIQUE HUMIDE, R. muqueux—*r. Caverneux, Gurgling*—*r. Bronchique sec, Râle Sonore*—*r. Cavernuleux,* see *Gurgling*—*r. Crépitant, Rhonchus crepitans,* see *Crepitant*—*r. des Mouvans,* see *Râle*—*r. c. Reduz,* see *Crepitant.*

RÂLE MUQUEUX, R. bronchique humide, Rhon'chus muc'o'sus, Hygrobronchiorrhon'chus, Mu'cous Rat'tle. The sound produced by the passage of air through mucus accumulated in the larger bronchi or trachea, or through softened tubercular matter. This *râle* occurs in catarrh, and in softened tubercle. When carried to a very high degree, it constitutes *gurg'ling, (F.) Gargouillement.* The *subcrepitant rhon'chus, (F.) Râle sous-crepitant,* is produced by the bubbling of air through liquid of variable consistence in the minute bronchial tubes. When heard at the base of both lungs, it is a sign of capillary bronchitis. When heard at the base of one only, it is generally connected with tubercles higher up.

RÂLE SIBILANT, Rhon'chus sib'ilans seu sib'ilans, Sib'ilant Rat'tle. A slight, though prolonged, whistling sound, occurring either at the commencement of inspiration or of expiration, owing to the presence of mucus, thin and viscid, but not abundant, which obstructs, more or less completely, the smaller bronchial ramifications.

It is seated in the small tubes, and occurs in the first stage of bronchitis. The *clicking* and *whistling* varieties differ, in the former being short, the latter prolonged. The former is a quick, sharp sibilus or whistle, which ceases almost instantaneously; the latter a prolonged sibilus, of less sharpness, lasting almost the whole time of the movement which it accompanies. To these may be added the *hissing* variety, the name sufficiently indicating its character.

RÂLE SONORE, *R. bronchique sec ou ronflant*, *Rhonchus sonorus*, *Sonorous Rattle*. A sound resembling at times the snoring—(F.) *Ronflement*—of a person asleep; at others the sound of the bass string of an instrument when rubbed with the finger; and not unfrequently the cooing—(F.) *Roucoulement*—of a dove. It seems to be caused by a contraction of the bronchial tubes, and is characteristic of chronic catarrh.

RÂLE SOUS-CRÉPITANT, see *Rôle muqueux*—*r. Trachéal*, see *Rôle*—*r. Vésiculaire*, see *Crepitation*.

RAMEAU, Ramus.

RAMBERGE, Mercurialis.

RAMEN'TA INTESTINORUM, (pl. of *ramentum*), *Scrapings of the Intestines*, (F.) *Rachures des Boyaux*. The shreds or scrapings, as it were, of the mucous membrane of the bowels, often discharged in malignant dysentery. The evacuation in which these are contained has been termed *Diachorema xymatodes*.

RAMENTUM, (contracted from *radimentum*, from *radere*, 'to scrape,') Fragment, Scabs.

RAMEX, gen. *Ramixis*, Hernia—*r. Varicosus*, Varicocele.

RAMIFICATION, *Ramificatio*, (*ramus*, and *facere*, 'to make,') A division of arteries, veins, or nerves into branches. Also, a branch itself.

RAMILLE, (from *ramus*), Ramusculus.

RAMOLLISSEMENT, (*re*, and *mollir*, (from [L.] *mollis*, 'soft,') 'to soften,') Mollities—*r. du Cerveau*, Mollities cerebri—*r. du Cœur*, Cardiomalacia—*r. de l'Estomac*, Gastromalacia—*r. du Foie*, Hepatomalacia—*r. Gris*, see Hepatization—*r. de l'Intestin*, Enteromalacia—*r. de la Moelle épinière*, Mollities medullæ spinalis—*r. des Os*, Mollities ossium—(F.) *Rouge*, see Hepatization.

RAMPANT, (*r.*), *ramper*, 'to climb,' from [L.] *repere*, 'to creep,') *Répens*, *Rep'tans*. That which creeps or crawls. The French *Bandage rampant* has the folds arranged spirally around the part, leaving intervals between them. It is a retentive bandage, but is easily displaced, and, therefore, not employed.

RAMPE, Scala.

RAMPES DU LIMAÇON, *Scalæ* of the cochlea.

RAMPHOS, (ῥαμφος, 'a beak,') Rostrum.

RAMULUS, (dim. of *ramus*), Ramusculus.

RAMUS, *Cladus*, Branch, (F.) *Rameau*. A division of an artery, vein, or nerve. Also, the male organ.

RAMUS DURIOR SEPTIMÆ CONJUGATIONIS, Facial nerve.

RAMUSCULUS, (dim. of *ramus*), *Ramusculus*, *Ramusculus*, *Surculus*, *Cladiscus*. A division of a ramus. (F.) *Ramuscule*, *Ramille*.

RANA, *Ranula*.

RANA ESCULENTA, 'esculent frog,' (F.) *Grenouille*. This frog and the bull-frog, *Rana Taurina*, are eaten as delicacies. They are nutritious, and easily digested.

RANCE, Rancid.

RAN'CID, *Ran'cidus*, *Sa'pros*, (F.) *Rance*, (*rancere*, 'to be stale,') An epithet given to fatty bodies, which have become acrid from age or the contact of air. Mild ointments may thus become very irritating, and unfit for the purposes for which they were originally intended.

RAN'DIA DUMETORUM, 'R. of thickets,'

(after J. Rand, a botanist of London,) *R. Spino'sa*, *Canthium Coronatum*, *Gardenia* seu *Posoqueria Dumetorum* seu *Spino'sa*. A Coromandel shrub; Ord. Cinchonaceæ; the fruit of which, bruised and thrown on the water, intoxicates, or even kills fish. In the form of powder it is a powerful emetic.

RANDIA SPINOSA, *R. dumetorum*.

RAN'INE, *Ran'ina*, (from *rana*, 'a frog,' 'shaped like a frog,') The portion of the lingual artery which advances horizontally between the genioglossus and lingualis muscles as far as the point of the tongue. The ranine vein follows the same course as the artery, and opens into the internal jugular or into the superior thyroid.

RANTREE, *Sorbus aucuparia*.

RAN'NULA, (dim. of *rana*, 'a frog,' so called from its resemblance,) *Dysphagia Ran'ula*, *Emphragma salivare*, *Batrachos*, *Batrachium*, *Hydroglossa*, *Hypoglossis*, *Hypoglossum*, *Hypoglossitis*, *Hypoglossium*, *Ra'na*, *Frog-tongue*, (F.) *Grenouillette*. A small, soft, fluctuating, and semitransparent tumour, which forms under the tongue, owing to the accumulation of saliva in Wharton's duct. In order to give exit to the accumulated fluid, and to establish a permanent opening for the discharge of the saliva, the cyst may be punctured, and the opening maintained by a minute cylinder, with a button at each extremity, as has been recommended by Dupuytren; or a portion of the cyst may be removed by the scissors, and the mouth be frequently washed by any astringent lotion. Sometimes the salivary duct is found plugged by a small calculus.

RANULA LAPIDEA, see *Calculi*, salivary.

RANUNCULUS, *Crowfoot*, (F.) *Renoncule*, (Sc.) *Yell'low grow'nd*, Ord. Ranunculaceæ; *Sec. Syst.* Polyandria Polygynia, (dim. of *rana*, 'a frog,' because found in marshy places where frogs abound.) A genus of plants, most of the species of which, when taken internally, act as acrid poisons. They are, also, acrid and vesicant when applied to the skin, and have accordingly been often used as counter-irritants. In the Pharmacopœia of the United States, *Ranunculus* is the cormus and herb of *Ranunculus bulbosus*. The following species answer to the above description.

RANUNCULUS ABORTIVUS, *Smallflowered Crowfoot*, *Chick'enpepper*.

RANUNCULUS ACRIS, *R. pratensis* seu *Steven'sii* seu *ru'fus* seu *ey'vaticus* seu *Sic'ulus*, *Up'right Meadow* or *Tall Crowfoot*, *Bul'ter Cups*, (Prov.) *Crai'sey*, *Guill'y Cups*, *Yell'lowweed*, *Blis'terweed*, *Pile'wort*, *Bur'wort*, *Mead'owbloom*, *Yell'wows*, (F.) *Bouton d'or*.

RANUNCULUS ALBUS, *Anemone nemorosa*—*r. Aquaticus*, *R. scleratus*.

RANUNCULUS BULBOSUS, *R. la'tus*, *Bul'bous* or *Bulbousrooted Crowfoot*, *Bul'ter Cups*, (F.) *Renoncule bulbeuse*, *Bassinet*.

RANUNCULUS DIGITATUS, *R. scleratus*—*r. Declinatus*, *R. flammula*.

RANUNCULUS FICARIA, *R. ver'nus* seu *pre'cox*, *Chelidonium* mi'nus, *Scrophularia* mi'nor, *Ficaria ranunculoïdes* seu *ver'na* seu *commu'nis*, *Chelidonia rotundifolia* mi'nor, *Cursu'ma hemorhoidalis* her'ba, *Egole'thon*, *Les'ser cell'andine*, *Pile'wort*, (F.) *Petite Chélidoine*, *Petite Éclairé*, *Ficaire*, *Herbe aux hémorrhôides*. The leaves and root have been used medicinally; the leaves as an antiscorbutic; the root as a cataplasm in piles.

RANUNCULUS FLAMMULA, *R. declinatus*, *Flam'mula*, *Surree'ta al'ba*, *Small'er Wa'ter Crowfoot*, *Less'er Spear'wort*, (F.) *Petite Douve*.

RANUNCULUS ILLYRICUS.

RANUNCULUS LINGUA.

RANUNCULUS LÆTUS, *R. bulbosus*—*r. Nemorosus*, *Anemone nemorosa*—*r. Palustris*, *R. scleratus*.

RANUNCULUS PENNSYLVANICUS, *Bristly Crow-foot*.

RANUNCULUS PRÆCOX, *R. ficaria*—1. *Pratensis*, *R. acris*.

RANUNCULUS PUBESCENS. Grows in ditches and marshes in South Africa. The expressed juice is used by the Cape Colonists in cancerous ulcers.

RANUNCULUS REPENS, *Creeping Crow-foot*.

RANUNCULUS RUPELUS, *R. acris*.

RANUNCULUS SCCLERATUS, *R. Palustris* seu *umbellatus* seu *digitatus* seu *aquatiscus*, *Rizus A'pium*, *Her'ba Sardo'nia*, *Her'ba Scclerata*, *Hecato'nia palustris*, *Marsh or Crush'ed or Cel'ery-leaved Crow-foot*.

RANUNCULUS SICULUS, *R. acris*—*r. Stevenii*, *R. acris*—*r. Umbellatus*, *R. sccleratus*—*r. Vernus*, *R. ficaria*—*r. Virginianus*, *Lobelia syphilitica*.

RAPA, *Rapum*, *Rapus*, *Rapys*, *Raphis*, *Raphus*, *Raphanus*, *Raphys*, *Rhaphis*, *Rhaphus*, *Rhaphanus*, *Rhaphys*, and *Rhaphys*. These terms were applied, of old, to the cabbage or turnip kind.

RAPA NAPUS, (*parvus*), *Brassica rapa*—*r. Rotunda*, *Brassica rapa*.

RAPE, (*rapa*), *Brassica napus*: also, *Stuprum*, (from *rapio*, 'to ravish'.)

RAPHANE'DON, (*ῥαφανιον*, from *ῥαφας*, 'a radish, like a radish.') *Rhaphanedon*. A transverse fracture of a long bone, leaving no splinters; what has been called, in France, *Fracture en rave*.

RAPHAN'IA, *Convulsio Raphania*, *Raphania*, *Mor'bus spasmodicus malignus* seu *popul'aris* seu *Silosis*, *Eclampsia typhoides*, *Synclonus Raphania*. Linnæus gives this name to a convulsive disease, not uncommon in Germany and Sweden, and which has been attributed to the seeds of the *Raphanus Raphanistrum* seu *sylvestris*, *Raphanistrum seg'etum* seu *arven'se*, *Raphistrum*, being mixed with the corn. The convulsions are seated in the limbs, and are attended with acute pain. See *Convulsio cerealis*.

RAPHANIA MAÏZTICA, *Pellagra*.

RAPHANIS, (*ῥαφανισ*.) *Raphanus hortensis*—*r. Magna*, *Cochlearia armoracia*.

RAPHANISTRUM ARVENSE, see *Raphania*.

RAPHANUS, *Rapa*, *R. hortensis*.

RAPHANUS HORTENSIS, *Raphanus*, *R. sativus* seu *n'ger*, *Raph'anis*, *Rhaph'anus*, *Radic'ula*, *Rha'phys*, *Ra'phys*, *Rhapa*, *Ra'pys*, *Rad'ish*, (*F.*) *Raifort*, *Radis*. *Ord. Cruciferae*. *Sez. Syst. Tetradynamia Siliquosa*. The root of this plant is a common esculent. It has, also, been given as a diuretic in calculous affections. The juice, made into a syrup, is exhibited to relieve hoarseness. Radishes have long been celebrated as antiscorbutics.

RAPHANUS MARINUS, *Cochlearia armoracia*—*r. Raphanistrum*, see *Raphania*—*r. Rusticanus*, *Cochlearia armoracia*—*r. Sylvestris*, see *Raphania*.

RA'PHÊ, *Rha'phê*, (*ῥαφή*, 'a suture.')

A name given to prominent lines resembling a raised stitch. Such is the *raphe* or *tramis* which divides the scrotum into two parts, and which extends from the anterior part of the anus to the extremity of the penis. The name *Ra'phê* seu *Ra'phê super'ior* seu *Lin'ea Median'a* seu *Sutu'ra corporis callosi ext'erna* seu *chor'da longitudinalis Lancisii* is, also, given to two prominent lines which are longitudinal and parallel, and are seen at the upper surface of the corpus callosum, extending from the anterior to the posterior extremity of that commissure.

RAPHEX, *Chenopodium vulvaria*.

RAPHIANKISTRON, (*ῥαφιον*, 'a needle,' and *αγκιστρον*, 'a hook.')

An instrument employed in the formation of artificial pupil.

RAPHION, (*ῥαφιον*, dim. of *ῥαφης*, 'a needle,') *Needle*.

RAPHIS, (*ῥαφης*.) *Needle*, see *Rapa*.

RAPHUS, *Rapa*.

RAPHYS, *Rapa*, *Raphanus hortensis*.

RAPISTRUM, (from *rapa*.) see *Raphania*.

RAPONTIC, *Rheum rhaiponticum*.

RAPPORT (*F.*), (*re*, and *portare*, 'to bring,') *Relat'io*, *Renunciat'io*, 'a report.' A deposition made by a physician or surgeon, to prove the condition of a person, the nature of a disease, the existence or non-existence of pregnancy, sudden or violent death, &c. Formerly, there were the *Rapports dénonciatifs*, made at the desire of the injured or their friends, and intended to point out to the judges the details of the crime or misdemeanor,—the *Rapports provisoires*, the object of which was to obtain provisions for the injured individual, both as regarded food and medicine, and to defray the expenses of prosecution,—and *Rapports mixtes*, or those which were both *dénonciatifs* and *provisoires*. At present, reports or depositions are generally divided into *judiciary*, or those which serve to throw light on the subject matter of any civil or criminal cause; and *administrative*, which afford information on subjects relating to any branch of the public administration, as on the nuisance, or otherwise, of certain establishments, the character of any epidemic, &c.

RAPPORT, *Eructation*, *Regurgitation*.

RAP'TUS, (*rapio*, *raptum*, 'to seize violently and suddenly.')

A violent and sudden attack or seizure.

RAPTUS HEMORRHAG'ICUS. A hemorrhagic attack or seizure.

RAPTUS NERVO'RUM. A sudden and violent nervous seizure. A spasm or cramp.

RAPTUS POSTERGANÆUS, (*rapitus*, *post*, 'behind,' and *tergum*, 'the back,') *Opisthotonus*.

RAPUM, *Rapa*—*r. Majus*, *Brassica rapa*.

RAPUNTIIUM SYPHILITICUM, (from *rapum*.) *Lobelia syphilitica*.

RAPURE, *Rasura*.

RAPUS, *Rapa*.

RAPYS, *Rapa*, *Raphanus hortensis*.

RAQUETTE, *Cactus opuntia*.

RARE (*F.*), *Rarus*, 'rare.' An epithet for the respiration and pulse, when less frequent than in health.

RAREFA'CIENS, *Rarifa'ciens*, *Aræol'icus*, (*F.*) *Raréfiant*, (*rarus*, 'rare, thin,' and *facere*, 'to make.')

Medicines were so termed, which were supposed to be possessed of the power of giving more bulk or extension to the blood or other humours.

RARÉFIANT, *Rarefaciens*.

RARIFACIENS, *Rarefaciens*.

RA'RITAS, (*F.*) *Raréte*. *Rarity*; scantiness; fewness.

RARITAS DEX'TIUM. A large space between the teeth, which appear to be, and may be, too few.

RASCATIO, *Excretion*.

RASCETA, *Carpus*, *Malleolus*—*r. Pedis*, *Tarsus*.

RASCHA, *Carpus*, *Malleolus*.

RASETA, *Carpus*, *Malleolus*.

RASETTA, *Carpus*, *Malleolus*.

RASH, *Exanthem*—*r. Canker*, see *Cynanche maligna*—*r. Fever*, *Scarlatina*—*r. Lichenous*, *Lichen*—*r. Mercurial*, *Eczema mercuriale*—*r. Mulberry*, see *Typhus*—*r. Nettle*, *Urticaria*—*r. Rose*, *Roseola*—*r. Summer*, *Lichen tropicus*—*r. Tooth*, *Strophulus*—*r. Wildfire*, *Ignis sylvaticus*.

RASION (*F.*), *Ra'sio*, (from *radere*, *rasum*, 'to shave, scrape.')

In pharmacy, the division of substances by the rasp or file.

RASOR, *Razor*.

RASORIAN, see *Controstimulus*.

RASO'RISM. The Italian medical doctrine of contro-stimulus, of which Rasori was the founder.

RASORIUM, (*radere, rasum*, 'to scrape,') *Rasporium*.

RASPATO'RIUM, (from *râper*, [old F.] *rasper*, 'to rasp,') *Rasso'rium*, *Rat'ula*, *Scal'prum raso'rium*, *Xys'ter*, *Xyste'rium*, *Xys'trum*, *Rad'ula*, *Runci'na*, *Runci'nula*, *Scal'prum Ras'patory*, (F.) *Rugine*, *Grattoir*. An instrument used for rasping bones, to detach the periosteum, — either in certain surgical operations, or for anatomical purposes.

RASPBERRY, *Rubus idæus* — r. Black, see *Rubus strigosus* — r. Ground, *Hydrastis Canadensis* — r. Wild, *Rubus strigosus*.

RASPING, Eructation.

RASPS, *Rubus idæus*.

RASSETA PEDIS, Tarsus.

RASTETA, *Carpus*, *Malleolus*.

RASU'RA, *Xys'ma*, *Xys'mus*, (F.) *Râpure*, (*radere, rasum*, 'to scrape,') A rasuro, or scratch, or erosion. Also, the rasping or shaving of any substance; as *Rasu'ræ Cor'ni Cer'vi*, *Harts'horn shavings*.

RASURA, Scabs.

RATAFIA, (pron. *Ratafe'a*, and *Rat'afee*,) (F.)

Alcoolé sucré, Saccharo-alcoolé. A liquor, prepared by imparting to ardent spirits the flavor of various kinds of fruits, adding sugar.

RATANHY, *Krameria ratanhia* — r. of the Antilles, *Krameria ixina*.

RATE, Spleen.

RATELEUX, Spleneticus.

RÂTELIER, (from *rateau*, 'a rack,') see *Dentier*.

RA'TIO, *Rationa'litas*, (*reor, ratus*, 'to think,') A reason, a cause, an explanation, a mode or manner; as *Ratio symptom'atum*, an explanation of the symptoms, — *Ratio meden'di*, *Ge'nus curatio'nis*, *Vit'a curan'di*; the reason, mode or manner of cure.

RATIO MEDENDI, see *Ratio* — r. *Symptomatum*, see *Ratio*.

RA'TION, (from *ratio*, 'a proportion,') *Dia'rium*. The daily allowance of necessaries, especially of food, to a soldier or sailor.

Diet rations are founded on principles laid down under *Dietary*; and in all civilized countries they are, in essential respects, the same. The following table is given by Professor Horsford of the rations of the more important armies in active service.

	American.	Aus- trian.	Bava- rian.	British in Cri- mea.	British in India.	French.	Prus- sian.	Russian in Cri- mea.
Flour.....	22 oz. or }		1.65 oz...					
Cornmeal.....	20 oz. or }							
White Bread.....	22 oz. or }		2.30 oz...	24 oz. or	16 oz. or			
Hard Bread.....	16 oz. or }			16 oz.....	16 oz.....	22.83 oz. and }	16 oz. or	
Black Rye Bread.....		32 oz.	17.16 oz.			8.52 oz. }	26.43 oz.	32 oz.....
Fresh Beef.....	20 oz. or.....	4 oz...	3.10 oz...	16 oz. or	16 oz. or	10.78 oz. or.....	8 oz. or.....	16 oz. or
Salt Beef.....	20 oz. or.....			16 oz. or	16 oz. or			16 oz. or
Salt Pork.....	12 oz.....			16 oz.....	16 oz.....	8.47 oz.....	4 oz.....	16 oz.
Potatoes.....	16 oz. 3 times weekly.....						20 oz. or	
Rice.....	1.6 oz.....			2 oz. or.....	4 oz.....	2.11 oz. or.....	2.66 or.....	
Barley.....				2 oz.....	other		4 oz. or.....	
Peas.....					vegeta-		8 oz. or.....	
Beans.....	0.64 gill.....				bles.	2.11 oz. or.....	8 oz. or.....	
Oatmeal.....					16 oz.....		5.3 oz.....	
Desiccated Vegeta- bles.....	or 1 oz. 2 times weekly.....			2 oz.				
Cabbage or Sauer- kraut.....				1 oz.....				3.5 gills.
Coffee, green.....	1.6 oz. or.....			1 oz.....		0.9 oz.		
Coffee, roasted and ground.....	1.28, or.....			0.25 oz...	1.43 or			
Tea.....	2.4 oz.....			0.5 gill...	.43 oz.			
Brandy.....						0.43 gill, or.....	2 gills...	1.5 gill.
Rum.....								
Wine.....						1.75 gill.		
Beer.....								1.1 qt.
Tobacco.....						1/3 oz.		
Sugar.....	2.4 oz.....			2 oz				
Vinegar.....	0.32 gill.....							1.75 gill.
Lime Juice.....				1 oz				quantity unkno'n
Mustard.....								3.86 gr.
Horsradish.....								3.86 gr.
Pepper.....	0.4 oz.....		0.05 oz...	0.31 oz...				0.75 oz.
Salt.....	0.6 oz.....		0.10 oz...	0.62 oz...	1 oz.....	0.9 oz.....	Supply...	
Candles.....	16 oz.) to 100 ra-							
Soap.....	64 oz.) tions.							
Wood.....					48 oz.			

The reduction of the 'marching ration' in bulk and weight, without impairing its nutritive power, has received much attention. Professor Horsford has proposed one consisting of roasted and ground wheat or self-raising flour,—prepared by intimately incorporating bicarbonate of soda and pulverulent acid phosphate of lime with flour, and

forming a dough with water,—and boiled or roasted whole beef, sausaged, dried and compressed, which, he says, is very palatable, imperishable, less expensive, of greatly reduced weight and bulk; requiring, therefore, less transportation, and "will subsist an army in health indefinitely long.

RATIONAL, *Rationalis*, (F.) *Rationnel*. That which is conformable to reason. This epithet is often opposed to *empirical*; as *rational treatment*. The French, also, say *Signes rationnels*, in opposition to *Signes sensibles*. See *Organic*.

RATIONALITAS, Ratio.

RATSBANE, Arsenious acid.

RATTLE, ([G.] *Rattel*), *Rôle*.

RATTLES, THE, see *Rôle*.

RATTLEBUSH, *Sophora tinctoria*.

RATTLESNAKE, *Crotalus horridus*—r. Leaf, *Goodyera pubescens*—r. Plantain, *Goodyera pubescens*—r. Root, *Nabulus albus*, *Polygala senega*, *Trillium latifolium*—r. Weed, *Hieracium venosum*.

RATTLESNAKE'S MASTER, *Agave Virginica*, *Liatris scariosa*, *Nabalus albus*.

RATTLEWEED, *Actæa racemosa*.

RATULA, *Raspatorium*.

RAUCE'DO, gen. *Rauce'dimis*, *Rau'citas*, *Paraphonia rau'ca*, *Asper'itas Arte'riæ as'peræ*, *Vox rau'ca*, *Brau'chus*, *Hoarseness*, (Sc.) *Roup*, (Prov.) *Raup*, (F.) *Enrouement*, *Raucité*, (*raucius*, 'hoarse.') A change in the voice, which loses its smoothness and becomes low and obscure. It is a common symptom in diseases of the larynx and trachea.

RAUCEDO CATARRH'ALIS. Hoarseness from cold.

RAUCITAS, *Raucedo*.

RAUCITÉ, *Raucedo*.

RAUCOUS, *Rau'cus*, (F.) *Rauque*, 'hoarse;' as a raucous voice—*Vox rauca seu raucis'ona*.

RAUQUE, *Raucous*.

RAVE, (from *rapa*), *Brassica rapa*.

RAVERY, *Delirium*.

RAWP, *Raucedo*.

RAY, *Diarrhœa*.

RAYONNÉ, (from [F.] *rayon*, [L.] *radius*, 'a ray,') *Radiated*.

RAYONS SOUS-IRIENS, *Ciliary processes*.

RA'ZOR, *Cul'ter tonsor'ius*, *Machæ'ris*, *Novacula*, *Xy'rism*, *Xy'rus*, (F.) *Rasoïr*, (*radere*, *rasum*, 'to shave.') An instrument for shaving. Used in surgery to remove hairs, where they would interfere with any operation.

RE (L.), in composition, 'back, in return, again.' Before vowels, at times; and before *h* and *d*, it becomes *red*.

REACHING, (from [Sax.] *pæcan*, 'to reach,') *Vomitition*.

REACTION, *Reäc'tio*, (*re*, 'again,' and *agere*, *actum*, 'to act.') An action of resistance to a power applied. A species of vital movement or excitement, tending to prevent or destroy the effects of any noxious power applied to the animal economy, and which, by some, has been attributed to the *vis medicatrix nature*. It is certainly dependent on the organic instinctive actions of the frame. It is the state of activity which succeeds the action on the nervous system of certain morbid influences.

READ, *Rennet*.

READY METHOD, *Marshall Hall's Method*. A mode of restoring respiration in cases of apnoea, proposed by Dr. Marshall Hall. It consists in placing the body in the prone position, making gentle pressure on the back; then removing the pressure, and turning the body on its side and a

little beyond, executing these movements sixteen times in the minute. It is also termed the method of *prone and postural respiration* in apnoea.

REAL'GAR, *Realgal*, *Arlada*, *Arladar*, *Auripigmentum rubrum*, *Arsenicum rubrum factitium*, *Abessi*, *Sandar'acha Græcorum*, *Red Sulphuret of Ar'senic*. A violent poison. See *Poison*.

REAM, *Cream*.

REASON, *Ra'tio*, *Lo'gos*, (F.) *Raison*. The faculty or property of the mind by means of which man perceives the distinction between right and wrong, in physics as well as in morals. *Reason* is a just conclusion resulting from the comparison of two ideas, by virtue of which we form a judgment.

REBOLEA, *Mummy*.

REBONA, *Mummy*. Also, dried faeces, *Ster'cus adustum*.

REBOUTEUR, (*re*, and *bouter*, 'to put,') *Renoueur*.

RECEIV'ER, *Ampul'la*, (F.) *Ballon*, *Recip'ient*, (*recipere*, *receptum*, (*re*, and *capere*, 'to take,') 'to take, to receive,') In pharmacy, a wide-necked globular vessel, the neck of which widens gradually outwards, to receive the tapering beak of the retort, in distillation. Like the retort, the receiver may be *plain* or *tubulated*.

A *quilled receiver* is one that has a tapering tube from its lower surface, which is inserted into a graduated bottle through a cork joint, when it is desirable to know accurately the amount of fluid that has passed over.

RECEPTA, *Prescription*.

RECEPTACULA DURÆ MATRIS, (from *receptare*, *receptatum*, 'to receive,') *Sinuses of the dura mater*.

RECEPTACULUM CHYLI, 'receptacle of the chyle;' *R. Pecquet'i seu Pecquetianum*, *Diversorium seu Stag'num chy'li*, *Sac'culus chy'lifer seu ro'rifer seu lac'teus*, *Al'veus ampullosus seu ampulles'ens*, *Utric'ulus lac'teus*, *Vesic'ula chy'li*, *Ampul'la chy'lifera seu chy'li*, *Cister'na lumb'aris seu Chy'li*, *Chylody's'tis*, *Chylodochi'um*, *Lat'ices lac'tei*, *Lum'bar cis'tern*, *Res'ervoir of Pec'quet*, (F.) *Réservoir du chyle*, *R. de Pecquet*, *Cisterna lombaire*. A considerable dilatation of the thoracic duct, in front of the lumbar region of the vertebral column. It is the commencement of the duct.

RECEPTACULUM COTUX'NI. A triangular space, formed by a separation of the two laminae of the dura mater of the internal ear, near the middle of the posterior surface of the pars petrosa of the temporal bone, where the aquæductus vestibuli terminates. It is always filled with the *Fluid of Cotugno*.

RECEPTACULUM PECQUETI, *R. Chyli*—r. *Sellæ equinae lateribus appositum*, *Cavernous sinus*.

RECEPTA'RII MED'ICI, (from *receptum*.) Physicians, according to *Langius*, who, without any reason, collect from pharmacopœias and chemical works masses of receipts (*recepta*) sufficient to load wagons, ("quibus plaustra onerare possent.")

RECEPTIV'ITY, *Receptiv'itas*, (F.) *Réceptivité*, (*recipere*, *receptum*, 'to receive,') A word used by *Tissot* to express the susceptibility of certain organs to receive morbid impressions.

RECEPTUM, *Prescription*.

RECESSUS, 'a retreat,' *Abscess*, see *Lingula*, *Sinus*—r. *Sulciformis*, see *Fovea vestibuli*.

RECETTE, *Prescription*.

RECHAMELCA, *Rachamelca*.

RECHUTE, (*re*, and *chute*, 'a fall,') see *Relapse*.

RECIDIVA MORBI, (*recidere*, (*re*, and *cadere*), 'to fall back.') Relapse.

RECIDIVE, see Relapse.

REC'IPĒ, (L.) (*re*, and *capere*, 'to take.') A Latin word which signifies 'take,' (F.) *Prenez*, and with which prescriptions are commenced. It is, often, represented by R, R̄, and was formerly by ʒ, the symbol of Jove. See Abbreviation, and R. Substantively, *Recipe* means a prescription or receipt.

RÉCIPIENT, Receiver.

RECIPROCATIÖN, VIBRATIONS OF, see Sound.

RECLINATIO, Reclination—r. Palpebrarum, Ectropion.

RECLINAT'ION, *Reclinat'io*, (*reclinare*, *reclinatum*, (*re*, and *clinare*, 'to bend,') 'to bend back,') A mode of operating for the cataract, which consists in applying the needle in a certain manner to the anterior surface of the cataract and depressing it into the vitreous humour, in such a way that the front surface of the cataract is the upper, and its back surface the lower one.

RECLUSOR PALPEBRARUM, (from *recludere*, *reclusum*, 'to open,') Levator palpebræ superioris.

RECOMPOSITION, Metasynerchris.

RECONVALESCENTIA, (*re*, and *convalescere*, 'to get well,') Convalescence.

RECORDATIO, (from *recordari*, *recordatus*, 'to remember,') Memory.

RECORDATUS, Memory.

RECORPORATIVUS, (*re*, and *corporare*, 'to form into a body,') Metasyneritic.

RECOVERY, (from *recuperare*, (*re*, and *capio*, 'to take,') 'to regain,') Restauratio.

RECREATIO, (*re*, and *creare*, 'to create,') Restauratio.

RECREMENT, *Recremen'tum*, (*re*, and *cernere*, *cretum*, 'to sift,') Of old, dross, excrement. With the moderns, a humour, which, after having been separated from the blood, is again returned to it; such as the saliva, the secretion of serous membranes, &c. Such fluids are said to be *recrementit'ial*, (F.) *Récrementeux*, *Récrementiel*, *Humores inquilini*.

RECREMENTA (pl. of *recrementum*) VESICÆ, Urine.

RÉCREMENTEUX, see Recrement.

RECREMENTITIAL FLUIDS, see Recrement.

RECREMEN'TOËXCREMENTIT'IAL. Applied to secreted fluids, as the saliva, bile, &c., which are in part reabsorbed and in part excreted.

RECREMENTUM, Recrement.

RECRUDESCENTIA, *Palinco'tesis*, *Recrudescence*, (*re*, and *crudus*, 'raw,') An increase of a disease after a temporary remission.

REC'TAL, *Recta'lis*. Appertaining, or relating, to the rectum,—as 'rectal tube.' See Tube, rectal.

RECTIFICA'TION, *Rectifica'tio*, (*rectum*, 'right,' and *facere*, 'to make,') An operation by which certain liquids are purified by distilling them, either singly or after admixture with other substances.

RECTITIS, (*rectum*, and *itis*), properly *architis* or *proctitis*. Inflammation of the rectum. *Proctalgia inflammatoria*.

REC'TO, in composition, relating or belonging to the rectum.

RECTOCELE, (*recto*, and *κηλη*, 'rupture,') Cystocele.

RECTOSTENOSIS, (*recto*, and *στενος*, 'contracted,') Stricture of the rectum.

RECTOÛTERINE, *Rectoûteri'nus*, (*recto*, and

uterus.) Appertaining, or relating, to the rectum and uterus.

RECTOÛTERINE CUL-DE-SAC, see Uterus—1. Follics. see Uterus—r. Pouch, see Uterus.

RECTOVAG'INAL, (*recto*, and *vagina*.) *Rectovagina'lis*. That which relates or belongs to the rectum and vagina.

RECTOVAGINAL CUL-DE-SAC or POUCH is formed by the reflection of the peritoneum from the upper third of the vagina to the rectum.

RECTOVAGINAL FIS'TULA is owing to a preternatural communication between the rectum and vagina through the above septum.

RECTOVAGINAL SEP'TUM, (F.) *Cloison rectovaginale*, is formed by the junction—back to back—(or by what the French term *adossement*) of the rectum and vagina. It separates these two canals from each other.

RECTOVES'ICAL, (*recto*, and *vesica*, 'the bladder,') *Rectovesica'lis*. Having relation to the rectum and bladder; as the '*rectovesical fold* or *pouch* or *cul-de-sac*' of the peritoneum, between the rectum and bladder. See Urinary Bladder.

RECTUM, from *rectus*, 'straight,' so called from a notion that it is straight, which it is not. *Intestinum Rectum*, *Apophys'menos*, *Enthys'ieron*, *Lon'gan*, *Lon'gas*, *Lon'ganon*, *Ar'chos*, *Cyl'saros*, *Prin'ceps*, *Monoco'lon* [?], *Bumgut*, (Prov.) *Jotgut*. The third and last portion of the great intestine. It forms the continuation of the sigmoid flexure of the colon, occupies the posterior part of the pelvis, and extends from the left side of the sacro-vertebral articulation to the coccyx, before which it opens outwards by the orifice called the *anus*. It is cylindrical, but *sacculated*, as it were, at its inferior portion, where it is consequently larger, and is attached to the sacrum by the mesorectum. Like the other portions of the canal it is formed of three membranes. Towards the lower extremity, some parallel and longitudinal wrinkles are observable on its inner surface: these are the *Columnæ*, *Columnæ car'næ of the Rectum* or of *Morgagni*, (F.) *Colonnes du rectum* or *de Morgagni*. They are merely folds of the mucous membrane, between which is a number of *Lacunæ*. The arteries of the rectum are numerous. They proceed from the inferior mesenteric, hypogastric, and internal pudic, and are called *hemorrhoidal*. The veins open, partly into the hypogastric vein, partly into the inferior mesenteric. The nerves proceed from the sciatic and hypogastric plexuses. The rectum receives the fecal matters, as they pass from the colon, which serves as a reservoir for them.

RECTUS, 'straight.' (F.) *Droit*. A term, used in anatomy for parts that pursue a straight course:—as

RECTUS ABDOM'INIS, (F.) *Pubio-sternal*, *Sterno-pubien* (Ch.), *Droit de l'abdomen*. A long flat muscle, situate at the anterior part of the abdomen, and separated from that of the opposite side by the *linea alba*. It is attached, above, to the cartilages of the last three true ribs; and, below, to the pubis by a very strong tendon, the outer edge of which is continuous with the fascia transversalis. This muscle is enclosed in an aponeurotic sheath; and, in its course, has three or four transverse aponeurotic intersections. It bends the chest on the pelvis, and conversely.

RECTUS ANTERIOR BREVIS, *Rectus capitis internus minor*—r. *Anterior longus*, R. *capitis internus major*—r. *Capitis anticus longus*, R. c. *internus major*.

RECTUS CAPITIS INTERNUS MA'JOR, R. *internus major* seu *anterior longus* seu *capitis anticus longus*, (F.) *Trachélo-basilaire*, *Grand droit antérieur de la tête*. This muscle is long in proportion to its breadth, flat, and broader above than below. It is situate at the anterior and lateral

part of the neck, and arises from the transverse processes of the third, fourth, fifth, and sixth cervical vertebrae. It is inserted into the basilar process of the occipital bone.

RECTUS CAP'ITIS INTER'NUS MI'NOR, *R. inter'nus mi'nor* seu *anterior brevis*, *Ren'ens*, *An'mens*, (F.) *Petit trachélo-basilaire*, *Petit droit antérieur de la tête*. This is situate deeper than the last. It is thin, flattened, irregularly quadrilateral, and passes from the anterior surface of the body of the first vertebra, near the origin of the transverse process, to be inserted into the basilar process.

RECTUS CAP'ITIS LATERA' LIS, *R. latera'lis* seu *lateralis Fallo'pii*, *Transversa'lis anti'cus pri'mus*, (F.) *Trachélo-atloïdo-basilaire*, *Atloïdo-sous-occipital* (Ch.), *Droit latéral de la tête*. There is a muscle of this name on each side of the head. It is flat, quadrilateral, and passes from the transverse process of the atlas to the occiput, behind the jugular fossa. It inclines the head to one side.

RECTUS CAP'ITIS POSTI'CUS MA'JOR, *R. Ma'jor* *Cap'itis posti'cus mi'nor* (Albinus), *Spini'axoïdo-occipita'lis*, (F.) *Axoïdo-occipitale* (Ch.), *Grand droit postérieur de la tête*. A flat, triangular muscle, situate at the posterior and superior part of the neck. It extends from the spinous process of the second cervical vertebra to the external surface of the lower semicircular ridge of the occipital bone. This muscle extends the head, inclines it to one side, and causes it to experience a rotary motion, which directs the face to its side.

RECTUS CAP'ITIS POSTI'CUS MI'NOR, *R. mi'nor*, (F.) *Tuber-atloïdo-occipital*, *Atloïdo-occipital* (Ch.), *Petit droit postérieur de la tête*. A flat, triangular muscle, situate beneath the last, which passes from the posterior arch of the atlas to the occipital bone. It extends the head.

RECTUS CAPITIS POSTICUS MINOR, *R. c. p. major* — *r. Cruris*, *R. femoris*.

RECTUS EXTER'NUS OC'ULI, *Abduc'tor oc'uli*, *Indignabundus*, *Indignatus*, *Inacundus*, *Oc'ulum mo'vens secundus*, (F.) *Droit externe*, *Inscissile* ou *Abduc'teur*, *Orbito-externo-scléroticien*. All the recti muscles of the eye, four in number, arise from the bottom of the orbit, and are inserted into the sclerotic coat of the eye. The *Rectus externus* is divided, at its posterior extremity, into two portions, one of which is attached to the outer part of the foramen opticum, and the other to the sphenoid bone.

RECTUS FEM'ORIS, *Rec'tus* seu *Grac'ilis anterior* seu *cru'ris*, (F.) *Ilio-rotulien*, *Droit antérieur de la cuisse*. A straight muscle, seated at the anterior part of the thigh. It extends from the anterior and inferior spine of the ilium and the cotyloid cavity of the patella. This muscle extends the leg upon the thigh, and conversely. It can, also, bend the thigh upon the pelvis or carry the pelvis forward.

RECTUS INFERIOR FEMORIS, *Gracilis*.

RECTUS INFE'RIOR OC'ULI, *Depres'sor oculi*, *Tim'idus*, *Dep'rimens*, *Hu'milis*, *Oc'ulum mo'vens quartus*, (F.) *Sous-optico-sphéno-scléroticien*, *Droit inférieur* ou *abaisseur de l'œil*, *Muscle humble*, arises below the optic foramen from the sphenoid bone. It depresses the eye.

RECTUS INTERNUS MAJOR, *R. capitis internus major* — *r. Internus minor*, *R. capitis internus minor*.

RECTUS INTERNUS OC'ULI, *Adduc'tus* seu *Adduc'tor oc'uli*, *Bibito'rius*, *Oc'uli ad'ductor*, *Oc'ulum mo'vens pri'mus*, (F.) *Orbito-intus-scléroticien*, *Droit interne de l'œil* ou *adducteur* ou *boueur*. It arises from the sphenoid at the inferior part of the foramen opticum. It turns the eye towards the nose.

RECTUS LATERALIS, *R. capitis lateralis* — *r. Lateralis Fallopii*, *R. capitis lateralis* — *r. Major*, *R. capitis posticus major* — *r. Minor*, *R. capitis posticus minor*.

RECTUS SUPERIOR OC'ULI, *Atto'lens oc'uli* seu *oc'ulum*, *Leva'tor* seu *Eleva'tor oc'uli*, *Super'bus*, *Oc'ulum mo'vens tertius*, (F.) *Superbe*, *Sus-optico-sphéno-scléroticien*, *Droit supérieur* ou *Élévateur de l'œil*, *Dédaigneux*, is attached to the circumference of the foramen opticum.

All the recti muscles are concerned in the voluntary motions of the eye. The oblique muscles have been supposed by Sir C. Bell to execute the involuntary.

RECUBITORIUM, (*recumbere*, *recubitum*, 'to recline,') *Anaclinterium*.

RECUPERATIO, *Restauratio*.

RECUPERATION, (from *recuperare*, *recuperatum*, 'to recover,') *Restauratio*.

RECUPERATIVE, *Recup'eratory*. *Restorative*. Recovering. Belonging to recovery.

RECUR'RENT, *Recur'rens*, (from *recur'rere*, (*re*, and *curre're*), 'to run back,') A name given to several arterial and nervous branches, which seem to re-ascend towards the origin of the trunk whence they emanate; also, to a disease that relapses.

RECURRENT ARTERIES. This name has been given to several arteries of the forearm, and to one of the leg. In the forearm, there are, 1. The *Radial recur'rent*, *Artère récurrente de l'épicondyle* (Ch.), which arises from the superior part of the radial, and ascends between the supinator longus, s. brevis, and the brachialis internus. 2. The *anterior cubital recur'rent*, *A. récurrente de l'épitrachlée* (Ch.), which arises from the upper and inner part of the cubital, and ascends between the pronator teres and brachialis internus. 3. The *posterior cubital recur'rent*, which arises from the preceding artery, or from the cubital itself, and ascends between the olecranon and internal tuberosity of the humerus. 4. The *posterior radial recur'rent*, *A. récurrente olécraniennne* (Ch.). It arises from the posterior interosseous, and ascends between the posterior cubital and anconeus muscles. In the leg, the *Arteria recur'rens tibialis*, *A. récurrente au genou* (Ch.), is furnished by the anterior tibial, and ascends to the tibialis anticus muscle.

RECURRENT BANDAGES are formed by convolutions of bandages, which make a kind of cap for the part—as the head—to which they are applied.

RECURRENT FEVER, see *Relapse*.

RECURVATIO, (*re*, and *curvus*, 'crooked,') see *Hump*.

RECUTITUS, (*re*, 'back,' and *cutis*, 'skin,') *Apella*.

REDBEAN, *Abrus precatorius*.

REDBERRY, *Arbutus uva ursi*, *Gaultheria*, *Panax quinquefolium* — *r. Bitter*, *Cornus Florida* — *r. Swamp*, *Vaccinium oxycoccos*.

REDGOWN, see *Strophulus*.

REDGUM, see *Strophulus*.

RED HEAD, *Asclepias Curassavica*.

REDMAILKES, *Papaver rhœas*.

REDROD, *Cornus sericea*.

RED ROOT, *Celastrus*, *Lachnanthes tinctoria*, *Sanguinaria Canadensis*.

REDSHANKS, *Polygonum persicaria*.

RED WATER TREE, see *Sassy Bark*.

RED WEED, *Phytolacca decandra*.

REDIBIA, *Paronychia*.

REDIMICULUM, ('a chaplet,') *Diadema*.

REDINGOTES ANGLAISES, 'English riding coats, frock coats,' see *Condom*.

REDINTEGRATIO, (*re*, and *integer*, 'entire,') *Regeneration*, *Restauratio*.

RÉDONDANCE, (*re*, and *unda*, 'a wave,') *Plethora*.

REDOUBLEMENT, (*re*, and *doubler*, 'to double,') *Exacerbation*, *Paroxysm*.

REDRESSEUR UTÉRIN, (*re*, and *dreuer*, 'to make straight,') *Pessary*, *intrauterine*.

REDS, (from the colour,) *Menstrues*.

REDUCTIO, (*reducere, reductum, (re, and ducere, ductum, 'to lead,') 'to bring back,'*) Reduction. Taxis—*r. Præputii impedita, Phimosis.*

REDUCTIO, (*Reductio, Repositio, Restitutio, Diapylaxis, Diaplasimus, Syntaxis.*) A surgical operation, the object of which is to restore displaced parts to their original situation. Thus, we say the reduction or *setting* of a fracture, when the displaced parts are again brought to their proper relative situation. The reduction of a hernia is the restoration of the protruded parts to the cavity of the abdomen. This is, also, called *Taxis*.

In pharmacy, the process of restoring oxides, chlorides, sulphurets, &c., to the metallic state.

REDUNDANTIA, (*re, and unda, 'a wave,'*) Plethora.

REDUVIA, Paronychia.

REED MACE, Typha latifolia.

REFECTIO, (*reficere, refectum, 'to restore,'* from *re, and facere, 'to make,'*) Restauratio.

REFECTIVA, Analeptica.

REFECTUS, Restauratio.

REFICIENTIA, Analeptica.

RÉFLECTIF, Reflex.

REFLECTION, (*re, and flexere, flexum, 'to bend,'*) *Reflexio*, (F.) *Reflexion*. A bending or turning back. Duplicateur. The act of turning the mind back on its actions; so as to constitute contemplation, deliberation.

REFLECTOR EPIGLOTTIDIS. A muscle, which arises from the arytenoid cartilage and inner part of the thyroid, and is inserted into the lateral edges of the epiglottis.

REFLEX, *Reflexus*, (F.) *Réflexe, Réflexif*, (same etymon as *Reflection*.) A term applied to an action, which consists in the reflection by an efferent nerve of an impression conveyed to a nervous centre by an afferent nerve; hence reflex movement, reflex paralysis, reflex epilepsy, &c.

A reflex action is generally regarded to be one executed without consciousness; like the 'reflex or excito-motory movements' of Dr. Marshall Hall.

REFLEX MOVEMENTS, see Motion—*r. System* of nerves, see Excitomotory, and Nerves.

REFLEXIO, Reflection—*r. Palpebrarum, Ectropion*.

RÉFLEXION, Reflection.

RÉFORME, Invaliding.

REFRACTION, *Refractio, Photocampsis, Diacelasis, (re, and frangere, fractum, 'to break,')* The change of direction which rays of light experience when they pass obliquely from one medium into another of different density. If the light passes from a rarer to a denser medium, it approaches the perpendicular raised at the point of immersion. The contrary occurs, when it goes from a denser to a rarer medium. The theory of lenses, telescopes, and of the eye, rests entirely on this property of light.

REFRIGÉRANT, see Alembic.

REFRIGÉRANTS, *Refrigerantia, Psyc'tica, Antither'ma, Frigefacientia*, (F.) *Rafrâchissants, Réfrigératifs, Desaltérants, (re, and frigere, 'to cool,')* Medicines which depress the morbid temperature of the body. The chief reputed refrigerants are the Potassæ Nitræs [?], the Sodæ Boras [?], but chiefly the abstraction of caloric by ice and beedinks, cold water, cold lemonades, sodawater, and effervescing draughts internally; and externally, cool air, cold water, evaporating lotions, and frigorific mixtures.

RÉFRIGÉRATIFS, Refrigerants.

REFRIGERATOR, see Alembic.

REGENERATIO, (*re, and generare, generatum, 'to beget,'*) Regeneration, Reproduction.

REGENERATION, *Regeneratio, Palingen'ia, Palingene'sia, Redintegratio*. Reproduc-

tion of a part of the body, which has been destroyed.

REG'IMEN, gen. *Regim'inis*, (F.) *Régime, (regere, 'to govern,')* The rational and methodical use of food, and of everything essential to life; both in a state of health and disease. It is often restricted, in its meaning, to *Diet*. It is sometimes used synonymously with hygiene.

Régime Sanitaire and *Système Sanitaire*, mean the measures and regulations adopted to prevent the development and propagation of diseases reputed to be pestilential,—especially plague, yellow fever, and cholera.

REGIM'INAL, *Regim'inalis*, (from *regimen*, gen. *regim'inis*.) Relating or appertaining to regimen,—as '*regiminal treatment*,' or treatment by regimen.

REGINA PRATI, 'queen of the meadow,' *Spiræa ulmaria*.

REGIO, Region.

REGIO AURICULA'RIS. The region of the ear.

REGIO BUCCA'LIS. The region of the cheek.

REGIO CARDI'ACA, Epigastric region.

REGIO CERVICA'LIS. The region of the neck. The cervical region.

REGIO EPIGASTRICA, Epigastrium.

REGIO FACIA'LIS. The facial region.

REGIO FRON'TALIS. The frontal region.

REGIO GASTRICA, Umbilical region—*r. Hypochondriaca, Hypochondre*—*r. Hypogastrica*, see Hypogastrium—*r. Iliaca, Iliac region*—*r. Inguinalis, Inguinal region*—*r. Ischiadica, Ischiadic region*—*r. Jugalis, Jugal region*—*r. Lumbalis, Lumbi*—*r. Mentalis, Mental region*—*r. Mesogastrica, Umbilical region*—*r. Nasalis, Nasal region*—*r. Nuchæ, Nuchal region*—*r. Nuchalis, Nuchal region*—*r. Occipitalis, Occiput*—*r. Occipitalis inferior, Nuchal region*—*r. Perinaei, Perinaeum*—*r. Præcordiaca, Præcordial region*—*r. Pubis, Pubic region*—*r. Stomachalis, Epigastric region*—*r. Stomachica, Epigastric region*—*r. Temporalis, Temple*—*r. Umbilicalis, Umbilical region*.

REG'ION, *Reg'io, (regere, 'to direct,')* Anatomists have called *regions* determinate spaces on the surface of the body or of different organs. Thus, the abdomen has been divided into several; to which different names have been given. See Abdomen, Epigastrium, Hypochondrium, Hypogastrium.

RÉGION OMBILICALE, Umbilical region.

REGIONAL ANATOMY, see Anatomy.

RÈGLES, (from *regula, 'a rule,'*) Menses—*r. Dèrives, Menstruation, vicarious*—*r. Difficiles, Dysmenorrhœa*.

RÉGLISSE, Glycyrrhiza.

RÈGNE ANIMAL, (from *regnum, 'a kingdom,'*) see Animal.

REG'ULAR, *Regula'ris*, (F.) *Régulier, (from regula, (regere, 'to govern,')* 'a rule,') According to rule. Applied to the types of a disease, and to the pulse, when according to rule.

REGULAR PRACIT'IONER. One who practises his profession according to the rules established by law or professional custom.

REG'ULUS, (dim. of *rex*, gen. *regis, 'a king,'*) A pure metal. One reduced from its ore.

REGULUS ANTIMONI, Antimonium.

REGURGITATION, *Regurgitatio, (re, and gurgere, gen. gur'gitis, 'a gulf,')* The act by which a canal or reservoir throws back substances accumulated in it. *Mitral regurgitation*, for example, means the reflux of blood through the left auriculo-ventricular opening, during the contraction of the left ventricle. Regurgitation is also applied to the *pu'king* or *pos'setting* of infants; and to the rising of solids or fluids into the mouth in the adult,—from *re, and gorge, (gurgere, 'the throat,')* (F.) *Rapport*. What is called "vomiting at pleasure" is regurgitation.

The substances brought into the mouth by regurgitation, the French call *Renvois*.

REIF, (from Sax. *hreoF*), Psora.

REIN, (from [L.] *ren*, gen. *renis*), Kidney.

REINE DES PRÉS, 'queen of the meadows,' *Spirea ulmaria*.

REINERZ, MINERAL WATERS OF. Rein-
erz is a small town in Silesia, three German miles from Glatz. The waters are acidulous chalybeates. The principal springs are the tepid drinking spring, the cold spring, and three bathing springs.

REINFORCEMENT, FASCICULUS OF, *F. Cuneatus*. A band of fibres from the *corpus innominatum* or lateral part of the anterolateral column of the medulla, which is continued up behind and around the corpus olivare, is lined on its inner or central surface with cineritious substance, enlarges as it ascends, passes over the upper surface of the pons Varolii, and is expanded into the optic thalamus. It thus forms, in its course, the anterior paries of the fourth ventricle, and is brought into view by brushing away the *tuberculum cinereum* or gray matter of the ventricle. This band has received the name owing to its constantly increasing as it ascends, and being mixed up with gray matter on its inner face, from which it seems to derive new fibres.

REINS, (from [L.] *ren*, pl. *renes*), Kidneys, Lumbi.

REINS SUCCENTURIÉS ou SUCCENTURIAUX, Capsules, renal.

REJECTIO, *Anagoge*, (*rejicere*, *rejectum*, (*re*, and *jacere*, 'to cast,') 'to cast back.')

Every excretion by the mouth,—spitting, expectoration, regurgitation, or vomiting.

REJECTIO SANGUINIS E PULMONIBUS, Hæmoptysis.

REJUVENESCENCE, *Rejuvenescencia*, (*re*, and *juvenescens*, 'growing young:' [*juvenis*, 'a youth.']) A renewal of youth. The state of being young again. The assumption of the characteristics of youth by the aged.

RELACHANT, Relaxant.

RELACHEMENT, Relaxation.

RELAPSE, *Hypotrophé*, *Epistrophé*, *Hypostrophé*, *Morb'bi Recidiva*, *Hypotropismus*, (*re*, and *labor*, *lapsus*, 'to fall down.')

The return of a disease during, or shortly after, convalescence. The French use *Rechute* in this sense; and *Récidive* (*Morb'bi recidivus*) for the return of a complaint, at a greater or less period after the complete recovery of health.

The terms *Relapsing* or *Recurrent fever*, *Short fever*, *Five-day fever*, *Seven-day fever*, *Bilious*

Remittent fever, *Remitting Icteric fever*, and *Mild Yellow fever*, *Typhus seu Febris recurrens*, (F.) *Fièvre à rechute*, have been given to a form of continued fever, which has occasionally appeared epidemically, as in Edinburgh, in 1843 and 1847, and was generally attended with icteric symptoms. The main phenomena disappeared at about the end of a week; but in many cases a relapse occurred about the end of a fortnight. Few died of the affection.

RELAPSING FEVER, see Relapse.

RELATIO, (*re*, and *ferre*, *latum*, 'to carry,')

Rapport.
RELATION, FUNCTIONS OF, see Function.

RELAXANT, *Laxans*, *Molacicum*, *Chalasticum*, *Pareticum*, (F.) *Relâchant*, (*re*, and *lazare*, 'to loosen.')

A medicine which diminishes the tension of organs. Mucilaginous and fatty substances are relaxants.

RELAXANTIA, Emollients.

RELAXATIO, Relaxation—r. *Cordis et arteriarum*, Diastole.

RELAXATION, *Relaxatio*, *Chalastis*, *Chalastimus*, (F.) *Relâchement*. In *Physiology*, relax-

ation is opposed to contraction. In *Pathology*, it means great looseness of a part.

RELÈVE PAUPIÈRE, ([L.] *re*, and *levare*, 'to raise.')

An instrument for raising the eyelid.

RELEVEUR DE L'ANUS, Levator ani—r. *de la Lèvre inférieure*, Levator labii inferioris—r. *du Menton*, Levator labii inferioris—r. *de l'Omoplate*, Levator scapulae—r. *de la Prostate*, Levator prostatae—r. *de l'Urètre*, Levator urethrae.

RELIGIOUS DISEASES, (F.) *Maladies religieuses*. Diseases affecting the nervous system, and arising from the excitement of religious emotions. See *Convulsivum*.

RELIQUAT (F.), (from [L.] *relinquere*, *relictum*, 'to leave.')

The remnant or remains of a disease.

REMAK FIBRES, (after Remak, the German physiologist), see *Fibres*, Remak.

REMANSIO MENSIUM, (*re*, and *maneo*, *mansum*, 'to stay,')

Amenorrhœa.

REMEDE, ([L.] *remedium*), Clyster, Medicament—r. *du Capuchin*, Liquor hydrargyri nitrici—r. *du Duc d'Antin*, Liquor hydrargyri nitrici—r. *de Durande*, Liquor anodynus terebinthinatus.

REMEDIAL, Medicinal.

REMEDiate, Medicinal.

REMEDILESS, Incurable.

REMEDium, (*re*, and *mederi*, 'to heal,')

Juans, Medicament—r. *Dentifricium*, Dentifrice.

REMEDY, Medicament—r. *English*, *Cinchona*.

REMETIC, Medicament.

REMISSIO, (*re*, and *mittere*, *missum*, 'to send,')

Decline, Remission—r. *Cordis et Arteriarum*, Diastole—r. *Morb'bi*, Remission—r. *Virium*, Atony.

REMISsion, *Remissio*, *Epanoësis*, *Anoësis*, *Submissio*, *Aphoësis*, *Eudosis*, *Meioësis*. In a more extensive signification, a temporary diminution of the symptoms of a disease, either acute or chronic, *Remissio morbi*. Diminution in the febrile symptoms, such as occurs in a remittent fever between the exacerbations.

REMITTENS, Remittent—r. *Icterodes*, Fever, yellow.

REMITTENT, *Remittens*, *Remitting*. Any disease which presents remissions.

REMITTENT OF CHILDREN, Fever, infantile, remittent.

REMITTENT FEVER, *Febris remittens*, *Epanoësis*, *Erae'ebating*, *paroxysmal*, *subcontinua*, *endem'ic*, *endem'ial* or *endem'ical* fever, (F.) *Fièvre rémittente*, is one which strikingly exacerbates and remits, but without intermission. The ordinary bilious fever of the United States is a simple remittent—*Epanoësis mitis*, (F.) *Fièvre rémittente simple*.

REMITTENT FEVER, BILIOUS, Remittent fever, see Relapse.

REMITTING ICTERIC FEVER, see Relapse.

REMORA, (*re*, and *mora*, 'delay.')

Stoppage, stagnation,—as of blood. Also, the name of two surgical instruments, intended to retain parts *in situ*. The one was used, formerly, in castration, to prevent the intestines from protruding at the inguinal ring: the other, called *Remora Hildani*, (F.) *Arrêt d'Hildan*, *Echeneis*, was employed to maintain fractures and luxations reduced. It is now not used.

REMORA ARATRI, *Ononis spinosa*—r. *Hildani*, see *Remora*.

REN, Kidney.

RE'NAL, *Renalis*, (from *ren*.) That which relates to the kidney.

RENAL ALTERNATIVES, see *Alternatives*.

RENAL ARTERIES, *Emulgent arteries*, are very large and short, and commonly two in number—one to each kidney. They arise from the sides of the abdominal aorta, forming with it a right angle. The left is commonly a little more anterior and elevated than the right. After a short

course, they arrive at the fissure of the kidney, where they divide, before entering it, into three or four very considerable branches.

RENAL CAPSULE, Capsule, renal—*r.* Nerve, see Splanchnic nerves.

RENAL PLEX'USES, *Emulj'ent Plex'uses*. These are two;—one on each side. They proceed from fasciculi of the solar and coeliac plexuses; from the outer side of the semilunar ganglion and the final expansion of the small splanchnic nerves. These plexuses begin by three or four ganglions, situate at the commencement of the renal artery; and they give off from their circumference many fine, straight filaments, which do not anastomose, and which enter the kidney with the divisions of the renal artery. These divisions they accompany.

RENAL VEINS, *Emulj'ent Veins*. These are very large. Their roots follow the same course as the arteries in the substance of the kidney. In the fissure or pelvic portion of the kidney, they unite into a single trunk, which passes transversely inwards, and opens at the side of the vena cava descendens.

RENCHUS, (*per'nos*.) Snoring. Stertor.

REN'culus, *Renic'ulus*, *Ren'ulus*, (dim. of *ren*.) A small kidney. A lobe of the kidney.

RENDERING, (as lard,) see Adeps preparatus.

RENDLES, Rennet.

RENEAL'MIA EXALTA'TA, (after P. and M. L. Rénéaulme, botanical writers of France,) *Alpin'ia exalta'ta* seu *tubula'ta*, (F.) *Rénéaulme*. A plant of British Guiana, which is possessed of diaphoretic and diuretic properties; and, in large doses, is emetic. It is much used there in dropsy, rheumatism, dysentery, &c. The bruised rhizoma is the part employed.

RENES SUCCENTURIATI, (*sucenturio*, 'a subcenturion,') Capsules, renal.

RENICULUS, (dim. of *ren*.) Reneulus.

RENIFORM, (*ren*, and *forma*, 'shape,') Nephroid.

RENNES. MIN'ERAL WA'TERS OF. Thermal salines in the department of Aube, France, containing carbonic acid and chloride of magnesium. Temperature, 103° to 121° of Fah. Season, from the 1st of May to October.

REN'NET, *Run'net*, *Cuig'ulum*, *Py'tia*, *Pyet'ia*, (from [G.] *gerinnen*, 'to coagulate' [?],) (F.) *Préure*, *Caille-lait*. When the fourth stomach or *Abomasus* of the calf is salted and dried, it possesses the property of coagulating milk, when a portion of it is soaked in water, and the infusion—*Ren'net*, *Ren'dles*, *Run'ning*, (Prov.) *Heslop*, *Cheeselope*, *Yernin*, *Earn'ing*, *Preaser*, *Read*, *Run'ning*, *Steeep*—is added to milk. Liquid rennet or *Rennet wine* may be made, according to Mr. Wm. Procter, Jr., of fresh *rennets*, (about three,) $\frac{3}{4}$ xxiv; chloride of sodium, three trouncings; alcohol, $\frac{1}{2}$ xvj; white wine, $\frac{1}{2}$ xvj; water, q. s.) A tablespoonful of this solution coagulates a quart of milk to a firm curd.

RENNET WHEY, see Serum lactis.

RENOCULE, *Ranunculus*—*r.* *Bulbouse*, *Ranunculus bulbosus*.

RENOUÉE, (*re*, and *nouer*, 'to knot,') Polygonum aviculare—*r.* *Acre*, Polygonum hydropiper.

RENOUEUR, (F.) *Rebouteur*, *Rhabilleur*, (*re*, and *nouer*, (*nodus*, 'a knot,') 'to tie or knot,') A bone-doctor:—a bonesetter. One who makes a trade of reducing fractures and dislocations.

RENOVATIO, (*re*, and *novare*, *novatum*, 'to make new,') Ananosis.

RENSELAERIA VIRGINICA, (after General Van Rensselaer, of Albany,) *Peltandra Virginica*.

RENUENS, (*renuo*, 'to deny by a motion of the head,') Rectus capitis internus minor.

RENULUS, (dim. of *ren*.) Reneulus.

RENUM ATONIA, 'atony of the kidneys,' Nephrotonia—*r.* Paralysis, Nephrotonia.

RENUNCIATIO, (*re*, and *nunciare*, *nunciatum*, 'to tell,') *Rapport*.

RENVERSEMENT DE LA LANGUE, (*re*, and *vertere*, *versum*, 'to turn,') Paraglossa—*r.* *de la Matrice*, Uterus, inversion of the—*r.* *des Pau-pières*, Ectropion—*r.* *du Rectum*, Proctocoele—*r.* *de la Vessie*, Exocyste, Extrophia of the bladder.

RENVOLIS, (*re*, and *envoyer*, 'to send,') see Regurgitation.

RÉOPHORE, (*jon*, 'a current,' and *porter*, 'to carry,') Electrode.

REPANDATIO, (*repandus*, (*re*, and *pandere*, 'to spread,') 'spread backwards,') Lordosis.

REPARATION, (*re*, and *parare*, *paratum*, 'to set in order,') Restauratio.

REPARATIVE, *Repar'atory*. Having the power of restoring or repairing.

REPELLENTIA, Repellents.

REPELLENTS, *Repellen'tia*, *Repercution'tia*, *Apoerust'ica*, *Stal'tica*, *Repercuss'ives*, (F.) *Répercussifs*, (*re*, and *pellere*, 'to drive,') Medicines which, when applied to a tumefied part, cause the fluids, that render it tumid, to recede, as it were, from it. Astringents, cold water, ice, &c., are repellents.

RÉPERCUSSIFS, (*re*, and *percutio*, *percussum*, (*per*, and *quater*, 'to shake or batter,') 'I strike back,') Repellents.

REPERCUSSIO, Contra-fissura, Repercussion.

REPERCUSSION, *Repercuss'io*, *Anac'lasis*. Disappearance of a tumour or cutaneous eruption, in consequence of the application of a repellent. The action of a repellent remedy. The French writers use the term *Délitescence*, *Délitescen'tia*, for the sudden disappearance of a tumour, inflammation, eruptive disease, or purulent collection, without the disease occurring in any other part of the body: in this respect *délitescence* differs from metastasis. *Délitescence* of an inflammation or 'sudden disappearance' is distinguished by certain French pathologists from *Resolution*. Repercussion also means *Ballottement*.

REPERCUSSIVES, Repellents.

REPERCUTIENTIA, Repellents.

REPLETIO, (*re*, 'again,' and *pleo*, 'to fill,') Fulness, Plethora.

REPLI LONGITUDINAL DE LA MÉNINGE, (*repli*, 'a fold,' from [L.] *re*, and *plicare*, 'to fold,') Falx cerebri.

REPOSITIO, (*re*, and *ponere*, *positum*, 'to place,') Anaplasia, Reduction, Taxis.

REPOUSSOIR, (*re*, and *pellere*, *pulsum*, 'to drive,') Punch—*r.* *d'Arêtes*, Probang.

REPRIMENS, (*re*, and *premere*, 'to press,') Epischeticus.

REPRODUCTION, *Reproduc'tio*, *Regenera'tio*, (*re*, and *producere*, *productum*, (*pro*, and *ducere*, 'to lead,') 'to bring forth,') The function by which living bodies produce bodies similar to themselves. See Generation. As a general remark, it will be found true, that the larger animals are parous; and the smaller, which are more exposed to destruction, multiparous.

REPRODUCTION MÉRISMATIQUE, see Generation.

REPRODUCTIVITAS, (F.) *Reproductivité*, Capability of reproducing.

REPULSORIUM, (*re*, and *pellere*, *pulsum*, 'to drive,') Punch.

REPURGATIO, *Anacathar'sis*, (*re*, and *purgare*, *purgatum*, 'to cleanse,') According to Castelli, purgation by transpiration or by expectoration. Medicines producing this effect are called *Repurgan'tia*.

REQUIES, gen. *Requie'tis*, (*re*, and *quies*, 'rest,') *Acinesia*.

REQUIETIO, (*re*, and *quiescere*, *quietum*, 'to become at rest,') *Acinesia*.

RES CON'TRA NATU'RAM, ('Things against nature,') (*F.*) *Chosæ contre-nature*, are those things which tend to destroy man;—as disease, and every thing connected with it.

RES CULINARIA, Culinary art.—*r.* Medica, Medicina.

RES NATURA'LES, 'natural things,') (*F.*) *Chosæ naturelles*. Those things, which, by their union, were conceived to constitute the nature of man; such as the elements, temperaments, humours, spirits, &c.

RES NON NATURA'LES, 'things not natural,') (?) (*F.*) *Chosæ non-naturelles*. Those things which, anciently, formed the matter of hygiene. See *Nonnaturals*.

RES VENEREA, Coition.—*r.* Vestitaria, Vestitus.

RESCISION, (*re*, and *scindere*, *scissum*, 'to cut asunder,') see *Amputation*.

RÉSEAU, (from *reticellum*, dim. of *rete*, 'a net,') *Rete*, *Reticulum*.—*r.* Admirable, *Rete mirabile*.—*r.* Intermédiaire, see *Corpus spongiosum urethrae*.—*r.* Merveilleux, *Rete mirabile*.

RESECTIO, Resection.—*r.* Articulorum, see *Resection*.

RESECT'ION, *Resectio*, (*resecare*, *resectum*, (*re*, and *secare*,) 'to cut off,') Act of cutting or paring off. A name, given, especially by the French surgeons, to operations in which the carious extremities of long bones, or the unconsolidated extremities of fractured bones forming irregular joints, are removed with the saw.—*Resectio articulo'rum*, *Decapitatio articulo'rum*.

RESEDA CRISPATA, (from *resedo*, (*re*, and *sedare*, 'to settle,') 'to calm or assuage,') *R. Luteola*—*r.* Lanceolata, *R. Luteola*.

RESE'DA LUTE'OLA, *R. lanceolata* seu *crispata*, *Lute'ola*, *Weld*, *Dyer's Weed*. A European plant naturalized in the United States, which was at one time used as a diaphoretic and diuretic.

RESERANS, (from *resero*, 'to open,') *Aperient*.

RESERVE AIR, see *Respiration*.

RÉSERVOIR DE LA BILE, ([*L.*] *re*, and *servare*, 'to keep,') Gallbladder.—*r.* du Chyle, *Receptaculum chyli*.—*r.* des Larmes, *Lachrymal sac*.—*r.* of Pecquet, *Receptaculum chyli*.—*r.* de Pecquet, *Receptaculum chyli*.—*r.* of the Thymus, see *Thymus*.

RÉSIDENTIA, (*resido*, 'to settle or sink down,') *Sediment*.

RESIDUAL AIR, see *Respiration*.

RESIMUS, 'turned up, bent back,') *Camus*.

RES'IN, *Resi'na*, *Rheti'nē*, (*F.*) *Résine*, (from *resu*, 'to flow,') A vegetable product, commonly dry and concrete, more or less brittle, inodorous or slightly odorous, insipid, or of an acrid warm taste; of a smooth, glassy fracture, heavier than water, inflammable, insoluble in water, soluble in alcohol, ether, and yolk of egg, and negatively electrifiable by friction. Many resins are used in medicine; the greater part are purgative and irritating. Some act like acrid poisons. See *Pinus sylvestris*.

RESIN, ACAROID, see *Xanthorrhœa*.—*r.* Botany Bay, see *Xanthorrhœa*.—*r.* Cayenne, *Caoutchouc*.—*r.* of New Holland, see *Xanthorrhœa*.—*r.* of the Spruce fir, see *Pinus abies*.—*r.* of Jalap, *Extractum Jalapæ resinosum*.—*r.* of Mayapple, *Resina Podophylli*.—*r.* of *Xanthorrhœa*, see *Xanthorrhœa*.

RESINA, Resin, see *Pinus sylvestris*.—*r.* Abietis, see *Pinus abies*, *Pinus picea*.—*r.* Abietis humida, see *Pinus abies*.—*r.* Abietis sicca, see *Pinus abies*.—*r.* Acaroides, see *Xanthorrhœa*.—*r.* Alba, see *Pinus sylvestris*.—*r.* Alba humida, see *Pinus abies*.—*r.* Benzoës, Benjamin.—*r.* Cayennensis, *Caoutchouc*.—*r.* Copal, *Copal*.—*r.* Elastica, *Caout-*

chouc.—*r.* Flava, see *Pinus sylvestris*.—*r.* Fluida Canadensis, see *Pinus balsamea*.—*r.* Jalapæ, *Extractum Jalapæ resinosum*.—*r.* Juniperi, *Sandaraç*.—*r.* Laricis, see *Pinus larix*.—*r.* Lentiscina, see *Pistachia lentiscus*.—*r.* Liquida pini balsameæ, see *Pinus balsamea*.—*r.* Liquida pini laricis, see *Pinus larix*.—*r.* Lutea Novi Belgii, see *Xanthorrhœa*.—*r.* Nigra, *Colophonia*.—*r.* Pini, see *Pinus sylvestris*.—*r.* Pini empyreumatica liquida, see *Pinus sylvestris*.—*r.* Pini abietis sponte concreta, see *Pinus abies*.

RESINA PODOPHYLLI, *Resin of May Apple*, (*Ph. U. S.* 1863.) *Podophyllin*. (Prepared from the powdered rhizoma of *Podophyllum* in the same manner as the *Resina Jalapæ* from *Jalap* root.) See *Podophyllum peltatum*. Dose, gr. j or gr. ij.

RESINA SCAMM'NI, *Extractum scammonii*.

RESINA STROBIL'NA, see *Pinus balsamea*, and *Pinus mughos*.—*r.* Terebinthi, see *Pistachia terebinthus*.

RESINATUM, *Rhetini'tes*. Wine impregnated with resin, used by the ancients as a stomachic.

RÉSINE, Resin.—*r.* de Cayenne, *Caoutchouc*.—*r.* Élastique, *Caoutchouc*.—*r.* de Gaïac, see *Guaiacum*.—*r.* Gomart, see *Bursera gummifera*.—*r.* de Lierre, see *Hedera helix*.

RÉSINÉINE, *Résinone*, *Résinéone*, (from *resin*,) A solid or liquid product obtained by the distillation of resin and lime or an alkali.

RESINOCE'NUM, *Rhetinoce'ron*, (*ῥητιν*, 'resin, and *κνος*, 'wax,') A mixture of resin and wax.

RÉSINONE, *Résinéine*.

RESINO'SA, in Pharmacy, resinous substances.

RÉSOLUTIFS, (*resolvere*, (*re*, and *solvere*, *solutum*,) 'to loosen,') *Discutients*.

RESOLUTIO, *Resolutio*.—*r.* et *Diabrosis Ventriculi*, *Gastromalacia*.—*r.* Membrorum, *Paralysis*.

RESOLUTIO MOR'BI. The resolution or disappearance of a disease; especially when without any critical evacuation.

RESOLUTIO NERVORUM, *Paralysis*.—*r.* Ventriculi atopeptica, *Gastromalacia*.—*r.* Virium, *Dialysis*.

RESOLUTION, *Resolutio*, *Discussio*. Removal or disappearance, as of a disease. Resolution is one of the terminations of inflammation, in which the inflamed part returns gradually, and without suppuration, to the healthy condition. Also, looseness, weakness.

RÉSOLUTION DES MEMBRES, *Paralysis*.

RESOLVENTIA, (from *resolvere*, 'to loosen, to dissipate,') *Discutients*.

RES'ONANCE, (*F.*) *Retentissement*, *Résonnance*, *Résonnement*, (*re*, and *sono*, 'I sound,') a return of sound; a resounding; a reverberation of sounds. A thrilling of the voice more loud than natural; or its existence in a part where it is not heard in health,—as detected by auscultation. A thickened and hardened state of the lungs, caused either by a mass of crude tubercles, or by inflammation, is generally considered to produce this phenomenon, by rendering the lung a better conductor of the murmur of the voice in the bronchi. Hence the origin of the term *Bronchophony*, (*bronchus*, and *φων*, 'voice,') (*F.*) *Voix bronchique*, *Voix tubaire*, *V. bourdonnante*.

Resonance is also used in the sense of resounding; as when the chest is said to be resonant on percussion.

RESONANCE, VESIC'ULOTYMPANIT'IC, see *Vesiculotympanic*.—*r.* Vibrations of, see *Sound*.

RES'ONANT, *Res'onans*, *Res'onus*, (*F.*) *Retentissant*, *Résonnant*. Resounding. See *Resonance*.

RESONITUS, *Contraffissura*.

RÉSONNANCE, *Resonance*.

RÉSONNEMENT, Resonance.

RESORBENS, (*re*, and *sorbere*, *sorptum*, 'to suck up,') Absorbent.

RESORPTIO, Absorption, Resorption—*r. Cutanea*, see Absorption—*r. Cutis*, see Absorption.

RESORPTION, *Resorptio*. Absorption of a fluid which has been previously deposited in a part, as of pus, blood, serum, &c.

RESPIRABLE, *Respirabilis*, (*respiratio*, and *habilis*, 'fit,') Capable of being breathed. Adapted for respiration.

RESPIRAMEN, (*respiro*, (*re*; and *spiro*, 'I take breath,') Respiration.

RESPIRAMENTUM, Respiration.

RESPIRATEUR, Respirator, Respiratory.

RESPIRATION, Respiration—*r. Abolita*, *Apnoea*—*r. Brevis*, *Dyspnoea*—*r. Difficilis*, *Dyspnoea*—*r. Stertens*, *Stertor*—*r. Sterterosa*, see *Stertor*—*r. Stertuosa*, *Stertor*.

RESPIRATION, *Respiratio*, *Anapnoë*, *Anapnoësis*, *Pneumosis*, *spiratio*, *Respiramen*, *Respiramentum*, *Respiratus*, *Respiritus*, *Spiritus*. A function proper to animals, the object of which is, to place the materials of the blood—the mixture of the venous blood with lymph and chyle—in contact with atmospheric air, in order that it may acquire the vivifying qualities which belong to arterial blood. The organs for executing this function are, in the mammalia, birds, and reptiles, the *lungs*. In man, the respiration consists of mechanical and chymical phenomena. The mechanical are *Inspiration* and *Expiration*. The evident chymical phenomena consist in the formation of a certain quantity of carbonic acid, the absorption of a part of the oxygen of the air, and the disengagement of a quantity of water in the state of vapour. In the healthy condition the respiration is easy, gentle, regular, and without noise. In man, the respirations are generally about 35 per minute in the first year of life; 25 during the second; 20 at puberty; and 18 in the adult age.

The air of respiration has been divided into *first*, the *residual air*, or that which cannot be expelled from the lungs, but remains after a full and forcible expiration, estimated at 120 cubic inches: *secondly*, the *supplementary or reserve air*, or that which can be expelled by a forcible expiration, after an ordinary outbreathing, valued at 130 cubic inches: *thirdly*, the *breath, tidal or breathing air*, valued at 26 cubic inches: and *fourthly*, the *complementary or complemental air*, or that which can be inhaled after an ordinary inspiration, which amounts to 100 cubic inches. This estimate gives 250 cubic inches as the average volume of air which the chest contains after an ordinary expiration.

RESPIRATION BRONCHIQUE, see *Murmur*, respiratory.

RESPIRATION, BRONCHOVESICULAR. A morbid modification of the respiratory murmur, having more or less of the characters of the bronchial and vesicular respiration combined.—(Flint.) It denotes incomplete solidification of the lung, as from the presence of tubercle.

RESPIRATION OF THE CELLS, *Murmur*, respiratory—*r. Coggedwheel*, *Respiration*, *jerking*—*r. Entrecoûpée*, *Respiration*, *jerking*—*r. Empêchée*, *Respiration*, *jerking*—*r. Exagérée*, see *puerile*—*r. Hypervésiculaire*, see *Puerile*.

RESPIRATION, JERKING, *Coggedwheel*, or *Wavy respiration*, (F.) *R. Saccadée* ou *Entrecoûpée* ou *Empêchée*. When the murmur of inspiration, in place of being continuous, is interrupted as by starts, it is called 'jerking.' It is a concomitant of incipient pleurisy, pleurodynia, spasmodic asthma, and tuberculosis of the lung with corresponding pleuritic adhesion.

RESPIRATION, LUCTUEUSE, see *Luc-*

tueur—*r. Metallic*, *Tintement métallique*—*r. Nulle*, see *Murmur*, respiratory.

RESPIRATION, POSTURAL, see *Ready method*—*r. Prone*, see *Ready method*.

RESPIRATION PROOF, *Dicimasia pulmonum*.

RESPIRATION, PUÉRILE, see *Puerile*—*r. Râpeuse*, see *Murmur*, respiratory—*r. Rude*, see *Murmur*, respiratory—*r. Saccadée*, *R. jerking*—*r. Silencieuse*, see *Murmur*, respiratory—*r. Soufflante*, *Murmur*, respiratory—*r. Supplémentaire*, see *Puerile*—*r. Tubal*, see *Murmur*, respiratory—*r. Wavy*, *Respiration*, *jerking*.

RESPIRATOIRE, Respiratory.

RESPIRATOR, (F.) *Respirateur*. Various instruments, under this name, have been devised for the purpose of tempering the air before it reaches the lungs in winter. One of the best is that of Jeffrey, which is made of several layers of fine wire tissue.

RESPIRATORIUM VENTRIS, Diaphragm.

RESPIRATORY, *Respiratorius*, (F.) *Respirateur*, *Respiratoire*. Appertaining to respiration. An epithet applied to the murmur heard by auscultation in a healthy adult, during inspiration and expiration. It marks the entrance of the air into the cells, and its passage out of them. See *Murmur*.

RESPIRATORY, EXTERNAL, NERVE, see *Thoracic nerves*.

RESPIRATORY GLANDIA, see *Encephalon*.

RESPIRATORY, INTERNAL, NERVES, *Diaphragmatic Nerves*—*r. Nerve of the face*, *Facial Nerve*—*r. Nerve, Superior*, *Spinal Nerve*.

RESPIRATORY PERIOD. The time from the commencement of one inspiration to the occurrence of the next. If this period be regarded as four seconds; a half second, it has been considered, may be occupied in health by inspiration; a half second in expiration, and two seconds in repose.

RESPIRATORY TRACT, *Tractus respiratorius*. A middle column of the spinal marrow, described by Sir Charles Bell as that whence the respiratory nerves originate. See *Nerves*.

RESPIRATUS, Respiration.

RESPIRITUS, Respiration.

RESSERRANT, (*re*, and *serre*, 'to lock up,') *Constipating*.

RESSERRÉ VENTRE, *Constipation*.

RESTA BOVIS, *Ononis spinosa*.

RESTAGNATIO, (*re*, and *stagnare*, *stagnatum*, 'to overflow,') *Eruetation*.

RESTAURANS, *Analeptic*, *Corroborant*.

RESTAURATIO, *Refectio*, *Refectus*, *Analepsis*, *Anatrepsis*, *Anacomidë*, *Anactes*, *Apocatasia*, *Recuperatio*, (*restaurare*, *restauratum*, 'to repair,') Restoration of strength, after disease or great fatigue; *Epanorthosis*, *Recreat*, *Redintegratio*, *Restauratio virium*. Return of the organs to their normal condition. *Recovery*, *Recuperation*, *Reparation*.

RESTAURATIO VIRIUM, see *Restauratio*.

RESTHARROW, *Ononis spinosa*.

RESTIFORM, *Restiformis*, (*restis*, 'a rope,' and *forma*, 'shape,') Shaped like a rope, as the 'corpora restiformia,' restiform bodies.

RESTIFORMIS PROCESSUS, *Coracoid*.

RESTITUTIO, (*restitu*, *restitutum*, (*re*, and *statuo*, 'to set up,') 'to set up again,') Reduction—*r. Organica*, *Morioplastice*—*r. Partium deperditarum*, *Morioplastice*.

RESTLESSNESS, *Agitation*.

RESTORATION, *Restauratio*.

RESTORATIVE, *Analeptic*, *Corroborant*.

RESTRICITIO INTESTINI RECTI seu *ANI*, (*re*, and *stringere*, *strictum*, 'to tighten,') *Stricture of the rectum*.

RESUMPTIVUS, *Resumptorius*, (*re* and *sumere*, *sumptum*, 'to take,') Causing to resume.

Resumptivum. Any agent calculated to restore flesh and strength to one who has been reduced.

RESUMPTORIUS, Resumptivus.

RESURRECTION MAN, Resurrectionist.

RESURRECTIONIST, *Resurrectio man, Body snatcher*, (*re*, and *surgere*, *surrectum*, 'to arise.') An individual who violates the sanctuary of the grave to obtain subjects for dissection.

RESUSCITATION, *Resuscitatio, Resuscitare*, (*re*, and *evocare* [*sursum citare se cernere*], 'to move up,' 'to stir up anew.') *Revivification*. The restoration of persons, apparently dead, to life. See Asphyxia and Suffocation.

RETCHING, ([Sax.] *pæcan*, 'to retch,') Vomiturition.

RETE, 'a net,' *Dictyon*, (F.) *Réseau*. A name given to the interlacing and decussation of blood-vessels, lymphatics, fibres, and nerves when they form a kind of network. Plexus.

RETE has also been applied to the *epiploon*.

RETE MAJUS, Epiploon, gastrocoelic.

RETE MINUS, Corpus mucosum.

RETE MIRABILE, (F.) *Réseau admirable, R. merveilleux*. A name given to the ramifications and anastomoses of the internal carotid and vertebral arteries at the base of the brain.

RETE MIRABILE seu RETE VASCULOSUM MALPIGHII is the plexus formed on the cells of the lungs by the minute ramifications of the pulmonary artery.

RETE MUCOSUM, Corpus mucosum.

RETE TESTIS, *Reté Vasculosum Testis* seu *Halleri*. An irregular plexus or network, formed by the communication of the *Vasa recta*, *Tubuli* seu *Ductuli recti*, *Tubuli seminiferi recti*, (F.) *Vaisseaux droits*, at the upper and back part of the testicle. These *Vasa recta* are sent out at the back part of the testicle, from the convoluted seminal tubes. The Rete Vasculosum Testis sends out from twelve to eighteen straight tubes, called *Vasa efferentia*, which convey the semen from the testis to the epididymus.

RETENTIO, Retention—*r. Aerea*, Dyspnoea—*r. Alvina*, Constipation—*r. Bilis*, Ischocolia.

RETENTIO FLUXUS HÆMORRHOÏDALIS. Suppression of the hemorrhoidal flux.

RETENTIO FŒTUS, Dystocia—*r. Lactis*, Ischogalactia—*r. Lochiorum*, Ischiolochia—*r. Mensium*, see Amenorrhœa—*r. Spermatidis*, Ischospemia—*r. Sutoris*, Hydroschesis.

RETENTION, *Retentio, Apolepsia, Apolepsia*, (*retinere*, *retentum*, (*re*, and *tenere*, 'to hold,') 'to hold back,') Accumulation of a solid or liquid substance in canals intended for its excretion, or in a reservoir intended to contain it for a short period.

RETENTION OF URINE, *Paruria Retentio nis vesicalis, Ischuria vesicalis, Retentio Urinæ*, (F.) *Rétention d'urine*. A disease in which the urine accumulated in the bladder cannot be evacuated; or, at least, cannot be passed without extreme difficulty. In the former case, the retention is said to be *complete*; in the latter, *incomplete*. Many writers have distinguished three degrees of this affection, to which they have given the names *Dysuria*, *Strangury*, and *Ischuria*. Retention of urine may depend upon loss of contractility, paralysis of the bladder, &c., or on some obstacle to the passing of the urine, as in cases of pressure of the womb on the bladder, of tumours in its vicinity, foreign bodies in its cavity, inflammation of the urethra, swelling of the prostate, stricture, &c. The pain is extremely violent, and the bladder may be found distended above the pubis. The treatment consists in introducing the catheter, or puncturing the bladder, and in combating the cause, which has produced the retention, by particular means adapted to each case.

RETENTISANT, (*re*, and *timere*, 'to tinkle,' to ring,) Resonant.

RETENTISSEMENT, Resonance—*r. Auto-phonique*, see Autophonia.

RETIA UTERI, (pl. of rete,) see Uterus.

RETICULAR, *Retiform, Reticularis, Retiformis*, *Amphiblastroïdes, Dictyôides, Retiformis*, (from *reticulum*,) (F.) *Réticulaire*. That which resembles a net. An epithet applied to many structures in the body.

RETICULAR BODY, Corpus mucosum—*r. Substance*, Cellular tissue.

RETICULARIA OFFICINALIS, Lichen pulmonarius.

RETICULUM, (dim. of rete,) The *Bonnet* or *Honeycomb bag* or *stomach*, (Sc.) *Kingshead*, (F.) *Bonnet, Réseau*—the second stomach of the ruminant animal. See also, Epiploon, and Rete.

RETICULUM CUTANEUM, Rete mucosum—*r. Mucosum, Rete mucosum*.

RETIFORM, (*rete*, and *forma*, 'shape,') Reticular.

RETINA, (from rete,) *Tunica retina seu reticularis seu retiformis seu intima seu nervosa oculi, Involucrum reti comparatum, Verricularis membrana, Amphiblastroïdes membrana*, (F.) *Rétine*. A soft, pulpy, grayish, semitransparent, very thin membrane; extending from the optic nerve to the crystalline, embracing the vitreous humour, and lining the choroid; without, however, adhering to either of those parts. It terminates by a defined edge—*marginis dentalis*—at the posterior extremities of the ciliary processes. It is constituted, according to most anatomists, by the expansion of the optic nerve. The retina appears to be formed of several laminae; so joined together, that it is difficult to discriminate them. One—the inner—is medullary, and pulpy; another—the outer—is stronger, and fibrovascular. The retina is the essential organ of vision; on it the images of objects are impressed. Both it and the optic nerve are devoid of general sensibility. They may be punctured or lacerated without pain being experienced. The nerve of general sensibility distributed to the eye is the fifth pair.

RETINACULA, see Retinaculum—*r. Morgagnii*, see Fræna of the valve of Bauhin.

RETINACULUM, (*retinere*, (*re*, and *tenere*, 'to hold,') 'to hold back,') A surgical instrument, formerly used in the operation of castration; and in that of hernia, to prevent the intestines from falling into the scrotum. *Retinacula* are also band-like extensions of a cellulomembranous structure, which seem to suspend the ovum in the Follicle of De Graaf. They appear to correspond to the chalazæ of the egg of the bird. See also *Bride*.

RETINACULUM TEN'DINUM PERONEORUM, *Vagina mulleolaris externa*. The ligamentous sheaths that retain in situ at the outer ankle the tendons of the peroneal muscles.

RETINAL, *Retinalis*. Relating or appertaining to the retina.

RÉTINE, Retina.

RETINITIS, *Inflammatio Retinæ, Amphiblastroïdis, Dictyôidis*, (F.) *Rétinite, vretina etitis*. Inflammation of the retina; an uncommon affection, except as an accompaniment of other forms of ophthalmia.

RÉTINOÏDÉ, see Rhétinolé.

RÉTINOLÉ, see Rhétinolé.

RETINOSCOPY, *Retinoscopia*, (*retina*, and *σκοπεω*, 'I view,') Examination of the retina.

RETINUS, (from rete,) Reticular.

RETOMBET, GROS, *Passerina tartonraira*.

RETORT, *Retorta, Balnea, Cornu, Cornuta, Cornes*, *Coramulæ*, (F.) *Cornes, retorques*, (*re*, and *torqueo, tortum*,) 'I bend back,') A vessel, commonly of glass or earthenware, employed in distillation. It may be plain, or have a tubulure for admitting substances whilst the distillation is

going on. In the latter case, it is said to be *tubulated*, (F.) *Cornu tubulé*.

RETRACT'ION, *Retract'io*, (*retrahere*, (re, and *trahere*, *tractum*), 'to draw back,') State of a part, when drawn towards the centre of the body or backwards:—as retraction of the testicles. Sometimes it is used synonymously with *shortening* or *deaving up*; as in *retraction* of the thigh.

RÉTRACTION DE LA MATRICE, *Retroversio uteri*.

RETRACTOR ANGULI ORIS, *Buccinator*.

RETRACTENS AU' RIS, (from *retrahere*, 'to draw back,') *Posterior seu Tri-ceps Au'ris*, *Tri-caudalis*, *Retractus Auriculæ seu Auriculi*, *Secundus proprius Auriculæ*, *Bicaudalis*, *Deprimens Auriculæ*, *Pro'prius Au'ris exterioris*, (F.) *Mastoido-conchiformis*, *Mastoido-orciculare* (Ch.), *Auriculare posterieur*. It consists of one or more small fasciculi, or fleshy, thin, somewhat irregular fibres, situated behind the ear. These are attached to the mastoid process, and pass horizontally forwards, to terminate at the convexity formed by the concha of the ear. If this muscle had any use, it would carry the ear backwards.

RETRAIT VENTRICULAIRE, Heart, hypertrophy of the.

RÉTRÉCISSEMENT, (re, and *étrécir*, [old F.] *estrécir*, (from [L.] *stringere*, *strictum*), 'to make narrow,') *Arctatio*, *Stricture*—*r. de l'Urèthre*, *Urethrostenosis*.

RETRIMENTA VESICÆ, Urine.

RETRIMENTUM, ('dregs,' from *re*, and *terere*, 'to cleanse as by rubbing,') *Excrement*.

RETRO (L), in composition, behind, backwards.

RETROCES'ION, *Retroces'sio*, *Retroces'sus*, *Retrogres'sus*, (*retrocedere*, *retrocessum*, (*retro*, and *cedere*, 'to give place,') The act of going back. A disappearance or metastasis of a tumour, eruption, &c., from the outer part of the body to the inner.

RETROCESSUS, *Retrocession*.

RETROFLEXIO UTERI, (*retro*, and *flectere*, *flezum*, 'to bend,') see *Retroversio uteri*.

RETROGRESSUS, (*retro*, and *gradior*, *gressus*, 'to step,') *Retrocession*.

RETROPHARYNGE'AL, *Retropharynge'us*, (*retro*, and *pharynx*, gen. *pharyngis*, 'the pharynx,') Relating to parts behind the pharynx,—as 'retropharyngeal abscess,—an abscess behind the pharynx.

RETROPULSION DE LA MATRICE, (*retro*, and *pellere*, *pulum*, 'to drive,') *Retroversio uteri*.

RETROU'TERINE, *Retrouteri'nus*, (F.) *Rétro-utérin*, (*retro*, and *uterus*.) Behind the uterus, as *retrouterine abscess*, (F.) *Abscès rétro-utérin*;—an abscess seated behind the uterus.

RETROVACCINATION, (*retro*, and *vaccination*.) Vaccination with matter obtained from the cow, after inoculating the animal with vaccine matter from the human subject.

RETROVER'SIO UTERI, (*retro*, and *vertere*, *versum*, 'to turn,') *Retrover'sion of the Uterus*, *Ædoptio'sis Uteri retrover'sa*, *Anapod'isis seu Anapod'is'mus Uteri*, (F.) *Rétroversion*, *Rétraction*, *Obliquité ou Rétroupsion de la Matrice*. A change in the position of the uterus, so that the fundus of the organ is turned towards the concavity of the sacrum, whilst the neck is directed towards the symphysis pubis. This displacement is commonly attended with constipation, tenesmus, and retention of urine. Its usual period of occurrence is between the third and fourth months of pregnancy; before the uterus has escaped above the superior aperture of the pelvis. In the treatment, the catheter must be regularly used twice a day, until the uterus, by its growth, rises above the pelvis. The catheter should be small, flat, and curved considerably more than common, in consequence of the distorted course of the ure-

thra. The bowels must be kept open: and absolute rest, in a recumbent posture, be enjoined. Should it be impracticable to draw off the urine, attempts must be made to replace the uterus. The woman being on her hands and knees, the fore and middle fingers of the accoucheur's left hand are to be passed up the rectum to the fundus uteri: which they must elevate, whilst the cervix uteri is carefully depressed by two fingers of the right hand in the vagina. Should the fingers employed to elevate the fundus not be long enough to effect this object, a piece of whalebone may be substituted, having a small piece of sponge attached to one extremity as a pad.

A simple inclination, or bending of the uterus, backwards, is termed *Retroflexion of the Uterus*, *Retroflex'io Uteri*: not unfrequently, however, this is used synonymously with *Retroversion of the Uterus*.

RÉTROVERSION DE LA MATRICE, *Retroversio uteri*.

REUNIO, (re, and *unire*, 'to unite,') *Synthesis*—*r. Partium separatarum*, see *Synthesis*—*r. per Primam intentionem*, see *Intention*—*r. per Secundam intentionem*, see *Intention*—*r. Vasorum*, *Anastomosis*.

RÉUNION PRIMITIVE, see *Intention*—*r. Secondaire*, see *Intention*,

REVACCINATION, see *Vaccina*.

REVALENTA ARABICA, see *Erym*.

RÉVASSERIE, (from *rêve*,) *Incoherence*; see *Somnium*.

RÊVE, *Somnium*.

RÊVEIL, (re, and *éveiller*, 'to awaken,') see *Sleep*—*r. en Sursaut*, see *Sleep*.

REVELENT, (re, and *tellere*, 'to pluck,') *Derivative*.

RÉVERIE, *Aphelexia*.

REVERY, *Aphelexia*.

REVIVIFICATION, (re, *vivere*, 'to live,' and *facere*, 'to make,') *Resuscitation*.

RÉVULSIF, (re, and *tellere*, *culsum*, 'to pluck,') *Derivative*.

REVUL'SION, *Revul'sio*, *Antis'pasis*. The act of turning the principle of a disease from the part in which it seems to have taken its seat. Rubefacients, vesicatories, bleeding in the foot, &c., are often used for this purpose. See *Derivation*.

REVULSIVE, *Derivative*—*r. Bleeding*, see *Bloodletting*.

REVULSORIUS, *Derivative*.

REX AMARORUM, 'king of bitters,' *Soula-men amara*—*r. Metallorum*, *Aurum*.

REYNOLDS'S SPECIFIC, see *Vinum colchici*.

REZZLE, *Wheeze*.

RHA, (*ῥα*, and *ῥη*, called, however, after the *Rha* or *Volga* in *Pontus*), *Rheum*.

RHABARBARUM, (*ῥα*, and *ῥαββαρος*, 'foreign,') *Rheum*—*r. Album*, *Convolvulus mechoacan*—*r. Antiquorum*, *Rheum Rhaponticum*—*r. Dioscoridis*, *Rheum Rhaponticum*—*r. Monachorum*, *Rumex patientia*—*r. Nigrum*, *Convolvulus jalapa*—*r. Pauperum*, *Thalictron*—*r. Plebeiorum*, see *Rhamnus frangula*—*r. Sibericum*, *Rheum*—*r. Tartaricum*, *Rheum*—*r. Turcicum*, *Rheum*.

RHABARBE, *Rheum*—*r. Blanche*, *Convolvulus mechoacan*—*r. Fœne*, *Thalictron*—*r. des Indes*, *Convolvulus mechoacan*—*r. des Moines*, *Rumex Alpinus*—*r. Sauvage*, *Rumex patientia*.

RHABDOIDES (SUTURA), (*ῥαβδαίος*, from *ῥαβδος*, 'a rod or staff,' and *αἶος*, 'resemblance,') *Sagittal suture*.

RHABILLEUR, (re, and *habiller*, 'to dress,') *Remoeur*.

RHACHETÆ, *Rhachitæ*.

RHACHETRUM, (*ῥαχητρον*, from *rhachia*,) The middle of the spine.

RHACHLEI, *Rhachitæ*.

RHACHIAGRA, *Rachisagra*.

RHACHIALGIA, Rachialgia, Rhachiodynia, Spinal irritation.

RHACHIALGITIS, (*rhachis*, αλγος, 'pain,' and *itis*.) Spinal irritation, Myelitis.

RHACHIDAGRA, Rachisagra, Rachialgia, Rhachiodynia.

RHACHIOCYPHOSIS, Gibbositas, Hump.

RHACHIODYN'IA, (*rhachis*, and *dύνη*, 'pain;') *Rhachial'gia*. Pain in the vertebral column.

RHACHIOMYELITIS, (*rhachis*, and *myelitis*.) Myelitis.

RHACHIOMYELOPHTHISIS, (*rhachis*, μυελος, 'marrow,' and *phthisis*.) Tubercular.

RHACHIOPARALYSIS, (*rhachis*, and *paralysis*.) Paraplegia.

RHACHIOPLEGIA, (*rhachis*, and *πληγη*, 'a stroke,') Paraplegia.

RHACHIOSCOLIOMA, (*rhachis*, and *scolioma*.) Scoliosis.

RHACHIOSCOLIOSIS, (*rhachis*, and *scoliosis*.) Scoliosis.

RHACHIOSTROPHO'SIS, (*rhachis*, and *στρέφω*, 'I turn;') *Contortio colum'næ vertebral'is*. Deviation of the vertebral column.

RHACHIOTOMUS, *Rhachiot'omum*, (*rhachis*, and *τομή*, 'incision,') An instrument for opening the vertebral column.

RHACHIPAG'IA, *Rachip'ia*, (*rhachis*, and *παγω*, 'I fasten,') A monstrous union of two foetuses, in which they are connected at the spine.

RHACHIPARALYSIS, (*rhachis*, and *paralysis*.) Paraplegia.

RHACHIPHYMA, Rachiphyma.

RHACHIRRHEUMA, gen. *Rhachirrhœum'atis*, (*rhachis*, and *ῥευμα*, 'a deluxion, rheumatism,') Lumbago.

RHACHIS, gen. *Rha'chidia*, (*ῥαχίς*, gen. *ῥαχιδός*.) Vertebral column—r. Nasi, see Nasus.

RHACHISAGRA, Rachisagra.

RHACHIT'Æ, (from *rhachis*.) *Rhachit'æ*, *Rachit'æ*, *Rhachet'æ*, or *Rachit'æ*. The muscles of the spine.

RHACHITES. Spinal; Medullaspinalis, (*ῥαχίτης νεύρος*.) see also Rachitis, (*ῥαχίτης vocos*.)

RHACHITICUS, Rachiticus.

RHACHITIS, Rachitis—r. Adultorum, Mollities ossium—r. Spinalis, Rachisagra.

RHACHITISMUS, (from *rhachis*.) Rachitis.

RHACHOSIS, Rachosis.

RHA'CHUS, (*ῥαχός*.) *Rha'cus*, *Rha'gus*, (from *ῥαγω*, or *ῥήγνυμι*, 'to break,') A wrinkle. A laceration of the skin. A lacerated ulcer—*ul'cus lac'erum*.

RHACUS, (*ῥαχός*.) Rhachus, Wrinkle.

RHÆBAUCHEN, (*ῥαβδος*, 'crooked,' and *αυχμη*, 'neck,') Torticollis.

RHÆBOCRANIA, (*ῥαβδος*, 'crooked,' and *κρανον*, 'skull,') Torticollis.

RHÆBODERE, (*ῥαβδος*, 'crooked,' and *δερν*, 'neck,') Torticollis.

RHÆBOSCELIA, (*ῥαβδος*, 'crooked,' and *σκελος*, 'leg,') see Rhæbosis.

RHÆBO'SIS, (from *ῥαβδος*, 'crooked,' and *osis*.) Crookedness, as of the legs,—*Rhæboscel'ia*, *Scolio-cambol'ia*.

RHAGADES, (*ῥαγᾶδες*.) pl. of *rhagus*, (*ῥαγας*, gen. *ῥαγᾶδος*.) *Rhagad'ia*, (from *ῥαγω*, or *ῥήγνυμι*, 'to break or rend,') *Scissur'æ*, *Fissur'æ*, *F'issæ*. Ruptures, chaps, or narrow and long ulcers, which form near the origin of mucous membranes, especially around the anus, and which are generally owing to the syphilitic virus.

RHAGADES, see Fissure.

RHAGE, *Rhag'ia*, *Ra'gē*, (*ῥαγη*.) Suffixes denoting violent rupture or discharge, as in *Hemorrhage*, *Menorrhagia*, &c. Also, a fissure.

RHAGES, (from *ῥαξ*, gen. *ῥαγος*, 'a grape,') *Uvæ passæ*.

RHAGICUS, (*ῥαγικός*.) Rhagoides.

RHAGOIDEITIS, (*rhagoides* (*tunica*.) and *itis*, *Nivitis*.)

RHAGOÏDES, *Rhago'des*, *Rhagoïdeus*, *Rha'giens*, (*ῥαγίδης*, from *ῥαξ*, gen. *ῥαγος*, 'a berry,' 'a grape,') Grape-like; as *Tunica Rhagoides*: the uvea. Also, the choroid coat of the eye.

RHAGUS, (*ῥαγος*.) Rhachus, Wrinkle.

RHAMNI BACCÆ, see Rhamnus—r. Succus, see Rhamnus.

RHAM'NUS, (*ῥαμνος*.) *R. catharticus*, *Sp'ina cervi'na* seu *domestica* seu *infectoria*, *Rhamnus solutivus*, *Cervip'ina*, *Buck'thorn*, *Purg'ing buck'thorn*, (F.) *Nerprun purgatif*, *Bourguépine*, *Épine de cerf*. Family, Rhamnaceæ. Ser. Syst. Pentandria Monogynia. The berries—*Rhamni Bucca*—have a faint and disagreeable odour; and bitterish, nauseous taste. Their juice—*Rhamni Succus*—is cathartic, and apt to excite griping, which must be obviated by the use of diluents.

RHAMNUS ALATER'SUS, *R. Hispanicus*, *Alater'us latifolius*, *Com'mon alater'us*, (F.) *Alaterne*. The leaves have been used as detergents and astringents, in the form of gargle especially.

RHAMNUS FRANGULA, *Black Al'der*, *Fran'gula Al'na*, *Al'na Nigra*, (F.) *Anne noir*, *Bourdane*, *Bourguène*. All the parts of this tree, as well as of the common alder, are astringent and bitter. The bark is most so. The inner, yellow bark of the trunk or root, once called *Rhabar'barum plebeio'rum*, acts as a cathartic. It has also been employed as an anthelmintic.

RHAMNUS HISPANICUS, *R. alaternus*—r. Jujuba, see Jujube—r. Lotus, see Jujube—r. Ziziphus, see Jujube.

RHANTER, (*ῥαντηρ*, from *ῥαίνω*, 'to besprinkle,') Canthus, (greater.)

RHAPHANEDON, Raphanodon.

RHAPHANIA, Raphania.

RHAPHANIS, Raphanus hortensis.

RHAPHANUS, Rapa, Raphanus hortensis.

RHAPHE, Raphe, Suture.

RHAPHIDAGOGUS, (*ῥαφίς*, gen. *ῥαφίδος*, 'a needle,' and *αγω*, 'to lead,') *Porte-aiguille*.

RHAPHION, (*ῥαφίον*, dim. of *ῥαφίς*.) Needle.

RHAPHIS, (*ῥαφίς*.) Needle. Also; Rapa.

RHAPHOSYMPH'YSIS, (*ῥαφη*, 'suture,' and *συνήνσις*, 'union,') Union by suture,—as of the bones of the cranium.

RHAPHUS, (*ῥαφος*.) Rapa.

RHAPHYS, (*ῥαφος*.) Rapa, Raphanus hortensis.

RHAPONTICUM, (*ῥα*, or *rhabarb*, of Pontus.) *Rheum Rhaponticum*—r. Vulgare, *Centaurea celtaurium*.

RHAPONTIQUE, *Rheum rhaponticum*.

RHAPS, (*ῥαπυς*.) Raphanus hortensis.

RHAPYS, (*ῥαπυς*.) Raphanus hortensis.

RHATANIA, *Krameria ratanhia*.

RHAX, (*ῥαξ*, gen. *ῥαγος*.) see *Vitis vinifera*.

RHEGE, (*ῥηγη*.) Cramp, Contusion, Laceration.

RHEGMA, gen. *Rheg'matis*, (*ῥήγμα*.) Cramp, Contusion, Laceration, Rhexis—r. Ligamentare, Apospasma.

RHEGMUS, (*ῥηγμος*.) Cramp, Contusion, Laceration.

RHEMBASMUS, (*ῥεμβασμος*, from *ῥεμβω*, 'I wander about,') Somnambulism.

RHEMB'Æ, *Rhem'bæ*, (*ῥεμβή*, 'a wandering about,' from *ῥεμβειν*, 'to wander about,') Irregularity in the return of febrile paroxysms.

RHEMBUS, (*ῥεμβος*.) Rheembe.

RHENCHUS, (*ῥεγχος*.) Rattle, Snoring.

RHENXIS, (*ῥεγξίς*.) Rattle, Snoring.

RHEON, Rheum.

RHÉOPHORE, (*ῥοη*, 'a current,') and *φerein*, 'to carry,') Electrode.

RHETINE, (*ῥητινη*, from *ῥεω*, 'to flow,') Resin.

RHETINITE'S, (*ῥητινίτης*.) Resinatum.

RHETINOCERON, (*rhétine*, and *κερος*, 'wax,') Resinatum.

RHÉTINOÏDES, see *Rhétinolé*.

RHÉTINOLÉ, (from *rhétine*.) A name given by M. Béral to a medicine which results from the

union of a resin with other medicinal substances. The *Rhétinolés* of Bérál differ from the *Rhétinoïdes* in containing but one resin, whilst in the latter there are several.

RHÉTINOTON, (from *rhétine*,) Resination.

RHETORANCHE, (*ῥητωρ*, 'a public speaker,' and *αἴχμη*, 'to strangle,') see Pharyngitis, follicular.

RHEUM, (pronounced *room*,) *Rheū'ma*, (F.) *Rhūme*; (*ῥέω*, from *ῥέω*, 'to flow,') Any thin watery discharge from mucous membranes or skin; as the thin discharge from the air passages arising from cold.

RHEUM, SALT. A popular name in the United States, like 'tetter,' for various cutaneous affections of the eczematous and herpetic forms more especially. It is at times applied to a kind of chronic psoriasis; and likewise to cutaneous affections in those who immerse the hands in metallic or acid solutions. It has been defined "a vague and indefinite popular name, applied to almost all the non-febrile cutaneous eruptions which are common among adults, except ringworm and itch."

RHEŪM, *Rhabar'barum*, *Rhe'on*, *Rha*, *Rheū'm*, *Lap'athum orientale* seu *Chinen'sē*, *Rhabar'barum vel'rum* seu *Tartar'icum* seu *Siber'icum* seu *Tur'cium*, *Rhū'barb*, (F.) *Rhabarbe*, *Rhubarbe*, From *Rha*, ('*Ῥα*, 'Pa,') now called *Volga*, a river in Russia from which it was first brought. *Ord.* Polygonaceæ. *Sex. Syst.* Enneandria Trigynia. The root—*Rheū'm*—is the part used in medicine. Its odour is aromatic, peculiar, and rather nauseous; taste, somewhat aromatic, subacid, bitterish, and astringent. It colours the saliva and the urine yellow. It is easily pulverized, and forms a powder of a bright buff-yellow colour. Both water and spirit extract its virtues. *Rhubarb* root is purgative, stomachic and astringent. [?] It is chiefly used for the properties first mentioned. Externally, the powder is sometimes sprinkled over ulcers, to assist their granulation. Dose, $\mathfrak{ss}$ to $\mathfrak{ss}$ of the powder. In a smaller dose, it is stomachic. Numerous species of *Rheum*, *Rh. palmatum*, *Rh. undulatum*, and *Rh. compactum*, yield the *rhubarb-root*.

The *rhubarbs* most used in the United States are the *Chinese'* or *India Rhū'barb*—*Rheū'm Sinen'sē* seu *Ind'icum*; the *Rus'sian*, *Tur'key* or *Bucharian Rhū'barb*—*Rheū'm Rus'sicum* seu *Tur'cium*; and *Europe'an Rhū'barb*.

RHEUM COMPACTUM, *Rheum*—*r.* Indicum, see *Rheum*—*r.* Palmatum, *Rheum*.

RHEUM RHAPONTICUM, *Rhapont'ic Rhū'barb*; *Rhapont'icum*, *Rhabar'barum Dioscor'idis* seu *Antiquo'rum*, (F.) *Rhapontique*, *Rhapontic*, *Rapontic*. The root of this seems to have been the *rhubarb* of the ancients. It is more astringent than that of the *rheum palmatum*, and less purgative; for the latter purpose, two or three drachms being required for a dose. *French*, *Rhapontic*, or *Krimea Rhubarb*, is obtained from this species, as well as from *Rh. undulatum*, and *Rh. compactum*.

Rheum rhaponticum is the *Pie rhubarb* or *Pie plant* cultivated for its fleshy esculent leafstalks.

RHEUM RUSSICUM, see *Rheum*—*r.* Sinese, see *Rheum*—*r.* Turcicum, see *Rheum*—*r.* Undulatum, see *Rheum*.

RHEŪ'MA, gen. *Rheū'matis*, (*ῥέω*, gen. *ῥέω*, from *ῥέω*, 'to flow,') Catarrh, Diarrhœa, *Rheum*, *Rheumatism*. Also, inflammation of a fibrous tissue, as in rheumatism and gout. In composition, a flow, a defluxion; also, rheumatism.

RHEUMA CATARRHALE, Bronchitis, (chronic)—*r.* Epidemicum, Bronchitis, chronic, Catarrh, epidemic—*r.* Pectoris, Catarrh—*r.* Uteri, Leucorrhœa.

RHEUMAPYRA, (*rheuma*, and *πῦρ*, 'fever,') *Rheumatism*, acute.

RHEUMARTHRTIS, (*rheuma*, and *arthritis*,) see *Rheumatism*, acute.

RHEUMARTHROSIS, (*rheuma*, and *arthrosis*, 'an affection of the joints,') see *Rheumatism*, acute.

RHEUMATALGIA, (*rheuma*, and *αλγος*, 'pain,') *Rheumatism*, chronic—*r.* Arthritica, see *Rheumatism*, acute.

RHEUMATIC, (*ῥευματικός*,) *Rheumat'icus*, *Rheumat'ic'nal*, (F.) *Rhumat'ismal*, *Rhumat'ique*. Belonging to rheumatism; as rheumatic fever. Also, applied to one affected with rheumatism; (F.) *Rhumat'isant*.

RHEUMATIC DIATHESIS, *Diath'esis rheumat'ica*. The special condition of the body that gives occasion to rheumatism. It may exist without presenting the usual phenomena in the joints; whilst the heart may be seriously affected.

RHEUMATICS, *Rheumatism*.

RHEUMATISM, *Catarrh'ea*, (from *rheuma*,) *Do'lor Rheumat'icus et arthrit'icus*, (Hoffman.) *Rheumat'osis*, *Myosot'is*, *Rheū'm*, *Myodyn'ia*, (vulg.) *Rheū'matiz*, *Rheumat'ics*, (F.) *Rhumatisme*. A kind of shifting phlegmasia or neuralgia, sometimes seated in the muscles, sometimes in the parts surrounding the joints; and at others, within them. Hence the names *Muscular*, *Articular*, and *Synov'ial*, which have been applied to it. The disease may be *acute*, or *chronic*.

RHEUMATISM OF THE ABDOMEN, *Cœliorrheuma*.

RHEUMATISM, ACUTE, *Rheumat'ismus acut'us* seu *articulo'rum acut'us* seu *univers'al'is febr'ilis* seu *inflammato'rius* seu *hypersthen'icus*, *Arthro'sia acut'a*, *Rheumatop'gra*, *Rheumap'gra*, *Fe'bris rheumat'ica inflammato'ria*, *Syn'ocha rheumat'ica*, *Myosot'is*, *Myi'tis*, *Cau'na* seu *Arthri'tis rheumat'ismus*, *Ar'thorrheumat'ismus*, *Synov'i'tis rheumat'ism'al'is*, *Hæmoarthri'tis*, *Rheumat'ic Fe'ver*, (F.) *Rhumatisme aigu*, *Rh. articulaire aigu*, *Fièvre rhumatismale*, usually comes on with the ordinary symptoms of fever; soon after which, or simultaneously, or even before the appearance of febrile signs, excruciating pains are felt in different parts of the body, particularly in the larger joints, which are more or less red and swollen; the pain shifting from one to the other, at times with great rapidity. The disease rarely terminates in less than six weeks, during the greater part of which period the febrile symptoms remain severe; and, what is peculiar to the disease, the skin may be covered daily with a profuse perspiration, although it feels extremely hot, and the pulse appears in no way modified by it. It is one of the essential symptoms of the affection, and consequently affords no relief. The only danger to be apprehended from acute rheumatism, notwithstanding the apparent severity of the symptoms, is the translation or extension of the disease—*metastat'ic rheumatism*—to some internal part, especially to the heart. This tendency must always be borne in mind. *Acute rheumatism* seldom terminates in *chronic*; they who are liable to the former are rarely so to the latter, and conversely. The disease will generally run its course in spite of treatment. The usual antiphlogistic plan had better be pursued during the first days of the disease. Bloodletting may be employed, if the vigour of the patient and of the disease seem to require it; but, although the blood is generally buffy, bleeding does not usually seem to afford much relief. The bowels must be kept open; effervescent draughts be administered, and opium be given in full doses to allay pain. This is, generally, all that is necessary during the first week; after which, advantage will usually be obtained from the exhibition of the sulphate of quinia with sulphuric acid. It does not augment the excitement, and will be found proper in almost all cases. *Rheumatic inflammation* is very different from other.

varieties, and this may account for the anomaly. After a few weeks the disease usually goes off, but leaves the patient very liable to a recurrence on slight exposures or errors of diet. At times, acute rheumatism of the joints puts on the appearance of gout, and seems to be a complication of the two affections. It is then called *Gouty* or *Arthritic rheumatism* or *Rheumatism of gout*, *Rheumatoid arthritis*, *Rheumatism of the joints*, *Rheumatism of the joints*, *Arthritis rheumatica* seu *juvenile*.

RHEUMATISM OF THE ARM, *Brachiorrheuma*—*r. Arthritis*, see *Rheumatism*, acute—*r. of the Bladder*, *Cystorrhœma*—*r. of the Bowels*, *Enterorrhœma*.

RHEUMATISM, CAP'SULAR, *Rheumatismus capsularis*. Rheumatism seated in the lining membrane of the joints and bursæ of the tendons. The parts most liable to its attacks are the feet and hands, where it is generally easily recognized by the enlargement of the joints; but the peculiar characters of the disease are, perhaps, most strikingly seen when it attacks the knee.

RHEUMATISM OF THE CHEST, *Pleurodynia*.

RHEUMATISM, CHRONIC, *Arthrosis chronica*, *Rheumatismus chronicus* seu *nonfebrilis* seu *vulgaris* seu *inveteratus* seu *habitualis* seu *frigidus*, *Rheumatoidia*, *Rheumatologia*, *Rheumatodynia*, *Arthrodynia*, *Arthritis arthrodynia*, (F.) *Rhumatisme chronique*, *Rhumatologie*, is attended with pains in the hips, shoulders, knees, and other large joints. These are at times confined to one joint; at others, shift from one to another, without occasioning inflammation or fever. In this manner the complaint often continues for a great length of time, and then goes off. There is no danger attendant upon it; but the patient may become lame, and is always liable to painful recurrences. Neither variety generally terminates in suppuration; but effusion of coagulable lymph is apt to occur, so as to occasion permanent thickening of the parts. The great preventive of chronic rheumatism, and one of the most valuable curative agents, is flannel, worn next the skin. The whole class of rubefacients may, also, be used with advantage. The warm bath, especially the natural warm bath, the temperature of which does not vary, is, perhaps, the most successful remedy of all.

RHEUMATISM OF THE EXTREMITIES, *Acorrhœma*—*r. of the Face*, *Prosoporrhœma*—*r. Gouty*, see *Rheumatism*, acute—*r. of the Heart*, *Cardiorrhœma*—*r. of the Hip*, see *Neuralgia femoropoplitea*—*r. Lead*, see *Lead rheumatism*—*r. of the Leg*, *Seclorrhœma*—*r. of a Limb*, *Merorrhœma*—*r. Local*, *Merorrhœma*—*r. of the Loins*, *Lumbago*—*r. Metastatic*, see *Rh. acute*—*r. of the Neck*, *Torticollis*—*r. Partial*, *Merorrhœma*.

RHEUMATISM, PRÆABDOMINAL, (F.) *Rhumatisme pré-abdominal*. Rheumatism of the muscles of the anterior and lateral portions of the abdomen; supposed by MM. Gênest and Requin to be the same as the nervous affection of the peritoneum of Gooch.

RHEUMATISM ROOT, *Jeffersonia Bartoni*.

RHEUMATISM OF THE SKIN, see *Dermalgia*—*r. Topical*, *Merorrhœma*—*r. of the Uterus*, *Metorrhœma*.

RHEUMATISM, VIS'CERAL. Rheumatism affecting the muscular or fibrous tissues of the viscera.

RHEUMATISM WEED, *Pyrola maculata*, *P. umbellata*.

RHEUMATISMAL, *Rheumatic*.

RHEUMATISM O'ID, *Rheumatismoides*, *Rheumatoid*, *Rheumatoides*, (*rheumatism*, and *oides*, 'resemblance,') Resembling rheumatism.

RHEUMATISMUS, *Rheumatismus*—*r. Abdominis*, *Coeliorrhœma*—*r. Acutus*, *Rheumatismus*, acute—*r. Articulatorum acutus*, *Rheumatismus*, acute—*r. Brachii*, *Brachiorrhœma*—*r. Cancerosus*, *Neuralgia faciei*—*r. Cervicis*, *Torticollis*—*r. Colli*, *Torti-*

collis—*r. Cordis*, *Cardiorrhœma*—*r. Cruris*, *Seclorrhœma*—*r. Dorsalis*, *Lumbago*—*r. Extremitatum*, *Acorrhœma*—*r. Faciei*, *Prosoporrhœma*—*r. Febrilis*, *Dengue*—*r. Inflammatorius* seu *hypersthenicus*, *Rheumatismus acutus*—*r. Intestinorum*, *Enterorrhœma*—*r. Intestinorum cum ulcere*, *Dysentery*—*r. Larvatus*, *Neuralgia faciei*—*r. Localis*, *Merorrhœma*—*r. Membrorum singulorum*, *Merorrhœma*—*r. Non febrilis*, *Rheumatismus*, acute—*r. Partialis*, *Merorrhœma*—*r. Pectoris*, *Pleurodynia*—*r. Phlegmonodes*, *Myositis*—*r. Spurius nervosus*, *Neuralgia*—*r. Universalis febrilis*, *Rheumatismus*, acute—*r. Uteri*, *Metorrhœma*—*r. Vesicæ urinariæ*, *Cystorrhœma*—*r. Vulgaris* seu *inveteratus* seu *habitualis* seu *frigidus*, *Rheumatismus*, chronic.

RHEUMATIZ, *Rheumatism*.

RHEUMATODYNIA, (*rheuma*, and *δύνη*, 'pain,') Rheumatism, chronic.

RHEUMATOID, *Rheumatismoid*.

RHEUMATOPHTHALMIA, *Rheumatophthalmia*, (*rheuma*, and *ophthalmia*,) Rheumatic ophthalmia.

RHEUMATOPYRA, (*rheuma*, and *πύρ*, 'fever,') Rheumatism, acute.

RHEUMATOSIS, (from *rheuma*, and *osis*,) Rheumatism.

RHEUMOPHTHALMIA, (*rheuma*, and *ophthalmia*,) Rheumatophthalmia, Scleritis.

RHEX'IS, *ρήξις*, from *ρήγνυμι*, 'to break,') *Rhegma*, *Dirup'tio*, *Rup'tio*, *Ruptura*, *Rupture*. Rupture of a vessel or organ. Spontaneous opening of an abscess.

RHEXIS OC'ULI. Bursting of the eye from any cause; spontaneous or excited.

RHICNOS'IS, (*ῥιχνωσις*, from *ῥιχνωμαι*, 'to become wrinkled,') *Cutis corrugatio*. Wrinkling of the skin from extenuation of the body; opposed to *επκασις*, or distension from repletion.

RHIGOS, (*ῥίγος*,) Rigor.

RHIN, gen. *Rhīnis*, (*ῥῆν*, gen. *ῥίνας*,) Nasus.

RHINAL'GIA, *Rhinodynia*, (*rhin*, and *αλγος*, 'pain,') Pain in the nose.

RHINAT'GIA AB-INTRU'SIS. Pain in the nose from foreign bodies.

RHINAL'GIA VERMINOSA. Pain in the nose from worms.

RHINAN'CHONĒ, *Rhinan'gia*, *Rhinostegnosia*, (*rhin*, and *αγχων*, 'narrowness,') Contraction of the nose. Snuffles.

RHINANGIA, *Rhinanchone*, Snuffles.

RHINENCEPHALIA, (*rhin*, εν, 'in,' and *κεφαλή*, 'head,') Cyclopia.

RHINEN'CHYSIS, *Rhinenchysis*, (*rhin*, εν, 'in,' and *χύω*, 'I pour,') An infusion or injection, made into the nostrils, with a syringe called *Rhinenchites*.

RHINES, (pl. of *rhin*,) Nares.

RHINI'TIS, *Nasi'tis*, (*rhin*, and *itis*,) Inflammation of the nose. Coryza.

RHINITIS ULCEROSA, *Ozæna*.

RHINOBLENNORRHOEA, *Rhinorrhœa*.

RHINOC'ACĒ, *Coryza scarlatino'sa*, (*rhin*, and *κακός*, 'evil,') The offensive discharge from the nose, and other symptoms indicating a morbid condition of the Schneiderian membrane, that accompany the worst forms of scarlatina.

RHINOCATARRHUS, *Coryza*.

RHINOCNES'MUS, (*rhin*, and *κνησις*, 'itching,') Itching of the nose.

RHINODACRYOLITHUS, (*rhin*, *δακρυον*, 'a tear,' and *λίθος*, 'a stone,') see *Dacryolite*.

RHINODYNIA, (*rhin*, and *δύνη*, 'pain,') Rhinialgia.

RHINOLITE, (*rhin*, and *λίθος*, 'a stone,') see *Dacryolite*.

RHINOPHO'NIA, (*rhin*, and *φωνή*, 'the voice,') A nasal voice, called, also, *Dyspho'nia immodulata nasalis*, *Poropho'nia nasalis* seu *res'onans*,

Pœllis' mus na'sitas, Mycteropho'nia, Na'sitas, Vox seu Parala'tia nasa'lis. Speaking through the nose.

RHINOPLASTY, *Rhinoplast'ia, Rhinoplast'ica* (ars), (F.) *Rhinoplastie*, (rhin, and πλασσω, 'I form.') The operation for forming a new nose, which dates far back. Three chief methods have been described: 1. The method of *Celsus*, which consists in bringing forward the skin of the cheeks, and forming a septum, if necessary, from the upper lip. 2. The *Italian* or *Taliacotian method* or *operation*, called, also, the *German method*, on account of the application made of it by Von Gräfe, which consists in paring the edges of the nostrils, and uniting them to a graft taken from the arm, but left attached to it, until union has taken place; and 3. The *Indian method*—practised for ages in the East Indies, where cutting off the nose has been a common punishment for criminals—now generally adopted, in which the graft is taken from the forehead.

RHINOPOLYPUS, (*rhin*, and *polypus*), Polypus of the nose.

RHINOP'TIA, (*rhin*, and *οπτοια*, 'I see.') The act of seeing over the nose. Squinting over the nose.

RHINORRHAGIA, (*rhin*, and *ρῆγν*, 'a breaking forth,') Epistaxis.

RHINORRHAPHY, *Rhinorrhaph'ia*, (*rhin*, and *ρῆγν*, 'a suture.') An operation for the removal of epicanthis, which consists in pinching up a longitudinal fold of the skin, including this fold in two elliptical incisions, removing it, and bringing together the edges of the wound by harelip suture.

RHINORRHŒA, *Rhinoblenorrhœ'a*, (*rhin*, and *ρῆω*, 'I flow.') A discharge of limpid mucus from the nose without any inflammatory symptom. A gleet from the nose.

RHINOSCOPE, *Rhinoscop'ium*, (*rhin*, and σκοπεω, 'I view.') An instrument of the speculum kind for inspecting the nasal passages. The operation is termed *Rhinoscopy*, *Rhinoscop'ia*.

RHINOSCOPY, see Rhinoscope.

RHINO'SIS, (*ρῆσις*, 'a skin, leather,'—or from ῥυκος, 'rugous.') The state of looseness and excess of skin observed in phthisis.

RHINOSTEGNOSIS, (*rhin*, and στεγνωσις, 'a constriction or obstruction,') Rhinanehene.

RHIPTASMUS, (*ρῆπτασμος*, from *ρῆπτω*, 'to throw,') Jactation.

RHIPTOLUSIA, (*ρῆπτω*, 'to throw,' and *λουω*, 'to wash,') Affusion.

RHIS, gen. *Rhi'nis*, (*ῆς*, gen. *ῥινος*), Nasus.

RHIZA, (*ρίζα*), Radix.

RHIZAGRA, (*ρίζαγρα*, from *rhiza*, and *αγρα*, 'seizure,') Punch.

RHIZODONT'TRYPY, *Rhizodontot'rypy*, (*rhiza*, *ὀδὸν*, gen. *ὀδοντος*, 'a tooth,' and *τρύπω*, 'I perforate,') Perforation of the root of a tooth for the discharge of fluid.

RHIZO'MA, *Rhi'zome*, (from *rhiza*.) The part of the root of a plant, which consists of wood or flesh, as in the carrot.

RHIZONYCHIA, (*rhiza*, and *ονυξ*, gen. *ονυχος*, 'a nail,') see Nail.

RHIZOPH'AGUS, (*ρίζοφάγος*, from *rhiza*, and *φαγω*, 'I eat,') One who lives on roots.

RHIZOTOMIST, (*ρίζοτομος*, from *rhiza*, and *τεμνειν*, 'to cut,') Radiseca.

RHIZOTOMUS, Apothecary.

RHOA, (*ρῶα*), *Punica granatum*.

RHODELÆ'ON, *Oleum rosarum*, (*rhodon*, and *ελαιον*, 'oil,') Oil of roses, or oil impregnated with roses.

RHO'DIA, *Rhodi'ola ro'sea, Ro'sea, Rose'wort*, (F.) *Rhodiola*, (*rhodon*, owing to its odour.) *Sex. Syst.* Diccia Octandria. The root, *Roseroot*, when dry, has a very pleasant smell, and is slightly astringent. It is not used.

RHO'DINUS, (*ρῶδινος*), *Rho'dius, Rosa'ceus, Ro'seus, Rosu'tus*, (F.) *Rosat*, Rosy. Formed of, or containing, roses.

RHODIOLA ROSEA, (dim. of *rhodia*), *Rhodia*.

RHODIOLE, *Rhodia*.

RHODITES VINUM, (*ρῶδιτης οἶνος*.) Wine in which roses have been macerated.

RHO'DIUM LIG'NUM, *Rho'dium* or *Rose'wood*, (from *rhodon*.) The wood or root of a tree supposed to be *Genis'ta Canariensis* of Linnaeus. (F.) *Bois de rose, Bois de Chypre*. *Ord.* Leguminosæ. *Sex. Syst.* Diadelphica Decandria. The essential oil is a perfume, and possesses cordial and tonic virtues. Its smell is attractive to fish, rats, &c.

RHODIUS, *Rhodinus*.

RHODODAPHNE, (*ρῶδοδάφνη*), *Nerium oleander*.

RHODODAPHNIS, *Nerium oleander*.

RHODODENDRON CHRYSAN'THEMUM, *R. chrysanthemum seu aur'eum seu officina'lē, Oleander, Rosebay, Yellow rhododendron, Mountain Lau'rel*, (F.) *Rosage, R. chrysanthe, Rose de Sibérie*, (*ρῶδοδένδρον*, from *rhodon*, and *δένδρον*, 'a tree,') *Nat. Ord.* Ericaceæ. *Sex. Syst.* Decandria Monogynia. This plant has been recommended in rheumatism, gout, and syphilis. In an overdose it produces acronarcotic symptoms.

RHODODENDRON MAX'IMUM, *Great Lau'rel, American Rosebay*. An indigenous shrub, *Ord.* Ericaceæ, which flowers in July. It is stimulant and astringent, and has been used by the Indians in gout and rheumatism.

RHODODENDRON PUNCTATUM. Indigenous; has similar properties.

RHODOMELA PALMA'TA, *Dulse, Sac'charine Fu'cus*. One of the Algæ, eaten in large quantities throughout the maritime countries of the north of Europe, and in the Grecian Archipelago.

RHODOMELI, (*rhodon*, and *μελι*, 'honey,') *Mel roseæ*.

RHODON, (*ρῶδον*), *Rosa, Rosa centifolia*.

RHODOSACCHARUM, (*rhodon*, and *σάκχαρον*, 'sugar,') Confectio rosæ gallicæ.

RHODOSTACTON, (*ρῶδοστακτον*), *Mel roseæ*.

RHODOSTAGMA, (*rhodon*, and *σταγμα*, 'that which has been dropped,') *Aqua rosæ*.

RHŒA, (*ρῶα*), *Punica granatum*.

RHŒAS, gen. *Rhœ'adis*, see Papaver rhœas.

RHŒBDESIS, (*ρῶιβδησις*, from *ρῶιβίω*, 'to sup up,') Absorption.

RHŒUM, *Rheum*.

RHOGMUS, (*ρῶγμος*), *Râle*.

RHO'ITES, (*ρῶιτης*), a wine impregnated with the pomegranate, (from *ρῶα*, the pomegranate.) A confection, made by boiling the juice of the pomegranate in honey.

RHOMBOÏDE'US, (*ρῶμβος*, 'a rhombus, whose sides are equal, with two obtuse and two acute angles.) *Rhomboideus major* and *minor, Rhomboides, Cervici-dorso-scapulaire*. A muscle situate at the posterior inferior part of the neck, and at the posterior part of the back. It has a rhomboidal shape, and is attached, by its inner edge, to the posterior cervical ligament, and to the spinous processes of the last cervical vertebrae, and of the first four or five dorsal. By its outer edge, it is attached to the spinal or posterior edge of the scapula. It is divided into two fasciculi, constituting the *Rhomboideus major* and *Rh. minor*, the latter being situate higher than the other. This muscle draws the scapula backwards and upwards, and impresses upon it a rotatory motion, which brings its inferior angle nearer to the spine; and, consequently, depresses the anterior angle and the shoulder.

RHOMBOS, (*ρῶμβος*.) A bandage mentioned

by Galen; so called on account of its rhomboidal shape.

RHONCHAL, *Rhoncha'lis*, (*rhonchus*, 'rattle') Relating or appertaining to rhonchus;—as '*rhonchal fluctuation*;' the fremitus that occurs through the influence of respiration as an attendant on cavernous rhonchi.

RHONCHUS, (*ῥῶνχος*) Rale, Snoring, Stertor—*r.* Cavernous, see Gurgling—*r.* Cavernulous, see Gurgling.

RHONCHUS, *Click'ing*. A rhonchus which has been considered positively indicative of the existence of tubercles in the lungs. [?]

RHONCHUS CREPITANS, *Râle crépissant*—*r.* Crepitans *redux*, see Crepitant—*r.* Mucous, see *Râle muqueux*—*r.* Sibilans, see *Râle crépissant*—*r.* Sibilus, *Râle sibilant*—*r.* Sonorus, *Râle sonore*—*r.* Subcrepitant, see *Râle muqueux*.

RHOPALISMUS, (*ῥοπαλίσμος*) Priapismus.

RHOPALOSIS, (*ῥοπαλῶσις*, 'a club,') The hair being matted like a club, and *osis*, 'Plica.

RHOPALUM, (*ῥοπαλόν*) Penis.

RHOPE, (*ῥοπή*, 'a moving forwards or downwards,') Congestion.

RHOTACISMUS, (*ῥωτακισμός*) Rotacism.

RHÖX, (*ῥῶξ*, gen. *ῥωγος*, 'a fissure,') Pupil, Uvea.

RHUBARB, *Rheum*—*r.* Chinese, see *Rheum*—*r.* French, *Rheum Rhaiponticum*—*r.* India, see *Rheum*—*r.* Crimea, *Rheum Rhaiponticum*—*r.* Monk's, *Rumex Alpinus*, *Rumex patientia*—*r.* Pie, *Rheum Rhaiponticum*—*r.* Poor Man's, *Thalictrum*—*r.* Rhaipontic, *Rheum Rhaiponticum*—*r.* Russia, see *Rheum*—*r.* Turkey, see *Rheum*—*r.* Wild, *Convolvulus panduratus*.

RHUBARBE, *Rheum*—*r.* *Fausse*, *Thalictrum*—*r.* *des Indes*, *Convolvulus mechoan*—*r.* *des Poures*, *Thalictrum*—*r.* *Savage*, *Rumex patientia*.

RHŪE, *Ruta*.

RHUM, *Rau*.

RHUMA, in composition, see *Rheuma*.

RHUMAPYRA, *Rheumatism*, acute.

RHUMATALGIA, *Rheumatism*, chronic.

RHUMATALGIE, *Rheumatism*, chronic.

RHUMATIQUE, *Rheumatic*.

RHUMATISANT, *Rheumatic*.

RHUMATISMAL, *Rheumatic*.

RHUMATISME, *Rheumatism*—*r.* *Aigu*, *Rheumatism*, acute—*r.* *Articulaire aigu*, *Rheumatism*, acute—*r.* *Chronique*, *Rheumatism*, chronic—*r.* *Præabdominale*, *Rheumatism*, præabdominal—*r.* *d. l'Utérus*, *Metrorrhœma*.

RHUME, *Rheum*—*r.* *de Cerveau*, *Coryza*—*r.* *de Poitrine*, *Catarrh*.

RHUS BELGICA, (gen. *Rhois*, *ῥῶς*) *Myrica gale*—*r.* *Copallina*, see *Rhus typhina*.

RHUS CORIARIA, *Adurion*, *Sumach*, *Shumach*, *Elm-leaved Sumach*, (F.) *Sumac*, *Sumac des Corroyeurs*, *Roure* ou *Rouvre*. *Ord.* *Anacardiaceæ*. *Ser. Syst.* *Pentandria Monogynia*. This is the only species of the genus *rhus* that is innocent. Both the leaves and berries have been used as astringents and tonics.

RHUS GLABRA, '*Smooth Sumach*.' Indigenous and official in the Pharmacopœia of the United States. An infusion of the fruit—*Rhus glabrum* [?] Ph. U. S.—which is acid and astringent, has been used as a cooling drink, and as a gargle. An infusion of the inner bark of the root has been recommended in mercurial ptyalism.

RHUS RADICANS, *R. vernix*, *Poison vine* (*America*). This plant is poisonous (see Poisons, Table of), and is extremely acrid, when applied to the skin, producing erysipelas and vesications, and has been used, like *Rh. toxicodendron*, in paralysis, and other nervous affections.

RHUS SUCCEDANEA, see *Wax*, *Japan*.

RHUS SYLVESTRIS, *Myrica gale*.

RHUS TOXICODENDRON, *Toxicodendron*, *Poison*

Oak (*America*), (F.) *Sumac venenose*. The leaves—*Toxicodendron* Ph. U. S.—have been used in paralytic affections and eruptions. It is poisonous, like *Rh. radicans*. Dose, gr. ss. to gr. ʒ.

RHUS TYPHINA, *Stag's horn Sumach*, and **RHUS COPALLINA**, *Mountain Sumach*, have similar properties with *Rhus glabra*.

RHUS VENENATA, *Poison Sumach*, and **RHUS PRUMILA**, have similar properties with *Rhus toxicodendron*.

RHUS VERNIX, *Rh. radicans*.

RHYAS, (*ῥῡας*, from *ῥέω*, 'I flow,') *Rhy'sis*. An ulceration of the lachrymal caruncle with a constant flow of tears.

RHYMA, (*ῥῡμα*) Medicament.

RHYPARIA, (*ῥῡπαρία*, from *ῥυπος*, 'dirt,') *Rupia*. *Sortes primarium viarum*.

RHYPIA, (*ῥῡπία*) *Rupia*.

RHYPUS, (*ῥῡπος*) *Sortes*.

RHYPTICA, (from *ῥῡπτω*,) *Detergents*.

RHYPTICUS, *Abstergeut*.

RHYSIS, (*ῥῡσις*) *Flux*, *Rhyas*.

RHYSEMATA, gen. *Rhysematis*, (*ῥῡσμητα*) *Wrinkle*.

RHYTHM, *Rhyth'mus*, (*ῥῡθμός*,) (F.) *Rhythme*, 'measured movement.' The order of proportion, as regards time, which reigns between the different movements of an organ, or of the organism in health and disease. Thus, rhythm is applied to the diurnal variations of the pulse; and to the paroxysmal movements of an intermittent.

RHYTHMICAL, (*ῥῡθμικός*) *Periodical*.

RHYTHMICAL CONTRACTIONS OF MUSCLES are those that succeed one another after regular intervals.

RHYTHMUS, (*ῥῡθμός*) *Rhythm*.

RHYTIDOSIS, (*ῥῡτιδῶσις*) *Rutidosis*.

RHYTIS, (*ῥῡτις*) *Wrinkle*.

RIB, ([Sax.] *rib*,) *Costa*—*r.* *False*, see *Costa*—*r.* *Floating*, see *Costa*—*r.* *Sternal*, see *Costa*—*r.* *True*, see *Costa*.

RIBES (of Arab. origin) **GLOSSULARIA**, *R. uva crispa*.

RIBES NIGRUM, *R. al'idum*, *Glossula'riani'gra*, *Cassia*, *Black Currant*, *Gazel*, (F.) *Groseillier noir*. *Ord.* *Grossulacææ*. *Ser. Syst.* *Pentandria Monogynia*. The berries—*Black currants*—have been recommended in sore-throat, and as diuretics. The leaves have been advised for the same purpose. They are chiefly used as preserves.

RIBES OFFICINALE, *R. rubrum*—*r.* *Odium*, *R. nigrum*.

RIBES RUBRUM, *R. officinale seu vulgare*, *Glossula'ria non spinosa seu rubra*, *Red currant*, (F.) *Groseillier rouge*; of which the *Whit* is only a variety. It is cultivated, and affords an agreeable fruit—*Red currants*, (F.) *Groseilles*. It possesses the qualities of the subacid fruits in general, and makes an excellent preserve.

RIBES UVA CRISPA, *R. grossula'ria*, *Common gooseberry*. The fruit—*Grossula'ria*, (F.) *Groseille à maquereau*, *Gooseberry*—is agreeable, and makes a good preserve.

RIBES VULGARE, *R. rubrum*.

RIBGRASS, *Plantago lanceolata*.

RIBS, NECK OF THE, *Collum costarum*—*r.* *Short*, see *Costa*.

RIBWORT, *Plantago Virginica*.

RICE, *Oryza*—*r.* *Canada*, *Zizania aquatica*.

RICE DISEASE, *Cholera*.

RICE, GROUND, see *Oryza*.

RICE JELLY, see *Oryza*.

RICE WATER, see *Oryza*. The evacuations of cholera have been termed—from their resemblance to ricewater—'*ricewater discharges*;' and in India, '*congee discharges*;'—*congee*, in Hindoostanee, meaning '*ricewater*.'

RICE, WILD, *Zizania aquatica*.

RICHARDSONIA, (after Richardson, an early English botanist,) see *Ipecacuanha*.

RICHLÉAF, *Collinsonia Canadensis*.

RICHWEED, *Actæa racemosa*, *Ambrosia trifida*, *Collinsonia Canadensis*, *Pilea pumila*.

RICIN, *Ricinus communis*.

RICINOIDES, (*ricinus*, and *aidos*, 'resemblance,') *Jatropha curcas*.

RICINUS COMMUNIS, (from *ricinus*, 'a tick,' which its seed resembles,') *Ricinus vulgaris* seu *chistidis*, *Palma Christi*, *Cataputia major*, *Faba purgatrix*, *Cerva*, *Kerva*, *Cici*, *Castor oil plant*, *F. Ricin*, *Fève purgatif*. Ord. Euphorbiaceæ. See *Syst. Monœcia Monadelphia*. Castor-oil seeds, whence the oil is obtained, are inodorous; taste acrid and slightly sweetish. One or two seeds—*Semina Cataputia majoris* seu *Ricinini vulgaris*—will act as a cathartic; but the oil is alone used. It is obtained by bruising the castor-seeds, previously decorticated, and then expressing the oil without the application of heat. Recently-drawn castor oil, *Oleum de kerva* seu *kervi-num*, *Alkerba*, *O. Ricinini* seu *Cicinum* seu *Palma liquidum* ou de *Palma Christi*, *Carapat*, *Karapat*,—in the United States often called simply Oil, (F.) *Huile de Ricin*—is inodorous and nearly insipid. It is cathartic, operating speedily, and may be used in all cases where powerfully stimulating purgatives would be injurious. Dose, fʒj to fʒiiss. Hufeland recommends that a drop of croton oil shall be added to an ounce of the oil of poppy, and that the mixture shall be called *Oleum Ricinini Officinale*.

At the Cape of Good Hope, the oil is obtained from *Ricinus lividus*.

A variety of *Ricinus* at Cape Verd is employed by the inhabitants to increase the secretion of milk. It is applied in the form of poultice, made with the green leaves to the mammae, or of a strong decoction, with which the mammae and the external parts of generation are washed.

RICINUS LIVIDUS, see *Ricinus communis*—r. Major, *Jatropha curcas*—r. Minor, *Jatropha manihot*—r. Viride, *R. communis*—r. *Vulgaris*, *R. communis*.

RICK, *Malleolus*.

RICKETS, (see *Rig*.) *Rachitis*.

RICKETY, *Rachiticus*.

RICTUS, (from *ringor*, *rietus*, 'to open the mouth wide,') Yawning—r. *Lupinus*, (*lupus*, 'a wolf,') see *Harelip*.

RIDE, *Wrinkle*.

RIDE', *Wrinkled*.

RIDING (of Bones). *Os sium superpositio*, (F.) *Chevauchement*. Displacement of the fragments of a bone, chiefly produced by the contraction of muscles, which occasions shortening of the limb; the fractured extremities riding over each other instead of being end to end.

RIDING HAG, *Incubus*.

RIEUR DE SANTORINI, *Risorius novus*.

RIFE, ([Sax. *rýpe*), (Prov.) *Brief*. Prevailing: as applied to diseases.

RIFT, (from *rive*, *rived*), *Erection*.

RIG, ([Sax.] *rycg* or *hyge*), *Dorsum*.

RIGIDITAS, *Rigidity*.

RIGIDITAS ARTICULORUM, *Ancylosis spuria*. Stiffness of the joints. False ancylosis.

RIGIDITÉ, *Rigidity*—r. *Cadavérique*, see *Rigidity*.

RIGIDITY, *Rigiditas*, *Rigor*, (*rigere*, *rycw*, 'to stiffen,') (F.) *Rigidité*, *Roideur*, *Raideur*. Great stiffness of fibre, or want of suppleness. The stiffness of the dead body, *Cadavérique rigidity*, (F.) *Roideur* ou *Rigidité cadavérique*, is one of the signs of the cessation of life. It may be removed, however, for a time, by the injection into the arteries of oxygenated defibrinated blood. See *Rigor mortis*.

RIGOR, *Rhigus*, *Algor*, *Coldness*, *Cheimia*, (F.) *Frisson*, (from *rycw*, *rygo*, 'to stiffen,' espe-

cially from cold.) *Rigidity*. Sensation of cold, with involuntary shivering or shaking of the whole body:—a symptom of fever. A slighter degree, and at times full rigor, is occasionally termed a chill. (Prov.) *crema*, *frigus tenué*, *pericrigatio*.

RIGOR ARTUUM, *Contractura*—r. *Cadaverosus*, *Rigor mortis*—r. *Corporis mortui*, *Rigor mortis*—r. *Emortualis*, *Rigor mortis*—r. *Maxillæ inferioris*, *Trismus*.

RIGOR MOR'TIS, *R. corporis mortui* seu *cadaverosus* seu *emortualis*, *Necronécéma*, (F.) *Roideur* ou *Raideur* ou *Rigidité cadavérique*. 'Stiffness of death.' Death stiffening. The rigidity of limbs that occurs on dissolution. See *Rigidity*.

RIGOR NERVORUM, *Tetanus*—r. *Nervosus*, *Tetanus*.

RIMA, *Schisma*, *Schiza*, 'a fissure, a cleft,' as *Rima* seu *Rimula Glottidis* seu *Laryngis*, the opening of the glottis. *Rima* seu *R. magnæ vulvæ*, *Introitus vaginæ*, *Sinus vulvæ*, *Scapha*, *Scaphus* seu *Fissura magna* seu *Scissura* seu *Schiza vulvæ*; the opening of the vulva. See *Vulva*. *Rima oris*, the opening of the mouth.

RIMA CANALIS ORBITARII, *Suborbital fissure*—r. *Clunium*, see *Nates*—r. *Glottidis*, see *Rima*—r. ad *Infundibulum*, *Vulva cerebri*—r. *Laryngis*, *Rima glottidis*—r. *Magna vulvæ*, see *Rima*—r. *Oris*, see *Rima*—r. *Transversa magna cerebri*, *Fissure*, *transverse*, of the brain—r. *Vulvæ*, see *Rima*.

RIMÆ, (pl. of *rima*), *Rhagades*.

RIMULA, (dim. of *rima*.) A small fissure or cleft.

RIMULA GLOTTIDIS, see *Rima*—r. *Laryngis*, *Rima glottidis*.

RINÆUS, (from *rhin*), *Compressor naris*.

RING, ([Sax.] *hring*, *ring*), *Circos*, (F.) *Anneau*. A name given to natural circular, or roundish apertures, with muscular or aponeurotic parietes, which serve for the passage of some vessel or part:—as the umbilical ring, inguinal ring, &c.

RINGS, CRAMP, see *Cramp rings*—r. *Festooned*, *Annuli fibrocartilaginei*—r. *Fibrocartilaginous*, *Annuli fibrocartilaginei*.

RINGFINGER, *Annular finger*.

RINGMAN, *Annular finger*.

RINGWORM, *Herpes circinatus*—r. of the Scalp, *Porriro seutulata*—r. *Ulcerative*, *Herpes exedens*—r. *Vesicular*, *Herpes præputii*.

RIO VINAIGRO, *Puñambio*. A stream of some size in New Granada, which is strongly impregnated with sulphuric acid.

RIRE ou RIS, *Risus*—r. *Canin*, *Canine laugh*—r. *Mogueur*, *Canine laugh*—r. *Sardonien*, *Canine laugh*, *Risus sardonius*—r. *Sardonique*, *Canine laugh*.

RISAGON, *Cassumuniar*.

RISIGALLUM, *Orpiment*.

RISING, *Abscess*, see *Regurgitation*—r. of the Lights, *Cynanche trachealis*.

RISING or ELEVATION OF THE PULSE, *Pulsus elevatio*. The pulse is said to rise, when it becomes more full and frequent, as occurs in the exacerbation of acute diseases.

RISORIOUS NOVUS, (from *ridere*, *risum*, 'to laugh,') *R. Santorini*, (F.) *Rieur de Santorini*, *Santorini* gives this name to a portion of the platysma myoides, which passes from the cheek towards the commissure of the lips.

RISSIGALLUM, *Orpiment*.

RISUS, (*ridere*, *risum*, 'to laugh,') *Gelus*, *Laughing*, *Laughing*, *Laugh*, (F.) *Rire* ou *Ris*. An involuntary movement of the muscles of the face, and of the lips in particular, accompanied with a sonorous and interrupted expiration, commonly indicating mirth and satisfaction.

RISUS APIUM, *Ranunculus sceleratus*—r. *Caninus*, *Canine laugh*—r. *Involuntarius*, *Canine*

as regards its physical qualities, is considered to

They are not used for any other purposes except for the distillation of rose water.

Oil of Roses, Oleum Rosæ, Attar, Otto or Essence of Roses, and the formation of a syrup.

ROSA DAMASCENA, *R. centifolia*.

ROSA GAL'LICA, *R. Austri'aca seu rubra*, *Red Rose*, *F. Rosier du Provens.* The petals of this variety are astringent; and are used in infusion, conserve, &c.

ROSA LEUCANTHA, *R. alba*—*r. Pallida*, *R. centifolia*—*r. Rubra*, *R. Gallica*—*r. Rubra vulgaris*, *R. canina*—*r. Saltaus*, *Essera*—*r. Senticosa*, *R. canina*—*r. Swartziana*, *R. canina*—*r. Sylvestris*, *R. canina*—*r. Usitatissima*, *R. alba*.

ROSACEA, *Roseola*.

ROSACEUS, (from *rosa*.) *Rhodinus*.

ROSAGE, *Nerium oleander*, *Rhododendron chrysanthemum*.

ROSAGO, gen. *Rosag'inis*, *Nerium oleander*.

ROSALIA, (from *rosa*, 'the rose,' from its colour.) *Scarlatina*.

ROSALIE, *Scarlatina*.

ROSAT, (from *rosa*, 'the rose,' from its colour.) *Rhodinus*.

ROSATUS, *Rhodinus*.

ROSE, *Erysipelas*, (from its colour,) *Rosa*—*r. Cabbage*, *Rosa centifolia*—*r. Campion*, *Lychnis githago*—*r. Christmas*, *Helleborus niger*—*r. Copper*, *Papaver rhoeas*—*r. Corn*, *Papaver rhoeas*—*r. Damask*, *Rosa centifolia*—*r. Dog*, *Rosa canina*—*r. Dog*, conserve or confection of, *Confectio rosæ caninæ*.

ROSE DROP, *Gutta rosæ*—*r. Du Chien*, *Rosa canina*.

ROSE, HUNDREDLEAVED, *Rosa centifolia*—*r. Marine*, *Rosmarinus*—*r. Pale*, *Rosa centifolia*.

ROSE RASH, *Roseola*.

ROSE, RED, *Rosa Gallica*—*r. Rock*, *Helianthemum Corymbosum*.

ROSE DE SIBÉRIE, *Rhododendron chrysanthemum*.

ROSE SPOTS, *Spots*, red.

ROSE, WHITE, *Rosa alba*.

ROSEA, *Rhodia*.

ROSEBALM, SCARLET, *Monarda coccinea*.

ROSEBAY, AMERICAN, *Rhododendron maximum*.

ROSÉE DU SOLEIL, 'sundew,' *Drosera rotundifolia*.

ROSELLINA, (from *rosa*, 'the rose,' on account of its colour.) *Roseola*.

ROSEMARY, *Rosmarinus*—*r. Marsh*, *Statice limonium*, and *S. Caroliniana*—*r. Wild*, *Eriocephalus umbellatus*.

ROSENHEIM, MIN'ERAL WATERS OF. *Rosenheim* is 34 miles S.E. from Munich, in Bavaria. The waters contain sulphuretted hydrogen, carbonic acid, carbonate of lime, chlorides of calcium and magnesium, carbonate of soda, and oxide of iron. They are used as tonics.

ROSENMÜLLER, ORGAN OF, *Parovarium*.

ROSEOLA, *Eczanth'e'is Rose'ola*, *Rose Rash*, (*F.*) *Rosule*, *Éruption anmale*, *Rosace*, *Fievre rouge*, *Efflorescence érysipélateuse*, (from *rosa*, 'the rose,' on account of the colour.) An efflorescence in blushing patches, gradually deepening to a rose-colour, mostly circular or oval; often alternately fading and reviving; sometimes, with a colourless nucleus; and chiefly on the cheeks, neck, or arms. It is frequently found, as a symptom, in dentition, dyspepsia, rheumatism, &c., and is of no consequence. See *Roseola*, *Rubeola*, and *Scarlatina*.

ROSEOLA ACNOSA, *Gutta rosæ*.

ROSEOLA ÆSTI'VA. The affection which prevails in summer. The eruption is of a darker colour, and is attended with more itching.

ROSEOLA ANNULA'TA. Characterized by patches in the form of rosecoloured rays, with central areas of the colour of the skin.

ROSEOLA FICOSA, *Sycosis*—*r. Typhosa*, *Spots*, red—*r. Variolosa*, *Roseola*.

ROSE'OLÆ, *Rose'ola*, *Rosell'ina*, *Rube'olaspuria*, *False or French measles*, (*F.*) *Rosule*, *Fausse varigole*. This name is given by some to an acute exanthem, midway between measles and scarlatina, and which belongs to neither one nor the other, as it affords no protection against either. It is, also, termed *Rube'ola* by some, and *Rube'oloid*, a term, which is applicable to any eruption resembling *rubeola*. An eruption which often precedes that of smallpox is termed *Roseola Variolosa*.

ROSÉOLE, *Roseola*, *Roseolæ*.

ROSE'PINK, *Chironia angularis*.

RO'SES, MILK OF. A cosmetic wash, formed, according to one method, of *Sulphate of potash*, gr. vj; *Oil of almonds*, ʒi; *Essence of bergamot*, ʒij; *Rose water*, ʒijj; *Orange flower water*, ʒij. Mix.

ROSEUS, (from *rosa*.) *Rhodinus*.

ROSEWOOD, *Rhodium lignum*.

ROSEWORT, *Rhodia*.

ROSIER BLANC, *Rosa alba*—*r. à Cent Feuilles*, *R. centifolia*—*r. de Provens*, *Rosa Gallica*.

ROSIN, BLACK, (from *resina*.) *Colephonia*—*r. Brown*, *Colephonia*.

ROSIN WEEB, *Silphium laciniatum*.

ROSIN, WHITE, see *Pinus sylvestris*—*r. Yellow*, see *Pinus sylvestris*.

ROSIO STOMACHI, (from *rodere*, *rosium*, 'to corrode,') *Cardialgia*—*r. Ventriculi*, *Cardialgia*.

ROSMARINUM, *Rosmarinus*.

ROSMARI'NUS, (*ros*, 'dew,' and *marinus*, 'marine,') *R. officinalis* seu *latifolius* seu *hortensis*, *Atchachi*, *Libanotis coronaria*, *Dendrolibanus*, *Rose'mary*, (*F.*) *Rosmarin*, *Rose marine*, *Encensier*; *Ord. Labiate*. *Sez. Syst.* *Diandra Monogynia*. The tops of this plant, *Rosmarin'us* (*Ph. U. S.*), *Herba Anthos*, have a fragrant, grateful smell; and an aromatic, warm, bitterish taste, which is dependent upon an essential oil, combined with camphor. Rosemary has been recommended in nervous headaches, in the form of infusion. It is now rarely used, except for its odour. The flower is called *Anthos*, *αἶθος*. Its oil—*O'leum Rosmarin'i*—is official. From two to six or ten drops are given in nervous complaints, rubbed up with sugar.

ROSMARINUS HORTENSIS, *Rosmarinus*—*r. Latifolius*, *Rosmarinus*—*r. Officinalis*, *Rosmarinus*—*r. Stoechadis facie*, *Teucrium creticum*—*r. Sylvestris*, *Ledum sylvestre*.

ROSSALIA, (from *rosa*, 'the rose,' from its colour.) *Scarlatina*—*r. Squamosa*, *Scarlatina*.

ROSSIGNOL, (pron. *Rosiniol*.) A name given, in France, to an affection to which pressmen are subject. It is a sprain, followed by tumefaction of the radiocarpal joint.

ROSSOLIO, see *Spirit*.

ROS'TRUM, *Ram'phos*, 'a beak.' (*F.*) *Bec*. A name given to several old forceps, on account of their resemblance to the beaks of different birds. The principal were the *Rostrum corvini*, *anati'*, *psittaci*, *vultur'*, *cygni* seu *alut'*, *grui*, *lucerti*, &c.

ROSTRUM EXTERNUM, *Olecranon*—*r. Percinum*, *Aeromion*—*r. Posterius*, *Olecranon*—*r. Sphenoidale*, *Azygos processus*.

ROSY BETTY, *Erigeron bellidifolius*—*r. Bush*, *Spiræa tomentosa*—*r. Drop*, *Gutta rosæ*—*r. Wheel*, *Gutta rosæ*.

ROT, (from [*L.*] *ructus*), *Eruetation*.

ROT, GRINDERS', *Asthma*, *grinders'*.

ROTACISM, *Rotacism'us*, *Rottacism'us*, properly *Rhotacism'us*, *Psellis'mus rhotacism'us*, (*F.*) *Grasseyement*. A vicious pronunciation of the Greek, *ρ*, *Rau*, common in the northern parts of England; especially near Newcastle. Called, there, the *Burr* or *Burth*.

ROTACISMUS, (*ῥωτακισμός*.) *Grasseyement*, *Rotacism*.

ROTATIO, (*rotare, rotatum*, 'to turn round,') Diarthrosis.

ROTA'TOR, (*rotare, rotatum, (rota, 'a wheel,')* 'to turn round,') (F.) *Rotateur*. A name given to several muscles that turn the parts to which they are attached upon their axes;—such as the oblique muscles of the eyes, the pronators, supinators, &c.

ROTATOR, Trochanter.

ROTATOR'ES DORS'I. Eleven small muscles on each side of the dorsal vertebræ, which arise from the point or upper edge of each transverse process, and pass to the lower edge of the arch of the vertebra above, as far as the base of the transverse process. Their action is to rotate the individual vertebræ on each other.

ROTATO'ES FEMORIS, Obturatores—r. Oculi, Oblique muscles of the eye.

ROTCOLL, Cochlearia armoracia.

ROTTACISMUS, Rotacism.

ROTTLE'RA SCHIMPERI, (after Dr. Rottler, a Danish missionary.) A large tree of Abyssinia, *Nat. Ord.* Euphorbiaceæ, the bark of which, called *Cortex muse'ne*, and formerly known as *Cortex Brayeræ anthelmin'ticæ*, mixed with koosso, is used in that country against tapeworm.

ROTTLE'RA TINCTO'RIA. A euphorbiaceous plant of India. A powder is attached to the fruit capsule of this plant, which is brushed off and collected when the fruit is dry. It is called *Kameyla*, *Rottle'ra*, (Ph. U. S., 1863,) or *Kameela*, *Kame'la* (Ph. B. 1864); is used in Northern India as a vermifuge for dogs; and has proved effectual against tapeworm in man. The dose is two or three drachms, which purges.

ROTULA, (dim. of *rota*, 'a wheel,') Patella, Trochiscus—r. Genu, Patella.

ROTULAD, see Rotular aspect.

ROT'ULAR, (from *rotula*.) Relating or appertaining to the patella.

ROTULAR AS'PECT. An aspect towards the side on which the rotula is situated.—(Barclay.) *Rot'ulad* is used by the same writer, adverbially, to signify 'towards the rotular aspect.'

ROTULE, Patella.

ROTU'LIAN, *Rotulianus*, (F.) *Rotulien*. Relating or belonging to the rotula or patella.

RÖTUNDUS, (from *rota*, 'a wheel,') Round.

ROUCOU, Terra Orleansa.

ROU'EN, MIN'ERAL WATERS OF. This water, known under the name, *Eau des fontaines de la Marégnerie*, is cold, and contains carbonate of iron, carbonic acid, carbonate of lime, chloride of calcium, &c. It is tonic.

ROUGE, 'red,' (from [L.] *rubeo*, 'to be red,') see Colcothar—r. d'Angleterre, Colcothar—r. de Prusse, Colcothar—r. du Seng, Hæmatin.

ROUGEOLE, (from [F.] *rouge*, 'red,') Rubella—r. Fausse, Roselæ.

ROUGEURS, (from [F.] *rouge*, 'red,') Flush.

ROUGH ROOT, Liatris.

ROUILLE, (originally from *rubigo*, 'rust,') Rubiginous.

ROULEMENT DU LAMBEAU, 'rolling of the flap,' see Flap.

ROUND, *Te'rea*, *Rotun'dus*, (F.) *Rond*, (from [L.] *rotundus*.) Anatomists have given this name to many organs whose fibres are collected in round fasciculi. See Teres.

ROUND FORA'MEN, *Fora'men rotund'um*. A foramen of the sphenoid bone, a little behind the *foramen-lacerum superius*, which gives passage to the second branch of the fifth pair of nerves, called *superior maxillary*.

ROUND HEART, *Thaspium atropurpureum*.

ROUND LIG'AMENTS OF THE U'TERUS, *Ligamen'ta rotund'a U'teri*, (F.) *Cordons sus-pubiens* (Ch.), *Cordons vasculaires*, *Ligaments ronds de l'utérus*. Two cords, which arise from the lateral and supe-

rior parts of the uterus, whence they proceed towards the abdominal rings, which they pass through, and terminate by vanishing in the areolar tissue of the groins, mons veneris, and labia majora. These cords are whitish, flattened, and narrower in the middle than at the extremities. According to Mr. Rainey, they are composed chiefly of the striped muscular fibre.

ROUTINE, *Routine*, Sorbus aucuparia.

ROUP, (like croup, by onomatopœia [?]), Cyranche trachealis, Raucedo.

ROU'E, Rhus coriaria.

ROUTINE PRACTITIONER, Routinist.

ROUTINIST, pronounced *routen'ist*; (F.) *Routinier*; ([F.] *routine*, 'a regular habit or practice, unvaried by circumstances,' from *route*, 'a road,') One who practises in such an unvaried manner:—a *routine practitioner*.

ROUVRE, Rhus coriaria.

ROVE, (from *rive*,) Scab.

ROVING, 'wandering,' Delirium.

ROWANTREE, Sorbus aucuparia.

ROWLAND'S SPRINGS, see Georgia, mineral waters of.

ROWTING, (from [Sax.] *hrutan*, 'to snore,') Snoring.

RUBBER, INDIAN, Caoutchouc.

RUBBING, ([G.] *reiben*, 'to rub,') Friction—r. Dry, see Friction—r. Moist, see Friction.

RUBBING SOUND, *Bruit de Frottement*.

RUBE'DO, gen. *Rube'dinis*, *Ru'dor*, 'redness,' (from *rubere*, 'to be red,') A diffused, but not spotted, redness on any part of the skin; like that which arises from blushing. Blushing.

RUBEFA'CIENT, *Erethis'ma*, *Erethis'ticum*, *Rubefa'ciens*, (ruber, 'red,' and *facio*, 'I make,') (F.) *Rubéfiant*. That which produces redness. A medicine which causes redness of the skin. The action is called *Rubefaction*.

RUBÉFIANT, Rubefacient.

RUBE'OLA, *Rubeola vulgar'is*, (from *rubeo*, 'I am red,') *Rose'ola*, *Exanth'e'xis*, *Rube'ola*, *Vetera'na*, *Blac'cia*, *Fer'se*, *Blac'tia*, *Bovill'e*, *Morbil'li*, *Scarlati'na pustulo'sa* seu *mil'ia'ris*, *Fe'bris morbil'lo'sa*, *Ty'phus morbil'lo'sus*, *Phenicis'mus*, *Ephy'mata*, *Measles*, (Sc.) *Mirles*, (Prov.) *Creas*, *Creds*, *Malls*, *Maules*, *Meas'lings*, (F.) *Rougeole*, *Fie'vre morbillense*. One of the major exanthemata; generally affecting individuals but once, and produced by specific contagion. The rash usually appears on the fourth, but sometimes on the third, fifth, or sixth day of a febrile disorder; and, after a continuance of four days, gradually declines with the fever. The disease generally commences from ten to fourteen days after the contagion has been received. The eruption first shows itself in distinct, red, and nearly circular spots, somewhat less than the ordinary areolæ of flea-bites. As these increase in number, they coalesce; forming small patches of an irregular figure, but approaching nearest to that of semicircles or crescents. These patches are intermixed with single, circular dots and with interstices of the natural colour of the skin. On the face they are slightly raised, so as to give the sensation of inequality of surface to the finger passed over the cuticle. The disappearance of the eruption is followed by desquamation of the cuticle. Measles is not dangerous of itself; but it is liable to induce pneumonia in winter, and dysentery in summer, which are, at times, very fatal. These are apt to come on at the time of, or soon after, the disappearance of the eruption. When they supervene, they must be treated as idiopathic affections. It demands a general antiphlogistic treatment. Willan has pointed out a kind of RUBEOLA SP'URIA, which he calls RUBEOLA SINE CATARRHO. In this the rash runs its regular course, with little fever or catarrhal affection; affording no certain security against the

common or regular disease. He also gives the name. RUBEOLA NIGRA or BLACK MEASLES, to an unusual appearance of the measles about the seventh or eighth day, when the rash becomes suddenly livid, with a mixture of yellow. It is devoid of inconvenience or danger; and is removed in a week or ten days by the mineral acids and tonics.

Epidemic measles has occasionally prevailed largely in the military service, and has been called *Camp Measles*.

RUBEOLA SPURIA, Roseolæ.

RUBEOLE, Roseolæ.

RUBEOLOID, (*rubeola*, and *oid*.) see Roseola.

RUBIA, (from *ruber*, 'red,') *R. tinctorum* seu *peregrina* seu *sykes'ia* seu *ma'jor*, *Erythrodanum*, *Ra'dix rubra*, *Dyer's Mad'der*, *Mad'der*, (F.) *Garance*, Family, Rubiaceæ. *Sex. Syst.* Tetrandria Monogynia. The root—*Ru'bia*, (Ph. U. S.)—has a bitterish, somewhat austere, taste, and a slight, not agreeable, smell. It was formerly considered deobstruent, detergent, and diuretic.

RUBIG'INOUS, *Rubigino'sus*, (from *rubigo*, gen. *rubig'inis*, 'rust,') (F.) *Rouillé*, Rusty. Having the colour of rust. An epithet given to the sputa in certain cases of pneumonia.

RUBINUS VERUS, (from *ruber*, 'red,') Anthrax.

RUBOR, 'redness,' Rubedo.

RUBRICA FABRILIS, (from *ruber*, 'red,') A red, heavy, earthy matter, formerly employed for making drying plasters; *Hard Ruddle*, *Red Chalk*.

RUBRICA SINOP'ICA. A heavy, compact, reddish earth from Sinope: formerly used in diarrhœa.

RUBRIN, (from *ruber*, 'red,') Hæmatin.

RUBUS, (from *ruber*, 'red,') The pharmacopœial name (Ph. U. S. 1863) of the root of *Rubus Canadensis* and *Rubus villosus*.

RUBUS AFFINIS, *R. fruticosus*.

RUBUS ARCTICUS, *Shrub'by Strawberry*, *Bac'cæ Norland'icæ*. Ord. Rosacæ. *Sex. Syst.* Icosandria Polygynia. The fruit is recommended by Linnæus as possessing antiseptic, refrigerant, and antiscorbutic qualities.

RUBUS BATES, *R. cæsius*.

RUBUS CÆSIUS, *R. Ba'tus*, *Doe'berry plant*, (F.) *Ronce bleue*. The fruit resembles the blackberry in appearance and properties. The bark of the root of *R. Tricla'lis* seu *Canaden'sis* seu *procumbens* seu *flagella'ris*, *American Dewberry*, *Low Blackberry*, is astringent.

RUBUS CANADENSIS, see *R. cæsius*.

RUBUS CHAMÆMO'RUS, *Cloud'berry Tree*, *Chamæmorus*, *C. Norve'gica*, *Chamæ'drys*, *Cloud'berries*, *Knolberries*, (F.) *Ronce de Montagne*. The ripe fruit is sometimes prepared as a jam, and used in fevers, &c., to allay thirst. It has been much extolled as an antiscorbutic.

RUBUS FLAGELLARIS, see *R. cæsius*—*r. Frambœianus*, *R. idæus*—*r. Frondosus*, *R. fruticosus*.

RUBUS FRUTICOSUS, *R. aff'nis* seu *paniculatus* seu *n'ger* seu *frondosus* seu *vilgus*, *Com'mon Bram'ble*, which affords *Black'berries*, *Bram'bleberries*, (Prov.) *Bun'blekites*, (F.) *Ronce noire*. The berries are eaten as a summer fruit, of which they form a poor variety. The bark of the root of the *Rubus villosus*, *American Blackberry*, *High or standing Blackberry*, *Hairy American Bramble*, is astringent, and has been used in the last stages of dysentery, and in cholera infantum.

RUBUS IDEUS, *R. frambœianus*, *Ba'tus*; *Rasp'berry*, (F.) *Framboise*, *Ronce*. The fruit of 'the best varieties—*Rasps*, (Prov.) *Hindberries*—is extremely agreeable; and is much used. It was, formerly, called *Buttison moron*, (*Барвиш морон*).

RUBUS NIGER, *R. fruticosus*—*r. Occidentalis*, see *R. strigosus*—*r. paniculatus*, *R. fruticosus*—*r. procumbens*, see *R. cæsius*.

RUBUS STRIGOSUS, *Wild red rasp'berry*, of the United States. The leaves are often used as a mild astringent, as well as those of *Rubus occidentalis*, *Black Rasp'berry*.

RUBUS TRIVIALIS, see *Rubus cæsius*—*r. Villosus*, see *Rubus fruticosus*—*r. Vulgaris*, *R. fruticosus*.

RUCKING (from *ructus*,) Eructation.

RUCTAMEN, (from *ructare*, *ructatum*, 'to belch,') Eructation.

RUCTATIO, Eructation.

RUCTUATIO, Eructation.

RUCTUOSITAS, see Eructation.

RUCTUS, Eructation—*r. Acidus*, *Oxyregmia*—*r. Nidorosus*, *Cnissoregmia*.

RUDBECK'IA LACINIA'TA, (after the Rudbecks, of Upsal,) *Coneflower*, *Thimbleweed*; indigenous: *Order*, *Compositæ*; is diuretic, and has been given in nephritic affections.

RUDDE, HARD, *Rubrica fabrilis*.

RUDIMENTAL, *Rudimenta'lis*, (F.) *Rudimenta're*, (*rudimentum*, (from *rudis*, 'rude,') 'the first beginning,') That which is but slightly developed.

RUDOLPHIA FRONDOSA, (after C. A. Rudolphi, a botanist of Jena,) *Butea frondosa*.

RUE, Ruta—*r. de Chèvre*, *Galega officinalis*—*r. Common*, *Ruta*—*r. Goat's*, *Galega officinalis*—*r. Goat's*, *Virginia*, *Galega Virginiana*—*r. Meadow*, *Thalictron*—*r. des Mursailles*, *Asplenium ruta muraria*—*r. des Prés*, *Thalictron*—*r. Sauvage*, *Ruta*.

RUELLES, Wrinkles.

RUFT, (from *rive*, *riued*,) Eructation.

RUGA, pl. *Rugæ*, Wrinkle.

RUGINE, (from *Runcina*,) *Raspatorium*.

RUGISSEMENT, *Rugitus*.

RUGITUS, *Brycheth'mus*, (*rugire*, *rugitum*, 'to roar,') (F.) *Rugissement*. A Latin word, which ordinarily expresses the roaring of the lion; and which is used, by some medical writers, to designate a particular species of borborygmus compared to it.

RUGOSE, (from *ruga*,) Wrinkled.

RUGOUS, Wrinkled.

RUM, *Tafia*, *Spir'itus Jamaicæ'sis*, *Molas'sea Spirit*, (F.) *Rhum*. A spirit distilled from sugar, or molasses. See Spirit.

RUMA, Mamma.

RUM, BAY, *Spiritus myrciæ*.

RUM BUDS, *Gutta rosea*.

RUMEX ACETOSELLA, gen. *Ru'micis*, (from *rumo*, 'to suck,' in allusion to the Romans sucking it to allay thirst [?]), *Sheep Sor'rel*, *Field Sorrel*; indigenous; has similar virtues with

RUMEX ACETO'SUS, *R. tuberosus* seu *intermedius*, *Lap'athum pratens'se*, *Ox'alis Aceto'sa*, *Aceto'sa vulg'aris* seu *pratens'sis* seu *nos'tras*, *Anac'yris*, *A. arven'sis*, *Sour dock*, *Sor'rel*, *Com'mon Sor'rel*, (Sc.) *Docken*, (F.) *Oseille*, *Vinette*, *Aigrette*, *Surelle*. Ord. Polygonacæ. *Sex. Syst.* Hexandria Trigynia. The leaves are sour, and are used in cookery, and also as a refrigerant.

RUMEX ACUTUS, *Sharp-pointed Wild dock*, *Oxylap'athum*, *Lap'athum*, *L. sylvestrê* seu *acutatum*, *Ru'mex*, *R. paludosus* seu *pursil'us*, *Lap'athus*, (F.) *Patience sauvage*. A decoction of the root has long been used in cutaneous affections. It is, sometimes, employed for the cure of the itch.

RUMEX ÆTENSIS, *R. scutatus*.

RUMEX ALPI'SUS, *Aceto'sa Alpi'na*. The systematic name of the plant that affords *Monk's Rhubarb*, (F.) *Rhabarbe des Moines*. See *Rumex patientia*.

RUMEX AQUATICUS, *Rumex hydrolapathum*—*r. Britannica*, *Rumex hydrolapathum*—*r. Confertus*, *R. patientia*—*r. Dignus*, *Oxyria reniformis*—*r. Glaucus*, *R. scutatus*—*r. Hastatus*, *R. scutatus*—*r. Hastifolius*, *R. scutatus*.

RUMEX HYDROLAPATHUM, *Water Dock*, (Old Eng.) *Brookcress*, *Hydrolapathum*, *Rumex aquaticus* *seu* *Britannica*, *Herba Britannica*, *Lapathum aquaticum*, (F.) *Parelle*, *Patience d'eau*. The leaves of this plant are subacid, and said to be laxative. The root has been used as a tonic, astringent, and antiscorbutic. It has also been employed externally in some cutaneous affections.

RUMEX INTERMEDIUS, *R. acetosus*.

RUMEX OBTUSIFOLIUS, *Lapathum sylvestris*, *Bluntleaved Dock*, *Bitter dock*. The root of this, as well as of *R. crispus*, *Curl'd*, or *Sour*, or *Yellow Dock*—*Rumex*, (Ph. U. S.)—has similar virtues to those of the other species. Both have been used in decoction in the same cases as sarsaparilla.

RUMEX OXYOTUS, *R. scutatus*—*r. Paludosus*, *R. acutus*.

RUMEX PATIENTIA, *R. confer'tus*, *Lapathum horten'se*, *Rhubarb*, *barum monacho'rum*, *Hippolapathum*, *Patientia*, *Gar'den Patience*, *Monk's Rhubarb*, (F.) *Patience*, *Rhubarbe sauvage*, *Parelle*. This root, which is supposed to possess the virtues of rhubarb, but in an inferior degree, is, according to Linnaeus, obtained from the *Rumex patientia*; according to Murray, from the *Rumex alpinus*.

RUMEX PUSILLUS, *R. acutus*.

RUMEX SANGUINEUS. The root of the *blood'y dock* or *bloody vein'd Dock*, has an austere and astringent taste; and has been employed in dysentery. It is called, also, *Lapathum sanguineum*, (F.) *Oseille* or *Patience Rouge*, *Sangdragon*.

RUMEX SCUTATUS, *R. glaucus* *seu* *hastifolius* *seu* *Athen'sis* *seu* *oxyotus* *seu* *hastatus*, *Acetosarotundifolia* *seu* *Romana* *seu* *Rotundifolia* *horten'sis* *seu* *scutata*, *Lapathum scutatum* *seu* *rotundifolia*, *Roman* or *Gar'den Sorrel*, *Green Sauce*, *French Sorrel*, (F.) *Oseille boucher*, *Oseille ronde*, *Petite Oseille*. Virtues the same as those of the common sorrel.

RUMEX TUBEROSUS, *R. acetosus*.

RUMINATION, *Ruminatio*, (*ruminare*, 'to chew again,' from *ruma* or *rumen*, 'the ingluviae.') *Merycis mus*. A function peculiar to ruminating animals, by which they chew a second time the food they have swallowed. An analogous phenomenon is sometimes seen in man. One who ruminates is termed, in French, *mérycole*.

RUMIS, *Mamma*.

RUMMET, *Pityriasis*.

RUMP, ([D. Romp, [G.] Rumpf.) *Croupon*—*r. Bone*, *Coccyx*.

RUNCHES, *Sinapis arvensis*.

RUNCINA, 'a plane,' *Raspatorium*.

RUNCINULA, (dim. of *runcina*.) *Raspatorium*.

RUNKILL, *Wrinkle*.

RUNKLE, *Wrinkle*.

RUNET, *Rennet*.

RUNNING, *Cur'vus*, (F.) *Courae*. A succession of leaps, executed with rapidity, to transport the body quickly from one place to another.

RUNNING, *Rennet*.

RUN ROUND, see *Paronychia*.

RUNSH, *Sinapis arvensis*.

RUPES, 'cliffs,' *Lips of the Vulva*. See *Lip*.

RUPIA, *Rhy'pia*, *U'vus atonicum*, *Ec'phlysis Rhy'pia*, *Rhypha'ria*, (from *rupes*, 'filth.') An eruption of large flattish blebs, which contain a fluid—at first serous, afterwards puriform, and often bloody, which rapidly concretes into crusts, at the base of which are ulcers of variable depths.

It requires the use of tonics internally, and of stimulating ointments externally.

RUPIA ESCHAROTICA, *Pem'phigus infanti'sis* *seu* *gangrenosus*; known in Ireland under the names—*white blisters*, *eating hives*, and *burnt holes*. An affection which bears a close similarity to pemphigus, particularly in the absence of a thick rugous crust, whilst in its chief feature, that of ulceration, it evidently belongs to *Rupia*.

RUPIAL, *Rupia'lis*. Appertaining or relating to *rupia*.

RUPTIBILITAS, (*ruptio* and *habilis*, 'able,') *Fragilitas*.

RUPTILITAS, *Fragilitas*.

RUPTIO, (from *rumpere*, *ruptum*, 'to break,') *Rhexis*.

RUPTORIUM, (from *rumpere*, *ruptum*, 'to break,') *Cauterium*—*r. Commune*, *Potassa fusa*.

RUPTURA, (from *rumpere*, *ruptum*, 'to break,') *Ecrexis*, *Hernia*, *Laceration*, *Rhexis*—*r. Corneae*, *Ceratorrhexis*—*r. Herniosa*, *Hernia*—*r. Ovarii*, *Ovariorrhexis*—*r. Uteri*, *Uterus*, *rupture of the*—*r. Vaginae*, *Colporrhexis*.

RUPTURE, (from *rumpere*, *ruptum*, 'to break or burst,') *Hernia*, *Rhexis*—*r. du Cœur*, *Cardiorrhexis*—*r. of the Groin*, *Bubonocoele*—*r. de la Mitrice*, *Uterus*, *rupture of the*—*r. Wort*, *Herniaria glabra*.

RUPTURED, see *Hernial*.

RUSCUS, *R. aculeatus*, *Bruscus*, *Oxymyrrhine*, *Oxymyrrhine*, *Myrtacantha*, *Catanagelus*, *Myacan'tha*, *Sco'pia re'gia*, *Wild Myrtle*, *Centromyrrinē*, *Butcher's Broom*, (Prov.) *Kneebolly*, (F.) *Petit houx*, *Fragon*. *Ord. Smilacaceae*. *Sez. Syst. Dioclea* *Monadelphica*. The root has been recommended as an aperient and diuretic.

RUSCUS ACULEATUS, *Ruscus*.

RUSCUS HYPOGLOSSEM, *Uvula'ria*, *Hypoglossum*, *Hypoglossum* [?], *Bisligna*, *Bonifolia*, *Laurus Alexandrina angustifolia*, (F.) *Laurier Alexandrin*, *Langue de cheval*. Formerly used against relaxed uvula.

RUSH, SWEET, ([Sax.] *ni'rc*, *ni'rc*, 'a reed.') *Acorus calamus*, *Juncus odoratus*.

RUSH, UTERINE, ([D.] *ruischen*, 'to murmur,') *Bruit placentaire*.

RUSMA, see *Orpiment*.

RUSPINI'S STYPTIC, (pron. *Ruspeen'i's*.) see *Styptic*, *Ruspini's*.

RUSPINI'S TINCTURE FOR THE TEETH. An empirical preparation, composed of *Florentine orris* ʒviij. *Cloves* ʒj. *Rectified spirit* ʒij. *Ambergis* ʒj.

RUT, (from *rugitus*, 'the exclamation of animals in heat,') *Heat*.

RUTA, (from *R. graveolens* *seu* *horten'sis* *seu* *altera*, *Peganum*, *Bosa'sa*, *Common Rue*, (F.) *Rue*, *Rhue*, *Rue sauvage*, *Herbe de grace*; *Ord. Rutaceae*. *Sez. Syst. Decandria Monogynia*. The plant has a strong, ungrateful odour, and bitter pungent taste. It is so acrid as to blister the skin. Its properties are reputed to be tonic, stimulant, antispasmodic, and emmenagogue. Dose, gr. xv to ʒj. The leaves are officinal—*Ru'ta*. (Ph. U. S.)

O'leum Ru'tae, *distilled Oil of Rue*, is antispasmodic; and, externally, rubefacient. Dose, gtt. ij to vj.

RUTA ALTERA, *Ruta*—*r. Capraria*, *Galega officinalis*—*r. Graveolens*, *Ruta*—*r. Hortensis*, *Ruta*—*r. Muraria*, *Asplenium ruta muraria*.

RUTABULUM, *Penis*.

RUTACEUM, (from *ruta*.) *Vinegar of rue*.

RUTIDOMA, gen. *Rutido'matis*, see *Rutidosia*.

RUTIDOSIS, *Rutido'sis*, *Rhytido'sis*, (*rutidom*, from *rutidom*, 'I wrinkle,' and *osis*.) The act of wrinkling; as *Rutido'ma* means a shrivelled or

wrinkled body. A destruction or atrophy of the eye. Corrugation and subsidence of the cornea.

RUYSCHIA'NA TU'NICA, (F.) *Lame* on *Membrane Ruyschienne*. According to Ruysch, (pron. *Risch*), the Dutch anatomist, and the greater part of his disciples, the choroid coat of the eye is formed of two laminae. His son gave to the inner the name *Tu'nica Ruyschia'na* or *Membra'na Ruyschia'na* seu *Ruysch'i'i*.

Tu'nica Cellulo'sa Ruyschia'na is the areolar substance under the peritoneal covering of the stomach. It is not numbered among the coats of that organ.

RYE, ([Sax.] *ryge*.) Secale—r. Horned, Ergot—r. Robbin's, *Polytrichum juniperinum*—r. Spurred, Ergot.

RYTIDOSIS, *Rutidosis*.

RYTIS, (*ρυτίς*.) Wrinkle.

S.

S, S. A. &c., see Abbreviation.

S. ROMANUM, Sigmoid flexure of the Colon.

SAAG, Urine.

SABADILLA, ([S.] *Cebadilla*.) *Veratrum sabadilla*, see *Veratrina*.

SABADILLIN, see *Veratrina*.

SABATILLA, *Veratrum sabadilla*.

SABBATIA, (after *Sabbati*, an Italian botanist.) *Chironia angularis*—s. *Angularis*, *Chironia angularis*.

SABINA, *Juniperus sabina*.

SABINE, *Juniperus sabina*.

SABLÉ, 'sanded,' (from *sabulum*, 'sand.') Having the appearance as if strewed with sand. The terms *sablé* and *piqueté* are sometimes applied to the brain, when a slice of it exhibits red points as if it had been strewed with red sand. Such is the case in cerebral hyperæmia.

SABOT DE VÉNUS JAUNE, 'yellow sabot of Venus,' *Cypripedium luteum*.

SABUCUS, *Sambucus*.

SABULOUS, *Sabulo'sus*, *Areno'sus*, *Psammo'des*, (*sabulum*, 'sand.') The sandy deposit often seen in the urine.

SABULUM CONARII, see Pineal gland.

SABURA, *Saburra*.

SABUR'RA, *Sabu'ra*, *Sor'des*, (F.) *Saburre*. This word signifies, by derivation, 'coarse sand, or ballast,' *sabulum*. In its common acceptation, it means *foulness of the stomach*, as it is sometimes called;—vitiated matters supposed to be retained and accumulated in the stomach—the result of imperfect digestion—and which have been sometimes considered as a morbid product of the mucous secretion of that organ, or of the biliary secretion; at others, as a residuum from alimentary substances badly digested. The *Sabur'ral state*, *Collu'ries gastrica*, *Crudita'tes ventric'uli*, *Gastris'mus*, (F.) *Embarras gastrique*, *État saburral*, is an accumulation of *saburrae* in the stomach, (F.) *Saburres gastriques*, which the Humorists considered to be the cause of a number of diseases.

SABURRA VERMINOSA, *Helminthiasis*.

SABUR'RAL, *Saburra'lia*. Relating or appertaining to *saburrae*,—as the 'saburral state.'

SABURRATIO, *Arenatio*.

SABURRE, *Saburra*.

SABURRES GASTRIQUES, see *Saburra*.

SAC, see *Sack*—s. *Herniaire*, *Hernial sac*—s. *Lachrymal*, see *Lachrymal puncta*—s. of the *Omentum*, see *Epiploon*, *gastrohepatic*.

SACCADE, see *Respiration*, *jerking*.

SACCATED, *Sacca'tus*, (from *saccus*, 'a sac,') Encysted. Also, ponched, as the Colon. *Sacculated*, *Saccula'tus*, are diminutives of *Saccated*, and *Saccatus*.

SACCHAR, (*σακχαρ*.) *Saccharum*.

SACCHARHYDROLE, *Syrup*.

SACCHARI FÆX, 'dregs of sugar,' *Melasses*.

SACCHARINA, *Saccharolés*.

SACCHARINE, *Sacchari'nia*, (F.) *Saccharin*, *Glucosique*, *Glycosique*, (from *saccharum*, 'sugar,') Having the nature of, or containing, sugar.

SACCHARO-ALCOOLÉ, *Ratafia*.

SACCHAROGUMMITA, (*saccharum*, and *gummi*.) see *Glycyrrhiza*.

SACCHAROLE LIQUIDE, *Mellitum*, *Syrup*—s. *Mou*, *Conserva*, *Electuarium*—s. *Solide*, see *Pastel*, and *Tabella*.

SACCHAROLÉS (F.), *Sacchari'na*, (from *saccharum*.) A name given by MM. Henry and Guibourt to medicines which have sugar or honey for excipient, or for predominant principle. M. Béal so calls an exact mixture of pulverized sugar with other substances also in powder, but in less quantity.

SACCHAROMETER, (F.) *Saccharomètre*, *Pèse-sirop*, (*saccharum*, and *μετρον*, 'measure,') A hydrometer adapted for determining the density of syrups. See *Areometer*.

SACCHAROMYCES, (*saccharum*, and *μυκός*, 'fungus,') *Torula cerevisiæ*.

SACCHARORRHŒA, (*saccharum*, and *ρῆμα*, 'to flow,') *Glycyrrhœa*—s. *Cutanea*, *Sudor dulcis*.

SACCHARORRHŒA LAC'TEA, *Galactorrhœa saccharata*. A secretion of unusually sweet milk.

SACCHARORRHŒA PULMONA' LIS, *Sputum dulcè* seu *saccharatum*. The copious expectoration of unusually sweet matter from the lungs.

SACCHARORRHŒA URINOSA, *Diabetes mellitus*.

SACCHARUM, (*σακχαρ*.) *Sac'char*, (*σακχαρ*), from Arabic, or rather Sanscrit, *Sakar*, 'white earth,') *Su'kar*, *Suc'char*, *Su'ter*, *Zu'char*, *Zu'caro*, *As'sakar*, *Ti'gala*, *Zac'charum*, *Zo'zar*, *Zu'charum*, *Zuc'carum*, *Zu'chra*, *Mel arundinaceum*, *Sugar*, (F.) *Sucre*. The produce of the juice of *Sac'charum officina'rum* seu *officina'le* of Linnaeus, *Cal'amus Indicus* seu *Sacchari'nis*, *Arund' saccharifera* of Sloane; *Sugar Cane*, (F.) *Canne à Sucre*, *Cannamelle*. Ord. Gramineæ; Sex. Syst. Triandria Digynia; and also of the beet, maple, *Chinese sugarcane*, (*Sorghum saccharatum*.) &c.

True Sugars are those with which alcoholic fermentation may be effected, by the addition of yeast and water. They are:—

Species.

Varieties.

- | | | |
|--------|---|--|
| 1..... | { | Hard sugar of the cane, maple, beet, chestnut, &c. |
| 2..... | | Liquid sugar, sugar of malt, of the sweet potato, molasses, honey, &c. |
| 3..... | { | Hard sugar of the grape, ripe fruits, starch, diabetes, &c. |
| 4..... | | Hard sugar of the mushroom. |

Imperfect sugars are those in which alcoholic fermentation cannot be effected by means of yeast and water.

posterior surface of the sacrum; which appear to terminate the vertebral grooves or gutters.

SACRAL NERVES, (F.) *Nerfs sacrés*, are generally six in number. They arise from the termination of the spinal marrow. The first, united with the great cord which descends from the lumbar plexus, concurs with the three others in the formation of the sacral or sciatic plexus. The third and fourth, with branches of the great sympathetic, form the hypogastric plexus. The fifth and sixth (*coccygeal nerve*) are distributed to the ischiococcygeus muscle, and to the sphincter and levator ani.

SACRAL PLEXUS, Sciatic plexus—the *sacral portion of the crural plexus* of Chaussier, constituted as just mentioned, is situate in front of the pyramidalis. It gives off, backwards, two branches, called the *pubic, (F.) honteux*, and the *lesser sciatic*. It terminates in the great sciatic nerve.

SACRÉ, Sacral.

SACRO, in composition, relating or belonging to the sacrum.

SACROCCOXYGEAL, Sacrococcygeus. Relating to the sacrum and coccyx.

SACROCCOXYGEAL ARTICULATION is that of the lower extremity of the sacrum with the superior *facette* of the coccyx. It is closed by two ligaments,—an *anterior* and a *posterior sacrococcygeal*.

SACROCASTALIS, (costa, 'a rib,') *Sacrolumbalis*—*s. Femoral, Gluteus major*.

SACROCOXAL'GIA, (sacro, coxa, 'the hip,' and algos, 'pain,') Some authors have called thus a rheumatic affection of the sacroiliac symphysis.

SACROÏLIAC, Sacroïli'acus. That which relates to the sacrum and ilium.

SACROÏLIAC ARTICULATION, SYN'PHYSIS or **SYNCHONDRO'SIS, Il'iosacral articulation, symphysis or synchondro'sis**, is the union of two surfaces,—two inclined planes, broader above than below; one formed by the sacrum, the other by the ilium. Both surfaces are incrustated by an articular cartilage. The bands which strengthen them are the two *sacrosciatic ligaments*, the *sacrospinal*, a *sacroïliac* ligament, and some irregular fibres. The *sacroïliac* ligament is a very thick assemblage of short, dense, irregular fibres, occupying the space which the sacrum and ilium leave between them, behind their articular surface. Some authors have termed some irregular fibres, situate in front of the sacroiliac articulation, *anterior sacroïliac ligaments*, in opposition to the preceding fasciæ, which they call *posterior sacroïliac*.

SACROÏSCHIATICUS, Sacrosiatic.

SACROLUMBAIRE, Sacrolumbalis.

SACROLUMBALIS, Sacrolumbar.

SACROLUMBAR, Sacrolumba'lis, Sacro-lombarre. That which belongs to the sacrum and loins.

SACROLUMBAR, Sacrolumba'lis, Sacrolumba'ris *Muscle, Sacrocostalis, Dorsotrachélien portion of the Sacrospinal*, of Chaussier, (F.) *Lombocosto-trachélien, Sacro-lombarre*, is seated at the posterior part of the trunk; is thick and triangular below; flat and thin in the remainder, and terminates in a point at its upper extremity. It is attached to the posterior surface of the sacrum; to the corresponding portion of the crista ili; to the extremities of the transverse processes of the lumbar vertebræ; to the angles of the eleven inferior ribs; to the tuberosity of the first, and the posterior tubercle of the transverse processes of the last five cervical vertebræ. The sacrolumbalis straightens the vertebral column when bent forwards, and maintains it straight. By means of its insertion into the ribs, it can elevate or depress these bones, according as it contracts from

above to below, or from below to above. It acts, consequently, sometimes as an inspiratory, at others as an expiratory muscle.

From six or eight of the lower ribs arises an equal number of fleshy portions, which terminate at the inner side of this muscle, and are called *Musculi Accessorii seu Additamentum ad Sacrolumba'lem, (F.) Accessoires du Sacro-lombarre*.

From the upper part of the sacrolumbalis a fleshy slip, called *Cervical'is descendens, Cervicalis descendens Diemerbroeckii, Transversalis collateralis colli, Accessorius ad Sacrolumba'lem, (F.) Transversaire grêle*, runs up, to be fixed to the transverse processes of the fourth, fifth, and sixth cervical vertebræ, by three distinct tendons. These last parts, though separated by some anatomists, belong to the sacrolumbalis.

SACROSCIATIC, Sacroïschiat'icus. That which belongs to the sacrum and ischium.

SACROSCIATIC LIGAMENTS are two ligaments which assist in securing the sacroiliac articulation. The *great or posterior sacrosciatic ligament* extends from the crista ili and the sides of the sacrum and coccyx to the tuberosity of the ischium. The *less or anterior sacrosciatic ligament* arises from the same point, and terminates in the spine of the ischium.

SACROSPINAL, Dorsotrachélien portion of the sacrolumbalis.

SACROSPINALIS. That which relates to the sacrum and spine. Under this name, Chaussier describes the sacrolumbalis, longissimus dorsi, transversalis, transversospinalis, and intertransversalis muscles. He divides it into three portions. It is the *Erector spinæ, Extensor dorsi communis, Lum'bocostalis*.

SACROTROCHANTÉRIEN, Pyramidalis.

SACROVERTEBRAL, Sacrovertebra'lis. That which relates to the sacrum and the vertebræ.

SACROVERTEBRAL ARTICULATION is the junction of the last vertebræ of the loins with the sacrum. The angle, which these two bones form at the anterior part, has been called the *Promontory of the Sacrum*, or the *Sacrovertebral Angle*.

SACROVERTEBRAL LIGAMENT is a strong ligament, which passes from the anterior part of each of the transverse processes of the last lumbar vertebra to be attached to the base of the sacrum. Its use is to maintain the articulation of these two bones *in situ*.

SACRUM, (neuter of sacer,) *Os sacrum seu basilarè seu Al'agas seu la'tum seu clu'nium, Vertebra magna, Albagia'zi, Hi'eron ost'eon, Clu'nis, Subvertebra, Subvertebrum, Vet'ula*, (from sacer, 'sacred;' because it contributes to protect the genital organs, which were considered sacred; or because it was offered in sacrifice.) The bone, which forms the posterior part of the pelvis, and is a continuation of the vertebral column. The sacrum is symmetrical and triangular, situate at the posterior part of the pelvis, and concave anteriorly. It has: 1. A *pelvic or anterior surface*, which corresponds to the pelvic cavity, and has four transverse furrows, that point out the parts where the primitive portions of the bone were separated from each other. On each side there are four foramina, called *anterior sacral*; these are oblique, and decrease in size from above downwards. Through them pass the anterior branches of the sacral nerves. 2. A *spinal or posterior surface*, which is convex, rough, and covered by muscles. Upon the median line are four horizontal, tubercular eminences, forming a continuation of the spinous processes of the vertebræ; and below, a triangular notch, where the *sacral canal* terminates. On each side of the surface are four foramina, called *posterior sacral*,

which are smaller than the anterior, and give passage to the posterior branches of the sacral nerves. 3. The *transverse surface* or *face*. Its greatest extent is transversely. It is articulated with the last lumbar vertebra. On it is seen the *rim of the sacral canal*, of a triangular shape, which decreases in width from above downward; constitutes the ventral canal; is lined by a prolongation of the cerebral membranes; contains the fasciculi of the sacral nerve; communicates, externally, by the sacral foramina; and terminates at the triangular notch on the posterior surface of the bone. At the sides of the upper orifice of the sacral canal are two articular processes, which receive the articular processes of the last lumbar vertebra. The projection, formed by the union of the base of the sacrum with the last lumbar vertebra, is called the *Promontory*. *From the Sacrum Derive Several Great Angles of the Sacrum or Sacrum's four Angles*. 4. A *conjugate face* or *apex*, (F.) *Face conjugata* on *Sacrum*, which is very narrow, and is articulated with the coryx. 5. The *basal coryx*, which have, above, a broad, oblique surface, articulated with the ilium.

The sacrum is developed by 30 to 36 points of

SADDLE, TURKISH. *Sella Turca.*

DEER, 2000.

SAFFLOW, *Carthamus tinctorius*.

SAFFLOWER, *Carchamus tinctorius*.

SAFFRON. *Arak safran*, 'yellow,' [?] — *s. Bastard*, *Carthamus tinctorius* — *s. Dyer's*, *Carthamus tinctorius*.

Saffron Flower. *Carthamus tinctorius*.

Saffron. Meadow, *Colchicum autumnale*.

S. AFRICA. Crocus — *s. Bataard*, *Carthamus tinctorius*, *Colchicum autumnale* — *s. de India*, *Carouma longa* — *s. de Mare agreste*, *Ferri sub-carthamus* — *s. de Mare arvensis*, *Colchicor* — *s. des Pêles*, *Colchicum autumnale* — *s. Racine de*, *Carouma longa*.

SAFRANUM. *Carthamus tinctorius*.

DATE -

SAGE. *Mullein.* In ancient Rome, the Sage exercised, also, the calling of magicians, procuresses, and perfumers: and were more especially called upon to induce abortion. Also, Sage.

SAGAPAN. Arins 12147-52.

[illegible]

SAGAPIN. *Sagapenum*.

SAGE. *Salvia* — s. Indian, *Eupatorium perfoliatum* — s. Jerusalem, *Palm. varia*.

Sage Tea. Infusum salviae.

SAGE TREE, LANTANA.

SAGE. Wild. *Salvia Africana*. S. *Lyrate*.
SAGE. *LEMME*.

SAGE-LEMMLE, 'sage or wise woman.' Mil-
wife.

SAGENA, *comp.* 'a seine.' English.

SAGESSE DES CHIRURGIENS, 'wisdom
of the surgeons.' Sisymbrium Sophia.

SAGIMEN VITRI. Soda, subcarbonate of.

SAGITTAL. *Sagittaria*, *Sagittaria*, (from
sagitta, 'an arrow.')

SAGITTAL FURROW, Sagittal Groove, [F.] *Groove*. *furca sagittalis*, is a channel on the median line of the inner surface of the skullcap. It is excavated in the os frontis, in the two parietal and occipital bones, and extends from the crista galli to the internal occipital protuberance. It is con-

the longitudinal sinus, which has been called the
great sinus.

SAGITTAL SUTURE. *Osteon saggittale seu Enigle*, *Sutura Vaginalis seu Jugalit seu Pabbat seu Rhachitica seu Rhachialis*, *Rhachidæna seu Epignathicæ seu Jugalis* *articularis*. The suture which unites the two parietal bones, and which extends from before to behind, on the median line, from the coronal suture to the lambdoidal. It is so called, from *sagitta*, ‘an arrow,’ because it seems to meet the coronal suture as an arrow meets the string of a bow.

SAGITTARIA VARIABILIS, from an arrow, an arrow-wing to the prevalent form of the leaves. *A. arifolia*, *A. cristata*, *A. lanceolata*, *Wapiti* of Oregon; *Nat. Qu.*, *Alismaceae*. Eminent, yielding fecula like arrow-root. The roots are cooling and subastringent, and are at times applied to the feet in cases of yaws and dropsical swellings of the legs. The leaves are applied to the breasts to discol milk.

SAGITTARIUM ALEXIPHARMACUM.
Melastomaceae. Common Indian. *Acanthaceae*. Indian.
A root cultivated in the West Indies, and supposed
to be a remedy for wounds by poisoned
arrows.

SAGITTATUS, *Sacchar*

SAGO, *Sagum*, *Sagum*, *Sagum*, *Sagum*, *Andropogon*
palustris (var.), *F. Sagum*, *Sagum*, *Sagum* is the Malay
name for various palms.) A fecula obtained from
the pith of the *Sagum Rumphii*, *Hydn. saccatum*,
E. candida, etc., growing in the Moluccas, Philip-
pine Isles, etc., and which is brought to us in
small grains. The same substance is also obtained
from the West Indies, but it is inferior to that
from the East. By boiling in water or milk,
sago becomes soft and transparent, and forms an
agreeable and nutritious food in febrile, calculous,
and other disorders. It is made palatable by
sugar, lemon-juice, or wine, where the last is not
contraindicated.

To make sage into a proper mucilage for the sick, a tablespoonful of it should be macerated in a pint of water for two hours, and then boiled for fifteen minutes, stirring assiduously. Care must be taken, that the grains are perfectly dissolved by long boiling and constant stirring.

Sassafras is made by putting ss. into water. Oij: boiling until the mucilage is formed: then rubbing ss. on the rim of a lemon, and putting it with ss. of ss. into them. Oss: adding this mixture to the mucilage, and boiling for five minutes. — It is a good restorative.

SAGE, FRENCH, see *Silene tuckerm.* and *Artemisia*, common—s. Portland Island, see *Artemisia maculatum*.

SAGOL, Sago.

SAGU. Saginaw.

SAGUE, Sage — *a. Paris*, see *Solanum tuberosum*.

SAGUERUS GOMATO, see Toddy—s. Rum-
phk. see Toddy—s. Saccharifer. see Toddy.

SALES, Sec'y, Boston, Mass.

SAMARA. Ins. m. 13.

SAIGNÉE. (from *saigner*, [L.] *sanguis*, "blood," or *bleed*.) Bloodletting — a. *Sanguinea*, *Spoliativa phlebotomia*.

SAIGNEMENT DU NEZ. Bleeding of the nose. Epistaxis.

SAILORS' DISEASES. *Ac.* **OF.** Sailors, by nature of their occupation, are liable to many diseases. Several of these may, however, be warded off by proper attention to cleanliness and ventilation.

SAIN BOIS. 'health wood.' *Daphne genkwa*
— *Douc.* (from *sapida*, 'fatness.'), *A. leys* praya-
talis.

SAINT AMAND, MINERAL WATERS OF. Sulphuretted waters, in the department of Du Nord, France. They contain carbonic and sulphohydric acids. The *Boue* or deposit contains sulphohydric acid, and salts of iron and magnesia. Temperature, 64° to 82° Fahrenheit. Season, from the 1st of June to the 1st of September.

SAINT ANTHONY'S FIRE, Erysipelas.

SAINT AUGUSTINE', CLIMATE OF. This sea-port of Florida is a frequent retreat during the winter season for the valetudinarian from the more northern parts of the United States. The climate is certainly comparatively mild and equable: but it is liable to frequent northeast storms. There are only a few days, however, in winter, in which the invalid cannot take exercise in the open air. Tampa Bay, and Pensacola, in Florida, and Mobile, in Alabama, which are sometimes visited by the consumptive, are less liable to the storms in question.

SAINT CATHERINE'S SPRINGS, see Canada, Mineral waters of.

SAINT CHARLES'S ROOT, Carlo Sancto (Radix).

SAINT CROIX, Santa Cruz.

SAINT JAMES'S WORT, Senecio Jacobææ, *Thlaspi bursa*.

SAINT JOHN'S DANCE, see Mania, dancing. —St. John's Wort, *Hypericum perforatum*.

SAINT KITTS, see West Indies.

SAINT LEONARD'S, CLIMATE OF. This place, which is situate about a mile to the westward of Hastings, in Sussex, England, is less sheltered than it, and is therefore a less favourable retreat, during the winter months, for the phthisical valetudinarian.

SAINT MYON, MINERAL WATERS OF. Some cold, acidulous waters at Saint Myon, near Riom, in the department of Puy-de-dôme. They contain free carbonic acid, subcarbonate of soda, chloride of sodium, and carbonate and sulphate of lime.

SAINT NECTAIRE, MINERAL WATERS OF. Springs in the department of Puy-de-Dôme, France. They contain carbonic acid, bicarbonate of soda, and chloride of sodium. Temperature, 75° to 103° Fah. Season, from June 15th to September 15th.

SAINT PARDAUX, MINERAL WATERS OF. Chalybeate springs in the department of Allier, France. They contain carbonic acid and iron.

SAINT SAUVEUR, MINERAL WATERS OF. Sulphurous waters in the department of Hautes-Pyrénées, France. Temperature, 76° to 93° Fah. The springs are frequented from May to October.

SAINT VINCENT'S, see West Indies.

SAINT VITUS'S DANCE, Chorea, see Mania, dancing —s. Vitus's Dance of the Voice Balbuties.

SAIR, (sore,) Uleer.

SAIRE, Essera.

SAISON, Season.

SAKI, see Spirit.

SAL, gen. *So'lis*, (ἀλς.) Salt —s. Absinthii, Potassæ subcarbonas, see Potash —s. Acetosellæ, Potassæ, oxalate of —s. Acetosus ammoniacalis, Liquor ammoniæ acetatis —s. Acidum benzoës, Benjamin —s. Acratus, see Potassæ carbonas —s. Allus, Soda, muriate of —s. Alembrothi, Alembroth salt —s. Alkali, Soda, subcarbonate of —s. Alkalinus fixus fossilis, Soda —s. Amarum, Magnesie sulphas —s. Ammoniac, volatile, Ammonie carbonas —s. Ammoniacum, Ammonie murias —s. Ammoniacum martiale, Ferrum ammoniatum —s. Ammoniacum secretum Glauberi, Ammonie sulphas —s. Ammoniacum tartareum, Ammonie tartaras —s. Ammoniacum vegetabile, Liquor ammoniæ acetatis —s. Ammoniacus nitrosus, Ammonie nitras —s. Ammonie, Ammonie murias —s. Anglicanum,

Magnesie sulphas —s. Anglicum, Magnesie sulphas —s. Antiepilepticus of Weissman, Cuprum ammoniatum —s. Antimonii, Antimonium tartarizatum —s. Armoniacum, Ammonie murias —s. Artis, Alembroth salt —s. Asphalitis, Bit noben —s. Catharticus amarus, Magnesie sulphas —s. Catharticus Glauberi, Soda, sulphate of —s. Communis, Soda, muriate of —s. Cornu cervi volatile, Ammonie carbonas —s. Culinaris, Soda, muriate of —s. Digestivus Sylvii, Potassæ murias —s. Diureticus, Potassæ acetas —s. de Duobus, Potassæ sulphas —s. Ebsdamense, Magnesie sulphas —s. Enixum, Potassæ supersulphas —s. Epsomensis, Magnesie sulphas —s. Essentiale Benzoës, Benjamin —s. Essentiale corticis Peruviani, Quinine —s. Febrifugus of Lemery, Potassæ supersulphas —s. Febrifugus Sylvii, Potassæ murias —s. Fontium, Soda, muriate of —s. Fossilis, Soda, muriate of —s. Fuliginis, Ammonie carbonas —s. Gemme, Soda, muriate of —s. Herbarum, Potassæ subcarbonas —s. Indus, Saccharum album —s. Lixiviosum, see Lixivium —s. Lucidum, Soda, muriate of —s. Marinus, Soda, muriate of —s. Martis, Ferri sulphas —s. Martis muriaticum sublimatum, Ferrum ammoniatum —s. Mirabile perlatum, Soda, phosphate of —s. Mirabilis Glauberi, Soda, sulphate of —s. Muriaticum baroticum, Baryta, muriate of —s. Nitrum, Potassæ nitras —s. Ossium, Ammonie carbonas —s. Panchrestum, Potassæ tartaras —s. Petræ, Potassæ nitras —s. Plumbi, Plumbi superacetas —s. Polychrestum, Potassæ sulphas —s. Polychrestum Seignetti, Soda, tartrate of —s. Polychrestum Glaseri, Potassæ sulphas cum sulphure —s. Prunelle, Potassæ nitras fusum sulphatis paucillo mixtus —s. Rupellensis, Soda, tartrate of —s. Sapientie, Alembroth salt, Potassæ sulphas —s. Saturni, Plumbi superacetas —s. Scientie, Alembroth salt —s. Secretus Glauberi, Ammonie sulphas —s. Sedativus Hombergi, Boracic acid —s. Sedlicensis, Magnesie sulphas —s. Seignetti, Soda, tartrate of —s. Sennerti, Potassæ acetas —s. Seydschützense, Magnesie sulphas —s. Sodæ, Soda, subcarbonate of —s. Sodomenus, Bit noben —s. Succini, Succinic acid —s. Tartari, Potassæ subcarbonas, see Potash —s. Tartari essentielle, Tartaric acid —s. Tartari sulphuratum, Potassæ sulphas —s. Urinæ, Microcosmicum Sal —s. Urinosum, Urea —s. Vegetabilis, Potassæ tartaras —s. Vitæ, Alembroth salt —s. Vitrioli, Zinci sulphas —s. Vitrioli narcoticum, Boracic acid —s. Volatile, Ammonie carbonas, Spiritus ammonie aromaticus —s. Volatile Benzoës, Benjamin —s. Volatile boracis, Boracic acid —s. Volatile oleosum Sylvii, Spiritus ammonie aromaticus —s. Volatile succini, Succinic acid —s. Volatilis salis ammoniaci, Ammonie carbonas —s. Vomitorium vitrioli, Zinci sulphas.

SALAAM or SALAM CONVULSION, see Convulsion, Salaam.

SALAB, Salep.

SALACITAS, Salacity, Satyriasis —s. Vulvæ, Nymphomania.

SALAC'ITY, *Salac'iousness, Salac'itas*, (F.) *Salacité, (salax, salac'is, 'lustful, lecherous.)* Lecherousness; strong desire for venery.

SALAD, (from [L.] *sal*, salt, [F.] *salé*, 'salted,') *Sallet*, (F.) *Salade*. Fresh herbs —as lettuce, dressed with salt, oil, vinegar, &c.

SALAD, CORN, Valeriana dentata.

SALAP, Salep.

SALCOMBE, CLIMATE OF. A small place —the Montpellier of Huxham —which is one of the warmest spots on the S. W. coast of England, and, therefore, in this respect adapted for the phthisical invalid as a winter retreat.

SALEP', *Salop', Saloop'* (usually so pronounced.) *Salab', Salap'.* The prepared bulb of the *Orchis mas'cula* and other species of Orchis. The salep of Cachmere is from a species of *Euloph'ia*. Salep is in small, irregular, hard masses. In its composition in relation to water, it is closely

like the quinia in medicinal efficiency. Dose, ʒi to ʒi of the powder. Dose of *Salicin* from 4 to 6 grains.

SALLENDERS, Malandria.

SALLIS, Adeps.

SALLOW, (from [F.] *sale*, 'dirty,') Lurid.

SALLY, Salix.

SALLYWITHY, Salix.

SALMIAC, contraction of *sal ammoniac*, Ammoniac maris.

SALOOP, Salep. Also, sassafras tea flavored with milk and sugar. Under this name a tea is sold in the streets of London which consists of an infusion of the wood of *Laurus Sassafras* with the addition of milk and sugar—*Sassafras Tea*. Sassafras tea has been long drunk as ordinary tea in different parts of the United States, where it is sometimes called *Sass tea*.

SALOP, Salep.

SALPETRA, (*sal*, and *petra*, 'rock,') Potassæ nitras. Soda nitrate of.

SALPINGEMPHRAXIS, (*salpingo*, and *emphrax*, 'obstruction,') Obstruction of the Eustachian tube,—a cause of deafness.

SALPINGITIS, (*salpingo*, and *itis*,) Inflammation of the Eustachian tube.

SALPINGO, (from *salpinge*,) in composition, the Eustachian, or the Fallopian, tube.

SALPINGOCATHETERISMUS, (*salpingo*, and *catheterisma*,) Catheterism of the Eustachian tube.

SALPINGOCYESIS, *Paragynesis tubarum*, *Gynagitis tubarum*, (*salpingo*, and *gynesis*, 'pregnancy,') Tubal pregnancy.—See Pregnancy, tubal.

SALPINGOMALLÉEN, Tensor tympani.

SALPINGOPHARYNGÆUS, (*salpingo*, and *pharynx*, 'the pharynx,') Valsalva and Douglas have given this name to a bundle of fibres, which passes from the Eustachian tube to the pharynx.

SALPINGOSTAPHYLINUS, Circumflex muscle—s. *Staphylinus internus*, Levator palati.

SALPINGOSTENOCHORIA, (*salpingo*, *stenos*, 'narrow,' and *choria*, 'space,') Narrowness or stricture of the Eustachian tube.

SALPINX, gen. *Salpingis*, *σαλπιγξ*, gen. *σαλπιγγος*, Tuba.

SALSIFY, *Tragopogon portifolium*.

SALSAMENTUM CRUDUM, ('salted fish,' from *salsus*, 'salted,') *Osmariobus*.

SALSAPARILLA GERMANICA, Sarsaparilla Germanica.

SALSEDQ, gen. *Salsedinis*, (from *salsus*, 'salt,') Muria—s. *Mucrum*, Potassæ nitras.

SALSEPAREILLE, *Smilax sarsaparilla*—s. *d'Allemagne*, Sarsaparilla Germanica—s. *du Brésil*, s. *du Pérou*, s. *de Honduras*, s. *de Pérou*, and s. *de Portugal*, see *Smilax sarsaparilla*.

SALSIFI, *Tragopogon portifolium*.

SALSILAGO, (from *salsus*, 'salt,') Muria.

SALSITUDO, gen. *Salsitudinis*, (from *salsus*, 'salt,') Muria.

SALSOLA DECUMBENS, s. Kali.

SALSOLA KALI, s. *Decumbens*, *Kali soda seu epinaeum coarctatum*, *Traquina sive Traquin Martiana*, *Prickly Saltwort*, Ord. Chenopodiaceæ; (from *salsus*, owing to its saline properties.) This plant, as well as *Salsola veru* and *Salsola soda seu longifolia*, *Kali inermis*, *Halba vitri*, produces the alkaline salt, commonly called *Bacilla*, *Soda*, or *Kelp*.

SALSOLA LONGIFOLIA, s. Kali—s. *Sativa*, &c., see s. Kali, and *Soda*—s. *Soda*, see s. Kali.

SALSGINOSUS, 'briny,' (from *salsus*,) Halmyrodes.

SALSUGO, gen. *Salsuginis*, (from *salsus*, 'salt,') Muria.

SALSUS, Saline.

SALT, *Sal*, *Hal*, *Hals*, (*sals*, [F.] *Sal*. A name formerly given to a considerable number of sub-

stances of very different nature. At present, chemists apply the term *Salis* generally, to the combination of an acid with one or more bases. *Neutral salts* are those in which the acid and base reciprocally saturate each other. *Acid salts* or *super-salts*, [F.] *Sur-sels*, are those in which the acid predominates over the base, and the prefixes *bi* and *ter* indicate the proportion of acid. *Alkaline salts* or *subsalts*, [F.] *Sous-sels*, those in which the base is predominant, when the prefixes *di* and *tre* indicate the proportion of base. The different denominations of the salts are compounded, 1. Of the name of the acid: the termination of which is changed into *ite*, when the name of the acid is in *ans*;—*ate*, when it is in *ic*. Thus the salts formed by the sulphurous acid are *sulphites*; those by the sulphuric, *sulphates*. 2. This name is followed up by that of the oxyd, which enters into the composition of the salt. Thus, the *sulphate of protoxyd of iron* is a combination of sulphuric acid with the protoxyd of that metal. Frequently, to shorten the name, the word *oxyd* is suppressed; and *proto*, *deuto*, *tritio*, &c., put before the acid:—as *protosulphate of iron*. *Simple salts* are those resulting from the combination of an acid with a single base; *double salts* with two different bases; *triple* with three, &c. A *biacid salt* is one constituted after the type of common salt, or which contains a metal and a *saltrical*, as chlorine, iodine, &c. An *oxyalt* is formed from the combination of an acid with an oxyd.

When *salt* is used in an abstract sense, it means the chloride of sodium.

SALT APERIENT, OF FREDERICK, Soda, sulphate of—s. Baker's, Ammoniac carbonas—s. of Barilla, Soda, subcarbonate of—s. Bay, Soda, muriate of—s. Bitter purging, Magnesiæ sulphas—s. Common, Soda, muriate of—s. Epsom, Magnesiæ sulphas—s. Glauber's, Soda, sulphate of—s. Glauber's, Lymington, see Soda, sulphate of—s. Preston, see Ammoniac carbonas—s. Rochelle, Soda, tartrate of—s. Rock, Soda, muriate of—s. Sea, Soda, muriate of—s. Secondary, Neutral salt—s. Seignette's, Soda, tartrate of—s. Smelting, Ammoniac carbonas—s. of Soda, Soda, subcarbonate of—s. Sorethroat, Potassæ nitras fusus sulphatis paucillo mixtus.

SALT OF TARTAR, see Potassæ subcarbonas, and Potash.

SALT, TASTELESS PURGING, Soda, phosphate of.

SALT OF URINE, Ammoniac carbonas.

SALT, WHITE, Soda, muriate of.

SALT OF WOODSORT, Ammoniac carbonas—s. of Wormwood, see Potash.

SALTANS ROSA, Urticaria.

SALTATIO, (*saltare*, *saltatum*, 'to dance,') Dancing—s. Sancti Viti, Chorea.

SALTERTON. A village about four miles to the eastward of Exmouth, England, frequented as a winter abode by invalids. It is in a small, open valley on the seashore, and is well protected from winds, particularly the northerly.

SALTISH, Halmyrodes.

SALTPETRE, (*sal*, and *petra*, 'a rock,') Potassæ nitras.

SALTUS, gen. *Saltus*, (*saltice*, *saltum*, 'to leap,') Leap, Vultus—s. Tendinum, *Salsultus tendinum*.

SALTWORT, PRICKLY, *Salsola kali*.

SALUBRIOUS, (from *salubris*,) Salutary.

SALUBRIOUSNESS, *Salubritas*.

SALUBRIS, (from *salus*,) Salutary.

SALUBRITY, *Sala'britas*, *Sala'brimness*, [F.] *Salubrité*, (from *salubris*,) Any thing which contributes to the health of individuals or of communities. Situations differ materially from each other in this respect; and the cause of the difference is often an interesting topic of inquiry.

SALUS, gen. *Salutis*, Sanitas.

SALUTAIRE, Salutary.

SALUTARY, *Salutaria*. *Salutar.* *Salubria*.
Hygienicus, *Hygienicus*, *Hygienus*, *Salubritas*.
(F.) *Salutaria*, *Salubre*, (from *salus*) Healthy.
That which is favourable to health. Some diseases have been considered salutary; but this is doubtful.

SALUTIFEROUS, *Salutifer*, *salvus*, and *fero*,
'I bear.' Bringing health; health-bearing.

SALVATELLA, *Vena salvatella*, is situated on the back of the hand, near its inner margin. It begins on the posterior surface of the fingers and dorsal surface of the hand, by a plexus, formed by a number of radicles, frequently anastomosing with each other; it then ascends to the inner part of the forearm, where it takes the name — *arteria cubitalis*. The ancients recommended this vein to be opened in certain diseases, as in melancholic and hypochondriacal affections; and they attributed to such abstraction of blood considerable efficacy in the cure of disease: hence its name, (from *salvere*, 'to be well.')

SALVE (Sax.) *resalp.* (H.) *Salbe.* Unguentum — s. Blistering. Unguentum lynte — s. Deschler's. Ceramum resinae compositum — s. Eye. Unguentum zolli etnei impuri — s. Eye. Singleton's. Unguentum hydragryi nitrooxydi — s. Healing. Cerate of ealamine.

SALVIA. (*From salveo, 'to be well,'* owing to its reputed healing qualities): *S. officinalis* seu *botanica* seu *salpicina* seu *salicaria*, *Eleutherophaca*, *Sage*, *Sage*, (F.), *Sauge*, *Peigne Sauge*, *Thé d'Europe* ou de France. *Ord. Labiatae. Gen. Sage*, *Dianthia Monogyna*. Sage leaves, *Salvia* (Ph. U. S.), have a peculiar, aromatic odour, and a warm, aromatic, bitterish taste. Their virtues depend upon an essential oil. They are tonic, stimulant, and carminative. In the form of a weak infusion, Sage is occasionally used as a diaphoretic, and as a wash for sore mouth. Dose, gr. xv to xxx of tincture.

SALVIA AFRICA'NA, *Will Sage*, a South African species, has the same medical properties as *Salvia*.

SALVIA COLORATA. Salvia.

SALVIA HERNI'NUM. *S. colorata*, *Herni'aria*.
H. colorata seu *nutum*, (F. *Sage-hernia*, of
the Pharmacopœias, is used in similar cases.

SALVIA HOERTEENSIS, *Salvia*.

SALVIA LYRATA, *Wild Sage, Cancerweed*; indigenous. The fresh leaves, bruised, are used to destroy warts. It has been supposed to be of use in cancer.

SALVIA MINOR. *Salvia*—*a.* *Officinalis, Salvia.*

SALVIA SOLA'BREA, *Sola'brea*, *Gallipetrum*, *Æthiopia*, *Blechnum*, *Collis Javia*, *Conium Cist.* (F.) *Sauge solvée*, *Grand sauge*, *Orcule ou Taché*. The leaves have been recommended as emollients and antispasmodics.

SALVIA VIRE, *Asplenium ruta muraria* — s.
Vulgaris, Salvia.

SAMADERA INDICA, *Orb.* Simarubaceæ, furnishes the *Nipa bark*, which is possessed of antiseptic properties.

SAMBO, see Mulatto.

Sambo, and *Zambo*, also signify the offspring of the Indian and Negro.

SAMBUCUS, (from *sambucus*, an ancient musical instrument, said to have been made of its wood.) *Sambucus*, *Sambucus nigra* and *Sambucus racemosa*, Auct. *Infelix hibernica*; *El der*, *E. verna*, *El-leutree*, *E. larnae*, (Sc.) *Bow-tree*, *Bow-tree*, *Bow-tree*, (Prov.) *Bow-tree*, *Bow-tree*, *El-ler*, *El-ler*, *El-ler*, (Scot.) *Sambucus racemosa*, *Sambucus*, *Orf.* *Caprifoliaceae*. *Sex.* *Syst.* *Pentandria Digynia* — (the flower and bark were formerly called *Corn-plum*). The flowers, *Sambucus*, Ph. U. S., berries, and bark of the elder have been used in medicine. The flowers have been reputed diaphoretic and discutient; the berries, aperient and sudorific; the bark, purgative, and in small doses, deob-

struent. The flowers are sometimes used in fermentations, and to form a cooling ointment.

SAMBUCUS ARBOREA, Sambucus — s. Chamæ-
actæ, S. ebulus.

SAMBUCUS EBULES, *S. Chamædore seu helion*
eu h'milla seu heri'cent, *Ed'aba*, *Chamaedore*,
Col'sea, *Dore' el'eb*, *Dumetum*, *F. Helle*, *Yelle*.
 The root, interior bark, leaves, flowers, berries,
 (*Bacca seu Grossa Aro.*) and seeds, have all been
 given in moderate doses as resolvents and cath-
 artics; and in large doses as hydragogues.

The berries of *Sambucus Canadensis*, the *Common Elder* of America, are made official in the Pharmacopœia of the United States—*Sambucus*. A distilled water of the flowers, *Aqua Sambæ*, of *Elder Water*, is official in the London and Edinburgh Pharmacopœias.

SAMBUCUS HELION. Sambucus ebulus — s. Her-
bacea, S. ebulus — s. Humilis, S. ebulus — s. Nigra,
Sambucus.

SAMBUCUS PUBENS, *S. pubescens* sensu *multicaulis*, *Rothschied Elder*; indigenous. The bark, in decoction, is said to be a valuable hydragogue cathartic.

SAMBUCUS RACEMOSA, S. pubens—a. Vulgaris,
Sambucus.

SAMI TER'RA, *Sa'mian Earth*. An argillaceous earth, found in the island of Samos, to which were ascribed virtues similar to those of the Terra Lemnii.

SA'MIUS LAPIS. *Sa'mian stone*. A stone found in the Isle of Samos, which the ancients believed to be astringent and refrigerant.

SAMPHIRE, *Crithmum maritimum*.

SAMPSON, BLACK, *Echinacea purpurea*.
SAMPSON, BLACK, *Echinacea purpurea*.
SAMPSON, BLACK, *Echinacea purpurea*.

SAMPSYCHINON. (*σαμψυχον*, *Sampsychi-*
nion. An ointment composed of *σαμψυχον*, which
was perhaps *majoram*.

SAMPSYCHUM, (σαμψυχον.) *Origanum majorana*.

SAN GIULIANO, MINERAL WATERS OF.
At the foot of the mountain San Giuliano, a league and a half from Pisa, are several mineral springs; temperature from 84° to 107° Fah.

The spring, *Pozzetto*, contains carbonic acid, sulphates of soda, lime, and magnesia; chlorides of sodium and magnesium; carbonates of lime and magnesia; argil and silica.

SANABILIS, *sanare*, 'to heal,' and *habilis*, 'able.' Curative.

SANABLE. Curable.

SANAMUNDA, *Geum urbanum*.

SANATIO, (*sannire, sanationem*, 'to cure,') Curation, Cure.

SANATIVE, Curative.

SANATORIUM, Sanitarium.

SANATORY, Cottage.

SAND, ([Sex.] razz,) see Gravel.

SANDBLIND, nearly blind. A defect of vision, in which small particles seem to fly before the eyes. See Metamorphopsia.

SAND, BRAIN, see Pineal gland — a. Pineal, see Pineal gland.

SANDALUM CÆRULEUM, Guilandina m.
ringa—s. Rubrum, Pterocarpus santalinus.

SAN'DARAC. *Sanda'racina*, *Gum Juniper*, or *San'darach*, *Pommé*, *Gum mi Juné'eri*, *San'darac*, *Véritable résine*, *Résine Juniper*, (F.) *Sandaracum*, *Gomme de Juné'rier*. A name given by the Arabs, to an odorous resin, that flows from the *Thuja oxyphloa*, Ord. Coniferae. See *Spr. Monocia Monadelphica*. It is stimulant, like all the resins; but is very little used.

SANDARACHA GRÆCORUM, (σανδαράχη.)
Realgar.

SANDIVER, *quint-de-verre*, 'loss of glass,' [?]
see Vitruv.

SANDIX, *Plumbi oxydum rubrum*.

SANDYX, (*sandyx*), *Plumbi oxydum rubrum*.

SANG, (from [L.] *sanguis*), Blood, *Panax quinquefolium*—s. *Arterial*, Arterial blood—s. *Dragon*, *Rumex sanguineus*, see *Calamus rotang*—s. *Globules du*, Globules of the blood—s. *Perte de*, Hæmorrhagia.

SANGAREE'. Wine and water, sweetened and spiced. First used in the West Indies.

SANGLOT, (from *singulus*), Sob.

SANGRA'DO, ([S.] *sangrador*, 'a bleeder.') A doctor, in Gil Blas, who was a heroic bleeder; and hence, popularly applied to such a one.

SANGSUE, (from *sanguisuga*), Hirudo.

SANGUIDUCTUS DURÆ MATRIS, (*sanguis*, 'blood,' and *ducere*, ductum, 'to lead,') Sinuses of the dura mater.

SANGUIFIATIO, (*sanguis*, 'blood,' and *facere*, 'to make,') Hæmotosis—s. *Debilior*, Hyphæmatosis.

SANGUIFLUXUS, gen. *Sanguifluxus*, (*sanguis*, 'blood,' and *fluere*, fluxum, 'to flow,') Hæmorrhagia—s. *Uterinus*, Metrorrhagia.

SANGUIMOTION, (*sanguis*, 'blood,' and *movere*, motum, 'to move,') Circulation.

SANGUIMOTORY, Circulatory.

SANGUIN (*Vaisseau*), Bloodvessel.

SANGUINALIS CORRIGIOLA, (from *sanguis*, because used to stop blood,) Polygonum aviculare.

SANGUINARIA, (from *sanguis*, 'blood,' owing to the colour of its juice.) *S. Canadensis* seu *acutalis*, Blood'root, Blood'wort, Puccoon', Red Puccoon', Red Root, Indian Paint, Turmeric. Nat. Ord. Papaveraceæ. Sez. Syst. Polyandria Monogynia. A North American plant, the rhizoma—*Sanguinaria* (Ph. U. S.)—and seeds of which are used in medicine. They are emetic. Dose of the powder, 8 to 20 grains. They are said to reduce the velocity of the pulse, like digitalis.

SANGUINARIA, Polygonum aviculare.

SANGUINE, *Sanguineus*, *Sanguineus*, *Hæmatious*, *Hæmato'des*, *Hæmate'rus*, *Hæmat'nus*, *Hæmo'des*, *Hæmo'des*, *Sanguine'ous*, *Hæmat'ic*, (F.) *Sanguin*, (from *sanguis*.) Bloody, bloodlike; having the colour of blood, as 'a sanguine countenance.' Relating to, or containing blood—as 'sanguine temperament.' Plethoric.

SANGUINEA, Potassæ nitras.

SANGUINENESS, Plethora.

SANGUINEOUS, Plethoric, Sanguine.

SANGUINEUS, Sanguine.

SANGUINIS ABSORPTIO, Hæmorrhophesis—s. Congestio, Congestion—s. Dissolutio, Hæmatolysis—s. Fluor, Hæmoptysis—s. Fluor vesicæ, Hæmorrhagia—s. Ictus, Apoplexy—s. Missio, Bloodletting—s. Profluvium copiosum, Hæmorrhagia—s. Profluvium ex ore, Stomatorrhagia—s. Stagnatio, Hæmostasis.

SANGUINOLENT, *Sanguinolentus*, (from *sanguis*.) Tinged with blood. An epithet applied, particularly, to the sputa when so tinged. Bloody.

SANGUINOSUS, Sanguine.

SANGUIPURGIUM, (*sanguis*, and *purgare*, 'to cleanse,') Purgation of the blood. A name given by some authors to mild fevers, from an idea that they purged the blood of noxious matters.

SANGUIS, gen. *Sanguinis*, Blood, Sperm—s. Concretus, Coagulum—s. Draconis, see *Calamus rotang*—s. Hirci siccatus, see Blood—s. Menstruus, see Menses—s. in Urinâ, Hæmaturia—s. Virginis primum menstruate, see Menses.

SANGUISOR'BA CANADENSIS, *Can'ada Burnett*, (*sanguis*, 'blood,' and *sorbere*, 'to absorb,' so called from its astringency.) A tall indigenous plant—Ord. Rosaceæ—with white, sometimes purple, flowers, which appear from August to October. It has the same virtues as

SANGUISOR'BA OFFICIN'ALIS, *S. car'nea* seu *rubra*, *Ital'ian Pimpinel*, *Pimpinel'la Ital'ica* seu *officin'al'is*, (F.) *Pimpinelle noire*. It was formerly used as an astringent and tonic, in hemorrhages, dysentery, &c.

SANGUISUGA, (*sanguis*, 'blood,' and *sugere*, 'to suck,') Hirudo.

SANGUISUGUM. A barbarous term, to denote too great an accumulation of blood in the heart.

SANICLE, Sanicula, *S. Marilandica*—s. American, *Heuchera cortusa*—s. de *Mariland*, Sanicula Marilandica—s. Yorkshire, *Pinguicula vulgaris*.

SANICULA, (from *sano*, 'I heal,') Saxifraga granulata.

SANIC'ULA, *S. Europæa* seu *officin'arum* seu *mas*, *Astrant'ia diapen'sia*, *Caucas'ia Sanic'ula*, *Cucullat'a*, *Dodecat'heon*, *Symph'ytum petra'rum*. *Diapen'sia*, *D. cortu'sa*, *San'icle*, Wood March, (F.) *Toute-saine*. Ord. Umbelliferae. Sez. Syst. Pentandria Digynia. This herb was formerly used as a mild astringent. It is bitterish, and somewhat austere.

SANICULA EBORACENSIS, *Pinguicula vulgaris*—s. *Europæa*, Sanicula.

SANICULA MARILAN'DICA, *San'icle*, *Ma'ryland San'icle*, *Black Snake'root*, (F.) *Sanicle de Mariland*. This indigenous plant grows in marshes, or low and shaded ground. It is possessed of tonic virtues; and the powdered root is employed in popular practice as a favourite remedy in intermittent fever. It has been recommended, also, in chorea.

SANICULA MONTANA, *Pinguicula vulgaris*—s. *Officinarium*, Sanicula.

SANIDO'DES, (*saidōdes*, 'resembling a board,' from *sais*, gen. *saidōs*, 'a board,' and *odes*.) One whose thorax is flat, like a board.

SANIE, Sanies.

SANIES, *P'chor*, *Ta'bun*, *V'irus*, *Pus malignum* seu *corrosivum*, (F.) *Sanie*, (corr. from *sanguis*, 'blood,') A thin, serous fluid, presenting some of the qualities of pus and blood, and commonly exhaled at the surface of ulcers.

SANIEUX, Ichorous.

SANIOSUS, Ichorous.

SANIOUS, Ichorous.

SANITAIRE, Sanitary.

SANITA'RIUM, *Sanato'rium*, (*sanitas*, 'health,') A retreat selected for valetudinarians, in consequence of its salubrity. Also, an establishment for the treatment of the sick.

SANITARY, *San'atory*, (F.) *Sanitaire*. That which relates to health, and especially to public health; as 'Sanitary commission.'

SANITAS, (*sonus*, 'healthy,') *Hygie'a*, *Hyge'a*, *Hyggian'sis*, *Hygi'asis*, *Integ'ritas*, *Prosper'itas*, *Valetu'dinis*, *Sal'us*, *Hygie'a*, *Valetu'do*, *Valitu'do*, *Valetu'do bona* seu *secun'da* seu *com'moda*, *Hyge'a*, *Euzoödyna'mia*, *Health*, (Sc.) *Heil*, *Heyle*, *Heal*, (F.) *Sanité*, (from *sonus*, 'sound,') Health has been defined, 'a state in which all the functions are exerted with regularity and harmony. A person may, however, be in health, although certain functions are not performed. A blind man, for example, has health when all his functions, except sight, proceed with regularity. *Sanity* is often used for soundness of mind.

SANITAS, Cure.

SANITY, Sanitas.

SANKIRA, Smilax China.

SANS PAIRE, 'pairless,' (*Veine*), Azygos vein.

SAN'TA CRUZ, CLIMATE OF. This West India island is frequently selected as a winter retreat for the physical valetudinarian from the United States; but it does not seem entitled to any preference, in such cases, over the other islands of the West Indies, whilst its general salubrity is said to be less than that of many.

SANTAL BLANC, *Santalum album*—s. Rouge, *Pterocarpus santalinus*.

SANTALUM, see *Pterocarpus santalinus*.

SANTALUM ALBUM, *S. citrinum* seu *pallidum* seu *myrtifolium*, *S. indicum myrtifolium*, *Yellow Saunders*, (F.) *Santal blanc*. Ord. Santalaceæ. See *Syst. Tetrandria Monogynia*. The virtues of the wood of the white saunders reside in a volatile oil and resinous principle. The odour is fragrant, and resembles a mixture of ambergris and roses. It has been used as a stimulant and sudorific, but is not now employed. The saunders woods are called, collectively, in French, *Santau*.

SANTALUM MYRTIFOLIUM, *S. album*—s. Rubrum, *Pterocarpus santalinus*.

SANTÉ, Sanitas.

SANTONICA, see *Artemisia santonica*.

SANTONICUM, (from *Santonia*, its native place), *Artemisia santonica*.

SANTONINE, *Santoninum*, *San'tonin*, *Ci'nin*. A peculiar crystalline substance obtained from *Artemisia Santonica*. It is directed, in the Ph. U. S. for 1863, to be prepared by digesting *santonica*, lime and alcohol together, and expressing; distilling off the alcohol and evaporating; adding acetic acid until in slight excess; washing the resulting crystalline mass, and drying it; boiling the dry residue with alcohol; and, having digested the tincture for several hours with animal charcoal, filtering, and setting aside to crystallize. *Santonine* possesses the anthelmintic properties of the plant. Dose, five or six grains.

SANTORINI, FIS/SURES OF, (F.) *Incisures ou Scissures de Santorini*, (after the Italian anatomist.) Clefts at the upper and back of the fibro-cartilaginous portion of the external ear.

SANTORINI, TUBERCLES OF. Small cartilaginous projections at the top of the arytenoid cartilages, which support the ligaments of the glottis. See *Corniculum Laryngis*.

SANYÉ, *Sinapis*.

SAPRIA, *Sauar'ja*, *Sau'ri*, (a Guiana name.) The ripe and dried fruit of *Moe'sa* seu *Bœob'otrys picea*, an Abyssinian tree, which is administered in its native country, in powder, in doses of from 32 to 44 grammes, as a tæniacuge.

SAP, ([Sax.] rap, [G.] Saft,) *Arborum Succus* seu *Lymph'a*, (F.) *Sève*. The circulating fluid of the vegetable, which corresponds to the blood of the animal.

SAPA, see Rob.

SAPHA'DA. Paracelsus has given this name to reddish scales adherent to the hair in some eruptions of the hairy scalp.

SAPHENA, *Saphena*.

SAPHATUM, *Asaphatum*.

SAPHENA, *Saphēna*, (*σαφης*, 'manifest,' 'evident,') (F.) *Saphène*. A name given to two subcutaneous veins of the lower limb. These are: 1. The GREAT SAPHENA VEIN, *S. inter'na*, *Tibio-malleolar* (Ch.), (F.) *Grande Veine Saphène ou Saphène interne*. Its radicles occupy the inner part of the great toe, and form a transverse arch on the back of the foot, which unites with the saphena externa, and receives at its convex part, which is turned forwards, a number of branches, coming from the toes. These radicles, when united, ascend before the inner ankle, and form a trunk, the size of which is farther increased by several branches that proceed from the region of the tarsus, and that of the metatarsus. This trunk ascends vertically at the inner part of the leg; passes behind the inner condyle of the femur; proceeds along the inner part of the thigh; receives other subcutaneous veins, such as the abdominal subcutaneous, the external pudic, &c., and passes through the inferior opening of the crural canal to empty itself into the femoral vein. 2. The LES'SER VE'NA SAPHENA or *Saphēna ex-*

ter'na, *Peroneo-malleolar Vein* of Chaussier, (F.) *Petite Veine Saphène ou Saphène externe*, commences by several roots, which are spread over the dorsum and outer side of the foot, and embrace the outer ankle. Behind this, they unite into a single trunk, which ascends obliquely, approaching the tendo Achillis, and proceeds vertically towards the ham to open into the popliteal vein.

SAPHÈNE, *Saphena*.

SAPHENOUS, *Saphenous*; (from *saphena*.) Of or belonging to the saphena.

SAPHENOUS NERVES are two in number, and are branches of the crural or femoral nerve. They are the short *saphenous*—*ner'vus cuta'neus internus*; and the long *saphenous*—*ner'vus cuta'neus internus longus*.

The external *saphenous*, formed by the union of the communicans poplitei and communicans peronei, passes down the leg by the side of the external saphenous vein of the foot.

SAPHENOUS OPENING. An oval, and somewhat oblique, opening in the fascia lata, a little below Poupart's ligament, for the transmission of the principal superficial vessels. Here, likewise, femoral hernia descends.

SAPID, *Sap'orous*, *Sap'idus*, (F.) *Savoureux*, (*sapere*, 'to taste,') Possessed of sapidity. Having the power of affecting the organs of taste.

SAPID'ITY, *Sap'idness*, *Sap'orosity*, *Sap'or*, (F.) *Sapidité*. The quality of bodies that gives them taste.

SAPIN COMMUN, ([L.] *Sapinus*), *Pinus picea*.

SAPIN'DUS, (*sapo*, and *Indus*, 'Indian soap,') *S. Saponaria*, *Saponaria*, *Soaptree*, (F.) *Savonnier commun*. Ord. Sapindacæ. The name of the plant that affords the *Soapnuts*, *Saponaria Nucula*, *Bac'æe Bermuden'ses*, *Soapberries*. A spherical fruit about the size of a cherry, the cortical part of which is yellow, glossy, and so transparent as to show the spherical black nut that rattles within, and includes a white kernel. The cortical part of this fruit, which has similar effects with soap in washing, has been recommended in chlorosis, and as a stomachic and deobstruent.

SAP'O, gen. *Sapo'nis*, (*σαπων*, [Sax.] *rape*), *Smegma*, (F.) *Sapon*. A product obtained by treating fatty bodies with caustic alkalies dissolved in water. In the Pharmacopœia of the United States, it means Soap made with soda and olive oil. See *Sapo medicinalis*.

YEL'LOW SOAP is formed of turpentine, oil, and alkali. A soap of subcarbonate of potassa and oil of turpentine—*Sapo terebin'thinæ*—was formerly used for indolent swellings, under the name of *Starkey's soap*.

Castile Soap, being made of fine olive oil, is generally used internally. The yellow and soft soaps are the most stimulating; and are, consequently best adapted for particular cutaneous eruptions.

Internally, soap is purgative and diuretic; externally, it is detergent and stimulant. Soapy water is used with the greatest success, in cases of poisoning by acids. It is mostly applied externally to bruises and sprains. In Pharmacy, it is used to favour the solution of resins.

SAP'O AC'IDUS, *Ac'id Soap*, is a combination of a fixed oil with an acid.

SAP'O AMMONIACALIS, *Linimentum ammoniæ fortius*—s. *Ammoniæ oleaceus*, *Linimentum ammoniæ fortius*—s. *Amygdalinus*, *Sapo medicinalis*—s. *Cacaotinus*, *Sapo coconeus*—s. *Calcarius*, *Linimentum aquæ calcis*—s. *Castiliensis*, see *Sapo medicinalis*.

SAP'O COCO'NEUS, *Sapo cacaōit'nis*, *Soap of Coconut Oil*, (F.) *Sapon de cacao*, has been much used in Germany in herpetic affections.

SAPO DURS, *S. medicinalis*—*s. Hispanicus*, see *S. medicinalis*—*s. Kalicus*, *S. Mollis*.

SAPO LIQ'VIDUS, *Liq'uid Soap*, *Lo'tio saponacea*, is a cosmetic, which is composed of *Olive oil* $\frac{3}{4}$ iv, *Ol. tartari per deliquium* $\frac{3}{4}$ ss. Rub together, and add *Aq. rosarum*, $\frac{3}{4}$ xij.

SAPO MEDICINALIS, *Sapo amygdalinus*, *Medicinal Soap*, (F.) *Sapon medicinalis*, is made of oil of sweet almonds and half its weight of caustic alkali, or like the *Sapo Hispanicus seu Castilienus*, Spanish or Castile soap, *Sapo durus*, *Sapo* (Ph. U. S.), (F.) *Sapon dur*, of olive oil and soda.

SAPO MOLLIS, *S. viridis seu niger seu kalicus*, *Common or Soft Soap*, (F.) *Sapon mou ou noir*, is made of potash and oil or tallow. The *Sapo vulgaris* or 'common soap' of the Pharm. U. S. is made with soda and animal oil.

SAPO NIGER, *Black Soap*, *Melanomagma*, is a composition of train oil and an alkali: and **GREEN SOAP**, *Sapo viridis*, of hempseed, linseed, or rape oil, with an alkali. See *Sapo mollis*.

SAPO TEREBINTHINÆ, see *Sapo*—*s. Vegetabilis*, *Pulvis gummosus alkalinus*—*s. Viridis*, *Sapo mollis*—*s. Vulgaris*, see *Sapo mollis*.

SAPONACEOUS, *Saponaceus*, (F.) *Saponact*, (from *sapo*.) Soapy. Containing soap, as *Linimentum saponaceum*, *Soap liniment*.

SAPONAIRE, *Saponaria*.

SAPONARIA, (from *sapo*,—the juice forming a lather with water.) *Sagewort*, *Brusewort*, *Bouncing Bet*, *Struthium*, *Lana'ria*, *Lych'nis officinalis seu sylvestris*, *Ibix'uma*, *Boot'ia vulgaris*, (F.) *Saponaire*, *Savonnère*. Ord. Caryophyllaceæ. *Sex. Syst.* Decandria Digynia. A decoction of the leaves of this plant has been recommended in scrofula, cancer, old syphilis, cutaneous eruptions, jaundice, visceral obstructions, &c. It is hardly ever used.

SAPONARIA, *Sapindus saponaria*.

SAPONARIE NUCULÆ, see *Sapindus saponaria*.

SAPONÉ. Soap, to which are added substances capable of communicating to it new properties, without affecting its own.

SAPO'NEA. A pectoral medicine, prepared with sweetened syrup of violets, and oil of sweet almonds.

SAPONULÉ. A tincture (*alcoolé*), sufficiently charged with soap to cause it to assume a gelatinous shape.

SAPONULES, *Sapon'uli*, (F.) *Savonules*. Imperfect soaps formed by combinations of volatile or essential oils with different bases. See *Opo-dodech*.

ACID SAPONULES are combinations of such oils with different acids.

SAPONURE. A medicine, formed of soap, in powder, and of resinous or extractive matters.

SAPOR, Sapidity, Savour.

SAPOROSITY, Sapidity.

SAPOROUS, Sapid.

SAPOTA, *S. Acharas*, *Ac'leras Sapo'ta seu australis seu Zapo'ta*, *El nispero*, *Onalfruited Sapota*, (F.) *Sapotier*, *Sapotillier*, *Sapotilla*, *Fam. Sapotaceæ*. *Sex. Syst.* Hexandria Digynia. A tree which grows in the Antilles, and particularly at St. Domingo. Its fruit is much esteemed. It resembles the marmalade of quinces; and hence has been called *natural marmalade*. The seeds make an emulsion, which has been given in nephritic affections. The bark has been recommended as a febrifuge.

SAPOTA ACHARAS, *Sapota*.

SAPOTIER, *Sapota*.

SAPOTILLA, *Sapota*.

SAPOTILLIER, *Sapota*.

SAPROPYRA, (*σαρπός*, 'putrid,' and *πύρ*, 'fever,') *Typhus gravior*.

SAPROS, (*σαρπός*) Putrid, Rancid.

SAPROS'TOMUS, (*sapros*, and *στόμα*, 'mouth,') One who has an offensive breath.

SAPROTÆS, (*σαρπότης*) Putrefaction.

SARAPUS, *Sarapous*, ('σαρῶ, 'I grin,' and *πούς*, 'the foot,') One who has a large foot; whose toes are largely separated, or whose feet are much asunder in walking. See *Kylosis*.

SARATOGA, **MINERAL WATERS OF**. The Saratoga Springs in New York are ten miles north-east from Ballston. They contain the same ingredients as those of the latter place; and are much frequented by visitors from every part of the United States. The following analysis has been given of the *Congress spring*. In a wine gallon—*gaseous contents*:—carbonic acid, 311 cubic inches; atmospheric air, 7. *Solid contents*: chloride of sodium, gr. 385; iodide of sodium, gr. 3.5; bicarbonate of soda, gr. 8.982; bicarbonate of magnesia, gr. 95.788; carbonate of lime, gr. 98.098; carbonate of iron, gr. 5.075; silica, gr. 1.5; bromide of potassium, a trace. Total, gr. 597.943.

The *Walton* or *Iodine spring* contains chloride of sodium, carbonate of magnesia, and iodide of sodium.

SARCEPIPOCELE, *Sarcoëpiplocele*.

SARCEPILOMPHALOCELE, *Sarcoëpiplomphalocèle*.

SARCEPILOMPHALUS, *Sarcoëpiplomphalus*.

SARCEUX, *Sarcous*.

SARCIDIUM, (*σαρκίδιον*, dim. of *σαρξ*, 'flesh,') *Caruncle*.

SARCINA, 'a pack, a burthen,' *Fœtus*.

SARCINA VENTRICULI, *S. Goodenii*, *Merismopœ'dia ventriculi*, (Robin.) A microscopic cryptogamous plant found in the fluid of water-brash.

SARCION, (*σαρκιον*), *Caruncle*.

SARCITES, (from *σαρξ*, 'flesh,') *Anasarca*—*s. Flatusos*, *Emphysema*.

SARCITIS, (*σαρξ*, 'flesh,' and *itis*), *Myositis*.

SARCO, (*σαρξ*, gen. *σαρκος*, 'flesh,') in composition, flesh.

SARCOCE'LE, *Her'nia carno'ea*, *Scirr'hus Testic'uli*, (*sarco*, and *κήλη*, 'a tumour,') *Sarco'ma scro'ti*, (F.) *Hernie charnue*. Scirrhus or cancer of the testicle. This disease affects adults particularly; and appears most commonly after an inflammatory swelling of the testicle. Sometimes it is dependent upon a blow; at others, it makes its appearance without any appreciable cause. It is well known by a hard, heavy, ovoid or spherical swelling of the testicle, which is, at first, slightly, or not at all painful, and merely causes an unpleasant traction on the spermatic cord. There is no heat or change of colour of the skin; the spermatic cord is swollen, and participates in the affection; very painful shootings occur; the lymphatic glands of the abdomen become swollen, and form a tumour, which may, at times, be felt through the abdomen; and the patient, at length, dies with every sign of the cancerous diathesis. The prognosis is very unfavourable. The only means, indeed, that can save life, is the extirpation of the testicle.

SARCOCOLLA, (F.) *Colle-chair*, (*σαρκοκόλλα*, from *sarco*, and *κόλλα*, 'glue,') A resinous matter, obtained from *Penæ'a Sarcocol'la*, *P. mucronata*, and other species of *Penæa*, an African shrub, which has been so called from a belief that it facilitates the consolidation of flesh. It is in the form of small, oblong, semitransparent globules, of a yellowish or reddish hue, and smells like aniseed. It was once employed as an astringent and detergent.

SARCOCOPTES, (*sarco*, and *κοπτω*, 'to wound,') see *Psora*.

SARCODES, (from *σαρξ*, 'flesh,' and *odes*), *Car-neous*.

SARCOËPIPOCELE, *Sarcepiploce*'le, (*sarco*, επιπλοον, 'the epiploon,' and κηλη, 'a tumour.') Epiplocele, complicated with sarcoma or sarcocele.

SARCOËPIPLOM'PHALUS, *Sar epiplo'm'pha-lus*, *Sarcepiplo'mphaloc*'le, (*sarco*, επιπλοον, 'the epiploon,' and ομφαλος, 'the navel.') Umbilical hernia, formed by scirrhus epiploon or complicated with sarcoma.

SARCOHYDROCELE, *Hydrosarcocele*.

SARCOID, *Sarcoï'des*, (*sarcoeides*, from *sarco*, and αἶδος, 'resemblance.') Resembling flesh.

SARCOLEM'MA, gen. *Sarcole'm'atis*, *Sarcole'm'a*, (*sarco*, and λεμμα, 'a coat.') The sheath which surrounds the fibrils of muscle that form a fibre. It is quite distinct from the areolar membrane that binds the fibres into fasciculi. See *Perymisium*.

SARCOLOGIA, (*sarco*, and λογος, 'a discourse,') Myology, Sarcology.

SARCOLOGY, *Sarcolog*'ia. The part of anatomy which treats of the soft parts. It comprises myology, angiology, neurology, and splanchnology.

SARCOMA, gen. *Sarco'm'atis*, (*sarcoma*, *Emphy'ma Sarco'm'a*, *Sarcem'atous tumour*, *Sarco'm'eis*, *Sarcophy'tis*, *Escarco'm'a*, *Escarco'm'a*, *Tu'mor car'neus*, *Escarcescent'ia carno'sa*, *Polyypus carno'sus*, (σαρξ, 'flesh,' and ομα.) Any species of excrescence having a fleshy consistence.

SARCOMA, **CARCINOMATOUS**, Scirrhus—s. *Cercotis*, see *Cercosis*—s. *Epulis*, *Epulis*—s. *Gelatinous*, see *Tumour*, *fibrocclular*—s. *Mastoid*, *Mammary sarcoma*—s. *Medullary*, see *Encephaloid*—s. *Medullary*, see *Encephaloid*, *Hæmatodes fungus*—s. *Pulpy*, *Hæmatodes fungus*—s. *Scroti*, *Sarcocele*.

SARCOMA, **COMMON VASCULAR OR ORGANIZED** of Abernethy, *Emphy'ma Sarco'm'a Vasculo'sum*, is vascular throughout: texture simple; when bulky, napped on the surface with arborescent veins. Found over the body and limbs generally.

SARCOMATEUX, *Sarcomatous*.

SARCOMATODES, *Sarcomatous*.

SARCOMATOSUS, *Sarcomatous*.

SARCOMATOUS, *Sarcomato'sus*, *Sarcoma-to'des*, (F.) *Sarcomateux*. Belonging to, or having the characters of sarcoma.

SARCOMPHALUM, (*sarκοφαλον*.) *Sarcomp'h'alus*.

SARCOMPHALUS, *Sarcom'ph'alum*, (*sarco*, and ομφαλος, 'navel.') A scirrhus or fleshy tumour, developed at the navel.

SARCOMYCES, (*sarco*, and μυκης, 'fungus,') *Sarcopongus*.

SARCOPHAGUS, (*sarκοφαγος*, from *sarco*, and φάω, 'to eat,') Carnivorous, Cathartic.

SARCOPHYIA, (*sarco*, and φωω, 'to grow,') Excrescence, Sarcoma.

SARCOPSYLLA PENETRANS, (*sarco*, and ψύλλα, 'a flea,') *Chique*.

SARCOPE, *Acarus*.

SARCOPTES SCABIEI, (*sarco*, and σκαπω, 'to wound,') see *Psora*.

SARCOPYODES, (*sarco*, πους, 'pus,' and οδεις.) Having the appearance of pus mixed with flesh. An epithet given to certain excreted matters, and particularly to the sputa of those labouring under consumption.

SARCO'SIS, (*sarκοσις*, from σαρξ, gen. *sarκος*, 'flesh,' and οσις.) The formation of flesh. Also, sarcoma.

SARCOSIS BULBI, *Eosphthal'mia fungo'sa seu sarcomat'ica*. A fleshy growth from the globe of the eye.

SARCOSIS UTERI, *Metrauxie*.

SARCOSPONGUS, *Sarcom'yces*, (*sarco*, and σπογγος, 'fungus,') A fleshy fungus.

SARCOSTOSIS, (*sarco*, and οσσις, 'formation of bone,') *Osteosarcoma*.

SARCOTHLASIS, *Sarcot'hl'as'm'a*, (*sarco*, and

θλαω, 'I bruise,') *Cu'ro conta'sa*. A bruise of the flesh.

SARCOTIC, (*σαρκωτικός*.) Incarnas.

SARCOTRIPTER, (*sarco*, and τριβω, 'to rub down,') *Écraseur*.

SARCOTRIPSIE, see *Écraseur*.

SARCOUS, (F.) *Sarceus*, (from σαρξ, gen. *sarκος*, 'flesh,') Of, or belonging to, flesh or muscle—as 'sarcous tissue.'

SARCOUS ELEMENTS, *Primitive particles*. The elementary particles, which, by their union, form the mass of muscular fibre.

SARDIASIS, (from Σαρδια, *Sardinia*, because the *Sardinia larva*, which grew there, was said to produce it,) Canine laugh, *Risus sardonius*.

SARDONIA HERBA, *Ranunculus sceleratus*.

SARDONIASIS, (same etymon as *Sardiasis*.) Canine laugh, *Risus sardonius*.

SARDONIUS RISUS, Canine laugh.

SARE, *Essera*.

SAROTHAMNUS SCOPARIUS, (*sarpos*, 'a broom,' and *scopos*, 'a shrub,') *Spartium scoparium*.

SARRACENIA, (after Dr. Sarrazin, of Quebec,) *Sidenaddle plant*, *Flytrap*. The root of *Sarracenia flvea* and *S. variol'ris* is bitter and astringent, and has been used in South Carolina, as a domestic remedy in dyspepsia. The root of *Sarracenia purpurea* or *pitcher plant*, *Indian* or *Huntsman's cap*, is used, in infusion, by the Indians of the North, as a remedy for smallpox; both in the way of prevention and cure.

SARRASIN, (after the Saracens,) *Polygonum fagopyrum*.

SARRETE, *Trismus infantum*.

SARRIETTE, *Satureia hortensis*—s. *de Crête*, *Satureia capitata*.

SARSA, *Smilax sarsaparilla*.

SARSAPARILLA, ([S.] *Zerza*, 'a briar,' and *parilla*, 'a little vine,' or, according to some, from *Parillo*, a physician who first employed it.) *Menispermum Canadense*, *Smilax S.*

SARSAPARILLA FALSE, *Aralia nudicaulis*.

SARSAPARILLA GERMANICA, *God'men maj'or seu ru'brum*, *Salsaparilla Germanica*, *Cu'res Are-na'ria*, (F.) *Loiche*, *L. des Sables*, *Salsapareille d'Allemagne*, *Chiendent rouge*. Ord. *Cyperaceæ*. Sec. *Sygt*. *Monoclea Triandria*. A plant, which grows plentifully on the seacoasts of Europe. It has been recommended in some mucous affections of the trachea, in rheumatism, gout, &c.

SARSAPARILLA INDIAN, *Hemidesmus Indicus*—s. *Wild*, *Aralia nudicaulis*.

SARTAPARILLA, *Smilax sarsaparilla*.

SARTORIUS, (*sartor*, 'a tailor,' because used in crossing the legs.) *Sut'orius*, *Longi'scimus Fem'oris*, *Foveal'is*, *F. Longus seu sut'orius*. (F.) *Ilio-cresti'al'is*, *Ilio-péit'al* (Ch.), *Conturvier*. A muscle situate at the anterior part of the thigh. It is long, small, and flattened, and extends obliquely, from the anterior and superior spine of the ilium to the superior and inner part of the tibia. This muscle can, also, bend the pelvis on the thigh, and conversely.

SARX, (σαρξ, gen. *sarκος*.) *Flesh*, *Pulp*.

SARZA, ([S.] *sarza*, 'a briar,') *Smilax sarsaparilla*.

SASSA GUM. A gum, occasionally carried into the European markets from the East, and answering exactly to Bruce's description of the product of a tree, which he calls *Sassa*. It has considerable resemblance to Gum Tragacanth.

SASSAFRAS, *Laurus sassafras*, *Noctandra cym-barum*—s. *Medulla*, see *Laurus sassafras*—s. *Nut*, see *Pichurin beans*—s. *Radiceis cortex*, see *Laurus sassafras*.

SASSAFRAS SWAMP, *Magnolia glauca*.

SASSAFRAS TEA, *Saloop*.

SASSY BARK. A poisonous bark of a large tree—Genus, *Erythrophle'm*, *E. Guineen'se*; *Nat. Ord. Fabaceæ*; *Radicater tree* of Sierra Leone—

growing in Western and Central Africa, used in an ordeal to which persons, suspected of witchcraft, secret murder, and other crimes, are subjected, to test their innocence or guilt. Hence it is called, in Congo, *ordeal bark*, and in Ashantee and Liberia, *doom bark*.

SAT'ELLITE, *Satel'les*, gen. *Satel'litis*. That which guards. That which is placed near.

SATELLITE VEINS are those that accompany arteries.

SATHE, (σαθη,) Penis.

SATIAs, Satiety.

SATIES, Satiety.

SATIETAS, Plesmone, Satiety,

SATIETY, *Sati'etas*, *Satu'ritas*, *Sat'ias*, *Sat'ies*, *Satura'tio*, *Asē*, *Ple'thos*, *Ples'monē*, (*satiare*, from *satis*, 'enough,' 'to give enough,') Disgust for food; commonly produced by repletion.

SATIN, *Lunaria* rediviva.

SATURANS, (*saturare*, *saturatum*, (*satis*, 'enough,') 'to satiate,') Absorbent (medicine.)

SATURATIO, Satiety, Saturation.

SATURATIO, *Satura'tio*. The state of a compound in which its elements are combined in such proportion that a fresh quantity of either cannot be added without producing excess. When an acid or an alkali is directed to be saturated, the saturation is determined by litmus and turmeric.

SATUREIA CAPITATA, (from Σατρυς, 'the Satyrs,' owing to its presumed power of exciting to venery.) *Cili'ated Sa'tory*, *Thy'mus*, *T. sylves'tris* seu *capita'tus* seu *Cre'ticus*, *Cuni'lla*, (F.) *Sarriette* de Crête. *Ord.* Labiate. *Sex. Syst.* Dilynamia Gymnospermia. Possesses similar properties to thyme.

SATUREIA HORTENSIS, *Satureia*, *S. sati'va*, *Cuni'lla sati'va* *Plin'ii*, *Thym'bra*, *Sum'mer Sa'tory*, (F.) *Sarriette*. An aromatic plant, cultivated for culinary purposes. It has been used, as an excitant, in anorexia, anaphrodisia, &c.

SATUREIA ORIGANOIDES, *Cunila mariana* — *S. Sativa*, *S. hortensis*.

SATURITAS, (*saturare*, 'to satiate,') Satiety.

SATURNI DULCEDO, *Plumbi superacetes*.

SATURNINE, *Saturni'nis*, (from *saturnus*),

Containing, or caused by, lead.

SATURNINE BREATH. The peculiar odour of the breath in one labouring under Saturnism.

SATURNISMUS, *Intoxica'tio* seu *Cachex'ia* seu *Dyscras'ia saturni'ni*, *Molybdocachex'ia*, (F.) *Maladie de plomb*, (from *saturnus*.) Poisoning by lead; lead-poisoning; saturnine cachexy.

SATURNUS, *Plumbum*.

SATYRIASIS, (*σατυριασις*, from Σατρυς, 'a Satyr,' because the Satyrs were greatly addicted to venery.) *S. fu'rens*, *Satyras'mus*, *Satyris'mus*, *Estroma'nia*, *Tenti'go vene'rea*, *Salac'itas*, *Brach'u'na*, *Lugne'sis Salac'itas*, *L. fu'ror masculi'nus*, *Lagne'ta*, *Lagni'ta*, *Laguo'sis*, *Lusciv'ia*, *Lusciv'itas*, *Tenti'go vere'tri*. An irresistible desire, in man, to have frequent connexion with the female, accompanied by the power of doing so without exhaustion. The causes are commonly obscure. Sometimes, the abuse of aphrodisiacs has occasioned it. The principal symptoms are:—almost constant erection; irresistible and almost insatiable desire for venery; frequent nocturnal pollutions. Cold lotions; the cold bath; a mild diet; active exercise, &c., are the only means that can be adopted for its removal.

SATYRIASIS FURENS, *Satyriasis*.

SATYRIASMUS, *Satyriasis*.

SATYRION, (*σατυριον*), *Orchis mascula*.

SATYRISMUS, (*σατυριασμος*), *Satyriasis*.

SATYRIUM, *Goodyera pubescens*.

SATARJA, *Saoria*.

SAUCE ALONE, *Alliaria*.

SAUCE, GREEN, *Rumex scutatus*.

SAUCH, ([Sax.] *ralh*), *Salix*.

SAUERKRAUT (G.), 'sour cabbage,' *Sour crout*, (F.) *Chou-croûte*. A culinary preparation, made with chopped cabbage, left to attain an incipient state of acetous fermentation. It is reputed by the Germans to be salubrious, easy of digestion, and eminently antiscorbutic.

SAUGE, *Salvia* — *s. des Bois*, *Teucrium scordium* — *s. Grande*, *Salvia sclarea* — *s. Hormin*, *Salvia horminum* — *s. de Jérusalem*, *Pulmonaria* — *s. Petite*, *Salvia* — *s. Sclérée*, *Salvia sclarea*.

SAUGH, ([Sax.] *ralh*), *Salix*.

SAULE, *Salix*.

SAUMURE, (from *sal*, and *muria*), *Muria*.

SAUNDERS, RED, *Pterocarpus santalinus* — *s. Tree*, *Pterocarpus santalinus* — *s. Yellow*, *Santalum album*.

SAURA, (*σαυρα*), *Lizard*.

SAURURUS CERNUUS, (*saura*, 'lizard,' and *ovpa*, 'a tail,') *Liz'ard's Tail*, *Breast weed*. Indigenous; *Order*, *Saururaceæ*; flowering in June. It has been used in lumbago. The bruised root is emollient, and is a popular application to inflamed breasts.

SAUSAGE POISON, *Allantotoxium*.

SAUT, ([L.] *saltus*), *Leap*.

SAUTEUSES, 'leapers,' see *Convulsionnaire*.

SAUVE-VIE, 'save life,' *Asplenium ruta muraria*.

SAVE, *Salvia*.

SAVEUR, *Savour*.

SAVINE, *Juniperus sabina*.

SAVINIER, *Juniperus sabina*.

SAVON, *Sapo* — *s. Ammoniacal*, *Linimentum ammoniac fortius* — *s. Cacao*, *Sapo coconus* — *s. Calcaire*, *Linimentum aquæ calcis* — *s. Dur*, *Sapo durus* — *s. Médicinal*, *Sapo medicinalis* — *s. Mou*, *Sapo mollis* — *s. Noir*, *Sapo mollis*.

SAVONNIER COMMUN, (from *savon*), *Sapindus saponaria*.

SAVONNIÈRE, *Saponaria*.

SAVONULE, *Saponulus*.

SAVONULUS, *Saponulus*.

SA'VOR, *Sa'por*, *Taste*, (F.) *Saveur*. Quality of bodies, by which they act upon the sense of taste. Chymists, at times, avail themselves of this quality to analyse an unknown substance.

SA'VORY, *Sap'idus*, (F.) *Savoureux*. An epithet given to bodies that have taste; and particularly to those that excite a very agreeable impression on the organ of taste.

SAVORY, *CILIATED*, *Satureia capitata* — *s. Summer*, *Satureia hortensis*.

SAVOREUX, *Savory*.

SAVOYAN, *Galium verum*.

SAW, *Pr'i'on*, *Ser'ra*, ([Da.] *Sawe*, [Sax.] *raga*), (F.) *Scie*. A surgical instrument, made like a common saw; and used for sawing the bones in amputations, or for removing exostoses, &c. Saws of different sizes and shapes are used in surgery. The crown of the trepan is nothing more than a species of circular saw.

SAW, AMPUTATION, *Ser'ra amputato'ria*. The saw used in amputation.

SAW, HEY'S. An instrument used by Mr. Hey, of Leeds, in fractures of the cranium. It consists of a long handle, to which a small saw with a convex or straight edge is attached, and by which a piece of bone of any shape may be removed. With the trephine, the saw must always be circular.

SAW PALMETTO, *Chamærops serratula* — *s. Wort*, *Liatris*.

SAXIFRAGA, (*saxum*, 'a stone,' and *frango*, 'I break,') *Pimpinella saxifraga* — *s. Anglica*, *Peucedanum silaus*. See *Calcifraga*.

SAXIFRAGA CRASSIFOLIA. *Ord.* Saxifragaceæ. *Sex. Syst.* Decandria Dignia. The root of this

species of saxifrage has been extolled by Pallas, as preëminently antiseptic.

SAXIFRAGA GRANULATA. *S. alba*, *Saxicula*, *Sedum*, *White saxifrage*. (F.) *Saxifraga granulata*. Recommended as diuretic and lithontripic; but not used.

SAXIFRAGA RUBRA, *Spiræa filipendula*—s. *Vulgaris*. *Peucedanum silaus*.

SAXIFRAGE. BURNET, *Pimpinella saxifraga*—s. *Granulata*, *Saxifraga granulata*—s. *Meadow*, *Peucedanum silaus*—s. *White*, *Saxifraga granulata*.

SAXIFRAGUS. (*saxum*, 'a stone,' and *frango*, 'I break,') *Lithontripic*.

SAXONICUS PULVIS. A powder, prepared with the roots of the wild and the cultivated angelica; the asclepias vincetoxicum, &c. It was considered formerly as an antidote.

SCAB, *Scabies*, *Crusta*, *Scall*, (Prov.) *Rove*, (F.) *Croûte*. An incrustation, which forms upon a sore, owing to the concretion of the fluid discharged from it. An eschar.

SCABBY or SCABBED. Covered or diseased with scabs.

SCABIES, (*scabere*, 'to scratch,') *Psora*—s. *Agria*, *Lichen*—s. *Capitis*, *Porrigio lupinosa*, *Porrigio scutulata*—s. *Capitis favosa*, *Porrigio favosa*—s. *Crustosa*, *Radzyge*—s. *Fera*, *Ecthyma*.

SCABIES FERINA, *Itch of animals*, *Mange*, (pronounced *Mänge*.) A cutaneous disease, which affects almost all domestic animals, but especially the horse, sheep, dog, and cow. It is said to have been transmitted to man; but this is questionable.

SCABIES FERINA, *Psoriasis*—s. *Norwegica* *Boeckii*, *Radzyge*—s. *Papuliformis*, *Prurigo*—s. *Sicca*, *Lichen*, *Psoriasis*—s. *Venerea contagiosa*, *Scherlievo*.

SCABIEUSE ORDINAIRE, *Scabiosa*.

SCABIOLA, *Scabies*.

SCABIOSA, (*scaber*, 'rough,' owing to its rough hairy surface,) *S. arven'sis*, *Triche'ra arven'sis*, *Field Scabious*, (Sc.) *Elaestingwort*. (F.) *Scabieuse ordinaire*. *Ord. Dipsacæ*. *Sec. Syst. Tetrandria Monogynia*. The plant is bitter and substringent, and was formerly used in the cure of leprous affections and diseased lungs.

SCABIOSA ARVEN'SIS, *Scabiosa*—s. *Carduifolia*, *Echinops*.

SCABIOSA SUCCI'SA, *Succi'sa*, *S. pratensis*, *Asteroccephalus succi'sa*, *Morsus Diaboli*, *Devil's bit*, (F.) *Mors du Diable*, has similar properties.

SCABIOSUS, *Erigeron Philadelphicus*—s. *Field*, *Scabiosa*—s. *Sweet*, *Erigeron heterophyllum*.

SCABISH, *Oenothera biennis*.

SCABRITIES, (*scaber*, 'rough,') *Lichen*, *Trachoma*.

SCÆVOLA, see *Bela-modagam*.

SCALA, (F.) *Rampe*, 'a ladder.' A machine formerly used for reducing dislocations of the humerus.

SCALE OF THE COCH'LEA, *Gy'ri seu Duc'tus spir'gellæ seu Cana'les coch'leæ*, (F.) *Rampes ou Échelles du Limaçon*—are the cavities in the cochlea, separated from each other by the spiral septum. The one—*Scala inter'na seu poste'rior seu infe'rior coch'leæ*, *S. tympani*—would communicate with the *cavitas tympani* by the *fenestra rotunda*, were it not for the membrane that closes it. The other—*Scala exter'na seu ante'rior seu supe'rior coch'leæ*, *S. vestibuli*—opens into the cavity of the vestibule. The *scalæ* communicate with each other by an opening at the top of the partition.

SCALA ANTERIOR COCHLEÆ, see *Scala*—s. *Clausæ*, see *Scala*—s. *Externa cochleæ*, see *Scala*—s. *Inferior cochleæ*, see *Scala*—s. *Interior cochleæ*, see *Scala*—s. *Posterior cochleæ*, see *Scala*

—s. *Superior cochleæ*, see *Scala*—s. *Tympani*, see *Cochlea*, and *Scala*—s. *Vestibuli*, see *Cochlea*, and *Scala*.

SCALDRICKS, *Sinapis arvensis*.

SCALE, (Sax. *scale*, from *scylan*, 'to divide, separate,') (G.) *Schale*, 'shell, peel,') *Squid'na*, *Le'pis*. (F.) *Écaille*, *Écaille*. An opaque and thickened lamina of cuticle, commonly produced by some degree of inflammation of the true skin, over which it is formed.

SCALE. DRY, *Psoriasis*.

SCALÈNE ANTÉRIEUR, *Scalenus anticus*—s. *Postérieur*, *Scalenus posticus*.

SCALE'NUS, (*σκαληνος*, 'irregular or unequal,') Geometers employ this word to designate a triangle whose three sides are unequal. Anatomists have given the name to two muscles. 1. SCALE'NUS ANTI'CUS. *Portion of the Costotrachelium* (Ch.), (F.) *Scalène antérieur*, is situate at the internal and inferior part of the neck. It is long and triangular; and is inserted, below, at the upper surface of the first rib; and, above, at the anterior tubercle of the transverse processes of the 3d, 4th, 5th, and 6th cervical vertebræ. This muscle bends the neck, and draws it to its side. It can also raise the first rib. 2. SCALE'NUS POSTI'CUS, (F.) *Scalène postérieur*, *Portion of the Costotrachelium* (Ch.), is situate behind the last. It is, also, long and triangular; but more bulky than the *S. anticus*. It is attached, below, to the outer surface of the first two ribs; and terminates, above, at the summit of the transverse processes of the last six vertebræ of the neck. This muscle bends the neck laterally, and can elevate the first two ribs. Some anatomists, as Albinus and Sömmerring, have described five *scaleni* muscles on each side, and have called them *Scalenus pri'or*, *S. mi'nimus*, *S. latera'lis*, *S. me'dius*, *S. posti'cus*. Riolan and Chaussier only describe one *scalenus* muscle on each side. Cowper and Douglas, and the generality of the English anatomists, admit three, *S. anti'cus*, *S. me'dius*, and *S. posti'cus*; or, *pri'mus*, *secun'dus*, and *ter'tius*; Winslow, Boyer, and many of the French anatomists,—two, as above described.

SCA'LING—same etymon as *Scale*—in dentistry, is an operation which consists in removing the tartar from the teeth.

SCALL, (*Scale* [?]) ([Sc.] *Skale*.) *Impetigo*, *Scab*—s. *Asbestos*, see *Eczema* of the hairy scalp—s. *Crusted*, *Impetigo*—s. *Dry*, *Psoriasis*—s. *Honeycomb*, *Porrigio favosa*—s. *Humid*, *Eczema*—s. *Milky*, *Porrigio larvalis*—s. *Papulous*, *Ecthyma*—s. *Running*, *Eczema*, *Impetigo*—s. *Vesicular*, *Eczema capitis*—s. *Washerwoman's*, *Psoriasis diffusa*.

SCALLARD, (from *scall*.) *Porrigio*.

SCALLED HEAD, *Porrigio*, *P. scutulata*.

SCALP, ([D.] *Schelp*, 'a shell,') (Old Eng.) *Scop*. The integuments covering the head. The hairy scalp is called, also, *Capilli'tium*, *Trichot'ion*, *Casa'ria*, (F.) *Cherelure*, *Cuir chevelu*.

SCAL'PEL, *Scalpellum*, *Scalpel*, *Cul'ter*, *Inciso'rium*, *Sni'tion*: (from *scalpo*, 'to cut,') A cutting instrument, formed of a blade of well-tempered steel, very sharp, of variable shape and size, and fixed solidly in a handle. Its use is to divide the soft parts in operations, dissection, &c.

SCALPELLUM, *Scalpel*—s. *Lenticularo*, *Phacopis*.

SCALPELLUS, *Bistouri*, *Scalpel*.

SCALPEUM, *Bistouri*.

SCALPRUM, (*scalpo*, 'to cut, to scrape,') *Rasatorium*—s. *Chirurgicum*, *Lancet*.

SCALPRUM DENTA'RIUM, *Li'na denta'ria*. A tooth file. A dental file, or file used by the dentist.

SCALPRUM RASORIUM, *Rasatorium*.

SCALY, (from *scale*.) *Squamous*.

SCALY DISEASES or SQUAMÆ, *Lepido'ses*, form the second order in Willan and Bateman's system of cutaneous diseases.

SCAMMA, (σκαμμα,) Fossa.

SCAMMONÉE D'ALEP, *Convolvulus scammonia*.

SCAMMONIA, see *Convolvulus Scammonia*.

SCAMMONIÆ RADIX, see *Convolvulus Scammonia*—s. *Resina*, *Extractum Scammonii*.

SCAMMONIUM, (σκαμνία, σκαμνον,) see *Convolvulus scammonia*—s. *Syriacum*, *Convolvulus scammonia*.

SCAMMONY, *Convolvulus scammonia*.

SCAMMONY, BOURBON. The concrete juice of *Periplaca Mauritiana*.

SCAMMONY, GERMAN, see *Convolvulus sepium*.

SCAMMONY OF MONTPELLIER, *Cynanchum Monspeliacum*.

SCAMNUM HIPPOCRATICUM, (*scammum*, 'a bench,') *Bathron Hippocraticum* of Galen, (F.) *Banc d'Hippocrate*. A machine invented by Hippocrates for reducing fractures. It was a sort of bed, six feet long, on which the patient was fixed. Straps were attached above and below the fracture or luxation, and extension and counterextension were effected by a winch. It is described by Galen, Oribasius, Scultetus, &c. See *Bathron*.

SCANDIX, gen. *Scandicis*, (σκανδξ,) *S. Cerefolium*.

SCANDIX BULBOCASTANUM, *Bunium bulbocastanum*.

SCANDIX CEREFOLIUM, *Scandix*, *Scandix*, *Pecten Venetis*, *Cerefolium*, *Daucus Seprinum*, *Chærophyllum*, Ch. *Sativum* seu *Cerefolium*, *Anthriscus cerefolium*, *Chærefolium*, *Chervil*, (F.) *Cerfeuil*. Ord. Umbelliferae. Ser. Syst. Pentandria Digynia. A culinary herb, which is slightly aromatic and grateful. It is said to be gently aperient and diuretic.

SCANDIX ODORATA, *Sweet Cicely*. Properties the same as the other. Also, *Chærophyllum odoratum*.

SCANDULARIUS MUSCULUS, (*scandere*, 'to climb,') *Paratenar*, *Transversus pedis*.

SCANDYX, (σκανδξ,) *Scandix cerefolium*.

SCAPHIA, (σκαφη,) *Scaphium*, (σκαφος, from σκαπτο, 'I make hollow,') The excavation or cavity of the external ear, between the helix and anthelix. The meatus auditorius externus. The rima vulvæ. Also the name of a bandage for the head, mentioned by Galen, and called, likewise, *Tholus Diacelus*.

SCAPHION, (σκαφιον, dim. of σκαφη, 'a trough,') *Cranium*, *Papaver* (Capsule).

SCAPHIUM OCULARE, *Pelvis Ocularis*, *Eyeglass*. (F.) *Bassin ou Gondole oculaire*, *Œillière*. A small porcelain, glass, or metallic vessel, used for applying lotions to the eye.

SCAPHOCARPOS UPERPHALANGIUS POLLICIS, *Abductor pollicis brevis*.

SCAPHOID, *Scaphoides*, (σκαφη, 'a skiff, and εidos, 'form,') A name given to several parts.

SCAPHOID BONE OF THE FOOT, *Os Scaphoides*, *Pedis* seu *navicularè* seu *cymbiformè*, *Boutlike bone*, (F.) *Scaphoïde du Pied*. This bone is situate at the forepart of the astragalus and inner part of the foot. The posterior surface forms a cavity, somewhat like that of a boat, for receiving the head of the astragalus. There is a prominence at the inner side of the bone, for the insertion of tendons, muscles, and strong ligaments. The forepart of the bone is convex, and divided into three articular surfaces for articulation with the ossa cuneiformia. Between the os naviculare and astragalus, the foot has its principal lateral and rotary motions; although each of the other joints of the tarsus contributes a little. Part of the *Abductor pollicis* is inserted into it.

SCAPHOID BONE OF THE HAND, *Os Scaphoides*, *Manus* seu *navicularè* seu *cymbiformè*, *Boutlike bone*, (F.) *Scaphoïde de la main*. The first bone of the first row of the carpus. The upper surface is convex, and articulated with the radius. The under and outer surfaces are also convex,—to be articulated with the trapezium and trapezoides. Between the upper and under cartilaginous surfaces there is a rough fossa for the insertion of the capsular ligament. The anterior and inner surface has an oval cavity, where it is articulated with the os magnum, which gives name to the bone. There is a process on the outer end of the bone, for the attachment of part of the anterior transverse ligament of the wrist.

SCAPHOID FOSSA, *Fossa Scaphoides*, (F.) *Enfoncement Scaphoïde*. A small cavity at the upper part of the internal ala of the pterygoid process, in which the peristaphylinus internus is attached.

SCAPHOÏDE DE LA MAIN, *Os scaphoides manus*—s. *du Pied*, *Os scaphoides pedis*.

SCAPHOÏDOÄSTRAG'ALAN, *Scaphoïdoästragalanus*. That which relates to the scaphoid bone and astragalus. The articulation of these bones is so called; and the ligament which strengthens it is termed the *Scaphoïdoästragalan ligament*.

SCAPHOÏDOCUBOID, *Scaphoïdocuboïdeus*. That which relates to the scaphoid and cuboid bones. The articulation of those bones is so called.

SCAPHULA, (dim. of *scapha*,) *Navicularis fossa*.

SCAPHUS, (σκαφος, 'a skiff, a trough,') *Auditory canal*, *external*, *Rima vulvæ*, *Scapha*.

SCAP'ULA, *Scap'ulum*, *Plata*, *Omo'plata*, *Homo'plata*, *Scop'tula*, *Epino'tion*, *Latitudo hu'meri*, *Omo'platè*, *Os latum hu'meri*, *Scutum thoracis* seu *cor'dis*, *Spa'tha*, *Spath'ula*, *Pteryg'ium*, *Chel'onium*, *Spa'tula*; *Bladebone*, *Shoulderblade*, (Prov.) *Shieldbone*, *Platebone*, *Spadebone*, (F.) *Omo'plate*. An irregular, broad, flat bone, of a triangular shape, situate at the posterior part of the shoulder. It has, 1. A posterior or dorsal surface or *Dorsum*, *Tes'tudo scap'ulæ*, divided transversely into two parts by a triangular process, called *Spine of the Scapula*; which terminates by a considerable eminence, called *Acromion*. Above the spine, is the *Fossa supraspinata*; below it, the *F. infraspinata*. 2. An anterior or costal surface or *venter*, forming the *Fossa subscapularis*. 3. A superior or cervical edge, which is thin, and terminates, anteriorly, by a strong, curved apophysis, called the *coracoid process*. 4. A posterior or vertebral edge, called, also, the *base*. 5. An outer or axillary or inferior edge, having a thick, truncated angle, in which is the *glenoid cavity*. With this is articulated the head of the humerus. The glenoid cavity is separated from the body of the bone by a narrow part called the *cervix* or *neck*. The edges or margins are, also, called *costæ*. The scapula has three angles—an inferior, superior, and anterior.

SCAPULA, *Humerus*.

SCAP'ULE NA'SI, *Par'tes laterales na'si*. The lateral parts of the nose.

SCAPULAIRE, *Scapular*.

SCAPULAR L'GIA, (*scapula*, 'the shoulderblade,' and *αλγος*, 'pain,') *Arthralgia of the scapula*.

SCAP'ULAR, *Scap'ulary*, *Scapularis*, (F.) *Scapulaire*, (from *scapula*.) That which relates or belongs to the scapula.

SCAPULAR APONEUROSIS. A broad, thin aponeurosis, with decussating fibres; which is attached, above, to the spine of the scapula; below, to a crest between the *teres major* and *infraspi-*

natus; within, to the spinal edge of the scapula; and which, at its middle, unites with the thin aponeurosis that covers a part of the external surface of the deltoid.

SCAPULAR ARTERIES. These are several. 1. The *Superior scapular*, *Superficial scapular* of Sömmerring, *Dorsalis scapula superior*, *Transversalis scapula*, *T. Humeri*. This arises from the subclavian; and often from the inferior thyroid, or from a trunk common to it and the posterior scapular or transverse scapular or cervical. It passes behind the clavicle, above the superior edge of the scapula, and buries itself between the dorsal surface of that bone and the muscles inserted into it. 2. The *Inferior scapular* or *common scapular* or *internal scapular* or *subscapular* artery, *A. Thoracica axillaris seu alaris*, is of considerable size. It arises from the inferior part of the axillary artery, behind the brachial plexus—gives off several branches to the armpit, and divides into two branches:—the one, *Scapularis interna*, and the smaller of the two, is distributed, almost entirely, to the serratus major anticus, and latissimus dorsi; the other, the larger of the two, *Dorsalis Scapulae inferior seu Circumflexa Scapularis*, which distributes its branches to the subscapular, supraspinal and infraspinal regions.

SCAPULAR NOTCH, see Notch, scapular.

SCAPULAR VEINS follow the same course as the arteries, which they accompany.

SCAPULARE, Scapulary.

SCAPULARY, *Scapularis*, *Fuscia Scapularis*. A broad, linen bandage, divided into two tails for three quarters of its length, the undivided extremity of which is attached to the posterior and middle part of a bandage carried round the body; the tails passing over the shoulders; and being attached to the anterior part of the body bandage, to prevent it from slipping down.

SCAPULO, in composition, the scapula.

SCAPULODYN'IA, (*scapula*, and *dyna*, 'pain.') Pain or rheumatism of the shoulder.

SCAPULO-HUMERAL. An epithet given to the *shoulderjoint*. It is an orbicular articulation, resulting from the reception of the head of the humerus into the glenoid cavity of the scapula. To keep the bones in connection, there are: 1. A very strong, conoidal capsular ligament; passing from the circumference of the glenoid cavity around the neck of the humerus. 2. A *fibrocartilaginous ring*, called the glenoid ligament, and a synovial capsule.

SCAPULO-HUMÉRAL, *Teres major*—*s. Huméro-ulcerant*, *Triceps extensor cubiti*—*s. Hyoidien*, *Omohyoideus*—*s. Radialis*, *Biceps flexor cubiti*—*s. Trochitérien*, *grand*, *Infraspinatus*.

SCAPUS, (*skapos*, 'a shaft, a stalk,') Penis, see Hair.

SCAR, (*εσκαπα*, [F.] *escarre*), Cicatrix.

SCAR, *Seam*. To mark with a cicatrix or seam:—as *scarred*, (*Sc.*) *arred*, means marked with a cicatrix or seam.

SCARABÆOLUS HEMISPHERICUS, (dim. of *scarabeus*), *Coccus cacti*.

SCARBOROUGH, MIN'ERAL WATERS OF. *Scarborough Spa*. A celebrated English chalybeate, at Scarborough, in Yorkshire. One of the springs is a simple, carbonated chalybeate, like that of Tunbridge: the other has an admixture of purging salt. Scarborough, having the additional advantage of seabathing, is much frequented by invalids.

SCARCHT, ([Sax.] *scritchta*), Hermaphrodite.

SCARDAMYG'MUS, (*σκαρδαμυγος*), *Scardamys*, *ie*, (F.) *Clignement*, (from *σκαρδαμυγος*, 'to wink.') Winking. Closure of the eyelids.

SCARFSKIN, Epidermis.

SCARIFICATEUR, Scarificator.

SCARIFICATION, *Scarificatio*, *Amyg'is*, *Schast'is*, *Catacas'mus*, *Amyg'mus*, *Schas'ans*, *Encharax'is*, (*scarificare*, *scarificatum*, *εσκαρ'ισμα*, 'to make a slight scratch.') The act of scarifying. Also, a small incision, *Amyg'hē*, made into the skin with a lancet, bistoury, or scarificator, for different therapeutical purposes,—as to draw blood, or to discharge some effused fluid. When the scarifications are very superficial, and do not go farther than the skin, they are termed by the French, *Mouchetures*.

SCARIFICATOR, *Scarificato'rrium*, *Schastel'rium*, (F.) *Scarificateur*. An instrument for making scarifications. It consists of a small brass box, of a cubical shape, in which 10 or 12 lancet points turn upon a common pivot. By means of a spring, all these blades can be made to issue at once, and make as many scarifications.

SCARIFICATORIUM, Scarificator.

SCARIOLA, *Cichorium endivia*, *Lactuca scariola*.

SCARIOLE, *Cichorium endivia*.

SCARLATA, *Scarlatina*.

SCARLATINA, ([L.] *scarlatte*, 'a deep red.')

Scarlat'a, *Eranthis'is Rosa'tia*, *Rosa'tia*, *Rosa'tia*, *R. Squam'osa*, *Rosa'tia*, *Rub'ella*, of some, *Morbilli' ignei seu confluent'es*, *Morbus scarlatino'sus*, *Fe'bris scarlatino'sa seu ru'bra*, *Gut'turis mor'bus epidem'icus*, *Ereth'i*, *Pur'pura*, *Ty'phus scarlatinus*, *Porphyris'mus*, *Porphyris'tia*, *Scar'lot* or *Rushfever*, (F.) *Fièvre rouge*, *F. purpurée*. The characteristic symptoms of scarlatina are:—a scarlet flush, appearing about the second day of fever on the face, neck, and fauces; and progressively spreading over the body; terminating about the 7th day. Two great varieties may be reckoned:—the *S. simplex seu febris seu benigna* seu *sine angina*, in which the fever is moderate and terminates with the efflorescence; prostration of strength being trifling;—and the *S. Anginosa seu Paristhmit'ica seu Cynanthe'tica seu mit'ior*, in which the fever is severe; the throat ulcerated; the eruption later in its appearance, and less extensive; often changing to a livid hue. *Scarlatina maligna seu gravior*, has been reckoned a variety of this, in its worst degree. It agrees with the *Cynanche maligna* of Cullen.

Scarlatina belongs to the *major exanthemata*, and is a disease, chiefly, of children. The eruption differs from that of measles, in being an efflorescence not raised above the cuticle. Measles, too, is attended with catarrhal symptoms, whilst the complication, in scarlatina, is cynanche.

The disease is generally regarded as eminently contagious, but doubt may and does exist on this point. To the zymotic principle, the name *Scarlatinin* has been given.

The treatment of simple scarlatina need not be much. It must be antiphlogistic. If the throat be very much ulcerated, acid gargles, and counter-irritants must be employed; and if the affection becomes manifestly typhoid, and the sore throat of a malignant character, the case must be treated like typhus gravior; with antiseptic gargles of bark, acid, &c. Anasarca sometimes supervenes on scarlatina, and requires attention. Purgatives are here demanded as well as the use of sorbifacients, such as mercury and squill, &c.

SCARLATINA ANGINOSA, see *Scarlatina*—*s. Benigna*, see *Scarlatina*—*s. Cynanchica*, see *Scarlatina*—*s. Febris*, see *Scarlatina*—*s. Gravior*, see *Scarlatina*—*s. Miliaris*, *Rubeola*—*s. Mitior*, see *Scarlatina*—*s. Paristhmitica*, see *Scarlatina*—*s. Pustulosa*, *Rubeola*—*s. Rheumatica*, *Dengue*—*s. Sine angina*, see *Scarlatina*—*s. Urticata*, *Urticaria*.

SCARLATININ, see *Scarlatina*.

SCARLATINOID, *Scarlatinoi'des*, (*scarlatina*, and *eidos*, 'resemblance.') Resembling scarlatina.

SCARLATINOUS or SCARLATINOUS. *Scarlatinal*, *Scarlatinosus*. Relating or appertaining to scarlatina.

SCARLET BERRY, *Solanum dulcamara* — s. Fever, *Scarlatina*.

SCARRED, see *Scar*.

SCATACRASIA, (σκωρ, gen. σκατος, 'excrement,' and κρσις, 'mixture,') *Scatocrasia*.

SCATCRATIA, (σκωρ, gen. σκατος, 'excrement,' and αρατία, 'want of power,') see *Scatocrasia*.

SCAW, *Sambucus*.

SCEAU DE SALOMON, 'seal of Solomon,' *Convallaria polygonatum*.

SCELAL'GIA, (*scelos*, and *αλγος*, 'pain,') Pain of the leg.

SCELALGIA ANTICA, Neuralgia femoropretibialis — s. Postica, Neuralgia femoropoplitea — s. Puerperarum, *Phlegmatia dolens*.

SCELETEUMA, *Skeleutema*, (σκελετευμα,) Mummy.

SCELETEU'SIS, *Skeleuten'sis*, (σκελετευω, 'to desiccate, to pickle,') Mummification. Also, the mode of making a skeleton, *Skeletope'ia*, *Syntar'ia*.

SCELETOGRAPHY, *Skeletography*.

SCELETOLOGIA, *Skeletology*.

SKELETON, *Skeleton*.

SCELETOPŒA, *Sceleteusis*, *Skeletope'ia*.

SCELETUM, *Skeleton* — s. Artificial, see *Skeleton* — s. Natural, see *Skeleton*.

SCOLONCUS, (*scelos*, and *ογκος*, 'a tumour,') A tumour of the leg. *Phlegmatia dolens*.

SCOLORRHEU'MA, *Rheumatism'us cru'ris*, (*scelos*, and *ρηνμα*, 'defluxion, rheumatism,') Rheumatism of the leg.

SCELOS, (σκελος,) Leg.

SCELOSCAMBOSIS, (*scelos*, and *σκαμβος*, 'crooked,') see *Rhæbosis*.

SCELOTYRBE, (σκελοτυρβη, from *scelos*, and τυρβη, 'tumult,') *Chorea*, *Porphyræ nautica* — s. Febrilis, see *Subsultus tendinum* — s. Festinans, *Paralysis agitans* — s. Tarantismus, *Chorea*.

SCENOMENIX, (σκηνος, 'a hut,' and *μνηγξ*, 'a membrane,') see *Cell*.

SCENOTHEORIA, (σκηνος, 'a hut,' and θεωρία, 'theory,') *Cell doctrine*.

SECAPRON, (σκαπαρον,) *Amnios*, *Ascia*.

SECAPASTERIUM, (from σκαπίζω, 'to enwrap,') *Couvrechef*.

SCEPASTRUM, *Couvrechef*.

SCEPTRUM, (σκαπτρον,) *Penis*.

SCHASIS, (σχασις,) *Scarification*.

SCHASMUS, (σχασμος,) *Scarification*.

SCHASTERION, (σχαστηριον,) *Fleam*, *Lancet*, *Scarificator*.

SCEBA ARABUM, *Artemisia santonica*.

SCHEDULA, (dim. of *scheda*, 'a leaf of paper,') Prescription.

SCHERBET', *Sherbet*, *Serbet*, *Tscherbet*, *Sorbetum*, *Sorbetum*, *Sorbetum*, *Sorbet*. A drink prepared by the Turks, Persians, Egyptians, &c., with sugar and the juice of acid fruits.

SCHERLIE'VO, *Skerljevo*, *Male di Scherlievo*, *Male di Breno*, *Male di Fiume*, *Male di Scarlievo*, *Male scabioso venereo*, *Margaretizza*, *Morb'us Croa'tus seu fluminen'sis*, *Frambo'sia Illyrica*, *Scab'ies vene'rea contagio'sa*, *Vene'real Itch*. A name given to a particular form of syphilis, observed in Illyria. It is said to be capable of being communicated without immediate contact; and that its principal symptoms are:—pains in the bones, ulceration of the fauces, pustules, and fungous growths in different parts of the body.

Affections, known by this name, as well as by those of *Male di Breno*, *Falcadi'na*, *Bouda* and *Frenja*, have prevailed in the frontier countries of the Austrian empire, and were generally regarded as contagious, of syphilitic origin, and as having been introduced from a surrounding country. The

terms appear, however, in the course of time, to have been applied to a variety of ulcers, eruptions, and other morbid conditions in no way connected with syphilis, so that they lost their former signification.

SCHERO'MA, gen. *Schero'matis*, (from σχεωσι, 'a shore, dry land,') A dryness of the eye from the want of lachrymal secretion.

SCHIAS, *Isehias*, *Neuralgia femoropoplitea*.

SCHIATICUS, *Sciatic*.

SCHIDAKE'DON, (σχιδακηδον,) *Schidace'dum*, *Schindace'dum*, *Schindal'amus*, *Schindal'mus*, (from σχιζω, 'I cleave,') A longitudinal fracture of a bone.

SCHINDACEDUM, *Schidakedon*.

SCHINDALAMUS, (σχινδαλαμω,) *Schidakedon*, *Schindylesis*.

SCHINDALESIS, *Schindylesis*.

SCHINDALMUS, (σχινδαλμω,) *Schidakedon*, *Schindylesis*.

SCHINDYLE'SIS, (σχινδυλησις,) *Schindale'sis*, *Schindal'amus*, *Schindal'mus*, (from σχινδύλω, 'I split into small pieces,') An articulation of bones — by furrowing, as it were, as in that of the vomer, which constitutes, in part, the septum narium. Also, the act of splitting into small pieces.

SCHINEL'ON, (σχινελων, from σχινω, 'the mastic tree,' and ελαιω, 'oil,') *Oleum lentis'cinum*. Oil of mastic.

SCHINOCEPHALUS, (*schinos*, 'the squill,' and κεφαλη, 'head,') *Scillocephalus*.

SCHINOS, (σχινω,) *Pistacia Lentiscus*, *Scilla*.

SCHINZ'NACH or HAPS'BURG, MINERAL WATERS OF. These springs are situate three leagues from Baden, and are amongst the most celebrated bathing-places in Switzerland. The waters are of a saline sulphureous kind; and have a high reputation in cutaneous and rheumatic affections, visceral obstructions, and glandular enlargements. They contain sulphate of lime, sulphate of soda, chloride of sodium, chloride of magnesium, sulphate of magnesia, sulphate of lime, carbonate of magnesia, and oxide of iron; sulphohydric acid, and carbonic acid gases. The season is from the 15th of May to the 15th of September.

SCHISMA, gen. *Schism'atis*, (σχισμα,) *Fissure*, *Rima*.

SCHISTOCEPH'ALUS, (σχιστος, 'cleft,' and κεφαλη, 'head,') A monster having some part of the head cleft or fissured.

SCHISTOCOR'MUS, (σχιστος, 'cleft,' and κορμος, 'trunk,') *Schistotho'rax*. A monster having part or the whole of the trunk cleft or fissured.

SCHISTOM'ELUS, (σχιστος, 'cleft,' and μελος, 'a limb,') A monster whose limbs are fissured.

SCHISTORRHACHIS, (σχιστος, 'cleft,' and ραχis, 'spine,') *Hydrorachis*.

SCHISTOSO'MUS, (σχιστος, 'cleft,' and σωμα, 'body,') A malformation of the abdomen, in which the fissure and evagination extend over the entire length of the abdomen; the lower extremities being absent, or very little developed, so that the body appears as if truncated inferiorly.

SCHISTOTHORAX, (σχιστος, 'cleft,' and thorax,) *Schistocormus*.

SCHIZA, (σχιζα,) *Rima* — s. *Vulvæ*, *Rima vulvæ*. SCHIZA TRICH'IA, *Trichos'chisis*, *Schizotrich'ia* (σχιζω, 'I split,' and τριχος, 'a hair,') The splitting of the hairs at their extremities.

SCHLANGENBAD or SCHLAGENBACH, MINERAL WATERS OF. *Schlangenbad* is a much frequented watering-place in Germany. The waters contain a little carbonic acid, and carbonate of soda. Temperature 86° of Fahrenheit.

SCHNAPPS, *SCHIEDAM*, see *Gin*.

SCHNEIDERIAN MEMBRANE, *Pituitary membrane*.

SCHÖENANTHUS, (*σχῶνος*, 'the common rush,' and *ανθος*, 'a flower,') *Juncus odoratus*.

SCHOLA, (*σχολή*), School.

SCHOLLERA OXYCOCCOS, *Vaccinium oxycoccos*.

SCHOOL, *Scho'la*, (F.) *École*. A place of education. In Europe, the large Medical Schools are usually attached to the universities. In England, however, there has been no medical school of celebrity at either of the universities of Oxford or Cambridge; owing greatly to their provincial situation. Of late, two schools have existed in London, attached to the London University—University College and King's College. Excellent schools are now connected with the hospitals there, and in the larger towns in the provinces. The medical schools of Europe which have been most celebrated, are those of Edinburgh, Leyden, Berlin, Halle, Tübingen, Paris, Montpellier, Bologna, Padua, Pavia, and Pisa. In the United States, the medical schools are very numerous. Those of Philadelphia and New York are the most largely attended.

SCHOOL'EY'S MOUNTAIN, MINERAL WATERS OF. The waters of this New Jersey Spring are a carbonated chalybeate. According to Dr. McNevin, they contain extractive, chloride of sodium, chloride of calcium, chloride of magnesium, carbonate of lime, sulphate of lime, carbonate of magnesia, silice, and carbonate, and oxide of iron.

SCHOUSBŒA COCCINEA, (after P. K. A. Schousboe, a Danish botanist,) *Cacoucia coccinea*.

SCHUBERTIA DISTICHA, (after G. H. von Schubert, Professor of Natural History at Munich,) *Taxodium distichum*.

SCHWAL'BACH or LANGENSCHWAL'BACH, MINERAL WATERS OF. Schwalbach is a town of Germany, seated on the Aa, 32 miles west of Frankfort. The waters contain chloride of sodium, carbonates of lime and magnesia, iron, sulphate of lime, carbonic acid, oxygen, &c. Employed as a tonic.

SCHWANN, WHITE SUBSTANCE OF, see Substance, white, of Schwann.

SCHWEN'DECK, MINERAL WATERS OF. These waters are about 5 leagues from Munich, in Bavaria. They contain carbonic acid; carbonate and sulphate of lime, chloride of calcium; chloride of magnesium; carbonate of soda, and oxide of iron, and are much used for cutaneous affections.

SCIAT'IC, *Sciati'cus*, *Ischiati'ic*, *Ischiati'icus*, *Sciati'icus*, (F.) *Sciatique*, (from *ισχίον*, 'the haunch,') Formed by contraction from *ischiatric*, which latter is still used. See Ischiatic. Used substantively for sciatica.

SCIATIC NERVE, GREAT, *Nervus ischiad'icus*, *Great fem'oro-poplite'al* (Ch.), (F.) *Grand nerf sciatique*, ought to be esteemed a continuation of the sacral plexus. It issues from the pelvis between the pyramidalis and superior gemellus; descends along the posterior part of the thigh; and after having given branches to the gemini, quadratus, obturator externus, glutæus maximus, semimembranosus, semitendinosus, biceps and third abductor muscles, it divides, about the inferior third of the thigh, into two branches; the one, named the *external popliteal*; the other, the *internal*.

SCIATIC NERVE, LES'SER, *Infe'rior glut'e'al branch of the sa'cral plex'us*, *Les'ser fem'oro-poplite'al nerve* (Ch.), (F.) *Nerf petit sciatique*, which seems especially formed by the 2d and 3d sacral, is given off from the lower part of the plexus, and escapes from the pelvis beneath the pyramidalis muscle. It gives branches—*nervi cut'anei clunium infe'iores*—to the glutæus maximus; a *sciat'ic branch or inf'rapel'vie cut'aneous or peri-*

ne'al cut'aneous, *Pudenda'lis lon'gus infe'rior*, which passes under the tuberosity of the ischium, and furnishes filaments to the glutæus maximus, and to the muscles of the perinæum, urethra, penis, scrotum, &c.; and a *posterior femoral cutaneous branch*—middle *posterior cutaneous nerve*—the filaments of which traverse the femoral aponeurosis, to be distributed on the skin of the posterior part of the thigh. The principal branch of the lesser sciatic nerve divides, near the ham, into two or three filaments, which descend under the integuments almost as far as the inferior part of the leg.

SCIATIC NOTCH, GREAT, *Sciatic For'a'men*, *Incis'u'ra ischiad'ica ma'ior*, (F.) *Grande échancrure sciatique ou grand trou sciatique*, is a large notch at the posterior edge of each os innominatum, below the posterior and inferior iliac spine, which is converted into a foramen by the sacrum and sacrosacral ligaments. The *les'ser sciatic notch*, *Incis'u'ra ischiad'ica mi'nor*, *Lu'na Albi'ni*, (F.) *Petite échancrure ischiatique*, is much smaller than the last, from which it is separated by the sciatic spine. It gives passage to the tendon of the obturator internus, and to the internal pudic vessels and nerves.

SCIATIC REGION, Ischiadic region.

SCIATIC SPINE, *Spine of the Is'chium*, *Sp'i'nous Proc'ess of the Is'chium*, is a short, flat, pyramidal eminence on the os innominatum, above the preceding foramen; which gives insertion to the lesser sacrosacral ligament.

SCIATICA, Neuralgia femoropoplitea.

SCIATIQUE, Neuralgia femoropoplitea, Sciatica.

SCIE, (from *secare*, 'to cut,') Saw.

SCIENTIA HERMETICA, 'Hermetic science,' Alchemy—s. Medendi, Medicina—s. Medica, Medicina.

SCIERO'PIA, (*σκιερὸς*, 'shady,' and *ὤψ*, 'the eye,') A defect of vision, in which all objects have a darker hue than natural.

SCIL'LA, (*σκιλλα*, from *σάω*, 'I cleave, I split,' or from *σκελλω*, 'I dry,') *Scil'la seu Squil'la marit'ima*, *Scyl'la*, *As'chil*, *Schi'nos*, *Oc'ulus Typho'nis*, *Ornithog'alum marit'imum seu squil'la*, *Stella'ris seu Urgan'ea scilla*, *Ce'pa mar'i'na*, *Squilla*, *Squill or sea onion*, *Eye of Typhon*, (F.) *Scille*, *Oignon marin*. Ord. Liliaceæ. Sex. Syst. Hexandria Monogynia. Native of Spain, Austria, &c. The bulb or root of the squill, *Scilla* (Ph. U. S.), has a bitter, nauseous taste, and is extremely acrid; inflaming the skin when rubbed on it. Its acrimony, on which its virtues depend, is destroyed by heat, drying, and keeping. It is extracted by vinegar, spirit, and water. In large doses, squill is emetic and purgative; in small doses, diuretic and expectorant. Its active principle has been called *scillitin*. Dose, gr. j to v of the dried root, united or not with mercury.

SCILLA ESCULEN'TA, *Phalan'gium esculen'tum*, *Eastern Quamash*, *Wild Hy'acinth*; indigenous; flowering in May. An article of diet amongst the Western Indians. It is made into bread and poultices for inflamed breasts.

SCILLIT'IC, *Scilliti'icus*, (F.) *Scillitique*, (from *scilla*.) Containing squill.

SCILLITIN, see Scilla.

SCILLOCEPH'ALUS, *Schinoceph'alus*, (*σκιλλα*, 'the squill,' and *κεφαλή*, 'head,') One who has a large bulbiform head.

SCINC'US, (*σκιγος* or *σκινγος*.) A species of *Saur'us* or *Lacert'a*, common in Egypt, and much extolled by the ancients as alexipharmic and aphrodisiac.

SCINTIL'LA VEN'ERIS, 'a spark, a relic of Venus.' A name given, by Paracelsus, to the impotence and loss of power in the limbs produced by syphilis.

SCINTILLEMENT DE L'ŒIL, (from *scintilla*, 'a spark.') 'sparkling of the eye.' Confusio.

SCIOM'ACHY, (σκιόμαχια, *Skio'ma'chia*, *Skio'ma'chia*, (σκια, 'a shade,' and πᾶν, 'a fight.') A gymnastic exercise with the ancients, which consisted in a mock encounter at boxing and jumping with one's own shadow.

SCIRRHUS, (σκίρρος,) *Scirrhus*.

SCIRPUS PALUSTRIS, (*scirpus*, 'a rush,') *Eleocharis palustris*.

SCIRRHE, *Scirrhus*.

SCIRRHOBLEPHARONCUS, (*scirrhus*, βλεφαρον, 'eyelid,' and ογκος, 'swelling,') see *Scleriasis*.

SCIRRHOCELE, (*scirrhus*, and κλην, 'rupture,') *Orchocele scirrhosa*.

SCIRRHODES, (σκιρῶδης, from *scirrhus*, and *odes*,) *Scirrhous*.

SCIRRHOGASTRIA, *Scirrhus ventriculi*, (F.) *Squirrogastrie*; (*scirrhus*, and γαστρ, 'stomach,') *Scirrhos* of the stomach.

SCIRRHOID, *Scirrhoïdes*, *Scirrho'des*, (*scirrhus*, and εἶδος, 'resemblance,') Resembling *scirrhus*.

SCIRRHOMA, gen. *Scirrho'matis*, (σκιρῶμα,) *Scirrhus*.

SCIRRHOPHTHALMIA, *Sclerophthal'mia*, (*scirrhus*, and ὀφθαλμος 'eye,') *Cancer oculi*. Cancer of the eye.

SCIRRHOPHTHALMIA PALPEBRARUM, see *Scleriasis*.

SCIRRHOSARCA NEONATORUM, (*scirrhus*, and σὰρξ, gen. *sarkos*, 'flesh,') Induration of the cellular tissue.

SCIRRHOSIS, (from *scirrhus*,) *Scirrhus*.

SCIRRHOUS, *Scirrhous*, *Scirrho'des*, (F.) *Squirrheux*, (from *scirrhus*,) Of or belonging to *scirrhus*.

SCIRRHUS SARCO'MA OF AB'ERNETHY, *Emphy'ma sarco'ma scirrho'sum*. Hard, rigid, vascular infarction of glandular follicles; indolent, insentient, glabrous; sometimes shrinking and becoming more indurated. Found in glandulous structures.

SCIRRHUS, *Scirrhus*, *Squirrhus*, *Skirrhus*, *Scirrho'ma*, *Scirrho'sis*, *Cancer scirrho'sus*, *Carcino'ma simplex* seu *fibro'sum*, *Induratio maligna*, *Sclero'sis*, *Pseudino'ma*, *Ido'ma*, *Carcinom'atous sarco'ma*, *Hard* or *Fibrous cancer*; (from σκίρρος, 'hard, indurated,') (F.) *Squirrhe*, *Squirre*, *Skirre*, *Skirre*, *Cancer dur* ou *squirrheux* ou *fibreux*. A disease, so called from the hardness that characterizes it. It is a state of induration, of a peculiar kind, affecting glandular structures generally, but capable of occurring in other textures. It usually precedes carcinoma, of which it may, indeed, be considered as the first stage. *Scirrhus* must not be confounded with the induration which succeeds inflammation, although the two states are often complicated, and may stand, occasionally, in the relation of cause and effect to each other. *Scirrhus* is ordinarily accompanied by violent shooting pains. It is also irregular on its surface; and when cut into has a bluish or grayish white colour. When the surgeon is satisfied of the existence of *scirrhus*, he had better remove it at once. No other treatment seems to possess much advantage.

Galen uses the term *Scirrho'sis* for a livid excrescence caused by inflammation.

SCIRRHUS ET CARCINOMA VENTRICULI, see *Gastrostenosis cardiaca* et *pylorica*—s. *Hepatis*, *Hepatoscirrhos*—s. *Intestinorum*, *Enteropathia cancerosa*—s. *Palpebrarum*, see *Scleriasis*—s. *Testiculii*, *Sarcocele*—s. *Uteri*, *Metroscirrhos*—s. *Ventriculii*, *Gastroscirrhos*.

SCISSION, (from *scindere*, *scissum*, 'to split,') see *Generation*.

SCISSIPARITÉ, (*scindere*, *scissum*, 'to split,' and *pario*, 'to bring forth,') see *Generation*.

SCISSORS, *Scissor*, a carver, (*scindere*, *scis-*

sum, 'to cut,') *For'fex*, *For'pex*, *Psallis*, *To'mis*, (F.) *Ciseaux*. A wellknown instrument in common use, and often employed in surgery. Scissors are composed of two cutting blades, crossing each other and moving upon an axis. They are straight or curved. Sometimes the curve is on the flat side, when they are called, in France, *Ciseaux à eniller*; at others, on their edges. The scissors, bent so that the blades make an obtuse angle with the handles, are also much used. They are called in France *Ciseaux coudés*. Scissors are used for dividing soft, loose, isolated parts, which it would be difficult to reach and to cut with the bistoury. They are employed, likewise, in the operation for harelip, and for removing warts, fungous excrescences, &c.

SCISSORS, *SMELLIE's*, see *Perforator*.

SCISSURA, (from *scindere*, *scissum*, 'to split,') Fissure—s. *Vulvæ*, see *Rima*.

SCISSURÆ, *Rhagades*.

SCISSURE, Fissure—s. *De Glaser*, Fissure—s. *Grand médiane du Cervelet*, Valley—s. *Grande du foie*, see *Liver*, Fissure, glenoid—s. *de Santorini*, *Santorini*, fissure of.

SCITTA, see *Dysentery*.

SCLAREA, *Salvia sclarea*.

SCLEREMA, gen. *Sclere'matis*, (from *scleros*,) Induration of the cellular tissue.

SCLÉRÈME, Induration of the cellular tissue.

SCLEREMIA, Induration of the cellular tissue.

SCLEREMUS, Induration of the cellular tissue.

SCLERENCEPHALIA, *Induratio cer'ebri*, (*scleros*, and εγκέφαλος, 'the brain,') (F.) *Endurcissement du cerveau*. Induration of the brain.

SCLERIA, (σκληρία, from *scleros*,) *Scleriasis*.

SCLERI'ASIS, (σκληρίασις, from *scleros*,) *Scleroma*, *Sclero'sis*, *Sclero'tis*, *Sclerys'mus*, *Seph'irus*. Any kind of hardness or induration. Sometimes, hardness of the eyelids, *Callositas* seu *Scirrhophthal'mia* seu *Scirrhus palpebrarum*, *Scirrhoblepharon'eus*; at others, hardness of the genital organs of the female; probably synonymous with *scirrhus*. Hardness of the bones, *Fragilitas Os'sium*.

SCLERITIS, *Sclerotitis*.

SCLEROCATARACTA, see *Cataract*.

SCLERODERMA, gen. *Scleroderm'atis*, (*scleros*, and δερμα, 'skin,') Induration of the areolar tissue.

SCLERODES, (σκληρῶδης, 'hard,' and *odes*,) *Sclerotic*.

SCLEROTITIS. Inflammation of the sclerotic and iris.

SCLEROMA, (from *scleros*,) Induration of the areolar tissue, *Scleriasis*.

SCLEROMENINX, gen. *Scleromenin'gis*, (*scleros*, and μενινξ, 'membrane,') *Dura mater*.

SCLERONYXIS, *Scleroticonyxis*.

SCLEROPHTHALMIA, (σκληροφθαλμία,) *Hordeolum*, *Schirophtalmia*.

SCLEROS, (σκληρος,) *Hard*.

SCLEROSARCO'MA, gen. *Sclerosarco'matis*, (*scleros*, and σαρκωμα, 'a fleshy tumour,') A hard, fleshy tumour seated on the gums, and resembling a cock's comb.

SCLEROSIS, (from *scleros*,) see *Induration* of the areolar tissue, and *Scleriasis*. With *Virchow*, thickening with condensation.

SCLEROSTENOSIS CUTANEA, (*scleros*, and στενός, 'contracted,') Induration of the areolar tissue.

SCLEROT'IC, *Sclerot'ica*, *Sclero'des*, (from *scleros*,) *Cor'nea opa'ca*, *Tu'nica du'ra* seu *alb'igin'ea* seu *duri'or* et *crassi'or* seu *ex'tima* seu *al'ba* seu *can'dida* seu *innomina'ta oculi*, *Sclero'tis*, (F.) *Sclerotique*. A hard, resisting, opaque membrane, of a pearly white colour and fibrous nature, which covers nearly the posterior four-fifths

of the globe of the eye, and has the form of a sphere truncated before. It is strengthened by the expansion of the muscles of the eye, to which expansion many anatomists have given the name *Tunica albuginea*. The subconjunctival sclerotic, (F.) *Blanc de l'œil*, forms the white of the eye, *Albu'men seu Albu'm seu Albu'go oculi*, *Lon'chades*, *Log'ades*, *Leu'cades*. The sclerotic is penetrated posteriorly by a round opening, for the passage of the optic nerve; and, anteriorly, it has a much larger opening, into which the transparent cornea is received. There are, besides, many small oblong apertures in it for the passage of nerves and vessels proceeding to the interior of the eye. Its inner surface is of a light brown colour, owing to a delicate areolar tissue containing brown pigment cells; and hence called *Membrana seu Lamina fusca*. In the fetus, the sclerotic may be separated into two very distinct laminae.

SCLEROTICA CERATOIDES, Cornea.

SCLEROTICECTOM'IA, *Sclerectomia*, (sclerotic, and *ektomē*, 'excision.') The operation for forming an artificial pupil by excision of a portion of the sclerotic.

SCLEROTICITIS, Sclerotitis.

SCLEROTICONYX'IS, *Scleronyxis*, (sclerotic, and *onyx*, 'puncturing.') Perforation of the sclerotic in the operation for cataract by depression.

SCLEROTICOTOM'IA, (sclerotic, and *tomē*, 'incision.') The operation for cataract by cutting through the sclerotic.

SCLEROTICUS, (from *scleros*.) Indurans.

SCLEROTIQUE, Sclerotic.

SCLEROTIS, (from *scleros*.) Sclerotic.

SCLEROTITIS, *Inflammatio Sclerotica*, *Leucitis*, *Lonchaditis*, *Logaditis*, *Scleritis*, *Sclerocitis*, *Rheumatic ophthalmia*, *Ophthalmia arthritica*, (sclerotic, and *itis*.) Inflammation of the sclerotic coat of the eye.

SCLEROTIUM CLAVUS, (from *scleros*.) see Ergot—s. Giganteum, see Lycoperdon tuber.

SCLEROTOMIA, Scleroticectomy.

SCLERUNTICUS, (σκληρυντικός.) Indurans.

SCLERUS, (σκληρος.) Induration, Scirrhus.

SCLERYMEN, Sclerotic.

SCLERYMENTIS, (sclerymen, and *itis*.) Inflammation of the sclerotic coat of the eye.

SCLERYNTICUS, (σκληρυντικός.) Indurans.

SCLERYS'MA, (σκληρυσμα, from *Scleros*.) Induration; also, induration of the liver.

SCLERYSMUS, (σκληρυσμος.) Scleriasis.

SCLOPETA'RIOUS, *Sclopeticus*, (from *sclopetum*, 'a gun.') Relating to a gun. Hence, *Aqua sclopetaria seu sclopetica*, a water for gunshot wounds. *Vuln'era sclopetaria seu sclopetica*, gunshot wounds, &c.

SCLOPETOPLAGÆ, (sclopetum, 'a gun,' and *plaga*, 'a wound,') Wounds, gunshot.

SCOBs, gen. *Scob'is*, *Pri's'ma*, *Xys'ma*, *Ramen'tum*, *Ras'u'ra*. A shaving; also, an alkali; and the scoria of any metal.

SCOLECESIS, (σκοληκσις.) Helminthiasis.

SCOLECIASIS, (σκοληκσις.) Helminthiasis.

SCOLECOLOGIA, (σκοληξ, 'a worm,' and *logos*, 'a discourse,') Helminthology.

SCOLEX, (σκοληξ.) *Ascaris lumbricoides*. *Scollex* was used by Müller, generically, for some imperfectly developed forms of *Tania*. The head of these being the most characteristic part, it was generally applied to the heads of all forms of cystic and tape worms. It has now no generic meaning, as the beings to which it was applied were immature forms of other genera. (Lankes-
cer.)

SCOLIOMA, (σκολιωμα.) Scoliosis.

SCOLIOSIS, gen. *Scol'e'is*, *Scolio'ma*, *Skolio'is*, *Rhachioscolio'ma*, (σκολιωσις, from *σκολιω*,

'crooked,' and *osis*.) A distortion of the spine to one side. See *Gibbosity* and *Hump*.

SCOLOPENDRA, (σκολοπένδρα.) *Asplenium scolopendrium*.

SCOLOPENDRIA, *Asplenium ceterach*, and *A. scolopendrium*.

SCOLOPENDRIUM LINGUA, *Asplenium scolopendrium*—s. *Officinatum*, *Asplenium scolopendrium*—s. *Phyllitis*, *Asplenium scolopendrium*—s. *Ruta muraria*, *Asplenium ruta muraria*—s. *Vulgare*, *Asplenium scolopendrium*.

SCOLOPOMACHÆRION, (σκολοπαξ, 'the woodcock,' and *μαχαίρα*, 'a knife,' so called because it is bent a little at the extremity like a woodcock's bill.) A sort of bluntpointed bistoury.

SCOLYMUS SATIVUS, (σκολυμος.) *Cynara scolymus*.

SCOOP, (Teut. *Schoepe*.) *Cochlea'rē*, (F.) *Curette*. A surgical instrument of the shape of a spoon, which is used for the extraction of certain foreign bodies. The scoop is often employed to extract balls impacted in the soft parts; to remove calculi from the urinary bladder in lithotomy, &c. A small scoop is sometimes used to extract foreign bodies from the meatus auditorius externus, nasal fossae, &c.

SCOP, (scalp.) Head, Scalp.

SCOPA REGIA, (*scopa*, 'a broom,') *Ruscus*.

SCOPARIN, see *Spartium scoparium*.

SCOPARIUS, (*scopa*, 'a broom,') see *Spartium scoparium*.

SCOPE, σκωπη, *Scop'y*, *Scop'ia*, *Scop'ium*, *Scopos*, (from *σκοπεω*, 'I look around.') A common suffix to words meaning 'view, inspection,' as *stethoscope*, *stethoscopy*, &c.

SCOPOLINA ATROPOIDES, (after J. A. Scopoli, Prof. at Pavia.) *Hyoscyamus scopolia*

SCOPTULA, Scapula.

SCOPULA, (dim. of *scopa*, 'a broom,') Brush.

SCOR, (σχωρ, gen. *σκαρος*.) Excrement.

SCORACRAT'IA, *Scoracrat'ia*, *Scatacrat'ia*, *Scatacrati'a*, *Copracrati'a*, *Copracrat'ia*, *Incontin'entia Al'vi*, (*scor*, and *ακρατια*, 'want of control.') Want of power to retain the faeces. Involuntary evacuation of the faeces.

Scoracrasia, *Scatacrasia*, and *Copracrasia*, although used in the same sense as *scoracratia*, really mean undue admixture of the excrement. See *Scatacrasia*.

SCORBUT, *Porphyra nautica*, *Purpura hæmorrhagica*—s. *des Alpes*, *Pellagra*—s. *de Terre*, *Purpura hæmorrhagica*.

SCORBUTIC, *Scorbuticus*, (F.) *Scorbutique*, (from *scorbutus*, 'scurvy.') That which belongs to scurvy. One affected with scurvy.

SCORBUTUS, ([G.] *Scharboeck*.) see *Purpura*—s. *Alpinus*, *Pellagra*—s. *Nauticus*, *Porphyra nautica*—s. *Oris*, *Cancer aquaticus*.

SCORDINEMA, (σκορδισμα.) *Carebaria*, *Pandiculation*.

SCORDINISMUS, (σκορδίνισμος.) *Carebaria*, *Pandiculation*.

SCORDIUM, (σκορδιον, perhaps from *scordon*, 'garlic,' from similarity of smell.) *Teucrium scordium*.

SCORDON, (σκορδον.) *Allium*.

SCORITH, Sulphur.

SCORIUM, Scrotum.

SCORODON, (σκοροδον.) *Allium*.

SCORODOSMINE, (*scorodon*, and *osmē*, 'odour,') *Oxyd*, cystic.

SCORPIODEN'IS, (σκαρπιος, 'scorpion,' and *ήξίς*, 'bite;'—more properly, *SCORPIOSTIG'MA*; (from *σκαρπιος*, and *στιγμα*, 'puncture.') The sting of the scorpion.

SCORPION, *Scor'pio*, *Scor'pinus*, (σκαρπιον and *σκαρπιος*, from *σκαρπιω*, 'I enrage.') A genus of insects having a sting at the extremity of their tail, with which they make dangerous wounds.

An Oil of *Scorpion* was once made from them, which is not now used.

SCORPIOSTIGMA, (*scorpius*, and σκorpion, 'a puncture,') Scorpiodexis.

SCORPIUS, (σκόρπιος,) Scorpion.

SCORTUM, 'a skin, leather,' Scrotum.

SCORZONE'RA, *S. hu'milis* seu *lana'ta* seu *nerve'ra*, *Scorzone'ra*, (from *skorzo*, old Spanish for a viper,) *Viperaria*, *Serpentaria Hispanica*, *Officinal viper's grass*, *Viper's grass*, *Goat's grass*. Family, Cichoraceæ. Sex. Syst. Syngenesia Polygamia. The roots have been used as alexipharmics, and in hypochondriacal disorders and obstructions of the viscera. The root of the *Scorzonera Hispanica* is esculent, but inefficacious as a drug.

SCOTAS'MA, gen. *Scotasm'atis*, *Scoto'ma*, *Scoto'dia*, *Scoto'sis*, *Scotos*; (from *skotos*, 'darkness,') Obscurity of vision, darkness, *Suffusio rad'ians*. Also *Scotodinia*.

SCOTCH FIDDLE, *Psora*.

SCOTODIA, (σκόροδια,) *Scotasma*, *Scotodinia*.

SCOTODIN'IA, *Scoto'ma*, *Di'nus* *Scoto'ma*, *Scotod'inē*, *Tenebrico'sa verti'go*, *Scot'os*, *Scotas'ma*, *Scoto'sis*, *Scot'omy*, (σκόροδινια, from *skotos*, 'darkness,' and *dineō*, 'I turn round,') Giddiness, with impaired sight, often succeeded by headache. See *Vertigo*.

SCOTOMA, gen. *Scotom'atis*, (σκότωμα,) *Scotasma*, *Scotodinia*.

SCOTOS, (σκότος,) *Scotasma*, *Scotodinia*.

SCOTOSIS, (σκότωσις,) *Scotodinia*.

SCOTT'S ACID BATH, *Nitromuriat'ic ac'id bath*. A bath of dilute aqua regia, employed by Dr. Scott, of India, as a remedy in hepatic diseases. Three pints by measure of muriatic acid, and two of nitric acid, are mixed to form the aqua regia. In preparing this for use, a pint of it is mixed with an equal quantity of water. The acid bath consists of three ounces of this dilute acid to every gallon of water.

SCOUR, ([Sax.] *scour*), *Diarrhœa*.

SCOURGING, *Mastigosis*—*s. Mania*, see *Masugosis*.

SCOURING, *Diarrhœa*.

SCRAT, ([Sax.] *scraþta*), *Hermaphrodite*; ([G.] *kratzen*, 'to scratch,') *Psora*.

SCRATCH, *Psora*.

SCREATUS, (*screeare*, *screatum*, 'to hawk,') *Excreation*.

SCRIBE PALSY, *Cramp*, writers'.

SCRIPULUM, *Sruple*.

SCRIPTIUM, *Sruple*.

SCRIPTULUM, *Sruple*.

SCRIPULUS, *Sruple*.

SCRIVENER'S CRAMP, *Cramp*, writers'.

SCROBIC'ULUS, (dim. of *scrobs*, 'a ditch,') *Scrobic'ulum*, (F.) *Scrobicule*, *Fossette*. A little ditch or furrow.

SCROBIC'ULUS COR'DIS, *Anticar'dium*, *Procar'dium*, (F.) *Fossette du cœur*, *Avant-cœur*, *Anti-cœur*. The pit of the stomach. The depression at the anterior part of the abdomen, below the xiphoid cartilage.

SCROBICULUS VARIOLE, *Pockmark*.

SCROFF'Æ, *Scrofula*.

SCROFULA, *Scroph'ula*, *Scroph'ulæ*, (from *scrofa*, 'a sow,' as *Cheras* is from *χαιρας*, 'a hog,' because swine were presumed to be subject to a similar complaint.) *Scroful'osis*, *Morbus scrophul'osus*, *Vit'ium scrophul'osum*, *Pedatroph'ia glandulo'sa*, *Stru'ma*, *Chas'ras*, *Co's'ras*, *Tu'ber Glandula'ris*, *Adeno'sis scrophulo'sa*, *Scroph'ie*, *King's evil*, *The Evil*, *Quince*, *Cruels*, and *Crenels* (*Scotticæ*), (F.) *Scrophules*, *Scrofulæ*, *Strumes*, *Écrouelles*, *Humeurs froides*, *Tubercules des glandes lymphatiques*, *Mal du Roi*. A morbid state of the system usually characterized by indolent, glandular tumours, chiefly in the neck; suppurating slowly

and imperfectly, and healing with difficulty; the disease ordinarily occurring in those of a sanguine temperament, with thick upper lip, &c. The tumours, after suppurating, degenerate into ulcers; which, in process of time, cicatrize, leaving scars. The internal organs are apt to be attacked in those disposed to scrofula; hence they are often the subjects of phthisis and mesenteric affections. Scrofula is hereditary; and is frequently excited by insufficient or improper regimen, and by a close, confined air during the first years of existence. The best treatment is:—to strengthen the system by animal diet; pure air, and exercise, cold bathing, or seawater bathing, &c.

SCROFULA ABDOMINALIS INTERNA, *Physconia strumosa*.

SCROFULAIRE, *Scrophularia nodosa*—*a. Aquatique*, *Scrophularia aquatica*.

SCROFULARIA, *Scrophularia*.

SCROFULES, *Scrofula*.

SCROFULÉUX, *Scrofulous*.

SCROFULIDE, *Scrophulide*. An affection of the skin or mucous membranes, dependent on scrofulosis.

SCROFULISM, see *Diathesis*.

SCROFULO'SIS, *Scrophulo'sis*, *Cacochym'ia* seu *Cachex'ia* seu *Dyscras'ia scrophulo'sa*, and *Scrofulous cachex'ia*, are used to signify the cachectic state of the system, which characterizes scrofula. See *Diathesis*.

SCROFULOUS, *Scrophulo'sus*, *Scrophula'rius*, *Stru'mous*, *Strumo'sus*, (F.) *Scrofuléux*, *Struméux*. Suffering from, or relating to, scrofula,—as *scrofulous cachex'ia*, see *Scrofula*.

SCROGGEGGED, *Cnemosecoliosis*.

SCROOBYGRASS, *Cochlearia officinalis*.

SCROOF, *Scurf*.

SCROPHULA, *Scrofula*—*s. Mesenterica*, *Tabes mesenterica*—*s. Molluccana*, *Frambœsia*.

SCROPHULÆ, *Scrofula*—*s. Gonorrhœica*, *Gonorrhœa impura*.

SCROPHULAR'IA, (from *scrofula*, against which it has been employed.) *S. Nodosa*.

SCROPHULAR'IA AQUAT'ICA, *Beton'ica aquat'ica*, *Fica'ria*, *Ferra'ria*, *Castran'gula*, *Great'er Water Fig'wort*, *Water Bel'ony*, (F.) *Scrofulaire aquatique*. Ord. *Scrophulariaceæ*. The leaves are celebrated as correctors of the bad flavour of senna. They were at one time regarded as eminently antiscrofulous, stomachic, and carminative.

SCROPHULARIA FETIDA, *Scrophularia nodosa*—*s. Lanceolata*, *S. Nodosa*—*s. Marilandica*, *S. Nodosa*—*s. Minor*, *Ranunculus ficaria*.

SCROPHULAR'IA NODO'SA, *S. fo'etida* seu *vulga'ris* seu *lanceola'ta* seu *Marilan'dica*, *Millemor'bia*, *Scrophula'ria*, *Fig'wort*, *Ken'nelwort*, *Holmes'weed*, *Heal'all*, (F.) *Scrofulaire*, *Herbe aux écouelles*. The root and leaves of this plant have been celebrated, both as an internal and external remedy, against inflammations, piles, scrofulous tumours, ulcers, &c.

SCROPHULARIA VULGARIS, *Scrophularia nodosa*.

SCROPHULARIUS, *Scrofulous*.

SCROPHULES, *Scrofula*—*s. Mésentériques*, *Tabes mesenterica*.

SCROPHULIDE, *Scrophulide*.

SCROPHULOSIS, see *Diathesis*.

SCROPHULOSUS, *Scrofulous*.

SCROTAL, *Scrota'lis*, (from *scrotum*.) Relating to the scrotum; as *Hernia scrota'lis*, *scrotal hernia*.

SCROTAL NERVE, see *Genitocrural Nerve*.

SCROTOCE'LE, *Oscheocece*, *Oschocele*, (*scrotum*, and *κῆλη*, 'a tumour,') *Hernia Scrota'lis*, *Scrotal her'nia*. Inguinal hernia, descending into the scrotum.

SCROTUM, (*scortum*, 'a skin, leather,') *Scot'rium*, *Os'cheë*, *Os'cheus*, *Os'chus*, *Os'cus*, *Os'cheon*, *Och'æus*, *Pe'r'in*, *Scortum*, *Co'tenus*, *Co'tens*, *Mar'supium*, *Bur'sula*, *Bur'sa* seu *Bur'sa tes'tium* seu

virilis, *Orchus*, *Folliculus genitalis*, *Hernia*, the *Purse*, the *Cud*, *Cods*, (F.) *Les bourses*. The integuments which cover the testes. These are nothing more than a prolongation of the skin of the inner part of the thighs, perinæum, and penis. The skin is remarkable for its brown colour, for the numerous rugæ, especially when the scrotum is contracted; for the great number of sebaceous follicles it contains; and for the long and few hairs which issue from it in the adult. The scrotum is divided into two equal parts by a raphe, which extends from the anus to the root of the penis. Its organization is similar to that of the skin of other parts of the body, except that the chorion is thinner.

SCROTUM CORDIS, Pericardium—s. Pendulum, Rachosis.

SCRUBIE, (*scurvy*), *Purpura nautica*.

SCRUBIE GRASS, *Cochlearia officinalis*.

SCRUPLE, *Diobolon*, *Granmarium*, *Servipulus*, *Servipulum*, *Scriptulus*, *Scriptulum*, *Scriptulum*, *Scriptulum*, *Scriptulum*, *Scriptulum*. A weight of twenty grains, Troy; twentyfour grains, Poids de marc.

SCUDAMORE'S MIXTURE, see Mixture, Scudamore's.

SCULL, (Dan. *skal*, or from *scutella*, 'a cup,' [?]) Cranium—s. Cap, Calvaria.

SCURF, ([Sax.] *scurf*,) (Prov.) *Scurf*; *Furfur*. Small exfoliations of the cuticle are so called.

SCURFY, *Furfuraceous*.

SCURVY, (from *scurf*, or from [G.] *Scharbock*), see *Purpura*—s. of the Alps, *Pellagra*—s. Button, see *Button scurvy*—s. Land, *Purpura hæmorrhagica*—s. Petechial, *Purpura simplex*—s. Sea, *Porphyra nautica*.

SCURVY GRASS, *Sisyrinchium Bermudianum*—s. Grass, common, *Cochlearia officinalis*—s. Grass, lemon, *Cochlearia officinalis*—s. Grass, wild, *Cochlearia coronopus*.

SCUTELLA, (dim. of *scutra*, 'a dish,') Cup.

SCUTELLARIA CAROLINIANA, (from *scutella*, 'a cup,' in allusion to the form of the calyx,) *S. integrifolia*.

SCUTELLARIA GALERICULA'TA, *Cassida galericulata*, *Tertiana'ra*, *Skull'cap*, (F.) *Casside bleue*, *Toque*. Ord. Labiatae. Sec. Syst. *Didynamia Angiospermia*. This plant has a bitter taste and a garlic smell. It has been esteemed especially serviceable in tertian ague.

SCUTELLARIA HYSSOPIFOLIA, *S. integrifolia*.

SCUTELLARIA INTEGRIFOLIA, *S. hyssopifolia* seu *Caroliniana*,—another indigenous variety,—is an intense bitter.

SCUTELLARIA LATERIFLO'RA, *Skull'cap*, *Mad-dog skull'cap*, *Mad'weed*, *Skull'wort*, *Blue Pimpernel*. An indigenous plant, *Scutellaria*, (Ph. U. S.) which grows in all parts of the United States in moist places. It was at one time esteemed a preventive of hydrophobia, but is probably possessed of little virtue of any kind. By some, however, it is regarded as a good nerveine.

SCUTICULA DIGITORUM MANÛS, Phalanges of the fingers.

SCUTER, Diarrhoea.

SCUTULÆ DIGITORUM MANÛS, (pl. of *scutula*, 'a cylinder,') Phalanges of the fingers.

SCUTUM, 'a shield,' Thyroid cartilage—s. Cordis, Scapula, Sternum—s. Genu, Patella—s. Pectoris, Sternum, Thorax.

SCUTUM STOMACH'ICUM. A large plaster, applied to the breast or stomach.

SCUTUM THORACIS, Scapula.

SCYB'ALA, (pl. of *σκυβαλον*, 'dung,') *Enterolithus Scybalum*, *Ecrementa Alvi sicciora*, *Fæces induratae*, (F.) *Scybales*. Hard fecal matters discharged in round lumps.

SCYLLA, *Scilla*.

SCYPHOPHORUS PYXIDATUS, (*σκυφος*, 'a cup,' and *φωρα*, 'to bear,') *Lichen pyxidatus*.

SCYPHUS AUDITORIUS, (*σκυφος*, 'a cup,') Infundibulum of the cochlea—s. *Cerebri*, Infundibulum of the brain—s. *Viussenii*, Infundibulum of the cochlea.

SCYROS, (*σκυρος*,) Callosity.

SCYTALIDES DIGITORUM MANÛS, (*σκυταλεις*, gen. *σκυταλεις*, 'a staff,') Phalanges of the fingers.

SCYTHICA (RADIX,) (from *Scythia*,) *Glycyrrhiza*.

SCYTHROPASMUS, (from *σκυθος*, 'gloomy,' *σχυρωπαζειν*, 'to look gloomy,') *Vultus tel'ricus et mac'us*. A gloomy, depressed countenance, of bad augury in serious diseases.

SCYTITIS, (*σκυτος*, 'a skin,' and *itis*,) Cytitis.

SCYTO DEPSIUM, (*σκυτοδεψειν*, 'to curry leather,') Tannin.

SEA AIR. The air in the neighbourhood of the sea contains saline particles commingled with it, which render it a valuable change for scrofulous and debilitated individuals. The sea and seacoast are usually, also, more equable in their temperature than places inland.

SEA SICKNESS, *Nausea marina*—s. Tangle, *Laminaria digitata*—s. Ware, see *Pila marina*—s. Water, see *Water*, sea—s. Wrack, *Pila marina*.

SEAL, GOLDEN, ([Sax.] *rigel*, [L.] *sigillum*,) *Hydrastis Canadensis*—s. Medicine, *Cachet*—s. Solomon's, *Convallaria polygonatum*—s. Solomon's, great, *Convallaria canaliculata*—s. Solomon's, smaller, *Convallaria pubescens*.

SEAM (Sax.), Cicatrix.

SEARCHER, Body searcher.

SEARCHING, ([F.] *chercher*, 'to seek,') Probing. Also, the operation of ascertaining, by the introduction of a metallic instrument, whether there is a stone in the bladder;—*Sound'ing*.

SEARCHCLOTH, ([Sax.] *rap*, 'a sore,') A cloth to cover a sore; a plaster.

SEAS'ON, *Tempes'tas*, (F.) *Saison*, (from *statio*, 'station,' [?]) One of the four great divisions of the year, reckoned by the return of the sun in its annual course,—Spring, Summer, Autumn, and Winter. It is to the seasons that we are indebted for the organic changes in the vegetable kingdom; and their mutation exerts a powerful influence on man.

SEASONING, (from [F.] *saison*, season,') Acclimation—s. Fever, Fever, stranger s.

SEAT, (from [L.] *sedeo*, 'to sit,') Anus, Nates—s. Bone, Ischion.

SEBACÆ, Sebaceous.

SEBACEOFOLLICULAR, see Sebaceous.

SEBA'CEOUS, *Seba'ceus*, (F.) *Sébacé*, (from *sebum*.) That which has the nature of suet,—as *sebaceous* or *sebaceofollicular tumours*.

SEBACEOUS FLUX, *Stearrhea*.

SEBACEOUS, SEBIP'AROUS or SEBIF'EROUS GLANDS, *Sebaceous Follicles* or *Crypts*, *Folliculi seba'cei*, *Cryptæ seba'ceæ*, Oil or *Miliary glands*, are small hollow organs, of a rounded or pyriform shape, seated in the substance of the skin, and opening on its surface by a small, excretory duct; furnishing a yellow, unctuous humour, *Smegma cutaneum*, *Sebum cutaneum*, having some analogy with suet, which is destined to lubricate the surface of the body. Follicles of a similar nature exist around the corona glandis of the male, and under the skin of the labia majora and nymphæ of the female. They secrete a sebaceous matter, which emits a peculiar odour; and hence, after the name of one who described them, have been called *Glandulæ odoriferæ Tysoni*, *G. Tysoni*, *Tyson's glands*, and, also, *Glandulæ odoriferæ* seu *seba'ceæ glandis*, *G. cor'onæ pe'nis*, *Folliculi seba'cei cor'onæ glandis*, *Glandulæ odoriferæ præputii*, *Cryptæ præputiales*.

SEBASTOMANIA, (σεβαστος, 'worshipped,' and *mania*.) Religious insanity, Demonomania.

SEBESTENA, *Cor'dia myxa seu sebestena seu Africa'na seu domestica seu obliqua, Sebestena officinalis, Sebesten, Sebesten, Myxa, Cor'vus sanguinea, Prunus sebestena, Smoothleaved Cor'dia, Asyri'ca plum, (F.) Sebestier myxa. Ord. Cordiaceae.* The dark, black fruit of the *Sebestena* possesses glutinous and aperient qualities, and is exhibited, in decoction, in various diseases of the chest.

SEBESTIER MYXA, Sebestena.

SEBIFEROUS, (*sebum*, 'suet, fat,' and *ferre*, 'to carry,') see Sebaceous.

SEBIPAROUS, (*sebum*, 'suet, fat,' and *pario*, 'to bring forth,') see Sebaceous.

SEBORRHAGIA, (*sebum*, 'suet, fat,' and *ρῆγη*, 'a breaking forth,') Stearrhœa.

SEBORRHOEA, (*sebum*, 'suet, fat,' and *ρῆω*, 'to flow,') Stearrhœa.

SEBSTEN, Sebestina.

SEBUM, Pinguedo, Sebum — s. Cutaneum, see Sebaceous glands — s. Palpebrale, *Chassie* — s. Præputiale, *Smegma præputii*.

SECALE, (*secare*, 'to cut,') [*?*] *S. cereale seu barbatum.* The Rye plant. (F.) *Seigle. Ord. Gramineæ.* Rye is chiefly used as an article of diet, particularly in the northern parts of Europe, where a spirit is also extracted from it. The grain is disposed to acnesency; and hence its internal use at first generally produces a laxative effect.

SECALE BARBATUM, Secale — s. Cornutum, Ergot — s. Luxurians, Ergot — s. Maternum turgidum seu tumulentum, Ergot.

SECALIS MATER, Ergot.

SÉCATEUR DES AMYGDALES, (from *secare*, 'to cut,') Amygdalotome — s. par Écrasement, Écraseur.

SECERNENT, (*secernere*, 'to separate,') Secreting.

SECERNING, Secreting — s. Substance of the kidney, see Kidney.

SECESSIO FÆCUM, (*secedere, secessum*, 'to secede,') Defecation.

SECESSUS, Vulva — s. Niger, Melæna.

SÊCHE, Sepia.

SECLUSORIUM, (*secludere, seclusum*, 'to shut up,') Sac.

SECONDARY, (*secundus*, 'the second,') (F.) *Secondaire.* Something that acts subordinately to another; as the *secondary symptoms* of a disease; that is, those which supervene on the primary. A *secondary amputation* is one performed after the constitutional effects of an injury have mainly passed away. See Fever, secondary, and Hæmorrhagia, for Secondary Hemorrhage.

SECONDINES, Secundines.

SECRETA, (pl. of *secretum*.) Same etymon as Secretion. Things or matters secreted or separated from the blood.

SÉCRÉTEUR, Secreting.

SECRETING, *Secreto'rius, Secre'ning, Sec'ernient, Secre'tory or Sec'retory, Excer'nent, (F.) Sécréteur, Sécrétoire.* That which secretes,—as a secreting organ.

SECRETIO, Secretion — s. Lactis, Galactosis — s. Lotii, Uropoiesis — s. Urinæ, Uropoiesis — s. Vicaria, see Vicarious.

SECRETION, *Secre'tio, Apoc'risis, (secernere, secretum, 'to separate,')* An organic function, which is chiefly executed in the glands, and consists in an elaboration or separation of the materials of the blood, at the very extremities of the arterial system, or rather of the vascular secretory system; and which differs in each organ according to its particular structure; hence the formation of different fluids—bile, saliva, urine, milk, &c.

The term is, also, applied to the thing secreted,—*Secre'tum*. The secretions are of three kinds,—*exhalant, follicular, and glandular*. Simple secretions are those which exist ready formed in the blood, and pass out of the vessels by an act of exosmosis, whilst the more complicated are formed from the liquor sanguinis by cell agency. The latter alone are, by some, called *Secretions*.

SÉCRÉTOIRE, Secreting.

SECRETOMOTORY, *Secretomotorius*. An epithet applied to nerves which influence secretion in the manner of excitomotory nerves.

SECRETORIUS, Secreting.

SECRETORY, Secreting.

SECRETUM, see Secretion.

SECT, METHODOICAL, (*secta*, from *secare*, *sectum*, 'to cut,') Methodists.

SECTIO, gen. *Seccio'nis, Sect'ion*. The act of cutting. A division.

SECTIO ALTA, see Lithotomy — s. Anatomica, Dissection — s. Cadaveris, Autopsia cadaverica — s. Cadaveris legalis, Autopsia cadaverica — s. Cæsarea, Cæsarean section — s. Francoiana, see Lithotomy — s. Hypogastrica, see Lithotomy — s. Lateralis, see Lithotomy — s. Legalis, Autopsia cadaverica legalis — s. Mariana, see Lithotomy — s. Nympharum, Nymphotomy — s. Perinealis, Perineal section — s. Rectovesicalis, see Lithotomy — s. Renalis, Nephrotomy — s. Tendinum, Tenotomy — s. Vaginovesicalis, Colpoecystotomia — s. Vesicalis, Lithotomy.

SECTION, *Seccio* — s. Sigaultian, Symphyseotomy.

SECUNDÆ, (*secundus*, 'the second,') Secundines.

SECUNDINÆ, Secundines.

SECUNDINES, *Hys'tera, Douterion, Me'træ, Secun'dæ, Secund'ina, Af'terbirth, Af'terburthen.* (F.) *Arrière-faix, Secundines, Délivré, (from secundus, 'second,')* All that remains in the uterus after the birth of the child, — viz., the placenta, a portion of the umbilical cord, and the membranes of the ovum. These are commonly not expelled till some time after the birth of the fetus; hence their name. See *Délivrance*.

SECUNDUS PROPRIORUM AURICULÆ, Retrahens auris.

SECUS, Sex.

SEDANTIA, Sedatives.

SEDATION, *Seda'tio, (sedare, sedatum, 'to settle or assuage,')* *Pau'sis, Catapau'sis.* The effect induced by sedatives.

SEDATIVES, *Seda'tiva, Pauan'ta, Seda'n'tia, Adec'ta, Catasta'tica, Deprimen'tia, Depress'ants, (F.) Sédatifs, Calmants, Tempéramts.* Medicines which directly depress the vital forces, and which are consequently employed whenever it is necessary to diminish preternaturally increased action. The chief reputed sedatives are Acidum Hydrocyanicum, Acidum Hydrosulphuricum, Tabacum, Missio Sanguinis, and certain gases by inhalation, as azote, carbonic acid, [*?*] carburetted and sulphuretted hydrogen.

SEDEM ATTOLLENS, 'raising the seat,' Levator ani.

SEDES, (*sedeo*, 'to sit,') Anus, Excrement — s. Cruentæ, Dysentery, Hæmatochezia — s. Lactescens, Colicæ flux — s. Procidua, Proctocele.

SEDHÆE, Banguæ.

SEDIMENT, *Sedimen'tum, Lem'ma, Residen'tia, (from sedeo, 'to sink down,')* (F.) *Dépôt.* A deposit formed by the precipitation of some one or more of the substances held in solution or suspension by a liquid. That which is formed in urine, at the time of cooling, has been called *Hypostasis, Hyposte'ma, Residentia, Subsidentia, (F.) Dépôt de l'urine.* This sediment or deposit varies in different states of the system. In calculous affections it is one of the most important objects of attention.

SEDIMENTUM, Sediment—s. Urine lateridium, see Lateritium—s. Urine pituitous, see Furfuraceous.

SEDILIA, (*sedere*, 'to sit,') Nates.

SEIDLITZ, MINERAL WATERS OF. *Seidlitz water*, 'Springs in Bohemia, near Prague, which are simple sodines. They contain sulphate and carbonate of magnesia, chloride of magnesium, sulphates of potassa, soda, and lime, carbonic acid and carbonate of lime. They are employed as purgatives in a multitude of diseases.

ARTIFICIAL SEIDLITZ POWDERS, in no respect, however, resembling the natural ingredients, are formed of *Soda et Potassa tartarica* ʒij, *Soda carbon.* ʒij, in one paper; *Acid. tartaric.* gr. xxxv, in another. For half a pint of water. They are called, also, *Rochelle powders*.

SÉDON BRÛLANT, Sedum—s. *Reprise*, Sedum telephium.

SEDUM, *S. acre* seu *glaciale* seu *minus* seu *vermicula'rè*, (*sedere*, 'I sit,' owing to its attachment to rocks and walls.) *Ille'ebra*, *Helles'ebra*, *Ille'ebrium*, *Vermicula'rè*, *Vermicula'ria*, *Piper mura'lè*, *Sempervivum acre*, *Ai'chryson*, *Er'vum*, Wall pepper, Jack of the Buttery, *Prick'et*, *Wid's beard*, *Stonecrop*, *Biting Stonecrop*, *Small House'leek*, (F.) *Junbarbe acre*, *Orpin brûlant*, *Vermiculaire brûlante*, *Sédon brûlant*, *Ord. Crassulacée*. *Scr. Syst. Decandria* Pentagynia. In its recent state it is very acrid, and proves both emetic and cathartic. Externally, in the form of cataplasm, it produces vesications and erosions. It has been recommended in cancerous and malignant ulcers.

SEDUM RHODIOLA and **S. AL'BUM**, (F.) *Triquetradome*, have similar properties.

SEDUM, *Saxifraga granulata*—s. Album, see Sedum—s. Glaciale, Sedum—s. Majus, *Sempervivum tectorum*—s. Minus, Sedum—s. Rhodiola, see Sedum.

SEDUM TELEPHIUM, *S. telephoi'des*, *Fa'ba cras'sa*, *Cras'sula*, *C. Ma'jar*, *Ille'ebra ma'jar*, *Tele'phium*, *Faba'ria cras'sula*, *Fa'ba cras'sa*, *Anacampt'era*, *A. al'bans* seu *triply'la* seu *vil'ga'ria*, *Orpine*, (F.) *Sédon reprise*, *Grand orpin*, *Fève épaisse*, *Junbarbe des vignes*, *Herbe à la coupure*, *H. aux Charpentiers*. Formerly used as a cataplasm in cuts, hemorrhoids, corns, whitlows, &c.

SEDUM TELEPHOIDES, *S. Telephium*—s. *Vermiculare*, Sedum.

SEED, ([Sax.] *rad.*) Semen. Sperm.

SEGMENTA CARTILAGINEA, (pl. of *segmentum*, 'a segment,') see Trachea.

SEGMENTATION, *Segmentatio*, (*segmentum*, 'a segment,' itself from *ecco*, 'to cut,') The act of being divided into segments or portions. (F.) *Segmentation*, *Fracti'onnement*, *Sill'onnement*. A term applied to the division of the vitellus or yolk into 'segments,' each containing a transparent vesicle, which may be a descendant of the primordial or germ cell.

SEGNIT'IA, *Segnit'ies*, (from *segnis*, 'sluggish,') Sluggishness or torpor in the exercise of a function:—as *SEGNITIES AL'VI*, Torpor of the bowels; *SEGNITIES VIR'UM*, Languor.

SEGNITIES, Segnitia.

SEGRAY, MINERAL WATERS OF. Near Pithiviers, in Gâtinais, in the French department of Loiret, there is a chalybeate spring, which has some reputation.

SEGU, Sago.

SEIDLITZ, Seidlitz.

SEIDSCHÜTZ, SEYDSCHÜTZ or SAID-SCHÜTZ, MINERAL WATERS OF. Bitter springs, in the neighbourhood of those of Püllna and Seidlitz, in Bohemia, which contain chiefly sulphate, nitrate, muriate, and carbonate of magnesia, sulphates of potassa, soda, and lime; carbonate of lime, iron and manganese. The water

is an excellent cathartic, and is well known in Europe.

SEIGLE, Secale—s. *Cornu*, Ergot—s. *Cornu'pus*, Ergot—s. *à Éperon*, Ergot—s. *Ergotée*, Ergot—s. *Ergotisée*, Ergot—s. *Fume*, Ergot.

SEIN, (from *seins*,) Uterus, see Matrua.

SEIRIASIS, (*seirias*,) Corp. de Suleit.

SEIZURE, Attack, see Raptus.

SEIZURES, Hidden, see Hidden seizures.

SEKISKAYAVODKA, see S, it.

SEL, (from *sol*,) Salt—s. *Adm. rille de Léman*, Magnesia sulphas—s. *Admirable puré*, Soda, phosphate of—s. *Ammoniac*, Ammonia maris—s. *Arménien*, Ammonia maris—s. *Arsténial de Moëquer*, Arseniate of protoxide of potassium—s. *de Bonting*, Mercury, iodide of chloride of—s. *de Quincé*, Soda, muriate of—s. *de Tricoune*, Nuxetina—s. *à détacher*, Potasse oxalas—s. *à Égry*, Magnesia sulphas—s. *à Égry*, Tin, muriate of—s. *de Lait*, Sugar of Milk—s. *Marin*, Soda, muriate of—s. *à Opium*, Nuxetina—s. *à Orsilla*, Potasse oxalas—s. *de Prunelle*, Potasse nitras fusus sulphatis panceillo mixtus—s. *de Satorne*, Plumbi superacetas—s. *de Verre*, see Vitrum—s. *Volatil d'Ammoniac*, Ammonia carbonas.

SELAGO, gen. *Selaginia*, Camphoresma Monspeliaca, Lycopodium selago.

SELENE, (*selene*,) Moon, see Nail.

SELENIACI, (*selenuci*, from *selene*,) see Lunatic.

SELENTIASIS, (from *selene*,) Somnambulism.

SELENIASMUS, Somnambulism.

SELENOBLETUS, (*selenoβletos*, from *selene*, and *βlētō*, 'to strike,') Lunatic.

SELENOGAMIA, (*selene*, and *γάμος*, 'marriage,') Somnambulism.

SELENOPLEGE, (*selene*, and *πληγή*, 'a stroke,') see Corp. de soleil.

SELENOPLEXIA, see Corp. de soleil.

SELFABUSE, Masturbation.

SELFHEAL, Prunella.

SELFLIMITED. An epithet applied to diseases which appear to run a definite course, but little modified by treatment,—small-pox, for example.

SELF POLLUTION, Masturbation.

SELIBRA, for *Semilibra*, *Semilibra*, 'half a pound.' Six ounces, Troy; eight Avoirdupois.

SELINITES, (*selinites*,) Wine impregnated with the seeds of the *celavus* or parsley.

SELINUM, (*selinos*,) Apium petroselinum—s. Angelica, Angelica Sylvestris—s. Galbanum, Dubon galbanum—s. Imperatoria, Imperatoria—s. Lutescens, Peucedanum palustre—s. Marsh, Peucedanum palustre—s. Opopanax, Pastinaca opopanax—s. Oreselinum, Athamanta aureoselinum—s. Ostruthium, Imperatoria—s. Palustre, Peucedanum palustre—s. Pastinaca, Pastinaca sativa—s. Peucedanum, Peucedanum—s. Putescens, Angelica sylvestris—s. Sylvestre, Angelica sylvestris, Peucedanum palustre—s. Thysselinum, Peucedanum palustre.

SELLA, 'a seat,' *S. Turcica*—s. Equina, *S. Turcica*—s. Familiaria, Lasanum—s. Obstetricia, Diphrys—s. Sphenoidalis, *S. Turcica*.

SELLA TURCICA, *Eph'ip'ium*, *Fos'sa Pituitaria*, *Sella*, *S. equina* seu *sphenoidalis*, *Pituitary fossa*, *Turkish Saddle*, (F.) *Selle Turcique* ou *du Turc*. A depression at the upper surface of the sphenoid bone, which is bounded, anteriorly and posteriorly, by the clinoid processes, and lodges the pituitary gland. It is so called from its resemblance to a Turkish saddle. The broad rough plate, behind, and somewhat overhanging the *Sella Turcica*, is the *Dorsum Eph'ip'ii*.

SELLE TURCIQUE, *Sella Turcica*—s. *du Turc*, *Sella Turcica*.

SELS NEUTRES, Neutral salts.

SELTZ or **SELTZER**, MINERAL WATERS OF. Cold, acidulous springs, the source of which

is at Seltz, nine leagues from Strashurg, and in the Department of the Bas-Rhin. They contain carbonates of lime, magnesia, and soda, chloride of sodium, and much carbonic acid. They are refrigerant, tonic, diuretic, and aperient. They are not frequented.

ARTIFICIAL SELTZER WATER may be made from *mariatric acid* gr. xxxv, *water* Oj, *white marble* gr. iij. Stop up till dissolved. Add *carbonate of Magnesia*, gr. v, and, after some time, *subcarbonate of soda*, gr. xxxij. Close it till used.

SEMBELLA, *Selibra*.

SEMECARPUS ANACARDIUM, (σημειον, 'a mark,' and καρμος, 'fruit,') *Avicennia tomentosa*.

SEMEIOGRAPHY, (σημειογραφία,) *Semeiographia*, *Semiography*, *Semiotographia*, *Symptomatology*, (σημειον, 'a symptom,' and γραφή, 'a description,') A description of symptoms or signs of disease.

SEMEIOLOGY, *Semeiologie*, *Semiology*, *Semioticé*, *Phænomenologie*, (σημειον, 'a symptom,' and λογος, 'a discourse,') (F.) *Sémiotique*. The branch of pathology whose object is the doctrine of the symptom or signs of disease.

SEMEION, (σημειον,) *Sign*, *Symptom*—s. *Boethematicum*, *Indication*.

SEMEIOSIS, (σημειωσις,) *Semeiosis*.

SEMEIOTICE, (σημειωτική,) *Semeiotique*.

SÉMÉIOTIQUE, *Semeiology*.

SEMEN, gen. *Seminis*, (σπορον, 'to sow,') That which is sown. *Seed* of plants and animals.

SEMEN, *Sperm*—s. *Badian*, *Illicium anisatum*—s. *Cine*, see *Artemisia santonica*—s. *Contra*, see *Artemisia santonica*—s. *Contra d'Alep*, *Artemisia santonica*—s. *Contra vermes*, *Artemisia santonica*—s. *Masculinum* seu *virile* seu *genitale*, *Sperm*—s. *Mulibre*, *Sperm* (of the female)—s. *Sanctum*, see *Artemisia santonica*—s. *Santonici*, see *Artemisia santonica*—s. *Zedoarie*, see *Artemisia santonica*.

SEMENCE, (from *semen*,) *Seed*, *Sperm*.

SEMI, (from ἡμιον, 'half,') *Semi* or *demi*, in composition, universally signifies 'half;' both in French and English.

SEMIANIMIS, *Semimortuus*, *Semivex*, *Semivivus*, (*semi*, and *animus*, 'life,') *Half-living*. *Half-dead*.

SEMI-BULB OF THE FEMALE, *Bulbus vestibuli*.

SEMICANALICULUS, (*semi*, and *canaliculus*, 'a small channel,') *Suleus*.

SEMICANALIS, (*semi*, and *canalis*, 'a channel,') *Suleus*.

SEMICEPHALUS, *Hemicephalus*.

SEMICIRCULI OSSEI, *Semicircular canals*.

SEMICIRCULAR, *Semicircula'ris*, (*semi*, and *circulus*, 'a circle,') (F.) *Demicirculaire*. That which represents the half of a circle.

SEMICIRCULAR CANALS, *Canales* seu *Ductus semicirculares* seu *tubiformes* seu *circulares*, *Semicirculi ossei*, *Funes* seu *Canaliculi semicirculares labyrinthi*, (F.) *Canaux demicirculaires*, are seated in the pars petrosa of the temporal bone, and open into the vestibule, behind which they are situate. They are three in number, of which one is *superior* or *vertical*,—*canalis semicircula'ris verticalis superior*; another *posterior* or *oblique*,—*canalis semicircula'ris verticalis posterior*, and the third *exterior* or *horizontal*,—*canalis semicircula'ris horizontalis*. Their parietes are formed of a bony, compact, hard plate. Their uses in audition are not known: at their termination they have the elliptical arrangements called *ampullæ*; and by both extremities they open into the vestibule, in the sac at its superior part—the *Sacculus Ellipticus*, *Alveus* seu *Utriculus* seu *Sinus communis*, *Sinus* seu *Alveus utriculus*, *Sacculus vestibuli* or *Median sinus*, *Utriculus*. In front of the S. ellipticus, nearer the cochlea, and opposite the foramen ovale, is the *Sacculus sphaericus*, or *Sacculus*.

SEMICIRCULUS EXSCULPTUS, *Semicirculus* cut out. Hiatus diaphragmatis aorticus.

SEMICONGIUS. Half a gallon, or four pints.

SEMICUBIUM, *Semicubium*, *Insectio*, *Insectus*, *Encathisma*, *Escathisma*, (F.) *Demicubium*, (*semi*, and *cubare*, 'to lie down,') A half-bath, or such as receives only the hips or extremities.

SEMICUPIUM, Bath, half, *Semicubium*.

SEMIFIBULÆUS, *Peroneus brevis*.

SEMIHOMO, (*semi*, 'half,' and *homo*, 'man,') *Atropa mandragora*.

SEMIINTEROSSEUS INDICIS, *Abductor indicis*.

SEMIINTEROSSEUS POLLICIS, *Opponens pollicis*.

SEMIILIBRA, *Selibra*.

SEMILLA DEL GUACHARO. Various kinds of hard and dry fruits, found in the stomach of the young Guacharo, a sort of nocturnal bird. The fruits are a celebrated South American remedy in Ague.

SEMILUNAR, *Semiluna'ris*, (F.) *Demilunaire* ou *Semilunaire*, (*semi*, and *luna*, 'the moon,') Having the shape of a halfmoon.

SEMILUNAR CARTILAGES, *Semilunar fibrocartilages*, *Sigmoid cartilages*, *Cartilagines sigmoides* seu *semilunares*, are two fibrocartilages, which exist between the condyles of the os femoris and the articular surfaces of the tibia. The innermost, which is longer from before to behind than transversely, is nearly semicircular. The outermost forms almost an entire circle; an arrangement, in both cases, corresponding to the different surfaces of the tibia. They are thicker at their outer circumference, which is convex, than at the inner, which is concave and very thin. Both are inserted, before and behind, into the spine of the tibia by means of fibrous fasciæ.

SEMILUNAR GANGLION,—*Ganglion abdominale* seu *splanchnicum* seu *transversum* seu *sola're*, *G. surrenal*, (Ch.), belongs to the great sympathetic. These ganglia present considerable variety, and are deeply situate in the abdomen, above and behind the suprarenal capsules. They correspond, posteriorly, to the pillars of the diaphragm and the aorta. They have the shape of a crescent reversed, often surrounded by other ganglia of a smaller size. From their periphery numerous filaments proceed, which go to the formation of the solar plexus.

SEMILUNAR NOTCH OF THE STERNUM, *Fourchette*—s. *Valves*, *Sigmoid valves*.

SEMILUNARE OS, *Lunare os*.

SEMILUNULA UNGUIUM, (*semi*, and *lunula*, dim. of *luna*, 'moon,') See *Nail*.

SEMIMAS, (*semi*, and *mas*, 'a male,') *Eunuch*.

SEMIMASCULATUS, *Eunuch*.

SEMIMASCULUS, (*semi*, and *masculus*, 'male,') *Eunuch*.

SEMIMEMBRANEOSUS, *Semimembranosus*.

SEMIMEMBRANOSUS, (*semi*, and *membrana*, 'membrane,') *Semimembraneus*, (F.) *Ischio-popliti-femoral*, *Ischio-popliti-tibial*, (Ch.) *Semimembraneus*. A muscle situate at the posterior part of the thigh. It is flat, thin, narrow, and aponeurotic in its upper third; broader and fleshy in the middle, and tendinous below. It is attached above to the tuberosity of the ischium; and below—by a tendon, which has three divisions—to the outer condyle of the femur, and to the posterior and inner part of the internal tuberosity of the tibia. This muscle is a rotator inwards, and a flexor of the leg. It can also bend the thigh upon the leg. In standing it maintains the pelvis in position, and can even draw it downwards.

SEMIMORTUUS, (*semi*, and *mortuus*, 'dead,') *Semianimis*.

SEMINA (pl. of *semen*, 'seed') CATAPUTIA MAJORIS, see *Ricinus communis*.

SEMINA FRIGIDA MAJORA. The greater cold seeds. The ancients gave this name to the emul-

five seeds of the cucumber, melon, gourd, and watermelon.

SEMINA FRIGIDA MINORA. The *lesser cold seeds*. The ancients gave this name to the seeds of the lettuce, purslane, endive, and cichory.

SEMINA RICINI VULGARIS, see *Ricinus communis*.

SEMINAL. [from *semen*.] Spermatic—s. Filaments, Spermatozoa—s. Fluid, Sperm—s. Granules, see *Granule*.

SEMINALE MEMBRUM, 'seminal member,' Penis.

SEMINALIS, Spermatic.

SEMINERVOSUS, (*semi*, and *nervus*, 'a sinew,') Semiteninosus.

SEMINEX, (*semi*, and *nex*, 'death,') Semianimis.

SEMINIA MORBORUM, (*seminium*, 'a seed store,') Predisposition.

SEMINIFEROUS, (*semen*, and *fero*, 'I carry,') Spermatorph'orous. An epithet given to the vessels which secrete and convey the seminal fluid.

SEMINIST, [from *semen*.] Animalculist. The term was, also, used for one, who believed, that the new being was formed by an admixture of the seed of the male with the supposititious seed of the female.

SEMINIUM, 'a seed store,' Sperm.

SEMOGRAPHY, Semeiology.

SEMOLOGY, Semeiology.

SEMON. (*semon*,) Sign.

SEMIORBITULARIS, see *Orbicular muscles*.

SEMIOTICE, Semeiotice—s. Faciel, Physiognomy.

SEMIPESTIS, Typhus.

SEMIPEGIA, (*semi*, and *pegyn*, 'a stroke,') Hemiplegia.

SEMISEXUM, Hemiection.

SEMISICHTICUS. A weight of one drachm or three scruples.

SEMISIDERATIO, (*semi*, and *eideratus*, 'planet struck,') Hemiplegia.

SEMISIDERATUS, see *Hemiplegia*.

SEMISOMNIS, (*semi*, and *somnus*, 'sleep,') Coma.

SEMISOPITUS, (*semi*, and *sopio*, *sopitum*, 'to put to sleep,') Coma.

SEMISOPORUS, (*semi*, and *sopor*, 'sleep,') Coma.

SEMISSPECULUM, (*semi*, and *speculum*, 'a mirror,') An instrument used for dilating the incision made into the neck of the bladder in the operation of lithotomy.

SEMISSPINALIS COLLI. *Semispinalis seu transversospinalis seu spinalis seu transversalis colli*, *Articulospinalis*, *Spinalis cervicis*, *Transverso-spinalis*, (F.) *Demi-épineux ou transversaire-épineux du cou*. This muscle arises from the transverse processes of the six uppermost vertebrae, by an equal number of distinct tendons, which run obliquely under the complexus, and is inserted into the spinous processes of all the cervical vertebrae, except the first and last. Its action is, to extend the neck obliquely backwards and to one side.

SEMISSPINALIS DORSI. *Semispinalis externus*, *Transverso-spinalis dorsalis*, *Semispinalis* (*Riolan*), *Transverso-spinalis*, (F.) *Demi-épineux du dos*, *Transverso-épineux du dos*. This muscle arises from the transverse processes of the seventh, eighth, ninth, and tenth dorsal vertebrae, by as many distinct tendons, which soon grow fleshy; and then again become tendinous, and are inserted into the spinous processes of the sixth or seventh uppermost dorsal, and two lowest cervical vertebrae, by as many tendons. Its action is, to extend the spine obliquely backwards.

SEMISSPINALIS EXTERNUS, *Semispinalis dorsi*—s. *Internus*, *Transversalis dorsi*—s. *Spinatus*, *Longissimus dorsi*, *Semispinalis dorsi*.

SEMISSPINATUS COLLI. *Semispinalis colli*.

SEMISSIS, *Se'mis*. The half of a pound, ounce, drachm, &c.

SEMITENDINOSUS. *Semiteninosus*, (F.) *Ischio-crètibial*, *Ischio-prètibial*, (Ch.) *Demi-tendineux*, *Demi-nerveux*. This muscle is situate at the posterior part of the thigh. It extends obliquely, downwards and inwards, from the tuberosity of the ischium—to which it is attached by a flat tendon, common to it and the biceps—as far as the posterior and inner part of the tibia, to which it is attached by means of another round tendon. Its fibres are commonly interrupted by an aponeurosis. Its uses are the same as those of the semimembranosus.

SEMITERTIANA, (*semi*, and *tertiana*, 'tertian,') Hemitritæa.

SEMIUNCIA, (*semi*, and *uncia*, 'an ounce,') Hemionchion.

SEMIVIR, (*semi*, and *vir*, 'a man,') Eunuch.

SEMIVIVUS, (*semi*, and *vivus*, 'alive,') Semianimis.

SEMOLA, *BUL'LOCK'S*: a farinaceous article of diet, which is recommended as highly nutritious and easily digestible, consequently well adapted for infants and invalids, is said to consist of the gluten of wheat, with a small proportion of starch. *Semola* and *Semolina* resemble each other greatly. They appear, indeed, to be identical.

SEMOLINA, *Man'na croup*. This may be made by grinding wheat in an apparatus, separating the flour from the middlings, dressing the latter four different times in a bolting mill, and sifting through parchment sieves, until the product is perfectly freed from bran. In Poland, it is called *Cra'covo grnatz*.

Semolina is generally imported into England from the Baltic, and in Russia is said to be manufactured from buckwheat. It has a granular form like some kinds of oatmeal, a dingy white colour, a farinaceous, somewhat insipid, taste, and is well adapted for the diet of children affected with derangement of the bowels. A recent analysis makes it consist of the gluten of wheat with a certain proportion of the starch, part of the latter having been removed.

SEMPERVIVUM ACRE, (*semper*, 'always, and *vivus*, 'alive, because always green,') *Sedum*.

SEMPERVIVUM TECTORUM, *S. majus seu tectorum*, *Aizoon*, *Beitahabalen*, *Euphthalmum*, *Bor'ba Jo'eis*, *Agrosia'nara*, *Andrachakura*, *Chrysomer'mum*, *Honzelek*, *Sen'green*, *Fenz*, *Fowet*, (Prov.) *Cyphe*, *Jupiter's Beard*, (F.) *Jombarte des toits*. The leaves have a mild, subacid austerity, and are often applied to bruises and old ulcers.

SEMPRESSSES' PALSY, see *Palsy*, *sempresses*.

SEMUDA, (*semuda*,) *Betula*.

SEMUNCIA, Hemionchion.

SENA, *Cassia senna*.

SÉNÉ, *Cassia senna*—s. *d'Amérique*, *Cassia Marilandica*—s. *Faux*, *Colutea*—s. *Indigène*, *Colutea*—s. *des Prés*, *Gratiola officinalis*.

SENEBIERA CORONOPUS, (after J. Senebier, the naturalist, of Geneva,) *Cochlearia coronopus*.

SENECA OIL, (after the Seneca Indians,) *Petroleum*, (New York.)

SENECIO, *S. vulgaris*, (*senex*, 'an old man,' in allusion to the grayish down on many species,) *Erig'erum*, *Cor'talon*, *Erig'erum*, *Ground'el*, *Fire'weed*, (So.) *Gravillea Siculica*, (F.) *Senequm canadense*. Ord. Composite. Soc. Syst. *Syngenesia Polygamia Superflua*. A common plant in Europe; frequently applied, bruised, to inflammations and ulcers, as a refrigerant and antiseptic.

SENECIO AU'REUS, *Golden rag'weed*, *Wild carle'rian*, *Life root*, *Squire root*; indigenous, is an

excitant, diaphoretic, and diuretic. The dried plant is used in decoction:— $\frac{3}{4}$ ss to Oj of water.

SENECIO HIERACIFOLIUS, *Erechtithis hieracifolius*.

SENECIO JACOBÆA, *Jacobæa*, *St. James's Wort*, *Ragwort*, (Sc.) *Buniede*, *Wesbo*, (F.) *Jacobée*. The leaves have a roughish, bitter, subacid, and extremely nauseous taste. A decoction has been used in dysentery. A poultice of the leaves has been recommended in rheumatic affections: and a decoction of the root has often been employed in cases of wounds and bruises.

SENECON COMMUN, *Senecio*.

SENECTA, *Senectus*.

SENECTA ANGIUM. The cast skin of a serpent. A decoction of it is said to cure deafness!

SENECTA DECREPITA, *Decrepitude*—s. *Extrema*, *Decrepitude*—s. *Summa*, *Decrepitude*—s. *Ultima*, *Decrepitude*.

SENECTUS, *Senect'a*, (from *senez*, 'an old person') *Ge'ras*, *Se'nium*, *Æ'tas sen'ilis* seu *pro'ecta* seu *ma'ta*, *Old age*, *Senility*, (F.) *Vielliesse*. The last period of life, commencing, according to some, at 60 years of age.—according to M. Flourens, at 70,—but varying according to numerous circumstances. It is characterized by progressive diminution of the physical and moral faculties.

SENECTUS ULTIMA, *Decrepitude*.

SENEGA, *Polygala senega*.

SENEGAL, GUM, *Gum'mi Senegalen'se* seu *Sen'cea* seu *Sen'ega* seu *Sen'ica*. The exudation from *Aca'cia* seu *Mimo'sa Sen'egal*.—*Ord.* Leguminosæ. *Sec. Syst.* *Polygama Monocia*—growing in the country of Africa through which the river Senegal runs. It is in loose or single drops, much larger than gum Arabic, or the gum which exudes from the cherry tree; and is much more difficult of solution than the former.

SENEGIN, see *Polygala senega*.

SENEGRAIN, *Trigonella fœnum*.

SENEKA, *Polygala senega*.

SENEVÉ, *Sinapis*.

SENGREEN, *Sempervivum tectorum*.

SENILE, *Sen'i'lis*, (F.) *Sénile*, (from *senez*, 'an old person.') Relating or belonging to old age:—as '*senile delirium*;' dotage.

SENILITY, *Senectus*.

SENNA, *Cassia senna*—s. *Alexandrina*, *Cassia senna*—s. *American*, *Cassia Marilandica*—s. *Bladder*, *Colutea arborescens*—s. *Essence* of, prepared, *Selway's* see *Infusum sennæ compositum*—s. *Germanica*, *Colutea*—s. *Italica*, *Cassia senna*—s. *Prairie*, *Cassia chamaecrista*—s. *Wild*, *Cassia chamaecrista*, *Cassia Marilandica*.

SENS, *Sense*,—s. *Genital*, *Appetite*, *venereal*.

SENSA'TION, *Sensa'tio*, (*sentire*, *sensum*, 'to feel;') *Æsthe'ma*. The consciousness or cognizance by the brain of an impression caused by an external body on the organs of the senses. (F.) *Sentiment*. It is not necessarily, however, confined to bodies external to us; for we can receive an impression by touching any part of our own body. The body which communicates the impressions needs but to be external to the part impressed. Sensations are divided into *external*, *internal*, and *morbid*. The *external* are communicated by the five organs of the senses. The *internal* are such as occur within the body, and arise from some alteration in the function of the part, for the time being. Hunger and thirst are internal sensations, as well as all the natural wants. *Morbid* sensations may be either internal or external. *Objective sensations* are produced by impressions on the peripheral nerves; as in vision, audition, &c. *Subjective sensations*, are such as originate centrally or in the encephalon,—as *tinuitus aurium*.

SENSATIONAL, *Sentient*.

SENSE, *Sen'sus*, *Sen'sio*, *Æsthe'sis*, (F.) *Sens*,

A faculty, possessed by animals, of appreciating impressions from external objects. The senses are five in number; sight, hearing, smell, taste and touch. Of these, the first two and the last administer more to the intellect: and, hence, have been called, by some, the *intellectual senses*; the other two seem more connected with the nutrition of the body: and, hence, have been termed *corporeal* or *nutritive senses*. Other senses have been suggested—as the intense sensation experienced during the venereal act; a sense of heat and cold; a muscular sense, a common sense or cœnæsthesia, &c.

SENSES, NERVOUS SYSTEM OF THE, *Medulla oblongata*.

SENSIBILITAS, *Sensibility*—s. *Anomala*, see *Irritable*—s. *Aucta*, see *Irritable*—s. *Morbosa*, see *Irritable*.

SENSIBILITY, *Sensibil'itas*, (*sentire*, *sensum*, 'to feel,' and *habilis*, 'able,') *Æsthe'ma*. The faculty of receiving and transmitting impressions, and having the consciousness of them. It thus includes three secondary properties, *Impressibility*, ([F.] *Impressionabilité*, *Impressibilité*), *Transmissibility*, and *Perceptibility*. Bichat defines it as the property possessed by living bodies, of receiving impressions, whether the individual be conscious of them or not. In the former case, where conscious, he calls the sensibility *animal*: in the latter, where not, he terms it *organic*. This last is common to vegetables and animals, and presides over nutrition, absorption, exhalation, secretion, &c. The other does not exist in vegetables: it is the origin of the sensations—olfaction, vision, gustation, audition, thirst, hunger, pain, &c. There are few parts of the animal body, which are not sensible—if not in health—in disease. The free extremities of the hair and nails, and the epidermis, are not so. See *Insensibility*.

SENSIFEROUS, (*sensus*, and *ferre*, 'to carry,') *Sentient*.

SENSIO, *Sense*.

SENSITIVE, *Sensiti'vus*, (F.) *Sensitif*. Having the faculty of feeling. Relating or appertaining to the senses, or to sensation.

SENSORIUM, *Sensorium*.

SENSORIVOLITIONAL, A term applied to nervous fibres which pass to and from the cerebro-spinal axis, and are respectively concerned in sensation and volition.

SENSORIAL, *Sentient*.

SENSORIAL POWER, according to Dr. Wilson Phillip, is composed of the two functions of sensation and volition. Dr. Darwin included in it the power of muscular contraction; but it ought not to be extended to acts in which there is not consciousness.

SENSORIUM, *S. commu'nē*, *Org'anism senso'rium inter'num*, *Sensito'rium*, *Empo'rium spiri'tuum*, *Æsthe'terion*, *Sen'sory*. The common centre of sensations.

SENSORY, *Sensorium*, *Sentient*.

SENSORY GANG'LIA. A name given by Dr. Carpenter to a series of ganglionic masses at the base of the brain, which are in direct communication with the nerves of sensation—as the *olfactory*, *optic*, *auditory*, and *gustatory*.

SENSORY NERVES, *Nerves of sensa'tion*, are so called in contradistinction to *Motor* nerves. There are *general sensory nerves*—as those connected with the posterior part of the spinal marrow, and the fifth pair; and *special sensory*, as those of the senses.

SENSUS EGREDITUDINIS, 'a feeling of sickness,' *Indisposition*—s. *Osmometricus*, *Olfaction*—s. *Perversus*, *Pseudæsthesia*.

SENTIENT, *Sen'tiens*, (from *sentire*, 'to feel,') *Sen'sory*, *Ideā'y'ismus*, *Sensorial*, *Sensational*, *Sensiferous*. Feeling; causing feeling.

SENTIENT EXTREMITIES OF NERVES are their minute terminations in the organs.

SENTIENT, see Sensation.

SENTINA, Pituitary gland.

SEPARATORIUM, (*separare, separatum*, 'to separate.') A surgical instrument, for separating the pericranium from the skull.

A *Septicatory*, *Vitrum Hypocreticum*, is a pharmacæutical vessel for separating fluids of different densities from each other.

SEPEDOGENESIS, *Sepeidogenesis*.

SEPEDON, (*σηπεδον*.) Putrefaction.

SEPEDONODES, (*σηπεδονωδης*.) Putrid.

SEPEDOGENESIS, *Sepeidogen'esis*, (*sepeidon*, and *genesis*, 'generation.') A septic disposition or tendency; such as is met with in typhus gravior.

SE'PES, *Se'pes*, 'a hedge,' 'a row,' as *Se'pes* seu *Se'ries* seu *Pec'ten dentium*:—a row of teeth.

SE'PES DENTIIUM, see *Sepes*.

SEPHIRUS, *Scleriosis*.

SE'PIA, (*sepi*, *S. officinalis*, *Se'pium*, Cuttlefish, (*Se*.) *Hussfish*, *Sliverfish*, (*F*.) *Sèche*. *Class*, Cephalopoda. The osseous part of the cuttlefish: (*F*.) *Os de Sèche*—*Præcipitans magnum*—has been used as an absorbent, and is often added to toothpowders.

SEPLASIA'RIUS. A name formerly given to one who sold perfumes, ointments, &c. (It is derived from *Seplasia*, the name of a public place at Capua, where such preparations were frequently sold.) A druggist.

SEPSICHY'MIA, *Septochymia*.

SEPSIS, (*σηψις*.) Putrefaction.

SEPTÆ'MIA, *Septha'mia*, *Septicæ'mia*, (*σηπτος*, 'putrid,' and *αἷμα*, 'blood;') *Putrid infection*. A morbid condition of the blood produced by septic or putrid matters.

SEPTAL, *Septa'lis*, (from *Septum*.) Relating, or appertaining to, a septum: as 'septal walls of the heart'—those that separate the cavities from each other.

SEPTANA FEBRIS, (*septem*, 'seven,') Fever, septan.

SEPTENARY, *Septena'rius*, (*F*.) *Septenaire*, (*septem*, 'seven,') 'Consisting of seven.' The septenary years of the old believers in critical periods were times at which, it was supposed, important changes might be expected to take place. The grand climacteric was fixed at 63, and it was considered that if a person passed that age, he had wellfounded expectations that his life might be protracted to 90.

SEPTENNAD, (*septem*, 'seven,') Climacteric, (years.)

SEPTERIUS, (*σηπτηριος*.) Septic.

SEPTIFOLI, UPRIGHT, (*septem*, 'seven,' and *folium*, 'a leaf,') *Tarmentilla*.

SEPTHÆ'MIA, *Septæmia*.

SEPTIC, *Septicus*, *Se'p'tus*, *Septe'rius*, (*F*.) *Septique*, (*σηπτος*, 'putrid,') That which produces putrefaction. A substance which corrodes and disorganizes the soft parts without causing much pain.

SEPTIC Poison, see *Poison*.

SEPTICÆ'MIA, *Septæmia*.

SEPTIMESTRIS FŒTUS, (*septem*, 'seven,' and *mensis*, 'a month,') see *Fœtus septimestris*.

SEPTINERVIA, (*septem*, 'seven,' and *nervus*, 'a nerve,') *Plantago*.

SEPTOCHY'MIA, *Sepeichymia*, (*σηπτος*, 'putrid,' and *χυμος*, 'juice,') Putrefaction or putrescency of the humours.

SEPTON, from *σηπτος*, 'putrid,') *Azote*.

SEPTOPYRA, (*σηπτος*, 'putrid,' and *πυρ*, 'fever,') Typhus gravior.

SEPTULUM, (dim. of *septum*.) A division between small spaces or cavities—as the *septula* or fibrous cords given off by the mediastinum testis to be inserted into the inner surface of the tunica albuginea.

SEPTUM, (*F*.) *Cloison*, 'a partition,' (from *sepio*, *septum*, 'I fence in,') A part destined to separate two cavities from each other, or to divide a principal cavity into several secondary cavities.

These septa are numerous in the human body. The chief are the following:—

SEPTUM ATRIORUM, *S. Auriculærum*. The septum or partition between the auricles of the heart.

SEPTUM ATRICULARUM, *S. Atriorum*—*s. Cerebelli*, *Falx cerebelli*—*s. Cerebri*, *Falx cerebri*—*s. Cochleæ auditoriæ*, *Lamina spiralis*.

SEPTUM COR'DIS, *Septum medium cordis*. The partition which separates the two ventricles of the heart.

SEPTUM CRURALE, see *Crural canal*—*s. Encephali*, *Tentorium*.

SEPTUM LUCIDUM seu *pellucidum* seu *tenue et pellucidum* seu *medium*, *Speculum*, *S. lucidum*, *La'pis specularis*, *Mediastinum* seu *Diaphragma ventriculorum interitium cerebri*, (*F*.) *Septum median* (*Ch*.), *Cloison transparente*. The soft portion or medullary substance, which separates the two lateral ventricles of the brain from each other. This septum is composed of two lamina; between which a small cavity exists, filled by a serous fluid, *ventriculus septi pellucidi*: see *Ventricles of the Brain*. The brothers Wenzel are of opinion that this cavity is lined by a peculiar membrane.

SEPTUM MEDIAN, *S. lucidum*—*s. Medium du cercelet*, *Falx cerebelli*—*s. Medium cerebri*, *S. lucidum*.

SEPTUM NARICUM, *Interseptum* seu *Dissepimentum* seu *Interitium* seu *Discrimen* seu *Ambræ seu Diaphragma naricum*, (*F*.) *Cloison des Fosses nasales*. The partition between the nares. It is formed by the vomer, the perpendicular plate of the ethmoid bone, and a cartilage of a triangular shape. These parts are lined by the pituitary membrane, which receives a considerable number of nerves and vessels.

SEPTUM NASI, see *Nares*.

SEPTUM PARVUM OCCIPITALE, *Falx cerebelli*—*s. Pectiniforme*, see *Cavernous bodies*—*s. Pellucidum*, *S. lucidum*—*s. Scala*, *Lamina spiralis*—*s. Staphylin*, *Velum pendulum palati*—*s. Tenue et pellucidum*, *S. lucidum*—*s. Thracis*, *Mediastinum*—*s. Transverse*, *Tentorium*—*s. Transversum*, *Diaphragm*—*s. Vestibuli nervosomembraneum*, see *Labyrinth*.

SEPTUS, (*σηπτος*.) Septicus.

SEPUM, *Sevum*.

SEQUESTRUM, (*sequestrare*, 'to separate from,') (*F*.) *Séquestre*. The portion of bone, in necrosis, which is dead, and separated from the living bone; acting, whilst retained, as an extraneous body. When the sequester is superficial and small, it is called *Exfoliation*. The portion of bone surrounding it is the *Capsula sequestralis*.

SERAPEUM, Syrup.

SERAPINON, *Sagapenum*.

SERAPIUM, Syrup.

SERBET, *Scherbet*.

SÉREUX, Serous.

SERICOCARPUS TORTIFOLIUS, (*σηρικός*, 'silken,' and *καρπος*, 'fruit,') *Unguis bœvæ*, *Aster tortifolius*; indigenous; *Order*, *Compositæ*: is said to be diuretic.

SERICUM, (*σηρικον*.) *Silk*: from *Sera*—the Thibetans probably—from whom it was first obtained. (*F*.) *Soie*. The thread spun by the *Bombyx Mori* or *Silkworm*. Silk is used by surgeons for ligatures, &c.

SEMICUM ANGLICUM, see *Sparadrapum adhæsivum*.

SERIDION, *Seriola*.

SERIES DENTIIUM, 'a row of teeth' seu *Sepes*.

SERIEUX, Serious.

SERINGUE, Syringe.

SERIOLEA, (*casus*, gen. *acidos*.) *Cichorium intybus*.

SERIOUS, *Sérîus*, (F.) *Sérieux*, Grave. A term applied to a disease. *Dyspathia*, *Morbus gravis*, and to a symptom that is attended with danger.

SERIS, (*sepis*.) *Cichorium intybus*.

SERMOUNTAIN, *Laserpitium album*.

SERO, in composition, Serous membrane.

SEROCOLITIS, see Colitis.

SEROCYSTIC, *Serocysticus*. An epithet for tumours of the female breast, which consist, in the first stage, of one or more membranous cysts, produced perhaps by dilatation of portions of the lactiferous tubes.

SEROENTERITIS, see Enteritis.

SEROHEPATITIS, see Hepatitis.

SEROLIN, *Serolina*, (*serum*, and *oleum*.) A peculiar fatty matter, found by Boudet, in the blood. According to Dr. Austin Flint, jr., normal faeces contain serolin or *sterolin* produced by a transformation of the cholesterol of the bile during the digestive act.

SEROSITE, Serum.

SEROSITY, Serum.

SEROUS, *Serous*, (F.) *Séieux*, (from *serum*.) Thin; watery. Relating to the most watery portion of animal fluids, or to membranes that secrete them. See Serum.

SEROUS LAYER, see *Tache embryonnaire*—s. Membranes, Membranes, serous.

SERPEDO, (*serpere*, 'to creep,') Psoriasis.

SERPENS, Serpent.

SERPENT, *Serpens*, *Anaïs*, *Ophis*, (from *serpere*, 'to creep,') A snake.

SNAKE BROTH was, at one time, recommended as a restorative.

Many serpents are venomous: and the bites of some fatal. See Poisons, table of.

SERPENT A LUNETTES, 'spectacled snake,' *Cobra di Capello*—s. à *Sonnettes*, *Crotalus horridus*.

SERPENTAIRE NOIRE, *Actæa racemosa*—s. *Rouge*, *Polygonum bistorta*.

SERPENTARIA, (from *serpens*, gen. *serpentis*, 'a snake,') 'snake root,' *Aristolochia serpentaria*—s. *Gallorum*, *Arum dracunculæ*—s. *Hispanica*, *Scorzonera*—s. *Minor*, *Arum maculatum*—s. *Nigra*, *Actæa racemosa*.

SERPENTARIE BRAZILIENSIS RADIX, *Cainca radix*.

SERPENTARIN, see *Aristolochia serpentaria*.

SERPENTIN, 'winding as a serpent,' see Alembic.

SERPENTINE, (from *serpens*, 'a serpent,') Ophites.

SERPENTINUS, Ophites.

SERPES, (*serpere*, 'to creep,') Herpes.

SERPIGINOUS, (from *serpere*, 'to creep,') (F.) *Serpigineux*. An epithet given to certain affections, which creep, as it were, from one part to another,—'*Serpiginous erysipelas*,' (F.) *Érysipèle serpiginoux*.

SERPIGO, gen. *Serpinis*, (from *serpere*, 'to creep,') Herpes circinatus, Lichen psoriasis.

SERPILLUM, (from *serpere*, 'to creep,') *Thymus serpyllum*.

SERPOLET, *Thymus serpyllum*.

SERPULLUM, (*εἰπωλλον*.) *Thymus serpyllum*.

SERPULLUM, (*εἰπωλλον*, from *εἰπειν*, *serpere*, 'to creep,') *Thymus serpyllum*—s. *Citratum*, see *Thymus serpyllum*.

SERRA, Saw—s. Amputatoria, Saw, amputation—s. Versatilis, Trepan.

SERRATED, *Serratus*, (from *serra*, 'a saw,')

Dentate, Sawshaped.

SERRATULA AMARA, dim. of *serratus*, 'sawshaped,' applied to the leaves owing to their

serrated shape. Family, *Cynaroecephalæ*. *Ser. Syst.* Syngenesia *Polygamia aequalis*. A species of *Sauvort*, recommended in ague.

SERRATULA BEHEN, *Centaurea behen*.

SERRATUS, (from *serra*, 'a saw,') Dentate.

SERRATUS, ANTIUS MINOR, Pectoralis minor.

SERRATUS MAGNUS. So called from its serrated appearance; (from *serra*, 'a saw,') *Serratus major anticus*, *S. major*, (F.) *Costo-lasi-scapulaire*, *Costo-scapulaire* (Ch.), *Grand dentelé*. A very broad (especially anteriorly), thin, flat, irregularly quadrilateral muscle, situate at the sides of the thorax. It is attached, before, to the external surface of the first 8 or 9 ribs, by as many distinct digitations; and, behind, to the spinal edge of the scapula. The superior fibres are almost horizontal; the lower become gradually more and more oblique upwards and backwards. This muscle carries the scapula forwards, and causes it to execute a movement of rotation, which directs its inferior angle forwards, and the anterior upwards. When the shoulder is fixed, it raises the ribs.

SERRATUS MAJOR, *Serratus magnus*—s. *Major anticus*, *Serratus magnus*.

SERRATUS POSTICUS INFERIOR, (F.) *Dorso-lombo-costal*, *Lombo-costal* (Ch.), *Petit dentelé postérieur et inférieur*. A broader and thinner muscle than the following, but nearly of the same shape. It is situate obliquely at the lower part of the back; and extends from the spinous processes of the last two or three dorsal vertebrae, and the first two or three lumbar, to the inferior margin of the last four false ribs. This muscle depresses the ribs, and thus concurs in expiration. It is an antagonist to the next.

SERRATUS POSTICUS SUPERIOR, (F.) *Cervico-dorso-costal*, *Dorso-costal*, (Ch.), *Petit dentelé supérieur*. A flat, thin, quadrilateral muscle; situate obliquely at the posterior inferior part of the neck, and the superior part of the back. It passes from the posterior cervical ligament, the spinous process of the last cervical vertebra; from the spinous processes of the first two or three dorsal, to the second, third, fourth, and fifth ribs; into which it is inserted by as many digitations. It raises the ribs, and is, consequently, an inspiratory muscle.

SERRE-ARTÈRE, (*serrer*, 'to press,' and *artère*, 'artery,') *Presse-artère*.

SERRE-COU (F.) (*serrer*, 'to press,' and *cou*, 'the neck,') An instrument, invented by Chabert, for compressing the jugular vein, in bleeding from that vessel. It surrounds the neck like a collar, and, by means of a pad, exerts the necessary pressure.

SERRE-FINE (F.), (*serrer*, 'to press,' and *fin*, 'fine,') A small spring forceps, invented by M. Vidal. It is formed of a wire of metal, turned in the middle into two spirals, one in front of the other. This constitutes the spring. Each branch describes nearly the letter S; and one extremity forms the spiral, the other lays hold of the parts to be united.

SERRE-NEUD (F.) (*serrer*, 'to press,' and *nœud*, 'a knot,') This name has been given to various instruments, employed by the surgeon for tightening a ligature passed round a pediculated tumour, or any part which has to be destroyed by ligature slowly and gradually. Thus, there is the *serre-nœud* of Roderic, the *constricteur* of Herbiniaux, the *serre-nœud* of Desault, &c.

SERRE-NEUD DE DESCHAMPS, *Presse-artère*.

SERTULA CAMPANA, (dim. of *serta*, 'a garland,') *Trifolium mellilotus*.

SERUM, 'whey,' (probably from *serus*, 'late,' because it remains after the butter and cheese have been separated from milk,) see Serum Lactis.

Or'chos, O'ros, Seros'ity. (F.) *Sérosité*. The most watery portion of animal fluids, exhaled by serous membranes. It is a constituent part of blood, milk, &c. Also, *Sperm*.

SERUM OF THE BLOOD, *Ha'my'dor, Or'chos, Sé'rum seu P'chor san'guinis*, (F.) *Sérum du Sang*, is the liquid which separates from the blood, when coagulated at rest. It is of a greenish-yellow colour; viscid, slightly coagulable by heat, acids, and alcohol. It is composed of water; chloride of sodium; certain phosphates; and albumen, constantly united to soda, almost in a saponaceous combination.

The fluid which exudes from the albumen of the serum of the blood, when coagulated by heat, is by some called *seros'ity*.

SERUM LAC'TIS, *A'qua Lac'tis; Serum of Milk, Whey*, (Prov.) *Whig*, (F.) *Sérum du lait, Petit lait*. That part of milk from which the butter and caseous matter have been separated. It is a transparent, citrine-coloured liquid, containing sugar of milk, mucilage, acetic acid, phosphate of lime, and some other saline substances. It is used as a slightly nutritious diluent and refrigerant. The French have magistral formulæ, the basis of which is whey. They are called *Petits-laits*, or 'wheys.'

SERUM LAC'TIS HOFFMAN'NI, (F.) *Petit-lait d'Hoffmann*, is a name given to the liquid resulting from treating with boiling water milk which has been evaporated to an almost solid consistence. It is not much used.

REN'NET WHEY is made thus: *milk* 2 pints; *rennet* 3ss, infused in a little hot water; mix and keep in a gentle heat for some hours, then strain.

SERUM DU SANG, S. of the blood.

SERUM SANGUINIS, Serum of the blood.

SERUM OF SE'ROUS MEMBRANES, *Sérum membr'anum serosum, Water of Drop'sy*, (F.) *Sérum des membranes séreuses, Eau des hydro-piques*. It resembles the serum of the blood, especially in cases of hydrocele. The water of hydrothorax and ascites contains much less albumen:—sometimes enough to coagulate on the application of heat: at others, not.

SERVICE, MOUNTAIN, *Sorbus aucuparia*.

SERVICE TREE, *Sorbus domestica*.

SERVIETTE EN CARRÉ, 'a napkin in square,' *Couvrechef*.

SÉSAME, *Sesamum orientale*.

SESAMOID, *Sesamoï'des, Sesamo'des, Sesamoï'deus*, (*σάσαυ*, 'a grain of sesamum,' and *είδος*, 'resemblance.')

SESAMOID BONES, *Os'sa seu Ossic'ula sesamoï-dea*, are small bones, situate in the substance of tendons, near certain joints. Their number is various. They are commonly more in the male than female. Two are constantly met with beneath the metatarsophalangeal articulations of the great toe. Sometimes, the articulations of other bones of the metatarsus with the phalanges of the toes have them likewise. In the hand, there are two before the metacarpophalangeal joint of the thumb. At times, they are met with in the other metacarpophalangeal articulations. Occasionally, two exist behind the condyles of the femur, in the tendons of the gastrocnemii muscles. A considerable os sesamoideum is also met with in the tendon of the peronæus longus. These bones are commonly round, and flattened only on one surface, which is incrustated with cartilage. Their size is very variable, according to the joints in which they exist. Traces of sesamoid bones are not observed in infants; in them, they are replaced by small, cartilaginous concretions. They are developed by a single point of ossification: and are, like the patella (which is a true sesamoid bone), composed of much cellular substance, covered by a thin layer of compact tissue. The

use of these bones is,—to allow the tendon to be inserted into the lever at a greater angle; and thus enable the power to act more advantageously.

Riolan has given the name *Ossa sesamoïdea* to two bony points sometimes met with;—the one at the outer side of the carotid canal of the temporal bone; and the other at the edge of the cavernous sinus, at the side of the internal carotid artery.

SESAMUM, (*σάσαυρον, σάσαυρη*), see *Sesamum orientale*.

SES'AMUM ORIENTA'LE, *S. edu'lē seu oleif'erum seu In'dicum* [?], *Ben'nē, Oil'y Grain*, (F.) *Sésame, Jugeoline*. Ord. Bignoniaceæ. The seeds of this African plant are cultivated in Georgia and South Carolina, where the negroes use them with Indian corn as an article of food. They yield a larger proportion of oil than any other vegetable; one hundred weight producing ninety pounds of oil, which is, at least, equal to Florence oil.

Ben'ne oil, O'leum Ses'ami, the oil of the seed, is used in the Southern States as a laxative. The leaves—*Sesami folium*, (Ph. U. S. 1863)—which are referred to *Sesamum orientale* and *S. Indicum*, afford, by infusion, an excellent mucilage.

SESCUN'CIA, *Sesquim'cia, Hemiol'ion*. An ounce and a half. See *Sesqui*.

SESELI, (*σέσλη*), *Laserpitium siler*—s. *Ægopodium*, *Ligustium podagraria*—s. *Annuum*, *S. tortuosum*—s. *Carum*, *Carum*—s. *Carvi*, *Carum*—s. *Creticum*, *Tordylium officinale*—s. *Feniculifolium*, *Sison ammi*—s. *Graveolens*, *Apium graveolens*—s. *Meum*, *Æthusa meum*—s. *Patens*, *S. tortuosum*—s. *Pratense*, *Peucedanum silaus*—s. *des Prés*, *Peucedanum silaus*.

SES'ELI TORTU'OSUM, *S. Massilien'sē seu annuum seu pa'tens, Enan'the striat'a rig'ida, Hartwort of Marseilles*. The seeds have a warm taste; and a greater degree of pungency than those of *Laserpitium siler*, which is the *Seseli* of the Pharmacopœias.

SESELIS, *Laserpitium siler*.

SES'QUI. This word, joined with any number, weight, measure, &c., signifies one and a half; as *sesquigranum*, 'a grain and a half.'

SESQUIUNCIA, *Sescuncia*.

SE'TA EQUI'NA, 'horsehair.' *Hair'worm, Helmin'thus Gor'dii*. A kind of worm, found in stagnant water, from 4 to 6 inches long, and twisted into various knots and contortions; colour pale-brown, with dark extremities. It is common in the intestines of the Laplanders; causing the *Col'ica Lappon'ica*, in which the gripings are, at times, exceedingly severe.

SETA'CEUM, *Se'ton*, (F.) *Séton*, (from *seta*, 'a bristle.')

A long strip of fine linen or cotton twist, passed through the skin and areolar membrane, to keep up an issue; or through suppurating cavities, between the fragments of fractured bones, &c., to fulfil certain indications. The term is, also, applied to the issue so established. The operation is simple. It consists in laying hold of a fold of skin, and passing a seton-needle through it, with the thread attached. Every day a fresh portion of the thread is drawn through the sore. Setons are established in many diseases;—in obstinate ophthalmia, cephalalgia; epilepsy, thoracic and abdominal affections, &c.; but they are not as much used as formerly.

SETARIA ROCCELLA, (from *seta*, 'a bristle,') *Lichen roccella*.

SETFAST, see *Furunculus*.

SETON, *Setaceum*.

SETERWORT, *Helleborus foetidus*.

SEVADILLA, ([S.] *Sebadilla*), *Veratrum sabadilla*.

SEVATIO, (from *sevom*.) *Stentoma*.

SÈVE, ([Sax.] *ræp*, [G.] *Saft*, 'juice,' *Sap*.

SEVENBARK, *Ilydrangea arborescens*.

SEVENDAYS' DISEASE, see Trismus.

SEVRAGE, (from [L.] *separare*, 'to separate,') Weaning.

SEVUM, *Se'bum*, *Se'pum*, *Ste'ar*, *A'deps ovil'i*, *Se'cum vervecinum*. Suet, mutton suet; Pinguedo.

SEVUM CETI, (*cetus*, 'whale,') Cetaceum.

SEVUM PREPARATUM, *Se'vum*, (Ph. U. S.),—*Ovil'i se'cum preparatum*. *Prepa'rad Su'et*, (F.) *Graisse de Mouton*, *Suif*, is formed by cutting suet in pieces; melting over a slow fire, and straining through linen. It is emollient, and is sometimes boiled in milk (Ξij to Oj), and is used in diarrhœa and dysentery. Its principal employment is in the formation of plasters.

SEVUM VERVECINUM, *Sevum*.

SEX, *Sex'us*, *Sex'us*. The physical difference between male and female in animals and vegetables:—as the *Male sex*, *Sex'us pot'ior* seu *for'tior* seu *vir'ilis*; and the *Female sex*, *Sex'us sequ'ior*, *S. mul'iebris*, *Al'ter Sex'us*, 'The Sex.'

SEXTANS, (from *sextus*, 'the sixth;') *Sex'ta pars li'bre*. The sixth part of a pound.

SEXTARIUS, (from *sextus*, 'the sixth,') An ancient measure for liquid and solid substances; the sixth part of a congius. See Weights and Measures.

SEX'TULA, (from *sextus*, 'the sixth,') *Hexa'gum*, *ἑξάγων*. The sixth part of an ounce.

SEX'UAL, *Sex'ual*, (F.) *Sexuel*, (*sexus*, 'sex,') That which relates to the sex; which characterizes the sex.

SEXUAL DISEASES, *Mor'bi sexua'les*, are the diseases of the genital organs; as SEXUAL FUNCTIONS are the genital functions.

SEXUAL INTERCOURSE, Coition—s. Organs, Genital organs.

SEXUS, gen. *Sexūs*, *Sex*—s. Alter, see Sex—s. Fortior, see Sex—s. Muliebris, see Sex—s. Potior, see Sex—s. Sequior, see Sex—s. Virilis, see Sex.

SEYDSCHÜTZ WATER, Seidschütz water.

SHACKLE, *Carpus*.

SHACKLEHAMMED, see Cnemosciosis.

SHAD'DOCK, *S. Or'ange*. The fruit of *Citrus decumana*, called after the captain of a vessel who first took it to the West Indies from China. It is large; but inferior to the orange in flavour. It keeps, however, longer; and, hence, is useful on long voyages.

SHAGBARK, see Hickory.

SHAKERS, see *Convulsionnaire*.

SHAKES, Intermittent.

SHAKING, Intermittent, see Rigor.

SHALLOT, *Echalotte*.

SHAM ABRAHAM, see Abraham man.

SHAMPOO'ING, *Cham'pooing*, *Mass'ing*, (F.)

Massage, *Massément*. A Hindoo process, which consists in pressing the body of the bather in a peculiar manner; cracking the various joints, and employing blows and friction. A variety of this was employed by the ancients in their baths. With the Romans, the operators were called *Fricatores* and *Tractatores*.

'Kneading' consists in pressing alternately upon a part of the body to arouse the muscles to more energetic action. It is employed at times in dyspepsia.

SHAMROCK, INDIAN, (Ir. *Seamrog* or *shamrog*.) *Trillium latifolium*—s. Water, *Menyanthes verna*.

SHANKER, *Chancre*.

SHAN'NONDALE SALINE' SPRINGS. These are within a few miles of Charlestown, Jefferson County, Virginia. They are mildly aperient and diuretic.

SHAPE, *Vulva*.

SHARON SPRINGS, NEW YORK. These springs, which are situate about 10 or 12 miles

south of Canajoharie, N. Y., and 45 miles west of Albany, are much frequented. They contain, according to the analysis of Dr. Chilton, sulphate of magnesia, sulphate of lime, chloride of sodium, chloride of magnesium, sulphohydrate of soda, sulphohydrate of lime, vegetable extractive matter, and free sulphohydric acid gas. Their temperature is 48° Fah.

SHEATH, ([Sax.] *reāse*.) Vagina.

SHEDDING OF THE TEETH, see Dentition.

SHEEP POISON, *Kalmia latifolia*.

SHELLBARK, see Hickory.

SHELL FLOWER, *Trillium latifolium*—s. Liquid, see *Calci murias*—s. Membrane, *Membrana testæ*.

SHEPHERD'S PURSE, *Thlaspi bursa*—s. Sun dial, *Anagallis arvensis*.

SHERBET, *Scherbet*—s. Orange, see *Syrupus aurantii*.

SHERMAN'S WORM LOZENGES, See Worm lozenges, *Sherman's*.

SHERRY, (after *Xeres*, in Spain,) Sack, see Wine.

SHIELDBONE, *Scapula*.

SHIELD, WATER, *Nelumbium luteum*.

SHILDER, (*shoulder*), *Humerus*.

SHIN, Sax. *reina*, *O'crea*, *Cris'ta tib'ie*. The epine, or anterior part of the tibia or leg. It is, also, called *Anticnemion*, and *O'crea*.

SHINBONE, *Tibia*.

SHINGLES, (*cingulum*, 'a girdle,') *Herpes zoster*.

SHINLEAF, *Pyrola*.

SHINLOCK, *Brassica eruca*.

SHIP FEVER, see Typhus.

SHITABED, *Leontodon taraxacum*.

SHIVERING, (from [G.] *Schauer*,) see Rigor.

SHOCK, (from [Sax.] *reacan*, *reacan*, 'to shake,') (F.) *Choc*, see Concussion.

SHOCK, NERVOUS, *Vi'tal Shock*, *Vi'tal Depression*, *Ner'vous Depression*, *Fu'tal Sink'ing*. Sudden or instantaneous depression of organic, nervous, or vital power, often with more or less perturbation of body and mind, passing either into reaction, or into fatal sinking, occasioned by the nature, severity or extent of an injury, or by an overwhelming moral calamity.—(Copland.)

SHOCK, VITAL, Shock, nervous.

SHOOT, see Crik-in-the-neck, and Pain.

SHORT, ([Sax.] *recept*, from *repan*, 'to cut, to shear,' *shored*, *shor'd*, *short*,) (F.) *Court*. A word used, by anatomists, in opposition to *long*; and to distinguish parts from each other that have otherwise the same name.

SHORT BONES, for example, are those in which the three dimensions—length, breadth, and thickness—are nearly alike; such are the vertebrae, bones of the carpus, tarsus, &c.

SHORT-OF-PUFF, Shortwinded—s. Ribs, see Costa.

SHORTSIGHTED, *Myopic*.

SHORTSIGHTEDNESS, *Myopia*.

SHORTWINDED, *Pursy*.

SHOTBUSH, *Aralia spinosa*.

SHOTT, (*shoot*), *Pleurodynia*.

SHOULDER, ([Sax.] *reuldræ*) *Humerus*—s. Blade, *Scapula*—s. Bone, *Humerus*.

SHOUP, *Haunch*.

SHOVEL PICKEREL WEED, *Unisema deltifolia*—s. Pond, *Unisema deltifolia*.

SHOW, see Parturition.

SHOWCHOO, see Spirit.

SHOWERBATH, see Bath.

SHRUB, (perhaps, from Arabic, *sharab*, 'syrup, or corr. from *syrup*.) A compound of spirits, lemonjuice, and sugar.

SHRUB, SWEET, *Calycanthus*—s. Sweetscented, *Calycanthus*.

SHRUMPSHOULDERED, see Hump.

SHUD'DERING, *Frem'tus*, (from [G.] Schaud'er, Schauer.) (F.) *Frémissement*. *Tremblement*. A peculiar sensation, felt either externally or internally; and which seems to be the result of a spasmodic movement of the parts in which it occurs. A slight feeling of cold sometimes accompanies it. It is, at times, the result of a moral cause, and is often the precursor of shivering.

SHULLBANE, *Humerus*.

SHUMAC, *Rhus coriaria*.

SIAGANTRITIS, (*siagon*, αντρον, 'a cavity,' and *itis*.) Inflammation of the lining membrane of the antrum of Highmore.

SIAGON, (*σάγων*.) Maxillary bone.

SIAGO'NAGRA, (*siagon*, and *αγρα*, 'a seizure,') *Arthritis maxilla'ris*. A gouty or rheumatic affection of the joint of the lower jaw.

SIALACHUS, (from *sialon*.) Salivation.

SIALADENITIS, (*sialon*, αδην, 'a gland,' and *itis*.) Inflammation of a salivary gland. At times, improperly written *Syaladenitis*.

SIALAGOGUE, *Sialogógus*, *Sial'ogogue*, *Ptyalagogue*, *Pty'smagogue*, *Salivans*, *Saliva'tum*, *Sialo'met'icus*, *Apophlegmat'icans per os*, (F.) *Salivant*; (*sialon*, and *αγω*, 'I drive,') That which provokes the secretion of saliva. Pyrethrum and mercury are sialogogues. Sialogogues may be of two kinds, those which act through the circulation, and those which act immediately on the salivary organs. Mercury belongs to the former,—pyrethrum to the latter class. The chief sialogogues are armoracia, calamus, mezereum, pyrethrum, tabacum, and zingiber.

SIALICA, (from *sialon*.) Medicines, which affect the salivary glands.

SIALISMUS, (*σιαλισμος*.) Salivation.

SIALOCHOUS, (*σιαλοχος*, from *sialon*, and *χω*, 'to pour,') Salivated.

SIALOCINETICUS, (*sialon*, and *κινεω*, 'to move,') Sialogogue.

SIALOLITHI, (*sialon*, and *λιθος*, 'a stone,') Calculi salivary.

SIALOLOG'IA, (*sialon*, and *λογος*, 'a discourse,') A treatise on saliva.

SIALON, (*σιαλον*.) Saliva.

SIALORRHEA, (*sialon*, and *ῥεω*, 'to flow,') Salivation.

SIALOS, (*σιαλος*.) Saliva.

SIALOS'CHESIS, (*sialon*, and *σχεσις*, 'retention,') Retention or suppression of the saliva.

SIALOSYRINGES, (*sialon*, and *συριγξ*, 'a pipe,') Salivary fistule.

SIALOZEMIA, (*sialon*, and *ζημια*, 'loss,') Salivation—s. Mercurialis, Salivation, mercurial.

SIAMESE TWINS, see *Twins*, Siamese.

SIBARE. Phrenitis.

SIB'BENS, *Sib'bens*, *Frambo'sia Sco'tica*. An infectious disease in the mountainous parts of Scotland, viewed to be of the venereal kind.

Sib'ens, in the Keltic, means raspberry; and it is said that, in this affection, fungi appear like raspberries. From *sib'ens*, the inhabitants have made *sib'ens*, *sibbens*, and *sibbens*. By some, this affection is regarded as a complication of the venereal and itch. In the Orkneys, *sibbens* means itch.

SIBILANT, *Sib'ilus*, *Sib'ilans*, (*sibilo*, 'I hiss, I whistle,') Making a hissing or whistling sound. See *Râle sibilant*.

SIBILISMUS AURIUM, 'hissing or whistling of the ears,' see *Tinnitus aurium*.

SIBILUS AURIUM, see *Tinnitus aurium*.

SICCANTIA, (*siccave*, *siccatum*, 'to dry,') *Xeranth'ica*, (F.) *Siccatis*. Drying medicines.

See *Desiccative*.

SICCATIFS, *Siccantia*.

SICCATIO, Drying.

SICCHA'SIA, (*σικχασια*, 'disgust,') A distress-

ing disgust for food; such as is experienced by women with child.

SICIL'CIUM, *Sic'lium*. A Roman weight of about two drachms.

SICK, ([Sax.] *reoc*.) *E'ger*, *Egro'tus*, *Nos'e-rus*, *Nos'o'des*, *Ja'cens*, *Labori'sus*, *Labo'rans*, (F.) *Malade*. Labouring under disease: *Diseased*, *Dis'tem'pered*, *Ill*, *Ail'ing*, *Disor'dered*. Such is its acceptance in old English, and generally in the United States. In most parts of the United States, if a patient is affected with a slight indisposition, he is said to be *sick*; if with one more severe, he is said to be *ill*; (Prov.) *bad*, *badly*. In England, it most commonly means,—affected with disorder of the stomach, or nausea.

A sick person, *E'ger*; *Egro'tus*, (F.) *Malade*, who is under the charge of a physician, is said to be a *pa'tient*, or the patient of the physician. At times, but rarely, patient is used for a sick person in the abstract, (from *patiens*, 'suffering,')

SICK STOMACH, Milk sickness, Nausea.

SICK'LY, *Morbo'sus*, *Valetudina'rius*, *Adym'a-tus*, *Nosac'erus*, *Nos'o'des*, *Mor'bose*, *Mor'bulent*, *Unhealth'y*, (Sc.) *Hough*, (F.) *Maladif*. Subject to be sick, or in feeble health. Ailing.

SICKNESS, Disease, Plague—s. Falling, Epilepsy—s. Mountain, Puna—s. River, Milk sickness—s. Sea, Nausea marina—s. of the Stomach, Vomiting—s. Swamp, Milk sickness.

SICLIUM, *Siciliu'm*.

SICUA, (*σικυα*.) Cuppingglass.

SICULA, (dim. of *sica*, 'a dagger,') Beta, Penis.

SICYEDON, (*σικυηδον*.) Caulledon.

SICYOS, (*σικυος*.) Cucumis sativus.

SIDA ABUTILON, *Abutilon Avicennæ*.

SIDE, PAIN IN THE, Pleurodynia.

SIDERA'TIO, *Sydera'tio*, *Sidera'tion*, (*sidus*, gen. *sid'ris*, 'a star,' because the condition was thought to be produced by the influence of the stars.) The state of one struck suddenly, without apparent cause, and as if by the influence of the stars or planets—*Plan'et struck*. The ancients comprised under this name, different morbid conditions, such as paralysis, apoplexy, and gangrene.

SIDERATIO, Apoplexy, Asphyxia, Phrenitis—s. Ossis, *Spina ventosa*.

SIDERATION, *Astrobolismus*, *Sideratio*.

SIDERION, (*σίδηριον*, 'an iron tool,') Ferramentum.

SIDERITES, (*σίδηρης*, from *sideros*.) Magnet.

SIDERITIS, Magnet, *Teucrum chamæpitys*.

SIDERO KRENÆ, (*sideros*, and *κρηνη*, 'a spring,') Water, mineral, (chalybeate.)

SIDEROPEGÆ, (*sideros*, and *πηγη*, 'a spring,') Water, mineral (chalybeate.)

SIDEROS, (*σίδηρος*.) Ferrum.

SIDEROXYLON, (*sideros*, and *ξύλον*, 'wood,') *Chrysophyllum cainito*.

SIDESADDLEPLANT, *Sarracenia*.

SIDHEE, see *Ganjah*.

SID'MOUTH, CLIMATE OF. Sidmouth is in Devonshire, England, on the seabeach, at the mouth of the Sid. It is much frequented by invalids, both during the summer and winter. The climate is, however, damp, and subject to sea-fogs.

SIÉGE, (originally from *sedes*, 'seat,') Anus, nates.

SIELISMUS, (*σιελισμος*.) Salivation.

SIELON, (*σιελον*.) Saliva.

SIEVE, ([Sax.] *ripe*, *ryfe*.) see *Cribration*—s. Drum, see *Cribration*.

SIFFLEMENT (F.), ([L.] *sibilare*, from *siffo*, 'to whistle,') Whistling.

SIFFLEMENT MODULÉ, *Chant des artères*, *Bruit musical*, 'modulated whistling, or music of the arteries.' *Musical sound*. A kind of sound heard on auscultation, resembling, at times, the

humming of certain insects. Various musical sounds, (F.) *Bruits musicaux*, are described from fancied resemblances:—as *Bruit de picage-mout*, resembling the clucking of the hen;—*Bruit de jappement*, resembling the barking or yelping of a young dog, *Bruit de rouet*, resembling the noise of a spinning wheel, &c.

The sound is chiefly met with in those labouring under chlorosis.

SIGAULTIAN SECTION, (after Dr. J. R. Sigault, a French physician,) Symphysectomy.

SIGE, (*ayn*), Taciturnity.

SIGH, ([Sax.] *gican*, 'to sigh,') *Suspirium*.

SIGHT, ([Sax.] *geþiht*,) *Vision*—s. *Askew*, *Dysopia lateralis*—s. *False*, *Metamorphopsia*, *Pseudoblepsia*—s. *Feebleness of*, *Amblyopia*.

SIGILLATION, *Sigillatio*, (from *sigillum*, 'a seal,') The mark of a cicatrix.

SIGILLUM SALAMONIS, (dim. of *signum*, 'a mark,' 'Solomon's seal,') *Convallaria polygonatum*—s. *Virginittas*, *Hymen*.

SIGMATOID, *Sigmoid*.

SIGMOID, *Sigmoidal*, *Sigmatoid*, *Sigmoides*, *Sigmoides*, *Sigmoides*; (*sigma*, and *eidōs*, 'resemblance,') That which has the form of the Greek Σ, or of C, an old form of the sigma.

SIGMOID CARTILAGES, *Semilunar cartilages*.

SIGMOID CAVITIES or FOSSE OF THE ULNA, (F.) *Cavités ou Fosses sigmoïdes*, are two notches at the upper part of that bone,—one for the reception of the humerus; the other for that of the radius.

SIGMOID FLEXURE OF THE COLON, *Flexura sigmoides* seu *iliaca*, *S. Romanum*, is a turn the colon takes immediately before its termination in the rectum. See *Colon*.

SIGMOID FOSSA (of the temporal bone), *Fossa sigmoidea*.

SIGMOID or SEMILUNAR VALVES, *Valvulæ Sigmoides* seu *semilunares*, *Processus Sigmoides*, (F.) *Valvulæ sigmoïdes* ou *semilunares*, are three valvular folds, at the commencement of the pulmonary artery and aorta. These valves have the form of a crescent, when applied against the parietes of the vessel. When the blood is sent from the ventricles by their contraction, the valves are applied against the sides of the vessel; but, during the diastole of the ventricle, they are depressed, and prevent the blood from returning to the heart. At the middle of the convexity of each valve is a small, hard, triangular granulum, called *Corpusculum* seu *Globulus Arantii*, which completely closes up the passage. It is, also, called *Corpusculum Morgagni* and *S. Semioideum*, (F.) *Globule d'Arantius*.

SIGN, *Signum*, *Signon*, (F.) *Signe*. Any present or past circumstance, afforded by the examination of a patient, or of matters concerning him, whence a conclusion may be drawn regarding the nature and seat of his disease. The phenomena which precede the disease are called *anæsthetic* or *commemorative signs*;—those which accompany it are termed *diagnostic*, if they reveal the nature or seat of the disease,—*prognostic*, when they indicate its probable duration and termination. See *Symptom*.

SIGNA ASSIDENTIA, see *Assidens*.

SIGNATURE, *Signatura*, *Signatio*, (*signum*, 'a sign,') This term was applied in the Eastern countries, in the middle ages, to mystic characters of good or bad augury, with which it was pretended that every one was marked by the star under which he was born.

SIGNATURES OF PLANTS meant certain peculiarities in their external formation or colour, which indicated that they were adapted for particular diseases;—as in the case of the *Lungwort*, *Liverwort*, &c.

SIGNE, *Nævus*—s. *Acant-coueurs*, *Precursory signs*—s. *Précurseurs*, *Precursory signs*.

SIGNUM, *Sign*, see *Parturition*—s. *Character isticum*, see *Pathognomonic*—s. *Conjunctum*, see *Pathognomonic*—s. *Diacriticum*, see *Diacritica signa*—s. *Morbi essentialia*, see *Pathognomonic*.

SILACH, *Silac*. An Arabic word, used to designate thickening of the eyelids, *Crassitie Palpebrarum*; and, also, *Alopexia*.

SILENCE, see *Murmur*, *Respiratory*.

SILENE CRASSIFOLIA, *Cucubalus behen*—s. *Inflata*, *Cucubalus behen*—s. *Pennsylvanica*, see *Silene Virginica*—s. *Thorei*, *Cucubalus behen*.

SILENE VIRGINICA, *Ground Pink*, *Wild Pink*, *Catch Fly*. *Ord. Caryophyllaceæ*. The root has been employed in decoction, as an efficacious anthelmintic.

Silene Pennsylvanica probably has the same properties.

SILER LANCIFOLIUM, *Laserpitium siler*—s. *Montanum*, *Laserpitium siler*.

SILITQUA, *Cera'tion*; the carat. A weight of about four grains. A pod.

SILIQUA ARABICA, *Tamarindus*—s. *Araci aromatici*, see *Vanilla*—s. *Banilla*, see *Vanilla*—s. *Dulcis*, *Ceratonium siliqua*—s. *Vaniglia*, see *Vanilla*—s. *Vanillia*, see *Vanilla*.

SILICUASTRUM PLINII, (from *siliqua*, 'a pod,') *Capsicum annuum*.

SILK, ([Sax.] *peole*,) *Sericum*.

SILK GRASS, *Yucca filamentosa*—s. *Weed*, *Asclepias Sullivantii*, *Asclepias tuberosa*—s. *Weed*, common, *Asclepias Syriaca*—s. *Weed*, flesh-coloured, *Asclepias incarnata*—s. *Weed*, swamp, *Asclepias incarnata*—s. *Worm*, see *Sericum*.

SILLOX, *Suleus*—s. *Horizontal de la veine ombilicale*, see *Liver*—s. *Transversale*, see *Liver*—s. *de la Veine cave inférieure*, see *Liver*—s. *de la Veine porte*, see *Liver*.

SILLONNEMENT, *Segmentation*.

SILLUS, (*σῖλος*,) *Camus*.

SILLO, (*σῖλος*,) *Camus*.

SILPHIUM, (*σῖλφιον*,) *Laserpitium*.

SILPHIUM LACINIATUM, *Rosinweed*, *Compass plant*, *Polar plant*;

SILPHIUM PERFORIATUM, *Cup Plant*, *Turpentine Sun Flower*, and

SILPHIUM TEREBINTHINACEUM, *Prairie Burdock*; *Nat. Order*, *Compositæ*; indigenous in Ohio and Michigan, and flowering from July to September; yield a fragrant gum which is esteemed stimulant and antispasmodic.

SILURUS GLANIS, (*σῖλουρος*,) see *Ichthyocolla*.

SILUS, (*σῖλος*,) *Camus*.

SILVER, ([Sax.] *peolfer*, [G.] *Silber*,) *Argentum*—s. *Ammoniochloride of*, see *Argentum*—s. *Chloride of*, see *Argentum*—s. *Chloruret of*, see *Argentum*—s. *Cyanide of*, see *Argentum*—s. *Cyanuret of*, see *Argentum*—s. *Iodide of*, see *Argentum*—s. *Ioduret of*, see *Argentum*.

SILVER LEAF, *Magnolia macrophylla*.

SILVER, *Muriate of*, see *Argentum*—s. *Nitrate of*, *Argenti nitras*—s. *Oxide of*, see *Argentum*.

SILVER AND AMMONIA, CHLORIDE OF, see *Argentum*—s. *and Ammonia*, *chloruret of*, see *Argentum*—s. *and Soda*, *hyposulphite of*, *Sodæ et argenti hyposulphitis*.

SILVESTER'S, (Dr. H. R.,) *METHOD FOR RESTORING SUSPENDED ANIMATION*. This consists in placing the patient on his back, pulling the tongue forward, and, then, drawing the arms slowly over the head, by which means the ribs are elevated by the pectoral muscles, and the chest is enlarged. The arms are now to be brought down to the side of the chest, which they must be made to compress slightly. These movements must be repeated so as to simulate respiration.

SILYBUM, *Carduus marianus*—s. *Maculatum*, *Carduus marianus*—s. *Marianum*, *Carduus marianus*.

SIMABA CEDRON. *Nat. Fam.* Simarubaceæ. A South American tree, the seeds of which—cedron seeds—have been used as an antidote for snake-bites. They have, also, been proposed in mental diseases and in epilepsy. The scrapings of the wood have been used as a substitute for sulphate of quinia.

SIMABA QUASSIOIDES, *Nima quassioides.*

SEMAROUBA, *Quassia simarouba*—s. Bitter, *Quassia simarouba*—s. *Faux*, *Malpighia mourella*.

SIMARUBA, *Quassia simarouba*—s. *Excoelsa*, see *Quassia*—s. *Officialis*, *Quassia simarouba*.

SIMILATEUR, *Malingerer.*

SIMILATIO, (*similis*, 'like,') *Assimilation.*

SIMILIA SIMILIBUS, 'like by like,' see *Homeopathy.*

SIMITIUM, *Plumbi subcarbonas.*

SIMO. (*σῆμος*,), *Camus.*

SIMONEA FOLLICULORUM, (after Simon, a German dermatologist), *Acarus folliculorum.*

SIMOTRACHELUS, (*σῆμος*, 'bent short,' and *τραχήλος*, 'neck,') *Bysauchen.*

SIMPLE, (opposite to compound.) A medicinal herb. One skilled in simples, is called a *Sim'plist* or *Simpler*. 'To *sim'ple*,' and 'to *cull simples*,' signify to gather medicinal herbs.

SIMPLER, see *Simple.*

SIMPLER'S JOY, *Verbena hastata.*

SIMPLIST, see *Simple.*

SIMULATED DISEASES, Feigned diseases.

SIMULATI MORBI, Feigned diseases.

SIMUS, (*σῆμος*,), *Camus.*

SINA LEVANTICA, *Artemisia santonica.*

SINAPELÆON, (*σινάπελαιον*), from *sinapis*, and *λαιον*, 'oil,' Oil impregnated with mustard.

SINAPI, (*σινάπι*), *Sinapis.*

SINAP'PIS, *S. n'gra*, *seu cer'nea seu Oliveria'na*, *Brassica n'gra*, *Na'pus*, *Eru'ca*, *Sina'pē*, *Sina'pi*, *Sine'pi*, *Napi'am*, *Na'py*, *Com'mon Black Mustard*, (F.) *Moltarde noire ou grise*, *Sénévé*, *Sanévé*. *Ord.* Cruciferae. *Sex. Syst.* Tetradynamia Siliquosa. The seeds are inodorous, when entire: but, when bruised, the odour is very pungent; taste bitterish and acrid. They yield their virtues to water, and afford a bland oil by expression. The seeds are stimulant, diuretic, emetic, and rubefacient. They are sometimes recommended to be swallowed whole, in dyspepsia, torpor of the bowels, &c. Externally, they are applied in the form of cataplasm. See *Sinapism.*

SINAPIS ALBA, *S. folio'ea*, *Bonan'nia officina'lis*, *Na'pus Leucosinapis*, *White Mustard*, also, is used medicinally. It is somewhat less pungent than the black. The distilled oil—*O'leum Sina'pis*—is a powerful stimulant and rubefacient. In the pharmacopœias of the United States and London, *Sinapis* is the seeds of *Sinapis nigra* and *Sinapis alba*.

White-head's Es'sence of Mustard consists of *al* of turpentine, camphor, and a portion of spirit of Rosemary: to which is added a small quantity of flour of mustard. It is a celebrated embrocation for rheumatism, &c.

White-head's Es'sence of Mustard Pills consist of yellow resin and balsam of tolu enough to make into pills.

Mustard Whey is made by taking bruised mustard seeds, 3ss, and milk, Oj. Boil together until the milk is curdled, and strain.

This whey is used in dropsy.

SINAPIS ARVEN'SIS, *S. Tinctoria'na*, *Char'lock*, *Wild Mustard*, *Gar'lock*, *Chad'lock*, *Rumsh*, (Sc.) *Runch'es*, *Scald'ricks*, *Skel'loch*, *Skel'lie*. The seeds are sometimes substituted for black mustard.

SINAPIS CER'NEA, *Sinapis*—s. *Eruca*, *Brassica eruca*—s. *Foliosa*, *S. alba*—s. *Nigra*, see *Sinapis*—s. *Oliveriana*, *Sinapis*—s. *Tamoriana*, *S. arvensis*—s. *Tuberosa*, *Brassica rapa*.

SINAPISM, *Sinapis'mus*, (from *sinapis*.) A

cataplasm, of which mustard forms the basis, which is used for exciting redness, and acting as a counterirritant. It is prepared by mixing flour of mustard and vinegar together for the due consistence. See *Cataplasma sinapis*.

SIN'CIPUT, gen. *Synceip'itis*, (from *semi*, 'half,' and *caput*, 'head,' or *sumum caput*, 'top of the head,') *Sincip'ital Re'gion*, *Syn'ciput*, *Hemicephal'eum*. The French use it synonymously with *Vertex*, *Mesocranium*, *Mesocranium*. By some, it is applied to the frontal region or forehead or forehalf of the cranium, *Hemicephalum*, *Hemicephalum*. The parietal bones have been called *sincip'ital bones*, *Ossa sincip'itis*, (F.) *Os du sinciput*.

SINDON, (*σινδών*, 'fine linen,') A small piece of rag, or round pledget, supported by a thread at the middle, and introduced into the opening in the cranium made by the trephine.

SINEPI, (*σινηπι*), *Sinapis*.

SINEW, ([Sax.] *ſinep*.) *Tendon.*

SINGLETON'S GOLDEN OINTMENT is formed of sulphuret of arsenic (realgar), and lard or spermaceti ointment.

SINGULT'OUS, *Lygmo'des*, *Lygo'des*, *Lyngo'des*, (F.) *Singultueux*, (from *singultus*.) Relating to or affected with hicough.

SINGULT'US, *Lynx*, *Lyg'mus*, *Spasmolgy'mus*, *Clo'nus seu Pneu'sis singult'us*, *Hic'cough*, *Hic'cup*, (old E.) *Hicket* or *Hickot*, *Hicquet*, (Prov.) *Heuks*, *Yez*, *Yux*, *Yoke*, (Sc.) *Es'kin*, *Ees'kin*, (F.) *Hoquet*. A noise made by the sudden and involuntary contraction of the diaphragm, and the simultaneous contraction of the glottis, which arrest the air in the trachea. It is a symptom of many morbid conditions, and especially of gangrene, but occurs, frequently, in persons otherwise in good health. Also, a sob.

SIN'ISTRAD, (*sinister*, 'to the left hand,') A term used by Dr. Barclay, adverbially, to signify 'towards the sinistral aspect.' See *Mesial*.

SINISTRAL ASPECT, see *Mesial*.

SINKING, FATAL, Shock, nervous.

SINUEUX, *Sinuoux.*

SINUSITAS, *Sinus.*

SIN'UOUS, *Sinu'ous*, (F.) *Sinueux*, (*sinuare*, 'to bow, to curve,') An epithet given to certain ulcers, and especially to certain fistulæ, which are tortuous.

SIN'US, gen. *Sinūs*, 'a bay or curve,' *Sinu'ositas*, *Abcon'sio*, *Eurycho'ria*, *Reces'sus*, (F.) *Golfe*. Any cavity, the interior of which is more expanded than the entrance. In this sense, anatomists have applied the term to cavities in certain bones of the head and face. It has been, also, applied to certain venous canals, into which a number of vessels empty themselves; such as the sinuses of the dura mater and of the spine; the sinus of the vena porta; uterine sinuses, &c.;—and to the bosom, and the vulva.

In *surgery*, it means a long, narrow, hollow track, leading from some abscess, diseased bone, &c., (F.) *Clapier*.

Sinuses of bones form part of the nasal cavities. They are the frontal, sphenoidal, maxillary, &c. The name has also been given, occasionally, to the ethmoid and mastoid cells.

Sinus Acusticus, Auditory canal, internal—s. *Ampullaceus*, *Cavitas elliptica*—s. *Aortici*, *Sinuses of Valsalva*—s. *Arteriosus*, *S. pulmonalis*—s. *Basilar*, *Sinus transversus*—s. *Cavernæ*, *Cavernous sinus*—s. *Cavernosi uteri*, *Sinus uterini*—s. *Cavernosus*, *Cavernous sinus*—s. *Cerebri*, *Ventricles of the brain*—s. *Choroidien*, *S. quartus*—s. *Circular*, of *Ridley*, *Sinus coronarius*—s. *Circularis*, *S. coronarius*—s. *Communis*, see *Semicircular canals*—s. *Coronaire du cœur*, see *Cardiac veins*.

SINUS CORONA'RIVS, *S. circula'ris seu circula'ris Ridlegi*, *Circular si'nus of Rid'ley*, *Corl*.

onary sinus. A venous canal, which surrounds, in a more or less regular manner, the pituitary fossa and gland, emptying itself into the corresponding cavernous sinus by each of its extremities.

SINUS, CORONARY, *Sinus coronarius*—*s. Coronary* of the heart, *Sinus coronary*, venous.

SINUS CORONARY, VE'NOS, *Coronary sinus of the heart*. A name given, on account of its great size, to that portion of the great coronary vein which embraces the left auriculoventricular furrow.

SINUS COXÆ, see *Cotyloid*.

SINUS DEX'TER COR'DIS, *S. vena'rum cava'rum seu venosus*, *Sac'cus vena'rum cava'rum*. The right auricle of the heart:—*Atrium cordis dextrum*. See *Auricula*.

SINUS DROIT, *S. quartus*.

SINUSES OF THE DU'RA MA'TER, *S. seu Fis'tulæ seu Cavita'tes seu Ventriculi seu Tubuli seu Sinquiductus seu Receptac'ula Du'ræ ma'tris*, are formed by a separation of the layers of the dura mater, which acts as a substitute for the outer membrane of veins. Bichat admits only four great sinuses, of which all the others seem to be dependencies;—the *two lateral sinuses*, the *straight sinus*, and the *longitudinal sinus*. To the lateral sinuses belong,—the *petrous sinuses*, superior and inferior; the *transverse sinus*; the *cavernous sinuses*; the *coronary sinus*, and the *occipital sinuses*. To the straight sinus belongs the *inferior longitudinal*. The longitudinal sinus, properly so called, has no sinus under its dependence.

SINUS FALCIFORMIS INFERIOR, *S. longitudinal*, inferior—*s. Falciformis superior*, *S. longitudinal*, superior—*s. Fourth*, *Torcular Herophili*—*s. Genæ pituitarius*, *Antrum of Highmore*—*s. of the Jugular vein*, see *Jugular veins*—*s. Lacrymalis*, *Lachrymal sac*—*s. Lactiferus*, see *Lactiferous vesels*.

SINUSES OF THE LARYNX, *Ventricles of the larynx*.

SINUSES, LAT'ERAL, *Sinus transversus*, *S. magnus*, *S. laterales*, (*F.*) *Sinus lateraux*. Each of these sinuses, distinguished into right and left, begins at the torcular Herophili or confluence of the sinuses, and terminates at the sinus of the jugular vein.

SINUS LATÉRAUX, *Sinuses, lateral*.

SINUSES, LONGITU'DINAL, *S. Longitudinales*. These are two in number. 1. **SUPERIOR LONGITU'DINAL SINUS**, *S. falciformis superior seu triangularis*, *Sinus median*, (*Ch.*) A long, triangular canal; convex above, concave below; which occupies the whole of the upper edge of the falx cerebri. It is broad behind, and much narrower before, where it commences by a *cul-de-sac*, at the crista galli of the ethmoid bone. It passes along the frontal or coronal crest, sagittal suture, and the vertical groove of the occipital bone. It receives several veins from the dura mater; and all those that are spread over the convex and plain surfaces of the two hemispheres of the brain. It discharges itself into the upper part of the torcular Herophili. 2. **INFERIOR LONGITU'DINAL SINUS**, *S. falciformis inferior*, *Vena fal'cis cer'ebr'i*. This is much narrower than the last, and occupies the inferior edge of the falx cerebri, from its anterior third to the tentorium cerebelli, where it terminates, commonly by two branches, in the straight sinus.

SINUS MAGNI, *Sinuses, lateral*—*s. Maxillary*, *Antrum of Highmore*—*s. Median*, *Sinus longitudinal superior*—*s. Median*, see *Semicircular canals*—*s. Muliebris*, *Vagina*.

SINUS, OCCIP'ITAL, *Sinus occipitalis*. These sinuses commence at the sides of the foramen magnum ossis occipitis, and ascend—becoming broader and nearer each other—into the sub-

stance of the falx cerebelli, where they frequently unite. Each opens into the inferior part of the torcular Herophili. They receive the veins of the falx cerebelli, of the dura mater which lines the fossæ cerebelli, and of the posterior part of the cerebellum. Haller and Sömmerring have given them the name of *posterior occipital sinuses*, to distinguish them from the *transverse sinus*, which they have called *anterior occipital sinus*.

SINUS OCCIPITALIS ANTERIOR, *Sinus transversus*—*s. Ophthalmic*, see *Cavernous sinus*—*s. Ovat*, see *Crista vestibuli*—*s. Perpendicularis*, *S. quartus*—*s. Pituitarii frontis*, *Frontal sinuses*—*s. Pocularis*, see *Ejaculatory ducts*—*s. Polymorphus*, *Cavernous sinus*—*s. Portarum*, see *Liver*—*s. Prostate*, see *Ejaculatory Ducts*—*s. Pudendi*, *Vagina*—*s. Pudoris*, *Vagina*, *Vulva*—*s. Pulmonales*, *Sinuses of Valsalva*.

SINUS PULMONALIS, *S. sinister cordis seu vena'rum pulmonalium seu arteriosus*, *Sac'cus vena'rum pulmonalium*, *Atrium cordis sinisterum*. The left auricle of the heart. See *Auricula*.

SINUS OF THE PULMONARY VEINS, *Auricle*, left. **SINUS QUARTUS**, *S. perpendicularis*, *Straight sinus*, (*F.*) *Sinus choroidien* (*Ch.*), *Sinus droit*. This sinus—triangular throughout; wide posteriorly; narrower anteriorly; slightly oblique downwards and backwards—passes along the base of the falx cerebri, from the termination of the inferior longitudinal sinus to the torcular Herophili. It receives the inferior longitudinal sinus, the veins of the lateral ventricles or *venæ Galeni*, the superior cerebellous veins, &c.

SINUS RENUM, *Pelvis of the kidney*—*s. Rhomboides*, see *Ventricles of the brain*—*s. Rotundus*, see *Crista vestibuli*—*s. of Santorini*, *Bulbus vestibuli*, and *Cavernous body of the vagina*—*s. of the Septum lucidum*, *Ventricle*, fifth—*s. Sinister cordis*, *S. pulmonalis*—*s. Sphenoidalis*, *Cavernous sinus*, see *Sphenoid bone*—*s. Straight*, *S. quartus*—*s. Superciliares*, *Frontal sinuses*—*s. Terminalis*, *Circulus venosus*—*s. Transversi*, *Sinuses, lateral*.

SINUS TRANSVER'SUS, *S. occipitalis anterior*, *Basilar sinus*. This sinus lies transversely at the anterior part of the basilar process of the occipital bone. It forms a communication between the two petrous sinuses, and the cavernous sinus of one side, and those of the other. Its breadth is commonly very great. It is situated between two layers of the dura mater, and presents, internally, a kind of cavernous tissue.

SINUS TRIANGULARIS, *S. longitudinal*, superior—*s. Urogenitalis*, *Urogenital sinus*.

SINUS UTERINI, *S. cavernosi Uteri*, *Uterine sinuses*. A name given to cavities, formed by the uterine veins in the substance of the uterine parietes. They are very large during pregnancy.

SINUS UTRICULOSUS, see *Semicircular canals*—*s. des Vaisseaux séminifères*, *Corpus Highmori*.

SINUSES OF VALSALVA, *Sinus Valsalvæ*, are three depressions in the aorta—*Sinūs aortici*,—and pulmonary artery—*Sinūs pulmonales*—immediately opposite the semilunar valves. Their use is to leave the edges of the valves free, so that they can be caught by the reflux blood. Where the aorta is about to send off the first of its large branches, at the top of the thorax, it is of great size, and has been called the *Large sinus of Valsalva*.

SINUS OF THE VENA CAVA, *Auricle*, right—*s. Venarum cavarum*, *S. dexter cordis*—*s. Venarum pulmonalium*, *S. pulmonalis*—*s. Venosus*, *S. dexter cordis*.

SINUSES, VERTEBRAL, (*F.*) *Sinus vertebraux*. Two great venous sinuses, different from those of the dura mater, which extend the whole length of the vertebral canal, from the occipital foramen to the end of the sacrum, behind the bodies of the vertebrae, before the dura mater, and at the sides of the posterior vertebral ligament. At their in-

ner side they communicate with each other by means of anastomoses, which form true *transverse sinuses*. They are situated at the middle of the body of each vertebra, passing under the posterior vertebral ligament. They receive, at their middle part, the veins that arise in the spongy tissue of the vertebrae. By their external side, the vertebral sinuses communicate with the posterior branches of the vertebral, intercostal, and lumbar veins. At their posterior side they receive the veins, which creep on the prolongation of the dura mater that envelops the marrow.

SINUS VULVÆ, Rima vulvæ.

SIPEERI, Bebeeru.

SIPEERINE, see Bebeeru.

SIPHAC, Peritoneum.

SIPHILIS, Syphilis.

SIPHITA, Somnambulism—s. Prava, Choreia.

SIPHON, Siphon—s. Auricularis, Otenthytes—

s. Urinæ, Diabetes.

SIPHON, *Sí'pho*, (σῖφων, 'a tube.') A bent tube, whose limbs are of unequal length, employed in pharmacy for transferring liquids from one vessel to another. With this view, the shorter limb is plunged into the liquid; and air is sucked from the extremity of the longer limb. The pressure of the atmosphere forces the fluid through the tube; and the flow will continue as long as the extremity of the longer limb is below the level of the fluid in which the shorter limb is immersed.

An exhausting syringe is, at times, attached to the longer extremity of the siphon, by means of which the air can be drawn from it.

SIPHON, PLUNG'ING. An instrument used in pharmacy for obtaining a specimen,—at times, a measured specimen,—of any liquid. With this view, the instrument is immersed into the liquid, and allowed to remain there until full. The opening at the top is then closed with the thumb; and the contents, thus removed, are allowed to flow out by removing the thumb and permitting the entrance of air. It is a form of the *pipette*.

SIPHONIA CAUCHU, (from σῖφων, 'a tube,' from the use made of it.) see Caoutchouc—s. Elastica, see Caoutchouc.

SIPHONO'MA, *Syphono'ma*, (F.) *Tumeur tubuleuse*, (from σῖφων, 'a tube.') A tumor of fibrous appearance and composed of parallel or intersecting tubes.—(Henle.)

SIPHUNCULUS, (dim. of σῖφων, 'a tube,') Syringe.

SIRA, Orpiment.

SIRENS, Symmeles.

SIRIASIS, (σῖρισις, a disease supposed to be produced by *Sirius*, 'the dog star' or by the sun.) *Coup de Soleil*, Insolation, Phrenitis—s. *Ægyptiaca*, *Coup de Soleil*.

SIRIUM MYRTIFOLIUM, Santalum album.

SIRIUS, (σῖριος,) Canicula.

SIROP, Syrop.

SIRUP, Syrup.

SISON AM'MI, (σῖσων,) *Æthusa am'mi*, *Onid'ium tenuifolium*, *Pimpinella laterifolia*, *Seseli fœniculifolium*. Ord. Umbelliferae. Sec. Syst. Pentandria Digynia. The plant that affords the *Am'mi ve'rum* of the shops; (F.) *Ammi des Boutiques*. The seeds—*Semen Am'mioa cret'ici* seu *ve'ri*—were once used as a carminative.

SISON AMOMUM, (F.) *Amome faux*, is considered to be excitant and diuretic. It is the *field honeywort*.

SISON ANISUM, *Pimpinella anisum*—s. *Podagraria*, *Ligusticum podagraria*.

SISTENS, (sistere, 'to still,') Episceticus.

SISYMBRIUM ALLIARIA, (σῖσμβριον,) *Alliaria*—s. *Barbarea*, *Erysimum barbarea*—s. *Erucastrum*, *Brassica eruca*—s. *Menthastrum*, *Mentha aquatica*.

SISYMBRIUM NASTURTIIUM, *Nasturtium aquat'icum* seu *officinale*, *La'ver odoratum*, *Cratæva-*

num, *Cressi*, *Cardamin' fontana* seu *nasturtium*, *Watercress*, (F.) *Cresson de fontaine*. Ord. Cruciferae. Sec. Syst. Tetradyamia Siliquosa. Watercresses are commonly used in salad. They have obtained a place in the *materia medica* for their antiscorbutic qualities.

SISYMBRIUM OFFICINALE, *Erysimum*—s. *Parviflorum*, S. *sophia*.

SISYMBRIUM SO'PHIA, *S. parviflorum*, *Herb Sophia*, *So'phia Chirurgorum*, *Flie or Flue-weed*, (F.) *Sagee des Chirurgiens*. It was once reputed to be anthelmintic, and valuable in hysterical affections. It is not used.

Other species of *Nasturtium*, as N. PALUSTRE, *Marsh watercress*; and N. AMPHIBIUM, *Water-radish*, possess similar virtues with *Nasturtium officinale*.

SISYRIN'CHIUM BERMUDIANUM, (σῖσσυριν'χον,) *Blue-eyed Lily*, *Physic* or *Scour'ey grass*; indigenous; Order, Iridaceae; flowering from June to August. Used as a cathartic by the Thomsonsians. A decoction of the root is acrid and cathartic.

SITAFISTE, see Furunculus.

SITHE, 'sigh,' *Suspirium*.

SITICULOSUS, (from *sitis*,) *Dipsodes*, see *Al* terative.

SITIENS, (*sitis*, 'to thirst,') *Dipsodes*.

SITIS, Thirst—s. *Intensa*, *Polydipsia*—s. *Morbosa*, *Polydipsia*.

SITKA, MIN'ERAL WATERS OF. See Archangel, New, mineral waters of.

SITOPHO'BIA, (σῖτος, 'food,' and φόβος, 'dread,') *Intense dread of food*.

SITOS, (σῖτος,) Aliment.

SITUS ALIENUS, 'a foreign position,' *Ectopia*—s. *Anomalus* seu *perversus* *intestinatorum*, *Splanchnectopia*—s. *Mutatus*, *Ectopia*, see *Monster*—s. *Perversus*, *Ectopia*.

SITZBATH, 'a seat bath,' see *Bath*, half.

SI'UM, (σῖον,) *S. nodiflorum* seu *angustifolium* seu *Berula* seu *erectum* seu *incisum*, *Apium si'um*, *Heliosciadium nodiflorum*, *Berula angustifolia*, *Berula*, *Creeping Waterparsnep*, (F.) *Berle nodiflore*. Ord. Umbelliferae. Sec. Syst. Pentandria Digynia. It is stimulant, and supposed to be diuretic and lithontriptic. It is, sometimes, employed in cutaneous eruptions.

SIUM ANGUSTIFOLIUM, *S. nodiflorum*—s. *Apium*, *Apium graveolens*—s. *Berula*, *S. nodiflorum*—s. *Bulbocastanum*, *Bunium bulbocastanum*—s. *Carvi*, *Carum*—s. *Erectum*, *S. nodiflorum*—s. *Graveolens*, *Apium graveolens*—s. *Incisum*, *S. nodiflorum*.

SIUM LATIFOLIUM is the common *Waterparsnep* of the United States. It is said to be poisonous.

SIUM LUNA'RE. The root is eaten by the Chepewyans. It has an agreeable nutty flavour. By the Canadians, it is called *Queue de Rat*. The poisonous roots of *Cicuta virosa*, *C. maculata*, and *C. bulbifera*, according to Sir J. Richardson, are often mistaken for it.

SIUM NIN'SI. The plant whose root is called *Ra'dix Nin'si*, *Nin'zin*, *Nind'sin*. This root was long supposed to be the same as ginseng, and it possesses similar, though weaker, properties. See *Panax schinseng*.

SIUM NODIFLORUM, *Sium*—s. *Silaus*, *Peucedanum silaus*.

SIUM SIS'ARUM, *Skir'ret*, (F.) *Chervi*, *Girole*, is used as a salad in Europe; and has been thought a useful diet in chest affections.

SIUM VULGARE, *Ligusticum podagraria*.

SIVVEN, (Gael.) *Rubus idæus*.

SIVVENS, *Sibbens*.

SIZY, (W. *Syth*, 'glue,') see *Corium phlogisticum* SKAW, Seal.

SKEELY, (from [G.] *scheelen*, 'to squint,') *Strabismus*.

SKELETEUMA, *Sceletuma*.

SKELETEUSIS, Mummification, Sceleteusis.

SKELETIA, (σκελετια,) Emaciation.

SKELETOGRAPHY, *Skeletograph'ia*, *Sceletograph'ia*, *Sceletograph'ia*, (skeletón, and γραφή, 'a description,') A description of the skeleton.

SKELETOL'OGY, *Skeletolog'ia*, *Sceletol'ogy*, *Sceletolog'ia*, (F.) *Squelettologie*, (skeletón, and λόγος, 'a discourse,') A treatise on the solid parts of the body. This part of anatomy comprehends *Osteology*, and *Syndesmology*.

SKELET'ON, *Scel'etum*, *Scel'etón*, *Cuda' ver exsiccata'*; (Old Eng. and Prov.) *At'omy*, *Anat'omy*, (Sc.) *Spwig*, (F.) *Squelette*, *Charpente osseuse*; (σκελετός, 'dried up,' from σκελλω, 'I dry,') A dried body. The aggregate of the hard parts of the body, or the bones. A skeleton is found in almost all animals, although it may not always be formed in the same manner. In some, as in the crustacea and testacea, it is wholly or partly external, *Ex'oskeleton*, *Derm'oskeleton*. In others, as in birds, the mammalia, &c., it is internal, *En'doskeleton*, *Nour'oskeleton*, (F.) *Endo-squelette*. It serves as a support for other organs. On it the general shape of the body and of its various parts is greatly dependent. When the bones are united by their natural ligaments, the skeleton is said to be natural, *Scel'etum natura'lē*; when articulated by means of wires, artificial or articulated, *Scel'etum artificia'lē*.

SKELETOPE'IA, *Sceletopœ'a*, *Sceleten'sis*, (skeletón, and ποίω, 'I make,') The part of practical anatomy which treats of the preparation of bones, and the construction of skeletons. Bones are prepared to exhibit their conformation, structure, chymical composition, mode of development, and changes; the different cavities formed by their union; and to demonstrate their connexions, relations, modes of union, &c.

SKELLIE, *Sinapis arvensis*, *Strabismus*.

SKELLOCH, *Sinapis arvensis*.

SKERLEJO, *Scherlievo*.

SKEVISCH, (from *Scabious*, [?]) *Erigeron Philadelphicum*.

SKEVISH, FLEABANE, *Erigeron Philadelphicum*.

SKESIGHT, 'oblique sight,' *Dysopia lateralis*.

SKIN, ([Sax.] *scin*), *Cutis*.

SKINBIND, Induration of the cellular tissue.

SKINBOUND DISEASE, Induration of the cellular tissue.

SKIN, BRONZED, see Capsule, renal—s. Scarf,

Epidermis.

SKINTIGHT, Induration of the cellular tissue.

SKIOMACHIA, *Sciomachy*.

SKIRRE, *Scirrhus*.

SKIRRET, *Sium sisarum*.

SKIRRIUS, *Scirrhus*.

SKITTER, (from [Sax.] *scetan*, 'to throw out,') *Diarrhœa*.

SKOKE, *Phytolacca decandra*.

SKOLEMORA FERNAMBUCENSIS, (σκωληξ, 'a worm,' and μωρω, 'to stupefy,') *Geoffrœa vermifuga*.

SKOLIOSIS, *Scoliosis*.

SKUESIGHT, 'oblique sight,' *Dysopia lateralis*.

SKULL, (Dan. *skal*), *Cranium*.

SKULLCAP, *Calva*, *Calvaria*, *Scutellaria galericulata*—s. Cap, maddog, *Scutellaria laterifolia*.

SKUNKCABBAGE, (from having a strong and offensive odour like that of the skunk,) *Dracontium foetidum*.

SKUNKWEED, *Dracontium foetidum*.

SLAB, *Torticiollis*.

SLABBERING, Slavering.

SLATE, IRISH, *Hibernicus lapis*

SLATERS, *Oniscæ aselli*.

SLATKAIA TRAVA, see Spirit.

SLAV'ERING, *Slab'bering*, *Slab'bering*, *Drin'elling*, (Old E.) *Pirt'ling*, (from *saliva*.) *Ptyalis'-mus in'ers*, (F.) *Beeve*. Involuntary flow of saliva, from sluggishness of deglutition, without increased secretion. It is seen in the infant, the aged, and the idiot.

SLEEP, (Sax. *slēp*), *Som'nus*, *Samp'nus*, *So'por*, *Dormit'io*, *Hyp'nos*, (F.) *Sommeil*. Temporary interruption of our relations with external objects. A repose of the organs of sense, intellectual faculties, and voluntary motion.

The act of quitting this condition is called *Awak'ing*, (F.) *Éveil*, *Réveil*. Sometimes this is progressive and natural; at others, sudden and startling; (F.) *Réveil en sursaut*,—as in nightmare, affections of the heart, hypochondriasis, &c.

SLEEP, MAGNETIC, *Hypno'sis biomagnet'ica*, is the sleep induced by the so-called animal magnetism. See Magnetism, animal.

SLEEP, NERVOUS, see Magnetism, animal.

SLEEP, WINTER, Hibernation.

SLEEPLESSNESS, *Insomnia*.

SLEEPWALKING, *Somnambulism*.

SLIACS, MIN'ERAL WA'TERS OF. *Sliacs* is a favourite bathing-place in Hungary, a few miles from Neusohl. Some of the waters are cold; others, thermal: the former are carbonated chalybeates; the latter are much used in the same cases as thermal springs in general.

SLIDING OF THE FLAP, see Flap.

SLIEVEFISH, *Sepia*.

SLING, ([G.] *Schlinge*), *Mittel'a*, *Mit'rium*, (F.) *Écharpe*. A bandage employed to keep the forearm bent on the arm, in cases of injury due to those parts, in fracture of the clavicle, injury of the hands, &c. A handkerchief is often sufficient for this purpose. It must be folded diagonally, and be tied, by the extremities, round the neck. The French surgeons have used many varieties of *Écharpes* or *Slings*; as the *Grande écharpe*, *Écharpe moyenne*, *Petite écharpe*, *Écharpe de J. L. Petit*, &c.

SLIPPERS, *Impatiens*.

SLIT AND TAIL BANDAGE, Invaginated bandage.

SLOBBERING, Slavering.

SLOE, ([Sax.] *sla*), *Viburnum prunifolium*.

SLOETREE, *Prunus spinosa*.

SLOKE, GREEN, *Ulva latissima*.

SLOPS, *Slop diet*. (Slop is, probably, the past participle of *slip*.) Liquid food, which is largely watery, and contains little that is nutritious.

SLOUGH, ([Sax.] *slōg*), *Eschar*.

SLOVEN WOOD, *Artemisia abrotanum*.

SLOWS, Milk-sickness.

SLUG, (D. *slak*), *Limax*—s. Sea. *Biche de Mer*.

SLUMBER, ([Sax.] *slumpean*, 'to slumber,') *Somnolency*.

SMALLAGE, *Apium graveolens*.

SMALLPOX, *Variola*—s. Modified, *Varioloid*.

SMALT, ([G.] *Schmalte*), *Protoc'ide of Cobalt*, *Smalt'm*, *Safre*, *Zaffre*, *A'zur*, *Blue Starch*. This oxyd has been given in rheumatism, in the dose of ten to twenty grains. A larger quantity excites vomiting.

SMARAG'DUS, (σμαραγδος,) (F.) *Émeraude*. The emerald. It was formerly regarded as a tonic.

SMARTWEED, *Polygonum hydropiper*, P. punctatum.

SMEARCASE, see Cheese.

SMECTICA, (σμηκτικός, 'cleansing,') Detergents.

SMECTIS, (σμηκτός,) *Cimolia terra*.

SMECTRIS, (σμηκτρίς,) *Cimolia terra*.

SMEGMA, gen. *Sme'gmat'is*, (σμηγμα, gen. σμηγματός, from σμηνω, 'to wipe, to cleanse,') Soap, grease, a salve, a liniment. Also, the secretion from the sebaceous follicles of the skin.

SMEGMA ARTICULARE, Synovia.—s. Cutaneum, see Sebaceous glands.

SMEGMA PRÆPUTII, *Se'būm præputiālē*. The fatty secretion from the sebaceous follicles of the glans penis.

SMEGMATORRHIŒA, (*smegma*, and *ῥέω*, 'to flow,') Stearrhœa.

SMEGMORRHIŒA, Stearrhœa.

SMELL, SENSE OF, Olfaction.

SMELLING, Olfaction.

SMELOOME'S OINTMENT FOR THE EYES, Ceratum resinæ.

SMILACINA RACEMOSA, (from *smilax*,) *False Spikenard*; indigenous; Ord. Liliaceæ; is said to be alterative, diuretic and diaphoretic.

SMILAX, gen. *Smilacis*, (σμιλαξ,) *Taxus baccata*—s. *Aspera Chinensis*, *Smilax China*—s. *Aspera Peruviana*, *Smilax sarsaparilla*—s. *Canellæfolia*, *S. pseudochina*.

SMILAX CHINA, *S. aspera Chinen'sis*. Ord. Smilacææ. Ser. Syst. Dioecia Hexandria. The root; *China*, *Ch. orientalis seu ponderosa seu ve'ra*, *Sankira*, *Gauquava*, *Chi'na root*, (F.) *Squine*, *Esquine*, *Racine de Chine*. Formerly much used in cutaneous and venereal affections.

SMILAX GLAUCA; indigenous; has similar virtues to those of *Sarsaparilla*.

SMILAX HEDEREFOLIA, *S. pseudochina*—s. *India Spinosa*, *Smilax pseudochina*—s. *Officinalis*, *Smilax sarsaparilla*.

SMILAX PSEUDOCINA, *S. Indica spinosa seu canellæfolia seu hederæfolia*, *Chi'na occidentalis seu spuriæ nodosa*, *American or West India China*. In scrofulous disorders, the root has been often preferred to the oriental kind. In other cases, it possesses similar virtues.

SMILAX SARSAPARILLA, *S. officinalis*, *Sarsaparilla*, *Sartaparilla*, *Smilax aspera Peruviana*, *Sar'za*, *Sar'sa*, *Caricilla'di*, *I'ra Pecan'ga*, *Macapatli*, *Zar'za*, *Zarzaparilla*, (F.) *Salsepareille*. The root of this and other species of *Smilax*, *Sarsaparilla*, (Ph. U. S.), is inodorous; taste bitterish, feculaceous, and fibrous; of a dark colour externally; white within. It is possessed of diuretic and demulcent properties, and has been often employed in the sequelæ of syphilis; when, after a mercurial course, nocturnal pains, ulcerations, &c., continue. It has, also, been advised in scrofula, chronic rheumatism, &c.; but its efficacy is, to say the least of it, a matter of doubt. Dose, ʒj to ʒj of the powder.

It is obtained from Honduras, Brazil, and Peru—the varieties being called respectively, *Honduras*, (F.) *Salsepareille de Honduras*; *Brazil*, (F.) *S. de Brésil ou de Portugal*, and *Peruvian Sarsaparilla*, (F.) *S. de Pérou ou caraque*.

SMILAX TAMNOIDES; indigenous; has similar properties.

SMILE, (σμιλη,) *A curved, two-edged bistoury. A knife.*

SMILION, (σμιλιον, dim. of *smile*,) Scalpel.

SMIT, (participle of *smite*,) Contagion.

SMITTLE, see Contagious.

SMYRNA, (σμύρνα,) Myrrha.

SMYRNISMUS, (from *σmyrna*,) Embalming.

SMYRNIUM OLUSATRUM, *Smyr'nium*, (from *σmyrna*; the odour of the seeds resembling that of myrrh:) *Hipposeli'num*, *Maceron'na*, *Macedon'sium*, *Hor'ba Alexandri'na*, *Gri'chem*, *Agrioceli'num*, *Alexan'ders*, *Aban'ders*. A plant formerly cultivated for culinary purposes. The seeds are bitter and aromatic, and the roots more powerfully bitter. They have been recommended as resolvents, diuretics, emmenagogues, &c.

SNAG. A tooth standing alone or projecting beyond the others.

SNAGGLETEETH, *Odontoparallaxis*.

SNAGREL, *Aristolochia serpentaria*.

SNAIL, ([Sax.] *nael*,) *Limax*—s. *Snail*, *Cochlea*.

SLAKEHEAD, *Chelone glabra*.

SNAKE, HOODED, *Cobra di Capello*—s. Spectacled, *Cobra di Capello*.

SNAKEROOT, *Garuleum bipinnatum*—s. Birthwort, *Aristolochia serpentaria*—s. Black, *Aetæa racemosa*, *Sanicula Marilandica*—s. Button, *Eryngium aquaticum*, *Liatriis picata*—s. Canada, *Asarum Canadense*—s. Dewitt, *Prenanthes*—s. Evergreen, *Polygala paucifolia*—s. Heart, *Asarum Canadense*—s. Virginiana, *Aristolochia serpentaria*—s. White, *Eupatorium ageratioides*.

SNAKEWEED, *Aristolochia serpentaria*, *Helenium autumnale*, *Polygonum bistorta*—s. American, *Cicuta maculata*.

SNAPS, Phaseolus.

SNEEZEWORT, *Achillea ptarmica*, *Helenium autumnale*.

SNEEZING, (Sax. *nieran*, [G.] *niesen*, 'to sneeze,' an example of onomatopœia,) *Clo'nus Sternutatio*, *Sternutatio*, *Sternutamentum*, *Sternumentum*, *Ptar'mus*, (Sc.) *Nees'ing*, (F.) *Êternuement*. A convulsive motion of the expiratory muscles, by which the air is driven rapidly, and rushes sonorously, through the nasal fossæ, carrying with it the mucus and foreign bodies which adhere to the pituitary membrane, and have occasioned the movement.

SNIRRELS, Nares.

SNOB, Nasal mucus.

SNO'RING, *Snore*, ([Sax.] *snora*,) *Rhen'chus*, *Ren'chus*, *Rhenx'is*, *Ron'chus*, *Rhon'chus*, *Ster'tor*, (Sc.) *Row'ing*, (F.) *Ronflement*. Noise made in the posterior part of the mouth and nasal fossæ during the movements of inspiration, particularly in sleep, both when the person is in health, and in disease, as in apoplexy. In certain affections of the chest—in some cases of pulmonary catarrh—a kind of snoring may be distinctly heard by applying the ear or the stethoscope to the chest.

SNOT, Nasal mucus.

SNOTH, Nasal mucus.

SNOUT, ([D.] *snuit*,) Nasus.

SNOW, (Sax. *snap*,) *Nix*, *Chi'on*, (F.) *Neige*. Water in a solid state, which falls from the atmosphere in white flakes. It is used externally as a tonic and discutient.

SNOWBALL, *Cephalanthus occidentalis*.

SNOWBERRY, *Symphoricarpos racemosus*.

SNOW BLINDNESS, *Niphableps'ia*. Blindness produced by the reflection of the solar light from snow. As a preventive, *Snow Eyes* are used by the Esquimaux. These are made of extremely light wood, with a bridge resting on the nose like spectacles, and a narrow slit for the passage of the light.

SNOWDROP TREE, *Chionanthus Virginica*.

SNOW EYES, see Snow blindness.

SNUFF, CEPHALIC, ([D.] *snuff*,) *Pulvis asari compositus*.

SNUFFLES, ([D.] *snuffelen*, 'to breathe hard through the nose,' probably an example of onomatopœia,) *Grave'lo neonat'rum*, *Rhinan'gia*, *Rhinanch'onē*. Breathing hard through the nose, often owing to accumulation of the mucous secretions,—an affection to which infants are liable, and which is relieved by oiling the interior of the nostrils.

SNURLE, *Coryza*.

SNURLES, Nares.

SOAP, ([Sax.] *rupe*,) *Sapo*—s. Acid, see *sapo*.

SOAP BERRIES, *Sapindus saponaria*.

SOAP, BLACK, see *Sapo*—s. Castile, *Sapo durus*.

SOAP OF THE COCOA-NUT OIL, see *Sapo*.

SOAP, COMMON OR SOFT, see *Sapo*—s. Liquid, see *Sapo*—s. Medicinal, see *Sapo*—s. Spanish, *Sapo durus*—s. Starkey's, see *Sapo*—s. Yellow, see *Sapo*.

SOAPWORT. *Saponaria*.

SOB. *Singultus*. ([Sax.] *peopan*, 'to grieve,' probably an example of onomatopoeia.) (F.) *Sanglot*. A spasmodic, sudden, and momentary contraction of the diaphragm, immediately followed by relaxation, by which the little air, that the contraction has caused to enter the chest, is driven out with noise. It is an evidence of corporeal or mental suffering.

SOBBY. Corpulent.

SOBOLES. 'offspring,' Epigone.

SOBRIETAS. Temperance.

SOCIA PAROTIDIS, 'companion of the parotid,' see Parotid.

SOCIETY, *Soci'etas*, (F.) *Société*, (*socius*, 'a companion.') An association for the promotion of some common object.

SOCIETIES, MEDICAL. Associations of medical gentlemen for the promotion of science have been numerous, and productive of valuable results. In the United States, societies exist in many of the States, possessing the power of examining into the qualifications of every one who may be desirous of practising within their particular jurisdiction. In some of the States, however, the qualified and unqualified are permitted to exercise their profession with equal privileges and immunities.

Amongst the medical associations that have issued from time to time useful transactions or publications are the following: IN AMERICA,—The Massachusetts Medical Society; the Physico-medical, and Medical Societies of the State of New York; the College of Physicians and the Pathological Society of Philadelphia, and the American Medical Association. IN EUROPE,—The Association of Fellows and Licentiates of the King's and Queen's College of Physicians of Ireland; the Medico-surgical and Phrenological Societies of Edinburgh; the College of Physicians, the Medico-surgical, the Medical, the Sydenham, New Sydenham, the Pathological, the Epidemiological, and Obstetrical Societies of London; the Associated Apothecaries and Surgeon Apothecaries of England and Wales, and the Provincial Medical and Surgical Association; the Royal Academy of Surgery, the Royal Society of Medicine, the Société Médicale d'Emulation, the Société de Médecine, the Royal Academy of Medicine, the Société Médicale d'Observation, and the Société de Biologie, of Paris; the Medico-surgical Academy of Vienna; the Medical Societies of Berlin and Copenhagen, &c., &c. IN ASIA,—The Medical and Physical Society of Calcutta.

Valuable medical essays are likewise contained in the published Transactions of the American Philosophical Society; the Royal Societies of London, Edinburgh, Göttingen, &c.; the Imperial Academy of Sciences of Paris; the Imperial Society of Sciences of St. Petersburg and Vienna; the Royal Academy of Sciences of Turin; the Academia Imperialis Leopoldina Naturæ Curiosorum, &c.

SOCORDIA, (*socors*, gen. *socordis*, from *se*, 'separation' and *cor*, gen. *cordis*, 'heart,') 'without intellect,' Hallucination.

SO'DA. (An Arabic word,) (F.) *Soude*, *Protoxide of sodium*, is obtained by adding caustic lime to a solution of carbonate of soda. The soda in solution may be obtained by evaporation as hydrate of soda. Its action is like that of potassa; but it is not official.

Soda impura, *Barilla*, *Bariglia*, *Barillor*, *So'da Hispanica*, *And'tron*, *Na'tron*, *And'ton*, *Na'tron Antiquorum*, *Kelp*, *Aphronitrum*, *Sal alkalinus fixus fossilis*, *Carbo'nes So'dæ impurus*, *Subcarbonas So'dæ impura*, *And'tron*; the mineral alkali is obtained from several sources:—naturally, in Egypt; artificially, from the incineration of marine plants; and from the decomposition of sulphate of soda, &c. It is principally,

however, derived from plants growing on the sea-coast:—as the *Sal'sola sativa*, *S. soda*, *S. tragus*, *Salicornia herbacea*, *Salicornia Arabica*, &c. The *Barilla ashes* are from the ashes of the *Salicornia Europæa*;—the *Kelp*, *Soude de Varec*, from the *Fucus vesiculosus* and other varieties;—the *Turkey Barilla*, *Roquette*, *Cendre du Levant*, *Rochetta Alexandrina*, from the *Mesembryanthemum Cyp'ticum*;—the *Alicant Barilla*, *Soda Alican'tina seu Alonen'sis*, from the *Mesembryanthemum nodiflorum*;—and the *Carthage'na Barilla*, from the *Salicornia* and *Sal'sola*.

Impure subcarbonate of soda, (F.) *Soude du commerce*, consists of subcarbonate of soda, potash, chloride of sodium, clay and other earthy substances. It is chiefly used for preparing the pure subcarbonate.

SODA, Cardialgia, Cephalalgia, Pyrosis.

SODA ACETA'TA, *Sodæ Acet'as*, *Ac'etate of Soda*, *Terra foliat'a minera'lis*, (F.) *Acétate de soude*. This salt is used in similar cases with the acetate of potass.

SODA AËRATA, *Sodæ carbonas*—*s. Alonensis*, *Soda*—*s. Bibo'rate of*, *Borax*—*s. Bicarbo'rate of*, *Sodæ carbonas*—*s. Bilate of*, *Picromel*—*s. Borate of*, *Borax*—*s. Boraxata*, *Borax*.

SODA, CAR'BO'NATE OF, *Sodæ Carbo'nas seu Bicarbo'nas*, *Carbo'nas Na'tricum*, *So'da Aërata*, *Carbonate or Bicarbonate of Soda*, *Carbonate of protoxide of So'dium*, (F.) *Carbonate de Soude*, *S. effervescente ou aëréé ou crayeuse*. (*Sodæ subcarb. lbj, Aquæ destill. Oijj.* Dissolve the subcarbonate of soda, and pass carbonic acid through the solution; set it aside to crystallize.) Its use is the same as the subcarbonate, than which it is less caustic, and better adapted for effervescing draughts.

SODA CAUS'TICA, *Protoxide of So'dium*, *Caustic Soda*, (F.) *Soude caustique*, *S. à la Chaux*, *Soude pure*. (The impure subcarbonate, treated by quicklime, and consequently deprived of its carbonic acid.) It may be used as a caustic like the *Potassa fusa*, and is even preferable to it, as it does not deliquesce and run.

SODA, CHLO'RIDE OF, *S. Chlorinat'a*, *So'dæ Chlo'ridum seu Chlo'metrum*, *Chlo'rinat'd Soda*, *Natrum Oxymuriat'icum*, *Oxymuriate of Soda*, (F.) *Chlorure de Soude*, *Chlorure d'oxide de Sodium*. This salt is used as a disinfecting agent; both to remove offensive smells in chambers, and to alter the condition of fetid ulcers. See *Liquor Sodæ Chlorinatæ*.

SODA, CHLORINATED, *Soda*, chloride of—*s. Hispanica*, see *Soda*—*s. Hydrochlorate of*, *Soda*, muriate of—*s. Hypophosphite of*, see *Calcis hypophosphis*.

SODA, HYPOSUL'PHITE OF, *Sodæ hyposul'phis*, (F.) *Hyposulfite de Soude*. This salt has been absurdly over extolled in chronic cutaneous, and scrofulous affections, and especially in tuberculosis of the lungs. Dose, gr. x and more. It is given in syrup.

SODA IMPURA, see *Soda*.

SODA, LITHATE OF, *Urate of Soda*—*s. Mephite of*, *Soda*, subcarbonate of.

SODA, MU'Riate OF, *S. muriat'a seu muriat'ica*, *Al'kali minera'le sali'num*, *So'dæ mu'rias*, *Sal communis seu culina'ris seu fixum seu gem'mas seu mari'nus seu al'bus seu fossilis seu li'gidum*, *Mu'ria*, *Na'tron muriat'um*, *Na'tron muriat'icum*, *Hydrochlo'rina Na'tricus*, *Chloruretum So'dii*, *So'dii Chlo'ridum*, *Com'mon salt*, *Sensalt*, *Rock salt*, *Bay-salt*, *White salt*, *Quadrichlo'ruet of So'dium*, *Hydrochlo'rate of protoxide of So'dium*, *Hydrochlo'rate of Soda*, *Chlo'ride of So'dium*, (F.) *Chlorure de Sodium*, *Sel marin*, *Sel de Cuisin*. In an impure state, this salt is one of the most abundant of natural productions. It is soluble in 3 parts of water, and is tonic, purgative, and anthelmintic. Externally, stimulant. It is a well-known and useful

culinary substance. 1. has been greatly recommended as an antiperiodic in the dose of $\frac{3}{4}$ ss or $\frac{3}{4}$ j.

SODA. NITRATE OF. *Soda Nitratum seu Acidum.* *Natronum seu Natrum nitricum.* *Nitron Nitricum.* *Alkali nitricum.* *Nitron Nitricum.* *Nitron Rhomboïdale seu cristallinum.* *Nitron Nitricum.* *Sulphate Nitrate of Soda.* *Protanitate of Sodium.* *Cubic Nitre.* Occurs native in Peru; but may be prepared by neutralizing carbonate of soda by nitric acid, evaporating and crystallizing.

Its virtues are similar to those of the nitrate of potassa, for which it has been substituted.

SODA. NITRITE OF. Made by heating a mixture of nitrate of soda and charcoal in a crucible to dull redness. When the salt has become white, raise the heat to melt it; pour it on a clean flag stone; break it into fragments, and preserve in a stoppered bottle.)

Used in the preparation of the Spiritus Ætheris Nitrosi of the British Pharmacopœia. (1864.)

SODA. OXYMURIATE OF. Soda, chloride of.

SODA. PHOSPHATE OF. *S. Phosphorata.* *Soda phosphos.* *Phosphas Natricum.* *Natronum Orthophosphos.* *Sol mineralis perlaticum.* *Alkali mineralis phosphoricum.* *Sulphate phosphate of protoxide of Sodium.* *Subphosphosphate of Sodium.* *Subphosphate of Soda.* *Protic phosphate of Soda.* *Tartaric purgating salt.* (F.) *Phosphate de Soude.* *Sol admissible pour l.* (Bone burnt to whiteness and powdered, 1b; *Sulphuric acid.* 1b; *Carbonate of soda.* q. s. Mix the bone with the acid in a covered vessel, add a gallon of water, and stir. Digest for three days, occasionally adding a little water to that lost by evaporation, and frequently stirring the mixture. Pour in now a gallon of boiling water, and strain through linen, gradually adding more boiling water, until the liquor passes nearly tasteless. Set the strained liquor by, that the dregs may subside: from which pour off the clear solution, and boil down to a gallon. To this solution, poured off from the dregs and heated in an iron vessel, add by degrees the carbonate of soda previously dissolved in hot water, until effervescence ceases, and the phosphoric acid is completely neutralized; then filter the liquor, and set it aside to crystallize. Having removed the crystals, add, if necessary, a small quantity of carbonate of soda to the liquor, so as to render it slightly alkaline; then alternately evaporate and crystallize so long as crystals are produced. Lastly, preserve the crystals in a well stopped bottle.—Ph. U. S.) Cathartic, in the dose of from $\frac{3}{4}$ ss to $\frac{3}{4}$ j.

SODA. PHOSPHATE OF, TRIBASIC. *S. Phosphate of.*

SODA POWDERS. see Powders, Soda.—s. Springs, see Water, Mineral (gaseous).

SODA. SUBCARBONATE OF. *Soda Sulcarbonas.* *S. Carbonas* (Ph. U. S.), *Natron purpuratum.* *Sol Soda.* *S. al'kali.* *Flos Salis.* *Sagittæ Vtri.* *Salt of Soda or of Borilla.* *Subcarbonate of protoxide of Sodium.* *Mophite of Soda.* (F.) *Subcarbonate de Soude.* The impure carbonate, dissolved in water: the solution filtered and crystallized. Soluble in two parts of water at 60°. It is much used as an antacid; and also as a 'deobstruent' in serofulous and other affections. Dose, gr. x to $\frac{5}{8}$ ss.

The dried Subcarbonate of Soda, *Soda Sulcarbonas exsiccata*, *Soda carbonas exsiccata*, *Carbonas Soda siccatum*, (F.) *Subcarbonate de Soude deséché*, is employed as an antacid and lithentriptic. Dose, gr. x to xv.

SODA. SUBPHOSPHATE OF. *S. phosphate of.*

SODA. SULPHATE OF. *Soda Sulphas.* *Natron vitriolatum.* *Sol catharticus seu mineralis Glauberi.* *Al'kali mineralis sulphuricum.* *Natron sulphuricum.* *Soda vitriolata.* *Sulphas natricus.* *Apo'ient Salt of Foodrich.* *Glauber's Salt.* (F.) *Sulfate de Soude.* Obtained from the salt that remains after the distillation of muriatic acid; the

superabundant acid being saturated with subcarbonate of soda.) It is soluble in three parts of water at 60°; is purgative, and, in small doses, diuretic. Dose, $\frac{5}{8}$ j to $\frac{5}{8}$ x.

The Lymington Glauber's Salt is a Sulphate of Magnesia and Soda, and is obtained from the mother liquor of sea-water, or by dissolving Epsom salt in a solution of Glauber's salt.

SODA. TARTRATE OF. *Tartrate of potassa and Soda.* *Tartarized Soda.* *Rochelle or Seignette's Salt.* *Soda Tartarizata.* *Sol Rapellen's seu Seignette's seu Polychoctum Seignette.* *Natron tartarizatum.* *Tartaris potassæ et Soda.* *Soda et Potassæ Tartras* (Ph. U. S. 1851.) *Potassa et Soda tartras*, (Ph. U. S. 1863.) *Soda tartarizans.* *Tartaris potassæ et Soda.* (F.) *Tartrate de potasse et de Soude.* (*Soda carb. bi.* *Potass. lithart.* in pulv. $\frac{3}{4}$ xvi, *Aq. bullient.* 1b.) Dissolve the carbonate in water; and add, gradually, the bitartrate. Filter the solution and crystallize.—Ph. U. S.) It is a gentle cathartic. Dose, $\frac{3}{4}$ ss to $\frac{3}{4}$ j.

SODA. URATE OF. *Urate of Soda.*—s. Valerianate of. *Soda valerianas.*—s. Vitriolata. S. sulphate of. **SODA WATER.** Acidulous water, simple.

SODA AND SILVER, HYPOSULPHITE OF. *Soda et Argenti Hyposulphitis.*

SODA ACETAS. *Soda,* acetate of—s. Arsenias, Arseniate of soda—s. Azotas, Soda, nitrate of—s. Bhoras, Borax—s. Bicarbonas, Soda, carbonate of—s. Boras, Borax—s. Boras alcalescens, Borax—s. Boras supersaturus, Borax—s. Carbonas, Soda, subcarbonate of—s. Carbonas exsiccatus, see Soda, subcarbonate of—s. Carbonas impurus, Soda—s. Chloras, Soda, chlorate of—s. Hyposulphitis, Hyposulphite of soda—s. Murias, Soda, muriate of—s. Nitras, Soda, nitrate of—s. Phosphas, Soda, phosphate of—s. Potassotartaras, Soda, tartrate of—s. Subboras, Borax—s. Subcarbonas impura, Soda.

SODA SULPHIS. *Sulphite of soda,* may be made by passing a stream of sulphurous acid gas into a strong solution of carbonate of soda, digesting the solution with sulphur at a gentle heat for several days, evaporating, and crystallizing.

SODA VALERIANAS. *Valerianate of soda,* is admitted into the Pharmacopœia of the United States for 1863. It is chiefly employed for obtaining the other valerianates.

SODA ET ARGENTI HYPOSULPHIS. *Hyposulphite of soda and silver.* (F.) *Hyposulphite de Soude et d'Argent.* (This salt is formed by pouring a solution of hyposulphite of soda over oxide of silver recently precipitated by potassa, until the oxide is entirely dissolved. Crystals are afforded by evaporation.) A solution of it is used in the same cases as one of the nitrate of silver.

SODA ET POTASSÆ TARTRAS. *Soda, tartrate of.*

SODII AUROTERTHLORIDUM. see Gold—s. Chloridum, Soda, muriate of—s. Chloruretum. Soda, muriate of.

SODII IODIDUM. *Iodide of sodium.* This salt has been prescribed in the same cases as the iodide of potassium.

SODIUM. *Natronum, Natronium,* (from *soda*.) The metallic basis of soda.

SODIUM. CARBONATE OF PROTOXIDE OF. *Soda carbonas*—s. Chloride of, Soda, muriate of—s. Chlorure de, Soda, muriate of—s. Chlorure d'oxide de, Soda, chloride of—s. Protinitrate of, Soda, nitrate of—s. Protoxide of, Soda—s. Protoxide of, hydrochlorate of, Soda, muriate of—s. Protoxide of, nitrate of, Soda, nitrate of—s. Protoxide of, phosphate of, Soda, phosphate of—s. Protoxide of, subcarbonate of, Soda, subcarbonate of—s. Quadrichloruret of, Soda, muriate of—s. Subborate of, Soda, muriate of—s. Subprotoborate of, Borax—s. Subprotophosphate of, Soda, phosphate of.

SODOMA GALLORUM, (*Sodom,* and *Galli,* 'the French') Syphilis.

SODOMIA, (from *Sodom*, 'the sin of Sodom,') Buggery.

SODOMY, Buggery.

SOFTENING, (from [Sax.] *roft*.) Mollities.

SOL-POUR-SOL, 'self for self,' Homœozygy.

SOIE, (originally from [L.] *seta*, 'a bristle,' [?]), Sericum.

SOIF, (from [L.] *sitis*.) Thirst—*s. Excessive*, Polydipsia.

SOL, 'the sun,' Gold.

SOLAIRE, Solar.

SOLLUNAR, *Solluna'ris*; (*sol*, 'the sun,' and *luna*, 'the moon,') An epithet applied to a fancied influence exerted by the sun and moon on man in a state of health and disease.

SOLAMEN INTESTINORUM, (from *solor*, 'to comfort,') *Pimpinella anisum*—*s. Scabiosorum*, Fumaria.

SOLANOID, *Solanoï'des*, (*solanum*, 'the potato,' and *oidos*, 'resemblance,') Resembling a potato; potato-like. An epithet applied to a form of cancer, which appears to be intermediate between scirrhus and encephaloid, increasing more rapidly than the former, yet approaching it in firmness.

SOLANUM, (from *solor*, 'to soothe,' [?]) *S. nigrum* seu *vulgarè* seu *vulgatum*, *Common* or *Gar'den Night'shade*, (F.) *Morelle à fruit noir*. Family, Solanaceæ. *Ser. Syst.* Pentandria Monogynia. A narcotic poison. The *Huile de Morelle* is made, by macerating in oil, for several days, the leaves and fruit of this plant. The oil is anodyne and discutient.

SOLANUM CAROLINENSE, *Horse nettle*; indigenous. The juice of the berries is said to have been employed successfully in idiopathic tetanus. [?]

SOLANUM DULCAMARA, *S. scan'dens* seu *ligu'osum*, *Dulcamara*, *D. flexu'osa*, *Glyciphy'eros*, *Amar'a dul'cis*, Σπυρος of Theophrastus, *Bit'tersweet*, *Woody Night'shade*, *Bit'tersweet Night'shade*, *Bit'tersweet Vine*, *Night'shade Vine*, *Violet bloom*, (F.) *Douce-amère*, *Morelle grimpeante*, *Vigne de Judée*. The taste of the stalks—*Dulcamara* (Ph. U. S.)—is bitter, followed by sweetness; hence its name. It has been used in many diseases, as a diuretic and sudorific, especially in chronic rheumatism, humoral asthma, dropsy, various skin diseases, scrofula, and jaundice; but is of doubtful efficacy. Dose, in powder, ℞j to ʒj.

SOLANUM ESCULENTUM, *S. tuberosum*—*s. Fœtidum*, *Datura stramonium*—*s. Furiosum*, *Atropa belladonna*—*s. Hortense nigrum*, *Atropa belladonna*—*s. Humboldtii*, *S. Lycopersicum*—*s. Lethale*, *Atropa belladonna*—*s. Lignosum*, *S. dulcamara*.

SOLANUM LYCOPERSICUM; *Lycopers'icum esculentum*, *L. po'mum amo'ris* seu *Humboldt'ii* seu *pseudolycopers'icum*, *Ma'tum lycopers'icum*, *Love-apple plant*. The fruit of this, called *Toma'to*, *Toma'to* or *Love-apple*, (F.) *Pomme d'Amour*, is much eaten in the United States; and, with the French, Spaniards, Portuguese, &c., forms an esteemed sauce.

SOLANUM MAGNUM VIRGINIANUM RUBRUM, *Phytolacca decandra*—*s. Maniacum*, *Atropa belladonna*, *Datura stramonium*—*s. Melanocerasus*, *Atropa belladonna*.

SOLANUM MELON'GENA, *Melon'gena*, *Egg-plant*, *Mad-Apple plant*; (F.) *Aubergine*. A native of Asia, Africa, and America. Its oblong, egg-shaped fruit, *Pomum Melon'genæ*, *Ma'tum insa'nium*, is prepared in soups and sauces, in those countries, like the tomato.

SOLANUM NIGRUM, *Solanum*.

SOLANUM NIVEUM. A South African plant, the leaves of which, applied to foul ulcers, cleanse them. The fresh juice of the berries and leaves, made into an ointment with lard or fat, is used by the farmers for the same purpose.

SOLANUM PARMENTERII, *S. tuberosum*—*s. Pseudolycopersicum*, *S. Lycopersicum*—*s. Quadrifolium*, Paris—*s. Racemosum Americanum*, *Phytolacca decandra*.

SOLANUM SANC'TUM; *Palestine Night'shade*. The fruit of this is globular, and is much eaten in the countries where it is a native.

SOLANUM SCANDENS, *S. Dulcamara*.

SOLANUM TUBEROSUM, *S. esculentum* seu *Parmenter'ii*, *Lycopers'icum tuberosum*, *Pa'pas Americ'anus*, *Pap'us Americ'anus*, *Americ'um Tuberosum*, *Batta'ta Virginia'na*, *Bata'tas*; the *Pota'to*, (F.) *Pomme de Terre*, *Parmentière*. A native of Peru. A nutritious and wholesome vegetable; one of the most useful, indeed, that exists. One hundred pounds of potatoes afford ten pounds of starch, which is equal to arrow-root, as a nutrient and demulcent. It is called *Pota'to Flour*, *P. Starch*, *French Sa'go*, *Petit Saque*, *Common Ar'row-root*, &c.

SOLANUM URENS, *Capsicum annuum*—*s. Vesicarium*, *Physalis*.

SOLANUM VIRGINIANUM; indigenous; is said to resemble *Solanum nigrum* in its properties.

SOLANUM VULGARE, *Solanum*.

SOLAR, *Sola'ris*, (F.) *Solaire*, (*sol*, gen. *solis*, 'the sun,') Relating to the sun; having rays like the sun.

SOLAR PLEX'US, *Plex'us sola'ris*, *Centrum communè*, *Cer'ebrium abdominalitè*. A collection of ganglions and filaments, interweaving and anastomosing with each other repeatedly, and belonging to the system of the great sympathetic. This vast plexus, lying on the vertebral column, the aorta, and the pillars of the diaphragm, has a very irregular shape. The ganglions and filaments composing it are soft, reddish, and almost all issue from the semilunar ganglions. It receives, also, branches from the pneumogastrics. It seems to be chiefly intended for the aorta, and accompanies all its branches; furnishing many secondary plexuses, which are called the *infra-diaphragmatic*, *cœliac*, *superior mesenteric*, *inferior mesenteric*, *renal* or *emulgent*, *spermatic*, &c.

SOLDANELLA, (from *solidare*, 'to make firm,') *Convolvulus soldanella*.

SOLDIER'S WEED, *Matico*.

SOLE, ([Sax.] *fol*.) *Solum*, *S. pe'dis*, *So'lea*, (from *solum*, 'the ground,') *Basis cor'poris*, *Ichnus*, *Superfici'es planta'ris pe'dis*, *Peti'na*, *Vestig'ium*, *Pe'dion*, *Pe'za*, *Pel'ma*, *Planta*, *Planta'ris superfici'es pe'dis*, *Faci'es concav'a seu inferior* seu *Pla'nium* seu *Concavum* seu *Par's inferior* seu *Plant'ies* seu *Plan'ta seu Vo'la pe'dis*, (F.) *Plante du Pied*. The sole of the foot; the under surface of the foot.

SOLEA, Sole—*s. Ipecacuanha*, see *Ipecacuanha*.

SOLÉAIRE, *Soleus*.

SOLEN, Canal, Cradle; see *Solenarium*, and *Vertebral column*.

SOLENA'RIMUM, (σωληνarium, from σωλην, 'a canal, a pipe,') An instrument of surgery, representing a kind of gutter, in the cavity of which the penis was placed, like a limb in a *cradle*. The word *Solen* itself—σωλην, *Cunilis*—was used, in antiquity, for a grooved channel or cradle, in which a fractured limb was placed.

SOLENOSTEMMA ARGEL, *Cynanchum olem-folium*.

SOLE'US, *Gastrocnemius inter'nus*, *Tibiocalcaneus* (Ch.), (F.) *Solaire*, (from *solen*, 'a sole of a shoe,' owing to its shape.) A muscle situate at the posterior part of the leg. It is broad, flat, and almost oval; and is attached, above, to the posterior edge of the fibula, to the oblique line on the posterior surface, and to the inner edge of the tibia. It terminates, below, by a large tendon, which unites to that of the gastrocnemius ex-

ternus, to form the ten to Achilles. This muscle extends the foot on the leg, and conversely.

SOLICATIO, (from *sol*, gen. *solis*, 'the sun,') Insolation.

SOLID, *Solidus*, *Stēreos*, (from *solum*, 'the ground,') A body whose integrant molecules are so united by the force of cohesion, that they will not separate by their own weight. The solids, in the human body, are the bones, cartilages, tendons, muscles, ligaments, arteries, veins, nerves, membranes, skin, &c. The anatomy, or rather study of the solids, is called *Stereology*, from *stēros*, 'solid,' and *logos*, 'a discourse.'

SOLIDA'GO, gen. *Solidag'inis*, (from *solidare*, 'to make firm,' in allusion to its properties as a vulnerary.) See *Solidago virgaurea*.

SOLIDAGO RIG'IDA, *Rig'id Goldērod*, *Bones' styptic*; indigenous; of the Composite family, flowering in August and September; is astringent and tonic.

SOLIDAGO SARACENICA, *S. virgaurea*.

SOLIDAGO VIRGAUREA, *S. vulgaris* seu *Saracēnica*, *Vir'ga au'rea*, *Hor'ba d'or'eā*, *Cmy'za co'ma au'rea*, *Sym'phytum Petra'm*, *El'ichrysum*, *Consol'ida Saracēnica*, *Gold'en rod*, (F.) *Verge d'or*, *Herbe des juis*. The leaves of *Solidago virgaurea*—**SOLIDA'GO** (Ph. U. S.)—*Fragrant* or *Sweet-scented Goldenrod*—and the flowers, have been recommended as aperients and tonics, in urinary diseases, and in internal hemorrhage.

SOLIDAGO VULGARIS, *S. virgaurea*.

SOL'IDISM, (from *solidus*, 'solid,') *Patholog'ia solida'ria*. The doctrine of those physicians who refer all diseases to alterations of the solid parts of the body. To the believers in this doctrine, the appellation *Solidists* has been given. The solidists think that the solids alone are endowed with vital properties; that they alone can receive the impression of morbid agents, and be the seat of pathological phenomena. Solidism has been the prevalent doctrine. It is scarcely necessary to say, that in all our investigations, the condition of both solids and fluids must be regarded.

SOLIUM, 'a bathing tub,' *Baignoire*; also, *Tænia solium*, (from *solus*, because considered to be found solitary or alone.)

SOLUBIL'ITY, *Solubil'itas*, (*solvere*, *solutum*, 'to dissolve,' and *habilis*, 'able,') That property of a body by virtue of which it can be dissolved in a men-struum; when it is said to be soluble.

SOLUBLE, *Solubilis*. Relaxing. Applied to the bowels when gently relaxed. See Solubility.

SOLUM, *Sole*—s. *Pedis*, *Metatarsus*, *Sole*.

SOLUTÉ, (from *solvere*, *solutum*, 'to dissolve,') Solution.

SOLUTIO, *Solution*—s. *Alexiteria Gaubiana*, see *Disinfection*—s. *Ammonia subcarbonatis*, *Liquor ammonia subcarbonatis*—s. *Arsenicalis*, *Liquor arsenicalis*—s. *Arsenicata*, *Liquor arsenicalis*—s. *Arsenitis kalica*, *Liquor arsenicalis*—s. *Calcis*, *Liquor calcis*—s. *Camphoræ æthereæ*, *Tinctura æthereæ camphorata*—s. *Camphoræ oleosa*, *Linimentum camphoræ*—s. *Continui*, *Solution of continuity*—s. *Cupri ammoniati*, *Liquor Cupri ammoniati*—s. *Ferri nitratis*, *Liquor Ferri nitratis*—s. *Guaiaei gummosa*, *Mistura guaiaei*—s. *Muriatis barytæ*, *Baryta*, *muriate of*—s. *Muriatis calcis*, see *Calcis muriatis*—s. *Muriatis ferriæ æthereæ*, *Tinctura seu alcohol sulfuricoæthereæ ferri*—s. *Muriatis hydrargyri oxygenati*, *Liquor hydrargyri oxy-muriatis*—s. *Potassii iodidi iodureta*, *Liquor iodinii compositus*—s. *Resinæ guaiaei aquosa*, *Mistura guaiaei*—s. *Solventis mineralis*, *Liquor Arsenici chloridi*—s. *Subcarbonatis ammoniaci spirituosus*, *Spiritus ammonia*.

SOLUTIO SULPHATIS CUPRI COMPOSITA, *A'qua citrili'ca ceru'lea*, *Syd'enhām's Styptic Wa'ter*, (*Sulph. cupri*, *Alum.* aa $\mathfrak{z}$ ij, *Acid. sulph.* aa $\mathfrak{z}$ iss, *Aquæ*, Oij: dissolve, and filter.) A styptic, and largely diluted, an astringent collyrium.

SOLUTION, *Solutio*, *Lysis*, (F.) *Dis'solution*. An operation which consists in dissolving a solid body in a menstruum. Likewise, the product of such operation—*Solutum*, (F.) *Soluté*. It means, also, with many, the termination of a disease:—with others, a termination accompanied by critical signs:—and, with others, again, it is synonymous with crisis.

SOLUTION OF ACETATE OF AMMONIA, *Liquor ammonia acetatis*—s. of *Ammonia*, beuzinated, see *Ammonia sulphas*—s. *Alkaline*, see *Fuligokali*—s. of *Alkaline iron*, *Liquor ferri alkalini*—s. of *Alum*, compound, *Liquor aluminis compositus*—s. of *subcarbonate of Ammonia*, *Liquor ammonia subcarbonatis*—s. of *Ammoniated copper*, *Liquor cupri ammoniati*—s. of *Arseniate of Soda*, see *Arseniate of soda*—s. *Arsenical*, *Liquor arsenicalis*—s. *Arsenical*, *De Valangin's*, *Liquor Arsenici chloridi*—s. of *Atropia*, *Liquor Atropiæ*—s. of *Carbonate of Soda*, *Liquor sodæ carbonatis*—s. of *Charcoal*, concentrated, see *Tinctura catechu*—s. of *Chloride of Arsenic*, *Liquor Arsenici chloridi*—s. of *Chloride of zinc*, see *Burnett's disinfecting liquid*—s. of *Chlorinated soda*, *Liquor sodæ chlorinata*—s. of *Chlorine*, see *Chlorine*—s. of *Citrate of Ammonia*, *Liquor ammonia citratis*—s. of *Citrate of Magnesia*, see *Magnesia citras*—s. of *Citrate of Potassa*, *Liquor Potassæ citratis*—s. of *Continuity*, see *Continuity*—s. of *Corrosive sublimate*, *Liquor hydrargyri oxy-muriatis*—s. *De Valangin's*, *Liquor arsenici chloridi*—s. *Donovan's*, see *Arsenic and Mercury*, *Iodide of*—s. *Fowler's*, *Liquor arsenicalis*—s. *Gannal's*, see *Alumina acetat*.

SOLUTION, **GOADBY'S**, called after the proposer, **Dr. Goadby**, is made of *baryt sol* $\mathfrak{z}$ iv, *corrosive sublimate* gr. iv, *alum* $\mathfrak{z}$ ij, and *boiling water* Oiv. It is employed to preserve animal substances. For most purposes it may be diluted with an equal bulk of water.

SOLUTION OF GUTTA PERCHA IN CHLOROFORM, see *Gutta percha*—s. *Heinecke's*, see *Arseniate of Soda*.

SOLUTION PAR LES HUILES, *Olea iac-dicinalia*.

SOLUTION OF HYDRIODATE OF POTASSA, *Liquor potassæ hydriodatis*—s. of *Iodide of Iron*, *Liquor ferri iodidi*—s. of *Iodine*, compound, *Liquor iodinii compositus*—s. of *Iron*, nitrated, *Liquor ferri nitratis*—s. *Labarraque's*, *Eau de Javelle*—s. of *Lime*, *Liquor calcis*—s. of *Lime*, saccharated, *Liquor calcis saccharatus*—s. *Lugol's*, *Liquor iodinii compositus*—s. of *Magnesia*, condensed, *Magnesia fluid*—s. *Monsel's*, see *Ferri sulphas*—s. of *Monro*, *Liquor of Monro*—s. of *Acetate of Morphia*, *Liquor morphiæ acetatis*—s. of *Hydrochlorate of Morphia*, *Liquor morphiæ muriatis*—s. of *Muriate of Morphia*, *Liquor morphiæ muriatis*—s. of *Sulphate of Morphia*, *Liquor morphiæ sulphatis*—s. of *Nitrate of iron*, *Liquor ferri nitratis*—s. of *Nitrate of mercury*, see *Hydrargyri nitratis*—s. of *Oxymuriate of mercury*, *Liquor hydrargyri oxy-muriatis*—s. *Pearson's*, see *Arseniate of soda*—s. of *Permanganate of Potassa*, *Liquor Potassæ Permanganatis*—s. of *Persulphate of iron*, see *Ferri sulphas*—s. *Plenck's*, see *Hydrargyri oxydum cinereum*—s. of *Potassa*, *Liquor potassæ*—s. *de Potasse*, *Liquor potassæ*—s. of *Sesquinitrate of Iron*, *Liquor ferri nitratis*—s. of *Soda*, *Liquor sodæ*—s. of *Strychnia*, see *Strychniæ muriatis*—s. of *Subacetate of lead*, *Liquor plumbi subacetatis*—s. of *Subacetate of lead*, diluted, *Liquor plumbi subacetatis dilutus*—s. of *Subcarbonate of potassa*, *Liquor potassæ subcarbonatis*—s. of *Subsulphate of iron*, see *Ferri sulphas*—s. of *Tartarized Antimony*, *Vinum antimonii tartarizati*—s. of *Terechloride of Antimony*, see *Antimonium muriatum*—s. of *Ter-nitrate of sesquioxide of iron*, *Liquor ferri nitratis*—s. of *Van Swieten*, *Liquor hydrargyri oxy-muriatis*.

SOLUTIVUS, *Laxative*.

SOLUTUM, Solution.

SOLVENS, (from *solvere*, 'to dissolve,') Solvent—s. Minerale, Liqueur arsenicalis.

SOLVENT, *Solvens*, (F.) *Fondant*. A medicine supposed to possess the property of dissolving or removing obstructions and extraneous substances. Thus we have solvents of the stone in the bladder, &c. See Dissolventia.

SOLVENT, MINERAL, Liqueur arsenicalis.

SOMA, (*soma*, gen. *somas*,) Body, Corpus.

SOMASCETICS, (*soma*, and *askein*, 'exercise,') Gymnastics.

SOMASCÉTIQUE, Gymnastics.

SOMATIC or SO'MATIC, (*somatikos*,) *Somat'icus*, (F.) *Somatique*, (from *soma*.) That which concerns the body;—as *somatic* death, or death of the body, in contradistinction to *molecular* death or the death of a part.

SOMATODYM'IA, (*soma*, and *δύω*, 'I enter into,') A double monstrosity in which the trunks are united. Of this, there are several varieties:—*Vertebrodym'ia*, where the union is by the vertebrae; *Ischiodym'ia*, by the ossa ischii; *Sternodym'ia*, by the sternum; *Supraömphalodym'ia*, (F.) *Sus-omphalodymie*, by the superumbilical regions; *Subpra et infra-omphalodym'ia*, by the superior and inferior umbilical regions; and *Sterno-omphalodym'ia*, where the union is by both the sternal and the umbilical regions.

SOMATOL'OGY, *Somatolog'ia*, (*soma*, and *λογος*, 'a discourse,') A treatise on the human body. Anatomy.

SOMATOTOMY, (*soma*, and *τεμνειν*, 'to cut,') Anatomy.

SOMMEIL, (from [L.] *sonnus*,) Sleep—s. *Hiberna*, Hibernation.

SOMMET, (from [L.] *summus*, 'highest,') Vertex.

SOMNAMBULATION, Somnambulism.

SOMNAMBULISM, *Somnambula'tio*, *Hypnobotate'sis*, *Hypnobotata'sis*, *Hypnobotat'ia*, *Hypnobotad'ies*, *Hypnobot'asis*, *Rhembas'mus*, *Lunatis'mus*, *Hypnonerig'ia*, *Selenogog'mia*, *Nyctobate'sis*, *Autonyctobat'ia*, *Selen'asis*, *Selenias'mus*, *Somnambulis'mus*, *Nyctob'asis*, *Noctambula'tio*, *Noctambulis'mus*, *Nyctobad'ia*, *Nyctobat'ia*, *Siph'ia*, *Syph'ia*, *Paroni'ia* *ambulans*, *Noctisur'gium*, *Oneirodyn'ia* *activa*, *Cenospu'dia*, *Som'novig'il*, *Somnambula'tion*, *Sleepwalking*, (*sonnus*, and *ambulare*, *ambulatium*, 'to walk,') A condition of the brain which occasions the individual to execute, during sleep, some of those actions that commonly take place when awake. It has been imagined that the somnambulist is guided solely by his subjective sensations; but there are many facts which induce the belief that the external senses are not always entirely suspended. The causes of somnambulism are very obscure; and the principal indication of treatment is,—so to arrange the bed-chamber of the somnambulist, that he cannot get out of it during sleep; and to avoid suppers, as well as all bodily and mental excitement.

SOMNAMBULISM, MAGNETIC, MESMERIC, or ARTIFICIAL, *Somnambulis'mus* *artificia'lis*, *Cataph'ora* *magnet'ica*, *Somnip'athy*, is that peculiar condition, into which individuals are thrown by the *Animal Magnetizer*. It occurs only in those of weak nervous temperament; in hysterical females, &c. See Magnetism, animal.

SOMNAMBULISMUS, Somnambulism—s. Artificialis, Somnambulism, magnetic.

SOMNAMBULIST, *Somnamb'ulus*, *Hybnob'ates*, *Hypnob'ates*, *Noctamb'ulus*, *Lunat'icus*, *Noctamb'ulo*, *Somnam'ulo*. A sleep-walker.

SOMNAMBULISTIC, *Somnam'bulosus*. Relating or appertaining to somnambulism,—as the '*somnambulis'tic* state.'

SOMNAMBULO, Somnambulist.

SOMNAMBULOUS, Somnambulist.

SOMNIA'TIO, (*sonnium*, 'a dream,') The act of dreaming.

SOMNIATIO MORBOSA, *Oncironosus*—s. *in* *Statu* *vigili*, Hallucination.

SOMNIFEROUS, *Som'nifer*, *Soporifer*, *Hypnot'icus*, *Hypnopa'us*, *Soporati'cus*, *Soporif'ice*, *Soporif'erosus*, (F.) *Hypnotique*, *Somnif'ere*, *Assoupissant*, *Soporatif*, *Soporif'ere*, *Soporifique*; an epithet for a medicine which causes sleep, (from *sonnus*, and *fero*, 'I carry,')

SOMNILOQUIST, *Somniloqu'us*, (F.) *Somniloque*, (from *sonnus*, 'sleep,' and *loquor*, 'I speak,') One who talks in his sleep.

SOMNILOQUIUM, *Somniloquis'mus*, *Somnil'ouism*. Talking in the sleep.

SOMNILOQUUS, Somniloquist.

SOMNIUM, (from *sonnus*,) *Eupn'ion*, *Insom'nium*, *Oneir'os*, *Oneir'on*, *Onar*, *Oneiropol'ia*, *a* *Dream*, (Sc.) *Sveuin*, *Sveeing*, *Sveeynyng*, *Sveenyng*, (F.) *Rêve*, *Songe*. A confused assemblage, or accidental and involuntary combination of ideas and images, which present themselves to the mind during sleep. Dr. S. Mitchell, of New York, used the word *Somnium* to signify the state between sleeping and waking, in which persons perform acts of which they are unconscious.

For 'waking dreams,' see Hallucination.

Those distressing dreams which occur during a state of half sleep, are called, by the French, *Rêveries*.

SOMNOLENCY, *Somnolen'tia*, *Hypnæthe'sia*, *Hypno'dia*, *Dormita'tio*, *Ca'rus* *Lethar'gus* *Cataph'ora*, *Cataph'ora*, *Co'ma*, *Somnolen'tium*, *Stum'ber*, *Heaviness*, (F.) *Assoupissement*; (from *sonnus*,) Sleepiness. Also a state intermediate between sleeping and waking. It is often a symptom of disease.

SOMNOPATHY, (*sonnus*, and *πάθος*, 'affection,') Somnambulism, magnetic.

SOMNOVIGIL, (*sonnus*, and *vigil*, 'awake,') Somnambulism.

SOMNUS, Sleep.

SOMP'NUS, Sleep.

SON, Furfur, (from *summus*, the last;—that which remains last in the sieve.) Also, Sound, (from *sonus*.)

SON HYDATIQUE, *Hydat'ic* sound, *Hydat'id* *frem'itus*. A sound, heard on percussion over tumours containing hydatids. The sensation experienced by the finger, is like that given by a watch struck on the surface opposite the glass.

SON'CHUS, (*σωνχος*,) *S. Olera'ceus*, *Hiera'cium* *oleraceum*, *Sonchist'le*, *Sonchist'le*, (F.) *Laiteron* *douz*. Ord. Cichoraceæ. Most of the species of *Sonchus* abound with a milky juice, which is very bitter, and said to possess diuretic virtues.

SONCHUS FLORIDANUS, *Mulgedium* *Floridanum*.

SONDE, Sound, Specillum—s. *Cannelée*, Director.

SONDE D'ANEL. A very fine stylet of silver, awl-shaped at one end. It is used for probing the lachrymal passages through the punctum lachrymale.

SONDE DE BELLOC. An instrument invented by Belloc for plugging the nasal fossæ, in cases of hemorrhage. It consists of a curved silver canula, open at both ends, and furnished with a button. The instrument is passed through the nares to the throat; and a plug is attached to the button, so that, when the instrument is drawn forwards through the nose, the posterior nostril is stopped.

SONDE BRISEE (F.), *Eyed probe*. A long, straight stylet, composed of two parts, which may be screwed to each other; blunt at one end, and furnished at the other with an eye, so that it may be used either as a probe for punctured wounds, or for a seton needle.

SONDE À CONDUCTEUR (F.). A modification of the ordinary catheter;—to facilitate the changing of catheters, without the fear of making false passages. It is a common catheter, open at its beak. The *Stylet* or *Mandrin* is one half

longer than that of the ordinary catheter. To use the instrument, the stylet is pushed into the cavity of the bladder through the catheter. The latter is then withdrawn by slipping it over the stylet, which is kept in its place, and serves as a guide to a fresh catheter.

SONDE À DARD, (*dard*, 'a dart,') see Lithotomy (high operation).

SONDE DE LAFOREST. A small, crooked canula, used for probing the nasal duct from below upwards, and for throwing injections into it.

SONDE ou PINCE DE HUNTER. A cylindrical silver canula; of the size of a common catheter; open at both extremities, and containing a wire of iron, one of the ends of which is split into two parts; each being terminated by a small silver scoop, and separating from each other by virtue of their elasticity, when pushed out of the canula, and again approximating when drawn in. It has been used for laying hold of small calculi in the urethra.

SONDE UTÉRINE, Sound, uterine—*s. de Vaquez*, Soda.

SONGE, Somnium.

SONITUS AURIUM, 'sound of the ears,' Tinnitus aurium—*s. Cardiacus*, see Heart, (sounds)—*s. Cordis*, see Heart, (sounds)—*s. Fluidi*, Hygrecinema—*s. Hepaticus*, Hepatechema—*s. Intestinalis*, Enterechema—*s. Stomachicus*, Gasterrechema.

SONOROUS RHONCHUS, *Rôle sonore*.

SONUS, Sound—*s. Vocis*, Accent.

SOOJA, see Dolichos.

SOOJIE, BASTER'S, a farinaceous article of diet, is said to consist of wheat flour sweetened with sugar.

SOOL, Gastroperiodynia.

SOOT, ([Sax.] *root*, *rot*.) Fuligo—*s. Tea*, see Fuligokali—*s. Wart*, Cancer, chimneysweepers.

SOPHIA, (*σοφία*, 'wisdom,' from its virtues,) Sisybrium sophia—*s. Chirurgorum*, Sisybrium sophia.

SOPHISTICATION, (from *σophistikos*, after the manner of the Sophists, 'captious, deceitful,') Falsification.

SOPHO'RA HEPTAPHYL'LA, *Anticholerica*. *Ord. Leguminosæ. Sex. Syst. Decandria Monogynia*. The root and seeds of this shrub are intensely bitter, and are said to have been found useful in cholera, colic, dysentery, &c.

SOPHORA TINCTORIA, *Baptis'ia seu Podalis'ia Tinctoria*, *Indigofera*, *Wild Indigo*, *Indigo Weed*, *Broom*, *Indigo Broom*, *Horsefly Weed*, *Yellow Broom*, *Clover Broom*, *Rattle Bush*, *Yellow Indigo*. An American plant, the taste of whose root is unpleasant, subacid, and nauseous—very similar to that of *ipeacacuanha*. In a large dose, the bark of the root acts both as an emetic and cathartic. It has been considered antiseptic and febrifuge, and has been used, in the form of fomentation or cataplasm, in phagedenic and gangrenous ulcers. A liniment, prepared by simmering the cortical part of the root in cream, has been found an efficacious application to sore nipples and to ulcerated breasts.

Baptis'ia Leucantha, *Tall white false Indigo*, an indigenous plant, has the same properties.

SOPHRONISTE' DENTES, (from *σωφρονιστος*, 'to act wisely,') see Dentition.

SOPHRONISTERES DENTES, see Dentition.

SOPHROSYNÉ, (*σωφροσύνη*) Temperance.

SOPIENS, (*sopio*, 'to put to sleep,') Paregoric.

SOPIO, Opium.

SO'POR, Sleep, and especially, a profound sleep, from which the person can be roused with difficulty. It is a symptom in many of the neurases.

SOPOR CAROTICUS, Carus.

SOPORALE ARTERIE, Carotids.

SOPORATIVUS, (from *sopor*.) Somniferous.

SOPORIFEROUS, (*sopor*, and *ferre*, 'to carry,') Somniferous.

SOPORIFIC, (*sopor*, and *facere*, 'to make,') Somniferous.

SO'POROUS, *So'porus*, (F.) *Soporeux*, (from *sopor*.) Sleepy: causing sleep.

SORA, Essera.

SORB TREE, *Sorbus domestica*.

SORBASTRELLA, *Pimpinella saxifraga*.

SORBEFA'CIENT, *Sorbefaciens*, (*sorbere*, 'to suck in,' and *facere*, 'to make,') A remedy that promotes absorption.

The following are the chief sorbefaciens:—*modifiers of nutritive action—entropics:—Acida Vegetabilia; Alkalia; Ammoniacum; Brominium; Galbanum; Hydrargyri Præparata; Iodinium; Liquor Calcii chloridi; Spongia usta; Compression, (methodical); Friction; Mental Sorbefaciens, (Imagination, Emotions.)*

SORBET, Scherbet.

SORBETHUM, Scherbet.

SORBETUM, Scherbet.

SORBETUM, Scherbet.

SORBIER DES OISELEURS, *Sorbus aucuparia*.

SORBIN, see *Sorbus domestica*.

SORBITIO, (*sorbere*, 'to drink,') Jus.

SORBIT'IO CARNIS seu EX CAR'NE. Broth or soup made of meat.

SORBITIUM, Jus.

SORBUM, see *Sorbus domestica*.

SORBUS, *Cratægus aria*, *S. domestica*—*s. Alpina*, *Cratægus aria*.

SORBUS AMERICA'NA, *Py'rus America'na*, *American Mountain Ash*. The bark resembles that of the Wild Cherry, and might be used in the same cases.

SORBUS ARIA, *Cratægus aria*.

SORBUS AUCUPARIA, *Mes'pilus*, *M. aucuparia*, *Mountain Ser'vice*, *Sorbus of the birdcatchers*, *Mountain Ash*, *Quick'en tree*, *Roan tree*, *Wit'chen*, *Witch hazel*, *Witchwood*, (Sc.) *Ro'den tree*, *Roun tree*, *Rouan tree*, *Rone*, (Prov.) *Keer*, *Rantree*, (F.) *Sorbier des Oiseleurs*, *Frêne sauvage*. *Ord. Rosaceæ*. The berries—(Sc.) *Rodens*—of this tree are employed for similar purposes as those of *Sorbus domestica*.

SORBUS OF THE BIRDCATCHERS, *S. Aucuparia*.

SORBUS CYDONIA, *Pyrus cydonia*.

SORBUS DOMESTICA, *Sor'bus*, *Py'rus domestica*, *Mes'pilus domestica*, *Oie*, *Ser'vice* or *Sorb Tree*, (F.) *Cormier*. *Family. Rosaceæ. Sex. Syst. Icosandria Trigynia*. The fruit of this tree, *Sorbum*, is astringent, and has been recommended in diarrhoea and dysentery. It is given in the form of rob. A kind of cider is made from it, and also brandy. A saccharine matter extracted from the fruit has been called *Sorbin*.

SORBUS MALUS, *Pyrus malus*.

SORDES, 'filth,' *Rhy'pus*. A dirty-looking sanies, discharged from ulcers. Also, accumulations of the secretions of the mouth, on the teeth in adynamic fevers more especially, and of foul matters in the stomach—*Sordes primæ'non vi'rum*, *Rhy'pus*. The French call an ulcer, exhibiting such an appearance, *sordide*. Also dirt of any kind.

SORDES, *Acatarsia*, *Ichor*, *Saburra*—*s. Acidæ*, *Acidities*—*s. Aurium*, *Cerumen*—*s. Primæ'non vi'rum*, *Embar'ras gastrique*, see *Sordes*—*s. Ventris*, *Excrement*.

SORDICULE AURIUM, (dim. of *sordes*, 'filth,') *Cerumen*.

SORE, ([Sax.] *rap*.) Ulcer.

SORE, *BAY*. A disease which Moseley considers to be a true cancer. It is endemic in the Bay of Honduras.

SORE MOUTH, *Stomatitis*—*s. m.* *Gangrenous*, *Cancer aquaticus*—*s. Swanalley*, see *Phagedenic*.

chaner—s. Throat, Cynanche—s. Throat, Clergyman's, Pharyngitis, follicular—s. Throat, inflammatory, Cynanche tonsillaris—s. Throat, putrid, Cynanche maligna—s. Throat, tubercular, Pharyngitis, follicular—s. Throat, ulcerous, Cynanche maligna.

SORENESS, *Parap'xis a'cris tenerit'u'do*. Painful uneasiness or tenderness, local or general, on being touched with a pressure that does not usually excite distressing feeling. Often, a febrile symptom. Also, the tenderness of a wound, ulcer, &c.

SORGHUM, *Panicum Italicum*—s. Saccharatum, see Saccharum.

SOROCO, *Puna*.

SOROCHE, *Puna*.

SORORIA'TIO, (from *sorores*, 'sisters,' *sororin*, 'to grow up or swell together,') The period at which the breasts of the female become developed. The act of becoming thus developed. A young maiden, whose mammae begin to show, —*sororian'tes mam'mæ*—was formerly called *sororians vir'go*.

SORREL, COMMON, ([F.] *Surelle*, from (Sax.) *rup*, 'sour,') *Rumex acetosus*—s. Field, *Rumex acetosella*—s. French, *Rumex scutatus*—s. Garden, *Rumex scutatus*—s. Mountain, *Oxalis acetosella*, *Oxyria reniformis*—s. Roman, *Rumex scutatus*—s. Salt of, Potass, oxalate of—s. Sheep, *Rumex acetosella*.

SORREL TREE, *Andromeda arborea*, *A. mariana*.

SORREL, WELSH, *Oxyria reniformis*—s. White, *Oxalis acetosella*—s. Wood, yellow, *Oxalis stricta*.

SORROCHE, *Puna*.

SOSTRUM, *So'trum*, (*σωστρον*, from *σῶζειν*, 'to save,' and *σωτρο*, 'a preserver,') A reward given to one who saves the life of another. A physician's fee or *honora'rium*, (F.) *Honoraire*.

SOTERIA DOCTRINA, (*σωτηριος*, 'preservative,') *Medicina*.

SOTERLÆ AQUÆ, Waters, mineral.

SOTERIOLOGY, *Soteriolog'ia*, (*σωτηριος*, 'preservative,' and *λογος*, 'a discourse,') Hygeology.

SOTIREL'LA. Ancient name of a medicine composed of opium and other narcotics, nutmeg, saffron, camphor, and saot. It was used in certain diseases of the teeth.

SOTRUM, (*σωτρον*,) *Sostrum*.

SOUBRELANGUE, *Ankyloglossia*.

SOUBRESAUT DES TENDONS, *Subsultus tendinum*.

SOUCHERLOON, *Bit noben*.

SOUCHE'T DES INDES, *Curcuma longa*—s. *Odorant*, *Cyperus longus*—s. *Rond*, *Cyperus rotundus*.

SOUCI (old F.), *Soulcie*, (from [L.] *solsequium*, like *heliotrope*, 'a follower of the sun,') *Calendula officinalis*, *Panophobia*—s. *des Alpes*, *Arnica montana*—s. *des Champs*, *Calendula arvensis*—s. *Ordnnaire*, *Calendula officinalis*.

SOUDE, *Soda*—s. *Acétate de*, *Soda*, acetate of—s. *Aéré*, *Sodæ carbonas*—s. *Borate de*, *Borax*—s. *Borate sursaturé de*, *Borax*—s. *Carbonate de*, *Sodæ carbonas*—s. *Caustique*, *Soda caustica*—s. *à la Chaux*, *Soda caustica*—s. *Chlorure de*, *Soda*, chloride of—s. *du Commerce*, *Soda*—s. *Crayeuse*, *Sodæ carbonas*—s. *Effervescente*, *Sodæ carbonas*—s. *Hyposulfite de*, *Soda*, *hyposulphite of*—s. *Phosphate de*, *Soda*, phosphate of—s. *et de Potasse*, *tartrate de*, *Soda*, tartrate of—s. *Pure*, *Soda caustica*—s. *Sousborate de*, *Borax*—s. *Sous-carbonate de*, *Soda*, subcarbonate of—s. *Sulfate de*, *Soda*, sulphate of.

SOUFFLE, (from *sufflare*, (*sub*, and *flare*,) 'to blow,') see Murmur, respiratory—s. *Amphorique*, see Cavernous respiration—s. *Bronchique*, see Murmur, respiratory—s. *Funic*, Bellows' sound, *funic*—s. *Métallique*, see Cavernous respiration—s. *Placentaire*, *Bruit placentaire*—s. *Trachéal*, see Murmur, respiratory—s. *Tubaire*, see Murmur,

respiratory—s. *Uterin*, *Bruit placentaire*—s. *Voilé* see Cavernous respiration.

SOUFFLE, WHISPERING. A bellows' sound heard over solidified lung and excavations when the patient whispers.

SOUFFRANCE, Suffering.

SOUFFRE, Sulphur—s. *Carburé*, *Carbonis sulphuretum*—s. *Iodure de*, *Sulphuris iodidum*—s. *Sublimé*, *Sulphur sublimatum*—s. *Végétal*, *Lycopodium*.

SOULLURE ([F.] *souiller*, *salir*, ([Sax.] *ryhan*, 'to soil,') Pollution.

SOULA'MEA AMA'RA, *Res amarorum*. A shrub of the Moluccas; *Order*, Polygalaceæ; all the parts of which have great bitterness—"horrenda amarities." It is employed by the Malays, and, it is said, with success, in cholera and pleurisy; and as a febrifuge.

SOUND, *Specill'um*, *Stylus*, *Explorato'rium*, *Rad'olus*, (F.) *Sonde*. An instrument used by surgeons to discover whether there be a stone in the bladder. It is usually made of highly polished steel, and is shaped like the catheter. The operation is termed *sounding*.

The French *Sonde* has, however, a more extensive signification. It means different instruments introduced into cavities of certain organs, or into wounds, fistulas, &c., to probe their condition, or to fulfil some therapeutical indication. See *Sonde*.

SOUND, *So'nus*, ([Sax.] *jon*,) *E'chos*, *Noise*, (F.) *Son*, *Bruit*. The sensation produced on the auditory nerve by the vibrations of a sonorous body. Sounds may be propagated in three modes.

1. By *reciproca'tion* or *con'sonance*, as when a sounding body, of a definite pitch, produces a musical tone when another body of the same pitch is sounded near it. 2. By *res'onance*, as when a sounding body is placed in connection with another, one or more of whose parts may be thrown into reciprocal vibration; and 3. By *conduc'tion*, as where the vibrations are transmitted through fluid, liquid, or solid media.

SOUND, ([L.] *sanus*, [Sax.] *junb*,) Healthy.

SOUND, Catheterize, Syncope—s. *Auricular*, *Apyromele*—s. Bellows', friction, rasp, saw, lancet, &c., see *Bruit*—s. Cracked pot, *Bruit de pot fêlé*—s. Crumpling, pulmonary, *Froissement pulmonaire*—s. Hydatid, *Son hydatique*—s. Laryngeal, *Laryngeche*—s. Musical, *Sifflement modulé*.

SOUND, U'tERINE, *U'terine bougie*, (F.) *Sonde utérine*. An instrument somewhat similar to a small male catheter, to be introduced into the cavity of the uterus, in the diagnosis of diseases of that viscus. When notched on the stem, it enables the size of the uterine cavity to be appreciated, and becomes a good *hysterom'eter*, (F.) *Hysteromètre*.

SOUNDING, Searching, see Sound—s. the Chest, see Auscultation.

SOUNDNESS OF MIND, Sanity.

SOUP, ([G.] *Suppe*,) see Broth.

SOUP'IR, (from [L.] *suspirm*,) Sigh.

SOUR BERRY, see *Oxyacantha* (Galen), *Vaccinium oxycoccos*—s. Croul, Sauer Kraut—s. Dock, boreal, *Oxyria reniformis*—s. Leaf, *Andromeda arborea*—s. Milk, Buttermilk—s. Sop, *Bromelia ananas*—s. Tree, *Andromeda arborea*—s. Wood, *Andromeda arborea*, *Andromeda mariana*.

SOURCIL, *Supercilium*.

SOURCILIER, *Corrugator supercilii*, Superciliary.

SOURD, (from [L.] *surdus*,) Deaf.

SOURDS-MUETS, 'deaf dumb,' see *Mutitas surdorum*.

SOURIS, *Nystagmus*.

SOURMILK, Buttermilk.

SOURSOP, *Bromelia ananas*.

SOUS (F.), in compound words, *Sub*.

SOUS-ACROMIO-CLAVI-HUMÉRAL, Deltoïd — *s. Atrodioid*, Subatloïdeus — *s. Acroïdien*, Subaxoïdeus — *s. Azote de Bismuth*, Bismuth, subnitrate of — *s. Clavier*, Subclavian — *s. Costaux*, see Intercostal muscles — *s. Cutané*, Subcutaneous — *s. Diaphragmatique*, Infradiaphragmatic — *s. Épineux*, Infraspinalis — *s. Lingual*, Sublingual — *s. Mammaire*, Submammary — *s. Maxillaire*, Submaxillary — *s. Maxillo-Labial*, Depressor anguli oris — *s. Mental*, Submental — *s. Métacarpolatéri-phalangien*, Prior annularis — *s. Occipital*, Suboccipital — *s. Optico-sphéno-scléroticien*, Rectus inferior oculi — *s. Orbitaire*, Suborbital — *s. Pubien*, Infrapubian — *s. Pubio-coecygien*, Levator ani — *s. Pubio-crété-tibial*, Gracilis — *s. Pubio-prétibial*, Gracilis — *s. Pubio-trochantérien-externe*, Obturator externus — *s. Pubio-trochantérien-interne*, Obturator internus — *s. Scapulaire*, Subscapularis — *s. Scapulo-trochantérien*, Subscapularis — *s. Sels*, see Salt.

SOUTH CAROLINA, MIN'ERAL WATERS OF. *Glenn's Springs*, in Spartanburg District, are highly charged with salts of lime, and, not far from them, is a chalybeate. *West's Spring*, and *Chick's Springs*, a few miles from Greenville, resemble Glenn's Springs.

SOUTHERNWOOD, *Artemisia abrotanum* — *s. Field*, *Artemisia campestris* — *s. Maritime*, *Artemisia maritima* — *s. Tartarian*, *Artemisia santonica*.

SOWBREAD, *Arthanita*, *Cyclamen*.

SOWENS, ([Sax.] *seape*, 'paste,') Flummery.

SOY, see *Dolichos soja*.

SOYMIDA FEBRIFUGA, *Swietenia febrifuga*.

SPA, MIN'ERAL WATERS OF. A town seven leagues southeast of Liège, where there are several springs, possessing a sharp acidulous taste; containing much carbonic acid, carbonates of iron, lime, and magnesia; carbonate of soda, and chloride of sodium. The water is much used as a tonic. Season, from the 1st of June to the 15th of October.

SPA WATER, ARTIFICIAL. (*Sodæ subcarb.*, gr. viij; *Magnesiæ carbon.*, ℥j; *Linat. ferri*, gr. iij; *Sodii chlorid.*, gr. j; *Aquæ*, Oijj. Impregnate with gas from *Marble powder* and *Sulph. acid.*, aā ℥x.)

SPACE, INTERPEDUNCULAR, see *Interpeduncular* — *s. Perforated*, anterior, *Locus perforatus anticus* — *s. Perforated*, posterior, see *Tarini pons*.

SPADEBONE, *Scapula*.

SPADO, (σπαδων,) *Castratus*, *Spasm*.

SPADON, (σπαδων,) *Castratus*, *Spasm*.

SPAGYRIA, *Chymistry*.

SPAGYRISTS, (σπαω, 'I separate,' and αγειω, 'I assemble,' because they reduced compounds into elements, and formed the latter into compounds [2].) A sect of physicians, who pretended to account for the changes that occur in the human body in health and disease, in the same manner as the chymists of their day explained those of the inorganic kingdom. *Medicina Paracelsistica* seu *Spagyrica*, was likewise called *Hermetica*, because it was believed that the means of cure adopted in it had been found in the books of Hermes.

SPAIG, *Skeleton*.

SPANÆMIA, (σπανος, 'poor,' and αἷμα, 'blood,') Poverty of the blood. Diminution in the quantity of red corpuscles of the blood, — as where bleeding has been carried beyond certain limits.

SPANÆMIC, *Spanæmicum* (remedium), *Dysplas'ticum*, *Plastily'ticum* et *erethily'ticum*, *Hæmatoly'ticum*, *Dyscrasy'ticum*. Relating to *spanæmia*.

A medicine whose protracted use is said to impoverish the blood, — as iodine, bromine, &c.

SPANDARAPUM, *Sparadrapum*.

SPANING, ([G.] *spān* n. 'to wean,') Weaning — *s. Brash*, *Brash*, weaning.

SPANISH FLY, *Cantharis* — *s. Juice*, *Extractum glycyrrhizæ*.

SPANNADRAPUM, *Sparadrapum*.

SPANAPOGON, (σπανος, 'rare,' and πωγων, 'beard,') One who has lost his beard, or has a thin beard.

The Greeks called those who had little beard, or who had lost it, *Spanopogones*.

SPARADRAPA, *Sparadrapum*.

SPARADRAPIER. The name of a machine for spreading sparadrap. A *Plaster machine*. It consists of a table, with two raised pieces, movable, and furnished with points, by which the cloth can be stretched, and of a lamina or blade of metal to extend the plaster over it.

SPARADRAPUM, *Sparadrapa*, *Sparadrapus*, *Sparadrap*, *Tela emplastrica* seu *emplastrica*, *Spandara'pum*, *Spaniadrappum*, (F.) *Sparadrap*. Any adhesive plaster spread upon linen or paper, (F.) *Papier médicamenté*. The chief sparadrapes are the following: —

SPARADRAPUM ADHÆSIVUM, *Adhesive Plaster*. A spread plaster of the *Emplastum adhæsivum*. It is also called *Strapp'ing*.

English Courtplaster is a *Sparadrap*, (F.) *Taffetas agglutinatif*, *T. gommé*, *Taffetas d'Angleterre*, *Ser'icum Anglieum*, *Emplastum Anglieum*, *E. Anglica'num*, *E. Adhæsivum Woodstock'i*, *E. Icthyocol'tæ tel'æ inductum*, *E. Glutin'osum*, *Tel'la Ichthyocol'tæ glutinosa*, *Isinglass Plas'ter*. (It is made by stretching black silk, and brushing it over with a solution of *isinglass* ℥i, in *spirit* ℥xij, to which *tinc. benzoin* ℥ij, are added. When dry, this is repeated five times; after which two coats are given it of a solution of *tereb. Chie* ℥iv, in *tinc. benz.* ℥vj, which render it less liable to crack.)

Lis'ton's Isinglass Plas'ter is made by spreading several coats of strong solution of isinglass in weak spirit over oiled silk, or, still better, over animal membrane previously prepared for the purpose from the peritoneal coat of the cæcum of the ox.

SPARADRAPUM PRO FONTICULIS, *Is'ne Plas'ter*, is sometimes made of *simple diachylon*, ℥ij; *Burgundy pitch* and *sarcocolla*, each ℥iv; *common turpentine*, ℥j. Spread upon linen and polished with a moistened calendering glass rubber.

SPARADRAPUM seu TEL'la GALTERI, *Defensive Plas'ter*, (F.) *Toile de Gantier*. (This is made of *Olive oil*, ℥ss; *Suet*, ℥iv; *Wax*, ℥x; *Litharge*, *Common turpentine*, *Thus*, and *Mastic*, aā ℥ij; *Bole armeniac*, *Flour*, aā ℥j. Pour it, while liquid, upon cloth, and spread.) Used for issues and to keep on dressings.

SPARADRAPUM VESICATORIUM. Several substances have been introduced as substitutes for blistering plaster, (see *Emplastum Lyttæ*), under the names, *Tel'la vessicatoria*, *Blister'ing Tis'sue*, &c., and other forms of which are the *Taffetas résicant* ou *épispastique*, *Papier épispastique*, ou *résicant*, *Charta vesicatoria*. They are made of an *etheral* or *alcoholic* extract of *cantharides*, or of *cantharidin*, mixed with *wax*, and spread, in a very fine layer, on silk or paper previously oiled or waxed. They are efficient agents.

SPARADRAPUM VIRIDÈ, *Cornplaster*. (This may be made of *Yellow wax*, ℥ij; *Burgundy pitch*, ℥xij; *Common turpentine*, ℥vj; *Verdigris*, ℥ij; spread on cloth and polished.)

Kennedy's Cornplaster is made of *Yellow wax*, ℥ij; *Venice turpentine*, ℥ij; *Verdigris*, ℥j.

SPARADRAPUS, *Sparadrapum*.

SPARAGMA, gen. *Sparagmatis*, (σπαρσσω, 'to tear,') *Laceration*.

SPARAGMUS, (σπαρσμος, 'tearing,') *Spasm*.

SPARAGRASS, *Asparagus*.

SPARAGUS, *Asparagus*.

SPARALLIUM, *Clyster uterinus*. An injection into the vagina.

SPAREDIA. A bandage covered with the white of egg.

SPARGANIUM RAMOSUM, (*σπαργάνιον*.) *Great Burweed*. (from *sparganon*, owing to the hand-like leaves.) Indigenous; Order, Typhaceæ; flowering in July and August. The roots are substringent, but esculent; yielding a fine fecula, similar to salep. They are sometimes made into a poultice for inflamed mammae.

SPARGANON, *Spar'ganum*, (*σπαργάνον*, from *σπαρνω*, 'I swathe,' 'I wrap.') 'Swathing clothes'; a kind of bandage, with which children were formerly surrounded. Also, a fascia or band.

SPARGANOSIS, *Spar'go'sis*, (*σπαργάνωσις*, from *σπαργνω*, 'I am full to bursting'.) Extreme distension, as of the breasts by milk; *Intumescencia lactea mammae*, *Mastodyn'ia polyg'ala*. Sparganosis also means the wrapping of a child in swathing clothes.

SPARGANOSIS PUERPERARUM, *Phlegmatia dolens*.

SPARGANUM, *Sparganon*.

SPARGOSIS, (*σπαργώσις*.) *Sparganosis*.

SPARROWGRASS, *Asparagus*.

SPARSUS, (*σπargere*, *sparsum*, 'to throw here and there,') *Sporadic*.

SPARTEIN, see *Spartium scoparium*.

SPARTIUM JUNCEUM, (*σπαρτίον*, 'a rope,'—the ancients used by their branches or bark to form a rope.) *Span'ish broom*. Ord. Leguminosæ. A small European shrub, cultivated in the gardens of the United States, on account of its yellow flowers. The seeds are diuretic and tonic, in small doses; emetic and cathartic in large. They have been used in dropsy—10 or 15 grains three times a day.

SPARTIUM SCOPARIUM, *Sarathamnus scoparius*, *Cyt'isus scoparius*, *Genis'ta*, *G. scoparia* seu *hirsuta*, *Cytisogenis'ta*, *Broom*, (F.) *Géné à balai*. *Broom Tops*, *Spar'tii Cacumina*, *Scoparius*, (Ph. U. S.) have a bitter taste, and are possessed of diuretic properties. They have been used in dropsies. Dose, of the powder, ℥j to ʒj. All the genistæ have similar virtues. A decoction of genista, along with the cauterization of the pustules under the tongue, has been recommended in hydrophobia. An organic liquid base—*Spar'tein*—has been separated from it, which has narcotic properties; and a yellow crystallized substance, *Scoparin*, which is diuretic.

SPARTIUM TINCTORIUM, *Genista tinctoria*.

SPASIS, (*σπασίς*.) *Spasm*.

SPASM, *Spas'mus*, *Spas'ma*, *Spas'is*, *Sparag'mus*, *Synol'cē*, *Spu'do*, *Spadon*, (F.) *Spasme*, (*σπασμος*, from *σπaw*, 'I draw'.) The Greeks gave this name to all kinds of convulsions. It is now usually applied to involuntary muscular contractions; and these, again, have been divided into *tonic* spasm, *Paraton'ia*, which consists in permanent rigidity and immobility of the muscles that are the seat of it (see *Tetanus*); and *clonic* spasm, which consists in alternate contractions and relaxations (see *Convulsion*).

Cullen has a class, *Spas'mi*—the *neurospas'mi* of Fuchs.

SPASM, CLONIC, *Convulsion*—s. *Cynic*, see *Canine laugh*.

SPASM OF THE GLOTTIS, *Asthma thymicum*—s. *Histrionic*, of the Face, see *Tic*—s. of the Larynx, *Asthma thymicum*—s. *Masticatory*, of the Face, *Trismus*—s. with Rigidity, *Tetanus*.

SPASM, WRITERS'. Involuntary spasms of the muscles of the thumb, or thumb and index finger, observed, at times, in those who are much accustomed to writing. See *Cramp*, writers'.

SPASMA, *Spasm*.

SPASMATIC, see *Spasmodicus*. Dr. Marshall Hall used the word *spasmat'ic* for one, in a convulsive disease, in the state of excitement. *Spasmatode* he employs for one unexcited.

SPASMATICUS, (from *spasma*.) *Spasmodicus*. SPASMATODES, (*σπασματώδης*, from *spasmus*, and *odes*.) *Spasmodicus*.

SPASME, *Spasm*—s. de la Glotte et du Thorax, *Asthma thymicum*—s. de la Vessie, *Cystospasmus*.

SPASMI, see *Spasm*.

SPASMODES, (*σπασμῳδης*.) *Convulsive*.

SPASMODIC, *Spasmodicus*.

SPASMOL'OGY, *Spasmolog'ia*, (*spasmus*, and *λογος*, 'a treatise'.) A treatise on spasm.

SPASMOLYGMUS, (*spasmus*, and *λυγμος*, 'hiccough'.) *Singultus*.

SPASMOPHIL'IA, *Convulsibil'itas*, *Convulsibil'ity*, (F.) *Convulsibilit'as*: (*spasmus*, and *φιλειω*, 'I love'.) Tendency to convulsions from unusual excitability of the motor nervous system.

SPASMOTICUS, *Spasmat'icus*, *Spasmatode's*, *Spas'ticus*, *Spas'tic*, *Spasmod'ic*, *Spasmat'ic*, *Spasmat'ic*, (F.) *Spasmodique*, *Spas'tique*, (from *spasmus*.) Anything relating to spasm. Also, an antispasmodic.

Morbi constrictorii, *M. spasmodici*, &c., are diseases accompanied with spasm.

SPASMUS, (*σπασμος*.) *Convulsion*, *Spasm*—s. *Aurium*, *Otalgia*—s. *Bronchialis*, *Asthma*—s. *Caninus*, see *Canine laugh*—s. *Clonicus*, *Convulsion*—f. *Cynicus*, see *Canine laugh*—s. *Facialis*, *Tic*—s. *Glottidis*, *Asthma thymicum*—s. *Linguae*, *Glossospasmus*—s. *Maxillae inferioris*, *Trismus*—s. *Muscularis*, *Cramp*—s. *Musculorum Faciei*, *Canine laugh*—s. *Nutans*, *Convulsion*, *salaam*—s. *Periphericus*, *Horrida cutis*—s. *Universalis*, *Synclonus*—s. *Ventriculi*, see *Cardialgia*—s. *Vesicae*, *Cystospasmus*.

SPAS'NIA, (same etymon as *Spasm*.[?]) A term used by Mercurialis to designate the lancinating pain produced, at times, in the chest by violent fits of coughing.

SPASTIC, (*σπαστικός*.) *Spasmodicus*, *Tonic spasm*.

SPASTICUM, *Tetanic*.

SPATHA, (*σπαθη*, a broad, flat instrument for stirring liquids, spreading plasters, &c.) *Hypaleipton scapula*.

SPATHES'TER, (from *σπaw*, 'to draw'.) A surgical instrument, used by P. Amman for drawing the prepuce over the glans, when too short.

SPATHOMELE, (*σπαθη*, and *μηλη*, 'a probe,') *Spatula*.

SPATHULA, *Scapula*, *Spatula*—s. *Fœtida*, *Iris fœtidissima*.

SPATHYEMA FÆTIDA, *Dracontium fœtidum*. SPATIA INTERLOBULARES, (pl. of *spatium*, 'a space,') *Interlobular fissures*.

SPAT'ILE, (*σπαρλή*, 'human excrement'.) A liquid fecal evacuation. *Excrement*.

SPATIUM ORIS, 'space of the mouth,' Mouth—s. *Trigonum*, *Lyra*.

SPATTERDOCK, *Nuphar advena*.

SPAT'ULA, *Spath'ula*, (dim. of *Spathr*.) *Spathome'le*, *Specill'um latum*. A flat instrument used for spreading plasters, stirring ointments, holding down the tongue, &c. Also, the *scapula*.

SPATULA PRO ORE, *Glossospatha*.

SPAUD, *Humerus*.

SPAUL, *Saliva*.

SPAUTBONE, *Humerus*.

SPAUBONE, *Humerus*.

SPAYING, (from *σπaw*, 'to draw'; [L.] *spado*, 'a castrated person,') see *Castration*.

SPEARWORT, *Ranunculus flammula*.

SPEAUTER, *Zinc*.

SPECIAL, see *Specific*.

SPEC'IALIST, (F.) *Spécialiste*, (from *species*,

'a peculiar sort.') One who devotes himself to a specialty; as to diseases of particular parts, as of the eye, ear, chest, &c.

SPECIES, (F.) *Espèces, Spécialité*. Official mixtures of substances cut or bruised. Sometimes used for compound powders, as it always appears to have been formerly. See *Pulvis*.

From *species* comes *spice*, (F.) *Épice*.

SPECIES AROMATICÆ, *Pulvis cinnamomi compositus*—s. *Diacinnamomi*, *Pulvis cinnamomi compositus*—s. *Diaceretæ*, *Pulvis cretæ compos.*—s. *Dialapæ*, *Pulvis jalapæ compositus*—s. *Diambræ sine odoratis*, *Pulvis cinnamomi compositus*—s. *Diatrageanthæ frigidæ*, *Pulvis tragacanthæ compositus*—s. *Hieræ pieræ*, *Pulvis aloes cum canellâ*—s. *Lactificantes Rhazis*, *Pulvis cinnamomi compositus*—s. e *Scordio cum opio*, *Pulvis cretæ compositus cum opio*—s. e *Scordio sine opio*, *Pulvis cretæ compositus*.

SPECIFIC, *Spécif'ique, Nosocrat'ic*, (F.) *Spécifique* (*species*, 'a peculiar sort,' and *facere*, 'to make.') A substance to which is attributed the special property of removing some particular disease. Probably no such remedy exists. *Mercury* in syphilis, and *sulphur* in the itch, have been regarded as the strongest examples. The word is also used adjectively in the sense of *special*; as "mercury has a *specific* or *special* action on the salivary glands." See *Specificity*, morbid.

SPECIFIC OF HERRENSCHWAND. A once celebrated German vermifuge. It is said to have consisted of 10 grains of *Gamboge*, with 20 of *Carbonate of Potassa*. It is affirmed, that mercury and arsenic have also been found in it.

SPECIFICITÉ MORBIDE, *Specificity*, morbid.

SPECIFICITY, MOR'BID, (F.) *Spécificité morbide*, is the occult quality of a disease which gives occasion to peculiarity in its nature, evolution and treatment. Such is said to be a *specific* disease.

SPECIFICUM PARACELSI, *Potassæ sulphas*.

SPECILUM, *Me'lè, Sty'lus, Explorato'rium*, (from *specio*, 'I examine,') *Specula'rium, a Probe*, (F.) *Stilet, Stylet, Sonde*. A surgical instrument, employed for examining wounds, fistulas, and for passing setons, &c. It is usually formed of silver, and is terminated, at one end, by an olive-shaped button. The *eyed probe* has an aperture at the other extremity.

The perforator of the Trocar is called a *Stylet'*; as well as the wire or metallic rod of the flexible catheter, which gives it shape and firmness.

SPECILLUM, *Everriculum, Hypaleiptron*, Sound—s. *Auricularium*, *Aprryomele*—s. *Cereum*, *Bougie*—s. *Excavatum*, *Stylus excavatus*—s. *Latum*, *Spatula*—s. *Minus*, *Melotis*—s. *Sulcatum*, see *Grooved*.

SPECIOLÉ, *Species*.

SPECKS OF THE CORNEA, see *Caligo*.

SPECLARION, (*σπεκλαριον*,) *Specillum*.

SPECTACLES, (*spectare*, 'to behold,') *Conspici'ula*, (Sc.) *Breels*, (F.) *Besicles, Lunettes ordinaires, Conserves*. Glasses to assist the sight; arranged so as to be adapted to both eyes. These glasses are more or less concave or convex, according as the sight is more or less short, (*myopic*), or long, (*presbyopic*.) When the glass is adapted to one eye, it is called an *Eyeglass*, *Conspici'ulum*, *Perspici'ulum*, *Vi'trum ocula'rè*.

SPECTRUM, *Spectre*, (F.) *Spectre*, (from *specio*, *spectrum*, 'to see,') An appearance real or imaginary.

SPECTRUM ANALYSIS, *Prismat'ic analysis*. A most delicate method of analysis, which consists in decomposing, by means of a prism, the coloured light given by the salts of metals to the colourless flames of alcohol or coal gas. It is a useful aid to quantitative analysis, as the coloured flame

of each metal, even in the minutest quantities, is found to give a well marked and characteristic spectrum. Bunsen estimated, that the amount of sodium which admitted of detection by prismatic analysis was the 195,000,000th part of a grain; of lithium the 70,000,000th; of potassium, the 60,000th; of barium the same; of strontian, the 1,000,000th; and of calcium the 100,000,000th of a grain. See *Poisons*.

SPECTRUM, OCULAR, *Spect'rum ocula'rè*, (F.) *Spectre oculaire*. A disturbance of vision, which consists in the appearance of objects after the objects themselves have been withdrawn. See *Colours*, accidental.

SPECTRUM, SOLAR, *Spect'rum Sola'rè*, (F.) *Spectre solaire*. The coloured image produced by white light, which has been decomposed by passing through a prism. It consists of seven different coloured fringes, arranged in the following order—red, orange, yellow, blue, green, indigo, and violet. Three of these—blue, red, and yellow—are simple; the others being mixtures of them.

SPECULARIS LAPIS. A transparent mineral, but of what nature is not clear, which was formerly employed in epilepsy. In old times it was used for glass.

SPECULUM, Catop'ter, Catop'tron, Diop'tron. (In Latin, 'a mirror,' from *specio*, 'to behold,') In surgery, it means different instruments for dilating cavities, and facilitating their examination. See *Dilator*. There are various instruments of this kind,—the *S. Ani*, *S. Auris*, *S. Vagina*, (*Colpeurynter*, *Elytreynter*,) *S. Matr'icis*, *S. Oculi*, *S. Oris* ([F.] *Bâillon*, *Pas d'Âne*,) *S. Gutturis*, *S. Vesicæ*.

SPECULUM CITRINUM, *Orpiment*—s. *Indicum*, *Ferri limatura*—s. *Larynx*, *Laryngoscope*—s. *Lucidum*, *Septum lucidum*—s. *Oculi*, see *Ophthalmoscopia*—s. *Oris*, *Glossocatochus*—s. *Veneris*, *Achillea millefolium*.

SPECUS, gen. *Spécus*, 'a cave,' *Vulva*—s. *Cor-dis*, *Ventricle of the Heart*—s. *Pro medullâ spinali*, see *Vertebral column*—s. *Vertebralis*, see *Vertebral column*.

SPEDALSKE, see *Radzyge*.

SPEDALSKEHD, see *Radzyge*.

SPEECH, ([Sax.] *præc*,) *Voice*, articulated.

SPEECHLESSNESS, *Mutitas*.

SPEEDIMAN'S PILLS, see *Pilulæ aloes et myrrhæ*.

SPEEDWELL, BROOKLIME, *Veronica beccabunga*—s. *Female*, *Antirrhinum elatine*—s. *Offi-cinal*, *Veronica*—s. *Purslain*, *Veronica peregrina*—s. *Water*, *Veronica beccabunga*.

SPELLBONE, *Fibula*.

SPELTRUM, *Zincum*.

SPERAGE, *Asparagus*.

SPERAGUS, *Asparagus*.

SERM, gen. *Sperm'atis*, *Sperma*, (*σπερμα*, gen. *σπερματος*, from *σπειρω*, 'I sow,') *Spermatic fluid* or *liquor*, *Sem'inal fluid*, *Seed*, *Se'men*, *S. viri'lè* seu *masculi'nium* seu *genita'lè*, *Semini'um*, *Genita'lè*, *Medull'æ Ros*, *San'guis*, *Se'rum*, *Hu'mor genita'lè* seu *semina'tis* seu *vene'reus*, *U'i'ma genita'lè*, *Genitu'ra*, *Sper'ma viri'lè*, *Spermi'um*, *Tho'rus*, *Lac ma'ris*, *Lagne'a*, *Lagni'a*, *Lagneu'ma*, *Ger'men*, *Vita'lè vi'rus*, *Male's milk*, *Prop'agatory* or *gen'ital liq'uor*, *Vit'al or quick'ening ven'um*, (F.) *Semence*, *Fluide séminal*. A whitish, viscid fluid, of a peculiar smell, secreted by the testicles, whence it is carried by the vasa deferentia to the vesiculæ seminales, to be thrown into the vagina, during coition, through the ejaculatory ducts and the urethra. It is the fecundating fluid, and must come into actual contact with the germ of the female. The *Aura sem'inis*, *Gonaur'a*, is incapable of effecting fecundation. The semen, at the time or emission, is composed of two different

fluids; the one liquid and transparent, which is considered to be the secretion of the prostate, — the other, white, and as thick as mucilage; the product of the testicular secretion. The sperm contains, according to Vauquelin, 900 parts of water, 50 of animal mucilage, 10 of soda, and 30 of calcareous phosphate. The animal matter is peculiar, and by some termed *spermatin*.

Microscopic observations show that it contains *spermatozo'a*, or more properly *spermatozo'ida*; for their animalcular nature is not demonstrated. They are produced in cells — *spermcells* — and have probably no more title to be considered animalcular, than the cilia of the ciliated epithelium. By careful examination, other minute, round, granulated bodies may almost always be detected, which are in all cases much less numerous than the spermatozoa. These bodies are the *seminal granules*, *granula seminalia*. Pure sperm, in its most perfect state, consists principally of spermatozooids and seminal granules; both of which are enveloped in a small quantity of fluid, *liquor seminalis*, *spermatic liquor*.

It has been imagined, but erroneously, that during coition there is a secretion of female sperm — *Semen mulieb're*, *Thelygonum*. The increased secretion that takes place is chiefly from the glands of Duverney.

Sperm also means spermaceti.

SPERM CELL, see Sperm.

SPERMA, Sperm — s. Mercurii, Hydrargyrum acetatus.

SPERMA RANA'RUM, *Sperni'ola* seu *Sperni'olum* seu *Spermi'ola* seu *Spermi'olum rana'rum*. Frog's spawn. Once used in medicine.

SPERMACETI, (*sperma*, and cetus, 'a whale,') Cetaceum — s. Whale, see Cetaceum.

SPERMACRASIA, (*sperma*, a, priv. and *κρasis*, 'mixture,') Spermatorrhœa.

SPERMATACRASIA, Spermatorrhœa.

SPERMATACRATIA, (*sperma*, a, priv. *κρatos*, 'power,') Spermatorrhœa.

SPERMAT'IC, *Spermat'icus*, *Seminal'is*, *Seminal*, (F.) *Spermatique*, (*σπερματικός*, from *sperma*.) That which relates to the sperm. A name given to different parts connected with the organs of generation.

SPERMATIC ARTERIES, *Arte'rie Spermat'ice* seu *preparan'tes*, (F.) *Artères testiculaires*, *A. de l'ovaire* (Ch.), are two in number — one on each side — and arise from the sides of the aorta, sometimes from the renal arteries. They descend, almost vertically, at the sides of the vertebral column, and are distributed differently in the two sexes. In man, the spermatic artery, situate at the side of the vas deferens, issues by the abdominal ring; gives numerous ramifications to the spermatic cord, and divides into fasciculi of branches, one of which goes to the testicle, — the other to the epididymis. In the female, the spermatic artery, *ova'rian* or *uterovarian artery*, dips into the pelvis, and passes to the ovary, Fallopian tube, and round ligament.

SPERMATIC CORD or CHORD, *Testicular Cord*, *Funiculus Spermat'icus*, *Spermatochord'a*, (F.) *Cordon spermaticum* ou *testiculaire*. The vascular and nervous chord, by which the testicle is suspended. It is composed of the spermatic artery and veins; of other small vessels; of lymphatics; of nervous filaments from the spermatic plexus, and from the genito-crural branch of the lumbosacral plexus; of the vas deferens, and, very often, of a fibrocellular cord, which unites the peritoneum to the upper part of the tunica vaginalis, and in which encysted hydrocele of the spermatic cord occurs. All these parts are united together by a very lax, areolar tissue, and surrounded by coats, which, reckoning from without, are: — 1. The skin and areolar membrane.

2. A fibrocellular membrane, formed by the fascia superficialis. 3. A very thin layer, formed by fibres of the cremaster muscle, united archwise before, and often also behind, the cord. 4. The proper sheath of the spermatic vessels, or the tubular prolongation furnished by the fascia transversalis to the chord, on a level with the superior orifice of the inguinal canal. The spermatic cord is commonly shorter on the right side than on the left; and of a different size in different individuals. It ascends almost vertically, from the superior margin of the testicle as far as the lower orifice of the inguinal canal; passes through this canal and enters the abdomen, crossing the epigastric artery. Here it forms an evident elbow, directing its course backward. At this part, also, the organs composing it separate from each other: — the vas deferens descending into the pelvis to pass behind the bladder; — the bloodvessels and lymphatics ascending towards the lumbar region, &c.

SPERMATIC FILAMENTS, Spermatozoa — s. Fluid, Sperm.

SPERMATIC GANG'LION. A large ganglion, formed by branches from the hypogastric ganglion, and from the spermatic plexus. It supplies the fundus uteri. Besides these ganglia, Dr. Robt. Lee describes *vesical* and *vag'inal ganglia*, and anterior and posterior *subperitoneal ganglia* and *plexuses*, which communicate with the preceding, and constitute an extensive nervous rete over the entire uterus.

SPERMATIC LIQUOR, Liqueur seminis, see Sperm. SPERMATIC PAS'SAGES or WAYS, *Vie Spermat'ice*, are the canals concerned in the excretion of semen. See Testicle.

SPERMATIC PLEX'USES of nerves, *Plex'us testicular'es*, are two in number, and are furnished by the renal plexuses. Their filaments, called *Spermatic nerves*, follow the arteries of the same name to the testicle in man; and to the ovary and Fallopian tube in the female, — *ova'rian nerves*. They cannot be traced into the substance of these organs.

SPERMATIC VEINS — *ve'næ spermat'ice* — are two or three in number on each side. They accompany the spermatic artery, and open — those of the right, into the vena cava inferior; those of the left, into the corresponding renal vein. These veins form, above the testicle, a kind of venous network, called, by some, the *Spermatic plex'us* and *corpus varicosum*; and another plexus before the psoas muscles, called the *Corpus pampiniforme*.

SPERMATIN, see Sperm.

SPERMATIQUE, Spermatic.

SPERMATIS'MUS, *Emis'sio seminalis*, *Spermob'ole*, (from *sperma*.) The emission of sperm.

Sperm'atism or *Animal'culism* is the doctrine which maintains that the embryo is produced by the so called spermatic animalcules.

SPERMATIST, Animalculist.

SPERMATOCE'LE, *Hernia seminalis scro'ti*, *Oscheoce'le seminalis*, *Gonocoe'le*, (*sperma*, and *κηλη*, 'a tumour,') The ancients gave this name to certain swellings of the testicle which were regarded as produced by an accumulation of sperm in the organ. Also, varicocele.

SPERMATOCHORDA, (*sperma*, and *chorda*.) Spermatic chord.

SPERMATOCLEMMMA, (*sperma*, and *κλεμνω*, 'to steal,') see Pollution.

SPERMATOCLEPSIS, see Pollution.

SPERMATOCYSTIDES, (*sperma*, and *κυστις*, 'a bladder,') Vesiculæ seminales.

SPERMATOCYSTIT'IS, (*spermatoecystis*, and *itis*.) Inflammation of a vesicula seminalis.

SPERMATOCYSTIDORRHAG'IA, *Caulorrhag'ia ejaculato'ria*, *Hæmat'ria ejaculato'ria* seu *seminalis*, (*spermatoecystides*, and *ῥαγη*, 'a

seu *einefor* me seu ca neō comparatū seu *pheno-*
noida' seu *multifor* me seu az'gyos seu *papilla-*
rē seu *polymer'*phon seu *paxilla'rē* seu *baçilla'rē*
seu *alæfor* mē seu *sphecoi'deum* seu *vespifor* mē
seu *inconjunctum* seu *pterygoideum* seu *abūm*
seu *colato'rii*, *Pterygoid bone*. An zygous bone,
situate on the median line, and at the base of the
cranium. It articulates with all the other bones
of that cavity; supports them, and strengthens
their union. Its form is singular, and resembles
a bat with its wings extended. It has, 1. *A*
inferior or *guttural surface*, on which is situate
the *crista*, that joins the vomer; a channel, which
concurs in forming the pterygopalatine foramen;
the pterygoid process; the pterygoid fossa; the
scaphoid depression; the Vidian or *-terygoid*

canal; the foramina—ovale, spinale, &c. 2. A *superior or cerebral surface*, on which are:—the clinoid processes; the pituitary fossa; the *foramina (ovale, rotundum, and spinale)*; the *Apophysis of Ingrassias or lesser wing*; the foramen opticum, &c. 3. An *occipital or posterior surface*, which is articulated with the basilar process of the occipital bone. 4. An *anterior or orbitary nasal surface*; having, anteriorly, a *crista* to unite with the ethmoid bone; and, on each side, a round aperture, which leads into two cavities in the substance of the bone, separated by a middle septum, and called the *sphenoidal sinuses* or cells. 5. Two *zygomatotemporal or external surfaces*, which correspond to the temporal and zygomatic fossæ.

Some divide the sphenoid into *body or middle portion*; and *ala*, which are four in number, and are subdivided into *great (Temporal Plates or Wings, Alæ magnæ)* and *little (Apophyses of Ingrassias, Alæ minores, Processus ensiformes)*.

The *Sphenoid suture* surrounds the bone.

SPHENOID, SPINOUS PROCESS OF THE, Sphenoid spine.

SPHENOIDAL, Sphenoïdalis. That which belongs or relates to the sphenoid bone.

SPHENOIDAL CELLS, see *Sphenoid bone*.

SPHENOIDAL CORNUA, Cornua Sphenoïdalis, Ossicula Bertini, Ossa triangulæria, Pyramids of Wislar, (F.) Cornets sphénoïdaux ou de Bertin, Osselets de Bertin. Two small, thin, and curved bones, situate between the sphenoid and ethmoid, with which they are confounded in the adult. They have the shape of a hollow pyramid, with the base turned backwards, and are developed by a single point of ossification. They are articulated with the sphenoid, ethmoid, palate bone, and vomer.

SPHENOIDAL OR SUPERIOR ORBITARY FISURE, Foramen lac'rum superius, (F.) Fente sphénoïdale, is a large fissure, situate between the great and little ala of the sphenoid. It is seen at the upper and back part of the orbit, between which and the cranium it is a means of communication, and transmits the third, fourth, and sixth pair of nerves, and the first branch of the fifth pair, to the eye and its appendages.

SPHENOIDAL SINUSES, see *Sphenoid*.

SPHENOIDAL SPINE, Spina seu Crista sphenoidalis, (F.) Épine sphénoïdale, Spinous Process of the Sphenoid;—1. A projecting crista at the inferior surface of the sphenoid bone, for articulation with the vomer. 2. A triangular process, *Apophyse sous-temporale (Ch.)*, met with near the posterior margin of the same bone, behind the foramen spinale. At the point of the spinous process, a *styloid process* is frequently met with.

SPHENOIDES, (σφηνοειδής.) Cuneiform, Sphenoid—s. Os, Sphenoid bone.

SPHENOMAXILLARY, Sphenomaxillaris. That which relates to the sphenoid and maxillary bones.

SPHENOMAXILLARY FISURE, Inferior orbitary fisure, (F.) Fente sphéno-maxillaire ou orbitaire inférieure; called, also, *Foramen lac'rum inf'rius seu sphenomaxillare*, is situate at the posterior part of the angle formed by the union of the internal and inferior parietes of the orbit. It is constituted, above, by the sphenoid bone; below, by the superior maxillary and palate bones; and, before, by the malar bone. It is narrower at the middle than at the extremities, and forms a communication between the orbital cavity and the zygomatic fossa.

SPHENOMAXILLARY FOS'SA is a name given by some anatomists to a depression at the union of the sphenomaxillary and pterygomaxillary fissures.

SPHENOÛRBITAR. A name given by Bé-

clard to the anterior part of the body of the sphenoid bone, which is developed by a variable number of points of ossification.

SPHENOPALATINE, Sphenopalatinus. That which relates to the sphenoid and palate bones.

SPHENOPALATINE ARTERY, Large lateral nasal A., is the termination of the internal maxillary. It enters at the posterior part of the superior meatus of the nose, through the sphenopalatine foramen, and spreads its numerous branches on the pituitary membrane covering the septum, the cornua, and the meatus.

SPHENOPALATINE FORAMEN is a round aperture, formed by the vertical portion of the os palati and the sphenoid. It establishes a communication between the nasal fossæ and the zygomatic fossa.

SPHENOPALATINE GANGLION, Ganglion of Meckel, Sphenoidal G. (Ch.) A small, nervous, cordiform, or triangular ganglion, of variable size, situate without the foramen sphenopalatinum, in the pterygomaxillary fissure. It seems suspended by several nervous filaments to the trunk of the superior maxillary nerve, and gives off *internal or sphenopalatine filaments, inferior or palatine filaments,* and a *posterior filament*, which is the *Vidian or pterygoid nerve*.

SPHENOPALATINE NERVES, Lateral nasal nerves, arise from the ganglion—just described—at its inner part, and enter the nasal fossæ by the sphenopalatine foramen. They are five or six in number, and distribute their filaments to the outer and inner parietes of the nasal fossæ. One of the most remarkable branches is the *Nasopalatine*.

SPHENOPALATINUS, Levator palati.

SPHENOPARIETAL, Sphenoparietalis. That which belongs or relates to the sphenoid and parietal bones.

SPHENOPARIETAL SUTURE is formed by the articulation of the extremity of the greater ala of the sphenoid with the anterior and inferior angle of the parietal bone.

SPHENOPTERYGOPALATINUS, Circumflexus—s. Salpingo-mallæen, Laxator tympani—s. Salpingostaphylinus, Circumflexus.

SPHENOSIS, (σφηνωσις.) see *Wedge*.

SPHENOSTAPHYLINUS, Levator palati.

SPHENOTEMPORAL, Sphenotemporalis. That which belongs to the sphenoid and temporal bones.

SPHENOTEMPORAL SUTURE is the suture at the articulation of the great ala of the sphenoid bone with the squamous portion of the temporal. Béclard gave the name *sphenotemporal* to the posterior part of the body of the sphenoid, which is developed by distinct points of ossification.

SPHEX, (σφήξ.) Wasp.

SPHINCTER, gen. Sphincteris, (σφιγκτήρ,) Constrictor, Musculus constrictorius, (from σφίγγω, 'I constrict.') A name given to several annular muscles, which constrict or close certain natural openings.

SPHINCTER ANI, Annularis Ani. Many anatomists have described two sphincter muscles of the anus:—1. The *S. exterius seu cutaneus, Aspidicus, Coccygio-entané-sphincter, Orbicularis Recti, Constrictor Ani, Coccygio-anal, (Ch.)* It is situate around the anus; is elliptical from before to behind; flattened, and pierced at its middle. Its fibres describe concentric arcs, which are attached, behind, to the extremity of the coccyx, by a dense, areolar substance; and are confounded, anteriorly, with the bulbo-cavernosi and transversi perinæi muscles. This muscle contracts and closes the anus. 2. The *in'ner or internal Sphincter Ani, Sphincter intestinalis* of Winslow, is by many anatomists considered as the termination of the circular fibres of the rectum. It is annular, and situate around the infe-

ral extremity of the rectum, to the extent of about a finger's breadth. It has the same uses as the other.

SPHINCTER GULÆ, Constrictores pharyngis—s. Ilei, Baubini, valve of—s. Intestinalis. Sphincter ani internus—s. Labiorum, Orbicularis oris—s. Oculi, Orbicularis palpebrarum—s. Palpebrarum, Orbicularis palpebrarum—s. Pylori, see Pylorus—s. Vaginae, Constrictores cunni—s. Ventriculi, Pylorus.

SPHINCTER VESTIGÆ, *Sphincter of the bladder*. Some anatomists have given this name to whitish, elastic, and circular fibres, which surround the neck of the bladder, but do not constitute a particular muscle. Morgagni has given the name **PSEUDOSPHINCTER** to the anterior fibres of the levator ani, which pass beneath the neck of the bladder, and, by their contraction, close that opening. See Compressor urethræ.

SPHONDYLIIUM, (σφονδύλιον,) Heracleum spondylium.

SPHONGUS, (σφονγος,) Fungus, Spongia.

SPHRAGIDONYCHARGOCOME'IA, (σφραγίς, 'a seal,' ονυχ, gen. ονυχος, 'a nail,' αργος, 'white,' or perhaps αργυρος, 'silver,' and κομω, 'I adorn.') A charlatan who adorned his fingers to the very nails with rings.

SPHYGMA, (σφυγμν,) Pulse.

SPHYG'MIC, *Sphygm'ic*, (F.) *Sphygmique*, (from *sphygma*.) Relating to the pulse; pulsatory.

SPHYGMIC ART, Sphygmologia.

SPHYGMICA DOCTRINA, Sphygmologia.

SPHYGMICUS, (σφυγμικός,) Throbbing.

SPHYGMOCEPHALUS, (σφυγμος, 'pulse,' and κεφαλή, 'head.') Cratyphæ.

SPHYGMODES, (σφυγμωδής, from *sphygmus*, and *odes*,) Throbbing.

SPHYG'MOGRAPH, (*sphygmus*, and γραφω, 'I describe,') An instrument which, when applied over an artery, indicates the character of the pulse as to force and extent of undulations.

SPHYGMOLEGIUM, (*sphygmus*, and *lego*, 'I read,') Pulsilegium.

SPHYGMOLOGE, (F.) *Sphygmolog'ium*, (*sphygmus*, and *λεγειν*, 'to indicate,') An instrument for showing the quickness or other qualities of the pulse.

SPHYGMOLOG'IA, *Sphygmology*, *Sphygmica Doctrina*, *Ars Sphygmica*, (*sphygmus*, and *λογος*, 'a description,') The doctrine of the pulse.

SPHYGMOMETER, Pulsilegium.

SPHYGMOSCOPE, (*sphygmus*, and σκοπεω, 'to examine,') Pulsilegium.

SPHYGMOSCOPIUM, Pulsilegium.

SPHYGMUS, (σφυγμος,) Pulsation, Pulse.

SPHYRA, (σφυρα,) Malleolus.

SPHYRON, (σφυρον,) Malleolus.

SPHYXIS, (σφυξίς,) Pulsation.

SPIC, Lavandula Spica.

SPICA, *Fas'cia re'pens*, *Spica ban'dage*, (F.) *Épi*. A bandage so called in consequence of its somewhat resembling a *spike* of barley. The turns of the bandage cross like the letter V;—each leaving a third of the roller uncovered. It is distinguished into *ascending* and *descending*. It may be applied over various parts of the body, and in a different manner in each case: thus, there is the *Spica seu Fas'cia inguinalis*, *Spica inguinalis duplex*, the *spica* for the shoulder, and another for the thumb.

SPICA, Lavandula Spica—s. Alpina, Valeriana Celtica—s. Celtica, Valeriana Celtica—s. Indica, Nardus Indica—s. Nardi, Nardus Indica.

SPICE, see Species.

SPICEBERRY, Gaultheria, Laurus Benzoin.

SPICEBUSH, Laurus Benzoin.

SPICES, FOUR, see Myrtus pimenta.

SPICEWOOD, Laurus Benzoin.

SPICILLUM, Spicillum.

SPIDER, (from *spin* [?]), see Araneæ tela.

SPIDER'S WEB, Araneæ tela.

SPIDERWORT, Liliago, Tradescantia Virginica.

SPIGELIA, (after Adrian Spigelius, the anatomist,) S. Marilandica.

SPIGELIA ANTHELMIA, *Demera'ra Pink'root*, (F.) *Spigélie anthelmintique*, Brinvilliers, Brinvillière, a native of South America and the West India Islands, has properties like the

SPIGELIA MARILAN'DICA, S. *Lonic'ra*, *Anthelmia*, *Lonic'ra Marilandica*, (F.) *Spigélie du Maryland*; *Perennial Wormgrass* or *Indian Pink*, *Carolina Pink*, *Star' bloom*, *Worm' root*, *Nat. Ord. Gentianaceæ*. *Class*, Pentandria. *Order*, Monogynia. *Indigenous*. The root—*Spigelia* (Ph. U. S.), *Pinkroot*—is celebrated as an anthelmintic, particularly in cases of lumbrici.

It is, also, asserted to have been found serviceable in remittent fever. It is a narcoticoâcid. Dose, gr. x to ʒss.

SPIGNEL, *Æthusa meum*.

SPIKENARD, (*spica*, 'a spike,' and *nardus*, 'nard,') Conyza squarrosa, Nardus Indica—s. American, Aralia racemosa—s. False, Smilacina racemosa—s. Small, Aralia nudicaulis—s. Tree, Aralia spinosa.

SPILANTHES, Spilanthus.

SPILANTHUS ACMEL'LA, S. *ciliata* seu *fimbriata*, *Spilanthes acmel'la*, (σπίλος, 'a spot,' and *ανθος*, 'a flower,' owing to its dotted or speckled flower,) *Bidens acmel'la*, *Achmel'la*, *Acme'lla*, A. *Mauritia'na*, *Verbesi'na acmella*, *Balm-leaved Spilanthus*. *Ord.* Composite. *Sex. Syst.* Syngenesia Polygamia æqualis. This plant possesses a glutinous, bitter taste, and fragrant smell. The herb and seed are said to be diuretic and emmenagogue. They have been used in dropsies, jaundice, fluor albus, and calculous complaints; given in infusion.

SPILANTHUS, BALM-LEAVED, Spilanthus acmella—s. Ciliata, S. acmella—s. Fimbriata, S. acmella.

SPILANTH'US OLERA'CEUS, *Spearleaved Spilanthus*, *Bidens fer'vida*, (F.) *Cresson de Para*. A tincture of the plant has been recommended in toothache, (F.) *Paraguay-rouze*.

SPILI, (σπίλος, 'a spot or stain,') see Nævus.

SPILOMA, (σπίλωμα,) see Nævus.

SPILOPLAXIE, (σπίλος, 'a spot,' and πλάξ, 'a plate,') Elephantiasis of India.

SPILOSIS, (from σπιδος, 'a spot,' and *osis*,) Epichrosis—s. Ephelis, Ephelides—s. Poliosis, Poliosis.

SPILS'BURY'S ANTISCORBU'TIC DROPS.

An empirical preparation, formed of *hydrarg. oxymer.*, *rad. gentian.*, *cort. aurant.*, *sicc. aa ʒij*; *antimon. crud.*, *aantal. rubr. aa ʒj*; *spiritus vini rectif.*, *aquæ, aa ʒviij*.

SPINA, 'a thorn,' *Spine*, (F.) *Épine*, old [F.] *Espine*. A process on the surface of a bone, which has been compared to the spines or thorns on certain vegetables. The chief processes of this name are:—the *nasal spine*, the *spine of the scapula*, the *spine of the ischium*, the *four iliac spines*, the *palatine spine*, the *maxillary*, the *sphenoid*, &c. The *spine of the back* is the collection of vertebrae constituting the *vertebral column*.

SPINA, Penis—s. *Acida*, *Oxyæantha Galeni*—s. *Ægyptiaca*, see *Acæcie gummi*—s. *Alba*, *Cardus marianus*, *Mespilus oxyæantha*, *Onopordon acanthium*—s. *Bifida*, see *Hydrorachis*—s. *Cervina*, *Rhamnus*—s. *Domestica*, *Rhamnus*—s. *Dorsi*, *Vertebral column*—s. *Dorsi introrsum flexa*, *Lordosis*—s. *Ferrea*, *Pin*—s. *Helmontii*, *Aiguillon*—s. *Hirci*, *Astragalus verus*—s. *Infectoria*, *Rhamnus*—s. *Nasi*, see *Nasus*—s. *Nodosa*, *Rachitis*—s. *Sphenoidalis*, *Sphenoidal spine*.

SPINA VENTOSA, *Spina ventositas*, *Tere'do*,

Ostathrocacæ, *Fungus seu Tu'mor fungosus articuli*, *Lu'pia juncturae*, *Hyperspin'gia*, *Flatus spinæ*, *Arthroacæ*, *Podarthrocacæ*, *Sidera'tio Os'sis*, *Can'cer seu Gangra'na Os'sis*, *Ecosto'sis*, *White Scell'ing* (of some). A term of no definite meaning, as is obvious from these various words having been considered its synonyms. By some, it is defined to be—a disease of the osseous system, in which the texture of the bone dilates, seeming to be distended with air, and constituting a variety of osteosarcoma. By others, it is considered to be a tumour arising from an internal cavity of a bone; occurring most frequently in the carpus or tarsus. The term itself is a translation from the Arabic of Rhazes. See, also, *Hydrarthrus*, and *Mollities Ossium*.

SPINA VERTEBRALIS, Vertebral column.

SPINACH, Spinacia.

SPINACHIA, Spinacia.

SPINA'CIA, *Spina'chia*, *Spina'cia olera'cea*, *Spin'age*, *Spin'ach*, (from *spina*, 'a thorn or prickle,') (F.) *Épinard*. Ord. Chenopodiaceæ. Sex. Syst. Dioecia Pentandria. A plant which resembles the cabbage in its dietetic powers. The leaves boiled, with the addition of oil, form a good emollient cataplasm. It has been used in phthisical complaints; but its medicinal properties, if it have any, are not now regarded.

SPINACIA OLERACEA, Spinacea.

SPINÆ, (pl. of *spina*.) Spinous processes—s. *Ventositas*, *Spina ventosa*.

SPINAGE, Spinacia.

SPI'NAL, *Spina'lis*, *Spino'sus*, *Spina'tus*, *Rhachi'tes*, *Rachid'ian*, (F.) *Épinière*, *Rachidien*, (from *spina*, 'spine,') That which relates to the vertebral column.

SPINAL ACCESSORY, Spinal nerve.

SPINAL ARTERIES, *Arte'rie spin'ales*, are two in number, viz.: 1. The posterior *spin'al*, *Artère médiane postérieure du Rachis*, (Ch.) It arises from the vertebral, near the corpora pyramidalia, and descends on the posterior surface of the spinal marrow, distributing its ramifications to it. 2. The anterior *spin'al artery*, *A. médiane antérieure*, (Ch.,) is larger than the last, and rises, also, from the vertebral. It descends, in a serpentine manner, upon the anterior surface of the marrow; furnishes ramusculi to it, and unites with that of the opposite side, opposite the foramen magnum occipitis. A very tortuous branch arises from this union, which descends as far as the inferior extremity of the marrow, to which it sends numerous divisions.

The term *Spin'al Ar'teries* or *Rachid'ian Ar'teries* is also given, in the abstract, to all the arteries of the spinal marrow and vertebral canal. The same may be said of the veins and nerves.

SPINAL CORD, *Medulla spinalis*.

SPINAL FORAM'INA, (F.) *Trous rachidiens*, in the abstract, are the foramina formed by every two contiguous vertebrae, through which the spinal nerves issue. See Vertebral. The term *Foramen Spina'lè seu spinosum* is especially applied, however, to a small foramen, in front of the spinous process of the sphenoid bone, through which the middle artery of the dura mater enters the cranium. It is, likewise, called *Foramen sphenospinosum*, (F.) *Trou Sphéno-épineux* ou *petit rond*, *Trou épineux*.

SPINAL GLANDION, see Encephalon.

SPINAL IRRITATION, *Rhachialgi'tis*, *Rhachial'gia*, *Neural'gia spina'lis*, *Notal'gia*. A modern pathological view, which refers most nervous diseases to irritation of the spinal cord. This irritation is presumed to be indicated by tenderness on pressure over the spinous process of one or more vertebrae or over the nerves proceeding from the cord and distributed to the parts at the sides of the spine. Such tenderness, however, by no

means indicates the pathological condition in question, as it is often met with in those enjoying perfect health. The treatment advised is cupping and counterirritation on each side of the spine, which may be beneficial in such diseases, no matter what part of the frame may be in a morbid state, by exciting a new and revealing impression on a very sensible portion of the cutaneous surface.

SPINAL MARROW, *Medulla spinalis*.

SPINAL NERVE, *Accessory of the Par va'gum* or *8th pair*, *Accessory nerve of Willis*, *Spin'al Accessory*, *Superior respiratory N.*, *Eleventh pair of encephalic nerves*, (F.) *Spin-eratio-trapézien*, *Trachélo-dorsal* (Ch.,) arises from the medulla spinalis, within the vertebral canal, between the anterior and posterior roots of the cervical nerves, at a greater or less distance from the cranium. The roots unite to form the nerve, which ascends into the cranium through the foramen magnum of the occipital bone, and issues by the foramen lacerum posterius, crossing the sternocleidomastoideus, to which it gives filaments, and losing itself entirely on the trapezius muscle. The pneumogastric and spinal accessory nerves together—*nervus vagus cum accessorio*—resemble the spinal nerves; the former, with its ganglion, being the posterior root; the latter the anterior.

The spinal accessory appears to be the nerve of voice—*vocal nerve*.

SPINAL NERVES, Vertebral nerves—s. *Prolongation*, *Medulla spinalis*.

SPINAL SYSTEM OF NERVES, see Nerves.

SPINALES COLLI MINORES, *Interspinales colli*—s. *et Transversales lumborum*, *Transversalis dors*.

SPINALIS CERVICIS, *Semispinalis colli*—s. *Colli*, *Semispinalis colli*.

SPI'NALIS DOR'SI, *Spina'lis dor'si ma'jor*, (F.) *Grand épineux du dos*. Winslow calls thus some fleshy fasciuli, which are situate on the lateral surfaces of the spinous processes; from the third dorsal vertebra to the first or second lumbar; and which form part of the transversospinalis of most authors. The same anatomist calls—*Spina'lis Dor'si mi'nor*, *Petit épineux du dos*—some small, fleshy fibres, situate on each side of the interspinal ligament. They are short, flat, and pass from one spinous process to another. Like the preceding, they form part of the transversospinalis. All these fleshy fasciuli strengthen the vertebral column.

SPINATI, *Interspinales colli*.

SPINATUS, Spinal.

SPINDLE TREE, *Eunonymus Americanus*.

SPINE, *Spina*, Vertebral column—s. *Curvature of the*, *Gibbositas*—s. *Hæmal*, *Sternum*—s. *Neural*, *Spinous process*.

SPINGEL, *Anethum*.

SPINIA'XOIDOOCIPITALIS, *Rectus capitis posticus major*—s. *Axoido-trachélio-atloïdien*. Obliquus inferior capitis.

SPINITIS, (*spina*, 'the spine,' and *itis*,) *Myelitis*.

SPINNICK, *Nanus*.

SPINODORSITIS, (*spina*, *dorsum*, 'the back,' and *itis*,) *Myelitis*.

SPINOLA, (from *spina*, 'the spine,') see *Hydrorachis*.

SPINOSUS, (from *spina*, 'the spine,') *Spinal*.

SPI'NOUS, *Spino'sus*, (F.) *Épineux*, (from *spina*, 'a thorn,') Having the shape of a spine or thorn.

SPINOUS PROCESSES or APOPH'YSES of the Vertebrae, *Acan'thæ*, *Sp'inae*, *Cynol'ophi*, *Neural Spines* of Owen, (F.) *Apophyses épineuses*, are situate at the posterior part of each vertebra, and afford attachment to the muscles, whose office it is to extend the spine. See Vertebra.

SPINTHEROPIA, (*σπινθηρ*, 'a spark, and *οπίς*, 'vision,') Confusio.

SPINULA, (dim. of *spina*, 'a thorn,') Pin.

SPIRACULA, (pl. of *spiraculum*,) (F.) *Spiracules*, (from *spira*, 'to breathe,') 'breathing holes.' Respiratory pores of the skin.

SPIRÆ CEREBRI, (*σπείρα*, *spira*, 'a fold, a winding,') Convolutions, cerebral.

SPIRÆA DENUATA, (from *spira*, 'a spire,') *S. ulmaria*.

SPIRÆA FILIPENDULA, *Filipendula*, *Saxifraga rubra*, *Dropwort*, (F.) *Filipendula*. Ord. Rosaceæ. Ser. Syst. Icosandria Pentagynia. The root of this plant is said to possess astringent and lithontriptic virtues.

SPIRÆA LOBATA, *S. Palma'ta*, *Queen of the Prairie*, *American Mead'owsweet*. An oil, resembling oleum gaultheriæ, is obtained from the roots and stems.

SPIRÆA OPULIFOLIA, *Opulus-leaved Spiræa*, *Nine Bark*, has similar properties.

SPIRÆA PALMATA, *S. lobata*.

SPIRÆA TOMENTOSA, *Hard'hack*, *Red mead'owsweet*, *Steel'plebush*, *Rosy bush*, *Whiteleaf*. The root of this indigenous species, *Spiræa* (Ph. U. S.), which is abundant in the northern States of the Union, is tonic and astringent; and is usually administered either in the form of extract or decoction, (*Spirææ*, $\frac{3}{4}$ ss; *Aque*, Oj.)

SPIRÆA TRIFOLIATA, *Gillenia trifoliata*.

SPIRÆA ULMARIA, *S. denudata*, *Ulmæria*, *U. palustris*, *Regina Præ'ti*, *Bar'ba ca'præ*, *Bar'bula capri'na*, *Mead'ow Sweet*, *Queen of the Mead'ows*, (F.) *Ulmæire*, *Spirée ulmaire*, *Reine des Prés*, *Vignette*, *Herbe aux Abeilles*. The leaves have been recommended as mild astringents and diuretics—the flowers as antispasmodics and diaphoretics.

SPIRAMENTA PULMONUM, 'airholes of the lungs,' (from *spirare*, 'to breathe,') see Pulmo.

SPIRAMENTUM ANIMÆ, 'airhole of the breath,' Pulmo.

SPIRAMINA PALPEBRARUM, 'pores of the eyelids,' *Lachrymal puncta*—s. Pulmonum, see Pulmo.

SPIRATIO, (*spirare*, *spiratum*, 'to breathe,') Respiration.

SPIRÉE ULMAIRE, *Spiræa ulmaria*.

SPIR'IT, *Spir'itus*, (*spirare*, 'to breathe, to exhale,') (F.) *Espirit*, *Alcoholat*. A name given to every liquid product of distillation. Spirits were formerly distinguished into—*inflammable*, *acid*, and *alkaline*; and consequently a number of substances were crowded together, which often resembled each other in no other property than in being volatile. The term is now confined to alcoholic liquors, of which the following are the chief:

Arrack'. Distilled from coarse palm sugar, named *jaggery*; fermented with the bark of the *Mind'sa leucophle'a*; also from rice and the fermented juice of the Palm; made in India. Its varieties are:—

1. *Mah'wah Arrack'*. Made in India from the flowers of the Madhuca tree, *Bas'sia butyra'cea*.

2. *Tuba*. Made from palm wine, in the Philippine Islands. See *Arack*.

Araka. Distilled by the Tartars, from koumis, fermented mares' milk.

Araki. Distilled from dates, by the Egyptians.

Arika. A variety of koumis, distilled from fermented mares' milk, in Tartary and Iceland.

Brand'y. Distilled from wine, figs, peaches, persimmon, apple, mulberries, and sometimes other fruits, in Europe, Asia, North and South America, wherever wine is made. The best brandy is that of Cognac; the next, that of Bordeaux and Rochelle. The varieties are—

1. *Aguardiente*, (S.) In Peru, the common brandy obtained from grapes is the *Aguardiente de Pisco*, so called, because shipped at the port of Pisco. Another kind, much dearer, and of excellent flavour, is made from Muscatel grapes, and is called *Aguardiente de Italia*. It is sometimes seen in the United States.

2. *Lau*. Made from rice, in Siam.

3. *Rakia*. Made in Dalmatia from the husks of grapes mixed with aromatics.

4. *Rossolio*. Made at Dantzic, from a compound of brandy, rossolis, and other plants.

5. *Troster*. Made on the Rhine, from the husks of grapes, fermented with barley and rye.

6. *Sekisakagvodka*. Made from the lees of wine and fruit at Scio.

Gene'va, *Holl'ands*, (*genièvre*, French for juniper.) Distilled from malted barley and rye, rectified on juniper berries, in Holland. Its variety is—

Gin. Made in England, from malted barley, rye, potatoes; rectified with turpentine.

Goldwasser. Distilled at Dantzic, from wheat, barley, and rye, rectified with aniseed, cinnamon, and other spices.

Kirschwasser. Distilled from the Malacherry, in Switzerland.

Maraschino, *Marasqui'no*. Distilled from the Macarska or Marasca cherry, at Zara, the capital of Dalmatia.

Rum, (supposed to be derived from the terminal syllable of the word *saccharum*: but this is questionable. Distilled, in the West Indies and South America, from canesugar and molasses, and in North America from maplesugar. Its variety is—

Sakee, *Saki*. A Japanese beverage, distilled from rice.

Stakkaia trava. Made in Kamtschatka, from a sweet grass.

Shonchoo. Distilled, in China, from the lees of mandarin, a wine made from boiled rice.

Whis'ky, (supposed to be derived from *usque*, the first two syllables of *usquebaugh*, the original name in Ireland,—itself from Irish *uisge* and *beatha*, 'life,' *Uisge beatha*, *Aqua vite*, 'Water of life.' (Distilled, in Scotland and Ireland, from malted and raw barley, rye, oats, and potatoes; and in the south of France, from sloes.) In Ireland it was called *builceann*, or madness of the head. The best Scotch whisky is *Glenlivet*, the best Irish, *Enniskow'en*.

Whisky, *Spir'itus Frumen'ti*, is official in the Pharmacopœia of the United States. It is the spirit obtained from fermented grain by distillation, and contains from 48 to 56 per cent. of absolute alcohol.

Y-ver-a. Distilled, in the Sandwich Islands, from the root of the Teeroot, baked, pounded, and fermented.

To these may be added an intoxicating liquor made by the Afghanistans, from ewes' milk; and that made in Kamtschatka, from a species of mushroom, named *muchumer*.

SPIRIT OF AMMONIA, *FETID*, *Spiritus ammoniæ fetidus*—s. of Ammonia, succinated, *Spiritus ammoniæ succinatus*—s. of Aniseed, *Spiritus anisi*.

SPIRIT, BATHING, FREEMAN'S, see Linimentum saponis compos.—s. Bathing, Jackson's, see Linimentum saponis compos.—s. Bezoardic of Bussius, see *Russii spiritus Bezoarticus*—s. Bone, *Liquor volatilis cornu cervi*.

SPIRIT OF BURRHUS, *Burhii spiritus matricialis*—s. of Cajeput, see *Cajeput oil*—s. of Camphor, *Spiritus Camphoræ*—s. of Caraway, *Spiritus carui*—s. of Chloroform, *Spiritus chloroformi*—s. of Colchicum, ammoniated, *Spiritus colchici ammoniatus*—s. of Cinnamon, *Spiritus cinnamomi*—s. of Ether, *Spiritus ætheris sulphurici*—s. of Ether, aromatic, *Spiritus ætheris aromaticus*.

SPIRIT, FUMING, OF BEGUIN, Ammoniae sulphuretum.—S. Fuming, of Boyle, Ammoniae sulphuretum.

SPIRIT OF HARTSHORN, Liquor volatilis cornu cervi.—S. of Horsradish, compound, Spiritus armoracae compositus.—S. of Juniper, compound, Spiritus juniperi compositus.—S. of Lavender, Spiritus lavandulae.—S. of Lavender, compound, Spiritus lavandulae compositus.—S. of Lemon, Spiritus Limonis.—S. of Mindererus, Liquor ammoniae acetatus.—S. of Myrica, Spiritus myricae.—S. of Nitre, sweet, Spiritus aetheris nitrici.—S. of Nitric ether, Spiritus aetheris nitrici.—S. of Pennyroyal, Spiritus pulegii.—S. of Peppermint, Spiritus menthae piperitae.—S. of Pimento, Spiritus pimentae.

SPIRIT, PROOF, Spiritus tenuior.—S. Pyroxylic, Spiritus Pyroxylicus.

SPIRIT OF ROSEMARY, Spiritus rosmarini.—S. of Salt, Muraticum acidum.—S. of Scourvygrass, golden, see Spiritus armoracae compositus.—S. of Spear-mint, Spiritus menthae viridis.—S. of Sulphuric ether, Spiritus aetheris sulphurici.—S. of Sulphuric ether, compound, Spiritus aetheris sulphurici compositus.—S. of Turpentine, Oleum terebinthinae rectificatum.

SPIRIT, VERDIGRIS, Aceticum acidum.

SPIRIT OF VITRIOL, Sulphuric acid.—S. of Wine, Spiritus rectificatus.—S. of Wine and camphor, Spiritus camphorae.

SPIRIT, WOOD, Spiritus Pyroxylicus.

SPIRITS, ANIMAL, Nervous fluid.

SPIRITUOUS, (from spiritus,) Alcoholic.

SPIRITUS, (spirare, 'to breathe,') Breath, Life, Respiration, Spirit.—S. Aetherus nitrosus, Spiritus aetheris nitrici.—S. Aetherus vitriolicus, Spiritus aetheris sulphurici.

SPIRITUS ÆTHERIS AROMATICUS, Æther Sulphuricus cum alcoho'le aromaticus, Elix'ir vitrioli dulcē, Sweet Elix'ir of Vitriol, Viga'n's Elix'ir of Vitriol, Aromatic Spirit of Æther. (Cinnam. cort. contus. ʒij, Cardam. sem. contus. ʒiiss, Piperis longi fruct. cont., Zingib. rad. concis. sing. ʒi, Spiritus aetheris sulphurici. Oj. Macerate, for 14 days, in a stopped glass vessel, and strain. Ph. L.) It is used as a stimulant in nervous affections. Dose, fʒss to fʒi.

SPIRITUS ÆTHERIS CHLORATI, Spiritus aetheris muriatici.—S. Aetheris compositus, Spiritus Aetheris sulphurici compositus.—S. Aetheris hydrochlorici, Spiritus Aetheris muriatici.

SPIRITUS ÆTHERIS MURIATICI, Sp. Æ. Hydrochlorici, Sp. Æ. chlorati, Sp. Æ. Muriatici, Sp. Muriatici aetheris, Sp. Salis dulcis, Æther Muriatici alcoholici seu chloratus alcoholici, Spirit of hydrochloric or muriatic ether, Dulcified marine ac'id. An old preparation, made by adding one part of Muriatic acid to three parts of Rectified spirit, and distilling. Its action is like that of the spiritus aetheris nitrici.

SPIRITUS ÆTHERIS NITRICI, (Ph. U. S. 1851.) Sp. Ætheris nitroci (Ph. U. S. 1863 and Ph. B. 1864) seu nitri dulcis seu Ætheris nitrosus, Æther nitricus alcoholatus, seu nitricæalcoholicus, Ac'idum Nitri dulcificatum, Sweet Spirit of Nitre, Nitre drops, Spirit of Nitric Ether, (F.) Æther azotens alcoholis, Liqueur anodine nitreuse. (Nitrate of Potassa, in coarse powder, ℥ij, Sulphuric acid, ℥iiss; Alcohol, O℥ss; Diluted Alcohol, Oj; Carbonate of Potassa, ʒi. Mix the nitrate of potassa and the alcohol in a large glass retort, and having gradually poured in the acid, digest with a gentle heat for two hours; then raise the heat, and distil a gallon. To the distilled liquor add the diluted alcohol and carbonate of potassa, and again distil a gallon.—Ph. U. S.) A modified form is given in the Ph. U. S. for 1863, s. g. 0.837, Ph. B. (1864), 0.843. In the latter it is distilled from a mixture of Sulphuric acid and Rectified spirit poured on Nitrite of

soda. It is refrigerant, diuretic, antispasmodic, and diaphoretic. Dose, gtt. xxv to fʒi. Specific gravity, 0.834.

SPIRITUS ÆTHERIS OLEOSUS, Spiritus Aetheris sulphurici compositus, Oleum æthereum.

SPIRITUS ÆTHERIS SULPHURICI, Sp. Ætheris, Æther Sulphuricus cum Alcoho'le, Li'quor Ætheris Sulphuricus, Spiritus Vitrioli dulcis, S. Ætheris vitriolici, Li'quor Sulphuricus alcoholatus, Spirit of Sulphuric Ether. (Æther. sulphuric. Oss; Sp. rect. Oj. Mix. Ph. L.) Stimulant, diaphoretic, diuretic, and antispasmodic. Dose, fʒss to fʒij.

SPIRITUS ÆTHERIS SULPHURICI COMPOSITUS, (Ph. U. S. 1842), S. ætheris vitriolici compositus, Æther hydricæ alcoholis, Æther sulfuricæ alcoholis, seu Ætheris compositus (Ph. U. S., 1863) seu Spiritus ætheris oleosus, Li'quor anodynus seu anodynus mineralis Hoffman'ni, Compound Spirit of Sulphuric Ether, Hoffman's Anodyne Li'quor, (F.) Esprit d'æther sulfurique, Liqueur anodine d'Hoffmann. (Æther. Oss; Alcohol, Oj; Æther. fʒij. Mix.) A stimulant and antispasmodic. Specific gravity, 0.816.

SPIRITUS ÆTHERIS VITRIOLICI COMPOSITUS, Spiritus aetheris sulphurici comp.

SPIRITUS AMMONIÆ, Al'kali ammoniacum spirituosum, Al'cohol ammoniacum, Sp. salis ammoniaci, Sp. salis ammoniaci vinosus seu dulcis seu dulcificatus, Ammoniacæ alcoholisatæ, Li'quor ammoniaci vinosus, L. ammoniacæ spirituosus, Lixivium ammoniacale vinosum, Solutio subcarbonatis ammoniaci spirituosæ, (F.) Esprit d'ammoniaque. (Ammon. muriat. in pulv. subtilissim., Calcis, aa ℥ij; Alcohol, fʒxx; Aquæ, Ovj. The water is employed to slake the lime; the muriate of ammonia is then well mixed with it, and by means of heat the ammonia is made to pass into the distilled water contained in a quart bottle. Ph. U. S.) Stimulant and antispasmodic. Dose, fʒss to fʒi.

SPIRITUS AMMONIÆ AROMATICUS, Al'cohol ammoniacum aromaticum, Spiritus salis volatilis oleosus seu volatilis aromaticus seu ammoniacæ compositus seu oleo'so'aromaticus Syll'vii, Sal volatilis oleosum Syll'vii, Tinctura aromatica ammoniacæ, Li'quor oleosus Syll'vii, L. Ammoniacæ aromatisatus seu ætheris seu oleo'so'aromaticus, Lixivium ammoniacale aromaticum, Aromaticæ ammoniacæ Al'cohol, Sal volatilis Drops. (Ammon. muriat. ʒv; Potass. carb. ʒviij; Cinnam. cont., Caryophyll. cont. aa ʒij; Limon. cort. ʒiv; Alcohol., Aquæ, aa, Ov. M. Distil Ovjss. In the Ph. U. S. for 1863, it is directed to be made of Ammon. carb. ʒi; Aq. ammon. fʒij; Ol. Limon. fʒiiss; Ol. myrist. ℥xl; Ol. Lavandul. ℥xv; Alcohol, Oiss; Aquæ, q. s. Dissolve the carbonate in the water of ammonia, mixed with fʒiv of water. Dissolve the oils in the alcohol; mix the two solutions, and add water sufficient to make the whole two pints.) Stimulant. Dose, fʒss to fʒi.

SPIRITUS AMMONIÆ COMPOSITUS, Spiritus aromaticus.

SPIRITUS AMMONIÆ FÆTIDUS, Sp. volatilis fætidus, Al'cohol ammoniacum fætidum, Tinctura assæfætidæ ammoniacæ, Fætid Spirit of Ammonia, Fit Drops. (Ammon. muriat. ʒx, Potass. carb. ʒxxv, Sp. rectificat., Aquæ, aa Oij, (Imp. meas.), Assæfætid. ʒv. Distil. Oij, (Imp. meas.) Ph. L.) Stimulant and antispasmodic. Dose, fʒss to ʒi.

SPIRITUS AMMONIÆ SUCCINATUS, Sp. Ammoniacæ cum Succino, Sp. Cornu cervi succinatus, Sp. Salis Ammoniaci buccinatus seu succinus, Tinctura ammoniacæ composita, Ammoniacum succinatum, A'qua Luleis, Li'quor ammoniaci succinici seu ex Ammoniacæ et Oleo Succini, L. ex citans, Ammo-

nium succinicum liquidum, *Succinated Spirit of Ammonia*, *Cammon Eau de Luce*. (Mastich. ʒij, Sp. rect. fʒix; Ol. Lavand. gtt. xiv; Ol. succin. gtt. iv; Lq. ammon. fʒx. Macerate the mastich in the alcohol, and pour off the clear tincture: add the rest and shake. Ph. L.) Stimulant and antispasmodic. Dose, gtt. x to fʒss or fʒi.

SPIRITUS ANIMALES, see Nervous fluid.

SPIRITUS ANISI, Sp. Anisi compositus, *Aqua anisi fortis*, A. Semivium Anisi composita, *Spirit of Aniseed*, (F.) *Esprit d'anis*. (Ol. Anis. fʒij; Alcohol. fort. fʒxv. Dissolve. Ph. U. S.) Carminative. Dose, fʒss to fʒss. In the compound spirit, angelica is usually an ingredient.

SPIRITUS ANTHOS, Spiritus rosmarini.

SPIRITUS ARMORACIÆ COMPOSITUS, Sp. Raphani compositus, *Aqua Raphani composita*, *Compound Spirit of Horse-radish*, (F.) *Esprit de raifort composé*. (Armorac. rad. recent. concis., Aurant. cort. exsic., sing. lbj, Myrist. nuc. contus. ʒss, Spirit. ten. cong., Aquæ, q. s. ad prohibendum empyreuma. Ph. L.) Stimulant and antiscorbutic. [?] Dose, fʒi to fʒiv.

The Gold'en Spirit of Scavvygrass—an empirical preparation—is formed by adding Gamboze, ʒviij, to one gallon of the Sp. armoraciæ compositus.

Greenough's Tincture for the teeth is formed of Amygd. amar. ʒij, Lig. Bresil., Cassiæ bacc. aa ʒiv, Irid. Florent. ʒij, Coccinell., Sul. acetosellæ, Alum. aa ʒj, Sp. vin. rect. Oij, Sp. armoraciæ comp. fʒss.

SPIRITUS BEGUINIS, Ammonia sulphuretum—s. Bellidis, see Osmitopsis asteriscoides—s. Cajuputi, see Cajuput oil.

SPIRITUS CAMPHORÆ, (Ph. U. S. 1863.) Sp. Camphoratus seu vinosus camphoratus, *Tinctura Camphoræ* (Ph. U. S. 1851), *Spirit of Wine and Camphor*, *Spirit of Camphor*, (F.) *Esprit de Camphre*, *Alcool camphré*, *Eau de Vie camphrée*. (Camphor. ʒiv, Alcohol. Oij. Dissolve.) Stimulant, anodyne and discutient. Used only externally.

SPIRITUS CARUI, Sp. Carui Carui, *Aqua Seminum Carui fortis* seu *Seminum Carui*, *Spirit of Caraway*, *Strong Carui Waters*, (F.) *Esprit de Carvi*. (Carui sem. cont. lbiss, Sp. tenuior. cong., Aquæ q. s. Distil a gallon. Ph. L.) Carminative. Dose, fʒi to fʒss.

SPIRITUS CHLOROFORMI, *Spirit of Chloroform*. (Chloroform. purif. ʒj, Alcohol. fort. fʒvj. Dissolve. Ph. U. S., 1863.) See Ether, chloric.

SPIRITUS CINNAMOMI, Sp. Lauri Cinnamomi, *Aqua Cinnamomi fortis* seu *Cinnam. spirituosus*, *Spirit of Cinamon*, *Strong Cinamon Water*, (F.) *Esprit de Cannelle*. (Ol. cinnam. fʒij, Alcohol. fort. fʒxv.) Stimulant and carminative. Dose, fʒi to fʒss.

SPIRITUS COLCHICI AMMONIATUS, *Tinctura Colchici Composita*, *Ammoniated Spirit of Colchicum*, *Compound Tincture of Colchicum*. (Sem. colchic. cont. ʒij, Sp. ammon. aromat. Oij. Ph. L.) Narcotic, cathartic, and diuretic. Used in gout, rheumatism, and dropsy. Dose, from gtt. xxx to fʒi.

SPIRITUS CORNI CERVI, *Liquor volatilis C. C.*—s. Cornu Cervi succinatus, *Spiritus ammoniæ succinatus*—s. Fortior, see Alcohol—s. Fumans Beguini, *Ammonia sulphuretum*—s. Genitalis, *Aura seminis*—s. Jamaicensis, Rum.

SPIRITUS JUNIPERI COMPOSITUS, *Aqua Juniperi composita*, *Compound Spirit of Juniper*, (F.) *Esprit de genévère composé*. (Ol. junip. ʒis, Ol. carui, Ol. fœnicul. aa ʒx, Alcohol dilut. cong. Solve. Ph. U. S.) Stimulant and diuretic. Dose, fʒi to fʒss.

SPIRITUS LAURI CINNAMOMI, *Spiritus Cinnamomi*.

SPIRITUS LAVANDULÆ, Sp. Lavan'dulæ spiritus seu *Lavand. sim'plex*, *Lavender Water*, (F.) *Es-*

prit de Lavande. (Lavand. recent. lbij, Alcohol. cong., Aquæ Oij. Distil a gallon. Ph. U. S.) Usually made by adding the oil to rectified spirit. Used as a perfume chiefly.

SPIRITUS LAVANDULÆ COMPOSITUS, *Tinctura Lavandulæ composita*, *Red Hart's horn*, *Lavender drops*, *Compound Spirit of Lavender*, (F.) *Esprit de Lavande composé*. (Sp. Lavand. Oij, Sp. rosamarin. Oj, Cinnam. contus. ʒj, Myristic. cont. ʒss, Caryophyll. cont. ʒij, Santali concis. ʒij. Macerate for 14 days, and filter through paper. Ph. U. S.) A modified form is given in the Ph. U. S. for 1863. Stimulant. Dose, gtt. xl to fʒij.

SPIRITUS LETHALIS, Carbonic acid.

SPIRITUS LIMO-NIS, *Spirit of Lemon*, *Essence of Lemon*, (Ol. Limon. fʒij, Cort. Lim. recente rasp. ʒj, Alcohol. fort. Oij. Dissolve the oil in the alcohol; add the lemon peel; macerate for 24 hours, and filter through paper. Ph. U. S., 1863.) To give odour and flavour.

SPIRITUS LUMBRICORUM, *Liquor volatilis cornu cervi*.

SPIRITUS MENTHÆ PIPERITÆ, *Tinctura O'lei Menthæ Piperitæ*, Sp. Menthæ piperitidis, *Aqua Menthæ piperitidis spirituosus*, *Spirit of Peppermint*, *Tincture of Oil of Peppermint*, *Essence of Mint*, (F.) *Esprit de Menthe poivrée*. (Olei menth. pip. ʒviss, Alcohol. fort. Oivss, Aquæ q. s. Distil a gallon;—or, Ol. menth. piperit. fʒij, Alcohol. Oj. Dissolve. Ph. U. S., 1851.)

The Spiritus Menthæ Piperitæ (Ph. U. S., 1863) is a slightly modified form. (Ol. menth. pip. fʒij, Menthæ piperit. in pulv. crass. gr. exx, Alcohol. fort. ʒxv. Dissolve the foil in the alcohol; add the peppermint; macerate for 24 hours, and filter. Dose, gtt. v—xx, as a carminative.

SPIRITUS MENTHÆ SATIVÆ, *Spiritus menthæ viridis*.

SPIRITUS MENTHÆ VIRIDIS, *Tinctura O'lei Menthæ viridis*, *Spiritus menthæ sativæ*, *Aqua Menthæ vulgaris spirituosus*, *Spirit of Spearmint*, *Tincture of Oil of Spearmint*, *Essence of Spearmint*, (F.) *Esprit de Menthe vert*.

The Spiritus Menthæ Viridis (Ph. U. S., 1863) is made like the Spiritus menthæ piperitæ, and has the same properties and dose.

SPIRITUS MILLEPEDARUM, *Liquor cornu cervi*—s. Mindereri, *Liquor Ammonia acetatis*—s. Muriatricus æthereus, *Spiritus Ætheris muratici*.

SPIRITUS MYRICÆ, *Spirit of Myrica*, *Bay rum*. A spirit obtained by distilling rum with the leaves of *Myrica acris*, *Engenia* seu *Myrtus acris*, *M. Caryophyllata*, *Bayberry Tree*. Order, Myrtaceæ, of the West Indies. Used externally like Cologne water, and in cases where whisky would be applied.

SPIRITUS MYRISTICÆ, Sp. Myristicæ Moschatæ seu *Nucis Moschatæ*, *Aqua Nucis Moschatæ* seu *Nephriticæ*, *Nutmeg Water*, (F.) *Esprit de Muscade*. (Myrist. cont. ʒij, Alcohol. dil. cong., Aquæ Oj. Distil a gallon.) Cordial and carminative. Dose, fʒi to fʒij.

SPIRITUS MYRTI PIMENTÆ, *Spiritus pimentæ*—s. Nitri acidus, Nitric acid—s. Nitri dulcis, *Spiritus ætheris nitrici*—s. Nitri duplex, *Nitric acid*—s. Nitri fumans, Nitric acid—s. Nitri Glauberi, Nitric acid—s. Nitri simplex, see Nitric acid—s. Nitri vulgaris, see Nitric acid—s. Nucis moschatæ, *Spiritus myristicæ*—s. Oleosaromaticus Sylvii, *Spiritus ammoniæ aromaticus*—s. Ophthalmicus Mindereri, *Liquor ammoniæ acetatis*.

SPIRITUS PIMENTÆ, Sp. Myrti Pimentæ seu *Piment*, *Aqua Mirabilis*, *Spirit of Pimento*. (Ol. Piment. fʒij, Alcohol. dilut. cong. Solve. Ph. U. S.) Cordial and carminative. Dose, fʒi, and more.

SPIRITUS PULEGI, *Aqua Pulegii spirituosus*, *Spirit of Pennyroyal*, *Spiritus Pen'nyroyal Water*, (F.) *Esprit de Poultier*. (Olei pulegii ʒvij, Sp. rect. Oivss, Aquæ q. s. Distil a gallon.

Ph. L.) Same as spirit of spearmint in properties and dose.

SPIRITUS PYROACETICUS, Acetone.

SPIRITES PYROXYLICUS, *Pyroxylic spirit*, *Wood spirit*, *Wood alcohol*, *Pyroxylic alcohol*, *Methyl alcohol*, *Hydrated oxide of methyle*, *Bihydrate of Methylene*, (F.) *Espirit pyroxylique*, *Espirit de bois*. This substance is by some supposed to have been the article given in phthisis by Dr. Hastings, under the names *Naphtha* and *Wood Naphtha*. It was probably, however, the *Pyrolicetic spirit*. See Acetone. When wood is subjected to destructive distillation, about one per cent. of an inflammable volatile liquid, along with other products, passes over, which, when separated and purified, constitutes pyroxylic spirit. It is given in the same cases and doses as Acetone.

SPIRITUS RAPHANI COMPOSITUS, *Spiritus aromaticæ compositus*.

SPIRITUS RECTIFICATUS, *Sp. Vini rectificatus seu vinosus rectificatus*, *Enostagma*, *Enelsum*, *Pyraenus*, *Alcohol*, *A. Vini*, *Spirit of Wine*, (F.) *Espirit de Vin rectifié*. Rectified spirit is much used in Pharmacy to dissolve resinous and other substances. It is seldom exhibited internally, although it exists in the diluted state in all vinous and spirituous liquors. Externally, it is employed as a discutient.—S. g., according to the British Pharmacopœia, 0·838; according to that of Dublin, 0·840; of Edinburgh and United States, 0·835 (see Alcohol), and of Paris, 0·819 to 0·827.

SPIRITUS RECTOR, *Aroma*—s. *Roriferus*, *Gaz sanguinis*.

SPIRITUS ROSMARINI, *Sp. Rosmarini officinalis*, *Sp. Anthos*, *Aqua Hungarica seu Regiæ Hungariæ*, *Spir'it of Rosemary*, *Hungary Water*, (F.) *Espirit de Romarin*. (*Ol. rosmarini* f 3iv, *Alcohol* cong. Solve. Ph. U. S.) Stimulant. Dose, f 3j to f 3iv.

SPIRITUS SALIS ACIDUS seu *FUMANS*, *Muriaticum acidum*—s. *Salis ammoniaci*, *Spiritus ammoniacæ*—s. *Salis ammoniaci aqueosus*, *Liquor ammoniacæ*—s. *Salis ammoniaci lactescens seu lacteus*, *Spiritus ammoniacæ succinatus*—s. *Salis ammoniaci sulphuratus*, *Ammoniacæ sulphuretum*—s. *Salis dulcis*, *Spiritus ætheris muriatici*—s. *Salis Glauberi*, *Muriaticum acidum*—s. *Salis marini*, *Muriaticum acidum*—s. *Salis marini coagulatus*, *Potasse murias*—s. *Salis volatilil oleosus*, *Spiritus ammoniacæ aromaticus*—s. *Sulphuricoæthereus camphoratus*, *Tinctura æthereæ composita*—s. *Sulphuricoæthereus martialis*, *Tinctura seu Alcohol sulphuricoæthereus ferri*—s. *Sulphuris*, *Sulphuric acid*, *Sulphurous acid*—s. *Sulphuris per campanam*, *Sulphurous acid*—s. *Sulphuris volatilil*, *Ammoniacæ sulphuretum*—s. *Sulphurous volatilil*, *Sulphurous acid*.

SPIRITUS TENUIOR, *Alcohol dilutum*, *Spir'itus vinosus tenuior*, *Proof Spirit*, (F.) *Espirit de vin délayé*. (Generally made for pharmaceutical purposes by mixing equal parts of *rectified spirit* and *distilled water*.) S. g. Ph. B., 0·920; Edinb., 0·935; U. S., 0·941. It is employed in the same cases as those in which alcohol is used:—chiefly, as a solvent of vegetable matters, in the form of tinctures.

SPIRITUS VENERIS, *Aceticum acidum*—s. *Vini Gallici*, *Brandy*—s. *Vini rectificatus*, *Spiritus rectificatus*—s. *Vinosus camphoratus*, *Spiritus camphoræ*—s. *Vinosus tenuior*, *Spiritus tenuior*—s. *Vitalis*, *Nervous fluid*—s. *Vitrioli*, *Sulphuric acid*—s. *Vitrioli acidus Vogleri*, *Sulphuricum acidum dilutum*—s. *Vitrioli coagulatus Mynsichti*, *Potasse sulphas*—s. *Vitrioli dulcis*, *Spiritus ætheris sulphurici*—s. *Vitrioli phlogisticatus*, *Sulphurous acid*—s. *Volatilil aromaticus*, *Spiritus ammoniacæ aromaticus*—s. *Volatilil foetidus*, *Spiritus ammoniacæ foetidus*.

SPIROCOLON, *Spyrocolon*, *Or'chida*, *Franzo*. A syphiloid affection, first known in Greece, during the war of freedom from 1820 to 1825.

SPIROID CANAL, (σπείρα, 'a winding,' and εὐδω, 'resemblance.') *Aquæductus Fallopii*.

SPIROMETER, *Pulmonometer*, (*spiro*, 'I breathe,' and *μετρον*, 'a measure:') properly *Anapnometer*, *Pneumeter*, *Pneusmoneter*, *Pneumometer*, *Pneumatometer*, *Pneumonometer*. Any instrument for measuring the quantity of air concerned in respiration, and consequently the capacity of the lungs. This mode of measurement has been called *Spirometry* and *Pneumetry*, as it was formerly called *Pulmonetry*. The term spirometer is now usually applied to a kind of gasometer introduced by Dr. Hutchinson to estimate the quantity of air contained in, or, rather, expelled from the lungs under different circumstances.

SPIROPTERA HOMINIS, (σπείρα, 'a spire,' and *πτερον*, 'a wing,') see Worms.

SPISSANTIA, (*epissare*, 'to thicken.') Incrassantia.

SPIT, ([Sax.] γρætan, γριτtan, 'to spit.') Sputum.

SPITAL, Hospital.

SPITAL, MINERAL WATERS OF. The village of Spital is situate on the south side of the Tweed, near Tweedmouth. The springs there contain chloride of calcium, sulphate of soda, sulphate and carbonate of lime. They act as saline cathartics.

SPITHAMA, (σπίδαμη) *Do'drans*. A measure of twelve fingers' breadth, or about nine inches.

SPITTING, (see Spit.) Exspuition.

SPITTLE, Hospital; Saliva, ([Sax.] rparl,)—s. Sweet, see Salivation.

SPLANCHNA, (σπλαγχνα, pl. of σπλᾱγχμα,) Entails.

SPLANCHNECTOPIA, *Splanchnodistasis*, *Situs perversus seu anomalous intestinum*, (*splanchna*, εκ, 'out of,' and *τοπος*, 'place,') Misplacement or ectopia of the intestines.

SPLANCHNELMINTHA, (*splanchna*, and *ἔλμης*, gen. *ἑλμινθος*, 'a worm,') Worms.

SPLANCHNEURYSMA, (*splanchna*, and *ευρυς*, 'I dilate,') Morbid dilatation of the intestines.

SPLANCHNIC, *Splanchnicus*, (σπλαγχνικός.) *Visceral*, (F.) *Splanchnique*, (from *splanchna*.) Relating or belonging to the viscera.

SPLANCHNIC CAVITIES are the three great cavities of the body;—those of the cranium, chest, and abdomen.

SPLANCHNIC NERVES belong to the great sympathetic, and are two in number on each side. They are distinguished into:—1. *The great splanchnic nerve*, (F.) *Grand splanchnique*, (Ch.) Its roots—four or five in number—come from the 6th, 7th, 8th, 9th, and sometimes the 10th, thoracic ganglia. They descend inwards, on the sides of the vertebral column, beneath the pleura, and unite into a single trunk, which enters the abdomen through a separation of the fleshy fibres of the pillars of the diaphragm; passes behind the stomach, and divides into several branches, which pass into the semilunar ganglion. 2. *The lesser splanchnic nerve*, *Re'nal nerve*, (F.) *Splanchnique accessoire* of Walther, *Petit urénal* (Ch.), is composed of two distinct branches, proceeding from the 10th and 11th thoracic ganglia, which unite on the 12th dorsal vertebra into a cord that pierces the diaphragm singly; enters the abdomen, and divides into two branches, one of which anastomoses with the greater splanchnic nerve, whilst the other sends its filaments to the renal and solar plexuses.

SPLANCHNICA. Medicines adapted to diseases of the bowels. Also, the 2d order of the class

Cœliaca of Good; defined,—diseases affecting the collatitious viscera, and characterized by disquiet or diseased action in the organs auxiliary to the digestive process, without primary inflammation.

SPLANCHNODIASTASIS, (*splanchna*, and *διαστασις*, 'displacement.') Splanchnectopia.

SPLANCHNODYNE, (*splanchna*, and *ὄδυνη*, 'pain.') Pain in the bowels.

SPLANCHNOGRAPHIA, Splanchnography.

SPLANCHNOGRAPHY, (*Splanchnographia*, (*splanchna*, and *γραφία*, 'a description.') The part of anatomy which treats of the viscera.

SPLANCHNOLITHIASIS, (*splanchna*, and *λίθος*, 'a stone.') The formation of a calculous concretion in any viscus.

SPLANCHNOLOGY, (*Splanchnologia*, (*splanchna*, and *λογία*, 'a discourse.') A treatise on the viscera; also, Splanchnography.

SPLANCHNON, (*σπλᾶγχνον*,) sing. of Splanchna.

SPLANCHNOPATHIA, (*splanchna*, and *πάθος*, 'suffering.') A disease of the intestines.

SPLANCHNOSCLEROSIS, (*splanchnon*, and *σκληρσις*, 'hard.') Induration of a viscus.

SPLANCHNOTOMY, (*Splanchnotomia*, (*Splanchnotomē*, (*splanchna*, and *τέμνω*, 'I cut.') Dissection or anatomy of the viscera.

SPLATFOOT, see Kyllosis.

SPLAYFOOT, (from *splay*, 'spread.') See Kyllosis.

SPLEEN, (*σπλήν*,) *Lien*, *He'par sinistrum seu adulteri'mum*, *Fo'mes ventriculi*, *Milt.* (Sc.) *Melt.* (F.) *Rate*. A soft, spongy, parenchymatous organ, of a more or less deep violet red, situate deeply in the left hypochondrium, below the diaphragm, above the colon, between the great tuberosity of the stomach and the cartilages of the false ribs, and above and anterior to the kidney. Its ordinary length is $4\frac{1}{2}$ inches; its thickness 2 $\frac{1}{2}$; and its weight 8 ounces. Its parenchyma, which is bathed in blood, contains a multitude of grayish, soft, and semi-transparent granulations—*splenic corpuscles* or *Malpighian bodies* of the spleen. It is covered by a serous membrane, furnished by the peritoneum; and by a proper coat, of a fibrous nature, which is intimately united with it, and transmits into its interior delicate, solid, and very elastic prolongations—*trabeculæ*. At its inner edge is a fissure—*Hilus seu Porta lienis*, by which the vessels and nerves enter the organ. The functions of the spleen are unknown. The best opinion is, that it appertains to hæmatisis, and acts, also, as a diverticulum to the vascular system.

SPLEEN, Hypochondriasis—s. Malpighian, bodies of the, see Spleen—s. Supernumerary, Lienculus.

SPLEENWORT, *Asplenium*, A. *filix femina*, *Asplenium scolopendrium*—s. Bush, *Comptonia asplenifolia*.

SPLEN, gen. *Sple'nis*, (*σπλήν*,) Spleen.

SPLENALGIA, *Neuralgia lienis*, *Sple'nicæ*, *Splenodyn'ia*, (*splen*, and *αλγος*, 'pain.') Pain in the spleen.

SPLENALGIA BENGALENSIS, see Cachexia, splenic—s. *phlegmonodea*, *Splenitis*—s. Subinflammatoria chronica, *Splenoncus*.

SPLENATROPHIA, *Atroph'ia Lienis*, (*splen*, and *atrophia*, 'atrophy.') Wasting or atrophy of the spleen.

SPLENAUXE, (*splen*, and *αὖξη*, 'increase.') *Splenoncus*.

SPLENECTAMA, gen. *Splenectam'atis*, *Splenoparectama*.

SPLENECTASIS, (*splen*, and *εκτασις*, 'dilatation.') *Splenoparectama*.

SPLENECTOMIA, *Splenectomē*, (*splen*, *εκ*, 'out of,' and *τέμνω*, 'I cut.') Extirpation of the spleen.

SPLENECTOPIA, *Dislocatio seu Ectop'ia lienis seu sple'nis*, *Anomospленотопie*, (*splen*, and

εκτοπος, (*εκ*, and *τοπος*,) 'out of place.') Dislocation of the spleen. Not a common occurrence.

SPLENEMPHRAX'IS, (*Splenemphras'ia*, (*splen*, and *εμφρασις*, 'I obstruct.') Obstruction of the spleen. See *Splenoncus*.

SPLENES, (pl. of *splen*,) *Hypochondriasis*.

SPLENETICUS, *Sple'ticus*, *Splenit'icus*, *Lienot'icus*, (F.) *Rateleux*. That which relates to the spleen: one labouring under diseased spleen. A medicine adapted for diseases of the spleen.

SPLENIC, *Sple'nicus*, *Sple'nus*, (F.) *Splénique*, (from *splen*.) That which relates to the spleen.

SPLENIC ARTERY. It arises from the cœliac; and passes from right to left, forming many windings along the upper edge of the pancreas, which lodges it in a special groove. It reaches the fissure of the spleen, and divides into several branches that enter the organ. In its course it furnishes the pancreatic, left gastro-epiploic, and the *vasa brevia*.

SPLENIC CACHEXIA, see Cachexia splenica—s. *Corpuscles*, see Spleen.

SPLENIC PLEX'US is a nervous network, which proceeds from the cœliac plexus, and accompanies the splenic artery, sending secondary plexuses to each division.

SPLENIC VEIN, *Vena sple'nicæ*, *V. lienalis*, arises from the spleen, and accompanies the splenic artery; proceeding from left to right, to unite with the superior mesenteric vein, and to form, with it, the abdominal vena porta. In its course, it receives veins corresponding to the *vasa brevia*, the right and left gastro-epiploic, the duodenal, pancreatic, coronaria ventriculi, and inferior mesenteric veins.

SPLENICA, *Splenalgia*. Also, medicines which affect the spleen.

SPLENICUS, (*σπληνικός*,) *Spleteticus*, *Splenic*.

SPLENIFICATION, (*splen*, and *σίο*, 'to become,') *Splenization*.

SPLENIOLA, (dim. of *splen*,) *Compress*.

SPLÉNIQUE, *Splenic*.

SPLENIS TUMOR, 'tumor of the spleen,' *Ague cake*.

SPLÉNISATION, *Splenization*.

SPLENISCUS, (*σπληνικός*,) *Compress*.

SPLENISIS, (from *splen*,) *Splenization*.

SPLENITICUS, *Splenic*.

SPLENITIS, (*splen*, and *itis*,) *Inflammation of the spleen*. *Empres'ma splenitis*, *Inflammatio lienis*, *I. sple'nis*, *Splenal'gia phlegmonodea*, *Lienit'is*, (F.) *Splénite*, *Inflammation de la Rate*. A rare disease, characterized by local pain, swelling, and heat, which sometimes terminates by suppuration. It generally affects the capsule—*Epi'splenit'is*, *Perisplenit'is*—and must be treated like other inflammations.

SPLENIUM, (*σπλήνιον*,) *Compress*—s. *Corporis callosi*, see *Corpus callosum*—s. *Cruciatum*, *Croix de Malte*.

SPLENIUS, *S. Cap'itis seu Col'li*, (F.) *Cervico-dorso-mastoidien et dorso-trachélien*, *Cervico-mastoidien* (Ch.), *Posterior mastoid muscle*. So called from its resemblance to the spleen of certain animals. A muscle situate at the posterior part of the neck, and upper part of the back. It is much broader above than below, where it terminates in a point. It is attached, by its inner edge, to the spinous processes of the first five dorsal vertebrae; to that of the 7th cervical, and to the inferior part of the posterior cervical ligament. By its upper extremity, it is attached to the transverse processes of the first two cervical vertebrae, to the mastoid process, and to the posterior surface of the occipital bone. Some have considered this muscle to be formed of two parts, which they have called *Splenius colli*, and *Splenius cap'itis*. The splenius extends the head, inclines it, and

rotates it. If the splēnii of each side act together, they extend the head.

SPLÉNIZ'ATION, *Spleniza'tio*, *Splenifica'tion*, *Splenifica'tio*, (F.) *Splénisation*, (from *splen*.) A term applied to the state of the lung in the first or second stage of pneumonia, in which its tissue resembles that of the spleen. To a conversion of the lung into a spleen-like substance in the yellow fever of Gibraltar, the term *splénistis*, (F.) *Splénis* was given by Louis and others.

SPLENOCE'LE, *Hernia liénal'is*, (*splen*, and *κῆλη*, 'a tumour.') Hernia formed by the spleen.

SPLENODES, (*splen*, and *odos*.) Splenoid.

SPLENODYNIA, (*splen*, and *ὄδυν*, 'pain,') Splenalgia.

SPLENOG'RAPHY, *Splenograph'ia*, (*splen*, and *γραφη*, 'a description.') A description of the spleen.

SPLENOHÆ'MIA, (F.) *Splénohémie*, (*splen*, and *αἷμα*, 'blood.') Congestion or hyperæmia of the spleen, a common phenomenon in many diseases, especially in typhoid and intermittent fevers.

SPLÉNOID, *Splēnoï'des*, *Splēno'des*, (*σπληνοειδής*, from *splen*, and *εἶδος*, 'resemblance,') Spleenlike. Having the appearance of the spleen.

SPLENOLOG'Y, *Splenolog'ia*, (*splen*, and *λογος*, 'a discourse.') A treatise on the spleen.

SPLENOMALACIA, *Lienomala'cia*, (*splen*, and *μαλακος*, 'soft,') Morbid softening of the spleen.

SPLENONCIE, Splenoncus.

SPLENON'CUS, *Megalo'sple'nia*, *Hypertroph'ia seu Supernutrit'io sple'nis seu lié'nis*, *Intumescent'ia seu Inflac'tus seu Physco'nia lié'nis*, *Splenal'gia subinflammato'ria chron'ica*, *Splenemphrax'is*, *Splenau'z*, *Hypersplenotroph'ia*, *Li'en in'gens*, (F.) *Splénioncie*, *Hypertrophie de la Rate*, (*splen*, and *γκως*, 'a tumour,') Tumefaction of the spleen. See *Aque cake*.

SPLENOPAREC'TAMA, gen. *Sple'noparec-tam'atis*, *Splēnec'tama*, *Splenoparec'tasis*, *Splēnec'tusis*, (*splen*, and *παρεκταμα*, 'excessive volume,') Excessive enlargement of the spleen.

SPLENOPARECTASIS, Splenoparectama.

SPLENOPHRAXIA, (*splen*, and *φρασσα*, 'to shut up,') Splenemphraxia.

SPLENORRHAG'IA, (*splen*, and *ῥαγη*, 'a bursting forth,') Hemorrhage from the spleen.

SPLENOSCIR'RHUS, (*splen*, and *σκιρῆς*, 'hardness,') Scirrhus or cancer of the spleen.

SPLENOT'OMY, *Splenotom'ia*, *Splenot'omē*, (*splen*, and *τεμνω*, 'I cut,') Dissection of the spleen.

SPLINT, ([D.] Splinter, from *splijten*, 'to split,') *Hastel'ia*, *Ass'ula*, *Asser'culum*, *Pla'gula*, *Nar'theax*, *Fer'ula*, (F.) *Atelle*, *Éclisse*. A flexible and resisting lamina of wood, used in the treatment of fractures, to keep the fragments in apposition, and prevent their displacement. Splints are made of iron, wood, bark of trees, leather, gutta percha, pasteboard, &c. They are generally padded, and are fixed by a roller or by tapes, according to circumstances.

SPLINTER, (F.) *Esquille*, (same etymon.) This term is sometimes applied to a small portion of wood, which occasionally enters the skin, and gives occasion to much irritation, until it is removed, — *Acul'cus lig'neus*, (F.) *Écharde*.

It is also applied to fragments that separate from a fractured or diseased bone.

SPLIT'CLOTH, *Lin'teum scis'sum*. A bandage for the head, consisting of a central part, and six or eight tails. The single split cloth has four tails. It is regarded as the most convenient bandage for the head, face, and jaws.

SPLITROCK, *Heuchera cortusa*.

SPOAT, (like *spit*.) Saliva.

SPODIUM GRÆCORUM (*σποδιον*, 'dross,') *Albium græcum*.

SPOLIARUM, (*spolio*, 'to deprive of covering,') *Apodyterium*.

SPO'LIATIVE, *Spoliat'iva*, '*spoliare*, *spoliatum*, 'to take away,') (F.) *Spoliatif*.

Phlebotom'ia spoliati'va, (F.) *Saignée spoliative*. Bloodletting, used for the purpose of diminishing the mass of blood.

SPOLIATORIUM, (*spolio*, 'to deprive of covering,') *Apodyterium*.

SPONDLES, (from *spondylus*.) Vertebrae.

SPONDYLAL'GIA, *Spondylodyn'ia*, (*spondylus*, and *αλγος*, 'pain,') Pain in the vertebrae. *Cacorrhachitis*. See *Vertebral Disease*.

SPONDYLARTHITIS, (*spondylus*, *arthron*, 'a joint, and *itis*.) Gouty or rheumatic inflammation of the spinal column.

SPONDYLARTHROCACE, (*spondylus*, *arthron*, 'a joint, and *κακος*, 'evil,') Vertebral disease.

SPONDYLI, (*σπονδυλοι*.) Vertebrae.

SPONDYLISTHE'SIS, (*spondylus*, and *ολισθησις*, 'a slipping or sliding,') Dislocation of the vertebrae, as of the last lumbar vertebrae forwards on the sacrum.

SPONDYLITIS, *Inflammat'io ver'tebrae*, (*spondylus*, and *itis*.) Inflammation of the vertebrae.

SPONDYLIUM, (*σπονδυλιον*.) *Heraclium spondylium*.

SPONDYLOCACE, (*spondylus*, and *κακος*, 'evil,') Vertebral disease.

SPONDYLODYNIA, (*spondylus*, and *ὄδυν*, 'pain,') Spondylalgia.

SPONDYLOPATHIA, (*spondylus*, and *παθος*, 'a disease,') A disease of the vertebral column.

SPONDYLOPYOSIS, (*spondylus*, and *πυωσις*, 'suppuration,') Vertebral disease.

SPOND'YLUS, (*σπονδυλος*.) A vertebra. Also, the vertex.

SPONGE, *Spongia*—s. *Bastard*, *Alcyonium*—s. Prepared, see *Spongia*—s. *Tent*, see *Spongia*.

SPONGI, (from *σπγγος*.) The tumefied glands of the neck.

SPONG'GIA, (*σπγγια*.) *S. officina'lis seu mari'na*, *Spon'gus*, *Sphon'gus*, *Sponge*, (F.) *Éponge*. An organized marine substance, of the *Class* *Porifera*, met with in many seas, but especially in the Mediterranean and Arabian. It is supple, elastic, porous, and capable of absorbing fluids in which it may be placed and of thus increasing in size. It is employed in surgery for cleansing or washing; for absorbing acrid discharges from ulcers; suppressing hemorrhage when the mouth of the vessel can be compressed by it, and for forming tents for dilating wounds; in which case, the sponge is immersed in melting wax, and cooled before being used. It is then called *Spongia cera'ta*, *Waxed*, *Cerated* or *Prepared Sponge* or *Sponge Tent*.

SPONGIA ASSA, *S. Usta*—s. *Cerata*, see *Spongia* —s. *Cynosbati*, *Bedegar*.

SPONGIA US'TA, *S. As'sea*, *Car'bo spon'gie*, *Burnt Sponge*, contains carbonate and phosphate of lime, carbonate of soda, charcoal, and iodine. It is used in bronchocoele, scrofulous affections, and herpetic eruptions; and its good effects seem to be dependent upon iodine. (It is made by cutting sponge into pieces, and heating it, that any extraneous matter may be separated; burning it in a close iron vessel until it becomes black and friable, and, lastly, rubbing it into very fine powder.—Ph. U.S.) Dose. ʒi to ʒiij.

SPONGIÆ LA'PIS. A name given to small friable stones found in sponge. They were formerly esteemed lithontriptic.

SPONGIEUX, see *Spongoid*.

SPONGIFORM, (*spongia*, and *forma*, 'form,') *Spongoid*.

SPONGIODES, *Spongoid*.

SPONGIOLE (F.), *Spongi'olum*, *Spongi'ola*, *Spongelet*, (from *σπγγιον*, 'a small sponge,') The soft succulent extremity of the fibrils or true roots

of a plant, by which it absorbs or sucks up fluid. Similar spongioles have been supposed to exist at the terminations of the absorbents in animals.

SPON'GION, (σπογγιον.) Ancient name of an epithem made of goats' dung: and also of a malagma, which was considered capable of absorbing the serum in dropsies, like a sponge.

SPON'GIOPILIN, *Spon'giopile*, (*spongia*, 'sponge,' and *pilus*, 'a hair.') A fabric, composed of sponge and wool, felted together in three layers, and coated, on one of its surfaces, with caoutchouc, which renders it impermeable. It is of great service as a cataplasm, fomentation, vehicle for lotions, liniments, &c.

SPONGIOSUM (OS), Ethmoid bone.

SPONGIOSUS, Spongioid.

SPONGODES, (*spongia*, and *odes*.) Spongioid.

SPONGOID, *Spongoi'des*, *Spongo'des*, (F.) *Spongiöle*, (σπογγιοειδής, σπογγώδης, from *spongia*, and *εἶδος*, 'resemblance.') Resembling sponge. Of the nature of sponge, *spongy*, *spongio'sus*, (F.) *Spongioux*.

SPONGOID INFLAMMATION, *Hæmatodes fungus*.

SPONGOID TISSUE, (F.) *Tissu Spongiöde*. An adventitious spongy tissue found in cases of rickets.

SPONGOS, (σπογγος,) Tonsil.

SPONGY, Spongioid—s. Tissue, see Bone.

SPONSA SOLIS, 'spouse of the sun,' *Calendula officinalis*.

SPONTA'NEOUS, *Sponta'neus*, (F.) *Spontanée*, (from *sponte*, 'voluntarily.') That which occurs of itself, or without any manifest external cause. Evacuations are termed *spontaneous*, which are not produced by medicine. *Sponta'neous las'situde* is that which is not the result of fatigue. Diseases are also *spontaneous* which supervene without apparent cause. *Sponta'neous Amputation* is the separation of a limb from the fœtus in utero.

SPONTANEOUS EVOLUTION, see Evolution—s. Version, Evolution, spontaneous.

SPONTÉPARITÉ, (*sponte*, 'voluntarily,' and *pario*, 'to bring forth,') Generation, spontaneous.

SPOON, (from its hollowness,) Umbilicus.

SPOONWOOD, *Kalmia latifolia*, *Tilia Americana*.

SPORAD'IC, *Sporad'icus*, (F.) *Sporadique*, (σποράδικος, 'scattered,' from σπείρω, 'I sow.') An epithet for diseases, *Spor'ades*, *Morbi sporad'ici*, *disper'si* seu *spar'si*, (F.) *Maladies sporadiques*, which supervene indifferently in every season and situation, from accidental causes, and independently of any epidemic or contagious influence.

SPORE, *Spor'ule*, *Spo'rum*, *Spor'ulum*, *Sporus*, (σπορος, from σπείρω, 'I sow.') The reproductive body in cryptogamous plants, which is analogous to the seed of phanerogamous plants.

SPORID'IUM, (from *sporus*.) The reproductive body in algaceous plants, which resembles the spore of other plants.

SPORULE, Spore.

SPORUM, Spore.

SPORUS, Spore.

SPOT OF SÖMMERING, see Foramen centrale, &c.

SPOTS, CEREBRAL, see *Tache cérébrale*—s. Germinal, see Molecule—s. Meningitic, see *Tache cérébrale*.

SPOTS, RED, (F.) *Taches rouges*, *Rose spots*, *Rose'ole* seu *Pap'ule typh'o'se*. Small red spots which appear in typhoid fever, from the 8th to the 15th day of the disease. They are most frequently seen on the lower and middle portions of the chest, and upper part of the abdomen.

SPOTS, ROSE, Spots, red.

SPRAIN, (corrupted from *strain* [?],) *Subluxa'tio*, *Distor'tio*, *Dias'trophê*, *Diastrem'ma*, *Luxa'tio*

imperf'cta, *Strem'ma*, *Thlas'ma Strem'ma*, *Wrench*, (Sc.) *Wramp*, (Prov.) *Wrick*, *Keuch*, *Keuk*, (F.) *Détorse*, *Entorse*, *Foulure*. A violent straining or twisting of the soft parts surrounding the joints. According to the degree of the sprain, the fibrous parts of the joint may be simply stretched, or they may be ruptured.

This affection, which has been regarded as incomplete luxation, occurs, particularly, in the gluglymoid joints;—as in the foot, wrist, knee, &c. The violence causes an afflux of blood; the serum of the blood is poured out, and often the blood itself, owing to the rupture of vessels; and inflammation arises, which ends in resolution or induration. Local, antiphlogistic applications are required, with perfect rest; and, when the inflammatory action has subsided, cold pumping and friction, with the application of a plaster and bandage, will afford support until the parts affected recover their due tone.

SPRAT, YELLOWBILLED, *Clupea thyrssa*.

SPRAYCURE. A variety of the *douche*, which consists in the application of water in the form of spray.

SPRUCE, (from Prussia [?],) see *Pinus abies*.

SPRUCE BEER, *White Spruce Beer*. A beer made as follows: To 10 gallons of water, add 6 pounds of sugar, and 4 oz. of essence of spruce. Add yeast; and bottle, when ready. It is grateful and antiscorbutic.

SPRUCE BEER. POW'DERS may be formed of white sugar, ʒi, and ʒij; *sodæ subcarb.*, gr. xxvj; essence of spruce, gr. x, in each blue paper; and acid of tartar, ʒss in each white paper. For half a pint of water.

SPUITIO, (*spuere*, *sputum*, 'to spit,') Exspuition.

SPUMA ARGENTI, 'froth of silver,' *Plumbi oxydum semivitreum*—s. *Cerevisiæ*, *Yest*—s. *Trium draconum*, *Antimonium muriatum*.

SPUMEUX, Frothy.

SPUMOSUS, (from *spuma*,) Frothy.

SPUNK, (Gael. *spong*, 'tinder,') *Boleus igniarius*.

SPUR, (Sax. *rpun*.) (F.) *Éperon des Artères*. The angle at which the arteries leave a cavity or trunk.

SPUR, Ergot.

SPURBLIND, (*Purblind*,) Myopic.

SPURGE, CAPER, (*Espurge*, [old F.] *Espurge*, from *e*, and *purgare*, 'to purge,') *Euphorbia lathyris*—s. *Cypress*, *Euphorbia cyparissias*—s. *Flax*, *Daphne gnidium*—s. Flowering, large, *Euphorbia corollata*—s. Garden, *Euphorbia lathyris*—s. Greater, *Euphorbia palustris*—s. *Ipecacuanha*, *Euphorbia ipecacuanha*.

SPURGE LAUREL, *Daphne laureola*.

SPURGE, MARSH, *Euphorbia palustris*.

SPURGE OLIVE, *Cneorum tricoceum*, *Daphne mezereum*.

SPURGE, PETTY, *Euphorbia peplus*—s. Sea, *Euphorbia paralias*—s. Spotted, *Euphorbia maculata*—s. Sun, *Euphorbia helioscopia*—s. Wartwort, *Euphorbia helioscopia*.

SPUTA, PRUNE JUICE, (pl. of *sputum*.) see Prune Juice.

SPUTAMEN, (*sputare*, *sputum*, 'to spit,') Saliva, Sputum.

SPUTAMENTUM, Saliva, Sputum.

SPUTATIO, Exspuition.

SPUTATION, Exspuition.

SPUTATORIUS, Expectoant.

SPU'TUM, *Apochrem'ma*, *Anachremptum*, *Chrem'ma*, *Emptys'ma*, *Sputa'men*, *Pty'clon*, *Ptyis'ma*, *Pty's'ma*, *Spit*, (*spuere*, *sputum*, 'to spit,') (F.) *Crachat*. The secretions ejected from the mouth in the act of spitting. It is composed of saliva, and of the mucus secreted by the mucous membrane of the nasal fossæ and fauces, and

often by the membrane of the larynx and bronchia. The sputa are, sometimes, called *cradginous*, when of a greenishyellow colour; *gummy*, when they resemble a thick solution of gum; *frothy*, (F.) *mousseux*, when covered with froth; *rubiculous*, (F.) *rubigineux ou rouillés*, when of a reddishyellow; and *streaked*, (F.) *stries*, when mixed with lines of blood. Also, saliva.

SPIRUM CREBENTEM, Hæmoptysis — s. Dulce, Saccharorrhœa pulmonalis — s. Oris, Salivation — s. Saccharatum, Saccharorrhœa pulmonalis — s. Sanguinis, Hæmoptysis.

SQUALLING, ([Sax.] *xiellan* or *gýllan*, 'to yell,') *Vagitus*.

SQUAMA, Scale, Squamous portion, Table.

SQUAMÆ, (pl. of *squama*), Scaly diseases.

SQUAMARIA, Plumbago Europea.

SQUAMEUX, Squamous.

SQUAMIFORMIS, (*squama*, 'a scale,' and *forma*, 'shape,') Squamous.

SQUAMOÛS, *Squamoïsus*, *Lepidoïdes*, *Lepidoïdes*, *Squamiformis*, *Squamiformis*, (F.) *Squameux*, *Lépidote*, *Sax.*, (from *squama*, 'a scale,') That which relates to or resembles a scale.

SQUAMOÛS PORTION of the temporal bone, (F.) *Portion squameuse ou écailleuse du Temporal*, is the upper part of the temporal, so called from a fancied resemblance to a scale. See Temporal Bone.

SQUAMOUS SUTURE, *Suturæ squamoïsa* seu *mendoïsa* seu *lepidoides* seu *lepidoides* seu *cradnii* *squamoïsa*, (F.) *Suture squameuse ou écailleuse*. The suture which unites the squamous portion of the temporal bone to the parietal.

Certain *scaly* cutaneous affections are also called *squamous*, (F.) *squameux*. See Scale.

SQUARE STALK, *Monarda coccinea*.

SQUARRA TONDENS, (*squarra*, 'sourfiness,' and *tondere*, 'to shave,') *Porrigio decalvans*.

SQUASH, ROUND or **PATTYPAN**, see *Cucurbita pepo* — s. *Warted* or *longnecked*, see *Cucurbita pepo*.

SQUATTING PILLS, (from [Prov.] *squat*, (from *quietus*, [?]) 'quiet,') *Pilulæ opiatæ*.

SQUAWMINT, *Hedeoma*.

SQUAWROOT, *Actæa racemosa*, *Caulophyllum thalictroides*, *Senecio aureus*.

SQUAWWEED, *Erigeron Philadelphicum*.

SQUEAMISHNESS, *Fastidium cibi*.

SQUEASINESS, *Nausea*.

SQUELETTE, *Skeleton*.

SQUELETTELOGIE, *Skeletology*.

SQUILL, *Scilla* — s. *Vinegar of*, *Acetum scillæ*.

SQUILLA, *Scilla*.

SQUINANCIE, *Cynanche*, *Cynanche tonsillaris*.

SQUINANCY, *COMMON*, *Cynanche tonsillaris*.

SQUINANTHIA, *Cynanche tonsillaris*.

SQUINANTHUS, (corr. from *schœnanthus* [?]), *Juncus odoratus*.

SQUINE, *Smilax China*.

SQUINSY, *Cynanche tonsillaris*.

SQUINT, ([D.] *Schuinte*, 'a slope,') *Strabismus*.

SQUINTEYED, see *Strabismus*.

SQUINTER, see *Strabismus*.

SQUINTING, *Strabismus*.

SQUINZEY, *Cynanche tonsillaris*.

SQUIRRE, *Scirrhus* — s. *en Cuirasse*, see *Ligineous*.

SQUIRREL CORN, *Dicentra Canadensis*.

SQUIRRHEUX, *Scirrhus*.

SQUIRRHE, *Scirrhus*.

SQUIRROGASTRIE, *Scirrhogastria*.

SQUIRROSARQUE, *Induration of the areolar or cellular tissue*.

SQUIRRUS, *Scirrhus*.

SQUIRT, *Syringe*.

STACHELBERG, MIN'ERAL WA'TERS OF. Stachelberg Springs are far down in the valley of the Linth, in Switzerland. The waters are sulphureous, and are much visited by gouty and rheumatic invalids.

STACHYS, (*σταχς*, gen. *σταχυς*, 'a spike,') A surgical bandage to the lower belly. The Hypogastrium [?].

STACHYS FÆTIDA, (*σταχυς*,) *Galeopsis*.

STACTE, (*στακτη*,) *Myrrha*.

STADE, (from *studium*,) *Stage* — s. *de la Chaleur*, see *Intermittent fever* — s. *de Froid*, see *Intermittent fever* — s. *de Sueur*, see *Intermittent fever*.

STADIUM, (*σταδιον*,) *Period*, *Stage*.

STADIUM AC'ES seu **STA'SEOS**. The period of the height of a disease.

STADIUM ANNIHILATIO'NIS seu **CONVALESCENT'ILE** seu **RECONVALESCENT'ILE**. The period of recovery from disease.

STADIUM AUGMEN'TI seu **INCREMEN'TI MOR'BI**. The period or stage of increase of a disease.

STADIUM CALORIS, *Stage*, (*hot*) — s. *Convalescentiæ*, *S. annihilationis*.

STADIUM DECREMEN'TI. The period or stage of decrease of a disease.

STADIUM FRIGORIS, *Stage*, (*cold*), see *Intermittent fever* — s. *Incrementi Morbi*, *S. Augmenti morbi* — s. *Latentis contagii*, *Prodromus* — s. *Opportunitatii*, *Prodromus* — s. *Prodromorum*, *Prodromus* — s. *Reconvalescentiæ*, *S. Annihilationis*.

STAFF, ([Sax.] *stæp*, *stæp*,) An instrument of great importance in lithotomy, — being the director for the gorget or knife. It is made of steel; and the handle is flat and rough, in order that it may be more securely held. The groove is on its convex surface.

STAFF TREE, CLIMBING, *Celastrus scandens*.

STAFFA, *Stapes*.

STAG'S PIZZLE, see *Cervus*.

STAGE, *Sta'dium*, (F.) *Estage*, *Étage*, *Stade*. The period or degree of a disease; especially the period of an intermittent: — as the *cold stage*, *sta'dium frig'oris* — *hot stage*, &c.

STAGGERBUSH, *Andromeda mariana*.

STAGGERWEED, *Delphinium*.

STA'GIUM. The sixth part of an ounce.

STAG'MA, gen. *Stag'matis*, (*σταγμα*, from *στέγω*, 'I distil,') *Stalay'ma*. Any distilled liquid or liquor.

STAGMA, *Gutta*.

STAGNA'TION, *Stagna'tio*, (*stagnare*, *stagnatum*, 'to form a pond,' *stagnum*,) Accumulation and retardation of fluids in any part of the body.

STAGNUM CHYL'I, 'pond of the chyle,' *Receptaculum chyl'i*.

STAHL'I'AN. One who follows the doctrine of Stahl.

STAHLIANISM, see *Animist*.

STALAGMA, gen. *Stalay'matist*, (*σταλαγμα*,) *Stagma*.

STALAGMI'TIS CAMBOGIOI'DES, *S. Cambogia*, (from *σταλαγμος*, 'a drop' — the gum exuding by drops.) Gamboge was long ascribed to this fancied Ceylonese plant. It is now known, that the name was given by Murray to specimens of two distinct trees of different genera. See *Cambogia*.

STALAGMUS, (*σταλαγμος*,) *Distillation*.

STALE, *Urine*.

STALICA, (*σταλικά*,) *Repellents*.

STAMMERING, (from [Sax.] *stæmen*, 'to stammer,') *Balbuties* — s. of the *Fingers*, *Cramp*, *writers*'.

STAMP, ([F.] *estampe*,) *Cachet* — s. *Medicine*, *Cachet* — s. *Oculist*, *Cachet*.

STAMPING MILL, see *Pulverization*.

STANG, ([Sax.] *reingan*, 'to sting,') see *Throbbing*.

by organized bodies in contradistinction to the organic or vital. Also, of old, a restraining or astringent remedy.

STATICA MEDICINA. That part of medical observation, which consists in the daily weighing the human body, and the various ingesta and egesta. A title to *Essays of Sanctorius and Keill*.

STATICA SEPTI NARIUM. *Appendix cutanea septi narium.* The cutaneous appendage to the septum narium.

STATICAL, Static.

STATICE CAROLINIANA. of the United States, (*statum*, from *statikos*, 'restraining,' owing to its astringency.) *American or Sea-side Thrift, Marsh Rosemary, Marshroot, Inkroot, Sea Lavender, Ord. Plumbaginaceae*, is considered to be a mere variety of the *Statice limonium*. The root is *Statice*, of the Ph. U. S.

STATICE LIMO'NIUM, S. Maritimum, Be'hen ru-brum, Limonium, L. majus seu vulgare, Be'hen, Sea Lavender, Red Be'hen, Marsh Rosemary, Lavender Thrift, (F.) Behen rouge. Rosmarin des marais, Lavande triste ou de mer. The roots have been esteemed astringent, tonic, and eminently antiseptic.

STATICE MARITIMUM, S. limonium—s. *Vulgare, S. limonium.*

STATICS, see *Static.*

STATION, Sta'tio, Sta'sis, Sta'tus, (stare, statum, 'to stand.') The act of standing.

STATIONARY, Stationarius, (F.) Stationnaire. A name given by Sydenham and Stoll to certain diseases, which depend upon a particular state of the atmosphere; and which prevail in a district for a certain number of years, and then give way to others.

STATISTICS, MEDICAL, Vi'tal Statistics, Biostatistics, (status, 'a state,') (F.) Statistique médicale. That part of medicine which relates to details—*medico-statistical*—connected with the mortality, salubrity, &c., of different states, countries and situations.

It has, also, been used in a wider sense to signify the same as *Numerical method, (F.) Méthode numérique*; which is applied to the systemization of medical facts, or to the use of numbers in the investigation of disease.

STATISTICS, VITAL, Statistics, medical.

STATURE, Statu'ra, Meg'ethos, Magnitu'do corporis, (F.) Taille, (from stare, statum, 'to stand.') The height of the human figure. The following table exhibits the mean height, in feet and decimals, of 1800 men, taken as they were entered upon the lists of recruits filed in the office of the Adjutant-General of the United States. The results are given for 100 men from each State; and no recruit under five feet five inches was received.

	Mean height. Feet.	Six feet and over.	Greatest height.
Indiana.....	5.7004	18	6.41½
Kentucky.....	5.7729	18	6.33½
Ohio.....	5.7537	15	6.31½
Tennessee.....	5.7779	18	6.3
Maine.....	5.7314	11	6.2
Vermont & New Hampshire.....	5.6951	6	6.1
Massachusetts and Connecticut.....	5.6821	5	6.3
North Carolina.....	5.7814	24	6.33½
Georgia.....	5.8272	30	6.61½
South Carolina.....	5.7729	15	6.41½
Alabama.....	5.7647	17	6.4
Virginia.....	5.7488	15	6.2
New York.....	5.6505	4	6.11½
Pennsylvania.....	5.6756	5	6.1
New Jersey and Delaware.....	5.6509	6	6.1
Maryland.....	5.7130	9	6.2
Illinois.....	5.7696	17	6.3
Missouri.....	5.7162	8	6.11½

The minimum height for the American soldier is now five feet three inches; whilst the British

standard is five feet five inches, and the French five feet one inch and a half.

It would seem, from the researches of Professor S. H. Dickson, that the stature of Americans is decidedly superior to that of Europeans, and that the same applies, but to a less degree, to the weight.

The following table exhibits the results of several admeasurements, with the heights of some of the monkey species, on the authority of Mr. Lawrence.

	STATURE.	
	Feet.	In.
An Englishman.....	6	4½
	5	00
	5	4
An Englishwoman.....	5	00
	5	8
European male skeleton.....	5	8
Do.....	5	5
A Negro at the Lunatic Hospital, Liverpool.....	5	10½
Another from Virginia.....	5	5½
Another from the Gold Coast.....	5	8
Another.....	5	00
Negro skeleton.....	4	11
Another.....	5	7½
A Lascar.....	5	4
Venus de Medici.....	5	00
Tyson's Chimpanzé (<i>Simia Troglodytes</i>).....	2	2
Mr. Abel's Orang Otang.....	2	7
Camper's Do.....	less than	30
Mr. White's Monkey.....	2	2

STATUS, (stare, statum, 'to stand.) Acme—s. *Cruditatis, Crudity*—s. *Erethiticus, Nervous diathesis*—s. *Gastricus, Embarras gastrique*—s. *Nervosus, Nervous diathesis*—s. *Parturiendi, Parturition*—s. *Sthenicus, Sthenia*—s. *Verminosus, Helminthiasis.*

STAVESACRE, Delphinium staphisagria.

STAXIS, (from σταξω, 'to drop,') Apostax'is. A slight epistaxis or nasal hemorrhage.

STEAMDOCTOR. A term applied to one who treats all, or most, diseases by steam.

STEAR, (στεαρ, gen. στεαρος,) see Fat, Sevm.

STÉARATÉ, Emplastrum.

STEARIN, see Fat.

STEARINUM CETACEUM, Cetaceum.

STÉAROLÉ, (from stear,) Pomatum—s. *Solide, Emplastrum.*

STEAROSIS, (from stear, and osis,) Steatosis.

STEARRHŒA, Steatorrhœ'a, Seborrhœ'a, Seborrhag'ia, Smeqatorrhœ'a, Smeqorrhœ'a, Fluxus seba'ceus, Seba'ceous Flux, (stear, and fluo, 'I flow.') Augmentation of secretion from the sebaceous follicles of the skin.

STEARRHŒA NIGRICANS. A term used by Dr. Neligan, of Dublin, for a black discoloration of the face, observed in young females affected with derangement of the catamenia, and which, in his opinion, is owing to a dark secretion from the cutaneous sebaceous glands.

STEATITES, (from stear,) Polysarcia.

STEATOCE'LE, (stear, and κηλη, 'a tumour.') A tumour formed by a collection of steatomatous matter in the scrotum.

STEATO'MA, gen. Steato'matis, Tu'mor lar'deus, (stear, and oma.) Seva'tio, Emphy'ma encystis steato'ma. An encysted tumour, whose contents are similar to fat.

STEATOPLY'GA, (stear, and πύγη, 'buttocks.') *Buttock-hump.* Fat buttocks; such as exist in the Hottentot female.

STEARRHŒA, Stearrhœa.

STEATO'SIS, Stearo'sis, (from stear and osis.) Fatty degeneration of a part, or unusual deposition and accumulation of fat in its adipose tissue.

STEATO'SIS COR'DIS, Degenera'tio adipo'sa cor'dis, Cardiomyoliposis and Cardiomyoliposis (Pi orry), Fatty heart. Unusual deposition of fat on the heart:—by some presumed to be occasionally

a true adipose degeneration of the substance of the heart.

STEATOSIS HEPATICA, Adiposis hepatica.

STEATOZOON FOLLICULORUM. (*steat*, and ζωον, 'an animal,') Acarus Folliculorum.

STEEL, ([Sax.] *stæle*.) Chalybs.

STEEP, Rennet.

STEEPLEBUSH, *Spiræa tomentosa*.

STEER'S OPODELDOCH, see Opodeldoch, Steer's.

STEGANOP'ODES, (*στεγανόδες*.) *Planip'edes*. They who have flat feet, or are webfooted, (from *στεγανός*, 'covered,' and *πους*, 'foot,')

STEGANUS, (*στεγανός*.) Constipated.

STEGMONT, see Parturient.

STEGNO'SIS, *Steno'sis*, *Constipatio*, *Stipa'tio*, *Condensatio*, *Obstruc'tio*, (*στεγνώσις*, from *στεγνώνω*, 'I constrict,') Constriction or narrowness of the pores and vessels. Stricture. Constipation. Suppression or stopping or stoppage of the evacuations. See Aretatio.

STEGNOTICA, (*στεγνωτικά*.) Astringents.

STEROSIS, (*στερωσις*.) Sterilitas.

STEL'LA, 'a star,' *Fus'cia stella'ta*, (F.) *Étoile*. A bandage, improperly compared to a star, because it is crossed like the letter X. This bandage is applied upon one or both shoulders. In the first case, it is called *simple*; in the second, *double*. It was employed in fractures of the sternum, clavicle, and scapula; in luxations of the humerus, &c.

STELLA TERRÆ, *Plantago coronopus*.

STELLARIA MEDIA, (*stella*, 'a star,') *Alsine media*—*s. Scilla*, *Scilla*.

STELLATA RUPINA, (from *stella*.) *Centaurea calcitrapa*.

STELL'ULE VERHEYEN'II, (*stella*, 'a star,' *stell'ula*, 'a little star,') *Substan'tia stella'ta serpenti'na* seu *radia'ta*, *Ellych'nia*. A stelliform plexus of veins at the surface of the kidney, which gives origin to the interlobular veins.

STELOCHITIS, *Osteocolla*.

STEMA, (*στυμα*.) Penis, (especially the glans.)

STENAGMUS, (*στεναιγμός*.) *Suspirium*.

STENAGMA, (*στεναιγμα*.) *Suspirium*.

STENO, (from *στενός*), in composition, Narrow.

STENO, DUCT OF, see Parotid.

STENOBREGMATE, (*steno*, and *βρεγμα*, 'the top of the head,') see Cranium.

STENOCARDIA, (*steno*, and *καρδία*, 'heart,') *Angina pectoris*, *Cardiostenosis*.

STENOCHORIA, *Stenochori'asis*, (*steno*, and *χωρα*, 'space,') *Angustia*, *Narrowness*, *Stricture*. Employed by some for contraction of the vagina especially.

STENOCHORIA INTESTINI RECTI, Stricture of the rectum—*s. Cæophagi*, *Dysphagia constricta*.

STENOMYCTE'RIA, (*steno*, and *μυκτηρ*, 'nose,') Contraction of the nasal passages.

STENON or STENO, DUCT OF, see Parotid.

STENOSIS, (*στενωσις*, *stenos* and *osis*), *Stegnosis*.

STENOSTEGNOSIS, (*steno*, and *stegnosis*), *Stenostenosis*.

STENOSTENO'SIS, *Stenostegno'sis*, (*steno*, and *στενωσις*, 'contraction,') *Obstruc'tio duct'us Stenonia'ni*. Narrowness or contraction of the duct of Steno.

STENOSTOM'IA, (*steno*, and *στομα*, 'mouth,') Contraction of the mouth.

STENOTHORAX, gen. *Stenothora'cis*, (*στενωθώραξ*, from *steno*, and *θώραξ*, gen. *θώρακος*, 'the chest,') One who has a narrow chest.

STENTOROPHO'NUS, (*στένωρ*, *Stentor*, a Greek of powerful voice in the army before Troy; and *φωνή*, 'voice,') One who has a strong voice.

STEP, ([Sax.] *stæp*), *Pas'sus*, (F.) *Pas*. The step is produced by the separation of the two lower limbs, and includes that space plus the length of the foot.

STEPHANE. (*στέφανος*.) Crown.

STEPHEN'S, MRS., REMEDY FOR THE STONE. This consisted of lime, prepared by calcining the shells of eggs and snails, made into pills with soap. A decoction was, also, administered, consisting of chamomile, fennel, parsley, and burdock, with a portion of Alicant soap. Its virtues were dependent upon the lime and the tonic properties of the decoction. Hales, Hartley, Kirkpatrick, Lobb, and others, wrote in favour of this nostrum; and it was ultimately bought by the British parliament for £5000—about \$25,000.

STEPHENSIA ELONGATA, *Matico*.

STEPMOTHER'S BLESSING, Hangnail.

STERCOLIN, (from *stercus*.) *Serolin*.

STERCORA'CEOUS, *Stercora'rius*, *Diachoret'icus*, *Ster'coral*, (from *stercus*.) *Fæ'cal*, *Fæ'cal*, (F.) *Stercoraire*, *Stercoral*.

The French use the term *Fistules stercoraires* for those which are kept up by the constant passage of fecal or *stercora'ceous* matter. The name *Stercoraires* is sometimes given, in derision, to those physicians who are fond of purging their patients: à posteriori has been similarly used.

STERCORAIRE, *Stercoraceous*.

STERCUS, gen. *Ster'coris*, Excrement, Fimus—*s. Anseris*, Fimus anseris—*s. Caninum album*, Album Græcum—*s. Diaboli*, *Asafetida*—*s. Equi non castrati*, Fimus.

STERCUS HUMA'NUM, *Homerda*. Human excrement.

STERCUS LACERTÆ, *Crocodilia*.

STEREOCYST'IS, (*stereus*, and *κυστις*, 'cyst,') A hard encysted tumour.

STEREOSCOPE, *Stereoscop'ium*, (*stereus*, and *σκοπεω*, 'I view,') A physical instrument by means of which plane pictures of an object are made on each retina, and so blended as to appear as one in relief. Also, an instrument for detecting a calculus in the bladder, and foreign bodies in the soft parts. It resembles a common catheter, having an expanded free extremity.

STEREOT'ICA, (from *stereus*.) Fortuitous lesions or deformities affecting the hard parts. The 2d order, class *Tychica* of Good.

STEREUS, (*στερεός*.) Solid.

STER'ILE, *Ster'ilis*, *A'gonos*, *Aph'orus*, *Aphor'icus*, *Ablast'es*, *Infecun'dus*, *Barren*, (F.) *Stérile*, (from *στερρός*, 'barren,') Not producing fruit; infecund.

STERILIS, *Sterile*.

STERILITAS, *Agon'ia*, *Steiro'sis*, *Stiro'sis*, *Agenn'e'sia*, *Agenne'sis*, *Infecun'ditas*, *Steril'ity*, *Acy'e'sis*, *Bar'renness*, *Infecun'dity*, *Apho'ria*, *Atoc'nia*, *Atoc'ia*, (F.) *Stérilité*. Quality of that which is barren. Sterility may be *absolute* or *relative*. The former depends upon some vicious conformation of the genital organs, apparent or concealed, and occasionally admits of cure. Sterility is *relative* when a female does not conceive with one individual, but does with another. Sterility in females is often dependent upon a condition of the uterus characterized by membranous menstruation.

STERNAD, see Sternal.

STER'NAL, *Sterna'lis*, (from *sternum*.) Relating or appertaining to the sternum or breast bone.

STERNAL ASPECT. The aspect of the body where the sternum is situate.—(Barelay.) *Sternad* is used by the same writer adverbially to signify towards the sternal aspect.

STERNALGIA, *Sternodyn'ia*, (*sterno*, and *αλγος*, 'pain,') Pain referred to the sternum. *Angina pectoris*.

STERNO, in composition, the sternum.

STERNOCLAVICULAR, *Sternoclavicula'ris*. That which relates to the sternum and clavicle.

STERNOCLAVICULAR ARTICULA'TION is the union of the inner extremity of the clavicle with a round

facette at each side of the superior extremity of the sternum. In this articulation there are, besides the incrusting cartilages, two radiated sternoclavicular, (F.) *Sterno-claviculaires rayonnées*; one anterior, and one posterior—an interclavicular ligament extending between the two clavicles, and two synovial capsules, separated by an interarticular fibrocartilage. The costoclavicular ligament may also be considered to belong to this articulation.

STERNOCLAVIOMASTOÏDIEN, Sternocleidomastoideus—s. Cleidobrachialis, Pectoralis major—s. Cleidohyoideus, Sternohyoideus.

STERNOCLEI'DOMASTOÏDE'US, *Sternomastoïde'us*, *Clei'domastoïde'us*, *Mastoïde'us*, *M. anterior*, *Nuta'tor cap'itis*; (F.) *Sternoclaviomastoïdien*, *Sternomastoïdien*; (*sterno*, κλεις, 'the clavicle,' and *μαστος*, 'the mastoid process.') A muscle situate at the anterior and lateral part of the neck. It is long and flattened; simple, above; and divided into two portions below; where it is inserted into the anterior and superior part of the sternum, and the inner quarter of the clavicle. Above, it terminates at the mastoid process, which it embraces, at the neighbouring part of the temporal bone, and at the outer third of the upper curved line of the occipital bone. It carries the head forward; inclines, and rotates it to one side. When both contract, they bend the head directly forwards.

STERNOCOSTAL, *Triangularis sterni*—s. *Costo-clavio-huméral*, *Pectoralis major*—s. *Huméral*, *Pectoralis major*.

STERNOHYOÏDE'US, *Sternocleidohyoïdeus*, *Sternohyoïdes*, *Hyoïdes pri'mus*; (from *στέρον*, 'the sternum,' and *ὡοειδής*, 'the os hyoides.') A muscle situate at the anterior part of the neck; inserted, above, at the inferior edge of the body of the hyoid bone; below, into the upper part of the posterior surface of the sternum;—at the posterior part of the sternoclavicular articulation, and sometimes into the cartilage of the first rib. It depresses the os hyoides.

STERNOMASTOÏDIEN, *Sternocleidomastoïdeus*—s. *Pubien*, *Rectus abdominis*.

STERNOTHYROÏDE'US, *Sternothyroïdes*, *Bronchius*. A muscle, situate at the anterior part of the neck. It is attached, above, to the oblique line seen on the outer surface of the thyroid cartilage; and terminates, below, at the upper part of the posterior surface of the sternum, and the cartilage of the second rib. When it contracts, it depresses the thyroid cartilage and os hyoides by the medium of the thyrohyoid membrane.

STERNODYNIA, (*sterno*, and *ὄδυνη*, 'pain,') *Sternalgia*—s. *Syncoalis*, *Angina pectoris*—s. *Syncoptica* et *Palpitans*, *Angina pectoris*.

STERNOÏMPHALODYMIA, (*sterno*, *ομφαλος*, 'navel,' and *δύω*, 'to mingle,') see *Somatodymia*.

STERNOPA'GIA, (*sterno*, and *παύω*, 'I fasten,') A monstrous union of fœtuses, in which they are attached at the sternum.

STERN'UM, *Ster'non*, (*στέρον*, from *στέγος*, 'solid,') *Pec'tus*, *Os as'ser* seu *pec'toris* seu *xiphoides* seu *gladi'oli* seu *ensifer'mē*, *Scu'tum cor'dis* seu *pec'toris*, *Breastbone*, *Hæ'mal Spine* of Owen. An azymous symmetrical bone, situate at the forepart of 'he chest. It is flat; broad above; narrower in the middle; and terminates, below, by a pointed cartilage—the xiphoid. It has an anterior or cutaneous surface, a posterior or mediastinal—a superior or clavicular extremity, and an inferior or abdominal. It is articulated with the clavicles and the seven upper ribs on each side, by means of their cartilages. According to Héclard, it is formed of six principal bones, which he names, from their position, 1. *Primi-sternal* or *-lari sternal*: 2. *Duo-sternal*: 3. *Tri-sternal*: 4.

Quarti-sternal: 5. *Quinti-sternal*: 6. *Ultimi-sternal* or *Easi-sternal*.

STERNUMENTUM, (*sternuo*, *sternutum*, 'to sneeze,') Sneezing.

STERNUTAMENTORIA, (*sternuto*, *sternutatum*, 'to sneeze,') *Achillea ptarmica*.

STERNUTAMENTUM, Sneezing.

STERNUTATIO, Sneezing.

STERNUTATOIRE, Sternutatory.

STERNUTATORY, *Sternutato'rius*, *Ster'nutory*, *Sternutamen'tum*, *Erri'hnum*, *Ptar'micum*, *Anarrhi'num*, (F.) *Sternutatoire*. A substance which provokes sneezing; as tobacco, &c.—See *Erhine*.

STERNUTORY, Sternutatory.

STERTEUR, Stertor.

STER'TOR, *Ron'chus*, *Rhon'chus*, *Ren'chus*, *Rhen'chus*, (F.) *Sterteur*, *Ronflement*, (*stertere*, 'to snore,') The deep snoring which accompanies inspiration in some diseases, particularly in apoplexy. Such respiration is said to be *ster'torous*, (F.) *Stertoreux*; *Respira'tio stertuo'sa* seu *stertens* seu *stertorosa*.

STERTOROUS, see Stertor.

STETHÆ'MIA, *Hyperæ'mia Pulmo'num* seu *Pec'toris*, *Conges'tio Pulmo'num* seu *Pec'toris*, (F.) *Hypérémie* ou *Congestion des Poumons*; (from *stetho*, and *αἷμα*, 'blood,') *Hyperæmia* of the lungs. Congestion or accumulation of blood in the pulmonary vessels.

STETHO, (from *στήθος*, 'chest,') in composition, 'the chest.'

STETHOCATHARSIS, (*stetho*, and *καθάρσις*, 'a purging,') Expectoration.

STETHOCHYSIS, (*stetho*, and *χύνω*, 'to pour out,') Hydrothorax.

STETHODESMIS, (*stetho*, and *δεσμός*, 'a bandage,') Corset.

STETHODESMIUM, Corset.

STETHODESMUS, Corset.

STETHOMETER, (*stetho*, and *μετρον*, 'measure,') An instrument, proposed by Dr. Richard Quain, for measuring the extent of movement of the walls of the chest, as a means of diagnosis in thoracic diseases. It consists of a flat case, not unlike a watch-case, having on its upper surface a graduated dial and index, which is acted on by a simple mechanism. A silk cord, long enough to surround one-half the chest, passes through an aperture in the case. The dial is graduated into fifty equal parts, each of which represents the 1-200th of an inch. In using the instrument, the dial is placed firmly against the spine, and the point of the tape over the sternum, or at any other spot according to the part to be examined. The extension of the string, as indicated by the dial, shows the amount of elasticity of the parietes of the thorax.

Under the name *Thoracom'eter*, Dr. Sibson has described an instrument of his invention, which is employed with a similar view as the stethometer of Dr. Quain.

STETHOPARALYSIS, *Paral'ysis nervo'rum thorac'is*, *phrenicorum*, &c.; (*stetho*, and *παράλυσις*, 'paralysis,') Palsy of the muscles of the chest.

STETHORRHEUMA, (*stetho*, and *ῥεύμα*, 'de-fluxion, rheumatism,') *Pleurodynia*.

STETHOSCOPE, *Stethoscop'ium*, *Stethos'copum*, *Echoscop'ium*, *Psophom'eter*, *Thoracoscop'ium*, (*stetho*, and *σκοπεω*, 'I examine,') An instrument invented by Laënnec, of Paris, for exploring the chest. The stethoscope, sometimes called *Pectoriloque*, is a cylinder of wood, from four inches to a foot long; pierced through and through by a longitudinal canal about a quarter of an inch in diameter. The longer are generally composed of two portions, that fit together by means of a screw, one of which is hollowed at the extremity, in the shape of a funnel. These two portions being screwed to each other, the

physician lays hold of the instrument, as he would of a pen, puts the funnel-shaped extremity on the chest of the patient, and applies his ear to the other. To explore the pulsations of the heart, the funnel is plugged up by a piece of the same kind of wood accurately adapted to it, and pierced by a canal of the same width as that in the body of the instrument. Stethoscopes are now made in the plainest and lightest manner and without any plug. Double stethoscopes have, likewise, been invented to enable both ears to be used at the same time.

Stethoscopes are sometimes flexible, like the flexible ear trumpet.

This mode of examining affections of the chest, *Stethoscopy*, *Stethoscop'ia*, &c., is what Laënnec terms *Auscultation médiate*, *Me'diate Auscultation*.

STETHOSCOPIA, see Auscultation, and Stethoscope.

STETHOSCOPI'IC, *Stethoscop'icus*, (F.) *Stéthoscopique*. Furnished by stethoscopy or auscultation,—as a 'stethoscopic sign.'

STETHOSCOPY, see Stethoscope.

STETHUS, (στήθος,) Thorax.

STEW, [old F.] *estuve*, [Sax.] *stēva*, Stove.

STHENIA, *Sthen'ia*, *Diath'esis sthen'ia*, *Stathus sthen'icus*, *Vis aucta*, (σθένος, 'strength,' 'power.') Excess of strength; excitement of the organic actions. A word used, especially by the Brunonians; as well as *Asthenia*, its antitheton. *Sthenic* or *dynamic diseases* are such as depend upon excessive excitement.

STHENICUS, (from σθένος, 'strength,') Active.

STHÉNIQUE, see Active.

STHENOPYRA, (from σθένος, 'strength,' and πυρ, 'fever,') see Synocha.

STIAN, ([Sax.] *stīgan*, from *stīgan*, 'to ascend, to rise,') Hordeolum.

STIBI, (στίβι,) Antimonium.

STIBIALIS, Antimonial.

STIBIALISM, *Stibialis'mus*, (from *stibium*, 'antimony.') The aggregate phenomena produced by antimonials, when given in large quantities,—antimonial intoxication or poisoning.

STIBIATUS TARTRIS LIXIVLÆ, Antimonium tartarizatum.

STIBII BUTYRUM, Antimonium muriatum—s. Deutomurias sublimatus, Antimonium muriatum—s. Hydrosulphuretum rubrum, Antimonii sulphuretum rubrum—s. Oxidum acido muriatico oxygenato paratum, Algaroth—s. Oxidum album mediante nitro confectum, Antimonium diaphoreticum—s. Oxidum præcipitando paratum, Algaroth—s. Oxodes album, Antimonium diaphoreticum—s. Oxodes sulphuretum, Antimonii sulphuretum præcipitatum—s. Oxydi sulphurati hydrosulphuretum luteum, Antimonii sulphuretum præcipitatum—s. Oxydulati sulphuretum, Antimonii sulphuretum præcipitatum—s. Oxydulum vitreum, Antimonii vitrum—s. Oxydum semivitreum, Antimonii vitrum—s. Oxydum sulphuratum, Oxydum stibii sulphuratum—s. Subhydrosulphas, Antimonii sulphuretum rubrum—s. Submuriat præcipitando paratum, Algaroth—s. Sulphurati hydrosulphuretum rubrum, Antimonii sulphuretum rubrum—s. Sulphuretum nigrum, Antimonium—s. Vitrum, Antimonii vitrum—s. et Potassæ deutotartaras, Antimonium tartarizatum.

STIBIODERMIC, see Unguentum antimonii.

STIBIOSUM HYDROSULPHURETUM CUM SULPHURE, Antimonii sulphuretum præcipitatum—s. Oxidum, Antimonium diaphoreticum.

STIBIUM, Antimonium.

STICHAS (Stechas) ARABICA, *Lavandula stoechas*.

STICKWORT, Agrimony.

STICTA PULMONACEA, (from στίκτος, 'spotted,') Lichen pulmonarius.

STIFF JOINT, MUSCULAR, *Contractura*

STIFF JOINTS, Milk sickness.

STIFFNECK, Torticollis.

STIGMA, (στίγμα, 'a small puncture,') *Cicatricula*.

STIG'MATA (pl. of *stigma*,) see *Nævus*. The breathing pores of insects are so called.

STIG'MATA OVARIO'RUM, *Cicatri'ces Ovario'rurum*. The cicatrices left in the ovary after the escape of ova.

STILBOMA, gen. *Stilbo'motis*, (στίλβωμα, from στίλβω, 'to make shining,') Cosmetic.

STILL, (from *stilla*,) Alembic.

STILET, (dim. of *stilus*,) Specillum.

STILLA, Gutta.

STILLATIO SANGUINIS E NARIBUS, (*stillare*, *stillatum*, 'to drop,') Epistaxis.

STILL'BORN, *Na'tus Mor'tuus*, (F.) *Mort-né*. Born still or lifeless,—dead born; a stillborn child,—*Neeroto'ion*.

STILLICIDIUM, *Substil'lum*, (*stilla*, 'a drop,' and *cadere*, 'to fall,') This term is often used synonymously with *Strangury*. It means the flow of any fluid—particularly the urine—drop by drop. Also a kind of embrocation, *Stillicidium su'pra par'tem*, effected by dropping a liquid upon a part.

STILLICIDIUM LACRYMARUM, Epiphora—s. *Narium*, *Coryza*—s. *Sanguinis e naribus*, Epistaxis—s. *Supra partem*, see *Stillicidium*.

STILLIN'GIA, (after Dr. Benj. Stillingfleet, an English botanist.) *S. sylvat'ica*, *Queen's Root*, *Queen's delight*, *Yew root*, *Mar'cory*, *Cockup hat*. An indigenous plant, which grows in pine barrens, from Virginia to Florida. *Sec. Syst.* Monœcia Monadelphica. *Nat. Ord.* Euphorbiacæ.

The root—*Stillingia*, Ph. U. S.—is said to be used in syphilis, obstinate cutaneous affections, &c., like sarsaparilla.

STILUS, Stylus.

STIMATO'SIS, *Stymato'sis*, *Hæmorrhag'ia Pe'nis*, *Stymatorrhag'ia*, *Phallorhag'ia*, *Caulorhag'ia*, (from στυπα, 'priapism,' itself from στυν, 'to erect,' and *osis*.) Hemorrhage from the male organ, accompanied or not with priapism.

STIMMI, (στίμμι,) Antimonium.

STIM'ULANT, *Stim'ulans*, *Exc'itans*, *Inc'itans*, *Incent'ivum*, *Irrit'ans*, *Exc'itant*, *Inc'itant*, *Hyperthén'ic*, *Pyretogenet'ic*, (from *stimulare*, 'to goad,') A medicine which has the power of exciting the organic action of the different systems of the economy. Some stimulants are *diffusible*,—that is, have a prompt but temporary action; others are *permanent* or *persistent*. The action of stimulants is called *Stimula'tion*, *Stimula'tio*, *Incita'tio*.

The chief stimulants are the following: *Æther Sulphureus*, *Ammonia*, *Asafetida*, *Brucia*, *Camp'hora*, *Capsicum*, *Castoreum*, *Cinnamomum*, *Copaiba*, *Cubeba*, *Guaiaicum*, *Hydrargyri Præparata*, *Iodinum*, *Nux Vomica*, *Olea Essentia*, *Opium* (in small dose), *Piper*, *Sinapis*, *Strychnia*, *Zingiber*,—*Alcohol*, *Caloric*, *Electricity*, *Mental Emotions*, (the *Impetuous*.)

STIM'ULUS, 'a goad.' *Incitamen'tum*, *Irritamen'tum*, *Potent'ia Irr'itans*. Any thing which excites the animal economy.

STINE, Hordeolum.

STINK POKE, *Dracontium fœtidum*—s. *Weed*, *Chenopodium anthelminticum*, *Datura stramonium*, *Polanisia graveolens*.

STIPATIO, (*stipare*, *stipatum*, 'to stop up,') *Stegnosis*.

STIPATIO TELÆ CELLULO'SÆ INFANT'UM. Induration of the cellular tissue of children.

STIROSIS, (στίρωσις, from στίρω, 'sterile,' and *osis*.) *Sterilitas*.

STITCH, (same root as *stick*, Sax. *stīcan*, 'to pierce,') (Sc.) *Stonnd*, (Prov.) *Day*, (F.) *Point*

(L.) *Punctum*. An acute lancinating pain, like that produced by the puncture of a sharp needle.

STITCH, COMMON, Suture, common — s. Royal, see Suture—s. in the Side, Pleurodynia.

STIZOLOBIUM. (σίζω, 'to prick,' and λοβιον, 'a small pod,') *Dolichos pruriens*.

STENGIS, (στέγγις,) Strigil.

STOBÆA RUBRICAULIS, (after D. Stobæus, of Sweden, a patron of Linnaeus,) A South African plant, *Nat. Ord.* Composite. A tincture of the bruised roots is diuretic, and of great service in gravel.

STOCK'ING, LA'CED, (F.) *Bas lissé*. A bandage used by surgeons, which consists of a stocking made of firm cloth or dogskin, admitting of being tightly laced anteriorly. It is made, also, of elastic gum cloth. It is employed in varices and ulcers of the legs.

STOCK'PORT, MINERAL WATERS OF, See New York, mineral waters of.

STECCHAS ARABICA, (στοιχας,) *Lavandula stoechas*.

STECCHIOL'OGY, *Stæchiolog'ia*, (F.) *Stæchiologie*; (στοιχεων, 'element,' and λογος, 'a discourse,') A treatise, or the theory, of elements; hence of the elementary parts of organized substances.

STOICHEION, (στοιχειον,) Element.

STOMA, gen. *Stom'atis*, (στομα, gen. στοματος,) Mouth.

STOMAC'ACĒ, (στομακαχη,) *Stomatoc'acē*, *Stomal'gia*, *Labriul'cium*, *Cheiloc'acē*, *Uloc'acē*, *Cheilonal'cia*, *Stomaton'oma*, *Stomatonecro'sis*, *Cancer O'ria*, *Can'ker*, (*stoma*, and *κακος*, 'evil,') Fœtor of the mouth with a bloody discharge from the gums. (F.) *Cancer aquaticus*; see Cancer Aquaticus. Also, Scurvy.

STOMACACE, *Porphyra nautica* — s. Gangrænomum, *Cancer aquaticus* — s. Universale, *Purpura hæmorrhagica*.

STOMACAL, Stomachal.

STOMACH, (στομαχος,) *Stom'achus*, *Ventriculus*, *Né'dys*, *Anocœ'lia*, *Gast'er*, *Gluttu'pateus*, *Maw*, (of animals,) (Sc.) *Kyte*, *Pechan*, (F.) *Estomac*, (from *stoma*.) One of the principal organs of digestion. It is a musculomembranous reservoir; continuous, on the one side, with the œsophagus; on the other, with the duodenum. It is situate beneath the diaphragm, between the liver and the spleen; and occupies the epigastrium and a part of the left hypochondrium. In it the food is formed into chyme. When viewed externally, the stomach has, 1. An anterior face, which looks a little upwards. 2. An inferior face, directed downwards. 3. An inferior or colic margin, which is convex and extensive, and is called the greater curvature, *curvaturæ major*, (F.) *Grand courbure*. It gives origin to the omentum majus. 4. A superior or diaphragmatic margin, which is shorter, concave, and is called the lesser curvature, *curvaturæ minor*, (F.) *Petit courbure*. The lesser omentum is attached to this. 5. A left or œsophageal orifice, called, also, the *cardia*, *Os ventriculi* or *upper orifice*. 6. A right or intestinal or inferior orifice, called the *pylorus*. 7. A considerable dilatation, situate to the left of the cardia and greater curvature—the great tuberosity or great cul-de-sac or fundus of the stomach; *fundus seu succus ventriculi*, and, 8. A less extensive dilatation, situate to the right of the greater curvature,—the lesser tuberosity or lesser cul-de-sac, *Antrum pylori seu pyloricum*, *Pars pylorica*. The inner surface of the stomach is of a reddishwhite colour, and has a marbled appearance. It is constantly covered by thick mucus, and is lined by a mucous membrane, which presents numerous wrinkles. The parietes of the stomach consist of three membranes in superposition. The outermost is serous,

and is an extension of the peritoneum. The middle coat is muscular,—some of its fibres running longitudinally; others, transversely, and others obliquely. The innermost membrane is of a mucous nature, *Crusta villosa ventriculi*, *Gastromyco'deris*, but not exactly a continuation of the membrane that lines the œsophagus. The mucous and muscular membranes form, at the pylorus, a valve, called the *Pyloric valve*. These three coats are united by a dense, close, areolar membrane; and, between the mucous and muscular coats, along the two curvatures especially, is a quantity of muciparous glands, called *Glands of Brunner*. The arteries of the stomach are very numerous, and proceed from the coronaria ventriculi, the pyloric, splenic, and right and left gastroëpiploic. The veins have the same name, and pursue the same course as the arteries. They pour their blood into the trunk of the vena porta. Its lymphatic vessels are very numerous, and pass into ganglia, situate along the two curvatures. The nerves of the stomach proceed from the pneumogastric, and three divisions of the cœliac plexus.

STOMACH CELL, see Favulus—s. Disease, Liosis.

STOMACH, FIRST, Inguivies.

STOMACH PUMP, *Gastren'chyta*, *Anili'a gas-trica*, (F.) *Pompe pour l'estomac*. A useful instrument for conveying fluids to the stomach when deglutition is impracticable, and for pumping out the contents of that organ. It consists of a forcing syringe, to the bottom and nozzle of which flexible tubes can be attached. When the object is to force fluids into the stomach, the stomach tube must be attached to the nozzle, and passed down the throat,—the tube connected with the bottom of the syringe being placed in the fluid to be injected. To empty the stomach, the stomach tube must be attached to the bottom of the instrument. This instrument has been of great service where poisons have been taken. The antidote, when any exists, and is at hand, should be administered immediately prior to its use.

The stomach pump was first employed in America by Dr. Physick, in the year 1812.

STOMACH, SECOND, Proventriculus—s. Sick, Milk sickness—s. Third, Ventriculus callosus.

STOMACH TUBE, see Tube, œsophageal.

STOMACHAL, *Stomach'ic*, *Stomach'ical*, *Cardiacus*, (στομαχος,) *Stomach'icus*, *Stomach'al'is*, *Viscer'al'is*, (F.) *Stomacal*, *Stomachique*. That which belongs to the stomach; that which is good for the stomach; which strengthens the stomach. A medicine that gives tone to the stomach.

STOMACHALGIA, (*stomach*, and *αλγος*, 'pain,') Cardialgia.

STOMACHI RESOLUTIO, 'weakness of stomach,' Dyspepsia.

STOMACHIC, Stomachal.

STOMACHICAL, Stomachal.

STOMACHIQUE, Stomachal.

STOMACHUS, (στομαχος, from *stoma*.) This word has been used in several different acceptations; for example, for the œsophagus, for the cardiac orifice of the stomach (see Cardia), and for the stomach itself. Hippocrates calls the neck of the bladder *στομαχος κυστεως*; and the neck of the uterus *της μητρας στομαχος*.

STOMAL'GIA, *Stomatal'gia*, *Stomatodyn'ia*, (*stoma*, and *αλγος*, 'pain,') Pain in the mouth.

STOMAPYRA, (*stoma*, and *πυρ*, 'fire, fever,') Aphthæ—s. Aphtha, Aphthæ.

STOMATALGIA, (*stoma*, and *αλγος*, 'pain,') Stomalgia.

STOMATEL'CIA, *Stomatelco'sis*, (*stoma*, and *εκος*, 'an ulcer,') Ulceration of the mouth.

STOMAT'IA, *Stomatopath'ia*, (from *stoma*.) Disease of the mouth.

STOMATIC, (στοματικός,) *Stomat'icus*, (F.) *Stomatique*. A medicine used in diseases, &c., of the mouth. Dentifrices, masticatories, &c., are stomatics.

STOMATITE APHTHEUSE, Stomatitis, aphthous—*s. Charbonneuse*, Cancer aquaticus—*s. Conuenneuse*, Stomatitis, pseudomembranous—*s. Crèmeuse pultacée*, Aphthæ—*s. Follicleuse*, Stomatitis, aphthous—*s. Gangréneuse*, Cancer aquaticus—*s. Mercurielle*, see Salivation, mercurial—*s. Ulcéreuse*, Stomatitis, aphthous.

STOMATITIS, (*stoma*, and *itis*.) (F.) *Inflammation de la Bouche*. Inflammation of the mouth.

STOMATITIS, APH'THOUS, *S. follicular* or *ulcéreux*, *Em'physis Aph'tha*, *A. Adultor'um*, (F.) *Stomatite*, *S. aphtheuse* ou *follicleuse* ou *ulcéreuse*. An inflammation of the follicles of the mouth, constituting the aphthæ of the adult, which is generally accompanied by cephalic, gastric, and general disturbance. It may be either discrete or confluent, and requires general and local treatment, adapted to the case.

STOMATITIS, ERYTHÉMATOUS. Simple inflammation of the mouth.

STOMATITIS, EXSUDATIVA, Aphthæ—*s. Follicular*, *S. aphthous*—*s. Gangrenous*, Cancer aquaticus—*s. Materna*, Stomatitis of nursing women.

STOMATITIS, MERCURIAL, *S. mercurialis*, *Hydrargyrostomatitis*, *Hydrargyrostomatitis*. Inflammation of the mouth induced by mercury.

STOMATITIS OF NURSING WOMEN, *S. Nutricum* seu *Mater'na*, appears to be a variety of *S. aphthous*.

STOMATITIS NUTRICUM, Stomatitis of nursing women.

STOMATITIS PSEUDOMEMBRANOUS, (F.) *Stomatite conuenneuse*, *Diphthérie buccale*, *Pseudomembranous inflammation of the mouth*;—in its worst form, *Stomat'iphus*, (F.) *Fégérite*. Inflammation of the mouth accompanied by the exudation of a false membrane,—a disease which is rarely seen extensively except in large foundling establishments.

STOMATITIS, ULCEROUS, *S. aphthous*—*s. Vesiculosa* infantum, Aphthæ.

STOMATOCACE, (*stoma*, and *κακος*, 'evil.') Stomacace.

STOMATODYNIA, (*stoma*, and *δύσιν*, 'pain.') Stomalgia.

STOMATOGASTRIC, *Stomatogas'tricus*, *Stomogas'tric*, *Stomogas'tricus*, (*stoma*, and *γαστήρ*, 'the stomach.') Appertaining to the mouth and stomach—as the *stomatogastric* ganglia of the lower animals. See Encephalon.

STOMATOMALACIA PUTRIDA, (*stoma*, and *μαλακία*, 'softness.') Cancer aquaticus.

STOMATONECROSIS, (*stoma*, and *νεκρωσις*, 'death.') Cancer aquaticus, Stomacace.

STOMATOXOMA, (*stoma*, and *σῶμα*, 'an eating ulcer.') Stomacace.

STOMATOPANUS, (*stoma*, and *πᾶς*, 'a glandular tumour.') *Stomatophy'ma glandulosum*, *Panus Fau'cium*. Tumefaction of the glands of the mouth.

STOMATOPHY'MA, gen. *Stomatophy'matis*, (*stoma*, and *φύμα*, 'a swelling.') A swelling of the mouth.

STOMATOPHYMA GLANDULOSUM, Stomatopanus.

STOMATOPLAS'TIC, *Stomatoplas'ticus*, (*stoma*, and *πλασσω*, 'I form.') An epithet for the operation of forming a mouth where the aperture has been contracted from any cause; *Stomatoplas'ty*.

STOMATORRHAG'IA, *Stomatorrhæ'a*, *Hæmorrhag'ia o'ris* seu *fau'cium*, *Sanguinis profluvium ex O'ri*, *Hæmorrhoides O'ris*, *Stomato'sis*, *Stomorrhag'ia*, (F.) *Hémorrhagie buccale*, (*stoma*, and *ῥήγναι*, 'I break out.') Hemorrhage from the mouth.

STOMATORRHAGIA ALVEOLORUM, Phatnorrhagia—*s. Gingivarum*, Ulorrhagia.

STOMATORRHEA, (*stoma*, and *ῥέω*, 'to flow.') Stomatorrhagia.

STOMATOSCOPE, (*stoma*, and *σκοπεω*, 'I view.') Any instrument for keeping the mouth open, so as to permit the parts within to be inspected; *Speculum o'ris*.

STOMATOSEPSIS, (*stoma*, and *σηψις*, 'putrefaction.') Cancer aquaticus.

STOMATOSIS, (from *stoma*.) Stomatorrhagia.

STOMATOSPASMUS, (*stoma*, and *spasmus*.) Trismus.

STOMATYPHUS, (*stoma*, and *τυφίς*.) 'Stomatitis, pseudomembranous.

STOMENCEPHALUS, Stomocephalus.

STOMOCEPH'ALUS, *Stomenceph'alus*, (F.) *Stomocéphale*, *Stomenceph'ale*, (*stoma*, and *κεφαλή*, 'head.') A genus of monsters having two contiguous eyes, or a double eye occupying the median line; the nasal apparatus atrophied and forming a tube (*trompe*); the jaws rudimental; and the mouth imperfect or wanting.—(I. G. St. Hilaire.)

STOMOMANICON, (*stoma*, 'face,' and *μανος*, 'thin' [?].) *Platysma myoides*.

STOMORRHAGIA, Stomatorrhagia.

STONE BINDER, Osteocolla—*s. In the Bladder*, Calculi vesicali—*s. Crop*, Sedum—*s. Crop*, biting, Sedum—*s. Mint*, Cunila mariana—*s. Moss*, *Tripe de roche*—*s. Poek*, Aene—*s. Root*, Collinsonia Canadensis—*s. Sweat*, see Hyraceum.

STONES, THE, Testes.

STOOL, ([Sax.] *rcol*.) Dejection.

STOOLS, CALOMEL, see Calomel.

STOPPAGE, Stegosis.

STOPPING, Filling, Stegosis—*s. of the Belly*, Bowels, Guts, or Stomach; Constipation.

STORAX, Styraç—*s. Liquida*, Liquidambar styraciflua.

STO'REY'S WORM CAKES. An empirical preparation formed of Calomel ʒj, Jalap ʒi, Zingib. ʒij, Sacch. ʒij, Cinnab. antim. q. s. to colour them; Syrup. simpl. q. s. to make into cakes.

STORKBILL, Geranium maculatum.

STOUND, ([Icel.] *stunde*, 'to be in pain,') Stitch.

STOVE, *Æstua'rium*, *Stu'pha*, Old French, *Estuve*; a *Stew*, (F.) *Étuve*. A limited space, warmed artificially, and in which the air is slowly renewed. It is used for drying various substances, as plants, extracts, conserves, &c., or for taking vapour baths. In this case the *stew* or *stove* is said to be *wet* or *humid*; in the opposite case it is said to be *dry*, (F.) *sèche*. The latter, which is used by the Turks, is the *Py'ria*, *Pyriate'rium*, *Lacon'ium*, *Calida'rium* or *Calda'rium*, *Suda'tio concamera'ta*, Turkish, Eastern, or Roman bath. In the true Turkish bath, however, the air of the bathing apartment is densely charged with vapor, which the so-called Turkish bath, of recent introduction into England, is generally not. The former is their *Tepida'rium* or *Vapora'rium*, which see. These kinds of baths greatly excite the cutaneous transpiration, and are valuable agents in rheumatic and other affections.

STOX'S MEDICINE FOR HYDROPHOBIA, see Anagallis.

STRAB, (σπαζοι,) see Strabismus.

STRABILISMUS, Strabismus.

STRABISM'US, (σπαζισμος.) *Strabilis'mus*, *Parap'is* Strabismus, (from σπαζοι, 'twisted; one who squints,') *Strabos'itas*, *Loxophthal'mus*, *Distor'sic Lus'citas*, *I'vlo*, *Ilo'sis*, *Ilo'sis*, *Helo'sis*, *Squint'ing*, *Squint*, *Cruiseye*, (F: v.) *Doyle*, *Glee'ing*, (Sc.) *Shel'lie*, *Ske'ly*, (F.) *Strabisme*, *Vue louchée*, *Louchement*. Want of concordance of the optic axes. It may be dependent upon natural or acquired inequality in the action of the motor muscles of the eye; on a convulsive state of one of

those muscles; a difference in the sensibility of the two eyes; or a cerebral affection. In the treatment, if the disease seems to depend upon an excess of strength in one of the motor muscles, we must endeavour to give tone to that which is too weak by placing before the eye a mask having a small aperture on the side to which it is necessary to direct the pupil; or by wearing glasses provided with a piece of looking-glass on the side to which the eye is vitiously turned, so that the reflection may be inconvenient, and occasion the organ to be turned in an opposite direction; or by putting a black patch on the angle opposite to that to which the eye is turned. If, owing to weakness, the organ must be strengthened by being exerted alone, and leaving the other at rest. If the disease be symptomatic of cerebral affection, attention must, of course, be directed to the latter.

Strabismus may be *single*, or affect one eye only, or *double*, affecting both eyes, and it may be *convergent* or *divergent*, that is, with the ball drawn towards the nose, or away from it.

Those affected with Strabismus are called *Strab'i*, *Strabo'nes*, *Strabo'ni*, *Streb'li*, *It'i*, *It'ides*, *Squint'ers*, *Squint'eyed*, (Sc.) *Gle'yed*, (F.) *Louchees*, *Bigles*. A slight deviation is generally called a *Cast in the Eye*, and, sometimes, a *Cock-eye*.

STRABISMUS, CONVERGENT, see Strabismus—s. Divergent, see Strabismus—s. Double, see Strabismus—s. Single, see Strabismus.

STRABONES, (στραβωνες,) see Strabismus.

STRABONI, see Strabismus.

STRABOSITAS, Strabismus.

STRABOTOMY, *Strabotom'ia*, (στραβος, 'one who squints,' and *τομή*, 'incision.') The operation for the removal of squinting by the division of the muscle or muscles that distort the eyeball. A *Strabotomist* is one who practises strabotomy.

STRAEOTHEATH, see Death.

STRAIN, (σπασμ, [L.] *stringere*, (F.) *estreindre*, 'to constrict or squeeze,') Sprain.

STRAIN, *Colare*, (F.) *conler*. To pass decoctions, infusions, &c., forcibly through linen; also, to exert an effort. This is accomplished by fixing firmly the parts where the muscles to be exerted originate, in order that their full power may be developed.

STRAINING, Nisus.

STRAITS OF THE PELVIS, see Pelvis.

STRAMOINE, *Datura stramonium*.

STRAMONIA, *Datura stramonium*.

STRAMONIUM, (compounded from *strychnos* and *μανικός*, 'maniacal,') *Datura stramonium*—s. Fetidum, *Datura stramonium*—s. Majus album, *Datura stramonium*—s. Spinousum, *Datura stramonium*—s. Vulgatum, *Datura stramonium*.

STRAMONY, *Datura stramonium*.

STRANGALIDES, (σπαγγαλίδες,) pl. of *strangalis*, (from *σπασσω*, 'I compress,') Small, hard tumours, which form as in the breast when the milk does not find issue.

STRANGALION, (σπαγγαλιον,) Tubercle.

STRANGALIS, (σπαγγαλικ,.) Tubercle.

STRANGALISMUS, (σπαγγαλισμος,) Strangulation.

STRANGERS' FEVER, see Fever, strangers'.

STRANGULATIO, (σπασμ, 'I compress,') Orthopnoea, Strangulation—s. Uterina, Angone—s. Vulvæ, Hysteria.

STRANGULATION, *Strangula'tio*, *Strang'ula'tio*, *Strang'ula'tio*, *Strang'ula'tio*, *Strang'ula'tio*, (F.) *Strangulation*, *Strangulation*. State of a part too closely constricted. Thus we say that there is *strangulation* of an intestinal hernia, when the opening that gives passage to the portion of protruded intestine seriously intercepts the continuity of the digestive canal. In *Legal Medi-*

cine it means the forcible obstruction of the air passages, by a ligature or by the hand, for criminal purposes. See Suffocation.

STRANGULATOR, see Lycoides.

STRANGULLION, Strangury.

STRANGURIA, Strangury—s. Cruenta, Cystorrhagia.

STRANGURIE, Strangury—s. Menstruelle, Dysmenorrhœa.

STRANGURIOSUS. Affected with, or of the nature of, strangury.

STRANGURY, (σπαγγουρία,) *Strangu'ria*, *Paru'ria stillatitia*, *Dysu'ria*, *Uri'ne Stillheid'ium* seu *Subst'itium*, *Strangull'ion*, *Strangull'ion*, (F.) *Strangurie*, (σπαγγυ, 'to squeeze,' and *ουρον*, 'urine,') Extreme difficulty in evacuating the urine, which issues only drop by drop, and is accompanied with heat, pain, tenesmus at the neck of the bladder, &c. See Ischuria, and Retention.

STRANX, (σπαγγξ,) Gutta.

STRAP'PING, ([Sax.] *ƿƿnopp*, 'a strap,') The dressing of ulcers by means of straps or strips of adhesive plaster, linen, &c. Also, the plaster thus applied.

STRAPPING, WATER, see Waterdressing—s. Wet, see Waterdressing.

STRATHPEFFER, MINERAL WATERS OF. A sulphuretted water, at Strathpeffer, a few miles west from Dingwall, in Rossshire, Scotland. It contains sulphate of soda, sulphate of lime, chloride of sodium, sulphate of magnesia, and sulphohydric acid.

STRATIOTES, (στρατιωτης,) Achillea millefolium.

STRATUM BACILLATUM seu *bacillosum*, 'bacillary layer,' (from *sternere*, *stratum*, 'to spread out,') see *Tunica Jacobi*—s. Corneum, Epidermis—s. Germinativum, Proligerous disc, *Tache embryonnaire*—s. Malpighii, *Corpus mucosum*—s. Pigmenti, see Choroid—s. Proligerum, Proligerous disc.

STRAWBERRY, (strewberry,) *Fragaria*—s. Bush, *Euonymus Americanus*—s. Shrubby, *Rubus arcticus*—s. Tree, *Euonymus Americanus*—s. Wild, *Fragaria Virginiana*.

STREAK, PRIMITIVE, (from [Sax.] *ƿƿrice*, 'a line,') Groove, primitive.

STREAT'HAM, MINERAL WATERS OF. Streatham is a village near London. The waters are saline and cathartic.

STREBLI, (στρεβλοι,) see Strabismus.

STREBLOSIS, (στρεβλος, 'twisted,') Luxation.

STREMA, (στρεμμα,) Luxation, Sprain.

STREPITUS MORBUS seu AFFECTUS, (F.) *Maladie bruyante*, 'noisy disease.' A kind of flatulent disease, said by Sorbait to be not uncommon in the Austrian Alps, in which emphysematous tumours form on different parts of the body,—accompanied by sonorous excretion of gas by the mouth and anus.

STREPITUS, (from *strepere*, *strepitum*, 'to make a noise,') *Bruit*—s. Aurium, 'noise of the ears,' Tinnitus aurium.

STRETCHING, ([Sax.] *ƿƿteccan*, 'to stretch,') Pandiculation.

STRIÆ, (pl. of *stria*, 'a furrow,') see Vibices—s. Ciliares, Ciliary processes—s. Corneæ, *Tænia semicircularis*—s. Longitudinales *Lancisii*, see *Corpus callosum*—s. Medullares, see *Processes teretes*—s. Retinæ subjectæ *ligamento ciliari*, *Halo signatus*—s. Semilunares, *Tænia semicircularis*—s. Terminales, *Tænia semicircularis*—s. Transversales *Willisii*, see *Corpus callosum*.

STRIATED, *Stria'tus*, (F.) *Strié*, (from *stria*, 'a channel or groove,') Marked with small channels or grooves, or with lines of another colour.

STRIATED MUSCULAR FIBRE, see Muscular fibre.

STRIATUS, (*striare, striatum*, 'to make grooves or furrows,') *Cannelé*, Grooved, Striated; see *Corpora striata*.

STRIBILIGO, gen. *Stribili'ginis*, Efflorescence.

STRICTURA, Stricture — s. Ani, Stricture of the rectum — s. Intestini recti, Stricture of the rectum — s. Intestini recti spasmodica, Stricture of the rectum, spasmodic — s. Œsophagi callosa, Dysphagia conscripta — s. Œsophagi spasmodica, Œsophagismus — s. Pharyngis seu Œsophagi vera, Dysphagia constricta — s. Præputii, Phimosis — s. Urethræ, Urethrostenosis — s. Ventriculi, Stricture of the stomach.

STRIC'TURE, *Strictu'ra*, (*stringere, strictum*, 'to draw tight,') *Coarcta'tio*, *Stegno'sis*, (F.) *Rétrécissement*. A diminution or contracted condition of some tube or duct, as the Œsophagus, rectum, urethra, &c. This must be dilated by appropriate bougies. Strictures may, also, occur in the intestines: they are unmanageable, with the exception of those of the rectum, which admit of topical applications.

STRUCTURE OF THE ŒSOPHAGUS, Dysphagia constricta — s. of the Pharynx, Dysphagia constricta.

STRUCTURE OF THE RECTUM, *Strictu'ra A'ni* seu *Intesti'ni Rect'i*, *Archostegno'ma*, *Rectostenosis*, *Euthyenterostenoma*, *Archostegno'sis*, *Archostenosis*, *Proctoënceli'sis*, *Angusta'tio et Restrictio intesti'ni rect'i* seu *a'ni*, *Stenochoria intesti'ni rect'i*, *Proctostenosis organica*, occurs organically, and is a very serious affection; being generally dependent upon scirrhus.

STRUCTURE OF THE RECTUM, SPASMODIC, *Strictu'ra Intesti'ni Rect'i spasmodica*, *Obstructio Rect'i spasica*. An affection occurring in the nervous especially, which subsides spontaneously, after a longer or shorter continuance.

STRUCTURE OF THE STOMACH, *Gastrostenosis*, *Coarcta'tio* seu *Strictu'ra ventriculi*. See *Gastrostenosis cardiaca*, &c.

STRUCTURE OF THE URETHRA, Urethrostenosis — s. of the Urinary bladder, Cystostenochoria — s. Vesicæ urinariæ, Cystostenochoria.

STRICTUROTOMY, Urethrotomy.

STRIDOR DENTII, 'grinding of the teeth,' *Brygma*.

STRIE, Striated.

STRIGA ('a furrow') **CARTILAGINOSA CORDIS**, Isthmus of the fossa ovalis.

STRIGIS, *Strig'ilis*, (from *stringere*, 'to touch lightly,') *Sten'gis*, *Stelen'gis*. A fleshbrush. Also, an instrument, anciently used in the baths, for scraping off the sweat. The process was called *Strigillatio*, *Strigillatio*.

STRIPED MUSCULAR FIBRE, see *Muscular fibre*.

STROBILITES, (*στροβιλίτης*, from *στροβίλος*, *strobilus*, 'a cone of the pine,') Wine impregnated with the cones of the pine.

STROEMIA, (after H. Stroem, a Norwegian naturalist,) *Cadaba*.

STROKE, APOPLECTIC, (from [Sax.] *aptri-can*, 'to strike,') An apoplectic seizure.

STROKE, BACK, OF THE HEART, Impulse, diastolic.

STROKE, PARALYTIC. A sudden attack of encephalospinal paralysis.

STROKE, WATER, see *Hydrocephalus internus*.

STROMA, gen. *Stromatis*, *Stragulum*, *Stramentum*, (*στρομα*, 'a bed or couch,' from *σπρᾶννω*, *σπρᾶννωμι*, 'to spread out; to cover,') The bed or foundation texture of an organ,—as the stroma of the ovary. Applied, also, to the bed or base of any formation,—as the 'amorphous stroma' of scrofulous formations.

STRONGLE, *Strongylus*—s. *Géant*, *Strongylus gigas*.

STRONGULLION, *Strangury*.

STRONGYLUS, (from *στρογγυλος*, 'round,')

Hippocrates, and others, mean the *Ascaris lumbricoïdes* by this term. The *Strongylus Gigas*, *Eustrongylus gigas*, *Ascaris viscera'lis* seu *renal'is*, *Fusa'ria viscera'lis* et *renal'is*, (F.) *Strongle*, *S. géant*, is, however, distinct. It has an obtuse, flat head; mouth surrounded with six flattish papillæ: the whole bursa of the male truncated: the tail of the female rounded. It is sometimes met with,—five inches, a foot, a foot and a half, and even three feet long,—and from two lines to half an inch in diameter. It is occasionally found in the human kidney; rarely in other viscera, and still more rarely in the intestinal tube.

STRONGYLUS, (*στρογγυλος*,) *Teres*.

STROPHOS, (*στροφος*, from *στρεφω*, 'to twist,') *Tormina*.

STROPHULE, *Strophulus*.

STROPH'ULUS, *Lichen'asis stroph'ulus*, *Ebullit'io*, *Ecanthe'ma* seu *Ecor'mia strophulus*, *Red Gum*, *Gumrash*, *Red Gown*, *Tooth Rash*, (F.) *Strophule*, *Feux de dents*. It consists of an eruption of red, or sometimes whitish pimples, *White Gum*, *Milksports*, occurring in early infancy, chiefly about the face, neck, and arms, surrounded by a reddish halo; or interrupted by irregular patches of cutaneous blush. All the varieties under this genus arise in consequence of the extreme vascularity and irritability of the skin at this period of life, when the constitution is liable to be disturbed by irritation, either in the alimentary canal, gums or other parts. None of these eruptions are of any importance, and no medical treatment is usually necessary. If the stomach seems at all disordered, a little rhubarb and magnesia may be administered occasionally. Willan and Bateman reckon five species,—*Strophulus intertine'tus*, *Red gum or gown*; *S. al'bidus*, *white gum*; *S. confer'tus*, *Rank red gum*, *Tooth rash*; *S. volat'icius* or *Erythe'ma volat'icum*, and *S. can'didus*.

STROPHULUS SYLVESTRIS, *Ignis sylvaticus* — s. *Volaticus*, *Ignis sylvaticus*.

STROPHUS, (*στροφος*, from *στρεφω*, 'to twist,') *Tormina*.

STRUCTIO, (from *struere, structum*, 'to build,') *Structure*.

STRUCTURAL, *Structura'lis*. Of, or belonging to, structure.

STRUCTURE, *Structu'ra*, *Struc'tio*, *Conformatio*, *Catascœ'e*. The arrangement of the different tissues or organic elements of which animals and vegetables are composed. See *Tissue*.

STRUMA, gen. *Strumæ*, *Bronchocele*, *Scrofula*. *STRUMA ADIPO'sa*, *Obes'itas colli*. Fatty neck. Prominence of the neck from obesity.

STRUMA FUNGOSA, *Encephaloid*.

STRUMES, *Scrofula*.

STRUMEUX, *Scrofulous*.

STRUMOSIS CEREBRI, *Encephalophymata* — s. *Pulmonum*, *Tubercles of the lungs*.

STRUMOSUS, *Scrofulous*.

STRUMOUS, *Scrofulous*.

STRUTHIOPHAGUS, (*στρονθιον*, 'a young ostrich,' and *φάγω*, 'I eat,') *Struthiophagous* tribes still exist in some parts of Africa.

STRUTHIUM, (*στρονθιον*,) *Saponaria*.

STRUVE'S LOTION FOR HOOPING-COUGH, see *Lotion*, *Struve's*, for *hooping-cough*.

STRYCHNIA, (from *strychnos*,) *Strychni'na*, *Strych'nine*, *Strych'nin*, *Strychni'num*, *Strych'nium*, *Teta'nium*, *Vauque'lin*, *Tet'anin*. An alkaline principle; solid, crystalline, inodorous, bitter, and excessively poisonous, which has been found in *Strych'nia nux vom'ica*, *S. Colubri'na*, *S. Tienté*, *Ignat'ia amara*, *Cerbera There'tia*, and *C. Aho'rai*. (Nucis vomice rasur. ßiv: Calcis pulv. ʒij; Acid. muri. t. ʒij; Alcohol., Acid. sulphuric. dil., Liquor. anemonie, Carbon. animal. purif., Aquæ, aa q. s. Digest the nux vomica in two

gallons of water, acidulated with a fluidounce of muriatic acid, for 24 hours: then boil for two hours, and strain with expression through a strong linen bag. Boil the residuum twice successively in the same quantity of acidulated water, each time straining as before. Mix the decoctions, and evaporate to the consistency of thin syrup; then add the lime previously mixed with a pint of water, and boil for ten minutes, frequently stirring. Pour the mixture into a double linen bag, and having washed the precipitate well with water, press, dry, and powder it. Treat the powder repeatedly with boiling alcohol, until deprived of its bitterness; mix the liquors, and distil off the alcohol by means of a waterbath. Mix the residue with water, and having applied heat, drop in sufficient diluted sulphuric acid to neutralize and dissolve the strychnia; then add purified animal charcoal; boil for a few minutes, filter, evaporate, and crystallize. Dissolve the crystals in water, and add sufficient solution of ammonia to precipitate the strychnia. Lastly, dry the precipitate on bibulous paper. Ph. U. S. 1851. A somewhat modified form is in Ph. U. S. 1863.) The action—therapeutical and toxic—of strychnia on man and animals is exactly like that of the alcoholic extract of nux vomica. It is, however, more active. An eighth of a grain is sufficient to kill a large dog; and a quarter of a grain produces marked effects on the human body, when in health. It has been given in paralysis and other cases like nux vomica. Dose, one-sixteenth to one-twelfth of a grain.

STRYCHNIA, ACETATE OF, *Strychniæ acetat*—s. Hydriodate of, *Strychniæ hydriodas*—s. Iodate of, *Strychniæ iodas*—s. Iodide of, *Strychniæ hydriodas*.

STRYCHNIA, I'ODIDE OF IODHY'DRATE OF, (F.) *Jodure d'Iodhydrate de Strychnine*, is made by pouring a solution of ioduretted iodide of potassium into a solution of a salt of strychnia; treating the precipitate with boiling alcohol, and crystallizing. Used in the same cases as strychnia. Dose, a tenth to a sixth of a grain.

STRYCHNIA, MURIATE OF, *Strychniæ murias*—s. Nitrate of, *Strychniæ nitras*—s. Sulphate of, *Strychniæ sulphas*.

STRYCHNIA AND ZINC, DOUBLE IODIDE OF, Zinc and Strychnia, double iodide of.

STRYCHNIE ACETAT, *Strychnium seu Strychninum aceticum*, *Acetas strychnii seu strychnicus*, *Ac'etate of Strychnia or Strychnine*. (This salt is formed by the direct combination of strychnia with acetic acid.) It is given in the same cases as pure strychnia.

STRYCHNIE HYDRIODAS, *S. Iodhyd'ras*, *Strychnium hydroiodicum*, *Hydriodas seu Iodhyd'ras strychnicus*, *Hydriodate or Iodide of Strychnia*, is prepared by mixing a solution of iodide of potassium with a concentrated solution of acetate of strychnia.

STRYCHNIE IODAS, *Strychniæ Iodas*, *Strychnina Iodata*, *Strychnium seu Strychninum Iod'icum*, *Iodas Strychnii seu Strychnicus*, *Iodate of Strychnia or Strychnine*. (This salt may be formed by adding a concentrated solution of iodic acid to strychnia; treating with boiling alcohol; filtering, and leaving it to spontaneous evaporation.)

STRYCHNIE MURIAS, *Muriate of Strychnia*. (The product of the union of muriatic acid and strychnia.) Used in the same cases as strychnia.

Liquor Strychnia, *Solution of Strychnia* (Ph. B. 1864), is composed of Strychnia, gr. iv; Acid, hydrochloric, dil. ℥vj; Sp. rectif. ʒij; Aq. destillat. fʒvj. Dose, grt iv—viii.

STRYCHNIE NITRAS, *Strychnium nitricum*, *Nitrate of Strychnia or Strychnine*, (F.) *Nitrate ou Azotate de Strychnine*. (The product of the union of nitric acid and strychnia.) Used in the same cases as strychnia.

STRYCHNIE SULPHAS, *Strychnium sulphuricum*, *Sulphate of Strychnia or Strychnine*. (The product of the union of strychnia and sulphuric acid is admitted into the Ph. U. S. for 1863.) Has the same properties.

STRYCHNINA, Strychnia.

STRYCHNINE, Strychnia—s. Azotate of Strychnie nitras.

STRYCHNINUM, Strychnia.

STRYCHNINUM, see Strychnos nux vomica.

STRYCHNIUM, Strychnia—s. Aceticum, Strychniæ acetat—s. Nitricum, Strychniæ nitras—s. Sulphuricum, Strychniæ sulphas.

STRYCHNOMANIA. With the ancients, the delirium caused by the Strychnos or Atropa belladonna.

STRYCHNOS, (στροχνος), *Atropa belladonna*, *Solanum dulcamara*—s. Colubrina, see Strychnia, and S. Nux vomica—s. Guianensis, see Curare—s. Ignatii, *Ignatia amara*.

STRYCHNOS NUX VOMICA, *Nux Vomica*, *Metella canina*, *Kooch'la tree*. Nat. Ord. Loganiaceæ. Sex. Syst. Pentandria Monogynia. (F.) *Vomiquier*. A tree of the family *Strychnoides*, which grows in India, and the seeds of which have been long sold in the shops, under the names, *Nux Vomica*, *Vom'ic* or *Poison nut*, *Bach'elor's buttons*, &c., (F.) *Noix vomique*. For a long time, these seeds were used only for poisoning rats; but subsequently an alcoholic extract was prepared from them, which was administered in paralytic affections, in small doses. The dose of the ALCOHOLIC EXTRACT, *Extractum nucis vomicæ spirituosum seu nucis vomicæ resinosum seu nucis vomicæ alcoholicum* (Ph. U. S. 1863), (F.) *Extrait alcoolique de noix vomique*, is one or two grains. It produces a kind of tetanic convulsion in the paralyzed parts, when given to the requisite extent. It has, also, been administered in chronic diarrhoea and dysentery. When taken to such an extent as to induce toxic phenomena—*Strychnism*—it causes, at first, or when given in a smaller dose, greatly augmented excitability of the spinal nervous system, with slight transient paroxysms or spasms of the limbs, and hurried respiration. In larger doses, the excitability is still more developed, with paroxysms of prolonged general tetanic convulsions, laryngismus or contraction of the glottis, violent efforts of respiration, and asphyxia.

The following is the formula for the preparation of the EXTRACTUM NUCIS VOMICÆ ALCOHOLICUM of the Pharm. U. S. Take of *Nux vomica*, lbj; Alcohol, a sufficient quantity. Expose the *Nux Vomica* to steam till it is softened; then, having sliced and dried it, grind it into powder. Introduce it into an apparatus for displacement, and pour alcohol upon it gradually, until the liquid passes without bitterness. Distil off the greater part of the alcohol from the filtered liquor, and evaporate the residue to the proper consistency.

The bark is the *false angustura* or *false cusparia bark*; Rohun. See Brucea antidysenterica.

The wood of STRYCHNOS COLUBRINA, *Modira canina*, an Indian tree, which contains strychnia, was formerly considered a specific against the bites of venomous animals; and was recommended by Boerhaave in intermittents. It is the *Lignum Colubrinum*, (F.) *Bois de Couleuvre*.

Under the names στροχνον, and στροχνος, different kinds of nightshade are described by Dioscorides.

STRYCHNOS POTATORUM, 'strychnos of drinkers,' *Clearing nut*. The fruit of a large tree of Silhet, which, when ripe, is sold in the markets of India, and has been introduced into those of London, under the name *Nirmallee* or *Nirmillies*. One of them is well rubbed for a minute or two around the inside of a vessel containing water, which it purifies by causing the impurities, which

the fluid contains, to be deposited, it is supposed, by its albumen and casein.

STRYCHNOS PSEUDOQUINA, *Quina do Campo*, of Brazil, is used by the natives as a substitute for cinchona.

STRYCHNOS TIEUTÉ, see *Strychnia*, and Upá—*s. Toxicaria seu Toxifera*, see *Curare*.

STRYPHNA, *Astringents*.

STRYPHNOS, (στυφνός, 'sour, harsh,') *Acerb.*

STUDIOSUS MEDICINÆ, ('fond of medicine,') *Philatros*.

STUDIUM INANE, 'empty study,' *Aphelxia otiosa*.

STUDY, BROWN, *Aphelxia otiosa*.

STUFF, ([D.] *stoffe*, [F.] *estoffe*, *éttoffe*,) see *Medicament*, and *Falsifications*, table of.

STUFFING, *Cynanche trachealis*, *Filling*.

STULTITIA, *Fatuitas*.

STUMP, ([G.] *Stumpf*, [D.] *Stomp*,) (F.) *Moignon*. The part remaining from which a limb or other part has been amputated or removed in any manner. Also, the portion of a tooth remaining in the alveolus, after the corona has been removed, (F.) *Chicot*.

STUMPFOOT, *Kyllosis*.

STUNNED. (Old French, *Estonné*, *Étonné*, [L.] *attonitus*, 'astonished,' [Sax.] *ƿerman*, 'to stun,') An epithet applied to one who, in consequence of a fall or other accident, has received such a concussion of the brain as to deprive him, for a time, of his mental manifestations.

STUPA, (στυπα,) *Stupha*, *Stuppa*, *Stupeion*. Tow, used in certain surgical apparatuses and dressings, (F.) *Étoupe*. Also, a *Stupe*, that is, cloth or tow used in fomentations. A flannel or other article wrung out of hot water, plain or medicated, applied to a part, is a *stupe*. The act of applying such stupe is called *stu'ping*.

STUPE, *Stupa*.

STUPEFACIENTIA, (*stupere*, 'to be benumbed,' and *facere*, 'to make,') *Narcotics*.

STUPEFACTIES, *Narcotics*.

STUPEFACTIO, *Narcosis*.

STUPEFIANTS, *Narcotics*.

STUPEION, (στυπιον,) *Stupa*.

STUPEUR, *Stupor*.

STUPHA, *Stove*, *Stupa*.

STUPIA, *Tin*.

STUPIDITAS, (from *stupere*, 'to be benumbed,') *Idiotism*, *Imbecility*.

STUPING, see *Stupe*.

STUPOR, (*stupere*, 'to be benumbed,') *Epilepsia*, *Epilepsia*, *Consternatio*, (Sc.) *Swarff*, *Swerf*, (F.) *Stupeur*. Diminished activity of the intellectual faculties, often amounting to lethargy. It occurs in many affections, especially in the neuroses.

STUPOR, *Narcosis*—*s. Artuum*, *Obdormitio*—*s. Dentium*, *Hæmodia*—*s. Formicans*, *Formication*—*s. Mentis*, *Fatuitas*—*s. Vigilans*, *Catalepsy*.

STUPPA, (στυππα,) *Stupa*.

STUPRATION, *Stuprum*.

STUPRUM, *Rape*, *Stupra'tion*, (F.) *Défloration*, *Viol.* Forcible and illicit enjoyment of a married or unmarried female. When committed upon the latter, which is most commonly the case, it is also called, *Défloration*, *Déflora'tio*, *Devirginitio*, *Vitatio*, *Apanthis'mus*, &c. In judging whether rape has been committed under such circumstances, the state of the genital organs; the age of the individual; and the possibility of accidents and diseases of the parts, will have to be considered. It will be necessary, also, to be acquainted with the evidences of virginity; and to determine,—if signs of virginity no longer exist,—whether its loss be attributable to copulation, or to the introduction into the vagina of any other body than the male organ, &c. *Recent deflowering* can, of course, be much more readily

distinguished than that which has taken place some time.

STUTTERING, ([D.] *stotteren*, 'to stutter,' formed by onomatopœia,) *Balbuties*.

STYAN, ([Sax.] *ƿtizen*, from *ƿtisan*, 'to ascend, to rise,') *Hordeolum*.

STYE, *Hordeolum*.

STYGMATES. M. J. Cloquet has proposed this word, or *Stigmates*, (from *στυγμα*, pl. *στυγματα*, 'a mark,') to designate the marks, in form of cicatrices, which remain on the peritoneum after the obliteration of the neck of a hernial sac. They are, commonly, radiated folds, of a whitish appearance, more or less thick, and of a fibrous or fibrocartilaginous nature.

STYLET, (from *stylus*,) see *Specillum*.

STYLI, (pl. of *stylus*,) *Styloid process*.

STYLIENS (MUSCLES), see *Bouquet anatomique de Riolan*.

STYLIFORM, (*stylus*, and *forma*, 'shape,') *Styloid*.

STYLO, (from *στυλος*, 'a style,') in composition, the styloid process of the temporal bone.

STYLOCERATOHYOÏDEUS, *Stylohyoideus*—*s. Ceratoides*, *Stylohyoides*.

STYLOCHONDROHYOÏDE'US. A name given by Douglas and Albinus, to a fleshy fasciculus which the stylohyoid muscle sometimes presents, and which is attached to the lesser cornu of the os hyoides. It is the same fasciculus which Santorini calls *Stylohyoïdes nōvus*, *Stylohyoïdeus al'ter*.

STYLOGLOSSUS. A muscle situate at the anterior and superior part of the neck. It is narrow behind, much broader before; is attached to the styloid process of the temporal bone, and to the stylomaxillary ligament; and is inserted into the side of the tongue. It raises the base of the tongue and carries it backwards.

STYLOHYOÏDE'US, *Stylocer'atohyoïdeus*, *Styloceratoïdes*, *Stylohyoïdes ma'jor*, (F.) *Stylohyoïdien*. A muscle, situate at the superior, anterior, and lateral part of the neck. It is thin and narrow,—especially posteriorly: anteriorly, it opens to give passage to the tendon of the digastricus. It is attached to the styloid process of the temporal bone, and to the body of the os hyoides. It raises the os hyoides, and carries it backwards.

STYLOHYOID LIG'AMENT is a fibrous flattened cord, which passes from the styloid process to the lesser cornu of the os hyoides.

STYLOHYOID NERVE is the second branch given off by the facial.

STYLOMASTOID, *Stylomastoïde'us*. That which relates to the styloid and mastoid processes.

STYLOMASTOID AR'TERY arises from the posterior auricular, and, in some subjects, from the occipital. It enters the stylomastoid foramen; passes along the aqueduct of Fallopius, and spreads its ramifications on the mucous membrane of the tympanum, and in the mastoid cells and semicircular canals; and terminates by anastomosing with a branch of the *Arteria meningea media*, which enters by the *Hiatus Fallopii*. Murray has given the name—*Stylomastoid*—to the posterior auricular artery itself.

STYLOMASTOID FORA'MEN, *Foramen Stylomastoïdeum seu Aqueduc'tus Fallo'pii*, (F.) *Trou anonyme*, is situate at the inferior surface of the petrous portion of the temporal bone, between the styloid and mastoid processes. It terminates the aqueduct of Fallopius, and gives passage to the facial nerve.

STYLOMAX'ILLARY, *Stylomaxilla'ris*. That which relates to the styloid processes and the jaw.

The STYLOMAXILLARY LIG'AMENT is a ligamentous, flat cord, extending between the styloid process and the angle of the jaw.

STYLOPHARYNGE'US, (F.) *Stylo-thyro-*

pharyngien. A muscle situate at the anterior and lateral part of the neck. It is slender; thin above; attached to the styloid process of the temporal bone, and terminates in the parietes of the pharynx, as well as at the posterior margin of the thyroid cartilage. It raises the pharynx and carries it backwards. Professor Chaussier has united, under the name *Stylo-pharyngien*, the three constrictor muscles of the pharynx, the stylopharyngeus, and palatopharyngeus.

STYLOID, *Styliform*, (*στυλοειδής*.) *Styloï'des*, *Graphioï'des*, *Graphoi'des*, *Graphioï'des*, *Graphidoï'des*, *Belonoï'des*, (from *στυλος*, 'a style,' 'a peg,' 'a pin,' and *εἶδος*, 'shape.') Shaped like a peg or pin.

STYLOID CORNUA, see *Hyoides* os.

STYLOID PHOC'ESSES, *Pencil-like Processes*, *Processus Belonides* seu *Belemnoides* seu *Belonoï'des* seu *Graphioï'des* seu *Graphoi'des*, *Styl'i*, are, 1. A long and slender process of the temporal bone, to which are attached the styloglossus, stylopharyngeus, and stylohyoides muscles, and the stylohyoid and stylomaxillary ligaments. 2. Two slender and pyramidal processes at the inferior extremities of the radius and ulna.

STYLO-THYRO-PHARYNGIEN, *Stylo-pharyngeus*.

STYLOPH'ORUM DIPHYLLUM, (*στυλος*, 'a style,' and *φέρω*, 'to bear,') *Meconopsis diphyl'la*, *Chelidonium diphyl'um*, *Celandine* or *Hornpoppy*, *Bruiersroot*. An indigenous plant, of the family *Papaveracea*, growing from western Pennsylvania to Wisconsin, which flowers from May to August. The juice is fetid and narcotic. Infused in wine, it has been given internally in gravel; and has been applied externally to contusions, &c.

STYLOSTIXIS, (*stylus*, 'a style,' and *στιξίς*, 'the act of puncturing,') *Acupuncture*.

STYLUS, (*στυλος*, 'a stake, a style,') *Stilus*, *Sound*.

STYLUS EXCAVATUS, *Specillum excavatum*. A hollow sound.

STYMA, gen. *Sty'matis*, (*στυμα*, from *στυνέω*, 'to stiffen,') *Erection*, *Priapismus*.

STYMATOSIS, (from *styma*, and *osis*.) *Stimatosi*s.

STYMATATA, (*στυμμα*, gen. *στυμματος*, 'that which constricts or thickens,') The most solid ointments. Also, the ingredients proper for increasing their consistence.

STYPSIS, (*στυψίς*.) *Astriction*, *Constipation*.

STYPTERIA, (*στυπτηρία*.) *Alumen*, *Styptics*.

STYPTIC, (*στυπτικός*.) see *Styptics*.

STYPTIC, BINELLI'S, *Aqua Binellii* — s. *Brocchieri's*, *Aqua Brocchieri*.

STYPTIC, E'ATON'S. A name first given to *Helictius's Styptic*, when introduced into Great Britain. It is now usually made of calcined green vitriol, ʒj; *proof spirit*, tinged with a little oak bark, Oij.

STYPTIC, HELVETIUS'S, see *Ferrum tartarizatum*, and *Styptic*, *Eaton's*.

STYPTIC, PAGLIARI'S, *Hæmostatic*, *Pagliari's*.

STYPTIC, RUSPI'NI'S, *Liquor Stypticus Ruspi'ni*. A nostrum, said to consist of *Gallic acid*, a small quantity of *Sulphate of zinc* and *Opium*, dissolved in a mixture of *Alcohol* and *Rosewater*, which has had much reputation as an internal and external astringent.

STYPTICS, *Contrahent'ia*, *Constricti'va Medicamenta*, *Constringen'tia*, *Anastalt'ica*, *Catastalt'ica*, *Stypte'ria*, (F.) *Styptiques*, (*στυπτικά*, from *στυφω*, 'I constringe,') Sometimes used synonymously with astringent, but generally applied to a substance employed to arrest hemorrhage, — *Enæ'mon*. A mechanical styptic is one that arrests the flow of blood, by being applied immediately to the mouth of the bleeding vessels, or by inducing a coagulum, mechanically, in it. A chymical styptic

is one which coagulates, chymically, the blood around a bleeding orifice.

STYRAC'INUM O'LEUM. Oil obtained by boiling storax in good olive oil.

STY'RAX, gen. *Sty'raxis*, (*στυράξ*, gen. *στυράκος*.) A resin obtained from *Styrax officina'lis*, and *S. calamita*. It is the solid storax; — *officina'lis* storax. There are two kinds of storax met with: 1. *Red Storax*, *Gum Storax*, *Thusa Judæarum*, *Styrax rubra*, *Sty'raxis balsamum*, *Balsamum sty'raxis officina'lis*, which is obtained by incision from the *Styrax officina'lis*, and perhaps from the *Liquidambar Orientalis*. The purest is in tears; but it has lost some of its smell in drying. 2. *Common Storax*, *Styrax calamita*. That which has been received in reeds or vessels, and has had sawdust added immediately to thicken it. The odour of storax is fragrant; the taste aromatic. It is considered, like the other balsams, to be stimulant and expectorant, but is rarely used long.

PURIFIED STORAX, *S. colata* seu *purificata*, *S. preparata*, *Extractum sty'raxis*, is prepared by dissolving storax in alcohol; straining the solution; then distilling off the alcohol with a gentle heat, until the storax acquires the proper consistence. — Ph. U. S. Dose, gr. x. to xxx.

STYRAX ALBA, see *Myroxylon Peruiferum* — s. *Benzoin*, see *Benjamin* — s. *Calamita*, see *Styrax* — s. *Liquida*, *Liquidambar styraciflua* — s. *Officinalis*, see *Styrax*.

STYSIS, (*στυσις*.) *Erection*, *Priapismus*.

SUAVEOLENS, (*suavis*, 'sweet,' and *olens*, 'smelling,') *Beneolens*.

SUB (L), in composition, 'beneath,' or 'a lower degree.' Before c it becomes *suc*; before f, *suf*; before g, *sug*; before p, *sup*; before r, *sur*; and before s, *sus*. In some words, however, it is *su*, as in *supspiro*. The French *sous*, as a prefix, has the same signification.

SUBACIDUS, *Acidulous*.

SUBAC'TIO, *Chiro'sis*, *Chieiro'sis*, (*subigo*, *subactum* (*sub*, and *ago*), 'to subject, to prepare,') The operation of making plasters, extracts, &c., with the hand, or with the pestle and mortar.

SUBAFFECTIO, *Hydropathia*.

SUBAGITATRIX, (*sub*, and *agitare*, 'to agitate,') *Triade*.

SUBATLOIDE'US, *Infraatloïdeus*, (F.) *Sous-atloïdien*. That which is situate beneath the atlas or first cervical vertebra. Chaussier has given this name to the second pair of cervical nerves.

SUBAXOIDE'US, *Infraaxoïdeus*, (F.) *Sous-axoïdien*. That which is below the axis or vertebra dentata. Chaussier has given this name to the third pair of cervical nerves.

SUBAZOTAS BISMUTHICUS, *Bismuth*, *subnitrate* of.

SUBBORAS NATRICUM, *Borax*.

SUBCARTILAGINEUM, (*sub*, and *cartilago*, 'cartilage,') *Hypochondre*.

SUBCHLORIS CALCICUS, *Calcis chloridum*.

SUBCLA'VIAN, *Subclavia'vis*, *Subclavicula'ris*, (F.) *Sousclavier*, (*sub*, and *clavie*, 'clavicle,') That which is under the clavicle.

SUBCLAVIAN ARTERIES, (F.) *Artères sousclavières*, are situate at the upper part of the chest, and the lateral and lower parts of the neck. The right arises at the corresponding side of the trachea, and comes off from the arteria innominata or brachiocephalic trunk. The left leaves the aorta at the end of the arch. Both extend as far as the superior surface of the first rib, in the space between the scapular muscles, beyond which they take the name *axillary arteries*. The subclavian arteries run, for some distance, without sending off any branch. In the neighbourhood of the first rib, however, they give off, 1. *Upwards*, the *vertebral* and *inferior thyroïdal*. 2. *Downwards*, the *internal mammary* and *superior intercostal*. 3. *Out-*

wards, the *transverse cervical* or *posterior scapular*, the *superior scapular*, and the *posterior cervical* or *profunda*.

SUBCLAVIAN or **SUBCLAVIUS MUSCLE**, *Subclaviarius*, (F.) *Costo-claviculaire* (Ch.) *Sous-clavier*, is situate at the superior and anterior part of the chest. It is attached, by its inner extremity, to the superior surface of the cartilage of the first rib; and by its superior margin and outer extremity to the inferior surface of the clavicle. This muscle depresses the clavicle, and carries it forward. When the shoulder is fixed, it can raise the first rib.

SUBCLAVIAN VEINS, (F.) *Veines sous-clavières*, are continuations of the axillary, and extend from the inferior extremity of the scalenus anticus, in front of which they pass, to the vena cava superior, which they form by their union. The right subclavian vein is very short; the left, longer and broader. These veins receive the mouths of the internal and external jugulars, vertebral and superior intercostal. The left subclavian receives two veins more than the right,—the left internal mammary and left inferior thyroid. The thoracic duct opens into the left subclavian:—the corresponding great lymphatic vessel into the right.

SUBCLAVICULA, (*sub*, and *clavicula*, 'clavicle.') The first rib.

SUBCLAVICULARIS, Subclavian.

SUBCLAVIUS, Subclavian muscle.

SUBCREPITANT, (*sub*, and *crepitant*), see *Râle muqueux*.

SUBCRUENTUS, 'somewhat bloody;' (*sub*, and *cruentus*, 'bloody;') *Hyphe'mos*. That which has the colour and nature of blood. An epithet given to certain excreted substances, which are mixed with blood, or have the appearance of blood.

SUBCUTANÆUS, see *Cruræus*.

SUBCUTANÆOUS, *Subcutaneus*, (*sub*, and *cutis*, 'the skin;') *Intercutaneus*, (F.) *Sous-cutané*. That which is placed immediately under the skin. See *Hypodermic*.

SUBCUTANEOUS SECTION, *Hypodermatomy*.

SUBCUTANEOUS COLLI, *Superficialis colli*—s. *Male* (Nervus), Orbital nerve.

SUBDELIRIUM, *Typhomania*.

SUB'DITA, *Subditi'ta*, *subdere*, (*sub*, and *dare*, 'to give;') 'to put under;') *Pros'thetet*. Medicines introduced into some one of the natural apertures;—as suppositories, pessaries, &c.

SUBDITI'TIA, *Subdita*.

SUBDUCTIO, (*sub*, and *ducere*, *ductum*, 'to lead;') *Dejection*.

SUB'BER, gen. *Su'beris*, *Cork*:—the bark of *Quercus Su'ber* or *Corktree*, (F.) *Liège*. *Family*, *Amentaceæ*. *Sex. Syst.* *Monœcia Polyandria*, *Cork*, when burnt, is applied as a mechanical styptic to bleeding piles, &c. Surgeons use it for making tents; and it has been recommended to be worn as an amulet to suppress the secretion of milk!

SUBETH, *Coma*.

SUBFRONTALIS SUTURA, (*sub*, and *frons*, gen. *frontis*, 'the forehead;') A suture resulting from the articulation of the os frontis with the nasal process of the superior maxillary bone, and the proper bones of the nose.

SUBGRONDATION, *Depression*.

SUBHUMERATIO, (*sub*, and *humerus*, 'the shoulder;') *Catomismus*.

SUBIGITATRIX, (*sub*, and *agitare*, 'to agitate;') *Tribas*.

SUBINFLAMMATIO, *Subinflammation*—s. *Hepatic*, *Hepatitis*, *chronic*.

SUBINFLAMMATION, *Subinflammatio*, (*sub*, and *inflammatio*,) *Inflammatio'cula*, *Hypophlegma'cia*. A mild degree of inflammation, so slight as hardly to deserve the name *inflammation*. Broussais understood by the term, an augmenta-

tion in the vital phenomena of the vessels that convey white blood. There are, however, no such vessels. Lymphatic engorgements, scrofula, herpes, and cancer he considered *subinflammations*.

SUBINTRANTES FEBRES, (*sub*, and *intrare*, 'to enter;') *Communicantes febres*. Fevers primarily intermittent, whose paroxysms approximate, so that one paroxysm begins before that which preceded it has gone through its stages.

SUBJECT, (*subjicere*, *subjectum*, (*sub*, and *jacere*,) 'to place under;') *Cadaver*.

SUBJECTIVE SENSATIONS, see *Sensation*.

SUBJEE, *Bangue*.

SUBLIGACULUM, (*sub*, and *ligare*, *ligatum*, 'to bind;') *Truss*.

SUBLIGATURA, *Truss*.

SUBLIMATE, **CORROSIVE**, *Hydrargyri oxy-murias*.

SUBLIMATIO, *Enœorema*, *Sublimation*.

SUBLIMATION, *Sublimatio*, *Meteorismus*, *Hypoe'sis*, (*sublimare*, *sublimatum*, 'to raise on high;') An operation by which dry and solid matters are volatilized and condensed at the upper part of a subliming apparatus. Calomel, sal ammoniac, &c., are commonly obtained by sublimation. The product of sublimation is a *Sub'imate*, *Sublimatum*, (F.) *Sublimé*.

SUBLIMATUM CORROSIVUM, *Hydrargyri oxy-murias*.

SUBLIMATUS CORROSIVUS, *Hydrargyri oxy-murias*.

SUBLIME, *Enœorema*.

SUBLIME, *Sublimis*.

SUBLIME, see *Sublimation*—s. *Corrosif*, *Hydrargyri oxy-murias*.

SUBLIMIS, (F.) *Sublime*, 'high, elevated, superficial.' Anatomists have given the name *sublimis* to certain muscles, which are situate more superficially than their kindred muscles. These last they have called *profundi*. Of the former, the *flexor sublimis digitorum* is an example.

The French use the term *respiration sublime* for the respiration which is accompanied by considerable elevation of the ribs, and with separation of the alae nasi at the time of inspiration.

SUBLINGUA, (*sub*, 'a lower degree;' and *lingua*, 'tongue;') *Uvula*.

SUBLINGUAL, *Sublingualis*, (F.) *Sublingual*, (*sub*, and *lingua*, 'the tongue;') That which is seated under the tongue.

SUBLINGUAL ARTERY, with some, is the *Lingual*. With others, it is a branch given off by the lingual opposite the genioglossus muscle, which is distributed particularly to the sublingual gland, to the mylohyoid, and genioglossus muscles, &c.

SUBLINGUAL GLAND, *Glandula sublingualis* seu *Borholinia'na* seu *Rivinia'na*, is seated in the substance of the inferior paries of the mouth, beneath the anterior part of the tongue. It is smaller than the submaxillary, of which it frequently seems to be only a prolongation. It is oblong, flattened, amygdaloid, and is covered by the mucous membrane of the mouth, beneath which it projects. This gland has 10 or 12 excretory ducts—*Ductus Rivinia'ni* seu *Waltheria'ni*—some of which open beneath the frænum lingue, joining Wharton's duct, whilst others pierce the mucous membrane of the mouth separately. It has the same structure as the parotid.

SUBLINGUIUM, *Uvula*.

SUBLINGUAL, *Epiglottis*.

SUBLOBULAR, *Sublobularis*, (*sub*, and *lobulus*, 'a little lobe;') That which is placed under a lobe.

SUBLOBULAR VEINS OF THE LIVER. The veins in which the intralobular veins terminate. By

their union, the sublobular veins form the hepatic veins.

SUBLUXATIO, (*sub*, and *luxare*, 'to dislocate,') *Subluxation*. An imperfect luxation of a bone, see Luxation. Not unfrequently, used for a sprain.

SUBMAMMARY, *Submammar'is*, (F.) *Sous-mammaire*, (*sub*, and *mamma*, 'the breast,') Under the mamma or breast:—hence, 'submammary inflammation,' inflammation of the areolar tissue beneath the mamma.

SUBMAXILLARY, *Submaxilla'ris*, *Inframaxilla'ris*, (F.) *Sous-maxillaire*, (*sub*, and *maxilla*, 'the jaw,') That which is seated beneath the jaw.

SUBMAXILLARY GANGLION is a small nervous ganglion, situate opposite the gland. It seems to be formed by the superior branch of the Vidian nerve, and communicates with the lingual nerve of the inferior maxillary. Its filaments form a plexus which penetrates the submaxillary gland.

SUBMAXILLARY GLAND, *Maxillary Gland*, *Hypognath'aden*, *Hypossal'aden*, is a salivary gland, less than the parotid, seated at the inner side of the ramus and body of the inferior maxillary bone, in the triangular space between the two bellies of the digastricus. Irregularly ovoid, and flattened on three surfaces, it has the same aspect and organization as the other salivary glands. Its excretory duct is called *Wharton's Duct*, and terminates at the side of the frenum lingue, by a very small orifice. It is also called *Ductus inferior*.

SUBMENTAL, *Submenta'lis*, (F.) *Sousmental*, (*sub*, and *mentum*, 'the chin,') Situate under the chin.

SUBMENTAL ARTERY is furnished by the facial, near the base of the jaw. It passes forwards along the attachment of the mylohyoids, to which it furnishes branches that cross it to anastomose with those of the sublingual. Near the median line it bifurcates, to be distributed to the chin and muscles of the infrahyoid region.

SUBMENTAL VEIN, which accompanies the artery, opens into the labial.

SUBMER'SION, *Submer'sio*, (*sub*, and *mergere*, *mersum*, 'to plunge,') The act of plunging, or being entirely plunged, in a liquid.

Asphyxia by submersion, *A. Immer'sorum*, *Drowning*, is not owing to a certain quantity of liquid being introduced into the alimentary or air-passages; but simply to the interception of air, and of the respiratory phenomena. It is a case of suffocation, the signs of which, on dissection, are equivocal, without circumstantial evidence.

SUBMISSIO, (*sub*, and *mittere*, *missum*, 'to send,') Remission—*s. Cordis*, *Systole*.

SUBMUCOUS, *Submuc'ous*, (*sub*, and *mucus*.) Under a mucous membrane,—as the 'submucous areolar tissue.'

SUBMURIAS AMMONTACOHYDRARGYRICUS, *Hydrargyrum præcipitatum*.

SUBMUSCULAR, *Submuscula'ria*, (*sub*, and *musculus*, 'a muscle,') Seated beneath muscles or a muscular layer.

SUBOCCIPITAL, *Suboccipita'lis*, (F.) *Sous-occipital*, (*sub*, and *occiput*.) That which is situated under the occiput.

SUBOCCIPITAL NERVE, *Occipital nerve*.

SUBORBITAR, *Suborbita'rius*, *Infræorb'itar*, *Infræorbita'lis*, *Infræorb'itar'ius*, (F.) *Sous-orbitaire*, (*sub*, and *orbita*, 'the orbit,') That which is seated beneath the orbitar cavity.

SUBORBITAR ARTERY, *Infræorb'itar A.*, proceeds from the internal maxillary, from which it separates near the anterior and superior part of the zygomatic fossa. It enters the infraorbital canal, and passes through it, giving branches to the anterior and superior dental canal; issues by the foramen infraorb'itar'um; and divides, in the fossa

canina, into a number of branches, which lose themselves in the neighbouring parts. The infraorb'itar vein follows the same course.

SUBORBITAR CANAL, *Infræorb'itar Canal*, (F.) *Canal ou Conduit sous-orbitaire*, is a small canal, which runs obliquely through the substance of the inferior paries of the orbit. It begins, behind, by a simple gutter, and divides, anteriorly, into two branches, one of which descends into the anterior paries of the maxillary sinus, whilst the other passes out, in the canine fossa, by an orifice, called *Foramen infraorb'itar'ium*, (F.) *Trou sous-orbitaire*. The infraorb'itar artery and nerve pass along it.

SUBORBITAR FIS'SURE, *Infræorb'itar Fis'sure*, *Fissu'ra infraorb'ita'lis*, *Ri'ma cana'lis orbita'rii*. A fissure in the superior maxillary bone; the superior commencement of the suborbitar canal.

SUBORBITAR FOSSA, *Canine fossa*.

SUBORBITAR NERVES, *Infræorb'itar Nerves*, seem to terminate the superior maxillary. They issue from the infraorb'itar canal by the foramen of the same name, and spread out in a radiated manner in the fossa canina, to divide into superior or palpebral filaments, inferior or labial, internal or nasal, and external or buccal.

SUPERITONEAL, *Subperitonea'lis*, (*sub*, and *peritoneum*.) That which lies under the peritoneum.

SUPERITONEAL APONEUROSIS, *Fascia, superitoneal*—*s. Ganglia*, see *Spermatic ganglion*.

SUBPLACENTA, *Decidua membrana*.

SUBLEUR'AL, *Subpleura'lis*, (*sub*, and *pleura*.) Situate under the pleura.

SUBPOPLITEUS, *Popliteus muscle*.

SUBPUBIC ARCH, *Ar'cus subpu'bicus*, (*sub*, and *pubis*.) A strong arch of the superior pelvic aponeurosis, which completes the posterior orifice of the obturator or subpubic canal.

SUBPUBIC MEMBRANE, *Obturator membrane*.

SUBPUBIOFEMORALIS, *Abductor brevis*—*s. Prostaticus*, *Compressor prostate*.

SUBPUTRIS, (*sub*, and *putris*, 'putrid,') *Hypopaprus*.

SUBSCAPULAR, *Subscapula'ria*, (F.) *Sous-scapulaire*, (*sub*, and *scapula*.) That which is beneath the scapula.

SUBSCAPULAR FOS'SA, *Fos'sa subscapula'ris*. A considerable excavation at the anterior surface or venter of the scapula, in which the subscapularis is seated.

SUBSCAPULAR MUS'CLE, *Subscapula'ris Mus'cle*, *Immer'sua*, *Infrascapula'ria*, (F.) *Sous-scapulo-trochinien* (Ch.), *Porte-feuille*, *Muscle-sous-scapulaire*. A muscle situate in the above-mentioned fossa. It is flat, broad, and triangular; is attached, by its base, to the subscapular fossa and to the anterior lip of the edge of the spine of the scapula, and is inserted into the lesser tuberosity of the humerus. It carries the arm in rotation inwards. It can also depress it, and bring it towards the body.

SUBSCAPULAR NERVES are two in number. One arises from the brachial plexus above the clavicle; the other from the posterior aspect of the plexus within the axilla. They are distributed to the subscapularis muscle.

SUBSE'ROUS, *Subser'ous*, (*sub*, and *serous*.) Under a serous membrane,—as the 'subserous areolar tissue.'

SUBSIDENTIA, (*sub*, and *sedere*, 'to settle,') *Sediment*.

SUBSTANCE ÉMAILLÉE DES DENTS, *Enamel of the teeth*—*s. Horny*, see *Tooth*.

SUBSTANCE, HYALINE, *Cytoblastema*—*s. Intercellular*, *Cytoblastema*—*s. Vitree des dents*, *Enamel of the teeth*.

SUBSTANCE, WHITE, of *Schwann*. A hollow cylinder within the tubular nervous fibre, which

differs in composition and refracting power from the matter that occupies the centre of the tube.

SUBSTANTIA ADAMANTINA DENTII, 'adamantine substance of the teeth,' Enamel.

SUBSTANTIA COMPACTA, see Bone.

SUBSTANTIA FERRUGINÆA, *Lo'cus cinereus*. 'An ashecoloured substance produced by the existence of well-marked pigmentary matter in the floor of the fourth ventricle or rhomboid fossa.

SUBSTANTIA FILAMENTOSA DENTII, Enamel—*s. Grisea centralis*, see *Medulla spinalis*—*s. Hyalina*, *Cytoblastema*—*s. Nigra pedunculi*, see *Peduncles of the brain*—*s. Ossea dentium*, see *Tooth*—*s. Osteoidea*, see *Tooth*—*s. Perforata anterior*, *Locus perforatus anticus*.

SUBSTANTIA PERFORATA MEDII. When the pia mater has been removed from the space between the crura cerebri, termed the *intercrural* or *interpeduncular*, the surface appears cribriform, from the perforations of the numerous minute vessels which penetrate it. This is the *substantia perforata media* of Vieq d'Azyr.

SUBSTANTIA PERFORATA POSTERIOR, see *Tarini pons*—*s. Pulposa dentis*, *Dental pulp*—*s. Radiata*, *Stellula Verheyenii*—*s. Rhomboidea*, *Corpus dentatum*—*s. Spongiosa*, see *Bone*—*s. Spongiosa urethræ*, *Corpus spongiosum urethræ*—*s. Stellata serpentina*, *Stellula Verheyenii*—*s. Vitrea*, *Cytoblastema*—*s. Vitrea dentium*, Enamel.

SUBSTANTIVE, *Substantivus*, (*substare*, (*sub*, and *stare*), 'to stand under or subsist.')

An epithet applied by Dr. Paris to aliments which are nutritious, in contradistinction to *adjective* aliments, which are themselves not possessed of nutritive virtues, but impart to the digestive organs greater energy, so as to enable them to extract more nutriment from the substantive aliments. Meat, for example, is a *substantive* aliment; the condiment, as mustard, eaten with it, an *adjective* aliment.

SUBSTILLUM, (*sub*, and *stilla*, 'a drop,') *Stillicidium*.

SUBSTITUTED PRESENTATION, see *Presentation*.

SUBSTITUTIVE, *Substitutivus*, (*sub*, and *statuere*, 'to place.')

That may be substituted or put in place of another. An agent is said to be 'substitutive,' which—as in the case of nitrate of silver applied to inflammation of a mucous membrane—substitutes a temporary irritation for one tending to be more permanent. Such a mode of treatment is termed *substitutive medication*, (*F.*) *Médication substitutive*.

SUBSULTIO, (*subsilio*, *subsultum*, (*sub*, and *salio*), 'to make short leaps,') *Palpitation*.

SUBSULTUS TENDINUM, *Sal'tus ten'di-num*, *Clo'nus subsul'tus*, *Subsul'tus*, *Carpholog'ia spasmod'ica*, *Myopal'mus*, *Twitch'ing of the Tendons*, (*F.*) *Soubresaut des tendons*. Twitching, communicated to the tendons by the involuntary and instantaneous contraction of the muscular fibres. It is more observable at the wrist than elsewhere, and is an evidence of great cerebral irritability and debility when it occurs at an advanced period of fever.

The muscular agitations or twitchings observed in febrile diseases, especially of children, have been termed *Convulsibil'itas*, *Sclerotyr'be febrilis*, and *Irritabil'itas morbo'sa*.

SUBSURDITAS, (*sub*, and *surdus*, 'deaf,') *Deafness*.

SUBUBERES, (*sub*, and *ubera*, 'the breasts,') *Infan'tes eugen'tes*, *Suck'ing chil'dren*. Infants at the breast, in contradistinction to *Exu'beres*, *am'yacroi*, or those which have been weaned.

SUBUNGUAL, *Subun'guis*, *Subun'guical*, (*F.*) *Sous-ungéale*, (*sub*, and *unguis*, 'a nail,') Belonging to parts under the nail—as *subunguial exostosis*, (*F.*) *Exostose sous-ungéale*, an exostosis under the nail of the great toe especially.

SUBVERSIO STOMACHI, (*sub*, and *vertere*, *versum*, 'to turn,') *Vomiturition*.

SUBVERTEBRA, (*sub*, and *vertebra*,) *Sacrum*.

SUBVERTEBRUM, *Sacrum*.

SUBVOLA, (*sub*, and *vola*, 'the palm of the hand,') *Hypothenar*.

SUC, *Succus*—*s. de Citron*, see *Citrus medica*—*s. Gastricæ*, *Gastric juice*—*s. de Kino*, *Kino*—*s. du Limon*, see *Citrus medica*—*s. Médullaire*, *Marrow*—*s. Nourricier*, *Liquor sanguinis*—*s. Pancre'tique*, see *Pancreas*.

SUC, as a prefix, *Sub*.

SUCCA'GO, gen. *Succag'ginis*, (from *succus*, 'juice,') The inspissated juice of plants, robs, jellies, &c.

SUCCÉDANÉ, *Succedaneum*.

SUCCEDANEUM, *Succida'neum*, *Substitu'tum*, *Antiballon'emum*, *Sirroga'tum*, *Antemballon'emum*, *Quid pro quo*, (*succedere*, (*suc*, and *cedere*,) 'to go under, to come in place of another,') (*F.*) *Succédané*. An epithet for medicines that may be substituted for others possessing similar properties. Also, used, in dentistry, for an amalgam for filling teeth.

SUCCENTURIA'TUS, (*succenturiare*, (*suc*, and *centuria*,) 'to supply a soldier to fill a vacancy,') and hence, 'to substitute or put in the room of,') That which is a substitute for another—as *Reues succenturiati*, *Ventriculus succenturiatus*, &c.

SUCCHAR, *Saccharum*.

SUCCIDANEUM, *Succedaneum*.

SUCCIN, *Succinum*.

SUCCINCTURA, (*suc*, and *cingere*, *cinctum*, 'to gird,') *Diaphragm*.

SUCCINCTUS, *Diaphragm*.

SUCCINI RESINA, *Musk*, artificial.

SUCCIN'IC or **SUC'CINIC ACID**, *Ac'idum Sue'cini*, *A. Succin'icum*, *A'cor Succin'eus*, *Sat Sue'cini*, *Sal volat'ile Sue'cini*, has been considered antispasmodic and diaphoretic, in the dose of from five to twenty grains.

SUC'CINUM, *Su'cinum*, *Elec'trum*, *Am'bra*, *Am'bram*, *Am'par*, *Beren'ice*, *Am'ber*, *Yel'low Am'ber*, (*F.*) *Succin*, *Ambre jaune*, (from *succus*, 'juice,' because at one time supposed to exude from a tree.) This kind of fossil resin, which is found on the shores of the Baltic, is composed of a resinous matter, of essential oil, and an acid *sui generis*. It is inodorous, except when heated or rubbed; insipid; in fragments of a pale golden-yellow colour; transparent, and has a shining lustre. *S. g.* 1.08; insoluble in water, and slightly acted on by alcohol. It is used for affording its essential oil, obtained by destructive distillation,—the *O'leum seu Bal'samum Sue'cini*, (*F.*) *Huile de Succin*. This oil possesses stimulant, antispasmodic, diuretic, and rubefacient properties.

The *O'leum Sue'cini rectificatum*, *Rectified Oil of Am'ber*, is made as follows:—*Oil of Amber*, Oj; *water*, Ovj. Mix in a glass retort, and distil until Oiv of the water have passed with the oil into the receiver; separate the oil from the water, and keep it in wellstopped bottles.—(*Ph. U. S.*)

The *White am'ber*, *Leucelectrum*, (*F.*) *Ambre blanc*, is a variety.

SUCCINUM CINEREUM, *Ambergris*—*s. Griseum*, *Ambergris*.

SUCCION, *Sucking*.

SUCCISA, (from *succidere*, *succisum*, 'to cut,') from its indented appearance.) *Scabiosa succisa*—*s. Pratensis*, *S. succisa*.

SUCCOCOLLATA, *Chocolate*.

SUCCOLATA, *Chocolate*.

SUCCORY, *WILD*, *Cichorium intybus*, *Chironia angularis*.

SUC'COTASH, *Suck'ataash*. A term of Indian derivation. A favorite dish in New England, formed of green Indian corn and beans boiled together.

SUCCUBA, see Succubus.

SUC'CUBUS, (*succo*, and *cubare*, 'to lie.') Some authors have used this word synonymously with nightmare. Others mean by it, and by *Succuba*, a female phantom, with which a man in his sleep may believe he has intercourse—as *Incubus* has been applied to the male phantom, with which a female may dream she is similarly situate. See Incubus.

SUC'CUS, *Su'cus*, *Chy'lus*, *O'pos*, *Opis'ma*, *Chylis'ma*, *Echyl'o'ma*, *Apothlin'ma*, *Juice*, (F.) *Sue*, *Jus*, (from *sugere*, *suctum*, 'to suck.') The fluid obtained by pressing plants, flesh, &c.; *Su'cus expressus*; *Opulé* (official), *Opulide* (magistral), according to Chéreau. It is also applied to certain animal fluids, as *S. Gastricus*, &c.

SUCCUS ACACIÆ GERMANICÆ INSPISSATUS, see *Prunus spinosa*—*S. Acaciæ nostratis*, see *Prunus spinosa*—*S. Arborum*, *Sap*—*S. Aurantii*, see *Citrus aurantium*—*S. Ciutæ spissatus*, *Extractum conii*—*S. Cyrenaicus*, *Laserpitium*.

SUCCUS ENTER'ICUS, *S. Intestina'lis*, *Li'quor intestina'lis* seu *enter'icus*, *Intestinal Juice*. The fluid copiously secreted from the lining membrane of the small intestines.

SUCCUS EXPRESSUS, *Apothlimma*—*S. Gastricus*, *Gastro juice*—*S. Glycyrrhizæ*, *Extractum glycyrrhizæ*—*S. Hypocistidis*, see *Cytinus*—*S. Intestinalis*, *Succus entericus*—*S. Japonicus*, *Catechu*—*S. Limonis*, see *Citrus medica*—*S. Liquiritiæ*, *Extractum glycyrrhizæ*—*S. Nervæus*, *Nervous fluid*—*S. Nutritius*, *Chyle*—*S. Orni concretus*, see *Fraxinus ornus*—*S. Pancreaticus*, see *Pancreas*—*S. Spissatus aconiti napelli*, *Extractum aconiti*—*S. Spissatus atropæ belladonnæ*, *Extractum belladonnæ*—*S. Spissatus conii maculati*, *Extractum ciutæ*—*S. Spissatus hyoscyami nigri*, *Extractum hyoscyami*—*S. Spissatus lactucæ sativæ*, *Extractum lactucæ*—*S. Thebaicus*, *Opium*.

SUCCUS'SION, *Succus'sio*, *Hippocrat'ic Succus'sion*, (*succutio*, *succussum*, (*suc*, and *quatio*), 'I shake from beneath.') A mode of ascertaining the existence of a fluid in the thorax (*fluctuation by succussion*), by giving the body one or more slight shakes.

SUCCUSSION, *Ballottement*.

SUCHOR, *Saccharum*.

SUCINUM, *Succinum*.

SUCKATASH, *Succotash*.

SUCK'ING, *Suc'tio*, *Suc'tus*, *Bda'lis*, *Myze'sis*, (F.) *Succion*, (from *sugere*, *suctum*, 'to suck.') The act of drawing liquid into the mouth by forming a vacuum with the tongue acting as a piston, during inspiration.

SUCKING BOTTLE. A bottle so formed that, when filled with milk, it can be sucked from instead of the breast.

SUCKING CHILD, *Lactens*. A suckling.

SUCK'LE, *Lacta'ræ*, (F.) *Allaiter* ou *Alaiter*. To give suck. To nurse.

SUCKLING, *Lactation*, *Lonicera periclymenum*, *Sucking child*.

SUCRE, *Saccharum*—*S. Candi*, *Saccharum candidum*—*S. du Foie*, see *Saccharum*—*S. de Lait*, *Sugar of milk*—*S. de Saturne*, *Plumbi superacetat*—*S. Toræ*, *Penidium*.

SUCTIO, (*sugere*, *suctum*, 'to suck,') *Exsuctio*, *Sucking*.

SUCT'ION POWER. The force presumed to be exerted on the blood in the veins by the active dilatation of the heart. See *Derivation*.

SUCTO'RIA, (*sugere*, *suctum*, 'to suck,') Animals provided with mouths for sucking. Also, organs for sucking or adhesion;—hence termed *suctorial*—as *suctorial oscula*.

SUCTORIAL, see *Suctoria*.

SUCTUS, *Sucking*.

SUCUS, *Succus*.

SUDAM'INA, pl. of *Suda'men*, *Hydro'ata*, *Hi'dro'ata*, *Pop'ulæ sudora'les*, *Suffersu'ræ*, *Atha'xæ*,

Plan'ta noct'is, *Bo'a*, (from *sudor*.) Small vesicles which appear upon the skin, especially in the summer time, in hot countries, and in diseases attended with much sweating. It is a miliary eruption. See *Miliary fever*.

SUDAMINA, *Desudatio*, *Échauboulu'res*, *Lichen tropicus*.

SUDARIUM, *Achieolum*.

SUDATIO, *Ephidrosis*.

SUDATIUNCULA, (dim. of *sudatio*,) *Ephidrosis*.

SUDATOIRE, *Sudatory*.

SUDATORIA, *Hidrosis*—*s. Maligna*, see *Sudor Anglicus*.

SUDATORIUM, *Achieolum*.

SUDATORY, *Achieolum*. As an adjective, (F.) *Sudatoire*, any thing that excites, or is accompanied by, sweating.

SUD'OR, gen. *Sudo'ris*, *Hi'dros*, (F.) *Sueur*, 'sweat.' The product of cutaneous transpiration when visible on the surface of the body. It is colourless; of an odour more or less strong and variable; and of a saline taste. It is composed, according to Berzelius, of water, lactic acid, lactate of soda united with a little animal matter, chloride of sodium, and a little chloride of potassium, an atom of earthy phosphate, and some oxyd of iron.

SUDOR, *Miliary fever*.

SUDOR AN'GLICUS, *S. Anglica'nus*, *Hidron'ous*, *Hidrop'gra*, *Hidrop'yetus*, *Fe'bris* seu *Milia'ris sudato'ria*, *Sudor milia'ris*, *Mor'bus sudatorius*, *Ephem'era maligna* seu *sudato'ria* seu *Britan'nica* seu *Anglica pestilens*, *Pes'tis Britan'nica*, *Sudato'ria maligna*, *Sweat'ing Sick'ness*, *Sweat'ing Fe'ver*, (F.) *Suette*, *S. miliare*, *S. épidémique*, *Sueur Anglaise*. A very severe epidemic disease, characterized by profuse sweating, which appeared in England in 1486, and recurred at different times until about the middle of the 16th century. It was accompanied with coldness, excessive prostration of strength, palpitation, frequency and inequality of the pulse, &c., and terminated favourably or unfavourably in the course of 24 hours. The French have given the name *SUETTE DE PICARDIE*, *Suette des Picards*, *Fièvre suante*, *Milia'ris sudato'ria*, *Su'dor Picard'icus*, to an epidemic disease, which has appeared several times in that province, the principal symptoms of which were profuse sweats and a miliary eruption. The disease recurred in 1821, and has been described at length by M. Rayer, who considers it to consist of a simultaneous state of inflammation of various tissues, and proposes to class it with variola, scarlatina, and measles.

A form of it in which the perspiration was of a black colour was called *Sudor Ang'licus n'ger*, *Melanephidro'is*, *Black English Sweating sickness* or *fever*.

SUDOR COLLIQUATIVUS, *S. Profusus*.

SUDOR CRUEN'TUS, *S. sanguin'eus*, *Hæmatidro'is*, *Hæmatididro'is*, *Ephidro'is cruen'ta*, *Hæmor'rhag'ia per Cut'em*, (F.) *Hæmatidrose*, *Sueur de Sang*; *Sweating of blood*, *Bloody sweat*, *Hæm'orrhage from the skin*. Cutaneous perspiration intermixed with blood;—a very uncommon affection.

SUDOR DUL'CIS, *Saccharorrhœ'a cuta'nea*, *Ephidro'is sacchara'ta*. A discharge of saccharine matter by perspiration.

SUDOR LOCALIS, *Meridrosis*—*s. Miliaris*, *Miliary fever*, *Sudor Anglicus*—*s. Partialis*, *Meridrosis*—*s. Picardicus*, see *Sudor Anglicus*.

SUDOR PROFUSUS, *S. Colliquati'vus*. A profuse or immoderate sweat.

SUDOR SANGUINEUS, *Sudor cruentus*—*s. Universalis*, *Panhidrosis*.

SUDORIFER, *Sudoriferous*, *Sudorific*.

SUDORIFEROUS, *Sudor'ifer*, *Hydroph'orus*,

(*sudor*, and *fero*, 'I carry.') That which carries sweat—as the *sudoriferous* ducts or glands.

SUDORIFIC, *Hidrot'ic*, *Sudorificum*, *Sudor'ifier*, *Hidrot'icum*, *Hidrot'erium*, *Hidrotop'um*, *Hidropo'ticum*, (F.) *Sudorifique*, *Hidrotique*, (*sudor*, and *facio*, 'I make.') A medicine which provokes sweating. See *Diaphoretic*.

FOUR **SUDORIFIC WOODS**, *Quat'uor ligna sudorifica*, were, formerly, *guaiacum*, *perfumed cherry*, *sarsaparilla*, and *sassafras*.

SUDORIPAROUS, (*sudor*, and *pario*, 'to produce,') see *Perspiration*.

SUDORIS NULLITAS, 'nonexistence of sweat,' *Anidrosis*—s. *Privatio*, *Anidrosis*—s. *Suppressio*, *Ischidrosis*.

SUET, *Sevum*.

SUETTE, *Sudor Anglicus*—s. *Epidémique*, *Sudor Anglicus*—s. *Miltaire*, *Sudor Anglicus*—s. *de Picardie*, *Sudor picardicus*—s. *des Picards*, *Sudor picardicus*.

SUEUR, *Sudor*—s. *Anglaise*, *Sudor Anglicus*—s. *d'Expression*, *Expression*, *sweat of*—s. *de Sang*, *Diapedesis*, *Sudor cruentus*—s. *Urineuse*, *Uridrosis*.

SUF, as a prefix, *Sub*.

SUFFERING, (F.) *Souffrance*, (*suf*, and *fero*, 'to bear,') Any painful feeling produced by a general or local pathological condition.

SUFFERSURÆ, (*suf*, and *fervere*, 'to be hot,') *Sudamina*.

SUFFIMEN, *Suffimentum*.

SUFFIMENTUM, *Hypocarpnis'ma*, *Suffi'men*, *Thymia'ma*, *Suffi'tus*, *Suffumiga'tio*, *Suffumiga'tion*, (F.) *Parfum*; (*suffio*, *suffitum*, 'I perfume,') A perfume; *suffumige*; *fumigation*;

SUFFITIO, *Fumigation*.

SUFFITUS, *Fumigation*, *Suffimentum*.

SUFFLATIO, (*suf*, and *flare*, *flatus*, 'to blow,') *Puffness*.

SUFFOCATIO, *Suffocation*, *Orthopnoëa*—s. *Hypochondriaca*, *Hypochondriasis*—s. *Hysterica*, *Angone*—s. *Mulierum*, *Hysteria*—s. *Stridula*, *Cyananche trachealis*—s. *Uterina*, *Angone*.

SUFFOCATION, *Suffoca'tio*, *Præfoca'tio*, *Apnoë'ia*, *Asphy'ia*, *Melan'ema*, (*suffocare*, *suffocatum*, 'to stop the breath,' from *suf*, and *facies*, 'the jaws.' [?]) Death, or suspended animation from impeded respiration, whether caused by the inhalation of noxious gases, drowning, hanging, strangling, or smothering. The principal morbid appearances in such cases are:—the lungs of a deep-blue colour, with the blood extravasated in the air-cells; right auricle and ventricle filled with dark blood, as well as the neighbouring veins; lividity of the countenance, turgescence, and, perhaps, rupture of the vessels of the brain.

Treatment of suspended animation by suffocation in general. The patient must be conveyed into a room not too warm. Bloodletting must be used, if at all, with caution;—friction must be employed with salt, or warm flannels; stimulating fluids, in a dilute state, be poured into the stomach by means of a tube, and attempts be made to inflate the lungs. Laryngotomy, if necessary.

For the treatment suggested by Dr. Marshall Hall, and by Dr. Silvester, see *Ready Method*, and *Silvester's method*.

SUFFOCATION, *Phigopho'bia*, (F.) *Étouffement*, is used, as *strangling* is, for threatened suffocation. It means the state of dyspnoea and oppression experienced when a sufficient quantity of air does not enter the lungs, or when the chemical phenomena of respiration are imperfectly executed; as in cases of asthma.

SUFFOCATION DE MATRICE, *Hysteria*—s. *Utrine*, *Angone*, *Hysteria*.

SUFFRAGO, *Poples*.

SUFFUMIGATION, (*suf*, and *fumigare*, 'to produce smoke,') *Suffimentum*.

SUFFUMIGE, *Suffimentum*.

SUFFUMIGIUM, *Fumigation*.

SUFFUMINATIO, *Fumigation*.

SUFFUSIO, (*suf*, and *fundere*, *fusum*, 'to pour,') *Pseudoblepsia*—s. *Auriginosa*, *Icterus*—s. *Bilis*, *Icterus*—s. *Colorans*, *Chromopsia*—s. *Dimidians*, *Hemiopia*—s. *Exclarians*, *Oxyopia*—s. *Fellis*, *Icterus*—s. *Lentis crystallinæ*, *Cataract*—s. *Metamorphopsis*, *Metamorphopsia*—s. *Multipli-cans*, see *Diplopia*—s. *Myodes*, *Metamorphopsia*—s. *Nigra*, *Amaurosis*—s. *Oculi*, *Cataract*, *Pseudoblepsia*—s. *Oculorum*, *Metamorphopsia*—s. *Radians*, *Scotasma*—s. *Sanguinis*, *Echymoma*.

SUG, as a prefix, *Sub*.

SUGAR, *Saccharum*—s. *Barley*, *Penidium*, *Saccharum hordeatum*—s. *Black*, *Extractum Glycyrrhizæ*.

SUGAR CANDY, *Saccharum candidum*—s. *Cane*, see *Saccharum*—s. *Cane*, *Chinese*, see *Saccharum*.

SUGARCOATED. Covered with a layer of refined sugar, generally mixed with starch. Sugarcoated pills are prepared like the sugarplums of the confectioners, and are much used in modern pharmacy, especially when they are composed of nauseous ingredients. See *Dragée*.

SUGAR, *DIABETIC*, *Glucose*—s. *Fruit*, *Glucose*—s. *Grape*, *Glucose*—s. *Hepatic*, see *Saccharum*—s. *Honey*, *Glucose*.

SUGAR OF LEAD, *Plumbi superacetates*.

SUGAR, *LIVER*, see *Saccharum*.

SUGAR, *MAGNETIC*, *Saccharum Magnet'icum*. A quack preparation, said to be composed of sugar and cannabis. It is given in nervous cases.

SUGAR, *MAPLE*, *Saccharum acernum*, see *Acer saccharinum*.

SUGAR OF MILK, *Saccharum lact'is*, *Galac'to-saccharum*, *Lac'tose*, *Lactine* or *Lactin*, (F.) *Sucre de Lait*, *Sel de Lait*, is obtained from whey by evaporation and crystallization, s. g. 1.500. It possesses the demulcent properties of the sugars.

SUGAR, *MUSCLE*, *Inosite*.

SUGAR *PINE*, see *Arrowroot*.

SUGAR, *STARCH*, *Glucose*—s. *Urinary*, *Glucose*.

SUGARS, *VARIOUS*, see *Saccharum*.

SUGARBERRY, *Celtis occidentalis*.

SUGGILLATION, *Sugilla'tio*, *Li'vor*, *Æde'ma cruentum*, *Sugillation*, *Bruise*; (*sugillare*, *sugillatum*, 'to beat black and blue,') By most authors used synonymously with *ecchymosis* and *ecchymoma*. It is, also, applied to the spots, or *ecchymoses* which occur in consequence of intrinsic causes,—in disease and in incipient putrefaction. It is common in dead bodies; *Li'vor seu macula emortua'lis*. To this last case it is, indeed, restricted by some medico-legal writers; and an interesting question sometimes arises:—Whether the appearance has been induced before death or afterwards? Whether it is a case of *ecchymosis* or of *sugillation*? In the former, the extravasated blood will usually be coagulated; not so in the latter. It is often, however, difficult to form a correct judgment without reference to the history of the case.

SUGGRUNDIUM, (*sug*, and *gradior*, 'to go,') The eave of a house.

SUGGRUNDIUM OCULI, *Superciliary arch*—s. *Superciliorum*, *Frontal protuberance*.

SUICIDA, *Suicide*.

SUICIDAL INSANITY, *Suicide*.

SUICIDE, *Suici'dium*, (*sui cædes*, 'murder of one's self,') The act of killing one's self. *Autochi'ria*, *Autoctonia*, *Idiotonia*, *Autophonia*, *Suicide* is very frequently the result of disease of the mind; the individual seeming, at times, to be irresistibly impelled to selfdestruction;—*Melancho'lia Suicid'ium seu autochi'ria*, *Suici'dal In-*

san'ity. Also, one who commits selfmurder; *Suicida*, *Autochir*.

SUIE, (Gael. *suithe*.) Fuligo.

SUIF, *Sovum*.

SUITES DE COUCHES, 'sequels of labour,' *Lochia*.

SULCATUS, (from *sulcus*.) *Cannelé*, Grooved.

SULCI, plural of *Sulcus*.

SULCI CILIARES, Halo signatus.

SULCULUS LABII SUPERIORIS, (dim. of *Sulcus*.) *Lacuna Labii superioris*.

SULCUS, (akin to *δακος*, 'a furrow,') *Semicanal'is*, *Semicanal'ulus*. A furrow. A groove. (F.) *Sillon*. Anatomists have given this name to grooves on the surface of bones and other organs. See *Anfractuosus*.

French writers on anatomy often use the term *Sillons* for the grooves which lodge the arteries and creep on the surface of bones; whilst those that receive veins are called *gouttières*, 'gutters.' Also, the vulva.

SULCUS ANTEROPOSTERIOR JECORIS, see *Liver* — *s. Labii superioris*, *Lacuna Labii superioris*.

SULCUS, LAT'ERAL, ANTE'RIOR, OF THE SPI'NAL MAR'ROW. A mere trace on the marrow, marked only by the attachment of the filaments of the anterior roots of the spinal nerves.

SULCUS, LAT'ERAL, POSTE'RIOR, OF THE SPI'RAL MAR'ROW, corresponds with the attachment of the posterior roots of the spinal nerves.

SULCUS RETROGLANDULARIS, see *Glans* — *s. Sinus lateralis*, *Fossa sigmoidea* — *s. Transversalis*, see *Liver* — *s. Umbilicalis*, see *Liver*.

SULFATE, see *Sulphuric Acid*.

SULFIDUM CARBONII, *Carbonis sulphuretum*.

SULFITE, see *Sulphurous acid*.

SULFUR, *Sulphur*.

SULFURATION, see *Vulcanite*.

SULFURE, see *Sulphur* — *s. d'Antimoine*, see *Antimonium* — *s. de Carbon*, *Carbonis sulphuretum*.

SULFUREUX, *Sulphureous*.

SULPHAS, see *Sulphuric acid* — *s. Aluminaris*, *Alumen* — *s. Ammoniacæ Cupratus*, *Cuprum ammoniacum* — *s. Cadmiceus*, *Cadmii Sulphas* — *s. Kalicoaluminicum*, *Alumen* — *s. Natriæ*, *Soda*, sulphate of — *s. Zincicum*, *Zinci sulphas*.

SULPHUR, gen. *Sulphuricis*, *Sul'fur*, *Sco'rith*, *The'lon*, *Thi'on*, *A'verich*, *Ter'ra folia'ta*, *Fu'mus cit'rinus*, *A'quila*, *Brim'stone*, (F.) *Soufre*. *Na'tive Sulphur* is called *Rock* and *Black Sulphur*, *Sul'phur vi'vum* seu *u'grum*. Sulphur, as we meet with it, is in rolls, *Sulphur in rot'ulis* seu *rotu'dum*, *Roll Brim'stone*. It is a volcanic production, and is obtained, in large quantities, from *Solfatara*, in Italy. *Roll Sulphur* is impure sulphur, melted, and run into moulds. It is unpleasantly inodorous, when heated or rubbed; insipid; solid, and brittle. S. g. 1.99. Fusible at 226°; volatilizable by heat, and condensing unchanged.

When sulphur is combined with a simple body or a compound radical it forms a *sulphuret*, (F.) *Sulfure*.

SULPHUR ANTIMONIATUM, *Antimonii sulphuretum præcipitatum* — *s. Caballinum*, see *Sulphur sublimatum* — *s. Carburet* of *Carbonis sulphuretum* — *s. Chloratum*, *Sulphur*, chloride of.

SULPHUR, CHLO'RIDE OF, *Sul'phuris Chlo'ridum* seu *Dichlo'ridum*, *Sulphur chlora'tum*, *Hypochlo'retum sulphuro'sum*, *Chlo'rum hypersulphura'tum*, *Dichlo'ride* or *Protechlo'ride* or *Hy'pochlo'ride* or *Subchlo'ride of Sulphur*, *Bisulph'uret of Chlo'rine*, is obtained by passing dry chlorine gas over washed and dried flowers of sulphur, until they are for the most part dissolved, and distilling the decanted fluid from the excess of dissolved sulphur. One drachm of the liquid to an ounce of hard has been used in lepra, psoriasis, and other

chronic cutaneous diseases. Ten drops, dissolved in ether, have been given in adynamic fever.

SULPHUR, DICHLORIDE OF, *Sulphur*, chloride of — *s. Hypochloride of Sulphur*, chloride of — *s. Iodatium*, *Sulphuris iodium* — *s. Iodide of Sulphuris iodium* — *s. Ioduret of Sulphuris iodium* — *s. Liver of Potassæ sulphuretum*.

SULPHUR LO'TUM, *Sulphur sublima'tum lo'tum*, *Wash'ed Sulphur*, *Magiste'rium Sul'phuris*, *Sulphuris Flo'res lo'ti*, is prepared by pouring boiling water on sublimed sulphur, so as to wash away any acid it may contain, and then drying it.

SULPHUR NIGRUM, *Sulphur vivum*, see *Sulphur sublimatum*.

SULPHUR PRÆCIPITA'TUM, *Lac Sul'phuris*, *Milk of Sulphur*, *Præcip'itated Sulphur*, (F.) *Crème de Soufre*, is prepared by boiling sublimed sulphur and lime together, and adding muriatic acid to precipitate the sulphur.

SULPHUR, PROTOCHLORIDE OF, *Sulphur*, chloride of — *s. Subchloride of Sulphur*, chloride of.

SULPHUR SUBLIMA'TUM, (Ph. U. S. 1863,) *Sulphur*, *Subl'med Sulphur*, *Flo'wers of Sulphur*, (F.) *Soufre sublimé*, *Fleurs de Soufre*, is the sulphur of commerce, sublimed in close vessels.

Sulphur is possessed of stimulant, laxative, and diaphoretic properties. As a laxative, it is used in hemorrhoidal and other affections. In the itch, applied externally and taken internally, it has been considered as a specific, and it is a valuable remedy in most cutaneous affections that are local. Dose, $\mathfrak{z}$ ss to $\mathfrak{z}$ j.

The faeces or dregs, left in the purification or sublimation of sulphur, are called *Horse Brim'stone*, *Dregs of Sulphur vi'vum*, *Sulphur caballi-num* seu *vivum*, *Sul'phuris vi'vi recremen'ta*. They are very impure, and are only used externally.

SULPHUR, VEGETABLE, *Lycopodium* — *s. Wort*, *Peucedanum*.

SULPHURATUS, *Sulphureous*, *Sulphurated*.

SULPHURO'US, *Sulphuro'sus*, *Sulphur'ius*, *Sulphura'tus*, *Sulph'uretted*, (F.) *Sulfureux*. Possessing the properties of, or relating to, or containing sulphur.

SULPHURET, see *Sulphur*.

SULPHURETTED, *Sulphureous* — *s. Hydrogen*, see *Hydrogen*, sulphuretted.

SULPHURETUM AMMONIACÆ, *Ammoniacæ sulphuretum* — *s. Ammoniacæ hydrogenatum*, *Liquor fumans Boyleii* — *s. Kalicum*, *Potassæ sulphuretum* — *s. Lixivæ*, *Potassæ sulphuretum*.

SULPHURIC or *SULPHURIC ACID*, *Ac'id'idum Sulphur'icum* seu *Vitri'ricum*, *O'leum Vitri'oli*, *Spir'itus Vitri'oli* seu *Vitri'oli for'tis*, *Ac'id'idum vitrioli'cum*, *A. Vitri'oli dephlogistica'tum*, *Oil of Vitriol*, *Spir'it of Vitriol*, *Vitrioli'c Ac'id*, *A'cor* seu *Spir'itus Sul'phuris*, (F.) *Acide Sulphurique* ou *Sulfurique*. Sulphuric acid is inodorous; has a strong acid taste, and is corrosive. S. g. 1.843, (Ph. U. S.) It is escharotic, stimulant, and rubefacient. Mixed with lard, it is sometimes used in local pains, and certain cutaneous affections. United with bases it forms *sulphates*, (F.) *Sulfates*.

SULPHURIC ACID, AROMAT'IC, *Ac'id'idum Sulphur'icum Aromat'icum*, *A. vitrioli'cum aromat'icum*, seu *vitriolicum alcoho'læ aromaticum*, *Elia'ir vitrioli seu vitrioli aromaticum* seu *vitrioli Mynsich'ti* seu *vitrioli Edinburgen'sium* seu *vitrioli cum tinctu'râ aromaticâ* seu *vitrioli dulci seu acidoaromat'ici*, *Alcohol cum aromat'ibus sulphuricatus*, *Tinctu'ra ac'id'id sulphurici*, is formed of *Sulphuric acid*, f $\mathfrak{z}$ ijss; *Ginger*, bruised, $\mathfrak{z}$ j; *Cinnamon*, bruised, $\mathfrak{z}$ iss; *Alcohol*, Oij. — (Ph. U. S. 1842.) In the last edition (1863), it is made of *Sulphuric acid*, f $\mathfrak{z}$ vj; *Ginger*, in coarse powder, $\mathfrak{z}$ j; *Cinnamon*, in coarse powder, $\mathfrak{z}$ iss; *Alcohol*, q. s. Mix the acid and alcohol, and allow them to cool. Mix the cinnamon and ginger and, in a percolator, pour alcohol gradually until a pint has passed. Then mix the diluted acid and the

tincture. Dose, grt. x to xxx. Its properties are like those of the next preparation, than which it is more frequently employed, being more agreeable.

SULPHURIC ACID, DILUTED, Ac'idum Sulphuricum Dilutum, A. vitriolatum dilutum, Elix' vitrioli, Spiritus vitrioli ac'idus Vaglerii, Elix'ir de Vitriol, (F.) Acide sulfurique diluée, is formed, according to the Pharmacopœia of the United States, (1842,) by adding *Aq. destillat.* f 3xliij. to *Acid. sulph.* f 3j. According to the last edition (1863), it consists of *Acid. sulphur.* 3ij; *Aqua destillat.* q. s., to make together a pint. It is tonic, astringent, and refrigerant. Dose, grt. x to xx, largely diluted. When used as a gargle, half a drachm may be added to 5vj of water.

Very dilute sulphuric acid was formerly termed *Phlegma vitrioli*.

SULPHURIS CARBURETUM, Carbonis sulphureum.

SULPHURIS CHLORIDUM, Sulphur, chloride of—a. Dichloridum, Sulphur, chloride of.

SULPHURIS IODIDUM, S. Ioduratum, Sulf'p'ur Ioduratum, Ioduret or Iodide of Sulphur, (F.) Soufre Ioduré, Iodure de Soufre. It is thus directed to be prepared in the Pharmacopœia of the U. S. (*Iodin.* 3iv; *Sulphur.* 3j.) Rub the iodine and sulphur together in a glass, porcelain, or marble mortar, until they are thoroughly mixed. Put the mixture into a matrass, close the orifice loosely, and apply a gentle heat, so as to darken the mass without melting it. When the colour has become uniformly dark throughout, increase the heat so as to melt the iodide; then incline the matrass in different directions, in order to return into the mass any portions of iodine which may have condensed on the inner surface of the vessel: lastly, allow the matrass to cool; break it, and put the iodide into bottles, which are to be well stoppered.

SULPHURIUS, Sulphureous.

SULPHUROSUS, Sulphureous.

SULPHUROUS, Sulphureous.

SULPHUROUS ACID, Ac'idum sulphuratum seu sulphuris colat'ile seu Sulphurosum seu Sulfuratum, Spiritus sulphuris seu sulphureus volatilis seu sulphuris per camp'nam seu vitrioli phlogisticatus, Gas sulphuris, (F.) Acide Sulfureuse, Air Acide Vitriolique. (Obtained by treating mercury with concentrated sulphuric acid, or by burning sulphur.) *Acidum sulphureum, Sulphurous acid* prepared by treating sulphuric acid with charcoal by means of heat: receiving the acid into distilled water, has been admitted into the Pharmacopœia U. S. for 1863. It is sometimes used in the gaseous,—at others, in the liquid, state. Sulphurous acid gas is used to disinfect clothes, confined air, letters coming from infected places, &c. It is also exhibited, with great success, in the form of fumigation, in the treatment of certain chronic cutaneous affections, and of sciatic and rheumatic pains in general. If respired pure, it acts in the manner of violent irritants; and is not long in producing death.

United with bases, it forms *sulphites*, (F.) *Sulfites*.

SULTZ-MALT, MINERAL WATERS OF. Sultz-malt is a village in the department of the Upper Rhine; where are several sources of cold, acidulous waters, which seem to contain carbonic acid, subcarbonate of soda, subcarbonate of lime, sulphate of lime, and a little bitumen. They are refrigerant, aperient, and alterative.

SUMAC, Rhus coriaria—s. des Corroyeurs, Rhus coriaria—s. Vénéneux, Rhus toxicodendron.

SUMACH, Rhus coriaria—s. Chinese, Ailanthus glandulosa—s. Mountain, Rhus copallina—s. Poison, Rhus venenata—s. Stag's horn, Rhus typhina.

SUMBUL. The name of a root, *Sumbuli ra-*

dia, Jitumani'si, Muskroot, presumed to belong to the Umbellifera family, much employed in Russia and Germany against cholera. Its exact botanical history is not determined. It contains a strongly odorous principle like that of musk; and is regarded as an antispasmodic.

SUMBULI RADIX, Sumbul root.

SUMEN, (Sanguin, from *sugere*, 'to suck;') formerly the breast,) Hypogastrium.

SUMMER COMPLAINT. A popular name in the United States for diarrhoea occurring in summer. It is often, also, made to include dysentery and cholera infantum. With some it means cholera infantum only.

SUMMER DISEASES, see Estival—s. Plant winter worm, *Sphæria Sinensis—s.* Rash, *Lichen tropicus.*

SUMMITAS CUTIS, (summa, for *superius*, from *superus*, 'highest,') Epidermis.

SUMMUS HUMERUS, Acromion.

SUN BURN, Ephelides—s. Dew, *Drosera rotundifolia—s.* Flower, false, *Helenium autumnale—s.* Flower, swamp, *Helenium autumnale—s.* Flower, tickweed, *Coreopsis trichosperma—s.* Flower, turpentine, *Silicium perfoliatum—s.* Pain, see *Hemierania—s.* Rash, *Lichen—s.* Stroke, *Coup de Soleil—s.* Stroke, Egyptian, *Coup de Soleil.*

SUP, as a prefix, Sub.

SUPALE, see Valeriana.

SUPELLECTILE DIOGENIS, Palm.

SUPELLEX ANATOMICA, (supellex, supellectilis, 'household goods,') Museum anatomicum—s. Diogenis, Palm.

SUPER, (L., in composition, 'above,')

SUPERBE, ([L.] *superbus*, 'proud,') Rectus superior oculi.

SUPERBUS, 'proud,') Rectus superior oculi.

SUPERCHLORIDUM FORMYLICUM, Chloroform.

SUPERCILIARIS, Corrugator supercillii, Superciliary.

SUPERCILIARY, Superciliaris, (from *supercilium*,) That which relates to the supercilia or eyebrows. (F.) *Sourcilier ou Supercilier.*

SUPERCILIARY ARCHES, Superciliary Ridges, Arcus superciliaris, Superciliaris, (F.) Arcades sourcilieres, are two slightly projecting apophyses, situate at the anterior surface of the os frontis, above the superior margin of the orbits. They are separated by the nasal fosse, and covered by the superciliaris muscle, to which they give attachment.

SUPERCILIARY RIDGES, Superciliary arches.

SUPERCILIUM, Ophrys, Ophrya, Falx, (F.) Sourcil, (super, and cilium, 'the tarsal cartilage,') The eyebrows, (De.) Brows, (Prov.) Eyebrows, are two arched eminences,—convex above, and more or less prominent, according to the individual,—which lie on the superciliary arch of the os frontis, above the orbits, and extend from the sides of the root of the nose to the temple. Their inner extremity or *head* is more prominent than the outer or *tail*. The eyebrows are covered with short and stiff hairs, which are directed, obliquely, from within outwards, and are, commonly, of the colour of the hair. The skin in which these hairs are implanted, rests on the *orbicularis palpebrarum*, the *frontalis*, and *superciliaris* muscles, which move the eyebrows in different directions. The eyebrows serve as an ornament to the countenance; and by covering the eye defend it from the impression of too strong a light. They direct the perspiration also from the eye.

SUPERCILIUM ACETABULI. The edge of the cotyloid cavity of the os innominatum.

SUPERCONCEPTION, see Superfætation.

SUPEREVACUATIO, Hypererisis.

SUPERFUNDATIO, see Superfætation.

SUPERFÉTATION, Superfætation.

SUPERFIBRINATION, Hyperinosis.

SUPERFICIALIS COLLI, (from *superficies*, 'surface,') *Sabota'neus Col'li*. A nerve formed by communicating branches from the second and third cervical nerves. It is distributed to the superficial parts of the neck.

SUPERFICIES ORBITALIS, (*super*, and *facies*, 'face,' 'the surface,') see Maxillary bone (superior)—s. *Plantaris Pedis*, Sole.

SUPERFECUNDATIO, Superfætation.

SUPERFETATIO, Superfætation.

SUPERFETATIO, *Superfætatio*, *Epiçye'sis*, *Hyperçye'sis*, *Hyperçye'ma*, *Superimprægna'tio*, *Epiç'one*, *Superfæcunda'tio*, *Superfæcunda'tion*, *Superimprægna'tion*, (F.) *Superfætation*, (*super*, and *fatere*, 'to breed,') Conception of a foetus—*epiçye'ma*—in a uterus which already contains one. The impregnation of a female already pregnant. The possibility of this has been denied; but there is abundant evidence to show, that it is possible at a very early period of pregnancy. Twin cases may be of this kind of *superconception*.

SUPERGEMINALIS, (*super*, and *gemini*, 'the testicles,') Epididymis.

SUPERGENUALIS, (*super*, and *genu*, 'the knee,') Patella.

SUPERHUMERALE, (*super*, and *humerus*, 'the shoulder,') Epomis.

SUPERIMPRÆGNATIO, Superfætation.

SUPERIOR AURIS, Attollens aurem—s. Rotundus clitoridis, Ischiocavernosus.

SUPERLABIA, (*super*, and *labia*, 'the labia pudendi,') Clitoris.

SUPERLIGAMEN, (*super*, and *ligare*, 'to bind,') Epidesmus.

SUPERLIGULA, (*super*, and *ligula*, dim. of *lingua*, 'a tongue,') Epiglottis.

SUPERNUMERARY FINGER, (*super*, and *numerus*, 'a number,') Digitus supernumerarius.

SUPERNUTRITIO, (*super*, and *nutrire*, *nutritum*, 'to nourish,') Hypertrophy—s. *Lienis*, *Splenoncus*—s. *Splenis*, *Splenoncus*.

SUPERNUTRITION, Hypertrophy.

SUPEROXALAS KALICUM, Potass, oxalate of.

SUPERPOSITUS MEDICORUM, (*super*, and *positus*, 'placed,') Archiater.

SUPERPURGATIO, (*super*, and *purgare*, *purgatum*, 'to purge,') Hypercatharsis.

SUPERSCAPULARIS, (*super*, and *scapula*,) Supraspinatus—s. Inferior, Infraspinatus.

SUPERSEDENT, *Super'sedens*, (*super*, and *sedere*, 'to sit,') A term used by Professor George B. Wood for a medicine whose operation on any part or organ is to displace or prevent diseased action in that part or organ. An-anti-periodic is a supersedent.

SUPERVACUATIO, Hypercrosis.

SUPIER, Sambucus.

SUPINATEUR, Supinator—s. *Court ou petit*, *Supinator radii brevis*—s. *Grand ou long*, *Supinator radii longus*.

SUPINATION, *Supina'tio*, (from *supinus*, 'lying on the back,') The movement in which the forearm and hand are carried outwards, so that the anterior surface of the latter becomes superior.

In Pathology, *Supination*, *Supina'tio*, *Hyptias'ma*, *Hyptias'mus*, means the horizontal position on the back, with the head thrown back, and the legs and arms extended. It is often a sign of great weakness in disease.

SUPINATOR, (F.) *Supinateur*. That which produces supination. A name given to two muscles of the forearm.

SUPINATOR BREVIS seu MINOR, *Supinator radii brevis*—s. *Longus seu major*, *Supinator radii longus*.

SUPINATOR RA'DII BRE'VIS, *S. bre'vis seu mi'nor*, (F.) *Épicondyl'o-radial*, (Ch.) *Supinateur court*

ou petit supinateur. A muscle seated at the outer and back part of the forearm. It is flattened, triangular, and curved upon itself in such a manner, that it embraces the superior extremity of the radius. It is attached, on the one hand, to the external tuberosity of the humerus, and to a small portion of the posterior surface of the ulna, and is inserted into the upper third of the outer and upper surfaces of the radius. It produces, by contracting, the motion of supination.

SUPINATOR RADII LONGUS, *S. longus sive ma'jor*, *Brachioradial'is*, (F.) *Huméro-sus-radial*, (Ch.) *Long ou grand supinateur*, is seated at the anterior and outer part of the forearm. It arises from the inferior part of the outer edge of the os humeri, and from the external intermuscular aponeurosis; and is inserted, by a long tendon, into the outer edge of the radius, near the base of the styloid process. It causes the supination of the forearm, which it can likewise bend on the arm.

SUPPEDANEÆ, (*sup*, and *pes*, gen. *pedis*, 'foot,') Hypopodia.

SUPPLANTALIA, (*sup*, and *planta*, 'the sole of the foot,') Hypopodia.

SUPPLEMENTÆ EPIPLOICA, (*sup*, and *pleo*, 'to fill,') Appendixæ pinguedinosæ.

SUPPLEMENTARY AIR, see Respiration.

SUPPORTER, ABDOMINAL, (*sup*, and *portare*, 'to carry,') Belt, Russian—s. *Uteroabdominal*, see Belt, Russian.

SUPPORTEUR ABDOMINAL, Belt, Russian.

SUPPOSITION DE PART, (*supponere*, *suppositum* (*sup*, and *ponere*, 'to put'), 'to counterfeited,') Pregnancy, pretended.

SUPPOSITOIRE, Suppositorium.

SUPPOSITORIUM, Suppository—s. *Uterinum*, *Pessary*.

SUPPOSITORY, *Supposito'rium*, *Hypoth'eton*, *Hypoth'ema*, *Bacil'lus*, *Bal'anus*, *Glands*, *G. subdidi'tia*, *Nod'ulus intestinal'is*, (*supponere*, *suppositum* (*sub*, and *ponere*, 'to put'), 'to put under,') (F.) *Suppositoire*. A solid medicine, in the form of a cone or cylinder, intended to be introduced into the rectum;—either for the purpose of favouring or of restraining evacuations, or to act as an anodyne. Purgative suppositories are made of soap, aloes, or any other irritating substance.

The vehicle commonly used for the active ingredients is cocoa-butter. It may require admixture with one eighth part of wax to give it the proper consistence. Tallow with the same proportion of wax is used at times.

Supposito'ria Ac'idi Tan'nici, *Tan'nin Supposito'ries*, and *Supposito'ria Mor'phia*, *Mor'phia Supposito'ries*, are officinal (Ph. B., 1864). The application of a suppository is termed *Balanis'mus*.

SUPPRESSIO ALVI, (*sup*, and *primere*, *pressum*, 'to press,') Constipation—s. *Lochiorum*, *Ischolochia*—s. *Lotii*, *Ischuria*—s. *Mensium*, *Amenorrhœa*—s. *Menstruationis*, *Amenorrhœa*—s. *Sudoris*, *Ischidrosis*.

SUPPRESSION DU FLUX MENSTRUËL, *Amenorrhœa*—s. *de Part*, see Parturition—s. *d'Urine*, *Ischuria*.

SUPPURANS, Suppurative.

SUPPURARE, Suppurate.

SUPPURATE, *Suppurat'æ*, *Matter*, *Beal*, (F.) *Suppurer*. To form or secrete pus.

SUPPURATIF, Suppurative.

SUPPURATION, *Suppurat'io*, *Eçpye'ma*, *Purulen'tia*, *Pyo'sis*, *Pye'sis*, *Diapye'ma*, *Diapye'sis*, *Pyogen'ia simplex*, *Pu'rulence*, *Pu'rulency*, *Gath'ering*, *Beal'ing*, (F.) *Suppuration*, *Aboutissement*. Formation or secretion of pus (see, also, *Pyogenia*), (from *sup*, and *pus*, gen. *puria*.) It is a frequent termination of inflammation, and may occur in almost any of the tissues. This termination is announced by slight chills; by remission of the pain, which, from being lancinating,

becomes heavy : by a sense of weight in the part, and, when the collection of pus can be easily felt, by fluctuation. When the collection of pus becomes prominent, the abscess is said to *point* or to *come to a head*. When pus is thus formed in the areolar membrane, and is collected in one or more cavities, it constitutes an *abscess*. If it be formed from a surface exposed to the air, it is an *ulcer* : and such ulcers we are in the habit of establishing artificially in certain cases of disease.

SUPPURATIVE, *Suppurans*, *Suppurativus*, *Diapye'ticus*, *Epye'ticus*, *Sup'purant*, (F.) *Suppuratif*, *Pyopuë'tique*. That which facilitates suppuration. Suppuratives are, generally, vesicants mixed with lard.

SUPPURER, *Suppurate*.

SUPRA (L.), in composition, above.

SUPRACOSTALES, *Levato' res Costa'rum*, (F.) *Surcostaux*, (from *supra*, and *costa*, 'a rib.') A name given to twelve small, fleshy fasciculi, which are flattened, triangular, and with radiating fibres. They pass from the transverse processes of the dorsal vertebrae to the superior margin of the rib beneath. Spigelius, Cowper, Boyer, Chaussier, and others, consider them to form part of the *intercostales externi*.

SUPRAHEPATIC VEINS, see Hepatic veins.

SUPRAÖMPHALODYMIA, (*supra*, *ομφαλος*, 'navel,' and *δω*, 'to mingle,') see Somatodymia.

SUPRA ET INFRAÖMPHALODYMIA, see Somatodymia.

SUPRAÖRBITAR, *Supraorbita'lis*, *Supraorbita'rius*, (F.) *Susorbitaire*, (*supra*, and *orbita*.) That which is situate above the orbit.

SUPRAÖRBITAR FORA'MEX, *F. Supraorbita'rium seu orbitarium superius*, (F.) *Trou sus-orbitaire ou orbitaire supérieur*, is a foramen or notch completed by a ligament at the inner third of the orbital arch. It gives passage to the *supraorbitar* or *superciliary artery*, a branch of the ophthalmic, which ascends the forehead, and is distributed thereto.

SUPRAPUBIAN, *Suprapu'bic*, *Suprapubia'nus*, (*supra*, and *pubes*.) That which is seated above the pubis.

SUPRAPUBIAN NERVE, (F.) *Nerf nupubien*, is the internal ramus of the inguino-cutaneous branch of the first lumbar nerve. It is the *genito-crural* branch of Bichat. It descends almost vertically before the *psaos muscle*; and, near the crural arch, divides into two branches. One of these accompanies the spermatic cord, and is distributed to the cremaster, serotum, and integuments on the inner part of the thigh. The other issues from the pelvis with the femoral filaments; traverses the aponeurosis of the thigh, and is distributed to the integuments of the upper and anterior part of the limb.

SUPRARENAL, *Suparena'lis*, (F.) *Surrénal*, (*supra*, and *ren*, 'the kidney.') Seated above the kidney—as the *suprarenal capsule*.

SUPRASCAPULAR, *Suprascapula'ris*, (*supra*, and *scapula*.) Seated above the scapula;—as

SUPRASCAPULAR NERVE. This arises from the fifth cervical; passes through the suprascapular notch, and is distributed to the supraspinatus and infraspinatus muscles.

SUPRASCAPULARIS, *Suprascapular*, and *Supraspinatus muscle*.

SUPRASEMIORBITULARIS, *Orbicularis oris*.

SUPRASPINAL, *Supraspinatus*.

SUPRASPINATUS, *Supraspino'eus*, *Supraspina'lis*, *Supraspi'nal*, (F.) *Sur-épineux*, *Sus-épineux*, (*supra*, and *spina*.) That which is seated above a spine.

SUPRASPINATA FOS'SA, (F.) *Fosse sus-épineuse*, is a triangular depression above the spine of the scapula.

SUPRASPINATUS MUS'CLE, *Suprascapula'ris*, *Su*

prascapularis, (F.) *Sus-spino-scapulo-trochitérien*, *Petit sus-scapulo-trochitérien* (Ch.), is situated in the fossa supraspinata. It is elongated, thick, and triangular, and is attached, on the one hand, to the two inner thirds of the fossa supraspinata; and terminates, on the other, at the anterior part of the great tuberosity of the humerus. This muscle raises the arm.

SUPRASPINOSA LIGAMENTA, *Supraspinal lig'aments*, are two in number. 1. The *Dorsolumbo-supraspinal ligament*, (F.) *Ligament sur-épineux dorso-lombaire*, extending above the spinous processes of the dorsal and lumbar vertebrae from the 7th cervical as far as the median crista of the sacrum. 2. *Cervical supraspinal ligament*, (F.) *Ligament sur-épineux cervical*, which extends above all the cervical spinous processes, and is attached, above, to the outer occipital protuberance.

SUR (F.), (L.) *Supra*, in composition, 'above.' **SURA**, *Gastrocnemë*, *Gastrocnem'ia*, *Gastrocnem'ium*; *Calf of the leg*, (F.) *Mollet*, *Gras de Jambe*. The projection, formed at the posterior part of the leg by the *gastrocnemii muscle*. Also, the fibula.

SUR'AL, *Sura'lis*, (from *sura*.) Appertaining or relating to the calf of the leg.

SURCILIER, (*sur*, and *cil*, 'the tarsal cartilage, an eyelash,') *Corrugator supercillii*, *Superciliary*.

SURCOSTAUX, (*sur*, and *costal*, 'belonging to the rib' [*costa*]), *Supracostales*.

SURCULATION, (from *surculus*;) see Generation.

SURCULI FELLEI, (pl. of *Surculus*;) *Pori bilarii*.

SURCULUS, 'a young branch,' *Ramusculus*.

SURDENT (F.), (*sur*, and *dent*, 'a tooth,') When a milktooth does not fall out at the period of the second dentition, but is merely forced to one side by the new tooth, it is called a *surdent*.

SURDI-MUTITÉ, *Mutitas surdorum*.

SURDITAS, (from *surdus*;) *Cophosis*, *Deafness*.

SURDITÉ, *Cophosis*, *Deafness*.

SURDOMUTITAS, (*surdus*, and *mutus*, 'mute,') *Mutitas surdorum*.

SURDUS, *Deaf*.

SUREAU COMMUN, *Sambucus*.

SURELLE, ([Sax.] *pup*, 'sour,') *Oxalis acetosella*, *Rumex acetosa*.

SUR-ÉPINEUX, (*sur*, and *épine*, 'a spine,') *Supraspinatus*.

SUR-EXCITATION, see Excitement, and Superexcitation.

SURFEIT, (*sur*, and *fait*, 'done,') *Acrasia*. *Colica crapulosa*.

SUR'GEON, *Chiru'geon*; *Chiro'nax*, *Chiru'gus*, *Vul'nerum medicus*, *Cheiria'ter*, *Chiria'ter*, *Iä'trus*, (F.) *Chirurgien*, old French, *Surgien*, (*χαιρ* 'the hand,' and *εργον*, 'work,') One who practises surgery. The surgeon, says Celsus, ought to be young, or, at all events, not advanced in life. His hand must be steady, expert, and never tremulous: he must use the right or left with equal facility: his sight must be clear and penetrating: his mind intrepid; and, although unmoved when he proceeds to operate, he must not be in haste, nor cut less than is necessary, but finish his operation, as if the cries of the patient made no impression upon him.

In the middle ages, the Universities of Italy and Spain sent out educated surgeons, as well as physicians, under the name *chiru'gi phys'ici*.

SURGEON ACCOUCHEUR, see Surgeon apothecary.

SURGEONAPOTHECARY. One who unites the practice of surgery with that of the apothecary. A general practitioner. He is the ordinary medical man for fairly medical attendant in Eng-

considerable depression. If accompanied with a vocal sound, it constitutes a *groan*.

When the respiration, on auscultation, resembles sighing, it is called, by some, *Suspirious*, (F.) *Suspiriosus*.

SUSPIRIUM CARDIACUM, Angina pectoris.

SUSTENTACULUM, (*oss*, and *tenuis*, *tendens*, 'to hold,') Aliment.

SUSTENTATOR CLITORIDIS, Erector clitoridis—s. Penis. Erector penis.

SUSURRATION, (*murmurare*, *murrationem*, 'to murmur, to whisper,') Whispering.

SUSURRUS, 'a murmur,' Bourdonnement—s. Aurium, Tinnitus aurium.

SUTELA, (*seare*, *sutum*, 'to sow,') Suture.

SUTERBERRY, Xanthoxylum fraxineum.

SUTORIUS, (*sutor*, 'a shoemaker,') Sartorius.

SUTTER, Saccharum.

SUTURA, gen. *Sutura*, Suture—s. Abdominalis, Gastroraphy—s. Arcualis, Coronal suture—s. Biparietalis, Sagittal suture—s. Clavata, Suture, quilled—s. Corporis callosi externa, see Raphe—s. Cranii squamosa, Squamous suture—s. Dentata, see Suture—s. Frontoparietalis, Coronal suture—s. Interparietalis, Sagittal suture—s. Jugalis, Sagittal suture—s. Lambdoidalis, Lambdoid suture—s. Lepidoides, Squamous suture—s. Limbosa, see Suture—s. Mentum, Squamous suture—s. Notha, see Suture—s. Obliqua, Sagittal suture—s. Ossium spuria, Harmony—s. Protra, Lambdoid suture—s. Punctata, Suture, common—s. Puppis, Coronal suture—s. Rhabdoides, Sagittal suture—s. Serrata, see Suture, Sagittal suture—s. Spuria, see Suture—s. Squamosa, see Suture—s. Vera, see Suture.

SUTURAL, *Suturalis*; (same etymon as the next.) Relating to a suture.

SUTURE, *Sutura*, *Sutula*, *Ar'mē*, *Rha'phē*, (*seare*, *sutum*, 'to stitch,') Donetail joint, (F.) *Engrenure*. A kind of immovable articulation, in which the bones unite by means of serrated edges, which are, as it were, dovetailed into each other. The articulations of the greater part of the bones of the skull are of this kind. The coronal, lambdoid, and sagittal sutures are called *true sutures*, *sutura vera*, the two temporal are called *false sutures*, *sutura notha* seu *spuria* seu *quadrata*.

A *dentated suture*—*Sutura dentata*—is one in which the processes are long and dentiform, as in the interparietal suture. In a *serrated suture*—*sutura seu agnata vera*—the indentations and processes are small and fine, like the teeth of a saw, as in the suture between the two portions of the frontal bone. In the *sutura limbosa* there is, along with the dentated margins, a degree of bevelling of one, so that one bone rests on the other, as in the occipitoparietal suture.

SUTURE, in *Surgery*, *Rha'phē*, is an operation which consists in stitching the lips of a wound to procure their union. The suture was, at one time, largely employed; but, in modern times, its use has been wisely restricted. There are few wounds in which the edges may not be brought together by adhesive plaster appropriately applied. This kind of union, by adhesive plaster simply, has been absurdly termed *Sutura sicca* or *dry suture*, in opposition to the *Sutura cruenta*, (F.) *Suture sanguinea*, or *Suture properly so called*. The suture is generally formed with waxed thread and straight or curved needles, according to the arrangement of parts.

The chief sutures are the *interrupted suture*, the *quilled suture*, the *glover's suture*, and the *twisted suture*. 1. The *INTERRUPTED SUTURE*, (F.) *Suture à points séparés*, *Suture entrecompie*, is formed by passing a needle, armed with a ligature, through the lips of the wound previously brought into contact; and then tying the

extremities of the thread. The other stitches are made in the same manner. They are separate or interrupted. It is the suture most commonly used. 2. The *QUILLED SUTURE*, *Sutura clavata*, (F.) *Suture échecaille* ou *empenée*, *Compound Suture*, is merely the interrupted suture, with this difference, that the ligatures are not tied over the face of the wound, but over two quills or rolls of plasters, or *boagies*, which are laid along the sides of the wound. It is used by some surgeons, on account of there being less danger of the stitches tearing out. It is not much employed. The *CLAMP SUTURE*, used by Dr. Sims, of New York in the treatment of vesicovaginal fistula, is a modification of this. It is formed by two cylinders of silver or lead, perforated at several points for the passage of small pieces of fine silver wire, which are to supply the places of thread, and are prevented from slipping by perforated shot carried down upon them, pressed against the cylinders, and kept in place by being firmly pinched with pliers.

3. The *GLOVER'S SUTURE*, *Continued or Uninterrupted Suture*, (F.) *Suture de Pelletier*, ou *en surjet*, is executed by introducing the needle first into one lip of the wound from within outwards, then into the other in the same way; and, in this manner, sewing up the whole extent of the wound. It has been chiefly used in wounds of the intestinal canal. It is now rarely employed except for stitching up dead bodies. The common stitch or suture, *Sutura punctata*, (F.) *Suture à points passés*, has been used in similar cases. 4. The *ROYAL STITCH* or *STITCH* is the name of an old operation for the cure of bubonocele. It consisted in putting a ligature under the hernial sac, close to the abdominal ring, and then tying that part of the sac, so as to render it impervious by the adhesive inflammation thus excited. 5. The *STAPLE SUTURE* consists of a piece of steel wire, shaped like a staple, the ends of which are ground to a triangular or glover's point. In using it, the points of the staple are to be brought through both margins of the wound, and then bent back. 6. The *TWISTED SUTURE*, (F.) *Suture entortillée*, is employed, chiefly, to unite wounds of the cheeks and of the free edges of the lips. To perform it, a pin or needle is passed through the edges of the wound, so as to keep them accurately in contact. A piece of thread is then wound accurately around the pin, from one side of the division to the other, in the form of the figure z. The number of pins and sutures must, of course, correspond with the extent of the wound. This is the suture used in cases of harelip. The *BUTTON SUTURE* of Dr. Bozeman, of Alabama, employed in vesicovaginal fistula, is a modification of this. 7. The *STITCH ANSE*, of Le Dran, has only been employed in cases of wounds of the intestines. For this purpose, the edges of the wound are brought together, and as many needles, with unwaxed threads, used as there are to be stitches. Each stitch is then made by passing a needle through the edges; and all the ligatures which belong to one side of the wound are tied together, and twisted so as to form a thread, which is fixed externally. 8. The *IMPLANTED SUTURE*, (F.) *Suture implantée*, is formed by pins arranged parallel to the edges of the wound, and implanted in their substance. By means of threads, the needles are approximated so as to bring the edges of the wound together. 9. The *PLASTIC SUTURE*, so called by Prof. Poncest, of Philadelphia, its proposer, is a suture by tongue and groove. The groove is made by an incision in the part, which is to receive the flap. The tongue is made on the flap by bevelling off the two surfaces, cuticular and adipose. The suture is made by passing a loop of thread, by two punctures, deep through

the inner lip of the groove. The two needles are passed through the base of the tongue, and again through the outer lip of the groove. The two ends of the thread are then tied over a small roll of adhesive plaster, by which the tongue is, of necessity, forced deep into the groove. Four raw surfaces are thus firmly held in connection, making the union of parts, by this suture, a very certain process.

SUTURE, BUTTON, see Suture, (twisted) — s. Clamp, see Suture, (quilled) — s. Compound, Suture, quilled — s. Continued, Suture, Glover's — s. *Écailleuse*, Squamous suture — s. *Empeignée*, Suture, quilled — s. *Emplumée*, Suture, quilled — s. *Enchevillée*, Suture, quilled — s. *Entortillée*, Suture, twisted — s. *Entrecoupée*, Suture, interrupted — s. False, Harmony, see Suture — s. *Implantée*, see Suture, (implanted) — s. *Occipitale*, Lambdoid suture — s. Occipitoparietal, Lambdoid suture — s. *de Pelletier*, Suture, Glover's — s. Perineal, Perineorrhaphy — s. Plastic, see Suture — s. *à Points passés*, Suture, common — s. *à Points séparés*, Suture, interrupted — s. Sagittal, see Sagittal suture — s. *Saigillante*, see Suture — s. *Squameuse*, Squamous suture — s. Superficial, Harmony — s. *en Surget*, Suture, Glover's — s. Twisted, see Suture — s. Uninterrupted, see Suture, Glover's.

SWAB, ([Sax.] *ꝛpebban*, 'to clean with a mop.') A piece of sponge or rag tied to a rod of whalebone or wood, for cleansing the mouth of the sick, or applying remedial agents to deep-seated parts. The process is termed *Swab'bing*.

SWADDLING CLOTHES, Swathing clothes.

SWAGBELLY, Physconia.

SWALLOW, *Hirundo* — s. Wort, *Asclepias vincetoxicum* — s. Wort, orange, *Asclepias tuberosa* — s. Wort, tuberous-rooted, *Asclepias tuberosa* — s. Wort, white, *Asclepias vincetoxicum*.

SWALLOW'S NEST, *Nidus hirundinis*.

SWALME, ([Sax.] *ꝛpam*, [G.] *Schwamm*.) Excrecence, Tumour.

SWAMP CABBAGE, *Dracontium foetidum* — s. Dogwood, *Petela trifoliata* — s. Sickness, Milk sickness — s. Wood, *Cephalanthus occidentalis*, *Dirca palustris*.

SWAN ALLEY SORE, see Phagedenic chancre.

SWARFF, Stupor, Syncope.

SWATHING CLOTHES, *Swad'dling Clothes*, *Incunabula*, *Fas'ciae*, *Pan'ni*, ([Sax.] *ꝛpeðan*, 'to bind.') (F.) *Maillot*. The kind of clothing or bandaging formerly applied to children during the first year of life. It is now, properly, laid aside, and every freedom allowed to muscular exertion.

SWEAMISH, see Syncope.

SWEAT, ([Sax.] *ꝛpar*.) Sudor — s. Bloody, Sudor cruentus — s. Glands, see Perspiration — s. Root, *Polemonium reptans* — s. Scab, *Crusta genu equinae*.

SWEATING, *Ehidrosis* — s. Bath, *Achicolum* — s. of Blood, see Diapedesis, and Sudor cruentus — s. House, *Achicolum* — s. Plant, *Eupatorium perfoliatum* — s. Room, *Achicolum* — s. Sickness, Sudor Anglicus.

SWEATING SICKNESS, MALWAH. A disease which appears to be allied to the worst form of cholera, and to bear a close relation to malignant congestive fever. As its name imports, it occurs in Malwah, in India.

SWEBBING, Syncope.

SWEET, (sway,) Vertigo.

SWEEMING, Syncope.

SWEEMISH, see Syncope.

SWEET, BITTER, *Solanum dulcamara*.

SWEET BITTER, *Triosteum* — s. Bread, *Pancreas*, Thymus — s. Bush, *Comptonia asplenifolia* — s. Fern, *Comptonia asplenifolia* — s. Flag, *Acorus calamus* — s. Pod, *Ceratonium siliqua* — s. Root, *Aralia nudicaulis*.

SWELL'ING, *Epanaste'ma*, *Eaar'ma*, *Exæde'is*, *Pre'sis*, *Pres'ma*, *On'cus*, *Extumefactio*, *Tu'mor*, *Tumefactio*, (Prov.) *Blaw'ing*, *Boist*, *Coul*, (Old Eng.) *Boche*, *Bol'tynge*, *Bol'ning*, (F.) *Enflure*, *Gonflement*, (from [Sax.] *ꝛpellan*, 'to swell.') Morbid increase of bulk in a part of the body. See Tumour.

SWELLING, *White*, *Hydrarthrus*.

SWELTING, see Syncope.

SWENSIE, (quinsy,) *Cynanche Tonsillaris*.

SWENYNG, *Somnium*.

SWERF, Stupor, Syncope.

SWERTIA CHIRAYITA, (after Emanuel Swert, a Dutch botanist,) *Gentiana chirayita* — s. Difformis, see *Calumba* — s. *Fraseri*, see *Calumba* — s. *Lutea*, *Gentiana lutea*.

SWEUIN, *Somnium*.

SWEVYNG, *Somnium*.

SWEVYNYNG, *Somnium*.

SWIETENIA FEBRIFUGA, (after Van Swieten,) *S. Smit'da*, *Soym'da* seu *Cedre'la febrif'uga*, *Feb'rifuge Swiete'nia*, (F.) *Mahogani febrifuge*. Ord. Meliaceæ. Ser. Syst. Decandria Monogynia. The bark of this tree, which is a native of the East Indies, called there *Kohena Bark*, has an aromatic odour and bitter taste. It is tonic and stomachic, and has been used in intermittents, and in diseases where bitter tonics are indicated. Dose, ʒj to ʒij of the powder.

SWIETENIA MAHOG'ANI, *Cedrus Mahog'ani*, *Mahag'oni*, *Maag'oni*, *Mahog'any Tree*, grows in the West Indies and tropical America, and is possessed of similar properties.

SWIETENIA SENEGALEN'SIS, *Khaya Senegalen'sis*. A common forest tree on the borders of the Gambia. The bark called *Cail-cedra*, and *Karson Khayi*, is bitter and antiperiodic.

SWIETENIA SOMIDA, *S. Febrifuga*.

SWIM, Air bladder.

SWIMBLADDER, Air bladder.

SWIMMING, *Natation* — s. Bladder, *Air bladder* — s. of the Head, *Vertigo*.

SWINACIE, *Cynanche tonsillaris*.

SWINEPOX, see *Varicella*.

SWINESAME, *Adeps*.

SWINETHISTLE, *Sonchus*.

SWINNYING, *Vertigo*.

SWOON, (from [Sax.] *arþpanan*, 'to languish,') Syncope.

SWOONING, Syncope.

SWOUNDING, Syncope.

SY, in composition, 'with,' 'together,' see *Sym*.

SYALADENTITIS, see *Sialadenitis*.

SYCAMINUS, (*συκαμινος*;) *Morus nigra*.

SYCAMINUM, (*συκαμινον*;) see *Morus nigra*.

SYCE, (*συκη*;) *Ficus*, *Ficus carica*.

SYCEA, (*συκεια*;) *Ficus*.

SYCEPH'ALUS, (*συ*, for *συν*, and *κεφαλη*, 'head.') A monster having two heads united together.

SY'CION, (*συκιον*, from *συκε*.) A decoction of dried figs.

SY'CIOS ANGULA'TUS, (*συκιος*, 'a gourd.') *One-seeded star cu'cumber*, *Wild Bry'ony*. The roots and seeds of this indigenous plant, Order Cucurbitaceæ, are bitter, purgative and diuretic. It is used in dropsy in the same cases as bryony.

SYCITES, (*συκιτης*, from *συκη*, 'a fig.') Made of figs, see *Catorechites*.

SYCOMA, (*συκωμα*, from *συκε*, and *αμα*;) *Ficus*.

SYCO'SIS, (*συκωσις*.) A tumor of the shape of a fig. Also, a fungous ulcer. A horny excrecence about the eyelids. Dr. Bateman defines it: — an eruption of inflamed but not very hard tubercles occurring on the bearded portion of the face and on the scalp, in adults; and usually clustering together in irregular patches; — *Phy'na Syco'sis*, *Rose'ola fico'ra*, *Ac'nē men'tagra*. Of this he has given two species: 1. *Syco'sis Men'ti*, *S. bar'ba*, *Men'tagra*, *Menti'go*, *Va'rus Men'tagra*,

SYMPATHY, CURE BY. *Medicina sympathetica*. *M. Magnetica*. A mode of treatment, at one time prevalent, which was based on the idea, that a mysterious consent exists between parts of the body and agents, for which there was no real

foundation. See Sympathetic powder, and Trial by Bier-right.

SYMPEPSIS, (*sym*, and *πέψις*, 'coction, digestion,') Coction.

SYMPEXIONS, see Sympexis.

SYMPEXIS, (*συμπίξις*), Concretion. M. Robin has applied the term *Sympexions* to solid concretions found in the vesicles of the thyroid gland, spleen, lymphatic ganglion, in the small cysts of the glands of the neck and body of the uterus, and, almost always, in the liquid of the vesiculae seminales. Little is known of their composition.

SYMPHORA, (*συμφωρα*, from *sym*, and *φωρ*, 'to carry,') Congestion.

SYMPHOREMA, gen. *Symphore'matis*, (*συμφωρημα*), Congestion.

SYMPHORESIS, (*συμφορησις*), Congestion.

SYMPHORICAR'PUS RACEMOSUS, *Synonberry*, (*συμφωρεω*, 'to bear together,' and *καρπος*, 'fruit,' owing to the clustered berries.) An indigenous plant, Order, Caprifoliaceae, Tribe, Lonicereae, whose berries are large, bright white, and remain till winter; flowering from June to September. The root is tonic and astringent, and has been used in ague; and by the Indians in syphilis.

SYMPHYIA, (*συμφυια*), Symphysis.

SYMPHYOCEPH'ALUS, (*Μετωπ'ος*, *Cephalop'os*, from *συμψυς*, (*sym*, and *ψω*, 'to grow,') 'united together,' and *κεφαλη*, 'head,') A monstrosity, in which twins are united by the head.

SYMPHYSE, Symphysis.

SYMPHYSEOT'OMY, *Symphysot'omy*, *Synchondrot'omy*, *Symphysiotom'ia*, *Synchondrotom'ia*, (from *symphyseis*, and *τεμνω*, 'I cut,') The operation of section of the *Symphysis*, *Synchondrian* operation or section. This operation is performed with the view of augmenting the diameter of the pelvis, in order to facilitate delivery in certain cases of faulty conformation of that cavity, with wedging of the head of the child therein. To perform the operation, (F.) *désynphyser*, the female is placed on her back; the pubes is shaved; and, with a strong scalpel, the soft parts are divided on the median line. The articulation is then cut. As soon as the fibrous parts, which hold the bones together, are divided, they separate to a certain distance from each other; and the delivery may be accomplished. The knife, with which the operation is performed, is called, by the French, *Couteau symphysectom*.

SYMPHYS'IA, (from *συμψυω*, 'to grow together,') A malformation consisting in the union of parts that are usually divided.

SYMPHYSIEN, see Symphyseotomy.

SYMPHYSIOTOMIA, Symphyseotomy.

SYMPHYSIS, *Symphys'ia*, *Coalescen'tia*, *Coall'itas*, *Coalit'io*, (F.) *Symphysie*, (*συμψυσις*, from *sym*, and *ψω*, 'I grow,') A union of bones. The bond of such union. The aggregate of means used for retaining bones *in situ* in the articulations. The name symphysis, (F.) *Symphysie*, has, however, been more particularly appropriated to certain articulations; as to the *symphysis pubis*, *maxillaris*, *symphysis*, &c. See Monster.

SYMPHYSIS, Coalescence, Commisura, Insertion — s. Cartilaginea, Synchondrosis — s. Ligamentosa, Syndesmosis — s. Menti, see Maxillary bone, inferior — s. Ossium muscularis, Syssarcosis — s. Pubis, Pubic articulation.

SYMPHYSODACTYL'IA, (*symphysis*, and *εκακτολος*, 'a finger,') An abnormal union of the fingers.

SYMPHYSOP'SIA, (*symphysis*, and *ωφ*, 'the eye,') A monstrous fusion of the two eyes into one.

SYMPHYSOTOMY, Symphyseotomy.

SYMPHYTES, (*συμφυτες*, 'grown or produced with,') Hereditary.

SYMPHYTUM, (*συμφυτον*), *Symphytum officinale* seu *album* seu *Bohemicum*, *Consolidila majus*, *Cons'froy*, *Alum*, *Al'us*, (Prov.) *Knitback*, (F.) *Grande Consolide*, *Oreille d'âne*. Ord. Bo-

raginaceae. See *Syst.* Pentandria Monogynia. The roots of this European plant are very mucilaginous, and are employed as emollients. They are used in decoction in hæmoptysis, dysentery, chronic diarrhæa, &c.

SYMPHYTUM, *Solidago virgaurea* — s. *Album*, *Symphytum* — s. *Bohemicum*, *Symphytum* — s. *Minimum*, *Bellis* — s. *Minus*, *Prunella* — s. *Officinale*, *Symphytum*.

SYMPHYTUM PETRÆ'UM, *Co'ris Montpellier'nis*, *Montpellier Co'ris*. An intensely bitter and nauseous plant, which is said to have been useful in syphilis. See also, *Sanicula*.

SYMPOLOCARPUS ANGUSTISPATHA, (from *συνπλοκον*, 'connection,' and *καρπος*, 'fruit,' from the coalescing ovaries,) *Dracontium angustipatha* — s. *Fetidus*, *Dracontium fetidum*.

SYMPOD'IA, *Monopod'ia*, (*sym*, and *πους*, gen. *podos*, 'a foot,') A monstrosity in which there is coalescence of the lower extremities.

SYMPTOM, *Symptom'a*, *Semel'ion*, *Ac'idens*, *Ca'usa*, (F.) *Symptôme*, (*συμπτωμα*, from *sym*, and *πτωμ*, 'a fall,' from *πτωω*, 'I fall,') Any change, perceptible to the senses in any organ or function, which is connected with morbid influence. It is by the aggregate and succession of symptoms that a disease is detected. The term *Symptoms of Symptoms* has been used for the effects which result from the symptoms of a disease; but which effects are not essentially connected with the disease itself. Thus, the debility arising from the frequency of evacuations in dysentery is a symptom of symptoms. *Symptom*, at one time, was generally used in the same sense as *sign*; but with many, perhaps most, of the present day, the former signifies a functional or vital phenomenon of disease, whilst the latter is applied to that which is more directly physical; and hence the expressions *functional or vital phenomena* or *symptoms*, in contradistinction to the *physical signs* afforded by auscultation, percussion, &c. See *Sign*.

SYMPTOMA, gen. *Symptom'atis*, *Symptom* — s. *Turpitudinis*, *Symphomania*.

SYMPTOMATIC, (*συμπτωματος*), *Symptomat'icous*, (F.) *Symptomatique*. That which is a symptom of some other affection. Thus, we speak of a symptomatic disease, in opposition to one that is idiopathic.

SYMPTOMATOGRAPHY, (*symptoma*, and *γραφειν*, 'a description,') *Semeiography*.

SYMPTOMATOLOG'Y, *Symptomatolog'ia*, *Symbol'ogy*, *Synbolog'ia*, (*symptomai*, and *λογος*, 'a treatise,') The branch of pathology, the object of which is the observation (*phænomenoscopy*) and knowledge of symptoms.

SYMPTOSIS, (*συμπτωσις*, from *sym*, and *πτωω*, 'I fall,') Depression or collapse, emaciation or atrophy, either of the whole body or of a part.

SYN, as a prefix, *Sym*.

SYNEREMA, gen. *Synærem'atis*, (*συναεμα*), *Corrugation*.

SYNETION, (*συναιτια*, from *syn*, and *αιτια*, 'a cause,') *Concusa*.

SYNANCHE, *Cynanche*, *Cynanche tonsillaris*.

SYNAN'GIE, *Cynanche*.

SYNAPHYMENTITIS, (*συναψη*, 'a conjunction,' *τιμη*, 'a membrane,' and *ιτις*,) see *Ophthalmia*.

SYNAPTASE, (*syn*, and *αεωω*, 'to join,') with amygdalin it forms hydrocyanic acid, is Amygdalin.

SYNARTHRODIA, *Synarthrosis*.

SYNARTHRO'DIAL, *Synarthrodia'lis*. That which belongs to synarthrosis; as *Synarthrodial articulation*.

SYNARTHROSIS, (*συνάρθρωσις*, from *syn*, and *άρθρσις*, 'articulation,') *Synarthrod'ia*, *Abartition'io*, *Quartulation'io*. An immoveable articulation. Synarthrosis includes three species: *Suture*, *Hermomy*, and *Gomphosis*.

have or hold,') 'I continue.') *Inflam'matory* or *Dynamic Fe'ver*, *Angiop'gra*, *Sthenop'gra*, *Fe'bris causo'des* seu *causato'des* seu *deu'rens* seu *ar'dens* seu *contin'ua* *sanguin'ea* seu *contin'ua* non *put'ris* seu *acnast'ica* seu *acut'a* seu *acut'a* *sanguin'ea* seu *contin'ens* seu *continens* non *put'rida* seu *angioten'ica* seu *sthen'ica* seu *sanguin'ea* seu *inflammato'ria*, *Hemato'nos*, *Syn'ochus* (Vogel), *S. impu'tris* (Galen), *Ene'cia* *cut'nea* seu *ton'ica*, *Philop'gra*, *Cu'd'ma*, *Calent'ua* *contin'ua* (F.) *Fièvre ardente* ou *continue* *inflammatoire* ou *sanguine* ou *irritative* ou *angioténique* ou *inflammatoire*, *Gastro-entérite intense*; *General inflammation*. A species of continued fever, characterized by increased heat; and by quick, strong, and hard pulse; urine highcoloured; disturbance of mind slight. It requires, of course, the most active treatment.

SYNOCHA RHEUMATICA, Rheumatism, acute.—s. *Vaccina*, *Vaccina*.

SYNOCHAL, *Synocha'lis*. Relating to, or having the characters of, synocha.

SYNOCHUS, (*συνος*, continued,) *Ene'cia syn'ochus*, *Fe'bris len'ta*, (Linnæus). Continued fever, compounded of synocha and typhus:—in its commencement often resembling the former; in its progress, the latter: *Mix'ed fe'ver*, *Com'mon contin'ued fe'ver*, *Syn'ochoid fe'ver*. See *Acnasticus*, *Continued fever*, *Fever*, and *Typhus*.

SYNOCHUS, Synocha.—s. *Biliosa*. Fever, gastric, Fever, bilious.—s. *Catarrhalis*, *Influenza*—s. *Ictero'des*, *Fever*, yellow.—s. *Impu'tris*, *Synocha*—s. *Miliaria*, *Miliary fever*—s. *Varicella*, *Varicella*.

SYNOLCE, (*συνολκη*, from *syn*, and *ελκω*, 'to draw,') Spasm.

SYNSTEOGRAPHY, *Synosteograph'ia*, (*syn*, *οστειον*, 'a bone,' and *γραφω*, 'I describe,') The part of anatomy which describes the joints.

SYNSTEOLOGY, *Synosteolog'ia*, (*syn*, *οστειον*, 'a bone,' and *λογος*, 'a treatise,') A treatise on the joints.

SYNSTEO'SIS, *Synosto'sis*, (*syn*, *οστειον*, 'a bone,' and *osis*,) Union by means of bone.

SYNSTEOTOMY, *Synosteotom'ia*, (*syn*, *οστειον*, 'a bone,' and *τεμνειν*, 'to cut,') Dissection of the joints.

SYNSTOSIS, *Synosteosis*.

SYNOTE, (*syn*, and *οτις*, *ωτις*, 'an ear,') A genus of double monster of Is. G. St. Hilaire, the second face of which is represented only by two ears close together, or a double median ear.

SYNOVIA, (*syn*, and *ων*, 'ovum, an egg,') *Mucila'go*, *Un'guen* seu *Unguen'tum* seu *Sney'ma articula're*, *Azun'gia articula'ria*, *Mucila'go articulo'rum* seu *junctura'rum*, *Hu'mor articula'ris*, *Hydrar'theris*, *Hu'mor seu Hu'midum nati'vum* seu *Glu'ten articulo'rum*, *Arthrohydry'nē*, (F.) *Synovie*. A fluid resembling the white of egg, which is exhaled by the synovial membranes surrounding the moveable articulations. The synovia has been found, on analysis, to consist of water, albumen, soda, chloride of sodium, phosphate of lime, and—Fourcroy fancied—uric acid. Its use is to lubricate the joints, and the sheaths in which tendons play.

SYNOVIAL, *Synovia'lis*. That which relates to the synovia.

SYNOVIAL CAPSULES, see *Membrane*.

SYNOVIAL GLANDS, *Hav'ers's Glands*, *Gland'ule articula'res* seu *Synovia'les* seu *Haversia'næ*, are the secretory fringes, which float in the interior of the synovial capsules (see *Capsules*.) They do not seem to be glandular. They have been called *Synovial fim'briae*, (F.) *Franges synoviales*.

SYNOVIAL MEMBRANES, see *Membrane*.

SYNOVITIS, (*Synovia*, and *itis*.) A term, badly compounded, applied at times to inflammation of the synovial membrane, *arthrophlog'osis synovia'lis*. When it affects the bursæ, it is sometimes called *Burs'itis*.

SYNOVITIS RHEUMATISMALIS, Rheumatism, (acute).

SYNTASIS, (*συντασις*, from *συντανω*, (*syn*, and *τεινω*, 'I stretch,') 'I strain,') Tension of parts: hence *συντατικα*, are medicines which render parts tense. *Syntex'is*, on the other hand, means *Colliquat'io*, or wasting of the body; from *συντενω*, 'I melt together.'

SYNTAXIS, (*συνταξις*, from *syn*, and *τασσειν*, 'to put in order,') Articulation, Reduction, Taxis—s. *Serrata*, see *Suture*.

SYNTECOPYRA, (*syntexis*, and *πυρ*, 'fever,') Hectic fever.

SYNTECTOCOPYRA, Hectic fever.

SYNTENO'SIS, (*syn*, *τεινω*, 'a tendon,' and *osis*.) A species of articulation in which two bones are joined by means of tendon; as the sesamoid bones with the toes, the rotula with the tibia, &c.

SYNTERESIS, (*συντηρησις*, from *syn*, and *τηρειν*, 'to keep,') Prophylaxis.

SYNTERETICUS, (*συντηρητικος*.) Prophylactic.

SYNTEXIS, (*συντηξις*.) Colliquation, Consumption. see *Synstasis*.

SYNTHENA. A term employed by Paracelsus to designate a species of epilepsy, accompanied by cardialgia and tormina.

SYNTHESIS, (*συνθεσις*,) *Reu'nio*, 'composition or putting together,' (from *syn*, and *τιθημι*, 'I place,') A surgical operation, which consists in uniting parts that are divided; and approximating such as are separated or distant from each other. *Reu'nio par'tium separatu'rum*, *Syn'thesis of continu'ity*, means the union of the edges of a wound, or the approximation of the extremities of a fractured bone. *Synthesis of contigu'ity* is the reduction of displaced organs; as in cases of hernia and luxations.

SYNTHESIS, Composition.

SYNTHETISM'US. The aggregate of operations and means for reducing a fracture, and maintaining it reduced:—comprising extension, counterextension, coaptation, and bandaging.

SYNTONIN, (*syn*, and *τονος*, 'a muscle,') see *Fibrin*.

SYNTRIMMA, gen. *Syntrimm'atis*, (*συντριμμα*, from *syn*, and *τριβω*, 'to rub,') Contritio.

SYNTRIPSIS, (*συντριψις*.) Contritio.

SYNTROPHUS, (*συντροφος*, from *syn*, and *τροφω*, 'to nourish,') Connutritus.

SYNULOSIS, (*συνουλωσις*, from *syn*, *ουλωω*, 'to cicatrize,' and *osis*.) Cicatrization.

SYNULOTICA, (*συνουλωτικά*.) Cicatrisantia.

SYNUSIA, (*συνουσια*.) Coition.

SYNYMEN'SIS, *Synhymen'esis*, *Synimen'esis*, (*syn*, and *υμην*, 'a membrane,') Union of bones by membrane, as in the case of the bones of the head in the fœtus. *Syndesmosis*.

SYPHAC, Peritoneum.

SYPHILIDES, (F.) *Syphilides*, *Dermo-syphilides*, *Dermatoses véroleuses*. (From *syphilis*.) The cutaneous eruptions that appertain to constitutional syphilis are so denominated by the French writers.

SYPHILIDATER, *Syphilidat'rus*, (*syphilis*, and *ιατρος*, 'a physician,') A pox doctor. One who occupies himself in treating syphilis.

SYPHILIDOGRAPHY, Syphilography.

SYPHILIDOLOGY, Syphilology.

SYPHILIDOMANIA, Syphilomania.

SYPHILIPHOBIA, (*syphilis*, and *φοβος*, 'dread,') A morbid dread of syphilis, giving rise to fancied symptoms of the disease; a form of hypochondriasis.

SYPHILIS, gen. *Syphil'idis*, *S. ven'e'ra* seu *malig'na*, *Siphilis*, *Syphilit'imus*, *Cachex'ia seu Cachex'ia Ven'e'ra*, *Mal'um ven'e'reum*, *Puden'dagra*, *Mer'curius*, *Lu'ës*, *L. ven'e'ra* seu *syphilis*, *Mor'bus Gu'tticus* seu *Ind'icus* seu *Hispan'icum* seu *Neapolita'nus* seu *Ind'icus* seu *aphrodis'ius*,

Pato'ca, Basiliscus, Bo's, Sod'oma Galla'eum; Venereal, Venereal Disease, Poz, French Poz, French Crot, French Dintem'per, Foul Disease, old cant term Gri'e'comes, (Prov.) Pip, (Sc.) Glengore, Glengour, Grandgore, (F.) Vérole ou Grosse Vérole, Maladie vénérienne, Mal de Naples, Mal Français, Maladie de Vénus. (The etymology of syphilis is unknown. Some consider it to proceed from *sus*, 'a hog,' and *phileo*, 'I love;' others, from *syn*, 'with,' and *phileo*, 'I love;' and others, with more probability, from *syphlos*,—formed, by contraction, from *syphlos*,—'a reproach,' &c.) It is an infectious disease, communicable by coition, or by the contact of parts that are only lined with a thin epidermis; as the lips, nipple, &c. It generally makes its first appearance by a chancre, followed sooner or later by a bubo. The chancre is a *primary* symptom, and there are other symptoms which are *secondary* or *constitutional*. These are:—ulcers in the throat; coppercoloured blotches on the skin; pains in the bones, nodes, &c.: the last are sometimes termed *tertiary* phenomena. The origin of syphilis is not known. It is believed to have been introduced into Europe from America, at the end of the 15th century [?]. Its progress is extremely variable, as well as its duration, which is often unlimited, if left to itself. Mercury may be esteemed the best remedy, but the disease admits of cure without it. Almost every antisypilitic composition, whose use is attended with success, probably contains mercury united with diaphoretics or other medicines. It is generally sufficient to keep the mouth sore for a fortnight for the cure of chancre: for venereal sore throat, and other recent constitutional symptoms, eight or ten weeks may be required; and, of course, a longer time, if these symptoms have persisted longer. Sometimes, after mercury has been given for a time, the symptoms continue stationary; and do not yield until the mercury has been discontinued, and tonics and eutrophics have been administered. The preparations of iodine have been much used in the revellent treatment of this disease.

SYPHILIS INDICA, Framboesia.

SYPHILIS PSEUDOSYPHILIS, *Lu'es Syphilitides, Pseudosyphilis*. A term given to ulcers or other affections that resemble syphilis, but are indeterminate in their character,—the symptoms being irregular in their appearance, and usually yielding spontaneously, or under the use of tonics, change of air, &c. The condition has not been at all well defined. *Sivens, Radezyge, &c.*, have been referred to this head.

SYPHILIS, SECONDARY, see Syphilis—s. Tertiary, see Syphilis.

SYPHILISATION, Syphilization.

SYPHILISMUS, Syphilis.

SYPHILITIC, *Syphiliticus*, (F.) *Syphilitique, Vérolique, Vérole*. Belonging, or relating to, or affected with, syphilis; (F.) *Syphilité*.

SYPHILIZATION, *Syphilisation*, (F.) *Syphilisation*. Saturation of the system by inoculation with the matter of syphilis. It has been used as a curative procedure in syphilis, as well as a preventive of that disease; and for the cure of other cachexiæ, as cancer.

SYPHILODERMA, gen. *Syphiloderm'ia*, (*syphilis*, and *derma*, 'skin,') A cutaneous eruption of a syphilitic character.

SYPHILOG'ENY, *Syphilogen'ia*, (*syphilis*, and *γενω*, 'to generate,') The origin or generation of syphilis.

SYPHILOG'RAPHER, (*syphilis*, and *γραφω*, 'I describe,') A writer on syphilis.

SYPHILOG'RAPHY, *Syphilograph'ia*, *Syphilid'ography, Syphilidograph'ia*, (*syphilis*, and *γραφω*, 'to describe,') A description of syphilis.

SYPHILOID, *Syphilo'id'es*, (*syphilis*, and *ειδός*,

'resemblance,') Resembling syphilis,—as 'a syphilitic affection.'

SYPHILOID'OGY, *Syphilid'og'ia, Syphilid'ology, Syphilidolog'ia*, (*syphilis*, and *λογος*, 'a discourse,') The doctrine of syphilis and its management. A treatise on syphilis.

SYPHILOMA, (*syphilis*, and *oma*,) A term given by Wagner to what he regards as a specific syphilitic new formation, having a definite structure of cells and nuclei; like as there is a specific carcinomatous new formation in cancer.

SYPHILOMA'NIA, *Syphilidoman'ia*,—vulgarly, *Nod'dle Poz*. A mania, with which some persons are affected, so that they subject themselves to antivenereal treatment, under the erroneous impression that they are affected with syphilis.

SYPHITA, Somnambulism.

SYPHON, see Siphon.

SYRIGMOPHO'NIA, (*σyrigma*, 'piping,' and *φωνη*, 'voice,') A piping state of the voice.

SYRIGMUS, (*σyrigma*,) Tinnitus aurium.

SYRINGA, (from *syrix*,) Eustachian Tube. *Fistula, Syringe, Trachea*—s. *Clysmatica, Syringe (clyster,)*

SYRINGA VULGA'RI, *Li'lac vulga'ris, Crum'mon Li'lac, Ord. Oleaceæ*. The leaves and fruit are bitter and somewhat acrid, and have been used as tonics and antiperiodics.

SYRINGE, *Syr'inx, Syrin'ga, Si'pho, Siphon'culus, Squirr*, (F.) *Seringue*, (from *σyrigē*, gen. *σyrigēs*, 'a pipe,') A familiar apparatus through which any liquor is injected.

Various forms of syringe are used in medicine and surgery.

SYRINGE, CLYSTER, *Syrin'ga Clysmat'ica, Clystant'um*, (F.) *Clyso Pompe*. A syringe for administering clysters.

SYRINGE, CUPPING, *Antlia sanguinis*.

SYRINGES AÉRIFERI, 'air tubes,' see Bronchia.

SYRINGICUS, (from *syrix*,) *Fistulosus*.

SYRINGITIS EUSTACHII, (*syringa*, and *itis*,) Inflammation of the Eustachian tube.

SYRINGMUS, (*σyrigma*,) Tinnitus aurium.

SYRINGODES, (*σyrigodes*,) *Fistulosus*.

SYRINGOTOME, *Syringot'oma, Syringot'omum, (syrix, 'fistula,' and τρυγω, 'to cut,')* A surgical instrument, formerly used in the operation for *fistula in ano*. It is a bistouri, concave at its edge, and terminated by a long, flexible, probe-pointed stylet. This stylet or probe was first introduced by the external opening of the fistula into the rectum, whence it was drawn externally through the anus, carrying with it the cutting edge of the instrument, which divided the intermediate parts.

SYRINGOT'OMY, *Syringotom'ia*. The operation for fistula in ano by incision.

SYRIX, gen. *Syring'is*, (*σyrigē*, gen. *σyrigēs*,) Eustachian tube, *Fistula, Syringe, Trachea*—s. *Hiera, Vertebral column*.

SYRMAISMUS, (from *συναίω*, 'I evacuate,') A gentle evacuation by vomiting or stool.

SYROP, Syrup—s. *d'Ail, Syrupus allii*—s. *d'Armoise composé, Syrupus armoisiæ*—s. *d'Capillaire*, see *Adiantum Capillus veneris*—s. *d'Chicorée composé, Syrupus de chicorée compositus*—s. *des Cinq racines, Syrupus e quinque radicibus*—s. *de Coquelicot, Syrupus rheodæ*—s. *de Coralline*, see *Corallina Corsicana*—s. *d'Citronier, Syrupus de sarsaparilla et sennæ compositus*—s. *d'Ether, Syrupus cum æthere sulphurico paratus*—s. *de Gingembre, Syrupus zingiberis*—s. *de Guimauve, Syrupus althææ*—s. *de Longue vie, Mellitum de mercuriali compositum*—s. *de Mercure, Syrupus de mercurio mediante gummi*—s. *de Mon de veau*, see *Syrupus zingiberis*—s. *de Mûres, Syrupus mori*—s. *de Nerprun, Syrupus rhamni*—s. *des Pavots blancs, Syrupus papaveris*—s. *de Pommes composé, see Syrupus sennæ*—s.

de Quinquina avec le Vin, Syrupus de kinâ kinâ cum vino—s. de *Roses pâles*, Syrupus rose—s. de *Roses rouges*, Syrupus rosæ—s. Simple, Mel despumatum—s. *Végétal de Velno*, Syrup, vegetable, of Velno—s. de *Vinaigre*, S. aceti.

SYR'UP, Sir'up, Syr'pus or Syr'upus Serap'um, Serap'um, (F.) Syrop, Syrop, Saccharolê liquide, Saccharhydrolê, (derived, according to some, from *apo*. 'I draw,' and *os*, 'juice;' but more probably, from the Arabic *Sirap*, *Sirab*, or *Schurab*, which signifies 'a potion.') A saturated solution of sugar in water, either simply, or united with some vegetable or other principle. Syrups, in general, require two pounds of sugar to the pint, and they are judged to be sufficiently boiled when a portion, taken up in a spoon, flows like oil. In the Pharmacopœia of the United States, it was directed, that syrups, whose density is not precisely determined by the process, should have the s. g. 1.261 when boiling, and about 1.319 at ordinary temperatures. Syrups should be kept in small bottles in a cool place. They are chiefly used to render medicines palatable. See Syrupus simplex.

SYRUP OF ALMONDS, Syrupus amygdalæ.

SYRUP, ANTISCORBUTIC, Syrupus antiscorbuticus—s. Balsamic, Syrupus toltanatus.

SYRUP OF BLACKBERRY ROOT, Syrupus Rubi—s. of Buckthorn, Syrupus rhamni—s. of Calf's lungs, see Syrupus zingiberis—s. of Capillaire, common, Syrupus simplex—s. of Cinchona with wine, Syrupus de kinâ kinâ cum vino—s. of Cinchonia, Syrupus cinchonise—s. Citric acid, Syrupus acidi citrici—s. of Clove July flower, Syrupus caryophylli rubri—s. of Clove pinks, Syrupus caryophylli rubri—s. of Cochineal, Syrupus coccii—s. of Emetine, Syrupus emetine—s. of Ether, Syrupus cum æthere sulphurico paratus—s. of Five roots, Syrupus e quinque radicibus—s. of Garlic, Syrupus allii—s. of Ginger, Syrupus zingiberis—s. of Gum Arabic, Syrupus acaciæ.

SYRUP, HIVE, Syrupus scillæ compositus.

SYRUP OF HONEY, Mellitum simplex—s. of Iodide of iron, Liquor Ferri iodidi—s. of Ipecacuanha, Syrupus ipecacuanhæ—s. of Jalap, Syrupus jalapæ—s. of Lactucarium, Syrupus Lactucarii—s. of Lemon, Syrupus limonis—s. of Marshmallow, Syrupus althææ—s. of Meadow saffron, Syrupus colchici autumnalis—s. of acetate of Morphia, Syrupus morphiæ acetatis—s. of muriate of Morphia, Syrupus morphiæ muratis—s. of Mugwort, compound, Syrupus aromaticus—s. of Mulberries, Syrupus mori—s. of Orange flower, Syrupus Aurantii florum—s. of Orange juice, see Syrupus aurantii—s. of Orange peel, Syrupus aurantii—s. of Orgeat, Syrupus amygdalæ—s. of Phosphate of iron, see Ferri phosphas—s. of Poppies, Syrupus papaveris—s. of Poppy, red, Syrupus rhæados—s. of Quinia, Syrupus quiniæ—s. of Rhatany, Syrupus krameris—s. of Rhubarb, Syrupus rhei—s. of Rhubarb, aromatic, Syrupus rhei aromaticus—s. of Rhubarb, spiced, Syrupus rhei aromaticus—s. of Roses, Syrupus rosæ—s. of Roses, red, Syrupus rosæ gallicæ—s. of Saffron, Syrupus croci—s. of Sarsaparilla, Syrupus sarsaparillæ—s. of Sarsaparilla and senna, compound, Syrupus de sarsaparillâ et sennâ compositus—s. of Sarsaparilla, compound, see Syrupus de sarsaparillâ et sennâ compositus—s. of Sarsaparilla, Indian, Syrupus hemidesmi—s. of Scammony, Syrupus de scammonio—s. of Seneka, Syrupus senegæ—s. of Senna, Syrupus sennæ—s. Simple, Syrupus simplex—s. of Squill, Syrupus scillæ—s. of Squill, compound, Syrupus scillæ compositus—s. of Succory, compound, Syrupus de cichorio compositus—s. of Tolu, Syrupus toltanatus.

SYRUP, VEGETABLE, OF VELNO, (F.) *Sirap végétal de Velno*. (Made of roots of burdock and dandelion, leaves of spearmint, senna, coriander, liquorice, sugar, and water.) It has been prescribed as

a eutrophic, especially in syphilis; but, like the syrups of sarsaparilla, is, perhaps, mainly indebted for its efficacy to the sugar. The corrosive chloride of mercury is often added to it.

SYRUP OF VINEGAR, Syrupus aceti—s. of Violets, Syrupus violæ—s. of Wild cherry bark, Syrupus Pruni Virginianæ.

SYRUPUS, S. simplex.

SYR'UPUS ACA'CIE, Syr'up of Gum Ar'abic, (Acacia, 3ij, Sacchar, 3xiv, Aq. f 3viij. Dissolve the gum in the water; add the sugar, and strain. Ph. U. S.) A good pectoral syrup. Dose, f 3j to f 3ij.

SYRUPUS ACETI, S. Ac'idi Aceto'si. Syr'up of Vin'egar, (F.) Syrop de Vinaigre. (Aceti, Oijss, Sacch, purif. lbjss. Boil to form a syrup. Ph. E.) It is refrigerant and antiseptic. Dose, f 3j to f 3ij.

SYRUPUS ACIDI ACETOSI, S. aceti.

SYRUPUS AC'IDI CIT'RICI, Syr'up of Cit'ric Ac'id. (Acid. citric. pulv. 3ij; Ol. Limon. Mjv; Syrup. Oij. Rub the acid and oil with an ounce of the syrup; add the rest of the syrup, and dissolve with a gentle heat. Ph. U. S.) Virtues the same as those of the Syrupus Limonis.

SYRUPUS CUM ÆTHERE SULPHURICO PARATUS, Syr'up of E'ther, (F.) Syrup d'ether. (Very Pure sugar, 1000 p.; Distilled water, 500 p.; dissolve in the cold, and strain; add Sulphuric ether, 48 p. Shake for five or six days, and draw off the limpid syrup. Ph. L.) Properties like those of ether.

SYRUPUS AL'LII, Syrup of Gar'lic, (F.) Syrop d'ail. (Allii recent. concis. 3vj, Acet. destillat, Oj, Sacchar. lbj. Macerate the garlic in the vinegar, in a glass vessel, for four days; then express the liquor and set it by, that the dregs may subside; lastly, add the sugar to the clear liquor, and proceed in the manner directed for syrup. Ph. U. S.) Use, same as that of garlic. Dose, f 3j to f 3iv.

SYRUPUS ALTHÆÆ, S. ex Althæâ seu de Althæâ seu Althææ officinalis, Syrup of Marsh-mallow, (F.) Syrop de Guinauve. (Althææ rad. recent. cont. lbss, Sacch. pur. lbj, Aquæ, Oiv. Boil the root in the water to one-half, and press out the liquor; when cold, defecate; add the sugar, and boil. Ph. L.) Emollient and demulcent. Dose, f 3j to f 3ij.

SYRUPUS AMOMI ZINGIBERIS, S. zingiberis.

SYRUPUS AMYGDALÆ, Syrup of Almonds, S. of Or'geat, Or'geat. (Amygdal. dulc. lbj, Amygdal. amar. 3iv, Aquæ, Oij, Sacchar. lbvj. Blanch the almonds, and rub them in a mortar to a very fine paste, adding, during the trituration, three fluid-ounces of the water, and a pound of the sugar. Mix the paste thoroughly with the remainder of the water; strain with strong expression; add the remainder of the sugar to the strained liquor, and dissolve it with the aid of a gentle heat. Strain the syrup through fine linen, and having allowed it to cool, put it into bottles, which must be well stopped, and kept in a cool place. Ph. U. S.) Demulcent.

SYRUPUS ANTICAUTOTICUS, see Anticaustotic.

SYRUPUS ANTISCORBUTICUS, S. de Armora'ciâ compos'itus, Antiscorbutic Syrup. (Leaves of Cochlearia, Buckbean, Water-cresses, Horse-radish root and Bitter oranges, each lbj, Cinnamon, 3iss. Cut, and pour on White wine, lbiv. Macerate for two days; distil over Oj in a sand-bath; add to this Sugar, lbj, and form a syrup. Again, take what remains in the cucurbit: pass it through cloth; let it rest; decant, and add Sugar, lbj; make into a syrup, and clarify with white of egg. Mix it with the first while yet warm. Ph. P.) Antiscorbutic, alterative, tonic, stomachic, &c. Dose, f 3j to f 3ij.

SYRUPUS DE ARMORACIÂ COMPOSITUS, S. antiscorbuticus.

SYRUPUS AROMATICUS, *S. de Artemisiâ compositus*, Compound Syrup of Mugwort, (F.) *Syrup d'arumaise composé*. (Tops of artemisia vulgaris, in flower, 3vj, Bulb root, Liquidum leuciscum, and Fennel, each 3ss, Pennyroyal, Catmint, Savine leaves, each 3vj, Marjoram, Hyssop, Pyrethrum, Rue, Basil, each 3ij, Aniseed and Cinnamon, each 3ix; bruise and macerate in Hydromel, lbviii; distil over Oss; add to this a pound of sugar. Again, take the liquid that remains in the cucurbit, strain it gently; and add Sugar, lbv. Then mix the two. Ph. P.) Tonic, emmenagogue, &c.

SYRUPUS DE ARTEMISIÂ COMPOSITUS, *S. antiscorbuticus*—s. Asparagi, see Asparagus.

SYRUPUS AURANTII, *S. Aurantiorum seu Citri Aurantii seu Corticis Aurantii seu Aurantii Corticis* (Ph. U. S.) seu *Corticibus Aurantiorum seu de Corticibus Aurantiorum*, Syrup of Orange peel. (Aurant. cort. cont. 3ij, Aq. ferr. Oj, Sacch. lbjss. Macerate the peel in the water for twelve hours. Add the sugar. Ph. U. S. A modified form is in Ph. U. S. 1863.) In the British Pharmacopœia, (1864,) it is made by mixing Tincture of orange peel, f3j with Syrup, f3vij. Stomachic.

Dose, f3j to f3ij.
A Syrup of Orange Juice, Orange Sherbet, Syrupus e succo Aurantiorum, may be made by adding white sugar, lbj to lbj of the juice of oranges, strained and clarified.

SYRUPUS AURANTII FLOREUM, Syrup of Orange Flowers, *S. Aurantii Floris* (Ph. B. 1864.) (Aq. Aurant. flor. f3vj; Sacchar. 3xxxvj; Aq. destillat. f3xv. Dissolve the sugar in the water and raise to the boiling point. When cold, mix with the orange flower water, and strain. Ph. U. S. 1863.)

SYRUPUS BALSAMICUS, *S. toluatanus*.

SYRUPUS CARYOPHYLLI RUTUM, *S. Infusio nis foetum caryophyllorum*, *S. Dianthi Caryophylli*, Syrup of Clove July flower or of Clove Pinks. (Petalarum dianthi, Caryoph. recent., unguibus resectis, lbj, Aquæ bull. Oiv, Sacch. purif. lbvij. Ph. E.) Aromatic.

SYRUPUS CASSIE SENNE, *S. senne*.

SYRUPUS DE CICHORIO COMPOSITUS, *S. de Rheo*, Compound Syrup of Succory, (F.) *Syrup de chicorée composé*. (Root of succory, 3vj, Leaves of succory, 3ix, Leaves of fumitory and scolopendrium, each 3ij, Alkeengi berries, 3ij. Boil in Water Oxiv to Oxij; pass through cloth, and dissolve in it White sugar, lbv. Boil till it has the consistence of syrup. Again, take Water, Oviij, Rhubarb, 3vj, Yellow sanders and Cinnamon, each 3ss. Infuse for twenty-four hours; pass through lnen, and add the preceding syrup. Ph. P.) It is laxative and stomachic. Dose, f3j to f3ss.

SYRUPUS CINCHONÆ CUM VINO, *S. de kinâ kinâ cum vino*.

SYRUPUS CINCHONÆ, Syrup of Cinchona. (Simple syrup, one pound; Sulphate of cinchonia, nine grains. Mix.) Six spoonfuls are commonly sufficient to arrest an intermittent.

SYRUPUS CITRI AURANTII, *S. aurantii*—s. Citri medicæ, *S. Limonium*.

SYRUPUS COCCII, Syrup of Cochineal. (Cocc. cont. 3iv, Aq. destillat. bullient. Oj (Imp. meas.), Sacchar. lbij, vel q. s. Sp. rectificat. f3iiss, vel q. s. Ph. L.) Chiefly used as a colouring agent.

SYRUPUS COLCHICI AUTUMNALIS, *S. Colchici*, Syrup of Meadow Saffron. (Rad. colch. autumn. recent. 3j, Acidi acetosi, f3xvi, Sacch. pur. 3xxvj. Macerate the root in the vinegar for two days, occasionally shaking the vessel; then strain with gentle expression. To the strained fluid add the sugar, and boil to a syrup. Ph. E.) Diuretic, antiarthritic. Dose, f3j to f3ss, or more.

SYRUPUS CORTICIS AURANTII, *S. aurantii*—s. e Succo citrionum, *S. limonium*.

SYRUPUS CROCI, Syrup of Saffron. (Croc-

stigmat. 3j, Aquæ ferrent. Oj, Sacch. purif. lbiss. Ph. L.) Cardiac. Chiefly used for colouring.

SYRUPUS DIACODION, *Diacodium*—s. *Dianthi caryophylli*, *S. caryophylli rubri*.

SYRUPUS EMETICUS, Syrup of Emetine. (Simple syrup, lbj, Emetine, 16 gr.) In small doses, expectorant; and in large, emetic.

SYRUPUS EMPYREUMATICUS, Melasses—s. Ferri iodidi, Liqueur Ferri iodidi—s. Ferri phosphatis, see Ferri phosphas—s. Fuscus, Melasses.

SYRUPUS HEMIDESMI, Syrup of Indian Sarsaparilla, (Hemidesm. cont. 3iv (avoirdup.), Aq. destillat. bullient. Oj (Imp. meas.), Sacchar. q. s. Ph. D.) A weak preparation of hemidesmus.

SYRUPUS HYDRARGYRI, Syrupus de mercurio mediante gummi—s. Infusio nis florum caryophyllorum, Syrupus caryophylli rubri.

SYRUPUS IPECACUANHÆ, Syrup of Ipecacuanha. (Ipecac. in pulv. crass. 3j, Alcohol. dilut. Oj, Sacchar. lbiss, Aquæ, q. s. Macerate the ipecacuanha in the alcohol for 14 days, and filter. Evaporate the filtered liquor to 6 fluidounces, again filter, and add sufficient water to make the liquid measure a pint; add the sugar, and evaporate by a water-bath to a proper consistence. In the Ph. U. S. 1863, it is directed to be made by mixing Ext. Ipecac. fluid. f3ij with Syrup, f3xxx.) It may also be made by displacement.

SYRUPUS DE JALAPÆ, Syrup of Jalap. (Pulv. Jalapæ, 3x, Sem. coriandr., Fœnicul. dulc. 3ss; Aquæ, f3xijss, Sacch. 3xxv. Ph. P. Cathartic. Dose, f3j to f3ss: for children.

SYRUPUS DE KINÂ KINÂ CUM VINO, *S. Cinchonæ cum vino*, Syrup of Cinchona with wine, (F.) Syrup de quinquina avec le vin. (Cort. cinchonæ officinalis, 3ij, Ext. cinchonæ, 3vj, Vini albi (Vin. Lunel, seu cupressi ei similis.) Oj, Alcohol (s. g. 985 to 915.) f3j, Sacchar. alb. lbiss. Bruise the cinchona; gradually add the alcohol, so as to form a soft mass; pour on the wine; macerate for two days, and strain. Dissolve the extract, add the sugar, and make a syrup. Ph. P.) Dose, f3ss to f3j.

SYRUPUS KRAMERIE, Syrup of Rhatany. (Ext. Kramer. 3ij, Aquæ, Oj, Sacchar. lbjss. Dissolve the extract in the water, and filter; then add the sugar, and form into syrup. Ph. U. S. It may also be made by displacement from Kramer. in pulv. crass. lbj, Sacchar. lbiss, Aq. q. s.) An adjunct to astringent mixtures; or given in diarrhœa, a teaspoonful to a child.

SYRUPUS LACTUCARI, Syrup of Lactucarium. (Lactuc. 3j, Syrup, f3xiv; Alcohol. dil. q. s. Rub the lactucarium with enough of diluted alcohol to give a syrupy consistence. Pass half a pint of tincture by percolation; evaporate, at a heat not above 160°, to f3ij; mix with the syrup previously heated, and strain while hot. Ph. U. S. 1863.) Anodyne: but not of much efficacy.

SYRUPUS LIMONIS, *S. Limonium*—s. Succo limonis, *S. Limonium*.

SYRUPUS LIMONUM, *S. Limonis*, (Ph. U. S.) seu citri medicæ seu e succo citrionum seu e succo limonium seu succo limonis, Syrup of Lemon, Lemon Syrup. (Limon. succ. colat. Oj, Sacchar. lbv, Aq. Oj.) Cooling and agreeable.

SYRUPUS MECONIS, *S. papaveris*—s. de Melle, Mellitum simplex.

SYRUPUS MERCURIO MEDIANTE GUMMI, *S. Hydrargyri*, Mercurius gummosus Planchetii. (F.) Syrup de mercure, Mercure gommeux de Planchet. (Hydrargyri pur. 3j, Gum. acacis. 3ij, Syrup diacod. 3ss. Rub together till the mercury is extinguished. Ph. P.) Possesses the properties of blue pill. Dose, gr. ij to viij, or more.

SYRUPUS MORI, *S. e succo mororum seu succo mori*, Syrup of Mulberries, (F.) Syrup de mûres. (Mori succ. colat. Oj, Sacchar. purif. lbj. Ph. L.) Gratefully refrigerant.

SYRUPUS E SUCCO MORORUM, *S. mori*.

SYRUPUS MORPH'Æ ACETAT'IS, *Syrup of acetate of morphia*. (*Liq. morphia acet. f 3j, Syrup, f 3xv. Ph. D.*) Dose, *f 3ss to f 3j*, as an opiate.

SYRUPUS MORPH'Æ MURIAT'IS, *Syrup of muriate of morphia*. (*Liq. morph. muriat. f 3j; Syrup, f 3xvj. M. Ph. D.*) Dose, *f 3ss to f 3j*, as an opiate.

SYRUPUS MORPH'Æ SULPHA'TIS may be made in the same way.

SYRUPUS NIGER, Melasses.

SYRUPUS PAPAVER'IS, *S. papaveris somniferi seu papaveris al'bi seu e meconio seu de meconio, Diacodium, Syrup of poppies*, (*F.*) *Syrup de papav. blancs ou Diacode. (Papav. capsul. exsicc. et contus. exemptis seminibus. 3xiv, Sacch. pur. lbj, Aq. fervent. cong. ijs.* Macerate the capsules in water for 12 hours; boil to a gallon, and express strongly. Boil the liquor to Oij; and strain while hot. Defecate by rest, and boil to Oj, adding the sugar so as to form a syrup. *Ph. L.*) One ounce contains about one grain of opium. Anodyne and demulcent.

The Dublin college has a SYRUPUS OPII, called *Common Syrup of Poppies*, which is made by dissolving 18 grains of the *Watery Extract of Opium* in 18 ounces of boiling water. Properties those of opium and syrup.

SYRUPUS DE PAPAVERE ERRATICO, *S. rhœados* — *s. de Pomis compositus*, see *S. sennæ* — *s. Proteæ*, see *Protea mellifera*.

SYRUPUS PRU'NI VIRGINIA'NÆ, *Syrup of Wild cherry bark. (Pruni Virginian. in pulv. crass. 3v, Sacchar. xxvij, Aquæ, q. s.* Moisten the bark with water; let it stand for 24 hours; transfer it to a percolator; pour water gradually upon it, until a pint of filtered liquor has passed. Add the sugar in a bottle, and dissolve by agitation. *Ph. U. S.*) Dose, as a demulcent expectorant, *f 3j to f 3ij*.

SYRUPUS E PULMONIBUS VITULINIS, see *Syrupus Zingiberis*.

SYRUPUS QUI'NÆ, *Syrup of Quinia*. (*Simple syrup, 2 pints, Sulphate of quinia, 64 grains.*) Six spoonfuls are commonly sufficient to arrest an intermittent.

SYRUPUS E QUIN'QUÈ RADICIBUS, *Syrup of five roots*, (*F.*) *Syrup des cinq racines. (Rad. Apii, Fœniculi, Apii petroselinii, 3v. Infuse for 24 hours in boiling water, Oiss, and strain. Rad. asparag., Ruscæ aculeati, 3v, Aquæ, Ovj. Boil and reduce to half; adding, towards the end, the residue of the preceding infusion. Boil again for some minutes; strain, and add White sugar, lbj. Ph. P.)* Aperient and diuretic. Used in obstructions of the liver, spleen, and mesentery; and in dropsy, &c. Dose, *f 3ss to f 3j*, or more.

SYRUPUS RHAM'NI, *S. Rhamni cathartici seu de Spina cervi' nâ seu spina cervi' nâ, Syrup of Buckthorn*, (*F.*) *Syrup de nerprun. (Rhamn. buccar. succ. recent. Oiv. Zingib. rad. concis., Pimentæ bacc. contrit., 3ss, Sacch. purif. lbiss.* Defecate the juice for 3 days, and strain. To a pint of this, add the ginger and pimento; macerate for 4 hours, and strain; boil what remains to Oiss. Mix the liquors and add sugar, so as to form a syrup. *Ph. L.*) Cathartic, but apt to gripe. Dose, *f 3iv to f 3ij*.

A factitious syrup has been made of aloes, dissolved in simple syrup.

SYRUPUS RHE'I, *Syrup of Rhu'barb. (Rhei, in pulv. crass., 3ij; Alcohol, Oss; Aquæ, Oiss; Sacchar. lbj.* Mix the alcohol and water; pour *f 3iv* on the rhubarb, mixed with an equal bulk of sand, and allow the whole to stand four hours; put the mass into a percolator, and gradually pour on the alcohol and water. By means of a water-bath, evaporate to *f 3xij*; add the sugar, and make a syrup. *Ph. U. S. 1851.*) In the *Ph. U. S.* for 1863, it is made by mixing *Ext. Rhei fluid. f 3ij*

with *Syrup, f 3xxix*. Dose to a child, as a laxative, one or two teaspoonfuls.

SYRUPUS RHEI AROMAT'ICUS, *Aromatic or Spiced Syrup of Rhu'barb. (Rhei cont. 3iiss; Caryophyl. cont., Cinnam. cont. 3ss; Myrist. cont. 3ij; Alcohol. dilut. Oij; Syrup, Ovj.* Macerate the rhubarb and aromatics in the diluted alcohol for 14 days, and strain; then, by means of a water-bath, evaporate the liquor to a pint, and while still hot, mix with the syrup previously heated. *Ph. U. S.*) It may also be made by the process of displacement.

The aromatic syrup of rhubarb is a gentle carminative cathartic. Dose to a child, *f 3j*; to an adult, *f 3j*.

SYRUPUS DE RHEO, *Syrupus de Cichorio compositus*.

SYRUPUS RHE'ADOS, *S. de papav'erè erratico seu papaveris erratici, Syrup of Red Pop'py*, (*F.*) *Syrup de coquelicot. (Rhead. petal. recent. lbj, Aq. fervent. f 3xvij, Sacch. purif. lbiss.* To the water, heated in a water-bath, add the petals. Remove the vessel; macerate for 12 hours; express; defecate, and add sugar to form a syrup. *Ph. L.*) Chiefly used to give colour.

SYRUPUS ROSÆ, *S. rosa'rum solutivus seu e rosa's sic'cia seu rosa'e centifoliae seu rosa'eus solutivus, Syrup of Ro'ses*, (*F.*) *Syrup de roses pâles. (Rosæ. centif. petal. exsicc. 3vij, Sacch. purif. lbvj, Aq. fervent. Oiv.* Macerate the petals for 12 hours. Evaporate the strained liquor to lbiss. Add the sugar. *Ph. L.*) Gently laxative. Dose, *f 3ij to f 3iss*, or more.

SYRUPUS ROSÆ GAL'LICÆ, *Syrup of Red Ro'ses*, (*F.*) *Syrup de roses rouges. (Rosæ Gallic. petal. exsicc. 3vij, Aquæ bull. Oij, Sacch. purif. lbvj; prepared like the last. Ph. E.)* In the *Ph. U. S.* for 1863, it is directed to be made by moistening red rose, in powder, with diluted alcohol, and, in a percolator, pouring on diluted alcohol until a certain portion of the tincture has passed; evaporating this to a certain point, and adding water; then, having added the sugar, dissolving it with a gentle heat. Mildly astringent. Added to stomachic infusions.

SYRUPUS RU'BI, *Syrup of Blackberry Root. (Rubi, in pulv. 3vij, Syrup. Oiss. Alcohol. dilut. q. s.* Moisten the powder with *f 3iv* of the alcohol; and, in a percolator, pour on the alcohol until a pint and a half of the tincture has passed. Evaporate this to half a pint, and mix, while hot, with the syrup previously heated, and strain. *Ph. U. S., 1863.*) Gently astringent, and used as an addition to prescriptions for diarrhoea of children.

SYRUPUS SACCHARI, *S. Simplex*.

SYRUPUS SARSÆ, *S. Sarsaparillæ*.

SYRUPUS SARSAPARIL'Æ, *S. Sar'sæ, S. Sar'æ, Syrup of Sarsaparill'a. (Rad. Sarsap. cont. lbj, Aquæ bullient. cong., Sacch. purif. lbj.* Macerate the root for 24 hours; boil to Oiv. Strain while hot; add the sugar, and boil. *Ph. L.*) Used in the same cases as the root. Dose, *f 3j to f 3iv*, and more.

SYRUPUS DE SARSAPARIL'LÂ ET SEN'NÂ COMPOSITUS, *Compound Syrup of Sarsaparill'a and Sen'na*, (*F.*) *Syrup de Cusinéier. (Rad. Sarsap. concis. lbj. Infuse for 24 hours in Warm water Oxij; boil for a quarter of an hour; pass with expression, and boil again the residuum in Ox of water. Reduce this to Ovj, and repeat the operation 2 or 3 times. Mix all the liquors and boil them together with Flowers of borage, White roses, Senna leaves, and Aniseed, each 3ij; reduce to half; pass, and add Honey lbj. Boil to a syrup. Ph. P.)* It is considered to be alterant, depurative, and sudorific. Dose, *f 3ss to f 3ij*. Corrosive chloride of mercury is sometimes added to it.

A similar syrup, under the name SYRUPUS SARSAPARIL'Æ COMPOSITUS, *Compound Syrup of Sarsaparill'a*, is directed in the Pharmacopœia

of the United States. (*Saccharill.* cont. *ibij.* *Lign. guaiac.* rasur. *℥ij*; *Rose centifol.* *Sennæ*, *Glycyrrhizæ* cont. *āā ℥ij*; *Ol. Sassafras*, *Ol. Anisi*, *āā ℥v*; *Ol. Gaultheriæ* m. *ij*; *Alcohol.* dilut. *Ox*; *Sacchar.* *℔vij*. Macerate the sarsaparilla, guaiacum wood, roses, senna, and liquorice root in the diluted alcohol for 14 days; express and filter. Evaporate the tincture in a warm bath to four pints; filter; add the sugar, and form into syrup. Lastly, having rubbed the oils with a small quantity of the syrup, mix them thoroughly with the remainder. Ph. U. S.)

It may also be made by the process of displacement.

SYRUPUS SARZÆ, S. Sarsaparillæ.

SYRUPUS DE SCAMMONIO, *Syr'up of Scammony*, *'Pulv. Scammon.* *℥ss*, *Sacch.* alb. *℥iv*, *Alcohol.* (s. g. '985 to '915) *f℥viii*. Heat the mixture moderately: then set fire to the alcohol with a lighted taper; agitate constantly, until the flame is extinguished. Suffer it to cool; pass through cloth, and add *Syrup of Violets*, *f℥iv*. Ph. P.) *℥j* contains 18 grains of scammony. Cathartic. Dose, *f℥j* to *f℥ss*.

SYRUPUS SCILLÆ, *S. scilliticus* seu *scill'æ maritimæ*, *Syr'up of Squills*. (Acet. *scill'æ* *Oj*, *Sacch.* *℔ij*, Ph. U. S.) Diuretic, expectorant, and emetic. Used in the same cases as the oxymel.

SYRUPUS SCILLÆ COMPOSITUS, *Ml Scill'æ Compositum*, *Compound Honey of Squill*, *Compound Syr'up of Squill*, *Hive Syrup*. (*Scillæ* cont.: *Senegæ* cont. *āā ℥iv*; *Antim.* et *Potassæ Tart.* gr. *xlviij*; *Aquæ Oiv*; *Sacchar.* *℔iiiss*. Pour the water upon the squills and seneka, and, having boiled to one-half, strain and add the sugar. Then evaporate to three pints, and while the syrup is still hot, dissolve in it the tartrate of antimony and potassa. It may also be made by the process of displacement. Ph. U. S.)

It is often given as an emetic in cases of croup, and in catarrh, especially in children. Dose, to infants, 10 drops to a drachm, repeated every 15 minutes, if vomiting be desirable.

SYRUPUS SENEGÆ, *Syrup of Seneka*. (*Senega* cont. *℥iv*; *Aquæ Oj*; *Sacchar.* *℔j*. Boil the water with the seneka to one half, and strain; then add the sugar, and form into syrup. Ph. U. S.)

It may also be formed by the process of displacement.

This is a favourite stimulant expectorant with many practitioners. Dose, *f℥j* or *f℥ij*.

SYRUPUS SENÆ, *S. cas'siæ sen'æ*, *Syr'up of Sen'na*. (*Sennæ* *℥ij*, *Sem. feniculi* cont. *℥j*, *Sacch.* pur. *℥xv*, *Aquæ bullient.* *Oj*. Macerate the senna leaves and fennel-seeds with a gentle heat for an hour. Strain: add the sugar, and make into a syrup. Ph. U. S.) Cathartic. Dose, *f℥ij* to *f℥iiss* or more.

The Parisian Codex has a *Syrupus de Sennâ* of a compound character, called *Syrupus de pomis compositus*, (F.) *Syrop de pommes composé*. It contains, also, borage, bugloss, &c. Dose, *f℥ss* to *f℥ij*.

SYRUPUS SIMPLEX, *Syrupus* (Ph. U. S.), *S. Sacchari*, *Saccharum clarificatum*, *Liquarium*, *Simple Syr'up*, *Syr'up*, *Common syrup of capillaire*. (*Sacch. purif.* *℔iiss*, *Aquæ Oj*. Dissolve the sugar: set aside for 24 hours; remove the scum, and pour off the clear liquid. A modified form is given in the Ph. U. S. for 1863.) Used to cover nauseous tastes, &c.

SYRUPUS DE SPINÂ CERVINÂ, *Syrupus rhamni* —s. de *Stœchade communis*, see *Lavandula*.

SYRUPUS TOLUTANUS, *S. Tolutani* seu *toluif'æ balsami* seu *balsamici*, *Balsamic syr'up*, *Syr'up of tolu*. (Tinct. *Tolut.* *f℥iiss*; *Aquæ Oj*; *Sacchar.* *℔iiss*. Mix the tincture with the sugar in coarse powder; expose it on a shallow dish to a gentle heat, until the alcohol has evaporated;

pour the water upon it: heat gradually until the sugar is dissolved, and strain. Ph. U. S.) Use—to give flavour.

SYRUPUS VIOLÆ, *S. violæ odoratæ*, *S. violæ-rum*, *S. e succo violæ-rum*, *Syr'up of Violets*. (Flor. recent. *Viol. odorat.* *℔ij*, *Aq. bull.* *Oviij*, *Sacch.* pur. *℔xv*. Macerate in a covered vessel, for 24 hours: pass, without expression, through linen; add the sugar so as to form a syrup. Ph. E.) Very gently laxative. Dose, *f℥j* to *f℥ij*.

SYRUPUS E SUCCO VIOLARUM, *S. violæ*.

SYRUPUS ZINGIBERIS, *S. amomi zingiberis*, *Syr'up of Ginger*, (F.) *Syrop de gingembre*. (Tinct. *Zingib.* *f℥iv*; *Aquæ Oiv*; *Sacchar.* *℔x*. Mix the tincture with four pounds of the sugar in coarse powder; and expose it in a shallow dish to a gentle heat until the alcohol has evaporated. Add the rest of the sugar, and then the water in a covered vessel; heat gradually until the sugar is dissolved, and strain. To the strained liquor add the sugar, and form a syrup. Ph. U. S.) Ex-citant. Dose, *f℥j* to *f℥ss*. A modified form is contained in the Ph. U. S. (1863.)

N. B. The Parisian Codex contained a few other compound syrups besides those enumerated; amongst them was a *Syrup of calf's lungs*, *Syrupus e pulmonibus vitulinis*, (F.) *Syrop de moule veau*, as an expectorant!

These syrups are made according to the usual rule, and possess the virtues of the substances after which they are named.

SYS, as a prefix, like *Sym*.

SYSDIENES, (*sys*, and *δυναμις*, 'body.') A family of double monsters, of Is. G. St. Hilaire, characterized by the union or fusion of two bodies surmounted by heads completely separate.

SYSPA'SIA, (*συσπαισις*, from *συσπαι* (*sys*, and *σπαι*, 'to draw'), 'to contract.') Clonic spasm, with diminished sensibility and inability of utterance. A genus in Good's Nosology, which includes convulsion, epilepsy, and hysteria.

SYSPASIA CONVULSIO, Convulsion.—s. *Epilepsia*, *Epilepsy*—s. *Hysteria*, *Hysteria*

SYSSARCO'SIS, (*sys*, *σάρξ*, gen. *σάρκως*, 'flesh,' and *osis*.) *Communi'ca os'sium car'neæ*, *Symph'yosis os'sium muscula'ris*, *Connex'io os'sium carno'sa*, *Carna'tio*, *Concarnatio*. Union of bones by means of muscles; such as the union of the os hyoides with the lower jaw, of the scapula with the ribs, &c.

SYSTALSIS, Systole.

SYSTATIC, (*συσταλτικός*) *Systalticus*, *Peristaltic*, (*συσταλῶ* (*sys*, and *στέλλω*), 'I contract.') (F.) *Systaltique*. A *systaltic motion* in a part is one in which there is alternate contraction and dilatation. See *Systole*.

SYSTATICA, (from *συνιστημι* (*sys*, and *ιστημι*, 'to place'), 'to associate.') Diseases affecting several or all of the sensorial powers simultaneously; irritation or inertness of mind, extending to the muscles, or external senses; or of the muscles, or external senses, extending to the mind; —the 4th order, class *Neurotica*, of Good.

SYSTEM, *Syste'ma*, (*συστημα*, from *sys*, and *στημι*, 'I place.') (F.) *Système*. This word, taken in a good sense, means the collection and arrangement of different matters having an analogy to each other, or working towards the same end. In natural history, it means a methodical arrangement of objects for the purpose of facilitating their study. System is, however, sometimes used in an unfavourable sense in physical science, and often means—a purely gratuitous supposition, by which the march of nature is attempted to be universally explained, and to which every thing is made to bend. In anatomy, system is an assemblage of organs, composed of the same tissues, and intended for similar functions; for instance, the nervous system, muscular system, &c.

SYSTEMA, gen. *Syste'matis*, System—s. Vaso-rum, Vascular system.

SYS'TEMATISM. A method, according to which every fact of a science is collected around an opinion, true or false. One who reduces things to such a method is called a *Sys'tematist*.

SYSTEMATIST, see Systematism.

SYSTEMATOLOGY, *Sys'tematolog'ia*, (*sys'te-ma*, and *logos*, 'a discourse.') The doctrine of, or a treatise on, systems.

SYSTÈME, System—s. *Vasculaire à sang noir*, see Vascular—s. *Vasculaire à sang rouge*, see Vascular.

SYSTEM'IC or SYS'TEMIC. Belonging to

the general system—as '*systemic circulation*' in contradistinction to the '*pulmonic*.'

SYS'TOLÈ, (*συστολή*, from *sys*, and *στέλλω*, 'to contract.') *Sys'tol'sis*, *Constrict'io*, *Contract'io* seu *Angusta'tio* seu *Submiss'io* seu *Mic'tio* *cor'dis*. The movement of constriction or contraction of the heart, to give impulse to the blood, or to cause its progression. It is opposed to the *Dias-tole* of the organ.

SYSTOLIC or SYS'TOLIC, *Sys'tol'icus*, (F.) *Systolique*. Relating to the systole of the heart,—as '*systolic murmur*.'

SYTHE, ([Sax.] *pean*, 'to sigh,') *Suspirium*.

SYZYGIIUM NERVORUM, (*συσζυγος*, 'united,' from *sys*, and *ζυγος*, 'a yoke,') *Par nervorum*.

T.

T BANDAGE, *T fas'cia*, *Fas'cia* *T form'is* seu *Helioid'ri*, *Cru'cial* *Ban'dage*. A bandage shaped like the letter T. It consists of a strip of linen, of variable length and breadth, the middle of which is attached, at right angles, to the extremity of another strip, also of variable length and breadth. When two such strips are attached, at a distance from each other, the bandage is called a *double T*. This bandage is used, principally, for supporting the dressings after the operation for fistula in ano, in diseases of the perinæum, groin, anus, &c.

T FASCIA, T Bandage.

TABAC, 'tobacco,' *Nicotiana tabacum*—t. *des Montagnes*, *Arnica montana*—t. *des Savoyards*, *Arnica montana*—t. *des Vosges*, *Arnica montana*.

TABACUM, 'tobacco,' *Nicotiana tabacum*.

TAB'ANUS, 'the horsefly.' A troublesome fly in the Hudson's Bay territory, called, by the *voyageurs*, *Bulldog*, which makes a perforation like a leechbite, from which it sucks the blood.

TABARDILLO (S.). A kind of highly adynamic fever, common in South America.

TABARZET, *Saccharum album*.

TAB'ASHEER, *Tab'aschir*, *Tab'axir*. A substance found in the knots of the bamboo, in the East Indies, as a consequence of a morbid condition of the part. It is distinguished for its great proportion of silicic acid, in combination with a small quantity of mineral and vegetable matters; as well as for its slight refractive property. It has been used in obstruction of the bowels. The sweet juice of the bamboo stalks has also been called *Tabasheer*.

TABEFACATION, (*tabes*, and *facere*, 'to make,') Emaciation.

TABEFY, Emaciate.

TABEL'LA, *Tab'ula*, *Tabula'tum*, *Lozan'ga*, *Lozan'gia*, *Mor'sulus*, *Morsell'us*, *Loz'enge*, *Tes-sell'la*, (F.) *Tablette*, *Lozange*. A solid medicine, composed of a powder, sugar, and mucilage, *Electuarium solidum*; and generally shaped like a disk. See *Pastil*, and *Troch*. The three comprise the *Saccharolés solidæ*.

TABELLA, (dim. of *tabula*, 'a table,') Table.

TABELLÆ ABSORBENTES, T. de *Magnesiâ*.

TABELLÆ AC'IDI OXAL'ICI, T. ad si'tem, *Loz-enges of Oxal'ic Ac'id*. (*Acid. oxalic*, 3j, *Sacch. alb.* 3ss, *Ol. limonis*, gtt. xvij seu gtt. xij. Make into lozenges with *Mucilage of gum tragacanth*. Ph. P.) Lozenges are made in the same way with the citric and tartaric acids. They are refrigerant.

TABELLÆ DE ALTHE'Â, *Loz'enges of Marsh'-mallows*. (*Rad. althææ pulv.*, 3iss, *Sacchar. alb.*

pulv. 3ivss; f. massa. Ph. P.) Demulcent; expectorant.

TABELLÆ ANTIMONIA'LES KUNCKE'LII, T. de *stib'ii sulphure'to*, *Loz'enges of Sulph'uret of Anti-mony*, *Mor'suli antimonial'es* seu *stib'ii seu stibial'ti*, *Trochis'ci stibiati*. (*Amygdal. dulc.* 3j, *Sacch. lbss*, *Sem. cardam.*, *cinnam. pulv.*, *Sulph. antim. præpar.*, aa 3j; fiant *tabellæ*. Ph. P. Each to weigh 10 grains.) Recommended in cutaneous diseases, blennorrhagia, arthritic pains, &c. Dose, 4 or more.

TABELLÆ AD ARDREM VENTRICULI, *Trochisci carbonatis calcis*—t. *Cardialgicæ*, *Trochisci carbonatis calcis*.

TABELLÆ DE CAT'ECHU ET MAGNE'SIÂ, *Loz-enges of Cat'echu and Magne'sia*, (F.) *Tablettes de cachou et de magnésie*. (*P. catech.* 3v, *P. magnès.* 3iv, *P. cinnam.* 3ij, *Sacch. alb. lbss*, *G. tragac.* gr. xij, *Aq. cinnam.* q. s. ut f. massa. Ph. P. Each to weigh 12 gr.) Antacid, absorbent, and stomachic. Dose, 5 to 10 lozenges.

TABELLÆ CINCHO'NÆ, T. de *k'i'nâ k'i'nâ*, *Bark Loz'enges*. (*Ext. cinchon. sicc.* 3ss, *Sacch. 3iv*, *Cinnam.* 3ss, *Mucilag. g. trag.* q. s. ut f. massa. Ph. P.) Tonic.

TABELLÆ DE CITRO ET DIACARTHAMO, T. de *Scammonio et Sennâ*.

TABELLÆ DE FER'RO, *Steel loz'enges*, *Tab'ulæ martial'es* seu *instaurant'es* seu *antichlorot'icæ*. (*Ferri limat. porphyriz.* 3ss, *P. cinnam.* 3j, *P. sacch.* 3v, *G. trag.* et *Aq. cinnam.* q. s., ut fiat massa. Each lozenge to weigh 12 gr. Ph. P.) Tonic, stomachic. Dose, v to x.

Aromat'ic loz'enges of steel. These consist of *sulphate of iron*, with a small proportion of the *Tincture of Cantharides*.

TABELLÆ DE KINÂ KINÂ, T. *cinchonæ*.

TABELLÆ DE MAGNE'SIA, T. *absorben'tes*, *Magne'sia loz'enges*. (*Magnès.* 3j, *Sacchar. alb.* 3iv, *Mucil. g. tragac.* q. s. Ph. P.) Absorbent and antacid.

TABELLÆ DE RHE'O, *Rhu'barb loz'enges*. (*P. rhej.* 3ss, *Sacch.* 3v, *Mucilag. gum. tragac.* q. s.; f. massa. Each to contain 12 grains. Ph. P.) Stomachic and laxative.

TABELLÆ DE SCAMMO'NIO ET SEN'NÂ COMPOS'ITÆ, T. de *c'i'tro et diacar'thamo*, *Compound loz-enges of scam'mony and sen'na*. (*Scammon.* 3ij, *Sennæ* 3ivss, *rhej.* 3iss, *Caryoph.* 3j, *Corticis limon. confect.* 3j, *Sacch.* 3vj; f. massa. Ph. P.) Purgative. Dose, 3j to 3vj, or more.

TABELLÆ AD SITEM, T. *acidi oxalici*—t. ad *Sodam*, *Trochisci carbonatis calcis*—t. de *stibii sulphure'to*, T. *antimoniales Kunckelii*.

TABELLÆ DE SULPH'URE SIM'PLICES, *Sim'plæ*

sulph'ur loz'enges. (*Sulphur* lot. $\frac{3}{4}$ ss, *Sacch.* $\frac{3}{4}$ iv, *Muc. g. tragac.* cum aquâ rosâ præpar. q. s. *Ph. P.*) Expectorant, demulcent, &c. Used in diseases of the chest, and in cutaneous affections. Dose, a lozenge.

TABELLÆ DE SULPHURÆ COMPOSITÆ. *Compound sulph'ur loz'enges.* (*Sulph.* lot. $\frac{3}{4}$ ij, *Acid. benzoic.* gr. xij, *Rad. irid. p.* $\frac{3}{4}$ ss, *Ol. anisi,* gtt. viij, *Sacch.* $\frac{3}{4}$ vss, *G. trag.* q. s. *Ph. P.*) Aperient, expectorant, and stimulant [?]. Dose, 4 a day.

TABERNA MEDICA, (*taberna*, 'a shop,') *Pharmacopolium.*

TABERNEMONTANA UTILIS, (after *Tabernemontanus*, a German botanist.) A tree of British Guiana, of the family *Apocynaceæ*, called, by the natives, *Hya-Hya*, the bark and pith of which yield a milk, which is used by the savages as a refreshing drink.

TABES, gen. *Ta'bis*, *Maras'mus Ta'bes*, (*tabere*, 'to decay;') *Te'kedon*, *Apotex'is*, *Atroph'ia*, *Tab'idness*, *Decline'*, *Consumption*, *Wasting*. Emaciation of the whole body, with general languor, hectic fever, and, for the most part, depressed spirits.

TABES, *Atrophy*, *Phthisis* — *t. Coxaria*, *Ischiophthisis* — *t. Diabetica*, *Diabetes mellitus* — *t. Diuretica*, *Diabetes mellitus*.

TABES DORSALIS, *T. Dorsu'al'is*, *Maras'mus ta'bes dorsa'lis*, *Phthi'sis no'tius seu dorsa'lis seu sic'ca*, *Myelophthi'sis*, *Rhachiomyelophthi'sis*, *Myelatrophi'ia*, *Myelanol'o'sis*, *Ta'bes os'sis sa'cri*. A wasting of the body, attended at first with a pain in the back or loins, and afterwards in the neck or head,—caused, it has been conceived, by too early or too frequent addition to venery.

TABES GLANDULARIS, *T. mesenterica*, *Scrofula* — *t. Ischiadica*, *Ischiophthisis* — *t. Lactea*, *Atrophia lactantium*, see *Galactia*.

TABES MENTERICA, *T. glandula'ris seu scrofulo'sa*, *Atroph'ia mesenter'ica seu infant'um seu glandula'ris seu infant'ilis seu dor'si*, *Parabys'ma mesenter'icum*, *Ma'cies Infant'um*, *Pedatrophi'ia*, *Scrof'ula seu Rachial'gia seu Physco'nia mesenterica*, *Myelophthi'sis sic'ca*, *Fe'bris hec'tica infant'um*, *Phthi'sis mesara'ica*, *Mesenterophthi'sis*, *Mesenter'ic Disease*, (F.) *Carreau*, *Chartre*, *Scrofulæ ou Écouelles mésentériques*, *Étiologie ou Rachialgie ou Atrophie ou Physaconie mésentérique*, *Entéro-mésentérique*, *Tubercules des ganglions mésentériques*, *Phthisie mésentérique*. An engorgement and tubercular degeneration of the mesenteric glands, followed by emaciation and general disorder of the nutritive functions. The disease is often owing to inflammation or irritation of the lining membrane of the intestines, giving occasion to enlargement of the glands of the mesentery,—as injury to the upper or lower extremities causes inflammation, and enlargement of the lymphatic glands of the axilla and groin. It occurs, particularly, in children of a serofulous diathesis; and in those who are weaned too early, or fed on indigestible substances. Diarrhœa, emaciation, anorexia, or immoderate appetite; hardness and swelling of the abdomen; and, towards the end, hectic fever, are the chief symptoms of this affection; the prognosis of which is unfavourable. On opening the bodies of those who die of the disease, the mesenteric glands are found disorganized, tubercular, or in a more or less advanced state of supuration. The treatment is the same as in scrophulosis. Frictions may be added over the abdomen and spine; and mercury, in small doses, so as to excite a new action in the system. The food must be easy of digestion, and the bowels be kept soluble.

TABES NUTRICUM, *Atrophia lactantium*, see *Galactia* — *t. Ossis sacri*, *Tabes dorsalis* — *t. Pectoræ*, *Rachitis* — *t. Pulmonalis*, *Phthisis pulmo-*

nalis — *t. Scrofulosa*, *T. mesenterica* — *t. Senum*, *Marasmus senilis*.

TABESCERE (*tabere*, 'to decay,') *Emaciate*.

TAB'ID, *Tab'idus*, (F.) *Tabide*. Relating or appertaining to tabes. Wasted by disease.

TABIDNESS, *Tabes*.

TABIF'IC, *Tabif'icus*, (F.) *Tabifique*, (*tabes*, and *facere*, 'to make,') That which causes consumption or phthisis.

TAB'LE, *Tab'ula*, *Tabell'a*, *Tabula'tum*, *Lam'ina*, *Pa'ries*, *Squa'ma*. A name given to the plates of compact tissue, which form the bones of the cranium. Of these, one is *external*; the other *internal*, and called *Tab'ula vi'trea*, (F.) *Lame vitrée*, on account of its brittleness.

TABLES DE MORTALITÉ, *Mortality*, bills of.

TABLET, 'a small table,' *Trochiscus* — *t. Vo'tive*, *Tabula votiva*.

TABELLETTE, 'tablet,' *Tabella*.

TABLETTES DE CACHOU ET DE MAGNÉSIE, *Tabellæ de catechu et magnesiâ*.

TABLIER, 'an apron,' *Ventrale cutaneum*.

TABOURET, *Thlasi bursa*.

TABULA, *Tabella*, *Table* — *t. Vitrea*, see *Table*.

TAB'ULA VO'TIVA, *Vo'tive tab'let*. A tablet, hung up as an offering — *anathe'ma* — in the ancient temples by the sick, descriptive of their diseases, and mode of cure; or by those who had been preserved from some calamity.

TABULÆ ANTICHLOROTICÆ, *Tabellæ de ferro* — *t. Instanturæ*, *Tabellæ de ferro* — *t. Martiales*, *Tabellæ de ferro*.

TABULATUM, (from *tabula*,) *Tabella*, *Table*.

TABUM, (allied to *tabes*,) *Ichor*, *Sanies*, *Plague*.

TAC, *Influenza*.

TACAMAHACA, *Fagara octandra* — *t. Caragna*, *Caranna*.

TACCA (Malay name) *OCEANICA*, see *Arrow-root*, (*Tahiti*.)

TACHE (F.), 'spot,' *Macula*, see *Taches*.

TACHE CÉRÉBRALE, *Tache méningitique*. A red mark or stain, observable, according to M. Trousseau, in a large number of hydrocephalic children, when pressure is made on the skin.

TACHE EMBRYONNAIRE, 'Embryonic spot,' *Cu'mulus*, *C. germinativus seu prolif'erus*, *A'rea germinati'va*, *Stratum germinativum*, *Blastotro'pha*, *Nu'cleus cicatric'ulæ seu blastoderm'itis*, *Ger'minal cu'mulus*. A name given by M. Coste to a spot seen on the blastoderm about the end of the first week, from which the development of the embryo proceeds.

According to Bischoff, the blastodermic vesicle consists here of a double layer, (F.) *Feuillet* — the upper one the *serous* or *animal*; the lower, the *mucous* or *vegetative*; and between these is afterwards formed the *vascular layer*.

TACHE GERMINATIVE, 'germinal spot,' see *Molecule* — *t. Jaune*, see *Foramen centrale retinae* — *t. Mélanienne*, see *Nævus* — *t. Méningitique*, *Tache cérébrale* — *t. de Naissance*, *Nævus* — *t. Pigmentaire*, *Nævus* — *t. Sanguine*, see *Nævus*.

TACHES (pl. of *tache*) *HÉPATIQUES*, *Chloasma*.

TACHES LAITEUSES, *Plaques laiteuses*, 'milky spots or patches.' White spots occasionally seen on the heart, and regarded by many pathologists as the results of previous pericarditis.

TACHES ROUGES, 'red spots,' — *t. ac Roussour*, *Ephelides*.

TACHETÉ DE ROUSSEURS, ('spotted with freckles,') see *Ephelides*.

TACHYIATER, (*ταχys*, 'quick,' and *ιατρος*, 'a physician,') *Tachyi'atrus*. One who cures speedily.

TACHYIATRIA, (ταχυς, 'quick,' and *iatria*, 'healing,') The art of curing quickly.

TACHYTOMY, *Tachytomyia*, (ταχυς, 'quick,' and *tomē*, 'incision,') The art of cutting or operating rapidly.

TACITURNITY, *Taciturnitas*, *Si'gē*, (from *tacere*, *tacitum*, 'to be silent,') State of a person who does not speak. Morbid taciturnity sometimes exists as a symptom of the neuroses, especially of melancholy.

TACT, *Tactus*, (from *tangere*, *tactum*, 'to touch,') A modification of the sense of touch. Passive judgment by the cutaneous nerves regarding bodies exterior to the skin. Tact is passive; Touch, active.

TACTILE, *Tactilis*. Relating or appertaining to tact or touch.

TACTION, Touch.

TACTUS, Tact, Touch — *t.* Dolorificus, Haptodysphoria — *t.* Eruditus, see Palpation.

TAE, 'toe,' *Digitus pedis*.

TÆDA, *Tæda*, *Dæs*, 'a torch.' A term used for a medicine employed under the form of fumigation, pessary, or troch.

TÆNIA, *Tænia*, *Fas'cia*, (*taenia*, 'a band, a fillet, a riband,') See Bandage. Also a genus of intestinal worms, — *Tape worms* — characterized by an elongated, flattened, and articulated body, with suckorial oscula on the head. Of this genus, two only inhabit the human body — the *Tænia lata*, (see *Bothriocephalus latus*), and *Tænia solium*.

TÆNIA ACEPHALA, *Bothriocephalus latus* — *t. à Anneaux courts*, *Bothriocephalus latus* — *t. à Anneaux longs*, *Tænia solium* — *t.* Articulus demittens, *Tænia solium* — *t.* Cucurbitina, *Tænia solium* — *t.* Dentata, *Bothriocephalus latus* — *t.* Grisea, *Bothriocephalus latus* — *t.* Hippocampi, *Corpora fibriata* — *t.* Humana armata, *Tænia solium* — *t.* Humana inermis, *Bothriocephalus latus* — *t.* Hydatigena, *Hydatid* — *t.* Lata, *Bothriocephalus latus* — *t.* à longues articulations, *Tænia solium* — *t.* Membranacea, *Bothriocephalus latus* — *t.* Nervosa Halleri, Ganglion of Gasser — *t.* Non armé, *Bothriocephalus latus* — *t.* Oculis lateralibus geminis, *Bothriocephalus latus* — *t.* Oculis lateralibus solitariis, *Bothriocephalus latus* — *t.* Oculis marginalibus solitariis, *Tænia solium* — *t.* Oculis superficialibus, *Bothriocephalus latus* — *t.* Prima, *Bothriocephalus latus* — *t.* Sans épine, *Tænia solium* — *t.* Secunda, *Tænia solium*.

TÆNIA SEMICIRCULARIS, *T. striata*, *Gem'inum centrum semicircularē*, *Lim'bus postērior cor'poris striat'is*, *Striæ cor'neæ seu semilunares seu terminales*, *Fra'nulum no'vum*, *Fas'cia semicircularis*, (F.) *Bandelette des éminences pyramiformes* (Ch.), *Bandelette*, *B. demicirculaire*, *Lame cornée*.

A medullary, grayish, semitransparent layer, situate in the groove that separates the thalamus opticus from the corpus striatum in the lateral ventricle.

TÆNIA SOLIUM, (from *solus*, 'alone,' because, at one time, supposed to be always alone.) *T. osculis marginalibus solitariis seu cucurbitina seu humana armata seu articulos demittens seu cucurbita*, *Hal'ysis solium*, *Ver'mis cucurbitinus*, *Lumbricus latus*, *Solium*, (F.) *Tænia sans épine ou à longues articulations ou à anneaux longs*, *Ver solitaire ou cucurbitin*, *Long Tapeworm*, has an almost hemispherical, discrete head; an obtuse beak; the neck full anteriorly; all the joints slightly obtuse, the anterior very short; the next almost square, and the rest oblong; the marginal foramina vaguely alternated. The *tænia solium* is commonly a few feet long: sometimes its length is enormous. It is said to have been met with 600 feet long. It principally inhabits the small intestines, and is not common. Brayera anthelmintica ('koussou'), pumpkin seeds, male fern, pomegranate root, are amongst the best remedies.

TÆNIA STRIATA, *T. semicircularis* — *t.* Tarini, *Lamina cornea* — *t.* Tenella, *Bothriocephalus latus* — *t.* Vulgaris, *Bothriocephalus latus*.

TÆNIACIDE, (*tænia*, and *cædere*, 'to kill,') *Tæniacide*.

TÆNIE COLI, (pl. of *tænia*, 'a band,') see Colon — *t.* Longitudinales, see Colon — *t.* Valsalvæ coli, see Colon.

TÆNIAFUGE, *Tæniifuge*, *Tæniaciide*, *Tæniicide*, *Tæniatugum*, (F.) *Téniifuge*, (*tænia*, and *fugo*, 'I drive off,') A remedy calculated to expel or destroy *tænia*.

TÆNICIDE, (*tænia*, and *cædere*, 'to kill,') A destroyer of *tænia* or tapeworm.

TÆNIFUGE, *Tæniacide*.

TÆNIIN, see *Hægenia Abyssinica*.

TÆNI'OLA, *Bandelette*; (dim. of *Tænia*.) Used by Gerber for a simple, bandlike, or flat fibre, *Filum tæniaform'e*, seen under the microscope.

TÆFFETAS AGGLUTINATIF, *Sparadrapum adhæsivum* — *t. d'Angleterre*, see *Sparadrapum adhæsivum* — *t.* *Épispatique*, *Sparadrapum vesicatorium* — *t.* *Gommé*, see *Sparadrapum adhæsivum* — *t.* *Vésicant*, *Sparadrapum vesicatorium*.

TAFIA, Rum.

TAGLIAOOZZI, *Opération de*, (after G. Tagliacozzi, of Bologna, died 1599,) see Rhinoplasty.

TAIE, (from *tegere*, 'to cover,') see Caligo.

TAILLADE, (from *tailler*, 'to cut,') Incision, see *Entaille*.

TAILLE, (from *tailler*, 'to cut,') Lithotomy, Stature — *t.* *Hypogastrique*, Hypogastric operation of lithotomy, see Lithotomy — *t.* *Postérieure*, see Lithotomy — *t.* *Recto-vésicale*, see Lithotomy — *t.* *par le Rectum*, see Lithotomy — *t.* *Sus-pubiéne*, see Lithotomy.

TAKAMAHAKA, *Fagara octandra*.

TAKE, Attack.

TAKEN DOWN, see Attack.

TAKING, see Contagious.

TAL. A word formerly employed for alkali. It also meant the dung of the peacock.

TALC EARTH, *Magnesia*.

TALCUM NEPHRITICUM, *Nephreticus lapis*.

TALENTUM, (ταλάνον.) An ancient weight equal to 54 pounds, 2 oz., 5 dr., and 24 grains, *Poids de Marc*. See Weights and Measures.

TALEOLUS, (dim. of *talus*,) *Malleolus*.

TALIAOTIAN OPERATION, (after *Taliacotius* or *Tagliacozzi*, see *Tagliacozzi*,) *Rhinoplasty*.

TALINUM UMBELLATUM, see *Mistura*.

TALIPES, (*talipede*, 'to be weak in the foot,') *Kyllosis* (Clubfoot) — *t.* *Calcaneus*, see *Kyllosis* — *t.* *Equinus*, see *Kyllosis* — *t.* *Varus*, see *Kyllosis*.

TALLAHATTA SPRINGS, see Alabama, mineral waters of.

TALLOW, Bayberry, see Wax, myrtle.

TALON, (from *talus*,) see *Calcaneum*.

TAL'PA, 'a mole,' *Topina'ria*, *Upina'ria*, *Talpa'ria*, *Testu'do*, (F.) *Taupe*. A tumour situate on the head, which has been supposed to burrow like a mole.

TALPARIA, *Talpa*.

TALUS, *Astragalus*, *Malleolus*, see *Calcaneum*. Pessary.

TAMARA CONGA, *Averrhoa carambola*.

TAMARÆA ZECLA, *Tamarindus*.

TAMARIN, *Tamarindus*.

TAMARIN'DI PUL'PA, *Pulp of Tam'arinds*. (*Tamarindi*, q. s. Digest them with a small quantity of water, until they become of a uniform consistence; then separate the seeds and filaments by pressing through a hair-sieve. Ph. U. S.)

TAMARINDS, PULP OF, *Tamarindi pulpa*.

TAMARINDUS, (from [Arab.] *tamar hendi*,

'Indian date.) *T. Indica*, *Thamari'ndus*, *Oxyphæ'nica*, *Sil'iqua Arab'ica*, *Balampul'i*, *Tamar'ia zec'la*, *Oxyphani'cia*, *Acu'cia Ind'ica*, *Tam'arind tree*, (F.) *Tamarin*. *Ord.* Leguminosæ. *Sex. Syst.* Triandria Digynia. The pulp of the tamarind with the seeds connected by numerous tough strings, *Tamarindus* (Ph. U. S.), is exported from the West Indies, preserved in syrup. It is refrigerant and laxative. *Tamarind whey* is made by boiling 3ij of the pulp with Biss of milk. When strained, it is used as a diluent in fever.

TAMARISCUS, (from *Tamaris*, now *Tambro*, on the borders of the Pyrenees.) *Tamarix Gallica*.

TAMARIX, gen. *Tamari'cis*, *T. Gal'lica* seu *Pentan'dra*, *Tamaris'cus*, *The Tamarisk tree*. *Ord.* Tamariscineæ. *Sex. Syst.* Pentandria Pentagynia. The bark, wood, and leaves were formerly employed medicinally;—the former for its aperient and corroborant virtues in obstructions of the liver; the latter in icterus, hæmoptysis, and some cutaneous affections.

TAMBOUR, *CAVITÉ DU*, 'drum, cavity of the,' Tympanum.

TAMNUS COMMUNIS, *Tamus communis*.

TAMPA BAY, see Saint Augustine.

TAMPAN. An insect belonging to the group of Acaridæ or Ticks, well known in South Africa, and described by Dr. Livingstone as dreaded by the inhabitants for its bite and the consequences.

TAMPON, ([F.] 'a bung, a plug,') see Plugging.

TAMPONNEMENT, Plugging.

TAMUS seu **TAMNUS COMMUNIS**, *Am'pe'los melæ'na* (Dioscorides), *Chiro'nia*, *Gynæcan'thè* seu *Ap'ro'nia* (Pliny), *Bryo'nia nig'ra*, *Com'mon Black Bry'ony*. The root is diuretic, and has been given as an antilithic. In the herbshops of Europe it is sold as a topical application for removing bruise marks, and in France is called *Herbe aux femmes battues*, 'herb for beaten or bruised women.'

TANACETUM, *T. vulga'rè*, *Tana'sia*, *Athana'sia*, *Parthen'ium mas*, *Hindheel*, *Tan'sy*, (F.) *Tana'ie vulgaire*, *Herbe aux vers*, *Coq des Jurdins*. *Ord.* Compositæ. *Sex. Syst.* Syngenesia Polygamia. The leaves and flowers of tansy, and the whole herb—*Tanacetum* (Ph. U. S.)—have a peculiar, strong odour, and warm bitter taste. They are tonic, 'deobstruent' and anthelmintic. Dose, ʒss to ʒj. The infusion is usually drunk as tea.

TANACETUM BALSAMI'TA, *T. horten'sè*, *Balsami'ta mas* seu *vulga'ris* seu *suave'olens* seu *ma'jor*, *Cost'us horto'rum*, *Men'tha Roma'na* seu *Saracen'ica*, *Cost'mary*, *A'lecost*, *Cockmint*, (F.) *Menthe coq*, *Herbe au coq*, *Grand Baume*. A fragrant-smelling herb, somewhat resembling mint, and formerly much esteemed as a corroborant, carminative, and emmenagogue.

TANACETUM HORTENSE, *T. balsamita*.

TANACETUM MULTIFLO'RUM. A South African plant, which is used as a tonic, antispasmodic, and anthelmintic. It is given in flatulency, gout, amenorrhæa, and dropsy; but is especially prescribed in cases of ascarides lumbricoides and other entozoa. It is administered in the form of powder or infusion; the latter acting also as a diaphoretic and diuretic. It is used likewise as a resolvent and anodyne fomentation.

TANASIE VULGAIRE, *Tanacetum*.

TANASIA, *Tanacetum*.

TAN'GÈ, (*ταγγη*, 'rancidity,') A putrescent tumour.

TANGHINIA, (Madagascar tanghin,) *Cerbera tanghin*.

TANGLES, BIG, *Pila marina*—t. Blue, *Gy-lussacia dumosa*.

TANNINUM, *Tanninum*.

TANNE, see Acne.

TANNIC ACID, *Tannin*.

TANNICUM PURUM, *Tannin*.

TANNIN, (F.) *tanner*, 'to tan,') *Tan'nicum pu'rum*, *Tannin'um*, *Tani'num*, *Principi'um Ad-strin'gens* seu *Scytodep'sicum* seu *Byrsodep'sicum* seu *Coria'ceum*, *Scytodep'sium*, *Mat'e'ria Scyto-deph'ica*, *Ac'idum Tan'nicum* seu *Scytodeph'icum* seu *Quercitan'nicum*, *Tan'nic acid*, (F.) *Acide tan-nique*. A principle obtained from oak bark and other astringent vegetables, so called from its forming the principal agent in the process of *tanning*. Pure tannin, obtained from *galls*, by the action of *ether* (Ph. U. S.), has been employed as an excellent and powerful astringent. Dose, two grains, repeated several times a day.

TANSY, *Tanacetum*—t. Wild, *Potentilla an-serina*.

TAPER, HIGH, *Verbascum nigrum*.

TAPES, (*ταπης*, 'a carpet,') *Tapetum*.

TAPE'TUM, *Ta'pes*, *Tape'te*, (F.) *Tapis*, 'tapestry, a carpet,' see *Corpus Callosum*.

TAPE'TUM CHOROÏDÆ. A shining spot on the outside of the optic nerve in the eyes of certain animals, which is owing to the absence of the pigmentum nigrum, occasioning the reflection of a portion of the rays from the membrana Ruy-schiana. Its use appears to be, by reflecting the rays, to cause a double impression on the retina, and thus add to the intensity of vision.

TAPE'TUM ALVE'OLI, *Membra'na exter'na den-tium*. The outer membrane of the teeth. The lining membrane or periosteum of the alveoli.

TAPEWORM, *Tænia*—t. Broad, *Bothriocephalus latus*—t. Long, *Tænia solium*.

TAPIOCA, see *Jatropha manihot*—t. Pearl, see *Jatropha manihot*.

TAPIS, (from *tapes*, 'a carpet,') *Tapetum*.

TAPPING, (from [Sax.] *tæppan*, 'to broach,' as a cask,) *Paracetasis*.

TAPSARIA, *Decoctum hordei*.

TAPUS BARBATUS, *Verbascum nigrum*.

TAR, ([Sax.] *tape*,) see *Pinus sylvestris*—t.

Barbados, Petroleum,

TAR BEER, see *Pinus sylvestris*.

TAR, MINERAL, *Pissasphaltum*.

TAR WATER, see *Pinus sylvestris*—t. Wile, see *Pinus sylvestris*.

TARA. An epidemicocontagious disease seen by Gmelin in Siberia, in the town of Tara, which appears to have resembled somewhat the Button Scurvy of Ireland.

TAR'ACHÈ, (*ταραχη*,) *Tarax'is*, *Tarag'ma*, *Tarag'mus*, 'disorder.' A disorder of sight which arises from a blow, compression of the eye, or the action of dust, smoke, or other external cause. By some it is understood to signify mild or incipient ophthalmia.

Ταραχη της κοιλίας, 'disturbance of the belly,' *Diarrhœa*.

TARAGMA, gen. *Tarug'matis*, (*ταραγμα*,) *Tarache*.

TARAGMUS, (*ταραγμος*,) *Tarache*.

TARAGON, *Artemisia dracuncul*.

TARAHU, *Peltandra Virginica*.

TARANTISM'US, *Tarentis'mus*, *Tarantalis'mus*, *Dinoma'nia*, *Chore'a Sane'ti Valenti'ni*, *Taren'tulism*, *Tar'antism*, (from *Tarentum* or *Taranto*, where it was observed.) A feigned or imaginary disease in Apulia, characterized by excessive avidity for dancing at the sound of instruments, and which was ascribed by the vulgar to the bite of the tarantula. A sort of tarantism, considered peculiar to Africa, and believed to be spontaneous, has been called *Tarentis'mus Tangita'mus*, *Janon-tarentis'mus*.

TARANTULA, (same etymon,) *Taren'tula*, (F.) *Tarentule*. A species of spider—the *Ara'nea Taren'tula* of Linnæus, *Lyc'o'sa Tarentula* of La-

treille—whose bite has been looked upon as dangerous. Music has been thought to cure it.

TARASCON, MINERAL WATERS OF. Chalybeate springs in the department of Arrière, France. They contain iron and carbonic acid.

TARAXACUM, (from *ταρασσειν*, 'to disturb or disorder,') *Leontodon taraxacum*—*t.* *Dens leonis*, *Leontodon taraxacum*.

TARAXIS, (*ταραξις*.) *Tarache*, see *Ophthalmia*.

TARCHON SYLVESTRIS, *Achillea ptarmica*.

TARANTISMUS, *Tarantismus*—*t.* *Tangitanus*, *Tarantismus*.

TARENTULA, *Tarantula*.

TARENTULISM, *Tarantismus*.

TARGAR, *Juniperus communis*.

TARI. Palm or cocoa wine, from which sugar was obtained, called *Jugre*. It was once used as a tonic.

TARINI PONS, (after the anatomist *Tarin*,) (*F.*) *Pont de Tarin*. A cineritious substance, between the corpora albicantia, which unites the crura, and assists in forming the bottom of the third ventricle. The fossa, at the bottom of which it is placed, is called *Lo'cus perforatus posticus*, *Posterior perforated space*, *Substantia seu Lamina perforata posterior*.

TARO, *Arum esculentum*.

TAR'SAL, *Tarsalis*, *Tar'seus*, (from *tarsus*,) (*F.*) *Tarsien*. That which relates to the tarsus.

TARSAL ARTICULATIONS result from the union of the bones of the tarsus with each other.

TARSAL CARTILAGES, see *Tarsus*.

TAR'SEA LA'TA (LIGAMENTA). Three ligaments, described by Winslow as stretching from the edge of the orbit to that of the tarsi, with which he thought they formed a complete layer of the eyelids. Zinn could not trace the ligaments all the way to the tarsi; whilst Haller doubted altogether their ligamentous texture, and Zeis regarded them as merely areolar substance.

TARSEUS, *Tarsal*.

TARSIEN, *Tarsal*.

TARSOMETATAR'SAL, *Tarsometatar'seus*. That which relates to tarsus and metatarsus.

TARSOMETARSAL ARTICULATIONS are those between the bones of the second row of the tarsus and the metatarsal bones. The term *Tarsometatarsal* is likewise applied to ligaments connected with these joints, some of which are *dorsal*, and others *plantar*.

TARSO-MÉTATARSI-PHALANGIEN DU POUCE, *Adductor pollicis pedis*—*t.* *Phalangienn du ponce*, *Flexor brevis pollicis pedis*—*t.* *Sous-phalangienn du petit orteil*, *Flexor minimi digiti pedis*—*t.* *Sous-phalangien du premier orteil*, *Flexor brevis pollicis pedis*.

TARSOPHY'MA, gen. *Tarsophy'matis*, (*tarsus*, and *φύμα*, 'a tumour,') A swelling of the tarsus.

TARSORRHAPHY, *Tarsorrhaph'ia*, (*tarsus*, and *ῥάφην*, 'suture,') An operation for diminishing the size of the opening between the eyelids when it is enlarged by surrounding cicatrices.

TARSOTOMY, *Tarsotom'ia*, (*tarsus*, and *τομή*, 'incision,') The section or removal of the tarsal cartilages.

TAR'SUS, (*ταρσος*,) *Plan'ta pri'ma seu Rasse'ta seu Rasse'ta pe'dis*, *Pe'dium*, *Claviculæ pe'dis no'dus*. 1. The posterior part of the foot. The tarsus consists of 7 bones arranged transversely in two rows, between which amputation is sometimes performed. The first or *Tib'ial row*, (*F.*) *Première rangée*, *R. jambière*, consists of the astragalus and os calcis; the second or *metatarsal*, (*F.*) *Seconde rangée*, *R. métatarsienne*, of the scaphoides, cuboides, and three ossa cuneiformia. 2. The *Tar'si*, *Tar'sal cartilages*, *Tar'sal fibrocartilages*, *Or'choi*, *Crepid'ines seu Coro'næ seu Che'læ seu Un'guis Palpebra'rum*, *Tar'si*, *Cil'ia*, are two small cartilaginous layers, seated in the substance of the free edge of each eyelid. The upper is

larger and broader than the lower. Both commence at the bifurcated extremity of the tendon of the orbicularis palpebrarum, and pass to the opposite commissure of the eyelids. They are broader in the middle than at their extremities.

TARTAR, (from *tart*, 'acid,' [?]) *Antimonium tartarizatum* (improperly), *Potassæ supertartas impurus*—*t.* *Calcareous*, *Potassæ subcarbonas*—*t.* *Crystals of Potassæ supertartas*.

TARTAR EMETIC, *Antimonium tartarizatum*.

TARTAR, SALT OF, see *Potash*, and *Potassæ subcarbonas*—*t.* *Soluble*, *Potassæ tartas*, see *Potash*.

TARTAR OF THE TEETH, *Odontolithus*.

TARTAREI MOR'BI. A term employed by Paracelsus to designate certain diseases which he attributed to tartar.

TARTARI CRYSTALLI, *Potassæ supertartas*.

TARTAR'IC ACID, *Ac'idum tartar'icum seu tartar'um seu tar'tari essential'e*, *A'cor tartar'icus*, *Sal tar'tari essential'e*, *Tartareus ac'id*, (*F.*) *Acide tartrique*. (*Potassæ bitar*, *Biess*, *Aq. destillat. ferv.*, cong. iij, *Cretæ ppt. lbj*, *Acidi sulph. lbj*. Boil the bitartrate with two gallons of the water, and gradually add the prepared chalk until no more bubbles appear; let the tartrate of lime subside; then pour off the fluid, and wash the tartrate of lime with distilled water until it is tasteless. Pour upon it the sulphuric acid, diluted with a gallon of boiling distilled water; and set the mixture aside for 24 hours, occasionally stirring it. Filter, and evaporate it to crystallize. *Ph. L.*) It is refrigerant and antiseptic, but is chiefly used for making effervescent draughts with soda.

TARTARUM, *Potassæ supertartas impurus*—*t.* *Solubile*, *Potassæ tartas*—*t.* *Vitriolatum*, *Potassæ sulphas*.

TARTARUS AMMONIÆ, *Ammonia tartas*—*t.* *Chalybeatus*, *Ferrum tartarizatum*—*t.* *Crudus*, *Potassæ supertartas impurus*—*t.* *Dentium*, *Odontolithus*—*t.* *Emeticus*, *Antimonium tartarizatum*—*t.* *Kalioferrius*, *Ferrum tartarizatum*—*t.* *Kalinus*, *Potassæ tartas*—*t.* *Lixivia*, *Potassæ tartas*—*t.* *Martialis*, *Ferrum tartarizatum*—*t.* *Potassæ ferruginosus*, *Ferrum tartarizatum*—*t.* *Stibiatus*, *Antimonium tartarizatum*—*t.* *Subpotassicus*, *Potassæ supertartas*—*t.* *Superpotassicus*, *Potassæ supertartas*—*t.* *Tartarizatus*, *Potassæ tartas*.

TARTON-RAIRE, *Passerina tartonraira*.

TARTRATE D'AMMONIAQUE, *Ammonia tartas*.

TARTRE BRUT, 'crude tartar,' *Potassæ supertartas impurus*—*t.* *Cru*, *Potassæ supertartas impurus*—*t.* *des Dents*, *Odontolithus*—*t.* *Émetique*, *Antimonium tartarizatum*—*t.* *Stibié*, *Antimonium tartarizatum*.

TASIS, (*τασις*,) *Extension*, *Tension*.

TASTE, (from [*F.*] *tāter*, [Old *F.*] *taster*, 'to taste, feel,' [*G.*] *tasten*, 'to feel,') *Gust'us*, *Gent'is*, *Gent'ia*, *Gent'ios*, (*F.*) *Goût*. One of the five external senses which instructs us of the savours of bodies, and of which the tongue is the principal organ.

Gustat'ion, *Gusta'tio*, is the exercise of this faculty. In disease the taste frequently becomes variously vitiated. The nerves of taste are probably the glossopharyngeal, and, perhaps, also, the lingual branch of the fifth pair.

TASTE, SAVOUR.

TATOUAGE, *Tattooing*.

TATTOO'ING, (*F.*) *Tatouage*. A practice with the South Sea Islanders of puncturing the skin, and forcing colouring matters into the punctures. Sailors in all countries are in the custom of practising it by rubbing or forcing gunpowder into the punctures. The figures thus formed remain through life.

TATZÉ. The fruit of *Myrsina Africa'na*, a native of Abyssinia, the Cape of Good Hope, the Azores, and Algeria, which is used as a tænia-fuge.

TAUPE, Talpa.

TAUROCOL'LA, (*ταυρος*, 'a bull,' and *κόλλα*, 'glue.')

 Glue made of the ears and genitals of the bull. It was formerly much esteemed.

TAURUS, (*ταυρος*,) Penis, Perineum. Of old, the lower part of the male organ with the rhaps extending to the anus.

TAX'IS, (*τάξις*, 'arrangement;') from *τάσσω*, 'I order,' 'I arrange;') *Repositio*, *Reductio*, *Syn-taxis*. A methodical pressure exerted by the hand on a hernial tumour, for the purpose of reducing it. This operation is generally easy in herniæ that are free from adhesions, of small size, and with a large opening into the sac; but it is very difficult and frequently impracticable in those that are adherent or strangulated. To perform the taxis, the patient should be put in the horizontal posture, and in one in which the parietes of the abdomen are relaxed. The manual procedure varies in each species of hernia. In all cases, too strong pressure of the parts must be avoided; as inflammation and gangrene might be induced. The displaced organs must be pushed into the abdomen, according to the direction of the axis of the hernial sac. The reduction is sometimes favoured by bloodletting *ad deliquium*; the warm bath; warm fomentations; tobacco; emollient glysters; and, at times, by cold applications, as ice, &c. In a case of strangulated hernia, should the taxis fail, an operation becomes necessary.

TAXODIUM DIS'TICHUM, (*taxis*, and *εἶδος*, 'resemblance,') *Cupressus disticha*, *Schubertia disticha*, *Cypress*, *American* or *Bald Cypress*; indigenous. *Nat. Ord.* Coniferæ. A balsam, obtained from it, is applied to cuts and wounds. The resin from the cones is diuretic and carminative.

TAXONOMY, (*τάξις*, 'order,' and *νομος*, 'law,') Biotaxy.

TAX'US BACCA'TA, (*ταξος*,) The *Yew tree*, *Smilax*, (F.) *If. Family*, Coniferæ. *Sex. Syst.* Dicoeca Monadelphica. The ancients believed the yew to be poisonous; but the fruit may be eaten without inconvenience. The leaves have been often recommended in epilepsy. The resin, obtained from the bark, has been called *albir*.

TAXUS CANADENS'IS, *American Yew*, *Ground Hemlock*; indigenous. The leaves are said to be sedative. [?]

TAXWAX, see Nucha.

TA YE, (from [L.] *tego*, 'to cover,') Caligo.

TAYLOR'S REMEDY FOR DEAFNESS, see Allium.

TEA, Thea.—t. Apple, see Apple tea.—t. Bahama, *Lantana camara*—t. Beef, see Beef tea.—t. Berried, *Gaultheria*.

TEA BERRY, *Gaultheria*.

TEA, **BLANKENHEIM**, see *Galeopsis grandiflora*—t. Brazil, *Lantana pseudothea*—t. Bushman's, *Methystophyllum glaucum*—t. Caffre, *Helichrysum nudifolium*—t. Chicken, see Chicken tea.—t. Flaxseed, *Infusum Lini compositum*—t. Ginger, *Infusum Zingiberis*—t. Hottentot's, *Helichrysum serpyllifolium*.

TEA IMPROVER, CHINESE, see Falsifications, table of.

TEA, **LABRADOR**, *Ledum latifolium*—t. Lemon-peel, see Citrus medica.—t. Lie, see Falsifications, table of.—t. Lye, Physick's, see Fuligokali.—t. Marsh, *Ledum palustre*—t. Mexico, *Chenopodium ambrosioides*—t. Mint, *Infusum menthæ*—t. Mountain, *Gaultheria*—t. Mutton, see Mutton tea.—t. New Jersey, *Celastrus*—t. Oswego, *Monarda coccinea*—t. Redberry, *Gaultheria*—t. Sage, *Infusum salvia*—t. Santa Fé, *Alstonia theaeformis*—t. Sass,

see Saloop.—t. Sassafras, Saloop.—t. Slippery elm, *Infusum Lini compositum*—t. Soot, see Fuligokali.—t. South sea, *Ilex vomitoria*—t. Spanish, *Chenopodium ambrosioides*—t. Veal, see Veal tea.

TEAK, *Paronychia*.

TEAR, (Sax. *tear*,) *Lacryma*, *Lachryma*, *Lacrima*, *Lacrimum*, *Dacry*, *Dacryma*, *Dacryon*, *Hu'mor lacrymalis*, *Eye'drop*, *Weepingtear*, (F.) *Larme*. An excrementitious humour, secreted by the lachrymal gland, and poured between the globe of the eye, and the eyelids, to facilitate the motions of those parts. The tears are saline, inodorous, and colourless. They turn the syrup of violets green; and contain chloride of sodium, with excess of base; phosphates of lime and soda, and albumen, and a special organic matter called *Lacrymine* and *Threnine*.

TEAR PUNCTA, *Lachrymal puncta*.

TEARS, **DEERS'**, Bezoar of the deer.

TEASEL, **CULTIVATED**, *Dipsacus sylvestris*.

TEASICK, (*phthisic*,) *Phthisis pulmonalis*.

TEAT, (*τιθήνη*, [Sax.] *tit*,) Nipple.

TEAZEL, ([Sax.] *tæzel*,) *Eupatorium perfoliatum*.

TECEDON, (*τηκεδών*,) *Phthisis*.

TECNOTONIA, (*τεκνοτονία*, from *τεκνω*, 'a child,' and *κτενω*, 'to kill,') Infanticide.

TECNOLOGY, *Tecology*, *Tecnolog'ia*, *Tecnolog'ia*, (*τεκνω*, and *τεκος*, 'a child,' and *λογος*, 'a discourse,') A treatise on children.

TECOLITHOS, (*τηκω*, 'to dissolve,' and *λίθος*, 'a stone,') *Judæus* (lapis.)

TECOLOGY, (*τεκος*, 'a child,' and *λογος*, 'a discourse,') *Tecology*.

TECOM'PA RADICANS, *Bigno'nia radicans*, *Trump'pet flower*, *Trump'pet creep'er*, *Virgin'ia creep'er*. Indigenous; the orange and scarlet flowers appearing in July. *Order*, *Bignoniaceæ*. The leaves are acrid.

TECTARIA CALAHUALA, see *Calagualæ radix*—t. Ferruginea, see *Galagualæ radix*.

TEDA, *Tæda*.

TEETH, see Tooth.—t. Artificial, see Artificial.—t. Barred, *Barrées* (*Dents*.)

TEETH, **BLOCK**. In dentistry, two or more artificial teeth carved from a piece of ivory, or formed from porcelain paste, afterwards baked and finished.

TEETH, **EYE**, Canine teeth—t. Incorruptible, see Artificial.—t. Jaw, Molar teeth—t. Milk, see Dentition.—t. Mineral, see Artificial—t. Neck of the, Collum dentium—t. Plate, Porrigo decalvans—t. Porcelain, see Artificial—t. Shedding of the, see Dentition—t. Vitreous substance of the, Enamel of the teeth—t. Vitrescent, see Artificial—t. Wisdom, see Dentition—t. Wit, see Dentition.

TEETHING, Dentition.

TEGENERA MEDICINALIS, *Cantharis*.

TEGIMEN, (*tegere*, *tectum*, 'to cover,') Integument.

TEGMEN, Integument.

TEGULA HIBERNICA, 'a tile,' *Hibernicus lapis*.

TEGUMEN, gen. *Tegu'minis*, (*tegere*, *tectum*, 'to cover,') Integument.

TEGUMENT, Integument.

TEGUMENT'ARY, *Tegumenta'rius*, (F.) *Tégumentaire*. Relating or appertaining to the integument.

TEGUMENTUM, Integument—t. *Αἰς*, see Tympanum—t. Ventriculorum cerebri, Centre, oval, of the brain.

TEIGNE, (*tinea*,) *Pityriasis*, *Porrigo*, *Porrigo lupinosa*—t. *Annulaire*, *Porrigo scutulata*—t. *en Anneau*, *Porrigo scutulata*—t. *Favense*, *Porrigo favosa*—t. *Furfuracée*, *Porrigo furfurans*—t. *Nemulaire*, *Porrigo scutulata*—t. *Felade*, *Porrigo decalvans*—t. *Tondante*, *Porrigo decalvans*—t. *Tonsurante*, *Porrigo decalvans*.

TEINESMUS, (*τενεσμος*,) *Tenesmus*.

TEINODYN'IA, (τεῖνυν, 'to extend, and ὀδυν, 'pain,') Pain of the sinews or tendons.

TEINOPHLOGO'SIS, (τεῖνυν, 'to extend,' and φλογωσις, 'inflammation,') Inflammation of the sinews or tendons.

TEINTURE, Tincture—t. Ἐθέρη, see Æthereal—t. Rubéfiante, Tinctura cantharidis.

TEKEDON, (τηκεδών,) Tabes.

TELA, 'a web,' Texture, Tissue—t. Accidentalis Medullaris, Encephaloid—t. Cavernosa, Cavernous texture—t. Cellulosa, Cellular tissue—t. Choroidea, Choroidea tela—t. Cornea, Corneous tissue—t. Emplastica, Sparadrapum—t. Erectilis, Erectile tissue—t. Fibrosa, see Fibrous—t. Galteri, Sparadrapum Galteri—t. Hippocratis cribrosa, Cellular tissue—t. Hymenochondrodes, Tissue, cartilaginous—t. Hymenochondroides, Tissue, cartilaginous—t. Ichthyocollæ glutinans, see Sparadrapum adhaesivum—t. Inodularis, Tissue, inodular—t. Mucosa, Cellular tissue—t. Organica, Histos—t. Vesicatoria, Sparadrapum vesicatorium.

TELAMON, (τελαμών,) Habena.

TELAMO'NES, (τελαμώνες, 'bandages,') Charpie, used for wounds; or linen employed for bandages.

TELANGECTASIS, Telangiectasia.

TELANGIECTA'SIA, *Telangiect'asis*, *Telangiect'osis*, *Angeiotellect'asia*, *Hæmatotelangi'osis*, *Hæmatotelangi'osis*, *Angiectangi'osis*, *Anastomosis aneurysmat'ica*, (τηλε, 'far,' αγγειον, 'vessel,' and εκτασις, 'dilatation,') Dilatation of far or capillary vessels. Nævus maternus. Aneurism by anastomosis. According to some, fungus hæmatodes. See Hæmatodes fungus.

TELANGIECTASIA OCULI, Cirsophthalmus.

TELEORGANIC, (τελος, 'end or object,') see Organic.

TELEPH'UUM, (after Τηλεφος, *Telephus*, son of Hercules,) *Ulcus telephium*, *Achille'um*. The ancients gave this name to any ulcer of a very bad kind, and difficult to cure, because the wound, which Achilles inflicted on Telephus, degenerated into a similar ulcer.

TELEPHIUM, Sedum telephium.

TELLURIUM, (F.) *Tellure*, (from *tellus*, gen. *telluris*, 'the earth,') A metal first met with in the gold mines of Transylvania. It is of a bluish white colour, very volatilizable; specific gravity 6.15; and oxidizable in the air and by heat.

The salts of tellurium occasion the exhalation of so disagreeable an alliaceous odour from the body, that they have not been much experimented with as therapeutical agents.

TELORASIS, (τηλε, 'far,' and ὁραω, 'to view,') Presbytia.

TELUM, 'a dart,' Penis.

TEMPE, Temple.

TEMPERAMENT, *Temperamentum*, *Temperat'ura*, *Temperat'io*, *Tempe'ries*, (from *temperare*, 'to moderate,' originally from *tempos*, gen. *temporis*, 'time,' the great *temperer*;) *Complex'io*, *Cra'sia*, *Eucra'sia*. A name given to the remarkable differences that exist between individuals, in consequence of the variety of relations and proportions between the constituent parts of the body. Thus, of old, it was supposed that, according to the predominance of any one of the four humours then generally admitted,—blood, lymph, bile, and atrabilis, or black bile,—the manifestations of the functions were *tempered* or modified so as to give rise to a *sanguine*, *lymphatic*, *choleric*, or *atrabilious* or *melancholic* predominance or temperament. The idea of the existence of certain of these humours has been banished; yet the influence of some, as of the blood, lymph, and bile, is still admitted by many. Too much influence has, doubtless, been assigned to temperament in the sense in which it was formerly, and still is, generally understood, although there can be no

question that difference of organization occasions a difference in the activity of organs.

TEMPERAMENT, BILIOUS, Choleric temperament. TEMPERANCE, *Temperant'ia*, *Sophros'ynê*, *Nephal'otes*, *Sobri'etas*, *Sobri'ety*. Moderation, opposed to gluttony and drunkenness. The advantage of temperance, in a hygienic point of view, requires no comment.

TEMPÉRANTS, (from *temperare*, 'to mitigate,') Sedatives.

TEMPERATIO, Temperament.

TEMPERATURA, Temperament.

TEMPERIES, Temperament.

TEMPESTAS, (from *tempus*, 'time,') Season.

TEMPLE, *Tem'pus*, *Re'gio Tempora'lis*, *Crot'aphus*, (F.) *Tempe*. A depression, on each side of the head, between the forehead and eye anteriorly, and the ear posteriorly. The *temples*, distinguished into *right* and *left*, correspond to the temporal fossa on each side. (The word is said to be from the Latin *tempus*, 'time,' because in this part the hair first begins to turn white, and to indicate age.)

TEMPORAL, *Tempora'lis*, (from *tempus*.) That which relates to the temples.

TEMPORAL APONEURO'SIS is a very strong aponeurosis, of a bluish, shining colour, which is attached to the whole of the curved line of the temporal bone; to the posterior edge of the malar bone; to the superior margin of the zygomatic arch; and forms an envelope for the temporal muscle.

TEMPORAL ARTERIES, *Crot'aphite Ar'teries*, are several:—I. The *Temporal Artery*, properly so called, arises from the outer side of the superior part of the external carotid. It ascends between the ramus of the jaw, the meatus auditorius externus, and the parotid gland, which covers it as far as the zygomatic arch. Above this it becomes subcutaneous; and, when it has attained the middle of the temporal region, it divides into two branches; the one anterior—*A. tempora'lis fronta'lis* seu *inter'na* seu *ante'rior*; the other posterior—the *A. tempora'lis occipita'lis* seu *exter'na* seu *superior*. The temporal artery, before its division, gives off, 1. Branches to the parotid; to the joint of the lower jaw; to the meatus auditorius, and to the masseter. 2. The *trans'verse artery of the face*. 3. The *mid'dle tempora'l*, which arises beneath the cheekbones, and traverses the temporal aponeurosis immediately beneath that bone. II. The *Deepseated Temp'orals*—*A. tempora'les profundi*. These are two in number. They arise from the internal maxillary, and have been distinguished, from their position, into *ante'rior* and *posterior*. They send their principal divisions to the temporal muscle.

TEMPORAL BONE, *Os tempora'le* seu *tempora'lè* seu *arcua'lè* seu *arma'lè* seu *parieta'lè inf'e'rius* seu *crot'aphum* seu *mement'o mo'ri* seu *menda'cium* seu *crotaphit'icum* seu *lapida'sum*, is seated at the lateral and inferior part of the cranium, of which it forms part; and contains, within it, the special organs of audition. It is usually divided into three portions—1. The *squamo'sa*, *Pars squamo'sa*, *Squa'ma*, *Lepi'sma*, *Os lepidoi'des* seu *squamo'sum* seu *squamifor'me*, (F.) *Portion écailleuse ou squameuse*. 2. The *mast'oid portion*, and 3. The *Pet'rous portion*, *Pet'rous Proc'ess*, *Pyr'amis trigo'na*, *Os lapide'um* seu *lapido'sum* seu *sac'e'um* seu *petro'sum* seu *lithoi'des* seu *du'rum* seu *præcip'ite ru'pi* *assimila'tum* seu *neva'lè*, *Pars petro'sa*, *P. pyramida'lis os'is temp'oris*, (F.) *Portion pierreuse*, *Rocher*, *Os pétreux*, *Apophyse pyramidale*, *A. pétée*.

It has, 1. an *auric'ular* or *exter'nal face*, on which is a surface that makes part of the temporal fossa; the *zygomatic process*; the *glenoid cavity*; the *fissure of Glaser*; the *meatus audito-*

rius externus; the mastoid process; the digastric groove; the canalis caroticus; the jugular fossa; the styloid process; the stylumastoid foramen, &c. 2. A cerebral or internal surface, which corresponds to the cavity of the cranium, and on which is the *Pars petrosa*, the hiatus Fallopii, the meatus auditorius internus, the fissure of the aqueductus vestibuli, a portion of the gutter for the lateral sinus, &c. 3. A circumference, on which are observed the orifice of the bony portion of the Eustachian tube, the aqueductus cochleæ, &c. The temporal bone is articulated with the sphenoid, occipital, parietal, malar, and inferior maxillary bones. It is developed from five points of ossification.

TEMPORAL FOS'SA, *Fos'sa temporalis*, is an excavation, observed at each side of the head. It is filled by the muscle whose name it bears, and is formed by the temporal and sphenoid, below; by the parietal and frontal bones, above; is separated, by a transverse crista, from the zygomatic fossa; and completed, before, by the cheek-bone.

TEMPORAL MUSCLE, *Tempor'al'is*, *Crotaph'ites*, (F.) *Arædi-temporo maxillaire*, *Temporo-maxillaire* (Ch.), is a broad, flattened, triangular muscle, which fills the temporal fossa. Its fibres are attached to the temporal aponeurosis and fossa; and all converge into a strong tendon, which passes under the zygomatic arch, and is inserted into the coronoid process of the lower jaw. This muscle raises the lower jaw, and applies the teeth against each other.

TEMPORAL NERVES. These are distinguished into the superficial temporal, furnished by the inferior maxillary branch of the fifth pair, behind the condyle of the jaw; and the temporales profundæ,—anterior and posterior—furnished by the same nerve at its exit from the cranium. Sömmering has given the name *temporal* to the divisions sent by the facial nerve to the temporal regions.

TEMPORAL VEINS were formerly called *Ætiol Phlebes*, *Aq'uile Venæ*.

TEMPORALIS, Temporal.

TEMPORO, (*tempus*, gen. *tem'poris*, 'temple,') in composition, the temple.

TEMPOROÏURICULAIRE, Attollens aurem.

TEMPOROFA'CIAL, *Tem'porofacia'lis*. Belonging to the temple and face.

TEMPOROFACIAL NERVE, *Nervus temporofacia'lis*. A branch of the facial nerve distributed to the face and temple.

TEMPOROMAXILLAIRE, Temporomaxillary. see Temporal muscle.

TEMPOROMAXILLARY, *Tem'poromaxilla'ris*, (F.) *Temporomaxillaire*. That which belongs to the temporal bone and lower jaw.

TEMPOROMAXILLARY ARTICULA'TION or joint of the lower jaw is seated between the glenoid cavity and transverse process of the temporal bone on the one hand, and the condyle of the lower jaw on the other. Two thin cartilages cover the surfaces of these bones. The one is common to the transverse process, and the portion of the glenoid cavity anterior to the fissure of Glaser; the other belongs to the condyle. This articulation is furnished with an interarticular fibrocartilage, two synovial capsules, and two lateral ligaments:—the one internal, and the other external.

TEMPOROMAXILLARY NERVES of Bichat are the divisions of the facial nerve, distributed to the temporal and maxillary regions.

TEMPUS, gen. *Tem'poris*, 'time,' Temple.—t. Intercalare. Apærexia, Intermision.

TEM'ULENCE, *Ebri'etas*, *Ebrius'itas*, *Methys'mus*, *Meth'ysis*, *Temulen'tia*, *Tem'ulency*, *Inebria'tion*, *Intoxica'tion*, *Drunk'ness*:—from wine, *Ænophyl'ia*, *Ænophyl'is*, *Vinolen'tia*, *Vino-*

lence; (F.) *Ivresse*; (from *tem'eum*, 'wine, any intoxicating drink,') Plater gave this name to delirium; Ettmüller, to an apoplectic condition, depending upon drunkenness; the *Apoplexia Temulen'tia* or dead drunkenness. Commonly *Temulentia* is used synonymously with drunkenness, *Me'thê*; and is often employed in the description of diseases to indicate a state resembling drunkenness. *Mania e Temulentia* is the same as *Delirium Tremens*. The French use the term *Ivresse* for drunkenness, *Ivrognerie* for habitual drunkenness.

TEMULENTIA, Temulency.

TENACULA. (pl. of *Tenaculum*.) Bonenippers—t. Volsella, Forceps (lithotomy).

TENAC'ULUM, (*tenax*, *tena'cis*, 'holding fast,' from *teneo*, 'I hold,') A kind of fine hook attached to a handle which is thrust through the parietes of a bloodvessel to draw it out and enable it to be tied.

TENACULUM, ASSALI'NI'S. An ingenious instrument, invented by Assalini. It consists of a forceps, or double tenaculum, the points of which are fine, sharp, and bifurcated, so as to be readily received into each other when the instrument is closed on the artery, which it is by a spring. It is of great use when the surgeon has no assistant at hand. The instrument is figured in S. Cooper's "First Lines."

TENAILLE INCISIVE, (from *tenacula*.) Bonenippers.

TÉNALGIE CRÉPITANTE, Aî.

TENASMUS, Tenesmus.

TENDINES CORONARIÆ, Coronary tendons.

TEN'DINOUS, *Tendino'sus*, *Tendin'eus*, *Sin'eus*, (F.) *Tendineux*, (*tendo*, gen. *ten'dinis*, 'a tendon,') Relating or appertaining to, or having the nature of tendons.

TENDO, gen. *Ten'dinis*, Tendon.—t. Achillis, Achillis tendo—t. Cordis venosus, Circulus callosus Halleri—t. Oculi, Ligament of Zinn.

TEN'DON, *Ten'do*, *Pronerra'tio*, *Ten'on*, (from *teno*, 'to stretch,') *Sin'eu*, *Thew*, *Leader*, (Prov.) *Guider*. A fibrous cord, more or less round, long, or flattened; of a white, pearl colour, and composed of very close parallel fibres. The tendons are attached to the bones, by one extremity; by the other, they receive the insertions of the fleshy fibres: sometimes they afford attachment to fleshy fibres at both ends:—as in the digastric muscle. The tendons must be considered as so many cords, for transmitting the motion of muscles to the bones or levers.

TENDON D'ACHILLE, Achillis tendo.

TENDON, CORDIFORM, OF THE DIAPHRAGM, Centre, phrenic—t. of Zinn, Ligament of Zinn.

TENDONS, TWITCHING OF THE, Subsalutis tendinum.

TENEOTOMIA, Tenotomy.

TÉNESME, Tenesmus—t. de l'Æsophage, Œsophagismus—t. Vésical, Bradysuria.

TENES'MUS, *Tenes'mus*, (*τενέσμος*.) *Tenas'mus* *Tines'mus*, (from *teno*, 'I stretch,') *Proc'tica* seu *Obstipa'tio* *Tenesmus*, *Proctospas'mus*, *Tinag'mus*, (F.) *Ténésme*, *Épreintes*. Frequent, vain and painful desires to evacuate;—one of the chief symptoms of inflammation of the lining membrane of the digestive tube, as of dysentery.

Used at times, also, for forcing down pains of the uterus and bladder.

TENESMUS VESICÆ, Bradysuria—t. Vesicæ mucosus, Cystirrhœa.

TENETTE, (from (F.) *tenir*, [L.] *teneo*, 'to hold,') see Forceps.

TÉNIA, Tænia.

TÉNIFUGE, Tæniafuge.

TENNESSEE', MIN'ERAL WATERS OF. On the waters of the French Broad River, in this State, is a large, clear, thermal spring. The heat

is so great that, on first going into the water, it is almost insupportable.

TENON. (τενον, gen. τενοντος,) Tendon.

TENONT'AGRA, (tenon, and αγρα, 'seizure,')

A variety of gout, which affects the tendinous parts of muscles, or the tendinous ligaments which strengthen certain articulations.

TENONTOTOMIA, Tenotomy.

TENONTOTROT'I, (tenon, and τρωσκω, 'I wound,') They who are wounded in the tendons.

TÉNOPHYTE, (tenon, and φυειν, 'to grow,') A growth of an osseous or cartilaginous nature on a tendon.

TENOR/RHAPHY, *Tenororrhaphia*, (τενον, 'a tendon,' and ραφη, 'a suture,') Union of a tendon by suture after it has been divided.

TENOSYNITIS, *Tenosynovitis* [?], (F.) *Ténosynite*. An imperfectly formed term. (tenon, syn, and itis.) Inflammation of the synovial sheath of a tendon. See *Ar.*

TÉNOTOME, *Tenotomus*, (from tenon, and τωμ, 'incision,') An instrument for dividing tendons, especially by the subcutaneous method.

TENOTOMY, *Tenotomia*, *Tenectomia*, *Tenontotomia*, *Secl'tio seu Dissectio tendinum*, (F.) *Ténotomy*. The operation of dividing a tendon, often practised for the removal of loxarthrus, or deviation of the joints—as in club-foot.

TENSEUR, Tensor — *t. de l'Aponérose fémoreale*, *Fascia lata muscle*.

TENSION, *Tensio*, *Tensura*, *Tu'sis*, *Diat'asis*, *Enton'ia*, (*Etonia* is, also used for overtension,) *Content'sio*, *Disten'sio*, (from *tendere*, *tensum*, 'to stretch,') State of a part, which is stretched, when its textures are distended by the afflux of fluids, or the accumulation of gas; or when its fibres are drawn, as it were, in an opposite direction, and are, consequently, more resisting than usual.

TENSIVE, *Tensivus*, *Tonodes*, (F.) *Tensif*. Accompanied by tension or by a feeling of tension—as a 'tensive pain.'

TEN'SOR, (F.) *Tenseur*. That which stretches or renders tense.

TENSOR CHOROIDEÆ, Ciliary muscle.—*t. Palati*, *Circumflexus musculus*.

TENSOR TAR'SI, *Lachrymal Muscle*. A small muscle, belonging to the inner commissure of the eyelids, well described by Professor Horner, of Philadelphia, and hence termed *Muscle of Horner*. It is about three lines broad and six long; arises from the posterior flat surface of the os unguis, near its junction with the ethmoid bone, and passes forwards and outwards, lying on the posterior face of the lachrymal ducts. As it approaches the commissure of the lids, it splits into two nearly equal parts, each of which is appropriated to a duct, and inserted along its course, almost to the punctum lacrymale. Its chief office seems to be, to influence the position of the punctum lacrymale, and to regulate the course of the tears through the lachrymal ducts.

TENSOR TYMPANI, *Inter'nus Aur'is*, *Internus Mal'lei*, (F.) *Salpingo-malleus*. A small muscle, which arises from the petrous portion of the temporal bone, and the cartilage of the Eustachian tube, and terminates by a tendon, which is reflected over the processus cochleariformis, and is inserted into the apophysis of the handle of the malleus.

TENSOR VAGINÆ FEMORIS, *Fascia lata muscle*.

TENSURA, (*tendere*, *tensum*, 'to stretch,') Tension.

TENSUS, (*tendere*, *tensum*, 'to stretch,') Penis.

TENT, *Ten'ta*, *Turru'da*, and its dim. *Turru'dula*, *Penic'ulus*, *Penic'il'us*, *Penic'il'um*, *Mol'tos*, *Lemnis'cus*, *Licil'ium*, (F.) *Tente*, *Mèche*, (*tendere*, *tensum* or *tentum*, 'to stretch,') In surgery, tents are small rolls of lint, of a cylindrical or pyramidal

shape, which are introduced into wounds and deep ulcers, to prevent them from closing before they are filled up from the bottom. Tents are sometimes, also, made of prepared sponge, gentian root, slippery elm, &c., &c. They are not so much used as they formerly were.

TENT, SPONGE, see *Spongia*.

TENTACULARIA, *Filaria hominis bronchialis*.

TENTACULUM CEREBELLI, (from *tendere*, *tensum* or *tentum*, 'to stretch,') Tentorium.

TENTA'MEN MED'ICUM. 'A medical trial,' (from *tentare*, 'to try,') An examination in medicine, in the Prussian universities, which is conducted both in writing and *visâ voce* before the Dean of the Medical Faculty.

TENTE, Tent—*t. du Cervelet*, Tentorium.

TENTIGO, gen. *Tentig'inis*, (from *tendere*, *tensum* or *tentum*, 'to stretch,') Priapism—*t. Venerea*, Nymphomania—*t. Veretri*, Satyriasis.

TENTIPEL'LUM, (*tendere*, *tensum* or *tentum*, 'to stretch,' and *pellis*, 'the skin,') Ancient name of a cosmetic, with which, it was pretended, wrinkles could be effaced.

TENTORIUM, *T. cerebelli superextensum*, *T. cerebelli*, *Tentaculum cerebelli*, *Septum encephali*, *Diaphragma cerebri*, *Intersep'ta horizontalia Pucchio'ni*, *Processus transversus duræ matriæ*, *Lat'eral Processes of the Duræ matriæ*, *Transverse septum*, (F.) *Tente du cervelet*, *Plancher du Cerveau*, *Septum transverse* (Ch.), (*tendere*, *tensum* or *tentum*, 'to stretch,') A process of the dura mater, which separates the cerebrum from the cerebellum. It extends from the internal horizontal spine of the occipital bone, directly forwards to the sella Turcica of the sphenoid.

TENTORIUM CEREBELLI, Tentorium.

TENTUM, (from *tendere*, *tensum* or *tentum*, 'to stretch,') Penis.

TENTWORT, *Asplenium ruta muraria*.

TENUIS MATER, *Pia mater*.

TENUITAS AQUOSA, 'watery thinness,' *Prædominium aquæ*.

TEPHRO'SIA APOLLINE'A, (τεφρος, 'ash coloured, gray,') *Gale'ga Apolline'a* of Egypt and Nubia. Order, Leguminosæ. The leaves are often mixed with senna.

TEPHROSIA PURPU'REA, *Gale'ga purpu'rea* of the coast of Coromandel. The root is bitter, and a decoction of it is prescribed by the native doctors in dysentery, lientery, and tympanites.

TEPHROSIA SENNA. The leaves are used for senna by the people of Popayan.

TEPHROSIA TOXICA'RIA, *Gale'ga toxica'ria* of Cayenne. The leaves are used for poisoning fish. It has been suggested as a substitute for digitalis.

TEPHROSIA VIRGINIANA, *Galega Virginiana*.

TEPIDARIUM, (*tepidus*, 'moderately warm,') see *Stove*.

TEPLITZ, Töplitz.

TERAS, gen. *Ter'atis*, (τερας, gen. τερασος,) Monster.

TERATOGEN'ENY, *Teratogen'ia*, (teras, and γενεα, 'generation,') The formation of monsters.

TERATOLOG'ICAL, *Teratolog'icus*, (F.) *Teratologique*. Same etymon as the next. Relating or appertaining to Teratology.

TERATOL'OGY, *Teratolog'ia*, (τερατολογία, from teras, and λογος, 'a discourse,') A treatise on monsters.

TERATOSIS, see *Monster*.

TERCERON, (from [F.] tierce, [L.] tertius, 'third,') see *Mulatto*—*t. Black*, see *Mulatto*.

TREBELLA, (from τερεω, 'to bore,') Trepan.

TREBELLUM. A name given by Dugès to a perforator, *Perce-crâne*, of his invention.

TÉRÉBINTHINE, Terebinthina — *t. de Bordeaux*, see *Pinus sylvestris*—*t. de Chic*, see *Pis-*

tacia terebinthus — *t. Commune*, see *Pinus sylvestris* — *t. de Copahu*, Copaiba — *t. d'Égypte*, see *Amyris opobalsamum* — *t. de Giléad*, see *Amyris opobalsamum* — *t. du Grand Caïre*, see *Amyris opobalsamum* — *t. de Judée*, see *Amyris opobalsamum* — *t. de Méléze*, see *Pinus larix* — *t. de Venise*, see *Pinus larix*.

TÉRÉBINTHINÉE, Terebinthinate.

TÉRÉBINTHINA, (from *τερεβινθος*, 'the turpentine tree.') *Tur'pentine*, *Bo'tin*, *Al'botim*, *Al'til'ibat*, *Bruti'no*, (F.) *Térébinthine*. A substance, of the consistence of honey, which flows from many trees of the terebinthaceæ and coniferae families. It is viscid; shining; more or less transparent; inflammable; of a warm, pungent taste; strong smell; and is entirely composed of resin and essential oil, without benzoic acid. It is soluble in alcohol. See *Turpentine*, *Pinus palustris* and *Pinus sylvestris*. All the turpentineæ are stimulant, diuretic, and cathartic. *Terebinthina*, (Ph. U. S.), is the concrete juice of *Pinus palustris* and other species of *Pinus*.

TÉRÉBINTHINA ARGENTORATENSIS, see *Pinus picea* — *t. Balsamea*, see *Pinus balsamea* — *t. Canadensis*, see *Pinus balsamea* — *t. Chia*, see *Pistacia terebinthus* — *t. Communis*, see *Pinus sylvestris* — *t. Cypria*, see *Pistacia terebinthus* — *t. Empyreumatica*, see *Pinus sylvestris* — *t. Laricina*, see *Pinus larix* — *t. Veneta*, see *Pinus larix* — *t. Vera*, see *Pistacia terebinthus* — *t. Vulgaris*, see *Pinus sylvestris*.

TÉRÉBINTHINATE, *Terebinthina'tus*, (F.) *Térébinthiné*, *Abiétique*. Having the qualities of or impregnated with turpentine.

TÉRÉBINTHUS, (*τερεβινθος*.) *Terminthus* — *t. Gummifera*, *Bursera gummifera* — *t. Lentiscus*, *Pistacia lentiscus* — *t. Vulgaris*, *Pistacia terebinthus*.

TÉREBRA, (*τερεω*, 'to bore,') 'a borer,' *Trepan*.

TÉRÉBRANT, see *Terebratio*.

TÉRÉBRATIO, (from *terebrare*, *terebratum*, 'to bore,') *Trepanning*. The French use the term *Térébration* for the puncturing of a bone by any instrument, and a pain is said to be *térébrant*, when it seems as if the part was penetrated or bored.

TÉRÉBRATION, see *Terebratio*.

TÉREDO OSSIIUM, (*τερεω*, 'to bore,') *Caries*, *Spina ventosa*.

TÉRES, gen. *Ter'etis*, ('long and round,' from *tero*, 'to rub;' as if 'rubbed smooth or rounded,') *Stron'gyllus*, *Cylindroi'des*. An epithet given to many organs, the fibres of which are collected into round fasciculi.

TÉRES LIGAMEN'TUM, (F.) *Ligament rond*. This name is given to the round ligament in the cotyloid cavity of the os innominatum, which is attached to the head of the bone, and to the bottom of the cavity. It is, also, given to a small fibrous fascia, which extends from the coronoid process of the ulna to below the bicipital tubercle of the radius.

TÉRES MA'JOR, (F.) *Anguli-scapulo-huméral*, *Scapulo-huméral*, (Ch.) *Grand rond*, is situate at the inferior and posterior part of the shoulder. It is attached, on the one hand, to the outer surface of the scapula; to the corresponding part of its axillary margin; and, on the other, to the posterior margin of the bicipital groove of the humerus. It carries the arm backwards and inwards, and carries it, also, inwards in rotation. When it acts with the longissimus dorsi and pectoralis major, it applies the arm against the lateral part of the chest, and keeps it forcibly there.

TÉRES MI'NOR, *No'nus hu'meri Placenti'ni*, (F.) *Le plus petit sus-scapulo-trochitérien* (Ch.), *Petit rond*, *Margini-sus-scapulo-trochitérien*, is situate at the posterior and inferior part of the shoulder; it is narrow and flattened, from above to below,

in its inner half; and, from before to behind, in the outer. It is attached, on the one hand, to the outer surface of the scapula, near its inferior angle; and, on the other, to the inferior part of the great tuberosity of the humerus. It depresses the arm, and causes it to turn on its axis from within outwards. It also carries it backwards.

TÉRESIS, (*τήρησις*,) *Observation*.

TÉRETRIUM, (*τερετριον*, from *τερεω*, 'to bore,') *Trepan*.

TÉRETRON, (*τερετρον*,) *Trepan*.

TERGAL, (from *tergum*, 'the back,') *Dorsal*.

TERGEMINI, *Trigemini*.

TERGUM, *Dorsum*, *Vertebral column*.

TERMINAL CAVITIES, see *Pulmo*.

TERMINALIA, see *Myrobalanus*.

TERMINOLOGY, (*termen*, gen. *term'inis*, 'a term,' and *logos*, 'a discourse,') *Nomenclature*.

TERMIN'THUS, *Terebin'thus*, (*τερεβινθος*, 'the turpentine tree,') *Phy'na An'thrac*, *Termin'thus*, *Ber'ry* or *Fungous car'buncle*. The ancients gave this name to a tumour, surmounted by a black pustule, and resembling the fruit of the turpentine tree.

TERMIN'THUS, *Pistacia terebinthus*.

TERMINUS ('limit') **SUPERIOR ASPERÆ ARTERIÆ**, *Larynx*.

TERMONOLOGY, (*termo*, gen. *termo'nis*, 'a term,' and *logos*, 'a discourse,') *Nomenclature*.

TERMS, (periods,) *Menses*.

TERRA ABSORBENS MINERALIS, 'absorbent mineral earth,' *Magnesia carbonas* — *t. Aluminis*, *Argilla pura* — *t. Amara aërata*, *Magnesia carbonas* — *t. Amara sulphurica*, *Magnesia sulphas* — *t. Argillacea hydrata*, *Argilla pura* — *t. Argillacea pura*, *Argilla pura* — *t. Bolaris*, *Argilla pura* — *t. Foliated*, *Sulphur* — *t. Foliated mineralis*, *Soda acetata* — *t. Foliated tartari*, *Potassa acetat* — *t. Fornacum*, see *Bricks* — *t. Fullonica*, *Cimolia purpurescens* — *t. Japonica*, *Catechu*, *Nauclea gambir*.

TERRA LEM'NIA, *Lemp'nias*, *Argil'la bo'lus fla'va*, *Lem'nian Earth*. A name given by the ancients to a solid, reddish, astringent, substance, prepared with the pulp of the fruit of the Baobab, an Africa tree, according to Prosper Alpinus. It seems, however, to be an argillaceous earth. It has been employed as an astringent.

TERRA LIVON'ICA. A *terra sigilla'ta* or *sealed earth* from Livonia, redder than that from Silesia, and very astringent.

TERRA MARITA, *Circumca longa*.

TERRA NOCERIA'NA. A whitish, soft, and astringent earth, found in the environs of Nocera, in Italy.

TERRA ORLEA'NA, *Pigmen'tum uru'cu*, *Uru'cu*, (F.) *Rocou*, *Roucou*, *Bichet*. A pigmental matter obtained from the seeds of *Bix'a Orlea'na* sea *Orella'na* seu *America'na*, *Orella'na*, *Orlea'no*. In Jamaica, it is recommended in dysentery, and is considered to possess astringent and stomachic qualities. It is called, according to its shape, *Flag*, *Roll*, or *Egg Annotto*.

Span'ish Annotto, *Bix'a*, is in small, oblong cakes. It is chiefly employed as a colouring matter.

TERRA OS'SEA, *Mate'ria os'sea*. Bony matter. The earthy portion of bones.

TERRA PONDEROSA, *Baryta* — *t. Ponderosa muriata*, *Baryta*, *muriate* of — *t. Ponderosa salita*, *Baryta*, *muriate* of.

TERRA PORTUGAL'ICA, *Bar'ros*, *Bucaros*. A reddish, astringent, styptic earth, obtained from Portugal.

TERRA SAPONARIA, *Cimolia purpurescens*.

TERRA SIGILLA'TA, (*sigillatus*, furnished with little figures — *sigilla*,) *Seal'd Earth*. A bole or earthy matter, made into little cakes or flat masses, and stamped with certain impressions, as with the head of the Grand Seigneur. To these

belong the *Terra Silesiaca*, *Terra Lemnia*, *T. Turcica*. See Bole, and Bolus alba.

TERRA SILESIACA, see *Terra sigillata*—*t. Talcosa oxyanthracodes*, *Magnesia carbonas*—*t. Turcica*, see *Terra sigillata*.

TERRA URINÆ. The earthy deposit in the urine.

TERRA VITRIOLI DULCIS, *Colcothar*.

TERRAPIN, *Emys palustris*.

TERRE BOLAIRE, 'bolar earth,' *Bolus*—*t. Magnésienne*, *Magnesia carbonas*—*t. Noie*. *Bunium bulbocastanum*—*t. Pesante*, *Baryta*—*t. Sigillée*, see *Bole*.

TERRETTE, *Glechoma hederacea*.

TERRITORY, *CELL*, see *Cell*.

TERTIAN FEVER, *Febris tertiæ*, *Anctus tertius*, *Tritæus*, *Tertian* or *Third Day Ague*, *Third Day Fever*, (F.) *Fièvre tierce*, (from *tertius*, 'the third.') An intermittent, whose paroxysms recur every third day, or every 48 hours. The mildest, and the most pernicious, intermittents belong to this head. As a general rule, it is the most manageable form of ague.

A double tertian, (F.) *Double tierce* (F.) *Fièvre*), has a paroxysm every day,—the paroxysms of alternate days being associated.

TERTIANARIA, (from its virtues in tertian fever,) *Scutellaria galericulata*.

TESSARA, (*τεσσαρα*, 'four,') *Cuboid*.

TESSELLA, (dim. of *tessera*, 'a small quadrangular stone for paving,' &c.,) *Tabella*.

TESSELLATED, *Tessellatus*, ('furnished with small quadrangular stones,') Having a quadrangular arrangement, like a pavement. See *Epithelium*.

TESSERA, 'a quadrangular stone, wood,' &c. *Cuboid*.

TEST, **DANIEL'S**, (from *testis*, 'a witness,') see *Docimasia pulmonum*—*t. Hydrostatic*, see *Docimasia pulmonum*—*t. Plouquet's*, see *Docimasia pulmonum*—*t. Static*, see *Docimasia pulmonum*.

TESTA, 'a shell,' see *Ostrea*—*t. Ovi*, see *Membrana putaminis*.

TESTA PRÆPARATA, *Prepared Oystershell*. Oystershell, freed from extraneous matter, washed with boiling water, reduced to powder, and prepared in the same manner as *Creta præparata*.

TESTACEOUS, *Testaceus*, (F.) *Testacé*, (from *testa*.) A powder, consisting of burnt shells. These contain carbonate of lime chiefly, and hence the term has been applied to cretaceous substances. See *Creta*.

TESTÆ FURNACEÆ, (*fornax*, gen. *fornacis*, 'a furnace,') *Bricks*—*t. Ostree*, see *Ostrea*.

TESTES, (pl. of *testis*, 'a witness,') The Testicles, see *Testicle*. Vulgarly, the *Stones*, *Ballocks*. A name, also, given to the inferior tubercles of the corpora quadrigemina, to distinguish them from the superior, called *Nates*.

TESTES CEREBRI, see *Quadrigenia tubercula*.

TESTICLE, *Testis*, *T. virilis*, *Testiculus*, *Orchis*, *Po'mum amoris*, *Didymus*, *Androarrium*, *Her'nia*, *Gemini* (pl.), *Gemelli* (pl.), *O'rium*, *Codling*, (F.) *Testicule*, (dim. of *testis*, because the testicles are 'witnesses' of virility.) The name of two glandular organs, contained in the scrotum, whose office is to secrete sperm. The substance of the testicle consists of numerous conical, flattened lobules—*lobuli testis*—whose bases are directed towards the surface of the organ, and the apices towards the corpus Highmorianum. They are formed of a gray and softish substance, composed of tortuous canals, called *Tubuli* seu *Canalesculi seminiferi* seu *seminales*, *Vasa seminalia*, *Vascula serpenti'na*, folded on each other, and of extreme fineness; the number of which, according to the estimate of Monro, is 62,500, and the total length 5,203 feet. All these canals, uniting in their course, pass through the corpus

Highmorianum, concur in forming the epididymis, and give origin to the vas deferens. The testicles are covered, immediately, by a fibrous membrane, of an opaque white colour, and very tough, called *Tunica Albuginea*, or, simply, *Albuginea*, *Peritestis*, (F.) *Pérididyme*, of which the corpus Highmorianum is only an enlargement. Over this, again, is the tunica vaginalis, the second envelope of the testis.

TESTICLE, **SWELLED**, *Hernia humoralis*.

TESTICONDUS, (*testis*, and *condere*, 'to hide,') *Cryptorchis*.

TESTICULAR, *Testicularis*, (F.) *Testiculaire*. Belonging or relating to the testicle.

TESTICULE, *Testicle*.

TESTICULUS, *Testicle*—*t. Accessorius*, *Epididymis*—*t. Caninus*, *Orchis mascula*.

TESTIMONIUM MATURITATIS. 'Evidence of fitness.' A certificate of complete education, required, in the Prussian universities, of a candidate, before he can be admitted to examination for the degree of Doctor of Medicine.

TESTIS, *Testicle*—*t. Femineus*, *Ovary*—*t. Irritabile*, *Orchidalgia*—*t. Minor*, *Prostate*—*t. Muliebris*, *Ovary*—*t. Virilis*, *Testicle*.

TESTITIS, (from *testis*, and *itis*,) *Hernia humoralis*.

TESTUDINATIO CRANII, 'arched like the back of a tortoise,' (from *testudo*,) *Camarosis*.

TESTUDO, gen. *Testudinis*, (from *testa*, 'a shell,') *Chelonæ*, (F.) *Tortue*. A genus of reptiles, including the turtle, the flesh of which is much esteemed as an article of diet.

Also, an encysted tumour, which has been supposed to resemble the shell of a testudo; *Emphy'ma encystis ganglion*. See *Talpa*.

TESTUDO CEREBRI, *Fornix*—*t. Scapulae*, see *Scapula*.

TETE, (teat,) *Nipple*.

TETANIC, *Tetanicum*, *Spas'ticum*, *Convulsivum*, *Convulsivæ*, (F.) *Tétanique*, (from *tetanus*.) A remedy which acts on the nerves, and, through them, on the muscles, occasioning, in large doses, convulsions. The chief agents of this class are *Nux Vomica*, *Strychnia*, *Brucia*, *Arnica*, and *Toxicodendron*.

Also, an epithet for a tonic convulsion.

TETANIN, *Strychnia*.

TETANIUM, *Strychnia*.

TETANIZATION, *Tetania'tio*. The production or condition of tetanic spasms, in which the patient is said to be *tet'itized*.

TETANOMATA, (*τετανωματα*,) *Tetanothra*.

TÉTANOS, *Tetanus*.

TETANOTHRA, (*τετανωθρα*,) *Tetanom'ata*, *Eru'gato'ria*, (from *τεταω*, 'stretched, free from wrinkles,') Medicines which remove wrinkles.

TETANUS, (*τεταως*, 'stretched; from *τετω*, 'to stretch,') *Spasm with rigidity*, *Convulsio Indica*, *Holoton'ia*, *Holoton'icus*, *Holotet'anus*, *Tet'anus universalis*, *Ri'gor nervosus* seu *nervor'um*, *Extensio* seu *Distensio*, *Ento'ia tet'anus*, *Cat'o'chus cervi'nis* seu *holoton'icus*, (F.) *Tétanos*, *Mal de cerf*. A disease which consists in a permanent contraction of all the muscles or merely of some, without alternations of relaxation. It is characterized by closure of the jaws; difficulty or impracticability of deglutition; rigidity and immobility of the limbs and trunk, which is sometimes curved forwards (*Emprosthotonus*), sometimes backwards (*Opisthotonus*), and sometimes to one side (*Pleurothotonus*). When tetanus is confined to the muscles of the jaws, it is called *Tris'mus*. It is a most formidable affection. The means of treatment are:—copious and repeated blood-letting; bathing, cold and warm; powerful doses of opium, and anæsthetics. In *Traumat'ic Tetanus*, or that which supervenes on a wound, every extraneous matter must, of course,

be removed from the wound, and soothing applications be made to it.

TETANUS ANTICUS, Emprosthotonus—t. Dolorificus, Cramp—t. Dorsalis, Opisthotonus—t. Lateralis, Pleurothotonus—t. Maxillæ inferioris, Trismus—t. Posterganeus, Opisthotonus—t. Posticus, Opisthotonus—t. Universalis, see Tétanus.

TETARTÆUS, (τεταρταῖος, 'occurring on the fourth day,' from τεταρτός, 'the fourth,') Quartan.

TETARTOPHYŪIA, (τεταρτος, 'the fourth,' and φῶω, 'I arise,') *Quartanus remittens*. A quartan, in which the intermission is inordinately short or imperfect.

TÊTE, (from [L.] *testa*, 'the skull,') Caput, Head—t. *de Veau*, Elephantiasis of the Greeks.

TETRA, (τετρα, 'four,') in composition, four.

TETRADRACHMON, (τετραδραχμον, from *tetra*, and δραχμη, 'a drachm,') A weight of four drachms or half an ounce.

TETRAGONUS, (τετραγωνος, from *tetra*, and γωνία, 'an angle,') *Platysma myoides*.

TETRAHIT LONGIFLORUM, *Galeopsis grandiflora*.

TETRAMASTHUS, (*tetra*, and μασθος, 'a breast,') *Quadrimummatus*.

TETRA'MYRON, (τετραμυρον, from *tetra*, and μυρον, 'an ointment,') An ointment of four ingredients.

TETRANGURIA, (*tetra*, and *anguria*,) *Cucurbita citrullus*.

TETRANTHERA PICHURIM, (*tetra*, and ανθηρα, 'anther,') *Pichurim* beans.

TETRAO COTUR'NIX, (after τετραξ, *Cotur'nix*, the Quail, (F.) *Caille*: *Class*, Aves; *Ord*, Gallinaceæ. The quail is an esteemed article of diet. The Greek name is ορτυξ, gen. ορτυγος; and the places, called *Ortygia*, are named after it. The excrements of the quail were formerly extolled in epilepsy, and the fat in specks of the eye.

TETRAPHAR'MACUM, (*tetra*, and φαρμακον, 'a medicine,') A medicine consisting of four ingredients.

TETRAS'CELUS, (*tetra*, and σκελος, 'leg,') A monster having four legs.

TETROB'OLON, (τετροβολον, from τετρα, and οβολος, *obolus*,) The weight of four oboli or two scruples.

TETROROS, (τετρορος,) *Astragalus*.

TETER, ([Sax.] *teeten*,) Herpes—t. Crusted, Impetigo—t. Honeycomb, Porrigo favosa—t. Humid, Eczema impetigo—t. Milky, Porrigo larvalis—t. Pustular, Impetigo—t. Running, Impetigo—t. Sealy, Psoriasis.

TETTERWORT, *Chelidonium majus*.

TEUCRIUM ÆGYPTIACUM, (τευκριον, after Teucer, T. capitatum—t. Belium, T. capitatum.

TEUCRIUM CAPITATUM, T. *Belium* seu *Egyptiacum*, *Po'lium Capitatum*, *Poley Mountain* of Montpellier, *Ord*, Labiata, is, generally, substituted for T. *chamædrys*. The common *Poley Mountain* is the *Teucrium montanum*; and the *Golden Poley Mountain*, the *Teucrium Po'lium*, *Po'lium*.

TEUCRIUM CHAMÆDRYS, T. *officinale*, *Chamædrys*, C. *minor* *repens* seu *vulgaris*, *Quer'cula minor* seu *Calanandrina*, *Trissa'go*, *Chamædrops*, *Trixa'go*, *Com'mon*, *Creeping* or *Small German'der*, *English Treacle*, (F.) *Petit chène*, *Chénette*, *Germandrée officinale*. This is possessed of aromatic and bitter properties. The dose of the dried powder is from ʒss to ʒj.

TEUCRIUM CHAMÆPITYS, (*Chamæpitys*, *Arthetica*, *Arthetica*, *Ajū'ga*, A. seu *Bulgula Chamæpitys*, *Ab'go*, *Chamæmorus*, *P'va arthritica*, *Holc'yon*, *Io'nia*, *Sideritis*, *Com'mon Ground'pine*, (F.) *Petite Ivette*. Properties like the last. The tops and leaves have been recommended as aperients and tonics; especially in female obstructions, and in paralytic disorders.

TEUCRIUM CRETICUM, T. *hyssopifolium* seu *Rosmarinifolium*, *Rosmarinus italicus* *fo'cie*, *Poley-mountain* of Candy, *Po'lium Creticum*. The tops and whole herb enter into the old compounds, *Mithridate* and *Theriaca*. It has a moderately aromatic smell; and a nauseous, bitter taste, and is placed among the aperients and corroborants.

TEUCRIUM FLAVUM has similar properties to T. *chamædrys*.

TEUCRIUM HYSSOPIFOLIUM, T. *Creticum*.

TEUCRIUM I'VA, *Chamæpitys moschata*, *I'va*, I. *moschata* *Monspelien'sium*, *Chamæpitys anthyllus*, *French Ground'pine*, (F.) *Ivette musquée*. It has similar virtues to T. *chamædrys*, but is weaker.

TEUCRIUM MARITIMUM, T. *Marum*.

TEUCRIUM MA'RUM, T. *maritimum*, *Ma'rum Syriacum* seu *Creticum* seu *verum* seu *Cortusi*, *Maj'ra'na Syriaca*, *Chamædrys incana* *maritima* seu *ma'rum*, *Orig'anum Syriacum*, *Ma'rum german'der*, *Syria'na herb mas'tich*, *Cut thyme*, (F.) *Marum* ou *Germandrée maritime*, *Herbe aux chats*. A very aromatic plant, of a camphorated smell; formerly much used in medicine as a tonic, antispasmodic, emmenagogue, &c.

TEUCRIUM OFFICINALE, T. *chamædrys*—t. *Palustre*, T. *scordium*—t. *Pyramidale*, *Ajuga*—t. *Rosmarinifolium*, T. *Creticum*.

TEUCRIUM SCORDIUM, T. *palustris*, *Scor'dium*, *Trissa'go palustris*, *Chamædrys palustris* seu *scor'dium*, *Allium redolens*, *Wa'ter German'der*, (F.) *Germandrée scorodone*, *Sauge des bois*, *Germandrée aquatique*. The leaves have a garlicky smell, and bitterish, slightly pungent taste. It has the tonic properties of the other *Teucria*.

TEXTUM INTERLOBULARE, (*texere*, *textum*, 'to weave,') Interlobular tissue.

TEXTURA, *Texture*—t. *Organica*, *Histos*.

TEXTURAL, *Texturalis*. Relating or appertaining to a texture.

TEXTURE, *Textura*, *Textus*, (from *texere*, *textum*, 'to weave,') *Te'la*, *Hy'pha*, *Hy'phæ*. The particular arrangement of the tissues that constitute an organ.

TEXTUS, *Texture*, *Tissue*—t. *Cellulosus*, *Cellular tissue*—t. *Desmosus*, *Desmoid tissue*—t. *Erectilis*, *Erectile texture*—t. *Nervorum*, *Plexus nervorum*—t. *Organicus*, *Cellular tissue*—t. *Papillaris*, *Corpus papillare*—t. *Parenchymalis*, *Cellular tissue*.

THÆRIA (*Thp*, 'a wild animal,') *Radzyge*.

THAL'AMUS, pl. *Thal'ami*, (*θαλαμος*,) (F.) *Couche*. 'A room or chamber; a bed.' The place in which a nerve originates, or has been considered to originate.

THALAMI NERVO'UM OPTICO'rum seu *Op'tici*, *Eminentie magnæ cer'ebrî*, *Gan'glia cer'ebrî postica*, *Cru'ra medullæ oblongatæ* (of some,) *Striata corpora super'na posteriora*, *Colliculi nervo'rum optico'rum*, *Optic thalami*, *Posterior cer'ebral gan'glion*, (F.) *Couches des nerfs oculaires* (Ch.), *C. des nerfs optiques*, *C. optiques*. Two rounded and irregular surfaces, which are seen exposed in the lateral ventricles of the brain, and in the third ventricle, the inferior surface of which presents two projections, called *Corpora geniculata*, that furnish several filaments to the optic nerves. The name was given from a belief that the optic nerves originate from them. They, however, arise more posteriorly from the corpora quadrigemina or optic lobes; and adhere merely to the inner margin of those bodies. Gall considers, that the thalami act as ganglions to the nerves: and, hence, he calls them *Grand ganglion cérébral inférieur*. They are, also, called *Posterior gan'glion of the cerebrum*. From the thalami and corpora striata fibres proceed upwards to constitute the convolutions of the brain, and the various bands that connect the different parts of

the brain together. The upper and inner parts of the thalami are so closely connected as to form one continued surface, called *Commissura mollis*. The posterior parts turn downwards and outwards; after which they are elongated to form the two white cords, termed *Tractus optici*.

THALAMUS REGALIS. Pericardium.

THALASSOLUTRUM, (θαλασσα, 'sea,' and λουτρον, 'a bath,') Bath, sea.

THALASSOMELI, (θαλασσα, 'sea,' and μελι, 'honey,') A cathartic medicine, composed of equal parts of sea water, honey, and rain water, exposed to the sun during the dog days in a pitched vessel.

THALICTRON, (θαλιτρον,) *Thalictrum*, *T. magnum* seu *flavum* seu *flavescentis* seu *pratensis* seu *nigrificans* seu *rugosum* seu *virginatum*, *Rhabarbarum purpureum*, *Meadow rue*, *Poor Man's Rhubarb*, (F.) *Pigamon jaunâtre*, *Rue des prés*, *Fausse-Rhabarbe*, *Rhabarbe des pauvres*. Ord. Ranunculaceæ. *Sex. Syst.* Polyandria Polygynia. The root resembles rhubarb in its properties.

THALICTRUM ANEMONOIDES, *Meadow Rue*, *Rue-leaved Anemone*; an indigenous plant, flowering in April and May.

THALLIUM, (from θαλλος, [L.] *thallus*, 'a green bud or shoot,') A metal discovered in 1861 by spectrum analysis in which it gives a very brilliant green ray. It is found in certain seleniferous and telluriferous deposits, in the native sulphur of Lipari, &c., and resembles lead physically, in its aspect, colour, density, &c., as well as its toxic action on the economy.

The salts of thallium, especially the nitrate, are very soluble.

THAMARINDUS, *Tamarindus*.

THAME, (thumb,) Pollex.

THANASIMUS, (θανασμος, from θανατος, 'death,') Mortal.

THANATICUS, (from *thanatus*.) Relating or appertaining to death. *Thanatici* (morbi) of Dr. Wm. Farr are fatal affections from violence.

THANATODES, (θανατωδης,) Mortal.

THANATOID, *Thanatoïdes*, (*thanatus*, and εidos, 'resemblance,') Resembling death. Apparently dead.

THANATOL'OGY, *Thanatolog'ia*, *Thnesecolog'ia*, (*thanatus*, and λογος, 'a discourse,') A treatise on, or the doctrine of, death.

THANATUS, (*Savatus*), Death.

THAP'SIA, (from the island Thapsus.) The deadly car'rot, *Thapsia asclepias*. Family, Umbelliferæ. *Sex. Syst.* Pentandria Digynia. The bark of the root operates violently, both upwards and downwards. It is not used.

THAPSIA SILPH'IUM. A North African plant, has similar properties. Six to ten grains of the powder produce several alvine evacuations.

THAPSIUM ACTÆIFOLIUM, (after the isle of Thapsus,) Ligusticum actæifolium.

THAPSIUM ATROPURPUREUM, *Th. cordatum*, *Roundheart*. Order, Umbelliferæ. The flowers, which appear in June, are of a dark purple.

THAPSIUM BARBINOÏDE, *Meadow Parsnep*. An indigenous plant, of the Northern and Western States, which flowers in June. Flowers, yellow.

Both these plants have been esteemed vulnerary, antisyphilitic, and diaphoretic; and as antidotes to the bite of a rattlesnake.

THAPSIUM CORDATUM, *T. atropurpureum*.

THAPSUS BARBATUS, *Verbascum nigrum*.

THARSI, see Tarsus.

THE, Thigh.

THÊ, *Thea*—*t. des Apalaches*, *Ilex vomitoria*—*t. du Canada*, *Gaultheria*—*t. d'Europe*, *Salvia veronica*—*t. de France*, *Salvia*—*t. de Jersey*, *Celastrus*—*t. de la Mer du Sud*, *Ilex vomitoria*—*t. du Mexique*, *Chenopodium ambrosioides*—*t. de Simon Pauli*, *Myrica gale*—*t. Suisse*, *Falltrank*.

THE'A, (Chinese tcha,) *Cha'a*, *Tea*, (F.) *Thê*

There are two principal varieties of teaplant; *Thea Sinen'sis*, the *Thea Bohe'a*; and *Thea viridis*; the black tea, and the green. Ord. Camelliæ. *Sex. Syst.* Polyandria Monogynia. In commerce, many kinds of tea are met with. Several of the differences between these appear to result from the age at which the leaves have been collected, and the mode of their desiccation.

Teadrinking was introduced into Europe about the year 1666; since which time its use has become almost universal; and, in spite of the assertions of medical terrorists, it is, except in particular idiosyncrasies, entirely harmless. It is not unfrequently adulterated. See Falsifications, table of.

By an analogical transference, very common in language, the word *Tea* has been used almost synonymously with infusion, as *Beef tea*, *Mint tea*, &c.

THEA GERMANICA, *Veronica*.

THEBE'SIUS, VEINS OF. A name given to supposititious branches of the coronary veins, which Christopher Adam Thebesius, a German anatomist, described as opening into the heart by small depressions observable in the right auricle; and which have been called *Foramina Thebesii*. No such veins can be demonstrated. The valve, at the orifice of the coronary vein, in the right ventricle, is called *Valvula Thebesii*, *V. Guiffartiana*, *V. venæ magne*.

THECA, (θηκη,) Case, Vagina, Sheath—*t. Cerebri*, Cranium—*t. Cordis*, Pericardium—*t. Folliculi*, see Folliculi Graafiani—*t. Vertebralis*, see Vertebral canal.

THE'CAL, *Thecalis*, (from *theca*, 'a sheath,') Relating or appertaining to a sheath,—as of a tendon.

THEC'I'TIS, (*theca*, and *itis*.) A term, of hybrid formation, signifying inflammation of the sheaths of tendons.

THEIN, (from *theca*.) see *Coffea Arabica*.

THEION, (θειον,) Sulphur.

THEIOPEGÆ, (*theion*, and *πηγη*, 'a spring,') Waters, mineral (sulphureous.)

THEIOTHERMÆ, (*theion*, and *θερμη*, 'heat,') Waters, mineral (sulphureous.)

THELASIS, (θελαιζω, 'to give suck,') Lactation.

THELASMUS, (θηλασμος,) Lactation.

THELASTRIA, (θηλαστρια,) Nurse.

THELE, (θηλη,) Mamma, Nipple.

THEL'ITIS, (F.) *Thélite*, (*thele*, and *itis*.) Inflammation of the nipple.

THELONCUS, (*thele*, and *ογκος*, 'a swelling,') Mastoncus.

THELORRHAG'IA, (*thele*, and *ρρηγη*, 'a rupture,') Hemorrhage from the nipple.

THELYGON'IA, (θηλυς, 'female,' and *γονη*, 'generation,') The part taken by the female in the act of generation. The procreation of female children. Also, nymphomania.

THELYGONUM, (θηλυς, 'female,' and *γονη*, 'sperm,') Sperm (of the female.)

THELYPTERIS, (θηλυς, 'female,' and *pterus*,) *Pteris aquilina*.

THENAD, *Thenal*.

THE'NAL, *The'nar*, (from *thenar*.) Relating or appertaining to the thenar.

THENAL AS'PECT. An aspect towards the side on which the thenar is situated.—(Barclay.) *Thenad* is used adverbially, by the same writer, to signify 'towards the thenal aspect.'

THE'NAR, (θεναρ, from *δενω*, 'I strike,' 'I push,') The palm of the hand, or sole of the foot.

THENAR, *Flexor brevis pollicis manus*, Palm.

THENAR or THENAL EM'INENCE is the projection at the anterior and outer part of the hand formed by the abductor brevis, opponens, and flexor brevis pollicis.

THENAR or **THENAL MUSCLE**. Riolan and Winslow give this name to the fleshy mass, formed of the abductor brevis, opponens pollicis, and the interior part of the flexor brevis pollicis. In the foot, Winslow gives the name *Thenal muscle* to the abductor and flexor brevis pollicis pedis.

THEOBROMA CACAO, (Θεός, 'God,' and βρώμα, 'food,') Cacao.

THEOMANIA, (Θεομανία, from Θεός, 'God,' and μανία, Demonomania.)

THEOPLEGIA, (Θεός, 'God,' and πλεγμα, 'a stroke,') Apoplexy.

THEOPLEXIA, (Θεός, 'God,' and πλεσσω, 'to strike,') Apoplexy.

THEORETICAL, (θεωρητικός,) *Theoreticus*, *Theoricus*, (θεωρω, 'I contemplate,') (F.) *Théorétique*, *Théorique*. That which is confined to theory, or is connected with it. An epithet also applied to a sect of physicians, who founded their doctrine chiefly on reasoning.

THEORIA, (θεωρία,) Theory.

THEORIQUE, Theoretical.

THEORY, (θεωρία,) *Theoria*, (F.) *Théorie*. The speculative part of a science. The connexion established in the mind between a general fact, or the least possible number of general facts, and all the particular facts dependent thereon: for example, the motions of the heavenly bodies, and the most important natural phenomena are connected with a single fact, previously known by observation; viz. that the force of gravity acts inversely according to the square of the distance. This constitutes the theory of universal gravitation. *Theory* must not be confounded with *system*. *Theory* regards nature as it is, and is a rigid deduction from facts. *System* is too often the creature of the imagination, to which nature is made to bend.

THEORY OF MEDICINE, *Institutes of Medicine*, *Theoretical Medicine*, is that part of the science which attempts philosophically to account for the various phenomena that present themselves during health as well as in disease. It is the philosophy of medicine. The *Institutes of Medicine* are generally considered to comprise Physiology and its applications to Pathology, Hygiene, and Therapeutics. By some they are considered to include General Pathology and General Therapeutics.

THERAPEIA, (θεραπεία,) Curation, Therapeutics.

THERAPEUSIS, Therapeutics.

THERAPEUTA, (θεραπευτής,) Therapist.

THERAPEUTER, (θεραπευτή,) Therapist.

THERAPEUTES, (θεραπευτής,) Therapist.

THERAPEUTICAL, *Therapeuticus*. Relating or appertaining to Therapeutics.

THERAPEUTICE, (θεραπευτική, [τεχνή,]) Therapeutics.

THERAPEUTICS, *Therapeutikē*, *Therapeutica*, *Therapēsis*, *Iatrotechnicē*, *Therapeia*, *Therapiā*, *Curatoria Methodus*, *Methodus Medendi*, *Practice of Physic*, (F.) *Thérapeutique*, *Thérapie*, (from θεραπεύω, 'I wait upon, I alleviate, I attend upon the sick,') That part of medicine, the object of which is the treatment of disease. With others, the department which comprises an explanation of the *modus operandi* of medicines.

THERAPEUTICUS, Therapeutical.

THERAPEUTIQUE, Therapeutics.

THERAPEUTIST, *Therapeutā*, *Therapeutēs*, *Therapeuter*. One who practices therapeutics. A practitioner of medicine. To be a good therapist, a man must be well versed in every department of medicine, and capable of observing and reasoning well. He may be a good observer, and yet a bad reasoner. He cannot practice well unless he is both. Hence, the comparatively small number of good therapists.

THERAPIA, (θεραπεία,) Therapeutics.

THERAPIE, Therapeutics.

THERIA, (θηρ, 'a wild animal,') Radzyge.

THERIAC, *Theriaca*—t. of Antipatri, Antipatri theriaca.

THERIACA, (θηριακά,) *Theriac*, (from θηρ, 'a ferocious or venomous animal,') *Theriacale*, *Melastase*, (F.) *Thériaque*. Also, a medicine, believed to be capable of curing or preventing the effects of the bite of a venomous animal. In this sense it is chiefly used by writers. *Theriac* and *Theriacal* have been used adjectively for medicinal.

THERIACA ANDROMACHI, *Venice Treacle*, is an ancient alexipharmic eluctary; consisting of a farrago of 61 different ingredients, which possessed the most opposite properties. It was invented by Andromachus of Crete, and prepared by order of Nero. It has received various modifications; and to the discredit of the *Faculté de Médecine* of Paris, long held its place in their Codex, with even an additional number of ingredients. The *Electuarium Opium polypharmacum* of the Codex has contained acid ingredients, 5; astringent, 5; bitter, 22; indigenous aromatics, 10; umbelliferous aromatics, 7; balsams and resinous substances, 8; fetid ingredients, 6; narcotics, 1; earthy substances, 1; gummy or amygdaceous, 4; saccharine, 3; total, 72! and one of these the flesh of the viper! A little more than a grain of opium is contained in each drachm of the compound.

THERIACA ARTICULORUM, *Colebicum*—t. Celestis, *Tinctura opii*—t. Communis, *Melasses*—t. *Edinensis*, *Confectio opii*.

THERIACA GERMANORUM. An extract or rob, prepared from juniper berries. Used as a stomachic.

THERIACA LONDINENSIS, *Cataplasma Cymini*. A cataplasmod of cummin seed, bay berries, germanander, snake-root, cloves, honey, and sometimes opium, or syrup of poppies.

THERIACA RUSTICORUM, *Allium*.

THERIAKI. In Turkey, *opium-eaters*, or they who indulge largely in the use of opium, are so called.

THÉRIQUE, *Theriaca*.

THERIATRICA, (θηρ, 'an animal,' and ιατρικά, 'medicine,') Veterinary art.

THERIODES, (θηριώδης, from θηρ, 'a venomous animal,' and οδός,) Ferine.

THERIOMA, gen. *Theriomatis*, *Therion*, (θηριον, from θηρ, 'a venomous animal,' and ομα,) Any extremely malignant ulcer. Sometimes confined to ulcers of the lungs.

THERION, (θηριον,) *Therionoma*.

THERIOTOMY, (θηρ, 'an animal,' and τεμνω, 'to cut,') Zootomy.

THERMA, (θερμή,) Heat—t. *Emphytum*, Animal heat.

THERMÆ, pl. of *therma*, (θερμα,) Warm baths or springs. See Waters (mineral).

THERMÆ FABIANÆ, *Pfeffers*, Mineral Waters of—t. *Piperine*, *Pfeffers*, Mineral Waters of—t. *Plumbaria*, *Plombières*, Mineral Waters of.

THERMAL, *Thermalis*, (from *therma*.) Relating or appertaining to heat. Hot, warm,—as 'a thermal mineral water.'

THERMANTICA, (θερμαντικά,) *Calefacients*.

THERMASMA, gen. *Thermasmatis*, (θερμασμα,) Fomentation.

THERMOLUSIA, (*therma*, and λυω, 'to wash,') Bath, warm.

THERMOMETER, (*therma*, and μετρον, 'measure,') see Heat.

THERMOTHERAPEIA, (*therma*, and *therapeia*, 'treatment of disease,') Treatment by means of heat, and especially of hot air.

THERMUM EMPHYTUM, (*therma*, and εμφυτον, 'innate,') Animal heat, Biolychnium.

THE'SIS, (*thesis*, 'a proposition,' from *τιθημι*, 'to put,') 'A position or proposition.' *Disputatio inauguralis*, *Inaugural Dissertation*. The name usually given to the essay composed by a candidate for graduation in medicine, which he is at times required to defend publicly. Often, however, it is a mere form, giving useless labour and trouble to the student, inasmuch as it is executed as a task, and never afterwards regarded by the preceptor or by others. Sandifort, Haller, Smellie, and Stoll, have published collections of these. In many universities, of Germany especially, the *Disputatio inauguralis* is the public dissertation; whilst the *Theses* are questions put to the candidate, and answered in writing, in the Latin language.

THEVE'TIA AHOUI, (after Andrew Thevet, a French-monk, who travelled in Brazil in the 16th century.) *Ahouai*. The kernels of the nut of this Brazilian tree are said to be a violent poison. In the Antilles its nuts are called *Noix de Serpent*; and they are used against the bites of serpents.

THEW, ([Sax.] *þeow*.) Tendon.

THEY, Thigh.

THICKLISTED, Pury.

THIGH, ([Sax.] *þeah*.) *Femur*, *Femur*, *Cnis*, *Me'rus*, *Me'rium*, (Sc.) *The*, *They*, (F.) *Cuisse*. (Of old, *Femur* meant the thigh generally, and especially the outer part,—*Femur*, the inner.) The part of the lower limb which extends from the pelvis to the leg. The thigh is larger at its superior than inferior part, and has the form of an inverted and truncated cone, slightly depressed from within outwards. *Above*, it is bounded, *anteriorly*, by the groin; *externally*, by the hip; *behind*, by the fold of the nates; and *within*, by the perineal region. *Below*, it is bounded, *anteriorly*, by the prominence of the knee; *posteriorly*, by the ham. It is formed of a considerable number of muscles, bloodvessels, lymphatics, nerves, &c.; and is covered by a strong aponeurosis.

THIGHBONE, *Femur*—t. Neck of the, *Collum femoris*.

THIMBLEWEED, *Rudbeckia laciniata*.

THION, (*Seow*.) Sulphur.

THIRST, ([Sax.] *þyrre*.) *Si'tis*, *Dip'sa*, *Potio'nis deside'rium*, (F.) *Soif*. A simple desire for, or absolute want of liquids. Physiologists are not entirely agreed regarding the seat of thirst; some place it in the fauces; others in the stomach. Its immediate cause is not known. It has been attributed to a dry condition of the nervous papillæ of the mouth and fauces from suppression of the salivary and mucous secretions. This is probably true; but, again, it is owing to the wants of the system,—a supply of fluid being required to compensate the numerous losses that are constantly taking place. Thirst is an *internal sensation*,—an *instinctive want*,—arising from organization, and inexplicable. It is an urgent symptom in many diseases, particularly in those of vascular excitement.

THIRST, EXCESSIVE, Polydipsia—t. Morbid, *Dipsosis*.

THIRSTY, *Dipsodes*.

THISTLE, BLESSED, ([Sax.] *þistel*.) *Centaurea benedicta*—t. Cotton, *Onopordium acanthium*—t. Globe, *Echinops*—t. Holy, *Centaurea benedicta*—t. Ladies', *Carduus Marianus*—t. Milk, common, *Carduus Marianus*—t. Pine, *Arctostaphylos gummifera*—t. Sow, *Sonchus oleraceus*—t. Star, *Centaurea calcitrapa*—t. Yellow, *Argemone Mexicana*.

THLADIAS, (*θλάς*, from *θλαω*, 'to crush,') *Eunuch*.

THLASIAS, (*θλασις*.) *Eunuch*.

THLASIS, (*θλασις*.) Contusion—t. *Depressio*, *Depression*.

THLASMA, gen. *Thlasm'atis*, (*θλασμα*.) *Contu-*

sion—t. *Concussio*, *Concussion*—t. *Stemma*, *Sprain*.

THLAS'PI, (*θλασις*.) *Pent'hyeross*, *Ord. Cruciferae*. See *Syst. Tetradynamia Siliculosa*. Two species of *thlaspi* are directed in some pharmacopœias for medical use:—*Thlaspi arven'se* or *Treacle mustard*; and *Thlaspi campest're* seu *hirsutum* seu *vulgatius*, *Lepid'ium campest're*, *Lep'ia* seu *Ibe'ris campest'ris*, *Mithridate mustard*. The pharmaceutical properties of both kinds resemble those of common mustard.

THLASPI BUR'SA, *Th. bursa'tum*, *Thlaspi* seu *Capwell'a* seu *Ibe'ris* seu *Nasturtium* seu *Rod-schied'ia bur'sa pasto'ris*, *Bur'sa pasto'ris*, *Shepherd's purse*, *Co'covert*, *St. James's wort*, *Poor man's Parmacit'y*, (F.) *Fleur de St. Jacques*, *Bourse à berger* ou à *pasteur*, *Tabouret*. A common European plant, introduced into this country. It is slightly astringent, but is little used.

THLASPI BURSATUM, *Thlaspi bursa*—t. *Nasturtium*, *Lepidium sativum*—t. *Sativum*, *Lepidium sativum*.

THLIBIAS, (*θλίβας*, from *θλιβω*, 'to crush,') *Eunuch*.

THLIPSENCEPHALIA, see *Derencephalia*.

THLIPSENCEPH'ALUS, (*θλίψις*, 'compression,' and *εγκεφαλος*, 'the encephalon,') A monster in whom the skull is open, not merely in the frontal and parietal, but also in the occipital region, a distinct fontanelle not existing.

THLIP'SIS, (*θλίψις*.) *Compres'sio*. Compression, and especially constriction of vessels by an external cause. Oppression.

THNESCOLOGIA, (*θνησκω*, 'to die,' and *λογος*, 'a discourse,') *Thanatology*.

THNETOBLASTE, (*θνητός*, 'mortal,' and *βλαστός*, 'germ,') A term proposed by M. Robin for the heteromorphous and anatomical element of the different tumours called 'cancerous.'

THNETOBLASTIQUE, *Cancerous*.

THOLUS, (*θολός*, 'a vaulted chamber,') *Achiculum*—t. *Diocleus*, *Scapha*.

THOMSO'NIAN. One who practises or believes in *Thomsonianism*.

THOMSONIANISM, *Thom'sonism*. A fanciful doctrine, of which a person of the name of Thomson, of New England, is the founder. One of its leading principles is, that the human body is composed of four elements [?], earth, air, fire, and water; and one of its apothegms, that metals and minerals are in the earth, and, being extracted from the depths of the earth, have a tendency to carry all down into the earth who use them; that the tendency of all vegetables is to spring up from the earth, and therefore to uphold mankind from the grave.

The Thomsonsians are Botanical Doctors. See *Herb doctor*.

THORACENTESIS, *Thoracentesis*.

THORACIQUE, *Thoracic*.

THORAC'IC, *Thorac'icus*, (from *thorax*,) (F.) *Thoracique* ou *Thorachique*. That which relates or belongs to the chest—as *Thoracic vis'cera*, &c.

THORACIC INFERIOR ARTERY, *Arte'ria Thorac'ica Exter'na Inferior* seu *Lon'ga* seu *Mamma'ria externa* seu *Thorac'ica inferior*, (F.) *Deuxième des Thoraciques* (Ch.), arises from the axillary, a little below the preceding, and descends vertically over the latter part of the thorax, and the serratus major anticus. It afterwards bends inwards, becomes subcutaneous, and divides into several branches, which embrace the breast. It gives numerous ramifications to the pectoralis major, serratus anticus, intercostals, ganglions of the axilla, mamma, &c.

THORACIC SUPERIOR ARTERY, *Arte'ria Thoracica Exter'na Superior*, (F.) *Arrière thoracique supérieure*, *Première des Thoraciques* (Ch.), arises from the axillary artery or from the acromial, and descends forwards between the *Pectoralis major*

and P. minor, to which it distributes itself by a great number of branches. In some subjects there are two or three *Arteria thoracica externa superiora*.

THORACIC DUCT, *Ductus thoracicus seu thoracicus posterior seu vertebralis seu chyliifer seu chyli seu lacteus seu rorifer seu Pecqueti seu Pecquetianus*, *Vena alba thoracica*, *Alveus ampullaceus*, *Distibutus lac-tea thoracica*, *Gallac'ia*, *Duct of Pecquet*, *Alimentary Duct*, (F.) *Canal ou Conduit Thoracique ou de Pecquet*, is the duct in which the lymphatics of the lower limbs, abdomen, left superior extremity, left side of the head, neck, and thorax terminate. It begins at the *receptaculum chyli*, which is formed by the union of five or six large lymphatic trunks,—themselves formed from the union of all the absorbent plexuses of the abdomen. The duct ascends into the chest through the pillars of the diaphragm, and by the side of the aorta and vena azygos. It contracts in dimension as far as the 6th dorsal vertebra, when it inclines towards the left hand; ascends behind the arch of the aorta; passes behind the left internal jugular vein, and opens at the posterior part of the subclavian vein of the same side. Its embouchure is furnished with two valves, which prevent the blood from passing from the vein into the duct.

THORACIC LIMBS are the upper limbs; so called because they are articulated with the lateral and upper parts of the chest.

THORACIC, LONG OR INFERIOR, Mammary inferior external artery.

THORACIC NERVES. The *short or anterior thoracic nerves* are two in number. They arise from the brachial plexus, and divide into an anterior and a posterior branch; the former distributed to the pectoralis major muscle, the latter uniting with a branch of the other to form a loop, from which numerous branches are given off to the pectoralis major and p. minor.

The *long thoracic nerve*, *Posterior thoracic*, *External respiratory* of Sir Charles Bell, is a long branch, which arises from the fourth and fifth cervical nerves, immediately after their escape from the intervertebral foramina, and passes downwards to be distributed upon the serratus magnus muscle.

THORACIC REGIONS are the different regions of the chest. Thus we say, *anterior, lateral, and superior thoracic*, &c.

THORACICA, (from *thorax*.) Pectorals.

THORACICS, FIRST OF THE, Mammary superior external artery.

THORACIQUE, Thoracic.

THORACO, (from *thorax*.) In composition, the thorax or chest.

THORACOCENTE/SIS, *Thoracenta'sis*, *Thoracotomy*, (thoraco, and *κέντρος*, 'puncture.') *Paracenta'sis thoracica*. Perforation of the chest to evacuate contained fluid—as in empyema.

THORACOCYSTIS, Thoracocystis.

THORACODID'YMU, (thoraco, and *ἄμμος*, 'a twin.') A monstrosity in which twins are united by the thorax.

THORACODYNE, (thoraco, and *ὄδυνη*, 'pain,') Pleurodynia.

THORACODYNIA, Pleurodynia.

THORACOFACIAL, *Platysma myoides*.

THORACOGASTRODID'YMU, *Did'ymus symphyothoracogastrius*, *Xiphodid'ymus*, (thoraco, *γαστήρ*, 'the belly,' and *ἄμμος*, 'a twin.') A monstrosity in which twins are united by the chest and abdomen.

THORACO-MAXILLO-FACIAL, *Platysma myoides*.

THORACOMETER, (thoraco, and *μετρον*, 'measure,') Stethometer.

THORACOMYODYNIA, (thoraco, and *μυοδυνία*.) Pleurodynia.

THORACOPATHIA, (thoraco, and *πάθος*, 'disease,') Disease or suffering in the chest.

THORACOSCOPIA, (thoraco, and *σκοπεω*, 'to inspect,') see Auscultation.

THORACOSCOPIUM, Stethoscope.

THORACOTOMY, (thoraco, and *τομή*, 'to cut,') Thoracocentesis.

THORACYST'IS, *Thoracocyst'is*, (thorax, and *κύστις*, 'a bladder,') Encysted dropsy of the chest. Hydatids in the chest.

THORAX, gen. *Thorac'is*, (θώραξ, gen. θωρακος, 'a cuirass, a coat of mail,') *Cith'arus*, *Venter medi'us*, *Pect'us*, *Steth'us*, *Scutum pectoris*, *Bir*, *Che'lyx*, *Oas'ea*, the *Breast*, the *Chest*, (F.) *Poitrine*. One of the splanchnic cavities, *Ca'vum thorac'is*, bounded, posteriorly, by the vertebrae; laterally, by the ribs and scapula; anteriorly, by the sternum; above, by the clavicle; and below, by the diaphragm. It is destined to lodge and protect the chief organs of respiration and circulation,—the lungs and the heart.

THORAX, Corset.

THORE, (Sperm.) Sperm.

THORN APPLE, *Datura stramonium*—t. Red, *Datura Sanguinea*.

THOROUGHSTEM, *Eupatorium perfoliatum*.

THOROUGHWAX, *Eupatorium perfoliatum*.

THOROUGHWORT, *Eupatorium perfoliatum*.

THOROW-WAX, *Bupleurum rotundifolium*.

THORULUS STRAMINEUS, 'a straw pad,' *Fanon*.

THRENEINE, ([G.] *Thräne*, 'a tear,') see Tears.

THREADWORM, *Draunculus*—t. Long, *Trichocephalus*.

THREPSIS, (θρεψις, from *τρέπω*, 'to nourish,') Assimilation, Nutrition.

THREPSOL'OGY, *Threpsolog'ia*, (threpsis, and *λογος*, 'a description,') The doctrine of, or a treatise on, the nutrition of organized bodies.

THRIDACE, (from *thridax*.) *Lactuearum*.

THRIDACINE, (θριδάκη.) *Lactuca*.

THRIDAX, gen. *Thri'dacis*, (θριδάξ, gen. θριδάκος,) *Lactuca*.

THRIFT, AMERICAN, *Statice Caroliniana*—t. *Lavender*, *Statice limonium*—t. *Sea Side*, *Statice Caroliniana*.

THRIX, (θρίξ, gen. *τριξ*.) *Capillus*, *Hair*.

THROAT, (θροτή, θροτα,) *Ju'gulum interstit'ium*, *Jugula'rē*, *Gut'ter*, *Spha'gē*, (Prov.) *Chokes*, *Droat*, *Hause*, *Hose*, (Sc.) *Halse*, *Hawse*, *Witter*. The anterior part of the neck. (F.) *Gorge*. Also, the pharynx.

THROATROOT, *Geum Virginianum*, *Liatris*.

THROATWORT, *Campanula trachelium*.

THROB'ING, *Pulsat'ivus*, *Sphygm'icus*, *Sphygm'o'des*, *Pul'satory*, (F.) *Pulsat'if*. A throbbing pain, (Prov.) *Stang*, (F.) *Douleur pulsative*, is a kind of pain which is, or seems to be, caused or augmented by the pulsation of arteries.

THROE, (from [Sax.] *þropan*, 'to suffer,') Agony, Pains, labour.

THROMBI LACTEI, (pl. of *thrombus*.) Infarctus mammae lacteus.

THROMBOCYST'IS, (thrombus, and *κύστις*, 'a cyst,') The cyst occasionally surrounding a clot of blood—as in encephalic hemorrhage.

THROMBOSIS, (θρομβωσις, from *thrombus*, and *osis*.) Coagulation. Applied especially to coagulation of blood in a blood-vessel.

THROMBUS, (θρομβος, 'a clot,') *Hæmato'ma*. A small, hard, round, bluish tumour, formed by an effusion of blood in the vicinity of a vein which has been opened in the operation of blood-letting. The thrombus is most commonly owing to the opening of the vein and that of the skin not corresponding; to excessive smallness of the cutaneous orifice; or to small, fatty granules, which prevent the discharge of the blood. Com-

presses, dipped in salt water, camphorated spirit, and slight compression, usually disperse it. See Blood.

THROMBUS NEONATORUM, Cephalæmatoma.

THROPILL, Trachea.

THROPPLE, Trachea.

THROTTLE, (dim. of *throat*.) Trachea.

THROW, (from [Sax.] *þropan*, 'to suffer,') Agony, Pains, labour.

THROWORT, *Leonurus cardiaca*.

THRUSH, *Aphthæ*—*t. Milk*, *Aphthæ*—*t. White*, *Aphthæ*.

THRUSTY, *Dipsodes*.

THYRSIS, (*θύρσις*, from *θρῦπω*, 'to break in pieces,') Communion.

THUJA OCCIDENTALIS, *Thuya occidentalis*.

THUMB, ([Sax.] *þuma*.) Pollex.

THUREA, (from *thus*.) *Juniperus lycia*—*t. Virga*, *Juniperus lycia*.

THUS, gen. *Thuris*, (from *þow*, 'to burn perfumes,') see *Pinus abies*—*t. Fœminium*, see *Pinus abies*—*t. Judæorum*, *Croton cascarilla*, *Styrax*, *Thymiana*—*t. Libanotis*, *Juniperus lycia*—*t. Masculam*, *Juniperus lycia*—*t. Verum*, *Juniperus lycia*—*t. Vulgare*, see *Pinus abies*.

THUYA APHYLLA, (*ῥύς*, 'incense,') see *Sandaracæ*.

THU'YA (*θῡα*, from *ῥύς*, 'incense') seu THU'JA OCCIDENTA'LIS, *T. obtusa*, *Cupressus Arbor Vitæ*, *American Arbor Vitæ*, *Tree of Life*. *Ord. Conifera*. The leaves and wood were formerly in high repute as resolvents, sudorifics, and expectorants, and were given in phthisical affections, intermittent fevers, and dropsies. The expressed juice has been applied to condylomata. The arrangement of the medullary matter of the cerebellum, termed *Arbor Vitæ*, is also called *Thuya*.

THYLACIÏTIS, (*θυλακίον*, 'a small sac,' and *itis*.) *Gutta rosea*.

THYM, *Thymus*.

THYMA, gen. *Thymatis*, *Eethyma*, *Thymion*.

THYMASTHMA, gen. *Thymasth'matis*, *Asthma thymicum*.

THYMBRA, (*θύμβρα*.) *Satureia hortensis*—*t. Hispanica*, *Thymus mastichina*.

THYME, *Thymus*—*t. Cat*, *Teucrium Marum*—*t. Common*, *Thymus*—*t. Garden*, *Thymus*—*t. Lemon*, see *Thymus serpyllum*—*t. Mother of*, *Thymus serpyllum*—*t. Virginia*, *Pycnanthemum linifolium*—*t. Wild*, *Thymus serpyllum*.

THYMELEA, (*θύμος*, 'thyme,' and *ελαία*, 'the olive tree,') *Daphne gnidium*—*t. Laureola*, *Daphne laureola*—*t. Mezereum*, *Daphne mezereum*—*t. Monspelica*, *Daphne gnidium*.

THYMELECO'SIS, (*thymus*, and *ἔλκος*, 'an ulcer,') Ulceration of the thymus gland.

THYMIAMA, gen. *Thymia'matis*, (*θυμίαμα*, from *þow*, 'to burn perfumes,') 'a perfume,') *Thus Judæorum*, *Muskwood*, (F.) *Narcaphthe*. A bark from Syria, Cilicia, &c., supposed to be the product of the liquid storax tree. It has an agreeable, balsamic smell; approaching that of liquid storax. See *Lacaphthon*, and *Narcaphthon*.

THYMIAMA, Fumigation, Suffimentum.

THYMIASIOTECHNIA, (*thymiasis*, and *τεχνη*, 'art,') *Thymiatechny*.

THYMIASIS, (*θυμίασις*.) Fumigation.

THYMIATECH'NY, *Thymiatechnia*, *Thymiasiotekhnia*, *Citrea fumigatoria*, (*θυμίαμα*, 'a perfume,' and *τεχνη*, 'art,') The art of employing perfumes in medicine.

THY'MIC, *Thymicus*, (F.) *Thymique*. Having relation to the thymus, as "thymic asthma."

THY'MION, (*θύμιον*.) *Thy'na*, *Thymus*, *Portulacum*, *Portulaca verrucosa rhagodia*, *Verrucosa minor*, (from *þymos*, 'thyme,') A small wart on the skin, resembling a bud of thyme.

THYMIOSIS, (from *thymion*.) *Framboesia*—*t. Indica*, *Framboesia*.

THYMIQUE, *Thymic*.

THYMITES, (*θυμίτης*.) Wine impregnated with thyme.

THYMITIS, (*θύμος*, 'thymus,' and *itis*.) Inflammation of the thymus gland.

THYMOPATHIA, *Psychopathia*, (*θύμος*, 'the mind,' and *πάθος*, 'affection,') A disease of the mind.

THYMOS, (*θύμος*.) Rage, *Thymus*,

THYMOXAL'ME, (*θύμος*, 'thyme,' *οξύς*, 'acid,' and *ἄλς*, 'salt,') A compound of thyme, vinegar, and salt.

THYMUM, *Thymus*.

THY'MUS, (*θύμος*.) *Glandula Thy'mus*, *Corpus Thymicum* seu *Thymicum*, *Glandium*, *Thy'mus gland*, *Corpus incomprehensibile* of Joseph Frank, — *Sweet's bread* of the butcher. An organ, concerned in hæmotosis, which is seated in the upper separation of the anterior mediastinum. The thymus has the appearance of a glandular body. It is oblong; bilobate; soft, and very variable in size and colour. In the fetus, it is very large, and contains in a central cavity — reservoir of the thymus — a milky fluid; but it gradually disappears, and in old age is scarcely discernible. The arteries, called *thy'mic*, are from the inferior thyroid, internal mammary, bronchial, and mediastinal. The veins have the same arrangement. It receives some nervous filaments from the pneumogastric nerves, the phrenic, and the inferior cervical ganglia.

THYMUS, (*θύμος*.) *Thy'mum*, *T. vulgaris* seu *temuifolius* seu *hortensis*, *Common* or *Garden Thyme*, (F.) *Thym*, *T. ordinaire*, *Farigoule*. *Ord. Labiata*. *Sec. Syst. Didynamia Gymnospermia*. This herb has an agreeable, aromatic smell; and a warm, pungent taste. It is reputed to be resolvent, emmenagogue, tonic, and stomachic. It is not much used. The volatile oil, *O'leum Thy'mi*, *Oil of Thyme*, is official in the Pharmacopœia of the United States.

THYMUS, (*θύμος*.) *Psyche*, *Satureia capitata*, *Thymion*—*t. Calaminthus*, *Melissa calamintha*—*t. Capitatus*, *Satureia capitata*—*t. Ciliatus*, *T. mastichina*—*t. Creticus*, *Satureia crepitata*—*t. Hortensis*, *Thymus*—*t. Includens*, *T. serpyllum*.

THYMUS MASTICH'NA, *T. Ciliatus*, *Ma'rum vulgare*, *Clinopodia*, *Mastich'na Gallo'rum*, *Thym'bra Hispan'ica*, *Ja'ca In'dica*, *Common herb Mastich*. A low, shrubby, Spanish plant, used as an errhine. It has a smell like mastich. Its virtues resemble those of the *Marum Syriacum*, but it is said to be less powerful.

THYMUS MULTIFLORUS, *Melissa nepeta*—*t. Nepeta*, *Melissa nepeta*.

THYMUS, RESERVOIR OF THE, see *Thymus*.

THYMUS SERPYLL'UM, *T. includens*, *Herpy'los*, *Herpy'los*, *Serpyll'um*, *Serpyll'um*, *Serpyll'um*, *Gila'rum*, *Serpyll'um vulgare* *serp' m'us*, *Moth'er of Thyme*, *Wild Thyme*, (F.) *Serpolet*. This plant has the same sensible properties as the garden thyme; but has a milder and rather more grateful flavour. *Lem'on Thyme*, *Serpyll'um citra'dum*, is merely a variety of *Thymus Serpyll'um*. It is very pungent; and has a particularly grateful odour, approaching that of lemons.

THYMUS SYLVATICUS, *Clinopodium vulgare*—*t. Sylvestris*, *Satureia capitata*—*t. Tenuifolius*, *Thymus*—*t. Vulgaris*, *Thymus*.

THYREMPHRAXIS, (*thyreo*, and *εμφραξις*, 'obstruction,') *Bronchocele*.

THYREO, *Thyro*, (from *þyeos*, 'a shield,') in composition, the thyroid cartilage or gland.

THYREOÄDENTIS, *Thyreitis*.

THYREOÄRYTENOID, *Thyroärytenoideus*, or *Thyroärytenoides*. That which relates to the thyroid and arytenoid cartilages.

THYREOÄRYTENOID LIG'AMENTS, *Inferior Lig'aments of the Lar'ynx*, *Lips of the Glottis*, *Vocal*

Cords, are two ligaments about two lines broad, formed of elastic and parallel fibres, which are contained in a duplicature of the mucous membrane of the larynx. They extend horizontally from the anterior projection at the base of each arytenoid cartilage to the middle of the angle of union of the thyroid cartilage. They are the essential organs of voice. See Glottis.

THYREOÄRYTENOID MUSCLES, *Thyreöärytenoidei*, are thin muscles, which arise from the middle and inferior part of the posterior surface of the thyroid cartilage; whence they proceed backwards and outwards, to be inserted into the outer part of the base of the arytenoid cartilage. They give the necessary tension to the ligaments of the glottis, in the production of the voice. Santorini has given the name *Thyreöärytenoideus obliquus* to a portion of the *arytenoideus* muscle.

THYREOCELE, (*thyreo*, and *κύη*, 'a tumor,') *Bronchocele*.

THYREOËPIGLOTTIC, *Thyreöëpiglottideus*, *Thyreopigloticus*. Sabatier and Santorini have given this name to the outer portion of the thyroarytenoid muscle; because it passes from the thyroid cartilage to the anterior part of the epiglottis.

THYREOHYOID, *Thyreohyoideus*, *Thyrophyoïdes*, *Thyrohyoid*, *Hyoëthyroïdes*. That which belongs or relates to the thyroid cartilage and os hyoides.

THYREOHYOID, **THYROHYOID**, or **HYOTHYROID MEMBRANE** is a very broad, yellowish, fibrous membrane, thicker at the middle than at the extremities, which is attached, above, to the posterior surface of the body and great cornu of the os hyoides; and, below, to the whole superior edge of the thyroid cartilage. The lateral *thyrohyoid ligaments* may be considered as the margins of the thyrohyoid membrane. They pass between the superior cornua of the thyroid cartilage to the extremities of the greater cornua of the os hyoides. A small cartilaginous or bony nodule—*cartilago triticea*—is sometimes found in each.

THYREOHYOIDEUS or *Hyoëthyroideus* *Muscle* is situate at the anterior and superior part of the neck. It is attached to the oblique line on the anterior surface of the thyroid cartilage; to the inferior edge of the body of the os hyoides, and to the anterior portion of its great cornu. It approximates the thyroid cartilage and os hyoides to each other, by raising the cartilage, and with it the whole larynx; or, it can depress the os hyoides.

THYREOID, *Thyroid*.

THYREOIDEUS, *Thyroideal*.

THYREOÏDIEN, *Thyroideal*.

THYREOÏTIS, *Thyreöidenitis*, *Thyreöiditis*, *Angina thyreöideä*, (*thyreo*, and *itis*.) Inflammation of the thyroid gland.

THYREONCUS, (*thyreo*, and *ὄγκος*, 'a tumor,') *Bronchocele*.

THYREOPHARYNGEUS, *Thyropharyngeus*. **THYREOPHRAXIA**, (*thyreo*, and *φρασσα*, 'to obstruct,') *Bronchocele*.

THYREPIGLOTTICUS, *Thyreöëpiglottic*.

THYRO, in composition, see *Thyreo*.

THYROÄRYTENOID, *Thyreöärytenoid*.

THYROCELE, (*thyro*, and *κύη*, 'a tumour,') *Bronchocele*.

THYROCICOPHARYNGEUS, see *Cricopharyngeal*.

THYROHYOID, *Thyreohyoid*.

THYROPHARYNGOSTAPHYLINUS, *Palatopharyngeus*.

THYROID, *Thyreoid*, *Thyroïdes*, (*Thyros*, from *θύρα*, 'a gate or folding-door,' or from *θυρεός*, 'a shield,' and *εἶδος*, 'form,') That which has the shape of a folding-door.

THYROID ARTERIES, *Thyroideal arteries*—*t. Artery of Neubauer*, see *Thyroidal arteries*.

THYROID CARTILAGE, *Cartilago Scutiformis* seu *Scutalis* seu *Clypealis*, *Coëpertonium*, *C. Peltalis* seu *peltatus*, *Scutum*, *Adami Moræus Os*, is the largest of the cartilages of the larynx, at the anterior part of which it is situate. It is larger transversely than vertically; broader above than below; and seems formed by the junction of the two quadrilateral plates, which produce, by their union, an angle that projects forwards. Its two posterior margins terminate above, and on each side, by an ensiform prolongation, called the *greater cornu of the thyroid cartilage*, *cornu superius seu longum*; and, below, by a less prominent eminence, the *lesser cornu*, *Cornu inferius seu brevè*; which is articulated with the cricoid.

THYROID GLAND, *Coëpus Thyreöideum*, *Glandula Thyreöidea*, (F.) *Gland on Corpus Thyreöide*, is an organ, the uses of which are not known. It covers the anterior and inferior part of the larynx, as well as the first rings of the trachea, and seems to be composed of two lobes, flattened from before to behind, which are united by a transverse prolongation of the same nature as themselves, called the *Isthmus*. The tissue of the thyroid is soft, spongy, and of a brownish colour; but its intimate structure is unknown. It is formed of several distinct lobules; collected in lobes of greater or less size. They are composed of granulations, some of which contain a yellowish or milky fluid. The thyroid gland receives four large arteries, called *Thyroidea*, as well as corresponding veins. Its nerves proceed from the pneumogastric, and from the cervical ganglia. No excretory duct has ever been found in it. Its uses are not known.

THYROIDAL or *Thyroideal*, *Thyreöideus*, *Thyreöideus*, (F.) *Thyroëdien*, *Thyreödien*. That which concerns the thyroid gland or cartilage.

THYROIDAL, or **THYROID ARTERIES** are two on each side. 1. The *superior Thyroideal*, *superior laryngeal*, *superior guttural*, arises from the anterior part of the external carotid, and proceeds to the upper part of the thyroid gland, after having given off a *laryngeal branch* and a *cricothyroid*. 2. The *inferior Thyroideal*, *inferior guttural*, much larger than the last, arises from the upper part of the subclavian. It passes, in a serpentine manner, to the inferior part of the thyroid gland, to which it is distributed, after having given a number of branches to the neighbouring parts, among which is the *cervicalis ascendens*. The inferior thyroideal artery is occasionally replaced by the *Thyroid artery of Neubauer*, *Artery of Neubauer*, which arises directly from the curvature of the aorta.

THYROIDAL VEINS are 1. A *superior thyroideal*, and several *middle thyroideal*, which open into the internal jugular vein. 2. An *inferior thyroideal*, much larger than the preceding, whose branches form—by anastomosing with those of the opposite side, in front of the trachea—a very remarkable venous plexus, which J. Cloquet calls the *infrathyroideal plexus*. The inferior thyroideal veins open,—the left into the corresponding subclavian,—the right, into the *vena cava superior*.

THYROIDIEN, *Thyroideal*.

THYROPHARYNGEUS, *Thyropharyngeus*. Valsalva, Morgagni, Santorini, and Winslow have given this name to the portion of the constrictor pharyngis inferior, which is attached to the thyroid cartilages.

THYROPHARYNGEUS, *Constrictor pharyngis*, see *Cricopharyngeal*.

THYROPHRAXIA, (*thyro*, and *φρασσα*, 'to obstruct,') *Bronchocele*.

THYRSUS, (*Thyrsos*.) *Penis*.

THYSSELINUM, *Peucedanum palustre*—*t. Palustre*, *Peucedanum palustre*.

TIARELLA CORDIFOLIA, (dim. of *tiara*, *tiara*, 'the turban of the Persian sovereigns': from the form of the capsules,) *Coolwort*. A tonic and diuretic.

TIBI-PÉRONÉO-TARSIEN, *Peronæus longus*.

TIB'IA, 'a flute or pipe,' *Che'mē*, *Procnemium*, *Fo'cile mal'jus*, *Arun'do* *crū Can'na ma'jor*, *Can'na ma'jor* seu *domestica eru'ris*, *Shin'bone*. The largest bone of the leg. A long, irregular, and triangular bone, situate on the inner side of the fibula. It has, 1. A *Superior or femoral extremity*, which is very large, rounded, and has two eminences at the sides, called *Tuberosities of the Tib'ia*, or *Tib'ial tuberosities*. The outermost is articulated with the fibula. On its head are two articular, oval, concave surfaces, separated by a process, called *Spine of the tibia*, which are articulated with the condyles of the *Os femoris*. 2. An *inferior or tarsal extremity*, having an articular surface beneath, which joins the astragalus; within, a triangular eminence, which forms the malleolus internus; and, on the outside, a triangular surface, which articulates with the fibula. 3. The body or shaft of the tibia has three faces, separated by three angles, the anterior of which is the most prominent, and is called the *spine* or *crista of the tib'ia* or *tib'ial spine*. It is the *shin*. The tibia is articulated with the femur, fibula, and astragalus. It is developed by three points of ossification, one for the body, and one for each extremity.

TIBIA MINIMA, Fibula.

TIBIAD, see Tibial aspect.

TIBIÆUS, Tibial.

TIB'IAL, *Tib'ialis*, *Tib'æus*, (F.) *Tib'ial*, *Jambier*. (From *tibia*.) That which relates to the tibia or to the leg.

TIBIAL APONEUROSIS, (F.) *Aponévrose Jambière*, surrounds the muscles of the leg. It is continuous, above, with the femoral aponeurosis, and arises, also, from the head of the fibula, and from several fibrous expansions which detach themselves from the tendons of the triceps, sartorius, gracilis, and semitendinosus. Thence it descends around the leg, attaching itself to the whole extent of the anterior and inner edges of the tibia. It sends, below, an expansion, which passes before the tendo Achillis, and is continuous with the anterior annular ligament of the tarsus. It is attached, also, to the sheath of the peronei muscles.

TIBIAL ARTERIES are two in number. 1. *Arteria Tib'ialis Ant'ica*, which arises from the popliteal, before its bifurcation; passes immediately through the opening in the upper part of the interosseous ligament, and reaches the anterior part of the leg; when it turns downwards, and descends obliquely, between the extensor communis digitorum pedis and the tibialis anticus, anterior to the interosseous ligament; glides under the anterior annular ligament of the tarsus, and takes the name *Dorsal'is tar'si*. Amongst the numerous branches, given off by the tibialis antica, may be distinguished—the *Tibialis recurvens*, *A. malleolar'is inter'na*, and *A. malleolar'is exter'na*. 2. *A. Tib'ialis post'ica*, arises from the bifurcation of the popliteal artery; and, slightly tortuous, descends between the two planes of the posterior muscles of the leg; until it arrives under the arch of the calcaneum, where it divides into two branches—the *Plantar arteries*.

TIBIAL ASPECT. An aspect towards the side on which the tibia is situated.—(Barclay.) *Tibiad* is used adverbially by the same writer, to signify, 'towards the tibial aspect.'

TIBIAL NERVES are two in number. 1. The *anterior tibial nerve*, *Prétibio-sus-plantaire*, (Ch.) One of the two branches in which the external popliteal terminates. It accompanies the arteria

tibialis antica. 2. The *posterior tibial nerve*, *Branche tibiale du nerf fémoro-poplitée* (Ch.), is the internal popliteal.

TIBIAL VEINS, ANTERIOR AND POSTERIOR, have the same arrangement as the arteries which they accompany.

TIBI'ALIS ANTICUS, *Catēnæ musculus*, (F.) *Tibio-sus-metatarsien*, *Tibio-sus-tarsien* (Ch.), *Jambier antérieur*. A muscle situate at the anterior part of the leg. Above, it has the shape of a triangular prism; below, it is slender and tendinous. It is attached, above, to the anterior part of the external tuberosity of the tibia; to the superior half of the outer surface of that bone; and to the anterior surface of the interosseous ligament. Below, its tendon terminates at the base of the first cuneiform bone, and at the posterior extremity of the first metatarsal bone. This muscle bends the foot on the leg, and directs its points inwards, at the same time that it raises its inner edge. It can, also, bend the leg on the foot, and prevent it from falling backwards in standing.

TIBIALIS GRACILIS, Plantar muscle.

TIBIALIS POSTICUS, (F.) *Tibio-tarsien*, *Tibio-sus-tarsien* (Ch.), *Jambier postérieur*. This muscle is situate at the posterior and deep part of the leg. It is thicker above than below; and is attached, above, to the posterior surface of the fibula; to the posterior surface of the tibia, and to the interosseous ligament. Its tendon terminates, below, at the tuberosity on the inferior extremity of the os scaphoides. This muscle extends the foot on the leg; at the same time raising its inner edge. It, likewise, extends the leg on the foot.

TIBIO-CALCANIEN, Soleus—*t. Phalangetien*, Flexor longus digitorum pedis profundus perforans—*t. Sous-tarsien*, Tibialis anticus—*t. Sus-metatarsien*, Tibialis anticus—*t. Sus-tarsien*, Tibialis anticus.

TIBIOTARSAL, *Tibiotar'seus*, (F.) *Tibio-tarsienne*. What relates to the tibia and tarsus.

TIBIOTARSAL ARTICULATION is the articulation of the foot with the leg. It is formed by the tibia and the astragalus; and is strengthened by lateral, and by anterior and posterior ligaments.

TIBIO-TARSIEN, Tibialis posticus.

TIC, (from [Sax.] *teogan*, [G.] *ziehen*, 'to draw,') *Twitch'ing*, *Vellic'tion*. A local and habitual convulsive motion of certain muscles; and, especially, of some of those of the face. It is, sometimes, called *Tic convulsif* on *Tic non douloureux*, *Histric'ic Spasm of the Face*, *Spas'mus facia'lis*, to distinguish it from *Tic douloureux* or *neur'al'gia fac'iei*, and has been termed a *par'tial chore'a* or form of *chron'ic chore'a*, *Chore'a fac'iei*.

TIC DOULEUREUX, 'painful tic,' Neuralgia, facial—*t. non douloureux*, see Tic.

TICK, ([F.] *Tique*, [G.] *Zecke*), *Ixodes ricinus*—*t. Wood*, *Ixodes ricinus*.

TICKLE WEED, *Veratrum viride*.

TICKLING, ([L.] *titillare*, [Sax.] *rinklan*, 'to tickle,') (Sc.) *Kit'ling*, *Kit'ling*, (F.) *Chatouillement*. This word, sometimes, means the action of tickling (*titilla'tio*, *titilla'tion*;) at others, the sensation produced by this action (*pruri'tus*). A vivid sensation, which commonly causes laughter, and a state of general spasm that may be dangerous if too long protracted. There are some parts of the body, where we are easier tickled than others; for example, the sole of the feet, and the hypochondriac regions.

TICKSEED SUNFLOWER, *Coreopsis trichosperma*.

TICKWEED, *Hedeoma*—*t. Sunflower*, *Coreopsis trichosperma*.

TICOREA FEBRIFUGA; a Brazilian plant,

Order, Rutaceæ, the bark of which is intensely bitter, astringent and antiperiodic.

TICOREA JASMINIFLORA. A decoction of the leaves is said to be drunk by the Brazilians as a cure for frambœsia.

TIDAL AIR, (from *tide*), see *Respiration*.

TIERCE, ([L.] *tertius*, 'third,') see *Tertian*.

TIGALA, Saccharum.

TIGE ('a stalk') **PITUITAIRE**, Infundibulum of the brain — *t. Sus-ephœnoidale*, Infundibulum of the brain.

TIGLIA *seu* **TIGLII GRANA**, see *Croton Tiglium*.

TIGRETIER, see *Mania, dancing*.

TIKIMMA, *Cacoucia coccinea*.

TILBURY, MIN'ERAL WA'TERS OF. A saline chalybeate at West Tilbury, in Essex, England. It is not much frequented.

TILIA, *T. Europæa* *seu* *grandiflora* *seu* *cordifolia* *seu* *pauciflora* *seu* *plataphylla*, (akin to *pyracæ*, 'the elm tree,') *Philyra*. The *Lime*, *Lind*, or *Linden tree*, (F.) *Tilleul*. *Ord.* Tiliaceæ. The flowers have been supposed to possess anodyne and antispasmodic virtues.

TILIA AMERICANA. *T. glabra* *seu* *Canadensis*, *Linden*, *Linn*, *Bass'wood*, *White'wood*, *Spoon'wood*. A mucilage prepared by macerating the inner bark in cold water, has been applied to burns. It may be made into a cataplasm by thickening it with bran.

TILIA CANADENSIS, *T. Americana* — *t. Glabra*, *T. Americana*.

TILLEUL, *Tilia*.

TILLICHERRY CORTEX, *Nerium antidysentericum*.

TILMA, gen. *Til'matis*, (τιλμα, from τιλλω, 'to pluck,') *Linteum*.

TILMUS, (τιλμος,) *Carphologia*, *Evlusion*.

TILY, OF THE INDIANS, *Viburnum dentatum*.

TIMAC. The name of an East India root, but of what vegetable is not clear. It is said to possess diuretic properties; and hence has been given in dropsies.

TIMBRE (F.), 'the sound of a bell, (τυμπανον, from τυπω, 'to strike' [?]) The quality of sound by which one voice, or one instrument, can be distinguished from another when producing the same note.

TIMIDUS, *Rectus inferior oculi*.

TIN, ([Sax.] *tin*,) *Stannum*, *Cassit'erus*, *Cattit'erus*, *Jupiter*, *Stal'gia*, *La'os*, *Plumb'um al'bum* *seu* *can'didum*, (F.) *Étain*. A solid metal; of a peculiar odour when rubbed; insipid; colour whitish; s. g. 7.291; giving a peculiar tinkling sound when struck. It is used, in medicine, only as a mechanical anthelmintic. Dose, $\mathfrak{z}\mathfrak{j}$ of the *Pulvis Stanni*, *Stannum granulat'um* or *Granulat'um*, in molasses. (This is made by melting tin in an iron vessel over the fire, and, while it is cooling, stirring until it is reduced to a powder, which is passed through a sieve.—Ph. U. S.)

TIN, BUTTER OF, Tin, muriate of — *t. Chloruret* of, Tin, muriate of — *t. Deutohydrochlorate* of, *T. muriate* of.

TINFOIL, *Stannum foliat'um*, *Stanni'olum*, is used for plugging carious teeth, &c.

TIN GLASS, Bismuth.

TIN, GRANULAR, see *Tin* — *t. Muriate* of, superoxygenated, *T. muriate* of.

TIN, MURIATE OF. *Fu'ming li'quor* of *Liba'vius*, *Mu'rias Stanni* *seu* *Oxyd'uli Stanni*, *Stanni chloridum*, *Stannum chlorat'um* *seu* *muriat'icum*, *Salt of Tin*, *Butter*, *Chloride* or *Deutohydrochlorate* or *Superoxygenated Mu'riate* of *Tin*, (F.) *Sel d'Étain*. (This is formed of tin, one part; concentrated muriatic acid, three parts. To be crystallized by the aid of heat.) A violent cathartic. Dose, gr. ij or iij.

TIN, SALT OF, Tin, Muriate of.

TIN, SULPHURET OF, *Aurum musivum*.

TINAGMUS, (τυναγμος,) *Concussion*.

TINASMUS, *Tenesmus*.

TINCTURA, (from *tingere*, *tinctum*, 'to dye,')

Tincture, *Essen'tia*, (F.) *Teinture Alcoolique*, *Alcoolé*, *Alcohol*, *Alcoholature*. The term *tincture* is generally restricted to spirituous solutions of vegetable, animal, and some saline substances. It corresponds, therefore, with the word *Quintessence*, in one of its old significations; and with the *Alcoholat'um* of the *Codex* of Paris. It is not unusual, however, to speak of *a'queous tincture*, *ethe'ral tincture* (*Éthérolature*, Béral), &c. Tinctures are usually made either with pure alcohol or proof spirit. The former are precipitated by water; and, therefore, are seldom used internally; the latter are common additions to infusions, decoctions, &c. They ought not, of course, to be united with any vehicle which can decompose them, or separate any thing from them in a palpable form; unless such decomposition is desired by the prescriber. In making tinctures, the ingredients should be reduced to a coarse powder; and the maceration be made in close vessels, exposed to a heat of about 80°, and frequently shaken. When completed, they must be filtered, and put away for use in close bottles. When the process of displacement is employed, great care must be taken, so that the substances treated may be, as far as possible, exhausted of their soluble principles, and a perfectly clear tincture be obtained. To those who are not familiar with the process, the plan of maceration is preferable.

TINCTURA ABSINTHII COMPOSITA, *Essen'tia absinthii composita* *seu* *amar'a*, *Compound Tincture of worm'wood*. (*Absinth.*, *Artem. pontic.*, *Caryoph.* $\mathfrak{aa}$ $\mathfrak{z}\mathfrak{ss}$. *Sacch.* $\mathfrak{z}\mathfrak{i}\mathfrak{j}$, *Alcohol*. *Oss.* Macerate for fifteen days. *Ph. P.*) Tonic, stomachic, vermifuge, and carminative. Dose, $\mathfrak{f}\mathfrak{z}\mathfrak{i}\mathfrak{j}$ to $\mathfrak{f}\mathfrak{z}\mathfrak{ss}$.

TINCTURA ACACIE CATECHU, *T. catechu* — *t. Acetatis ferri cum alcohole*, see *T. ferri acetatis* — *t. Acidi sulphurici*, *Sulphuricum acidum aromaticum*.

TINCTURA ACONITI, (Ph. U. S. 1842.) *Tinctura Aconiti foliorum*, (Ph. U. S. 1851.) *T. A. Folii*, (Ph. U. S. 1863.) *Tincture of Aconite*, (*Aconit. folior.* $\mathfrak{z}\mathfrak{i}\mathfrak{j}$, *Alcohol. dilut.* *Oij.* Macerate for 14 days, express, and filter through paper. It may also be made by displacement. Ph. U. S.) Dose, grt. xx or xxx. It is rubbed on the seat of neuralgia.

TINCTURA ACONITI RADICIS, *Tincture of Aconite root*, (*Aconit. radices contus.* $\mathfrak{f}\mathfrak{ij}$, *Alcohol. Oij.* Macerate for 14 days; express and filter. It may also be prepared by displacement. Ph. U. S.) Uses same as the last.

TINCTURA ÆTHEREÆ CAMPHORATÆ, *Solutio camphoræ æthereæ*, *Li'quor neri'us Bang'ii*, *Spiritus sulphurico-æthereus camphorat'us*, *Naph'tha vitri'oli camphorat'a*. (*Camphor.* p. i. *Æther sulphuric.* p. ii.) Stimulant in atonic diseases, arthritic cardialgia, and spasm. Dose, 20 to 30 drops in white wine.

TINCTURA ÆTHEREÆ FERRI, *Alcohol (seu Tinctura) sulphurico-æthereus ferri* — *t. Alcoholica Chinæ composita*, *T. cinchonæ composita* — *t. Alcoholica corticum aurantiorum Whyttii*, *Tinctura cinchonæ amara* — *t. Alexipharmaca Huxhami*, *Tinctura cinchonæ composita*.

TINCTURA AL'OEËS, *T. Al'oeis Sacotori'næ*, *Essen'tia Aloes*, *Al'cohol cum Al'oe perfoliat'a*, *Tincture of Aloes*. (*Aloes*, pulv. $\mathfrak{z}\mathfrak{j}$. *Ext. glycyrrh.* $\mathfrak{z}\mathfrak{i}\mathfrak{j}$, *Aque Oiss.* *Alcohol. Oss.* Macerate for 14 days, and filter. Ph. U. S.) Properties, those of aloes. Dose, $\mathfrak{f}\mathfrak{z}\mathfrak{ss}$ to $\mathfrak{f}\mathfrak{z}\mathfrak{i}\mathfrak{j}$.

TINCTURA AL'OEËS ÆTHEREÆ, *T. Aloes vitriolata*, *Æthe'ral Tincture of Al'oes*. (*Aloes Sacot.*, *Myrrh.* $\mathfrak{aa}$ $\mathfrak{z}\mathfrak{i}\mathfrak{j}$, *Croc.* $\mathfrak{z}\mathfrak{i}\mathfrak{j}$. *Sp. ætheris sulph.* *Oj.* Digest the myrrh in the ether for 4 days; then

add the aloes and saffron. *Ph. E.*) Stimulant and cathartic. Dose, fʒi to fʒij.

TINCTURA AL'OE'S COMPOSITA, *Eliz'ir Proprieta'tis seu Proprieta'tis Paracel'si seu aperiti'um seu Al'oe's, Tinctura Al'oe's cum My'r'hā seu Al'oe's et My'r'hā* (*Ph. U. S.*) seu *Al'oe's et My'r'hā crocata, Eliz'ir of long-life, Compound Tincture of Al'oes*, (*F.*) *Eliz'ir de longue vie, Baume de vie de Lelierre*, (*Aloës*, in pulv., ʒijj. *Croc.*, ʒi. *Tinct. Myrrhæ*, Oij. Macerate for 14 days, and filter.) Purgative and stomachic. Dose, fʒi to fʒij.

Baur'haari's Eliz'ir consisted of aloes, myrrh, saffron, and tartrate of potassa, digested in alcohol and water.

Rad'cliff's Eliz'ir is formed of *Aloes socotr.* ʒvj. *Cort. cinam.*, *Rad. zolatr.* ʒā ʒss. *Rad. rhei* ʒj. *coccinell.* ʒss. *Syrup. rhummi* ʒij. *Sp. tenuior.* Oj. *Aqua purā* fʒv.

TINCTURA ALOES ET MYRRHÆ, *T. aloes composita* — t. Aloes et myrrhæ crocata, *T. A. composita* — t. Aloes vitriolata, *T. A. aetherea* — t. Amara, *T. gentianæ composita* — t. Ammoniac composita, *Spiritus ammoniac succinatus* — t. Amomi repentis, *T. cardamomi* — t. Amomi zingiberis, *T. zingiberis*.

TINCTURA ANGUSTURÆ, *T. Cuspa'ria, Tincture of Angustura*, (*Cort. cuspar.* in pulv. crass. redact. ʒij. *Sp. vin. ten.* Oij. Digest. *Ph. D.*) Dose, fʒi to fʒj.

TINCTURA ANTIFEBRILIS WARBURGI, see *Bebeeru*.

TINCTURA ANTIODONTALGICA, Tooth'ach Tincture, (*Tomlin.* ʒj. *Mustich.* gr. v. *Ether.* ʒij. *M.*) To be applied on cotton wool to the tooth previously dried.

TINCTURA ARISTOLOCHIE SERPENTARIE, *T. serpentariae*.

TINCTURA AR'NICÆ, *Tincture of Ar'nica*, (*Arnic.* ʒvj. *Alcohol.* Oiss. *Aq. Oss.* *Alcohol. dilut.* q. s. Mix the alcohol and water, and bruise the arnica, slightly moistened with the mixture, thoroughly in a mortar; pack it firmly in a percolator and pour on first the remainder of the mixture, and then sufficient Diluted Alcohol to measure two pints. *Ph. U. S.* 1863.) Dose, gtt. v. to x. Applied, diluted with an equal proportion of water, by means of lint, as an excitant to bruises, and in cases of neuralgia. *Tinctura Arnicæ* of the British Pharmacopœia (1864), made by percolation, from the root — the U. S. preparation, being from the flowers — is much weaker.

TINCTURA AROMATICA, *T. cinnamomi composita* — t. *Aromaticæ ammoniata, Spiritus ammoniac aromaticus*.

TINCTURA ASSAFETIDÆ, *T. Fer'ula Assafetida, T. fo'tida seu assafetida, Tincture of Assafetida*, (*Assafetid.* ʒiv. *Alcohol.* Oij. Macerate for 14 days and filter. *Ph. D.*) Dose, gtt. xv to fʒj.

TINCTURA ASSAFETIDÆ AMMONIATA, *Spiritus Ammoniac foetidus*.

TINCTURA AURANTII, *T. Car'ticis Aurantii, Tincture of Or'ange Peel*, (*Aurant. cort. recent.* ʒijj. *Sp. rect.* Oij. Digest for 14 days.) Stomachic. Used as an adjunct to stomachic draughts. Dose, fʒss to fʒij.

TINCTURA AUREA, *Alcohol (seu tinctura) sulphuricoethereus ferri*.

TINCTURA BELLADONNÆ, *Tincture of Belladonna*, (*Belladonna.* ʒiv. *Alcohol. dilut.* Oij. Macerate for 14 days; express, and filter through paper. It may, also, be prepared by displacement. *Ph. U. S.* Dose, gtt. xv to xxx.

Tinctura Belladonnæ of the British Pharmacopœia (1864), made by percolation, is half the strength of the above.

TINCTURA BENZOINI COMPOSITA, *T. Ben'zoës composita, Balsamum Catholiceum seu Per'sicum seu trauaticeum, Eliz'ir trauaticum, Balsamum Commendato'ris, Compound Tincture of Ben'zoïn,*

Fri'ar's or Ver'vain's Bal'sam, Wade's or Jesuit's drops, Commander's or Wound Balsam, Balsam for Cuts, (*Benzoin.* ʒijj. *Syrac. purif.* ʒij. *Balsam. tolu.* ʒj. *Aloes*, in pulv. ʒss. *Alcohol.* Oij. Macerate for 14 days and filter. *Ph. U. S.*) Stimulant. Used chiefly to wounds and ulcers.

The basis of *Tur'lington's Bal'sam of Life, Tur'lington's Bal'sam*, is the Compound Tincture of Benzoin. The following form is recommended by a committee of the Philadelphia College of Pharmacy: — *Alcohol.* Ovij. *Benzoin.* ʒxij. *Syrac. liquid.* ʒiv. *Aloes Socotr.* ʒj. *Balsam. Peruvian.* ʒij. *Myrrhæ.* ʒj. *Rad. Angelic.* ʒss. *Balsam. Tolu.*, *Ext. Glycyrrh.* ʒā ʒiv. Digest for 10 days, and strain.

The *Es'sence of Colts'foot* consists of equal parts of the Compound Tincture of Benzoin and Balsam of Tolu, to which is added double the quantity of Rectified Spirit of Wine. It is sold as a pectoral.

TINCTURA BUCHU, *T. Buchu, T. Bucco, Tincture of Buchu*, (*Buch.* ʒv. *Alcohol dilut.* Oij. *Imp. meas.*) Digest for seven days and filter. It may be made, also, by percolation. *Ph. E.*) Dose, fʒi to fʒiv. See *Diosma*.

TINCTURA BUCKU, *Tinctura Buchu*.

TINCTURA COLUMBÆ, (*Ph. U. S.* 1863.) *T. Colum'ba seu Colum'ba* (*Ph. U. S.* 1851). *Tincture of Colum'ba, Colum'ba Rit'tera*, (*Columb. rad. cont.* ʒiv. *Alcohol. dilut.* Oij. Macerate for 14 days; express, and filter. *Ph. U. S.*) Dose, fʒss to fʒiv. It may, also, be made by displacement.

TINCTURA CAMPHORÆ, *Spiritus camphoræ*.

TINCTURA CAMPHORÆ COMPOSITA, *T. O'pii Camphora'ta* (*Ph. U. S.*), *T. Cam'phoræ cum o'pio, Compound Tincture of Camphorated Tincture of O'pium, Eliz'ir Paregoriceum, Paregoric Eliz'ir* (*O'pii*, in pulv., *Acid. Benzoi.* ʒā ʒj. *Ol. anisi.* fʒj. *Mellis despumat.* ʒij. *Camphor.* ʒij. *Alcohol. dilut.* Oij. Macerate for 14 days, and filter through paper. *Ph. U. S.*) One fluidounce contains nearly gr. ij of opium. Anodyne. Dose, fʒi to fʒijj.

Squire's Eliz'ir, an empirical carminative preparation, is thus made, according to one formula. (*O'pii.* ʒiv. *Camphor.* ʒj. *Coccinell.* ʒj. *Fenicul. dulc.* ʒij. *Tinct. serpent.* Oj. *Sp. anisi.* cong. ij. *Aquo.* Oij. *Auri musci.* ʒvj.

TINCTURA CAN'NABIS INDICÆ, *Tincture of Indian Hemp*, (*Ext. Cannab. purif.* ʒvj. *Alcohol.* Oj. Dissolve. *Ph. U. S.*) Dose, about 40 drops.

TINCTURA CANTHARIDIS, *T. C. Vesicato'ris, T. Ly'tta seu M'la's vesicato'ris, Tincture of Cantharides, Tincture of Blistering Flies, T. of Span'ish Flies*, (*F.*) *Tincture rubefaciens*, (*Cantharid.* pulv. ʒj. *Alcohol. dilut.* Oij. Macerate for 14 days; express, and filter. It may also, be prepared by displacement. *Ph. U. S.*) Stimulant and diuretic, but not often used internally. Externally, stimulant. Dose, gtt. x—xx.

Matthews's Injec'tion, a once celebrated nostrum for fistula in ano, consisted wholly of a dilute Tincture of Cantharides.

TINCTURA CAP'SICI, *Tincture of Cap'sicum or Cayenne' Pop'per*, (*Capsic.* ʒj. *Alcohol. dil.* Oij. Macerate for 14 days, and filter. It may, also, be prepared by displacement. *Ph. U. S.*) Stimulant. Dose, fʒss to fʒj.

TINCTURA CARDAMOMI, *T. Amo'mi repentis, Tincture of Cardamoms*, (*Cardam.* cont. ʒiv. *Alcohol. dilut.* Oij. Macerate for 14 days, strain, and filter. It may, also, be made by displacement. *Ph. U. S.*) Use, the same as the seeds. Dose, fʒi to fʒiv.

Solomon's Balm of Gil'ead, an empirical preparation, seems to be an aromatic tincture, — of which Cardamoms form a leading ingredient, — made with brandy. Some affirm that it contains cantharides.

TINCTURA CARDAMOMI COMPOSITA, *Tinctura Stomach'ica, Compound Tincture of Cardamoms,*

Stomach Tincture. (Cassianum, cont. 5vj. Card. cont. 5j. Cloves, cont. 5v. One penny druggist's shilling. 3s. Card. cont. 5j. Alcohol. 5lb. Oils. Moderate for 14 days; express and filter. Ph. U. S. Prepared, also, by displacement.) Stomachic and carminative. Dose, (5j) to (5ss).

TINCTURA CASSIARINÆ SYLVÆ. *Alcoholica Cassiarinæ seu Salicis, Cassiæ albae Eliæris of Sylicis.* (Rad. aspicis 4 p. Rad. Imperatoris, Galbani. 25 6 p. Fol. menthae, Melissa, Ruta, Basilic. 25 48 p. Baccas. Laur. noli. 12 p. Sem. aspicis, Ligust. lavat. Anis. 25 18 p. Zingib. Nuc. maculat. 25 8 p. Cloves, 12 p. Caryoph. Card. 1000. 55 4 p. Add Alcohol. 1000 p. Macerate and distil. 1000 p. Ph. P.) A warm stomachic, carminative, &c. Dose, (5ss) to (5ss).

TINCTURA CASIABILLÆ. *T. Cassiæ Eliæris.* *Alcoholica Cassiæ seu Cassiæ Cinnamomifoliae, Tincture of Cassia-bill.* (Cinnam. cont. 5iv. Sp. ten. 6j. Ph. L.) Dose, (5j) to (5v).

TINCTURA CASTORÆ. *T. Castoreæ Rostri seu Castoreæ Canadensis, Essentia Castoreæ, Alcohol castoreæ, Tincture of Castor.* (Castor. cont. 5ij. Alcohol. 6j. Macerate for 7 days; express, and filter. Ph. U. S.) Tonic and antispasmodic. Dose, gr. xx to (5j), or more.

TINCTURA CASTORÆ COMPOSITA. *T. C. communis, Castoreæ Tincture of Castor.* (Cast. 5j. Asafoetida. 5ss. Alcohol. ammoniac. 6j. Ph. E.) Antispasmodic. Dose, (5j) to (5iv). Called, also, *Elixir Asafetide*. *Tinctura castoreæ factitia seu fictitia ammoniac.*

TINCTURA CATÆCHÆ. *Tinctura Japonica seu Matico Catæchæ seu Asiatica Catæchæ, Tincture of Catæch.* (Catæch. 5vj. Cloves, contus. 5j. Alcohol. dilut. 6j. Macerate for 14 days; express and filter. Ph. U. S. Prepared also by displacement.) Astringent. Dose, (5j) to (5ij).

The *Concentrated Solution of Catæch*, sold for cleaning the teeth, seems to be little more than the tincture of catæch.

TINCTURA CHIRETÆ. *T. Chiretæ* (Ph. B. 1844). *Tincture of Chiretæ.* (Chiret. cont. 3v. (steird.) Alcohol. dilut. 6j. (Imp. meas.) Macerate for 14 days. Ph. B.) Dose, (5j), or (5ss), as a tonic.

TINCTURA CISTÆ. *T. cistæ maculati.*

TINCTURA CINCHONÆ. *T. Corticis Peruviani Simplicis seu Cort. Peruviani, Tincture of Cinchonæ or of Bark.* (Cinchon. flou. in pulv. 5vj. Alcohol. dilut. 6j. Macerate for 14 days; express, and filter through paper. It may, also, be made by displacement. Ph. U. S.) Dose, (5j) to (5v), or more.

TINCTURA CINCHONÆ ETHERÆA COMPOSITA. *T. Kina Kina ætheræa composita, Elixir seu Astringens Iuxta de Chamber, Chamber's Astringent Elixir.* (Cinchon. offic. 5j. Cassia-cinn. 5ss. Cloves. 5j. Card. 5ss. Sassa. alb. 5xxxvj. Put these bruised into a mattress, and add Vin. alb. Hygroma. vel Vin. mentac. Alcohol. 55 6j. Macerate for two days, and add Sulphuric ether. (5ss. Ph. P.) Tonic, stimulant, and antiseptic. Dose, (5ss) to (5j).

TINCTURA CINCHONÆ AMARÆ seu alcoholica corticis peruviani simplicis Wigori seu Kibbæ canadensis, Elixir antispasmodicæ seu rubicæ Wigori seu Wigori seu cinchonæ spirit. camph. *Essentia Corticis Peruviani Composita.* (Cinch. p. vjij. Gentian. Rad. Card. Aurant. p. ij. Alcohol. 1000) 90 p. Digest. Dose, one or two teaspoonfuls.

TINCTURA CINCHONÆ AMMONIATA. *T. Corticis Peruviani rubicæ, Ammoniacæ or Valerianæ Tincture of Bark.* (Cinchon. lavat. cont. cont. 5iv. Sp. ammon. aromat. 6j. Ph. L.) In dyspepsia, combined with acidity and languor.

TINCTURA CINCHONÆ COMPOSITA. *T. Corticis Peruviani Composita seu Simplici Iuxta de Barki seu Abstrahat' mæ Hæsh'ni seu Alcoholica Ch'na Composita, Essentia Ch'na seu*

Corticis Peruviani Abstrahat' mæ Hæsh'ni seu Simplici Hæsh'ni, Concentrated Tincture of Cinchonæ or Bark, Hæsh'ni's Tincture of Bark. (Cinchon. rub. pulv. 5v. Aurant. amar. cont. cont. 5ij. Sassa. cont. 5j. Card. Sassa. 55 5j. Alcohol. dilut. 15xx. Macerate for 14 days; express, and filter. Ph. U. S.) It may, also, be made by displacement. More grateful than the simple tincture. Dose, (5j) to (5ij).

TINCTURA CINNAMOMI. *T. Lauri Cinnamomifolia, Essentia Cinnamomifolia, Tincture of Cinnamon.* (Cinnam. cont. 5ij. Alcohol. dilut. 6j. Macerate for 14 days; express, and filter. It may, also, be prepared by displacement. Ph. U. S.) Stomachic. Dose, (5j) to (5j).

TINCTURA CINNAMOMI COMPOSITA. *T. seu Essentia Amomum, Alcoholica Amomum Composita, Concentrated Tincture of Cinnamon.* (Fol. Eux de Borneo in d'Amomum. (Cinnam. cont. 5j. Cardam. cont. 5ss. Zingib. cont. 5j. Alcohol. dilut. 6j. Macerate for 14 days; express, and filter. Ph. U. S.) It may, also, be prepared by displacement.) Use, same as the last. Dose, (5j) to (5ij).

TINCTURA CINCHONÆ CACTI. *T. Cacti, Tincture of Cacti wood.* (Cacti pulv. 5ij (crusid.) Alcohol. dilut. 6j. (Imp. meas.) Ph. B.) Chiefly a coloring ingredient. At times, given in nervous affections. Dose, gr. xij to (5j).

TINCTURA DE COCHLEARIS. *Alcoholica de Cochlearis seu antiscorbutica.* (Pul. cochlear. recent. 2000 p. Rad. ascoruticæ. 200 p. Alcohol. (20° to 32° Beaumé) 1000 p. Well off 2000 p. Ph. P.) Antiscorbutic. Dose, (5j) to (5j).

TINCTURA COLCHICÆ. (Ph. U. S. 1842.) *T. C. Semina.* (Ph. U. S. 1851). *Tincture of Colchicum Seed.* (Colchic. sem. cont. 5iv. Alcohol. dilut. 6j. Macerate for 14 days; express, and filter. It may, also, be prepared by displacement. Ph. U. S.) Dose, ten to sixty drops.

TINCTURA COLOCYTHÆ COMPOSITA. *Spiritus colocythæ ammoniacus—t. Colocythæ seminis.* *T. colocythæ—t. Columba.* *T. Columba.*

TINCTURA CORYL. *T. C. Maculati.* *T. Cistæ, Tincture of Hazelnut.* (Cist. fol. 5iv. Alcohol. dil. 6j. Macerate for 14 days; express, and filter. It may also be made by displacement. Ph. U. S.) Use—the same as that of the leaves. Dose Max to (5j). A *Tincture of Hazelnut Fruit*. *Tinctura Corticis fructus*, formed of *Hazelnut fruit* or seeds bruised, 5ss, to *Pung' quic*, 6j, made by percolation, is official in the British Pharmacopæia &c 1864. Dose, gr. xxi.

TINCTURA CUSCUMACULATI. *T. cusci—t. Cuscutæ Jalapæ.* *T. Jalapæ—t. Corticis aurantii.* *T. aurantii—t. Corticis Peruviani composita.* *T. cinchonæ comp.—t. Corticis Peruviani simplicis.* *T. cinchonæ—t. Corticis Peruviani rubicæ.* *T. cinchonæ ammoniacæ.*

TINCTURA CUSCUMACULATI. *T. Cusci, Tincture of Saffron.* (Cusci. cont. 5j. Alcohol. dilut. 5xxv. Ph. E.) Slightly stomachic (?). Dose, (5j) to (5j).

TINCTURA DE CUSCUMACULATI. *Elixirium de Cusci, Alcoholatum de Cusci Compositum, Concentrated Tincture of Saffron, Elixir of Saffron.* (Aloe Sassa. 120 p. Myrr. 64 p. Card. 12 p. Cloves. Nuc. maculat. Caryoph. 25 16 p. Alcohol. 1000 p. Ap. flou. current. 2000 p. Digest for two days; distil 4000 p. and add Symp. of capillaire, 1000 p. Ph. P.) Aromatic, carminative, stomachic. It is also called *Elixir de cardus* & *stomachicum*, and *E. aurantii* & *cinchonæ*.

TINCTURA CUSCUMACULATI. *Tincture of Cusci.* Under this name Mr. E. Wilson recommends "a most valuable cutaneous stimulant," made by adding four fluidounces of Alcohol to one ounce of the bruised seeds of *Desmodium*.

TINCTURA CUSCUMACULATI ELEUTHERIE. *T. cusci-rubæ.*

TINCTURA CUBEÆ, *Tincture of Cubebs*. (Cubebs. cont. $\frac{3}{4}$ iv, Alcohol. dilut. Oij. Macerate for 14 days, express, and filter. It may, also, be prepared by displacement. Ph. U. S.) Dose, one or two fluidrachms.

TINCTURA CUSPARIÆ, *Tinctura Augusturæ*.

TINCTURA DIGITALIS, *T. Digitalis purpurea*, *Tincture of Digitalis*. (Digital. $\frac{3}{4}$ iv, Alcohol. dilut. Oij. Macerate for 14 days, express, and filter. It may, also, be prepared by displacement. Ph. U. S.) Use, same as that of the leaves. Dose, gtt. x.

TINCTURA ERGOTÆ, *Tincture of Ergot*. (Ergot. in pulv. crass. $\frac{3}{4}$ vij (avoid.), Sp. tenuior. Oij. (Imp. meas.) Macerate for 14 days, express, and filter. Ph. D.) Dose, f $\frac{3}{4}$ ij to f $\frac{3}{4}$ ij.

The formula for the Tincture of the British Pharmacopœia (1864), made by percolation, is *Ergot*, $\frac{3}{4}$ iv, to *Proof spirit*, Oj. Dose, $\mathcal{M}$ xv to xxx.

TINCTURA ERGOTÆ ÆTHEREÆ, *Ethereal Tincture of Ergot*. (Ergot. contus. $\frac{3}{4}$ xv, Æther. Oij. (Imp. meas.) Macerate for 7 days, express, and filter. Ph. L.) Dose, 60 to 90 minims.

TINCTURA FERRIFUGA DOCTORIS HUXHAM, *T. cinchonæ composita*.

TINCTURA FERRI ACETATIS, *Tincture of Acetate of Iron*. (Potass. acetat. $\frac{3}{4}$ ij, Sulph. ferri, $\frac{3}{4}$ j, Sp. rectif. Oij. Rub the acetate and sulphate into a soft mass. Dry and rub with the spirit. Digest for 7 days, and pour off. Ph. D.) Tonic and astringent. Dose, gtt. xx to f $\frac{3}{4}$ j. The Dublin college directs a *Tinctura Acetatis Ferri cum Alcoholæ*, which differs from the other, in having half the quantity, only, of the Acetate of Potassa.

TINCTURA FERRI AMMONIATÆ, *T. Ferri Ammoniacalis seu Florum martialis seu Martis Myrsichiti*, *Tincture of Ammoniated Iron*. (Ferri ammon. $\frac{3}{4}$ iv, Sp. ten. Oj. Ph. L.) Use—same as that of the ammoniated iron. Dose, f $\frac{3}{4}$ ss to f $\frac{3}{4}$ ij.

TINCTURA FERRI AMMONIOCHLORIDI, *Tincture of Ammoniochloride of Iron*. (Ferri ammoniochloridi. $\frac{3}{4}$ iv, Alcohol. dilut., Aquæ destillat. $\frac{3}{4}$ ij (Imp. meas.) Dissolve and filter. Ph. L.) Dose, gtt. x to xx.

TINCTURA FERRI CHLORIDI, *T. ferri muriatis*.

TINCTURA FERRI MURIATIS, *T. Ferri chloridi*, (Ph. U. S.) seu *sesquichloridi seu Martis in Spiritu Sælis seu Martis cum Spiritu Sælis seu Ferri Muriati*, *Liquor Ferri Muriatis*, *Tincture of Muriate of Iron*, *Tincture of Steel*, *Alcohol ferratus*. (Ferri subcarb. $\frac{3}{4}$ ss, Acid. muriat. Oj, Alcohol. Oij. Add the acid to the subcarbonate in a glass vessel, and shake during three days. Pour off and add the spirit. Another form is given in the Ph. U. S., 1863, in which iron, in the form of wire, is used in place of the subcarbonate. S. g. 0990.) Tonic and styptic. It has been given, also, with advantage in erysipelas and scarlatina in the dose of $\mathcal{M}$ x to $\mathcal{M}$ xx every two or three hours.

TINCTURA FERRI PERCHLORIDI, *Tincture of Perchloride of Iron*, of the British Pharmacopœia, is made of *Solution of perchloride of iron* f $\frac{3}{4}$ iv, and *rectified spirit* f $\frac{3}{4}$ xv. S. g. 0992.

Bestucheff's Tincture or *Kla'proth's Tincture* is an ethereal tincture of chloride of iron. It is composed of dry perchloride of iron, one part; *Hoffmann's anodyne liquor*, seven parts.

The *Antivenereal Drops*, at one time so celebrated at Amsterdam, were examined by Scheele, and found to be a tincture of chloride of iron and corrosive sublimate.

TINCTURA FERRI SESQUICHLORIDI, *T. Ferri muriatis*—*t. Ferulæ Assafoetidæ*, *T. Assafoetidæ*—*t. Florum martialium*, *T. ferri ammoniata*—*t. Fœtida*, *T. assafoetidæ*—*t. Fœtida ammoniata*, *T. castorei composita*—*t. Fuliginis*, see *Fuligo*.

TINCTURA GALBANI, *Tincture of Galbanum*. (Galb. $\frac{3}{4}$ ij, Sp. tenuior. Oij. Ph. D.) Stimulant and antispasmodic. Dose, f $\frac{3}{4}$ j to f $\frac{3}{4}$ ij.

TINCTURA GALLARUM, *T. Gallæ*. (Ph. U. S.) *Tincture of Galls*. (Gall. contus. $\frac{3}{4}$ iv, Alcohol. dilut. Oij. Macerate for 14 days, express, and filter. It may also be prepared by displacement. Ph. U. S.) Astringent. Dose, f $\frac{3}{4}$ j to f $\frac{3}{4}$ ij.

TINCTURA GENTIANÆ COMPOSITA, *Tinct. Amaræ, Eliæir Stomachicæ*, *Compound Tincture of Gentian*, *Spirit'it Bitters*. (Gent. concis. $\frac{3}{4}$ ij, Aurant. cort. $\frac{3}{4}$ j, Cardam. cont. $\frac{3}{4}$ ss, Alcohol. dilut. Oij. Macerate for 14 days, express, and filter. It may likewise be prepared by displacement. Ph. U. S.) Tonic and stomachic. Dose, f $\frac{3}{4}$ j to f $\frac{3}{4}$ ij.

Brodm's Nervous Cordial consists of the *Tinctures of Gentian*, *Columba*, *Cardamom*, and *Bark*, with the *Compound Spirit of Lavender* and *Wine of Iron*.

Stoughton's Eliæir is a compound tincture of gentian. The *Elixirium Docto'ris Stoughton* seu *Tinctura Amara* of the Parisian Codex is prepared as follows:—*Absinth. summit.* exsiccât., *Teneri chamædrys*, *Rad. gentian.*, *Cort. aurant.*, $\frac{3}{4}$ ss, *Cascarilla* $\frac{3}{4}$ j, *Rhei* $\frac{3}{4}$ ss, *Aloes socotr.* $\frac{3}{4}$ j, Alcohol. Oij. Digest. Dose, f $\frac{3}{4}$ j to f $\frac{3}{4}$ ij.

The *Elixir Viceratæ Hoffmanni* does not differ essentially from these preparations.

TINCTURA GUAIACI, *T. Guaiacæ Officinæ*, *Tincture of Guaiacum*. (Guaiac. $\frac{3}{4}$ vj, Alcohol. Oij. Macerate for 14 days, and filter. Ph. U. S.) Stimulant and sudorific. Used especially in rheumatic and arthritic cases. Nearly the same as *Hill's Essence of Bardana* or *Burdock*.

TINCTURA GUAIACI AMMONIATÆ, *T. Guaiacæ Volatilis*, *T. Guaiacæ, Eliæir Guaiacæ Volatilis*, *Alcohol cum Guaiaco officina'le ammoniatum*, *A. ammoniæ et Guaiacæ*, *Ammoniated Tincture of Guaiacum*, *Volatile Tincture of Guaiacum*. (Guaiac. pulv. $\frac{3}{4}$ vj, Sp. ammon. arom. Oij. Macerate for 14 days, and filter. Ph. U. S.) Properties same as the last. Dose, f $\frac{3}{4}$ j to f $\frac{3}{4}$ ij.

Hatfield's Tincture—a nostrum—consists of *Guaiacum* and *Soap*, $\frac{3}{4}$ ij, *rectified spirit*, Oiss.

TINCTURA HELLEBORI, *T. Hellebori Nigri*, *T. Melampodii*, *Tincture of Black Hellebore*. (Helleb. nigr. cont. $\frac{3}{4}$ iv, Alcohol. dilut. Oij. Macerate for 14 days, express, and filter. It may also be prepared by displacement. Ph. U. S.) Dose, f $\frac{3}{4}$ ss to f $\frac{3}{4}$ j.

TINCTURA HIERA, *Vinum aloes*.

TINCTURA HUMULI, *T. Lupuli*, *Tincture of Hops*. (Humuli $\frac{3}{4}$ v, Alcohol. dilut. Oij. Prepared also by displacement.) Tonic and sedative. Dose, f $\frac{3}{4}$ ss to f $\frac{3}{4}$ ij.

TINCTURA HYOSCYAMI, *T. Hyoscyami nigri*, *Tincture of Henbane*. (Hyoscyam. fol. $\frac{3}{4}$ iv, Alcohol. dilut. Oij. Macerate for 14 days, express, and filter. It may also be prepared by displacement. Ph. U. S.) Narcotic, anodyne. Dose, gtt. xx to f $\frac{3}{4}$ j.

TINCTURA IODINI (Ph. U. S., 1842), *T. Iodinæ* (Ph. U. S. 1851), *Alcohol Iodii*, *Tincture of Iodine*. (Iodin. $\frac{3}{4}$ j, Alcohol. Oj. Dissolve. Ph. U. S.) The tincture spoils by keeping, or at least, deposits the iodine. It has been much used in goitre, &c. Dose, ten drops three times a day.

Tinctura Iodi (Ph. B., 1864), is made of *Iodine* $\frac{3}{4}$ ss, *Iodide of potassium* $\frac{3}{4}$ ij, *Rectified spirit* Oj. Dose, $\mathcal{M}$ x to xxx.

An ethereal and a chloroform tincture of iodine have been used, to affect the system with iodine by inhalation.

TINCTURA IODINI COMPOSITA (Ph. U. S., 1842), *T. Iodinæ Composita* (Ph. U. S., 1851), *Compound Tincture of Iodine*. (Iodin. $\frac{3}{4}$ ss, Potassii iodid. $\frac{3}{4}$ j, Alcohol. Oj. Dissolve. Ph. U. S.) Dose, 10 to 30 drops.

TINCTURA JALAPÆ, *T. Jalapæ* (Ph. U. S.) seu *Convolvuli Jalapæ*, *Voligæ*, *Tincture of Jalap*. (Jalap. pulv. $\frac{3}{4}$ vi, Alcohol. dilut. Oij. Macerate

for 14 days, express, and filter. It may also be made by displacement. *Ph. U. S.*) Cathartic. Dose, f3j to f3iv.

TINCTURA JALAPII COMPOSITA, *Eliz'ir Jalap'æ Compos'itum* seu pur'gans seu anthelminticum. *Succo'rum, Essen'tia cathol'ica pur'gans Roth'ii, (F.) Eau de vie Allemande. (Jalap. Ibs, Scammon. 3j, Alcohol. (229) Ovj.* Macerate for eight days. *Ph. P.*) Dose, 60 to 100 drops.

TINCTURA JAPONICA, Tinctura catechu—t. Kinæ kinæ ætherea composita, Tinctura cinchonæ ætherea composita—t. Kinkinæ amara, Tinctura cinchonæ amara.

TINCTURA KI'NO, *Tincture of Ki'no. (Kino, in pulv. 3vj, Alcohol. dilut. q. s.* Put the kino, mixed with an equal bulk of sand, into a percolator, and gradually pour on diluted alcohol until f3viij of filtered liquor pass. The tincture should be renewed frequently, and kept in closely-stopped bottles. *Ph. U. S.*) Astringent. Dose, f3j to f3ij.

TINCTURA KRANE'RIÆ, *Tincture of Rhat'omy. (Kramer. in pulv. 3vj, Alcohol. dilut. Oij.* Macerate for 14 days, express, and filter. It may also be prepared by displacement. *Ph. U. S.*) Dose, f3j to f3ij.

TINCTURA LACTUCÆRII, *Tincture of Lactuca'rium. (Lactucar. in pulv. 3iv, Sp. tenuior. Oij. (Imp. meas.)* Best prepared by percolation. *Ph. E.*) Dose, 30 minims to f3j.

TINCTURA LAURI CINNAMOMI, Tinctura cinnamomi—t. Lavandulæ composita, Spiritus lavand. comp.

TINCTURA LIMO'NIS, *Tincture of Lem'on peel. (Limon. cort. 3iiss, Sp. tenuior. Oij. (Imp. meas.)* Macerate for 7 days, express, and filter. *Ph. D.*) Dose, f3j to f3ij.

TINCTURA LOBE'LIÆ, *Tincture of Lobe'lia. (Lobel. 3iv, Alcohol. dilut. Oij.* Macerate for 14 days, express, and filter. It may also be prepared by displacement. *Ph. U. S.*) Dose, f3j to f3ij.

TINCTURA LOBE'LIÆ ÆTHE'REA, *Ethe'real Tincture of Lobe'lia. (Lobel. pulv. 3v, Sp. æther. sulph. Oij. (Imp. meas.) Ph. E. L. and B.)* Dose, same as the last.

TINCTURA LUPULI, Tinctura Humuli, T. Lupulinæ.

TINCTURA LUPULI'NÆ, T. Lu'puli, (*Ph. E.*) *Tincture of Lu'pulin. (Lupulin. 3iv, Alcohol. Oij.* Macerate for 14 days, and filter. *Ph. U. S.*) Dose, f3j to f3ij.

TINCTURA LYTTE, Tinctura cantharidis—t. Martis Glauberi, see Ferrum tartarizatum—t. Martis Mynsichti, Tinctura ferri ammoniata—t. Martis in Spiritu salis, Tinctura ferri muriatis—t. Martis vinosæ, Vinum ferri.

TINCTURA MATI'CO, *Tincture of Mati'co. (Matico in pulv. crass. 3viij (avoird.), Sp. tenuior. Oij. (Imp. meas.)* Macerate for 14 days, express, and filter. *Ph. D.*) Dose, f3j to f3ij.

TINCTURA MELAMPODII, Tinctura hellebori nigri—t. Meloes vesicatorii, Tinctura cantharidis—t. Metallorum, Lilium Paracelsi—t. Mimosæ catechu, Tinctura catechu.

TINCTURA MOS'CHI, *Tincture or Es'sence of Musk. (Mosch. in pulv. redact. 3ij, Sp. vini rect. Oj.)* Dose, f3j to f3iv.

TINCTURA MYR'RHÆ, *Tincture of Myrrh, Sim'ple Tincture of Myrrh. (Myrrhæ contusæ 3iij, Alcohol Oij.* Macerate for 14 days, and filter. Prepared also by displacement. *Ph. U. S.*) Tonic, 'deobstruent', antiseptic, and detergent. It is chiefly used in gargles, and applied to foul ulcers, spongy gums, &c.

Had'son's Preser'vative for the Teeth and Gums consists of Tinct. myrrh, Tinct. cinch., Ag. cinnam. 3ss. 3ij, Eau d'arquebuse 3j, Alo. gum. Arb. 3ss. M.

TINCTURA NERVINA BESTUCHEFFII, Tinctura

(seu alcohol) sulphuricoætherea ferri—t. Nervina Halensis, Tinctura (seu alcohol) sulphuricoætherea ferri—t. Nervinotonica, Lanotte's, Alcohol (seu Tinctura) sulphuricoæthereus ferri.

TINCTURA NUCIS VOMICÆ, *Tincture of Nuc Vom'ica. (Alcohol. s. g. 837, f3j, Dry Extract of Nuc Vomica, 3 gr.)* Dose, gtt. v to xxx, in cases where the nux vomica is indicated.

It is directed in the *Ph. U. S.* (1851), to be prepared as follows:—(*Nucis vomica rasur. 3viij, Alcohol Oij.* Macerate for 14 days, express, and filter. It may also be prepared by the process of displacement.) Dose, gtt. v to xx.

TINCTURA OLEI MENTHÆ PIPERITÆ, Spiritus menthæ piperitæ.

TINCTURA OLEI MENTHÆ VIRIDIS, Spiritus menthæ viridis.

TINCTURA O'PII, T. Theba'ica seu Sedati'va, *Ther'aca cales'tis, Al'cohol cum O'pio, Tincture of O'pium, Liq'uid Laudanum. (Op'ii pulv. 3ijss, Alcohol. dilut. Oij.* Macerate for 14 days, express, and filter. Prepared also by displacement. *Ph. U. S.*) Nineteen minims contain about one grain of opium. Properties, those of opium. Dose, gtt. xx to lx, and more. By macerating the dregs remaining after making the Tinctura Op'ii in a solution of Tartaric Acid, a solution is obtained which is said to be devoid of the exciting and constipating properties of the tincture of opium. Gray calls it *Haden's Li'quor Op'ii Sedati'vus*.

TINCTURA O'PII ACETA'TA, *Ac'etated Tincture of O'pium. (Op'ii 3ij, aceti f3xij, Alcohol. Oss.* Rub the opium with the vinegar, add the alcohol, and having macerated for 14 days, express, and filter. *Ph. U. S.*) Twenty drops are equivalent to a grain of opium.

TINCTURA O'PII AMMONIA'TA, *Amo'niated Tincture of O'pium, Ed'inburgh Paregor'ic Eliz'ir. (Flor. benz., croci, 3ss, 3iij, O'ii 3ij, Ol. anisi 3ss, Alcohol. ammoniat. f3xvj. Digest. Ph. E.)* f3j contains one grain of opium. Used like the Tinctura Camphoræ Composita.

TINCTURA OPII CAMPHORATA, Tinctura camphoræ composita—t. Plumbosæ, Liquor Plumbi subacetatis dilutus.

TINCTURA O'PII DEODORA'TA, *Deo'dorized Tincture of O'pium. (A solution of opium, 3iiss, in water, shaken with ether; and, the ethereal solution having been poured off, evaporating until all traces of ether have disappeared; mixing with water, and filtering; and adding water through the filter sufficient to make the liquid a pint and a half; adding, lastly, half a pint of alcohol, and mixing. Ph. U. S., 1863.)* By treating the opium with water in place of alcohol, the resinous and odorous principles are sparingly taken up. Its strength is that of Tinctura Op'ii.

TINCTURA QUAS'SIÆ, T. Quas'siæ excel'sæ, *Tincture of Quas'sia. (Quassie rasur. 3ij, Alcohol dilut. Oij.* Macerate for 14 days, express, and filter. It may also be prepared by displacement. *Ph. U. S.*) Tonic. Dose, f3j to f3ij.

TINCTURA QUAS'SIÆ COMPOS'ITA, *Com'pound Tincture of Quas'sia. (Cardam. sem. contus., Cucci contus. 3ss, Cinnam., Quassia, 3vj, Uvar. passar. 3viij, Sp. tenuior. Oij. (Imp. meas.)* Digest for 7 days, express, and filter. May be prepared also by percolation. *Ph. E.*) Dose, as a tonic, f3j or f3ij.

TINCTURA QUI'NÆ COMPOS'ITA, *Com'pound Tincture of Qui'na. (Quina sulph. 3v and 3j, Tinct. auranti, Oij, (Imp. meas.)* Digest for 7 days, and filter. *Ph. L.*) f3j contains a grain of the sulphate.

TINCTURA QUI'NÆ SULPHA'TIS, *Tincture of Sul'phate of Qui'na. (Sulphate of Quinia gr. vj, Alcohol (847) f3j. M.)* Dose, f3j to f3iv, in the day.

Tinctura Cinchoſiſe Sulphatis may be made in a ſingular manner.

TINCTURA RHABARBARI, *Tinctura rhei*—*t. Rhabarbari composita*, *Tinctura rhei composita*—*t. Rhabarbari vinosa*, *Vinum rhei palmati*.

TINCTURA RHEI, *T. Rhabarbari* ſeu *Rhabarbari ſpirituſa* ſeu *Rhei* ſeu *Rhei palmati*, *Tincture of Rhu'barb.* (*Rhei contus.* $\mathfrak{Z}$ ij, *Cardam. cont.* $\mathfrak{Z}$ ss, *Alcohol. dilut.* Oij. Macerate for 14 days, expreſs, and filter. It may alſo be prepared by diſplacement. *Ph. U. S.*) Stomachic or purgative, according to the doſe employed— $\mathfrak{f}\mathfrak{Z}$ i to $\mathfrak{Z}$ j as a ſtomachic; $\mathfrak{f}\mathfrak{Z}$ iv to $\mathfrak{Z}$ j as a purgative.

TINCTURA RHEI COMPOSITA, *T. Rhabarbari composita*, *Compound Tincture of Rhu'barb.* (*Rhei rad. concis.* $\mathfrak{Z}$ ij, *Glycyrrhiz. rad. cont.* $\mathfrak{Z}$ ss, *Zingib. rad. concis.*, *Croci stigmat.* aa $\mathfrak{Z}$ ij, *Aque* $\mathfrak{f}\mathfrak{Z}$ xij, *Sp. ten. Oj.* *Ph. L.*) Properties ſame as the laſt.

TINCTURA RHEI ET AL'OEIS, *Elis'ir ſacrum*, *Tincture of Rhu'barb and Aloes.* (*Rhei concis.* $\mathfrak{Z}$ x, *Aloes pulv.* $\mathfrak{Z}$ vj, *Cardam. cont.* $\mathfrak{Z}$ ss, *Alcohol. dilut.* Oij. Macerate for 14 days, expreſs, and filter. *Ph. U. S.*) Properties like the laſt.

Rymer's Cardiac Tincture conſiſts of an infusion of *Capsicum*, *Camphor*, *Cardamom ſeeds*, *Rhubarb*, *Aloes*, and *Caſtor* in *Proof Spirit*, with a very ſmall quantity of *Sulphuric Acid*.

TINCTURA RHEI ET GENTIANÆ, *Tinctura Rhei Amara*, *Tincture of Rhu'barb with Gentian.* (*Rhei cont.* $\mathfrak{Z}$ ij, *Gentian. cont.* $\mathfrak{Z}$ ss, *Alcohol. dilut.* Oij. Macerate for 14 days, expreſs, and filter. *Ph. U. S.* It may alſo be prepared by diſplacement.) Properties ſame as the laſt.

TINCTURA RHEI ET SEN'NÆ, *Tincture of Rhu'barb and Sen'na*, *Warner's Gout Cordial.* (*Rhei cont.* $\mathfrak{Z}$ j, *Sennæ* $\mathfrak{Z}$ ij, *Coriandr. cont.*, *Fenicul. cont.* aa $\mathfrak{Z}$ j, *Santal. raſur.* $\mathfrak{Z}$ ij, *Croci*, *Ext. glycyrrhiz. rad.* aa $\mathfrak{Z}$ ss, *Uvar. paſſar.* acinis exemptis, *lbss*, *Alcohol. dilut.* Oij. Macerate for 14 days, expreſs, and filter. *Ph. U. S.*) Doſe, $\mathfrak{f}\mathfrak{Z}$ ss to $\mathfrak{f}\mathfrak{Z}$ ij.

TINCTURA RHEI, *Tinctura rhei*—*t. Rhei amara*, *Tinctura rhei et gentianæ*—*t. Sacra*, *Vinum aloes*.

TINCTURA SABI'NÆ, *Tincture of Sav'ine.* (*Sav'ine*, dried and bruised, $\mathfrak{Z}$ iiss, *Proof ſpirit* Oj. Made by percolation. *Ph. B.*, 1864.) Doſe, $\mathfrak{M}$ xv to xxx.

TINCTURA SANGUINARIÆ, *Tincture of Blood-root.* (*Sanguinar. cont.* $\mathfrak{Z}$ iv, *Alcohol. dilut.* Oij. Macerate for 14 days, expreſs, and filter. It may alſo be prepared by diſplacement. *Ph. U. S.*) Doſe, 30 to 60 drops.

TINCTURA SAPONIS CAMPHORATA, *Linimentum saponis camphoratum*—*t. Saponis et opii*, *Linimentum saponis et opii*.

TINCTURA SCILL'Æ, *Tincture of Squill.* (*Scille* $\mathfrak{Z}$ iv, *Alcohol. dilut.* Oij. Macerate for 14 days, expreſs, and filter. It may alſo be made by diſplacement. *Ph. U. S.*) Doſe, gtt xv to lx.

TINCTURA SEDATIVA, *Tinctura opii*.

TINCTURA SEN'NÆ, *T. Sen'ne Composita*, *Elis'ir Salutis*, *Daf'ly's Elis'ir*, *Tincture of Sen'na.* (*Sennæ fol.* $\mathfrak{Z}$ ij, *Carui ſem.* cont. $\mathfrak{Z}$ ij, *Cardam. ſem.* cont. $\mathfrak{Z}$ j, *Uvarum paſſarum demptis acinis*, $\mathfrak{Z}$ iv, *Sp. ten. Oij.* *Ph. L. & D.*) Stomachic, carminative and cathartic. Doſe, $\mathfrak{f}\mathfrak{Z}$ ij to $\mathfrak{f}\mathfrak{Z}$ j.

TINCTURÆ SEN'NÆ ET JALAP'Æ, *Tincture of Sen'na and Jalap.* (*Sennæ* $\mathfrak{Z}$ ij, *Jalap. pulv.* $\mathfrak{Z}$ j, *Coriandr. cont.*, *Carui cont.*, aa $\mathfrak{Z}$ ss, *Cardam. cont.* $\mathfrak{Z}$ ij, *Sacch.* $\mathfrak{Z}$ iv, *Alcohol. dilut.* Oij. Macerate for 14 days, expreſs, and filter. *Ph. U. S.* It may alſo be prepared by diſplacement.) Doſe, $\mathfrak{f}\mathfrak{Z}$ ij to $\mathfrak{f}\mathfrak{Z}$ j.

TINCTURA SERPENTARIÆ, *T. Serpentariæ* ſeu *Virginianæ* ſeu *Aristolochiæ Serpentariæ*, *Tincture of Snake-root.* (*Rad. serpent.* $\mathfrak{Z}$ iv, *Alcohol. dilut.* Oij. Macerate for 14 days, expreſs, and

filter. It may alſo be prepared by diſplacement. *Ph. U. S.*) Tonic. Doſe, $\mathfrak{f}\mathfrak{Z}$ ss to $\mathfrak{f}\mathfrak{Z}$ ij.

TINCTURA STOMACHICA, *Tinctura cardamomi composita*.

TINCTURA STRAMO'NII, *Tincture of Stramonium.* (*Stramon. ſem.* cont. $\mathfrak{Z}$ iv, *Alcohol. dilut.* Oij. Macerate for 14 days, expreſs, and filter. It may alſo be prepared by diſplacement. *Ph. U. S.*) Doſe, ten to twenty minims.

TINCTURA STRYCH'NIÆ, *Tincture of Strychnia.* (*Alcohol. s. g.* 837, Oj, *Strychniæ*, gr. ij.) Doſe, 6 drops to 24. Properties, thoſe of ſtrychnia.

TINCTURA (ſeu AL'COHOL) SULPHURICOÆTH'EREA FER'RI, *Gul'tæ nervinæ*, *Liquor Anodynus Martialis*, *Al'cohol æthereus ferratus*, *Naph'tha Vitrioli Martialis*, *Solutio Murialis fer'ricæ æthereæ*, *Spiritus sulphuricoæthereus martialis*, *Tinctura nervina Halen'sis* ſeu *Tonicum nervinum Halen'sis* ſeu *au'rea* ſeu *nervina Bestucheff's* ſeu *nervinotonica* (*Lamotte's*) ſeu *æthereæ fer'ri*, *Æther Martialis*, *De La Motte's Golden Drops*, *Bestucheff's Nervous Tincture*, (*F.*) *Elis'ir d'Or de M. le Général De La Motte.* (An ethereal ſolution of chloride of iron.) It is much uſed in gout, hypochondriaſis, &c.

TINCTURA SULPHURIS VOLATILIS, *Liquor fumans Boylii*—*t. Thebaica*, *Tinctura opii*, *Vinum opii*.

TINCTURA TOLUIF'ERÆ BAL'SAMI, *T. Bal'sami Tolutani*, *T. Tolutani* (*Ph. U. S.*, 1842), *T. Tolutana* (*Ph. U. S.*, 1851 and 1863), *Tincture of Balsam of Tolu'*, (*Bals. Tolutan.* $\mathfrak{Z}$ ij, *Alcohol.* Oij.) Reputed to be expectorant and corroborant. Doſe, $\mathfrak{f}\mathfrak{Z}$ ss to $\mathfrak{f}\mathfrak{Z}$ j, or more.

TINCTURA TONICA NERVINA HALENSIS, *Tinctura (ſeu Alcohol) sulphuricoætherea ferri*.

TINCTURA VALERIANÆ, *Tincture of Valerian.* (*Valerian. cont.* $\mathfrak{Z}$ iv, *Alcohol. dilut.* Oij. Macerate for 14 days, expreſs, and ſtrain. It may alſo be prepared by diſplacement. *Ph. U. S.*) Stimulant and antispasmodic. Doſe, $\mathfrak{f}\mathfrak{Z}$ ss to $\mathfrak{f}\mathfrak{Z}$ ij.

TINCTURA VALERIANÆ AMMONIAT'Æ, *T. Valerianæ Volatilis*, *Ammoniated or Volatile Tincture of Valerian.* (*Valerianæ cont.* $\mathfrak{Z}$ iv, *Sp. ammon. arom.* Oij. Macerate for ſeven days, expreſs, and filter. It may alſo be made by diſplacement. *Ph. U. S.*) Like the former. Doſe, $\mathfrak{f}\mathfrak{Z}$ ss to $\mathfrak{f}\mathfrak{Z}$ ij.

TINCTURA VERA'TRI, *T. Veratri al'bi*, *Tincture of White Hel'lebores.* (*Rad. veratri alb.* cont. $\mathfrak{Z}$ vij, *Alcohol. dilut.* Oijss. *Ph. E.*) Emetic and cathartic; but not very manageable. Doſe, gtt. v to x.

TINCTURA VERA'TRI VIRIDIS, *Tincture of American Hel'lebores.* (*Veratr. virid. pulv.* $\mathfrak{Z}$ xvj, *Alcohol. q. s.* to produce by percolation two pints. *Ph. U. S.* 1863.) Sedative. Doſe, gtt. vij, every three hours until the pulſe is reduced.

TINCTURA ZINGIB'ERIS, *T. Amo'mi Zingib'eris*, *Tincture of Gin'ger.* (*Zingib. cont.* $\mathfrak{Z}$ vij, *Alcohol.* Oij. Macerate for 14 days, expreſs, and filter. It may alſo be made by diſplacement. *Ph. U. S.*) Stimulant and carminative. Doſe, $\mathfrak{f}\mathfrak{Z}$ ss to $\mathfrak{f}\mathfrak{Z}$ j.

TINCTURÆ, *Tinctura*—*t. of Aconite*, *Tinctura aconiti*—*t. of Aloes*, *Tinctura aloes*—*t. of Aloes*, compound, *Tinctura aloes composita*—*t. of Aloes*, ethereal, *Tinctura aloes æthereæ*—*t. of Ammonia*, compound, *Spiritus ammoniæ succinatus*—*t. of Angustura*, *Tinctura angusturæ*—*t. of Arnica*, *Tinctura arnicæ*—*t. of Assaſetida*, *Tinctura assaſetidæ*—*t. of Balsam of Tolu*, *Tinctura toluiferæ balsami*—*t. of Bark*, *Tinctura cinchonæ*—*t. of Bark*, ammoniated, *Tinctura cinchonæ ammoniata*—*t. of Bark*, compound, *Tinctura cinchonæ composita*—*t. of Bark*, *Huxham's*, *Tinctura cinchonæ composita*—*t. of Bark*, volatile, *Tinctura cinchonæ ammoniata*—*t. of Belladonna*, *Tinctura belladonnæ*—*t. of Benzoin*, compound, *Tinctura benzoini composita*—*t. Bestucheff's*, ſee *Tinctura ferri*

muriatis—t. of Blistering fly, *Tinctura cantharidis*—t. of Bloodroot, *Tinctura sanguinaria*—t. of Buchu, *Tinctura buchu*—t. of Camphor, compound, *Tinctura camphoræ composita*—t. of Cantharides, *Tinctura cantharidis*—t. of Capsicum, *Tinctura capsici*—t. of Cardamoms, *Tinctura cardamomi*—t. of Cardamoms, compound, *Tinctura cardamomi composita*—t. Cardiac, Rymer's, *Tinctura rhei et alicæ*—t. of Cascarella, *Tinctura cascariellæ*—t. of Castor, *Tinctura castorei*—t. of Castor, compound, *Tinctura castorei composita*—t. of Catechu, *Tinctura catechu*—t. of Cayenne pepper, *Tinctura capsici*—t. of Chiretta, *Tinctura chirette*—t. of Cinchona, *Tinctura cinchonæ*—t. of Cinchona, compound, *Tinctura cinchonæ composita*—t. of Cinnamon, *Tinctura cinnamomi*—t. of Cinnamon, compound, *Tinctura cinnamomi composita*—t. of Cochineal, *Tinctura coccii cacti*—t. of Colchicum, compound, *Spiritus colchici ammoniatus*—t. of Colchicum seeds, *Tinctura colchici*—t. of Columbo, *Tinctura calumbæ*—t. of Croton, *Tinctura crotonis*—t. of Cubebs, *Tinctura cubebæ*—t. of Digitalis, *Tinctura digitalis*—t. of Ergot, *Tinctura ergotæ*—t. of Ergot, ethereal, *Tinctura ergotæ æthereæ*—t. of Galbanum, *Tinctura galbani*—t. of Galls, *Tinctura gallarum*—t. of Gentian, compound, *Tinctura gentianæ composita*—t. of Ginger, *Tinctura zingiberis*—t. for the Gout, Dr. Wilson's, see *Colchicum autumnale*—t. of Guaiacum, *Tinctura guaiaci*—t. of Guaiacum, ammoniated, *Tinctura guaiaci ammoniata*—t. of Guaiacum, volatile, *Tinctura guaiaci ammoniata*—t. Hatfield's, see *Tinctura guaiaci ammoniata*—t. of Hellebore, black, *Tinctura hellebori*—t. of American Hellebore, *Tinctura veratri viridis*—t. of Hellebore, white, *Tinctura veratri*—t. of Hemlock, *Tinctura conii*—t. of Henbane, *Tinctura hyoscyami*—t. of Hops, *Tinctura humuli*—t. of Indian hemp, *Tinctura cannabis Indicæ*—t. of Iodine, *Tinctura iodini*—t. of Iodine, compound, *Tinctura iodini composita*—t. of Iron, acetate of, *Tinctura ferri acetatis*—t. of Iron, ammoniated, *Tinctura ferri ammoniata*—t. of Iron, ammoniochloride of, *Tinctura ferri ammoniochloridi*—t. of Iron, muriated, *Tinctura ferri muriatis*—t. of Jalap, *Tinctura jalapii*—t. of Kino, *Tinctura kino*—t. Klaproth's, see *Tinctura ferri muriatis*—t. of Lactucarium, *Tinctura lactucarii*—t. of Lemon peel, *Tinctura limonis*—t. of Litmus, see *Lichen roccella*—t. of Lobelia, *Tinctura lobeliæ*—t. of Lobelia, ethereal, *Tinctura lobeliæ æthereæ*—t. of Ludwig, see *Ferrum tartarizatum*—t. of Lupulin, *Tinctura lupulinæ*—t. of Matico, *Tinctura matico*—t. of Musk, *Tinctura moschi*—t. of Myrrh, *Tinctura myrrhæ*—t. Nervous, Bestucheff's, *Tinctura (seu alcohol) sulphuricoæthereæ ferri*—t. of Nux vomica, *Tinctura nucis vomicæ*—t. of Oil of Peppermint, *Tinctura olei menthæ piperitæ*—t. of Oil of Spearmint, *Tinctura olei menthæ viridis*—t. of Opium, *Tinctura opii*—t. of Opium, acetated, *Tinctura opii acetata*—t. of Opium, ammoniated, *Tinctura opii ammoniata*—t. of Opium, camphorated, *Tinctura camphoræ composita*—t. of Opium, deodorized, *Tinctura opii deodorata*—t. of Orange peel, *Tinctura aurantii*—t. of Quassia, *Tinctura quassiæ*—t. of Quassia, compound, *Tinctura quassiæ composita*—t. of Quinia, compound, *Tinctura quiniæ composita*—t. of Quinia, sulphate of, *Tinctura quiniæ sulphatis*—t. of Rhatany, *Tinctura kramerie*—t. of Rhubarb, *Tinctura rhei*—t. of Rhubarb compound, *Tinctura rhei composita*—t. of Rhubarb and aloes, *Tinctura rhei et aloes*—t. of Rhubarb and gentian, *Tinctura rhei et gentianæ*—t. of Rhubarb and senna, *Tinctura rhei et sennæ*—t. of Saffron, *Tinctura croci sativi*—t. of Saffron, compound, *Tinctura de croco composita*—t. of Senna, *Tinctura sennæ*—t. of Senna and jalap, *Tinctura sennæ et jalapæ*—t. of Snake root, *Tinctura serpentariæ*—t. of Soap, camphorated, *Linimentum*

saponis—t. of Spanish fly, *Tinctura cantharidis*—t. of Spearmint, *Tinctura olei menthæ viridis*—t. of Squill, *Tinctura scillæ*—t. of Steel, *Tinctura ferri muriatis*—t. Stomach, *Tinctura cardamomi composita*—t. of Stramonium, *Tinctura stramonii*—t. of Strychnia, *Tinctura strychniæ*—t. for the Teeth, Greenough's, see *Spiritus armoraciæ compositus*—t. Toothache, *Tinctura antiodontalgica*—t. of Valerian, *Tinctura valerianæ*—t. of Valerian, ammoniated, *Tinctura valerianæ ammoniata*—t. of Valerian, volatile, *Tinctura valerianæ absinthii*—t. of Wormwood, compound, *Tinctura absinthii composita*.

TINDER, ([Sax.] *tyndre*, *tenðre*, from *tyñan*, 'to kindle,') *Boletus igniarius*.

TINEA, 'a gnawing worm,' Porriño—t. Capitis, Porriño—t. Ciliaris, see *Ophthalmia tarsi*—t. of the Eyelids, see *Ophthalmia tarsi*—t. Faciei, Porriño larvalis—t. Favosa, Porriño favosa—t. Ficiosa, Porriño scutulata—t. Furfuracea, Porriño furfurans—t. Granulata, Porriño scutulata, Porriño lupinosa—t. Lactea, Porriño larvalis—t. Lupina, Porriño lupinosa—t. Mucosa, Porriño lupinosa—t. Porriginosa, Porriño furfurans—t. Tarsi, see *Ophthalmia tarsi*—t. Tendens, Porriño decalvans—t. Tonsurans, Porriño decalvans—t. Vera, Porriño favosa.

TINEÆ, pl. of Tinea, Worms.

TINESMUS, Tenesmus.

TING'LING, (*tinuire*, 'to jingle,') (Sc.) *Dirling*. A sharp, purring or thrilling sensation, not always painful.

TINKER'S WEED, *Triosteum*.

TINKLING, METALLIC, (like tingling,) *Tintement métallique*.

TINNIMENTUM METALLICUM, (from *tinnio*, 'to tinkle,') *Tintement métallique*.

TINNITUS AURIUM, (from *tinnio*, *tinnitus*, 'to tinkle,') 'ringing of the ears.' *Fluctuatio seu Sonitus seu Susurrus seu Sibilus seu Streptitus seu Sibilismus aurium*, *Paracusis imaginaria*, *Otclusus*, *Syrigimus*, *Syringismus*, *Sibilus clangorinus*, *E'chos*, *Eneche'ma*, *Bom'bus*, *Bombitatio*, *Mur'mur*, (F.) *Tintement*, *T. d'Oreille*, *Tintouin*, *Bourdonnement*, *Bruissement*. An imaginary sound, like the ringing of a bell, the noise of wind, the murmur of water, &c., heard in health and in sickness. It is, often, an accompaniment of cerebral disorder.

TINTEMENT, Tinnitus aurium—t. *Bullaire*, *T. métallique*.

TINTEMENT MÉTALLIQUE (F.), *Cliquetis métallique*, *Tintement bullaire*, *Tinnimentum metallicum*, *Ca'nor stethoscopicus*, *Metal'lic Respiration*, *Metal'lic Voice*, *Metal'lic Tink'ling*. A particular noise, heard by the stethoscope when applied to the chest, and which, according to Laënnec, resembles that caused by striking glass, metal, or porcelain, with a pin; but is, perhaps, more like the sound of the keys of a musical snuffbox. The *tintement métallique* or *metallic tinkling* is heard on causing the patient to speak or breathe; but is more distinct when he coughs. When the phenomenon is not so strongly marked, it produces only the *metal'lic resonance*. The 'metallic tinkling' is a pathognomonic symptom of a communication between the bronchia and cavity of the chest; in other words, of *Pneumothorax*.

TINTEMENT D'OREILLE, Tinnitus aurium.

TINTINNABULUM, 'a bell,' (from *tintinnio*, 'to ring,') *Uvula*.

TINTOUIN, Tinnitus aurium.

TIPFOOT, see *Kyllosis*.

TIPSARIA, Decoctum hordei.

TIQUE (F.), (G.) *Zecke*, *Ixodes ricinus*.

TIRE-BALLE, (*tirer*, 'to draw,' and *balle*, 'a ball,') Forceps (bullet.)

TIRE-FOND (F.), (*tirer*, 'to draw,' and *fond*, 'bottom.') A surgical instrument, formerly used to elevate the piece of bone sawed off by the trephine.

TIRE-PUS, Pyulcon.

TIRE-TÊTE (F.), (*tirer*, 'to draw,' and *tête*, 'head,') *Tractor*. A name given to different instruments used for extracting the head of the child when left in the uterus, and in certain cases of difficult labour. The crochets are one of those, as well as the *Tire-tête à biseau* of Levret, &c.

TIRES, Milk sickness.

TIRETOIR (F.), *Tirtoir*, (from *tirer*, 'to draw,') An instrument used by dentists for extracting the incisors of the lower jaw.

TISANE (F.), *Ptisana* or *Ptisana*, *Ptisana*, from *πτισάνω*, 'pearl barley,' itself from *πτισσω*, 'I decocticate.' (F.) *Hydroë*. A name given to aqueous medicines, containing but little, if any, medicinal agent. The ancients gave the name particularly to the decoction of barley.

TISANE COMMUNE, Decoctum hordei.

TISANE DE FELTZ. A celebrated anti-syphilitic, formed of *sarsaparilla*, *isinglass*, *sulphuret of antimony* and *water*, boiled together. According to some, bichloride of mercury; and, to others, arsenic enters into its composition. M. Rayer, supposing that it acts by the arsenic it contains, proposes to replace it by arseniate of potassa; and Guibourt, who is of the same opinion, by arsenious acid, in order to ensure certainty of action.

TISIC, Phthisis.

TISICAL, Phthisicus.

TISSICK, see Phthisic.

TISSU, (from *texere*, *textum*, 'to weave,') *Tissue*—*t.* *Accidental*, see *Accidental*, and *Tissue*, *accidental*—*t.* *Aréolaire*, Cellular tissue—*t.* *Caverneux*, Cavernous texture—*t.* *Cellulaire*, Cellular tissue—*t.* *Celluleux des os*, Cancelli—*t.* *Conjunctif*, Cellular tissue—*t.* *Cribleux*, Cellular tissue—*t.* *Inodulaire*, Tissue, inodular—*t.* *Jaune*, see *Fibrous*—*t.* *Lamineux*, Cellular tissue—*t.* *Lardacé*, see *Lardaceous*—*t.* *Muqueux*, Cellular tissue—*t.* *Poreux*, Cellular tissue—*t.* *Réticulé*, Cellular tissue—*t.* *Spongoïde*, Spongoid tissue—*t.* *Unitif*, Cellular tissue.

TISSUE, *Textus*, *Te'la*, (F.) *Tissu*. By this term, in anatomy, is meant the various parts, which, by their union, form the organs; and are, as it were, their anatomical elements. Histological anatomy is the anatomy of the tissues, which are the seat of the investigations of the pathological anatomist. The best division, indeed, of diseases would be according to the tissues mainly implicated. For the elementary tissues, see *Fibre*. The compound tissues, which, by associating or combining variously, form every organ of the body, have been variously classed; but every division is imperfect and liable to objection.

TISSUE, ACCIDENTAL OR ADVENTITIOUS, consists of every substance—foreign to the primitive organization of the economy, but yet organized and living—which may be developed in the interior or at the surface of organs. Laënnec divides the accidental tissues into two sections: 1. Those that resemble others in the animal economy, or *analogous accidental tissues*; and, 2. Those that have nothing analogous in the body, and which are always the result of a morbid process—the *heterologous* or *heterolite accidental tissues*. To the first class belong, ossifications; accidental fibrous, fibrocartilaginous, cartilaginous, areolar, and corneous tissues; hairs; the serous membranes of certain encysted tumours; the mucous membranes of fistulous canals, and accidental synovial membranes. To the second belong tubercles, scirrhi, melanosis, &c.

TISSUE, AREOLAR, Cellular tissue.

TISSUE, CARTILAGINOUS, *Hymenochondroïdes* seu *Hymenochondroïdes* (*Textura seu Te'la*.)

TISSUE, CELLULAR, Cellular tissue—*t.* *Connective*, Cellular tissue—*t.* *Cribriform*, Cellular tissue—*t.* *Fibrous*, see *Fibrous*—*t.* *Filamentous*, Cellular tissue.

TISSUE FORMATION, Histogeny.

TISSUE, INODULAR, *Te'la inodulaire*, (F.) *Inodule*, *Tissu inodulaire*. A name given by Delpech to an accidental fibrous tissue developed in suppurating wounds, which is the principal agent in cicatrization. It has, at first, the appearance of a reddish cellulofibrous layer; but soon loses its vascularity; and its fibres, which pass, in all directions, become of a dull white colour, and of a consistence and hardness, that may be compared to those of the strongest articular ligaments. It is well seen in the cicatrix left after burns.

TISSUE, LAMINATED, Cellular tissue—*t.* *Ligamentous*, Desmoid tissue—*t.* *Mucous*, Cellular tissue—*t.* *Porous*, Cellular tissue—*t.* *Reticulated*, Cellular tissue.

TIT, (τιτθ, [Sax.] τιτ,.) Nipple.

TITHENE, (τιθηνη,) Nurse.

TITHYMALUS CYPARISSIAS, (τιθυμαλος,) *Euphorbia cyparissias*—*t.* *Lathyrus*, *Euphorbia lathyrus*—*t.* *Latifolius*, *Euphorbia lathyrus*—*t.* *Palustris*, *Euphorbia palustris*—*t.* *Paralias*, *Euphorbia paralias*.

TITILLAMENTUM, Gargarism.

TITILLATION, (*titillare*, *titillatum*, 'to tickle,') Tickling.

TITTHE, (τιτθ,) Mamma, Nipple.

TITTHION, (τιτθιον, dim. of *titthos*,) Nipple.

TITTHIS, (τιτθις,) Mamma, Nipple.

TITTHOS, (τιτθος,) Mamma, Nipple.

TITTY, Mamma, Nipple.

TITUBANTIA, 'a staggering,' *Bredouillement*.

TITUBATIO, (*titubare*, *titubatum*, 'to stagger or totter,') Fidgets, Vacillatio—*t.* *Linguae*, Balbuties.

TJETTEK, Upas tieuté.

TMETICA, (τμητικά, from *τεμαίνω*, 'to cut,') Incidentia.

TO-AND-FRO-SOUND, *Bruit de frottement*.

TOADFLAX, *Antirrhinum linaria*—*t.* *Bastard*, *Comandra umbellata*.

TOASTWATER, *A'qua to'ste pan'is*, (F.) *Eau panée*. (Toast well half a slice of a stale quarter loaf; put it into a pitcher, and pour over it a quart of water. After two hours, decant the water from the bread.) It is a common drink in febrile affections.

TOBACCO, (from *tabacos*, the Caribbean name for the pipe in which it was smoked [?]), *Nicotiana tabacum*—*t.* *English*, *Nicotiana rustica*—*t.* *Indian*, *Lobelia inflata*—*t.* *Poison*, *Hyoseyamus niger*.

TOBACCO ROOT, see *Valeriana*.

TOBACCO, WILD, *Lobelia inflata*, *Nicotiana rustica*.

TOCETUS, (τοκετος,) Parturition.

TODODOMYCODORITIS MALIGNA VAGINALIS, (*tocus*, *mucus*, 'mucus,' *ôpa*, 'skin,' and *itis*,) Colpoeace puerperarum.

TOCOLOGY, (*tocus*, and *logos*, 'a discourse,') Obstetrics.

TOCUS, (τοκος,) Parturition.

TODDY. A name given, by the English, to sweet, delicious and refreshing liquors procured in India by wounding the *spathæ* of certain palms. The best is obtained from the *Cocos nucifera*.

It is obtained in large quantities from *Saguë-rus saccharifer* seu *Rum'phii*, *Pal'ma Indica vinaria* seu *Indica*, *Saguë-rus* seu *Gomm'us Gomm'us*, *Aren'ga saccharifera*, *Gomm'us palm*. It is called, also, *Ne'va*, and, by boiling, yields a coarse kind

of sugar, called *Jugjery*. The inhabitants of the Moluccas used, in their wars, a liquor obtained by macerating the fruit, which the Dutch called *Hell water*, *Aqua infernalis*.

Toddy is, also, the common name for a mixture of spirit and water sweetened, and usually spiced with nutmeg.

TODDY TREE, Mamei.

TOE, ([Sax.] *ta*.) *Digitus pedis*—*t. Supernumerary*, see *Monster*.

TOEPLITZ, see *Töplitz*.

TOFTS, *Tophus*.

TOILE D'ARAIGNÉE, 'spider's web,' *Aranea tela*—*t. Choroidienne*, see *Choroid*—*t. de Gautier*, *Sparadrapum Galteri*.

TOKAS, (*tokas*, gen. *tokados*.) *Puerpera*.

TOKEN, PLAGUE, see *Plague token*.

TOKOLOGY, (*tokos*, and *logos*, 'a discourse,') *Obstetrics*.

TOLA, *Tonsil*.

TOL'ERANCE, *Tolerantia*, (*tolerare*, 'to bear.') The power of bearing. A word used by the Italian school of *Rasori*, to signify the power of bearing large doses of certain potent remedies, as tartrate of antimony and potassa. See *Contra-stimulus*.

TOLES, *Tonsil*.

TOLIUM, *Tonsil*.

TOLLES, *Tonsil*.

TOLLUM, *Tonsil*.

TOLU, (from *Tolu* in South America,) see *Toluifera balsamum*.

TOLUIFERA BAL'SAMUM, (*tolu*, and *ferre*, 'to bear,') *Myracrylon* seu *Myrospermum Toluiferum*. *Ord. Leguminosæ. Sex. Syst. Decandria Monogynia*. The name of the tree which affords the *Tolu*, *Bal'sam of Tolu*, *Bal'samum Tolutanum* (Ph. U. S.), *Tolutanum*, *Toluiferæ Bal'sami Bal'samum*, *Goac'onaz*, *Red Bal'sam of Peru*, (F.) *Baume de Tolu*, *B. de Carthagène ou d'Amérique ou de Saint Thomas*. It has a very fragrant odour, and warm, sweetish taste; is thick, and of a reddishyellow colour. It is stimulant and expectorant (?), and is used in coughs. It is chiefly, however, employed on account of its flavour.

TOLUTANUM, see *Toluifera balsamum*.

TOM OF BEDLAM, see *Abraham man*.

TOMACINUM, (*τομήν*, 'a cutting up,') *Isicium*.

TOMACLUM, (same etymon,) *Isicium*.

TOMACULUM, (same etymon,) *Isicium*.

TOMATA, (Indian,) *Solanum lycopersicum*.

TOMATO, *Solanum lycopersicum*.

TOMÉ, *To'mus*, (*τομήν*, *τομός*, 'incision,') A common suffix, as in *Lithotomy*, *Bronchotomy*, &c.

TOMÉ, *Cut*, *Incision*.

TOMEION, (*τομεῖον*.) *Knife*.

TOMENTUM CER'EBRI, (*tomentum*, 'a flock of wool,') The infinite number of small vessels which pass between the pia mater and convolutions of the brain. They enter the substance of the brain at right angles.

TOMEUS, (*τομεύς*.) *Knife*.

TOMIS, (*τομῆς*.) *Scissors*.

TOMOTOCIA, (*τομή*, 'section,' and *τομός*, 'parturition,') *Cæsarean section*.

TOM TOE, *Pollex*, see *Digitus pedis*.

TONE, *To'nus*, (*τόνος*, from *τονός*, 'I brace or give vigour,') The state of tension or firmness proper to each organic tissue. It is the effect of tonicity.

TONE, see *Elasticity*.

TONGA, see *Datura sanguinea*.

TONGRES, MIN'ERAL WATERS OF. The town of Tongres is about a league from Maestricht in the Low Countries. The springs, of which there are several, are chalybeate containing carbonates of iron and magnesia.

TONGUE, (Sax. *tung*, [D.] *Tong*.) *Lin'gua*, *Glos'sa*, *Glot'ta*, (F.) *Langue*. The tongue is a muscular, symmetrical, and very mobile organ: situate in the mouth, from the os hyoides and epiglottis to the incisor teeth. Its particular use is, to procure the sensation of savours; and it aids in the acts of sucking, mastication, deglutition, pronunciation, and expulsion. It has the shape of a pyramid, flattened above and below; rounded at its angles, and terminated, anteriorly, by a blunt point. Its upper surface, called the *dorsum*, is free, flat, and divided into two portions by a median furrow, *lin'ea median'a*; at the posterior extremity of which is the *fora'men cæcum* or *Fora'men Morgagni*. The inferior surface is free, and covered by the mucous membrane on its anterior third and sides: but behind and in the middle it is attached to the lower jaw by means of the *genio-glossus*; and to the hyoides by the *hyoglossus*. Its margins are rounded; thick behind, thinner anteriorly. Its tip, or point, or dental extremity is rounded and free:—the root, base, or hyoid extremity, *Rad'ice lin'gue*, is continuous with the epiglottis and velum palati. It is very thick at the *foramen cæcum*; but thinner as it approaches the hyoid bone. The tongue is composed of muscles, and covered by a mucous membrane—*Periglott'is*, *Invol'crum lin'gue*, *Membrana seu Tu'nica vagina'lis seu Cut'is seu Epider'mis seu Tu'nica seu Invol'crum mucosum seu Crus'ta villosa lin'gue*—which forms also the frænum beneath the tongue. On this membrane are seen the different papillæ. The nerves of the organ proceed from the inferior maxillary, *glossopharyngeal* and *hypoglossal*. The first and second are connected with taste; the last with the motions of the organ. Some recent observers, however, consider the *glossopharyngeal* to be the special nerve of gustation. The arteries are the *lin'qual*, from the external carotid; and the *palatine* and *tonsillar*, from the labial. The veins are the *superficialis lin'gue*, *rani'na*, *lingual'is*, and *submentalis*. They open into those of the pharynx and larynx. Its lymphatic vessels pass into ganglia, situate at the edge of the *hyoglossi* muscles.

TONGUE, BLACK. A fever which prevailed in the Western States in the winter of 1842-3; probably typhoid fever. According to some, an epidemic erysipelas.

TONGUE, HERNIA OF THE, *Glossocèle*.

TONGUE HOLDER, *Glossocatochus*.

TONGUE, INFLAMMATION OF THE, *Glossitis*—*t. Red*, fever, see *Typhus*.

TONGUE SCRAPER, *Cure-langue*—*t. Tie*, see *Ankyloglossum*.

TON'IC, (*τόνος*.) *Ton'icus*, (F.) *Tonique*. Relating to tone.

Tonic, *Ton'icus*, *Tono'ticus*, *Antasthen'icus*, (F.) *Tonique*, *Fortifiant*, in Therapeutics, means a medicine, which has the power of exciting slowly, and by insensible degrees, the organic actions of the different systems of the animal economy, and of augmenting their strength in a durable manner. Bitter, vegetable substances which are not associated with an acid or narcotic principle; preparations of iron; cold water, &c., act as tonics. They are also called *Comfortan'tia*, *Confortativa*, *Carroboran'tia*, *Confirman'tia*, *Roboran'tia*, &c.

The following are the chief Tonics:—*Acidum Arseniosum*, *Acidum Nitricum*, *Acidum Sulphuricum*, *Argentum Nitras*, *Bismuthi Subnitras*, *Carbo Ligni*, *Cascarilla*, *Cinchona*, *Cinchonia*, *Colombia*, *Cornus Florida*, *Cupri Acetas*, *Cupri Sulphas*, *Cuprum ammoniatum*, *Cusparia*, *Eupatorium perfoliatum*, *Ferri Preparata*, *Gentiana*, *Liriodendron*, *Lupulina*, *Piperina*, *Prunus Virginiana*, *Quassia*, *Quinia*, *Salix*, *Salicina*, *Serpentaria*, *Zinci*

præparata: Aquæ minerales acidulæ et ferruginæ; and Mental Tonics, (Confidence, Hope, Amusement.)

TONIC POWER, Irritability.

TONIC SPASM, *Spasmus tonicus*, *Convulsio tonica*, *Conductio*, *Spas'ticus*, *Tonus*, *Entonia*, *Entasis*, *Eutasia*. A rigid contraction of muscles without relaxation, as in trismus, tetanus, &c. See Tetanus and Spasm.

TONICITY, *Tonicitas*. The faculty that determines the general tone of the solids. Excessive tonicities causes *erethism* or *crispness*; deficient tonicities, *atony* or *weakness*. Tonicity has been used in the same sense as irritability.

TONIQUE, Tonic.

TONKA BEAN, *Dipterix odorata*.

TONODES, (*τονωδης*.) Tensive.

TONOTICUS, (*τονωτικός*.) Tonic.

TONSIL, *Tonsilla*, *Amygdala*, *Paristh'mia*, *Paristh'mium*, *Amphibranchium*, *Glandula colli*, *To'la*, *To'les*, *To'les*, *To'lium*, *To'lum*, *Spong'os*, *An'tius*, pl. *Ant'ides*, *Almonds of the ear*, *Almonds or Bulbs of the throat*, (F.) *Amygdale*, *Tonsille*, (Prov.) *Earkecker*. An ovoid collection of follicles, about half an inch in length, similar in shape to an almond, and situate, on each side, between the pillars of the velum palati. The inner surface, projecting into the isthmus of the throat, is covered by the mucous membrane, in which are the orifices of a dozen cells, whence a transparent, viscid mucus oozes on pressure. This mucus is destined to facilitate deglutition, by lubricating the isthmus faucium; and is forced out during the passage of the alimentary bolus.

TONSIL-GULLOTTINE, *Amygdalotome*.

TONSILLA, pl. *Tonsillæ*, *Tonsil*.

TONSILLÆ PESTILENTES, *Cynanche maligna*.

TON'SILLAR, *Tonsillaris*, (F.) *Tonsillaire*. Appertaining or relating to the tonsils.

TONSILLARUM GANGRÆNA, *Cynanche maligna*.

TÖNSILLE, *Tonsil*.

TONSILLIA, *Cynanche tonsillaris*.

TONSILLITIC, *Tonsilliticus*, (better *Tonsilliv.*) Belonging or relating to the tonsils; as the 'tonsillitic branches of the glossopharyngeal nerve.'

TONSILLITIS, (*tonsilla*, and *itis*.) *Cynanche tonsillaris*.

TONSILLITOME, (*tonsilla*, and *τεμνω*, 'to cut,') *Amygdalotome*.

TONSORIUS CULTER, (*tonsor*, 'a barber,' and *cultus*, 'a knife,') *Razor*.

TONUS, (*τονος*.) Tonic spasm—t. Trismus, Trismus.

TOODLE, *Tooth*.

TOOTH, (Sax. *toth*.) *Dens*, *Mor'dex*, *O'dous*, (Prov.) *Toodle*, (F.) *Dent*. The teeth, *Den'tes*, *Mord'ices*, are small, hard bodies fixed in the alveoli of each jaw, and serving to seize, and to cut, tear, and triturate, alimentary substances. True teeth are found only in man, the mammalia, reptiles, and fish. The teeth have, in general, the shape of an irregular cone, the base of which is towards the interior of the mouth; and the apex, single or divided, towards the bottom of the alveoli. Each tooth has a part situate out of the alveolus, and covered by a vitriform substance. This part is called the *crown* or *corona*. It is bounded by a narrower portion, called the *cervix*, (F.) *collet*, which separates it from the portion in the alveolus—the *fang* or *root*, *Odon'torrhiza*, (F.) *racine*. In the adult, the teeth are 32 in number; 16 to each jaw: these consist of 4 *incisors*, 2 *canine*, 4 *bicuspidati*, and 6 *molars*. (See these various terms.) The teeth enclose a vascular and nervous pulp, in a central cavity, *Cavitas Pul'pæ*, *Ca'vum Den'tis*, *An'trum den-*

ta'le. They are formed of a very hard, compact tissue; and the corona is covered by a vitreous substance called *Enam'el*. The incisor and canine teeth are developed by a single point of ossification; the lesser molars, by two; and the larger, by four or five. See Dentition. Researches by Retzius, Müller and others, have led them to believe that the ivory of the tooth, *Den'tine* or *Dentin*, *E'bur*, *Substantia os'sea* of Malpighi, *proper tooth substance*, *bone of the tooth*, *os'seum substance of tooth*, *toothbone*, consists of tubular fibres, which contain organic deposits of calcareous salts; and of intertubular substance, in which are observed corpuscles or osseous cells of the tooth, called, by Professor Owen, *calcig'erous cells*, which also contain earthy salts. At the part where the enamel terminates at the cervix of the tooth, the *crus'ta petro'sa*, *cæmentum*, *cementum* or *cortical substance*, commences in an extremely thin stratum, passes down and gradually increases in thickness towards the extremity of the root, where it is generally the thickest. It is said by Nasmyth to have been traced on the enamel, and Purkinje and Fränkel thought that it lined the cavitas pulpæ. They gave it the name *Substantia os'toidea*. It is the *mate'ria tartare'a* of Malpighi. It has been found, however, that the hard substance deposited on the inner surface of the dentine, at the age of twenty years or later, and which encroaches on the cavity, has a distinct structure from the cement, and resembles in part dentine, and in part bone. It is the *horn'y substance* of Blumenbach, the *osteodentine* of Owen, and the *secondary dentine* of Tomes. In the compound teeth of the herbivorous animal, a deposition is found on the surface of the enamel, which has been called *cementum*. Müller, however, regards it to be a deposit from the salts and the saliva, and to be essentially the same as what is called tartar in the human subject.

TOOTHACHE, *Odontalgia*—t. *Bush*, *Xanthoxylum fraxineum*—t. *Tree*, *Aralia spinosa*, *Xanthoxylum clava Herculis*.

TOOTHBONE, see *Tooth*—t. *Edge*, *Agacement des dents*, *Hæmodia*—t. *Paste*, *Dentrificium*—t. *Rake*, *Denticulapium*—t. *Rash*, *Strophulus*—t. *Shell*, *dog-like*, *Dentalium*.

TOPHACEOUS, *Tophaceus*, (F.) *Tophacé*; (from *tophus*.) Relating or belonging to tophus, —as 'a *tophaceous* concretion.'

TOPHACEOUS CONCRETION, *Tophus*.

TOPHUS, (*τοφος*.) *To'fus*, 'a porous or friable stone.' *Osteo'ophus*, *Tophaceous Concretion*. A collection of hard, calcareous matter, which forms, particularly, at the surface of joints affected with gout, *Arthro'ophus*, (see *Calculi*, *arthritic*), and occasionally, in the interior of organs, around the teeth, &c. It likewise meant gravel, and chazaza.

TOPIC, *Top'icus*, (from *τοπος*, *locus*, 'a place,') A local application; (F.) *Topique*. See *Local*.

TOPICUS, (*τοπίκος*.) *Local*, *Topic*.

TOPINARIA, *Talpa*.

TOPIQUE, *Topic*.

TÖPLITZ or TEP'LITZ, MIN'ERAL WATERS OF. Töplitz is a town of Bohemia, celebrated for its numerous thermal springs. The water contains chloride of sodium; carbonate, and sulphate of soda; and carbonates of lime and iron, &c. Temp. 114° to 122° Fah. Season, from the 1st of June to the 15th of September.

TOPOGRAPHICAL ANATOMY, (*τοπος*, 'a place or region,' and *γραφω*, 'to describe,') see *Anatomy*.

TOQUE, (a kind of hat,) *Scutellaria galericulata*.

TORCULAR, (from *torqueo*, 'to twist,') *Tourniquet*. A press for wine, oil, or cider.

TORCULAR HEROPH'ILL, *Press of Heroph'illus*,

Lecheu'on, Le'nos, Le'num, Fourth Sinus, (F.) Confluent des Sinus, Pressoir d'Hérophile. A smooth and polished cavity, of irregular shape, in which several sinuses of the dura mater meet. It is situate before the internal occipital protuberance, at the union of the three great duplicatures of the dura mater. It has six openings into it;—one above, that of the superior longitudinal sinus; two below, answering to the occipital sinuses; one anterior, belonging to the straight sinus; and two lateral, and very broad, which lead into the lateral sinuses; and convey from the cavity the blood that has been poured into it by the others.

TORCULUM, (from *torqueo*, 'to twist,') Tourniquet.

TORDYLUM, (τορδύλιον.) *T. Officinale*, *Tordylum*, (τορδύλον,) *Seseli Creticum* seu *Montanum*, (F.) *Siseli de Crète.* Family, Umbelliferae. *Sex. Syst.* Pentandria Digynia. The seeds and roots of this Southern European plant are considered carminative and diuretic. They enter into the theriaca.

TORÉNIA ASIATICA, (after Olaf Toren, of Gothenburg, a pupil of Linnæus,) *Caladula*. A small Malabar plant, the juice of whose leaves, sweetened with sugar, is said to cure gonorrhœa.

TORMENTIL, *Geranium maculatum*, *Tormentilla*.

TORMENTILLA, (from *tormina*, for which it was used,) *T. erecta* seu *sylvestris* seu *officinalis* seu *tuberosa*, *Fragaria tormentilla* *officinalis*, *Potentilla tormentilla*, *Heptaphyllum*, *Consolida rubra*, *Comamon Tormentil*, *Upright Septfoil*, (F.) *Tormentille.* *Ord.* Rosaceæ. *Sex. Syst.* Icosandria Polygynia. The root, *Tormentilla* (Ph. U. S.), has a slightly aromatic odour; and austere, styptic taste. It is astringent. Dose, of the powder, gr. x to ʒi.

TORMENTUM, 'pain, suffering,' (from *torqueo*, 'to twist,') *Ileus*, *Intussusceptio*.

TORMINA, pl. of *Tormen*, *Strophus*, *Cæliatigæ*, *Encæliatigæ*, *Anilema*, *Anilesis*, *Gripes*, *Mulligrubs*, *Fretting*, (F.) *Tranchées*, (from *torqueo*, 'I twist,') Acute colicky pains. *Dysentery*.

TORMINA CELSI, *Dysentery*—t. *Parturientium*, Pains, labour—t. *Post partum*, see Pains, labour.

TORNACULUM, (from *torquere*, 'to twist,') Tourniquet.

TOROSITAS, (from *torosus*, 'brawny,'—*torus*, 'brawn,') *Corpulentia carnosæ*, *Habitus athleticus*. Muscular strength.

TOROSUS, Muscular.

TORPEDO, (*torpeo*, 'to be torpid,') Narcosis.

TORPEFACIO UNIVERSALIS, (*torpeo*, 'to be torpid,' and *facere*, 'to make,') *Holonanarcosis*.

TORPEUR, Torpor.

TORPID, *Torpidus*, (F.) *Torpidæ*, *Engourdi*, (from *torpeo*, 'to be sluggish,') Numb. Incapable of exertion and of feeling. Dull, stupid, inactive.

TORPIDITAS, Torpor.

TORPIDUS, Narcotized, Torpid.

TORPOR, gen. *Torporis*, *Torpiditas*, *Nothotes*, *Parapsis ex'pers*, *Torpidity*, (F.) *Engourissement*. State of a part which becomes numb, and almost incapable of motion. *Torpor*, *Torpidity*, *Torpidness*, *Torpidude* are also applied to a debilitated, sluggish condition of a part, or of the whole of the body; *Ær'gia*, (F.) *Accablement*.

TORPOR, Narcosis—t. of the Colon, see Colon, torpor of the—t. *Intestinorum*, *Constipation*—t. *Soporificus*, *Nocar*.

TORQUAY, (pron. *Torkee*.) A town in Devonshire, England, which, on account of the mildness of its climate, and its sheltered situation from the bleak winds, is a frequent retreat for the phthisical invalid.

TOR'SION, *Tor'sio*, (from *torquere*, *tortum*, 'to twist,') The act of twisting. *Torsion* of the ex-

tremities of bleeding vessels—*Angio'trophæ*—until the blood ceases to flow, is sometimes used to arrest hemorrhage.

TORSK, see *Oleum jecoris aselli*.

TORTELLE, *Erysimum*.

TORTICOLLIS, (*torquere*, *tortum*, 'to twist,' and *collum*, 'the neck,') *Torticollis*, *Caput seu Collum obliquum*, *Obliqua cervicis*, *Obstipitata capitis* seu *Collis*, *Trachel'agra*, *Entonia Loxia*, *Lox'ius*, *Céphalobol'ia*, *Ancylob'erè*, *Ancylob'eris*, *Rhabocra'nia*, *Rhaban'chen*, *Rhabod'erè*, *Auchenorrh'ema*, *Rheumatismus collis* seu *cervicis*, *Stiff-neck*, *Wry'neck*, (Prov.) *Slab*, (F.) *Obstipité*. A variety of rheumatism, seated in the muscles of the neck, which prevents the motion of the head, and causes the patient to hold it inclined to the side affected. It is, commonly, of short duration; usually disappearing in a few days. Rube-faciants are the best topical applications. The term is also applied to permanent contraction of the muscles of the neck, which causes the head to be held to one side. The remedy is the division of the contracted muscles.

TORTOISE, FRESHWATER, GREAT, (*torquere*, *tortum*, 'to twist, to wind,') *Hydraspis expansa*.

TORTUE, 'tortoise,' *Testudo*.

TORTURA ORIS, (from *torqueo*, 'to twist,') Canine laugh, *Neuralgia faciei*—t. *Oris* paralytica *Linnæi*, *Paralysis*, *Bell's*.

TOR'ULA CEREVISIÆ, (from *torus*, 'a knot,') *T. Ace'ti*, *Saccharomyces* seu *Mycoderma cerevisiæ*, *Cryptococcus fermentum* seu *cerevisiæ*, (F.) *Champignon du ferment*, *Yeast plant*. One of the simplest forms of vegetation, capable, under favourable circumstances, of vegetating rapidly, and assisting in producing the fermentation of saccharine substances. The plant has been found in vomited fluids, and in fæces.

TORULUS GLANDIS, (dim. of *torus*, 'a fleshy part,') Crown of the glands.

TORUS, 'a fleshy part,' Muscle—t. *Manûs*, *Metacarpus*.

TOTA BONA, 'all good,' *Chenopodium bonum*, *Henricus*.

TOTSU KASO, *Sphæria Sinensis*.

TOTTY, *Vertiginous*.

TOTUM CARNOSUM, 'all fleshy,' *Pancreas*.

TOUCH, *Tactus*, *Tactio*, *Hap'hæ*, *Hap'ais*, *Tactio*, *Palpation*, (F.) *Toucher*, (from [L.] *to care*, 'to touch,') One of the five senses, and that which makes known to us the palpable qualities of bodies, such as their consistence, quality, &c. It is seated, particularly, in the hands; and differs from tact in being active. See *Esaphe*, and *Palpation*.

'To touch the mouth with mercury' is to produce the constitutional effects of that agent, as evinced by the falling away of the gums, &c.

TOUCH-ME-NOT, *Impatiens fulva*.

TOUCH ROYAL. It was at one time almost universally believed, in England, that the royal touch would remove *Scurfula* or *King's Evil* (as it was thence called.) This superstition is now entirely exploded. Edward, the Confessor, was the first English king who touched for the Evil. Medals of gold or silver were often suspended around the necks of the persons touched. These were called *touching* or *healing pieces*.

TOUCHER, 'touch,' *Esaphe*, *Touch*—t. *Vaginal*, *Esaphe*.

TOUCHING PIECE, see *Touch*, royal.

TOUCHWOOD, *Boletus ignarius*.

TOULOUOUNIN, see *Carapa touloucouna*.

TOUR DE MAÎTRE, 'a master turn or feat,' *Coup de maître*.

TOURBILLON VASCULAIRE, (from *turbo*, gen. *turb'inis*, 'a whirlpool,') *Vasa vasorum*.

TOURNESOL, ([F.] *turner*, 'to turn,') and

sol, 'the sun.'—a translation of ἡλιοσποριον, *heliotropium*.) *Heliotropium Europæum*.

TOURNIOLE, (from *tourner*, 'to turn,' like *Rumex* and *Paronychia*.)

TOURNIQUET, (pron. *toorniket* or *turniket*.) *Torcular*, *Torculum*, *Tornaculum*, *Pra'lium*, *Pra'lium*, *Pra'lium* seu *Prelum arteria'le*, *Verticil'um*, *Fascia tortilis*, (from [F.] *tourner*, 'to turn.') A surgical instrument for stopping the course of blood in the limbs, by exerting a strong compression upon the principal artery. It was invented by a French surgeon, Morel, in the middle of the 17th century; and was modified subsequently, by Nuck, Verdus, Monro, Petit, and Louis. It is used to suspend, for a time, the circulation in a limb, during great operations; to arrest arterial hemorrhage in cases of wounds; to compress certain aneurismal and other tumours, &c.

TOUS-LES-MOIS, 'every month.' *Am'ylum cornu'cum*, *Canna* (Ph. U. S.), *Canna Starch*. A fecula obtained from the root of *Canna edu'lis*, which flowers every month. It resembles potato starch, and may be substituted for arrowroot. It is obtained from St. Kitts.

TOUT, Nates.

TOUTE BONNE, 'all good,' *Salvia selarea*.

TOUTE ÉPICE, 'all spice,' see *Myrtus pimenta*—*t. Sainte*, *Sanicula*.

TOUX, Tussis—*t. Bleue*, *Pertussis*—*t. Bronchique*, *Bronchial cough*, *Cough tubal*—*t. Coverneuse*, see *Cavernous respiration*—*t. Convulsive*, *Pertussis*—*t. Quinteuse*, *Pertussis*—*t. Tubaire*, *Bronchial cough*, *Cough tubal*.

TOW, ([Sax.] *τορ*), *Stupa*.

TOXÆMIA, *Toxiœhæmia*.

TOXÆMIC, *Toxiœhm'ic*, *Toxiœ'mic*, *Hæmatœ'ic*, *Hæmatotox'ic*, *Hæmatœ'ic*, *Toxæ'micus*, *Toxiœhm'icus*, *Toxæ'micus*, *Hæmatœ'icus*, *Hæmatotox'icus*, *Hæmatœ'icus*. Relating or appertaining to a poisoned state of the blood.

TOXÉMIE, *Toxiœhæmia*.

TOXICÆMIA, *Toxiœhæmia*.

TOXICARIA MACASSARIENSIS, see *Upas*.

TOXICAL, *Poisonous*.

TOXICATIO, (from *toxicum*), *Poisoning*.

TOXICÉMIE, *Toxiœhæmia*.

TOXICODENDRON, (*toxicum*, and δένδρον, 'a tree,') *Rhus toxicodendron*.

TOXICOHÆMIA, *Toxiœhm'ia*, *Toxæ'mia*, (F.) *Toxiœhémie*, *Toxiœmie*, *Toxémie*, (*toxicum*, and *αἷμα*, 'blood,') *Poisoning of the blood*.

TOXICOHÉMIE, *Toxiœhæmia*.

TOXICOÏD, *Toxiœ'ides*, (*toxicum*, and εἶδος, 'resemblance,') *Resembling poison*.

TOXICOL'OGY, *Toxicolog'ia*, (*toxicum*, and *λογος*, 'a description,') *A treatise on poisons*.

TOXICOSES, (G.) *Toxikosen*, (from *toxi-cum*, 'an arrow,' originally 'arrow poison,' and *οἷα*.) A family of diseases, according to the classification of Fuchs, caused by the reception of poisons into the system.

TOXICUM, (*τοξικον*, from *τοξον*), *Poison*, *Venom*.

TOXIQUE, *Poisonous*.

TRABECULA, (dim. of *trabs*, gen. *trabis*, 'a beam,') (F.) *Trabécule*, 'a small beam.' Applied to prolongations of fibrous membrane, for example, which form septa, as in the corpora cavernosa; to the medullary fibres, which form the commissures; and to the fleshy projections, which form the columnæ carneæ, &c.

TRABECULA CEREBRI, *Corpus callosum*.

TRABECULÆ CARNEÆ, *Columnæ carneæ*—1. *Cordis*, *Columnæ carneæ*—*t. Corporum cavernosorum*, see *Cavernous bodies*—*t. Splenis*, see *Spleen*—*t. Willisii*, *Chordæ Willisii*.

TRABES CORDIS, (pl. of *trabs*), *Columnæ carneæ*.

TRABS, 'a beam,' *Penis*—*t. Cerebri*, *Corpus callosum*—*t. Medullaris*, *Corpus callosum*.

TRACE, *PRIMITIVE*, *Groove*, *primitive*.

TRACHÉA, (*τραχεια*, fem. of *τραχης*, 'rough,' *απρηγία*, 'understood,') (generally, however, accented *Trachéa*;) *Trachéa* seu *Traché'a artéria* (*τραχεια απρηγία*), *Traché'us*, *Bronchus*, *Aspera artéria*, *Eistula spirital'is* seu *pulmonalis*, *Cun'na*, *Syrin'ga*, *Syr'inx*, the *Wind'pipe*, *Wo'sen*, *Weas'and*, *Wizen*, *Wee'sel*, *Throl'tle*, (Sc.) *Throp'ill*, (Prov.) *Keck'corn*, *Keck'er*, *Throp'ple*, (F.) *Trachée*, *Trachée artère*; (composed of *αἶρ*, 'air,' and *τραχεα*, 'to keep,') The trachea is a cylindrical, fibrocartilaginous and membranous tube, flattened behind, and situate on the median line, before the vertebral column, from the inferior part of the larynx, as far as the third dorsal vertebra. There it divides into two branches, the *bronchia*, which separate, one to go to each lung. The trachea, the function of which is to convey air to the lungs during respiration, is composed, 1. Of cartilaginous rings, *An'uli cartilagin'ci*, *Segmenta cartilagin'ca*, *Orbea cartilagin'o'si*, incomplete behind; situate one above the other, and kept *in situ* by a fibrous membrane. 2. Of a mucous membrane, which lines it internally, and contains numerous mucous follicles. 3. Of transverse muscular fibres, not well-marked, which occupy its posterior surface. 4. Of vessels, which proceed from the superior and inferior thyroideals; and, 5. Of nerves, that are given off by the pneumogastric and the cervical ganglia.

The breathing tubes of insects are termed *Trachéæ*.

TRACHEA ARTERIA, *Trachea*.

TRACHEÆ, see *Trachea*.

TRACHÉAL, *Traché'alis*, (F.) *Trachéal*. Appertaining to the trachea. An epithet applied to respiration as heard through the stethoscope opposite the trachea, larynx, and root of the bronchia; the air appearing as if sucked in from the cylinder during inspiration, and expelled during expiration.

TRACHEAL GLANDS. Mucous follicles on the posterior surface of the trachea. They are small, flattened ovoid bodies.

TRACHEALIA, (from *trachea*), *Cynanche trachealis*.

TRACHEALIS. The fibres of unstriped muscle, which pass from one portion of each of the tracheal cartilages to the other, at the posterior part of the trachea, have been so called by some anatomists. When they contract, they approximate the extremities of the cartilage; diminish the calibre of the trachea; and thus increase the velocity of the air forced through it, as in coughing; and facilitate the removal of any irritating substance in contact with the lining membrane.

TRACHÉE, *Trachea*—*t. Artère*, *Trachea*.

TRACHEITIS, (*trachea* and *itis*), *Cynanche trachealis*.

TRACHELAGRA, *Torticollis*, (*τραχηλας*, 'neck,' and *αγρα*, 'seizure,') Also, *gout in the neck*.

TRACHELIAN, (from *trachelus*), *Cervical*.

TRACHELISM'US, (from *trachelus*.) A term proposed by Dr. Marshall Hall, for a fancied spasmodic action of the muscles of the neck, which he esteems to be the cause of many morbid phenomena, by inducing compression of the veins of the neck and an impeded return of blood from the head.

TRACHELITIS, (*trachelus*, and *itis*), *Cynanche trachealis*.

TRACHELIUM AMERICANUM, (from *trachelus*, having been used in throat affections), *Lo-belia cardinalis*.

TRACHÉLO-ATLOÏDO-BASILAIRES, *Rectus capitis lateralis*—*t. Atloïdo-occipital*, *Obliquus superior capitis*—*t. Basilaire*, *Rectus capitis in-*

ternus major—*t. Basilaire*, *petit*, Rectus capitis internus minor.

TRACHELODIAPHRAGMATIC, *Trachelo-diaphragmaticus*. The fourth pair of cervical nerves, whence the phrenic nerve chiefly arises.

TRACHELODYN'IA, (*trachelus*, and *δύνη*, 'pain,') Pain in the neck.

TRACHÉLO-MASTOÏDIEN, Complexus minor—*t. Occipital*, Complexus—*t. Scapulaire*, Levator scapulae.

TRACHELUS, Collum, Trachea.

TRACHEOBRONCHITIS. Inflammation of the trachea and bronchia.

TRACHEOCELE, (*trachea*, and *κῆλη*, 'a tumor,') Bronchocele.

TRACHEOPHONY, (*trachea*, and *φωνή*, 'voice,') Laryngophony.

TRACHEOPHTHISIS, see Phthisis laryngea.

TRACHEOPHYMA, (*trachea*, and *φύμα*, 'a swelling,') Bronchocele.

TRACHEOPYRA, (*trachea*, and *πύρ*, 'fever,') Cynanche trachealis.

TRACHEORRHAG'IA, *Hæmoptoë* seu *Hæmoptysis laryngea* et *trachealis*, (*trachea*, and *ῥήγνυμι*, 'to break forth,') Hæmorrhage from the trachea.

TRACHEOSTENO'SIS, (*trachea*, and *στενωσις*, 'contraction,') Contraction or narrowness of the trachea.

TRACH'EOTOME, (*trachea*, and *τέμνω*, 'to cut,') An instrument of the trocar kind, for opening the trachea.

TRACHEOTOMY, see Bronchotomy.

TRACHITIS, (*trachea*, and *ίτις*,) Cynanche trachealis.

TRACHO'MA, gen. *Tracho'matis*, (from *τράχυν*, 'rough,') *Aspre'do*, *Aspritu'do*, *Pladar'otes*, *Asperitudo*, *Scabrit'ies*, *Gron'ular conjunct'iva*, *Gron'ular eye'tid*, (F.) *Aspérité des Paupières*. A roughness of the inner surface of the eyelids. A variety of ophthalmia, of which three kinds have been designated:—*T. sabulo'sum*, from sand, &c., getting between the eyelids; *T. carunculo'sum*, *Pladar'otes*, *Pladaro'sis*, *Morum pul'pebræ inter'næ*, from fleshy excrescences; and *T. herpet'icum*, from hard pustules on the inner surface of the eyelids. This last is, also, called *Fico'sis*, and *Pul'pebræ fico'sa*.

TRACHYPHON'IA, (*τράχυν*, 'rough,' and *φωνή*, 'voice,') Roughness of the voice.

TRACT, *Trac'tus*, (from *trahere*, *tractum*, 'to draw out lengthwise,') A track or course; a region.

TRACT, OPTIC, see Optic nerves—*t. Respiratory*, see Respiratory tract.

TRACTATIO MANUARIA, (*tractare*, *tractatum*, 'to handle,') Diacheirismus, Surgery.

TRACTORATION, (*trahere*, *tractum*, 'to attract, influence,') Perkinism, see Tractors, metallic.

TRACTATORES, (*tractare*, *tractatum*, 'to handle,') see Shampooing.

TRACTOR, (*trahere*, *tractum*, 'to draw,') Lever, *Tire-tête*.

TRACTORS, METAL'LIC. Metallic agents, introduced by Dr. Elisha Perkins, of Connecticut, about the end of the last century, to cure diseases. The parts were rubbed with these tractors, and hundreds of thousands of cases were reported to have been cured by the *Tractor'ation*. The affections, in which they were chiefly used, were local inflammations, and pains in different parts. The good effects were, doubtless, owing to the influence exerted by the mind over the body. The bubble did not exist long. See Perkinism.

TRACTUS, Tract—*t. Intestinorum*, Intestinal Tube—*t. Opticus*, see Optic nerves—*t. Respiratorius*, Respiratory tract.

TRACTUS SPIR'ALIS FORAMINULENTUS. Minute openings at the base of the modiolus, arranged

in a spiral manner, which transmit the filaments of the cochlear nerve. The central canal of the Tractus is longer than the rest, and is called *Tubulus central'is mod'oli*.

TRADESCANT'IA VIRGIN'ICA, (after John Tradescant, gardener to Charles I.,) *Spider'wort*; indigenous; Order, Commelynaceæ; flowering from May to August. The roots are demulcent.

TRAGACANT'HA, (*τραγος*, 'a goat,' and *κανθα*, 'a horn,' owing to its hirsute boughs.) *Gummi Tragacanthæ* seu *Astrag'ali Tragacanthæ*, *Gum Tragacanth*, *Gum Drag'on*. The concrete juice of *Astrag'alus Tragacantha* or *Astragalus Fémus*, a native of Persia. Ord. Leguminosæ. *Sex. Syst.* Diadelphia Decandria. (F.) *Gomme Adragant*, *Adragant*, *Adragante*, *Adraganthe*. This gum is inodorous; nearly insipid; of a whitish colour; semitransparent and striated. It is in thin, vermiform pieces; and does not form a smooth, uniform mucilage with water. It is demulcent, but is rarely used alone.

TRAGEA, *Tragea*—*t. Aromatica*, Pulvis cinnamomi compositus.

TRAGE'MA, gen. *Trage'matis*, (*τραγῆμα*, 'a sweetmeat,') Of old, in the plural, — *Trage'mata*, — it meant a dessert. Afterwards, like *Tragea*, it meant an aromatic powder, containing much sugar.

TRAG'ICUS, *Tragien* (Ch.), (from *Tragus*.) A small, flat, triangular muscle, which almost wholly covers the outer surface of the tragus, arising from its base, and terminating at the top of the eminence.

TRAGIEN, *Tragicus*.

TRAGOMASCH'IA, (*Tragomas'chalē*, (*τραγος*, 'a goat,' and *μασχαλή*, 'axilla,') The strong, unpleasant odour of the armpits.

TRAGOPHON'IA, (*τραγος*, 'a goat,' and *φωνή*, 'voice,') Egophony.

TRAGOPO'GON, *Ban'ba hir'ci*, (*τραγος*, 'a goat,' and *παγων*, 'beard,') A genus of plants; Ord. Cichoraceæ. *Sex. Syst.* Syngenesia Polygamia Æqualis. (F.) *Barbe de Bouc*; of which the roots of the *Tragopo'gon porrifo'lum*, *Sal'sifi*, *Sal'sify*, *Sal'sify*, *Oysterplant*, *Oysterroot*, *Veg'etable oyster*, (F.) *Salsify* ou *Cercifis*; and the young shoots of the *T. pratens'is*, *Mead'ow Sal'sify*, *Com'mon Goat's Beard*, (F.) *Salsify des prés*, are eaten as food. The root of the latter has been used as a diuretic.

TRAGOSELINUM, (*τραγος*, 'a goat,' and *σελίον*, 'parsley,') *Pimpinella saxifraga*—*t. Angelica*, *Ligusticum podagraria*—*t. Magnus*, *Pimpinella magna*—*t. Majus*, *Pimpinella magna*—*t. Saxifraga*, *Pimpinella magna*.

TRAG'US, (*τραγος*, 'a goat,' from its being furnished, in some persons, with hair, like the beard of a goat.) *Hir'cus*, *Hir'quus*. A small, flattened, triangular nipple, situate before the meatus auditorius externus, which it appears to conceal. It is covered with hair in old people, whence its name. Also, the peculiar goatlike smell of the axillæ.

TRAGUS MATTHIOLI, *Salsola kali*.

TRAINASSE, (from *trainer*, 'to trail,') Polygonum aviculare.

TRAINING, ([F.] *trainer*, from *trahere*, 'to draw,') (F.) *Entraînement*. A system of treating man and animals, so as to develop their full physical powers. It is practised especially as regards the prize-fighter and the race-horse, and appertains to what have been termed by the French *Organoplasties*.

TRAITEMENT, ([L.] *tractare*, [F.] *traiter*, 'to treat,') Curatio—*t. Palliatif*, see Palliation.

TRAME, (*τραμή*,) Perinæum.

TRAMIS, (*τραμς*,) Perinæum.

TRANCE, Catalepsy, Ecstasis—*t. Death*, see Asphyxia.

TRANCHÉES, (from *transcendere*, 'to cut across,') *Tormina*—*t. Uterine*, Pains, after.

TRANS, *n* composition, through, across, over, or beyond.

TRANSFIGURATIO, (*trans*, and *figurare*, *figuratum*, 'to form,') Transformation.

TRANSFORMATION, *Transformatio*, *Transfiguratio*, *Transmutatio*, *Metabolē*, (*trans*, and *forma*, 'form,') Change of form.

TRANSFORMATIONS DE TISSU (F.). The French pathologists designate thus the change that occurs in an organ, the tissue of which becomes similar to that of another organ. The chief transformations are cartilaginous, osseous, and adipous. See Tissue.

TRANSFUSION, *Transfusio*, *Metach'ysis*, (from *transfundere* (*trans*, and *fundere*, *fusum*, 'to pour,') To pour from one vessel into another.

TRANSFUSION OF BLOOD, *Transfusio Sanguinis*, *Cul'ra Medea'na*, *Methemachym'ia*, *Hæmatometach'ysis*, *Hæmatometach'ysis*, *Diach'ysis*, *Transplantatio medic'ia no'ra*, *Chirur'gia transfus'ria*, *Ecotichæmato'sis*. The act of passing the blood of one animal into the veins of another; an operation which was formerly used to fulfil various therapeutical intentions. It has been revived in cases of uterine hemorrhage; and, apparently, with some success.

TRANSLATION, (*trans*, and *ferre*, *latum*, 'to carry,') Metastasis.

TRANSMISSIBILITY, (*trans*, and *mittere*, *missum*, 'to send,') see Sensibility.

TRANSMUTATIO, (*trans*, and *mutare*, *mutatum*, 'to change,') Transformation.

TRANSPIRATIO, (*trans*, and *spirare*, *spiratum*, 'to breathe,') Perspiration—*t*. Unilatera, Hemidiaphoresis.

TRANSPIRATOIRE, Perspiratory.

TRANSPLANTATIO, (*trans*, and *plantare*, *plantatum*, 'to plant,') *Metaphyte'ia*. A pretended method of curing diseases, devised by Paracelsus—by making them pass from one individual to another, either animal or vegetable.

TRANSPLANTATIO, Morioplæsticæ.

TRANSPLANTATIO DENTIS, *Insit'io dentis*. The act of transplanting a tooth from one person to another.

TRANSPLANTATIO MEDICA NOVA, Transfusion.

TRANSPORT, (*trans*, and *portare*, 'to carry,') Delirium.

TRANSPPOSITIO, (*trans*, and *ponere*, *positum*, 'to place,') Metathesis, Transposition.

TRANSPPOSIT'ION, *Transposit'io*, *Metath'esie*. Change of situation.

TRANSPOSITION OF THE VIS'CERA, *Polype'ria promisc'ua translati'va*, *In'tus in'er'cus*, is a congenital vice of conformation, which consists in the viscera being found out of the situations they ordinarily occupy: the heart, for example, being on the right side; the liver on the left, &c.

TRANSUDATIO, (*trans*, and *sudare*, *sudatum*, 'to sweat,') Diapedesis, Transudation.

TRANSUDATION, *Transudatio*, (F.) *Transudation*. The passage of a fluid through the tissue of any organ, which may collect in small drops on the opposite surface, or evaporate from it.

TRANSUDATION, Exosmose.

TRANSVERSAIRE, Transverse—*t*. *Épineux*, *Transversalis dorsi*—*t*. *Épineux du cou*, *Semispinalis colli*—*t*. *Épineux du dos*, *Semispinalis dorsi*—*t*. *Grêle*, see *Sacrolumbalis*.

TRANSVERSAL DE LA CONQUE, *Transversalis auris*—*t*. *du Nez*, *Compressor nasi*—*t*. *des Orteils*, *Transversus pedis*.

TRANSVERSALIS, (from *transverse*,) Transverse.

TRANSVERSA' LIS ABDOM'INIS, (F.) *Lombo-abdominal*, (Ch.) *Transverse de l'Abdomen*. This muscle is seated, deeply, on the lateral parts of the abdomen. It is flat, thin, and broader before

than behind. It is attached, above, to the cartilage of the last true rib, and to those of every false rib; and, below, to the inner lip of the crista of the ilium; to the two outer thirds of the crural arch, and to the upper part of the pubis. Its fibres, moreover, are inserted, behind, by means of broad aponeuroses, into the top of the transverse and spinous processes of the first four lumbar vertebrae; and, before, into the linea alba, and the ensiform cartilage. This muscle, all of whose fibres have a transverse direction, constricts the belly, and diminishes the base of the chest, by drawing inwards the ribs to which it is attached.

TRANSVERSALIS ANTICUS PRIMUS, *Rectus capitis lateralis*—*t*. *Collateralis colli*, see *Sacrolumbalis*.

TRANSVERSALIS COL'LI, *Transversalis Major colli*, *Portion of the Sacrospinal* (Ch.), (F.) *Transversaire*. A muscle, seated at the posterior and lateral part of the neck and upper part of the back. It is attached to the transverse processes of the five or six lowest vertebrae of the neck; and to those of the four or five first dorsal. It extends the vertebrae of the neck, and inclines them to its side. See *Semispinalis colli*.

TRANSVERSALIS DORSI, *Transversospinalis*, *Multif'idus Spi'ne*, *Lumbocervical portion of the Spinal* (Ch.), (*Transverso-spinalis Lumborum*, *Transverso-spinalis Dorsi*, *Transverso-spinalis Col'li*), *Musculus sac'cer*, *Spinal'es et transversales lumborum*, *Semispinalis inter'na* sive *Transverso-spinalis col'li pars inter'na* (Winslow), *Lombodorsi-spinal*, &c., (F.) *Transversaire épineux*. This muscle, usually called, by English anatomists, *Multif'idus spi'ne*, and including all the *transversospinales*, is situate in the vertebral gutters, extending from the posterior part of the sacrum to the second cervical vertebra. It is thicker in the neck and loins than in the back and behind the sacrum. Its use is to straighten the vertebral column, and to impress upon it a slight rotary motion.

TRANSVERSALIS NASI, *Compressor naris*.

TRANSVERSARIUS, Transverse.

TRANSVERSE, *Transver'sus*, *Transversa'lis*, *Transversarius*, (F.) *Transverse*, *Transversaire*, (from *trans*, 'across,' and *vertere*, *versum*, 'to turn,') That which runs across: also, that which relates to the transverse processes of the vertebrae.

TRANSVERSE ARTERY OF THE FACE, arises from the temporal; passes transversely across the face, in front of the condyle of the lower jaw, and gives its branches to the different muscles of the cheek.

TRANSVERSE FISSURE, see Liver—*t*. Fossa, see Liver—*t*. Ligament of the Atlas, Annular ligament.

TRANSVERSE PERINEAL ARTERY, *Ure'throbulbar* (Ch.), is given off from the upper branch of the internal pudic; and passes inwards and forwards above the transversus perinæi muscles, as far as the bulb of the urethra, into which it dips, subdividing into numerous branches.

TRANSVERSE PROC'ESSES OF THE VERTEBRÆ, *Diapoph'yses* of Owen, (F.) *Apophyses transversæ des vertèbres*, are the bony eminences that jut out transversely and posteriorly from the sides of the vertebrae.

TRANSVERSE SUTURE, *Sutur'a transversa'lis*, runs across the face, and sinks into the orbit; joining the bones of the skull to those of the face, but with so many irregularities and interruptions, that it can scarcely be recognised as a suture.

TRANSVERSE, Transverse—*t*. *de l'Abdomen*, *Transversalis abdominis*—*t*. *de l'Oricula*, *Transversus auris*—*t*. *du Périnée*, *Transversus perinæi*.

TRANSVERSOSPINAL, *Semispinalis colli*, *Semispinalis dorsi*—*t*. *Spinalis*, *Transversalis*

dorsi—t. *Spinalis colli*, *Semispinalis colli*—t. *Spinalis dorsi*, *Semispinalis dorsi*.

TRANSVERSUS, Transverse.

TRANSVERSUS AURIS, *Transversus Auriculae*, (F.) *Transverse de l'Oreille*, (Ch.) *Transversal de la Conque*, is often wanting. It is formed of some transverse fibres, situate transversely, behind the projection of the helix which divides the concha into two portions.

TRANSVERSUS PEDIS, *Scandula rius*, (F.) *Transversal des Orteils*, *Métatarso-sous-phalangien transversal du premier orteil* (Chaussier). A muscle, which arises by fleshy slips from the heads of the metatarsal bones of the three outer toes. Its tendon is inserted into the base of the first phalanx of the great toe, being blended with that of the adductor pollicis.

TRANSVERSUS PERINÆI, *Levator Ani parvus*, (F.) *Ischio-perinéal*, (Ch.), *Transverse du Périnée*, is placed at the posterior part of the perineum. It is thin, triangular, and situate transversely. Its external extremity is attached to the ramus and tuberosity of the ischium: its inner extremity is confounded with its fellow of the opposite side, and with the anterior part of the sphincter ani and posterior part of the bulbocavernosus. The use of this muscle is,—to compress the urethra, and to support the rectum and bladder. To a fasciculus of this muscle, Santorini has given the name *Urethrae Elevator seu Ejaculator*.

TRANSVERSUS PERINÆI ALTER, *Prostat'icus inferior*. A small muscle which occasionally accompanies the last.

TRAPA NA'TANS, *Tribulus*, (F.) *Macre flottante*, *Noix ou Châtaigne d'Eau*. Ord. Onagraceæ. Sex. Syst. Tetrandria Monogynia. The plant which affords the *Nu'ces Aquat'icæ*, *Tribuli aquatici*, *Caltrops*. The nut is considered nutrient and demulcent, and is said to be useful in diarrhœa from abrasion of the bowels, and in calculus. A poultice is sometimes made of it to discuss hard and indolent tumours.

TRAPEZA, (*τραπέζα*.) Mensa.

TRAPEZE, Trapezium, Trapezus.

TRAPEZIUM OS, (*τραπέζιον*, 'a trapezium.') So called from its shape. *Os multan'gulum seu polygonum majus*, (F.) *Os trapezium*. The first bone of the second row of the carpus. It is articulated above with the scaphoides; below, with the first bone of the metacarpus; within, with the trapezoides and second metacarpal bone. Anteriorly, posteriorly, and externally, it gives attachment to ligaments.

TRAPEZIUS, *Cuculla'ris*, (F.) *Dorso-sus-acromion*, (Ch.), *Trapèze*, *Capuchon*. A muscle,

seated at the posterior part of the neck and shoulder, and at the upper part of the back. It has the shape of a trapezium; is attached, on the one hand, to the inner third of the upper curved line of the occipital bone; to the posterior cervical ligament; to the spinous process of the 7th cervical vertebra; and to those of all the dorsal vertebrae; and, on the other hand, to the spine of the scapula; the acromion and the outer third of the posterior margin of the clavicle. Its upper fibres are situate obliquely downwards and outwards; the transverse and inferior, obliquely outwards and inwards. This muscle elevates the shoulder; carries it backwards, or depresses it, according as its upper, middle, or lower fibres contract separately. It straightens the head, also, and inclines it to one side.

TRAPEZOID, (*τραπέζιον*, 'a trapezium,' and *oidos*, 'shape or resemblance,') Trapezoides.

TRAPZOID LIGAMENT, *Ligamentum trapezoid'es*. The anterior part of the coracoclavicular ligament. It is a very strong, fibrous fasciculus, which has the shape of a trapezium, and is situate obliquely between the acromion process and clavicle.

TRAPEZOÏDES OS, *Os multan'gulum seu polygonum minus seu pyramida'le carpi*. The second bone of the second row of the carpus—so called from its shape. It is smaller than the trapezium, on the inside of which it is situate. It is articulated, above, with the os naviculare; below, with the second metacarpal bone; on the outside, with the trapezium; and internally, with the magnum. Anteriorly and posteriorly, it affords attachment to ligaments.

TRAUBENKUR, Grape cure.

TRAULISMUS, (*τραυλισμος*), see Balbuties, and Blæstias.

TRAULOTES, (*τραυλότης*), see Blæstias.

TRAUMA, gen. *Trau'matis*, (*τραυμα*, gen. *τραυματος*.) Wound.

TRAUMAT'IC, *Traumat'icus*, *Vulnera'rius*, *Vul'nerary*, (F.) *Traumatique*, (*τραυματικός*, from *trauma*.) Anything relating to a wound.

TRAUMATICA, Detergents, Vulneraries.

TRAUMATICINE, (from *trauma*.) Liquor Guttaperchæ.

TRAUMATICUS, Traumatic, Vulnerary.

TRAUMATISM, *Traumat'ismus*. The condition of the organism occasioned by a grave wound.

TRAUMATOCOMI'UM, (*trauma*, and *κομω*, 'I take care of,') An hospital for the reception of the wounded.

TRAUMATOP'YRA, *Febris traumat'ica*, (*trauma*, and *πυρ*, 'fever,') A wound fever, or fever consecutive to a wound.

TRAVAIL, ([F.] *travailler*, 'to labor,') Parturition.

TRAVAIL D'ENFANT, 'child labor,' Parturition.

TRAVELLER'S JOY, *Clematis vitalba*.

TREACLE, (from *Σκαρα*.) (F.) *Thériaque*, (old E.) *Trincle*, Melasses, Theriaca—t. English, Teucrium chamædrys—t. Venice, Theriaca.

TREAD, *Cicatricula*—t. of the Cock, Mole-cule.

TREATMENT, (from [L] *tractare*, [F.] *traiter*, 'to treat,') Curatio.

TREATMENT, DRV. One in which the patient is almost wholly, if not wholly, restricted from the alimentary use of fluids; a valuable *modus medendi* in inflammatory and hemorrhagic conditions.

TREE, ELK, *Andromeda arborea*—t. of Heaven, *Ailanthus glandulosa*—t. of Life, *Thuya occidentalis*—t. Sour, *Andromeda arborea*.

TRÉFLE DE CASTOR, (from [L] *trifolium*, from *tri*, and *folium*, 'a leaf,') *Menyanthes trifoliata*—t. d'Eau, *Menyanthes trifoliata*—t. des Marais, *Menyanthes trifoliata*.

TREFOIL, *Hepatica triloba*—t. Bean, stinking, *Anagyris*—t. Marsh, *Menyanthes verna*—t. Shrubby, *Ptelea trifoliata*—t. Sour, *Oxalis acetosella*—t. Water, *Menyanthes trifoliata*.

TREMA, (*τρημα*.) Foramen, Vulva.

TREMBLEMENT, Tremor—t. *Métallique*, see Tremor.

TREMLES, Milksickness.

TREMBLING, Tremor—t. Fevers, Intermittent fever.

TREMELLA AURICULA JUDÆ, (from *tremere*, 'to tremble,') *Peziza auricula*.

TRE'MOR, *Trepidat'io*, *Syn'clonus Tre'mor*, *Tro'mospas'mus*, *Tro'mus*, *Trem'bling*, (F.) *Tremblement*, (from *tremere*, 'to tremble,') Involuntary agitation of the body, or of some part of it, without any obstacle to voluntary motion. It depends, generally, upon debility of the muscular, or of the nervous system; and, hence, is observed in convalescence, and in typhoid affections. It occurs, also, in old people; in hard drinkers; workers in mercury, lead, &c.: in the last case, it is called by the French, *Tremblement*.

métallique; Tremor metallicus. Tremor seems to resemble paralysis more than convulsion. See Paralysis agitata, and Pavitatio.

TREMOR CORDIS, Hippus, Palpitation—t. Metallicus, see Tremor—t. Potatorium, see Delirium tremens—t. Purring, *Frémissement catinaire.*

TREPAN', *Trep'anium*, (*τρῑπανον*) *Try'panon*, *Trip'anon*, *Tereb'ila*, *Peretor'ium*, *Peretor'ium*, *Sar'va versat'ilis*, *Cham'icis*, *Cham'ic'ion*, *Ter'ebra*, *Ter'etron*, *Ter'etrium*, *Vertib'ulum*, (F.) *Trep'an*, (from *τρῑνω*, 'to turn,' or *τρῑνω*, 'to perforate,') A surgical instrument resembling a wimble, and worked in the same manner. It is used for removing portions of bone, and particularly of the bones of the skull. The term *trep'an* is given more particularly to the part of the instrument that makes the perforation. The handle is so constructed as to receive different bits,—as the crown, the *Trep'an perforatif* or *Perite'ri'on*, the *T. exfoliatif* or *Desquamato'rium trep'anum*, and the *T. adapt'ita*. 'To *trep'an*' is to use the trepan.

In Great Britain and the United States this instrument has been superseded by the trephine, which is more manageable.

TREPANATIO, Trepanning.

TRÉPANATION, Trepanning.

TREPANG, *Biche de Mer.*

TREPANNING, *Terebra'tio*, *Anatre'sis*, *Cephalotryp'e'sis*, *Craniotryp'e'sis*, *Trepana'tio*, *Tryp'e'sis*, *Tre'sis*, *Pri'sis*, *Pria'mus*, *Fora'tio*, (F.) *Trepanation*. The methodical application of the trepan.

TREPANUM, Trepan.

TREPHINE', (from *τρῑνω*, 'to turn,') The instrument which has replaced the trepan in some countries. It consists of a simple, cylindrical saw; with a handle placed transversely like that of a gimlet, and from the centre of the circle, described by the teeth of the saw, a sharp little perforator projects, called the *Cent'trepin*. The centrepin is capable of being removed, at the surgeon's option, by means of a key for the purpose. It is used to fix the instrument until the teeth of the saw have made a circular groove, sufficiently deep for it to work steadily. The pin must then be removed. Sometimes the pin is made to slide up and down, and to be fixed in any position, by means of a screw. See Trepan. 'To *trephine*' is to use the trephine.

TREPIDATIO, (*trepidare*, *trepidatum*, 'to be agitated,') Tremor—t. Cordis, Cardiotromus.

TRÉPIED (tripod), *CELLAQUE*, see Cœliac artery—t. Vital, Tripod, vital.

TREPON'DO, (*tres*, 'three,' and *pondus*, 'a pound,') A weight of three pounds.

TRE'SIS, (*τρῑσις*, from *τρῑνω*, 'I bore,') Properly, a perforation. A wound. A forcible solution of continuity in a soft part, commencing externally. A genus in the nosology of Good.

TRESIS, Trepanning, Wound—t. Causis, Burn—t. Punctura, Puncture—t. Vulus, Wound—t. Vulus laceratum, see Laceration—t. Vulus simplex, Cut.

TRESSAILLEMENT, (*trans*, 'beyond,' and *salto*, 'to leap,') Shuddering.

TRESSO'RIA, (low Latin,) *Episi'on*. A term formerly applied to the hair covering the mons veneris, the absence of which has, by the vulgar, been esteemed a matter of reproach.

TRI, (*τρῑ*), in composition, three.

TRIACLE (old E.), Melasses.

TRIAL BY BIER'RIGHT. An ordeal, founded on a superstition formerly prevalent, that the wounds of a murdered person would bleed afresh if the body were touched ever so lightly, in any part, by the murderer.

TRIANGLE MÉDULLAIRE, (*tri*, and *angulus*, 'an angle,') Fornix.

TRIANGLE VESTICAL, see U'tinary Bladder. TRIANGLES. CERVICAL, see Cervical.

TRIANGULAR, *Trigo'nus*, *Triangula'ris*, (F.) *Triangula'ire*. That which has three angles.

TRIANGULAR or INFRAPUBIAN LIGAMENT. *Ligamentum triangula're seu in'frapubia'nium*, is a ligamentous fascia with short fibres, situate transversely beneath the symphysis pubis, which it strengthens. Also, a ligament extending from the coracoid to the acromial process of the scapula.

TRIANGULARIS, *Nasori anguli oris*—t. Coccygis, Coccygeus—t. Nasi, Compressor naris.

TRIANGULA'RI'S STERNI, *Sternocostal*, (Ch.), *Pectora'lis inter'nus*. A muscle situate at the anterior and inner part of the chest, behind the cartilages of the ribs. It has the shape of a lengthened triangle, the base of which is directed downwards. It is attached to the posterior, lateral, and inferior part of the sternum, whence its fibres pass upwards and outwards, and terminate by as many distinct digitations at the cartilages of the 3d, 4th, 5th, and 6th true ribs. This muscle depresses the ribs and contributes to expiration.

TRIBALUS, (*τρίβαλος*.) *Trapa natans*.

TRIBAS, gen. *Tri'badis*, (*τρίβας*) (F.) *Tribade*, (from *τρίβας*, 'I rub,') *Fric'trix*, *Frica'trix*, *Futu'trix*, *Mas'cula*, *Subagita'trix*, *Subigita'trix*, *Eteris'tria*. A female whose clitoris is so large as to cause her to be regarded as a hermaphrodite. One who acts as a male with another female.

TRIBASIC, (from *tri*, and *basia*), see Salt.

TRIBULI AQUATICI, (pl. of *tribulus*, from *τρίβαλος*.) see *Trapa natans*.

TRIBUTUM LUNARE, 'lunar or monthly tribute,' Menses—t. Menstruum, Menses.

TRICE, (from *τρίς*, gen. *τρίχης*, 'a hair,') *Plica*—t. Incuborum, *Plica*—t. Seroforum, *Plica*.

TRICAGO, *Teucrium chamaedrys*.

TRICAUDALIS, (*tri*, and *cauda*, 'a tail,') *Retrahens auris*.

TRICEPH'ALUS, (*tri*, and *κεφαλη*, 'head,') A monster with three heads.

TRICEPS, gen. *Tricip'itis*, (*tri*, and *caput*, 'head,') A name given to muscles that have three fleshy bundles at one of their extremities.

TRICEPS ADDUC'TOR FEM'ORIS. Under this appellation are comprehended three distinct muscles; *Adductor brevis*, *A. longus*, and *A. magnus*.

TRICEPS AURIS, *Retrahens auris*—t. *Brachialis*, *Triceps extensor cubiti*.

TRICEPS CRURAL, *Triceps cruris*.

TRICEPS CRU'RI'S, *Femora'lis*, (F.) *Trifémoro-rotulien*, (Ch.) *Crural*, *Trifémoro-tibi-rotulien*, *Triceps crural* or *de la cuisse*, is situate at the anterior, inner, and outer part of the thigh. It is formed of three bundles at its superior extremity, viz.: 1. A middle fasciculus (the *crural'is* of most anatomists). 2. An external fasciculus (the *vastus externus*); and 3. An inner fasciculus (the *vastus internus*). It is attached, above, to the anterior, inner, and outer surfaces of the femur, and to the two edges of the linea aspera, from the base of the trochanter to within four fingers' breadth of the knee. Below, it terminates by a large tendon, which is attached to the base and edges of the patella, as well as to the inner and outer tuberosities of the tibia. The triceps extends the leg on the thigh, and conversely.

TRICEPS DE LA CUISSE, *Triceps cruris*.

TRICEPS EXTEN'SOR CU'BITI, *Tri'ceps brachia'lis*, (F.) *Scapulo-huméro-olecrani'en*, (Ch.) *Triceps brachial*. This muscle, which occupies all the posterior part of the os humeri, is described as two muscles by Douglas, and as three by Winslow. It arises by three heads. The first or long head,—long head of the *biceps externus* of Douglas, *Anconeus major* of Winslow,—arises from the anterior extremity of the inferior costa of the

scapula, near its neck, and below the origin of the upper muscle. The second head, or short head of the *longus externus* of Douglas, *Assomus externus* of Winslow, arises from the upper and outer part of the os humeri, at the base of the great tuberosity; and the third head, — *Brachialis externus* of Douglas, *Assomus internus* of Winslow, — the shortest of the three, arises from the back part of the os humeri, behind the flat tendon of the latissimus dorsi. These three portions unite about the middle of the arm, so as to form one thick and powerful muscle, which is inserted at the upper part of the olecranon. It extends the forearm on the arm, and conversely.

TRICEPS MAGNIS. *Adductor magnus* — t. Minor, *Adductor longus femoris* — t. Secundus, *Adductor brevis*.

TRICHE INCUBORUM, (*ῥαχίς*, gen. *ῥαχός*, 'a hair,' and *incubis*, 'the Fawns,' [?]) *Plica*.

TRICHANGIA, (*tricho*, and *αγγία*, 'vessels,') *Capillary vessels*.

TRICHANGIECTASIA, *Angiectasia*, *Trichangia*, (*tricho*, *αγγίον*, 'a vessel,' and *εκτασις*, 'dilatation,') Morbid dilatation of the capillary vessels.

TRICHAUXË, (*tricho*, and *αύξη*, 'increase,') Increase in the quantity and size of the hair, without change of texture.

TRICHERA ARVENSIS, (from *ῥαχίς*, gen. *ῥαχός*, 'hair,') *Scabiosa*.

TRICHIA, (from *ῥαχίς*, gen. *ῥαχός*, 'hair,') Entropion.

TRICHIASIS, *Trichiasis*, *Trichiasis*, (from *ῥαχίς*, gen. *ῥαχός*, 'hair,') *Mal'ina pil'aris*. This name has been given to several affections. 1. To a disease of the kidneys or bladder, in which filamentous substances, resembling hairs, are passed in the urine. This is also called *Pilinictio*. It must be regarded as a variety of *Calculus vesicæ*. 2. To a painful swelling of the breasts, in child-bed women, when the milk is excreted with difficulty: a disease attributed, by Aristotle, to a hair swallowed by accident, and carried to the breasts through the circulation; and 3dly, To inversion of the eyelashes. See Entropion.

TRICHIASIS LACTEA, *Infarctus mammae lacteus*.

TRICHILIA EMETICA, *Eionja*. A tree, *Order*, *Meliaceæ*, common on the mountains of Yemen and in Senegal. The fruit is used by the Arabs as an emetic. The ripe seeds are formed with sesamum oil into an ointment for the itch.

TRICHILIA TRIPOLITANA a bush of Curaçoa, a decoction of the roots of which is used to promote abortion.

TRICHINA SPIRALIS, (from *ῥαχίς*, gen. *ῥαχός*, 'hair,' from its small size,) *Pseudolium trichina*. A small species of entozoa, *Fam.* *Nematodea*, discovered in the muscles of voluntary motion. *Trichinae* consist of very minute cysts, of an oblong figure, having considerable resemblance, in size and colour, to young pediculi, and are somewhat allied to the cels found in paste and vinegar. Küchenmeister regarded them as the primitive state of *Trichocephalus dispar*; but this has not been confirmed. When taken with the food, they multiply in the intestines and then migrate to the muscles. No symptoms have, as yet, been observed to be pathognomonic of their presence in the muscles. It would appear, however, that they may be the cause of the morbid phenomena caused by eating the flesh of animals in which they existed, and may be the cause of sausage poisoning, see *Allantotoxium*, and of the poisoning produced by eating pork and ham. The disease thus induced has been called *Trichinosis* or *Trichina disease*.

TRICHINIASIS, see *Trichina spiralis*.

TRICHIOSIS, (from *ῥαχίς*, gen. *ῥαχός*, 'hair,') *Trichosis* — t. *Distrix*, *Distrix*.

TRICHISMUS, *Capillismus*, *Fim'ca pil'aris*. A capillary fracture, fissure, or crack.

TRICHIURIA, (*tricho*, and *ουρα*, 'tail,') *Trichocephalus*.

TRICHO, (*ῥαχίς*, gen. *ῥαχός*, 'a hair,') in composition, hair.

TRICHOCEPHALE DE L'HOMME, *Trichocephalus*.

TRICHOCEPHALUS, *Trichocephalus dispar*, (*tricho*, and *κεφαλή*, 'head,') *Trichina*, *T. pulgaris seu intestinalis*, *Ascaris tricharia*, *Tricharia*, *Trichocephalus seu Mustip'ides humilis*, *Long thread worm*, (F.) *Trichocephalus de l'homme*. A worm, from an inch and a half to two inches long; the head acute; the body spirally invaginated in the male, almost straight in the female. The *trichocephalus dispar* generally inhabits the cæcum and colon, and is rarely met with in the small intestines. It gives rise to no peculiar symptoms, and requires the usual anthelmintic treatment.

TRICHOCIRSUS, (*tricho*, and *κυρς*, 'a dilated vessel,') *Trichangiectasia*.

TRICHOG'ENOUS, (*tricho*, and *γενος*, 'I generate,') Hair-forming; as a '*trichogenous*' remedy, — one that promotes the growth of hair.

TRICHO LABIS, (*trichos*, and *λαβω*, 'to seize hold of,') *Madisterium*.

TRICHO LABIUM, (*trichos*, and *λαβω*, 'I seize hold of,') *Madisterium*.

TRICHOLOGIA, (*tricho*, and *λογία*, 'I collect or pluck,') *Caryologia*. See *Trichology*.

TRICHOLOGY, *Trichologia*, (*tricho*, and *λογος*, 'a description,') A treatise on the hair.

TRICHOMA, gen. *Trichomata*, (from *ῥαχίς*, gen. *ῥαχός*, 'hair,') *Capillamentum*, *Plica*.

TRICHOMANES, (*tricho*, and *μανος*, 'thin,') *Asplenium trichomanoides* — t. *Crenata*, *Asplenium trichomanoides*.

TRICHOMAPHYTE, *Trichomophyton*, (*trichoma*, 'plica,' and *φυτον*, 'a vegetable,') A cryptogamous growth, found in plica, and said, by some, to be its cause. *Mycoderma plicæ*.

TRICHOMATION, (*τρυχωματισμος*,) *Capillamentum*.

TRICHOMATOSE HAIR, (from *trichoma*,) *Plica*.

TRICHOM'ONAS, *Trichomonad*, (*tricho*, and *μοναδ*,) An animalcule found by M. Donné in the mucus of the vagina, where cleanliness had not been attended to, and to which he has given the name *Trichomonas vaginitis*. By some it is considered to be nothing more than ciliated epithelium separated from the uterus.

TRICHOMYCES DECALVANS, (*tricho*, and *μυκός*, 'fungus,') see *Porrigo decalvans* — t. *Tonsurans*, see *Porrigo decalvans*.

TRICHONOSIS, *Porrigo* — t. *Furfuracea*, *Porrigo decalvans* — t. *Plica*, *Plica* — t. *Pollitosis*, *Pollitosis*.

TRICHON'OSUS, (*tricho*, and *νοσος*, 'disease,') A disease of the hair.

TRICHOPHY'IA, *Trichophytica*, (*tricho*, and *φυειν*, 'to grow,') Remedies that promote the growth of the hair.

TRICHOPHYTON, (*tricho*, and *φυτον*, 'a vegetable,') see *Plica* — t. *Decalvans*, see *Porrigo decalvans* — t. *Sporuloides*, see *Fungus* — t. *Tonsurans*, see *Fungus*, and *Porrigo decalvans*.

TRICHOSANTHES AMARA, (*tricho*, and *ανθη*, 'a flower,' from the ciliated corollæ,) *Order*, *Cucurbitaceæ*; a plant of India; the seeds are bitter and astringent, sometimes emetic. In Jamaica, it is considered to be a poison, and is used for killing rats.

TRICHOSANTHES CODONATA. The root is used by the natives of India as a substitute for *Calumba root*.

TRICHOSANTHES CUCUMERINA. The fruit is considered in India to be anthelmintic.

TRICHOSANTHES INCISA, *Gashed Trichosanthos*, of the forests of India. The fruit is esteemed to be poisonous. Pounded and intimately blended with cocoa nut oil, it is issued in India as an application to offensive sores of the ears, and to ozena.

TRICHOSANTHES VILLOSA, of Java. The fruit acts like colocynth.

TRICHOSCHISIS, (*tricho*, and σχίζω, 'to split,') Schizotrichia.

TRICHOSIS, *Pila'rē ma'lum*, (from *Spiz*, gen. *τριχος*, 'hair,' and *osis*.) Morbid organization or deficiency of hair. A genus in the system of Good.

Trichosis, Entropion, Trichiasis.—*t. Area*, Porrigo decalvans.—*t. Athrix*, Alopecia.—*t. Distrix*, Distrix.—*t. Furfuracea*, Porrigo furfurans.—*t. Hirsuties*, Hirsuties.—*t. Plica*, Plica.—*t. Poliosis*, Poliosis.—*t. Setosa*, Hystriacis.

TRICHOTON, (*τριχωτος*, 'covered with hair,') Scalp.

TRICHURIS, (*tricho*, and *ουρα*, 'a tail,') Trichocephalus.

TRICLISPERMA GRANDIFLORA, (*Spiz*, 'hair,' and *σπέρμα*, 'seed,') Polygala paucifolia.

TRICUSPID, *Tricus'pis*, *Tricuspidal'is*, *Tricuspidatus*, (*tri*, and *cuspis*, gen. *cuspidis*, 'a cusp, a point,') That which has three points.

TRICUSPID VALVES, *Val'vulæ Tricus'pidæ* seu *triglo'chinæ* seu *tricusul'cæ*, are three triangular valvular duplicatures, formed by the inner membrane of the right cavities of the heart, around the orifice by which the auricle communicates with the ventricle. The elongated and pointed top of each of these valves is continuous with the chordæ tendinæ, which set out from the columnæ carneæ of the heart. These valves are depressed to allow the blood to flow from the auricle into the ventricle; and are raised to prevent its reflux into the auricle.

TRICUSPIS, gen. *Tricus'pidis*, Tricuspid.

TRIENCEPHALUS, *Triceph'alus*, (F.) *Trien'céphale*, *Triocéphale*, (*tri*, and *εγκεφαλος*, 'the encephalon,') A monster wanting three cephalic organs of sense,—smell, hearing, and vision.

TRIENS, *Un'ciæ quatuor*. (From *tri*, 'three,') The third part of a pound. Four ounces, Troy.

TRIFACIAL, Trigemini.

TRIFÉMORO-ROTULIEN, *Triceps'eruris*.—*Tibi-rotulien*, *Triceps' cruris*.

TRIFOLIUM ACETOSUM, (*tri*, and *folium*, 'a leaf,') *Oxalis acetosella*.—*t. Aquaticum*, *Menyanthes trifoliata*.—*t. Aureum*, *Hepatica triloba*.—*t. Cervinum*, *Eupatorium cannabinum*.—*t. Fibrinum*, *Menyanthes trifoliata*.—*t. Hepaticum*, *Hepatica triloba*.

TRIFO'LIUM MELLO'TUS OFFICINA'LIS, *T. offic'ina'lē* seu *odora'tum* seu *cuballi'nium*; (from *tri*, and *folium*, 'a leaf,') *Melilo'tus*, *Melilo'tus cit'rina*, *Lo'tus sylves'tris*, *Ser'tula Camp'na*, *Coro'na re'gia*, *Common mel'ilot*, (F.) *Mé'ilot*. Ord. Leguminosæ. Ser. Syst. Diadelphia Decandria. The infusion or distilled water of this plant has been employed in cases of ophthalmia. It has also been used in fomentations, gylsters, &c.

TRIFOLIUM OFFICINALE, *T. melilotus officinalis*.—*t. Paludosum*, *Menyanthes trifoliata*.—*t. Palustre*, *Menyanthes trifoliata*.

TRIGAS'TRIC, *Trigas'tricus*, *Triven'ter*, (*tri*, and *γαστρ*, 'belly,') That which has three bellies. An epithet for muscles so circumstanced.

TRIGEM'INI, *Tergem'ini*, (*tri*, and *gem'inus*, 'double,' 'threefold,' 'triplets,') The fifth pair of nerves, *Ner'vi d'ic'ei* seu *gustato'rii*, *Ner'vus quin'tus* seu *trémel'tus* seu *mix'tus* seu *sympathet'icus* *me'dius* seu *sympath'icus* *me'dius* seu *anon'ymus* seu *innominat'us*, *Par trigem'inum* seu *quin'tum* *ner'vorum* *cerebra'lium* seu *tri'tum* *funiculo'rum*, *Tri'facial*, (F.) *Nerf gustatif innominé*, *Nerf*

à trois cordes, *Nerf trijumeau*. This nerve arises below the tuber annulare by one large posterior root from the corpora restiformia, and by two small anterior roots from the corpora olivaria, whence, proceeding forwards to the side of the sella Turcica, the filaments composing the large root form a ganglion—the Gasserian. Emerging from this ganglion, the filaments, of which the large root consists, form, with those constituting the small roots, and which had not passed through the ganglion, two flattened trunks, each of which is soon divided into three branches, in such a manner, however, that the first or ophthalmic branch, and the second or upper maxillary, are composed exclusively of the filaments that arose from the corpora restiformia and passed through the Gasserian ganglion; whilst the third or lower maxillary branch is composed in part of those filaments, and in part of those that arose from the corpora olivaria, and passed on one side of the ganglion without going through it. The whole of the first and second branches, and a part of the third are sensiferous nerves, whilst the remaining part of the third—the manducatory or mas'ticatory—is wholly motiferous, and passes to the pterygoid and other muscles concerned in mastication.

TRIGONE CÉRÉBRAL, (*τριγωνον*, 'a triangle,') Fornix.—*t. Vésical*, see Urinary Bladder.

TRIGONELLA FENUM, *T. Fé'nium* *Græ'cum*, (dim. of *τριγωνον*, 'a triangle,' from the shape of its flower.) *Fén'ugreek*, *Fæ'nugreek*, *Fæ'nium* *Græ'cum*, *Bu'ceras*, *B. Fé'nium* *Græ'cum*, *Car'phos*, *Ægoc'eras*, (F.) *Fénugrec*, *Sénegrain*. Ord. Leguminosæ. Ser. Syst. Diadelphia Decandria. A native of the south of France. The seeds have a strong, disagreeable smell, and an unctuous, farinaceous taste; accompanied by a slight bitterness. They are chiefly used in poultices.

TRIGONOS, (*τριγωνος*,) Triangular.

TRIGONUM OLFACTORIUM, 'olfactory triangle,' see Mammary eminence.

TRIGONUM VESICÆ, see Urinary bladder.

TRIGONUS LIUTAUDI, (*τριγωνος*,) see Urinary bladder.—*t. Vesicæ*, see Urinary bladder.

TRIJUMEAUX, (*tri*, and *jumeau*, 'a twin,') Trigemini.

TRILLIUM LATIFOLIUM, (from *trilix*, 'triple,' its parts being chiefly in threes.) *Broad-leaf Bethroot*, *Beth'root*, *Rattle-snake root*, *Wake'robin*, *Cough'root*, *Indian Balm*, *Ground Lil'y*, *Jews' Hurd*, *Indian Sham'rock*, *Pariscourt*, *True-love*. Ord. Trilliaceæ. Ser. Syst. Hexandria Trigynia. A plant peculiar to North America, which blossoms in the spring. The root is astringent.

There are many species of Trillium, all of which possess the same properties.

TRINTANELLE MALHERBE, *Passerina tartonraira*.

TRIOCÉPHALE, *Trioccephalus*.

TRIOCEPHALIA, Aprosopia, see Triencephalus.

TRIOCEPHALUS, Triencephalus.

TRIOR'CHIS, (*tri*, and *ορχις*, 'a testicle,') One who has three testicles.

TRIOSTOSPERMUM, Triosteum.

TRIOS'TEUM, *T. Perfoliatum* seu *ma'jus*, (*tri*, and *οστέον*, 'a bone,' in allusion to the three bony nuts;) *Triosteospermum*, *Bastard Ipecacuan'ha*, *Fever root*, *Fever'wort*, *Tink'er's weed*, *Horse gin'seng* or *gent'ian*, *Ip'ecac*, *Wild Cof'fee*, *White gin'seng*, *Gent'ian*, *White Gent'ian*, *Sweet Bit'ter*. Indigenous; Ord. Caprifoliaceæ. In very large doses, *Triosteum* (Ph. U. S.) proves emetic. It is a good cathartic, in doses of twenty or thirty grains. It sometimes operates as a diuretic.

TRIOSTEUM AUGUSTIFOLIUM, *T. minus*, has similar virtues.

TRIPLE DE ROCHE, 'rock tripe.' Different lichens—*Rock l'ichens*, *Stone moss*—of the species

Gyrophora, inhabiting the Arctic regions of America, bear this name. They are eaten as food by the hunters.

TRIPHARMACUM, (τρίφάρμακον, from *tri*, and *pharmakon*, 'a medicine.') A medicine composed of three ingredients.

TRIPLET, (*triplex*, 'triple.') One of three children born at the same accouchement or gestation. Also, relating to triplets—as a 'triplet conception.'

TRIPLET MONSTERS, *Mim'stra trip'lica* seu *trigem'ina*, are such as have parts of the body tripled. They are rare.

TRIPPOD, VITAL, (τρίπους, from *tri*, and *pous*, gen. *πόδος*, 'foot.') (F.) *Triépied vital*. The heart, lungs, and brain have been so called, as their united action is necessary for the maintenance of life.

TRIPSIS, (τρίψις,) Contusio, Trituration.

TRIPUDIATIO SPASTICA, (*tripudiare*, *tripudiatum*, 'to leap, to dance.') Chorea.

TRIPUS, ('tripod,') CÆLIACUS seu HALLERII, see Cæliac artery.

TRIQUE-MADAME, *Sedum album*.

TRIQUETRA OSSA, (*triquetrus*, 'three cornered,') Wormiana ossa.

TRIQUESTRUM, (OS,) Cuneiform (bone.)

TRISMUS, (τρίσμος, from *τρί*, 'I gnash,') *Entasia* seu *To'vus Trismus*, *Capsit'rum*, *Masticatory spasm of the face*, *Lock'ed Jaw*, *Spas'mus seu Ri'gor seu Tr'ismus maxill'æ inferio'ris*, *Gnathospas'mus*, *Stomatospas'mus*, (F.) *Mal de Mchoire*. Spastic closure of the under jaw, a partial tetanus, arising from erethism of the masticatory or motor branch of the fifth pair of nerves. Cullen makes two varieties—*T. nascentium* seu *neonatorum*, *Nineday fits*, (F.) *Sarrête*, attacking infants during two weeks from birth; and *T. traumaticus*, originating from a wound.

In the West Indies and in South America, *Trismus nascentium* is called *Seven days' disease*, (F.) *Mal de sept Jours*.

TRISMUS CAPISTRATUS, *Brédisure*—t. Catarrhalis maxillaris, Neuralgia maxillaris—t. Clonicus, Neuralgia faciei—t. Cynicus, see Canine laugh—t. Diastrophe Sauvagesii, Paralysis, Bell's—t. Dolorificus, Neuralgia faciei—t. Maxillaris, Neuralgia faciei—t. Nascentium, see Trismus—t. Neonatorum, see Trismus—t. Sardonius, Risus sardonius.

TRISPERMUM, (*tri*, and *σπερμα*, 'a grain or seed.') An ancient cataplasm, consisting of three seeds—*cumin*, *bay*, and *smallege*.

TRISPLANCHNIA, (*tri*, and *σπλᾶγχνον*, 'viscus,') Cholera.

TRISPLANCHNIC, *Trisplanch'nicus*, (F.) *Trisplanchnique*. That which relates to the viscera of the great cavities.

TRISPLANCHNIC NERVE, *Nerv'vus trisplanch'nicus* seu *sympathetic'us mag'nus* seu *sympath'icus seu ganglion'icus seu vasomotor'ius*, (F.) *Nerf sympathique ou grand sympathique*, *Great sympathetic*, *Intercostal*, *Ganglion'ic* or *Ver'tebral nerve*, *Nerve of organic life*;—named by Bichat the organic nervous system, in opposition to the other nervous parts, called the animal nervous system. Gall terms it, the *nerv'vus syst'em of the automat'ic functions*. It has also been called the *ganglion'ic nerv'vus syst'em*. It is termed *Trisplanch'nic* because it distributes its branches to the organs in the three great splanchnic cavities—the chest, abdomen, and pelvis. It is composed of a series of ganglia, united by intermediate branches, and passing along the side of the spine to the lowest part of the trunk; communicating, by a branch, with each of the thirty pairs of spinal nerves and several of the cerebral nerves; and detaching nerves from its several ganglia, which accompany the arteries, and are distributed, particularly, to the organs of involuntary func-

tions. At its upper part it is concealed in the canalis caroticus and cavernous sinus, and is joined there by a branch of the 6th pair of nerves, and by a twig from the Vidian nerve of the 5th pair. In passing down by the side of the spine, it forms three ganglia in the neck:—the *superior*, *middle*, (*thy'roid ganglion*), and *inferior*, (*vertebral ganglion*); 12 in the back—the *thorac'ic*; 5 in the loins—the *lum'bar*; and three or four *sac'ral*. When it reaches the coccyx it unites with its fellow from the opposite side, forming a small ganglion—the *coccyge'al*, *gan'gion im'par*, *az'ygous gan'gion*. Meckel divides it into two portions—1. The *central ganglions* or *central part*, composed of the numerous ganglionic plexuses in the abdomen: the centre of which is the *semilunar* and *solar plexuses*. 2. The *external ganglions*, *limitrophes* or *external part*, which comprise the series of ganglions situate at the sides of the spine, from the base of the cranium to the sacrum. The great sympathetic seems to form a distinct nervous system, destined for the organs of involuntary motion; for although communicating with both brain and spinal marrow, it does not seem to be immediately under the influence of either. Its special functions are not yet, however, well understood, and are the subject—as they have always been—of much interesting investigation.

TRISPLANCHNITIS, (*trisplanchnic*, and *itis*,) Cholera (spasmodic).

TRISSAGO, gen. *Trissag'inis*, *Teucrium chamaedrys*—t. *Palustris*, *Teucrium scordium*.

TRISTEMANIA, (*tristis*, 'sad,' and *mania*,) Melancholy.

TRISTERNAL. Bécclard has given this name to the third bony portion of the sternum;—that which corresponds to the third intercostal space.

TRISTITIA, (from *tristis*, 'sad,') Athymia, Mœror.

TRITÉOPHYA AMERICANA, Fever, yellow.

TRITÉOPHYES, *Tritéoph'ya*, (τρίτεως, from *tritus*, and *φυν*, 'form,') A fever whose type approaches the tertian. Sauvages gave the name *Tritéophya Vratislavien'sis* to an epidemic which broke out in the Prussian army at Breslaw, in the middle of the last century.

TRITÉUS, (τρίτεως,) Tertian fever.

TRITICUM ÆSTIVUM, (from *terere*, *tritum*, 'to thrash,') *T. cereal'è* seu *Cevallos* seu *compac'tum* seu *erind'ecum* seu *Gaertnerianum* seu *Linnæum* seu *pruin'osum*, *Sum'merheat*; and *TRIT'ICUM HYBERN'UM* seu *glau'ecum* seu *Muticum* seu *Sic'ulum*, *Win'terheat*, (F.) *Froment*. Ord. Gramineæ. Sex. Syst. Triandria Digynia. From these plants our wheat flour is obtained, by grinding the seeds—Wheat, *Pyrus*, *Fromentum*—and to them we are indebted for our bread and pastry. The nutritive properties of wheat flour are so well known as not to need comment.

Bread, *Pa'nis*, (F.) *Pain*, mixed with milk, constitutes the ordinary emollient poultice;—and the crumbs of bread, *M'œa pa'nis*, are a common excipient for pills and boluses.

TRITICUM ARVENSE, *T. repens*—t. *Cereale*, *T. æstivum*—t. *Cevallos*, *T. æstivum*—t. *Compactum*, *T. æstivum*—t. *Erinaceum*, *Triticum æstivum*—t. *Faginum*, *Polygonum fagopyrum*—t. *Gaertnerianum*, *T. æstivum*—t. *Glaucum*, *T. hybernum*, and *T. repens*—t. *Hybernum*, see *T. æstivum*—t. *Linnæanum*, *T. æstivum*—t. *Muticum*, *T. hybernum*—t. *Pruinosum*, *T. æstivum*.

TRIT'ICUM RE'PENS, *T. arven'sè* seu *glau'ecum*, *Agropyrum larissimum*, *Brom'us gl'aber*, *Gr'amen* seu *Gr'amen can'dinum* seu *Dioscor'idis* seu *re'pens*, *Lol'ium radi'cè repen'tè*, *Grass*, *Day'grass*, *Com'grass*, *Quick'grass*, *Trit'ch'grass*, (Se.) *Ran'nachs*, (F.) *Chiaudent*. The roots are sweet, and

possess aperient properties. They are said also to be diuretic.

TRITICUM SICULUM, T. hybernium.

TRITIO, (terere, tritium, 'to rub, to grind,') Contritio, Trituration.

TRITON PALUSTRIS, (from *Triton*, 'the sea deity,') see Ectozoon.

TRITORIUM, (from *terere*, *tritum*, 'to wear away,') Infundibulum.

TRITURA, Contritio, Trituration.

TRITURATIO, *Trituratio*, *Tritura*, *Tritus*, *Tritio*, *Tritipis*, (terere, tritum, 'to rub, to grind,') The act of reducing a substance to powder. See Contritio.

TRITUS, gen. *Tritus*, Contritio, Trituration.

TRIVENTER, (*tri*, and *enter*, 'belly,') Tri-gastric.

TROCAR, *Trochar*, *A'cus paracent'ica* seu *paracent'ica* seu *tri'quetra vulg'is*, *Paracent'ica*, *Paracent'ica*. (Corrupted from the (F.) *Troisquarts* or *Trocart*, 'three quarters,') so called from the triangular shape of the point—*Tri'gu'rum*. An instrument used for evacuating fluids from cavities, particularly in ascites, hydrocele, &c. A trocar consists of a perforator or stylet and a canula, which is so adapted to the perforator, that when the puncture is made, both enter the wound with facility; after which, the stylet being withdrawn, the canula remains in the wound, and affords the fluid a ready passage outwards.

TROCART, Trocar.

TROCH, (from τροχός, 'a wheel,') Trochiscus.

TROCHAN'TER, gen. *Trochant'ria*, *Trochan'ter*, *Rota'tor*, (from τροχάω, 'to run or roll,') Anatomists have given the names *great* and *little trochanter* to two processes at the upper extremity of the femur. The great *trochan'ter*, *glu'tus*, is the one on the outside; the *less* is lower down and more internally. Both afford attachment to rotator muscles of the thigh, and hence their name. *Chaussier*, by the word *trochanter*, means the larger process; the smaller he calls *trochantin*.

TROCHANTER'IAN, *Trochanteria'nus*, *Trochant'ian*. That which belongs or relates to the great trochanter.

TROCHANTIN'IAN, *Trochantin'ian*, *Trochantinia'na*. That which belongs or relates to the trochantin or less trochanter.

TROCHAR, Trocar.

TROCHES (pl. of troch) OF CARBONATE OF LIME, Trochisci carbonatis calcis—t. of Chalk, Trochisci carbonatis calcis—t. of Cubels, Trochisci cubelæ—t. Escharotic, Trochisci escharotici—t. of Ginger, Trochisci zingiberis—t. Gum, Trochisci gummosi—t. of Gum Arabic, Trochisci acacia—t. of Ipecacuanha, Trochisci ipecacuanhæ—t. of Subcarbonate of Iron, Trochisci Ferri subcarbonatis—t. of Lactucarium, Trochisci lactucarii—t. Liqueurice, Trochisci glycyrrhizæ—t. Liqueurice with opium, Trochisci glycyrrhizæ cum opio—t. of Magnesia, Trochisci magnesiæ—t. of Morphina, Trochisci morphiæ—t. of Morphina and ipecacuanha, Trochisci morphiæ et ipecacuanhæ—t. Nitre, Trochisci nitratis potassæ—t. of Peppermint, Trochisci menthæ piperitæ—t. of Tartaric acid, Trochisci acidi tartarici.

TROCHIA, (τροχία, 'the rut of a wheel,') Orbit.

TROCHILIA, (τροχίλια, 'a wheel,') Trochlea.

TROCHIN, (from τροχάω, 'to turn,') *Chaussier* has given this name to the smaller of the tuberosities at the upper extremity of the os-humeri; because it gives attachment to one of the rotator muscles of the arm—the subscapularis.

TROCHIN'IAN, *Trochin'ian*. That which belongs or relates to the trochin.

TROCHIS'CUS, (τροχίσκος, *Phtho'ia*, *Artis'eus*, *Col'icis*, *Col'icis*, *Phtho'icis*, *Rotula*, *Cyclid'ius*, dim. of τροχός, 'a wheel,') 'any thing circular,') A *troch* or round table, (F.) *Trochisque*. See Pas-

til, and *Tabella*. A solid medicine, prepared of powders, incorporated by means of mucilage, crumb of bread, juices of plants, &c. In French nomenclature, the *trochisque* differs from the *tablette*,—the former containing no sugar. This form of preparation is adapted for the purpose of allowing the medicines of which the troch is composed to dissolve slowly in the mouth, and to pass gradually into the stomach.

The Parisian Codex has two formulæ for troches for external use.

TROCHISCI ACA'CIÆ, *Troches of Gum Arabic*, (Acacia, $\tilde{\text{S}}\text{iv}$; Amyli, $\tilde{\text{S}}\text{ij}$; Sacerchar. Rij . Make into a mass with rose water, and form lozenges.—Ph. E.) To allay cough.

TROCHISCI AC'IDI TAN'SICI, *Tan'nin Loz'enges*, (Acid. tannic. gr. ceelx; Tinct. tolut. f $\tilde{\text{S}}\text{ss}$; Sacchar. purif. $\tilde{\text{S}}\text{xxv}$; Acacia pulv. $\tilde{\text{S}}\text{ij}$; Mucilag. Acacia, f $\tilde{\text{S}}\text{ij}$; Ag. destillat. bullient. f $\tilde{\text{S}}\text{ij}$. To be divided into 720 lozenges. Mix. Ph. B., 1864.) Each lozenge contains half a grain of tannic acid.

TROCHISCI AC'IDI TARTAR'ICI, *Troches of Tartar'ic Ac'id*, (Acid. Tartar. $\tilde{\text{S}}\text{ij}$; Sacchar. $\tilde{\text{S}}\text{viij}$; Ol. Limon. Mx . Form into troches with mucilage.—Ph. E.) Refrigerant and demulcent.

TROCHISCI BECHICI NIGRI, T. glycyrrhizæ glabrae.

TROCHISCI BISMUTHI, *Bis'muth Loz'enges*, (Bismuth. alb. gr. 1440; Magnes. carb. $\tilde{\text{S}}\text{iv}$; Calcin carb. princip. $\tilde{\text{S}}\text{vj}$; Sacchar. purif. $\tilde{\text{S}}\text{xxx}$; Acacia pulv. $\tilde{\text{S}}\text{ij}$; Ag. destillat. f $\tilde{\text{S}}\text{vj}$; Ol. Cinnamon. f $\tilde{\text{S}}\text{ss}$. Mix. To be divided into 720 lozenges. Ph. B., 1864.)

TROCHISCI CARBONATIS CAL'IS, *T. e Cre'ta*, *T. Cre'tæ* (Ph. U. S.), *Troches of Carbonate of lime or chalk*, *Loz'enges for the heartburn*, *Tabellæ ad ardo'rem ventriculi seu cardia'gicæ* seu *ad so'dam*. (Cret. ppt. $\tilde{\text{S}}\text{iv}$; Acacia, in pulv. $\tilde{\text{S}}\text{ij}$; Myrist. in pulv. $\tilde{\text{S}}\text{ij}$; Sacch. in pulv. $\tilde{\text{S}}\text{vj}$. M. Form into troches with water.) Antacid and absorbent.

TROCHISCI CAT'ECHEU, *Cat'echu Loz'enges*, (Cat'echu, in pulv. $\tilde{\text{S}}\text{ij}$; Sacchar. purif. Rij ; Acacia pulv. $\tilde{\text{S}}\text{ij}$; Tinct. capsic. f $\tilde{\text{S}}\text{ss}$; Ag. destillat. q. s. Mix, and divide into 720 lozenges. Ph. B., 1864.)

TROCHISCI E CRE'TA, T. carbonatis calcis—t. Cretæ, T. carbonatis calcis.

TROCHISCI CUBEB'Æ, *Troches of Cubels*, (Oleoresin. cubeb. f $\tilde{\text{S}}\text{ij}$; Ol. assaf. f $\tilde{\text{S}}\text{ij}$; Glycyrrhiz. pulv., Acacia pulv., Sacchar. pulv. aa $\tilde{\text{S}}\text{ij}$; Syrup. Tolutan. q. s. To be divided into troches, each weighing ten grains. Ph. U. S., 1863.) Excitant; in relaxed sore throat, &c.

TROCHISCI seu PASTILLI EMETIN'Æ PECTORALES, *Pectoral Loz'enges of Emetine*, (Sugar, $\tilde{\text{S}}\text{iv}$; Emetine, 32 grains. Form into lozenges of nine grains each.) One is a dose.

TROCHISCI ESCHAROT'ICI, *Escharot'ic Troches*, (Hydrag. oxymer. p. 8; Amyli, p. 16; Mucilag. g. tragac. q. s. Make into troches, to which a few drops of laudanum may be added.—Ph. P.) Used in foul ulcers, to remove excrescences, &c.

TROCHISCI ESCHAROT'ICI DE MIN'IO, (Oxyd. plumbi. rubr. p. 16; Hydrag. oxymer. p. 32; Muc. panis. p. 128; Ag. rosæ, q. s.—Ph. P.) Used like the last.

TROCHISCI FERRI SUBCARBONATIS, *Troches of Subcarbonate of Iron*, (Ferri subcarb. $\tilde{\text{S}}\text{v}$; Vanill. gr. lx; Sacch. in pulv. $\tilde{\text{S}}\text{xv}$; Mucilag. tragac. q. s. To be divided into troches, each weighing 20 grains. Ph. U. S., 1863.) Tonic.

TROCHISCI GLYCYRRHIZ'Æ, *T. G. Glu'bræ* seu *Be'chicæ nigri*, *Liquorice troches*, *Black pectoral loz'enges*, (Ext. glyc. glubr., Gum. acac. aa p. j; Sacch. purif. p. ij. Dissolve in warm water; strain; evaporate, and form into troches.—Ph. E.) Demulcent. To allay cough.

TROCHISCI GLYCYRRHIZ'Æ CUM O'PIO, *T. G. et o'pii* (Ph. U. S.) seu *Thoba'ici* seu *sedati'vobal'amici* seu *pectorales re'gii Dama'rum*, *Liquorice Troches with O'pium*, *Wis'tar's Loz'enges*. (Opii,

in pulv. $\mathfrak{Z}$ ss; *Glycerrhiz.* in pulv., *Sacchar.* in pulv., *Acacia*, in pulv., aa $\mathfrak{Z}$ x; *Ol. anisi*, f $\mathfrak{Z}$ j. Make into troches, each weighing six grains.—Ph. U. S.) Demulcent and anodyne. An analogous formula is given in the British Pharmacopœia (1864), under the name *Trochisci opii*, *Opium Lozenges*.

Spirita's Lozenges for coryza, hoarseness, and sore-throat, are composed as follows:—(*Cubeb.* recent, in pulv. $\mathfrak{Z}$ j; *Extract. glycyrrhiz.* $\mathfrak{Z}$ j; *Myrrorh.* $\mathfrak{Z}$ j; *Acacia*, $\mathfrak{Z}$ iv; *Syrup*, q. s. Make into troches of ten grains each.)

TROCHISCI GUMMO'SI, *Gum Troches.* (*G. acac.* p. v; *Amyli*, p. i; *Sacch. pur.* p. xij. Make up the troches with rose-water.—Ph. E.) Demulcent.

TROCHISCI IPECACUAN'HÆ, *Troches of Ipecacuan'ha.* (*Ipecacuanh.* in pulv. $\mathfrak{Z}$ ss; *Sacchar.* in pulv. $\mathfrak{Z}$ xiv; *Maranta*, in pulv. $\mathfrak{Z}$ iv; *Mucilag. tragacanth.* q. s. Make into troches, each weighing ten grains.—Ph. U. S.)

TROCHISCI LACTUCA'RII, *Troches of Lactuca'rium.* (Prepared with *Lactucarium* in the same proportion and manner as the opium lozenges.—Ph. E.) Each lozenge contains between the fifth and sixth of a grain of lactucarium.

TROCHISCI MAGNE'SIÆ, *Troches of Magnesia.* (*Magnes.* $\mathfrak{Z}$ iv; *Sacchar.* lbj; *Myrist.* pulv. $\mathfrak{Z}$ j; *Mucilag. tragacanth.* q. s. Make into troches, each weighing ten grains.—Ph. U. S.) Useful in acidity of the stomach, with constipation.

TROCHISCI MENTHÆ PIPERITÆ, *Troches of Peppermint.* (*Ol. menth. piperit.* f $\mathfrak{Z}$ j; *Sacchar.* in pulv. lbj; *Mucilag. tragacanth.* q. s. Make into troches, each weighing ten grains.—Ph. U. S.) Used as a carminative.

TROCHISCI MORPH'Æ, *Troches of Morphia.* (*Morph. Muriat.* $\mathfrak{Z}$ j; *Tinct. Tolut.* $\mathfrak{Z}$ ss; *Sacchar.* $\mathfrak{Z}$ xv. Make into lozenges, with mucilage, weighing fifteen grains each.—Ph. E.) Each lozenge contains about one-fortieth of a grain of muriate of morphia. Used to allay cough.

TROCHISCI MORPH'Æ ET IPECACUAN'HÆ, *Troches of Morphia and Ipecacuan'ha.* (*Morph. mur.* $\mathfrak{Z}$ j; *Ipecac.* pulv. $\mathfrak{Z}$ j; *Tinct. Tolut.* f $\mathfrak{Z}$ ss; *Sacchar.* $\mathfrak{Z}$ xv. Make into a mass with mucilage, and form lozenges, fifteen grains each.—Ph. E.) Each lozenge contains about one-fortieth of a grain of the muriate. Used to allay cough.

TROCHISCI NITRA'TIS POTAS'SÆ, *Nitre Troches.* (*Potassæ nitrat.* p. j; *Sacch. pur.* p. iij. Make into troches with mucilage of tragacanth.—Ph. E.) In slight inflammatory sorethroat and fever.

TROCHISCI OPII, see T. *Glycyrrhizæ cum opio*.

TROCHISCI PECTORALES REGIS DANORUM, T. *glycyrrhizæ cum opio*—t. *Sedativobalsamica*, T. *glycyrrhizæ cum opio*.

TROCHISCI SO'DÆ BICARBONATIS, *Troches of Bicarbonate of Soda.* (*Sodæ bicarb.* $\mathfrak{Z}$ iv; *Sacchar.* in pulv. lbj; *Mucilag. tragacanth.* q. s. To be divided into troches, each weighing ten grains.—Ph. U. S.) Antacid.

The *Pastilles de Vichy*, *Pastilles digestives d'Hauterive-Vichy*, *Pastilli digestivi Darcel'ii*, composed of Bicarbonate of soda, Oil of peppermint, and Sugar, are much used in France, like the Vichy waters, in cardialgia, gout, and lithiasis.

TROCHISCI STIBIATI, *Tabellæ antimoniales Kunkeli*—t. *Thebaici*, T. *glycyrrhizæ cum opio*.

TROCHISCI ZINGIB'ERIS, *Troches of Ginger.* (*Tinct. Zingib.* f $\mathfrak{Z}$ j; *Tragac.* pulv. gr. cxx; *Sacchar.* $\mathfrak{Z}$ xij; *Syrup.* *Zingib.* q. s. Make into troches, each weighing twenty grains.)

TROCHISQUE, *Trochiscus*.

TROCHITE'RI, gen. *Trochite'ris*, (from *τροχαις*, 'to turn.') The larger of the two tuberosities at the upper extremity of the os humeri; so called because it affords insertion to rotator muscles.

TROCHITE'RIAN, in the language of Chaus-

sier, means any thing belonging or relating to the trochiter.

TROCH'LEA, (*τροχλία*.) *Trochil'ia*, (F.) *Trochleë* ou *Trochlée*, (*τροχλῆς*, a pulley; (from *τροχαις*, 'to turn.')—for example, the articular surface at the lower extremity of the os humeri; so called from its forming a kind of pulley on which the ulna moves, in flexion and extension of the forearm. Also, the cartilaginous pulley over which the tendon of the trochlearis muscle passes, at the upper and inner part of the orbit. Pulleys are needed as mechanical powers in the reduction of dislocations where much force is required.

TROCHLEA LABYRINTHI, *Cochlea*.

TROCHLEARIS, *Obliquus superior oculi*.

TROCHLÉATEUR, *GRAND*, *Obliquus superior oculi*—t. *Petit*, *Obliquus superior oculi*.

TROCHLÉE, *Trochlea*.

TRO'CHOID, (*τροχικός*, from *τροχος*, 'a wheel,' and *αίς*, 'resemblance.') *Trochoï'des*, *Axis commissura*. An articulation, in which one bone turns upon another, like a wheel upon its axle; as the *Atloïdoïxoid articulation*. See *Axis commissura*.

TROÈNE, *Ligustrum vulgare*.

TROISQUARTS, (*trois*, 'three,' and *quarts*, pl. of *quart*, 'a fourth,') *Trocar*.

TROKANTER, *Trochanter*.

TROKANTERIAN, *Trochanterian*.

TROKANTINIAN, *Trochanterian*.

TROKLÉE, *Trochlea*.

TROLLIUS LAX'US, *T. America'nus*, *Globe flower*; Order, *Ranunculacæ*; indigenous; has the same medicinal properties as the *Ranunculus*.

TROMOMANIA, (*tromus*, and *mania*.) *Delirium tremens*.

TROMOPARANCEA, (*tromus*, and *parancea*.) *Delirium tremens*.

TROMOS, (*τρομος*.) *Paralysis agitans*, *Tremor*.

TROMOSPASMUS, (*tromus*, and *spasmus*.) *Tremor*.

TROMPE, 'a trumpet,' *Tuba*—t. *d'Eustache*, *Eustachian tube*—t. *de Fallope*, *Tuba Fallopiana*—t. *Uterine*, *Tuba Fallopiana*.

TROMUS, (*τρομος*.) *Paralysis agitans*, *Tremor*.

TROMYLE, (*τρομος*, 'trembling,' and *ἐλη*, 'matter,') see *Cilia*.

TRONC, *Trunk*—t. *Basilaire*, *Basiliary artery*—t. *Brachial*, *Brachial artery*.

TROPE'OLUM MAJUS, (from *τροπαιον*, 'a trophy,' the leaves resembling a buckler,) *Nasturtium Indicum* seu *Peruvia'num*, *Acrivola*, *Flos sanguineus monard'i*, *Cardaminum minus*, *Indian Cress*. Ord. *Geraniacæ*. *Sex. Syst.* *Ocandria Monogynia*. (F.) *Cresson des Indes*, *Cresson du Mexique*, *Capucine*. An antiscorbutic.

TROPÆOLUM TUBERO'SUM, *Ulluco*. A tuberous plant, cultivated in the Sierra, Peru. It is smaller than the potato.

TROPHALIS, (from *τροφή*, 'nourishment,') *Colostrum*.

TROPHE, (*τροφή*.) *Aliment*, *Cibatio*, *Pabulum*.

TROPHI, (from *τροφή*, 'nourishment,') In insects, the parts of the mouth that are employed in receiving and preparing the food.

TROPHICAL, *Trophicus*, (from *trophe*,) (F.) *Trophique*. Relating to nourishment or nutrition.

TROPICAL NERVES. The organic nerves or nerves of the sympathetic system have been so termed, from the belief that they are concerned in nutrition.

TROPHIMUS, (*τροφίμος*.) *Nutrition*.

TROPHODES, (*τροφικός*, from *τροφή*, 'nourishment,') and *odes*,) *Nutritious*.

TROPHONEURO'SES, (*τροφή*, 'nourishment,' and *neurosis*.) Morbid conditions of the process of nutrition, owing to modified nervous influence.

TRYPHONOSI, *Trophonosi*, (*τροφος*, 'nourishment,' and *νσος*, 'a disease,') *Diseases of nutrition*.

TROPHY, *Troph'ia*, (τροφία, a suffix denoting 'nourishment or nutrition:') as in *hypertrophy*, 'excess of nourishment or nutrition.'

TROPONUSI, *Morbi tropici*, (τροπή, 'the solstitial or tropical point where the sun turns or alters its course,' and νόσος, 'a disease,') Diseases that prevail in the tropics.

TROSTER, ([G.] 'a comforter,') see Spirit.

TROU, Foramen—*t. Anonyme*, Foramen stylo-mastoideum—*t. Avenge* ou *Borgue*, Cæcum foramen—*t. Borgue* of the Medulla oblongata, Foramen cæcum of the medulla oblongata—*t. de Botal*, Botal foramen, Ovale foramen—*t. Déchiré antérieur*, Lacerum anterius foramen—*t. Déchiré postérieur*, Lacerum posterius foramen—*t. Épineux*, Spinale foramen—*t. de Galien*, Botal foramen—*t. Meunier*, Mental foramen—*t. Optique*, Foramen opticum—*t. Orbitaire supérieur*, Orbital foramen superior—*t. Ovale*, Ovale foramen—*t. Rond*, *petit*, Spinale foramen—*t. Sous-orbitaire*, see Suborbital canal—*t. Sous-pubien*, Obturatorium foramen—*t. Sphéno-épineux*, Spinale foramen—*t. Sus-orbitaire*, Foramen supraorbitarium.

TROUS (pl. of trou) **CAROTIDIENS**, Carotica foramina—*t. Condyloldiens*, Condylolidea foramina—*t. de Conjugaison*, Conjugationis foramina, see Vertebrae—*t. Olfactifs*, Olfactory foramina—*t. Rachidiens*, Spinal foramina.

TROUSSE, Case (pocket).

TROUSSE-GALANT, (*trousser*, 'to despatch,' and *galant*, 'a gallant,') Cholera morbus.

TROUSSEAU, 'a bundle,' Fasciculus.

TROUSSEAUX ARRIÈRE-MÉSENTÉRIQUES (F.) A name given by Winslow to the nervous plexus situate between the two mesenteric plexuses, of which it is a dependence, and which descends into the pelvis to concur in the formation of the hypogastric plexus.

TRUBS, Lycoperdon tuber.

TRUCK, see Medicament.

TRUCULENTUS, (from *trux*, gen. *trucis*, 'ferocious,') Homicidal.

TRUDGEBACKED, see Hump.

TRUELOVE, Paris, Trillium latifolium.

TRUFFE, Lycoperdon tuber.

TRUFFLE, Lycoperdon tuber—*t. Hart's*, *Ela-phomyces granulatus*.

TRUMPET CREEPER, Tecoma radicans—*t. Flower*, Tecoma radicans—*t. Weed*, Eupatorium purpureum.

TRUNCUS, Trunk.

TRUNK, *Trunc'us*, (F.) *Tronc*. The principal part of the body, to which the limbs are articulated. The trunk has been divided into three parts:—*a superior, the head; a middle, the thorax, and an inferior, the pelvis*. These three regions contain the great splanchnic cavities. They are united by a common *stalk*—the vertebral column. To the *middle* are articulated the superior extremities; and, to the *inferior*, the lower extremities.

The *trunk* of an artery, vein, nerve, or lymphatic, is its largest part, which has not yet given off branches. It is, also, applied to the parent vessel, after branches have been sent off.

TRUSS, ([F.] *trousser*, 'to tie up,') *Brach'erium*, *Brach'erium*, *Brach'il'e*, *Brach'rotum*, *Brach'er'otum*, *Am'na*, *Hon'ma*, *Sublig'ulum*, *Subligat'ura*, (F.) *Brayer*. A hernial bandage for keeping hernia reduced. The ancient trusses were inelastic. They consisted simply of a pad and strap. At the present day, elastic bandages are alone employed. They possess the advantage of making an equable and continuous pressure on the aperture of the sac, and of yielding to the changes of form in the abdominal parietes. The most important part of these bandages is a piece of narrow steel, flat, and adapted to the shape of the body. This is the *spring*, (F.) *Ressort*. It embraces the body on the side affected; extends

from beyond the spine, and terminates, *opposite* the ruptured part, by an iron plate to which it is riveted. The posterior surface of this plate is furnished with a convex *pad*, (F.) *Pelote*, adapted in shape and size to the aperture which it has to close. The spring is covered with leather, and its inner surface is padded; a strap extends from its posterior extremity; passes round the sound side, and is attached to the plate of the pad. This strap is pierced with numerous holes, which allow the patient to tighten or slacken it at pleasure. Trusses have been variously constructed; and different patents obtained for the modifications. They all act upon the above principle. Sometimes they have a double pad.

TRYGODES, (τρυγώδης, from τρυγέ, gen. τρυγέ, 'lees, and odes,') Fecculent.

TRYPANON, (τρυπανόν,) Trepan.

TRYPESIS, (τρύπησις, from τρυπώ, 'to bore,') Trepanning.

TRYPSIS, (τρυψις, 'to rub,') Friction.

TSCHERBET, Sherbet.

TSETSE, Glossina morsitans.

TSHETTIK, Upas tieuté.

TSIANA, Costus.

TSITLA. A South African root of the Araceæ family, which contains a very large quantity of sweet tasted starch. According to Dr. Livingstone, when dried, pounded into meal, and allowed to ferment, it is not an unpleasant article for food.

TSJAMPACA, Michelia champaca.

TUBA, 'a trumpet,' Tube—*t. Aristotelica*, Eustachian tube—*t. Cæca*, Tuba Fallopiana—*t. Uterina*, Tuba Fallopiana—*t. Vocalis*, Tube, vocal.

TUBAGE, see Catheterismus.

TUBAIRE, Tubal.

TUBAL, *Tubal'is*, (F.) *Tubaire*, (from *tuba*, and *tubus*,) Relating to a tube or tubes, and especially to the Fallopian.

TUBAL PREGNANCY, see Pregnancy, tubal—*t. Respiration*, see Murmur, respiratory.

TUBE. A name given to some parts which are hollow, and which have, at times, the shape of a trumpet. *Tu'ba*, *Tu'bus*, *Sal'pinx*, (F.) *Trompé*, *Tube*.

TUBE, ALIMENTARY, Canal, alimentary.

TUBE CASTS. A term applied to the casts or moulds of the uriniferous tubes of the kidney, observed in the urine by the microscope in diseases of those organs. The casts are fatty, desquamative, hyaline, granular, &c., and are observed especially in Bright's disease.

TUBE, EUSTACHIAN, Eustachian tube.

TUBE, FALLOPIAN, *Tu'ba Fallopi'na* seu *Fallo'pii* seu *uteri'na* seu *cæ'ca*, *Me'trosal'pinx*, *Hys'terosal'pinx*, *Oviductus muli'ebri's*, *Vas def'eren's muli'eris*, *Ductus vario'sus* seu *Proces'sus latera'lis* seu *Mea'tus semina'lis* seu *Me'atus semina'rius u'teri*, *O'viduct*, *Vec'tor canal'*, (F.) *Trompe de Fallope* ou *uté'rine*, is a canal floating in the abdomen, and extending on each side from the superior angles of the uterus as far as the sides of the brim of the pelvis. They are 4 or 5 inches long. Near the uterus, they are straight and very narrow; but, in their course, they widen, become tortuous, and terminate by a free, expanded, and *fin'brated exten'sity*; called, also, *Folia'cium ornamen'tum*, *Mor'sus diab'oli*, *Infin'dib'ulum* seu *Digitatio'nes* seu *Fin'brice* seu *Vex'il'la* seu *Lacin'iæ tuba'rum Fallo'pii*, (F.) *Morceau frangé*, *Parillon de la trompe*, *Morceau ou Morsure du diable*. Within, the tubes are hollow and lined by a mucous membrane: the tissue exterior to which is spongy and erectile. The use of the Fallopian tubes is to conduct the sperm to the ovarium, and the fecundated ovum back to the uterus. They establish a communication between the cavity of the peritoneum and that of the uterus, the abdominal termination being termed *Os'tium abdomina'le*; the uterine *Os'tium*

uterinum. When the tube embraces the ovary, a funnel-shaped cavity—*Corolla infundibuliformis*—is formed at its extremity, into which the ovule passes.

TUBE, INTESTINAL, Canal, alimentary.

TUBE, ŒSOPHAGEAL, *Stomach Tube*. A long elastic gum tube, capable of being passed into the œsophagus or stomach.

TUBE, POLLEN, see Pollen.

TUBE, RECTAL, *Defecation tube*. An elastic gum tube, similar to the last, which is sometimes passed into the colon, to remove air from that intestine, or to enable enmata to be thrown up into it.

TUBE, STOMACH, see Tube, œsophageal.

TUBE, VO'CAL, *Tu'ba vocal'is*. The part of the air passages above the inferior ligaments of the larynx, including the passages through the nose and mouth.

TUBER, ([L.], 'a bump, a protuberance,') **Hump**, *Lycoperdon tuber*, *Tubercle*—t. Annulare, *Pons varolii*—t. *Atloïdo-occipital*, *Rectus capitis posticus minor*—t. *Cervinum*, *Elaphomyces granulatus*—t. *Cibarius*, *Lycoperdon tuber*.

TU'BER CINE'REUM, *T. C. cum Infundib'ulo*. A grayish tubercle, seen at the base of the brain behind the commissure of the optic nerves, which is continuous with the infundibulum.

TUBER CORPORIS CALLOSI, see *Corpus callosum*.

TUBER FRONTALE, *Frontal protuberance*—t. *Gulosorum*, *Lycoperdon tuber*—t. *Gutturosum*, *Bronchocele*—t. *Ischii*, see *Ischiatic*—t. *Ischiotrochantrien*, *Quadratus femoris*—t. *Nigrum*, *Lycoperdon tuber*—t. *Parietale*, see *Parietal Bones*—t. *Pellagra*, *Pellagra*—t. *Tympani*, *Promontory of the tympanum*.

TUBERA TERRÆ, *Lycoperdon tuber*.

TUBERCLE, *Tuber'culum*, (dim. of *tuber*,) *Tu'ber*, *Strangalion*, *Stran'galia*, *Tuber'culous deposit*, (vulg.) a *Ker'nel*, a *Knot*, (F.) *Tubercle*. A tumour in the substance of organs, from the production of new matter;—sensation null; growth sluggish. In *Pathological Anatomy*, the term is generally given to a species of degeneration which consists of an opaque matter, of a pale yellow colour: this, in its crude condition, has a consistence analogous to that of concrete albumen: it subsequently becomes soft and friable, and gradually acquires a consistence and appearance analogous to those of pus. Tubercles may be developed in different parts of the body; but they are most frequently observed in the lungs and mesentery. Pulmonary tubercles, *Pneumophymata*, *Tuber'cula Pulmo'num*, (F.) *Tubercules pulmonaires*, characterize *Tuber'cular Phthi'sis*. *Mesenteric tubercles* are often met with in the same affection, but particularly in *Tabes Mesenterica*. Laënnec classes tubercles among the accidental tissues, which have no resemblance to the natural tissues, and which never exist except in consequence of morbid action. Others consider them as a serofulous degeneration. Broussais conceives that they are produced exclusively by inflammation of the white vessels, whose existence, however, is more than questionable. When tubercles in any organ are few in number, they may pass to the state of permanent induration without danger to the patient; but when they are very numerous, they usually cause serious mischief. Dr. Baron considers that they are produced by hydatids. The morbid condition which gives rise to the development of tubercles, *Tuberculization*, is termed *Tuberculo'sis*, *Morbus tuberculo'sus*, *Strum'o'sis* seu *Phymato'sis pulmo'num*, and *Dyscrasia Tuberculo'sa*. Tubercles are seen of various forms,—to wit, in fine points, (F.) *Poussière tuberculeuse*, gray and yellow granulations, miliary tubercles; and gray and yellow tubercular masses, softened, and cretaceous.

TUBERCLE, BLACK, Melanosis—t. Painful subcutaneous, see *Neuroma*.

TUBERCLES OF THE AREOLA, see *Mamma*—t. *Cretaceous*, see *Cretaceous*—t. *Miliary*, see *Granulation*, and *Tubercle*.

TUBERCLES, PEARLY, *T. sebaceus miliary*, *Follicular elevations*. Small collections of sebaceous matter in the sebaceous glands or ducts, often seen in the thin skin of the lower eyelids, where they sometimes attain an inconvenient size.

TUBERCLES, SEBACEOUS MILIARY, *T. pearly*.

TUBER'CU'LA (pl. of *tuberculum*) *ARAN'TII*, *Cor'pora* seu *Corpus'cula Aran'tii*, *Tu'bercles of Aran'tius or Aran'zi*. Small eminences at the middle part of the convex edge of the semilunar valves of the pulmonary artery and aorta.

TUBERCULA ARTHRITICA, *Calculi*, *arthritis*—t. *Cerebri*, *Encephalophymata*—t. *Gonorrhoea*, *Gonorrhoea impura*—t. *Gummata*, see *Gumma*—t. *Hepatis*, *Hépatostromosie*—t. *Intestinarum*, *Enterophymata*—t. *Nervorum nodosa*, *Ganglions*, *nervous*—t. *Parva duræ matris*, *Glandula Pæchioni*—t. *Peritonei*, *Cœliophymata*—t. *Pulmonum*, see *Tubercles (of the lungs)*—t. *Quadrigenina*, *Quadrigenina Tubercula*.

TUBER'CU'LAR, *Tuber'culous*, *Tuberculo'sus*, *Tuber'culate*, *Tuber'culated*, (F.) *Tuberculeux*, *Tuberculé*. That which relates to tubercles, or which is formed by tubercles.

TUBERCULAR or TUBERCULOUS CONSTITUTION or *DIATHESIS*, *Phthisio'sis*, is the habit of body which predisposes to tubercular phthisis; as *Tuberculous Cachexia*, *Tuberculo'sis*, *Morbus tuberculo'sus*, *Dyscrasia tuberculo'sa*, is the constitutional condition of one affected with tubercles.

TUBERCULAR MATTER. A morbid production, contained in cysts, or loose in the tissue of organs. It varies, in its consistence, at the different periods of the tubercular affection: being, at first, compact, and yellowish; at times, calcareous; afterwards pulaceous, semifluid, and curdy. See *Tubercle*.

TUBERCULAR PHTHISIS is that produced by the development of tubercles in the lungs.

TUBERCULAR SORE THROAT, *Pharyngitis*, *follicular*.

TUBERCULATE, *Tubercular*.

TUBER'ULATE SARCO'MA of *AB'ERNETHY*, *Emphy'ma Sarco'ma tuberculo'sum*, *Tu'mor tuberculo'sus*. Tubercles, firm, round, and clustering; pea-sized or bean-sized; yellowish or brownish-red; when large, disposed to ulcerate, and produce a painful, malignant, and often fatal sore. Found chiefly in the lymphatic glands of the neck; often, simultaneously, in other glands and organs.

TUBERCULATED, *Tubercular*—t. *Liver*, *Cirrhosis*.

TUBERCULE, *Tubercle*.

TUBERCULÉ, *Tubercular*.

TUBERCULES BIJUMEAUX, *Quadrigenina tubercula*—t. *Crétacés*, see *Cretaceous*—t. *du Foie*, *Hépatostromosie*—t. *des Ganglions Mésentériques*, *Tabes mesenterica*—t. *des Glandes lymphatiques*, *Scrofula*—t. of the *Larynx* and *Fauces*, *Pharyngitis*, *follicular*—t. *Miliary*, see *Granulation*—t. *Pisiformes*, *Mamillary tubercles*—t. *Pulmonaires*, see *Tubercle*—t. *Quadrigenaux*, *Quadrigenina corpora*.

TUBERCULEUX, *Tubercular*.

TUBERCUL'TIS, (*tubercle*, and *itis*.) *Inflammation of a tubercle or of tubercular masses*.

TUBERCULIZATION, see *Tubercle*.

TUBERCULOSIS, see *Tubercle*, and *Tubercular Cachexia*—t. *Laryngis et Tracheæ*, *Phthisis laryngea*—t. *Pulmonum*, *Phthisis pulmonalis*—t. *Vertebrarum*, *Vertebral disease*.

TUBERCULOSUS, *Tubercular*.

TUBERCULOUS, *Tubercular*—t. *Deposit*, *Tu-*

herec—t. Disease of the Lungs, Phthisis pulmonalis—t. Dust, *Poussière tuberculeuse*.

TUBERCULUM, (dim. of *tuber*), Tubercle.

TUBERCULUM CINE/REUM. *Fasci'ola cine'rea*. A mass of cineritious substance at the top of the calamus scriptorius, which is continuous below with the posterior horn of cineritious substance in the cord, and on the sides with the corpus restiforme.

TUBERCULUM LACRYMALE, see Lachrymal puncta—t. Lower, Lower, tubercle of.

TUBEROSITAS, Tuberosity—t. Tympani, Promontory of the tympanum.

TUBÉROSITÉ BICIPITALE, Bicipital tubercle—t. *Sciaticque*, see Ischiatic.

TUBEROSITY, *Tuberos'itas*, (F.) *Tubérosité*, (from *tuber*, 'a bump or protuberance.') Anatomists have given this name to an eminence or process, the surface of which is unequal and rough; and which gives attachment to muscles or ligaments.

TUBES OF FERREIN, see Uriniferous Tubes.

TUBI MEMBRANACEI, (pl. of *tubus*, 'a tube,') see Villous membranes.

TUBULAR, *Tubula'ris*, *Tubulo'us*, (F.) *Tubuleux*, *Tubulaire*, (from *tubule*.) Relating to or having the form of a small tube. An epithet applied to glands—*Glandulæ tubulo'sæ*, which, like those of the stomach, have the shape of small tubes.

TUBULAR BLOWING SOUND, see Murmur, respiratory—t. Neurine, see Neurine—t. Substance of the Kidney, see Kidney.

TUBULE, *Tu'bulus*, (dim. of *tubus*, 'a tube,') A small tube or tubelet.

TUBULEUX, Tubular.

TUBULI (pl. of *tubulus*) BELLINI, Uriniferous tubes—t. Contorti, see Uriniferous tubes—t. Duræ matris, Sinuses of the dura mater—t. Ferreini, see Uriniferous tubes—t. Nervi, Nerve fibres—t. Pelvis renum, see Calix—t. Recti, see Rete testis and Uriniferous tubes—t. Seminales, see Testicle—t. Seminiferi, see Testicle—t. Seminiferi recti, see Rete testis—t. Uriniferi Bellini, Uriniferous tubes.

TUBULOSUS, Tubular.

TUBULUS, (dim. of *tubus*, 'a tube,') Canula, Tube—t. Centralis Modiolii, see Tractus spiralis foraminulentus.

TUBUS, Tube—t. Acusticus, Ear trumpet—t. Alimentarius, Canal, alimentary—t. Intestinalis, Canal, alimentary—t. Nervorum, Nerve, Vermicularis cæci, Appendix vermiformis cæci.

TUCKAHO, Lycopodon tubum.

TUE-CHIEN, 'kill dog,' Colchicum autumnale, Veratrum album—t. *Loup*, 'kill wolf,' Aconitum napellus.

TULBAG'HIA ALLIACEA, (after Tulbagh, a Dutch governor of the Cape of Good Hope.) A South African plant, *Nat. Ord.* Asphodelæ, the bulbs of which, boiled in milk, are recommended in phthisis as an anthelmintic.

Tulbaghia cepa'cea and *T. viola'cea*, serve the same purpose.

TULIP, CAPE, *Homeria collina*—t. Tree, *Liriodendron*.

TULIPIER, *Liriodendron*.

TULIPIFERA LIRIODENDRON, (*tulipa*, 'tulip,' and *ferre*, 'to bear,') *Liriodendron tulipifera*.

TUMEFACATIO, (*tumere*, 'to swell,' and *facere*, 'to make,') Swelling—t. Mollis, Puffiness.

TUMESCENTIA PITUITOSA, (from *tumescere*, 'to swell,') Leucophlegmatia.

TUMEUR, Tumour—t. *Anomale*, see *Hæmatodes fungus*—t. *Blanche*, *Hydrarthrus*—t. *Cavernuse*, see *Hæmatodes fungus*—t. *Encéphaloïde*, *Encephaloid*—t. *Érectile*, see *Hæmatodes fungus*

—t. *Fonguïde*, *Encephaloid*—t. *Fongueuse*, *Encephaloid*—t. *Gommeuse*, *Gumma*—t. *Hémorrhagiale circonscrite*, see *Aneurism*—t. *Lymphatique des articulations*, *Hydrarthrus*—t. *Ossinore*, *Encephaloid*—t. *Pulpeuse*, *Encephaloid*—t. *Spléniforme*, *Encephaloid*—t. *Spongieuse*, *Encephaloid*—t. *Variqueuse*, see *Hæmatodes fungus*.

TUMOR, gen. *Tumo'ris*, (from *tumere*, 'to swell,') Swelling, Tumour—t. *Albus*, *Hydrarthrus*—t. *Anomalus*, *Hæmatodes fungus*—t. *Artium*, *Obdormitio*—t. *Capitis Sanguineus*, *Cephalæmatoma*—t. *Carneus*, *Sarcoma*—t. *Cysticus*, *Encystis*—t. *Cysticus serosus*, *Hygroma*—t. *Desmoides*, *Tumour*, *fibrous*—t. *Fibrosus*, *Tumour*, *fibrous*—t. *Ficosus*, *Ficus*—t. *Flatulentus*, *Emphysema*—t. *Fugax*, see *Fugacious*—t. *Fungosus articuli*, *Spina ventosa*—t. *Fungosus sanguineus*, *Hæmatodes fungus*—t. *Gangliiformis*, *Ganglion*, *nervous*—t. *Genarum*, *Meloneus*—t. *Genu albus*, *Genocele*—t. *Glandulæ parotidæ*, *Parotidoneus*—t. *Gummatosus*, *Gumma*—t. *Lacteus*, *Galactocoele*—t. *Lardeus*, *Steatoma*—t. *Malarum*, *Meloneus*—t. *Ovarii*, *Oarioncus*—t. *Parotidis*, *Parotidoneus*—t. *Prostatæ*, *Prostatoneus*—t. *Renalis*, *Nephroncus*—t. *Sanguineus*, *Hæmatoma*—t. *Squamiformis carnosus*, *Lepidosarcoma*—t. *Tuberculosus*, *Tuberculate sarcoma*—t. *Tunicatus*, *Encystis*—t. *Uteri*, *Hysteroneus*.

TUMOUR, (*tumere*, 'to swell,') *Tu'mor*, *On'cos*, *On'cus*, *Epar'ma*, *Epar'sis*, *Emphy'ma*, *Excrescentia* (Sauvages), *Ecphy'ma*, (Sc.) *Sic'almæ*, (F.) *Tu'meur*. A rising or prominence, of greater or less size, developed by a morbid cause in some part of the body. Cullen defines a tumour to be "a partial swelling without inflammation." Sauvages restricts the term to a rising formed by the congestion of a fluid; and he calls *excrescences* those which have a fleshy or osseous consistence. Boyer defines it "any preternatural eminence developed on any part of the body." Good uses it synonymously with tubercle. Boyer's is the best general definition. Tumours differ greatly from each other; according to their seat, the organs interested, their nature, &c.

The French use the term *Tumeur blanche* synonymously with *White Swelling*;—see *Hydrarthrus*. The term is, however, applied by the generality of surgeons to swellings of the greater joints, without change of colour of the skin, and of a more or less firm consistence, which are dependent upon disease of the osseous, or of the soft, parts about the joint.

Scrofula is the most common cause of these affections.

TUMOUR, CELLULOFIBROUS, Tumour fibrocellular—t. Epithelial, see Epithelial.

TUMOUR, ERECTILE. A tumour produced by the development of a soft, vascular tissue, which is susceptible of elevation and depression. See *Erectile*.

TUMOUR, FIBROCELLULAR, *Cellulofib'rous tu'mour*. A soft and elastic form of fibrous tumour, a variety of the *gelatinous sarcoma* of Rokitsansky of which the softer kind of polypus of the nose is an example.

TUMOUR, FIBROPLASTIC, Sarcoma.

TUMOUR, FIBROUS, *Fib'rous growth*, *Fibro'ma*, *Fib'roid*, *Ino'ma*, *Tu'mor fib'rous seu demoi'des*, *Inohylo'ma*. A tumour or growth formed of fibrous tissue.

TUMOUR, LAMINATED NACREOUS FATTY, Cholesteatoma—t. Milklike, *Encephaloid*—t. Phantom, see *Fantom*.

TUMOUR, VARICOSE, *Tu'mor varico'sus*. A circumscribed, bluish tumour, formed by the dilatation and turgescence of the capillary vessels of a part.

TUMOUR, VASCULAR, see *Hæmorrhoids* and *Nævus*.

TUNAS, see *Cactus opuntia*.

TUNBELLY. *Physconia*.

TUN'BRIDGE, MIN'ERAL WATERS OF. A celebrated acidulous chalybeate in Kent, England, a few miles distant from the village of Tunbridge, and thirty-six miles south of London, which is much frequented. It contains carbonic acid, carbonate of iron, and sulphate of magnesia.

TUNDA OCULI, *Conjunctiva membrana*.TUNDUS CORDIS, *Mucro cordis*.

TUN'IC, *Tunica*, *Chiton*, *Coat*, (F.) *Tunique*. An envelope. A name given to different membranes, which envelop organs; as the tunics or coats of the eye, stomach, bladder, &c.

TUNICA, Tunic—*t. Acinialis*, Uvea—*t. Aciniformis*, Uvea—*t. Acinosa*, Uvea—*t. Adnata*, Conjunctiva—*t. Adventitia*, see Artery—*t. Agnata*, Conjunctiva—*t. Alba oculi*, Sclerotic—*t. Albuginea*, Albuginea, see Kidney—*t. Albuginea oculi*, Sclerotic—*t. Allantoidea*, Allantois—*t. Arachnoides*, Arachnoid membrane—*t. Aranea*, Arachnoid membrane—*t. Caduca*, Decidua—*t. Candida oculi*, Sclerotic—*t. Carnosa*, Panniculus carnosus—*t. Cellularis*, see Artery—*t. Conjunctiva*, Conjunctiva—*t. Costalis*, Pleura—*t. Crassa*, Decidua—*t. Crystallina*, Arachnoid membrane—*t. Dartos*, see Dartos—*t. Dura oculi*, Sclerotic—*t. Durior et crassa*, Sclerotic—*t. Elastica*, see Artery—*t. Elytroides*, Vaginal coat—*t. Erythroides*, Cremaster—*t. Exterior ovi*, Decidua—*t. Externa*, see Artery—*t. Extima oculi*, Sclerotic—*t. Farcinialis*, Allantois—*t. Fibrosa*, see Folliculi Graafiani—*t. Filamentosa*, Decidua—*t. Glabra*, see Artery.

TUNICA GRANULO'SA, 'granular coat.' An investment of the ovum exterior to the zona pellucida, described by Dr. Barry, but the existence of which, as a structure, distinct and independent from the membrana granulosa, is denied.

TUNICA HORTENSIS, *Dianthus caryophyllus*—*t. Innominata oculi*, Sclerotic—*t. Interna oculi*, Retina—*t. Intima*, see Artery—*t. Intima oculi*, Retina.

TUNICA JACOBI, *Membrana Jacobi*, *Jacob's coat* or *membrane*. A highly delicate serous membrane, interposed between the retina and the choroid coat, discovered by Dr. Jacob, of Dublin. Its structure is curious, being composed of cylindrical, transparent, highly refractive staffshaped bodies, *Bacilli*,—hence called *Stratum bacillosum* seu *bacillatum*, *Bacillar layer of the retina*,—which are arranged perpendicularly to the surface of the retina, their outer extremities being imbedded, to a greater or less depth, in a layer of the pigmentum nigrum, and are continued into the more internal layers of the retina, by means of thin processes—the *filaments* or *fibræ* of Müller.

TUNICA MEDIA, 'middle coat,' see Artery—*t. Muscularis*, Dartos—*t. Nervea oculi*, Retina—*t. Nervorum*, Neurilemma—*t. Perforata*, Uvea—*t. Prætensa abdominis*, Peritonæum—*t. Propria*, see Kidney—*t. Reticularis*, Retina—*t. Retiformis*, Retina—*t. Rhagoides*, Choroidea tunica, Uvea—*t. Rubicunda scroti*, Dartos—*t. Rubra*, *Dianthus caryophyllus*—*t. Ruyschiana*, see Choroid—*t. Subcostalis*, Pleura—*t. Thoracis*, Corset—*t. Uvulis*, Uvea—*t. Uvea*, Uvea—*t. Uviformis*, Uvea—*t. Vaginalis*, see Vaginal—*t. Vaginalis linguæ*, see Tongue—*t. Vaginalis propria*, Vaginal coat of the Testicle—*t. Vaginalis reflexa*, Vaginal coat of the Testicle—*t. Vaginalis testis*, see Testicle—*t. Vasculosa oculi*, Choroid, see Eye.

TUNICA VASCULO'SA TESTIS, 'vascular coat of the testis,' *Pia mater testis*. An extremely delicate membrane, consisting of minute ramifications of the spermatic vessels united by areolar tissue, which separates the lobules of the testis from each other. It is situate immediately within the tunica albuginea, and encloses the substance of the gland. It sends processes inward that separate the lobules, in the same manner as the pia

mater is reflected between the convolutions of the brain.

TUNICA VITREA, Hyaloid membrane.

TUNICULA, (dim. of *tunica*), *Tunicle*, (F.) *Tunicule*. A fine membrane, coat, or skin. Clitoris.

TUNIQUE, Tunic—*t. Albuginée*, Albuginea—*t. Commune de Richat*, Endangium.

TUN'NELLING. A term, sometimes employed for the treatment of stricture of the urethra, by making pressure upon it from within so as to promote its removal by absorption.

TUNNYFISH, SALTED, *Omotrichos*.

TUPHLOËNTERITIS, Typhloënteritis.

TUPHUS, Typhus.

TUPINARIA, Talpa.

TURBA HÆMORRHOIDALIS, 'hemorrhoidal disorder,' see Hæmorrhoids.

TURBATIONES ANIMI, (*turbare*, *turbatum*, 'to disorder,') Affections of the mind.

TUR'BINATED BONES, *Ossa turbinate*. Bones shaped like a top; (from *turba*, gen. *turbinis* or *turben*, 'a top;') *Os sa spongio'sa*, *O. tertii maxillæ superioris*, *Cor'na*, *Con'chæ na'rium seu ren'eris*, *Tur'bines na'si seu na'rium*, *Conchyl'is*, *Bucci'na*, *Lam'ine spongio'sæ na'si*, (F.) *Cornets*. Very thin, bony plates, rolled up in the form of horns, and situate in the nasal fossa. 1. *Os spongio'sum super'ius* or *Cornet de Morgagni*;—the highest. 2. *Os spongiosum me'dium*, the middle *cor'na* of French anatomists,—the *ethmoid'al cor'nu* or *Os spongio'sum super'ius seu turbinate*, of English anatomists, *Os convolutum super'ius*, *Con'cha*, *C. na'rium super'ior seu Morgagniana*, (F.) *Cornet moyen*. This bone projects inwards and downwards into the cavity of the nose, from the ethmoid cells at the side of the nasal lamella, and serves to enlarge the organ of smell. It is convex towards the septum, and concave outwards. 3. *Os spongio'sum seu turbinate infer'ius*, *Con'cha infer'ior*, *Convolutum infer'ius os*, *Os sous-ethmoid'al* (Ch.), projects at the under part of the side of the nose, is convex towards the septum, and concave outwards. It is connected to the os maxillare, os palati, and os unguis. 4. *Sphenoid'al cornu*.

All the turbinated bones extend the olfactory surface.

TURBINES NASI SEU NARIUM, (pl. of *turbo*, gen. *turbinis*, 'a top,') Turbinated bones.

TURBITH BLANC, Globularia alypum—*t. Mineral*, Hydrargyrus vitriolatus—*t. Montpelier*, Globularia alypum—*t. Noir*, see Hydrargyri oxydum cinereum—*t. Plant*, *Convolvulus turpe-thum*.

TURBO CEREBRI, Pineal gland.

TURGENTIA, Turgescence.

TURGES'CE, *Turges'cency*, *Turgidity*, *Tur'gidness*, *Turgescentia*, *Turgen'tia*, *Orgas'mus*, (from *turgescere*, 'to begin to swell, turgere, 'to swell,') Superabundance of humours in a part. The term *Turges'cence* of Bile was formerly used to denote the passage of that fluid into the stomach, and its discharge by vomiting. A vital action of turgescence or expansibility—*Turgor vitalis*—has been presumed to exist in the capillaries of certain organs, and in erectile tissues; but the generality of physiologists do not admit it.

TURGESCENTIA, Turgescence.

TURGESCENTIA VESICULE FELLEÆ, *Ascites Hep'atocysticus*, *Physconia seu Cystocœlia biliaris*, *Hep'at'gia Petiti'na*, *Hydrocholecystis*, *Hydrops vesicæ fellæ*, (F.) *Hydropisie de la Vésicule du Fiel*, *Distension de la Vésicule du Fiel*, *Cholécystectasie*. Distension of the gall-bladder by bile.

TURGIDITY, Turgescence.

TURGIDNESS, Turgescence.

TURGOR VITALIS, see Turgescence.

TURMERIC, *Curcuma longa*, *Sanguinaria Canadensis*.

TURN OF LIFE, see *Menses*.

TURN'ING, *Ver'sio*, (from [F.] *tourner*, 'to turn.') (F.) *Version*.

TURNING OF THE FŒTUS, *Ver'sio Fœtus*. Generally, the operation of bringing down the feet, or some part of the lower extremity, when the presentation of the child is such that it cannot be delivered by the natural efforts. This has been termed *podalic version*, in contradistinction to *cephalic version*, which consists in bringing the head down,—in shoulder-presentations, for example.

TURNIP, (*turn*, and [Sax.] *næpe*, 'navew,') *Brassica rapa*—t. *Dragon*, *Arum triphyllum*—t. *Indian*, *Arum triphyllum*, *Psoralea esculenta*—t. *Pepper*, *Arum triphyllum*—t. *Prairie*, *Camassia esculenta*.

TURNS, *Menses*.

TURPENTINE, *Terebinthina*—t. *Bordeaux*, see *Pinus sylvestris*—t. *Briançon*, see *Pinus cembra*—t. *Chio*, see *Pistacia terebinthus*—t. *Common*, see *Pinus sylvestris*—t. *Common*, *American*, see *Pinus sylvestris*—t. *Cyprus*, see *Pistacia terebinthus*—t. *Damarra*, see *Pinus damarra*—t. *Dombeya*, see *Dombeya excelsa*—t. *Horse*, see *Pinus sylvestris*—t. *Strasbourg*, see *Pinus picca*.

TURPENTINE TREE, *Pistacia terebinthus*.

TURPENTINE, VENICE, see *Pinus larix*—t. *Venice*, true, see *Pistacia terebinthus*—t. *White*, see *Pinus sylvestris*.

TURPETHUM, *Convolvulus turpethum*—t. *Minerale*, *Hydrargyrum vitriolatus*—t. *Nigrum*, *Hydrargyri oxydum cinereum*.

TURQUETTE, (from *Turkey*,) *Herniaria glabra*.

TURQUOISE, (from (I.) *turchino*, 'greenish-blue,') This stone has been long imagined to change its colour, according as the wearer is at the time in good or bad health: a superstitious notion.

TURTLE BLOOM, *Chelone glabra*—t. *Green*, *Chelonia mydas*—t. *Head*, *Chelone glabra*.

TURTUR, gen. *Turturis*, *Penis*.

TURUNDA, *Tent*.

TURUNDÆ ITALICÆ. Pastes made of the finest and most glutinous wheat—known under the names of *Macaroni*, *Vermicelli* and *Italian* or *Cagliari paste*. Similar preparations, made in Pennsylvania of flour and eggs, are called *noodles*, from the German *Nudel* n, 'pastry.'

TURUNDULA, (dim. of *turunda*), *Tent*.

TUS, see *Pinus abies*.

TUSCARORA ACID SPRING, see *Canada*, mineral waters of.

TUSSACA RETICULATA, (after G. R. De Tussac, a French botanist,) *Goodyera pubescens*.

TUSSEDO, *Tussis*.

TUSSICULA, see *Tussis*.

TUSSICULAR, *Tussicula'ris*, *Tussiculo'sus*, (from *tussicula*.) Relating to a cough, or a slight cough.

TUSSICULOSUS, *Tussicular*.

TUSSILA'GO, gen. *Tussilag'inis*, *T. Far'fura* seu *vulga'ris*, (from *tussis*, which it relieves;) *Bel-chium*, *Bel'chion*, *Cal'ceum equi'nium*, *Chamaele'n'ce*, *Chamaeg'e'ron*, *Fil'ius an'te' patrem*, *Farfarella*, *Gulionan'chus*, *Far'fura*, *F. Bel'chium*, *Un'gula calah'llina*, *Colts'foot*, *Foal'foot*, (F.) *Pus d'âne*, *Herbe de Saint Quirin*. *Ord. Compositæ*. *Sec. Syst.* Syngenesia Polygamia Superflua. It is demulcent and expectorant, and has been used in coughs, and pulmonary affections in general, and in some cutaneous diseases.

TUSSILAGO FARFARA, *Tussilago*—t. *Hybrida*, *T. petasites*.

TUSSILAGO PETASI'TES, *T. hybrida* seu *petasites* hermaphrodi'ta, *Petasites*, *P. ma'jor* seu *vir'bens* seu *officinalis* seu *hybrida* seu *vulga'ris*, *But'terbur*, *Pest'ilentwort*, (F.) *Pétasite*, *Herbe aux*

teigneux, *Chapelière*. The roots of this plant have been recommended as aperient and alexipharmic. They have a strong smell; and a bitterish, acrid taste, of the aromatic kind, but not agreeable.

TUSSILAGO PETASITES HERMAPHRODITA, *T. petasites*—t. *Vulgaris*, *Tussilago*.

TUSSIS; dim. *Tussicula*; *Beg'ma*, *Bex*, *Tuss'ida*, *Bex'is*, *Tussis nervosa*, *Catarr'chus*, *Puer'is*, *Tussis*, *Cough*, *Cough'ing*, *Haustr*, *Houstr*, (Prov.) *Conf.* (Sc.) *Host*, *Hoiat*, (F.) *Toux*. A violent, sonorous, frequent, and short expiration, by means of which the air, in passing rapidly through the bronchia and trachea, carries along with it the mucus of the parts, which forms the sputa. The cough is said to be *dry*, when unaccompanied by expectoration. It is symptomatic of many diseases. See *Bronchitis*.

TUSSIS AMPHEMERINA, *Pertussis*—t. *Asinina*, *Pertussis*—t. *Bronchica*, *Cough*, *tubal*, see *Bronchitis*—t. *Canina*, *Pertussis*—t. *Catarrhalis Simplex*, *Catarrh*—t. *Clamosa*, *Pertussis*—t. *Clangosa*, *Pertussis*—t. *Convulsiva*, *Pertussis*—t. *Delassans*, *Pertussis*—t. *Ferina*, *Pertussis*—t. *Pueros strangulans*, *Pertussis*—t. *Quinta*, *Pertussis*—t. *Senilis*, *Bronchitis* (chronic)—t. *Spasmodica*, *Pertussis*—t. *Stomachalis*, *Pertussis*—t. *Suffocans*, *Pertussis*—t. *Suffocativa*, *Pertussis*—t. *Tussiculosa*, *Pertussis*.

TUSSIVE, *Tussiv'ens*, (from *tussis*.) Belonging or relating to cough—as *tussive* vibration; the vibration of the parietes of the chest, caused by coughing.

TUTAM'INA, pl. of *tuta'men*, 'a defence,' 'a protection,' (from *tutare*, 'to defend.') Parts are so called which defend or protect others.

TUTAMINA CER'EBRI are the scalp, pericranium, bones of the skull, and the meninges.

TUTAMINA OC'ULI are the eyebrows, eyelids, and lachrymal apparatus.

TUTENAG, *Zincum*.

TUTHIA, *Tutia*.

TUT'IA, *Pom'pholyx*, *Cadmia*, *Alfasa*, *Mesera*, *Capni'tis*, *Tu'thia*, *Ox'ydum zin'ci impu'rum*, *Tutty*. The oxyd of zinc that attaches itself to the chimneys of furnaces in which ores of zinc are smelted. It is in the form of gray incrustations; and is sometimes used for making an eye-salve. It is prepared, for this purpose, by levigation.

TUTTERING, *Stuttering*.

TUTTY, *Tutia*.

TWIGG'S HAIR DYE, see *Hair dye*.

TWIN, ([Sax.] *twā*, *twā*, 'two,') *Gemellus*.

TWINGE, (from [D.] *dwingen*, 'to force,') see *Pain*.

TWINKLING OF THE EYE, *Nictation*.

TWINLEAF, *COMMON*, *Jeffersonia Bartoni*.

TWINROOT, *Orchis*.

TWINS, SIAMESE'. Two brothers, Chang and Eng, connected by means of a ligamentous band passing from the epigastrium of one to that of the other, who were exhibited in various parts of Europe and America, about the year 1830. They have lived in North Carolina, are married, and have children. See *Xiphopages*.

TWITCHGRASS, *Triticum repens*.

TWITCHING, (from [Sax.] *twiccan*, 'to wrest,') *Tic*.

TYCH'ICA, (from *τυχ*, 'accident,') Fortuitous lesions or deformities. The 7th class of Good's *Nosology*.

TYLE, (τυλη,) *Callosity*.

TYLOMA, (τυλωμα, from *tylus*, and *oma*,) *Callosity*.

TYLOPH'ORA ASTHMAT'ICA, (*tylus*, and *φerein*, 'to bear,' from the fleshiness of the leaflets of the corolla.) *Ascle'pias asthmatica*, *Cynan'chem Ipecacuan'ha* seu *vomito'rum* seu *viridiflorum* seu *tomento'sum*, *Ipecacuan'ha* of the *Isle*

of France, (F.) *Ipecacuanha blanc de l'Île de France*. *Order*, Aselepiadaceæ. A native of various parts of the East Indies, both of the continent and islands. The root is acrid, and is used on the coast of Coromandel as a substitute for ipecacuanha. In large doses, it is emetic; in smaller, often repeated, it is cathartic. It is recommended, also, as a diaphoretic, and especially in humeral asthma.

TYLOSIS, (τυλῶσις, from *tylus*, and *osis*), Callosity.

TYLUS, (τύλος,) Callosity.

TYMMA, gen. *Tym'matis*, (τυμμά,) Wound.

TYMPAN, Tympanum—t. *Membrane du*, see Tympanum.

TYMPANAL, Tympanic.

TYMPANAL, *Ouvre du tympan*.

TYMPANIA, Tympanites.

TYMPANIAS, Tympanites.

TYMPANIC or **TYMPANIC**, *Tympan'icus*, *Tym'panal*, (F.) *Tympanique*, (from *tympanum*.) Relating to the tympanum, as 'tympanic canal.'

TYMPANIC NERVE, see Petrous ganglion.

TYMPANIQUE, Tympanic.

TYMPANISM, *Tympanian'us*, (from *tympanum*.) State of parts affected with tympanites.

TYMPANITA, Tympanites.

TYMPANITE, Tympanites.

TYMPANITES, (τυμπανιτις,) *Emphyse'ma seu Pneumato'sis Abdom'inis*, *Hy'drops Tympanites*, *Affectio tympanit'ica*, *Ectympano'sis*, *Hy'drops abdom'inis aë'reus*, *Aëro'sis*, *Aëro'enterecto'sia*, *Aë'renterecto'sia*, *Enterærecta'sia*, *Physocæ'lia*, *Tympano'sia*, *Physe'ma*, *Physe'sis*, *Hy'drops sic'cus et flatulen'tus*, *Tympan'ia*, *Tympan'ias*, *Tympan'ia*, *Tympanit'is*, *Emphyse'ma Tympanites*, *E. Tympanit'icum*, *Meteorism'us*, *Tym'pany*, *Me'teorism*, *Drum belly*, *Wind drop'sy*, (F.) *Tympanite*, *Pneumatose péritonéale*, *Ballonnement*, (from *τυμπαον*, 'a drum,' so called because the abdomen is distended with wind, and sounds like a drum when struck.) A swelling of the abdomen, caused by accumulation of air in the intestinal tube or in the peritoneum. Tympanites may be idiopathic or symptomatic. The former depends upon exhalation of air from the inner surface of the intestine, or from decomposition of substances contained in it: the latter is the result of some organic affection, and often depends upon obliteration of the digestive tube, which prevents the gas from escaping. Idiopathic tympanites may be more readily cured. The symptomatic is usually fatal. On dissection, in such cases, the stomach and intestines are found enormously distended; with strangulation or disorganization of some part of the mucous membrane of the intestine. Carminatives may be proper when tympanites depends upon atony of the digestive organs: but, in cases of the symptomatic kind, they cannot be productive of advantage, and may do harm. Sometimes the air is contained in the cavity of the peritoneum—constituting *Tympan'ites abdom'inis*, *Aëropéritonie* of some:—the accumulation in the intestines being called *Tympanit'is intestinalis*.

TYMPANITES, UTERINE, *Physometra*.

TYMPANITIS, *Inflammatio tym'pani*, (*tympanum*, and *itis*.) Inflammation of the lining membrane of the middle ear. Also, tympanites.

TYMPANOSIS, Tympanites.

TYMPANUM, (τυμπαον, 'a drum,') The *Drum* or *bar'el of the ear*, *Ear drum*. The names *Cav'ity of the Tym'panum* or of the *Drum*, *Cav'itas seu Cavum Tym'pani*, *An'trum seu cav'itas antro'sa Au'ris*, (F.) *Tympan*, *Cavité du tympan*, *Caisse du tambour* ou du *Tympan*—have been given to a cavity of an irregular shape, which constitutes the middle ear, and has been compared to a drum. It is situate in the pars petrosa of the temporal bone, between the meatus audito-

rius externus and the labyrinth. This cavity is lined by a mucous membrane; and communicates, externally, by means of the Eustachian tube, which is situate between it and the pharynx. The tympanum has six parietes. 1. An *external*, which is almost entirely formed of the *Mem'brane of the Drum*, *Membrana tym'pani*, *Diaphrag'ma seu Mediast'ium seu Tegumen'tum au'ris*, *Myri'n-ga*, *My'rinc*, (F.) *Membrane du tympan*—a species of thin, transparent, circular septum; convex within, and concave without, which closes the inner extremity of the meatus auditorius. 2. An *internal*, on which are the *Fenes'tra ovalis*, *Protr'ontory*, and *Fora'men rotundum*. 3. A *superior*. 4. An *inferior*, on which is the *glenoid fissure*. 5. A *posterior*, on which are the opening of the mastoid cells, and the pyramid. 6. An *anterior*, on which are the *Process'us cochleariformis* and the inner orifice of the *Eusta'chian tube*. The cavity of the tympanum is crossed by a series of four small bones, articulated with each other by diarthrosis, moved by certain muscles, and representing a kind of bent lever extending from the membrana tympani to the fenestra ovalis. They are called the *Bones* or *Oss'icles of the Ear*, and are the malleus, incus, or orbicular, and stapes.

For the nervous plexus of the tympanum, see Petrosal Ganglion, and Otic Ganglion.

TYMPANUM MIXUS, see Fenestra—t. *Secundarium*, see Fenestra.

TYMPANY, Tympanites—t. of the Womb, *Physometra*.

TYPE, *Ty'pus*, *Peri'odus mor'bi*, (τυπος, 'a stamp,' itself from τυπτω, 'I strike,') The *type* is the order in which the symptoms of a disease exhibit themselves, and succeed each other. It may be continued, intermittent, or remittent.

TYPHA ANGUSTIFOLIA, (τυφή,) has the same properties as

TYPHA AROMATICA, *Acorus calamus*, and

TYPHA LATIFOLIA, *Cattail*, *Reed-mace*, *Greater Reed-mace*, *Bull-segg*, (F.) *Massette*, *Massé* ou *Quenouille d'Eau*. A plant common in marshes and ponds, in this country and Europe. *Ord.* Typhaceæ. *Sex. Syst.* Monœcia triandria. Its roots have been used in dysentery and blennorrhagia. The young shoots, before they reach the surface of the water, when boiled, taste like asparagus.

The down, (F.) *Aigrette*, *Poils*, which accompanies the flowers, has been applied like cotton to burns.

TYPHICUS, (from *typhus*), Typhoid.

TYPHIQUE, Typhoid, Typhus.

TYPHLITIS, (*typhlo*, and *itis*), Typhloënteritis.

TYPHLO, (from τυφλος, *cæcus*, 'blind,') in composition, the cæcum.

TYPHLODIDIDITIS, (*typhlo*, διδκς, gen. διδκδος, 'a double door, a valve,' and *itis*.) Inflammation of the ileocæcal valve.

TYPHLOËNTERITIS, *Typhloënterit'is*, *Typhlit'is*, *Inflammatio Cæci*, *Phleg'monous Inflammation of the Cæcum*, (F.) *Inflammation du Cæcum*, *Cæcite* (Piorry): (*typhlo*, and *enteritis*, inflammation of the intestines.) Inflammation of the cæcum, occasionally leading to perforative ulceration.

TYPHLOENTERUM, (*typhlo*, and *enteron*, 'intestine,') Cæcum.

TYPHLOPS, (*typhlo*, and *ωψ*, 'eye,') Cæcum.

TYPHLOSIS, (τυφλωσις,) Cæcitas.

TYPHLOTERTIS, (*typhloterum*, and *itis*), Typhloënteritis.

TYPHLOTERUM, Cæcum.

TYPHLOTES, (τυφλωτης,) Cæcitas.

TYPHO, (from τυφος, 'stupor,') in composition, typhus.

TYPHODES, (τυφωδης, affected with τυφος, 'stupor,') Typhous.

TYPHOËMIE, Typhobæmia—t. *Entérique*, see Typhus.

TYPHOEMIA, (*typhus*, and *αἷμα*, 'blood,') see Putrefaction.

TYPHOID, *Typhoides*, (*τυφώδης*, 'stupefied,') *Typhicus*, *Typhosus*, *Lactica*, (F.) *Typhique*, *Typhoide*, (*typho*, and *εἶδος*, 'resemblance,') Appertaining to or resembling typhus; as *Typhoid Fever*. Also, according to some, *Typhous Fever*, typhus. Fuchs has employed the word *Typhoide* for a family of diseases, to distinguish them from *Phlogoses*; of this family, *Stomacoe*, *Pharyngocoe*, and *Parinacoe* are examples.

TYPHOID AFFECTION, see Typhus—t. Fever of India, Cholera.

TYPHOMANIA, (*τυφομανία*, from *typho*, and *μανία*, 'delirium,') *Delirium* with stupor, *Subdelirium*. The kind of delirium common in typhus.

TYPHOMANIA, Coma vigil, Insomnia.

TYPHONIA, (from *typhus*,) Coma vigil.

TYPHOSEPTES, Typhoseptes.

TYPHOSEPTOSES, *Typhosep'ses*, (*typho*, and *εἶδος*, 'putrefaction,') Typhous affections.

TYPHOSIS, (*typhus*, and *osis*,) The typhous condition or diathesis.

TYPHOSUS, (from *typhus*,) Typhoid, Typhous.

TYPHOUS, *Typhosus*, (F.) *Typhique*, *Typhode*. Belonging or relating to typhus.

TYPHOUS DEPOSIT. A peculiar substance of new formation found in the areolar membrane, between the mucous and muscular coats of the patches of Peyer in typhoid fever.

TYPHOUS FEVER, see Typhoid.

TYPHUS, *Semipestis*, *Typhus*, *Febris Typhoides* seu *asthenica* seu *adynamica*, *Asthenopyra*, *Eucia typhus*, (F.) *Fièvre adynamique* ou *ataxique* ou *adynamique ataxique*, *Typhus d'Europe*; (from *τυφος*, 'stupor,') A fever characterized by small, weak, and unequal, but usually frequent pulse, with great prostration of strength, and much cerebral disturbance; its duration being generally from a fortnight to three weeks or longer. It is continued fever, accompanied with great cerebral irritation and prostration. (See Fever.) By most writers, this disease was formerly divided into two varieties—*Typhus mitior* and *Typhus gravior*.

TYPHUS MITIOR seu *simples* seu *nervosus* seu *comatosus*, *Febris lenta* *nervosa* seu *putrida nervosa* seu *hectica maligna nervosa*, *Blethropyra*, *Neuropyra*, *Neuropyretus*, *Nervous Fever*, (F.) *Fièvre nerveuse*, is characterized by slight shiverings; heavy, vertiginous headach; great oppression, peculiar expression of anxiety, nausea, sighing, despondency, and coma or quiet delirium.

TYPHUS GRAVIOR seu *carcerum* seu *castrensis* seu *anthropophthoracis* seu *contagiosis exanthematicis* seu *nostrias* seu *Europæus* seu *communis* seu *bellicus* seu *exanthematicus* seu *Hibernicus* seu *petechialis*, *Febris nervosa epidemica* seu *nervosa exanthematica* seu *maligna cum sopore* seu *typhoides* seu *nervosa petechialis* seu *pestilentialis* *Europæ* seu *punctularis* seu *saprophyra* seu *carcerum* et *nosocomium* seu *putrida* seu *continens putrida* seu *continua putrida* seu *pestilens maligna* seu *nautica pestilentialis* seu *petechialis*, *Septopyra*, *Porphyropythys*, *Sapropythys*, *Pulicularis morbus*, *Pestis bellica*, *True* or *Exanthematic* or *contagious typhus*, *Nervous fever* with *exanthematicous eruption*, *Putrid*, *Jail*, *Hospital* or *Ship Fever*, (F.) *Fièvre des Camps*, *F. nautique*, *F. d'hôpital*, *F. nosocomiale*, *F. des Prisons*, *F. lenticulaire*, *Typhus contagieux*, is attended with rigors and heat, alternating; little or no perspiration; pulse, tense and hard, usually quick, but fluttering; pain over the forehead and vertex; a 'mulberry rash,' appearing on the fifth to the seventh day of the disease, and lasting till the end; delirium succeeded by stupor, signs of incipient putrescence,—as true petechiæ, vibices,

hemorrhages, &c., *Pete'chial*, *Mac'ulated* or *Spotted Fever*, *Febris maculo'sa*. Typhus, under particular circumstances, seems to be communicable; but if proper attention be paid to cleanliness and ventilation, there is not much danger of communication. The general principles of treatment, laid down under the head of Fever, are applicable here. All the varieties of continued fever resemble each other in the cardinal points, although they may differ in many of their characters. Ventilation and cleanliness; warm or cold ablution or sponging; keeping the bowels free; and attending to symptoms as they may rise, with the avoiding of all irritation of every kind, are the main indications. When symptoms of great prostration occur, the strength must be supported by wine; and tonics be exhibited, with the mineral acids. Of the tonics, the sulphate of quinia is perhaps as good as any, whilst it is free from the objection which applies to most others,—of being too bulky. In the whole management of this affection, however, the physician has to discriminate carefully between oppression and prostration. The former may require evacuates; the latter cannot.

It is now maintained that there is an essential difference between the two varieties, typhoid fever and typhus; that the former is essentially an abdominal affection, and dependent upon, or connected with, an inflamed or ulcerated condition of the follicles or glands of Peyer; hence it has received the names *Fièvre mésentérique*, *F. typhoide*, *Maladie* ou *Affection typhoide*, *F. entéro-mésentérique*, *Entérite typho-hémique*, *Typhémie entérique*, *Entéro-mésentérite typhoide*, *Exanthème intestinale*, *Dothinentérite*, *Dothinentérite*, &c., *Neodididite*, *Typhodididite*, *Entérite folliculeuse*, *Gastro-entérite* with nervous affection of the brain, *Follicular gastroenteritis*, *F. enteritis*, *Enteritis folliculo'sa*, *Typhus abdominalis*, *T. intestinalis*, *T. Sporadicus*, *Enterotyphus*, *T. ganglionaris abdominalis*, *Febris intestinalis ulcero'sa*, *F. nervosa gastrica*, *F. nervosa enterica*, *F. enterica*, *F. nervosa mesenterica*, *Enterhelcosis nervosa*, *Typhus entericus*, *Ileopyra*, *Ileotyphus*, *Typhoid affection*, *Enteric fever*, *Pythogen'ic fever*. Accompanying this form of fever there is generally pain or gurgling or pressing the right iliac fossa, over the seat of the intestinal lesion; with more or less diarrhœa, and a typhoid eruption, consisting of rose-red papule, (see Spots, red,) which appears mostly on the abdomen in the second week of the disease. They are distinct from true petechiæ, as they can be removed by pressure; and distinct also from sudamina, which are perhaps the universal accompaniments of sweating, and are consequently a variety of miliaria. The common continued fever or synochus of Great Britain, and of this country, belongs to this variety, and the *Red Tongue Fever* of Kentucky was considered by Professor Bartlett to be typhoid fever.

Certain modern French writers have considered the Typhus of the European continent, *Typhus of Europe*, *pestilential fever*, *contagious typhus*, *Hungary fever*, *Hospital fever*, *camp fever*, *jail fever*, *lenticular petechial fever*, (F.) *Peste d'Europe*, &c., to be identical with the typhoid affection. They admit but one fever on the European continent—the typhoid. Typhus, they maintain—but by no means establish the position—to be peculiar to England.

TYPHUS ABDOMINALIS, see Typhus—t. *d'Amé'rique*, Fever, yellow—t. *Anthraxicus*, *Plague*—t. *Anthropophthoracis*, Typhus—t. *Aphthoides*, *Aphthæ*—t. *Bellicus*, Typhus—t. *Bengalensis*, *Cholera*—t. *Bubonicus*, *Plague*—t. *Carcerum*, Typhus gravior—t. *Castrensis*, Typhus gravior—t. *Cerebral*, *Meningitis*, *cerebrospinal*—t. *Cerebrospinal*, *Meningitis*, *cerebrospinal*—t. *Comatosus*, Typhus mitior—t. *Communis*, Typhus—t. *Conta-*

gieux, Typhus—*t.* Contagiosus exanthematicus, Typhus—*t.* Contagious, Typhus—*t.* Convulsivo-cerebral, Chorea, electric—*t. d'Europe*, Typhus—*t.* of Europe, see Typhus—*t.* Exanthematicus, Typhus—*t.* Ganglionaris abdominalis, see Typhus—*t.* Gravisimus, Plague—*t.* Hibernicus, *T.* gravior—*t.* Icterodes, Fever, yellow—*t.* Intestinalis, see Typhus—*t.* Jaune, Fever, yellow—*t.* Miasmaticque ataxique putride jaune, Fever, yellow—*t.* Morbillosus, Rubecula—*t.* Nervosus, Typhus mitior—*t.* Nostras, Typhus—*t.* d'Orient, Plague—*t.* Pestilentialis, Plague—*t.* Pestis, Plague—*t.* Petechialis, Typhus gravior—*t.* Puerpéral, see Peritonitis—*t.* Recurrens, see Relapse—*t.* Scarlatinus, Scarlatina—*t.* Simplex, Typhus mitior—*t.* Sporadicus, see Typhus—*t.* Tropicus, Fever, yellow—*t.* True, Typhus—*t.* Vesicularis, Pemphigus.

TYP'IC, *Typ'ical*, *Typ'icus*, (τυπικός,) (F.) *Typique*. Characterized by a particular type, as a 'typical fever.'

TYPOSES, (from *typus*), see Periodicity.

TYPUS, (τυπος,) Type.

TYRANNUS, 'a tyrant,' see Critical days.

TYRBASIA, (τυρβασία,) Agitation.

TYRBE, (τυρβη,) Agitation.

TYREMESIS, *Tyrot'emesis*, (tyros, and εμεσις, 'vomiting;') *Tyro'sis*, *Tyren'sis*. Vomiting of curdy matter, in infants especially.

TYREUSIS, (τυρεωσις, 'cheese-making,') Tyremesis, Tyrosis.

TYRIA, Porrigo decalvans.

TYRIASIS, Elephantiasis Arabica.

TYRINE, (from *tyrus*), Casein.

TYRODES, (τυροδης, from *tyrus* and *odes*), Cheesy.

TYROMA, gen. *Tyro'matis*, Porrigo decalvans.

TYRUS, (τυρος,) Cheese.

TYROSIS, (from *tyrus*, and *osis*), Tyremesis. Also, the curdling of milk in the stomach; *Tyreu'sis*.

TYROTEMESIS, (τυρωσις, 'made into cheese,' and εμεσις, 'vomiting,') Tyremesis.

U.

UBER, see Mamma.

UBERIS APEX, 'apex of the breast,' Nipple, Papilla.

UDDER, (Sax. *uðer*), see Mamma.

ULA, (ουλα, from ουλος, 'solid,') Gingivæ.

ULÆMORRHAGIA, Ulorrhagia.

ULATROPH'IA, (ουλον, 'the gum,' and *atrophia*, 'atrophy,') Shrinking of the gums; falling away of the gums.

UL'CER, (έλκος,) *Ul'cus*, *Hel'cos*, *Elco'sis*, *Elco'ma*, *Hel'cus*, *Helco'ma*, a *Sore*, (Sc.) *Sair*, (Prov.) *Kile*, (F.) *Ulcère*. A solution of continuity in the soft parts, of longer or shorter standing, and kept up by some local disease or constitutional cause. Richerand makes four great distinctions between a wound and an ulcer. 1. A wound arises from the action of an extraneous body:—the cause of an ulcer is inherent in the economy. 2. A wound is always idiopathic:—an ulcer is always symptomatic. 3. A wound has essentially a tendency to heal, because the action of its cause has been momentary:—an ulcer, on the contrary, has a tendency to enlarge, because its cause persists. 4. The treatment of a wound is purely surgical; that of an ulcer is medical as well. The immediate cause of an ulcer is perverted nutrition; the formative cells losing their plastic power. Ulcers have been variously divided, according to their nature;—into *simple*, *sinuous*, *fistulous*, *fungous*, *gangrenous*, *scorbutic*, *syphilitic*, *cancerous*, *scirrhus*, *epithelial*, *encephaloid*, *invertebrate*, *scrofulous*, *phagedenic*, *virulent*, *cacoëthic*, *sordid*, *carious*, *varicose*, &c. The treatment consists in removing, by appropriate means, the internal and the local causes which keep up the ulceration, and in exciting the vessels of the part, where necessary, by appropriate bandaging, &c.

ULCER, **ADEN**, see Beriberi—*u.* Delhi, see Delhi—*u.* Jacob's, Lupus—*u.* Malignant, Hospital gangrene—*u.* Putrid, Hospital gangrene—*u.* Warty of Marjolin, see Warty Tumor of Cicatrices—*u.* Yemen, see Beriberi.

ULCERA INTESTINALIA, (pl. of *ulcus*), *Gastroeclosis*—*u.* Laryngis, Phthisis laryngea—*u.* Serpentina oris, Aphthæ—*u.* Uteri, see Metrophthisis—*u.* Ventriculi, Gastroeclosis—*u.* Vesicæ, Cystophthisis.

UL'CERATED, *Ulcera'tus*, *Exulcera'tus*, *Helco'des*, (F.) *Ulcéré*; (*ulcerare*, *ulceratum*, 'to make

sore,' from *ulcus*.) In the state of an ulcer. Affected with ulcers.

ULCERATIO, Ulceration—*u.* Intestinalis, Enterolecosis—*u.* Ventriculi, Gastroeclosis.

ULCERATION, *Ulcera'tio*, *Exulcera'tio*, *Pyrogen'ia corrosiv'a*. A superficial ulcer. The formation of an ulcer—*Exulc'a'sis*, *Elco'sis*, and *Helco'sis*.

UL'CERATIVE, *Ulcera'tivus*, (F.) *Ulcératif*. Relating to ulceration,—as an 'ulcerative process.'

ULCÈRE, Ulcer—*u.* du *Larynx*, Phthisis laryngea.

ULCÈREUX, Ulcerous.

UL'CEROUS, *Ulcero'sus*, *Helco'des*, *Helcoï'des*, (from *ulcus*), (F.) *Ulcéreur*. Having the character of an ulcer. Affected with ulcers.

ULCERS, **EGYPTIAN**, *Ægyptia* ulcera.

ULCUS, gen. *Ul'ceris*, (έλκος,) Ulcer—*u.* Atonicum, Rupia—*u.* Cacoëthes, see Cacoëthes—*u.* Cancerosum, Cancer, Chancre—*u.* Dacryodes, see Dacryodes—*u.* Depascens, Phagedenic ulcer—*u.* Ferinum, Phagedenic ulcer—*u.* Fistulosum, Hypophora—*u.* Hypupum, see Hypupus—*u.* Narium foetens, Ozæna—*u.* Obambulans, Phagedenic ulcer—*u.* Pulmonum, Phthisis pulmonalis—*u.* Sinuosum, Hypophora—*u.* Syriacum, Cynanche maligna—*u.* Telephium, Telephium—*u.* Tuberculosum, Lupus—*u.* Uteri, Hyperlecosis, Leucorrhœa.

ULCUSCULA PRÆPUTII, (pl. of *ulcusculum*), Herpes præputii.

ULCUSCULUM, (dim. of *ulcus*), Ulcer—*u.* Cancerosum, Chancre.

ULE, (υλη,) Cicatrix.

ULETICUS, (from *ulon*), Gingivalis.

ULICUS, Gingivalis.

ULITIS, (*ulon*, and *itis*), *Ouli'tis*, *Inflammat'io gingivæ*, *Periodont'itis gingiva'rum*, (F.) *Ulite*, *Gingivite*, *Inflammation des gencives*. Inflammation of the gums.

ULITIS SEPTICA, Cancer aquaticus.

ULLUCO, *Tropæolum tuberosum*.

ULMAIRE, *Spiræa ulmaria*.

ULMARIA, (from *ulmus*), *Spiræa ulmaria*—*u.* Palustris, *Spiræa ulmaria*.

ULM'US, *U. campestris* seu *glabra* seu *montana* seu *n'da* seu *sati'va* seu *sco'bra*, *Pte'lea*, *Com'mon Elm*, (F.) *Orme*, *Ormeau*, *Orme pyramidale* ou *champêtre*. *Ord. Ulmaceæ*. *Sex. Syst.* Pentandria Digynia. The inner, rough bark—*Ulmus*, (Ph. B.)—of this tree is inodorous; and has a

bitter, austere, and mucilaginous taste. It is considered tonic, alterative, and diuretic; and has been used, chiefly, in lepra and other cutaneous affections.

ULMUS ALA'TA, *Lynn Wahoo'*. The bark of the root has been used successfully as a cataplasm in phlegmonous inflammation.

ULMUS AMERICA'NA, *U. fulva* seu *rubra*, *Rough-leaved, Red, or Slippery Elm*. The inner bark—*Ulmus fulva*, (Ph. U. S. 1863.)—is highly mucilaginous, and is used in coughs, diarrhoeas, and dysenteries; and, as a poultice for tumours, lacerated and contused wounds, &c. It may be employed, also, as a tent for dilating openings or passages.

ULMUS CAMPESTRIS, Ulmus—*u. Fulva*, *U. Americana*, *U. rubra*—*u. Glabra*, Ulmus—*u. Montana*, Ulmus—*u. Nuda*, Ulmus—*u. Sativa*, Ulmus—*u. Scabra*, Ulmus.

UL'NA, (ὤληνη,) *O'lenē*, *Foc'ilē inf'e'rius* seu *ma'jus*, *Can'na Bra'chii*, *Os cu'biti inf'e'rius*, *Os pro'eubita'le*, *Cu'bitus*, *Arui'do bra'chii ma'ior*, *Cu'bit*. Anatomists have given this name to the bone of the forearm which forms the prominence of the elbow, during the flexion of that joint. It is the longer and larger of the two bones, and is inserted at the inner side. Its upper and larger extremity has, behind, a considerable process, called *olecranon*; and, before, a smaller one, called *coronoid*. These two eminences are separated from each other by the greater *sig'moid* or *semilun'ar fos'sa* or *olecrano'id cavity*, which receives the articular trochlea of the humerus. The tubercle of the ulna is a small, rough spot, under the root of the coronoid process, for the insertion of the *Brachialis internus*. On the outside of this extremity of the bone is the *les'ser sig'moid cavity*. It is articulated with the head of the radius. The inferior extremity of the cubitus, which is much smaller than the superior, and articulated with the radius, has a surface that corresponds to the fibrocartilage which separates it from the os cuneiforme; and a pointed process called *sty'loid*. The body of the ulna is manifestly triangular. It is articulated with the os humeri and radius, and, mediately, with the os cuneiforme. It is developed by three points of ossification. Also the elbow.

ULNAD, see Ulnar aspect.

ULNAIRE, Ulnar.

ULNAR, Cubital.

UL'NAR AS'PECT. An aspect towards the side on which the ulna is situated.—(Barclay.) *Ulnad* is used adverbially by the same writer, to signify 'towards the ulnar aspect.'

ULNARIS, (from *ulna*,) Cubital—*u. Externus*, see Cubital muscles—*u. Gracilis*, *Palmaris longus*—*u. Internus*, see Cubital muscles.

ULOCACE, (*ulon*, and *κακος*, 'evil,') Cancer aquticus, Stomacace.

ULON, (ὤλον,) Gingiva.

ULON'CUS, (F.) *Uloncie*, (*ulon*, and *γκος*, 'tumour,') Swelling of the gums; epulis.

ULORRIAG'IA, *Ulamorrhag'ia*, *Ondorrhag'ia*, *Stomatorrhag'ia*, *Hæmorrhag'ia gingiva'rum*, *Ulorrhœ'a*, (*ulon*, and *ρηννυ*, 'I break forth,') Bleeding from the gums.

ULORRHEA, (*ulon*, and *ῥεω*, 'to flow,') Ulorrhagia.

ULOSIS, (*ule*, and the suffix *osis*,) Cicatrization.

ULOTICA, (from *ule*,) Cicatrissantia.

UL'TIMISTERNAL. Professor Bérard gave this appellation to the sixth or last bony portion of the sternum. He calls it also *Ensi'ernal*.

UL'IMUM MO'RIENS, 'last dying.' A term given to the part of the economy which dies last.

ULUM, (ὤλον,) Gingiva.

ULVA CRISPA, Fucus erispus.

UL'VA Latis'sima, Broad Green La'ver, Green Stake, *Oys'ter green*; Order, Algæ; is used at

table cooked in the same way as the next, to which it is greatly inferior.

ULVA UMBILICA'LIS, *Por'phyra lacinia'ta* seu *umbilica'tis*, *Lacinia'ted Pur'ple La'ver*, abundant on the shores of Great Britain, is pickled with salt and sold in London as *La'ver*. It is eaten at table, stewed with pepper, butter or oil and lemonjuice or vinegar. It has been suggested, also, as an antiscorbatic.

UMBEL, YELLOW, *Cypripedium luteum*.

UMBILIC, Umbilical, Umbilicus.

UMBILICAL, *Um'bilic*, *Umbilica'tis*, (F.) *Ombilical*, (from *umbilicus*.) That which belongs or relates to the navel.

UMBILICAL ARTERIES, which exist only in the fœtus, seem, as it were, continuations of the primitive iliacs. They clear the umbilical ring, and proceed to the placenta, to which they carry the residuum of the blood sent to the fœtus by the umbilical vein. As soon as respiration is established, the blood ceases to pass by the umbilical arteries, which become obliterated, as well as the vein, and are transformed into two very strong ligamentous cords,—the vein forming the *Ligamentum rotun'dum* of the liver.

UMBILICAL CORD, Funiculus umbilicalis.

UMBILICAL RE'GION, *Re'gio umbilica'tis* seu *gas'trica seu me'sogas'trica*, *Mesogast'rium*, (F.) *Région ombilicale*, is the middle region of the abdomen, in which the umbilicus is placed. The sides of this are called the *flanks* or *lum'bar re'gions*. In the umbilical region are the omentum majus; the inferior extremity of the duodenum; the jejunum; and the greater part of the mesentery; the aorta; vena cava; the trunks of the renal arteries and veins; the origin of the spermatic arteries, &c.

UMBILICAL RING, *An'nulus umbilica'tis*, (F.) *Anneau ombilical*, is a fibrous ring which surrounds the aperture of the umbilicus, and through which umbilical hernia occurs in children.

UMBILICAL VESICLE, *Vesicula umbilicalis*.

UMBILICAL VES'SELS, (F.) *Vois'seaux ombilicaux*, include the two arteries, and umbilical vein. The umbilical vein arises from the placenta, and terminates at the fissure on the inferior surface of the liver of the fœtus, to which it conveys the blood necessary for its nutrition.

UMBILICALIS, Umbilical.

UMBILICATED, *Umbilica'tus*, (F.) *Ombiliqué*. Depressed in the centre like the umbilicus,—as an 'umbilicated pustule.'

UMBILICUS, *Umbili'cum*, *Um'bo*, *Om'phalos*, *Ra'dix seu Me'dium ven'tris*, *Mesom'phalum*, *Mesom'phalum*, *Protome'sis* (navel of a newborn child); the *um'bilic*, *na'vel*, (Prov.) *Spoon*; (from *umbo*, 'the button or prominence in the middle of a buckle,' or from *ὀμφαλος*, or rather *ὀμφακος*, which signifies the same, as well as the navel;) (F.) *Ombilie*, *Nombril*. A round cicatrix, about the median line of the abdomen. It is in the situation of an aperture, which, in the fœtus, affords passage to the vessels of the umbilical cord.

UMBILICUS MARI'NUS, *Cotyle'don mari'na*, *Andro'sacé*, *Acetab'ulum mari'num*, *Andro'sacé Mat'hé'olt*, *Fun'gus petræus mari'nus*. A submarine production, found on the rocks and the shells of fishes, in the south of France. It is reputed to be anthelmintic and diuretic.

UMBO, Elbow, Umbilicus.

UMBRELLA TREE, (dim. of *umbra*, 'ashade,') *Magnolia tripetala*.

UMSCHLAG, ([G.] *um*, 'around,' and *schlagen*, 'to throw,') see Compress.

UNBOWELING, Exenterismus.

UNCARIA GAMBIR, *Nauclea Gambir*.

UNCAS, INDIAN, *Veratrum viride*.

UNCIA, (ὀνγία,) Ounce.

UN'CIFORM, *Uncifor'mis*, *Uncina'tus*, (*uncus*,

and *forma*, 'shape.') That which has the shape of a hook.

UNCIFORM BONE, *Os unciforme seu hamatum seu uncinatum*, (F.) *Os crochu*. The fourth bone of the second row of the carpus. Its shape is very irregular. Inwards and forwards, it has a considerable eminence, which is curved upon itself, and gives attachment to the annular ligament of the carpus. It is articulated with the os semilunare, os magnum, os cuneiforme, and the fourth and fifth metacarpal bones. It is developed from a single point of ossification.

UNCIFORM EMINENCE, Hippocampus minor.

UNCIFORM PROC'ESS, *Processus uncinatus seu Blumenbachii*, Great Proc'ess of the Eth'moid. The most considerable of the thin irregularly twisted laminae on the inferior surface of the ethmoid bone, which narrow the opening of the maxillary sinuses.

UNCINATUS, (from *uncinus*,) Unciform.

UNCINUS, Hook.

UNCTION, (*ungere, unctum*, 'to anoint,') Inunction, Unguentum, Unguentum hydrargyri.

UNC'TOR, (same etymon as Unction.) The anointer in ancient bathing establishments.

UNCTORIUM, Elæothesion.

UNCTUARIUM, Elæothesion.

UNCUS, gen. *Unci*, Hook.

UNDERCLIFF, ISLE OF WIGHT, (CLIMATE OF.) The Undercliff comprehends a tract of country from Dunnose to St. Catherine's Hill, on the southeast coast, about six miles in length, and from a quarter to half a mile in breadth. It is well protected from the colder winds; and the climate is remarkably equable, as well as mild and dry, so that there are not many days during the winter in which the invalid cannot take exercise in the open air. It is an excellent climate for the pulmonary valetudinarian.

UNDIM'IA. A name given by Paracelsus to a tumour filled with a gelatinous matter, like the white of egg.

UNDOSUS, (from *unda*, 'a wave,') Cymatodes.

UNDULATIO, (from *unda*, 'a wave,') Fluctuation.

UNEDO, gen. *Unedo*'nis, *Arbutus unedo*—u. Papyracea, *Arbutus unedo*.

UN'GUEAL, *Unquæ'lis*, *Un'qual*, *Unqua'lis*, (from *unguis*, 'a nail,') Belonging or relating to the nails of the hands or feet.

UNGUEAL MATRIX, see Nail.

UNGUEN, (from *ungere, unctum*, 'to anoint,') Unguentum—u. Articulare, Synovia.

UNGUENT, ARMATORY, Hoplochrysa.

UNGUENTARIA, see Myristica Moschata.

UNGUENTARIUS, *Myrep'sus*, *Myropæ'us*, *Myropo'les*, *Myropo'tus*, *Pimenta'rius*, (from *unguentum*. One who makes and sells ointments. A perfumer.

UNGUENTUM, *Un'guen*, *Diachrym'a*, *My'ron*, *Un'ction*, *Salve*, (F.) *Onguent*, (from *ungere, unctum*, 'to anoint,') Ointment. Ointments are topical applications, of a consistence more or less analogous to that of lard. They are used chiefly as local applications to ulcers and wounds; but are sometimes rubbed upon a part in cutaneous affections; and, especially, where it is desired that the constituents of the ointment shall be absorbed. Ointments are of less consistence than cerates, although the terms are often used indiscriminately. See Pomatum, (F.) *Pommade*.

UNGUENTUM AC'IDI NITRO'SI, *Oint'ment of Nit'rous Acid'id*, *Poma'tum seu Unguentum Nit'ricum seu Oxygena'tum*, (F.) *Pommade oxygénée*, *Graisse oxygénée*, *Pommade d'Algon*. (*Adip. suilli*, lbj; *Acid nitros*. ʒvj. Melt the fat, and rub in the acid gradually, till cold. Ph. E.) Stimulant. Used in itch and other cutaneous eruptions, and in foul ulcers.

UNGUENTUM AC'IDI TAN'NICI, *Ointment of Tan-*

nic Acid. (*Acid. Tannic*. gr. xxx; *Aq. fʒij*; *Adipis*. ʒi. Rub the acid with the water, and mix with the lard. Ph. U. S., 1863.) Used like Unguentum Gallæ.

UNGUENTUM ACONI'TIÆ, *Ointment of Aconit'ia*. (*Aconit'ie*, gr. viij; *Sp. rectif*, fʒss; *Adipis præparat*. ʒj M.) In neuralgic pains.

UNGUENTUM ADIPIS, U. Ceræ.

UNGUENTUM ADIPOCERÆ CETORUM, *Ceratum cetacei*—u. *Ægyptiacum*, *Linimentum æruginis*—u. *Ægyptium album*, *Crinomyron*—u. *Æruginis*, U. subacetatis cupri—u. *Album*, *Unguentum cetacei*—u. *Album Resolvens*, *Linimentum ammoniæ fortius*.

UNGUENTUM DE ALTHÆ'Æ, *U. de Terebinthinâ et cerâ seu citrinum seu resinæ pi'ni sylvestris compositum seu resumptivum seu lenitivum*, *Ceratum de Althæ'a*, *Emplastum mucilaginosum seu emolliens seu citrinum seu flavum seu de Althæ'a*, *Unguentum flavum*, *Marshmallow Oint'ment*, (F.) *Onguent de guimauve*, *O. d'althæa*. (*Olei de mucilagibus*, 1000 p., *Ceræ flavæ*, 250 p., *Resinæ Terebinthin.*, ʒā 125 p. Liquefy; strain and stir till cold. Ph. P.) Common Olive Oil might be substituted for the Oil of Mucilages.

UNGUENTUM AMMONIACALE GONDRETI, *Pommade de Gondret*—u. *Amygdalinum*, *Ceratum Galeni*.

UNGUENTUM ANTIMONI', *U. Antimonii'le*, *U. A. Tartarizati*, *U. A. Potassiotartaratis*, *U. tartari emetici*, *Oint'ment of tartarized antimony*, *Antimo'nial Oint'ment*, *Tartar emetic oint'ment*. (*Antimon. et potass. tartrat. in pulv. subtiliss*. ʒij, *Adipis*. ʒi. Ph. U. S.) If a drachm of this ointment be rubbed upon any part, night and morning, it will excite, in a few days, a painful pustular eruption. Hence, it is employed as a counterirritant. This preparation is also called *Ung. Tartari stibiat'i seu emetici*, *Ung. e Tartaro stibiat'o*, *A'deps Tartaræ stib'ii medicat'us*, *Ung. Tartra'tis Potass'æ stibiat'i*, (F.) *Pommade d'Autenrieth*, *Pommade stibiée*. This method of employing the tartrate of antimony and potassa, as a revellent, is, at times, called *stib'iodermic*, (F.) *La méthode stibio-dermique*.

UNGUENTUM ANTIMONII POTASSIOTARTRATIS, *Unguentum antimonii*—u. *Antimonii tartarizati*, *U. antimonii*—u. *Antipernium*, see *Antipernium*—u. *Antipersorium Werlhofii*, *Unguentum hydrargyri præcipitai albi*.

UNGUENTUM A'QUE RO'SÆ, *Oint'ment of Rose water*, *Cold Cream*. (*Aq. Rosæ*, fʒij; *Ol. amygdal.* ʒiiiss; *Cetacei*, ʒi; *Ceræ albæ*, ʒi. Melt together, in a water bath, the oil, spermaceti, and wax; then add the rose-water, and stir until it is cold. Ph. U. S.) Cooling to irritated surfaces.

UNGUENTUM ARMARII, Hoplochrysa—u. Arthanite, see Arthanita—u. Articulare, Synovia.

UNGUENTUM ATROP'IE, *Ointment of Atrop'ia*. (*Atropiæ*, gr. viij; *Spir. rectif.* fʒss; *Adipis præparat*. ʒj M. Ph. B., 1864.) Smeared around the eye to dilate the pupil.

UNGUENTUM BASILICON FLATUM, *Ceratum resinæ*—u. *Basilicon nigrum*, *Unguentum resinæ nigre*—u. *Basilicon viride*, *Unguentum subac-tatis cupri*.

UNGUENTUM BELLADON'NÆ, *Oint'ment of Belladonna*. (*Ext. belladonnæ*, ʒj; *Aq. fʒss*; *Adipis*, ʒj; M. Ph. U. S.) An anodyne application to irritable ulcers, &c.

UNGUENTUM BENZO'INI, *Ointment of Ben'zoin*, *Ben'zoated Lard*. (*Benzoin*. ʒj; *Adipis*, ʒxvj. Heat them together; strain without pressing, and stir while cooling. Ph. U. S., 1863.) A substitute for lard in seasons when the latter may become rancid.

UNGUENTUM CALOMEL'ANOS, *Ointment of Cal'omel*. (*Calomel*, gr. lxxx; *Prepared Lard*, ʒi. Ph. B., 1864.) In chronic cutaneous affections.

UNGUENTUM CALOMELANOS COMPOSITUM, *Com' pound Oint'ment of Cal'omel*, *Ban'yer's Oint'ment*.

(Calomel, $\mathfrak{z}\text{ij}$; Burnt alum, $\mathfrak{z}\text{iv}$; Carbonate of lead, $\mathfrak{z}\text{iv}$; Venice turpentine, $\mathfrak{z}\text{vj}$; Spermaceti ointment, $\mathfrak{z}\text{iss}$.) In scalled head and other chronic cutaneous affections.

UNGUENTUM CANTHARIDIS, Unguentum lyttæ.

UNGUENTUM CERÆ (*al'be vel fla'væ*), *Unguentum simplex* (Ph. U. S., 1851), *Unguentum ad'ipis*, *Linimentum simplex*, *Wax Ointment*, *Ointment of Lard*, *Oil and Bees' wax*, (F.) *Onguent de cire*. Ceræ, $\mathfrak{z}\text{ij}$; Adipis, $\mathfrak{z}\text{viii}$ (Ph. U. S.):—or—Ceræ, $\mathfrak{z}\text{ij}$, *Ol. oliv.* $\mathfrak{z}\text{v}$. Emollient. A mild dressing. Also, the basis of most of the compound ointments.

UNGUENTUM CERATUM, *Ceratum Galeni*—u. *Cerussæ*, *Unguentum oxidi plumbi albi*, *Unguentum plumbi subcarbonatis*—u. *Cerussæ acetatis*, *Ceratum plumbi superacetatis*, *U. plumbi superacetatis*.

UNGUENTUM CETA'CEI, *U. Sper'matia Ceti seu al'bum*, *Linimentum al'bum*, *Spermace'ti oint'ment*, (F.) *Onguent de blanc de baleine*. (Cetacci, $\mathfrak{z}\text{vj}$, Ceræ albæ, $\mathfrak{z}\text{iv}$, *Ol. oliv.* $\mathfrak{f}\mathfrak{z}\text{ij}$. Ph. L.) The ordinary dressing for blisters and excoriations. See *Ceratum cetacei*.

UNGUENTUM CITRINUM, *U. de althæa*, *U. hydrargyri nitratiss*.

UNGUENTUM COC'CVLI, *Oint'ment of Cocculus Indicus*. (Take any quantity of *Cocculus Indicus*; separate and preserve the kernels; beat them well in a mortar, first alone, and then with a little lard; then add lard till it amounts altogether to five times the weight of the kernels. Ph. E.) Used to destroy vermin, and to cure scabies, and ringworm of the scalp.

UNGUENTUM CONI'LI, *Ointment of Hem'lock*. (Coni'fol. recent.; Adipis, $\mathfrak{aa}$ $\mathfrak{lbj}$. Boil the hemlock in the lard till it becomes crisp, and express through linen. Ph. L.) Used to irritable tumours and ulcers.

UNGUENTUM CONTRA SCABIEM WERLHOFFI, *Unguentum hydrargyri præcipitati albi*.

UNGUENTUM CREASOTI, *Oint'ment of Creasote*. (Creasot. $\mathfrak{f}\mathfrak{z}\mathfrak{ss}$; Adipis, $\mathfrak{z}\text{ij}$. Add the creasote to the lard previously melted by a moderate heat, and stir constantly till cold. Ph. U. S.) Used in chronic cutaneous affections.

It is double the strength in the Ph. B., 1864.

UNGUENTUM CUPRI SUBACETATIS, *Unguentum subacetatis cupri*—u. *Diapompholygos*, *Unguentum oxidi zinei impuri*.

UNGUENTUM DIGESTIVUM SIM'PLEX, *U. Terebinthinæ et ovorum vitell'i*, *Simple digestive oint'ment*, (F.) *Onguent digestif simple*. (Terebinthin. pur. 64 p. Vitell. ovorum, No. 2, vel 32 p. Mix, for a long time, in a glass mortar, and add, gradually, enough of the Oil of Hypericum (Mil'lepertuis) to make a soft ointment. Ph. P.) Detergent. It may be rendered more so by adding a little of the Acetate of copper.

UNGUENTUM EL'EMI, *Oint'ment of El'emi*. (Elemi, $\mathfrak{z}\mathfrak{z}$; Simple ointment, $\mathfrak{z}\text{ij}$. Melt, strain through flannel, and stir until the ointment solidifies. Ph. B., 1864.) Uses, the same as the next.

UNGUENTUM EL'EMI COMPOSITUM, *Unguentum e gum'mi el'emi seu de terebinthinâ et adip'ibus seu El'emi*, *Linimentum seu Bal'samum Arcæ'i*, *Compound Oint'ment of El'emi*, *Balsam of Arcæus*, (F.) *Baume d'Arcæus*. (Elemi, $\mathfrak{lbj}$, Terebinthinæ vulgaris, $\mathfrak{z}\mathfrak{x}$, *Sevi præparati* $\mathfrak{lbj}$, *Olivæ olei*, $\mathfrak{f}\mathfrak{z}\text{ij}$. Melt the elemi with the suet; remove it from the fire; and mix in the turpentine and oil. Strain the mixture through a linen cloth. Ph. L.) Stimulant and digestive. To keep open issues, &c.

UNGUENTUM EPISPASTICUM DE DAPH'NE GNID'IO, *A'deps Cortic'is daph'næ gnid'ii medicatus*, (F.) *Pommade de Garou*. (Adip. præpar. 320 p., ceræ, 32 p., Cort. daphn. gnid. 128 p. Melt the fat and wax; and add the bark softened in water. Boil until the water is evaporated; then pass through cloth. Ph. P.) To keep open blisters, issues, &c. See *Unguentum Mezerei*.

UNGUENTUM EPISPASTICUM FORTIUS, *Cerate of cantharides*—u. *Epispasticum mitius*, *Unguentum lyttæ*—u. *Epispasticum viride*, *Unguentum lyttæ medicatum*—u. *Flavum*, *Unguentum de althæa*.

UNGUENTUM GAL'LÆ, *Oint'ment of Galls*, (*Gallæ* in pulv. $\mathfrak{z}\text{ij}$; Adipis, $\mathfrak{z}\text{vij}$. M. Ph. U. S.) As an astringent in piles, indolent ulcers, &c.

UNGUENTUM GAL'LÆ COMPOSITUM, *U. Gal'læ et O'pii*, *U. Gallæ cum O'pio*, *Compound Oint'ment of Galls*. (Gall. in pulv. subtil. $\mathfrak{z}\text{vj}$, Adipis, $\mathfrak{z}\text{vj}$, *Op'ii pulv.* $\mathfrak{z}\text{ss}$. M. Ph. L.) Used in painful hemorrhoids. $\mathfrak{z}\mathfrak{ss}$ or $\mathfrak{z}\text{ij}$ of camphor is sometimes added.

UNGUENTUM E GUMMI ELEMI, *Unguentum compositum*—u. *Hellebori albi*, *Unguentum veratri*.

UNGUENTUM HYDRARGYRI, *U. cæru'lem seu Neapolita'num seu mercuria'le*, *Ceratum mercuria'le*, *A'deps Hydrargyri medicatus*, *Mercurial Oint'ment*, *Blue oint'ment*, *Uction*, (F.) *Onguent mercuriel ou Napolitain*.

The Strong Mercurial Oint'ment, *Unguentum Hydrargyri fortius* of the London Pharmacopœia, *U. Hydrargyri* (Ph. U. S.), is thus made—Hydrarg. $\mathfrak{z}\mathfrak{xxiv}$, Adipis, *Sevi*, $\mathfrak{aa}$ $\mathfrak{z}\mathfrak{xij}$. Rub till the globules entirely disappear.

The Milder Mercurial Oint'ment, *U. Hydrargyri Mit'ius*, is made by adding, to one pound of the strong, two pounds of lard.

Mercurial ointment is antisyphilitic and discutient. It is rubbed on the inside of the thigh, in venereal affections; especially in those cases where mercury, given internally, runs off by the bowels. The stronger ointment is used for this purpose. The weaker is employed as a dressing, or as an application to cutaneous eruptions.

UNGUENTUM HYDRARGYRI AMMONIATI, *Unguentum hydrargyri præcipitati albi*—u. *Hydrargyri calcis albæ*, *Unguentum hydrargyri præcipitati albi*—u. *Hydrargyri compositum*, *Unguentum hydrargyri nitratiss*—u. *Hydrargyri deutoxydi nitratiss*, *Unguentum hydrargyri nitricooxydi*.

UNGUENTUM HYDRARGYRI IO'DIDI, *Oint'ment of Iodide of Mercury*. (Hydrarg. Iodid. $\mathfrak{z}\text{ij}$, Ceræ alb. $\mathfrak{z}\text{ij}$, Adipis $\mathfrak{z}\text{vj}$. Ph. L.)

UNGUENTUM HYDRARGYRI IO'DIDI RU'BRI, *Oint'ment of Red Iodide of Mercury*. (Hydrarg. Iodid. Rubr. $\mathfrak{z}\text{ij}$, Ung. ceræ albæ $\mathfrak{z}\text{vij}$. Ph. D.) Both these ointments may be applied to scrofulous and indolent ulcers.

UNGUENTUM EX HYDRARGYRO MITIUS DICTUM CINEREUM, *Unguentum oxidi hydrargyri cinereum*.

UNGUENTUM HYDRARGYRI NITRA'TIS, *U. hydrargyri nitra'ti seu supernitrat'is hydrargyri seu hydrar'gyri compositum seu mercuria'le citrinum seu citrinum ad sca'biem*, *Bal'samum mercuria'le*, *A'deps nitrat'is hydrar'gyri medicatus*, *Oint'ment of Nitrate of Mercury*, *Citrine Oint'ment*, (F.) *Onguent de nitrate de mercure*, *O. citrin contre la gale*. (Hydrarg. $\mathfrak{z}\text{iss}$, Acid. nitric. $\mathfrak{z}\text{iiiss}$, *Olei bubuli* $\mathfrak{z}\mathfrak{xij}$, Adipis $\mathfrak{z}\text{ivss}$. Dissolve the mercury in the acid; and, while hot, add the oil (for which butter might be substituted) and melted lard. Ph. U. S.)

A milder ointment, *Ung. nitrat'is Hydrargyri mit'ius*, is made with triple the quantity of oil and lard. It is stimulant and detergent, and is much used as an application to herpes, porrigo, and cutaneous eruptions. The weaker ointment is sometimes used in ophthalmia tarsi, &c.

UNGUENTUM HYDRARGYRI NI'TRICOOX'YDI, *U. subnitrat'is hydrargyri seu Ox'yd'i Hydrar'gyri rub'ri*, *U. H. ox'idi rub'ri* (Ph. U. S.), *U. deutoxydi hydrarg.* *nitrat'is seu Hydrargyri rub'ri seu ex Hydrar'gyro oxyda'to ru'bro seu mercur'ii ru'brum seu ophthalmicum ru'brum seu subnitrat'is hydrargyri'ci*, *Bal'samum ophthalmicum ru'brum*, *Oint'ment of Nitric Ox'yd of Mercury*, *Red Precipitate Oint'ment*, *Gold'en Oint'ment*, *Sin'gleton's Eye Salve*. (Hydrarg. oxid. rubr. in pulv.

subtiliss. $\mathfrak{Zj}$; Ung. adipis $\mathfrak{Zj}$. Add the oxyd, reduced to a fine powder, to the melted ointment. Ph. U. S.) Stimulant and escharotic. Applied to foul ulcers, to inflammation of the tarsi, &c. &c.

Belleville's Ce'rate, which has been long used in cases of scalled head, and of chronic cutaneous diseases in general, is made as follows:—*Plumbi acet.* $\mathfrak{Zj}$; *Hydrarg. acid. rubr.* $\mathfrak{Zss}$; *Hyd. chlorid. mit.* $\mathfrak{Zij}$; *Cere albæ* $\mathfrak{Ziv}$; *Ol. oliv.* $\mathfrak{Zxj}$. Melt the last two, and add the others in fine powder.

UNGUENTUM HYDRARGYRI OXIDI RUBRI, Unguentum hydrargyri nitricooxydi—u. ex Hydrargyro oxydato rubro, Unguentum hydrargyri nitricooxydi.

UNGUENTUM HYDRARGYRI PRÆCIPITATI ALBI, U. Hydrargyri ammoniati (Ph. U. S.) seu submuriatis hydrargyri ammoniati seu antipsoricum Werlhofii seu ad sca'biem Zelleri seu mercuriale album seu mundificans Zelleri seu e mercurio præcipitato seu calcis hydrargyri albi, Ointment of white precipitate. (*Hyd. ammon.* $\mathfrak{Zj}$, Ung. simpl. $\mathfrak{Ziss}$. Melt the fat, and add the precipitate.) Detergent. Used incutaneous eruptions, to destroy vermin, &c.

UNGUENTUM HYDRARGYRI RUBRI, U. Hydrargyri nitricooxydi—u. Hydrargyri submuriatis ammoniati, U. Hydrargyri præcipitati albi—u. Hydrargyri subnitratis, U. hydrargyri nitricooxydi—u. Hydrargyri supernitratis, U. hydrargyri nitratis—u. Hydrargyri subnitratis, U. hydrargyri nitricooxydi—u. Infusi meloes vesicatorii, U. lyttæ.

UNGUENTUM IO'DINI, U. Iodin'ii, Ointment of Iodine. (*Iodin.* $\mathfrak{Hj}$; *Potass. ioidid.* gr. iv; *Aquæ Mvi*; *Adipis.* $\mathfrak{Zj}$. Rub the iodine and iodide first with the water until liquefied, and then with the lard until mixed. Ph. U. S.) Used as a local application to goutrous and other swellings.

UNGUENTUM IO'DINI COMPOSITUM, U. Iodin'ii compos'itum, U. Io'di compos'itum, Compound Ointment of Iodine. (*Iodin.* gr. xv; *Potassii ioidid.* gr. xxx; *Aq. M.* xxx; *Adipis.* $\mathfrak{Zj}$. Rub the iodine and iodide first with the alcohol and then with the lard until they are well mixed. Ph. U. S.)

UNGUENTUM IRRITANS, U. lyttæ.

UNGUENTUM LAURI'NUM, O'leum lauri'num, A'deps lauro medica'tus, Bay Ointment, (F.) O'nguent on Huile de laurier. (*Fol. lauri.* 500 p., *Adip. suilli.* 1000 p. Bruise in a marble mortar; evaporate all humidity by boiling; and add Bay berries, in powder, 500 p. Digest and pass through linen. Ph. P.) Reputed to be corroborant and nervine. It is simply emollient.

UNGUENTUM LENITIVUM, U. de althæa—u. Liliaceum, Crinomyron.

UNGUENTUM LYT'TÆ, Ung. canthar'idis seu vesicatorium seu irritans seu ad vesica'ta seu infu'si mel'oës vesicatorii seu epispasticum mit'tus, Blist'ering Salve, Ointment of Spanish Flies, Ointment of Infusion of Spanish Flies, Milder blist'ering Ointment. (*Cantharid.* in pulv. $\mathfrak{Zij}$; *Aquæ destillat.* Oss; *Cerat. resin.* $\mathfrak{Zviij}$. Boil down the water with the Spanish flies to one half, and strain; then mix the cerate with the strained liquor, and evaporate to the proper consistence. Ph. U. S.) It is stimulant and epispastic; and is chiefly used for keeping up the discharge from a blistered surface.

UNGUENTUM LYT'TÆ MEDICA'TUM, A'deps canthar'idibus medica'tus, Poma'tum seu Unguen'tum epispasticum vir'idè stim'ulans ac'ribus, Green blist'ering Ointment, (F.) Pommade épispastique verte. (*Pomati seu unguenti cantharid.* 64 p., *Unguent populei* 168 p., *Cere albæ* 256 p., *Capri acetat.*, *Ext. opii*, $\mathfrak{aa}$ 24 p. Melt the unguentum populeum, and wax, and add, as they cool, the oxyd of copper, cantharides, and opium. Ph. P.) Use, same as the last.

UNGUENTUM LYT'TÆ MEDICA'TUM A'LIUD, A'deps canthar'idibus medica'tus a'lius, Poma'tum epispasticum fla'vum et mit'tus, Yellow Blist'ering Ointment, (F.) Pommade épispastique jaune. (*Pulv. cantharid.* 120 p., *Adip. præp.* 1680 p., *Aquæ* 250 p. Melt the lard; mix the cantharides, and add the water at the same time; place the whole on a gentle fire; stir continually for two hours, and add a little water to replace that which has been evaporated. Pass through linen and strain. Keep it in a water-bath, and add powdered Turmeric, 8 p. Let it cool till the superfluous liquid separates; pour off and add Yellow wax, 250 p., Oil of Lemon, 8 p. Ph. P.) A complex preparation. Use, same as the last.

An ointment, not very philosophically combined, was used by Dr. Physick, which is known by the name of "Dr. Physick's Is'sue Ointment." It is made by boiling half an ounce of powdered Cantharides in two ounces of Rosewater, and evaporating until onehalf of the fluid is driven off, having previously added 15 grains of Tartar emetic. The decoction is to be strained, and three ounces of Olive oil, an ounce and a half of White wax, and one ounce of Spermaceti added to it, and the mixture then boiled till all the water is evaporated.

UNGUENTUM MELANICUM CAUSTICUM, Causticum Æthiopicum.

UNGUENTUM MERCURIALE, Unguentum hydrargyri—u. Mercuriale album, Unguentum hydrargyri præcipitati albi—u. Mercuriale citrinum, Unguentum hydrargyri nitratis—u. e Mercurio præcipitato, Unguentum hydrargyri præcipitati albi—u. Mercurii rubrum, Unguentum hydrargyri nitricooxydi.

UNGUENTUM MEZE'REI, Ointment of Meze'reon. (*Mezer. incis. transversim.* $\mathfrak{Ziv}$; *Adipis.* $\mathfrak{Zxiv}$; *Cere albæ.* $\mathfrak{Zij}$; moisten the meze'reon with a little alcohol, and beat in an iron mortar, until it is reduced to a fibrous mass; then digest it with the lard, in a saltwater bath, for 12 hours; strain forcibly, and allow the strained liquor to cool slowly, so that any undissolved matters may subside; from these separate the medicated lard; melt it with the wax at a moderate heat, and stir constantly till they are cold. Ph. U. S.) To keep open blisters, issues, &c.

UNGUENTUM MUNDIFICANS ZELLERI, Unguentum hydrargyri præcipitati albi.

UNGUENTUM MURIA'TIS HYDRAR'GYRI OXYGENA'to MEDICA'TUM, A'deps muriat'æ hydrar'gyri oxygena'to medica'tus, Cyrrillo's Ointment, (F.) Pommade de Cyrrillo. (*Hydrar. oxyzur.* 4 p.; *Adipis suilli præp.* 32 p.—Ph. P.) Antivenereal and alterative. Used in friction, &c. at a time.

UNGUENTUM NARDINUM, see Nardus Indica—u. Neapolitanum, Unguentum hydrargyri—u. Nervinum, Unguentum Sambuci—u. Nitricum seu oxygenatum, Unguentum acidi nitrosi—u. Nutritum, Unguentum plumbi superacetatis.

UNGUENTUM OPHTHALMICUM, A'deps ox'ido hydrargyri ru'bro et plum'bi acetat'è medica'tus, Re'gent's Pommade ophthalmique, (F.) Pommade ophthalmique. (*Butyr. recent.* *Aq. rosar.* lavat. p. 68; *Camphor.* p. 0.3; *Oxyd. hydrarg. rubr.*, *Acetatis plumbi*, $\mathfrak{aa}$ p. 4. Mix intimately.—Ph. P.) In chronic inflammation of the conjunctiva or tarsi.

UNGUENTUM OPHTHALMICUM, Unguentum oxidi zinci impuri—u. Ophthalmicum rubrum, Unguentum hydrargyri nitricooxydi.

UNGUENTUM O'PII, Ointment of O'pium. (*Opii pulv.* $\mathfrak{Hj}$; *Adipis.* $\mathfrak{Zj}$; *M.*—Ph. L.) To be used where the local application of opium by inunction is needed.

UNGUENTUM OX'IDI HYDRARGYRI CINE'REI, A'deps seu Unguen'tum ex hydrar'gyro mit'tus die'tum cine'reum, Ointment of gray oxyde of mer'cury, (F.) O'nguent gris. (*Oxyd. hydrarg. ciner.* p. j; *Adipis suilli.* p. iij.—Ph. E.) Properties

like the mercurial ointment. It is not much used.

UNGUENTUM OX'IDI PLUM'BI AL'BI, *U. cerus'se seu al'bum*, Ointment of white oxide of lead, *White ointment*. (*Ung. simplicis*, p. v.; *Oxyd. plumb. alb.* p. j.—Ph. E. and P.) Cooling and desiccative. Applied to excoriated and burnt surfaces.

UNGUENTUM OX'IDI PLUMBI CARBONATI, *U. Plumbi subcarbonatis*—*u. Oxidi zinci*, Unguentum zinci.

UNGUENTUM OX'IDI ZIN'CI IMPU'RI, *U. tūt'ie seu ophthalmicum seu diopompholygos*, *A'deps ox'idi zin'ci medica'tus*, *Eyesalve*, *Tūt'ly ointment*, Ointment of impure oxide of zinc. (*Linim. simpl.* p. v; *Oxyd. zinc. impur. præp.* p. j.—Ph. E.) Generally used in ophthalmia tarsi and atonic ophthalmia.

UNGUENTUM PI'CIIS LIQ'UIDÆ, *U. Pi'cis seu e Pi'cæ*, *Tur Ointment*. (*Picis liquidæ*, *Seri*, sing. f*bj.* Melt, and stir till cold.—Ph. U. S.) Stimulant and detergent. Used in tinea capitis, and other cutaneous eruptions.

UNGUENTUM PICIS NIGRÆ, Unguentum resinæ nigrae—*u. Plumbi acetatis*, Ceratum plumbi superacetatis—*u. Plumbi carbonatis*, Unguentum plumbi subcarbonatis.

UNGUENTUM PLUM'BI COMPOS'ITUM, *Compositum Ointment of Lead*. (*Emp. Plumbi*, f*ijj*; *Ol. oliv. f*℥*viiij*; *Crete. præpar.* ℥*vj*; *Acid. acet. dil. f*℥*vj*.) Dissolve the plaster in the oil with a slow fire; then add the chalk, and afterwards the acid, constantly stirring until they become cold.—(Ph. L.) Used as a dressing to flabby, indolent ulcers.

UNGUENTUM PLUM'BI IO'DIDI, Ointment of Iodide of Lead. (*Plumb. iodic.* ℥*j*; *Adipis*, ℥*viiij*; *M.*—Ph. L.) Used as a eutrophic in chronic, glandular, and other swellings.

UNGUENTUM PLUMBI OXYDI CARBONATI, Unguentum plumbi subcarbonatis.

UNGUENTUM PLUMBI SUBCARBONA'TIS, *U. plum'bi carbonat'is* (Ph. U. S.) seu *al'bum seu al'bum simp'lex seu carbonat'is plum'bici seu cerus'se seu ox'idi plum'bi carbonat'is*, *Cera'tum de cerus'sâ*, (F.) *Onguent blanc de Rhazes*, *O. de cêruse*, *O. de Tornamira*. (*Plumbi carbonat. in pulv. subtiliss. gr. lxxx*, *Unguent. simpl.* ℥*j*.) Add the carbonate of lead to the ointment previously softened over a gentle fire, and mix.—(Ph. U. S.) Chiefly used as a dressing for burns.

UNGUENTUM PLUM'BI SUPERACETA'TIS, *U. acetat'is plum'bi seu saturni'num seu nutrit'um seu cerus'se acetat'is*, *Bal'samum universa'le*, Ointment of sugar of lead, *Sat'urnine ointment*. (*Olive oil*, f*ss*; *White wax*, ℥*ss*; *Superacetate of lead*, ℥*ij*.) Cooling, desiccative. Called, also, *Bal'samum saturni'num*, *Buty'rum satur'n*, *Linimen'tum plum'ba'tum*, *Poma'tum satur'ni*, *Ung. triphar'macum*.

UNGUENTUM PLUMBICI CARBONATIS, Unguentum plumbi subcarbonatis.

UNGUENTUM POPULE'UM, *A'deps popav'eræ*, *hyoscyamo*, *et belladonnâ medica'tus*. (Fresh buds of the *Populus nigra*, 500 p., macerated in *Adeps euillus præp.* melted, 1500 p., for 24 hours. Preserve this. Take Black poppy leaves, *Belladonna*, *Hyoscyamus*, Black nightshade, aâ 128 p.; bruise in a mortar with the former preparation; boil until all the humidity is evaporated, and strain.—(Ph. P.) Anodyne. Applied to local inflammations, hemorrhoids, sore nipples, cancer, &c. Green elder ointment is often sold for this.

UNGUENTUM POPULEUM, see *Populus*.

UNGUENTUM POTASSÆ HYDRIDA'TIS, *U. potas'sii io'didi* (Ph. U. S.), Ointment of hydi'odate of potass. (*Potass. iodic. in pulv. subtil.* ℥*j*; *Aq. f*℥*j*; *Adipis*, ℥*j*.) Dissolve the iodide in the water, and mix the solution with the lard.—(Ph. U. S.) Half a drachm of this ointment is rubbed on scrofulous glands, goitre, &c.

UNGUENTUM POTASSII IO'DIDI, Unguentum Po-

tassæ Hydi'odatis—*u. Pulveris meloes vesicatorii*, Cerate of cantharides—*u. Resinæ flavæ*, Ceratum resinæ flavæ.

UNGUENTUM RESINÆ NI'GRÆ, *U. resi'næ nigrum seu basil'icon nigrum seu Pi'cis nigrae seu Pi'cis tetrapharm'acum*, Black basil'icon, Black resin ointment, Ointment of pitch, Ointment of black pitch. (*Resin nigræ*, *Cere flavæ*, *Resinæ flavæ* aâ ℥*ix*; *Olei olivæ*, O*j.*—Ph. L.) Uses, the same as those of Ceratum resinæ or Yellow basilicon.

UNGUENTUM RESINÆ PINI SYLVESTRIS COMPOSITUM, Unguentum de Althæa—*u. Resinosum*, Ceratum resinæ flavæ—*u. Resumptivum*, Unguentum de althæa.

UNGUENTUM SABINÆ, Ceratum Sabinæ.

UNGUENTUM SAMBU'CI, *U. sambuci'num seu nervi'num*, Green ointment, *El'der ointment*, (F.) *Onguent de sureau*. (*Sambuci flor. f*ijj**; *Adipis præp. f*ijj.** Boil.—Ph. L.) Emollient.

UNGUENTUM SATURNI'UM, Unguentum plumbi superacetatis—*u. ad Scabiem Zelleri*, Unguentum hydrargyri præcipitatis albi—*u. Simplex*, Ceratum Galeni, Unguentum ceræ—*u. Solidum* de cicutâ, Emplastrum cicutæ—*u. Spermat'is ceti*, Unguentum cetacei.

UNGUENTUM STRAMO'NII, Ointment of Stramonium. (*Ext. stramon. fol.* ℥*j*; *Aq. f*℥*ss*; *Adipis*, ℥*j*.) Rub the extract with a little water until uniformly soft, and then with the lard.—(Ph. U. S.) An anodyne ointment to irritable ulcers, hemorrhoids, &c.

UNGUENTUM SUBACETA'TIS CU'PRI, *U. cu'pri subaceta'tis* (Ph. U. S.) seu *deter'gens seu basil'icm vir'idè seu æri'ginis*, *Bal'samum seu Cera'tum seu Unguent'um vir'idè*, Ointment of subac'etate of copper, *Ver'digris ointment*. (*Cu'pri subacet. in pulv. subtiliss.* ℥*j*; *Ung. simpl.* ℥*xv*.) Add the subacetate to the ointment previously melted with a moderate heat, and stir till cold.—(Ph. U. S.) Detergent and escharotic.

UNGUENTUM SULPHURA'TUM AD SCA'BIEM, *A'deps sulphurè et ammo'niæ muriat'è medica'tus*. (*Adipis*, 120 p.; *Sulph. lot.* 60 p.; *Ammonizæ muriat.*, *Aluminae sulphat.* aâ 4 p.; fiat unguentum.—Ph. P.) Antipruric.

Bail'ey's Itch Ointment is composed of olive oil, lard, nitre, alum, sulphate of zinc, and cinnamon; scented with oil of aniseed, oil of origanum, and oil of spike, and coloured with alkane root.

UNGUENTUM SULPHURA'TUM ALCALI'UM AD SCA'BIEM, *A'deps sulph'urè et carbonat'è potas'se medica'tus*, *Hel'merich's Al'caline Sulphurat'ed Itch Ointment*, *Pommade antiprurique de Helmerich*, *Pommade de Helmerich*. (*Adip. præp.* 800 p.; *Sulphur. lot.* 200 p.; *Carbonat. potass. pur.* 100 p.; fiat ung.—Ph. P.) Use, same as the last.

Bate'man's Itch Ointment resembles this. It is coloured with cinnamon, and scented with essence of bergamot.

UNGUENTUM SULPHURIS, *U. e sulph'urè*, *Sulphur ointment*. (*Sulph.* ℥*j*; *Adip.* ℥*ij*—Ph. U. S.) In Ph. B. (1864) sulphur, ℥*j*; lard, ℥*v*. Used chiefly in itch, and some herpetic affections.

UNGUENTUM SULPHURIS COMPOS'ITUM, *U. Antipruricum*, *Compound Sulphur Ointment*, *Itch Ointment*, *Jack'son's Itch Ointment*, (F.) *Pommade Antiprurique de Jasser*. (*Sulph. sublim. f*ss**; *Vetri' rad. contus.* ℥*ij*; *Potassæ nitrat.* ℥*j*; *Saponis mollis*, f*ss*; *Adipis*, f*ss*; *Ol. bergamot.* ℥*xxx*—Ph. L.) THE UNGUENTUM SULPHURIS COMPOS'ITUM of the Pharm. U. S. is made as follows:—*Sulphur.* ℥*j*; *Hydrargyri ammoniat.*, *Acid. benzoic.* aâ ℥*j*; *Ol. bergamot.*, *Acid. Sulphuricæ.* aâ f*℥j*; *Potassæ nitrat.* ℥*ij*; *Adipis*, f*ss*. To the melted lard add the other ingredients, and stir till cold. Use, the same as the last; but it is more stimulating.

UNGUENTUM SULPHURIS IO'DIDI, Ointment of Iodide of Sulphur. (*Sulphur. iodic.* ℥*ss*; *Adipis*, ℥*j*.) Rub the iodide with a little of the lard add

the remainder, and mix.—Ph. U. S.) An excellent application in chronic cutaneous affections.

UNGUENTUM SERSINUM, Crinomyon.

UNGUENTUM TABACI, *Tobacco Ointment*. (*Tabac, recent, concis.* ʒj; *Adipis*, lbj. Boil the tobacco in the lard, over a gentle fire, till it becomes friable; then strain through linen.—Ph. U. S. 1851.) In the Ph. U. S. 1863, it is directed to be made as follows, (*Tabac* in pulv. ʒss; *Adipis*, ʒviij; *Aq. q. s.* Moistens the tobacco with water, and, by percolation, pass water through it until four fluidounces are obtained. Evaporate to the consistence of a soft extract, and mix with the lard.) Applied to irritable ulcers, tinea capitis, &c.

UNGUENTUM TARTARI EMETICI, U. antimonii—u. Tartari stibiat, U. antimonii—u. e Tartaro potassæ stibiat, U. antimonii—u. Tartratis potassæ stibiat, U. antimonii.

UNGUENTUM TEREBINTHINÆ, *Ointment of Turpentine*. (*Oil of Turpentine*, fʒj; *Resin*, in coarse powder, gr. lx; *Yellow wax*, ʒss; *Prepared lard*, ʒss. Melt together. Ph. B. 1864.) Uses, the same as *Ceratum Resinæ* or *Yellow Basilicon*.

UNGUENTUM DE TEREBINTHINÆ ET ADIPIS, U. elemi compositum—u. de Terebinthinæ et ceræ, U. de althæa—u. Terebinthinæ et ovorum vitelli, U. digestivum simplex—u. Tetrapharmacum, Basilicon, U. resinæ nigre—u. Tripharmacum, U. plumbi superacetatis—u. Tutie, U. oxidi zinci impuri.

UNGUENTUM VERATRÆ, *U. Veratræ al'bi* (Ph. U. S.) seu *helleb'ori al'bi*, *White hellebore ointment*. (*Veratr. pulv.* ʒij; *Adipis*, ʒviij; *Olei limon.* mxx.) Used in cases of itch, where sulphur is objected to. It is not equally efficacious.

The principal ingredients in the EDINBURGH OINTMENT are white hellebore and muriate of ammonia.

UNGUENTUM VERATRÆ, *Ointment of Veratris*, (*Veratr. gr. xx; Adipis*, ʒj. Mix. Ph. U. S. 1863.) In the British Pharmacopœia (1864) it is directed to be made of *Veratris*, gr. viij; *Prepared lard*, ʒj; *Olive oil*, fʒss; well mixed together. To be rubbed on the part in neuralgic and other pains.

UNGUENTUM AD VESICATAM, U. lyttæ—u. ad Vesicatoria, Cerate of cantharides—u. Vesicatorium, Unguentum lyttæ—u. Viride, Unguentum subacetatis cupri.

UNGUENTUM ZINCI, *U. oxidi zinci seu zin'ci oxidi* (Ph. U. S.), *Zinc ointment*. (*Zinci oxid.* gr. lxxx; *Adipis*, ʒi.) Use, the same as that of the *Unguentum tutie*.

UNGUENTUM ZINCI OXIDI, U. zinci.

UNGUIS, 'a nail,' Hippocampus minor, Nail, Onyx, Pterygion—u. Avis, Hippocampus minor—u. Halleri, Hippocampus minor—u. Odoratus, Blatta Byzantia.

UNGUIS OS. *Os lacrymale*. *Lachrymal bone*. A small, quadrilateral, very thin, and semi-transparent bone, which has been compared to a human nail, and is situate at the anterior and inner part of the orbit. It aids in the formation of the lachrymal gutter and the nasal duct. It is articulated with the frontal, superior maxillary, ethmoid, and inferior spongy bones, and is developed by a single punctum ossificationis.

UNGULA CABALLINA, 'horse's hoof,' Tus-silago.

UNGULÆ PALPEBRARUM, 'hoofs of the eyelids,' see Tarsus.

UNHEALTHY, Sickly.

UNICORN ROOT, Chamælerium luteum.

UNILOCULAR, *Unilocularis*, (unus, 'one,' and locus, (diminutive of locus,) 'a cell.') Having one cell or cavity; as *bilocular* means having two cells or cavities; *multilocular*, having many.

UNIO, gen. *Unio'nis*, Pearl.

UNIOCLUS, (unus, 'one,' and oculus, 'eye,') Cyclops, Monoculus.

UNION BY THE FIRST AND SECOND INTENTIONS, see Intention.

UNIPARIENS, Uniparus.

UNIP'AROUS, *Unipa'rient*, *Unipa'riens*, (unus, 'one,' and pario, 'I bring forth,') Producing one at a birth.

UNIPOLAR, (unus, 'one,' and polus, 'pole,') see Neurine.

UNISE'MA DELTIFO'LIA, (unus, 'one,' and semen, 'seed' [?], *Pontede'ria corda'ta*, *Shovel Pick'erehweed*, *Pond Shorel*, *Shovellett*, *Water Plantain*. An indigenous plant, Ord. Pontederiaceæ, *Sex. Syst.* Hexandria Monogynia, which grows in water, and flowers from June to August. The root is emollient and astringent.

UNITAIRES (MONSTRES) (F.), (from unitas, 'oneness,') A class of monsters in which are found only the complete or incomplete elements of a single individual; in contradistinction to *Compound monsters*, (F.) *Monstres composés*, in which there are the complete or incomplete elements of more than one.

UNITING BANDAGE, (from unire, unitum, (unus, 'one,') 'to join into one,') (F.) *Bandage unissant*. A bandage for bringing the edges of a wound into approximation. The arrangement of such a bandage differs according as the wound is transverse or longitudinal.

UNIVOCAL GENERATION, (unus, 'one,' and vox, gen. vocis, 'voice,') see Generation.

UNOCULUS, (unus, 'one,' and oculus, 'eye,') *Borgne*, Cyclops, Monoculus.

UNORGANIZED, Inorganic.

UNSTRIPED MUSCULAR FIBRE, see Muscular fibre.

UNWELL, see Indisposition. 'To be unwell,'—and, in Scotland, 'to alter,'—is, popularly, to menstruate.

UPAS. A word used in the Archipelagos of Molucca and Sunda, and signifying *Veg'etuable Poi'son*; *I'po*, *Bo'hon*, *Po'hon*, or *Boa* or *Bo'on Upas*, *Po'hon Antiar*, *Ants'jar*. A substance used by the natives of those places for poisoning their arrows. The smallest quantity is capable of destroying the largest animal. The tree which produces it is said to be the *Antiaris* seu *I'po Toxicaria*, *Toxicaria Macassarien'sis*. Another species of upas, *U. Tienté*, *Tshettik* or *Tjették*, is the produce of a *Strychnos*, *S. Tienté*. *Upas antiar*, like corroval and vao, the South American arrow poisons, appears to kill by paralyzing the heart; whilst *U. tienté* causes the most violent convulsions. Many idle stories are told regarding the upas tree, most of which are fabulous.

URACHUS, *U'racus*, *Urinaculum*, (uron, and εχω, 'to contain,') (F.) *Ouraque*. The urachus, in certain animals, is a long membranous canal, which arises from the bladder, makes its exit from the abdomen by the umbilicus, and terminates in the pouch called *allantois*. According to Sabatier, this is the arrangement of the urachus in the human fetus; but Bichat and other anatomists consider, that when the canal exists in it, it is a malformation. They regard it as a kind of suspensory ligament of the bladder, extending from the top of that organ to the umbilicus; and, at the umbilicus, confounded with the abdominal aponeurosis.

URACRATIA, (uron, a priv., and κραα, 'power,') Enuresis.

URACUS, Urachus.

URÆMIA, *Uræ'mia*, (F.) *Urémie*, *Uroémie*, (uron, and αμα, 'blood,') A condition of the blood in which it contains, or is presumed to contain, urea, and to give rise to sundry morbid phenomena implicating the nervous centres more especially.

URÆMIC, *Uræ'micus*, (F.) *Urémique*. Relating or appertaining to uræmia

URALI, Curare.

URANA, (from *uron*.) Ureter.

URANE, (*ουρανν*.) Urinal.

URANISCONITIS, (*uranisens*, and *itis*.) Palatit'is, (F.) Palatite. Inflammation of the palate.

URANIS'COPLASTY, *Uraniscoplas'tiōē*, (*uranisens*, and *πλαστικός*, 'forming.') The operation of engraving in case of deficiency of the soft palate.

URANISCORRHAPHIA, (*uraniscus*, and *ῥαφν*, 'a suture,') Staphylorrhaphy.

URANISCUS, (*ουρανικός*, 'a round ceiling or canopy,') Palate.

URANOPLASTY, Staphyloplasty.

URANORRHAPHIA, Staphylorrhaphy.

URANUS, (*ουρανός*, 'heaven,') Palate.

URAPOSTEMA, gen. *Uraposte'matis*, *Abcess'us urino'sus*, *Uræde'ma*, (*uron*, and *αποστημα*, 'an abscess,') A tumefaction produced by the escape of urine. An abscess containing urine.

URARI, Curare.

URAS, Urate.

URATE, *U'ras*, *Lith'ate*, *Li'thas*, (from *uron*.)

A generic name for salts, formed by the combination of uric or lithic acid with different bases. *U'rate* or *Lith'ate* of *So'da*, for example, is found in arthritic and certain urinary concretions:—*U'rate* or *Lith'ate* of *Ammo'nia* in certain urinary deposits.

URE'A, or U'REA, *Ure'um*, *U'ren*, *Uren'ium*, *U'ricum*, *Mate'ria urino'sa*, *Principi'um seu Sal urino'sum*, *Neph'rin*, (F.) *Urée*, (from *uron*.) A white, pearly, brilliant substance, in transparent plates, inodorous, and of a cool taste; very soluble in water; deliquescent, and soluble in alcohol. It exists, in great quantity, in the urine; and is an essential part of it. It has been used as a diuretic. Dose, gr. xv to ℥j. Nitrate of urea has been used in like cases, in doses of 2 to 10 grains in the day, in solution or in pill.

URE'AL or U'REAL, *Ureal'is*. Relating or appertaining to urea.

UREC'HYSIS, *Urech'ysis*, (*uron*, and *εκχυσις*, 'effusion,') Effusion of urine into the areolar membrane.

URED'O, gen. *Ure'dinis*, (*urere*, 'to burn,') Urticaria.

URÉE, Urea.

UREMA, gen. *Ure'matis*, (*ουρημα*.) Urine.

URÉVIE, Uræmia.

URÉMIQUE, Uræmic.

UREN, Urea.

URENIUM, Urea.

URENS, (from *urere*, 'to burn,') Caustic.

UREORRHŒA, (*uron*, and *ῥεω*, 'to flow,') Diabètes.

URESIÆSTHESIS, (*uresis*, and *αἰσθησις*, 'feeling,') Diuresiæsthesia.

URESIS, (*ουρησις*.) Micturition.

URETER, (*ουρητηρ*.) *U'ra*, *Ve'na al'ba re'num*, *Canal'is ner'veus fistulo'sus re'num*, *Duct'us seu Duct'or Uri'næ*, *Vas urina'rium*, (F.) *Urêtère*. A long, excretory canal—membranous and cylindrical—which conveys the urine from the kidney to the bladder. It extends, obliquely, between the pelvis of the kidney, with which it is continuous, and the *bas-fond* of the bladder, into which it opens. It enters at the posterior and inferior part, for some distance between the coats, and opens into the cavity of the bladder, at the posterior angle of the *Trigone vésical*. The ureter is formed of an exterior, white, opaque, and fibrous membrane; and of an inner, which is mucous.

URETERAL'GIA, (*ureter*, and *αλγος*, 'pain,') Pain in the course of the ureter, —as from calculus, *Ureteral'gia calculo'sa*.

URETERALGIA CALCULOSA, see Ureter'algia—u. Inflammatoria, Ureteritis.

URÊTÈRE, Ureter.

URETER'ICUS, (from *ureter*.) An epithet for ischury, &c., arising from an affection of the ureter.

URETERIT'IS, *Ureteral'gia inflammato'ria*, *Ureterophlegma'sia*, (*ureter*, and *itis*.) Inflammation of the ureter: generally caused by calculus.

URETEROLITH'ICUS, (*ureter*, and *λίθος*, 'a stone,') An epithet applied to ischury or other affection caused by a stone in the ureter.

URETEROLITHUS. Calculus in the ureter.

URETEROPHLEGMASIA, (*ureter* and *phlegmasia*, 'inflammation,') Ureteritis.

URETEROPHLEGMA'TICUS, (*ureter*, and *φλεγμα*, 'phlegm or mucus,') Any affection caused by mucus accumulated in the ureter.

URETEROPY'ICUS, (*ureter*, and *πυρ*, 'pus,') An epithet applied to affections caused by the presence of pus in the ureter.

URETERORRHAG'IA, (*ureter*, and *ῥαγν*, 'rupture,') Hemorrhage from the ureter.

URETEROSTOMAT'ICUS, (*ureter*, and *στομα*, 'orifice,') That which is dependent upon obstruction of the orifice of the ureter.

URETEROTHROMBOÏ'DES, (*ureter*, *θρομβος*, 'grumous blood,' and *εἶδος*, 'resemblance,') Ischury from grumous blood in the ureter.

URE'THRA, (*ουρητρα*, from *ουρεω*, 'to pass the urine,') *Canal'is seu Meatus seu Duct'us urina'rius*, *U'rias*, *Ure'tra*, *Fist'ula urina'ria*, *Pter urina'rium seu Uri'næ*; (F.) *Urêthre*, *Urêtère*. The excretory ducts for the urine in both sexes; and for the sperm in man. In the latter, it is 8 or 10 inches long; and extends from the neck of the bladder, *os'tium vesic'al'e*, to the extremity of the penis, in which its external aperture, *os'tium cuta'neum*, is situate. In the male urethra, three portions are distinguished:—1. A *prostat'ic por'tion*, near the bladder, about an inch and a quarter in length, which traverses the prostate gland: 2. A *mem'branous por'tion*, *Isth'mus ure'thræ*, *Pars ure'thræ membranacea*, from 8 to 10 lines long, which is separated from the rectum by the areolar tissue only:—3. A *spongy por'tion*, which expands, anteriorly, to form the glands, and which commences at the *bulb of the urethra*. Before opening externally the canal has a dilatation, called the *Fos'sa seu Fo'vea seu Fossic'ula navicular'is seu Morgagni'i*. On the interior of the urethra two white lines are perceptible—the one above, the other below. The latter commences at the *verumontanum*; and at the surface of this, the *ejaculatory ducts*, and *prostatic ducts* open. The urethra is lined, in its whole extent, by a mucous membrane; on which are seen, particularly at the bulb, the orifices of a number of follicular ducts, *Lacunæ* or *Sinusces of Morgagni*, (F.) *Lacunes de Morgagni*, proceeding from or constituting the *Glands of Morgagni* or of *Littre*, (F.) *Glandes de Morgagni* ou de *Littre*, *Glandule Littria'næ*. At about an inch and a half from the opening of the meatus, one of these lacunæ is generally found, much larger than the rest, and is named *Lacuna Mag'na*. The arteries of the urethra are numerous, and proceed from the internal pudic,—the largest branches entering the bulb. The veins follow the course of the arteries. The lymphatics pass into the inguinal and hypogastric plexuses. The nerves come from the pudic and gluteal. In the female, the urethra is only about an inch long. It is wider than in man, and susceptible of great dilatation. It is very wide at its commencement; and descends obliquely forwards, to terminate at the top of the external aperture of the vagina by the *meatus urina'rius*, *os'tium vagina'l'e ure'thræ*. In its course, it is slightly curved, the concavity being forwards.

URETHRA, BLIND DUCTS OF THE, see Cæcus—u. Mucous lacunæ of the, see Cæcus.

URETHRAL, or **URETHRAL**, *Urethra'lis*, *Ur'ibicus*. Relating to the urethra.

URETHRAL'GIA, (*urethra*, and *algos*, 'pain.') Pain in the urethra. Gonorrhœa.

URETHRE, *Urethra*.

URETHREMPHRAXIS, *Urethrophraxis*.

URETHRITIS, (*urethra*, and *itis*.) Inflammation of the urethra. Gonorrhœa.

URETHROBLENNORRHŒA, (*urethra*, *βλανα*, 'mucus,' and *ῥευ*, 'I flow.') Gonorrhœa; gleet.

URETHROCYSTAUCHENOTOMIA, (*urethra*, *cystauchen*, 'neck of the bladder,' and *τομήν*, 'to cut,') see Lithotomy.

URETHROCYSTEOANEURYSMATOTOMIA, (*urethra*, *κυστίς*, 'the bladder,' *ανευρύσμα*, 'dilatation,' and *τομήν*, 'to cut,') see Lithotomy.

URETHROCISTEOTRACHELOTOMIA, (*urethra*, *κυστίς*, 'the bladder,' *τραχηλός*, 'neck,' and *τομήν*, 'to cut,') see Lithotomy.

URETHROILEMORRHAGIA, (*urethra*, *αἷμα*, 'blood,' and *ῥήγνυμι*, 'to break forth,') *Urethrorrhagia*.

URETHROPHRAX'IS, *Urethrophraxis* (*urethra*, and *φρασσα*, 'I obstruct,') Obstruction of the urethra.

URETHROPLAS'TIC, *Urethroplasticus*, (*urethra*, and *πλασσω*, 'I form,') An epithet given to the operation, *Urethrophlasty*, which supplies defects in the canal of the urethra through which the urine escapes.

URETHRORAPHY, *Urethrorrhaphy*.

URETHRORRHAGIA, *Urethrorrhœa*, *Urethrorrhœa*, *Caulorrhœa* seu *Hæmatu'ria stillatit'ia*, (*urethra*, and *ῥήγνυμι*, 'I break out,') Hemorrhage from the urethra.

URETHRORRHAPHY, *Urethrorrhap'ia*, *Urethrorraphy*, (*urethra*, and *ῥαφή*, 'a suture,') 'suture of the urethra.' The operation of closing a urethral fistula by suture.

URETHRORRHŒA, (*urethra*, and *ῥευ*, 'I flow,') A discharge of any kind from the urethra.

URETHROSCOPE, *Urethroscop'ium*, (*urethra*, and *σκοπεω*, 'I examine,') An instrument on the principle of the ophthalmoscope, laryngoscope, &c., for inspecting the interior of the urethra.

URETHROSTENOSIS, *Strictu'ra*, *ure'thræ*, *Urostenosis*, *Urethrosten'ia*, (F.) *Uréthrosténie*, *Rétrécissement de l'Urèthre*, (*urethra*, and *στενωσις*, 'narrowness,') Stricture of the urethra.

URETHROTOME, *Urethrot'omus*, (F.) *Uréthrotome*, (*urethra*, and *τομήν*, 'I cut,') An instrument so called by Le Cat, with which he divided the skin, and laid open the urethra, in the operation for lithotomy. Also, an instrument for dividing strictures of the urethra.

URETHROTOMY, *Urethrotom'ia*, *Stricturot'omy*. An operation, which consists in opening the urethra with the *Urethrotome*, for the removal of stricture. See *Boutonnière*.

URETHRYMENO'DES, (*urethra*, *ὑμν*, 'a membrane,' and *odes*.) That which is caused by a membrane formed in the urethra.

URETICUS, (*ουρητικός*.) Diuretic, Urinary.

URETRA, *Urethra*.

URÈTRE, *Urethra*.

URETRIS, (*ουρητρίς*.) Urinal.

UREUM, *Urea*.

URGINEA SCILLA, *Scilla*.

URHÆMIA, *Uræmia*.

URIAGE, MIN'ERAL WA'TERS OF. Simple sulphurous springs in the department of Isère, France, which are visited from the 15th of May to the 15th of September.

URIAS, gen. *Uriadis*, (from *uron*,) *Fistula* (urinary), *Urethra*.

URIASIS, (from *uron*,) *Lithia*, *Uresis*.

URIC ACID, *Ac'idum uricum* seu *urolith'icum* seu *lith'icum*, *Lith'ic*, *Uryl'ic*, *Cal'culous*,

Bezoâr'dic, *Lithin'sic* or *Lithen'ic Acid*, (F.) *Acide urique* ou *lithique* ou *lithénique* ou *lithiasique* ou *de l'urine*. An acid which exists in human urine, chiefly, if not wholly, in the form of urate of soda or urate of ammonia, and which in combination constitutes the greater part of urinary calculi. It is very sparingly soluble, and often deposited in combination from the urine in health. See *Calculi*, *urinary*.

URIC OX'IDE, *Ox'idum uricum*, *Xan'thic ox'ide*, *Xan'thine*, *U'rous ac'id*. A substance not discovered in healthy urine, although probably bearing some relation to the yellow colouring matter. It is a very rare ingredient of calculous concretions, and little is known either of its chemical or pathological history.

URICUM, *Urea*.

URICUS, (from *uron*,) *Urinary*.

URIDRO'SIS, (*uron*, and *ιδρωσ*, 'sweat,') *Urinous sweating*; (F.) *Sueur urinaire*.

URINA, *Urine*—*u. Cerebella*, see *Cerebella*—*u. Chyli*, see *Urine*—*u. Cibi*, see *Urine*—*u. Diabetica*, *Urine*, *diabetic*—*u. Febrilis*, *Urine*, *febrile*—*u. Furfuracea*, see *Furfuraceus*—*u. Galactodes*, see *Galactodes*—*u. Genitalis*, *Sperm*—*u. Jumentosa* seu *Jumentaria*, *Urine*, *jumentous*—*u. Mucosa*, *Cystirrhœa*—*u. Nephelodes*, see *Nepheloid*—*u. Oroboides* seu *Oroboides*, see *Oroboides*—*u. Pericardii*, see *Pericardium*—*u. Potûs*, see *Urine*—*u. Sanguinis*, see *Urine*—*u. Vaccæ*, *Urine*, *cows'*.

URINACULUM, *Urachus*, *Urinal*.

URINÆ PRÔFLUVIUM, 'a flux of urine,' *Diabetes*—*u. Stillicidium*, *Strangury*—*u. Substillum*, *Strangury*—*u. Suppressio*, *Ischuria*.

URINAIRE, *Urinary*.

URINAL, *Urinato'rium*, *Chernib'ium*, *U'rand*, *Ure'tris*, *Urinac'ulum*, *Urodochi'um*, *Urod'ochê*, *Urodoche'um*, *Matra'cium*, *Mat'ula*, *Matel'la*, (from *urina*.) A name given, 1. To certain vessels, used to pass the urine into. 2. To reservoirs, of different shapes and character, which are adapted to the penis, in cases of incontinence of urine, and which are intended to receive the urine as it flows off.

URINALIS, *Diuretic*, *Urinary*.

URINARIA, *Antirhinum linaria*, *Leontodon taraxacum*.

URINARY, *Urina'rius*, *Urina'lis*, *Urino'sus*, *U'rius*, *Uret'icus*, (F.) *Urinaire*, (from *urina*.) That which relates to the urine.

URINARY APPARATUS, *Urinary organs*.

URINARY BLAD'DER, *The Blad'der*, *Vesi'ca Urina'ria*, *Cys'tis*, *Urocys'tis*, (F.) *Vessie*. A musclemembranous reservoir, intended for the reception of the urine, until the accumulation of a certain quantity solicits its excretion. The bladder is situate in the hypogastric region, between the rectum and pubis in man; and between the pubis and vagina in the female. Its upper region is not covered, except by the peritoneum, posteriorly. From its centre arises the *urachus*. Its lower region is bounded, anteriorly, by the prostate, in man; and rests on the vesiculæ seminales and rectum. In the female, it corresponds to the vagina. The lowest portion of this region bears the name *Bas-fond*. The anterior surface of the organ is not covered by peritoneum. It is attached to the symphysis pubis by a fibrocellular expansion called the *anterior ligament*. The *posterior ligaments* or *rectovesical folds*—*pliœ semilunares*—are duplicatures of the peritoneum on its posterior surface, which extend from this surface to the rectum in the male, and to the uterus in the female. The bladder is composed of a serous membrane, formed by the peritoneum;—of a muscular membrane; of a somewhat thick layer of areolar tissue; and is lined on its inner surface by a mucous membrane, which is sometimes raised up, here and there, in long promi-

nences or *columns* by the muscular fibres beneath. To this condition, the French anatomists give the term—*vesicæ à colonnes*. In the *bas fond* the two ureters open, and the urethra commences; and the triangular space, which the apertures of these three canals leave between them, is called, by the French, the *Trigone vésical* or *Vésical triangle*, *Trigonus seu Trigōnum vesi-cæ seu Lien-taud'i*, *Corpus trigonum*. In the orifice of the urethra, which is also called the neck of the bladder, there is, at its lower part, a more or less prominent tubercle, which is formed by the anterior angle of the *Trigone vésical*, and is called the *Vesical nodule*, *Uvula vesi-cæ*, (F.) *Luetta vésicale*.

The arteries of the bladder proceed from the hypogastric, umbilical, sciatic, middle hemorrhoidal, and internal pudic. Its veins, which are more numerous than the arteries, open into the hypogastric venous plexus. Its nerves emanate from the sciatic and hypogastric plexus; and its lymphatic vessels pass into the hypogastric ganglia.

URINARY CALCULI. The calculi which form in the urinary passages. See Calculi, urinary.

URINARY ORGANS, Urinary apparatus, Or-gana uropœitica, Apparatus uropœiticus. The organs concerned in the formation of urine—the kidneys. The common acceptance includes, also, the urinary passages.

URINARY PASSAGES, (F.) Voies urinaires. The aggregate of canals and cavities intended to contain the urine and convey it externally. These ways are composed of excretory ducts, which form the tubular portion of the kidney; of calices; pelvis; ureters; bladder; and urethra.

URINATION, (from *urina*), Micturition. M. Béraud elaborately defines it after Ch. Robin,—a function, the result of which is the expulsion of matters principally solid, but held in solution, which have become improper for nutrition.

URINATORIUM, (from *urina*), Urinal.

URINE, *Urina*, (ουρον), *Uron*, *Ure-ma*, *Lo-tium*, *Lot*, *La-pis aureus*, *Omich'ma*, *Recremen-ta seu Retrimen-ta vesi-cæ*, *Aqua pe'dum*, *Aqua*, *Water*, (Old Eng.) *Stale*, *Lee*, *Chamber-lee*, (Prov.) *Herrin*, *Netting*, *Saag*. An excremental fluid; secreted by the cortical part of the kidney; filtered through the tubular portion; poured *guttatim* from the apices of the tubular papillæ into the pelvis of the kidney; and transmitted by it to the ureters; which convey it slowly, but in a continuous manner, into the bladder, where it remains deposited, until its accumulation excites a desire to void it. The excretion of the fluid takes place through the urethra; and is caused by the action of the abdominal muscles and diaphragm and the contraction of the fibrous coat of the bladder. Urine is transparent; of a citron-yellow colour; of a peculiar odour, and of an acid, saline, and slightly bitter taste. That which is passed some time after taking fluid, is less coloured, and less odorous and dense than that which is voided 7 or 8 hours after eating. The first is the *Urina potûs*, (F.) *Urine de la Boisson* ou des *Boissons*, *Urine of drink*—the latter *Urina sanguinis*, (F.) *Urine de la digestion*, *Urine de la coction*, *Urine cuite ou du Chyle*, *Coc'ted u'rine*, *Urine of digestion*. Three distinct varieties of urine may, according to Dr. Golding Bird, be recognised: *First*, That passed some little time after drinking freely of fluids, which is generally pale, and of low specific gravity—1.003 to 1.009—*Urina potûs*. *Secondly*, That secreted after the digestion of a full meal; s. g. 1.020 to 1.028, or even 1.030—*Urina chyli seu cibi*. *Thirdly*, That secreted independently of the immediate stimulus of food and drink, as after a night's rest, which is usually of average density—1.015 to 1.025, and presents the essential characters of urine—*Urina sanguinis*.

The following are the ordinary constituents of healthy human urine:—urea, uric acid, [hippuric acid;] extractive matters, embracing alcohol extract, spirit extract, and water extract, with their respective constituents; mucus; brown colouring matter of the urine, (hæmaphæin; red colouring matter of the urine, (uroerythrin; carbonic, lactic, hydrochloric, sulphuric, phosphoric, silicic, and hydrofluoric acids; soda; potassa; ammonia; lime; magnesia; and peroxide of iron. The proportion of solid matters in 1000 parts of urine may vary from 30 to 70 and upwards. The following table expresses the relative amounts of the different components in every 100 parts of the solid matter, according to the analysis of different observers:

	Berzelius.	Lehmann.	Simon.	Marchand.
Urea,	45.10	49.68	33.80	48.91
Uric Acid,	1.50	1.61	1.40	1.59
Extractive matter,				
Ammonia Salts,				
and Chloride of Sodium.	36.30	28.95	42.60	32.49
Alkaline Sulphates,	10.30	11.58	8.14	10.18
Alkaline Phosphates,	6.88	5.96	6.50	4.57
Phosphates of lime and magnesia.	1.50	1.97	1.59	1.81

Its average specific gravity is about 1.015. The quantity passed by the adult in the 24 hours varies. On the average, it may amount to two pounds and a half.

Dr. Golding Bird obtained the following results on examining 1000 parts of urine passed in health after 10 hours' fasting; and after dinner in the evening.

	Urina sanguinis.	Urina cibi
Specific gravity	1.016	1.030
Water, ...	962.72	930.10
Solids, ...	37.28	69.90
Urea, ...	14.30	24.40
Uric acid, ...	0.23	1.33
Fixed salts, chiefly chlorides, sulphates, and phosphates.	5.10	9.90
Organic matter, creatin, creatinin, colouring matter, and volatile salts.	17.80	34.27

The urine, when first passed, is generally acid, and this has been ascribed to uric acid, hippuric acid or lactic acid. It is, doubtless, however, owing to the phosphatic supersalts, and especially perhaps to the superphosphate or biphosphate of soda.

Sometimes, the urine, in health and disease, exhibits appearances that have received special names. Thus, that which is characterized by a deficiency of solid matters generally, has been called *Hydruria*, (ὕδωρ, 'water;') by a deficiency of urea, *Anazoturia*, (αζω, privative, and *azote*;) by a superabundance of urea, *Azoturia*. Urine in which the deposit consists of lithic acid, and the lithates, has been termed *Lithuria*; and that in which the deposit consists of the earthy and earthy alkaline phosphates, *Phosphuria*, *Ceramuria*, (κεραμα, 'potters' earth,') &c. &c. At times, a pellicle forms on the surface (*Cremor urinx*) which consists of mucus and some salts: at others, there is a cloud, (*Nube-cula*, *Nubes*), towards its upper part, which is called *Emore-ma* when it is suspended, as it were, about the middle of the liquid. Occasionally, there is a sediment, to which the names *Hypostasis*, *Sedimentum*, &c., have been given. (See these various words.) The urine is deserving of attention in disease; but it has often been considered a matter of more importance than it merits. Some empirics have presumed to pronounce on the nature of disease by simply inspecting it. See Uroscopy.

Human urine was, at one time, considered aperient; and was given in jaundice, in the dose of one or two ounces. Cows' urine, *Urina vac-cæ*,

All flower water, was once used, warm from the cow, as a purge, in the dose of Oss.

Urine has been variously designated by pathologists, according to its appearance, &c.;—for example:—

URINE, ANÆMIC, *Uri'na anæ'mica*. A term used, but not felicitously, for the form of urine that occurs in anæmia, chlorosis, &c.

URINE ARDENTE, Ardent urine—*u. Bloody*, Hæmaturia—*u. de la Boisson*, Urine, hysterical, see Urine—*u. Briqueté*, see Lateritious—*u. de la Coction*, see Urine.

URINE, CRUDE, (F.) *Urine crue*. Urine that is thin, transparent, and but little coloured, affording neither cloud nor deposit.

URINE CRUE, Urine, crude—*u. Cuite*, see Urine.

URINE, CYSTIN'IC, *Cystin'uria*. Urine containing cystine or cystic oxide in solution, or as a deposit.

URINE, DIABET'IC, *Uri'na diabé'tica*. Urine passed in *Diabé'tes mell'itus*, *Melitu'ria*, *Saccharine urine*. It contains a large quantity of sugar analogous to that of the grape, which may be separated in the form of crystals. In other respects, the urine is very abundant and very limpid in that affection.

URINE DE LA DIGESTION, see Urine.

URINE, DROP'SICAL, is an ammoniacal urine, often containing much albumen and scarcely any urea.

URINE, DYSPEP'TIC, has been found to precipitate tannin abundantly, and to putrefy soon.

URINE ÉPAISSE, Urine, mucilaginous.

URINE, FE'BRILE, *Uri'na febr'ilis*. Urine discharged in inflammation and inflammatory fever.

URINE, FILAMENTOUS, see Filamentous.

URINE, FLOCCULENT, (F.) *Urine floconneuse*, is when it is troubled by flocculi suspended in it.

URINE FLOCONNEUSE, Urine, flocculent—*u. des Goutteux*, Urine, gouty—*u. Furfuraceus*, see Furfuraceous.

URINE, GOUT'Y, (F.) *Urine des Goutteux*, contains much phosphate of lime, and after severe attacks of gout, precipitates a mixture of the uric and rosacic acids.

URINE HUILEUSE, Urine, oily.

URINE, HYSTER'ICAL, is clear, colourless, and similar to the *Urine de la Boisson*.

URINE, ICTER'ICAL, contains bile.

URINE, INCONTINENCE OF, Enuresis.

URINE OF INFLAMMATORY DISEASES. This is, generally, small in quantity, and very high-coloured.

URINE, INVOLUNTARY DISCHARGE OF, Enuresis.

URINE, JUMEN'TOUS, *Uri'na jumentosa seu jumenta'ria*, (F.) *Urine Jumentouse*. A term sometimes used for ammoniacal urine, rendered turbid by a substance similar to small grains of dust, which causes it to resemble the urine of herbivorous animals.

URINE LACTESCENTE, U. milky—*u. Laitieuse*, Urine, milky.

URINE, MIL'KY, (F.) *Urine lactescente ou laitieuse*. White and turbid urine.

URINE, MUCILAG'INOUS, (F.) *Urine épaisse*. Urine containing much mucus.

URINE, NER'VOUS. Thin and very liquid urine; such as is passed shortly after the attack of nervous diseases.

URINE OF NER'VOUS FE'VERS is generally very thin; and often deposits a mixture of uric and rosacic acids.

URINE, OIL'Y, (F.) *Urine huileuse*, is that which pours like oil, or which has an oily pellicle at its surface.

URINE, OXAL'IC, *Oxalu'ria*. Urine containing a salt of the oxalic acid.

URINE, PHOSPHORES'CENT, *Phosphoru'ria*, *Phos-*

phuria. Urine which is luminous or phosphorescent.

URINE, PURPURIC, see Porphyruria.

URINE, RICK'ETY, contains much phosphate of lime.

URINE, SACCHARINE, Urine, diabetic—*u. Sandy*, Arenosa urina—*u. Stoppage of*, Ischuria—*u. Ténue*, Urine, thin.

URINE, THIN, (F.) *Urine ténue*. Urine that is transparent, slightly coloured, and but little dense.

URINEUX, Urinous.

URINIF'EROUS, *Urinifer*, (F.) *Urinifère*, (from *urina*, and *fero*, 'I carry.') That which carries urine.

URINIFEROUS SUBSTANCE OF THE KIDNEY, see kidney.

URINIFEROUS TUBES OR DUCTS OF BELLINI, *Tu'buli seu ductus uriniferi Belli'ni*, vel *Bellini'ni*, *Tu'buli Belli'ni seu rec'ti*, *Bellini'ni Ductus*, *Bellini'na Fis'tule*. The collection of small tubes, converging from the cortical part of the kidney to the apices of the papillæ. The tortuous uriniferous ducts are called *Tubes of Ferrein*, *Tu'buli contorti* seu *Ferrein'i*.

URINIP'AROUS, (F.) *Urinipare*, (*νεμα*, and *parere*, 'to produce.') An epithet for tubes in the cortical portion of the kidney, which prepare the urine.

URINODES, (from *urina*, and *odes*.) Urinous.

URINOM'ETER, (from *urina*, and *μετρον*, 'measure.') A small hydrometer for taking the specific gravity of urine. The urinometer suggested by Dr. Prout is divided into 60 degrees, the zero being the point at which it floats in distilled water. The numbers on the scale added to 1000, the specific gravity of water, give the particular specific gravity: thus, supposing the number cut by the surface of the liquid to be 30, it would indicate a s. g. of 1.030. The letters H. S. on the back of the scale mean "healthy standard," which ranges from 10° to 20°.

U'RINOUS, *Urinous*, *Urinodes*, (F.) *Urineux*, (from *urina*.) Having the character of urine. Similar to urine.

URINOVAGINAL, Vesicovaginal.

UROCE'LE, *Edéma scro'ti urino'sum*, *Uroscheoce'le*, *Oscheoce'le urina'lis*, (*uron*, and *κηλη*, 'a rupture.') A tumour of the scrotum, owing to the effusion of urine into the same, and into the areolar tissue.

UROCHES'IA, *Urochez'ia*, (*uron*, and *χεζειν*, 'to go to stool.') A discharge of urine by the bowels.

UROCRIS'IA, (*uron*, and *κρινω*, 'I judge.') *Urocrite'riion*. A judgment formed either of the urine, or of disease after inspecting the urine. Also, the crisis of a disease by uresis. *Urocriterion* is the symptom by which such judgment is formed.

UROCRITERION, (*uron*, and *κριτηριον*, 'criterion.') see Urocrisia.

UROCYSTIS, (*uron*, and *κυστις*, 'a bladder,') Urinary bladder.

UROCYSTITIS, (*urocystie*, and *itis*.) Cystitis.

UROCYSTOCATARRHUS, Cystirrhœa.

URODIAL'YSIS, (*uron*, and *διαλυσις*, 'dissolution, loss of strength.') A partial and temporary suspension of the function of the kidney, such as occurs in the course of most acute diseases. A condition of the function of the kidney and of the urine, similar to that observed in acute diseases, but without local mischief or preceding inflammatory fever, has been termed, when occurring in children, *Urodial'ysis Neonatorum*; in the aged, *Urodial'ysis Senum*.

URODOCHIUM, (*ουροδοχιον*, *ουροδοχιον*, from *urine*, and *δοχωμαι*, 'to receive,') Urinal.

URODYN'IA, (*uron*, and *δυνειν*, 'pain') Pain during the excretion of the urine.

URŒDEMA, gen. *Urœde'matis*, (uron, and ὤδημα, 'a swelling,') Urapostema.

UROËMIE, Uremia.

UROËRYTHRIN, (uron, and ερυθρος, 'red,') see Porphyria.

UROGENITAL, *Urogenita'lis*, *Gen'itoû'ri-nary*, *Gen'itoû'rina'rius*. Relating or appertaining to the urinary and genital apparatus—as the 'urogenital organs,' *Org'ana urogenita'lia*.

UROGENITAL S'INUS, *Sinus urogenita'lis*. A cavity or canal in the embryo of the mammalia opening externally, in which the excretory ducts of the Wolffian bodies, the ureters and the efferent parts of the generative apparatus, terminate internally. It is prolonged into the urachus. Subsequently it becomes separated by a process of division into a *pars urina'ria* and a *pars genita'lis*. The former, extending towards the urachus, is converted into the urinary bladder; whilst from the latter are formed the *vesiculæ seminales* in the male, and the middle portion of the uterus in the female.

UROGLAUCIN, (uron, and γλαυκος, 'blue,') see Uroxanthin.

UROHÆMATIN, Hæmaphæin.

UROLITHI, (uron, and λιθος, 'a stone,') Calculi, urinary.

UROLITHIASIS, Lithia.

UROLITHOLOG'IA, (uron, λιθος, 'a stone,' and λογος, 'a description,') A description of, or treatise on, ordinary calculi.

UROLITHOTOMIA, Lithotomy.

UROLOGY, Uronology.

UROMACNY, *Uromanti'a*, (from uron, and μαντεία, 'divination,') The art of divining diseases by simple inspection of the urine; *Uroscop'ia*, *Uroscopy*, *Ouroscop'ia*, *Ou'roscopy*. One professing to be able to do this is called *Uroman'tes*. Vulgarly, a *water-doctor*, *water-caster*.

UROM'ELES, (F.) *Uromèle*, (ουρα, 'a tail,' and μέλος, 'a limb,') A genus of monsters in which the two lower limbs are united, very incomplete, and terminated by a single foot, which is almost always imperfect, and the sole of which is turned forwards.

UROMPH'ALUS, (uron, and ομφαλος, 'navel,') *Hēr'nia Ura'chi*. A monstrosity, which consists in the protrusion of the urachus at the umbilicus.

URON, (ουρον,) Urine.

URON'CUS, (uron, and οων, 'a swelling,') A urinary swelling.

URONOL'OGY, *Uronolog'ia*, *Ourolog'ia*, *Ou'rology*, *Urol'ogy*, *Urolog'ia*, (uron, and λογος, 'a discourse,') The part of medicine which treats of the urine.

UROPHTHISIS, (uron, and phthisis,) Diabetes mellitus.

UROPLA'NIA, (uron, and πλανη, 'wandering,') The presence of urine in other parts than the urinary organs.

UROPOE'SIS, *Uropoë'sis*, (uron, and ποιεω, 'I make,') *Secre'tio uri'næ seu lo'tii*. The secretion of urine by the kidney.

UROPOËT'IC, *Uropoët'icus*, (F.) *Uropoëtique*. Concerning or favouring the production of urine.

UROPOIESIS, Uropoesis.

UROPYGION, (ουρα, 'tail,' and πυγη, 'rump,') Coccyx.

UROPYGIUM, *Cronpion*.

URORRHAGIA, (uron, and ραγη, 'a breaking forth,') Diabetes.

URORRHODIN, (uron, and ρόδον, 'a rose,') see Porphyria.

URORRHŒA, (uron, and ρεω, 'to flow,') Eneuræsis, Diabetes.

UROSACIN, (uron, and rosa,) see Porphyria.

UROSCHEOCELE, (uron, σσχων, 'scrotum,' and κηλη, 'rupture,') Urocele.

UROSCOPY and UROSCOPIA, (uron, and σκοπω, 'I examine,') see Uromantia.

URO'SES, (from uron, and osis,) Diseases of the urinary organs.

UROST'EALITH, *Urosteal'ithus*, (uron, στεαρ, 'suet,' and λιθος, 'stone,') A peculiar fatty ingredient of urinary calculi, discovered by Heller.

UROSTENOSIS, Urethrostenosis.

UROUS ACID, Uric oxide.

UROXAN'THIN, *Uroxan'thine*, (uron, and ξανθος, 'yellow,') A yellow colouring matter found in the urine by Heller, which is decomposable into a red colouring matter, *ur'rhodine*, and a blue, *uroglau'cin*,—the former having the same composition as indigo red; the latter as indigo blue.

UROZEMIA (uron, and ζημια, 'loss,') Diabetes —u. Albuminosa, Kidney, Bright's disease of the —u. Mellita, Diabetes mellitus.

URRHODIN, (uron, and ρόδον, 'rose,') see Porphyria.

UROSACIN, (uron, and rosa, 'rose,') see Porphyria.

URTICA, (from urere, 'to burn,') *U. Dioica* seu *ma'jor seu his'pida*, *Cni'de*, *Py'r'ethrum*, *Ad'icè*, *Com'mon sting'ing net'tle*, (Prov.) *El'tle*, (F.) *Ortie*. Ord. Urticacæ. Ser. Syst. Monœcia Tetrandria. This plant has been long used for medical and culinary purposes. The young shoots, in the spring, are considered to possess diuretic and antiscorbutic properties. The pain produced by its sting is well known, but it is not to be compared to the torture occasioned by *Urtica stim'ulans*, and *U. crenula'ta* of India; and, still less, to *Urtica urentis'sima* of Timor, which is called by the natives *Daoun Setan*, 'Devil's leaf,' on account of the persistence and severity of the pain.

URTICA BALEARICA, *U. pilulifera* — u. *Crenulata*, see Urtica — u. *Dioica*, Urtica — u. *Hispida*, Urtica — u. *Iners* magna foetidissima, Galeopsis — u. *Major*, *U. dioica* — u. *Minor*, *U. urens* — u. *Mortua*, *Lamium album*.

URTICA PILULIF'ERA, *U. Pilulo'sa* seu *Roma'na* seu *Balea'rica*, *Pillbearing Net'tle*, (F.) *Ortie Romaine*. The seed was formerly given in diseases of the chest.

URTICA PILULOSA, *U. pilulifera* — u. *Romana*, *U. pilulifera* — u. *Stimulans*, see Urtica.

URTICA U'RENS, *U. mi'nor*, *Diwarf net'tle*, (F.) *Ortie brulante*. A lesser variety of the nettle, possessing similar virtues.

URTICA URENTISSIMA, see Urtica.

Nettles have been used as stimulants to paralytic limbs.

URTICAIRE, Urticaria.

URTICA'RIA, *Enanthe'sis urtica'ria*, *Urd'edo*, *Scarlati'na* seu *Fe'bris* seu *Pur'pura urtica'ra*, *Cnido'sis*, *Urtica'ria porcel'lana*, *Eranthe'ma urtica'tum*, *Epinyctis prurigin'o'sa*, *Fe'bris ru'bra prurigin'o'sa*, *Sall'ians ro'sa*, (F.) *Urticaire*, *Fidère Ortie*, *Porcelaine*, *Nettlerash*, (Prov.) *Nettlesprings*; vulgarly *Hives*, *Bald Hives*; (from urtica, because the eruption resembles that produced by the stings of a nettle.) A disease distinguished by those elevations of the cuticle called *wheals*. They have a white top, and are often surrounded by a diffuse redness. They excite intolerable itching, especially when the person is warm in bed. The eruption frequently goes and comes once or more in the 24 hours, and is generally dependent upon gastric derangement. In some people, nettlerash is induced by shellfish, eggs, nuts, &c. It is not usually a disease of any consequence, and its treatment is limited to the use of absorbent laxatives, and simple general means. Willan has described six varieties of the complaint:—*Urtica'ria febrilis*, *U. euan'tida*, *U.*

per'stans, *U. confer'ta*, *U. subcuta'nea*, and *U. tubero'sa*.

URTICARIA PORCELLANA, Essera, Urticaria.

URTICATIO, *Urticatio*, *Urtica'tion*, (from *urtica*.) A sort of flagellation, with nettles;—used with the intention of exciting the skin.

URUCU, Terra Orleansa.

URYLIC ACID, Uric acid.

USNEA CRANII HUMANI, Lichen saxatilis — *u. Florida hirta*, Lichen plicatus — *u. Hirta*, Lichen plicatus — *u. Plicata*, Lichen plicatus.

USSACU, Hura Brasiliensis.

USQUEBAUGH, Whisky, see Spirit.

USTILAGO, (same etymon as *ustio*.) Ergot.

USTILAGINOUS, *Ustilagin'ea*, (F.) *Ustilagineus*. Relating or belonging to ergot, as *Gangrena ustilaginea*; see Ergotism.

USTIO, (*ureo*, *ustum*, 'to burn,') Burn, Ustion.

USTION, *Us'tio*, *Cous'is*. The act of burning or of applying the actual cautery. Also, the effect of cauterization; a burn.

USTULATIO, *Ustula'tion*, (dim. of *ustio*.) A pharmaceutical operation by which a substance is dried by heat. It is often restricted to the roasting of ores to drive off the volatile matters, as in the case of arsenic.

USUALIA, (pl. of *usualis*, 'usual, ordinary,' from *usus*, 'use,') Officials.

USURE DES DENTS, ('wearing out of the teeth' from *use*.) Odontotripsis.

USUS, 'use,' Habit.

UTER, gen. *U'trie*, 'a leather bottle,' Uterus.

UTERAL'GIA, A hybrid word, (from *uterus*, and *algos*, 'pain,') Metralgia.

UTERI ADSCENSUS, Hysteria — *u. Anteversio*, Hysteroloxia anterior — *u. Carcinoma*, Metrocarcinoma — *u. Convulsivus morbus*, Lorind matricis — *u. Coryza*, Leucorrhœa — *u. Exulceratio*, Hysteroleosis — *u. Flexio*, Hysteroloxia — *u. Inclination*, Hysteroloxia — *u. Obliquitas*, Hysteroloxia — *u. Pronatio*, Hysteroloxia anterior — *u. Pruritus*, Nymphomania — *u. Reflexio completa*, Hysteroloxia posterior — *u. Retroversio*, Hysteroloxia posterior — *u. Rheuma*, Leucorrhœa — *u. Ulcus*, Hysteroleosis — *u. Versio incompleta*, Hysteroloxia.

UTERINE, *Uteri'nus*, (F.) *Uterin*, (from *uterus*.) That which belongs or relates to the womb.

UTERINE ARTERY, *Ar'teria uteri'na hypoga'strica*. Its size is always in a ratio with the greater or less state of development of the uterus. It arises from the hypogastric, either separately or with the umbilical, or from the internal pudic; passes upon the lateral parts of the vagina, to which it gives branches; and afterwards ascends on the sides of the uterus, and divides into a considerable number of very tortuous branches, which enter the tissue of the organ.

UTERINE BOUGIE, Sound, uterine.

UTERINE BROTHER, *Uteri'nus fra'ter*. One born of the same mother, but of a different father.

UTERINE ELEVATOR, see Pessary, intrauterine — *u. Glands*, Utricular glands — *u. Murmur*, *Bruit placentaire*.

UTERINE MUSCLE OF RUYSCHE consists of some fleshy fibres at the fundus of the uterus, which he regarded as forming a particular muscle.

UTERINE RUSH, *Bruit placentaire* — *u. Sound*, Sound, uterine.

UTERINUS FRATER, Uterine brother.

UTEROABDOMINAL SUPPORTER. See Belt, Russian.

UTEROGESTATION, (*uterus*, and *gestare*, 'to carry,') Pregnancy.

UTEROMANIA, Nymphomania.

UTEROOVARIAN ARTERY, see Spermatie artery.

UTERUS, (*borrea*), *U'ter*, *Utric'ulus*, *Vul'va*, *Lo'ci*, *L. mul'tebres*, *Ven'ter*, *Vis'cera*, *Concep'taculum*, *Mat'rix*, *Ar'eum* seu *A'ger natu'ra*, *Hys'tera*, *Me'tra*, *Eugi'um*, *Eugi'on*, *An'gus*, *Del'phys*, *Al'vus*, *Epompha'tion*, *Gus'ter*, *Ma'ter*, *Me'ter*, *Ne'dys*, the Womb, (Sc. and Prov.) *Wayme*, *Wame*, *Wemm*, *Wem*, (F.) *Matrice*, *Sein*. The uterus, destined to lodge the fœtus from the commencement of conception till birth, is a hollow symmetrical organ, having the shape of a truncated conoid: it is situate in the pelvis, between the bladder and the rectum, above the vagina, and below the convolutions of the small intestine. The uterus is flattened from before to behind, and is nearly an inch in thickness. It is two inches broad at its highest part, and becomes narrower towards the vagina; terminating in a contracted portion, called the *cer'vix* or *neck*, to distinguish it from the rest of the organ, called the *body*. From its *fundus* or upper portion, the Fallopian tubes pass off. The mouth is called *Os uteri*, *Os tin'ce*, *Os u'teri extern'um*, *Orific'i'um vagi'na'le*. At the point where the body of the uterus is continuous, below, with the neck, the cavity is slightly constricted, forming what is sometimes named the *inter'nal orifice*, *Os u'teri intern'um*, *Os intern'um*, *Isth'mus* seu *os'tium u'teri*, *Orific'i'um uteri'num*. Between these is the *Cava'lis cer'vi'cis*.

The cavity of the uterus is very small, and its parietes are thick. The portion of the cavity corresponding with the body is triangular and flattened. Its superior angles present the extremely fine orifices of the Fallopian tubes. The cavity of the neck is slightly dilated before opening into the vagina. The uterus is composed, 1. Of a *serous membrane*. This is external, and furnished by the peritoneum; which is reflected upon the uterus from the posterior surface of the bladder and the anterior surface of the rectum. These folds of the peritoneum are sometimes called, respectively, the *anterior*, and the *posterior*, ligaments. The former are also named the *vesico-uterine*, and the latter the *recto-uterine* folds or *pli'ce semiluna'res*, (F.) *Plis de Douglas*, — the *pouch*, thus formed, being termed *Douglas's cul-de-sac*. It forms, also, the broad ligaments of the uterus, *Ligamen'ta la'ta*, *A'ix espermatio'nis*, *Re'tia U'teri*. These contain, above, the Fallopian tube; beneath and anteriorly, the round ligaments; and behind, the ovary. They are two membranous productions or duplicatures of the peritoneum, sent from the edges of the uterus and posterior extremity of the vagina, in a transverse direction, to be fixed to, and line, the side of the pelvis. 2. Of a *mucous membrane*, which is a prolongation of that of the vagina, and extends into the tubes. 3. Of a *proper tissue*, whose thickness is considerable. It is of a close texture; is grayish and elastic; and resists the instrument that divides it. Its intimate nature is but little known. During gestation it is truly muscular. The arteries of the uterus proceed from the spermatic and hypogastric. Its veins follow the same course, and bear the same name as the arteries. They are very tortuous, and form large sinuses in the parietes of the organ, called *U'terine Si'nuses*. Its nerves proceed from the sciatic and hypogastric plexuses. Its lymphatic vessels are very multiplied; and during pregnancy acquire enormous dimensions. The changes experienced by the uterus during gestation, and after delivery, are very numerous and important to be known. They regard its situation, direction, size, shape, texture, connexions, vital properties, &c.

UTERUS, Abdomen — *u. Anteversion* of the Hysteroloxia anterior.

UTERUS BIFIDUS, 'a bifid uterus.' A double uterus. One separated into two cavities.

UTERUS DUPLEX, 'a double uterus,' *Dihysteria*—u. Hourglass contraction of the, *Chaton*—u. *Inclinatus*, *Hysteroloxia*.

UTERUS, INVERSION OF THE, *Inversio Uteri*, *Metranstrophè*, *Adopto'sis Uteri inversa*, (F.) *Inversion ou Rversement de la Matrice*. The uterus, displaced and turned inside out. This can only happen through want of caution on the part of the medical practitioner. It is a fatal accident, unless speedily reduced. This must be done by gradually returning the superior part, by grasping it with the hand, and making the rest follow.

UTERUS, IR'RITABLE, *Neuralgia of the Uterus*. A disease characterized by deepseated pain in the lower part of the abdomen, and in the back and loins; generally diminished by lying down, and increased by exercise, and more severe for a few days preceding and during menstruation. On examination *per vaginam*, the uterus is found to be tender on pressure, *Metriperæsthesis*; and the cervix and body slightly swollen, but not usually indurated. It is an obstinate affection, but frequently ends of itself after years of suffering. The treatment must be,—the horizontal posture; local blood-letting to the uterus; emollients; soothing injections; narcotics, warm bath, &c. See *Hysteralgia*.

UTERUS MASCULINUS, see Ejaculatory ducts—u. *Obliquus*, *Hysteroloxia*—u. *Retroversion of the*, *Hysteroloxia posterior*.

UTERUS, RUPTURE OF THE, *Metrorrhæsis*, *Hæmorrhæsis*, *Ruptura seu Perforatio Uteri*, *Laceration of the Womb*, (F.) *Rupture ou Déchirure de la Matrice*. This dreadful accident occurs during labour, and is known to have taken place, when the child's head suddenly recedes during a pain, with vomiting, sinking of the pulse, cold sweats, &c. When it is evident that this has occurred, the hand must be passed into the uterus, and the feet of the child be brought down.

Cases are on record, in which partial lacerations of the uterus have healed, and the patient has done well. This is a rare event, and not to be anticipated.

UTRICLE, *Utriculus*, (dim. of *uter*, gen. *utris*), 'a small bag or bottle,' see Ejaculatory ducts, Semicircular canals, Uterus.

UTRICLE, PRIMORDIAL, *Utriculus primordia'lis*. A delicate membrane, of protein composition, which lines the cell wall of vegetable structures, and from the exterior of which the layer of cellulose is secreted.

UTRICULAR, *Utricula'ris*. Relating to or resembling a small bag or bottle.

UTRICULAR GLANDS, *Glandulæ utricula'res seu uteri'nae*. The glands or follicles in the interior of the uterus, which are concerned in the formation of the decidua.

UTRICULUS, *Utricle*, see Ejaculatory ducts, Semicircular canals, Uterus—u. *Communis*, see Semicircular canals—u. *Lacteus*, *Receptaculum chyl'i*—u. *Primordialis*, *Utricle*, *primordial*—u. *Prostaticus*, see Ejaculatory ducts.

UVA, 'a grape, or a bunch of grapes,' *Uvula*, see *Vitis vinifera*—u. *Inversa*, *Paris*—u. *Lupina*, *Paris*—u. *Passa*, see *Vitis vinifera*—u. *Ursi*, see *Arbutus uva ursi*—u. *Vulpina*, *Paris*.

UVÆ CORINTHIACÆ, see *Vitis Corinthiaca*.

UVÆ PAS'SÆ, 'Dried Grapes,' *Rhag'es*, *Raisins*, see *Vitis vinifera*.

UVÆ PASSÆ MAJORES, see *Vitis vinifera*—u. *Passæ minores*, see *Vitis Corinthiaca*—u. *Passæ sole siccata*, see *Vitis vinifera*.

UVARIA TRILOBA, (from *uva*, 'a grape,') see *Carica papaya*.

UVATIO, (from *uva*, 'a grape,') *Ceratocela*, *Staphyloma*.

UVEA, (from *uva*, 'a grape,') *Tu'nica acinosa seu aciniformis seu uva'lis seu acina'tis seu uviformis seu rhagoi'des seu u'væ seu perforata*, *Membra'na u'væ*, *Rhagoi'des*, *Rhor*, (F.) *Uvée*. Some anatomists have given this name to the choroid coat of the eye; others, to the posterior layer of the iris, on account of the black and very thick varnish that covers it. See *Iris*.

UVEA, COMMISSURE OF THE, *Ciliary ligament*.

UVÉE, *Uvea*.

UVEITIS, (F.) *Uvéite*, (*uvea*, and *itis*), properly, *Rhagoidei'tis*. Inflammation of the uvea. See *Iritis*.

UVIGENA, *Uvula*.

UVIGERA, *Uvula*.

UVULA, (dim. of *uva*, 'a grape,') *Pinnaculum for'nicis guttura'lis*, *U'væ*, *Staphylæ*, *Tintinnabulum*, *Columell'a*, *C'ion*, *C'ionis*, *Garga'reon*, *Colum'na o'ris*, *Gurgulio*, *Interseptum*, *Processus seu Cartila'go Uvifer*, *Uvig'ena*, *Uvig'era*, *H'ymas*, *Plect'rum*, *Subling'ua*, *Sublinguim*, *Pap of the Throat*, (Sc.) *Pap of the Hass*, (F.) *Luette*. A fleshy appendix or prolongation, which hangs from the middle or free edge of the velum palati. The uvula has a conical shape, and is of a greater or less size in individuals. It is formed, especially, by the mucous membrane of the velum palati; and contains within it the azygos uvulae muscle. Its use is not clear.

Also, a small prominence or lobule in the portion of the cerebellum that forms the posterior boundary of the fourth ventricle, *U'vula cerebelli*. It is on the median line, and behind the nodulus.

UVULA, see Vermiform process inferior—u. *Cerebelli*, see *Uvula*—u. *Decidua*, *Staphylœdema*—u. *Relaxata*, *Staphylœdema*—u. *Vesicæ*, see *Urinary bladder*—u. *Vesical*, see *Urinary bladder*.

UVULAR, *Uvula'ris*, (F.) *Uvulaire*, (from *uvula*.) That which belongs to the uvula.

UVULAR GLANDS are small follicles, belonging to the mucous membrane covering the uvula.

UVULARIA, (so called from the flowers hanging like the uvula, [?]) *Ruscus hypoglossum*.

UVULARIA PERFOLIA'TA, *Smaller Bellwort*; indigenous; *Order*, *Melanthaceæ*; flowering in May and June. The root, when fresh, is subacid and mucilaginous. A decoction of the plant is used in sore mouth. It is said to cure the bites of rattlesnakes.

UVULARIA GRANDIFLO'RA and *U. Sessilifolia* have similar properties.

UVULITIS, (*uvula*, and *itis*), *Angi'na uvula'ris*, *Staphyli'tis*, *Cioni'tis*, *Inflamma'tio u'vulae*, (F.) *Inflammation de la Luette*, *Falling down of the Palate*. Inflammation of the uvula. When to a slight degree, *Hypostaphyli'tis*.

V.

VACCIN, see Vaccina.

VACCINA, *V. primigenia* seu *vaccinatoria* seu *antivariolosa*, *Vaccinia*, *Variola vaccina*, *V. vaccinia* seu *tutoria* seu *tutrix*, *Eranthema vaccina* seu *antivariolsum*, *Synocha vaccina*, *Eriophysis vaccina* *inserita*, *Vacciola*, *V. scutellata*, *V. vera*, *Variola tutoria*, *Cowpox*, *Cowpock*, *Kine Pock*, *Inoculated Cowpox*, (from *vacca*, 'a cow,') (F.) *Vaccine*. The cowpox is a disease of the cow, arising spontaneously, which, if transmitted to man by means of inoculation, may preserve him from smallpox contagion. The promulgation of this valuable property of the vaccine virus is due to Dr. Jenner. When inoculating for smallpox in Gloucestershire in 1775, he was surprised to meet with a number of persons in the same district, on whom the insertion of the virus produced no effect, although they had never passed through smallpox. On investigation, he found that all these persons had previously been affected by a pustular eruption of the cow; which they had taken by milking the animal; and which eruption they called *Cowpox*. This discovery induced him to institute experiments, and he found that such preservative effect actually existed. The *Vaccine Virus*, *Virus Vaccinum*, (F.) *Vaccin*, the fluid of the eruption, is not now usually taken immediately from the cow; experience having demonstrated that it is perhaps as efficacious when obtained from the human vesicle. The characteristic of cowpox eruption is:—a semitransparent, pearlecoloured vesicle, with a circular or somewhat oval base; its upper surface, until the end of the 8th day, being more elevated at the margin than in the centre; and the margin itself being turgid, shining, and rounded, so as often to extend a little over the line of the base. This vesicle is filled with clear lymph, contained in numerous little cells, that communicate with each other. After the 8th or 9th day from the insertion of the virus, it is surrounded by a bright red, circumscribed areola, which varies in its diameter, in different cases, from a quarter of an inch to two inches; and is usually attended with a considerable tumour and hardness of the adjoining areolar membrane. This areola declines on the 11th or 12th day; the surface of the vesicle then becomes brown in the centre; and the fluid in the cells gradually concretes into a hard, rounded scab or crust, of a reddish-brown colour, which at length becomes darker, contracted, and dry, but is not detached till three weeks after vaccination. It leaves a permanent, circular cicatrix, about five lines in diameter, and a little depressed; the surface being marked with very minute pits or indentations, denoting the number of cells of which the vesicle had been composed. This is the progress of the *genuine cowpox*. At times, fever occurs about the 9th or 11th day; but, at others, not the slightest indisposition is felt. Aberrations from this progress occur, which must be attended to; as the same preservative influence is not exerted by *irregular* or *spurious cowpox*. If the eruption runs speedily to a height, has an areola too early, is irregular at its edges, and contains pus instead of a limpid fluid, it cannot at all be depended upon.

Smallpox occurs, occasionally, as an epidemic after vaccination. For a while, the enthusiastic advocates of vaccination would not admit the affection to be smallpox; and, accordingly, they gave it the name Varioloid. It is unquestionably smallpox, but modified, and greatly divested of its terrors by previous vaccination; and it is

said to happen less frequently after vaccination than after smallpox inoculation. Vaccination is now practised everywhere, except among those in whom ignorance and prejudice exclude the lights of reason and philanthropy. There is ground, however, for the belief, that for full protection revaccination may be occasionally necessary.

VACCINA ANTIVARIOLOSA, *Vaccina*—v. *Primigena*, *Vaccina*—v. *Tutoria*, *Vaccina*—v. *Tutrix*, *Vaccina*—v. *Vaccinatoria*, *Vaccina*.

VACCINAL, *Vaccine*.

VACCINATION, *Vaccinatio*, (from *vacca*, 'a cow;') *Cowpox inoculation*, *Jennerian inoculation*. An operation which consists in inserting the vaccine virus under the cuticle, so that it may enter into the absorbents. It is extremely simple, and requires merely that the point of the lancet should be held under the cuticle for a short time. The best time for taking the matter from the *Cowpox vesicle*, for vaccination, is about the 7th or 8th day, before the appearance of the areola, when it ought to be limpid and transparent. The most certain method is, to pass it from arm to arm; but this is not always convenient; and, hence, it is taken upon points or pieces of quill, or between glasses. It must, of course, be softened for use. This is done by means of a little water, which does not destroy its efficiency. The scab, too, which falls off about the end of three weeks—sooner or later—is capable of communicating the affection, if dissolved in water; and this is the most common method of transmitting the vaccine virus in some countries.

VACCINATOR, *Vaccinist*. One who inoculates for the cowpox.

VACCINE, *Vaccinus*, *Vaccinallis*, (F.) *Vaccinal*; (same etymon as Vaccination.) Relating or appertaining to vaccination, as the '*vaccine vesicle*,' 'a vaccine physician.'

VACCINEA, *Vaccina*.

VACCINELLA, (dim. of *vaccina*.) A term employed by some to designate spurious cowpox. *Vaccinola*, *Vaccinola spuria* seu *leprosa*, (F.) *Vaccinelle*, *Vaccinoides*.

VACCINIA, *Vaccina*—v. *Palustris*, *Vaccinium oxycoccus*.

VACCINIST, *Vaccinator*.

VACCINIUM MACROCARPON, see *V. oxycoccus*.

VACCINIUM MYRTILLUS, (either corrupted from *hyacinthus*, or, and more probably, from *baccinum*; from *bacca*, 'berries,' in allusion to its fruit.) *Vitis idæa myrtillus*, *Vaccinia myrtillus*, *Vaccinium nigrum*, *Myrtleberry*, *Bilberry*, *Blueberry*, *Whortleberry*, *Black Whorts*, (Sc.) *Blacberry*, (Prov.) *Harleberry*, *Whimberry*, (F.) *Airelle Anglaise*, *Raisin de Bois*. Ord. *Vaccinææ*. Sex. Syst. *Ocandria Monogynia*. The berries—*Baccæ Myrtillorum*—are esteemed to be antiscorbutic, when prepared with vinegar. They are also said to possess antiseptic and subastrigent properties when dried. They are not used in medicine; but are frequently eaten with milk and sugar.

VACCINIUM NIGRUM, *V. myrtillus*.

VACCINIUM OXYCOCOS, *Oxycoccus*, *O. palustris*, *Schollera oxycoccus*, *Vaccinia palustris*, *Vitis Idæa palustris*, *Moortberry*, *Cranberry*, *Mossberry*, *Swamp Redberry*, *Sourberry*, (F.) *Canneberge*. The berries are pleasantly acid and cooling. They make an agreeable preserve. The common American Cranberry is from *Vaccinium macrocarpon*, *Oxycoccus macrocarpus*.

VACCINIUM PENNSYLVANICUM, *Low Blueberry*. The leaves and bark of the root are astringent.

VACCINIUM PUNCTATUM, *V. vitis idæa*—*v. Resinosum*, *Gaylussacia resinosa*.

VACCINIUM STAMINÆUM, *Buckberry*, *Squaw Huckleberry*, *Deerberry*. The leaves of these plants are astringent.

VACCINIUM VITIS IDÆA, *V. punctatum*, *Vitis Idæa punctata*, *Am'pelos Idæa*, *Red Bilberry*, (Sc.) *Braw'lins*, *Al'pine Cran'berry*, *Whorl'leberry*, *Cue'berry*, (F.) *Airelle* ou *Canneberge ponctuée*. A decoction of the leaves has been used in calculous affections. They are astringent. The ripe berries are pleasantly acid, and refrigerant. *Whortleberries* are furnished by different species of *Vaccinium*.

VACCINOÏDE, (*vaccina*, and εἶδος, 'resemblance,') *Vaccinella*.

VACCINOIS, *Vaccinella*.

VACCINUS, *Vaccine*.

VACCIOLOA, *Vaccina*—*v. Leprosa*, *Vaccinella*—*v. Scutellata*, *Vaccina*—*v. Spuria*, *Vaccinella*—*v. Vera*, *Vaccina*.

VACILLATIO, *Titubatio*, (from *vacillare*, *vacillatum*, 'to waver,') *Staggering*; *waddling*—*claudicatio antica*.

VACILLATIO DENTUM, *Odontoseisis*.

VACUOLE, (dim. of (L.) *vacuus*, 'empty,') *Areola*.

VAGIN, *Vagina*.

VAGINA, 'a sheath,' *V. muliebri* seu *uteri*—*na* seu *Uteri* seu *Pe'nis*, *Sinus muliebri* seu *Pudoris* seu *Puden'ti*, *Col'leus*, *Pe'ris*, *Pe'rin*, *Colpus*, *Vulvot'lerine canal*, (F.) *Vagin*, *Conduit* ou *Canal de la pudeur*. A cylindrical canal, five or six inches long; situate within the pelvis, between the bladder and rectum. It communicates by one extremity, *Aul'us* seu *Intro'itus*, with the vulva; by the other, *For'nix* seu *Fim'dus*, with the womb, the *Pars* seu *Por'tio vagina'lis u'teri*, the neck of which it embraces. The vagina is lined, internally, by a mucous membrane; and is partly closed, in young females, by the hymen, and carunculae myrtiformes. Around the mucous membrane is a layer of spongy, erectile tissue; and a cellulovascular membrane. A constrictor muscle; numerous vessels furnished by the vaginal artery, particularly; veins and nerves, proceeding from the sciatic plexus, also enter into the composition of the organ.

The term *Vagina* or *Sheath*, *The'ca*, (F.) *Gaine*, is extended to many parts, which serve as envelopes to others. Thus, it is sometimes applied to the fascia enveloping the limbs; to areolar membrane, enveloping the muscles, and entering between their fibres; to that which surrounds the bloodvessels; to the synovial membranes surrounding tendons, and lining the gutters of bones, &c.

VAGINA CORDIS, *Pericardium*—*v. Femoris*, *Fascia lata aponeurosis*—*v. Funiculi umbilicalis*, see *Funiculus umbilicalis*—*v. Glissonii*, *Capsule of Glisson*—*v. Malleolaris externa*, *Retinaeculum tendinum perinæorum*—*v. Muliebris*, *Vagina*—*v. Mucularis*, *Perimysium*—*v. Nervorum*, *Neurilemma*—*v. Penis*, *Vagina*—*v. Pili*, see *Hair*—*v. Portæ*, *Capsule of Glisson*—*v. Uterina*, *Vagina*.

VAGINA OF THE VE'NA POR'TA, *Sheath of the Vena Porta*, (F.) *Gaine de la Veine-porte*, is the areolar membrane that accompanies the divisions of the vena porta into the substance of the liver.

VAGINÆ SYNOVIALES, 'synovial sheaths,' *Bursæ mucosæ*.

VAG'INAL, or **VAG'NAL**, *Vagina'lis*, (from *vagina*, 'a sheath,') *Relating* or *pertaining* to a sheath.

VAGINAL ARTERY. This exists, as a matter of course, only in the female. It arises from the hypogastric, or from the uterine, vesical, internal pudic, umbilical, &c., and sends its branches,

first to the lateral parts of the vagina, and afterwards to the anterior and posterior surfaces.

VAGINAL BILIARY PLEX'US. A plexus of ramifications of the hepatic duct through the capsule of Glisson.

VAGINAL BRANCHES OF THE VE'NA POR'TA are those which have to pass through the sheath of the capsule of Glisson, previous to entering the interlobular spaces. In this course, they form the *vaginal plex'us*. There are, also, *vaginal branches*, and a *vaginal plexus* formed by the hepatic artery.

VAGINAL COAT OF THE EYE, see *Eye*.

VAGINAL COAT OF THE TESTICLE, *Tu'nica Vagina'lis* seu *Elytro'ides* seu *Elytro'des*, is the serous membrane which envelops the testicle. Before the descent of the testicle, the tunica vaginalis does not exist. It is formed by a prolongation, furnished by the peritoneum to the testicle when it issues from the abdomen. The tunica vaginalis forms a close sac, investing the testicle and epididymis, without containing them in its cavity, and reflected so as to form a bag around the circumference of the testis; hence it is divided into *Tu'nica vagina'lis pro'pria* and *Tu'nica vagina'lis reflex'a*. Between the two layers, the inner surface is smooth, and lubricated by a serous fluid.

Inflammation of the tunica vaginalis has been called *Vaginali'tis*, (F.) *Vaginalite*.

VAGINAL GANGLIA, see *Spermatic ganglion*—*v. Plexus*, see *Vaginal branches of the vena porta*.

VAGINAL PROC'ESS OF THE TEMPORAL BONE, (F.) *Apophyse vaginale* ou *engainante*, *Gaine de l'apophyse styloïde*, *Chaton*, is a bony crista, which embraces the base of the styloid process.

VAGINAL PULSE, *Pulse*, *vaginal*.

VAGINALITIS, see *Vaginal coat of the testicle*.

VAGINISM'US. A name given by Dr. J. Marion Sims to spasm of the sphincter muscle of the vagina, with hyperæsthesia of the surrounding parts, and altered structure, which requires division by the knife.

VAGINITIS, (*vagina*, and *itis*.) Inflammation of the vagina, see *Leucorrhœa* (vaginal.)

VAGINODYNIA, (*vagina*, and *odyni*, 'pain,') *Colpalgia*.

VAGINOVESICAL, *Vesicovaginal*.

VAGISSEMENT, (from *vagitus*,) *Cry*, *Vagitus*.

VAGITUS, (*vagire*, *vagitum*, 'to cry,') (F.) *Vagissement*. The cry of the newborn child. *Squall'ing*.

VAGITUS UTERI'NUS. The cry of the child in utero.

VAGUS (NERVUS), 'wandering (nerve,)' *Pneumogastric*.

VAIRON (F.), (from *varius*, 'of different colours,') *Dispar ocul'us*. One who has eyes of different colors. It also means a *wall-eyed animal*, or one whose iris is surrounded by a whitish circle.

VAISSEAU, (from *vas*, dim. *vasculum*,) *Vessel*—*v. Sanguin*, *Bloodvessel*.

VAISSEAU'X (pl. of *vaisseau*) **CAPILLAIRES**, *capillary vessels*—*v. Chylifères*, *Chyliferous vessels*—*v. Courts*, *Brevia vasa*—*v. Droits*, see *Rete testis*—*v. Efférents*, see *Efferent*—*v. Lactés*, *Chyliferous vessels*—*v. Lactifères*, *Lactiferous vessels*—*v. Omphalicaux*, *Umbilical vessels*—*v. Tourbillonnés*, *Vasa vorticosa*—*v. des Vaisseaux*, *Vasa vasorum*.

VALANIDA, *Fagus sylvatica*.

VALANTIA APARINE, (after S. Vaillant, a French botanist,) *Galium aparine*.

VALENTINIAN VESICLES, (after Valentin, the German physiologist,) see *Molecule*.

VALERIAN, AMERICAN, (from *valere*, 'to have efficacy' [?]), *Cypripedium luteum*, *Valeriana pauciflora*—*v. Garden*, *Valeriana phu*—*v. Greek*, *Polemonium reptans*—*v. Official*, *Valeriana*—*v. Wild*, *Senecio aureus*, *Valeriana*, *V. pauciflora*.

VALERIA'NA, *V. officinalis* seu *mi'nor*, *Leucolach'anum*, *Wild* or *Offic'inal Valerian*. *Ord. Valerianaceæ*. *Sex. Syst.* Triandria Monogynia. The root, *Valeria'na* (Ph. U. S.), has been long extolled as an efficacious remedy in epilepsy, and in a variety of nervous complaints. It has been also advised as a tonic, antispasmodic, and emmenagogue. It has a strong, fetid odour; bitterish, subacid, and warm taste; and its virtues are extracted by water and alcohol. The distilled oil is official in the Pharmacopœia of the United States.

The *Kooyah* root or *Kooyahs* of the Indians in Oregon, used by them for making a bread called *Supate*, is said to be that of *Valeriana officinalis* or *Valeriana edulis*, probably the same as that sometimes written *Kous*. It is frequently called *Tobacco* root. It has to be buried in the ground for two days to deprive it of its noxious properties. The bread has an offensive taste to those not accustomed to it.

VALERIANA *CAPEN'SIS*, a South African species, has similar virtues.

VALERIANA *CELTICA*, *V. Saxatilis*, *Spica Celtica* seu *Celtica Dioscoridis* seu *Alpi'na*, *Nardus Celtica*, *Salum'ca*, *Nardum Gallicum*, *Celtic Nard*, (F.) *Nard Celtique*. The roots of this Alpine plant have been recommended as a stomachic, carminative, and diuretic. It has a moderately strong smell, and a warm, bitterish, subacid taste.

VALERIANA *DENTA'TA*, *V. locus'ta*, *Corn Sal'ad*. A wholesome, succulent plant, cultivated in Europe as a salad. It is antiscorbutic and gently aperient.

VALERIANA *EDULIS*, see *Valeriana*—v. *Locusta*, *V. dentata*—v. *Major*, *V. phu*—v. *Minor*, *Valeriana*—v. *Officinalis*, *Valeriana*.

VALERIANA *PAUCIFLO'RA*, *American* or *Wild Valerian*, and *V. sylvatica*, have the properties of the other *Valerians*.

VALERIANA *PHU*, *V. maj'or*, *Gar'den Valerian*. (F.) *Grande Valériane*. The root has been recommended in rheumatism; especially in sciatica, and in epilepsy.

VALERIANA *SAXATILIS*, *V. celtica*—v. *Spica*, *Nardus Indica*—v. *Sylvatica*, see *V. pauciflora*.

VALÉRIANE, *GRANDE*, *Valeriana phu*.

VALET À PATIN (F.), *Volcel'la Pati'ni*. The common ligature forceps is a modification of this instrument invented by one Patin. It consists of two branches, united at the middle by a hinge, which may be separated or approximated by means of a screw or running ring. It is used for seizing hold of vessels that are to be tied.

VALETUDINAIRE, *Valetudinary*.

VALETUDINARIUM, (from *valetudo*,) Hospital.

VALETUDINARIUS, Sickly, *Valetudinary*.

VALETU'DINARY, *Valetudina'rius*, *Valetudina'rius*, *Inval'idus*, (F.) *Valetudinaire*. One of delicate health. One subject to frequent diseases. An *Inval'id*.

VALETUDO, gen. *Valetu'dinis*, 'health, whether good or bad,' *Sanitas*—v. *Adversa*, Disease—v. *Bona*, *Sanitas*—v. *Commoda*, *Sanitas*—v. *Secunda*, *Sanitas*.

VALGUS, 'bowlegged,' see *Cnemosciosis* and *Kyllosis*.

VALIGA, *Tinctura jalapii*.

VALITUDO, gen. *Valit'udinis*, *Sanitas*.

VALLECULA, (dim. of *vallis*,) Valley—v. *Unigu*, see *Nail*.

VAL'LEX, *Vallis*, dim. *Vallec'ula*, *V. Reil'ii*, (F.) *Grand Scissure Médiane du Cervelet*. The deep fissure on the inferior part of the cerebellum, which divides it into two symmetrical portions.

VALLIS, Valley—v. *Alarum*, 'valley of the armpits,' *Axilla*—v. *Femorum*, *Valva*.

VALLUM, 'a pallsaded rampart.' The eyebrow or supercilium. Also, a species of bandage.

VALS, MIN'ERAL WATERS OF. A French acidulous chalybeate at Vals, six leagues from Viviers, in the department of Ardèche. There are five springs, containing carbonic acid, sulphate of iron and alum. They are employed in passive hemorrhages, leucorrhœa, &c. Season, from the 1st of June to the 15th of September.

VALVA, Valve.

VALVE, *Valva*, dim. *Valvula*, *Di'clis*, (F.) *Valvule*, (*valvè*, 'folding-doors,' itself from *volvo*, 'to fold up.')

A small door. Any membrane or duplicature of a membrane, which prevents a reflux of humours or other matters in the vessels and canals of the animal body. There are some valves, whose functions appear to be,—to retard or to modify the course of substances along canals, &c. See *Mitral*, *sigmoid*, *tricuspid*, *ileocecal valve*, &c.

VALVE OF FALLOPIUS, *Bauhin*, valve of—v. *Ileocecal*, *Bauhin*, valve of—v. *Ileocolic*, *Bauhin*, valve of—v. of *Tulpius*, *Bauhin*, valve of—v. of *Varolius*, *Bauhin*, valve of.

VALVES, KERKRINGIAN, *Valvula conniventes*—v. *Semilunar*, *Sigmoid valves*.

VALVULA, pl. *valvulae*, (dim. of *valva*,) Valve—v. *Cæci*, *Bauhin*, valve of—v. *Cerebelli*, *Valvula Viesssenii*—v. *Coli*, *Bauhin*, valve of—v. *Foraminis ovalis*, *Ovalis fossa*—v. *Guiffartiana*, see *Thebesius*, veins of—v. *Ilei*, *Bauhin*, valve of—v. *Magna cerebri*, *V. Viesssenii*.

VALVULA *SEMILUNA'RI*, *Plu'ca Semiluna'ris* seu *luna'ta*, *Membran'ula semiluna'ris conjunctivæ*. A small doubling of the tunica conjunctiva, which lies between the caruncula lacrymalis and the ball of the eye. In birds, this is large, and called *Membrana nictitans* seu *Palpebra ter'tia*.

VALVULA *TARINI*, *Velum medullare posterius*—v. *Thebesii*, see *Thebesius*, veins of—v. *Tulpii*, *Bauhin*, valve of—v. *Vaginæ*, *Hymen*—v. *Venæ magnæ*, see *Thebesius*, valve of.

VALVULA *VIÉSSENII*, *V. cerebelli* seu *Willisiana* seu *magna cerebri*, *Velum Viesssenii* seu *interjectum cerebelli*, *V. apoph'ysi vermiform'i obtentum*, *V. medullarè seu medullarè anti'um seu ante'rius*, *Lacu'nar ventriculi quatuor'ti en'terior*, (F.) *Valvule de Viesssens*, *Voile médullaire supérieure*. A thin, medullary lamina, over the under end of the Aquæductus Sylvii and upper part of the fourth ventricle. At the sides of the valves are two medullary tracts, called *Processus ad Testes*, *Colum'næ Valvulæ Viesssenii* or *Oblique intercerebral Commissure*.

VALVULA *WILLISIANA*, *Valvula Viesssenii*.

VALVULÆ, pl. of *Valvula*—v. *Bicuspidalæ*, *Mitral valves*.

VALVULÆ *CONNIVENT'ES*, 'self-closing valvulæ,' *V. C. Kerkrin'gii*, *Plu'cæ conniven'tes*, *Kerkrin'gian valves*. Numerous circular folds in the intestinal canal from the pyloric orifice through the greater part of the small intestine. They are folds of the mucous membrane; and their chief use appears to be, to retard the course of the alimentary mass, and to afford a larger surface for the origin of chyliferous vessels.

VALVULÆ *CORDIS*, *Ostiola cordis*—v. *Cordis mitrals*, *Mitral valves*—v. *Semilunares*, *Sigmoid valves*—v. *Triglochinæ*, *Tricuspid valves*—v. *Trisulcæ*, *Tricuspid valves*.

VALVULE, Valve—v. *d'Eustache*, *Eustachian valve*—v. *de Pyllore*, see *Pylorus*—v. *de Viesssens*, *Valvula Viesssenii*.

VALVULES *ÉPISCOPALES*, *Mitral valves*—v. *Sigmoides* ou *Semilunaires*, *Sigmoid valves*.

VANDELLIA *DIFFUSA*, (after *Vandelli*, professor of botany at Coimbra.) A native of Brazil, Guiana and Isle of France; *Order*, *Scrophulariaceæ*; is used there as an antibilious emetic and

antiperiodic; and is regarded as an efficacious remedy in malignant fever and dysentery, especially in cases connected with a disordered state of the liver.

VANELLOE, Vanilla.

VANILLA, *Bani'l'ia*, *Bani'l'as*, *Bani'l'oes*, *Bani'lla*, *Benzo'enil*, *Ar'acius aromati'cus*, (F.) *Vanille*, *Baume de Vanille*, ([S.] *vainilla*, dim. of *vaina*, 'a sheath, a pod.') The *Vanel'loe*—*Sil'igna vanill'e* seu *Bani'll'e* seu *Vanigli'e* seu *vanill'ie* seu *Ar'aci aromati'ci*—is a long, flattish pod; containing, under a wrinkled, brittle shell, a reddish-brown pulp, with small, shining, black seeds. The parasitic plant which affords this pod is the *Epidendrum Vanill'a*, *Vanill'a aromati'ca* seu *Mexica'na*, and other species. *Family*, Orchidaceæ. *Sex. Syst.* Gynandria Monandria. Vanelloes have an unctuous, aromatic taste, and a fragrant smell, like that of some of the finer balsams, heightened with musk. Although chiefly used as perfumes, they are reputed to possess aphrodisiac virtues.

VANILLA AROMATICA, see Vanilla.

VAO, see Curare.

VAPEURS, 'vapors,' Hypochondriasis, Hysteria—*v. de Rate*, Hypochondriasis.

VAPOR SANGUINIS, 'vapour of the blood,' Gaz sanguinis.

VAPORARIUM, *Atmist'e'rium*, *Atmolu'tron*, *Lac'on'icum*, *Bal'neum vapo'ris*, *Vapour bath*, (F.) *Bain de vapeur*, (from *vapor*, 'steam.') A remedy in rheumatic and cutaneous affections; and wherever it is important to impress the cutaneous surface. In the *Russian Vapour Bath*, the vapour is produced in Russia by throwing water over hot stones, and the bather passes, at once, from the heated condition to cool or cold water, not only with impunity, but with positive pleasure. Its temperature is from 122° to 145°.

VAPORARIUM, Hypocaustum, see Stove.

VAPORATIO, (*vaporare*, *vaporatum*, 'to emit steam or vapour,') Evaporation.

VAPORES UTERINI, Hysteria.

VAPORS, Hypochondriasis, Hysteria.

VAPOUR, Flatulence—*v. Bath*, Vaporarium.

VAPOURISH, Hypochondriac, Hysterical.

VAPOUROUS, Windy.

VAPOURY, Hypochondriacal, Hysterical.

VARA, *Barre*.

VARAIRE, *Veratrum album*.

VAREC, 'wrack,' Kelp—*v. Vésiculeux*, *Fucus vesiculosus*.

VARENI, Ambulofatulentus, Gout, (wandering.)

VARICE, Varix.

VARICELLA, (dim. of *Variola*), *Em'phlysis* seu *E'xanth'e'ma* seu *Syn'oehus varicell'a*, *Vari'ola lymphati'ca* seu *spu'ria* seu *syloes'tria*, *Vari'ole* *no'the* seu *illegiti'mæ* seu *volati'cæ* seu *pucell'æ*, *Pseudovaricell'æ*, *Æol'ion*, *Æol'tium*, *Chick'enpox*, (F.) *Varicelle*, *Pétite vérole volante*, *Vérolète*, *Vérette*, *Fausse variole*. A disease characterized by vesicles scattered over the body, which are glabrous, transparent, and about the size of peas. They appear in successive crops, are covered by a thin pellicle, and, about the third, fourth, or fifth day from their appearance, burst at the top, and concrete into small, puckered scabs, which rarely leave a pit in the skin. Three varieties have, sometimes, been designated. 1. The *V. lentiform'es* or *lentic'ular chick'enpox*, which are irregular circular, and flattened on the top; the fluid being at first whitish, and afterward straw-coloured. This is the common *chickenpox*. 2. The *Varicella conifor'mis*, *Pem'phigus variolo'des*, *Vari'ola lymphati'ca*, *Hydrach'nia*, *Conoid'al Chick'enpox*, *Suinepox*, *Waterpox*, *Waterjags*, in which the vesicles are generally acuminate, and the fluid pellucid throughout; and, 3. The *Varicell'a globu'laris* or *Hives*, in which the vesicles are globular and larger, the fluid being, at first, whey-coloured,

and afterwards yellowish. These divisions are, however, very arbitrary, for they are generally confounded.

Hornpox is a popular variety of varicella.

Variola has been, sometimes, mistaken for this disease, and conversely. A little attention to the history of the eruption in the two affections will prevent error. Varicella is not contagious. At times, it prevails epidemically. It cannot be propagated by inoculation. The treatment is extremely simple; rest, abstinence, and the antiphlogistic regimen, being all that is generally required. See Varioloid.

VARICELLA VARIOLOIDEA, Varioloid.

VARICELLAR FEVER, see Varioloid.

VARICIFOR'MES PARAS'TATÆ, (*varix*, pl. *varices*, and *forma*, 'form.') With some authors, the ducts of the epididymis, because they seem knotty and varicose.

VARICOBLEPH'ARON, (*varix*, and *βλεφαρον*, 'eyelid.') A varicose tumour of the eyelid.

VARICOCE'LE, (*varix*, and *κηλη*, 'a tumour;') *Osteocele'le varico'sa*, *Her'nia varico'sa*, *Ra'mæ varico'sus*, *Spermatocele'le*, *Cirrhocele'le*. A varicose dilatation of the veins of the scrotum and spermatic cord; usually, by English and American writers, restricted to the latter, like *Cirrhocele*. This latter variety of the disease is more common on the left side than the right. It appears under the form of a soft, doughy, unequal, knotty, compressible, and indolent tumour, situate in the course of the cord, and increasing from below upwards. It is less when the individual is in the recumbent posture, and increases when he is erect. It is, ordinarily, an affection of no moment, and is very common, requiring merely the use of a proper suspensory bandage. It has been advised, in bad cases, to tie the enlarged vein, but the operation is not free from danger induced by phlebitis. Sir A. Cooper has proposed the removal of a portion of the scrotum, so as to render the remainder of the scrotum a natural bandage. This plan has been found effectual.

VARICOM'PHALUS, (*varix*, and *ομφαλος*, 'the umbilicus.') Varicose tumour of the umbilicus; *circs'omphalus*.

VARICOSE, *Varico'sus*, *Var'icous*, (F.) *Variqueux*, (from *varix*.) Affected with *varix*: or belonging to, or connected with, *varix*;—as a *varicose vein*; a *varicose ulcer*, one kept up by *varices*.

VARICOSITAS, see Varix—*v. Conjunctivæ*, *Cirsophthalmia*.

VARICOSITY, see Varix.

VARIC'ULA, dim. of *Varix*. Also, a varicose swelling of the veins of the tunica conjunctiva.

VARI'OLA, *Vari'olæ*, *V. ve'ræ*, *Vari'oli*, *Em'pye'sis Vari'ola*, *Pes'tis* seu *Fe'bris variolo'sa*, *Æolec'thyma*, *Smallpox*, *Pox* (North of England); (from *varius*, 'spotted,' or from *vari*, 'pimples;') (F.) *Variolæ*, *Petite Vérole*, *Picote*. A disease, now of somewhat less interest than before the discovery of vaccination. It is of a very contagious nature, and is supposed to have been introduced into Europe from Asia, at an early period of the middle ages. It is characterized by fever, with pustules appearing from the third to the fifth day, and suppurating from the eighth to the tenth, and it possesses all the distinctive properties of the major exanthemata. It is capable of being produced by inoculation, but this inoculated *smallpox*—*Vari'ola inser'ta*—communicates the disease as readily through the air as the *natural smallpox*, or that received without inoculation.

Smallpox is distinguished into two classes, according to the character of the eruption, the *discrete* or *distinct*, and the *confluent*. 1. In *VARIOLA DISCRE'TA*, *Em'pye'sis vari'ola discre'ta*, *Va-*

Var'ola discre'ta benigna, *Var'iolæ regula' res seu discre'tæ*, *Distinct' Smallpox*, the pustules are usually of the size of peas; distinct, distended, and circular; the intervening spaces being red; the fever inflammatory throughout, and ceasing when the eruption is complete. 2. In *Var'iola Con'fluens*, *Empy'e'is var'iolæ con'fluens*, *Var'iolæ regula' res con'fluentes*, *Vari'olæ Di'cæ Bar'baræ*, *Vésicules de Sainte-Barbe*, *Con'fluent Smallpox*, the pustules are confluent, or run together; flaccid and irregularly circumscribed; the intervening spaces being pale, and the accompanying fever typhoid. In children, diarrhœa, and, in adults, ptialism, with swelling of the hands and feet, generally appear, towards the period of the *sec'ondary fe'ver*, which occurs from the tenth to the thirteenth day. The fever that precedes and accompanies the eruption is called the *Eruptive fe'ver*. The prognosis is favourable in the distinct kind, if properly managed. The confluent is always dangerous,—the unfavourable symptoms being—flattening of the pustules, or subsidence of the eruption; the breathing becoming much obstructed or oppressed; or marks of inflammatory and congestive affections occurring in the different viscera. The treatment of the distinct kind is very simple: the antiphlogistic regimen being all that is necessary. The confluent kind requires the same management as typhus fever.

When the pimples are confluent in patches, the patches being, however, separated by intervals of unaffected skin; it constitutes the clustered, coherent, or corymbose variety—*Var'iola corymbo'sa*.

VARIOLA AMBOINENSIS, *Frambœsia*—v. *Corymbosa*, see *Var'iola*—v. *Inserta*, *Var'iola* (by inoculation)—v. *Lymphatica*, *Varicella*—v. *Mitigata*, *Var'ioloid*—v. *Spuria*, *Varicella*—v. *Vaccina*, *Vaccina*—v. *Vaccinatorum*, *Var'ioloid*.

VARIOLA sine VARIOLIS, *Var'iolous Fe'ver*. These terms are applied to cases occasionally observed during the prevalence of *variola*, in which the constitutional phenomena are present without the cutaneous eruption.

VARIOLÆ, *Var'iola*—v. *Equinæ*, see *Grease*—v. *Illegitima*, *Varicella*—v. *Modificate*, *Var'ioloid*—v. *Nothæ*, *Varicella*—v. *Puellæ*, *Varicella*—v. *Sylvestris*, *Varicella*—v. *Tutoriæ*, *Vaccina*—v. *Voræ*, *Var'iola*—v. *Volaticæ*, *Varicella*.

VARIOLATION, *Inoculation* (*Smallpox*.)

VARIOLE, *Var'iola*.

VARIOLEUX, *Var'iolous*.

VARIOLI, *Var'iola*.

VARIOLIC, *Var'iolous*.

VARIOLIQUE, *Var'iolous*.

VARIOLOID, *Var'iol'ois*, *Var'ioloi'des*, *Var'iolæ modifi'cata seu mitiga'ta seu vaccinato'rum Varicell'la* (of some), *Varicell'la Var'ioloidæ*, (*variola*, 'smallpox,' and *ubs*, 'form,')—resembling smallpox. This is, really, smallpox, modified by previous inoculation or vaccination; and hence it has been properly called *modified smallpox*. That it is smallpox is proved by the fact, that matter, taken from a varioloid pustule, will communicate smallpox to one who has never been vaccinated. It is, almost always, a milder disease than smallpox; and this circumstance, with its shorter duration, exhibits the salutary effects of previous vaccination or inoculation. It has appeared epidemically.

Occasionally, when varioloid is epidemic, *varioloid* or *varicell'lar fe'ver* occurs without the eruption. This is the *varicell'la sine varicell'lis* of Mr. Erasmus Wilson, who calls varioloid, *varicella*.

VARIOLOID FEVER, see *Var'ioloid*.

VARIOLIS, *Var'ioloid*.

VARI'OLOUS, *Var'iolous*, *Var'iolic*. (F.) *Var'ioleux*, *Var'iolique*. Relating or belonging to *variola*.

VARIQUEUX, (from *varix*.) *Varicose*.

VARIX, gen. *Var'icis*, *Phlebeury'sma*, *Phlebec'tasia*, *Her'nia seu Ecta'sia vena'rum*, *Angiecta'sia vena'ra*, *Ix'ia*, *Cir'sus*, *Ced'ma*, *Var'icose Vein*, (F.) *Varice*, is considered, by some, to come from the verb *variare*, 'to turn,' 'to twist,'—on account of the sinuosities of varicose veins. Dilatation of a vein; *Varicos'itas*, *Varicosity*. Varices are owing to local retardation of the venous circulation; and, in some cases, to relaxation of the parietes of the veins. They are very common in the superficial veins of the lower limbs, especially of pregnant females. The tumour or tumours, formed by varices, are soft, knotty, unequal, indolent, and livid; without pulsation, and yielding readily to the impression of the finger; but returning as soon as the compression is discontinued. Sometimes, the vein bursts and gives rise to hemorrhage. The treatment is usually palliative, and consists in exerting a uniform and constant pressure upon the part, by means of an appropriate bandage. The radical cure has, also, been obtained by extirpation, division of the vein or ligature. The second course is apt to be followed by phlebitis.

VAROLI, *BRIDGE OF*, *Pons Varolii*.

VARUS, (*varius*, 'spotted,') A spot or pimple on the face: *Acne Ionthus*. Also, *Entogonyancon* and *Kylosis*, (from *varus*, 'having the legs bent,')

VARUS MENTAGRA, *Sycosis*—v. *Punctatus*, see *Acne*.

VAS, gen. *Va'sis*, *Vessel*, *Penis*—v. *Aberrans Halleri*, *Vasculum aberrans*—v. *Deferens*, see *Deferens*—v. *Deferens mulieris*, *Tuba Fallopiana*—*Urinarium*, *Ureter*.

VASA (pl. of *Vas*) *ABERRAN'TIA*. Peculiar ramifications of the biliary duct in the liver are so called by E. H. Weber.

VASA ABSORBENTIA, *Absorbent vessels*—v. *Afferentia*, see *Afferent* and *Efferent*—v. *Brevia*, see *Brevia vasa*—v. *Capillaria*, *Capillary vessels*—v. *Chylifera seu Chylofera*, *Chyliferous vessels*.

VASA CHYLOPOËTICA URINIFERA. Blood-vessels, which, according to Lippi, convey drinks from the chyliferous vessels to the renal veins.

VASA EFFERENTIA, see *Efferent*—v. *Exhalantia*, *Exhalant vessels*—v. *Graafiana*, see *Efferent*—v. *Hydragoga*, *Lymphatic vessels*—v. *Inferentia*, see *Afferent* and *Efferent*—v. *Lactea*, *Chyliferous vessels*, *Lactiferous vessels*—v. *Lymphatica*, *Lymphatic vessels*—v. *Meningea*, *Meningeal vessels*—v. *Nutrientia*, *Vasa vasorum*—v. *Nutritia*, *Vasa vasorum*—v. *Recta*, see *Rete testis*—v. *Resorbentia*, *Absorbent vessels*, *Lymphatic vessels*—v. *Sedalia*, *Hæmorrhoidal vessels*—v. *Serosa*, see *Capillary vessels*.

VASA VASO'RUM, 'vessels of vessels;' *V. nutrit'ia seu nutrit'ia*, (F.) *Vaisseaux des vaisseaux*. Small vessels that supply larger with blood for their nutrition. The cardiac or coronary arteries and veins may be regarded as the *vasa vasorum* of the heart. However minutely a vessel is traced, its parietes will be found supplied with blood for their nutrition by a smaller vessel.

VASA VORTICO'SA, (*vortex*, 'a whirlpool,') (F.) *Vaisseaux tourbillonnés*, *Tourbillon vasculaire*. The contorted vessels which creep on the choroid coat of the eye: *ciliary veins*.

VASCO'A AMPLEXICAUL'IS, (after *Vaseo* de Gama.) A South African shrub, *Nat. Ord. Leguminosæ*, the roots of which taste like liquorice. They are used, in decoction, in catarrh and phthisis, and are a good substitute for liquorice.

VASCULA SERPENTINA, see *Testicle*.

VAS'ULAR, *Vascula'ria*, *Vasculo'sus*, *Angi'at*, (F.) *Vasculaire*, *Vasculenz*, (fr. *en vas*, 'a ves-

sel,' *vasculum*, 'a little vessel.') That which belongs or relates to vessels;—arterial, venous, or lymphatic; but generally restricted to bloodvessels only. Full of vessels. Bichat gave the name *Vascular sys'tem*, *Système vasorum*, *Hæmatologica*, to the blood-vessels, and of this he made two divisions. 1. The *Arterial sys'tem*, or *Sys'tem of red blood*, (F.) *Système vasculaire à sang rouge*, which commences with the radicles of the pulmonary veins; and includes the pulmonary veins, the left cavities of the heart, and the aorta with all its ramifications. The blood, in this system, is distributed to the general capillary system, where it loses its arterial quality and becomes venous. 2. The *Venous sys'tem* or *Vascular sys'tem of black blood*, (F.) *Système vasculaire à sang noir*, carries the blood from every part of the system to the lungs. It takes its origin in the general capillary system; continues in the veins, right auricle, and ventricle of the heart and the pulmonary artery and its branches, and is finally lost in the capillary system of the lungs. See Circulation.

VASCULAR GLANDS, see Ganglion—v. Layer, see *Tache embryonnaire*—v. Radicles, Radical vessels—v. Substance of the Kidney, see Kidney—v. System, see Vascular—v. System, intermediate, Capillary system—v. System, peripheral, Capillary system—v. Tumour, see Hemorrhoids.

VASCULARITY, *Vascular'itas*, (F.) *Vascularité*. The state of being vascular, or largely supplied with vessels.

VASCULEUX, Vascular.

VASCULOMOTOR, Vasomotor.

VASCULUM, (dim. of *vas*, 'a vessel,') Penis, Vessel.

VAS'CULUM ABER'RANS, (*aberrare*, 'to wander from,') *Vas aberrans Halleri*, *Appen'dix to the Epidid'yms*. A small convoluted duct, of variable length, generally connected with the duct of the epididymis, immediately before the commencement of the vas deferens.

VASE À SAIGNER, 'bleeding vessel,' Cup.

VASOMOTOR, *Vasculomotor*, (F.) *Vasomoteur*, *Vasculo-moteur*, (from *vas*, 'vessel,' and *motor*, 'mover,') That which causes movement in the vessels. Fibres of the great sympathetic are 'vasomotor.'

VASTUS. That which is *vast* or has a great extent.

VASTUS EXTERNUS and VASTUS INTERNUS are two considerable, fleshy masses, which form part of the *Triceps cruris*.

VASUM, Vessel.

VA'TER, AMPUL/LA OF, (F.) *Ampoule de Vater*, (called after Vater, a German anatomist.) An ampulla formed by the orifice of the pancreatic duct in the duodenum.

VATER, CORPUSCLES OF, Corpuscles, Pacinian.

VATERIA INDICA, (after Abraham Vater, a physician and botanist of Wittenberg,) see Copal.

VATICINIUM CHIROMANTICUM, (*vaticinor*, 'to foretell,') Chiromancy.

VAULT, (old F. *Voute*, 'a round or turn; from *volvere*, *volutum*, 'to turn;') *For'nix*, *Cam'era*, (F.) *Voute*. A name given by anatomists to several parts which are rounded superiorly; concave and arched inferiorly,—as the *Vault of the cranium* or the upper part of the skull,—the *Palatine vault* or roof of the mouth, &c. &c.

VAUQUELINE, (after Vauquelin, the French chemist,) Strychnia.

VEAL TEA. This may be made in the same manner as beef tea, by using a pound of fillet of veal, free from fat and sliced, and a pint and a half of boiling water; and boiling for half an hour. It may also be made with the same quantity of the fleshy part of a knuckle of veal.

VECORDIA, 'want of reason,' Idiocy.

VECTIS, (*vehere*, *vectum*, 'to carry,') Lever—

v. Elevatorius, Elevator—v. Triploides, Elevator, triploid.

VECTOR CANAL, (*vehere*, *vectum*, 'to carry,') Tube, Fallopian.

VEG'ETABLE, *Vegetab'ilis*, (F.) *Végétal*, (from *vegetus*, 'lively,' and *vegeo* or *vigeo*, 'I flourish,') Appertaining or relating to plants. A vegetable, *Phyton*, (*φύτον*), is an organized being devoid of sensation and voluntary motion, which receives its nutriment, essentially, by the roots. The study of vegetables is *Phytology*, *Phytolog'ia*.

VÉGÉTAL, Vegetable.

VEGETALITY, *Vegetal'itas*, (F.) *Végétalité*, (from *vegeo* or *vigeo*, 'to be active,') The aggregate of physiological phenomena, which exist alone in the vegetable, but are common to both plants and animals. See Animality.

VEGETARIAN, see Vegetarianism.

VEGETARIANISM. Same etymon as Vegetation. A modern term, employed to designate the view, that man, for his full mental and corporeal development, ought to subsist on the direct productions of the vegetable kingdom, and totally abstain from flesh and blood. An embracer and practiser of this doctrine is called a *vegetarian*.

VÉGÉTATIF, Vegetative.

VEGETATION, *Vegeta'tio*, 'growing like a plant,' (from *vegeo*, or *vigeo*, 'to be active,') A morbid production, which rises as an excrescence on an organ or part, as on the valves of the heart. The term is also applied to the fleshy granulations or *Hypersarcosis*, which sometimes arises at the surfaces of wounds or ulcers.

VEG'ETATIVE, *Vegetati'vus*, *Phyt'ions*, (F.) *Végétatif*. Having relation to growth or nutrition; hence, the *vegetative* or nutritive functions.

VEGETATIVE LAYER, see *Tache embryonnaire*.

VEGETATIVUS, Vegetative.

VEHICLE, *Vehic'ulum*, 'a carriage,' *Oche'ma*, *Consti'tuens*, (F.) *Véhicule*, (from *vehere*, 'to carry,') Anything which serves to carry; thus air is the *vehicle* of sound; the arteries are the *vehicles* of the blood, &c. In *Pharmacy*, the name *vehicle* is given to any excipient; or to substances which serve as the media of administration for any medicine.

VÉHICULE, Vehicle.

VEILED PUFF, see Cavernous respiration.

VEIN, *Ve'na*, dim. *Ve'nula*, *Phlebs*, *Phleps*, (F.) *Veine*. The veins are vessels for the conveyance of blood from every part of the body to the heart. They are found wherever there are arteries, and, altogether, form the *venous system*, which may be subdivided into two distinct secondary systems. 1. The *Gen'eral ve'nous sys'tem*, which commences in all the organs, by very minute radicles; and terminates in the heart by the cava and the coronary vein. 2. The *Abdom'inal ve'nous sys'tem*, which is limited to the abdominal cavity; commences, also, by a great number of branches, and terminates in the liver by a single trunk, which subdivides in that organ. It is called, also, the *Sys'tem of the Ve'na Porta* or the *Portal Sys'tem*. See Porta. The veins form two sets, one *deepseated*, *Ve'næ satel'lites*, *V. concom'ites*, which accompany the arteries; the other *superf'ial* or *subcuta'neous*. Their parietes, which are thinner than those of the arteries, are composed of three coats; the outermost, of dense areolar tissue; the middle coat elastic, with parallel, longitudinal fibres—the *proper membrane*; the other, the *common* or *inner membrane*, which is thin, polished, and has numerous duplicatures or folds forming parabolic valves. These valves are most numerous in the veins of the lower extremities. The coats of the veins are supplied with small arteries or *vasa vasorum*, veins, and lymphatics. They have not as many nervous filaments as the arteries. These filaments,—*vasomotor*,—are from the great sympathetic.

TABLE OF THE VEINS.

1. VEINS WHICH FORM THE SUPERIOR VENA CAVA.

1. Subclavians.	Receive the 1. Axillary which	1. Receives the 1. Basilic, formed of the 2. Cephalic, forming the 3. Circumflex veins. 4. Inferior scapular. 5. Long thoracic. 6. Superior thoracic. 7. Acromial veins.	1. Posterior ulnar. 2. Anterior ulnar. 3. Median basilic. 1. Superficial radial. 2. Median cephalic.
	2. External Jugular.	1. Receives the 1. Internal maxillary, composed of the 2. Superficial Temporal, composed of the 3. Posterior Auricular. 4. Cervical cutaneous. 5. Tracheloscapular, &c.	1. Pterygoid. 2. Sphenopalatine. 3. Alveolar. 4. Infraorbital. 5. Mental. 6. Inferior dental. 7. Deep temporal. 1. Middle temporal. 2. Anterior auriculars. 3. Transverse of the face.
	3. Internal Jugular.	1. Receives 1. Superior Cerebral Veins. 2. Veins of the Corpus Striatum. 3. Veins of the Choroid plexus. 4. Superior Cerebellar Veins. 5. Inferior Cerebellar Veins. 6. Lateral and inferior Cerebral Veins. 7. Ophthalmic Vein, composed of the 8. Facial vein, called <i>angular</i>, near the eye, receiving the 9. Lingual and Pharyngeal Veins. 10. Superior thyroid. 11. Occipital, and 12. Veins of the Diploë.	1. Lachrymal. 2. Central of the retina. 3. Infraorbital. 4. Ciliary. 5. Ethmoidal. 6. Palpebral, and 7. Nasal Veins. 1. Palpebral, and 2. Superciliary Veins. 3. Dorsals of the nose. 4. Superior Coronaries 5. Inferior } of the lip. 6. Several Buccal, and 7. Masseteric veins. 8. Ranine. 9. Submental, and 10. Inferior Palatine.

2. Right Internal Mammary Vein.
3. Inferior Thyroid Vein, opening into the Vena Cava, between the two Subclavians.
4. Vena Azygos, which receives the
- 1. Right Bronchial.
 - 2. Intercoastal Veins.
 - 3. Semiazygos.

2. VEINS WHICH FORM THE INFERIOR VENA CAVA.

1. Common Iliacs.	Receive the 1. External Iliacs, formed by the—	1. Femoral or Crural, which is a continuation of the 2. Popliteal, commencing by 3. veins which accompany the fibular arteries, and receiving the—	1. External abdominal veins. 2. Circumflex Iliac, and 3. External pudic veins.
	2. Internal Iliacs which receive the—	1. Vesical Veins commencing with the 2. Sacrolateral Veins.	1. Dorsal veins of the penis of the male. 2. Veins of the clitoris, in the female.

3. Middle Sacral Vein.
4. Lumbar Veins—four in number on each side, commencing by an—
- 1. Abdominal branch.
 - 2. Dorsal branch.
5. Spermatic Veins, commencing with the
- 1. Spermatic plexus in the male.
 - 2. Those of the Ovarium, Fallopian tube, &c., in the female.
6. Renal Veins.
7. Capsular and Adipose Veins.
8. Hepatic Veins.
9. Middle, and
10. Left, and
11. Right, } Hepatic Veins
12. Inferior Diaphragmatic Veins, two in number.

3. VEINS OF THE HEART.

1. Great right Coronary.
2. Small right Coronary.
3. Left Coronary Veins.

4. VEINS WHICH FORM THE VENA PORTA

1. Splenic Vein.
2. Superior Mesenteric Vein.
- Receives the
- 1. Veins which correspond to the vasa brevia.
 - 2. Right and left gastroepiploic.
 - 3. Duodenal, and
 - 5. Pancreatic veins.
 - 5. Coronary vein of the stomach, and
 - 6. Small mesenteric vein.

VEIN OF BRESCHET, see Diploic—v. Bronchial, left, see Azygos—v. Coronary, Cardiac veins—v. Demiazygous, see Azygous—v. Diploetic, see Diploic—v. Frontal, see Facial vein—v. of Galen, see Cardiac veins, and Choroides venæ—v. Intralobular, Hepatic veins—v. Palatolabial, Facial vein—v. Peroneomalleolar, Saphena minor—v. Semiazygous, see Azygous vein—v. Superior intercoastal, left, see Azygous vein—v. Suprahepatic Hepatic veins—v. Tibiomalleolar, Saphena, great—v. Varicose, Varix.

VEINE, Vein—v. Cave, Cava vena—v. Céphalique, Jugular vein, internal—v. Céphalique du pouce, Cephalic vein of the thumb—v. Cubitale cutanée, Basilic vein—v. Fessière, Gluteal vein—v. Hypogastrique, Hypogastric vein—v. de Méline, Dracunculus—v. Porte, Porta vena—v. Prélombothoracique, Azygous vein—v. Préparate, Facial vein—v. Radiale cutanée, Cephalic vein—v. Sans paire, Azygous vein—v. Trachélo-sous-cutanée, Jugular vein, external.

VEINES ARTÉRIELLES, Arterial veins—v. Ciliaires, Ciliary veins—v. Sous-clavières, Subclavian veins—v. Sus-hépatiques, Hepatic veins.

VEINEUX, Venous.

VEINSTONE Phlebolite.

VEINULE, (dim. of *veine*), Venula.

VEJUDO DE MAVACURE, see Curare.

VELAMEN ('a covering') NATIVUM, Integument.

VELAMENTA (pl. of *velamentum*, 'a covering') BOMBYCINA, Villous membranes—v. Cerebralia, Meninges—v. Infantis, Membranes of the foetus.

VELAMENTUM ('a covering') ABDOMINALE, Peritonæum—v. Corporis commune, Integument—v. Linguae, Glossoepiglottic ligament—v. Nativum, Integument.

VELAR, Erysimum.

VELLICATION, (*vellicare*, *vellicatum*, 'to twitch,') Tic.

VELOSYNTHESIS, (*velum*, and *σύνθεσις*, 'a putting together,') Staphylorrhaphy.

VELUM, 'a veil, a cover,' Velum pendulum palati—v. Apophysi vermiformi obtentum, Valvula Viuessenii—v. Interjectum cerebelli, Valvula Viuessenii—v. Medullare anticum, Valvula Viuessenii.

VELUM MEDULLARE INFANTIS, (F.) Voile ml.

dullaire inférieur. A name given, by Reil, to the medullary band or strip, by which the cerebellum communicates with the medulla.

VELUM MEDULLÆ POSTERIORIS seu **POSTICUS** seu **SEMILUNA'RE**, *Val'vula Tari'ni.* The small posterior valve of the cerebellum; a thin medullary plate, situate between the medulla oblongata and vermis inferior, in front of the nodulus.

VELUM PALATI, Velum pendulum palati—v. Palatinum, Velum pendulum palati.

VELUM PENDULUM PALAT'I, 'pendulous veil of the palate.' *Ve'tum*, *V. Palati'mum* seu *Palat'i* seu *Pendulum* seu *Staphyl'inum*, *Palat'um pendulum* seu *mol'le* seu *mob'ile*, *Pendulum* seu *Clastrum Palat'i*, *Palat'um Mol'le*, *Soft Pal'ate*, (F.) *Septum Staphylin* (Ch.), *Voile du Palais*. A movable, soft, and broad septum, attached to the posterior extremity of the palatine arch, and separating the mouth from the pharynx. Its form is nearly quadrilateral. Its anterior surface corresponds to the mouth; the posterior to the pharynx. Its superior edge is fixed to the palatine arch; the inferior is free, floating, and has an appendix at its middle—the *uvula*. The lateral margins of the velum are held down by two folds, separated from each other by a triangular space; these are called the *pillars*; the anterior passes to the base of the tongue, the posterior to the sides of the pharynx. The velum palati is formed of a mucous coat, which contains numerous follicles, and of a muscular stratum, in which are the peristaphylini interni and p. externi muscels, the glossostaphylini, the pharyngostaphylini and the palatostaphylinus. The arteries of the velum palati are furnished by the internal maxillary, the labial and the superior pharyngeal. Its veins open into the internal jugular. Its nerves are sent off from the ganglion of Meckel, and come from the palatine branches. The glossopharyngeal nerve, likewise, sends it some filaments.

VELUM PUPILLÆ, Pupillary membrane—v. Staphylinum, Velum pendulum palati—v. Vienssenii, Valvula Vienssenii.

VELVET LEAF, Abutilon Avicennæ, Pareira.

VENA, Vein, Penis, — v. Alba renum, Ureter—v. Alba thoracis, Thoracic duct—v. Apoplectica, Jugular vein, internal—v. Arteriosa, Porta vena, Pulmonary artery—v. Capitis, Cephalic vein—v. Cava, Cava vena—v. Cubiti interior, Basilic vein—v. Demiazygos, see Azygos—v. Diploeticæ, see Diploic—v. Falcis cerebri, Sinus longitudinalis inferior—v. Galeni, see Cardiac veins, and Choroideæ venæ—v. Hemiazyga, see Azygous vein—v. Lactea, Porta vena—v. Lienalis, Splenic vein—v. Magna, Porta vena—v. Medina Arabum, Dracunculus—v. Medinensis, Dracunculus—v. Ostiaria, Porta vena—v. Pari carens, Azygos, (vena)—v. sine Pari, Azygos, (vena)—v. Portalis, Porta vena—v. Portarum, Porta vena—v. ad Portas, Porta vena—v. quæ ad Portas est, Porta vena—v. Præparata, Facial vein—v. Ramalis, Porta vena—v. Splenica, Splenic vein—v. Stelechica, Porta vena—v. Subalaris, Vena axillaris—v. Terminalis, Circulus venosus.

VENÆ, pl. of Vena—v. Apoplecticæ, Jugular veins—v. Cavæ hepatis, Hepatic veins—v. Centrales, Intralobular veins—v. Concomites, see Vein—v. Galeni, Choroid, (veins)—v. Inominate, see Inominatum—v. Intralobulares, Intralobular veins—v. Lactæe, Chyliferous vessels—v. Lymphaticæ, Lymphatic vessels—v. Micantes, (arteries,) see Artery—v. Pulsatiles, (arteries,) see Artery—v. Satellites, see Vein—v. Soporales, Jugular veins—v. Spermaticæ, Spermatic veins—v. Sphagitides, Jugular veins.

VENÆ VORTICOSÆ. The veins which principally form the external or venous layer of the choroid coat of the eye; so called from their peculiar arrangement; (from *vertex* seu *vortex*, 'a whirlpool'.)

VENÆSECTIO, (*vena*, 'a vein,' and *secare*, *sectum*, 'to cut,') see Bloodletting—v. Revulsoria, see Bloodletting.

VENAL, Venous.

VENEFICIUM, (*venenum*, and *facere*, 'to make,') Poisoning.

VENENATION, (from *venenum*,) Poison, Poisoning, Venom.

VENENATUS, Venomous.

VÉNÉNEUX, Poisonous.

VENENOSE, Venomous.

VENENUM, Poison, Venom.

VENÉREAL, *Venereus*, *Aphrodisi'acus Venerean*, (F.) *Vénérien*, (from *Venus*, gen. *Veneris*, 'the goddess of pleasure,') That which relates to the pleasures of love. In a more restricted sense, the word *venereal*—as well as *aphroditic*—is synonymous with *syphilitic*, and also with *syphilis*. Some authors have proposed that those diseases which are produced by excess in venery should be called *venereal*; and that *syphilitic* should be limited to those which are the result of impure connexion. It is, also, used in the same sense as *aphrodisiac*,—adapted to excite venereal desire.

VENEREAL DESIRE, Libido—v. Diseases, Syphilis.

VÉNÉRIEN, Venereal. Also, one labouring under syphilis; as *Hôpital des vénériens*.

VENERIVAGUS, (*venus*, gen. *veneris*, 'the pleasures of love,' and *vagare*, 'to wander,') Libidinous.

VENERY, (from *venus*, gen. *venereis*,) The pleasures of love. Sexual communication. Coition.

VENESECTIO, (*vena*, 'a vein,' and *secare*, *sectum*, 'to cut,') see Bloodletting.

VENIMEUX, Venomous.

VENIN, (from *venenum*,) Venom.

VENO BENO, see Falsification, table of.

VENOM, *Venenum*, *Tox'icum*, *Venenat'ion*, (F.) *Venin*. A poisonous fluid secreted by certain animals, as the viper, in a state of health, and which they preserve in a particular reservoir, to use as a means of attack or defence. It differs, according to some, from *virus*, as the latter is always the result of a morbid process, which may be produced by the individuals who receive it, in their turn, and be transmitted to others.

VENOM, QUICKENING, Sperm—v. Vital, Sperm.

VENOMOUS, *Venenat'us*, *Ven'ense*, (F.) *Venimeux*. An epithet applied to animals which have a secretion of venom, as the viper, rattlesnake, &c., as well as to the venom itself; and, by some, to liquids in the animal body, which have been so perverted by previous disease, that their contact occasions serious mischief in sound individuals; as happens in hydrophobia. Virus is more commonly used in this case.

VENOSITAS, (from *vena*, 'a vein,') Venosity—v. Sanguinis, Prædominium sanguinis venosi.

VENOSITY, *Venos'itas*, *Ven'ousness*, *Venous state* or *cachex'ia*, *Melaném'ia*, *Melano'sis universalis*, (F.) *Mélanémie*, (from *vena*, 'a vein,') A condition in which, it has been supposed, the blood is moved slowly; is more venous; and the venous blood itself in greater quantity. This state is said to be chiefly observed in hemorrhoids, gout, hypochondriasis, melænholia, &c. In general, it signifies abundance of veins in a part; afflux of blood into the veins.

VENOUS, *Venal*, *Venous*, (F.) *Veineux*, (from *vena*, 'a vein,') Appertaining to a vein, or to the blood of the veins; as *venous cachexia*, *venous pulse*.

VENOUS CACHEXIA, Venosity—v. Hum, *Bruit de Diable*—v. State, Venosity—v. System, see Vascular—v. System, abdominal, Porta vena.

VENOUS SYSTEM of JACOBSON. A system of vessels in the three classes of vertebrate ani-

imals other than the mammalia, which forms a communication between the vena porta and the vena cava.

VENOUSNESS, Venosity.

VENT, ([L.] *ventus*.) Wind.

VENTER, gen. *Ventris*, Abdomen, Belly, Uterus. In law, venter designates the maternal parentage of children. Thus, if A were B's child by his first wife, he would be described in law as 'by the first venter.' 'A died seised, leaving two infant daughters by different venters.'

VENTER IMUS, 'lowest belly,' Abdomen, Hypogastrium—v. Inferius, Abdomen—v. Infimus, Abdomen—v. Magnus, Inguities—v. Medius, Thorax—v. Musculi, see Muscle—v. Parvus, Hypogastrium—v. Propendens, Hysteroloxia anterior—v. Renum, Pelvis of the kidney—v. Supremus, Cavum crani.

VENTEUX, (from *ventus*, 'wind,') Windy.

VENTILA'TION, *Ventila'tio*, (from *ventilare*, *ventilatum*, 'to fan,') (F.) *Ventilation*, *Aérage*, *Aération*. The act of renewing the air of vessels, hospitals, and apartments, or of situations where many people are collected together. The utility of such renewal must be evident, when it is considered that atmospheric air loses its oxygen during respiration, and carbonic acid supplies its place. Stagnant air, also, becomes loaded with numerous impurities, and where numbers are crowded together, a large amount of zymotic or fermentable matter is exhaled from the lungs, and taken in again during respiration. This is, indeed, a great cause of the mortality which occurs in camps and cities, and hence one of the greatest hygienic improvements, of modern times, has been a proper attention to circulation of air, and various arrangements have been made for this purpose in hospitals and similar establishments. They are almost all, however, based on the renewal of air by suction, forcing, or both.

Ventilation is one of the most valuable additions to our stock of therapeutical agencies in many diseases.

Instruments used for the purpose of renewing the air, in ships, apartments, &c., have been called *ventilators*.

VENTOSITÉ, Flatulence.

VENTOSITY, (from *ventus*, 'wind,') Flatulence.

VENTOUSES SCARIFIÉES, (from *ventosus*, 'full of wind,') Cupping with the scarificator.

VENTOUSES SÈCHES, Cupping dry.

VENTRAL, (from *venter*, 'the belly,') Abdominal.

VENTRAL HER'NIA, *Her'nia ventra'lis*. Direct hernia occurring in any part of the abdomen. It is most frequently found between the recti muscles.

VENTRALE, 'a belly band,' Belt (Russian.)

VENTRALE CUTA'NEUM, 'cutaneous apron.' (F.) *Tablier*. An elongation of the nymphæ in the Bosjesman female, which hangs down before the vulva like an apron.

VENTRE, (from *venter*, 'the belly,') Abdomen—v. *Inférieur*, Abdomen—v. *Relâché*, see *Diarrhœa*—v. *Resserré*, Constipation.

VENTRE INSPICIENDO DE, see *De ventre inspiendo*.

VENTRICLE, *Ventriculus*, (dim. of *venter*, 'a belly,') (F.) *Ventricule*. Literally, a little belly. A name given, in anatomy, to various parts.

VENTRICLE OF ARAN'TIUS. A small cavity, lined with gray substance, at the point of the calamus scriptorius.

VENTRICLES OF THE BRAIN, *Ventriculi seu Cavita'tes seu Sinus Cerebri*, (F.) *Ventricules du cerveau*, are five cavities in the interior of that organ, which are distinguished into the *lateral*

ventricles, *mid'dle ventricle*, *fourth ventricle* or *ventricle of the cerebellum*, and *fifth ventricle*. The *mid'dle ventricle* or *third ventricle*, (F.) *Ventricule moyen du cerveau*, is situate in front of the pineal gland. It is bounded, *above*, by the tela choroidea and the fornix; *below*, by a portion of cerebral substance, which separates it from the base of the cranium; *laterally*, by the thalami nervorum opticorum; *anteriorly*, by a medullary cord, called the *anterior commissure*; and, *posteriorly*, by another medullary cord, called the *posterior commissure*. It communicates with the lateral ventricles by two openings, situate behind the anterior pillar or crus of the fornix. The *Lateral ventricles*, *Ventriculi laterales seu majores seu majores seu priores seu superiores seu tricornes*, (F.) *Ventricules latéraux*, are two in number. They are large, and, owing to their considerable curvature, occupy a great part of the cerebral centre. They are bounded, *above*, by the inferior surface of the corpus callosum; *anteriorly*, by the curved portion of the same body; *posteriorly*, by a considerable mass of cerebral substance; and, *internally*, they are separated from each other by the septum of the ventricles, — a medullary plate, which is given off from the lower part of the corpus callosum. The lateral ventricles have, in their upper and anterior portion, the *Corpora striata*, *Thalami optici*, and *Tentoria semicircularia*: — their inferior portion contains the *corpus fimbria'tum*, *corpus Ammonis*, &c. Each of the ventricles has a *body* or *centre* — *cella media* — and *horns*, *cornua*. The *Fourth ventricle*, *Ventriculus cerebelli seu quartus seu Arantii*, *Sinus rhomboideus*, (F.) *Quatrième ventricle*, is situate beneath the aqueduct of Sylvius, by which it communicates with the third ventricle. It extends from this aqueduct to the upper part of the medulla; and is seated in the substance of the tuber annulare. The *Fifth ventricle*, *Ventriculus septi pellucidii*, *Incisura septi*, (F.) *Cinquième ventricle*, is the small serous cavity between the laminae of the septum lucidum; called, also, *Sinus of the septum lucidum*, *S. of the median septum* (Ch.).

VENTRICLES OF THE HEART, *Ventriculi cordis*, *Spicus cordis*, (F.) *Ventricules du cœur*, are two in number. The *Right ventricle*, *Pulmonary ventricle*, *Ventriculus anterior seu primus seu pulmonalis*, (F.) *Ventricule droit ou pulmonaire*, is situate at the fore-part of the heart, the base and apex corresponding to those of the heart. In it are seen the *Columnae carneae* and *Corde tendineae*. The right ventricle communicates with the right auricle by the *Ostium venosum*, and from it the *Pulmonary artery* arises. Around the auricular orifice is a tendinous margin or ring, from which the tricuspid valve is given off. The use of the ventricle is to send the blood, by its contraction, along the pulmonary artery to the lungs. The *Left ventricle*, *Ventriculus aërticus seu posterior seu secundus*, (F.) *Ventricule gauche ou aortique*, is situate at the posterior and left part of the heart. Its parietes are thrice thicker and stronger than those of the right, owing to its having to send the blood over the whole system. The inner surface has the same general appearance as that of the right. The *Columnae carneae* and *Corde tendineae* are, however, much stronger. It communicates with the left auricle by means of the *Mitral valve*; and from it proceeds the largest bloodvessel of the body — the aorta.

VENTRICLES OF THE LARYNX, *Ventriculi seu Sacculi laryngis*, *Ventriculi Morgagni*, *Sinus of the larynx*, (F.) *Ventricules ou Sinus du larynx*, are two depressions in the larynx, comprised between the superior and inferior ligaments.

VENTRICLE, PULMONARY, V. right.

VENTRICOUS, *Ventricosus*, *Ventricosus*, (F.) *Ventru*. Bellied. Having a large belly.

VENTRICULAR, *Ventriculæ ris*, (F.) *Ventriculaire*. Relating to a ventricle.

VENTRICULE, *Ventricule*.

VENTRICULE AORTIQUE, *Ventriele*, left—*v. Cinquième*, *Ventriele*, fifth—*v. Droit*, *Ventriele*, right—*v. Gauche*, *Ventriele*, left—*v. Moyer du Cerveau*, *Ventriele*, middle, of the brain—*v. Pulmonaire*, *Ventriele*, right—*v. Quatrième*, *Ventriele*, fourth.

VENTRICULES DU CERVEAU, *Ventricles* of the brain—*v. du Cœur*, *Ventricles* of the heart—*v. Latéraux*, *Ventricles*, lateral.

VENTRICULI CORDIS, (pl. of *ventriculus*), *Ventricles* of the heart—*v. Dilatatio*, *Gastrectasis*—*v. Dissolutio*, *Gastromalacia*—*v. Duræ matris*, *Sinuses* of the dura mater—*v. Emollitio*, *Gastromalacia*—*v. Exulceratio*, *Gastreleosis*—*v. Fluxus*, *Gastrorrhœa*—*v. Inflammatio*, *Gastritis*—*v. Laryngei*, *Ventricles* of the larynx—*v. Laterales*, see *Ventricles*, lateral, of the brain—*v. Magni*, *Ventricles*, lateral, of the brain—*v. Majores*, *Ventricles*, lateral—*v. Metamorphosis gelatiniformis*, *Gastromalacia*—*v. Morgagnii*, *Ventricles* of the larynx—*v. Perforatio*, *Gastrobrosis*—*v. Priores*, *Ventricles*, lateral—*v. Pseudophlogosis resolutiva et colligativa*, *Gastromalacia*—*v. Superiores*, *Ventricles*, lateral—*v. Tricornes*, *Ventricles*, lateral—*v. Ulceratio*, *Gastreleosis*.

VENTRICULUS, (dim. of *venter*, 'belly,') *Stomach*, *Ventriele*—*v. Anterior*, *Ventriele*, right—*v. Aorticus*, *Ventriele*, left—*v. Arantii*, see *Ventricles* of the brain.

VENTRICULUS CALLOSUS, 'callous stomach.' The gizzard or third stomach of the gallinaceous bird. (F.) *Gésier*. It consists of a doublebellied or *digastric* muscle; is lined by a firm cuticle, and forms a kind of pouch, depending from the alimentary tube, in which the second stomach terminates, and from which the duodenum or *Ventriculus succenturiatus* issues. In the interior of the gizzard there are numerous pebbles, which seem to be destined to aid in the trituration of the food. The gizzard is an organ of mastication, and is possessed of great muscular power.

VENTRICULUS CEREBELLI, see *Ventricles* of the brain—*v. Intestinalis*, *Abomasus*—*v. Posterior*, *Ventriele*, left—*v. Primus*, *Ventriele*, right—*v. Pulmonalis*, *Ventriele*, right—*v. Quartus*, see *Ventricles* of the brain—*v. Secundus*, *Ventriele*, left—*v. Septi Pellucidi*, see *Ventricles* of the brain—*v. Succenturiatus*, *Duodenum*; see *Ventriculus Callosus*.

VENTRIÈRE, (from *venter*, 'belly, uterus,') *Midwife*.

VENTRILOQUE, (*venter*, 'belly, and loquor

'to speak,') *Engastrimythus*.

VENTRILOQUISM, see *Engastrimyth*.

VENTRILOQUIST, see *Engastrimyth*.

VENTRILOQUY, see *Engastrimyth*.

VENTRIOSUS, (from *venter*,) *Abdominous*.

VENTRIPOTENT, (*venter*, and *potens*, 'powerful,') *Abdominous*.

VENTROSITAS, (from *venter*,) *Physconia*.

VENTROSUS, *Abdominous*.

VENTS, (pl. of *vent*, [L.] *ventus*, 'wind,') *Flatulence*.

VENTUS, *Wind*.

VE'NULA, (dim. of *vena*,) *Phleb'ion*. See *Vein*. (F.) *Veinule*, *Vénule*. A small vein. Hippocrates seems to use it in the same sense as *Arteria*.

VENULÆ CENTRALES, *Intralobular veins*—*v. Intralobulares*, *Interlobular veins*.

VÉNULE, *Venula*.

VENUS, gen. *Veneris*, 'the goddess of love,' *Clitoris*, *Coition*, *Cuprum*, *Venery*—*v. Crystals* of, *Cupri subacetis*—*v. Crystaux de*, *Cupri subacetis*.

VER, (from *vermis*,) *Worm*—*v. Cucurbitin*, *Tœnia solium*—*v. Cutané*, *Dracunculus*—*v. Filaire*, *Dracunculus*—*v. de Guinée*, *Dracunculus*—*v. Luissant*, *Cicindela*—*v. de Médine*, *Dracunculus*—*v. Sanguin*, *Hæmatozoon*—*v. Solitaire*, *Bothriocephalus latus*, *Tœnia solium*.

VÉRAIRE BLANC, *Veratrum album*.

VÉRATRE BLANC, *Veratrum album*.

VERATRIA, *Veratrina*.

VERATRINA, *Vera'tria*, *Veratri'num*, *Ver'atrin*, *Ver'atrine*, (from *veratrum*.) A vegetable alkali, found in the seeds of *Vera'trum sabadill'a*, *V. officina'le*, *Helo'nias officina'lis*, and *Asagra'a officina'lis*, which are known in commerce under the names *Cebadill'a*, *Cevadill'a*, and *Sabadill'a*, and are imported from Mexico. They usually occur, in commerce, mixed with the fruit of the plant, are two or three lines long, of a black colour, and are shining, flat, shrivelled, winged, and elastic seeds. *Veratrine* is solid, white, pulverulent, and inodorous; of a very acrid taste; decomposable by heat; very soluble in water, and soluble in alcohol. When prepared in the usual manner, it still contains other substances,—*Sabadill'in*, *Ver'atrin*, gum resin of the *sabadilla*, and black greasy matter. It is extremely poisonous, and but little used internally in medicine. Magendie gave it as an emetic and drastic in very small doses. It has been recommended to be applied in the way of friction, or endermically, (from gr. v to xx to ʒj of lard or cerate,) in various neuralgic affections. The following is the formula for its preparation in the Pharmacopœia of the United States:—Take of *Ceradilla*, lbij; Alcohol, 2 gallons; Sulphuric acid, *Solution of ammonia*, *Purified animal charcoal*, and *Magnesia*, each a sufficient quantity. Boil the *cevadilla* in a gallon of the alcohol, in a retort with a receiver attached, for an hour, and pour off the liquor. To the residue add another gallon of the alcohol, together with the portion recently distilled; again boil for an hour, and pour off the liquor. Repeat the boiling a third time with the remaining alcohol, and with that distilled in the previous operation. Press the *cevadilla*, mix and strain the liquors, and, by means of a waterbath, distil off the alcohol. Boil the residue three or four times in water acidulated with sulphuric acid, mix and strain the liquors, and evaporate to the consistence of syrup. Add *magnesia* in slight excess, shake the mixture frequently, then express and wash what remains. Repeat the expression and washing two or three times, and having dried the residue, digest it with a gentle heat several times in alcohol, and strain after each digestion. Distil off the alcohol from the mixed liquor, boil the residue for fifteen minutes in water with a little sulphuric acid and purified animal charcoal, and strain. Having thoroughly washed what remains, mix the washings with the strained liquor, evaporate with a moderate heat to the consistence of syrup, and then drop in as much solution of ammonia as may be necessary to precipitate the *veratria*. Lastly, separate and dry the precipitate.

An analogous form is given in the British Pharmacopœia for 1864.

VERATRINE, *Veratrina*.

VER'ATRINE or VERA'TRIA, SUL'PHATE OF, *Vera'trix sul'phas*, (F.) *Sulfate de Véatrine*, possesses the same virtues as *veratrine*. A solution of one grain in two ounces of distilled water has been suggested as a substitute for the *Enu médicina'le d'Husson*—a celebrated gout remedy—in the dose of fʒj to fʒij.

VERATRINUM, *Veratrina*.

VERA'TRUM AL'BUM, (*verē atrum*, 'truly black,' from the colour of the roots [?]). *Helleb'orus al'bus*, *Elleb'orum al'bum*, *Elleb'orus al'bus*,

Cher'bachem, White *hell'ebore*, *Lang'wort*, (F.) *Vérase blanc*, *Hellébore on Ellébore blanc*, *Véraise blanc*, *Vuraire*, *Tue-chien*. *Sex. Syst.* Polygamia Monœcia. *Nat. Ord.* Melanthaceæ (Lindley). The root or rhizoma of this common European plant is indebted to the veratrine for its properties. It is inodorous; has a bitterish, acrid, nauseous taste, and is violently emetic and purgative, even when applied to an ulcerated surface. It is, also, possessed of emetic properties, and acts, externally, as a stimulant. It is rarely exhibited internally. Sometimes it is given as a *sneezing powder*; but it operates with great violence. It forms, also, an itch ointment. See Unguentum Veratri.

VERATRUM LUTEUM, *Chamælorium luteum*.

VERATRUM SABADILLÆ, *Sabadilla*, *Sabatilla*, *Carnis Interfector*, *Caus'ticum America'num*, *Cevadilla Hispanorum*, *Seradilla*, *Hor'deum caus'ticum*, (F.) *Cévadille*. The seed—dried fruit—of this plant, *Sabadilla*, *Cevadilla*, (Ph. U. S. and Ph. B.) have been used mostly as anthelmintics. Dose, gr. ij to a child four years old. It possesses the same general properties as the veratrum album.

VERATRUM VIRIDÈ, *American Hell'ebore*, *Swamp hell'ebore*, *Indian poke*, *Indian Un'cas*, *Poke'weed*, *Bear'weed*, *Ich'weed*, *Tickleweed*, *Earth'gall*, *Dev'ilbit*, *Wolf'bane*, *Pup'pet root*, is indigenous in the swampy districts of almost every part of the United States. The properties of the rhizoma are like those of the genus. It has been brought forward, however, in recent times, in the Southern States, as a powerful sedative in inflammatory diseases. Dose, eight drops of a saturated tincture, every three hours, until its sedative effects are induced.

VERBASCULUM, *Primula veris*.

VERBASCUM ALATUM, (supposed to be a corruption from *barbacum*, 'bearded,' owing to its hairy coat; *barba*, 'a beard,' [?]) *V. thapsus*—*v. Album*, *V. thapsus*—*v. Collinum*, *V. thapsus*—*v. Crassifolium*, *V. thapsus*—*v. Elongatum*, *V. thapsus*—*v. Lychnitis*, *V. thapsus*—*v. Montanum*, *V. thapsus*.

VERBASCUM NIGRUM, *Cande'la re'gia*, *Thapsus barbat'us*, *Thapsus barbat'us*, *Candela'ria*, *Lan'aria*, *Black Mul'lein*; and VERBASCUM THAPSUS, *V. thapsoides seu alatum seu collinum seu album seu crassifolium seu elongatum seu lychnitis seu montanum seu simplex*, *Phlog'mus*, *Phlo'mus*, *Phlox*, *Great broadleaved or yellow Mul'lein*, *High Ta'per*, *Cow's Lung'wort*, (F.) *Molène*, *Bouillon blanc*, *Bonhomme*, *Cierge de notre dame*, *Herbe Saint-Fiacre*. *Ord.* Scrophulariaceæ, *Sex. Syst.* Pentandria Monogynia,—especially the latter,—have been used as emollients, applied externally in the form of fomentation. The large leaves have been chosen for this purpose. Occasionally, the mucilage which they contain has been extracted by decoction or infusion, and administered in pectoral affections.

VERBASCUM SIMPLEX, *V. thapsus*—*v. Thapsoides*, *V. thapsus*—*v. Thapsus*, see *V. nigrum*.

VERBENA HASTATA, *V. paniculata*, *Blue ver'vain*, *Simpler's Joy*, an American species, is more bitter than the European, and said to be emetic.

VERBENA OFFICINALIS, *Verbena*, *Dichro'mus*, *Verben'aca*, *Perist'ricum*, *Hierobot'anz*, *Her'ba sa'era*, *Ver'vain*, *Pur'vain*, (Prov.) *Ber'bine*, *Juno's Tears*, (F.) *Verveine*, *Herbe à tous les maux*. *Ord.* Verbenaceæ. *Sex. Syst.* Diandria Monogynia. A plant formerly held sacred; used in sacrifices, and suspended round the neck as an amulet. In the form of cataplasm, it is said to have relieved obstinate headaches, and to have been of service in scrofula. It is not used.

VERBENA PANICULATA, *V. hastata*.

VERBENA URTICIFOLIA, an indigenous spe-

cies, has been used in poisoning from *Rhus Toxicodendron*.

VERBENACA, *Verbena officinalis*.

VERBESINA ACME'LLA, (changed from *verbena*, [?]) *Spilanthus Acmella*.

VERBESINA VIRGINICA, *Crown Beard*. Indigenous. *Order*, Compositæ. A decoction of the roots is used by the Indians as a diaphoretic and depurative.

VERCOQUIN, (ver, 'a worm,' and coquin, 'a rascal,') *Phrenitis vermino'sa*. A form of phrenitis, attributed to the presence of a worm in the brain!

VERDET, (from verd, [L.] *viridis*, 'green,') *Cupri subacetas*.

VERDIGRIS, (from [F.] *verd*, [L.] *viridis*, 'green,' and *gris*, 'gray,') *Cupri subacetas*.

VERENDA, (from *vereor*, 'to reverence,') Genital organs.

VERETRUM, (from *vereor*, [?]) Penis.

VÉRÉTTE, *Varicella*.

VERGE, (from [L.] *virga*, 'a rod,') Penis—*v. d'Or*, *Solidago vigaurea*.

VERGERETTE DE PHILADELPHIE, *Erigeron Philadelphicum*.

VERGETTURES, (from *virga*, 'a rod,') *Vibices*.

VERJUICE, *Om'phacè*, *Ompha'cion*, *Ompha'cium*, (F.) *Verjus*, *Verdjus*, (*verd*, 'green,' and *jus*, 'juice,') 'The juice of green fruits.' A sour kind of grape. Also, an acid liquor, prepared from very sour grapes or crabapples. It is principally used in culinary preparations; although, occasionally, an ingredient in medicinal compounds. See *Vitis vinifera*.

VERJUS, *Verjuice*.

VERMES, (pl. of *vermis*,) Worms—*v. Cerebri*, *Choroid plexus*—*v. Intestini*, see Worms.

VERMICELLI (I.), (from [L.] *vermiculus*, dim. of *vermis*, 'a worm,' from its resemblance,) see *Turundæ Italicæ*.

VERMICULAIRE BRÛLANTE, *Sedum*.

VERMICULAR, (dim. of *vermis*, 'a worm,') Vermiform. Also, wormlike; as the *vermicular* or wormlike motion of peristole.

VERMICULARIS, *Lumbricalis*, *Sedum*.

VERMICULI SPERMATICI, (pl. of *vermiculus*, dim. of *vermis*, 'a worm,') *Spermatozoa*.

VERMICULUM, *Vermiculum*. An elixir; a tincture, according to Ruland and Johnson.

VERMICULUS CAPILLARIS, 'hair worm,' *Draconculus*.

VERMIFORM, *Vermicular*, *Vermicularis*, *Vermiformis*, (F.) *Vermiculaire*, *Vermiforme*, (*vermis*, 'a worm,' and *forma*, 'form,') That which resembles a worm.

VERMIFORM PROCESSES, *Processus seu Protuberantia seu appendices vermiformes*, *Vermes*, *Monticuli cerebelli*, Worms, (F.) *Eminences vermiformes du cerebel*, are two medullary projections at the surface of the cerebellum. The superior vermiform process, *Vermis superior cerebelli*, is an elongated projection at the anterior and middle part of the superior surface of the cerebellum; and the inferior vermiform process, *Vermis inferior cerebelli*, *Pyramid of Malacarne*, is a somewhat larger projection, situate in the depression at the inferior surface of the same organ. The inferior vermiform process is usually described as consisting of three portions,—the pyramid, the uvula, and the nodule. The anterior portion of the superior vermiform process is elevated, and termed *Monticulus*.

VERMIFUGE, (*vermis*, 'a worm,' and *fugare*, 'to drive away,') Anthelmintic.

VERMIFUGE, MATHIEU'S. This consisted of two distinct electuaries; the one for killing, the other for expelling tapeworm. The former is

made of an ounce of *tin filings*, and six drachms of *ferri root*.

VERMIFUGUS, (Vermifuge,) Anthelminthic.

VERMILION, (from *ver*, 'a worm,' the colour being like that from the cochineal insect. It also means cochineal,) Hydrargyri sulphuretum rubrum.

VERMILLON DE PROVENCE, Antimonii sulphuretum rubrum.

VERMINATIO, (from *vermis*, 'a worm,') Helminthiasis, Vermination.

VERMINATION, Helminthiasis—*v.* Cutaneous, Malis.

VERMINEUX, Verminous.

VERMINOUS, *Vermineus*, (F.) *Vermineux*, (from *vermis*, 'a worm,') That which is caused by worms; as 'a *verminous affection*.'

VERMIS, (from *vertere*, 'to wind about,' [?]) see Vermiform processes—*v.* Bombycinus, Cornu ammonis—*v.* Cerebelli, Vermiform process—*v.* Cerebri, Fever, Hungary—*v.* Cucurbitinus, Tania solium—*v.* Digni, Paronychia.

VERMITIO, (from *vermis*, 'a worm,') Helminthiasis.

VERNIX CASEOSA, 'cheesy varnish,' (F.) *Vernis Caséux*. The fatty varnish, or deposit, often found on the fetus, which is secreted by the sebaceous follicles of the skin.

VERNIX SICCA, 'dry varnish,' Sandarac.

VERNONIA ANTHELMINTICA, (after William Vernon, an English botanical traveller in North America,) *Ascaricida anthelmintica*, *Calag'rah*, *Calag'ri*. A plant which is indigenous in India. *Ord. Compositæ. Sex. Syst. Syngenesia Polygamia superflua*. It has great celebrity in the East Indies, and is a bitter tonic; but the seeds are chiefly used as an anthelmintic in cases of ascarides and lumbrici in children.

VERNONIA NOVEBORACENSIS, *Iron weed*, *Flat top*; an indigenous plant, whose purple flowers appear in August and September, is reported to possess cathartic properties.

VÉROLE, (from *varius*, 'spotted,' [?]) Syphilis—*v.* Grande, Syphilis—*v.* Grosse, Syphilis—*v.* Petite, Variola—*v.* Petite volante, Varicella.

VÉROLÉ, Syphilitic.

VÉROLETTE, (dim. of *vérole*,) Varicella.

VÉROLIQUE, Syphilitic.

VERONICA or **VERONICA**, after St. Veronica, *V. officinalis* seu *Tournefortii* seu *mas*, *The'a Germanica*, *Betonica* *Pau'li*, *Chamædrys*, *Male Veronica*, *Fluel'tin*, *Offic'inal Speed'well*, *Eye'bright*, *Pau'ls Bet'ony*, (F.) *Véronique*, *Thé d'Europe*, *Herbe aux ladres*. *Ord. Scrophulariaceæ. Sex. Syst. Diandria Monogynia*. This plant was once used as a pectoral.

VERONICA AQUATICA, *V. beccabunga*.

VERONICA BECCABUNGA, *Beccabunga*, *Anagallis* seu *Veronica aquatica*, *La'ver Germanicum*, *Copa'a*, *Wa'ter Pin'pernel*, *Brook'time*, *Brook'time Speed'well*, *Wa'ter Speed'well*, *Neck'weed*, *Wa'ter Pur'slain*, (Sc.) *Wa'terpur'pie*, (F.) *Véronique cressonnée*. This plant was formerly used in a variety of diseases. It is, however, almost insipid and inodorous, and has been long banished from practice.

VERONICA MAS, *Veronica*—*v.* *Officinalis*, *Veronica*.

VERONICA PEREGRINA, *Neck'weed*, *Pur'slain*, *Speed'well*; indigenous; flowering from April to June, has been given internally in scrofulosis, and is applied as a wash to scrofulous tumours.

VERONICA PURPUREA, *Betonica officinalis*—*v.* *Tournefortii*, *Veronica*—*v.* *Virginica*, *Leptandra Virginica*.

VÉRONIQUE, *Veronica*—*v.* *Cressonæ*, *Veronica beccabunga*.

VERPA, *Penis*.

VERPUS, *Circumcisis*, *Medius digitus*.

VERRE, *Vitrum*—*v.* *d'Antimoine*, *Actimonii vitrum*.

VERRUCA, dim. *Verrucula*, *Ecphy'ma verrucosa*, *Phymatosis verrucosa*, an *Excrescence*, a *Wart*, (Prov.) *Wret*, (F.) *Verrue*, *Poireau*. A small, hard, mamillary tumour, which forms at the surface of the skin, and particularly on that of the hands and face. Warts are generally rugous at the surface, and broad at the base; their roots being implanted in the derma by whitish filaments, which are dense, semifibrous, and very numerous.

Warts may be destroyed by caustic, or removed by excision. Their vitality is, generally, not great.

VERRUCA CARNOSA, *Condyloma*—*v.* *Formicaria*, *Formica*.

VERRUCA MINOR, *Thymion*—*v.* *Palpebrarum*, see *Hydatid*—*v.* *Pensilis*, *Acrochordon*—*v.* *Rhagoidea*, *Thymion*.

VERRUCA NECROGENICA. A name proposed for productions of morbid epithelium, caused, in dissecting, by the acrid fluids of the dead body.

VERRUCARIA, (from *verruca*,) *Calendula officinalis*, *Heliotropium Europæum*.

VERRUCOSE, *Warty*.

VERRUCOUS, *Warty*.

VERRUCULA, (dim. of) *Verruca*.

VERRUE, *Verruca*.

VERRUQUEUX, *Warty*.

VERS, pl. of *Ver*, *Worms*.

VERSE CHARMS, see *Charm*.

VERSIO, (*vertere*, *versum*, 'to turn,') *Turning*—*v.* *Fœtus*, *Turning*—*v.* *Spontanea*, *Evolution*, *spontaneous*—*v.* *Uteri*, *Hysteroloxia*.

VERSION, *Turning*—*v.* *Cephalic*, see *Turning*—*v.* *Podalic*, see *Turning*.

VERT-DE-GRIS, ([F.] *vert*, 'green,' and *gris*, 'gray,') *Cupri subacetatis*.

VERTEBRA DENTATA, 'toothed vertebra,' *Axis*—*v.* *Dorsi* *Prima*, *Lophia*—*v.* *Lumborum* *Prima*, *Asphaltites*—*v.* *Magna*, *Sacrum*—*v.* *Proëminens*, see *Vertebra*.

VERTEBRÆ, pl. of *Vertebra*, *Spondyli*, *Spondyles*, *Whet'lebones*, (F.) *Vertèbres*, (from *vertere*, 'to turn,') This name has been given to the bones which by their union form the vertebral or spinal column. The vertebrae are short, thick, very angular bones, twenty-four in number, and situate one above another. The *body*—*centrum*, of Owen—of the vertebra is the middle and anterior portion. Above and below it is articulated, by means of a fibrocartilage, with the contiguous vertebrae. The *Proc'esses*, (F.) *Masses apophysaires*, of each vertebra are:—1. The *epi'nois proc'ess*, situate behind, and in the median line. 2. The two *trans'verse proc'esses*, standing off from each side. 3. The four *artic'ular or oblique proc'esses*—*zygapophyses*, of Owen—two *superior*, and two *inferior*—which serve to unite the vertebrae with each other. These transverse and articulating processes are made continuous with the lateral and posterior parts of the body of the vertebra by narrow bony portions or pedicles, on each of which are notches. These, by their union with similar notches in the contiguous vertebra, form the lateral foramina, (F.) *Trous de conjugaison*, through which the spinal nerves leave the vertebral canal. The different processes are united with each other, so as to form a kind of *ring* of the lateral and posterior parts of the vertebra; and the foramen within this,—that is, between the body and processes, *fora'men vertebra'le rachid'ium*,—concurs in forming the vertebral canal. The *arch* or *lam'ina* of a vertebra, *neurapoph'ysis*, is the part from which the several

processes project. The vertebræ have particular characters, according to the regions they occupy. They are divided into seven *cervical*, twelve *dorsal*, and five *lumbar*. The first cervical is called *atlas*; the second, *axis* or *vertebra densata*; and the seventh, *vertebra prom'inens*. The vertebræ are articulated with each other. Those of the back are joined, moreover, to the ribs by their articulating surfaces—*parapophyses*. The first or atlas is articulated with the occipital bone; and the last lumbar vertebra of the sacrum. The cervical vertebræ are developed by nine points of ossification; the dorsal and lumbar by eight.

VERTEBRAL, Vertebra'lis. That which relates or belongs to the vertebræ.

VERTEBRAL ARTERY, Posterior cer'ebral artery (Ch.), is the largest of the branches of the subclavian. It arises from the posterior part of that vessel; is lodged in a canal, (F.) *Petit canal vertébral*, formed by the foramina in the transverse processes; enters the cranium, and proceeds to anastomose with its fellow of the opposite side, and to form the *Bas'iliary artery*. Besides the branches sent to the neck, the vertebral artery gives off the *anterior* and *posterior spi'nal arteries*, and the *inferior cerebel'lous*, in the cavity of the cranium.

VERTEBRAL CANAL, see Vertebral column.

VERTEBRAL COLUMN, Rac'his, Rho'chis, No'tus, Acon'es'tis, Cari'na, C. hom'inis, Spi'na, S. dor'si seu vertebra'lis, Colum'na spi'na'lis seu vertebra'lis seu dor'si, Dor'sum, Ter'gum, Compa'ges Vertebra'rurum, Acan'tha, Sa'cra fistu'la, Hi'era Sy'rinx, Spine, Backbone, (Sc.) Écalsprent, (F.) Colonne vertébrale, Épine du dos, Échine. A kind of bony column, situate at the posterior and central part of the trunk; extending from the head to the sacrum; flexible in every direction, yet very solid, and hollowed by a canal called *vertebral*, which gives it lightness without diminishing its strength, and runs through its whole length. The vertebral column is formed by the superposition of the vertebræ. It represents a pyramid, whose base is below. Anteriorly, it is convex in the neck; concave in the back, and convex, again, in the loins. Behind, the curvatures are of course the reverse of these. There is also in the dorsal region a lateral curvature, the concavity of which is towards the left side. Considered as a whole, the vertebral column has an *anterior* or *prespi'nal* surface; a *posterior* or *spi'nal*; two *lateral*; a *base*; and an *apex* or top. In the nomenclature of Chaussier, the anterior surface, in the neck, is called *trache'lian*; in the back, *predor'sal*; and in the loins, *prelum'bar*. The dorsal surface has at its middle the spinous processes; and at its sides the *vertebral gut'ters*, (F.) *Gouttières vertébrales*, formed by the union of the *vertebral plates*, (F.) *Lames vertébrales*, which, by their junction, constitute the spinous processes. These are filled by the sacrospinalis muscle. The sides or lateral surfaces present the foramina resulting from the union of the *vertebral not'ches*, (F.) *Échancrures vertébrales*. These are called *vertebral foram'ina*, (F.) *Trous de conjugaison*, and give passage to the *vertebral nerves*. The base of the vertebral column is oblique, to unite with the sacrum, and forms with it an angle called *sacro-vertebral angle* or *prom'ontory*. The *apex* or top of the vertebral column is articulated with the occipital bone. Sometimes the term *vertebral column* includes the whole of the column from the occipital bone to the extremity of the coccyx. The *Vertebral* or *Sp'i'nal* or *Rachid'ian Canal*, *Sol'en, Cana'lis vertebra'lis seu medull'ie Spi'na'lis, Spe'cus pro medull'a Spi'na'li seu vertebra'lis*, exists through the whole of the vertebral column, whose curvatures it follows. It is continuous,

above, with the cranium; below, with the sacral canal, if we consider the vertebræ to terminate at the sacrum. It is lined by a prolongation of the meninges, forming the *The'ca vertebra'lis*, and lodges the spinal marrow.

The vertebral column, which unites lightness with solidity, serves to support the head and chest. It is the seat of all the movements of the trunk, the weight of which it transmits to the pelvis. It lodges and protects the spinal marrow, and gives passage to the vertebral nerves, and to numerous vessels.

VERTEBRAL DISEASE', Rachial'gia (of some), *Ca'ries of the Ver'tebræ, Spondylarthroc'acé, Spondyl'oc'acé, Kypho'sis inflammato'ria seu paraly'tica, Gibbos'itas cario'sa, Ca'ries vertebra'rurum, Spondylal'gia, Spondyl'oppy'osis, Tuberculo'sis Vertebra'rurum*. The spinal column is liable to distortion (see Hump), at times, without the vertebræ being much, if at all, diseased. At other times, a more formidable affection is induced; the vertebræ becoming carious, and the intervertebral cartilages ulcerated; followed by a more or less complete loss of power over the lower extremities. In such case, on examining the spine, one or more of the spinous processes is found to project beyond the rest. The disease was first well described by Mr. Percival Pott; and hence it has been called by the French, *Maladie de Pott, Malum seu Gib'bus Pott'ii, Pott's disease* or *curv'ature*. It usually occurs in scrofulous subjects, and the treatment consists in establishing a counterirritation by blisters, issues, or the moxa, opposite the transverse processes of the diseased vertebræ. The sore must be suffered to remain open, and the patient kept in the horizontal posture until the use of the limbs is recovered; employing tonics, the free admission of fresh air, &c., at the same time.

VERTEBRAL LIG'AMENTS, (F.) Ligaments vertébraux, Surtouts ligamenteux de la colonne vertébrale, are distinguished into *anterior* and *posterior*. They consist of long, ligamentous coverings; the *anterior* occupying the anterior part of the vertebral column; and the *posterior* seeming to arise from the occipito'axoid ligament, and seated behind the bodies of the vertebræ, from the axis to the sacrum.

VERTEBRAL NERVE, Trisplanchnic.

VERTEBRAL NERVES, Spi'nal Nerves, (F.) Nerfs rachidiens, spring successively from the spinal marrow, from its origin to its termination. These nerves—30 pairs in all—arise by two roots; one *anterior*, from the *anterior* or *anterolateral column* of the spinal marrow, and the other *posterior*, from the *posterior* or *posterolateral column*, and afterwards issue by the vertebral and sacral foramina. Each of these roots seems to have a distinct function; the *posterior* being destined to convey sensation to parts; the *anterior*, motion. On the posterior root, before its junction with the anterior, there is a ganglion. Of the 30 pairs, 7 are *cervical* nerves, 12 *dorsal*, 5 *lumbar*, and 6 *sacral*. See Nerves.

VERTEBRATED, Vertebra'tus, (F.) Vertébré. Provided with vertebræ.

VERTÈBRES, Vertebrae.

VERTEBRODY'MIA, (vertebra, and dy'u, 'to mingle,') see Somatody'mia.

VERTEBROÏLIAC LIGAMENT, Ileolumbar ligament.

VERTEX, gen. Ver'ticis, Cor'yphè, Corys, Acro'ria, Spon'dylus. The top or highest part or crown of the head, (F.) *Sommet; (Se.) Can'tel, Now, Pal'lat, Pal'ad.* See Sinciput.

VERTEX CORDIS, Muero Cordis—v. Cubiti Olecranon—v. Palpitans, Fontanella.

VERTIBULUM, (from vertere, 'to turn,') The rounded head of a bone; a joint; trepan,

VERTICULA, (from *vertere*, 'to turn,') Articulation.

VERTICULUM, Articulation.

VERTICULUS, Articulation.

VERTIGE, Vertigo.

VERTIG'INOUS, *Vertigino'sus*, (Old Eng. and Prov.) *Totty*, (F.) *Vertigineux*. Relating or pertaining to vertigo; *Gid'dy*, *Diz'zy*.

VERTIGO, gen. *Vertig'inis*, (from *vertere*, 'to turn,') *Di'nos*, *Ilin'gos*, *Di'nus* seu *Hallucina'tio* seu *Autal'gia verti'go*, *Circumgyra'tio*, *Gid'diness*, *Diz'ziness*, *Swim'ming of the head*, (Prov.) *Sieec*, *Swim'nying*, (F.) *Vertige*, *Etourdissement*. A state in which it seems that all objects are turning round, or that the individual himself is performing a movement of gyration. Two varieties have been designated;—*simple verti'go*, where the objects are seen; and *scotodyn'ia*, in which the sight is obscured, and the individual often falls. Vertigo is dependent upon the condition of the brain as affected by the circulation, and often announces an attack of apoplexy or epilepsy.

VERTIGO TENEBRICOZA, Scotodynia.

VERUGAS, (S.) (from *verruca*, 'a wart,') A peculiar disease, affecting certain localities of Peru, which manifests itself by sorethroat, pains in the bones, and other febrile symptoms. In the course of a few days, an eruption of red pimples or boils appears. These sometimes increase in magnitude till they become nearly as large as an egg, and discharge blood copiously. It is ascribed to the water of certain springs in the valleys, and especially in that of Surea, called *Aqua de Veruga*.

VERU MONTANUM, 'a mountain spit,' Gallinaginis caput.

VERVAIN, *Verbena officinalis*—v. Blue, *Verbena hastata*.

VERVEINE, *Verbena officinalis*.

VERVE'NIA, *Vesv'nix*, (from *ve*, negative, and *venus*, 'sound,') 'madness.' Derangement of the intellectual and moral faculties, without coma or fever. Many nosologists have used this as a generic term, under which they have included the different species of mental alienation, hypochondriasis, somnambulism, and some even hydrophobia. See Nosology.

VESANIA MANIA, Mania.

VESANIE, Insanity, Vesania.

VESICA, 'a bladder,' Urinary bladder—v. Biliaria, Gallbladder—v. Cordis, see Pericardium—v. Fellea, Gallbladder—v. Intestinalis, Vesicula umbilicalis—v. Natatoria, Airbladder—v. Prostatica, see Ejaculatory ducts—v. Saccata, Cystostomochoria—v. Urinaria, Urinary bladder.

VESICÆ IRRITABILITAS, Cysterethismus—v. Unguinosæ tendinum, Bursæ mucosæ.

VESICAL, *Vesical'is*, (F.) *Vésical*, (from *vesica*.) That which belongs or relates to the bladder.

VESICAL ARTERIES vary in number and origin. The umbilical artery always sends off two or three: the middle hemorrhoidal, internal pudic, and obturator furnish others, which ramify and anastomose in the parietes of the bladder; the hypogastric furnishes one which is somewhat larger, the *Vesicoprostatic* (Ch.), and proceeds to the *bas-fond* of the bladder, sending numerous ramifications to it as well as to the surrounding parts.

VESICAL GANGLIA, see Spermatie Ganglion—v. Ligaments, see Urinary bladder—v. Oxyd, Oxyd, cystic—v. Triangle, see Urinary bladder.

VESICAL VEINS, much more numerous than the arteries, open into the hypogastric plexus.

VESICANT, (from *vesica*.) Blister.

VESICARIUM, Ovarium.

VESICA'TION, *Vesica'tio*, (from *vesica*.) The action of a vesicant. The formation of blisters.

VÉSICATION VOLANTE (F.), 'a flying vesication.' A temporary vesication produced by art.

VÉSICATOIRE, Blister—v. *Magistral*, Blister, *magistral*—v. *Volant*, see Blister.

VESICATORIUM, Blister.

VESICLE, (dim. of *vesica*.) Vesicula—v. Allantoid, Allantoids—v. Blastodermic, see Molecule—v. Cowpox, see Vaccination.

VESICLE OF BAER, *Vesic'ula Buerii*, (F.) *Vésicule de Baer*. The ovule in the follicle of De Graaf; which was well described by Baer in 1827.

VESICLE OF DEVELOPMENT, *V. of Evolution*. A vesicle or cyst concerned in histogeny.

VESICLE OF EVOLUTION, *V. of Development*—v. Germinal, see Molecule—v. Purkinjean, see Molecule—v. Umbilical, Vesicula umbilicalis—v. Valentinian, see Molecule—v. Wagner's, see Molecule.

VESICLES, ASCHERSO'NIAN. When fluid fat and fluid albumen are shaken together, the resulting fat globules become surrounded by an albuminous layer. The vesicles, thus formed, are called after Ascherson, their describer.

VESICLES OF DE GRAAF, Folliculi Graafiani—v. Lung, see Pulmo—v. Malpighian, Cellulæ, bronchic, see Pulmo—v. Seminal, Vesiculæ seminales.

VESICOCE'LE, (*vesica*, 'a bladder, and *κηλη*, 'a rupture,') Cystocoele.

VESICOPROSTATIC, see Vesical.

VESICOÛ'RACHAL, *Vesi'coû'racha'lis*. Relating or appertaining to the bladder and urachus, —as a 'vesicoû'rachal fistula.'

VESICOÛ'TERINE, *Vesi'coû'teri'nus*. Appertaining, or relating, to the bladder and uterus.

VESICOÛTERINE FOLDS, see Uterus.

VESICOVAG'INAL, *Vesi'covagina'lis*, *Vag'ino-ves'ical*, *U'rino-vag'inal*. Relating to the bladder and the vagina; as *vesicovaginal fistula*.

VESIC'ULA, *Cyst'inæ*, *Ves'icle*, *Blis'ter*, (F.) *Vésicule*, (dim. of *vesica*.) A small bladder or cyst.

VESICULA, *Ves'icle*, *Ec'phlyis*, *Cyst'inæ*, *Pomphus*, in pathology, means a small orbicular elevation of the cuticle containing lymph, which is sometimes clear and colourless, but often opaque and whitish or pearlecoloured. It is succeeded, either by a scurf, or by a laminated scab.

VESICULA ENIGMATOSA, Vesicula umbilicalis—v. Alba, Vesicula umbilicalis—v. Bilis, Gall bladder—v. Blastodermatica, see Molecule—v. Blastodermica, see Molecule—v. Chyli, Receptaculum chyli—v. Erythroidea, Erythroid vesicle—v. Fellis, Gallbladder—v. Gangranescens, see Anthrax—v. Germinativa, see Molecule—v. Omphalomesenterica, Vesicula umbilicalis—v. Ovalis, Vesicula umbilicalis—v. Prolifera, see Molecule—v. Prostatica, see Ejaculatory ducts.

VESICULA UMBILICA' LIS, *V. al'ba* seu *om'phalomesenterica* seu *enigmatosa* seu *ovalis*, *Ves'ica intestinal'is*, *Vitell'iculus*, *Umbil'ic'ul* or *Intestinal Ves'icle*, *Vit'elline Pouch*, *Vitell'icle*, (F.) *Vésicule ombilicale*. A small, pyriform, round, or spheroidal sac, which, about the fifteenth or twentieth day after fecundation, is of the size of a common pea, and probably acquires its greatest dimensions in the course of the third or fourth week. After the seventh week, it becomes shrivelled and disappears insensibly. It seems to be situate between the chorion and amnion, and is commonly adherent either to the outer surface of the amnion, or to the inner surface of the chorion, but at times is loose between them. The *vitelline ped'icle*, which attaches the vesicle to the embryo, is continuous with the intestinal canal; and, up to twenty or thirty days of embryonic life, is hollow,—*Ductus om'phalomesenter'icus* seu *vitello-intestina'lis* seu *vitell'arius*, *Vit'elline Duct*. Its arteries are the omphalomesenteric. The *Vit'*

elline fluid, contained in the vesicle—in the *Vitelline* or *Yolk membrane*—has been compared, from analogy, to the *vitellus* or *yolk* of birds; and, like it, is an oleaginous and albuminous substance, which is inservient to the nutrition of the embryo.

VESICULÆ, pl. of *vesicula*, constitute the 6th order of Bateman's classification of *Cutaneous Diseases*.

VESICULÆ, see Villous membranes, *Vesiculæ aereæ*, Cellules, bronchie—v. *Bronchiæ*, Cellules, bronchie—v. *Divæ Barbaræ*, Variola confluens—v. *Gingivarum*, Aphthæ—v. *Graafianæ*, Folliculi Graafiani—v. *Malpighianæ*, Cellules, bronchie—v. *Pulmonales*, Cellules, bronchie—v. *Rotundæ cervicis uteri*, Nabothi glandulæ—v. *Sanguinis*, Globules of the blood.

VESICULÆ SEMINALES, *V. Semina'rix* seu *spermatice*, *Cap'culæ seminales*, *Conceptac'ula seminaria*, *Gonocyst'ides*, *Spermatozyst'ides*, *Seminal vesicles*, (F.) *Vésicules séminales*, *Gardouches*, *Greniers*, are two small membranous sacs, which serve as reservoirs for the sperm, and probably secrete a fluid to be mixed with it. They are situate beneath the bladder, and above the rectum, behind the prostate and to the outer side of the vasa deferentia. They are of an irregular conoidal shape; flattened from above to below; embossed on their surface, and of a grayish tint. Their *posterior extremity* or *fundus* terminates by a round cul-de-sac. Their *anterior extremity* or *neck* is narrow and elongated, and is continuous with the excretory duct of the vesicula. This duct joins the vas deferens, and forms with it the ejaculatory duct. The interior of the vesiculæ exhibits several cells, which communicate with each other, and are lined by a very thin, mucous membrane.

VESICULÆ SEMINALES MULIERUM, Nabothi glandulæ—v. *Spermatice*, *Vesiculæ seminales*.

VESICULAR, *Vesicularis*, (F.) *Vésiculaire*. Of or belonging to a vesicle:—full of or containing small vesicles or cells. The cortical or gray matter of the brain is sometimes termed *vesicular*. See *Cortex cerebri* and *Neurine*.

VESICULAR FEVER, *Pemphigus*.

VESICULE, *Vesicula*—v. *Biliare*, Gallbladder—v. *du Fiel*, Gallbladder—v. *Germinative*, see *Molecule*—v. *Ombilicale*, *Vesicula umbilicalis*—v. *Ovulifère*, see *Folliculi Graafiani*.

VESICULÆ DE DE GRAAF, Folliculi Graafiani—v. *de Naboth*, Nabothi glandulæ—v. *de Sainte Barbe*, Variola confluens—v. *Séminales*, *Vesiculæ seminales*.

VESICULOUS, *Vesicul'ous*, (F.) *Vésiculeux*, (from *vesica*.) Shaped like, or characterized by, vesicles.

VESICULOTYMPANITIC, *Vesiculotympaniticus*. An epithet for a sound or resonance heard on percussion, which consists of the vesicular and tympanic sounds combined. It occurs, according to Dr. Flint, when the parenchyma of the lungs is incompletely solidified.

VESPA, Wasp.

VESSEL, *Vas*, *Va'sum*, dim. *Vas'culum*; *Ang'ion*, *Ang'us*, *Conceptac'ulum* ([L.] *vas*, *vasculum*.) (F.) *Vaisseau*. A canal, more or less elastic, formed by the superposition of membranes, and distinguished, according to its uses and general arrangement, into *Artery*, *Vein*, and *Lymphatic*. See, also, *Vascular*.

VESSELOUP, 'wolf fizzle,' *Lycoperdon*.

VESSIE, (from *vesica*, 'a bladder,') Urinary bladder—v. *Col de la*, Collum vesicæ—v. *Natoire*, Airbladder.

VESSIE A COLONNES, see Urinary bladder.

VESTIBULAR, *Vestibularis*, (F.) *Vestibulaire*. Relating or appertaining to a vestibule.

VESTIBULAR NERVE, see Auditory nerve.

VESTIBULE, *Vestibulum*, 'from *vestio*, 'to

clothe or cover.')

The porch or entrance to a house.
VESTIBULE, *Vestibulum*, *V. Vaginæ*, *Prona'us*, *Atrium Vaginæ*, *Am'bitus genit'alis mul'iebris*, is a smooth cavity between the perinæum and nymphæ in the female, which leads to two passages—to the urethra above, and to the vagina below.

VESTIBULE OF THE EAR, *Vestibulum*, *V. labyrinthi*, (F.) *Vestibule*. A cavity of an irregular shape, which forms part of the labyrinth or internal ear. The vestibule is situate on the inner side of the tympanum; on the outer side of the meatus auditorius internus; before the semicircular canals, and behind the cochlea. It is lined by a particular membrane, and contains the liquor of Cotunnus and branches of the auditory nerve. There is also another membrane, constituting the *membranous vestibule*, but it is not an exact imitation of the osseous cavity. It is composed of two distinct sacs, which open into each other—the *sacculus vestibuli*, and the *sacculus*. There are three contiguous cavities in the vestibule; one of which, the *semi'oval*, is situate above; another, the *hemispher'ical*, below; and the third, or *sul'ciformal*, which is the orifice of the aquæductus vestibuli, behind.

VESTIBULUM LABYRINTHI, Vestibule of the ear—v. *Vaginæ*, Vestibule.

VESTIGE (F.), *Vestig'ium*, 'a vestige,' 'a trace.' A kind of fracture of flat bones, consisting of a simple incision having the mark of the instrument that caused it. The term is also applied to a part of the body, which may have no evident use in man, and yet may be greatly employed in animals. For example, the muscles of the external ear in man are almost devoid of use, or are *vestiges* of what they are in certain animals.

VESTIGIUM, Sole, *Vestige*—v. *Foraminis ovalis*, *Ovalis fossa*—v. *Pedis*, *Metatarsus*.

VESTIMENTUM, Vestitus.

VESTITUS, *Vestimentum*, *Esthe'ma*, *Esthe'sis*, *Res vestia'ria*, *Vest'ment*, *Clo'thing*, *Dress*, *Ves'ture*, (F.) *Vêtement*, (from *vestire*, *vestitum*, 'to clothe,') Any substance used for the purpose of covering nakedness, or preserving from atmospheric temperature by being worn on the body. The best clothing in winter is composed of articles that are bad conductors of caloric, and do not admit of ready evaporation of the matter of perspiration. In this respect, woollen vestments deserve the first place, and cotton the second. Colour has likewise to be regarded—the darkest hues transmitting solar heat most readily through them. It has been found that the same rule applies to odours—the darker colours retaining a greater amount of odorous particles than the lighter. Hence the necessity of white walls, white bedclothes, &c., in the wards of hospitals; for it is probable that the like law applies to contagious miasms.

VESTMENT, Vestitus.

VESTURE, Vestitus.

VETA, Puna.

VÊTEMENT, ([L.] *Vestimentum*.) Vestitus.

VETERANA, Rubeola.

VETERINARIUS, see Hippiater and Veterinary.

VETERINARY, *Veterina'rius*, (F.) *Vété'rinaire*, (from *veterinus*, 'that which bears burdens; itself, perhaps, from *vehiterinus*, from *vehere*, 'to carry,') That which appertains to beasts of burden: hence, *veterinary surgeon*, (see *Farrier*), *veterinary college*, *veterinary medicine*, &c.

VETERINARY ANATOMY, see Anatomy.

VETERINARY ART, *Ars seu Medi'cina veterina'ria*, *Zoô'i'asis*, *Zoiatri'a*, *Theriat'ria*, *M'lo'medici'na*, *Zoôtherapi'a*, *Zoôtherapeu'ticæ*, *Ars zoiat'*

vica, Farriery. The art of treating the diseases, &c., of cattle.

VETERNOSITAS, (from *veternus*, 'aged;' because old persons are subject to somnolency,) *Coma vigil*.

VETERNUS, *Coma vigil*.

VÉTIVER, *Andropogon muricatus*.

VETIVERIA ODORATA, *Andropogon muricatus*.

VETTIVERT, *Andropogon muricatus*.

VETTONICA, (after the *Vettones* of Spain,) *Betonica officinalis*.

VETULA, *Sacrum*.

VETYVER, *Andropogon muricatus*.

VEXILLA (pl. of *vexillum*) **TUBARUM FALLOPII**, see Tube, Fallopian.

VEXILLUM, 'a flag,' *Drapeau*.

VIA, Way—*v. Curandi*, Ratio medendi—*v. Stomachi et ventris*, Œsophagus.

VIABILITY, see *Viable*.

VIABLE, *Viabilis*, (from *vita*, [F.] *vie*, 'life,' and *habilis*, 'able,') A word introduced with great advantage from the French into our language. It is an epithet applied to a fœtus whose organs are properly formed, and so developed as to permit its continued existence. Every fœtus is not *viable*. Anencephali have lived ten or twelve days, yet they are not *viable*. The older the fœtus, the more *viable* is it. It is an idle prejudice that a child, born at the end of eight months, is less likely to live than one born at the end of seven. The signs of *viability* are drawn from the weight, length, external conformation, &c., of the fœtus. It may be laid down as a general rule, that no fœtus born before the end of the seventh month is *viable*.

VIE (pl. of *via*) **CHYLIFERÆ**, Chyliferous vessels.

VIE CLANDESTINÆ, *Clandestine Pas'sages*. Supposititious secret ways or passages or 'royal roads,' to account, for example, for the rapid transmission of substances taken into the stomach to the kidneys or bladder.

VIE EXTRAORDINARIÆ LACTIS, Galactoplasia—*v. Lacrymales*, Lachrymal passages—*v. Primæ*, Ways, digestive—*v. Secundæ*, Ways, second.

VIAL, *Phiala*.

VIANDES FAISANDÈES, ([L.] *vivenda*, from *vivere*, 'to live,') see *Hyposaprus*.

VIBICES, (pl. of *vibex*, 'the mark of a stripe,') *Mol'opes*, *Stri'æ*, (F.) *Vergetures*. Large purple spots, like the marks produced by the strokes of a whip, which appear under the skin in certain malignant fevers. They are unfavourable indications, as they denote great prostration.

VIBRAMEN, (*vibrare*, 'to quiver,') Oscillation.

VIBRANS, *Vibratory*.

VIBRATILE, (from *vibrare*, *vibratum*, 'to quiver,') *Vibratory*.

VIBRATILITY, *Vibratilitas*, (F.) *Vibratilité*. Capability of being thrown into vibration.

VIBRATIO, Oscillation, *Vibration*—*v. Cardiacæ*, Palpitation—*v. Cordis*, Palpitation.

VIBRATION, *Vibratio*—*v. Pectoral*, see *Frenitus*—*v. Tactile*, see *Frenitus*—*v. Tussivo*, see *Tussive*.

VIBRATIONS OF SOUND, see *Sound*.

VIBRATOR, *Deferens vas*.

VIBRATORY, *Vibrans*, *Vibrative*, (F.) *Vibratile*, *Vibratoire*. An epithet applied to the pain that accompanies some neuralgic, which seems to the patient to vibrate among the nerves.

VIBRATORY MOTION, see *Cilia*.

VIBRATUS, *Oscillation*.

VIBRIO, gen. *Vibrio'nis*, (from *vibrare*, 'to quiver,') A genus of infusory animalcules, of which different species are met with,—for example, *Vibrio prolifer* in almost all putrescent fluids containing protein, and in the pus of chan-

cre; in which last the *V. line'ola* has likewise been found.

VIBRIS'Æ, (from *vibrare*, 'to quiver,') The hairs which grow at the entrance of the nostrils and of other apertures. In the feline tribe, the whiskers.

VIBURNUM ACERIFOLIUM, *Ma'pleaved*, *Ar'rowwood*, (from *vire*, 'to bind;' several species having long flexible twigs.) *Dockmackie*; *Ord.* *Caprifoliaceæ*. A plant used by the Indians, and afterwards by the Dutch, as an external application to tumours.

VIBURNUM CASSINOIDES, *Ilex Paraguensis*.

VIBURNUM DENTA'TUM, *Ar'row Wood*, *Meal'y Tree*, *Tily* of the Indians. *Ord.* *Caprifoliaceæ*. A high indigenous shrub, of the same family as *V. lentago*; the bark of which has been used by the Indians as a diuretic.

VIBURNUM EDU'LE flourishes in the Hudson's Bay territory. The fruit has an orange colour, and is less acid, more fleshy, and more agreeable to the taste than the cranberry.

VIBURNUM LÆVIGATUM, *Ilex Paraguensis*.

VIBURNUM LENTA'GO, *Sweet Vibur'num*, *Nann'y-berry*. A handsome indigenous tree, which flowers in May and June. It has been used as an antiperiodic.

VIBURNUM PRUNIFOLIUM, *Black Haw*, *Sloe-leaved Viburnum*, *Sloe*. Indigenous: flowering in May. The leaves have been used as tea.

VIBURNUM, SLOWLEAVED, *V. prunifolium*—*v. Sweet*, *V. lentago*.

VIC-LE-COMTE, MIN'ERAL WATERS OF. *Vie-le-Compte* is a small village in the department of Puy de Dôme, three leagues from Issoire; where there is an acidulous chalybeate, which is aperient.

VICA'RIOUS, *Vica'rius*, (from *vici*, 'change,') Taking the place of another:—as 'vicarious secretion,' *Secre'tio vica'ria*; a secretion which takes place in one part instead of another.

VICAR'S BRAY, MIN'ERAL WATERS OF. This spring is situate in Perthshire, Scotland, about two miles from Dollor, and near to Blarngone. The water is one of the strongest chalybeates known.

VICE, (*vitium*, 'a defect,' 'a fault,') *Cacosis*—*v. Secret*, *Masturbation*.

VICE DE CONFORMATION. Faulty formation of a part of the body.

VICHY, MIN'ERAL WATERS OF. Carbonated alkaline chalybeates, the sources of which are at Vichy, six leagues from Moulins, in the department of Allier. The springs are seven in number. Almost all are thermal; give off bubbles of carbonic acid gas, and have a slightly sulphureous odour. They deposit carbonates of lime and magnesia, and a little oxide of iron; and afford, on analysis, carbonates of lime, soda, iron, and magnesia, carbonic acid, sulphate of soda, and chloride of sodium. They are employed as tonics, particularly in chronic affections of the abdominal viscera, in old catarrhs, diseases of the joints, &c. The season is from the 15th of May to the 15th of September.

VICHY WATER, *FACTU'IOUS*, *A'qua Vicien'sis*, (F.) *Eau de Vichy*, is made of simple *Acidulous water*, impregnated with twice its bulk of *Carbonic acid*, $\frac{3}{4}$ xxss; *Subcarbonate of soda*, gr. xxxij; *Sulphate of soda*, gr. xvj; *Chloride of sodium*, gr. iv; *Subcarbonate of magnesia*, gr. ss; *Chloride of iron*, gr. $\frac{1}{2}$. *Mix.* *Ph. P.*

A **VICHY WATER POWDER**, (F.) *Poudre pour Eau de Vichy*, is made as follows:—*Sodæ bicarb.* 5 gram.; *Sodii chlorid.* 20 centigram.; *Sodæ sulph.* 50 centigram.; *Magnes. sulph.* 15 centigram.; *Ferri sulph.* 1 centigram.; *Aquæ*, 625 gram. *Mix.* To render it gaseous, add *Acid. citric*, 3 gram.

VICIA ERVILIA, *Ervum*.

VIC'IA FA'BA, Fa'ba, *F. major* seu *vulgaris*, *Orabus* fa'ba, *Com'mon Gar'debenn*, (F.) Fève ou Fêve des marais. *Ord. Leguminosæ. Ser. Syst.* Diodaphnia Decandria. A native of Egypt, of which there are many varieties. Beans are not easy of digestion. They are proverbially flatulent, especially when old. When young, they are more digestible and wholesome. Bean meal was one of the four resolvent meals of the ancients.

VICINITRACTUS, (vicinus, 'near,' and *trahere*, *tractum*, 'to draw,') Erysipelas.

VICINITRAHA, Erysipelas.

VICIS'SITUDE, *Vicissitudo*, (from *vicis*, 'change,') Constant change. The vicissitudes of the atmosphere, as regards its barometric, thermometric, and hygrometric condition, have to be guarded against by the delicate; and yet, by the new impressions they produce, they are probably a cause of the greater mental and corporeal activity of the inhabitants of the temperate regions of the globe.

VICTORIALIS FEMINEA, *Gladiolus vulgaris*—v. *Lunga*, *Allium victorale*—v. *Rotunda*, *Gladiolus vulgaris*.

VICTUS RATIO, 'order of victuals,' (from *vivere*, *victum*, 'to live,') Dietary.

VIDANGES, (from *vider*, 'to empty,') Lochia.

VIDIAN CANAL, (after *Vidus Vidius*), Pterygoid canal—v. *Nerve*, Pterygoid nerve.

VIE, (from *vita*), Life—v. *Durée de la*, see *Longevity*—v. *Moyenne*, see *Longevity*.

VEILLEESSE, (from [L.] *vetulus*, 'an old person,') Senectus.

VIELLOTE, *Colchicum autumnale*.

VIENNA PASTE AND POWDER, see *Powder*, *Vienna*.

VERGE, ([L.] *virgo*), Virgin.

VIF ARGENT, 'quicksilver,' *Hydrargyrum*.

VIGILANCE, *Insomnia*.

VIGILANTIA, (from *vigil*, 'on the watch,') *Egrogorsis*.

VIGILATIO, *Egrogorsis*.

VIGILIA, *Egrogorsis*.

VIGILÆ, *Egrogorsis*—v. *Nimix*, *Coma vigil*.

VIGILIUM, *Egrogorsis*.

VIGNE, ([L.] *vineæ*), *Vitis vinifera*—v. *Blanche*, *Bryonia alba*—v. *de Judée*, *Solanum dulcamara*—v. *du Nord*, *Humulus lupulus*—v. *Vierge*, *Bryonia alba*.

VIGNETTE, (dim. of *vigne*), *Spiræa ulmaria*.

VIGNOLE, ([L.] *vineæ*, 'vine,') *Mercurialis*.

VIGOR, 'activity, force,' *Acme*.

VIL'LA FRANCA, **CLIMATE OF**. This little town is situate immediately to the eastward of Montalbano, which separates the bay of Villa Franca from that of Nice. It is on the southern declivity of a steep and lofty range of mountains; and is more effectually sheltered from the north and northwest winds than Nice, with the climate of which that of Villa Franca closely corresponds.

VILLI, (pl. of *villus*, 'a tuft of hair,') see *Villous membranes*.

VILLOSITÉS, see *Villous membranes*.

VILLOUS, *Villosus*, (F.) *Villeux*, (from *villus*, 'a tuft of hair,') Furnished with villi.

VILLOUS MEMBRANES OR COATS, *Velamen'ta Bombyc'ina*, (F.) *Membranes villenses*, *Membranes velontés*, are such as are covered with fine delicate prolongations, *Papillæ* or *Vil'li*, *Floc'ci*, *Floc'culi*, *Projections*, *Cylin'dri*, *Tu'bi*, *Membrana'cei*, (F.) *Villosités*, constituted of blood-vessels, nerves, and absorbents; or such as are covered with fine down, so as to cause them to feel soft and velvety. The term is, however, chiefly restricted to the former. Chaussier gives the name *sim'ple villous membranes* to the serous membranes; and that of *com'plicated or follicular villous membranes* to the mucous membranes.

VIN, ([L.] *vinum*), Wine—v. *de Bordeaux*, *Claret*—v. *de Poulet*, see *Poulet*—v. *de Quinquina composé*, *Vinum de kinâ kinâ compositum*—v. *Rouge d'Oporto*, *Vinum Portugallicum*—v. *Sec*, *Sack*.

VINAIGRE, (*vin*, 'wine,' and *aigre*, 'sharp or sour,') *Acetum*, see *Acetium acidum*—v. *Aromaticum*, *Acetum aromaticum*—v. *de Bois*, *Pyroligneous acid*—v. *Distillé*, *Acetium acidum dilutum*—v. *de Quatre voleurs*, *Acetum aromaticum*—v. *Scillitique*, *Acetum scillæ*.

VINAIGRES MÉDICINAUX, *Acetia*.

VIN'CA MINOR, *V. pervin'ca*, (from *vincio*, 'I bind,' because of its usefulness in making bands [?]), *Clem'atis daphnoï'des major*, *Les'er per'icinkle*, (F.) *Pervenche*, *Violette des Sorciers*. It has bitter and astringent properties, and has been used in gargles and diseases of the chest.

VINCETOXICUM, (*vincere*, 'to overcome,' and *toxicum*, 'poison,') *Asclepias vincetoxicum*.

VINCTURA, (*vincire*, *vinctum*, 'to bind,') *Fascia*.

VIN'CU'LA (pl. of *vinculum*) **ACCESSORIA**. On the theca of the fingers several small tendinous fasciculi are seen, which pass between the phalanges and the edges of the tendons, and are so called.

VINCULUM, (*vincire*, *vinctum*, 'to bind,') Bandage, *Fascia*, *Lien*—v. *Caninum*, *Frænum penis*—v. *Linguae*, *Frænum linguae*—v. *Præputii*, *Frænum penis*—v. *Umbilicale*, *Funiculus umbilicalis*.

VINE, ([L.] *vineæ*), *Vitis vinifera*—v. *Grape*, *Vitis vinifera*—v. *Poison*, *Rhus radicans*—v. *Wild*, *Paraira*.

VIN'EGAR, (F.) *Vinaigre*, (itself from *vin*, [L.] *vinum*, 'wine,' and *aigre*, [L.] *acris*, 'sharp or sour,') *Acetum*—v. *Aromatic spirit of*, see *Acetium acidum*.

VINEGAR OF CANTHARIDES, *Acetum cantharidis*.

VINEGAR, COMMON, *Acetum*—v. *Distilled*, see *Acetium acidum*—v. *Estragon*, see *Artemisia dracunculæ*—v. *Marseilles*, *Acetum aromaticum*.

VINEGAR OF MEADOW SAFFRON, *Acetum colchici*—v. of *Opium*, *Guttæ nigre*.

VINEGAR, RADICAL, *Acetium acidum*.

VINEGAR OF SQUILLS, *Acetum scillæ*—v. of the *Four Thieves*, *Acetum aromaticum*.

VINEGAR, THIEVES', (so called, because, under its use, four thieves—it was said—plundered, with impunity, the bodies of those dead of the plague in London [?]), *Acetum aromaticum*—v. *Whcy*, see *Acetum*.

VINEGAR OF WOOD, see *Acetium acidum*.

VINETTE, *Rumex acetosa*.

VINETTIER, *Oxyacantha Galeni*.

VINEUX, *Vinous*.

VINOLENTA, (from *vinum*), see *Temulentia*.

VINOLENTIA, *Temulentia*.

VINOSUS, *Vinous*.

VINOUS, *Vinosus*, *Æno's*, *Æno'sides*, (F.) *Vineux*, (from *vinum*), Relating to wine. Having the character of, or resembling, wine. Made of wine.

VINUM, (vinox), Wine, Sherry wine. See *Wine*. A medicinal wine, (F.) *Vin médicinal*, *Onolè*, *Onolature*.

VINUM ABSINTHITES, *Wine of Worm'wood*. (Fol. *absinth. essicc.*, *Absinth. Pontic.*, 3ʒj, *Vini adhi Oiv.* Macerate for 24 hours, and strain. *Ph. P.*) Stimulant, tonic, sebrifuge, vermifuge.

VINUM ADESTUM, *Brandy*—v. *Album*, *Sherry wine* (Ph. U. S. 1851), see *Wine*—v. *Album Hispanicum*, see *Wine*—v. *Album*, *Montanum*, see *Wine*.

VINUM ALOËS, *V. aloës Socotori'næ* seu *aloët'icum* seu *aro'moalœot'icum*, *Tinctu'ra hi'era*, *Tinctu'ra sa'era*, *Wine of Aloës*. (Aloës in pulv. ʒj, *Cardam. cont.*, *Zingib. cont.*, 3ʒj; *Vini alb. Oj.* Macerate for 14 days, and filter. *Ph. U. S.*) Pur-

gative and stomachic, according to the dose. Dose, f 3j to f 5ij.

VINUM AMARUM, V. Gentianæ comp.—v. Amarum scillicetum, &c., v. diureticum amarum—v. Amarum cum spiritu vini, v. gentianæ compositum—v. Antimoniale, v. antimonii tartarizati—v. Antimoniatum Huxhami, v. antimonii tartarizati—v. Antimonii potassiotartratis, v. antimonii tartarizati.

VINUM ANTIMONII, V. antimonii tartarizati seu tartaricis antimonii seu antimonii potassiotartratis seu benedictum seu stibiatum seu stibiatum Huxhami seu antimoniatum seu antimoniatum Huxhami seu antimoniatum seu emeticum, Exsultia antimonii seu stibii, A'qua benedicta Rulandi, Solution of tartarized antimony, Antimoniai wine. (Ant. et Potass. tart. gr. xxxij, Aq. destillat. sufficit. f 3j). Vini albi, q. s. Dissolve the tartarized antimony in the distilled water, and while hot add the wine to make a pint. Ph. U. S. 1863.) Each fluidounce contains two grains of the tartar.

The ordinary Antimonial wine was formerly made with tincture of antimony, 3j, Sherry, Oiss.

Ward's Red Drop consists of a strong vinous solution of tartarized antimony.

VINUM ANTISCORBUTICUM, Antiscorbutic wine. (Rad. amomacis recent, 3j, Fol. cochlearis recent, Nasturtii aquaticæ, Trifol. aquaticæ, Semin. sinap., 3ss, Ammonia muriat, 3ij, Vin. albi Oij. Macerate for 36 hours; filter, and add Tincturæ cochlear. 3ss, Ph. P.) Stimulant and tonic. Dose, f 3ss to f 3ij.

VINUM AROMATICUM, Aromatic wine. (Species aromaticæ, 3iv, Vin. rubr. Oij. Macerate for 6 days and filter. Ph. P.) Stimulant and tonic. Used, externally, in fomentations, embrocations, &c.

VINUM AROMALOECHICUM, V. aloes—v. Benedictum, V. antimonii tartarizati.

VINUM CAMPA'NUM, V. Campanaceum. Champagne wine. A well-known sparkling wine of France. See Wine.

VINUM CANARINUM, see Wine—v. Chalybeatum, V. ferri—v. Cinchonæ compositum, V. de kinâ kinâ comp.

VINUM COL'CHICI, V. Col'chici Radicis, Wine of Colchicum, Wiant's Eau d'Huison. (Colchic. rad. bene contus. 3ij, Vini albi, Oij. Macerate for 24 days, with occasional agitation, and then express strongly, and filter. It may also be prepared by displacement. Ph. U. S.) Diuretic, sedative, and purgative. Chiefly used in gout. Dose, gtt. 40 to f 3j.

Reynold's Specific for gout and rheumatism consisted of the fresh bulb of Colchicum, 3vij, Sherry wine, f 3xvj. The colchicum was macerated for eight or ten days in the wine, at a gentle heat; coloured with Syrup of poppies, and flavoured with Rum. Reynolds is said to have killed himself by taking an over-dose of it.

VINUM COLCHICI RADICIS, V. colchici.

VINUM COLCHICI SEMINIS, Wine of Colchicum seed. (Colchic. sem. cont. 3iv; Vini albi, Oij. Macerate for 14 days, with occasional agitation, then express and filter. Ph. U. S.) Dose, f 3j to f 3ij.

VINUM CREMATUM, Brandy.

VINUM DIURETICUM AMARUM, V. amarum scillicetum compositum in Charitatis Nosocomio usurpatum, Bitter diuretic wine. (Cort. cinchon., C. Winteran., Limon., 3ss, 3ij, Rhi toxicendendr., Scilla Cæcamin., Angelica, 3ss, Fol. absinth., Melissa, 3ss, 3ij, Baccar. junip., Macia, 3ss, 3ss, Vin. albi, Ovij. Macerate and filter. Ph. P.) Stimulant, tonic, and diuretic. Dose, f 3ss to f 3ss.

VINUM EMETICUM, V. antimonii tartarizati.

VINUM ER'GOTE, Wine of Er'got. (Ergots cont., 3iv; Vini albi, Oij. Macerate for 14 days, with occasional agitation; then express and filter. It may be prepared, also, by displacement. Ph. U.

S.) Dose, ʒxx to f 3j to produce the peculiar effects of ergot.

VINUM FERRATUM, V. Ferri.

VINUM FERRI, V. Chalybeatum seu ferratum seu Martiatum seu Martis, Tinctura Martis vino'sa seu Martis cum vino, Steel wine, Wine of Iron. (Iron wire, 3j; Sherry wine, Oij (Imp. meas.). Digest for thirty days and filter. Ph. L.) Tonic. Dose, f 3j to f 3iv.

It is sometimes made by adding two pints of Wine to two ounces of Iron filings; macerating and filtering.

VINUM GENTIANÆ, V. G. Compositum seu Amarum seu Amarum cum spiritu Vini, Infusum amarum vino'sum, Compound wine of gentian, Wine bitter. (Rad. gentian. luteæ, 3ss, Cort. cinchon. offic. 3j, Cort. aurant. 3ij, Cnelliæ alb. 3j. Alcohol. dilut. f 3iv, Vini albi, 3ss, Hispanicæ. Oijss. Ph. E.) Dose, f 3j to f 3vj.

VINUM HIPPOCRATICUM, Claret—v. Hispanicum, Vinum, see Wine—v. Hordeaceum, Cerevisia—v. Hydatodes, see Hydatoid.

VINUM IPECACUAN'HE, Ipecacuan'ha wine. (Ipecacuan. cont. 3ij, Vini albi, Oij.) Macerate for 14 days, express, and filter. It may be prepared, also, by displacement. Ph. U. S.) Emetic and diaphoretic. Dose, as an emetic, f 3iv to f 3x.

VINUM DE KINÂ KINÂ COMPOSITUM, V. cinchonæ compositum, Compound bark wine, (F.) Vin de quinquina composé, (Cort. cinchon. pulv. ʒss, Quassia, Cort. Winteran., Cort. aurant. sicc. 3ss, Alcohol. (12° to 22° Beaumé) Oj. Macerate for 24 hours; and add Vini albi, Ovj. Macerate for four days, and filter. Ph. P.) Tonic, febrifuge, and stomachic. Dose, f 3ss to f 3ij.

VINUM MARTIATUM, V. Ferri—v. Martis, V. ferri—v. Medicatum, see Claret.

VINUM MEDICINA'LE, Medicated Wine. Wine, holding in solution one or more medicines. Medicinal wines are obtained by macerating or digesting medicinal substances in wine. They differ, of course, according to the nature of the medicines employed, the strength of the wine, &c. In a late Pharmacopœia of the London College, all the wines were changed into weak tinctures, but old names were left unaltered. Sherry is the wine usually employed. The following are medicated wines. See Vinum.

VINUM MERTUM, Acriatum—v. Nicotianæ tabaci, V. tabaci—v. Opiatum fermentatione paratum, Laudanum abbatiss Rousseau.

VINUM O'PII, V. de Opio compositum, Laudanum liguidum Sydenhami seu Hoffmanni, Tinctura Thebaica, Wine of Opium, Sydenhami Laudanum, Ford's Laudanum, (Opium pulv. 3ij; Cinnamon. cont., Caryophyll. cont. 3ss, 3j, Vini albi, Oj. Macerate for 14 days; express and filter. It may be made, also, by displacement. Ph. U. S.) Narcotic. Dose, gtt. xv to f 3j.

VINUM DE OPIO COMPOSITUM, V. opii—v. Picatum, Pissites.

VINUM PORTUGAL'LICUM, V. rubrum, (Ph. U. S. 1831.) V. Portense, (Ph. U. S. 1863.) (F.) Vin rouge d'Oporto. Wine of Portugal or of Oporto. Port wine. A strong astringent wine, not much drunk in the United States, but greatly used in England.

VINUM QUINIE, Wine of Quinia. (Wine, ʒij, Sulphate of quinia, gr. x. Mix.) Dose, f 3ss or more; according as it is used as a stomachic, or to arrest ague. A Vinum Cinchoniz may be made in the same way.

VINUM RHABARBARI, V. rhei palmati.

VINUM RHEI, V. R. palmati, V. Rhabarbari, Tinctura Rhabarbari Vinosæ, Elixi'r Rhei dulci, E. R. Compositum, Rhabarbar Wine. (Rhei cont. 3ij, Cnelliæ cont. 3j, Alcohol. dilut. f 3ij, Vini albi, 3xiv. Macerate for 14 days; express, and filter. It may be made, also, by displacement.

Ph. U. S.) Laxative and stimulant. Dose, f $\overline{3}$ iv to f $\overline{3}$ iss.

VINUM RHEI PALMATI, V. Rhei—v. Rhenanum, see Wine—v. Rubrum, V. Portugallicum—v. Rubrum Portugallicum, see Wine—v. Stibiatum, V. Antimonii.

VINUM TABACI, *Vinum Nicotianæ Tabaci*, *Tobacco Wine*. (*Tubac*, incisi. $\overline{3}$ j, *Vini albi*, Oj. Macerate for 14 days; express and filter. *Ph. U. S.*) Narcotic, diuretic, and antispasmodic. In dropsical cases, colica pictorum, and ileus. Not much used. Dose, gtt. xx to xxxvj.

VINUM VERATRI, V. V. *albi*, *Wine of White Hellebore*. (*Veratri* cont. $\overline{3}$ iv; *Vini albi*, Oj. Macerate for 14 days, express and filter.—*Ph. U. S.*) In cutaneous affections, and in gout. Dose, ten minims.

VINUM VERATRI ALBI, V. Veratri—v. Xericum, see Vinum.

VIOL, (from *violare*, 'to defile,') Stuprum.

VIOLA, (after *Ion*, *mw*, 'the violet,') *V'iola odorata seu mar'tia*, *Pon*, *Matrona'lis*, *Viola'ria*, *Sweet Violet*, (F.) *Violette à bouquets*, *V. odorante*. Ord. Violacæ. The flowers of the sweet violet have a pleasant, peculiar odour, and scarcely any taste. They are slightly laxative.

VIOLA CANINA, *V. sylvestris*, *Dog Violet*. The root of this plant is emetic and cathartic. Dose, $\overline{3}$ j.

VIOLA IPECACUANHA, see Ipecacuanha—v. Lutea, Cheiranthus cheiri—v. Martia, Viola—v. Odorata, Viola—v. Palustris, Pinguicula vulgaris.

VIOLA PEDA'TA, *V'iola* (*Ph. U. S.*) This indigenous violet, which grows from New England to Carolina, is regarded by some as a useful demulcent in pectoral affections.

VIOLA SYLVESTRIS, V. Canina.

VIOLA TRI'COLOR, *Jac'e'a tri'color*, *Her'ba seu Flos Trinitatis*, *Heart's ease*, *Pan'sy*, *Pen'sy*, *Pan'sies*, *Johnny Juniper*, (Prov.) *Beedy's eyes*, *Biddy's eyes*, (F.) *Pensée*, *P. sauvage*. A decoction of this beautiful little plant has been recommended in diseases of the skin, &c. Its roots are, also, emetic and cathartic.

VIOLARIA, Viola.

VIOLET, Viola—v. Dog, Viola canina—v. Dog's Tooth, Erythronium Americanum.

VIOLET BLOOM, Solanum dulcamara.

VIOLET, SWEET, Viola.

VIOLETTE À BOUQUETS, Viola—v. *Odorante*, Viola—v. *des Sorciers*, Vinca minor.

VIOLIER JAUNE, Cheiranthus cheiri.

VIORNA, Clematis vitalba.

VIPER, EGYPTIAN, (see Vipera,) Asp—v. Ten inch long, Echidna ocellata.

VIPER GRASS, Scorzonera.

VIPERA, ([L.] *vipera*, for *vivip'ara*, from *vi-*ens, 'alive,' and *pario*, 'to bring forth,') Coluber berus.

VIPERARIA, (from *vipera*), Aristolochia serpentaria, Scorzonera.

VIPERINA VIRGINIANA, (from *vipera*), Aristolochia serpentaria.

VIPÉRINE DE VIRGINIE, Aristolochia serpentaria.

VIPER'S DANCE, Chorea.

VIR, gen. *V'ris*, 'a man,' Aner, Penis—v. Ef-fœminatus, Gynæcaner.

VIRA, Woman.

VIRAGO, (from *vir*, 'a man,') see Pogoniasis.

VIREUX, Virous.

VIRGA, 'a rod,' Penis—v. Cereæ, *Bougie*—v. Cerebri, Pineal gland.

VIRGILIA, (after the poet Virgil,) Cladrastis tinctoria.

VIR'GIN, *Vir'go*, gen. *Vir'ginis*, *Par'thenus*, *Ex'pers nuptia'rum*, (F.) *Vierge*, *Pucelle*. A female who has never had sexual intercourse. Ap-

plied, also, occasionally to the male so circum-stanced.

VIRGIN MARY THISTLE, *Centaurea benedicta*.

VIR'GINAL, (from *virgo*.) The external genital organs, of the virgin especially.

VIRGINAL MEMBRANE, Hymen.

VIRGINALE CLAUSTRUM, 'a virginal fence,' Hymen.

VIRGINIA, MIN'ERAL WA'TERS OF. There are several celebrated mineral springs in Virginia, on the western side of the Blue Ridge. Two Thermal springs are in Bath county. The Warm Spring, as it is called, issues with a bold stream, sufficient to turn a mill, and to keep the water of its basin at the temperature of 97°. The Hot Spring is a few miles from the warm. Its temperature is 107°. They are both slightly sulphureous. The Sweet Springs are at the eastern foot of the Allegheny, in Botetourt county; 42 miles from the Warm Springs. The water has a temperature of 70°, and is highly impregnated with carbonic acid. The White Sulphur, as well as the Red Sulphur, and Salt Sulphur Springs, are also much visited. The Red Spring is affirmed to exert a sedative influence on the pulse, diminishing the number of its beats. Prof. W. B. Rogers analyzed the water of the White Sulphur, and found it to be composed as follows:—Gaseous contents in a wine gallon; Sulphuretted hydrogen, 2.5 cubic inches, carbonic acid, 2; oxygen, 1.448; nitrogen, 3.552.—Total, 9.5. Solid contents in a pint; sulphate of magnesia, 5.588 grains; sulphate of lime, 7.744; carbonate of lime, 1.150; chloride of calcium, 0.204; chloride of sodium, 0.180; oxide of iron, a trace; loss, 0.410. Total, 15.276 grains.

The Rock'bridge Alum Springs are situated 17 miles west of Lexington, and are in much favour. Their solid constituents are said to be,—sulphate of potassa, 1.765 gr.; sulphate of lime, 3.263; sulphate of magnesia, 1.763; protoxide of iron, 4.863; alumina, 17.905; crenate of ammonia, 0.700; chloride of sodium, 1.008; silicic acid, 2.840; free sulphuric acid, 15.224; free carbonic acid, 7.536—water 58315.133. They are used in dyspepsia, scrofula, chronic diarrhoea, and cutaneous affections. The Bath Alum Springs are about 17 miles west of the Rockbridge Springs, in Bath county. They consist of free sulphuric acid, 7.878 grains; carbonic acid, 3.846; sulphate of potassa, 0.258; magnesia, 1.282; lime, 2.539; protoxide of iron, 21.776; alumina, 12.293; crenate of ammonia, 1.776; silicate of soda, 3.150; water, 58317.202. They are more chalybeate than the last.

About 15 miles from Staunton are *Strib'ling's Springs*—a sulphuretted water; and near Warrenton, in Fauquier county, there is a sulphureous water, called *Lee's Springs*, which is much frequented. Berkley or Bath, and Ca'pon Springs are likewise celebrated.

VIRGINIS VITIATIO, 'violation of a virgin,' Defloration.

VIRGIN'ITY, *Virgin'itas*, *Parthen'e'a*, *Integ'ritas*, *Maid'enhood*, *Maid'enhead*, (F.) *Virginité*, *Pucelage*, (from *virgo*.) The state of one who has never had carnal commerce.

VIRGIN'S BOWER, COMMON, Clematis vitalba, Clematis virginica—v. Bower, upright, Clematis recta—v. Sweet-scented, Clematis flammula.

VIR'GIN'S MILK, *Lac Vir'ginis* seu *Virginal'e* seu *Virgin'eum*, (F.) *Lait virginal*. A tincture of benjamin or of any other balsam, precipitated by the addition of water, which gives it a milky appearance. Also, subacetate of lead precipitated by water. It has been used as a cosmetic to remove freckles, &c.

VIRGINITAS, Virginity—v. Deflorata. Deflo-ration.

VIRGO, gen. *Virginis*, Virgin—*v. Aurea*, *Solidago virgaurea*.

VIRGULA, (dim. of *virga*, 'a rod,') Penis.

VIRIDEÆRIS, 'green of brass,' Cupri sub-nocuas—*v. Montanum*, Cuprum.

VIRILE, (from *vir*, 'a man,') Penis.

VIRILE, *Virilis*, (from *vir*, 'a man,') Relating or appertaining to the adult age.

VIRILE AGE, see Age.

VIRILES'CENCE, (from *virilis* (*vir*, 'a man'), 'manly,') The state of the aged female in which she assumes certain of the characteristics of the male.

VIRILIA, (from *vir*, 'a man,') Pudibilia.

VIRILITAS, Adult age, see Age.

VIRILITY, Adult age, see Age.

VIROSE, Virous.

VIROUS, *Virose*, *Virens*, (F.) *Viroux*, (from *virus*,) Possessed of noxious properties. Also, substances having a peculiar nauseous taste and odour.

VIROSUS, Virous.

VIRTUE, Faculty.

VIRTUS, 'power,' (allied to *vir*, 'a man,') Faculty, Property—*v. Formatrix*, Plastic force—*v. Plastica*, Plastic force.

VIRULENT, *Virulentus*, (from *virus*,) That which is of the nature of a virus or which is caused by a virus. Highly noxious.

VIRUS. A Latin word, which signifies poison: but which, in medicine, has a somewhat different acceptation. By it is understood a principle, unknown in its nature and inappreciable by the senses, which is the agent for the transmission of infectious diseases. Thus we speak of the *varicolic*, the *rubeoline*, and the *syphilitic viruses*. *Virus* differs from *venum* in the latter being a secretion natural to certain animals, whilst the former is always the result of a morbid process,—a morbid poison.

VIRUS, Ichor, Poison, Sanies—*v. Vaccine*, see *Vaccina*—*v. Vitale*, Sperm.

VIS, gen. *Visis*, (allied to *vir*, 'a man,') Force—*v. Adhesionis*, Cohesion (force of)—*v. Attractionis*, Cohesion (force of)—*v. Aucta*, Sthenia—*v. Cohesionis*, Cohesion (force of)—*v. Conservatrix*, *V. mediatrix* nature—*v. Elastica*, Elasticity—*v. Essentialis*, Plastic force.

VIS EXPEL'TRIX. The organic force by which, it was supposed, the useless and effete was removed from the body.

VIS FORMATIVA, Plastic force.

VIS A FRON'TE, 'force from the front,' An action which is exerted from before: as that of the capillaries in the circulation of the blood.

VIS GENERATRIX, Potentia generandi—*v. Imminuta*, Asthenia—*v. Insita*, Irritability—*v. Irritabilitas*, Irritability.

VIS MEDICATRIX NAT'URE, *V. conservatrix*, *Autocrati'a*, *Autonomi'a*, *Autocrati'ria*, *Autogygiasis*, *Autotherapi'a*, *Physiatrice*, *Physiocratilla*, *Physiocrati'a*, *Physiatherapi'a*, (F.) *Forces médiatrice*. A term, employed by some, to express that instinctive healing power, in an animal or vegetable, by virtue of which it can repair injuries inflicted upon it, or remove disease.

VIS METABOLICA, Metabolic force.

VIS MOR'TUA. Under this name Haller included the mechanical properties of membrane,—flexibility, extensibility, elasticity, &c.—which are totally independent of the vital properties. They remain some time after the complete extinction of life in all its functions: seem to be connected with the mechanical arrangement of particles and the chymical composition of the substance in which they reside, and not to be affected until the progress of decomposition has become sensible.

VIS MUSCULARIS, Myodynamia—*v. Nervorum*, Nervous power—*v. Nervosa*, Nervous power—*v.*

Plastica, Plastic force—*v. Reproductiva*, Plastic force.

VIS A TER'GO, 'force from behind,' An action of impulsion or repulsion, as that of the heart in the circulation of the blood.

VIS VITÆ, Irritability.

VIS VITA'LIS, *Vis Vita*, *Vis seu Facultas Zōtica*, (F.) *Force vitale*. The vital force—Brod of Von Reichenbach—which distinguishes living matter from the dead. Also, Irritability.

VIS ZOTICA, Vis vitalis.

VISAGE, (from [L.] *videre*, *visum*, 'to see,') Countenance.

VISCAGO BEHEN, (*viscum*, 'birdlime,') Cucubalus behen.

VISCERA, (pl. of *viscus*, 'an internal organ,') Entrails, Uterus.

VISCERAL, Splanchnic.

VISCERALIS, Splanchnic, Stomachal.

VISCERATION, Exenterismus.

VISCERES, ([L.] *viscera*,) Entrails.

VISCONIA, Physconia.

VISCUM ALBUM, *Is'ia*, *Is'ia*, *Da'bach*, *Mis'tetoe*, (F.) *Gai*, *Gai de Chêne*, *Gilim*. A parasitic plant, which grows on a number of trees.

Order, Lorantheæ. *Sec. Syst.* Diœcia Tetrandria. That of the oak—*Viscum quercinum*—has been most celebrated, but it does not differ from the others. It was, at one time, much used in epilepsy and other neuroses; the wood or leaves being given in powder, in the dose of ʒss or ʒi. The berries, also called *Pommes hémorrhoidales*, from their supposed virtues, are drastic; and, when applied externally, are maturative.

VISCUM CAPES'Æ. A South African parasitic shrub, which grows on the stems of several species of *Rhus* and *Euclea*.

VISCUM FLAVESCENS, *V. verticillatum*, *Yel'low* or *Yel'lowish Mis'tetoe*, indigenous, has been given in similar cases as the *Viscum album*.

VISCUM VERTICILLATUM, *V. flavescens*.

VISCUS, gen. *Visceris*. Any internal organ of the body.

VISIO, Pupil, Vision.

VISION, *Sight*, *Vis'io*, *Vi'sus*, (from *videre*, *visum*, 'to see,') *Op'sis*, *Om'ma*, (F.) *Vue*. The action of seeing. The sight is one of the five senses, and the eye is its organ. By it we distinguish colours, and appreciate most of the physical qualities of external objects. The eye is composed of different transparent media, whose curvatures and refractive powers are so combined as to render insensible the aberrations of sphericity and refrangibility, and to concentrate the numerous rays proceeding from more or less distant objects. These rays *traverse*, successively, the transparent cornea, the aqueous humour, the crystalline, the vitreous humour, and strike upon the retina, on which they impress the figure of the object whence they have proceeded; and, in order that no obscurity may arise from the reflections, which might otherwise occur, the anterior of the eye is lined by a membrane—the choroid—which is itself covered internally by a dark pigment. This absorbs the light, like the dark pigment in the interior of telescopes, and thus obviates confusion. As the rays proceeding from an object must cross each other in their course to the retina, it follows that those proceeding from the upper part of an object must fall upon the lower part of the retina, and conversely: so that the object must be impressed upon the retina in an inverted position: yet experience teaches us to appreciate the position accurately.

VISION, BINOCULAR, see BINOCULAR—*v. Coloured*, Chromopsia—*v. Double*, Diplopia—*v. Iridescent*, Chromopsia—*v. Manifest*, Polyopsia—*v. Multiple*, Polyopsia—*v. Phantasmatum*, Metamorphopsia—*v. Single*, see BINOCULAR.

VISNOMY, Physiognomy.

VISORUM ORGANUM, (from *videre*, *visum*, 'to see,' Eye.

VISORIUS, Visual.

VISUAL, *Visua'lis*, *Visu'alis*, (F.) *Visual*. That which concerns the sight or belongs to vision.

VISUAL ANGLE is the angle formed by the crossing of two rays, proceeding from opposite points of any body, in their passage through the pupil. It is by the size of this angle that we judge, to a certain extent, of the dimensions of objects.

VISUEL, Visual.

VISUS, (from *videre*, *visum*, 'to see,') Vision—*v.* Aerior, Nyctalopia—*v.* Aeris, Oxydercia—*v.* Coloratus, Chromopsia—*v.* Debilitas, Asthenopia—*v.* Decolor, Achromatopsia—*v.* Defiguratus, Metamorphopsia—*v.* Dimidiatus, Hemipia—*v.* Divinus, Hemeralopia—*v.* Duplicitatus, Diplopia—*v.* Hebetudo, Amblyopia—*v.* Juvenum, Myopia—*v.* Lucidus, Photopsia—*v.* Muscarum, see Metamorphopsia—*v.* Nebulosus, Nebula—*v.* Nocturnus, Nyctalopia—*v.* Senilis, Presbytia.

VITA, Life—*v.* Propria, Irritability.

VITAL, *Vita'lis*, *Vital'is*, *Vital'is*, (F.) *Vital*, (from *vita*.) Relating or appertaining to life.

VITAL CAPAC'ITY, *Pulmon'is* or *Lung capac'ity*. A term employed by Dr. Hutchinson to designate the number of cubic inches of air, which a person can forcibly expire after a full inspiration. He found, that for every inch of height—from five feet to six—eight additional cubic inches of air can be given out by a forced expiration. The quantity, however, is influenced, also, by weight and age. See Respiration.

VITAL CENTRE, *Centrum vitale*—*v.* Dynamics, see Dynamic—*v.* Force, *Vis vitalis*—*v.* Point, see *Centrum vitale*.

VITAL PRINCIPLE, *Flamm'ula* seu *Au'ra vita'lis*, *Microcosm'us*, (F.) *Flamme vitale*, *Vital Flame*. The principle which, in association with matter, as in organized bodies, controls its manifestations and properties, and distinguishes organized matter from the inorganic. We know nothing of this principle, except as a force in connexion with organization.

VITAL PROPERTIES, see Properties, vital—*v.* Statistics, Statistics, medical.

VITALBA, (*vitis*, 'vine,' and *albus*, 'white,') Clematis vitalba.

VITALISM, *Vitalism'us*, (from *vita*.) The doctrine of the *vitalist*—*vitalis'ta*—who ascribes every function of the organism to the direct agency of a vital principle or force.

VITALIST, see Vitalism.

VITALITY, *Vital'itas*, (F.) *Vitalité*. The vital principle: also, used in the sense of vital action or movement.

VITALIZE, To endow with life.

VITALS, *Vita'lia*. Parts that are necessary for life—the heart, lungs, and nervous centres more especially.

VITAROB'ORANT, **PALM'ER'S**, (*vita*, and *roborare*, 'to strengthen,') A farinaceous preparation, which appears to consist of a mixture of the red or Arabian lentil and wheat flour, the latter in considerable amount, sweetened with sugar. It bears considerable resemblance to Revalenta.

VITELLARY, Vitelline—*v.* Membrane, *Zona pellucida*.

VITELLINE, *Zona pellucida*.

VITELLINE, *Vitel'linus*, *Vit'ellary*, *Lecith'odea*, (from *vitellus*, 'the yolk of an egg,') Appertaining to or resembling the vitellus or yolk. Also, the yolk membrane. See Vesicula umbilicalis.

VITELLINE DISC, Proliferous Disc—*v.* Fluid, see Vesicula umbilicalis.

VITELLINE PED'ICLE or **APOPH'YSIS** is the pedi-

cle which attaches the umbilical vesicle to the embryo. See Vesicula umbilicalis.

VITELLINE POUCH, *Vesicula umbilicalis*—*v.* Vessels, Omphalomesenteric.

VITELLOMESENTERIC, Omphalomesenteric.

VITELLUM OVI, 'the yolk of an egg,' see Ovum.

VITELLUS OVI, see Ovum.

VITEX, gen. *Viticia*, (from *vicio*, 'I bind' (?) : *V. Agnus castus* seu *verticillatus*, (F.) *Guaiiver*, *Petit Poivre*, *Poirée de main* ou *marma*, Ord. Verbenaceae. See *Syst. Didynamia Angiospermia*. The *Choate Tree*. The seeds, when fresh, have a fragrant smell, and an acrid, aromatic taste. Formerly, they were celebrated as antaphrodisiacs.

VITI SALTUS, 'Vitus's dance,' Chorea.

VITIA (pl. of *vitium*) **PRIME CONFORMATIONIS**, see Monster.

VITIATIO, (*vitare*, *vitatum*, 'to violate,') Stuprum.

VITILIGINES, Ephelides.

VITILIGO, gen. *Vitilig'ina*, (from *vitulus*, 'a calf,' hence 'calfskin,') A disease of the skin, so called from its resembling the vitiligo of Willan. It appears under two forms; either as tubercles—*V. tuberosa*—varying from the size of a pin's head to that of a large pea. Isolated or confluent; or as yellow patches of irregular outline, slightly elevated, and with but little hardness—*V. plana*.

VITIS ALBA SYLVESTRIS, (*vitis*, 'to bind,') from its numerous tendrils,) *Bryonia alba*—*v.* *Apyrena*, see *V. Corinthaica*.

VITIS CORINTH'ACA, Ord. Vitaceae. See *Syst. Pentandria Monogynia*. This plant affords the *Uva Corinthaica* *pas'sa* seu *min'ra*, *Pas'sa Corinthaica* seu *Pas'sula min'ra*, *Corinthaica*, *Cor'caica*, (Prov.) *Cura'barria*, (F.) *Raisin de Damas*; which are the dried fruit of the *Vitis Corinthaica* or *Vitis opuscula*; and are similar, in their properties, to the raisin.

VITIS IDÆA MYRTILLUS, *Vaccinium myrtillus*—*v.* *Idæa palustris*, *Vaccinium oxycoccus*—*v.* *Idæa punctata*, *Vaccinium vitis idæa*—*v.* *Sativa*, *V. vinifera*.

VITIS VINIF'ERA, *V. sat'va*, *Au'pela*, *A. Enoph'ora*, *Grapevine*, *Vine*, (F.) *Vigne*. Vine leaves and the tendrils have an astringent taste, and were formerly used in diarrhoea, hemorrhage, &c. The juice or sap, called *Za'ryna*, has been recommended in calculous disorders and in cases of chronic ophthalmia. The juice of the unripe fruit is called *Vin'juice*. The *Grape*, *Uva*, *Bol'trys*, *Staph'yli*, *Rhax*, (F.) *Raisin*, when ripe, is a wholesome, delicious fruit; and, when dried, forms the *Uva pas'sa* *major*, *Pas'sula major*, *Uva pas'sa* *co'le siccata*, *Uva pas'sa* (Ph. U. S.) *Zibeb*, *Raisin*. These are inodorous; and to the taste, subsaccharine, sweet, and mucilaginous. They are demulcent and nutritive.

VITIUM, Disease—*v.* Caducum, Epilepsy.

VITIUM CONFORMATIONIS, A disease or fault of conformation.

VITIUM DIVINUM, Epilepsy—*v.* Heretulum, Epilepsy—*v.* *Seraphulosum*, *Serofula*.

VITRARIA, *Parietaria*.

VITRE, (from *vitrum*.) Hyaline.

VITREOCAPSULITIS, (*vitreo*, 'appertaining to the vitreous humor,' and *capulitis*, 'inflammation of a capsule,') Hyalitis.

VITREOUS, (from *vitrum*.) Hyaline—*v.* Humour, *Corpus vitreum*—*v.* Humour of the ear, *Vitrine auditive*.

VITREUS, Hyaline.

VITREUX, Hyaline.

VITRINA OCULARIS. Corpus vitreum.

VITRINE AUDITIVE (F.). *Humeur vitrée*, (from *vitrum*.) *Aquula vitrea auditiva*, *Endolymph'a*, *Endolymph*, *Liquor* or *Fluid of Scarpa*, *Vitreous Humour of the Ear*. The fluid contained in the membranous labyrinth of the ear, so called from its resemblance to the *vitreous* humour of the eye.

VITRIOL, (from *vitrum*, 'glass,') *Ferri sulphas* — v. Blue, *Cupri sulphas* — v. Green, *Ferri sulphas* — v. Roman, *Cupri sulphas* — v. White, *Zinci sulphas*.

VITRIOLUM ALBUM, *Zinci sulphas* — v. *Ammoniacale*, *Ammonie sulphas* — v. *Caruleum*, *Cupri sulphas* — v. *Cupri*, *Cupri sulphas* — v. *Cyprium*, *Cupri sulphas* — v. *Ferri*, *Ferri sulphas* — v. *Goslariense*, *Zinci sulphas* — v. *Martis*, *Ferri sulphas* — v. *Romanum*, *Cupri sulphas* — v. *Veneris*, *Cupri sulphas* — v. *Viride*, *Ferri sulphas* — v. *Zinci*, *Zinci sulphas*.

VITRUM, *Hy'alos*, *Glass*, *Bran'cia*, (F.) *Verre*. Glass, coarsely powdered, has been sometimes used to remove specks of the cornea.

Powdered glass has been regarded, but erroneously, as a poison. It acts only mechanically.

The saline scum, which swims on glass when first made, has been used in toothpowders. It is called *Sau'diver*, *Sel de verre*, *Glass Gall*, *Fel vitri*, &c.

VITRUM ANTIMONI, see *Antimonii vitrum* — v. *Hypoclepticum*, *Separatorium* — v. *Oculare*, see *Spectacles* — v. *Sublimatorium*, *Aludel*.

VITTA, *Bandelette*, *Caul*.

VITTARIA CETERACH, (*vitta*, 'a band, a fillet,') *Asplenium ceterach*.

VITTIE VAYR, *Andropogon muricatus*.

VIVACE, (from *vivax*, gen. *viva'cia*, (from *vivere*, 'to live,') 'living long, tenacious of life.' An epithet given by *Levet* to uterine polypi, which he considers to be, commonly, vegetations of some ulcer in the interior of the uterus, and incurable.

VIVÆSECTIO, (*vivus*, 'alive,' and *secare*, *sectum*, 'to cut,') *Vivisection*.

VIVERRA CIVETTA, see *Civetta*.

VIVIPARE, *Viviparous*.

VIVIPARITÉ, see *Viviparous*.

VIVIPAROUS, (F.) *Vivipare*, (*vivus*, 'alive,' and *pario*, 'to bring forth,') An epithet applied to animals whose young are brought forth detached, without requiring to be hatched, as in the *Oviparous*. The condition may be termed *Viviparousness*, (F.) *Viviparité*. See *Generation*.

VIVIPAROUSNESS, see *Viviparous*.

VIVISECTION, *Vivisection*, *Vivisection*, *Biotomia*, (*vivus*, 'alive,' and *secare*, *sectum*, 'to cut,') The act of opening or dissecting living animals. One who performs vivisections is a *Vivisector*.

VOCAL, (from *vox*), *Oral* — v. *Cords*, see *Cords*, *vocal* — v. *Ligaments*, see *Cords*, *vocal* — v. *Tube*, see *Tube*, *vocal*.

VOCALIS, *Vowel*.

VOCE DI PETTO, 'chest voice,' see *Voice* — v. *di Testa*, 'head voice,' see *Voice*.

VOICE, *Vox*, *Au'dē*, *Pho'nē*, (F.) *Voix*, (from *vocare*, 'to name, to call,') An appreciable sound produced by the air, when driven from the lungs, throwing the inferior ligaments of the glottis into vibration. The function by which this is effected has been called *Phonation*. It is a function of *animal* life; and, in animals, is limited to the production of the *simple* or *instinctive* voice, as well as in the idiot; but, in intellectual man, becomes much more complicated and articulated, — an evidence of his intellectual superiority.

VOICE, ARTICULATED, *Speech*, *Loque'la*, *Glos'sa*, *Glott'a*, *La'lia*, *Phra'sis*, (F.) *Parole*, *Voiz articulée*, is the voice as produced by the glottis, modi-

fied by the motions of the tongue, lip, and other parts of the vocal tube. Most physiologists agree in regarding the glottis, or the space between the thyroarytenoid ligaments plus the ligaments themselves, as the part where vocal sounds are formed; but the mechanism of this formation is a matter of dispute. The greater part of individuals regard the glottis and its ligaments, with the *vocal tube*, or the space which the sound has to traverse after it is produced in the glottis, as a reed instrument, like the clarinet, the inferior ligaments of the glottis constituting the reed. In speech, as ordinarily effected, the tongue, lips, teeth, &c., are concerned, but there are some authentic cases on record, where it has been accomplished after the tongue has been removed. In such cases, the articulated voice must have been produced in the glottis alone, or in it aided by an obscure action of the parts above it.

The singing voice is not limited to the larynx; the pharynx would seem to be concerned also. The notes that are produced in the latter have been termed *supralarynge'al*, or *notes of the second register*. They constitute the *pharynge'al voice*, *falsetto* or *fauvette* voice or *voce di testa* (I.), *Voix de tête*, *Voix de fausset* (F.). The *larynge'al voice* or *voice of the first register*, *Chest voice*, is the *Voce di Petto* (I.), *Voix de Poitrine* (F.). The *laryngeal* voice, which is always more elevated, by an octave, in the female than in the male, furnishes the types called *Sopra'no*, *Al'to* or *Contra'to*, *Tenore* and *Basso*. The first two belong to the female voice; the last two to the male. The *Bar'itone* is between the bass and tenor.

VOICE, BLEATING, *Egophony* — v. *Cavernous*, *Pectoriloquy* — v. *Chest*, see *Voice*, articulated.

VOICE, CHOLERIC, *Vox Choleric'a*, *Cholero-phonia*, *Cholaphonia*. The peculiar husky voice of those affected with cholera.

VOICE, CONVULSIVE, *Vox convulsiva*, (F.) *Voix convulsive*. A neurosis of the voice, consisting in the production of discordant sounds — acute and grave — which cannot be brought to the natural tones by any effort of volition. It seems to be caused by disordered contraction of the muscles of the larynx.

VOICE, FALSETTO, see *Voice* — v. *Goat's*, *Egophony* — v. *Laryngeal*, *Laryngophony*, see *Voice* — v. *Pharyngeal*, see *Voice* — v. *Tracheal*, *Laryngophony*.

VOIE, ([L.] *via*), *Way*.

VOIES DIGESTIVES, *Ways*, *digestive* — v. *Lacrymales*, *Lachrymal passages* — v. *Premières*, *Ways*, *digestive* — v. *Secondes*, *Ways*, *second* — v. *Urinaires*, *Urinary passages*.

VOILE ([L.] *velum*) DU PALAIS, *Velum pendulum palati* — v. *Médullaire inférieure*, *Velum medullare inferius* — v. *Médullaire supérieure*, *Valvula Viesseni*.

VOIX, ([L.] *Vox*), *Voice* — v. *Aigre*, see *Acid* — v. *Articulée*, *Voice*, articulated, *Pectoriloquy* — v. *Bourdonnante*, see *Resonance* — v. *Bronchique*, *Resonance* — v. *Caverneuse*, *Pectoriloquy* — v. *Chévroutante*, *Egophony* — v. *Égophonique*, *Egophony* — v. *de Fausset*, see *Voice*, articulated — v. *de Poitrine*, see *Voice*, articulated — v. *de Polichinelle*, *Egophony* — v. *Sénile*, *Egophony* — v. *de Tête*, see *Voice*, articulated — v. *Tubaire*, see *Resonance*.

VOLA, *Palm* — v. *Mands*, *Palm* — v. *Pedis*, *Sola*.

VOLANT D'EAU, *Nymphæa alba*.

VOLATICA, (*volare*, 'to fly,') *Lichen*.

VOLATILIZATION, *Volatilisa'tio*, *Volatifa'tio*, (from *volare*, 'to fly,') The operation of reducing volatilizable substances into gas or vapour.

VOLATISATIO, (*volare*, 'to fly,') *Volatilization*.

VOLCANISATION, see *Vulcanite*.

VOLIT'ION, *Volit'io*, (from *volo*, 'to will,') The act of willing. Hence:

VOLIT'IONAL, *Relating to volition*. Thus

the muscles may be moved by a volitional or by an emotional impulse.

VOLNUS, Wound.

VOLONTAIRE, Voluntary.

VOLONTÉ, Voluntas.

VOLSELLA, (*vellere*, 'to pluck,') *Acanthabobus*, Forceps, *Madisterium*—*v. Patini*, *Valet à Patin*.

VOLTAISM, (after *Volta*,) Galvanism.

VOLTUS, Countenance, Face.

VOLUMETRIC, *Volumetricus*, (F.) *Volumétrique*, (from *volume*, and *meter*.) Relating to the measurement of volumes; as the *volumetric* method of analysis for the urine, the principle of which consists in adding to a known quantity of urine the exact amount of a test liquid required to precipitate the whole of any one ingredient contained in it.

It is usually employed only for the determination of the chlorine, urea, phosphoric and sulphuric acids, and sugar of the urine.

VOLUNTARY, *Hecusius*, *Voluntarius*, (F.) *Volontaire*, (from *voluntas*, 'will,') Relating to the will; as 'voluntary muscles,' muscles which act in obedience to the will.

VOLUNTAS, gen. *Voluntatis*, *Volentia*, *Arbitrium*, *Cupiditas*, *Boule'sis*, *Bule'sis*, *Thelema*, (F.) *Volonté*, (from *volo*, 'to will,') Will or desire.

VOLUNTAS seu CUPIDITAS DESEDEND'I, *Lubi'do intestini*. Desire to go to stool.

VOLVA, Vulva.

VOLVULUS, (from *volvere*, 'to roll,') *Ileus*—*v. Intestinorum*, *Intussusceptio*.

VOMENDI CONAMEN INANE, 'vain effort at vomiting,' *Vomituration*.

VO'MER, gen. *Vo'meris*, 'a ploughshare,' *Os vo'meris*, *Os maxillæ superioris undecimūm*, *Ara'trum*. One of the bones of the face, so called from its resemblance. It is an azygous bone, forming the posterior part of the septum narium. It is thin, flat, and quadrilateral; and is articulated, below, to the superior maxillary and palatine bones; and above, to the sphenoid, ethmoid, and cornets of Bertin. It is developed by a single point of ossification. Also, the penis.

VOMICA, gen. *Vom'icæ*, *Pyocystis*, (from *vomere*, 'to vomit,' 'to cast up,') (F.) *Vomique*, *Crachement de pus*. A term used by some authors for any encysted collection of purulent matter in the interior of a viscus. It is commonly, however, restricted to a purulent collection in the lungs, encysted or not, which may be discharged by the mouth, by breaking into the bronchial tubes. Two kinds of vomices are distinguished; the one caused by the softening or breaking down of pulmonary tubercles; the other, of much more rare occurrence, formed by a circumscribed abscess of the lungs. Instead of opening into the bronchia, the abscess is sometimes discharged into the cavity of the pleura, causing empyema.

VOMICINE, Brucine.

VOMIQUE, Vomica.

VOMIQUEUR, *Strychnos nux vomica*.

VOMISSEMENT, (from *vomir*, [L.] *vomere*, 'to vomit,') Vomiting—*v. de Sang*, *Hæmatemesis*.

VOMIT, (*vomere*, *vomitum*, 'to vomit,') Emetic—*v. Black*, Fever, yellow, Melæna.

VOMIT, DRY, OF MAR'RIOTT. A once celebrated emetic, called 'dry,' from its having been given without solution. It consisted of equal portions of tartrate of antimony and potassa, and sulphate of copper.

VOMITIF, Emetic.

VOMITINE, Emetine.

VOMITING, *Vomititus*, *Vomit'io*, *Vomit'ium*, *Vomitum*, *Emomit'io*, *Ana'tropé*, *Em'esis*, *Em'etus*, *Pal'mus vom'itus*, *Ægritu'do ventriculi*, *Spew'ing*, *Pu'king*, *Sick'ness of the Stom'ach*, (Old Eng.)

Par'breaking, (Sc.) *Bok'king*, *Bock'ing*, *Bra'king*, *Brocht*, (Prov.) *Hull'up'ing*, (vulg.) *Casca'd'ing*, *Cast'ing*, (F.) *Vomissement*. The act by which solids and liquids, contained in the stomach, are ejected by the mouth. Vomiting is the effect of some special irritation of the nerves of the stomach, calling, by reflex action, on the appropriate muscles to expel the contents of the viscus. It is not accomplished solely by the contraction of the stomach, as was for a long time believed. That organ, indeed, assists in the process; but the main cause is the compression exerted upon the parietes of the viscus by the diaphragm and abdominal muscles. Vomiting is a symptom common to numerous diseases. At times, it is sympathetic, as in affections of the kidney, uterus, brain, &c. At others, it is symptomatic, as in gastritis, peritonitis, &c. When very troublesome, it may often be palliated by effervescent draughts, aromatics, sinapisms, or blisters to the epigastric region, &c.

VOMITING AT PLEASURE, Regurgitation—*v. of Blood*, *Hæmatemesis*—*v. Stercoraceous*, *Copremesis*.

VOMITIO, Vomiting—*v. Sanguinis*, *Hæmatemesis*.

VOMITIUM, Vomiting.

VOMITO NEGRO, 'black vomit,' Fever, yellow—*v. Prieto*, Fever, yellow.

VOMITORIUM, Emetic.

VOMITORY, Emetic.

VOMITUM, Vomiting.

VOMITURIT'ION, *Vomiturit'io*, *Em'esia*, *Dysemes'ia*, *Subver'sio stom'achi*; ineffectual efforts to vomit; *Vomen'di cona'men ina'næ*, *Retch'ing*, *Reach'ing*, *Heav'ing*, (Old Eng.) *Bolke*, (Sc.) *Bok*, *Bock*, *Braking*. Some authors mean, by this term, a vomiting of but little matter, or that which is effected almost without effort. The first acceptance is the most general.

VOMITUS, (from *vomere*, *vomitum*, 'to vomit,') Vomiting—*v. Cruentus*, *Hæmatemesis*—*v. Fæculentus*, *Copremesis*—*v. Marinus*, *Nausea marina*—*v. Navigantium*, *Nausea marina*—*v. Niger*, Fever, yellow—*v. Pituitosus*, *Blennemesis*—*v. Profusus*, *Hyperemesis*—*v. Purulentus*, *Pyemesis*—*v. Sanguinis*, *Hæmatemesis*—*v. Stercoris*, *Copremesis*.

VOORARA, Curare.

VOTIVE TABLET, (from *votum*, 'a vow,') *Tabula votiva*.

VOUACAPOUA AMERICANA, *Geoffræa inermis*.

VOUSSURE PRÉCORDIALE, (*vouter*, 'to arch,' from [L.] *volvere*, *volutum*, 'to curve,') 'præcordial arching or prominence.' The elevation or arching of the parietes of the thorax, witnessed in certain thoracic affections, as in hypertrophy of the heart.

VOÛTE, (from [L.] *volvere*, *volutum*, 'to curve,') Vault—*v. Médullaire*, *Corpus callosum*—*v. à Trois Piliers*, *Fornix*.

VOWEL, *Voca'lis*, (from *vox*, gen. *vo'cis*, 'the voice,') (F.) *Voyelle*. Physiologically, a continuous breath or sound, produced in the glottis, but more or less modified by the form of the vocal tube, through which it has to pass. In the English language, *a*, *e*, *i*, *o*, *u*, *w*, and *y*, are vowels.

VOX, gen. *Vo'cis*, Voice—*v. Cholericæ*, Voice, choleric—*v. Clandestina*, Whispering—*v. Clangosa*, *Oxyphonia*—*v. Convulsiva*, *Voix Convulsive*—*v. Nasalis*, *Rhinophonia*—*v. Rauca*, *Raucedo*, *Raucous voice*—*v. Rancisona*, *Raucous voice*.

VOYELLE, Vowel.

VUE, (*voir*, 'to see,' *vu*, 'seen,') Vision—*v. Basse*, *Myopia*—*v. Courte*, *Myopia*—*v. Diurne*, *Hemeralopia*—*v. Faible*, *Amblyopia*—*v. Longue*, *Presbytia*—*v. Louche*, *Strabismus*—*v. Nocturne*, *Nyctalopia*.

VULCANITE, DENTAL, (from *Vulcanus*, *Vulcan*, 'the god of fire.') A compound of caoutchouc and sulphur or sulphur compounds, used in the formation of plates for artificial teeth, &c. The formation of vulcanite is termed *Vulcanisation*, (F.) *Vulcanisation*, *Sulfuration*.

VULGA, *Vulva*.

VULNERA SCLOPETARIA, (pl. of *vulnus*, and *sclopetum*, 'a gun,') see *Sclopetarius* and *Wound*—v. *Sclopetica*, see *Sclopetarius* and *Wound*—v. a *Sclopetis*, see *Sclopetarius* and *Wound*.

VULNÉRAIRE, *Vulnerary*—v. *Suisse*, see *Falltranck*.

VULNERARIUS, *Traumatic*.

VULNERARY, *Vulnerarius*, *Traumat'icus*, *Traumat'ic*, (from *vulnus*, gen. *vul'neris*, 'a wound,') (F.) *Vulnéraire*. A name applied by the ancients to medicines, which they considered capable of favouring the consolidation of wounds. The property has been ascribed to various plants.

VULNERATIO NERVI, (*vulnerare*, *vulneratum*, 'to wound,') *Neurotrosis*.

VULNERUM VINCTURA, 'binding of wounds,' *Deligation*.

VULNUS, gen. *Vul'neris*, (from *vellere*, 'to pluck or tear,') *Wound*—v. *Laceratum*, see *Laceration*—v. *Sclopetarium*, *Wound*, *gun-shot*—v. *Simplex*, *Cut*.

VULNUSCULUM, (dim. of *vulnus*,) *Wound*.

VULPIS MORBUS, 'disease of the fox,' *Alopacia*.

VULTUEUX, (F.) *Vultuosus*, 'full of expression,' (from *vultus*,) *Corvisart* has used the expression, *Face vultueuse*, *Fa'cies vultu'osa*, for the face when it is more florid and swollen than in the natural state. This appearance, which is observed in active aneurism of the heart, according to *Corvisart*, is owing to the afflux of blood into the arterial capillaries, and hence differs from the livid and swollen appearance caused by the accumulation of blood in the venous capillaries, constituting the *Face injectée*.

VULTUS, *Countenance*, *Face*—v. *Tetricus* et *Mæstus*, *Scythrospasmus*.

VULVA, (from *volvere*, 'to enfold,' or from *volve*, 'folding doors,') (F.) *Uterus*, especially of animals; see *Vulva*.

VULVA, *Vulva*, *Hor'tus*, *Cun'us*, *Puden'dum mul'eb're*, *Pudenda mul'eb'ria*, *Sig'nus pud'o'ris*, *Le'tar*, *Charus*, *As'eum*, *Buc'ea*, *Bul'ga*, *Cadu'cus*, *Cus'tos*, *Fe'men*, *An'ulus*, *Fe'mur sum'mum*, *Follic'ulus*, *Fun'dus*, *Gre'mium*, *Hia'tus*, *Lanu'rium*, *Navis*, *Oppid'ulum*, *Spe'cus*, *Vul'ga*, *Cym'ba*, *Gyn'esthus*, *Sal'tus*, *Clitor'ium*, *Os'tium*, *Sul'cus*, *Con'cha*, *Por'tus*, *Por'ta*, *Fores* (of virgins), *Interfemin'eum*, *Mul'eb'ria*, *Fossa magna mul'eb'ris*, *Epise'i'on*, *Es'chara*, *Tre'ma*, *Hor'tulus cupid'inis*, *Byssos*, *Ca'va*, *Caver'na*, *Del'phys*, *Del'ta*, *Fo'vea*, *Gynæce'um*, *Secess'us*, *Vallis fem'o'rum*, *Eugi'um*, *Eugi'on*, *Lo'ci mul'ic'b'res*, *Fe'male puden'dum*, *Fe'male or'gans of generation*, *Fe'male parts*, *Shape*, (F.) *Vulve*. A longitudinal opening between the projecting parts of the external organs of generation in the female, extending from the mons veneris to the perineum, and which is called, also, *Fossa magna*. Some anatomists mean, by this term, the whole of the external genital organs of the female:—mons veneris; labia majora, and the cleft or vestibule separating them; the clitoris; nymphæ; meatus urinarius; entrance of the vagina; with the hymen or the carunculae myrtiliformes, fossa navicularis and fourchette.

VULVA, *Fora'men commu'nē ante'rius*, *It'er seu Ad'itus ad infundib'ulum*, I. seu *Ad'itus ad tertium ventric'ulum*, *Ri'ma ad infundib'ulum*, *Aper'tura ante'rior ventric'uli tertii cereb'ri*. An opening situate at the junction of the optic thalami, in the third ventricle of the brain, and immediately beneath the anterior crus of the fornix.

VULVAIRE, *Chenopodium vulvaria*, *Vulvar*.

VULVAR, *Vul'varis*, (F.) *Vulvaire*, (from *vulva*,) That which relates to the vulva. *Chausier* gives this name to the external pudic arteries, because they are distributed to the vulva.

VULVARIA, (from its use in diseases of the vulva [?],) *Chenopodium vulvaria*.

VULVE, *Vulva*.

VULVITIS, (*vulva* and *itis*,) *Inflammation of the vulva*.

VULVOÛTERINE, *Vulvoûteri'nus*, (F.) *Vulvoûtéri'n*. Relating or appertaining to the vulva and uterus.

VULVOUTERINE CANAL, *Vagina*.

VULVOVAG'INAL, *Vulvovagina'tis*. Relating or appertaining to the vulva and vagina.

VULVOVAGINAL GLANDS, *Cowper's glands in the female*.

W.

WABRAN or **WAYBREAD**, ([Sax.] *pæg-hwæpe*,) *Plantago*.

WAD, ([Sax.] *wad*,) *Graphites*.

WAFERTREE, *Ptelea trifoliata*.

WAGNER, CORPUSCLES OF, *Corpuscula tactis*—w. *Vesicles of*, see *Molecule*.

WAGNERIAN SPOT, see *Molecule*.

WAHOO, see *Euonymus*.

WAIR, see *Pila marina*.

WAIST'COAT, STRAIT, *Indu'cula*, (F.) *Camisole*, *Gilet de force*. A dress, used for restraining maniacs, or those labouring under violent delirium. It has long sleeves, which are tied behind the body, so that the arms cannot be extricated from them. It ought, of course, to be made of very resisting materials.

WAKEROBIN, *Arum maculatum*, *Trillium latifolium*.

WALE, ([Sax.] *palan*, 'marks of stripes,') *Wheal*.

WALK'ING, *Gres'sus*, *Inces'sus*, *Deambula'tio*, *Ambula'tio*, *Bad'ieis*, (F.) *La Marche*, *Progression*. ([Sax.] *pealcan*, 'to walk,') The action by which we change place, by means of a succession of steps in the same direction. In walking forwards, the centre of gravity is constantly altered; and a series of small, parabolic curves is described on the heads of the thighbones, the extent of which is regulated by volition, under the guidance of vision.

WALLA'CHIA, MIN'ERAL WATERS OF. There are many excellent sulphureous springs in this country, especially those at *Bobotsch*, *Finceschti*, *Sibitschiudi Suz*, *Bräsa*, *Otschin*, *Serbo-neschte*, *Kimpalungi*, *Kosia*, and *Oloneschti*. *Chalybeate* and saline springs have, also, been found at *Sibitschiudi Suz*.

WALL EYED, (of uncertain etymology.) Applied to a horse—not perfectly white or cream-coloured—which has a white iris. (Prov.) Having eyes with too much white. In the north of England persons having the white of the eye very large and to one side, are so called. Applied, also, to an eye with little or no sight, "all white like a plastered wall."

WALLFLOWER, (probably *Gaulflower*), *Cheiranthus cheiri*.

WALLRUE, *Asplenium ruta muraria*.

WALLTEETH, Molar teeth.

WALNUT, (probably *Gaulnut*), *Juglans regia*—w. Black, *Juglans regia*—w. White, *Juglans cinerea*.

WALTHER, DUCT OF, see Sublingual gland.

WALTHERIA DOURADINHA, (after A. F. Walther, medical professor at Leipsic.) A plant of Brazil, *Order*, Sterculiaceæ, which is used there in diseases of the chest, and in venereal affections. It is a demulcent.

WALTON, MINERAL WATERS OF. A mineral spring near Tewkesbury, in Gloucestershire, England. It is an acidulous chalybeate.

WAME, (womb.) Belly, Uterus.

WAMPEE, *Peltandra Virginica*.

WANG, ([Sax. *þang*, *þeng*, 'the cheek,') Gena.

WANG TEETH, Molar teeth.

WANT, (from past participle of Sax. *þanian*, 'to wane,' 'to fall away;') 'need of that which is necessary.' (F.) *Besoin*. This word is sometimes used to express the imperious necessity which impels us to take food, drink, &c. (See Hunger and Thirst.) In a more general sense, it means the impressions transmitted by the organs to the brain, when it is necessary that they should act: these wants are called *instinctive desires* and *internal sensations* or *stimuli*. Such are the desires or wants to eat, drink, void the urine, breathe (*besoin de respirer*), &c.

WANT, in the sense of indigence, (L.) *Egestas*, *Indigentia*, is the cause of a number of diseases.

WAPS, Wasp.

WARBURG'S FEVER DROPS, see Bebeern.

WARD'S ESSENCE FOR THE HEADACH, Linimentum camphoræ compositum—w. White Drops, see Hydrargyri nitras.

WARE, see Pila marina.

WARK, ([Sax. *þærce*), Pain.

WARMBRUNN, MINERAL WATERS OF. Much frequented springs in Silesia, about a league from Hirschberg. They are thermal, 95° to 100° Fah., and are chiefly used as baths. The impregnating materials are carbonate of soda, sulphate of soda, sulphate of lime, chloride of sodium, carbonate of magnesia, and sulphohydric acid gas.

WARM SPRINGS, see Georgia, Mineral waters of, and Virginia, Mineral waters of.

WARNER CANADENSIS, (after Mr. Richard Warner, of Woodford, England,) *Hydrastis Canadensis*.

WARRENTON SPRINGS, see Virginia, mineral waters of.

WART, (Sax. *þeart*), *Verruca*—w. Corneous, Cornu—w. Moist, *Acrothymion*.

WARTY, *Verrucose*, *Verrucosus*, *Verrucosus*, (F.) *Verruquenz*. Full of warts. Resembling, relating, or belonging to warts.

WARTY TUMOUR OF CICATRICES. A name given by Mr. Howship to the tumour, which sometimes appears on an old scar, many years, perhaps, after the injury that caused it. The ulcer, formed under such circumstances, has been called the *warty ulcer of Marjolin*—having been well described by that surgeon.

WASH, BLACK, see Hydrargyri submurias—w. Phagedenic, see Hydrargyri submurias—w. Preventive, Lotion, Hannay's—w. Red, Lotion rubra, and Lotion, camphorated, of Bates—w. White, Liqueur plumbi subacetatis dilutus—w. Yellow, see Hydrargyri oxymurias.

WASHERWOMAN'S SCALL, Psoriasis diffusa.

WASHITAW SPRINGS, see Arkansas, mineral waters of.

WASP, *Waps*, ([Sax. *þæpp*, *þæppr*), (Prov.) *Whamp*, *Ves'pa*, *Sphæx*, (F.) *Guêpe*. A genus of gregarious insects, like the bee and the ant. They are armed with a sting, which pours a poisonous fluid into the puncture made by it. (See Poisons, Table of.) The best application is the sp. ammoniæ aromaticus, or some preparation of ammonia.

WASSERBURG, MINERAL WATERS OF. Wasserburg is a town of Bavaria, seated on the Inn, 28 miles east of Munich. The water, called, also, *Aqua Agatii*, contains carbonic acid; carbonates of lime and magnesia; sulphates of lime and magnesia; chloride of sodium, carbonate of soda and oxide of iron.

WASSERKUR (G.), 'water cure,' Hydro-sudotherapeia.

WASTING, ([Sax. *þertan*, 'to waste,') Consumption, Tabes.

WASSERSCHLAG (G.), 'water stroke,' see Hydrocephalus internus.

WATER, ([Sax. *þæter*, [G.] *Wasser*), *Aqua*, *Hy'dor*, (F.) *Eau*, *Protoc'hde* of hydrogen. Water was at one time regarded as a simple element. It is now known to consist of eight parts by weight of oxygen, and one of hydrogen. It freezes at 32°, and boils at 212° of Fahrenheit; and is taken as the unit in all tables of specific gravity of solid and liquid substances. Water, as we meet with it, is very different in its state of purity; generally, it contains small quantities of saline matter; at other times, as in seawater and mineral waters, it is largely impregnated. The characters of water, as met with under ordinary circumstances, are well compared by Celsus, "*Aqua levissima pluvialis est; deinde fontana; tum ex flumine; tum ex puteo; post huc ex nive, aut glacie; gravior his ex lacu; gravissima ex palude.*" Pure water—*Aqua p'u'ra*—is one of the best diluents that can be used. When cold, and especially when iced, it is one of the most useful refrigerants; and, in the form of the cold, tepid, and warm bath, it is serviceable in numerous diseases.

WATER, Urine—w. of Aix-la-Chapelle, see Aix-la-Chapelle—w. Allflower, Urine, cows'—w. Alum, Bates's, Liqueur aluminis compositus—w. of Ammonia, stronger, see Liqueur ammoniæ—w. of carbonate of Ammonia, Liqueur ammoniæ subcarbonatis—w. Angel, Angel water—w. Anise, see Pimpinella anisum—w. Apple, Apple tree—w. Barley, Decoctum hordei—w. Bath, see Bath, water.

WATER BED, *Hydrostatic bed*. A bed, contrived by Dr. Arnott, which, in consequence of its being placed on water, is well adapted for the bedridden.

WATER, BINELLI, *Aqua Binellii*.

WATER OF BITTER ALMONDS, *Aqua Amygdalarum concentrata*.

WATER BLEBS, Pemphigus—w. Brash, Pyrosis.

WATER, BROCCHERI, *Aqua Brocccherii*—w. Camphor, Mistura Camphoræ.

WATER CANKER, Cancer aquaticus.

WATER, CARAWAY, see Carum carui—w. Caraway, strong, Spiritus carui—w. Carmelita, see Melissa—w. of Cassia, see Laurus Cassia.

WATER CASTER, see Uromantia.

WATER, CHERRY LAUREL, see *Prunus lauro-cerasus*—w. Chicken, Chickentea—w. Chlorine, see Chlorine—w. Cinnamon, *Aqua cinnamomi*—w. Cinnamon, strong, *Spiritus cinnamomi*—w. Cologne, *Eau de Cologne*.

WATER, CREASOTE, see Creasote.

WATER CRESS, *Sisymbrium nasturtium*—w. Cure, *Hydrosudotherapia*.

WATER, DILL, see *Anethum graveolens*.

WATER, DISTILLED, *A'qua destilla'ta*, *H'o'ly Water*, (F.) *Eau distillée*, *Hydrolat*, is generally ordered in extemporaneous prescriptions, but is rarely used. In some cases, however, it is absolutely necessary;—the water, containing any saline impregnation—hard water, for example—decomposing certain substances. Distilled water has been recommended as a diet-drink in cancerous affections; and it is used in the religious ceremonies of the Catholic church. In the Pharmacopœia of the United States, it is directed to be made as follows:—Take of water, ten gallons. First distil two pints, and throw them away; then distil eight gallons. Keep the distilled water in glass bottles.

Distilled waters, (F.) *Hydrolats*, consist of common water, charged by distillation with the volatile principles of medicinal agents. See *Aqua destillata*.

WATERDOCTOR, see Uromantia.

WATERDRESS'ING. The treatment of wounds and ulcers by the application of water. It generally consists in dipping folds of lint in water, and placing them on the part:—the whole being covered with oiled silk, elastic gum, or spongopiline. *Water strap'ping* or *wet strap'ping* means the treatment of ulcers, and certain cutaneous affections of the extremities, &c., with strips of linen or cotton saturated with water.

WATER, ELDER, see *Sambucus*—w. Epileptic, of Langius, *Langii aqua epileptica*—w. Fennel, *Aqua fœniculi*.

WATER FLAG, YELLOW, *Iris pseudacorus*.

WATER, FLORIDA, —a celebrated perfume, —may be made of *Ol. Bergami*, f3iv; *Tinct. benzoin. comp.* f3ij; *Alcohol. cong.*

WATERFRIGHT, *Hydrophobia*.

WATER, GOULARD, *Liquor plumbi subacetatis dilutus*.

WATER GRUEL, see Avena, and Grits.

WATER, HARD. Common water, which decomposes and curdles soap; in contradistinction to *soft water*, which is readily miscible with soap. The hardness of water depends upon the presence in it of earthy salts, the most common of which is sulphate of lime.

WATER IN THE HEAD, *Hydrocephalus*—w. Hell, see Toddy.

WATER, HOLY, Water, distilled—w. Honey, see Mel.

WATER, HUNGARY, *Spiritus rosmarini*.

WATER JAGS, see Variicella.

WATER, LAKE, *A'qua ex lacu*, (F.) *Eau de lac*, is apt to be impure, from decayed animal and vegetable matters. A great deal will, however, depend upon the magnitude of the collection, and the degree of stagnation.

WATER, LAUREL, see *Prunus laurocerasus*—w. Lavender, *Spiritus lavandulæ*—w. Lead, *Liquor plumbi subacetatis dilutus*—w. Lemonpeel, see Lemonpeel Tea—w. Lime, *Liquor calcis*—w. Lime, compound, *Liquor calcis compositus*—w. Magnesia, aerated, *Magnesia, fluid*—w. Magnesia, carbonated, *Magnesia, fluid*.

WATER, MARSH, *A'qua paludo'sa*, (F.) *Eau de Marais*, is the most impure of all. It is generally more stagnant, and of course more loaded with decomposing animal and vegetable matters.

WATER, MEDICATED, see *Aqua medicatæ*.

WATER, MIN'ERAL, *A'qua mineralis seu salu'bris seu medica'ta seu sote'ria*, *Fons salu'taris seu medica'tus seu sote'rius*, (F.) *Eau minérale ou médicinale naturelle*. Water, holding in solution different saline and gaseous substances in sufficient quantity to be possessed of medicinal properties, or of a temperature different from that of the ordinary springs of the country.

Various mineral waters are referred to under their appropriate heads; and a table of the fixed constituents, which enter into the composition of some of the more celebrated springs abroad, is given by Dr. Pereira, in his "Elements of Materia Medica and Therapeutics." There is, also, a "Table of some of the principal Mineral Waters," at home and abroad, in the "Author's General Therapeutics and Materia Medica," Vol. 2, p. 460, 6th edit., Philadelphia, 1857.

Mineral waters may be divided into four classes:—1. *Ga'zeous or Acid'u'ous*. 2. *Chalybeate*. 3. *Saline*. 4. *Sulphu'reous*. These may be *ther'mal*, or *cold*,—*nat'ural*, or *artific'ial*. Many of these divisions, however, run into each other; some of the substances, which they hold dissolved, belonging to one class as well as to others.

1. *Ga'zeous, Acid'u'ous or Carbonated Min'eral Waters*, *So'da or Beer Springs*, *A'qua mineral'es acid'ule*, (F.) *Eaux minérales gazeuses ou acidules*. The waters referred to this class are those that contain carbonic acid gas in such quantity as to communicate to it certain sensible qualities. Waters impregnated with free carbonic acid, sparkle when drawn from the spring, or when poured into a glass. They have a sharp, acidulous taste; but become rapid from exposure to the air. Along with the carbonic acid, there are generally present portions of saline, earthy or metallic matter, chiefly carbonates of lime, magnesia, and iron. Waters, highly impregnated with carbonic acid gas, are grateful to the stomach; increase the appetite, and are diuretic; hence, their utility in dyspepsia, hypochondriasis, and gout. Their properties are, of course, modified by the saline matter that may be also contained in them. The most celebrated amongst the acidulous waters, are those of Bar, Chateldon, St. Myon, Mont d'Or, Langeac, Seltzer, Schlangenbad, Sultzmat, Pyrmont, Spa, Carlsbad, Cheltenham, Scarborough, Saratoga, Ballston, and the Sweet Springs of Virginia.

2. *Chalybeate or Ferru'ginous Min'eral Waters*, *A'qua mineral'es ferrugine'as seu Chalybeat'es seu Martia'les seu Ferro'sæ*, *Chalybop'gæ*, *Chalybokre'næ*, *Siderop'gæ*, *Siderokre'næ*, (F.) *Eaux minérales ferrugineuses*, contain iron—sulphate, chloride or carbonate, generally the latter—held in solution by an excess of acid. Chalybeate waters have a peculiar styptic taste. They are transparent when taken from the spring; but, when exposed for some time to the air, a pellicle forms on the surface, and a deposit of the iron takes place. Chalybeate waters are used as tonics in debility of all kinds; in all cases, in fact, where iron is considered to be indicated. They are the most numerous of all the classes of mineral waters. In this dictionary, those only are specified which have some celebrity; yet there are about one hundred; more than sixty of these being saline, and forty sulphureous. The most celebrated chalybeates are those of Tunbridge, Scarborough, Spa, Bussang, Forges, Vichy, Pyrmont, Passy, Provins, and Vals.

3. *Saline Min'eral Waters*, *A'qua mineral'es salinæ*, *Am'na alcaliza'ta* (Paracelsus), *Hydrat'æ*, (F.) *Eaux minérales salines*. Waters, holding in solution different saline substances in greater quantity than the acidulous waters. They differ in properties, according to the salts con-

tained in them. The salts usually present are sulphates, chlorides, and carbonates; and the bases, with which the acids forming these are combined, are soda, magnesia, and lime. Saline mineral waters are usually aperient; and the most noted are those of Seltzer, Sedlitz, Balaruc, Bourbonne-les-Bains, Baden, Epsom, Cheltenham, &c.

To this class may also be added *Sea water*.

4. *Sulphureous Mineral Waters*, *A'qua mineral'ia sulphurea*; when warm, termed *Theiother'ma*, *Thiopo'ga*, *Thiather'ma*, *Fon'tes sulphu'ri cal'id'i*, (F.) *Eaux minérales sulfureuses ou sulfurées ou sulfureuses ou hépatiques*, &c. These waters owe their distinguishing character to an impregnation of sulphuretted hydrogen, and are at once recognised by their peculiar fetid smell, resembling that of rotten eggs. They usually contain saline substances, which modify their powers. From the action of the sulphuretted hydrogen, they are useful in cutaneous affections; and, from the combined action of this and the saline matter, they are valuable agents in diseases of the digestive organs. They are also employed in cutaneous eruptions; and the warm sulphur baths have been especially celebrated in such cases, as well as in rheumatic affections. The most renowned sulphureous waters are those of Baréges, Bagnères-de-Luchon, Cauterets, Bagnès, Aix in Savoy, Aix-la-chapelle, Enghien, Harrogate, Moffat, Virginia Springs, &c.

Some springs, as those of Bath, Buxton, Dax, Matlock, Warm and Hot Springs of Virginia, &c., are almost pure *Ther'mal Mineral Waters*. They are valuable agents in rheumatic affections; the warmth being equable during the whole period the individual is immersed in them; which cannot be the case in the artificial thermal bath.

WATER, MINERAL, Acidulous water, simple.

WATERS, MINERAL, ARTIFICIAL, (F.) *Eaux minérales artificielles ou factices*. These are imitations of the natural; and some of them—as the factitious Cheltenham water, and Sedlitz water—answer the purpose of the natural water tolerably well. The acidulous and chalybeate waters are, however, most easily imitated.

WATER, NUTMEG, *Spiritus myristice*—w. Orange flower, see *Citrus aurantium*—w. Oxygenated, Hydrogen, peroxide of.

WATER PANG, *Pyrosis*—w. Parsnep, creeping, Sium.

WATER, PENNYROYAL, SPIRITUOUS, *Spiritus pulegi*—w. Peppermint, *Aqua menthae piperitæ*—w. Pimento, see *Myrtus pimenta*.

WATER POX, see *Varicella*.

WATER, PROTOXIDE OF NITROGEN, *Aqua nitrogenii protoxydi*.

WATER PURPLE, *Veronica beccabunga*—w. Qualm, *Pyrosis*.

WATER, RAIN, *A'qua pluvi'alis seu plu'via seu im'brium*, (F.) *Eau de pluie*, when collected at a distance from houses or other elevated objects, is the purest natural water, and has the least specific gravity. The only bodies which it usually holds in solution are carbonic acid, and minute traces of carbonate of lime and chloride of calcium.

WATER, RICE, see *Oryza*.

WATER, RIVER, *A'qua fluvial'is*, is derived from the conflux of numerous springs and rain water. It is, generally, pretty pure.

WATER, ROSE, *Aqua rosæ*.

WATER, SEA, *A'qua Mari'na*, *Alma*, *Hu'mor Do'ridis*, (F.) *Eau de Mer*, *Eau salée*. This contains chlorides of sodium, magnesium, and calcium, and sulphate of magnesia. It is cathartic, and forms the usual glyster at sea. It makes an

excellent tonic bath; superior in most cases—especially in serofula—to the freshwater bath.

WATER, SEARLE'S PATENT OXYGENOUS AERATED, *Aqua Nitrogenii protoxydi*—w. Seydewitz, Sedlitz water.

WATER SHIELD, *Brasenia hypopeltis*.

WATER, SNOW, *A'qua niva'is*, (F.) *Eau de neige*, has usually been deemed unwholesome. It exactly resembles rain water in composition, and is equally salubrious.

WATER, SODA, Acidulous water, simple—w. Soft, see Water, hard—w. Spearmint, *Aqua menthae viridis*.

WATER, SPRING, *A'qua fonta'na*, *Hydrope'gæ*, (F.) *Eau de fontaine*, contains, in addition to the substances detected in rain water, more or less sulphate of lime. When this is to such an extent as to curdle soap, the water is said to be *hard*; if not, *soft*. Hard water is, of course, inferior to soft, for domestic and medicinal purposes.

WATER STRAPPING, see Water dressing—w. Stroke, see *Hydrocephalus internus*.

WATER, STYPTIC, SYDENHAM'S, *Solutio sulphatis cupri composita*—w. Tar, see *Pinus sylvestris*.

WATER TARGET, *Brasenia hypopeltis*.

WATER, TOAST, see Toast water.

WATER, WELL, *A'qua putea'lis seu putea'na seu ex puteo*, (F.) *Eau de puit ou de source*, is the same as spring water, but liable to impregnation, owing to the land springs filtering into the wells, and conveying impurities into them.

WATERS, THE, Liqueur Amnii—w. Bag of, see Liqueur Amnii—w. Discharged or broken, *Profusio aquæ*—w. Distilled, *Aqua destillatæ*—w. First, *Primitiæ*—w. Green, see *Lochia*.

WATTWEILER, MINERAL WATERS OF. Wattweiler is a small town in the department of Haut-Rhin, at the foot of the Vosges. The waters are acidulous chalybeates, and contain carbonates of iron, lime, and soda, chloride of sodium, and free carbonic acid.

WAX, ([Sax.] *Þæxe*,) *Cera*—w. Ear, *Cerumen*.

WAX, JAPAN. Obtained from *Rhus succedanea* of Japan. It is of a medium quality between beeswax and the ordinary vegetable tallow.

WAX, MYRTLE, *Bayberry tallow*. A wax obtained from the fruit of *Myrica cerifera*, *Wax myrtle*, *Candleberry myrtle*; an indigenous plant, which yields the bayberry bark, possessing tonic, stimulant and astringent properties. It has been prescribed in dysentery.

WAX, YELLOW AND WHITE, *Cera flava et alba*.

WAXING KER'NELS, *Wax'enkernels*, *Crescentia*, (F.) *Croissances*, *Croissants*, *Glandes de croissance*. A popular term for small tumours, formed by an enlargement of the lymphatic glands—in the groins of children particularly. They have been looked upon as connected with the growth of the body,—hence their name.

WAY, ([Sax.] *Þæg*,) *O'dos*, *Via*, *Passage*, (F.) *Voie*. A name given to different canals.

WAYBREAD, ([Sax.] *Þæg bræde*,) *Plantago major*.

WAYME, Womb, Belly, Uterus.

WAYS, DIGESTIVE, *Digestive Passages*, *Primæ Viæ*, (F.) *Voies digestives*, *Premières voies*, are the series of hollow organs of digestion, composed of mouth, œsophagus, stomach, and small and large intestines. The term is often restricted to the first three of these.

WAYS, SECOND, *Second Passages*, *Secundæ viæ*, (F.) *Secondes voies*, comprise the chyliferous, lymphatic and blood-vessels.

WEAKSIGHTEDNESS, *Asthenopia*.

WEAL, ([Sax.] *palan*, 'marks of stripes,') *Wheal*.

WEAM, (Womb,) Belly, Uterus.

WEAM ILL, Colic.

WEAN'ING, (from [Saxon] *penan*, 'to wean;')

Abductio, *Apogalactismus*, (Prov.) *Sp'aning*, (F.) *Sévrage*. The act of depriving the infant, permanently, of the breast, in order to feed it on other and more solid nourishment.

WEANING BRASH, see Brash, weaning.

WEASAND, ([Sax.] *pærand*), Trachea.

WEARING, (from [Sax.] *penan*, 'to wear,') Consumption.

WEATHERCOCKS, Impatiens.

WEB, ([Sax.] *web*), Caligo, Pterygium—w. Muscular, Panniculus carnosus.

WEBEYE, Caligo, Pterygium.

WEBBED FINGERS, Dactylon.

WEBERIAN CORPUSCLES, (after Weber, a German anatomist,) see Ejaculatory Ducts—w. Organ, see Ejaculatory Ducts.

WED'GED, ([Sax.] *pæc*, 'a wedge,') *Inclu'sus*, *Inhærens*, (F.) *Enclavé*. The head of the foetus is said to be *wedged*, *locked* or *impacted* in the pelvis, when it remains fixed, notwithstanding the uterine efforts.

The condition is called *Incumectio* seu *Inclavatio foetus*, *Spheno'sis*, (F.) *Enclavement*. See Paramphosis.

WEEBO, Senecio Jacobæa.

WEED, (from [Sax.] *wean*, 'to rage,') see Mastitis.

WEED, ARROW, ([Sax.] *weow*), *Sagittaria variabilis*—w. Beetle, *Galaxaphylla*—w. Breast, *Saururus cernuus*—w. in the Breast, see Mastitis—w. Butter, *Erigeron Canadense*—w. Cancer, *Goodyera pubescens*—w. Dyer's, *Reseda luteola*—w. Field, *Anthemis cotula*—w. Horse, *Erigeron Canadense*—w. Horsefly, *Sophora tinctoria*—w. Milk, *Apoecynum androsæmifolium*—w. Phthisis, *Ludwigia palustris*—w. Pride, *Erigeron Canadense*—w. Red, *Phytolacca decandra*—w. Silver, *Potentilla anserina*—w. Soldiers', *Matico*—w. Thimble, *Rudbeckia laciniata*—w. Wind, *Asclepias tuberosa*.

WEEPING, (from [Sax.] *weopan*, 'to weep,') *Fletus*—w. Tear, Tear.

WEESEL, Trachea.

WEID, (Weed,) see Mastitis.

WEIGHT, (from [Sax.] *pægan*, 'to weigh;') (F.) *Pesanteur*, see Stature. A sensation of heaviness or pressure over the whole body, or over a part—the stomach or head, for example.

WEIGHTS AND MEASURES, *Pondera et Mensuræ*, (F.) *Poids et Mesures*. The importance of possessing a uniform system of weights and measures has impressed the scientific of all countries, and numerous endeavours have been made to accomplish the object. It is, however, a matter of considerable difficulty, and one not likely to be attained. The new French measures are upon decidedly the best footing, but they are not yet generally adopted. The greatest diversity, indeed, prevails in the measures, both of weight and capacity. Some of the following tables will show, that every subdivision of the pound, as well as the pound itself, differs in England and in France.

WEIGHTS.

Troy Weights, as used by the British Apothecaries, —Pondus pharmaceuticum seu medicinale.

The pound lb.	} contains	12 ounces.
The ounce		8 drachms.
The drachm		3 scruples.
The scruple		20 grains.
The grain gr.		

These, and the signs by which they are denoted,

have been the same in all the British Pharmacopœias, as well as in the American.

lb.	Oz.	Drachms.	Scrups.	Grs.
1 =	12 =	96 =	288 =	5760
	1 =	8 =	24 =	480
		1 =	3 =	60
			1 =	20

Poids de Marc.

The pound	} contains	16 ounces.
The ounce		8 drachms.
The drachm		3 scruples.
The scruple		24 grains.
The grain		

Avoirdupois.

Pound.	Ounces.	Drachms.	Troy grains.
lb. 1. =	16 =	256 =	7000
	1 =	16 =	437.5
		1 =	27.34375

The avoirdupois drachm is sometimes divided into three scruples, and the scruple into ten grains. The pound of 7680 grains avoirdupois=7000 grains troy, and hence 1 grain troy = .97 grain avoirdupois.

The *Poids de Marc* is that employed by the French Pharmacians, when the new weights are not. The Avoirdupois has been used by the Dublin College.

To avoid the danger of mistakes by confounding the troy and the avoirdupois pounds, the term *pound* has been discontinued in the formulæ of the Ph. U. S. 1863—the desired weight being expressed in ounces, each containing 480 grains. The ounce is always written *troyounce*; and the *drachm* and *scruple* are disused, their equivalents being expressed in grains. On the other hand, the General Medical Council of Great Britain resolved (1863) that "the weights used in the British Pharmacopœia shall be the Imperial or avoirdupois pound, ounce, and grain; and that the terms *drachm* and *scruple*, as designating specific weights, shall be discontinued." The following are the weights and measures of that Pharmacopœia with their symbols.

WEIGHTS.

The Avoirdupois pound =	16 oz. =	7000 grs.
	1 oz. =	437.5 grs.
	1 gr. =	1 gr.

MEASURES.

C. 1 gallon	=	8 pints	weighing 10 pounds.
O. 1 pint	=	20 fluid ounces	" 1 1/4 "
fl. oz. 1 fluid ounce	=	8 fluid drachms	" 437.5 grs.
fl. dr. 1 fluid drachm	=	60 minims	" 54.68 "
℥. 1 minim	=		" .91 "

The gallon holds 10 1/2 avoirdupois of distilled water, at a temperature of 60° Fahrenheit.

The measures in the Ph. U. S., 1863, have the same names as in the British; but their value is different; the pint weighing 16 oz. 291.2 grs. avoirdupois, and the fluidounce 455.7 grs. In the formulæ, the Acids and the Oils are ordered by weight; other liquids by measure. The gallon is not used, that measure being always expressed in pints. In Continental Europe all medicines are weighed, and the gramme is becoming universal—used.

The following tables exhibit the relative value of the old French and English weights:

Poids de Marc.	Troy. Wt.	Avoird.	Troy gra.
1 lb. (livre) =	1.31265 lb.	1.080143 lb.	7561
1 oz. (once) =	.984504 oz.	1.080143 oz.	472.5625
1 dr. (gros.) =	.954504 dr.		59.0703125
1 gr. =			.820421

<i>Troy.</i>	<i>Poids de Marc.</i>	<i>French grains.</i>
1 pound =	0.76180 lb.	= 7561
1 ounce =	1.01574 once	= 585.083
1 drachm =	1.01574 gros	= 73.135
1 grain		= 1.219
<i>Avoirdupois.</i>	<i>Poids de Marc.</i>	<i>French grains.</i>
1 pound =	0.925803 lb.	= 8532.3
1 ounce =	0.925803 once	= 533.27

To convert	French grains into Troy grains, divide by	1.2189
"	Troy grains into French grains, multiply by	
"	French ounces into Troy ounces, divide by	1.015734
"	Troy ounces into French ounces, multiply by	
"	French pounds (<i>poids de marc</i>) into Troy pounds, multiply by	1.31268
"	Troy pounds into French pounds, divide by	

<i>Troy grain.</i>	<i>Poids de Marc or French grain.</i>
1	= 1.219
2	= 2.438
3	= 3.657
4	= 4.876
5	= 6.095
6	= 7.314
7	= 8.533
8	= 9.752
9	= 10.971

<i>French grain.</i>	<i>Troy grain.</i>
1	= 0.8203
2	= 1.6406
3	= 2.4609
4	= 3.2812
5	= 4.1015
6	= 4.9218
7	= 5.7421
8	= 6.5624
9	= 7.3827

New French Decimal or Metrical Weights.

	<i>Troy grains.</i>	
Milligramme =	.0154	
Centigramme =	.1543	
Decigramme =	1.5434	
Gramme =	15.4340	
Decagramme =	154.3402	= 0 0 2 34.4
Hectogramme =	1543.4023	= 0 3 1 44.4
Kilogramme =	15434.0234	= 2 8 1 24
Myriagramme =	154340.2344	= 26 9 6 0

Comparative Value of the French Medicinal Pound and that of other Places.

	<i>lbs. oz. dr. gr.</i>
Germany.....	0 15 4 48
Amsterdam.....	1 00 0 42
Belgium.....	1 00 0 42
Berne.....	1 00 0 00
Copenhagen.....	0 15 3 20 $\frac{1}{4}$
Florence.....	0 11 0 50
Genoa.....	0 10 5 60
Geneva.....	1 00 0 18
Hamburg.....	0 15 2 15
Lisbon.....	0 15 7 68
London.....	0 12 3 12
Madrid.....	0 15 0 16
Milan.....	0 9 3 00
Monaco.....	0 15 2 23
Naples.....	0 10 7 54
Paris.....	1 00 0 00
Rome.....	0 11 0 50
Stockholm.....	0 13 7 8
United States.....	0 12 3 12
Warsaw.....	1 10 4 24
Venice.....	0 8 6 00
Vienna.....	1 2 2 32

MEASURES OF CAPACITY.

English Measure of Fluids.

The gallon (congius).	0 $\frac{7}{8}$ } contains {	8 pints.
The pint (octarius) ..		16 fluidounces.
The fluidounce.....		8 fluidrachms.
The fluidrachm.....		60 minims.
The minim.....	1	

Proportions of the Apothecaries' and Wine Gallon.

<i>Gal. Pints. Fluidoz. Fluidr. Minims. Cub. inches</i>
1 = 8 = 128 = 1024 = 61440 = 231
1 = 16 = 128 = 7680 = 2887 $\frac{1}{2}$
1 = 8 = 480 = 18047
1 = 60 = 2256

Imperial Measure, adopted by the London College in their Pharmacopæia of 1836.

<i>Gallon. Pints. Fluidounces. Fluidr. Minims.</i>
1 = 8 = 160 = 1280 = 76800
1 = 20 = 160 = 9600
1 = 8 = 480
1 = 60

Comparative Value of the Proportions of the Wine and Imperial Gallons.

<i>Wine.</i>		<i>Imperial.</i>			
	<i>Pints.</i>	<i>Fluidoz.</i>	<i>Fluidr.</i>	<i>Minims.</i>	
1 gallon	= 6	13	2	23	
1 pint	=	16	5	18	
1 fluidounce	=	1	0	20	
1 fluidrachm	=		1	2½	

<i>Imperial.</i>		<i>Wine.</i>			
	<i>Gallon.</i>	<i>Pint.</i>	<i>Fluidoz.</i>	<i>Fluidr.</i>	<i>Minims.</i>
1 gallon	= 1	1	9	5	8
1 pint		1	3	1	38
1 fluidounce				7	41
1 fluidrachm					58

French Measures of Capacity.

<i>English cubic inches.</i>	<i>Wine Measure.</i>
Millilitre =	.061028 = 16.2318 minims.
Centilitre =	.610280 = 2.7053 fluidrachms.
Decilitre =	6.102800 = 3.3516 fluidounces.
Litre =	61.028000 = 2.1135 pints.
Decalitre =	610.280000 = 2.6419 gallons.
Hectolitre =	6102.800000
Kilolitre =	61028.000000
Myrialitre =	610280.000000

Approximate comparison between the ancient French Measures of Capacity, and the new, and conversely.

	<i>Grammes.</i>	<i>Litres.</i>	<i>Decilitres.</i>	<i>Centilitres.</i>
1 poisson =	125 or 0	1	2	
1 demi-setier =	250 or 0	2	5	
1 chopine =	500 or 0	5	0	
1 pinte =	1000 or 1	0	0	

<i>Grammes. lbs. oz. dr. gr.</i>
1 centilitre = 10 or 0 0 2 $\frac{1}{2}$ 00
1 decilitre = 100 or 0 3 2 00
1 litre = 1000 or 2 0 3 36 = 2.113 Eng. wine

Measures of Length.

	<i>Inches.</i>
1 line, the 12th part of an inch.	1.000
3 barleycorns	
A palm or hand's breadth (Scripture measure).....	3.648
A hand (horse measure).....	4.000
A span (Scripture measure).....	10.944
A foot.....	12.000
A cubit (Scripture measure for common purposes).....	18.000
A cubit (Scripture measure for sacred purposes).....	21.888
A Flemish ell.....	27.000
A yard.....	3 ft 00
An English ell.....	3 " 09
A fathom or twice.....	6 " 00

New French Measures of Length.

<i>English inches.</i>	<i>yd.</i>	<i>ft.</i>	<i>inch.</i>
Millimètre 0.039			
Centimètre 0.393			
Decimètre 3.937			
Mètre 39.371 =	1	0	3.37

VALUES OF THE GRECIAN, ROMAN, AND ARABIAN
WEIGHTS AND MEASURES IN POIDS DE MARC.

1. *Weights of the Ancient Greeks.*

	lbs.	oz.	dr.	gr.
The talent (ταλαντον).....	54	2	5	24
The mina (μινᾶ).....	14	3	40	
The drachm (δραχμη).....	1	11		
The obolus (οβολος).....			13	
The ceration (κερατιον).....			4	
The chalcus (χαλκος).....			2	
The septem (σεπτον).....			1/2	

2. *Weights of the Ancient Romans.*

	oz.	dr.	gr.
The pound (libra).....	10	6	48
The ounce (uncia).....		7	16
The duella.....		2	29
The sicilicus.....		1	58
The sextula.....		1	14
The consular denier, denarius.....		1	2
The imperial denier or drachm, drachma.....			65
The victorialis.....			37
The scriptulum or scruple.....			21
The obolus.....			11
The siliqua.....			4

The Greeks divided their obolus into *chalcæ* and *lepta*: some divided it into 6 *chalcæ*, and each *chalcus* into 7 *lepta*; others into 8 *chalcæ*, and every *chalcus* into 8 *lepta* or *minuta*.

Dr. Milligán, in his edition of Celsus, gives the following table, exhibiting the Troy weight of measures of capacity and of weight in use amongst the Romans.

	Urna.	Libra.	Uncia.	Denarius.	Scrupulus.	Sextans.	Chalcus.	Grana.
Amphora.....	2	= 80	= 960	= 6720	= 20160	= 40320	= 403200	= 420480
Urna.....	1	= 40	= 480	= 3360	= 10080	= 26160	= 210600	= 210240
Congius.....	1/4	= 10	= 120	= 840	= 2540	= 5040	= 50400	= 52920
Sextarius.....		1 1/2	= 20	= 140	= 420	= 840	= 8400	= 8760
Libra.....		1	= 12	= 84	= 252	= 504	= 5040	= 5256
Hemina.....			8 4/7	= 60	= 180	= 360	= 3600	= 3759
Acetabulum.....			2 1/7	= 15	= 45	= 90	= 900	= 939
Sesqui-cyathus.....			2 1/7	= 15	= 45	= 90	= 900	= 939
Cyathus.....			1 3/7	= 10	= 30	= 60	= 600	= 626
Sesuncia.....			1 1/2	= 10 1/2	= 31 1/2	= 63	= 630	= 657
Uncia.....			1	= 7	= 21	= 42	= 420	= 438
Coehleure.....				2 1/2	= 7 1/2	= 15	= 150	= 156
Drachma.....				1	= 3	= 6	= 60	= 62 4/7
Denarius.....				1	= 3	= 6	= 60	= 62 4/7
Scrupulus.....					1	= 2	= 20	= 20 2/3
Scrupulus dimidiatus.....					1/2	= 1	= 10	= 10 1/3
Obolus.....						1	= 10	= 10 1/3
Sextans.....						1	= 10	= 10 1/3
Chalcus.....							1	= 11 3/4

He gives also the following 'Carmen Mnemonicum,' which exhibits the analogies of the Roman and British imperial weights.

To the congius and gallon, each, ten pounds allow,

On the bushel and amphora eighty bestow;

Rome's pound, as in Troy weight, twelve ounces obtains,

But her ounce is Absoedupois, strictly, in grains:

Denarii drams, scruples scrupls define,

Septarius answers to our bottle of wine,

And the mina's a short pint—fourteen ounces in fine.

3. *Weights said to be of the Arabians, Modern Greeks, and Latins of the barbarous periods of the Middle Ages.*

	oz.	dr.	gr.
The alchemion.....	14	3	40
The mines or ominos.....	10	6	28
Suerus, aghien, adar, assatil.....		7	16
The great or royal nut.....		3	44
The sextarius, stater.....		3	44
The lesser nut.....		2	50
Alomanus.....		2	29
Ayruus, Alcoholus.....		2	14
The hazel nut, bendacate, holca, alchi, dar-chimi, atogochilos, ologinat, nabach.....		1	11
The acorn, lupine, Egyptian or Syrian bean, the bushil.....			42
The Alexandrian bean or tremessis.....			30
The Greek bean or gramme, the kermel, gornu, harni, grachus.....			21
The ring, cumulus, seminet, onoloch, onolosal.....			11
The danish.....			8
The kirat, alkilat, karat.....			4

1. *Greek Measures.*

	lbs.	oz.	dr.	gr.
The metretes (μετρητης).....	84	4	3	00
The chus, choa, congius (χους).....		7	0	2 66
The zestes (ζεστης).....		1	1	7 44
The cotyla (κοτυλη).....			8	7 58
The tetarton (τεταρτον).....			4	3 65
The oxybaphon (οξυβαφον).....			2	1 69
The cyathos (κυαθος).....			1	4 00
The comcha (κογχη).....				6 00
The mystron (μυστρον).....				3 00
The chama minor (χημη).....				2 00
The cochlarian (κοχλιαριον).....				1 00

2. *Roman Measures of Capacity.*

	lbs.	oz.	dr.	gr.
The amphora or cadus.....	56	2	7	24
The urna.....	28	1	3	48
The congius.....		7	0	2 66
The sextarius.....		1	4	7 44
The hemina.....			10	1 18
The quartarius.....			5	0 45
The acetabulum.....			2	4 23
The cyathus or small glass.....			1	5 30
The ligula or spoonful.....				3 24

3. *Measures said to be of the Arabian, Arabian, and Latin Physicians of the Middle Ages.*

	lbs.	oz.	dr.	gr.
The missohacs weighed.....	3	8	1	33
Aben, kirt, ejub, eberia, or Roman mina.....	1	6	0	00
The phial, hufilius, or hassilinus.....		10	1	18
The caliz or rejelat.....			6	0 44
The handful, pugilum, cornusum.....			3	2 68
The hassuf, aessuse, or anesime.....			2	4 20
The comos or coatus, alcantur or alumesi, briale cubus.....			1	5 34
The lesser bachates.....				5 66
The largest spoonful.....				4 44
A spoonful.....				1 52
The colanos or reclanarium.....				1 28
A small spoonful or flagerina, or cyanes.....				1 11
The smallest spoonful or fahatiei.....				42

A good section on weights and measures is contained in the edition of "The Seven Books of Paulus Aegineta," by Mr. Francis Adams, published by the Sydenham Society of London, vol. iii. p. 609, London, 1847.

Besides the weights and measures above mentioned, employed by the moderns, there are modes of estimating the quantities of substances by approximation. For example:—

A *glassful* or *cupful*, ([F.] *Verre*), is reckoned to contain 4 or 5 fluidounces.
 A *wineglassful*, $1\frac{1}{2}$ ounce or two fluidounces.
 A *tablespoonful*, ([F.] *Cuiller à bouche*), about half a fluidounce.
 A *coffee* or *dessertspoonful*, about 3 fluidrachms.
 A *teaspoonful*, a fluidrachm.
 A *handful*, *manipulus*, ([F.] *Poignée*), as much as can be held in the hand.
 The *Pugillus* ([F.] *Pincée*) is as much as can be held by the three fingers.

These last quantities are occasionally prescribed by the French practitioners. Their weights must, of course, vary, according to the article. The authors of the Parisian codex have valued them, as regards certain substances.

	French.	oz.	dr.
A <i>Manipulus</i> of barley weighs.....	3	$2\frac{1}{2}$	
“ linseed	1	4	
“ linseed meal	3	3	
“ dried mallow leaves.....	1	3	
“ dried cichory leaves.....	1	0	
“ flowers of the tilia.....	1	$2\frac{1}{2}$	
	dr.	gr.	
A <i>Pugillus</i> of chamomile flowers weighs.....	2		
“ arnica	1	48	
“ marshmallow.....	1	24	
“ mallow.....		50	
“ fennel seeds.....	1	60	
“ anised.....	1	12	

A *hen's egg*, newly laid, weighs about 2 ounces and 2 drachms: when deprived of its shell, 2 ounces. The *white* weighs 1 ounce, 2 drachms, and 57 grains: the *yolk*, 5 drachms and 15 grains.

WEILBACH, MINERAL WATERS OF. The village of Weilbach is in the plain between the Maine and the southern extremity of the Taunus hills. It has in its neighbourhood a cold sulphureous alkaline spring, the water of which is bottled and exported, although not of great efficacy. Its use is at times combined with that of the saline springs in Nassau, and of Baden-Baden, in various chronic abdominal and thoracic affections.

WEINTRAUBENKUR, Grape cure.

WELCOME TO OUR HOUSE, Euphorbia cyparissias.

WELD, *Reseda luteola*.

WELK, *Whelk*, (like *Wheal*.) An inequality; a protuberance; a cutaneous eruption of this character. Acne.

WELTER'S BITTER, Acid, carbazotic.

WEM, (Womb,) Belly, Uterus.

WEM'DING, MINERAL WATERS OF. These waters are situate about four leagues from Donauwörth, in Bavaria. They contain sulphuretted hydrogen; carbonates of lime, magnesia, and soda; sulphates of lime and magnesia; a little chloride of calcium, and oxide of iron; and are much used in asthenic diseases.

WEN, ([Sax.] *pen*), *Ln'pia*, (F.) *Loup*. A name given to circumscribed, indolent tumours, without inflammation or change of colour of the skin. They may be met with in almost every part of the body. Their magnitude, as well as nature, is very variable. The term is sometimes given to an encysted tumour, and to bronchocele.

WESTERN ISLANDS, Azores.

WEST INDIES, CLIMATE OF. The climate of the West Indies has long been regarded as favourable to the consumptive. In confirmed consumption, it appears, however, to be positively injurious. As a prophylactic for those predisposed to consumption, it is generally favourable. The nature of the patient's constitution, in regard to the effects of elevated temperature, has always to be taken into account. In diseases that are benefited by warm climates, the West Indies afford an appropriate residence. The islands in which the invalid can best obtain all the advantages of the climate, are Jamaica, Barbadoes, St. Vincents, Antigua, St. Kitts, and Santa Cruz.

WEST'S SPRING, see South Carolina, mineral waters of.

WET STRAPPING, see Water dressing.

WHAHOO, see Eucynismus.

WHAMP, Wasp.

WHARTON'S DUCT, (after Dr. Thomas Wharton, an English anatomist,) Ductus salivaris inferior.

WHEAL, ([Sax.] *Palan*, 'marks of stripes,') *Weal*, *Wale*, *Pom'phus*. A ridge, or elevation of the skin, produced by a rod or whip; or as if produced in that manner. Such elevations are seen in urticaria.

WHEAT, ([Sax.] *hpeate*), *Triticum*—w. Indian, Zea mays—w. Turkey, Zea mays.

WHEEZE, ([Sax.] *hpeoran*), (Prov.) *Re'zale*, (formed probably, like *whiz* and *whisper*, from the sound it expresses.) To breathe with difficulty and noise, as in asthma.

WHELK, (like *wheal*), Acne, Welk—w. Chin, Sycosis—w. Rosy, Gutta rosea.

WHETLEBONES, Vertebrae.

WHEY, ([Sax.] *hpæg*), Serum lactis.

WHEY CURE, (G.) *Molkenkur*, (F.) *Cure de petit lait*. In the Kursaal of many of the German watering places, a space is set apart for whey drinkers. The whey is a pleasant diluent, and the saccharine matter it contains may act as a eutrophic; but the main effect on the valetudinarian is probably produced by the new impressions made upon him by his visit to the watering place.

WHEY, MUSTARD, see Sinapis—w. Rennet, see Serum lactis—w. Sack, Winewhey—w. Tamarind, see Tamarindus—w. Vinegar, see Acetum—w. Wine, Winewhey.

WHICKFLAW, Whitlow.

WHIFFING MURMUR, see Murmur, respiratory.

WHIG, ([Sax.] *hpæg*), Buttermilk, Serum lactis.

WHIMBERRY *Vaccinium myrtillus*.

WHIRLING CHAIR, see Chair.

WHISKY, see Spirit.

WHISKY LIVER, Liver, nutmeg.

WHISPER, CAVERNOUS, see Cavernous.

WHISPERING, *Vox elandest'na*, (from [Sax.] *hpiþþian*, 'to whisper.') An example of onomatopœia. *Susurra'tio*, *Susurra'tion*. Articulation of the air sent through the vocal tube without any action of the glottis.

WHISPERING SOUFFLE, see Souffle.

WHISTLING, (a case of onomatopœia, from [Sax.] *hpiþþan*, 'to whistle,') *Sifflement*, see *Râle sibilant*.

WHITE BALL, *Cephalanthus occidentalis*—w. Blisters, *Rupia escharotica*.

WHITE, FLAKE, *Plumbi subcarbonas*.

WHITE LEAF, *Pyrola maculata*, *Spiraea tomentosa*—w. Leg, *Phlegmatia dolens*—w. Matter of the brain, see Cerebrum—w. Root, *Angelica lucida*, *Asclepias tuberosa*, *Ligusticum actæifolium*.

WHITE, SPANISH, Bismuth, subnitrate of.

WHITE SUBSTANCE OF SCHWANN, see Nerve fibre—w. Sulphur, see Virginia, mineral waters of—w. Swelling, Hydrarthrus—w. Swelling of

lying-in women, Phlegmatia dolens—w. Weed, Chrysanthemum leucanthemum—w. Wood, Liriodendron, Tilia Americana.

WHITENESS OF COMPLEXION, Paleness.

WHITES, THE, Leucorrhœa.

WHITLEYA STRAMONIFOLIA, (after Whitley, an English botanist,) Anisodus luridus.

WHITLOW, ([Sax.] *hþix*, 'white,' and *low*, 'a flame,') Paronychia.

WHOOPINGCOUGH, (from the attendant whoop,) Pertussis.

WHORLYWORT, Leptandria purpurea.

WHORTLEBERRY, Vaccinium myrtillus, Vaccinium vitis idæa—w. Bears', Arbutus uva ursi.

WHORTS, BLACK, Vaccinium myrtillus.

WHYTTS DISEASE, (after Dr. Robt. Whytt, of Edinburgh,) Hydrocephalus internus.

WIDOWWAIL, Cneorum tricoccum, Daphne Alpina.

WIESBADEN, MINERAL WATERS OF, see Wiesbaden.

WIESSAU, MINERAL WATERS OF. These waters are situate about four leagues from the convent of Waldsassen in Bavaria. They contain carbonic acid, carbonates of lime and magnesia, chlorides of calcium, magnesium, and aluminium, carbonate of soda, and much oxide of iron. The Bavarians consider them to resemble the waters of Pyrmont.

WILD, ([Sax.] *þilo*.) An epithet given to the countenance, when not in harmony with the condition of the individual, and indicating strong mental emotion;—a *wild look*, (F.) *Physiognomie égarée*.

WILD/BAD, MINERAL WATERS OF. Wildbad is a small town of Wurtemberg, three German miles from Baden-Baden. It is much frequented as a watering-place. The water is thermal, from 95° to 100° of Fah., and is much employed in cases in which thermal waters are indicated. It is used internally, a pint not containing more than a grain of solid ingredients, which consist of salts of soda and lime. The proportion of carbonic acid gas is very small.

WILDFIRE, Erysipelas.

WILDFIRE RASH, Ignis sylvaticus.

WILDUNGEN, MINERAL WATERS OF. Wildungen is situate a few miles from Cassel in Germany. In its neighborhood are several mineral springs, the chief of which, according to Stucke, contain a bituminous matter, chloride of sodium and sulphate of soda, carbonate and sulphate of lime, &c. They are used chiefly as refrigerants.

WILL, ([Sax.] *þilla*.) Voluntas.

WILLAN, Salix.

WILLOW, ([Sax.] *þelix*.) Salix—w. Bedford, Salix fragilis—w. Broudeaved, Salix latifolia—w. Crack, Salix fragilis—w. Great roundleaved, Salix caprea—w. Herb, common, Lythrum salicaria—w. Herb, great, Epilobium angustifolium—w. Herb, purple, Lythrum salicaria—w. Herb, purple veined, Epilobium coloratum.

WILLOW, Low Bush, Salix humilis—w. Red, Cornus sericea—w. Rose, Cornus sericea—w. Speckled, Salix humilis—w. Sweet, Myrica gale—w. White, Salix alba.

WIND, ([Sax.] *þind*.) (an example of onomatopœia—*oind*.) *Ventus*, *An'emos*, (F.) *Vent*. Winds exert considerable influence on the animal economy; acting by their temperature, which necessarily modifies that of the circumambient air, as well as by their moisture or dryness, and by the emanations of different kinds which they transport to greater or less distances. The winds which blow over a considerable extent of country are dry and serene: those which come from the ocean are moist and chilly.

WIND, Breath, see Flatulence.

WIND OF A BALL OR SHOT. A term applied to

the compression of air, supposed to be produced by the passage of a ball near a part of the body, so as to occasion what has been called a *wind contusion*.

WIND CONTUSION, see Wind of a Ball—w. Dropsy, Emphysema, Tympanites—w. Flower, Anemone—w. Pipe, Trachea—w. Root, Asclepias tuberosa—w. Weed, Asclepias tuberosa.

WINDINESS, Flatulence.

WIND'Y, *Flatulentus*, *Flatulent*, *Vaporosus*, (F.) *Venteux*, *Flatueux*. That which causes wind or flatulence; also, that which is caused by wind or flatulence; as *windy food*, *windy colic*, &c. Also, one affected with flatulence, or who is 'troubled with the wind.'

WINE, ([Sax.] *þin*, [G.] *Wein*, originally, perhaps, from *oivos*, *Œnos*, *Vinum*, *Meth'u*, (F.) *Vin*. A name given by chymists to all liquors that have become spirituous by fermentation. The term is generally, however, restricted to the fermented juice of the grape, or of fruits, the product of whose fermentation resembles, in almost all respects, that of the juice of the grape. Wine is composed of alcohol, æthanolic ether, saccharine matter, malic acid, tartaric acid, bitartrate of potassa, acetic acid, an extractive colouring matter more or less bitter and partly resinous, and of an aromatic substance. The extractive colouring matter is chiefly met with in red wines. All these constituents, except the alcohol, are found ready formed in the grape. The alcohol proceeds from the decomposition of the saccharine matter. A part of the acetic acid is also formed during fermentation. In all wines, too, there is more or less of an odorous principle, partly derived directly from the grape, and partly formed during fermentation, which has the characters of an essential oil, constitutes the perfume or *bouquet* of the wine, and is said not to exceed the forty-thousandth part of the wine. It is regarded as a true ether—a combination of oxide of ethyle with æthanolic acid.

The following table, drawn up by Mr. Brande, exhibits the quantity of spirit in different kinds of wine and liquors. It is proper, however, to remark, that many of these wines are prepared for the London market, and are more branded or "reinforced" than the same varieties sold in the United States. This is strikingly the case with port. Dr. Henderson, too, has remarked that some of the wines analyzed by Mr. Brande were mixed with a considerable quantity of adventitious alcohol. Dr. Henderson's additions and corrections have the letter H. affixed.

PROPORTION OF ALCOHOL, S. G. 0.825, IN ONE HUNDRED PARTS BY MEASURE OF THE FOLLOWING WINES, AND MALT AND SPIRITUOUS LIQUORS.

1. Lissa.....	26.47
Do.....	24.35
	Average, (a) 25.41
2. Raisin Wine.....	26.40
Do.....	25.77
Do.....	23.20
	Average, 25.12
3. Marsala.....	26.03
Do.....	25.05
	Average, (b) 25.09
4. Port—average of six kinds.....	23.43
Do.—highest.....	25.83
Do.—lowest.....	21.40
5. Madeira.....	24.42
Do.....	23.93
Do. (Sercial).....	21.45
Do.....	19.24
	Average, 22.27
6. Currant Wine.....	20.55
7. Sherry.....	19.81
Do.....	19.83
Do.....	18.79
Do.....	18.25
	Average, 19.17
8. Teneriffe.....	19.79

(a) 15° 90 H.

(b) 18° 40 H.

9. Colares	19-75
10. Lacryma Christi	19-70
11. Constantia—white	19-75
" red	18-92 (c)
12. Lisbon	18-94
13. Malaga	18-94
14. Bucelas	18-49
15. Red Madeira	22-30
Do	18-40
Average, ...	20-35
16. Cape Muscat	18-25
17. Cape Madeira	22-94
Do	20-50
Do	18-11
Average, ...	20-51
18. Grape Wine	18-11
19. Calcevilla	19-20
Do	18-10
Average, ...	18-65
20. Vidonia	19-25
21. Alba Flora	17-26
22. Malaga	17-26
23. White Hermitage	17-43
24. Roussillon	19-00
Do	17-26
Average, ...	18-13
25. Claret	17-11
Do	16-32
Do	14-08
Do	12-91
Average, (d)	15-10
26. Malmsey Madeira	16-40
27. Lunel	15-52
28. Scheraaz	15-52
29. Syracuse	15-28
30. Sauterne	14-22
31. Burgundy	16-60
Do	15-22
Do	14-53
Do	11-95
Average, ...	14-57
32. Hock	14-37
Do	13-00
Do. (old in cask)	8-88
Average, ...	12-08
Rudesheimer (1811)	H. 10-72
Do. (1800)	H. 12-22
Average, H.	11-47
Johannisberger	H. 8-71
33. Nice	14-63
34. Barsac	13-86
35. Tent	13-30
36. Champagne (still)	13-80
Do. (sparkling)	12-80
Do. (red)	12-66
Do.	11-30
Average, ...	12-61
37. Red Hermitage	12-32
38. Vin de Grave	13-94
Do.	12-80
Average, ...	13-37
39. Frontignac	12-79
40. Côte-Rôtie	12-32
41. Gooseberry Wine	11-54
42. Orange Wine—average of six samples made by London manufacturer	11-26
43. Tokay	9-88
44. Elder Wine	9-87
45. Rhenish Wine	H. 8-71
46. Cider—highest average	9-87
Do. lowest	5-21
47. Perry—average of four samples	7-26
48. Mead	7-32
49. Ale (Burton)	8-83
Do. (Edinburgh)	6-20
Do. (Dorchester)	5-56
Average, ...	6-87
50. Brown Stout	6-80
51. London Porter—average	4-20
Do. Small Beer—average	1-28
52. Brandy	53-39
53. Rum	53-68
54. Gin	51-60
55. Scott's Whisky	54-32
56. Irish do.	53-90

The only wine recommended in some pharmacopœias is *Sher'ry*, *Vi'num*, *V. al'bum*, *V. al'bum Hispan'icum*, *V. Xer'icum*, *Leuco'nus*. It is a *dry wine*; the least variable in its properties, and agrees best with the dyspeptic. The *brown sherries* differ from the *pale* in the addition of a colouring liquid to the former, termed *Arrope*, which

is prepared by boiling sherry down to a syrup; but they frequently owe their hue to burnt sugar or caramel. Other wines used officially are,—*Candy* or *Sack wine*, *Vi'num Canari'um*; *Mountain wine*, *Vi'num al'bum montanum*; *Tent wine*, *Vi'num tinto*; *Rhen'ish wine*, *Vi'num Rhena'num*; and *Port wine*, *Vi'num ru'brum*, *V. ru'brum Portugal'icum*, *V. Porten'sæ*.

The wines habitually drunk are almost innumerable. Those that are sweet, or contain a large portion of free saccharine matter, are decidedly the least wholesome to the dyspeptic.

When wine is good, and of a proper age, it is tonic and nutritive; when new, flatulent and cathartic, disagreeing with the stomach and bowels. In medicine, it is a valuable tonic in the last stage of typhus, when the skin is not too hot and dry. Its use, however, requires great caution; and when once commenced under proper circumstances, it cannot be suddenly dropped without mischief. It is, perhaps, the best permanent stimulus in the catalogue of the *materia medica*.

Mul'led Wine is made as follows:—Take of bruised *Cinnamon* $\mathfrak{z}\text{ij}$, half a *Nutmeg*, grated, and ten bruised *Cloves*. Infuse in *boiling Water* Oss for an hour; strain, and add *Sugar* $\mathfrak{z}\text{ij}$. Pour the whole into a pint of hot *Port* or *Sherry*. Useful in adynamic conditions.

WINE, AMIXÆAN, *Aminæum vinum*.

WINE OF ALOES, *Vinum aloes*.

WINE, ANTIMONIAL, *Vinum antimonii*—w. Antiscorbutic, *Vinum antiscorbuticum*—w. Aromatic, *Vinum aromaticum*—w. Bark, compound, *Vinum de kinâ kinâ compositum*—w. Barley, *Cerevisia*—w. Bitter, diuretic, *Vinum diureticum amarum*.

WINE BITTERS, *Vi'num gentianæ compositum*. The formulæ for these may be various. The following is as good and as agreeable as any. (*Rad. gentian.* $\mathfrak{Rj}$, *Cort. aurant.* $\mathfrak{z}\text{ss}$, *Sem. cardam.* cont. $\mathfrak{z}\text{iv}$, *Cinnam.* cort. $\mathfrak{z}\text{iv}$, *Wine*, foreign or domestic, three gallons and a half.) Tonic and stomachic.

WINE, CEDAR, *Cedrinum vinum*—w. Champagne, *Vinum campanum*—w. Chian, *Chium vinum*.

WINE OF COLCHICUM, *Vinum colchici*—w. of Colchicum seed, *Vinum colchici seminis*—w. of Ergot, *Vinum ergotæ*—w. of Gentian, compound, *Vinum gentianæ compositum*—w. of Hellebore, white, *Vinum veratri*.

WINE, IPECACUANHA, *Vinum Ipecacuanhæ*.

WINE OF IRON, *Vinum ferri*.

WINE, MEDICATED, *Vinum medicinale*—w. Mulled, see *Wine*.

WINE OF OPIUM, *Vinum opii*.

WINE, PORT, *Vinum Portugallicum*—w. Pullet, *Poulet, vin de*.

WINE OF QUINIA, *Vinum quiniæ*.

WINE, RHUBARB, *Vinum rhei*—w. Sherry, *Vinum*—w. Steel, *Vinum ferri*—w. Tar, see *Pinus sylvestris*—w. Tobacco, *Vinum tabaci*.

WINE WHEY, *White wine whey*, (Prov.) *Sack'whey*. Take of good milk two-thirds of a pint, and add water to make a pint. Take of sherry, or any other good white wine, two glasses, and of sugar a dessertspoonful. Place the milk and water in a deep pan on the fire, and the moment it boils, pour into it the wine and sugar. Stir assiduously for 12 or 15 minutes, whilst it boils. Lastly, strain through a sieve.

It is a good mode of giving wine in adynamic states.

WINE, WHITE, see *Wine*.

WINE OF WORMWOOD, *Vinum absinthites*.

WINGSEED, *Ptelea trifoliata*.

WINKERS, *Cilia*.

WINKING, ([Sax.] *pinician*, 'to wink,') *Niotation*, *Scardamygmus*.

WINTER BERRY, SMOOTH, *Prinos lævigatus*—w. Berry, *Virginia*, *Prinos*—w. Berry

(c) 14-50 H.

(d) 12-91 H.

whorled, Prinos—w. Bloom, Hamamelis Virginiana—w. Cough, Bronchitis, chronic—w. Green, Gaultheria—w. Green, roundleaved, Pyrola—w. Green, spotted, Pyrola maculata.

WINTER'S BARK TREE, *Wintera aromatica*.

WINTERA, (after Captain Winter, who first carried it to Europe,) see *Wintera aromatica*.

WINTERA AROMATICA, *Drîmÿs Winteri*, *Wintera aromatica*, *Winter's bark Tree*; *Oril. Magnoliaceæ*. The bark, *Wintera* (Ph. U. S.), *Cortex Winteranus* seu *Winteranus Magellanicus*, *C. Magellanicus*, *Cinnamomum Magellanicum*, (F.) *Écorce de Winter*, *Cannelle poivrée*, is very much allied in its properties to *Canella alba*.

WINTERANA AROMATICA, *Wintera aromatica*.

WIRSUNG'S DUCT, see *Pancreas*.

WIRTSPRINGS, Hangnails.

WISBADEN, MINERAL WATERS OF.

Wisbaden is a town of Germany, about 6 miles north of Mainz, and 22 west of Frankfurt. It has been long celebrated for its hot springs, and is much frequented. There are numerous cold and thermal springs:—the former containing sulphuric acid; the latter being saline and acidulous. The temperature of the hot springs varies from 117° to 160° Fah.

WISMUTHUM, Bismuth.

WITCHEN, *Sorbus aucuparia*.

WITCHHAZEL, *Hamamelis Virginiana*, *Sorbus aucuparia*.

WITCHMEAL, *Lycopodium*.

WITCHWOOD, *Sorbus aucuparia*.

WIT TEETH, see *Dentition*.

WITTER, (corr. from *guttur*), Throat.

WIZEN, (Weasand), *Trachea*.

WOAD, ([Sax.] *paaw*), *Isatis tinctoria*.

WOD, ([Sax.] *pow*), Insane.

WODNES, Insanity, Mania.

WOLFBANE, *Veratrum viride*.

WOLFSBANE, *Aconitum*—w. Wholesome, *Aconitum anthora*.

WOLFSCLAW, *Lycopodium*.

WOLFSJAW, see *Harelip*.

WOMAN, pl. *Women*, *Gy'ne*, *Mul'ier*, *Fæ'mina*, *Fî'ra*, (F.) *Femme*, (Sax. *pyrman*, 'wife-man,' 'woofman.') The adult female of the human race.

WOMANHOOD, *Mulieritas*.

WOMB, ([Sax.] *pamb*), Uterus—w. Falling down of the, *Prolapsus uteri*.

WOMB GRAIN, *Ergot*.

WOMB, LACERATION OF THE, Uterus, rupture of the—w. Tympany of the, *Physometra*.

WONGE, ([Sax.] *pænge*), Cheek.

WOOD, BRAZIL, *Cæsalpinia echinata*—w. Elk, *Andromeda arborea*.

WOOD LICE, *Onisci aselli*—w. March, *Sanicle*.

WOOD, NICARAGUA, see *Cæsalpinia*—w. Peach, see *Cæsalpinia*—w. Pernambuco, *Cæsalpinia echinata*—w. Sappan, *Cæsalpinia sappan*—w. Sappan, *Cæsalpinia sappan*.

WOOD SORREL, *Oxalis acetosella*—w. Sorrel, yellow, *Oxalis stricta*.

WOOD, SOUR, *Andromeda arborea*—w. Waxen, *Genista tinctoria*—w. Witch, *Sorbus aucuparia*.

WOODBINE, COMMON, *Lonicera periclymenum*.

WOODROOF, SWEETSCENTED, *Asperula odorata*.

WOODWEX, *Genista tinctoria*.

WOODS, SUDORIFIC, (F.) *Bois sudorifiques*. This term is applied, collectively, to *guaiacum*, *sassafras*, *china*, and *sarsaparilla*, which are often used together to form the sudorific decoction.

WOORALI, *Curare*.

WOORARA, *Curare*.

WORM, see *Alembic*, *Vermiform process* and *Worms*—w. Bark tree, *Geoffraea inermis*—w. Cakes, *Story's*, see *Cakes*, *worm*, *Story's*—w. Disease, *Helminthiasis*—w. Goosefoot, *Chenopodium anthelminticum*—w. Grass, perennial, *Spigelia Marilandica*.

WORM, GUINEA, *Dracunculus*.

WORM, LOZ'ENGES, CHING'S. Empirical preparations, which consist of yellow and brown lozenges. The former are directed to be taken in the evening; the latter on the following morning. The *Yellow Lozenges*. (*Saffron*, $\frac{3}{4}$ ss, *Water*, Oj; boil, and strain; and add of *White panacea of mercury*, (calomel washed in spirit of wine,) $\frac{1}{2}$ ij, *White sugar*, 28 lbs, *Mucilage of gum tragacanth* q. s. to make a mass. Divide, so that each lozenge may contain a grain of the panacea.) The *Brown Lozenges*. (*Panacea* $\frac{3}{4}$ vij, *Resin of Jalap* $\frac{1}{2}$ ijss, *White sugar* $\frac{1}{2}$ ix, *Mucilage of gum tragac.* q. s. Each lozenge to contain gr. ss of the panacea.)

SHER'MAN'S WORM LOZENGES, are said to be composed of calomel, gamboge, and sugar.

Peters's Worm Lozenges are said to be composed in a similar manner.

WORM ROOT, *Spigelia Marilandica*—w. Seed, *Artemisia santonica*, *Chenopodium anthelminticum*—w. Seed, goosefoot, *Chenopodium anthelminticum*—w. Weed, Corsican, *Corallina Corsicana*, *Polanisia graveolens*—w. Weed, white, *Corallina*—w. Wood, biennial, *Artemisia biennis*—w. Wood, Canada, *Artemisia canadensis*—w. Wood, common, *Artemisia absinthium*—w. Wood, creeping, *Artemisia rupestris*—w. Wood, lesser, *Artemisia Pontica*—w. Wood, Roman, *Artemisia Pontica*—w. Wood, sea, *Artemisia maritima*—w. Wood, silky, *Artemisia glaciata*.

WORMIA'NA OSSA SEU OSSIC'ULA, *O. epactalia* seu *raphogeminantia* seu *triq'uetra* seu *triangula'ria* *Blavii* seu *sutura'rum*, *Clav'es calvarie*, *Wormian bones*, (F.) *Os Wormiens* ou *épactaux* ou *surnuméraires* (Ch.) ou *intercalés* ou *triangulaires* ou *Cléfs du crâne*. Small bones, which are sometimes observable in the sutures of the bones of the cranium, of which they form a part. They are called *Wormia'na*, from Olaus Wormius, who is said to have first described them. They had been previously, however, mentioned by G. Andernach, a physician at Strasburg. The *Ossa Wormiana* exist more commonly in the sutures of the vault of the skull; especially in the lambdoidal, sagittal, and squamous. They are uncommon at the base of the skull. Their size is very variable, and shape irregular. Their structure and development are similar to those of the other bones of the cranium; and, like them, they are bounded by sutures.

WORMIT, *Artemisia absinthium*.

WORMS, originally from (L.) *Vermes*; (F.) *Vers*; *Entozoa*, *Entozoa'ria*, *Enterozoa*, *Endozoa*, *Entelmint'h*, *Entelmint'h'es*, *Tin'æ*, *Helminth'i*, *Entoparasites*, (F.) *Entozoaires*. Animals whose common character is that of existing only in other animals; (hence their name *entozoa*, from *εντος*, 'within,' and *ζωον*, 'animal.') They are met with, not only in the natural cavities, but even in the tissue of the organs, but chiefly in the intestines—*Ver'mes intesti'ni*, *Helminthia al'vi* seu *pod'icia*, *Splanchnelminth'ia*, *Intestinal worms*, (F.) *Vers intestinaux*, *Angiobremelminth'ia* of Piorry. Cuvier, Rudolphi, Brera, Bremser, Laënnec, Küchenmeister, and others, have endeavoured to classify them,—some according to their shape; others, according to their anatomical characters.

Worms form the second class of the first group of Animal Parasites according to the classification of Küchenmeister. They are subdivided by him as follows:

VERMES, HELMINTHA.

A. Platyelmia, Flat worms, (πλατύς, 'flat,' and ἔλμινξ, 'a worm.')

First Subclass.—Cestodea, cestode or tape worms, (κεστός, 'a band or girdle,' and ὄδω, 'resemblance.')

First Order—Bothriocephali.

Bothriocephalus latus.

Second Order—Tæniæ.

I. Tæniæ, which occur in the mature state.

1. Tænia solium.

2. Tænia mediocancellata.

3. Tænia nana.

II. Immature Tæniæ.

1. Cysticercus cellulosæ.

2. Cysticercus tenuicollis.

3. Echinococci.

Appendix.—Acephalocysts.

4. Cysticercus vesicæ hominis.

Second Subclass.—Trematoda, (trema, and odes.)

Family I. Monostoma.

1. Monostoma lentis.

Family II. Distoma.

1. Distoma hepaticum.

2. Distoma lanceolatum.

3. Distomum heterophyes.

4. Distomum hæmatobium.

5. Distomum ophthalmobium.

B. Nematelmia, Round or thread worms, (νῆμα, 'a thread,' and ἔλμινξ, 'a worm.')

1. Tricocephalus dispar.

Trichina spiralis.

2. Oxyuris vermicularis.

3. Strongylus gigas.

4. S. longevaginatus.

5. Ancylostomum duodenale.

6. Ascaris lumbricoides.

The following table exhibits most of the entozoa which have been met with in the human body, and their usual habitats:—

Entozoa.	Where found.
Trichoceph'alus dispar,	Intestines.
Oxyu'ris vermicula'ris,	
As'caris ala'ta,	
As'caris lumbricoi'des,	
Ancyelos'toma duodenale,	
Bothrioceph'alus latus,	" and Liver.
Tæ'nia so'lium,	
Pentas'toma constrictum,	Intestines and Ve-
Tæ'nia nana,	
Distoma hæmato'bium,	na porta.
Ditrachy'eras rudis, (?)	Intestines.
Diplos'o'ma crena'tum,	
Spirop'tera hom'inis,	Urinary bladder.
Dactyl'i'us aculea'tus,	
Dis'toma hepaticum,	Gall bladder.
Dis'toma lanceola'tum,	Hepatic duct.
Stron'gylus gigas,	Kidney.
Tetras'toma rena'lē,	
Fila'ria oc'uli,	Eye.
Monos'toma lentis,	Crystalline.
Dis'toma oc'uli huma'ni,	Capsule of crystal-
Acephalocyst'is endog'ena,	line.
Echinoco'cus hom'inis,	Liver.
Polys'toma pingu'ola,	Liver, Spleen, and
Hexathyrid'ium pingu'ola	
Polys'toma vena'rum seu	Ovary.
sanguic'ola,	Veins.
Hexathyrid'ium venarum,	

Fila'ria bronchia'lis, }
 Trich'ina spira'lis, }
 Cysticer'cus cellulo'sæ, }
 Acephalocyst'is multifi'da, }
 Fila'ria medinen'sis, }
 Bronchial glands.
 Muscles.
 Brain.
 Areolar texture.

The most common of these are:—the Oxyuris vermicularis, Ascaris lumbricoides, and Tæniæ, which are found in the intestines. The origin of these worms is extremely singular, and more favourable than any other fact to the hypothesis of spontaneous generation in the lowest tribes of animated nature. They are certainly not identical with any worms out of the body. They are most commonly met with in children improperly fed; but their presence cannot be positively detected by any symptoms, except that of the discharge of the worms themselves in the evacuations. They may, of course, give rise to every kind of irritation in the intestinal tube, and to sympathetic disturbance in most of the functions; but all these signs may be produced by intestinal irritation resulting from other causes. See Anthelmintics.

WORMS, Acne punctata.

WORSUM, (from [Sax.] wǽr, 'pus,') Pus.

WORT, Must'um, Infu'sum By'nes seu Br'a'sii seu Malt'i, (F.) Moût de la Bière. An infusion of malt. This has been recommended in scurvy. One measure of ground malt is added to three equal measures of boiling water. The mixture must be well stirred, and left to stand covered three or four hours. Dose, one to four pints, daily. It has, also, been recommended in other cases, where a strong putrescent disposition appeared to prevail in the fluids, as in cancerous and phagedenic ulcers. It is not now employed.

WORT, (Sax. wǽrt,) Herb.

WORTWALE, Hangnail.

WOSEN, (Weasand,) Trachea.

WOUND, (Sax. wund,) Vul'nus, (diminutive Vulnus'culum), Vol'nus, Pla'ga, Traum'a, Tro'ma, Tre'sis, Tre'sis Vul'nus, Bla'bē, Tym'ma, (F.) Plaie, Blessure. A solution of continuity in the soft parts, produced by some mechanical agent. Worms present innumerable differences, as regards their situation; the parts interested; their direction; size; shape; the nature of the instrument or agent by which they are produced; their more or less simple or complex character, duration, &c. A wound is called incised, (F.) Coupure, Incision, when made by a cutting instrument;—punctured, Nyg'ma, (F.) Pique, when made by a pointed instrument;—lacerated, (F.) Déchirure, Plaie par arrachement, when the parts are lacerated or torn by the wounded body; poi'soned, Celluli'tis venena'ta, Necu'sia, (F.) Plaie emvenimée, when some virulent or venomous substance has been introduced;—and contu'sed, (F.) Plaie contuse, when produced by a blunt body. Gun'shot wounds, Scolopetopla'ge, Vul'nera scolopeta'ria seu à scolopet'is seu Scolopet'ica, (F.) Plaies d'armes à feu ou d'arquebuse ou d'arquebuse, Coup de feu, belong to the last division.

WOURALI, Curare.

WOURARI, Curare.

WOURARU, Curare.

WOURSUM, ([Sc.] from [Sax.] wǽr, 'pus,') Pus.

WRACK BLADDER, YELLOW, (F.) Varce, Fucus vesiculosus—w. Sea, Fucus vesiculosus, Pila marina.

WRAMP, Sprain.

WRAPPING UP, (of the hydropathists,) Émaillottage.

WRATAK, Natus.

WRENCH, (from [Sax.] wringan, 'to wring,') Sprain.

WRET, 'Wart,' Verruca.

WRETWEED, Euphorbia palustris.

WRICK, Sprain.

WRIGHTIA ANTIDYSENTERICA, (after Dr. Wright, a physician and botanist of Jamaica.) Nerium antidysentericum.

WRINK'LE, (Sax. *rynkle*.) *Ru'ga*, *Rhy'tis*, *Ry'tis*, *Rhu'cus*, *Rhu'chus*, *Rhu'gus*, *Rhy'sse'ma*, *Corru'gio cu'tis*, *Ruelles*, (Sc.) *Ru'n'kle*, *Ru'n'kill*, (Prov.) *Crin'kleeran'kle*, (F.) *Ride*. A furrow or fold in the skin of the face, forehead, vagina, &c. See Corrugation.

WRINKLED, *Rugo'sus*, *Ru'gous*, *Ru'gose*, (F.) *Ride*. Full of wrinkles.

WRIST, ([Sax. *þýrr*, from *þnærtan*, 'to wrest,') Carpus.

WRISTDROP. Paralysis of the muscles of the forearm, induced by the poison of lead.

WRITER'S SPASM, see Spasm, writers'.

WRONG HEIR, Cancellus.

WURALI, Curare.

WURSTGIFT, 'sausage poison,' Allontoxicum.

WRYNECK, Torticollis.

X.

X LEG. Entogonyancon.

XALAPPA, *Convolvulus jalapa*.

XANTHIC OXIDE, Uric oxide

XANTHINE, Uric oxide.

XANTHIUM, *X. Strumarium* seu *Oriental'e*, seu *Vulga'rè*, *Barda'na mi'nor*, *Lap'pa mi'nor*, *Les'ser Bur'dock*, *Bur'weed*, *Bur'this'tle*, *Clot'bur*, *Cockle'bur*, (F.) *Lampourde*, *Petit Glouteron*, *Petite Bardane*, (from *xanthos*, because of the colour the plant yields.) This herb, as one of its names imports, was once esteemed for the cure of scrofula. The seeds have been administered in cutaneous affections.

XANTHIURIA. Xanthuria.

XANTHOPIGORIN and XANTHOPICRITE, (*xipos*, 'bitter,') see *Xanthoxylum clava Herculis*.

XANTHOP'SIA, (from *xanthos*, and *opsis*, 'vision,') Yellow vision, — as sometimes occurs in jaundice.

XANTHORRHIZA, (*xanthos*, and *ρίζα*, 'root,') *Xanthorrhiza apiifolia*.

XANTHORRHIZA APIIFOLIA, *Xanthorrhiza*, *X. simplicis'sima* seu *Morbo'ica*, *Xanthorrhiza tinctoria*, *Yel'lowroot*, *Par'aley-lea'ved Yel'lowroot*, *Yel'lowroot*, *Shrub yel'lowroot*. The root of this American plant — *Xanthorrhiza*, (Ph. U. S.) — is a strong and pleasant bitter; and, in the dose of two scruples, sits easy upon the stomach.

XANTHORRHEA, (*xanthos*, and *ῥεω*, 'to flow,') from its yellow resin.) *Grasstree*. A genus of trees in Australia, *Nat. Order*, Liliaceæ, of which there are several species. Two resins are obtained from them; one the *yel'low res'in of Xanthorrhæa* or of *New Hol'land Res'in lu'tea No'vi Bel'gii*, *Bot'any Bay res'in* or gum, *Ac'aroid res'in* or gum, *Res'in a vel Gum'mi acaroi'des*, which probably resembles *tolu* and *storax* in medicinal properties; the other the *red resin of Xanthorrhæa*, *Blackboy gum*.

XANTHOS, (*ξανθός*.) Yellow.

XANTHOSIS, *Xanthose*, (from *xanthos*, and *osis*.) A term applied to the yellow discoloration often observed in cancerous tumours — especially in encephaloid of the testicle.

XANTHOXYLIN, see *Xanthoxylum clava Herculis*.

XANTHOXYLUM, (*xanthos*, and *ξύλον*, 'wood,') *X. Frazin'eum*, *X. America'num*, *X. fraxinifolium*, *X. ramiflorum*, *X. tricar'pum*, *Prick'ly ash*, *Shrub'by prick'ly ash*, *Tooth'ache bush*, *Tooth'ache tree*, *Pell'itory*, *Yel'lowwood*, *Su'terberry*. *Ord.* Xanthoxylaceæ. A spirituous infusion of the bark — *Xanthoxylum* — Ph. U. S. — has been highly recommended in cases of violent *o'lio*. The fresh juice, expressed from the root of *xanthoxylum*, is said to afford relief in

the dry bellyache of the West Indies. A decoction of the bark is sometimes used as a wash to foul ulcers.

XANTHOXYLUM CAROLINIANUM has similar properties; but is more acrid.

XANTHOXYLUM CLAVATA HERCULIS of the West Indies, (F.) *Clavaliér à feuilles de frêne*, *Frêne épineux*, is a very strong stimulant and powerful sialogogue; and has been given internally in rheumatism. A neutral, very bitter, and astringent substance has been obtained from it, called *Xanthopierin*, *Xanthopierite*, and *Xanthox'glin*.

XANTHOXYLUM FRAXINEUM, *Xanthoxylum* — x. *Fraxinifolium*, *Xanthoxylum*.

XANTHOXYLUM OCTANDRA, *Fagara octandra* — x. *Piperitum*, *Fagara piperita* — x. *Ramiflorum*, *Xanthoxylum* — x. *Tricar'pum*, *Xanthoxylum*.

XANTHURIA, *Xanthiuria*, (from *xanthos*, and *ουρον*, 'urine,') A condition of the system and of the urine, in which xanthic oxide is deposited from the urine.

XANTOLINA, *Artemisia santonica*.

XENODOCEUM, (*ξενόδοceον*, from *ξενος*, 'a stranger,' and *δοceομαι*, 'to receive,') Hospital.

XENODOCHEUM, (*ξενόδοceον*.) Hospital.

XERANSIS, (*ξηρανσις*, from *ξηρος*, 'dry,') Arefaction, Drying.

XERANTIA, (from *ξηρος*, 'dry,') Siccantia.

XERA'SIA, (from *ξηρος*, 'dry,') A disease of the hairs, which become dry, cease to grow, and resemble down covered with dust.

XERION, (from *ξηρος*, 'dry,') Cataplasma.

XERO, (from *ξηρος*, 'dry,') in composition, dry.

XEROCOLLYRIUM, (*zero*, and *κολλυριον*, 'a collyrium,') *Collyrium sic'cum*. A dry collyrium.

XERODERMA, gen. *Xeroder'matis*, (*zero*, and *δερμα*, 'skin,') Diminution of secretion of the sebaceous glands.

XEROMA, gen. *Xero'matis*, (*zero*, and *oma*.) Xerophthalmia — x. *Conjunctival*, *Xerophthalmia*.

XEROMA, LACHRYMAL. Suppression of the lachrymal secretion. See Xerophthalmia.

XEROMYCTER, Xeromycteria.

XEROMYCTERIA, *Xeromycter*, (*zero*, and *μυκτηρ*, 'the nose,') Dryness of the nose.

XEROMYRON, *Unguentum sic'cum*, (from *zero*, and *μυρον*, 'an ointment,') A dry ointment.

XEROPHAGIA, (*zero*, and *φαγω*, 'I eat,') Excessive use of dry aliment. A kind of regimen to which the ancient athletes subjected themselves, with the view of increasing their strength.

XEROPHTHALMIA, *Ophthalmia sic'ca*, *Ophthalmoxero'sis*, *Xero'ma*, (*zero*, and *οφθαλμία*, 'inflammation of the eye,') An inflammation of the eye, without discharge. The term — as well as *Conjunctival* and *Lachrymal Xero'ma*, and *Ocular Conjunctiva* — has been given to cases

in which the conjunctiva is so changed, that it presents more of the characters of skin than of mucous membrane. It is incapable of radical cure. Also, lippitudo.

XEROTRIBIA, (*xero*, and *τριβω*, 'to rub,') see Friction.

XEROTRIPSIS, Friction, dry.

XIPHISTERNUM, (*xipho*, and *sternum*,) Xiphoid.

XIPHO, (from *ξίφος*, 'a sword,') in composition, the ensiform cartilage.

XIPHODES, Xiphoid.

XIPHODYMUS, (*xipho*, and *δύω*, 'to mingle,') Thoracogastrodidymsus.

XIPHOID, *Xiphoides*, *Xiphoides*, *Ensiformis*, *xipho*, and *εἶδος*, 'shape,') *Cartilago Xiphoides* seu *Xiphoides* seu *Scutalis* seu *Ensiformis* seu *Peltalis* seu *Peltatus*, *Furculā inferior*, *Mucronatum* os, *Mucronata Cartilago*, *Chondrus*, *Processus icphosternalis* seu *xiphoides* seu *Sterni xiphoides*, *Xiphisternum*, *Ensiform cartilage*, (F.) *Appendice ou cartilage xiphoides*, *C. Mucroné*, *Appendice sous-sternale*, *Palette*. The name of the appendix which terminates the sternum beneath.

XIPHOID LIG'AMENT, *Costoxiphoid L.* (F.) *Ligament xiphoidien ou costo-xiphoidien*, is a small, very thin, ligamentous fasciculus, which passes from the cartilage of prolongation of the 7th rib

to the anterior surface of the xiphoid cartilage, into which it is inserted by its fibres decussating with those of the opposite ligament.

XIPHOP'AGES, *Didymus Symphyoepigastrius*, (*xipho*, and *πηνωμ*, 'I fix,') A monstrosity, in which twins are united by the epigastrum, as in the case of the Siamese twins.

XYLOALOES, (*ξύλον*, 'wood,' and *aloes*,) *Agalochum*—*x.* Balsamum, see *Amyris opobalsamum*—*x.* Cassia, *Laurus cassia*—*x.* Cinnamonum, *Laurus cinnamomum*.

XYMPATHIA, Sympathy.

XYRIS, (from *ξύρον*, 'a razor,' from the shape of its leaves,) *Iris foetidissima*.

XYRIS BULBO'SA, *X. Indica* seu *Caroliniana*, *Yellow-eyed Grass*; indigenous; *Order*, *Xyridaceæ*; flowering from July to September. The roots and leaves are said to be used by the Hindoos in lepra, and chronic cutaneous diseases in general.

XYRUM, (*ξύρον*,) *Novacula*.

XYRUS, (*ξύρος*,) *Novacula*.

XYSMA, (*ξύσμα*,) *Lintheum*, *Rasura*, *Scobs*.

XYSMUS, (from *ξύω*, 'to scrape,') *Rasura*.

XYSTER, (*ξύστης*,) *Raspatorium*.

XYSTOS, (*ξύστος*, 'scraped,') *Lintheum*.

XYSTRUM, (*ξύστρον*,) *Raspatorium*—*x.* *Ophthalmicum*, *Ophthalmoxyster*.

Y.

YAM. This esculent root is principally obtained from three species of *Dioscorea*,—*alata bulbifera*, and *sativa*. They grow spontaneously in both the Indies, and their roots are eaten promiscuously, as the potato is with us. Their taste is somewhat like that of the potato.

YAM ROOT, WILD, *Dioscorea villosa*.

YARD, ([Sax.] *ýrn*,) *Penis*.

YARROW, COMMON, ([Sax.] *gearn*), *Achillea millefolium*.

YAPON, *Ilex vomitoria*.

YAVA SKIN, *Elephantiasis Arabum*.

YAWN'ING, (from [Sax.] *geonan*, 'to yawn,') *Chas'nē*, *Chas'mus*, *Hiat'us*, *Ric'tus*, *Osce'do*, *Oscitatio*, *Clo'nus pandiculatio maxilla'rum*, *Ga'ping*, (F.) *Baillement*. Yawning consists of a deep inspiration, with considerable separation of the jaws, executed slowly and in an involuntary manner, and followed by a prolonged and more or less sonorous expiration. This respiratory movement is preceded by a feeling of oppression in the epigastric region, and in the muscles of respiration and mastication. Yawning has been conceived to be owing to torpor in the pulmonary circulation; the causes producing it being commonly *emut*, fatigue, sleepiness, monotonous sounds, hunger, sympathy, &c. Yawning often precedes the invasion of certain intermittents, attacks of hysteria, &c., and it may, by its frequency and obstinacy, become a true disease.

YAW ROOT, *Stillingia*.

YAWS, *Framboesia*.

YEAST, *Yest*—*y.* Plant, *Torula cerevisiæ*.

YEBLE, (from *Ebulus*,) *Sambucus ebulus*.

YECKE, (*Yuke*), *Psora*.

YELK, ([Sax.] *gealep*, 'yellow,') see *Ovum*.

YELLOWOT, (*Yellow*), *Icterus*.

YEL'LOW, ([Sax.] *gealep*,) *Fla'vus*, *Lu'teus*, *Xan'thus*, (from past participle of Sax. *geaelan*, 'to burn,') One of the seven primitive colours.

YELLOW ASH, *Cladrastis tinctoria*—*y.* Berry, *Podophyllum montanum*—*y.* Fever, see *Fever*,

yellow—*y.* Fever, mild, see *Relapse*—*y.* Jack, *Fever*, *yellow*.

YELLOW LIG'AMENTS, (F.) *Ligaments jaunes*, occupy the spaces between the vertebral plates, from the interval, which separates the second vertebra from the third, to that which separates the last vertebra from the sacrum. They are so called in consequence of their colour.

YELLOW PAINT, *Hydrastis Canadensis*—*y.* Root, *Jeffersonia Bartoni*—*y.* Root, *parsley-leaved*, *Xanthorrhiza apiifolia*—*y.* Root, shrub, *Xanthorrhiza apiifolia*.

YELLOW SPRING, see *Ohio*, mineral waters of.

YELLOW SPRINGS. These are in Chester county, Pennsylvania. The temperature of the chief spring is 52° Fah. The water contains no impregnation.

YELLOW STAR, *Helenium autumnale*—*y.* Weed, *Ranunculus acris*—*y.* Wood, *Xanthoxylum fraxineum*—*y.* Wort, *parsley-leaved*, *Xanthorrhiza apiifolia*.

YELLOWOWS, *Cypripedium luteum*, *Icterus*, see *Leucorrhœa*, *Ranunculus acris*.

YEMEN ULCER, see *Beriberi*.

YERBA DEL SOLDADA, 'soldier's herb,' *Matico*—*y.* de *Huaca*, *Datura sanguinea*.

YERNIN, *Rennet*.

YEST, *Yeast*, ([Sax.] *geȝt*, from *ȝȝt*, 'a storm,' *ȝȝtȝ*, 'stormy,') *Cerevisiæ Fermentum*, *Sputum Cerevisiæ*, *Barm*, (F.) *Levûre*, *Levûre de la Bière*. A peculiar product which collects on the surface of beer whilst fermenting. *Yest* is antiseptic and tonic. It has been much extolled in putrid fevers, a tablespoonful being given as a dose, in porter, wine, &c. It is also applied in the form of cataplasm to foul ulcers. In all these cases it acts by virtue of the generation of carbonic acid gas.

YEUX ARDENS, Ardent eyes—*y.* d'Écrevisse, *Cancerum chelæ*.

YEW TREE, ([Sax.] *ip*,) *Taxus baccata*—*y.* American, *Taxus Canadensis*.

YEWER, see Mamma.
 YEX, (by onomatopœia.) Singultus.
 YLECH, Ilech.
 YLIATER, Ileen.
 YOKE, (by onomatopœia.) Singultus.
 YOLK, (Yelk,) see Ovum, and Vesicula umbilicalis—y. Membrane, see Vesicula umbilicalis.
 YOPON, Ilex vomitoria.
 YORK SPRINGS. These are in Adams county, Pennsylvania. One of them—a saline, containing sulphates of magnesia and lime, and chloride of sodium—is a mild cathartic and diuretic; the other is a strong chalybeate.
 Y O U K, ([G.] jucken, 'to itch,') Itching, Psora.
 YOUNG, Mr. JAMES, DISINFECTANT OF, see Manganese, sulphate of.

YOUTH, ([Sax. *yeorgu*.) Adolescence.
 YPSELOGLOSSUS, Basioglossus.
 YRIDES, Orpiment.
 YSAMBRA, *Pedigan'bra*. An ancient Spanish poison, of which hellebore formed the greatest part.
 YSOPE, Hyssopus.
 YUCCA, (Peruvian name,) *Jatropha manihot*.
 YUC'CA FILAMENTO'SA, *Bear's grass*, *Silkgrass*, indigenous. A tincture of the root has been used in gonorrhœa.
 Y U K E, ([G.] jucken, 'to itch,') Itching, Psora.
 YUX, (by onomatopœia.) Singultus.
 Y-WER-A, see Spirit.

Z.

ZACCHARUM, Saccharum.
 ZAEL, Borosail.
 ZAFFRAN, Crocus.
 ZAFFRE, Smalt.
 ZAMBO, see Mulatto, and Sambo.
 ZAMIA INTEGRIFOLIA, see Arrowroot—z. Pumila, see Arrowroot.
 ZAN'NA. A kind of Armenian bole, used by the ancients as a dessicative. It was also called *Zernacha*.
 ZANTHORRHIZA APIIFOLIA, *Xanthorrhiza apiifolia*.
 ZANTHOXYLUM, *Xanthoxylum*.
 ZAPPANIA NODIFLORA, (after P. A. Zappa, an Italian botanist.) *Anacoluppa*.
 ZARNACHA, Zanna.
 ZARSA, *Smilax sarsaparilla*.
 ZARZAPARILLA, *Smilax sarsaparilla*.
 ZEA MAYS, (*Ze* or *zea*, was the ancient name of a sort of corn, from *zaw*, 'to live,') *Z. vulgaris*, *Mays*, *M. ze'a* seu *America'na*, *A'dor*, *In'dian Corn*, *Corn*, *Turk'ey* or *In'dian Wheat*, *Maize*, *Mayz*, (F.) *Mais*, *Graine de Turquie*, *Blé d'Inde* ou *d'Espagne* ou *d'Italie* ou *de Turquie*. *Ord.* Gramineæ. A native of America, and much used over the whole of the continent as an article of diet. It is ground for the same purposes as wheat, and forms an excellent bread. The fecula or starch—*corn starch*, *maize'na*—makes an agreeable aliment; four tablespoonfuls of the starch to a quart of milk; boiled for a few minutes. It may be sweetened and flavoured, and eaten like *Mannamanger*.
 ZÉDOAIRE ROND, *Kæmpferia rotunda*.
 ZEDOARIA, *Kæmpferia rotunda*—z. Longa, see *Kæmpferia*—z. Rotunda, see *Kæmpferia*.
 ZEDOARY, *Kæmpferia rotunda*.
 ZELOSIS, (*ζηλωσις*), *Zelotypia*.
 ZELOTYP'IA, (from *zelos*, 'love,' and *τυπος*, 'form,' 'model,') *Zelo'sis*. It has been proposed to apply this name to the violent mental affection produced by the acknowledged or presumed infidelity of a beloved object; and to the consequences of such affection;—melancholy, mania, &c.
 ZEMA, gen. *Zem'atis*, (*ζεμα*.) Decoction.
 ZEN'ICON, (*ζενικον*.) Ancient name of a Keltic poison, used for poisoning arrows.
 ZEOCRITON DISTICHUM, (*ζεα*, 'corn,' and *πρην*, 'barley,') *Hordeum distichon*.
 ZEPHACH, *Peritonæum*.
 ZERMA, Zerna.
 ZER'NA, *Zerna*. Amongst the barbarous

Latin writers it signified ulcerated impetigo, *Impetigo ulcera'ta*.
 ZERUMBET, Cassumuniar.
 ZESIS, (*ζεσις*.) Effervescence, Decoction.
 ZEST, see Citrus medica.
 ZESTOLUSIA, (*ζεστολουσια*, from *ζεστος*, 'hot,' and *λουω*, 'to wash,') Bath. hot.
 ZIBEBÆ, see *Vitis vinifera*.
 ZIBETHUM, Civetta.
 ZIGNITES, Zingrites.
 ZINC, *Zink*, *Zin'eum*, *Zin'kum*, *Spel'trum*, *Spean'ter*, *Tin'tenag*, *Zin'chum*, *Zint*, *Stan'num Indicum*. (Perhaps formed etymologically from (G.) *Zin*, 'tin,') A metal obtained from calamine and blende. Its colour is bluishwhite; lustre of the fresh surface considerable, but soon dulled, from the facility with which it oxidates; hard; texture striated. S. g. 7.190. Melts at 700° Fah., burns with a bright flame in a higher temperature, and is volatilized in the form of a white, flocculent oxyd. Its use, in medicine, is in the formation of the following preparations.
 ZINC, ACÉTATE DE, *Zinci acetas*—z. Butter of, *Zinci chloridum*—z. Chloride of, *Zinci chloridum*—z. *Chlorure* de, *Zinci chloridum*—z. *Cyanure* de, *Zinci cyanuretum*—z. Ferrohdrocyanate of, *Zinci cyanuretum*—z. Flowers of, *Zinci oxydum*—z. Hydrocyanate of, *Zinci cyanuretum*—z. Iodide of, *Zinci Iodidum*—z. Ioduret of, *Zinci Iodidum*.
 ZINC AND MORPH'IA, DOUBLE I'ODIDE OF, (F.) *Iodure double de Zinc et de Morphine*. (This salt is prepared by boiling iodide of iodhydrate of morphia with water and zinc, after they have acted on each other for some days; filtering and crystallizing.) It is considered to combine the properties of morphia and the salts of zinc, and has been prescribed as an anodyne and antispasmodic. Dose, a quarter of a grain.
 ZINC, OXIDE DE, *Zinci oxydum*—z. Oxyd of, *Zinci oxydum*.
 ZINC AND STRYCH'NIA, DOUBLE I'ODIDE OF. This salt is prepared by digesting for several days in hot water, iodide of iodhydrate of strychnia and zinc; filtering and crystallizing. It has been proposed in the same cases as strychnia.
 ZINC, SULFATE DE, *Zinci sulphas*—z. Sulphate of, *Zinci sulphas*—z. Valerianate of, *Zinci Valerianas*.
 ZINCUM, *Zincum*.
 ZIN'CI ACE'TAS, *Ac'etate* of *Zinc*, (F.) *Ac'tate* de *Zinc*. (*Plumbi acet. 3ij*, *Zinc. granul. 3ix*, *Aquæ destillat. Oij*. Dissolve the acetate

of lead in the water, and filter. Add the zinc to the solution, and agitate them occasionally together, in a stopped bottle, for five or six hours, or until the liquor yields no precipitate with a solution of iodide of potassium. Filter the liquor, evaporate it with a moderate heat to one fifth, and set it aside to crystallize. Pour off the liquid, and dry the crystals on bibulous paper.

Should the crystals be coloured, dissolve them in distilled water, and, having heated the solution, drop into it, while hot, recently prepared carbonate of zinc until a little of the liquid, on being filtered, passes colourless; then filter the liquor, acidulate it with a few drops of acetic acid, evaporate, and crystallize.—*Ph. U. S.*)

Frequently used to form a gently astringent wash in ophthalmia and other local inflammations; and as an injection in gonorrhoea.

ZINCI CARBONAS IMPURUS, Calamina.

ZINCI CARBONAS PRÆCIPITATUS seu PRÆCIPITATA (Ph. U. S., 1863), *Præc'ipitated carbonate of Zinc*. (Zinci sulphat., Sodæ carbonat. aa lbj; Aq. bullient. cong. Dissolve the sulphate of zinc and carbonate of soda respectively in Oiv of water. Mix the solutions; stir, and set the mixture by, that the powder may subside. Having poured off the liquid, wash the precipitated carbonate of zinc with hot water, until the fluid of the washings is nearly tasteless, and dry with a gentle heat.—*Ph. U. S.*) Used to prepare the Zinci oxidum.

ZINCI CARBONAS PRÆPARATUS, Calamina præparata—z. Chloridi liquor, see Burnett's Disinfecting Liquid—z. Chloretum, zinci chloridum.

ZINCI CHLORIDUM, *Z. chloruræ* tum seu *Chloræ* tum, *Zinci cum chloræ* tum seu *muratæ* tum, *Chloride* of Zinc, (F.) *Chlorure* de Zinc. (Zinc. in frustulis, ℥ijss ; Acid. nitricæ, *Cretæ præparat.* aa ℥j ; Acid. muriat., *Aquæ* q. s. To the zinc, in the porcelain vessel, add gradually sufficient muriatic acid to dissolve it; strain, add the nitric acid, and evaporate to dryness. Dissolve the dry mass in water, add the chalk, allow the mixture to stand for 24 hours, filter, and again evaporate to dryness.—*Ph. U. S.*) It is very deliquescent in the air, forming the *Bufter* of Zinc, *Butyrum Zinci*.

Chloride of zinc is a powerful escharotic, and has been lately employed as an effective antiseptic. See Burnett's Disinfecting Liquid.

ZINCI CHLORURETUM, Z. chloridum.

ZINCI CYANURETUM, *Z. cyanidum* seu *Hydrocyanas* seu *Prus'ias*, *Zinci cum Cyanogenæ* tum seu *Borus'icum* seu *Zoët'icum*, *Hydrocyanate* of Zinc, (F.) *Cyanure* de Zinc. (This salt is formed by adding sulphate of zinc to hydrocyanate of potassa so long as any precipitate is thrown down. When dried and calcined at a low heat, the product is a mixture of cyanuret of zinc and of potassium.) It has been recommended in nervous cardialgia. Dose, from one-sixteenth to one-twelfth of a grain.

It is chiefly used as a caustic; one part of the chloride being mixed with four, three, or two parts of flour, according to the desired strength, and a few drops of water being added to form a paste.

ZINCI FERROHYDROCYANAS, *Zinci ferrocyanure* tum, *Cyanure* tum *Ferrozin'icum*, *Zinci cum Ferrohydrocyan'icum*, *Ferrohydrocyanate* of Zinc. (Made by the mutual decomposition of boiling-hot solutions of sulphate of zinc and ferrocyanate of potassa.) It has been recommended in the same cases as the last preparation.

ZINCI HYDROCYANAS, Zinci cyanuretum.

ZINCI IODIDUM, *Z. Ioduræ* tum, *Zinci cum Iodæ* tum, *Iodæ* tum *Zinci* seu *Zinci cum*, *Protoioduræ* tum *Zinci*, *Iodide* or *Ioduret* of Zinc, (F.) *Iodure* de Zinc. (This salt is made by boiling together iodine and zinc to dryness, and subliming.) An

ointment, composed of a drachm to an ounce of lard, has been used in cases where the external use of iodide of potassium is indicated.

ZINCI IODURETUM, Z. iodidum—z. Nix, Zinci oxydum.

ZINCI OXYDUM seu OXYDUM, *Zinci cum calcina* tum, *Nihil Al'bum*, *Lut'na Philosoph'orum* seu *Philosoph'ica*, *Nix Zinci*, *Nihil Gri'æum*, *Pompholyx*, *Oxyd* of Zinc, *Flow'ers* of Zinc, (F.) *Oxide* ou *Blanc* de Zinc. (Procured by burning zinc in a long, deep crucible, placed so as to collect the sublimate. Also, by exposing precipitated carbonate of zinc to a strong heat, so as to drive off the carbonic acid.—*Ph. U. S.*) It is said to be tonic and antispasmodic; externally, detergent and desiccative. Employed in epilepsy, chorea, &c. Usual dose, gr. ij to x; but it is probably inert. For its external use, see Ung. Zinci.

ZINCI OXYDUM IMPURUM, Tutia—z. Prussias, Zinci cyanuretum.

ZINCI SULPHAS, *Sulphate* of Zinc, *Zinci cum Vitriola* tum, *Vitriolum Al'bum* seu *Goslarien'se* seu *Zinci*, *Sulphas Zinci* cum, *Gil'la Vitrioli* seu *Theophrasti*, *Sol Vitrioli* seu *Vomitivum Vitrioli*, *Chalcum Vitriolum Al'bum*, *White Vitriol*, *White Cop'peras*, (F.) *Sulfate* de Zinc, *Couperose blanc*. (Zinc. in frustulis, ℥iv ; Acid. sulphuricæ, ℥vj ; *Aquæ destillat.* Oiv. To the zinc and water, previously introduced into a glass vessel, add by degrees the sulphuric acid, and, when the effervescence shall have ceased, filter the solution through paper; then boil it down till a pellicle begins to form, and set aside to crystallize.—*Ph. U. S.*) Sulphate of zinc is inodorous; taste, styptic. It is in white, semitransparent, efflorescent crystals; soluble in three parts of water at 60°. It is emetic, tonic, antispasmodic, and, externally, astringent. A nostrum, sold under the name *Antipertussis*, and, as its title imports, given in whooping-cough, contains this salt as its chief ingredient. It is emetic. Dose, as an emetic, gr. x to ℥ss ; as a tonic, gr. j to ij. Used externally as a collyrium, gr. j to water f. ℥j . A solution of the salt, to which a little sulphate of copper has been added, is said to constitute the *Disinfecting Liquid* of Larnaudès.

ZINCI VALERIANAS, *Zinci valerian'icum*, *Valerianate* of Zinc. (A salt formed by saturating valerianic acid with freshly prepared carbonate of zinc, or by the double decomposition of valerianate of soda and sulphate of zinc.—*Ph. U. S.*, 1863, and *Ph. B.*, 1864.) It is in pure white, pearly scales, and has been recommended as a tonic in neuralgic diseases, epilepsy, &c. The dose is one or two grains several times a day. It is officinal in the Dublin Pharmacopœia.

ZINCUM, Zinc—z. Borussicum, Zinci cyanuretum—z. Calcinatum, Zinci oxydum—z. Chloratum, Zinci chloridum—z. Cyanogenatum, Zinci cyanuretum—z. Ferrohydrocyanicum, Zinci ferohydrocyanas—z. Iodatatum, Zinci iodidum—z. Muraticum, Zinci chloridum—z. Valerianicum, Zinci valerianas—z. Vitriolatatum, Zinci sulphas—z. Zooticum, Zinci cyanuretum.

ZINGIBER, gen. *Zingib'eris*, (Zingib'ep). Amomum zingiber—z. Cassumuniar, see Cassumuniar—z. Germanicum, Arum maculatum—z. Officinale, Amomum zingiber—z. Purpureum, see Cassumuniar—z. Spurius, see Cassumuniar—z. Zerumbet, see Cassumuniar.

ZINGIBERIN, Piperoid.

ZINGRITES, *Zigni'tes*. A stone, resembling glass; to which the ancients attributed marvelous properties.

ZINK, Zinc.

ZINKUM, Zinc.

ZINT, Zincum.

ZINZIBER, (Zingib'ep), Amomum zingiber.

ZIPHAC, Peritonæum.

ZIRBUS, Epiploon—*z. Adipinus*, Epiploon, gastrocolic.

ZIZA'NIA AQUAT'ICA, (*Zizaniav* was the ancient name of some wild grain.) *Wild Rice*, *Can'-ada Rice*, *Water Oats*. *Ord.* Gramineæ; called by the Indians, *Menomene*, grows abundantly on the marshy margins of the northern lakes and waters of the upper branches of the Mississippi. The grain resembles oats. It furnishes the northern savages and the Canadian traders and hunters with their annual supplies of grain.

ZIZYPHUM, (*Arab.*), Jujube.

ZIZYPHUS JUJUBA, see Jujube—*z. Nitida*, see Jujube.

ZIZYPHUS CENOP'LIA, *Z. Nape'ca*; *Ord.* Rhamnaceæ; common all over India. The fruit is eaten by the natives. It is of a pleasant acid taste. A decoction of the bark of the fresh root is used to promote the healing of fresh wounds.

ZIZYPHUS SATIVA, see Jujube—*z. Sylvestris*, see Jujube.

ZOA'GRIA, (*ζωη*, 'life,' and *αγρω*, 'I catch up.') The saving and preservation of life. *Zoô'ois*. A reward for preserving life; *Sos'trum*. A votive offering hung up in the temples for a life saved.

ZOANTHRO'PIA, *Melanchol'ia zoanthro'pia*, (*zoon*, and *ανθρωπος*, 'a man.') A species of monomania, in which the patient believes himself transformed into an animal. Lycanthropia and Cynanthropia belong to this species of monomania.

ZOARA, Insomnia.

ZOE, (*ζωη*), Life.

ZOGRAPHIA, Zoographia.

ZOIATRIA, (*zoon*, and *ιατρεία*, 'medical treatment,') Veterinary art.

ZOIATRUS, (*zoon*, and *ιατρος*, 'a physician,') Hippiastr.

ZOISMUS, (from *zoon*,) Animality.

ZOMIDIUM, (*ζωμίδιον*, dim. of *ζωμος*), Jus.

ZOMOS, (*ζωμος*,) *Consommé*, Jus. See Broth.

ZONA, (*ζωνη*, 'a girdle,') *Herpes zoster*—*z. Castitatis*, *Hymen*—*z. Ciliaris*, Ciliary zone—*z. Ignea*, *Herpes zoster*.

ZONA PELLU'CIDA, *Oolem'ma pellucidum*, *Vitellary membrane*, *Cho'rion* (of some), (*F.*) *Vitelline*, *Membrane vitelline*. The membrane that surrounds the yolk in the mammalia is so called on account of its breadth and peculiar transparency.

ZONA SERPIGINOSA, *Herpes zoster*.

ZONA TENDINO'SA. The whitish circle around each auriculoventricular orifice of the heart, see *Circulus callosus Halleri*.

ZONA VIRGINITATIS, *Hymen*.

ZONE, *Cingulum*.

ZONULA CILIARIS, (dim. of *zona*,) see Ciliary, and Ciliary zone—*z. Cochleæ*, see *Lamina spiralis*—*z. Hildani*, *Cingulum Hildani*—*z. Zinnii*, see Ciliary zone.

ZOOBIOLOGIA, (*zoon*, *βίος*, 'life,' and *λογος*, 'a treatise,') see Physiology.

ZOOCHEMIA, Chymistry, animal.

ZOOCHEMICAL, *Zoöchym'ical*, *Zoöchem'icus*, *Zoöchym'icus*. Appertaining or having relation to animal chymistry.

ZOOCHEMY, Chymistry, animal.

ZOOCHYMY, Chymistry, animal.

ZOOGENIES, see Zoögeny. Bouchut has applied the term *Zoögenies* to animal parasites infesting the body.

ZOODES, (*ζωοδής*, from *zoon*, and *odes*,) Animal.

ZOÖG'ENY, *Zoögen'ia*, *Zoögen'esia*, (*zoon*, and *γενναιω*, 'to beget.') The doctrine of animal formation.

ZOÖG'RAPHY, *Zoögra'phia*, *Zogra'phia*, (*zoon*, and *γραφη*, 'a description.') A description of animals. The natural history of animals.

ZOÖHEMATINE, *Hæmatine*.

ZOOHYLAS, (*zoon*, and *βλην*, 'matter,') Animal matter.

ZOOHY'LICA CONCREMEN'TA. Concretions formed of animal matter.

ZOÖIASIS, (*zoon*, and *ιασις*, 'curation,') Veterinary art.

ZO'OID, *Zoöi'des*, (*ζωοειδής*, from *zoon*, and *ειδός*, 'resemblance,') Resembling or belonging to an animal. Every living cell may be regarded as a Zooid.

ZOÖLOG'ICAL, *Zoölog'icus*, (*F.*) *Zoologique*, (*zoon*, and *λογος*, 'a description,') Relating or appertaining to the natural history of animals.

ZOÖL'OGIST, *Zoölogis'ta*, *Zoö'l'ogus*. One skilled in the natural history of animals.

ZOÖL'OGY, *Zoölog'ia*. The natural history of animals.

ZOÖMAGNETISMUS, Magnetism, animal.

ZOÖMYCUS, (*zoon*, and *μυκος*, 'mucus,') Mucus.

ZOÖMYXA, (*zoon*, and *μυξα*, 'mucus,') Mucus.

ZOÖN, (*ζωον*,) Animal.

ZOÖNOM'IA, *Organozoönom'ia*, *Zoö'nomy*, (*zoon*, and *νομος*, 'law,') The law which governs the organic actions of animals in general:—*Animal Physiology*, *Zoöphysiolog'ia*.

ZOÖNOSOLOGY, *Zoonosolog'ia*, (*zoon*, *νοσος*, 'a disease,' and *λογος*, 'a description,') The nosology of animals.

ZOÖPATHOL'OGY, *Zoöpatholog'ia*, (*zoon*, *παθος*, 'a disease,' and *λογος*, 'a description,') The pathology of animals.

ZOÖPH'AGOUS, *Zoöph'agus*, (*F.*) *Zoöphage*, (*ζωοφαγος*, from *zoon*, and *φαγω*, 'I eat,') Feeding on animals. Carnivorous.

ZOÖPTHALMUS, (*ζωοφθαλμος*,) *Hydrophthalmia*.

ZOÖPHYSIOLOGY, *Zoöphysiolog'ia*, (*zoon*, and *physiologia*,) Physiology, animal, *Zoonomia*.

ZOÖPHYTE, *Zoöph'ytem*, (*zoon*, and *φυτον*, 'a plant,') *Phytozoö'n*, (*F.*) *Phytozoöaire*, *Animal plant*. A name given to the lowest class of animals, which seem to form the transition between the animal and the vegetable kingdoms.

ZOOSIS, (from *zoon*,) Animation, *Zoagria*.

ZOÖSPERMATA, *Spermatozoa*.

ZOÖSPERMES, *Spermatozoa*.

ZOÖTHAP'SIS, (*ζωη*, 'life,' and *θάπτω*, 'I inter,') Burial alive. Premature interment.

ZOÖTOMIST, *Zootom'icus*, (*zoon*, and *τεμνω*, 'I cut,') One who dissects animals.

ZOÖTOMY, *Zoötom'ia*, *Anatom'ia animal'ia* seu *comparat'ia* seu *comparat'a*, *Theriot'omy*, *Compar'ative Anat'omy*. The anatomy of animals. Animal anatomy.

ZOPISSA, (*ζωπισσα*, from *ζωος*, 'living,' and *πισσα*, 'pitch,') see *Pinus sylvestris*.

ZOPYRON, (*ζωπυρον*, from *ζωω*, 'to be alive,' and *πυρ*, 'fire,') Fomites.

ZOSTER, (*ζωστρον*, 'a girth,') *Herpes zoster*.

ZOSTERA MARINA, *Pila marina*—*z. Martima*, *Pila marina*.

ZOSTÈRE, *Pila marina*.

ZUCINETTE, *Narcissus pseudonarcissus*.

ZOZAR, *Saccharum*.

ZUCARO, *Saccharum*.

ZUCCARUM, *Saccharum*.

ZUCHAR, *Saccharum*.

ZUCHARUM, *Saccharum*.

ZUCHRA, Saccharum.

ZULAPIUM, Julep.

ZYGAPOPHYSES, (ζυγαν, 'a yoke,' and *apophysis*,) see Vertebrae.

ZYGOMA, (ζυγμα, from ζυγον, 'a yoke,')

Malæ os, see Zygomatic process.

ZYGOMATIC, *Zygomaticus*, *Juga'lis*, *Ju'gal*, (F.) *Zygomatique*. That which relates to the *zygoma* or cheekbone.

ZYGOMATIC ARCH, see Zygomatic process — z. Bone, Malæ os.

ZYGOMATIC FOS'SA is the space included between the posterior margin of the outer ala of the pterygoid process and the crista which descends from the malar tuberosity to the superior alveolar margin.

ZYGOMATIC MUS'LES are two in number. 1. *Zygomaticus major*, *Distort'or O'ris*, (F.) *Zygomatico-labial* (Ch.), *Muscle grand zygomatique*, which is situate obliquely at the fore-part and sides of the face. It is a small, rounded muscle; arises from the outer surface of the os malæ, and is inserted at the commissure of the lips, which it raises and draws outwards. It acts principally in laughing. 2. *Zygomaticus minor*, (F.) *Petit Zygomatico-labial*, (Ch.), *Petit Zygomatique*, does not always exist. It is situate on the inner side of the last; arises from the outer surface of the os malæ, and is inserted at the corner of the mouth, or rather into the upper lip, which it raises and draws outwards.

ZYGOMATIC PROC'ESS, *Processus zygomaticus*, *Apoph'ysis Zygomatica*, *Ma'lar* or *Ju'gal proc'ess*, (F.) *Apophyse Zygomatique*, arises from the outer surface of the temporal bone by two roots, one of which is transverse, (*Condyle du temporal*, (Ch.,) and the other longitudinal: between them is the glenoid cavity. The process passes for-

wards, and is articulated with the posterior angle of the os malæ, with which it forms a bony bridge, called the *Zygomatic arch*, *Ar'cus* seu *Pons zygomaticus*, (F.) *Arcade zygomatique*. The cavity beneath the bridge is sometimes called *Zygo'ma*.

ZYGOMATIC SUT'URE is placed between the zygomatic parts of the temporal and cheek bones; and slants obliquely downwards and backwards.

ZYGOMATIQUE, Zygomatic — z. *Grand*, Zygomaticus major — z. *Petit*, Zygomaticus minor.

ZYGOMATO-AURICULAIRE, Anterior auris — z. *Labial*, Zygomaticus major — z. *Labial*, *petit*, Zygomaticus major — z. *Maxillaire*, *Masseter* — z. *Oriculaire*, Anterior auris.

ZYGOPHYLLUM FABA'GO, (ζυγαν, 'yoke, a pair,' and φυλλον, 'a leaf,') (F.) *Fabagelle*. *Ord.* Zygophyllaceæ. A Syrian and Mauritanic plant, possessed of an acrid and bitter taste. It has vermifuge properties. The Arabs believe that the recently expressed juice of *Zygophyllum simplex*, is capable of removing specks from the cornea.

ZYMA, (ζυμη,) Ferment.

ZYMOMA, gen. *Zymo'matis*, (ζυμομα,) Ferment. ZYMOSIS, (ζυμωσις, from *zyma*, and *osis*,) Fermentation, see Zymotic.

ZYMOTIC, *Zymo'ticus*, (F.) *Zymotique*, (from *zyma*.) Relating or appertaining to fermentation. A *zymotic disease* is any epidemic, endemic, contagious or sporadic affection, which is produced by some morbid principle acting on the organism similar to a ferment — as the major exanthemata. *Zymoses* has been used by some to signify such affections.

ZYTHOGALA, (*zythus*, and γαλα, 'milk.')

Posset.

ZYTHUM, Cerevisia.

ZYTHUS, (ζυθος,) Cerevizia.

HENRY C. LEA'S

(LATE LEA & BLANCHARD'S)

CLASSIFIED CATALOGUE OF MEDICAL AND SURGICAL PUBLICATIONS.

In asking the attention of the profession to the works contained in the following pages, the publisher would state that no pains are spared to secure a continuance of the confidence earned for the publications of the house by their careful selection and accuracy and finish of execution.

The printed prices are those at which books can generally be supplied by booksellers throughout the United States, who can readily procure for their customers any works not kept in stock. Where access to bookstores is not convenient, books will be sent by mail post-paid on receipt of the price, but no risks are assumed either on the money or the books, and no publications but my own are supplied. Gentlemen will therefore in most cases find it more convenient to deal with the nearest bookseller.

An ILLUSTRATED CATALOGUE, of 64 octavo pages, handsomely printed, will be forwarded by mail, postpaid, on receipt of ten cents.

HENRY C. LEA.

Nos. 706 and 708 SANSON ST., PHILADELPHIA, Sept. 1870.

ADDITIONAL INDUCEMENT FOR SUBSCRIBERS TO

THE AMERICAN JOURNAL OF THE MEDICAL SCIENCES.

THREE MEDICAL JOURNALS, containing over 2000 LARGE PAGES,

Free of Postage, for SIX DOLLARS Per Annum.

TERMS FOR 1870:

THE AMERICAN JOURNAL OF THE MEDICAL SCIENCES, and	} Five Dollars per annum, in advance.
THE MEDICAL NEWS AND LIBRARY, both free of postage,	

OR,

THE AMERICAN JOURNAL OF THE MEDICAL SCIENCES, published quarterly (1150 pages per annum), with	} Six Dollars per annum in advance.
THE MEDICAL NEWS AND LIBRARY, monthly (384 pp. per annum), and	
THE HALF-YEARLY ABSTRACT OF THE MEDICAL SCIENCES, published Feb. and August (600 pages per annum), all free of postage.	

SEPARATE SUBSCRIPTIONS TO

THE AMERICAN JOURNAL OF THE MEDICAL SCIENCES, subject to postage when not paid for in advance, Five Dollars.

THE MEDICAL NEWS AND LIBRARY, free of postage, in advance, One Dollar.

THE HALF-YEARLY ABSTRACT, Two Dollars and a Half per annum in advance. Single numbers One Dollar and a Half.

It is manifest that only a very wide circulation can enable so vast an amount of valuable practical matter to be supplied at a price so unprecedentedly low. The publisher, therefore, has much gratification in stating that the rapid and steady increase in the subscription list promises to render the enterprise a permanent one, and it is

with especial pleasure that he acknowledges the valuable assistance spontaneously rendered by so many of the old subscribers to the "JOURNAL," who have kindly made known among their friends the advantages thus offered and have induced them to subscribe. Relying upon a continuance of these friendly exertions, he hopes to be able to maintain the unexampled rates at which these works are now supplied, and to succeed in his endeavor to place upon the table of every reading practitioner in the United States a monthly, a quarterly, and a half-yearly periodical at the comparatively trifling cost of SIX DOLLARS *per annum*.

These periodicals are universally known for their high professional standing in their several spheres.

I.

THE AMERICAN JOURNAL OF THE MEDICAL SCIENCES,

EDITED BY ISAAC HAYS, M. D.,

is published Quarterly, on the first of January, April, July, and October. Each number contains nearly three hundred large octavo pages, appropriately illustrated, wherever necessary. It has now been issued regularly for over FORTY years, during nearly the whole of which time it has been under the control of the present editor. Throughout this long period, it has maintained its position in the highest rank of medical periodicals both at home and abroad, and has received the cordial support of the entire profession in this country. Among its Collaborators will be found a large number of the most distinguished names of the profession in every section of the United States, rendering the department devoted to

ORIGINAL COMMUNICATIONS

full of varied and important matter, of great interest to all practitioners. Thus, during 1869, articles have appeared in its pages from one hundred and fifteen gentlemen of the highest standing in the profession throughout the United States.*

Following this is the "REVIEW DEPARTMENT," containing extended and impartial reviews of all important new works, together with numerous elaborate "ANALYTICAL AND BIBLIOGRAPHICAL NOTICES" of nearly all the medical publications of the day.

This is followed by the "QUARTERLY SUMMARY OF IMPROVEMENTS AND DISCOVERIES IN THE MEDICAL SCIENCES," classified and arranged under different heads, presenting a very complete digest of all that is new and interesting to the physician, abroad as well as at home.

Thus, during the year 1869, the "JOURNAL" furnished to its subscribers One Hundred and Forty-seven Original Communications, Eighty-eight Reviews and Bibliographical Notices, and Two Hundred and Eighty-seven articles in the Quarterly Summaries, making a total of over FIVE HUNDRED articles emanating from the best professional minds in America and Europe.

To old subscribers, many of whom have been on the list for twenty or thirty years, the publisher feels that no promises for the future are necessary; but gentlemen who may now propose for the first time to subscribe may rest assured that no exertion will be spared to maintain the "JOURNAL" in the high position which it has so long occupied ("one of the best of its kind,"—*London Lancet*, Aug. 20, 1870), as a national exponent of scientific medicine, and as a medium of intercommunication between the profession of Europe and America—in the words of the "London Medical Times" (Sept. 5th, 1868) "almost the only one that circulates everywhere, all over the Union and in Europe"—to render it, in fact, necessary to every practitioner who desires to keep on a level with the progress of his science.

The subscription price of the "AMERICAN JOURNAL OF THE MEDICAL SCIENCES" has never been raised, during its long career. It is still FIVE DOLLARS *per annum*; and when paid for in advance, the subscriber receives in addition the "MEDICAL NEWS AND LIBRARY," making in all about 1500 large octavo pages *per annum*, free of postage.

II.

THE MEDICAL NEWS AND LIBRARY

is a monthly periodical of Thirty-two large octavo pages, making 384 pages *per annum*. Its "NEWS DEPARTMENT" presents the current information of the day, with Clinical Lectures and Hospital Gleanings; while the "LIBRARY DEPARTMENT" is devoted to publishing standard works on the various branches of medical science, pagéd

* Communications are invited from gentlemen in all parts of the country. Elaborate articles inserted by the Editor are paid for by the Publisher.

separately, so that they can be removed and bound on completion. In this manner subscribers have received, without expense, such works as "WATSON'S PRACTICE," "TODD AND BOWMAN'S PHYSIOLOGY," "WEST ON CHILDREN," "MALGAIGNE'S SURGERY," &c. &c. And with January, 1870, is commenced a series of monographs on Diseases of the Spinal Column and of the Nerves from "REYNOLDS'S SYSTEM OF MEDICINE," rendering this a very eligible period for the commencement of new subscriptions.

As stated above, the subscription price of the "MEDICAL NEWS AND LIBRARY" is ONE DOLLAR per annum in advance; and it is furnished without charge to all advance paying subscribers to the "AMERICAN JOURNAL OF THE MEDICAL SCIENCES."

III.

THE HALF-YEARLY ABSTRACT OF THE MEDICAL SCIENCES

is issued in half-yearly volumes, which will be delivered to subscribers about the first of February, and first of August. Each volume contains about 300 closely printed octavo pages, making about six hundred pages per annum.

"RANKING'S ABSTRACT" has now been published in England regularly for more than twenty years, and has acquired the highest reputation for the ability and industry with which the essence of medical literature is condensed into its pages. It purports to be "*A Digest of British and Continental Medicine, and of the Progress of Medicine and the Collateral Sciences,*" and it is even more than this, for America is largely represented in its pages. It draws its material not only from all the leading American, British, and Continental journals, but also from the medical works and treatises issued during the preceding six months, thus giving a complete digest of medical progress. Each article is carefully condensed, so as to present its substance in the smallest possible compass, thus affording space for the very large amount of information laid before its readers. The volumes of 1869, for instance, have contained

THIRTY-FOUR ARTICLES ON GENERAL QUESTIONS IN MEDICINE.
ONE HUNDRED AND ELEVEN ARTICLES ON SPECIAL QUESTIONS IN MEDICINE.
NINETEEN ARTICLES ON FORENSIC MEDICINE.
NINETY-NINE ARTICLES ON THERAPEUTICS.
THIRTY-SEVEN ARTICLES ON GENERAL QUESTIONS IN SURGERY.
ONE HUNDRED AND SIXTY-SEVEN ARTICLES ON SPECIAL QUESTIONS IN SURGERY.
NINETY-FIVE ARTICLES ON MIDWIFERY AND DISEASES OF WOMEN AND CHILDREN.

Making in all over five hundred and fifty articles in a single year. Each volume, moreover, is systematically arranged, with an elaborate Table of Contents and a very full Index, thus facilitating the researches of the reader in pursuit of particular subjects, and enabling him to refer without loss of time to the vast amount of information contained in its pages.

The subscription price of the "ABSTRACT," mailed free of postage, is Two DOLLARS AND A HALF per annum, payable in advance. Single volumes, \$1 50 each.

As stated above, however, it will be supplied in conjunction with the "AMERICAN JOURNAL OF THE MEDICAL SCIENCES" and the "MEDICAL NEWS AND LIBRARY," the whole free of postage, for SIX DOLLARS PER ANNUM IN ADVANCE.

For this small sum the subscriber will therefore receive three periodicals costing separately Eight Dollars and a Half, each of them enjoying the highest reputation in its class, containing in all over two THOUSAND PAGES of the choicest reading, and presenting a complete view of medical progress throughout both hemispheres.

In this effort to bring so large an amount of practical information within the reach of every member of the profession, the publisher confidently anticipates the friendly aid of all who are interested in the dissemination of sound medical literature. He trusts, especially, that the subscribers to the "AMERICAN MEDICAL JOURNAL" will call the attention of their acquaintances to the advantages thus offered, and that he will be sustained in the endeavor to permanently establish medical periodical literature on a footing of cheapness never heretofore attempted.

* * * Gentlemen desiring to avail themselves of the advantages thus offered will do well to forward their subscriptions at an early day, in order to insure the receipt of complete sets for the year 1870, as the constant increase in the subscription list almost always exhausts the quantity printed shortly after publication.

☞ The safest mode of remittance is by postal money order, drawn to the order of the undersigned. Where money order post-offices are not accessible, remittances for the "JOURNAL" may be made at the risk of the publisher, by forwarding in REGISTERED letters. Address,

HENRY C. LEA,
Nos. 706 and 708 SANSON ST., PHILADELPHIA, PA.

DUNGLISON (ROBLEY), M. D.,*Professor of Institutes of Medicine in Jefferson Medical College, Philadelphia.*

MEDICAL LEXICON; A DICTIONARY OF MEDICAL SCIENCE: Containing a concise explanation of the various Subjects and Terms of Anatomy, Physiology, Pathology, Hygiene, Therapeutics, Pharmacology, Pharmacy, Surgery, Obstetrics, Medical Jurisprudence, and Dentistry. Notices of Climate and of Mineral Waters; Formulæ for Official, Empirical, and Dietetic Preparations; with the Accentuation and Etymology of the Terms, and the French and other Synonymes; so as to constitute a French as well as English Medical Lexicon. Thoroughly Revised, and very greatly Modified and Augmented. In one very large and handsome royal octavo volume of 1048 double-columned pages, in small type; strongly done up in extra cloth, \$6 00; leather, raised bands, \$6 75.

The object of the author from the outset has not been to make the work a mere lexicon or dictionary of terms, but to afford, under each, a condensed view of its various medical relations, and thus to render the work an epitome of the existing condition of medical science. Starting with this view, the immense demand which has existed for the work has enabled him, in repeated revisions, to augment its completeness and usefulness, until at length it has attained the position of a recognized and standard authority wherever the language is spoken. The mechanical execution of this edition will be found greatly superior to that of previous impressions. By enlarging the size of the volume to a royal octavo, and by the employment of a small but clear type, on extra fine paper, the additions have been incorporated without materially increasing the bulk of the volume, and the matter of two or three ordinary octavos has been compressed into the space of one not unhandy for consultation and reference.

It would be a work of supererogation to bestow a word of praise upon this Lexicon. We can only wonder at the labor expended, for whenever we refer to its pages for information we are seldom disappointed in finding all we desire, whether it be in accentuation, etymology, or definition of terms.—*New York Medical Journal*, November, 1865.

It would be mere waste of words in us to express our admiration of a work which is so universally and deservedly appreciated. The most admirable work of its kind in the English language. As a book of reference it is invaluable to the medical practitioner, and in every instance that we have turned over its pages for information we have been charmed by the clearness of language and the accuracy of detail with which each abounds. We can most cordially and confidently commend it to our readers.—*Glasgow Medical Journal*, January, 1866.

A work to which there is no equal in the English language.—*Edinburgh Medical Journal*.

It is something more than a dictionary, and something less than an encyclopædia. This edition of the well-known work is a great improvement on its predecessors. The book is one of the very few of which it may be said with truth that every medical man should possess it.—*London Medical Times*, Aug. 26, 1865.

Few works of the class exhibit a grander monument of patient research and of scientific lore. The extent of the sale of this lexicon is sufficient to testify to its usefulness, and to the great service conferred by Dr. Robley Dunglison on the profession, and indeed on others, by its issue.—*London Lancet*, May 13, 1865.

The old edition, which is now superseded by the new, has been universally looked upon by the medical profession as a work of immense research and great value. The new has increased usefulness; for medicine, in all its branches, has been making such progress that many new terms and subjects have recently been introduced: all of which may be found fully defined in the present edition. We know of no other dictionary in the English language that can bear a comparison with it in point of completeness of subjects and accuracy of statement.—*N. Y. Druggists' Circular*, 1865.

For many years Dunglison's Dictionary has been the standard book of reference with most practitioners in this country, and we can certainly commend this work to the renewed confidence and regard of our readers.—*Cincinnati Lancet*, April, 1865.

It is undoubtedly the most complete and useful medical dictionary hitherto published in this country.—*Chicago Med. Examiner*, February, 1865.

What we take to be decidedly the best medical dictionary in the English language. The present edition is brought fully up to the advanced state of science. For many a long year "Dunglison" has been at our elbow, a constant companion and friend, and we greet him in his replenished and improved form with especial satisfaction.—*Pacific Med. and Surg. Journal*, June 27, 1865.

This is, perhaps, the book of all others which the physician or surgeon should have on his shelves. It is more needed at the present day than a few years back.—*Canada Med. Journal*, July, 1865.

It deservedly stands at the head, and cannot be surpassed in excellence.—*Buffalo Med. and Surg. Journal*, April, 1865.

We can sincerely commend Dr. Dunglison's work as most thorough, scientific, and accurate. We have tested it by searching its pages for new terms, which have abounded so much of late in medical nomenclature, and our search has been successful in every instance. We have been particularly struck with the fulness of the synonymy and the accuracy of the derivation of words. It is as necessary a work to every enlightened physician as Worcester's English Dictionary is to every one who would keep up his knowledge of the English tongue to the standard of the present day. It is, to our mind, the most complete work of the kind with which we are acquainted.—*Boston Med. and Surg. Journal*, June 22, 1865.

We are free to confess that we know of no medical dictionary more complete; no one better, if so well adapted for the use of the student; no one that may be consulted with more satisfaction by the medical practitioner.—*Am. Jour. Med. Sciences*, April, 1865.

The value of the present edition has been greatly enhanced by the introduction of new subjects and terms, and a more complete etymology and accentuation, which renders the work not only satisfactory and desirable, but indispensable to the physician.—*Chicago Med. Journal*, April, 1865.

No intelligent member of the profession can or will be without it.—*St. Louis Med. and Surg. Journal*, April, 1865.

It has the rare merit that it certainly has no rival in the English language for accuracy and extent of references.—*London Medical Gazette*.

HOBLYN (RICHARD D.), M. D.

A DICTIONARY OF THE TERMS USED IN MEDICINE AND THE COLLATERAL SCIENCES. A new American edition, revised, with numerous additions, by ISAAC HAYS, M.D., Editor of the "American Journal of the Medical Sciences." In one large royal 12mo. volume of over 500 double-columned pages; extra cloth, \$1 50; leather, \$2 00.

It is the best book of definitions we have, and ought always to be upon the student's table.—*Southern Med. and Surg. Journal*.

NEILL (JOHN), M.D., and SMITH (FRANCIS G.), M.D.,

Prof of the Institutes of Medicine in the Univ. of Penna.

AN ANALYTICAL COMPENDIUM OF THE VARIOUS BRANCHES OF MEDICAL SCIENCE; for the Use and Examination of Students. A new edition, revised and improved. In one very large and handsomely printed royal 12mo. volume, of about one thousand pages, with 374 wood cuts, extra cloth, \$4; strongly bound in leather, with raised bands, \$4 75.

The Compend of Drs. Neill and Smith is incomparably the most valuable work of its class ever published in this country. Attempts have been made in various quarters to squeeze Anatomy, Physiology, Surgery, the Practice of Medicine, Obstetrics, Materia Medica, and Chemistry into a single manual; but the operation has signally failed in the hands of all up to the advent of "Neill and Smith's" volume, which is quite a miracle of success. The outlines of the whole are admirably drawn and illustrated, and the authors are eminently entitled to the grateful consideration of the student of every class.—*N. O. Med. and Surg. Journal.*

There are but few students or practitioners of medicine unacquainted with the former editions of this unassuming though highly instructive work. The whole science of medicine appears to have been sifted, as the gold-bearing sands of El Dorado, and the pre-

cious facts treasured up in this little volume. A complete portable library so condensed that the student may make it his constant pocket companion.—*Western Lancet.*

In the rapid course of lectures, where work for the students is heavy, and review necessary for an examination, a compend is not only valuable, but it is almost a *sine qua non*. The one before us is, in most of the divisions, the most unexceptionable of all books of the kind that we know of. Of course it is useless for us to recommend it to all last course students, but there is a class to whom we very sincerely commend this cheap book as worth its weight in silver—that class is the graduates in medicine of more than ten years' standing, who have not studied medicine since. They will perhaps find out from it that the science is not exactly now what it was when they left it off.—*The Stethoscope.*

HARTSHORNE (HENRY), M.D.,

Professor of Hygiene in the University of Pennsylvania.

A CONSPECTUS OF THE MEDICAL SCIENCES; containing Handbooks on Anatomy, Physiology, Chemistry, Materia Medica, Practical Medicine, Surgery, and Obstetrics. In one large royal 12mo. volume of 1000 closely printed pages, with over 300 illustrations on wood, extra cloth, \$4 50; leather, raised bands, \$5 25. (Now Ready.)

The ability of the author, and his practical skill in condensation, give assurance that this work will prove valuable not only to the student preparing for examination, but also to the practitioner desirous of obtaining within a moderate compass, a view of the existing condition of the various departments of science connected with medicine.

This work is a remarkably complete one in its way, and comes nearer to our idea of what a Conspectus should be than any we have yet seen. Prof. Hartshorne, with a commendable forethought, intrusted the preparation of many of the chapters on special subjects to experts, reserving only anatomy, physiology, and practice of medicine to himself. As a result we have every department worked up to the latest date and in a refreshingly concise and lucid manner. There are an immense amount of illustrations scattered throughout the work, and although they have often been seen before in the various works upon general and special subjects, yet they will be none the

less valuable to the beginner. Every medical student who desires a reliable refresher to his memory when the pressure of lectures and other college work crowds to preclude him from having an opportunity to drink deeper in the larger works, will find this one of the greatest utility. It is thoroughly trustworthy from beginning to end; and as we have before intimated, a remarkably truthful outline sketch of the present state of medical science. We could hardly expect it should be otherwise, however, under the charge of such a thorough medical scholar as the author has already proved himself to be.—*N. York Med. Record,* March 15, 1869.

LUDLOW (J. L.), M.D.,

A MANUAL OF EXAMINATIONS upon Anatomy, Physiology, Surgery, Practice of Medicine, Obstetrics, Materia Medica, Chemistry, Pharmacy, and Therapeutics. To which is added a Medical Formulary. Third edition, thoroughly revised and greatly extended and enlarged. With 370 illustrations. In one handsome royal 12mo. volume of 816 large pages, extra cloth, \$3 25; leather, \$3 75.

The arrangement of this volume in the form of question and answer renders it especially suitable for the office examination of students, and for those preparing for graduation.

TANNER (THOMAS HAWKES), M.D., &c.

A MANUAL OF CLINICAL MEDICINE AND PHYSICAL DIAGNOSIS. Third American from the Second London Edition. Revised and Enlarged by TILBURY FOX, M.D., Physician to the Skin Department in University College Hospital, &c. In one neat volume small 12mo., of about 375 pages, extra cloth, \$1 50. (Now Ready.)

This favorite little work has remained out of print for some years in consequence of the pressing engagements which have prevented the author from giving it the thorough revision which it required. The great advance which has taken place of late in the means and appliances for observation and diagnosis has necessitated a very considerable enlargement of the work, so that it now contains about one-half more matter than the last edition. The Laryngoscope, Ophthalmoscope, Sphygmograph, and Thermometer have received special attention. The chapter on the diagnostic indications afforded by the Urine has been much enlarged, and a section has been inserted on the administration of Chloroform. Special attention has been given to the medical anatomy of regions and organs, and much has been introduced relative to pericardial, endocardial, abdominal, and cerebro-spinal diseases. On every subject coming within its scope such additions have been made as seemed essential to bring the book on a level with the most advanced condition of medical knowledge; and it is hoped that it will continue to merit the very great favor with which it has hitherto been received.

GRAY (HENRY), F. R. S.,*Lecturer on Anatomy at St. George's Hospital, London*

ANATOMY, DESCRIPTIVE AND SURGICAL. The Drawings by H. V. CARTER, M. D., late Demonstrator on Anatomy at St. George's Hospital; the Dissections jointly by the AUTHOR and DR. CARTER. A new American, from the fifth enlarged and improved London edition. In one magnificent imperial octavo volume, of nearly 900 pages, with 465 large and elaborate engravings on wood. Price in extra cloth, \$6 00; leather, raised bands, \$7 00. (*Just Ready.*)

The author has endeavored in this work to cover a more extended range of subjects than is customary in the ordinary text-books, by giving not only the details necessary for the student, but also the application of those details in the practice of medicine and surgery, thus rendering it both a guide for the learner, and an admirable work of reference for the active practitioner. The engravings form a special feature in the work, many of them being the size of nature, nearly all original, and having the names of the various parts printed on the body of the cut, in place of figures of reference, with descriptions at the foot. They thus form a complete and splendid series, which will greatly assist the student in obtaining a clear idea of Anatomy, and will also serve to refresh the memory of those who may find in the exigencies of practice the necessity of recalling the details of the dissecting room; while combining, as it does, a complete Atlas of Anatomy, with a thorough treatise on systematic, descriptive, and applied Anatomy, the work will be found of essential use to all physicians who receive students in their offices, relieving both preceptor and pupil of much labor in laying the groundwork of a thorough medical education.

Notwithstanding its exceedingly low price, the work will be found, in every detail of mechanical execution, one of the handsomest that has yet been offered to the American profession; while the careful scrutiny of a competent anatomist has relieved it of whatever typographical errors existed in the English edition. A few notices of previous editions are subjoined.

Thus it is that book after book makes the labor of the student easier than before, and since we have seen Blanchard & Lea's new edition of Gray's Anatomy, certainly the finest work of the kind now extant, we would fain hope that the bugbear of medical students will lose half its horrors, and this necessary foundation of physiological science will be much facilitated and advanced.—*N. O. Med. News.*

The various points illustrated are marked directly on the structure; that is, whether it be muscle, process, artery, nerve, valve, etc. etc.—we say each point is distinctly marked by lettered engravings, so that the student perceives at once each point described as readily as if pointed out on the subject by the demonstrator. Most of the illustrations are thus rendered exceedingly satisfactory, and to the physician they serve to refresh the memory with great readiness

and with scarce a reference to the printed text. The surgical application of the various regions is also presented with force and clearness, impressing upon the student at each step of his research all the important relations of the structure demonstrated.—*Cincinnati Lancet.*

This is, we believe, the handsomest book on Anatomy as yet published in our language, and bids fair to become in a short time THE standard text-book of our colleges and studies. Students and practitioners will alike appreciate this book. We predict for it a bright career, and are fully prepared to endorse the statement of the *London Lancet*, that "We are not acquainted with any work in any language which can take equal rank with the one before us." Paper, printing, binding, all are excellent, and we feel that a grateful profession will not allow the publishers to go unrewarded.—*Nashville Med. and Surg. Journal.*

SMITH (HENRY H.), M.D., and HORNER (WILLIAM E.), M.D.,
Prof. of Surgery in the Univ. of Penna., &c. Late Prof. of Anatomy in the Univ. of Penna., &c.

AN ANATOMICAL ATLAS, illustrative of the Structure of the Human Body. In one volume, large imperial octavo, extra cloth, with about six hundred and fifty beautiful figures. \$4 50.

The plan of this Atlas, which renders it so peculiarly convenient for the student, and its superlative execution, have been already pointed out. We must congratulate the student upon the completion of this Atlas, as it is the most convenient work of

the kind that has yet appeared; and we must add, the very beautiful manner in which it is "got up" is so creditable to the country as to be flattering to our national pride.—*American Medical Journal.*

HARTSHORNE (HENRY), M.D.,
Professor of Hygiene, etc., in the University of Pennsylvania.

A HANDBOOK OF HUMAN ANATOMY AND PHYSIOLOGY, for the use of Students, with 176 illustrations. In one volume, royal 12mo. of 312 pages; extra cloth, \$1 75. (*Now Ready.*)

SHARPEY (WILLIAM), M.D., and QUAIN (JONES & RICHARD).

HUMAN ANATOMY. Revised, with Notes and Additions, by JOSEPH LEIDY, M.D., Professor of Anatomy in the University of Pennsylvania. Complete in two large octavo volumes, of about 1300 pages, with 511 illustrations; extra cloth, \$6 00.

The very low price of this standard work, and its completeness in all departments of the subject, should command for it a place in the library of all anatomical students.

ALLEN (J. M.), M.D.

THE PRACTICAL ANATOMIST; OR, THE STUDENT'S GUIDE IN THE DISSECTING ROOM. With 266 illustrations. In one very handsome royal 12mo volume, of over 600 pages; extra cloth, \$2 00.

One of the most useful works upon the subject ever written.—*Medical Examiner.*

WILSON (ERASMUS), F.R.S.

A SYSTEM OF HUMAN ANATOMY, General and Special. A new and revised American, from the last and enlarged English edition. Edited by W. H. GOBRECHT, M. D., Professor of General and Surgical Anatomy in the Medical College of Ohio. Illustrated with three hundred and ninety-seven engravings on wood. In one large and handsome octavo volume, of over 600 large pages; extra cloth, \$4 00; leather, \$5 00.

The publisher trusts that the well-earned reputation of this long-established favorite will be more than maintained by the present edition. Besides a very thorough revision by the author, it has been most carefully examined by the editor, and the efforts of both have been directed to introducing everything which increased experience in its use has suggested as desirable to render it a complete text-book for those seeking to obtain or to renew an acquaintance with Human Anatomy. The amount of additions which it has thus received may be estimated from the fact that the present edition contains over one-fourth more matter than the last, rendering a smaller type and an enlarged page requisite to keep the volume within a convenient size. The author has not only thus added largely to the work, but he has also made alterations throughout, wherever there appeared the opportunity of improving the arrangement or style, so as to present every fact in its most appropriate manner, and to render the whole as clear and intelligible as possible. The editor has exercised the utmost caution to obtain entire accuracy in the text, and has largely increased the number of illustrations, of which there are about one hundred and fifty more in this edition than in the last, thus bringing distinctly before the eye of the student everything of interest or importance.

BY THE SAME AUTHOR.

THE DISSECTOR'S MANUAL; OR, PRACTICAL AND SURGICAL ANATOMY. Third American, from the last revised and enlarged English edition. Modified and rearranged by WILLIAM HUNT, M. D., late Demonstrator of Anatomy in the University of Pennsylvania. In one large and handsome royal 12mo. volume, of 582 pages, with 154 illustrations; extra cloth, \$2 00.

HEATH (CHRISTOPHER), F.R.C.S.,
Teacher of Operative Surgery in University College, London.

PRACTICAL ANATOMY: A Manual of Dissections. From the Second revised and improved London edition. Edited, with additions, by W. W. KEEN, M. D., Lecturer on Pathological Anatomy in the Jefferson Medical College, Philadelphia. In one handsome volume, with over 200 illustrations. (In Press.)

HODGES, (RICHARD M.), M.D.,
Late Demonstrator of Anatomy in the Medical Department of Harvard University.

PRACTICAL DISSECTIONS. Second Edition, thoroughly revised. In one neat royal 12mo. volume, half-bound, \$2 00. (Just Issued.)

The object of this work is to present to the anatomical student a clear and concise description of that which he is expected to observe in an ordinary course of dissections. The author has endeavored to omit unnecessary details, and to present the subject in the form which many years' experience has shown him to be the most convenient and intelligible to the student. In the revision of the present edition, he has sedulously labored to render the volume more worthy of the favor with which it has heretofore been received.

MACLISE (JOSEPH).

SURGICAL ANATOMY. By JOSEPH MACLISE, Surgeon. In one volume, very large imperial quarto: with 68 large and splendid plates, drawn in the best style and beautifully colored, containing 190 figures, many of them the size of life; together with copious explanatory letter-press. Strongly and handsomely bound in extra cloth. Price \$14 00.

As no complete work of the kind has heretofore been published in the English language, the present volume will supply a want long felt in this country of an accurate and comprehensive Atlas of Surgical Anatomy, to which the student and practitioner can at all times refer to ascertain the exact relative positions of the various portions of the human frame towards each other and to the surface, as well as their abnormal deviations. Notwithstanding the large size, beauty and finish of the very numerous illustrations, it will be observed that the price is so low as to place it within the reach of all members of the profession.

We know of no work on surgical anatomy which can compete with it.—*Lancet*.

The work of MacLise on surgical anatomy is of the highest value. In some respects it is the best publication of its kind we have seen, and is worthy of a place in the library of any medical man, while the student could scarcely make a better investment than this.—*The Western Journal of Medicine and Surgery*.

No such lithographic illustrations of surgical regions have hitherto, we think, been given. While the operator is shown every vessel and nerve where an operation is contemplated, the exact anatomist is

refreshed by those clear and distinct dissections, which every one must appreciate who has a particle of enthusiasm. The English medical press has quite exhausted the words of praise, in recommending this admirable treatise. Those who have any curiosity to gratify, in reference to the perfectibility of the lithographic art in delineating the complex mechanism of the human body, are invited to examine our specimen copy. If anything will induce surgeons and students to patronize a book of such rare value and everyday importance to them, it will be a survey of the artistic skill exhibited in these fac-similes of nature.—*Boston Med. and Surg. Journal*.

HORNER'S SPECIAL ANATOMY AND HISTOLOGY. Eighth edition, extensively revised and modified.

In 2 vols. 8vo., of over 1000 pages, with more than 300 wood-cuts; extra cloth, \$6 00.

MARSHALL (JOHN), F. R. S.*Professor of Surgery in University College, London, &c.***OUTLINES OF PHYSIOLOGY, HUMAN AND COMPARATIVE.**

With Additions by FRANCIS GURNEY SMITH, M. D., Professor of the Institutes of Medicine in the University of Pennsylvania, &c. With numerous illustrations. In one large and handsome octavo volume, of 1026 pages, extra cloth, \$6 50; leather, raised bands \$7 50. (Just Issued.)

In fact, in every respect, Mr. Marshall has presented us with a most complete, reliable, and scientific work, and we feel that it is worthy our warmest commendation.—*St. Louis Med. Reporter*, Jan. 1869.

This is an elaborate and carefully prepared digest of human and comparative physiology, designed for the use of general readers, but more especially serviceable to the student of medicine. Its style is concise, clear, and scholarly; its order perspicuous and exact, and its range of topics extended. The author and his American editor have been careful to bring to the illustration of the subject the important discoveries of modern science in the various cognate departments of investigation. This is especially visible in the variety of interesting information derived from the departments of chemistry and physics. The great amount and variety of matter contained in the work is strikingly illustrated by turning over the copious index, covering twenty-four closely printed pages in double columns.—*Silliman's Journal*, Jan. 1869.

We doubt if there is in the English language any compend of physiology more useful to the student than this work.—*St. Louis Med. and Surg. Journal*, Jan. 1869.

It quite fulfils, in our opinion, the author's design of making it truly educational in its character—which is, perhaps, the highest commendation that can be asked.—*Am. Journ. Med. Sciences*, Jan. 1869.

We may now congratulate him on having completed the latest as well as the best summary of modern

physiological science, both human and comparative, with which we are acquainted. To speak of this work in the terms ordinarily used on such occasions would not be agreeable to ourselves, and would fail to do justice to its author. To write such a book requires a varied and wide range of knowledge, considerable power of analysis, correct judgment, skill in arrangement, and conscientious spirit. It must have entailed great labor, but now that the task has been fulfilled, the book will prove not only invaluable to the student of medicine and surgery, but serviceable to all candidates in natural science examinations, to teachers in schools, and to the lover of nature generally. In conclusion, we can only express the conviction that the merits of the work will command for it that success which the ability and vast labor displayed in its production so well deserve.—*London Lancet*, Feb. 22, 1868.

If the possession of knowledge, and peculiar aptitude and skill in expounding it, qualify a man to write an educational work, Mr. Marshall's treatise might be reviewed favorably without even opening the covers. There are few, if any, more accomplished anatomists and physiologists than the distinguished professor of surgery at University College; and he has long enjoyed the highest reputation as a teacher of physiology, possessing remarkable powers of clear exposition and graphic illustration. We have rarely the pleasure of being able to recommend a text-book so unreservedly as this.—*British Med. Journal*, Jan. 25, 1868.

CARPENTER (WILLIAM B.), M. D., F. R. S.*Examiner in Physiology and Comparative Anatomy in the University of London.***PRINCIPLES OF HUMAN PHYSIOLOGY; with their chief appli-**

cations to Psychology, Pathology, Therapeutics, Hygiene and Forensic Medicine. A new American from the last and revised London edition. With nearly three hundred illustrations. Edited, with additions, by FRANCIS GURNEY SMITH, M. D., Professor of the Institutes of Medicine in the University of Pennsylvania, &c. In one very large and beautiful octavo volume, of about 900 large pages, handsomely printed; extra cloth, \$5 50; leather, raised bands, \$6 50.

With Dr. Smith, we confidently believe "that the present will more than sustain the enviable reputation already attained by former editions, of being one of the fullest and most complete treatises on the subject in the English language." We know of none from the pages of which a satisfactory knowledge of the physiology of the human organism can be as well obtained, none better adapted for the use of such as take up the study of physiology in its reference to the institutes and practice of medicine.—*Am. Journ. Med. Sciences*.

We doubt not it is destined to retain a strong hold on public favor, and remain the favorite text-book in our colleges.—*Virginia Medical Journal*.

The above is the title of what is emphatically the great work on physiology; and we are conscious that it would be a useless effort to attempt to add anything to the reputation of this invaluable work, and can only say to all with whom our opinion has any influence, that it is our authority.—*Atlanta Med. Journal*.

BY THE SAME AUTHOR.**PRINCIPLES OF COMPARATIVE PHYSIOLOGY.** New American, from the Fourth and Revised London Edition. In one large and handsome octavo volume, with over three hundred beautiful illustrations. Pp. 752. Extra cloth, \$5 00.

As a complete and condensed treatise on its extended and important subject, this work becomes a necessity to students of natural science, while the very low price at which it is offered places it within the reach of all.

KIRKES (WILLIAM SENHOUSE), M. D.,**A MANUAL OF PHYSIOLOGY.** A new American from the third and improved London edition. With two hundred illustrations. In one large and handsome royal 12mo. volume. Pp. 586. Extra cloth, \$2 25; leather, \$2 75.

It is at once convenient in size, comprehensive in design, and concise in statement, and altogether well adapted for the purpose designed.—*St. Louis Med. and Surg. Journal*.

The physiological reader will find it a most excel-

lent guide in the study of physiology in its most advanced and perfect form. The author has shown himself capable of giving details sufficiently ample in a condensed and concentrated shape, on a science in which it is necessary at once to be correct and not lengthened.—*Edinburgh Med. and Surg. Journal*.

DALTON (J. C.), M.D.,*Professor of Physiology in the College of Physicians and Surgeons, New York, &c.*

A TREATISE ON HUMAN PHYSIOLOGY. Designed for the use of Students and Practitioners of Medicine. Fourth edition, revised, with nearly three hundred illustrations on wood. In one very beautiful octavo volume, of about 700 pages, extra cloth, \$5 25; leather, \$6 25. (Just Issued.)

From the Preface to the New Edition.

"The progress made by Physiology and the kindred Sciences during the last few years has required, for the present edition of this work, a thorough and extensive revision. This progress has not consisted in any very striking single discoveries, nor in a decided revolution in any of the departments of Physiology; but it has been marked by great activity of investigation in a multitude of different directions, the combined results of which have not failed to impress a new character on many of the features of physiological knowledge. . . . In the revision and correction of the present edition, the author has endeavored to incorporate all such improvements in physiological knowledge with the mass of the text in such a manner as not essentially to alter the structure and plan of the work, so far as they have been found adapted to the wants and convenience of the reader. . . . Several new illustrations are introduced, some of them as additions, others as improvements or corrections of the old. Although all parts of the book have received more or less complete revision, the greatest number of additions and changes were required in the Second Section, on the Physiology of the Nervous System."

The advent of the first edition of Prof. Dalton's Physiology, about eight years ago, marked a new era in the study of physiology to the American student. Under Dalton's skillful management, physiological science threw off the long, loose, negatively garments of probability and surmise, in which it had been arrayed by most artists, and came among us smiling and attractive, in the beautifully tinted and closely fitting dress of a demonstrated science. It was a stroke of genius, as well as a result of erudition and talent, that led Prof. Dalton to present to the world a work on physiology at once brief, pointed, and comprehensive, and which exhibited plainly in letter and drawings the basis upon which the conclusions arrived at rested. It is no disparagement of the many excellent works on physiology, published prior to that of Dalton, to say that none of them, either in plan of arrangement or clearness of execution, could be compared with his for the use of students or general practitioners of medicine. For this purpose his book has no equal in the English language.—*Western Journal of Medicine*, Nov. 1867.

A capital text-book in every way. We are, therefore, glad to see it in its fourth edition. It has already been examined at full length in these columns, so that we need not now further advert to it beyond remarking that both revision and enlargement have been most judicious.—*London Med. Times and Gazette*, Oct. 19, 1867.

No better proof of the value of this admirable work could be produced than the fact that it has already reached a fourth edition in the short space of eight years. Possessing in an eminent degree the

merits of clearness and condensation, and being fully brought up to the present level of Physiology, it is undoubtedly one of the most reliable text-books upon this science that could be placed in the hands of the medical student.—*Am. Journal Med. Sciences*, Oct. 1867.

Prof. Dalton's work has such a well-established reputation that it does not stand in need of any recommendation. Ever since its first appearance it has become the highest authority in the English language; and that it is able to maintain the enviable position which it has taken, the rapid exhaustion of the different successive editions is sufficient evidence. The present edition, which is the fourth, has been thoroughly revised, and enlarged by the incorporation of all the many important advances which have lately been made in this rapidly progressing science.—*N. Y. Med. Record*, Oct. 15, 1867.

As it stands, we esteem it the very best of the physiological text-books for the student, and the most concise reference and guide-book for the practitioner.—*N. Y. Med. Journal*, Oct. 1867.

The present edition of this now standard work fully sustains the high reputation of its accomplished author. It is not merely a reprint, but has been faithfully revised, and enriched by such additions as the progress of physiology has rendered desirable. Taken as a whole, it is unquestionably the most reliable and useful treatise on the subject that has been issued from the American press.—*Chicago Med. Journal*, Sept. 1867.

DUNGLISON (ROBLEY), M.D.,*Professor of Institutes of Medicine in Jefferson Medical College, Philadelphia.*

HUMAN PHYSIOLOGY. Eighth edition. Thoroughly revised and extensively modified and enlarged, with five hundred and thirty-two illustrations. In two large and handsomely printed octavo volumes of about 1500 pages, extra cloth. \$7 00.

LEHMANN (C. G.)

PHYSIOLOGICAL CHEMISTRY. Translated from the second edition by GEORGE E. DAY, M.D., F.R.S., &c., edited by R. E. ROGERS, M.D., Professor of Chemistry in the Medical Department of the University of Pennsylvania, with illustrations selected from Funke's Atlas of Physiological Chemistry, and an Appendix of plates. Complete in two large and handsome octavo volumes, containing 1200 pages, with nearly two hundred illustrations, extra cloth. \$6 00.

BY THE SAME AUTHOR.

MANUAL OF CHEMICAL PHYSIOLOGY. Translated from the German, with Notes and Additions, by J. CHESTON MORRIS, M.D., with an Introductory Essay on Vital Force, by Professor SAMUEL JACKSON, M.D., of the University of Pennsylvania. With illustrations on wood. In one very handsome octavo volume of 336 pages extra cloth. \$2 25.

TODD (ROBERT B.), M.D. F.R.S., and BOWMAN (W.), F.R.S.

THE PHYSIOLOGICAL ANATOMY AND PHYSIOLOGY OF MAN. With about three hundred large and beautiful illustrations on wood. Complete in one large octavo volume of 950 pages, extra cloth. Price \$4 75.

BRANDE (WM. T.), D. C. L., and TAYLOR (ALFRED S.), M. D., F. R. S.

CHEMISTRY. Second American edition, thoroughly revised by Dr. TAYLOR. In one handsome 8vo. volume of 764 pages, extra cloth, \$5 00; leather, \$6 00. (*Lately Issued.*)

FROM DR. TAYLOR'S PREFACE.

"The revision of the second edition, in consequence of the death of my lamented colleague, has devolved entirely upon myself. Every chapter, and indeed every page, has been revised, and numerous additions made in all parts of the volume. These additions have been restricted chiefly to subjects having some practical interest, and they have been made as concise as possible, in order to keep the book within those limits which may retain for it the character of a Student's Manual."—*London*, June 29, 1867.

A book that has already so established a reputation, as has Brande and Taylor's Chemistry, can hardly need a notice, save to mention the additions and improvements of the edition. Doubtless the work will long remain a favorite text-book in the schools, as well as a convenient book of reference for all.—*N. Y. Medical Gazette*, Oct. 12, 1867.

For this reason we hail with delight the republication, in a form which will meet with general approval and command public attention, of this really valuable standard work on chemistry—more particularly as it has been adapted with such care to the wants of the general public. The well known scholarship of its authors, and their extensive researches for many years in experimental chemistry, have been long appreciated in the scientific world, but in this work they have been careful to give the largest possible amount of information with the most sparing use of technical terms and phraseology, so as to furnish the reader, "whether a student of medicine, or a man of the world, with a plain introduction to the science and practice of chemistry."—*Journal of Applied Chemistry*, Oct. 1867.

This second American edition of an excellent treatise on chemical science is not a mere republication from the English press, but is a revision and enlargement of the original, under the supervision of the surviving author, Dr. Taylor. The favorable opinion expressed on the publication of the former edition of this work is fully sustained by the present revision, in which Dr. T. has increased the size of the volume, by an addition of sixty-eight pages.—*Am. Journ. Med. Sciences*, Oct. 1867.

THE HANDBOOK IN CHEMISTRY OF THE STUDENT.—For clearness of language, accuracy of description, extent of information, and freedom from pedantry and mysticism, no other text-book comes into competition with it.—*The Lancet*.

The authors set out with the definite purpose of writing a book which shall be intelligible to any educated man. Thus conceived, and worked out in the most sturdy, common-sense method, this book gives in the clearest and most summary method possible all the facts and doctrines of chemistry.—*Medical Times*.

ODLING (WILLIAM),

Lecturer on Chemistry, at St. Bartholomew's Hospital, &c.

A COURSE OF PRACTICAL CHEMISTRY, arranged for the Use of Medical Students. With Illustrations. From the Fourth and Revised London Edition. In one neat royal 12mo. volume, extra cloth. \$2. (*Just Issued.*)

As a work for the practitioner it cannot be excelled. It is written plainly and concisely, and gives in a very small compass the information required by the busy practitioner. It is essentially a work for the physician, and no one who purchases it will ever regret the outlay. In addition to all that is usually given in connection with inorganic chemistry, there are most valuable contributions to toxicology, animal and or-

ganic chemistry, etc. The portions devoted to a discussion of these subjects are very excellent. In no work can the physician find more that is valuable and reliable in regard to urine, bile, milk, bone, urinary calculi, tissue composition, etc. The work is small, reasonable in price, and well published.—*Richmond and Louisville Med. Journal*, Dec. 1869.

BOWMAN (JOHN E.), M. D.

PRACTICAL HANDBOOK OF MEDICAL CHEMISTRY. Edited by C. L. BLOXAM, Professor of Practical Chemistry in King's College, London. Fifth American, from the fourth and revised English Edition. In one neat volume, royal 12mo., pp. 351, with numerous illustrations, extra cloth. \$2 25. (*Now Ready.*)

The fourth edition of this invaluable text-book of Medical Chemistry was published in England in October of the last year. The Editor has brought down the Handbook to that date, introducing, as far as was compatible with the necessary conciseness of such a work, all the valuable discoveries in the science

which have come to light since the previous edition was printed. The work is indispensable to every student of medicine or enlightened practitioner. It is printed in clear type, and the illustrations are numerous and intelligible.—*Boston Med. and Surg. Journal*.

BY THE SAME AUTHOR.

INTRODUCTION TO PRACTICAL CHEMISTRY, INCLUDING ANALYSIS. Fifth American, from the fifth and revised London edition. With numerous illustrations. In one neat vol., royal 12mo., extra cloth. \$2 25. (*Now Ready.*)

One of the most complete manuals that has for a long time been given to the medical student.—*Athenæum*.

We regard it as realizing almost everything to be desired in an introduction to Practical Chemistry.

It is by far the best adapted for the Chemical student of any that has yet fallen in our way.—*British and Foreign Medico-Chirurgical Review*.

The best introductory work on the subject with which we are acquainted.—*Edinburgh Monthly Jour.*

GRAHAM (THOMAS), F. R. S.

THE ELEMENTS OF INORGANIC CHEMISTRY, including the Applications of the Science in the Arts. New and much enlarged edition, by HENRY WATTS and ROBERT BRIDGES, M. D. Complete in one large and handsome octavo volume, of over 800 very large pages, with two hundred and thirty-two wood-cuts, extra cloth. \$5 50.

KNAPP'S TECHNOLOGY; or Chemistry Applied to the Arts, and to Manufactures. With American additions, by Prof. WALTER R. JOHNSON. In two

very handsome octavo volumes, with 500 wood engravings, extra cloth, \$6 00.

FOWNES (GEORGE), Ph. D.

A MANUAL OF ELEMENTARY CHEMISTRY; Theoretical and Practical. With one hundred and ninety-seven illustrations. A new American, from the tenth and revised London edition. Edited by ROBERT BRIDGES, M. D. In one large royal 12mo. volume, of about 850 pp., extra cloth, \$2 75; leather, \$3 25. (*Just Issued.*)

Some years having elapsed since the appearance of the last American edition, and several revisions having been made of the work in England during the interval, it will be found very greatly altered, and enlarged by about two hundred and fifty pages, containing nearly one half more matter than before. The editors, Mr. Watts and Dr. Bence Jones, have labored sedulously to render it worthy in all respects of the very remarkable favor which it has thus far enjoyed, by incorporating in it all the most recent investigations and discoveries, in so far as is compatible with its design as an elementary text-book. While its distinguishing characteristics have been preserved, various portions have been rewritten, and especial pains have been taken with the department of Organic Chemistry in which late researches have accumulated so many new facts and have enabled the subject to be systematized and rendered intelligible in a manner formerly impossible. As only a few months have elapsed since the work thus passed through the hands of Mr. Watts and Dr. Bence Jones, but little has remained to be done by the American editor. Such additions as seemed advisable have however been made, and especial care has been taken to secure, by the closest scrutiny, the accuracy so essential in a work of this nature.

Thus fully brought up to a level with the latest advances of science, and presented at a price within the reach of all, it is hoped that the work will maintain its position as the favorite text-book of the medical student.

This work is so well known that it seems almost superfluous for us to speak about it. It has been a favorite text-book with medical students for years, and its popularity has in no respect diminished. Whenever we have been consulted by medical students, as has frequently occurred, what treatise on chemistry they should procure, we have always recommended Fownes', for we regarded it as the best. There is no work that combines so many excellences. It is of convenient size, not prolix, of plain perspicuous diction, contains all the most recent discoveries, and is of moderate price.—*Cincinnati Med. Repertory*, Aug. 1869.

Large additions have been made, especially in the department of organic chemistry, and we know of no other work that has greater claims on the physician, pharmacist, or student, than this. We cheerfully recommend it as the best text-book on elementary chemistry, and bespeak for it the careful attention of students of pharmacy.—*Chicago Pharmacist*, Aug. 1869.

The American reprint of the tenth revised and corrected English edition is now issued, and represents the present condition of the science. No comments are necessary to insure it a favorable reception at the hands of practitioners and students.—*Boston Med. and Surg. Journal*, Aug. 12, 1869.

It will continue, as heretofore, to hold the first rank as a text-book for students of medicine.—*Chicago Med. Examiner*, Aug. 1869.

This work, long the recognized Manual of Chemistry, appears as a tenth edition, under the able editorship of Bence Jones and Henry Watts. The chapter on

the General Principles of Chemical Philosophy, and the greater part of the organic chemistry, have been rewritten, and the whole work revised in accordance with the recent advances in chemical knowledge. It remains the standard text-book of chemistry.—*Dublin Quarterly Journal*, Feb. 1869.

There is probably not a student of chemistry in this country to whom the admirable manual of the late Professor Fownes is unknown. It has achieved a success which we believe is entirely without a parallel among scientific text-books in our language. This success has arisen from the fact that there is no English work on chemistry which combines so many excellences. Of convenient size, of attractive form, clear and concise in diction, well illustrated, and of moderate price, it would seem that every requisite for a student's hand-book has been attained. The ninth edition was published under the joint editorship of Dr. Bence Jones and Dr. Hofmann; the new one has been superintended through the press by Dr. Bence Jones and Mr. Henry Watts. It is not too much to say that it could not possibly have been in better hands. There is no one in England who can compare with Mr. Watts in experience as a compiler in chemical literature, and we have much pleasure in recording the fact that his reputation is well sustained by this, his last undertaking.—*The Chemical News*, Feb. 1869.

Here is a new edition which has been long watched for by eager teachers of chemistry. In its new garb, and under the editorship of Mr. Watts, it has resumed its old place as the most successful of text-books.—*Indian Medical Gazette*, Jan. 1, 1869.

ATTFIELD (JOHN), Ph. D.

Professor of Practical Chemistry to the Pharmaceutical Society of Great Britain, &c.

CHEMISTRY, GENERAL, MEDICAL, AND PHARMACEUTICAL; Including the Chemistry of the British Pharmacopœia. Adapted to the Requirements of American Practice of Medicine and Pharmacy. In one handsome volume. (*Preparing.*)

It contains a most admirable digest of what is specially needed by the medical student in all that relates to practical chemistry, and constitutes for him a sound and useful text-book on the subject. . . . We commend it to the notice of every medical, as well as pharmaceutical, student. We only regret that we had not the book to depend upon in working up the subject of practical and pharmaceutical chemistry for the University of London, for which it seems to us that it is exactly adapted. This is paying the book a high compliment.—*The Lancet*.

Dr. Attfield's book is written in a clear and able manner; it is a work *sui generis* and without a rival; it will be welcomed, we think, by every reader of the 'Pharmacopœia,' and is quite as well suited for the medical student as for the pharmacist.—*The Chemical News*.

A valuable guide to practical medical chemistry, and an admirable companion to the 'British Pharmacopœia.' It is rare to find so many qualities combined, and quite curious to note how much valuable information finds a mutual interdependence.—*Medical Times and Gazette*.

It is almost the only book from which the medical student can work up the pharmacopœial chemistry required at his examinations.—*The Pharmaceutical Journal*.

At page 350 of the current volume of this journal, we remarked that "there is a sad dearth of [medical] students' text-books in chemistry." Dr. Attfield's volume, just published, is rather a new book than a second edition of his previous work, and more nearly realizes our ideal than any book we have before seen on the subject.—*The British Medical Journal*.

The introduction of new matter has not destroyed the original character of the work, as a treatise on pharmaceutical and medical chemistry, but has simply extended the foundations of these special departments of the science.—*The Chemist and Druggist*.

We believe that this manual has been already adopted as the class-book by many of the professors in the public schools throughout the United Kingdom. . . . In pharmaceutical chemistry applied to the pharmacopœia, we know of no rival. It is, therefore, particularly suited to the medical student.—*The Medical Press and Circular*.

PARRISH (EDWARD),*Professor of Materia Medica in the Philadelphia College of Pharmacy.*

A TREATISE ON PHARMACY. Designed as a Text-Book for the Student, and as a Guide for the Physician and Pharmaceutist. With many Formulæ and Prescriptions. Third Edition, greatly improved. In one handsome octavo volume, of 850 pages, with several hundred illustrations, extra cloth. \$5 00.

The immense amount of practical information condensed in this volume may be estimated from the fact that the Index contains about 4700 items. Under the head of Acids there are 312 references; under Emplastrum, 36; Extracts, 159; Lozenges, 25; Mixtures, 55; Pills, 56; Syrups, 131; Tinctures, 138; Unguentum, 57, &c.

We have examined this large volume with a good deal of care, and find that the author has completely exhausted the subject upon which he treats; a more complete work, we think, it would be impossible to find. To the student of pharmacy the work is indispensable; indeed, so far as we know, it is the only one of its kind in existence, and even to the physician or medical student who can spare five dollars to purchase it, we feel sure the practical information he will obtain will more than compensate him for the outlay.—*Canada Med. Journal*, Nov. 1864.

The medical student and the practising physician will find the volume of inestimable worth for study and reference.—*San Francisco Med. Press*, July, 1864.

When we say that this book is in some respects the best which has been published on the subject in the English language for a great many years, we do

not wish it to be understood as very extravagant praise. In truth, it is not so much the best as the only book.—*The London Chemical News*.

An attempt to furnish anything like an analysis of Parrish's very valuable and elaborate *Treatise on Practical Pharmacy* would require more space than we have at our disposal. This, however, is not so much a matter of regret, inasmuch as it would be difficult to think of any point, however minute and apparently trivial, connected with the manipulation of pharmaceutical substances or appliances which has not been clearly and carefully discussed in this volume. Want of space prevents our enlarging further on this valuable work, and we must conclude by a simple expression of our hearty appreciation of its merits.—*Dublin Quarterly Jour. of Medical Science*, August, 1864.

STILLÉ (ALFRED), M.D.,*Professor of Theory and Practice of Medicine in the University of Penna.*

THERAPEUTICS AND MATERIA MEDICA; a Systematic Treatise on the Action and Uses of Medicinal Agents, including their Description and History. Third edition, revised and enlarged. In two large and handsome octavo volumes of about 1700 pages, extra cloth, \$10; leather, \$12. (*Just Issued.*)

Dr. Stillé's splendid work on therapeutics and materia medica.—*London Med. Times*, April 8, 1865.

Dr. Stillé stands to-day one of the best and most honored representatives at home and abroad, of American medicine; and these volumes, a library in themselves, a treasure-house for every studious physician, assure his fame even had he done nothing more.—*The Western Journal of Medicine*, Dec. 1868.

We regard this work as the best one on Materia Medica in the English language, and as such it deserves the favor it has received.—*Am. Journ. Medical Sciences*, July 1868.

We need not dwell on the merits of the third edition of this magnificently conceived work. It is the work on Materia Medica, in which Therapeutics are primarily considered—the mere natural history of drugs being briefly disposed of. To medical practitioners this is a very valuable conception. It is wonderful how much of the riches of the literature of Materia Medica has been condensed into this book. The references alone would make it worth possessing. But it is not a mere compilation. The writer exercises a good judgment of his own on the great doctrines and points of Therapeutics. For purposes of practice, Stillé's book is almost unique as a repository of information, empirical and scientific, on the actions and uses of medicines.—*London Lancet*, Oct. 31, 1868.

Through the former editions, the professional world is well acquainted with this work. At home and

abroad its reputation as a standard treatise on Materia Medica is securely established. It is second to no work on the subject in the English tongue, and, indeed, is decidedly superior, in some respects, to any other.—*Pacific Med. and Surg. Journal*, July, 1868.

Stillé's Therapeutics is incomparably the best work on the subject.—*N. Y. Med. Gazette*, Sept. 26, 1868.

Dr. Stillé's work is becoming the best known of any of our treatises on Materia Medica. . . . One of the most valuable works in the language on the subjects of which it treats.—*N. Y. Med. Journal*, Oct. 1868.

The rapid exhaustion of two editions of Prof. Stillé's scholarly work, and the consequent necessity for a third edition, is sufficient evidence of the high estimate placed upon it by the profession. It is no exaggeration to say that there is no superior work upon the subject in the English language. The present edition is fully up to the most recent advance in the science and art of therapeutics.—*Leavenworth Medical Herald*, Aug. 1868.

The work of Prof. Stillé has rapidly taken a high place in professional esteem, and to say that a third edition is demanded and now appears before us, sufficiently attests the firm position this treatise has made for itself. As a work of great research, and scholarship, it is safe to say we have nothing superior. It is exceedingly full, and the busy practitioner will find ample suggestions upon almost every important point of therapeutics.—*Cincinnati Lancet*, Aug. 1868.

GRIFFITH (ROBERT E.), M.D.

A UNIVERSAL FORMULARY, Containing the Methods of Preparing and Administering Official and other Medicines. The whole adapted to Physicians and Pharmaceutists. Second edition, thoroughly revised, with numerous additions, by ROBERT P. THOMAS, M.D., Professor of Materia Medica in the Philadelphia College of Pharmacy. In one large and handsome octavo volume of 650 pages, double-columns. Extra cloth, \$4 00; leather, \$5 00.

Three complete and extended Indexes render the work especially adapted for immediate consultation. One, of DISEASES AND THEIR REMEDIES, presents under the head of each disease the remedial agents which have been usefully exhibited in it, with reference to the formulæ containing them—while another of PHARMACEUTICAL AND BOTANICAL NAMES, and a very thorough GENERAL INDEX afford the means of obtaining at once any information desired. The Formulary itself is arranged alphabetically, under the heads of the leading constituents of the prescriptions.

We know of none in our language, or any other, so comprehensive in its details.—*London Lancet*.

One of the most complete works of the kind in any language.—*Edinburgh Med. Journal*.

We are not cognizant of the existence of a parallel work.—*London Med. Gazette*.

PEREIRA (JONATHAN), M.D., F.R.S. and L.S.

MATERIA MEDICA AND THERAPEUTICS; being an Abridgment of the late Dr. Pereira's Elements of Materia Medica, arranged in conformity with the British Pharmacopœia, and adapted to the use of Medical Practitioners, Chemists and Druggists, Medical and Pharmaceutical Students, &c. By F. J. FARRE, M.D., Senior Physician to St. Bartholomew's Hospital, and London Editor of the British Pharmacopœia; assisted by ROBERT BENTLEY, M.R.C.S., Professor of Materia Medica and Botany to the Pharmaceutical Society of Great Britain; and by ROBERT WARINGTON, F.R.S., Chemical Operator to the Society of Apothecaries. With numerous additions and references to the United States Pharmacopœia, by HORATIO C. WOOD, M.D., Professor of Botany in the University of Pennsylvania. In one large and handsome octavo volume of 1040 closely printed pages, with 236 illustrations, extra cloth, \$7 00; leather, raised bands, \$8 00. (*Latest Published.*)

The task of the American editor has evidently been no sinecure, for not only has he given to us all that is contained in the abridgment useful for our purposes, but by a careful and judicious embodiment of over a hundred new remedies has increased the size of the former work fully one-third, besides adding many new illustrations, some of which are original. We unhesitatingly say that by so doing he has proportionately increased the value, not only of the condensed edition, but has extended the applicability of the great original, and has placed his medical countrymen under lasting obligations to him. The American physician now has all that is needed in the shape of a complete treatise on materia medica, and the medical student has a text-book which, for practical utility and intrinsic worth, stands unparalleled. Although of considerable size, it is none too large for the purposes for which it has been intended, and every medical man should, in justice to himself, spare a place for it upon his book-shelf, resting assured that the more he consults it the better he will be satisfied of its excellence.—*N. Y. Med. Record*, Nov. 15, 1866.

It will fill a place which no other work can occupy in the library of the physician, student, and apothecary.—*Boston Med. and Surg. Journal*, Nov. 8, 1866.

Of the many works on Materia Medica which have appeared since the issuing of the British Pharmacopœia, none will be more acceptable to the student and practitioner than the present. Pereira's Materia Medica had long ago asserted for itself the position of being the most complete work on the subject in the English language. But its very completeness stood in the way of its success. Except in the way of reference, or to those who made a special study of Materia Medica, Dr. Pereira's work was too full, and its perusal required an amount of time which few had at their disposal. Dr. Farre has very judiciously availed himself of the opportunity of the publication of the new Pharmacopœia, by bringing out an abridged edition of the great work. This edition of Pereira is by no means a mere abridged re-issue, but contains many improvements, both in the descriptive and therapeutical departments. We can recommend it as a very excellent and reliable text-book.—*Edinburgh Med. Journal*, February, 1866.

The reader cannot fail to be impressed, at a glance, with the exceeding value of this work as a compend of nearly all useful knowledge on the materia medica. We are greatly indebted to Professor Wood for his adaptation of it to our meridian. Without his emendations and additions it would lose much of its value to the American student. With them it is an American book.—*Pacific Medical and Surgical Journal*, December, 1866.

ELLIS (BENJAMIN), M.D.

THE MEDICAL FORMULARY: being a Collection of Prescriptions derived from the writings and practice of many of the most eminent physicians of America and Europe. Together with the usual Dietetic Preparations and Antidotes for Poisons. The whole accompanied with a few brief Pharmaceutic and Medical Observations. Twelfth edition, carefully revised and much improved by ALBERT H. SMITH, M.D. In one volume 8vo. of 376 pages, extra cloth, \$3 00. (*Now Ready.*)

This work has remained for some time out of print, owing to the anxious care with which the Editor has sought to render the present edition worthy a continuance of the very remarkable favor which has carried the volume to the unusual honor of a TWELFTH EDITION. He has sedulously endeavored to introduce in it all new preparations and combinations deserving of confidence, besides adding two new classes, Antemetics and Disinfectants, with brief references to the inhalation of atomized fluids, the nasal douche of Thudichum, suggestions upon the method of hypodermic injection, the administration of anæsthetics, &c. &c. To accommodate these numerous additions, he has omitted much which the advance of science has rendered obsolete or of minor importance, notwithstanding which the volume has been increased by more than thirty pages. A new feature will be found in a copious Index of Diseases and their remedies, which cannot but increase the value of the work as a suggestive book of reference for the working practitioner. Every precaution has been taken to secure the typographical accuracy so necessary in a work of this nature, and it is hoped that the new edition will fully maintain the position which "ELLIS' FORMULARY" has long occupied.

CARSON (JOSEPH), M.D.,

Professor of Materia Medica and Pharmacy in the University of Pennsylvania, &c.

SYNOPSIS OF THE COURSE OF LECTURES ON MATERIA MEDICA AND PHARMACY, delivered in the University of Pennsylvania. With three Lectures on the Modus Operandi of Medicines. Fourth and revised edition, extra cloth, \$3 00. (*Just Issued.*)

DUNGLISON'S NEW REMEDIES, WITH FORMULÆ FOR THEIR PREPARATION AND ADMINISTRATION. Seventh edition, with extensive additions. One vol. 8vo., pp. 770; extra cloth. \$4 00.

ROYLE'S MATERIA MEDICA AND THERAPEUTICS. Edited by JOSEPH CARSON, M.D. With ninety-eight illustrations. 1 vol. 8vo., pp. 700, extra cloth. \$3 00.

CHRISTISON'S DISPENSATORY. With copious additions, and 213 large wood-engravings. By R. EGGESFIELD GRIFFITH, M.D. One vol. 8vo., pp. 1000; extra cloth. \$4 00.

CARPENTER'S PRIZE ESSAY ON THE USE OF ALCOHOLIC LIQUORS IN HEALTH AND DISEASE. New edition, with a Preface by D. F. CONDIE, M.D., and explanations of scientific words. In one neat 12mo. volume, pp. 178, extra cloth. 60 cents.

DE JONGH ON THE THREE KINDS OF COD-LIVER OIL, with their Chemical and Therapeutic Properties. 1 vol. 12mo., cloth. 75 cents.

MAYNE'S DISPENSATORY AND THERAPEUTICAL REMEMBRANCER. By R. E. GRIFFITH, M.D. In one 12mo. volume, 300 pp., extra cloth. 75 cents.

GROSS (SAMUEL D.), M. D.,
Professor of Surgery in the Jefferson Medical College of Philadelphia.

ELEMENTS OF PATHOLOGICAL ANATOMY. Third edition, thoroughly revised and greatly improved. In one large and very handsome octavo volume of nearly 800 pages, with about three hundred and fifty beautiful illustrations, of which a large number are from original drawings; extra cloth. \$4 00.

The very beautiful execution of this valuable work, and the exceedingly low price at which it is offered, should command for it a place in the library of every practitioner.

To the student of medicine we would say that we know of no work which we can more heartily commend than Gross's Pathological Anatomy.—*Southern Med. and Surg. Journal.*

The volume commends itself to the medical student; it will repay a careful perusal, and should be upon

the book-shelf of every American physician.—*Charleston Med. Journal.*

It contains much new matter, and brings down our knowledge of pathology to the latest period.—*London Lancet.*

JONES (C. HANDFIELD), F.R.S., and SIEVEKING (ED. H.), M.D.,
Assistant Physicians and Lecturers in St. Mary's Hospital.

A MANUAL OF PATHOLOGICAL ANATOMY. First American edition, revised. With three hundred and ninetyseven handsome wood engravings. In one large and beautifully printed octavo volume of nearly 750 pages, extra cloth, \$3 50.

Our limited space alone restrains us from noticing more at length the various subjects treated of in this interesting work; presenting, as it does, an excellent summary of the existing state of knowledge in relation to pathological anatomy, we cannot too strongly urge upon the student the necessity of a thorough acquaintance with its contents.—*Medical Examiner.*

We have long had need of a hand-book of pathological anatomy which should thoroughly reflect the present state of that science. In the treatise before us this desideratum is supplied. Within the limits of a moderate octavo, we have the outlines of this great department of medical science accurately defined,

and the most recent investigations presented in sufficient detail for the student of pathology. We cannot at this time undertake a formal analysis of this treatise, as it would involve a separate and lengthy consideration of nearly every subject discussed; nor would such analysis be advantageous to the medical reader. The work is of such a character that every physician ought to obtain it, both for reference and study.—*N. Y. Journal of Medicine.*

Its importance to the physician cannot be too highly estimated, and we would recommend our readers to add it to their library as soon as they conveniently can.—*Montreal Med. Chronicle.*

GLUGE'S ATLAS OF PATHOLOGICAL HISTOLOGY.

Translated, with Notes and Additions, by JOSEPH LEIDY, M.D. In one volume, very large imperial quarto, with 320 copper-plate figures, plain and colored, extra cloth. \$4 00.

SIMON'S GENERAL PATHOLOGY, as conducive to the Establishment of Rational Principles for the Prevention and Cure of Disease. In one octavo volume of 212 pages, extra cloth. \$1 25.

WILLIAMS (CHARLES J. B.), M.D.,
Professor of Clinical Medicine in University College, London.

PRINCIPLES OF MEDICINE. An Elementary View of the Causes, Nature, Treatment, Diagnosis, and Prognosis of Disease; with brief remarks on Hygienics, or the preservation of health. A new American, from the third and revised London edition. In one octavo volume of about 500 pages, extra cloth. \$3 50.

No work has ever achieved or maintained a more deserved reputation.—*Virginia Med. and Surg. Journal.*

There is no work in medical literature which can fill the place of this one. It is the *Primer* of the young practitioner, the *Koran* of the scientific one.—*Stethoscope.*

A text-book to which no other in our language is comparable.—*Charleston Med. Journal.*

The absolute necessity of such a work must be evident to all who pretend to more than mere empiricism. We must conclude by again expressing our high sense of the immense benefit which Dr. Williams has conferred on medicine by the publication of this work. We are certain that in the present state of our knowledge his Principles of Medicine could not possibly be surpassed.—*London Jour. of Medicine.*

HARRISON'S ESSAY TOWARDS A CORRECT THEORY OF THE NERVOUS SYSTEM. In one octavo volume of 292 pp. \$1 50.

SOLLY ON THE HUMAN BRAIN; its Structure, Physiology, and Diseases. From the Second and much enlarged London edition. In one octavo volume of 500 pages, with 120 wood-cuts; extra cloth. \$2 50.

LA ROCHE ON YELLOW FEVER, considered in its Historical, Pathological, Etiological, and Therapeutical Relations. In two large and handsome octavo volumes, of nearly 1500 pages, extra cloth, \$7 00.

LA ROCHE ON PNEUMONIA; its Supposed Connection, Pathological, and Etiological, with Autumnal

Fevers, including an Inquiry into the Existence and Morbid Agency of Malaria. In one handsome octavo volume, extra cloth, of 500 pages. Price \$3 00.

BUCKLER ON FIBRO-BRONCHITIS AND RHEUMATIC PNEUMONIA. In one octavo vol., extra cloth, pp. 150. \$1 25.

FISKE FUND PRIZE ESSAYS.—LEE ON THE EFFECTS OF CLIMATE ON TUBERCULOUS DISEASE. AND WARREN ON THE INFLUENCE OF PREGNANCY ON THE DEVELOPMENT OF TUBERCLES. Together in one neat octavo volume extra cloth. \$1 00.

BARCLAY (A. W.), M.D.

A MANUAL OF MEDICAL DIAGNOSIS; being an Analysis of the Signs and Symptoms of Disease. Third American from the second and revised London edition. In one neat octavo volume of 451 pages, extra cloth. \$3 50.

A work of immense practical utility.—*London Med. Times and Gazette.*

The book should be in the hands of every practical man.—*Dublin Med. Press.*

FLINT (AUSTIN). M. D..*Professor of the Principles and Practice of Medicine in Bellevue Med. College, N. Y.*

A TREATISE ON THE PRINCIPLES AND PRACTICE OF MEDICINE: designed for the use of Students and Practitioners of Medicine. Third edition, revised and enlarged. In one large and closely printed octavo volume of 1002 pages; handsome extra cloth, \$5 00; or strongly bound in leather, with raised bands, \$7 00. (*Just Issued.*)

From the Preface to the Third Edition.

Since the publication, in December, 1866, of the second edition of this treatise, much time has been devoted to its revision. Recognizing in the favor with which it has been received a proportionate obligation to strive constantly to increase its worthiness, the author has introduced in the present edition additions, derived from his clinical studies, and from the latest contributions in medical literature, which, it is believed, will enhance considerably the practical utility of the work. A slight modification in the typographical arrangement has accommodated these additions without materially increasing the bulk of the volume.

NEW YORK, October, 1868.

At the very low price affixed, the profession will find this to be one of the cheapest volumes within their reach.

This work, which stands pre-eminently as the advance standard of medical science up to the present time in the practice of medicine, has for its author one who is well and widely known as one of the leading practitioners of this continent. In fact, it is seldom that any work is ever issued from the press more deserving of universal recommendation.—*Dominion Med. Journal*, May, 1869.

The third edition of this most excellent book scarcely needs any commendation from us. The volume, as it stands now, is really a marvel: first of all, it is excellently printed and bound—and we encounter that luxury of America, the ready-cut pages, which the Yankees are 'cute enough to insist upon—nor are these by any means trifles; but the contents of the book are astonishing. Not only is it wonderful that any one man can have grasped in his mind the whole scope of medicine with that vigor which Dr. Flint shows, but the condensed yet clear way in which this is done is a perfect literary triumph. Dr. Flint is pre-eminently one of the strong men, whose right to do this kind of thing is well admitted; and we say no more than the truth when we affirm that he is very nearly the only living man that could do it with such results as the volume before us.—*The London Practitioner*, March, 1869.

This is in some respects the best text-book of medicine in our language, and it is highly appreciated on the other side of the Atlantic, inasmuch as the first edition was exhausted in a few months. The second edition was little more than a reprint, but the present has, as the author says, been thoroughly revised. Much valuable matter has been added, and by making the type smaller, the bulk of the volume is not much increased. The weak point in many American works is pathology, but Dr. Flint has taken peculiar pains on this point, greatly to the value of the book.—*London Med. Times and Gazette*, Feb. 6, 1869.

Published in 1866, this valuable book of Dr. Flint's has in two years exhausted two editions, and now we gladly announce a third. We say we gladly announce it, because we are proud of it as a national representative work of not only American, but of

cosmopolitan medicine. In it the practice of medicine is young and philosophical, based on reason and common sense, and as such, we hope it will be at the right hand of every practitioner of this vast continent.—*California Medical Gazette*, March, 1869.

Considering the large number of valuable works in the practice of medicine, already before the profession, the marked favor with which this has been received, necessitating a third edition in the short space of two years, indicates unmistakably that it is a work of more than ordinary excellence, and must be accepted as evidence that it has largely fulfilled the object for which the author intended it. A marked feature in the work, and one which particularly adapts it for the use of students as a text-book, and certainly renders it none the less valuable to the busy practitioner as a work of reference, is brevity and simplicity. The present edition has been thoroughly revised, and much new matter incorporated, derived, as the author informs us, both from his own clinical studies, and from the latest contributions to medical literature, thus bringing it fully up with the most recent advances of the science, and greatly enhancing its practical utility; while, by a slight modification of its typographical arrangement, the additions have been accommodated without materially increasing its bulk.—*St. Louis Med. Archives*, Feb. 1869.

If there be among our readers any who are not familiar with the treatise before us, we shall do them a service in persuading them to repair their omission forthwith. Combining to a rare degree the highest scientific attainments with the most practical common sense, and the closest habits of observation, the author has given us a volume which not only sets forth the results of the latest investigations of other laborers, but contains more original views than any other single work upon this well-worn theme within our knowledge.—*N. Y. Med. Gazette*, Feb. 27, 1869.

Practical medicine was at sea when this book appeared above the horizon as a safe and capacious harbor. It came opportunely and was greeted with pleasurable emotions throughout the land.—*Nashville Med. and Surg. Journal*, May, 1869.

DUNGLISON, FORBES, TWEEDIE, AND CONOLLY.

THE CYCLOPEDIA OF PRACTICAL MEDICINE: comprising

Treatises on the Nature and Treatment of Diseases, Materia Medica and Therapeutics, Diseases of Women and Children, Medical Jurisprudence, &c. &c. In four large super-royal octavo volumes, of 13254 double-columned pages, strongly and handsomely bound in leather, \$15; extra cloth, \$11.

*** This work contains no less than four hundred and eighteen distinct treatises, contributed by sixty-eight distinguished physicians.

The most complete work on practical medicine extant, or at least in our language.—*Buffalo Medical and Surgical Journal*.

For reference, it is above all price to every practitioner.—*Western Lancet*.

One of the most valuable medical publications of

the day. As a work of reference it is invaluable.—*Western Journal of Medicine and Surgery*.

It has been to us, both as learner and teacher, a work for ready and frequent reference, one in which modern English medicine is exhibited in the most advantageous light.—*Medical Examiner*.

BARLOW'S MANUAL OF THE PRACTICE OF MEDICINE. With Additions by D. F. CONDIE, M. D. 1 vol. 8vo., pp. 609, cloth. \$2 50.

HOLLAND'S MEDICAL NOTES AND REFLECTIONS. From the third and enlarged English edition. In one handsome octavo volume of about 500 pages, extra cloth. \$3 50.

HARTSHORNE (HENRY), M.D.,
Professor of Hygiene in the University of Pennsylvania.

ESSENTIALS OF THE PRINCIPLES AND PRACTICE OF MEDICINE. A handy-book for Students and Practitioners. Second edition, revised and improved. In one handsome royal 12mo. volume of 450 pages, clearly printed on small type, cloth, \$2 38; half bound, \$2 63. (*Just Issued.*)

The very cordial reception with which this work has met shows that the author has fully succeeded in his attempt to condense within a convenient compass the essential points of scientific and practical medicine, so as to meet the wants not only of the student, but also of the practitioner who desires to acquaint himself with the results of recent advances in medical science.

As a strikingly terse, full, and comprehensive embodiment in a condensed form of the essentials in medical science and art, we hazard nothing in saying that it is incomparably in advance of any work of the kind of the past, and will stand long in the future without a rival. A mere glance will, we think, impress others with the correctness of our estimate. Nor do we believe there will be found many who, after the most cursory examination, will fail to possess it. How one could be able to crowd so much that is valuable, especially to the student and young practitioner, within the limits of so small a book, and yet embrace and present all that is important in a well-arranged, clear form, convenient, satisfactory for reference, with so full a table of contents, and extended general index, with nearly three hundred formulas and recipes, is a marvel.—*Western Journal of Medicine*, Aug. 1867.

The little book before us has this quality, and we can therefore say that all students will find it an invaluable guide in their pursuit of clinical medicine. Dr. Hartshorne speaks of it as "an unambitious effort to make useful the experience of twenty years of private and hospital medical practice, with its attendant study and reflection." That the effort will prove successful we have no doubt, and in his study, and at the bedside, the student will find Dr. Hartshorne a

safe and accomplished companion. We speak thus highly of the volume, because it approaches more nearly than any similar manual lately before us the standard at which all such books should aim—of teaching much, and suggesting more. To the student we can heartily recommend the work of our transatlantic colleague, and the busy practitioner, we are sure, will find in it the means of solving many a doubt, and will rise from the perusal of its pages, having gained clearer views to guide him in his daily struggle with disease.—*Dub. Med. Press*, Oct. 2, 1867.

This work of Dr. Hartshorne must not be confounded with the medical manuals so generally to be found in the hands of students, serving them at best but as blind guides, better adapted to lead them astray than to any useful and reliable knowledge. The work before us presents a careful synopsis of the essential elements of the theory of diseased action, its causes, phenomena, and results, and of the art of healing, as recognized by the most authoritative of our professional writers and teachers. A very careful and candid examination of the volume has convinced us that it will be generally recognized as one of the best manuals for the use of the student that has yet appeared.—*American Journal Med. Sciences*, Oct. 1867.

WATSON (THOMAS), M.D., &c.

LECTURES ON THE PRINCIPLES AND PRACTICE OF PHYSIC. Delivered at King's College, London. A new American, from the last revised and enlarged English edition, with Additions, by D. FRANCIS CONDIE, M.D., author of "A Practical Treatise on the Diseases of Children," &c. With one hundred and eighty-five illustrations on wood. In one very large and handsome volume, imperial octavo, of over 1200 closely printed pages in small type; extra cloth, \$6 50; strongly bound in leather, with raised bands, \$7 50.

Believing this to be a work which should lie on the table of every physician, and be in the hands of every student, every effort has been made to condense the vast amount of matter which it contains within a convenient compass, and at a very reasonable price, to place it within reach of all. In its present enlarged form, the work contains the matter of at least three ordinary octavos, rendering it one of the cheapest works now offered to the American profession, while its mechanical execution makes it an exceedingly attractive volume.

DICKSON'S ELEMENTS OF MEDICINE; a Compendious View of Pathology and Therapeutics, or the History and Treatment of Diseases. Second edition, revised. 1 vol. 8vo. of 750 pages, extra cloth. \$4 00.

WHAT TO OBSERVE AT THE BEDSIDE AND AFTER DEATH IN MEDICAL CASES. Published under the authority of the London Society for Medical Obser-

vation. From the second London edition. 1 vol. royal 12mo., extra cloth. \$1 00.

LAYCOCK'S LECTURES ON THE PRINCIPLES AND METHODS OF MEDICAL OBSERVATION AND RESEARCH. For the use of advanced students and junior practitioners. In one very neat royal 12mo. volume, extra cloth. \$1 00.

FULLER (HENRY WILLIAM), M.D.,
Physician to St. George's Hospital, London.

ON DISEASES OF THE LUNGS AND AIR-PASSAGES. Their Pathology, Physical Diagnosis, Symptoms, and Treatment. From the second and revised English edition. In one handsome octavo volume of about 500 pages, extra cloth, \$3 50. (*Just Issued.*)

Dr. Fuller's work on diseases of the chest was so favorably received, that to many who did not know the extent of his engagements, it was a matter of wonder that it should be allowed to remain three years out of print. Determined, however, to improve it, Dr. Fuller would not consent to a mere reprint, and

accordingly we have what might be with perfect justice styled an entirely new work from his pen, the portion of the work treating of the heart and great vessels being excluded. Nevertheless, this volume is of almost equal size with the first.—*London Medical Times and Gazette*, July 26, 1867.

BRINTON (WILLIAM), M.D., F.R.S.

LECTURES ON THE DISEASES OF THE STOMACH; with an Introduction on its Anatomy and Physiology. From the second and enlarged London edition. With illustrations on wood. In one handsome octavo volume of about 300 pages, extra cloth. \$3 25. (*Just issued.*)

Nowhere can be found a more full, accurate, plain, and instructive history of these diseases, or more rational views respecting their pathology and therapeutics.—*Am. Journ. of the Med. Sciences*, April, 1865.

The most complete work in our language upon the diagnosis and treatment of these puzzling and important diseases.—*Boston Med. and Surg. Journal*, Nov. 1865.

FLINT (AUSTIN), M. D.,*Professor of the Principles and Practice of Medicine in Bellevue Hospital Med. College, N. Y.***A PRACTICAL TREATISE ON THE DIAGNOSIS, PATHOLOGY, AND TREATMENT OF DISEASES OF THE HEART.** Second revised and enlarged edition. In one octavo volume of 550 pages, with a plate, extra cloth, \$4. (*Now Ready.*)

The author has sedulously improved the opportunity afforded him of revising this work. Portions of it have been rewritten, and the whole brought up to a level with the most advanced condition of science. It must therefore continue to maintain its position as the standard treatise on the subject.

Dr. Flint chose a difficult subject for his researches, and has shown remarkable powers of observation and reflection, as well as great industry, in his treatment of it. His book must be considered the fullest and clearest practical treatise on those subjects, and should be in the hands of all practitioners and students. It is a credit to American medical literature. —*Amer. Journ. of the Med. Sciences*, July, 1860.

We question the fact of any recent American author in our profession being more extensively known, or more deservedly esteemed in this country than Dr. Flint. We willingly acknowledge his success, more particularly in the volume on diseases of the heart, in making an extended personal clinical study avail-

able for purposes of illustration, in connection with cases which have been reported by other trustworthy observers. —*Brit. and For. Med.-Chirurg. Review*.

In regard to the merits of the work, we have no hesitation in pronouncing it full, accurate, and judicious. Considering the present state of science, such a work was much needed. It should be in the hands of every practitioner. —*Chicago Med. Journ.*

With more than pleasure do we hail the advent of this work, for it fills a wide gap on the list of textbooks for our schools, and is, for the practitioner, the most valuable practical work of its kind. —*N. O. Med. News*.

BY THE SAME AUTHOR. (Just Issued.)**A PRACTICAL TREATISE ON THE PHYSICAL EXPLORATION OF THE CHEST AND THE DIAGNOSIS OF DISEASES AFFECTING THE RESPIRATORY ORGANS.** Second and revised edition. In one handsome octavo volume of 595 pages, extra cloth, \$4 50.

Premising this observation of the necessity of each student and practitioner making himself acquainted with auscultation and percussion, we may state our honest opinion that Dr. Flint's treatise is one of the most trustworthy guides which he can consult. The style is clear and distinct, and is also concise, being free from that tendency to over-refinement and unnecessary minuteness which characterizes many works on the same subject. —*Dublin Medical Press*, Feb. 6, 1867.

In the invaluable work before us, we have a book of facts of nearly 600 pages, admirably arranged, clear, thorough, and lucid on all points, without prolixity; exhausting every point and topic touched; a monument of patient and long-continued observation, which does credit to its author, and reflects honor on

American medicine. —*Atlanta Med. and Surg. Journal*, Feb. 1867.

The chapter on Phthisis is replete with interest; and his remarks on the diagnosis, especially in the early stages, are remarkable for their acumen and great practical value. Dr. Flint's style is clear and elegant, and the tone of freshness and originality which pervades his whole work lend an additional force to its thoroughly practical character, which cannot fail to obtain for it a place as a standard work on diseases of the respiratory system. —*London Lancet*, Jan. 19, 1867.

This is an admirable book. Excellent in detail and execution, nothing better could be desired by the practitioner. Dr. Flint enriches his subject with much solid and not a little original observation. —*Ranking's Abstract*, Jan. 1867.

PAVY (F. W.), M. D., F. R. S.*Senior Asst. Physician to and Lecturer on Physiology, at Guy's Hospital, &c.***A TREATISE ON THE FUNCTION OF DIGESTION; its Disorders and their Treatment.** From the second London edition. In one handsome volume, small octavo, extra cloth, \$2 00. (*Just Ready.*)

The work before us is one which deserves a wide circulation. We know of no better guide to the study of digestion and its disorders. —*St. Louis Med. and Surg. Journal*, July 10, 1869.

A thoroughly good book, being a careful systematic

treatise, and sufficiently exhaustive for all practical purposes. —*Leavenworth Med. Herald*, July, 1869.

A very valuable work on the subject of which it treats. Small, yet it is full of valuable information. —*Cincinnati Med. Repertory*, June, 1869.

CHAMBERS (T. K.), M. D.,*Consulting Physician to St. Mary's Hospital, London, &c.***THE INDIGESTIONS; or, Diseases of the Digestive Organs Functionally Treated.** Third and revised Edition. In one handsome octavo volume of 333 pages, extra cloth. \$3 00. (*Now Ready.*)**AUTHOR'S PREFACE.**

Since publishing my first edition, I have inserted upwards of ten dozen cases, and have rearranged, indeed in part rewritten, the commentary upon them. A third edition is not yet required in England, so I send the MS. for publication to America. I have faith in the kindly feeling with which it will be received there.

LONDON, December, 1869.

We look upon this chapter as a most valuable guide to physicians, and warning to patients concerning transgressions against the established physiological conduct of life. The advice as to medical treatment proper, is also most serviceable. It should be read by every medical man in the country, and he should read his lessons to his patients from out its pages. We

only regret that we have no room for a more thorough analysis of its contents. —*N. Y. Medical Journal*, March, 1868.

The work should be in the hands of every practicing physician. —*Boston Med. and Surg. Journal*, Nov. 21, 1867.

HABERSHON ON DISEASES OF THE ALIMENTARY CANAL, ŒSOPHAGUS, STOMACH, CÆCUM, AND INTESTINES. With illustrations on wood. One vol. 8vo., 312 pages, extra cloth. \$2 50.

CLYMER ON FEVERS; THEIR DIAGNOSIS PA-

THOLOGY AND TREATMENT. In one octavo volume of 600 pages, leather. \$1 75.

TODD'S CLINICAL LECTURES ON CERTAIN ACUTE DISEASES. In one neat octavo volume, of 320 pages, extra cloth. \$2 50.

ROBERTS (WILLIAM), M. D.*Lecturer on Medicine in the Manchester School of Medicine, &c.***A PRACTICAL TREATISE ON URINARY AND RENAL DISEASES, including Urinary Deposits. Illustrated by numerous cases and engravings. In one very handsome octavo volume of 516 pp., extra cloth. \$4 50. (Just Issued.)**

In carrying out this design, he has not only made good use of his own practical knowledge, but has brought together from various sources a vast amount of information, some of which is not generally possessed by the profession in this country. We must now bring our notice of this book to a close, regretting only that we are obliged to resist the temptation of giving further extracts from it. Dr. Roberts has already on several occasions placed before the profession the results of researches made by him on various points connected with the urine, and had thus led us to expect from him something good—in which expectation we have been by no means disappointed. The book is, beyond question, the most comprehensive

work on urinary and renal diseases, considered in their strictly practical aspect, that we possess in the English language.—*British Medical Journal*, Dec. 9, 1865.

We have read this book with much satisfaction. It will take its place beside the best treatises in our language upon urinary pathology and therapeutics. Not the least of its merits is that the author, unlike some other book-makers, is contented to withhold much that he is well qualified to discuss in order to impart to his volume such a strictly practical character as cannot fail to render it popular among British readers.—*London Med. Times and Gazette*, March 17, 1866.

BASHAM (W. R.), M. D.*Senior Physician to the Westminster Hospital, &c.***RENAL DISEASES: a Clinical Guide to their Diagnosis and Treatment. With illustrations. In one neat royal 12mo. volume of 304 pages. \$2 00. (Just Ready.)**

It is with the view of promoting a practical and clinical knowledge of a class of diseases which are not without their difficulties in diagnosis that the present work has been prepared, with the hope that both student and young practitioner may by it be assisted in their clinical observations.—*Author's Preface.*

MORLAND ON RETENTION IN THE BLOOD OF THE ELEMENTS OF THE URINARY SECRETION. 1 vol. 8vo., extra cloth. 75 cents.

JONES (C. HANDFIELD), M. D.*Physician to St. Mary's Hospital, &c.***CLINICAL OBSERVATIONS ON FUNCTIONAL NERVOUS DISORDERS. Second American Edition. In one handsome octavo volume of 348 pages, extra cloth, \$3 25. (Just Issued.)**

Taken as a whole, the work before us furnishes a short but reliable account of the pathology and treatment of a class of very common but certainly highly obscure disorders. The advanced student will find it a rich mine of valuable facts, while the medical practitioner will derive from it many a suggestive hint to aid him in the diagnosis of "nervous cases," and in determining the true indications for their amelioration or cure.—*Amer. Journ. Med. Sci.*, Jan. 1867.

We must cordially recommend it to the profession of this country as supplying, in a great measure, a deficiency which exists in the medical literature of the English language.—*New York Med. Journ.*, April, 1867.

The volume is a most admirable one—full of hints and practical suggestions.—*Canada Med. Journal*, April, 1867.

SLADE (D. D.), M. D.**DIPHTHERIA; its Nature and Treatment, with an account of the History of its Prevalence in various Countries. Second and revised edition. In one neat royal 12mo. volume, extra cloth. \$1 25. (Just issued.)**

SMITH ON CONSUMPTION; ITS EARLY AND REMEDIABLE STAGES. In one neat octavo volume of 254 pages, extra cloth. \$2 25.

SALTER ON ASTHMA; its Pathology, Causes, Consequences, and Treatment. In one volume octavo, extra cloth. \$2 50.

WALSHE'S PRACTICAL TREATISE ON THE DISEASES OF THE HEART AND GREAT VESSELS. Third American, from the third revised and much enlarged London edition. In one handsome octavo volume of 420 pages, extra cloth. \$3 00.

HUDSON (A.), M. D., M. R. I. A.,*Physician to the Meath Hospital.***LECTURES ON THE STUDY OF FEVER. In one vol. 8vo., extra Cloth, \$2 50. (Just Issued.)**

As an admirable summary of the present state of our knowledge concerning fever, the work will be as welcome to the medical man in active practice as to the student. To the hard-worked practitioner who wishes to refresh his notions concerning fever, the book will prove most valuable. . . . We heartily commend his excellent volume to students and the profession at large.—*London Lancet*, June 22, 1867.

The truly philosophical lectures of Dr. Hudson add

much to our previous knowledge, all of which they, moreover, analyze and condense. This well-conceived task has been admirably executed in the lectures, illustrative cases and quotations being arranged in an appendix to each. We regret that space forbids our quotation from the lectures on treatment, which are, in regard to research and judgment, most masterly, and evidently the result of extended and mature experience.—*British Medical Journal*, Feb. 22, 1868.

LYONS (ROBERT D.), K. C. C.**A TREATISE ON FEVER; or, Selections from a Course of Lectures on Fever. Being part of a Course of Theory and Practice of Medicine. In one neat octavo volume, of 362 pages, extra cloth. \$2 25.**

BUMSTEAD (FREEMAN J.), M.D.,*Professor of Venereal Diseases at the Col. of Phys. and Surg., New York, &c.***THE PATHOLOGY AND TREATMENT OF VENEREAL DISEASES.** Including the results of recent investigations upon the subject. Third edition, revised and enlarged, with illustrations. In one large and handsome octavo volume of over 700 pages, extra cloth. (*Nearly Ready.*)

Well known as one of the best authorities of the present day on the subject.—*British and For. Med.-Chirurg. Review*, April, 1866.

A regular store-house of special information.—*London Lancet*, Feb. 24, 1866.

A remarkably clear and full systematic treatise on the whole subject.—*Lond. Med. Times and Gazette*.

The best, completest, fullest monograph on this subject in our language.—*British American Journal*.

Indispensable in a medical library.—*Pacific Med. and Surg. Journal*.

We have no doubt that it will supersede in America

every other treatise on Venereal.—*San Francisco Med. Press*, Oct. 1864.

A perfect compilation of all that is worth knowing on venereal diseases in general. It fills up a gap which has long been felt in English medical literature.—*Brit. and Foreign Med.-Chirurg. Review*, Jan., '65.

We have not met with any which so highly merits our approval and praise as the second edition of Dr. Bumstead's work.—*Glasgow Med. Journal*, Oct. 1864.

We know of no treatise in any language which is its equal in point of completeness and practical simplicity.—*Boston Medical and Surgical Journal*, Jan. 30, 1864.

CULLERIER (A.), and*Surgeon to the Hôpital du Midi.***BUMSTEAD (FREEMAN J.),***Professor of Venereal Diseases in the College of Physicians and Surgeons, N. Y.***AN ATLAS OF VENEREAL DISEASES.** Translated and Edited by FREEMAN J. BUMSTEAD. In one large imperial 4to. volume of 328 pages, double-columns, with 26 plates, containing about 150 figures, beautifully colored, many of them the size of life; strongly bound in extra cloth, \$17 00; also, in five parts, stout wrappers for mailing, at \$3 per part. (*Just Ready.*)

Anticipating a very large sale for this work, it is offered at the very low price of THREE DOLLARS A Part, thus placing it within the reach of all who are interested in this department of practice. Gentlemen desiring early impressions of the plates would do well to order it without delay.

A specimen of the plates and text sent free by mail, on receipt of 25 cents.

We wish for once that our province was not restricted to methods of treatment, that we might say something of the exquisite colored plates in this volume.—*London Practitioner*, May, 1869.

As a whole, it teaches all that can be taught by means of plates and print.—*London Lancet*, March 13, 1869.

Superior to anything of the kind ever before issued on this continent.—*Canada Med. Journal*, March, '69.

The practitioner who desires to understand this branch of medicine thoroughly should obtain this, the most complete and best work ever published.—*Dominion Med. Journal*, May, 1869.

This is a work of master hands on both sides. M. Cullerier is scarcely second to, we think we may truly say is a peer of the illustrious and venerable Ricord, while in this country we do not hesitate to say that Dr. Bumstead, as an authority, is without a rival. Assuring our readers that these illustrations tell the whole history of venereal disease, from its inception to its end, we do not know a single medical work, which for its kind is more necessary for them to have.—*California Med. Gazette*, March, 1869.

The most splendidly illustrated work in the language, and in our opinion far more useful than the French original.—*Am. Journ. Med. Sciences*, Jan. '69.

HILL (BERKELEY),*Surgeon to the Lock Hospital, London.***ON SYPHILIS AND LOCAL CONTAGIOUS DISORDERS.** In one handsome octavo volume; extra cloth, \$3 25. (*Just Issued.*)

Bringing, as it does, the entire literature of the disease down to the present day, and giving with great ability the results of modern research, it is in every respect a most desirable work, and one which should find a place in the library of every surgeon.—*California Med. Gazette*, June, 1869.

Considering the scope of the book and the careful attention to the manifold aspects and details of its subject, it is wonderfully concise. All these qualities render it an especially valuable book to the beginner, to whom we would most earnestly recommend its study; while it is no less useful to the practitioner.—*St. Louis Med. and Surg. Journal*, May, 1869.

The author, from a vast amount of material, with all of which he was perfectly familiar, has undertaken to construct a new book, and has really succeeded in producing a capital volume upon this subject.—*Nashville Med. and Surg. Journal*, May, 1869.

The most convenient and ready book of reference we have met with.—*N. Y. Med. Record*, May 1, 1869.

Most admirably arranged for both student and practitioner, no other work on the subject equals it; it is more simple, more easily studied.—*Buffalo Med. and Surg. Journal*, March, 1869.

LALLEMAND AND WILSON.**A PRACTICAL TREATISE ON THE CAUSES, SYMPTOMS, AND TREATMENT OF SPERMATORRHOEA.** By M. LALLEMAND. Fifth American edition. To which is added—ON DISEASES OF THE VESICULÆ SEMINALES. By MARRIS WILSON, M.D. In one neat octavo volume, of about 400 pp., extra cloth, \$2 75.

WILSON (ERASMUS), F.R.S.

ON DISEASES OF THE SKIN. With Illustrations on wood. Seventh American, from the sixth and enlarged English edition. In one large octavo volume of over 800 pages, \$5. (*Just Issued.*)

A SERIES OF PLATES ILLUSTRATING "WILSON ON DISEASES OF THE SKIN;" consisting of twenty beautifully executed plates, of which thirteen are exquisitely colored, presenting the Normal Anatomy and Pathology of the Skin, and embracing accurate representations of about one hundred varieties of disease, most of them the size of nature. Price, in extra cloth, \$5 50.

Also, the Text and Plates, bound in one handsome volume. Extra cloth, \$10.

From the Preface to the Sixth English Edition.

The present edition has been carefully revised, in many parts rewritten, and our attention has been specially directed to the practical application and improvements of treatment. And, in conclusion, we venture to remark that if an acute and friendly critic should discover any difference between our present opinions and those announced in former editions, we have only to observe that science and knowledge are progressive, and that we have done our best to move onward with the times.

The industry and care with which the author has revised the present edition are shown by the fact that the volume has been enlarged by more than a hundred pages. In its present improved form it will therefore doubtless retain the position which it has acquired as a standard and classical authority, while at the same time it has additional claims on the attention of the profession as the latest and most complete work on the subject in the English language.

Such a work as the one before us is a most capital and acceptable help. Mr. Wilson has long been held as high authority in this department of medicine, and his book on diseases of the skin has long been regarded as one of the best text-books extant on the subject. The present edition is carefully prepared, and brought up in its revision to the present time. In this edition we have also included the beautiful series of plates illustrative of the text, and in the last edition published separately. There are twenty of these plates, nearly all of them colored to nature, and exhibiting with great fidelity the various groups of diseases treated of in the body of the work.—*Cincinnati Lancet*, June, 1863.

No one treating skin diseases should be without a copy of this standard work.—*Canada Lancet*. August, 1863.

We can safely recommend it to the profession as the best work on the subject now in existence in the English language.—*Medical Times and Gazette*.

Mr. Wilson's volume is an excellent digest of the actual amount of knowledge of cutaneous diseases; it includes almost every fact or opinion of importance connected with the anatomy and pathology of the skin.—*British and Foreign Medical Review*.

These plates are very accurate, and are executed with an elegance and taste which are highly creditable to the artistic skill of the American artist who executed them.—*St. Louis Med. Journal*.

The drawings are very perfect, and the finish and coloring artistic and correct; the volume is an indispensable companion to the book it illustrates and completes.—*Charleston Medical Journal*.

BY THE SAME AUTHOR.

THE STUDENT'S BOOK OF CUTANEOUS MEDICINE and DISEASES OF THE SKIN. In one very handsome royal 12mo. volume. \$3 50. (*Lately Issued.*)

NELIGAN (J. MOORE), M.D., M.R.I.A.

A PRACTICAL TREATISE ON DISEASES OF THE SKIN. Fifth American, from the second and enlarged Dublin edition by T. W. Belcher, M.D. In one neat royal 12mo. volume of 462 pages, extra cloth. \$2 25. (*Just Issued.*)

Fully equal to all the requirements of students and young practitioners. It is a work that has stood its ground, that was worthy the reputation of the author, and the high position of which has been maintained by its learned editor.—*Dublin Med. Press and Circular*, Nov. 17, 1869.

Of the remainder of the work we have nothing beyond unqualified commendation to offer. It is so far the most complete one of its size that has appeared, and for the student there can be none which can compare with it in practical value. All the late discoveries in Dermatology have been duly noticed, and their value justly estimated; in a word, the work is

fully up to the times, and is thoroughly stocked with most valuable information.—*New York Med. Record*, Jan. 15, 1867.

This instructive little volume appears once more. Since the death of its distinguished author, the study of skin diseases has been considerably advanced, and the results of these investigations have been added by the present editor to the original work of Dr. Neligan. This, however, has not so far increased its bulk as to destroy its reputation as the most convenient manual of diseases of the skin that can be procured by the student.—*Chicago Med. Journal*, Dec. 1866.

BY THE SAME AUTHOR.

ATLAS OF CUTANEOUS DISEASES. In one beautiful quarto volume, with exquisitely colored plates, &c., presenting about one hundred varieties of disease. Extra cloth, \$5 50.

The diagnosis of eruptive disease, however, under all circumstances, is very difficult. Nevertheless, Dr. Neligan has certainly, "as far as possible," given a faithful and accurate representation of this class of diseases, and there can be no doubt that these plates will be of great use to the student and practitioner in drawing a diagnosis as to the class, order, and species to which the particular case may belong. While looking over the "Atlas" we have been induced to examine also the "Practical Treatise," and we are

inclined to consider it a very superior work, combining accurate verbal description with sound views of the pathology and treatment of eruptive diseases.—*Glasgow Med. Journal*.

A compend which will very much aid the practitioner in this difficult branch of diagnosis. Taken with the beautiful plates of the Atlas, which are remarkable for their accuracy and beauty of coloring, it constitutes a very valuable addition to the library of a practical man.—*Buffalo Med. Journal*.

HILLIER (THOMAS), M.D.,

Physician to the Skin Department of University College Hospital, &c.

HAND-BOOK OF SKIN DISEASES, for Students and Practitioners. Second American Edition. In one royal 12mo. volume of 358 pp. With Illustrations. Extra cloth, \$2 25. (*Now Ready.*)

SMITH (J. LEWIS), M. D.,*Professor of Morbid Anatomy in the Bellevue Hospital Med. College, N. Y.***A COMPLETE PRACTICAL TREATISE ON THE DISEASES OF CHILDREN.** In one handsome octavo volume of 620 pages, extra cloth, \$4 75; leather, \$5 75. (*Just Issued.*)

We have no work upon the Diseases of Infancy and Childhood which can compare with it.—*Buffalo Med. and Surg. Journal*, March, 1869.

The description of the pathology, symptoms, and treatment of the different diseases is excellent.—*Am. Med. Journal*, April, 1869.

So full, satisfactory, and complete is the information to be derived from this work, that at no time have we examined the pages of any book with more pleasure. The diseases incident to childhood are treated with a clearness, precision, and understanding that is not often met with, and which must call forth the approval of all who consult its pages.—*Cincinnati Med. Repository*, May, 1869.

The author of this volume is well known as a valued contributor to the literature of his specialty. The faithful manner in which he has worked in the public institutions with which he has been connected, the conscientious regard for truth which has for years characterized all his researches, the great amount of experience which he has been enabled to acquire in the treatment of infantile diseases, and the care which he has accustomed himself to take in the study of the significant facts relating to the pathological anatomy

of the diseases of childhood, eminently fit him for the task which he has taken upon himself. The remarkable faculty of bringing out salient points and stating concisely other less important facts, enables him to crowd within a small compass a vast amount of practical information. The attention given to the treatment of the various maladies, as well as the presentation of all the recently accepted pathological views, make it one of the most valuable treatises, within its present compass, that can be placed in the hands of any seeker after truth. The volume as a whole will still further establish for the writer a permanent and enviable reputation as a careful observer, an impartial interpreter, a safe and trustworthy adviser, and a modest and untrifling student.—*N. Y. Med. Record*, March 15, 1869.

We have perused Dr. Smith's book with not a little satisfaction; it is indeed an excellent work; well and correctly written; thoroughly up to the modern ideas; concise, yet complete in its material. We cannot help welcoming a work which will be worthy of reliance as a text-book for medical students and younger physicians in their investigation of disease in children. *Boston Med. and Surg. Journal*, March 4, 1869.

CONDIE (D. FRANCIS), M. D.**A PRACTICAL TREATISE ON THE DISEASES OF CHILDREN.**

Sixth edition, revised and augmented. In one large octavo volume of nearly 800 closely-printed pages, extra cloth, \$5 25; leather, \$6 25. (*Lately Issued.*)

Dr. Condie has been one of those who have performed such a service satisfactorily, and, as a result, his popular, comprehensive, and practical work has received that high compliment of approval on the part of his brethren, which several editions inconceivably set forth. The present edition, which is the sixth, is fully up to the times in the discussion of all those points in the pathology and treatment of infantile diseases which have been brought forward by the German and French teachers. As a whole, however, the work is the best American one that we have, and in its special adaptation to American practitioners it certainly has no equal.—*New York Med. Record*, March 2, 1868.

No other treatise on this subject is better adapted to the American physician. Dr. Condie has long stood

before his countrymen as one peculiarly pre-eminent in this department of medicine. His work has been so long a standard for practitioners and medical students that we do no more now than refer to the fact that it has reached its sixth edition. We are glad once more to refresh the impressions of our earlier days by wandering through its pages, and at the same time to be able to recommend it to the youngest members of the profession, as well as to those who have the older editions on their shelves.—*St. Louis Med. Reporter*, Feb. 15, 1868.

We pronounced the first edition to be the best work on the diseases of children in the English language, and, notwithstanding all that has been published, we still regard it in that light.—*Medical Examiner*.

WEST (CHARLES), M. D.,*Physician to the Hospital for Sick Children, &c.***LECTURES ON THE DISEASES OF INFANCY AND CHILDHOOD.** Fourth American from the fifth revised and enlarged English edition. In one large and handsome octavo volume of 656 closely-printed pages. Extra cloth, \$4 50; leather, \$5 50. (*Lately issued.*)

Of all the English writers on the diseases of children, there is no one so entirely satisfactory to us as Dr. West. For years we have held his opinion as judicial, and have regarded him as one of the highest living authorities in the difficult department of medical science in which he is most widely known.—*Boston Med. and Surg. Journal*, April 26, 1866.

Dr. West's volume is, in our opinion, incomparably the best authority upon the maladies of children that the practitioner can consult.—*Cincinnati Jour. of Medicine*, March, 1866.

We have long regarded it as the most scientific and practical book on diseases of children which has yet appeared in this country.—*Buffalo Medical Journal*.

SMITH (EUSTACE), M. D.,*Physician to the Northwest London Free Dispensary for Sick Children.***A PRACTICAL TREATISE ON THE WASTING DISEASES OF INFANCY AND CHILDHOOD.** 1 vol. 8vo. Second and revised edition. (*Preparing.*)

In this brief treatise, the author has made one of the most valuable contributions to medical literature that has been given to our profession for many years. To supply the want of information on this subject is the task which Dr. Smith has set himself, and admirably has he performed it. Keeping steadily in view

a purpose of clinical usefulness, he has succeeded in producing a treatise on the causes of chronic wasting so complete that but little could be added, and yet so concise that it would be almost impossible to give a synopsis of his views in fewer words than the book itself contains.—*N. Y. Med. Gazette*, April 2, 1870.

DEWEES (WILLIAM P.), M. D.**A TREATISE ON THE PHYSICAL AND MEDICAL TREATMENT OF CHILDREN.** Eleventh edition, with the author's last improvements and corrections. In one octavo volume of 548 pages. \$2 80.

THOMAS (T. GAILLARD), M. D.,

Professor of Obstetrics, &c. in the College of Physicians and Surgeons, N. Y., &c.

A PRACTICAL TREATISE ON THE DISEASES OF WOMEN. Second edition, revised and improved. In one large and handsome octavo volume of 650 pages, with 225 illustrations, extra cloth, \$5; leather, \$6. (*Now Ready.*)

From the Preface to the Second Edition.

In a science so rapidly progressive as that of medicine, the profession has a right to expect that, when its approbation of a work is manifested by a call for a new edition, the author should respond by giving to his book whatever of additional value may be derivable from more extended experience, maturer thought, and the opportunity for correction. Fully sensible of this, the author of the present volume has sought by a careful revision of the whole, and by the addition of a chapter on Chlorosis, to render his work more worthy of the favor with which it has been received.—**NEW YORK, March, 1869.**

If the excellence of a work is to be judged by its rapid sale, this one must take precedence of all others upon the same, or kindred subjects, as evidenced in the short time from its first appearance, in which a new edition is called for, resulting, as we are informed, from the exhaustion of the previous large edition. We deem it scarcely necessary to recommend this work to physicians as it is now widely known, and most of them already possess it, or will certainly do so. To students we unhesitatingly recommend it as the best text-book on diseases of females extant.—*St. Louis Med. Reporter*, June, 1869.

Of all the army of books that have appeared of late years, on the diseases of the uterus and its appendages, we know of none that is so clear, comprehensive, and practical as this of Dr. Thomas', or one that we should more emphatically recommend to the young practitioner, as his guide.—*California Med. Gazette*, June, 1869.

If not the best work extant on the subject of which it treats, it is certainly second to none other. So short a time has elapsed since the medical press teemed with commendatory notices of the first edition, that it would be superfluous to give an extended review of what is now firmly established as the American text-book of Gynecology.—*N. Y. Med. Gazette*, July 17, 1869.

This is a new and revised edition of a work which we recently noticed at some length, and earnestly commended to the favorable attention of our readers. The fact that, in the short space of one year, this second edition makes its appearance, shows that the general judgment of the profession has largely confirmed the opinion we gave at that time.—*Cincinnati Lancet*, Aug. 1869.

It is so short a time since we gave a full review of the first edition of this book, that we deem it only necessary now to call attention to the second appearance of the work. Its success has been remarkable, and we can only congratulate the author on the brilliant reception his book has received.—*N. Y. Med. Journal*, April, 1869.

We regard this treatise as the one best adapted to serve as a text-book on gynecology.—*St. Louis Med. and Surg. Journal*, May 10, 1869.

The whole work as it now stands is an absolute indispensable to any physician aspiring to treat the diseases of females with success, and according to the most fully accepted views of their etiology and pathology.—*Leavenworth Medical Herald*, May, 1869.

We have seldom read a medical book in which we found so much to praise, and so little—we can hardly say to object to—to mention with qualified commendation. We had proposed a somewhat extended review with copious extracts, but we hardly know where we should have space for it. We therefore content ourselves with expressing the belief that every practitioner of medicine would do well to possess himself of the work.—*Boston Med. and Surg. Journal*, April 29, 1869.

The number of works published on diseases of women is large, not a few of which are very valuable. But of those which are the most valuable we do not regard the work of Dr. Thomas as second to any. Without being prolix, it treats of the disorders to which it is devoted fully, perspicuously, and satisfactorily. It will be found a treasury of knowledge to every physician who turns to its pages. We would like to make a number of quotations from the work of a practical bearing, but our space will not permit. The work should find a place in the libraries of all physicians.—*Cincinnati Med. Repository*, May, 1869.

No one will be surprised to learn that the valuable, readable, and thoroughly practical book of Professor Thomas has so soon advanced to a second edition. Although very little time has necessarily been allowed our author for revision and improvement of the work, he has performed it exceedingly well. Aside from the numerous corrections which he has found necessary to make, he has added an admirable chapter on chlorosis, which of itself is worth the cost of the volume.—*N. Y. Med. Record*, May 15, 1869.

CHURCHILL (FLEETWOOD), M. D., M. R. I. A.

ON THE DISEASES OF WOMEN; including those of Pregnancy and Childbed. A new American edition, revised by the Author. With Notes and Additions, by D. FRANCIS CONDIE, M. D., author of "A Practical Treatise on the Diseases of Children." With numerous illustrations. In one large and handsome octavo volume of 768 pages, extra cloth, \$4 00; leather, \$5 00.

BY THE SAME AUTHOR.

ESSAYS ON THE PUERPERAL FEVER, AND OTHER DISEASES PECULIAR TO WOMEN. Selected from the writings of British Authors previous to the close of the Eighteenth Century. In one neat octavo volume of about 450 pages, extra cloth. \$2 50.

ASHWELL (SAMUEL), M. D.,

Late Obstetric Physician and Lecturer at Guy's Hospital.

A PRACTICAL TREATISE ON THE DISEASES PECULIAR TO WOMEN. Illustrated by Cases derived from Hospital and Private Practice. Third American, from the Third and revised London edition. In one octavo volume, extra cloth, of 528 pages. \$3 50.

BROWN ON SOME DISEASES OF WOMEN ADMITTING OF SURGICAL TREATMENT. With handsome illustrations. One volume 8vo., extra cloth, pp. 274. \$1 60.

DEWEES'S TREATISE ON THE DISEASES OF FEMALES. With illustrations. Eleventh Edition,

with the Author's last improvements and corrections. In one octavo volume of 536 pages, with plates, extra cloth, \$3 00.

RIGBY ON THE CONSTITUTIONAL TREATMENT OF FEMALE DISEASES. In one neat royal 12mo. volume, extra cloth, of about 250 pages. \$1 00.

HODGE (HUGH L.), M.D.,
Emeritus Professor of Obstetrics, &c., in the University of Pennsylvania.

ON DISEASES PECULIAR TO WOMEN; including Displacements of the Uterus. With original illustrations. Second edition, revised and enlarged. In one beautifully printed octavo volume of 531 pages, extra cloth. \$4 50. (*Just Issued.*)

In the preparation of this edition the author has spared no pains to improve it with the results of his observation and study during the interval which has elapsed since the first appearance of the work. Considerable additions have thus been made to it, which have been partially accommodated by an enlargement in the size of the page, to avoid increasing unduly the bulk of the volume.

From PROF. W. H. BYFORD, of the Rush Medical College, Chicago.

The book bears the impress of a master hand, and must, as its predecessor, prove acceptable to the profession. In diseases of women Dr. Hodge has established a school of treatment that has become world-wide in fame.

Professor Hodge's work is truly an original one from beginning to end, consequently no one can peruse its pages without learning something new. The book, which is by no means a large one, is divided into two grand sections, so to speak: first, that treating of the nervous sympathies of the uterus, and, secondly, that which speaks of the mechanical treatment of displacements of that organ. He is disposed, as a non-believer in the frequency of inflammations of the

uterus, to take strong ground against many of the highest authorities in this branch of medicine, and the arguments which he offers in support of his position are, to say the least, well put. Numerous woodcuts adorn this portion of the work, and add incalculably to the proper appreciation of the variously shaped instruments referred to by our author. As a contribution to the study of women's diseases, it is of great value, and is abundantly able to stand on its own merits.—*N. Y. Medical Record*, Sept. 15, 1868.

In this point of view, the treatise of Professor Hodge will be indispensable to every student in its department. The large, fair type and general perfection of workmanship will render it doubly welcome.—*Pacific Med. and Surg. Journal*, Oct. 1868.

WEST (CHARLES), M.D.

LECTURES ON THE DISEASES OF WOMEN. Third American, from the Third London edition. In one neat octavo volume of about 550 pages, extra cloth. \$3 75; leather, \$4 75. (*Just Issued.*)

The reputation which this volume has acquired as a standard book of reference in its department, renders it only necessary to say that the present edition has received a careful revision at the hands of the author, resulting in a considerable increase of size. A few notices of previous editions are subjoined.

The manner of the author is excellent, his descriptions graphic and perspicuous, and his treatment up to the level of the time—clear, precise, definite, and marked by strong common sense.—*Chicago Med. Journal*, Dec. 1861.

We cannot too highly recommend this, the second edition of Dr. West's excellent lectures on the diseases of females. We know of no other book on this subject from which we have derived as much pleasure and instruction. Every page gives evidence of the honest, earnest, and diligent searcher after truth. He is not the mere compiler of other men's ideas, but his lectures are the result of ten years' patient investigation in one of the widest fields for women's diseases—St. Bartholomew's Hospital. As a teacher, Dr. West is simple and earnest in his language, clear and comprehensive in his perceptions, and logical in his deductions.—*Cincinnati Lancet*, Jan. 1862.

We return the author our grateful thanks for the vast amount of instruction he has afforded us. His valuable treatise needs no eulogy on our part. His graphic diction and truthful pictures of disease all speak for themselves.—*Medico-Chirurg. Review*.

Most justly esteemed a standard work. . . . It bears evidence of having been carefully revised, and is well worthy of the fame it has already obtained.—*Dub. Med. Quar. Jour.*

As a writer, Dr. West stands, in our opinion, second only to Watson, the "Macanlay of Medicine;" he possesses that happy faculty of clothing instruction in easy garments; combining pleasure with profit, he leads his pupils, in spite of the ancient proverb, along a royal road to learning. His work is one which will not satisfy the extreme on either side, but it is one that will please the great majority who are seeking truth, and one that will convince the student that he has committed himself to a candid, safe, and valuable guide.—*N. A. Med.-Chirurg. Review*.

We must now conclude this hastily written sketch with the confident assurance to our readers that the work will well repay perusal. The conscientious, painstaking, practical physician is apparent on every page.—*N. Y. Journal of Medicine*.

We have to say of it, briefly and decidedly, that it is the best work on the subject in any language, and that it stamps Dr. West as the *facile princeps* of British obstetric authors.—*Edinburgh Med. Journal*.

We gladly recommend his lectures as in the highest degree instructive to all who are interested in obstetric practice.—*London. Lancet*.

We know of no treatise of the kind so complete, and yet so compact.—*Chicago Med. Journal*.

BY THE SAME AUTHOR.

AN ENQUIRY INTO THE PATHOLOGICAL IMPORTANCE OF ULCERATION OF THE OS UTERI. In one neat octavo volume, extra cloth. \$1 25.

MEIGS (CHARLES D.), M.D.,
Late Professor of Obstetrics, &c. in Jefferson Medical College, Philadelphia.

WOMAN: HER DISEASES AND THEIR REMEDIES. A Series of Lectures to his Class. Fourth and Improved edition. In one large and beautifully printed octavo volume of over 700 pages, extra cloth, \$5 00; leather, \$6 00.

BY THE SAME AUTHOR.

ON THE NATURE, SIGNS, AND TREATMENT OF CHILD BED FEVER. In a Series of Letters addressed to the Students of his Class. In one handsome octavo volume of 365 pages, extra cloth. \$2 00.

SIMPSON (SIR JAMES Y.), M.D.

CLINICAL LECTURES ON THE DISEASES OF WOMEN. With numerous illustrations. In one octavo volume of over 500 pages. Second edition, *preparing*.

HODGE (HUGH L.), M.D.,*Emeritus Professor of Midwifery, &c. in the University of Pennsylvania, &c.***THE PRINCIPLES AND PRACTICE OF OBSTETRICS.** Illustrated with large lithographic plates containing one hundred and fifty-nine figures from original photographs, and with numerous wood-cuts. In one large and beautifully printed quarto volume of 550 double-columned pages, strongly bound in extra cloth, \$14. (*Lately published.*)

The work of Dr. Hodge is something more than a simple presentation of his particular views in the department of Obstetrics; it is something more than an ordinary treatise on midwifery: it is, in fact, a cyclopædia of midwifery. He has aimed to embody in a single volume the whole science and art of Obstetrics. An elaborate text is combined with accurate and varied pictorial illustrations, so that no fact or principle is left unstated or unexplained.—*Am. Med. Times*, Sept. 3, 1864.

We should like to analyze the remainder of this excellent work, but already has this review extended beyond our limited space. We cannot conclude this notice without referring to the excellent finish of the work. In typography it is not to be excelled; the paper is superior to what is usually afforded by our American cousins, quite equal to the best of English books. The engravings and lithographs are most beautifully executed. The work recommends itself for its originality, and is in every way a most valuable addition to those on the subject of obstetrics.—*Canada Med. Journal*, Oct. 1864.

It is very large, profusely and elegantly illustrated, and is fitted to take its place near the works of great obstetricians. Of the American works on the subject it is decidedly the best.—*Edinb. Med. Jour.*, Dec. '64.

*** Specimens of the plates and letter-press will be forwarded to any address, free by mail, on receipt of six cents in postage stamps.

We have examined Professor Hodge's work with great satisfaction; every topic is elaborated most fully. The views of the author are comprehensive, and concisely stated. The rules of practice are judicious, and will enable the practitioner to meet every emergency of obstetric complication with confidence.—*Chicago Med. Journal*, Aug. 1864.

More time than we have had at our disposal since we received the great work of Dr. Hodge is necessary to do it justice. It is undoubtedly by far the most original, complete, and carefully composed treatise on the principles and practice of Obstetrics which has ever been issued from the American press.—*Pacific Med. and Surg. Journal*, July, 1864.

We have read Dr. Hodge's book with great pleasure, and have much satisfaction in expressing our commendation of it as a whole. It is certainly highly instructive, and in the main, we believe, correct. The great attention which the author has devoted to the mechanism of parturition, taken along with the conclusions at which he has arrived, point, we think, conclusively to the fact that, in Britain at least, the doctrines of Nægele have been too blindly received.—*Glasgow Med. Journal*, Oct. 1864.

TANNER (THOMAS H.), M.D.**ON THE SIGNS AND DISEASES OF PREGNANCY.** First American from the Second and Enlarged English Edition. With four colored plates and illustrations on wood. In one handsome octavo volume of about 500 pages, extra cloth, \$4 25. (*Just Issued.*)

The very thorough revision the work has undergone has added greatly to its practical value, and increased materially its efficiency as a guide to the student and to the young practitioner.—*Am. Journ. Med. Sci.*, April, 1868.

With the immense variety of subjects treated of and the ground which they are made to cover, the impossibility of giving an extended review of this truly remarkable work must be apparent. We have not a single fault to find with it, and most heartily commend it to the careful study of every physician who would not only always be sure of his diagnosis of pregnancy, but always ready to treat all the numerous ailments that are, unfortunately for the civilized women of to-day, so commonly associated with the function.—*N. Y. Med. Record*, March 16, 1868.

We have much pleasure in calling the attention of our readers to the volume produced by Dr. Tanner, the second edition of a work that was, in its original

state even, acceptable to the profession. We recommend obstetrical students, young and old, to have this volume in their collections. It contains not only a fair statement of the signs, symptoms, and diseases of pregnancy, but comprises in addition much interesting relative matter that is not to be found in any other work that we can name.—*Edinburgh Med. Journal*, Jan. 1868.

In its treatment of the signs and diseases of pregnancy it is the most complete book we know of, abounding on every page with matter valuable to the general practitioner.—*Cincinnati Med. Repository*, March, 1868.

This is a most excellent work, and should be on the table or in the library of every practitioner.—*Humboldt Med. Archives*, Feb. 1868.

A valuable compendium, enriched by his own labors, of all that is known on the signs and diseases of pregnancy.—*St. Louis Med. Reporter*, Feb. 15, 1868.

MONTGOMERY (W. F.), M.D.,*Professor of Midwifery in the King's and Queen's College of Physicians in Ireland.***AN EXPOSITION OF THE SIGNS AND SYMPTOMS OF PREGNANCY.** With some other Papers on Subjects connected with Midwifery. From the second and enlarged English edition. With two exquisite colored plates, and numerous wood-cuts. In one very handsome octavo volume of nearly 600 pages, extra cloth. \$3 75.**MILLER (HENRY), M.D.,***Professor of Obstetrics and Diseases of Women and Children in the University of Louisville.***PRINCIPLES AND PRACTICE OF OBSTETRICS, &c.;** including the Treatment of Chronic Inflammation of the Cervix and Body of the Uterus considered as a frequent cause of Abortion. With about one hundred illustrations on wood. In one very handsome octavo volume of over 600 pages, extra cloth. \$3 75.

RIGBY'S SYSTEM OF MIDWIFERY. With Notes and Additional Illustrations. Second American edition. One volume octavo, extra cloth, 422 pages. \$2 50.

DEWEES'S COMPREHENSIVE SYSTEM OF MIDWIFERY. Twelfth edition, with the author's last improvements and corrections. In one octavo volume, extra cloth, of 600 pages. \$3 50.

MEIGS (CHARLES D.), M.D.,*Lately Professor of Obstetrics, &c., in the Jefferson Medical College, Philadelphia.***OBSTETRICS: THE SCIENCE AND THE ART.** Fifth edition, revised. With one hundred and thirty illustrations. In one beautifully printed octavo volume of 760 large pages. Extra cloth, \$5 50; leather, \$6 50. (Just Issued.)

The original edition is already so extensively and favorably known to the profession that no recommendation is necessary; it is sufficient to say, the present edition is very much extended, improved, and perfected. Whilst the great practical talents and unlimited experience of the author render it a most valuable acquisition to the practitioner, it is so condensed as to constitute a most eligible and excellent text-book for the student.—*Southern Med. and Surg. Journal*, July, 1867.

It is to the student that our author has more particularly addressed himself; but to the practitioner we believe it would be equally serviceable as a book of reference. No work that we have met with so thoroughly details everything that falls to the lot of the accoucheur to perform. Every detail, no matter

how minute or how trivial, has found a place.—*Canada Medical Journal*, July, 1867.

This very excellent work on the science and art of obstetrics should be in the hands of every student and practitioner. The rapidity with which the very large editions have been exhausted is the best test of its true merit. Besides, it is the production of an American who has probably had more experience in this branch than any other living practitioner of the country.—*St. Louis Med. and Surg. Journal*, Sept. 1867.

He has also carefully endeavored to be minute and clear in his details, with as little reiteration as possible, and beautifully combines the relations of science to art, as far as the different classifications will admit.—*Detroit Review of Med. and Pharm.*, Aug. 1867.

RAMSBOTHAM (FRANCIS H.), M.D.**THE PRINCIPLES AND PRACTICE OF OBSTETRIC MEDICINE AND SURGERY**, in reference to the Process of Parturition. A new and enlarged edition, thoroughly revised by the author. With additions by W. V. KEATING, M.D., Professor of Obstetrics, &c., in the Jefferson Medical College, Philadelphia. In one large and handsome imperial octavo volume of 650 pages, strongly bound in leather, with raised bands; with sixty-four beautiful plates, and numerous wood-cuts in the text, containing in all nearly 200 large and beautiful figures. \$7 00.

We will only add that the student will learn from it all he need to know, and the practitioner will find it, as a book of reference, surpassed by none other.—*Stethoscope*.

The character and merits of Dr. Ramsbotham's work are so well known and thoroughly established, that comment is unnecessary and praise superfluous. The illustrations, which are numerous and accurate, are executed in the highest style of art. We cannot too highly recommend the work to our readers.—*St. Louis Med. and Surg. Journal*.

To the physician's library it is indispensable, while to the student, as a text-book, from which to extract the material for laying the foundation of an education on obstetrical science, it has no superior.—*Ohio Med. and Surg. Journal*.

When we call to mind the toil we underwent in acquiring a knowledge of this subject, we cannot but envy the student of the present day the aid which this work will afford him.—*Am. Jour. of the Med. Sciences*.

CHURCHILL (FLEETWOOD), M.D., M.R.I.A.**ON THE THEORY AND PRACTICE OF MIDWIFERY.** A new American from the fourth revised and enlarged London edition. With notes and additions by D. FRANCIS CONDIZ, M.D., author of a "Practical Treatise on the Diseases of Children," &c. With one hundred and ninety-four illustrations. In one very handsome octavo volume of nearly 700 large pages. Extra cloth, \$4 00; leather, \$5 00.

In adapting this standard favorite to the wants of the profession in the United States, the editor has endeavored to insert everything that his experience has shown him would be desirable for the American student, including a large number of illustrations. With the sanction of the author, he has added, in the form of an appendix, some chapters from a little "Manual for Midwives and Nurses," recently issued by Dr. Churchill, believing that the details there presented can hardly fail to prove of advantage to the junior practitioner. The result of all these additions is that the work now contains fully one-half more matter than the last American edition, with nearly one-half more illustrations; so that, notwithstanding the use of a smaller type, the volume contains almost two hundred pages more than before.

These additions render the work still more complete and acceptable than ever; and with the excellent style in which the publishers have presented this edition of Churchill, we can commend it to the profession with great cordiality and pleasure.—*Cincinnati Lancet*.

Few works on this branch of medical science are equal to it, certainly none excel it, whether in regard to theory or practice, and in one respect it is superior to all others, viz., in its statistical information, and therefore, on these grounds a most valuable work for the physician, student, or lecturer, all of whom will find in it the information which they are seeking.—*Brit. Am. Journal*.

The present treatise is very much enlarged and amplified beyond the previous editions but nothing

has been added which could be well dispensed with. An examination of the table of contents shows how thoroughly the author has gone over the ground, and the care he has taken in the text to present the subjects in all their bearings, will render this new edition even more necessary to the obstetric student than were either of the former editions at the date of their appearance. No treatise on obstetrics with which we are acquainted can compare favorably with this, in respect to the amount of material which has been gathered from every source.—*Boston Med. and Surg. Journal*.

There is no better text-book for students, or work of reference and study for the practising physician than this. It should adorn and enrich every medical library.—*Chicago Med. Journal*.

SWAYNE (JOSEPH GRIFFITHS), M.D.,*Physician-Accoucheur to the British General Hospital, &c.***OBSTETRIC APHORISMS FOR THE USE OF STUDENTS COMMENCING MIDWIFERY PRACTICE.** From the Fourth and Revised London Edition, with Additions by E. R. HUTCHINS, M.D. With Illustrations. In one neat 12mo. volume. Extra cloth, \$1 25. (Just Ready.)

GROSS (SAMUEL D.), M.D.,*Professor of Surgery in the Jefferson Medical College of Philadelphia.*

A SYSTEM OF SURGERY: Pathological, Diagnostic, Therapeutic, and Operative. Illustrated by upwards of Thirteen Hundred Engravings. Fourth edition, carefully revised, and improved. In two large and beautifully printed royal octavo volumes of 2200 pages, strongly bound in leather, with raised bands. \$15 00.

The continued favor, shown by the exhaustion of successive large editions of this great work, proves that it has successfully supplied a want felt by American practitioners and students. Though but little over six years have elapsed since its first publication, it has already reached its fourth edition, while the care of the author in its revision and correction has kept it in a constantly improved shape. By the use of a close, though very legible type, an unusually large amount of matter is condensed in its pages, the two volumes containing as much as four or five ordinary octavos. This, combined with the most careful mechanical execution, and its very durable binding, renders it one of the cheapest works accessible to the profession. Every subject properly belonging to the domain of surgery is treated in detail, so that the student who possesses this work may be said to have in it a surgical library.

It must long remain the most comprehensive work on this important part of medicine.—*Boston Medical and Surgical Journal*, March 23, 1865.

We have compared it with most of our standard works, such as those of Erichsen, Miller, Fergusson, Syme, and others, and we must, in justice to our author, award it the pre-eminence. As a work, complete in almost every detail, no matter how minute or trifling, and embracing every subject known in the principles and practice of surgery, we believe it stands without a rival. Dr. Gross, in his preface, remarks "my aim has been to embrace the whole domain of surgery, and to allot to every subject its legitimate claim to notice;" and, we assure our readers, he has kept his word. It is a work which we can most confidently recommend to our brethren, for its utility is becoming the more evident the longer it is upon the shelves of our library.—*Canada Med. Journal*, September, 1865.

The first two editions of Professor Gross' System of Surgery are so well known to the profession, and so highly prized, that it would be idle for us to speak in praise of this work.—*Chicago Medical Journal*, September, 1865.

We gladly indorse the favorable recommendation of the work, both as regards matter and style, which we made when noticing its first appearance.—*British and Foreign Medico-Chirurgical Review*, Oct. 1865.

The most complete work that has yet issued from the press on the science and practice of surgery.—*London Lancet*.

This system of surgery is, we predict, destined to take a commanding position in our surgical literature, and be the crowning glory of the author's well earned fame. As an authority on general surgical subjects, this work is long to occupy a pre-eminent place, not only at home, but abroad. We have no hesitation in pronouncing it without a rival in our language, and equal to the best systems of surgery in any language.—*N. Y. Med. Journal*.

Not only by far the best text-book on the subject, as a whole, within the reach of American students, but one which will be much more than ever likely to be resorted to and regarded as a high authority abroad.—*Am. Journal Med. Sciences*, Jan. 1865.

The work contains everything, minor and major, operative and diagnostic, including mensuration and examination, venereal diseases, and uterine manipulations and operations. It is a complete Thesaurus of modern surgery, where the student and practi-

tioner shall not seek in vain for what they desire.—*San Francisco Med. Press*, Jan. 1865.

Open it where we may, we find sound practical information conveyed in plain language. This book is no mere provincial or even national system of surgery, but a work which, while very largely indebted to the past, has a strong claim on the gratitude of the future of surgical science.—*Edinburgh Med. Journal*, Jan. 1865.

A glance at the work is sufficient to show that the author and publisher have spared no labor in making it the most complete "System of Surgery" ever published in any country.—*St. Louis Med. and Surg. Journal*, April, 1865.

The third opportunity is now offered during our editorial life to review, or rather to indorse and recommend this great American work on Surgery. Upon this last edition a great amount of labor has been expended, though to all others except the author the work was regarded in its previous editions as so full and complete as to be hardly capable of improvement. Every chapter has been revised; the text augmented by nearly two hundred pages, and a considerable number of wood-cuts have been introduced. Many portions have been entirely re-written, and the additions made to the text are principally of a practical character. This comprehensive treatise upon surgery has undergone revisions and enlargements, keeping pace with the progress of the art and science of surgery, so that whoever is in possession of this work may consult its pages upon any topic embraced within the scope of its department, and rest satisfied that its teaching is fully up to the present standard of surgical knowledge. It is also so comprehensive that it may truthfully be said to embrace all that is actually known, that is really of any value in the diagnosis and treatment of surgical diseases and accidents. Wherever illustration will add clearness to the subject, or make better or more lasting impression, it is not wanting; in this respect the work is eminently superior.—*Buffalo Med. Journal*, Dec. 1864.

A system of surgery which we think unrivalled in our language, and which will indelibly associate his name with surgical science. And what, in our opinion, enhances the value of the work is that, while the practising surgeon will find all that he requires in it, it is at the same time one of the most valuable treatises which can be put into the hands of the student seeking to know the principles and practice of this branch of the profession which he designs subsequently to follow.—*The Brit. Am. Journ., Montreal*.

BY THE SAME AUTHOR.

A PRACTICAL TREATISE ON THE DISEASES, INJURIES, AND MALFORMATIONS OF THE URINARY BLADDER, THE PROSTATE GLAND, AND THE URETHRA. Second edition, revised and much enlarged, with one hundred and eighty-four illustrations. In one large and very handsome octavo volume of over nine hundred pages, extra cloth. \$4 00.

BY THE SAME AUTHOR.

A PRACTICAL TREATISE ON FOREIGN BODIES IN THE AIR-PASSAGES. In one handsome octavo volume, extra cloth, with illustrations. pp. 468. \$2 75.

MALGAIGNE'S OPERATIVE SURGERY. With numerous illustrations on wood. In one handsome octavo volume, extra cloth, of nearly 600 pp. \$2 50.

SKYE'S OPERATIVE SURGERY. In one very handsome octavo volume, extra cloth, of over 650 pages with about 100 wood-cuts. \$3 25.

ERICHSEN (JOHN),
Senior Surgeon to University College Hospital.

THE SCIENCE AND ART OF SURGERY; being a Treatise on Surgical Injuries, Diseases, and Operations. From the Fifth enlarged and carefully revised London Edition. With Additions by JOHN ASHHURST, Jr., M. D., Surgeon to the Episcopal Hospital, &c. Illustrated by over six hundred Engravings on wood. In one very large and beautifully printed imperial octavo volume, containing over twelve hundred closely printed pages: cloth, \$7 50; leather, raised bands, \$8 50. (*Just Issued.*)

This volume having enjoyed repeated revisions at the hands of the author has been greatly enlarged, and the present edition will thus be found to contain at least one-half more matter than the last American impression. On the latest London edition, just issued, especial care has been bestowed. Besides the most minute attention on the part of the author to bring every portion of it thoroughly on a level with the existing condition of science, he called to his aid gentlemen of distinction in special departments. Thus a chapter on the Surgery of the Eye and its Appendages has been contributed by Mr. Streatfeild; the section devoted to Syphilis has been rearranged under the supervision of Mr. Berkeley Hill; the subjects of General Surgical Diseases, including Pyæmia, Scrofula, and Tumors, have been revised by Mr. Alexander Bruce; and other professional men of eminence have assisted in other branches. The work may thus be regarded as embodying a complete and comprehensive view of the most advanced condition of British surgery; while such omissions of practical details in American surgery as were found have been supplied by the editor, Dr. Ashhurst.

Thus complete in every respect, thoroughly illustrated, and containing in one beautifully printed volume the matter of two or three ordinary octavos, it is presented at a price which renders it one of the cheapest works now accessible to the profession. A continuance of the very remarkable favor which it has thus far enjoyed is therefore confidently expected.

The high position which Mr. Erichsen's Science and Art of Surgery has for some time attained, not only in this country, but on the Continent and in America, almost limits the task of the reviewer, on the appearance of a new edition, to the mere announcement. Elaborate analysis and criticism would be out of place; and nothing remains to be done except to state in general terms that the author has bestowed on it that labor which such a work required in order to be made a representative of the existing state of surgical science and practice. Of the merits of the book as a guide to the "Science and Art of Surgery" it is not necessary for us to say much. Mr. Erichsen is one of

those enlightened surgeons of the present day, who regard an acquaintance with the manual part of surgery as only a portion of that knowledge which a surgeon should possess.—*British Medical Journal*, Jan. 2, 1869.

Thus the work bears in every feature a stamp of novelty and freshness which will commend it to those who are making its acquaintance for the first time, whilst those who have found it a safe guide and friend in former years will be able to refer to the new edition for the latest information upon any point of surgical controversy.—*London Lancet*, Jan. 23, 1869.

BY THE SAME AUTHOR. (*Just Issued.*)

ON RAILWAY, AND OTHER INJURIES OF THE NERVOUS SYSTEM. In small octavo volume. Extra cloth, \$1 00.

MILLER (JAMES).
Late Professor of Surgery in the University of Edinburgh, &c.

PRINCIPLES OF SURGERY. Fourth American, from the third and revised Edinburgh edition. In one large and very beautiful volume of 700 pages, with two hundred and forty illustrations on wood, extra cloth. \$3 75.

BY THE SAME AUTHOR.

THE PRACTICE OF SURGERY. Fourth American, from the last Edinburgh edition. Revised by the American editor. Illustrated by three hundred and sixty-four engravings on wood. In one large octavo volume of nearly 700 pages, extra cloth. \$3 75.

It is seldom that two volumes have ever made so profound an impression in so short a time as the "Principles" and the "Practice" of Surgery by Mr. Miller, or so richly merited the reputation they have

acquired. The author is an eminently sensible, practical, and well-informed man, who knows exactly what he is talking about and exactly how to talk it.—*Kentucky Medical Recorder*.

PIRRIE (WILLIAM), F. R. S. E.,
Professor of Surgery in the University of Aberdeen.

THE PRINCIPLES AND PRACTICE OF SURGERY. Edited by JOHN NEILL, M. D., Professor of Surgery in the Penna. Medical College, Surgeon to the Pennsylvania Hospital, &c. In one very handsome octavo volume of 780 pages, with 316 illustrations, extra cloth. \$3 75.

SARGENT (F. W.), M. D.

ON BANDAGING AND OTHER OPERATIONS OF MINOR SURGERY. New edition, with an additional chapter on Military Surgery. One handsome royal 12mo. volume, of nearly 400 pages, with 184 wood-cuts. Extra cloth, \$1 75.

Exceedingly convenient and valuable to all members of the profession.—*Chicago Medical Examiner*, May, 1862.

The very best manual of Minor Surgery we have seen.—*Buffalo Medical Journal*.

We cordially commend this volume as one which the medical student should most closely study; and to the surgeon in practice it must prove itself instructive on many points which he may have forgotten.—*Brit. Am. Journal*, May, 1862.

DRUITT (ROBERT), M.R.C.S. &c.

THE PRINCIPLES AND PRACTICE OF MODERN SURGERY.

A new and revised American, from the eighth enlarged and improved London edition. Illustrated with four hundred and thirty-two wood-engravings. In one very handsome octavo volume, of nearly 700 large and closely printed pages. Extra cloth, \$4 00; leather, \$5 00.

All that the surgical student or practitioner could desire.—*Dublin Quarterly Journal*.

It is a most admirable book. We do not know when we have examined one with more pleasure.—*Boston Med. and Surg. Journal*.

In Mr. Drutt's book, though containing only some seven hundred pages, both the principles and the practice of surgery are treated, and so clearly and perspicuously, as to elucidate every important topic. The fact that twelve editions have already been called for, in these days of active competition, would of itself show it to possess marked superiority. We have examined the book most thoroughly, and can say that this success is well merited. His book, moreover, possesses the inestimable advantages of having the subjects perfectly well arranged and classified, and of being written in a style at once clear and succinct.—*Am. Journal of Med. Sciences*.

Whether we view Drutt's Surgery as a guide to operative procedures, or as representing the latest

theoretical surgical opinions, no work that we are at present acquainted with can at all compare with it. It is a compendium of surgical theory if we may use the word, and practice in itself, and well deserves the estimate placed upon it.—*Brit. Am. Journal*.

Thus enlarged and improved, it will continue to rank among our best text-books on elementary surgery.—*Columbus Rev. of Med. and Surg.*

We must close this brief notice of an admirable work by recommending it to the earnest attention of every medical student.—*Charleston Medical Journal and Review*.

A text-book which the general voice of the profession in both England and America has commended as one of the most admirable "manuals," or, "*caie mecum*," as its English title runs, which can be placed in the hands of the student. The merits of Drutt's Surgery are too well known to every one to need any further eulogium from us.—*Nashville Med. Journal*.

HAMILTON (FRANK H.), M.D.,

Professor of Fractures and Dislocations, &c. in Bellevue Hosp. Med. College, New York.

A PRACTICAL TREATISE ON FRACTURES AND DISLOCATIONS. Third edition, thoroughly revised. In one large and handsome octavo volume of 777 pages, with 294 illustrations, extra cloth, \$5 75. (Just Issued.)

In fulness of detail, simplicity of arrangement, and accuracy of description, this work stands unrivalled. So far as we know, no other work on the subject in the English language can be compared with it. While congratulating our trans-Atlantic brethren on the European reputation which Dr. Hamilton, along with many other American surgeons, has attained, we also may be proud that, in the *mother tongue*, a classical work has been produced which need not fear comparison with the standard treatises of any other nation.—*Edinburgh Med. Journal*, Dec. 1866.

The credit of giving to the profession the only complete practical treatise on fractures and dislocations in our language during the present century, belongs to the author of the work before us, a distinguished

American professor of surgery; and his book adds one more to the list of excellent practical works which have emanated from his country, notices of which have appeared from time to time in our columns during the last few months.—*London Lancet*, Dec. 15, 1866.

These additions make the work much more valuable, and it must be accepted as the most complete monograph on the subject, certainly in our own, if not even in any other language.—*American Journal Med. Sciences*, Jan. 1867.

This is the most complete treatise on the subject in the English language.—*Ranking's Abstract*, Jan. 1867.

A mirror of all that is valuable in modern surgery.—*Richmond Med. Journal*, Nov. 1866.

CURLING (T. B.), F.R.S.

Surgeon to the London Hospital, President of the Hunterian Society, &c.

A PRACTICAL TREATISE ON DISEASES OF THE TESTIS, SPERMATIC CORD, AND SCROTUM. Second American, from the second and enlarged English edition. In one handsome octavo volume, extra cloth, with numerous illustrations. pp. 420. \$2 00.

BRODIE'S CLINICAL LECTURES ON SURGERY. 1 vol. 8vo., 350 pp.; cloth, \$1 25.

COOPER'S LECTURES ON THE PRINCIPLES AND PRACTICE OF SURGERY. In one very large octavo volume, extra cloth, of 750 pages. \$2 00.

GIBSON'S INSTITUTES AND PRACTICE OF SURGERY. Eighth edition, improved and altered. With thirty-four plates. In two handsome octavo volumes, about 1000 pp.; leather, raised bands. \$6 50.

MACKENZIE ON DISEASES AND INJURIES OF THE EYE. 1 vol. 8vo., 1027 pp., extra cloth. \$6.

ASHTON (T. J.).

ON THE DISEASES, INJURIES, AND MALFORMATIONS OF THE RECTUM AND ANUS; with remarks on Habitual Constipation. Second American, from the fourth and enlarged London edition. With handsome illustrations. In one very beautifully printed octavo volume of about 300 pages. \$3 25. (Just Issued.)

We can recommend this volume of Mr. Ashton's in the strongest terms, as containing all the latest details of the pathology and treatment of diseases connected with the rectum.—*Canada Med. Journ.*, March, 1866.

One of the most valuable special treatises that the physician and surgeon can have in his library.—*Chicago Medical Examiner*, Jan. 1866.

The short period which has elapsed since the appearance of the former American reprint, and the numerous editions published in England, are the best arguments we can offer of the merits, and of the usefulness of any commendation on our part of a book already so favorably known to our readers.—*Boston Med. and Surg. Journal*, Jan. 25, 1866.

MORLAND (W. W.), M.D.

DISEASES OF THE URINARY ORGANS; a Compendium of their Diagnosis, Pathology, and Treatment. With illustrations. In one large and handsome octavo volume of about 600 pages, extra cloth. \$3 50.

WELLS (J. SOELBERG).*Professor of Ophthalmology in King's College Hospital, &c.*

A TREATISE ON DISEASES OF THE EYE. First American Edition, with additions; illustrated with 216 engravings on wood, and six colored plates. Together with selections from the Test-types of Jaeger and Snellen. In one large and very handsome octavo volume of about 750 pages: extra cloth, \$5 00; leather, \$6 00. (Just Ready.)

A work has long been wanting which should represent adequately and completely the present aspect of British Ophthalmology, and this want it has been the aim of Mr. Wells to supply. The favorable reception of his volume by the medical press is a guarantee that he has succeeded in his undertaking, and in reproducing the work in this country every effort has been made to render it in every way suited to the wants of the American practitioner. Such additions as seemed desirable have been introduced by the editor, Dr. I. Minis Hays, and the number of illustrations has been more than doubled. The importance of test-types as an aid to diagnosis is so universally acknowledged at the present day that it seemed essential to the completeness of the work that they should be added, and as the author recommends the use of those both of Jaeger and of Snellen for different purposes, selections have been made from each, so that the practitioner may have at command all the assistance necessary. The work is thus presented as in every way fitted to merit the confidence of the American profession.

His chapters are eminently readable. His style is clear and flowing. He can be short without over-condensing, and accurate without hair splitting. These merits appear in a remarkable degree when he comes to treat of the more abstruse departments of his subject, and contrast favorably with the labored obscurity which mars the writings of some greater authorities in the same line. We congratulate Mr. Wells upon the success with which he has fulfilled his ideal, as

represented in the preface, in producing "an English treatise on the diseases of the eye, which should embrace the modern doctrines and practice of the British and Foreign Schools of Ophthalmology." The new school of Ophthalmology may also be congratulated in having found an exponent who is neither a bigoted partisan of everything new, nor a scold at everything old.—*Glasgow Med. Journal*, May, 1889.

TOYNBEE (JOSEPH), F. R. S.,*Aural Surgeon to and Lecturer on Surgery at St. Mary's Hospital.*

THE DISEASES OF THE EAR: their Nature, Diagnosis, and Treatment. With one hundred engravings on wood. Second American edition. In one very handsomely printed octavo volume of 440 pages; extra cloth, \$4.

The appearance of a volume of Mr. Toynbee's, therefore, in which the subject of aural disease is treated in the most scientific manner, and our knowledge in respect to it placed fully on a par with that which we possess respecting most other organs of the body, is a matter for sincere congratulation. We may reasonably hope that henceforth the subject of this treatise will cease to be among the *apropria* of medical science.—*London Medical Review*.

The work, as was stated at the outset of our notice, is a model of its kind, and every page and paragraph of it are worthy of the most thorough study. Considered all in all—as an original work, well written, philosophically elaborated, and happily illustrated with cases and drawings—it is by far the ablest monograph that has ever appeared on the anatomy and diseases of the ear, and one of the most valuable contributions to the art and science of surgery in the nineteenth century.—*N. Am. Med.-Chirurg. Review*.

LAURENCE (JOHN Z.), F. R. C. S.,*Editor of the Ophthalmic Review, &c.*

A HANDY-BOOK OF OPHTHALMIC SURGERY, for the use of Practitioners. Second Edition, revised and enlarged. With numerous illustrations. In one very handsome octavo volume, extra cloth, \$3 00. (Now Ready.)

No book on ophthalmic surgery was more needed. Designed, as it is, for the wants of the busy practitioner, it is the *ne plus ultra* of perfection. It epitomizes all the diseases incidental to the eye in a clear and masterly manner, not only enabling the practitioner readily to diagnose each variety of disease, but affording him the more important assistance of proper treatment. Altogether this is a work which ought certainly to be in the hands of every general practitioner.—*Dublin Med. Press and Circular*, Sept. 12, '66.

We cordially recommend this book to the notice of our readers, as containing an excellent outline of modern ophthalmic surgery.—*British Med. Journal*, October 13, 1866.

Not only, as its modest title suggests, a "Handy-Book" of Ophthalmic Surgery, but an excellent and well-digested *résumé* of all that is of practical value in the specialty.—*New York Medical Journal*, November, 1866.

This object the authors have accomplished in a highly satisfactory manner, and we know no work we can more highly recommend to the "busy practitioner" who wishes to make himself acquainted with the recent improvements in ophthalmic science. Such a work as this was much wanted at this time, and this want Messrs. Laurence and Moon have now well supplied.—*Am. Journal Med. Sciences*, Jan. 1867.

LAWSON (GEORGE), F. R. C. S., Engl.,*Assistant Surgeon to the Royal London Ophthalmic Hospital, Moorfields, &c.*

INJURIES OF THE EYE, ORBIT, AND EYELIDS: their Immediate and Remote Effects. With about one hundred illustrations. In one very handsome octavo volume, extra cloth, \$3 50. (Now Ready.)

This work will be found eminently fitted for the general practitioner. In cases of functional or structural diseases of the eye, the physician who has not made ophthalmic surgery a special study can, in most instances, refer a patient to some competent practitioner. Cases of injury, however, supervene suddenly and usually require prompt assistance, and a work devoted especially to them cannot but prove essentially useful to those who may at any moment be called upon to treat such accidents. The present volume, as the work of a gentleman of large experience, may be considered as eminently worthy of confidence for reference in all such emergencies.

It is an admirable practical book in the highest and best sense of the phrase.—*London Medical Times and Gazette*, May 18, 1867.

WALES (PHILIP S.), M. D., Surgeon U. S. N.

MECHANICAL THERAPEUTICS: a Practical Treatise on Surgical Apparatus, Appliances, and Elementary Operations: embracing Minor Surgery, Bandaging, Orthopraxy, and the Treatment of Fractures and Dislocations. With six hundred and forty-two illustrations on wood. In one large and handsome octavo volume of about 700 pages: extra cloth, \$5 75; leather, \$6 75. (*Just Issued.*)

A Naval Medical Board directed to examine and report upon the merits of this volume, officially states that "it should in our opinion become a standard work in the hands of every naval surgeon;" and its adoption for use in both the Army and Navy of the United States is sufficient guarantee of its adaptation to the needs of every-day practice.

The title of this book will give a reasonably good idea of its scope, but its merits can only be appreciated by a careful perusal of its text. No one who undertakes such a task will have any reason to complain that the author has not performed his duty, and has not taken every pains to present every subject in a clear, common-sense, and practical light. It is a unique specimen of literature in its way, in that, treating upon such a variety of subjects, it is as a whole so completely up to the wants of the student and the general practitioner. We have never seen any work of its kind that can compete with it in real utility and extensive adaptability. Dr. Wales perfectly understands what may naturally be required of him in the premises, and in the work before us has bridged over a very wide gap which has always heretofore existed between the first rudiments of surgery and practical surgery proper. He has emphatically given us a comprehensive work for the beginner; and when we say of his labors, that in their particular sphere they leave nothing to be desired, we assert a great deal to recommend the book to the attention of those specially concerned. In conclusion, we would state, at the risk of reiteration, that this is the most comprehensive book on the subject that we have seen; is the best that can be placed in the hands of the stu-

dent in need of a first book on surgery, and the most useful that can be named for such general practitioners who, without any special pretensions to surgery, are occasionally liable to treat surgical cases.—*N. Y. Med. Record*, March 2, 1868.

It is certainly the most complete and thorough work of its kind in the English language. Students and young practitioners of surgery will find it invaluable. It will prove especially useful to inexperienced country practitioners, who are continually required to take charge of surgical cases, under circumstances precluding them from the aid of experienced surgeons.—*Pacific Med. and Surg. Journal*, Feb. 1868.

The title of the above work is sufficiently indicative of its contents. We have not seen for a long time (in the English language) a treatise equal to this in extent, nor one which is better adapted to the wants of the general student and practitioner. It is not to the surgeon alone that this book belongs; the physician has frequent opportunities to fill an emergency by such knowledge as is here given. Every practitioner should make purchase of such a book—it will last him his lifetime.—*St. Louis Med. Reporter*, Feb. 1868.

BIGELOW (HENRY J.), M. D.,

Professor of Surgery in the Massachusetts Med. College.

ON THE MECHANISM OF DISLOCATION AND FRACTURE OF THE HIP. With the Reduction of the Dislocation by the Flexion Method. With numerous original illustrations. In one very handsome octavo volume. Cloth. \$2 50. (*Now Ready.*)

We cannot too highly praise this book as the work of an accomplished and scientific surgeon. We do not hesitate to say that he has done much to clear up the obscurities connected with the mechanism of dislocation of the hip-joint, and he has laid down most valuable practical rules for the easy and most successful management of these injuries. The mono-

graph is largely illustrated with exquisitely executed woodcuts, after photographs, which help to elucidate the admirable subject-matter of the text. We cordially commend the "Hip," by Dr. Bigelow, to the attention of surgeons.—*Dublin Quarterly Journal of Medical Science*, Feb. 1870.

THOMPSON (SIR HENRY),

Surgeon and Professor of Clinical Surgery to University College Hospital.

LECTURES ON DISEASES OF THE URINARY ORGANS. With illustrations on wood. In one neat octavo volume, extra cloth. \$2 25. (*Now Ready.*)

These lectures stand the severe test. They are instructive without being tedious, and simple without being diffuse; and they include many of those practical hints so useful for the student, and even more valuable to the young practitioner.—*Edinburgh Med. Journal*, April, 1869.

Very few words of ours are necessary to recommend these lectures to the profession. There is no subject

on which Sir Henry Thompson speaks with more authority than that in which he has specially gathered his laurels; in addition to this, the conversational style of instruction, which is retained in these printed lectures, gives them an attractiveness which a systematic treatise can never possess.—*London Medical Times and Gazette*, April 24, 1869.

BY THE SAME AUTHOR. (Nearly Ready.)

ON THE PATHOLOGY AND TREATMENT OF STRICTURE OF THE URETHRA AND URINARY FISTULÆ. With plates and wood-cuts. From the third and revised English edition. In one very handsome octavo volume, extra cloth, \$3 50. (*Now Ready.*)

This classical work has so long been recognized as a standard authority on its perplexing subjects that it should be rendered accessible to the American profession. Having enjoyed the advantage of a revision at the hands of the author within a few months, it will be found to present his latest views and to be on a level with the most recent advances of surgical science.

With a work accepted as the authority upon the subjects of which it treats, an extended notice would be a work of supererogation. The simple announcement of another edition of a work so well and favor-

ably known by the profession as this before us, must create a demand for it from those who would keep themselves well up in this department of surgery.—*St. Louis Med. Archives*, Feb. 1870.

TAYLOR (ALFRED S.), M.D.,*Lecturer on Med. Jurisp. and Chemistry in Guy's Hospital.***MEDICAL JURISPRUDENCE.** Sixth American, from the eighth and revised London edition. With Notes and References to American Decisions, by CLERMONT B. PENROSE, of the Philadelphia Bar. In one large octavo volume of 776 pages, extra cloth, \$4 50; leather, \$5 50. (*Just Issued.*)

Considerable additions have been made by the editor to this edition, comprising some important sections from the author's larger work, "The Principles and Practice of Medical Jurisprudence," as well as references to American law and practice. The notes of the former editor, Dr. Hartschorne, have likewise been retained, and the whole is presented as fully worthy to maintain the distinguished position which the work has acquired as a leading text-book and authority on the subject.

A new edition of a work acknowledged as a standard authority everywhere within the range of the English language. Considering the new matter introduced, on trichiniasis and other subjects, and the plates representing the crystals of poisons, etc., it may fairly be regarded as the most compact, comprehensive, and practical work on medical jurisprudence which has issued from the press, and the one best fitted for students.—*Pacific Med. and Surg. Journal*, Feb. 1867.

The sixth edition of this popular work comes to us in charge of a new editor, Mr. Penrose, of the Philadelphia bar, who has done much to render it useful, not only to the medical practitioners of this country, but to those of his own profession. Wisely retaining the references of the former American editor, Dr. Hartschorne, he has added many valuable notes of his own. The reputation of Dr. Taylor's work is so well established, that it needs no recommendation. He is now the highest living authority on all matters con-

nected with forensic medicine, and every successive edition of his valuable work gives fresh assurance to his many admirers that he will continue to maintain his well-earned position. No one should, in fact, be without a text-book on the subject, as he does not know but that his next case may create for him an emergency for its use. To those who are not the fortunate possessors of a reliable, readable, interesting, and thoroughly practical work upon the subject, we would earnestly recommend this, as forming the best groundwork for all their future studies of the more elaborate treatises.—*New York Medical Record*, Feb. 15, 1867.

The present edition of this valuable manual is a great improvement on those which have preceded it. It makes thus by far the best guide-book in this department of medicine for students and the general practitioner in our language.—*Boston Med. and Surg. Journal*, Dec. 27, 1866.

WINSLOW (FORBES), M.D., D.C.L., &c.**ON OBSCURE DISEASES OF THE BRAIN AND PSYCHIC DISORDERS OF THE MIND;** their incipient Symptoms, Pathology, Diagnosis, Treatment, and Prognosis. Second American, from the third and revised English edition. In one handsome octavo volume of nearly 600 pages, extra cloth. \$4 25. (*Just Issued.*)

Of the merits of Dr. Winslow's treatise the profession has sufficiently judged. It has taken its place in the front rank of the works upon the special department of practical medicine to which it pertains.—*Cincinnati Journal of Medicine*, March, 1866.

It is an interesting volume that will amply repay for a careful perusal by all intelligent readers.—*Chicago Med. Examiner*, Feb. 1866.

A work which, like the present, will largely aid the practitioner in recognizing and arresting the first insidious advances of cerebral and mental disease, is one of immense practical value, and demands earnest attention and diligent study on the part of all who have embraced the medical profession, and have

thereby undertaken responsibilities in which the welfare and happiness of individuals and families are largely involved. We shall therefore close this brief and necessarily very imperfect notice of Dr. Winslow's great and classical work by expressing our conviction that it is long since so important and beautifully written a volume has issued from the British medical press.—*Dublin Medical Press*.

It is the most interesting as well as valuable book that we have seen for a long time. It is truly fascinating.—*Am. Jour. Med. Sciences*.

Dr. Winslow's work will undoubtedly occupy an unique position in the medico-psychological literature of this country.—*London Med. Review*.

LEA (HENRY C.).**SUPERSTITION AND FORCE: ESSAYS ON THE WAGER OF LAW, THE WAGER OF BATTLE, THE ORDEAL, AND TORTURE.** Second Edition, Enlarged. In one handsome volume royal 12mo. of nearly 500 pages; extra cloth, \$2 75. (*Just Ready.*)

The copious collection of facts by which Mr. Lea has illustrated his subject shows in the fullest manner the constant conflict and varying success, the advances and defeats, by which the progress of humane legislation has been and is still marked. This work fills up with the fullest exemplification and detail the wise remarks which we have quoted above. As a book of ready reference on the subject it is of the highest value.—*Westminster Review*, Oct. 1867.

When—half in spite of himself, as it appears—he sketches a scene or character in the history of legalized error and cruelty, he betrays so artistic a feeling, and

a humor so fine and good, that he makes us regret it was not within his intent, as it was certainly within his power, to render the whole of his thorough work more popular in manner.—*Atlantic Monthly*, Feb. '67.

This is a book of extraordinary research. Mr. Lea has entered into his subject *con amore*; and a more striking record of the cruel superstitions of our unhappy Middle Ages could not possibly have been compiled. . . . As a work of curious inquiry on certain outlying points of obsolete law, "Superstition and Force" is one of the most remarkable books we have met with.—*London Athenæum*, Nov. 3, 1866.

BY THE SAME AUTHOR. (Just Issued.)**STUDIES IN CHURCH HISTORY—THE RISE OF THE TEMPORAL POWER—BENEFIT OF CLERGY—EXCOMMUNICATION.** In one large royal 12mo. volume of 516 pp. extra cloth. \$2 75.

Altogether, the book is a useful addition to the popular literature of a most important and too little known department of mediæval history.—*London Saturday Review*, Feb. 26, 1870.

They are careful studies by a thorough scholar in the most interesting of all historical fields, made without passion or prejudice, and recorded with honesty. The whole volume is of the deepest interest; the style is masculine and animated, and great skill

is shown in weaving in anecdote and picturesque stories, without impairing the flow of the relation or the proper dignity of the composition.—*Hartford Courant*, Jan. 22, 1870.

We recommend the book as a highly instructive discussion of matters which are always of interest to scholars, and which are just now clothed with a special importance.—*N. Y. Nation*, Feb. 3, 1870.

INDEX TO CATALOGUE.

	PAGE		PAGE
Allen's Dissector and Practical Anatomist	6	Lea's Superstition and Force	31
American Journal of the Medical Sciences	1	Lea's Studies in Church History	31
Abstract, Half-Yearly, of the Med. Sciences	3	Lallemand and Wilson on Spermatorrhœa	19
Anatomical Atlas, by Smith and Horner	6	La Roche on Yellow Fever	14
Ashton on the Rectum and Anus	28	La Roche on Pneumonia, &c.	14
Attfield's Chemistry	11	Laurence and Moon's Ophthalmic Surgery	29
Ashwell on Diseases of Females	22	Lawson on the Eye	29
Basham on Renal Diseases	18	Laycock on Medical Observation	16
Brinton on the Stomach	16	Lehmann's Physiological Chemistry, 2 vols.	9
Bigelow on the Hip	30	Lehmann's Chemical Physiology	9
Barclay's Medical Diagnosis	14	Ludlow's Manual of Examinations	5
Barlow's Practice of Medicine	15	Lyons on Fever	18
Bowman's (John E.) Practical Chemistry	10	MacClise's Surgical Anatomy	7
Bowman's (John E.) Medical Chemistry	10	Malgaigne's Operative Surgery, by Brittan	26
Brande & Taylor's Chemistry	10	Marshall's Physiology	8
Brodie's Clinical Lectures on Surgery	28	Mayne's Dispensatory and Formulary	13
Brown on the Clinical Diseases of Women	22	Mackenzie on Diseases of the Eye	28
Buckler on Bronchitis	14	Medical News and Library	2
Bucknill and Tuke on Insanity	14	Meigs's Obstetrics, the Science and the Art	25
Bumstead on Venereal	19	Meigs's Lectures on Diseases of Women	23
Bumstead and Cullerier's Atlas of Venereal	19	Meigs on Puerperal Fever	23
Carpenter's Human Physiology	8	Miller's System of Obstetrics	24
Carpenter's Comparative Physiology	8	Miller's Practice of Surgery	27
Carpenter on the Use and Abuse of Alcohol	13	Miller's Principles of Surgery	27
Carson's Synopsis of Materia Medica	13	Montgomery on Pregnancy	24
Chambers on the Indigestions	17	Morland on Urinary Organs	25
Christison and Griffith's Dispensatory	13	Morland on Uræmia	18
Churchill's System of Midwifery	25	Neill and Smith's Compendium of Med. Science	5
Churchill on Diseases of Females	22	Neligan's Atlas of Diseases of the Skin	20
Churchill on Puerperal Fever	22	Neligan on Diseases of the Skin	20
Clymer on Fevers	17	Odling's Practical Chemistry	10
Condie on Diseases of Children	21	Pavy on Digestion	17
Cooper's (B. B.) Lectures on Surgery	28	Prize Essays on Consumption	12
Cullerier's Atlas of Venereal Diseases	19	Parrish's Practical Pharmacy	14
Curling on Diseases of the Testis	29	Pirrie's System of Surgery	17
Cyclopedia of Practical Medicine	15	Pereira's Mat. Medica and Therapeutics, abridged	23
Dalton's Human Physiology	9	Quain and Sharpey's Anatomy, by Leidy	6
De Jongh on Cod-Liver Oil	13	Ranking's Abstract	3
Dewees's System of Midwifery	24	Roberts on Urinary Diseases	18
Dewees on Diseases of Females	22	Ramsbotham on Parturition	25
Dewees on Diseases of Children	21	Rigby on Female Diseases	22
Dickson's Practice of Medicine	16	Rigby's Midwifery	24
Druitt's Modern Surgery	28	Rokitansky's Pathological Anatomy	14
Dunglison's Medical Dictionary	4	Royle's Materia Medica and Therapeutics	13
Dunglison's Human Physiology	9	Salter on Asthma	18
Dunglison on New Remedies	13	Swayne's Obstetric Aphorisms	25
Ellis's Medical Formulary, by Smith	13	Sargent's Minor Surgery	27
Erichsen's System of Surgery	27	Sharpey and Quain's Anatomy, by Leidy	6
Erichsen on Nervous Injuries	27	Simon's General Pathology	14
Flint on Respiratory Organs	17	Simpson on Females	23
Flint on the Heart	17	Skey's Operative Surgery	26
Flint's Practice of Medicine	15	Slade on Diphtheria	18
Fownes's Elementary Chemistry	11	Smith (J. L.) on Children	21
Fuller on the Lungs, &c.	16	Smith (H. H.) and Horner's Anatomical Atlas	6
Gibson's Surgery	28	Smith (Edward) on Consumption	18
Gluge's Pathological Histology, by Leidy	14	Smith on Wasting Diseases of Children	21
Graham's Elements of Chemistry	10	Solly on Anatomy and Diseases of the Brain	14
Gray's Anatomy	6	Stillé's Therapeutics	12
Griffith's (R. E.) Universal Formulary	12	Tanner's Manual of Clinical Medicine	5
Gross on Urinary Organs	26	Tanner on Pregnancy	24
Gross on Foreign Bodies in Air-Passages	26	Taylor's Medical Jurisprudence	31
Gross's Principles and Practice of Surgery	26	Thomas on Diseases of Females	22
Gross's Pathological Anatomy	14	Thompson on Urinary Organs	30
Hartshorne's Essentials of Medicine	16	Thompson on Stricture	30
Hartshorne's Conspectus of the Medical Sciences	5	Todd and Bowman's Physiological Anatomy	9
Hartshorne's Anatomy and Physiology	6	Todd on Acute Diseases	17
Habershon on Alimentary Canal	17	Toynbee on the Ear	29
Hamilton on Dislocations and Fractures	28	Wales on Surgical Operations	30
Harrison on the Nervous System	14	Walshe on the Heart	18
Heath's Practical Anatomy	7	Watson's Practice of Physic	16
Hoblyn's Medical Dictionary	4	Wells on the Eye	29
Hodge on Women	23	West on Diseases of Females	23
Hodge's Obstetrics	24	West on Diseases of Children	21
Hodge's Practical Dissections	7	West on Ulceration of Os Uteri	23
Holland's Medical Notes and Reflections	15	What to Observe in Medical Cases	16
Horner's Anatomy and Histology	7	Williams's Principles of Medicine	14
Hudson on Fevers	18	Wilson's Human Anatomy	7
Hill on Venereal Diseases	19	Wilson's Dissector	7
Hillier's Handbook of Skin Diseases	20	Wilson on Diseases of the Skin	20
Jones and Sieveking's Pathological Anatomy	14	Wilson's Plates on Diseases of the Skin	20
Jones (C. Handfield) on Nervous Disorders	18	Wilson's Handbook of Cutaneous Medicine	19
Kirkes's Physiology	5	Wilson on Spermatorrhœa	20
Knapp's Chemical Technology	11	Winslow on Brain and Mind	31

Accession no.

Author **Dunglison, R.**
Medical lexicon.

19th cent

Call no. **R121**
D85
1868

